SELF-ABSORPTION OF HANKEL SYSTEMS ON MONOIDS – a seemingly universal property

YONG HAN, YANQI QIU, AND ZIPENG WANG

ABSTRACT. Given any cancellative monoid \mathcal{M} , we study the Hankel system determined by its multiplication table (see (1.1)). We prove that the Hankel system admits self-absorption property (Definition 1) provided that the monoid \mathcal{M} has the local algebraic structure:

$$(ax = by, cx = dy, az = bw \text{ in } \mathcal{M}) \Longrightarrow (cz = dw \text{ in } \mathcal{M}).$$

Our result holds for all group-embeddable monoids and goes beyond. In particular, it works for all cancellative Abelian monoids and most common non-Abelian cancellative monoids such as

$$\mathrm{SL}_d(\mathbb{N}) := \{ [a_{ij}]_{1 \le i,j \le d} \in \mathrm{SL}_d(\mathbb{Z}) | a_{ij} \in \mathbb{N} \}.$$

The Hankel system determined by the multiplication table of a monoid is further generalized to that determined by level sets of any abstract two-variable map (see (5.2)). We introduce an algebraic notion of lunar maps (Definition 4) and establish a stronger hereditary self-absorption property (Definition 5) for the corresponding generalized Hankel systems. As a consequence, we prove the self-absorption property for arbitrary spatial compression of the regular representation system $\{\lambda_G(g)\}_{g\in G}$ of any discrete group *G*, as well as the Hankel system $\{\Gamma_\ell^{\Phi}\}$ determined by the level sets of any rational map of the form $\Phi(x, y) = ax^m + by^n$ with $a, b, m, n \in \mathbb{Z}^*$:

$$\Gamma^{\Phi}_{\ell}(x,y) = \mathbb{1}(ax^m + by^n = \ell), \quad x, y \in \mathbb{N}^*, \, \ell \in \Phi(\mathbb{N}^* \times \mathbb{N}^*).$$

Our result is already new in the classical setting of the additive monoid \mathbb{N} of non-negative integers. The self-absorption property is applied to the study of completely bounded Fourier multipliers between Hardy spaces. In the case of endpoint spaces, we obtain the full characterizations:

$$(H^1, BMOA)_{cb} = BMOA$$
 and $(H^1, H^2)_{cb} = (H^2, BMOA)_{cb} = BMOA^{(2)}$.

Then, using complex interpolation, we obtain, for any $1 \le p \le 2 \le q < \infty$,

$$\mathrm{BMOA}^{(u)} \subset (H^p, H^q)_{cb} \quad \text{with } \frac{1}{u} = \frac{1}{p} - \frac{1}{q}.$$

Here BMOA^(*u*) is a new Banach space of analytic functions on the unit circle \mathbb{T} with

$$\left\|\varphi\right\|_{\mathrm{BMOA}^{(u)}} \approx \left|\widehat{\varphi}(0)\right| + \sup_{n \ge 1} \left\{\sum_{k=1}^{\infty} \left(\sum_{kn \le j < (k+1)n} |\widehat{\varphi}(j)|^{u}\right)^{2}\right\}^{1/2u}$$

Further applications are: i) exact complete bounded norm of the Carleman embedding in any dimension; ii) mixed Fourier-Schur multiplier inequalities with critical exponent 4/3; iii) failure of hyper-complete-contractivity for the Poisson semigroup.

CONTENTS

2 3

Foreword: What does absorption mean and why it is natural ? 1. Introduction

2020 Mathematics Subject Classification. Primary 46L07, 47B35, 47A80; Secondary 42A45, 30H10, 43A20.

Key words and phrases. self-absorption property; tensor products; Hankel systems on monoids; complete bounded Fourier multipliers; Hardy and BMOA spaces.

2.	The simplest SAP monoid: an illustration	10
3.	Self-absorption property: general results	16
4.	Hankel systems on monoids	20
5.	Hankel systems of lunar maps	21
6.	Applications	28
Re	ferences	43

FOREWORD: WHAT DOES ABSORPTION MEAN AND WHY IT IS NATURAL ?

I. Fell's absorption in group representation theory. The well-known Fell's aborption principle states that, the left regular representation λ_G of any discrete countable group *G* absorbs all its unitary representations. In other words, for any unitary representation $\pi : G \to \mathcal{U}(H)$, there exists a unitary operator $U \in \mathcal{U}(\ell^2(G) \otimes_2 H)$ such that

$$\lambda_G(g) \otimes \pi(g) = U[\lambda_G(g) \otimes Id_H]U^{-1}, \quad \forall g \in G.$$

This means that all operators $\lambda_G(g) \otimes \pi(g)$ can be simultaneously block-diagonalized, with all block-diagonal entries being $\lambda_G(g)$.

II. A new absorption in operator-space theory. We say that a family of Hilbert-space operators $\{x_i : i \in I\} \subset B(H)$ contractively absorbs another family $\{y_i : i \in I\} \subset B(K)$ if there exist two contractive operators U, V on $H \otimes_2 K$, such that all operators $x_i \otimes y_i$ simultaneously factor through block-diagonalized operators as:

(0.1)
$$x_i \otimes y_i = U(x_i \otimes Id_K)V, \quad \forall i \in I$$

The main focus of this paper is the self-absorption property. Specifically, we consider the problem when the family $\{x_i : i \in I\}$ contractively absorbs itself. In fact, from the proofs of our main results (Theorem A in §1 and Theorem B in §5), we obtain (0.1) for very general Hankel systems with U and V even being partial isometries (see (5.13) and (5.18)).

The absorption property mentioned above is an algebraic property of the Hankel systems. However, for simplicity, we shall state our main results in the framework of the operator-space theory in a slightly weaker version. Indeed, the simultaneous factorization (0.1) implies the following contractivity inequalities:

(0.2)
$$\|\sum_{i} c_i \otimes x_i \otimes y_i\| \le \|\sum_{i} c_i \otimes x_i\|,$$

where $(c_i)_i$ is any finitely supported family in an arbitrary C^* -algebra. Conversely, the fundamental factorization/extension theorem in operator-space theory (consequence of the Stinespring and Arveson's dialation theorems) says that (0.2) is equivalent to the existence of a C^* -representation $\pi : B(H) \to B(\widehat{H})$ and two contractive operators $U, V \in B(\widehat{H}, H \otimes_2 K)$ with

$$x_i \otimes y_i = U \pi(x_i) V^*, \quad \forall i \in I.$$

The inequality (0.2) has close connections to harmonic analysis: it appears to be related to the well-known open problem of the existence of Nehari Theorem on higher dimensional torus (see Problem 1 in §1), and it will be applied to the study of completely bounded Fourier multipliers between Hardy spaces (see Section 6). For instance, we show that, within the framework of operator-space theory, it is surprising that Janson's hypercontractivity fails on Hardy spaces (see Corollary 6.17).

Finally, it is worth mentioning the following universal property for an arbitrary family of unit norm Boolean operators $\{x_i\}$ (see §3.2 for its definition) : under an appropriate positivity condition on the coefficients c_i , we always have (see Proposition 3.2)

$$\left\|\sum_{i}c_{i}\otimes x_{i}^{\otimes m}\right\|=\left\|\sum_{i}c_{i}\otimes x_{i}\right\|.$$

III. Relation with Graph-theory. For any family $\{x_i : i \in I\}$ of unit norm Boolean operators with disjoint supports (see §3.2), one can naturally associate an oriented colored graph \mathcal{G} . Similarly, an associated oriented colored graph $\widehat{\mathcal{G}}$ is constructed from $\{x_i \otimes x_i : i \in I\}$. In a subsequent paper, we show that, if all connected components of $\widehat{\mathcal{G}}$ (after certain natural operations) can be reduced to saturated subgraphs of \mathcal{G} , then $\{x_i : i \in I\}$ contractively absorbs itself. This realization allows us, on the one hand, to verify the self-absorption property for various specific examples of families of Boolean operators, and on the other hand, provides a conjectured equivalent graph-theory criterion of the self-aborption property in the context of Boolean operators.

1. INTRODUCTION

This paper is mainly devoted to the study of Hankel systems on abstract monoids. The key ingredients are the notion of self-absorption property (Definition 1), lunar monoids (Definition 2), SAP monoids (Definition 3) and the coupled foliations (Figure 1).

We begin by introducing the definition of the Hankel systems on monoids. Let \mathcal{M} be a cancellative monoid (not necessarily Abelian). The associated Hankel system $\{\Gamma_t^{\mathcal{M}}\}_{t \in \mathcal{M}} \subset B(\ell^2(\mathcal{M}))$ is a collection of operators defined as follows:

(1.1)
$$\Gamma_t^{\mathcal{M}}(s_1, s_2) = \mathbb{1}(s_1 s_2 = t), \quad s_1, s_2 \in \mathcal{M}$$

Here, we recall that a cancellative monoid \mathcal{M} refers to a semigroup with a unit element in which the implication $(abc = adc) \Longrightarrow (b = d)$ holds in \mathcal{M} . Note that \mathcal{M} is cancellative if and only if $\|\Gamma_t^{\mathcal{M}}\|_{B(\ell^2(\mathcal{M}))} = 1$ for all $t \in \mathcal{M}$.

For instance, if $\mathcal{M} = \mathbb{N}^d$ (resp. $\mathbb{N}^{(\infty)}$) is the free Abelian monoid of *d* generators (resp. countably infinite generators), then the operator $\Gamma_t^{\mathcal{M}}$ is unitarily equivalent, via Fourier transform, to an elementary small Hankel operator on the Hardy space of the *d*-dimensional torus \mathbb{T}^d (resp. infinite dimensional torus $\mathbb{T}^{\mathbb{N}}$). Our work on self-absorptioin property of the Hankel system on \mathbb{N}^d is closely related to the open problem of the existence of higher dimensional Nehari-Sarashon-Page Theorem, see Problem 1 and Problem 2 and the discussions there.

The Hankel operators or Hankel matrices associated with the system $\{\Gamma_t^{\mathcal{M}}\}_{t \in \mathcal{M}}$ are defined as follows:

(1.2)
$$\sum_{t\in\mathcal{M}}c_t\otimes\Gamma_t^{\mathcal{M}}=[c_{s_1s_2}]_{s_1,s_2\in\mathcal{M}},$$

where c_t is either in \mathbb{C} or belongs to a C^* -algebra. In the classical setting of the monoid \mathbb{N} , the study of Hankel operators is closely related to the theory of one-dimensional discrete time Gaussian processes (see [Pel03, Chapters 8 and 9]). In order to investigate Gaussian processes on abstract monoids, one needs to study positive definite Hankel matrices of the form (1.2) or similar forms, see [LM71, Sch77, GS03] for details.

In the case of free monoids (or more general trace monoids), Fliess [Fli74] studied the Hankel matrices of the form (1.2) with coefficients c_t belonging to an abstract semiring. These Hankel

matrices on free monoids also play a role in quantized calculus for non-commutatative geometry. Connes [Con94, Section 4.5] conjectured a non-commutative analogue of the well-known Kronecker theorem on the relation between finite rank Hankel matrices and rational functions. Duchamp and Reutenauer [DR97] confirmed Connes' conjecture and their proof is based on Hankel matrices on the free monoids (see [BR11, Thm. 2.1.6]). See also [Miy22] for similar phenomenon of free semicircular elements.

In this paper, the Hankel systems on monoids are further generalized to the non-classical ones determined by level sets of abstract two-variable maps. More precisely, given three sets $\mathcal{A}, \mathcal{X}, \mathcal{L}$ and a map $\Phi : \mathcal{A} \times \mathcal{X} \to \mathcal{L}$, we consider the non-classical Hankel system $\{\Gamma_{\ell}^{\Phi}\}_{\ell \in \mathcal{L}}$ with Γ_{ℓ}^{Φ} the matrices defined by

(1.3)
$$\Gamma^{\Phi}_{\ell}(a,x) = \mathbb{1}(\Phi(a,x) = \ell), \quad (a,x) \in \mathcal{A} \times \mathcal{X}$$

The associated Hankel matrices are then defined as

(1.4)
$$\sum_{\ell \in \mathcal{L}} c_{\ell} \otimes \Gamma_{\ell}^{\Phi} = [c_{\Phi(a,x)}]_{a \in \mathcal{A}, x \in \mathcal{X}},$$

where c_{ℓ} is either in \mathbb{C} or belongs to a C^* -algebra.

The consideration of such non-classical Hankel systems $\{\Gamma_{\ell}^{\Phi}\}_{\ell \in \mathcal{L}}$ is not merely a pure mathematical curiosity, but rather it is essential for proving the hereditary self-absorption property (Definition 5) for the Hankel system on the simplest monoid \mathbb{N} , for the big Hankel system on the higher dimensional torus \mathbb{T}^d or the infinite dimensional torus $\mathbb{T}^{\mathbb{N}}$ and for the regular representation system of any discrete group (see Corollaries 5.1, 5.2 and 5.3). In particular, our result implies a recent result of Katsoulis [Kat23, Cor. 5.3]. Note also that the Hankel matrices studied in [LM71] are associated to the monoids with an involution and they are of the form (1.4) for a suitable map Φ .

For simplicity, in the remaining part of the introduction, we mainly concentrate on the Hankel systems on monoids. The results concerning non-classical hankel systems will be given in §5.

I. Self-absorption property. The following definition is inspired by Pisier's operator Hilbert space *OH*. Here we recall that *OH* is the Hilbert space $\ell^2 = \ell^2(\mathbb{N})$ equipped with the unique *operator space structure* (abbrev. *o.s.s.*, see [Pis03, Chapters 1-5] and [Rua88, ER22] for the background of *o.s.s.*) such that for any orthonormal vectors $(v_i)_{i \in I}$ in *OH* and any finitely supported sequence $(a_i)_{i \in I}$ in any *C**-algebra *A*,

(1.5)
$$\left\|\sum_{i\in I}v_i\otimes a_i\right\|_{OH\otimes_{min}A} = \left\|\sum_{i\in I}a_i\otimes\overline{a}_i\right\|_{A\otimes_{min}\overline{A}}^{1/2},$$

where \otimes_{min} denotes the standard minimal tensor product and \overline{A} denotes the complex conjugate of the *C*^{*}-algebra *A* (see [Pis20, Section 2.3]).

Definition 1 (Self-absorption property). Given $\kappa \ge 1$. A family (also called a system) $\mathcal{F} = \{x_i\}_{i\in I}$ in a C^* -algebra A or in an operator space is said to have κ -self-absorption property (abbrev. κ -SAP) if the following inequalities hold for any finitely supported family $\{b_i\}_{i\in I}$ in any other C^* -algebra B:

(1.6)
$$\frac{1}{\kappa} \Big\| \sum_{i \in I} b_i \otimes x_i \Big\|_{B \otimes_{\min} A} \leq \Big\| \sum_{i \in I} b_i \otimes x_i \otimes \overline{x}_i \Big\|_{B \otimes_{\min} A \otimes_{\min} \overline{A}} \leq \kappa \Big\| \sum_{i \in I} b_i \otimes x_i \Big\|_{B \otimes_{\min} A}.$$

For simplicity, we denote 1-SAP just by SAP.

Definition 2 (Lunar monoids). We say that a monoid \mathcal{M} is a *lunar monoid* (or we say \mathcal{M} is lunar) if it is cancellative and satisfies the algebraic lunar condition :

(1.7)
$$(ax = by, cx = dy, az = bw \text{ in } \mathcal{M}) \Longrightarrow (cz = dw \text{ in } \mathcal{M}).$$

Our terminology "lunar condition" comes from the theory of group-embeddability of monoids ([Mal39, Mal40] and [Lam51]). Indeed, the condition (1.7) is the first-order one of a series of conditions (called lunar conditions there) in Jackson's thesis with his thesis supervisor Halperin (see [Bus71]).

Theorem A. For any lunar monoid \mathcal{M} , the Hankel system $\{\Gamma_t^{\mathcal{M}}\}_{t\in\mathcal{M}}$ has SAP.

Theorem A follows from a stronger result in Theorem 4.1, which states that any lunar monoid induces a Hankel system with the stronger *hereditary SAP* (see Definition 5). With extra efforts, our formalism can be adapted to operators on ℓ^p -spaces or on more general symmetric sequence-spaces.

Note that the lunar condition of a monoid is a *local algebraic condition*: to check whether a monoid is lunar or not, it suffices to study the configurations of all 4×4 blocks in its multiplication table. Theorem A means that, such local algebraic condition of a monoid implies a global analytic property –the SAP–for the associated Hankel system. Furthermore, we conjecture that, the above local algebraic condition is equivalent to the global analytic one. See Problem 3 and the more tractable Problem 6, Problem 7 for related discussions.

It can be checked directly that the lunar condition holds true for all group-embeddable monoids. Therefore, we obtain for instance the SAP for the Hankel systems associated with the following monoids: the free monoids (submonoids of free groups), the submonoid $SL_d(\mathbb{N})$ of $SL_d(\mathbb{Z})$ defined by

$$\mathrm{SL}_d(\mathbb{N}) := \big\{ [a_{ij}]_{1 \le i,j \le d} \in \mathrm{SL}_d(\mathbb{Z}) \big| a_{ij} \in \mathbb{N} \big\}.$$

Note also that, any cancellative Abelian monoid is lunar, since it can be embedded into its Grothendieck group (see, e.g., [BG09, Section 2.A, page 52]).

Corollary 1.1. The Hankel system on any cancellative Abelian monoid has SAP.

As a consequence, we obtain the SAP (and in fact the much stronger hereditary SAP in the sense of Definition 5) for both the system of small Hankel operators and that of big Hankel operators on the higher dimensional torus \mathbb{T}^d ($d \ge 1$), as well as on the infinite dimensional torus $\mathbb{T}^{\mathbb{N}}$. Note that, by the unique factorization integers theorem, the small Hankel operators on $\mathbb{T}^{\mathbb{N}}$ can be transformed to multiplicative Hankel operators associated to the multiplicative monoid of positive integers $\mathbb{N}^* = \mathbb{N} \setminus \{0\}$ (see [Hel10, OCS12, BOCS21]).

It should be noted that Theorem A goes beyond the class of group-embeddable monoids. In fact, Bush [Bus71] showed that the lunar condition is not sufficient for the group-embeddability.

Theorem A holds in a much more general abstract setting (see Theorem B below). Namely, by introducing a key algebraic notion of *lunar maps* (see Definition 4), we obtain SAP for non-classical Hankel systems (1.3) associated with any lunar map. Here we only mention two concrete such examples:

- Arbitrary spatial compression of the regular representation system of any discrete group. See Corollaries 5.1 and 5.2.
- The Hankel system $\{\Gamma_{\ell}^{\Phi}\}$ determined by the level sets of any rational map $\Phi(x,y) = ax^m + by^n$ with $a, b, m, n \in \mathbb{Z}^* = \mathbb{Z} \setminus \{0\}$:

$$\Gamma^{\mathbf{\Phi}}_{\ell}(x,y) = \mathbb{1}(ax^m + by^n = \ell), \quad x, y \in \mathbb{N}^* = \mathbb{N} \setminus \{0\}, \, \ell \in \Phi(\mathbb{N}^* \times \mathbb{N}^*).$$

See Example 5.1 for details. It seems to be a natural problem of determine all the lunar rational maps or lunar polynomial maps, see Problem 8 in §5.2.

We conjecture that lunar condition on a map is equivalent to the SAP of its associated Hankel system, see Problem 7. For a concrete example of non-lunar map inducing a non-SAP Hankel system, see the map induced by a multicolor checkerboard in Example 3.5.

II. Harmonic analysis applications of SAP. In the classical setting of the additive monoid of non-negative integers $\mathbb{N} = \{0, 1, 2, \dots\}$, as well as all the more general free Abelian monoids \mathbb{N}^d , the self-absorption property is applied to the study of completely bounded Fourier multipliers between Hardy spaces (see Section 6). The reader is referred to [AMZ22, AMZ23, CAGPPT23] for recent works about completely bounded multipliers.

All notations below are classical and will be recalled in §6.1. All subspaces or quotient spaces of L^p are equipped with their standard *o.s.s.*, see [Pis03, Chapters 1-5] and [Pis98, Chapters 1-2]. For simplifying notation, throughout the paper, by writing L^p, H^p etc, we mean $L^p(\mathbb{T}), H^p(\mathbb{T})$ respectively. We identify $L^{\infty}/\overline{H_0^{\infty}}$ with BMOA. All spaces of completely bounded Fourier multipliers (such as $(H^p, H^q)_{cb}, (H^p, \text{BMOA})_{cb}$ etc.) are equipped with their natural *o.s.s.*

Based on Grothendieck's work [Gro53], Pisier [Pis01] obtained a striking description of $(H^1, H^1)_{cb}$ as follows: $\varphi \in (H^1, H^1)_{cb}$ if and only if there exist $v, w \in \ell^{\infty}(\ell^2)$ such that the matrix $[\langle v_m, w_n \rangle]_{m,n \in \mathbb{N}}$ is of Hankel type and

(1.8)
$$\widehat{\varphi}(m+n) = \langle v_m, w_n \rangle_{\ell^2} \text{ for all } m, n \in \mathbb{N}.$$

Very recently, in the same spirit, Arnold, Le Merdy and Zadeh [AMZ22, AMZ23] described $(H^p(\mathbb{R}), H^p(\mathbb{R}))_{cb}$ for $1 \le p < \infty$. However, in general, it seems difficult to determine pairs (v, w) of vectors in $\ell^{\infty}(\ell^2)$ such that $[\langle v_m, w_n \rangle]_{m,n \in \mathbb{N}}$ is of Hankel type. Therefore, it would be interesting to give an effective method to determine when φ satisfies (1.8).

On the other hand, the question of characterizing $(H^p, H^q)_{cb}$ seems open for any other non-trivial distinct pairs of (p,q) for $1 \le p,q < \infty$.

To state our new findings, we introduce analytic function spaces BMOA^(*p*) on \mathbb{T} as follows. For any $1 \le p < \infty$, define

(1.9)
$$\operatorname{BMOA}^{(p)} := \left\{ \varphi = \sum_{n=0}^{\infty} \widehat{\varphi}(n) e^{in\theta} \Big| \|\varphi\|_{\operatorname{BMOA}^{(p)}} = \left\| \sum_{n=0}^{\infty} |\widehat{\varphi}(n)|^p e^{in\theta} \right\|_{\operatorname{BMOA}}^{1/p} < \infty \right\}.$$

By a well-known result of Fefferman on BMOA functions with non-negative Fourier coefficients (see, e.g., [JVA16, Thm. 6.4.3]),

$$\|\boldsymbol{\varphi}\|_{\mathrm{BMOA}^{(p)}} \approx |\widehat{\boldsymbol{\varphi}}(0)| + \sup_{n \ge 1} \Big\{ \sum_{k=1}^{\infty} \Big(\sum_{kn \le j < (k+1)n} |\widehat{\boldsymbol{\varphi}}(j)|^p \Big)^2 \Big\}^{1/2p}.$$

By convention, we set

(1.10)
$$\operatorname{BMOA}^{(\infty)} := \left\{ \varphi = \sum_{n=0}^{\infty} \widehat{\varphi}(n) e^{in\theta} \Big| \|\varphi\|_{\operatorname{BMOA}^{(\infty)}} = \sup_{n \ge 0} |\widehat{\varphi}(n)| < \infty \right\}.$$

Corollary 1.2 (see Proposition 6.2 for its complete isometric version). *We have the following isometric isomorphisms:*

$$(H^1, BMOA)_{cb} = BMOA \text{ and } (H^1, H^2)_{cb} = (H^2, BMOA)_{cb} = BMOA^{(2)}.$$

We then prove in Lemmas 6.5 and 6.6 that, BMOA^(p) are Banach spaces and the family {BMOA^(p) : $1 \le p \le \infty$ } forms a complex interpolation scale.

Corollary 1.3 (see Theorem 6.1). *For any* $1 \le p \le 2 \le q < \infty$,

$$\mathsf{BMOA}^{(u)} \xleftarrow{bounded}_{inclusion} (H^p, H^q)_{cb} \quad with \ \frac{1}{u} = \frac{1}{p} - \frac{1}{q}.$$

Further applications are as follows.

1) Fourier-Schur multiplier inequalities with critical exponent 4/3 (Corollary 6.9 and Proposition 6.10): For any ℓ^2 -column-vector-valued function $f \in H^1(\ell^2)$,

$$\left\| \left(\int_{\mathbb{T}} \left[(f * \varphi) (f * \varphi)^* \right] dm \right)^{1/2} \right\|_{S_{4/3}} \le \|\varphi\|_{\mathrm{BMOA}^{(2)}} \cdot \|f\|_{H^1(\ell^2)}$$

The exponent 4/3 is critical: it can not be replaced by any exponent p < 4/3.

Exact complete bounded norm of the Carleman embedding (see Corollary 6.13 for the one-dimensional case and Corollary 6.14 for its higher dimensional analogue): For any *f* ∈ *H*¹(*S*₁) and *g* ∈ ℂ[*z*] ⊗ *S*∞ (the constant √π below is optimal),

$$\left|\int_{\mathbb{D}} \operatorname{Tr}(f(z)\overline{g(z)}) dA(z)\right| \leq \sqrt{\pi} \|f\|_{H^{1}(S_{1})} \left\|\int_{\mathbb{D}} \left[g(z) \otimes \overline{g(z)}\right] dA(z) \right\|_{B(\ell^{2} \otimes 2^{\ell^{2}})}^{1/2}$$

3) Failure of hyper-complete-contractivity for the Poisson semigroup (Corollary 6.17): Consider the Poisson convolution P_r on \mathbb{T} . Janson's hyper-contractivity [Jan83] states that P_r is contractive from H^1 to H^2 if and only if $r \le \sqrt{1/2}$. However,

$$||P_r||_{(H^1, H^2)_{cb}} = (1 - r^4)^{-1/2} > 1$$
 for any $r \in (0, 1)$.

An inequality for Hankel operators (see Corollary 6.8 for its higher dimensional analogue): Let φ ∈ H¹ and set φ[†](e^{iθ}) = φ(e^{-iθ}), then for any mutually orthogonal functions (f_k)_{k≥1} in H²,

$$\Big|\sum_{k=1}^{\infty} (\Gamma_{f_k * \varphi})^* \Gamma_{f_k * \varphi} \Big\|_{B(\overline{H^2})}^{1/2} \le \|\Gamma_{\varphi * \varphi^{\dagger}}\|_{\operatorname{Hank}(\overline{H^2}, H^2)}^{1/2} \Big(\sum_{k=1}^{\infty} \|f_k\|_{H^2}^4\Big)^{1/4},$$

where $\text{Hank}(\overline{H^2}, H^2)$ denotes the space of Hankel operators and Γ_{ψ} denotes the corresponding Hankel operator with symbol ψ (see §6.1.2 for the notation).

III. Coupled foliations. The main ingredient in the proof of Theorem A is the construction and analysis of the *coupled foliations* illustrated in Figure 1 for the following two subsets in \mathcal{M}^2 (in Abelian case, they both equal \mathcal{M}^2 , the roles of these two subsets will be explained soon):

$$(\mathcal{M}^2)_{\text{left}} := \{(x, y) \in \mathcal{M}^2 : \exists (a, b), s.t. ax = by\};\$$

 $(\mathcal{M}^2)_{\text{right}} := \{(a, b) \in \mathcal{M}^2 : \exists (x, y), s.t. ax = by\}.$

The lunar condition (1.7) is crucial in our construction. Under this lunar condition, we may define an equivalence relation on $(\mathcal{M}^2)_{\text{right}}$ by

$$(a,b) \sim (c,d) \stackrel{def}{\iff} \exists (x,y), s.t. ax = by \& cx = dy.$$



FIGURE 1. The coupled foliation structures.

Denote by $[\mathcal{M}^2]_{\text{right}} = (\mathcal{M}^2)_{\text{right}}/\sim$ the quotient space and by $[a, b] \in [\mathcal{M}^2]_{\text{right}}$ the equivalence class of $(a, b) \in (\mathcal{M}^2)_{\text{right}}$. Then the coupled foliation decompositions are given by

(1.11)
$$(\mathcal{M}^2)_{\text{left}} = \bigsqcup_{[a,b] \in [\mathcal{M}^2]_{\text{right}}} \blacklozenge_{[a,b]} \text{ and } (\mathcal{M}^2)_{\text{right}} = \bigsqcup_{[a,b] \in [\mathcal{M}^2]_{\text{right}}} \clubsuit_{[a,b]}$$

with $\blacklozenge_{[a,b]} := \{(x,y) \in \mathcal{M}^2 : ax = by\}$ and $\clubsuit_{[a,b]} := \{(c,d) \in \mathcal{M}^2 : (c,d) \sim (a,b)\}$. The lunar condition ensures that the definition of $\blacklozenge_{[a,b]}$ depends only on the equivalence class of (a,b).

A key property shared by all these leaves $\blacklozenge_{[a,b]}, \clubsuit_{[a,b]}$ is the "*one-dimensionality*". More precisely, let π_1, π_2 be the usual coordinate projections on \mathcal{M}^2 , then the lunar condition combined with the cancellativity implies that all the following maps

$$\pi_1|_{\bigstar_{[a,b]}}, \quad \pi_2|_{\bigstar_{[a,b]}}, \quad \pi_1|_{\bigstar_{[a,b]}}, \quad \pi_2|_{\bigstar_{[a,b]}}$$

are injective for any $[a,b] \in [\mathcal{M}^2]_{\text{right}}$.

Remark 1.1. Consider the dual equivalence relation on $(\mathcal{M}^2)_{\text{left}}$ defined by: $(x, y) \sim' (z, w)$ if and only if there exists (a, b) such that ax = by & az = bw. The relation \sim' induces a natural decomposition of $(\mathcal{M}^2)_{\text{left}}$. One can show that, under the lunar condition on \mathcal{M} , this decomposition coincides with the one given in (1.11). However, it should be emphasized that, the one-to-one correspondance given by (1.11) between the leaves in $(\mathcal{M}^2)_{\text{left}}$ and $(\mathcal{M}^2)_{\text{right}}$ is important for us.

IV. How coupled foliations lead to SAP ? The central step to SAP is the *simultaneous block-diagonalizations* of operators $\Gamma_t^{\mathcal{M}} \otimes \Gamma_t^{\mathcal{M}}$ for all $t \in \mathcal{M}$. Such simultaneous block-diagonalizations will be explicitly realized by the coupled foliations in (1.11).

Now, we explain the roles of $(\mathcal{M}^2)_{\text{left}}$ and $(\mathcal{M}^2)_{\text{right}}$: they induce two subspaces $\ell^2((\mathcal{M}^2)_{\text{left}})$ and $\ell^2((\mathcal{M}^2)_{\text{right}})$ of $\ell^2(\mathcal{M}^2)$. Moreover, using the canonical identification of $\ell^2(\mathcal{M}^2)$ with $\ell^2(\mathcal{M}) \otimes_2 \ell^2(\mathcal{M})$ and hence the canonical identifications of their subspaces, one can show that

(i) the orthogonal complement of $\ell^2((\mathcal{M}^2)_{\text{left}})$ in $\ell^2(\mathcal{M}^2)$ lies in the kernel of $\Gamma_t^{\mathcal{M}} \otimes \Gamma_t^{\mathcal{M}}$ simultaneously for all $t \in \mathcal{M}$:

$$[\ell^2((\mathcal{M}^2)_{\text{left}})]^{\perp} = \ell^2(\mathcal{M}^2) \ominus \ell^2((\mathcal{M}^2)_{\text{left}}) \subset \bigcap_{t \in \mathcal{M}} \ker(\Gamma_t^{\mathcal{M}} \otimes \Gamma_t^{\mathcal{M}});$$

(ii) the images of $\ell^2(\mathcal{M}^2)$ under all the maps $\Gamma_t^{\mathcal{M}} \otimes \Gamma_t^{\mathcal{M}}$ are contained in $\ell^2((\mathcal{M}^2)_{\text{right}})$:

$$\overline{\operatorname{span}}\big\{\bigcup_{t\in\mathcal{M}}\operatorname{Im}(\Gamma^{\mathcal{M}}_t\otimes\Gamma^{\mathcal{M}}_t)\big\}\subset\ell^2((\mathcal{M}^2)_{\operatorname{right}}).$$

Next we explain how the coupled foliations (1.11) are used to construct simultaneous blockdiagonalization of all $\Gamma_t^{\mathcal{M}} \otimes \Gamma_t^{\mathcal{M}}$. More precisely, the coupled foliations induce the following coupled orthogonal decompositions:

$$\ell^{2}((\mathcal{M}^{2})_{\text{left}}) = \bigoplus_{[a,b] \in [\mathcal{M}^{2}]_{\text{right}}} \ell^{2}(\clubsuit_{[a,b]}) \text{ and } \ell^{2}((\mathcal{M}^{2})_{\text{right}}) = \bigoplus_{[a,b] \in [\mathcal{M}^{2}]_{\text{right}}} \ell^{2}(\clubsuit_{[a,b]}).$$

We further show that

$$(\Gamma_t^{\mathcal{M}} \otimes \Gamma_t^{\mathcal{M}})(\ell^2(\blacklozenge_{[a,b]})) \subset \ell^2(\clubsuit_{[a,b]}), \text{ for all } t \in \mathcal{M}.$$

Finally, by the "one-dimensional structure" of all the leaves in the foliations (1.11), for each pair of the coupled leaves $(\bigstar_{[a,b]}, \bigstar_{[a,b]})$, the family $\{(\Gamma_t^{\mathcal{M}} \otimes \Gamma_t^{\mathcal{M}})|_{\ell^2(\bigstar_{[a,b]})}\}_{t \in \mathcal{M}}$ is shown to satisfy the following simultaneous commutative diagram (as before, here " simultaneous" means that the contractive interwining operators $R^{[a,b]}$ and $S^{[a,b]}$ only depend on the equivalence class $[a,b] \in [\mathcal{M}^2]_{\text{right}}$ and are both independent of $t \in \mathcal{M}$):

with $R^{[a,b]}, S^{[a,b]}$ two explicitly constructed operators such that

$$||\mathbf{R}^{[a,b]}|| \le 1$$
 and $||\mathbf{S}^{[a,b]}|| \le 1$.

As a result, all operators $\Gamma_t^{\mathcal{M}} \otimes \Gamma_t^{\mathcal{M}}$ are simultaneously block-diagonalized, with notably simplified diagonal-blocks. In notation, we establish the decomposition

$$\Gamma_t^{\mathcal{M}} \otimes \Gamma_t^{\mathcal{M}} = \left([\ell^2((\mathcal{M}^2)_{\text{left}})]^{\perp} \xrightarrow{0} 0 \right) \oplus \bigoplus_{[a,b] \in [\mathcal{M}^2]_{\text{right}}} \left(\ell^2(\clubsuit_{[a,b]}) \xrightarrow{S^{[a,b]} \Gamma_t^{\mathcal{M}} R^{[a,b]}} \ell^2(\clubsuit_{[a,b]}) \right).$$

V. Open problems and further discussions. The SAP for the Hankel system on \mathbb{N}^d established in this paper (Corollary 1.1) is closely related to the well-known open problem on the existence of a higher dimensional Nehari-Sarason-Page Theorem on \mathbb{T}^d ($d \ge 2$). Indeed, by the SAP for the Hankel system on \mathbb{N}^d , a positive answer to the following Problem 1 would follow from the existence of such a higher dimensional NSP theorem.

Problem 1. Let $d \ge 2$ be an integer and consider the operator space $L^{\infty}(\mathbb{T}^d)/\overline{H_0^{\infty}(\mathbb{T}^d)}$ (equipped with its natural quotient *o.s.s.*). Does the family $\{e^{in\cdot\theta}\}_{n\in\mathbb{N}^d} \subset L^{\infty}(\mathbb{T}^d)/\overline{H_0^{\infty}(\mathbb{T}^d)}$ have κ -SAP for a finite κ ?

We say that a multiplier T from $H^1(\mathbb{T}^d)$ to $L^{\infty}(\mathbb{T}^d)/\overline{H_0^{\infty}(\mathbb{T}^d)}$ is *liftable* if it satisfies the commutative diagram:

$$L^{1}(\mathbb{T}^{d}) \xrightarrow{K} L^{\infty}(\mathbb{T}^{d})$$

$$\mathcal{I} \stackrel{\uparrow}{\text{inclusion}} Q \downarrow quotient$$

$$H^{1}(\mathbb{T}^{d}) \xrightarrow{T} L^{\infty}(\mathbb{T}^{d}) / \overline{H_{0}^{\infty}(\mathbb{T}^{d})}$$

where *K* is a bounded convolution operator.

Using similar arguments in the proof of Corollary 6.15 for the one-dimensional case, one can show that Problem 1 is equivalent to the following

Problem 2. Is every multiplier in $(H^1(\mathbb{T}^d), L^{\infty}(\mathbb{T}^d)/\overline{H_0^{\infty}(\mathbb{T}^d)})_{cb}$ liftable ?

Definition 3 (SAP monoids). A monoid is called a *SAP monoid* if its associated Hankel system has SAP. In such situation, we also say that the monoid has SAP.

The SAP of a monoid can be seen as an analytic property (the operator space structure) of an algebraic object (the associated Hankel system), while the lunar condition for a monoid is a pure algebraic property. Theorem A says that, this algebraic property implies the analytic one.

Problem 3. Does the class of SAP monoids coincide with that of lunar ones ?

It is not even known to us whether SAP is preserved by taking submonoids.

Problem 4. Can SAP always be inherited by submonoids ?

The following table provides informations of some classical operations preserving the class of lunar monoid and that of SAP monoid respectively.

	Cartesian product	free product	submonoid
lunar monoid (algebraic property)	\checkmark	\checkmark	\checkmark
SAP monoid (analytic property)	\checkmark	√ ^a	?

We believe that not all cancellative monoids have SAP. However, at this moment, we are not aware of such examples.

Problem 5. Do there exist non-SAP cancellative monoids ?

Acknowledgements. This work is supported by the National Natural Science Foundation of China (No.12288201). YH is supported by the grant NSFC 12131016 and 12201419, ZW is supported by NSF of Chongqing (CSTB2022BSXM-JCX0088, CSTB2022NSCQ-MSX0321) and FRFCU (2023CDJXY-043).

2. The simplest SAP monoid: An illustration

To illustrate our main idea, we include an algebraic proof of the SAP for the simplest additive monoid $(\mathbb{N}, +)$ of non-negative integers. See Proposition 2.1 and Corollary 2.2.

By identifying the Hankel operators on $\ell^2(\mathbb{N})$ with the standard Hankel operators on the Hardy spaces of \mathbb{T} , one can prove the SAP for the Hankel system on \mathbb{N} in a pure analytic method by applying the celebrated Nehari-Sarason-Page Theorem (which is briefly recalled in §6.1.2).

However, the Nehari-Sarason-Page Theorem is not available in general situations. Our algebraic proof is self-contained and can be easily adapted to all cancellative Abelian monoids and more general non-Abelian monoids (see §4 and §5). Moreover, even in the setting of \mathbb{N} , this algebraic approach leads to a stronger hereditary SAP (see Definition 5, Theorem 4.1 and Corollary 5.1) and it can even be adapted to operators on ℓ^p -spaces or on more general symmetric sequence-spaces, both of which clearly do not follow from the Nehari-Sarason-Page Theorem.

^aThe proof that free product preserves the class of SAP monoids requires much more technical details, so we leave it to a separate work.

2.1. The simplest SAP monoid $(\mathbb{N}, +)$. Throughout the paper, $B(\mathcal{H}, \mathcal{K})$ denotes the bounded linear operators from a Hilbert space \mathcal{H} to another one \mathcal{K} and $B(\mathcal{H}) = B(\mathcal{H}, \mathcal{H})$ and $\mathcal{H} \otimes_2 \mathcal{K}$ denotes the Hilbertian tensor product of \mathcal{H} and \mathcal{K} .

For simplifying notation, we always write $\ell^2 = \ell^2(\mathbb{N})$. The standard Hankel matrix associated to a complex sequence $a = (a_i)_{i>0} \in \mathbb{C}^{\mathbb{N}}$ is defined by

$$\Gamma_a = [a_{i+j}]_{i,j\geq 0}.$$

Let $\operatorname{Hank}(\ell^2)$ denote the collection of all bounded $\Gamma_a \in B(\ell^2)$. More generally, for any sequence $(x_n)_{n\geq 0}$ in $B(\mathscr{H})$, we define a block Hankel matrix (which a priori does not represent an element in $B(\mathscr{H} \otimes_2 \ell^2)$):

$$\Gamma_x = \Gamma_{(x_n)_{n \in \mathbb{N}}} := [x_{i+j}]_{i,j \ge 0}.$$

Informally, we may write $\Gamma_x = \sum_{n=0}^{\infty} x_n \otimes \Gamma_n$, where the operator Γ_n (for any non-negative integer $n \ge 0$) is defined by

(2.1)
$$\Gamma_n(i,j) = \mathbb{1}(i+j=n), \quad i,j \ge 0.$$

In particular, for any $\Gamma_a \in \text{Hank}(\ell^2)$, define a block Hankel matrix by the informal series:

$$J_2(\Gamma_a) := \Gamma_{(a_n \Gamma_n)_{n \in \mathbb{N}}} = \sum_{n=0}^{\infty} a_n \Gamma_n \otimes \Gamma_n.$$

Proposition 2.1. Let $\Gamma_a \in \text{Hank}(\ell^2)$. Then $J_2(\Gamma_a) \in B(\ell^2 \otimes_2 \ell^2)$. Moreover, the linear map $J_2 : \text{Hank}(\ell^2) \to B(\ell^2 \otimes_2 \ell^2)$ is a complete isometric embedding.

Corollary 2.2. *The monoid* $(\mathbb{N}, +)$ *has SAP.*

Remark 2.1. Let $C^*({\{\Gamma_n\}_{n\in\mathbb{N}}}) \subset B(\ell^2)$ (resp. $C^*({\{\Gamma_n\otimes\}_{n\in\mathbb{N}}}) \subset B(\ell^2\otimes_2\ell^2)$) denote the C^* -algebra generated by the family ${\{\Gamma_n\}_{n\in\mathbb{N}}}$ (resp. ${\{\Gamma_n\otimes\Gamma_n\}_{n\in\mathbb{N}}}$). Then one may check that the map $\Gamma_n \mapsto \Gamma_n \otimes \Gamma_n(\forall n \in \mathbb{N})$ can not be extended to a C^* -representation from $C^*({\{\Gamma_n\}_{n\in\mathbb{N}}})$ to $C^*({\{\Gamma_n\otimes\Gamma_n\}_{n\in\mathbb{N}}})$. Indeed, by direct computation, one obtains

$$\|2\Gamma_0 \otimes \Gamma_0 - 2\Gamma_1 \otimes \Gamma_1 - (\Gamma_1 \otimes \Gamma_1)^2\| = 3 > \|2\Gamma_0 - 2\Gamma_1 - \Gamma_1^2\| = \sqrt{5}.$$

The same computation also implies that the system $\{Id\} \cup \{\Gamma_n\}_{n \in \mathbb{N}}$ has no SAP since

$$||2\Gamma_0 \otimes \Gamma_0 - 2\Gamma_1 \otimes \Gamma_1 - Id \otimes Id|| = 3 > ||2\Gamma_0 - 2\Gamma_1 - Id|| = \sqrt{5}.$$

2.2. **Proof of Proposition 2.1.** Let $\{e_i : i \in \mathbb{N}\}$ be the standard orthonormal basis of ℓ^2 . For any integer $k \ge 0$, define closed subspaces \mathcal{H}_k and \mathcal{H}_{-k} of $\ell^2 \otimes_2 \ell^2$ as

$$\mathcal{H}_k = \overline{\operatorname{span}} \{ e_i \otimes e_{i+k} : i \in \mathbb{N} \} \text{ and } \mathcal{H}_{-k} = \overline{\operatorname{span}} \{ e_{i+k} \otimes e_i : i \in \mathbb{N} \}.$$

Clearly, \mathcal{H}_{ℓ} are mutually orthogonal and

(2.2)
$$\ell^2 \otimes_2 \ell^2 = \bigoplus_{\ell \in \mathbb{Z}} \mathcal{H}_\ell.$$

Under the above orthogonal decomposition, any operator $A \in B(\ell^2 \otimes_2 \ell^2)$ has a block-matrix representation:

$$A = [A_{\ell,\ell'}]_{\ell,\ell'\in\mathbb{Z}} ext{ with } A_{\ell,\ell'} \in B(\mathcal{H}_{\ell'},\mathcal{H}_{\ell}).$$

In particular, an operator $D \in B(\ell^2 \otimes_2 \ell^2)$ (resp. $D' \in B(\ell^2 \otimes_2 \ell^2)$) is called of block diagonal form (resp. of block anti-diagonal form), if

$$D = \begin{bmatrix} \ddots & & & & \\ & D_{-1} & & & \\ & & D_0 & & \\ & & & D_1 & \\ & & & & \ddots \end{bmatrix}, \quad D' = \begin{bmatrix} & & & & \ddots \\ & & D'_{-1} & & \\ & D'_0 & & & \\ & D'_1 & & & \\ & \ddots & & & & \end{bmatrix},$$

where $D_{\ell} \in B(\mathcal{H}_{\ell}, \mathcal{H}_{\ell})$ and $D'_{\ell} \in B(\mathcal{H}_{\ell}, \mathcal{H}_{-\ell})$. For simplification, we write

(2.3)
$$D = \operatorname{diag}(D_{\ell})_{\ell \in \mathbb{Z}} \text{ and } D' = \operatorname{adiag}(D'_{\ell})_{\ell \in \mathbb{Z}}$$

Lemma 2.3. The subspace $\mathcal{H}_0 \subset \ell^2 \otimes_2 \ell^2$ is $\Gamma_n \otimes \Gamma_n$ -invariant for all $n \in \mathbb{N}$. Moreover, we have the following commutative diagram:

(2.4)
$$\begin{array}{c} \mathcal{H}_{0} \xrightarrow{\Gamma_{n} \otimes \Gamma_{n}} \mathcal{H}_{0} \\ U \downarrow \simeq & \simeq \downarrow U \\ \ell^{2} \xrightarrow{\Gamma_{n}} \ell^{2} \end{array}$$

where U is the unitary operator such that

(2.5)
$$U: \begin{array}{ccc} \mathcal{H}_0 & \to & \ell^2 \\ e_i \otimes e_i & \mapsto & e_i, & i \in \mathbb{N}. \end{array}$$

In other words, for all $n \in \mathbb{N}$ *,*

(2.6)
$$(\Gamma_n \otimes \Gamma_n)|_{\mathcal{H}_0} = U^{-1} \Gamma_n U.$$

Proof. Fix $n \in \mathbb{N}$. By the definition of Γ_n , we have

(2.7)
$$\Gamma_n e_i = \mathbb{1}(i \le n) e_{n-i}, \quad i \in \mathbb{N}.$$

Hence, for any $i \in \mathbb{N}$,

$$(\Gamma_n \otimes \Gamma_n)(e_i \otimes e_i) = \mathbb{1}(i \le n)e_{n-i} \otimes e_{n-i}$$

It follows that \mathcal{H}_0 is $\Gamma_n \otimes \Gamma_n$ -invariant. Now by (2.5), for any $i \in \mathbb{N}$,

$$\Gamma_n U(e_i \otimes e_i) = \Gamma_n e_i = \mathbb{1}(i \le n) e_{n-i},$$
$$U(\Gamma_n \otimes \Gamma_n)(e_i \otimes e_i) = U(\Gamma_n e_i \otimes \Gamma_n e_i) = \mathbb{1}(i \le n) e_{n-i}.$$

The desired equality $\Gamma_n U = U(\Gamma_n \otimes \Gamma_n)$ then follows.

For any integer $k \ge 0$, define operators $W_{\pm k} : \mathcal{H}_{\pm k} \to \mathcal{H}_0$ and $V_{\pm k} : \mathcal{H}_0 \to \mathcal{H}_{\pm k}$ as follows: for any $i \in \mathbb{N}$,

(2.8)
$$\begin{cases} W_k(e_i \otimes e_{i+k}) = e_{i+k} \otimes e_{i+k}, \\ W_{-k}(e_{i+k} \otimes e_i) = e_{i+k} \otimes e_{i+k}, \\ V_k(e_i \otimes e_i) = e_i \otimes e_{i+k}, \\ V_{-k}(e_i \otimes e_i) = e_{i+k} \otimes e_i. \end{cases}$$

Note that $W_{\pm k}$ are isometric embeddings and $V_{\pm k}$ are unitary operators. In particular, $W_0 = V_0 = I_{\mathcal{H}_0}$, where $I_{\mathcal{H}_0}$ denotes the identity map on \mathcal{H}_0 .

Remark 2.2. The operation rules in (2.8) are easy to remember. Indeed, these operations share a common rule of increasing only one of the indices of a tensor $e_i \otimes e_j$ (even for the operators with negative indices as W_{-k} and V_{-k}) in an appropriate way.

Lemma 2.4. For any $\ell \in \mathbb{Z}$ and any $n \in \mathbb{N}$, the operator $\Gamma_n \otimes \Gamma_n$ maps \mathcal{H}_ℓ into $\mathcal{H}_{-\ell}$:

$$(\Gamma_n \otimes \Gamma_n)|_{\mathcal{H}_\ell} \in B(\mathcal{H}_\ell, \mathcal{H}_{-\ell}).$$

Moreover, we have the following commutative diagram:

$$egin{array}{cccc} \mathcal{H}_\ell & \stackrel{\Gamma_n \otimes \Gamma_n}{\longrightarrow} \mathcal{H}_{-\ell} \ W_\ell & & \uparrow V_{-\ell} \ \mathcal{H}_0 & \stackrel{\Gamma_n \otimes \Gamma_n}{\longrightarrow} \mathcal{H}_0 \end{array}$$

In other words,

(2.9)
$$(\Gamma_n \otimes \Gamma_n)|_{\mathcal{H}_\ell} = V_{-\ell} \big[(\Gamma_n \otimes \Gamma_n)|_{\mathcal{H}_0} \big] W_\ell.$$

Consequently, with respect to the orthogonal decomposition (2.2), the operator $\Gamma_n \otimes \Gamma_n$ is of block anti-diagonal form. According to the notation (2.3), we can write

$$\Gamma_n \otimes \Gamma_n = \operatorname{adiag} \left(V_{-\ell} \left[(\Gamma_n \otimes \Gamma_n) |_{\mathcal{H}_0} \right] W_{\ell} \right)_{\ell \in \mathbb{Z}}.$$

Proof. The case $\ell = 0$ is trivial. Now take any positive integer $k \ge 1$. Then for all $i \in \mathbb{N}$,

$$(\Gamma_n \otimes \Gamma_n)(e_i \otimes e_{i+k}) = \mathbb{1}(i+k \le n)e_{n-i} \otimes e_{n-i-k} \in \mathcal{H}_{-k}$$
$$(\Gamma_n \otimes \Gamma_n)(e_{i+k} \otimes e_i) = \mathbb{1}(i+k \le n)e_{n-i-k} \otimes e_{n-i} \in \mathcal{H}_k.$$

It follows that $(\Gamma_n \otimes \Gamma_n)(\mathcal{H}_\ell) \subset \mathcal{H}_{-\ell}$ for any integer $\ell \in \mathbb{Z} \setminus \{0\}$. By the definitions of $W_{\pm k}, V_{\pm k}$ in (2.8) and Remark 2.2, we have

$$V_{-k}(\Gamma_n \otimes \Gamma_n) W_k(e_i \otimes e_{i+k}) = V_{-k}(\Gamma_n \otimes \Gamma_n)(e_{i+k} \otimes e_{i+k})$$

= $V_{-k}(\mathbb{1}(i+k \le n)e_{n-i-k} \otimes e_{n-i-k})$
= $\mathbb{1}(i+k \le n)e_{n-i} \otimes e_{n-i-k}$

and

$$V_k(\Gamma_n \otimes \Gamma_n) W_{-k}(e_{i+k} \otimes e_i) = V_k(\Gamma_n \otimes \Gamma_n)(e_{i+k} \otimes e_{i+k})$$

= $V_k(\mathbb{1}(i+k \le n)e_{n-i-k} \otimes e_{n-i-k})$
= $\mathbb{1}(i+k \le n)e_{n-i-k} \otimes e_{n-i}.$

Therefore, for all $i \in \mathbb{N}$,

$$V_{-k}(\Gamma_n \otimes \Gamma_n) W_k(e_i \otimes e_{i+k}) = (\Gamma_n \otimes \Gamma_n)(e_i \otimes e_{i+k}),$$

$$V_k(\Gamma_n \otimes \Gamma_n) W_{-k}(e_{i+k} \otimes e_i) = (\Gamma_n \otimes \Gamma_n)(e_{i+k} \otimes e_i).$$

The desired equality (2.9) follows.

Let $\bigoplus_{\ell \in \mathbb{Z}} \mathcal{H}_0$ denote the Hilbert space

$$\bigoplus_{\ell \in \mathbb{Z}} \mathcal{H}_0 = \left\{ (v_\ell)_{\ell \in \mathbb{Z}} \middle| v_\ell \in \mathcal{H}_0 \text{ for all } \ell \in \mathbb{Z} \text{ and } \sum_{\ell \in \mathbb{Z}} ||v_\ell||^2 < \infty \right\}$$

carrying the natural inner product. The unitary operator $U : \mathcal{H}_0 \to \ell^2$ defined in (2.5) induces a faithful C^* -representation:

(2.10)
$$\begin{array}{ccc} \mathcal{R}_U : & B(\ell^2) & \xrightarrow{faithful} & B\bigl(\bigoplus_{\ell \in \mathbb{Z}} \mathcal{H}_0\bigr) \\ & T & \mapsto & \mathcal{R}_U(T) = (U^{-1}TU)^{\oplus \mathbb{Z}} \end{array}$$

where the operator $(U^{-1}TU)^{\oplus\mathbb{Z}}$ is given by $(U^{-1}TU)^{\oplus\mathbb{Z}}((v_{\ell})_{\ell\in\mathbb{Z}}) = (U^{-1}TUv_{\ell})_{\ell\in\mathbb{Z}}$. Recall the orthogonal decomposition in (2.2): $\ell^2 \otimes_2 \ell^2 = \bigoplus_{\ell\in\mathbb{Z}} \mathcal{H}_{\ell}$.

Proposition 2.5. For any $n \in \mathbb{N}$, the operator $\Gamma_n \otimes \Gamma_n$ admits the factorization:

(2.11)
$$\Gamma_n \otimes \Gamma_n = \mathcal{VFR}_U(\Gamma_n)\mathcal{W}_s$$

where \mathcal{V}, \mathcal{W} are defined by (recall the definitions of $W_{\pm k}, V_{\pm k}$ in (2.8))

$$\bigoplus_{\ell \in \mathbb{Z}} \mathcal{H}_0 \xrightarrow{\mathcal{V} = \bigoplus_{\ell \in \mathbb{Z}} V_\ell} \bigoplus_{\ell \in \mathbb{Z}} \mathcal{H}_\ell, \quad \bigoplus_{\ell \in \mathbb{Z}} \mathcal{H}_\ell \xrightarrow{\mathcal{W} = \bigoplus_{\ell \in \mathbb{Z}} W_\ell} \bigoplus_{\ell \in \mathbb{Z}} \mathcal{H}_0$$

and $\mathcal{F}: \bigoplus_{\ell \in \mathbb{Z}} \mathcal{H}_0 \to \bigoplus_{\ell \in \mathbb{Z}} \mathcal{H}_0$ is the unitary operator defined by $\mathcal{F}((v_\ell)_{\ell \in \mathbb{Z}}) = (v_{-\ell})_{\ell \in \mathbb{Z}}$. *Proof.* By Lemmas 2.3 and 2.4,

$$\Gamma_n \otimes \Gamma_n = \operatorname{adiag} \left(V_{-\ell} \left[(\Gamma_n \otimes \Gamma_n) |_{\mathcal{H}_0} \right] W_\ell \right)_{\ell \in \mathbb{Z}}$$

= $\operatorname{adiag} \left(V_{-\ell} U^{-1} \Gamma_n U W_\ell \right)_{\ell \in \mathbb{Z}}.$

Take an arbitray $\ell_0 \in \mathbb{Z}$ and $\xi \in \mathcal{H}_{\ell_0}$. First of all,

$$(\Gamma_n \otimes \Gamma_n)(\xi) = V_{-\ell_0} U^{-1} \Gamma_n U W_{\ell_0}(\xi) \in \mathcal{H}_{-\ell_0}.$$

On the other hand, by the definitions of $\mathcal{V}, \mathcal{F}, \mathcal{R}_U$ and \mathcal{W} ,

$$\begin{split} [\mathcal{VFR}_{U}(\Gamma_{n})\mathcal{W}](\xi) &= \left[\left(\bigoplus_{\ell \in \mathbb{Z}} V_{\ell} \right) \circ \mathcal{F} \circ (U^{-1}\Gamma_{n}U)^{\oplus \mathbb{Z}} \circ \left(\bigoplus_{\ell \in \mathbb{Z}} W_{\ell} \right) \right] (\xi) \\ &= \left[\left(\bigoplus_{\ell \in \mathbb{Z}} V_{\ell} \right) \circ \mathcal{F} \right] (\cdots, 0, \underbrace{U^{-1}\Gamma_{n}UW_{\ell_{0}}(\xi)}_{\text{at the }\ell_{0}\text{-th position}}, 0, \cdots) \\ &= \left(\bigoplus_{\ell \in \mathbb{Z}} V_{\ell} \right) (\cdots, 0, \underbrace{U^{-1}\Gamma_{n}UW_{\ell_{0}}(\xi)}_{\text{at the }(-\ell_{0})\text{-th position}}, 0, \cdots) \\ &= (\cdots, 0, \underbrace{V_{-\ell_{0}}U^{-1}\Gamma_{n}UW_{\ell_{0}}(\xi)}_{\text{at the }(-\ell_{0})\text{-th position}}, 0, \cdots) \\ &= V_{-\ell_{0}}U^{-1}\Gamma_{n}UW_{\ell_{0}}(\xi) \in \mathcal{H}_{-\ell_{0}}. \end{split}$$

Hence

$$(\Gamma_n \otimes \Gamma_n)(\xi) = [\mathcal{VFR}_U(\Gamma_n)\mathcal{W}](\xi).$$

Since $\ell_0 \in \mathbb{Z}$ and $\xi \in \mathcal{H}_{\ell_0}$ are chosen arbitrarily, the desired equality (2.11) follows.

Now we are ready to prove Proposition 2.1. We need the following elementary equality.

Lemma 2.6. For any Hilbert space \mathscr{H} and any bounded sequence $(x_n)_{n\geq 0}$ in $B(\mathscr{H})$,

(2.12)
$$\left\|\sum_{n=0}^{\infty} x_n \otimes \Gamma_n\right\|_{B(\mathscr{H} \otimes_2 \ell^2)} = \sup_{0 < r < 1} \left\|\sum_{n=0}^{\infty} r^n x_n \otimes \Gamma_n\right\|_{B(\mathscr{H} \otimes_2 \ell^2)} \in [0,\infty].$$

Proof. For any $r \in (0,1)$, define a block diagonal operator $D_r = \text{diag}(I_{\mathscr{H}}, rI_{\mathscr{H}}, r^2I_{\mathscr{H}}, \cdots)$ on $\mathscr{H} \otimes_2 \ell^2 = \ell^2(\mathscr{H})$, where $I_{\mathscr{H}}$ is the identity map on \mathscr{H} . Clearly, $||D_r|| = 1$. Hence

$$\sup_{0 < r < 1} \|\Gamma_{(r^n x_n)_{n \in \mathbb{N}}}\| = \sup_{0 < r < 1} \|D_r \Gamma_{(x_n)_{n \in \mathbb{N}}} D_r\| \le \|\Gamma_{(x_n)_{n \in \mathbb{N}}}\|$$

Conversely, by the standard approximation,

$$\|\Gamma_{(x_n)_{n\in\mathbb{N}}}\| = \sup_{L\geq 0} \sup_{0< r<1} \|[r^{i+j}x_{i+j}]_{0\leq i,j\leq L}\| \leq \sup_{0< r<1} \|[r^{i+j}x_{i+j}]_{i,j\geq 0}\| = \sup_{0< r<1} \|\Gamma_{(r^nx_n)_{n\in\mathbb{N}}}\|.$$

The desired equality then follows.

Remark 2.3. The equality (2.12) applied to $(x_n \otimes \Gamma_n)_{n \in \mathbb{N}}$ in $B(\mathscr{H} \otimes_2 \ell^2)$ yields

$$\left\|\sum_{n=0}^{\infty} x_n \otimes \Gamma_n \otimes \Gamma_n\right\|_{B(\mathscr{H} \otimes_2 \ell^2 \otimes_2 \ell^2)} = \sup_{0 < r < 1} \left\|\sum_{n=0}^{\infty} r^n x_n \otimes \Gamma_n \otimes \Gamma_n\right\|_{B(\mathscr{H} \otimes_2 \ell^2 \otimes_2 \ell^2)}.$$

Note that if one side of the above equality is finite, then the sequence $(x_n)_{n \in \mathbb{N}}$ must be bounded.

Proof of Proposition 2.1. Consider the following linear subspace of $Hank(\ell^2)$ (which is clearly not norm-closed nor norm-dense):

$$\operatorname{Hank}^{(\infty)}(\ell^2) = \left\{ \sum_{n=0}^{\infty} a_n \Gamma_n \Big| a_n \in \mathbb{C} \text{ and } \sum_{n=0}^{\infty} |a_n| < \infty \right\} \subset \operatorname{Hank}(\ell^2)$$

By Lemma 2.6 and Remark 2.3, we only need to show that the restriction of J_2 on $\text{Hank}^{(\infty)}(\ell^2)$ is completely isometric. By Proposition 2.5, restricted on $\text{Hank}^{(\infty)}(\ell^2)$,

$$J_2\Big(\sum_{n=0}^{\infty}a_n\Gamma_n\Big)=\sum_{n=0}^{\infty}a_n\Gamma_n\otimes\Gamma_n=\sum_{n=0}^{\infty}a_n\mathcal{VFR}_U(\Gamma_n)\mathcal{W}=\mathcal{VFR}_U\Big(\sum_{n=0}^{\infty}a_n\Gamma_n\Big)\mathcal{W}.$$

The operator \mathcal{R}_U defined in (2.10) is a C^* -representation, hence $\|\mathcal{R}_U\|_{cb} \leq 1$. Since $\|\mathcal{F}\| \leq 1, \|\mathcal{V}\| \leq 1$ and $\|\mathcal{W}\| \leq 1$, we have (see, e.g., [Pis03, Thm. 1.6])

$$\|J_2: \operatorname{Hank}^{(\infty)}(\ell^2) \to B(\ell^2 \otimes_2 \ell^2)\|_{cb} \le \|\mathcal{VF}\| \|\mathcal{R}_U\|_{cb} \|\mathcal{W}\| \le 1.$$

Conversely, by the equality (2.6),

$$\sum_{n=0}^{\infty} a_n \Gamma_n = U \Big[\Big(\sum_{n=0}^{\infty} a_n \Gamma_n \otimes \Gamma_n \Big) \Big|_{\mathcal{H}_0} \Big] U^{-1} = U \Big[J_2 \Big(\sum_{n=0}^{\infty} a_n \Gamma_n \Big) \Big|_{\mathcal{H}_0} \Big] U^{-1}$$

It follows that the map

$$J_2(\operatorname{Hank}^{(\infty)}(\ell^2)) \xrightarrow{J_2^{-1}} \operatorname{Hank}^{(\infty)}(\ell^2)$$
$$J_2\left(\sum_{n=0}^{\infty} a_n \Gamma_n\right) \mapsto \sum_{n=0}^{\infty} a_n \Gamma_n$$

is completely contractive and we complete the whole proof.

3. Self-absorption property: general results

Recall the notion of SAP introduced in Definition 1. In this section, we give several general properties of SAP:

- SAP is preserved by tensor product (Proposition 3.1).
- Any system of Boolean operators has a restricted-version SAP (Proposition 3.2).
- In general, SAP is neither preserved by direct sum (Example 3.2) nor preserved by spatial compression (Example 3.3).
- The standard Fell's absorption principle implies that the regular representation system of any group has SAP (Example 3.4). However, it is quite non-trivial to prove that any spatial compression of the regular representation system of a group also has SAP (Corollary 5.2). Note that our result implies in particular a recent result of Katsoulis [Kat23, Cor. 5.3].

3.1. Tensor product preserves SAP. Given two families $\mathcal{F}_1 = \{x_i\}_{i \in I}$ and $\mathcal{F}_2 = \{y_j\}_{j \in J}$ in two *C**-algebras A_1 and A_2 respectively, slightly abusing the notation \otimes_{min} , we denote

$$\mathcal{F}_1 \otimes_{\min} \mathcal{F}_2 = \{x_i \otimes y_j\}_{(i,j) \in I \times J} \subset A_1 \otimes_{\min} A_2.$$

Denote by $\kappa_{SA}(\mathcal{F})$ the best constant in the defining inequality (1.6) of κ -SAP of a family \mathcal{F} . Hence \mathcal{F} has SAP if and only if $\kappa_{SA}(\mathcal{F}) = 1$.

Proposition 3.1. For any two families $\mathcal{F}_1 \subset A_1$ and $\mathcal{F}_2 \subset A_2$ in C^* -algebras A_1 and A_2 ,

$$\kappa_{SA}(\mathcal{F}_1 \otimes_{\min} \mathcal{F}_2) \leq \kappa_{SA}(\mathcal{F}_1)\kappa_{ST}(\mathcal{F}_2).$$

In particular, if both \mathcal{F}_1 and \mathcal{F}_2 have SAP, then so does $\mathcal{F}_1 \otimes_{min} \mathcal{F}_2$.

Proof. Write $\mathcal{F}_1 = \{x_i\}_{i \in I}$ and $\mathcal{F}_2 = \{y_j\}_{j \in J}$. Take any finitely supported sequence $\{b_{i,j}\}$ in a C^* -algebra B, by using the canonical C^* -isomorphism between $C_1 \otimes_{min} C_2$ and $C_2 \otimes_{min} C_1$ for any given C^* -algebras C_1, C_2 , we have (the norms are taken in the corresponding C^* -algebras)

$$\left\|\sum_{i\in I, j\in J} b_{i,j} \otimes x_i \otimes y_j \otimes \overline{x}_i \otimes \overline{y}_j\right\| = \left\|\sum_{j\in J} \left(\sum_{i\in I} b_{i,j} \otimes x_i \otimes \overline{x}_i\right) \otimes y_j \otimes \overline{y}_j\right\|$$

$$\leq \kappa_{SA}(\mathcal{F}_2) \left\|\sum_{j\in J} \left(\sum_{i\in I} b_{i,j} \otimes x_i \otimes \overline{x}_i\right) \otimes y_j\right\| = \kappa_{SA}(\mathcal{F}_2) \left\|\sum_{i\in I} \left(\sum_{j\in J} b_{i,j} \otimes y_j\right) \otimes x_i \otimes \overline{x}_i\right\|$$

$$\leq \kappa_{SA}(\mathcal{F}_2) \kappa_{SA}(\mathcal{F}_1) \left\|\sum_{i\in I} \left(\sum_{j\in J} b_{i,j} \otimes y_j\right) \otimes x_i\right\| = \kappa_{SA}(\mathcal{F}_1) \kappa_{SA}(\mathcal{F}_2) \left\|\sum_{i\in I, j\in J} b_{i,j} \otimes x_i \otimes y_j\right\|.$$

The converse inequality is proved similarly.

3.2. **SAP–an restricted version.** An operator $a \in B(\ell^2)$ is called Boolean if its standard matrix representation $[a(i, j)]_{i,j\in\mathbb{N}}$ satisfies $a(i, j) \in \{0, 1\}$ for all i, j. The support of a Boolean operator is defined by

$$supp(a) = \{(i, j) \in \mathbb{N}^2 : a(i, j) \neq 0\}.$$

Note that any non-zero element in a SAP system has unit norm. Proposition 3.2 suggests that any family of unit norm Boolean operators has a restricted-version SAP if we require certain positivity condition on the coefficients. This result is probably known, but we have not found it in the literature.

Proposition 3.2. Fix any family \mathcal{F} of unit-norm Boolean operators in $B(\ell^2)$. Then for any integer $m \ge 2$ (the norms below are taken in the corresponding C^* -algebras), the equality

$$\left\|\sum_{x\in\mathcal{F}}c_x\otimes x^{\otimes m}\right\| = \left\|\sum_{x\in\mathcal{F}}c_x\otimes x\right\| \quad \text{with } x^{\otimes m} = \underbrace{x\otimes\cdots\otimes x}_{m \text{ times}}$$

holds for any finitely supported family $(c_x)_{x \in \mathcal{F}}$ of $N \times N$ matrices satisfying any one of the following conditions:

- (c1) all matrices c_x have non-negative coefficients;
- (c2) $N = k^2$ and all matrices c_x have the form $c_x = b_x \otimes \overline{b}_x$ with b_x an $k \times k$ matrix;
- (c3) $N = k^2$ and all matrices c_x have the form $c_x = b_x \otimes b_x^*$ with b_x an $k \times k$ matrix.

The following lemma is elementary.

Lemma 3.3. A Boolean matrix $a = [a(i, j)]_{i,j \in \mathbb{N}}$ has operator norm ||a|| = 1 if and only if there exists a bijection $\sigma : I_{\sigma} \to J_{\sigma}$ between two subsets $I_{\sigma}, J_{\sigma} \subset \mathbb{N}$ such that

$$a(i,j) = \mathbb{1}(i \in I_{\sigma})\mathbb{1}(j = \sigma(i)).$$

Proof. It suffices to notice that a non-zero Boolean matrix has operator norm 1 if and only if it has at most one non-zero entry in every row and in every column. \Box

The unit norm Boolean matrix a in Lemma 3.3 will be denoted by a_{σ} and it is given by

(3.1)
$$a_{\sigma} = \sum_{i \in I_{\sigma}} e_{i,\sigma(i)} = \sum_{j \in J_{\sigma}} e_{\sigma^{-1}(j),j}$$

where $e_{i,j}$ denotes the Boolean matrix whose all entries are zero except the (i, j)-entry equals 1. Let $S_{bool}(\ell^2)$ denote the set of all unit norm Boolean operators in $B(\ell^2)$. The following lemma is simple and we omit its proof (the unit norm assumption is important).

Lemma 3.4. If $a_{\sigma}, a_{\tau} \in S_{bool}(\ell^2)$, then $a_{\sigma}^* \in S_{bool}(\ell^2)$ and $a_{\sigma}a_{\tau} \in S_{bool}(\ell^2) \cup \{0\}$.

Proof of Proposition 3.2. By a standard truncation-approximation argument, we may assume that \mathcal{F} is a family of unit norm Boolean matrices of a fixed finite size $d \times d$. Then, by Lemma 3.3, each Boolean matrix in \mathcal{F} corresponds to a bijection $\sigma : I_{\sigma} \to J_{\sigma}$ with I_{σ}, J_{σ} two subsets of $\{1, 2, \dots, d\}$. Hence we can write $\mathcal{F} = \{a_{\sigma}\}_{\sigma}$. Denote

$$T_m := \left\| \sum_{x \in \mathcal{F}} c_x \otimes x^{\otimes m} \right\| = \left\| \sum_{\sigma} c_{\sigma} \otimes a_{\sigma}^{\otimes m} \right\|.$$

Our goal is to prove the equality

$$T_m = T_1$$

By the classical formula for spectral radius, we have

$$T_{m} = \lim_{n \to \infty} \left\{ \operatorname{Tr} \left[\left(\sum_{\sigma} c_{\sigma} \otimes a_{\sigma}^{\otimes m} \right)^{*} \left(\sum_{\sigma} c_{\sigma} \otimes a_{\sigma}^{\otimes m} \right) \right]^{n} \right\}^{1/2n} \\ = \lim_{n \to \infty} \left\{ \sum_{\substack{\sigma_{1}, \cdots, \sigma_{n}, \\ \tau_{1}, \cdots, \tau_{n}}} \operatorname{Tr} (c_{\sigma_{1}}^{*} c_{\tau_{1}} \cdots c_{\sigma_{n}}^{*} c_{\tau_{n}}) \cdot [\operatorname{Tr} (a_{\sigma_{1}}^{*} a_{\tau_{1}} \cdots a_{\sigma_{n}}^{*} a_{\tau_{n}})]^{m} \right\}^{1/2n}$$

By Lemma 3.4, for any $n \ge 1$ and any $\sigma_1, \dots, \sigma_n, \tau_1, \dots, \tau_n$, either $a_{\sigma_1}^* a_{\tau_1} \dots a_{\sigma_n}^* a_{\tau_n} = 0$ or $a_{\sigma_1}^* a_{\tau_1} \dots a_{\sigma_n}^* a_{\tau_n} = a_{\sigma}$ for a certain bijection $\sigma : I_{\sigma} \to J_{\sigma}$. Therefore,

$$\operatorname{Tr}(a_{\sigma_1}^*a_{\tau_1}\cdots a_{\sigma_n}^*a_{\tau_n})\in\{0,1,\cdots,d\}.$$

Under any one of the conditions (c1), (c2) and (c3), we have

$$\mathrm{Tr}(c_{\sigma_1}^*c_{\tau_1}\cdots c_{\sigma_n}^*c_{\tau_n})\geq 0.$$

Consequently, using $\lim_{n\to\infty} d^{(m-1)/2n} = 1$, we obtain

$$T_m \leq \limsup_{n \to \infty} \left\{ d^{m-1} \sum_{\substack{\sigma_1, \cdots, \sigma_n, \\ \tau_1, \cdots, \tau_n}} \operatorname{Tr}(c^*_{\sigma_1} c_{\tau_1} \cdots c^*_{\sigma_n} c_{\tau_n}) \cdot \operatorname{Tr}(a^*_{\sigma_1} a_{\tau_1} \cdots a^*_{\sigma_n} a_{\tau_n}) \right\}^{1/2n}$$
$$= \limsup_{n \to \infty} \left\{ \sum_{\substack{\sigma_1, \cdots, \sigma_n, \\ \tau_1, \cdots, \tau_n}} \operatorname{Tr}(c^*_{\sigma_1} c_{\tau_1} \cdots c^*_{\sigma_n} c_{\tau_n}) \cdot \operatorname{Tr}(a^*_{\sigma_1} a_{\tau_1} \cdots a^*_{\sigma_n} a_{\tau_n}) \right\}^{1/2n} = T_1.$$

Conversely, consider the diagonal subspace $D_m := \operatorname{span}\{e_i^{\otimes m} : 1 \leq i \leq d\} \subset (\mathbb{C}^d)^{\otimes m}$ and the corresponding orthogonal projection $P_m : (\mathbb{C}^d)^{\otimes m} \to D_m$. Let $U_m : \mathbb{C}^d \to D_m$ be the natural unitary operator sending each e_i to $e_i^{\otimes m}$. Take any unit norm Boolean matrix $a_{\sigma} \in \mathcal{F}$ associated to a bijection $\sigma : I_{\sigma} \to J_{\sigma}$. Using (3.1), we obtain

$$a_{\sigma}^{\otimes m}(e_i^{\otimes m}) = (a_{\sigma}e_i)^{\otimes m} = (\mathbb{1}(i \in J_{\sigma})e_{\sigma^{-1}(i)})^{\otimes m} = \mathbb{1}(i \in J_{\sigma}) \cdot e_{\sigma^{-1}(i)}^{\otimes m}, \quad i = 1, \cdots, d.$$

Hence $U_m^{-1}P_m(a_{\sigma}^{\otimes m})P_mU_m = a_{\sigma}$ and

$$\sum_{\sigma} c_{\sigma} \otimes a_{\sigma} = \sum_{\sigma} c_{\sigma} \otimes U_m^{-1} P_m(a_{\sigma}^{\otimes m}) P_m U_m = (Id \otimes U_m^{-1} P_m) \left(\sum_{\sigma} c_{\sigma} \otimes a_{\sigma}^{\otimes m} \right) (Id \otimes P_m U_m).$$
follows immediately that $T_m > T_1$.

It follows immediately that $T_m \ge T_1$.

3.3. Examples and counter-examples.

Example 3.1. Let $e_{i,j} \in B(\ell^2)$ denote the operator corresponding to the elementary matrix whose all entries are zero except the (i, j)-entry equals 1. It is a standard fact that $\{e_{i,j} : (i, j) \in \mathbb{N}^2\} \subset$ $B(\ell^2)$ has SAP.

Example 3.2 (SAP is not preserved by direct sum). Given any two operators x, y, define $x \oplus y =$ $\begin{bmatrix} x & 0 \\ 0 & y \end{bmatrix}$. Consider two families with SAP: $\{x_1 = 1, x_2 = 1, x_3 = 0\} \subset \mathbb{C}$ and $\{y_1 = 1, y_2 = 0, y_3 = 0\}$ 1} $\subset \mathbb{C}$. Note that the family

(3.2)
$$\left\{ z_1 = x_1 \oplus y_1 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, z_2 = x_2 \oplus y_2 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, z_3 = x_3 \oplus y_3 = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}$$

does not have SAP, since $0 = ||z_1 - z_2 - z_3|| \neq ||z_1 \otimes z_1 - z_2 \otimes z_2 - z_3 \otimes z_3|| = 1$.

Example 3.3 (SAP is not preserved by spatial compression). Consider the family

(3.3)
$$\{X_1 = \operatorname{diag}(1,1,1,1), X_2 = \operatorname{diag}(1,0,1,0), X_3 = \operatorname{diag}(0,1,1,0)\}.$$

It has SAP since for any b_1, b_2, b_3 in a C^* -algebra B,

$$\left\|\sum_{i=1}^{3}b_{i}\otimes X_{i}\right\|=\left\|\sum_{i=1}^{3}b_{i}\otimes X_{i}\otimes X_{i}\right\|=\max_{\varepsilon,\varepsilon'\in\{0,1\}}\|b_{1}+\varepsilon b_{2}+\varepsilon' b_{3}\|.$$

Note that the non-SAP family (3.2) is a spatial compression of the SAP family (3.3).

Example 3.4 (*Regular representation system has SAP*). Let *G* be a discrete group and $\lambda_G : G \to B(\ell^2(G))$ be its left regular representation. We call $\{\lambda_G(g)\}_{g\in G}$ the regular representation system of *G*. The standard Fell's absorption principle (see [Pis03, Prop. 8.1]) implies that $\{\lambda_G(g)\}_{g\in G}$ has SAP. Recall the definition (1.1) for the Hankel operators. One can check that $\Gamma_g^G = \lambda_G(g)U_{flip}$ with U_{flip} the unitary operator in $B(\ell^2(G))$ sending each δ_s to $\delta_{s^{-1}}$ for all $s \in G$. It follows that $\{\Gamma_g^G\}_{g\in G}$ has SAP.

Example 3.5 (*A non-SAP Hankel system inducing by a multicolored checkerboard*). The multicolored checkerboard in Figure 2 has the property that every color appears at most once in each column and each row and it corresponds to a coordinatewise injective (see Definition 4 below) map

 $\Phi: \{1,2,3\} \times \{1,2,3\} \rightarrow \{\text{red, orange, blue, purple, grey}\}.$



FIGURE 2. A multicolored checkerboard.

This colored checkerboard induces a Hankel system in the C^* -algebra $M_3(\mathbb{C})$ of all 3×3 complex matrices:

 $\{\Gamma_c^{\Phi} | c \in \{\text{red, orange, blue, purple, grey}\}\} \subset M_3(\mathbb{C}).$

For instance, $\Gamma^{\Phi}_{\text{orange}} = e_{1,2} + e_{2,3}$, $\Gamma^{\Phi}_{\text{blue}} = e_{1,3} + e_{2,1}$, etc. Note that the coordinatewise injectivity of Φ implies that all Γ^{Φ}_c are of unit norm. The self-tensorized system

$$\{\Gamma_c^{\Phi}\otimes\Gamma_c^{\Phi}\}_c\subset M_3(\mathbb{C})\otimes M_3(\mathbb{C})\simeq M_9(\mathbb{C})$$

is induced by the multicolored checkerboard in Figure 3. To verify that $\{\Gamma_c^{\Phi}\}_c$ is non-SAP, we



FIGURE 3. The tensorized multicolored checkerboard.

consider the following substitution

(red, orange, blue, purple, grey) \rightarrow (4,2,-1,0,0).

More precisely, set

$$A := 4\Gamma_{\text{red}}^{\Phi} + 2\Gamma_{\text{orange}}^{\Phi} - \Gamma_{\text{blue}}^{\Phi} = \begin{bmatrix} 4 & 2 & -1 \\ -1 & 4 & 2 \\ 0 & 0 & 4 \end{bmatrix}.$$

Then the tensorized matrix is given by

$$B := 4\Gamma^{\Phi}_{\text{red}} \otimes \Gamma^{\Phi}_{\text{red}} + 2\Gamma^{\Phi}_{\text{orange}} \otimes \Gamma^{\Phi}_{\text{orange}} - \Gamma^{\Phi}_{\text{blue}} \otimes \Gamma^{\Phi}_{\text{blue}} \xrightarrow{column-row}_{permutations} \begin{bmatrix} A & 0 & 0 \\ 0 & A_1 & 0 \\ 0 & 0 & A_2 \end{bmatrix},$$

where

$$A_1 = \begin{bmatrix} 4 & 2 & -1 \\ 0 & 4 & 0 \\ 0 & 0 & 4 \end{bmatrix}, \quad A_2 = \begin{bmatrix} 4 & 0 & 0 \\ -1 & 4 & 2 \\ 0 & 0 & 4 \end{bmatrix}.$$

It can be checked that $||A|| = 3\sqrt{3}$ and $||A_1|| = ||A_2|| = \sqrt{\frac{1}{2}} \left(\sqrt{345} + 37\right) > 3\sqrt{3}$. Hence $||B|| = \max\{||A||, ||A_1||, ||A_2||\} > ||A||$. This implies that $\{\Gamma_c^{\Phi}\}_c$ does not have SAP.

4. HANKEL SYSTEMS ON MONOIDS

All monoids below will be assumed to be cancellative. For the reader's convenience, we recall the following definitions and give a new definition of *hereditary SAP monoids*:

• Lunar monoids (Definition 2): a monoid \mathcal{M} is called a lunar monoid if it satisfies the lunar condition :

$$(ax = by, cx = dy, az = bw \text{ in } \mathcal{M}) \Longrightarrow (cz = dw \text{ in } \mathcal{M}).$$

• Hankel system on a monoid \mathcal{M} : by the Hankel system on \mathcal{M} , we mean the family $\{\Gamma_t^{\mathcal{M}}\}_{t \in \mathcal{M}} \subset B(\ell^2(\mathcal{M}))$ consisting of Hankel operators $\Gamma_t^{\mathcal{M}}$ defined by

$$\Gamma_t^{\mathcal{M}}(s_1, s_2) = \mathbb{1}(s_1 s_2 = t), \quad s_1, s_2 \in \mathcal{M}.$$

- SAP monoids (Definition 3): a monoid *M* is called a SAP monoid if its associated Hankel system {Γ_t^M}_{t∈M} ⊂ B(ℓ²(M)) has SAP.
 Hereditary SAP monoids: we say that a monoid *M* is a hereditary SAP monoid if for
- Hereditary SAP monoids: we say that a monoid *M* is a hereditary SAP monoid if for any subsets S₁, S₂ ⊂ *M*, the compressed system

$$\{P_{S_1}\Gamma_t^{\mathcal{M}}P_{S_2}\}_{t\in\mathcal{M}}$$

has SAP, where P_{S_1} , P_{S_2} are othogonal projections from $\ell^2(\mathcal{M})$ onto $\ell^2(S_1)$ and $\ell^2(S_2)$ respectively (see Definition 5 for the definition of hereditary SAP of a general system). In particular, any hereditary SAP monoid is a SAP monoid.

One of our main results-Theorem A-follows from the following

Theorem 4.1. Any lunar monoid is a hereditary SAP monoid.

The proof of Theorem 4.1 is postponed to §5.7.

In general, it seems to be quite non-trivial to determine whether a given monoid is a SAP monoid or not. However, we have the following stability result.

Proposition 4.2. Cartesian product preserves the class of SAP monoids.

Proof. Observe that $\Gamma_{(t_1,t_2)}^{\mathcal{M}_1 \times \mathcal{M}_2} = \Gamma_{t_1}^{\mathcal{M}_1} \otimes \Gamma_{t_2}^{\mathcal{M}_2}$. The result follows from Proposition 3.1.

Remark 4.1. In the sequel to this paper, we shall prove the following more involved fact:

free product preserves the class of SAP monoids.

In Problem 3, we asked whether the class of SAP monoids coincides with that of lunar ones. A more tractable open problem is

Problem 6. Does the class of hereditary SAP monoids coincides with that of lunar ones ?

Recall that the lunar condition of a monoid \mathcal{M} is a local condition: to check whether \mathcal{M} satisfies the lunar condition, it suffices to check the configurations of all 4×4 blocks in the multiplication table of \mathcal{M} . Therefore, a positive answer to Problem 6 would follow from a positive answer to Problem 7 in §5.

5. HANKEL SYSTEMS OF LUNAR MAPS

As explained in the introduction of this paper, the definition of Hankel systems determined by the multiplication tables of monoids can be extended to the ones determined by level sets of abstract two-variable maps. This section is devoted to such non-classical Hankel systems.

By generalizing the definition of lunar monoids (Definition 2), we introduce a notion of lunar maps (Definition 4). Then we prove that the Hankel system associated to any lunar map has SAP (see Theorem B) and in fact has a stronger hereditary SAP (see Definition 5 and Corollary 5.1).

Given any lunar map, we are able to construct a coupled discrete foliation decomposition (see the equalities (5.8) below). And this coupled foliation structure allows us to obtain simultaneous block-diagonalization of all the tensorized operators of the associated Hankel system (see the discussion in the beginning of §5.3).

Definition 4 (Lunar maps). Consider three sets $\mathcal{A}, \mathcal{X}, \mathcal{L}$. A two-variable map $\Phi : \mathcal{A} \times \mathcal{X} \to \mathcal{L}$ is called a *lunar map* if

- (i) Φ is *coordinatewise injective*, that is, the map $\Phi(a, \cdot)$ is injective on \mathcal{X} for each $a \in \mathcal{A}$ and so is the map $\Phi(\cdot, x)$ on \mathcal{A} for each $x \in \mathcal{X}$.
- (ii) Φ satisfies the *generalized lunar condition*: for any $a, b, c, d \in A$ and $x, y, z, w \in X$, the following implication holds:

(5.1)
$$\left(\Phi(a,x) = \Phi(b,y), \Phi(c,x) = \Phi(d,y), \Phi(a,z) = \Phi(b,w)\right) \Longrightarrow \left(\Phi(c,z) = \Phi(d,w)\right).$$

Given a coordinatewise injective map $\Phi : \mathcal{A} \times \mathcal{X} \to \mathcal{L}$, by the Hankel system associated to the map Φ , we mean the family $\{\Gamma_{\ell}^{\Phi}\}_{\ell \in \mathcal{L}} \subset B(\ell^2(\mathcal{X}), \ell^2(\mathcal{A}))$ consisting of operators Γ_{ℓ}^{Φ} :

(5.2)
$$\Gamma^{\Phi}_{\ell}(a,x) = \mathbb{1}(\Phi(a,x) = \ell), \quad (a,x) \in \mathcal{A} \times \mathcal{X}$$

Note that Φ is coordinatewise injective if and only if the operators $\Gamma_{\ell}^{\Phi} \in B(\ell^2(\mathcal{X}), \ell^2(\mathcal{A}))$ are of unit norm $\|\Gamma_{\ell}^{\Phi}\| = 1$ for all ℓ in the image of Φ .

Theorem B. For any lunar map $\Phi : \mathcal{A} \times \mathcal{X} \to \mathcal{L}$, the Hankel system $\{\Gamma_{\ell}^{\Phi}\}_{\ell \in \mathcal{L}}$ has SAP.

One may wonder whether the lunar condition on a two-variable map Φ is also necessary for the Hankel system $\{\Gamma_{\ell}^{\Phi}\}_{\ell}$ to have SAP. However, even for maps on finite sets of very small size, we are not able to answer this question.

Problem 7. Consider any the map Φ : {1,2,3,4} × {1,2,3,4} $\rightarrow \mathcal{L}$. Assume that the Hankel system of associated to Φ has SAP. Does it follow that Φ is a lunar map ?

5.1. **Hereditary SAP.** In general, the SAP is not preserved by compression (see Example 3.3). Therefore, we introduce a stronger notion of SAP in the following

Definition 5 (Hereditary SAP). A system $\{x_i\}_{i \in I} \subset B(\ell^2)$ is said to have hereditary SAP if for any subsets $S_1, S_2 \subset \mathbb{N}$, the compressed system

$$\{P_{S_1}x_iP_{S_2}\}_{i\in I}$$

has SAP. Here P_{S_1}, P_{S_2} are othogonal projections onto $\ell^2(S_1)$ and $\ell^2(S_2)$ respectively. More generally, the hereditary SAP for systems in $B(\ell^2(\mathcal{X}), \ell^2(\mathcal{A}))$ is defined in a similar way.

Corollary 5.1. For any lunar map $\Phi : \mathcal{A} \times \mathcal{X} \to \mathcal{L}$, the Hankel system $\{\Gamma_{\ell}^{\Phi}\}_{\ell \in \mathcal{L}}$ has hereditary SAP.

As a consequence, we prove that the regular representation system of any countable discrete group (see Example 3.4) has hereditary SAP. We mention that, it does not follow from the standard Fell's absorption principle. In particular, it implies [Kat23, Cor. 5.3].

Corollary 5.2. The regular representation system $\{\lambda_G(g)\}_{g\in G} \subset B(\ell^2(G))$ of any countable discrete group G has hereditary SAP.

Recall that the big Hankel operators on the Hardy spaces over higher dimensional or infinite dimensional torus \mathbb{T}^d , where *d* is an integer $d \ge 2$ or $d = \infty$ and \mathbb{T}^{∞} means $\mathbb{T}^{\mathbb{N}}$ (all the notation is classical and will be recalled in §6 below):

$$\Gamma^{\text{big}}_{\varphi}: H^2(\mathbb{T}^d) \xrightarrow{M_{\varphi}} L^2(\mathbb{T}^d) \xrightarrow{\text{orthogonal}} L^2(\mathbb{T}^d) \ominus H^2(\mathbb{T}^d),$$

where M_{φ} is initially densely defined by $M_{\varphi}(f) = \varphi f$ with $\varphi \in L^2(\mathbb{T}^d)$. By the big Hankel system, we mean

(5.3)
$$\{\Gamma^{\text{big}}_{e^{in\cdot\theta}}\}_{n\in\mathbb{Z}^d\setminus\mathbb{N}^d}\subset B\big(H^2(\mathbb{T}^d),L^2(\mathbb{T}^d)\ominus H^2(\mathbb{T}^d)\big),$$

when $d = \infty$, the index set $\mathbb{Z}^d \setminus \mathbb{N}^d$ should be replaced by $\mathbb{Z}^{(\mathbb{N})} \setminus \mathbb{N}^{(\mathbb{N})}$.

Corollary 5.3. For any integer $d \ge 2$ or $d = \infty$, the big Hankel system (5.3) has SAP.

5.2. **Constructions of lunar maps.** Here we give a convenient reformulation of the generalized lunar condition.

For any pair $(a,b) \in \mathcal{A}^2$, define the solution set of the equation $\Phi(a,x) = \Phi(b,y)$ by

(5.4)
$$\operatorname{Sol}_{\Phi}(a,b) := \{(x,y) \in \mathcal{X}^2 | \Phi(a,x) = \Phi(b,y) \}$$

Lemma 5.4. A map Φ satisfies the generalized lunar condition (5.1) if and only if for any $a, b, c, d \in A$, either $Sol_{\Phi}(a, b) = Sol_{\Phi}(c, d)$ or $Sol_{\Phi}(a, b) \cap Sol_{\Phi}(c, d) = \emptyset$.

The proof of Lemma 5.4 is direct and is omitted. This reformulation provides an algorithm for checking the generalized lunar condition.

We now turn to basic constructions of lunar maps.

Example 5.1 (Rational maps). Given $a, b, m, n \in \mathbb{Z}^*$, define $\Phi : \mathbb{N}^* \times \mathbb{N}^* \to \mathbb{Q}$ by

$$\Phi(x,y) = ax^m + by^n$$

Then Φ is a lunar map.

Example 5.2 (Tensor products). If two maps $\Phi_i : A_i \times X_i \to L_i$ (i = 1, 2) are lunar, then so is their tensor product $\Phi_1 \otimes \Phi_2$ given by

$$(\Phi_1 \otimes \Phi_2)((a_1, a_2), (x_1, x_2)) := (\Phi_1(a_1, x_1), \Phi_2(a_2, x_2))$$

Example 5.3 (Refinement of level sets). Given two maps defined on the same product space $\Phi_i : \mathcal{A} \times \mathcal{X} \to \mathcal{L}_i (i = 1, 2)$, define $\Phi : \mathcal{A} \times \mathcal{X} \to \mathcal{L}_1 \times \mathcal{L}_2$ by $\Phi(a, x) = (\Phi_1(a, x), \Phi_2(a, x))$. If Φ_1 and Φ_2 are lunar maps, then so is Φ .

Note that the level sets of Φ are refinement of level sets of both Φ_1 and Φ_2 and the Hankel system $\{\Gamma^{\Phi}_{(\ell_1,\ell_2)}\}_{(\ell_1,\ell_2)}$ is obtained as Hadamard-Schur products of operators from the two systems $\{\Gamma^{\Phi_1}_{\ell_1}\}_{\ell_1}$ and $\{\Gamma^{\Phi_2}_{\ell_2}\}_{\ell_2}$:

$$\Gamma^{\Phi}_{(\ell_1,\ell_2)}(a,x) = \Gamma^{\Phi_1}_{\ell_1}(a,x)\Gamma^{\Phi_2}_{\ell_2}(a,x).$$

Example 5.4 (Restrictions). Let $\Phi : \mathcal{A} \times \mathcal{X} \to \mathcal{L}$ be a lunar map. For any subsets $\mathcal{A}_1 \subset \mathcal{A}$ and $\mathcal{X}_1 \subset \mathcal{X}$, the restriction $\Phi|_{\mathcal{A}_1 \times \mathcal{X}_1} : \mathcal{A}_1 \times \mathcal{X}_1 \to \mathcal{L}$ is again a lunar map.

Example 5.5 (Transposition). Given a map $\Phi : \mathcal{A} \times \mathcal{X} \to \mathcal{L}$, its transposition Φ^t is defined by $\Phi^t(x,a) = \Phi(a,x)$. It can be checked that Φ is a lunar map if and only if so is Φ^t .

Example 5.6 (Group operations). For any group G, let $\Phi_1, \Phi_2 : G \times G \to G$ be defined by

(5.5)
$$\Phi_1(g_1,g_2) = g_1g_2 \text{ and } \Phi_2(g_1,g_2) = g_1g_2^{-1}, \quad \forall g_1,g_2 \in G.$$

Then both Φ_1 and Φ_2 are lunar maps.

Example 5.7 (Restriction of group operations). By the constructions in Examples 5.4 and 5.6, for any $S_1, S_2 \subset G$, the corresponding restricted maps $\Phi_1|_{S_1 \times S_2}, \Phi_2|_{S_1 \times S_2} : S_1 \times S_2 \to G$ of Φ_1, Φ_2 defined in (5.5) are lunar maps.

Example 5.8 (Lunar monoids). Let \mathcal{M} be a monoid and define $\Phi_{\mathcal{M}} : \mathcal{M} \times \mathcal{M} \to \mathcal{M}$ by

$$\Phi_{\mathcal{M}}(x,y) = xy.$$

It can be checked directly that $\Phi_{\mathcal{M}}$ is a lunar map if and only if \mathcal{M} is a lunar monoid in the sense of Definition 2.

Example 5.9 (Restriction of monoid operations). Let \mathcal{M} be a lunar monoid and let $S_1, S_2 \subset \mathcal{M}$. Then, by the constructions in Examples 5.4 and 5.8, the restriction onto the subset $S_1 \times S_2$ of the map $\Phi_{\mathcal{M}}$ defined in (5.6) is a lunar map. In particular, the class of lunar monoids is closed under the operation of taking the submonoids.

Example 5.10 (Group-embeddable monoids). By the constructions in Examples 5.4, 5.6 and 5.8, any group-embeddable monoid is a lunar monoid in the sense of Definition 2. Therefore, any cancellative Abelian monoid is a lunar monoid, since it can be embedded into its Grothendieck group (see, e.g., [BG09, Section 2.A, page 52]).

For non-lunar maps, we have

Example 5.11 (Non-lunar polynomials). The polynomial $P(x, y) = x^2 + y^2 + xy$ defines a non-lunar map on $\mathbb{N}^* \times \mathbb{N}^*$.

Therefore, it is natural to consider

Problem 8. Determine all integer-coefficient polynomials $P \in \mathbb{Z}[x, y]$ such that their restrictions on $\mathbb{N} \times \mathbb{N}$ (or $\mathbb{Z} \times \mathbb{Z}$) are lunar maps.

5.3. **Proof of Theorem B.** Fix a lunar map $\Phi : \mathcal{A} \times \mathcal{X} \to \mathcal{L}$ in the sense of Definition 4.

The proof is divided into the following ten steps.

Step 1. An equivalence relation on \mathcal{A}^2 . Recall the notation introduced in (5.4). The map Φ induces an equivalence relation on \mathcal{A}^2 given by

(5.7)
$$(a,b) \sim_{\Phi} (c,d) \iff \operatorname{Sol}_{\Phi}(a,b) = \operatorname{Sol}_{\Phi}(c,d).$$

Let $[\mathcal{A}^2] = \mathcal{A}^2 / \sim_{\Phi}$ be the corresponding quotient space whose general elements will be denoted as [a,b]. Note that all pairs (a,b) with $Sol_{\Phi}(a,b) = \emptyset$ are equivalent under the relation \sim_{Φ} . The corresponding equivalent class will be denoted by $[\star] := \{(a,b) \in \mathcal{A}^2 | \text{Sol}_{\Phi}(a,b) = \emptyset\}$. Set

$$[\mathcal{A}^2]^{\times} = [\mathcal{A}^2] \setminus \{[\star]\}$$

By Lemma 5.4, if $Sol_{\Phi}(a,b) \neq \emptyset$, then $(a,b) \sim_{\Phi} (c,d)$ if and only if $Sol_{\Phi}(a,b) \cap Sol_{\Phi}(c,d) \neq \emptyset$. Step 2. Coupled foliation decompositions. Define

$$(\mathcal{X}^2)_{\text{left}} := \{ (x, y) \in \mathcal{X}^2 | \exists (a, b) \, s.t. \, \Phi(a, x) = \Phi(b, y) \}.$$
$$(\mathcal{A}^2)_{\text{right}} := \{ (a, b) \in \mathcal{A}^2 | \exists (x, y) \, s.t. \, \Phi(a, x) = \Phi(b, y) \}.$$

Then we have the following coupled foliation decompositions:

$$(\mathcal{X}^2)_{\text{left}} = \bigsqcup_{[a,b] \in [\mathcal{A}^2]^{\times}} \text{Sol}_{\Phi}(a,b),$$
$$(\mathcal{A}^2)_{\text{right}} = \bigsqcup_{[a,b] \in [\mathcal{A}^2]^{\times}} [a,b] = \bigsqcup_{[a,b] \in [\mathcal{A}^2]^{\times}} \left\{ (c,d) \in \mathcal{A}^2 \big| (c,d) \sim_{\Phi} (a,b) \right\}.$$

The two decompositions (5.8) are coupled in the sense that their components are in natural one-to-one correspondance. This correspondance will be useful later.

Step 3. Coupled orthogonal decompositions. The coupled decompositions in (5.8) induce coupled orthogonal decompositions of $\ell^2(\mathcal{X}) \otimes_2 \ell^2(\mathcal{X})$ and $\ell^2(\mathcal{A}) \otimes_2 \ell^2(\mathcal{A})$ as follows. For any equivalence class $[a,b] \in [\mathcal{A}^2]^{\times}$, define a closed subspace of $\ell^2(\mathcal{X}) \otimes_2 \ell^2(\mathcal{X})$ by

$$\mathcal{H}_{\Phi}([a,b]) := \overline{\operatorname{span}}\{e_x \otimes e_y | (x,y) \in \operatorname{Sol}_{\Phi}(a,b)\} \subset \ell^2(\mathcal{X}) \otimes_2 \ell^2(\mathcal{X})$$

Clearly, if $[a,b] \neq [c,d]$ in $[\mathcal{A}^2]^{\times}$, then $\mathcal{H}_{\Phi}([a,b]) \perp \mathcal{H}_{\Phi}([c,d])$. Therefore,

$$\mathcal{H}_{\Phi} := \ell^2((\mathcal{X}^2)_{\mathrm{left}}) = igoplus_{[a,b] \in [\mathcal{A}^2]^{ imes}} \mathcal{H}_{\Phi}([a,b]) \subset \ell^2(\mathcal{X}) \otimes_2 \ell^2(\mathcal{X}).$$

Here $\bigoplus_{[a,b]\in[\mathcal{A}^2]^{\times}} \mathcal{H}_{\Phi}([a,b])$ denotes the closed subspace of $\ell^2(\mathcal{X}) \otimes_2 \ell^2(\mathcal{X})$ given by

$$\bigoplus_{[a,b]\in[\mathcal{A}^2]^{\times}}\mathcal{H}_{\Phi}([a,b]) = \overline{\operatorname{span}}\Big\{\bigcup_{[a,b]\in[\mathcal{A}^2]^{\times}}\mathcal{H}_{\Phi}([a,b])\Big\}$$

Denoting by $\mathcal{H}_{\Phi}^{\perp}$ the orthogonal complement of the subspace \mathcal{H}_{Φ} in $\ell^2(\mathcal{X}) \otimes_2 \ell^2(\mathcal{X})$, we obtain the orthogonal decomposition:

(5.9)
$$\ell^{2}(\mathcal{X}) \otimes_{2} \ell^{2}(\mathcal{X}) = \mathcal{H}_{\Phi}^{\perp} \oplus \mathcal{H}_{\Phi} = \mathcal{H}_{\Phi}^{\perp} \oplus \bigoplus_{[a,b] \in [\mathcal{A}^{2}]^{\times}} \mathcal{H}_{\Phi}([a,b]).$$

Similarly, for any $[a,b] \in [\mathcal{A}^2]^{\times}$, define a closed subspace of $\ell^2(\mathcal{A}) \otimes_2 \ell^2(\mathcal{A})$ by

(5.10)
$$\mathcal{K}_{\Phi}([a,b]) := \overline{\operatorname{span}}\{e_c \otimes e_d | (c,d) \sim_{\Phi} (a,b)\} \subset \ell^2(\mathcal{A}) \otimes_2 \ell^2(\mathcal{A}).$$

Then

$$\mathcal{K}_{\Phi} := \ell^2((\mathcal{A}^2)_{\mathrm{right}}) = \bigoplus_{[a,b] \in [\mathcal{A}^2]^{\times}} \mathcal{K}_{\Phi}([a,b]) = \overline{\mathrm{span}} \Big\{ \bigcup_{[a,b] \in [\mathcal{A}^2]^{\times}} \mathcal{K}_{\Phi}([a,b]) \Big\} \subset \ell^2(\mathcal{A}) \otimes_2 \ell^2(\mathcal{A})$$

(5.8)

and we obtain the orthogonal decomposition:

(5.11)
$$\ell^{2}(\mathcal{A}) \otimes_{2} \ell^{2}(\mathcal{A}) = \mathcal{K}_{\Phi}^{\perp} \oplus \mathcal{K}_{\Phi} = \mathcal{K}_{\Phi}^{\perp} \oplus \bigoplus_{[a,b] \in [\mathcal{A}^{2}]^{\times}} \mathcal{K}_{\Phi}([a,b]).$$

Step 4. The space $\mathcal{H}_{\Phi}^{\perp}$ is in the simultaneous kernel of all $\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi}$. We claim that

(5.12)
$$\mathcal{H}_{\Phi}^{\perp} \subset \bigcap_{\ell \in \mathcal{L}} \ker(\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi})$$

Indeed, note that

$$\mathcal{H}_{\Phi}^{\perp} = \overline{\operatorname{span}}\Big\{e_x \otimes e_y\Big|(x,y) \in \mathcal{X}^2 \setminus \bigcup_{[a,b] \in [\mathcal{A}^2]} \operatorname{Sol}_{\Phi}(a,b)\Big\}.$$

Fix any pair $(x,y) \in \mathcal{X}^2 \setminus \bigcup_{[a,b] \in [\mathcal{A}^2]} \operatorname{Sol}_{\Phi}(a,b)$. Assume by contradiction that there exists $\ell \in \mathcal{L}$ such that $(\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi})(e_x \otimes e_y) \neq 0$. Then by the definition (5.2) of Γ_{ℓ}^{Φ} , there exists $(a,b) \in \mathcal{A}^2$, such that $\Phi(a,x) = \Phi(b,y)$ and $(\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi})(e_x \otimes e_y) = e_a \otimes e_b$. But this implies that $(x,y) \in \operatorname{Sol}_{\Phi}(a,b)$ and we obtain a contradiction.

Step 5. Simultaneous block-diagonalization of all $\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi}$. We claim that, with respect to the orthogonal decompositions (5.9) and (5.11), all operators $\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi}$ for $\ell \in \mathcal{L}$ are simultaneously block-diagonalized. In notation, we write this simultaneous block-diagonalization as

(5.13)
$$\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi} = \left(\mathcal{H}_{\Phi}^{\perp} \xrightarrow{0} \mathcal{K}_{\Phi}^{\perp}\right) \oplus \bigoplus_{[a,b] \in [\mathcal{A}^2]^{\times}} \left(\mathcal{H}_{\Phi}([a,b]) \xrightarrow{\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi}} \mathcal{K}_{\Phi}([a,b])\right)$$

Here by simultaneous block-diagonalization, we mean that the orthogonal decompositions (5.9) and (5.11) are independent of $\ell \in \mathcal{L}$.

Indeed, by (5.12), we have

$$(\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi})|_{\mathcal{H}_{\Phi}^{\perp}} \equiv 0 \quad \text{for all } \ell \in \mathcal{L}.$$

It remains to show that for any $[a,b] \in [\mathcal{A}^2]^{\times}$,

(5.14)
$$(\Gamma^{\Phi}_{\ell} \otimes \Gamma^{\Phi}_{\ell})(\mathcal{H}_{\Phi}([a,b])) \subset \mathcal{K}_{\Phi}([a,b]) \quad \text{for all } \ell \in \mathcal{L}.$$

Equivalently, we need to show that for any $(x, y) \in Sol_{\Phi}(a, b)$,

$$(\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi})(e_x \otimes e_y) \in \mathcal{K}_{\Phi}([a,b]) \quad \text{ for all } \ell \in \mathcal{L}$$

But by the definition (5.2) of Γ_{ℓ}^{Φ} , either $(\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi})(e_x \otimes e_y) = 0$ (the zero-vector 0 clearly belongs to $\mathcal{K}_{\Phi}([a,b])$) or there exists $(c,d) \in \mathcal{A}^2$ such that $\Phi(c,x) = \Phi(d,y) = \ell$ and

$$(\Gamma^{\Phi}_{\ell} \otimes \Gamma^{\Phi}_{\ell})(e_x \otimes e_y) = e_c \otimes e_d.$$

Now the condition $\Phi(c,x) = \Phi(d,y) = \ell$ implies that $(x,y) \in \operatorname{Sol}_{\Phi}(c,d)$ and hence $(x,y) \in \operatorname{Sol}_{\Phi}(a,b) \cap \operatorname{Sol}_{\Phi}(c,d)$. Therefore, $\operatorname{Sol}_{\Phi}(a,b) \cap \operatorname{Sol}_{\Phi}(c,d) \neq \emptyset$. By assumption, Φ satisfies the generalized lunar condition, hence by Lemma 5.4, $\operatorname{Sol}_{\Phi}(a,b) = \operatorname{Sol}_{\Phi}(c,d)$ and then, by the definition (5.7) of the equivalence relation \sim_{Φ} , we obtain $(a,b) \sim_{\Phi} (c,d)$. Consequently, by the definition (5.10), we have $e_c \otimes e_d \in \mathcal{K}_{\Phi}([a,b])$. That is,

$$(\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi})(e_x \otimes e_y) = e_c \otimes e_d \in \mathcal{K}_{\Phi}([a,b])$$

This completes the proof of the desired relation (5.14).

Step 6. Restriction of $\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi}$ onto the diagonal subspace. Define the diagonal subspace of $\ell^2(\mathcal{X}) \otimes_2 \ell^2(\mathcal{X})$ and that of $\ell^2(\mathcal{A}) \otimes_2 \ell^2(\mathcal{A})$ as

$$\mathcal{H}^{\text{diag}} := \overline{\text{span}} \{ e_x \otimes e_x | x \in \mathcal{X} \} \subset \ell^2(\mathcal{X}) \otimes_2 \ell^2(\mathcal{X});$$

$$\mathcal{K}^{\text{diag}} := \overline{\text{span}} \{ e_a \otimes e_a | a \in \mathcal{A} \} \subset \ell^2(\mathcal{A}) \otimes_2 \ell^2(\mathcal{A}).$$

Let $U : \mathcal{H}^{\text{diag}} \to \ell^2(\mathcal{X})$ be the unitary operator such that $U(e_x \otimes e_x) = e_x$ for all $x \in \mathcal{X}$. Similarly, define a unitary operator $V : \mathcal{K}^{\text{diag}} \to \ell^2(\mathcal{A})$ such that $V(e_a \otimes e_a) = e_a$ for all $a \in \mathcal{A}$. Then we have the following simultaneous commutative diagram for all $\ell \in \mathcal{L}$ (here by simultaneous commutative diagram, we mean that the interwining operators U, V are independent of ℓ):

(5.15)
$$\begin{array}{c} \mathcal{H}^{\text{diag}} \xrightarrow{\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi}} \mathcal{K}^{\text{diag}} \\ U \downarrow \simeq \qquad \simeq \downarrow V \\ \ell^{2}(\mathcal{X}) \xrightarrow{\Gamma_{\ell}^{\Phi}} \ell^{2}(\mathcal{A}) \end{array}$$

Indeed, if $\Gamma_{\ell}^{\Phi}(e_x) = 0$, then $V(\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi})(e_x \otimes e_x) = 0 = (\Gamma_{\ell}^{\Phi}U)(e_x \otimes e_x)$. If $\Gamma_{\ell}^{\Phi}(e_x) \neq 0$, then there exists a unique $a \in \mathcal{A}$ such that $\Phi(a, x) = \ell, \Gamma_{\ell}^{\Phi}(e_x) = e_a$ and

$$[V(\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi})](e_x \otimes e_x) = V(e_a \otimes e_a) = e_a = \Gamma_{\ell}^{\Phi}(e_x) = (\Gamma_{\ell}^{\Phi}U)(e_x \otimes e_x)$$

Thus the commutative diagram (5.15) follows.

Remark 5.12. Note that the coordinatewise injectivity of Φ implies $Sol_{\Phi}(a, a) = \{(x, x) | x \in \mathcal{X}\}$ for any $a \in \mathcal{A}$ and $Sol_{\Phi}(a, b) \cap Sol_{\Phi}(a, a) = \emptyset$ for any distinct $a, b \in \mathcal{A}$. It follows that

$$\mathcal{H}^{\mathrm{diag}} = \mathcal{H}_{\Phi}([a,a]) \text{ and } \mathcal{K}^{\mathrm{diag}} = \mathcal{K}_{\Phi}([a,a]).$$

Step 7. Definitions of interwining operators for $\mathcal{H}_{\Phi}([a,b])$ and $\mathcal{K}_{\Phi}([a,b])$. For any $[a,b] \in [\mathcal{A}^2]^{\times}$, define a bounded linear operator $P^{[a,b]} : \mathcal{H}_{\Phi}([a,b]) \to \ell^2(\mathcal{X})$ such that

$$P^{[a,b]}(e_x \otimes e_y) = e_x \quad \text{for all } (x,y) \in \operatorname{Sol}_{\Phi}(a,b).$$

And define also a bounded linear operator $Q^{[a,b]}: \ell^2(\mathcal{A}) \to \mathcal{K}_{\Phi}([a,b])$ such that

(5.16)
$$Q^{[a,b]}(e_c) = \begin{cases} e_c \otimes e_d, & \text{if there exists } d \in \mathcal{A} \text{ such that } (a,b) \sim_{\Phi} (c,d) \\ 0, & \text{otherwise} \end{cases}$$

We need to check that the above $Q^{[a,b]}(e_c)$ is well-defined. Indeed, assume that $d_1, d_2 \in \mathcal{A}$ such that $(a,b) \sim_{\Phi} (c,d_1) \sim_{\Phi} (c,d_2)$. Since $[a,b] \in [\mathcal{A}^2]^{\times}$, we have $\operatorname{Sol}_{\Phi}(c,d_1) = \operatorname{Sol}_{\Phi}(c,d_2) = \operatorname{Sol}_{\Phi}(a,b) \neq \emptyset$. Take any $(x_0,y_0) \in \operatorname{Sol}_{\Phi}(c,d_1) = \operatorname{Sol}_{\Phi}(c,d_2) = \operatorname{Sol}_{\Phi}(a,b)$, then $\Phi(c,x_0) = \Phi(d_1,y_0)$ and $\Phi(c,x_0) = \Phi(d_2,y_0)$. Hence $\Phi(d_1,y_0) = \Phi(d_2,y_0)$. Consequently, by the coordinatewise injectivity assumption on Φ , we must have $d_1 = d_2$.

Clearly, both operators $P^{[a,b]}$ and $Q^{[a,b]}$ are partial isometries (unitary operators are also considered as partial isometries). In particular, their operator norms satisfy

(5.17)
$$||P^{[a,b]}|| \le 1 \text{ and } ||Q^{[a,b]}|| \le 1.$$

Step 8. Restriction of $\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi}$ onto $\mathcal{H}_{\Phi}([a,b])$. We claim that, for any $[a,b] \in [\mathcal{A}^2]^{\times}$, simultaneously for all $\ell \in \mathcal{L}$, the following diagram commutes (here again, we emphasize that, the

interwining operators $P^{[a,b]}$ and $Q^{[a,b]}$ are both independent of $\ell \in \mathcal{L}$):

(5.18)
$$\begin{array}{c} \mathcal{H}_{\Phi}([a,b]) \xrightarrow{\Gamma^{\Phi}_{\ell} \otimes \Gamma^{\Phi}_{\ell}} \mathcal{K}_{\Phi}([a,b]) \\ P^{[a,b]} \downarrow & \uparrow \mathcal{Q}^{[a,b]} \\ \ell^{2}(\mathcal{X}) \xrightarrow{\Gamma^{\Phi}_{\ell}} \ell^{2}(\mathcal{A}) \end{array}$$

Indeed, take any $\ell \in \mathcal{L}$. Then for any $(x, y) \in Sol_{\Phi}(a, b)$, one and only one of the following three cases happens:

(i) There does not exist $c \in A$ such that $\Phi(c,x) = \ell$. In this case, $\Gamma_{\ell}^{\Phi}(e_x) = 0$. Hence, on the one hand,

$$(\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi})(e_x \otimes e_y) = (\Gamma_{\ell}^{\Phi} e_x) \otimes (\Gamma_{\ell}^{\Phi} e_y) = 0$$

and on the other hand,

$$Q^{[a,b]}\Gamma^{\Phi}_{\ell}P^{[a,b]}(e_x\otimes e_y)=Q^{[a,b]}\Gamma^{\Phi}_{\ell}(e_x)=0.$$

(ii) There exists $c \in A$ with $\Phi(c, x) = \ell$ but there does not exist $d \in A$ with $\Phi(d, y) = \ell$. In this case, $\Gamma_{\ell}^{\Phi}(e_x) = e_c$ and $\Gamma_{\ell}^{\Phi}(e_y) = 0$. Moreover, there is no element $d \in A$ such that $(a,b) \sim_{\Phi} (c,d)$. Hence by the definition (5.16) of $Q^{[a,b]}$, we must have $Q^{[a,b]}(e_c) = 0$. Therefore, on the one hand,

$$(\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi})(e_x \otimes e_y) = (\Gamma_{\ell}^{\Phi} e_x) \otimes (\Gamma_{\ell}^{\Phi} e_y) = 0$$

and on the other hand,

$$Q^{[a,b]}\Gamma^{\Phi}_{\ell}P^{[a,b]}(e_x\otimes e_y)=Q^{[a,b]}\Gamma^{\Phi}_{\ell}(e_x)=Q^{[a,b]}(e_c)=0.$$

(iii) There exist $c, d \in A$ such that $\Phi(c, x) = \Phi(d, y) = \ell$ (in particular, it implies that $(a, b) \sim_{\Phi} (c, d)$). In this case, $\Gamma_{\ell}^{\Phi}(e_x) = e_c, \Gamma_{\ell}^{\Phi}(e_y) = e_d$ and $Q^{[a,b]}(e_c) = e_c \otimes e_d$. Consequently, on the one hand,

$$(\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi})(e_x \otimes e_y) = (\Gamma_{\ell}^{\Phi} e_x) \otimes (\Gamma_{\ell}^{\Phi} e_y) = e_c \otimes e_d$$

and on the other hand,

$$Q^{[a,b]}\Gamma^{\Phi}_{\ell}P^{[a,b]}(e_x\otimes e_y)=Q^{[a,b]}\Gamma^{\Phi}_{\ell}(e_x)=Q^{[a,b]}(e_c)=e_c\otimes e_d.$$

Clearly, in all these three cases, $(\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi})(e_x \otimes e_y) = Q^{[a,b]}\Gamma_{\ell}^{\Phi}P^{[a,b]}(e_x \otimes e_y)$. Since $\ell \in \mathcal{L}$ and $(x,y) \in \operatorname{Sol}_{\Phi}(a,b)$ are chosen arbitrarily, we prove that the diagram (5.18) commutes.

Step 9. The proof of the inequality $\|\sum_{\ell} a_{\ell} \otimes \Gamma_{\ell}^{\Phi}\| \leq \|\sum_{\ell} a_{\ell} \otimes \Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi}\|$. Take any finitely supported sequence $(a_{\ell})_{\ell \in \mathcal{L}}$ in an arbitrary unital C^* -algebra B with unit 1_B . By the simultaneous commutative diagram (5.15), we have

$$\begin{split} \sum_{\ell} a_{\ell} \otimes \Gamma_{\ell}^{\Phi} &= \sum_{\ell} a_{\ell} \otimes V[(\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi})|_{\mathcal{H}^{\text{diag}}}]U^{-1} \\ &= (1_{B} \otimes V) \Big(\sum_{\ell} a_{\ell} \otimes (\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi})|_{\mathcal{H}^{\text{diag}}} \Big) (1_{B} \otimes U^{-1}). \end{split}$$

Consequently,

$$\left\|\sum_{\ell} a_{\ell} \otimes \Gamma_{\ell}^{\Phi}\right\| = \left\|\sum_{\ell} a_{\ell} \otimes (\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi})|_{\mathcal{H}^{\text{diag}}}\right\| \leq \left\|\sum_{\ell} a_{\ell} \otimes \Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi}\right\|$$

Step 10. The proof of the converse inequality $\|\sum_{\ell} a_{\ell} \otimes \Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi}\| \leq \|\sum_{\ell} a_{\ell} \otimes \Gamma_{\ell}^{\Phi}\|$. Take any finitely supported sequence $(a_{\ell})_{\ell \in \mathcal{L}}$ in an arbitrary unital C^* -algebra B with unit 1_B . The simultaneous block-diagonalization (5.13) together with the commutative diagram (5.18) and the contractivity of all the interwining operators $P^{[a,b]}, Q^{[a,b]}$ in (5.17) imply that

$$\begin{split} \left\| \sum_{\ell} a_{\ell} \otimes \Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi} \right\| &= \sup_{[a,b] \in [\mathcal{A}^2]^{\times}} \left\| \sum_{\ell} a_{\ell} \otimes \left(\mathcal{H}_{\Phi}([a,b]) \xrightarrow{\Gamma_{\ell}^{\Phi} \otimes \Gamma_{\ell}^{\Phi}} \mathcal{K}_{\Phi}([a,b]) \right) \right\| \\ &= \sup_{[a,b] \in [\mathcal{A}^2]^{\times}} \left\| \sum_{\ell} a_{\ell} \otimes (\mathcal{Q}^{[a,b]} \Gamma_{\ell}^{\Phi} P^{[a,b]}) \right\| \\ &= \sup_{[a,b] \in [\mathcal{A}^2]^{\times}} \left\| (\mathbf{1}_B \otimes \mathcal{Q}^{[a,b]}) \left(\sum_{\ell} a_{\ell} \otimes \Gamma_{\ell}^{\Phi} \right) (\mathbf{1}_B \otimes P^{[a,b]}) \right\| \\ &\leq \left\| \sum_{\ell} a_{\ell} \otimes \Gamma_{\ell}^{\Phi} \right\|. \end{split}$$

This completes the proof of the converse inequality.

We thus complete the whole proof of Theorem **B**.

5.4. **Proof of Corollary 5.1.** Corollary 5.1 follows immediately from Theorem B by using Example 5.4: the lunar map is preserved by restriction.

5.5. **Proof of Corollary 5.2.** Corollary 5.2 follows from Corollary 5.1. Indeed, for any group *G*, the map $\Phi : G \times G \to G$ defined by $\Phi(a, x) = ax^{-1}$ is a lunar map. Moreover, the corresponding operator Γ_g^{Φ} coincides with $\lambda_G(g)$ for any $g \in G$.

5.6. **Proof of Corollary 5.3.** For any integer $d \ge 2$, by using the standard orthonormal basis $\{e^{in\cdot\theta}\}_{n\in\mathbb{Z}^d}$ of $L^2(\mathbb{T}^d)$, one can easily show that the big Hankel system (5.3) is unitarily equivalent to the compression of the regular representation of the group \mathbb{Z}^d . When $d = \infty$, we only need to replace T^d by $\mathbb{T}^{\mathbb{N}}$ and replace \mathbb{Z}^d by the countable group $\mathbb{Z}^{(\mathbb{N})}$. Hence Corollary 5.3 follows from Corollary 5.2.

5.7. **Proof of Theorem 4.1.** By the construction of lunar maps via lunar monoids (see Example 5.8), Theorem 4.1 follows immediately from Corollary 5.1.

6. APPLICATIONS

6.1. Notation and preliminaries.

6.1.1. *Notation.* Let $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ be the unit disk endowed with the normalized area measure $dA(z) = \frac{dxdy}{\pi}$ and $\mathbb{T} = \partial \mathbb{D}$ be the unit circle endowed with the normalized Haar measure *dm*. Recall that all the classical function spaces such as $L^p(\mathbb{T},m), H^p(\mathbb{T}), BMOA(\mathbb{T})$ on \mathbb{T} are simply denoted by:

$$L^p = L^p(\mathbb{T}, m), \quad H^p = H^p(\mathbb{T}), \quad \text{BMOA} = \text{BMOA}(\mathbb{T}).$$

Given any pair of Banach spaces $X, Y \subset H^1$ and a complex sequence $m = (m_n)_{n=0}^{\infty}$, the associated Fourier multiplier T_m (a priori densely defined) is the linear map:

$$X \ni \sum_{n=0}^{\infty} \widehat{f}(n) e^{in\theta} = f \xrightarrow{T_m} T_m f = f * \varphi_m = \sum_{n=0}^{\infty} m_n \widehat{f}(n) e^{in\theta} \in Y,$$

here the associated formal Fourier series $\varphi_m = \sum_{n \ge 0} m_n e^{in\theta}$ is called the symbol of T_m and will always be identified with T_m . The space of all bounded Fourier multipliers from X to Y is a Banach space with respect to the natural norm and will be denoted by

(X,Y).

If X and Y are both endowed with an o.s.s., then the space of all completely bounded (abbrev. cb) Fourier multipliers from X to Y, denoted by

$$(X,Y)_{cb},$$

is equipped with a natural o.s.s. through isometries

$$M_d((X,Y)_{cb}) = (X,M_d(Y))_{cb} \quad \text{for all } d \ge 1.$$

Here $M_d(Z)$ denotes the space of $d \times d$ matrices with entries in Z.

We shall also use the following notation:

- $M_m = M_m(\mathbb{C})$ for the set of complex matrices of size $m \times m$;
- $\ell_n^2 = \ell^2(\{1, 2, \cdots, n\});$
- $H_0^p = \{ f \in H^p | \widehat{f}(0) = 0 \};$
- $\overline{H^p} = \{\overline{f} | f \in H_p\}$ and similar notation like $\overline{H_0^{\infty}}$;
- S_p denotes the *p*-Schatten-von-Neumann class of operators on ℓ^2 for $1 \le p < \infty$ and S_{∞} denotes the class of compact operators on ℓ^2 ; in the case of $n \times n$ matrices, the corresponding spaces will be denoted by S_p^n . In particular, $S_{\infty}^n = M_n$;
- $H^1(S_1)$ denotes the space of S_1 -vector-valued H^1 -functions and similar notation like $L^{\infty}(B(\ell^2)), H^p(S_p)$ etc;
- The discrete non-commutative vector-valued L_p-spaces (introduced by Pisier in [Pis98, Chapter 1]) are denoted by S_p[E] (or Sⁿ_p[E]), where E is some operator space; recall Pisier's descriptions ([Pis98, Prop. 2.3]) on the complete bounded norms: let u : E → F be a map between two operator spaces, then, for any 1 ≤ p ≤ ∞,

(6.1)
$$||u||_{cb} = ||Id \otimes u : S_p[E] \to S_p[F]|| = \sup_{n \ge 1} ||Id \otimes u : S_p^n[E] \to S_p^n[F]||.$$

- For any two operator spaces E, F, we denote by CB(E, F) the space of completely bounded operators from E to F.
- For any *C**-algebra *A*, denoted by \overline{A} its complex conjugate. Under the non-canonical identification $\ell^2 \simeq \overline{\ell^2}$, we have $\overline{B(\ell^2)} \simeq B(\ell^2)$. In particular, if $a = [a_{ij}]_{i,j \in \mathbb{N}} \in B(\ell^2)$, we define $\overline{a} = [\overline{a}_{ij}]_{i,j \in \mathbb{N}} \in B(\ell^2)$. See [Pis20, Section 2.3].

6.1.2. *Nehari-Sarason-Page Theorem*. Let $\text{Hank}(\overline{H^2}, H^2) \subset B(\overline{H^2}, H^2)$ denote the set of all bounded Hankel operators obtained by

$$\Gamma_{\varphi}: \overline{H^2} \xrightarrow{M_{\varphi}} L^2 \xrightarrow{\mathcal{R}_+} H^2,$$

where M_{φ} is initially densely defined by $M_{\varphi}(f) = \varphi f$ with $\varphi \in L^2$ and \mathcal{R}_+ is the classical Riesz projection. With respect to the orthonormal basis $\{e^{-in\theta}\}_{n\geq 0}$ and $\{e^{in\theta}\}_{n\geq 0}$ for $\overline{H^2}$ and H^2 respectively, any operator in $\operatorname{Hank}(\overline{H^2}, H^2)$ has a matrix representation (here we slightly abuse the notation Γ):

$$\Gamma_a = [a_{i+j}]_{i,j \ge 0}$$
, where $a = (a_i)_{i \ge 0} \in \mathbb{C}^{\mathbb{N}}$ is a complex sequence.

In particular, we use identification: $\Gamma_{\varphi} = \Gamma_{(\widehat{\varphi}(n))_{n\geq 0}} = [\widehat{\varphi}(i+j)]_{i,j\geq 0}$. Recall that we denote by Hank (ℓ^2) the collection of all bounded $\Gamma_a \in B(\ell^2)$.

The Nehari-Sarason-Page Theorem [Neh57, Sar67, Pag70] asserts that, by equipping the space $L^{\infty}/\overline{H_0^{\infty}}$ with the quotient *o.s.s.*, the map $L^{\infty}/\overline{H_0^{\infty}} \ni \varphi \mapsto \Gamma_{\varphi} \in \text{Hank}(\overline{H^2}, H^2)$ is a complete isometric isomorphism. Throughout the whole paper, we will use the following equivalent norm on BMOA:

$$\|\varphi\|_{\text{BMOA}} := \|\varphi\|_{L^{\infty}/\overline{H_0^{\infty}}}.$$

This choice of the equivalent norm on BMOA will allow us to compute the precise cb-norms (not just the equivalent ones) of multipliers in various situations.

Thus, by equipping BMOA with the *o.s.s.* induced by $L^{\infty}/\overline{H_0^{\infty}}$, we have complete isometric identifications:

(6.3)
$$\begin{array}{cccc} \operatorname{BMOA} \simeq L^{\infty}/\overline{H_{0}^{\infty}} & \xrightarrow{\simeq} & \operatorname{Hank}(\overline{H^{2}}, H^{2}) & \xrightarrow{\simeq} & \operatorname{Hank}(\ell^{2}) \\ \varphi & \mapsto & \Gamma_{\varphi} & \mapsto & [\widehat{\varphi}(i+j)]_{i,j\geq 0} \end{array}$$

6.2. The multipliers in $(H^s, H^r)_{cb}$. Recall the definition of BMOA^(p) $(1 \le p \le \infty)$ introduced in (1.9) and (1.10).

Theorem 6.1. *For any* $q \in [2, \infty]$ *,*

$$BMOA^{(q')} \xrightarrow[inclusion]{contractive} (H^1, H^q)_{cb} = (H^{q'}, BMOA)_{cb} \quad with \ \frac{1}{q} + \frac{1}{q'} = 1.$$

More generally, for any $1 \le r \le 2 \le s < \infty$ *,*

(6.4)
$$\operatorname{BMOA}^{(u)} \xleftarrow{bounded}_{inclusion} (H^r, H^s)_{cb} \quad with \ \frac{1}{u} = \frac{1}{r} - \frac{1}{s}$$

Remark 6.1. The inclusion (6.4) can not be generalized to the case $s = \infty$ since

BMOA⁽²⁾
$$\not\subset H^2 = (H^2, H^\infty)_{cb} = (H^2, H^\infty).$$

6.2.1. *The end point situations:* $(H^1, BMOA)_{cb}$ and $(H^1, H^2)_{cb}$. Using the self-absorption property introduced in Definition 1 and Proposition 2.1, we may fix an *o.s.s.* on BMOA⁽²⁾ by requiring (for any integer $d \ge 1$)

(6.5)
$$\left\|\sum_{n=0}^{\infty}a_{n}\otimes e^{in\theta}\right\|_{M_{d}\otimes_{min}\mathrm{BMOA}^{(2)}} := \left\|\sum_{n=0}^{\infty}a_{n}\otimes\overline{a}_{n}e^{in\theta}\right\|_{L^{\infty}(\mathbb{T};M_{d}\otimes\overline{M}_{d})/\overline{H_{0}^{\infty}}(\mathbb{T};M_{d}\otimes\overline{M}_{d})}^{1/2},$$

where $(a_n)_{n>0}$ is any finitely supported sequence in M_d .

Proposition 6.2. We have the following complete isometric equalities:

$$(H^1, BMOA)_{cb} = BMOA \text{ and } (H^1, H^2)_{cb} = (H^2, BMOA)_{cb} = BMOA^{(2)}.$$

(i) Characterization of $(H^1, BMOA)_{cb}$. We proceed to the proof of the complete isometric equality $(H^1, BMOA)_{cb} = BMOA$ in Proposition 6.2. Namely, we shall prove

(6.6)
$$(H^1, M_d(BMOA))_{cb} \xrightarrow{isometric} M_d(BMOA) \text{ for all } d \ge 1.$$

Using the complete isometric isomorphism $(H^1)^* =$ BMOA and the definining property of *o.s.s.* on operator space dual (see [Pis03, Section 2.3]), we have

(6.7)
$$\operatorname{BMOA} \otimes_{\min} M_d(\operatorname{BMOA}) \xrightarrow[\text{isometric}]{\operatorname{complete}} CB(H^1, M_d(\operatorname{BMOA})).$$

We shall use the following Lemma 6.3 to reduce the computation of the cb-norm of any $T_m \in (H^1, M_d(BMOA))_{cb} \subset CB(H^1, M_d(BMOA))$ to the computation of the cb-norms of associated Fourier multipliers lying in the image of the embedding (6.7).

Given any sequence $m = (m_n)_{n=0}^{\infty}$ in M_d and any 0 < r < 1, let $m(r) = (m_n(r))_{n=0}^{\infty}$ denote the dilated sequence

$$(6.8) m_n(r) := r^n m_n, \quad n \in \mathbb{N}.$$

Lemma 6.3. For any multiplier $T_m : H^1 \to M_d(BMOA)$,

$$||T_m||_{(H^1, M_d(\text{BMOA}))_{cb}} = \sup_{0 < r < 1} ||T_{m(r)}||_{(H^1, M_d(\text{BMOA}))_{cb}}.$$

Proof. By (6.1) and the non-commutative Fubini theorem (see [Pis98, Prop. 2.1])

(6.9)
$$S_p[H^p] \xrightarrow{isometric} H^p(S_p), \quad 1 \le p < \infty,$$

we have

(6.10)
$$\|T_m\|_{(H^1, M_d(BMOA))_{cb}} = \|Id \otimes T_m : S_1[H^1] \to S_1[M_d(BMOA)]\|$$
$$= \sup\left\{ \left\| \sum_{n=0}^{\infty} m_n \otimes \widehat{f}(n) e^{in\theta} \right\|_{S_1[M_d(BMOA)]} \right\| \|f\|_{H^1(S_1)} < 1 \right\}.$$

Note that for any $g \in H^1(S_1)$ with $||g||_{H^1(S_1)} < 1$,

$$\lim_{r\to 1^-} \left\|\sum_{n=0}^{\infty} r^n \widehat{g}(n) e^{in\theta} - \sum_{n=0}^{\infty} \widehat{g}(n) e^{in\theta}\right\|_{H^1(S_1)} = 0.$$

Hence the following subset is dense in the open unit ball of $H^1(S_1)$:

$$\bigcup_{0 < r < 1} \Big\{ P_r * g \Big| \|g\|_{H^1(S_1)} < 1 \Big\} \stackrel{\text{dense}}{\subset} \Big\{ f \Big| \|f\|_{H^1(S_1)} < 1 \Big\},$$

where $P_r * g$ is the vector-valued version of the standard Poisson convolution $P_r : L^1 \to L^1$ on \mathbb{T} given by

(6.11)
$$(P_r * h)(e^{i\theta}) = \sum_{n \in \mathbb{Z}} r^{|n|} \widehat{h}(n) e^{in\theta}$$

Therefore, the equality (6.10) implies

$$||T_m||_{(H^1, M_d(BMOA))_{cb}} = \sup_{0 < r < 1 \text{ and } ||g||_{H^1(S_1)} < 1} \left\| \sum_{n=0}^{\infty} m_n \otimes r^n \widehat{g}(n) e^{in\theta} \right\|_{S_1[M_d(BMOA)]}$$

For any 0 < r < 1, the equality (6.10) applied to m(r) implies that

$$\sup_{\|g\|_{H^{1}(S_{1})} < 1} \left\| \sum_{n=0}^{\infty} m_{n} \otimes r^{n} \widehat{g}(n) e^{in\theta} \right\|_{S_{1}[M_{d}(\text{BMOA})]} = \|T_{m(r)}\|_{(H^{1}, M_{d}(\text{BMOA}))_{cb}}.$$

We thus complete the whole proof.

Proof of the isometric equality (6.6). Clearly, if $T_m \in (H^1, M_d(BMOA))_{cb}$, then the sequence $m = (m_n)_{n \in \mathbb{N}}$ is bounded and hence

(6.12)
$$\sum_{n=0}^{\infty} \|m_n(r)\| < \infty \quad \text{for all } 0 < r < 1.$$

Thus $T_{m(r)}$ coincides with the image of following tensor under the embedding (6.7):

$$\sum_{n=0}^{\infty} e^{in\theta} \otimes m_n(r) \otimes e^{in\theta} \in \text{BMOA} \otimes_{\min} M_d(\text{BMOA}),$$

where $e^{in\theta}$ denotes the function $e^{i\theta} \mapsto e^{in\theta}$. It follows that

$$\|T_{m(r)}\|_{(H^1, \text{BMOA})_{cb}} = \left\|\sum_{n=0}^{\infty} e^{in\theta} \otimes m_n(r) \otimes e^{in\theta}\right\|_{\text{BMOA} \otimes_{min} M_d(\text{BMOA})}$$
(by (6.7))

$$= \left\| \sum_{n=0}^{\infty} \Gamma_n \otimes m_n(r) \otimes \Gamma_n \right\|_{\operatorname{Hank}(\ell^2) \otimes_{\min} M_d(\operatorname{Hank}(\ell^2))}$$
(by (6.3))

$$= \left\| \sum_{n=0}^{\infty} m_n(r) \otimes \Gamma_n \right\|_{M_d(\text{Hank}(\ell^2))}$$
(by Proposition 2.1)
$$= \left\| \sum_{n=0}^{\infty} m_n(r) \otimes e^{in\theta} \right\|_{M_d(\text{BMOA})}$$
(again by (6.3)).

Now by Lemma 2.6 and Lemma 6.3,

$$\|T_m\|_{(H^1, M_d(\text{BMOA}))_{cb}} = \left\|\sum_{n=0}^{\infty} m_n \otimes e^{in\theta}\right\|_{M_d(\text{BMOA})}$$

Conversely, if $\sum_{n=0}^{\infty} m_n \otimes e^{in\theta} \in M_d(BMOA)$, then $(m_n)_{n=0}^{\infty}$ is also bounded. We may repeat the above argument and obtain the desired isometric equality (6.6).

(ii) Characterization of $(H^1, H^2)_{cb}$. We now turn to the proof of the second complete isometric equality of Proposition 6.2:

$$(H^1, H^2)_{cb} = (H^2, \text{BMOA})_{cb} = \text{BMOA}^{(2)}$$

where the *o.s.s.* on H^2 is Pisier's operator Hilbert space determined by (1.5). In other words, we shall prove the following isometric equalities:

(6.13)
$$(H^1, M_d(H^2))_{cb} = (H^2, M_d(BMOA))_{cb} = M_d(BMOA^{(2)})$$
 for all $d \ge 1$,

where the norm on $M_d(BMOA^{(2)})$ is determined by (6.5).

Lemma 6.4. For any multiplier $T_m: H^1 \to M_d(H^2)$,

$$||T_m||_{(H^1, M_d(H^2))_{cb}} = \sup_{0 < r < 1} ||T_m(r)||_{(H^1, M_d(H^2))_{cb}}.$$

Proof. The proof follows almost verbatim that of Lemma 6.3.

Proof of the isometric equality (6.13). Fix any integer $d \ge 1$. By the duality $(H^1)^* = BMOA$, the equality $(H^1, M_d(H^2))_{cb} = (H^2, M_d(BMOA))_{cb}$ holds isometrically. Hence it suffices to prove the isometric equality $(H^1, M_d(H^2))_{cb} = M_d(BMOA^{(2)})$. Following along the same lines of the proof of the equality (6.6), by Proposition 2.1, we only need to prove that for any bounded sequence $m = (m_n)_{n=0}^{\infty}$ in M_d and any 0 < r < 1,

$$\|T_{m(r)}\|_{(H^1,M_d(H^2))_{cb}} = \left\|\sum_{n=0}^{\infty} m_n(r) \otimes \overline{m_n(r)} \otimes e^{in\theta}\right\|_{M_d \otimes_{min} \overline{M}_d \otimes_{min} \overline{B} MOA}^{1/2}$$

Now fix any bounded sequence $m = (m_n)_{n=0}^{\infty}$ in M_d and any 0 < r < 1. The condition (6.12) implies $T_{m(r)} \in (H^1, M_d(H^2))_{cb}$. It also implies that, under the complete isometric embedding

(6.14) BMOA
$$\otimes_{\min} M_d(H^2) = (H^1)^* \otimes_{\min} M_d(H^2) \xrightarrow{complete isometric}_{embedding} CB(H^1, M_d(H^2)),$$

the multiplier $T_{m(r)}$ corresponds to the tensor

$$\sum_{n=0}^{\infty} e^{in\theta} \otimes m_n(r) \otimes e^{in\theta} \in \text{BMOA} \otimes_{\min} M_d \otimes_{\min} H^2 = \text{BMOA} \otimes_{\min} M_d(H^2)$$

Therefore, we have

$$\|T_{m(r)}\|_{(H^1,M_d(H^2))_{cb}} = \left\|\sum_{n=0}^{\infty} e^{in\theta} \otimes m_n(r) \otimes e^{in\theta}\right\|_{\operatorname{BMOA}\otimes_{\min}M_d(H^2)}$$
(by (6.14))

$$= \left\| \sum_{n=0}^{\infty} \Gamma_n \otimes m_n(r) \otimes e^{in\theta} \right\|_{\operatorname{Hank}(\ell^2) \otimes_{\min} M_d(H^2)}$$
(by (6.3))

$$= \left\| \sum_{n=0}^{\infty} m_n(r) \otimes \overline{m_n(r)} \otimes \Gamma_n \otimes \Gamma_n \right\|_{M_d \otimes_{\min} \overline{M}_d \otimes_{\min} \operatorname{Hank}(\ell^2) \otimes_{\min} \operatorname{Hank}(\ell^2)}^{1/2} \qquad (by (1.5))$$

$$\| \sum_{n=0}^{\infty} m_n(r) \otimes \overline{m_n(r)} \otimes \Gamma_n \|_{M_d \otimes_{\min} \overline{M}_d \otimes_{\min} \operatorname{Hank}(\ell^2) \otimes_{\min} \operatorname{Hank}(\ell^2)} \qquad (by (1.5))$$

$$= \left\| \sum_{n=0}^{\infty} m_n(r) \otimes \overline{m_n(r)} \otimes \Gamma_n \right\|_{M_d \otimes_{\min} \overline{M}_d \otimes_{\min} \operatorname{Hank}(\ell^2)}$$
(by Proposition 2.1)
$$= \left\| \sum_{n=0}^{\infty} m_n(r) \otimes \overline{m_n(r)} \otimes e^{in\theta} \right\|_{M_d \otimes_{\min} \overline{M}_d \otimes_{\min} \operatorname{BMOA}}^{1/2}$$
(again by (6.3)).

This completes the whole proof.

6.2.2. *General situations*. For using the complex interpolation method, we now prove in Lemmas 6.5 and 6.6 that, the quantity $\|\cdot\|_{BMOA^{(p)}}$ given in (1.9) defines a norm on $BMOA^{(p)}$ for any $1 \le p < \infty$ and the family $\{BMOA^{(p)} : 1 \le p \le \infty\}$ forms a complex interpolation scale.

Lemma 6.5. Let $1 \le p < \infty$ and $q = p/(p-1) \in (1,\infty]$. Then for any $\varphi \in BMOA^{(p)}$ and $\psi \in BMOA^{(q)}$, we have $\|\varphi * \psi\|_{BMOA} \le \|\varphi\|_{BMOA^{(p)}} \|\psi\|_{BMOA^{(q)}}$. Moreover,

$$\|\boldsymbol{\varphi}\|_{\mathrm{BMOA}^{(p)}} = \sup\left\{\|\boldsymbol{\varphi} \ast \boldsymbol{\psi}\|_{\mathrm{BMOA}} : \|\boldsymbol{\psi}\|_{\mathrm{BMOA}^{(q)}} \le 1\right\}.$$

In particular, $\|\cdot\|_{BMOA^{(p)}}$ defines a norm on $BMOA^{(p)}$.

Proof. We give the proof for 1 , the proof for <math>p = 1 is similar. It suffices to show that for any sequences $(a_n)_{n=0}^{\infty}, (b_n)_{n=0}^{\infty}$ of complex numbers,

(6.15)
$$\left\|\sum_{n=0}^{\infty}a_{n}b_{n}e^{in\theta}\right\|_{\text{BMOA}} \leq \left\|\sum_{n=0}^{\infty}|a_{n}|^{p}e^{in\theta}\right\|_{\text{BMOA}}^{1/p} \cdot \left\|\sum_{n=0}^{\infty}|b_{n}|^{q}e^{in\theta}\right\|_{\text{BMOA}}^{1/q}$$

and

(6.16)
$$\left\|\sum_{n=0}^{\infty}|a_n|^p e^{in\theta}\right\|_{\text{BMOA}}^{1/p} = \sup\left\{\left\|\sum_{n=0}^{\infty}a_nb_ne^{in\theta}\right\|_{\text{BMOA}}: \left\|\sum_{n=0}^{\infty}|b_n|^q e^{in\theta}\right\|_{\text{BMOA}} \le 1\right\}.$$

By Nehari's Theorem, the inequality (6.15) is equivalent to

$$\|[a_{i+j}b_{i+j}]\| \le \|[|a_{i+j}|^p]\|^{1/p} \cdot \|[|b_{i+j}|^q]\|^{1/q},$$

where $[c_{i+j}] = [c_{i+j}]_{i,j\geq 0}$ denotes the infinite Hankel matrix and $\|\cdot\|$ denotes the operator norm of $B(\ell^2)$. Since $\|[a_{i+j}b_{i+j}]\| \le \|[|a_{i+j}b_{i+j}|]\|$, we may assume $a_n \ge 0$ and $b_n \ge 0$ for all $n \ge 0$. Now for non-negative coefficients, the operator $[a_{i+j}b_{i+j}]$ is self-adjoint and

$$||[a_{i+j}b_{i+j}]|| = \sup \sum_{i,j\geq 0} a_{i+j}b_{i+j}v_iv_j$$

where the supremum runs over the set of all vectors $v \in \ell^2$ with non-negative coefficients $v_n \ge 0$ and $||v||_{\ell^2} \le 1$. But for any such vector $v \in \ell^2$, by Hölder's inequality,

$$\sum_{i,j\geq 0} a_{i+j} b_{i+j} v_i v_j \leq \left(\sum_{i,j\geq 0} a_{i+j}^p v_i v_j\right)^{1/p} \left(\sum_{i,j\geq 0} b_{i+j}^q v_i v_j\right)^{1/q} \leq \|[a_{i+j}^p]\|^{1/p} \|[b_{i+j}^q]\|^{1/q}.$$

This completes the proof of the inequality (6.15).

Finally, by considering $b_n = \frac{\bar{a}_n}{|a_n|} |a_n|^{p-1} \mathbb{1}(a_n \neq 0)$, we obtain the equality (6.16)

Lemma 6.6. Let $1 \le p_0, p_1 \le \infty$ and $\alpha \in (0, 1)$. Set $1/p_{\alpha} = (1 - \alpha)/p_0 + \alpha/p_1$, then

$$(BMOA^{(p_0)}, BMOA^{(p_1)})_{\alpha} = BMOA^{(p_{\alpha})}$$
 with equal norms

Proof. Assume that $p_0 \neq p_1$ and hence $p_{\alpha} \in (1, \infty)$. Let $S = \{z \in \mathbb{C} | 0 < \Re(z) < 1\}$. Assume that $\|\sum_{n\geq 0} |m_n|^{p_{\alpha}} e^{in\theta}\|_{BMOA} < 1$. For any $z \in \overline{S}$, set

$$m_n(z) = \frac{m_n}{|m_n|} |m_n|^{p_\alpha(1-z)/p_0 + p_\alpha z/p_1} \mathbb{1}(m_n \neq 0), \quad n \ge 0.$$

Then $m_n(\alpha) = m_n$. Clearly, for any $t \in \mathbb{R}$, we have

$$\left\|\sum_{n\geq 0}m_n(it)e^{in\theta}\right\|_{\mathrm{BMOA}^{(p_0)}} < 1 \text{ and } \left\|\sum_{n\geq 0}m_n(1+it)e^{in\theta}\right\|_{\mathrm{BMOA}^{(p_1)}} < 1$$

The contractive embedding BMOA^{(p_α)} \subset (BMOA^{(p_0)}, BMOA^{(p_1)})_{α} then follows.

Conversely, assume that $\|\sum_{n\geq 0} m_n e^{in\theta}\|_{(BMOA^{(p_0)}, BMOA^{(p_1)})_{\alpha}} < 1$. Then there exist functions $m_n: \overline{S} \to \mathbb{C}$, analytic on S with $m_n(\alpha) = m_n$, and

$$\sup_{t\in\mathbb{R}}\left\|\sum_{n\geq 0}|m_n(it)|^{p_0}e^{in\theta}\right\|_{\text{BMOA}}<1 \text{ and } \sup_{t\in\mathbb{R}}\left\|\sum_{n\geq 0}|m_n(1+it)|^{p_1}e^{in\theta}\right\|_{\text{BMOA}}<1.$$

Now take any sequence $(a_n)_{n\geq 0}$ in \mathbb{C} with $\|\sum_{n\geq 0} |a_n|^{q_\alpha} e^{in\theta}\|_{BMOA} < 1$ and $q_\alpha = p_\alpha/(p_\alpha - 1)$. By the previous argument, we can construct functions $a_n : \overline{S} \to \mathbb{C}$, analytic on S with $a_n(\alpha) = a_n$, and

$$\sup_{t\in\mathbb{R}}\left\|\sum_{n\geq0}|a_n(it)|^{q_0}e^{in\theta}\right\|_{\text{BMOA}}<1 \text{ and } \sup_{t\in\mathbb{R}}\left\|\sum_{n\geq0}|a_n(1+it)|^{q_1}e^{in\theta}\right\|_{\text{BMOA}}<1,$$

where $q_0 = p_0/(p_0 - 1), q_1 = p_1/(p_1 - 1)$. Therefore, by the inequality (6.15),

$$\sup_{z\in\partial S}\left\|\sum_{n=0}^{\infty}a_n(z)m_n(z)e^{in\theta}\right\|_{\text{BMOA}}<1$$

It follows immediately that

$$\left\|\sum_{n=0}^{\infty}a_{n}m_{n}e^{in\theta}\right\|_{\text{BMOA}}<1.$$

Then, by applying the equality (6.16), we obtain $\|\sum_{n=0}^{\infty} m_n e^{in\theta}\|_{BMOA^{(p_\alpha)}} < 1$. The desired inverse contractive embedding $(BMOA^{(p_0)}, BMOA^{(p_1)})_{\alpha} \subset BMOA^{(p_\alpha)}$ now follows.

Proof of Theorem 6.1. Observe that, for any $1 \le r, s \le \infty$, the equality

$$(H^r, H^s)_{cb} = (H^r, L^s)_{cb}$$

holds completely isometrically. By Proposition 6.2, $BMOA^{(2)} = (H^1, H^2)_{cb} = (H^1, L^2)_{cb}$ with equal norms (in fact, the equalities hold completely isometrically). On the other hand,

$$BMOA^{(1)} \xrightarrow{contractive} BMOA = (H^1, L^{\infty})_{cb}$$

Thus, by the standard complex interpolation method, we obtain that

$$(BMOA^{(1)}, BMOA^{(2)})_{\theta} \xrightarrow{contractive} (H^1, (L^{\infty}, L^2)_{\theta})_{cb}$$
 for all $\theta \in [0, 1]$.

Therefore, by Lemma 6.6, for any $2 \le q \le \infty$, we have

(6.17)
$$\operatorname{BMOA}^{(q')} \xrightarrow{contractive} (H^1, L^q)_{cb} = (H^1, H^q)_{cb} \text{ with } \frac{1}{q} + \frac{1}{q'} = 1.$$

This completes the first assertion of Theorem 6.1.

Now we proceed to the proof of the bounded inclusion (6.4). By interpolating (6.17) with $BMOA^{(2)} = (H^2, BMOA)_{cb}$, we obtain

(6.18)
$$(BMOA^{(2)}, BMOA^{(q')})_{\theta} \xrightarrow{contractive} ((H^2, H^1)_{\theta}, (BMOA, H^q)_{\theta})_{cb}.$$

Observation: for any $1 < r < \infty$ and any $\theta \in (0, 1)$, we have

(6.19)
$$H^{r_{\theta}} \xrightarrow{complete isomorphism} (H^{1}, H^{r})_{\theta}$$

and its dual form

(6.20)
$$H^{r'_{\theta}} \xrightarrow{complete \ isomorphism} (BMOA, H^{r'})_{\theta},$$

where

$$r' = \frac{r}{r-1}, \quad \frac{1}{r_{\theta}} = \frac{1-\theta}{1} + \frac{\theta}{r} \text{ and } r'_{\theta} = \frac{r_{\theta}}{r_{\theta}-1}$$

Lemma 6.6 combined with (6.18), (6.19) and (6.20) implies

$$\mathsf{BMOA}^{(p_{\theta})} \xrightarrow[inclusion]{bounded}} (H^{r_{\theta}}, H^{s_{\theta}})_{cb}$$

where $\theta \in (0,1)$ and

(6.21)
$$\frac{1}{p_{\theta}} = \frac{1-\theta}{2} + \frac{\theta}{q'}, \quad \frac{1}{r_{\theta}} = \frac{1-\theta}{2} + \frac{\theta}{1}, \quad \frac{1}{s_{\theta}} = \frac{1-\theta}{\infty} + \frac{\theta}{q}$$

with $2 \le q \le \infty$ and $q' = q/(q-1) \in [1,2]$.



FIGURE 4. $(\frac{1}{r}, \frac{1}{s}) = (1 - \theta)(\frac{1}{2}, 0) + \theta(1, \frac{1}{q})$ with $q \in [2, \infty], \theta \in (0, 1)$.

Now from Figure 4, it is clear that for any pair of exponents (r, s) satisfying

(6.22)
$$\frac{1}{r} \ge \frac{1}{s} + \frac{1}{2} \text{ and } 1 < r < 2 < s < \infty,$$

we can find

$$\theta \in (0,1), r_{\theta} = r, s_{\theta} = s, \frac{1}{p_{\theta}} = \frac{1}{r} - \frac{1}{s} \text{ and } q \in [2,\infty],$$

satisfying the equation system (6.21). The desired bounded inclusion (6.4) follows for any pair of exponents (r, s) satisfying (6.22).

Note also that for r = 1 and any $2 \le s \le \infty$, the bounded inclusion (6.4) holds. Finally, by interpolating the above result with the trivial isometric equality $(H^2, H^2)_{cb} = \text{BMOA}^{(\infty)}$, we extend the bounded inclusion (6.4) to all pairs (r, s) with $1 \le r \le 2 \le s < \infty$.

Proof of the Observation. Indeed, by [Xu90, Thm. 4.3] (see also [BX91, Remark 5.3]),

(6.23)
$$(H^1(S_1), H^r(S_r))_{\theta} \xrightarrow{equivalent}_{norms} H^{r_{\theta}}(S_{r_{\theta}}).$$

On the other hand, by [Pis98, Cor. 1.4, formula (1.5)] and the equality (6.9), we have

(6.24)
$$(H^1(S_1), H^r(S_r))_{\theta} = (S_1[H^1], S_r[H^r])_{\theta} = S_{r_{\theta}}[(H^1, H^r)_{\theta}]$$
 with equal norms.

The relations (6.23) and (6.24) together imply

$$H^{r_{\theta}}(S_{r_{\theta}}) = S_{r_{\theta}}[H^{r_{\theta}}] = S_{r_{\theta}}[(H^{1}, H^{r})_{\theta}]$$
 with equivalent norms.

Consequently, by (6.1), we obtain (6.19).

For the dual form (6.20), since $H^r(S_r) = S_r[H^r]$ is reflexive for any $1 < r < \infty$, we may apply the duality theory for complex interpolation [Pis16, Thm. 8.37] to obtain the dual of $(S_1^n[H^1], S_r^n[H^r])_{\theta}$. Thus, by [Pis98, Cor. 1.4, formula (1.5)], for any $n \ge 1$,

$$S_{r_{\theta}'}^{n}[(\mathsf{BMOA}, H^{r'})_{\theta}] = (S_{\infty}^{n}[\mathsf{BMOA}], S_{r'}^{n}[H^{r'}])_{\theta} = \left((S_{1}^{n}[H^{1}], S_{r}^{n}[H^{r}])_{\theta}\right)^{*} \text{ with equal norms.}$$

Then by (6.23) and (6.24) and the classical result on the dual of $H^p(S_p)$, we have

$$S^{n}_{r'_{\theta}}[(\mathsf{BMOA}, H^{r'})_{\theta}] \xrightarrow{isomorphism} (H^{r_{\theta}}(S^{n}_{r_{\theta}}))^{*} \xrightarrow{isomorphism} H^{r'_{\theta}}(S^{n}_{r'_{\theta}}) = S^{n}_{r'_{\theta}}[H^{r'_{\theta}}],$$

where the norm equivalence constants in the above isomorphisms are uniformly bounded. Therefore, by (6.1), we complete the proof of the complete isomorphism (6.20).

This completes the whole proof of the theorem.

6.3. **Bounded Hankel operators induce cb-Fourier multipliers.** The second complete isometric equality in Proposition 6.2 has a higher dimensional analogue. Here we only state the isometric counterpart, the completely isometric counterpart requires a definition of the *o.s.s.* on the corresponding space and will not be treated here.

For any scalar function φ in $H^1(\mathbb{T}^d)$, set

$$\boldsymbol{\varphi}^{\dagger}(z_1,\cdots,z_d):=\overline{\boldsymbol{\varphi}(z_1,\cdots,z_d)}.$$

Corollary 6.7. A function $\varphi \in H^1(\mathbb{T}^d)$ induces a completely bounded Fourier multiplier in $(H^2(\mathbb{T}^d), \operatorname{Hank}(\overline{H^2(\mathbb{T}^d)}, H^2(\mathbb{T}^d)))_{cb}$ if and only if $\Gamma_{\varphi * \varphi^{\dagger}} \in \operatorname{Hank}(\overline{H^2(\mathbb{T}^d)}, H^2(\mathbb{T}^d))$. Moreover,

$$\|\varphi\|_{(H^2(\mathbb{T}^d),\operatorname{Hank}(\overline{H^2(\mathbb{T}^d)},H^2(\mathbb{T}^d)))_{cb}} = \|\Gamma_{\varphi\ast\varphi^\dagger}\|_{\operatorname{Hank}(\overline{H^2(\mathbb{T}^d)},H^2(\mathbb{T}^d))}^{1/2}$$

Proof. The proof follows almost verbatim that of the second isometric equality in Proposition 6.2, where the SAP of \mathbb{N} is replaced by the SAP of \mathbb{N}^d .

Corollary 6.8. Assume that $\varphi \in H^1(\mathbb{T}^d)$ with $\Gamma_{\varphi * \varphi^{\dagger}} \in \text{Hank}(\overline{H^2(\mathbb{T}^d)}, H^2(\mathbb{T}^d))$. Then for any sequence $(f_k)_{k \ge 1}$ in $H^2(\mathbb{T}^d)$, we have

$$\left\|\sum_{k=1}^{\infty} (\Gamma_{f_k \ast \varphi})^* \Gamma_{f_k \ast \varphi}\right\|_{B(\overline{H^2(\mathbb{T}^d)})}^{1/2} \leq \left\|\Gamma_{\varphi \ast \varphi^\dagger}\right\|_{\operatorname{Hank}(\overline{H^2(\mathbb{T}^d)}), H^2(\mathbb{T}^d))}^{1/2} \left(\sum_{k,l=1}^{\infty} |\langle f_k, f_l \rangle|^2\right)^{1/4}.$$

In particular, if $(f_k)_{k>1}$ are mutually orthogonal, then

$$\Big\|\sum_{k=1}^{\infty} (\Gamma_{f_{k}*\varphi})^{*} \Gamma_{f_{k}*\varphi} \Big\|_{B(\overline{H^{2}(\mathbb{T}^{d})})}^{1/2} \leq \|\Gamma_{\varphi*\varphi^{\dagger}}\|_{\operatorname{Hank}(\overline{H^{2}(\mathbb{T}^{d})},H^{2}(\mathbb{T}^{d}))}^{1/2} \Big(\sum_{k=1}^{\infty} \|f_{k}\|_{H^{2}(\mathbb{T}^{d})}^{4} \Big)^{1/4}.$$

Proof. Take any $\varphi \in H^1(\mathbb{T}^d)$ with $\Gamma_{\varphi * \varphi^{\dagger}} \in \text{Hank}(\overline{H^2(\mathbb{T}^d)}, H^2(\mathbb{T}^d))$. By Corollary 6.8, the Fourier multiplier T_{φ} belongs to $(H^2(\mathbb{T}^d), \text{Hank}(\overline{H^2(\mathbb{T}^d)}, H^2(\mathbb{T}^d)))_{cb}$ and we have

$$\left\| C \otimes_{\min} H^2(\mathbb{T}^d) \xrightarrow{Id \otimes T_{\varphi}} C \otimes_{\min} \operatorname{Hank}(\overline{H^2(\mathbb{T}^d)}, H^2(\mathbb{T}^d)) \right\| \leq \left\| \Gamma_{\varphi * \varphi^{\dagger}} \right\|_{\operatorname{Hank}(\overline{H^2(\mathbb{T}^d)}, H^2(\mathbb{T}^d))}^{1/2}.$$

Recall that for any operator space *E*, the tensor product $C \otimes_{min} E$ coincides with the Haagerup tensor product $C \otimes_h E$. Since $H^2(\mathbb{T}^d)$ is equipped with *OH* structure, then by applying the complex interpolation (see [Pis03, Thm. 5.22 and Cor. 7.11]), we have

$$C \otimes_h H^2(\mathbb{T}^d) \xrightarrow{complete} C \otimes_h (R, C)_{\frac{1}{2}} \xrightarrow{complete} (C \otimes_h R, C \otimes_h C)_{\frac{1}{2}}.$$

In the setting of Banach spaces, we have the isometric isomorphism

$$(C \otimes_h R, C \otimes_h C)_{\frac{1}{2}} \xrightarrow{isometric} S_4.$$

The above isometric isomorphism leads to the equality:

$$\|(f_1, f_2, \cdots, f_n, \cdots)^T\|_{C\otimes_{\min}H^2(\mathbb{T}^d)} = \left(\sum_{k,l=1} |\langle f_k, f_l \rangle|^2\right)^{1/4}$$

and we complete the whole proof.

6.4. Fourier-Schur multiplier inequalities with critical exponent 4/3.

Corollary 6.9. Let $\varphi \in BMOA^{(2)}$. Then for any ℓ^2 -column-vector-valued $f \in H^1(\ell^2)$,

$$\left\| \left(\int_{\mathbb{T}} \left[(f * \varphi) (f * \varphi)^* \right] dm \right)^{1/2} \right\|_{S_{4/3}} \le \|\varphi\|_{\mathrm{BMOA}^{(2)}} \cdot \|f\|_{H^1(\ell^2)}$$

Remark 6.2. The inequalities obtained in Corollary 6.9 are called Fourier-Schur multiplier inequalities since they can be rewritten as

$$\left\|\left(\widehat{\varphi}(0)\widehat{f}(0),\widehat{\varphi}(1)\widehat{f}(1),\cdots,\widehat{\varphi}(n)\widehat{f}(n),\cdots\right)\right\|_{\mathcal{S}_{4/3}} \leq \|\varphi\|_{\mathrm{BMOA}^{(2)}} \cdot \|f\|_{H^{1}(\ell^{2})}.$$

The exponent 4/3 in Corollary 6.9 is optimal, this can be seen from

Proposition 6.10. Let p > 0. Assume that there exist $q \in [1, \infty)$ and a numerical constant c > 0 such that for any $\varphi \in BMOA^{(2)}$ and any $f \in H^q(\ell^2)$,

$$\left\|\left(\int_{\mathbb{T}}\left[(f*\varphi)(f*\varphi)^*\right]dm\right)^{1/2}\right\|_{S_p} \leq c \|\varphi\|_{\mathrm{BMOA}^{(2)}} \cdot \|f\|_{H^q(\ell^2)}.$$

Then $p \ge 4/3$.

Proof of Corollary 6.9. Let *C* (resp. *R*) denote the column (resp. row) Hilbert space, which by definition, consists of all bounded operators in $B(\ell^2)$ whose all but the first column (resp. row) entries vanish.

Take any $\varphi \in \text{BMOA}^{(2)}$. By Proposition 6.2, the Fourier multiplier $T_{\widehat{\varphi}}$ associated with the sequence $(\widehat{\varphi}(n))_{n\geq 0}$ belongs to $(H^1, H^2)_{cb}$. Hence, by using the Haagerup tensor product \otimes_h (see [Pis03, Chapter 5]), we have

$$\|H^1 \otimes_h C \xrightarrow{T_{\widehat{\varphi}} \otimes Id} H^2 \otimes_h C\| \le \|T_{\widehat{\varphi}}\|_{(H^1, H^2)_{cb}} = \|\varphi\|_{\mathrm{BMOA}^{(2)}}.$$

By noticing the following relations (see [Pis98, p.18, formula (1.1) and Prop. 2.1])

$$H^1 \otimes_h C \xrightarrow{isometric}_{embedding} R \otimes_h H^1 \otimes_h C \xrightarrow{isometric}_{\simeq} S_1[H^1] \xrightarrow{isometric}_{\simeq} H^1(S_1),$$

we obtain an isometric isomorphism (as Banach spaces):

$$H^1 \otimes_h C \xrightarrow{isometric} H^1(\ell^2)$$

Now since H^2 is equipped with the *OH* structure, by applying the complex interpolation (see [Pis03, Thm. 5.22 and Cor. 7.11]), we have

$$H^2 \otimes_h C \xrightarrow{complete}_{isometric} (R,C)_{\frac{1}{2}} \otimes_h C \xrightarrow{complete}_{isometric} (R \otimes_h C, C \otimes_h C)_{\frac{1}{2}}.$$

In the setting of Banach spaces, we have the following isometric isomorphisms (in fact, the first one is completely isometric):

$$R \otimes_h C \xrightarrow{isometric} S_1$$
 and $C \otimes_h C \xrightarrow{isometric} S_2$,

hence

$$(R \otimes_h C, C \otimes_h C)_{\frac{1}{2}} \xrightarrow{isometric} (S_1, S_2)_{\frac{1}{2}} = S_{4/3}.$$

It follows that

$$\left\| H^{1}(\ell^{2}) \xrightarrow{isometric} H^{1} \otimes_{h} C \xrightarrow{T_{\widehat{\varphi}} \otimes Id} H^{2} \otimes_{h} C \xrightarrow{isometric} S_{4/3} \right\| \leq \left\| \varphi \right\|_{\mathrm{BMOA}^{(2)}}.$$

In other words, the following map

$$\begin{array}{cccc} H^{1}(\ell^{2}) & \longrightarrow & S_{4/3} \\ f & \mapsto & (\widehat{\varphi}(0)\widehat{f}(0), \cdots, \widehat{\varphi}(n)\widehat{f}(n), \cdots) \end{array}$$

is bounded with norm no larger than $\|\varphi\|_{BMOA^{(2)}}$. This completes the whole proof.

For proving Proposition 6.10, we need the following classical results on the lacunary Fourier series.

Lemma 6.11 (see, e.g., [BPc91, Prop. 3.1]). For any $q \in [1, \infty)$, there exists a numerical constant $c \ge 1$ such that for any sequence $(v_n)_{n=0}^{\infty}$ of vectors in ℓ^2 ,

$$\frac{1}{c} \Big(\sum_{n=0}^{\infty} \|v_n\|_{\ell^2}^2 \Big)^{1/2} \le \Big\| \sum_{n=0}^{\infty} v_n e^{i2^n \theta} \Big\|_{H^q(\ell^2)} \le c \Big(\sum_{n=0}^{\infty} \|v_n\|_{\ell^2}^2 \Big)^{1/2}.$$

Lemma 6.12 (see, e.g., [Pav14, Thm. 6.12]). There exists a numerical constant c > 0 such that for any complex sequence $(a_n)_{n=0}^{\infty}$,

$$\left\|\sum_{n=0}^{\infty}a_ne^{i2^n\theta}\right\|_{\mathrm{BMOA}}\leq c\left(\sum_{n=0}^{\infty}|a_n|^2\right)^{1/2}.$$

Proof of Proposition 6.10. Fix any $q \in [1, \infty)$. By assumption,

$$\left\| \left(\sum_{n=0}^{\infty} |\widehat{\varphi}(n)|^2 \widehat{f}(n) \widehat{f}(n)^* \right)^{1/2} \right\|_{S_p} \le c \left\| \sum_{n=0}^{\infty} |\widehat{\varphi}(n)|^2 e^{in\theta} \right\|_{\mathrm{BMOA}}^{1/2} \|f\|_{H^q(\ell^2)}.$$

In the above inequality, by taking $\varphi \in BMOA^{(2)}$ and $f \in H^q(\ell^2)$ with Fourier supports on the set of dyadic integers, we obtain

$$\left\| \left(\sum_{n=0}^{\infty} |d_n|^2 v_n v_n^* \right)^{1/2} \right\|_{S_p} \le c \left\| \sum_{n=0}^{\infty} |d_n|^2 e^{i2^n \theta} \right\|_{\text{BMOA}}^{1/2} \left\| \sum_{n=0}^{\infty} v_n e^{i2^n \theta} \right\|_{H^q(\ell^2)}$$

Then by Lemmas 6.11 and 6.12, there exists a numerical constant c' > 0 such that

$$\left\| \left(\sum_{n=0}^{\infty} |d_n|^2 v_n v_n^* \right)^{1/2} \right\|_{S_p} \le c' \left(\sum_{n=0}^{\infty} |d_n|^4 \right)^{1/4} \left(\sum_{n=0}^{\infty} \|v_n\|_{\ell^2}^2 \right)^{1/2}.$$

Hence, by setting $v_n = \lambda_n e_n$ with $\lambda_n \in \mathbb{C}$ and $e_n \in \ell^2$ the natural basis vector of ℓ^2 , we have

$$\left(\sum_{n=0}^{\infty} |d_n \lambda_n|^p\right)^{1/p} \le c' \left(\sum_{n=0}^{\infty} |d_n|^4\right)^{1/4} \left(\sum_{n=0}^{\infty} |\lambda_n|^2\right)^{1/2}.$$

Clearly, the above inequality holds for all $(d_n)_{n>0}$ and $(\lambda_n)_{n>0}$ only if $p \ge 4/3$.

6.5. Exact complete bounded norms of Carleman embeddings. Let $L^2_a(\mathbb{D})$ be the Bergman space on \mathbb{D} with respect to the normalized area measure $dA(z) = \frac{dxdy}{\pi}$. Recall the famous Carleman inequality: for any $f \in H^1$, we have $||f||_{L^2_a(\mathbb{D})} \leq ||f||_{H^1}$. Namely, the Carleman embedding

$$i: H^1 \to L^2_a(\mathbb{D})$$

is contractive (see [GK89] for an elementary proof).

The space $L^2_a(\mathbb{D})$ carries the *o.s.s.* inherited from that of $L^2(\mathbb{D})$. Since the natural *o.s.s.* on $L^2(\mathbb{D})$ is the Pisier's operator Hilbert space structure *OH*, by the uniqueness of *OH*, we have

(6.25)
$$\begin{array}{ccc} M: & L^2_a(\mathbb{D}) & \xrightarrow{completely \ isometric} & H^2 \\ & \sqrt{n+1}z^n & \xrightarrow{\simeq} & e^{in\theta} \end{array}$$

Corollary 6.13. $||i: H^1 \to L^2_a(\mathbb{D})||_{cb} = \sqrt{\pi}.$

Remark 6.3. Applying (6.1) to the Carleman embedding $i : H^1 \to L^2_a(\mathbb{D})$ and using the duality $(S_{\infty}[L^2_a(\mathbb{D})])^* = S_1[L^2_a(\mathbb{D})]$ together with the non-commutative Fubini theorem (6.9), we can reformulate Corollary 6.13 as follows: for any $f \in H^1(S_1)$ and $g \in \mathbb{C}[z] \otimes S_{\infty}$ (with $\sqrt{\pi}$ optimal):

(6.26)
$$\left|\int_{\mathbb{D}} \operatorname{Tr}(f(z)\overline{g(z)}) dA(z)\right| \leq \sqrt{\pi} \|f\|_{H^{1}(S_{1})} \left\|\int_{\mathbb{D}} \left[g(z) \otimes \overline{g(z)}\right] dA(z)\right\|_{B(\ell^{2} \otimes 2^{\ell^{2}})}^{1/2}$$

Proof of Corollary 6.13. By (6.25), $||i: H^1 \to L^2_a(\mathbb{D})||_{cb} = ||M \circ i: H^1 \to H^2||_{cb}$. Clearly, $M \circ i$ coincides with the Fourier multiplier corresponding to the sequence $(1/\sqrt{n+1})_{n=0}^{\infty}$. Hence, by the second isometric equality in Proposition 6.2, we have

(6.27)
$$\|i: H^1 \to L^2_a(\mathbb{D})\|_{cb} = \left\|\sum_{n=0}^{\infty} \frac{1}{\sqrt{n+1}} e^{in\theta}\right\|_{\mathrm{BMOA}^{(2)}} = \left\|\left[\frac{1}{i+j+1}\right]_{i,j\geq 0}\right\|_{B(\ell^2)}^{1/2} = \sqrt{\pi},$$

where the operator norm of the Hilbert matrix $[1/(i+j+1)]_{i,j\geq 0}$ is determined by Schur and can be found in, e.g., [HLP52, p.226] or [Mag50].

For any integer $d \ge 2$, write $L^2_a(\mathbb{D}^d) = L^2_a(\mathbb{D}^d, dA^{\otimes d})$ with $dA^{\otimes d}(z) = dA(z_1)\cdots dA(z_d)$. Let $i_d: H^1(\mathbb{T}^d) \to L^2_a(\mathbb{D}^d)$ denote the canonical embedding. The higher dimensional analogue of Corollary 6.13 is

Corollary 6.14. For any integer $d \ge 2$, we have $\|i_d : H^1(\mathbb{T}^d) \to L^2_a(\mathbb{D}^d)\|_{cb} = \pi^{d/2}$.

Proof. By the norm equality (6.27) of the Hilbert matrix and the Nehari Theorem, there exists a function $\psi \in L^{\infty}(\mathbb{T})$ with $\|\psi\|_{L^{\infty}(\mathbb{T})} = \pi$ such that $\widehat{\psi}(n) = \frac{1}{n+1}$ for all $n \ge 0$. Define $\psi_d(z) = \psi(z_1) \cdots \psi(z_d)$. Then $\|\psi_d\|_{L^{\infty}(\mathbb{T}^d)} = \pi^d$. Write $e_n(z_j) = z_j^n$ for $n \ge 1$ and $e_n(z_j) = \overline{z_j}^{|n|}$ for n < 0. For any $\alpha = (\alpha_1, \cdots, \alpha_d) \in \mathbb{Z}^d$, denote $|\alpha| = |\alpha_1| + \cdots + |\alpha_d|$ and $z^{\alpha} = e_{\alpha_1}(z_1) \cdots e_{\alpha_d}(z_d)$. Since $L^2_a(\mathbb{D}^d)$ carries Pisier's operator Hilbert space structure and the operator space dual of $H^1(\mathbb{T}^d)$ is given by $(H^1(\mathbb{T}^d))^* = L^{\infty}(\mathbb{T}^d)/H^1(\mathbb{T}^d)^{\circ}$ (where $H^1(\mathbb{T}^d)^{\circ}$ is the annihilator of $H^1(\mathbb{T}^d)$ in $L^{\infty}(\mathbb{T}^d)$), we have

$$\begin{split} \|i_{d}: H^{1}(\mathbb{T}^{d}) &\to L^{2}_{a}(\mathbb{D}^{d})\|_{cb} \\ = \sup_{0 < r < 1} \|\sum_{\alpha \in \mathbb{N}^{d}} \prod_{j=1}^{d} \frac{r^{|\alpha|}}{\alpha_{j} + 1} z^{\alpha} \otimes \overline{z^{\alpha}} \|_{(L^{\infty}(\mathbb{T}^{d})/H^{1}(\mathbb{T}^{d})^{\circ}) \otimes_{min}(\overline{L^{\infty}(\mathbb{T}^{d})/H^{1}(\mathbb{T}^{d})^{\circ})} \\ &\leq \sup_{0 < r < 1} \|\sum_{\alpha \in \mathbb{Z}^{d}} r^{|\alpha|} \widehat{\psi_{d}}(\alpha) z^{\alpha} \otimes \overline{z^{\alpha}} \|_{L^{\infty}(\mathbb{T}^{d}) \otimes_{min} \overline{L^{\infty}(\mathbb{T}^{d})}}^{1/2} \\ &= \sup_{0 < r < 1} \|\sum_{\alpha \in \mathbb{Z}^{d}} r^{|\alpha|} \widehat{\psi_{d}}(\alpha) z^{\alpha} \overline{w^{\alpha}} \|_{L^{\infty}(\mathbb{T}^{d} \times \mathbb{T}^{d})}^{1/2} \leq \|\psi_{d}\|_{L^{\infty}(\mathbb{T}^{d} \times \mathbb{T}^{d})}^{1/2} = \pi^{d/2}. \end{split}$$

Conversely, by Corollary 6.13, for any $\varepsilon > 0$, there exist functions $F \in S_1[H^1(\mathbb{T})] = H^1(\mathbb{T}; S_1)$ and $G \in S_{\infty}[L^2_a(\mathbb{D})]$ such that

$$||F||_{S_1[H^1(\mathbb{T})]} = 1, \quad ||\int_{\mathbb{D}} G(z_1) \otimes \overline{G(z_1)} dA(z_1)||_{B(\ell^2 \otimes_2 \ell^2)} = 1$$

and

$$\int_{\mathbb{D}} \operatorname{Tr}(F(z_1)\overline{G(z_1)}) dA(z_1) \geq \pi^{1/2} - \varepsilon.$$

Now we define an $S_1(\ell^2 \otimes_2 \cdots \otimes_2 \ell^2)$ -valued function $F_d(z) := F(z_1) \otimes \cdots \otimes F(z_d)$ and an $S_{\infty}(\ell^2 \otimes_2 \cdots \otimes_2 \ell^2)$ -valued function $G_d(z) := G(z_1) \otimes \cdots \otimes G(z_d)$. Then,

$$\|F_d\|_{H^1(\mathbb{T}^d;S_1(\ell^2 \otimes_2 \cdots \otimes_2 \ell^2))} = \int_{\mathbb{T}^d} \|F(z_1)\|_{S_1} \cdots \|F(z_d)\|_{S_1} dm(z_1) \cdots dm(z_d) = 1$$

and

$$\begin{split} &\|\int_{\mathbb{D}^d} G_d(z) \otimes \overline{G_d(z)} dA^{\otimes d}(z)\|_{B(\ell^2 \otimes_2 \cdots \otimes_2 \ell^2)} \\ &= \|\left[\int_{\mathbb{D}} G(z_1) \otimes \overline{G(z_1)} dA(z_1)\right] \otimes \cdots \otimes \left[\int_{\mathbb{D}} G(z_d) \otimes \overline{G(z_d)} dA(z_d)\right]\|_{B(\ell^2 \otimes_2 \cdots \otimes_2 \ell^2)} = 1. \end{split}$$

On the other hand,

$$\begin{aligned} \|F_d\|_{S_1[L^2_a(\mathbb{D}^d)} &\geq \int_{\mathbb{D}^d} \operatorname{Tr} \left(F_d(z) \overline{G_d(z)} \right) dA^{\otimes d}(z) \\ &= \left(\int_{\mathbb{D}} \operatorname{Tr} \left(F(z_1) \overline{G(z_1)} \right) dA(z_1) \right)^d \geq (\pi^{1/2} - \varepsilon)^d \end{aligned}$$

Since $\varepsilon > 0$ is arbitrary, this completes the proof of the inverse inequality.

6.6. Lifting property of Fourier multipliers in $(H^1, BMOA)_{cb}$.

Corollary 6.15. A Fourier multiplier T_m belongs to $(H^1, BMOA)_{cb}$ if and only if there exists a function $\psi \in L^{\infty}$ such that T_m satisfies the following commutative diagram :

(6.28)
$$\begin{array}{c} L^{1} \xrightarrow{K_{\Psi}} L^{\infty} \\ \mathcal{I} \uparrow inclusion \qquad \mathcal{Q} \downarrow quotient \\ H^{1} \xrightarrow{T_{m}} L^{\infty} / \overline{H_{0}^{\infty}} \simeq BMOA \end{array}$$

where $K_{\psi}(f) = f * \psi$. Moreover, $||T_m||_{cb} = \inf \{ ||\psi||_{\infty} | \psi \text{ satisfies (6.28)} \}.$

The above lifting property (6.28) of Fourier multipliers in $(H^1, BMOA)_{cb}$ implies

Corollary 6.16. $(H^1, BMOA)_{cb} = (H^1, H^{\infty})$ with equal norms.

Remark 6.4. Corollary 6.16 can also be proved by combining Proposition 6.2 and the equality $(H^1, H^\infty) = BMOA$. It is worthwhile to mention that a general multiplier in $(H^1, BMOA)$ need not to be liftable to a bounded one from H^1 to H^∞ , since

$$(H^1, H^\infty) \xrightarrow{isometric} BMOA \neq \mathcal{B} \xrightarrow{isomorphic} (H^1, BMOA)$$

where \mathcal{B} is the Bloch space on \mathbb{D} (see [HKZ00, Section 1.3]) and the last isomorphic equality is due to Mateljević and Pavlović ([MP90] and [Pav14, Thm. 11.10]).

Proof of Corollary 6.15. For any $T_m \in (H^1, BMOA)_{cb}$, by Proposition 6.2, we have

$$\|T_m\|_{(H^1, \text{BMOA})_{cb}} = \left\|\sum_{n=0}^{\infty} m_n e^{in\theta}\right\|_{\text{BMOA}}$$

Therefore, by recalling our choice (6.2) of the norm on BMOA, for any $\varepsilon > 0$, we can find $\psi \in L^{\infty}$ such that

$$\|\Psi\|_{L^{\infty}} \leq \|T_m\|_{(H^1, \operatorname{BMOA})_{cb}} + \varepsilon \text{ and } \mathcal{R}_+(\psi)(e^{i\theta}) = \sum_{n=0}^{\infty} m_n e^{in\theta},$$

where \mathcal{R}_+ is the Riesz projection. Define $K_{\psi}: L^1 \to L^{\infty}$ by $K_{\psi}(f) = f * \psi$ and the commutative diagram (6.28) follows immediately.

Conversely, assume that T_m satisfies (6.28). Since both the embedding map $H^1 \to L^1$ and the quotient map $L^{\infty} \to L^{\infty}/\overline{H_0^{\infty}}$ are completely contractive, by the property for minimal *o.s.s.* on L^{∞} (or the maximal *o.s.s.* on L^1) (see [Pis03, Chapter 3]), we have

$$||T_m||_{(H^1, \text{BMOA})_{cb}} \le ||K_{\psi}||_{CB(L^1, L^{\infty})} = ||K_{\psi}||_{B(L^1, L^{\infty})} = ||\psi||_{L^{\infty}}.$$

This completes the whole proof.

Proof of Corollary 6.16. Fix any Fourier multiplier T_m defined on H^1 . Since the subspace $H^{\infty} \subset L^{\infty}$ carries the mininal *o.s.s.* (see [Pis03, Chapter 3]), we have

$$|T_m||_{(H^1,H^\infty)} = ||T_m||_{(H^1,H^\infty)_{cb}}$$

The quotient map $Q: L^{\infty} \to BMOA = L^{\infty}/\overline{H_0^{\infty}}$ is completely contractive. Hence, by factorizing $T_m: H^1 \to BMOA$ as

$$H^1 \xrightarrow{T_m} H^\infty \xrightarrow{Q|_{H^\infty}} BMOA$$

we obtain

$$||T_m||_{(H^1, \text{BMOA})_{cb}} \le ||T_m||_{(H^1, H^\infty)_{cb}} \cdot ||Q|_{H^\infty} : H^\infty \to \text{BMOA}||_{cb} \le ||T_m||_{(H^1, H^\infty)}.$$

Conversely, assume that $||T_m||_{(H^1, BMOA)_{cb}} < 1$. By Corollary 6.15, there exists a function ψ on \mathbb{T} with $||\psi||_{L^{\infty}} < 1$ and

(6.29)
$$\mathcal{R}_{+}(\psi)(e^{i\theta}) = \sum_{n=0}^{\infty} m_{n}e^{in\theta},$$

such that $T_m: H^1 \to BMOA$ factorizes as

$$H^1 \xrightarrow{\mathcal{I}} L^1 \xrightarrow{K_{\psi}} L^\infty \xrightarrow{Q} BMOA.$$

Observe that $(K_{\psi} \circ \mathcal{I})(H^1) \subset H^{\infty}$. By (6.29), the multiplier $T_m : H^1 \to H^{\infty}$ coincides with the map $K_{\psi} \circ \mathcal{I} : H^1 \to H^{\infty}$. Consequently,

$$\|T_m\|_{(H^1,H^\infty)} \leq \|K_{\psi} \circ \mathcal{I} : H^1 \to H^\infty\| \leq \|\psi\|_{L^\infty} < 1.$$

This completes the whole proof.

6.7. Failure of hyper-complete-contractivity for the Poisson semigroup. Recall the Poisson convolution P_r on \mathbb{T} given by (6.11). The hypercontractivity result due to Janson (see [Jan83] or [Wei80]) says that $P_r : H^1 \to H^2$ is contractive if and only if $r \le \sqrt{1/2}$. However, Proposition 6.2 implies that there is no completely contractive version of hypercontractivity for Poisson semigroup.

Corollary 6.17. $||P_r||_{(H^1, H^2)_{cb}} = (1 - r^4)^{-1/2} > 1$ for any $r \in (0, 1)$.

Proof. Given any $r \in (0, 1)$, the Poisson convolution $P_r : H^1 \to H^2$ is just the Fourier multiplier associated with the sequence $(r^n)_{n=0}^{\infty}$. Therefore, by Proposition 6.2,

$$\|P_r\|_{(H^1, H^2)_{cb}} = \left\|\sum_{n=0}^{\infty} r^n e^{in\theta}\right\|_{\text{BMOA}^{(2)}} = \left\|[r^{2i+2j}]_{i,j\geq 0}\right\|_{B(\ell^2)}^{1/2}.$$

Since $[r^{2i+2j}]_{i,j\geq 0} = v_r v_r^*$ with $v_r = (1, r^2, r^4, \cdots, r^{2i}, \cdots)$ a row vector, we have

$$\left\| [r^{2i+2j}]_{i,j\geq 0} \right\|_{B(\ell^2)} = \left\| v_r \right\|_{\ell^2}^2 = \sum_{n=0}^{\infty} r^{4n} = \frac{1}{1-r^4}$$

This completes the whole proof.

REFERENCES

- [AMZ22] Loris Arnold, Christian Le Merdy, and Safoura Zadeh. S^1 -bounded fourier multipliers on $H^1(\mathbb{R})$ and functional calculus for semigroups. *arXiv: 2203.16829, to appear in J.Anal. Math.*
- [AMZ23] Loris Arnold, Christian Le Merdy, and Safoura Zadeh. Hankel operators on $L^p(\mathbb{R}_+)$ and their *p*-completely bounded multipliers. *arXiv: 2301.09481*.
- [BR11] Jean Berstel and Christophe Reutenauer. *Noncommutative rational series with applications*, volume 137 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2011.
- [BPc91] Oscar Blasco and Aleksander Pekzyński. Theorems of Hardy and Paley for vector-valued analytic functions and related classes of Banach spaces. *Trans. Amer. Math. Soc.*, 323(1):335–367, 1991.
- [BX91] Oscar Blasco and Quan Hua Xu. Interpolation between vector-valued Hardy spaces. J. Funct. Anal., 102(2):331–359, 1991.
- [BOCS21] Ole Fredrik Brevig, Joaquim Ortega-Cerdà, and Kristian Seip. Idempotent Fourier multipliers acting contractively on H^p spaces. Geom. Funct. Anal., 31(6):1377–1413, 2021.
- [BG09] Winfried Bruns and Joseph Gubeladze. *Polytopes, rings, and K-theory*. Springer Monographs in Mathematics. Springer, Dordrecht, 2009.
- [Bus71] George C. Bush. The embeddability of a semigroup—Conditions common to Malcev and Lambek. *Trans. Amer. Math. Soc.*, 157:437–448, 1971.
- [CAGPPT23] José M. Conde-Alonso, Adrián M. González-Pérez, Javier Parcet, and Eduardo Tablate. Schur multipliers in Schatten-von Neumann classes. Annals of Mathematics, 198(3):1229 – 1260, 2023.
- [Con94] Alain Connes. Noncommutative geometry. San Diego, CA, Academic Press Inc., 1994.

- [DR97] Gérard Duchamp and Christophe Reutenauer. Un critère de rationalité provenant de la géométrie non commutative. *Invent. Math.*, 128(3):613–622, 1997.
- [ER22] Edward G. Effros and Zhong-Jin Ruan. *Theory of operator spaces*. AMS Chelsea Publishing, Providence, RI, 2022.
- [Fli74] Michel Fliess. Matrices de Hankel. J. Math. Pures Appl. (9), 53:197–222, 1974.
- [GK89] Theodore W. Gamelin and Dmitry Khavinson. The isoperimetric inequality and rational approximation. *Amer. Math. Monthly*, 96(1):18–30, 1989.
- [GS03] Valerie Girardin and Rachid Senoussi. Semigroup stationary processes and spectral representation. *Bernoulli*, 9(5):857–876, 2003.
- [Gro53] Alexander Grothendieck. Résumé de la théorie métrique des produits tensoriels topologiques. *Bol. Soc. Mat. São Paulo*, 8:1–79, 1953.
- [HLP52] Godfrey H. Hardy, John E. Littlewood, and George Pólya. *Inequalities*. Cambridge, at the University Press, 1952. 2d ed.
- [HKZ00] Haakan Hedenmalm, Boris Korenblum, and Kehe Zhu. *Theory of Bergman spaces*, volume 199 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2000.
- [Hel10] Henry Helson. Hankel forms. *Studia Math.*, 198(1):79–84, 2010.
- [Jan83] Svante Janson. On hypercontractivity for multipliers on orthogonal polynomials. *Ark. Mat.*, 21(1):97–110, 1983.
- [JVA16] Miroljub Jevtić, Dragan Vukotić, and Miloš Arsenović. *Taylor coefficients and coefficient multipliers of Hardy and Bergman-type spaces*, volume 2 of *RSME Springer Series*. Springer, Cham, 2016.
- [Kat23] Elias Katsoulis. Fell's absorption principle for semigroup operator algebras. *arXiv: 2302.12809, to appear in J. Noncommutative Geometry.*
- [Lam51] Joachim Lambek. The immersibility of a semigroup into a group. Canad. J. Math., 3:34-43, 1951.
- [LM71] Robert J. Lindahl and Peter H. Maserick. Positive-definite functions on involution semigroups. Duke Math. J., 38:771–782, 1971.
- [Mag50] Wilhelm Magnus. On the spectrum of Hilbert's matrix. Amer. J. Math., 72:699-704, 1950.
- [Mal39] Anatoly I. Malcev. Über die Einbettung von assoziativen Systemen in Gruppen. *Rec. Math. [Mat. Sbornik] N.S.*, 6 (48):331–336, 1939.
- [Mal40] Anatoly I. Malcev. Über die Einbettung von assoziativen Systemen in Gruppen. II. *Rec. Math. [Mat. Sbornik] N.S.*, 8 (50):251–264, 1940.
- [MP90] Miodrag Mateljević and Miroslav Pavlović. Multipliers of *H^p* and BMOA. *Pacific J. Math.*, 146(1):71–84, 1990.
- [Miy22] Akihiro Miyagawa. A characterization of rationality in free semicircular operators. *Journal of Functional Analysis*, 283(8):109609, 2022.
- [Neh57] Zeev Nehari. On bounded bilinear forms. Ann. of Math. (2), 65:153–162, 1957.
- [OCS12] Joaquim Ortega-Cerdà and Kristian Seip. A lower bound in Nehari's theorem on the polydisc. J. Anal. Math., 118(1):339–342, 2012.
- [Pag70] Lavon B. Page. Bounded and compact vectorial Hankel operators. Trans. Amer. Math. Soc., 150:529– 539, 1970.
- [Pav14] Miroslav Pavlović. Function classes on the unit disc: An introduction, volume 52 of De Gruyter Studies in Mathematics. De Gruyter, Berlin, 2014.
- [Pel03] Vladimir V. Peller. *Hankel operators and their applications*. Springer Monographs in Mathematics. Springer-Verlag, New York, 2003.
- [Pis96] Gilles Pisier. The operator Hilbert space OH, complex interpolation and tensor norms. Mem. Amer. Math. Soc., 122(585):viii+103, 1996.
- [Pis98] Gilles Pisier. Non-commutative vector valued L_p -spaces and completely *p*-summing maps. *Astérisque*, (247):vi+131, 1998.
- [Pis01] Gilles Pisier. Similarity problems and completely bounded maps, volume 1618 of Lecture Notes in Mathematics. Springer-Verlag, Berlin, expanded edition, 2001. Includes the solution to "The Halmos problem".
- [Pis03] Gilles Pisier. Introduction to operator space theory, volume 294 of London Mathematical Society Lecture Note Series. Cambridge University Press, Cambridge, 2003.
- [Pis16] Gilles Pisier. Martingales in Banach spaces, volume 155 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2016.

- [Pis20] Gilles Pisier. Tensor products of C^{*}-algebras and operator spaces—the Connes-Kirchberg problem, volume 96 of London Mathematical Society Student Texts. Cambridge University Press, Cambridge, 2020.
- [Rua88] Zhong-Jin Ruan. Subspaces of C*-algebras. J. Funct. Anal., 76(1):217–230, 1988.
- [Sar67] Donald Sarason. Generalized interpolation in H^{∞} . Trans. Amer. Math. Soc., 127:179–203, 1967.
- [Sch77] Walter Schempp. On functions of positive type on commutative monoids. *Math. Z.*, 156(2):115–121, 1977.
- [SZ67] Elias M. Stein and Antoni Zygmund. Boundedness of translation invariant operators on Hölder spaces and *L^p*-spaces. *Ann. of Math.* (2), 85:337–349, 1967.
- [Wei80] Fred B. Weissler. Logarithmic Sobolev inequalities and hypercontractive estimates on the circle. J. *Funct. Anal.*, 37(2):218–234, 1980.
- [Xu90] Quan Hua Xu. Applications du théorème de factorisation pour des fonctions à valeurs opérateurs. *Studia Math.*, 95(3):273–292, 1990.

YONG HAN: SCHOOL OF MATHEMATICAL SCIENCES, SHENZHEN UNIVERSITY, SHENZHEN 518060, GUANGDONG, CHINA

Email address: hanyong@szu.edu.cn

YANQI QIU: SCHOOL OF FUNDAMENTAL PHYSICS AND MATHEMATICAL SCIENCES, HIAS, UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, HANGZHOU 310024, CHINA; INSTITUTE OF MATHEMATICS, AMSS, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, CHINA

Email address: yanqi.qiu@hotmail.com, yanqiqiu@ucas.ac.cn

ZIPENG WANG: COLLEGE OF MATHEMATICS AND STATISTICS, CHONGQING UNIVERSITY, CHONGQING 401331, CHINA

Email address: zipengwang2012@gmail.com