

ON MODULI SPACES OF CANONICAL THREEFOLDS WITH SMALL GENERA AND MINIMAL VOLUMES

MENG CHEN, YONG HU, AND CHEN JIANG

ABSTRACT. We prove that the canonical model of a 3-fold of general type with geometric genus 2 and with minimal canonical volume $\frac{1}{3}$ must be a hypersurface of degree 16 in $\mathbb{P}(1, 1, 2, 3, 8)$, which gives an explicit description of its canonical ring. This implies that the coarse moduli space $\mathcal{M}_{\frac{1}{3}, 2}$, parametrizing all canonical 3-folds with canonical volume $\frac{1}{3}$ and geometric genus 2, is an irreducible variety of dimension 189. Parallel studies show that $\mathcal{M}_{1, 3}$ is irreducible as well and is of dimension 236, and that $\mathcal{M}_{2, 4}$ is irreducible and is of dimension 270. As being conceived, every member in these 3 families is simply connected. Additionally, our method yields the expected Noether inequality $\text{Vol} > \frac{4}{3}p_g - \frac{10}{3}$ for 3-folds of general type with $p_g = 5$.

1. INTRODUCTION

Within birational geometry, perhaps the last phase of classifying projective varieties is to construct and to study the geometry of the concrete moduli space for each designated class. This is corroborated by the theory for curves and surfaces.

In higher dimensions, the birational classification of smooth projective varieties of general type splits into two steps: the first step is to find a good representative in each birational equivalence class, namely, the canonical model, and this step is completed by the minimal model program (see [21, 25, 1]); the second step is to study the geometry of canonical models and the moduli space of canonical models with given discrete numerical invariants (see [26, 33, 24]). While the general moduli theory is well-established, the explicit geometry of certain concrete moduli spaces is not yet well-understood.

This paper aims to investigate the birational geometry and moduli spaces of several special families, of 3-folds of general type, with small genera and attaining minimal canonical volumes.

We start by recalling the following theorem:

Theorem 1 ([7, Theorem 1.4, Theorem 1.5]). *Let W be a smooth projective 3-fold of general type. Denote by $\text{Vol}(W)$ the canonical volume of W . Then*

Date: July 2, 2024.

2020 Mathematics Subject Classification. 14J30, 14J10, 14D22.

Key words and phrases. canonical threefold, moduli space, canonical volume, weighted hypersurface.

- (1) when $p_g(W) \geq 2$, $\text{Vol}(W) \geq \frac{1}{3}$;
- (2) when $p_g(W) \geq 3$, $\text{Vol}(W) \geq 1$;
- (3) when $p_g(W) \geq 4$, $\text{Vol}(W) \geq 2$.

Furthermore, the above three lower bounds $\frac{1}{3}$, 1 and 2 are all optimal and the minimal volume is attained only if $p_g = 2, 3, 4$, respectively.

It is a natural question to study the classification problem for those 3-folds attaining the above minimal volumes, as analogue to curves with small genus or surfaces with small c_1^2 . Also, it is expected that the classification of such 3-folds with extreme numerical values will benefit the development of explicit classification theory of higher dimensional algebraic varieties.

To study the birational geometry of a variety W of general type, one classical method is to study its *canonical ring* $R(W) := \bigoplus_{k \geq 0} H^0(W, kK_W)$ which originates from the work of Zariski and Mumford ([34]). Reid proposed a program to study the canonical ring from the point of view of commutative algebra which predicts the construction of W (as a subvariety in a weighted projective space) from the algebraic structure of $R(W)$, see [29, 31]. However, to the best of our knowledge, the explicit algebraic structure of the canonical ring of a variety is only known in the case that the given variety has a very clear geometric structure (e.g. a complete intersection in a weighted projective space). In this paper, we will give explicit descriptions for the canonical rings of smooth projective 3-folds of general type with $2 \leq p_g(X) \leq 4$ attaining minimal volumes.

For a positive rational number a and a nonnegative integer b , denote by $\mathcal{M}_{a,b}$ the coarse moduli space parametrizing all projective 3-folds X with canonical singularities such that K_X is ample, $K_X^3 = a$, and $p_g(X) = b$ in the sense of Viehweg [33, §8.5] and Kollár–Shepherd-Barron [26, §5.4] (see also [24]). When $\mathcal{M}_{a,b}$ is nonempty, one would like to know its geometric properties as detailed as possible, for example, the number of its irreducible or connected components and the dimension of each component. A further task is to construct an ideal compactification for $\mathcal{M}_{a,b}$, and one successful way is the KSB moduli theory via the minimal model program (see [26, 24]). To the best of our knowledge, for the explicit geometry of moduli spaces of 3-folds of general type, very few is known in the literature. In this paper, we mainly study the irreducibility and dimensions of $\mathcal{M}_{\frac{1}{3},2}$, $\mathcal{M}_{1,3}$ and $\mathcal{M}_{2,4}$. It is not out of expectation that every member in these families will be proven to be simply connected.

Our main results are as follows.

Theorem 1.1. *Let W be a smooth projective 3-fold of general type with $p_g(W) \geq 2$ and $\text{Vol}(W) = \frac{1}{3}$. Then*

- (1) W has the same plurigenera as those of a general hypersurface of degree 16 in $\mathbb{P}(1, 1, 2, 3, 8)$;

- (2) the canonical model of W is a hypersurface in $\mathbb{P}(1, 1, 2, 3, 8)$ defined by a weighted homogeneous polynomial f of degree 16, where

$$f(x_0, \dots, x_4) = x_4^2 + f_0(x_0, x_1, x_2, x_3)$$

in suitable homogeneous coordinates $[x_0 : \dots : x_4]$ of $\mathbb{P}(1, 1, 2, 3, 8)$; in other words, the canonical ring of W is

$$R(W) \simeq \mathbb{C}[x_0, \dots, x_4]/(f),$$

where $\mathbb{C}[x_0, \dots, x_4]$ is viewed as a weighted polynomial ring with $\text{wt}(x_0, \dots, x_4) = (1, 1, 2, 3, 8)$;

- (3) W is simply connected;
 (4) the moduli space $\mathcal{M}_{\frac{1}{3}, 2}$ is irreducible and $\dim \mathcal{M}_{\frac{1}{3}, 2} = 189$.

Theorem 1.2. *Let W be a smooth projective 3-fold of general type with $p_g(W) \geq 3$ and $\text{Vol}(W) = 1$. Then*

- (1) W has the same plurigenera as those of a general hypersurface of degree 12 in $\mathbb{P}(1, 1, 1, 2, 6)$;
 (2) the canonical model of W is a hypersurface in $\mathbb{P}(1, 1, 1, 2, 6)$ defined by a weighted homogeneous polynomial f of degree 12, where

$$f(x_0, \dots, x_4) = x_4^2 + f_0(x_0, x_1, x_2, x_3)$$

in suitable homogeneous coordinates $[x_0 : \dots : x_4]$ of $\mathbb{P}(1, 1, 1, 2, 6)$; in other words, the canonical ring of W is

$$R(W) \simeq \mathbb{C}[x_0, \dots, x_4]/(f),$$

where $\mathbb{C}[x_0, \dots, x_4]$ is viewed as a weighted polynomial ring with $\text{wt}(x_0, \dots, x_4) = (1, 1, 1, 2, 6)$;

- (3) W is simply connected;
 (4) the moduli space $\mathcal{M}_{1, 3}$ is irreducible and $\dim \mathcal{M}_{1, 3} = 236$.

Theorem 1.3. *Let W be a smooth projective 3-fold of general type with $p_g(W) \geq 4$ and $\text{Vol}(W) = 2$. Then*

- (1) W has the same plurigenera as those of a general hypersurface of degree 10 in $\mathbb{P}(1, 1, 1, 1, 5)$;
 (2) the canonical model of W belongs to either of the following 2 types:
 (a) **Type I:** a hypersurface in $\mathbb{P}(1, 1, 1, 1, 5)$ defined by a weighted homogeneous polynomial f of degree 10, where

$$f(x_0, \dots, x_4) = x_4^2 + f_0(x_0, x_1, x_2, x_3)$$

in suitable homogeneous coordinates $[x_0 : \dots : x_4]$ of $\mathbb{P}(1, 1, 1, 1, 5)$; in other words, the canonical ring of W is

$$R(W) \simeq \mathbb{C}[x_0, \dots, x_4]/(f),$$

where $\mathbb{C}[x_0, \dots, x_4]$ is viewed as a weighted polynomial ring with $\text{wt}(x_0, \dots, x_4) = (1, 1, 1, 1, 5)$;

(b) **Type II:** a subvariety in $\mathbb{P}(1, 1, 1, 1, 2, 5)$ defined by 2 weighted homogeneous polynomials q and f of degrees 2 and 10 respectively, where

$$q(x_0, x_1, x_2, x_3) = x_3^2 + q_0(x_0, x_1, x_2),$$

$$f(x_0, \dots, x_5) = x_5^2 + f_0(x_0, \dots, x_4)$$

in suitable homogeneous coordinates $[x_0 : \dots : x_5]$ of $\mathbb{P}(1, 1, 1, 1, 2, 5)$; in other words, the canonical ring of W is

$$R(W) \simeq \mathbb{C}[x_0, \dots, x_5]/(q, f),$$

where $\mathbb{C}[x_0, \dots, x_5]$ is viewed as a weighted polynomial ring with $\text{wt}(x_0, \dots, x_5) = (1, 1, 1, 1, 2, 5)$;

furthermore, a canonical 3-fold of type II is a specialization of canonical 3-folds of type I;

(3) W is simply connected;

(4) the moduli space $\mathcal{M}_{2,4}$ is irreducible and $\dim \mathcal{M}_{2,4} = 270$.

As a by-product of this paper, we obtain the following result which solves one remaining case ($p_g = 5$) of the Noether inequality in the previous work of Chen–Chen–Jiang [4, 5].

Theorem 1.4 (=Theorem 9.5). *Let X be a minimal 3-fold of general type with $p_g(X) = 5$. Then $K_X^3 \geq \frac{52}{15}$. In particular, the inequality $K_X^3 > \frac{4}{3}p_g(X) - \frac{10}{3}$ holds in the case of $p_g(X) = 5$.*

We briefly explain the main methods and difficulties of this paper. Consider Theorem 1.1 for instance, which is the most difficult one. The key part is to show that the canonical model can be embedded into $\mathbb{P}(1, 1, 2, 3, 8)$. As in [19, 20], we have an effective criterion on how to embed a given polarized variety into a certain weighted projective space. In the context of Theorem 1.1, to apply this criterion, we need to know all the plurigenera of W and the behavior of pluricanonical systems of W . Usually plurigenera can be computed by Reid’s Riemann–Roch formula on a minimal model X of W , however, in order to apply this formula, only knowing the geometric genus and the canonical volume is not sufficient, one has to know the singularities of X , as well as the holomorphic Euler characteristic of X . We put our main effort to get such geometric information.

In the study of explicit geometry of 3-folds of general type, one of the most challenging cases is when X admits a pencil $|F_X|$ which is not free. One key observation (see §3) is, by applying MMP to a log resolution of the pair (X, F_X) in question, to obtain a relatively minimal model (X', F) over X on which F is free. Such a birational model can be viewed as a minimal partial resolution for the pencil $|F_X|$. With the help of this new birational model, we can efficiently analyze the geometry of X , and get the information on the singularities of X . Then we analyze the behavior of pluricanonical systems of X which gives restrictions on the plurigenera of X . Together

with Reid's Riemann–Roch formula, such restrictions will eventually lead us to the precise value of the holomorphic Euler characteristic of X and all plurigenera of X .

This paper is organized as the following. In §2, we recall some basic knowledge and background. In §3, we construct a relative minimal model resolving a pencil by the minimal model program. In §4–§6, we study the geometry, of minimal 3-folds of general type with $2 \leq p_g \leq 4$ and with minimal canonical volume, such as their singularities, pluricanonical systems and plurigenera. In §7, we recall the criterion in [20] in a slightly more general form. In §8, we prove our main results upon the preparation of previous sections. In §9, we prove the Noether inequality for $p_g = 5$ as an application of the birational model in §3.

2. PRELIMINARIES

Throughout this paper, we work over the complex number field \mathbb{C} meanwhile all results can be easily generalized to any algebraically closed field of characteristic 0.

We adopt the standard notation and definitions in [21, 25], and will freely use them.

A \mathbb{Q} -divisor is said to be \mathbb{Q} -effective if it is \mathbb{Q} -linearly equivalent to an effective \mathbb{Q} -divisor.

2.1. Singularities.

A *log pair* (X, B) consists of a normal projective variety X and an effective \mathbb{Q} -divisor B on X such that $K_X + B$ is \mathbb{Q} -Cartier.

Let $f : Y \rightarrow X$ be a log resolution of the log pair (X, B) , write

$$K_Y = f^*(K_X + B) + \sum a_i D_i,$$

where $\{D_i\}$ are distinct prime divisors and the sum runs through all irreducible components in the union of f -exceptional divisors and the strict transform of B . The log pair (X, B) is called

- (a) *kawamata log terminal* (*klt*, for short) if $a_i > -1$ for all i ;
- (b) *purely log terminal* (*plt*, for short) if $a_i > -1$ for all f -exceptional divisors F_i and all f ;
- (c) *terminal* if $a_i > 0$ for all f -exceptional divisors D_i and all f ;
- (d) *canonical* if $a_i \geq 0$ for all f -exceptional divisors D_i and all f .

Usually we write X instead of $(X, 0)$ in the case $B = 0$.

2.2. Minimal model and canonical model.

A normal projective variety X is said to be *minimal* if X has \mathbb{Q} -factorial terminal singularities and K_X is nef.

A normal projective variety X with canonical singularities is called *of general type* if K_X is big.

According to the minimal model program (see [1]), a smooth projective variety W of general type is always birational to a *minimal model* W_{\min}

which is a minimal variety of general type, and to a unique *canonical model* W_{can} which is a normal projective variety with canonical singularities and $K_{W_{\text{can}}}$ is ample. In this case, the *canonical volume* of W is defined to be the self-intersection number of the canonical divisor of a minimal model or the canonical model, i.e., $\text{Vol}(W) := K_{W_{\text{min}}}^{\dim W} = K_{W_{\text{can}}}^{\dim W}$. The *geometric genus* of W is defined by $p_g(W) := h^0(W, K_W) = h^0(W_{\text{min}}, K_{W_{\text{min}}})$.

In the study of birational geometry of a smooth projective variety of general type, we often freely replace it by its minimal model or canonical model.

For two integers $m > 0$ and $n \geq 0$, an (m, n) -*surface* is a smooth projective surface S of general type with $\text{Vol}(S) = m$ and $p_g(S) = n$.

2.3. Rational maps defined by linear systems.

Let X be a normal projective variety on which D is a \mathbb{Q} -Cartier Weil divisor with $h^0(X, D) \geq 2$. Take a linear subspace $V \subseteq H^0(X, D)$ with $\dim V \geq 2$ and consider the corresponding sublinear system $\Lambda \subseteq |D|$. We can consider the rational map defined by Λ , say $\Phi_\Lambda : X \dashrightarrow \mathbb{P}^{\dim V - 1}$, which is not necessarily well-defined everywhere. By Hironaka's big theorem, we can take successive blow-ups $\pi : W \rightarrow X$ such that:

- (i) W is smooth and projective;
- (ii) corresponding to Λ , the sublinear system $\Lambda_W \subseteq |\pi^*(D)|$ has free movable part $\text{Mov}(\Lambda_W)$ and, consequently, the rational map $\gamma = \Phi_\Lambda \circ \pi$ is a morphism.

Let $W \xrightarrow{\psi} \Sigma' \xrightarrow{s} \Sigma$ be the Stein factorization of γ with $\Sigma = \gamma(W) \subseteq \mathbb{P}^{\dim V - 1}$. There is a commutative diagram:

$$\begin{array}{ccc}
 W & \xrightarrow{\psi} & \Sigma' \\
 \pi \downarrow & \searrow \gamma & \downarrow s \\
 X & \xrightarrow{\Phi_\Lambda} & \Sigma
 \end{array}$$

- (1) If $\dim(\Sigma') \geq 2$, then a general member of $\text{Mov}(\Lambda_W)$ is a smooth projective variety by Bertini's theorem, and is called a *general irreducible element* of $\text{Mov}(\Lambda_W)$ (or Λ_W).
- (2) If $\dim(\Sigma') = 1$, then a general fiber S of ψ is a smooth projective variety by Bertini's theorem. In this case, we say that Λ is *composed with a pencil* and S is called a *general irreducible element* of $\text{Mov}(\Lambda_W)$ (or Λ_W). Λ is said to be *rational* if $\Sigma' \simeq \mathbb{P}^1$. A general member of $\text{Mov}(\Lambda_W)$ is of the form $\sum_{i=1}^a S_i$, where S_i is a smooth fiber of ψ for each i . It is clear that $a = \dim V - 1$ if $\Sigma' \simeq \mathbb{P}^1$, and $a \geq \dim V$ if $\Sigma' \not\simeq \mathbb{P}^1$.

We will frequently use the following setting.

Setting 2.1. Let X be a minimal 3-fold of general type with $p_g(X) \geq 2$. Take a birational modification $\pi : W \rightarrow X$ as in §2.3 with respect to the canonical system $|K_X|$ and keep the same notation. Write

$$\begin{aligned} K_W &= \pi^*(K_X) + E_\pi, \\ \pi^*(K_X) &= M + Z_\pi, \end{aligned}$$

where $|M| = \text{Mov } \lfloor \pi^*(K_X) \rfloor$, E_π and Z_π are effective \mathbb{Q} -divisors. Let S be a general irreducible element of $|M|$. After taking some further blow-up, we may further assume that the following hold:

- (1) $\text{Mov } |K_S|$ is base point free;
- (2) for any integers $0 < m \leq 4$ and $0 \leq j \leq m$, the linear system

$$|M_{m,-j}| := \text{Mov } \lfloor \pi^*(mK_X) - jS \rfloor$$

is base point free or empty;

- (3) $\text{Supp}(Z_\pi + E_\pi)$ is of simple normal crossing.

Consider the natural restriction map

$$\theta_{m,-j} : H^0(W, M_{m,-j}) \rightarrow H^0(S, M_{m,-j}|_S),$$

and set $u_{m,-j} = \dim \text{Im } \theta_{m,-j}$.

For two linear systems Λ_1 and Λ_2 , we write $\Lambda_1 \succcurlyeq \Lambda_2$ if there exists an effective divisor N such that $\Lambda_1 \supseteq \Lambda_2 + N$.

2.4. Reid's Riemann–Roch formula.

A *basket* is a collection of pairs of integers (permitting weights).

Let X be a projective terminal 3-fold. According to Reid [30], there is a basket of orbifold points (called the *Reid basket*)

$$B_X = \left\{ (b_i, r_i) \mid i = 1, \dots, s; 0 < b_i \leq \frac{r_i}{2}; b_i \text{ is coprime to } r_i \right\}$$

associated to X , which comes from locally deforming the singularities of X into cyclic quotient singularities, where a pair (b_i, r_i) corresponds to a (virtual) orbifold point Q_i of type $\frac{1}{r_i}(1, -1, b_i)$.

By Reid's Riemann–Roch formula [30, Corollary 10.3], for any positive integer m ,

$$\begin{aligned} &\chi(X, \mathcal{O}_X(mK_X)) \\ &= \frac{1}{12}m(m-1)(2m-1)K_X^3 - (2m-1)\chi(\mathcal{O}_X) + l(m), \end{aligned} \quad (2.1)$$

where $l(m) = \sum_i \sum_{j=1}^{m-1} \frac{\overline{jb_i(r_i - \overline{jb_i})}}{2r_i}$ and the first sum runs over all orbifold points in the Reid basket. Here $\overline{jb_i}$ means the smallest nonnegative residue of $jb_i \bmod r_i$.

For a positive integer m , the m -th plurigenus of X is denoted by $P_m(X) := h^0(X, mK_X)$, and it can be computed by Reid's Riemann–Roch formula and the Kawamata–Viehweg vanishing theorem when X is minimal of general type and $m \geq 2$.

2.5. Hodge index theorem.

We recall the following well-known result which is a consequence of the Hodge index theorem.

Lemma 2.2. *Let D_1, D_2 be \mathbb{Q} -Cartier \mathbb{Q} -divisors on a normal projective surface S .*

- (1) *If D_1, D_2 are nef, $D_1^2 = D_2^2 > 0$, and $D_1 - D_2$ is effective, then $D_1 = D_2$.*
- (2) *If D_1, D_2 are effective, $(D_1 \cdot D_2) = 0$, and $D_1 + D_2$ is nef and big, then either $D_1 = 0$ or $D_2 = 0$.*

Proof. Recall that the Hodge index theorem states that if H_1 and H_2 are \mathbb{Q} -Cartier \mathbb{Q} -divisors on S such that $H_1^2 > 0$ and $(H_1 \cdot H_2) = 0$, then $H_2^2 \leq 0$; moreover, the equality holds if and only if $H_2 \equiv 0$. Here recall that if H_2 is effective, then $H_2 \equiv 0$ if and only if $H_2 = 0$.

(1) Denote $D = D_1 - D_2 \geq 0$. Since $0 = D_1^2 - D_2^2 = ((D_1 + D_2) \cdot D)$ and D_1, D_2 are nef, we have $(D_1 \cdot D) = (D_2 \cdot D) = 0$. In particular, $D^2 = 0$. On the other hand, $(D_1 + D_2)^2 > 0$. So by the Hodge index theorem, $D = 0$.

(2) Note that $D_1^2 + D_2^2 = (D_1 + D_2)^2 > 0$. So either $D_1^2 > 0$ or $D_2^2 > 0$. Without loss of generality, we may assume that $D_1^2 > 0$. Then $(D_1 \cdot D_2) = 0$ implies that $D_2^2 \leq 0$ by the Hodge index theorem. On the other hand, $D_2^2 = (D_2 \cdot (D_1 + D_2)) \geq 0$ as $D_1 + D_2$ is nef. Hence $D_2^2 = 0$ and $D_2 = 0$ by the Hodge index theorem. \square

3. A RELATIVELY MINIMAL MODEL RESOLVING A PENCIL

In this section, we construct a birational model which resolves a given pencil by using the minimal model program.

Proposition 3.1. *Let X be a normal \mathbb{Q} -factorial projective variety on which a linear system Λ is composed with a pencil. Then there exists a projective variety X' with a birational morphism $g : X' \rightarrow X$ having the following properties:*

- (1) *X' is \mathbb{Q} -factorial terminal;*
- (2) *the movable part of the linear system Λ' , on X' , corresponding to Λ is base point free and, after taking the Stein factorization, Λ' induces a fibration $f : X' \rightarrow \Gamma$ onto a curve Γ ;*
- (3) *for a general fiber F of f , $K_{X'} + F$ is g -nef;*
- (4) *$g^*(K_X + g_*F) - K_{X'} - F$ is an effective g -exceptional \mathbb{Q} -divisor.*

Proof. Take a birational modification $\pi : W \rightarrow X$ and a fibration $\psi : W \rightarrow \Gamma$ as in §2.3 induced by Λ . Take S_1, S_2 to be two general fibers of ψ , then S_1 and S_2 are disjoint and smooth. The pair $(W, \frac{1}{2}S_1 + \frac{1}{2}S_2)$ is klt and terminal. By [1, Corollary 1.4.2], we may run a $(K_W + \frac{1}{2}S_1 + \frac{1}{2}S_2)$ -MMP over X which terminates at a model $g : X' \rightarrow X$ such that $K_{X'} + \frac{1}{2}F_1 + \frac{1}{2}F_2$ is nef over X , where F_1 and F_2 are strict transforms of S_1 and S_2 on X' . We claim that X' has all required properties.

As S_1 and S_2 are movable, they are not contracted by the MMP, so $(X', \frac{1}{2}F_1 + \frac{1}{2}F_2)$ is terminal by [25, Corollary 3.42, Corollary 3.43]. In particular, X' is \mathbb{Q} -factorial terminal.

We claim that $F_1 \cap F_2 = \emptyset$. Suppose that $F_1 \cap F_2 \neq \emptyset$, then there exists a subvariety $Q \subseteq F_1 \cap F_2$ of codimension 2 in X' as X' is \mathbb{Q} -factorial. On the other hand, X' is smooth at the generic point of Q by [25, Corollary 5.18]. So by [25, Lemma 2.29], the blowing up along Q induces an exceptional prime divisor E with discrepancy $a(E, X', \frac{1}{2}F_1 + \frac{1}{2}F_2) \leq 1 - \frac{1}{2} - \frac{1}{2} = 0$, but this contradicts the fact that $(X', \frac{1}{2}F_1 + \frac{1}{2}F_2)$ is terminal. So $F_1 \cap F_2 = \emptyset$.

If $\Gamma \simeq \mathbb{P}^1$, then the free linear system $|F_1|$ induces a natural morphism $f : X' \rightarrow \Gamma$. On the other hand, if $g(\Gamma) > 0$, then all steps of the MMP are relative over Γ since negative extremal rays are generated by rational curves, and hence $W \rightarrow \Gamma$ descends to a natural morphism $f : X' \rightarrow \Gamma$. Clearly in either case, this morphism is defined by the linear system Λ' , on X' , corresponding to Λ .

Finally consider

$$G = g^*(K_X + g_*F) - K_{X'} - F.$$

Then G is exceptional over X and

$$-G \equiv_X K_{X'} + F \equiv K_{X'} + \frac{1}{2}F_1 + \frac{1}{2}F_2$$

is nef over X by our construction. Hence $G \geq 0$ by the negativity lemma [25, Lemma 3.39]. \square

As a consequence of Proposition 3.1, we will frequently use the following modification for a minimal 3-fold of general type with nontrivial canonical map.

Setting 3.2. Let X be a minimal 3-fold of general type with $p_g(X) \geq 2$. Take a general sublinear system $\Lambda \subseteq |K_X|$ which is composed with a pencil. Applying Proposition 3.1 to Λ , we get a projective birational morphism $\mu : X' \rightarrow X$ with a surjective fibration $f : X' \rightarrow \Gamma$ such that X' is \mathbb{Q} -factorial terminal and

$$\mu^*(K_X + F_X) - K_{X'} - F = G \tag{3.1}$$

is an effective μ -exceptional \mathbb{Q} -divisor, where F is a general fiber of f and $F_X = \mu_*F$. Here G is independent of F by the negativity lemma [25, Lemma 3.39]. We may write

$$\begin{aligned} K_X &= F_X + Z_X, \\ \mu^*(K_X) &= F + Z, \\ K_{X'} &= \mu^*(K_X) + E_\mu, \end{aligned}$$

where Z_X , E_μ and Z are effective \mathbb{Q} -divisors and E_μ is μ -exceptional. Note that F is a smooth projective surface of general type. Denote by $\sigma : F \rightarrow F_0$ the contraction to its minimal model.

Lemma 3.3. *Keep the notation in Setting 3.2. Suppose that $|K_X|$ is composed with a rational pencil, then*

$$p_g(X)\mu^*(K_X)|_F - (p_g(X) - 1)\sigma^*(K_{F_0})$$

is effective. More precisely, there exists an effective divisor $D \in |K_X|$ and an effective divisor $D_0 \in |K_{F_0}|$ such that

$$p_g(X)\mu^*(D)|_F - (p_g(X) - 1)\sigma^*(D_0)$$

is an effective \mathbb{Q} -divisor.

Proof. To simplify the notation we denote $d = p_g(X) - 1$. Clearly Λ and $|K_X|$ are composed with the same pencil. So we may write $K_X = dF_X + \Delta$, where Δ is the fixed part of $|K_X|$. Choose a general fiber F' of f different from F , denote $F'_X = \mu_*F'$. Then we can take $D = dF'_X + \Delta$.

We first explain the construction of D_0 . As $D \in |K_X|$, we have $\mu^*(D) + E_\mu \in |K_{X'}|$. Then $\mu^*(D)|_F + E_\mu|_F \in |K_F|$, where the restriction is well-defined as $F \not\subset \text{Supp}(\mu^*(D) + E_\mu)$. Then we just take $D_0 := \sigma_*(\mu^*(D)|_F + E_\mu|_F) \in |K_{F_0}|$. Clearly $\mu^*(D)|_F + E_\mu|_F - \sigma^*(D_0) = K_F - \sigma^*(K_{F_0})$ is effective.

Applying (3.1) to $D \in |K_X|$ and F' , one has

$$(d+1)\mu^*(D) \geq d\mu^*(D + F'_X) = d(\mu^*(D) + E_\mu + F' + G) \geq d(\mu^*(D) + E_\mu).$$

Then we get the conclusion by restricting on F . \square

Lemma 3.4. *Keep the notation in Setting 3.2. Suppose that $(F, G|_F)$ is klt.*

(1) *Then (X, F_X) is plt, F_X is normal and klt, and*

$$K_F + G|_F = (\mu|_F)^*(K_{F_X}).$$

(2) *Suppose that $Z_X = 0$. For a non-Gorenstein singularity P of X , denote by r_P the Cartier index of K_X at P . Then $P \in F_X$ and r_P equals to the Cartier index of $K_X|_{F_X}$ (as a well-defined Weil divisor on F_X) at P . Furthermore, if P is a Gorenstein point on F_X , then $r_P = 2$.*

Proof. (1) Recall that by (3.1),

$$K_{X'} + F + G = \mu^*(K_X + F_X).$$

Since $(F, G|_F)$ is klt, by [25, Theorem 5.50], $(X', F + G)$ is plt in a neighborhood of F . Then by [25, Theorem 5.48], (X, F_X) is plt in a neighborhood of F_X . This implies that (X, F_X) is plt as X is terminal. In particular, F_X is normal by [25, Proposition 5.51]. The remaining assertions follow from the adjunction formula $(K_X + F_X)|_{F_X} = K_{F_X}$ and [25, Theorem 5.50] as X has only isolated singularities.

(2) Since $K_X \sim F_X$, all non-Gorenstein singularities of X are contained in F_X . Recall that $K_X|_{F_X}$ is a well-defined Weil divisor on F_X since X and F_X both have only isolated singularities and F_X is general. Then r_P equals to the Cartier index of $K_X|_{F_X}$ at P by [14, Theorem A.1]. For the last assertion, note that $2K_X|_{F_X} \sim (K_X + F_X)|_{F_X} = K_{F_X}$ by the adjunction formula. So if K_{F_X} is Cartier at P , then $r_P = 2$. \square

Lemma 3.5. *Keep the notation in Setting 3.2. Then $Z_X = 0$ if and only if $\mu^*(Z_X)|_F \equiv 0$.*

Proof. We only need to show the “if” part. Take a general very ample divisor H on X , by the projection formula,

$$(F_X|_H \cdot Z_X|_H) = (H \cdot Z_X \cdot F_X) = (\mu^*(H) \cdot \mu^*(Z_X) \cdot F) = 0.$$

On the other hand, $K_X|_H = F_X|_H + Z_X|_H$ is a nef and big divisor on H . Clearly $F_X|_H \neq 0$, so $Z_X|_H = 0$ by Lemma 2.2(2), which implies that $Z_X = 0$. \square

4. BIRATIONAL GEOMETRY OF 3-FOLDS WITH $p_g = 2$ AND $\text{Vol} = 1/3$

In this section, we study the birational geometry of a minimal 3-fold X of general type with $p_g(X) = 2$ and $K_X^3 = \frac{1}{3}$.

4.1. Geometric structure and non-Gorenstein singularities.

Since $p_g(X) = 2$, $|K_X|$ is composed with a pencil. Take a birational modification $\mu : X' \rightarrow X$ as in Setting 3.2 and keep the same notation. Since $h^1(\mathcal{O}_X) = 0$ by [7, Theorem 1.5(3)], we have $\Gamma \simeq \mathbb{P}^1$, in other words, $|K_X|$ is composed with a rational pencil. By [7, Theorem 1.4(2)], F is a $(1, 2)$ -surface.

We recall the following lemma from [7, §2.15].

Lemma 4.1. *Let X be a minimal 3-fold of general type with $p_g(X) = 2$ and $K_X^3 = \frac{1}{3}$. Keep the notation in Setting 3.2. Then*

$$(\mu^*(K_X)|_F)^2 = \frac{1}{3}, \quad (\mu^*(K_X)|_F \cdot \sigma^*(K_{F_0})) = (\mu^*(K_X)|_F \cdot \text{Mov} |\sigma^*(K_{F_0})|) = \frac{2}{3}.$$

Proof. For the convenience of readers who are not familiar with [7], we briefly recall the proof. Denote by $\xi = (\mu^*(K_X)|_F \cdot \text{Mov} |\sigma^*(K_{F_0})|)$ (which coincides with the definition in [7, §2.9]). Then by the first 3 lines of [7, §2.15], one has $\xi \geq \frac{2}{3}$. By Lemma 3.3, $2\mu^*(K_X)|_F - \sigma^*(K_{F_0})$ is effective. Then

$$\begin{aligned} K_X^3 &= \mu^*(K_X)^3 \geq (\mu^*(K_X)^2 \cdot F) = (\mu^*(K_X)|_F)^2 \\ &\geq \frac{1}{2}(\mu^*(K_X)|_F \cdot \sigma^*(K_{F_0})) \geq \frac{\xi}{2} \geq \frac{1}{3}. \end{aligned}$$

Then by assumption, all the above inequalities are equalities. \square

Now we will analyze the geometry of X in more details.

Lemma 4.2. *Let X be a minimal 3-fold of general type with $p_g(X) = 2$ and $K_X^3 = \frac{1}{3}$. Keep the notation in Setting 3.2. Then there exists a unique μ -exceptional prime divisor E_0 such that $(\sigma^*(K_{F_0}) \cdot E_0|_F) > 0$. Moreover,*

- (1) $(\sigma^*(K_{F_0}) \cdot E_0|_F) = 1$;
- (2) $\text{coeff}_{E_0} E_\mu = \frac{1}{3}$, $\text{coeff}_{E_0} Z = \frac{2}{3}$;
- (3) $\mu(E_0)$ is a point.

Proof. By the adjunction formula, we have

$$K_F = \mu^*(K_X)|_F + E_\mu|_F.$$

By Lemma 4.1 and the equality $(K_F \cdot \sigma^*(K_{F_0})) = K_{F_0}^2 = 1$, we have

$$(E_\mu|_F \cdot \sigma^*(K_{F_0})) = \frac{1}{3}. \quad (4.1)$$

Thus there exists a μ -exceptional prime divisor $E_0 \subseteq \text{Supp } E_\mu$ such that $(E_0|_F \cdot \sigma^*(K_{F_0})) > 0$, which means that $(E_0|_F \cdot \sigma^*(K_{F_0})) \geq 1$. Hence

$$((K_{X'} - E_0)|_F \cdot \sigma^*(K_{F_0})) = (K_F \cdot \sigma^*(K_{F_0})) - (E_0|_F \cdot \sigma^*(K_{F_0})) \leq 0. \quad (4.2)$$

On the other hand,

$$K_{X'} = F + Z + E_\mu$$

is an effective integral divisor whose support contains all μ -exceptional divisors. So $K_{X'} - E_0$ is an effective integral divisor. As $\sigma^*(K_{F_0})$ is nef, we conclude from (4.2) that

$$((K_{X'} - E_0)|_F \cdot \sigma^*(K_{F_0})) = 0; \quad (4.3)$$

$$(E_0|_F \cdot \sigma^*(K_{F_0})) = 1. \quad (4.4)$$

In particular, (4.3) implies that $\text{coeff}_{E_0}(Z + E_\mu) = 1$ and E_0 is the unique μ -exceptional prime divisor such that $(E_0|_F \cdot \sigma^*(K_{F_0})) > 0$. Furthermore, (4.1) and (4.4) imply that $\text{coeff}_{E_0} E_\mu = \frac{1}{3}$ and hence $\text{coeff}_{E_0} Z = \frac{2}{3}$.

Finally, since X has only isolated singularities and $\text{coeff}_{E_0} E_\mu \notin \mathbb{Z}$, we deduce that $\mu(E_0)$ is a point. \square

Lemma 4.3. *Let X be a minimal 3-fold of general type with $p_g(X) = 2$ and $K_X^3 = \frac{1}{3}$. Keep the notation in Setting 3.2 and Lemma 4.2. Then*

- (1) $E_0|_F = C_0$ is an integral (-3) -curve;
- (2) $G|_F = \frac{1}{3}C_0$;
- (3) $Z_X = 0$.

Proof. By Lemma 4.2(1), there is a unique integral curve C_0 on F with $\text{coeff}_{C_0} E_0|_F = 1$ such that $(\sigma^*(K_{F_0}) \cdot C_0) = 1$. In particular, this implies that C_0 is not contracted by σ and hence

$$(K_F \cdot C_0) \geq (\sigma^*(K_{F_0}) \cdot C_0) = 1. \quad (4.5)$$

Note that in Setting 3.2,

$$Z - E_\mu = \mu^*(2K_X) - K_{X'} - F = \mu^*(Z_X) + G \quad (4.6)$$

is an effective \mathbb{Q} -divisor. Hence

$$(Z - E_\mu)|_F = \mu^*(2K_X)|_F - K_F \quad (4.7)$$

is an effective \mathbb{Q} -divisor with

$$\text{coeff}_{C_0}(Z - E_\mu)|_F = \text{coeff}_{E_0}(Z - E_\mu) = \frac{1}{3}$$

by Lemma 4.2(2). Hence by (4.7),

$$\frac{1}{3}C_0^2 \leq ((Z - E_\mu)|_F \cdot C_0) = -(K_F \cdot C_0). \quad (4.8)$$

Here we used the fact that C_0 is contracted by μ by Lemma 4.2(3). Combining (4.5) and (4.8), we have

$$(K_F \cdot C_0) + C_0^2 \leq -2(K_F \cdot C_0) \leq -2,$$

According to the genus formula, we deduce that $(K_F \cdot C_0) + C_0^2 = -2$ and $C_0^2 = -3$, which means that C_0 is a (-3) -curve.

Since $\sigma^*(K_{F_0})$ is nef and

$$\left(\sigma^*(K_{F_0}) + \frac{1}{3}C_0 \right) \cdot C_0 = 1 - 1 = 0,$$

we know that $\sigma^*(K_{F_0}) + \frac{1}{3}C_0$ is nef. Moreover, $(\sigma^*(K_{F_0}) + \frac{1}{3}C_0)^2 = \frac{4}{3}$. On the other hand, from (4.7) we have

$$\begin{aligned} \mu^*(2K_X)|_F &= K_F + (Z - E_\mu)|_F \geq K_F + \frac{1}{3}E_0|_F \\ &\geq K_F + \frac{1}{3}C_0 \geq \sigma^*(K_{F_0}) + \frac{1}{3}C_0. \end{aligned} \quad (4.9)$$

Here $(\mu^*(2K_X)|_F)^2 = \frac{4}{3}$ by Lemma 4.1. Hence by Lemma 2.2(1), all inequalities in (4.9) become equalities. In particular, $(Z - E_\mu)|_F = \frac{1}{3}E_0|_F = \frac{1}{3}C_0$. Moreover, $(\mu^*(Z_X) + G)|_F = \frac{1}{3}C_0$ by (4.6), so $\mu^*(Z_X)|_F$ and $G|_F$ are multiples of C_0 . Since C_0 is contracted by μ by Lemma 4.2(3), we have $\mu^*(Z_X)|_F = 0$ and $G|_F = \frac{1}{3}C_0$. The last assertion follows from Lemma 3.5. \square

Lemma 4.4. *Let X be a minimal 3-fold of general type with $p_g(X) = 2$ and $K_X^3 = \frac{1}{3}$. Suppose that P is a non-Gorenstein singularity of X . Then $r_P \in \{2, 3\}$ and there is exactly one such P with $r_P = 3$. Here r_P is the Cartier index of K_X at P .*

Proof. By Lemma 4.3, all conditions of Lemma 3.4 are satisfied.

We claim that F_X has exactly one non-Gorenstein singularity which is of type $\frac{1}{3}(1, 1)$. Indeed, by Lemmas 4.3 and 3.4,

$$K_F + \frac{1}{3}C_0 = (\mu|_F)^*(K_{F_X}).$$

So outside the point $P_0 = \mu|_F(C_0)$, F_X has at worst Du Val singularities which are clearly Gorenstein. The fiber of $\mu|_F$ over P_0 has simple normal crossing support as F_X has only rational singularities, so the fiber of $\mu|_F$ over P_0 is exactly C_0 , otherwise there exists an exceptional curve C' connecting C_0 with

$$(K_F \cdot C') = \frac{1}{3}(-C_0 \cdot C') = -\frac{1}{3},$$

which is absurd. Since C_0 is a (-3) -curve, it is clear that C_0 is contracted to a cyclic quotient singularity of type $\frac{1}{3}(1, 1)$.

Now to conclude the lemma, by Lemma 3.4(2), we only need to consider the case that $P = P_0$ is the unique non-Gorenstein singularity of F_X . Then $r_P = 3$ as $3K_X|_{F_X}$ is Cartier at P . \square

4.2. Geometry of pluricanonical systems.

In this subsection, we study the pluricanonical maps of X .

Let $\pi : W \rightarrow X$ be as in Setting 2.1 and keep the same notation. We may further assume that π factorizes through the modification $\mu : X' \rightarrow X$ in §4.1. In this case, S is a $(1, 2)$ -surface. Denote by $\sigma_S : S \rightarrow S_0 = F_0$ the morphism to its minimal model. Let C be a general member of $\text{Mov } |K_S| = \text{Mov } |\sigma_S^* K_{S_0}|$. Then C is a smooth projective curve of genus 2. Here for a good reference on geometry of $(1, 2)$ -surfaces, we refer to [16, §2].

Recall that

$$(\pi^*(K_X)|_S)^2 = \frac{1}{3}, \quad (4.10)$$

$$(\pi^*(K_X)|_S \cdot C) = \frac{2}{3} \quad (4.11)$$

by Lemma 4.1 and the projection formula.

Lemma 4.5. *Let X be a minimal 3-fold of general type with $p_g(X) = 2$ and $K_X^3 = \frac{1}{3}$. Then $P_2(X) \leq 4$.*

Proof. Keep the notation in Setting 2.1.

Step 1. We claim that $u_{2,0} \leq 2$.

Suppose that $u_{2,0} \geq 3$. If $|M_{2,0}|_S$ and $|C|$ are composed with the same pencil, then $M_{2,0}|_S \geq 2C$ since $u_{2,0} \geq 3$. Thus $2\pi^*(K_X)|_S \geq 2C$, which implies that

$$K_X^3 \geq (\pi^*(K_X)|_S)^2 \geq (\pi^*(K_X)|_S \cdot C) = \frac{2}{3},$$

which is a contradiction. On the other hand, if $|M_{2,0}|_S$ and $|C|$ are not composed with the same pencil, then $(M_{2,0}|_S \cdot C) \geq 2$ as $|M_{2,0}|$ induces a nontrivial movable linear system on C . Hence

$$(\pi^*(K_X)|_S \cdot C) \geq \frac{1}{2}(M_{2,0}|_S \cdot C) \geq 1,$$

which contradicts (4.11). We conclude that $u_{2,0} \leq 2$.

Step 2. We claim that $u_{2,-1} \leq 1$.

Suppose that $u_{2,-1} \geq 2$. If $|M_{2,-1}|_S$ and $|C|$ are composed with the same pencil, then $M_{2,-1}|_S \geq C$ and hence

$$\begin{aligned} 2K_X^3 &\geq (\pi^*(K_X)^2 \cdot S) + (\pi^*(K_X)^2 \cdot M_{2,-1}) \\ &\geq (\pi^*(K_X)^2 \cdot S) + (\pi^*(K_X)|_S \cdot M_{2,-1}|_S) \\ &\geq (\pi^*(K_X)^2 \cdot S) + (\pi^*(K_X)|_S \cdot C) \\ &\geq 1, \end{aligned}$$

which is a contradiction. On the other hand, if $|M_{2,-1}|_S$ and $|C|$ are not composed with the same pencil, then $(M_{2,-1}|_S \cdot C) \geq 2$. Hence

$$(\pi^*(K_X)|_S \cdot C) \geq \frac{1}{2}(M_{2,-1}|_S \cdot C) \geq 1,$$

which contradicts (4.11). Thus $u_{2,-1} \leq 1$.

Step 3. We claim that $h^0(W, M_{2,-2}) \leq 1$.

Otherwise, $|M_{2,-2}|$ is a nontrivial movable linear system. Since K_X is nef and big, we have $(\pi^*(K_X)^2 \cdot M_{2,-2}) > 0$. Recall that $2\pi^*(K_X) - 2S \geq M_{2,-2}$ by definition. Then by (4.10),

$$K_X^3 \geq (\pi^*(K_X)|_S)^2 + \frac{1}{2}((\pi^*(K_X))^2 \cdot M_{2,-2}) > \frac{1}{3},$$

which is a contradiction. Thus $h^0(W, M_{2,-2}) \leq 1$.

Step 4. The conclusion that $P_2(X) \leq 4$.

This follows from

$$P_2(X) = h^0(W, \lfloor 2\pi^*(K_X) \rfloor) = u_{2,0} + u_{2,-1} + h^0(W, M_{2,-2})$$

and previous steps. \square

Lemma 4.6. *Let X be a minimal 3-fold of general type with $p_g(X) = 2$ and $K_X^3 = \frac{1}{3}$. Then*

- (1) $P_3(X) \geq P_2(X) + 3$;
- (2) $|3K_X|$ defines a generically finite map;
- (3) $P_4(X) \geq P_3(X) + 4$.

Proof. Keep the notation in Setting 2.1.

By the Kawamata–Viehweg vanishing theorem, the natural restriction map

$$H^0(W, K_W + \lceil \pi^*(K_X) \rceil + S) \rightarrow H^0(S, K_S + \lceil \pi^*(K_X)|_S \rceil)$$

is surjective. Here in the last term, the restriction on S commutes with the roundup as S is general. So

$$|3K_W|_S \succcurlyeq |K_W + \lceil \pi^*(K_X) \rceil + S|_S \succcurlyeq |K_S + \lceil \pi^*(K_X)|_S \rceil. \quad (4.12)$$

Note that $2\pi^*(K_X)|_S \geq \sigma_S^*(K_{S_0})$ by Lemma 3.3 as π factorizes through μ . By [2, Theorem 3.2], we have $h^0(S, K_S + \lceil \pi^*(K_X)|_S \rceil) \geq 3$. Then $u_{3,0} \geq 3$, which implies that

$$P_3(X) = h^0(W, \lfloor 3\pi^*(K_X) \rfloor - S) + u_{3,0} \geq P_2(X) + 3.$$

Assertion (1) is proved.

Next, we claim that $|K_S + \lceil \pi^*(K_X)|_S \rceil$ defines a generically finite map on S . Otherwise, it is composed with the same pencil with $|C|$ and thus $K_S + \lceil \pi^*(K_X)|_S \rceil \geq 2C$ since $h^0(S, K_S + \lceil \pi^*(K_X)|_S \rceil) \geq 3$. By comparing

the movable parts of both sides of (4.12), we know that $3\pi^*(K_X)|_S \geq 2C$. Thus by (4.11),

$$K_X^3 \geq (\pi^*(K_X)|_S)^2 \geq \frac{2}{3}(\pi^*(K_X)|_S \cdot C) = \frac{4}{9},$$

which is a contradiction. So $|K_S + \lceil \pi^*(K_X)|_S \rceil$ defines a generically finite map on S . By (4.12), $|3K_X|$ defines a generically finite map. Assertion (2) is proved.

By [10, Theorem 2.4(1)], the natural restriction map

$$H^0(W, 2(K_W + S)) \rightarrow H^0(S, 2K_S)$$

is surjective. Note that $P_2(S) = 4$ and $|4K_W| \gtrsim |2(K_W + S)|$. We deduce that $u_{4,0} \geq 4$, which implies that $P_4(X) \geq P_3(X) + 4$. \square

Proposition 4.7. *Let X be a minimal 3-fold of general type with $p_g(X) = 2$ and $K_X^3 = \frac{1}{3}$. Then*

- (1) $\chi(\mathcal{O}_X) = -1$, $P_2(X) = 4$;
- (2) the Reid basket of X is $\{2 \times (1, 2), (1, 3)\}$;
- (3) the Hilbert series of X is given by

$$\sum_{k \geq 0} h^0(X, kK_X)q^k = \frac{1 - q^{16}}{(1 - q)^2(1 - q^2)(1 - q^3)(1 - q^8)}.$$

All the above data are the same as those of a general hypersurface of degree 16 in $\mathbb{P}(1, 1, 2, 3, 8)$.

Proof. By Lemma 4.4 and [30, (6.1), (6.4)], the basket B_X of X is $\{a \times (1, 2), b \times (1, 3)\}$, for some integers $a \geq 0$ and $b \geq 1$. By Reid's Riemann–Roch formula (2.1) and the Kawamata–Viehweg vanishing theorem,

$$\begin{aligned} P_2(X) &= \frac{1}{6} - 3\chi(\mathcal{O}_X) + \frac{a}{4} + \frac{b}{3}; \\ P_3(X) &= \frac{5}{6} - 5\chi(\mathcal{O}_X) + \frac{a}{4} + \frac{2b}{3}; \\ P_4(X) &= \frac{7}{3} - 7\chi(\mathcal{O}_X) + \frac{a}{2} + \frac{2b}{3}. \end{aligned}$$

Then it is clear that

$$2P_4(X) - P_3(X) - 3P_2(X) = \frac{10}{3} - \frac{b}{3} \leq 3.$$

On the other hand, by Lemmas 4.6 and 4.5,

$$2P_4(X) - P_3(X) - 3P_2(X) = P_4(X) - P_3(X) + P_4(X) - P_2(X) - 2P_2(X) \geq 3.$$

Then all the above inequalities become equalities, which means that $b = 1$, $P_2(X) = 4$, $P_3(X) = 7$, $P_4(X) = 11$. Then we can solve the equations and get $\chi(\mathcal{O}_X) = -1$ and $a = 2$. Assertions (1) and (2) are proved.

Note that Reid's Riemann–Roch formula only depends on K_X^3 , $\chi(\mathcal{O}_X)$ and B_X . These data are the same as those of a general hypersurface of degree

16 in $\mathbb{P}(1, 1, 2, 3, 8)$ (see [18, Table 3, No. 12]). So the Hilbert series of them also coincide and are certainly computable (see [13, Theorem 3.4.4]). \square

Proposition 4.8. *Let X be a minimal 3-fold of general type with $p_g(X) = 2$ and $K_X^3 = \frac{1}{3}$. Then*

- (1) $|2K_X|$ is not composed with a pencil;
- (2) $|3K_X|$ defines a generically finite map;
- (3) $|8K_X|$ defines a birational map.

Proof. If $|2K_X|$ is composed with a pencil, then it is the same pencil as $|K_X|$ and $2\pi^*(K_X) \geq 3S$ since $P_2(X) = 4$. Thus by (4.10),

$$K_X^3 \geq \frac{3}{2}(\pi^*(K_X)|_S)^2 \geq \frac{1}{2},$$

which is a contradiction. This proves Assertion (1).

Assertion (2) follows from Lemma 4.6. Assertion (3) follows from [7, Theorem 1.4]. \square

5. BIRATIONAL GEOMETRY OF 3-FOLDS WITH $p_g = 3$ AND $\text{Vol} = 1$

In this section, we study the birational geometry of a minimal 3-fold X of general type with $p_g(X) = 3$ and $K_X^3 = 1$.

5.1. Singularities.

In this subsection, we study the Gorenstein index of X .

Take a birational modification $\mu : X' \rightarrow X$ as in Setting 3.2 and keep the same notation. By [7, Theorem 1.5(4)], the canonical image of X is \mathbb{P}^2 . By [8, Lemma 5.2], F is a (4, 4)-surface.

The geometry of X is simpler than that in §4.

Lemma 5.1. *Let X be a minimal 3-fold of general type with $p_g(X) = 3$ and $K_X^3 = 1$. Keep the notation in Setting 3.2. Then*

- (1) $\mu^*(K_X)|_F \equiv \frac{1}{2}K_F \equiv \frac{1}{2}\sigma^*(K_{F_0})$; in particular, F is minimal;
- (2) $G|_F = 0$, $Z_X = 0$.

Proof. By (3.1), we have

$$K_F + G|_F = \mu^*(K_X + F_X)|_F.$$

Since G is effective, we deduce that

$$2\mu^*(K_X)|_F = K_F + G|_F + \mu^*(Z_X)|_F \geq K_F \geq \sigma^*(K_{F_0}). \quad (5.1)$$

Then $(\mu^*(K_X)|_F)^2 \geq \frac{1}{4}K_{F_0}^2 = 1$. On the other hand, $(\mu^*(K_X)|_F)^2 \leq K_X^3 = 1$. Hence $(\mu^*(K_X)|_F)^2 = \frac{1}{4}K_{F_0}^2 = 1$. By Lemma 2.2(1), inequalities in (5.1) become equalities. In particular, Assertion (1) holds and $G|_F = \mu^*(Z_X)|_F = 0$. The last assertion follows from Lemma 3.5. \square

Lemma 5.2. *Let X be a minimal 3-fold of general type with $p_g(X) = 3$ and $K_X^3 = 1$. Suppose that P is a non-Gorenstein singularity of X . Then $r_P = 2$. Here r_P is the Cartier index of K_X at P .*

Proof. By Lemma 5.1, all conditions of Lemma 3.4 are satisfied. Moreover, $G|_F = 0$ implies that $K_F = (\mu|_F)^*(K_{F_X})$. So F_X has at worst Du Val singularities which are clearly Gorenstein, which implies that $r_P = 2$ by Lemma 3.4(2). \square

5.2. Geometry of pluricanonical systems.

Take a birational modification $\pi : W \rightarrow X$ as in Setting 2.1 and keep the same notation. We may further assume that π factorizes through the modification $\mu : X' \rightarrow X$ in §5.1. In this case, S is a $(4, 4)$ -surface. By [7, §3.2], $|S|_S$ is composed with a free rational pencil and $S|_S \sim C$, where C is a general fiber of $\psi : W \rightarrow \Sigma'$.

Lemma 5.3. *Let X be a minimal 3-fold of general type with $p_g(X) = 3$ and $K_X^3 = 1$. Then $P_2(X) \leq 7$. Moreover, the equality holds only if $|2K_X|$ defines a generically finite map of degree 2.*

Proof. Recall that $\Phi_{|2K_X|}$ denotes the rational map defined by $|2K_X|$. Denote by Σ_2 the image of $\Phi_{|2K_X|}$. Since the canonical image of X is \mathbb{P}^2 by [7, Theorem 1.5(4)], it is clear that $\dim \Sigma_2 \geq 2$.

If $\dim \Sigma_2 = 2$, then $|2K_X|$ induces a fibration $\psi_2 : W \rightarrow \Sigma'_2$ which factors through $\psi : W \rightarrow \Sigma'$ and Σ'_2 is birational to Σ' . In particular, we can identify a general fiber of ψ_2 with C . So we may write $M_2|_{M_2} \equiv d_2 C$, where M_2 is a general element in $\text{Mov } |2\pi^*(K_X)|$ and $d_2 \geq P_2(X) - 2$. Then

$$4K_X^3 \geq (\pi^*(K_X) \cdot M_2 \cdot M_2) \geq d_2(\pi^*(K_X) \cdot C) \geq d_2 \geq P_2(X) - 2,$$

where the third inequality follows from [7, §3.2] as $(\pi^*(K_X) \cdot C) = \xi \geq 1$. We deduce that $P_2(X) \leq 6$ when $\dim \Sigma_2 = 2$.

If $\dim \Sigma_2 = 3$, then by [7, Theorem 1.5(4)], $\Phi_{|4K_X|}$ is not birational. So $\deg(\Phi_{|2K_X|}) \geq 2$. Then

$$8K_X^3 \geq \deg(\Phi_{|2K_X|}) \deg(\Sigma_2) \geq 2(P_2(X) - 3),$$

where $\deg(\Sigma_2) \geq P_2(X) - 3$. We conclude that $P_2(X) \leq 7$. Moreover, when $P_2(X) = 7$, the above inequalities become equalities. In particular, $\deg(\Phi_{|2K_X|}) = 2$. \square

Lemma 5.4. *Let X be a minimal 3-fold of general type with $p_g(X) = 3$ and $K_X^3 = 1$. Then $P_3(X) \geq P_2(X) + 6$.*

Proof. Keep the notation in Setting 2.1. By the Kawamata–Viehweg vanishing theorem, we have

$$|3K_W||_S \succcurlyeq |K_W + [\pi^*(K_X)] + S||_S \succcurlyeq |K_S + [\pi^*(K_X)]_S|.$$

Since $\pi^*(K_X)|_S \geq S|_S = C$, we have

$$|3K_W||_S \succcurlyeq |K_S + C|. \quad (5.2)$$

By [15, Theorem 3.4], $h^1(\mathcal{O}_S) = 0$. By the exact sequence

$$0 \rightarrow \mathcal{O}_S(K_S) \rightarrow \mathcal{O}_S(K_S + C) \rightarrow \mathcal{O}_C(K_C) \rightarrow 0,$$

we deduce that $h^0(S, K_S + C) = p_g(S) + g(C) \geq 6$. Thus (5.2) implies that

$$P_3(X) \geq h^0(3K_W - S) + h^0(S, K_S + C) \geq P_2(X) + 6.$$

The proof is completed. \square

Proposition 5.5. *Let X be a minimal 3-fold of general type with $p_g(X) = 3$ and $K_X^3 = 1$. Then*

- (1) $\chi(\mathcal{O}_X) = -2$, $P_2(X) = 7$;
- (2) the basket of X is $B_X = \{2 \times (1, 2)\}$;
- (3) the Hilbert series of X is given by

$$\sum_{k \geq 0} h^0(X, kK_X)q^k = \frac{1 - q^{12}}{(1 - q)^3(1 - q^2)(1 - q^6)}.$$

All the above data are the same as those of a general hypersurface of degree 12 in $\mathbb{P}(1, 1, 1, 2, 6)$.

Proof. By Lemma 5.2 and [30, (6.1), (6.4)], the basket B_X of X is $\{a \times (1, 2)\}$ for some $a \geq 0$. By Reid's Riemann–Roch formula (2.1) and the Kawamata–Viehweg vanishing theorem, we have

$$\begin{aligned} P_2(X) &= \frac{1}{2} - 3\chi(\mathcal{O}_X) + \frac{a}{4}, \\ P_3(X) &= \frac{5}{2} - 5\chi(\mathcal{O}_X) + \frac{a}{4}. \end{aligned}$$

Then by Lemmas 5.4 and 5.3,

$$14 \geq 2P_2(X) = 3P_3(X) - 3P_2(X) - 5 + \frac{a}{2} \geq 13 + \frac{a}{2}.$$

So $P_2(X) = 7$ and $a \leq 2$. Then it is not hard to see that $a = 2$ and $\chi(\mathcal{O}_X) = -2$ from the equation of $P_2(X)$. Assertions (1) and (2) are proved.

Note that Reid's Riemann–Roch formula only depends on K_X^3 , $\chi(\mathcal{O}_X)$ and B_X . These data are the same as those of a general hypersurface of degree 12 in $\mathbb{P}(1, 1, 1, 2, 6)$ (see [18, Table 3, No. 7]). So the Hilbert series of them also coincide and can be computed by [13, Theorem 3.4.4]. \square

Proposition 5.6. *Let X be a minimal 3-fold of general type with $p_g(X) = 3$ and $K_X^3 = 1$. Then*

- (1) $|K_X|$ is not composed with a pencil;
- (2) $|2K_X|$ defines a generically finite map of degree 2;
- (3) $|6K_X|$ defines a birational map.

Proof. Assertion (1) follows from [7, Theorem 1.5(4)]. Assertion (2) follows from Lemma 5.3 and $P_2(X) = 7$ in Proposition 5.5. Assertion (3) follows from [6, Theorem 1.2]. \square

6. BIRATIONAL GEOMETRY OF 3-FOLDS WITH $p_g = 4$ AND $\text{Vol} = 2$

In this section, we study the birational geometry of a minimal 3-fold X of general type with $p_g(X) = 4$ and $K_X^3 = 2$.

Recall that Σ denotes the canonical image of X in \mathbb{P}^3 . By [7, Theorem 1.5(5)], we have $\dim \Sigma \geq 2$.

When $\dim \Sigma = 3$, the geometry of X is well-understood by Kobayashi [22]. We summarize the result with more details as the following.

Proposition 6.1. *Let X be a minimal 3-fold of general type with $p_g(X) = 4$ and $K_X^3 = 2$. Suppose that the canonical map of X is generically finite. Then*

- (1) X is Gorenstein;
- (2) the canonical map is a generically finite morphism to \mathbb{P}^3 of degree 2;
- (3) the canonical model of X is a hypersurface of degree 10 in $\mathbb{P}(1, 1, 1, 1, 5)$;
- (4) the Hilbert series of X is given by

$$\sum_{k \geq 0} h^0(X, kK_X)q^k = \frac{1 - q^{10}}{(1 - q)^4(1 - q^5)}.$$

Proof. Assertions (1) and (2) are due to [22, Proposition 2.5]. Denote by Y the canonical model of X and $\phi : Y \rightarrow \mathbb{P}^3$ the canonical map which is a morphism defined by $|K_Y|$. Then ϕ is finite as K_Y is ample. Moreover, $\deg(\phi) = 2$. Denote by $D \subseteq \mathbb{P}^3$ the branch locus, then $K_Y = \phi^*(K_{\mathbb{P}^3} + \frac{1}{2}D)$ by the ramification formula. By computing the self-intersection number, we get $D \in |\mathcal{O}_{\mathbb{P}^3}(10)|$. So Y can be expressed as a hypersurface of degree 10 in $\mathbb{P}(1, 1, 1, 1, 5)$ defined by $x_4^2 + f_0(x_0, x_1, x_2, x_3) = 0$, where $[x_0 : \cdots : x_4]$ are coordinates of $\mathbb{P}(1, 1, 1, 1, 5)$ and f_0 is the defining equation of $D \subseteq \mathbb{P}^3$. Finally, to compute the Hilbert series, we can use either [13, 3.4.2] or Reid's Riemann–Roch formula (2.1). Here recall that $K_X^3 = 2$ and $B_X = \emptyset$, and $\chi(\mathcal{O}_X) = -3$ which can be computed from the double cover construction. \square

Thus we only need to treat the case when $\dim \Sigma = 2$. We recall the following properties from [7, §3.2].

Lemma 6.2. *Let X be a minimal 3-fold of general type with $p_g(X) = 4$ and $K_X^3 = 2$. Suppose that the canonical map of X is not generically finite. Let $\pi : W \rightarrow X$ be as in Setting 2.1 and keep the same notation. Denote by C a general fiber of $\psi : W \rightarrow \Sigma'$. Then*

- (1) $(\pi^*(K_X) \cdot C) = 1$;
- (2) C is a smooth projective curve of genus 2;
- (3) $S|_S \sim 2C$ which is a rational pencil;
- (4) $\Sigma' \simeq \Sigma \subseteq \mathbb{P}^3$ is a nondegenerate surface of degree 2, in particular, Σ is either $\mathbb{P}(1, 1, 2)$ or $\mathbb{P}^1 \times \mathbb{P}^1$.

Proof. Assertions (1)–(3) are directly from [7, §3.2]. For Assertion (4), just note that $S|_S \equiv \deg(s) \deg(\Sigma)C$ and $\deg(\Sigma) \geq 2$ as it is nondegenerate.

So $\deg(\Sigma) = 2$ and $\deg(s) = 1$ by Assertion (3). By [28, §10], Σ is either $\mathbb{P}(1, 1, 2)$ or $\mathbb{P}^1 \times \mathbb{P}^1$. In particular, Σ is normal and hence s is an isomorphism. \square

By adopting the idea in [17, Proof of Proposition 2.1], we have the following lemma.

Lemma 6.3. *Let X be a minimal 3-fold of general type with $p_g(X) = 4$ and $K_X^3 = 2$. Suppose that the canonical map of X is not generically finite. Then there exists a minimal 3-fold X_1 birational to X , admitting a fibration $f_1 : X_1 \rightarrow \mathbb{P}^1$ whose general fibers are $(1, 2)$ -surfaces.*

Proof. By Lemma 6.2, Σ is either $\mathbb{P}(1, 1, 2)$ or $\mathbb{P}^1 \times \mathbb{P}^1$.

Case 1. $\Sigma \simeq \mathbb{P}(1, 1, 2)$.

In this case, there is a contraction $r : \mathbb{F}_2 \rightarrow \mathbb{P}^3$ from the second Hirzebruch surface induced by the linear system $|s + 2\ell|$ such that $\Sigma = r(\mathbb{F}_2)$. Here ℓ is the ruling of the natural fibration $p : \mathbb{F}_2 \rightarrow \mathbb{P}^1$ and s is the unique negative section.

Replacing W by its birational model, we may assume that there is a morphism $\varphi : W \rightarrow \mathbb{F}_2$ such that $\psi = r \circ \varphi$. Thus we obtain a fibration $f_W := p \circ \varphi : W \rightarrow \mathbb{P}^1$ with a general fiber $F_W = \varphi^*(\ell)$. Let $\zeta : W \dashrightarrow X_1$ be the contraction to a relative minimal model over \mathbb{P}^1 , where we have a relatively minimal fibration $f_1 : X_1 \rightarrow \mathbb{P}^1$ with a general fiber denoted by F_1 . Up to a further birational modification of W , we may assume that ζ is a morphism. Thus $\sigma_1 := \zeta|_{F_W} : F_W \rightarrow F_1$ is just the contraction to its minimal model. We claim that F_1 is a $(1, 2)$ -surface and X_1 is minimal.

Note that by definition,

$$\pi^*(K_X) \geq \psi^*\mathcal{O}_\Sigma(1) \geq \varphi^*(2\ell) = 2F_W.$$

By [4, Corollary 2.3], $\pi^*(K_X)|_{F_W} - \frac{2}{3}\sigma_1^*(K_{F_1})$ is \mathbb{Q} -effective. Note that $S|_{F_W} \sim C$. By Lemma 6.2(1), we have

$$(\sigma_1^*(K_{F_1}) \cdot C) \leq \frac{3}{2}(\pi^*(K_X)|_{F_W} \cdot C) = \frac{3}{2}.$$

Then $(\sigma_1^*(K_{F_1}) \cdot C) = 1$. Since C is a moving curve on F_W , we conclude that F_1 is a $(1, 2)$ -surface by [3, Lemma 2.4]. Note that $K_{F_W} \geq \pi^*(K_X)|_{F_W}$ and by Lemma 6.2(1),

$$(\pi^*(K_X)|_{F_W})^2 \geq (\pi^*(K_X)|_{F_W} \cdot S|_{F_W}) = (\pi^*(K_X)|_{F_W} \cdot C) = 1 = (\sigma_1^*(K_{F_1}))^2.$$

We deduce that $\pi^*(K_X)|_{F_W} = \sigma_1^*(K_{F_1})$ by considering the Zariski decomposition of K_{F_W} and Lemma 2.2(1). Thus X_1 is minimal by [4, Lemma 3.2]. We finish the proof in this case.

Case 2. $\Sigma \simeq \mathbb{P}^1 \times \mathbb{P}^1$.

Denote by p_1 and p_2 the two natural projections of $\mathbb{P}^1 \times \mathbb{P}^1$. For $i = 1, 2$, write $f_{W,i} = p_i \circ \gamma : W \rightarrow \mathbb{P}^1$. Then $f_{W,i}$ is a fibration for each i . Let T_i be

a general fiber of $f_{W,i}$. Then

$$\pi^*(K_X) \geq \psi^*(\mathcal{O}_\Sigma(1)) = T_1 + T_2. \quad (6.1)$$

Since $K_X^3 = 2$, we may assume that $(\pi^*(K_X)|_{T_1})^2 \leq 1$. Same as in Case 1, we get a contraction $\zeta : W \rightarrow X_1$ to a relative minimal model of $f_{W,1}$, where we denote by $f_1 : X_1 \rightarrow \mathbb{P}^1$ the relatively minimal fibration and by F_1 its general fiber. Thus $\sigma_1 := \zeta|_{T_1} : T_1 \rightarrow F_1$ is just the contraction to its minimal model. We claim that F_1 is a $(1,2)$ -surface and X_1 is minimal.

For sufficiently large m , we have

$$|2mK_W|_{T_1} \succcurlyeq |m(K_W + T_1 + T_2)|_{T_1} \succcurlyeq |m(K_{T_1} + T_2|_{T_1})|,$$

where the first inequality is by (6.1) and the second is by [10, Theorem 2.4(2)]. Hence by considering the movable parts of both sides, we deduce that $\pi^*(2K_X)|_{T_1} - \sigma_1^*(K_{F_1}) - T_2|_{T_1}$ is \mathbb{Q} -effective. Note that $T_2|_{T_1} = C$. Combining with Lemma 6.2(1) and taking the intersection with $\pi^*(K_X)|_{T_1}$, we get

$$(\pi^*(K_X)|_{T_1} \cdot \sigma_1^*(K_{F_1})) \leq 2(\pi^*(K_X)|_{T_1})^2 - (\pi^*(K_X)|_{T_1} \cdot C) \leq 1.$$

Since $\pi^*(K_X)|_{T_1} \geq C$ by (6.1), we have $(C \cdot \sigma_1^*(K_{F_1})) \leq 1$. By [3, Lemma 2.4], F_1 is a $(1,2)$ -surface. Furthermore, by Lemma 6.2(1),

$$1 \geq (\pi^*(K_X)|_{T_1})^2 \geq (\pi^*(K_X)|_{T_1} \cdot C) = 1.$$

Thus $(\pi^*(K_X)|_{T_1})^2 = 1$. By considering the Zariski decomposition of K_{T_1} and Lemma 2.2(1), we have $\pi^*(K_X)|_{T_1} = \sigma_1^*(K_{F_1})$. By [4, Lemma 3.2], X_1 is minimal. The proof is completed. \square

Proposition 6.4. *Let X be a minimal 3-fold of general type with $p_g(X) = 4$ and $K_X^3 = 2$. Suppose that the canonical map of X is not generically finite. Then*

- (1) $h^1(\mathcal{O}_X) = h^2(\mathcal{O}_X) = 0$; in particular, $\chi(\mathcal{O}_X) = -3$;
- (2) X is Gorenstein;
- (3) the Hilbert series of X is given by

$$\sum_{k \geq 0} h^0(X, kK_X)q^k = \frac{1 - q^{10}}{(1 - q)^4(1 - q^5)};$$

- (4) $|2K_X|$ defines a generically finite map of degree 2;
- (5) $|5K_X|$ defines a birational map.

Proof. By Lemma 6.3, after replacing by a birational model, we may assume that X admits a fibration $f : X \rightarrow \mathbb{P}^1$ with a general fiber F a $(1,2)$ -surface. Now we are in the setting of [17, §3]. Assertion (1) follows from [17, Lemma 3.5].

Recall Reid's Riemann–Roch formula (2.1) for $P_2(X)$:

$$P_2(X) = \frac{1}{2}K_X^3 - 3\chi(\mathcal{O}_X) + l_2(X).$$

Here $l_2(X) \geq 0$ and $l_2(X) = 0$ if and only if X is Gorenstein. So $P_2(X) \geq 10$. On the other hand, since $K_X^3 < \frac{4}{3}p_g(X) - \frac{17}{6}$, by [17, Proposition 3.8], we have

$$P_2(X) \leq \lfloor 2K_X^3 \rfloor + \lfloor 2K_X^3 - \frac{5(p_g(X) - 1)}{3} \rfloor + 7 = 10.$$

So $P_2(X) = 10$. Thus $l_2(X) = 0$ and X is Gorenstein. Assertion (2) is proved.

For Assertion (3), note that Reid's Riemann–Roch formula only depends on K_X^3 , $\chi(\mathcal{O}_X)$ and B_X . These data are the same as those of a general hypersurface of degree 10 in $\mathbb{P}(1, 1, 1, 1, 5)$ (see [18, Table 3, No. 5]). So the Hilbert series of them also coincide and can be computed by [13, Theorem 3.4.4].

For Assertion (4), note that the inequality in [17, Proposition 3.8(1)] is an equality now. The inequality in Step 1 of [17, Proof of Proposition 3.8] is also an equality. In particular, $|2K_W|_F$ defines a generically finite map of degree at most 2. Thus $|2K_X|$ defines a generically finite map of degree at most 2. However, $|2K_X|$ does not define a birational map as $|2K_F|$ does not define a birational map on a $(1, 2)$ -surface F by [16, Lemma 2.1]. So $|2K_X|$ defines a generically finite map of degree 2.

Assertion (5) follows from [6, Theorem 1.2]. \square

7. CHARACTERIZATION OF WEIGHTED HYPERSURFACES

The following theorem is a refinement of [20]. It provides a criterion for embedding a polarized variety into a weighted projective space by numerical and geometric properties of this variety.

Theorem 7.1. *Let Y be a normal projective 3-fold and let H be a \mathbb{Q} -Cartier Weil ample divisor on Y . Suppose that there exists a well-formed quasi-smooth weighted hypersurface*

$$X_d \subseteq \mathbb{P}(1, a_1, a_2, a_3, a_4)$$

with $1 \leq a_1 \leq a_2 \leq a_3 < a_4$ and $d = 2a_4$ such that

- (1) $H^3 = (\mathcal{O}_{X_d}(1))^3 = \frac{2}{a_1 a_2 a_3}$ and
- (2) $h^0(Y, kH) = h^0(X_d, \mathcal{O}_{X_d}(k))$ for any nonnegative integer k .

Suppose further that

- (3) $|a_2 H|$ is not composed with a pencil;
- (4) $|a_3 H|$ defines a generically finite map, while either the degree of this map is greater than 1, or Y is not rational;
- (5) $|a_4 H|$ defines a birational map.

Then Y is a hypersurface in $\mathbb{P}(1, a_1, a_2, a_3, a_4)$ defined by a weighted homogeneous polynomial f of degree d , where

$$f(x_0, \dots, x_4) = x_4^2 + f_0(x_0, x_1, x_2, x_3)$$

in suitable homogeneous coordinates $[x_0 : \cdots : x_4]$ of $\mathbb{P}(1, a_1, a_2, a_3, a_4)$; in other words, there is an isomorphism between graded \mathbb{C} -algebras

$$\bigoplus_{k \geq 0} H^0(Y, kH) \simeq \mathbb{C}[x_0, \dots, x_4]/(f),$$

where $\mathbb{C}[x_0, \dots, x_4]$ is viewed as a weighted polynomial ring with

$$\text{wt}(x_0, \dots, x_4) = (1, a_1, a_2, a_3, a_4).$$

Remark 7.2. In Theorem 7.1, by [13, 3.4.2], we have the following formula:

$$\sum_{k \geq 0} h^0(X_d, \mathcal{O}_{X_d}(k))q^k = \frac{1 - q^d}{(1 - q)(1 - q^{a_1})(1 - q^{a_2})(1 - q^{a_3})(1 - q^{a_4})}.$$

We will always use this formula to compute $h^0(Y, kH)$. In practice, we do not need to calculate the precise value of $h^0(Y, kH)$, we will often identify the value with the cardinality of a certain set in a combinatorial way instead.

Lemma 7.3. *Keep the notation in Theorem 7.1. For $0 \leq i \leq 4$, take distinct general elements $g_i \in H^0(Y, a_i H) \setminus \{0\}$ (where $a_0 = 1$). For a positive integer k , denote*

$$S_k = \{g_0^{s_0} g_1^{s_1} g_2^{s_2} g_3^{s_3} \mid s_0, \dots, s_3 \in \mathbb{Z}_{\geq 0}, s_0 + a_1 s_1 + a_2 s_2 + a_3 s_3 = k\}.$$

Then

- (1) the set $\{g_0, g_1, g_2, g_3\}$ is algebraically independent in the graded algebra $R(Y, H) = \bigoplus_{k \geq 0} H^0(Y, kH)$;
- (2) $H^0(Y, a_4 H)$ is spanned by a basis $S_{a_4} \cup \{g_4\}$.

Here we remark that $R(Y, H)$ is viewed as a subalgebra of $\mathbb{C}(Y)[u]$, where the variable u is from the grading of $R(Y, H)$. So a rational function $g \in H^0(Y, kH) \subseteq \mathbb{C}(Y)$ is identified with $gu^k \in R(Y, H)$. But to simplify the notation, we will just write $g \in R(Y, H)$ as long as the grading of g is clear in the context.

Proof. First we show the following claim.

Claim 7.4. *The set $\{g_0, g_1, g_2\}$ is algebraically independent in $R(Y, H)$.*

Proof. As $h^0(Y, a_1 H) \geq 2$, $\{g_0, g_1\}$ is algebraically independent. By assumption, $|a_2 H|$ is not composed with a pencil. So by the generality of g_2 , the transcendental degree of $\mathbb{C}(g_0, g_1, g_2)$ as a subfield of the fractional field of $R(Y, H)$ is greater than 2, which implies that $\{g_0, g_1, g_2\}$ is algebraically independent. \square

By Claim 7.4,

$$S'_k = \{g_0^{s_0} g_1^{s_1} g_2^{s_2} \mid s_0, s_1, s_2 \in \mathbb{Z}_{\geq 0}, s_0 + a_1 s_1 + a_2 s_2 = k\}$$

is linearly independent in $H^0(Y, kH)$ for any positive integer k . On the other hand,

$$h^0(Y, a_3 H) = h^0(X_d, \mathcal{O}_{X_d}(a_3)) = |S_{a_3}| = |S'_{a_3}| + 1.$$

Then $S_{a_3} = S'_{a_3} \cup \{g_3\}$ is a basis of $H^0(Y, a_3H)$ by the generality of g_3 . By assumption, $|a_3H|$ defines a generically finite map, so the transcendental degree of $\mathbb{C}(g_0, g_1, g_2, g_3)$ as a subfield of the fractional field of $R(Y, H)$ is greater than 3, which implies that $\{g_0, g_1, g_2, g_3\}$ is algebraically independent. This proves Assertion (1).

In particular, S_k is linearly independent in $H^0(Y, kH)$ for any positive integer k . So Assertion (2) follows from the computation that

$$h^0(Y, a_4H) = h^0(X_d, \mathcal{O}_{X_d}(a_4)) = |S_{a_4}| + 1$$

and the generality of g_4 . \square

Proof of Theorem 7.1. Keep the notation in Lemma 7.3. We can define 3 rational maps:

$$\begin{aligned} \Psi_{a_2} : Y &\dashrightarrow \mathbb{P}(1, a_1, a_2); \\ P &\mapsto [g_0(P) : g_1(P) : g_2(P)]; \\ \Psi_{a_3} : Y &\dashrightarrow \mathbb{P}(1, a_1, a_2, a_3); \\ P &\mapsto [g_0(P) : g_1(P) : g_2(P) : g_3(P)]; \\ \Psi_{a_4} : Y &\dashrightarrow \mathbb{P}(1, a_1, a_2, a_3, a_4); \\ P &\mapsto [g_0(P) : \cdots : g_4(P)]. \end{aligned}$$

We claim that they have the following geometric properties.

- Claim 7.5.** (1) Ψ_{a_2} is dominant; Ψ_{a_3} is dominant and generically finite of degree 2;
 (2) Ψ_{a_4} is birational onto its image;
 (3) let Y' be the closure of $\Psi_{a_4}(Y)$ in $\mathbb{P}(1, a_1, a_2, a_3, a_4)$, then Y' is defined by a weighted homogeneous polynomial f of degree d , where

$$f(x_0, \dots, x_4) = x_4^2 + f_0(x_0, x_1, x_2, x_3)$$

in suitable homogeneous coordinates $[x_0 : \cdots : x_4]$ of $\mathbb{P}(1, a_1, a_2, a_3, a_4)$.

Proof. (1) By Lemma 7.3, $\{g_0, g_1, g_2, g_3\}$ is algebraically independent, so Ψ_{a_2} and Ψ_{a_3} are dominant. In particular, Ψ_{a_3} is generically finite by dimension reason. The degree of Ψ_{a_3} is the number of points in the fiber over a general point in $\mathbb{P}(1, a_1, a_2, a_3)$. After taking a resolution $\pi : W \rightarrow Y$ such that $N_k = \text{Mov} |[\pi^*(a_kH)]|$ is free for $k = 1, 2, 3$, the degree is just the intersection number $(N_1 \cdot N_2 \cdot N_3)$. So

$$\deg(\Psi_{a_3}) = (N_1 \cdot N_2 \cdot N_3) \leq a_1 a_2 a_3 H^3 = 2.$$

On the other hand, it is clear that $\deg(\Psi_{a_3}) > 1$ by the assumption that either $|a_3H|$ defines a generically finite map of degree greater than 1 or Y is not rational.

(2) By assumption, $|a_4H|$ defines a birational map. As g_4 is general, it can separate two points in a general fiber of Ψ_{a_3} , so Ψ_{a_4} is birational onto its image.

(3) Note that $d = 2a_4$ and $h^0(Y, dH) = |S_{2a_4}| + |S_{a_4}|$. On the other hand,

$$S_{2a_4} \sqcup (S_{a_4} \cdot g_4) \sqcup \{g_4^2\} \subseteq H^0(Y, dH).$$

So $S_{2a_4} \sqcup (S_{a_4} \cdot g_4) \sqcup \{g_4^2\}$ is linearly dependent in $H^0(Y, dH)$. In other words, there exists a weighted homogeneous polynomial $f(x_0, \dots, x_4)$ of degree d with $\text{wt}(x_0, \dots, x_4) = (1, a_1, a_2, a_3, a_4)$ such that $f(g_0, \dots, g_4) = 0$. So Y' is contained in $(f = 0) \subseteq \mathbb{P}(1, a_1, a_2, a_3, a_4)$. Note that Y' is a hypersurface in $\mathbb{P}(1, a_1, a_2, a_3, a_4)$ by dimension reason.

We claim that $Y' = (f = 0)$ and x_4^2 has nonzero coefficient in f . Otherwise, either Y' is defined by a weighted homogeneous polynomial of degree at most $d - 1$, or x_4^2 has zero coefficient in f . In either case, Y' is defined by a weighted homogeneous polynomial \tilde{f} of the form

$$\tilde{f}(x_0, \dots, x_4) = x_4 \tilde{f}_1(x_0, x_1, x_2, x_3) + \tilde{f}_2(x_0, x_1, x_2, x_3).$$

Here $\tilde{f}_1 \neq 0$ as $\{g_0, g_1, g_2, g_3\}$ is algebraically independent. Then Y' is birational to $\mathbb{P}(1, a_1, a_2, a_3)$ under the rational projection map

$$\begin{aligned} \mathbb{P}(1, a_1, a_2, a_3, a_4) &\dashrightarrow \mathbb{P}(1, a_1, a_2, a_3); \\ [x_0 : \dots : x_4] &\mapsto [x_0 : \dots : x_3]. \end{aligned}$$

But this contradicts the fact that $\deg(\Psi_{a_3}) = 2$. So $Y' = (f = 0)$ and x_4^2 has nonzero coefficient in f . After a suitable coordinate transformation, we may assume that $f = x_4^2 + f_0(x_0, x_1, x_2, x_3)$. \square

Now go back to the proof of Theorem 7.1. By Claim 7.5, f is the only algebraic relation on g_0, \dots, g_4 . Denote \mathcal{R} to be the graded sub- \mathbb{C} -algebra of $R(Y, H)$ generated by $\{g_0, \dots, g_4\}$. Then there is a natural isomorphism between graded \mathbb{C} -algebras

$$\mathcal{R} \simeq \mathbb{C}[x_0, \dots, x_4]/(x_4^2 + f_0(x_0, x_1, x_2, x_3))$$

by sending $g_i \mapsto x_i$ ($0 \leq i \leq 4$) and the right-hand side is exactly the coordinate ring of Y' . Write $\mathcal{R} = \bigoplus_{k \geq 0} \mathcal{R}_k$, where \mathcal{R}_k is the homogeneous part of degree k . Then, by [13, 3.4.2],

$$\sum_{k \geq 0} (\dim_{\mathbb{C}} \mathcal{R}_k) q^k = \frac{1 - q^d}{(1 - q)(1 - q^{a_1})(1 - q^{a_2})(1 - q^{a_3})(1 - q^{a_4})}.$$

So $\dim_{\mathbb{C}} \mathcal{R}_k = h^0(X_d, \mathcal{O}_{X_d}(k)) = h^0(Y, kH)$ for any $k \in \mathbb{Z}_{\geq 0}$, and hence the inclusion $\mathcal{R} \subseteq R(Y, H)$ is an identity. This implies that

$$Y \simeq \text{Proj } R(Y, H) = \text{Proj } \mathcal{R} \simeq Y'.$$

This finishes the proof. \square

8. PROOFS OF THEOREM 1.1, THEOREM 1.2 AND THEOREM 1.3

8.1. Plurigenera.

Proof of Theorems 1.1(1), 1.2(1), and 1.3(1). These are directly from Propositions 4.7, 5.5, 6.1, and 6.4. \square

8.2. Weighted embedding.

Proof of Theorems 1.1(2) and 1.2(2). We may replace W by its canonical model Y and apply Theorem 7.1 to $H = K_Y$. Then Theorem 1.1(2) is a direct consequence of Propositions 4.7 and 4.8; Theorem 1.2(2) is a direct consequence of Propositions 5.5 and 5.6. \square

Proof of Theorem 1.3(2). We may replace W by its canonical model Y .

Suppose that the canonical map of Y is generically finite, then we get (2a) by Proposition 6.1.

Suppose that the canonical map of Y is not generically finite. Then we are going to embed Y into $\mathbb{P}(1, 1, 1, 1, 2, 5)$. Theorem 7.1 is not directly applicable here as we need 2 defining equations, but the proof is similar to that of Theorem 7.1 with some essential modification.

Since $p_g(Y) = 4$, we may take $\{g_0, g_1, g_2, g_3\}$ as a basis of $H^0(Y, K_Y)$. Take general elements $g_4 \in H^0(Y, 2K_Y)$ and $g_5 \in H^0(Y, 5K_Y)$. We regard g_i as elements in $R(Y) = \bigoplus_{k \geq 0} H^0(Y, kK_Y)$.

By Lemma 6.3, the canonical image of Y in \mathbb{P}^3 is a surface of degree 2, so g_0, g_1, g_2, g_3 satisfy a unique quadratic algebraic relation $q(g_0, g_1, g_2, g_3) = 0$. Without loss of generality, we may assume that q is of the form

$$q(x_0, x_1, x_2, x_3) = x_3^2 + q_0(x_0, x_1, x_2).$$

In particular, $\{g_0, g_1, g_2\}$ is algebraically independent and $\mathbb{C}(g_0, \dots, g_3)$ is a field extension of $\mathbb{C}(g_0, g_1, g_2)$ of degree 2. Then, for any $k \geq 0$,

$$S_k = \{g_0^{s_0} g_1^{s_1} g_2^{s_2} g_3^{s_3} \mid s_0, \dots, s_3 \in \mathbb{Z}_{\geq 0}, s_0 + s_1 + s_2 + s_3 = k \text{ and } s_3 \leq 1\}$$

is linearly independent in $H^0(Y, kK_Y)$. By Proposition 6.4(3),

$$H^0(Y, 2K_Y) = 10 = |S_2| + 1,$$

so $S_2 \sqcup \{g_4\}$ forms a basis of $H^0(Y, 2K_Y)$. Since $|2K_Y|$ defines a generically finite map of degree 2 by Proposition 6.4(4), $\{g_0, g_1, g_2, g_4\}$ is algebraically independent. Then $S_5 \sqcup (S_3 \cdot g_4) \sqcup (S_1 \cdot g_4^2)$ is linearly independent in $H^0(Y, 5K_Y)$. By Proposition 6.4(3),

$$H^0(Y, 5K_Y) = 57 = |S_5| + |S_3| + |S_1| + 1,$$

hence $S_5 \sqcup (S_3 \cdot g_4) \sqcup (S_1 \cdot g_4^2) \sqcup \{g_5\}$ forms a basis of $H^0(Y, 5K_Y)$.

Now we can consider 2 rational maps:

$$\begin{aligned} \Psi_2 : Y &\dashrightarrow \mathbb{P}(1, 1, 1, 1, 2); \\ P &\mapsto [g_0(P) : \dots : g_4(P)]; \\ \Psi_5 : Y &\dashrightarrow \mathbb{P}(1, 1, 1, 1, 2, 5); \\ P &\mapsto [g_0(P) : \dots : g_5(P)]. \end{aligned}$$

Then Ψ_2 is generically finite of degree 2 as $|2K_Y|$ defines a generically finite map of degree 2; moreover, as $|5K_Y|$ defines a birational map by Proposition 6.4(5), g_5 separates fibers of Ψ_2 and hence Ψ_5 is birational onto its image.

By Proposition 6.4(3),

$$h^0(Y, 10K_Y) = 342 = \sum_{i=0}^5 |S_{10-2i}| + \sum_{j=0}^2 |S_{5-2j}|.$$

On the other hand,

$$\bigsqcup_{i=0}^5 (S_{10-2i} \cdot g_4^i) \sqcup \bigsqcup_{j=0}^2 (S_{5-2j} \cdot g_4^j g_5) \sqcup \{g_5^2\} \subseteq H^0(Y, 10K_Y).$$

So there is a weighted homogeneous polynomial f of degree 10 such that $f(g_0, \dots, g_5) = 0$.

Let Y' be the closure of $\Psi_5(Y)$ in $\mathbb{P}(1, 1, 1, 1, 2, 5)$. Then clearly $Y' \subseteq (q = f = 0)$. We claim that $Y' = (q = f = 0)$ and x_5^2 has nonzero coefficient in f . Otherwise, Y' is contained in $(q = \tilde{f} = 0)$, where \tilde{f} of the form

$$\tilde{f}(x_0, \dots, x_5) = x_5 \tilde{f}_1(x_0, \dots, x_4) + \tilde{f}_2(x_0, \dots, x_4).$$

Here \tilde{f} is not divisible by q , which implies that $\tilde{f}_1 \neq 0$. But then Y' is birational to $\Psi_2(Y)$ under the rational projection map

$$\mathbb{P}(1, 1, 1, 1, 2, 5) \dashrightarrow \mathbb{P}(1, 1, 1, 1, 2),$$

which contradicts the fact that Ψ_2 is not birational. So $Y' = (q = f = 0)$ and x_5^2 has nonzero coefficient in f . After a suitable coordinate transformation, we may assume that $f = x_5^2 + f_0(x_0, \dots, x_4)$.

Now q and f are the only algebraic relations on g_0, \dots, g_5 . Denote \mathcal{R} to be the graded sub- \mathbb{C} -algebra of $R(Y)$ generated by $\{g_0, \dots, g_5\}$. Then there is a natural isomorphism between graded \mathbb{C} -algebras

$$\mathcal{R} \simeq \mathbb{C}[x_0, \dots, x_5]/(q, f)$$

by sending $g_i \mapsto x_i$ ($0 \leq i \leq 5$) and the right-hand side is exactly the coordinate ring of Y' . Write $\mathcal{R} = \bigoplus_{k \geq 0} \mathcal{R}_k$, where \mathcal{R}_k is the homogeneous part of degree k . Then, by [13, 3.4.2],

$$\sum_{k \geq 0} (\dim_{\mathbb{C}} \mathcal{R}_k) q^k = \frac{(1 - q^2)(1 - q^{10})}{(1 - q)^4(1 - q^2)(1 - q^5)} = \frac{1 - q^{10}}{(1 - q)^4(1 - q^5)}.$$

So $\dim_{\mathbb{C}} \mathcal{R}_k = h^0(Y, kK_Y)$ for any $k \in \mathbb{Z}_{\geq 0}$ by Proposition 6.4(3), and hence the inclusion $\mathcal{R} \subseteq R(Y)$ is an identity. This implies that

$$Y \simeq \text{Proj } R(Y) = \text{Proj } \mathcal{R} \simeq Y'.$$

This finishes the proof of (2b).

Finally, for a 3-fold Y defined by $(q = f = 0)$ in $\mathbb{P}(1, 1, 1, 1, 2, 5)$, we can view it as a specialization of Y_t ($t \in \mathbb{C}^*$) which is defined by

$$(q + tx_4 = f + tg = 0),$$

where g is a general weighted polynomial in x_0, \dots, x_3, x_5 of degree 10. In particular, $(g = 0) \subseteq \mathbb{P}(1, 1, 1, 1, 5)$ is smooth. Then Y_t turns out to be a 3-fold defined by

$$f(x_0, x_1, x_2, x_3, -\frac{1}{t}q, x_5) + tg(x_0, x_1, x_2, x_3, x_5) = 0$$

in $\mathbb{P}(1, 1, 1, 1, 5)$ with coordinates $[x_0 : x_1 : x_2 : x_3 : x_5]$, which is actually smooth for general $t \neq 0$. \square

8.3. Simply connectedness.

Proof of Theorems 1.1(3), 1.2(3), and 1.3(3). Let W be as in the assumption. By (2) of each theorem, the canonical model Y of W is a specialization of general hypersurfaces in a certain weighted projective space. Namely, there is a flat morphism $\mathcal{Y} \rightarrow C$ to a smooth curve such that the central fiber \mathcal{Y}_o at $o \in C$ is isomorphic to Y and a general fiber \mathcal{Y}_t for $t \neq o$ is a general weighted hypersurface. By [23, Lemma 2.8.1], after shrinking C , we may assume that $\mathcal{Y} \setminus \mathcal{Y}_o \rightarrow C \setminus \{o\}$ is a topological fiber bundle. Note that \mathcal{Y} is normal by [27, Theorem 23.9]. Hence by [23, Theorem 2.12], there is a surjective map $\pi_1(\mathcal{Y}_t) \rightarrow \pi_1(\mathcal{Y}_o)$ for general $t \in C$. Such \mathcal{Y}_t is quasismooth by [18] and hence is simply connected by [13, Theorem 3.2.4]. Hence Y is also simply connected. Since fundamental group is a birational invariant among projective klt varieties by [32, Theorem 1.1], we conclude the simply connectedness of W . \square

8.4. Irreducibility and dimensions of moduli spaces.

Proof of Theorem 1.1(3). Write $\mathbb{P} = \mathbb{P}(1, 1, 2, 3, 8)$. By Theorem 1.1(2), it is clear that $\mathcal{M}_{\frac{1}{3}, 2}$ is irreducible. Furthermore,

$$\dim \mathcal{M}_{\frac{1}{3}, 2} = h^0(\mathbb{P}, \mathcal{O}_{\mathbb{P}}(16)) - 1 - \dim \text{Aut } \mathbb{P}.$$

By an easy computation, $h^0(\mathbb{P}, \mathcal{O}_{\mathbb{P}}(16)) = 246$. By [11, §4] and [12, Proposition 4.3], we have

$$\begin{aligned} \dim \text{Aut}(\mathbb{P}) &= 2h^0(\mathcal{O}_{\mathbb{P}}(1)) + h^0(\mathcal{O}_{\mathbb{P}}(2)) + h^0(\mathcal{O}_{\mathbb{P}}(3)) + h^0(\mathcal{O}_{\mathbb{P}}(8)) - 1 \\ &= 56. \end{aligned}$$

Therefore, $\dim \mathcal{M}_{\frac{1}{3}, 2} = 189$. \square

Proof of Theorem 1.2(3). Write $\mathbb{P} = \mathbb{P}(1, 1, 1, 2, 6)$. By Theorem 1.2(2), it is clear that $\mathcal{M}_{1, 3}$ is irreducible. Furthermore,

$$\dim \mathcal{M}_{1, 3} = h^0(\mathbb{P}, \mathcal{O}_{\mathbb{P}}(12)) - 1 - \dim \text{Aut } \mathbb{P}.$$

By an easy computation, $h^0(\mathbb{P}, \mathcal{O}_{\mathbb{P}}(12)) = 303$. By [11, §4] and [12, Proposition 4.3], we have

$$\dim \text{Aut}(\mathbb{P}) = 3h^0(\mathcal{O}_{\mathbb{P}}(1)) + h^0(\mathcal{O}_{\mathbb{P}}(2)) + h^0(\mathcal{O}_{\mathbb{P}}(6)) - 1 = 66.$$

Therefore, $\dim \mathcal{M}_{1, 3} = 236$. \square

Proof of Theorem 1.3(4). Write $\mathbb{P} = \mathbb{P}(1, 1, 1, 1, 5)$. By Theorem 1.3(2), it is clear that $\mathcal{M}_{2,4}$ is irreducible. Furthermore,

$$\dim \mathcal{M}_{2,4} = h^0(\mathbb{P}, \mathcal{O}_{\mathbb{P}}(10)) - 1 - \dim \text{Aut } \mathbb{P}.$$

By an easy computation, $h^0(\mathbb{P}, \mathcal{O}_{\mathbb{P}}(10)) = 343$. By [11, §4] and [12, Proposition 4.3], we have

$$\dim \text{Aut}(\mathbb{P}) = 4h^0(\mathcal{O}_{\mathbb{P}}(1)) + h^0(\mathcal{O}_{\mathbb{P}}(5)) - 1 = 72.$$

Therefore, $\dim \mathcal{M}_{2,4} = 270$. \square

9. A LOWER BOUND FOR CANONICAL VOLUMES OF 3-FOLDS OF GENERAL TYPE WITH $p_g = 5$

Throughout this section, we always assume that X is a minimal 3-fold of general type with $p_g(X) \geq 5$.

Setting 9.1. Let $\pi : W \rightarrow X$ be as in §2.3 with respect to $|K_X|$ and keep the same notation. We have the following commutative diagram:

$$\begin{array}{ccc} W & \xrightarrow{\psi} & \Sigma' \\ \pi \downarrow & \searrow \gamma & \downarrow s \\ X & \xrightarrow{\Phi_{|K_X|}} & \Sigma \end{array}$$

We may write

$$\begin{aligned} K_W &= \pi^*(K_X) + E_\pi, \\ \pi^*(K_X) &= M + Z_\pi, \end{aligned}$$

where $|M| = \text{Mov } |\pi^*(K_X)|$, E_π, Z_π are effective \mathbb{Q} -divisor.

Let S be a general irreducible element of $|M|$. Then S is a smooth projective surface of general type. Denote by $\sigma_S : S \rightarrow S_0$ the contraction to its minimal model.

After taking some further blow-up, we may further assume that the following hold:

- (1) $\text{Mov } |K_S|$ is base point free;
- (2) $\text{Supp}(Z_\pi + E_\pi)$ is of simple normal crossing;
- (3) π factorizes through the modification $\mu : X' \rightarrow X$ in Setting 3.2.

We first consider the case when $|K_X|$ is not composed with a pencil of $(1, 2)$ -surfaces.

Proposition 9.2. *Let X be a minimal 3-fold of general type with $p_g(X) \geq 5$. Suppose that $|K_X|$ is not composed with a pencil of $(1, 2)$ -surfaces. Then*

$$K_X^3 \geq \frac{1}{4} \left[\frac{8}{3} (2p_g(X) - 5) \right].$$

Proof. Keep the notation in Setting 9.1.

If $\dim \Sigma = 3$, then by [22, Theorem 2.4], we have $K_X^3 \geq 2p_g(X) - 6$.

If $\dim \Sigma = 1$, since $|K_X|$ is not composed with a pencil of $(1, 2)$ -surfaces, then by [4, Theorem 4.4, Theorem 4.5], we have $K_X^3 > 2p_g(X) - 6$.

If $\dim \Sigma = 2$ and a general fiber C of ψ is of genus at least 3, then by [4, Theorem 4.1], we have $K_X^3 \geq 2p_g(X) - 4$.

From now on, suppose that $\dim \Sigma = 2$ and a general fiber C of ψ is of genus 2. Since $\dim \Sigma = 2$, we may write $S|_S \equiv dC$, where $d = \deg(s) \deg(\Sigma) \geq p_g(X) - 2$. By the adjunction formula,

$$\begin{aligned} K_S &= (K_{X'} + S)|_S = (\pi^*(K_X) + S + E_\pi)|_S \\ &= 2S|_S + (E_\pi + Z_\pi)|_S \equiv 2dC + (E_\pi + Z_\pi)|_S. \end{aligned}$$

It is clear that $p_g(S) \geq p_g(X) - 1 \geq 3$. By [4, Proposition 2.9], $K_{S_0}^2 \geq \frac{8}{3}(2d - 1)$. Since $K_{S_0}^2$ is an integer, we deduce that

$$K_{S_0}^2 \geq \lceil \frac{8}{3}(2p_g(X) - 5) \rceil.$$

By [4, Corollary 2.3] (take $D = K_X$ and $\lambda = 1$), $\pi^*(K_X)|_S - \frac{1}{2}\sigma_S^*(K_{S_0})$ is \mathbb{Q} -effective. Hence

$$K_X^3 \geq (\pi^*(K_X)|_S)^2 \geq \frac{1}{4}K_{S_0}^2 \geq \frac{1}{4} \lceil \frac{8}{3}(2p_g(X) - 5) \rceil.$$

The proof is completed. \square

From now on, we consider the case that $|K_X|$ is composed with a pencil of $(1, 2)$ -surfaces.

Proposition 9.3. *Let X be a minimal 3-fold of general type with $p_g(X) \geq 5$. Suppose that $|K_X|$ is composed with a pencil which is not rational. Then $K_X^3 \geq p_g(X)$.*

Proof. Keep the notation in Setting 9.1. Then we may write

$$\pi^*(K_X) \equiv dS + Z_\pi,$$

where $d \geq p_g(X)$ as the pencil is not rational. Then by [3, Lemma 2.1(i)],

$$(\pi^*(K_X)|_S)^2 = \text{Vol}(S) \geq 1.$$

Then

$$K_X^3 \geq d(\pi^*(K_X)|_S)^2 \geq p_g(X).$$

The proof is completed. \square

Proposition 9.4. *Let X be a minimal 3-fold of general type with $p_g(X) \geq 5$. Suppose that $|K_X|$ is composed with a rational pencil of $(1, 2)$ -surfaces. Then*

$$K_X^3 \geq \frac{p_g(X) - 1}{m + 1} \left(\frac{2p_g(X) - 2}{p_g(X)} + m - 1 \right),$$

where $m = \lfloor \frac{7p_g(X) - 11}{10} \rfloor$.

Proof. Keep the notation in Setting 9.1. We may write

$$\pi^*(K_X) = (p_g - 1)S + Z_\pi.$$

Here we simply write $p_g = p_g(X)$.

Take m to be the maximal integer such that

$$\left(m - \frac{1}{p_g - 1}\right) \cdot \frac{p_g - 1}{p_g} > m - 1 + \frac{3}{10},$$

or equivalently, $m = \lfloor \frac{7p_g(X) - 11}{10} \rfloor$. By the Kawamata–Viehweg vanishing theorem,

$$H^1(W, K_W + \lceil m\pi^*(K_X) - \frac{1}{p_g - 1}Z_\pi - S \rceil) = 0,$$

which implies that the natural restriction map

$$\begin{aligned} & H^0(W, K_W + \lceil m\pi^*(K_X) - \frac{1}{p_g - 1}Z_\pi \rceil) \\ & \rightarrow H^0(S, K_S + \lceil m\pi^*(K_X)|_S - \frac{1}{p_g - 1}Z_\pi|_S \rceil) \end{aligned}$$

is surjective. Here in the last term, the restriction on S commutes with the roundup as S is general. Therefore,

$$|(m + 1)K_W|_S \succcurlyeq |K_S + \lceil L \rceil + (m - 1)\sigma_S^*(K_{S_0})|, \quad (9.1)$$

where

$$L = m\pi^*(K_X)|_S - \frac{1}{p_g - 1}Z_\pi|_S - (m - 1)\sigma_S^*(K_{S_0}).$$

Recall that by our construction, π factorizes through μ in Setting 3.2. Hence by Lemma 3.3,

$$\begin{aligned} L & \geq \left(m - \frac{1}{p_g - 1}\right) \pi^*(K_X)|_S - (m - 1)\sigma_S^*(K_{S_0}) \\ & \geq \left(m - \frac{1}{p_g - 1}\right) \frac{p_g - 1}{p_g} \sigma_S^*(K_{S_0}) - (m - 1)\sigma_S^*(K_{S_0}) > \frac{3}{10} \sigma_S^*(K_{S_0}). \end{aligned}$$

Recall that S is a $(1, 2)$ -surface by assumption. By [2, Theorem 3.2], $h^0(S, K_S + \lceil L \rceil) \geq 3$.

Denote $|H| = \text{Mov } |K_S + \lceil L \rceil|$, then by comparing the movable parts of both sides of (9.1), we have

$$\pi^*((m + 1)K_X)|_S \geq H + (m - 1)C, \quad (9.2)$$

where $C \in \text{Mov } |K_S| = \text{Mov } |\sigma_S^*(K_{S_0})|$ is a general element. Recall that by [7, §3.7],

$$(\pi^*(K_X) \cdot C) = \xi \geq 1. \quad (9.3)$$

If $|H|$ is composed with a pencil, then it is composed with the same pencil as $|K_S|$ and hence $H \geq 2C$ as $h^0(S, H) \geq 3$. Then (9.2) implies that $\pi^*(K_X)|_S - C$ is \mathbb{Q} -effective. In this case, by (9.3),

$$K_X^3 \geq (p_g - 1)(\pi^*(K_X)|_S)^2 \geq (p_g - 1)(\pi^*(K_X) \cdot C) \geq p_g - 1.$$

If $|H|$ is not composed with a pencil, then $(H \cdot C) \geq 2$. On the other hand, by Lemma 3.3, $\pi^*(K_X)|_S - \frac{p_g-1}{p_g}C$ is \mathbb{Q} -effective. Hence $(\pi^*(K_X)|_S \cdot H) \geq \frac{2p_g-2}{p_g}$. Then, by (9.2) and (9.3),

$$\begin{aligned} (\pi^*(K_X)|_S)^2 &\geq \frac{1}{m+1}((\pi^*(K_X)|_S \cdot H) + (m-1)(\pi^*(K_X)|_S \cdot C)) \\ &\geq \frac{1}{m+1} \left(\frac{2p_g-2}{p_g} + m-1 \right). \end{aligned}$$

Then

$$K_X^3 \geq (p_g-1)(\pi^*(K_X)|_S)^2 \geq \frac{p_g-1}{m+1} \left(\frac{2p_g-2}{p_g} + m-1 \right).$$

The proof is completed. \square

By Propositions 9.2, 9.3, and 9.4, we have the following theorem.

Theorem 9.5. *Let X be a minimal 3-fold of general type with $p_g(X) = 5$. Then $K_X^3 \geq \frac{52}{15}$.*

Remark 9.6. (1) We remark that the lower bound in Theorem 9.5 might not be sharp. The minimal volume among known examples of minimal 3-folds of general type with $p_g = 5$ is $\frac{7}{2}$ in [9, Table 10, No. 2]. However, in this example, the canonical image is a surface, and this example justifies the optimality of Proposition 9.2. On the other hand, the minimal volume among known examples of minimal 3-folds of general type with $p_g = 5$ and with canonical system composed with a pencil is $\frac{109}{30}$ in [9, Table 10, No. 3].

(2) By Propositions 9.2, 9.3, and 9.4, we can also give lower bounds for minimal 3-folds of general type with $6 \leq p_g(X) \leq 10$. However, the lower bounds are not sufficient to conclude the expected Noether inequality, so we just collect the resulting lower bounds in the following table. Our lower bounds are in the second line while the lower bounds of the Noether inequality are in the third line.

$p_g(X)$	6	7	8	9	10
Our bound	55/12	39/7	133/20	208/27	87/10
Expected bound	14/3	6	22/3	26/3	10

Acknowledgments

This work was supported by National Key Research and Development Program of China #2023YFA1010600, #2020YFA0713200, and NSFC for Innovative Research Groups #12121001. Chen was supported by NSFC grant #12071078. Hu was supported by NSFC grant #12201397. Chen and Jiang are members of the Key Laboratory of Mathematics for Nonlinear Sciences, Fudan University.

We are grateful to Jungkai Chen, Stephen Coughlan, Roberto Pignatelli, Mingchen Xia and Tong Zhang for discussions.

REFERENCES

- [1] C. Birkar, P. Cascini, C. D. Hacon, J. McKernan, *Existence of minimal models for varieties of log general type*, J. Amer. Math. Soc. **23** (2010), no. 2, 405–468.
- [2] F. Catanese, M. Chen, D.-Q. Zhang, *The Noether inequality for smooth minimal 3-folds*, Math. Res. Lett. **13** (2006), no. 4, 653–666.
- [3] J. A. Chen, M. Chen, *Explicit birational geometry for 3-folds and 4-folds of general type, III*, Compos. Math. **151** (2015), no. 6, 1041–1082.
- [4] J. A. Chen, M. Chen, C. Jiang, *The Noether inequality for algebraic 3-folds*, with an appendix by J. Kollár, Duke Math. J. **169** (2020), no. 9, 1603–1645.
- [5] J. A. Chen, M. Chen, C. Jiang, *Addendum to “The Noether inequality for algebraic 3-folds”*, Duke Math. J. **169** (2020), no. 11, 2199–2204.
- [6] M. Chen, *Canonical stability of 3-folds of general type with $p_g \geq 3$* , Internat. J. Math. **14** (2003), no. 5, 515–528.
- [7] M. Chen, *A sharp lower bound for the canonical volume of 3-folds of general type*, Math. Ann. **337** (2007), no. 4, 887–908.
- [8] M. Chen, Y. Hu, M. Penegini, *On projective 3-folds of general type with small positive geometric genus*, Electron. Res. Arch. **29** (2021), no. 3, 2293–2323.
- [9] M. Chen, C. Jiang, B. Li, *On minimal varieties growing from quasismooth weighted hypersurfaces*, J. Differential Geom. **127** (2024), no. 1, 35–76.
- [10] M. Chen, Z. Jiang, *A reduction of canonical stability index of 4 and 5 dimensional projective varieties with large volume*, Ann. Inst. Fourier (Grenoble) **67** (2017), no. 5, 2043–2082.
- [11] D. A. Cox, *The homogeneous coordinate ring of a toric variety*, J. Algebraic Geom. **4** (1995), no. 1, 17–50.
- [12] D. A. Cox, *Erratum to “The homogeneous coordinate ring of a toric variety”*, J. Algebraic Geom. **23** (2014), no. 2, 393–398.
- [13] I. Dolgachev, *Weighted projective varieties*, In: Group actions and vector fields (Vancouver, B.C., 1981), pp. 34–71, Lecture Notes in Math., **956**, Springer, Berlin, 1982.
- [14] J. Han, J. Liu, V. V. Shokurov, *ACC for minimal log discrepancies of exceptional singularities*, arXiv:1903.04338v2.
- [15] E. Horikawa, *Algebraic surfaces of general type with small c_1^2 , I*, Ann. of Math. (2) **104** (1976), no. 2, 357–387.
- [16] E. Horikawa, *Algebraic surfaces of general type with small c_1^2 , II*, Invent. Math. **37** (1976), no. 2, 121–155.
- [17] Y. Hu, T. Zhang, *Algebraic threefolds of general type with small volume*, arXiv:2204.02222v1.
- [18] A. R. Iano-Fletcher, *Working with weighted complete intersections*, In: Explicit birational geometry of 3-folds, pp. 101–173, London Math. Soc. Lecture Note Ser., **281**, Cambridge University Press, Cambridge, 2000.
- [19] C. Jiang, *Characterizing terminal Fano threefolds with the smallest anti-canonical volume*, In: Birational geometry, Kähler–Einstein metrics and degenerations, pp. 355–362, Springer Proc. Math. Stat., **409**, Springer, Cham, 2023.
- [20] C. Jiang, *Characterizing terminal Fano threefolds with the smallest anti-canonical volume, II*, to appear in Higher dimensional algebraic geometry, A volume in honor of V. V. Shokurov, London Math. Soc. Lecture Note Ser., **489**, arXiv:2207.03832v1.
- [21] Y. Kawamata, K. Matsuda, K. Matsuki, *Introduction to the minimal model problem*, In: Algebraic geometry, Sendai, 1985, pp. 283–360, Adv. Stud. Pure Math., **10**, North-Holland, Amsterdam, 1987.
- [22] M. Kobayashi, *On Noether’s inequality for threefolds*, J. Math. Soc. Japan **44** (1992), no. 1, 145–156.
- [23] J. Kollár, *Shafarevich maps and automorphic forms*, M. B. Porter Lectures, Princeton University Press, Princeton, NJ, 1995.

- [24] J. Kollár, *Families of varieties of general type*, Cambridge Tracts in Math., **231**, Cambridge University Press, Cambridge, 2023.
- [25] J. Kollár, S. Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Math., **134**, Cambridge University Press, Cambridge, 1998.
- [26] J. Kollár, N. I. Shepherd-Barron, *Threefolds and deformations of surface singularities*, Invent. Math. **91** (1988), no. 2, 299–338.
- [27] H. Matsumura, *Commutative ring theory*, Cambridge Stud. Adv. Math., **8**, Cambridge University Press, Cambridge, 1986.
- [28] M. Nagata, *On rational surfaces. I. Irreducible curves of arithmetic genus 0 or 1*, Mem. Coll. Sci. Univ. Kyoto Ser. A. Math. **32** (1960), 351–370.
- [29] M. Reid, *Canonical 3-folds*, In: Journées de Géométrie Algébrique d’Angers, juillet 1979/Algebraic Geometry, Angers, 1979, pp. 273–310, Sijthoff & Noordhoff, Alphen aan den Rijn-Germantown, Md., 1980.
- [30] M. Reid, *Young person’s guide to canonical singularities*, In: Algebraic geometry, Bowdoin, 1985 (Brunswick, Maine, 1985), pp. 345–414, Proc. Sympos. Pure Math., **46**, Part 1, American Mathematical Society, Providence, RI, 1987.
- [31] M. Reid, *Constructing algebraic varieties via commutative algebra*, In: European Congress of Mathematics, pp. 655–667, European Mathematical Society (EMS), Zürich, 2005.
- [32] S. Takayama, *Local simple connectedness of resolutions of log-terminal singularities*, Internat. J. Math. **14** (2003), no. 8, 825–836.
- [33] E. Viehweg, *Quasi-projective moduli for polarized manifolds*, Ergeb. Math. Grenzgeb. (3), **30**, Springer-Verlag, Berlin, 1995.
- [34] O. Zariski, *The theorem of Riemann–Roch for high multiples of an effective divisor on an algebraic surface*, Ann. of Math. (2) **76** (1962), 560–615.

School of Mathematical Sciences & Shanghai Center for Mathematical Sciences, Fudan University, Shanghai 200433, China

Email address: `mchen@fudan.edu.cn`

School of Mathematical Sciences, Shanghai Jiao Tong University, Shanghai 200240, China

Email address: `yonghu@sjtu.edu.cn`

Shanghai Center for Mathematical Sciences & School of Mathematical Sciences, Fudan University, Shanghai 200438, China

Email address: `chenjiang@fudan.edu.cn`