

(k, a) -GENERALIZED FOURIER TRANSFORM WITH NEGATIVE a

TATSURO HIKAWA

*Dedicated to Professor Toshiyuki Kobayashi
with admiration for his profound and creative works*

ABSTRACT. The (k, a) -generalized Fourier transform $\mathcal{F}_{k,a}$ introduced by Ben Saïd–Kobayashi–Ørsted is a deformation family of the classical Fourier transform \mathcal{F} with the Dunkl parameter k and the parameter $a > 0$ that interpolates minimal representations of two different simple Lie groups. In the present paper, we focus on the case $a < 0$. As a main result, we find a unitary transform that intertwines the known case $a > 0$ and the new case $a < 0$.

CONTENTS

1. Introduction	1
1.1. Background	1
1.2. Result of the paper	2
1.3. Organization of the paper	3
1.4. Notation	3
2. Review of the theory of (k, a) -generalized Fourier transforms	3
2.1. Preliminaries on the Dunkl theory	3
2.2. \mathfrak{sl}_2 -triple of differential-difference operators	4
2.3. L^2 -theory for the \mathfrak{sl}_2 -triple	5
2.4. Lifting to a unitary representation $\Omega_{k,a}$	7
2.5. (k, a) -generalized Laguerre semigroup	8
2.6. (k, a) -generalized Fourier transform	8
3. (k, a) -generalized Fourier transform with negative a	9
3.1. A unitary transform that intertwines the cases $a > 0$ and $a < 0$	9
3.2. L^2 -theory for the \mathfrak{sl}_2 -triple with negative a	11
3.3. Lifting to a unitary representation $\Omega_{k,a}$ with negative a	12
3.4. (k, a) -generalized Laguerre semigroup with negative a	13
3.5. (k, a) -generalized Fourier transform with negative a	14
Acknowledgements	15
References	15

1. INTRODUCTION

1.1. **Background.** A minimal representation is an infinite-dimensional unitary representation with minimum Gelfand–Kirillov dimension. On the other hand, we can consider a minimal representation as a manifestation of “large symmetry” of the space acted by the group, and

hence it expected to control well global analysis on the space. This is the idea of “global analysis of minimal representations” initiated by T. Kobayashi [Kob2011], which caused a transition from algebraic representation theory to analytic representation theory, spawning a active research area that continues to the present day.

The Fourier transform in the classical harmonic analysis on \mathbb{R}^N can be interpreted as the unitary inversion operator appeared in the Schrödinger model of Weil representation, which is a unitary representation of the metaplectic group $Mp(N, \mathbb{R})$ on the Hilbert space $L^2(\mathbb{R}^N)$ (see [Fol1989] for more details), and decomposes into two minimal representations. This is a classical example of global analysis controlled well by a minimal representation.

Kobayashi–Ørsted [KØ2003a, KØ2003b, KØ2003c] analyzed the minimal representation of the indefinite orthogonal group $O(p, q)$ from various viewpoint, and in particular, they constructed a realization of the minimal representation on $L^2(C)$, or the Schrödinger model. Here, C is a $p+q-3$ -dimensional light cone in $\mathbb{R}^{p-1, q-1}$. Based on the idea of “global analysis of minimal representations”, Kobayashi–Mano [KM2005, KM2007a, KM2007b, KM2011] built a new theory of harmonic analysis by using the Schrödinger model $L^2(C)$ of the minimal representation of $O(p, q)$ instead of that of $Mp(N, \mathbb{R})$. In particular, they introduced the *Fourier transform on the light cone*, a counterpart of the classical Fourier transform. The special case $(p, q) = (N+1, 2)$, where the Hilbert space $L^2(C_+)$ (C_+ denotes the “future part” of the light cone) is isomorphic to $L^2(\mathbb{R}^N, |x|^{-1} dx)$, is studied in [KM2005, KM2007a].

After that, Ben Saïd–Kobayashi–Ørsted [BKØ2009, BKØ2012] introduced a family of \mathfrak{sl}_2 -triples

$$\mathbb{H}_{k,a}, \quad \mathbb{E}_{k,a}^+, \quad \mathbb{E}_{k,a}^-$$

of differential operators indexed by two parameters k and a (see Section 2.2), where k is a Dunkl parameter, and a is a deformation parameter. By using this \mathfrak{sl}_2 -triple, they introduced the (k, a) -generalized Laguerre semigroup $(\mathcal{F}_{k,a}(z))_{\operatorname{Re} z \geq 0}$ (see Section 2.5), and its special value, the (k, a) -generalized Fourier transform $\mathcal{F}_{k,a}$ (see Section 2.6). The (k, a) -generalized Fourier transform $\mathcal{F}_{k,a}$ includes some known transforms:

- the classical Fourier transform $\mathcal{F} = \mathcal{F}_{0,2}$,
- the Hankel transform $\mathcal{F}_{0,1}$, or the Fourier transform on the light cone when $(p, q) = (N+1, 2)$ [KM2007a]
- the Dunkl transform $\mathcal{D}_k = \mathcal{F}_{k,2}$ [Dun1992].

In particular, the parameter a interpolates continuously the minimal representation of two simple Lie groups $Mp(N, \mathbb{R})$ and $O(N+1, 2)$.

In recent years, the theory of the (k, a) -generalized Fourier transform has been actively researched from the perspectives of real analysis and probability theory, beyond the framework of representation theory.

1.2. Result of the paper. In the present paper, we find a unitary transform that intertwines the known case $a > 0$ and the new case $a < 0$ (Theorems 3.2 and 3.3). Although the \mathfrak{sl}_2 -triple $(\mathbb{H}_{k,a}, \mathbb{E}_{k,a}^+, \mathbb{E}_{k,a}^-)$ is defined for any $a \in \mathbb{C} \setminus \{0\}$, L^2 -theory is only available for $a > 0$ in the previous studies. This intertwining operator allows us to extend the results of [BKØ2009, BKØ2012] to the case $a < 0$. In particular, we naturally extend the definition of the (k, a) -generalized Laguerre transform and Fourier transform to the case $a < 0$ (Sections 3.4 and 3.5).

Note that our intertwining operator transfers a holomorphic discrete series representation to a holomorphic one, not an anti-holomorphic one.

1.3. Organization of the paper. In Section 2, we briefly review the theory of (k, a) -generalized Fourier transforms by [BKØ2009, BKØ2012]. This section contains no new results. In Section 3, we show a unitary transform that intertwines the cases $a > 0$ and $a < 0$, which is used to extend the results stated in Section 2 to the case $a < 0$.

1.4. Notation.

- $\mathbb{N} = \{0, 1, 2, \dots\}$, $\mathbb{R}_{>0} = \{x \in \mathbb{R} \mid x > 0\}$.
- We write $\langle -, - \rangle$ for the Euclidean inner product, and $|\cdot|$ for the Euclidean norm.
- Function spaces, such as C^∞ spaces and L^2 spaces, are understood to consist of complex-valued functions.

2. REVIEW OF THE THEORY OF (k, a) -GENERALIZED FOURIER TRANSFORMS

In this section, we briefly review the theory of (k, a) -generalized Fourier transform by [BKØ2009, BKØ2012]. This section contains no new results. Symbols generally follow from [BKØ2012], but for convenience, new or slightly modified symbols are used in a few places.

2.1. Preliminaries on the Dunkl theory. In this subsection, we explain some basic definitions and facts on the Dunkl theory [Dun1988, Dun1989] as in [BKØ2012, Section 2].

For $\alpha \in \mathbb{R}^N \setminus \{0\}$, we write r_α for the orthogonal reflection with respect to the hyperplane $(\mathbb{R}\alpha)^\perp$, namely,

$$r_\alpha(x) = x - \frac{2\langle \alpha, x \rangle}{|\alpha|^2} \alpha \quad (x \in \mathbb{R}^N).$$

Let \mathcal{R} be a reduced root system on \mathbb{R}^N in the sense that

- (R0) \mathcal{R} is a finite subset of $\mathbb{R}^N \setminus \{0\}$,
- (R1) $r_\alpha(\mathcal{R}) = \mathcal{R}$ for all $\alpha \in \mathcal{R}$, and
- (R2) $\mathcal{R} \cap \mathbb{R}\alpha = \{\alpha, -\alpha\}$ for all $\alpha \in \mathcal{R}$.

Note that we do not impose crystallographic conditions on roots and do not require that \mathcal{R} spans \mathbb{R}^N .

The subgroup \mathfrak{C} of $O(N)$ generated by all the reflections r_α is called the *finite Coxeter group associated with \mathcal{R}* . We say that a function $k: \mathcal{R} \rightarrow \mathbb{C}$ is a *multiplicity function* if it is constant on every \mathfrak{C} -orbit. We usually write k_α instead of $k(\alpha)$. We say that a multiplicity function k is *non-negative* if $k_\alpha \geq 0$ for all $\alpha \in \mathcal{R}$. The *index* of a multiplicity function k is defined as

$$\langle k \rangle = \sum_{\alpha \in \mathcal{R}^+} k_\alpha.$$

For a (not necessarily non-negative) multiplicity function k , the *Dunkl Laplacian* Δ_k (see [Dun1988] and [Dun1989, 1.1 Definiton]) is defined by

$$\Delta_k f(x) = \Delta f(x) + \sum_{\alpha \in \mathcal{R}^+} k_\alpha \left(\frac{2\langle \nabla f(x), \alpha \rangle}{\langle \alpha, x \rangle} - |\alpha|^2 \frac{f(x) - f(r_\alpha x)}{\langle \alpha, x \rangle^2} \right),$$

where $\Delta = \sum_{i=1}^N \left(\frac{\partial}{\partial x_i}\right)^2$ is the classical Laplacian and $\nabla = \left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_N}\right)$ is the gradient operator. When $k = 0$, Δ_k reduces to the classical Laplacian Δ .

Let $\mathcal{P}^m(\mathbb{R}^N)$ denote the space of homogeneous polynomials of degree m . We define the space of k -harmonic polynomials of degree m as

$$\mathcal{H}_k^m(\mathbb{R}^N) = \{p \in \mathcal{P}^m(\mathbb{R}^N) \mid \Delta_k p = 0\},$$

and the space of k -spherical harmonics of degree m as

$$\mathcal{H}_k^m(S^{N-1}) = \{p|_{S^{N-1}} \mid p \in \mathcal{H}_k^m(\mathbb{R}^N)\}.$$

These spaces are \mathbb{C} -stable. When $k = 0$, these are reduced to the space $\mathcal{H}^m(\mathbb{R}^N)$ of classical harmonic polynomials and the space $\mathcal{H}^m(S^{N-1})$ of classical spherical harmonics, respectively, which are $O(N)$ -stable.

We have the following Hilbert sum decomposition analogous to the classical case ($k = 0$).

Fact 2.1 ([Dun1988, pp. 37–39]). *For a non-negative multiplicity function k , we have the Hilbert sum decomposition*

$$L^2(S^{N-1}, w_k(\omega) d\omega) = \sum_{m \in \mathbb{N}}^{\oplus} \mathcal{H}_k^m(S^{N-1}),$$

where the weight function w_k with respect to the standard measure $d\omega$ on S^{N-1} is defined by

$$w_k(\omega) = \prod_{\alpha \in \mathcal{R}^+} |\langle \alpha, \omega \rangle|^{2k_\alpha} \quad (\omega \in S^{N-1}).$$

Henceforth, we fix a reduced root system \mathcal{R} on \mathbb{R}^N .

2.2. \mathfrak{sl}_2 -triple of differential-difference operators. For a (not necessarily non-negative) multiplicity function k and $a \in \mathbb{C} \setminus \{0\}$, the differential-difference operators on $\mathbb{R}^N \setminus \{0\}$ introduced by [BKØ2009, BKØ2012] are

$$\begin{aligned} \mathbb{H}_{k,a} &= \frac{N - 2 + 2\langle k \rangle + a}{a} + \frac{2}{a}E, \\ \mathbb{E}_{k,a}^+ &= \frac{i}{a}|x|^a, \\ \mathbb{E}_{k,a}^- &= \frac{i}{a}|x|^{2-a}\Delta_k, \end{aligned}$$

where $E = \sum_{i=1}^N x_i \frac{\partial}{\partial x_i}$ is the Euler operator on \mathbb{R}^N . They form an \mathfrak{sl}_2 -triple [BKØ2012, Theorem 3.2], and hence yield a Lie algebra representation

$$\omega_{k,a}: \mathfrak{sl}(2, \mathbb{R}) \rightarrow \text{End}_{\mathbb{C}}(C^\infty(\mathbb{R}^N \setminus \{0\})).$$

We again write $\omega_{k,a}$ for its complexification.

For the above parameters k , a and $m \in \mathbb{N}$, we consider the following differential-difference operators on $\mathbb{R}_{>0}$:

$$\begin{aligned} \mathbb{H}_{k,a}^{(m)} &= \frac{N - 2 + 2\langle k \rangle + a}{a} + \frac{2}{a}\vartheta, \\ \mathbb{E}_{k,a}^{+(m)} &= \frac{i}{a}r^a, \\ \mathbb{E}_{k,a}^{-(m)} &= \frac{i}{a}r^{-a}(\vartheta - m)(\vartheta + N - 2 + 2\langle k \rangle + m), \end{aligned}$$

where $\vartheta = r \frac{d}{dr}$ is the Euler operator on $\mathbb{R}_{>0}$. They are the radial parts of $\mathbb{H}_{k,a}$, $\mathbb{E}_{k,a}^+$, and $\mathbb{E}_{k,a}^-$ respectively in the following sense.

Proposition 2.2. *Let k be a (not necessarily non-negative) multiplicity function, $a \in \mathbb{C} \setminus \{0\}$, and $m \in \mathbb{N}$. For $p \in \mathcal{H}_k^m(S^{N-1})$ and $f \in C^\infty(\mathbb{R}_{>0})$, we have*

$$\mathbb{H}_{k,a}(p \otimes f) = p \otimes \mathbb{H}_{k,a}^{(m)} f,$$

$$\mathbb{E}_{k,a}^+(p \otimes f) = p \otimes \mathbb{E}_{k,a}^{+(m)} f,$$

$$\mathbb{E}_{k,a}^-(p \otimes f) = p \otimes \mathbb{E}_{k,a}^{-(m)} f,$$

where $p \otimes f$ denotes the function $r\omega \mapsto p(\omega)f(r)$ on $\mathbb{R}^N \setminus \{0\}$.

Proof. The first and second equations are obvious. The last equation follows from the second equation in [BKØ2012, Lemma 3.6]. \square

In particular, $\mathbb{H}_{k,a}^{(m)}$, $\mathbb{E}_{k,a}^{+(m)}$, and $\mathbb{E}_{k,a}^{-(m)}$ also form an \mathfrak{sl}_2 -triple, and hence yield a Lie algebra representation

$$\omega_{k,a}^{(m)} : \mathfrak{sl}(2, \mathbb{R}) \rightarrow \text{End}_{\mathbb{C}}(C^\infty(\mathbb{R}_{>0})).$$

We again write $\omega_{k,a}^{(m)}$ for its complexification.

2.3. L^2 -theory for the \mathfrak{sl}_2 -triple. For a non-negative multiplicity function k and $a \in \mathbb{R}$, we consider a weight function

$$w_{k,a}(x) = |x|^{a-2} \prod_{\alpha \in \mathcal{R}^+} |\langle \alpha, x \rangle|^{2k_\alpha} \quad (x \in \mathbb{R}^N \setminus \{0\}), \quad (1)$$

and the Hilbert space $L^2(\mathbb{R}^N, w_{k,a}(x) dx)$. By the polar decomposition $w_{k,a}(x) dx = w_k(\omega) d\omega \otimes r^{N-3+2\langle k \rangle + a} dr$ and Fact 2.1, we have the Hilbert sum decomposition

$$\begin{aligned} L^2(\mathbb{R}^N, w_{k,a}(x) dx) &= L^2(S^{N-1}, w_k(\omega) d\omega) \widehat{\otimes} L^2(\mathbb{R}_{>0}, r^{N-3+2\langle k \rangle + a} dr) \\ &= \sum_{m \in \mathbb{N}}^{\oplus} \mathcal{H}_k^m(S^{N-1}) \otimes L^2(\mathbb{R}_{>0}, r^{N-3+2\langle k \rangle + a} dr). \end{aligned} \quad (2)$$

Henceforth, for $\lambda \in \mathbb{C}$ and $l \in \mathbb{N}$, we write the Laguerre polynomial as

$$L_l^{(\lambda)}(t) = \sum_{j=0}^l \frac{(-1)^j}{j!} \binom{\lambda+l}{l-j} t^j = \sum_{j=0}^l \frac{(-1)^j}{j!(l-j)!} \frac{\Gamma(\lambda+l+1)}{\Gamma(\lambda+j+1)} t^j \quad (t \in \mathbb{R}_{>0}).$$

Proposition 2.3 ([BKØ2012, Proposition 3.15]). *Let k be a non-negative multiplicity function, $a > 0$, and $m \in \mathbb{N}$, and set*

$$\lambda_{k,a,m} = \frac{N-2+2\langle k \rangle+2m}{a}.$$

For $l \in \mathbb{N}$, we define a function $f_{k,a,m;l}$ on $\mathbb{R}_{>0}$ by

$$f_{k,a,m;l}(r) = \left(\frac{2^{\lambda_{k,a,m}+1} \Gamma(l+1)}{a^{\lambda_{k,a,m}} \Gamma(\lambda_{k,a,m}+l+1)} \right) r^m L_l^{(\lambda_{k,a,m})} \left(\frac{2}{a} r^a \right) \exp\left(-\frac{1}{a} r^a\right) \quad (r \in \mathbb{R}_{>0}).$$

If $\lambda_{k,a,m} > -1$, $(f_{k,a,m;l})_{l \in \mathbb{N}}$ is an orthonormal basis for $L^2(\mathbb{R}_{>0}, r^{N-3+2\langle k \rangle + a} dr)$. \square

Let

$$\mathbf{h} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \mathbf{e}^+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \mathbf{e}^- = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$

be the standard basis for $\mathfrak{sl}(2, \mathbb{R})$, and

$$\begin{aligned} \mathbf{k} &= \text{Ad} \left(\frac{1}{\sqrt{2}} \begin{pmatrix} -i & -1 \\ 1 & i \end{pmatrix} \right) \mathbf{h} = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \\ \mathbf{n}^+ &= \text{Ad} \left(\frac{1}{\sqrt{2}} \begin{pmatrix} -i & -1 \\ 1 & i \end{pmatrix} \right) \mathbf{e}^+ = \frac{1}{2} \begin{pmatrix} i & -1 \\ -1 & -i \end{pmatrix}, \\ \mathbf{n}^- &= \text{Ad} \left(\frac{1}{\sqrt{2}} \begin{pmatrix} -i & -1 \\ 1 & i \end{pmatrix} \right) \mathbf{e}^- = \frac{1}{2} \begin{pmatrix} -i & -1 \\ -1 & i \end{pmatrix} \end{aligned}$$

be the Cayley transforms of them.

Proposition 2.4. *Let k be a non-negative multiplicity function, $a > 0$, and $m \in \mathbb{N}$ with $\lambda_{k,a,m} = \frac{N-2+2\langle k \rangle + 2m}{a} > -1$. For $l \in \mathbb{N}$, we have*

$$\begin{aligned} \omega_{k,a}^{(m)}(\mathbf{k}) f_{k,a,m;l} &= (\lambda_{k,a,m} + 2l + 1) f_{k,a,m;l}, \\ \omega_{k,a}^{(m)}(\mathbf{n}^+) f_{k,a,m;l} &= i \sqrt{(l+1)(\lambda_{k,a,m} + l + 1)} f_{k,a,m;l+1}, \\ \omega_{k,a}^{(m)}(\mathbf{n}^-) f_{k,a,m;l} &= i \sqrt{l(\lambda_{k,a,m} + l)} f_{k,a,m;l-1}, \end{aligned}$$

where we regard $f_{k,a,m;-1} = 0$.

Proof. By Proposition 2.2, the assertion is essentially the same as [BKØ2012, Theorem 3.21]. \square

In particular, we determine the spectra of the differential-difference operators $\omega_{k,a}^{(m)}(\mathbf{k})$ and $\omega_{k,a}(\mathbf{k})$, which play a fundamental role for the definitions of (k, a) -generalized Laguerre semigroups and Fourier transforms (Sections 2.5 and 2.6).

Corollary 2.5. *Let k be a non-negative multiplicity function, $a > 0$, and $m \in \mathbb{N}$ with $\lambda_{k,a,m} = \frac{N-2+2\langle k \rangle + 2m}{a} > -1$. The differential-difference operator*

$$\omega_{k,a}^{(m)}(\mathbf{k}) = \frac{1}{a} (r^a - r^{-a} (\vartheta - m)(\vartheta + N - 2 + 2\langle k \rangle + m))$$

restricted to $W_{k,a}^{(m)} = \text{span}_{\mathbb{C}} \{f_{k,a,m;l} \mid l \in \mathbb{N}\}$ is essentially self-adjoint on $L^2(\mathbb{R}_{>0}, r^{N-3+2\langle k \rangle + a} dr)$ and diagonalized by the orthonormal basis $(f_{k,a,m;l})_{l \in \mathbb{N}}$. The discrete spectrum corresponding to $f_{k,a,m;l}$ is $\lambda_{k,a,m} + 2l + 1$.

Proof. It follows from Proposition 2.3 and Proposition 2.4. \square

Corollary 2.6 ([BKØ2012, Corollary 3.22]). *Let k be a non-negative multiplicity function and $a > 0$ with $\lambda_{k,a,0} = \frac{N-2+2\langle k \rangle}{a} > -1$. The differential-difference operator*

$$\omega_{k,a}(\mathbf{k}) = \frac{1}{a} (|x|^a - |x|^{2-a} \Delta_k)$$

restricted to $W_{k,a} = \bigoplus_{m \in \mathbb{N}} \mathcal{H}_k^m(S^{N-1}) \otimes W_{k,a}^{(m)}$ is essentially self-adjoint on $L^2(\mathbb{R}^N, w_{k,a}(x) dx)$, has no continuous spectra, and has the set of discrete spectra

$$\begin{cases} \{\lambda_{k,a,m} + 2l + 1 \mid m, l \in \mathbb{N}\} & (N \geq 2) \\ \{\frac{2\langle k \rangle \pm 1}{a} + 2l + 1 \mid l \in \mathbb{N}\} & (N = 1). \end{cases}$$

Proof. It follows from Corollary 2.5 combined with the Hilbert sum decomposition (2) and Proposition 2.2. (Note that $\mathcal{H}_k^m(S^0) \neq 0$ only for $m = 0, 1$ when $N = 1$.) \square

2.4. Lifting to a unitary representation $\Omega_{k,a}$. In this subsection, we write $\widetilde{SL}(2, \mathbb{R})$ for the universal covering Lie group of $SL(2, \mathbb{R})$, and K for the connected Lie subgroup of $SL(2, \mathbb{R})$ with Lie algebra $\mathfrak{so}(2) = \mathbb{R}i\mathbf{k}$.

To clarify the assertions below, we pin down some basic definitions and facts about $(\mathfrak{sl}(2, \mathbb{C}), K)$ -modules as in [BKØ2012]. Let (ϖ, V) be a $(\mathfrak{sl}(2, \mathbb{C}), K)$ -module. We say that $v \in V \setminus \{0\}$ is a *lowest weight vector of weight $\mu \in \mathbb{C}$* if

$$\varpi(\mathbf{k})v = \mu v \quad \text{and} \quad \varpi(\mathbf{n}^-)v = 0.$$

We say that (ϖ, V) is a *lowest weight module of weight μ* if V is generated by such v . It is known that, for each $\lambda \in \mathbb{C}$, there exists a unique irreducible lowest weight module of weight $\lambda + 1$ (up to isomorphism), for which we write $\pi_K(\lambda)$ [BKØ2012, p. 28]. Moreover, for $\lambda \geq -1$, $\pi_K(\lambda)$ lifts to a unique irreducible unitary representation $\pi(\lambda)$ of $\widetilde{SL}(2, \mathbb{R})$, which is infinite-dimensional except for the one-dimensional trivial representation $\pi(-1)$ [BKØ2012, Fact 3.27, 1), 2), 7)].

Theorem 2.7. *Let k be a non-negative multiplicity function, $a > 0$, and $m \in \mathbb{N}$ with $\lambda_{k,a,m} = \frac{N-2+2\langle k \rangle + 2m}{a} > -1$. We set*

$$W_{k,a}^{(m)} = \text{span}_{\mathbb{C}}\{f_{k,a,m;l} \mid l \in \mathbb{N}\}.$$

Then, $(\omega_{k,a}^{(m)}, W_{k,a}^{(m)})$ equips a natural $(\mathfrak{sl}(2, \mathbb{C}), K)$ -module structure, and is isomorphic to $\pi_K(\lambda_{k,a,m})$. Moreover, the $(\mathfrak{sl}(2, \mathbb{C}), K)$ -module $(\omega_{k,a}^{(m)}, W_{k,a}^{(m)})$ lifts to a unique unitary representation $\Omega_{k,a}^{(m)}$ of $\widetilde{SL}(2, \mathbb{R})$ on the Hilbert space $L^2(\mathbb{R}_{>0}, r^{N-3+2\langle k \rangle + a} dr)$, which is unitarily equivalent to $\pi(\lambda_{k,a,m})$.

Theorem 2.8. *Let k be a non-negative multiplicity function and $a > 0$ with $\lambda_{k,a,0} = \frac{N-2+2\langle k \rangle}{a} > -1$. We set*

$$W_{k,a} = \bigoplus_{m \in \mathbb{N}} \mathcal{H}_k^m(S^{N-1}) \otimes W_{k,a}^{(m)}.$$

Then, $(\omega_{k,a}, W_{k,a})$ equips a natural $\mathfrak{C} \times (\mathfrak{sl}(2, \mathbb{C}), K)$ -module structure, where \mathfrak{C} denotes the Coxeter group. Moreover, the $\mathfrak{C} \times (\mathfrak{sl}(2, \mathbb{C}), K)$ -module $(\omega_{k,a}, W_{k,a})$ lifts to a unique unitary representation $\Omega_{k,a}$ of $\mathfrak{C} \times \widetilde{SL}(2, \mathbb{R})$ on the Hilbert space $L^2(\mathbb{R}, w_{k,a}(x) dx)$, which decomposes as

$$L^2(\mathbb{R}^N, w_{k,a}(x) dx) = \sum_{m \in \mathbb{N}}^{\oplus} \mathcal{H}_k^m(S^{N-1}) \otimes L^2(\mathbb{R}_{>0}, r^{N-3+2\langle k \rangle + a} dr).$$

Here, \mathfrak{C} acts on $\mathcal{H}_k^m(S^{N-1})$, and $\widetilde{SL}(2, \mathbb{R})$ acts on $L^2(\mathbb{R}_{>0}, r^{N-3+2\langle k \rangle+a} dr)$ via the unitary representation $\Omega_{k,a}^{(m)}$.

Proof of Theorems 2.7 and 2.8. Theorem 2.8 is nothing but [BKØ2012, Theorems 3.28, 3.30, 3.31], in the proof of which Theorem 2.7 is essentially proven by using [BKØ2012, Fact 3.27, 7)]. \square

2.5. (k, a) -generalized Laguerre semigroup. Let k be a non-negative multiplicity function, $a > 0$, and $m \in \mathbb{N}$ with $\lambda_{k,a,m} = \frac{N-2+2\langle k \rangle+2m}{a} > -1$. For $z \in \mathbb{C}$, we define

$$\mathcal{J}_{k,a}^{(m)}(z) = \exp(-z\omega_{k,a}^{(m)}(\mathbf{k})) = \exp\left(\frac{z}{a}(r^{-a}(\vartheta - m)(\vartheta + N - 2 + 2\langle k \rangle + m) - r^a)\right).$$

By Corollary 2.5, each operator $\mathcal{J}_{k,a}^{(m)}(z)$ on $L^2(\mathbb{R}_{>0}, r^{N-3+2\langle k \rangle+a} dr)$ is diagonalized by the orthonormal basis $(f_{k,a,m;l})_{l \in \mathbb{N}}$ and satisfies

$$\mathcal{J}_{k,a}^{(m)}(z) f_{k,a,m;l} = e^{-z(\lambda_{k,a,m}+2l+1)} f_{k,a,m;l}. \quad (3)$$

In particular, $\mathcal{J}_{k,a}^{(m)}(z)$ is a Hilbert–Schmidt operator when $\operatorname{Re} z > 0$, and a unitary operator when $\operatorname{Re} z = 0$.

Now, let k be a non-negative multiplicity function and $a > 0$ with $\lambda_{k,a,0} = \frac{N-2+2\langle k \rangle}{a} > -1$. For $z \in \mathbb{C}$, we define

$$\mathcal{J}_{k,a}(z) = \exp(-z\omega_{k,a}(\mathbf{k})) = \exp\left(\frac{z}{a}(|x|^{2-a}\Delta_k - |x|^a)\right),$$

which decomposes as

$$\mathcal{J}_{k,a}(z) = \sum_{m \in \mathbb{N}}^{\oplus} \operatorname{id}_{\mathcal{H}_k^m(S^{N-1})} \otimes \mathcal{J}_{k,a}^{(m)}(z).$$

The operator $\mathcal{J}_{k,a}(z)$ is a Hilbert–Schmidt operator when $\operatorname{Re} z > 0$, and a unitary operator when $\operatorname{Re} z = 0$. The operator semigroup $(\mathcal{J}_{k,a}(z))_{\operatorname{Re} z \geq 0}$ is introduced and named the (k, a) -generalized Laguerre semigroup by [BKØ2012].

For more details on the (k, a) -generalized Laguerre semigroup, see [BKØ2012, Section 4].

2.6. (k, a) -generalized Fourier transform. Let k be a non-negative multiplicity function, $a > 0$, and $m \in \mathbb{N}$ with $\lambda_{k,a,m} = \frac{N-2+2\langle k \rangle+2m}{a} > -1$. We define a unitary operator $\mathcal{F}_{k,a}^{(m)}$ on $L^2(\mathbb{R}_{>0}, r^{N-3+2\langle k \rangle+a} dr)$ by

$$\begin{aligned} \mathcal{F}_{k,a}^{(m)} &= e^{\frac{i\pi}{2}(\lambda_{k,a,0}+1)} \mathcal{J}_{k,a}^{(m)}\left(\frac{i\pi}{2}\right) \\ &= e^{\frac{i\pi}{2}(\lambda_{k,a,0}+1)} \exp\left(\frac{i\pi}{2a}(r^{-a}(\vartheta - m)(\vartheta + N - 2 + 2\langle k \rangle + m) - r^a)\right). \end{aligned}$$

As a special case of Equation (3), $\mathcal{F}_{k,a}^{(m)}$ is diagonalized by the orthonormal basis $(f_{k,a,m;l})_{l \in \mathbb{N}}$ and satisfies

$$\mathcal{F}_{k,a}^{(m)} f_{k,a,m;l} = e^{-i\pi(\frac{m}{a}+l)} f_{k,a,m;l}.$$

Now, let k be a non-negative multiplicity function and $a > 0$ with $\lambda_{k,a,0} = \frac{N-2+2\langle k \rangle}{a} > -1$. We define a unitary operator $\mathcal{F}_{k,a}$ on $L^2(\mathbb{R}^N, w_{k,a}(x) dx)$ by

$$\mathcal{F}_{k,a} = e^{\frac{i\pi}{2}(\lambda_{k,a,0}+1)} \mathcal{J}_{k,a} \left(\frac{i\pi}{2} \right) = e^{\frac{i\pi}{2}(\lambda_{k,a,0}+1)} \exp \left(\frac{i\pi}{2a} (|x|^{2-a} \Delta_k - |x|^a) \right)$$

which decomposes as

$$\mathcal{F}_{k,a} = \sum_{m \in \mathbb{N}}^{\oplus} \text{id}_{\mathcal{H}_k^m(S^{N-1})} \otimes \mathcal{F}_{k,a}^{(m)}.$$

The unitary operator $\mathcal{F}_{k,a}$ is introduced and named the (k, a) -generalized Fourier transform by [BKØ2012].

For more details on the (k, a) -generalized Fourier transform, see [BKØ2012, Section 5].

3. (k, a) -GENERALIZED FOURIER TRANSFORM WITH NEGATIVE a

3.1. A unitary transform that intertwines the cases $a > 0$ and $a < 0$. For $\alpha \in \mathbb{R} \setminus \{0\}$ and $\beta \in \mathbb{R}$, we define a transform $\kappa_{\alpha,\beta}$ of functions on $\mathbb{R}^N \setminus \{0\}$ by

$$\kappa_{\alpha,\beta} F(x) = |x|^\beta F(|x|^{\alpha-1} x) \quad (x \in \mathbb{R}^N \setminus \{0\}).$$

We also write, for a function f on $\mathbb{R}_{>0}$,

$$\kappa_{\alpha,\beta} f(r) = r^\beta f(r^\alpha) \quad (r \in \mathbb{R}_{>0}).$$

Lemma 3.1. For $\alpha \in \mathbb{R} \setminus \{0\}$ and $\beta \in \mathbb{R}$, the transform $\kappa_{\alpha,\beta}$ of functions on $\mathbb{R}_{>0}$ satisfies the following properties.

- (1) The transforms $\kappa_{\alpha,\beta}$ yield an action of the one-dimensional affine group $\text{Aff}(1, \mathbb{R}) = \left\{ \begin{pmatrix} \alpha & 1 \\ 0 & \beta \end{pmatrix} \mid \alpha \in \mathbb{R} \setminus \{0\}, \beta \in \mathbb{R} \right\}$, namely, $\kappa_{\alpha,\beta} \circ \kappa_{\alpha',\beta'} = \kappa_{\alpha\alpha', \beta+\alpha\beta'}$.
- (2) $\vartheta \circ \kappa_{\alpha,\beta} = \kappa_{\alpha,\beta} \circ (\alpha\vartheta + \beta)$, where $\vartheta = r \frac{d}{dr}$ is the Euler operator on $\mathbb{R}_{>0}$.
- (3) For $d \in \mathbb{C}$, $r^{\alpha d} \circ \kappa_{\alpha,\beta} = \kappa_{\alpha,\beta} \circ r^d$, where r^d and $r^{\alpha d}$ denote the multiplication operators.
- (4) For $d \in \mathbb{R}$, the transform $\kappa_{\alpha,\beta}$ defines a unitary operator from $L^2(\mathbb{R}_{>0}, r^d dr)$ onto $L^2(\mathbb{R}_{>0}, |\alpha| r^{\alpha d + \alpha - 2\beta - 1} dr)$.

Proof. Confirmed by straightforward calculations. □

Recall from Section 2.3 that we write $(\mathbf{h}, \mathbf{e}^+, \mathbf{e}^-)$ for the standard basis for $\mathfrak{sl}(2, \mathbb{R})$. We define an automorphism τ of $\mathfrak{sl}(2, \mathbb{R})$ by

$$\tau(\mathbf{h}) = \mathbf{h}, \quad \tau(\mathbf{e}^+) = -\mathbf{e}^+, \quad \tau(\mathbf{e}^-) = -\mathbf{e}^-. \quad (4)$$

We again write τ for its complexification.

Here are our main theorems, which state that the unitary transform $\kappa_{-1, -(N-2+2\langle k \rangle)}$ intertwines the cases $a > 0$ and $a < 0$.

Theorem 3.2. Let k be a (not necessarily non-negative) multiplicity function, $a \in \mathbb{C} \setminus \{0\}$, and $m \in \mathbb{N}$. The transform $\kappa_{-1, -(N-2+2\langle k \rangle)}$ of functions on $\mathbb{R}_{>0}$ satisfies the following properties.

- (1) $\kappa_{-1, -(N-2+2\langle k \rangle)}^2 = \text{id}$.

(2) We have the following intertwining relations:

$$\begin{aligned}\kappa_{-1, -(N-2+2\langle k \rangle)} \circ \mathbb{H}_{k,a}^{(m)} &= \mathbb{H}_{k,-a}^{(m)} \circ \kappa_{-1, -(N-2+2\langle k \rangle)}, \\ \kappa_{-1, -(N-2+2\langle k \rangle)} \circ \mathbb{E}_{k,a}^{+(m)} &= -\mathbb{E}_{k,-a}^{+(m)} \circ \kappa_{-1, -(N-2+2\langle k \rangle)}, \\ \kappa_{-1, -(N-2+2\langle k \rangle)} \circ \mathbb{E}_{k,a}^{-(m)} &= -\mathbb{E}_{k,-a}^{-(m)} \circ \kappa_{-1, -(N-2+2\langle k \rangle)}.\end{aligned}$$

Thus, for $X \in \mathfrak{sl}(2, \mathbb{C})$, we have

$$\kappa_{-1, -(N-2+2\langle k \rangle)} \circ \omega_{k,a}^{(m)}(X) = \omega_{k,-a}^{(m)}(\tau(X)) \circ \kappa_{-1, -(N-2+2\langle k \rangle)}.$$

(3) If k is non-negative and $a \in \mathbb{R} \setminus \{0\}$, the transform $\kappa_{-1, -(N-2+2\langle k \rangle)}$ defines a unitary operator from $L^2(\mathbb{R}_{>0}, r^{N-3+2\langle k \rangle+a} dr)$ onto $L^2(\mathbb{R}_{>0}, r^{N-3+2\langle k \rangle-a} dr)$.

Proof. (1) It is a special case of Lemma 3.1 (1).

(2) By Lemma 3.1 (2) and (3),

$$\begin{aligned}\kappa_{-1, -(N-2+2\langle k \rangle)} \circ \vartheta &= (-\vartheta - (N - 2 + 2\langle k \rangle)) \circ \kappa_{-1, -(N-2+2\langle k \rangle)}, \\ \kappa_{-1, -(N-2+2\langle k \rangle)} \circ r^d &= r^{-d} \circ \kappa_{-1, -(N-2+2\langle k \rangle)}.\end{aligned}$$

Thus, we have

$$\begin{aligned}\kappa_{-1, -(N-2+2\langle k \rangle)} \circ \mathbb{H}_{k,a}^{(m)} &= \kappa_{-1, -(N-2+2\langle k \rangle)} \circ \left(\frac{N - 2 + 2\langle k \rangle + a}{a} + \frac{2}{a} \vartheta \right) \\ &= \left(\frac{-(N - 2 + 2\langle k \rangle) + a}{a} - \frac{2}{a} \vartheta \right) \circ \kappa_{-1, -(N-2+2\langle k \rangle)} \\ &= \mathbb{H}_{k,-a}^{(m)} \circ \kappa_{-1, -(N-2+2\langle k \rangle)}, \\ \kappa_{-1, -(N-2+2\langle k \rangle)} \circ \mathbb{E}_{k,a}^{+(m)} &= \kappa_{-1, -(N-2+2\langle k \rangle)} \circ \frac{i}{a} r^a \\ &= \frac{i}{a} r^{-a} \circ \kappa_{-1, -(N-2+2\langle k \rangle)} \\ &= -\mathbb{E}_{k,-a}^{+(m)} \circ \kappa_{-1, -(N-2+2\langle k \rangle)}, \\ \kappa_{-1, -(N-2+2\langle k \rangle)} \circ \mathbb{E}_{k,a}^{-(m)} &= \kappa_{-1, -(N-2+2\langle k \rangle)} \circ \frac{i}{a} r^{-a} (\vartheta - m)(\vartheta + N - 2 + 2\langle k \rangle + m) \\ &= \frac{i}{a} r^a (-\vartheta - (N - 2 + 2\langle k \rangle + m))(-\vartheta + m) \circ \kappa_{-1, -(N-2+2\langle k \rangle)} \\ &= -\mathbb{E}_{k,-a}^{-(m)} \circ \kappa_{-1, -(N-2+2\langle k \rangle)}.\end{aligned}$$

(3) It is a special case of Lemma 3.1 (4). \square

Theorem 3.3. Let k be a (not necessarily non-negative) multiplicity function and $a \in \mathbb{C} \setminus \{0\}$. The transform $\kappa_{-1, -(N-2+2\langle k \rangle)}$ of functions on $\mathbb{R}^N \setminus \{0\}$ satisfies the following properties.

- (1) $\kappa_{-1, -(N-2+2\langle k \rangle)}^2 = \text{id}$.
- (2) The transform $\kappa_{-1, -(N-2+2\langle k \rangle)}$ commutes the translations by the natural action of $O(N)$ (in particular, ones by the natural action of the Coxeter group \mathfrak{C}).

(3) The transform $\kappa_{-1, -(N-2+2\langle k \rangle)}$ satisfies the following intertwining relations:

$$\begin{aligned}\kappa_{-1, -(N-2+2\langle k \rangle)} \circ \mathbb{H}_{k,a} &= \mathbb{H}_{k,a} \circ \kappa_{-1, -(N-2+2\langle k \rangle)}, \\ \kappa_{-1, -(N-2+2\langle k \rangle)} \circ \mathbb{E}_{k,a}^+ &= -\mathbb{E}_{k,-a}^+ \circ \kappa_{-1, -(N-2+2\langle k \rangle)}, \\ \kappa_{-1, -(N-2+2\langle k \rangle)} \circ \mathbb{E}_{k,a}^- &= -\mathbb{E}_{k,-a}^- \circ \kappa_{-1, -(N-2+2\langle k \rangle)}.\end{aligned}$$

Thus, for $X \in \mathfrak{sl}(2, \mathbb{C})$, we have

$$\kappa_{-1, -(N-2+2\langle k \rangle)} \circ \omega_{k,a}(X) = \omega_{k,-a}(\tau(X)) \circ \kappa_{-1, -(N-2+2\langle k \rangle)}.$$

(4) If k is non-negative and $a \in \mathbb{R} \setminus \{0\}$, the transform $\kappa_{-1, -(N-2+2\langle k \rangle)}$ defines a unitary operator from $L^2(\mathbb{R}^N, w_{k,a}(x) dx)$ onto $L^2(\mathbb{R}_{>0}, w_{k,a}(x) dx)$.

Proof. The assertion (2) is obvious. Since $\mathbb{H}_{k,a}^{(m)}$, $\mathbb{E}_{k,a}^{+(m)}$, and $\mathbb{E}_{k,a}^{-(m)}$ are the radial parts of $\mathbb{H}_{k,a}$, $\mathbb{E}_{k,a}^+$, and $\mathbb{E}_{k,a}^-$ respectively (Proposition 2.2) and we have the Hilbert sum decomposition (2), the assertions (1), (3), and (4) follow from Theorem 3.2. \square

Remark 3.4. When $k = 0$, the transform in Theorem 3.3 is

$$\kappa_{-1, -(N-2)} F(x) = \frac{1}{|x|^{N-2}} F\left(\frac{x}{|x|^2}\right),$$

which is known as the Kelvin transform [Hel2009, p. 39].

3.2. L^2 -theory for the \mathfrak{sl}_2 -triple with negative a . Recall from Section 2.3 that we set

$$\lambda_{k,a,m} = \frac{N-2+2\langle k \rangle+2m}{a}.$$

Substituting a by $-a$, we get

$$\lambda_{k,-a,m} = \frac{N-2+2\langle k \rangle+2m}{-a} = -\lambda_{k,a,m}.$$

Proposition 3.5. Let k be a non-negative multiplicity function, $a > 0$, and $m \in \mathbb{N}$. For $l \in \mathbb{N}$, we define a function $f_{k,-a,m;l}$ on $\mathbb{R}_{>0}$ by

$$\begin{aligned}f_{k,-a,m;l}(r) &= \kappa_{-1, -(N-2+2\langle k \rangle)} f_{k,a,m;l}(r) \\ &= \left(\frac{2^{-\lambda_{k,-a,m}+1} \Gamma(l+1)}{a^{-\lambda_{k,-a,m}} \Gamma(-\lambda_{k,-a,m}+l+1)} \right) r^{-(N-2+2\langle k \rangle+m)} L_l^{(-\lambda_{k,-a,m})} \left(\frac{2}{a} r^{-a} \right) \exp\left(-\frac{1}{a} r^{-a}\right) \quad (r \in \mathbb{R}_{>0}).\end{aligned}$$

If $\lambda_{k,-a,m} < 1$, $(f_{k,-a,m;l})_{l \in \mathbb{N}}$ is an orthonormal basis for $L^2(\mathbb{R}_{>0}, r^{N-3+2\langle k \rangle-a} dr)$.

Proof. It follows from the fact that $(f_{k,a,m;l})_{l \in \mathbb{N}}$ is an orthonormal basis for $L^2(\mathbb{R}_{>0}, r^{N-3+2\langle k \rangle+a} dr)$ (Proposition 2.3) and that $\kappa_{-1, -(N-2+2\langle k \rangle)} : L^2(\mathbb{R}_{>0}, r^{N-3+2\langle k \rangle+a} dr) \rightarrow L^2(\mathbb{R}_{>0}, r^{N-3+2\langle k \rangle-a} dr)$ is a unitary operator (Theorem 3.2 (3)). \square

Recall from Section 2.3 that we write $(\mathbf{n}, \mathbf{k}^+, \mathbf{k}^-)$ for the Cayley transform of the standard basis $(\mathbf{h}, \mathbf{e}^+, \mathbf{e}^-)$ for $\mathfrak{sl}(2, \mathbb{R})$. Note that the automorphism τ defined by Equation (4) satisfies

$$\tau(\mathbf{k}) = -\mathbf{k}, \quad \tau(\mathbf{n}^+) = -\mathbf{n}^-, \quad \tau(\mathbf{n}^-) = -\mathbf{n}^+.$$

Proposition 3.6. *Let k be a non-negative multiplicity function, $a > 0$, and $m \in \mathbb{N}$ with $\lambda_{k,-a,m} = \frac{N-2+2\langle k \rangle + 2m}{-a} < 1$. For $l \in \mathbb{N}$, we have*

$$\begin{aligned}\omega_{k,-a}^{(m)}(\mathbf{k})f_{k,-a,m;l} &= (\lambda_{k,-a,m} - 2l - 1)f_{k,-a,m;l}, \\ \omega_{k,-a}^{(m)}(\mathbf{n}^+)f_{k,-a,m;l} &= -i\sqrt{l(-\lambda_{k,-a,m} + l)}f_{k,-a,m;l-1}, \\ \omega_{k,-a}^{(m)}(\mathbf{n}^-)f_{k,-a,m;l} &= -i\sqrt{(l+1)(-\lambda_{k,-a,m} + l + 1)}f_{k,-a,m;l+1},\end{aligned}$$

where we regard $f_{k,-a,m;-1} = 0$.

Proof. Apply $\kappa_{-1, -(N-2+2\langle k \rangle)}$ to the both sides of the equations in Proposition 2.4, and then use Theorem 3.2 (2). \square

In the same way as Corollaries 2.5 and 2.6, we obtain the following spectral properties.

Corollary 3.7. *Let k be a non-negative multiplicity function, $a > 0$, and $m \in \mathbb{N}$ with $\lambda_{k,-a,m} = \frac{N-2+2\langle k \rangle + 2m}{-a} < 1$. The differential-difference operator*

$$\omega_{k,-a}^{(m)}(\mathbf{k}) = -\frac{1}{a}(r^{-a} - r^a(\vartheta - m)(\vartheta + N - 2 + 2\langle k \rangle + m))$$

restricted to $W_{k,-a}^{(m)} = \text{span}_{\mathbb{C}}\{f_{k,-a,m;l} \mid l \in \mathbb{N}\}$ is essentially self-adjoint on $L^2(\mathbb{R}_{>0}, r^{N-3+2\langle k \rangle - a} dr)$ and diagonalized by the orthonormal basis $(f_{k,-a,m;l})_{l \in \mathbb{N}}$. The discrete spectrum corresponding to $f_{k,-a,m;l}$ is $\lambda_{k,-a,m} - 2l - 1$. \square

Corollary 3.8. *Let k be a non-negative multiplicity function and $a > 0$ with $\lambda_{k,-a,0} = \frac{N-2+2\langle k \rangle}{-a} < 1$. The differential-difference operator*

$$\omega_{k,-a}(\mathbf{k}) = \frac{1}{-a}(|x|^{-a} - |x|^{2+a}\Delta_k)$$

restricted to $W_{k,-a} = \bigoplus_{m \in \mathbb{N}} \mathcal{H}_k^m(S^{N-1}) \otimes W_{k,-a}^{(m)}$ is essentially self-adjoint on $L^2(\mathbb{R}^N, w_{k,-a}(x) dx)$, has no continuous spectra, and has the set of discrete spectra

$$\begin{cases} \{\lambda_{k,-a,m} - 2l - 1 \mid m, l \in \mathbb{N}\} & (N \geq 2) \\ \{\frac{2\langle k \rangle \pm 1}{-a} - 2l - 1 \mid l \in \mathbb{N}\} & (N = 1). \end{cases} \quad \square$$

3.3. Lifting to a unitary representation $\Omega_{k,a}$ with negative a . Recall from Section 2.4 that we write $\widetilde{SL}(2, \mathbb{R})$ for the universal covering Lie group of $SL(2, \mathbb{R})$, and K for the connected Lie subgroup of $SL(2, \mathbb{R})$ with Lie algebra $\mathfrak{so}(2) = \mathbb{R}i\mathbf{k}$.

In Section 2.4, we use a lowest weight $(\mathfrak{sl}(2, \mathbb{C}), K)$ -module $\pi_K(\lambda)$ of weight $\lambda + 1$, and its lifting $\pi(\lambda)$. Correspondingly, in this subsection, we use a highest weight $(\mathfrak{sl}(2, \mathbb{C}), K)$ -module and its lifting.

Let (ϖ, V) be a $(\mathfrak{sl}(2, \mathbb{C}), K)$ -module. We say that $v \in V \setminus \{0\}$ is a *highest weight vector of weight $\mu \in \mathbb{C}$* if

$$\varpi(\mathbf{k})v = \mu v \quad \text{and} \quad \varpi(\mathbf{n}^+)v = 0.$$

We say that (ϖ, V) is a *highest weight module of weight μ* if V is generated by such v . As with the case of lowest weight modules, for each $\lambda \in \mathbb{C}$, there exists a unique irreducible highest weight module of weight $\lambda - 1$ (up to isomorphism), for which we write $\pi'_K(\lambda)$. Moreover,

for $\lambda \leq 1$, $\pi'_K(\lambda)$ lifts to a unique irreducible unitary representation $\pi'(\lambda)$ of $\widetilde{SL}(2, \mathbb{R})$, which is infinite-dimensional except for the one-dimensional trivial representation $\pi'(1)$.

Remark 3.9. *With the automorphism τ of $\mathfrak{sl}(2, \mathbb{R})$ defined by (4) and its lifting $\tilde{\tau}$ to an automorphism of $\widetilde{SL}(2, \mathbb{R})$, we have*

$$\begin{aligned}\pi_K(\lambda) \circ \tau &= \pi'_K(-\lambda), \\ \pi(\lambda) \circ \tau &= \pi'(-\lambda).\end{aligned}$$

Theorem 3.10. *Let k be a non-negative multiplicity function, $a > 0$, and $m \in \mathbb{N}$ with $\lambda_{k,-a,m} = \frac{N-2+2\langle k \rangle + 2m}{-a} < 1$. We set*

$$W_{k,-a}^{(m)} = \text{span}_{\mathbb{C}}\{f_{k,-a,m;l} \mid l \in \mathbb{N}\}.$$

Then, $(\omega_{k,-a}^{(m)}, W_{k,-a}^{(m)})$ equips a natural $(\mathfrak{sl}(2, \mathbb{C}), K)$ -module structure, and is isomorphic to $\pi'_K(\lambda_{k,-a,m})$. Moreover, the $(\mathfrak{sl}(2, \mathbb{C}), K)$ -module $(\omega_{k,-a}^{(m)}, W_{k,-a}^{(m)})$ lifts to a unique unitary representation $\Omega_{k,-a}^{(m)}$ of $\widetilde{SL}(2, \mathbb{R})$ on the Hilbert space $L^2(\mathbb{R}_{>0}, r^{N-3+2\langle k \rangle - a} dr)$, which is unitarily equivalent to $\pi'(\lambda_{k,-a,m})$.

Proof. It follows from the result for positive a (Theorem 2.7), the intertwining property of $\kappa_{-1, -(N-2+2\langle k \rangle)}$ (Theorem 3.2), and Remark 3.9. \square

Theorem 3.11. *Let k be a non-negative multiplicity function and $a > 0$ with $\lambda_{k,-a,0} = \frac{N-2+2\langle k \rangle}{-a} < 1$. We set*

$$W_{k,-a} = \bigoplus_{m \in \mathbb{N}} \mathcal{H}_k^m(S^{N-1}) \otimes W_{k,-a}^{(m)}.$$

Then, $(\omega_{k,-a}, W_{k,-a})$ equips a natural $\mathfrak{C} \times (\mathfrak{sl}(2, \mathbb{C}), K)$ -module structure, where \mathfrak{C} denotes the Coxeter group. Moreover, the $\mathfrak{C} \times (\mathfrak{sl}(2, \mathbb{C}), K)$ -module $(\omega_{k,-a}, W_{k,-a})$ lifts to a unique unitary representation $\Omega_{k,-a}$ of $\mathfrak{C} \times \widetilde{SL}(2, \mathbb{R})$ on the Hilbert space $L^2(\mathbb{R} \setminus \{0\}, w_{k,-a}(x) dx)$, which decomposes as

$$L^2(\mathbb{R}^N, w_{k,-a}(x) dx) = \sum_{m \in \mathbb{N}}^{\oplus} \mathcal{H}_k^m(S^{N-1}) \otimes L^2(\mathbb{R}_{>0}, r^{N-3+2\langle k \rangle - a} dr).$$

Here, \mathfrak{C} acts on $\mathcal{H}_k^m(S^{N-1})$, and $\widetilde{SL}(2, \mathbb{R})$ acts on $L^2(\mathbb{R}_{>0}, r^{N-3+2\langle k \rangle - a} dr)$ via the unitary representation $\Omega_{k,-a}^{(m)}$.

Proof. It follows from the result for positive a (Theorem 2.8) and the intertwining property of $\kappa_{-1, -(N-2+2\langle k \rangle)}$ (Theorem 3.3). \square

3.4. (k, a) -generalized Laguerre semigroup with negative a . In this subsection, we extend the definition of the (k, a) -generalized Laguerre semigroup to the case $a < 0$.

Let k be a non-negative multiplicity function, $a > 0$, and $m \in \mathbb{N}$ with $\lambda_{k,-a,m} = \frac{N-2+2\langle k \rangle + 2m}{-a} < 1$. For $z \in \mathbb{C}$, we define

$$\mathcal{F}_{k,-a}^{(m)}(z) = \exp(-z\omega_{k,-a}^{(m)}(\mathbf{k})) = \exp\left(\frac{z}{-a}(r^a(\vartheta - m)(\vartheta + N - 2 + 2\langle k \rangle + m) - r^{-a})\right).$$

By Corollary 3.7, each operator $\mathcal{F}_{k,-a}^{(m)}(z)$ on $L^2(\mathbb{R}_{>0}, r^{N-3+2\langle k\rangle-a} dr)$ is diagonalized by the orthonormal basis $(f_{k,-a,m;l})_{l \in \mathbb{N}}$ and satisfies

$$\mathcal{F}_{k,-a}^{(m)}(z) f_{k,-a,m;l} = e^{-z(\lambda_{k,-a,m}-2l-1)} f_{k,-a,m;l}. \quad (5)$$

In particular, $\mathcal{F}_{k,-a}^{(m)}(z)$ is a Hilbert–Schmidt operator when $\operatorname{Re} z < 0$, and a unitary operator when $\operatorname{Re} z = 0$.

Now, let k be a non-negative multiplicity function and $a > 0$ with $\lambda_{k,-a,0} = \frac{N-2+2\langle k\rangle}{-a} < 1$. For $z \in \mathbb{C}$, we define

$$\mathcal{F}_{k,-a}(z) = \exp(-z\omega_{k,-a}(\mathbf{k})) = \exp\left(\frac{z}{-a}(|x|^{2+a}\Delta_k - |x|^{-a})\right),$$

which decomposes as

$$\mathcal{F}_{k,-a}(z) = \sum_{m \in \mathbb{N}}^{\oplus} \operatorname{id}_{\mathcal{H}_k^m(S^{N-1})} \otimes \mathcal{F}_{k,-a}^{(m)}(z).$$

The operator $\mathcal{F}_{k,-a}(z)$ is a Hilbert–Schmidt operator when $\operatorname{Re} z < 0$, and a unitary operator when $\operatorname{Re} z = 0$. We call the operator semigroup $(\mathcal{F}_{k,-a}(z))_{\operatorname{Re} z \geq 0}$ the $(k, -a)$ -generalized Laguerre semigroup.

Remark 3.12. By the intertwining property of $\kappa_{-1, -(N-2+2\langle k\rangle)}$ (Theorems 3.2 and 3.3), we have

$$\begin{aligned} \kappa_{-1, -(N-2+2\langle k\rangle)} \circ \mathcal{F}_{k,a}^{(m)}(z) &= \mathcal{F}_{k,-a}(-z) \circ \kappa_{-1, -(N-2+2\langle k\rangle)}, \\ \kappa_{-1, -(N-2+2\langle k\rangle)} \circ \mathcal{F}_{k,a}(z) &= \mathcal{F}_{k,-a}(-z) \circ \kappa_{-1, -(N-2+2\langle k\rangle)}. \end{aligned}$$

3.5. (k, a) -generalized Fourier transform with negative a . In this subsection, we extend the definition of the (k, a) -generalized Fourier transform to the case $a < 0$.

Let k be a non-negative multiplicity function, $a > 0$, and $m \in \mathbb{N}$ with $\lambda_{k,-a,m} = \frac{N-2+2\langle k\rangle+2m}{-a} < 1$. We define a unitary operator $\mathcal{F}_{k,-a}^{(m)}$ on $L^2(\mathbb{R}_{>0}, r^{N-3+2\langle k\rangle-a} dr)$ by

$$\begin{aligned} \mathcal{F}_{k,-a}^{(m)} &= e^{\frac{i\pi}{2}(\lambda_{k,-a,0}+1)} \mathcal{F}_{k,-a}^{(m)}\left(\frac{i\pi}{2}\right) \\ &= e^{\frac{i\pi}{2}(\lambda_{k,-a,0}+1)} \exp\left(\frac{i\pi}{-2a}(r^a(\vartheta - m)(\vartheta + N - 2 + 2\langle k\rangle + m) - r^{-a})\right). \end{aligned}$$

As a special case of Equation (5), $\mathcal{F}_{k,-a}^{(m)}$ is diagonalized by the orthonormal basis $(f_{k,-a,m;l})_{l \in \mathbb{N}}$ and satisfies

$$\mathcal{F}_{k,-a}^{(m)} f_{k,-a,m;l} = -e^{i\pi(\frac{m}{a}+l)} f_{k,-a,m;l}.$$

Now, let k be a non-negative multiplicity function and $a > 0$ with $\lambda_{k,-a,0} = \frac{N-2+2\langle k\rangle}{-a} < 1$. We define a unitary operator $\mathcal{F}_{k,-a}$ on $L^2(\mathbb{R}^N, w_{k,-a}(x) dx)$ by

$$\mathcal{F}_{k,-a} = e^{\frac{i\pi}{2}(\lambda_{k,-a,0}+1)} \mathcal{F}_{k,-a}\left(\frac{i\pi}{2}\right) = e^{\frac{i\pi}{2}(\lambda_{k,-a,0}+1)} \exp\left(\frac{i\pi}{-2a}(|x|^{2+a}\Delta_k - |x|^{-a})\right)$$

which decomposes as

$$\mathcal{F}_{k,-a} = \sum_{m \in \mathbb{N}}^{\oplus} \operatorname{id}_{\mathcal{H}_k^m(S^{N-1})} \otimes \mathcal{F}_{k,-a}^{(m)}.$$

We call the unitary operator $\mathcal{F}_{k,-a}$ the $(k, -a)$ -generalized Fourier transform.

Remark 3.13. As a special case of Remark 3.12, we have

$$\begin{aligned} \kappa_{-1,-(N-2+2\langle k \rangle)} \circ \mathcal{F}_{k,a}^{(m)} &= -(\mathcal{F}_{k,-a}^{(m)})^{-1} \circ \kappa_{-1,-(N-2+2\langle k \rangle)}, \\ \kappa_{-1,-(N-2+2\langle k \rangle)} \circ \mathcal{F}_{k,a} &= -\mathcal{F}_{k,-a}^{-1} \circ \kappa_{-1,-(N-2+2\langle k \rangle)}. \end{aligned}$$

ACKNOWLEDGEMENTS

The author is deeply grateful to his supervisor Professor Toshiyuki Kobayashi for his kind and patient guidance. His many insightful suggestions served as a guiding light for the author's research. It is an honor to dedicate the present paper to him. The author also thanks Kazuki Kannaka, whose comments led the author to consider (k, a) -generalized Fourier transform with negative a .

The author warmly thanks Professor Ali Baklouti, his colleagues, and Professor Hideyuki Ishi for their kindness during "7th Tunisian-Japanese Conference: Geometric and Harmonic Analysis on Homogeneous Spaces and Applications in Honor of Professor Toshiyuki Kobayashi".

This work was supported by World-leading Innovative Graduate Study for Frontiers of Mathematical Sciences and Physics (WINGS-FMSP).

REFERENCES

- [BKØ2009] S. Ben Saïd, T. Kobayashi, B. Ørsted, "Generalized Fourier transforms $\mathcal{F}_{k,a}$ ", *Comptes Rendus Mathématique* **347**.19–20 (2009), pp. 1119–1124.
DOI: [10.1016/j.crma.2009.07.015](https://doi.org/10.1016/j.crma.2009.07.015)
- [BKØ2012] S. Ben Saïd, T. Kobayashi, B. Ørsted, "Laguerre semigroup and Dunkl operators", *Compositio Mathematica* **148**.4 (2012), pp. 1265–1336.
DOI: [10.1112/S0010437X11007445](https://doi.org/10.1112/S0010437X11007445)
- [Dun1988] C. F. Dunkl, "Reflection groups and orthogonal polynomials on the sphere", *Mathematische Zeitschrift* **197** (1988), pp. 33–60.
DOI: [10.1007/BF01161629](https://doi.org/10.1007/BF01161629)
- [Dun1989] C. F. Dunkl, "Differential-difference operators associated to reflection groups", *Transactions of the American Mathematical Society* **311**.1 (1989), pp. 167–183.
DOI: [10.2307/2001022](https://doi.org/10.2307/2001022)
- [Dun1992] C. F. Dunkl, "Hankel Transforms Associated to Finite Reflection Groups", *Hypergeometric Functions on Domains of Positivity, Jack Polynomials, and Applications* American Mathematical Society, 1992, pp. 123–138.
DOI: [10.1090/conm/138/1199124](https://doi.org/10.1090/conm/138/1199124)
- [Fol1989] G. B. Folland, *Harmonic Analysis in Phase Space*, Princeton University Press, 1989.
- [Hel2009] L. L. Helms, *Potential Theory*, 2nd edition, Springer, 2009.
DOI: [10.1007/978-1-4471-6422-7](https://doi.org/10.1007/978-1-4471-6422-7)
- [Kob2011] T. Kobayashi, "Algebraic Analysis of Minimal Representations", *Publications of the Research Institute for Mathematical Sciences* **47**.2 (2011), pp. 585–611.
DOI: [10.2977/PRIMS/45](https://doi.org/10.2977/PRIMS/45)
- [KØ2003a] T. Kobayashi, B. Ørsted, "Analysis on the minimal representation of $O(p, q)$ I. Realization via conformal geometry", *Advances in Mathematics* **180**.2 (2003), pp. 486–512.
DOI: [10.1016/S0001-8708\(03\)00012-4](https://doi.org/10.1016/S0001-8708(03)00012-4)
- [KØ2003b] T. Kobayashi, B. Ørsted, "Analysis on the minimal representation of $O(p, q)$ II. Branching laws", *Advances in Mathematics* **180**.2 (2003), pp. 513–550.
DOI: [10.1016/S0001-8708\(03\)00013-6](https://doi.org/10.1016/S0001-8708(03)00013-6)

- [KØ2003c] T. Kobayashi, B. Ørsted, “Analysis on the minimal representation of $O(p, q)$ III. Ultrahyperbolic equations on $\mathbb{R}^{p-1, q-1}$ ”, *Advances in Mathematics* **180.2** (2003), pp. 513–550.
DOI: [10.1016/S0001-8708\(03\)00014-8](https://doi.org/10.1016/S0001-8708(03)00014-8)
- [KM2005] T. Kobayashi, G. Mano, “Integral Formulas for the Minimal Representation of $O(p, 2)$ ”, *Acta Applicandae Mathematicae* **86** (2005), pp. 103–113.
DOI: [10.1007/s10440-005-0464-2](https://doi.org/10.1007/s10440-005-0464-2)
- [KM2007a] T. Kobayashi, G. Mano, “The inversion formula and holomorphic extension of the minimal representation of the conformal group”, *Harmonic Analysis, Group Representations, Automorphic Forms and Invariant Theory: In honor of Roger E. Howe*, World Scientific, 2007, pp. 151–208.
DOI: [10.1142/9789812770790_0006](https://doi.org/10.1142/9789812770790_0006)
- [KM2007b] T. Kobayashi, G. Mano, “Integral formula of the unitary inversion operator for the minimal representation of $O(p, q)$ ”, *Proceedings of the Japan Academy, Series A, Mathematical Sciences* **83.3** (2007), pp. 27–31.
DOI: [10.3792/pjaa.83.27](https://doi.org/10.3792/pjaa.83.27)
- [KM2011] T. Kobayashi, G. Mano, *The Schrödinger Model for the Minimal Representation of the Indefinite Orthonormal Group $O(p, q)$* , American Mathematical Society, 2011.
DOI: [10.1090/s0065-9266-2011-00592-7](https://doi.org/10.1090/s0065-9266-2011-00592-7)

(Tatsuro Hikawa) GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, 3-8-1 KOMABA, MEGURO-KU, TOKYO, 153-8914, JAPAN