ON RANKIN-SELBERG INTEGRAL STRUCTURES AND EULER SYSTEMS FOR $GL_2 \times GL_2$.

Alexandros Groutides

Mathematics Institute, University of Warwick

ABSTRACT. We study how Rankin-Selberg periods interact with integral structures in spherical Whittaker type representations. Using this representation-theoretic framework, we show that the local Euler factors appearing in the construction of the motivic Rankin-Selberg Euler system for a product of modular forms are integrally optimal; i.e. any construction of this type with any choice of integral input data would give local factors appearing in tame norm relations at p which are integrally divisible by the Euler factor $\mathcal{P}'_p(\operatorname{Frob}_p^{-1})$ modulo p-1.

1. INTRODUCTION

1.1. Overview of the results. In [LSZ21b], Loeffler-Skinner-Zerbes introduce a new, local representation theoretic, approach to Euler system tame norm relations. In this paper, we adapt their techniques and further develop the underlying local *integral* theory of this approach for the Rankin-Selberg case of $G = \text{GL}_2 \times \text{GL}_2$. We show that *any* family of integral input data with determinant level in the sense of [LSZ21b], gives rise to cohomology classes which are norm-compatible in the strongest possible integral sense, i.e. as classes in motivic cohomology whose local factors appearing in tame norm relations at p are integral operator multiples of the Euler factor $\mathcal{P}'_p(\text{Frob}_p^{-1})$ modulo (p-1). It is important to note that our results hold for non-trivial coefficient sheaves as well. However, we avoid doing this here to ease exposition. We refer the reader to [LSZ21b], [LSZ21a] and [Gro20] for the formalisms required to extend this to non-trivial motives.

The integral structures with determinant level $G_p^{\circ}[p]$ introduced in [LSZ21b] arise from global cohomological considerations. They can be briefly described locally, through certain lattices $\mathcal{I}_0(G_p/G_p^{\circ}[p], \mathbb{Z}[1/p])$ inside the $\mathrm{GL}_2(\mathbb{Q}_p)$ -coinvariants of $\mathcal{S}_0(\mathbb{Q}_p^2) \otimes C_c^{\infty}(G_p)$, which are given by

$$\mathcal{I}_{0}(G_{p}/G_{p}^{\circ}[p],\mathbf{Z}[1/p]) := \operatorname{span}_{\mathbf{Z}[1/p]} \left\{ \phi \otimes \operatorname{ch}(gG_{p}^{\circ}[p]) \mid g \in G_{p}, \phi \text{ valued in } \frac{1}{\operatorname{vol}(\operatorname{Stab}(\phi) \cap gG_{p}^{\circ}[p]g^{-1})} \mathbf{Z}[1/p] \right\}.$$

Of course, the local setup of $S_0(\mathbf{Q}_p^2) \otimes C_c^{\infty}(G_p)$ is the natural one for our case and should resemble the classical $\operatorname{GL}_2 \times \operatorname{GL}_2$ Rankin-Selberg philosophy of "an Eisenstein series and two cusp forms".

Theorem A (Theorem 5.2.7, Corollary 5.2.9). Let $G = \operatorname{GL}_2 \times \operatorname{GL}_2$. Let S be a finite set of primes containing 2, and \mathscr{Z}_S the set of square-free positive integers coprime to S. For any family $\underline{\delta} = (\delta_p)_{p\notin S}$ of integral data with determinant level, that is either the "trivial" family $\underline{\delta}_0$ or a family in $\prod_{p\notin S} \mathcal{I}_0(G_p/G_p^\circ[p], \mathbf{Z}[1/p])$, there exists a collection of motivic cohomology classes

$$\Xi_{\mathrm{mot},n}(\underline{\delta}) \in H^3_{\mathrm{mot}}(Y_G(K_S) \times_{\mathbf{Q}} \mathbf{Q}(\mu_n), \mathbf{Q}(2)), \ n \in \mathscr{Z}_S$$

where $\Xi_{\text{mot},n}(\underline{\delta})$ depends on δ_p for p|n, that satisfy the following norm relations:

(1) For all $n, m \in \mathscr{Z}_S$ with $\frac{m}{n} = p$ prime

$$\operatorname{norm}_{\mathbf{Q}(\mu_n)}^{\mathbf{Q}(\mu_m)} (\Xi_{\operatorname{mot},m}(\underline{\delta})) = \mathcal{P}_{\operatorname{Tr}(\delta_p)}^{\operatorname{cycl}} \cdot \Xi_{\operatorname{mot},n}(\underline{\delta})$$

with

$$\mathcal{P}_{\mathrm{Tr}(\delta_p)}^{\mathrm{cycl}} \in \left(p-1, \mathcal{P}_p'(\mathrm{Frob}_p^{-1})\right) \subseteq \mathcal{H}_{G_p}^{\circ}(\mathbf{Z}[1/p])[\mathrm{Frob}_p^{\pm 1}]$$

The local factor $\mathcal{P}_{\mathrm{Tr}(\delta_p)}^{\mathrm{cycl}}$ is canonical and $\mathrm{Frob}_p \in \mathrm{Gal}(\mathbf{Q}(\mu_n)/\mathbf{Q})$ is arithmetic Frobenius at p. Additionally, $\mathcal{P}'_p(x) \in \mathcal{H}_{G_p}^{\circ}(\mathbf{Z}[1/p])[x]$ is the polynomial over the integral-away-from-p spherical Hecke algebra which interpolates local L-factors of unramified principal-series G_p -representations $\pi_{p,1} \boxtimes \pi_{p,2}$.

(2) Specialising $\underline{\delta}$, we have for all $n, m \in \mathscr{Z}_S$ with $\frac{m}{n} = p$ prime

$$\operatorname{norm}_{\mathbf{Q}(\mu_n)}^{\mathbf{Q}(\mu_m)}(\Xi_{\operatorname{mot},m}(\underline{\delta})) = \begin{cases} (p-1) \cdot \Xi_{\operatorname{mot},n}(\underline{\delta}) , & \text{if } \underline{\delta} = \underline{\delta}_0 \\ \mathcal{P}'_p(\operatorname{Frob}_p^{-1}) \cdot \Xi_{\operatorname{mot},n}(\underline{\delta}) , & \text{if } \underline{\delta} = \underline{\delta}_1 \end{cases}$$

We refer the reader to Definition 2.2.2 and Section 3.3 for more details and properties of the lattices \mathcal{I}_0 , and to Theorem 5.2.7 for the definition of the families $\underline{\delta}_0$ and $\underline{\delta}_1$. Part (2) of the theorem is a special instance of part (1) and it is the corresponding version of Theorem A in [LSZ21b], in our setup. The generality of part (1) shows that the motivic norm relations for $\underline{\delta}_1$ are optimal out of all possible choices of integral data, in the sense that the Euler factor appearing is integrally as "small" as possible without being trivial modulo p-1. Vice versa, part (2) shows that the ideal present in part (1) is optimal, in the sense that it cannot be shrunk any further, and any local factor appearing in norm relations for any integral input data is divisible integrally by the Euler factor $\mathcal{P}'_p(\operatorname{Frob}_p^{-1})$ modulo (p-1). This is also in line with the different approach of [LLZ14] and [LLZ18]: The classes $\Xi_{\mathrm{mot},n}(\underline{\delta}_1)$ are not exactly the same as the corresponding ones in *op.cit*, (e.g [LLZ18] 3.5.3, split case) which satisfy the analogous norm relations (Theorem A (2)) at p, only modulo p-1, which for applications is usually fine due to a result of [Rub00].

Additionally, fixing a prime ℓ and combining Theorem A with integral work of [LSZ21b], we see that for any family $\underline{\delta}$ as in Theorem A, one can obtain *integral* classes in étale cohomology $_{c}\Xi_{\acute{e}t,n}(\underline{\delta})$ (for appropriate integers c and n) for which the norm relations of Theorem A hold *integrally*. In particular for fixed c, and for n|m in an appropriate family of square-free integers, we have:

$$\operatorname{norm}_{\mathbf{Q}(\mu_n)}^{\mathbf{Q}(\mu_m)}(c\Xi_{\acute{e}t,m}(\underline{\delta})) = \left(\prod_{p\mid\underline{m}} \mathcal{P}'_p(\operatorname{Frob}_p^{-1})\right) \cdot c\Xi_{\acute{e}t,n}(\underline{\delta}) \quad \text{in} \quad H^3_{\acute{e}t}\left(Y_G(K_S) \times_{\mathbf{Q}} \mathbf{Q}(\mu_n), \mathbf{Z}_{\ell}(2)\right) / \ell$$

up to the action of $\prod_{p|\frac{m}{n}} \mathcal{H}^{\circ}_{G_p}(\overline{\mathbf{Z}}_{\ell})[\operatorname{Frob}_p^{\pm 1}]$ on the right. We refer the reader to Section 5.2.5 for more details.

Our techniques are representation theoretic. The local theory developed to show that the tame norm relations are integrally optimal, contains several novel stand-alone results from the viewpoint of representation theory of *p*-adic groups and period integrals. Our representation theoretic results hold for arbitrary finite extensions F/\mathbf{Q}_p , however we stick with \mathbf{Q}_p in order to avoid unnecessary notation shifts when we are dealing with applications to Euler systems or modular forms.

Our Hecke module approach to norm relations and the integral structures arising from global considerations in cohomology, have deep connections to the integral behaviour of the Rankin-Selberg periods of [JS81] and [JPSS83], attached to a product of modular forms.

Theorem B (Theorem 6.2.1, Theorem 6.4.5). Let $G = \operatorname{GL}_2 \times \operatorname{GL}_2$ and $H = \operatorname{GL}_2 \subseteq G$ diagonally. Let ℓ be a fixed prime and fix $\mathbf{C} \simeq_{\iota} \overline{\mathbf{Q}}_{\ell}$. Let $f_1, f_2 \in S_k(\Gamma_1(N)), k \geq 2$, be normalized cuspidal Hecke eigenforms. Write ϵ_i for their respective Nebentype, and π_{f_i} for their associated automorphic representations. Let \mathbf{L}_{f_1,f_2} be the smallest ℓ -adic number field containing the composite of the number fields of f_1 and f_2 under ι , and let S be the set of primes $\{\infty, p | \ell N\}$. Let $\varphi_i \simeq \otimes_p W_{i,p} \in \pi_{f_i}$ be decomposable cusp forms with $W_{i,p} = W_{\pi_{f_i},p}^{\operatorname{sph}}$ for $p \notin S$, $g = (g_1, g_2) \in G(\mathbf{A}^{\infty})$ and $\Phi = \otimes_p \Phi_p \in \mathcal{S}(\mathbf{A}^2)$ a decomposable Schwartz function. Finally, write

$$\varepsilon_{\ell,f_1,f_2}(\Phi,g_1,g_2) := \lim_{s \to 0} \frac{I(s,\Phi,g_1\varphi_1,g_2\varphi_2)}{L^S(\pi_{f_1} \boxtimes \pi_{f_2},s) \cdot \prod_{p \in S} Z(\Phi_p,g_{1,p}W_{p,1},g_{2,p}W_{p,2},s)} \in \overline{\mathbf{Q}}_{\ell}.$$

The following are true:

(1) If Φ_p is valued in $\operatorname{vol}_{H_p}(\operatorname{Stab}_{H_p}(\Phi_p) \cap g_p G_p^{\circ} g_p^{-1})^{-1} \mathcal{O}_{\mathbf{L}_{f_1, f_2}}$ for $p \notin S$, then

$$\varepsilon_{\ell,f_1,f_2}(\Phi,g_1,g_2) \in \mathcal{O}_{\mathbf{L}_{f_1,f_2}}$$

(2) Suppose that the assumption of part (1) holds and let S_0 be a finite set of places away from S for which: $p \equiv 1 \mod \ell$, $(\epsilon_1 \epsilon_2)(p) \neq 1$ and Φ_p is valued in $\operatorname{vol}_{H_p}(\operatorname{Stab}_{H_p}(\Phi_p) \cap g_p G_p^\circ[p] g_p^{-1})^{-1} \mathcal{O}_{\mathbf{L}_{f_1, f_2}}$ (note the addition of the determinant level $G_p^\circ[p]$ instead of G_p°). Then,

$$\left(\prod_{\substack{p \in S_0 \text{ s.t.} \\ \Phi_p(0,0) \neq 0}} L(\epsilon_1 \epsilon_2, 0)^{-1}\right) \cdot \varepsilon_{\ell, f_1, f_2}(\Phi, g_1, g_2) \in \prod_{p \in S_0} \left(p - 1, L_p^S(\pi_{f_1} \boxtimes \pi_{f_2}, 0)^{-1}\right) \subseteq \mathcal{O}_{\mathbf{L}_{f_1, f_2}}$$

In particular, if the level of f_1, f_2 satisfies $(\#(\mathbf{Z}/N\mathbf{Z})^{\times}, \ell) = 1$, then the containment holds without the bracketed product on the left.

Here I denotes the global Rankin-Selberg integral of [JS81], Z denotes the local Rankin-Selberg integral of [JPSS83] and [Jac90], L^S denotes the partial L-function, L_p^S denotes its local p-component and vol_{H_p} denotes the normalized Haar measure on $H_p = \operatorname{GL}_2(\mathbf{Q}_p)$ which gives $\operatorname{GL}_2(\mathbf{Z}_p)$ volume 1.

It is important to note that Theorem B (and more generally Section 6.2, Section 6.3, Section 6.4) treats *arbitrary* integral test vectors as per Definition 2.2.2. There are related results in the literature, mostly resembling Section 6.2, but only for *specific* nice integral input data (see for example [Gro20] Section 4). Overall, it does not seem feasible to directly study these periods for arbitrary integral test vectors and expect these general integrality results. This is mostly due to the lack of compatibility between the additive structure of the local space of Schwartz functions, and the normalisation volume factors of integral data arising from global considerations. The way we work around this issue is by translating our problem to an integral problem in $\operatorname{Ind}_{P_p}^{G_p} \mathbf{1}$ where P is the mirabolic subgroup of GL₂. This is also crucial in proving Theorem A since in order to realize the local factors for general integral input data and study them, one essentially needs to closely examine how local Rankin-Selberg periods interact with integral structures in unramified principal-series representations of GL₂ × GL₂.

The Rankin-Selberg case which we study here, has lots of similarities with the Asai case ([LLZ18], [Gro20]). As such, we expect that with some work, one could formulate and prove in the same manner the corresponding integral results in the Asai case by also attacking the local problem at non-split primes. Of course our setup is also related to the more technical GU(2, 1) case of [LSZ21b] and we expect similar optimal integrality results using the tools introduced here if one is keen to work through the calculations and technicalities. Finally, we believe that the interplay between local representation theory, period integrals and Euler systems will be fruitful in many other contexts.

1.2. Outline of the paper. After setting up some preliminaries in Section 2; Section 3 and Section 4 are mainly dedicated to reproducing the original local construction of [LSZ21b] and then "embedding it" into a different setup. This transition captures and realizes integrality in a more natural and approachable manner. It also bypasses the problem mentioned above regarding the local Rankin-Selberg zeta integral evaluated on arbitrary integral elements. There are a few main ingredients to this: Firstly, we verify that a general cyclicity result of [Sak08] can be applied to our setup, and proceed to link the two as in [LSZ21b]. Then, we construct and study a Hecke equivariant morphism to a certain space of compactly supported functions that are invariant under left translations by the diagonal mirabolic subgroup $P \subseteq GL_2$. Afterwards, using a splitting of measures inspired by [KM15], together with certain local zeta integrals, we construct nonzero Hecke equivariant forms from this space to $(\Pi_n^{\vee})^{G_p^{\circ}}$ for a sufficiently dense family of Whittaker type principal-series Π_n . These are easier to work with at an integral level and with general input data. Finally, we prove a main integrality result (Theorem 4.4.5) within this *P*-invariant setting. We also establish a (integral) structure result for the aforementioned space which even though is not crucial for norm relations it offers a more complete version of the theory and it links our work with integral structures in unramified Whittaker type G-representations distinguished by the diagonal mirabolic, and with branching laws for ℓ -modular representations (Section 6.1).

In Section 5.1, by "increasing the level" in our previous constructions and using certain trace maps, we transfer our integral results back to the original Rankin-Selberg setting. This is how we obtain the *local abstract integral analogue* of Theorem A (Theorem 5.1.1). The rest of Section 5 is dedicated to introducing the setup of Theorem A and defining the Rankin-Eisenstein map \mathcal{RE} , following [LSZ21b]. The next step is to apply our integral local result to this sophisticated example. Doing this, we obtain the integrally optimal norm-compatibility of our classes in motivic cohomology attached to general integral data.

In the final section (Section 6), we link our local Hecke theoretic results to the integral behavior of local Rankin-Selberg type periods which generate certain one-dimensional Hom-spaces. We finally link this to the global Rankin-Selberg integral of [JS81] for a product of modular forms, which culminates in Theorem B.

1.3. Acknowledgments. I am vastly grateful to my PhD supervisor David Loeffler, without whom this paper would not have been possible. In particular, I thank him for his constant support and invaluable mathematical input while this work was taking place. Additionally, I would like to thank Ju-Feng Wu, Robert Kurinczuk, Dipendra Prasad, Yiannis Sakellaridis and Johannes Girsch for useful discussions.

This research was funded by the ERC (Grant No. 101001051 — Shimura varieties and the Birch–Swinnerton-Dyer conjecture)

Contents

2. Preliminaries

2.1. Group schemes and Hecke algebras. Throughout, p will always denote a prime number. We fix once and for all the algebraic groups

$$G := \operatorname{GL}_2 \times \operatorname{GL}_2, \ H := \operatorname{GL}_2.$$

We identify H with a subgroup of G via the diagonal embedding $h \mapsto (h, h)$. For a group Γ in $\{H, G\}$, we write Γ_p for the locally pro-finite group $\Gamma(\mathbf{Q}_p)$ and Γ_p° for the maximal open compact subgroup $\Gamma(\mathbf{Z}_p)$. We fix a **Q**-valued Haar measure $d\gamma$ on Γ_p , normalized to give Γ_p° volume one. We write $\mathcal{H}(\Gamma_p) := C_c^{\infty}(\Gamma_p)$ for the full Hecke algebra of smooth, compactly supported **C**-valued functions on Γ_p , under the usual convolution,

which we denote by

$$(\theta_1 \cdot_{\Gamma} \theta_2)(x) := \int_{\Gamma_p} \theta_1(\gamma) \theta_2(\gamma^{-1}x) \ d\gamma, \ x \in \Gamma_p, \theta_1, \theta_2 \in \mathcal{H}(\Gamma_p)$$

The subscript Γ will be used when it's necessary to distinguish between different subgroups, but when it is clear from context, it will be omitted. We write $\mathcal{H}_{\Gamma_p}^{\circ} := C_c^{\infty}(\Gamma_p^{\circ} \setminus \Gamma_p/\Gamma_p^{\circ})$ for the spherical (or unramified) Hecke algebra of (smooth), compactly supported, Γ_p° -bi-invariant, **C**-valued functions on Γ_p , under the same convolution as before, and we also write $\mathcal{H}_{\Gamma_p}^{\circ}(\mathbf{Z}[1/p])$ for the subalgebra of functions valued in $\mathbf{Z}[1/p]$. It is well known that $\mathcal{H}_{\Gamma_p}^{\circ}$ is a commutative **C**-algebra of Weyl group invariant polynomials in the Satake parameters. Explicitly, we have:

$$\begin{aligned} \mathcal{H}_{H_p}^{\circ} \simeq \mathbf{C}[\mathcal{S}_p^{\pm 1}, \mathcal{T}_p] & \longleftrightarrow \mathcal{H}_{G_p}^{\circ} \simeq \mathbf{C}[\mathcal{S}_{p,1}^{\pm 1}, \mathcal{T}_{p,1}, \mathcal{S}_{p,2}^{\pm 1}, \mathcal{T}_{p,2}] \\ \mathcal{H}_{H_p}^{\circ}(\mathbf{Z}[1/p]) \simeq \mathbf{Z}[1/p][\mathcal{S}_p^{\pm 1}, \mathcal{T}_p] & \longleftrightarrow \mathcal{H}_{G_p}^{\circ}(\mathbf{Z}[1/p]) \simeq \mathbf{Z}[1/p][\mathcal{S}_{p,1}^{\pm 1}, \mathcal{T}_{p,1}, \mathcal{S}_{p,2}^{\pm 1}, \mathcal{T}_{p,2}] \\ \mathcal{S}_p & \longmapsto \mathcal{S}_{p,1}\mathcal{S}_{p,2} \\ \mathcal{T}_p & \longmapsto \mathcal{T}_{p,1}\mathcal{T}_{p,2} \end{aligned}$$

where

$$\begin{split} \mathcal{S}_p &:= \operatorname{ch}\left(\left[\begin{smallmatrix}p\\p\end{smallmatrix}\right] H_p^\circ\right) & \mathcal{T}_p &:= \operatorname{ch}\left(H_p^\circ\left[\begin{smallmatrix}p\\1\end{smallmatrix}\right] H_p^\circ\right) \\ \mathcal{S}_{p,1} &:= \operatorname{ch}\left(\left(\left[\begin{smallmatrix}p\\p\end{array}\right], 1\right) G_p^\circ\right) & \mathcal{T}_{p,1} &:= \operatorname{ch}\left(G_p^\circ\left(\left[\begin{smallmatrix}p\\1\end{smallmatrix}\right], 1\right) G_p^\circ\right) \\ \mathcal{S}_{p,2} &:= \operatorname{ch}\left(\left(1, \left[\begin{smallmatrix}p\\p\end{array}\right]\right) G_p^\circ\right) & \mathcal{T}_{p,1} &:= \operatorname{ch}\left(G_p^\circ\left(1, \left[\begin{smallmatrix}p\\1\end{smallmatrix}\right]\right) G_p^\circ\right) \end{split}$$

We will denote by (-)' the automorphism of $\mathcal{H}_{\Gamma_p}^{\circ}$ induced by inversion in Γ_p , i.e. for any $\theta \in \mathcal{H}_{\Gamma_p}^{\circ}$, we have $\theta'(\gamma) := \theta(\gamma^{-1})$ for all $\gamma \in \Gamma_p$. For an open subgroup U_p of Γ_p , we write $ch(U_p)$ for the characteristic function of U_p . All representations considered will be **C**-linear (or $\overline{\mathbf{Q}}_{\ell}$ -linear after fixing $\mathbf{C} \simeq \overline{\mathbf{Q}}_{\ell}$) unless otherwise stated (e.g Section 6.1), and every smooth Γ_p -representation will also be regarded as a $\mathcal{H}(\Gamma_p)$ -module in the usual way, via

$$\theta \cdot v := \int_{\Gamma_p} \theta(\gamma) \ \gamma \cdot v \ d\gamma, \ \theta \in \mathcal{H}(\Gamma_p), v \in V.$$

In particular, if $v \in V$ is U_p -invariant, then for any $g \in \Gamma_p$, $\operatorname{ch}(gU_p) \cdot v = \operatorname{vol}_{\Gamma_p}(U_p, d\gamma)g \cdot v$.

2.2. Whittaker models, equivariant maps and volume factors. We write "Ind" (resp. "c-Ind") for unnormalized (resp. compact unnormalized) induction, and "ind" for the usual normalized induction (e.g. [Cas80] Section 2). We will usually denote by B_p the upper triangular Borel of H_p , N_p its unipotent radical of upper unitriangular matrices, and by T_p the diagonal torus. Then $B_p = T_p \ltimes N_p$ and we also have the well-known Iwasawa decomposition $H_p = B_p H_p^{\circ}$. Let $\psi : \mathbf{Q}_p \to \mathbf{C}^{\times}$ be an additive character which we regard as a character of N_p in the usual way by identifying N_p with \mathbf{Q}_p through its top right entry. As in [Loe21], we say that a smooth H_p -representation π_p is of Whittaker type if it is either irreducible and generic or a twist of the reducible principle series, which contains the Steinberg representation as a co-dimension one submodule. By a classic vet deep theorem of Gel'fand-Kazhdan (in a more general setting, [GK72]) such a representation is isomorphic to a space of C-valued functions on H_p transforming by ψ under left translations by N_p , i.e. $\pi_p \hookrightarrow \operatorname{Ind}_{N_p}^{H_p} \psi$. This space is unique and we call it the Whittaker model of π_p , denoted by $\mathcal{W}(\pi_p, \psi)$. We will thus identify such a representation with its Whittaker model. Additionally, we say that such a representation is unramified if it has a non-zero H_p° -fixed vector. It is well-known that for such representation the space $(\pi_p)^{H_p^{\circ}}$ is one-dimensional and is generated by a canonical spherical vector $W_{\pi_p}^{\rm sph}$ such that $W_{\pi_p}^{\rm sph}(1) = 1$. Thus, the spherical Hecke algebra of H_p acts on this space via a character which we call the *spherical Hecke eigensystem* of π_p and denote it by Θ_{π_p} .

We write $S(\mathbf{Q}_p^2)$ for the space of **C**-valued Schwartz functions (i.e locally constant and compactly supported) on \mathbf{Q}_p^2 , regarded as a smooth H_p -representation under right translations. We also write $S_0(\mathbf{Q}_p^2)$ for the subrepresentation of $S(\mathbf{Q}_p^2)$ consisting of functions that vanish at (0, 0).

Definition 2.2.1 ([LSZ21b] 3.1.1). Let V be a smooth G_p -representation. A linear map

$$\mathfrak{Z}: \mathcal{S}(\mathbf{Q}_p^2) \otimes \mathcal{H}(G_p) \longrightarrow V$$

is $(G_p \times H_p)$ -equivariant, if it is equivariant with respect to the following actions:

- (1) G_p acts on the left hand side by $g \cdot (\phi \otimes \xi) := \phi \otimes (\xi((-)g))$ and on the right-hand side by its assigned action on V.
- (2) H_p acts on the left-hand side by $h \cdot (\phi \otimes \xi) := \phi((-)h) \otimes \xi(h^{-1}(-))$ and trivially on the right-hand side

The family of such equivariant maps can be canonically identified with $\operatorname{Hom}_{H_p}(\mathcal{S}(\mathbf{Q}_p^2) \otimes V^{\vee}, \mathbf{1})$, where V^{\vee} denotes the smooth dual of V ([LSZ21a] Section 4.9). The latter is often non-zero and very interesting to study. For an open subgroup $U_p \subseteq G_p^{\circ}$, we write $\mathcal{I}(G_p/U_p)$, for the H_p -coinvariants of $\mathcal{S}(\mathbf{Q}_p^2) \otimes C_c^{\infty}(G_p/U_p)$, and [·] for the class of an element in this space of coinvariants. We will eventually drop the [·] notation once it's no longer necessary. It follows from the definitions that for every such pair (V, \mathfrak{Z}) , we get an induced map $\mathcal{I}(G_p/U_p) \to V^{U_p}$. We equip $\mathcal{S}(\mathbf{Q}_p^2) \otimes C_c^{\infty}(G_p/G_p^{\circ})$ with the structure of an $\mathcal{H}_{G_p}^{\circ}$ -module via the G_p -action defined above. It can be easily checked that this induces a well-defined action of $\mathcal{H}_{G_p}^{\circ}$ on $\mathcal{I}(G_p/G_p^{\circ})$. A formal unravelling of definitions, gives the following two identities as in [LSZ21b]:

- Let $\phi \in \mathcal{S}(\mathbf{Q}_p^2)$ and $\xi_1, \xi_2 \in \mathcal{H}(G_p)$. Then $[\phi \otimes (\xi_1 \cdot_G \xi_2')] = \xi_2 \cdot [\phi \otimes \xi_1]$.
- Let $\phi \in \mathcal{S}(\mathbf{Q}_p^2), \xi_1 \in \mathcal{H}(H_p)$ and $\xi_2 \in \mathcal{H}(G_p)$. Then $[(\xi_1 \cdot \phi) \otimes \xi_2] = [\phi \otimes (\xi_1' \cdot H \xi_2)].$
- These two identities will be used throughout without further mention.

Definition 2.2.2. We introduce the following integral lattices in $\mathcal{I}(G_p/G_p^\circ)$ as in [LSZ21b] 3.2.1. Let U_p be an open subgroup of G_p° . The integral lattice (at level U_p) $\mathcal{I}_{(0)}(G_p/U_p, \mathbb{Z}[1/p])$ consists of all $\mathbb{Z}[1/p]$ -linear combinations of elements of the form $[\phi \otimes ch(gU_p)]$, where $\phi \in \mathcal{S}_{(0)}(\mathbb{Q}_p^2)$, $g \in G_p$, and ϕ is valued in

$$\frac{1}{\operatorname{vol}_{H_p}\left(\operatorname{Stab}_{H_p}(\phi) \cap gU_pg^{-1}\right)}\mathbf{Z}[1/p].$$

Here $S_{(0)}$ is shorthand notation to mean either S or S_0 and $\mathcal{I}_{(0)}$ is then defined accordingly. Additionally, vol_{H_p} denotes the normalized \mathbf{Q} -valued Haar measure which gives H_p° volume 1.

Remark 2.2.3. At first glance this definition seems a bit odd. However from a global viewpoint as pointed out in [LSZ21b], this is precisely the data which gives integral classes in the étale cohomology of our Shimura variety. As we will see later on, these elements do not only possess global integral properties, but also deep *local* integral properties.

Of particular interest to us, will be the lattices $\mathcal{I}_{(0)}(G_p/G_p^\circ, \mathbf{Z}[1/p])$ and $\mathcal{I}_{(0)}(G_p/G_p^\circ[p], \mathbf{Z}[1/p])$, where $G_p^\circ[p]$ is the open level subgroup of G_p° given by

$$G_p^{\circ}[p] := \{ (g_1, g_2) \in G_p^{\circ} \mid \det(g_2) \equiv 1 \mod p \}.$$

Remark 2.2.4. One might think that there is a certain asymmetry in this definition in the sense that we've imposed the determinant condition on the second factor. However, this choice is irrelevant. The results are identical if one imposes the condition on the first factor instead. In fact, the existence of this choice can be eliminated all together if one works with the larger group $G \times GL_1$ and then imposes this determinant condition on the GL_1 factor. We do not do that here to avoid unnecessarily complicating the notation everywhere else.

The inclusion $G_p^{\circ}[p] \subseteq G_p^{\circ}$ induces trace maps which descend to an integral level ([LSZ21b] Proposition 3.2.3):

$$\begin{array}{c}
\mathcal{I}_{(0)}(G_p/G_p^{\circ}[p]) \xrightarrow{\operatorname{Tr}_{G_p^{\circ}}^{G_p^{\circ}[p]}} \mathcal{I}_{(0)}(G_p/G_p^{\circ}) \\
\uparrow & \uparrow \\
\mathcal{I}_{(0)}(G_p/G_p^{\circ}[p], \mathbf{Z}[1/p]) \xrightarrow{\operatorname{Tr}_{G_p^{\circ}}^{G_p^{\circ}[p]}} \mathcal{I}_{(0)}(G_p/G_p^{\circ}, \mathbf{Z}[1/p])
\end{array}$$

where $\operatorname{Tr} := \operatorname{Tr}_{G_p^\circ}^{G_p^\circ[p]}(\phi \otimes \xi) := \phi \otimes \sum_{\gamma \in G_p^\circ/G_p^\circ[p]} \xi((-)\gamma).$

1

3. Structure theorems

3.1. Connection to spherical varieties and a result of Sakellaridis. We introduce the space of C-valued functions $C_c^{\infty}(H_p^{\circ}\backslash G_p/G_p^{\circ})$, which we regard as a module over $\mathcal{H}_{G_p}^{\circ} \otimes \mathcal{H}_{H_p}^{\circ}$ via

$$\left(\left(\theta_1\otimes\theta_2\right)\cdot\xi\right)(x):=\int_{G_p}\int_{H_p}\theta_1(g)\theta_2(h)f(h^{-1}xg)\ dg\ dh,\ \theta_1\otimes\theta_2\in\mathcal{H}^\circ_{G_p}\otimes\mathcal{H}^\circ_{H_p},\ f\in C^\infty_c(H^\circ_p\backslash G_p/G^\circ_p).$$

The goal of this section is to prove that the module $C_c^{\infty}(H_p^{\circ}\backslash G_p/G_p^{\circ})$ is cyclic over $\mathcal{H}_{G_p}^{\circ} \otimes \mathcal{H}_{H_p}^{\circ}$, and generated by the characteristic function of G_p° . To do this, we heavily rely on an application of a general theorem of Sakellaridis ([Sak13]) regarding function spaces on spherical varieties. Sakellaridis' result ([Sak13] Corollary 8.0.4) shows that for any split reductive group \mathscr{G} over \mathbf{Z}_p , and any homogeneous affine spherical \mathscr{G} -variety \mathscr{X} that satisfies a long list of conditions, the module $C_c^{\infty}(\mathscr{X}(\mathbf{Q}_p))^{\mathscr{G}_p^{\circ}}$, of \mathscr{G}_p° -invariant Schwartz functions on $\mathscr{X}(\mathbf{Q}_p)$, is cyclic over $\mathcal{H}_{\mathscr{G}_p}^{\circ}$, generated by the characteristic function of $\mathscr{X}(\mathbf{Z}_p)$. In fact, Sakellaridis' results also give us the annihilator of this action ([Sak13] Theorem 8.0.2), but we won't be concerned with that here.

Theorem 3.1.1. The module $C_c^{\infty}(H_p^{\circ}\backslash G_p/G_p^{\circ})$ is cyclic over $\mathcal{H}_{G_p}^{\circ} \otimes \mathcal{H}_{H_p}^{\circ}$, generated by the characteristic function $ch(G_p^{\circ})$.

Proof. In order to be able to apply Sakellaridis' result, we need some preparation. We write \mathscr{G} for the group scheme $G \times H = \operatorname{GL}_2^3$, and \mathscr{H} for the diagonal copy of H in \mathscr{G} . We define three different Borel subgroups of GL_2 . We write B_1 for the upper triangular Borel, B_2 for the lower triangular Borel, and B_3 for nB_2n^{-1} , where $n = \begin{bmatrix} 1 & 1 \\ 1 \end{bmatrix}$. Then $\mathscr{B} := B_1 \times B_2 \times B_3$ is a Borel of \mathscr{G} . With this choice, the \mathscr{B} -orbit of the identity is the unique open \mathscr{B} -orbit on the homogeneous affine variety $\mathcal{X} := \mathscr{H} \setminus \mathscr{G}$. We write \mathcal{Y} for this open orbit. We write T for the diagonal split torus of GL_2 , and $\mathscr{T} := T \times T \times nTn^{-1} \subseteq \mathscr{B}$. Write w_1, w_2, w_3 for the simple reflections associated to each copy of GL_2 in \mathscr{G} . The Weyl group W, of \mathscr{G} is given by $\langle w_1 \rangle \times \langle w_2 \rangle \times \langle w_3 \rangle \simeq (\mathbb{Z}/2\mathbb{Z})^3$. For each simple reflection we write \mathscr{P}_i for the associated parabolic with respect to our fixed choice of $\mathscr{T} \subseteq \mathscr{B}$ (e.g $\mathscr{P}_1 = \operatorname{GL}_2 \times B_2 \times B_3$). In [Kno95], Knopp defines an action of the Weyl group of \mathscr{G} on the set of Borel orbits (over $\overline{\mathbb{Q}}_p$) which is summarized in Section 2.2 of [Sak08]. We want to compute the action of each simple reflection on the open orbit \mathcal{Y} . We do this for i = 1 and the other two cases are handled in the same fashion. For this, we once again follow [Sak08] Section 2.2. Consider the quotient

$$\mathscr{P}_1 \to \mathrm{PGL}_2 = \mathrm{Aut}(\mathbf{P}_1) = \mathrm{Aut}(\mathscr{B} \setminus \mathscr{P}_1).$$

The image of $\operatorname{Stab}_{\mathscr{P}_1}(1)$ (where we regard 1 as an element of $\mathscr{H}\backslash\mathscr{G}$) in PGL₂ is a spherical subgroup. It is readily seen that this image is given by the non-trivial torus $B_2 \cap B_3$ in PGL₂. Thus it is of "type T" as per the classification in [Sak08] Section 2.1.1. From the description of Knopp's action, we deduce that w_1 fixes the open orbit \mathcal{Y} . Repeating this for w_2 and w_3 , we see that W fixes the open orbit \mathcal{Y} . Now, in our case, the standard parabolic $P(\mathcal{X}) := \{g \in \mathscr{G} \mid \mathcal{Y} \cdot g = \mathcal{Y}\}$ is simply equal to \mathscr{B} , and hence the Weyl group and modular quasi-character of its Levi are trivial. Since we've shown that \mathcal{Y} is a fixed point for the Knopp action, it follows from [Sak08] Theorem 2.2.2 (originally due to [Kno95]) that the "little Weyl group", $W_{\mathcal{X}}$, of \mathcal{X} (originally introduced by [Bri90]) coincides with the whole Weyl group W, of \mathscr{G} . We now turn our attention to verifying the assumptions in Section 8 of [Sak13]. Using, the discussion on simple root types above, and [Sak13] Section 7, we see that the assumptions of [Sak13] Theorem 7.2.1 are satisfied. Furthermore, it is clear that $\mathcal{Y}(\mathbf{Q}_p)$ has a unique $\mathscr{B}(\mathbf{Q}_p)$ -orbit, since $\operatorname{Stab}_{\mathscr{B}}(1) = Z_{\operatorname{GL}_2}^{\operatorname{diag}}$ is smooth and by [BG14] Proposition 1, the set of such orbits is in bijection with the kernel of the map

$$H^1\left(\operatorname{Gal}(\overline{\mathbf{Q}}_p/\mathbf{Q}_p), \operatorname{Stab}_{\mathscr{B}}(1)(\overline{\mathbf{Q}}_p)\right) \longrightarrow H^1\left(\operatorname{Gal}(\overline{\mathbf{Q}}_p/\mathbf{Q}_p), \mathscr{B}(\overline{\mathbf{Q}}_p)\right)$$

But the left-most term is trivial by an application of Hilbert 90 (in fact both terms are). We write $\mathscr{T}_{\mathcal{X}}$ for $\mathscr{T}/(\mathscr{T} \cap \mathscr{H}) = \mathscr{T}/Z_{\mathrm{GL}_2}^{\mathrm{diag}}$. The little Weyl group $W_{\mathcal{X}} = W$ acts on $X_*(\mathscr{T}_{\mathcal{X}})$ and it remains to verify that the natural map

(1)
$$\mathbf{C}[X_*(\mathscr{T})]^W \longrightarrow \mathbf{C}[X_*(\mathscr{T}_{\mathcal{X}})]^{W_{\mathcal{X}}}$$

is surjective. But $W_{\mathcal{X}} = W$, and $\mathbb{C}[X_*(\mathscr{T})]$ is a semisimple $\mathbb{C}[W]$ -module. Thus, surjectivity of (1) follows from surjectivity of the map $\mathbb{C}[X_*(\mathscr{T})] \to \mathbb{C}[X_*(\mathscr{T}_{\mathcal{X}})]$ which is simply the quotient from $\mathbb{C}[x_1, y_1, x_2, y_2, x_3, y_3]$ down to $\mathbb{C}[x_1, y_1, x_2, y_2, x_3, y_3]/(1 - x_1y_1x_2y_2x_3y_3)$. Finally, we are in a position to apply [Sak13] Corollary 8.0.4 which tells us that $C_c^{\infty}(\mathcal{X}(\mathbb{Q}_p))^{\mathscr{G}_p^{\circ}}$ is a cyclic $\mathcal{H}_{\mathscr{G}_p}^{\circ}$ -module generated by $ch(\mathcal{X}(\mathbb{Z}_p))$, where the Hecke action is the one induced from regarding $C_c^{\infty}(\mathcal{X}(\mathbb{Q}_p))$ as a smooth \mathscr{G}_p -representation via right translations. Finally, an unravelling of definitions, shows that we have a canonical isomorphism

$$C_c^{\infty}(\mathcal{X}(\mathbf{Q}_p))^{\mathscr{G}_p^{\circ}} \simeq C_c^{\infty}(H_p^{\circ}\backslash G_p/G_p^{\circ}) , F \mapsto (f_F : g \mapsto F(g, 1))$$

which is equivariant with respect to the Hecke action of $\mathcal{H}_{\mathscr{G}_p}^{\circ} \simeq \mathcal{H}_{G_p}^{\circ} \otimes \mathcal{H}_{H_p}^{\circ}$, and maps $ch(\mathcal{X}(\mathbf{Z}_p))$ to $ch(G_p^{\circ})$. This concludes the proof of the theorem. 3.2. Cyclicity of $\mathcal{I}(G_p/G_p^\circ)$. We will now use Theorem 3.1.1 to determine the structure of $\mathcal{I}(G_p/G_p^\circ)$ over the spherical Hecke algebra of G_p . Our approach draws heavily from [LSZ21b] Section 4.3.

Theorem 3.2.1. Let $\phi_0 = \operatorname{ch}(\mathbf{Z}_p^2)$ and $\xi_0 = \operatorname{ch}(G_p^\circ)$. The module $\mathcal{I}(G_p/G_p^\circ)$ is cyclic over $\mathcal{H}_{G_p}^\circ$ generated by $[\phi_0 \otimes \xi_0]$.

Proof. The proof is essentially translating Lemma 4.3.3 and Proposition 4.3.4 in [LSZ21b], and then applying the formalism of [LSZ21b] Theorem 4.3.6 together with Theorem 3.1.1. But since it's quite involved and combines quite a few results, we give the details here. We write A for the rank one torus given by Z_H . We have embeddings of algebras

$$\begin{aligned} &\mathcal{H}_{A_p}^{\circ} \xrightarrow{\Delta_H} \mathcal{H}_{H_p}^{\circ}, \ \mathrm{ch}([\begin{smallmatrix}p\\p\end{bmatrix} A_p^{\circ}) \mapsto \mathrm{ch}([\begin{smallmatrix}p\\p\end{bmatrix} H_p^{\circ}) \\ &\mathcal{H}_{A_p}^{\circ} \xrightarrow{\Delta_G} \mathcal{H}_{G_p}^{\circ}, \ \mathrm{ch}([\begin{smallmatrix}p\\p\end{bmatrix} A_p^{\circ}) \mapsto \mathrm{ch}(([\begin{smallmatrix}p\\p\end{bmatrix}, [\begin{smallmatrix}p\\p\end{bmatrix}) G_p^{\circ}). \end{aligned}$$

We write $\phi_0 = \operatorname{ch}(\mathbf{Z}_p^2)$. By [LSZ21b] Lemma 4.3.3, there exists an algebra homomorphism $\mathcal{H}_{H_p}^{\circ} \xrightarrow{\zeta_H} \mathcal{H}_{A_p}^{\circ}$, such that $\theta \cdot \phi_0 = (\Delta_H \circ \zeta_H)(\theta) \cdot \phi_0$ for every $\theta \in \mathcal{H}_{H_p}^{\circ}$. The definition of this map can be found in *op.cit*, and it is not hard to check that it satisfies the aforementioned property. We now consider an arbitrary element in $\mathcal{I}(G_p/G_p^{\circ})$ of the form $[\delta] = [\phi \otimes \xi]$. Up to a non-zero scalar multiple, such an element is equal to $[(\operatorname{ch}(A_p^{\circ}) \cdot_A \phi) \otimes \xi]$ in $\mathcal{I}(G_p/G_p^{\circ})$, and thus we may assume that $[\delta] = [\phi \otimes \xi]$ with $\phi \in (\mathcal{S}(\mathbf{Q}_p^2))^{A_p^{\circ}}$. By the proof of [LSZ21b] Proposition 4.3.4, the $\mathbf{C}[H_p]$ -module $\mathcal{S}(\mathbf{Q}_p^2)^{A_p^{\circ}}$ is cyclic, generated by ϕ_0 . Since the vector ϕ_0 is unramified, we may assume that $[\delta] = [(\theta \cdot \phi_0) \otimes \xi]$ with $\theta \in C_c^{\infty}(H_p/H_p^{\circ})$. But this is nothing more than $[\phi_0 \otimes (\theta' \cdot_H \xi)]$. Since ξ is right G_p° -invariant and $\theta' \in \mathcal{H}_{H_p}^{\circ}$, the convolution $\theta' \cdot_H \xi$ is an element of $C_c^{\infty}(H_p^{\circ}/G_p/G_p^{\circ})$. By Theorem 3.1.1, we can express $\theta' \cdot_H \xi$ as a linear combination of elements of the form $(\theta_1 \otimes \theta_2) \cdot \operatorname{ch}(G_p^{\circ})$ with $\theta_1 \otimes \theta_2 \in \mathcal{H}_{G_p}^{\circ} \otimes \mathcal{H}_{H_p}^{\circ}$. A quick integral computation shows that $(\theta_1 \otimes \theta_2) \cdot \operatorname{ch}(G_p^{\circ})$ is equal to $\theta_2 \cdot_H \theta_1'$. Thus, we may assume that $[\delta] = [\phi_0 \otimes (\alpha \cdot_H \beta)]$ where $\alpha \in \mathcal{H}_{H_p}^{\circ}$ and $\beta \in \mathcal{H}_{G_p}^{\circ}$. As in the proof of [LSZ21b] Theorem 4.3.6, we have the chain of equalities

$$\begin{split} [\delta] &= [\phi_0 \otimes (\alpha \cdot_H \beta)] \\ &= [(\alpha^{'} \cdot \phi_0) \otimes \beta] \\ &= [(\Delta_H(\alpha_1) \cdot \phi_0) \otimes \beta] \\ &= [\phi_0 \otimes (\Delta_H(\alpha_1)^{'} \cdot_H \beta)] \\ &= [\phi_0 \otimes (\Delta_G(\alpha_1)^{'} \cdot_G \beta)] \end{split}$$

where $\alpha_1 := \zeta_H(\alpha') \in \mathcal{H}_{A_n}^{\circ}$, and the last equality follows from the fact that

$$\Delta_H(\operatorname{ch}([{}^p {}_p] A_p^{\circ})) \cdot_H \beta = \beta([{}^p {}_p]^{-1} (-)) = \Delta_G(\operatorname{ch}([{}^p {}_p] A_p^{\circ})) \cdot_G \beta.$$

If we set $\theta_3 := \Delta_G(\alpha_1)' \cdot_G \beta$, which is an element of $\mathcal{H}^{\circ}_{G_p}$, we see that $[\delta]$ is equal to $[\phi_0 \otimes (\xi_0 \cdot_G \Lambda)]$, which is in turn equal to $\Lambda' *_G [\phi_0 \otimes \xi_0]$. This concludes the proof.

Remark 3.2.2. Later on, we will actually show that $\mathcal{I}(G_p/G_p^\circ)$ is free over $\mathcal{H}_{G_p}^\circ$. But that will require a different approach, so we postpone it for now.

We now showcase how the structure of $\mathcal{I}(G_p/G_p^\circ)$ can be used to recover, unify and adapt certain results of Jacquet, Shalika, Piatetski-Shapiro and Prasad to unramified representations of Whittaker type.

3.2.1. Relation to results of Jacquet, Piatetskii-Shapiro, Shalika.

Theorem 3.2.3 (Jacquet, Piatetskii-Shapiro, Shalika). For unramified H_p -representations $\pi_{p,1}, \pi_{p,2}$ of Whittaker type, we have

dim Hom_{$$H_p$$} ($\mathcal{S}(\mathbf{Q}_p^2) \otimes \pi_{p,1} \otimes \pi_{p,2}, \mathbf{1}$) ≤ 1

and every non-zero such linear form, is non-vanishing on $\phi_0 \otimes W_{\pi_n,1}^{\mathrm{sph}} \otimes W_{\pi_n,2}^{\mathrm{sph}}$.

A "generic" version of this was proved in [JPSS83] using invariant distributions to bound the dimension, and (even though not explicitly stated) one can use the local Rankin-Selberg zeta integral of [JPSS83] to realize a normalized linear form in this space and show that it takes the value 1 on $\phi_0 \otimes W_{\pi_p,1}^{\text{sph}} \otimes W_{\pi_p,2}^{\text{sph}}$ (see Section 6.4.1). If we put the existence of good test vectors aside, this was also proved "generically" in [KM15] using a purely algebraic approach, in particular Bernstein-Zelevinski derivatives, to show both existence and uniqueness. Below, we give a short algebraic proof of Theorem 3.2.3, which gives uniqueness, and existence of good test vectors, for unramified Whittaker type representations. This proof only relies on the structure of $\mathcal{I}(G_p/G_p^\circ)$ as a module over the spherical Hecke algebra. This Hecke approach is what will allow us to study the *integral* behaviour of this period later on.

Proof. There exists a surjective linear map from $\mathcal{S}(\mathbf{Q}_p^2) \otimes C_c^{\infty}(G_p/G_p^{\circ})$ to $\mathcal{S}(\mathbf{Q}_p^2) \otimes \pi_{p,1} \otimes \pi_{p,2}$ that is given by sending $\phi \otimes \xi$ to $\phi \otimes \left(\xi \cdot (W_{\pi_{p,1}}^{\mathrm{sph}} \otimes W_{\pi_{p,2}}^{\mathrm{sph}})\right)$. This clearly induces a map from $\mathcal{I}(G_p/G_p^{\circ})$ to $\left(\mathcal{S}(\mathbf{Q}_p^2) \otimes \pi_{p,1} \otimes \pi_{p,2}\right)_{H_p}$. By Theorem 3.2.1, every element of $\mathcal{I}(G_p/G_p^{\circ})$ is of the form $\theta * [\phi_0 \otimes \xi_0] = [\phi_0 \otimes \theta']$ for $\theta \in \mathcal{H}_{G_p}^{\circ}$. Hence the space $\left(\mathcal{S}(\mathbf{Q}_p^2) \otimes \pi_{p,1} \otimes \pi_{p,2}\right)_{H_p}$ is at most one dimensional and is generated by $[\phi_0 \otimes W_{\pi_{p,1}}^{\mathrm{sph}} \otimes W_{\pi_{p,2}}^{\mathrm{sph}}]$. This concludes the proof.

3.2.2. Relation to results of Prasad.

Theorem 3.2.4 (Prasad). Let $\pi_{p,1}, \pi_{p,2}, \pi_{p,3}$ be unramified H_p -representations of Whittaker type. Then

dim Hom_{H_p} $(\pi_{p,1} \otimes \pi_{p,2} \otimes \pi_{p,3}, \mathbf{1}) \leq 1$

and every non-zero such linear form is non-vanishing on $W^{\mathrm{sph}}_{\pi_{p,1}} \otimes W^{\mathrm{sph}}_{\pi_{p,2}} \otimes W^{\mathrm{sph}}_{\pi_{p,3}}$

For $\pi_{p,1}, \pi_{p,2}, \pi_{p,3}$ irreducible, the uniqueness is (part of) Theorem 1.1 in [Pra90] and is proved using invariant distributions. For the cases where some of the $\pi_{p,i}$ are reducible of Whittaker type, the uniqueness is treated in three results of [HS01]. For good test vectors in the unramified case, this is Theorem 1.3 in [Pra90] and in this case as well, the proof distinguishes between different cases. Below we give a short unified proof of this result that takes care of all unramified $\pi_{p,i}$ of Whittaker type at once. In fact later on we also remark how one can obtain the equality of dimension which is also part of [Pra90] and [HS01], for all unramified Whittaker type representations at once (see Remark 6.1.4). Our proof Theorem 3.2.4 also depends on the structure of $\mathcal{I}(G_p/G_p^{\circ})$ and a construction of [LSZ21b]. Once again, this approach will later allow us to get *integral* results for the non-zero period of this Hom-space using results for the lattice $\mathcal{I}(G_p/G_p^{\circ}, \mathbb{Z}[1/p])$.

Proof. By twisting, we may assume that $\pi_{p,1}$ is the normalized induction $I(\chi_p, |\cdot|_p^{-1/2})$ with χ_p an unramified quasi-character of \mathbf{Q}_p^{\times} with $I(\chi_p, |\cdot|_p^{-1/2})$ of Whittaker type. Here $I(\chi_p, |\cdot|_p^{-1/2})$ denotes the representation $\operatorname{ind}_{B_p}^{G_p} \begin{bmatrix} \chi_p & |\cdot|_p^{-1/2} \end{bmatrix}$ where B_p is the upper triangular Borel, and the diagonal character acts trivially on the unipotent radical. Using Proposition 3.2.2 and Lemma 3.2.5 of [LSZ21a], and Proposition 1.5.2 in [Gro20], one has a non-zero H_p -equivariant map $S(\mathbf{Q}_p^2) \to \pi_{p,1}$. Since $\pi_{p,1}$ is unramified of Whittaker type, this map is surjective and maps ϕ_0 to $W_{\pi_{p,1}}^{\operatorname{sph}}$. This induces a surjective H_p -equivariant map $S(\mathbf{Q}_p^2) \otimes C_c^{\infty}(G_p/G_p^\circ) \to \pi_{p,1} \otimes \pi_{p,2} \otimes \pi_{p,3}$, which in turn induces a surjective linear map $\mathcal{I}(G_p/G_p^\circ) \to (\pi_{p,1} \otimes \pi_{p,2} \otimes \pi_{p,3})_{H_p}$. The result then follows at once in the same fashion as the proof of Theorem 3.2.3 given above, once again utilizing Theorem 3.1.1.

Remark 3.2.5. We would now like to realise the cyclicity of the Hecke module $\mathcal{I}(G_p/G_p^\circ)$ and study its properties when we specialise to integral data as in Definition 2.2.2. The right way forward as mentioned in the introduction is *not* to directly use the local Rankin-Selberg zeta integral as in [LSZ21b]. Indeed, it does not seem possible to treat *general* integral test vectors that way and obtain meaningful results at an integral level. Instead, we proceed as in the following sections.

3.3. Rephrasing using the mirabolic subgroup. From now on, we will drop the notation $[\cdot]$ on elements of $\mathcal{I}(G_p/G_p^{\circ})$. We denote by P the mirabolic subgroup of $H = \operatorname{GL}_2$, given by $\{[* \ 1 \ 1]\}$, and we will also regard it as a subgroup of G through the diagonal embedding $H \hookrightarrow G$. As usual, we write P_p for its \mathbf{Q}_p -points and P_p° for its \mathbf{Z}_p -points. There's an identification

$$P_p \setminus H_p \xrightarrow{\simeq} \mathbf{Q}_p^2 - \{(0,0)\}, \ h \mapsto (0,1)h.$$

This induces an embedding of H_p -representations (where recall that Ind denotes unnormalized, non-compact induction)

$$\mathcal{S}(\mathbf{Q}_p^2) \hookrightarrow \operatorname{Ind}_{P_p}^{H_p} \mathbf{1} , \ \phi \mapsto (f_\phi : h \mapsto \phi((0,1)h))$$

under which $\mathcal{S}_0(\mathbf{Q}_p^2)$ gets identified with c-Ind $_{P_p}^{H_p} \mathbf{1}$.

Definition 3.3.1. We define the linear map

$$\Xi: \mathcal{S}(\mathbf{Q}_p^2) \otimes C_c^{\infty}(G_p/G_p^\circ) \longrightarrow C^{\infty}(P_p \backslash G_p/G_p^\circ) = \left(\mathrm{Ind}_{P_p}^{G_p} \mathbf{1}\right)^{G_p^\circ}$$
$$\phi \otimes \xi \mapsto \left(\Xi_{\phi \otimes \xi}: g \mapsto \int_{H_p} \xi(h^{-1}g) f_{\phi}(h) \ dh\right)$$

A note should be made on the "convergence" of the integral used to define the map Ξ , which follows from the compactness of $H_p \cap g_1 G_p^{\circ} g_2$ in H_p , for any $g_1, g_2 \in G_p$. It follows by construction, that $\Xi_{h(\phi \otimes \xi)} = \Xi_{\phi \otimes \xi}$ for any $h \in H_p$, and hence Ξ factors through $\mathcal{I}(G_p/G_p^{\circ})$. An unraveling of definitions also shows that this map is Hecke equivariant with respect to $\mathcal{H}_{G_p}^{\circ}$, where the action on the right is given by the natural action of $\mathcal{H}_{G_p}^{\circ}$ on $(\operatorname{Ind}_{P_p}^{H_p} \mathbf{1})^{G_p^{\circ}}$.

Once again, we emphasize that for $\phi \otimes \xi$ with $\phi \in S_0(\mathbf{Q}_p^2)$, the corresponding function $\Xi_{\phi \otimes \xi}$ belongs in $C_c^{\infty}(P_p \setminus G_p/G_p^{\circ})$, i.e. it is already compactly supported. This will be important later on. We write $C^{\infty}(P_p \setminus G_p/G_p^{\circ}, \mathbf{Z}[1/p])$ for the $\mathcal{H}_{G_p}^{\circ}(\mathbf{Z}[1/p])$ -submodule of $C^{\infty}(P_p \setminus G_p/G_p^{\circ})$ consisting of $\mathbf{Z}[1/p]$ -valued functions, and similarly for compactly supported functions.

Proposition 3.3.2. The image of $\mathcal{I}(G_p/G_p^\circ, \mathbb{Z}[1/p])$ under the map Ξ , is contained in $C^{\infty}(P_p \setminus G_p/G_p^\circ, \mathbb{Z}[1/p])$.

Proof. Let $\phi \otimes \operatorname{ch}(gG_p^\circ)$ be an element of $\mathcal{I}(G_p/G_p^\circ, \mathbb{Z}[1/p])$. Recall that this means by definition that ϕ is valued in

$$\frac{1}{\operatorname{vol}_{H_p}\left(\operatorname{Stab}_{H_p}(\phi) \cap gG_p^{\circ}g^{-1}\right)}\mathbf{Z}[1/p]$$

For ease of notation, write $\delta = \phi \otimes \operatorname{ch}(gG_p^\circ)$. For $x \in G_p$, we have by construction that

(2)
$$\Xi_{\delta}(x) = \int_{H_p} \left(h \cdot \operatorname{ch}(gG_p^{\circ}) \right)(x) \ (h \cdot \phi)(0,1) \ dh$$

Let K be the open compact of H_p given by $\operatorname{Stab}_{H_p}(\phi) \cap gG_p^{\circ}g^{-1}$. We can split the integral in (2) into a finite sum of the form

(3)
$$\Xi_{\delta}(x) = \sum_{\gamma} \int_{\gamma K} \left(h \cdot \operatorname{ch}(gG_p^{\circ}) \right)(x) \ (h \cdot \phi)(0,1) \ dh = \operatorname{vol}_{H_p}(K) \sum_{\gamma} \operatorname{ch}(gG_p^{\circ})(\gamma^{-1}x) \ \phi((0,1)\gamma).$$

By the assumption on the values of ϕ , (3) is an element of $\mathbb{Z}[1/p]$ which concludes the proof.

3.4. Mapping to compact induction. We first start by noting that the element $\Xi_0 = \Xi(\operatorname{ch}(\mathbf{Z}_p^2) \otimes \operatorname{ch}(G_p^\circ))$, even though it is unramified in the sense that is G_p° -fixed, it does not coincide with the trivial unramified element in c-Ind $_{P_p}^{G_p} \mathbf{1} \subseteq \operatorname{Ind}_{P_p}^{G_p} \mathbf{1}$ given by $\operatorname{ch}(P_p G_p^\circ)$. For $i \in \mathbf{Z}$, we write $Z_{p,i}$ for the central subset of H_p given by $\left\{ \begin{bmatrix} p^n \\ p^n \end{bmatrix} \mid n \geq i \right\}$.

Lemma 3.4.1. The function Ξ_0 is equal to the characteristic function $\operatorname{ch}(Z_{p,0}P_pG_p^\circ)$ in $C^{\infty}(P_p\backslash G_p/G_p^\circ)$.

Proof. For an element $g = (g_1, g_2) \in G_p$, we have

$$\Xi_0(g) = \int_{H_p \cap gG_p^\circ} \operatorname{ch}(\mathbf{Z}_p^2)((0,1)h) \ dh = \begin{cases} 0, & \text{if } g_1 H_p^\circ \neq g_2 H_p^\circ \\ \operatorname{ch}(\mathbf{Z}_p^2)((0,1)g_1), & \text{if } g_1 H_p^\circ = g_2 H_p^\circ. \end{cases}$$

The result then follows from the decomposition $H_p = \bigsqcup_{n \in \mathbb{Z}} \begin{bmatrix} p^n & \\ & p^n \end{bmatrix} P_p H_p^{\circ}$.

Proposition 3.4.2. There exists $\Psi \in \operatorname{End}_{\mathcal{H}_{G_p}^\circ}(C^{\infty}(P_p \setminus G_p/G_p^\circ))$ for which the composition $\Xi_c := \Psi \circ \Xi$ gives rise to the following well-defined commutative diagram, where the top map is $\mathcal{H}_{G_p}^\circ$ -equivariant.

Finally, the map Ξ_c maps $\operatorname{ch}(\mathbf{Z}_p^2) \otimes \operatorname{ch}(G_p^\circ)$ to $\operatorname{ch}(P_p G_p^\circ)$.

Proof. We write $S_p = S_{p,1}S_{p,2}$ for the Hecke operator in $(\mathcal{H}_{G_p}^{\circ})^{\times}$ associated to the element $([p_p], [p_p])$ in G_p . We define the $\mathcal{H}_{G_p}^{\circ}$ -equivariant endomorphism Ψ to be the map

$$\Psi: C^{\infty}(P_p \backslash G_p / G_p^{\circ}) \to C^{\infty}(P_p \backslash G_p / G_p^{\circ}), \ \xi \mapsto (1 - \mathcal{S}_p^{-1}) \cdot \xi$$

As in the statement of the proposition, we put $\Xi_c = \Psi \circ \Xi$. By Hecke equivariance and Theorem 3.2.1, it suffices to show that Ξ_c maps $\operatorname{ch}(\mathbf{Z}_p^2) \otimes \operatorname{ch}(G_p^\circ)$ to an element in the compact induction. In particular we will show that $\operatorname{ch}(\mathbf{Z}_p^2) \otimes \operatorname{ch}(G_p^\circ)$ is mapped to $\operatorname{ch}(P_p G_p^\circ)$. By Lemma 3.4.1, $\Xi_c \left(\operatorname{ch}(\mathbf{Z}_p^2) \otimes \operatorname{ch}(G_p^\circ)\right)$ is given by $\Psi \left(\operatorname{ch}(Z_{p,0} P_p G_p^\circ)\right)$. This in turn is given by the function

$$\Psi\left(\operatorname{ch}(Z_{p,0}P_pG_p^{\circ})\right)(g) = \operatorname{ch}(Z_{p,0}P_pG_p^{\circ})(g) - \int_{G_p} \mathcal{S}_p^{-1}(\gamma) \operatorname{ch}(Z_{p,0}P_pG_p^{\circ})(g\gamma) d\gamma$$
$$= \operatorname{ch}(Z_{p,0}P_pG_p^{\circ})(g) - \operatorname{ch}(Z_{p,0}P_pG_p^{\circ})\left(g\left[\begin{smallmatrix}p\\p\end{smallmatrix}\right]^{-1}\right).$$

From this it is clear that $\Xi_c \left(\operatorname{ch}(\mathbf{Z}_p^2) \otimes \operatorname{ch}(G_p^\circ) \right) = \operatorname{ch}(P_p G_p^\circ)$ as claimed. Finally, the fact that Ξ_c induces a map on integral level follows immediately from Proposition 3.3.2.

Our constructions so far can be summarised in the following commutative diagrams, where the square on the left, is equivariant with respect to the action of the spherical Hecke algebra $\mathcal{H}_{G_p}^{\circ}$ and $\mathcal{I}(G_p/G_p^{\circ})$ is cyclic over $\mathcal{H}_{G_p}^{\circ}$ generated by $\delta_0 := \operatorname{ch}(\mathbf{Z}_p^2) \otimes \operatorname{ch}(G_p^{\circ})$.

$$\begin{array}{cccc} \delta_0 & \mathcal{I}(G_p/G_p^\circ) \xrightarrow{\Xi} C^{\infty}(P_p \backslash G_p/G_p^\circ) & \mathcal{I}(G_p/G_p^\circ, \mathbf{Z}[1/p]) \xrightarrow{\Xi} C^{\infty}(P_p \backslash G_p/G_p^\circ, \mathbf{Z}[1/p]) \\ & \downarrow_{\Xi_c} & \downarrow_{\Xi_c} & \cdot (1-\mathcal{S}_p^{-1}) \downarrow & \downarrow_{\Xi_c} & \cdot (1-\mathcal{S}_p^{-1}) \downarrow \\ & ch(P_p G_p^\circ) & C_c^{\infty}(P_p \backslash G_p/G_p^\circ) & \subset C^{\infty}(P_p \backslash G_p/G_p^\circ) & C_c^{\infty}(P_p \backslash G_p/G_p^\circ, \mathbf{Z}[1/p]) & \hookrightarrow C^{\infty}(P_p \backslash G_p/G_p^\circ, \mathbf{Z}[1/p]) \end{array}$$

4. Hecke equivariant forms on $C_c^{\infty}(P_p \setminus G_p/G_p^{\circ})$

We fix once and for all an additive character $\psi : \mathbf{Q}_p \to \mathbf{C}^{\times}$ of conductor one. It is standard that such a character exists (e.g [RV13] Chapter 7). We consider the family of G_p -representations $\pi_{p,1} \boxtimes \pi_{p,2}$ where $\pi_{p,1}$ and $\pi_{p,2}$ are unramified H_p -representations of Whittaker type. As it is often the case, we identify $\pi_{p,1}$ with its Whittaker model $\mathcal{W}(\pi_{p,1},\psi)$ and similarly, $\pi_{p,2}$ with $\mathcal{W}(\pi_{p,2},\psi^{-1})$. For further theory surrounding Whittaker models for this family of representations, we refer the reader to [Bum97].

4.1. Splitting of measures. It follows from a standard computation, that the modular quasi-character of the mirabolic subgroup P_p is simply given by $\delta_P\left(\begin{bmatrix}a&b\\1\end{bmatrix}\right) = |a|_p$. We write $d^R x$ for the normalized right Haar measure on P_p . Then $\delta_P^{-1} d^R x$ is the normalized left Haar measure on P_p . We write $C_c^{\infty}(P_p \setminus G_p; \delta_P)$ for the space of smooth functions on G_p , that are compactly supported modulo P_p on the left, transform by δ_P under left P_p -translations, and are fixed under right translations by some open compact subgroup of G_p° . Following ([KM15] 2.2) and ([Vig96] I 2.8), there exists a surjective map

(4)
$$\mathcal{H}(G_p) \longrightarrow C_c^{\infty}(P_p \backslash G_p; \delta_P), \ f \mapsto \left(f^P : g \mapsto \int_{P_p} f(xg) \delta_P(x)^{-1} \ d^R x \right)$$

(

and a unique, up to scalars, linear form $d_{P_p \setminus G_p} g$ on $C_c^{\infty}(P_p \setminus G_p; \delta_P)$, that is right invariant by G_p . We write

$$d_{P_p \setminus G_p} g(f) := \int_{P_p \setminus G_p} f(g) \ d_{P_p \setminus G_p}, \ f \in C_c^{\infty}(P_p \setminus G_p; \delta_P) g.$$

Up to correct normalization of $d_{P_p \setminus G_p} g$, we have

(5)
$$\int_{G_p} h(g) \, dg = \int_{P_p \setminus G_p} h^P(g) \, d_{P_p \setminus G_p} g, \ h \in \mathcal{H}(G_p)$$

where dg is the normalized Haar measure on G_p . We want to be able to unfold the integral $\int_{P_p \setminus G_p} f(g) d_{P_p \setminus G_p} g$ for $f \in C_c^{\infty}(P_p \setminus G_p / G_p^\circ; \delta_P)$.

Lemma 4.1.1. Let f be a function in $C_c^{\infty}(P_p \setminus G_p/G_p^{\circ}; \delta_P)$. We write Z_p for the center of H_p embedded diagonally in G_p . Then,

$$\int_{P_p \setminus G_p} f(g) \ d_{P_p \setminus G_p} g = \int_{Z_p} \int_{\mathrm{GL}_2(\mathbf{Q}_p)} f(z(\gamma, 1)) \ dz \ d\gamma.$$

Proof. By (4), we can write f as h^P for some h in $\mathcal{H}(G_p)$. It is clear that G_p splits as a semidirect product of $\operatorname{GL}_2(\mathbf{Q}_p) \times 1$ and H_p . Hence by a slightly modified version of [KM15] Lemma 2.4, using unimodularity, and (5), we see that the integral in question is equal to $\int_{H_p} \int_{\operatorname{GL}_2(\mathbf{Q}_p)} h(x(\gamma, 1)) dx d\gamma$. Using Cartier's argument and *op.cit* one more time, this is nothing more than

$$\int_{P_p} \int_{Z_p} \int_{H_p^\circ} \int_{\mathrm{GL}_2(\mathbf{Q}_p)} h\left(yzk(\gamma,1)\right) \ d^L y \ dz \ dk \ d\gamma = \int_{P_p} \int_{Z_p} \int_{H_p^\circ} \int_{\mathrm{GL}_2(\mathbf{Q}_p)} h\left(yzk(\gamma,1)\right) \delta_P(y)^{-1} \ d^R y \ dz \ dk \ d\gamma.$$

By definition of h^P , this reduces to

$$\int_{Z_p} \int_{\mathrm{GL}_2(\mathbf{Z}_p)} \int_{\mathrm{GL}_2(\mathbf{Q}_p)} f(z(\kappa\gamma,\kappa)) \ dz \ d\kappa \ d\gamma = \int_{Z_p} \int_{\mathrm{GL}_2(\mathbf{Q}_p)} f(z(\gamma,1)) \ dz \ d\gamma$$

where the last equality follows from the fact that f is right G_p° -invariant, together with a change of variables.

4.2. The linear form Λ_{Π_p} . The goal of this section is to construct, for each unramified G_p -representation Π_p of Whittaker type, a linear form Λ_{Π_p} on $C_c^{\infty}(P_p \backslash G_p/G_p^{\circ})$, satisfying a certain Hecke equivariance condition. This will allow us to "realize", in a way, the cyclicity of the module $\mathcal{I}(G_p/G_p^{\circ})$.

Definition 4.2.1. Let $\pi_{p,1}$ and $\pi_{p,2}$ be two unramified H_p -representations of Whittaker type and let $\Pi_p = \pi_{p,1} \boxtimes \pi_{p,2}$. For $f \in C_c^{\infty}(P_p \setminus G_p/G_p^{\circ})$ and $g \in G_p$, we define

$$\mathscr{W}_{\Pi_{p}}(s;g,f) := f(g) \mid \det(g_{2})|_{p}^{s} \int_{\mathbf{Q}_{p}^{\times}} \left(W_{\pi_{p,1}}^{\mathrm{sph}} \boxtimes W_{\pi_{p,2}}^{\mathrm{sph}} \right) \left(\begin{bmatrix} a \\ 1 \end{bmatrix} g \right) \mid a|_{p}^{s-1} d^{\times}a \in \mathbf{C}(p^{s},p^{-s})$$

where g_2 denotes the projection onto the second factor.

For fixed f and g and $\Re(s)$ large enough, (independently of f and g), using one of the main results of [JPSS83], the integral appearing in the definition of $\mathscr{W}_{\Pi_p}(s; g, f)$ converges and admits unique meromorphic continuation as a rational function of p^s . It is also compactly supported modulo P_p on the left by construction. For $x = \begin{bmatrix} x_1 & x_2 \\ 1 \end{bmatrix} \in P_p$, we have

(6)
$$\mathscr{W}_{\Pi_{p}}(s;xg,f) = f(xg) |\det(xg_{2})|_{p}^{s} \int_{\mathbf{Q}_{p}^{\times}} \left(W_{\pi_{p,1}}^{\mathrm{sph}} \boxtimes W_{\pi_{p,2}}^{\mathrm{sph}} \right) \left(\begin{bmatrix} ax_{1} & ax_{2} \\ 1 \end{bmatrix} g \right) |a|_{p}^{s-1} d^{\times}a$$
$$= f(g) |\det(xg_{2})|_{p}^{s} |x_{1}|_{p}^{1-s} \int_{\mathbf{Q}_{p}^{\times}} \left(W_{\pi_{p,1}}^{\mathrm{sph}} \boxtimes W_{\pi_{p,2}}^{\mathrm{sph}} \right) \left(\begin{bmatrix} a \\ 1 \end{bmatrix} g \right) |a|_{p}^{s-1} d^{\times}a$$
$$= \delta_{P}(x) \mathscr{W}_{\Pi_{p}}(s;g,f).$$

Definition 4.2.2. For $f \in C_c^{\infty}(P_p \setminus G_p/G_p^{\circ})$ and notation as in Definition 4.2.1, we define

$$\Lambda_{\Pi_p}(s;f) := \int_{P_p \setminus G_p} \mathscr{W}_{\Pi_p}(s;g,f) \ d_{P_p \setminus G_p}g \in \mathbf{C}(p^s,p^{-s})$$

Proposition 4.2.3. For $f \in C_c^{\infty}(P_p \setminus G_p/G_p^{\circ})$, the integral $\Lambda_{\Pi_p}(s; f)$ converges for $\Re(s)$ large enough and admits unique meromorphic continuation as a rational function of p^s , in the fractional ideal $L(\Pi_p, s)\mathbf{C}[p^s, p^{-s}]$.

Proof. For $\Re(s)$ sufficiently large, we can unfold the integral using Lemma 4.1.1. The result then follows from [JPSS83].

$$\Lambda_{\Pi_p}: C_c^{\infty}(P_p \backslash G_p / G_p^{\circ}) \longrightarrow \mathbf{C}, \ \Lambda_{\Pi_p}(f) := \lim_{s \to 0} \frac{\Lambda_{\Pi_p}(s; f)}{L(\Pi_p, s)}$$

which is well-defined by Proposition 4.2.3.

4.2.1. Equivariance properties. As shown in the following proposition, the linear form Λ_{Π_p} becomes $\mathcal{H}_{G_p}^{\circ}$ equivariant if we interpret **C** as the one dimensional subspace of G_p° -fixed vectors in the smooth dual of Π_p .

Proposition 4.2.5. Let $\Pi_p = \pi_{p,1} \boxtimes \pi_{p,2}$, with $\pi_{p,i}$ unramified H_p -representations of Whittaker type. We write ω_{Π_p} for its central character and Θ_{Π_p} for its spherical Hecke eigensystem.

(1) The linear form Λ_{Π_p} defines an element of

$$\operatorname{Hom}_{\mathcal{H}_{G_p}^{\circ}}\left(C_c^{\infty}(P_p \backslash G_p / G_p^{\circ}), \left(\Pi_p^{\vee}\right)^{G_p^{\circ}}\right).$$

(2) For $f_0 = ch(P_p G_p^\circ)$, we have $\Lambda_{\Pi_p}(f_0) = L(\omega_{\Pi_p}, 0)^{-1} = \Theta_{\Pi_p}(1 - \mathcal{S}_p)$.

Proof. We firstly prove the first part. Let $f \in C_c^{\infty}(P_p \setminus G_p)$ and $\mathcal{P} \in \mathcal{H}_{G_p}^{\circ}$.

$$\begin{split} \Lambda_{\Pi_p}(s; \mathcal{P} \cdot f) &= \int_{P_p \setminus G_p} \int_{\mathbf{Q}_p^{\times}} (\mathcal{P} \cdot f)(g) \mid \det(g_2) \mid_p^s \left(W_{\pi_{p,1}}^{\mathrm{sph}} \boxtimes W_{\pi_{p,2}}^{\mathrm{sph}} \right) \left(\begin{bmatrix} a \\ 1 \end{bmatrix} g \right) \mid a \mid_p^{s-1} d^{\times} a \ d_{P_p \setminus G_p} g \\ &= \int_{P_p \setminus G_p} \int_{\mathbf{Q}_p^{\times}} \int_{G_p} \mathcal{P}(\gamma) f(g\gamma) \mid \det(g_2) \mid_p^s \left(W_{\pi_{p,1}}^{\mathrm{sph}} \boxtimes W_{\pi_{p,2}}^{\mathrm{sph}} \right) \left(\begin{bmatrix} a \\ 1 \end{bmatrix} g \right) \mid a \mid_p^{s-1} d\gamma \ d^{\times} a \ d_{P_p \setminus G_p} g \end{split}$$

From Section 4.1, $d_{P_p \setminus G_p} g$ is right G_p -invariant, hence the above is equal to

$$\int_{P_p \setminus G_p} \int_{\mathbf{Q}_p^{\times}} \int_{G_p} \mathcal{P}(\gamma) f(g) \mid \det(g_2 \gamma_2^{-2}) \mid_p^s \left(W_{\pi_{p,1}}^{\mathrm{sph}} \boxtimes W_{\pi_{p,2}}^{\mathrm{sph}} \right) \left(\begin{bmatrix} a \\ 1 \end{bmatrix} g \gamma^{-1} \right) \mid a \mid_p^{s-1} d\gamma \ d^{\times} a \ d_{P_p \setminus G_p} g.$$

Since G_p is unimodular, the Haar measure $d\gamma$ is invariant under the change of variables $\gamma \rightsquigarrow \gamma^{-1}$. Thus, this is equal to

$$\int_{P_p \setminus G_p} \int_{\mathbf{Q}_p^{\times}} f(g) |\det(g_2)|_p^s \left(\int_{G_p} |\det(\gamma_2)|_p^s \mathcal{P}'(\gamma) \left(W_{\pi_{p,1}}^{\mathrm{sph}} \boxtimes W_{\pi_{p,2}}^{\mathrm{sph}} \right) \left(\begin{bmatrix} a \\ 1 \end{bmatrix} g \gamma \right) d\gamma \right) |a|_p^{s-1} d^{\times} a d_{P_p \setminus G_p} g.$$

The first part then follows upon using Cartan decomposition to express \mathcal{P} as a linear combination of G_p° double coset operators. For the second part, using Lemma 4.1.1, we have

$$\begin{split} \Lambda_{\Pi_p}(s;f_0) &= \int_{Z_p} \int_{\mathrm{GL}_2(\mathbf{Q}_p)} \mathscr{W}_{\Pi_p}(s;(z\gamma,z),f_0) \ dz \ d\gamma. \\ &= \int_{Z_p} \int_{\mathrm{GL}_2(\mathbf{Q}_p)} \int_{\mathbf{Q}_p^{\times}} f_0((z\gamma,z)) |\det(z)|_p^s \left(W_{\pi_{p,1}}^{\mathrm{sph}} \boxtimes W_{\pi_{p,2}}^{\mathrm{sph}} \right) \left(\begin{bmatrix} a \\ 1 \end{bmatrix} (z\gamma,z) \right) \ |a|_p^{s-1} \ d^{\times}a \ d\gamma \ dz. \end{split}$$

Now, $f_0((z\gamma, z))$ vanishes for z outside Z_p° , and is equal to $f_0((\gamma, 1))$ for $z \in Z_p^{\circ}$. Similarly, $f_0((\gamma, 1))$ vanishes for γ outside $\operatorname{GL}_2(\mathbf{Z}_p)$ and is identically equal to one on $\operatorname{GL}_2(\mathbf{Z}_p)$. Thus, the triple integral above collapses to give

$$\int_{\mathbf{Q}_p^{\times}} \left(W_{\pi_{p,1}}^{\mathrm{sph}} \boxtimes W_{\pi_{p,2}}^{\mathrm{sph}} \right) \left(\begin{bmatrix} a \\ 1 \end{bmatrix} \right) \ |a|_p^{s-1} \ d^{\times} a.$$

It is well-known that the latter is given by $L(\Pi_p, s)L(\omega_{\Pi_p}, 2s)^{-1}$ and thus the proof of the second part is complete.

Remark 4.2.6. It is clear from Proposition 4.2.5 that the linear form $\Lambda_{\pi_{p,1}\boxtimes\pi_{p,2}}$ is non-zero for $\omega_{\pi_{p,1}} \neq \omega_{\pi_{p,2}}^{-1}$.

Corollary 4.2.7. The Hecke equivariant map Ξ_c gives a canonical isomorphism onto its image, of free rank one $\mathcal{H}_{G_p}^{\circ}$ -modules.

Proof. Suppose that some $\mathcal{P} \in \mathcal{H}_{G_p}^{\circ}$ annihilates $\operatorname{ch}(\mathbf{Z}_p^2) \otimes \operatorname{ch}(G_p^{\circ})$ in $\mathcal{I}(G_p/G_p^{\circ})$. By the Hecke equivariance of Ξ_c , \mathcal{P} also annihilates $\Xi_c(\operatorname{ch}(\mathbf{Z}_p^2) \otimes \operatorname{ch}(G_p^{\circ})) = \operatorname{ch}(P_pG_p^{\circ})$. Now, applying Λ_{Π_p} , and using Proposition 4.2.5 part (2), we obtain

(7)
$$\Theta_{\Pi_p} \left(\mathcal{P}'(1-\mathcal{S}_p) \right) = 0.$$

The equality in (2) holds for every unramified $\Pi_p = \pi_{p,1} \boxtimes \pi_{p,2}$ with $\pi_{p,i}$ of Whittaker type. This family of representations is dense in $\text{Spec}(\mathcal{H}_{G_p}^{\circ})$ and thus $\mathcal{P}'(1-\mathcal{S}_p)$ is identically zero in $\mathcal{H}_{G_p}^{\circ}$. From here it follows instantly that $\mathcal{P} = 0$. This combined with Theorem 3.2.1, shows that $\mathcal{I}(G_p/G_p^{\circ})$ is free of rank one and that Ξ_c is injective, completing the proof.

Definition 4.2.8. Let $\delta_0 = \operatorname{ch}(\mathbf{Z}_p^2) \otimes \operatorname{ch}(G_p^\circ)$. For $\delta \in \mathcal{I}(G_p/G_p^\circ)$, the Hecke operator \mathcal{P}_{δ} is the unique element of $\mathcal{H}_{G_p}^\circ$ that satisfies $\mathcal{P}_{\delta} \cdot \delta_0 = \delta$ in $\mathcal{I}(G_p/G_p^\circ)$.

Remark 4.2.9. Equivalently, the Hecke operator \mathcal{P}_{δ} is the unique element of $\mathcal{H}_{G_p}^{\circ}$ for which $\mathcal{P}_{\delta} \cdot \operatorname{ch}(P_p G_p^{\circ})$ is equal to $\Xi_c(\delta)$.

4.3. Explicit formulas for Λ_{Π_p} in terms of *L*-factors and Satake parameters. Now that we have constructed the Hecke equivariant linear form Λ_{Π_p} , we want to directly compute the integrals involved in its definition, for general input data.

Definition 4.3.1. The polynomial $\mathcal{P}_p \in \mathcal{H}_{G_p}^{\circ}[x]$, is characterized by the property

$$\Theta_{\Pi_p}(\mathcal{P}_p)(p^{-s}) = L(\Pi_p, s)^-$$

for every unramified G_p -representation $\Pi_p = \pi_{p,1} \boxtimes \pi_{p,2}$, where Θ_{Π_p} denotes the spherical Hecke eigensystem of Π_p and $L(\Pi_p, s)$ is the usual Rankin-Selberg local L-factor attached to such an unramified principal-series representation Π_p (e.g [JL06]).

We note that such a polynomial does exists as the local L-factor can be expressed as a Weyl-group invariant polynomial in the Satake parameters of Π_p .

Definition 4.3.2. Let $\Pi_p = \pi_{p,1} \boxtimes \pi_{p,2}$ with $\pi_{p,i}$ unramified H_p -representations of Whittaker type. Write $\alpha_{\pi_{p,i}}$ and $\beta_{\pi_{p,i}}$ for the Satake parameters of $\pi_{p,i}$. For $r \in \mathbb{Z}_{\geq 1}$, we define

$$L_{1}^{(r)}(\omega_{\Pi_{p}},s)^{-1} := \frac{(\alpha_{\pi_{p,1}}^{r} - \beta_{\pi_{p,1}}^{r}) - \alpha_{\pi_{p,1}}\beta_{\pi_{p,1}}(\alpha_{\pi_{p,2}} + \beta_{\pi_{p,2}})(\alpha_{\pi_{p,1}}^{r-1} - \beta_{\pi_{p,1}}^{r-1})p^{-s} + (\alpha_{\pi_{p,2}}\beta_{\pi_{p,2}}\alpha_{\pi_{p,1}}^{2}\beta_{\pi_{p,1}}^{2})(\alpha_{\pi_{p,1}}^{r-2} - \beta_{\pi_{p,1}}^{r-2})p^{-2s}}{\alpha_{\pi_{p,1}} - \beta_{\pi_{p,1}}}$$

$$L_{2}^{(r)}(\omega_{\Pi_{p}},s)^{-1} := \frac{(\alpha_{\pi_{p,2}}^{r} - \beta_{\pi_{p,2}}^{r}) - \alpha_{\pi_{p,2}}\beta_{\pi_{p,2}}(\alpha_{\pi_{p,1}} + \beta_{\pi_{p,1}})(\alpha_{\pi_{p,2}}^{r-1} - \beta_{\pi_{p,2}}^{r-1})p^{-s} + (\alpha_{\pi_{p,1}}\beta_{\pi_{p,1}}\alpha_{\pi_{p,2}}^{2}\beta_{\pi_{p,2}}^{2})(\alpha_{\pi_{p,2}}^{r-2} - \beta_{\pi_{p,2}}^{r-2})p^{-2s}}{\alpha_{\pi_{p,2}} - \beta_{\pi_{p,2}}}$$

Remark 4.3.3. The L-factors considered above are natural generalisations of $L(\omega_{\Pi_p}, s)$ and will show up in our computation of Λ_{Π_p} . In fact, one has $L_1^{(1)}(\omega_{\Pi_p}, s) = L_2^{(1)}(\omega_{\Pi_p}, s) = L(\omega_{\Pi_p}, 2s)$.

Lemma 4.3.4. For $r \in \mathbb{Z}_{\geq 1}$, there exist polynomials $\mathcal{P}_{p,1}^{(r)}(x)$ and $\mathcal{P}_{p,2}^{(r)}(x)$ in $\mathcal{H}_{G_p}^{\circ}[x]$ such that

$$\Theta_{\Pi_p}\left(\mathcal{P}_{p,1}^{(r)}\right)(p^{-s}) = L_1^{(r)}(\omega_{\Pi_p}, s)^{-1} \text{ and } \Theta_{\Pi_p}\left(\mathcal{P}_{p,2}^{(r)}\right)(p^{-s}) = L_2^{(r)}(\omega_{\Pi_p}, s)^{-1}$$

for every Π_p as in Definition 4.3.2.

Proof. The inverse L-factors $L_i^{(r)}(\omega_{\Pi_p}, s)^{-1}$ are polynomials in p^{-s} with coefficients that are themselves Weyl-group invariant polynomials in the Satake parameters of Π_p . The result then follows at once.

Lemma 4.3.5. For $n \in \mathbb{Z}_{\geq 0}$ and with notation as above, we have

$$\int_{\mathbf{Q}_{p}^{\times}} \left(W_{\pi_{p,1}}^{\mathrm{sph}} \boxtimes W_{\pi_{p,2}}^{\mathrm{sph}} \right) \left(\begin{bmatrix} ap^{n} \\ 1 \end{bmatrix}, \begin{bmatrix} a \\ 1 \end{bmatrix} \right) \ |a|_{p}^{s-1} \ d^{\times}a = L(\Pi_{p}, s) \cdot p^{\frac{-n}{2}} L_{1}^{(n+1)}(\omega_{\Pi_{p}}, s)^{-1}$$

$$\int_{\mathbf{Q}_{p}^{\times}} \left(W_{\pi_{p,1}}^{\mathrm{sph}} \boxtimes W_{\pi_{p,2}}^{\mathrm{sph}} \right) \left(\begin{bmatrix} a \\ 1 \end{bmatrix}, \begin{bmatrix} ap^{n} \\ 1 \end{bmatrix} \right) \ |a|_{p}^{s-1} \ d^{\times}a = L(\Pi_{p}, s) \cdot p^{\frac{-n}{2}} L_{2}^{(n+1)}(\omega_{\Pi_{p}}, s)^{-1}$$

Proof. The proof is a direct algebraic computation in the same style as [Gro20] Lemma 3.3.2, once again again using [Shi76] to express the values of $W_{\pi_{p,i}}^{\text{sph}}([a_1])$ as polynomials in the Satake parameters. \Box

We now have all the tools needed to evaluate the linear form Λ_{Π_p} on $f \in C_c^{\infty}(P_p \setminus G_p/G_p^{\circ})$. By Iwasawa decomposition, we may clearly assume that f is given by a characteristic function ch $(P_p z_0(\gamma_0, 1)G_p^{\circ})$ with $z_0 \in Z_p$ and $\gamma_0 \in \operatorname{GL}_2(\mathbf{Q}_p)$. We recall that $\mathcal{S}_{p,1}$ and $\mathcal{T}_{p,1}$ denote the spherical Hecke operators in $\mathcal{H}_{G_p}^{\circ}$, corresponding to the elements $([p_p], 1)$ and $([p_1], 1)$ respectively. Similarly, $\mathcal{S}_{p,2}$ and $\mathcal{T}_{p,2}$ denote the Hecke operators corresponding to $(1, [p_p])$ and $(1, [p_1])$ respectively. Additionally, \mathcal{S}_p will once again denote $\mathcal{S}_{p,1}\mathcal{S}_{p,2}$. We write $\mathfrak{s}_n(x, y)$ for the Schur polynomial $(x^{n+1} - y^{n+1})/(x - y)$, and $\mathfrak{s}_n^{\circ}(X, Y)$ for the polynomial that satisfies

$$\Theta_{\pi_{p,i}}\left(\mathfrak{s}_{n}^{\circ}(\mathcal{S}_{p,i},\mathcal{T}_{p,i})\right) = \mathfrak{s}_{n}\left(\alpha_{\pi_{p,i}},\beta_{\pi_{p,i}}\right), \ i \in \{1,2\}$$

Finally, we put $f_{z_0,\gamma_0} := \operatorname{ch}\left(P_p z_0(\gamma_0,1)G_p^\circ\right)$ in $C_c^\infty(P_p \setminus G_p/G_p^\circ)$, with

$$z_0 = \begin{bmatrix} p^{r_0} \\ p^{r_0} \end{bmatrix} \in Z_p, \ \gamma_0 = \begin{bmatrix} p^{r_1} \\ p^{r_1} \end{bmatrix} \begin{bmatrix} p^m & y \\ 1 \end{bmatrix} k \in \operatorname{GL}_2(\mathbf{Q}_p) = Z_p P_p \operatorname{GL}_2(\mathbf{Z}_p)$$

Proposition 4.3.6. With notation as above, we have

$$\Lambda_{\Pi_p}(f_{z_0,\gamma_0}) = \Theta_{\Pi_p}\left(\mathscr{V}_{z_0,\gamma_0}\left[\mathscr{P}_{\gamma_0} + \mathscr{Q}_{\gamma_0}\mathcal{P}_p(1)\right]\right)$$

where:

$$\begin{split} \mathscr{V}_{z_{0},\gamma_{0}} &:= \operatorname{vol}_{\operatorname{GL}_{2}(\mathbf{Q}_{p})}(P_{p}^{\circ}\gamma_{0}\operatorname{GL}_{2}(\mathbf{Z}_{p})) \cdot \mathcal{S}_{p}^{r_{0}}\mathcal{S}_{p,1}^{r_{1}} \\ \mathscr{P}_{\gamma_{0}} &:= p^{\frac{-m}{2}} \begin{cases} \mathcal{P}_{p,1}^{(m+1)}(1), & \text{if } m \geq 0 \\ \mathcal{P}_{p,2}^{(-m+1)}(1), & \text{if } m < 0 \end{cases} \\ \mathscr{Q}_{\gamma_{0}} &:= p^{\frac{-m}{2}} \begin{cases} \sum_{i=0}^{-v_{p}(y)-1} \varepsilon_{i}(y)\mathfrak{s}_{i+m}^{\circ}(\mathcal{S}_{p,1},\mathcal{T}_{p,1})\mathfrak{s}_{i}^{\circ}(\mathcal{S}_{p,2},\mathcal{T}_{p,2}), & \text{if } m \geq 0, v_{p}(y) < 0 \\ \sum_{i=0}^{-v_{p}(y)-1} \varepsilon_{i}(y)\mathfrak{s}_{i}^{\circ}(\mathcal{S}_{p,1},\mathcal{T}_{p,1})\mathfrak{s}_{i-m}^{\circ}(\mathcal{S}_{p,2},\mathcal{T}_{p,2}), & \text{if } m < 0, v_{p}(y) < 0 \\ 0, & \text{if } v_{p}(y) \geq 0. \end{cases} \\ \varepsilon_{i}(y) &:= \begin{cases} -1, & \text{if } 0 \leq i < -v_{p}(y) - 1 \\ \frac{-1}{p-1}, & \text{if } i = -v_{p}(y) - 1. \end{cases} \end{split}$$

Proof. Using Lemma 4.1.1, we once again unfold $\Lambda_{\Pi_p}(s; f_{z_0,\gamma_0})$ as

where the second equality follows from the following fact: As a function of $z \in Z_p$, $f_{z_0,\gamma_0}((z\gamma, z))$ is equal to $f_{1,\gamma_0}(\gamma, 1)$ on $z_0Z_p^{\circ}$, and zero everywhere else. Similarly, as a function of $\gamma \in \operatorname{GL}_2(\mathbf{Q}_p)$, $f_{1,\gamma_0}(\gamma, 1)$ is equal to 1 on $P_p^{\circ}\gamma_0 \operatorname{GL}_2(\mathbf{Z}_p)$, and zero everywhere else. We now claim that the innermost integral in (8) is independent of $\gamma \in P_p^{\circ}\gamma_0 \operatorname{GL}_2(\mathbf{Z}_p)$ and is completely determined by its value on γ_0 . Indeed, let γ be an element of $P_p^{\circ}\gamma_0 \operatorname{GL}_2(\mathbf{Z}_p)$ written as $\begin{bmatrix} x_1 & x_2 \\ & 1 \end{bmatrix} \gamma_0 k$, where by definition, $x_1 \in \mathbf{Z}_p^{\times}$ and $x_2 \in \mathbf{Z}_p$. Then the innermost integral in (8) is given by

$$\int_{\mathbf{Q}_p^{\times} \cap \mathbf{Z}_p} \psi(ax_2) \left(W_{\pi_{p,1}}^{\mathrm{sph}} \boxtimes W_{\pi_{p,2}}^{\mathrm{sph}} \right) \left(\begin{bmatrix} ax_1 \\ 1 \end{bmatrix} \gamma_0, \begin{bmatrix} a \\ 1 \end{bmatrix} \right) \ |a|_p^{s-1} \ d^{\times} a.$$

Using the fact that ψ has conductor one, $x_2 \in \mathbf{Z}_p$, $x_1 \in \mathbf{Z}_p^{\times}$ and a change of variables $a \rightsquigarrow ax_1$, we see that the claim holds. We thus have

(9)

$$\Lambda_{\Pi_{p}}(s; f_{z_{0}, \gamma_{0}}) = \omega_{\Pi_{p}}(z_{0}) |\det(z_{0})|_{p}^{s} \operatorname{vol}_{\mathrm{GL}_{2}(\mathbf{Q}_{p})} \left(P_{p}^{\circ} \gamma_{0} \operatorname{GL}_{2}(\mathbf{Z}_{p})\right) \int_{\mathbf{Q}_{p}^{\times}} \left(W_{\pi_{p,1}}^{\mathrm{sph}} \boxtimes W_{\pi_{p,2}}^{\mathrm{sph}}\right) \left(\begin{bmatrix}a_{1}\\1\end{bmatrix}\gamma_{0}, \begin{bmatrix}a_{1}\\1\end{bmatrix}\right) |a|_{p}^{s-1} d^{\times}a.$$

The integral appearing in (9) is treated in a very similar way to ([Gro24] 6.3.1), once again using [Shi76] to evaluate the spherical Whittaker functions. Writting $C_{\gamma_0} := \omega_{\Pi_p} \left(\begin{bmatrix} p^{r_1} \\ p^{r_1} \end{bmatrix}, 1 \right)$, this approach together

with Lemma 4.3.5, allows us to split the integral into a "main" term of the form

$$\begin{cases} C_{\gamma_0} L(\Pi_p, s) \cdot p^{\frac{-m}{2}} L_1^{(m+1)}(\omega_{\Pi_p}, s)^{-1}, & \text{if } m \ge 0\\ \\ C_{\gamma_0} L(\Pi_p, s) \cdot p^{\frac{-m}{2} + ms} L_2^{(-m+1)}(\omega_{\Pi_p}, s)^{-1}, & \text{if } m < 0 \end{cases}$$

and an "error" term of the form

$$\begin{cases} C_{\gamma_0} \sum_{i=0}^{-v_p(y)-1} \left(\int_{p^i \mathbf{Z}_p^{\times}} \psi(ay) - 1 \ d^{\times}a \right) \cdot p^{\frac{-m}{2}} \mathfrak{s}_{i+m}(\alpha_{\pi_{p,1}}, \beta_{\pi_{p,1}}) \mathfrak{s}_i(\alpha_{\pi_{p,2}}, \beta_{\pi_{p,2}}) p^{-is}, & \text{if } m \ge 0\\ C_{\gamma_0} \sum_{i=0}^{-v_p(y)-1} \left(\int_{p^i \mathbf{Z}_p^{\times}} \psi(ay) - 1 \ d^{\times}a \right) \cdot p^{\frac{-m}{2} + ms} \mathfrak{s}_i(\alpha_{\pi_{p,1}}, \beta_{\pi_{p,1}}) \mathfrak{s}_{-m+i}(\alpha_{\pi_{p,2}}, \beta_{\pi_{p,2}}) p^{-is}, & \text{if } m < 0. \end{cases}$$

All it remains to do is to use the expression found in ([Gro24] 6.3.1) for the integrals involving ψ , combine these two terms with (9), apply $\lim_{s\to 0} \frac{(-)}{L(\Pi_p,s)}$ and pass to the spherical Hecke algebra through Θ_{Π_p} .

Lemma 4.3.7. With notation as in Proposition 4.3.6, the elements $\mathscr{V}_{z_0,\gamma_0}\left[\mathscr{P}_{\gamma_0} + \mathscr{Q}_{\gamma_0}\mathcal{P}_p(1)\right]$ are contained in $\mathcal{H}^{\circ}_{G_p}(\mathbf{Z}[1/p])$ for any f_{z_0,γ_0} .

Proof. There is a number of things we need to check. Firstly, we need to make sure that the operators \mathscr{P}_{γ_0} and \mathscr{Q}_{γ_0} involve no odd powers of $p^{\frac{-1}{2}}$. Indeed, this follows from a simple parity consideration on m (and i for \mathscr{Q}_{γ_0}), using the expansion of the polynomials $\mathfrak{s}_n(x,y)$ found in [Gro24] Section 6.3, together with the well-known fact that $\Theta_{\Pi_p}(\mathcal{S}_{p,i}) = \alpha_{\pi_{p,i}}\beta_{\pi_{p,i}}$ and $\Theta_{\Pi_p}(\mathcal{T}_{p,i}) = p^{\frac{1}{2}}(\alpha_{\pi_{p,i}} + \beta_{\pi_{p,i}})$. Hence \mathscr{P}_{γ_0} and \mathscr{Q}_{γ_0} are elements of $\mathcal{H}^{\circ}_{G_p}(\mathbf{Z}[1/p])$ and $\mathcal{H}^{\circ}_{G_p}(\frac{1}{p-1}\mathbf{Z}[1/p])$ respectively. It follows from a direct computation that the Euler factor $\mathcal{P}_p(1)$ is also an element of $\mathcal{H}^{\circ}_{G_p}(\mathbf{Z}[1/p])$. We now claim that for every γ_0 ,

$$\operatorname{vol}_{\operatorname{GL}_2(\mathbf{Q}_p)}(P_p^{\circ}\gamma_0\operatorname{GL}_2(\mathbf{Z}_p))\mathscr{Q}_{\gamma_0}$$

is an element of $\mathcal{H}_{G_p}^{\circ}(\mathbf{Z}[1/p])$. By Cartan decomposition, γ_0 is contained in $z_1 \operatorname{GL}_2(\mathbf{Z}_p) \begin{bmatrix} p^{\lambda} \\ 1 \end{bmatrix} \operatorname{GL}_2(\mathbf{Z}_p)$, with $z_1 \in \mathbb{Z}_p$ and $\lambda \in \mathbf{Z}_{\geq 0}$, uniquely determined by γ_0 . We now recall the decomposition

(10)
$$\operatorname{GL}_{2}(\mathbf{Z}_{p})\begin{bmatrix}p^{\lambda}\\ & 1\end{bmatrix}\operatorname{GL}_{2}(\mathbf{Z}_{p}) = \left(\bigsqcup_{\beta \in \mathbf{Z}/p^{\lambda}\mathbf{Z}}\begin{bmatrix}p^{\lambda}&\beta\\ & 1\end{bmatrix}\operatorname{GL}_{2}(\mathbf{Z}_{p})\right) \sqcup \left(\bigsqcup_{\substack{0 < i < \lambda\\\beta \in (\mathbf{Z}/p^{i}\mathbf{Z})^{\times}}}\begin{bmatrix}p^{i}&\beta\\ & p^{\lambda-i}\end{bmatrix}\operatorname{GL}_{2}(\mathbf{Z}_{p})\right) \sqcup \begin{bmatrix}1\\ & p^{\lambda}\end{bmatrix}\operatorname{GL}_{2}(\mathbf{Z}_{p}).$$

If γ_0 is contained in $z_1 \begin{bmatrix} p^{\lambda} & \beta_0 \\ 1 \end{bmatrix} \operatorname{GL}_2(\mathbf{Z}_p)$ or $z_1 \begin{bmatrix} 1 \\ p^{\lambda} \end{bmatrix} \operatorname{GL}_2(\mathbf{Z}_p)$, then by Proposition 4.3.6, we see that $\mathscr{Q}_{\gamma_0} = 0$ hence the claim holds trivially. If on the other hand γ_0 is contained in $z_1 \begin{bmatrix} p^i & \beta_0 \\ p^{\lambda-i} \end{bmatrix} \operatorname{GL}_2(\mathbf{Z}_p)$ with β_0 in $(\mathbf{Z}/p^i \mathbf{Z})^{\times}$, then $\gamma_0 = z_1 \begin{bmatrix} p^{\lambda-i} \\ p^{\lambda-i} \end{bmatrix} \begin{bmatrix} p^{2i-\lambda} & \beta_0 p^{i-\lambda} \\ 1 \end{bmatrix} k$ with $i - \lambda < 0$ and $k \in \operatorname{GL}_2(\mathbf{Z}_p)$. Thus in this case, Proposition 4.3.6 shows that \mathscr{Q}_{γ_0} is non-zero and contained in $\mathcal{H}^o_{G_p}(\frac{1}{p-1}\mathbf{Z}[1/p])$. However in this case, it is straightforward to see that

$$P_p^{\circ}\gamma_0\operatorname{GL}_2(\mathbf{Z}_p) = z_1 P_p^{\circ} \begin{bmatrix} p^i & \beta_0 \\ p^{\lambda-i} \end{bmatrix} \operatorname{GL}_2(\mathbf{Z}_p) = \bigsqcup_{\beta \in (\mathbf{Z}/p^i \mathbf{Z})^{\times}} z_1 \begin{bmatrix} p^i & \beta \\ p^{\lambda-i} \end{bmatrix} \operatorname{GL}_2(\mathbf{Z}_p).$$

Thus $\operatorname{vol}_{\operatorname{GL}_2(\mathbf{Q}_p)}(P_p^{\circ}\gamma_0\operatorname{GL}_2(\mathbf{Z}_p)) = \#(\mathbf{Z}/p^i\mathbf{Z})^{\times}$, which is a multiple of p-1.

Corollary 4.3.8. Let $\Pi_p = \pi_{p,1} \boxtimes \pi_{p,2}$ with $\pi_{p,i}$ unramified H_p -representations of Whittaker type with $\omega_{\pi_{p,1}} \neq \omega_{\pi_{p,2}}^{-1}$. Suppose that Θ_{Π_p} restricts to a morphism $\mathcal{H}^{\circ}_{G_p}(\mathcal{R}) \to \mathcal{R}$ for some $\mathbb{Z}[1/p]$ -algebra $\mathcal{R} \subseteq \mathbb{C}$. Then Λ_{Π_p} induces a non-zero Hecke equivariant morphism

$$\Lambda_{\Pi_p}: C_c^{\infty}(P_p \backslash G_p / G_p^{\circ}, \mathcal{R}) \longrightarrow \mathcal{R} \cdot W^{\mathrm{sph}}_{\Pi_p^{\vee}}$$

Proof. Hecke equivariance is part of Proposition 4.2.5 using the assumption on Π_p . The result then follows at once using Lemma 4.3.7 and the assumption on Θ_{Π_p} .

4.4. Inverting $1 - S_p$. The Hecke action by $1 - S_p$ induces an injective endomorphism of $C_c^{\infty}(P_p \setminus G_p/G_p^{\circ})$. Thus, the localisation $C_c^{\infty}(P_p \setminus G_p/G_p^{\circ})_{1-S_p}$ is canonically isomorphic to

$$C_c^{\infty}(P_p \backslash G_p / G_p^{\circ})[\frac{1}{1 - \mathcal{S}_p}] := \operatorname{coker}(C_c^{\infty}(P_p \backslash G_p / G_p^{\circ})[x] \xrightarrow{1 - (1 - \mathcal{S}_p)x} C_c^{\infty}(P_p \backslash G_p / G_p^{\circ})[x])$$

where $C_c^{\infty}(P_p \setminus G_p/G_p^{\circ})[x]$ denotes the natural $\mathcal{H}_{G_p}^{\circ}[x]$ -module of formal polynomials in a single variable x with coefficients in $C_c^{\infty}(P_p \setminus G_p/G_p^{\circ})$, or equivalently the tensor product $\mathcal{H}_{G_p}^{\circ}[x] \otimes_{\mathcal{H}_{G_p}^{\circ}} C_c^{\infty}(P_p \setminus G_p/G_p^{\circ})$. For this reason, it is clear that $C_c^{\infty}(P_p \setminus G_p/G_p^{\circ})$ embeds into its localisation $C_c^{\infty}(P_p \setminus G_p/G_p^{\circ})[\frac{1}{1-S_p}]$. As usual, we consider representations $\Pi_p = \pi_{p,1} \boxtimes \pi_{p,2}$ where $\pi_{p,i}$ are unramified of Whittaker type. From Proposition 4.3.6, and the density of this family of representations in $\operatorname{Spec}(\mathcal{H}_{G_p}^{\circ})$, we can deduce that for each function f in $C_c^{\infty}(P_p \setminus G_p/G_p^{\circ})$, there exists a unique Hecke operator which we call \mathcal{Q}_f in $\mathcal{H}_{G_p}^{\circ}$, for which $\Lambda_{\Pi_p}(f) = \Theta_{\Pi_p}(\mathcal{Q}_f)$ for every such Π_p . Proposition 4.2.5 then implies that

(11)
$$\Lambda_{\Pi_p}((1-\mathcal{S}_p)\cdot f) = \Theta_{\Pi_p}(-\mathcal{S}_p^{-1}(1-\mathcal{S}_p)\mathcal{Q}_f)) = \Lambda_{\Pi_p}(-\mathcal{S}_p\mathcal{Q}_f'\cdot f_0).$$

Theorem 4.4.1. The module $C_c^{\infty}(P_p \setminus G_p/G_p^{\circ})[\frac{1}{1-S_p}]$ is free of rank one over $\mathcal{H}_{G_p}^{\circ}[\frac{1}{1-S_p}]$, generated by the characteristic function $f_0 := \operatorname{ch}(P_p G_p^{\circ})$. In particular, for $f \in C_c^{\infty}(P_p \setminus G_p/G_p^{\circ})$, we have $f = \frac{-S_p Q_f'}{1-S_p} \cdot f_0$, where Q_f is the Hecke operator described above.

Proof. The proof is long and quite technical, so we give it in Appendix A in order to ease exposition. \Box

Remark 4.4.2. Theorem 4.4.1, strictly speaking, is not necessary for our proof of optimal integral norm relations, however it definitely makes it cleaner and more complete. In addition, it showcases the obstruction regarding the invertibility of the operators $1 - S_p$. In fact, later on we will see that Theorem 4.4.1 is simply not true without the localisation. We also like to stress the fact that Theorem 4.4.1 by no means replaces the need for Theorem 3.2.1 and the two results are not equivalent.

Corollary 4.4.3. The module $C_c^{\infty}(P_p \setminus G_p/G_p^{\circ}, \mathbb{Z}[1/p])[\frac{1}{1-S_p}]$ is free of rank one over $\mathcal{H}_{G_p}^{\circ}(\mathbb{Z}[1/p])[\frac{1}{1-S_p}]$ generated by $\operatorname{ch}(P_p G_p^{\circ})$.

Proof. This follows from (11), Lemma 4.3.7 and Theorem 4.4.1.

Definition 4.4.4. Let $f_0 = \operatorname{ch}(P_p G_p^\circ)$. For $f \in C_c^\infty(P_p \setminus G_p/G_p^\circ)[\frac{1}{1-S_p}]$, the Hecke operator $\mathcal{P}_{f,\operatorname{loc}}$ is the unique element of $\mathcal{H}_{G_p}^\circ[\frac{1}{1-S_p}]$ that satisfies $\mathcal{P}_{f,\operatorname{loc}} \cdot f_0 = f$.

Such a unique Hecke operator clearly exists by Theorem 4.4.1. We put

$$C_c^{\infty,1}(P_p \backslash G_p / G_p^{\circ}, \mathbf{Z}[1/p]) = \left\{ f \in C_c^{\infty}(P_p \backslash G_p / G_p^{\circ}, \mathbf{Z}[1/p]) \mid f \text{ is valued in } \begin{cases} (p-1)\mathbf{Z}[1/p], \text{ on } Z_p(T_p \times \{1\}) \\ \mathbf{Z}[1/p], \text{ everywhere else.} \end{cases} \right\}$$

This very much resembles the definition of the lattice $\mathcal{L}_1^{H_p,\text{unv}}$ in [Gro24] which plays a similar role in the setup of an unramified maximal torus H embedded in GL₂. We also consider a non-compactly supported version of this which we denote by $C^{\infty,1}(P_p \setminus G_p/G_p^\circ, \mathbf{Z}[1/p])$ and is defined in the same way, simply by dropping the compact support condition.

We write $C_c^{\infty,1}(P_p \setminus G_p/G_p^\circ, \mathbf{Z}[1/p])[\frac{1}{1-S_p}]$ for the $\mathcal{H}_{G_p}^\circ(\mathbf{Z}[1/p])[\frac{1}{1-S_p}]$ -submodule generated by the lattice $C_c^{\infty,1}(P_p \setminus G_p/G_p^\circ, \mathbf{Z}[1/p])$ in $C_c^\infty(P_p \setminus G_p/G_p^\circ, \mathbf{Z}[1/p])[\frac{1}{1-S_p}]$. We now obtain the following main integrality result, on the P_p -invariant side of things.

Theorem 4.4.5. Let f be an element of $C_c^{\infty,1}(P_p \setminus G_p/G_p^\circ, \mathbf{Z}[1/p])[\frac{1}{1-S_p}]$. Then $f = \mathcal{P}_{f,\text{loc}} \cdot f_0$ and the Hecke operator $\mathcal{P}_{f,\text{loc}}$ is contained in the ideal

$$\mathfrak{h}_p = \left(p - 1, \mathcal{P}_p(1)'\right) \subseteq \mathcal{H}^{\circ}_{G_p}(\mathbf{Z}[1/p])[\frac{1}{1 - \mathcal{S}_p}].$$

If moreover $\frac{f}{1-S_p}$ is an element of $C_c^{\infty,1}(P_p \setminus G_p/G_p^{\circ}, \mathbb{Z}[1/p])$, then the containment holds without inverting $1-S_p$.

Proof. We may assume that $f \in C_c^{\infty,1}(P_p \setminus G_p/G_p^\circ, \mathbf{Z}[1/p])$ and that is is given by a characteristic function $c_{z,\gamma} \operatorname{ch}(P_p z(\gamma, 1)G_p^\circ)$ with $z \in Z_p$ and $\gamma \in H_p$. Then $\mathcal{P}_{f,\operatorname{loc}} = c_{z,\gamma} \frac{-\mathcal{S}_p \mathcal{Q}'_I}{1-\mathcal{S}_p}$ where $\mathcal{Q}_f = \mathscr{V}_{z,\gamma} [\mathscr{P}_\gamma + \mathscr{Q}_\gamma \mathcal{P}_p(1)]$. We now proceed in the same style as in the final step of the proof of Lemma 4.3.7. If γ is contained in $y \begin{bmatrix} p^{\lambda} & \beta \\ 1 \end{bmatrix} \operatorname{GL}_2(\mathbf{Z}_p)$ with $y \in Z_p$, $\lambda \geq 0$ and $\beta \in \mathbf{Z}/p^{\lambda}\mathbf{Z}$, then $\mathscr{Q}_\gamma = 0$ and $\operatorname{ch}(P_p z(\gamma, 1)G_p^\circ)$ coincides with $\operatorname{ch}(P_p z(y \begin{bmatrix} p^{\lambda} & \\ 1 \end{bmatrix}, 1)G_p^\circ)$. Thus, by the assumption on the values of f, we must have $c_{z,\gamma} \in (p-1)\mathbf{Z}[1/p]$ and $\mathcal{Q}_f \in (p-1)\mathcal{H}_{G_p}^\circ(\mathbf{Z}[1/p])$. If γ is contained in $y \begin{bmatrix} 1 & \\ p^{\lambda} \end{bmatrix} \operatorname{GL}_2(\mathbf{Z}_p)$ for some $y \in Z_p$ and $\lambda \geq 0$, then the situation is practically the same. Finally if γ is contained in $y \begin{bmatrix} p^i & \beta \\ p^{\lambda-i} \end{bmatrix} \operatorname{GL}_2(\mathbf{Z}_p)$ for some $y \in Z_p$, $0 < i < \lambda, \lambda > 0$ and $\beta \in (\mathbf{Z}/p^i\mathbf{Z})^{\times}$, then as in the last part of the proof of Lemma 4.3.7, using that in this case $\operatorname{vol}_{\operatorname{GL}_2(\mathbf{Q}_p)}(P_p^\circ \gamma \operatorname{GL}_2(\mathbf{Z}_p))$ is a multiple of p-1, we deduce that $\mathscr{V}_{z,\gamma}[\mathscr{P}_\gamma + \mathscr{Q}_\gamma \mathcal{P}_p(1)]$ is an element of $\mathfrak{h}_p \subseteq \mathcal{H}_{G_p}^\circ(\mathbf{Z}[1/p])$ as required. Moreover, if $f = (1 - \mathcal{S}_p) \cdot f_0$ with $f_0 \in C_c^{\infty,1}(P_p \setminus G_p/G_p^\circ, \mathbf{Z}[1/p])$ as in the statement of the theorem, then by freeness, $\mathcal{P}_{f,\operatorname{loc}}$ is given by $(1 - \mathcal{S}_p) \cdot \frac{-\mathcal{S}_p \mathcal{Q}'_{f_0}}{1-\mathcal{S}_p}$ and thus we are done. \Box

5. Euler systems for $GL_2 \times GL_2$

We now finally have all the necessary tools in order to state and prove our optimal abstract norm relations at a local integral level and use them in order to obtain Theorem A of the introduction.

5.1. Local abstract integral norm relations. Let V be a smooth G_p -representation and let $\mathfrak{Z} : \mathcal{S}(\mathbf{Q}_p^2) \otimes \mathcal{H}(G_p) \to V$ be any $(G_p \times H_p)$ -equivariant map as in Definition 2.2.1. Consider the induced diagram

$$\mathcal{I}(G_p/G_p^{\circ}[p], \mathbf{Z}[1/p]) \xrightarrow{\mathrm{Tr}\downarrow} \mathcal{I}(G_p/G_p^{\circ}, \mathbf{Z}[1/p]) \xrightarrow{3} V$$

It follows straight from the definitions that the horizontal map is equivariant with respect to $\mathcal{H}_{G_p}^{\circ}$. Furthermore, for $\delta \in \mathcal{I}(G_p/G_p^{\circ}[p], \mathbb{Z}[1/p])$, we have (using the "norm" notation of [LSZ21b])

$$\mathfrak{Z}\left(\mathrm{Tr}\left(\delta\right)\right) = \mathrm{norm}_{G_{p}^{\circ}}^{G_{p}^{\circ}[p]}\left(\mathfrak{Z}\left(\delta\right)\right) := \sum_{\gamma \in G_{p}^{\circ}/G_{p}^{\circ}[p]} \gamma \cdot \mathfrak{Z}\left(\delta\right).$$

Finally, we write δ_0 for the unramified vector $\operatorname{ch}(\mathbf{Z}_p^2) \otimes \operatorname{ch}(G_p^\circ)$, which implies that $\operatorname{norm}_{G_p^\circ}^{G_p^\circ[p]}(\mathfrak{Z}(\delta_0))$ is nothing more than $(p-1) \cdot \mathfrak{Z}(\delta_0)$. Recall that $\mathcal{S}_0(\mathbf{Q}_p^2)$ denotes the space of Schwartz functions that vanish at the origin and $\mathcal{I}_0(G_p/G_p^\circ)$ denotes the submodule of $\mathcal{I}(G_p/G_p^\circ)$ given by the image of $\mathcal{S}_0(\mathbf{Q}_p^2) \otimes C_c^\infty(G_p/G_p^\circ)$ down to the H_p -coinvariants. Similarly, recall the definitions of $\mathcal{I}_0(G_p/G_p^\circ, \mathbf{Z}[1/p])$ and $\mathcal{I}_0(G_p/G_p^\circ[p], \mathbf{Z}[1/p])$ given in Section 2.2

Theorem 5.1.1 (Integral abstract norm-relations). Let V and \mathfrak{Z} be as above. Then:

- (1) For $\delta \in \mathcal{I}(G_p/G_p^\circ, \mathbb{Z}[1/p])$, we have $\mathcal{P}_{\delta} \in \mathcal{H}_{G_n}^\circ(\mathbb{Z}[1/p])$ and $\mathfrak{Z}(\delta) = \mathcal{P}_{\delta} \cdot \mathfrak{Z}(\delta_0)$.
- (2) For $\delta \in \mathcal{I}_0(G_p/G_p^{\circ}[p], \mathbf{Z}[1/p])$, we have

$$\mathcal{P}_{\mathrm{Tr}(\delta)} \in \left(p-1, \mathcal{P}_p(1)'\right) \subseteq \mathcal{H}^{\circ}_{G_p}(\mathbf{Z}[1/p]) \quad and \quad \mathrm{norm}_{G^{\circ}_p}^{G^{\circ}_p[p]}(\mathfrak{Z}(\delta)) = \mathcal{P}_{\mathrm{Tr}(\delta)} \cdot \mathfrak{Z}(\delta_0).$$

(3) For $\delta \in \mathcal{I}(G_p/G_p^{\circ}[p], \mathbb{Z}[1/p])$, we have $\mathcal{P}_{\mathrm{Tr}(\delta)} \in \left(p-1, \mathcal{P}'_p(1)\right) \subseteq \mathcal{H}^{\circ}_{G_p}(\mathbb{Z}[1/p])[\frac{1}{1-\mathcal{S}_p}]$ and $\mathcal{P}_{\mathrm{Tr}(\delta)}$ has degree at most one as a polynomial in $\frac{1}{1-\mathcal{S}_p}$. If furthermore $1-\mathcal{S}_p$ acts invertibly as a linear endomorphism of $V^{G_p^{\circ}}$, then the norm relation of part two is also satisfied.

This theorem can be interpreted as the local integral essence of the tame norm relations in motivic cohomology for integral test data, as presented later on in Theorem 5.2.7

Proof. Let $\delta \in \mathcal{I}(G_p/G_p^{\circ}, \mathbb{Z}[1/p])$. By Proposition 3.3.2 and Theorem 3.2.1,

$$\mathcal{P}_{\delta} \cdot \operatorname{ch}(P_p G_p^{\circ}) = \Xi_c(\delta) \in C_c^{\infty}(P_p \backslash G_p / G_p^{\circ}, \mathbf{Z}[1/p]).$$

By Corollary 4.4.3, we deduce that $\mathcal{P}_{\delta} = \mathcal{P}_{\Xi_c(\delta), \text{loc}} \in \mathcal{H}^{\circ}_{G_p}(\mathbf{Z}[1/p])[\frac{1}{1-S_p}] \cap \mathcal{H}^{\circ}_{G_p}$, and thus in particular that $\mathcal{P}_{\delta} \in \mathcal{H}^{\circ}_{G_p}(\mathbf{Z}[1/p])$. This proves the first part when combined with Hecke equivariance of \mathfrak{Z} . For the other two parts, we need some more work. One checks that we obtain a commutative diagram

- r 1

$$\begin{split} \mathcal{I}(G_p/G_p^{\circ}[p]) & \xrightarrow{\Xi[p]} C^{\infty}(P_p \setminus G_p/G_p^{\circ}[p]) & \phi \otimes \xi \longmapsto \longrightarrow \int_{H_p} \xi(h(-)) f_{\phi}(h) \ dh \\ & \underset{Tr}{} \downarrow & \downarrow \\ \mathcal{I}(G_p/G_p^{\circ}) \xrightarrow{\Xi} C^{\infty}(P_p \setminus G_p/G_p^{\circ}) & \phi \otimes \sum_{\gamma \in G_p^{\circ}/G_p^{\circ}[p]} \xi((-)\gamma) \longmapsto \int_{H_p} \sum_{\gamma \in G_p^{\circ}/G_p^{\circ}[p]} \xi(h(-)\gamma) f_{\phi}(h) \ dh \end{split}$$

where $\Xi[p]$ is defined in the same way as Ξ , simply by replacing G_p° with $G_p^{\circ}[p]$, and the trace maps are the usual ones with respect to $G_p^{\circ}[p] \subseteq G_p^{\circ}$. We now take $\delta \in \mathcal{I}(G_p/G_p^{\circ}[p], \mathbf{Z}[1/p])$. The proof of Proposition 3.3.2 goes through in the same way at level $G_p^{\circ}[p]$ and thus $\Xi[p](\delta) \in C^{\infty}(P_p \setminus G_p/G_p^{\circ}[p], \mathbf{Z}[1/p])$. It is easy to check that for an element in $C^{\infty}(P_p \setminus G_p/G_p^{\circ}[p], \mathbf{Z}[1/p])$, its image under the trace map to $C^{\infty}(P_p \setminus G_p/G_p^{\circ}, \mathbf{Z}[1/p])$ is valued in $(p-1)\mathbf{Z}[1/p]$ on $Z_p(T_p \times \{1\})$. This is because for $g \in Z_p(T_p \times \{1\})$, $P_pgG_p^{\circ}[p] = P_pgG_p^{\circ}$ and hence Tr is simply multiplication by $[G_p^{\circ}: G_p^{\circ}[p]] = p-1$ on such a characteristic function. Thus $\mathrm{Tr}(\Xi[p](\delta))$ is contained in $C^{\infty,1}(P_p \setminus G_p/G_p^{\circ}, \mathbf{Z}[1/p])$. By commutativity of the diagram above, we deduce that $\Xi(\mathrm{Tr}(\delta))$ belongs in $C^{\infty,1}(P_p \setminus G_p/G_p^{\circ}, \mathbf{Z}[1/p])$ and hence $\Xi_c(\mathrm{Tr}(\delta))$ belongs in $C_c^{\infty,1}(P_p \setminus G_p, G_p^{\circ}, \mathbf{Z}[1/p])$ and hence $\Xi_c(\mathrm{Tr}(\delta))$ belongs in $C_c^{\infty,1}(P_p \setminus G_p, G_p^{\circ}, \mathbf{Z}[1/p])$ and hence $\Xi_c(\mathrm{Tr}(\delta))$ belongs in $C_c^{\infty,1}(P_p \setminus G_p, G_p^{\circ}, \mathbf{Z}[1/p])$. Thus, by Theorem 4.4.1 and Theorem 4.4.5, $\mathcal{P}_{\mathrm{Tr}(\delta)} = \mathcal{P}_{\Xi_c(\mathrm{Tr}(\delta)),\mathrm{loc}}$ is an element of the ideal $\left(p-1, \mathcal{P}_p(1)'\right) \subseteq \mathcal{H}_{G_p}^{\circ}(\mathbf{Z}[1/p])[\frac{1}{1-\mathcal{S}_p}]$ and has degree at most one as a polynomial in $\frac{1}{1-\mathcal{S}_p}$. This proves the third part of the theorem. For the second part, if $\delta \in \mathcal{I}_0(G_p/G_p^{\circ}, \mathbf{Z}[1/p])$ then $\mathrm{Tr}(\delta) \in \mathcal{I}_0(G_p/G_p^{\circ}, \mathbf{Z}[1/p])$. We have already remarked that for an element in the latter, its image under Ξ is already compactly supported. Hence $\Xi_c(\mathrm{Tr}(\delta))$ is divisible by $1-\mathcal{S}_p$ in $C_c^{\infty,1}(P_p \setminus G_p/G_p^{\circ}, \mathbf{Z}[1/p])$. The second part of the theorem now follows from the "moreover" part of Theorem 4.4.5.

5.2. Cohomology classes and norm compatibility. We now showcase how we may use our representation theoretic results to obtain norm-compatible classes in the cohomology of the infinite level Shimura variety Y_G . This approach was introduced in [LSZ21b] and thus we draw heavily from *op.cit*.

5.2.1. Modular varieties and motivic cohomology. We follow the setup and conventions of [LSZ21a] and we write $Y_{\text{GL}_2} = \varprojlim_U Y_{\text{GL}_2}(U)$ for the infinite level modular (Shimura) curve regarded as a pro-variety over \mathbf{Q} with a right action of $\text{GL}_2(\mathbf{A}_f)$, whose **C**-points are $\text{GL}_2^+(\mathbf{Q}) \setminus (\text{GL}_2(\mathbf{A}_f) \times \mathbf{H})$ (as usual **H** denotes the upper-half plane). Similarly, we define Y_G as $Y_{\text{GL}_2} \times Y_{\text{GL}_2}$ which is the Shimura variety associated to G. The compatibility of our Shimura data, induces a morphism of **Q**-varieties

$$Y_{\mathrm{GL}_2}(\mathrm{GL}_2(\mathbf{A}_f) \cap gUg^{-1}) \xrightarrow{\iota_{gU}} Y_G(U)$$

$$\uparrow \simeq$$

$$Y_G(gUg^{-1})$$

for any $U \subseteq G(\mathbf{A}_f)$ open compact and $g \in G(\mathbf{A}_f)/U$. If we temporarily let Γ denote any of the groups $\{\mathrm{GL}_2, G\}$, then the motivic cohomology of Y_{Γ} is given by

$$H^{i}_{\mathrm{mot}}(Y_{\Gamma}, \mathbf{Q}(j)) := \varinjlim_{U \text{ open compact}} H^{i}_{\mathrm{mot}}(Y_{\Gamma}(U), \mathbf{Q}(j))$$

where at each finite level U the groups $H^i_{\text{mot}}(Y_{\Gamma}(U), \mathbf{Q}(j))$ are defined in the sense of [Voe00] and [FV00]. The spaces $H^i_{\text{mot}}(Y_{\Gamma}, \mathbf{Q}(j))$ are naturally smooth **Q**-linear, $\Gamma(\mathbf{A}_f)$ -representations.

Remark 5.2.1. It is worth noting that everything in this section extends to the case of "non-trivial coefficient sheaves". See for example [LSZ21b], [LSZ21a] and [Gro20] for some of the formalisms required to extend the theory.

By functoriality of motivic cohomology (with a shift in bi-degree) the map ι_{qU} induces a morphism

$$\iota_{gU,*}: H^1_{\mathrm{mot}}(Y_{\mathrm{GL}_2}(\mathrm{GL}_2(\mathbf{A}_f) \cap gUg^{-1}), \mathbf{Q}(1)) \longrightarrow H^3_{\mathrm{mot}}(Y_G(U), \mathbf{Q}(2)).$$

In the same way as ([LSZ21b] 9.1) and ([LSZ21a] 8.2), we have:

$$\iota_*: H^1_{\mathrm{mot}}(Y_{\mathrm{GL}_2}, \mathbf{Q}(1)) \otimes_{\mathbf{Q}} \mathcal{H}(G(\mathbf{A}_f), \mathbf{Q}) \longrightarrow H^3_{\mathrm{mot}}(Y_G, \mathbf{Q}(2))$$

characterized as follows: If $U \subseteq G(\mathbf{A}_f)$ is an open compact subgroup, $g \in G(\mathbf{A}_f)$ and x is an element of $H^1_{\text{mot}}(Y_{\text{GL}_2}(\text{GL}_2(\mathbf{A}_f) \cap gUg^{-1}), \mathbf{Q}(1))$, then

$$\iota_*(x \otimes \operatorname{ch}(gU)) = \operatorname{vol}_{\operatorname{GL}_2(\mathbf{A}_f)} \left(\operatorname{GL}_2(\mathbf{A}_f) \cap gUg^{-1} \right) \cdot \iota_{gU,*}(x).$$

5.2.2. Eisenstein classes and the Rankin-Eisenstein map.

Theorem 5.2.3. There exists a canonical $GL_2(\mathbf{A}_f)$ -equivariant map

$$S_0(\mathbf{A}_f^2, \mathbf{Q}) \longrightarrow H^1_{\mathrm{mot}}(Y_{\mathrm{GL}_2}, \mathbf{Q}(1)) = \mathcal{O}(Y_{\mathrm{GL}_2})^{\times} \otimes \mathbf{Q}, \ \phi \mapsto g_{\phi}$$

characterized by the following: If $\phi = ch((\alpha, \beta) + N\mathbf{\hat{Z}}^2)$ for some $N \in \mathbf{Z}_{\geq 1}$ and $(\alpha, \beta) \in \mathbf{Q}^2 - N\mathbf{Z}^2$, then $g_{\phi} = g_{\alpha/N,\beta/N}$ is the Siegel unit in the notation of ([Kat04] 1.4).

Proof. This is Theorem 1.8 in [Col04].

Definition 5.2.4. We define the Rankin-Eisenstein map

$$\mathcal{RE}: \mathcal{S}_0(\mathbf{A}_f^2, \mathbf{Q}) \otimes_{\mathbf{Q}} \mathcal{H}(G(\mathbf{A}_f), \mathbf{Q}) \longrightarrow H^3_{\mathrm{mot}}(Y_G, \mathbf{Q}(2)), \ \phi \otimes \xi \mapsto \iota_*(g_\phi \otimes \xi)$$

where ι_* is the map of Proposition 5.2.2.

It is not hard to see that the map \mathcal{RE} is $(\operatorname{GL}_2(\mathbf{A}_f) \times G(\mathbf{A}_f))$ -equivariant in the sense of Definition 2.2.1. The detailed argument in ([Gro20] 6.1) works verbatim.

5.2.3. The classes and norm compatibility. Let S be a fixed finite set of primes containing 2 and write $\mathbf{Q}_S = \prod_{p \in S} \mathbf{Q}_p$. Let $\delta_S \in \mathcal{I}_0(G(\mathbf{Q}_S)/K_S, \mathbf{Z})$ where K_S is some open compact subgroup. These will form the local data at the "bad" primes and will be fixed throughout.

Definition 5.2.5. For the data at primes $p \notin S$, we take any family $\underline{\delta} = (\delta_p)_{p\notin S}$ contained in the product $\prod_{p\notin S} \left(S(\mathbf{Q}_p^2, \mathbf{Q}) \otimes_{\mathbf{Q}} \mathcal{H}(G_p/G_p^\circ, \mathbf{Q}) \right)$. For a square free integer $n \in \mathbf{Z}_{\geq 1}$ coprime to S and $p \notin S$, we define the "n-truncated" element:

$$\delta[n]_p = \begin{cases} \operatorname{ch}(\mathbf{Z}_p^2) \otimes \operatorname{ch}(G_p^\circ), & \text{if } p \nmid n \\ \delta_p, & \text{if } p | n. \end{cases}$$

and we set $\delta[n] := \delta_S \bigotimes_{p \notin S} \delta[n]_p$.

Finally, for such integer n, we write $K_S[n]$ for the open compact subgroup given by the following product: $K_S \times \{g = (g_1, g_2) \in G(\hat{\mathbf{Z}}^S) \mid \det(g_2) \equiv 1 \mod n\}$. We are now ready to define a family of cohomology classes which will depend on $\underline{\delta}$, and implicitly δ_S , however to ease notation, we omit the dependence on δ_S since we fix it throughout. We write \mathscr{Z}_S for the set of square-free positive integers coprime to S.

Definition 5.2.6. Similarly to [LSZ21b], we set

$$\mathcal{Z}_{\text{mot},n}(\underline{\delta}) := \mathcal{RE}(\delta[n]) \in H^3_{\text{mot}}(Y_G(K_S[n]), \mathbf{Q}(2)).$$

We note that for a fixed family $\underline{\delta}$, the class $\mathcal{Z}_{\text{mot},n}(\underline{\delta})$, for any $n \in \mathscr{Z}_S$, only depends on the elements δ_p for primes p|n.

Theorem 5.2.7. For positive integers n|m| let pr_n^m denote the natural map $Y_G(K_S[m]) \to Y_G(K_S[n])$.

(1) If $\underline{\delta} \in \prod_{p \notin S} \mathcal{I}(G_p/G_p^\circ, \mathbb{Z}[1/p])$, then for any $n, m \in \mathscr{Z}_S$ with $\frac{m}{n} = p$ prime:

$$\mathcal{Z}_{\mathrm{mot},m}\left(\underline{\delta}\right) = \mathcal{P}_{\delta_{p}} \cdot \mathcal{Z}_{\mathrm{mot},n}\left(\underline{\delta}\right), \text{ with } \mathcal{P}_{\delta_{p}} \in \mathcal{H}_{G_{p}}^{\circ}(\mathbf{Z}[1/p]).$$

(2) If $\underline{\delta} \in \prod_{p \notin S} \mathcal{I}_0(G_p/G_p^{\circ}[p], \mathbf{Z}[1/p])$, then for any $n, m \in \mathscr{Z}_S$ with $\frac{m}{n} = p$:

(12)
$$(\operatorname{pr}_{n}^{m})_{*}(\mathcal{Z}_{\operatorname{mot},m}(\underline{\delta})) = \mathcal{P}_{\operatorname{Tr}(\delta_{p})} \cdot \mathcal{Z}_{\operatorname{mot},n}(\underline{\delta}), \text{ with } \mathcal{P}_{\operatorname{Tr}(\delta_{p})} \in \left(p-1, \mathcal{P}_{p}^{'}(1)\right) \subseteq \mathcal{H}_{G_{p}}^{\circ}(\mathbf{Z}[1/p]).$$

(3) Specialising $\underline{\delta}$, we have for all $n, m \in \mathscr{Z}_S$ with $\frac{m}{n} = p$ prime

$$(\mathrm{pr}_{n}^{m})_{*}\left(\mathcal{Z}_{\mathrm{mot},m}\left(\underline{\delta}\right)\right) = \begin{cases} (p-1) \cdot \mathcal{Z}_{\mathrm{mot},n}\left(\underline{\delta}\right), & \text{if } \underline{\delta} = \underline{\delta}_{0} \\ \mathcal{P}_{p}^{'}(1) \cdot \mathcal{Z}_{\mathrm{mot},n}\left(\underline{\delta}\right), & \text{if } \underline{\delta} = \underline{\delta}_{1} \end{cases}$$

where
$$\underline{\delta}_0 = \left((p-1) \cdot \operatorname{ch}(\mathbf{Z}_p^2) \otimes \operatorname{ch}(G_p^{\circ}[p])\right)_{p \notin S}$$
, and $\underline{\delta}_1 = \left(n_p \phi_{p,2} \otimes \left(\operatorname{ch}(G_p^{\circ}[p]) - \operatorname{ch}\left(\left(1, \begin{bmatrix} 1 & 1/p \\ 1 & \end{bmatrix}\right) G_p^{\circ}[p]\right)\right)\right)_{p \notin S}$,
 $n_p = \frac{1}{(p-1)^2(p+1)}$ and $\phi_{p,2} = \operatorname{ch}(p^2 \mathbf{Z}_p \times (1+p^2 \mathbf{Z}_p))$

Proof. Firstly note that the data $\delta[np]$ and $\delta[n]$ only differ at the prime p and coincide everywhere else. Write V for $H^3_{\text{mot}}(Y_G, \mathbf{Q}(2))$ regarded as a smooth G_p -representation. It is also clear that we can check the equalities after tensoring with \mathbf{C} and thus we will omit it from our notation. By the global equivariance of the Rankin-Eisenstein map, we can observe that after fixing the input data away from p, the composite map at p

$$\mathcal{RE}_{\delta[m]^{(p)}}: \mathcal{S}(\mathbf{Q}_p^2) \otimes \mathcal{H}(G_p) \longrightarrow \mathcal{S}_0(\mathbf{A}_f^2) \otimes_{\mathbf{Q}} \mathcal{H}(G(\mathbf{A}_f)) \xrightarrow{\mathcal{RE}} V$$

is $(G_p \times H_p)$ -equivariant, where the first map is given by $x_p \mapsto \delta[m]^{(p)} \otimes x_p$. Thus if $\underline{\delta}$ is as in part one,

$$\mathcal{Z}_{\mathrm{mot},m}\left(\underline{\delta}\right) = \mathcal{RE}_{\delta[m]^{(p)}}(\delta_p) = \mathcal{P}_{\delta_p} \cdot \mathcal{RE}_{\delta[m]^{(p)}}(\mathrm{ch}(\mathbf{Z}_p^2) \otimes \mathrm{ch}(G_p^\circ)), \ \mathcal{P}_{\delta_p} \in \mathcal{H}_{G_p}^{\circ}(\mathbf{Z}[1/p]).$$

This follows from the first part of Theorem 5.1.1. But since (p, nS) = 1, $\mathcal{RE}_{\delta[m]^{(p)}}$ evaluated at the unramified vector $\operatorname{ch}(\mathbf{Z}_p^2) \otimes \operatorname{ch}(G_p^\circ)$ is nothing more than $\mathcal{Z}_{\operatorname{mot},n}(\underline{\delta})$ and the first part of the theorem follows. For the second part, we have commutative diagram

where τ_n, τ_m are the natural maps, and the leftmost vertical map is again given by $x_p \mapsto \delta[m]^{(p)} \otimes x_p$ which is well-defined since we are now assuming that $\underline{\delta}$ is as in the second part of the theorem. The commutativity of the right-most square follows from the fact that pushforward acts on cohomology via coset representatives. Thus, by the $(G_p \times H_p)$ -equivariance property of $\mathcal{RE}_{\delta[m]^{(p)}}$ and the second part of Theorem 5.1.1, we have for $\underline{\delta}$ as in the second statement of the theorem,

$$\operatorname{norm}_{G_{p}^{\circ}}^{G_{p}^{\circ}[p]}\left(\mathcal{RE}_{\delta[m]^{(p)}}(\delta_{p})\right) = \mathcal{P}_{\operatorname{Tr}(\delta_{p})} \cdot \mathcal{RE}_{\delta[m]^{(p)}}(\operatorname{ch}(\mathbf{Z}_{p}^{2}) \otimes \operatorname{ch}(G_{p}^{\circ})), \ \mathcal{P}_{\operatorname{Tr}(\delta_{p})} \in \left(p-1, \mathcal{P}_{p}^{'}(1)\right) \subseteq \mathcal{H}_{G_{p}}^{\circ}(\mathbf{Z}[1/p]).$$

The proof of the second statement in the theorem is now complete by following the commutative diagram above and unravelling the definitions. The third and final part follows by specialising our general treatment in Section 4.3 to these special cases of integral elements, with the $\underline{\delta}_0$ case being trivial. One also checks quite easily, that these two families are indeed integral in the sense of Definition 2.2.2.

5.2.4. Base change to cyclotomic fields. Following [LSZ21b] Section 8.2, we can give an alternative description of the classes $\mathcal{Z}_{\text{mot},n}(\underline{\delta})$. We can regard G as a subgroup of $G \times \mathbf{G}_m$ via the embedding $i : g = (g_1, g_2) \mapsto (g, \det(g_2))$. For every $n \in \mathscr{Z}_S$, this gives an open and closed embedding (which we still denote by i)

$$i: Y_G(K_S[n]) \hookrightarrow Y_G(K_S) \times_{\operatorname{Spec}(\mathbf{Q})} \operatorname{Spec}(\mathbf{Q}(\mu_n)).$$

The pushforward along this map gives

$$i_*: H^3_{\text{mot}}\left(Y_G(K_S[n]), \mathbf{Q}(2)\right) \longrightarrow H^3_{\text{mot}}\left(Y_G(K_S) \times_{\text{Spec}(\mathbf{Q})} \text{Spec}(\mathbf{Q}(\mu_n)), \mathbf{Q}(2)\right).$$

which satisfies certain intertwining properties. In particular, for every prime $p \nmid n$, the map i_* satisfies the following intertwining properties with respect to the action of $\mathcal{H}^{\circ}_{G_p}(\mathbf{Z}[1/p])$ on the left and the action of $\mathcal{H}^{\circ}_{G_p}(\mathbf{Z}[1/p])[\operatorname{Gal}(\mathbf{Q}(\mu_n)/\mathbf{Q})]$ on the right: For $x \in H^3_{\operatorname{mot}}(Y_G(K_S[n]), \mathbf{Q}(2))$, we have

(13)
$$i_*(\theta \cdot x) = \begin{cases} \theta \cdot i_*(x) &, \text{ if } \theta = \mathcal{S}_{p,1} \\ \theta \cdot i_*(x) &, \text{ if } \theta = \mathcal{T}_{p,1} \\ (\theta \operatorname{Frob}_p^2) \cdot i_*(x) &, \text{ if } \theta = \mathcal{S}_{p,2} \\ (\theta \operatorname{Frob}_p) \cdot i_*(x) &, \text{ if } \theta = \mathcal{T}_{p,2} \end{cases}$$

where Frob_p denotes arithmetic Frobenius at p, regarded as an element of $\operatorname{Gal}(\mathbf{Q}(\mu_n)/\mathbf{Q})$. This follows from [LSZ21b] Section 8.2 using the same normalization for the global Artin reciprocity map, and our definition of i.

Remark 5.2.8. As remarked earlier, at first glance it might appear that there's a certain asymmetry lingering in the background since we've made a choice in which slot to apply the determinant map to \mathbf{G}_m . However the results are independent of this choice. In fact as we pointed out earlier, it can be eliminated all together if one works with the larger group $G \times \mathbf{G}_m$ to begin with. However, this would make notation somewhat awkward in other sections and that's why we choose not to.

The polynomial $\mathcal{P}_p(x)$ of Definition 4.3.1 can be computed explicitly. It is given by

$$\mathcal{P}_p(x) = 1 - \frac{1}{p}\mathcal{T}_{p,1}\mathcal{T}_{p,2}x + \left(\frac{1}{p}\mathcal{S}_{p,1}\mathcal{T}_{p,2}^2 + \frac{1}{p}\mathcal{S}_{p,2}\mathcal{T}_{p,1}^2 - 2\mathcal{S}_{p,1}\mathcal{S}_{p,2}\right)x^2 - \frac{1}{p}\mathcal{S}_{p,1}\mathcal{S}_{p,2}\mathcal{T}_{p,1}\mathcal{T}_{p,2}x^3 + \mathcal{S}_{p,1}^2\mathcal{S}_{p,2}x^4.$$

Combining this with (13) it follows that i_* intertwines the action of $\mathcal{P}'_p(1)$ on $H^3_{\text{mot}}(Y_G(K_S[n]), \mathbf{Q}(2))$ with that of $\mathcal{P}'_p(\operatorname{Frob}_p^{-1})$ on $H^3_{\text{mot}}(Y_G(K_S) \times_{\operatorname{Spec}(\mathbf{Q})} \operatorname{Spec}(\mathbf{Q}(\mu_n)), \mathbf{Q}(2))$.

For data as in Definition 5.2.6, we write $\Xi_{\text{mot},n}(\underline{\delta})$ for the image of $\mathcal{Z}_{\text{mot},n}(\underline{\delta})$ under i_* . Then we can re-state Theorem 5.2.7 for the classes $\Xi_{\text{mot},n}(\underline{\delta})$:

Corollary 5.2.9. (1) If
$$\underline{\delta} \in \prod_{p \notin S} \mathcal{I}_0(G_p/G_p^\circ[p], \mathbf{Z}[1/p])$$
, then for any $n, m \in \mathscr{Z}_S$ with $\frac{m}{n} = p$

$$\operatorname{norm}_{\mathbf{Q}(\mu_n)}^{\mathbf{Q}(\mu_m)}(\Xi_{\operatorname{mot},m}(\underline{\delta})) = \mathcal{P}_{\operatorname{Tr}(\delta_p)}^{\operatorname{cycl}} \cdot \Xi_{\operatorname{mot},n}(\underline{\delta}), \text{ with } \mathcal{P}_{\operatorname{Tr}(\delta_p)}^{\operatorname{cycl}} \in \left(p-1, \mathcal{P}_p'(\operatorname{Frob}_p^{-1})\right) \subseteq \mathcal{H}_{G_p}^{\circ}(\mathbf{Z}[1/p])[\operatorname{Frob}_p^{\pm 1}].$$

(2) In the special cases $\underline{\delta} = \underline{\delta}_0$ and $\underline{\delta} = \underline{\delta}_1$ we have

$$\operatorname{norm}_{\mathbf{Q}(\mu_{n})}^{\mathbf{Q}(\mu_{m})}(\Xi_{\operatorname{mot},m}(\underline{\delta})) = \begin{cases} (p-1) \cdot \Xi_{\operatorname{mot},n}(\underline{\delta}), & \text{if } \underline{\delta} = \underline{\delta}_{0} \\ \mathcal{P}_{p}^{'}(\operatorname{Frob}_{p}^{-1}) \cdot \Xi_{\operatorname{mot},n}(\underline{\delta}), & \text{if } \underline{\delta} = \underline{\delta}_{1} \end{cases}$$

Proof. Theorem 5.2.7 and the discussion above

Remark 5.2.10. The classes $\Xi_{\text{mot},n}(\underline{\delta}_1)$ are not exactly the same as the corresponding ones in [LLZ14], [LLZ18], (e.g [LLZ18] 3.5.3 for split case) which satisfy the analogous norm relations (Corollary 5.2.9 (2)) at p, only modulo p-1, which is in line with the "optimal" aspect of both parts of Corollary 5.2.9.

5.2.5. Étale realisation and a "modulo- ℓ " norm relation. In this section we fix an prime $\ell \notin S$. By [Hub00] we have étale regulator maps

$$r_{\acute{e}t}: H^3_{\mathrm{mot}}(Y_G, \mathbf{Q}(2)) \longrightarrow H^3_{\acute{e}t}(Y_G, \mathbf{Q}_\ell(2))$$

defined at each finite level. Let $n \in \mathbb{Z}_{\geq 1}$ be a square-free integer coprime to ℓS . As in [LSZ21b], but with different notation, for each integer c > 1 and coprime to 6nS, we define classes

$${}_{c}\mathcal{Z}_{\acute{\mathrm{et}},n}\left(\underline{\delta}\right) := \left(c^{2} - \prod_{p|c} \mathcal{S}_{p}^{v_{p}(c)}\right) \cdot r_{\acute{\mathrm{et}}}\left(\mathcal{Z}_{\mathrm{mot},n}\left(\underline{\delta}\right)\right)$$
$${}_{c}\Xi_{\acute{\mathrm{et}},n}(\underline{\delta}) := \left(c^{2} - \prod_{p|c} \mathcal{S}_{p}^{v_{p}(c)} \mathrm{Frob}_{p}^{2}\right) \cdot r_{\acute{\mathrm{et}}}\left(\Xi_{\mathrm{mot},n}\left(\underline{\delta}\right)\right)$$

Of course by (13) these definitions are compatible with one another. These normalisation factors arise from the integral variant of the map of Theorem 5.2.3, a description of which can be found in ([LSZ21a] Section 7). If $\underline{\delta}$ is as in Theorem 5.2.7, then by the argument of ([LSZ21b] 9.5.2) but simpler since we are dealing with trivial coefficients, the classes $_{c}Z_{\acute{e}t,n}(\underline{\delta})$ actually lie in $H^{3}_{\acute{e}t}(Y_{G}, \mathbf{Z}_{\ell}(2))$ (and similarly for the $_{c}\Xi_{\acute{e}t}$ -classes). Furthermore, by Theorem 5.2.7, the Euler factors appearing are also defined over \mathbf{Z}_{ℓ} for $p \neq \ell$. Thus the norm relations of Theorem 5.2.7 hold *integrally* over \mathbf{Z}_{ℓ} . In particular letting $\mathscr{Z}_{c,\ell,S}$ denote the set of square-free positive integers coprime to $c\ell S$ having prime factors congruent to 1 modulo ℓ :

Corollary 5.2.11. Let $\underline{\delta}$ be as in the second statement of Theorem 5.2.7 and fix some $c \in \mathbb{Z}_{>1}$ coprime to 6S. Then for all $n, m \in \mathscr{Z}_{c,\ell,S}$, n|m:

$$(\mathrm{pr}_{n}^{m})_{*} (_{c}\mathcal{Z}_{\acute{e}t,m}(\underline{\delta})) = \left(\mathcal{Q}_{m/n} \prod_{p \mid \frac{m}{n}} \mathcal{P}_{p}^{'}(1)\right) \cdot _{c}\mathcal{Z}_{\acute{e}t,n}(\underline{\delta}) \mod \ell$$
$$\operatorname{norm}_{\mathbf{Q}(\mu_{n})}^{\mathbf{Q}(\mu_{m})} (_{c}\Xi_{\acute{e}t,m}(\underline{\delta})) = \left(\mathcal{Q}_{m/n}^{\operatorname{cycl}} \prod_{p \mid \frac{m}{n}} \mathcal{P}_{p}^{'}(\operatorname{Frob}_{p}^{-1})\right) \cdot _{c}\Xi_{\acute{e}t,n}(\underline{\delta}) \mod \ell$$

for some $\mathcal{Q}_{m/n} \in \prod_{p \mid \frac{m}{n}} \mathcal{H}_{G_p}^{\circ}(\overline{\mathbf{Z}}_{\ell})$ and $\mathcal{Q}_{m/n}^{\operatorname{cycl}} \in \prod_{p \mid \frac{m}{n}} \mathcal{H}_{G_p}^{\circ}(\overline{\mathbf{Z}}_{\ell})[\operatorname{Frob}_p^{\pm 1}].$

Proof. Immediate from the above discussion and the second part of Theorem 5.2.7.

Remark 5.2.12. As we said earlier, corresponding statements can also be made for non-trivial coefficient sheaves and the proofs go through.

6. INTEGRALITY FOR RANKIN-SELBERG TYPE PERIODS

6.1. The space $\operatorname{Hom}_{P_p}(\pi_{p,1} \otimes \pi_{p,2}, \delta_P)$ and an integral lattice in $\pi_{p,1} \otimes \pi_{p,2}$. The space $\operatorname{Hom}_{P_p}(\pi_{p,1} \otimes \pi_{p,2}, \delta_P)$, up to twist, has been studied in [JPSS83] in the more general setting of $\operatorname{GL}(n)$ and for (possibly ramified) Whittaker type representations, where a "generic" bound (≤ 1) on the dimension of this space is established, using vastly different methods. This step is crucial for the proof of the functional equation for the local zeta integrals of *op.cit*. To link this to our work we will relate the module $C_c^{\infty}(P_p \setminus G_p) \subset C_c^{\infty}(P_p \setminus G_p)$ to a certain module of P_p -coinvariants. The natural object to consider here is the module of δ_P -twisted P_p -coinvariants of $C_c^{\infty}(G_p/G_p^{\circ})$ given by

$$C_c^{\infty}(G_p/G_p^{\circ})_{P_p,\delta_P} := C_c^{\infty}(G_p/G_p^{\circ})/\langle x \cdot \xi - \delta_P(x)\xi \mid x \in P_p, \xi \in C_c^{\infty}(G_p/G_p^{\circ})\rangle.$$

This is again a module over $\mathcal{H}_{G_n}^{\circ}$ under the usual Hecke action induced by right translation.

Proposition 6.1.1. There is a Hecke equivariant isomorphism

$$\Phi: C_c^{\infty}(P_p \backslash G_p / G_p^{\circ}) \xrightarrow{\simeq} C_c^{\infty}(G_p / G_p^{\circ})_{P_p, \delta_P}, \ \operatorname{ch}(P_p g G_p^{\circ}) \mapsto \frac{1}{\operatorname{vol}_{P_p}\left(P_p \cap g G_p^{\circ} g^{-1}\right)} \operatorname{ch}(g G_p^{\circ})$$

where vol_{P_p} denotes the volume with respect to right or left normalised Haar measure and is independent of such choice.

Proof. Firstly, we show that the volume factor in question is independent of the choice of left or right normalised Haar measure. Indeed,

$$\begin{aligned} \operatorname{vol}_{P_p}(P_p \cap gG_p^{\circ}g^{-1}, d^R x) &= \int_{P_p \cap gG_p^{\circ}g^{-1}} d^R x \\ &= \int_{P_p \cap gG_p^{\circ}g^{-1}} \delta_P(x) \ d^L x = \operatorname{vol}_{P_p}(P_p \cap gG_p^{\circ}g^{-1}, d^L x) \end{aligned}$$

where the last equality follows from the fact that δ_P is trivial on $P_p \cap gG_p^{\circ}g^{-1}$. We now check that the map Φ is well-defined. To do this, it clearly suffices to show that $\Phi(\operatorname{ch}(P_p gG_p^{\circ}))$ agrees with $\Phi(\operatorname{ch}(P_p ygG_p^{\circ}))$ for $y \in P_p$. Indeed, $\operatorname{ch}(P_p ygG_p^{\circ})$ gets mapped to

(14)
$$\frac{1}{\operatorname{vol}_{P_p}(P_p \cap ygG_p^\circ g^{-1}y^{-1}, d^R x)}\operatorname{ch}(ygG_p^\circ) = \frac{1}{\operatorname{vol}_{P_p}(P_p \cap ygG_p^\circ g^{-1}, d^R x)}\operatorname{ch}(ygG_p^\circ)$$
$$= \frac{\delta_P(y)}{\operatorname{vol}_{P_p}(P_p \cap ygG_p^\circ g^{-1}, d^R x)}\operatorname{ch}(gG_p^\circ).$$

Now, $\operatorname{vol}_{P_p}(P_p \cap ygG_p^{\circ}g^{-1}, d^Rx)$ is given by $\int_{P_p \cap ygG_p^{\circ}g^{-1}} d^Rx = \int_{P_p \cap ygG_p^{\circ}g^{-1}} \delta_P(x)d^Lx$ which is in turn equal to $\delta_P(y) \operatorname{vol}_{P_p}(P_p \cap gG_p^{\circ}g^{-1})$. From (14) we now see that Φ is well-defined. The map Φ is a priori a morphism of **C**-vector spaces. Showing directly that Φ is Hecke equivariant using its current definition appears to

be very hard. We will instead construct an inverse morphism to Φ which can be readily seen to be Hecke equivariant. The result will then follow at once. There is a natural morphism

$$\Psi: C_c^{\infty}(G_p/G_p^{\circ}) \longrightarrow C_c^{\infty}(P_p \backslash G_p/G_p^{\circ}), \ \xi \mapsto \left(f_{\xi}: g \mapsto \int_{P_p} \xi(xg) \ d^Rx\right)$$

which is Hecke equivariant by construction. Now, for $y \in P_p$

$$f_{y \cdot \xi}(g) = \int_{P_p} \xi(y^{-1}xg) \ d^R x = \int_{P_p} \xi(y^{-1}xg)\delta_P(x) \ d^L x = \delta_P(y) \int_{P_p} \xi(xg)\delta_P(x) \ d^L x = \delta_P(y)f_{\xi}(g).$$

Thus, the map Ψ factors through $C_c^{\infty}(G_p/G_p^{\circ})_{P_p,\delta_P}$. Finally, for $g \in G_p$, the function $f_{\operatorname{ch}(gG_p^{\circ})}$ is supported on $P_pgG_p^{\circ}$ and $f_{\operatorname{ch}(gG_p^{\circ})}(g) = \operatorname{vol}_{P_p}(P_p \cap gG_p^{\circ}g^{-1})$. Hence $\Psi(\operatorname{ch}(gG_p^{\circ})) = \operatorname{vol}_{P_p}(P_p \cap gG_p^{\circ}g^{-1})\operatorname{ch}(P_pgG_p^{\circ})$ which shows that Φ and Ψ are mutually inverse as required. \Box

Definition 6.1.2. (1) The lattice $\mathcal{J}(G_p/G_p^\circ, \mathbf{Z}[1/p])_{P_p,\delta_P}$ inside $C_c^\infty(G_p/G_p^\circ)_{P_p,\delta_P}$ consists of $\mathbf{Z}[1/p]$ -linear combinations of elements of the form

$$\frac{1}{\operatorname{vol}_{P_p}\left(P_p \cap gG_p^\circ g^{-1}\right)}\operatorname{ch}(gG_p^\circ), \ g \in G_p$$

(2) The lattice $\mathcal{J}_1(G_p/G_p^\circ, \mathbf{Z}[1/p])_{P_p,\delta_P}$ inside $C_c^\infty(G_p/G_p^\circ)_{P_p,\delta_P}$ consists of $\mathbf{Z}[1/p]$ -linear combinations of elements of the form

$$\frac{1}{\operatorname{vol}_{P_p}\left(P_p(1) \cap gG_p^\circ g^{-1}\right)}\operatorname{ch}(gG_p^\circ), \ g \in G_p$$

where $P_p(1)$ is the open subgroup of P_p given by $\{x \in P_p \mid \det(x) \in \mathbf{Z}_p^{\times}, \det(x) \equiv 1 \mod p\}$.

Lemma 6.1.3. The map Φ identifies the lattices

$$C_c^{\infty}(P_p \backslash G_p / G_p^{\circ}, \mathbf{Z}[1/p]) \simeq_{\Phi} \mathcal{J}(G_p / G_p^{\circ}, \mathbf{Z}[1/p])_{P_p, \delta_P}$$
$$C_c^{\infty, 1}(P_p \backslash G_p / G_p^{\circ}, \mathbf{Z}[1/p]) \simeq_{\Phi} \mathcal{J}_1(G_p / G_p^{\circ}, \mathbf{Z}[1/p])_{P_p, \delta_P}.$$

Proof. The first identification follows at once from the definitions and Proposition 6.1.1. Thus we deal with the second one. It is straightforward to show that every double coset $P_p g G_p^{\circ}$ is equal to $P_p(z_1 \begin{bmatrix} p^m & 1 \end{bmatrix}, z_2) G_p^{\circ}$ with $m \in \mathbb{Z}$ and $z_i \in Z_p$, or $P_p(z_1 \begin{bmatrix} p^m & p^{-n} \\ 1 \end{bmatrix}, z_2) G_p^{\circ}$ with $n \in \mathbb{Z}_{\geq 1}, m \in \mathbb{Z}, m > -n$ and $z_i \in Z_p$. By definition of the two lattices and the map Φ , it suffices to show that

$$\frac{1}{p-1} \cdot \operatorname{vol}_{P_p} \left(P_p \cap gG_p^{\circ}g^{-1} \right) = \operatorname{vol}_{P_p} \left(P_p(1) \cap gG_p^{\circ}g^{-1} \right), \ g = \left(z_1 \begin{bmatrix} p^m \\ 1 \end{bmatrix}, z_2 \right)$$
$$\operatorname{vol}_{P_p} \left(P_p \cap gG_p^{\circ}g^{-1} \right) = \operatorname{vol}_{P_p} \left(P_p(1) \cap gG_p^{\circ}g^{-1} \right), \ g = \left(z_1 \begin{bmatrix} p^m & p^{-n} \\ 1 \end{bmatrix}, z_2 \right).$$

We firstly treat the first case, in which we have $P_p \cap gG_p^{\circ}g^{-1} = P_p^{\circ} \cap \begin{bmatrix} p^m \\ 1 \end{bmatrix} H_p^{\circ} \begin{bmatrix} p^m \\ 1 \end{bmatrix}^{-1}$. A quick matrix calculation shows that this is nothing more than P_p° if $m \leq 0$ and $\begin{bmatrix} \mathbf{Z}_p^{\times} & p^m \mathbf{Z}_p \\ 1 \end{bmatrix}$ if m > 0. The former has volume 1 and the latter has volume $\frac{1}{p^m}$. On the other hand, if $m \leq 0$ then $P_p(1) \cap gG_p^{\circ}g^{-1} = P_p(1) \cap P_p^{\circ} = \begin{bmatrix} 1+p\mathbf{Z}_p & \mathbf{Z}_p \\ 1 \end{bmatrix}$ which clearly has volume $\frac{1}{p-1}$. If m > 0 then $P_p(1) \cap gG_p^{\circ}g^{-1} = \begin{bmatrix} 1+p\mathbf{Z}_p & p^m\mathbf{Z}_p \\ 1 \end{bmatrix}$ which has volume $\frac{1}{(p-1)p^m}$ as required. For the second case, note that m + n is strictly greater than zero. Another quick matrix computation shows that in this case the two subgroups are identical and given by $\begin{bmatrix} 1+p^{n+m}\mathbf{Z}_p & \mathbf{Z}_p \\ 1 \end{bmatrix}$ if $m \leq 0$ and $\begin{bmatrix} 1+p^{n+m}\mathbf{Z}_p & p^m\mathbf{Z}_p \\ 1 \end{bmatrix}$ if m > 0.

We now let $\pi_{p,1}, \pi_{p,2}$ be unramified Whittaker type H_p -representations with $\omega_{\pi_{p,1}} \neq \omega_{\pi_{p,2}}^{-1}$. One then has a surjective linear map by acting on the spherical vector

$$C_c^{\infty}(G_p/G_p^{\circ})_{P_p,\delta_P}[\frac{1}{1-\mathcal{S}_p}] \longrightarrow (\pi_{p,1} \otimes \pi_{p,2})_{P_p,\delta_P}$$

where the object on the right denotes the δ_P -twisted P_p -coinvariants of the representation $\pi_{p,1} \boxtimes \pi_{p,2}$ defined in a similar fashion as $C_c^{\infty}(G_p/G_p^{\circ})_{P_p,\delta_P}$. Proposition 6.1.1 and Theorem 4.4.1 then imply that

dim Hom_{Pp}
$$(\pi_{p,1} \otimes \pi_{p,2}, \delta_P) \leq 1$$

and a non-zero linear form in this space is non-vanishing on the spherical vector $W_{\pi_{p,1}}^{\text{sph}} \otimes W_{\pi_{p,2}}^{\text{sph}}$. In our setup we can actually explicitly write down a non-zero element of this space using zeta integrals. The construction is similar to that of Section 4.2. Indeed, the assignment

(15)
$$\mathscr{D}_{\Pi_p}: \Pi_p := \pi_{p,1} \otimes \pi_{p,2} \longrightarrow \mathbf{C}, \ W_1 \otimes W_2 \mapsto \lim_{s \to 0} \frac{\int_{\mathbf{Q}_p^{\times}} \left(W_1 \boxtimes W_2\right) \left(\begin{bmatrix} a \\ 1 \end{bmatrix}\right) |a|_p^{s-1} d^{\times} a}{L(\pi_{p,1} \boxtimes \pi_{p,2}, s)}$$

is well-defined and gives an element of $\operatorname{Hom}_{P_p}(\pi_{p,1} \otimes \pi_{p,2}, \delta_P)$, which takes the value $L(\omega_{\pi_{p,1}\boxtimes \pi_{p,2}}, 0)^{-1}$ on the normalized spherical vector. This is clearly non-zero since $\omega_{\pi_{p,1}} \neq \omega_{\pi_{p,2}}^{-1}$.

Remark 6.1.4. It is worth noting that one can now obtain equality in Theorem 3.2.4 for all unramified Whittaker type H_p -representations, $\pi_{p,1}, \pi_{p,2}, \pi_{p,3}$ with the product of their central characters being trivial. This revolves around using Frobenius reciprocity and a property of the mirabolic subgroup of GL₂: Let $\pi_{p,1}, \pi_{p,2}, \pi_{p,3}$ be unramified Whittaker type H_p -representations with the product of their central characters being trivial. As usual, we write $I(\chi_{p,1}, \chi_{p,2})$ for the normalized induction of $\begin{bmatrix} \chi_{p,1} & \chi_{p,2} \end{bmatrix}$. We may assume after twisting that $\pi_{p,3}$ has trivial central character. Hence, $\omega_{\pi_{p,1}}\omega_{\pi_{p,2}} = 1$ and $\pi_{p,3}$ is isomorphic to $I(|\cdot|_p^{-a}, |\cdot|_p^a)$ with $a \in \mathbb{C}$ different to $\frac{1}{2}$. However, using the fact that $B_p = Z_p \times P_p$, one can check that

$$\pi_{p,3}^{\vee} \simeq I(|\cdot|_p^a, |\cdot|_p^{-a}) \simeq \left(\operatorname{Ind}_{P_p}^{H_p} \left(\delta_P^{a+1/2} \right) \right)_{\mathbb{T}}$$

where the subscript on the right denotes the eigenspace of the trivial character with respect to the action of the center. Thus, we have a chain of isomorphisms

$$\operatorname{Hom}_{H_p}(\pi_{p,1} \otimes \pi_{p,2} \otimes \pi_{p,3}, \mathbf{1}) \simeq \operatorname{Hom}_{H_p}(\pi_{p,1} \otimes \pi_{p,2}, I(|\cdot|_p^a, |\cdot|_p^{-a}))$$
$$\simeq \operatorname{Hom}_{P_p}\left(\pi_{p,1} \otimes \pi_{p,2}, \delta_P^{a+1/2}\right)$$
$$\simeq \operatorname{Hom}_{P_p}\left(\pi_{p,1} \otimes \pi_{p,2} \otimes \delta_P^{-a+1/2}, \delta_P\right)$$

where the second isomorphism follows by Frobenius reciprocity for the non-compact induction (e.g [Ren10] III 2.5) and the fact that $\omega_{\pi_{p,1}}\omega_{\pi_{p,2}} = 1$. However, as a P_p -representation, $\pi_{p,1} \otimes \pi_{p,2} \otimes \delta_P^{-a+1/2}$ is isomorphic to $\pi_{p,1} \otimes \sigma_{p,2}$ where $\sigma_{p,2}$ is the unramified Whittaker type H_p -representation given by $\pi_{p,2} \otimes |\det|_p^{-a+1/2}$. Finally, as a quasi-character of \mathbf{Q}_p^{\times} , we have $\omega_{\pi_{p,1}}\omega_{\sigma_{p,2}} = |\cdot|_p^{-2a+1}$ and this non-trivial since $a \neq \frac{1}{2}$. Thus we have shown that

$$\operatorname{Hom}_{H_p}(\pi_{p,1}\otimes\pi_{p,2}\otimes\pi_{p,3},\mathbf{1})\simeq\operatorname{Hom}_{P_p}(\pi_{p,1}\otimes\sigma_{p,2},\delta_P)$$

where $\pi_{p,1}$ and $\sigma_{p,2}$ are unramified Whittaker type H_p -representations with $\omega_{\pi_{p,1}}\omega_{\sigma_{p,2}} \neq 1$. But for each such pair, as we already noted, the explicit integral representation (15) above, gives a non-zero element of the Hom-space on the right. Unlike [Pra90] and [HS01], this approach does not distinguish between different cases of representations within this family. Of course it does not deal with ramified representations, but it does however give a unified proof for the whole of family of unramified Whittaker type representations, making full use of these two properties.

It is also worth noting that Λ_{Π_p} and \mathscr{D}_{Π_p} are related through the following commutative diagram

$$C_{c}^{\infty}(G_{p}/G_{p}^{\circ})_{P_{p},\delta_{P}}[\frac{1}{1-\mathcal{S}_{p}}] \longrightarrow (\Pi_{p})_{P_{p},\delta_{P}} \xrightarrow{\mathfrak{D}\Pi_{p}} \mathbf{C} \simeq (\Pi_{p}^{\vee})^{G_{p}^{\circ}}$$
$$\simeq_{\Phi} \uparrow$$
$$C_{c}^{\infty}(P_{p}\backslash G_{p}/G_{p}^{\circ})[\frac{1}{1-\mathcal{S}_{p}}] \xrightarrow{\Lambda_{\Pi_{p}}}$$

Passing to an integral level, we have the following result.

Theorem 6.1.5. Let $\Pi_p := \pi_{p,1} \otimes \pi_{p,2}$ be as above, and assume that the spherical Hecke eigensystem Θ_{Π_p} takes values in \mathcal{R} when restricted to $\mathcal{H}^{\circ}_{G_p}(\mathcal{R})$, for some $\mathbb{Z}[1/p]$ -algebra \mathcal{R} . Let \mathcal{Z} be a non-zero linear form in $\operatorname{Hom}_{P_p}(\Pi_p, \delta_P)$, normalized to take the value $L(\omega_{\Pi_p}, 0)^{-1}$ on $W^{\mathrm{sph}}_{\pi_{p,1}} \otimes W^{\mathrm{sph}}_{\pi_{p,2}}$. Then:

(1) The linear form \mathcal{Z} is valued in \mathcal{R} on the lattice

$$\operatorname{span}_{\mathcal{R}}\left\{\frac{1}{\operatorname{vol}_{P_p}(P_p \cap gG_p^{\circ}g^{-1})} g \cdot (W_{\pi_{p,1}}^{\operatorname{sph}} \otimes W_{\pi_{p,2}}^{\operatorname{sph}}) \mid g \in G_p\right\} \subseteq \pi_{p,1} \otimes \pi_{p,2}.$$

(2) The linear form \mathcal{Z} is valued in the ideal $(p-1, L(\Pi_p, 0)^{-1}) \subseteq \mathcal{R}$ on the lattice

$$\operatorname{span}_{\mathcal{R}}\left\{\frac{1}{\operatorname{vol}_{P_p}(P_p(1)\cap gG_p^\circ g^{-1})} g\cdot (W_{\pi_{p,1}}^{\operatorname{sph}}\otimes W_{\pi_{p,2}}^{\operatorname{sph}}) \mid g\in G_p\right\}\subseteq \pi_{p,1}\otimes \pi_{p,2}.$$

Proof. By multiplicity one described above, \mathcal{Z} is already completely determined and coincides with \mathscr{D}_{Π_p} . The value of \mathscr{D}_{Π_p} on an element of the lattice in question can be expressed as an \mathcal{R} -linear combination of elements of the form $\mathscr{D}_{\Pi_p}\left(\xi \cdot (W_{\pi_{p,1}}^{\mathrm{sph}} \otimes W_{\pi_{p,2}}^{\mathrm{sph}})\right)$ for some $\xi \in \mathcal{J}(G_p/G_p^\circ, \mathbb{Z}[1/p])_{P_p,\delta_P}$. Then ξ corresponds to some f in $C_c^{\infty}(P_p \setminus G_p/G_p^\circ, \mathbb{Z}[1/p])$ via Φ . The first part then follows from Corollary 4.3.8 and the commutative diagram above. For the second part, using the same argument as for the first part, the value of \mathscr{D}_{Π_p} evaluated on an element of the lattice in question is an \mathcal{R} -linear combination of elements of the form $\Lambda_{\Pi_p}(f)$ where now $f \in C_c^{\infty,1}(P_p \setminus G_p/G_p^\circ, \mathbb{Z}[1/p])$. But by Theorem 4.4.1, and Theorem 4.4.5, the corresponding Hecke operator $\mathcal{P}_{f,\mathrm{loc}}$, is given by $\frac{-S_p \mathscr{Q}'_f}{1-S_p}$ where $\mathcal{Q}_f \in (p-1, \mathcal{P}_p(1)) \subseteq \mathcal{H}_{G_p}^\circ(\mathbb{Z}[1/p])$. Hence by Proposition 4.2.5, $\Lambda_{\Pi_p}(f) = \Lambda_{\Pi_p}(\mathcal{P}_{f,\mathrm{loc}} \cdot f_0) = \Theta_{\Pi_p}(\mathcal{Q}_f)\Theta_{\Pi_p}(1-S_p)^{-1}\Lambda_{\Pi_p}(f_0) = \Theta_{\Pi_p}(\mathcal{Q}_f)$. This gives the result. \Box

6.1.1. Necessity of inverting $1 - S_p$ via an Ext-vanishing result of Prasad. Recall Theorem 4.4.1 which states that the module $C_c^{\infty}(P_p \setminus G_p/G_p^{\circ})[\frac{1}{1-S_p}]$ is free of rank one over $\mathcal{H}_{G_p}^{\circ}[\frac{1}{1-S_p}]$, generated by $\operatorname{ch}(P_p G_p^{\circ})$. A natural thing to ask is whether the localisation at the Hecke operator $1 - S_p$ is optimal and necessary. In other words is the module $C_c^{\infty}(P_p \setminus G_p/G_p^{\circ})$ generated by $\operatorname{ch}(P_p G_p^{\circ})$ over $\mathcal{H}_{G_p}^{\circ}$. As the title of this section suggests, the answer is no. Indeed, if $C_c^{\infty}(P_p \setminus G_p/G_p^{\circ})$ was cyclic as a $\mathcal{H}_{G_p}^{\circ}$ -module, then our earlier argument would show that

dim
$$\operatorname{Hom}_{P_p}(\pi_{p,1} \otimes \pi_{p,2}, \delta_P) \leq 1$$

for every unramified Whittaker type $\pi_{p,1}, \pi_{p,2}$. In particular,

$$\dim \operatorname{Hom}_{P_p}(\pi_p \otimes \pi_p^{\vee}, \delta_P) \leq 1$$

for every unramified irreducible principal-series π_p with $\omega_{\pi_p} = 1$. By the same argument as the one in Remark 6.1.4, one can deduce that the Hom-space above is nothing more than

$$\operatorname{Hom}_{H_p}\left(\pi_p\otimes\pi_p^{\vee}\otimes\operatorname{ind}_{B_p}^{H_p}\delta_B^{-1/2},\mathbf{1}\right)=\operatorname{Hom}_{\operatorname{PGL}_2(\mathbf{Q}_p)}\left(\pi_p\otimes\pi_p^{\vee}\otimes\operatorname{ind}_{B_p}^{H_p}\delta_B^{-1/2},\mathbf{1}\right).$$

It is well-known that the non-Whittaker type principal-series $\operatorname{ind}_{B_p}^{H_p} \delta_B^{-1/2}$ fits into a short exact sequence of $\operatorname{PGL}_2(\mathbf{Q}_p)$ -modules

$$0 \longrightarrow \mathbf{1} \longrightarrow \operatorname{ind}_{B_p}^{H_p} \delta_B^{-1/2} \longrightarrow \operatorname{St} \longrightarrow 0$$

Tensoring with π_p and taking $\operatorname{Ext}^*_{\operatorname{PGL}_2(\mathbf{Q}_p)}(-,\pi_p)$, we obtain a long exact sequence

(16)
$$0 \longrightarrow \operatorname{Hom}_{\operatorname{PGL}_{2}(\mathbf{Q}_{p})} \left(\pi_{p} \otimes \pi_{p}^{\vee} \otimes \operatorname{St}, \mathbf{1} \right) \longrightarrow \operatorname{Hom}_{\operatorname{PGL}_{2}(\mathbf{Q}_{p})} \left(\pi_{p} \otimes \pi_{p}^{\vee} \otimes \operatorname{ind}_{B_{p}}^{H_{p}} \delta_{B}^{-1/2}, \mathbf{1} \right)$$
$$\longrightarrow \operatorname{Hom}_{\operatorname{PGL}_{2}(\mathbf{Q}_{p})} \left(\pi_{p} \otimes \pi_{p}^{\vee}, \mathbf{1} \right) \longrightarrow \operatorname{Ext}_{\operatorname{PGL}_{2}(\mathbf{Q}_{p})}^{1} \left(\pi_{p} \otimes \operatorname{St}, \pi_{p} \right).$$

By ([Pra17] Proposition 2.6) and self-duality of St, the Ext-group appearing above is naturally isomorphic to $\operatorname{Ext}^{1}_{\operatorname{PGL}_{2}(\mathbf{Q}_{p})}(\pi_{p} \otimes \pi_{p}^{\vee}, \operatorname{St})$. By the Schneider-Stuhler duality theorem ([Pra17] Theorem 8.1) and the fact that $\operatorname{Ext}^{i}_{\operatorname{PGL}_{2}(\mathbf{Q}_{p})}(\pi_{p} \otimes \pi_{p}^{\vee}, \operatorname{St}) = 0$ for i > 1 (which is the split rank of PGL₂), we have

$$\operatorname{Ext}^{1}_{\operatorname{PGL}_{2}(\mathbf{Q}_{p})}\left(\pi_{p}\otimes\pi_{p}^{\vee},\operatorname{St}\right)\simeq\operatorname{Hom}_{\operatorname{PGL}_{2}(\mathbf{Q}_{p})}\left(\mathbf{1},\pi_{p}\otimes\pi_{p}^{\vee}\right).$$

However, this space vanishes by ([Pra17] Proposition 8.1). The space $\operatorname{Hom}_{PGL_2(\mathbf{Q}_p)}(\pi_p \otimes \pi_p^{\vee}, \mathbf{1})$ is at least one dimensional and $\operatorname{Hom}_{PGL_2(\mathbf{Q}_p)}(\pi_p \otimes \pi_p^{\vee} \otimes \operatorname{St}, \mathbf{1})$ is precisely one dimensional, where the second fact follows from the main result of [Pra90]. Thus going back to (16), we finally deduce that

dim Hom_{PGL2}(
$$\mathbf{Q}_p$$
) $\left(\pi_p \otimes \pi_p^{\vee} \otimes \operatorname{ind}_{B_p}^{H_p} \delta_B^{-1/2}, \mathbf{1}\right) \ge 2$

Of course this contradicts the bound dim $\operatorname{Hom}_{P_p}(\pi_p \otimes \pi_p^{\vee}, \delta_P) \leq 1$. Thus the $\mathcal{H}_{G_p}^{\circ}$ -module $C_c^{\infty}(P_p \setminus G_p / G_p^{\circ})$ is indeed not generated by $\operatorname{ch}(P_p G_p^{\circ})$ without localising at the Hecke operator $1 - \mathcal{S}_p$.

6.1.2. Relation to branching laws for ℓ -modular representations. We fix a prime $\ell \neq p$ and we identify the residue field of $\overline{\mathbf{Z}}_{\ell}$ with $\overline{\mathbf{F}}_{\ell}$. We fix a choice of p^i in $\overline{\mathbf{F}}_{\ell}$ for each $i \in \mathbf{Z}$, compatible with the projection from $\overline{\mathbf{Z}}_{\ell}$ to $\overline{\mathbf{F}}_{\ell}$. The character δ_P is still given (using different notation) by $\delta_P([a\ 1\ 1]) = p^{-v_p(a)}$. We have algebra and module structures on $\mathcal{H}^{\circ}_{G_p}(\overline{\mathbf{F}}_{\ell}), C^{\infty}_c(P_p \backslash G_p^{\circ}, \overline{\mathbf{F}}_{\ell}), C^{\infty}_c(G_p/G_p^{\circ}, \overline{\mathbf{F}}_{\ell})_{P_p, \delta_P}$ by lifting to the corresponding $\overline{\mathbf{Z}}_{\ell}$ -points and then projecting back down. Clearly this is independent of the choice of lifts and thus well-defined.

As in ([Gro24] Section 3.3) we say that an unramified $\overline{\mathbf{F}}_{\ell}$ -linear H_p -representation π_p is non-degenerate if dim $(\pi_p)^{H_p^{\circ}} = 1$ and $(\pi_p)^{H_p^{\circ}}$ generates π_p . For such a representation we still write $W_{\pi_p}^{\text{sph}}$ for its spherical vector. The irreducible $\overline{\mathbf{F}}_{\ell}$ -linear unramified principal-series are indeed non-degenerate. This is part of [MS14], but for GL₂, it's been known before that as mentioned in [BL95]. For the reducible cases, it is a bit more subtle compared to the zero characteristic case (see for example [Vig89]).

Theorem 6.1.6. Suppose $\ell \nmid p(p-1)$ and let $\pi_{p,1}, \pi_{p,2}$ be unramified non-degenerate $\overline{\mathbf{F}}_{\ell}$ -linear H_p -representations with $\omega_{\pi_{p,1}}(p) \neq \omega_{\pi_{p,2}}^{-1}(p)$ in $\overline{\mathbf{F}}_{\ell}^{\times}$. Then

 $\dim_{\overline{\mathbf{F}}_{\ell}} \operatorname{Hom}_{P_p}(\pi_{p,1} \otimes \pi_{p,2}, \delta_P) \leq 1$

and any such non-zero linear form is non-vanishing on $W^{\mathrm{sph}}_{\pi_{p,1}} \otimes W^{\mathrm{sph}}_{\pi_{p,2}}$

Proof. By fixing an isomorphism $\mathbf{C} \simeq \overline{\mathbf{Q}}_{\ell}$, it follows from Corollary 4.4.3 that $C_c^{\infty}(P_p \setminus G_p/G_p^{\circ}, \overline{\mathbf{Z}}_{\ell})[\frac{1}{1-\mathcal{S}_p}]$ is cyclic as a $\mathcal{H}_{G_p}^{\circ}(\overline{\mathbf{Z}}_{\ell})[\frac{1}{1-\mathcal{S}_p}]$ -module generated by $\operatorname{ch}(P_p G_p^{\circ})$. From this, it is clear that $C_c^{\infty}(P_p \setminus G_p/G_p^{\circ}, \overline{\mathbf{F}}_{\ell})[\frac{1}{1-\mathcal{S}_p}]$ is also cyclic as a $\mathcal{H}_{G_p}^{\circ}(\overline{\mathbf{F}}_{\ell})[\frac{1}{1-\mathcal{S}_p}]$ -module generated by $\operatorname{ch}(P_p G_p^{\circ})$. We have a commutative square

$$C_{c}^{\infty}(P_{p}\backslash G_{p}/G_{p}^{\circ},\overline{\mathbf{Z}}_{\ell}) \stackrel{\Phi}{\longrightarrow} C_{c}^{\infty}(G_{p}/G_{p}^{\circ},\overline{\mathbf{Z}}_{\ell})_{P_{p},\delta_{P}}$$

$$\downarrow^{\nu_{\ell}} \qquad \tau_{\ell}\downarrow$$

$$C_{c}^{\infty}(P_{p}\backslash G_{p}/G_{p}^{\circ},\overline{\mathbf{F}}_{\ell}) \stackrel{\tilde{\Phi}}{\longrightarrow} C_{c}^{\infty}(G_{p}/G_{p}^{\circ},\overline{\mathbf{F}}_{\ell})_{P_{p},\delta_{P}}$$

where Φ is the map from Proposition 6.1.1 and $\tilde{\Phi}$ is defined by first lifting through ν_{ℓ} and then applying Φ and τ_{ℓ} , which is clearly well-defined. From the volume calculations in the proof of Lemma 6.1.3, the definition of Φ and the assumption on $\ell \nmid p(p-1)$, it follows that $\tilde{\Phi}$ is surjective. A diagram chase using injectivity of Φ also shows that $\tilde{\Phi}$ is injective, and finally $\mathcal{H}_{G_p}^{\circ}(\overline{\mathbf{F}}_{\ell})$ -equivariance follows from the definitions. Thus, $C_c^{\infty}(G_p/G_p^{\circ}, \overline{\mathbf{F}}_{\ell})_{P_p,\delta_P}[\frac{1}{1-S_p}]$ is a cyclic $\mathcal{H}_{G_p}^{\circ}(\overline{\mathbf{F}}_{\ell})[\frac{1}{1-S_p}]$ -module generated by $ch(G_p^{\circ})$. Of course, the linear map $C_c^{\infty}(G_p/G_p^{\circ}, \overline{\mathbf{F}}_{\ell}) \to \Pi_p = \pi_{p,1} \boxtimes \pi_{p,2}$ can be defined in a completely algebraic manner via $ch(gG_p^{\circ}) \mapsto g \cdot W_{\Pi_p}^{\mathrm{sph}}$. Then the linear map

$$C_c^{\infty}(G_p/G_p^{\circ}, \overline{\mathbf{F}}_{\ell})_{P_p, \delta_P}[\frac{1}{1-\mathcal{S}_p}] \longrightarrow (\Pi_p)_{P_p, \delta_P}, \ \frac{\operatorname{ch}(gG_p^{\circ})}{(1-\mathcal{S}_p)^i} \mapsto \frac{g \cdot W_{\Pi_p}^{\operatorname{sph}}}{(1-\omega_{\Pi_p}(p))^i}, \ g \in G_p, i \in \mathbf{Z}_{\geq 0}$$

is surjective. By the cyclicity mentioned above and the fact that for such modular representations the space $(\Pi_p)^{G_p^{\circ}}$ is still one dimensional, its image is generated by the spherical vector and the result follows. \Box

Corollary 6.1.7. Suppose $\ell \nmid p(p-1)$. Let $\pi_{p,1}, \pi_{p,2}$ be unramified non-degenerate $\overline{\mathbf{F}}_{\ell}$ -linear representations of H_p and let $\pi_{p,3}$ be an unramified normalized parabolic induction $\operatorname{ind}_{B_p}^{H_p} \begin{bmatrix} \chi_{p,1} & \chi_{p,2} \end{bmatrix}$ with $(\chi_{p,1}/\chi_{p,2})(p) \neq p$ in $\overline{\mathbf{F}}_{\ell}^{\times}$. Then

 $\dim_{\overline{\mathbf{F}}_{\ell}} \operatorname{Hom}_{H_p}(\pi_{p,1} \otimes \pi_{p,2} \otimes \pi_{p,3}, \mathbf{1}) \leq 1.$

If furthermore $\pi_{p,3}$ is also non-degenerate then any such non-zero linear form does not vanish on the spherical vector $W_{\pi_{p,1}}^{\mathrm{sph}} \otimes W_{\pi_{p,2}}^{\mathrm{sph}} \otimes W_{\pi_{p,3}}^{\mathrm{sph}}$.

Proof. Again by twisting we may assume that $\pi_{p,3}$ has trivial central character, i.e. $\chi_{p,2} = \chi_{p,1}^{-1}$ and $\omega_{\pi_{p,1}}\omega_{\pi_{p,2}} = 1$. in the same manner as Remark 6.1.4, one can then relate the Hom-space in question to $\operatorname{Hom}_{P_p}(\pi_{p,1} \otimes \sigma_{p,2}, \delta_P)$ where $\sigma_{p,2} = \pi_{p,2} \otimes \left[(\delta_P^{1/2} \chi_{p,1})(\det(-)) \right]$ as an H_p -representation. By assumption, we have $\chi_{p,1}(p)^2 \neq p$ and thus the product of central characters $(\omega_{\pi_{p,1}}\omega_{\sigma_{p,2}})(p) = \chi_{p,1}(p)^2 \cdot p^{-1} \neq 1$. Then the bound on the dimension follows from Theorem 6.1.6. Finally, to show that if such a non-zero linear form exists then it doesn't vanish on the spherical vector (assuming $\pi_{p,3}$ is non-degenerate), one needs to trace through the various isomorphisms of Hom-spaces in Remark 6.1.4 (Frobenius reciprocity and duality) while using [Vig89] Lemme 18 part (3) and its proof, and the corresponding result of Theorem 6.1.6.

6.2. Integral lattice in $S(\mathbf{Q}_p^2) \otimes \pi_{p,1} \otimes \pi_{p,2}$. As usual, we let $\pi_{p,1}, \pi_{p,2}$ be unramified H_p -representations of Whittaker type. By Theorem 3.2.3 we know that a non-zero H_p -invariant period in $\operatorname{Hom}_{H_p}(S(\mathbf{Q}_p^2) \otimes \pi_{p,1} \otimes \pi_{p,2}, \mathbf{1})$ is unique up to scalars and is non-vanishing on the unramified vector $\operatorname{ch}(\mathbf{Z}_p^2) \otimes W_{\pi_{p,1}}^{\operatorname{sph}} \otimes W_{\pi_{p,2}}^{\operatorname{sph}}$. We can thus rescale it so it maps $\operatorname{ch}(\mathbf{Z}_p^2) \otimes W_{\pi_{p,1}}^{\operatorname{sph}} \otimes W_{\pi_{p,2}}^{\operatorname{sph}}$ to 1 and we call such a period normalized. In fact we will see later, when we discuss an application of this to global Rankin-Selberg integrals, that such a non-zero period always exists and we will explicitly write it down. For now though, we need only consider an abstract period of this form, without any sort of realization of it.

Theorem 6.2.1. Let \mathcal{Z} be the non-zero normalized H_p -invariant period in $\operatorname{Hom}_{H_p}(\mathcal{S}(\mathbf{Q}_p^2) \otimes \pi_{p,1} \otimes \pi_{p,2}, \mathbf{1})$. Assume that the spherical Hecke eigensystem of $\pi_{p,1} \boxtimes \pi_{p,2}$ restricts to a morphism $\mathcal{H}^{\circ}_{G_p}(\mathcal{R}) \to \mathcal{R}$ for some $\mathbf{Z}[1/p]$ -algebra $\mathcal{R} \subseteq \mathbf{C}$. Then:

(1) The period \mathcal{Z} is valued in \mathcal{R} on the lattice

$$\operatorname{span}_{\mathcal{R}}\left\{\phi\otimes g\left(W_{\pi_{p,1}}^{\operatorname{sph}}\otimes W_{\pi_{p,2}}^{\operatorname{sph}}\right) \mid g\in G_p \text{ and } \phi\in\mathcal{S}(\mathbf{Q}_p^2) \text{ valued in } \frac{1}{\operatorname{vol}_{H_p}\left(\operatorname{Stab}_{H_p}(\phi)\cap gG_p^\circ g^{-1}\right)}\mathcal{R}\right\}$$

(2) The period \mathcal{Z} is valued in the ideal $(p-1, L(\pi_{p,1} \boxtimes \pi_{p,2}, 0)^{-1}) \subseteq \mathcal{R}$ on the lattice

$$\operatorname{span}_{\mathcal{R}}\left\{\phi\otimes g\left(W_{\pi_{p,1}}^{\operatorname{sph}}\otimes W_{\pi_{p,2}}^{\operatorname{sph}}\right) \mid g\in G_{p} \text{ and } \phi\in\mathcal{S}_{0}(\mathbf{Q}_{p}^{2}) \text{ valued in } \frac{1}{\operatorname{vol}_{H_{p}}\left(\operatorname{Stab}_{H_{p}}(\phi)\cap gG_{p}^{\circ}[p]g^{-1}\right)}\mathcal{R}\right\}$$

(3) If $\omega_p := \omega_{\pi_{p,1}} \omega_{\pi_{p,2}} \neq 1$, the period \mathcal{Z} satisfies

$$L(\omega_p, 0)\mathcal{Z}(-) \in (p-1, L(\pi_{p,1} \boxtimes \pi_{p,2}, 0)^{-1}) \subseteq \mathcal{R}$$

on the lattice

$$\operatorname{span}_{\mathcal{R}}\left\{\phi\otimes g\left(W_{\pi_{p,1}}^{\operatorname{sph}}\otimes W_{\pi_{p,2}}^{\operatorname{sph}}\right) \mid g\in G_{p} \text{ and } \phi\in\mathcal{S}(\mathbf{Q}_{p}^{2}) \text{ valued in } \frac{1}{\operatorname{vol}_{H_{p}}\left(\operatorname{Stab}_{H_{p}}(\phi)\cap gG_{p}^{\circ}[p]g^{-1}\right)}\mathcal{R}\right\}$$

Proof. The proof relies on the three parts of Theorem 5.1.1, in particular the integral behavior of the Hecke operators \mathcal{P}_{δ} attached to integrals elements δ . To obtain the result one combines this with Proposition 4.4.1 of [LSZ21b], whose proof is formal and adapts to our setup in an identical fashion.

Remark 6.2.2. Notice how parts (2) and (3) compared to part (1), of Theorem 6.2.1, have the determinant level condition $G_p^{\circ}[p]$. Also, note that part (3) compared to part (2) allows for Schwartz functions which might not vanish at the origin, at the expense of a potential Dirichlet *L*-factor popping up.

Remark 6.2.3. The assumption on the spherical Hecke eigensystem of $\pi_{p,1} \boxtimes \pi_{p,2}$ is a natural one to make from a global viewpoint as we will see later when we discuss an application of this to global Rankin-Selberg integrals for product of modular forms.

6.3. Integral lattice in $\pi_{p,1} \otimes \pi_{p,2} \otimes \pi_{p,3}$. For notational convenience we now let $\pi_{p,3}$ be an unramified Whittaker type H_p -representation of the form $I(\chi_p, |\cdot|_p^{-1/2})$ such that $\omega_{\pi_{p,1}}\omega_{\pi_{p,2}}\omega_{\pi_{p,3}} = 1$. From Theorem 3.2.4 we know that a non-zero invariant period in $\operatorname{Hom}_{H_p}(\pi_{p,1} \otimes \pi_{p,2} \otimes \pi_{p,3}, 1)$ is unique up to scalars and is non vanishing on the spherical vector $W_{\pi_{p,1}}^{\mathrm{sph}} \otimes W_{\pi_{p,2}}^{\mathrm{sph}} \otimes W_{\pi_{p,3}}^{\mathrm{sph}}$. We can thus rescale it to map this spherical vector to 1 and call such a period normalized. In fact by Remark 6.1.4 (or by [HS01]) such a period always exists. Once again we can make use of the H_p -equivariant map

$$\mathcal{S}(\mathbf{Q}_p^2) \to \pi_{p,3}, \ \phi \mapsto f_{\hat{\phi},\chi_p}$$

constructed in [LSZ21a], which we describe in this section for completeness: For a Schwartz function $\phi \in \mathcal{S}(\mathbf{Q}_p^2)$, we write $\hat{\phi}$ for its Fourier transform

$$\hat{\phi}(x,y) := \int_{\mathbf{Q}_p} \int_{\mathbf{Q}_p} e_p(xv - yu)\phi(u,v) \, dudv$$

where e_p is the standard additive character of \mathbf{Q}_p mapping $1/p^n$ to $\exp(2\pi i/p^n)$ (this is an explicit description of the character ψ fixed at the beginning of Section 4) and du, dv denote the usual additive Haar measure on \mathbf{Q}_p . Since $\pi_{p,3}$ is unramified, the character χ_p is of the form $|\cdot|_p^a$ for some $a \in \mathbf{C}$. Let $b = a + \frac{1}{2}$, then by ([LSZ21a] 3.2.2),

$$f_{\phi,\chi_p}(h) := \lim_{s \to 0} \frac{|\det(h)|_p^{s+a+\frac{1}{2}}}{L\left(|\cdot|_p^b, 2s+1\right)} \int_{\mathbf{Q}_p^{\times}} \phi((0,x)h) |x|_p^{2s+b+1} \ d^{\times}x, \ h \in H_p$$

is well-defined and gives an element of $I(\chi_p, |\cdot|_p^{-1/2})$. Furthermore, the assignment $\phi \mapsto f_{\hat{\phi}, \chi_p}$ is surjective, H_p -equivariant and by ([LSZ21b] 3.2.5) it maps ch(\mathbf{Z}_p^2) to $W_{\pi_{p,3}}^{\mathrm{sph}}$.

Corollary 6.3.1. Let \mathcal{Z} be the non-zero normalized invariant period in $\operatorname{Hom}_{H_p}(\pi_{p,1} \otimes \pi_{p,2} \otimes \pi_{p,3}, \mathbf{1})$. Assume that the spherical Hecke eigensystem of $\pi_{p,1} \boxtimes \pi_{p,2}$ restricts to a morphism $\mathcal{H}^{\circ}_{G_p}(\mathcal{R}) \to \mathcal{R}$ for some $\mathbf{Z}[1/p]$ -algebra $\mathcal{R} \subseteq \mathbf{C}$. Then:

(1) The period \mathcal{Z} is valued in \mathcal{R} on the lattice

$$\operatorname{span}_{\mathcal{R}}\left\{g\left(W_{\pi_{p,1}}^{\operatorname{sph}}\otimes W_{\pi_{p,2}}^{\operatorname{sph}}\right)\otimes f_{\hat{\phi},\chi_{p}} \mid g \in G_{p} \text{ and } \phi \in \mathcal{S}(\mathbf{Q}_{p}^{2}) \text{ valued in } \frac{1}{\operatorname{vol}_{H_{p}}\left(\operatorname{Stab}_{H_{p}}(\phi) \cap gG_{p}^{\circ}g^{-1}\right)}\mathcal{R}\right\}$$

(2) The period \mathcal{Z} is valued in the ideal $(p-1, L(\pi_{p,1} \boxtimes \pi_{p,2}, 0)^{-1}) \subseteq \mathcal{R}$ on the lattice

$$\operatorname{span}_{\mathcal{R}}\left\{g\left(W_{\pi_{p,1}}^{\operatorname{sph}}\otimes W_{\pi_{p,2}}^{\operatorname{sph}}\right)\otimes f_{\hat{\phi},\chi_{p}}\mid g\in G_{p} \text{ and } \phi\in\mathcal{S}_{0}(\mathbf{Q}_{p}^{2}) \text{ valued in } \frac{1}{\operatorname{vol}_{H_{p}}\left(\operatorname{Stab}_{H_{p}}(\phi)\cap gG_{p}^{\circ}[p]g^{-1}\right)}\mathcal{R}\right\}$$

(3) If $\omega_p := \omega_{\pi_{p,1}} \omega_{\pi_{p,2}} \neq 1$, the period Z satisfies

$$L(\omega_p, 0)\mathcal{Z}(-) \in (p-1, L(\pi_{p,1} \boxtimes \pi_{p,2}, 0)^{-1}) \subseteq \mathcal{R}$$

on the lattice

$$\operatorname{span}_{\mathcal{R}}\left\{g\left(W_{\pi_{p,1}}^{\operatorname{sph}}\otimes W_{\pi_{p,2}}^{\operatorname{sph}}\right)\otimes f_{\hat{\phi},\chi_{p}} \mid g \in G_{p} \text{ and } \phi \in \mathcal{S}(\mathbf{Q}_{p}^{2}) \text{ valued in } \frac{1}{\operatorname{vol}_{H_{p}}\left(\operatorname{Stab}_{H_{p}}(\phi) \cap gG_{p}^{\circ}[p]g^{-1}\right)}\mathcal{R}\right\}$$

Proof. This follows from Theorem 6.2.1 and the map induced by $f_{(-),\chi_p}$ at the level of Hom-spaces.

6.4. Rankin-Selberg integral for product of modular forms.

6.4.1. The local integrals. We once again fix an additive character $\psi : \mathbf{Q}_p \to \mathbf{C}^{\times}$ of conductor one. As before, we consider the family of G_p -representations $\pi_{p,1} \boxtimes \pi_{p,2}$ where $\pi_{p,1}$ and $\pi_{p,2}$ are unramified H_p -representations of Whittaker type, and we identify $\pi_{p,1}$ with $\mathcal{W}(\pi_{p,1}, \psi)$ and $\pi_{p,2}$ with $\mathcal{W}(\pi_{p,2}, \overline{\psi})$.

Definition 6.4.1 ([JPSS83]). For $W_1 \in \mathcal{W}(\pi_{p,1}, \psi), W_2 \in \mathcal{W}(\pi_{p,2}, \overline{\psi})$ and $\phi \in \mathcal{S}(\mathbf{Q}_p^2)$, the zeta integral $Z(W_1, W_2, \phi, s)$ is given by

(17)
$$Z(\phi, W_1, W_2, s) = \int_{N_p \setminus H_p} W_1(h) W_2(h) \phi((0, 1)h) |\det(h)|_p^s dh$$

Proposition 6.4.2. (1) The zeta integral of Definition 6.4.1 converges for $\Re(s)$ large enough and admits unique meromorphic continuation as a rational function of p^s .

(2) As W_1, W_1 and ϕ vary, the $Z(W_1, W_2, \phi, s)$ generate the fractional ideal $L(\pi_{p,1} \boxtimes \pi_{p,2}, s) \mathbf{C}[p^s, p^{-s}]$.

Proof. This is one of the main results of [JPSS83] where it is stated and proved in greater generality.

Proposition 6.4.3. Let $\phi_0 = \operatorname{ch}(\mathbf{Z}_p^2)$. Then,

$$Z(\phi_0, W^{\text{sph}}_{\pi_{p,1}}, W^{\text{sph}}_{\pi_{p,1}}, s) = L(\pi_{p,2} \boxtimes \pi_{p,2}, s)$$

Proof. This is well know and the strategy is that of [Gro20] 2.2.2, using [Shi76] to evaluate the spherical Whittaker functions in terms of Satake parameters. \Box

Definition 6.4.4. Let $\pi_{p,1}, \pi_{p,2}$ be as above. The linear form $\mathcal{Z} : \mathcal{S}(\mathbf{Q}_p^2) \otimes \pi_{p,1} \otimes \pi_{p,2} \to \mathbf{C}$ is given by

$$\mathcal{Z}(\phi \otimes W_1 \otimes W_1) = \lim_{s \to 0} \frac{Z(\phi, W_1, W_2, s)}{L(\pi_{p,1} \boxtimes \pi_{p,2}, s)}.$$

By Proposition 6.4.2, the linear form \mathcal{Z} is well defined. Additionally, Proposition 6.4.3 shows that \mathcal{Z} takes the value 1 on $\phi_0 \otimes W^{\text{sph}}_{\pi_{p,1}} \otimes W^{\text{sph}}_{\pi_{p,2}}$. A standard change of variables also shows that \mathcal{Z} is in fact H_p invariant. Putting everything together, the linear form \mathcal{Z} gives a non-zero invariant period in the space $\operatorname{Hom}_{H_p}(\mathcal{S}(\mathbf{Q}_p^2) \otimes \pi_{p,1} \otimes \pi_{p,2}, \mathbf{1})$, which by Theorem 3.2.3, is in fact a basis.

6.4.2. The global integral. We fix a prime ℓ and an isomorphism $\iota : \mathbf{C} \simeq \overline{\mathbf{Q}}_{\ell}$. We let f_1 and f_2 be two normalised cuspidal Hecke eigenforms in $S_k(\Gamma_1(N))$ with $k \ge 2$ and we write ϵ_i for the Nebentype of f_i . To these we can associate two irreducible admissible automorphic representations of $H(\mathbf{A})$ ([Gel06]), which we call $\pi_1 = \pi_{\iota,f_1} = \bigotimes_p' \pi_{p,1}$ and $\pi_2 = \pi_{\iota,f_2} = \bigotimes_p' \pi_{p,2}$, with normalisation as in [LW12]. It is well-known that each local component is generic, and it is unramified if and only if p doesn't divide ∞ or N. We write ω_1 and ω_2 for their respective central characters. We also write $L = L_{f_1,f_2}$ for the number field associated to f_1 and f_2 , given by adjoining the Hecke eigenvalues of f_1 and f_2 to \mathbf{Q} . We write $\mathbf{L} = \mathbf{L}_{\iota,f_1,f_2}$ for a choice of ℓ -adic number field containing the image of L under ι .

To a Schwartz function $\Phi \in \mathcal{S}(\mathbf{A}^2)$, one can associate an Eisenstein series using Godement-Jacquet theory ([JL06]), which is given by

$$E(g,s;\Phi) = \sum_{\gamma \in B(\mathbf{Q}) \setminus H(\mathbf{Q})} |\det(\gamma g)|_p^s \int_{\mathbf{A}^{\times}} \Phi((0,x)\gamma g)\omega_1(x)\omega_2(x)|x|_p^{2s} d^{\times}x.$$

For cusp forms $\varphi_1 \in \pi_1$ and $\varphi_2 \in \pi_2$, one has the global Rankin-Selbger integral as in [JS81], which is given by

$$I(s, \Phi, \varphi_1, \varphi_2) = \int_{Z(\mathbf{A})H(\mathbf{Q})\backslash H(\mathbf{A})} \varphi_1(g)\varphi_2(g)E(g, s; \Phi) \ dg$$

By [JS81], this is a meromorphic function of $s \in \mathbf{C}$ and satisfies a functional equation. We let S denote the finite set of places $S = \{\infty, p | \ell N\}$, and write $L^S(\pi_1 \boxtimes \pi_2, s)$ for the partial L-function

$$L^{S}(\pi_{1} \boxtimes \pi_{2}, s) = \prod_{p \notin S} L(\pi_{p,1} \boxtimes \pi_{p,2}, s)$$

where the local L-factors are as in Proposition 6.4.2. Finally, we note that the local zeta integral

$$Z(\Phi_p, W_{p,1}, W_{p,2}, s)$$

of Definition 6.4.1 still exists for the infinite place and satisfies similar properties ([Jac09]).

Theorem 6.4.5. Let $\varphi_i \simeq \otimes_p W_{p,i}$ be decomposable cusp forms, with $W_{p,i} = W_{\pi_{p,i}}^{\text{sph}}$ for $p \notin S$. Let $g = (g_1, g_2) \in G(\mathbf{A}^{\infty})$, and suppose that $\Phi = \otimes_p \Phi_p$ is also decomposable. Write

$$\varepsilon_{\ell,f_1,f_2}\left(\Phi,g_1,g_2\right) := \lim_{s \to 0} \frac{I(s,\Phi,g_1\varphi_1,g_2\varphi_2)}{L^S(\pi_1 \boxtimes \pi_2, s) \cdot \prod_{p \in S} Z(\Phi_p,g_{1,p}W_{p,1},g_{2,p}W_{p,2},s)} \in \overline{\mathbf{Q}}_{\ell}$$

The following are true:

(1) If Φ_p is valued in $\operatorname{vol}_{H_p}(\operatorname{Stab}_{H_p}(\Phi_p) \cap g_p G_p^\circ g_p^{-1})^{-1} \mathcal{O}_{\mathbf{L}}$ for $p \notin S$, then

$$\varepsilon_{\ell,f_1,f_2}(\Phi,g_1,g_2) \in \mathcal{O}_{\mathbf{L}}$$

(2) Suppose that the assumption of part (1) holds and let S_0 denote a finite set of places away from Sfor which: $p \equiv 1 \mod \ell$, $(\epsilon_1 \epsilon_2)(p) \neq 1$ and Φ_p is valued in $\operatorname{vol}_{H_p}(\operatorname{Stab}_{H_p}(\Phi_p) \cap g_p G_p^\circ[p] g_p^{-1})^{-1} \mathcal{O}_{\mathbf{L}}$ (note the addition of the determinant level $G_p^\circ[p]$ instead of G_p°). Then,

$$\left(\prod_{\substack{p\in S_0 \text{ s.t:}\\ \Phi_p\notin\mathcal{S}_0(\mathbf{Q}_p^2)}} L(\epsilon_1\epsilon_2, 0)^{-1}\right) \cdot \varepsilon_{\ell, f_1, f_2}(\Phi, g_1, g_2) \in \prod_{p\in S_0} \left(p-1, L(\pi_{p,1}\boxtimes \pi_{p,2}, 0)^{-1}\right) \subseteq \mathcal{O}_{\mathbf{L}}.$$

In particular, if the level of f_1, f_2 satisfies $(\#(\mathbf{Z}/N\mathbf{Z})^{\times}, \ell) = 1$, then this holds without the bracketed product on the left.

Proof. Firstly note that by the works of [JS81], [JPSS83] and [Jac09], the function inside the limit admits unique analytic continuation as an entire function of s, and thus the limit makes sense. We write S' for the finite set of places outside S, consisting of p such that $g_p \notin G_p^\circ$, or $\Phi_p \neq \operatorname{ch}(\mathbf{Z}_p^2)$. By Section 4 of [JS81], the limit in question coincides with

$$\prod_{p \in S'} \left(\lim_{s \to 0} \frac{Z(\Phi_p, g_{1,p} W_{p,1}, g_{2,p} W_{p,2}, s)}{L(\pi_{p,1} \boxtimes \pi_{p,2}, s)} \right) = \prod_{p \in S'} \mathcal{Z}(\Phi_p \otimes g_{1,p} W_{p,1} \otimes g_{2,p} W_{p,2})$$

where \mathcal{Z} is as in Definition 6.4.4. We are now in a position to apply Theorem 6.2.1 at each prime $p \in S'$, with the $\mathbb{Z}[1/p]$ -algebra \mathcal{R} taken to be $\mathcal{O}_{\mathbf{L}}$. Thus, it suffices to show that the local condition on the spherical Hecke eigensystem of each local component $\pi_{p,1} \boxtimes \pi_{p,2}$ is satisfied. Indeed, this follows similarly to Theorem 8.2.1 in [Gro24] using as mentioned before, the normalizations of [LW12]. If the last assumption on the level of f_1, f_2 holds then the bracketed integral on the left is a unit in $\mathcal{O}_{\mathbf{L}}$ which concludes the proof.

APPENDIX A. STRUCTURE OF $C_c^{\infty}(P_p \setminus G_p / G_p)[\frac{1}{1-S_r}]$

This appendix is dedicated to the proof of Theorem 4.4.1. As such we refer the reader back to Section 4 for notation and definitions.

Theorem A.0.1. The module $C_c^{\infty}(P_p \setminus G_p/G_p^{\circ})[\frac{1}{1-S_p}]$ is free of rank one over $\mathcal{H}_{G_p}^{\circ}[\frac{1}{1-S_p}]$, generated by the characteristic function $f_0 := \operatorname{ch}(P_p G_p^{\circ})$. In particular, for $f \in C_c^{\infty}(P_p \setminus G_p/G_p^{\circ})$, we have $f = \frac{-S_p Q_f'}{1-S_p} \cdot f_0$, where Q_f is the Hecke operator described in Section 4.4.

Proof. We firstly show that the module in question is cyclic with f_0 as a generator. By (11), it suffices to show that the intersection of the kernels of the linear forms Λ_{Π_p} as we vary the representation Π_p , is trivial. For an element in this intersection, we have $\Lambda_{\Pi_p}(f) = \Theta_{\Pi_p}(\mathcal{Q}_f) = 0$ for every Π_p . By the usual density argument, this implies that \mathcal{Q}_f is identically zero in $\mathcal{H}^c_{G_p}$. We now claim that $\mathcal{Q}_f = 0$ implies that f = 0. Firstly, for $m, n \in \mathbb{Z}_{\geq 0}$ and $r, t \in \mathbb{Z}$, we put $f_{m,n}^{r,t} := \operatorname{ch}(P_p p^r (p^t \begin{bmatrix} p^m \ p^{-n} \\ 1 \end{bmatrix}, 1)G_p^\circ)$ in $C_c^{\infty}(P_p \backslash G_p / G_p^\circ)$. We can always write f as

$$f = \left(\sum_{i=1}^{N} a_i f_{m_i,0}^{r_i,t_i}\right) + \left(\sum_{j=1}^{M} b_j f_{\mu_j,\nu_j}^{\rho_j,\tau_j}\right)$$

where:

• $\nu_j > 0$ for all j

1

- $|m_0| \le |m_1| \le \cdots \le |m_N|$; if $|m_i| = |m_{i+1}|$ then $r_i \le r_{i+1}$; if $|m_i| = |m_{i+1}|$ and $r_i = r_{i+1}$ then $t_i \le t_{i+1}$.
- $|\mu_0| + \nu_0 \le |\mu_1| + \nu_1 \le \cdots \le |\mu_M| + \nu_M$; if $|\mu_j| + \nu_j = |\mu_{j+1}| + \nu_{j+1}$ then $|\mu_j| \le |\mu_{j+1}|$; if $|\mu_j| + \nu_j = |\mu_{j+1}| + \nu_{j+1}$ and $|\mu_j| = |\mu_{j+1}|$ then $\rho_j \le \rho_{j+1}$; if $|\mu_j| + \nu_j = |\mu_{j+1}| + \nu_{j+1}$, $|\mu_j| = |\mu_{j+1}|$ and $\rho_j = \rho_{j+1}$ then $\tau_j \le \tau_{j+1}$.

By proposition 4.3.6 we have

(18)
$$\mathcal{Q}_f = \left(\sum_{i=1}^N a_i \mathscr{V}_{r_i, t_i} \mathscr{P}_{m_i}\right) + \left(\sum_{j=1}^M b_j \mathscr{V}_{\rho_j, \tau_j} \left(\mathscr{P}_{\mu_j} + \mathscr{Q}_{\mu_j, \nu_j} \mathcal{P}_p(1)\right)\right) = 0$$

in $\mathcal{H}_{G_p}^{\circ} = \mathbf{C}[\mathcal{S}_{p,1}^{\pm 1}, \mathcal{T}_{p,1}, \mathcal{S}_{p,2}^{\pm 1}, \mathcal{T}_{p,2}]$. We write (18) as

(19)
$$\left(\sum_{i=1}^{N} a_i \mathscr{V}_{r_i, t_i} \mathscr{P}_{m_i}\right) + \left(\sum_{j=1}^{M} b_j \mathcal{V}_{\rho_j, \tau_j} \mathscr{P}_{\mu_j}\right) + \left(\sum_{j=1}^{M} b_j \mathscr{V}_{\rho_j, \tau_j} \mathscr{Q}_{\mu_j, \nu_j} \mathcal{P}_p(1)\right) = 0$$

If the rightmost sum in (19) is non-zero, then $\mathcal{P}_p(1)$ divides $\left(\sum_{i=1}^N a_i \mathscr{V}_{r_i,t_i} \mathscr{P}_{m_i}\right) + \left(\sum_{j=1}^M b_j \mathscr{V}_{\rho_j,\tau_j} \mathscr{P}_{\mu_j}\right)$. We now claim that this cannot happen. In particular, we show that $\mathcal{P}_p(1)$ never divides a non-zero finite sum

of the form $\sum c \ (\mathcal{S}_{p,1}\mathcal{S}_{p,2})^s \mathcal{S}_{p,1}^k \mathscr{P}_{\lambda}$. Consider an arbitrary sum of this form, which we write as (20)

$$\Delta := \left(p^{-n/2} \mathcal{P}_{p,1}^{(n)}(1) \sum_{i,j} c_{i,j} (\mathcal{S}_{p,1} \mathcal{S}_{p,2})^{s_i} \mathcal{S}_{p,1}^{k_j} \right) + \left(p^{-n/2} \mathcal{P}_{p,2}^{(n)}(1) \sum_{i,j} d_{i,j} (\mathcal{S}_{p,1} \mathcal{S}_{p,2})^{\tilde{s}_i} \mathcal{S}_{p,1}^{\tilde{k}_j} \right) + \text{terms of this form involving } \mathcal{P}_{p,1}^{(e_1)}(1), \mathcal{P}_{p,2}^{(e_2)}(1) \text{ with } e_1, e_2 < n.$$

It is straightforward to compute, using the explicit description of the spherical Hecke eigensystem of the representations Π_p , that

• $\mathcal{P}_{p}(1) = 1 - \frac{1}{p}\mathcal{T}_{p,1}\mathcal{T}_{p,2} - \frac{1}{p}\mathcal{S}_{p,1}\mathcal{S}_{p,2}\mathcal{T}_{p,1}\mathcal{T}_{p,2} + \frac{1}{p}\mathcal{S}_{p,1}\mathcal{T}_{p,2}^{2} + \frac{1}{p}\mathcal{S}_{p,2}\mathcal{T}_{p,1}^{2} - 2\mathcal{S}_{p,1}\mathcal{S}_{p,2} + \mathcal{S}_{p,1}^{2}\mathcal{S}_{p,2}^{2}$ • $\frac{1}{p^{n/2}}\mathcal{P}_{p,1}^{(n)}(1) = \frac{1}{p^{\lfloor n/2 \rfloor}}\mathcal{T}_{p,1}^{n-1} - \frac{1}{p^{\lfloor n/2 \rfloor}}\mathcal{S}_{p,1}\mathcal{T}_{p,2}\mathcal{T}_{p,1}^{n-2} + O(\mathcal{T}_{p,1}^{n-3})$ • $\frac{1}{p^{n/2}}\mathcal{P}_{p,2}^{(n)}(1) = \frac{1}{p^{\lfloor n/2 \rfloor}}\mathcal{T}_{p,2}^{n-1} - \frac{1}{p^{\lfloor n/2 \rfloor}}\mathcal{S}_{p,2}\mathcal{T}_{p,1}\mathcal{T}_{p,2}^{n-2} + O(\mathcal{T}_{p,2}^{n-3})$

where $O(\mathcal{T}_{p,i}^e)$ denotes a linear combination of terms of the form $\mathcal{S}_{p,1}^a \mathcal{S}_{p,2}^b \mathcal{T}_{p,i}^c$ with $c \leq e$. From this, by looking at the degrees of the $\mathcal{T}_{p,i}$ terms, we can assume that that $n \geq 3$ in (20), since otherwise the non-divisibility claim holds trivially. It is also important to note that Δ has no monomials containing $\mathcal{T}_{p,1}^a \mathcal{T}_{p,2}^b$ with both $a, b \geq 2$. If $\mathcal{P}_p(1)$ divides Δ in $\mathcal{H}_{G_p}^\circ$, then it is not hard to see that we can write

$$\Delta = \left(A_1 \mathcal{T}_{p,1}^{n-3} + A_2 \mathcal{T}_{p,1}^{n-4} \mathcal{T}_{p,2} + \dots + A_{n-3} \mathcal{T}_{p,1} \mathcal{T}_{p,2}^{n-4} + A_{n-2} \mathcal{T}_{p,2}^{n-3} + C\right) \mathcal{P}_p(1)$$

where $A_i \in \mathbf{C}[\mathcal{S}_{p,1}^{\pm 1}, \mathcal{S}_{p,2}^{\pm 1}]$ and C is a $\mathbf{C}[\mathcal{S}_{p,1}^{\pm 1}, \mathcal{S}_{p,2}^{\pm 1}]$ -linear combination of monomials containing only $\mathcal{T}_{p,1}^a \mathcal{T}_{p,2}^b$ with a + b < n - 3.

Comparing coefficients of $\mathcal{T}_{p,1}^{n-1}$ and $\mathcal{T}_{p,2}^{n-1}$ in Δ , we get the relations

(22)
$$A_{1} = \frac{1}{p^{\lfloor n/2 \rfloor - 1}} \sum_{i,j} c_{i,j} \mathcal{S}_{p,1}^{s_{i}+k_{j}} \mathcal{S}_{p,2}^{s_{i}-1} \text{ and } A_{n-2} = \frac{1}{p^{\lfloor n/2 \rfloor - 1}} \sum_{i,j} d_{i,j} \mathcal{S}_{p,1}^{\tilde{s}_{i}+\tilde{k}_{j}-1} \mathcal{S}_{p,2}^{\tilde{s}_{i}}$$

The monomials that contribute to $\frac{1}{p^{\lfloor n/2 \rfloor}} \mathcal{S}_{p,1} \mathcal{T}_{p,2} \mathcal{T}_{p,1}^{n-2}$ in Δ are $A_1 \mathcal{T}_{p,1}^{n-3}$ and $A_2 \mathcal{T}_{p,1}^{n-4} \mathcal{T}_{p,2}$. Similarly, the monomials contributing to $\frac{1}{p^{\lfloor n/2 \rfloor}} \mathcal{S}_{p,2} \mathcal{T}_{p,1} \mathcal{T}_{p,2}^{n-2}$ in Δ are $A_{n-2} \mathcal{T}_{p,2}^{n-3}$ and $A_{n-3} \mathcal{T}_{p,1} \mathcal{T}_{p,2}^{n-4}$. Comparing coefficients of $\mathcal{T}_{p,2} \mathcal{T}_{p,1}^{n-2}$ and $\mathcal{T}_{p,1} \mathcal{T}_{p,2}^{n-2}$, we get the relations

(23)
$$-\frac{1}{p}(1+\mathcal{S}_{p,1}\mathcal{S}_{p,2})A_1 + \frac{1}{p}\mathcal{S}_{p,2}A_2 = \frac{-1}{p^{\lfloor n/2 \rfloor}}\sum_{i,j}c_{i,j}\mathcal{S}_{p,1}^{s_i+k_j+1}\mathcal{S}_{p,2}^{s_i}$$
$$\frac{1}{p}(1+\mathcal{S}_{p,1}\mathcal{S}_{p,2})A_{n-2} + \frac{1}{p}\mathcal{S}_{p,1}A_{n-3} = \frac{-1}{p^{\lfloor n/2 \rfloor}}\sum_{i,j}d_{i,j}\mathcal{S}_{p,1}^{\tilde{s}_i+\tilde{k}_j}\mathcal{S}_{p,2}^{\tilde{s}_i+1}.$$

Combining (22) and (23) and simplifying, we obtain the relations

(24)
$$A_1 = S_{p,2}A_2$$
 and $A_{n-2} = S_{p,2}A_{n-3}$

We observe that for n = 3 or 4, (24) already implies our claim that $\mathcal{P}_p(1)|\Delta$ if and only if $\Delta = 0$. Thus we may assume that $n \geq 5$. Comparing coefficients of $\mathcal{T}_{p,1}^a \mathcal{T}_{p,2}^b$ with a + b = n - 1 and $a, b \geq 2$ (which as we remarked earlier, they vanish), we obtain the following system of equations:

(25)

$$S_{p,1}A_1 - (1 + S_{p,1}S_{p,2})A_2 + S_{p,2}A_3 = 0$$

$$S_{p,1}A_2 - (1 + S_{p,1}S_{p,2})A_3 + S_{p,2}A_4 = 0$$

$$\vdots$$

$$S_{p,1}A_{n-4} - (1 + S_{p,1}S_{p,2})A_{n-3} + S_{p,2}A_{n-2} = 0.$$

A lengthy but elementary substitution exercise, together with (24), yields that

$$A_2 = \mathcal{S}_{p,1}\mathcal{S}_{p,2}A_2 \text{ for } n = 5$$

$$A_2 = S_{p,2}^{n-5} A_{n-3}$$
 and $A_{n-3} = S_{p,1}^{n-5} A_2$ for $n \ge 6$.

Combining this with (24) and (25), we deduce that $A_1 = \cdots = A_{n-2} = 0$, which finishes the proof of our first claim.

Thus, (19) now implies that $S := \sum_{j=1}^{M} b_j \mathscr{V}_{\rho_j,\tau_j} \mathscr{Q}_{\mu_j,\nu_j}$ is identically zero. We claim that $b_M = 0$, and thus

33

inductively, all the $b_i = 0$. To show this, we use the explicit expression of \mathscr{V} and \mathscr{Q} found in proposition 4.3.6. The leading monomial of $\mathscr{V}_{\rho_M,\tau_M} \mathscr{Q}_{\mu_M,\nu_M}$ is given up to a scalar multiple, by

$$\begin{cases} (\mathcal{S}_{p,1}\mathcal{S}_{p,2})^{\rho_M} \mathcal{S}_{p,1}^{\tau_M} \mathcal{T}_{p,1}^{\mu_M + \nu_M - 1} \mathcal{T}_{p,2}^{\nu_M - 1} &, \text{ if } \mu_M \ge 0\\ (\mathcal{S}_{p,1}\mathcal{S}_{p,2})^{\rho_M} \mathcal{S}_{p,1}^{\tau_M} \mathcal{T}_{p,1}^{\nu_M - 1} \mathcal{T}_{p,2}^{|\mu_M| + \nu_M - 1} &, \text{ if } \mu_M < 0. \end{cases}$$

Firstly, we assume that $\mu_M = 0$. If $|\mu_{M-1}| + \nu_{M-1} < |\mu_M| + \nu_M = \nu_M$, then the result follows at once. by considering powers of $\mathcal{T}_{p,1}$. If $|\mu_{M-1}| + \nu_{M-1} = |\mu_M| + \nu_M = \nu_M$, then we must have $|\mu_{M-1}| \leq |\mu_M| = 0$ and hence $\mu_{M-1} = 0$ which the implies that $\nu_{M-1} = \nu_M$. The result then follows by considering ρ_M, τ_M and ρ_{M-1}, τ_{M-1} since in this case, the equality $(|\mu_M| + \nu_M, |\mu_M|, \rho_M, \tau_M) = (|\mu_{M-1}| + \nu_{M-1}, |\mu_{M-1}|, \rho_{M-1}, \tau_{M-1})$ implies that $f_{\mu_M,\nu_M}^{\rho_M,\tau_M} = f_{\mu_{M-1},\nu_{M-1}}^{\rho_{M-1},\tau_{M-1}}$. We now assume that $|\mu_M| > 0$. If $|\mu_{M-1}| + \nu_{M-1} < |\mu_M| + \nu_{M-1}$ then the result follows again by considering the exponents of $\mathcal{T}_{p,1}$ or $\mathcal{T}_{p,2}$ depending on the parity of μ_M . Assume instead that $|\mu_{M-1}| + \nu_{M-1} = |\mu_M| + \nu_{M-1}$, in which case we have $|\mu_{M-1}| \leq |\mu_M|$. Then

$$\nu_M = \nu_{M-1} + |\mu_{M-1}| - |\mu_M| \le \nu_{M-1}$$

Let $K \in \{1, ..., M-1\}$ be the smallest integer for which $|\mu_K| + \nu_K = |\mu_{K+1}| + \nu_K = \cdots = |\mu_{M-1}| + \nu_{M-1} = |\mu_M| + \nu_M$. Then

 $|\mu_K| \leq \cdots \leq |\mu_{M-1}| \leq |\mu_M|$ and $\nu_M \leq \nu_{M-1} \leq \cdots \leq \nu_K$.

Without loss of generality, assume that $\mu_M > 0$. We now show that we may also assume $b_M \mathscr{V}_{\rho_M,\tau_M} \mathscr{Q}_{\mu_M,\nu_M}$ is the only term of S with a monomial which is a scalar multiple of

$$(\mathcal{S}_{p,1}\mathcal{S}_{p,2})^{\rho_M}\mathcal{S}_{p,1}^{\tau_M}\mathcal{T}_{p,1}^{\mu_M+\nu_M-1}\mathcal{T}_{p,1}^{\nu_M-1}$$

and hence $b_M = 0$: Any other term containing such monomial, has to come from some $\mathscr{Q}_{\mu_L,\nu_L}$, with $L \in \{K, \ldots, M-1\}$ and $\mu_L \geq 0$. Let $L_1 > L_2 > \cdots > L_d$ be all such values of L. The leading monomial of $b_{L_d} \mathscr{V}_{\rho_{L_d},\tau_{L_d}} \mathscr{Q}_{\mu_{L_d},\nu_{L_d}}$ is a scalar multiple of

$$(\mathcal{S}_{p,1}\mathcal{S}_{p,2})^{\rho_{L_d}}\mathcal{S}_{p,1}^{\tau_{L_d}}\mathcal{T}_{p,1}^{\mu_{L_d}+\nu_{L_d}-1}\mathcal{T}_{p,2}^{\nu_{L_d}-1}.$$

Since in this case $f_{\mu_{L_i},\nu_{L_i}}^{\rho_{L_i},\tau_{L_i}}$ is completely determined by $(|\mu_{L_i}| + \nu_{L_i}, |\mu_{L_i}|, \rho_{L_i}, \tau_{L_i})$, or equivalently $(|\mu_{L_i}| + \nu_{L_i}, \nu_{L_i}, \tau_{L_i})$, it follows that $b_{L_d} \mathscr{V}_{\rho_{L_d},\tau_{L_d}} \mathscr{Q}_{\mu_{L_d},\nu_{L_d}}$ is the only term in S containing a monomial which is a scalar multiple of $(\mathcal{S}_{p,1}\mathcal{S}_{p,2})^{\rho_{L_d}}\mathcal{S}_{p,1}^{\tau_{L_d}}\mathcal{T}_{p,1}^{\mu_{L_d}+\nu_{L_d}-1}\mathcal{T}_{p,2}^{\nu_{L_d}-1}$. Hence $b_{L_d} = 0$ and continuing like this we conclude that $b_{L_1} = \cdots = b_{L_d} = 0$. Thus our assumption is justified. But then $b_M = 0$ which is what we initially wanted to prove. This proves our second claim, i.e $b_j = 0$ for all $j = 1, \ldots, M$. Thus (19) now implies that

$$T := \sum_{i=1}^{N} a_i \mathscr{V}_{r_i, t_i} \mathscr{P}_{m_i} = 0$$

Our last claim is that the coefficients a_i are also zero: We do this by showing that $a_N = 0$. Recall that

$$\mathscr{P}_{m_i} = \frac{1}{p^{m_i}} \begin{cases} \mathcal{P}_{p,1}^{(m_i+1)}(1), & \text{if } m_i \ge 0\\ \mathcal{P}_{p,2}^{(|m_i|+1)}(1), & \text{if } m_i < 0 \end{cases}$$

Suppose first that $|m_N| \geq 2$. Looking at the explicit expressions for $\mathcal{P}_{p,1}^{(n)}(1)$ and $\mathcal{P}_{p,2}^{(n)}(1)$ it follows that $a_N \mathscr{V}_{r_N,t_N} \mathscr{P}_{m_N}$ is the only term of T that contains a monomial which is a scalar multiple of

$$(\mathcal{S}_{p,1}\mathcal{S}_{p,2})^{r_N}\mathcal{S}_{p,1}^{t_N}\mathcal{T}_{p,1}^{m_N}$$

and hence $a_N = 0$. Now suppose that $|m_N| = 1$. If $0 = |m_{N-1}| < 1$ we are clearly done by considering once again exponents of $\mathcal{T}_{p,1}$ or $\mathcal{T}_{p,2}$. If $|m_{N-1}| = 1$, we let $K \in \{1, \ldots, N-1\}$ be the smallest integer for which $|m_K| = |m_{K+1}| = \cdots = |m_{N-1}| = |m_N| = 1.$

Then, we have

$$T = \left(\sum_{i=1}^{K-1} a_i (\mathcal{S}_{p,1} \mathcal{S}_{p,2})^{r_i} \mathcal{S}_{p,1}^{t_i}\right) \mathscr{P}_0 + \left(\sum_{i=K}^N a_i (\mathcal{S}_{p,1} \mathcal{S}_{p,2})^{r_i} \mathcal{S}_{p,1}^{t_i} \mathscr{P}_{m_i}\right), \text{ where } m_i = \pm 1$$
$$= \left(\sum_{i=1}^{K-1} a_i (\mathcal{S}_{p,1} \mathcal{S}_{p,2})^{r_i} \mathcal{S}_{p,1}^{t_i}\right) \mathscr{P}_0 + \left(\sum_{i\in I_1} a_i (\mathcal{S}_{p,1} \mathcal{S}_{p,2})^{r_i} \mathcal{S}_{p,1}^{t_i}\right) \mathscr{P}_1 + \left(\sum_{i\in I_{-1}} a_i (\mathcal{S}_{p,1} \mathcal{S}_{p,2})^{r_i} \mathcal{S}_{p,1}^{t_i}\right) \mathscr{P}_{-1}.$$

where $I_1 \cup I_{-1} = \{K, \ldots, N\}$, $i \in I_1$ if and only if $m_1 = 1$ and $i \in I_{-1}$ if and only if $m_i = -1$. We write

$$C_1 := \sum_{i \in I_1} a_i (\mathcal{S}_{p,1} \mathcal{S}_{p,2})^{r_i} \mathcal{S}_{p,1}^{t_i} \text{ and } C_{-1} := \sum_{i \in I_{-1}} a_i (\mathcal{S}_{p,1} \mathcal{S}_{p,2})^{r_i} \mathcal{S}_{p,1}^{t_i}$$

Recall that

$$\mathscr{P}_0 = 1 - \mathcal{S}_{p,1}\mathcal{S}_{p,2}, \ \mathscr{P}_1 = -\frac{1}{p}\mathcal{S}_{p,1}\mathcal{T}_{p,2} + \frac{1}{p}\mathcal{T}_{p,1}, \ \mathscr{P}_{-1} = \frac{1}{p}\mathcal{S}_{p,2}\mathcal{T}_{p,1} + \frac{1}{p}\mathcal{T}_{p,2}.$$

Since T = 0, we must have

$$-\frac{C_1}{p}\mathcal{S}_{p,1}\mathcal{T}_{p,2} + \frac{C_1}{p}\mathcal{T}_{p,1} - \frac{C_{-1}}{p}\mathcal{S}_{p,2}\mathcal{T}_{p,1} + \frac{C_{-1}}{p}\mathcal{T}_{p,2} = 0$$

From this, we deduce that

$$C_1 = C_{-1} \mathcal{S}_{p,2}$$
 and $C_{-1} = C_1 \mathcal{S}_{p,1}$.

It follows at once that $C_1 = C_{-1} = 0$. Since for $i \in I_1$, $f_{m_i,0}^{r_i,t_i} = f_{1,0}^{r_i,t_i}$ is completely determined by (r_i, t_i) we conclude that $a_i = 0$ for every $i \in I_1$ and in an identical fashion $a_i = 0$ for every $i \in I_{-1}$. This deals with the case where $|m_N| = 1$. Finally, if $m_N = 0$, then

$$T = \left(\sum_{i=1}^{N} a_i (\mathcal{S}_{p,1} \mathcal{S}_{p,2})^{r_i} \mathcal{S}_{p,1}^{t_i}\right) \mathscr{P}_0 = 0$$

and $f_{m_i,0}^{r_i,t_i} = f_{0,0}^{r_i,t_i}$ is once again determined by (r_i,t_i) . Thus the claim follows. Putting everything together, we have shown that $\mathscr{Q}_f = 0$ implies that f = 0 and as a result, we deduce that

$$\bigcap_{\Pi_p} \ker(\Lambda_{\Pi_p}) = 0.$$

This is exactly what we wanted to prove in order to establish the cyclicity statement. For freeness, note that $\Lambda_{\Pi_p}: C_c^{\infty}(P_p \setminus G_p/G_p^{\circ}) \to (\Pi_p^{\vee})^{G_p^{\circ}}$, for $\omega_{\pi_{p,1}} \neq \omega_{\pi_{p,2}}^{-1}$, can be extended naturally to a Hecke equivariant map

$$C_c^{\infty}(P_p \backslash G_p / G_p^{\circ})[\frac{1}{1 - \mathcal{S}_p}] \longrightarrow (\Pi_p^{\vee})^{G_p^{\circ}}, \ \frac{f}{(1 - \mathcal{S}_p)^r} \mapsto \frac{\Lambda_{\Pi_p}(f)}{\Theta_{\Pi_p}(1 - \mathcal{S}_p^{-1})^r}$$

We just proved that every such element $\frac{f}{(1-S_p)^r}$ is given by $\frac{-S_p Q'_f}{(1-S_p)^{r+1}} \cdot f_0$. Thus freeness follows from Hecke equivariance and the fact that this slightly shrinked family of representations Π_p is still dense in $\text{Spec}(\mathcal{H}^{\circ}_{G_p})$.

References

- [BG14] Manjul Bhargava and Benedict H Gross. Arithmetic invariant theory. Symmetry: Representation Theory and Its Applications: In Honor of Nolan R. Wallach, pages 33–54, 2014.
- [BL95] Laure Barthel and Ron Livné. Modular representations of GL(2) of a local-field: The ordinary, unramified case. Journal of Number Theory, 55(1):1–27, 1995.
- [Bri90] Michel Brion. Vers une généralisation des espaces symétriques. Journal of Algebra, 134(1):115–143, 1990.
- [Bum97] Daniel Bump. Automorphic Forms and Representations. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 1997.
- [Cas80] William Casselman. The unramified principal series of p-adic groups. i. the spherical function. Compositio Mathematica, 40(3):387–406, 1980.
- [Col04] Pierre Colmez. La conjecture de Birch et Swinnerton-Dyer p-adique. Asterisque-societe mathematique de France, 294:251–320, 2004.
- [FV00] Eric M Friedlander and Vladimir Voevodsky. Bivariant cycle cohomology. Cycles, transfers, and motivic homology theories, 143:138–187, 2000.
- [Gel06] Stephen Gelbart. Automorphic forms and Artin's conjecture. In Modular Functions of One Variable VI: Proceedings International Conference, University of Bonn, Sonderforschungsbereich Theoretische Mathematik July 2–14, 1976, pages 241–276. Springer, 2006.
- [GK72] Izrail Moiseevich Gel'fand and David A Kazhdan. On the representation of the group GL(n, K) where K is a local field. Functional Analysis and Its Applications, 6(4):315–317, 1972.

- [Gro20] Giada Grossi. On norm relations for Asai-Flach classes, 2020.
- [Gro24] Alexandros Groutides. Integral structures in smooth $\operatorname{GL}_2(\mathbf{Q}_p)$ -representations and zeta integrals, 2024.
- [HS01] Michael Harris and Anthony J Scholl. A note on trilinear forms for reducible representations and Beilinson's conjectures. *Journal of the European Mathematical Society*, 3(1):93–104, 2001.
- [Hub00] Annette Huber. Realization of Voevodsky's motives. Journal of Algebraic Geometry, 9(4):755, 2000.
- [Jac90] Hervé Jacquet. Rankin-Selberg convolutions: archimedean theory. (No Title), page 125, 1990.
- [Jac09] Hervé Jacquet. Archimedean Rankin-Selberg integrals. Contemporary Mathematics, 14:57, 2009.
- [JL06] Hervé Jacquet and Robert P Langlands. Automorphic forms on GL(2): Part 1, volume 114. Springer, 2006.
- [JPSS83] Hervé Jacquet, Ilya I Piatetskii-Shapiro, and Joseph A Shalika. Rankin-Selberg convolutions. American journal of mathematics, 105(2):367–464, 1983.
 - [JS81] Hervé Jacquet and Joseph A Shalika. On Euler products and the classification of automorphic representations i. *American Journal of Mathematics*, 103(3):499–558, 1981.
- [Kat04] Kazuya Kato. p-adic Hodge theory and values of zeta functions of modular forms. Astérisque, 295:117–290, 2004.
- [KM15] Robert Kurinczuk and Nadir Matringe. Rankin-Selberg local factors modulo ℓ , 2015.
- [Kno95] Friedrich Knop. On the set of orbits for a Borel subgroup. *Commentarii Mathematici Helvetici*, 70(2):285–309, 1995.
- [LLZ14] Antonio Lei, David Loeffler, and Sarah Zerbes. Euler systems for Rankin-Selberg convolutions of modular forms. Annals of Mathematics, 180(2):653–771, September 2014.
- [LLZ18] Antonio Lei, David Loeffler, and Sarah Livia Zerbes. Euler systems for Hilbert modular surfaces. In Forum of Mathematics, Sigma, volume 6. Cambridge University Press, 2018.
- [Loe21] David Loeffler. Gross-Prasad periods for reducible representations. Forum Mathematicum, 33(5):1169–1177, 2021.
- [LSZ21a] David Loeffler, Chris Skinner, and Sarah Livia Zerbes. Euler systems for GSp(4), 2021.
- [LSZ21b] David Loeffler, Christopher Skinner, and Sarah Livia Zerbes. An Euler system for GU(2, 1). Mathematische Annalen, 382(3-4):1091–1141, jul 2021.
 - [LW12] David Loeffler and Jared Weinstein. On the computation of local components of a newform. Mathematics of Computation, 81(278):1179–1200, 2012.
 - [MS14] Alberto Mínguez and Vincent Sécherre. Unramified l-modular representations of $GL_n(F)$ and its inner forms. International Mathematics Research Notices, 2014(8):2090–2118, 2014.
 - [Pra90] Dipendra Prasad. Trilinear forms for representations of GL(2) and local epsilon factors. Compositio Mathematica, 75(1):1–46, 1990.
- [Pra17] Dipendra Prasad. Ext-analogues of branching laws, 2017.
- [Ren10] David Renard. *Représentations des groupes réductifs p-adiques*. Société mathématique de France Paris, 2010.
- [Rub00] Karl Rubin. Annals of Mathematics Studies. Princeton University Press, 2000.
- [RV13] Dinakar Ramakrishnan and Robert J Valenza. Fourier analysis on number fields, volume 186. Springer Science & Business Media, 2013.
- [Sak08] Yiannis Sakellaridis. On the unramified spectrum of spherical varieties over p-adic fields. Compositio Mathematica, 144(4):978–1016, July 2008.
- [Sak13] Yiannis Sakellaridis. Spherical functions on spherical varieties, 2013.
- [Shi76] Takuro Shintani. On an explicit formula for class-1 "Whittaker Functions" on GL(n) over p-adic fields. Proceedings of the Japan Academy, 52(4):180–182, 1976.
- [Vig89] Marie-France Vignéras. Représentations modulaires de GL(2, F) en caractéristique l, f corps p-adique, $p \neq l$. Compositio Mathematica, 72(1):33–66, 1989.
- [Vig96] Marie France Vignéras. Représentations l-modulaires d'un groupe réductif p-adique avec $l \neq p$. (No Title), 1996.
- [Voe00] Vladimir Voevodsky. Triangulated categories of motives over a field. Cycles, transfers, and motivic homology theories, 143:188–238, 2000.

Mathematics Institute, Zeeman Building, University of Warwick, Coventry CV4 7AL, England.

Email address: Alexandros.Groutides@warwick.ac.uk