# IRREDUCIBLE SMOOTH REPRESENTATIONS IN DEFINING CHARACTERISTIC WITHOUT CENTRAL CHARACTER

### DANIEL LE

ABSTRACT. Let p > 3 and F be a non-archimedean local field with residue field a proper finite extension of  $\mathbb{F}_p$ . Let E be an algebraically closed countable field extension of the residue field of F. In this short note, we explain how the methods from [Le19, GLS23] can be used to construct irreducible smooth representations of  $\operatorname{GL}_n(F)$ , n > 1, over E without a central character.

#### 1. INTRODUCTION

Let p be a prime number. Smooth irreducible representations of p-adic reductive groups over fields of characteristic distinct from p have long been known to be admissible [Vig96, II.2.8]. One favorable feature of admissible representations over an algebraically closed field is that they satisfy Schur's lemma, namely their endomorphisms are scalars. In particular, these representations have central characters, as the center acts by endomorphisms. Turning to characteristic p representations, [AHHV17b, Question 1] asks if every irreducible smooth representation of a p-adic reductive group over a field of characteristic p is admissible. [AHHV17b, Question 2] asks if such a representation over an algebraically closed field necessarily has a central character. One may also ask whether every endomorphism of such a representation over an algebraically closed field is necessarily scalar. These questions have a positive answer for  $GL_2(\mathbb{Q}_p)$ [BL94, Bre03, Ber12]. One reason for the significance of central characters in characteristic p is that any representation with a Hecke eigenvalue for a Serre weight has central characters and the converse is true for GL<sub>2</sub> by [BL94, Theorem 33]. Hecke eigenvalues play an essential role in the classification theory of [AHHV17a].

Let p > 3, n > 1, and F a nonarchimedean local field with residue field a proper extension of  $\mathbb{F}_p$ . Recently, [GLS23], building on [Le19, GS20], constructed examples of mod p irreducible smooth representations of  $\operatorname{GL}_n(F)$  which are *not* admissible. In this short note, we use similar methods to give examples of irreducible smooth  $\operatorname{GL}_n(F)$ -representations without central characters and with non-scalar endomorphisms.

**Theorem 1.1.** Let n > 1 and p > 3 be a prime. Let F be a nonarchimedean local field with residue field a proper extension of  $\mathbb{F}_p$ . Let E be an extension of the residue field of F of countable cardinality.

- (1) There exists an irreducible smooth  $\operatorname{GL}_n(F)$ -representation over E which does not have a central character.
- (2) Let E/E be a field extension of countable cardinality. There exists an irreducible smooth  $GL_n(F)$ representation over E which has a central character and whose endomorphism algebra contains an E-subalgebra isomorphic to  $\tilde{E}$ .

**Remark 1.2.** (1) If E is algebraically closed, any representation satisfying the properties in Theorem 1.1(1) is necessarily not admissible.

- (2) If E/E is an infinite degree extension, any representation satisfying the properties in Theorem 1.1(2) is necessarily not admissible.
- (3) Taking E to be the residue field of F and E the algebraic closure of E recovers [GLS23, Theorem 1.2].

#### DANIEL LE

- (4) The hypothesis that the residue field of F is a proper extension of  $\mathbb{F}_p$  cannot be completely removed given that every irreducible smooth representation of  $\operatorname{GL}_2(\mathbb{Q}_p)$  is admissible.
- (5) If E is algebraically closed and uncountable, then every irreducible smooth representation of a p-adic reductive group satisfies Schur's lemma and thus has a central character. Indeed, a non-scalar endomorphism  $\varphi$  of such a representation  $\pi$  defines an action of a function field  $E(\varphi)$  in one variable which is of uncountable dimension over E. On the other hand, the dimension of  $\pi$ , which is the quotient of a compact induction of a finite-dimensional representation of a compact open subgroup, is of countable dimension over E.
- (6) As mentioned above, Theorem 1.1(1) implies the existence of an irreducible smooth  $\operatorname{GL}_n(F)$ representation over E without Hecke eigenvalues, answering [AHHV17b, Question 8] negatively. One might still ask whether irreducible smooth  $\operatorname{GL}_n(F)$ -representations over algebraically closed characteristic p fields with central characters have Hecke eigenvalues. As mentioned before, this is true for  $\operatorname{GL}_2(F)$  with F a nonarchimedean local field.

1.1. Acknowledgments. The key ideas in this note essentially appeared in [Le19], and we embarrassingly did not notice the application to Theorem 1.1 at that time. We thank Benjamin Schraen again for asking about endomorphism algebras years ago. We thank Eknath Ghate and Mihir Sheth for comments on previous drafts of this paper and for our earlier collaboration. The author was supported by the National Science Foundation under agreement DMS-2302623 and a start-up grant from Purdue University.

### 2. Some irreducible diagrams

Let p > 3 be a prime number. Let F be a nonarchimedean local field with residue field a proper extension of  $\mathbb{F}_p$  and uniformizer  $\varpi$ . We let  $G = \operatorname{GL}_2(F)$ ,  $K = \operatorname{GL}_2(\mathcal{O}_F)$ , and  $Z \subset G$  be the center of G. Let E be a field extension of the residue field of F. Let  $\tilde{E}$  be a field extension of E of countable degree. To construct the desired representation in the generality of Theorem 1.1, we will use constructions (and notation) from [GLS23] rather than [Le19, GS20]. The K-representation  $D_0$  in [GLS23, §2] has a model over the residue field of F and thus a model over E. This E[K]-representation is rigid of Loewy length 2 and has multiplicity free socle and cosocle. Its E[K]-socle has a decomposition

$$\sigma \oplus \sigma_1 \oplus \cdots \oplus \sigma_{l-1} \oplus \sigma'_1 \oplus \cdots \oplus \sigma'_{l-1}$$

into simple E[K]-modules. In this article, we will denote by  $D_0$  the tensor product of this *E*-model with  $\tilde{E}$  over *E*. By an abuse, We also denote by  $\sigma$ , etc. the E[K]-module which is the natural image of  $\sigma$ , etc. in  $D_0$ . Fix an  $\tilde{E}$ -valued character  $\xi$  of  $\mathbb{Z} \cong (\overline{\mathfrak{T}} \ 0 \ \varpi)^{\mathbb{Z}}$ . We extend the action on  $D_0$  to  $KZ \cong K \times (\overline{\mathfrak{T}} \ 0 \ \varpi)^{\mathbb{Z}}$  so that  $(\overline{\mathfrak{T}} \ 0 \ \varpi)^{\mathbb{Z}}$  acts by  $\xi$ .

Let  $I \subset K$  be the upper triangular Iwahori subgroup,  $I_1 \subset I$  the normal pro-p Sylow subgroup, and  $\Pi \in N_G(I)$  be  $\begin{pmatrix} 0 & 1 \\ \varpi & 0 \end{pmatrix}$ . We let  $D_1 = D_0^{I_1}$  which has a natural IZ-action. We extend this to an  $N_G(I)$ -action as follows. Let  $S_1 = (\operatorname{soc}_K D_0)^{I_1}$  and  $Q_1 = (\operatorname{cosoc}_K D_0)^{I_1}$  which are semisimple multiplicity-free  $\widetilde{E}[IZ]$ -representations. Moreover,  $D_1 \cong S_1 \oplus Q_1$ , and the twisted IZ-representation  $Q_1^{(\Pi)} = \{v^{(\Pi)} \mid v \in Q_1\}$ , with IZ-action given by  $g \cdot v^{(\Pi)} = (\Pi g \Pi^{-1} \cdot v)^{(\Pi)}$  for  $g \in IZ$  and  $v \in Q_1$ , is isomorphic to  $S_1$ . Choosing an  $\widetilde{E}[IZ]$ -isomorphism  $\psi : S_1 \cong Q_1^{(\Pi)}$ , there is a unique extension of the IZ-action on  $D_1$  to an  $N_G(I)$ -action such that  $v \mapsto (\Pi v)^{(\Pi)}$  is the map  $\psi$ .

Let  $D_0(\infty) = \bigoplus_{i \in \mathbb{Z}} D_0(i)$  be the smooth KZ-representation with component-wise KZ-action, where there is a fixed isomorphism  $D_0(i) \cong D_0$  of  $\widetilde{E}[KZ]$ -representations for every  $i \in \mathbb{Z}$ . We denote the natural inclusion  $D_0 \xrightarrow{\sim} D_0(i) \hookrightarrow D_0(\infty)$  by  $\iota_i$ , and write  $v_i = \iota_i(v)$  for  $v \in D_0$  for every  $i \in \mathbb{Z}$ . Let  $D_1(\infty) :=$  $D_0(\infty)^{I(1)} \cong \bigoplus_{i \in \mathbb{Z}} (S_1 \oplus Q_1)$ . We define a  $\Pi$ -action on  $D_1(\infty)$  as follows. Let  $\lambda = (\lambda_i) \in \prod_{i \in \mathbb{Z}} \widetilde{E}^{\times}$ . For all integers  $i \in \mathbb{Z}$ , define

$$\Pi v_{i} := \begin{cases} (\Pi v)_{i-1} & \text{if } v \in S_{1}^{\chi(\sigma_{1})}, \\ \lambda_{i}(\Pi v)_{i+1} & \text{if } v \in S_{1}^{\chi(\sigma'_{1})}, \\ (\Pi v)_{i} & \text{if } v \in S_{1}^{\chi} \text{ for } \chi \in \{\chi(\sigma), \chi(\sigma_{2}), \dots, \chi(\sigma_{l-1}), \chi(\sigma'_{2}), \dots, \chi(\sigma'_{l-1})\}. \end{cases}$$

This uniquely determines a smooth  $N_G(I)$ -action on  $D_1(\infty)$  such that  $\left( \begin{smallmatrix} \overline{0} & 0 \\ \overline{0} & \infty \end{smallmatrix} \right)^{\mathbb{Z}}$  acts by the character  $\xi$ . Thus we get a basic 0-diagram  $D(\lambda) := (D_0(\infty), D_1(\infty), \operatorname{can})$  (in the sense of [BP12, §9]) with the above actions where can is the canonical inclusion  $D_1(\infty) \hookrightarrow D_0(\infty)$ .

**Proposition 2.1.** If  $\lambda_0 \in E$ ,  $\lambda_i \neq \lambda_0$  for all  $i \neq 0$ , and the *E*-span of  $\{\prod_{j=0}^{m-1} \lambda_j \mid m \geq 1\}$  is  $\widetilde{E}$ , then the basic 0-diagram  $D(\lambda)$  is irreducible after restricting scalars to *E*.

*Proof.* The proof is a fiber product of the proofs of [GLS23, Proposition 3.1] and [Le19, Theorem 3.4]. Let  $W \subset D_0(\infty)$  be a non-zero E[KZ]-subrepresentation such that  $\Pi$  stabilizes  $W^{I(1)}$ . We claim that  $W = D_0(\infty)$ . We begin with a series of lemmata.

**Lemma 2.2.** If 
$$(c_i) \in \bigoplus_{i \in \mathbb{Z}} E$$
 such that  $(\sum_i c_i \iota_i) (\sigma) \subseteq W$ , then  $(\sum_i c_i \iota_{i-m}) (\sigma) \subseteq W$  and  
 $\left(\sum_i c_i \left(\prod_{j=0}^{m-1} \lambda_{i+j}\right) \iota_{i+m}\right) (\sigma) \subseteq W$ 

for all m > 0.

*Proof.* The II-action on  $(\sum_i c_i \iota_i) (S_1^{\chi(\sigma)})$  gives that  $(\sum_i c_i \iota_i) (Q_1^{\chi(\sigma)^s}) \subseteq W^{I(1)}$  which implies that

$$\left(\sum_{i} c_{i}\iota_{i}\right)\left(M(\sigma^{[s]})\right) \subseteq W$$

because  $M(\sigma^{[s]})$  is indecomposable. Hence

(2.3) 
$$\left(\sum_{i} c_{i}\iota_{i}\right)(\sigma_{1}\oplus\sigma_{1}')\subseteq W.$$

Now the  $\Pi$ -action on  $(\sum_i c_i \iota_i) (S_1^{\chi(\sigma_1)})$  and  $(\sum_i c_i \iota_i) (S_1^{\chi(\sigma'_1)})$  gives respectively

$$\left(\sum_{i} c_{i}\iota_{i-1}\right) \left(Q_{1}^{\chi(\sigma_{1})^{s}}\right) \subseteq W^{I(1)} \text{ and } \left(\sum_{i} \lambda_{i}c_{i}\iota_{i+1}\right) \left(Q_{1}^{\chi(\sigma_{1}')^{s}}\right) \subseteq W^{I(1)}.$$

Hence

$$\left(\sum_{i} c_{i}\iota_{i-1}\right) \left(E(\sigma_{2}, \sigma_{1}^{[s]})\right) \subseteq W \text{ and } \left(\sum_{i} \lambda_{i}c_{i}\iota_{i+1}\right) \left(E(\sigma_{2}', \sigma_{1}'^{[s]})\right) \subseteq W$$

so that

$$\left(\sum_{i} c_{i}\iota_{i-1}\right)(\sigma_{2}) \subseteq W \text{ and } \left(\sum_{i} \lambda_{i}c_{i}\iota_{i+1}\right)(\sigma_{2}) \subseteq W.$$

Acting by  $\Pi$  again and iterating a similar process shows that

$$\left(\sum_{i} c_{i}\iota_{i-1}\right)(\sigma) \subseteq W \text{ and } \left(\sum_{i} \lambda_{i}c_{i}\iota_{i+1}\right)(\sigma) \subseteq W$$

The lemma now follows from induction.

Lemma 2.4.  $\operatorname{Hom}_{E[K]}(\sigma, W \cap \iota_0(D_0)) \neq 0.$ 

#### DANIEL LE

Proof. We have  $\operatorname{Hom}_{E[K]}(\tau, W) \neq 0$  for some  $\tau \in \operatorname{soc}_{E[K]}D_0$ . If  $\tau = \sigma_m$  (resp.  $\tau = \sigma'_m$ ) with  $1 \leq m < l-1$ , then we have  $\operatorname{Hom}_{E[K]}(\kappa, W) \neq 0$  for  $\kappa = \sigma_{m+1}$  (resp.  $\kappa = \sigma'_{m+1}$ ) as in the proof of Lemma 2.2. Similarly, if  $\tau = \sigma_{l-1}$  or  $\sigma'_{l-1}$ , then  $\operatorname{Hom}_{E[K]}(\sigma, W) \neq 0$ . Without loss of generality, we can assume that  $\tau = \sigma$ . Thus there exists a non-zero  $(c_i) \in \bigoplus_{i \in \mathbb{Z}} \widetilde{E}$  such that

(2.5) 
$$\left(\sum_{i} c_{i}\iota_{i}\right)(\sigma) \subset W.$$

Suppose that  $\#(c_i) := \#\{i \in \mathbb{Z} \mid c_i \neq 0\}$  is minimal amongst nonzero  $(c_i) \in \bigoplus_{i \in \mathbb{Z}} \tilde{E}$  satisfying (2.5). By Lemma 2.2 we can assume without loss of generality that  $c_0$  is nonzero. Lemma 2.2 implies that  $(\sum_i \lambda_i c_i \iota_i)(\sigma) \subset W$ . Using that  $\lambda_0 \in E$ , we have that  $(\sum_i (\lambda_i - \lambda_0)c_i \iota_i)(\sigma) \subset W$ . Since  $\lambda_i - \lambda_0 \neq 0$  for all  $i \neq 0$ ,  $\#((\lambda_i - \lambda_0)c_i) < \#(c_i)$ . Minimality of  $\#(c_i)$  implies that  $c_i = 0$  for  $i \neq 0$ .

**Lemma 2.6.** W contains the  $\sigma$ -isotypic part of  $\iota_0(D_0)$ .

*Proof.* We have that  $c_0\iota_0(\sigma) \subset W$  for some  $c_0 \in \widetilde{E}^{\times}$ . By Lemma 2.2, we also have that  $(\prod_{j=0}^{m-1} \lambda_j)c_0\iota_0(\sigma) \subset W$  for all  $m \geq 1$ . Since the *E*-span of  $\{(\prod_{j=0}^{m-1} \lambda_j)c_0 \mid m \geq 1\}$  is  $\widetilde{E}$ , we are done.  $\Box$ 

Finally, we have that  $\iota_i(D_0) \subset W$  for all  $i \in \mathbb{Z}$  arguing as in the proof of Lemma 2.2 so that  $W = D_0(\infty)$ .

## 3. Proof of Theorem 1.1

**Theorem 3.1.** There is an E[G]-representation  $\pi$  and an injection of diagrams  $(D_0(\infty), D_1(\infty), \operatorname{can}) \hookrightarrow (\pi|_{KZ}, \pi|_{N_G(I)}, \operatorname{can})$  such that the map  $D_0(\infty) \hookrightarrow \pi|_{KZ}$  induces an isomorphism on K-socles and  $D_0(\infty)$  generates  $\pi$  as an E[G]-representation. Moreover, such a  $\pi$  is an irreducible smooth E[G]-representation (by restriction of scalars).

Proof. The first part is proved as in [Le19, Theorem 3.2], noting that the E[G]-representation generated by  $D_0(\infty)$  is the  $\widetilde{E}[G]$ -representation generated by  $D_0(\infty)$  as  $D_0(\infty)$  is an  $\widetilde{E}$ -vector space. If  $\pi' \subset \pi$  is a nonzero E[G]-subrepresentation, then  $(\pi'|_{KZ} \cap D_0(\infty), \pi'|_{KZ} \cap D_1(\infty), \operatorname{can}) \subset (D_0(\infty), D_1(\infty), \operatorname{can})$  is a nonzero subdiagram over E (restricting scalars). Indeed,  $\pi'|_{KZ} \cap D_0(\infty)$  contains the E[K]-socle of  $\pi'|_K$ . Then we have that  $\pi'|_{KZ}$  contains  $D_0(\infty)$  by Proposition 2.1 so that  $\pi'$  contains the E[G]-representation generated by  $D_0(\infty)$ , which is  $\pi$ .

**Theorem 3.2.** Let n > 1 and p > 3 be a prime. Let F be a nonarchimedean local field with residue field a proper extension of  $\mathbb{F}_p$ . Let  $E \subset \widetilde{E}$  be extensions of the residue field of F of countable cardinality. Then there exists an irreducible smooth  $\operatorname{GL}_n(F)$ -representation over E whose endomorphism algebra contains an E-subalgebra isomorphic to  $\widetilde{E}$ . Moreover, there exists such a representation with a central character, and if  $E \subsetneq \widetilde{E}$ , there exist such representations without a central character.

Proof. We can choose

$$(\lambda_i)_{i\in\mathbb{Z}}\in\prod_{i\in\mathbb{Z}}\widetilde{E}^{\times}$$

so that  $\lambda_0 \in E$ ,  $\lambda_i \neq \lambda_0$  for all  $i \neq 0$ , and the *E*-span of  $\{\prod_{j=0}^{m-1} \lambda_j \mid m \geq 1\}$  is  $\tilde{E}$ . For example, we can choose an *E*-basis  $(\mu_i)_{i\geq 0}$  for  $\tilde{E}$  with  $\mu_0 = 1$  and set  $\lambda_0 = 1$ ,  $\lambda_i = \mu_i/\mu_{i-1}$  for all i > 0, and  $\lambda_i = -1$  for all i < 0. Let  $\rho$  be as in Theorem 3.1. Then  $\rho$  is a smooth irreducible E[G]-representation. Let  $\overline{P}$  denote the lower block triangular parabolic subgroup of  $\operatorname{GL}_n$  with blocks of size 2, 1, ..., 1. By the proof of [GLS23, Lemma 4.5], there is a character of  $(F^{\times})^{n-2} \to E^{\times}$  such that  $\pi = \operatorname{Ind}_{\overline{P}(F)}^{\operatorname{GL}_n(F)} \rho \otimes \chi$  is a smooth irreducible  $\operatorname{GL}_n(F)$ -representation. Then  $\tilde{E}$  acts on  $\rho$  by E[G]-endomorphisms, and thus on  $\pi$  by  $E[\operatorname{GL}_n(F)]$ -endomorphisms by functoriality. If  $\xi ( {\mathbb{T}}_0 {\mathbb{T}}_\infty ) \in E^{\times}$ , then  $\rho$  and thus  $\pi$  has a central character. If  $\xi ( {\mathbb{T}}_0 {\mathbb{T}}_\infty ) \in \tilde{E}^{\times}$  is not in  $E^{\times}$ , then  $\varpi I_n \in \operatorname{GL}_n(F)$  does not act on  $\pi$  by a scalar in E.

IRREDUCIBLE SMOOTH REPRESENTATIONS IN DEFINING CHARACTERISTIC WITHOUT CENTRAL CHARACTER5

Proof of Theorem 1.1. Theorem 1.1(2) follows immediately from Theorem 3.2. To prove Theorem 1.1(1), we need only find a proper extension  $\tilde{E}$  of E which is countable. For example, one could take the field of rational functions in one variable over E.

**Remark 3.3.** One can show that the endomorphism ring of the representation  $\pi$  in the proof of Theorem 3.2 is  $\tilde{E}$ .

### References

- [AHHV17a] N. Abe, G. Henniart, F. Herzig, and M.-F. Vignéras, A classification of irreducible admissible mod p representations of p-adic reductive groups, J. Amer. Math. Soc. 30 (2017), no. 2, 495–559. MR 3600042
- [AHHV17b] Noriyuki Abe, Guy Henniart, Florian Herzig, and Marie-France Vigneras, Questions on mod p representations of reductive p-adic groups, 2017, available at https://arxiv.org/abs/1703.02063.
- [Ber12] Laurent Berger, Central characters for smooth irreducible modular representations of  $GL_2(\mathbf{Q}_p)$ , Rend. Semin. Mat. Univ. Padova **128** (2012), 1–6 (2013). MR 3076828
- [BL94] L. Barthel and R. Livné, Irreducible modular representations of GL<sub>2</sub> of a local field, Duke Math. J. 75 (1994), no. 2, 261–292. MR 1290194
- [BP12] Christophe Breuil and Vytautas Paškūnas, Towards a modulo p Langlands correspondence for GL<sub>2</sub>, Mem. Amer. Math. Soc. 216 (2012), no. 1016, vi+114. MR 2931521
- [Bre03] Christophe Breuil, Sur quelques représentations modulaires et p-adiques de  $GL_2(\mathbf{Q}_p)$ . I, Compositio Math. **138** (2003), no. 2, 165–188. MR 2018825
- [GLS23] Eknath Ghate, Daniel Le, and Mihir Sheth, Non-admissible irreducible representations of p-adic  $GL_n$  in characteristic p, Represent. Theory **27** (2023), 1088–1101. MR 4664337
- [GS20] Eknath Ghate and Mihir Sheth, On non-admissible irreducible modulo p representations of  $GL_2(\mathbb{Q}_{p^2})$ , C. R. Math. Acad. Sci. Paris **358** (2020), no. 5, 627–632. MR 4149863
- [Le19] Daniel Le, On some nonadmissible smooth irreducible representations for GL<sub>2</sub>, Math. Res. Lett. 26 (2019), no. 6, 1747–1758. MR 4078693
- $[Vig96] Marie-France Vignéras, Représentations l-modulaires d'un groupe réductif p-adique avec l \neq p, Progress in Mathematics, vol. 137, Birkhäuser Boston, Inc., Boston, MA, 1996. MR 1395151$