Bases for Washington's cyclotomic units of real cyclotomic fields and totally deployed fields

SOUANEF Rafik *

SOUANEF RafikJuly 3, 2024AbstractWe aim to present families of generators with minimal cardinality - we call such families bases - of
the free abelian group $Was(\mathbb{K})/\mathbb{Z}(\mathbb{K})$ whenever \mathbb{K} is a real cyclotomic field $\mathbb{Q}(\zeta_n)^+$ or \mathbb{K} is a totally
deployed abelian number field. Here, $Was(\mathbb{K})$ refers to the group of Washington's cyclotomic units
of \mathbb{K} and $\mathbb{Z}(\mathbb{K})$ refers to the group of roots of unity lying in \mathbb{K} .Introductioncyclotomic units are of interest because of their link with the theory of \mathbb{Z}_p -extensions. For instance,
the main conjecture of Iwasawa's theory can be vaguely formulated in this way: a module that one can
construct using cyclotomic units has the same characteristic ideal as the standard *p*-ramified module of

 \sum construct using cyclotomic units has the same characteristic ideal as the standard *p*-ramified module of \eth wasawa's theory that one can construct with some extensions that satisfy some ramification related con-Litions (see [1], proposition 4.5.7). Another reason why cyclotomic units are of interest is to approximate the whole group of units of any abelian number field. There are many different types of cyclotomic units (see [7]) but in this article we will deal with two types only that are the cyclotomic units of Washington tand those of Sinnott. Washington's cyclotomic units are defined through Galois invariants and Sinnott's circular units are defined by explicit generators that generate a subgroup of the group given by Washington: the drawback of having a smaller group - so that we may expect it to be a worse approximation of > the group of units - is balanced by a better knowledge of the elements of this group. These two types of Exception of the constructed through two processes that have a common starting point that is to define in the same way cyclotomic units of cyclotomic fields and then deduce a definition for cyclotomic units of all abelian number fields. A crucial article in the study of Sinnott's circular units is [12] in which it is proven that the group of Sinnott's circular units has finite index in the group of units and that this index is somehow linked to a real class number. On the other side, the group of Washington's cyclotomic

^{*}Université de Franche-Comté, CNRS, LmB (UMR 6623), 16 route de Gray, 25000 Besançon, France Email: rafik.souanef@ens-rennes.fr, Url: https://perso.eleves.ens-rennes.fr/people/rafik.souanef/

units remains quite mysterious. There are two articles ([5] and [15]) in which these last units has been studied under some hypothesis on the considered number fields. These last two articles and the work that we present today aim to give us a better understanding of this group by giving explicit system of generators with no Z-relations modulo roots of unity (we then say "basis"). What is new in our work is that we use different hypothesis. For example, we do not consider real fields only. In our work, we consider totally deployed abelian number fields, that is fields of the form

$$\mathbb{K} = \mathbb{K}_1 \cdots \mathbb{K}_n$$

with $\mathbb{K}_i \subset \mathbb{Q}(\zeta_{p_i^{e_i}})$ for some prime numbers p_i and some integers e_i .

More precisely, let \mathbb{K} be a number field. Suppose it is abelian, that is \mathbb{K}/\mathbb{Q} is Galois and its Galois group is abelian. Then, recall that Kronecker-Weber theorem states there is an integer n such that $\mathbb{K} \subset \mathbb{Q}(\exp(2i\pi/n))$. Define the conductor of \mathbb{K} to be the least integer n that satisfies this last condition. From now on, suppose \mathbb{K} is an abelian number field with conductor n. Let $\zeta_n = \exp(2i\pi/n)$.

Let $\mathbf{Z}(\mathbb{K})$ denote the group of roots of unity of \mathbb{K} . Let $\mathbf{Was}(\mathbb{K})$ denote the Galois module of Washington's cyclotomic units and let $\mathbf{Sin}(\mathbb{K})$ denote the Galois module of Sinnott's circular units (we will recall their definition later).

By abuse of language, we will rather talk about bases of $Was(\mathbb{K})$ instead of $Was(\mathbb{K})/\mathbb{Z}(\mathbb{K})$.

If $\mathbb{K} = \mathbb{Q}(\zeta_n)$ is a cyclotomic field, recall Gold and Kim have given bases of $\mathbf{Was}(\mathbb{Q}(\zeta_n)) = \mathbf{Sin}(\mathbb{Q}(\zeta_n))$ (see [2], theorem 2).

Based on this work, we state theorems 6, 8 and 14 that all describe bases of $\mathbf{Was}(\mathbb{K})$ under different hypotheses on \mathbb{K} . The first two theorems are easy consequences of proposition 5 that is itself a consequence of proposition 1 more or less and these two theorems give bases assuming $\mathbb{K} = \mathbb{Q}(\zeta_n)^+$ is a real cyclotomic field. In theorem 14, we suppose \mathbb{K} is totally deployed and the principal idea in the proof of this theorem is to show we have a basis by proving the group that is generated by the elements we consider is a direct factor of $\mathbf{Was}(\mathbb{Q}(\zeta_n))$. This idea has also been used in [5] and [15].

Divisibility relations arise from this last basis (see corollary 21).

2 Notation and preliminaries

2.1 On units

Let A be a Galois module, that is A is an abelian group with some $\operatorname{Gal}(\mathbb{K}/\mathbb{Q})$ acting \mathbb{Z} -linearly on it (we consider extensions of \mathbb{Q} only). Suppose \mathbb{K}/\mathbb{Q} is an abelian extension, so that complex conjugation is well defined as an element of $\operatorname{Gal}(\mathbb{K}/\mathbb{Q})$. We let A^+ denote the Galois submodule of A that consists of all the elements A on which the complex conjugation acts trivially. Later on, we will consider $A = \mathcal{O}_{\mathbb{K}}^{\times}$ the group of units of the ring of integers of \mathbb{K} .

If $x \in A$, then $u \in \mathbb{Z}[\operatorname{Gal}(\mathbb{K}/\mathbb{Q})]$ acts on x and we denote by ux or u(x) or x^u the image of x under u.

Let $\zeta_n = \exp(2i\pi/n)$ for all $n \in \mathbb{N}^*$. For now on, let $n \ge 2$ satisfying $n \ne 2 \mod 4$ (with no loss of generality because $\mathbb{Q}(\zeta_n) = \mathbb{Q}(\zeta_{2n})$ if n is odd). If $p \in \mathbb{P}$ is a prime number, let e_p denote the p-valuation of n and, more generally, let $v_p(k)$ denote that of any integer k. Let $n = \prod_{j=1}^r p_j^{e_j}$ and let $q_j = p_j^{e_j}$.

We now recall that if n is not a prime power, then $1 - \zeta_n$ is a unit of the ring of integers of $\mathbb{Q}(\zeta_n)$ (see [14] proposition 2.8). Now, if n is a prime power, then $1 - \zeta_n$ is no longer a unit but $\frac{1-\zeta_n^{\sigma}}{1-\zeta_n}$ is a unit for all $\sigma \in \operatorname{Gal}(\mathbb{Q}(\zeta_n)/\mathbb{Q})$ (lemma 1.3 [14]).

Let \mathbb{K} be an abelian number field of conductor n.

We say \mathbb{K} is totally deployed when $\operatorname{Gal}(\mathbb{K}/\mathbb{Q})$ is the direct product of its inertia subgroups (see the introduction of [3]). This condition is equivalent to writting

$$\mathbb{K} = \mathbb{K}_1 \cdots \mathbb{K}_r$$

with $\mathbb{K}_j \subset \mathbb{Q}(\zeta_{q_j})$.

Let $\mathbf{E}(\mathbb{K})$ be the group of units of the ring of integers $\mathcal{O}_{\mathbb{K}}$ of \mathbb{K} . Let \mathcal{C}_n be the Galois module generated by the roots of unity of $\mathbb{Q}(\zeta_n)$ and by the $1 - \zeta_d$ for $d \mid n$. Let $\mathbf{Was}(\mathbb{K}) = \mathbf{E}(\mathbb{K}) \cap \mathcal{C}_n$ and let $\mathbf{Sin}(\mathbb{K})$ be the intersection of $\mathbf{E}(\mathbb{K})$ with the Galois module generated by the roots of unity lying in \mathbb{K} and the

$$N_{\mathbb{Q}(\zeta_d)/\mathbb{K}\cap\mathbb{Q}(\zeta_d)}(1-\zeta_d)$$

where $d \in \mathbb{N}$.

One can show (see [6]) $Sin(\mathbb{K})$ is generated by :

-the roots of unity of \mathbb{K} , which form a group that we will denote by $\mathbf{Z}(\mathbb{K})$

-the $N_{\mathbb{Q}(\zeta_d)/\mathbb{K}\cap\mathbb{Q}(\zeta_d)}(1-\zeta_d^{\sigma})$ with $d \mid n$ such that d is not a prime power and $d \wedge (n/d) = 1$ and $\sigma \in Gal(\mathbb{Q}(\zeta_d)/\mathbb{Q})$

-the $N_{\mathbb{Q}(\zeta_d)/\mathbb{K}\cap\mathbb{Q}(\zeta_d)}(1-\zeta_d)^{1-\sigma}$ with d being a prime power dividing n such that $d \wedge (n/d) = 1$ and with $\sigma \in \text{Gal}(\mathbb{Q}(\zeta_d)/\mathbb{Q})$.

It is known that both $Was(\mathbb{K})$ and $Sin(\mathbb{K})$ have finite index in $E(\mathbb{K})$, that is they both have maximal rank as \mathbb{Z} -submodules of $E(\mathbb{K})$ and that their index is linked to the class number of the maximal real subfield \mathbb{K}^+ of \mathbb{K} (see [12]).

When the situation makes it clear, we will omit writting \mathbb{K} . For example, we will write **Was** instead of writting **Was**(\mathbb{K}).

We now recall the following relations (see [13] lemma 2.1):

$$1 - \zeta_n^a = -\zeta_n^a (1 - \zeta_n^{-a})$$
 (1)

$$N_{\mathbb{Q}(\zeta_n)/\mathbb{Q}(\zeta_d)}(1-\zeta_n) = \left(\prod_{\substack{p|n\\p \nmid d}} (1-\operatorname{Frob}(p,d)^{-1})\right) (1-\zeta_d)$$

where $d \mid n$, the integer p is prime and $\operatorname{Frob}(p, d)$ denotes the Frobenius of $\mathbb{Q}(\zeta_d)$ that is defined by $\zeta_d \mapsto \zeta_d^p$. We will refer to this second relation as "norm relation".

We recall a property of Hasse's index.

Proposition 1. We have

$$[\mathbf{E} : \mathbf{ZE}^+] \in \{1; 2\}.$$

Moreover, if $\mathbb{K} = \mathbb{Q}(\zeta_n)$, this index is 1 if and only if n is a prime power.

Proof. See [14] theorem 4.12 and corollary 4.13.

We also recall Dirichlet's units theorem.

Theorem 2. The group $\mathbf{E}(\mathbb{K})$ is isomorphic to the product of $\mathbf{Z}(\mathbb{K})$ and a free abelian group of rank $r_1 + r_2 - 1$ where r_1 denotes the number of real embeddings of \mathbb{K} and r_2 denotes the number of complex embeddings of \mathbb{K} up to conjugation.

We now introduce the notation we will use to work with bases of **Was** (most of this notation comes from [2] and [15]).

Recall $n = \prod_{j=1}^{r} p_j^{e_j}$ and $q_j = p_j^{e_j}$. If p_j is odd, let σ_j be a generator of $\operatorname{Gal}(\mathbb{Q}(\zeta_{q_j})/\mathbb{Q})$. If $p_j = 2$, let J_j be the complex conjugation considered as an element of $\operatorname{Gal}(\mathbb{Q}(\zeta_{q_j})/\mathbb{Q})$ and let $\tilde{\sigma}_j$ be so that this last Galois group is generated by $\tilde{\sigma}_j$ and J_j . Let

$$\sigma_j^k = \begin{cases} \widetilde{\sigma}_j^k & \text{if } 0 \leqslant k < 2^{e_j - 2} \\ \widetilde{\sigma}_j^k J_j & \text{if } 2^{e_j - 2} \leqslant k < 2^{e_j - 1} \end{cases}$$

See the elements of $\operatorname{Gal}(\mathbb{Q}(\zeta_{q_j})/\mathbb{Q})$ as elements of $\operatorname{Gal}(\mathbb{Q}(\zeta_n)/\mathbb{Q})$ by letting them acting trivially on $\mathbb{Q}(\zeta_{n/q_j})$. Let J_j be the complexe conjugation of $\operatorname{Gal}(\mathbb{Q}(\zeta_{q_j})/\mathbb{Q})$ considered as an element of $\operatorname{Gal}(\mathbb{Q}(\zeta_n)/\mathbb{Q})$. Let $J = J_1 \cdots J_r$ be the complex conjugation considered as an element of $\operatorname{Gal}(\mathbb{Q}(\zeta_n)/\mathbb{Q})$.

Let $a \in \mathbb{Z}$. Define (see [14] lemma 8.1)

$$\xi_{q_j,a} = \sqrt{\zeta_{q_j}^{1-\sigma_j^a}} \frac{1-\zeta_{q_j}^{\sigma_j^a}}{1-\zeta_{q_j}} \in \mathbf{Was}^+(\mathbb{Q}(\zeta_{q_j})).$$

We recall the basis of $\mathbf{Was}(\mathbb{Q}(\zeta_n))$ Gold and Kim described in [2] (theorem 2). **Theorem 3.** For all $\Omega = \{i_1, \ldots, i_s\} \subset [\![1, r]\!]$, let

$$n_{\Omega} = \prod_{j \in \Omega} p_j^{e_j}$$
$$\zeta_{\Omega} = \zeta_{n_{\Omega}}.$$

For all $\Omega = \{j\} \subset \llbracket 1, r \rrbracket$, let

$$X_{\Omega}^{1} = \{a \in \mathbb{N} : 1 \leq a < \varphi(q_{j})/2\}$$
$$B_{\Omega} = \{\xi_{q_{j},a} : 1 \leq a < \varphi(q_{j})/2\}.$$

For all $\Omega = \{i_1, \ldots, i_s\} \subset [\![1, r]\!]$ that is neither empty nor a singleton and such that $i_1 < \cdots < i_s$ and for all $0 \leq k \leq s$ let X_{Ω}^k be the set of all tuples $(a_1, \ldots, a_s) \in \mathbb{N}^s$ such that

$$\begin{cases} 1 \leqslant a_j < \varphi(q_{i_j}) & \text{if } j \in \llbracket 1, k \llbracket \\ 1 \leqslant a_k < \frac{\varphi(q_{i_k})}{2} \\ a_j = 0 & \text{if } j \in \llbracket k, s \rrbracket. \end{cases}$$

Let

$$B_{\Omega}^{k} = \left\{ \left(\prod_{j=1}^{s} \sigma_{i_{j}}^{a_{j}}\right) (1 - \zeta_{\Omega}) : (a_{1}, \dots, a_{s}) \in X_{\Omega}^{k} \right\}.$$

For k = 0 let

$$X_{\Omega}^{k} = \{(a_{1}, \dots, a_{s}) \in \mathbb{N}^{s} : \forall j \in [[1, s]], a_{j} = 0\}$$

and let B^k_Ω be defined in a similar way. Then, let

$$B_{\Omega} = \begin{cases} \bigcup_{\substack{k=0 \\ s}}^{s} B_{\Omega}^{k} & \text{if } 2 \mid |\Omega| \\ \bigcup_{k=1}^{k} B_{\Omega}^{k} & \text{if } 2 \nmid |\Omega|. \end{cases}$$

The family $\cup_{\Omega} B_{\Omega}$ - where Ω runs over the set of all non empty subsets of $\llbracket 1, r \rrbracket$ - is a basis of $Was(\mathbb{Q}(\zeta_n))$. *Proof.* See [2] (see theorem 2).

We will keep in mind that there is a one-to-one correspondence between all the elements of Gold and Kim's basis and all the tuples $(\Omega, a_1, \ldots, a_s)$. We say Ω or n_{Ω} is the level of x.

In the proof of theorem 14, we will order those tuples in the following way:

$$(\Omega_1, a_1, \dots, a_{s_1}) \leqslant (\Omega_2, b_1, \dots, b_{s_2}) \iff \begin{cases} (\Omega_1 \not\subset \Omega_2 \text{ and } \Omega_2 \not\subset \Omega_1 \text{ and } |\Omega_1| \geqslant |\Omega_2|) & \text{or} \\ (\Omega_2 \subsetneq \Omega_1) & \text{or} \\ (\Omega_1 = \Omega_2 \text{ and } (a_1, \dots, a_{s_1}) \leq_L (b_1, \dots, b_{s_2})) \end{cases}$$

where \leq_L denotes the inverse lexicographic order associated to the natural order on \mathbb{N} - that is we compare integers starting from last index and ending with first index. Here, let $s = s_1 = s_2$ so we may have $a_s < b_s$ or $a_s = b_s$ and $a_{s-1} < b_{s-1}$ or ... In particular, note this relation is not a partial order.

In what's next, we will need the following notation. Let

$$\mathbb{L}_n = \mathbb{Q}(\zeta_{q_1})^+ \cdots \mathbb{Q}(\zeta_{q_r})^+.$$

There is a root of unity $\eta \in \mathbb{Q}(\zeta_n)$ (see [15] 2-ii) such that

$$\eta_n := \eta \operatorname{N}_{\mathbb{Q}(\zeta_n)/\mathbb{Q}(\zeta_{q_1})^+ \cdots \mathbb{Q}(\zeta_{q_{r-1}})^+ \mathbb{Q}(\zeta_{q_r})} (1-\zeta_n) \in \mathbb{Q}(\zeta_{q_1})^+ \cdots \mathbb{Q}(\zeta_{q_r})^+.$$

For all $\mathbb{L} \subset \mathbb{L}_n$ of conductor n, let

$$e_{\mathbb{L}} = \mathcal{N}_{\mathbb{L}_n/\mathbb{L}}(\eta_n) \in \mathbb{L}.$$

We also define similar objects by swapping n with any of its divisors.

2.2 On convolution product

Through this section, we recall - in the needed context only - several facts that are stated in a more general context in [10] and [4] and that deal with Möbius functions.

Let E be a finite set. Define $\mathcal{F}(E)$ to be the set of functions

$$f: \mathcal{P}(E) \longrightarrow \mathbb{C}$$
.

This set has a law of addition and a convolution product definied in the following way

$$\forall f,g \in \mathcal{F}(E), \ \forall \Omega \subset E, \quad f \ast g(\Omega) = \sum_{X \subset \Omega} f(X)g(\Omega \setminus X).$$

One can show $(\mathcal{F}(E), +, *)$ is a ring whose identity element is the function that maps \emptyset to 1 and all subset $\Omega \neq \emptyset$ to 0.

Denote by 1 the element of $\mathcal{F}(E)$ that maps all $\Omega \subset E$ to 1.

One can show 1 is a unit and we let μ denote its inverse. We have (see [4] equation 3.3)

$$\forall \Omega \subset E, \quad \mu(\Omega) = (-1)^{|\Omega|}.$$

In particular, we have the following theorem.

Theorem 4. Let $f, g \in \mathcal{F}(E)$. We have

$$\left(\forall \Omega \subset E, \ \sum_{X \subset \Omega} f(X) = g(\Omega)\right) \Longleftrightarrow \left(\forall \Omega \subset E, \ f(\Omega) = \sum_{X \subset \Omega} (-1)^{|\Omega| - |X|} g(X)\right).$$

Proof. See [10] proposition 2.

Later, we will use this convolution product with $E = \llbracket 1, r \rrbracket$.

3 From nonreal fields to real fields

In this section, we aim to give bases of $\mathbf{Was}^+(\mathbb{Q}(\zeta_n))$ (recall we talk about $\mathbf{Was}^+(\mathbb{Q}(\zeta_n))$ instead of talking about the quotient $\mathbf{Was}^+(\mathbb{Q}(\zeta_n))/\mathbb{Z}^+(\mathbb{Q}(\zeta_n))$). More precisely, for any abelian number field \mathbb{K} , we give a way to construct a basis of $\mathbf{Was}^+(\mathbb{K})$ given a basis of $\mathbf{Was}(\mathbb{K})$ (proposition 5) and we then apply this method when \mathbb{K} is a cyclotomic field (theorems 6 and 8).

3.1 Abelian fields

Proposition 5. Let \mathbb{K} be an abelian number field. Let (x_1, \ldots, x_r) be a \mathbb{Z} -basis of $\mathbf{Was}(\mathbb{K})$. With no loss of generality, suppose there is $r' \in [0; r]$ such that $x_1, \ldots, x_{r'}$ have order 2 in the quotient group \mathbf{E}/\mathbf{ZE}^+ and $x_{r'+1}, \ldots, x_r$ have order 1. Then, the family $(|x_1||x_1|, \ldots, |x_1||x_{r'}|, |x_{r'+1}|, \ldots, |x_r|)$ is a basis of $\mathbf{Was}(\mathbb{K}^+)$.

Proof. First, if $x \in \mathbf{ZE}^+$, observe we have $|x| \in \mathbf{E}^+$ and, if we also suppose $x \in \mathbf{Was}$, then $|x| \in \mathbf{Was}^+$. Indeed, write $x = zu \in \mathbf{ZE}^+$. Then, we have $|x| = \pm u$ and this proves the first statement. Now, suppose we also have $x \in \mathbf{Was}$. Then, we have $u = z^{-1}x \in \mathbf{Was} \cap \mathbf{E}^+$ and this proves the second statement. Hence, the family $(|x_1||x_1|, \ldots, |x_1||x_{r'}|, |x_{r'+1}|, \ldots, |x_r|)$ is made of elements of \mathbf{Was}^+ . Now, let us show these elements are generators of \mathbf{Was}^+ .

Let $x \in \mathbf{Was}^+$ and write

$$x = \zeta \prod_{i=1}^{r} x_i^{a_i}$$

for some $a_i \in \mathbb{Z}$, $\zeta \in \mathbb{Z}$. In particular, we have $x \in \mathbb{E}^+$ so there is an even number of elements of the form x_i with $i \leq r'$, that is we have:

$$\sum_{i=1}^{r'} a_i \in 2\mathbb{Z}.$$

Let

$$A = a_1 - \sum_{i=2}^{r'} a_i \in 2\mathbb{Z}.$$

Therefore, we have

$$x = \zeta x_1^A \left(\prod_{i=2}^{r'} (x_1 x_i)^{a_i} \right) \left(\prod_{i>r'} x_i^{a_i} \right)$$
$$= \zeta' |x_1|^A \left(\prod_{i=2}^{r'} |x_1 x_i|^{a_i} \right) \left(\prod_{i>r'} |x_i|^{a_i} \right)$$

for some root of unity ζ' . As we have $x \in \mathbf{E}^+$ and $|x_i| \in \mathbb{R}$, we have $\zeta' = \pm 1$, which proves the considered family is a generating family. It is a basis because of Dirichlet's theorem.

3.2 Cyclotomic field with odd conductor

Theorem 6. Suppose *n* is odd. A basis of $Was^+(\mathbb{Q}(\zeta_n))$ is given by:

-the $|\xi_{p^{e_p},a}|$'s with $1 \leq a < \varphi(p^{e_p})/2$ and p running over the set of all prime divisors of n

-the $|1 - \zeta_n^{\sigma_1}| |x|$'s with $\Omega \subset [[1, r]], |\Omega| \ge 2$ and $x \in B_{\Omega}$.

Proof. We will apply proposition 5 to the basis of $Was(\mathbb{Q}(\zeta_n))$ given in theorem 3.

We have $\xi_{p^{e_p},a} \in \mathbf{Z}(\mathbb{Q}(\zeta_{p^{e_p}}))\mathbf{E}^+(\mathbb{Q}(\zeta_{p^{e_p}}))$ from proposition 1. For the other generators, observe we have, for all divisor $d \mid n$ and for all $a \in \mathbb{Z}$

$$1 - \zeta_d^a = \zeta_{2d}^a (\zeta_{2d}^{-a} - \zeta_{2d}^a).$$

As d is odd, we have $\zeta_{2d} \in \mathbb{Q}(\zeta_d)$, so that the previous decomposition takes place in $\mathbb{Q}(\zeta_d)$. Moreover, we have $(\zeta_{2d}^{-a} - \zeta_{2d}^{a}) = \pm i |1 - \zeta_d^{a}|$. We will keep the following decomposition in mind

$$1 - \zeta_d^a = \pm i \zeta_{2d}^a |1 - \zeta_d^a|.$$
⁽²⁾

which shows that $1 - \zeta_d^a$ has order 2 in the quotient group $\mathbf{E}(\mathbb{Q}(\zeta_n))/\mathbf{Z}(\mathbb{Q}(\zeta_n))\mathbf{E}^+(\mathbb{Q}(\zeta_n))$ whenever a is prime to d, otherwise we would have $i \in \mathbb{Q}(\zeta_n)$ and that is not the case.

Remark 7. We can understand the fact that there are products of pairwise cyclotomic units only when the conductor n is not a prime a power through proposition 1.

3.3 Cyclotomic field with even conductor

Theorem 8. Suppose *n* is even and write $n = 2^{e_1} p_2^{e_2} \cdots p_r^{e_r} \in \mathbb{N}$. Let $m = p_2^{e_2} \cdots p_r^{e_r}$. A basis of $\mathbf{Was}^+(\mathbb{Q}(\zeta_n))$ is given by:

-the $|\xi_{p^{e_p},a}|$'s with $1 \leq a < \varphi(p^{e_p})/2$ and p being a prime divisor of n (we will say these generators have type 0)

-the |x|'s where $\Omega \subset \llbracket 1, r \rrbracket$ satisfies $|\Omega| \ge 2$ and $x \in B_{\Omega}$ has odd level d (those have type 1)

-the $|1 - \zeta_n^{\sigma_1}||x|$'s where $\Omega \subset [\![1, r]\!]$ satisfies $|\Omega| \ge 2$ and $x \in B_\Omega$ has even level d, that is $v_2(d) = e_1$ (those have type 2).

Proof. We just apply proposition 5 to the basis of $Was(\mathbb{Q}(\zeta_n))$ given in theorem 3.

For generators having type 0, see our previous proof.

For generators having type 1, such an element x can be written as $1 - \zeta_d^a$ for some $d \mid n$ with d being odd and $a \wedge d = 1$. Moreover, equation (2) we have $1 - \zeta_d^a \in \mathbf{Z}(\mathbb{Q}(\zeta_n))\mathbf{E}^+(\mathbb{Q}(\zeta_n))$ as desired. For generators having type 2, such an element x can be written as $1-\zeta_d^a$ for some $d \mid n$ satisfying $v_2(d) = e_1$ and $a \wedge d = 1$. The same equation shows we have $1-\zeta_d^a \notin \mathbb{Z}(\mathbb{Q}(\zeta_n))\mathbb{E}^+(\mathbb{Q}(\zeta_n))$, otherwise we would have some primitive 2^{1+e_1} th root of unity lying in $\mathbb{Q}(\zeta_n)$, which is not the case.

4 Totally deployed fields

Through this section, we aim go give a basis of $Was(\mathbb{K})$ given \mathbb{K} is a totally deployed abelian number field (see our introduction). For now, we will suppose \mathbb{K} is a totally deployed abelian number field, with conductor n and we write

$$\mathbb{K} = \mathbb{K}_1 \cdots \mathbb{K}_r$$

where $\mathbb{K}_j \subset \mathbb{Q}(\zeta_{q_j})$.

To this end, we will consider a family of elements of \mathbb{K} that has $\operatorname{rg}_{\mathbb{Z}}(\operatorname{Was}(\mathbb{K})) = r_1 + r_2 - 1$ elements and that generates a direct factor of $\operatorname{Was}(\mathbb{Q}(\zeta_n))/\mathbb{Z}(\mathbb{Q}(\zeta_n))$. It is not hard to see that this property makes this family generate $\operatorname{Was}(\mathbb{K})/\mathbb{Z}(\mathbb{K})$ so that this family is a basis. More precisely, we will construct a basis of $\operatorname{Was}(\mathbb{K})$ that can be completed with Gold and Kim's basis to form a basis of $\operatorname{Was}(\mathbb{Q}(\zeta_n))$. This idea has already been used in [15], [5]. Actually, in order to prove proposition 2 from [15], the author proves the following fact.

Lemma 9. Let \mathbb{L} be an abelian number field with conductor n. If H is a group such that $\mathbf{Z}(\mathbb{L}) \subset H \subset \mathbf{Was}(\mathbb{L})$ is a direct factor of $\mathbf{Was}(\mathbb{Q}(\zeta_n))$ and H has the same \mathbb{Z} -rank as $\mathbf{Was}(\mathbb{L})$, then we have $H = \mathbf{Was}(\mathbb{L})$.

Proof. See the proof of proposition 2 from [15].

We should also mention that the fact one can state more results when \mathbb{K} is totally deployed appears in [12], [15] and [5].

We now constructs some sets and set notation to state our next theorem. To make it easier to understand, we tried to divide it into many definitions. The reader may not understand the following definition items as independent definitions but instead think of this separation as a help to read the following more easily.

We will refer to this notation as the notation of theorem 14.

Definition 10. Let d_j denote the degree of \mathbb{K}_j/\mathbb{Q} . Let

$$\widetilde{d}_j = \begin{cases} d_j & \text{if } \mathbb{K}_j \text{ is real} \\ \frac{d_j}{2} & \text{if not.} \end{cases}$$

With no loss of generality, suppose $\mathbb{K}_1, \ldots, \mathbb{K}_{t-1}$ are reals and $\mathbb{K}_t, \ldots, \mathbb{K}_r$ are not. For all $\Omega = \{i_1, \ldots, i_s\} \subset \llbracket 1, r \rrbracket$ that is non-empty, let

$$\mathbb{K}_{\Omega} = \mathbb{K}_{i_1} \cdots \mathbb{K}_{i_s}.$$

For all $\Omega = \{j\} \subset \llbracket 1, r \rrbracket$, let

$$B_{\Omega}(\mathbb{K}) = \left\{ \mathbb{N}_{\mathbb{Q}(\zeta_{q_j})^+ / \mathbb{K}_j^+}(\xi_{q_j, a}) : 1 \leqslant a < \widetilde{d}_j \right\}.$$

Definition 11. For all $X \subset \mathbb{N}$, let $\mathbb{1}_X : \mathbb{N} \to \mathbb{N}$ denote the indicator function of X.

For all $\Omega = \{i_1, \ldots, i_s\} \subset [\![1, r]\!]$ with $s \ge 2$, such that $i_1 < \cdots < i_s$ and \mathbb{K}_{i_s} is non-real (that is \mathbb{K}_{Ω} decomposes with at least 1 non-real field), let t_{Ω} be the integer such that $\mathbb{K}_{i_1}, \ldots, \mathbb{K}_{i_{t_{\Omega}-1}}$ are real and $\mathbb{K}_{i_{t_{\Omega}}}, \ldots, \mathbb{K}_{i_s}$ are non-real. For all $t_{\Omega} \le k < s$, let $X_{\Omega}^k(\mathbb{K})$ be the set of all tuples $(a_1, \ldots, a_s) \in \mathbb{N}^s$ such that

$$\begin{cases} \mathbbm{1}_{2\mathbb{Z}}(s-k)\frac{\varphi(q_{i_j})}{2} < a_j < d_{i_j} + \mathbbm{1}_{2\mathbb{Z}}(s-k)\frac{\varphi(q_{i_j})}{2} & \text{if } j \in [\![1,k[\![\\ \mathbbm{1}_{2\mathbb{Z}}(s-k)\frac{\varphi(q_{i_k})}{2} < a_k < \frac{d_{i_k}}{2} + \mathbbm{1}_{2\mathbb{Z}}(s-k)\frac{\varphi(q_{i_k})}{2} \\ a_j = \frac{d_{i_j}}{2} + \mathbbm{1}_{2\mathbb{Z}}(s-j)\frac{\varphi(q_{i_j})}{2} & \text{if } j \in [\!]k,s]\!]. \end{cases}$$

If k = s, let $X_{\Omega}^{k}(\mathbb{K})$ be the set of all tuples $(a_{1}, \ldots, a_{s}) \in \mathbb{N}^{s}$ such that

$$\left\{ \begin{array}{ll} 0 < a_j < d_{i_j} & \text{ if } j \in [\![1,s[\![\\ 0 < a_s < \frac{d_{i_s}}{2}. \end{array} \right. \end{array} \right.$$

If k = 0 (which will be useful if $t_{\Omega} = 1$), let

$$X_{\Omega}^{k}(\mathbb{K}) = \left\{ (a_{1}, \dots, a_{s}) \in \mathbb{N}^{s} : \forall j \in [\![1, s]\!], a_{j} = \frac{d_{i_{j}}}{2} + \mathbb{1}_{2\mathbb{Z}}(s - j)\frac{\varphi(q_{i_{j}})}{2} \right\}$$

and if $k = t_{\Omega} - 1$ (assuming $t_{\Omega} > 1$), let $X_{\Omega}^{k}(\mathbb{K})$ be the set of all tuples $(a_{1}, \ldots, a_{s}) \in \mathbb{N}^{s}$ such that

$$\begin{cases} \mathbb{1}_{2\mathbb{Z}}(s-k)\frac{\varphi(q_{i_j})}{2} < a_j < d_{i_j} + \mathbb{1}_{2\mathbb{Z}}(s-k)\frac{\varphi(q_{i_j})}{2} & \text{if } j \in [\![1, t_{\Omega}[\![\\a_j = \frac{d_{i_j}}{2} + \mathbb{1}_{2\mathbb{Z}}(s-j)\frac{\varphi(q_{i_j})}{2} & \text{if } j \in [\![t_{\Omega}, s]\!]. \end{cases}$$

Definition 12. Let $\Omega_{\mathbb{C}} = \Omega \cap \llbracket t, r \rrbracket$ and $\Omega_{\mathbb{R}} = \Omega \cap \llbracket 1, t - 1 \rrbracket$.

Under any of the previous assumptions on k, we let

$$B_{\Omega}^{k}(\mathbb{K}) = \left\{ \left(\prod_{j=1}^{s} \sigma_{i_{j}}^{a_{j}}\right) \mathbb{N}_{\mathbb{Q}(\zeta_{\Omega})/\mathbb{K}_{\Omega}}(1-\zeta_{\Omega}) : (a_{1},\ldots,a_{s}) \in X_{\Omega}^{k}(\mathbb{K}) \right\}$$

$$B_{\Omega}(\mathbb{K}) = \begin{cases} \bigcup_{\substack{k=0 \\ s}}^{s} B_{\Omega}^{k}(\mathbb{K}) & \text{if } \Omega_{\mathbb{R}} = \emptyset \text{ and } 2 \mid |\Omega| \\ \bigcup_{\substack{k=1 \\ s}}^{k} B_{\Omega}^{k}(\mathbb{K}) & \text{if } \Omega_{\mathbb{R}} = \emptyset \text{ and } 2 \nmid |\Omega| \\ \bigcup_{\substack{k=t_{\Omega}-1 \\ s}}^{k} B_{\Omega}^{k}(\mathbb{K}) & \text{if } \Omega_{\mathbb{R}} \neq \emptyset \text{ and } 2 \mid |\Omega_{\mathbb{C}}| \\ \bigcup_{\substack{k=t_{\Omega}}}^{s} B_{\Omega}^{k}(\mathbb{K}) & \text{if } \Omega_{\mathbb{R}} \neq \emptyset \text{ and } 2 \nmid |\Omega_{\mathbb{C}}|. \end{cases}$$

Definition 13. Recall $e_{\mathbb{L}}$ is defined in our introduction.

If \mathbb{K}_{i_s} is real (that is \mathbb{K}_{Ω} decomposed with real fields only), let

$$B_{\Omega}(\mathbb{K}) = \left\{ \left(\prod_{j=1}^{s} \sigma_{i_{j}}^{a_{j}} \right) (e_{\mathbb{K}_{\Omega}}) : 0 < a_{1} < d_{i_{1}}, \dots, 0 < a_{s} < d_{i_{s}} \right\}.$$

Theorem 14. The family $B(\mathbb{K}) = \bigcup_{\Omega} B_{\Omega}(\mathbb{K})$ where Ω runs over the set of all non-empty subsets of $[\![1, r]\!]$ is a basis of **Was**(\mathbb{K}).

Moreover, as a group, $\operatorname{Was}(\mathbb{K})/\mathbb{Z}(\mathbb{K})$ (which is naturally isomorphic to $\operatorname{Was}(\mathbb{K})\mathbb{Z}(\mathbb{Q}(\zeta_n))/\mathbb{Z}(\mathbb{Q}(\zeta_n)))$ is a direct factor of $\operatorname{Was}(\mathbb{Q}(\zeta_n))/\mathbb{Z}(\mathbb{Q}(\zeta_n))$. In what's next, to make it simple to read, we will talk about $\operatorname{Was}(\mathbb{Q}(\zeta_n))$ instead of $\operatorname{Was}(\mathbb{Q}(\zeta_n))/\mathbb{Z}(\mathbb{Q}(\zeta_n))$. More precisely, the family $B(\mathbb{K})$ can be completed to a basis of $\operatorname{Was}(\mathbb{Q}(\zeta_n))$ using Gold and Kim's basis.

Lemma 15. For all non-empty subset $\Omega \subset [\![1, r]\!]$, let

$$f_{\mathbb{C}}(\Omega) = \frac{1}{2} \left(\prod_{i \in \Omega} d_i - 1 \right) + \frac{(-1)^{|\Omega|}}{2}$$
$$f_{\mathbb{R}}(\Omega) = \left(\prod_{i \in \Omega} d_i - 1 \right)$$

$$g_{\mathbb{C}}(\Omega) = \frac{1}{2} \prod_{i \in \Omega} d_i$$
$$g_{\mathbb{R}}(\Omega) = \prod_{i \in \Omega} d_i.$$

If $\Omega = \emptyset$, say each of these functions maps Ω to 1. We have

$$\sum_{\substack{\Omega \subset \llbracket 1, r \rrbracket\\ \Omega \neq \emptyset}} f_{\mathbb{C}}(\Omega) = g_{\mathbb{C}}(\llbracket 1, r \rrbracket) - 1$$
(3)

$$\sum_{\substack{\Omega \subset \llbracket 1, r \rrbracket \\ \Omega \neq \emptyset}} f_{\mathbb{R}}(\Omega) = g_{\mathbb{R}}(\llbracket 1, r \rrbracket) - 1.$$
(4)

Proof. Let us prove the lemma first.

<u>Case 1</u> We may start with equation (4).

We have to prove

$$\mathbf{1} * f_{\mathbb{R}}([\![1,r]\!]) = g_{\mathbb{R}}([\![1,r]\!])$$

but instead, we will show we have,

$$\forall \Omega \subset \llbracket 1, r \rrbracket, \quad f_{\mathbb{R}}(\Omega) = \mu * g_{\mathbb{R}}(\Omega)$$

and the desired result will then be proven.

We have

$$\mu * g_{\mathbb{R}}(\Omega) = \sum_{X \subset \Omega} (-1)^{|\Omega| - |X|} g_{\mathbb{R}}(X)$$

=
$$\sum_{X \subset \Omega} (-1)^{|\Omega| - |X|} \prod_{i \in X} d_i$$

=
$$(-1)^{|\Omega|} + \sum_{k=1}^{|\Omega|} (-1)^{|\Omega| - k} \sum_{\substack{i_1, \dots, i_k \in \Omega \\ i_1 < \dots < i_k}} d_{i_1} \cdots d_{i_k}.$$

Using Vieta's formulas, we can see that this last expression matches the evaluation of the polynomial $(-1)^{|\Omega|} \prod_{i \in \Omega} X - d_i$ at X = 1, hence

$$\mu * g_{\mathbb{R}}(\Omega) = (-1)^{|\Omega|} \prod_{i \in \Omega} 1 - d_i = f_{\mathbb{R}}(\Omega).$$

<u>Cas 2</u> In a similar way, we now consider equation (3). We have

$$\mu * g_{\mathbb{C}}(\Omega) = \sum_{X \subset \Omega} (-1)^{|\Omega| - |X|} g_{\mathbb{C}}(X)$$

= $(-1)^{\Omega} + \sum_{\substack{X \subset \Omega \\ X \neq \emptyset}} (-1)^{|\Omega| - |X|} \frac{1}{2} \prod_{i \in X} d_i$
= $(-1)^{\Omega} + \frac{1}{2} \sum_{k=1}^{|\Omega|} (-1)^{|\Omega| - k} \sum_{\substack{i_1, \dots, i_k \in \Omega \\ i_1 < \dots < i_k}} d_{i_1} \cdots d_{i_k}.$

Using Vieta's formulas, we can see that this last term on the right side matches the evaluation of the polynomial

$$\frac{(-1)^{|\Omega|}}{2} \left(\left(\prod_{i \in \Omega} X - d_i \right) - X^{|\Omega|} \right)$$

at X = 1, hence

$$\mu * g_{\mathbb{C}}(\Omega) = (-1)^{\Omega} + \frac{(-1)^{|\Omega|}}{2} \left(\left(\prod_{i \in \Omega} 1 - d_i \right) - 1 \right) = f_{\mathbb{R}}(\Omega).$$

We may now prove the previously stated theorem.

Proof. First, we may prove $B(\mathbb{K})$ has cardinality $r_1 + r_2 - 1$, that is

$$|B(\mathbb{K})| = \begin{cases} \frac{1}{2} \left(\prod_{i \in \llbracket 1, r \rrbracket} d_i \right) - 1 & \text{if } |\llbracket 1, r \rrbracket_{\mathbb{C}}| \neq 0 \\ \left(\prod_{i \in \llbracket 1, r \rrbracket} d_i \right) - 1 & \text{if not.} \end{cases}$$

This can also be stated in the following way. For all non-empty subset $\Omega \subset [\![1, r]\!]$, we denote by $f(\Omega)$ the number of elements of $B_{\Omega}(\mathbb{K})$ and we let

$$g(\Omega) = \begin{cases} \frac{1}{2} \prod_{i \in \Omega} d_i & \text{if } |\Omega_{\mathbb{C}}| \neq 0\\ \prod_{i \in \Omega} d_i & \text{if not.} \end{cases}$$

 \Box

Also, say these functions both map $\Omega = \emptyset$ to 1. Then, we have to show

$$\mathbf{1} * f([\![1, r]\!]) = g([\![1, r]\!]).$$

Again, we rather show

$$\forall \Omega \subset \llbracket 1, r \rrbracket, \quad f(\Omega) = \mu * g(\Omega).$$

If $|\Omega| \leqslant 1$, there is nothing to be proven so we may suppose $|\Omega| \ge 1$. We separate three cases.

Suppose we have $\Omega_{\mathbb{C}} = \emptyset$.

Then, lemma 15 gives

$$\mu * g(\Omega) = \sum_{X \subset \Omega} (-1)^{|\Omega| - |X|} g(X)$$
$$= \sum_{X \subset \Omega} (-1)^{|\Omega| - |X|} \prod_{i \in X} d_i$$
$$= \prod_{i \in \Omega} (d_i - 1)$$

and it remains to observe

$$f(\Omega) = \prod_{i \in \Omega} (d_i - 1)$$

since we supposed $\Omega_{\mathbb{C}} = \emptyset$.

Now suppose $\Omega_{\mathbb{R}} = \emptyset$. Again, lemma 15 gives

$$\mu * g(\Omega) = f_{\mathbb{C}}(\Omega).$$

If $|\Omega|$ is odd, we have

$$f(\Omega) = \sum_{k=1}^{r} |B_{\Omega}^{k}(\mathbb{K})| = \sum_{k=1}^{r} (\frac{1}{2}d_{i_{k}} - 1)(d_{i_{k-1}} - 1) \cdots (d_{i_{1}} - 1)$$

and by induction on $N \in [\![1,r]\!],$ one can show

$$\sum_{k=1}^{N} (\frac{1}{2}d_{i_k} - 1)(d_{i_{k-1}} - 1) \cdots (d_{i_1} - 1) = \frac{1}{2}(d_{i_N} - 1) \cdots (d_{i_1} - 1) - \frac{1}{2}$$

Taking N = r, we get

$$\mu * g(\Omega) = f_{\mathbb{C}}(\Omega) = f(\Omega).$$

In the same way, if $|\Omega|$ is even, we have

$$f(\Omega) = 1 + \sum_{k=1}^{r} |B_{\Omega}^{k}(\mathbb{K})| = 1 + \sum_{k=1}^{r} (\frac{1}{2}d_{i_{k}} - 1)(d_{i_{k-1}} - 1) \cdots (d_{i_{1}} - 1)$$

and we get the same conclusion.

Now suppose we have $\Omega_{\mathbb{C}} \neq \emptyset$ and $\Omega_{\mathbb{R}} \neq \emptyset$. We have

$$\begin{split} \mu * g(\Omega) &= \sum_{X \subset \Omega} (-1)^{|\Omega| - |X|} g(X) \\ &= \sum_{\substack{X_1 \subset \Omega_{\mathbb{R}} \\ X_2 \subset \Omega_{\mathbb{C}} \\ X_2 \neq \emptyset}} (-1)^{|\Omega| - |X_1| - |X_2|} \frac{1}{2} \prod_{i \in X_1 \cup X_2} d_i + \sum_{X_1 \subset \Omega_{\mathbb{R}}} (-1)^{|\Omega| - |X_1|} \prod_{i \in X_1} d_i \\ &= \sum_{\substack{X_1 \subset \Omega_{\mathbb{R}} \\ X_2 \subset \Omega_{\mathbb{C}}}} (-1)^{|\Omega| - |X_1| - |X_2|} g_{\mathbb{C}}(X_1 \cup X_2) - \sum_{X_1 \subset \Omega_{\mathbb{R}}} (-1)^{|\Omega_{\mathbb{C}}| + |\Omega_{\mathbb{R}}| - |X_1|} g_{\mathbb{C}}(X_1) \\ &+ \sum_{X_1 \subset \Omega_{\mathbb{R}}} (-1)^{|\Omega_{\mathbb{C}}| + |\Omega_{\mathbb{R}}| - |X_1|} g_{\mathbb{R}}(X_1) \\ &= f_{\mathbb{C}}(\Omega) - (-1)^{|\Omega_{\mathbb{C}}|} f_{\mathbb{C}}(\Omega_{\mathbb{R}}) + (-1)^{|\Omega_{\mathbb{C}}|} f_{\mathbb{R}}(\Omega_{\mathbb{R}}) \\ &= \frac{1}{2} \prod_{i \in \Omega} (d_i - 1) + \frac{(-1)^{|\Omega_{\mathbb{C}}|}}{2} \prod_{i \in \Omega_{\mathbb{R}}} (d_i - 1). \end{split}$$

Separate cases depending on whether $|\Omega_{\mathbb{C}}|$ is even or not and one can show (using a similar induction argument as before) we have

$$f(\Omega) = \frac{1}{2} \prod_{i \in \Omega} (d_i - 1) + \frac{(-1)^{|\Omega_{\mathbb{C}}|}}{2} \prod_{i \in \Omega_{\mathbb{R}}} (d_i - 1).$$

This conclude the proof of the fact $B(\mathbb{K})$ has the expected cardinality.

We may now show the elements of $B(\mathbb{K})$ generate a direct factor of $Was(\mathbb{Q}(\zeta_n))$.

To make it easier to comprehend, we may assume n is odd and, at the end, we will explain what to do if n is even.

In order to show what we want, we will actually make every element x of $B_{\Omega}(\mathbb{K})$ correspond to a tuple $(a_1, \ldots, a_s) \in \bigcup_{k=0}^s X_{\Omega}^k$. Most of the time, this tuple will be the least tuple - ordering tuples with inverse lexicographic order (see our introduction) - such that

$$\left(\prod_{j=1}^{s} \sigma_{i_j}^{a_j}\right) \left(1 - \zeta_{\Omega}\right)$$

appears in the decomposition of x in Gold and Kim's basis of $\mathbf{Was}(\mathbb{Q}(\zeta_n))$. Moreover, we will show this element appears with exponent ± 1 and also these tuples are pairwise distinct. Later on, we will say we have made correspond the tuple (a_1, \ldots, a_s) to x. All this will conclude because, if we consider the matrix whose colums are the elements of $B(\mathbb{K})$ that are decomposed in Gold and Kim's basis, ordering all these elements correctly make this matrix be triangular with coefficients ± 1 over the diagonal.

More precisely, we may order the columns in the following way. Let $x \in B_{\Omega_1}(\mathbb{K})$ and $y \in B_{\Omega_2}(\mathbb{K})$ such that they correspond to (a_1, \ldots, a_{s_1}) and (b_1, \ldots, b_{s_2}) by the method we just described. The column that is associated to x is to the left of the one associated to y if we have

$$(\Omega_1, a_1, \ldots, a_{s_1}) \leqslant (\Omega_2, b_1, \ldots, b_{s_2})$$

according to what we have defined in our introduction. Note we then have to arbitrarily choose a way to order the subsets Ω 's according to this last binary relaton.

Then, we add columns to the right - in an arbitrary order - that corresponds to all the elements of Gold and Kim's basis that are not associated to the tuples $(\Omega, a_1, \ldots, a_s)$'s we have just talked about.

We may also order rows in the following way: starting from the top and ending at the bottom, we order in an ascending order, with the binary relation we just recalled - all the elements of Gold and Kim's basis that are associated to the tuples $(\Omega, a_1, \ldots, a_s)$'s we just mentionned. Then, we add - at the bottom - all the elements of Gold and Kim's basis that were left by this last ordering according to the same arbitrary order we have chosen when ordering the last columns.

We will use the same technics as Gold and Kim use in [2] to prove their theorem 1.

Again, to make it easier to comprehend, we will consider $x \in B_{[\![1,r]\!]}(\mathbb{K})$ but one can argue in the same way for any subset $\Omega \subset [\![1,r]\!]$. We will suppose *n* is not a prime power because this case has already been considered in the proof of proposition 2 from [15] and in the proof of theorem 2.1 from [5].

Finally, observe all groups $\operatorname{Gal}(\mathbb{Q}(\zeta_{q_i})/\mathbb{K}_i)$ are finite, cyclic and generated by $\sigma_i^{d_i}$.

We now separate cases, just like in the definition of $B_{[\![1,r]\!]}(\mathbb{K})$.

Suppose $\mathbb{K}_1, \ldots, \mathbb{K}_r$ are real fields. This case has already been considered in [15] (see proposition 2 and remark 4) but the author proved his result only in a special case (and stated it in the general case), we

may prove it now. In this case, we may show that, for any $0 < a_1 < d_1, \ldots, 0 < a_r < d_r$, we can make correspond (a_1, \ldots, a_s) to

$$x = \left(\prod_{j=1}^r \sigma_j^{a_j}\right)(e_{\mathbb{K}}).$$

First, observe we have $d_j | \varphi(q_j)/2$ since every \mathbb{K}_j is real. Modulo roots of unity of $\mathbb{Q}(\zeta_n)$, we have

$$x = \mathcal{N}_{\mathbb{L}_n/\mathbb{K}}(\eta_n^{\sigma_1^{a_1}\cdots\sigma_r^{a_r}})$$

=
$$\prod_{j_1=0}^{\varphi(q_1)/(2d_1)-1}\cdots\prod_{j_r=0}^{\varphi(q_r)/(2d_r)-1}\prod_{\varepsilon_1,\dots,\varepsilon_{r-1}\in\{0;1\}}1 - \zeta_n^{J_1^{\varepsilon_1}\sigma_1^{a_1+j_1d_1}\cdots J_{r-1}^{\varepsilon_{r-1}}\sigma_{r-1}^{a_{r-1}+j_{r-1}d_{r-1}}\sigma_r^{a_r+j_rd_r}}$$

and this is the decomposition of x in the basis of theorem 3. Indeed, we have $a_r + j_r d_r \in [0, \varphi(q_r)/2[$ and

$$J_i^{\varepsilon_i} \sigma_i^{a_i + j_i d_i} = \sigma_i^{\varepsilon_i \frac{\varphi(q_i)}{2} + a_i + j_i d_i}$$

$$\varepsilon_i \frac{\varphi(q_i)}{2} + a_i + j_i d_i \in]\!]0; \varphi(q_i)[\![$$

As we wanted, observe we can make correspond (a_1, \ldots, a_r) to x. Suppose $\mathbb{K}_1, \ldots, \mathbb{K}_{t-1}$ are real and $\mathbb{K}_t, \ldots, \mathbb{K}_r$ are not, for some $t \ge 2$ and $r \ge t$. Observe we have

$$\forall i \ge t, \quad \varphi(q_i)/2 = d_i/2 \mod d_i$$

since \mathbb{K}_i is not real, so that

$$\forall i \ge t, \quad J_i = \sigma_i^{d_i/2 + kd_i}$$

holds for $k = \varphi(q_i)/(2d_i) - 1/2$. Now, consider

$$x = \left(\prod_{j=1}^{r} \sigma_j^{a_j}\right) \mathcal{N}_{\mathbb{Q}(\zeta_n)/\mathbb{K}}(1-\zeta_n)$$

with $(a_1, \ldots, a_r) \in X^k_{[\![1,r]\!]}(\mathbb{K})$ for some $t \leq k \leq r$.

We may show by induction on $l \ge k$ we can make correspond the following tuple

$$(a_1 - \varepsilon_k \frac{\varphi(q_1)}{2}, \dots, a_k - \varepsilon_k \frac{\varphi(q_k)}{2}, 0, \dots, 0)$$

to the following element

$$x_{l} = \prod_{j_{1}=0}^{\varphi(q_{1})/d_{1}-1} \cdots \prod_{j_{l}=0}^{\varphi(q_{l})/d_{l}-1} 1 - \zeta_{n}^{J_{1}^{\varepsilon_{l}} \sigma_{1}^{a_{1}+j_{1}d_{1}} \cdots J_{l}^{\varepsilon_{l}} \sigma_{l}^{a_{l}+j_{l}d_{l}}}$$

with

$$\varepsilon_l = \mathbb{1}_{2\mathbb{Z}}(r-l).$$

<u>Base case with l = k</u>: Let

$$A_i = a_i - \varepsilon_k \frac{\varphi(q_i)}{2} \in]]0; d_i[]$$

Observe we have $A_k \in]0; d_k/2[$. Modulo roots of unity of $\mathbb{Q}(\zeta_n)$, we have

$$x_{k} = \prod_{j_{1}=0}^{\varphi(q_{1})/d_{1}-1} \cdots \prod_{j_{k-1}=0}^{\varphi(q_{k-1})/d_{k-1}-1} \prod_{j_{k} \leqslant \varphi(q_{k})/(2d_{k})-1/2} 1 - \zeta_{n}^{\sigma_{1}^{A_{1}+j_{1}d_{1}} \cdots \sigma_{k}^{A_{k}+j_{k}d_{k}}} \prod_{j_{1}=0}^{\varphi(q_{1})/d_{1}-1} \cdots \prod_{j_{k-1}=0}^{\varphi(q_{k-1})/d_{k-1}-1} \prod_{j_{k} \geqslant \varphi(q_{k})/(2d_{k})-1/2} 1 - \zeta_{n}^{\sigma_{1}^{A_{1}+j_{1}d_{1}} \cdots \sigma_{k}^{A_{k}+j_{k}d_{k}}}.$$

The first group of terms is already decomposed in the basis of theorem 3: all exponents $A_i + j_1 d_1$ lie in $[]0, \varphi(q_i)[$ whenever i < k and we have $A_k + j_k d_k \in]]0, \varphi(q_k)/2[$. We can see the tuple we are looking for appears with exponent 1.

The second group of terms can be treated just like the third one in the following induction step and then we can see the tuple we are looking for does not appear since this second group of terms decomposes with tuples from $\bigcup_{j>k} X_{[\![1,r]\!]}^j$ and tuples from $X_{[\![1,r]\!]}^k$ which have their kth coefficient congruent to $a_k + d_k/2$ in modulus d_k .

Induction step: Let l > k and suppose the result holds for previous cases. Let

$$A_i = a_i - \varepsilon_l \frac{\varphi(q_i)}{2}.$$

Observe we have

$$\forall i \leq k, \quad A_i \neq 0 \mod d_i$$
$$A_l = d_l/2.$$

Write

$$\begin{aligned} x_{l} &= \left(\prod_{j_{1}=0}^{\varphi(q_{1})/d_{1}-1} \cdots \prod_{j_{l-1}=0}^{\varphi(q_{l-1})/d_{l-1}-1} \prod_{j_{l}<\varphi(q_{l})/(2d_{l})-1/2} 1 - \zeta_{n}^{\sigma_{1}^{A_{1}+j_{1}d_{1}} \cdots \sigma_{l}^{A_{l}+j_{l}d_{l}}}\right) \\ & \left(\prod_{j_{1}=0}^{\varphi(q_{1})/d_{1}-1} \cdots \prod_{j_{l-1}=0}^{\varphi(q_{l-1})/d_{l-1}-1} \prod_{j_{l}=\varphi(q_{l})/(2d_{l})-1/2} 1 - \zeta_{n}^{\sigma_{1}^{A_{1}+j_{1}d_{1}} \cdots \sigma_{l}^{A_{l}+j_{l}d_{l}}}\right) \\ & \left(\prod_{j_{1}=0}^{\varphi(q_{1})/d_{1}-1} \cdots \prod_{j_{l-1}=0}^{\varphi(q_{l-1})/d_{l-1}-1} \prod_{j_{l}>\varphi(q_{l})/(2d_{l})-1/2} 1 - \zeta_{n}^{\sigma_{1}^{A_{1}+j_{1}d_{1}} \cdots \sigma_{l}^{A_{l}+j_{l}d_{l}}}\right). \end{aligned}$$

The first group of terms is almost decomposed in the basis of theorem 3: whenever i < l, we may reduce $A_i + j_i d_i$ modulo $\varphi(q_i)$. Then, zeros might appear and we are now going to explain how to decompose these terms in the basis of theorem 3. Consider

$$1-\zeta_n^{\sigma_1^{b_1}\dots\sigma_l^{b_l}}$$

with $b_l \in]0, \varphi(q_l)/2[$ and, for all i < l, suppose $b_i \in [0; \varphi(q_i)[$. Let i_1, \ldots, i_B be all the indexes i satisfying $b_i = 0$. Norm relations for $\mathbb{Q}(\zeta_{n/q_i})$ allow us to write

$$1-\zeta_n^{\sigma_1^{b_1}\dots\sigma_l^{b_l}}$$

with terms that have lower level and terms of the form

$$1 - \zeta_n^{\sigma_1^{c_1} \dots \sigma_l^{c_l}}$$

with $c_i = b_i$ for all $i \neq i_1$ and $c_{i_1} \in]0, \varphi(q_i)[$. By iterating this process, we see we can decompose

$$1 - \zeta_n^{\sigma_1^{b_1} \dots \sigma_l^{b_l}}$$

with terms that have lower level and terms of the form

$$1 - \zeta_n^{\sigma_1^{c_1} \dots \sigma_l^{c_l}}$$

with $c_l = b_l$ and $c_i \in]0, \varphi(q_i)[$ for all i < l.

Now, come back to x_l . From what preceds, we can say the tuple we are looking for does not appear in the first group of terms.

We may now consider the third group of terms. We may observe these terms satisfy $A_l + j_l d_l > \varphi(q_l)/2$. Using norm relations for $\mathbb{Q}(\zeta_{n/q_{l+1}})$, we can decompose these terms with terms that have lower level and terms of the form

$$1 - \zeta_n^{\sigma_1^{A_1 + j_1 d_1} \dots \sigma_l^{A_l + j_l d_l} \sigma_{l+1}^{B_{l+1}}}$$

with $B_{l+1} \in]0, \varphi(q_{l+1})[$ and with exponent ± 1 . The same considerations as those we used for the first group of terms show that if $B_{l+1} < \varphi(q_{l+1})/2$, then the following term

$$1 - \zeta_n^{\sigma_1^{A_1 + j_1 d_1} \dots \sigma_l^{A_l + j_l d_l} \sigma_{l+1}^{B_{l+1}}}$$

decomposes in the basis of theorem 3 with terms that have lower level and terms from $X_{[l,r]}^{l+1}$. Hence, these terms do not involve the tuple we are looking for. Then, we may now consider terms with $B_{l+1} \ge \varphi(q_{l+1})/2$. To this aim, we use norm relations for $\mathbb{Q}(\zeta_{n/q_{l+2}})$ and this leads to consider terms of the form

$$1 - \zeta_n^{\sigma_1^{A_1 + j_1 d_1} \dots \sigma_l^{A_l + j_l d_l} \sigma_{l+1}^{B_{l+1}} \sigma_{l+2}^{B_{l+2}}}$$

with $B_{l+2} \in]0, \varphi(q_{l+2})[$. Again, separate cases depending on whether we have $B_{l+2} < \varphi(q_{l+2})/2$. Now iterate this process and we are now lead to consider terms of the form

$$1 - \zeta_n^{\sigma_1^{A_1+j_1d_1}\dots\sigma_l^{A_l+j_ld_l}\sigma_{l+1}^{B_{l+1}}\dots\sigma_r^B}$$

with $\varphi(q_i)/2 \leq B_i < \varphi(q_i)$ for all $l < i \leq r$. Equation (1) allows us to transform these terms into terms of the form

$$1-\zeta_n^{\sigma_1^{C_1+j_1d_1}\ldots\sigma_l^{C_l+j_ld_l}\sigma_{l+1}^{D_{l+1}}\ldots\sigma_r^{D_r}}$$

with $0 \leq D_i < \varphi(q_i)/2$ for all i > l and $C_i = A_i - \varphi(q_i)/2$ for all $i \leq l$, so that

$$C_l + j_l d_l \in]\!]d_l/2, \varphi(q_l)/2[\![$$

Now, we just have to do the same manipulations as those for the first group of terms to see these last terms can be decomposed in the basis of theorem 3 with tuples from $\bigcup_{j \ge l} X_{[\![1,r]\!]}^j$ so that the third group of terms does not involve the tuple we are looking for.

We may now consider the second group of terms. This can be done through a similar process just like for the third group of terms. That is we are lead to consider terms of the form

$$1-\zeta_n^{\sigma_1^{C_1+j_1d_1}\cdots\sigma_l^{C_l+j_ld_l}\sigma_{l+1}^{D_{l+1}}\cdots\sigma_r^{D_r}}$$

with $0 \leq D_i < \varphi(q_i)/2$ for all i > l and $C_i = A_i - \varphi(q_i)/2$ for all $i \leq l$ but in this case we now have

$$C_l + j_l d_l = 0.$$

For the same reasons as before, terms such that there is i > l satisfying $D_i \neq 0$ can be decomposed in the basis of theorem 3 with tuples from $\bigcup_{j>l} X^j_{[1,r]}$ (so that the tuple we are looking for is not involved). On the other side, if we gather all terms such that $D_i = 0$ for all i > l, this leads us to consider the following product of terms instead of the second group of terms

$$\left(\prod_{j_1=0}^{\varphi(q_1)/d_1-1} \cdots \prod_{j_{l-1}=0}^{\varphi(q_{l-1})/d_{l-1}-1} 1-\zeta_n^{J_1\sigma_1^{A_1+j_1d_1}\cdots J_{l-1}\sigma_{l-1}^{A_{l-1}+j_{l-1}d_{l-1}}}\right)^{\pm 1}$$

and we can use induction hypothesis to put an end to this induction proof.

All this concludes if r - t + 1 is odd, that is we have made correspond any element $x \in B(\mathbb{K})$ to a tuple (a_1, \ldots, a_s) as wanted. Now, suppose r - t + 1 is even. The case k = t - 1 is the only case that has not been treated before. Let

$$x = \left(\prod_{j=1}^{r} \sigma_j^{a_j}\right) \mathcal{N}_{\mathbb{Q}(\zeta_n)/\mathbb{K}}(1-\zeta_n)$$

with $(a_1, \ldots, a_r) \in X^k_{[\![1,r]\!]}(\mathbb{K})$. The same proof shows we can make correspond $(a_1, \ldots, a_{t-1}, d_t/2, 0, \ldots, 0)$ to x (but in this case, this tuple is not the least among all the tuples that appears in the decomposition of x in the basis of theorem 3).

Indeed, we have

$$\begin{split} & \left(\prod_{j_{1}=0}^{\varphi(q_{1})/d_{1}-1} \cdots \prod_{j_{t-1}=0}^{\varphi(q_{t-1})/d_{t-1}-1} 1 - \zeta_{n}^{\sigma_{1}^{a_{1}+j_{1}d_{1}} \cdots \sigma_{t-1}^{a_{t-1}+j_{t-1}d_{t-1}}} \right) \\ & = \prod_{j_{1}=0}^{\varphi(q_{1})/d_{1}-1} \cdots \left(\prod_{j_{t-1}<\varphi(q_{t-1})/2d_{t-1}} 1 - \zeta_{n}^{\sigma_{1}^{a_{1}+j_{1}d_{1}} \cdots \sigma_{t-1}^{a_{t-1}+j_{t-1}d_{t-1}}} \right) \\ & \prod_{j_{1}=0}^{\varphi(q_{1})/d_{1}-1} \cdots \left(\prod_{j_{t-1}\geqslant\varphi(q_{t-1})/2d_{t-1}} 1 - \zeta_{n}^{\sigma_{1}^{a_{1}+j_{1}d_{1}} \cdots \sigma_{t-1}^{a_{t-1}+j_{t-1}d_{t-1}}} \right) \end{split}$$

and we can see the tuple we are looking for is not involed in the first group of terms as this is already decomposed in the basis of theorem 3. To show this is also the case for the second one, we may follow the same steps as in the previous induction proof and pay more attention. Using norm relations for $\mathbb{Q}(\zeta_{n/q_t})$, we are lead to consider the following terms only (what we have done during the previous induction proof allows us not to consider the other terms that appear when using norm relations straight forward because they do not involve the tuple we are looking for)

$$\left(1 - \zeta_n^{\sigma_1^{a_1+j_1d_1}\cdots\sigma_{t-1}^{i_{t-1}+j_{t-1}d_{t-1}}\sigma_t^{d_t/2}}\right)^{-1} \\ \left(1 - \zeta_n^{\sigma_1^{a_1+j_1d_1}\cdots\sigma_{t-1}^{i_{t-1}+j_{t-1}d_{t-1}}J_t}\right)^{-1}.$$

This first term make the tuple we are looking for appear when $j_1 = \cdots = j_{t-1} = 0$ only and it appears with exponant -1. The second term can be treated using norm relations again. We are then lead to consider the following terms (with exponant +1)

$$\left(1-\zeta_n^{\sigma_1^{a_1+j_1d_1}\dots\sigma_{t-1}^{i_{t-1}+j_{t-1}d_{t-1}}J_tJ_{t+1}}\right)^{+1}$$

By iterating this process, we are then lead to consider the following terms

$$\left(1 - \zeta_n^{\sigma_1^{a_1+j_1d_1}\dots\sigma_{t-1}^{i_{t-1}+j_{t-1}d_{t-1}}J_t\dots J_r}\right)^{+1}$$

with exponant +1 because r - t + 1 is even. Equation (1) transforms these terms to the following ones

$$\left(1-\zeta_n^{J_1\sigma_1^{a_1+j_1d_1}\cdots J_{t-1}\sigma_{t-1}^{i_{t-1}+j_{t-1}d_{t-1}}\sigma_t^{d_t/2}}\right)^{+1}.$$

These terms involve the tuple we are looking for when $j_1 = \varphi(q_1)/(2d_1), \ldots, j_{t-1} = \varphi(q_{t-1})/(2d_{t-1})$ and it appears with exponant +1.

At the end, the tuple we are looking for is not involed in the decomposition of the following element

$$\prod_{j_1=0}^{\varphi(q_1)/d_1-1} \cdots \prod_{j_{t-1}=0}^{\varphi(q_{t-1})/d_{t-1}-1} 1 - \zeta_n^{\sigma_1^{a_1+j_1d_1} \dots \sigma_{t-1}^{a_{t-1}+j_{t-1}d_{t-1}}}$$

Then, we can show the tuple we are looking for appears with exponant +1 in the decomposition of the following term

$$\prod_{j_1=0}^{\varphi(q_1)/d_1-1} \cdots \prod_{j_t=0}^{\varphi(q_t)/d_t-1} 1 - \zeta_n^{\sigma_1^{a_1+j_1d_1}\cdots\sigma_t^{d_t/2+j_td_t}}$$

The same induction proof shows we can make correspond $(a_1, \ldots, a_{t-1}, d_t/2, 0, \ldots, 0)$ to x. We may observe this tuple is not involved in the decomposition of any other element of $B_{[\![1,r]\!]}(\mathbb{K})$. Suppose $\mathbb{K}_1, \ldots, \mathbb{K}_r$ are not real.

What we have done just before allows us to make correspond the same tuple

$$(a_1 - \varepsilon_k \frac{\varphi(q_1)}{2}, \dots, a_k - \varepsilon_k \frac{\varphi(q_k)}{2}, 0, \dots, 0)$$

to

$$x = \left(\prod_{j=1}^{r} \sigma_j^{a_j}\right) \mathcal{N}_{\mathbb{Q}(\zeta_n)/\mathbb{K}}(1-\zeta_n)$$

for any $(a_1, \ldots, a_r) \in X^k_{[\![1,r]\!]}(\mathbb{K})$ and $0 < k \leq r$ and we can also go through this same proof if k = 0 and r is even.

All this concludes the proof of the theorem when n is odd.

Now, suppose n is even and let i_0 be such that $p_{i_0} = 2$. The little difference with the case when n is odd comes from the fact that the Galois group of $\mathbb{Q}(\zeta_n)/\mathbb{K}$ and the norm operator are no longer caracterised by the d_i 's only. More precisely, the Galois group of $\mathbb{Q}(\zeta_{q_{i_0}})/\mathbb{K}_{i_0}$ is no longer caracterised by d_{i_0} . Moreover, we no longer have $d_{i_0} \mid \varphi(q_{i_0})/2$ if and only if \mathbb{K}_{i_0} is real. Recall σ_{i_0} has been constructed with J_{i_0} and $\tilde{\sigma}_{i_0}$. We may now explain how to adapt our previous proof. When all the \mathbb{K}_j 's are real, we can argue just like before and observe the Galois group of $\mathbb{Q}(\zeta_{q_{i_0}})^+/\mathbb{K}_{i_0}$ is generated by $\tilde{\sigma}_{i_0}^{d_{i_0}}$. Then, we may write

$$\mathcal{N}_{\mathbb{L}_n/\mathbb{K}} = \prod_{j_1=0}^{\varphi(q_1)/d_1-1} \cdots \prod_{j_r=0}^{\varphi(q_r)/d_r-1} \sigma_1^{j_1d_1} \cdots \widetilde{\sigma}_{i_0}^{j_{i_0}d_{i_0}} \cdots \sigma_r^{j_rd_r}$$

and write the same proof as before.

If \mathbb{K}_{i_0} is real, we may write things differently. Observe J_{i_0} does not lie in $\operatorname{Gal}(\mathbb{Q}(\zeta_{q_{i_0}})/\mathbb{K}_{i_0})$. We have

$$\mathcal{N}_{\mathbb{Q}(\zeta_n)/\mathbb{K}} = \prod_{j_1=0}^{\varphi(q_1)/d_1-1} \cdots \prod_{g \in \mathrm{Gal}(\mathbb{Q}(\zeta_{q_{i_0}})/\mathbb{K}_{i_0})} \cdots \prod_{j_r=0}^{\varphi(q_r)/d_r-1} \sigma_1^{j_1d_1} \cdots g \cdots \sigma_r^{j_rd_r}$$

and having this in mind, we can use the same proof again after writting the elements g as "powers" of σ_{i_0} . In particular, we may take caution with the following case that is when

$$x = \left(\prod_{j=1}^{r} \sigma_j^{a_j}\right) \mathcal{N}_{\mathbb{Q}(\zeta_n)/\mathbb{K}}(1-\zeta_n)$$

with $(a_1, \ldots, a_r) \in X^{i_0}_{[\![1,r]\!]}$. Since J_{i_0} does not appear in $N_{\mathbb{Q}(\zeta_n)/\mathbb{K}}$, we are sure that we can make correspond the same tuple as before to such an element x. More precisely, the manipulations we have operated before show that, in order for (a_1, \ldots, a_r) to appear, we have two choices : either it appears directly or (J_1a_1, \ldots, J_ra_r) appears but this second case cannot occure as J_{i_0} does not appear in $N_{\mathbb{Q}(\zeta_n)/\mathbb{K}}$.

If \mathbb{K}_{i_0} is real but there is some j such that \mathbb{K}_j is not real, then we may write $N_{\mathbb{Q}(\zeta_n)/\mathbb{K}}$ as we just did and make the same proof as in the case of an odd integer n by writting the elements g as powers of σ_{i_0} .

Remark 16. Equation (3) from lemma 15 shows Gold and Kim's basis has the cardinality it should have to be a basis (this has been stated with no proof in [2]).

Corollary 17. Under the hypotheses and notation of theorem 14, suppose \mathbb{K}_r is not real and $\mathbb{K}_1, \ldots, \mathbb{K}_{r-1}$ are real, then

$$\mathbf{Was}(\mathbb{K}) = \mathbf{Z}(\mathbb{K}) \, \mathbf{Was}^+(\mathbb{K}).$$

Proof. This results of the fact we have

$$\mathbf{Z}(\mathbb{K}) \operatorname{Was}^+(\mathbb{K}) \subset \operatorname{Was}(\mathbb{K})$$

and $\operatorname{Was}^+(\mathbb{K})$ is a direct factor of $\operatorname{Was}(\mathbb{Q}(\zeta_n))$ as

$$\mathbf{Was}^+(\mathbb{K}) = \mathbf{Was}(\mathbb{K}^+) = \mathbf{Was}(\mathbb{K}_1\mathbb{K}_2\dots\mathbb{K}_r^+)$$

(see lemma 9).

Corollary 18. Under the hypotheses and notation of theorem 14, the quotient group $\operatorname{Was}(\mathbb{K})/\operatorname{Sin}(\mathbb{K})$ is isomorphic to $(\mathbb{Z}/2\mathbb{Z})^a$ for some $a \in [0; [\mathbb{K}_{[1,r]_{\mathbb{R}}} : \mathbb{Q}][$.

Proof. Indeed, all generators that are (mentionned in the previous theorem and) associated to some $\Omega \subset [\![1, r]\!]$ such that $|\Omega_{\mathbb{C}}| \ge 1$ are already elements of $\mathbf{Sin}(\mathbb{K}_{\Omega})$ and all other generators that make use of $e_{\mathbb{K}_{\Omega}}$ have order 2 in the quotient group $\mathbf{Was}(\mathbb{K}_{\Omega})/\mathbf{Sin}(\mathbb{K}_{\Omega})$ (see [15], equation 11 and corollary 3).

Hence, the quotient group $Was(\mathbb{K})/Sin(\mathbb{K})$ is an elementary 2-group and we have bounded its rank with the following number

$$\sum_{\substack{\Omega \subset [\![1,t]\![] \\ \Omega \neq \emptyset}} \prod_{i \in \Omega} (d_i - 1) = (d_1 \cdots d_{t-1} - 1)$$

by lemma 15.

Remark 19. Werl Milàn stated and proved in a special case (see [15] remark 4) we have $a = [\mathbb{K} : \mathbb{Q}] - 1$ if $\mathbb{K}_1, \ldots, \mathbb{K}_r$ are real.

We may also observe if $\mathbb{K}_1, \ldots, \mathbb{K}_r$ are not real, we have $\mathbf{Was}(\mathbb{K}) = \mathbf{Sin}(\mathbb{K})$ so that the previous theorem gives a basis of $\mathbf{Sin}(\mathbb{K})$. A different basis of $\mathbf{Sin}(\mathbb{K})$ has been given in [8] but both bases are quite similar.

Corollary 20. Under the hypotheses and notation of theorem 14, we have

$$[\mathbf{E}(\mathbb{K}):\mathbf{Was}(\mathbb{K})]=h^+(\mathbb{K})2^x$$

for some $x \in \mathbb{N}$.

Proof. This results of the previous corollary and the formula Sinnott has given for the index of $Sin(\mathbb{K})$ in $E(\mathbb{K})$ (see [12] theorem 4.1, proposition 5.1 and theorem 5.4).

Corollary 21. Under the hypotheses and notation of theorem 14, let A_1, \ldots, A_k be disjoint subsets of $[\![1, r]\!]$. We have a canonical injective map

$$\prod_{j=1}^{k} \operatorname{E}(\mathbb{K}_{A_{j}}) / \operatorname{Was}(\mathbb{K}_{A_{j}}) \hookrightarrow \operatorname{E}(\mathbb{K}) / \operatorname{Was}(\mathbb{K})$$

)

In particular, if we let $h_p^+(\mathbb{K})$ denote the *p*-part of the class number of \mathbb{K}^+ , we have for all odd prime *p*

$$\prod_{j=1}^{k} h_p^+(\mathbb{K}_{A_j}) \mid h_p^+(\mathbb{K}).$$

Proof. Let $x = x_1 \cdots x_k \in \mathbf{Was}(\mathbb{K})$ with $x_j \in \mathbf{E}(\mathbb{K}_{A_j})$. We have to show $x_j \in \mathbf{Was}(\mathbb{K}_{A_j})$. There is an integer N such that we have $x_j^N \in \mathbf{Was}(\mathbb{K}_{A_j})$. Modulo roots of unity of \mathbb{K} , we have

$$x = \prod_{u \in B(\mathbb{K})} u^{x_u}$$
$$x_j^N = \prod_{u \in B(\mathbb{K}_{A_j})} u^{x_{j,u}}$$

that is the decomposition of x and the x_j^N 's in the basis we gave in the previous theorem. This theorem shows the following group is a direct factor of $\mathbf{Was}(\mathbb{K})$

$$\prod_{j=1}^k \operatorname{Was}(\mathbb{K}_{A_j}).$$

Now, we may identify the exponents of x^N so that we get

 $\forall j \in \llbracket 1, k \rrbracket, \ \forall u \in B(\mathbb{K}_{A_j}), \quad Nx_u = x_{j,u}$

then we have

$$x_j^N = \left(\prod_{u \in B(\mathbb{K}_{A_j})} u^{x_u}\right)^N$$

hence $x_j \in \mathbf{Was}(\mathbb{K}_{A_j})$.

It turns out we can prove this last result on class numbers through class field theory. We have found no reference related to the proof of this next proposition so far and that is why we give a proof.

Proposition 22. Suppose

$$\mathbb{K} = \mathbb{K}_1 \cdots \mathbb{K}_r$$

with $\mathbb{K}_i \subset \mathbb{Q}(\zeta_{q_i})$. Let A_1, \ldots, A_k be a partition of $[\![1; r]\!]$. We have

$$\prod_{j=1}^{k} h(\mathbb{K}_{A_j}) \mid h(\mathbb{K}).$$

Proof. Class field theory gives the following commutative diagram (see what is before theorem 5 in the appendix of [14])

where **H** denotes the Hilbert class field, $\mathcal{I}_{\mathbb{L}}$ denotes the group of all fractionnal ideals of \mathbb{L} , $\mathcal{P}_{\mathbb{L}}$ denotes the group of all principal ideals of \mathbb{L} , the map res is given by the restriction maps that arise with $\mathbf{H}_{\mathbb{K}_i} \subset \mathbf{H}_{\mathbb{K}}$ and the map Norm is given by the norm maps

$$\mathcal{I}_{\mathbb{K}}/\mathcal{P}_{\mathbb{K}} \to \mathcal{I}_{\mathbb{K}_i}/\mathcal{P}_{\mathbb{K}_i}$$

We may prove the map Norm is surjective to conclude. To this aim, we will show res is surjective.

First, observe the $\mathbf{H}_{\mathbb{K}_i}$'s form a free compositum, that is we have

$$\mathbf{H}_{\mathbb{K}_i} \cap \left(\prod_{j
eq i} \mathbf{H}_{\mathbb{K}_j}
ight) = \mathbb{Q}.$$

Indeed, each prime number ramifies in $\mathbf{H}_{\mathbb{K}_i}$ if and only if it ramifies in \mathbb{K}_i and \mathbb{Q} has no unramified extension (see [9] theorem 2.18).

Let x_1, x_2 be such that

$$\mathbf{H}_{\mathbb{K}_{A_1}} = \mathbb{Q}(x_1)$$
$$\prod_{i>1} \mathbf{H}_{\mathbb{K}_{A_i}} = \mathbb{Q}(x_2).$$

Let N be the degree of x_2 over \mathbb{Q} and observe x_2 has degree N over $\mathbf{H}_{\mathbb{K}_{A_1}}$. Indeed, its minimal polynomial over $\mathbf{H}_{\mathbb{K}_{A_1}}$ has coefficients in $\mathbf{H}_{\mathbb{K}_{A_1}}$ and it also has coefficient in $\mathbb{Q}(x_2)$ because the $\mathbf{H}_{\mathbb{K}_i}$'s are Galois since the \mathbb{K}_i 's themselves are Galois. Therefore, what is before proves the minimal polynomial of x_2 over $\mathbf{H}_{\mathbb{K}_{A_1}}$ is the same over \mathbb{Q} . As a result, Galois theory gives a surjective map (by restriction)

$$\operatorname{Gal}\left(\prod_{i\geq 1}\mathbf{H}_{\mathbb{K}_{i}}/\mathbb{K}\right)\twoheadrightarrow\operatorname{Gal}\left(\mathbf{H}_{\mathbb{K}_{A_{1}}}/\mathbb{K}_{A_{1}}\right)\times\operatorname{Gal}\left(\left(\prod_{i>1}\mathbf{H}_{\mathbb{K}_{A_{i}}}\right)/\left(\prod_{i>1}\mathbb{K}_{A_{i}}\right)\right)$$

and by induction

$$\operatorname{Gal}\left(\prod_{i\geq 1}\mathbf{H}_{\mathbb{K}_{i}}/\mathbb{K}\right)\twoheadrightarrow\prod_{i\geq 1}\operatorname{Gal}(\mathbf{H}_{\mathbb{K}_{A_{i}}}/\mathbb{K}_{A_{i}}).$$

Galois theory also shows the following map is surjective

$$\operatorname{Gal}(\mathbf{H}_{\mathbb{K}}/\mathbb{K}) \twoheadrightarrow \operatorname{Gal}\left(\prod_{i \ge 1} \mathbf{H}_{\mathbb{K}_i}/\mathbb{K}\right).$$

Remark 23. We can adapt this last proof to get a similar result with h^+ instead of h.

We may use proposition 5 and theorem 14 to obtain a basis of $\mathbf{Was}(\mathbb{K}_1 \cdots \mathbb{K}_{t-1}(\mathbb{K}_t \cdots \mathbb{K}_r)^+)$.

Next lemma 24 allows us to state corollary 20 with \mathbb{K}^+ replacing \mathbb{K} .

We can also observe $\mathbf{Was}^+(\mathbb{Q}(\zeta_n))$ is not a direct factor of $\mathbf{Was}(\mathbb{Q}(\zeta_n))$ because, if it was a direct factor, we would get $\mathbf{Was}(\mathbb{Q}(\zeta_n)) = \mathbf{Z}(\mathbb{Q}(\zeta_n)) \mathbf{Was}^+(\mathbb{Q}(\zeta_n))$ by lemma 9.

Lemma 24. Let \mathbb{K} be an abelian number field. We have

$$[\mathbf{E}^+(\mathbb{K}):\mathbf{Was}^+(\mathbb{K})] = [\mathbf{E}(\mathbb{K}):\mathbf{Was}(\mathbb{K})] \frac{[\mathbf{Was}(\mathbb{K})\mathbf{E}^+(\mathbb{K}):\mathbf{Z}(\mathbb{K})\mathbf{E}^+(\mathbb{K})]}{[\mathbf{E}(\mathbb{K}):\mathbf{Z}(\mathbb{K})\mathbf{E}^+(\mathbb{K})]}$$

Proof. As the index is multiplicative, we get

$$\begin{split} [\mathbf{E}(\mathbb{K}):\mathbf{Was}^+(\mathbb{K})] &= [\mathbf{E}(\mathbb{K}):\mathbf{Was}(\mathbb{K})][\mathbf{Was}(\mathbb{K}):\mathbf{Z}(\mathbb{K})\,\mathbf{Was}^+(\mathbb{K})][\mathbf{Z}(\mathbb{K})\,\mathbf{Was}^+(\mathbb{K}):\mathbf{Was}^+(\mathbb{K})] \\ &= [\mathbf{E}(\mathbb{K}):\mathbf{Z}(\mathbb{K})\mathbf{E}^+(\mathbb{K})][\mathbf{Z}(\mathbb{K})\mathbf{E}^+(\mathbb{K}):\mathbf{E}^+(\mathbb{K})][\mathbf{E}^+(\mathbb{K}):\mathbf{Was}^+(\mathbb{K})]. \end{split}$$

It remains to see the second isomorphism theorem gives

$$[\mathbf{Z}(\mathbb{K})\mathbf{E}^+(\mathbb{K}):\mathbf{E}^+(\mathbb{K})] = \frac{|\mathbf{Z}(\mathbb{K})|}{2} = [\mathbf{Z}(\mathbb{K})\mathbf{Was}^+(\mathbb{K}):\mathbf{Was}^+(\mathbb{K})]$$

and

$$\operatorname{Was}(\mathbb{K})\mathrm{E}^+(\mathbb{K})/\mathbf{Z}(\mathbb{K})\mathrm{E}^+(\mathbb{K}) \simeq \operatorname{Was}(\mathbb{K})/\mathbf{Z}(\mathbb{K})\operatorname{Was}^+(\mathbb{K}).$$

Proposition 25. Suppose \mathbb{K} ramifies at p. The universal norms Galois module $\mathbf{Was}_{\infty}^{0}$ is generated - as a Galois module - by the x's such that $x \in B_{\Omega}(\mathbb{K})$ with $1 \in \Omega$.

References

- John Coates and R. Sujatha. Cyclotomic fields and zeta values. Springer Monogr. Math. Berlin: Springer, 2006. doi:10.1007/978-3-540-33069-1.
- [2] Robert Gold and Jaemoon Kim. Bases for cyclotomic units. Compos. Math., 71(1):13–27, 1989.
- [3] Georges Gras. Class field theory. From theory to practice. Springer Monogr. Math. Berlin: Springer, 2003.
- [4] Curtis Greene. The Moebius function of a partially ordered set. Ordered sets, Proc. NATO Adv. Study Inst., Banff/Can. 1981, 555-581 (1982)., 1982.
- Jae Moon Kim and Jado Ryu. Construction of a certain circular unit and its applications. J. Number Theory, 131(4):737-744, 2011. doi:10.1016/j.jnt.2010.11.002.
- [6] Radan Kučera. A note on Sinnott's definition of circular units of an abelian field. J. Number Theory, 63(2):403-407, 1997. doi:10.1006/jnth.1997.2094.
- [7] Radan Kučera. Circular units and class groups of abelian fields. Ann. Sci. Math. Québec, 28(1-2):121-136, 2004.
- [8] Radan Kučera. The circular units and the Stickelberger ideal of a cyclotomic field revisited. Acta Arith., 174(3):217–238, 2016. doi:10.4064/aa8009-4-2016.
- [9] Jürgen Neukirch. Algebraic number theory. Transl. from the German by Norbert Schappacher, volume 322 of Grundlehren Math. Wiss. Berlin: Springer, 1999.
- [10] Gian-Carlo Rota. On the foundations of combinatorial theory. I: Theory of Möbius functions. Z. Wahrscheinlichkeitstheor. Verw. Geb., 2:340–368, 1964. doi:10.1007/BF00531932.
- [11] W. Sinnott. On the Stickelberger ideal and the circular units of a cyclotomic field. Ann. of Math. (2), 108(1):107-134, 1978. doi:10.2307/1970932.
- [12] W. Sinnott. On the stickelberger ideal and the circular units of an abelian field. Inventiones mathematicae, 62:181-234, 1980/81. URL: http://eudml.org/doc/142770.
- [13] David Solomon. Galois relations for cyclotomic numbers and p-units. J. Number Theory, 46(2):158–178, 1994. doi:10.1006/jnth.1994.1010.
- [14] Lawrence C. Washington. Introduction to cyclotomic fields., volume 83 of Grad. Texts Math. New York, NY: Springer, 2nd ed. edition, 1997.
- [15] Milan Werl. On bases of Washington's group of circular units of some real cyclic number fields. J. Number Theory, 134:109–129, 2014. doi:10.1016/j.jnt.2013.07.016.