# GROUP SCHEMES OVER LG-RINGS AND APPLICATIONS TO CANCELLATION THEOREMS AND AZUMAYA ALGEBRAS

PHILIPPE GILLE AND ERHARD NEHER

ABSTRACT. We prove several results on reductive group schemes over LG-rings, e.g., existence of maximal tori and conjugacy of parabolic subgroups. These were proven in [\[SGA3\]](#page-63-0) for the special case of semilocal rings. We apply these results to establish cancellation theorems for hermitian and quadratic forms over LG-rings and show that the Brauer classes of Azumaya algebras over connected LG-rings have a unique representative and allow Brauer decomposition.

## **CONTENTS**



2010 Mathematics Subject Classification. 20G35; 11E39, 13A20, 13H99, 14L15.

Date: July 3, 2024.

Key words and phrases. LG-rings, reductive group schemes, parabolic subgroup schemes, maximal tori, cancellation theorems, Azumaya algebras, Wedderburn property, Brauer decomposition.

The first author was supported by the project "Group schemes, root systems, and related representations" founded by the European Union - NextGenerationEU through Romania's National Recovery and Resilience Plan (PNRR) call no. PNRR-III-C9-2023- I8, Project CF159/31.07.2023, and coordinated by the Ministry of Research, Innovation and Digitalization (MCID) of Romania.

#### 2 P. GILLE AND E. NEHER

### <span id="page-1-0"></span>**INTRODUCTION**

In volume III of the fundamental treatise [\[SGA3\]](#page-63-0) on group schemes over a scheme S, several results are only proven if  $S = \operatorname{Spec}(R)$  for R a semilocal ring. In this paper we extend most of these results to the case  $S = \text{Spec}(R)$ for  $R$  an LG-ring.

LG-rings, an abbreviation of "local-global" rings, axiomatize a basic property of a semilocal ring R. By definition, a ring R is an LG-ring if a polynomial f in several variables represents a unit of  $R$  as soon as for every maximal ideal  $\mathfrak{m} \triangleleft R$  the canonical image  $f_{R/\mathfrak{m}}$  represents a unit over  $R/\mathfrak{m}$ . Besides semilocal rings, the class of LG-rings include rings that are von Neumann regular modulo their radical and the ring of algebraic integers, see [1.2](#page-4-0) for more examples and an historical background.

Among the results we prove for a reductive group scheme  $G$  over an  $LG$ ring  $R$ , we highlight the following:

- <span id="page-1-1"></span>(1) Every parabolic subgroup of  $G$  admits a maximal  $R$ –torus, in particular this holds for G itself  $(2.6, 2.8)$  $(2.6, 2.8)$ .
- <span id="page-1-5"></span>(2) Two parabolic subgroups of the same type are conjugate under an element of  $G(R)$  [\(3.1\)](#page-12-1).
- <span id="page-1-2"></span>(3) If R is a connected LG-ring, then  $G(R)$  acts transitively on the set of minimal parabolic subgroups of  $G$ , and if  $G$  is semisimple, then  $G(R)$  also acts transitively on the set of maximal split subtori of G  $(4.7).$  $(4.7).$

An important ingredient in the proofs of  $(1)$ – $(3)$  is a geometric characterization of LG-rings:

<span id="page-1-3"></span>(4) If M is a finite locally free module over an LG-ring R and U is an open quasi-compact subscheme of the scheme  $\mathbf{W}(M)$  associated with M, then  $U(R) \neq \emptyset \iff U(R/\mathfrak{m}) \neq \emptyset$  for every maximal ideal  $\mathfrak{m} \triangleleft R$ , see [1.4](#page-6-0)[\(a\).](#page-6-1)

With  $(4)$  in place, our proof strategy often is to recast a claim in terms of an open quasi-compact subscheme of an affine space and then follow the approach of [\[SGA3\]](#page-63-0).

Our applications are twofold. First, in §[5](#page-26-0) we prove several cancellation theorems, based on the cohomological cancellation principle [5.1](#page-26-1) which says that certain embeddings of reductive group schemes over LG-rings induce injective maps in cohomology. Specializing the groups involved, we easily derive cancellation of modules and Azumaya algebras in tensor products [5.2,](#page-26-2) cancellation of hermitian forms in [5.4](#page-29-0) and Witt cancellation of quadratic forms in [5.5.](#page-29-1)

<span id="page-1-4"></span>Second, we consider Azumaya  $R$ –algebras  $A$  in §[6](#page-31-0) and extend several results from R connected semilocal to R being connected LG. In particular, we show:

(5) Each two indecomposable finite projective A–modules are isomorphic. Equivalently, the Brauer class of A contains, up to isomorphism, a unique algebra B with idemp $(B) = \{0, 1\}$ , where idemp $(B)$ is the set of idempotents of  $B(6.7)$  $B(6.7)$ .

(6) Every A with idemp $(A) = \{0, 1\}$  has a Brauer decomposition [\(6.9\)](#page-35-0). We note that  $(5)$  says that connected LG-rings have the Wedderburn prop-

erty in the sense of  $[AW17]$  (or see  $[Fe, §7.6]$ ).

Organization of the paper. We begin with a short introduction to LGrings in §[1,](#page-3-0) list examples, recall some immediate consequences and prove the important characterization [\(4\)](#page-1-3) of LG-rings, stated above. Our investigation of group schemes over LG-rings starts in  $\S$ [2,](#page-7-0) where we prove the crucial result [\(1\)](#page-1-1). We finish this section by discussing quasi-split and split reductive groups over LG-rings.

We study parabolic subgroups of reductive group schemes over LG-rings in §[3.](#page-12-0) Besides [\(2\)](#page-1-5) above and some immediate corollaries on three or two parabolic subgroups  $(3.2, 3.6)$  $(3.2, 3.6)$ , we prove in  $3.8(b)$  $3.8(b)$  that there exists a unique smallest element  $t_{\min}$  in the set of types of parabolic subgroups of a reductive  $R$ –group scheme G for R an LG-ring. This allows us to introduce the Tits index in case  $G$  is semisimple and  $R$  is connected (no new indices occur, [3.9\)](#page-19-3).

We focus on minimal parabolic subgroups and maximal split tori in  $\S4$ , in particular prove  $(3)$  above. As a consequence, it makes sense to define the anisotropic kernel of a reductive  $R$ –group scheme  $G, R$  connected LG, as the derived group of a minimal Levi subgroup, [\(4.9.1\)](#page-24-0).

Finally, §[5](#page-26-0) and §[6](#page-31-0) are devoted to the applications mentioned above.

Notation. We use standard notation and terminology, but note here that  $R$ -alg denotes the category of unital associative commutative  $R$ -algebras. An R–scheme is a scheme over  $Spec(R)$ . A finite locally free R–module is the same as a finitely generated projective R–module, often abbreviated as finite projective R–module. If M is such a module,  $\mathbf{W}(M)$  denotes the R–scheme representing the R–functor  $T \mapsto M \otimes_R T$ ,  $T \in R$ -alg. For any unital associative algebra  $S$  we write  $S^{\times}$  for the set of invertible elements of S. Given an Azumaya algebra A, over a scheme or over a ring  $R$ , we follow [\[CF,](#page-62-1) 2.4.2.2 and 2.4.2.2] and denote by  $\mathbf{GL}_1(A)$  and  $\mathbf{PGL}(A)$  the group schemes of invertible elements and of automorphisms of A. Regarding cohomology,  $H^1(R, G) = H^1(\text{Spec}(R), G)$  is fppf-cohomology.

References to [\[SGA3\]](#page-63-0). Our paper has many references to [\[SGA3\]](#page-63-0). For better readability, a reference to [\[SGA3\]](#page-63-0) will simply be written by specifying the exposé in Roman numbers and the result in arabic numbers, but leaving out  $[\text{SGA3}]$ . For example,  $[\text{SGA3}, \text{XIV}, 3.20] = [\text{XIV}, 3.20]$ .

Acknowledgements. We thank Skip Garibaldi and Holger Petersson for having aroused our interest in LG-rings and for comments on an earlier version of this paper. We also thank Laurent Moret-Bailly for correspondence regarding Example [2.13.](#page-11-0)

#### <span id="page-3-0"></span>4 P. GILLE AND E. NEHER

## 1. LG-rings

This section gives an introduction to LG-rings (definition, examples, immediate consequences of the definition, preliminary results). In particular, we prove the important geometric characterization [1.4](#page-6-0) of LG-rings and of rings satisfying the primitive condition.

<span id="page-3-3"></span>1.1. LG-rings (definition and some known facts). For  $S \in R$ -alg we say that a polynomial  $g \in S[X_1, \ldots, X_n]$  represents a unit over S if there exist  $s_1, \ldots, s_n \in S$  such that  $g(s_1, \ldots, s_n) \in S^{\times}$ . We apply this notion for a polynomial  $f \in R[X_1, \ldots, X_n]$  by viewing f as a polynomial over S using the structure homomorphism  $R \to S$ . Observe that if  $f \in R[X_1, \ldots, X_n]$ represents a unit over R, then it represents a unit over every  $S \in R$ -alg. LG-rings are rings for which also the converse holds. Namely, we call  $R$  an LG-ring if for every  $n \in \mathbb{N}$  and every  $f \in R[X_1,\ldots,X_n]$  the polynomial f represents a unit over R if and only if one of the following obviously equivalent conditions hold:

- (i) f represents a unit over every localization  $R_{\mathfrak{m}}$ ,  $\mathfrak{m}$  a maximal ideal of R;
- (ii) f represents a unit over every field  $R/\mathfrak{m}$ ,  $\mathfrak{m}$  a maximal ideal of R;
- (iii) f represents a unit over every field  $F \in R$ -alg;
- (iv) f represents a unit over every  $S \in R$ -alg.

An LG-ring is sometimes called a "local-global ring", hence the short form LG, or a *ring with many units*. The following facts will be helpful:

<span id="page-3-4"></span>(a) (Direct products) Let  $R_1, \ldots, R_n$  be rings. Then the direct product  $R_1 \times \cdots \times R_n$  is an LG-ring if and only if every  $R_i$  is an LG-ring.

<span id="page-3-5"></span>(b) (*Characterization*) Recall that  $Jac(R)$  denotes the Jacobson radical of a ring R, i.e., the intersection of all maximal ideals of R. The following are equivalent for R:

<span id="page-3-1"></span>(i)  $R$  is an LG-ring;

<span id="page-3-2"></span>(ii)  $R/\mathrm{Jac}(R)$  is an LG-ring.

The equivalence [\(bi\)](#page-3-1)  $\iff$  [\(bii\)](#page-3-2) holds because the maximal ideals of R and of  $R/\mathrm{Jac}(R)$  are in obvious bijection.

(c) (*Finite modules*) Let  $M$  be a finitely presented and let  $N$  be a finitely generated module over an LG-ring R. Then  $M \cong N$  if and only if  $M_{\mathfrak{m}} \cong$  $N_{\rm m}$  for all maximal ideals  $\mathfrak{m} \triangleleft R$  [\[EG,](#page-62-2) Thm. 2.6]. In particular, any finite projective R–module of constant rank is free [\[EG,](#page-62-2) Thm. 2.10], [\[MW,](#page-63-1) II, Thm.], and therefore

<span id="page-3-6"></span>(1.1.1) 
$$
Pic(R) = \{0\}.
$$

We will give a quick proof, different from the published ones, of the last statement in Corollary [1.6](#page-7-1)[\(b\)](#page-7-2).

<span id="page-4-0"></span>1.2. Examples of LG-rings and some history. (a) Clearly,  $\{0\}$  is an LG-ring and so is every field. It then follows from  $1.1(a)$  $1.1(a)$  and  $1.1(b)$  $1.1(b)$  that every semilocal ring is an LG-ring.

<span id="page-4-4"></span>(b) If R is an LG-ring and  $R' \in R$ -alg is an integral extension, then R' is also an  $LG\text{-ring}$  by [\[EG,](#page-62-2) Cor. 2.3]. In particular, every finite R-algebra of an LG-ring R is itself an LG-ring. (Recall that an  $R$ –algebra S is finite if and only if S is integral and of finite type as  $R$ –algebra [\[St,](#page-64-0) Tag 00GN].

(c) Condition [1.1](#page-3-3)?? implies that every homomorphic image of an  $LG$ -ring is an LG-ring. Similarly, any direct limit of LG-rings is an LG-ring  $[EG,$  $|2|$ .

<span id="page-4-1"></span>(d) If  $R/Jac(R)$  is von Neumann regular, then R is an  $LG$ -ring  $\langle \text{MW}, \rangle$ I, Prop.]. Recall [\[B:AC1,](#page-62-3) I, §2, Exc. 17] that a commutative ring A is von Neumann regular if and only if A is absolutely flat, i.e., all A–modules are flat.

(e) A zero-dimensional ring is an  $LG\text{-}ring$  [\[MW,](#page-63-1) I, Cor.]. Indeed, a ring R is zero-dimensional if all its prime ideals are maximal. For such a ring the Jacobson radical Jac $(R)$  equals the nil radical, and so  $R/\text{Jac}(R)$  is absolutely flat = von Neumann regular by  $[B:AC1, II, §4, Exc. 16(d)]$  or by  $[St, Tag]$ 092F. Therefore R is LG by  $(d)$ .

<span id="page-4-3"></span>(f) A polynomial in  $R[X_1, \ldots, X_n]$  is called *primitive* if its coefficients generate  $R$  as ideal. One says that a ring  $R$  satisfies the primitive criterion [\[EG,](#page-62-2) [MW\]](#page-63-1) if the following equivalent conditions hold:

- (I) for every primitive polynomial  $P \in R[X]$  there exists  $r \in R$  such that  $P(r) \in R^{\times}$ ;
- (II) for every primitive  $Q \in R[X_1, \ldots, X_n]$  there exists  $(r_1, \ldots, r_n) \in R^n$ such that  $Q(r_1, \ldots, r_n) \in R^{\times};$
- <span id="page-4-2"></span>(III) R is LG and all residue fields are infinite.

An example of a ring satisfying the primitive criterion, is the ring  $S^{-1}R[X]$ where  $R$  is arbitrary and  $S$  is the multiplicative subset of all primitive polynomials in the polynomial ring  $R[X]$ , [\[vdK,](#page-64-1) 1.13].

Another interesting example is the ring of all algebraic integers or of all real algebraic integers is an LG-ring. That these rings are LG-rings, is shown in [\[Da,](#page-62-4) [EG\]](#page-62-2). Since they do not have homomorphic images of finite characteristic, [\(III\)](#page-4-2) proves our claim.

(g) Non-examples: The rings  $\mathbb Z$  and  $R[X]$ , R an integral domain, are not LG-rings [\[GPR2,](#page-62-5) Exc. 11.42].

(h) Some history of LG-rings. It seems that the concept of an LG-ring goes back to the paper [\[MW\]](#page-63-1) by McDonald and Waterhouse, where the authors refer to such a ring as a "ring in which every polynomial with local unit values has unit values". The motivation of  $[MW]$  comes from  $K$ -theory, in particular the study of  $GL_2(R)$  and  $Aut (GL_2(R))$ . The name "LG-ring" was introduced in the paper [\[EG\]](#page-62-2) by Estes and Guralnick, which is the first in-depth investigation of LG-rings. Later on, LG-rings were used by Dias [\[Di\]](#page-62-6) in her Ph.D. thesis with the goal of extending some classical theorems on quadratic forms over fields to quadratic forms over LG-rings (her result on Witt cancellation of quadratic forms is generalized in [5.5\)](#page-29-1). The subject seem to have fallen in oblivion until it was recently resurrected by Garibaldi-Petersson-Racine in their paper [\[GPR1\]](#page-62-7) and the forthcoming book [\[GPR2\]](#page-62-5).

<span id="page-5-0"></span>Our first goal is to give a more geometric approach to LG-rings, see [1.3](#page-5-0) and [1.4.](#page-6-0)

1.3. Lemma. Let R be an LG-ring and let  $n \geq 1$  be an integer. We further assume that U is an open quasi-compact subscheme of  $\mathbb{A}^n_R = X$  and let I  $\triangleleft$  $R[X_1, \ldots, X_n]$  be the radical ideal such that  $U = X \setminus \text{Spec}(R[X_1, \ldots, X_n]/I)$ . Hence  $U = \bigcup_{f \in I} X_f$ .

<span id="page-5-1"></span>(a) Denoting by  $I_{\text{prim}}$  the set of primitive polynomials in I, we have

$$
\bigcup_{f \in I_{\text{prim}}} X_f(R) = \bigcup_{f \in I} X_f(R) = U(R).
$$

<span id="page-5-3"></span>(b) Moreover, if  $U(R/\mathfrak{m}) \neq \emptyset$  for every maximal ideal  $\mathfrak{m} \triangleleft R$ , then also  $U(R) \neq \emptyset$ .

*Proof.* [\(a\)](#page-5-1) The first equality is obvious since  $X_f(R) = \emptyset$  for  $f \in I \setminus I_{\text{prim}}$ . The inclusion  $\bigcup_{f\in I}X_f(R)\subset U(R)$  is obvious. For the converse we are given  $u \in U(R)$  and need to find  $f \in I$  such that  $u \in X_f(R)$ . Since U is quasi-compact,the ideal I is finitely generated ([\[EGA-I,](#page-63-2) I,  $(1.1.4)$ ]), say by  $f_1, \ldots, f_d$ . Let us introduce the auxiliary linear polynomial in the variables  $Y_1, \ldots, Y_d$ :

$$
P(Y_1, ..., Y_d) = f_1(u)Y_1 + \dots + f_d(u)Y_d.
$$

We claim:

<span id="page-5-2"></span>(1.3.1) for every maximal ideal  $\mathfrak{m} \triangleleft R$ , P represents a unit in  $(R/\mathfrak{m})^{\times}$ .

Indeed, fix a maximal ideal  $\mathfrak{m} \triangleleft R$  and put  $\kappa = R/\mathfrak{m}$ . We have  $U(\kappa) =$  $\bigcup_{g\in I\otimes_{R}\kappa}X_g(\kappa)$ . Hence  $u\otimes 1_{\kappa}\in X_g(\kappa)$  for some  $g\in I$ , i.e.,  $g(u\otimes 1_{\kappa})\in \kappa^{\times}$ . We can write g in the form  $g = f_1b_1 + \cdots + f_d b_d$  with suitable  $b_i \in R$  and then have  $P(b_1, \ldots, b_d) = f_1(u)b_1 + \cdots + f_d(u)b_d = g(u) \otimes 1_{\kappa} \in \kappa^{\times}$ , establishing the claim  $(1.3.1)$ .

The LG-property now applies and says that  $P$  represents a unit in  $R$ , say  $P(a_1, \ldots, a_d) \in R^{\times}$  for suitable  $a_i \in R$ . Putting

$$
(1.3.2) \t\t f = a_1 f_1 + \dots + a_d f_d \in I \subset R[X_1, \dots, X_n],
$$

we have  $f(u) = a_1 f_1(u) + \cdots + a_d f_d(u) = P(a_1, \ldots, a_d) \in R^{\times}$ , i.e.,  $u \in$  $X_f(R)$ .

[\(b\)](#page-5-3) We consider the polynomial

<span id="page-5-4"></span>
$$
Q = f_1(X_1, ..., X_n)Y_1 + \dots + f_d(X_1, ..., X_n)Y_d
$$
  
\n
$$
\in R[X_1, ..., X_n, Y_1, ..., Y_d].
$$

We have seen in [\(a\)](#page-5-1) that the assumption of [\(b\)](#page-5-3) implies that  $Q$  represents a unit in  $R/\mathfrak{m}$  for every maximal ideal  $\mathfrak{m} \triangleleft R$ . Hence, since R is LG, the

polynomial Q represents a unit in R, i.e., there exist  $(u_1, \ldots, u_n) \in R^n$  and  $(a_1, \ldots, a_d) \in R^d$  such that

$$
Q(u_1,\ldots,u_n,a_1,\ldots,a_d)\in R^{\times}.
$$

It then follows that  $u \in X_f(R)$  for the polynomial  $f \in I$  defined in [\(1.3.2\)](#page-5-4), in particular  $U(R) \neq \emptyset$ .

<span id="page-6-1"></span><span id="page-6-0"></span>1.4. Proposition. Let R be a LG-ring, let M be a finite locally free Rmodule, and let  $U \subset \mathbf{W}(M)$  be an open quasi-compact subscheme.

- <span id="page-6-2"></span>(a)  $U(R) \neq \emptyset \iff U(R/\mathfrak{m}) \neq \emptyset$  for every maximal ideal  $\mathfrak{m} \triangleleft R$ .
- (b) If R satisfies the primitive condition as in  $1.2(f)$  $1.2(f)$  and U is R-dense, then  $U(R) \neq \emptyset$ .

Proof. [\(a\)](#page-6-1) Of course, only the implication from right to left is to show. To do so, we take a presentation  $M \oplus N = R^n$ , hence  $\mathbf{W}(M) \times_R \mathbf{W}(N) = \mathbb{A}^n_R$ . It then follows from [\[EGA,](#page-63-3) IV<sub>1</sub>, 1.1.2] (or [\[St,](#page-64-0) Tag 01K5]) that  $V = U \times_R \mathbf{W}(N)$ is an open quasi-compact R-subscheme of  $\mathbb{A}_R^n$ . Moreover,  $V(R/\mathfrak{m}) \neq \emptyset$  for each maximal ideal m of R since  $\mathbf{W}(N)(R/\mathfrak{m}) = N \otimes_R (R/\mathfrak{m}) \neq 0$ . Now [1.3\(](#page-5-0)[b\)](#page-5-3) shows that  $U(R) \times N = V(R) \neq \emptyset$ , so that also  $U(R) \neq \emptyset$ .

[\(b\)](#page-6-2) Let m be a maximal ideal of R, and put  $k = R/m$ . By R-denseness, the k–variety  $U_k$  is an open dense subvariety of the affine k–space  $\mathbf{W}(M_k)$ , in particular  $U_k$  is a rational non-empty subvariety of  $\mathbb{A}_k^n$ . Since k is an infinite field by [1.2](#page-4-0)[\(f\)](#page-4-3), it follows that  $U_k(k) = U(k) \neq \emptyset$ . Now [\(a\)](#page-6-1) implies that  $U(R) \neq \emptyset$ .

Remark. Proposition [1.4](#page-6-0)[\(a\)](#page-6-1) characterizes LG-rings. More precisely, if R is any base ring such that for every finite locally free  $R$ -module and every open quasi-compact  $U \subset \mathbf{W}(M)$  we have the equivalence [\(a\),](#page-6-1) then R is an LG-ring.

Indeed, it suffices to evaluate the condition for a free R-module M of finite rank and a principal open affine  $U \subset U(M)$ .

<span id="page-6-3"></span>1.5. Faithfully projective modules. Recall that an  $R$ –module M is faithful if the structure map  $R \to \text{End}_R(M)$ ,  $r \mapsto r \text{Id}_M$ , is injective. It is a standard fact in commutative algebra that the following conditions are equivalent for a finite projective  $R$ –module  $P$ :

- $(i)$  P is faithful;
- (ii) every localization  $P_{\mathfrak{p}}$ ,  $\mathfrak{p} \in \text{Spec}(R)$ , is non-zero;
- (iii)  $P_{R/\mathfrak{m}} \neq \{0\}$  for every maximal ideal  $\mathfrak{m} \triangleleft R$ ;
- (iv)  $P$  is faithfully flat;

(v) there exists an R–module Q such that  $P \otimes_R Q \cong R^n$  for some  $n \in \mathbb{N}_+$ .

In this case, P is called *faithfully projective*.

The following corollary is a first application of Proposition [1.4.](#page-6-0) An element m of a finite projective R–module M is called unimodular if  $m \otimes_R$  $1_{\kappa(\mathfrak{p})}\neq 0$  for all  $\mathfrak{p} \in \text{Spec}(R)$ , equivalently,  $R \cdot m$  is a free R-module of rank 1 and a direct summand of  $M$  ([\[GPR2,](#page-62-5) 9.17], [\[Lo1,](#page-63-4) 0.3]).

#### <span id="page-7-1"></span>8 P. GILLE AND E. NEHER

<span id="page-7-3"></span>1.6. Corollary. Let R be an LG-ring and let M be a finite locally free  $R$ module.

- <span id="page-7-2"></span>(a) If M is faithfully projective, M contains a unimodular element.
- (b) If  $M$  has constant rank, then  $M$  is free.

*Proof.* [\(a\)](#page-7-3) We consider the open subset  $U = W(M) \setminus \text{Spec}(R)$ , that is, the punctured affine space associated with the R-module M. Thus  $U(R)$ consists of the unimodular elements of  $\mathbf{W}(M)(R) = M$ . We have  $\mathbf{W}(M) =$ Spec(Sym $(M^*)$ ) where Sym $(M^*)$  is the symmetric algebra of M, and  $R \cong$  $Sym(M^*)/I$  where I is the ideal generated by finitely many linear forms  $\lambda_1, \ldots, \lambda_n$  spanning  $M^*$ . Hence, by [\[EGA-I,](#page-63-2) I, (1.1.4)], U is quasi-compact. Clearly,  $U(R/\mathfrak{m}) \neq \emptyset$ . Therefore Proposition [1.4](#page-6-0)[\(a\)](#page-6-1) shows that  $U(R)$  is not empty.

[\(b\)](#page-7-2) We prove the statement by induction on the rank d of  $M$ , starting with the obvious case  $d = 0$ . In the following we will assume that  $d \geq 1$ . Then M is faithfully projective and therefore contains a unimodular element  $m \in M$ . Thus  $M = Rm \oplus M'$  with M' being a finite projective R-module of rank  $d-1$ . By the induction hypothesis, M' is free of rank  $d-1$ , so M is free of rank d.  $\Box$ 

#### 2. Existence of maximal tori

<span id="page-7-0"></span>In this section our goal is to generalize Grothendieck's Theorem on the existence of maximal tori in reductive group schemes  $[\text{SGA3}, \text{XIV}, 3.20(*)]$ from the semilocal case to the LG case, see Theorem [2.6.](#page-9-0) Our proof is a mild modification of the original proof, and is based on the exposés XIII and XIV of [\[SGA3\]](#page-63-0).

Throughout this section  $G$  is a reductive group scheme defined over an arbitrary ring R, unless specified otherwise. We abbreviate  $\mathfrak{g} = \text{Lie}(G)(R)$ , which is a Lie R–algebra whose underlying module is finite projective.

2.1. The conditions  $(C_0)$ ,  $(C_1)$ ,  $(C'_1)$  and  $(C_2)$  of  $[{\rm XIV},$  2.9] hold for Lie(G). The exposes XIII and XIV of  $[SGA3]$  are written in a more general setting than the reductive case. For example, several results of loc. cit. assume the conditions  $(C_0)$ ,  $(C_1)$ ,  $(C'_1)$  and/or  $(C_2)$  of [XIV, 2.9]. We claim: These conditions are fulfilled in the reductive case. Indeed, the nilpotent rank equals the reductive rank in this case, which is locally constant by  $[XIX, 2.6]$ , i.e.,  $(C_0)$  holds. Moreover, the proof of  $[XIV, 3.7]$  shows that quasi-regular sections are regular for Lie algebras of reductive groups, which proves  $(C_2)$ , and thereby also  $(C_1)$  and  $(C'_1)$  in view of [XIV, 2.9].

<span id="page-7-4"></span>2.2. Lemma. For  $A \in R$ -alg and  $\mathfrak{g}_A = \mathfrak{g} \otimes_R A$  we denote by  $\mathfrak{g}_A^{\text{reg}}$  $A^{\text{reg}}$  the set of regular elements of the Lie A–algebra  $\mathfrak{g}_A$ , defined in [XIV, 2.5]. Then the R–  $\mathit{functor}\ A \mapsto (\mathfrak{g}_A)^\mathrm{reg}$  is representable by an open quasi-compact  $R$ –subscheme U of  $\mathbf{W}(\mathfrak{g})$ . Furthermore, U is R-dense in  $\mathbf{W}(\mathfrak{g})$ .

*Proof.* Since the condition  $(C_0)$  holds for g, the representability of the given R–functor by an open subscheme of  $W(g)$  follows from [XIV, 2.10].

Because  $\mathbf{W}(\mathfrak{g})$  is an affine scheme, the structure morphism  $\mathbf{W}(\mathfrak{g}) \rightarrow$  $Spec(R)$  is quasi-compact in the sense of  $[St, Tag 01K3]$ . Hence, if the immersion  $\iota: U \to \mathbf{W}(\mathfrak{g})$  is also a quasi-compact morphism, then so is the structure morphism  $U \to \text{Spec}(R)$  since quasi-compact morphisms respect composition [\[St,](#page-64-0) Tag 01K6]. It is therefore sufficient to show that  $\iota$  is a quasi-compact morphism.

To do so, we will apply noetherian reduction, writing  $R = \varinjlim_{i \in I} R_i$  as the direct limit of its finitely generated, hence noetherian Z–subalgebras. It follows from [VI<sub>B</sub>, 10.1, 10.3] and [\[Co1,](#page-62-8) 3.1.11] that there exists  $i \in I$  and a reductive  $R_i$ -group scheme  $G_i$  such that  $G = G_i \times_{R_i} R$ . Let  $\mathfrak{g}_i = \text{Lie}(G_i)(R_i)$ and let  $U_i$  be the open subscheme of  $\mathbf{W}(\mathfrak{g}_i)$  representing regular elements. Since the formation of U commutes with base change, we can assume  $U =$  $U_i \times_{R_i} R$ . Furthermore, the open immersion  $\iota$  is obtained by base change from the open immersion  $\iota_i: U_i \to \mathbf{W}(\mathfrak{g}_i)$ . Because quasi-morphisms respect base change [\[St,](#page-64-0) 01K5], it is now sufficient to show that  $\iota_i$  is a quasi-compact morphism. This is indeed true: Since  $R_i$  is noetherian,  $\mathbf{W}(\mathfrak{g}_i)$  is a noetherian scheme and the open immersion  $\iota_i$  is q quasi-compact morphism by [\[St,](#page-64-0) Tag 01OX]. Thus, U is indeed quasi-compact.

Finally, since Lie algebras of reductive groups over infinite fields admit regular elements [XIV, 2.11(b)], U is R-dense in  $\mathbf{W}(\mathfrak{g})$ .

<span id="page-8-0"></span>2.3. Cartan subalgebras and regular elements. Let  $S = \text{Spec } R$ . A *Cartan subalgebra* of the Lie algebra  $Lie(G)$  over S is a Lie subalgebra D of Lie(G) which is locally a direct summand and whose geometric fibres  $D_{\bar{s}}$ are Cartan subalgebras of Lie( $G_{\bar{s}}$ ) for all  $s \in S$  [XIV, 2.4]. The concept of a Cartan subalgebra of a finite-dimensional Lie algebra over an infinite field, is defined in [XIII, after 4.5]. In this special case, they are the nil spaces of regular elements, and therefore always exist since regular elements exist.

One can use the equivalence of quasi-coherent  $\mathscr{O}_S$ -modules and R-modules to translate the definition of a Cartan subalgebra of  $Lie(G)$  to that of a Cartan subalgebra of the R-Lie algebra  $\mathfrak{g} = \text{Lie}(G)(R)$ . In particular, a Cartan subalgebra of  $\mathfrak g$  is always a direct summand of the finite projective Lie algebra g.

The R–functor, associating with  $A \in R$ -alg the set of Cartan algebras of  $\mathfrak{g}_A$ , is representable by a quasi-projective finitely presented R-scheme  $\mathscr{D}$ [XIV, 2.16]. Also, the R-subfunctor X of  $\mathscr{D} \times_R W(\mathfrak{g})$ , defined by

$$
X(A) = \{ (\mathfrak{h}, y) \in \mathscr{D}(A) \times \mathfrak{g}_A : y \in \mathfrak{h} \}
$$

for each R–algebra A, is representable by a quasi-projective R–scheme  $\mathscr X$ . Furthermore, the first projection  $p_1 : \mathcal{X} \to \mathcal{D}$  is a vector bundle.

We denote by  $\psi : \mathscr{X} \to W(\mathfrak{g})$  the second projection, and recall that the restriction  $\psi^{-1}(U) \to U$  is an isomorphism, where  $U \subset \mathbf{W}(\mathfrak{g})$  is the open

scheme of [2.2.](#page-7-4) In particular, this implies that a regular element of  $\mathfrak{g}_A$  is contained in a unique Cartan subalgebra of  $g_A$ .

<span id="page-9-3"></span><span id="page-9-1"></span>2.4. **Theorem.** We assume that  $R$  is an  $LG$ -ring and that one of the following conditions hold:

- <span id="page-9-2"></span>(i) R satisfies the primitive criterion;
- (ii) G is adjoint.

Then g admits a regular element and hence a Cartan R–subalgebra.

Proof. By [2.3,](#page-8-0) a regular element is contained in a (unique) Cartan subalgebra of  $\mathfrak g$ . It is therefore enough to prove that  $\mathfrak g$  contains regular elements.

Case [\(i\):](#page-9-1) Regular elements always exist in finite-dimensional Lie algebras over infinite fields [XIII, 4.2], i.e.,  $U(R/\mathfrak{m}) \neq \emptyset$  for all infinite residue fields of R. Since  $U \subset W(\mathfrak{g})$  is open and quasi-compact by [2.2,](#page-7-4) Proposition [1.4](#page-6-0)[\(b\)](#page-6-2) shows that  $U(R) \neq \emptyset$ .

*Case* [\(ii\):](#page-9-2) Applying again Proposition  $1.4(b)$  $1.4(b)$  and Lemma [2.2,](#page-7-4) it suffices to show that  $U(R/\mathfrak{m})\neq \emptyset$  for every maximal ideal  $\mathfrak{m} \triangleleft R$ . As mentioned in [\(i\),](#page-9-1) this is always true if  $R/\mathfrak{m}$  is infinite. But it is also true if  $R/\mathfrak{m}$  is finite and [\(ii\)](#page-9-2) holds, according to a result of Chevalley-Serre [XIV, App.].  $\Box$ 

<span id="page-9-4"></span>2.5. Groups of type (C) and Cartan subalgebras. Let  $S = Spec(R)$ . We recall that a smooth R–subgroup  $H \subset G$  with connected geometric fibers is a subgroup of type (C) if  $Lie(H)$  is a Cartan  $\mathcal{O}_S$ -Lie subalgebra of Lie(G). The map  $H \mapsto \text{Lie}(H)(R)$  is a bijection between subgroups of type (C) of G and the R–Cartan subalgebras of  $Lie(G)(R)$ , [XIV, 3.9]. The inverse map is given by  $\mathfrak{h} \mapsto \text{Norm}_G(\mathfrak{h})^0$ . We shall use that R–subgroups of type (C) are precisely the maximal tori of  $G$  when  $G$  is adjoint,  $[XIV, 3.18]$ .

<span id="page-9-0"></span>2.6. Theorem. Let R be an LG-ring and let G be a reductive  $R$ -group scheme. Then G admits a maximal R–torus.

*Proof.* Let  $\mathfrak{Z}(G)$  be the (schematic) centre of G. The flat quotient  $G/\mathfrak{Z}(G)$  is represented by a semisimple adjoint R–group scheme  $G^{ad}$ , [XXII, 4.3.5(ii)] or [\[Co1,](#page-62-8) 3.3.5]. Moreover,  $T \mapsto T/\mathcal{Z}(G)$  defines a bijective correspondence between the set of maximal tori of  $G$  and the set of maximal tori of  $G^{\text{ad}}$ . Thus, without loss of generality we can assume that  $G$  is adjoint. Then Theorem [2.4](#page-9-3)[\(ii\)](#page-9-2) provides a Cartan R-algebra of  $\mathfrak g$ , which by [2.5](#page-9-4) "integrates" to a maximal torus of  $G$ .

<span id="page-9-7"></span>2.7. Lemma (Tori in subgroups of type  $(RC)$ ). Let G be a reductive group scheme over a ring R, let  $H \subset G$  be a subgroup scheme of G of type (RC) in the sense of [XXII, 5.11.1] and let rad<sup>u</sup>(H) be the unipotent radical of H, [XXII, 5.11.4]. Suppose:

- <span id="page-9-5"></span>(i) every reductive R–group scheme admits a maximal torus, e.g., assume that R is an LG-ring, and
- <span id="page-9-6"></span>(ii)  $H^1(R, \text{rad}^u(H)) = 0$ , e.g., assume that there exists a parabolic subgroup scheme  $P \subset G$  such that  $\text{rad}^u(H) = \text{rad}^u(P) \cap H$ .

#### Then G admits a maximal torus contained in H.

Proof. (Modelled after the proof of [XXVI, 4.2.7(ii)]) The unipotent radical  $U = rad<sup>u</sup>(H)$  is a smooth, finitely presented, and normal subgroup scheme of H, whose geometric fibres are connected and unipotent. Furthermore, by [XXII, 5.11.4], we have an exact sequence of  $R$ –group schemes

$$
1\longrightarrow U\longrightarrow H\stackrel{f}\longrightarrow M\longrightarrow 1
$$

where M is reductive. The subgroup H is in particular of type  $(R)$ , so that H and G have the same rank by [XXII, 5.2.2(b)]. By the properties of  $U$ , this is then also the rank of M.

The assumption [\(i\)](#page-9-5) provides us with a maximal  $R$ –torus  $T$  of  $M$ . Let  $N = f^{-1}(T)$  be its pre-image. We thus have an induced exact sequence

$$
1\longrightarrow U\longrightarrow N\stackrel{f}\longrightarrow T\longrightarrow 1
$$

of R–group schemes. The properties of U and T, together with  $[V]_B$ , 9.2(viii), imply that N is a smooth and finitely presented subgroup scheme of G. Moreover, by [XVII, 5.1.1(i)(a)], its geometric fibres  $N_{\bar{s}}$ ,  $s \in \text{Spec}(R)$ , are connected, solvable and contain a maximal torus of  $G_{\bar{s}}$ . Thus, N is a subgroup scheme of G of type (R). We have rank  $N = \text{rank } M = \text{rank } G$ , and by [XXII, 5.6.9(ii)] also  $\text{rad}^u(N) = \text{rad}^u(H) = U$ .

By  $[XXII, 5.6.13]$ , the functor of maximal tori of N is representable by an R–scheme  $\mathscr{T}$ , which is a U–torsor. It then follows from assumption [\(ii\)](#page-9-6) that  $\mathcal{T}(R) \neq \emptyset$ , i.e., N admits a maximal torus T, which is a maximal torus of G because rank  $N = \text{rank } G$ . Hence  $T \subset N \subset H$  fulfills our claim.

That  $H^1(R, \text{rad}^u(H)) = 0$  in case there exists a parabolic subgroup scheme  $P \subset G$  such that  $rad^{u}(H) = rad^{u}(P) \cap H$ , is shown in [XXVI, 2.11 (use [XXVI, 2.5]).

<span id="page-10-0"></span>2.8. Corollary (Tori in parabolic subgroups). Let R be an arbitrary ring satisfying the condition [2.7](#page-9-7)[\(i\)](#page-9-5), e.g., suppose that R is an LG-ring, and let G be a reductive  $R$ -group scheme. Then every parabolic subgroup of  $G$  contains a maximal torus of G.

*Proof.* A parabolic subgroup P of G is a group of type  $(RC)$  by  $[XXVI, 1.5]$ . The condition [2.7](#page-9-7)[\(ii\)](#page-9-6) holds by [XXVI, 2.2]. Hence Lemma 2.7 with  $H = P$ proves the claim.

Corollary [2.8](#page-10-0) is the special case  $P = Q$  of Proposition [3.4,](#page-16-0) whose proof requires however much more work. We will present an immediate application of Corollary [2.8](#page-10-0) to quasi-split groups.

2.9. Quasi-split reductive groups. We refer the reader to [XXIV, 3.9] for the definition of a quasi-split reductive group over an arbitrary scheme S. It simplifies greatly if S is an LG-scheme, i.e.,  $S = \text{Spec}(R)$  for R an LG-ring:

<span id="page-11-1"></span>(a) If S is an LG-scheme, a reductive  $S$ -group G is quasi-split if and only if G admits a Borel subgroup.

Indeed, it is explained in  $[XXIV, 3.9]$  that a reductive group G over a scheme S with  $Pic(S) = 0$  is quasi-split if and only if it contains a Killing couple, i.e., a pair  $(B, T)$  consisting of a Borel subgroup B of G and a maximal torus T of G contained in T. But, by Corollary [2.8,](#page-10-0) every Borel subgroup contains a maximal torus of  $G$ . We note that [\(a\)](#page-11-1) is the definition of a quasi-split reductive  $S$ –group in  $[Co1, 5.2.10]$ .

Recall( $[Co1, §7.2]$  $[Co1, §7.2]$  or  $[CF, 2.2.4.9]$ ) that an *inner form* of a reductive S–group scheme  $G$  is a twisted form of  $G$  under a torsor in the image of  $H^1(S, G/\mathcal{Z}(G))$  in  $H^1(S, \text{Aut}(G)).$ 

<span id="page-11-2"></span>(b) For an LG-scheme S, up to isomorphism every reductive S–group admits a unique quasi-split inner form.

This is proven in  $\lbrack \text{Co1, 7.2.12} \rbrack$  for semilocal S, but as  $\lbrack \text{Co1, 7.2.13} \rbrack$  states, [\(b\)](#page-11-2) holds whenever  $Pic(S') = 0$  for any finite étale cover S' of S. By [1.2\(](#page-4-0)[b\)](#page-4-4), such a cover is again an LG-scheme, so that  $(1.1.1)$  establishes the condition  $Pic(S')=0.$ 

<span id="page-11-3"></span>2.10. Split reductive groups ([XXIV, 2.14]). Let S be a scheme for which  $Pic(S) = 0$ , and let G be a reductive S-group scheme of constant type. Then G is split if and only if G contains a split maximal torus.

The result applies to  $S = \text{Spec}(R)$  for R an LG-ring since by [\(1.1.1\)](#page-3-6) we know that  $Pic(R) = 0$ . Moreover, by Theorem [2.6](#page-9-0) we know that a reductive group scheme G over an LG-ring R contains a maximal torus T.

2.11. **Isotrivial reductive groups.** A reductive group scheme G over S is *isotrivial* if there exists a finite étale cover (= finite étale surjective)  $S' \rightarrow S$ such that  $G_{S'}$  is split [\[Gi1,](#page-62-9) 4.4.1]. By [Gi1, 2.3], an isotrivial reductive S–group scheme  $G$  is necessarily of constant type. If  $G$  is of constant type with underlying Chevalley group scheme  $G_0$  (over  $\mathbb{Z}$ ), then G is isotrivial if and only if the Aut $(G_0)$ -torsor  $\underline{\mathrm{Isom}}(G_{0,S}, G)$  is isotrivial.

<span id="page-11-4"></span>2.12. Corollary. Let R be an  $LG$ -ring and let G be a reductive  $R$ -group scheme of constant type which is linear, e.g., G is semisimple [\[Gi2,](#page-62-10) Cor. 4.3]. Then  $G$  is isotrivial. In particular, if  $R$  is connected and simply connected  $(=$  does not admit non-trivial finite étale covers), then G is split.

*Proof.* Theorem [2.6](#page-9-0) provides a maximal  $R$ –torus  $T$  of  $G$ . Since  $T$  is linear (and of constant rank),  $T$  is isotrivial according to a result of Grothendieck, see [\[Gi2,](#page-62-10) Th. 3.3]. In other words, there exists a finite étale cover  $R'$  of R such that  $T_{R'}$  is split. Since  $R'$  is an LG-ring as well, we conclude that  $G_{R'}$  is split in view of  $2.10$ . Assume now that  $R$  connected and simply connected. Then  $R' = R \times \cdots \times R$ , so that T is already split.

<span id="page-11-0"></span>2.13. **Example.** The ring  $\overline{\mathbb{Z}}$  of algebraic integers is an LG-ring by [1.2](#page-4-0)[\(f\)](#page-4-3). It is also well-known that  $\overline{Z}$  is simply-connected. Hence, Corollary [2.12](#page-11-4) shows that any semisimple group scheme over  $\overline{Z}$  is split. In particular, this holds for the semisimple  $\overline{\mathbb{Z}}$ –group G of type G<sub>2</sub>, which is known to be the automorphism group scheme of an octonion algebra  $\mathcal O$  over  $\mathbb Z$  ([\[Co2,](#page-62-11) B.15] or [\[GPR2,](#page-62-5) 55.4]). As a consequence, O is split.

Moret-Bailly told us that this is a direct consequence of a theorem of Rumely on the existence of points for "reasonable" schemes over  $\overline{\mathbb{Z}}$  [\[Ru,](#page-63-5) Th. 1] (see also [\[MB,](#page-63-6) Th. 1.3]). For  $H$  a flat affine group scheme of finite type over  $\overline{\mathbb{Z}}$  whose generic fiber is (smooth) connected, Rumely's Theorem implies that any H-torsor has a  $\overline{\mathbb{Z}}$ -point, so is trivial. Since  $\overline{\mathbb{Z}}$  is simply connected a semisimple  $\overline{\mathbb{Z}}$ -group G is an inner form of its Chevalley form  $G_0$ , that is, is the twist of  $G_0$  by a  $G_{0,ad}$ -torsor E where  $G_{0,ad}$  is the adjoint group of  $G_0$ . Since  $E$  is trivial, it follows that  $G$  is split.

Coming back to the example of octonions, Rumely's theorem directly does the job, since the automorphism group  $G_2$  of the split octonions is connected. Of course, one could also appeal to the strong approximation theorem to show that  $H^1(\overline{\mathbb{Z}}, G_2)$  vanishes. Indeed, a class in  $H^1(\overline{\mathbb{Z}}, G_2)$ arises from  $\gamma \in H^1(R, G_2)$  where R is the ring of integers of a number field *F*. Furthermore we can assume that  $\gamma_F = 1$  since  $H^1(\overline{\mathbb{Q}}, G_2) = 1$ . But then [\[Ha,](#page-63-7) Satz 3.3] proves  $\gamma = 1$ .

Yet another way to see that any octonion algebra over the LG-ring  $\overline{Z}$  is split, has been communicated to us by Skip Garibaldi and works for any simply connected LG-ring  $R:$  By [\[GPR2,](#page-62-5) 19.16(b)], any octonion algebra over R contains a quadratic étale subalgebra. It is split, since R is simply connected. By  $[GPR2, 22.9]$ , the algebra  $\Theta$  is therefore reduced. But then it is split by  $[GPR2, 22.16]$  and  $1.6(b)$  $1.6(b)$ .

# <span id="page-12-0"></span>3. Parabolic subgroups (conjugacy, relative positions)

A fundamental result of Borel-Tits proves conjugacy of parabolic subgroups of the same type in reductive groups over a field  $[BT, Th. 4.13(a)].$ This was extended by Demazure to reductive group schemes over a semilocal ring [\[SGA3,](#page-63-0) XXVI, 5.2, 5.10]. In Theorem [3.1](#page-12-1) we prove this result for reductive groups over an LG-ring. In [3.2](#page-14-0) and [3.6](#page-19-0) we apply this to conjugate parabolic subgroups to a transversal or osculating position.

We use the notion of an opposite parabolic subgroup of a reductive group scheme as defined in [XXVI, 4.3.1]. Over an affine base, every parabolic subgroup admits an opposite parabolic subgroup  $[XXVI, 4.3.5(i)]$ . We will also employ the concept of the  $type(P) \in Of(Dyn(G))(R)$  of a parabolic subgroup  $P$  of  $G$ , defined in [XXVI, 3.2].

<span id="page-12-1"></span>3.1. Theorem. Let R be an  $LG\text{-ring}$ , and let G be a reductive R-group scheme with a pair  $(P, P')$  of opposite parabolic subgroup schemes of  $G$ . We abbreviate the unipotent radicals of P and P' by  $U = rad<sup>u</sup>(P)$  and  $U' = rad<sup>u</sup>(P')$  respectively.

<span id="page-12-2"></span>(a) Any parabolic subgroup  $Q$  of  $G$  of the same type as  $P$  has the form  $Q = u^u P$  for suitable  $u \in U(R)$  and  $u' \in U'(R)$ . In particular, the group  $G(R)$  acts transitively on  $(G/P)(R)$ .

- (b) The map  $H^1(R, P) \to H^1(R, G)$ , induced by the inclusion  $P \subset G$ , is injective.
- <span id="page-13-1"></span>(c) We have a decomposition  $G(R) = U(R) U'(R) P(R)$ .

*Proof.* [\(a\)](#page-12-2) By [XXVI, 3.6], we have an isomorphism  $G/P \longrightarrow \text{Par}_{\text{type}(P)}(G)$ where  $Par_{type(P)}(G)$  stands for the smooth projective R-scheme of parabolic subgroups of type type(P). Hence it suffices to prove the first part of [\(a\).](#page-12-2)

We consider the R-subfunctor  $\text{Opp}(P)$  of  $\text{Par}_{\text{type}(P)}(G)$  of parabolic subgroups which are opposite to  $P$ . By [XXVI, 2.5] (or [\[Co1,](#page-62-8) 5.4.3]), we have  $U \cong W(E)$  for a locally free R–module of finite type. (That E has finite type, is not explicitly stated in [XXVI, 2.5], but follows for example from the representability of  $\mathbf{W}(E)$ . Hence U is an affine smooth R-scheme. Furthermore, according to [XXVI, 4.3.6], we have a commutative diagram



This implies that  $Opp(P)$  is representable by an affine smooth R-scheme, since this is so for  $\overline{U}$ .

We consider the R-subfunctor  $\text{Opp}(P) \cap \text{Opp}((Q)$  of  $\text{Par}_{\text{type}(P')}$ ; it is representable by an open  $R$ -subscheme  $V$  of  $U$ . We claim that it is also quasi-compact. Indeed, we have a commutative diagram



in which the top rectangle is cartesian and c and  $d \circ b$  are quasi-compact morphisms since  $\text{Opp}(P)$  and  $\text{Opp}(Q)$  are affine schemes. As d is a pro-jective,hence separated morphism, cancellation ([\[EGA,](#page-63-3) IV<sub>1</sub>, 1.1.2] or [\[St,](#page-64-0) Tag  $03GI$ ) says that b is quasi-compact morphism. By base change, a is quasi-compact and so  $c \circ a$  is quasi-compact too, i.e., V is a quasi-compact scheme.

For every R-ring A, the set  $V(A)$  is the set of A–parabolic subgroups of  $G_A$  which are opposite to P and to Q. For a residue field  $R/\mathfrak{m}$  of R, two parabolic subgroups of  $G_{R/\mathfrak{m}}$  of the same type are conjugate by [\[BT,](#page-62-12) 4.3], so that  $[BT, 6.27]$  says that  $V(R/\mathfrak{m})$  is non-empty for each maximal ideal m of R. Since R is a LG-ring, Proposition [1.4](#page-6-0) implies that  $V(R) \neq \emptyset$ . In

<span id="page-13-0"></span>

other words,  $P$  and  $Q$  admit a common opposite  $R$ -parabolic subgroup  $Q'$ . We now use the isomorphism of R-functors [XXVI, 4.3.5.(i)]

$$
Opp(/P) \xrightarrow{\sim} \underline{\text{Lev}}(/P), \quad H \mapsto H \cap P
$$

between parabolic subgroups of  $G$  opposite to  $P$  and Levi subgroups of P, and note that it is P–equivariant. We consider the two Levi subgroups  $L = P' \cap P$  and  $M = Q' \cap P$  of P. Since P is a subgroup of type (RC), [XXVI, 1.8] says that  $M = uL$  for an unique  $u \in U(R)$ , so that  $Q' = uP'$  by using the above bijection. It follows that  $P' = u^{-1}Q'$  is opposite to P and to  $u^{-1}Q$ . Similarly, there exists a unique  $u' \in U'(R)$  such that  $u^{-1}P \cap P' = u'(P \cap P')$ and  $u^{-1}Q = u'P$ . Thus  $Q = u^{-1}u'P$ .

[\(b\)](#page-13-0) According to [\[Gir,](#page-62-13) III, 3.3.1], transitivity in [\(a\)](#page-12-2) implies that the map  $H^1(R, P) \to H^1(R, G)$  has trivial kernel. The classical twisting argument applies and yields the injectivity of the map  $H^1(R, P) \to H^1(R, G)$ , see for example the proof of [XXVI, 5.10].

[\(c\)](#page-13-1) Let  $g \in G(R)$  and apply [\(a\)](#page-12-2) to  $Q := {}^{g}P$ . Thus, there exists  $u \in U(R)$ , and  $u' \in U'(R)$  such that  $Q = u^2P$ . It follows that  $(uu')^{-1}g \in N_G(P)(R) =$  $P(R)$ . Thus  $g \in P(R) U'(R) U(R)$ . By applying this to  $g^{-1}$ , we obtain the desired decomposition  $G(R) = U(R) U'(R) P(R)$ .

<span id="page-14-0"></span>We can also extend [XXVI, 5.3] to LG-rings.

3.2. Proposition. Let R be an LG-ring, and let G be a reductive  $R$ -group scheme. Let  $P$ ,  $P'$ ,  $Q$  be three parabolic  $R$ -subgroups of  $G$ . Then there exists  $g \in G(R)$  such that  ${}^g\!Q$  and P as well as  ${}^g\!Q$  and P' are in transversal position.

*Proof.* We denote by  $Gen(Q/P)$  the open subscheme of G representing the subfunctor which assigns to an S-scheme S' the set of elements  $g \in G(S')$ such that  ${}^gQ_{S'}$  and  $P_{S'}$  are in transversal position [XXVI, 4.2.4(iii)]. We have to show that  $X = \text{Gen}(Q/P) \cap \text{Gen}(Q/P') \subset G$  has an R-point.

We claim that  $Gen(Q/P) \rightarrow G$  is quasi-compact. Indeed, since quasicompact morphisms allow (fpqc) descent (see for example [\[GW,](#page-63-8) 14.51]) and since  $(G, P)$  is splitable étale-locally [XXVI, 1.14], we can assume that G is split and that  $P = P_I$  is a standard parabolic subgroup. In this case,  $Gen(Q/P) \rightarrow G$  arises by base change from the analogous morphism over  $\mathbb{Z}$ . But the latter is quasi-compact since  $Spec(\mathbb{Z})$  is noetherian [\[St,](#page-64-0) Tag 01OX]. Since quasi-compact morphism allow base change [\[St,](#page-64-0) Tag 01K5], we are done. (Alternatively, one can use noetherian reduction as in the proof of [2.2,](#page-7-4) to prove that  $Gen(Q/P) \rightarrow G$  is quasi-compact.) In any case, since  $Gen(Q/P)$  and  $Gen(Q/P')$  are quasi-compact, so is their fibre product X.

We pick a parabolic subgroup  $Q'$  of G which is opposite to  $Q$ , (XXVI, 4.3.5(i)]), put  $U = rad^u(Q)$  and  $U' = rad^u(Q')$ , and let  $\iota: U \times_R U' \to G$ ,  $(u, u') \mapsto u u'$ , be the immersion obtained by restricting the open immersion

 $U \times_R Q' \to G$  ([XXVI, 4.3.2.b(vi)]). We then consider the fibre product



.

The right immersion is open and quasi-compact, hence so is  $V \hookrightarrow U \times_R U'$ . Moreover, by [XXVI, 2.5],  $U = W(M)$  for some locally free R-module M of finite rank. The analogous fact then holds for  $U \times_R U'$ .

With the aim of applying Proposition [1.4](#page-6-0)[\(a\)](#page-6-1) to  $V \subset U \times_R U' = W(M)$ , let  $\mathfrak{m} \triangleleft R$  be a maximal ideal of R, and put  $k = R/\mathfrak{m}$ . By [XXVI, 5.3] we know that there exists  $g \in G(k)$  such that g fulfills the claim of the proposition for  $R = k$ . By [3.1](#page-12-1)[\(c\)](#page-13-1) we can write g in the form  $g = uu'q$  with  $u \in U(k)$ ,  $u' \in U'(k)$  and  $q \in Q(k)$ . But then  $uu' \in V(k)$ . Hence, by [1.4](#page-6-0)[\(a\),](#page-6-1) we get  $V(R) \neq \emptyset$ . Thus  $X(R) \neq \emptyset$ , as desired.

Corollary [2.8](#page-10-0) is the special case  $P = Q$  of the following Proposition [3.4.](#page-16-0)

<span id="page-15-6"></span>3.3. Parabolic subgroups in standard position [XXVI, 4.5.1]. Let S be a scheme and let G be a reductive  $S$ -group scheme. Two parabolic subgroups  $P$  and  $Q$  of  $G$  are said to be in *standard position*, if they satisfy the following equivalent conditions  $(i)$ – $(iv)$ :

- <span id="page-15-4"></span><span id="page-15-0"></span>(i)  $P \cap Q$  is smooth.
- (ii)  $P \cap Q$  is a subgroup of type  $(R)$ .
- <span id="page-15-5"></span><span id="page-15-3"></span>(ii)'  $P \cap Q$  is a subgroup of type (RC).
- <span id="page-15-1"></span>(iii)  $P \cap Q$  contains (fpqc)-locally a maximal torus of G.
- (iv)  $P \cap Q$  contains Zariski-locally a maximal torus of G.

<span id="page-15-2"></span>Moreover, if  $S = \operatorname{Spec}(R)$  for a semilocal ring, then [\(i\)](#page-15-0)[–\(iv\)](#page-15-1) is equivalent to (v)  $P \cap Q$  contains a maximal torus.

If  $S$  is the spectrum of a field, every pair of parabolic subgroups is in standard position by  $[XXVI, 4.1.1]$  or  $[BT, 4.5]$ .

Our goal is to generalize condition [\(v\)](#page-15-2) to  $S = \text{Spec}(R)$  for R an LG-ring, see  $3.4(c)$  $3.4(c)$ . In order to do so, we need to retake part of the proof of [XXVI, 4.5.1]; at the same time we will add more details.

Regarding the equivalence of the conditions above, we note that the im-plications [\(ii\)](#page-15-4)<sup>'</sup>  $\implies$  (ii)  $\implies$  [\(i\)](#page-15-0) and [\(iv\)](#page-15-1)  $\implies$  [\(iii\)](#page-15-5) are trivial. The implication [\(i\)](#page-15-0)  $\implies$  [\(ii\)](#page-15-3)' is easy: because of [XXVI, 4.1.1] and [\[BT,](#page-62-12) 4.5], every geometric fibre  $K_{\bar{s}}$  of  $K = P \cap Q$  is connected and contains a maximal torus  $T_{\overline{s}}$  of  $G_{\overline{s}}$  such that the roots of  $K_{\overline{s}}$  with respect to  $T_{\overline{s}}$  are a closed subset of the roots of  $(G_{\overline{s}}, T_{\overline{s}})$ . Thus, if K is smooth, it is a subgroup of type (RC) by definition. We will show the remaining implications [\(iii\)](#page-15-5)  $\implies$ [\(i\)](#page-15-0) and [\(iii\)](#page-15-5)  $\implies$  [\(iv\)](#page-15-1) in the proof of [3.4,](#page-16-0) since they easily follow from the arguments in the proof of [3.4.](#page-16-0) The statements  $3.4(a)$  $3.4(a)$  and  $3.4(b)$  $3.4(b)$  below are

used in the proof of [\(iii\)](#page-15-5)  $\implies$  [\(iv\)](#page-15-1) in [XXVI, 4.5.1], which is the main point of the proof the equivalences in [XXVI, 4.5.1]

Regarding [\(3.4.1\)](#page-16-4) we note that  $P \cap Q$  is a subgroup of type (RC) by [\(ii\)](#page-15-3)' and hence has a unipotent radical rad<sup>u</sup> $(P \cap Q)$ , defined in [XXII, 5.11.4].

<span id="page-16-0"></span>3.4. Proposition. Let  $P$  and  $Q$  be two parabolic subgroups of a reductive S–group scheme G such that  $P \cap Q$  contains a maximal torus of G (fpqc)locally, cf. [3.3](#page-15-6). Then the following hold.

<span id="page-16-2"></span>(a) The (fpqc)–image sheaf of the multiplication morphism

 $f: \text{rad}^u(P) \rtimes (P \cap Q) \to G, \quad (u, h) \mapsto uh$ 

is representable by a parabolic subgroup  $P'$  of  $G$ , satisfying

<span id="page-16-4"></span> $P' = (P \cap Q) \cdot rad^u(P).$ 

<span id="page-16-3"></span>(b) The induced map  $f'$ :  $rad^u(P) \rtimes (P \cap Q) \rightarrow P'$  is smooth and

<span id="page-16-1"></span>(3.4.1) 
$$
\operatorname{rad}^u(P \cap Q) = \operatorname{rad}^u(P') \cap (P \cap Q).
$$

(c) If  $S = \text{Spec}(R)$  with R an LG–ring, then  $P \cap Q$  contains a maximal S-torus of G.

*Proof.* We put  $K = P \cap Q$  and prove [\(a\)](#page-16-2) and [\(b\)](#page-16-3) together. These statements are local with respect to the (fpqc)-topology. Hence, after passing to a suitable cover, we can assume that  $K$  contains a split maximal torus  $T$  and that  $P$  and  $Q$  have constant type.

We let R be the root system of  $(G, T)$ , choose an order on R and denote by  $U_r$ ,  $r \in R$ , the associated root subgroups. First some general reminders. For each closed subset  $R_0$  of R, we have an S-subgroup  $H_{R_0}$  of type  $(R)$ , which is characterized by its Lie algebra:  $\text{Lie}(H_{R_0}) = \text{Lie}(T) \oplus \bigoplus_{r \in R_0} \text{Lie}(U_r)$ , [XXII, 5.4.2, 5.4.7]. The multiplication map

$$
\prod_{r \in R_0 \cap R_+} U_r \times_S T \times_S \prod_{r \in R_0 \cap R_+} U_r \longrightarrow H_{R_0}
$$

is an open immersion, [XXII, 5.4.4], whose image we denote by  $\Omega_{R_+,R_0}$ , the so-called big cell of  $H_{R_0}$ . We claim that  $\Omega_{R_+,R_0}$  is schematically dense in  $H_{R_0}$ . Indeed, each fibre  $H_{R_0,s}$ ,  $s \in S$ , is an integral scheme in which  $\Omega_{R_+,R_0} \cap H_{R_0,s}$  is open and nonempty, hence dense, so that the reference  $[{\rm EGA, IV}_3, 11.10.10]$  proves our claim  $^{(1)}$  $^{(1)}$  $^{(1)}$ .

Coming back to the situation at hand, in view of [XXVI, 1.14] we can assume that  $P = H_{R_1}$  and  $Q = H_{R_2}$  for parabolic subsets  $R_1$  and  $R_2$  of  $R$ . Since according to  $[BT, Cor. 4.5]$ , the group K has geometrically connected fibers, it is shown in [XXII, 5.4.5] that

<span id="page-16-6"></span>
$$
(3.4.2) \t\t K = H_{R_1 \cap R_2},
$$

and that K is S–smooth and of type (R). Since  $R_1 \cap R_2$  is a closed subset of  $R$ , it follows from the definition of groups of type  $(RC)$  that  $K$  is such a

<span id="page-16-5"></span> $(1)$ This argument is taken from the web version of [XXII, 5.6.7, N.D.E. (38)]

group. Note that our arguments so far show the implication [\(iii\)](#page-15-5)  $\implies$  [\(i\)](#page-15-0) of [3.3](#page-15-6) (as well as  $(iii) \implies (ii)'$  $(iii) \implies (ii)'$  $(iii) \implies (ii)'$  $(iii) \implies (ii)'$ , which is however not needed).

We denote by  $R_1^s = R_1 \cap -R_1$  the symmetric part of  $R_1$  and let  $R_1^a =$  $R_1 \setminus R_1^s$  so that  $U_{R_1^a} = \text{rad}^u(P)$ , [XXVI, 1.12]. Thus, using the notation established so far, the map  $f$  of  $(a)$  is defined on

(3.4.3) 
$$
\text{rad}^u(P) \rtimes K = U_{R_1^a} \rtimes H_{R_1 \cap R_2}.
$$

We put

<span id="page-17-0"></span>
$$
R' = (R_1 \cap R_2) \cup R_1^a = (R_1^s \cap R_2) \cup R_1^a.
$$

One easily shows, see [\[BT,](#page-62-12) 4.4], that  $R'$  is a closed subset of R satisfying  $R' \cup (-R') = R$ , i.e., R' is a parabolic subset of R. Therefore  $P' = H_{R'}$  is a parabolic subgroup of  $G$ ,  $[XXVI, 1.4]$ . Keeping in mind  $(3.4.3)$ , we now claim that

f factorizes through an S-homomorphism

<span id="page-17-1"></span>(3.4.4) 
$$
f': U_{R_1^a} \rtimes H_{R_1 \cap R_2} \to P', \quad (u, h) \mapsto u h.
$$

In other words, we contend that the induced map  $f_{\sharp}: U_{R_1^a} \rtimes H_{R_1 \cap R_2} \to G/P'$ is the trivial map. Since  $R_1^a \subset R'$ , this is equivalent to  $\hat{f}_b: H_{R_1 \cap R_2} \to G/P'$ being the trivial map. This is the case when we restrict  $f_{\flat}$  to the big open cell  $\Omega_{R_+,R_1\cap R_2}$  of  $H_{R_1\cap R_2}$  which is a schematically dense in  $H_{R_1\cap R_2}$ . Hence, by [\[St,](#page-64-0) Tag 01RH], it follows that  $f_{\flat}$  and therefore also  $f_{\sharp}$  is trivial, finishing the proof of  $(3.4.4)$ . Note that  $(3.4.4)$  is part of the claim  $(a)$  (surjectivity will be established later). Next we show the first part of  $(b)$ , namely

$$
(3.4.5) \t\t\t f' \t\t is \ smooth.
$$

It follows from [\(3.3\)](#page-17-0) and the definition of  $U_{R_1^a}$ ,  $H_{R_1 \cap R_2}$  and  $P'$  that  $\text{Lie}(f')$ : Lie $(U_{R_1^a} \rtimes H_{R_1 \cap R_2}) \to \text{Lie}(P')$  admits a section (as  $\mathcal{O}_S$ -module map), so is in particular surjective. The reference  $[EGA, IV_4, 17.11.1(d)]$  then shows that  $f'$  is smooth along the unit section, so is smooth everywhere. We have thus proven the first part of [\(b\).](#page-16-3)

It follows, see e.g. [\[EGA,](#page-63-3) IV<sub>4</sub>, 17.5.1], that  $\text{Ker}(f')$  is smooth and hence in particular flat, so that the quotient  $(U_{R_1^a} \rtimes H_{R_1 \cap R_2})/Ker(f')$  is representable by an  $S$ -group scheme  $H'$  which is locally of finite presentation and equipped with a monomorphism  $h' : H' \to P'$ , [XVI, 2.3]. We consider the commutative diagram



The quotient map  $U_{R_1^a} \rtimes H_{R_1 \cap R_2} \to H'$  is smooth surjective and  $f'$  is smooth, hence  $h' : H' \to P'$  is smooth in view of [\[EGA,](#page-63-3) IV<sub>4</sub>, 17.7.7] (or see [\[St,](#page-64-0) Tag 02K5]). According to [\[EGA,](#page-63-3) IV<sub>4</sub>, 17.9.1], the smooth monomorphism  $h'$  is an open immersion. Since  $P'$  has smooth connected fibers,  $h'$  is surjective and we conclude that  $h'$  is an isomorphism. This then proves  $(a)$  in full.

To finish the proof of [\(b\),](#page-16-3) it remains to show  $(3.4.1)$ , i.e., rad<sup>u</sup> $(K)$  = rad<sup>u</sup>( $P'$ ) ∩ K. In view of [\(3.4.2\)](#page-16-6) and the definition of  $P'$ , this boils down to the equality

$$
(R_1 \cap R_2)^a = R'^a \cap R_1 \cap R_2,
$$

which is a special case of Lemma [3.5.](#page-18-0)

Once [\(b\)](#page-16-3) established, the implication [\(iii\)](#page-15-5)  $\implies$  [\(iv\)](#page-15-1) of [3.3](#page-15-6) follows: we apply  $[XXVI, 2.11]$  to the parabolic subgroup  $P'$  and its subgroup K of type (RC) (recall  $3.3(iii) \implies 3.3(ii)$  $3.3(iii) \implies 3.3(ii)$  $3.3(iii) \implies 3.3(ii)$  $3.3(iii) \implies 3.3(ii)$ , and get that K admits a Levi subgroup, hence contains Zariski-locally a maximal torus of G by [XIV, 3.20].

[\(c\)](#page-16-1) By [\(b\),](#page-16-3) the parabolic subgroup scheme  $P'$  and its subgroup K of type (RC) satisfy the assumptions of Lemma [2.7,](#page-9-7) so that [\(c\)](#page-16-1) follows by applying that lemma.  $\Box$ 

The notation of the following Lemma [3.5](#page-18-0) is the same as that in the proof of [3.4,](#page-16-0) except that we consider subsets of an arbitrary free  $\mathbb{Z}-$ module M: For a subset  $N \subset M$  we put  $N^s = N \cap (-N)$  and  $N^a = N \setminus N^s = N \setminus (-N)$ .

<span id="page-18-0"></span>3.5. Lemma. Let A and B be subsets of a free  $\mathbb{Z}-$ module M. We consider  $C = (A \cap B) \cup A^a = (A^s \cap B) \cup A^a$ . Then

<span id="page-18-1"></span>
$$
C^a \cap A \cap B = (A \cap B)^a.
$$

Proof. We will first establish several auxiliary statements, starting with

$$
(3.5.1)\qquad (A^s \cap B)^a = A^s \cap B^a
$$

Indeed, since  $A^s = -A^s$  we have  $x \in (A^s \cap B)^a \iff x \in A^s \cap B$  and  $-x \notin (A^s \cap B) \iff x \in A^s \cap B \text{ and } -x \notin B \iff x \in A^s \cap B^a$ . Next, we claim

.

<span id="page-18-3"></span>(3.5.2) 
$$
C^{a} = (A^{s} \cap B)^{a} \cup A^{a} = (A^{s} \cap B^{a}) \cup A^{a}.
$$

Because of [\(3.5.1\)](#page-18-1) we only need to prove the first equality. By definition of  $C^a$ , we have

<span id="page-18-2"></span>(3.5.3) 
$$
x \in C^{a} \iff x \in (A^{s} \cap B) \cup A^{a}
$$

$$
and -x \notin A^{s} \cap B \text{ and } -x \notin A^{a}.
$$

This easily implies the inclusion  $A^a \subset C^a$ : if  $x \in A^a$ , then  $x \in A$  and  $-x \notin A = A^s \cup A^a$ , in particular  $-x \notin A^s \cap B$  and  $-x \notin A^a$ . We also see that  $(A^s \cap B)^a = A^s \cap B^a \subset C^a$  since  $x \in A^s \cap B^a \iff x \in A^s \cap B$  and  $-x \notin B$ . Moreover  $-x \in A^s$ , so that  $-x \notin A^a$  also holds. We have now proved  $(A^s \cap B)^a \cup A^a \subset C^a$ . For the proof of the other inclusion, consider  $x \in A^s \cap B$ , but  $-x \notin A^s \cap B$  and  $-x \notin A^a$ , cf. [\(3.5.3\)](#page-18-2). Since  $A^s = \pm A^s$  we get  $x \notin B$ , thus  $x \in A^s \cap B^a = A^s \cap B^a$  by [\(3.5.1\)](#page-18-1). This finishes the proof of [\(3.5.2\)](#page-18-3). Next we observe

<span id="page-18-4"></span>(3.5.4) 
$$
(A \cap B)^a = (A^a \cap B) \cup (A \cap B^a)
$$

since  $x \in (A \cap B)^a \iff x \in A \cap B$ , but  $-x \notin A$  or  $-x \notin B \iff x \in A^a \cap B$ or  $x \in A \cap B^a$ . Finally, by [\(3.5.2\)](#page-18-3) and [\(3.5.4\)](#page-18-4),

$$
C^{a} \cap A \cap B = ((A^{s} \cap B^{a}) \cup A^{a}) \cap (A \cap B)
$$
  
= ((A^{s} \cap B^{a}) \cup A^{a}) \cap B = (A^{s} \cap B^{a}) \cup (A^{a} \cap B)  
= (A \cap B^{a}) \cup (A^{a} \cap B) = (A \cap B)^{a}.

With Propositions [3.2](#page-14-0) and [3.4](#page-16-0) in place, we can now also extend [XXVI, 5.4] to LG-rings.

<span id="page-19-0"></span>3.6. Corollary. Let R be an  $LG\text{-ring}$ , let G be a reductive R-group scheme, and let P, Q be two parabolic subgroups of G. Then there exists  $q \in G(R)$ such that  ${}^{g}P \cap Q$  is a parabolic subgroup of G.

Proof. Our proof follows the proof of [XXVI, 5.4]. We include it for the convenience of the reader.

Let  $P'$  be an opposite parabolic R-subgroup of G. Proposition [3.2,](#page-14-0) applied to the triple  $(P', P, Q)$  provides an R-parabolic R-subgroup  $P'_1$  of G of same type as  $P'$  such that  $P'_1$  and  $P$  (resp.  $P'_1$  and  $Q$ ) are in transversal position, hence in particular in standard position. By Proposition [3.4](#page-16-0)[\(c\),](#page-16-1) there exists a maximal torus T of G such that  $T \subset P'_1 \cap Q$ . Let  $P_1$  be the opposite parabolic  $R$ -subgroup of  $P'_1$  related to T, [XXVI, 4.3.3]. It follows that  $P_1 \cap Q$  is an R–parabolic subgroup [XXVI, 4.4.5]. Since P and  $P_1$ are both opposite to  $P'_1$ , they have same type, so that  $P_1 = {}^{g}P$  for some  $g \in G(R)$ , according to Theorem [3.1](#page-12-1)[\(a\).](#page-12-2) Thus  ${}^gP \cap Q$  is an R-parabolic subgroup of G as desired.

Having established Corollary [3.6](#page-19-0) for LG-rings, we also get [XXVI, 5.5 and 5.7] for LG-rings replacing semilocal rings in loc. cit..

3.7. Corollary ( $[XXVI, 5.5(i)]$  for semilocal rings). Let G be a reductive group scheme over an LG-ring, and let P and Q be two parabolic subgroups of G such that  $\mathbf{t}(P) \subset \mathbf{t}(Q)$ . Then there exists  $g \in G(S)$  such that  ${}^g P \subset Q$ .

<span id="page-19-1"></span>3.8. Corollary ( $[XXVI, 5.7(i)]$  for semilocal rings). Let G be a reductive group scheme over an LG-ring.

(a) Let  $t, t' \in \mathrm{Of}(\mathrm{Dyn}(G))(S)$ . If there exist parabolic subgroups of type t and t', then there also exists a parabolic subgroup of type  $t \cap t'$ .

<span id="page-19-2"></span>(b) There exists a smallest element  $\mathbf{t}_{\min}$  in the set of types  $\mathbf{t}(P)$ , P any parabolic subgroup of G.

<span id="page-19-3"></span>3.9. The Tits index. Let G be a semisimple group scheme over a  $con$ nected LG-ring. In view of  $3.8(b)$  $3.8(b)$ , we can define the Tits index of G as  $(Dyn(G), t_{min})$ , following [\[Ti\]](#page-64-2) for fields and the generalization to the semilocal case [\[PS,](#page-63-9) p. 202-203]. Moreover, as the diligent reader will verify, all the facts used in [\[PS\]](#page-63-9) for semilocal rings hold in fact for LG-rings. In particular, [\[PS,](#page-63-9) Thm. 3] extends to connected LG-rings: The Tits index of G is one of those listed in Tits' original table [\[Ti,](#page-64-2) Table II], reproduced in [\[PS,](#page-63-9) App.].

#### <span id="page-20-0"></span>4. Minimal parabolic subgroups, maximal split subtori

The topic of this section is minimal parabolic subgroups, their Levi subgroups and maximal split tori in reductive group schemes over a connected scheme S. We investigate the relations between these subjects in Proposi-tion [4.3](#page-21-0) and in Lemma [4.6](#page-23-1) for arbitrary connected S. In case  $S = \text{Spec}(R)$ for  $R$  a connected LG-ring and  $G$  a reductive  $R$ -group scheme, we show in Theorem [4.7](#page-23-0) that the group  $G(R)$  acts transitively on the minimal parabolic subgroups and on the maximal split tori of G. This allows us to define the anisotropic kernel of G in [4.9.](#page-24-1)

For perspective we note here that Appendix [B](#page-44-0) contains results on parabolic and Levi subgroups based on the dynamic method.

We start by reviewing the concepts used in Proposition [4.3.](#page-21-0)

<span id="page-20-2"></span>4.1. Isotropic and irreducible reductive groups. Let  $G$  be a reductive group scheme over an arbitrary scheme S.

Generalizing the well-known concept of an (an)isotropic reductive group over a field  $[BT, 4.23]$  or over a connected semilocal scheme  $[XXVI, 6.13]$ , we call G isotropic if it admits a subgroup isomorphic to  $\mathbb{G}_{m,S}$ ; otherwise, G is said to be anisotropic, [\[Gi1,](#page-62-9) 7.1.1]. Because of [B.3,](#page-45-0) these concepts coincides with those defined in  $[PS, §5]$  for semisimple group schemes over a connected semilocal scheme.

Following [\[Gi1,](#page-62-9) 3.5.1], we say that  $G$  is *reducible* (as reductive group scheme) if G admits an everywhere proper parabolic subgroup P, i.e.,  $P_{\overline{s}} \subsetneq$  $G_{\bar{s}}$  for all  $s \in S$ , and P admits a Levi subgroup; otherwise, G is said to be *irreducible.* The two notions are related by  $|Gi1, Thm. 7.3.1(2)|$ :

<span id="page-20-1"></span>(a) A reductive  $S$ -group scheme G is isotropic if and only if G is reducible or the central torus  $rad(G)$  is isotropic. In particular, a semisimple S–group scheme is isotropic if and only if it is reducible.

If S is affine, the notion of reducibility of  $G$  is equivalent to the existence of an everywhere proper parabolic subgroup [XXVI, 2.3], so the definition here agrees with the terminology of [\[GP,](#page-62-14) 3.2].

We will generalize the concepts of (an)isotropic and (ir)reducible reductive group schemes in  $B.2$ , and the criterion  $(a)$  in  $B.3$ .

4.2. Maximal split subtori, minimal parabolics, and faithful representations. A *maximal split subtorus* of  $G$  is a split subtorus of  $G$  which is maximal among all split subtori of  $G$ . By  $[XXIV, 2.11]$ , such a torus always exists if S is a scheme with trivial Picard group, e.g.,  $S$  is an LGscheme, and G has constant type. Existence is also assured in the setting of Proposition [4.3.](#page-21-0)

A parabolic subgroup of a reductive  $S$ -group G is a minimal parabolic subgroup if for all parabolic subgroups Q of G with  $Q \subset P$  we have  $Q = P$ , [XXVI, 5.6].

We use the term *faithful linear representation* of an  $S$ –group scheme G in the sense of [\[Gi2\]](#page-62-10), i.e., it is a group monomorphism  $G \to \mathbf{GL}(E)$  where E is a locally free  $\mathcal{O}_S$ -module of finite rank.

Now that we have explained all the concepts used in the following Proposition [4.3,](#page-21-0) we can finally state it. In fact, Proposition [4.3](#page-21-0) is a refinement of [\[Gi1,](#page-62-9) Prop. 7.4.1], which proves the implications [\(4.3.1\)](#page-21-1).

<span id="page-21-0"></span>4.3. Proposition. Let S be a connected scheme, G a reductive S-group scheme, and  $T_0$  a split subtorus of G. Moreover, let P be a parabolic subgroup of G for which  $C_G(T_0) := \text{Cent}_G(T_0)$  is a Levi subgroup of P; such a parabolic subgroup exists by [XXVI, 6.2]. We consider the following assertions:

- <span id="page-21-3"></span><span id="page-21-2"></span>(i)  $T_0$  is a maximal split S-subtorus of G;
- <span id="page-21-4"></span>(ii)  $T_0$  is a maximal split S-subtorus of  $C_G(T_0)$ ;
- <span id="page-21-5"></span>(iii) The reductive S-group  $C_G(T_0)/T_0$  is anisotropic;
- <span id="page-21-6"></span>(iv) The reductive S-group  $C_G(T_0)/T_0$  is irreducible;
- <span id="page-21-7"></span>(v) The reductive S-group  $C_G(T_0)$  is irreducible;
- <span id="page-21-1"></span>(vi)  $P$  is a minimal parabolic subgroup of  $G$ .

We then have the implications

$$
(4.3.1) \t(i) \Leftrightarrow (ii) \Leftrightarrow (iii) \implies (iv) \Leftrightarrow (v) \Leftrightarrow (vi)
$$

Furthermore, if G admits a faithful linear representation, then  $(ii) \iff (iii)$  $(ii) \iff (iii)$  $(ii) \iff (iii)$ .

*Proof.* It suffices to prove the implication [\(ii\)](#page-21-3)  $\implies$  [\(iii\)](#page-21-4) under the assumption that G admits a faithful linear representation. Let  $T_0$  be a maximal split S-subtorus of  $C<sub>G</sub>(T<sub>0</sub>)$ . We argue by contradiction and therefore assume that  $C_G(T_0)/T_0$  contains a subtorus  $\mathbb{G}_{m,S}$ . Denoting by  $E \subset C_G(T_0)$  the preimage of this subtorus  $\mathbb{G}_{m,S}$  under the quotient map, we get an exact sequence of S–group schemes

<span id="page-21-9"></span>
$$
(4.3.2) \t 1 \longrightarrow T_0 \longrightarrow E \longrightarrow \mathbb{G}_{m,S} \longrightarrow 1.
$$

By  $[XVII, 7.1.1], E$  is of multiplicative type. Clearly, it is also of finite type and it has a faithful linear representation. Therefore, by [\[Gi2,](#page-62-10) 3.3], the group scheme  $E$  is isotrivial, i.e., it is split by a finite étale cover, which we can assume to be a (connected) Galois cover  $S' \to S$ , whose Galois group we denote by  $\Gamma$  <sup>[\(2\)](#page-21-8)</sup>. According to the dictionary [X, 1.1], the extension [\(4.3.2\)](#page-21-9) corresponds to an extension

$$
0 \longrightarrow \mathbb{Z} \longrightarrow \widehat{E}(S') \longrightarrow \mathbb{Z}^{r_0} \longrightarrow 0
$$

<span id="page-21-8"></span><sup>(2)</sup> We use here: Given a finite étale cover  $T \rightarrow S$  of a connected scheme S, there exists a connected Galois cover  $S' \to S$  dominating  $T \to S$ . Proof: Since finite étale covers form a Galois category with respect to the base change functor induced by any geometric point of S [\[St,](#page-64-0) Tag 0BNB], the axioms of a Galois category [\[St,](#page-64-0) Tag 0BMZ] imply that  $T \to S$ is a product of finite étale connected covers. The existence of a Galois cover as claimed then follows from [\[St,](#page-64-0) Tag 0BN2], see the discussion in [\[St,](#page-64-0) Tag 03SF].

of  $\Gamma$ -modules, where  $\mathbb{Z}^{r_0} = \text{Hom}_{S'-gr}(E, \mathbb{G}_{m,S'})$ . The  $\Gamma$ -module  $\mathbb Z$  is the trivial module. Hence  $H^1(\Gamma, \mathbb{Z}) = \text{Hom}_{\text{Groups}}(\Gamma, \mathbb{Z}) = 0$  because  $\Gamma$  is finite. Thus, this extension splits, so that E is a split S–subtorus of  $C_G(T_0)$ . This contradicts the maximality of  $T_0$ .

4.4. **Remarks.** (a) The implication [\(ii\)](#page-21-3)  $\implies$  [\(iii\)](#page-21-4) is false without an additional assumption. Indeed, according to [X, 1.6], there exists a nonsplit rank 2 torus T over a singular projective connected complex curve S such that T admits a split subtorus  $T_0 \cong \mathbb{G}_m$  and an S–exact sequence  $1 \to T_0 \to T \to \mathbb{G}_m \to 1$ . Thus,  $T_0$  is a maximal split subtorus of the reductive S-group  $T = C_T(T_0)$ , yet  $C_T(T_0)/T_0$  is isotropic. We note that T is locally split in the sense of [A.1.](#page-38-1)

(b) The implication [\(iv\)](#page-21-5)  $\implies$  [\(iii\)](#page-21-4) is already false for  $G = \mathbb{G}_m = T$  and the obvious split subtorus  $T_0 = \{1\}$ . One may then be tempted to conjec-ture that [\(iv\)](#page-21-5)  $\implies$  [\(iii\)](#page-21-4) holds at least in the semisimple case. But this is false too: indeed, let  $G = SL_3$  and let  $T_0$  be the image of  $\mathbb{G}_{m,S}$  under the diagonal homomorphism  $t \mapsto diag(t, t^2, t^{-3})$ . The centralizer  $C_G(T_0)$  is the standard diagonal maximal torus of G of rank 2. Hence  $C_G(T_0)/T_0 \cong \mathbb{G}_{m,S}$ is irreducible, but also isotropic.

(c) The reader may have noticed that our proof of [\(ii\)](#page-21-3)  $\implies$  [\(iii\)](#page-21-4) in [4.3](#page-21-0) only uses that  $C<sub>G</sub>(T<sub>0</sub>)$  has a faithful linear representation. This is however equivalent to our assumption on G: if  $C<sub>G</sub>(T<sub>0</sub>)$  has a faithful linear representation, then so does rad(G), the unique maximal torus of the centre  $\mathcal{Z}(G)$ of  $G$ , and therefore also  $G$  by [\[Gi2,](#page-62-10) Thm. 4.1].

<span id="page-22-0"></span>4.5. The dynamic description of parabolic and Levi subgroups. Let S be a scheme and let H be a reductive  $S$ –group scheme. We recall the "dynamic method" in the sense of [\[Co1,](#page-62-8) 4.1, 5.2].

We will refer to a homomorphism  $\lambda: \mathbb{G}_{m,S} \to H$  of S–groups as a cocharacter (or a 1–parameter subgroup  $[Co1, Def. 4.1.6]$  $[Co1, Def. 4.1.6]$ ). Any cocharacter gives rise to a conjugation action of  $\mathbb{G}_m$  on H, and so for a fixed  $h \in H(S)$  to an orbit map orb<sub>h</sub>:  $\mathbb{G}_m \to H$ , which assigns to an S–scheme T and  $t \in \mathbb{G}_m(T)$ the element  $\lambda(t)h|_{T}\lambda(t)^{-1} \in H(T)$ . If orb<sub>h</sub> extends to a morphism  $\mathbb{G}_a \to H$ , i.e., an element of  $H(\mathbb{G}_a)$ , such a morphism is unique (since H is separated) and we will abbreviate this by " $\lambda(t)h\lambda(t)^{-1} \in H(\mathbb{G}_a)$ " (the abbreviation " $\lim_{t\to 0} \lambda(t)h\lambda(t)^{-1}$  exists" is used in [\[Co1,](#page-62-8) §4.1]). One then defines an S–functor  $P_H(\lambda)$  which assigns to an S–scheme T the group

$$
\underline{\mathbf{P}}_H(\lambda)(T) = \left\{ h \in H(T) : \lambda(t) \, h_{\mathbb{G}_{m,T}} \, \lambda(t)^{-1} \in H(\mathbb{G}_{a,T}) \right\}
$$

It is known that  $P_H(\lambda)$  is represented by a parabolic subgroup scheme of H, denoted  $P_H(\lambda)$ . The centralizer  $Cent_H(\lambda) = C_H(\lambda)$  of  $\lambda$  is a Levi subgroup of  $P_H(\lambda)$ ; its unipotent radical is given by those elements for which the limit is  $1$  [\[Co1,](#page-62-8) 4.1.7].

If S is a connected scheme, and  $(P, L)$  is a pair consisting of a parabolic subgroup of G and a Levi subgroup L of P, then there exists a cocharacter  $\lambda$ 

#### 24 P. GILLE AND E. NEHER

such that  $(P, L) = (P_G(\lambda), C_G(\lambda)),$  [\[Gi1,](#page-62-9) Thm. 7.3.1(1)]. We will refine this result in [B.1.](#page-44-1) We point out that even over a connected  $S$  there may exist parabolic subgroups which do not contain Levi subgroups and hence are not amenable to a dynamic description, [\[Co1,](#page-62-8) Ex. 5.4.9]. However, this cannot happen if S is affine  $[XXVI, 2.3]$  (or  $[Co1, 5.4.8]$ ). Thus, over an affine base every parabolic subgroup P has the form  $P = P_H(\lambda)$  for a suitable cocharacter  $\lambda$  ([\[Gi1,](#page-62-9) 7.3.2]).

In [4.6,](#page-23-1) we will use the dynamic method to describe minimal parabolic subgroups and their Levi subgroups over a connected base.

<span id="page-23-2"></span><span id="page-23-1"></span>4.6. Lemma (Minimal parabolic subgroups and their Levi subgroups). Let G be a reductive group scheme over a connected scheme S.

- <span id="page-23-3"></span>(a) G admits a minimal parabolic subgroup.
- (b) Let L be a Levi subgroup of a minimal parabolic R-subgroup  $P$  of  $G$ , and let  $T_0$  be a maximal split S-subtorus of the torus rad(L); it is unique by Lemma [A.6](#page-41-0)[\(b\)](#page-41-1). Then  $L = \text{Cent}_G(T_0)$ .
- <span id="page-23-4"></span>(c) We assume furthermore that G admits a faithful linear representation, and let  $T_0$  be a maximal split subtorus of G. Then  $Cent_G(T_0)$ is a Levi subgroup of a minimal parabolic subgroup of G.

*Proof.* Since S is connected, the type of G is constant  $[XXII, 2.8]$ , and so is the type of any parabolic subgroup of G.

[\(a\)](#page-23-2) The length of a strictly increasing chain of parabolic subgroups of G is bounded by the relative dimension of  $G$ . So it is obvious that a minimal  $S$ -parabolic subgroup of  $G$  exists.

[\(b\)](#page-23-3) According to  $[G_1, Thm. 7.3.1]$ , there exists an  $S$ -group homomorphism  $\lambda: \mathbb{G}_{m,S} \to G$  such that  $P = P_G(\lambda)$  and  $L = C_G(\lambda)$  (notation of [4.5\)](#page-22-0). We can suppose that  $\lambda$  is non-trivial. Hence, by [A.3,](#page-40-0)  $\mathbb{G}_{m,S}/\text{Ker}(\lambda) \cong \mathbb{G}_{m,S}$ . Without loss of generality, we can then assume that  $\lambda$  is a monomorphism. In particular,  $\lambda$  factors through the unique maximal split S-subtorus  $T_0$  of rad(L). We have  $L \subset C_G(T_0) \subset C_G(\lambda) = L$  so that  $L = C_G(T_0)$  as desired. [\(c\)](#page-23-4) This follows from Proposition [4.3,](#page-21-0) [\(i\)](#page-21-2)  $\implies$  [\(vi\).](#page-21-7)

<span id="page-23-0"></span>4.7. **Theorem** (Transitivity of  $G(R)$  on minimal parabolic subgroups and maximal split tori). Let R be a connected  $LG$ -ring, and let G be a reductive R-group scheme.

- <span id="page-23-5"></span>(a) The group  $G(R)$  acts transitively on the minimal R-parabolic subgroups of G.
- <span id="page-23-6"></span>(b) We assume furthermore that G admits a faithful linear representation (e.g. G is semisimple). Then the group  $G(R)$  acts transitively on the maximal split R–subtori of G.

*Proof.* [\(a\)](#page-23-5) Let P and Q be minimal parabolic subgroups of G. Corollary [3.6](#page-19-0) shows that there exists  $g \in G(R)$  such that  ${}^g P \cap Q$  is an R-parabolic subgroup of G. By minimality, we have  ${}^{g}P \cap Q = Q = {}^{g}P$ .

[\(b\)](#page-23-6) Let  $T_1$  and  $T_2$  be maximal split R-subtori of G. We want to show that  $T_1$  and  $T_2$  are  $G(R)$ –conjugate. According to Lemma [4.6](#page-23-1)[\(c\),](#page-23-4) the centralizers  $L_i = C_G(T_i)$ ,  $i = 1, 2$ , are Levi subgroups of minimal parabolic subgroups  $P_i$  of G. Applying [\(a\),](#page-23-5) reduces to the case  $P_1 = P_2$ . Since Levi subgroups of a fixed parabolic subgroup are  $G(R)$ –conjugate by [XXVI, 1.8], we can further assume that  $L_1 = L_2$ . According to Proposition [4.3,](#page-21-0) [\(i\)](#page-21-2)  $\implies$  [\(ii\),](#page-21-3)  $T_1$  and  $T_2$  are maximal R-split tori of  $L_1 = L_2$  and a fortiori a maximal split tori of the torus  $rad(L_1)$ . Since we have uniqueness for maximal split subtori of a given subtorus (Lemma [A.6](#page-41-0)[\(b\)\)](#page-41-1), we conclude that  $T_1 = T_2$ .  $\Box$ 

Theorem [4.7](#page-23-0)[\(b\)](#page-23-6) is proven in [XXVI, 6.16] for a reductive  $R$ -group scheme over a connected semilocal R.

4.8. Remarks (Witt-Tits decomposition). Let R be a connected LG-ring, and let G be a reductive  $R$ -group scheme. Then the conclusions of [\[Gi1,](#page-62-9) 4.3.1, 4.4.3, 5.2.1] regarding the Witt-Tits decomposition of  $H^1(\text{Spec}(R), G)$ hold.

Indeed, the reader will easily check that the proofs of the quoted references, stated in  $[G1]$  for semilocal rings, only require that R is a connected ring for which  $G(R)$  acts transitively on the set of minimal parabolic subgroups of G. But this is  $4.7(a)$  $4.7(a)$ .

<span id="page-24-1"></span>4.9. The anisotropic kernel. Let R be a connected LG-ring and let G be a reductive R–group scheme.

By  $4.6(a)$  $4.6(a)$ , G admits minimal parabolic subgroups and by  $4.7(a)$  $4.7(a)$ , they are all conjugate. Let us fix one of them, say  $P_{\text{min}}$ . Levi subgroups of  $P_{\text{min}}$  exist by [XXVI, 2.3], and they are all conjugate by [XXVI, 1.8]. Let us fix one of them, say  $L_{\text{min}}$ . The derived subgroup  $\mathscr{D}(L_{\text{min}})$  of  $L_{\text{min}}$  is a semisimple  $R$ –group scheme. It follows from  $[XXVI, 1.20]$  and  $[Gi1, Lem. 3.2.1(2)]$  that  $\mathscr{D}(L_{\text{min}})$  is irreducible, hence anisotropic by [4.1](#page-20-2)[\(a\).](#page-20-1) Summarizing, up to isomorphism, there exists up to isomorphism a unique semisimple anisotropic  $R$ –subgroup of  $G$ , defined as

<span id="page-24-0"></span>
$$
(4.9.1) \t Gan = \mathscr{D}(Lmin)
$$

and called the *anisotropic kernel of G*. The terminology follows Tits [\[Ti\]](#page-64-2) for the case of a field and Petrov-Stavrova [\[PS\]](#page-63-9) for the case of semilocal rings.

<span id="page-24-5"></span>4.10. **Example.** Let G be a reductive group scheme over a connected LGring, and suppose  $T_0$  is a split subtorus of G for which  $Cent_G(T_0)/T_0$  is anisotropic. Then

- <span id="page-24-3"></span><span id="page-24-2"></span>(i)  $T_0$  is a maximal split subtorus of  $G$ ;
- (ii) there exist parabolic subgroups of G admitting  $Cent_G(T_0)$  as Levi subgroup, and any such parabolic subgroup is minimal;
- <span id="page-24-4"></span>(iii)  $\mathscr{D}(\text{Cent}_G(T_0))$  is an anisotropic kernel of  $\tilde{G}$ .

Indeed, Proposition [4.3](#page-21-0) implies [\(i\)](#page-24-2) and [\(ii\),](#page-24-3) while [\(iii\)](#page-24-4) follows from the definition [\(4.9.1\)](#page-24-0) of an anisotropic kernel. We will specialize this example in [6.13](#page-37-0) to determine the anisotropic kernel of  $GL_1(A)$ , A an Azumaya R-algebra.

In the remainder of this section we investigate the interplay between arbitrary parabolic subgroups of a reductive group over a connected LG-ring, Levi subgroups and split subtori.

<span id="page-25-0"></span>4.11. Proposition (XXVI, 6.8) for R semilocal connected). Let R be a connected LG-ring, let G be a reductive R–group scheme, let  $P \subset G$  be a parabolic subgroup scheme and let  $L$  be a Levi subgroup of  $P$ , which exists by [XXVI, 2.3]. We put  $Q = \text{rad}(L)$  and let  $Q_0$  the maximal split subtorus of  $Q$ , which exists by  $A.6(d)$  $A.6(d)$ . Then

$$
(4.11.1) \t\t\t L = \text{Cent}_G(Q_0).
$$

The proof of [4.11](#page-25-0) for a connected LG-scheme is the same as the proof of [XXVI, 6.8], up to replacing the reference [XXVI, 3.20] by [2.6.](#page-9-0) We leave the details to the reader.

Recall  $[XXVI, 6.2]$ : If S is any scheme, and Q a split subtorus of a reductive S–group scheme, then there exists a parabolic subgroup of G for which  $Cent_G(Q)$  is a Levi subgroup.

Combining this result with [4.11,](#page-25-0) we get the following corollary, which is [XXVI, 6.9] in the semilocal case.

4.12. Corollary. Let S be a connected  $LG\text{-}scheme, G$  a reductive  $S\text{-}group,$ and T a critical subtorus in the sense that  $T = \text{rad}(\text{Cent}_G(T))$ . We denote the maximal split subtorus of  $T$  by  $T_0$ . Then the following are equivalent:

- (i) Cent<sub>G</sub>(T) is a Levi subgroup of a parabolic subgroup of G;
- (ii)  $\text{Cent}_G(T) = \text{Cent}_G(T_0);$
- (iii)  $\text{Lie}(G)^T = \text{Lie}(G)^{T_0}.$

More corollaries of Proposition [4.11](#page-25-0) are established in [XXVI, 6.10–6.12] for reductive groups over a connected semilocal scheme. Their proofs carry over without change to groups over a connected LG-scheme. We state these corollaries below for the reader's convenience.

4.13. Corollary (XXVI, 6.10) for R semilocal connected). Let S be a connected LG-scheme, and let G be a reductive S–group. Then the following conditions are equivalent for a subgroup scheme  $L$  of  $G$ :

- (i) There exists a parabolic subgroup with Levi subgroup L.
- (ii) There exists a split subtorus of  $G$  whose centralizer is  $L$ .
- (iii) There exists a cocharacter  $\mathbb{G}_{m,S} \to G$  whose centralizer is L.

4.14. Corollary (XXVI, 6.11) for R semilocal connected). Let S be a connected  $LG$ -scheme, and let G be a reductive  $S$ -group. Given a torus  $T$ , we denote by  $T_0$  its unique maximal split subtorus, whose existence is guaranteed by  $A.6(d)$  $A.6(d)$ . Then the maps

 $L \mapsto \text{rad}(L)_0, \qquad Q \mapsto \text{Cent}_G(Q)$ 

are order-reversing inverse bijections between the set of Levi subgroups of parabolic subgroups of G and the set of split subtori Q of G satisfying  $Q =$ rad( $\text{Cent}_G(Q)$ )<sub>0</sub>.

## 5. Cancellation Theorems

<span id="page-26-0"></span>In this section we prove several cancellation theorems that are classically known over fields, in some cases even over semilocal rings, but that we establish here over LG-rings: cancellation of modules and Azumaya algebras in tensor products [5.2,](#page-26-2) cancellation of hermitian forms in [5.4](#page-29-0) and cancellation of quadratic forms in [5.5.](#page-29-1) All of these cancellation results are applications of the cohomological injectivity result of Theorem  $3.1(b)$  $3.1(b)$ . We start with presenting the principle [5.1](#page-26-1) that describes the basis of cancellation.

<span id="page-26-1"></span>5.1. Cancellation Principle. Let  $R$  be an LG-ring. We consider the diagram  $(5.1.1)$  of R-group schemes and R-group homomorphisms

<span id="page-26-3"></span>(5.1.1) 
$$
G \xrightarrow{\Delta} L \xrightarrow{\text{inc}_L} P \xrightarrow{\text{inc}_P} H
$$

where G and H are reductive  $R$ -group schemes, P is a parabolic subgroup of H with Levi subgroup L, the map  $\Delta$  is split as group homomorphism, and where  $\text{inc}_{L}$  and  $\text{inc}_{P}$  are the natural inclusions. Then the canonical map

$$
(5.1.2) \qquad \alpha: H^1(R, G) \longrightarrow H^1(R, H),
$$

induced by the composition of the maps in  $(5.1.1)$ , is injective.

*Proof.* The sequence [5.1.1](#page-26-3) of group homomorphisms induces maps in cohomology:

.

<span id="page-26-4"></span>(5.1.3)  
\n
$$
H^{1}(R, G) \xrightarrow{\Delta^{*}} H^{1}(R, L)
$$
\n
$$
\cong \begin{cases}\n\operatorname{inc}_{L}^{*} \\
\operatorname{inc}_{L}^{*} \\
H^{1}(R, H) \xleftarrow{\operatorname{inc}_{P}^{*}} H^{1}(R, P)\n\end{cases}
$$

In the diagram [\(5.1.3\)](#page-26-4) the map  $\Delta^*$  is injective since  $\Delta$  is a split group homomorphism, the map  $inc<sub>L</sub><sup>*</sup>$  is a bijection by [XXVI, 2.3] and the map inc<sup>\*</sup><sub>*p*</sub> is injective by Theorem [3.1](#page-12-1)[\(b\).](#page-13-0)  $\Box$ 

As a first application of the Cancellation Principle [5.1,](#page-26-1) we prove a cancellation result for finite projective modules and for Azumaya algebras over LG-rings. Cancellation of Azumaya algebras was proven for semilocal rings by Knus [\[Kn1,](#page-63-10) Thm. 3.3] and again by Ojanguren-Sridharan in [\[OS,](#page-63-11) Cor. 1], see also  $[Kn2, III, 5.2.3(b)].$ 

<span id="page-26-2"></span>5.2. Application (Cancellation of modules and Azumaya algebras). Let R be an LG-ring.

<span id="page-26-5"></span>(a) Let  $M_1$  and  $M_2$  be finite projective R-modules, and let N be a faithfully projective R–module. Then

<span id="page-26-6"></span>
$$
(5.2.1) \t\t M_1 \otimes_R N \cong M_2 \otimes_R N \implies M_1 \cong M_2
$$

(isomorphism of R–modules). In particular, for any  $n \in \mathbb{N}_+$ ,

<span id="page-27-1"></span>
$$
(5.2.2) \t\t M_1^{(n)} \cong M_2^{(n)} \implies M_1 \cong M_2.
$$

<span id="page-27-3"></span><span id="page-27-0"></span>(b) Let  $A$ ,  $B$  and  $C$  be  $Azumaya R-algebras$ . Then

(5.2.3) 
$$
A \otimes_R C \cong B \otimes_R C \implies A \cong B.
$$

In particular, for  $n \in \mathbb{N}_+$ ,

(5.2.4) 
$$
M_n(A) \cong M_n(B) \implies A \cong B.
$$

Proof. [\(a\)](#page-26-5) We apply the Principle [5.1](#page-26-1) with the following choices:

- <span id="page-27-2"></span>(i)  $G = GL_d$  is the R-group scheme representing the R-functor which associates with  $T \in R$ -alg the group  $GL_d(T)$  of invertible matrices in  $M_d(T)$ ;
- (ii)  $H = GL_{nd};$
- (iii) L the subgroup scheme of  $GL_{nd}$ , representing the R-functor which associates with  $T \in R$ -alg the subgroup

$$
L(T) = \{ \text{diag}(x_1, \ldots, x_n) : x_i \in \text{GL}_d(T) \}
$$

of  $\mathbf{GL}_{nd}(T);$ 

- (iv) P is the product of L and upper triangular matrices in  $GL_{nd}$ ;
- (v)  $\Delta$  is the diagonal group homomorphism

$$
\Delta: \mathbf{GL}_d \to L, \quad x \mapsto \mathrm{diag}(x, \dots, x)
$$

which is split by the projection onto the first factor

$$
\mathrm{pr}_1\colon L\to \mathbf{GL}_d,\quad \mathrm{diag}(x_1,\ldots,x_n)\mapsto x_1.
$$

We first prove  $(5.2.2)$ . In view of the rank decomposition of finite pro-jective modules and of [1.1\(](#page-3-3)[a\)](#page-3-4), it is no harm to assume that  $M_1$  and hence also  $M_2$  have constant rank, say they are both of rank d. Thus, they represent cohomology classes  $[M_1]$  and  $[M_2]$  in  $H^1(R,\mathbf{GL}_d)$ . The assumption in  $(5.2.2)$  is that  $\alpha([M_1]) = \alpha([M_2]) \in H^1(R, GL_{nd}).$  Therefore  $[M_1] = [M_2]$ by injectivity of  $\alpha$ .

We now prove  $(5.2.1)$  as a consequence of  $(5.2.2)$ . Let Q be an R-module such that  $N \otimes_R Q$  is free of finite rank, say  $N \otimes_R Q \cong R^n$ , see [1.5.](#page-6-3) Observe that  $M_1 \otimes_R N \otimes_R Q \cong M_1 \otimes_R R^{(n)} \cong M_1^{(n)}$  $1^{(n)}$ . Therefore, the assumption in  $(5.2.1)$  implies  $M_1^{(n)}$  $\chi_1^{(n)} \cong M_2^{(n)}$  $\chi_2^{(n)}$ , which forces  $M_1 \cong M_2$  by  $(5.2.2)$ .

[\(b\)](#page-27-1) The proof of [\(b\)](#page-27-1) is a straightforward modification of the proof of [\(a\)](#page-26-5). Quotioning the sequence

$$
\mathbf{GL}_d \xrightarrow{\Delta} L \xrightarrow{\mathrm{inc}_L} P \xrightarrow{\mathrm{inc}_P} \mathbf{GL}_{nd}
$$

of [\(a\)](#page-26-5) by the central  $\mathbb{G}_m$ , we obtain the sequence of R-group schemes

$$
\mathbf{PGL}_d \stackrel{\bar{\Delta}}{\longrightarrow} \overline{L} = L/\mathbb{G}_m \xrightarrow{\text{inc}_{\bar{L}}} \overline{P} = P/\mathbb{G}_m \xrightarrow{\text{inc}_{\bar{P}}} \mathbf{PGL}_{nd}
$$

which satisfies the assumptions of the Principle [5.1,](#page-26-1) cf. [\[Gi1,](#page-62-9) 3.2.1]. Thus the canonical map  $\bar{\alpha}$ :  $H^1(R, \text{PGL}_d) \to H^1(R, \text{PGL}_{nd})$  is injective. The cohomology set  $\mathrm{H}^1(R,\mathbf{PGL}_d)$  represents the isomorphism classes of Azumaya

algebras of constant degree d. The map  $\bar{\alpha}$  sends the class of an Azumaya R-algebra D of degree d to the class  $[M_n(D)] \in H^1(R, \mathbf{PGL}_{nd})$ .

For the proof of  $(5.2.4)$  we can assume that A and B have constant rank, say rank d. Then [\(5.2.4\)](#page-27-2) follows from injectivity of the map  $\bar{\alpha}$  above. To prove [\(5.2.3\)](#page-27-3), we can suppose that C has constant rank n. Then  $C \otimes_R C^{\text{op}} \cong$  $\text{End}_R(C)$ , where on the right hand side we view C as projective R-module of rank n. Since  $C \cong R^n$  by [1.6](#page-7-1)[\(b\)](#page-7-2), we get  $C \otimes_R C^{\text{op}} \cong M_n(R)$ . Now [\(5.2.3\)](#page-27-3) follows:

$$
A \otimes_R C \cong B \otimes_R C \implies A \otimes_R C \otimes_R C^{\text{op}} \cong B \otimes_R C \otimes_R C^{\text{op}}
$$
  

$$
\implies M_n(A) \cong M_n(B) \implies A \cong B.
$$

Based on [5.2\(](#page-26-2)[b\)](#page-27-1), we will be able to say more about Azumaya algebras over LG-rings in §[6.](#page-31-0)

5.3. Remark (Additive Cancellation of modules). The additive version of [\(5.2.1\)](#page-26-6) is known to be true too, even in a more general setting than modules over LG-rings [\[EG,](#page-62-2) Thm. 2.5]: Let R be an LG-ring, let  $S \in R$ -alg be a direct limit of finite R-algebras, and let  $M_1$ ,  $M_2$  and N be S-modules. If N is finitely generated, then

<span id="page-28-0"></span>(5.3.1) 
$$
M_1 \oplus N \cong M_2 \oplus N \quad \Longrightarrow \quad M_1 \cong M_2.
$$

The proof of loc. cit. establishes that  $\text{End}_S(N)$  has 1 in the stable range of  $\text{End}_{R}(N)$  and then applies a result of Evans [\[Ev,](#page-62-15) Thm. 2] to show [\(5.3.1\)](#page-28-0). Additive cancellation goes back to the classical results of Bass [\[Bas,](#page-61-2) 6.6, 9.3]; in the semilocal case it is proven in  $[Kn2, VI; (1.3.3)].$ 

The point of this remark is that the special case  $S = R$  and  $M_1, M_2$ and N finite projective R–modules of  $(5.3.1)$  can easily be proven using the Cancellation Principle  $5.1$ . Indeed, there exists a finite projective  $R$ -module  $N'$  such that  $M \oplus N' \cong R^n$  for some  $n \in \mathbb{N}_+$ . Therefore  $M_1 \oplus R^n \cong M_2 \oplus R^n$ . By induction on *n*, it suffices to prove the case  $n = 1$ ,

(5.3.2) 
$$
M_1 \oplus R \cong M_2 \oplus R \implies M_1 \oplus M_2.
$$

It follows that  $\operatorname{rank}_RM_1 = \operatorname{rank}_RM_2$ . Applying the standard rank decomposition of finite projective modules and  $1.1(a)$  $1.1(a)$ , we can assume that  $M_1$  and  $M_2$  have constant rank r.

Let  $L$  be the Levi subgroup of the parabolic subgroup  $P$  of the reductive R–group scheme  $H = GL_{r+1}$ , given with obvious meaning by

<span id="page-28-2"></span><span id="page-28-1"></span>
$$
L = \begin{pmatrix} \mathbf{GL}_r & 0 \\ 0 & \mathbb{G}_m \end{pmatrix}, \qquad P = \begin{pmatrix} \mathbf{GL}_r & * \\ 0 & \mathbb{G}_m \end{pmatrix}
$$

.

We can apply the Cancellation Principle [5.1](#page-26-1) with  $\Delta: G = GL_r \to L$  the obvious embedding and  $L \subset P \subset GL_{r+1} = H$  as above. Hence, we get an injective map

(5.3.3) 
$$
H^1(R,\mathbf{GL}_r) \hookrightarrow H^1(R,\mathbf{GL}_{r+1}),
$$

which sends the isomorphism class of a finite projective  $R$ –module  $Q$  of rank r to the isomorphism class of  $Q \oplus R$ . Thus, injectivity of [\(5.3.3\)](#page-28-1) yields a proof of  $(5.3.2)$ .

The final two results of this section concern cancellation of hermitian and quadratic forms. For the convenience of the reader, Appendix [C](#page-47-0) reviews the concepts needed for [5.4](#page-29-0) and [5.5](#page-29-1) and their proofs.

<span id="page-29-0"></span>5.4. Application (Cancellation of hermitian forms). Let R be an  $LG$ -ring, and let  $S/R$  be a quadratic étale extension with standard involution  $\sigma$ . Then cancellation holds for  $(S, \sigma)$ –hermitian spaces: if  $h_1$ ,  $h_2$  and  $h_3$  are hermitian spaces such that  $h_1 \perp h_3$  and  $h_2 \perp h_3$  are isometric, then already  $h_1$  and  $h_2$ are isometric:

<span id="page-29-2"></span>(5.4.1) 
$$
h_1 \perp h_3 \cong h_2 \perp h_3 \implies h_1 \cong h_2.
$$

*Proof.* The form  $h_3 \perp (-h_3)$  is hyperbolic [\[Kn2,](#page-63-12) I; 3.7.3]. Since the assumption implies that  $h_1 \perp (h_3 \perp -h_3) \cong h_2 \perp (h_3 \perp -h_3)$ , it is no harm to assume that  $h_3$  is hyperbolic, say  $h_3 = \mathbb{H}(U_3)$ . We can then apply the Cancellation Principle [5.1](#page-26-1) with the following choices:

- (i)  $G = U(h_1)$  and  $H = U(h_1 \perp h_3)$  are the unitary R-group schemes of the regular hermitian forms  $h_1$  and  $h_1 \perp h_3$  respectively, which are reductive  $R$ –group schemes by  $C.6(d)$  $C.6(d)$ .
- (ii)  $L = \mathbf{U}(h_1) \times_R \mathfrak{R}_{S/R}(\mathbf{GL}(U_3));$
- (iii) P is the stabilizer of  $U_3$  in H, which by [C.7](#page-52-1) is a parabolic subgroup of  $H$  with Levi subgroup  $L$ .
- (iv)  $\Delta: U(h_1) \to U(h_1) \times_R U(h_3)$  is the canonical embedding.

Thus the natural map  $H^1(R, U(h_1)) \rightarrow H^1(R, U(h_1 \perp h_3))$  is injective. Taking the bijection of  $C.6(e)$  $C.6(e)$  as identification, it sends the isometry class of a hermitian space  $h'$  to the isometry class of  $h' \perp h_3$ , which implies  $(5.4.1).$  $(5.4.1).$ 

**Remark.** For a semilocal ring R, a more general result than  $5.4$  is proven in  $[Kn2, VI; (5.7.5)]$  — it concerns cancellation of unitary (= hermitian) spaces over unitary rings.

<span id="page-29-1"></span>5.5. Application (Cancellation of quadratic forms). Let R be an  $LG\text{-}ring$ , let  $q_1$  and  $q_2$  be nonsingular quadratic forms and let  $q_3$  be a regular quadratic form. Then

$$
(5.5.1) \t q_1 \perp q_3 \cong q_2 \perp q_3 \implies q_1 \cong q_2.
$$

*Proof.* Applying the rank decomposition  $C.8(g)$  $C.8(g)$  of the quadratic forms  $q_i$ ,  $i = 1, 2, 3$ , together with [1.1\(](#page-3-3)[a\)](#page-3-4), we can assume that the  $q_i$  have constant rank. We can further suppose that all three ranks are positive, since otherwise the claim is obvious. Moreover, because the quadratic form  $q_3 \perp -q_3$ is hyperbolic by  $[Bae, I, (4.7)],$  we can assume that  $q_3$  is hyperbolic. In particular,  $q_3$  is regular, so that  $q := q_1 \perp q_3$  is nonsingular by [\(C.8.3\)](#page-54-1).

At this point, the reader might be inclined to apply the principle [5.1](#page-26-1) to  $\mathbf{O}(q_1)$  and  $\mathbf{O}(q)$ , which by [C.15](#page-59-0) describe the isometry classes of nonsingular quadratic forms. This is however not possible because  $O(\cdot)$  is not reductive in general. A way out of this problem is to use  $SO(q_1)$  and  $SO(q)$  instead, which are indeed reductive group schemes by [C.16,](#page-60-0) and then investigate the relation between  $H^1(R, SO(\cdot))$  and  $H^1(R, O(\cdot))$ .

Following this approach, we apply [5.1](#page-26-1) with  $G = SO(q_1)$  and  $H = SO(q)$ , using the parabolic and Levi subgroups of  $H$  exhibited in [C.17.](#page-61-4) Hence

$$
\alpha \colon \mathrm{H}^1(R, \mathbf{SO}(q_1)) \to \mathrm{H}^1(R, \mathbf{SO}(q))
$$

is injective. To link  $H^1(R, \mathbf{SO}(\cdot))$  and  $H^1(R, \mathbf{O}(\cdot))$ , we use that there exist an  $R$ –group scheme  $A$ , an  $R$ –group homomorphism  $D$  and an exact sequence of  $R$ –group schemes

(5.5.2) 
$$
1 \longrightarrow SO(q) \xrightarrow{i} O(q) \xrightarrow{D} A \longrightarrow 1
$$

where  $i$  is the canonical inclusion and where

<span id="page-30-0"></span>
$$
A = \begin{cases} \mu_{2,R}, & \text{if } q \text{ has odd rank,} \\ \mathbb{Z}/2\mathbb{Z}_R, & \text{if } q \text{ has even rank.} \end{cases}
$$

Indeed, this follows from  $C.16(c)$  $C.16(c)$  in the odd rank case and from  $(C.16.5)$  in the even rank case.

Part of the long exact cohomology sequence associated with [\(5.5.2\)](#page-30-0) is the sequence of pointed sets

$$
\mathrm{O}(q) \xrightarrow{D(R)} A(R) \xrightarrow{\delta} \mathrm{H}^1(R, \mathbf{SO}(q)) \xrightarrow{i^*} \mathrm{H}^1(R, \mathbf{O}(q)),
$$

in which  $D(R)$  is surjective by [C.13\(](#page-58-0)[b\)](#page-58-1), implying that  $i^*$  has trivial kernel. Since the parities of the ranks of  $q_1$  and q agree, we have the analogous exact sequence  $(5.5.2)$  for  $q_1$  replacing q, with the same A. Moreover, we get a commutative diagram of pointed sets

$$
H^{1}(R, \mathbf{SO}(q_{1})) \xrightarrow{i_{1}^{*}} H^{1}(R, \mathbf{O}(q_{1})) \xrightarrow{D^{*}} H^{1}(R, A)
$$
\n
$$
\downarrow^{\alpha} \qquad \qquad \downarrow^{\beta} \qquad \qquad \parallel^{\beta} \qquad \qquad \parallel
$$
\n
$$
H^{1}(R, \mathbf{SO}(q)) \xrightarrow{i^{*}} H^{1}(R, \mathbf{O}(q)) \xrightarrow{D^{*}} H^{1}(R, A)
$$

where  $i_1^*$  is associated to the inclusion  $i_1: SO(q_1) \to O(q_1)$  and where  $\beta$ maps the isometry class of  $q_2$  to that of  $q_2 \perp q_3$  (recall [C.15\)](#page-59-0). Our claim then is that  $\beta$  has trivial kernel. But this follows from a simple diagram chase, using that  $\alpha$  is injective and that  $i^*$  has trivial kernel.  $\Box$ 

Remarks. For regular quadratic forms over fields of characteristic not 2, the result [5.5](#page-29-1) goes back to Witt, and is therefore referred to as Witt cancellation, even in more general settings.

Witt cancellation is proven in [\[EKM,](#page-62-16) Thm. 8.4] for quadratic forms over arbitrary fields, in [\[Bae,](#page-61-3) III, Cor. 4.3] for regular quadratic forms  $q_1$ ,  $q_2$  and

#### 32 P. GILLE AND E. NEHER

 $q_3$  over semilocal rings, and in [\[Di,](#page-62-6) II, 6.4] for LG-rings, again for regular forms, following Baeza's approach which in turn goes back to Knebusch.

<span id="page-31-0"></span>Witt cancellation is not true if all  $q_i$  are merely nonsingular, see [\[EKM,](#page-62-16) p. 49] for a counterexample.

## 6. Azumaya algebras over LG-rings

In this section we consider Azumaya algebras over rings. One of its goals is Corollary [6.9](#page-35-0) on the Brauer decomposition of Azumaya algebras over connected LG-rings. This is a consequence of Theorem [6.7](#page-34-0) which says that indecomposable finite projective modules of Azumaya algebras over a connected LG-ring are isomorphic. Another highlight of this section is Proposition [6.2](#page-32-0) proving that an Azumaya algebra A of constant degree over an LG-ring admits a splitting ring which is a maximal étale subalgebra of  $A$ .

We start this section with Corollary [6.1,](#page-31-1) which is an application of cancellation of tensor products of Azumaya algebras and establishes "Hilbert 90". In this corollary,  $Br(R)$  denotes the Brauer group of a ring R as defined in [\[Fo,](#page-62-0) §7.3], which is the Brauer-Azumaya group of [\[C-TS,](#page-62-17) 3.1.3].

<span id="page-31-2"></span><span id="page-31-1"></span>6.1. Corollary. Let R be an  $LG\text{-}ring$ , and let A be an Azumaya R–algebra.

(a) If B is another Azumaya R-algebra with  $\deg A = \deg B$ , then A and B are isomorphic as R–algebras if and only if their Brauer classes agree,

<span id="page-31-3"></span>(6.1.1) 
$$
A \cong B \iff [A] = [B] \in \text{Br}(R).
$$

(b) ("Hilbert 90")  $H^1(R,\mathbf{GL}_1(A)) = \{1\}.$ 

*Proof.* [\(a\)](#page-31-2) It suffices to prove that  $[A] = [B] \implies A \cong B$ . The equality  $[A] = [B]$  means that there exist faithfully projective R-modules P and Q such that  $A \otimes_R \text{End}_R(P) \cong B \otimes_R \text{End}_R(Q)$  as R–algebras. Using the rank decomposition of  $P$  and  $Q$ , we can easily reduce to the case that both  $P$  and Q have constant rank. Hence, by [1.6](#page-7-1)[\(b\)](#page-7-2), there exist  $m, n \in \mathbb{N}_+$  such that  $M_m(A) \cong M_n(B)$  (isomorphism of R–algebras). Comparing degrees, we get  $m = n$ , hence  $A \cong B$  by  $(5.2.4)$ .

[\(b\)](#page-31-3) Let  $\textbf{PGL}(A)$  be the automorphism group scheme of A, [\[CF,](#page-62-1) 2.4.4.2]. The standard exact sequence of  $Spec(R)$ –group schemes,

$$
1 \to \mathbb{G}_m \to \mathbf{GL}_1(A) \to \mathbf{PGL}(A) \to 1
$$

gives rise to a long exact sequence of pointed cohomology sets, part of which is

$$
H^1(R, \mathbb{G}_m) \to H^1(R, GL_1(A)) \xrightarrow{\alpha} H^1(R, \mathbf{PGL}(A)) \xrightarrow{\delta} H^2(R, \mathbb{G}_m).
$$

On the left end,  $H^1(R, \mathbb{G}_m) = Pic(R) = 0$  by  $(1.1.1)$ , which implies that  $\alpha$ has trivial kernel. The set  $H^1(R, \mathbf{PGL}(A))$  classifies twisted forms of A. If B is such a twisted form, it has the same degree as  $A$ . By  $[\text{Gir}, 4.2.12\text{..}(\text{iii})]$ , the map  $\delta$  sends the isomorphism class of B to  $[A] - [B] \in Br(R) \subset H^2(R, \mathbb{G}_m)$ . Hence, part [\(a\)](#page-31-2) implies that  $\delta$  has trivial kernel, i.e.,  $\alpha$  has trivial image. The standard torsion argument then proves  $(b)$ . <span id="page-32-1"></span><span id="page-32-0"></span>6.2. Proposition. Assume that R is LG-ring. Let A be an Azumaya  $R$ algebra of constant degree d.

- (i) A admits a maximal étale  $R$ -subalgebra S of finite degree d and  $A \otimes_R S \cong M_d(S)$ .
- <span id="page-32-2"></span>(ii) If R is connected, then A admits a connected finite étale  $R$ –subalgebra  $S'$  such that  $A \otimes_R S' \cong M_d(S')$ .

*Proof.* [\(i\)](#page-32-1) According to Theorem [2.6,](#page-9-0) the reductive R-group scheme  $GL_1(A)$ admits a maximal torus T. By  $[Gi2, Prop. 3.2], T occurs from an embedding$  $S \hookrightarrow A$  of a maximal finite étale algebra S of degree d and  $T = GL_1(S)$ . The canonical S–module A is finite projective of constant rank d and  $A \otimes_R B$  $S \cong \text{End}_{S}(A)$  $S \cong \text{End}_{S}(A)$  $S \cong \text{End}_{S}(A)$  as Azumaya S–algebras ([\[Fo,](#page-62-0) 7.4.2], [\[KO,](#page-63-13) III, Prop. 6.1]). In particular,  $A \otimes_R S$  is a neutral Azumaya S–algebras of degree d. Next, S is an LG-ring by Example [1.2](#page-4-0)[\(b\)](#page-4-4). Since  $A \otimes_R S$  and  $M_d(S)$  are both neutral Azumaya  $S$ -algebras of the same degree, Corollary [6.1](#page-31-1)[\(a\)](#page-31-2) enables us to conclude that  $A \otimes_R S \cong M_d(S)$ .

[\(ii\)](#page-32-2) Because R is connected, we have a decomposition  $S = S_1 \times \cdots \times S_c$ with  $S_i$  finite étale connected. Since  $A \otimes_R S \cong M_d(S)$ , we obtain that  $A \otimes_R S' \cong M_d(S')$  for  $S' = S_1$ .

<span id="page-32-4"></span>6.3. Indecomposable finite projective modules. Let R be an arbitrary ring and let A be an Azumaya R-algebra. Recall that an  $A$ -module M is called *decomposable* if there exists a family  $(M_i)_{i\in I}$  of submodules  $M_i$  of M such that  $M = \bigoplus_{i \in I} M_i$  and every  $M_i \neq 0$ . Otherwise, it is called *indecomposable.* Hence  $M = \{0\}$  is decomposable by taking  $I = \emptyset$ , and an A–module M is indecomposable if and only if  $M \neq 0$  and idemp  $(\text{End}_A(M)) = \{0, 1\}.$ Here idemp $(B) = \{b \in B : b^2 = b\}$  is the set of idempotents of an R-algebra B.

Following [\[DI,](#page-62-18) V, p. 131], we define an equivalence relation  $\sim$  on the set of indecomposable finite projective left A–modules by  $P \sim Q \iff$  there exists an invertible R–module E such that  $P \cong Q \otimes_R E$  as (left) A–modules. Obviously, if  $Pic(R) = \{0\}$ , then "equivalence" reduces to "isomorphism". We will use the following result.

<span id="page-32-5"></span>6.4. **Theorem** ([\[DI,](#page-62-18) V; Thm. 1.1]). Let A be an Azumaya algebra over a connected ring R. Then

<span id="page-32-3"></span>
$$
P \mapsto \operatorname{End}_A(P)^{\rm op}
$$

induces a bijection between the set of equivalence classes of indecomposable finite projective A–modules and the set of isomorphism classes of Azumaya R–algebras B satisfying

(6.4.1) 
$$
A \sim_{\text{Br}} B \quad and \quad \text{idemp}(B) = \{0, 1\},
$$

where  $\sim_{\text{Br}}$  indicates Brauer equivalence.

The inverse map assigns to B satisfying  $(6.4.1)$  the module P constructed as follows: By definition of  $A \sim_{\text{Br}} B$  there exists a faithfully projective  $R$ –module  $P$  satisfying

<span id="page-33-0"></span>(6.4.2) 
$$
A \otimes_R B^{\text{op}} \cong \text{End}_R(P).
$$

We use the canonical R–algebra homomorphism  $A \to A \otimes_R B^{op}$  and  $(6.4.2)$ to endow  $P$  with an  $A$ –module structure. It is finite projective as  $A$ –module by [\[Fo,](#page-62-0) 4.4.1]. Viewing B as a subalgebra of  $\text{End}_R(P)$  the double centralizer theorem inmplies

$$
(6.4.3) \t\t B \cong \text{End}_A(P).
$$

Hence idemp  $(End_A(P)) = \{0,1\}$  which says that P is an indecomposable A–module by the characterization of indecomposability mentioned in [6.3.](#page-32-4)

6.5. Corollary. Let R be a connected ring. Then every Brauer class  $\alpha \in$  $Br(R)$  has a representative B satisfying idemp $(B) = \{0, 1\}.$ 

*Proof.* By Theorem [6.4,](#page-32-5) it suffices to show that for every Azumaya  $R$ –algebra A there exists an indecomposable finite projective A–module. This can be seen by decomposing the finite "projective" left  $A$ –module  $_A A$ .

<span id="page-33-8"></span><span id="page-33-2"></span>6.6. Lemma. Let R be a connected ring and let A be an Azumaya  $R$ –algebra. Then the following conditions are equivalent:

- <span id="page-33-1"></span>(i) Any two indecomposable finite projective A–modules are equivalent.
- (ii) Up to isomorphism, the Brauer class of A contains a unique B satisfying idemp $(B) = \{0, 1\}.$

<span id="page-33-3"></span>If these conditions hold, then we have for  $B$  as in [\(ii\)](#page-33-1) that

- <span id="page-33-5"></span>(iii)  $deg(B) = ind([A]),$  in particular  $deg(B) | deg(A)$ .
- (iv) Up to equivalence, the left B-module  $_B B$  is the unique indecomposable finitely generated projective left B–module.
- <span id="page-33-7"></span>(v) In addition to [\(i\)](#page-33-2) and [\(ii\)](#page-33-1), suppose  $Pic(R) = \{0\}$ , and let M and N be finite projective R–modules satisfying  $\operatorname{rank}_R M \ge \operatorname{rank}_R N$ . Then N is a homomorphic image of M.

*Proof.* The equivalence [\(i\)](#page-33-2)  $\iff$  [\(ii\)](#page-33-1) is a special case of Theorem [6.4.](#page-32-5)

[\(iii\)](#page-33-3) We can write  $A$  as a finite sum of indecomposable finitely generated projective A–modules, say  $A = P_1 \oplus \cdots \oplus P_n$ . By [\(i\)](#page-33-2) we can assume that there exists an indecomposable finite projective A–module P such that  $P_i =$  $P \otimes_R E_i$  for some invertible R-module  $E_i$ . Thus,  $\text{rank}_R P_i = \text{rank}_R P$  and we get

<span id="page-33-4"></span> $(6.6.1)$  rank<sub>R</sub>  $A = n$  rank<sub>R</sub>  $P$  and rank<sub>R</sub>  $A \cdot \text{rank}_R B = (\text{rank}_R P)^2$ 

where the second equation in  $(6.6.1)$  follows from  $(6.4.2)$ . The two equations in  $(6.6.1)$  imply

<span id="page-33-6"></span>(6.6.2)  $\operatorname{rank}_R P = n \operatorname{rank}_R B, \operatorname{rank}_R A = n^2 \operatorname{rank}_R B.$ 

The second equation then forces  $\deg B | \deg(A)$ .

Let  $[A] = \alpha \in \text{Br}(R)$ . By definition  $\text{ind}(\alpha) = \text{gcd}\{\text{deg }A' : A' \sim_{\text{Br}} A\}.$ The equivalence [\(i\)](#page-33-2)  $\iff$  [\(ii\)](#page-33-1) shows that the condition (i) also holds for A'. Since  $A \sim_{\text{Br}} A' \implies B \sim_{\text{Br}} A'$ , it then follows that  $deg(B)| deg(A')$ , and therefore  $deg(B) = ind(\alpha)$ .

 $(iv)$  As observed in [\(iii\),](#page-33-3) the condition [\(i\)](#page-33-2) holds for B replacing A. Hence we get  $(6.6.2)$  for  $A = B$ , which forces  $n = 1$  and therefore [\(iv\).](#page-33-5)

[\(v\)](#page-33-7) Since  $Pic(R) = 0$ , "equivalence" in [\(i\)](#page-33-2) becomes "isomorphism". We fix an indecomposable finite projective A–module K, and then get  $M \cong K^a$ and  $N \cong K^b$  for positive integers a and b. The rank assumption on M and N implies that  $a \geq b$ . The claim then follows.

<span id="page-34-0"></span>6.7. **Theorem** ([\[DeM,](#page-62-19) Thm. 1] for R semilocal). Let R be a connected  $LG$ ring. Then the condition  $6.6(i)$  $6.6(i)$  holds with "equivalence" replaced by "iso-morphism". Hence the conditions [\(ii\)](#page-33-1)–[\(v\)](#page-33-7) of [6.6](#page-33-8) are satisfied too.

*Proof.* Let P and Q be indecomposable finite projective  $A$ -modules. Since R is connected,  $\text{rank}_{R} P$  and  $\text{rank}_{R} Q$  are constant, so that Corollary [1.6](#page-7-1) says that  $P$  and  $Q$  are free as  $R$ –modules. Without loss of generality we can therefore suppose that  $\operatorname{rank}_R P \ge \operatorname{rank}_R Q$ . We claim:

<span id="page-34-1"></span> $(6.7.1)$  Q is a homomorphic image of P.

Assuming [\(6.7.1\)](#page-34-1) for a moment, we can quickly finish the proof. Indeed, because Q is projective, any epimorphism  $P \to Q$  splits. But P is indecomposable. Therefore  $P \cong Q$ .

It remains to prove  $(6.7.1)$ . This can be done by observing that  $6.6(i)$  $6.6(i)$  and hence [6.6](#page-33-8)[\(v\)](#page-33-7) holds for the localization  $R_{\mathfrak{m}}$  in a maximal ideal  $\mathfrak{m}\triangleleft R$ , thanks to [\[DeM,](#page-62-19) Thm. 1]. Applying then [\[EG,](#page-62-2) Thm. 2.6(i)], proves the claim. A more direct proof of  $(6.7.1)$  goes as follows. Since P and Q are free of finite rank as R-modules,  $\text{Hom}_R(P,Q)$  is an affine space; it contains the open quasicompact subscheme  $U' = \{ \varphi \in \text{Hom}_R(P, Q) : \varphi \text{ is surjective } \}$  (note that the complement of  $U'$  is given by the vanishing of finitely many minors). The A–linear maps  $\text{Hom}_{A}(P,Q)$  is a subspace of  $\text{Hom}_{R}(P,Q)$ , hence is again an affine space. It contains  $U = \{ \varphi \in \text{Hom}_A(P,Q) : \varphi \text{ is surjective} \} =$  $Hom_A(P,Q) \cap U'$  as an open quasi-compact subscheme. If  $\mathfrak{m} \triangleleft R$  is any maximal ideal of R, it is well-known that  $6.6(i)$  $6.6(i)$  and hence  $6.6(v)$  $6.6(v)$  hold for R replaced by the field  $R/\mathfrak{m}$ , i.e.,  $U(R/\mathfrak{m}) \neq \emptyset$ . But then Proposition [1.4](#page-6-0)[\(a\)](#page-6-1) shows that  $U(R) \neq \emptyset$ , which is [\(6.7.1\)](#page-34-1).

6.8. The Wedderburn property. Condition [\(ii\)](#page-33-1) of [6.6](#page-33-8) and Theorem [6.7](#page-34-0) say that a connected LG-ring R has the Wedderburn property in the sense of [\[AW17\]](#page-61-1) or [\[Fo,](#page-62-0) §7.6], i.e., for every Azumaya R-algebra A over a connected ring  $R$  there is up to isomorphism a unique representative  $B$  in the Brauer class of A such that idemp(B) = {0,1} and  $B^{\rm op} \cong \text{End}_A(M)$  for some Aprogenerator module M. As far as we know, this was previously only known for connected semilocal rings [\[DeM,](#page-62-19) Cor. 1].

Examples of rings that do not have the Wedderburn property are given in [\[AW14,](#page-61-6) Ex. 1.5], [\[Bas,](#page-61-2) page 46] and [\[Ch\]](#page-62-20).

<span id="page-35-0"></span>6.9. Corollary (Brauer Decomposition for LG-rings). Let R be a connected LG-ring, let A be an Azumaya R-algebra with idemp $(A) = \{0, 1\}$ , and let  $\text{ind}([A]) = p_1^{m_1} \dots p_c^{m_c}$  be the prime factor decomposition of  $\text{ind}([A])$ . Then there exist Azumaya R–algebras  $B_1, \ldots, B_c$  satisfying

<span id="page-35-1"></span>(6.9.1) 
$$
\deg(B_i) = p_i^{m_i} \text{ and } \text{idemp}(B_i) = \{1, 0\}
$$

for  $i = 1, \ldots, c$ , as well as

<span id="page-35-2"></span>(6.9.2) 
$$
A \cong B_1 \otimes_R \cdots \otimes_R B_c.
$$

The conditions [\(6.9.1\)](#page-35-1) and [\(6.9.2\)](#page-35-2) determine the family  $(B_1, \ldots, B_c)$  up to isomorphism.

*Proof.* Because Br(R) is an abelian torsion group, we can write  $\alpha = [A] \in$  $Br(R)$  as  $\alpha = \alpha_1 + \cdots + \alpha_c$  where  $\alpha_i \in Br(R)$  has order  $p_i^{m_i}$  for  $1 \le i \le c$ . We now use Gabber's Theorem [\[Ga,](#page-62-21) II, Thm. 1] for each class  $\alpha_i$ . It says that there exist Azumaya R-algebras  $A_i$  satisfying  $\alpha_i = [A_i]$  and  $\deg(A_i) = p_i^{\ell_i}$ for some  $\ell_i \in \mathbb{N}_+$ . Next, by [6.7,](#page-34-0) we can apply [6.6.](#page-33-8) Thus, there exist Azumaya algebras  $B_i$  satisfying  $[B_i] = [A_i]$  and idemp $(B_i) = \{0, 1\}$  for  $i = 1, \ldots, c$ . Observe that  $deg(A) = ind(A)$  and  $deg(B_i) = ind(\alpha_i)$  by [6.6](#page-33-8)[\(iii\).](#page-33-3) We then get

$$
\deg(B_1 \otimes_R \cdots \otimes_R B_c) = \prod_{i=1}^c \deg(B_i)
$$

$$
= \prod_{i=1}^c \text{ind}(\alpha_i) \stackrel{(*)}{=} \text{ind}(\alpha) = \deg(A)
$$

where (∗) holds by [\[AW17,](#page-61-1) Thm. 3]. Similarly,

$$
[A] = \alpha = \alpha_1 + \dots + \alpha_c = [A_1] + \dots + [A_c]
$$

$$
= [B_1] + \dots + [B_c] = [B_1 \otimes_R \dots \otimes_R B_c],
$$

so that Corollary  $6.1(a)$  $6.1(a)$  yields the isomorphism  $(6.9.2)$ .

Unicity: Let  $(B'_1, \ldots, B'_c)$  be a family of Azumaya R-algebras satisfying [\(6.9.1\)](#page-35-1) and [\(6.9.2\)](#page-35-2). By [6.6](#page-33-8)[\(iii\)](#page-33-3) we then get  $ind([B'_i]) = deg(B'_i) = p_i^{m_i}$ , thus  $\text{per}([B_i]) = p_i^{\ell_i}$  for some  $\ell_i \leq m_i$ . It follows that  $[A] = [B'_1] + \cdots$  $[B'_{c}]$  is the decomposition of  $[A]$  into primary components. Hence  $[B'_{i}]$  =  $[B_i]$  for  $i = 1, \ldots, c$ . As  $\deg(B'_i) = p_i^{\tilde{m_i}} = \deg(B_i)$ , another application of Corollary [6.1\(](#page-31-1)[a\)](#page-31-2) yields  $B'_i \cong B_i$ .

The results in [6.4–](#page-32-5)[6.9](#page-35-0) show the importance of Azumaya algebras A satisfying idemp(A) =  $\{0, 1\}$ . While this is a purely algebraic condition, it is perhaps not surprising that these algebras also have a natural group theoretic characterization, which in fact holds beyond the case of connected LG-rings. We will prove this in [6.12,](#page-37-1) using the following Lemmas [6.10](#page-36-0) and [6.11](#page-36-1) which (in our view) are of independent interest.

For an Azumaya algebra A over a scheme S, the S–group scheme  $GL_1(\mathcal{A})$ is for example defined in [\[CF,](#page-62-1) 2.4.2.2].

<span id="page-36-2"></span><span id="page-36-0"></span>6.10. **Lemma.** Let  $A$  be an Azumaya algebra over a connected scheme  $S$ , and let e be an idempotent of  $A = A(S)$ .

(a) Then

<span id="page-36-5"></span>(6.10.1) 
$$
\lambda_e: \mathbb{G}_{m,S} \longrightarrow \mathbf{GL}_1(\mathcal{A}) =: G,
$$

$$
t(\in \mathbb{G}_{m,S}(T)) \longrightarrow te_T + t^{-1}(1_{\mathcal{A}(T)} - e_T)
$$

 $(T$  an  $S$ -scheme) is a cocharacter such that

<span id="page-36-4"></span>(6.10.2) 
$$
\lambda_e \text{ is central } \iff e \in \{0, 1\} \iff G = P_G(\lambda) = C_G(\lambda).
$$

<span id="page-36-3"></span>(b)  $e = 0 \iff e_T = 0$  for some S-scheme T; analogously for  $e = 1$ .

*Proof.* [\(a\)](#page-36-2) It is immediate that  $\lambda_e$  is a cocharacter, which is central if  $e \in$  $\{0,1\}$ . In general, let  $\mathcal{A}_{ij}$ ,  $i, j \in \{1,0\}$  be the Peirce spaces of A with respect to the orthogonal idempotents  $e_1 = e$  and  $e_0 = 1 - e$ . Thus  $A_{ij} = e_i A e_j$ . The centralizer of  $\lambda_e$  in A is  $C_{\mathcal{A}}(\lambda_e) = \mathcal{A}_{11} \oplus \mathcal{A}_{00}$ .

Assume now that  $\lambda_e$  is central. Then  $\mathcal{A} = C_{\mathcal{A}}(\lambda) = \mathcal{A}_{11} \times \mathcal{A}_{00}$ . It follows that  $\mathcal{A}_{11}$  and  $\mathcal{A}_{00}$  are ideals of  $\mathcal{A}$ . They induce ideals  $\mathcal{I}_1$  and  $\mathcal{I}_0$  of  $\mathcal{O}_S$  by  $\mathcal{I}_i = \mathcal{A}_{ii} \cap \mathcal{O}_S$ ,  $i = 1, 0$ , which satisfy  $\mathcal{A}_{ii} = \mathcal{I}_i \mathcal{A}$ . Since  $\mathcal{A} = \mathcal{A}_{11} \times \mathcal{A}_{00}$  we have  $\mathcal{O}_S = \mathcal{I}_1 \times \mathcal{I}_0$ . Because S is connected, either  $\mathcal{I}_i = \mathcal{O}_S$ , i.e.,  $e = 1$ , or  $\mathcal{I}_1 = 0$ , i.e.,  $e = 0$ .

Clearly,  $\lambda_e$  is central  $\iff G = C_G(\lambda)$ . Since  $C_G(\lambda) \subset P_G(\lambda) \subset G$  in general, this proves the second equivalence.

[\(b\)](#page-36-3) The parabolic subgroup  $P_G(\lambda_e)$  has constant type since S is connected [XXVI, 3.3]. Hence, if  $e_T = 0$ , then  $P_G(\lambda_e)(T) = G(T)$ . Because  $P_G(\lambda_e)$ has constant type, we get  $P_G(\lambda) = G$  and therefore  $e \in \{0,1\}$  by  $(6.10.2)$ . The assumption  $e = 1$  contradicts  $e_T = 0$ . Thus  $e = 0$ . The other direction is obvious.  $\Box$ 

<span id="page-36-1"></span>6.11. **Lemma.** Let A be an Azumaya algebra over a connected scheme  $S$ , and suppose  $\lambda: \mathbb{G}_{m,S} \to GL_1(\mathcal{A})$  is a non-central cocharacter. Then  $\{0,1\} \subsetneq$ idemp $(A(S))$ .

*Proof.* Let  $\mathcal{B} = \mathcal{A}^{\lambda}$  be the subalgebra of  $\mathcal{A}$  centralizing the image of  $\lambda$ , and let C be the centre of B. By [\[Gi2,](#page-62-10) Prop. 3.2] we know that C is a finite étale  $\mathcal{O}_S$ -algebra of positive rank. We will eventually show that  $\mathcal{C}(S)$  contains a nontrivial idempotent  $e$  defined by using the decomposition of  $C$  into its connected components. But first we need to eliminate the case that  $\mathcal C$  is connected.

By construction of C, the image of  $\lambda$  lies in C, and this also holds for the central cocharacter  $\Delta$ . Thus  $\lambda$ ,  $\Delta \in \text{Hom}_{S-\text{gr}}(\mathbb{G}_{m,S}, GL_1(\mathcal{C}))$ . To compare the two cocharacters, let  $C$  be the scheme associated with the finite étale  $\mathcal{O}_S$ -algebra C and let  $\mathfrak{R}_{C/S}(\cdot)$  be the Weil restriction. Then  $GL_1(\mathcal{C})$  =  $\Re_{C/S}(\mathbb{G}_m, C)$ ,  $(\mathbb{G}_{m,S})_C = \mathbb{G}_{m,C}$  and the fundamental identity of the Weil restriction  $\Re_{C/S}$  becomes (after a canonical identification)

$$
\mathrm{Hom}_{S-\mathrm{gr}}\left(\mathbb{G}_{m,S}, \mathfrak{R}_{C/S}(\mathbb{G}_{m,C})\right) = \mathrm{Hom}_{C-\mathrm{gr}}(\mathbb{G}_{m,C}, \mathbb{G}_{m,C}).
$$

Suppose that C is connected. Then  $\text{Hom}_{C-\text{gr}}(\mathbb{G}_{m,C}, \mathbb{G}_{m,C}) \cong \mathbb{Z}$  with basis  $\Delta$ . Therefore  $\lambda = \Delta^n$  for some  $n \in \mathbb{Z}$ , in particular  $\lambda$  is central, contradiction. Therefore  $C$ , equivalently  $C$ , is not connected.

Let  $\mathcal{C}_1 \times \cdots \times \mathcal{C}_m$  be the decomposition of C corresponding to the decomposition of C into its connected components, and let  $e_1, \ldots, e_m$  be the identity elements of the algebras  $C_i(S)$ . Then the  $e_i$  are non-zero idempotents of A since  $C_i(S) \neq 0$ , and  $e_1 \neq 1$  because  $m \geq 2$ .

We use the concepts of (ir)reducible and (an)isotropic group schemes as defined in [4.1.](#page-20-2) For an Azumaya algebra  $A$  over a scheme  $S$  the  $S$ –group scheme  $SL_1(\mathcal{A})$  is for example defined in [\[CF,](#page-62-1) 3.5.0.91]; it is semisimple by [\[CF,](#page-62-1) 3.5.0.92]. The S–group scheme  $\textbf{PGL}(\mathcal{A})$  is a semisimple S–group scheme by [\[CF,](#page-62-1) 3.0.5.82].

<span id="page-37-2"></span><span id="page-37-1"></span>6.12. Proposition. Let A be an Azumaya algebra over a connected scheme S. The following are equivalent:

- <span id="page-37-3"></span>(i)  $SL_1(\mathcal{A})$  is isotropic,
- <span id="page-37-6"></span>(ii)  $SL_1(\mathcal{A})$  is reducible,
- <span id="page-37-4"></span>(iii)  $GL_1(\mathcal{A})$  is reducible,
- <span id="page-37-5"></span>(iv)  $\textbf{PGL}(\mathcal{A})$  is reducible,
- <span id="page-37-8"></span>(v)  $\textbf{PGL}(\mathcal{A})$  is isotropic,
- (vi) idemp  $(\mathcal{A}(S)) \supsetneq \{0,1\}.$

*Proof.* By [4.1](#page-20-2)[\(a\),](#page-20-1) the equivalences [\(i\)](#page-37-2)  $\iff$  [\(ii\)](#page-37-3) and [\(iv\)](#page-37-4)  $\iff$  [\(v\)](#page-37-5) hold since  $SL_1(\mathcal{A})$  and  $PGL(\mathcal{A})$  are semisimple S-group schemes. The equivalences [\(ii\)](#page-37-3)  $\iff$  [\(iii\)](#page-37-6)  $\iff$  [\(iv\)](#page-37-4) are special cases of [\[Gi1,](#page-62-9) 3.5.3(b)<sup>[\(3\)](#page-37-7)</sup> and 3.2.1(2)].

If e is an idempotent as in [\(vi\),](#page-37-8) then  $\lambda_e$  is not central by [\(6.10.1\)](#page-36-5). Hence  $G = GL_1(\lambda_e) \neq C_G(\lambda_e)$  and a fortiori  $G \neq P_G(\lambda_e)$ , which proves [\(iii\).](#page-37-6)

Conversely, if  $(iii)$  holds, then G contains a proper parabolic subgroup P with a Levi subgroup L. By [\[Gi1,](#page-62-9) 7.3.1(1)] we have  $(P, L) = (P_G(\lambda), C_G(\lambda))$ for a non-central cocharacter  $\lambda$ . Thus [\(vi\)](#page-37-8) follows from Lemma [6.11.](#page-36-1)  $\Box$ 

<span id="page-37-0"></span>6.13. Corollary (Anisotropic kernel of  $GL_1(A)$ ). Let R be a connected LGring and let A be an Azumaya R-algebra. By  $6.4$  and  $6.7$  there exist unique  $n \in \mathbb{N}$  and an Azumaya algebra B with idemp $(B) = \{0, 1\}$  such that  $A =$  $M_n(B)$ . Then a minimal parabolic subgroup  $P_{\min}$  of  $G = GL_1(A)$  and a minimal Levi subgroup  $L_{\text{min}}$  are

$$
P_{\min} = \begin{pmatrix} \mathbf{GL}_1(B) & * & * \\ 0 & \ddots & * \\ 0 & 0 & \mathbf{GL}_1(B) \end{pmatrix}, \quad L_{\min} = \begin{pmatrix} \mathbf{GL}_1(B) & 0 & 0 \\ 0 & \ddots & 0 \\ 0 & 0 & \mathbf{GL}_1(B) \end{pmatrix}.
$$

<span id="page-37-7"></span> $(3)$ The reference refers to the web version of the article: https://hal.science/hal-01063601v2/document. The published version does not distinguish between  $H$  and  $H^{\rm ss} = H/\text{rad}(H).$ 

A maximal split torus  $T_0$  of G and an anisotropic kernel  $G_{\text{an}}$  of G are

$$
T_0 = \begin{pmatrix} \mathbb{G}_m & 0 & 0 \\ 0 & \ddots & 0 \\ 0 & 0 & \mathbb{G}_m \end{pmatrix}, \qquad G_{\text{an}} = \begin{pmatrix} \mathbf{SL}_1(B) & 0 & 0 \\ 0 & \ddots & 0 \\ 0 & 0 & \mathbf{SL}_1(B) \end{pmatrix}.
$$

*Proof.* It is clear that  $T_0$  as displayed above is a split torus whose centralizer in G has the form of the matrix group  $L_{\text{min}}$ . The quotient  $L_{\text{min}}/T_0$  can be identified with

$$
\begin{pmatrix}\n\mathbf{PGL}(B) & 0 & 0 \\
0 & \ddots & 0 \\
0 & 0 & \mathbf{PGL}(B)\n\end{pmatrix}
$$

which is anisotropic by [6.12.](#page-37-1) Example [4.10](#page-24-5) can therefore be applied and yields that the matrix groups displayed above are a minimal parabolic subgroup, a minimal Levi subgroup, a maximal split torus and an anisotropic kernel respectively.

# <span id="page-38-0"></span>Appendix A. Maximal split subtori of groups of multiplicative **TYPE**

In this appendix, we study maximal split and maximal locally split Ssubtori of groups of finite multiplicative type. We prove their existence in [A.6,](#page-41-0) and show in the Example [A.9\(](#page-42-0)b) that the two notions are different in general.

<span id="page-38-1"></span>A.1. Basic notions, review. We mainly use the terminology and notation of [\[SGA3,](#page-63-0) VIII, IX], as before abbreviated by [VIII] and [IX], see also [\[Co1,](#page-62-8) App. B and  $[Oe]$ . Throughout this appendix, S is an arbitrary scheme.

Given an abelian group M, one denotes by  $D_S(M) = Spec(\mathcal{O}_S[M])$ the abealin S–group scheme representing the functor of characters of the constant S–group  $M_S$ . An S–group scheme G is called *split*, equivalently diagonalizable, if G is isomorphic to an  $S$ -group  $D_S(M)$  for some abelian group  $M$ . A possible confusion with the notion of a split reductive  $S$ -group cannot occur, if the reader takes the context into account. A split S–torus is an S-group T isomorphic to  $D_S(\mathbb{Z}^r)$  for some  $r \in \mathbb{N}_+$ . We call an S-group scheme G locally split if there exists a family  $(S_i)_{i\in I}$  of open subschemes  $S_i$ of S such that  $S = \bigcup_{i \in I} S_i$  and every  $G|_{S_i} = G \times_S S_i$  is diagonalizable. If in this case all  $G|_{S_i}$  are split tori, we call T a locally split torus.

An  $S$ -group G has *multiplicative type* if G is locally diagonalizable for the fpqc topology, i.e., for every  $s \in S$  there exists an open affine neighbourhood U of s in G and a finite family  $(X_i)_{i\in I}$  of affine schemes together with flat morphisms  $f_i: X_i \to U$  such that  $U = \bigcup_i f_i(X_i)$  and every  $G|_{X_i}$  is diagonalizable. In this case, we call G a *torus*, if all  $G_{X_i}$  are split tori. A group of finite multiplicative type is a group of multiplicative type which is also of finite type.

Let G be a group of multiplicative type. For  $s \in S$  let  $G_s$  be the fibre of s. There exists an extension k of the residue field of s such that  $(G_s)_k$  is diagonalizable, say  $\cong$  D<sub>S</sub>( $M_s$ ) for some abelian group  $M_s$ . The isomorphism class of  $M_s$  is independent of the choice of k, and called the type of G at s, [IX; 1.4]. The function on the underlying topological space  $S_{\text{top}}$  of S, associating with  $s \in S$  the type of G in s, is locally constant. Assume now that G has finite multiplicative type. For every  $s \in S$  the type of G is a finitely generated abelian group  $M_s$ . Hence we get a well-defined, locally constant function rank $(G): S_{top} \to \mathbb{N}$ , associating with  $s \in S$  the rank of the finitely generated abelian group  $M_s$ . It follows that we obtain a partition  $S = \bigsqcup_{r \geq 0} S_r$  such that every  $S_r$  is an open and closed subscheme of S and  $G|_{S_r}$  is of constant rank r. If  $G=T$  is a torus, this is the so-called partition by type,  $[Oe, 5.4]$ .

A subtorus of a group G of multiplicative type is a monomorphism  $T \to G$ where T is torus. By [IX; 2.5] or [\[Co1,](#page-62-8) B.1.3] in the finite type case,  $T \to G$ is a closed immersion. Obviously,  $T$  is called a *split subtorus* or a *locally* split subtorus, if T is a split torus or a locally split torus. We say T is a maximal split subtorus or maximal locally split subtorus, if the image of T in G is maximal with respect to inclusion among all split subtori or locally split subtori respectively.

<span id="page-39-3"></span><span id="page-39-0"></span>A.2. **Lemma.** Let  $T$  be a torus and let  $Q$  be the quotient of  $T$  by a subgroup of multiplicative type. Then Q is a torus. Moreover,

- <span id="page-39-2"></span>(i) if  $T$  is locally split, then  $Q$  is a locally split torus.
- (ii) If S is connected and T is a split torus, then  $Q$  is a split torus with rank  $Q \leq \text{rank } T$ .

Proof. We first prove [\(i\).](#page-39-0) By definition of "locally split", it suffices to show:

<span id="page-39-1"></span>
$$
(A.2.1) \t\t If T is a split torus, then Q is a locally split torus.
$$

For the proof of [\(A.2.1\)](#page-39-1) we write T in the form  $T = D<sub>S</sub>(M)$  with  $M \cong \mathbb{Z}^r$ and observe that any torus is a group of finite multiplicative type, so that we can apply  $[IX; 2.11(i)]$ : the group Q is a locally split group of multiplicative type. Hence, Zariski-locally  $Q = D<sub>S</sub>(N)$  for some abelian group N. Cartier duality [VIII; §3] provides a monomorphism  $N_S \rightarrow M_S$  of constant group schemes, so that N is torsion free and finitely generated. Therefore,  $N \cong \mathbb{Z}^{r'}$ with  $r' \leq r$ , in particular Q is a split S-torus, proving  $(A.2.1)$  and thus also [\(i\).](#page-39-0)

We next show  $(ii)$  by modifying the proof of  $(A.2.1)$ . Indeed, the reference [IX; 2.11(i)] also says that  $Q$  is a split group of multiplicative type if  $S$  is connected. Hence, as the proof of  $(i)$  shows,  $Q$  is a split torus.

Finally, we can prove the general case: If  $T$  is a torus, then  $Q$  is a torus. Being a torus is local for the fpqc topology. We can therefore assume that T is a split torus. Then  $Q$  is locally split by [\(i\),](#page-39-0) which is all we needed to show.  $\square$ 

<span id="page-40-0"></span>A.3. Example. By [\[Co1,](#page-62-8) B.3.3], any fppf-closed subgroup of a group G of multiplicative type is a group of multiplicative type. Hence, Lemma [A.2](#page-39-3) holds for fppf-closed subgroups of tori.

For example, by [A.2](#page-39-3)[\(ii\),](#page-39-2) if S is a connected scheme and if  $\lambda: \mathbb{G}_{m,S} \to H$ is a non-trivial group homomorphism, then  $\mathbb{G}_{m,S}/\text{Ker}(\lambda) \cong \mathbb{G}_{m,S}$ .

The following lemma is the first step towards the existence of a maximal locally split subtorus.

<span id="page-40-7"></span><span id="page-40-1"></span>A.4. **Lemma.** Let G be an  $S$ -group of finite multiplicative type.

(a) Then the family of locally split  $S$ -subtori of G is a directed poset with respect to inclusion.

<span id="page-40-3"></span>(b) If S is connected, the family of split  $S$ -subtori of G is a directed poset with respect to inclusion.

<span id="page-40-2"></span>*Proof.* [\(a\)](#page-40-1) It suffices to show:

(i) If  $E_1$  and  $E_2$  are locally split subtori of G, then there exists a locally split subtorus of G containing  $E_1$  and  $E_2$ .

To prove this, we consider the S–group scheme  $E_1 \times_S E_2$  and claim that  $E_1 \times_S E_2$  is a locally split torus. Indeed, Zariski-locally  $E_1$  and  $E_2$  have the form  $D<sub>S</sub>(M<sub>1</sub>)$  and  $D<sub>S</sub>(M<sub>2</sub>)$  for free abelian groups of finite type. Since for arbitrary abelian groups  $N_1$  and  $N_2$  we have

<span id="page-40-4"></span>
$$
(A.4.1) \tD_S(N_1) \times_S D_S(N_2) \cong D_S(N_1 \times N_2),
$$

see e.g. [\[Oe,](#page-63-14) 5.1], it follows that  $E_1 \times_S E_2$  is a locally split torus.

Let  $h: E_1 \times_S E_2 \to G$  be the group homomorphism given by multiplication. By  $[IX; 2.7]$ , its kernel  $Ker(h)$  is a subgroup of multiplicative type, so that  $E_3 := (E_1 \times E_2)/\text{Ker}(h)$  is a locally split torus by [A.2](#page-39-3)[\(i\).](#page-39-0) Moreover, again by [IX; 2.7], the canonical map  $E_3 \rightarrow G$  is a monomorphism. Clearly,  $E_1 \cap \text{Ker}(h) = \{0\}.$  Therefore,  $E_1 \rightarrow E_3$  is a closed immersion, and the same holds for  $E_2 \rightarrow E_3$ . Thus, [\(i\)](#page-40-2) holds.

The proof of [\(b\)](#page-40-3) is a straightforward modification of the proof of [\(a\)](#page-40-1), see [XXVI, 6.5]: If  $E_1$  and  $E_2$  are split tori, then so is  $E_1 \times_S E_2$  by [\(A.4.1\)](#page-40-4). The quotient  $E_3$  is then a split torus by [A.2](#page-39-3)[\(ii\).](#page-39-2)

<span id="page-40-8"></span><span id="page-40-5"></span>A.5. Lemma. (a) A monomorphism  $T \hookrightarrow T'$  between S-tori with the same rank functions is an isomorphism.

<span id="page-40-6"></span>(b) Assume that  $S$  is connected and that  $G$  is an  $S$ -group of finite multiplicative type. Then every family of  $S$ -subtori of  $G$ , which is a directed poset with respect to inclusion, admits a unique maximal element.

*Proof.* We can assume that  $S$  is non-empty. All statements are local for the fpqc topology, allowing us to deal with split S-tori.

[\(a\)](#page-40-5) We are given a monomorphism  $f: T = D_S(M) \rightarrow D_S(M') = T'$ , where  $M$  and  $M'$  are free  $\mathbb{Z}-$ modules of rank r. Cartier duality provides an epimorphism  $\hat{f}: M'_{S} \to M_{S}$  of constant S-schemes. For each point s of S,

we obtain a surjective map  $\hat{f}_{\overline{s}}$ :  $M'_{\overline{s}} \to M_{\overline{s}}$  between free Z-modules of same finite rank. The map  $\hat{f}_{\overline{s}}$  is then an isomorphism, and hence so are  $\hat{f}$  and f.

[\(b\)](#page-40-6) Let  $(T_i)_{i\in\mathcal{T}}$  be a family as in the statement of [\(b\)](#page-40-6). Since S is connected, G has of constant rank, say rank r. By the same reason, each  $T_i$  has constant rank, say rank  $T_i = r_i$ . Note  $r_i \leq r$ . Let  $r_j$  be maximal among all  $r_i$ ,  $i \in I$ . We claim that  $T_j$  is a maximal element of the given family. Indeed, fix  $i \in I$  and let  $T_k$  be a member of the family such that  $T_i \subset T_k$  and  $T_j \subset T_k$ . We have  $r_j = r_k$  by maximality of  $r_j$ . Applying [\(a\)](#page-40-5) yields  $T_j = T_k$ , hence  $T_i \subset T_j$ . Thus  $T_i$  is a maximal element; unicity is obvious.

<span id="page-41-0"></span>We can now prove the existence of maximal (locally) split subtori.

<span id="page-41-3"></span>A.6. Proposition. Let G be an  $S$ -group of finite multiplicative type.

- <span id="page-41-1"></span>(a) If E is a maximal locally split S-subtorus of G, then E is unique.
- <span id="page-41-4"></span>(b) If S is connected, G admits a unique maximal locally split S–subtorus.
- <span id="page-41-2"></span>(c) G admits a maximal split S–subtorus.
- (d) If S is connected, G admits a unique maximal split S–subtorus.

*Proof.* Lemma  $A.4(a)$  $A.4(a)$  tells us that the family of locally split S-subtori of G is a directed poset, implying  $(a)$ , while Lemma  $A.5(b)$  $A.5(b)$  shows the existence of a maximal element in case  $S$  is connected. Hence  $(b)$  holds.

[\(c\)](#page-41-4) We use the partition  $S = \bigsqcup_{r \geq 0} S_r$  by rank, see [A.1.](#page-38-1) Up to localizing at some  $S_r$ , we can then assume that G has constant rank r and that  $S \neq \emptyset$ . Since an ascending chain of split tori of bounded rank is stationary, it follows from Lemma  $A.5(a)$  $A.5(a)$  that T admits a maximal split S-subtorus.

[\(d\)](#page-41-2) The reasoning is the same as for  $(b)$  $(b)$ , using this time Lemma  $A.4(b)$  $A.4(b)$ .  $\Box$ 

Proposition [A.6](#page-41-0) implies that a torus  $T$  over a connected  $S$  admits both a maximal split subtorus  $T_0$  and a maximal locally split subtorus  $T_{l0}$ . Clearly,  $T_0 \subset T_{10}$ . The following Proposition [A.7](#page-41-5) gives a sufficient criterion for the two subtori to coincide, while the Examples [A.9](#page-42-0) show that in general the two tori do not coincide.

In Proposition [A.7](#page-41-5) we denote by  $X_*(T) = \text{Hom}_{S-\text{gr}}(\mathbb{G}_m, T)$  the commutative  $S$ -group of cocharacters of a torus  $T$ ; it is a locally constant  $S$ -group scheme. We also recall [\[St,](#page-64-0) Tag 033N]: a connected, normal locally noetherian scheme is integral and hence has a field of fractions.

<span id="page-41-5"></span>A.7. Proposition. Assume that S is connected, normal and locally noether-ian with field of fractions K. By Lemma [A.6](#page-41-0)[\(d\)](#page-41-2), the  $S$ -torus T has a unique maximal split  $S$ -subtorus, denoted  $T_0$ .

- <span id="page-41-7"></span><span id="page-41-6"></span>(a) We have  $X_*(T_0)(K) = X_*(T_0)(S) = X_*(T)(S) = X_*(T)(K)$ .
- <span id="page-41-8"></span>(b)  $T_0$  is the unique maximal locally split S-subtorus of T.
- (c) The formation of  $T_0$  commutes with Zariski localization.

*Proof.* We shall use that T is isotrivial  $[X, 5.16]$ , that is, there exists a finite étale cover  $S'$  of S which splits T. Grothendieck's Galois theory [\[SGA1,](#page-63-15) V,  $\S8$ ] (or see footnote [\(2\)\)](#page-21-8) allows us to assume that the cover S' is a connected Galois cover over S. We denote the Galois group of  $S'/S$  by  $\Gamma$ ; this is also the Galois group of  $K'/K$ , where  $K'$  is the fraction field of  $S'$ .

[\(a\)](#page-41-6) We have  $X_*(T)(S') = X_*(T)(K')$ ; by taking Γ-invariants, we obtain  $X_*(T)(S) = X_*(T)(K)$  and, similarly,  $X_*(T_0)(S) = X_*(T_0)(K)$ . We have the obvious inclusion  $X_*(T_0)(S) \subset X_*(T)(S)$ ; so it remains to establish the other inclusion. To do so, we are given  $\lambda : \mathbb{G}_{m,S} \to T$ . Its image E is a split S-torus, so that Lemma [A.4\(](#page-40-7)[b\)](#page-40-3) implies that  $E \subset T_0$ . It follows that  $\lambda$ factorizes through  $T_0$ , so that  $\lambda$  comes from  $X_*(T_0)(S)$ .

[\(b\)](#page-41-7) Let  $T_{l0}$  be the maximal locally S-split subtorus of T. Clearly  $T_0$  is the maximal S-split subtorus of  $T_{l0}$ . Since  $T_{l0,K}$  is split, applying [\(a\)](#page-41-6) to  $T_{l0}$ , shows that  $X_*(T_0)(K) = X_*(T_0)(K)$ . It follows that  $T_0$  and  $T_{l0}$  have the same rank, so that  $T_0 = T_{l0}$  in view of Lemma [A.5\(](#page-40-8)[a\)](#page-40-5).

[\(c\)](#page-41-8) This follows immediately from [\(a\).](#page-41-6)  $\Box$ 

A.8. **Remarks.** (a) That  $X_*(T)(S) = X_*(T)(K)$  can be also proven by applying [X, 8.4].

(b) The argument of Proposition [A.7](#page-41-5)[\(b\)](#page-41-7) can be generalized as follows. Suppose that S is an integral scheme with field of fractions  $K$  and that  $T$ is an S-torus. Let  $T_0$  and  $T_{l0}$  be the maximal split and locally maximal split subtorus of T. If  $T_{0,K} = T_{l0,K}$  (this is the crucial assumption), then  $T_0 = T_{l0}.$ 

<span id="page-42-0"></span>A.9. Examples. Example [\(a\)](#page-42-1) below shows that [A.7](#page-41-5)[\(c\)](#page-41-8) is wrong without the normality assumption, while Example  $(b)$  shows that  $A.7(b)$  $A.7(b)$  is wrong without the normality assumption. These examples are taken from  $[Co1,$ Exc. 2.4.12].

Let k be an algebraically closed field of characteristic zero. For each  $n \geq 1$ , let  $X_n = \text{Spec}(A_n)$  be the k-scheme obtained by gluing  $2^n$  affine lines in a loop, with 0 on the *i*-th line glued to 1 on the  $(i + 1)$ -th line  $(i \in \mathbb{Z}/2^n\mathbb{Z})$ . This means that

$$
A_n = \{ (P_i) \in \prod_{i \in \mathbb{Z}/2^n \mathbb{Z}} k[x_i] \mid P_i(0) = P_{i+1}(1) \ \forall \ i \in \mathbb{Z}/2^n \mathbb{Z} \}.
$$

We have then a canonical  $k$ -map  $f_n : \widetilde{X}_n = \bigsqcup_{i \in \mathbb{Z}/2^n \mathbb{Z}} \mathbf{A}_k^1 \to X_n$ , obtained by inclusion  $A_n \subset \prod_{i \in \mathbb{Z}/2^n \mathbb{Z}} k[x_i]$ .

<span id="page-42-1"></span>(a) The group  $\mathbb{Z}/2\mathbb{Z}$  acts on  $X_1$  by permuting  $x_0$  and  $x_1$ . Since  $\mathbb{Z}/2\mathbb{Z}$  acts freely on  $X_1(k) = k \sqcup k' \sim$ , where the equivalence relation  $\sim$  identifies 0 of each summand with 1 of the following summand, the action of  $\mathbb{Z}/2\mathbb{Z}$  on  $X_1$  is free [\[DG,](#page-62-22) Cor. III, 2.2.5]. The algebraic group  $\mathbb{Z}/2\mathbb{Z} \cong \mu_2$  is diagonalizable, so that the fppf quotient  $X = X_1/(\mathbb{Z}/2\mathbb{Z})$  is representable by the affine scheme  $X = \text{Spec}(A)$  where  $A = (A_1)^{\mathbb{Z}/2\mathbb{Z}} = \{ P \in k[x] \mid P(0) = P(1) \},\$ [VIII.5.1]. Furthermore, the quotient map  $q: X_1 \to X$  is a  $\mathbb{Z}/2\mathbb{Z}$ -torsor.

We observe that  $X$  is a nodal affine curve. We denote by  $\alpha$  its nodal point and note that  $X \setminus \{a\} \cong \mathbb{G}_m$ . Summarizing, we have a cartesian diagram of  $\mathbb{Z}/2\mathbb{Z}$ -torsors



.

Clearly the  $\mathbb{Z}/2\mathbb{Z}$ -torsor  $q: X_1 \to X$  is trivial once restricted to  $X \setminus \{a\}.$ We claim:

<span id="page-43-1"></span>(A.9.1) The  $\mathbb{Z}/2\mathbb{Z}$ -torsor  $q: X_1 \to X$  is not trivial at the local ring  $\mathcal{O}_{X,q}$ .

Assume to the contrary that there exists a splitting  $s_a: Spec(\mathcal{O}_{X,a}) \to X_1$ of q. It extends to a splitting  $s: U \to X_1$  of q defined on an affine open neighborhood U of a in X. Let V be the inverse image of U in  $A_k^1$ ; this is an affine open neighborhood of 0 and 1. Then s induces a section  $\widetilde{s}$  of  $V \sqcup V \rightarrow V$ . Since  $\mathbf{A}_k^1$  is irreducible, V is connected, hence  $\widetilde{s}$  is the first (or the second) component map. It follows that  $\tilde{s}(V)$  contains the two nodal points  $a_0, a_1$  of  $X_1$ , so that  $s(a) = a_0 = a_1$ . This is a contradiction, proving the claim  $(A.9.1)$ .

We consider the isotrivial X-torus  $T = \Re_{X_1/X}(\mathbb{G}_m)/\mathbb{G}_m$ , that is, the quotient of the Weil restriction  $\mathfrak{R}_{X_1/X}(\mathbb{G}_m)$  by the diagonal  $\mathbb{G}_m$ . Its isomorphism class is determined by  $[X_1] \in H^1(X, \mathbb{Z}/2\mathbb{Z})$ . Then  $T|_{X \setminus \{a\}}$  is split, but  $T \times_S \text{Spec}(\mathcal{O}_{X,a})$  is not split according to [\(A.9.1\)](#page-43-1).

<span id="page-43-0"></span>(b) We now assume  $n \geq 2$ . The map  $\mathbb{Z}/2^{n+1}\mathbb{Z} \to \mathbb{Z}/2^n\mathbb{Z}$ ,  $[i] \mapsto [i]$  induces a morphism

$$
\widetilde{g}_n: \bigsqcup_{i \in \mathbb{Z}/2^{n+1}\mathbb{Z}} \mathbf{A}^1_k \to \bigsqcup_{i \in \mathbb{Z}/2^n\mathbb{Z}} \mathbf{A}^1_k
$$

which is a  $\mathbb{Z}/2\mathbb{Z}$ -torsor, where  $\mathbb{Z}/2\mathbb{Z}$  is the kernel of  $\mathbb{Z}/2^{n+1}\mathbb{Z} \to \mathbb{Z}/2^n\mathbb{Z}$ and acts by permuting the factors. This map induces a  $\mathbb{Z}/2\mathbb{Z}$ -torsor  $g_n$ :  $X_{n+1} \to X_n$  fitting in the cartesian diagram



We claim

<span id="page-43-2"></span>(A.9.2) The 
$$
\mathbb{Z}/2\mathbb{Z}
$$
-torsor  $g_n: X_{n+1} \to X_n$  is locally trivial  
for the Zariski topology, but is not trivial.

Since a section of  $g_n$  must come from a section of  $\tilde{g}_n$ , the argument of [\(a\)](#page-42-1) shows that such a section does not exist. For each  $i \in [0, 2<sup>n</sup> - 2]$ , we denote

by  $\tilde{U}_i$  (resp.  $\tilde{V}_i$ ) the open subset of  $\tilde{X}_n$  (resp.  $\tilde{X}_{n+1}$ ) which is  $\mathbf{A}_k^1 \sqcup \mathbf{A}_k^1$  located at i and  $i + 1$ . Also, we denote by  $U_i$  (resp.  $V_i$ ) the fiber product



.

.

(resp. with  $k[V_n] \subset A_{n+1}$ ). We then get a cartesian of isomorphisms



Since the  $U_i$ 's cover  $X_n$ , we obtain that  $g_n: X_{n+1} \to X_n$  admits sections for the Zariski topology, establishing the claim [\(A.9.2\)](#page-43-2).

We consider the  $X_n$ -torus  $T_n = \Re_{X_{2n}/X_n}(\mathbb{G}_m)/\mathbb{G}_m$  of rank one, whose isomorphism class is determined by  $[X_{2n}] \in H^1(X_n, \mathbb{Z}/2\mathbb{Z})$ . Then  $T_n$  is isotrivial and  $(A.9.2)$  shows that  $T_n$  is locally split of rank 1, but is not split.

# <span id="page-44-0"></span>Appendix B. Parabolic subgroups of reductive groups via the dynamic method

In this appendix we consider parabolic and Levi subgroups of a reductive group  $H$  over a *connected* scheme  $S$  that can be described by the dynamic method, reviewed in [4.5.](#page-22-0) The appendix complements §[4](#page-20-0) on minimal parabolic subgroups and their Levi subgroups.

We start with an improvement of [\[Gi1,](#page-62-9) Thm. 7.3.1]. According to part (2) of that theorem, a reductive  $S$ -group scheme H over a connected S is isotropic in the sense of [4.1](#page-20-2) if and only if its radical torus  $rad(H)$  is isotropic or  $H$  is reducible, also defined in [4.1.](#page-20-2) This characterization is a consequence of part (1) of [\[Gi1,](#page-62-9) Thm. 7.3.1] which proves Theorem [B.1](#page-44-1) without the important invariance of  $\lambda$  under the S–group scheme Aut $(H, P, L)$ , the automorphisms of  $H$  normalizing  $P$  and  $L$ .

<span id="page-44-1"></span>B.1. **Theorem.** Let S be a connected scheme and let P be a parabolic subgroup of the reductive S–group H, equipped with a Levi subgroup  $L \subset P$ . Then there exists a cocharacter  $\lambda : \mathbb{G}_m \to H$  which is fixed by  $Aut(H, P, L)$ and which allows a dynamic description of P and L as  $P = P_H(\lambda)$  and  $L = C_H(\lambda)$ .

*Proof.* Our proof is a refinement of the proof of [\[Gi1,](#page-62-9) Thm. 7.3.1].

Because S is connected, the reductive group  $H$  has constant type  $[XXII]$ , 2.8], so it is a form of a Chevalley S-group G equipped with a Killing couple  $(B, T)$  and defined over  $\mathbb{Z}$ . This gives rise to a root datum and a Dynkin diagram  $\Delta$ . For  $I \subset \Delta$  we let  $P_I$  and  $L_I$  be the standard parabolic and Levi subgroup associated with I. Since triples of type  $(H, P, L)$  allow descent, there exists  $I \subset \Delta$  such that  $(H, P, L)$  is the twist of  $(G, P_I, L_I)$  under the sheaf torsor  $E = \underline{\text{Isom}}((G, P_I, L_I), (H, P, L)).$ 

We now move to the adjoint quotients  $H^{\,\mathrm{ad}}=H/\operatorname{Z}_H$  and  $G^{\,\mathrm{ad}}=G/\operatorname{Z}_G$ where  $Z_H$  and  $Z_G$  are the centres of H and G respectively. By [\[Gi1,](#page-62-9) Lem. 3.2.1], the pairs  $(P^{ad}, L^{ad}) = (P/Z_H, L/Z_H)$  and the analogously defined  $(P_I^{\text{ad}}, L_I^{\text{ad}})$  are pairs of parabolic subgroups and Levi subgroups of  $H^{\text{ad}}$  and  $G^{\text{ad}}$  respectively. Using the last line of [\[Gi1,](#page-62-9) Ex. 7.2], the original proof of [B.1](#page-44-1) shows that  $(P_I^{\text{ad}}, L_I^{\text{ad}})$  has a dynamic description with respect to a cocharacter  $\lambda_I^{\text{ad}}$ :  $\mathbb{G}_m \to G^{\text{ad}}$ , which is fixed by  $\text{Aut}(G^{\text{ad}}, P_I^{\text{ad}}, L_I^{\text{ad}})$ . Since  $(H<sup>ad</sup>, P<sup>ad</sup>, L<sup>ad</sup>)$  is a twisted form of  $(G<sup>ad</sup>, P<sub>I</sub><sup>ad</sup>, L<sub>I</sub><sup>ad</sup>)$ , the pair  $(P<sup>ad</sup>, L<sup>ad</sup>)$ has a dynamic description with respect to the twist  $\lambda^{\text{ad}}$ :  $\mathbb{G}_m \to H^{\text{ad}}$  of  $\lambda_I^{\text{ad}}$ . By construction,  $\lambda^{ad}$  is fixed by

$$
Aut(Had, Pad, Lad) = Aut(Gad, PIad, LIad)E.
$$

Next, we consider the derived groups  $\mathscr{D}(H)$  and  $\mathscr{D}(G)$  and recall that H<sup>ad</sup> and  $G^{\text{ad}}$  are quotients of  $\mathscr{D}(H)$  and  $\mathscr{D}(G)$  respectively. Let n be an integer annihilating the finite diagonalizable S-group scheme  $\mathbb{Z}_{\mathscr{D}(G)}$ . Then *n* also annihilates the centre  $\mathbb{Z}_{\mathscr{D}(H)}$  of  $\mathscr{D}(H)$ . Hence the cocharacter  $(\lambda^{ad})^n \colon \mathbb{G}_m \to$ H<sup>ad</sup> uniquely lifts to a cocharacter  $\lambda: \mathbb{G}_m \to \mathscr{D}(H)$ . The original proof of [B.1](#page-44-1) shows that  $(P, L)$  has a dynamic description with respect to  $\lambda$ . It remains to be shown that  $\lambda$  is fixed by  $Aut(H, P, L)$ . This follows from the unicity of the lifting of  $(\lambda^{ad})^n$ , since the action of  $Aut(H, P, L)$  on  $Hom_{S-gp}(\mathbb{G}_m, H^{ad})$ factorizes through the map  $\text{Aut}(H^{\text{ad}}, P^{\text{ad}}, L^{\text{ad}})$ .

<span id="page-45-1"></span>B.2. Irreducible and anisotropic actions. We extend the notion of a reducible or isotropic reductive group scheme [\(4.1\)](#page-20-2) to the setting where an S-group scheme M acts on a reductive  $S$ -group scheme H by automorphisms of H. In this situation we say that the action is reducible if it normalizes a pair  $(P, L)$  consisting of an everywhere proper parabolic subgroup P of H in the sense of [4.1](#page-20-2) and a Levi subgroup  $L$  of  $P$ . Otherwise, the action is called *irreducible*. If S is connected, as in  $B.3$  and  $B.4$ , then P is everywhere proper if and only if  $P$  is proper because the type of a parabolic subgroup is locally constant [XXVI, 3.2].

Similarly, we say that the action of  $M$  on  $H$  is *isotropic*, if it centralizes an S-subgroup of H isomorphic to  $\mathbb{G}_{m,S}$ . Otherwise, the action is called anisotropic.

<span id="page-45-0"></span>For S the spectrum of a field and  $M = \mathbb{G}_{m,S}$ , Corollary [B.3](#page-45-0) is [\[BT,](#page-62-12) 4.23].

<span id="page-45-2"></span>B.3. Corollary. Assume that  $S$  is connected and that an  $S$ -group scheme M acts on the reductive  $S$ -group H. Then the following are equivalent:

- <span id="page-45-3"></span>(i) The action of M on H is isotropic.
- (ii) The action of M on  $rad(H)$  is isotropic or the action of M on H is reducible.

*Proof.* [\(i\)](#page-45-2)  $\implies$  [\(ii\):](#page-45-3) Suppose the image of the non-trivial map  $\lambda: \mathbb{G}_{m,S} \to$  $H$  is centralized by the action of  $M$ , and consider the parabolic subgroup  $P = P_H(\lambda)$  together with its Levi subgroup  $L = C_H(\lambda)$ , see [4.5.](#page-22-0) If  $P = H$ , then  $\lambda$  is central and takes values in rad(H), so that the action of M on rad(H) is isotropic. Otherwise, P is a proper parabolic subgroup. Since M normalizes  $(P, L)$ , its action on H is reducible.

 $(ii) \implies (i)$  $(ii) \implies (i)$ : If the action of M on rad(H) is isotropic, so is its action on H. Assume now that the action of M on H is reducible, that is, there exists a pair  $(P, L)$  consisting of a proper parabolic subgroup P of H and a Levi subgroup  $\subset P$ , such that  $(P, L)$  is normalized by M. Theorem [B.1](#page-44-1) provides a group homomorphism  $\lambda: \mathbb{G}_{m,S} \to H$  such that  $(P, L) = (\mathbb{P}_H(\lambda), \mathbb{C}_H(\lambda))$ and such that  $\lambda$  is fixed by Aut $(H, P, L)$ . In particular,  $\lambda$  is non-central. After quotioning by  $\text{Ker}(\lambda)$ , we can assume that  $\lambda$  is a monomorphism, [A.3.](#page-40-0) Since M acts on H through  $Aut(H, P, L)$ , the image of  $\lambda$  is fixed by M. Thus, the action of M on H is isotropic.  $\Box$ 

<span id="page-46-0"></span>The following Proposition [B.4](#page-46-0) is a complement to Theorem [B.1.](#page-44-1)

B.4. Proposition. Assume that  $S = Spec(R)$  is affine and connected, that M is a flat affine  $R$ -group scheme whose geometric fibers are linearly reductive, e.g., M is of multiplicative type, and that M acts on the reductive group  $H$ . If  $M$  normalizes a parabolic subgroup  $P$  of  $H$ , there exists an M–invariant group homomorphism  $\lambda: \mathbb{G}_{m,S} \to H$  such that  $P = P_H(\lambda)$  and such that  $C_H(\lambda)$  is a Levi subgroup of P which is normalized by M.

*Proof.* Let U be the unipotent radical of P which, according to [XXVI, 2.1], is Aut(P)–linearizable, that is, U admits a composition series  $1 = U_n \subset$  $U_{n-1} \subset \cdots \subset U_1 = U$  which is Aut(P)-stable and such that  $U_i/U_{i+1} =$  $W(E_i)$  where each  $E_i$  is a locally free R-module of finite type on which the action of  $Aut(H, P)$  is linear. A fortiori, the action of M on U is linearizable.

By [XXVI, 2.3 and 1.9], the set of sections of  $P \to X := P/U$  is non-empty and is a principal homogeneous space under  $U(R)$ . For each R-ring A, the group  $M(A)$  acts on the A-sections of  $P \to X$ . Let  $s : X \to P$  be such a section. For each  $m \in M(A)$ , there exists a unique  $u_A(m) \in U(A)$  such that  ${}^m s_A = {}^{u_A(m)^{-1}} s_A$ . For  $m_1, m_2 \in M(A)$ , we apply  $m_1$  to  ${}^{m_2} s_A = {}^{u_A(m_2)^{-1}} s_A$ and get

$$
{}^{m_1m_2}s_A = {}^{m_1(u_A(m_2))^{-1}}({}^{m_1}s_A) = {}^{m_1(u_A(m_2))^{-1}}({}^{u_A(m_1)^{-1}}s_A)
$$
  

$$
= {}^{m_1(u_A(m_2))^{-1}}{}^{u_A(m_1)^{-1}}(s_A),
$$

so that  $u_A(m_1m_2) = u_A(m_1) \cdot {}^{m_1}(u_A(m_2))$ . The map  $u_A : M(A) \to U(A)$ ,  $m \mapsto u_A(m)$ , is therefore a 1-cocycle for the action on  $M(A)$  on  $U(A)$ , so that the data of the  $u_A$ 's for A running over all R-rings define an Hochschild 1cocycle (or crossed homomorphism)  $u \in \mathcal{Z}^1_{\text{coc}}(M,U)$ , as defined by Demarche [\[Dem,](#page-62-23) §3.2]. Thus, we obtain a cohomology class  $[u] \in H^1_{\text{coc}}(M, U)$ . We now claim

<span id="page-46-1"></span>(B.4.1) 
$$
H_{\text{coc}}^1(M, U) = 1.
$$

#### 48 P. GILLE AND E. NEHER

We postpone the proof of  $(B.4.1)$  for the moment. Assuming  $(B.4.1)$ , we get an element  $v \in U(R)$  such that  $u_A(m) = v_A^{-1}$  $_{A}^{-1}$ <sup>m</sup> $v_A$  for all  $m \in U(A)$ . It follows that  ${}^m s_A = {}^{m_v} v_A^{-1} v_A s_A$ , so that  $s' = {}^{v_s} s$  is an *M*-invariant section of  $P \to X$ . In particular,  $s(X)$  is a Levi subgroup of P which is M-invariant. Theorem [B.1](#page-44-1) then shows that there exists an  $M$ –invariant homomorphism  $\lambda: \mathbb{G}_m \to H$  such that  $(P, s(X)) = (P_H(\lambda), C_H(\lambda))$  as desired.

We now come to the proof of  $(B.4.1)$ . According to  $[Dem, Prop. 3.2.8]$ , we have an exact sequence of pointed sets

$$
1 \to \mathrm{H}^1_{\mathrm{coc}}(M,U) \to \mathrm{H}^1_0(M,U) \to \mathrm{H}^1(R,U)
$$

where  $H_0^1(M, U)$  is the set of isomorphism classes of M-U-torsors over R [\[Dem,](#page-62-23) 3.2.6]. We know from [XXVI, 2.2] that  $H^1(R, U) = 1$ , so that it is enough to show that  $H_0^1(M, U) = 1$ .

According to Grothendieck for  $M$  of multiplicative type  $[IX, 3.1]$  and Margaux in general [\[Ma,](#page-63-16) Th. 1.2], we have  $H_{\text{coc}}^1(M, U_i/U_{i+1}) = 0$  for  $i =$  $1, \ldots, n-1$  so that  $H_0^1(M, U_i/U_{i+1}) = 0$  by the exact sequence above applied to  $U_i/U_{i+1}$ . On the other hand, by [\[Dem,](#page-62-23) Prop. 3.5.1] for  $i=1,\ldots,n-1$ we have an exact sequence of pointed sets

$$
H_0^1(M, U_{i+1}) \to H_0^1(M, U_i) \to H_0^1(M, U_i/U_{i+1}),
$$

so that we conclude by "dévissage" that  $H_0^1(M, U) = 1$ , finishing the proof of  $(B.4.1)$ .

B.5. Remarks. (a) The proof of Proposition [B.4](#page-46-0) is clearly inspired by Demarche's paper [\[Dem\]](#page-62-23) and by McNinch's approach [\[Mc\]](#page-63-17).

(b) Let k be a field, let H be a reductive k–group and let  $M \subset H$  be a subgroup of H. Following Serre  $[Se1, Se2]$  $[Se1, Se2]$ , see also  $[BMR]$ , the subgroup M is called H–completely reducible, if whenever  $M \subset P$ , where P is a parabolic subgroup of H, then  $M \subset L$  for some Levi subgroup of P.

Provisionally, we will use the completely analogous terminology for reductive groups over a scheme S. Specializing Proposition [B.4](#page-46-0) to the case of M being a subgroup of  $H$  acting on  $H$  by inner automorphisms, the proposition implies that  $M$  is  $H$ -completely reducible. This generalizes [J<sub>a</sub>, 11.24], proven there for algebraically closed fields and restated in [\[BMR,](#page-61-7) Lem 2.6]. In characteristic 0, the result is due to Mostow.

## <span id="page-47-0"></span>Appendix C. Hermitian and quadratic forms over rings

In this appendix we review the concepts needed for Propositions [5.4](#page-29-0) and [5.5,](#page-29-1) but not more. This appendix is not an introduction to the topics of the section heading, like for example the book  $[Kn2]$ , nor did the authors strive for the most general setting, i.e., considering forms over schemes instead of base rings. Despite this limitation, the appendix will hopefully still serve as a useful quick introduction to regular hermitian and nonsingular quadratic forms over rings, their associated isometry groups and the link to their flat cohomology sets.

Throughout this appendix,  $R$  is an arbitrary base ring,  $S$  is an algebra in R-alg, equipped with an R-linear involution  $\sigma$ . This will later be specialized to the *hermitian case*, where  $S$  is a quadratic étale extension with standard involution  $\sigma$ , and to the *quadratic case*, where  $S = R$  and  $\sigma = \text{Id}_R$ . The general setting avoids a duplication of definitions.

<span id="page-48-0"></span>C.1. Sesquilinear forms. (a) (Modules) Unless specified otherwise, all modules considered will be finite projective S–modules. Given such a module M, we denote by  $\sigma M$  its  $\sigma$ -conjugate, i.e., the S-module with the same additive group as M, but with the S–action given by  $s \star (\sigma m) = \sigma(s)m$  for  $s \in S$  and  $m \in M$ , using the S-action of M on the right-hand side.

Instead of the usual dual space  $\text{Hom}_S(M, S)$  we will use its twisted version

$$
M^* := \sigma \operatorname{Hom}_S(M, S).
$$

One has an isomorphism of S–modules  $M^* \stackrel{\sim}{\longrightarrow} \text{Hom}_S({}_\sigma M, S)$ , given by  $\varphi \mapsto \big(\sigma m \mapsto (\sigma \circ \varphi)(m)\big)$  ([\[Kn2,](#page-63-12) I; Lem. (2.1.1)]).

(b) (Sesquilinear forms) A sesquilinear form, also called an  $(S, \sigma)$ –sesquilinear form is a bi-additive map  $h: M \times M \rightarrow S$ , which is defined on a finite projective  $S$ -module  $M$  and which satisfies

$$
h(m_1s_1, m_2s_2) = \sigma(s_1) h(m_1, m_2) s_2
$$

for all  $m_i \in M$  and  $s_i \in S$ . We will denote such a sesquilinear form by  $(M, h)$  or simply by h.

Given two sesquilinear forms  $(M_1, h_1)$  and  $(M_2, h_2)$ , an S-linear map  $f: M_1 \to M_2$  is an *isometry* if  $\hat{f}$  is bijective and if  $h_2(f(m_1), f(m'_1)) =$  $h_1(m_1, m'_1)$  holds for all  $m_1, m'_1 \in M_1$ . In this case, we say that  $(M_1, h_1)$ and  $(M_2, h_2)$  are *isometric*, abbreviated by  $(M_1, h_1) \cong (M_2, h_2)$  or simply by  $h_1 \cong h_2$ .

(c) (*Hermitian and symmetric forms*) A sesquilinear form  $(M, h)$  is called  $(S, \sigma)$ -hermitian or simply hermitian, if  $h(m, m') = \sigma(h(m', m))$  holds for all  $m, m' \in M$ . In the quadratic case, i.e.,  $(S, \sigma) = (R, Id)$ , we say h is symmetric instead of hermitian.

<span id="page-48-1"></span>(d) (Adjoints, regularity) A sesquilinear form  $(M, h)$  gives rise to the S– linear map  $h^*: M \to M^*, m \mapsto h(m, -)$ , called the *adjoint of*  $(M, h)$ . Mapping  $h$  to  $h^*$  gives rise to a bijection between the S-module of sesquilinear forms on M and  $\text{Hom}_S(M, M^*)$ . We call h regular if  $h^*$  is a bijection, hence an isomorphism of S–modules.

<span id="page-48-2"></span>(e) (*Orthogonality*) Given two sesquilinear forms  $(M_1, h_1)$  and  $(M_2, h_2)$ , their *orthogonal sum* is the sesquilinear form  $h_1 \perp h_2$  defined on  $M =$  $M_1 \oplus M_2$  by

$$
(h_1 \perp h_2) (m_1 + m_2, m_1' + m_2') = h_1(m_1, m_1') + h(m_2, m_2')
$$

for  $m_1, m'_1 \in M_1$  and  $m_2, m'_2 \in M_2$ . The sesquilinear form  $h_1 \perp h_2$  is regular ifand only if  $h_1$  and  $h_2$  are regular ([\[Kn2,](#page-63-12) I, (3.6.2.2)]). The analogous statement holds for the property "hermitian".

(f) (*Base change*) Let  $T \in R$ -alg. Then  $S_T = S \otimes_R T$  is an object in R-alg with a T-linear involution  $\sigma_T = \sigma \otimes 1_T$ , and  $S_T/T$  satisfy the assumptions of this appendix. Given a finite projective  $S$ -module M we put  $M_T = M \otimes_R T = M \otimes_S S_T$ , which is canonically a finite projective  $S_T$  module. The twisted dual respects this base change: the canonical map  $\omega: M^* \otimes_R T \stackrel{\sim}{\longrightarrow} (M_T)^*$  is an isomorphism because M is finite projective.

Let now  $(M, h)$  be an  $(S, \sigma)$ -sesquilinear form. Then  $(M, h)$  uniquely extends to an  $(S_T, \sigma_T)$ -sesquilinear form  $h_T$  on  $M_T$  by requiring

<span id="page-49-0"></span>
$$
h_T(m \otimes t, m' \otimes t') = h(m, m') \otimes tt'
$$

for  $m, m' \in M$  and  $t, t' \in T$ . The adjoint maps  $(h_T)^*$  and the base change  $h^* \otimes 1_R$  are related by the commutative diagram



<span id="page-49-4"></span>Hence, if  $(M, h)$  is regular, then so is  $(M_T, h_T)$ .

(g) (Hyperbolic forms  $\text{[Kn2, I, (3.5)]}$  $\text{[Kn2, I, (3.5)]}$  $\text{[Kn2, I, (3.5)]}$ ) Let U be a finitely generated projective S-module. One defines a hermitian module  $\mathbb{H}(U) = (U \oplus U^*, h_{\mathbb{H}(U)})$ by  $h_{\mathbb{H}(U)}(u_1 + \varphi_1, u_2 + \varphi_2) = \varphi_1(u_2) + \sigma(\varphi_2(u_1))$  for  $u_i \in U$  and  $\varphi_i \in U^*$ . One calls  $(M, h)$  hyperbolic if it is isometric to  $\mathbb{H}(U)$  for some U.

A hyperbolic space  $(M, h) = \mathbb{H}(U)$  is equipped with a natural action of the R-group scheme  $\Re_{S/R}(\mathbf{GL}_S(M))$ ,  $\Re(\cdot)$  = Weil restriction, as follows. For each R-ring T, each  $g \in \mathfrak{R}_{S/R}(\mathbf{GL}_S(M))(T) = \mathrm{GL}_{S\otimes_R T}(M\otimes_R T)$  and each pair  $(u, f) \in (U \otimes_R T) \oplus (M^* \otimes_R T)$ , we have  $g.(u, f) = (g.u, f \circ g^{-1})$ . It follows that we have a closed embedding

(C.1.2) 
$$
\mathfrak{R}_{S/R}(\mathbf{GL}_S(U)) \hookrightarrow \mathbf{U}(M,h)
$$

of  $R$ –group schemes.

(h) (Totally isotropic submodules) Let  $(M, h)$  be a hermitian module. A submodule  $U \subset M$  is called *totally isotropic* if U is complemented and  $U \subset U^{\perp}$ , where, in general,

<span id="page-49-3"></span>
$$
U^{\perp} = \{ m \in M : h(m, U) = 0 \}.
$$

A vector  $m \in M$  is *isotropic* if m is unimodular and  $h(m, m) = 0$ . In case  $(S, \sigma) = (R, \text{Id})$  we also require  $q(m) = 0$ .

(i) A sesquilinear form  $(M, h)$  is *diagonalizable* if there exists a basis  $(e_1, \ldots, e_n)$  of the S-module M such that  $h(e_i, e_j) = 0$  whenever  $i \neq j$ .

(j) The free S-module  $M = S^n$ ,  $n \in \mathbb{N}_+$ , carries the *split hermitian form*  $h_{0,n}$ , defined by

<span id="page-49-1"></span>(C.1.3) 
$$
h_{0,n}\left(\sum_{i=1}^n e_i s_i, \sum_{j=1}^n e_j s'_j\right) = \sum_{i=1}^n \sigma(s_i) s'_i.
$$

<span id="page-49-2"></span>

The term "split" will be justified in Lemma [C.5](#page-51-1) where we consider regular hermitian forms.

<span id="page-50-0"></span>We characterize regularity of sesquilinear forms in the following lemma.

<span id="page-50-2"></span>C.2. Lemma. Assume  $S \in R$ -alg is finite projective as R-module. Then the following are equivalent for a sesquilinear form h.

- <span id="page-50-3"></span>(i) h is regular;
- <span id="page-50-4"></span>(ii)  $h_T$  is regular for all  $T \in R$ -alg;
- <span id="page-50-6"></span>(iii)  $h_K$  is regular for all algebraically closed fields  $K \in R$ -alg;
- <span id="page-50-5"></span>(iv)  $h_{R_m}$  is regular for all maximal  $\mathfrak{m} \in \text{Spec}(R)$ ;
- (v) there exists a Zariski cover  $(f_1, \ldots, f_n)$  of R such that  $h_{R_{f_i}}$  is regular for all i;
- <span id="page-50-7"></span><span id="page-50-1"></span>(vi)  $h_{R/\mathfrak{m}}$  is regular for all maximal  $\mathfrak{m} \in \text{Spec}(R)$ ;
- (vii)  $h_T$  is regular for some faithfully flat  $T \in R$ -alg.

*Proof.* Let  $h = (M, h)$ . We will use that the adjoint map  $h^*: M \to M^*$ is bijective as  $S$ –linear map if and only if the underlying  $R$ –linear map is bijective.Moreover, by transitivity of "finite projective" ( $[Fe, 1.1.8]$ ), M and  $M^*$  are finite projective as R–modules. We will use the following scheme to prove [C.2.](#page-50-0)



The implications [\(i\)](#page-50-2)  $\implies$  [\(ii\)](#page-50-3) and (i)  $\implies$  [\(iv\)](#page-50-6) follow from the commutative diagram  $(C.1.1)$ , while [\(ii\)](#page-50-3)  $\implies$  [\(iii\)](#page-50-4) and (ii)  $\implies$  [\(vi\)](#page-50-7) are trivial. The implication [\(iii\)](#page-50-4)  $\implies$  [\(ii\)](#page-50-3) follows again from [\(C.1.1\)](#page-49-0), for  $(R, S, T)$  replaced by the triple  $(F, S \otimes_R F, K)$ . For [\(iv\)](#page-50-6)  $\iff$  [\(vi\)](#page-50-7) we need  $h_{R_m}^*$  bijective  $\iff h_{R/\mathfrak{m}}^*$  bijective, which is [\[B:AC1,](#page-62-3) II, §3.2, Cor. de la Prop. 6].

[\(iv\)](#page-50-6)  $\implies$  [\(v\):](#page-50-5) By [\[B:AC1,](#page-62-3) II, §5.1, Prop. 2(ii)] there exists for every  $\mathfrak{m} \in$ Spec(R) a  $g_{\mathfrak{m}} \in R$  such that  $h_{g_{\mathfrak{m}}}^*$  is bijective. The existence of a Zariski cover with the stated property then follows as usual:  $\{g_m : m \in \text{Spec}(R) \text{ maximal}\}\$ generates R as an ideal; one takes a generating set  $\{f_1, \ldots, f_n\}$  of this ideal. [\(v\)](#page-50-5)  $\implies$  [\(vii\):](#page-50-1) By [\[B:AC1,](#page-62-3) II, §5.1, Prop. 3] the algebra  $S = R_{f_1} \times \cdots \times R_{f_n}$ is faithfully flat and  $h_S^* = \prod_i h_{f_i}^*$  is bijective. Finally, the implication [\(vii\)](#page-50-1)  $\implies$  [\(i\)](#page-50-2) follows from [\[B:AC1,](#page-62-3) I, §3.1, Prop. 2].

<span id="page-50-8"></span>[C.3](#page-50-8). **Hermitian spaces.** In C.3 – [C.7](#page-52-1) we consider the "hermitian case", i.e.,

<span id="page-50-9"></span> $(C.3.1)$  S is a quadratic étale R–algebra with standard involution  $\sigma$ .

A hermitian sesquilinear form will be hermitian with respect to  $(S, \sigma)$ . A hermitian space  $(M, h)$  is a hermitian module  $(M, h)$  with h a regular hermitian form; a *hermitian space of rank r* is a hermitian space  $(M, h)$  for which  $M$  is a locally free S-module of constant rank  $r$ .

<span id="page-51-4"></span>C.4. **Lemma** (LG diagonalizability, hermitian case). Let R be an  $LG$ -ring, let  $(S, \sigma)$  be a quadratic étale R–algebra with standard involution  $\sigma$ , and let  $(M, h)$  be a hermitian space with M having constant positive rank. Then  $(M, h)$  is diagonalizable, in particular

<span id="page-51-2"></span>(C.4.1) 
$$
\{h(m, m) : m \in M\} \cap R^{\times} \neq \emptyset.
$$

*Proof.* This is  $[GPR2, Exc. 21.21(a)]$ ; its proof is given in  $[GPR3]$ : one can assume that M has constant rank; one then shows  $(C.4.1)$  and finally proves diagonalizability by induction on the rank of M (recall [\[Kn2,](#page-63-12) I,  $(3.6.2.1)$ ] and  $1.6(b)$  $1.6(b)$ ).

<span id="page-51-3"></span><span id="page-51-1"></span>C.5. Lemma. Let R be arbitrary and let  $(S, \sigma)$  be a quadratic étale R-algebra with standard involution  $\sigma$ .

(a) (Zariski-diagonalizability) Let  $(M, h)$  be a hermitian space with M being faithfully projective. Then  $(M, h)$  is Zariski-locally diagonalizable, *i.e.*, there exists a Zariski cover  $(f_1, \ldots, f_m)$  of R such that for every  $g =$  $f_1, \ldots, f_m$  the base change  $(M,h)_{R_g} := (M_{R_g}, h_{R_g})$  is the split hermitian form [\(C.1.3\)](#page-49-1) for appropriate  $n \in \mathbb{N}_+$ , possibly varying with g.

- <span id="page-51-6"></span><span id="page-51-5"></span>(b) The following are equivalent for a hermitian form  $(M, h)$ .
	- (i)  $M$  is locally free of rank n and h is regular.
- <span id="page-51-7"></span>(ii) There exists a flat cover  $T \in R$ -alg such that  $(M,h)_T$  is isometric to the split form  $(C.1.3)$ .

*Proof.* [\(a\)](#page-51-3) Let  $\mathfrak{m} \triangleleft R$  be a maximal ideal of R. By Lemma [C.4,](#page-51-4)  $(M, h)$ is diagonalizable over the local ring  $R_{\mathfrak{m}}$ . Therefore, since  $R_{\mathfrak{m}} \cong \underline{\lim}_{f \notin \mathfrak{m}} R_f$ , there exists  $f \notin \mathfrak{m}$  such that  $h_{R_f}$  is diagonalizable. Hence the ideal generated by  $\{f \in R : h|_{R_f}$  is diagonalizable} is all of R, from which [\(a\)](#page-51-3) easily follows.

[\(b\)](#page-51-5) Assume [\(i\).](#page-51-6) Let  $(f_1, \ldots, f_m)$  be a Zariski cover as in [\(a\)](#page-51-3), and put  $T' = R_{f_1} \times \cdots \times R_{f_m}$ . Then T' is a flat cover of R and  $(M,h)_T \cong (S^n,h')$ with a diagonalizable  $h'$ . Since a flat cover  $T''$  of  $T'$  is a flat cover of  $R$ , it suffices to prove [\(ii\)](#page-51-7) under the assumption that  $M \cong S^n$  and that h is diagonalizable with respect to the standard basis  $(e_1, \ldots, e_n)$  of  $S^n$ . Note  $h(e_i, e_i) = r_i \in R^\times$  since h is supposed to be hermitian and regular. Let  $R[X_1, \ldots, X_n]$  be the polynomial ring in *n* variables. Then

$$
T = R[X_1, \ldots, X_r]/(X_1^2 - r_1, \ldots, X_n^2 - r_n)
$$

is a flat cover of R such that  $h_T = h_{0,n}$ .

Conversely, if [\(ii\)](#page-51-7) holds, then M is locally free of rank n ([\[B:AC1,](#page-62-3) II,  $\S 5.3$ , Prop. 4) and h is regular by Lemma [C.2.](#page-50-0)

<span id="page-51-0"></span>C.6. Proposition (Unitary groups). Let  $(S, \sigma)$  be quadratic étale with standard involution  $\sigma$ , and let  $(M, h)$  be a hermitian space. Then the following hold.

<span id="page-51-8"></span>(a)  $A := \text{End}_S(M)$  is an Azumaya S-algebra.

<span id="page-52-3"></span>(b) There exists a unique involution  $\tau_h$  on A, the adjoint involution satisfying

<span id="page-52-5"></span>(C.6.1) 
$$
h(m_1, f(m_2)) = h(\tau_h(f)(m_1), m_2)
$$

for all  $m_i \in M$  and  $f \in A$ . The triple  $(S, A, \tau_h)$  is an Azumaya algebra with involution of second kind in the sense of [\[CF,](#page-62-1) 2.7.0.33].

<span id="page-52-4"></span>(c) We define  $U(M, h) := U(A, \tau_h)$  where  $U(A, \tau_h)$  is the unitary R-group scheme of [\[CF,](#page-62-1) 3.5.0.84]. Then, for every  $R$ -ring T,

$$
\mathbf{U}(M,h)(T) = \{ f \in \mathrm{GL}_{S\otimes_R T}(M\otimes_R T) : f \text{ is an isometry of } h_T \}
$$

<span id="page-52-0"></span>where  $h_T$  is the base change of h in the sense of [C.1](#page-48-0)[\(f\)](#page-49-2).

(d) The group  $U(M, h)$  is a reductive R-group scheme. If M has constant rank n, then  $\mathbf{U}(M,h)$  has type  $A_{n-1}$ .

<span id="page-52-2"></span>(e) Fix  $n \in \mathbb{N}_+$ . The map  $[(M,h)] \mapsto [\underline{\text{Isom}}((S^n,h_{0,n}), (M,h))]$  is a bijection between the set of isometry classes of hermitian spaces over  $(S, \sigma)$ of rank n and  $H^1(R, U(S^n, h_{0,n}))$ . Equivalently, for every hermitian space  $(M, h)$  of rank n, the set of twisted forms of  $(M, h)$  is in bijection with  $\mathrm{H}^1(R,\mathbf{U}(M,h))$ .

*Proof.* For [\(a\)](#page-51-8) see for example [\[Fo,](#page-62-0) 7.1.10]. In case S is a field, [\(b\)](#page-52-3) is proven in [\[KMRT,](#page-63-19) 4.1]; the same proof works in our setting. [\(c\)](#page-52-4) Since  $h((\tau_h(f) \circ$  $f(m_1), m_2$  =  $h(f(m_1), f(m_2))$  by [\(C.6.1\)](#page-52-5), the equation  $\tau_h(f) \circ f = \text{Id}_M$ , defining  $\mathbf{U}(M,h)(T)$ , is equivalent to  $h \circ (f \times f) = h$ , i.e., f is an isometry. [\(d\)](#page-52-0) follows from [\[CF,](#page-62-1) 3.5.0.87].

[\(e\)](#page-52-2) Using  $C.5(b)$  $C.5(b)$ , it is straightforward (but somewhat technical) to define the gerbe of hermitian spaces of rank  $n$  over the big affine fppf site of Spec $(R)$ . The claim then becomes a special case of  $[CF, 2.2.4.5]$ . Alternatively (and more down-to-earth), one can easily adjust the notion of a tensor system of  $\left[GPR2, 54.7\right]$  to apply to hermitian spaces, cf. [C.1\(](#page-48-0)[d\)](#page-48-1). The claim then becomes a consequence of the Descent Theorem [\[GPR2,](#page-62-5) 54.15].  $\Box$ 

**Remark.**Since  $U(M, h)$  is smooth, it is known ([\[Mil1,](#page-63-20) III, Remark 4.8] or [\[CF,](#page-62-1) 2.2.5.15]) that the canonical map  $H^1_{\text{\'et}}(R, \mathbf{U}(M, h)) \xrightarrow{\sim} H^1(R, \mathbf{U}(M, h))$ is a bijection. Thus, one can replace twisted forms with respect to the flat topology in  $C.6(e)$  $C.6(e)$  by twisted forms in the étale topology. The analogous remark applies to  $C.5(b)$  $C.5(b)$ .

<span id="page-52-1"></span>C.7. **Proposition** (Parabolic and Levi subgroups of  $U(M, h)$ ). Assume  $(C.3.1)$ , and let  $(M, h)$  be a hermitian space over  $(S, \sigma)$  containing a totally isotropic submodule  $U \subset M$ .

<span id="page-52-7"></span>(a) Then there exists a totally isotropic submodule V and a submodule  $M'$ such that

<span id="page-52-6"></span>(C.7.1) 
$$
M = (U \oplus V) \perp M', \quad (U \oplus V, h|_{U \oplus V}) \cong \mathbb{H}(U),
$$

in particular  $(M',h|_{M'})$  is a hermitian space.

(b) The subgroup scheme P of  $G = U(M, h)$  which stabilizes U is a parabolic subgroup of the reductive  $R$ -group scheme  $G$ .

<span id="page-53-2"></span>(c) Consider a decomposition  $(M, h) = (U \oplus V) \perp M'$  as in  $(C.7.1)$ , and the embedding  $\mathfrak{R}_{S/R}(\mathbf{GL}(U)) \hookrightarrow G$  defined in [\(C.1.2\)](#page-49-3). Then the R-group scheme  $L = \Re_{S/R}(\mathbf{GL}(U)) \times_R \mathbf{U}(M',h|_{M'})$  is a Levi subgroup of P.

*Proof.* [\(a\)](#page-52-7) Since the trace map  $S \to R$ ,  $s \mapsto s + \sigma(s)$  is surjective [\[Kn2,](#page-63-12) III,  $(4.2.1)$ ], it follows from  $\overline{[Kn2, I, (3.1.1)]}$  that h is an even hermitian module in the sense of  $[Kn2, I, (3.1)]$ . The lemma is therefore a consequence of  $[Kn2, I, (3.1)]$ . I, (3.7.1)]. That V can be chosen totally isotropic follows from the proof of [\[Kn2,](#page-63-12) I, (3.7.1)].

[\(b\)](#page-53-1) We will use the dynamic method of [4.5.](#page-22-0) It is easily verified that

$$
\lambda \colon \mathbb{G}_m \to \mathbf{U}(M, h), \quad t \mapsto t \operatorname{Id}_U + t^{-1} \operatorname{Id}_V + 1_{M'}
$$

defines a cocharacter. Writing a  $q \in U(M, h)(T)$ ,  $T \in R$ -alg, as a matrix  $g = (g_{ij})_{1 \leq i,j \leq 3}$  with respect to the submodules  $(U, V, M')$ , one finds

$$
\lambda(t) (g_{ij}) \lambda^{t})^{-1} = \begin{pmatrix} g_{11} & t^2 g_{12} & tg_{13} \\ t^{-2} g_{21} & g_{22} & t^{-1} g_{23} \\ t^{-1} g_{31} & tg_{32} & g_{33} \end{pmatrix}.
$$

Hence,  $g \in P_G(\lambda)(T) \iff 0 = g_{21} = g_{31} = g_{23}$ , i.e., g stabilizes U and  $U^{\perp} = U \oplus M'$ . Since g is an isometry, the latter conditions are equivalent to  $g$  stabilizing  $U$ .

[\(c\)](#page-53-2) The centralizer of the cocharacter  $\lambda$  above consists of those g that are diagonal with respect to  $(U, V, M')$ . These are easily seen to be the isometries in  $L(T)$ . The claim then follows from the dynamic method.  $\square$ 

**Remarks.** (1) That P in [C.7](#page-52-1)[\(b\)](#page-53-1) is parabolic with Levi subgroup L as in [\(c\)](#page-53-2), is not surprising and likely folklore. It is in the spirit of Appendix T of Conrad's course [\[Co3\]](#page-62-25), which identifies the parabolic subgroups for symplectic and special orthogonal groups over fields.

(2) Suppose that M is free of rank  $n \geq 2$  and that U has rank i for some  $1 \leq i < n/2$ . Then P is of type  $A_{n-1} \setminus \{i\}$ , while the derived group  $\mathscr{D}(L)$ has absolute type  $A_{i-1} \times A_{n-1-i}$  for  $i \geq 2$  and type  $A_{n-2}$  for  $i = 1$ .

<span id="page-53-0"></span>C.8. Quadratic forms. From now on until the end of this appendix we consider quadratic forms. Thus, we specialize [C.1](#page-48-0) to  $(S, \sigma) = (R, Id)$ . Unless stated otherwise, R is arbitrary and, we recall, M is a finite projective  $R$ module. Below is a quick reminder of the basics of quadratic forms over rings; proofs and more can be found in  $\left[\text{Bae}\right]$  and  $\left[\text{Kn2}, I, II\right]$ .

(a) A quadratic form (over R) is a pair  $(M, q)$  consisting of a finite projective R-module M and a map  $q: M \to R$  satisfying  $q(rm) = r^2q(m)$ for all  $r \in R$  and  $m \in M$  and for which the polar form  $b_q$ , defined by  $b_q(m, m') = q(m + m') - q(m) - q(m')$  for  $m, m' \in M$ , is a (symmetric) bilinear form on M. We often abbreviate  $(M, q) = q$  and refer to  $(M, q)$ as a quadratic module. We call  $(M, q)$  a *faithful* quadratic R-module if M

<span id="page-53-1"></span>

is a faithfully projective R–module, [1.5.](#page-6-3) We say that  $(M, q)$  is a quadratic module of rank  $n \in \mathbb{N}$ , if M has constant rank n.

An isometry  $f: (M_1, q_1) \to (M_2, q_2)$  is an R-linear isomorphism  $f: M_1 \to$  $M_2$  satisfying  $q_2 \circ f = q_1$ . If  $(M_1, q_1) = (M_2, q_2) = (M, q)$ , the isometries of  $(M, q)$  form a group  $O(M, q) = O(q)$ , the *orthogonal group of q*.

(b) (Regularity) By definition, a quadratic form q is regular, if  $b<sub>q</sub>$  is regular in the sense of  $C.1(d)$  $C.1(d)$ .

(c) (*Base change*) Let  $T \in R$ -alg and let  $(M, q)$  be a quadratic form. Analogous to [C.1](#page-48-0)[\(f\)](#page-49-2) we consider the T–module  $M_T = M \otimes_R T$ . There exists a quadratic form  $(M_T, q_T)$  uniquely determined by the condition  $q_T(m \otimes t)$  =  $q(m)t^2$  for all  $m \in M$  and  $t \in T$  ([\[GPR2,](#page-62-5) 11.5], [\[Sa,](#page-64-5) Thm. 1]). The polar of  $q_T$  is the base change of the polar  $b_q$  of q, i.e.,  $b_{q_T} = (b_q)_T$ . In particular, if q is regular, then so is  $q_T$ .

(d) (Radical) The *radical* of a quadratic module  $(M, q)$  is the submodule  $rad(q) = \{m \in M : q(m) = 0 = b_q(m, M)\}\$  of M. Given  $T \in R$ -alg, clearly rad(q)  $\otimes_R T \subset \text{rad}(q_T)$ . This inclusion is in general not an equality, cf. [\(e\)](#page-54-2).

<span id="page-54-2"></span>(e) (*Nonsingularity*) A quadratic form  $(M, q)$  is called *nonsingular* if  $rad(q_F) = 0$  for all fields  $F \in R$ -alg. In this case, we call  $(M, q)$  a quadratic space. We point out that a "quadratic space" in the sense of [\[Bae\]](#page-61-3) or [\[GPR2\]](#page-62-5) is a quadratic module  $(M, q)$  with a regular q.

<span id="page-54-3"></span>If q is regular, then  $rad(q) = 0$  and hence [C.2](#page-50-0)[\(ii\)](#page-50-3) implies

(C.8.1) 
$$
q \text{ regular} \implies q \text{ nonsingular}.
$$

The converse of  $(C.8.1)$  is not true. For example, any  $u \in R^{\times}$  gives rise to a nonsingular quadratic form  $\langle u \rangle: R \to R$ , given by  $\langle u \rangle(r) = u r^2$  for  $r \in R$ . Its polar form is the symmetric bilinear form  $b_{\langle u \rangle}(r_1, r_2) = 2ur_1r_2$ . Hence,  $\langle u \rangle$  is regular if and only if  $2 \in R^{\times}$ .

However, if  $2 \in R^{\times}$ , then  $\text{rad}(q) = \{m \in M : b_q(m, M) = 0\}$  for any quadratic form q and so rad(q) is stable under base change. Hence,

(C.8.2) if 
$$
2 \in R^{\times}
$$
, then q is nonsingular  $\iff$  q is regular.

(f) (*Orthogonality*) Given two quadratic forms  $(M_1, q_1)$  and  $(M_2, q_2)$ , their *orthogonal sum* is the quadratic form  $q_1 \perp q_2$ , defined on  $M = M_1 \oplus M_2$  by  $(q_1 \perp q_2)(m_1, m_2) = q_1(m_1) + q_2(m_2)$  for  $m_1 \in M_1$  and  $m_2 \in M_2$ . The polar form of  $q_1 \perp q_2$  is the orthogonal sum  $b_{q_1 \perp q_2} = b_{q_1} \perp b_{q_2}$  defined in [C.1\(](#page-48-0)[e\)](#page-48-2).

By [C.1\(](#page-48-0)[e\)](#page-48-2), the quadratic form  $q = q_1 \perp q_2$  is regular if and only if  $q_1$  and  $q_2$  are regular. Regarding nonsingularity, one easily sees:

<span id="page-54-1"></span> $q_1 \perp q_2$  nonsingular  $\implies q_1$  and  $q_2$  nonsingular, (C.8.3)  $q_1$  regular and  $q_2$  nonsingular  $\implies$   $q_1 \perp q_2$  nonsingular.

<span id="page-54-0"></span>(g) (Direct products) Let  $R = R_0 \times \cdots \times R_n$  be a direct product. Every R-module M uniquely decomposes  $M = M_0 \times \cdots \times M_n$  as a direct product of  $R_i$ -modules  $M_i = R_i M$ . The R-module M is finite projective if and only if every  $R_i$ -module  $M_i$  is finite projective.

Let  $(M, q)$  be a quadratic R-module. Then  $b_q(M_i, M_j) = 0$  for  $i \neq j$  and  $(M_i, q_i)$  with  $q_i = q|_{M_i}$  is a quadratic  $R_i$ -module. Thus

 $(M, q) = (M_0, q_0) \perp ... \perp (M_n, q_n), \quad q_i = q|_{M_i}.$ 

The quadratic R–module  $(M, q)$  is regular (nonsingular respectively) if and only if every  $(M_i, q_i)$  is a regular (nonsingular respectively) quadratic  $R_i$ module.

A standard way to obtain the situation considered here occurs by letting  $M = M_0 \times \cdots \times M_n$  be the *rank decomposition* of a finite projective Rmodule M for which  $M_i$ ,  $0 \leq i \leq n$ , is a finite projective  $R_i$ -module of constant rank i.

(h) (*Hyperbolic spaces*) Let U be a finite projective R-module. The associated *hyperbolic space*  $\mathbb{H}(U)$  is the quadratic module  $(U^* \oplus U, \text{hyp})$  with quadratic form  $hyp_U(\varphi \oplus u) = \varphi(u)$ , where  $\varphi \in U^*$  and  $u \in U$ . The quadratic form hyp<sub>U</sub> is regular, hence nonsingular by  $(C.8.1)$ . The polar form of hyp is the hyperbolic symmetric bilinear form of  $C.1(g)$  $C.1(g)$ . In general, a hyperbolic space is a quadratic module  $(M, q)$  isometric to some  $\mathbb{H}(U)$ .

<span id="page-55-1"></span>(i) (Split quadratic forms) Let  $m \in \mathbb{N}$ . The quadratic form  $q_{0,2m}$  is the hyperbolic form associated with the free  $R$ –module  $R^m$ . After identifying  $R^{m*} = R^m$ , it is given on  $R^{2m}$  by

<span id="page-55-6"></span>(C.8.4) 
$$
q_{0,2m}(r_{-m},\ldots,r_{-1},r_1,\ldots,r_m)=\sum_{i=1}^m r_ir_{-i},
$$

It is regular, hence also nonsingular by  $(C.8.1)$ . The quadratic form  $q_{0,2m+1} =$  $\langle 1 \rangle \perp q_{0,2m}$  on  $R^{2m+1}$ , defined by

$$
(C.8.5) \tqquad q_{0,2m+1}(r_{-m},\ldots,r_{-1},r_0,r_1\ldots,r_m)=r_0^2+\sum_{i=1}^m r_ir_{-i},
$$

is nonsingular, e.g. by the even rank case, by  $(C.8.3)$  and by nonsingularity of  $\langle 1 \rangle$ , see [\(e\)](#page-54-2). We will refer to  $q_{0,n}$  for n even or odd as the split quadratic forms, see Proposition [C.9](#page-55-0) for a justification for this terminology.

<span id="page-55-2"></span><span id="page-55-0"></span>C.9. **Proposition.** For a faithful quadratic module  $(M, q)$  the following are equivalent:

- <span id="page-55-4"></span>(i) q is nonsingular;
- <span id="page-55-5"></span>(ii)  $q_T$  is nonsingular for all  $T \in R$ -alg;
- <span id="page-55-7"></span>(iii)  $q_{R/\mathfrak{m}}$  is nonsingular for all maximal ideals  $\mathfrak{m} \triangleleft R$ ;
- (iv) there exists a flat cover  $(R_1, \ldots, R_n)$  such that each  $M \otimes_R R_i$  is a free  $R_i$ -module of finite rank  $r_i$  and  $q_{R_i}$  is the split quadratic form  $q_{0,r_i}$  over  $R_i$  defined in  $C.8(i)$  $C.8(i)$ ;
- <span id="page-55-3"></span>(v)  $q_S$  is nonsingular for some faithfully flat  $S \in R$ -alg;

<span id="page-55-9"></span>If M has constant rank  $n \in \mathbb{N}_+$ , then [\(i\)](#page-55-2)–[\(v\)](#page-55-3) are equivalent to

- (vi) q is regular if n is even and q is semiregular in the sense of  $[Kn2,$ IV,  $(3.1)$  if n is odd;
- <span id="page-55-8"></span>If  $R$  is a field, then  $q$  is nonsingular if and only if
	- (vii) q is nondegenerate in the sense of  $[EKM, (7.17)]$  $[EKM, (7.17)]$ , i.e., one of the following two conditions hold:

- (a) q is regular, or
- (b) char(R) = 2, rad(q) = 0, dim<sub>R</sub>{ $m \in M : b_q(m, M) = 0$ } = 1, and  $\dim_R M$  is odd.

*Proof.* [\(i\)](#page-55-2)  $\iff$  [\(ii\)](#page-55-4) is easy. It is shown in [\[Sw,](#page-64-6) Prop. 1.1, Cor. 1.3] that (i)  $\iff$  [\(iii\)](#page-55-5)  $\iff$  [\(iv\)](#page-56-0)<sub>ét</sub>, defined as

<span id="page-56-0"></span>(iv)<sub>ét</sub> there exists an étale cover  $(R_1, \ldots, R_n)$  of R such that each  $M \otimes_R R_i$ is a free  $R_i$ -module of finite rank  $r_i$  and  $q_{R_i}$  is hyperbolic as in [\(C.8.4\)](#page-55-6) if  $r_i$  is even, while  $q_{R_i}$  is the orthogonal sum of a hyperbolic form and a 1-dimensional form  $\langle a_i \rangle$  with  $a_i \in R_i^{\times}$  in case  $r_i$  is odd.

One shows  $(iv)_{\text{\'et}} \implies (iv)$  $(iv)_{\text{\'et}} \implies (iv)$  as in the proof of  $C.5(b)(i) \implies (ii)$  $C.5(b)(i) \implies (ii)$  $C.5(b)(i) \implies (ii)$  $C.5(b)(i) \implies (ii)$ . If  $(iv)$ holds, then [\(v\)](#page-55-3) is satisfied with  $S = R_1 \times \cdots \times R_n$ . For (v)  $\iff$  [\(i\)](#page-55-2) see [\[Lo2,](#page-63-21) Lem. 6.2]. Thus, we proved that  $(i) - (v)$  $(i) - (v)$  $(i) - (v)$  are all equivalent using the diagram

[\(ii\)](#page-55-4) ks +3[\(i\)](#page-55-2) ks +3 KS [\(iv\)](#page-55-7)´et ks +3 [\(iii\)](#page-55-5) [\(v\)](#page-55-3) [\(iv\)](#page-55-7) ks

If R is a field, then [\(i\)](#page-55-2)  $\iff$  [\(vii\)](#page-55-8) by [\[EKM,](#page-62-16) 7.16]. Again over a field, [\(vii\)](#page-55-8) implies that q is semiregular in the odd rank case by  $[Kn2, IV, (3.1.7)].$ Finally, if M has constant rank over an arbitrary  $R$ , then  $q$  is regular respectively semiregular if and only if it is so over all residue fields  $R/\mathfrak{m}$  ([\[Kn2,](#page-63-12) IV;  $(3.1.5)$  for odd rank, [C.2](#page-50-0) for even rank). Thus, to prove that [\(vi\)](#page-55-9) is equivalent to  $q$  being nonsingular, we can assume that  $R$  is a field, in which case the equivalence [\(vi\)](#page-55-9)  $\iff$  [\(vii\)](#page-55-8) is easily established using [\[Kn2,](#page-63-12) IV,  $(3.1.7)$  for a semiregular q.

<span id="page-56-1"></span>C.10.Quadratic algebras. We recall ( $[Kn2, I, (1.3.6),$  $[Kn2, I, (1.3.6),$  and III, §4) that a quadratic  $R$ -algebra is an algebra S in R-alg whose underlying R-module is projective of rank 2. Properties:

(a) A quadratic R-algebra S has a standard involution  $\sigma_S$ . If S is free with basis  $\{1, z\}$  and hence  $z^2 = az + b$  with  $a, b \in R$ , it is given by  $\sigma_S(z) = a - z$ . In general,  $\sigma_S$  is defined by Zariski-descent.

(b) (*Automorphisms*) We denote by  $\mathbb{Z}/2\mathbb{Z}_R$  the constant group scheme of locally constant functions with values in  $\mathbb{Z}/2\mathbb{Z} = \{0, 1\}$ . By |Kn2, III,  $(4.1.2)$ , there exists a natural R–group homomorphism

<span id="page-56-2"></span>
$$
(C.10.1) \t\t \psi: \mathbb{Z}/2\mathbb{Z}_R \longrightarrow \text{Aut}_R(S),
$$

<span id="page-56-3"></span>whichis an isomorphism if S is étale ( $[Kn2, III, (4.1.2)].$  $[Kn2, III, (4.1.2)].$ 

(c) (*Example*) Let  $S = S_0 \oplus S_1 \in R$ -alg be a  $\mathbb{Z}/2\mathbb{Z}$ -graded R-algebra. We denote by  $Aut(S, S_1)$  the group of R-linear automorphisms of the R-algebra S, i.e., the automorphisms  $\alpha$  of S satisfying  $\alpha(S_1) = S_1$ , and by

$$
(C.10.2) \qquad \mathbf{Aut}(S, S_1)
$$

the R–group scheme representing the R–functor  $T \mapsto \text{Aut}(S_T, (S_1)_T)$ .

<span id="page-57-0"></span>Suppose now that  $S = S_0 \oplus S_1$  satisfies

<span id="page-57-1"></span>(i)  $R \xrightarrow{\sim} S_0, r \mapsto r1_S$ , and

(ii)  $\theta: S_1 \otimes_R S_1 \xrightarrow{\sim} S_0$ ,  $s_1 \otimes s_1' \mapsto s_1 s_1'$  (isomorphism of R-modules).

Thus S is a quadratic R-algebra,  $(S_1, \theta)$  is a discriminant module in the sense of [\[Kn2,](#page-63-12) III, §3], and its standard involution  $\sigma_S$  is the grading automorphism,

(C.10.3) 
$$
\sigma_S(s_i) = (-1)^i s_i, \quad s_i \in S_i, \quad (i = 0, 1,
$$

which follows from the free case by localization. We have isomorphisms of R–group schemes

<span id="page-57-6"></span>(C.10.4) 
$$
\mathbf{Aut}(S, S_1) \xrightarrow{\sim} \mathbf{Aut}(S_1, \theta) \xleftarrow{\sim} \mu_{2,R},
$$

where the first (obvious) isomorphism is obtained by restriction and where the second isomorphism is  $x \mapsto x \, \text{Id}_{S_1}$  [\[Kn2,](#page-63-12) III, (3.2.1)].

<span id="page-57-3"></span>C.11. Discriminant algebras. Let  $(M, q)$  be a faithful quadratic R-space and let  $\mathcal{C}\ell(M,q) = \mathcal{C}\ell(q)$  be its Clifford algebra, [\[Kn2,](#page-63-12) IV, §1]. It is a (Z/2Z)– graded R–algebra:  $\mathcal{C}\ell(q) = \mathcal{C}\ell_0(q) \oplus \mathcal{C}\ell_1(q)$ . The discriminant algebra  $Dis(q)$ of  $(M, q)$  is the subalgebra of  $\mathcal{C}\ell(q)$  centralizing  $\mathcal{C}\ell_0(q)$ :

$$
Dis(q) = \mathcal{C}\ell(q)^{\mathcal{C}\ell_0(q)}.
$$

<span id="page-57-4"></span>Some facts that we will use (see [\[Kn2,](#page-63-12) IV, §4]).

(a) Discriminant algebras respect base change and direct products of base rings.

(b)  $\mathcal{D} := \mathcal{D}is(q)$  inherits the  $\mathbb{Z}/2\mathbb{Z}$ -grading of  $\mathcal{C}\ell(q)$ ,

$$
\mathcal{D} = \mathcal{D}_0 \oplus \mathcal{D}_1, \quad \mathcal{D}_j = \mathcal{D} \cap \mathcal{C} \ell_j(q), \ j = 0, 1.
$$

It is a quadratic  $R$ –algebra in the sense of [C.10.](#page-56-1)

(c) (The group homomorphism Dis) By the universal property of the Clifford algebra  $\mathcal{C}\ell(q)$ , every  $g \in O(q)$  induces an automorphism  $\mathcal{C}\ell(q)$  of the algebra  $\mathcal{C}\ell(q)$  stabilizing  $\mathcal{C}\ell_0(q)$  and  $\mathcal{C}\ell_1(q)$ , and hence an automorphism of the  $\mathbb{Z}/2\mathbb{Z}$ -graded algebra  $\mathcal{D} = \mathcal{D}_0 \oplus \mathcal{D}_1$ . Thus, we get a homomorphism of groups,

<span id="page-57-7"></span><span id="page-57-2"></span>(C.11.1) 
$$
\mathcal{D}is: O(q) \longrightarrow Aut(\mathcal{D}, \mathcal{D}_1), \quad g \mapsto \mathcal{C}\ell(g)|_{\mathcal{D}} =: \mathcal{D}is(g).
$$

(d) (*Even rank*) Assume  $M$  has constant even rank. Thus  $q$  is regular by [C.9](#page-55-0)[\(vi\).](#page-55-9) In this case  $\mathcal D$  is the centre of  $\mathcal{C}\ell_0(q)$ , in particular  $\mathcal D_1=0$ , and  $\mathcal D$ a quadratic étale  $R$ –algebra. Hence, by  $(C.10.1)$ ,

(C.11.2) 
$$
\mathbf{Aut}(\mathcal{D}, \mathcal{D}_1) = \mathbf{Aut}(\mathcal{D}) \stackrel{\sim}{\longleftarrow} (\mathbb{Z}/2\mathbb{Z})_R.
$$

The standard involution  $\sigma_{\mathcal{D}}$  of the quadratic R–algebra D is (of course) the standard involution of the quadratic étale  $R$ –algebra  $\mathcal D$ .

<span id="page-57-5"></span>(e) (Odd rank) Assume M has constant odd rank. Then the discriminant algebra D is the centre of  $Cl(q)$ :  $\mathcal{D} = \mathbb{Z}(\mathcal{C}\ell(q))$ . Moreover,  $\mathcal{D} = \mathcal{D}_0 \oplus \mathcal{D}_1$  is

a  $\mathbb{Z}/2\mathbb{Z}$ -graded R-algebra satisfying the conditions [\(i\)](#page-57-0) and [\(ii\)](#page-57-1) of [C.10\(](#page-56-1)[c\)](#page-56-3). Thus

(C.11.3) 
$$
\mathbf{Aut}(\mathcal{D}, \mathcal{D}_1) \cong \boldsymbol{\mu}_{2,R}.
$$

C.12. **Lemma** (Realizing the standard involution of  $Dis(q)$ ). Let  $(M, q)$  be a faithful quadratic space, let  $x \in M$  with  $q(x) \in R^{\times}$ , and let  $\rho_x$  be the associated reflection, given by  $\rho_x(m) = m - b_q(m, x)q(x)^{-1}x$  for  $m \in M$ . Then the automorphism  $Dis(\rho_x) \in Aut(\mathcal{D}, \mathcal{D}_1), \mathcal{D} = Dis(q)$ , is the standard involution of D:

<span id="page-58-2"></span>(C.12.1) Dis(ρx) = σD.

*Proof.* The element  $x \in M \subset \mathcal{C}\ell_1(q)$  is invertible in  $\mathcal{C}\ell(q)$  with inverse  $x^{-1} = q(x)^{-1}x$ . We will use the well-known formula relating  $\rho_x(m)$ ,  $m \in M$ , with the inner automorphism of  $\mathcal{C}\ell(q)$  induced by x:

(C.12.2) 
$$
\rho_x(m) = -xmx^{-1}
$$

which follows from  $xm^{\text{-}1} = (xm)(xq(x)^{-1}) = (-mx+b_q(m,x))(xq(x)^{-1}) =$  $-m + b_q(m, x)q(x)^{-1}x = -\sigma_x(m)$ . It implies

$$
Dis(\rho_x)(c_j) = (-1)^j x c_j x^{-1}, \qquad (c_i \in \mathcal{C}\ell_j(q), j = 0, 1).
$$

For the proof of  $(C.12.1)$ , we can without loss of generality assume that M has constant rank.

Suppose M has constant even rank. By [\[Kn2,](#page-63-12) IV, (4.3.1.4)],  $\sigma_{\mathcal{D}}(d)x = xd$ holds for  $d \in \mathcal{D}$ . Since  $\mathcal{D} \subset \mathcal{C}\ell_0(q)$  we get  $\mathcal{D}is(\rho_x)(d) = xdx^{-1} = \sigma_{\mathcal{D}}(d)$ .

Suppose M has constant odd rank. Since then  $\mathcal{D} = \mathcal{Z}(\mathcal{C}\ell(q))$ , we obtain for  $d_j \in \mathcal{D}_j$ ,  $j = 0, 1$ , that  $Dis(\rho_x)(d_j) = (-1)^j x d_j x^{-1} = (-1)^j d_j x x^{-1} =$  $(-1)^{j}d_{j} = \sigma_{\mathcal{D}}(d_{j})$  by [C.10\(](#page-56-1)[c\)](#page-56-3).

<span id="page-58-3"></span><span id="page-58-0"></span>C.13. **Lemma.** Let  $(M, q)$  be a quadratic space over an LG-ring R.

- <span id="page-58-4"></span>(a) The following are equivalent:
	- (i)  $q(M) \cap R^{\times} \neq \emptyset$ ;
	- (ii) M is faithfully projective.
- <span id="page-58-5"></span><span id="page-58-1"></span>(b) If M is faithfully projective, the group homomorphism

$$
Dis: O(q) \to Aut(\mathcal{D}, \mathcal{D}_1)
$$

of [\(C.11.1\)](#page-57-2) is surjective.

*Proof.* [\(a\)](#page-58-3) The equivalence [\(ai\)](#page-58-4)  $\iff$  [\(aii\)](#page-58-5) is proven in [\[GPR2,](#page-62-5) Lem. 11.26] for a regular q. The proof in the nonsingular case is essentially the same. We reproduce it here for the convenience of the reader.

Assuming [\(ai\)](#page-58-4), it follows that  $q(M) \not\subset \mathfrak{m}$  for every maximal ideal  $\mathfrak{m} \triangleleft R$ . In particular,  $M_{R_m} \neq 0$ , which, by [\(1.5\)](#page-6-3), implies [\(aii\)](#page-58-5).

Conversely, suppose that M is faithfully projective. We can view  $q$  as a quadratic polynomial on the affine scheme  $\mathbf{W}(M)$  and define  $U = \mathbf{W}(M)_{q}$ , the principal open subscheme determined by  $q$ . Let  $\mathfrak m$  be a maximal ideal of R, put  $k = R/\mathfrak{m}$  and observe that  $M_k \neq 0$  by [1.5,](#page-6-3) while  $\text{rad}(q_k) = 0$  by

nonsingularity of q, in particular  $q_k \neq 0$ . Hence  $U(k) = \{m \in M_k : q_k(m) \neq 0\}$  $0\} \neq \emptyset$ , and so  $U(R) \neq \emptyset$  by [1.4](#page-6-0)[\(a\),](#page-6-1) i.e., [\(ai\)](#page-58-4) holds.

[\(b\)](#page-58-1) Applying the rank decomposition of quadratic modules  $C.8(g)$  $C.8(g)$  and  $C.11(a)$  $C.11(a)$ , we see that we can assume that M has constant rank r.

Assume that r is odd. By [C.11](#page-57-3)[\(e\)](#page-57-5) and [\(C.10.4\)](#page-57-6), an automorphism  $q \in$  $\text{Aut}(\mathcal{D}, \mathcal{D}_1)$  has the form  $g|_{\mathcal{D}_0} = \text{Id}$  and  $g|_{\mathcal{D}_1} = x \text{Id}_{\mathcal{D}_1}$  for some  $\det(g) = x \in$  $\mu_2(R)$ . Observe  $x \text{Id}_M \in \text{O}(q)$ . By construction of  $\mathcal{C}\ell(q)$ , the orthogonal map x Id<sub>M</sub> acts on  $\mathcal{C}\ell_0(q)$  as identity, and on  $\mathcal{C}\ell_1(q)$  as x Id. Thus  $Dis(x \text{ Id}_M) = g$ .

Assume that r is even. By [C.11](#page-57-3)[\(d\)](#page-57-7) and the proof of  $[Kn2, III, (4.1.2)],$  to every  $g \in Aut(\mathcal{D}, \mathcal{D}_1) = Aut(\mathcal{D})$  one can associate a unique complete system  $(\epsilon_0, \epsilon_1 = 1 - \epsilon_0)$  of orthogonal idempotents  $\epsilon_i$ , such that g stabilizes  $\epsilon_i \mathcal{D}$ and satisfies  $g|_{\epsilon_0, \mathcal{D}} = \text{Id}, g|_{\epsilon_1, \mathcal{D}}$  is the standard involution of the quadratic  $\epsilon$ tale  $\varepsilon_1R$ –algebra  $\varepsilon_1$  D. By [1.1](#page-3-3)[\(a\)](#page-3-4),  $\varepsilon_1R$  is an LG–ring. Hence, by [C.8\(](#page-53-0)[g\)](#page-54-0), we can assume that  $R = \varepsilon_1 R$ , and have to show that there exists  $f \in O(q)$ such that  $Dis(f) = \sigma_{\mathcal{D}}$ . Since  $q(M) \cap R^{\times} \neq \emptyset$  by [\(a\)](#page-58-3), this follows from  $(C.12.1)$ .

<span id="page-59-1"></span>C.14. Orthogonal group schemes. Let  $(M, q)$  be a quadratic form over R. The R–group functor  $O(q)$ , assigning to  $T \in R$ -alg the group  $O(q)(T) =$  $O(q_T)$  is represented by an affine finitely presented R–group scheme  $O(q)$ ,  $[CF, Def. 4.1.0.2]$  $[CF, Def. 4.1.0.2]$  or  $[Col, page 364]$ . Properties that we will use:

<span id="page-59-4"></span>(a) (Direct products) Let  $R = R_0 \times \cdots \times R_n$  be a direct product. Orthogonal groups respect the decomposition of  $C.8(g)$  $C.8(g)$  as follows:

$$
O(q) = O(q_0) \times \cdots \times O(q_n),
$$
  
\n
$$
O(M, q) \cong p_{0*} (O(M_0, q_0)) \times \cdots \times p_{n*} (O(M_n, q_n))
$$
  
\n(isomorphism of *R*-group schemes)

where  $p_i$ :  $Spec(R_i) \rightarrow Spec(R)$  is the morphism associated with the canonical projection  $R \to R_i$ .

<span id="page-59-2"></span>(b) $\mathbf{O}(q)$  is in general not reductive. For example ([\[Co1,](#page-62-8) Thm. C.1.5]), if  $(M, q)$  has constant positive rank n, then  $\mathbf{O}(q)$  is smooth if and only if either *n* is even or *n* is odd and  $2 \in R^{\times}$ .

(c) The homomorphism  $Dis$  of  $(C.11.1)$  respects base change and thus defines a homomorphism of  $R$ –group schemes

(C.14.1) 
$$
\mathcal{D}is: \mathbf{O}(q) \to \mathbf{Aut}(\mathcal{D}, \mathcal{D}_1), \quad \mathcal{D} = \mathcal{D}is(q),
$$

<span id="page-59-0"></span>assigning to  $T \in R$ -alg the map  $Dis(T): \mathcal{O}(q_T) \to \text{Aut}(\mathcal{D}_T, \mathcal{D}_{1,T}).$ 

C.15. **Lemma** (Cohomology). Fix  $n \in \mathbb{N}_+$ . The map

<span id="page-59-3"></span>
$$
[(M,q)] \mapsto [\underline{\text{Isom}}((R^n, q_{0,n}), (M,q))]
$$

is a bijection between the set of isometry classes of quadratic spaces over R of rank n and  $H^1(R, O(q_{0,n}))$ . Equivalently, for every quadratic space  $(M, q)$  of rank n, the set of twisted forms of  $(M, q)$  is in bijection with  $H^1(R, O(M, q))$ .

*Proof.* This can be proven in the same way as  $C.6(e)$  $C.6(e)$ , replacing the reference  $C.5(b)$  $C.5(b)$  used there by [C.9.](#page-55-0)

It may be appropriate to remind the reader that we are using fppfcohomology which does in general not coincide with étale cohomology, see [C.14](#page-59-1)[\(b\)](#page-59-2). In this respect, the description of the Galois cohomology set  $H_{Gal}^1(R, O(q))$  $H_{Gal}^1(R, O(q))$  $H_{Gal}^1(R, O(q))$  in characteristic 2 ([\[KMRT,](#page-63-19) p. 408]) is instructive.

<span id="page-60-0"></span>C.16. Special orthogonal group scheme  $SO(q)$ . Let  $(M, q)$  be a faithful quadratic R–space, and let  $\mathcal{D} = Dis(q)$  be its discriminant algebra. We define the R–group scheme as the kernel of the homomorphism  $Dis$  of  $(C.14.1):$  $(C.14.1):$ 

(C.16.1) 
$$
\mathbf{SO}(q) = \text{Ker}(\mathcal{D}is).
$$

Thus, for  $T \in R$ -alg we have  $SO(q)(T) = \{g \in O(q_T) : \mathcal{D}is(g) = \mathrm{Id}_{\mathcal{D}is(q_T)}\}.$ Since the inclusion inc:  $\text{Aut}(\mathcal{D}, \mathcal{D}_1) \to \text{Aut}(\mathcal{D})$  is a monomorphism, this definition coincides with the one of [\[Kn2,](#page-63-12) IV,  $(5.1)$ ], where  $SO(q)$  is defined as the kernel of inc  $\circ$  Dis. The discussion below shows that it also agrees with the definition used in [\[Co1,](#page-62-8) App. C]. However, it coincides with the definition of  $SO(q)$  in [\[CF,](#page-62-1) §4.3] only in the even rank case. Following is a list of properties of  $SO(q)$  that we will use.

(a) (Direct products) Let  $R = R_0 \times \cdots \times R_n$  be a direct product of rings, and let  $(M, q) = (M_0, q_0) \times \cdots \times (M_n, q_n)$  be the corresponding decomposition into a direct product of quadratic  $R_i$ –modules, [C.8](#page-53-0)[\(g\)](#page-54-0). Analogously to  $C.14(a)$  $C.14(a)$ , we have

(C.16.2) 
$$
\mathbf{SO}(M,q) \cong p_{1*}(\mathbf{SO}(M_1,q_1)) \times \cdots \times p_{n*}(\mathbf{SO}(M_n,q_n))
$$

into a direct product of  $R$ –group schemes. It will follow from this and the discussion below that, in general,  $SO(q)$  is a reductive R–group scheme which is semisimple if rank  $M \geq 3$ .

<span id="page-60-2"></span>(b) (**SO** and determinants) By  $\overline{[Kn2, IV, (5.1.1)]}$  we have a group homomorphism

(C.16.3) 
$$
\det: \mathbf{O}(q) \longrightarrow \boldsymbol{\mu}_{2,R}, \quad g \mapsto \det(g).
$$

We always have  $SO(q) \subset \text{Ker}(\text{det})$ ; equality holds if M has constant odd rank or if  $2 \in R^{\times}$ . This is proven in [\[Kn2,](#page-63-12) IV, (5.1.1)], but note the misprint in (3) of loc. cit., where "⊂" should be "=", as one can see from the proof.

<span id="page-60-1"></span>(c)  $(Odd~rank)$  Let  $(M, q)$  be a quadratic space of odd rank. Then the morphism  $z_M : \mu_{2,R} \longrightarrow \mathbf{O}(q), x \mapsto x \, \mathrm{Id}_M$ , is a section of det. Hence, by [\(b\)](#page-60-2),

(C.16.4) 
$$
\mathbf{O}(q) \cong \boldsymbol{\mu}_2 \times_R \mathbf{SO}(q).
$$

If M has constant rank 1, then  $SO(q) = \{ \star \}$ , and if M has constant odd rank  $2n + 1 \geq 3$ , then  $SO(q)$  is an adjoint semisimple R-group scheme of type  $B_n$   $(B_1 = A_1 \text{ for } n = 1)$  [\[Co1,](#page-62-8) Prop. C.3.10].

(d) (*Even rank*) Let  $(M, q)$  be a quadratic space of even rank  $2n \geq 2$ . Then  $q$  is regular by [C.9,](#page-55-0) and the following hold.

(i) Since the map  $\psi$  of [\(C.10.1\)](#page-56-2) is an isomorphism, the *Dickson map* 

Dick :=  $\psi^{-1} \circ \mathcal{D}is \colon \mathbf{O}(q) \to \mathbb{Z}/2\mathbb{Z}_R$ 

is a group homomorphism satisfying  $SO(q) = \text{Ker}(\text{Dick})$ . Moreover, by  $[Co1, C.2.8]$  or  $[Kn2, IV, (5.2.2)],$  the sequence

(C.16.5) 
$$
1 \longrightarrow SO(q) \longrightarrow O(q) \xrightarrow{Dick} \mathbb{Z}/2\mathbb{Z}_R \longrightarrow 1
$$

<span id="page-61-5"></span>is exact.

- (ii) If M has constant rank 2, then  $SO(q)$  is a rank one torus. Indeed, by descent, we are reduced to the hyperbolic case  $q = xy$  where  $SO(q) = \mathbb{G}_m$  [\[Kn2,](#page-63-12) IV, 5.1.3 and V, (2.6.3)].
- (iii) If M has constant rank  $2n \geq 4$ , then  $SO(q)$  is a semisimple R–group scheme of type  $D_n$ .

(e) Suppose  $M = U \oplus V$  such that  $q(U) = 0 = q(V)$  and  $(M, q) \longrightarrow \mathbb{H}(U)$ under an isometry identifying V with  $U^*$ . Then, associating with  $g \in GL(U)$ the map  $M \to M$ ,  $(u, v) \mapsto (gu, g^{-1}v)$ , gives rise to a closed embedding of R–group schemes

<span id="page-61-4"></span>(C.16.6) GL(U) −→ SO(q).

C.17. Proposition (Parabolic and Levi subgroups of  $SO(q)$ ). Let  $(M, q)$  be a quadratic space, let U and V be faithfully projective submodules and let  $M' \subset M$  be a submodule such that

<span id="page-61-8"></span>
$$
M = (U \oplus V) \perp M', \quad q(U) = 0 = q(V), \quad (U \oplus V, q|_{U \oplus V}) \cong \mathbb{H}(U).
$$

(a) The subgroup scheme P of  $SO(q)$ , which stabilizes U, is a parabolic subgroup of  $SO(q)$ .

(b) Moreover,  $GL(U) \times SO(q|_{M'})$  is a Levi subgroup of P, where the first factor embeds into P using the embedding  $GL(U) \to SO(q|_{U \oplus V}) \to SO(q)$ of  $(C.16.6)$ .

*Proof.* This can be shown in the same way as parts [\(b\)](#page-53-1) and [\(c\)](#page-53-2) of [C.7,](#page-52-1) using the same cocharacter.  $\Box$ 

#### <span id="page-61-0"></span>**REFERENCES**

- <span id="page-61-6"></span>[AW14] B. Antieau and B. Williams, Unramified division algebras do not always contain Azumaya maximal orders, Invent. Math. 197 (2014), 47-56.
- <span id="page-61-1"></span>[AW17]  $\_\_\_\_\_\_\$  *Prime decomposition for the index of a Brauer class, Ann. Sc. Norm. Su*per. Pisa Cl. Sci.(5) 17 (2017), 277–285.
- <span id="page-61-3"></span>[Bae] R. Baeza, Quadratic Forms over Semilocal Rings, Lecture Notes in Mathematics 655 (1978).
- <span id="page-61-2"></span>[Bas] H. Bass, K-theory and stable algebra, Publ. Math. Inst. Hautes Etudes Sci. 22 (1964), 5–60.
- <span id="page-61-7"></span>[BMR] M. Bate, B. Martin, and G. Roehrle, A geometric approach to complete reducibility, Invent. Math. 161 (2005), 177–218.
- <span id="page-62-12"></span>[BT] A. Borel and J. Tits, *Groupes réductufs*, Publ. Math. I.H.É.S.,  $27$  (1965), 55–150.
- <span id="page-62-3"></span>[B:AC1] N. Bourbaki, *Algèbre commutative*, Ch. 1 à 4, Masson, Paris 1985.
- <span id="page-62-1"></span>[CF] B. Calmès and J. Fasel, Groupes classiques, in Autour des schémas en groupes. A celebration of  $SGA3$ , vol II, Panoramas et Synthèses 46, 1-133, Société Mathématique de France, Paris, 2015.
- <span id="page-62-20"></span>[Ch] L. N. Childs, On projective modules and automorphisms of central separable algebras, Can. J. Math. 21 (1969), 44–53.
- <span id="page-62-17"></span>[C-TS] J.-L. Colliot-Thélène and A. N. Skorobogatov, The Brauer-Grothendieck group, Ergebnisse der Mathematik und ihrer Grenzgebiete, 3. Folge, 71 (2021), Springer, Cham.
- <span id="page-62-8"></span>[Co1] B. Conrad, Reductive group schemes, in Autour des schémas en groupes, vol. I, Panoramas et Synthèses 42-43, Soc. Math. France 2014.
- <span id="page-62-11"></span> $[Co2]$  , Non-split reductive groups over  $\mathbb{Z}$ , in Autour des schémas en groupes, vol II, Panoramas et Synthèses  $46$ , Soc. Math. France 2015.
- <span id="page-62-25"></span>[Co3] , Algebraic Groups II, AMS Open Math Notes, version January 26, 2024.
- <span id="page-62-4"></span>[Da] E. C. Dade, Algebraic integral representations by arbitrary forms, Mathematika 10 (1963), 96–100; A correction, Mathematika 11 (1964), 89–90.
- <span id="page-62-23"></span>[Dem] C. Demarche, Cohomologie de Hochschild non Abélienne et extensions de Faisceaux en groupes, Autours des schémas en groupes. Vol. II, 255-292. Panor. Synthèses 46 (2015) Soc. Math. France, Paris.
- <span id="page-62-22"></span>[DG] M. Demazure, P. Gabriel, Groupes algébriques, North-Holland (1970).
- <span id="page-62-19"></span>[DeM] F. R. DeMeyer, Projective modules over central separable algebras, Canadian J. Math. 21 (1969), 39–43.
- <span id="page-62-18"></span>[DI] F. R. DeMeyer and E. Ingraham, Separable Algebras over Commutative Rings, Lecture Notes in Mathematics vol. 181, Springer-Verlag, Berlin 1971.
- <span id="page-62-6"></span>[Di] I. Dias, Formas quadráticas sobre LG-anéis, Ph.D. thesis, Universidade Estadual de Campinas, Campinas, SP, Brazil (1988).
- <span id="page-62-16"></span>[EKM] R. Elman, N. Karpenko, and A. Merkurjev, The algebraic and geometric theory of quadratic forms, Amer. Math. Soc. Colloq. Publ. 56, Amer. Math. Soc., Providence, RI, 2008.
- <span id="page-62-2"></span>[EG] D. R. Estes and R. M. Guralnick, Module equivalences: Local to global when primitive polynomials represent units, J. Algebra 77 (1982), 138-–157.
- <span id="page-62-15"></span>[Ev] E. G. Evans, Krull-Schmidt and cancellation over local rings, Pacific J. Math. 46 (1973), 115–121.
- <span id="page-62-0"></span>[Fo] T. J. Ford, Separable Algebras, Graduate Studies in Mathematics 183, Amer. Math. Soc, Provident, RI, (2017).
- <span id="page-62-21"></span>[Ga] O. Gabber, Some theorems on Azumaya algebras, in: Michel Kervaire, Manuel Ojanguren (Eds.), Groupe de Brauer (Sem., les Plans-sur-Bex, 1980), in: Lect. Notes Math., vol. 844, Springer, Berlin, 1981, pp. 129–209.
- <span id="page-62-7"></span>[GPR1] S. Garibaldi, H. P. Petersson and M. L. Racine, Albert algebras over Z and other rings, Forum Math. Sigma  $11$  (2023), Paper No. 18, 38 pp.
- <span id="page-62-5"></span>[GPR2] , Albert algebras over commutative rings, New Mathematical Monographs vol. 48, Cambridge University Press 2024 (to appear).
- <span id="page-62-24"></span>[GPR3] , Solutions to the exercises from the book [\[GPR2\]](#page-62-5),  $arXiv:2406.02933$ .
- <span id="page-62-9"></span>[Gi1] P. Gille, Sur la classification des groupes semi-simples, Autour des schémas en groupes (III), Panoramas et Synthèses 47, Société Mathématique de France 2015.
- <span id="page-62-10"></span>[Gi2] P. Gille, When is a reductive group scheme linear?, Michigan Math. J. 72 (2022), 439–448.
- <span id="page-62-14"></span>[GP] P. Gille and A. Pianzola, Galois cohomology and forms of algebras over Laurent polynomial rings, Mathematische Annalen 338 (2007), 497-543.
- <span id="page-62-13"></span>[Gir] J. Giraud, *Cohomologie non-abélienne*, Springer (1970).
- <span id="page-63-3"></span> $[EGA]$  A. Grothendieck (avec la collaboration de J. Dieudonné), Eléments de Géométrie  $Alg\'ebrique$ , Publications mathématiques de l'I.H.É.S. no. 4, 8, 11, 17, 20, 24, 28, 32, 1960–1967.
- <span id="page-63-2"></span>[EGA-I] A. Grothendieck and J. Dieudonné, Éléments de géométrie algébrique: I. Le langage des schémas, Grundlehren der Mathematischen Wissenschaften 166 (2nd) ed.). Berlin; New York: Springer-Verlag, 1971.
- <span id="page-63-8"></span>[GW] U. Görtz and T. Wedhorn, *Algebraic Geometry I*, second edition, Springer Fachmedien Wiesbaden, 2020.
- <span id="page-63-7"></span>[Ha] G. Harder, Halbeinfache Gruppenschemata ¨uber Dedekindringen, Invent. Math. 4 (1967), 165-191.
- <span id="page-63-18"></span>[Ja] J.-C. Jantzen, Nilpotent Orbits in Representation Theory, in: Lie Theory, Lie Algebras and Representations. Prog. Math., vol. 228, ed. by J.-P. Anker, B. Orsted. Boston, Birkhäuser 2004.
- <span id="page-63-10"></span>[Kn1] M.-A. Knus, Algebres d'Azumaya et modules projectifs, Comment. Math. Helv. 45 (1970), 372–383.
- <span id="page-63-12"></span>[Kn2] , Quadratic and Hermitian Forms over Rings, Grundlehren der mathematischen Wissenschaften 294 (1991), Springer.
- <span id="page-63-19"></span>[KMRT] M.-A. Knus, A. Merkurjev, M. Rost and J.-P. Tignol, The Book of Involutions, Amer. Math. Soc. Colloquium Publ., 44, Amer. Math. Soc., Providence, RI (1998).
- <span id="page-63-13"></span>[KO] M.-A. Knus, M. Ojanguren, Théorie de la Descente et Algèbres d'Azumaya, Lecture Notes in Mathematics 389 (1974), Springer.
- <span id="page-63-4"></span>[Lo1] O. Loos, *Generically algebraic Jordan algebras over commutative rings*, J. Algebra 297 (2006), 474–529.
- <span id="page-63-21"></span>[Lo2]  $\qquad \qquad \qquad$ , Cubic and symmetric compositions over rings, manuscripta math. 124 (2007), 195–236.
- <span id="page-63-16"></span>[Ma] B. Margaux, Vanishing of Hochschild cohomology for affine group schemes and rigidity of homomorphisms between algebraic groups, Documenta Math. 14 (2009), 653–672.
- <span id="page-63-1"></span>[MW] B. R. McDonald and W. C. Waterhouse, Projective modules over rings with many units, Proc. Amer. Math. Soc. 83(3) (1981), 455--458.
- <span id="page-63-17"></span>[Mc] G. J. McNinch Levi decompositions of a linear algebraic group, Transf. Groups (Morozov centennial issue) 15 (2010), 937–964.
- <span id="page-63-20"></span>[Mil1] J. S. Milne, *Étale cohomology*, Princeton Mathematical Series 33 (1980), Princeton University Press, Princeton, N.J.
- <span id="page-63-6"></span>[MB] L. Moret-Bailly, Applications of local-global principles to arithmetic and geometry, Hilbert's tenth problem: relations with arithmetic and algebraic geometry (Ghent, 1999), Contemp. Math. 270 (2000), Amer. Math. Soc., 169–186.
- <span id="page-63-14"></span>[Oe] J. Oesterl´e, Sur les sch´emas en groupes de type multiplicatif, in Autour des  $sch{émas en groupes, vol. I, Panorems et Syntheses 42-43, Soc. Math. France}$ 2014.
- <span id="page-63-11"></span>[OS] M. Ojanguren and R. Sridharan, Cancellation of Azumaya algebras, J. Algebra 18 (1971), 501–505.
- <span id="page-63-9"></span>[PS] V. Petrov and A. Stavrova, The Tits indices over semilocal rings, Transform. Groups 16 (2011), 193–217.
- <span id="page-63-5"></span>[Ru] R. Rumely, Arithmetic over the ring of all algebraic integers, J. Reine Angew. Math. 368 (1986), 127–133.
- <span id="page-63-15"></span>[SGA1] A. Grothendieck, Séminaire de géométrie algébrique I, Lecture Notes in Mathematics 224, Springer-Verlag 1971; réédition as Documents Mathématiques 3, Société Mathématique de France (2011).
- <span id="page-63-0"></span>[SGA3] Séminaire de Géométrie algébrique de l'I.H.E.S., 1963-1964, schémas en groupes, dirigé par M. Demazure et A. Grothendieck, Lecture Notes in Math. 151-153. Springer (1970).
- <span id="page-64-5"></span>[Sa] C. H. Sah, Symmetric bilinear forms and quadratic forms, J. Algebra 20 (1972), 144-160.
- <span id="page-64-3"></span>[Se1] J.-P. Serre, The notion of complete reducibility in group theory, in Moursund Lectures Part II (Eugene, 1998), Notes by W.E. Duckworth, [http://darkwing.uoregon.edu/math/serre/index.html](http://darkwing.uoregon.edu/∼math/serre/index.html).
- <span id="page-64-4"></span>[Se2] , Complète réductibilité, Séminaire Bourbaki, Vol. 2003/2004, Astérisque 299 (2005), Exp. No. 932, 195–217.
- <span id="page-64-0"></span>[St] The Stacks Project Authors, Stacks project, <http://stacks.math.columbia.edu/>
- <span id="page-64-6"></span>[Sw] R. G. Swan, K-theory of quadric hypersurfaces, Ann. of Math. 122 (1985), 113– 153.
- <span id="page-64-2"></span>[Ti] J. Tits, Classification of algebraic simple groups, in "Algebraic Groups and Discontinuous Subgroups", Proc. Symp. Pure Math., 9, Amer. Math. Soc. (1966), pp. 33–62.
- <span id="page-64-1"></span> $[vdK]$  W. van der Kallen, The  $K_2$  of rings with many units, Ann. Sci. Ecole Norm. Sup. (4) 10 (1977), 473–515.

UMR 5208 DU CNRS - INSTITUT CAMILLE JORDAN - UNIVERSITÉ CLAUDE BERNARD Lyon 1, 43 boulevard du 11 novembre 1918, 69622 Villeurbanne cedex, France and Institute of Mathematics "Simion Stoilow" of the Romanian Academy, 21 Calea Grivitei Street, 010702 Bucharest, Romania

Email address: gille@math.univ-lyon1.fr

Department of Mathematics and Statistics, University of Ottawa, Ottawa, Ontario, Canada, K1N 6N5

Email address: neher@uottawa.ca