## SARD PROPERTIES FOR POLYNOMIAL MAPS IN INFINITE DIMENSION

ANTONIO LERARIO, LUCA RIZZI, AND DANIELE TIBERIO

ABSTRACT. Motivated by the Sard conjecture in sub-Riemannian geometry, and more specifically on Carnot groups, we consider the set of maps defined on a Hilbert space H:

$$\mathscr{P}_d^m(H) = \left\{ f: H \to \mathbb{R}^m \ \Big| \ \forall E \subset H, \ \dim(E) < \infty \implies f|_E \text{ is a polynomial map of degree } d \right\}$$

Example of maps belonging to this class are: the map  $f: \ell^2 \to \mathbb{R}$  constructed by Kukpa [Kup65] as a counterexample for the Sard theorem in infinite dimension; the Endpoint maps of Carnot groups.

We prove that, for infinite-dimensional subspaces  $V \subset H$  satisfying quantitative assumptions on their Kolmogorov *n*-width, the restriction of f on V has the Sard property, i.e. the set of critical values of  $f|_V$  has measure zero in  $\mathbb{R}^m$ . We prove various quantitative versions of this result, allowing to study the Sard property also for the case V = H. By constructing suitable examples, we prove the sharpness of our *n*-width assumptions.

We apply these results to sub-Riemannian geometry, proving that on Carnot groups the Sard property holds true for the restriction of the Endpoint map on the set of real-analytic controls with large enough radius of convergence. This is a more functional-analytic approach to the Sard conjecture that, unlike previous ones, does not resort to reduction to finite-dimensions.

# Contents

1. Introduction	2
1.1. The Sard property for polynomial maps on infinite-dimensional spaces	2
1.2. Quantitative semialgebraic geometry	5
1.3. Applications to Endpoint maps of Carnot groups	6
1.4. Closing thoughts on the Sard conjecture	8
1.5. Acknowledgements	9
2. Table of notations	9
3. Quantitative variations estimates	9
3.1. Semialgebraic sets and maps	9
3.2. Definable choice	11
3.3. Variations and their behaviour under polynomial maps	12
3.4. Variations of almost-critical values of polynomial maps	16
4. Sard properties for infinite-dimensional maps	
4.1. Sard-type theorem for well approximated maps	18
4.2. Sard-type theorems and Kolmogorov <i>n</i> -width	21
4.3. Counterexamples to the Sard theorem in infinite dimension	22
4.4. Sard threshold theorems	24
4.5. A class of maps with the global Sard property	26
5. Applications to the Endpoint maps of Carnot groups	30
5.1. Carnot groups	30

SISSA, VIA BONOMEA 265, 34136 TRIESTE, ITALY

*E-mail addresses*: alerario@sissa.it, lrizzi@sissa.it, dtiberio@sissa.it. *Date*: July 3, 2024.

5.2.	Endpoint maps	31
5.3.	Sard property for real-analytic controls	33
5.4.	Surjectivity of the Endpoint map on finite-dimensional spaces of controls	35
References		38

## 1. INTRODUCTION

1.1. The Sard property for polynomial maps on infinite-dimensional spaces. The classical Sard theorem [Mor39, Sar42] states that the set of critical values of a smooth map  $f : N \to M$ , where N and M are finite dimensional manifolds, has measure zero in M. Smale [Sma65] proved a version of this result in the case both M and N are infinite dimensional Banach manifolds and the map f is Fredholm: in this case the conclusion is that the set of regular values of f is residual in N.

When N is infinite-dimensional and M is finite dimensional, f cannot be Fredholm and Smale's result cannot be applied. In fact there are smooth surjective maps without regular points from any infinite-dimensional Banach space to  $\mathbb{R}^2$ , see [Bat93]. Even under the hypothesis that f is a polynomial map, the set of its critical values can be of positive measure: Kupka [Kup65] constructed an example of a smooth map  $f: \ell^2 \to \mathbb{R}$ , whose restriction to each finite dimensional space is a polynomial of degree 3, and with the property that the set of its critical values is the segment [0, 1]. On the other hand, using a quantitative version of the classical Sard theorem [Yom83], Yomdin has shown that if a map  $f: \ell^2 \to \mathbb{R}$  can be approximated well-enough with finite dimensional polynomials, still the set of its critical values has measure zero, see [Yom88]. (Kupka's map does not have this property.) In Yomdin's proof, the fact that the codomain is one-dimensional is essential and new technical difficulties arise for maps with values in  $\mathbb{R}^m$ , m > 1.

In this paper we investigate the Sard property for polynomial maps defined on Hilbert spaces and with values in  $\mathbb{R}^m$ , for general  $m \geq 1$ .

**Definition 1.** Given a Hilbert space H and  $d, m \in \mathbb{N}$ , we define the class of maps

$$\mathscr{P}_d^m(H) := \bigg\{ f: H \to \mathbb{R}^m \ \Big| \ \dim(E) < \infty \implies f|_E \text{ is a polynomial map of degree } d \bigg\},$$

where by polynomial map we mean that each component of  $f|_E$  is a polynomial of degree d. We require that each element of  $\mathscr{P}_d^m(H)$  is of class  $\mathcal{C}^1$  (in the Fréchet sense), and that its differential  $Df: H \to \mathcal{L}(H, \mathbb{R}^m)$  is weakly continuous and locally Lipschitz.

The map constructed by Kupka belongs to this family, as well as the Endpoint maps for horizontal path spaces on Carnot groups (see Section 1.3).

We need to introduce a notion of quantitative compactness for subsets of Hilbert spaces. More precisely, given  $K \subset H$  and  $n \in \mathbb{N}$ , we denote by  $\Omega_n(K, H)$  its Kolmogorov n-width:

(1) 
$$\Omega_n(K,H) := \inf_{\dim(E)=n} \sup_{u \in K} \inf_{v \in E} ||v-u||,$$

where  $\|\cdot\|$  denotes the norm of the Hilbert space H, and the infimum ranges over all vector subspaces  $E \subset H$  of dimension n (see Definition 27). A set  $K \subset H$  is compact if and only if it is bounded and its n-width goes to zero as n goes to infinity. For a compact set  $K \subset H$  we also define the quantity:

(2) 
$$\omega(K,H) := \limsup_{n \to \infty} \Omega_n(K,H)^{1/n} \in [0,1].$$

Roughly speaking, smaller  $\omega(K, H)$  means that K is "more compact" in H, and better approximated by finite-dimensional subspaces.

Our results are stated in terms of entropy dimension of the set of  $\nu$ -critical values of a map  $f: H \to \mathbb{R}^m$ , namely  $f(\operatorname{Crit}_{\nu}(f))$ , where  $\operatorname{Crit}_{\nu}(f)$  is the set of points where Df has rank at most  $\nu \leq m-1$ . The classical set of critical values corresponds to setting  $\nu = m-1$ , and it is denoted by  $f(\operatorname{Crit}(f))$ , where  $\operatorname{Crit}(f)$  is the set of critical points. We refer to Definition 25 of entropy dimension, noting that it is larger than the Hausdorff one so that in the forthcoming estimates one can replace the former with the latter for simplicity.

Our first theorem is a sufficient condition for the validity of the Sard property (see Theorem 32). We denote by  $\mu$  the Lebesgue measure on  $\mathbb{R}^m$ .

**Theorem A** (Sard under *n*-width assumptions). Let  $d, m \in \mathbb{N}$ . There exists  $\beta_0 = \beta_0(d, m) > 0$  such that the following holds. Let H be a Hilbert space,  $f \in \mathscr{P}_d^m(H)$  and  $K \subset H$  be a compact set such that

$$\omega(K,H) = \limsup_{n \to \infty} \Omega_n(K,H)^{1/n} \le q^{-1} \in (0,1).$$

Then, for every  $\nu = 1, \ldots, m-1$  we have

$$\dim_e \left( f \big( \operatorname{Crit}_{\nu}(f) \cap K \big) \right) \le \nu + \frac{\ln \beta_0}{\ln q}.$$

In particular, if  $q > \beta_0$ , then the Sard property holds on K:

$$\mu\bigg(f\big(\operatorname{Crit}(f)\cap K\big)\bigg)=0.$$

Remark 2. The constant  $\beta_0(d, m)$  of Theorem A comes from semialgebraic geometry, and its origin is discussed in Section 1.2. For this reason, we sometimes call it the *semialgebraic constant*.

By building upon Kupka's counterexamples [Kup65], we prove that the quantitative assumption  $q > \beta_0$  in Theorem A cannot be dispensed of. The following result corresponds to Theorem 36. We denote by  $B_X(r)$  the closed ball of radius r in a Banach space X.

**Theorem B** (Counterexamples to Sard). Let  $d, m \in \mathbb{N}$ , with  $d \ge 3$ , and set  $q := (d-1)^{1/d}$ . There exist a Hilbert space H, and  $f \in \mathscr{P}_d^m(H)$  such that  $K = \operatorname{Crit}(f) \cap B_H(r)$  is compact for all r > 0, with

$$\omega(K,H) = \limsup_{n \to \infty} \Omega_n(K,H)^{1/n} \le q^{-1},$$

and  $f: H \to \mathbb{R}^m$  does not verify the Sard property, namely  $\mu(f(\operatorname{Crit}(f) \cap K)) > 0$ . Therefore, the semialgebraic constant  $\beta_0(d,m)$  of Theorem A satisfies

$$\beta_0(d,m) \ge (d-1)^{1/d}, \qquad \forall m \in \mathbb{N}, d \ge 3.$$

Remark 3. The case d = 2, which is left out from Theorem B, is special. This is related to the fact that the complexity of semialgebraic sets defined by quadratic equations has a different behaviour compared to the case  $d \ge 3$ . In this case we currently expect that any  $f \in \mathscr{P}_2^m(H)$  satisfies the Sard property globally on H. This will be the subject of a future work.

As a consequence of Theorems A and B, we see that there is a quantitative threshold on the *n*-width of compact sets for the validity of the Sard property. Indeed, we can define  $\omega_0(d, m)$  as the supremum of the set of  $\omega \in [0, 1]$  such that for every  $f \in \mathscr{P}_d^m(H)$  and for every compact set  $K \subset H$  with  $\omega(K, H) = \omega$  it holds  $\mu(f(\operatorname{Crit}(f) \cap K)) = 0$ . We obtain the following statement (see Theorem 38).

**Theorem C** (Sard threshold on compacts). For all  $d, m \in \mathbb{N}$ , there exists  $\omega_0(d, m) \in (0, 1]$  such that

(i) for every  $f \in \mathscr{P}_d^m(H)$  and for every compact set  $K \subset H$  with  $\omega(K, H) < \omega_0(d, m)$ ,

$$\mu\bigg(f\left(\operatorname{crit}(f)\cap K\right)\bigg)=0;$$

(ii) for every  $\omega > \omega_0(d, m)$ , with  $\omega \in (0, 1]$ , there exist  $f \in \mathscr{P}_d^m(H)$  and a compact set  $K \subset H$  with  $\omega(K, H) = \omega$  and such that

$$\mu\bigg(f\left(\operatorname{crit}(f)\cap K\right)\bigg)>0.$$

*Remark* 4. By construction, and Theorem A,  $\omega_0 \ge \beta_0^{-1}$ . Furthermore, if  $d \ge 3$ , Theorem B yields  $\omega_0 < 1$ , so that Item (ii) is non-vacuous in these cases.

We can strengthen the conclusions from Item (i) of Theorem C as follows (see Theorem 39).

**Theorem D** (Sard threshold on linear subspaces). Let  $f \in \mathscr{P}_d^m(H)$  and  $K \subset H$  be a compact set such that  $\omega(K, H) < \omega_0(d, m)$ . Consider the linear subspace

$$V := \operatorname{span}(K).$$

Then the restriction  $f|_V: V \to \mathbb{R}^m$  satisfies the Sard property:

$$\mu\bigg(f\big(\operatorname{Crit}(f|_V)\big)\bigg) = 0$$

In particular,  $\mu(f(\operatorname{Crit}(f) \cap V)) = 0.$ 

Remark 5. Since  $V \subset H$  is a (possibly non-closed) linear subspace, the restriction  $f|_V : V \to \mathbb{R}^m$  is a  $\mathcal{C}^1$  map in the Fréchet sense from the normed vector space  $(V, \|\cdot\|)$  to  $\mathbb{R}^m$ , with  $D_u(f|_V) = (D_u f)|_V$  for all  $u \in V$ .

Example 6. Let  $\eta \in (1, \infty)$  with  $\eta > \omega_0(d, m)^{-1}$ , where  $\omega_0(d, m)$  is the number in Theorems C and D, for  $d, m \in \mathbb{N}$ . Let  $H = \ell^2$  with the usual norm. Consider the subset  $K \subset B_H(1)$  given by

$$K = \left\{ u \in \ell^2 \left| \sum_{j=1}^{\infty} |u_j|^2 \eta^{2j} \le 1 \right\}.$$

One can easily verify that K is compact. Furthermore, the linear space  $V = \operatorname{span}(K)$  is dense in H. We can estimate the *n*-width of K as follows. For  $n \in \mathbb{N}$  consider the *n*-dimensional subspaces  $E_n = \{x \in \ell^2 \mid x_j = 0, \forall j \ge n+1\} \subset H$ . From (1), we obtain

$$\Omega_n(K,H)^2 \le \sup_{u \in K} \inf_{v \in E_n} ||u - v||^2$$
  
=  $\sup_{u \in K} \sum_{j=n+1}^{\infty} |u_j|^2$   
 $\le \eta^{-2(n+1)} \sup_{u \in K} \sum_{j=1}^{\infty} |u_j|^2 \eta^{2j} \le \eta^{-2(n+1)}$ 

Hence  $\omega(K, H) = \limsup_n \Omega_n(K, H)^{1/n} \le \eta^{-1} < \omega_0(d, m)$ . Hence, by Theorem D it holds

$$\mu\bigg(f\big(\operatorname{Crit}(f|_V)\big)\bigg) = \mu\bigg(f\big(\operatorname{Crit}(f) \cap V\big)\bigg) = 0, \qquad \forall f \in \mathscr{P}_d^m(H)$$

Note that even if  $V \subset H$  is dense, the unrestricted map  $f : H \to \mathbb{R}^m$  may not have the Sard property (see e.g. the Kupka counterexamples in Section 4.3).

Theorems A, C and D are deduced from a more general result for maps that are "well-approximated" by polynomials. This can be regarded as our main result concerning the Sard property, and does not make use of the concept of n-width. We report here the statement (see Theorem 26).

**Theorem E** (Sard criterion for well-approximated maps). Let  $d, m \in \mathbb{N}$ . There exists a constant  $\beta_0 = \beta_0(d, m) > 0$  such that the following holds. Let H be a Hilbert space, and let  $f : H \to \mathbb{R}^m$  be a  $C^1$  map such that its differential  $Df : H \to \mathcal{L}(H, \mathbb{R}^m)$  is weakly continuous. Let  $K \subset H$  be a bounded set with this approximation property: there exist a sequence  $E_n \subset H$  of linear subspaces,  $\dim(E_n) = n$ , and polynomial maps  $f_n : E_n \to \mathbb{R}^m$  with uniformly bounded degree:

$$\sup_{n\in\mathbb{N}} \deg f_n \le d < \infty,$$

such that for some q > 1,  $c \ge 0$ , and all large enough n it holds

(3) 
$$\sup_{x \in K} \left( \|f(x) - f_n \circ \pi_{E_n}(x)\| + \|(D_x f)|_{E_n} - D_{\pi_{E_n}(x)} f_n\|_{\mathrm{op}} \right) \le cq^{-n}.$$

Then

$$\dim_e \left( f \big( \operatorname{Crit}_{\nu}(f) \cap K \big) \right) \le \nu + \frac{\ln \beta_0}{\ln q}, \qquad \forall \nu = 1, \dots, m-1.$$

In particular, if  $q > \beta_0$ , then f satisfies the Sard property on K:

$$\mu\bigg(f(\operatorname{Crit}(f)\cap K)\bigg)=0.$$

Remark 7. The constant  $\beta_0(d, m)$  of Theorem E is the same appearing in Theorem A, and comes from semialgebraic geometry, see Section 1.2.

By means of the general Theorem E we single out a class of maps satisfying the Sard property on the whole domain of definition. This is the content of Theorem 43, that we report here. The case m = 1 corresponds to [YC04, Thm. 10.12].

**Theorem F** (Global Sard for special maps). Let H be a separable Hilbert space. For all  $k \in \mathbb{N}$ , let  $p_k : E_k \to \mathbb{R}^m$  be polynomial maps with  $\sup_{k \in \mathbb{N}} \deg p_k \leq d$  for some  $d \in \mathbb{N}$ , and such that

$$\sup_{x \in B_{E_k}(1)} \|p_k(x)\| \le q^{-k}, \qquad \forall k \in \mathbb{N},$$

for some q > 1. Then the map  $f : H \to \mathbb{R}^m$  defined by

$$f(x) := \sum_{k=1}^{\infty} p_k(x_1, \dots, x_k), \qquad \forall x \in H,$$

is well-defined,  $f \in \mathscr{P}_d^m(H)$  (see Definition 1), and for all  $\nu = 1, \ldots, m-1$  and r > 0 it holds

$$\dim_e \left( f\left(\operatorname{Crit}_{\nu}(f) \cap B_H(r)\right) \right) \leq \nu + \frac{\ln \beta_0}{\ln q},$$

where  $\beta_0 = \beta_0(d, m)$  is the same constant given by Theorem E. In particular, if  $q > \beta_0$ , then f satisfies the Sard property globally on H:

$$\mu\bigg(f\big(\operatorname{Crit}(f)\big)\bigg) = 0.$$

1.2. Quantitative semialgebraic geometry. The proof of Theorem E needs some fine properties of semialgebraic sets. The strategy of the proof is conceptually similar to Yomdin's [Yom88], and uses the theory of *variations*, introduced by Vitushkin [Vit55, Vit61] and developed in [YC04].

Let us denote with  $V_i(S)$  the *i*-th variation of a semialgebraic set, which is a sort of *i*-dimensional volume, (see Definition 15). Furthermore, recall that for a  $C^1$  map  $F : \mathbb{R}^n \to \mathbb{R}^m$ , and  $\Lambda = (\Lambda_1, \ldots, \Lambda_m) \in \mathbb{R}^m_+$  the set of almost-critical values of F is defined by

$$C^{\Lambda}(F) := \left\{ x \in \mathbb{R}^n \, \middle| \, \sigma_i(D_x F) \le \Lambda_i, \quad \forall \, i = 1, \dots, m \right\},\$$

where  $\sigma_1(D_xF) \geq \cdots \geq \sigma_m(D_xF)$  are the singular values of  $D_xF$ , see Definition 17 (here we assume  $n \geq m$ ). In [YC04, Cor. 7.4] a quantitative estimate on the variations of the almost-critical values of polynomial maps has been obtained. In that estimate the dimensional parameter n does not appear explicitly. We obtain Theorem G below, which makes this dependence explicit (see Theorem 24).

**Theorem G** (Quantitative variations estimates). Let  $n \ge m$ , and  $p : \mathbb{R}^n \to \mathbb{R}^m$  be a polynomial map with components of degree at most d. For i = 0, ..., m,  $\Lambda = (\Lambda_1, ..., \Lambda_m) \in \mathbb{R}^m_+$  and r > 0, we have

$$V_i(p(C^{\Lambda}(p) \cap B_{\mathbb{R}^n}(r))) \leq \operatorname{cst}(m,r)n^m \beta_0^n \Lambda_0 \cdots \Lambda_i$$

where  $\beta_0 = \beta_0(d, m)$  depends only on d and m,  $\operatorname{cst}(m, r)$  depends only on m, r, and we set  $\Lambda_0 = 1$ .

A key technical step in the proof of Theorem G is a quantitative substitute for the *definable choice* theorem in semialgebraic geometry, which we prove in [LRT24] (see Theorem 14).

1.3. Applications to Endpoint maps of Carnot groups. We discuss now the implications of Theorem C and Theorem D in the context of sub-Riemannian geometry.

Recall that an *m*-dimensional *Carnot group* of step  $s \in \mathbb{N}$  is a connected and simply connected Lie group  $(\mathbb{G}, \cdot)$  of dimension *m*, whose Lie algebra  $\mathfrak{g}$  admits a stratification of step *s*, that is

$$\mathfrak{g} = \mathfrak{g}_1 \oplus \cdots \oplus \mathfrak{g}_s$$

where  $\mathfrak{g}_i \neq \{0\}$ ,  $\mathfrak{g}_{i+1} = [\mathfrak{g}_1, \mathfrak{g}_i]$  for all  $i = 1, \ldots, s-1$  and  $[\mathfrak{g}_1, \mathfrak{g}_s] = \{0\}$ . The group exponential map yields an identification  $\mathbb{G} \simeq \mathbb{R}^m$ , and the first stratum  $\mathfrak{g}_1$  of the Lie algebra defines a smooth, totally non-holonomic distribution  $\Delta \subseteq T\mathbb{R}^m$  of rank  $k := \dim \mathfrak{g}_1$ .

Fix a global trivialization of  $\Delta$ :

$$\Delta = \operatorname{span}\{X_1, \ldots, X_k\},\$$

where each  $X_i$  is a left-invariant vector field. Let I := [0, 1] be the unit interval and  $H := L^2(I, \mathbb{R}^k)$ . We say that a curve  $\gamma : I \to \mathbb{G}$  is *horizontal* if it is absolutely continuous and there exist  $u \in H$ , called *control* such that for a.e.  $t \in I$  it holds

(4) 
$$\dot{\gamma}(t) = \sum_{i=1}^{k} u_i(t) X_i(\gamma(t)).$$

Furthermore, for any given  $u \in H$  there exists a unique  $\gamma_u : I \to \mathbb{G}$  satisfying (4) and such that  $\gamma_u(0) = e$  (the identity  $e \in \mathbb{G}$ , identified with  $0 \in \mathbb{R}^m$ ). Note that the class of horizontal curves does not depend on the choice of the global trivialization of  $\Delta$ .

The Endpoint map is the map that sends a control u to the the corresponding final point  $\gamma_u(1)$ :

$$\mathcal{E}: L^2(I, \mathbb{R}^k) \to \mathbb{G}.$$

The Sard conjecture is the conjecture that Endpoint map has the Sard property:

Sard conjecture: the set  $\mathcal{E}(\operatorname{Crit}(\mathcal{E}))$  has zero measure.

The conjecture was introduced by Zhitomirskii and Montgomery, see [Mon02, Sec. 10.2] for general sub-Riemannian structures, where the Endpoint map can be defined in a similar way, on a suitable domain of horizontal paths, which has the structure of a Hilbert manifold. For the specific case of Carnot groups, it is known to be true for step  $\leq 2$ , see [AGL15, LDMO<sup>+</sup>16]. Furthermore, the conjecture has been verified in [BNV22] for filiform Carnot groups, in [BV20] for Carnot groups of rank 2 and step 4, and rank 3 and step 3, and for a handful of other specific examples described in [LDMO<sup>+</sup>16].

Assume that  $\Delta$  is also endowed with a left-invariant norm. This choice induces a length structure on the space of horizontal curves, and correspondingly, a distance, which gives  $\mathbb{G}$  the structure of *sub-Finsler Carnot group* (or *sub-Riemannian Carnot group* if the norm is induced by a scalar product). In this case, within the set critical points of  $\mathcal{E}$  (also called singular horizontal paths), the energy-minimizing ones play a significant role. At the corresponding critical values, the distance loses regularity: it is never smooth and it can even lose local semiconcavity [BR19, Sec. 4.2]. For this reason, the understanding of the "abundance" of such critical values is crucial. Denoting with  $H_{\min} \subseteq L^2(I, \mathbb{R}^k)$  the set of minimizing horizontal paths starting from the identity, the *minimizing Sard conjecture* can be formulated as follows:

## Minimizing Sard conjecture: the set $\mathcal{E}(\operatorname{Crit}(\mathcal{E}) \cap H_{\min})$ has zero measure.

The most general result to date is that the above set is a closed nowhere dense set [Agr09, RT05], which of course does not imply the minimizing Sard conjecture. The above conjecture is one of the main open problems in sub-Riemannian geometry [Agr14, Prob. 3], [RT05, Conj. 1]. In general, the problem is settled in the following cases:

- Thanks to the work of Agrachev-Sarychev [AS96] and Agrachev-Lee [AL09], in absence of socalled Goh singular minimizing paths, all sub-Riemannian minimizing paths (resp. sub-Finsler, under suitable smoothness assumptions on the norm) are solution of a finite-dimensional Hamiltonian flow so that the corresponding set of critical values has zero measure by the finite-dimensional Sard theorem. Recently, Rifford proved that, more in general, the minimizing Sard conjecture holds for structures where all non-trivial Goh paths have Goh-rank  $\leq 1$  almost everywhere, see [Rif23]. For Carnot groups, these results can be applied for example when the distribution is *pre-medium fat*, see [Rif23, (1.2)]. In passing, we remark that for generic (so, typically not Carnot) sub-Riemannian structures with distribution of rank  $\geq 3$ , there are no non-trivial singular Goh minimizing paths, see [CJT06, Cor. 2.5] and [AG01, Thm. 8]. As a consequence the minimizing Sard conjecture holds true. Thanks to [Rif23, Cor. 1.4], the latter result extends the case of rank  $\geq 2$ .

- The minimizing Sard conjecture is true also for Carnot groups of step  $\leq 3$ , see [LDMO<sup>+</sup>16]. In this case, the problem is reduced (in a non-trivial way) to a finite-dimensional one. First, by noting that singular minimizing curves are non-singular in some proper subgroup, and then exploiting the fact that Carnot subgroups are parametrized by a finite-dimension manifold. Also in this case, the Goh condition plays a fundamental role.

Finally, we refer to [ZZ95, BdSR18, BdSFPR22, BdSPR23, BdSPR22, OV19, RT05] for further details and discussions of various forms of the Sard conjecture on general sub-Riemannian structures.

The following result (Theorem 46) connects this framework to the previous sections.

**Theorem H** (Polynomial properties of the Endpoint map). Let  $\mathbb{G}$  be a Carnot group of topological dimension m, step s, and rank k. Then the Endpoint map  $\mathcal{E} \in \mathscr{P}_s^m(H)$ , for  $H = L^2(I, \mathbb{R}^k)$ 

As a consequence, we can apply Theorem D to the study of Endpoint maps of Carnot groups. We record here a first immediate corollary (see Theorem 47).

**Theorem I** (Sard criterion for Endpoint maps). Let  $\mathbb{G}$  be a Carnot group of topological dimension m, step s, and rank k. Let  $K \subset L^2(I, \mathbb{R}^k)$  be a compact subset with

$$\omega(K, L^2(I, \mathbb{R}^k)) < \omega_0(m, s).$$

Then, letting  $V := \operatorname{span}(K)$ , the restriction  $\mathcal{E}|_V : V \to \mathbb{G}$  has the Sard property, namely

$$\mu\Big(\mathcal{E}\big(\operatorname{Crit}(\mathcal{E}|_V)\big)\Big) = 0.$$

In particular  $\mu(\mathcal{E}(\operatorname{Crit}(\mathcal{E}) \cap V)) = 0.$ 

We can apply Theorem I to find large sets on which the Sard property holds for the Endpoint map of Carnot groups. This is a more functional-analytic approach to the Sard conjecture that, unlike previous ones, does not resort to reduction to finite-dimensional cases.

An interesting case in which the assumptions of Theorem I can be effectively checked is given by sets of controls that are "sufficiently regular". We introduce the notation to state our result, see Section 5.3 for additional details. Given r > 0, and a closed interval  $I \subset \mathbb{R}$ , consider the set:

$$\mathcal{C}^{\omega}(I, \mathbb{R}^k; r) := \bigg\{ u : I \to \mathbb{R}^k \, \Big| \, u \text{ is real-analytic with radius of convergence} > r \bigg\},$$

endowed with the sup norm over the *r*-neighbourhood of *I* in  $\mathbb{C}$ . More in general, let  $\mathcal{C}^{\omega}(I, \mathbb{R}^k; r, \ell)$  be the set of piecewise analytic controls:

$$\mathcal{C}^{\omega}(I,\mathbb{R}^k;r,\ell) = \bigg\{ u: I \to \mathbb{R}^k \ \bigg| \ u|_{I_j} \in \mathcal{C}^{\omega}(I_j,\mathbb{R}^k;r), \text{ for all } j = 1,\dots,\ell \bigg\},\$$

where  $I_j = \inf I + \left[\frac{(j-1)|I|}{\ell}, \frac{j|I|}{\ell}\right]$ , for all  $j = 1, \ldots, \ell$ , which we equip with a suitable norm (see Section 5.3). In Theorem 48 we prove that the unit ball K of  $\mathcal{C}^{\omega}(I, \mathbb{R}^k; r, \ell)$ , with I = [0, 1], is compact in H and its *n*-width satisfies:

$$\Omega_n(K,H) \le \frac{(k\ell)^{1/2}}{\ln r} \left(\frac{1}{r}\right)^{\lfloor \frac{n}{k\ell} \rfloor}, \qquad \forall n \in \mathbb{N}, \quad r > 1.$$

By applying Theorem I we can therefore deduce the following result (see Theorem 49).

**Theorem J** (Sard property on piecewise real-analytic controls). Let  $\mathbb{G}$  be a Carnot group of topological dimension m, step s, and rank k. Given  $\ell \in \mathbb{N}$ , there exists  $r = r(m, s, k, \ell) > 0$  such that, letting  $V = \mathcal{C}^{\omega}(I, \mathbb{R}^k; r, \ell)$ , with I = [0, 1], it holds

$$\mathcal{E}(V) = \mathbb{G}$$
 and  $\mu(\mathcal{E}(\operatorname{Crit}(\mathcal{E}|_V)) = \mu(\mathcal{E}(\operatorname{Crit}(\mathcal{E}) \cap V)) = 0.$ 

Namely, the Sard property holds on the space of piecewise real-analytic controls with radius of convergence > r and with  $\ell$  pieces.

*Remark* 8 (Regularity and Sard). Sussmann established that for real-analytic sub-Riemannian structures (such as Carnot groups), minimizing horizontal paths are real-analytic on an open and dense set of their interval of definition [Sus14]. In light of that result, Theorem J hints at a unexpected link between the regularity problem of sub-Riemannian geodesics and the minimizing Sard conjecture.

Remark 9 (Surjectivity of the Endpoint map). Of course a subset of controls  $V \subset L^2(I, \mathbb{R}^k)$  must satisfy  $\mathcal{E}(V) = \mathbb{G}$  to be relevant for the theory. By the Rashevskii-Chow theorem,  $\mathcal{E} : L^2(I, \mathbb{R}^k) \to \mathbb{R}^m$ is surjective. We observe that  $\mathcal{E}$  is surjective also on  $\mathcal{C}^{\omega}(I, \mathbb{R}^k; r, \ell)$  if  $\ell \geq 4m$ , since from the proof of the Rashevskii-Chow theorem any pair of points can be joined by a concatenation of 4m horizontal paths with constant controls, see [ABB20, Sec. 3.2]. Actually, the Endpoint map is surjective when restricted on any set that is dense in  $L^2(I, \mathbb{R}^k)$ , see e.g. [Bel96, Sec. 2.5]. It follows that  $\mathcal{E}$  is surjective on  $\mathcal{C}^{\omega}(I, \mathbb{R}^k; r)$ , since the latter contains all (restrictions on I of) polynomial controls.

Finally, we prove in Theorem 53 that the Endpoint map on Carnot groups is surjective already when restricted to polynomial controls of large enough degree (depending on  $\mathbb{G}$ ). This is a particular case of the following more general result (see Theorem 54).

**Theorem K** (Quantitative surjectivity). Let  $\mathbb{G}$  be a Carnot group with topological dimension m and rank k. Let  $S \subset L^2(I, \mathbb{R}^k)$  be a dense set. Then there exist  $u_0, u_1, \ldots, u_m \in S$  such that

$$\mathcal{E}(\operatorname{span}\{u_0, u_1, \dots, u_m\}) = \mathbb{G}.$$

In other words, one can find an (m + 1)-dimensional vector subspace  $E \subset L^2(I, \mathbb{R}^k)$  of controls (depending on G) such that the restriction of the Endpoint map to E is surjective. Furthermore, E can be assumed to have generators in any prescribed dense set  $S \subset L^2(I, \mathbb{R}^k)$ . Theorem K can be seen as a more refined version (for Carnot groups) of the well-known fact that the Endpoint map is surjective when restricted on any set of controls that is dense in  $L^2(I, \mathbb{R}^k)$ , see e.g. [Bel96, Sec. 2.5].

1.4. Closing thoughts on the Sard conjecture. We do not venture in a guess in favor of Sard conjecture on the whole  $H = L^2(I, \mathbb{R}^k)$ , instead we propose a direction of future investigation that can be approached with our techniques. Recall that  $H_{\min} \subset H$  is the set of energy-minimizing controls. It is known that  $H_{\min}$  is boundedly compact in H i.e., the intersection  $H_{\min} \cap B_H(r)$  is compact for all r > 0, see [ABB20, Thm. 8.66], [Agr98]. In fact, from the very recent result by Lokutsievskiy and Zelikin [LZ23, Cor. 1 ( $\mathfrak{G}$ )], one can deduce the following estimate for the *n*-width on Carnot groups

(5) 
$$\Omega_{2n+1}(H_{\min} \cap B_H(1), H) \le Cn^{-\frac{1}{2s}}, \quad \forall n \in \mathbb{N},$$

for some C > 0 depending on G. Estimate (5) can be understood as a quantitative compactness property for  $H_{\min}$ . Unfortunately, the polynomial decay (w.r.t. n) of (5) is too weak to apply Theorem A, since the latter requires an exponential one.

If one could prove that, for some  $q^{-1} < \omega_0(m, s)$ , it holds

(6) 
$$\Omega_n(H_{\min} \cap B_H(1), H) \le q^{-n}, \quad \text{as } n \to \infty$$

then Theorem C would settle the minimizing Sard conjecture on Carnot groups. At present, however, producing an estimate better than (5) requires a finer analysis than the one in [LZ23].

We also note that it would be sufficient to prove (6) for the set of the so-called *strictly abnormal* energy-minimizing controls  $H_{\min}^{\text{str.abn}} \subsetneq H_{\min}$ . Unfortunately, we were not able to produce a direct estimate of the *n*-width of (bounded subsets of)  $H_{\min}$  or  $H_{\min}^{\text{str.abn}}$ , so that we have no further evidence to support this idea towards a proof of the minimizing Sard conjecture.

In the opposite direction, and to conclude, our results can be used to identify spaces where the (general) Sard conjecture can be violated. If, for a given Carnot group  $\mathbb{G}$ , there is a linear subset of controls  $V \subset H$  such that  $\mu(\mathcal{E}(\operatorname{Crit}(\mathcal{E}) \cap V)) > 0$ , then the following necessary condition must hold:

$$\limsup_{n \to \infty} \Omega_n (V \cap B_H(r), H)^{1/n} \ge \omega_0(m, s).$$

1.5. Acknowledgements. This project has received funding from (i) the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 945655); (ii) the PRIN project "Optimal transport: new challenges across analysis and geometry" funded by the Italian Ministry of University and Research; (iii) the Knut and Alice Wallenberg Foundation. The authors also acknowledge the INdAM support. The authors also wish to thank Yosef Yomdin for helpful discussions that improved the presentation of the paper.

# 2. TABLE OF NOTATIONS

$\mathbb{R}^m_+$	<i>m</i> -tuples of positive real numbers
$B_X(x,r)$	closed ball of radius $r > 0$ centered at x of a Banach space X
$B_X(r)$	closed ball of radius $r > 0$ centered at 0 of a Banach space X
$B_X$	closed unit ball centered at 0 of a Banach space $X$
$\mathcal{L}(X,Y)$	linear continuous maps between banach spaces $X, Y$
·    <sub>op</sub> · · · · · · · · · · · · ·	operator norm on $\mathcal{L}(X,Y)$ , see (23)
$\ \cdot\ _X$	norm for a Banach space $X$ (the $X$ is omitted when there is no confusion)
$\mathcal{B}(X,Y)$	continuous bounded maps between Banach spaces $X, Y$
$\mathcal{C}^{\omega}(I,\mathbb{R}^k;r,\ell)$	piece-wise real-analytic functions, Section 5.3
$\mu$	Lebesgue measure
$\Omega_n(K,H)$	Kolmogorov <i>n</i> -width of $K \subset H$ , Definition 27
$\omega(K,H)$	asymptotic Kolmogorov width of $K \subset H$ , Eq. (2)
$\mathcal{P}_d^m(H)$	set of well-approximable maps, Definition 1
$\beta_0(d,m)$	semialgebraic constant, Remark 2
$\omega(d,m)$	threshold constant, Theorem C
$\mathcal{H}^i(A)$	i-dimensional Hausdorff measure of a rectifiable Borel set $A$
$V_i(A)$	Vitushkin variation of a bounded semialgebraic set $A$ , Definition 15
$b_0(S)$	number of connected components of a set $S$
$\sigma_k(L)$	$k\text{-th}$ singular value of a linear map $L:H\to \mathbb{R}^m$ on a Hilbert space, ordered
	in decreasing order $\sigma_1(M) \geq \cdots \geq \sigma_m$ , Definition 17
$\operatorname{Crit}(f)$	critical points of a map, Eq. (22)
$\operatorname{Crit}_{\nu}(f)$	$\nu$ -critical points of a map, Eq. (22)
$C^{\Lambda}(f)$	$\Lambda$ -critical (or almost-critical) points of a map, Definition 20
$M(\epsilon, S)$	$\epsilon$ -entropy of a set $S \subset \mathbb{R}^m$ , Definition 25
$\dim_e(S)$	entropy dimension of a set $S \subset \mathbb{R}^m$ , Definition 25
$\mathcal{U}_{\varepsilon}(S)$	$\epsilon$ -neighbourhood of a subset $S \subset \mathbb{R}^n$
$\operatorname{dist}_H$	Hausdorff distance
$\beta_{\rm tm}$	Thom-Milnor bound constant, Theorem $13$
$\beta_{\rm dc}$	definable choice constant, Theorem 14
$\beta_{\rm yc}$	Yomdin-Comte constant, Theorem 19
D(S)	diagram of a semialgebraic set $S$ , Definition 11
$\mathbb{G}$	Carnot group (of step $s$ , rank $k$ , dimension $m$ ), Section 5.1
<i>E</i>	Endpoint map, Definition $45$

# 3. QUANTITATIVE VARIATIONS ESTIMATES

3.1. Semialgebraic sets and maps. The goal of this section is to prove Theorem G. We begin by recalling some basic notions from semialgebraic geometry.

**Definition 10** (Semialgebraic sets and maps). We say that a set  $S \subset \mathbb{R}^n$  is semialgebraic if

(7) 
$$S = \bigcup_{i=1}^{a} \bigcap_{j=1}^{b_i} \{x \in \mathbb{R}^n \mid \operatorname{sign}(p_{ij}(x)) = \sigma_{ij}\},$$

for some finite set of polynomials  $p_{ij} \in \mathbb{R}[x_1, \ldots, x_n]$  and  $\sigma_{ij} \in \{0, +1, -1\}$ , where

$$\operatorname{sign}(r) := \begin{cases} +1 & r > 0\\ -1 & r < 0\\ 0 & r = 0 \end{cases}$$

We call the description (7) a representation of S.

A map  $f : A \to B$  between semialgebraic sets  $A \subset \mathbb{R}^n, B \subset \mathbb{R}^\ell$  is said to be semialgebraic if its graph is a semialgebraic set in  $\mathbb{R}^n \times \mathbb{R}^\ell$ .

The representation of a semialgebraic set S as in (7) is not unique. However, a representation is useful to quantify the complexity of S, using the following notion.

**Definition 11** (Diagram of a semialgebraic set). Let  $S \subset \mathbb{R}^n$  be a semialgebraic set represented as in (7). We will say that the triple

$$\left(n, a \cdot \max_{i} \{b_i\}, \max_{i,j} \{\deg(p_{ij})\}\right) \in \mathbb{N}^3$$

is a diagram for S. Below, the equation "D(S) = (m, c, d)" will mean that there exists a representation of S as in (7) with  $n \le m$ ,  $a \cdot \max_i \{b_i\} \le c$  and  $\max_{i,j} \{\deg(p_{ij})\} \le d$ .

Remark 12. There are several ways of quantifying the complexity of a semialgebraic set, depending on the way it is presented. Essentially, these notions should contain three pieces of information: the number of variables, the "combinatorics" of the presentation, and the degrees of the defining polynomials. For instance, every semialgebraic set  $S \subset \mathbb{R}^n$  can be presented as

$$S = \bigcup_{i=1}^{a} \bigcap_{j=1}^{b_i} E_{ij}$$

where each  $E_{ij}$  has the form  $\{x \in \mathbb{R}^n \mid p_{ij}(x) \leq 0\}$  or  $\{x \in \mathbb{R}^n \mid p_{ij}(x) < 0\}$ , where  $p_{ij}$  is a polynomial with  $\deg(p_{ij}) \leq d$  for all i, j. Given a presentation of this type, it is immediate to see that S also admits a diagram D(S) = (n, c, d) (i.e. it can be presented as in (7) with  $c = c(a, b_1, \ldots, b_a)$ .

3.1.1. Dimension and stratifications. Every semialgebraic subset of  $\mathbb{R}^n$  can be written as a finite union of semialgebraic sets, each of them semialgebraically homeomorphic to an open cube  $(0,1)^m \subset \mathbb{R}^m$ , for some  $m \leq n$  ([BCR98, Thm. 2.3.6]). This allows to define the *dimension* of a semialgebraic set as the maximum of the dimensions m of these cubes. Every semialgebraic set can be written as a finite union of smooth, semialgebraic disjoint manifolds called *strata* ([BCR98, Prop. 9.1.8]).

Notice that the dimension of a semialgebraic set is preserved by semialgebraic homeomorphisms and behaves naturally under product structures:

$$\dim(A \times B) = \dim(A) + \dim(B).$$

Moreover, if  $f: A \to B$  is a continuous semialgebraic map, then

$$\dim(f(A)) \le \dim(A).$$

3.1.2. Semialgebraic triviality. Continuous semialgebraic maps  $f : A \to B$  are "piecewise" trivial fibrations: there exists a partition of B into finitely many semialgebraic sets

$$B = \bigsqcup_{j=1}^{b} B_j$$

and, for every j = 1, ..., b there exist fibers  $F_j := f^{-1}(b_j)$ , with  $b_j \in B_j$  and a semialgebraic homeomorphism  $\psi_j : f^{-1}(B_j) \to B_j \times F_j$  that make the following diagram commutative:



This result is called *semialgebraic triviality*, see [BPR06, Thm. 5.46].

3.1.3. Projections. The image of a semialgebraic set  $S \subset \mathbb{R}^n \times \mathbb{R}^\ell$  under the projection map  $\pi : \mathbb{R}^n \times \mathbb{R}^\ell \to \mathbb{R}^n$  is semialgebraic ([BCR98, Thm. 2.2.1]). Moreover, it follows from [BPR06, Thm. 14.16, Notation 8.2]<sup>1</sup> that there exists  $C_1 > 0$  such that if  $(n + \ell, c, d)$  is a diagram for S, we can write

$$\pi(S) = \bigcup_{i \in I} \bigcap_{j \in J_i} \bigcup_{k \in N_{ij}} \left\{ x \in \mathbb{R}^n \mid \operatorname{sign}(p_{ijk}(x)) = \sigma_{ijk} \right\},\$$

with

$$#I \le c^{(n+1)(\ell+1)} d^{C_1 \ell n}, \quad #J_i \le c^{\ell+1} d^{C_1 \ell}, \quad #N_{ij} \le d^{C_1 \ell}$$

and with

$$\deg(p_{ijk}) \le d^{C_1\ell}.$$

In particular, there exists  $C_2 > 0$  such  $\pi(S)$  admits a diagram with:

$$D(\pi(S)) = \left(n, (cd)^{C_2\ell n} (d^{C_2\ell})^{(cd)^{C_2\ell}}, d^{C_2\ell}\right).$$

In particular, the diagram of the projection of a semialgebraic set is controlled explicitly by the diagram of the original set.

3.1.4. *Thom-Milnor bound*. We will need the following qualitative bound on the number of connected components of a semialgebraic set, that follows from [BPR06, Thm. 7.50].

**Theorem 13** (Thom-Milnor). Let  $c, d \in \mathbb{N}$ . There exists a constant  $\beta_{tm} = \beta_{tm}(c, d) > 1$ , such that for any semialgebraic set  $S \subset \mathbb{R}^n$  with diagram D(S) = (n, c, d) it holds

$$b_0(S) \le \beta_{\mathrm{tm}}^n$$

where  $b_0(S)$  denotes the number of connected components of S.

3.2. **Definable choice.** We recall from [LRT24] the following *quantitative approximate definable choice* result, which will be a key tool in the proof of Theorem G.

Let us explain the name and the context of this result. Given a polynomial map  $F : \mathbb{R}^n \to \mathbb{R}^\ell$ and a compact semialgebraic set  $A \subset \mathbb{R}^n$ , for every  $y \in F(A)$  one would like to choose a point  $x(y) \in F^{-1}(y) \cap A$  so that the resulting set  $C := \{x(y)\}_{y \in F(A)}$  is semialgebraic and with controlled dimension. In particular, the "choice" of such x(y) has to be "definable". In this way C would be of dimension  $\ell$ . The fact that this can be done follows from semialgebraic triviality (see Section 3.1.2). However, in this way there is no control on the geometry of C (in particular, the number of its connected components) in terms of the data defining A (namely, its diagram, see Definition 11).

For many geometric applications (as in this paper) one does not really need that C is a choice over A, but it is enough that it is close to it. More precisely, given  $\epsilon > 0$ , denote by  $\mathcal{U}_{\epsilon}(A)$  the Euclidean  $\epsilon$ -neighbourhood of A. Then one can relax the requirements for the definable choice and ask, given  $\epsilon > 0$ , for a set  $C_{\epsilon} \subset \mathcal{U}_{\epsilon}(A)$ , of dimension  $\leq \ell$ , whose image under F is  $\epsilon$ -close to F(A) (in the Hausdorff metric, denoted by dist<sub>H</sub>), and with a control the geometry of  $C_{\epsilon}$ .

<sup>&</sup>lt;sup>1</sup>Note that the role of k and  $\ell$  for us is swapped with respect to [BPR06, Thm. 14.16]: we eliminate  $\ell$  variables from a list of  $n + \ell$ , they eliminate k variables from a list of  $k + \ell$ .

**Theorem 14** (Quantitative approximate definable choice [LRT24]). For every  $c, d, \ell \in \mathbb{N}$  there exists  $\beta_{dc} > 1$  satisfying the following statement. Let  $n \in \mathbb{N}$  and let  $A \subset \mathbb{R}^n$  be a closed semialgebraic set contained in the ball  $B_{\mathbb{R}^n}(r)$  with diagram

$$D(A) = (n, c, d).$$

Let also  $F = (F_1, \ldots, F_\ell) : \mathbb{R}^n \to \mathbb{R}^\ell$  be a polynomial map with components of degree bounded by d. Then for every  $\epsilon \in (0, r)$  there exists a closed semialgebraic set  $C_\epsilon \subset \mathbb{R}^n$  such that

- (i) dim $(C_{\epsilon}) \leq \ell$ ;
- (*ii*)  $C_{\epsilon} \subset \mathcal{U}_{\epsilon}(A)$ ;
- (*iii*) dist<sub>H</sub>( $F(C_{\epsilon}), F(A)$ )  $\leq L(F, r) \cdot \epsilon$ , where  $L(F, r) := 3 + \operatorname{Lip}(F, B_{\mathbb{R}^n}(2r))$ ;
- (iv) for every e = 1, ..., n and every affine space  $\mathbb{R}^e \simeq E \subset \mathbb{R}^n$ , the number of connected components of  $E \cap C_{\epsilon}$  is bounded by

$$b_0(E \cap C_\epsilon) \le \beta_{\mathrm{dc}}^e$$

3.3. Variations and their behaviour under polynomial maps. We recall the definition of variations, introduced by Vitushkin [Vit55, Vit61] and developed in the semialgebraic context by Comte and Yomdin [YC04]. The variations encode the size of a set, through an integral-geometric approach. Given a polynomial map  $p : \mathbb{R}^n \to \mathbb{R}^m$  and a semialgebraic set  $A \subset \mathbb{R}^n$ , it is possible to estimate the variations of the set p(A), in a quantitative way, see [YC04, Thm. 7.2]. However, these estimates are not quantitative with respect to the dimensional parameter n. The main result of this section is Theorem 19, where we prove a version of [YC04, Thm. 7.2], where the dependence on n is explicit, thanks to the use of Theorem 14.

For  $i \leq n$ , we denote by  $G_i(\mathbb{R}^n)$  the Grassmannian of all *i*-dimensional linear subspaces of  $\mathbb{R}^n$ . We endow it with the standard probability measure  $\gamma_{i,n}$  defined through the action of the orthogonal group of  $\mathbb{R}^n$ . For any  $Z \in G_i(\mathbb{R}^n)$  we the denote the orthogonal projection on Z by

$$\pi_Z: \mathbb{R}^n \to Z$$

**Definition 15** (Vitushkin variations). Let  $A \subset \mathbb{R}^n$  be a bounded semialgebraic set. We define the 0-th variation of A as

$$V_0(A) := b_0(A),$$

where  $b_0(A)$  denotes the number of connected components of A. For i = 1, ..., n we define the *i*-th variation of A as

(8) 
$$V_i(A) := c(n,i) \int_{G_i(\mathbb{R}^n)} \left( \int_Z b_0(A \cap \pi_Z^{-1}(x)) \mathcal{H}^i(dx) \right) \gamma_{i,n}(dZ),$$

where  $\mathcal{H}^i$  denotes the *i*-dimensional Hausdorff measure.

(9) 
$$c(n,i) := \frac{\Gamma(\frac{1}{2})\Gamma(\frac{n+1}{2})}{\Gamma(\frac{i+1}{2})\Gamma(\frac{n-i+1}{2})}.$$

We note that the integrand in (8) is measurable, see [YC04, Sec. 3].

A list of properties of variations is given in [YC04, page 35]. Here we prove the following one.

**Lemma 16.** If  $A \subset \mathbb{R}^n$  is a bounded semialgebraic set of dimension  $\ell$ , then the  $\ell$ -variation coincides with the  $\ell$ -dimensional Hausdorff volume:

$$V_{\ell}(A) = \mathcal{H}^{\ell}(A).$$

We also have  $V_i(A) = 0$  for all  $i > \ell$ .

*Proof.* We recall a classical result in Integral Geometry, called Cauchy–Crofton formula [Fed47, Thm. 5.14]: if A is a *i*-rectifiable and Borel subset of  $\mathbb{R}^n$ , for  $i \leq n$ , then it holds

(10) 
$$\mathcal{H}^{i}(A) = c(n,i) \int_{G_{i}(\mathbb{R}^{n})} \left( \int_{Z} \#(A \cap \pi_{Z}^{-1}(x)) \mathcal{H}^{i}(dx) \right) \gamma_{i,n}(dZ).$$

If  $A \subset \mathbb{R}^n$  is semialgebraic, the set A can be written as a finite union of smooth, semialgebraic disjoint manifolds called strata (see Section 3.1.1). We can relabel the strata so that

$$A = \bigsqcup_{j=0}^{\iota} A_j,$$

where each  $A_j$  is a finite union of smooth *j*-dimensional manifolds. For any fixed  $Z \in G_{\ell}(\mathbb{R}^N)$  and for almost every  $x \in Z$  the set  $\pi_Z^{-1}(x)$  does not intersect<sup>2</sup>  $A_j$  for  $j < \ell$ . Hence,  $V_0(A \cap \pi_Z^{-1}(x)) =$  $V_0(A_{\ell} \cap \pi_Z^{-1}(x))$  for almost every  $x \in Z$ . Furthermore, for almost every  $x \in Z$  the intersection  $A_{\ell} \cap \pi_Z^{-1}(x)$  is transverse, and dim  $A + \dim \pi_Z^{-1}(x) = n$ , so that  $A_{\ell} \cap \pi_Z^{-1}(x)$  has dimension zero. Thus, its cardinality coincides with its number of connected components. We conclude by (10).

Let M be a real  $m \times m$  symmetric and positive semidefinite matrix. We denote by

$$\lambda_1(M) \ge \cdots \ge \lambda_m(M) \ge 0,$$

its eigenvalues, ordered in decreasing order.

**Definition 17** (Singular values). Let H be a Hilbert space with dim  $H \ge m$ , and let  $L : H \to \mathbb{R}^m$  be a linear and continuous map. For any k = 1, ..., m the k-th singular value of L is

$$\sigma_k(L) = \lambda_k(LL^{\top})^{\frac{1}{2}}.$$

In finite dimension, the following Weyl inequality holds for the singular values of the difference of two matrices (see [Tao12, Ex. 1.3.22 (iv)]): for  $n \ge m$  and linear maps  $L_1, L_2 : \mathbb{R}^n \to \mathbb{R}^m$  we have

$$|\sigma_k(L_1) - \sigma_k(L_2)| \le ||L_1 - L_2||_{\text{op}}, \quad \forall k = 1, \dots, m.$$

The Weyl inequality generalizes to infinite-dimensional spaces. Since we are not able to find a reference, we provide below a self-contained statement and its proof in the form that we will need later.

**Lemma 18.** If  $L_1, L_2 \in \mathcal{L}(H, \mathbb{R}^m)$ , then  $|\sigma_k(L_1) - \sigma_k(L_2)| \le ||L_1 - L_2||_{op}$  for all k = 1, ..., m.

*Proof.* Let  $L: H \to \mathbb{R}^m$  be linear and continuous. Let  $W \subset H$  a finite dimensional subspace such that  $(\ker L)^{\perp} \subseteq W$ , and denote by  $L_W: W \to \mathbb{R}^m$  the restriction of L to W. More precisely,  $L_W = L \circ i_W$  where  $i_W: W \hookrightarrow H$  is the inclusion. We now prove that for all  $k \leq m$ 

(11) 
$$\sigma_k(L) = \sigma_k(L_W).$$

Indeed, since the adjoint  $i_W^{\top}$  coincides with the orthogonal projection  $\pi_W : H \to W \subset H$ , we have

$$\lambda_k(L_W \circ L_W^{\top}) = \lambda_k(L \circ i_W \circ i_W^{\top} \circ L^{\top}) = \lambda_k(L \circ \pi_W \circ L^{\top}) = \lambda_k(L \circ L^{\top})$$

In the last equality, we used the fact that  $L \circ \pi_W = L$ , indeed  $L(v) = L(\pi_W v + \pi_{W^{\perp}} v) = L(\pi_W v)$ since  $W^{\perp} \subseteq \ker L$ . Now given  $L_1, L_2 \in \mathcal{L}(H, \mathbb{R}^m)$  let  $W := (\ker L_1)^{\perp} + (\ker L_2)^{\perp}$ , which is a finite dimensional linear subspace of H. By (11) and the finite dimensional Weyl inequality, we have

$$|\sigma_k(L_1) - \sigma_k(L_2)| = |\sigma_k((L_1)_W) - \sigma_k((L_2)_W)| \le ||(L_1)_W - (L_2)_W||_{\text{op}} \le ||L_1 - L_2||_{\text{op}},$$

concluding the proof.

In the next result, given a semialgebraic set  $A \subset \mathbb{R}^n$  and a polynomial map  $p : \mathbb{R}^n \to \mathbb{R}^m$ , we bound the *i*-th variation  $V_i(p(A))$  in terms of the singular values of p and the diagram of A. Its proof follows the blueprint of [YC04, Thm. 7.2]. The main novelty (which is key for our work) is that the dependence on the dimension n is explicit.

**Theorem 19.** For every  $c, d, m \in \mathbb{N}$  there exists  $\beta_{yc} = \beta_{yc}(c, d, m) > 1$  with the following property. For  $n \ge m$ , let  $A \subset \mathbb{R}^n$  be a closed and bounded semialgebraic set with  $A \subset B_{\mathbb{R}^n}(r)$  and diagram D(A) = (n, c, d). Let  $p : \mathbb{R}^n \to \mathbb{R}^m$  be a polynomial map with components of degrees at most d. For all i = 1, ..., m set

(12) 
$$\overline{\sigma}_i := \sup_{x \in A} \left( \sigma_1(D_x p) \cdots \sigma_i(D_x p) \right), \qquad \overline{\sigma}_0 := 1.$$

 $<sup>^{2}</sup>$ We note that in this step of the proof it is enough to use the Semialgebraic Sard's theorem [BCR98, Thm. 9.6.2], which is logically independent from the result in the smooth category.

Then, for all  $i = 0, \ldots, m$  it holds

$$V_i(p(A)) \le \operatorname{cst}(m, r) n^m \beta_{\mathrm{vc}}^n \overline{\sigma}_i$$

where cst(m, r) > 0 is a constant that depends only on m, r.

Remark 1 (Origin of  $\beta_{yc}$ ). An inspection of the proof shows that it holds  $\beta_{yc} = \beta_{tm}^2 \beta_{dc}$  where  $\beta_{tm} = \beta_{tm}(c(m+1), d)$  is the constant from the Thom-Milnor-type bound of Theorem 13, and  $\beta_{dc} = \max_{i \leq m} \beta_{dc}(c, d, i)$  is the definable choice constant from Theorem 14.

*Remark* 2. In the course of the proof, see (18), we obtain the following estimate, reminiscent of an area formula for variations, which is an improved version of [YC04, Thm. 7.2], with the explicit dependence of the constants on n: there exist a constant  $\beta_{yc} = \beta_{yc}(c, d, m)$ , such that

$$V_i(p(A)) \le c(m,i)\beta_{\mathrm{vc}}^n \overline{\sigma}_i V_i(A), \qquad i=1,\ldots,m_i$$

where c(m, i) is the constant in (9).

*Proof.* By Definition 15, for i = 1, ..., m, we have

$$V_i(p(A)) = c(m,i) \int_{G_i(\mathbb{R}^m)} \left( \int_Z b_0(p(A) \cap \pi_Z^{-1}(x)) \mathcal{H}^i(dx) \right) \gamma_{i,m} (dZ).$$

For any  $x \in Z$ , the set  $A \cap (\pi_Z \circ p)^{-1}(x)$  is semialgebraic with controlled diagram (more precisely,  $D(A \cap (\pi_Z \circ p)^{-1}(x)) = (n, c(1+m), d)$ . Using Theorem 13 we obtain

(13) 
$$b_0(p(A) \cap \pi_Z^{-1}(x)) \le b_0(A \cap (\pi_Z \circ p)^{-1}(x))) \le \beta_{\rm tm}^n$$

where  $\beta_{tm} = \beta_{tm}(c(1+m), d)$  depends only on c, d, m. Note that, since by definition  $V_0 = b_0$ , (13) yields the thesis of the theorem for the case i = 0. We continue then assuming  $i = 1, \ldots, m$ . Denoting by  $\mathbb{1}_S$  the characteristic function of a set S, we get

(14)  

$$V_{i}(p(A)) \leq c(m,i)\beta_{\mathrm{tm}}^{n} \int_{G_{i}(\mathbb{R}^{m})} \left( \int_{Z} \mathbb{1}_{\pi_{Z} \circ p(A)}(x) \mathcal{H}^{i}(dx) \right) \gamma_{i,m}(dZ)$$

$$= c(m,i)\beta_{\mathrm{tm}}^{n} \int_{G_{i}(\mathbb{R}^{m})} \mathcal{H}^{i}(\pi_{Z} \circ p(A)) \gamma_{i,m}(dZ).$$

We would like to estimate  $\mathcal{H}^i(\pi_Z \circ p(A))$  applying area formula to the map

 $p_Z := \pi_Z \circ p : \mathbb{R}^n \to \mathbb{R}^i.$ 

However, exactly as in [YC04, Thm. 7.2], we cannot apply directly the area formula to A, since the dimension of A can be larger than i. In [YC04, Thm. 7.2] this problem is solved using [YC04, Ex. 4.11]. Here we overcome the obstacle using Theorem 14, replacing the set A with a suitable approximation.

For sufficiently small  $\epsilon > 0$  let  $C_{\epsilon}$  the semialgebraic set obtained by applying Theorem 14 to the set A and the polynomial map  $F = p_Z : \mathbb{R}^n \to \mathbb{R}^i$ . We then proceed using the area formula to estimate  $\mathcal{H}^i(\pi_Z \circ p(C_{\epsilon}))$ . (We will see at the end of the proof how  $\mathcal{H}^i(\pi_Z \circ p(C_{\epsilon})) \to \mathcal{H}^i(\pi_Z \circ p(A))$  as  $\epsilon \to 0$ ). By Item (i) of Theorem 14, dim $(C_{\epsilon}) \leq i$ . We can assume without loss of generality that  $C_{\epsilon} \subset \mathbb{R}^n$  is a smooth embedded submanifold with dimension equal to i, since the other strata give zero contribution to  $\mathcal{H}^i$ . Denoting with  $\bar{p}_Z : C_{\epsilon} \to \mathbb{R}^i$  the restriction, we obtain

$$\begin{aligned} \mathcal{H}^{i}(p_{Z}(C_{\epsilon})) &\leq \int_{C_{\epsilon}} |\det(D_{x}\bar{p}_{Z})| \mathcal{H}^{i}(dx) \qquad \qquad \text{by the area formula} \\ &\leq \int_{C_{\epsilon}} \sigma_{1}(D_{x}\bar{p}_{Z}) \cdots \sigma_{i}(D_{x}\bar{p}_{Z}) \mathcal{H}^{i}(dx). \qquad \qquad \text{by definition of singular values} \end{aligned}$$

Note that, denoting with  $j: C_{\epsilon} \hookrightarrow \mathbb{R}^n$  the inclusion, we have

$$D_x \bar{p}_Z : T_x C_\epsilon \xrightarrow{D_x j} \mathbb{R}^n \xrightarrow{D_x p} \mathbb{R}^m \xrightarrow{D_{p(x)} \pi_Z} \mathbb{R}^i.$$

Thus, since  $D_{p(x)}\pi_Z$  is an orthogonal projection and  $D_x j$  is a linear inclusion, we have

$$\langle z, (D_x \bar{p}_Z)(D_x \bar{p}_Z)^\top z \rangle \le \langle z, (D_x p)(D_x p)^\top z \rangle, \qquad \forall z \in Z = \mathbb{R}^i.$$

It follows that  $\sigma_k(D_x\bar{p}_Z) \leq \sigma_k(D_xp)$  for all  $k = 1, \ldots, i$ . Continuing the above inequalities we obtain

$$\mathcal{H}^{i}(p_{Z}(C_{\epsilon})) \leq \int_{C_{\epsilon}} \sigma_{1}(D_{x}p) \cdots \sigma_{i}(D_{x}p) \mathcal{H}^{i}(dx)$$

$$\leq \sup_{x \in \mathcal{U}_{\epsilon}(A)} \sigma_{1}(D_{x}p) \cdots \sigma_{i}(D_{x}p) \mathcal{H}^{i}(C_{\epsilon}) \qquad \text{by Item (ii) of Theorem 14}$$

$$\leq (\overline{\sigma}_{i} + \eta(\epsilon)) \mathcal{H}^{i}(C_{\epsilon}), \qquad \text{by continuity of singular values}$$

where  $\overline{\sigma}_i$  is defined in (12), and  $\eta(\epsilon) \to 0$  as  $\epsilon \to 0$ . Since  $C_{\epsilon}$  is a bounded semialgebraic set of dimension *i*, then  $\mathcal{H}^i(C_{\epsilon}) = V_i(C_{\epsilon})$  by Lemma 16. Hence

$$\begin{aligned} \mathcal{H}^{i}(C_{\epsilon}) &= c(n,i) \int_{G_{i}(\mathbb{R}^{n})} \left( \int_{Z} b_{0}(C_{\epsilon} \cap \pi_{Z}^{-1}(x)) \mathcal{H}^{i}(dx) \right) \gamma_{i,n} (dZ) \\ &\leq c(n,i) \beta_{\mathrm{dc}}^{n-i} \int_{G_{i}(\mathbb{R}^{n})} \left( \int_{Z} \mathbb{1}_{\pi_{Z}(C_{\epsilon})}(x) \mathcal{H}^{i}(dx) \right) \gamma_{i,n} (dZ) \qquad \text{by Item (iv) of Theorem 14} \\ &\leq c(n,i) \beta_{\mathrm{dc}}^{n-i} \int_{G_{i}(\mathbb{R}^{n})} \left( \int_{Z} \mathbb{1}_{\pi_{Z}(\mathcal{U}_{\epsilon}(A))}(x) \mathcal{H}^{i}(dx) \right) \gamma_{i,n} (dZ) \qquad \text{by Item (ii) of Theorem 14} \\ &\leq c(n,i) \beta_{\mathrm{dc}}^{n-i} \int_{G_{i}(\mathbb{R}^{n})} \left( \int_{Z} b_{0}(\mathcal{U}_{\epsilon}(A) \cap \pi_{Z}^{-1}(x)) \mathcal{H}^{i}(dx) \right) \gamma_{i,n} (dZ) \\ &= \beta_{\mathrm{dc}}^{n-i} V_{i}(\mathcal{U}_{\epsilon}(A)), \end{aligned}$$

where  $\beta_{dc} = \beta_{dc}(c, d, i) > 1$  is the constant from Theorem 14, which (up to taking the maximum for  $i \leq m$ ) depends only on c, d, m. Summing up, we have proved that for all  $i = 1, \ldots, m$  it holds

(15) 
$$\mathcal{H}^{i}(p_{Z}(C_{\epsilon})) \leq (\overline{\sigma}_{i} + \eta(\epsilon))\beta_{\mathrm{dc}}^{n}V_{i}(\mathcal{U}_{\epsilon}(A)).$$

In order to take the limit in (15), we recall two properties of semialgebraic sets. The first one is the continuity of the Lebesgue measure in the Hausdorff topology in the semialgebraic category (see [YC04, Thm. 5.10]): if  $\{S_{\epsilon}\}_{\epsilon \geq 0} \subset \mathbb{R}^{i}$  is a one-parameter family of bounded semialgebraic sets then

(16) 
$$\lim_{\epsilon \to 0} \operatorname{dist}_H(S_{\epsilon}, S_0) = 0 \qquad \Longrightarrow \qquad \lim_{\epsilon \to 0} \mathcal{H}^i(S_{\epsilon}) = \mathcal{H}^i(S_0).$$

The second property is the behaviour of variations for  $\epsilon$ -neighbourhoods (see [YC04, Thm. 5.11]): if  $A \subset \mathbb{R}^n$  is a bounded semialgebraic set, then for all i = 1, ..., n it holds

(17) 
$$\lim_{\epsilon \to 0} V_i(\mathcal{U}_{\epsilon}(A)) \le V_i(A).$$

Therefore, using (16) (and recalling that by Item (iii) of Theorem 14 it holds  $p_Z(C_{\epsilon}) \rightarrow p_Z(A)$  in the Hausdorff topology) and (17) we can pass to the limit in (15) and obtain

$$\mathcal{H}^{i}(p_{Z}(A)) \leq \overline{\sigma}_{i}\beta_{\mathrm{dc}}^{n}V_{i}(A).$$

Hence from (14) we get for all i = 1, ..., m

(18) 
$$V_i(p(A)) \le c(m,i)\beta_{\rm tm}^n \beta_{\rm dc}^n \overline{\sigma}_i V_i(A).$$

To conclude, we estimate  $V_i(A)$ . Similarly as in (14), and recalling that  $A \subset B_{\mathbb{R}^n}(r)$  we have

(19) 
$$V_i(A) \le c(n,i)\beta_{\mathrm{tm}}^n \int_{G_i(\mathbb{R}^n)} \mathcal{H}^i(\pi_Z(A))\gamma_{i,n}(dZ) \le c(n,i)\beta_{\mathrm{tm}}^n \mathcal{H}^i(B_{\mathbb{R}^i}(r)) = c(n,i)\beta_{\mathrm{tm}}^n \frac{\pi^{i/2}}{\Gamma\left(\frac{i}{2}+1\right)}r^i.$$

Putting together (18) and (19), we obtain

$$V_i(p(A)) \le \left[ c(m,i)c(n,i)\frac{\pi^{i/2}}{\Gamma\left(\frac{i}{2}+1\right)} \right] (\beta_{\rm tm}^2\beta_{\rm dc})^n \overline{\sigma}_i r^i$$

Set  $\beta_{\rm yc} = \beta_{\rm tm}^2 \beta_{\rm dc}$ . Using the form of the constants c(m, i), c(n, i) in (9) and elementary estimates we obtain that for all  $i = 0, \ldots, m$  there exists a constant  $\operatorname{cst}(i) > 0$  such that

$$V_i(p(A)) \le \operatorname{cst}(i)n^i \beta_{\mathrm{yc}}^n \overline{\sigma}_i r^i$$

Taking the maximum over i = 1, ..., m we obtain the result with  $\operatorname{cst}(m, r) = \max_{i=0,...,m} \operatorname{cst}(i)r^i$ .

3.4. Variations of almost-critical values of polynomial maps. In this section we recall the notion of almost-critical points  $C^{\Lambda}(p)$  of a polynomial map  $p : \mathbb{R}^n \to \mathbb{R}^m$ . The set  $p(C^{\Lambda}(p))$  is the set of its almost-critical values. The main result of this section is Theorem 24, which provides a version of [YC04, Cor. 7.4] where the dependence of the parameters with respect to n is explicit. This result is based on Theorem 19 and on the specific semialgebraic structure of almost-critical points of polynomial maps, that we prove in Proposition 23.

**Definition 20.** Let  $f : \mathbb{R}^n \to \mathbb{R}^m$  be a  $C^1$  map with  $n \ge m$ . Given  $\Lambda = (\Lambda_1, \ldots, \Lambda_m) \in \mathbb{R}^m_+$ , the set of almost-critical points of f is defined as

$$C^{\Lambda}(f) := \left\{ x \in \mathbb{R}^n \middle| \sigma_i(D_x f) \le \Lambda_i, \, \forall \, i = 1, \dots, m \right\}.$$

The following lemma is fundamental to study the semialgebraic structure of the set of almost-critical points of a polynomial map.

**Lemma 21.** Let  $A \subset \mathbb{R}^s$  be a semialgebraic set and let  $f : A \to \mathbb{R}$  be a semialgebraic function. Then, for any  $t \in \mathbb{R}$ , the sublevel set  $\{x \in A \mid f(x) \leq t\}$  admits a semialgebraic description whose diagram does not depend on t.

*Proof.* The graph of f is semialgebraic, hence we can write it as

$$\operatorname{graph}(f) = \bigcup_{i=1}^{a} \bigcap_{j=1}^{b_i} E_{ij},$$

where  $E_{ij}$  is of the form  $\{p_{ij} < 0\}$  or  $\{p_{ij} \le 0\}$  and  $p_{ij} : \mathbb{R}^s \times \mathbb{R} \to \mathbb{R}$  is a polynomial. Denoting by  $\pi_1 : \mathbb{R}^s \times \mathbb{R} \to \mathbb{R}^s$  the projection, we have

(20) 
$$\{x \in A \mid f(x) \le t\} = \pi_1 \Big( \operatorname{graph}(f) \cap \{(x, y) \in \mathbb{R}^s \times \mathbb{R} \mid y - t \le 0\} \Big)$$

The set graph $(f) \cap \{(x, y) \in \mathbb{R}^s \times \mathbb{R} \mid y - t \leq 0\}$  is semialgebraic, with a description given by

graph(f) 
$$\cap \{(x,y) \mid y-t \le 0\} = \bigcup_{i=1}^{a} \bigcap_{j=1}^{b_i+1} E_{i,j},$$

where  $E_{i,b_i+1} = \{(x, y) \in \mathbb{R}^s \times \mathbb{R} \mid y-t \leq 0\}$ . In particular, this set admits a diagram that does not depend on t, since the degree of the polynomial y-t is equal to 1. As explained in Section 3.1.3, the projection of a semialgebraic set is semialgebraic and its diagram is controlled explicitly by the diagram of the original set only. In particular, the diagram of the projection (20) admits a description with a diagram (s, I, J), that does not depend on t.

Let  $\operatorname{Sym}_m^+$  denote the set of positive semidefinite matrices of size m. This is clearly a semialgebraic subset of all  $m \times m$  real matrices. The following lemma is elementary and we omit its proof.

**Lemma 22.** The functions  $\lambda_i : \operatorname{Sym}_m^+ \to \mathbb{R}$  and are continuous and semialgebraic.

In the following proposition we provide an estimate on the diagram of a description of the (semial-gebraic) set of almost-critical points of polynomial maps.

**Proposition 23.** Let  $d, m \in \mathbb{N}$ . Then there are c' = c'(d, m) and d' = d'(d, m) such that for all  $n \geq m$  and any polynomial map  $p : \mathbb{R}^n \to \mathbb{R}^m$  with components of degrees at most d, and all  $\Lambda = (\Lambda_1, \ldots, \Lambda_m) \in \mathbb{R}^m_+$ , the set  $C^{\Lambda}(p)$  is closed, semialgebraic, with

$$D(C^{\Lambda}(p)) = (n, c', d')$$

(In particular, the diagram does not depend on  $\Lambda$  and it depends on n only in the number of variables.)

*Proof.* By Lemma 22, for every k = 1, ..., m, the function  $\lambda_k : \operatorname{Sym}_m^+ \to \mathbb{R}$  is semialgebraic, hence from Lemma 21 applied to  $A = \operatorname{Sym}_m^+$  and the semialgebraic functions  $\lambda_k : A \to \mathbb{R}$ , we deduce the following fact: for any  $t \in \mathbb{R}$  we can write

$$\{M \in \operatorname{Sym}_{m}^{+} \mid \lambda_{k}(M) \leq t^{2}\} = \bigcup_{i=1}^{a} \bigcap_{j=1}^{b_{i}} \{M \in \mathbb{R}^{m \times m} \mid P_{k,i,j,t}(M) \leq 0\}$$

for some polynomials  $P_{k,i,j,t} : \mathbb{R}^{m \times m} \to \mathbb{R}, a, b_1, \ldots, b_a \in \mathbb{N}$ , and with  $a \max_i b_i$  and  $\max_{i,j} \deg(P_{k,i,j,t})$  depending only on m and on k (and not on t). We can assume that a does not depend on k, and all the  $b_i$ 's do not depend on a, k, simply by adding empty or trivial sets.

Now, for any  $k = 1, \ldots, m$ 

$$\{N \in \mathbb{R}^{m \times n} \mid \sigma_k(N) \le t\} = \{N \in \mathbb{R}^{m \times n} \mid \lambda_k(NN^{\top}) \le t^2\},\$$

therefore we can write

$$\{x \in \mathbb{R}^n \mid \sigma_k(D_x p) \le t\} = \bigcup_{i=1}^a \bigcap_{j=1}^{b_i} \{x \in \mathbb{R}^n \mid P_{k,i,j,t}(D_x p D_x p^\top) \le 0\}.$$

Observe that  $N \to NN^{\top}$  is polynomial with components of degree 2. Therefore, the map  $q : \mathbb{R}^n \to \mathbb{R}^{m \times m}$ , given by  $q(x) := D_x p D_x p^{\top}$  is polynomial, with components of degree at most 2*d*. Hence, the polynomial

$$P_{k,i,j,t} \circ q : \mathbb{R}^n \to \mathbb{R}$$

has degree at most  $2d \max_{i,j} \deg P_{k,i,j,t}$ , and in particular its degree has a bound that depends on only d, m, k (and not on t). The set of almost-critical points of p can be described as

$$C^{\Lambda}(p) = \bigcap_{k=1}^{m} \{ x \in \mathbb{R}^{n} \mid \sigma_{k}(D_{x}p) \le \Lambda_{k} \} = \bigcap_{k=1}^{m} \bigcup_{i=1}^{a} \bigcup_{j=1}^{b_{i}} \{ x \mid P_{k,i,j,\Lambda_{k}}(D_{x}pD_{x}p^{\top}) \le 0 \}.$$

It is clear that  $C^{\Lambda}(p)$  is closed. Note that  $d' := \max_{k,i,j} \deg(P_{k,i,j,\Lambda_k} \circ q)$  depends only on d, m. Using the distributivity of intersection, we can rewrite  $C^{\Lambda}(p)$  in the form (7), where the number of unions and intersections depends only on m, and the maximum degree of the polynomials is c' = c'(d, m), concluding the proof.

Now we state and prove the main result of this section (Theorem G in the Introduction).

**Theorem 24.** Let  $n \ge m$ , and  $p : \mathbb{R}^n \to \mathbb{R}^m$  be a polynomial map with components of degree at most d. For  $i = 0, \ldots, m$ ,  $\Lambda = (\Lambda_1, \ldots, \Lambda_m) \in \mathbb{R}^m_+$  and r > 0, we have

(21) 
$$V_i(p(C^{\Lambda}(p) \cap B_{\mathbb{R}^n}(r))) \le \operatorname{cst}(m, r)n^m \beta_0^n \Lambda_0 \cdots \Lambda_i$$

where  $\beta_0 = \beta_0(d,m)$  depends only on d and m,  $\operatorname{cst}(m,r)$  depends only on m, r, and we set  $\Lambda_0 = 1$ .

Proof. By Proposition 23 the set  $A = C^{\Lambda}(p) \cap B_{\mathbb{R}^n}(r)$  is a closed and bounded semialgebraic set whose diagram is D(A) = (n, c', d') where c' = c'(d, m), d' = d'(d, m) (which in particular do not depend on  $n, \Lambda$ ). Furthermore, for the polynomial map  $p : \mathbb{R}^n \to \mathbb{R}^m$  it holds, by construction

$$\overline{\sigma}_i := \sup_{x \in A} \left( \sigma_1(D_x p) \cdots \sigma_i(D_x p) \right) \le \Lambda_0 \cdots \Lambda_i.$$

We can then apply Theorem 19, yielding (21) with  $\beta_0 = \beta_{yc}(c', \max\{d, d'\}, m)$ .

# 

#### 4. SARD PROPERTIES FOR INFINITE-DIMENSIONAL MAPS

Let H be a Hilbert space. We consider a map  $f: U \to \mathbb{R}^m$  where U is an open subset of H. If f is  $C^1$  we denote by  $\operatorname{Crit}(f)$  the set of its critical points. For a fixed  $\nu \in \mathbb{N}$ , we also consider the set

(22) 
$$\operatorname{Crit}_{\nu}(f) = \{ x \in U \mid \operatorname{rank}(D_x f) \le \nu \}.$$

Given a relatively compact  $S \subset \mathbb{R}^m$ , for any  $\epsilon > 0$  we denote with  $M(\epsilon, S)$  the  $\epsilon$ -entropy of S, which is the minimum number of closed balls of radius  $\epsilon$  that we need to cover S.

**Definition 25** (Entropy dimension). The entropy dimension of S is defined as

$$\dim_e(S) = \limsup_{\epsilon \to 0^+} \frac{\ln M(\epsilon, S)}{\ln(\frac{1}{\epsilon})}.$$

In [YC04, Ch. 2], it is proved that

 $\dim_{\mathcal{H}}(S) \le \dim_{e}(S).$ 

The inequality can be strict, since there are sequences of real numbers with positive entropy dimension.

For Banach spaces X, Y and linear continuous maps  $L \in \mathcal{L}(X, Y)$ , we denote by

(23) 
$$||L||_{\rm op} := \sup_{\|x\|_X \le 1} \|Lx\|_Y$$

the operator norm. With some abuse of notation the same symbol  $||L||_{\text{op}}$  is used for different X and Y, but there should be no confusion as the domain and codomain of L are clear from the context. We also use without risk of confusion the symbol  $|| \cdot ||$  to denote the usual Euclidean norm of  $\mathbb{R}^m$ .

4.1. Sard-type theorem for well approximated maps. The purpose of this section is to prove the following theorem, which is our main Sard-type result (Theorem E in the Introduction).

**Theorem 26.** Let  $d, m \in \mathbb{N}$ . There exists a constant  $\beta_0 = \beta_0(d, m) > 0$  such that the following holds. Let H be a Hilbert space, and let  $f : H \to \mathbb{R}^m$  be a  $C^1$  map such that its differential  $Df : H \to \mathcal{L}(H, \mathbb{R}^m)$ is weakly continuous. Let  $K \subset H$  be a bounded set with this approximation property: there exist a sequence  $E_n \subset H$  of linear subspaces,  $\dim(E_n) = n$ , and polynomial maps  $f_n : E_n \to \mathbb{R}^m$  with uniformly bounded degree:

$$\sup_{n \in \mathbb{N}} \deg f_n \le d < \infty,$$

such that for some q > 1,  $c \ge 0$ , and all large enough n it holds

(24) 
$$\sup_{x \in K} \left( \|f(x) - f_n \circ \pi_{E_n}(x)\| + \|(D_x f)|_{E_n} - D_{\pi_{E_n}(x)} f_n\|_{\mathrm{op}} \right) \le cq^{-n}.$$

Then

$$\dim_e \left( f \big( \operatorname{Crit}_{\nu}(f) \cap K \big) \right) \le \nu + \frac{\ln \beta_0}{\ln q}, \qquad \forall \, \nu = 1, \dots, m-1.$$

In particular, if  $q > \beta_0$ , then f satisfies the Sard property on K:

$$\mu\bigg(f(\operatorname{Crit}(f)\cap K)\bigg)=0.$$

Remark 3. As it will be clear from the proof, the upper bound  $cq^{-n}$  in the r.h.s. of (24) can be replaced by any  $C_n \ge 0$ , such that  $\limsup_{n\to\infty} C_n^{1/n} = q^{-1}$ , yielding the same results.

For the benefit of the reader we provide a general outline of the strategy of the proof. Our goal is to estimate the "size" of the set  $f(\operatorname{Crit}_{\nu}(f)\cap K)$ . More precisely, we bound the  $\epsilon$ -entropy of  $f(\operatorname{Crit}_{\nu}(f)\cap K)$  for any  $\epsilon > 0$ , and then pass to the limit  $\epsilon \to 0$ .

We observe that the  $\epsilon$ -entropy of  $f(\operatorname{Crit}_{\nu}(f) \cap K)$  is controlled with the one of the set of almostcritical values  $f(C^{\Lambda}(f) \cap K)$ , see Definition 17. In turn, this is controlled by the  $\epsilon$ -entropy of suitable almost-critical values of the approximating maps  $f_n$ , namely  $f(C^{\Lambda_{\epsilon}}(f_n) \cap K)$ , see (29).

To proceed, we use the relation between the  $\epsilon$ -entropy of sets in  $\mathbb{R}^m$  and their Vitushkin variations (Definition 15), provided by [YC04, Thm. 3.5].

Finally, we estimate the Vitushkin variations of  $f_n(C^{\Lambda_{\epsilon}}(f_n)\cap K)$  applying Theorem G, exploiting the hypothesis that the maps  $f_n$  are polynomial. This is the connection with quantitative semialgebraic geometry from Section 3. The crucial point of Theorem G is that the bound is quantitative with respect to parameters  $\Lambda = \Lambda_{\epsilon}$  and n. This allows to chose a large enough  $n = n_{\epsilon}$ , see (35), to ensure convergence when passing to the limit  $\epsilon \to 0$ .

*Proof.* We divide the proof in steps.

Step 1. In the first step we show that for any  $\nu = 1, \ldots, m-1$  the set  $\operatorname{Crit}_{\nu}(f) \cap K$  is contained in a suitable set of almost-critical points of f. By assumption, K is weakly pre-compact. Since  $Df : H \to \mathcal{L}(H, \mathbb{R}^m)$  is weakly continuous, and by Lemma 18 the function  $\sigma_i : \mathcal{L}(H, \mathbb{R}^m) \to \mathbb{R}$  is continuous for every  $i = 1, \ldots, m$ . Therefore we have

$$\Sigma_i := \sup_K \sigma_i(D_x f) < \infty.$$

Hence, defining  $\Lambda := (\Sigma_1, \ldots, \Sigma_\nu, 0 \ldots, 0) \in \mathbb{R}^m$ , we get

(25) 
$$\operatorname{Crit}_{\nu}(f) \cap K \subset C^{\Lambda}(f) \cap K.$$

Step 2. We now relate critical points of f to almost-critical points of the approximating polynomials  $f_n$ . Fix  $\epsilon > 0$ . Recall that, by Lemma 18, the singular values are 1-Lipschitz with respect to the operator norm. Then for any sufficiently large  $n \in \mathbb{N}$  such that  $cq^{-n} \leq \epsilon$  and for all  $x \in K$  we have

$$|\sigma_i (D_x f|_{E_n}) - \sigma_i (D_{\pi_{E_n}(x)} f_n)| \le || (D_x f) |_{E_n} - D_{\pi_{E_n}(x)} f_n ||_{\text{op}} \le \epsilon.$$

Hence, for all  $x \in K \cap C^{\Lambda}(f)$  we have the following estimate for all  $i = 1, \ldots, m$ 

(26) 
$$\sigma_i \left( D_{\pi_{E_n}(x)} f_n \right) \le \epsilon + \sup_{y \in C^{\Lambda}(f) \cap K} \sigma_i \left( D_y f |_{E_n} \right) \le \epsilon + \sup_{y \in C^{\Lambda}(f) \cap K} \sigma_i \left( D_y f \right) \le \epsilon + \Lambda_i,$$

where, in the second inequality, we used that singular values are monotone with respect to restriction to subspaces. Hence, defining

$$\Lambda^{\epsilon} := (\Sigma_1 + \epsilon, \dots, \Sigma_{\nu} + \epsilon, \epsilon, \dots, \epsilon),$$

by (26) we get

(27) 
$$\pi_{E_n}\left(C^{\Lambda}(f)\cap K\right)\subset C^{\Lambda^{\epsilon}}(f_n)\cap\pi_{E_n}(K)\subset C^{\Lambda^{\epsilon}}(f_n)\cap B_{E_n}(r),$$

where r > 0 is such that  $K \subseteq B_H(r)$ , and thus  $B_{E_n}(r) = B_H(r) \cap E_n$  denotes the ball in  $E_n$  defined by the restriction of the Hilbert norm.

Step 3. Now we use the inclusion (27) to estimate the entropy dimension of  $f(\operatorname{Crit}_{\nu}(f) \cap K)$  with the one of almost-critical values of  $f_n$ . By (24), for sufficiently large n, we have

(28) 
$$\sup_{x \in K} \|f(x) - f_n \circ \pi_{E_n}(x)\| \le \epsilon$$

Finally, we get

$$M(2\epsilon, f(\operatorname{Crit}_{\nu}(f) \cap K) \leq M(2\epsilon, f(C^{\Lambda}(f) \cap K)) \qquad \text{by (25)}$$
  
$$\leq M\left(2\epsilon, \mathcal{U}_{\epsilon}\left(f_{n} \circ \pi_{E_{n}}\left(C^{\Lambda}(f) \cap K\right)\right)\right) \qquad \text{by (28)}$$
  
$$\leq M\left(\epsilon, f_{n} \circ \pi_{E_{n}}\left(C^{\Lambda}(f) \cap K\right)\right) \qquad \text{by definition of } \mathcal{U}_{\epsilon}$$
  
$$\leq M(\epsilon, f_{n}(C^{\Lambda^{\epsilon}}(f_{n}) \cap B_{E_{n}}(r))), \qquad \text{by (27)}$$

(29) 
$$\leq M(\epsilon, f_n(C^{\Lambda^{\epsilon}}(f_n) \cap B_{E_n}(r))), \qquad \text{by (27)}$$
  
where we recall that  $M(\varepsilon, A)$  is the minimal number of closed balls of  $\mathbb{R}^m$  of radius  $\epsilon$  needed to cover

 $A \subset \mathbb{R}^m$ , and  $\mathcal{U}_{\epsilon}(A)$  denotes the  $\epsilon$  neighbourhood of A. This concludes the proof of step 3.

The rest of the proof consists in estimating (29). We will use the results from Section 3.4, for this reason in the next step we discuss how to bring the problem to the Euclidean space  $\mathbb{R}^n$ .

**Step 4.** We fix a basis for  $E_n$  which is orthonormal with respect to the inner product of H. This yields a linear isometry of Hilbert spaces  $\ell_n : \mathbb{R}^n \to E_n$ . In particular, for the transpose, it holds  $\ell_n^{\top} = \ell_n^{-1}$ . Consider then the polynomial map  $\tilde{f}_n : \mathbb{R}^n \to \mathbb{R}^m$  defined by  $\tilde{f}_n := f_n \circ \ell_n, n \ge m$ . We have

$$\sigma_i(D_v \tilde{f}_n) = \lambda_i (D_{\ell_n(v)} f_n \circ \ell_n \circ \ell_n^\top \circ D_{\ell_n(v)} f_n^\top)^{1/2} = \sigma_i (D_{\ell_n(v)} f_n), \qquad \forall v \in \mathbb{R}^n.$$

It follows that, for sufficiently large n,

$$\ell_n(C^{\Lambda^{\epsilon}}(\tilde{f}_n)) = C^{\Lambda^{\epsilon}}(f_n).$$

Hence, we have

(30) 
$$f_n(C^{\Lambda^{\epsilon}}(f_n) \cap B_{E_n}(r)) = f_n(\ell_n(C^{\Lambda^{\epsilon}}(\tilde{f}_n)) \cap B_{E_n}(r)) = \tilde{f}_n(C^{\Lambda^{\epsilon}}(\tilde{f}_n) \cap B_{\mathbb{R}^n}(r)).$$

By [YC04, Thm. 3.5] we have

(31) 
$$M(\epsilon, \tilde{f}_n(C^{\Lambda^{\epsilon}}(\tilde{f}_n) \cap B_{\mathbb{R}^n}(r))) \le C(m) \sum_{i=0}^m \frac{1}{\epsilon^i} V_i \bigg( \tilde{f}_n(C^{\Lambda^{\epsilon}}(\tilde{f}_n) \cap B_{\mathbb{R}^n}(r)) \bigg),$$

where C(m) > 0 is a constant that depends only on m. We now apply the results from Section 3.4 in order to estimate the right-hand side of the last inequality.

**Step 5.** Take *n* sufficiently large so that it also satisfies  $n \ge m$ . Then, by Theorem G, for every i = 0, ..., m we have

(32) 
$$V_i\bigg(\tilde{f}_n(C^{\Lambda^{\epsilon}}(\tilde{f}_n) \cap B_{\mathbb{R}^n}(r))\bigg) \leq \operatorname{cst}(m,r)n^m\beta_0^n\Lambda_0^{\epsilon}\cdots\Lambda_i^{\epsilon},$$

where we recall  $\beta_0 = \beta_0(d, m)$  and  $\Lambda_0^{\epsilon} = 1$  by definition. Now, we have for all  $i = 1, \ldots, m$ 

(33)  

$$\Lambda_{1}^{\epsilon} \cdots \Lambda_{i}^{\epsilon} = (\Lambda_{1} + \epsilon) \cdots (\Lambda_{i} + \epsilon)$$

$$= \sum_{h=0}^{i} \epsilon^{i-h} \sum_{\substack{0=j_{0} < j_{1} < \cdots < j_{h} \leq i}} \Lambda_{j_{0}} \cdots \Lambda_{j_{h}}$$

$$\leq \sum_{h=0}^{i} \epsilon^{i-h} \sum_{\substack{0=j_{0} < j_{1} < \cdots < j_{h} \leq i}} \Lambda_{0} \cdots \Lambda_{h}$$

$$= {i \choose h} \sum_{h=0}^{i} \epsilon^{i-h} \Lambda_{0} \cdots \Lambda_{h}.$$

Therefore, by (31), (32) and (33), for large n we have

$$M(\epsilon, \tilde{f}_n(C^{\Lambda^{\epsilon}}(\tilde{f}_n) \cap B_{\mathbb{R}^n}(r))) \leq \operatorname{cst}(m, r)C(m)n^m \beta_0^n \sum_{i=0}^m \frac{i!}{\epsilon^i} \sum_{h=0}^i \epsilon^{i-h} \Lambda_0 \cdots \Lambda_h$$
$$\leq \widetilde{\operatorname{cst}}(m, r)n^m \beta_0^n \sum_{h=0}^\nu \frac{\Lambda_0 \cdots \Lambda_h}{\epsilon^h},$$

where we note that, by definition,  $\Lambda_h = 0$  for all  $h > \nu$ , and  $\widetilde{\operatorname{cst}}(m, r)$  denotes a constant that depends only on m, r. From this we deduce that

(34) 
$$\ln M(\epsilon, \tilde{f}_n(C^{\Lambda^{\epsilon}}(\tilde{f}_n) \cap B_{\mathbb{R}^n}(r))) \le \ln(\widetilde{\operatorname{cst}}(m, r)) + n \ln \beta_0 + m \ln n + \ln \left(\sum_{h=0}^{\nu} \frac{\Lambda_0 \cdots \Lambda_h}{\epsilon^h}\right),$$

for all  $\epsilon > 0$  and  $n \in \mathbb{N}$  such that  $cq^{-n} \leq \epsilon$  and  $n \geq m$ .

**Step 6.** We now show how to use (34) in order to get the estimate on  $\dim_e(f(\operatorname{Crit}_{\nu}(f) \cap K))$ . For  $\epsilon > 0$  we choose  $n = n_{\epsilon}$  where

(35) 
$$n_{\epsilon} := \left\lceil \log_q \frac{c}{\epsilon} \right\rceil,$$

in such a way that (29), (30) and (34) hold when  $\epsilon \to 0$ . By (29) and (30) we have

$$\dim_{e} f(\operatorname{Crit}_{\nu}(f) \cap K) \leq \limsup_{\epsilon \to 0} \frac{\ln M(\epsilon, \tilde{f}_{n_{\epsilon}}(C^{\Lambda^{\epsilon}}(\tilde{f}_{n_{\epsilon}}) \cap B_{\mathbb{R}^{n_{\epsilon}}}(r)))}{\ln \frac{1}{2\epsilon}}$$

Hence now we estimate the right-hand side through (34). As  $\epsilon$  goes to zero we have

$$\limsup_{\epsilon \to 0} \frac{n_{\epsilon} \ln \beta_0}{\ln \frac{1}{2\epsilon}} = \lim_{\epsilon \to 0} \frac{\left\lceil \log_q \frac{c}{\epsilon} \right\rceil \ln \beta_0}{\ln \frac{1}{2\epsilon}} = \frac{\ln \beta_0}{\ln q}.$$

We also have, taking into account (35), that

$$\lim_{\epsilon \to 0} \frac{m \ln n_{\epsilon}}{\ln(\frac{1}{2\epsilon})} = 0.$$

Furthermore

$$\lim_{\epsilon \to 0} \frac{1}{\ln(\frac{1}{2\epsilon})} \ln\left(\sum_{h=0}^{\nu} \frac{\Lambda_0 \cdots \Lambda_h}{\epsilon^h}\right) = \nu.$$

Therefore,

 $\dim_e f(\operatorname{Crit}_\nu(f) \cap K) \leq \nu + \frac{\ln \beta_0}{\ln q},$ 

concluding the proof.

4.2. Sard-type theorems and Kolmogorov *n*-width. In this section, we recall the definition of Kolmogorov *n*-width, and we show its role in Sard-type theorems for maps from a Hilbert space to a Euclidean one.

**Definition 27** (Kolmogorov *n*-width). Let X be a normed space and  $S \subset X$  be a subset. For  $n \in \mathbb{N}$ , the Kolmogorov *n*-width of S in X is

$$\Omega_n(S, X) = \inf_{\dim E = n} \sup_{u \in S} \inf_{y \in E} ||u - y||,$$

where the infimum is taken over all n-dimensional linear subspaces E of X.

The asymptotics of n-width measures in a quantitative way the compactness of a set, in fact the following holds (see [Pin85, Prop. 1.2]).

**Proposition 28.** S is compact if and only if S is bounded and  $\lim_{n\to\infty} \Omega_n(S) = 0$ .

For various properties of the *n*-width we refer the reader to [Pin85]; we recall here those that we will need in the sequel, see [Pin85, Thm. 1.1].

**Theorem 29.** Let X be a normed space and  $S \subset X$ . Then for all  $n \in \mathbb{N}$ :

- (i)  $\Omega_n(S, X) = \Omega_n(\bar{S}, X)$ , where  $\bar{S}$  is the closure of S;
- (ii) For every  $\alpha \in \mathbb{R}$

 $\Omega_n(\alpha S, X) = |\alpha|\Omega_n(S, X);$ 

(iii) Let co(S) be the convex hull of S. Then

$$\Omega_n(\operatorname{co}(S), X) = \Omega_n(S, X);$$

(iv) Let  $b(S) = \{\alpha x \mid x \in S, |\alpha| \le 1\}$  be the balanced hull of S. Then

$$\Omega_n(\mathbf{b}(S), X) = \Omega_n(S, X).$$

Remark 30. As a consequence, in all our results, up to enlarging S, one can assume without loss of generality that S is convex and centrally symmetric, without changing its n-width.

We establish now a connection between *n*-width of compact sets and locally Lipschitz functions.

**Lemma 31.** Let H be a Hilbert space,  $(Y, \|\cdot\|_Y)$  a Banach space, and let  $f : H \to Y$  be locally Lipschitz. Let  $K \subset H$  be a compact subset. Then there exists  $n_0 \in \mathbb{N}$  such that for all  $n \ge n_0$  there exist a linear subspace  $E_n \subset H$  of dimension n such that

$$\sup_{x \in K} \|f(x) - f(\pi_{E_n}(x))\|_Y \le c(f, K)\Omega_n(K, H).$$

Proof. Let  $\{B_H(x_i, r/2)\}_{i \in I}$  a finite cover of K by balls, with centers  $x_i \in K$  and radii r/2 > 0. We can assume that f is L-Lipschitz on each  $B_H(x_i, r)$ , for some L > 0. Since K is compact  $\Omega_n(K, H) \to 0$  as  $n \to \infty$ , hence there exists  $n_0 \in \mathbb{N}$  such that for all  $n \ge n_0$  we have  $\Omega_n(K, H) < r/4$ . By definition of n-width, for all  $n \ge n_0$  there exists a n-dimensional subspace  $E_n \subset H$  such that

(36) 
$$\sup_{x \in K} \|x - \pi_{E_n}(x)\| < 2\Omega_n(K, H) < \frac{7}{2}$$

Let  $x \in K$ . Then by construction  $x \in B_H(x_i, r/2)$  for some  $i \in I$ . By (36) we have  $\pi_{E_n}(x) \in B_H(x_i, r)$ . Since f is L-Lipschitz on every  $B_H(x_i, r)$  we have

$$\sup_{x \in K} \|f(x) - f(\pi_{E_n}(x))\|_Y \le L \sup_{x \in K} \|x - \pi_{E_n}(x)\| \le 2L\Omega_n(K, H),$$

where we used again (36). This proves the result, with c(f, K) = 2L.

We can now prove the following Sard-type theorem, corresponding to Theorem A in the Introduction.

**Theorem 32.** Let  $d, m \in \mathbb{N}$ . There exists  $\beta_0 = \beta_0(d, m) > 0$  such that the following holds. Let H be a Hilbert space,  $f \in \mathscr{P}_d^m(H)$  and  $K \subset H$  be a compact set such that

$$\omega(K,H) = \limsup_{n \to \infty} \Omega_n(K,H)^{1/n} \le q^{-1} \in (0,1).$$

Then, for every  $\nu = 1, \ldots, m-1$  we have

$$\dim_e \left( f \left( \operatorname{Crit}_{\nu}(f) \cap K \right) \right) \le \nu + \frac{\ln \beta_0}{\ln q}$$

In particular, if  $q > \beta_0$ , then the Sard property holds on K:

$$\mu\bigg(f\big(\operatorname{Crit}(f)\cap K\big)\bigg)=0$$

*Proof.* Let  $q_{\epsilon} \in (0,q)$ . By assumption,  $\Omega_n(K,H) \leq q_{\epsilon}^{-n}$  for sufficiently large n. We prove that f satisfies the hypothesis of Theorem E. Since  $f \in \mathscr{P}_d^m(H)$  (see Definition 1), the map

$$(f, Df): H \to \mathbb{R}^m \times \mathcal{L}(H, \mathbb{R}^m)$$

is locally Lipschitz. The codomain  $Y = \mathbb{R}^m \times \mathcal{L}(H, \mathbb{R}^m)$  is a Banach space equipped with the norm  $||(v, A)||_Y = ||v|| + ||A||_{\text{op}}$  for  $v \in \mathbb{R}^m$  and  $A \in \mathcal{L}(H, \mathbb{R}^m)$ . Hence we can apply Lemma 31 to get  $n_0 \in \mathbb{N}$  such that for all  $n \ge n_0$  there exists a *n*-dimensional linear subspace  $E_n \subset H$  such that

$$\sup_{x \in K} \|f(x) - f(\pi_{E_n}(x))\| + \|D_x f - D_{\pi_{E_n(x)}} f\|_{\text{op}} \le c(f, K) q_{\epsilon}^{-n},$$

where  $\pi_{E_n} : H \to E_n$  denotes the orthogonal projection. Let  $f_n : E_n \to \mathbb{R}^m$  defined by the restriction  $f_n := f|_{E_n}$ , which are polynomial maps with degree uniformly bounded above by d. It holds

$$\|(D_x f)|_{E_n} - D_{\pi_{E_n}(x)} f_n\|_{\text{op}} = \|(D_x f)|_{E_n} - (D_{\pi_{E_n}(x)} f)|_{E_n}\|_{\text{op}} \le \|D_x f - D_{\pi_{E_n}(x)} f\|_{\text{op}}.$$

As a consequence of the above two estimates, assumption (3) of Theorem E holds, yielding

$$\dim_e \left( f \big( \operatorname{Crit}_{\nu}(f) \cap K \big) \right) \le \nu + \frac{\ln \beta_0}{\ln q_{\epsilon}}$$

Letting  $q_{\epsilon} \uparrow q$  we obtain the thesis.

4.3. Counterexamples to the Sard theorem in infinite dimension. In this section we provide examples of maps  $f \in \mathscr{P}_d^m(H)$  as in Theorem A for which there exists a compact set  $K \subset H$  such that the set  $f(\operatorname{Crit}(f) \cap K)$  has not measure zero, however it holds  $\Omega_n(K, H) \leq cq^{-n}$ . This shows the necessity of the quantitative assumption  $q > \beta_0(m, d)$ . As a consequence, we get lower bounds on the semialgebraic constant  $\beta_0$ .

We start with the following example, which is a minor modification of Kupka's one [Kup65].

*Example* 33 (Kupka revisited). Let q > 1. Let  $f : \ell^2 \to \mathbb{R}$  defined by

$$f(x) = \sum_{k=1}^{\infty} \frac{1}{2^k} \phi(q^{k-1}x_k),$$

where  $\phi$  is the Kupka polynomial as in [YC04, Sec. 10.2.3]. Namely  $\phi$  has degree 3, with  $\phi(0) = \phi'(0) = \phi(1) - 1 = \phi'(1) = 0$ . If we assume  $q^3/2 < 1$  then f is  $C^{\infty}$ . All critical points have the form

$$\operatorname{Crit}(f) = \left\{ \left( x_1, \frac{x_2}{q}, \dots, \frac{x_k}{q^{k-1}}, \dots \right) \middle| x_i \in \{0, 1\} \right\},\$$

and the set of critical values is [0,1]. Furthermore, given  $y = \left(x_1, \frac{x_2}{q}, \ldots, \frac{x_k}{q^{k-1}}, \ldots\right) \in \operatorname{Crit}(f)$ , we consider its projection  $y_n$  on  $\operatorname{span}\{e_1, \ldots, e_n\}$ , that is

$$y_n = \left(x_1, \frac{x_2}{q}, \dots, \frac{x_n}{q^{n-1}}, 0, 0, 0, \dots\right).$$

We have

$$||y - y_n||_{\ell^2}^2 \le \sum_{k=n+1}^\infty \frac{1}{q^{2(k-1)}} \le \int_n^\infty \frac{1}{q^{2t}} = \frac{q^{-2n}}{2\ln q}$$

Hence,

$$\Omega_n(\operatorname{Crit}(f),\ell^2) \le \sup_{y \in \operatorname{Crit}(f)} \|y - y_n\|_{\ell^2} \le \frac{q^{-n}}{\sqrt{2\ln q}} \qquad \Longrightarrow \qquad \omega(\operatorname{Crit}(f),\ell^2) \le q^{-1}.$$

Since f and  $\operatorname{Crit}(f)$  satisfy all the hypothesis of Theorem A (provided that  $q^3/2 < 1$ ), we get a consequence that

$$\beta_0(1,3) \ge 2^{1/3}$$

We now get the lower bound on  $\beta_0(1, d)$  for  $d \ge 3$  by improving on the above construction.

*Example* 34 (Kupka revisited, degree  $d \ge 3$ ). We need the following classical fact, which follows from [Kuh69]: for any  $d \ge 2$  there exists a polynomial  $\psi : \mathbb{R} \to \mathbb{R}$  of degree d such that

(37) 
$$\psi(\operatorname{Crit}(\psi)) = \{0, \dots, d-2\}.$$

(See also [Kri84, Thm. 1] which gives a version of this result taking into account also possible multiplicities.) We fix  $d \ge 3$  and we consider a polynomial  $\psi$  of degree d such that (37) holds. For q > 1 we consider the map  $f_d : \ell^2 \to \mathbb{R}$ 

$$f_d(x) := \sum_{k=1}^{\infty} \frac{1}{(d-1)^k} \psi(q^{k-1}x_k).$$

If  $q \in (1, (d-1)^{1/d})$ , then the map  $f_d$  is  $C^{\infty}$ . The set of critical points of  $f_d$  is

$$\operatorname{Crit}(f_d) = \left\{ \left( y_1, \frac{y_2}{q}, \dots, \frac{y_k}{q^{k-1}}, \dots \right) \in \ell^2 \, \middle| \, \psi'(y_k) = 0, \, \forall \, k \in \mathbb{N} \right\}.$$

Hence we have

$$f_d(\operatorname{Crit}(f_d)) = \left\{ \sum_{k=1}^{\infty} \frac{1}{(d-1)^k} c_k \, \middle| \, c_k \in \{0, \dots, d-2\} \right\} = [0, 1].$$

Let  $\zeta = \max\{|z| \mid \psi'(z) = 0\}$ . Exactly as in Example 33 we get for every  $n \in \mathbb{N}$ 

$$\Omega_n(\operatorname{Crit}(f_d), \ell^2) \leq \frac{\zeta}{\sqrt{2\ln q}} q^{-n} \qquad \Longrightarrow \qquad \omega(\operatorname{Crit}(f_d), \ell^2) \leq q^{-1}.$$

For any  $q \in (1, (d-1)^{1/d})$  the function  $f_d$  and the set  $K = \operatorname{Crit}(f_d)$  satisfy the hypothesis of Theorem 32, but  $f_d(\operatorname{Crit}(f_d)) = [0, 1]$ . It follows that

$$\beta_0(1,d) \ge (d-1)^{1/d}, \quad \forall d \ge 3.$$

In the next example we extend the construction to arbitrary dimension of the codomain.

Example 35 (Kupka revisited,  $d \ge 3$ , arbitrary codomain). Let  $f_d$  be as in Example 34, and consider  $g_d : \mathbb{R}^{m-1} \times \ell^2 \to \mathbb{R}^m$  defined as

$$g_d(x,y) := (x, f_d(y)).$$

We have

$$\operatorname{Crit}(g_d) = \mathbb{R}^{m-1} \times \operatorname{Crit}(f_d).$$

Taking  $E_{n+m-1} = \text{span}\{(e_i, 0), (0, e_j) \mid i = 1, \dots, m-1, j = 1, \dots, n\} \subset \mathbb{R}^{m-1} \times \ell^2$ , for all  $n \in \mathbb{N}$ , we can estimate the *n*-width of  $\text{Crit}(g_d)$  as

$$\Omega_{n+m-1}(\operatorname{Crit}(g_d), \mathbb{R}^{m-1} \times \ell^2) \le \frac{\zeta}{\sqrt{2\ln q}} q^{-n} \qquad \Longrightarrow \qquad \omega(\operatorname{Crit}(g_d), \mathbb{R}^{m-1} \times \ell^2) \le q^{-1}.$$

We observe, provided that  $q \in (1, (d-1)^{1/d})$ , the function  $g_d$  satisfies all the hypothesis of Theorem 32, but the set  $g_d(\operatorname{Crit}(g_d)) = \mathbb{R}^{m-1} \times [0,1]$  has not measure zero. Hence

$$\beta_0(d,m) \ge (d-1)^{1/d}, \qquad \forall m \in \mathbb{N}, \, d \ge 3.$$

We collect the content of Examples 33 to 35 in a unified statement (see Theorem B).

**Theorem 36.** Let  $d, m \in \mathbb{N}$ , with  $d \geq 3$ , and set  $q := (d-1)^{1/d}$ . There exist a Hilbert space H, and  $f \in \mathscr{P}_d^m(H)$  such that  $K = \operatorname{Crit}(f) \cap B_H(r)$  is compact for all r > 0, with

$$\omega(K,H) = \limsup_{n \to \infty} \Omega_n(K,H)^{1/n} \le q^{-1},$$

and  $f: H \to \mathbb{R}^m$  does not verify the Sard property, namely  $\mu(f(\operatorname{Crit}(f) \cap K)) > 0$ . Therefore, the semialgebraic constant  $\beta_0(d,m)$  of Theorem E satisfies

$$\beta_0(d,m) \ge (d-1)^{1/d}, \qquad \forall m \in \mathbb{N}, \, d \ge 3.$$

In all the examples above we have taken K to be set of all critical points of a given function in a ball. We now show a different construction, where the set K is strictly contained in the set of critical points, more precisely is the set of points where the differential has rank zero.

*Example* 37 (Rank zero counterexample to Sard). We consider  $f_d: \ell^2 \to \mathbb{R}$  as in Example 34, and  $m \in \mathbb{N}$ . We define  $h: (\ell^2)^m \to \mathbb{R}^m$  as

$$h(x^1, \dots, x^m) := (f_d(x^1), \dots, f_d(x^m)).$$

For any  $x = (x^1, \ldots, x^m) \in (\ell^2)^m$  we have

$$\operatorname{im}(D_x h) = \operatorname{im}(D_{x^1} f_d) \times \cdots \times \operatorname{im}(D_{x^m} f_d).$$

Hence  $x \in \operatorname{Crit}_{\nu}(h)$  (the set of points where the rank of the differential is  $< \nu$ ) if and only if at least  $m-\nu$  components of x are critical points of  $f_d$ . In particular, we consider the compact set K of all critical points of h of rank zero, namely

$$K := \operatorname{Crit}_0(h) = \operatorname{Crit}(f_d)^m.$$

We have  $K \subsetneq \operatorname{Crit}(h)$ , and

$$h(\operatorname{Crit}(h) \cap K) = h(\operatorname{Crit}_0(h)) = [0, 1]^m$$

As in the previous examples, K is compact with exponential n-width. By taking finite-dimensional spaces  $E_{nm} = \operatorname{span}\{\underbrace{(0,\ldots,e_i,\ldots,0)}_{j\text{-th component}} \in (\ell^2)^m \mid i = 1,\ldots,n, j = 1,\ldots,m\}$ , for  $n \in \mathbb{N}$  we have

$$\Omega_{nm}(K,(\ell^2)^m) \le m\Omega_n(\operatorname{Crit}(f_d),\ell^2) \le \frac{m\zeta}{\sqrt{2\ln q}}q^{-n} \qquad \Longrightarrow \qquad \omega(K,(\ell^2)^m) \le q^{-1/m},$$

where  $\zeta$  is the same constant appearing in Example 34.

4.4. Sard threshold theorems. We can now prove Theorem C, of which we recall the statement.

**Theorem 38.** For every  $d, m \in \mathbb{N}$ , there exists  $\omega_0(d, m) \in (0, 1]$  such that

(i) for every  $f \in \mathscr{P}_d^m(H)$  and for every compact set  $K \subset H$  with  $\omega(K, H) < \omega_0(d, m)$ ,

$$\mu\bigg(f\left(\operatorname{crit}(f)\cap K\right)\bigg)=0;$$

(ii) for every  $\omega > \omega_0(d,m)$ , with  $\omega \in (0,1]$ , there exist  $f \in \mathscr{P}_d^m(H)$  and a compact set  $K \subset H$ with  $\omega(K, H) = \omega$  and such that

$$\mu\bigg(f\left(\operatorname{crit}(f)\cap K\right)\bigg)>0.$$

*Proof.* By definition, for any compact set K, it holds  $\omega(K, H) \in [0, 1]$ , see (2). We define  $\omega_0(d, m)$  as the supremum of the set of  $\omega \in [0, 1]$  such that for every  $f \in \mathscr{P}_d^m(H)$  and for every compact set  $K \subset H$  with  $\omega(K, H) = \omega$  it holds:

$$\mu\bigg(f\left(\operatorname{Crit}(f)\cap K\right)\bigg)=0.$$

By Theorem A this set is non-empty and  $\omega_0(d,m) \ge \beta_0(d,m)^{-1}$ . Hence  $\omega_0(d,m) \in (0,1]$ . Both items of Theorem 38 follow. Note that by Theorem B (more specifically, the construction of Example 35), provided that  $d \ge 3$ , one must have  $\omega_0(d,m) \le (d-1)^{-1/d} < 1$ , so that Item (ii) is non-vacuous.  $\Box$ 

Next, we prove Theorem D, of which we recall the statement.

**Theorem 39.** Let  $f \in \mathscr{P}_d^m(H)$  and  $K \subset H$  be a compact set such that  $\omega(K, H) < \omega_0(d, m)$ . Consider the linear subspace

$$V := \operatorname{span}(K).$$

Then the restriction  $f|_V: V \to \mathbb{R}^m$  satisfies the Sard property:

$$\mu\bigg(f\big(\operatorname{Crit}(f|_V)\big)\bigg) = 0$$

In particular,  $\mu(f(\operatorname{Crit}(f) \cap V)) = 0.$ 

*Proof.* Let H denote the closure of V in H. Observe first that

$$\Omega_n(K,H) = \Omega_n(K,\tilde{H}).$$

The inequality  $\Omega_n(K, H) \leq \Omega_n(K, \tilde{H})$  is clear since every *n*-dimensional subspace of  $\tilde{H}$  is also a subspace of H. For the other inequality, since  $\tilde{H}$  is closed in H, we can write  $H = \tilde{H} \oplus \tilde{H}^{\perp}$  and denote by  $\pi: H \to \tilde{H}$  the orthogonal projection. Then, using that  $K \subset V \subset \tilde{H}$ , we obtain

$$\begin{aligned} \Omega_n(K,H) &= \inf_{\substack{Y \subset H \\ \dim Y = n}} \sup_{\substack{u \in K}} \inf_{y \in Y} \|u - y\| \ge \inf_{\substack{Y \subset H \\ \dim Y = n}} \sup_{u \in K} \inf_{y \in Y} \|u - \pi(y)\| \\ &= \inf_{\substack{Y \subset H \\ \dim Y = n}} \sup_{u \in K} \sup_{z \in \pi(Y)} \|u - z\| \\ &= \inf_{\substack{Z \subset \tilde{H} \\ \dim Z \le n}} \sup_{u \in K} \inf_{z \in Z} \|u - z\| \\ &= \inf_{\substack{Z \subset \tilde{H} \\ \dim Z = n}} \sup_{u \in K} \inf_{z \in Z} \|u - z\| = \Omega_n(K, \tilde{H}). \end{aligned}$$

We can apply Item (i) of Theorem C to the map  $\tilde{f} \in \mathscr{P}_d^m(\tilde{H})$ , defined by  $\tilde{f} := f|_{\tilde{H}}$ , with  $f \in \mathscr{P}_d^m(H)$ , and the compact set  $K \subset \tilde{H}$ , which satisfies  $\omega(K, \tilde{H}) < \omega_0(d, m)$ , obtaining

$$\mu\left(\tilde{f}(\operatorname{Crit}(\tilde{f})\cap K)\right) = 0.$$

Observe now that if u is a critical point for  $g := f|_V$ , then u is also critical for  $\tilde{f} = f|_{\tilde{H}}$ . In fact,  $D_u g = (D_u f)|_V$  for all  $u \in V$ . Thus we have  $\lambda \in (\mathbb{R}^m)^* \setminus \{0\}$  such that  $0 \equiv \lambda \circ D_u g = (\lambda \circ D_u \tilde{f})|_V$ and, since V is dense in  $\tilde{H}$ , it follows that  $\lambda \circ D_u \tilde{f} \equiv 0$ . This means that

$$\operatorname{Crit}(g) \cap K \subseteq \operatorname{Crit}(\tilde{f}) \cap K$$

In particular, since on  $K \subset V$  we have  $\tilde{f} = g$  by definition, this implies that

(38) 
$$0 = \mu\left(\tilde{f}(\operatorname{Crit}(\tilde{f}) \cap K)\right) = \mu\left(g(\operatorname{Crit}(g) \cap K)\right)$$

Let K' be the balanced and convex hull of K. It holds:

$$V = \operatorname{span}(K) = \operatorname{span}(K') = \bigcup_{j=1}^{\infty} jK'.$$

By Theorem 29 the *n*-width of the compact, convex, centrally symmetric sets jK' satisfy

$$\Omega_n(jK',H) = j\Omega_n(K',H) = j\Omega_n(K,H).$$

Thus  $\omega(jK', H) = \omega(K, H)$  for all  $j \in \mathbb{N}$ . Then, (38) holds also for jK' in place of K. Therefore,

$$\mu\left(g(\operatorname{Crit}(g))\right) \leq \sum_{j\geq 1} \mu\left(\tilde{f}\left(\operatorname{Crit}(\tilde{f}) \cap jK'\right)\right) = 0.$$

Finally, since  $\operatorname{Crit}(f) \cap V \subset \operatorname{Crit}(g)$ , and  $g = f|_V$ , it also holds  $\mu(f(\operatorname{Crit}(f) \cap V)) = 0$ .

4.5. A class of maps with the global Sard property. Using Theorem E, we study a class of maps from a Hilbert space to  $\mathbb{R}^m$  for which the Sard property holds true. The case m = 1 is not new, and it was proved in [Yom88], see also [YC04]. We need the following preliminary facts.

**Lemma 40** (Markov inequality). Let  $p : \mathbb{R}^n \to \mathbb{R}^m$  be a polynomial map with components of degree at most d. Then for every r > 0 it holds

$$\sup_{x \in B_{\mathbb{R}^n}(r)} \|D_x p\|_{\text{op}} \le \frac{\sqrt{md^2}}{r} \sup_{x \in B_{\mathbb{R}^n}(r)} \|p(x)\|.$$

*Proof.* From [Kel28, Thm. VI] we get the thesis for r = 1. Then, we conclude considering the rescaled polynomial  $x \to p(rx)$ .

**Lemma 41** (Estimates on balls at different radii). Let  $p : \mathbb{R}^n \to \mathbb{R}^m$  be a polynomial map with components of degree at most d. Then for all r > 0 it holds

$$\sup_{x \in B_{\mathbb{R}^n}(r)} \|p(x)\| \le \alpha(d,m) n^d (1+r)^d \sup_{x \in B_{\mathbb{R}^n}(1)} \|p(x)\|,$$

where  $\alpha(d,m) = m^{d/2} d^{2d} (d+1)$  is a constant depending only on d and m.

*Proof.* We can write the polynomial map as

$$p(x) = \sum_{|\alpha| \le d} c_{\alpha} x^{\alpha}$$

where  $c_{\alpha} \in \mathbb{R}^{m}$ , and the sum is over the multi-indices  $\alpha \in \mathbb{N}^{n}$  with  $|\alpha| = \sum_{i} \alpha_{i} \leq d$ . Since  $|\alpha|!c_{\alpha} = \partial_{x^{\alpha}}p|_{x=0}$ , we can iterate Markov inequality (for r = 1, see Lemma 40) and obtain

$$\|c_{\alpha}\| \leq \frac{1}{|\alpha|!} m^{|\alpha|/2} d^{2|\alpha|} \sup_{x \in B_{\mathbb{R}^n}(1)} \|p(x)\| \leq m^{|\alpha|/2} d^{2|\alpha|} \sup_{x \in B_{\mathbb{R}^n}(1)} \|p(x)\|.$$

Therefore, for r > 1, we have

$$\sup_{x \in B_{\mathbb{R}^{n}}(r)} \|p(x)\| = \sup_{x \in B_{\mathbb{R}^{n}}(1)} \|p(rx)\|$$
$$= \sup_{x \in B_{\mathbb{R}^{n}}(1)} \left\| \sum_{i=1}^{d} r^{i} \left( \sum_{|\alpha|=i} c_{\alpha} x^{\alpha} \right) \right\|$$
$$\leq (1+r)^{d} \left( \sum_{|\alpha|\leq d} m^{|\alpha|/2} d^{2|\alpha|} \right) \sup_{x \in B_{\mathbb{R}^{n}}(1)} \|p(x)\|$$
$$\leq (1+r)^{d} m^{d/2} d^{2d} (d+1) n^{d} \sup_{x \in B_{\mathbb{R}^{n}}(1)} \|p(x)\|,$$

where we used the fact that the number of multi-indices  $\alpha \in \mathbb{N}^n$  with  $|\alpha| \leq d$  is equal to  $(1 + n + n^2 + \cdots + n^d) \leq (d+1)n^d$ 

For the rest of this section, we assume that the Hilbert space H is separable, and we fix a Hilbert basis  $\{e_i\}_{i\in\mathbb{N}} \subset H$ . For  $k \in \mathbb{N}$ , we set  $E_k := \operatorname{span}\{e_1, \ldots, e_k\}$  and we denote by  $\pi_k : H \to E_k$  the corresponding orthogonal projection, that is if  $x = \sum_i x_i e_i$ , then  $\pi_k(x) = (x_1, \ldots, x_k)$ . We now define a special class of maps in this setting.

**Proposition 42** (Construction of special maps). Let H be a separable Hilbert space. For all  $k \in \mathbb{N}$ , let  $p_k : E_k \to \mathbb{R}^m$  be polynomial maps with  $\sup_{k \in \mathbb{N}} \deg p_k \leq d$  for some  $d \in \mathbb{N}$ , and such that

(39) 
$$\sum_{k=1}^{\infty} \sup_{x \in B_{E_k}(r)} \|p_k(x)\| < \infty, \qquad \forall r > 0.$$

Then the map  $f: H \to \mathbb{R}^m$  given by

(40) 
$$f(x) := \sum_{k=1}^{\infty} p_k(x_1, \dots, x_k), \quad \forall x \in H,$$

is well-defined,  $f \in \mathscr{P}_d^m(H)$  (see Definition 1) and its differential is given by

(41) 
$$D_x f = \sum_{k=1}^{\infty} D_x (p_k \circ \pi_k)$$

*Proof.* For a Banach spaces  $(X, \|\cdot\|_X)$ ,  $(Y, \|\cdot\|_Y)$ , and a closed set  $U \subseteq X$ , we denote by  $\mathcal{B}(U, Y)$  the space of all continuous and bounded functions  $f: U \to Y$ . It is a Banach space when endowed with the norm  $\sup_{x \in U} \|f(x)\|_Y$ .

We first note that the assumption (39) implies that the sequence of maps  $f_n : H \to \mathbb{R}^m$  given by  $f_n := \sum_{k=1}^n p_k \circ \pi_k$  is Cauchy in  $\mathcal{B}(B_H(r), \mathbb{R}^m)$  for all r > 0. In particular the right hand side of (40) converges locally uniformly. For any  $n \in N$  we have

(42) 
$$D_x f_n = \sum_{k=1}^n D_x (p_k \circ \pi_k).$$

Furthermore,  $x \mapsto D_x f_n \in \mathcal{B}(B_H(r); \mathcal{L}(H, \mathbb{R}^m))$  for all  $n \in \mathbb{N}, r > 0$ . We now prove that it is a Cauchy sequence. Indeed, using Lemma 40, for any  $n_2 \ge n_1$ 

$$\sup_{x \in B_H(r)} \|D_x f_{n_2} - D_x f_{n_1}\|_{\text{op}} \le \frac{\sqrt{m}d^2}{r} \sum_{k > n_1} \sup_{x \in B_{E_k}(r)} \|p_k(x)\| \to 0,$$

as  $n_2, n_1 \to \infty$ . Therefore,  $\{Df_n\}_{n \in \mathbb{N}}$  has a limit in  $\mathcal{B}(B_H(r), \mathcal{L}(H, \mathbb{R}^m))$ , for all r > 0. It follows that f is  $\mathcal{C}^1$  and (41) holds, see e.g. [Lan93, Thm. 9.1].

Thanks to (41), we prove that  $Df : H \to \mathcal{L}(H, \mathbb{R}^m)$  is weakly continuous. Let us consider a sequence  $\{x^j\}_{j\in\mathbb{N}}$  weakly convergent to  $x \in B_H(r)$ ; we prove that  $D_{x^j}f \to D_xf$  as  $j \to \infty$ . Indeed, given  $\epsilon > 0$  we fix  $N_{\epsilon} \in \mathbb{N}$  such that  $\sum_{k>N_{\epsilon}} \sup_{B_{E_k}(r)} \|p_k\| \leq \epsilon$  and, by Lemma 40, we get

$$\begin{split} \|D_{x^{j}}f - D_{x}f\|_{\mathrm{op}} &\leq \sum_{k=1}^{\infty} \|D_{x^{j}}(p_{k} \circ \pi_{k}) - D_{x}(p_{k} \circ \pi_{k})\|_{\mathrm{op}} \\ &= \sum_{k=1}^{\infty} \|(D_{\pi_{k}(x^{j})}p_{k}) \circ \pi_{k} - (D_{\pi_{k}(x)}p_{k}) \circ \pi_{k}\|_{\mathrm{op}} \\ &= \sum_{k=1}^{\infty} \|D_{\pi_{k}(x^{j})}p_{k} - D_{\pi_{k}(x)}p_{k}\|_{\mathrm{op}} \\ &\leq \sum_{k=1}^{N_{\epsilon}} \|D_{\pi_{k}(x^{j})}p_{k} - D_{\pi_{k}(x)}p_{k}\|_{\mathrm{op}} + \sum_{k>N_{\epsilon}} \|D_{\pi_{k}(x^{j})}p_{k}\|_{\mathrm{op}} + \sum_{k>N_{\epsilon}} \|D_{\pi_{k}(x)}p_{k}\|_{\mathrm{op}} \\ &\leq \sum_{k=1}^{N_{\epsilon}} \|D_{\pi_{k}(x^{j})}p_{k} - D_{\pi_{k}(x)}p_{k}\|_{\mathrm{op}} + \frac{\sqrt{md^{2}}}{r} 2\epsilon, \end{split}$$

where with some abuse of notation we denoted by the same symbol  $\|\cdot\|_{\text{op}}$  either the norm in  $\mathcal{L}(H, \mathbb{R}^m)$ or the one for the finite-dimensional subspaces  $\mathcal{L}(E_k, \mathbb{R}^m)$ . Since  $\pi_k(x^j) \to \pi_k(x)$  as  $j \to \infty$  for all  $k \in \mathbb{N}$ , we conclude by the continuity of  $Dp_k$  on  $E_k$ . We prove now that  $Df : H \to \mathcal{L}(H, \mathbb{R}^m)$  is locally Lipschitz. We will prove that the sequence  $\{Df_n\}_{n\in\mathbb{N}}$  defined by (42) is uniformly Lipschitz on balls. For  $k \in \mathbb{N}$ , fix  $z, w \in E_k$ . Then it holds

(43) 
$$D_z p_k - D_w p_k = D_z p_k - D_z \tilde{p}_k = D_z (p_k - \tilde{p}_k),$$

where  $\tilde{p}_k : E_k \to \mathbb{R}^m$ , defined by  $\tilde{p}_k(\cdot) := p_k(\cdot - z + w)$ , is polynomial depending on the fixed choices of z, w, with degree bounded by d. Thus for all fixed  $z, w \in B_{E_k}(r), r > 0$ , it holds

$$\begin{split} \|D_z p_k - D_w p_k\|_{\text{op}} &\leq \sup_{a \in B_{E_k}(r)} \|D_a(p_k - \tilde{p}_k)\|_{\text{op}} & \text{by (43)} \\ &\leq \frac{\sqrt{m}d^2}{r} \sup_{a \in B_{E_k}(r)} \|p_k(a) - \tilde{p}_k(a)\| & \text{by Lemma 40} \end{split}$$

$$\leq \frac{\sqrt{md^2}}{r} \sup_{a \in B_{E_k}(r)} \|p_k(a) - p_k(a - z + w)\| \qquad \text{by definition of } \tilde{p}_k$$

$$\leq \left(\frac{\sqrt{md^2}}{r} \sup_{a \in B_{E_k}(3r)} \|D_a p_k\|_{\text{op}}\right) \|z - w\| \qquad p_k \text{ is Lipschitz}$$

(44) 
$$\leq \left(\frac{md^4}{r^2} \sup_{a \in B_{E_k}(3r)} \|p_k(a)\|\right) \|z - w\|.$$
 by Lemma 40

Note that if  $x \in B_H(r)$  then  $\pi_k(x) \in B_{E_k}(r)$ . Thus, for all  $x, y \in B_H(r)$  it holds

$$\|D_x f_n - D_y f_n\|_{\text{op}} \le \sum_{k=1}^n \|D_x (p_k \circ \pi_k) - D_y (p_k \circ \pi_k)\|_{\text{op}}$$
 by (42)

$$\leq \sum_{k=1}^{n} \|D_{\pi_k(x)} p_k - D_{\pi_k(y)} p_k\|_{\text{op}} \qquad \text{by Leibniz rule}$$

$$\leq \frac{md^4}{r^2} \sum_{k=1}^n \sup_{a \in B_{E_k}(3r)} \|p_k(a)\| \|\pi_k(x) - \pi_k(y)\| \qquad \text{by (44)}$$

$$< \frac{md^4}{r} \|r - y\| \sum_{k=1}^\infty \sup_{a \in B_{E_k}(3r)} \|p_k(a)\| = \pi_k \text{ is 1-Lipschitz}$$

$$\leq \frac{md^{*}}{r^{2}} \|x - y\| \sum_{k=1} \sup_{a \in B_{E_{k}}(3r)} \|p_{k}(a)\| \qquad \pi_{k} \text{ is 1-Lipschitz}$$
  
$$\leq c(r) \|x - y\|,$$

where c(r) > 0 does not depend on n, and in the last inequality we used (39) for the convergence of the series. Therefore the sequence  $\{Df_n\}_{n\in\mathbb{N}}$ , is uniformly Lipschitz on every ball  $B_H(r)$ . Thus its limit  $Df: H \to \mathcal{L}(H, \mathbb{R}^m)$  is Lipschitz on any ball  $B_H(r)$ , and a fortiori locally Lipschitz.

Finally, we show that for every finite dimensional space  $E \subset H$ , the map  $f|_E$  is a polynomial map with components of degree bounded by d. Let  $n = \dim(E)$  and fix a linear isometry  $L : \mathbb{R}^n \to E$ . The statement is equivalent to show that  $f \circ L$  is a polynomial map with components of degree bounded by d. Denoting by  $(t_1, \ldots, t_n)$  the standard coordinates in  $\mathbb{R}^n$ , we have

$$f(L(t_1,...,t_n)) = \sum_{k=1}^{\infty} p_k(x_1(t_1,...,t_n),...,x_k(t_1,...,t_n))$$
  
=  $\sum_{k=1}^{\infty} \tilde{p}_k(t_1,...,t_n),$ 

where each  $\tilde{p}_k$  is a polynomial in  $(t_1, \ldots, t_n)$  of degree at most d (since the  $x_i$  depend linearly on the  $t_j$ ). Condition (39) guarantees that, for each monomial  $t_1^{\alpha_1} \cdots t_n^{\alpha_n}$ , its coefficient in the expansion  $\sum_{k \leq s} \tilde{p}_k$  converges as  $s \to \infty$ . Note that there are only a finite number of such coefficients, determined by the upper bound d on the degree and the dimension n. We conclude by noting that the uniform limit of polynomials with degree bounded by d is a polynomial with degree bounded by d.

With the following we exhibit a special class of maps, obtained via the construction of Proposition 42, for which the Sard property holds true *globally*. This is the content of Theorem F in the introduction, of which we recall the statement here.

**Theorem 43.** Let H be a separable Hilbert space. For all  $k \in \mathbb{N}$ , let  $p_k : E_k \to \mathbb{R}^m$  be polynomial maps with  $\sup_{k \in \mathbb{N}} \deg p_k \leq d$  for some  $d \in \mathbb{N}$ , and such that

(45) 
$$\sup_{x \in B_{E_k}(1)} \|p_k(x)\| \le q^{-k}, \qquad \forall k \in \mathbb{N},$$

for some q > 1. Then the map  $f : H \to \mathbb{R}^m$  defined by

$$f(x) := \sum_{k=1}^{\infty} p_k(x_1, \dots, x_k), \qquad \forall x \in H,$$

is well-defined,  $f \in \mathscr{P}_d^m(H)$  (see Definition 1), and for all  $\nu = 1, \ldots, m-1$  and r > 0 it holds

$$\dim_e \left( f\left(\operatorname{Crit}_{\nu}(f) \cap B_H(r)\right) \right) \le \nu + \frac{\ln \beta_0}{\ln q},$$

where  $\beta_0 = \beta_0(d, m)$  is the same constant given by Theorem E. In particular, if  $q > \beta_0$ , then f satisfies the Sard property globally on H:

$$\mu\bigg(f\big(\operatorname{Crit}(f)\big)\bigg) = 0.$$

*Proof.* We show first that the map f is well-defined. The upper bound (45), together with Lemma 41 imply that for all r > 0

(46) 
$$\sup_{x \in B_{E_k}(r)} \|p_k(x)\| \le \alpha(d,m)(1+r)^d \frac{k^d}{q^k}, \qquad \forall k \in \mathbb{N},$$

with q > 1. In turn, this implies that

$$\sum_{k=1}^{\infty} \sup_{x \in B_{E_k}(r)} \|p_k(x)\| < \infty, \qquad \forall r > 0.$$

We can then apply Proposition 42 obtaining that  $f \in \mathscr{P}_d^m(H)$ .

Fix r > 0. We now prove that f satisfies the hypothesis of Theorem **E** with  $K = B_H(r)$  and polynomial approximating maps  $f_n : E_n \to \mathbb{R}^m$ 

$$f_n(x) := \sum_{k=1}^n p_k(x_1, \dots, x_k), \qquad \forall x \in H$$

Note that, by our assumptions,  $\sup_{n \in \mathbb{N}} \deg f_n \leq d$ . It remains to check the validity of (3). Firstly, we have for all  $n \in \mathbb{N}$ 

$$\sup_{K} \|f(x) - f_n \circ \pi_{E_n}(x)\| \le \alpha(d, m)(1+r)^d \sum_{k=n+1}^{\infty} \frac{k^d}{q^k}$$

where we used (46).

We now estimate the derivatives. We have

$$\sup_{x \in K} \left\| (D_x f) |_{E_n} - D_{\pi_{E_n}(x)} f_n \right\|_{\text{op}} \leq \sup_{x \in K} \left\| D_x f - D_x (f_n \circ \pi_{E_n}) \right\|_{\text{op}} \leq \sup_{x \in K} \sum_{k=n+1}^{\infty} \left\| D_x (p_k \circ \pi_k) \right\|_{\text{op}} \quad \text{by (41)} \\
= \sup_{x \in K} \sum_{k=n+1}^{\infty} \left\| (D_{\pi_k(x)} p_k) \circ \pi_k \right\|_{\text{op}} \\
\leq \sum_{k=n+1}^{\infty} \sup_{x \in B_{E_k}(r)} \left\| D_x p_k \right\|_{\text{op}}.$$

Note that, in the last line, the supremum is taken on a finite-dimensional Euclidean ball. To proceed, by Lemma 40 and (46) we have

$$\sup_{x \in B_{E_k}(r)} \left\| D_x p_k \right\|_{\text{op}} \le \frac{d^2 \sqrt{m}}{r} \alpha(d, m) (1+r)^d \frac{k^d}{q^k}$$

Therefore, continuing the previous estimate, we obtain

$$\sup_{x \in K} \left\| (D_x f) \right\|_{E_n} - D_{\pi_{E_n}(x)} f_n \right\|_{\text{op}} \le \frac{d^2 \sqrt{m}}{r} \alpha(d, m) (1+r)^d \sum_{k=n+1}^{\infty} \frac{k^d}{q^k}.$$

We have therefore proved that for all  $n \in \mathbb{N}$ , and r > 0 it holds

$$\sup_{x \in K} \left( \|f(x) - f_n \circ \pi_{E_n}(x)\| + \|(D_x f)\|_{E_n} - D_{\pi_{E_n}(x)} f_n\|_{\mathrm{op}} \right) \le \left(1 + \frac{d^2 \sqrt{m}}{r}\right) \alpha(d, m)(1+r)^d \sum_{k=n+1}^{\infty} \frac{k^d}{q^k}.$$

To conclude, note that by elementary estimates it holds

$$\left(1+\frac{d^2\sqrt{m}}{r}\right)\alpha(d,m)(1+r)^d\sum_{k=n+1}^{\infty}\frac{k^d}{q^k}\leq cq^{-n},$$

for sufficiently large n, where  $c \ge 1$  is a constant depending only on the fixed parameters r, d, m. Therefore, the main assumption (3) of Theorem E is satisfied, yielding

$$\dim_e \left( f \left( \operatorname{Crit}_{\nu}(f) \cap B_H(r) \right) \right) \leq \nu + \frac{\ln \beta_0}{\ln q}, \qquad \forall \nu = 1, \dots, m-1.$$

In particular, if  $q > \beta_0$  then f satisfies the Sard property on  $B_H(r)$ , namely

$$\mu\bigg(f\big(\operatorname{Crit}(f)\cap B_H(r)\big)\bigg)=0$$

Since r > 0 is arbitrary, we obtain that  $\mu(f(\operatorname{Crit}(f))) = 0$ .

# 5. Applications to the Endpoint maps of Carnot groups

We apply in this last section the results from the previous sections to the study of the Sard property for Endpoint maps for Carnot groups.

5.1. Carnot groups. An *m*-dimensional Carnot group of step  $s \in \mathbb{N}$  is a connected and simply connected Lie group  $(\mathbb{G}, \cdot)$  of dimension *m*, whose Lie algebra of left-invariant vector fields  $\mathfrak{g}$  admits a stratification of step *s*, that is

$$\mathfrak{g}=\mathfrak{g}_1\oplus\cdots\oplus\mathfrak{g}_s,$$

where  $\mathfrak{g}_i \neq \{0\}$ ,  $\mathfrak{g}_{i+1} = [\mathfrak{g}_1, \mathfrak{g}_i]$  for all  $i = 1, \ldots, s - 1$  and  $[\mathfrak{g}_1, \mathfrak{g}_s] = \{0\}$ . We set  $k_i = \dim \mathfrak{g}_i$ ,  $a_i = k_1 + \cdots + k_i$ , and  $k = k_1$ , which is called the *rank* of the Carnot group. As usual, we identify elements of  $\mathfrak{g}$  with vectors of  $T_e \mathbb{G}$ . Since  $\mathfrak{g}$  is nilpotent and  $\mathbb{G}$  is simply connected, the group exponential map  $\exp_{\mathbb{G}} : \mathfrak{g} \to \mathbb{G}$  is a smooth diffeomorphism.

The stratification (47) allows the definition of a family of automorphisms  $\delta_{\lambda} : \mathbb{G} \to \mathbb{G}$ , for  $\lambda > 0$ , called *dilations* such that

$$\delta_{\lambda}(\exp_{\mathbb{G}}(v)) = \exp_{\mathbb{G}}(\lambda^{i}v), \quad \forall v \in \mathfrak{g}_{i}, \quad i = 1, \dots, s$$

Fix an *adapted basis* of  $\mathfrak{g}$ , namely a set of vectors  $v_1, \ldots, v_m \in \mathfrak{g}$  such that for all  $i = 1, \ldots, s$  the list of vectors  $v_{k_{i-1}+1}, \ldots, v_{k_i}$  is a basis of  $\mathfrak{g}_i$ , with the convention  $k_0 = 0$ . For any  $j = 1, \ldots, m$  the *weight* of  $v_j$  is the unique  $w_j \in \{1, \ldots, s\}$  such that  $v_j \in \mathfrak{g}_{w_j}$ . We define a diffeomorphism  $\varphi_{\mathbb{G}} : \mathbb{R}^m \to \mathbb{G}$ , by

$$\varphi_{\mathbb{G}}(x_1,\ldots,x_m) := \exp_{\mathbb{G}}\left(\sum_{i=1}^m x_i v_i\right).$$

The map  $\varphi_{\mathbb{G}}$  defines global coordinates on  $\mathbb{G}$ , called *exponential coordinates* (of first kind). For any  $x \in \mathbb{G}$  we denote by  $\tau_x : \mathbb{G} \to \mathbb{G}$  the left translation on  $\mathbb{G}$ , that is  $\tau_x(y) = x \cdot y$ . We denote by

 $X_1, \ldots, X_m$  the left-invariant vector fields on  $\mathbb{G}$  corresponding to the adapted basis. They are given by  $X_j(x) = D_e \tau_x(v_j)$  for any  $x \in \mathbb{G}$ , where  $D_e \tau_x$  denotes the differential of  $\tau_x$  at the unit element.

We recall the definition of (homogeneous) polynomials on Carnot groups. We say that  $P : \mathbb{G} \to \mathbb{R}$  is a polynomial if for some (and then any) set of exponential coordinates  $\varphi_{\mathbb{G}}$ , the map  $P \circ \varphi_{\mathbb{G}} : \mathbb{R}^m \to \mathbb{R}$ is a polynomial. Any polynomial  $P : \mathbb{G} \to \mathbb{R}$  can be written as

$$P(\varphi_{\mathbb{G}}(x_1,\ldots,x_m)) = \sum_{\alpha \in \mathbb{N}^m} c_{\alpha} x_1^{\alpha_1} \cdots x_m^{\alpha_m}, \qquad \forall x \in \mathbb{R}^m,$$

for some  $c_{\alpha} \in \mathbb{R}$ , which are non-zero only for a finite set of multi-indices. For  $\alpha \in \mathbb{N}^m$ , we denote by

$$|\alpha|_{\mathbb{G}} := \sum_{i=1}^m w_i \alpha_i,$$

the weighted degree of the multi-index  $\alpha$ . The weighted degree of a polynomial P is

$$\deg_{\mathbb{G}}(P) := \max\left\{ |\alpha|_{\mathbb{G}} \mid c_{\alpha} \neq 0 \right\}$$

A polynomial P is homogeneous of weighted degree w if  $P(\delta_{\lambda} x) = \lambda^w P(x)$  for any  $x \in \mathbb{G}$  and  $\lambda > 0$ .

Remark 4. In other words, P is homogeneous of weighted degree w if and only if

$$P(\varphi_{\mathbb{G}}(x_1,\ldots,x_m)) = \sum_{\substack{\alpha \in \mathbb{N}^m \\ |\alpha|_{\mathbb{G}} = w}} c_{\alpha} x_1^{\alpha_1} \cdots x_m^{\alpha_m}.$$

In particular any such polynomial depends only on the variables  $x_i$  with  $w_i \leq w$ , namely  $x_1, \ldots, x_{k_w}$ .

It is easy to see that the concepts we just introduced do not depend on the choice of exponential coordinates. For this reason, in the following, we fix such a choice, and we omit  $\varphi_{\mathbb{G}}$  from the notation, identifying  $\mathbb{G} \simeq \mathbb{R}^m$ . Note that  $e \in \mathbb{G}$  is identified with  $0 \in \mathbb{R}^m$ .

We recall from [Bel96, Prop. 5.18] the following result on the structure of the vector fields  $X_j$ .

**Proposition 44.** In exponential coordinates  $(x_1, \ldots, x_m) \in \mathbb{R}^m$ , the vector fields  $X_j$ , for  $j = 1, \ldots, k$  have the following form:

$$X_j(x_1,\ldots,x_m) = \partial_j + \sum_{h>k} Q_{jh}(x_1,\ldots,x_{k_{w_h-1}})\partial_h,$$

where  $Q_{jh}: \mathbb{G} \simeq \mathbb{R}^m \to \mathbb{R}$  is a homogeneous polynomial of weighted degree  $w_h - 1$ .

5.2. Endpoint maps. In this section we introduce the Endpoint map associated to a Carnot group. In this section we work with the separable Hilbert space

$$H := L^2(I, \mathbb{R}^k),$$

for some fixed interval I, say I = [0, 1], and where k is the rank of  $\mathbb{G}$ . For any  $u \in H$  with  $u = (u_1, \ldots, u_k)$  we consider the following Cauchy problem for a curve  $\gamma : I \to \mathbb{G}$ :

(48) 
$$\dot{\gamma}(t) = \sum_{j=1}^{k} u_j(t) X_j(\gamma(t)), \qquad \gamma(0) = e.$$

It is well-known that (48) admits a unique absolutely continuous solution,  $\gamma_u: I \to \mathbb{G}$ , for any  $u \in H$ .

**Definition 45.** Let  $\mathbb{G}$  be a Carnot group. The Endpoint map of  $\mathbb{G}$  is the function

$$\mathcal{E}: H \to \mathbb{G},$$

given by  $\mathcal{E}(u) := \gamma_u(1)$ . We call the function  $u \in H$  a control. We use the same notation to denote the Endpoint map  $\mathcal{E} : H \to \mathbb{R}^m \simeq \mathbb{G}$  with an identification in exponential coordinates. Thanks to the special form of the vector fields  $X_j$  given by Proposition 44, we can rewrite the system (48) in exponential coordinates. Adopting the notation  $\gamma = (\gamma_1, \ldots, \gamma_m)$ , we have

(49) 
$$\begin{cases} \gamma_i(0) = 0 & i = 1, \dots, m, \\ \dot{\gamma}_i(t) = u_i(t) & i = 1, \dots, k, \\ \dot{\gamma}_i(t) = \sum_{j=1}^k u_j(t) Q_{ji}(\gamma_1(t), \dots, \gamma_{k_{w_i-1}}(t)) & i = k+1, \dots, m, \end{cases}$$

for a.e.  $t \in I$ . Similarly, we write  $\gamma_u = (\gamma_{u,1}, \ldots, \gamma_{u,m})$  for the solution to (49).

The next result (Theorem H in the Introduction) connects this framework to the previous sections.

**Theorem 46.** Let  $\mathbb{G}$  be a Carnot group of topological dimension m, step s, and rank k. Then the Endpoint map  $\mathcal{E} \in \mathscr{P}^m_s(H)$ , for  $H = L^2(I, \mathbb{R}^k)$ .

*Proof.* The fact that  $\mathcal{E}$  is  $\mathcal{C}^1$  (in fact smooth) and that the differential  $D\mathcal{E}$  is locally Lipschitz follows from [ABB20, Prop. 8.5]. The fact that the  $D\mathcal{E}$  is weakly continuous is proved in [BL17, Thm. 23]. It remains to show that for all finite dimensional linear subspaces E of H, the restriction  $\mathcal{E}|_E : E \to \mathbb{R}^m$  is a polynomial map of degree at most s.

We actually prove a stronger claim: for any  $n \in \mathbb{N}$ , for any *n*-dimensional linear subspace  $E \subset L^2(I, \mathbb{R}^k)$ , for any  $t \in I$  and  $i = 1, \ldots, m$ , the function on E defined by  $E \ni u \mapsto \gamma_{u,i}(t)$  is a polynomial of degree  $w_i$ , and its coefficients (when u is written in terms of some basis of  $E \simeq \mathbb{R}^n$ ) are continuous functions of  $t \in I$ . We prove the claim by induction on w. We prove the base case, that is  $w_i = 1$ . In this case  $i = 1, \ldots, k$ . For these values of i, by (49) we obtain

$$\gamma_{u,i}(t) = \int_0^t u_i(s) \, ds,$$

which is clearly a polynomial function of  $u \in E$ , of degree 1. Moreover its coefficients (when u is written in terms of a basis of  $E \simeq \mathbb{R}^n$ ) are continuous functions of  $t \in I$ . This concludes the base case.

Let then  $1 \leq \theta < s$ . The induction assumption is that for any  $t \in I$  the function  $E \ni u \to \gamma_{u,i}(t)$  is a polynomial of degree  $w_i$  for all *i* such that  $w_i \leq \theta$ , with coefficients that depends continuously on *t*.

Now we prove that the above property holds for any *i* such that  $w_i = \theta + 1$ . In particular i > k. In this case, by (49), the curve  $\gamma_{u,i} : I \to \mathbb{R}$  is determined by

(50) 
$$\gamma_{u,i}(t) = \sum_{j=1}^{k} \int_{0}^{t} u_j(s) Q_{ji}(\gamma_{u,1}(s), \dots, \gamma_{u,k_{w_i-1}}(s)) \, ds$$

Recall that  $Q_{ji}$  is a homogeneous polynomial of weighted degree  $w_i - 1$  and as already highlighted in the notation, depends only on the functions  $\gamma_{u,\ell}(s)$  with  $w_{\ell} \leq w_i - 1 = \theta$ . By the induction assumption, all the functions  $E \ni u \mapsto \gamma_{u,\ell}(s)$  are polynomials of degree  $w_{\ell} \leq \theta$ , depending continuously on s. By using the definition of weighted degree (see Remark 4), it follows that the function

$$E \ni u \mapsto Q_{ji}(\gamma_{u,1}(s),\ldots,\gamma_{u,k_{w_i-1}}(s)),$$

is a polynomial of degree  $w_i - 1$ , with coefficients depending continuously on  $s \in I$ . Then, the integrand in (50), namely the function

$$E \ni u \mapsto \sum_{j=1}^{k} u_j(s) Q_{ji}(\gamma_{u,1}(s), \dots, \gamma_{u,k_{w_i-1}}(s)),$$

is a polynomial of degree  $w_i = \theta + 1$ , with coefficients depending continuously on  $s \in I$ . We conclude easily using (50).

Thanks to Theorem H we can apply the results of Section 4 to Endpoint maps of Carnot groups, obtaining the following statement (see Theorem I in the Introduction).

**Theorem 47.** Let  $\mathbb{G}$  be a Carnot group of topological dimension m, step s, and rank k. Let  $K \subset L^2(I, \mathbb{R}^k)$  be a compact subset with

$$\omega(K, L^2(I, \mathbb{R}^k)) < \omega_0(m, s)$$

Then, letting  $V := \operatorname{span}(K)$ , the restriction  $\mathcal{E}|_V : V \to \mathbb{G}$  has the Sard property, namely

$$\mu\Big(\mathcal{E}\big(\operatorname{Crit}(\mathcal{E}|_V)\big)\Big) = 0.$$

In particular  $\mu(\mathcal{E}(\operatorname{Crit}(\mathcal{E}) \cap V)) = 0.$ 

*Proof.* Since  $\mathcal{E} \in \mathscr{P}_m^d(H)$  by Theorem **H**, we can apply Theorem **D**.

5.3. Sard property for real-analytic controls. We specialize the result in Theorem I to some concrete set of controls K, estimating its *n*-width.

Fix a closed interval  $I \subset \mathbb{R}$  and let  $\mathcal{D}(r, I) := \{z \in \mathbb{C} \mid d(z, I) \leq r\}$ . The radius of convergence of a real-analytic function  $u: I \to \mathbb{R}$  is the largest  $r \in (0, \infty]$  such that u extends to a (unique) holomorphic function on  $\mathcal{D}(r, I)$ . For fixed r > 0 we denote by  $\mathcal{C}^{\omega}(I, \mathbb{R}^k; r) \subset L^2(I, \mathbb{R}^k)$  the set of real-analytic controls whose components have radius of convergence strictly greater than r:

 $\mathcal{C}^{\omega}(I,\mathbb{R}^k;r) := \Big\{ u \in L^2(I,\mathbb{R}^k) \, \Big| \, u \text{ is real-analytic with radius of convergence strictly greater than } r \Big\},$ 

which we endow with the supremum norm:

$$\|u\|_{\mathcal{C}^{\omega}(I,\mathbb{R}^k;r)} := \sup_{z\in\mathcal{D}(r,I)} \|u(z)\|.$$

Similarly, we consider finite concatenations of real-analytic controls as above. More precisely, for  $\ell \in \mathbb{N}$  divide the interval I in equispaced sub-intervals:

(51) 
$$I = I_1 \cup \dots \cup I_\ell, \qquad I_a = \inf I + \left[\frac{(a-1)|I|}{\ell}, \frac{a|I|}{\ell}\right],$$

and we let  $\mathcal{C}^{\omega}(I, \mathbb{R}^k; r, \ell) \subset L^2(I, \mathbb{R}^k)$  be the set of piecewise analytic controls  $u: I \to \mathbb{R}^k$  that are real-analytic on each set  $I_1, \ldots, I_\ell$ , with radius of convergence strictly greater than r, namely

$$\mathcal{C}^{\omega}(I,\mathbb{R}^k;r,\ell) = \left\{ u \in L^2(I,\mathbb{R}^k) \, \middle| \, u|_{I_a} \in \mathcal{C}^{\omega}(I_a,\mathbb{R}^k;r) \subset L^2(I_a,\mathbb{R}^k), \text{ for all } a = 1,\ldots,\ell \right\},\$$

endowed with the norm

(52) 
$$\|u\|_{\mathcal{C}^{\omega}(I,\mathbb{R}^k;r,\ell)} = \max_{a=1,\dots,\ell} \|u\|_{\mathcal{C}^{\omega}(I_a,\mathbb{R}^k;r,\ell)}$$

**Theorem 48.** Fix I = [0,1],  $\ell \in \mathbb{N}$  and r > 1. Let  $V = \mathcal{C}^{\omega}(I, \mathbb{R}^k; r, \ell)$ . Then the set  $K_V := \{u \in V \mid ||u||_V \leq 1\}$  is compact in  $L^2(I, \mathbb{R}^k)$ , and for its n-width it holds:

$$\Omega_n(K_V, L^2(I, \mathbb{R}^k)) \le \frac{(k\ell)^{1/2}}{\ln r} \left(\frac{1}{r}\right)^{\lfloor \frac{n}{k\ell} \rfloor}.$$

*Proof.* We start by proving the case  $\ell = 1$ . Fix  $u = (u_1, \ldots, u_k) \in K_V$ . Since r > 1, any  $u_j$  for  $j = 1, \ldots, k$  has an expansion in power series on the complex unit ball centered at the origin, denoted by  $B_{\mathbb{C}}(1)$ . In particular, we have for any  $t \in I$ 

$$u_j(t) = \sum_{h=0}^{\infty} \frac{u_j^{(h)}(0)}{h!} t^h.$$

For any  $N \ge 1$  we consider the truncated sum, given for any  $t \in I$  as

$$u_{j,N}(t) = \sum_{h=0}^{N-1} \frac{u_j^{(h)}(0)}{h!} t^h.$$

By the Cauchy integral formula, and since r > 1, we have

$$u_{j}^{(h)}(0) = \frac{h!}{2\pi i} \int_{\partial B_{\mathbb{C}}(r)} \frac{u_{j}(z)}{z^{h+1}} \, dz,$$

where  $B_{\mathbb{C}}(r)$  is the complex unit ball of radius one centred at zero. Therefore,

$$|u_j^{(h)}(0)| \le \frac{h!}{r^h}.$$

Hence we have

$$\|u_j - u_{j,N}\|_{L^2(I)} \le \|u_j - u_{j,N}\|_{L^\infty(I)} \le \sum_{h=N}^\infty \frac{|u_j^{(h)}(0)|}{h!} \le \sum_{h=N}^\infty \frac{1}{r^h} \le \int_N^\infty \frac{1}{r^x} \, dx = \frac{r^{-N}}{\ln r}$$

(1)

It follows that for all  $u \in K_V$  it holds

(53) 
$$\|u - u_N\|_{L^2(I,\mathbb{R}^k)} \le \left(\sum_{j=1}^k \|u_j - u_{j,N}\|_{L^2(I)}^2\right)^{1/2} \le \frac{k^{1/2}r^{-N}}{\ln r}$$

Thanks to this result we can find an approximating subspace. Consider the standard basis  $e_1, \ldots, e_k$  of  $\mathbb{R}^k$ , and for any  $k \in \mathbb{N}$  we denote by  $f_h \in L^2(I)$  the function  $f_h(t) = t^h$ . We have

$$u_N = \sum_{h=0}^{N-1} \frac{u^{(h)}(0)}{h!} f_h = \sum_{j=1}^k \sum_{h=0}^{N-1} \frac{u_j^{(h)}(0)}{h!} f_h \otimes e_j.$$

In particular,  $u_N \in V_N := \operatorname{span} \{ f_h \otimes e_j \mid j = 1, \dots, k, h = 0, \dots, N-1 \}$  and  $\dim(V_N) = kN$ . Hence from (53) it follows that for any  $N \in \mathbb{N}$ 

(54) 
$$\Omega_{kN}(K_V, L^2(I, \mathbb{R}^k)) \le \frac{k^{1/2}}{\ln r} \left(\frac{1}{r}\right)^N$$

For  $n \in \mathbb{N}$  we consider  $N = \lfloor \frac{n}{k} \rfloor$ . By Definition 27, the *n*-width is non-increasing as a function of  $n \in \mathbb{N}$ , hence by (54) we have

$$\Omega_n(K_V, L^2(I, \mathbb{R}^k)) \le \Omega_{kN}(K_V, L^2(I, \mathbb{R}^k)) \le \frac{k^{1/2}}{\ln r} \left(\frac{1}{r}\right)^{\lfloor \frac{n}{k} \rfloor}$$

This proves the estimate on the *n*-width per  $\ell = 1$ .

We sketch the argument for general  $\ell$ . In this case, for any  $u \in \mathcal{C}^{\omega}(I, \mathbb{R}^k; r, \ell)$  we can write

$$u = \sum_{a=1}^{\ell} u|_{I_a} \mathbb{1}_{I_a},$$

where  $I_a$  are the intervals of the decomposition (51), and  $\mathbb{1}_{I_a}$  are the corresponding characteristic functions. By definition, each  $u|_{I_a}$  is the restriction to  $I_a$  of a real-analytic function with radius of convergence  $r_a \ge r > 1$ . We can expand each  $u|_{I_a}$  in Taylor series centered at the lower bound of  $I_a$ . Then, in order to obtain the finite-dimensional approximation, we truncate the series as in the previous case, repeating analogous estimates for the remainder (taking into account the length of the intervals  $|I_a| = 1/\ell$ ). This concludes the estimate on the *n*-width for general  $\ell$ .

To prove the compactness assertion, observe that there is a linear immersion (given by the inclusion)  $\mathcal{C}^{\omega}(I, \mathbb{R}^{k}; r, \ell) \to L^{2}(I, \mathbb{R}^{k})$  and it holds  $\|u\|_{L^{2}(I, \mathbb{R}^{k})} \leq \|u\|_{C^{\omega}(I, \mathbb{R}^{k}; r, \ell)}$  for all  $u \in C^{\omega}(I, \mathbb{R}^{k}; r, \ell)$ . In particular  $K_{V}$  is bounded in  $L^{2}(I, \mathbb{R}^{k})$  and since its *n*-width tends to zero as  $n \to \infty$  it is also compact by Proposition 28.

We can now prove Theorem J in the Introduction, of which we recall the statement.

**Theorem 49.** Let  $\mathbb{G}$  be a Carnot group of topological dimension m, step s, and rank k. Given  $\ell \in \mathbb{N}$ , there exists  $r = r(m, s, k, \ell) > 0$  such that, letting  $V = \mathcal{C}^{\omega}(I, \mathbb{R}^k; r, \ell)$ , with I = [0, 1], it holds

$$\mathcal{E}(V) = \mathbb{G}$$
 and  $\mu \left( \mathcal{E}(\operatorname{Crit}(\mathcal{E}|_V)) = \mu \left( \mathcal{E}(\operatorname{Crit}(\mathcal{E}) \cap V) \right) = 0.$ 

Namely, the Sard property holds on the space of piecewise real-analytic controls with radius of convergence > r and with  $\ell$  pieces.

Proof. The surjectivity of  $\mathcal{E}$  on  $\mathcal{C}^{\omega}(I, \mathbb{R}^k; r, \ell)$ , for any value of  $r, \ell$ , is discussed in Section 5.4, where we prove in particular that  $\mathcal{E}$  is surjective when restricted on polynomial controls (see Theorem 53). To prove the Sard property, observe that  $\mathcal{E} \in \mathscr{P}_s^m(L^2(I, \mathbb{R}^k))$  by Theorem H. Let  $V = \mathcal{C}^{\omega}(I, \mathbb{R}^k; r, \ell)$ , and  $K_V$  be its unit ball with respect to the norm (52). By Theorem 48, for any r > 1 we have

$$\omega(K_V, L^2(I, \mathbb{R}^k)) := \limsup_{n \to \infty} \Omega_n(K_V, L^2(I, \mathbb{R}^k))^{1/n} \le r^{-\frac{1}{k\ell}},$$

which is smaller than  $\omega_0(m, s)$  if r > 1 is large enough. The result follows now from Theorem D, observing that  $\operatorname{span}(K_V) = V$ .

5.4. Surjectivity of the Endpoint map on finite-dimensional spaces of controls. The Endpoint maps of sub-Riemannian manifolds are surjective when restricted to piecewise constant controls: this follows from the proof of the Rashevskii-Chow theorem in [ABB20]. In this section, we prove that in Carnot groups the Endpoint maps are surjective also when they are restricted to the space of controls which consists in the set of polynomial maps of some large enough fixed degree (which depends on the Carnot group). The proof is obtained by using a quantitative version of the inverse function theorem, as it can be found in [Cla76]. For completeness we present and prove in our setting the statements we need.

Given  $M \in \mathcal{L}(\mathbb{R}^m, \mathbb{R}^m)$  we denote by  $\sigma(M)$  the smallest singular value of M, namely

$$\sigma(M) := \inf_{\|v\|=1} \|Mv\|.$$

From the definition it follows that for all  $M_1, M_2 \in \mathcal{L}(\mathbb{R}^m, \mathbb{R}^m)$  it holds

(55) 
$$|\sigma(M_1) - \sigma(M_2)| \le ||M_1 - M_2||_{\text{op}}.$$

From [Cla76, Lemma 3] we obtain the following technical lemma.

**Lemma 50.** Let  $M_0 \in \mathcal{L}(\mathbb{R}^m, \mathbb{R}^m)$  be invertible. Then, for every  $v \in \mathbb{R}^m$  with ||v|| = 1 there exists  $w \in \mathbb{R}^m$  with ||w|| = 1 such that

$$\langle w, Mv \rangle \ge \frac{\sigma(M_0)}{2},$$

for all M such that  $||M - M_0||_{\text{op}} \leq \frac{\sigma(M_0)}{2}$ .

*Proof.* Given  $v \in \mathbb{R}^m$  with ||v|| = 1, the set  $C_v \subset \mathbb{R}^m$ 

$$C_v := \left\{ Mv \in \mathbb{R}^m \, \middle| \, \|M - M_0\|_{\text{op}} \le \frac{\sigma(M_0)}{2} \right\},\$$

is convex. We now prove that its distance from 0 is at least  $\frac{\sigma(M_0)}{2}$ . Indeed, by (55), for any M with  $\|M - M_0\|_{\text{op}} \leq \frac{\sigma(M_0)}{2}$  we obtain  $\sigma(M) > \frac{\sigma(M_0)}{2}$ . Therefore,

$$||Mv|| \ge \inf_{\|v\|=1} ||Mv|| = \sigma(M) > \frac{\sigma(M_0)}{2}$$

Hence, we have proved that  $C_v$  is separated from the ball  $B_{\mathbb{R}^m}\left(\frac{\sigma(M_0)}{2}\right)$ . We conclude the proof thanks to the usual separation theorem for convex sets. Indeed, it directly provides  $w \in \mathbb{R}^m$  with ||w|| = 1 such that for all M with  $||M - M_0||_{\text{op}} \leq \frac{\sigma(M_0)}{2}$  we have

$$\langle w, Mv \rangle \ge \frac{\sigma(M_0)}{2},$$

concluding the proof.

From [Cla76, Lemma 4] we obtain the following "quantitative injectivity" lemma.

**Lemma 51.** Let  $W \subset \mathbb{R}^m$  be an open set, and  $f \in \mathcal{C}^1(W, \mathbb{R}^m)$ . Assume  $x_0 \in W$  is such that  $D_{x_0}f$  is invertible. Set  $\sigma_{f,x_0} := \sigma(D_{x_0}f) > 0$ . There exists  $r_{f,x_0} > 0$  such that for any  $g \in C^1(W, \mathbb{R}^m)$  with

(56) 
$$\sup_{z \in W} \left\| D_z f - D_z g \right\|_{\text{op}} < \frac{\sigma_{f, x_0}}{4},$$

and all  $x, y \in B_{\mathbb{R}^m}(x_0, r_{f,x_0})$  it holds

$$||g(x) - g(y)|| \ge \frac{\sigma_{f,x_0}}{2} ||x - y||$$

*Proof.* Since f is  $\mathcal{C}^1$  there exists  $r = r_{f,x_0} > 0$  such that  $B_{\mathbb{R}^m}(x_0,r) \subset W$  and

$$\sup_{z \in B_{\mathbb{R}^m}(x_0, r)} \|D_z f - D_{x_0} f\|_{\text{op}} < \frac{\sigma_{f, x_0}}{4}.$$

Let  $g \in \mathcal{C}^1(W, \mathbb{R}^m)$  as in the statement. By (56) we get

(57) 
$$||D_z g - D_{x_0} f||_{\text{op}} < \frac{\sigma_{f,x_0}}{2}$$

for all  $z \in B_{\mathbb{R}^m}(x_0, r)$ . Now we fix  $x \neq y$  in  $B_{\mathbb{R}^m}(x_0, r)$ , and we have

$$g(x) - g(y) = ||x - y|| \int_0^1 D_{(1-t)y+tx}g \cdot \frac{x - y}{||x - y||} dt.$$

We define  $v_{x,y} := \frac{x-y}{\|x-y\|} \in \mathbb{R}^m$ , and we apply Lemma 50 to  $v = v_{x,y}$  and  $M_0 = D_{x_0}f$ , which is invertible by hypothesis. We denote by  $w_{x,y} \in \mathbb{R}^m$  the provided unit vector, such that

$$\left\langle w_{x,y}, D_z g \cdot \frac{x-y}{\|x-y\|} \right\rangle \ge \frac{\sigma_{f,x_0}}{2},$$

for all  $z \in B_{\mathbb{R}^m}(x_0, r)$ . We conclude the proof by the Cauchy-Schwarz inequality, which gives

$$\|g(x) - g(y)\| \ge \|x - y\| \left\langle w_{x,y}, \int_0^1 D_{(1-t)y + tx}g \cdot \frac{x - y}{\|x - y\|} dt \right\rangle \ge \frac{\sigma_{f,x_0}}{2} \|x - y\|,$$
  
he proof

concluding the proof.

We conclude with the following inclusion, analogous to [Cla76, Lemma 5], providing the "quantitative surjectivity" counterpart of the previous statement. We report the proof for completeness.

Lemma 52. In the same setting of Lemma 51, the following inclusion holds

$$g(B_{\mathbb{R}^m}(x_0, r_{f, x_0})) \supset B_{\mathbb{R}^m}\left(g(x_0), \frac{r_{f, x_0}\sigma_{f, x_0}}{8}\right).$$

*Proof.* Set  $r = r_{f,x_0}$  and  $\sigma = \sigma_{f,x_0}$ . Let  $y \in B_{\mathbb{R}^m}\left(g(x_0), \frac{r\sigma}{8}\right)$ . Let x be a minimum of  $||y - g(\cdot)||^2$  on  $B_{\mathbb{R}^n}(x_0, r)$ . We claim that  $x \in \operatorname{int} B_{\mathbb{R}^n}(x_0, r)$ . Otherwise, using Lemma 51, we have

$$\begin{aligned} \frac{r\sigma}{8} &\geq \|y - g(x_0)\| \\ &\geq \|g(x) - g(x_0)\| - \|y - g(x)\| \\ &\geq \frac{\sigma}{2} \|x - x_0\| - \|y - g(x)\| \\ &\geq \frac{\sigma r}{2} - \|y - g(x_0)\| \\ &\geq \frac{\sigma r}{2} - \frac{\sigma r}{8} = \frac{3r\sigma}{8}, \end{aligned}$$

which is a contradiction. Thus x yields a local minimum for the function  $||y - g(\cdot)||^2$ , and consequently  $\nabla_x ||y - g(\cdot)||^2 = -2D_x g \cdot (y - g(x)) = 0$ . Note that by (57) and the fact that the singular values are 1-Lipschitz (see Lemma 18) it follows that  $\sigma(D_x g) > \frac{\sigma}{2}$ , hence  $D_x g$  is invertible. Hence y = g(x).  $\Box$ 

We can now state and prove the surjectivity property of the Endpoint maps of Carnot groups.

**Theorem 53.** For every Carnot group  $\mathbb{G}$  there exists  $d_{\mathbb{G}} \in \mathbb{N}$  such that

 $\mathcal{E}\left(\left\{u \in L^2(I, \mathbb{R}^k) \mid u \text{ is a polynomial map with components of degrees at most } d_{\mathbb{G}}\right\}\right) = \mathbb{G}.$ 

*Proof.* The classical proof of the Rashevskii-Chow theorem provides  $u_0 \in L^2(I, \mathbb{R}^k)$  such that  $D_{u_0}\mathcal{E}$  is surjective and  $\mathcal{E}(u_0) = 0$  (see e.g. [ABB20, Sec. 3.2]). In particular there exist  $w_1, \ldots, w_m \in \ker(D_{u_0}\mathcal{E})^{\perp}$  such that the vectors  $D_{u_0}\mathcal{E}(w_1), \ldots, D_{u_0}\mathcal{E}(w_m)$  are linearly independent. Now we consider the map  $f : \mathbb{R}^m \to \mathbb{G} \simeq \mathbb{R}^m$  defined as

$$f(s_1,\ldots,s_m) = \mathcal{E}\left(u_0 + \sum_{i=1}^m s_i w_i\right).$$

The differential  $D_0 f$  is invertible by construction, in particular f covers a neighbourhood of f(0) = 0. The idea of the proof is to construct a perturbation  $g : \mathbb{R}^m \to \mathbb{G} \simeq \mathbb{R}^m$  of the form

$$g(s_1,\ldots,s_m) = \mathcal{E}\left(q_0 + \sum_{i=1}^m s_i p_i\right),$$

where  $q_0, p_1, \ldots, p_m \in L^2(I, \mathbb{R}^k)$  are suitably chosen polynomial maps, in such a way that the image of g stills contains a neighbourhood of 0. This will be done by applying Lemma 52, as we now explain.

Given  $\epsilon_1, \eta_1 > 0$  by the density of polynomial maps in  $L^2(I, \mathbb{R}^k)$  and by the continuity of  $\mathcal{E}$  and  $D\mathcal{E}$ , there exists a polynomial map  $q_0 \in L^2(I, \mathbb{R}^k)$  such that

$$\|\mathcal{E}(q_0)\| < \epsilon_1$$
 and  $\|D_{u_0}\mathcal{E} - D_{q_0}\mathcal{E}\|_{\text{op}} < \eta_1$ 

Now we prove that we can find polynomial maps  $p_1, \ldots, p_m$  and a neighbourhood  $W \subset \mathbb{R}^m$  of 0 such that, setting  $\sigma_{f,0} = \sigma(D_{x_0}f)$ , it holds

(58) 
$$\sup_{W} \left\| D_s f - D_s g \right\|_{\text{op}} \le \frac{\sigma_{f,0}}{4}.$$

By the density of polynomial maps, given  $\eta_2 > 0$  we consider  $p_1, \ldots, p_m$  such that for any  $i = 1, \ldots, m$ 

$$\|w_i - p_i\| < \eta_2$$

We get the following estimate

$$\begin{split} \|D_{0}f - D_{0}g\|_{\mathrm{op}} &\leq \sum_{i=1}^{m} \|D_{u_{0}}\mathcal{E}(w_{i}) - D_{q_{0}}\mathcal{E}(p_{i})\| \\ &\leq \sum_{i=1}^{m} \|D_{u_{0}}\mathcal{E}(w_{i}) - D_{q_{0}}\mathcal{E}(w_{i})\| + \|D_{q_{0}}\mathcal{E}(w_{i}) - D_{q_{0}}\mathcal{E}(p_{i})\| \\ &\leq \|D_{u_{0}}\mathcal{E} - D_{q_{0}}\mathcal{E}\|_{\mathrm{op}} \sum_{i=1}^{m} \|w_{i}\| + \|D_{q_{0}}\mathcal{E}\|_{\mathrm{op}} \sum_{i=1}^{m} \|w_{i} - p_{i}\| \\ &\leq \eta_{1} \sum_{i=1}^{m} \|w_{i}\| + \|D_{q_{0}}\mathcal{E}\|_{\mathrm{op}} m\eta_{2}. \end{split}$$

Given  $\eta_3 > 0$ , by the continuity of Df and Dg there exists a neighbourhood  $W \subset \mathbb{R}^m$  of 0 such that

$$\sup_{s \in W} \|D_s f - D_0 f\|_{\text{op}} + \|D_s g - D_0 g\|_{\text{op}} < \eta_3.$$

Hence, by the triangle inequality we get the following estimate

$$\sup_{s \in W} \|D_s f - D_s g\|_{\text{op}} \le \eta_3 + \eta_1 \sum_{i=1}^m \|w_i\| + \|D_{q_0} \mathcal{E}\|_{\text{op}} m\eta_2.$$

To obtain (58) it is enough to choose  $\eta_1, \eta_2, \eta_3$  small enough. We have thus found g in such a way  $||f(0) - g(0)|| < \epsilon_1$  and g satisfies the hypothesis of Lemma 52. By the latter, we find  $r_{f,0} > 0$  such that

$$g(B_{\mathbb{R}^m}(r_{f,0})) \supset B_{\mathbb{R}^m}\left(g(0), \frac{r_{f,0}\sigma_{f,0}}{8}\right).$$

Since  $||g(0)|| < \epsilon_1$  by construction, taking  $\epsilon_1 = \frac{r_{f,0}\sigma_{f,0}}{16}$  we also get

$$g(B_{\mathbb{R}^m}(r_{f,0})) \supset B_{\mathbb{R}^m}\left(\frac{r_{f,0}\sigma_{f,0}}{16}\right)$$

Let then  $d_{\mathbb{G}}$  be the maximum degree of the polynomial maps  $q_0, \ldots, p_m$ . We have proved that

(59)  $B_{\mathbb{R}^m}\left(\frac{r_{f,0}\sigma_{f,0}}{16}\right) \subset \mathcal{E}\left(\left\{u \in L^2(I,\mathbb{R}^k) \mid u \text{ is a polynomial map of degree at most } d_{\mathbb{G}}\right\}\right).$ 

In other words, the Endpoint map is surjective on a small ball when restricted to polynomial controls of degree  $\leq d_{\mathbb{G}}$ . Now recall that, for Carnot groups, dilations have the following property:

(60) 
$$\delta_{\lambda}(\mathcal{E}(u)) = \mathcal{E}(\lambda u), \qquad \forall \lambda > 0, \quad u \in L^{2}(I, \mathbb{R}^{k}).$$

From (59) and (60) it follows that  $\mathcal{E}$ , when restricted to polynomial controls of degree  $\leq d_{\mathbb{G}}$  is surjective on the whole  $\mathbb{G} \simeq \mathbb{R}^m$ .

With the same proof we can obtain the following statement, corresponding to Theorem K in the Introduction, of which Theorem 53 is a special case.

**Theorem 54.** Let  $\mathbb{G}$  be a Carnot group with topological dimension m and rank k. Let  $S \subset L^2(I, \mathbb{R}^k)$  be a dense set. Then there exist  $u_0, u_1, \ldots, u_m \in S$  such that

$$\mathcal{E}(\operatorname{span}\{u_0, u_1, \dots, u_m\}) = \mathbb{G}.$$

### References

- [ABB20] A. Agrachev, D. Barilari, and U. Boscain. A comprehensive introduction to sub-Riemannian geometry, volume 181 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2020. From the Hamiltonian viewpoint, With an appendix by Igor Zelenko.
- [AG01] A. Agrachev and J.-P. Gauthier. On the subanalyticity of Carnot-Caratheodory distances. Ann. Inst. H. Poincaré C Anal. Non Linéaire, 18(3):359–382, 2001.
- [AGL15] A. Agrachev, A. Gentile, and A. Lerario. Geodesics and horizontal-path spaces in Carnot groups. Geom. Topol., 19(3):1569–1630, 2015.
- [Agr98] A. Agrachev. Compactness for sub-Riemannian length-minimizers and subanalyticity. volume 56, pages 1–12 (2001). 1998. Control theory and its applications (Grado, 1998).
- [Agr09] A. Agrachev. Any sub-Riemannian metric has points of smoothness. Dokl. Akad. Nauk, 424(3):295–298, 2009.

[Agr14] A. Agrachev. Some open problems. In Geometric control theory and sub-Riemannian geometry, volume 5 of Springer INdAM Ser., pages 1–13. Springer, Cham, 2014.

- [AL09] A. Agrachev and P. Lee. Optimal transportation under nonholonomic constraints. Trans. Amer. Math. Soc., 361(11):6019–6047, 2009.
- [AS96] A. Agrachev and A. V. Sarychev. Abnormal sub-Riemannian geodesics: Morse index and rigidity. Ann. Inst. H. Poincaré C Anal. Non Linéaire, 13(6):635–690, 1996.
- [Bat93] S. M. Bates. On smooth rank-1 mappings of Banach spaces onto the plane. J. Differential Geom., 37(3):729– 733, 1993.
- [BCR98] J. Bochnak, M. Coste, and M.-F. Roy. Géométrie algébrique réelle (Second edition in english: Real Algebraic Geometry), volume 12 (36) of Ergebnisse der Mathematik und ihrer Grenzgebiete [Results in Mathematics and Related Areas]. Springer-Verlag, Berlin, 1987 (1998).
- [BdSFPR22] A. Belotto da Silva, A. Figalli, A. Parusiński, and L. Rifford. Strong Sard conjecture and regularity of singular minimizing geodesics for analytic sub-Riemannian structures in dimension 3. *Invent. Math.*, 229(1):395–448, 2022.
- [BdSPR22] A. Belotto da Silva, A. Parusiński, and L. Rifford. Abnormal subanalytic distributions and minimal rank sard conjecture. arXiv preprint, 2022.
- [BdSPR23] A. Belotto da Silva, A. Parusiński, and L. Rifford. Abnormal singular foliations and the sard conjecture for generic co-rank one distributions. *arXiv preprint*, 2023.
- [BdSR18] A. Belotto da Silva and L. Rifford. The Sard conjecture on Martinet surfaces. Duke Math. J., 167(8):1433– 1471, 2018.
- [Bel96] A. Bellaïche. The tangent space in sub-Riemannian geometry. In Sub-Riemannian geometry, volume 144 of Progr. Math., pages 1–78. Birkhäuser, Basel, 1996.
- [BL17] F. Boarotto and A. Lerario. Homotopy properties of horizontal path spaces and a theorem of Serre in subriemannian geometry. Comm. Anal. Geom., 25(2):269–301, 2017.
- [BNV22] F. Boarotto, L. Nalon, and D. Vittone. The Sard problem in step 2 and in filiform Carnot groups. ESAIM Control Optim. Calc. Var., 28:Paper No. 75, 20, 2022.
- [BPR06] S. Basu, R. Pollack, and M.-F. Roy. Algorithms in real algebraic geometry, volume 10 of Algorithms and Computation in Mathematics. Springer-Verlag, Berlin, second edition, 2006.
- [BR19] D. Barilari and L. Rizzi. Sub-Riemannian interpolation inequalities. *Invent. Math.*, 215(3):977–1038, 2019.
- [BV20] F. Boarotto and D. Vittone. A dynamical approach to the Sard problem in Carnot groups. J. Differential Equations, 269(6):4998–5033, 2020.
- [CJT06] Y. Chitour, F. Jean, and E. Trélat. Genericity results for singular curves. J. Differential Geom., 73(1):45– 73, 2006.

F. H. Clarke. On the inverse function theorem. Pacific J. Math., 64(1):97-102, 1976. [Cla76] H. Federer. The  $(\varphi, k)$  rectifiable subsets of n-space. Trans. Amer. Math. Soc., 62:114–192, 1947. [Fed47] [Kel28] O. D. Kellogg. On bounded polynomials in several variables. Math. Z., 27(1):55–64, 1928. [Kri84] G. K. Kristiansen. Characterization of polynomials by means of their stationary values. Arch. Math. (Basel), 43(1):44-48, 1984. [Kuh69] H. Kuhn. Interpolation vorgeschriebener Extremwerte. J. Reine Angew. Math., 238:24-31, 1969. [Kup65] I. Kupka. Counterexample to the Morse-Sard theorem in the case of infinite-dimensional manifolds. Proc. Amer. Math. Soc., 16:954-957, 1965. [Lan93] S. Lang. Real and functional analysis, volume 142 of Graduate Texts in Mathematics. Springer-Verlag, New York, third edition, 1993. [LDMO<sup>+</sup>16] E. Le Donne, R. Montgomery, A. Ottazzi, P. Pansu, and D. Vittone. Sard property for the endpoint map on some Carnot groups. Ann. Inst. H. Poincaré C Anal. Non Linéaire, 33(6):1639-1666, 2016. [LRT24] A. Lerario, L. Rizzi, and D. Tiberio. Quantitative approximate definable choices. arXiv preprint, 2024. [LZ23] L. Lokutsievskiv and M. Zelikin. Derivatives of sub-Riemannian geodesics are  $L_p$ -Hölder continuous. ESAIM Control Optim. Calc. Var., 29:Paper No. 70, 30, 2023. R. Montgomery. A tour of subriemannian geometries, their geodesics and applications, volume 91 of [Mon02] Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2002. [Mor39] A. P. Morse. The behavior of a function on its critical set. Ann. of Math. (2), 40(1):62–70, 1939. [OV19] A. Ottazzi and D. Vittone. On the codimension of the abnormal set in step two Carnot groups. ESAIM Control Optim. Calc. Var., 25:Paper No. 18, 17, 2019. [Pin85] A. Pinkus. n-widths in approximation theory, volume 7 of Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]. Springer-Verlag, Berlin, 1985. [Rif23] L. Rifford. Subdifferentials and minimizing Sard conjecture in sub-Riemannian geometry. J. Éc. polytech. Math., 10:1195-1244, 2023. [RT05] L. Rifford and E. Trélat. Morse-Sard type results in sub-Riemannian geometry. Math. Ann., 332(1):145–159, 2005.[Sar42]A. Sard. The measure of the critical values of differentiable maps. Bull. Amer. Math. Soc., 48:883-890, 1942[Sma65] S. Smale. An infinite dimensional version of Sard's theorem. Amer. J. Math., 87:861-866, 1965. [Sus14] H. J. Sussmann. A regularity theorem for minimizers of real-analytic subriemannian metrics. In 53rd IEEE Conference on Decision and Control, pages 4801-4806. 2014. [Tao12] T. Tao. Topics in random matrix theory, volume 132 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2012. [Vit55] A. G. Vituškin. O mnogomernyh variaciyah. Gosudarstv. Izdat. Tehn.-Teor. Lit., Moscow, 1955. [Vit61] A. G. Vituškin. Theory of the transmission and processing of information. Pergamon Press, New York-Oxford-London-Paris, 1961. Translated from the Russian by Ruth Feinstein; translation editor A. D. Booth. [YC04] Y. Yomdin and G. Comte. Tame geometry with application in smooth analysis, volume 1834 of Lecture Notes in Mathematics. Springer-Verlag, Berlin, 2004. [Yom83] Y. Yomdin. The geometry of critical and near-critical values of differentiable mappings. Math. Ann., 264(4):495-515, 1983.[Yom88] Y. Yomdin. Approximational complexity of functions. In Geometric aspects of functional analysis (1986/87), volume 1317 of Lecture Notes in Math., pages 21-43. Springer, Berlin, 1988. [ZZ95] I. Zelenko and M. Zhitomirskii. Rigid paths of generic 2-distributions on 3-manifolds. Duke Math. J.,

79(2):281-307, 1995.