

Berndt-type Integrals: Unveiling Connections with Barnes Zeta and Jacobi Elliptic Functions

Zachary P. Bradshaw¹ and Christophe Vignat^{2,3}

¹Naval Surface Warfare Center, Panama City Division, USA

²LSS, CentraleSupélec, Université Paris Saclay, France

³Tulane University, Department of Mathematics, New Orleans, USA

Abstract

We address a class of definite integrals known as Berndt-type integrals, highlighting their role as specialized instances within the integral representation framework of the Barnes-zeta function. Building upon the foundational insights of Xu and Zhao, who adeptly evaluate these integrals using rational linear combinations of Lambert-type series and derive closed-form expressions involving products of $\Gamma^4(1/4)$ and π^{-1} , we uncover direct evaluations of the Barnes-zeta function. Moreover, our inquiry leads us to establish connections between Berndt-type integrals and Jacobi elliptic functions, as well as moment polynomials investigated by Lomont and Brillhart, a relationship elucidated through the seminal contributions of Kuznetsov. In this manner, we extend and integrate these diverse mathematical threads, unveiling deeper insights into the intrinsic connections and broader implications of Berndt-type integrals in special function and integration theory.

1 Introduction

Since its inception, the theory of integration has captivated practitioners of mathematics. Unlike the derivative operation, which relies solely on local information and adheres to a finite set of rules applicable to any differentiable function, definite integration hinges on non-local data spanning an entire interval. As a result, integration theory encompasses a diverse array of techniques, each tailored to tackle specific challenges, and yet no single combination suffices to evaluate every conceivable integral. This inherent complexity renders the theory fertile ground for exploration, replete with unsolved problems which the practitioner may spend a lifetime resolving.

Here we concern ourselves with the class

$$I_{\pm}(s, p) = \int_0^{\infty} \frac{x^{s-1}}{(\cosh(x) \pm \cos(x))^p} dx \quad (1)$$

of *Berndt-type integrals* of order p [6], a class with notable historical roots dating back to the pioneering work of the esteemed mathematician S. Ramanujan in 1916 [22]. These integrals have also featured prominently in the context of moment problems studied by Ismail and Valent [14], and the ongoing effort to evaluate them has been advanced by Xu and Zhao [28, 29]. Central to the methodology for evaluating these integrals is the application of Cauchy's residue theory, a formidable tool in one's integration toolkit. In this way, Berndt-type integrals of varying orders reveal their equivalence to series involving hyperbolic trigonometric functions, and by leveraging results attributed to Ramanujan, explicit closed-form expressions for these sums emerge.

In this work, we give an alternative evaluation of the Berndt-type integrals in terms of the Barnes zeta function [3, 2, 23, 12] defined by

$$\zeta_N(s, w | a_1, \dots, a_N) = \sum_{n_1 \geq 0, \dots, n_N \geq 0} \frac{1}{(w + n_1 a_1 + \dots + n_N a_N)^s}$$

and of its alternating version

$$\tilde{\zeta}_N(s, w | a_1, \dots, a_N) = \sum_{n_1 \geq 0, \dots, n_N \geq 0} \frac{(-1)^{n_1 + \dots + n_N}}{(w + n_1 a_1 + \dots + n_N a_N)^s}.$$

While the Lambert series representation given by Xu and Zhao allows for a closed form evaluation in many cases, the Barnes zeta representation presented here gives the evaluation of the integral for *any* choice of s, p in the domain

of convergence but does not produce a closed form in terms of more elementary functions. Combining this with the results of Xu and Zhao produces closed form evaluations of the Barnes zeta function. Our results are easily generalized to the case of Dirichlet type analogs of the Barnes zeta function. We also relate Berndt-type integrals to Jacobi elliptic functions through the generating function methods of Kuznetsov [16] and further connect them to a class of moment polynomials [20]. The work of Lomont and Brillhart then produces recurrence relations for Berndt-type integrals.

In Section 2, we evaluate Berndt-type integrals using the Barnes zeta function and observe that this procedure generalizes to Dirichlet type analogs of this function. In the appendix, we take an alternative approach involving Euler-Barnes and Bernoulli-Barnes polynomials [4, 10, 15] and identify the Bernoulli-Barnes polynomials as an analytic continuation of Berndt-type integrals to negative powers. In Section 3 we review the work of Kuznetsov connecting Berndt-type integrals to Jacobi elliptic functions before using it to draw connections to a class of moment polynomials defined by a recurrence relation. Kuznetsov's direct evaluation of $I_+(s, 1)$ is then extended to the case of $I_-(s, 1)$ in Section 4. Finally, we give concluding remarks in Section 5

2 Connections with the Barnes Zeta Function

We begin with a classic integral representation of the Barnes zeta function, which appears as [23, (3.2)]. Its proof, here reproduced for completeness, follows as a consequence of Euler's integral.

Proposition 1. *Let $\Re(s) > N$, $\Re(w) > 0$, and $\Re(a_j) > 0$ for $j = 1, \dots, N$. Then*

$$\zeta_N(s, w|a_1, \dots, a_N) = \frac{1}{\Gamma(s)} \int_0^\infty u^{s-1} e^{-wu} \prod_{j=1}^N (1 - e^{-a_j u})^{-1} du.$$

The alternating version is

$$\tilde{\zeta}_N(s, w|a_1, \dots, a_N) = \frac{1}{\Gamma(s)} \int_0^\infty u^{s-1} e^{-wu} \prod_{j=1}^N (1 + e^{-a_j u})^{-1} du.$$

Proof. Since $\Re(s) > N$, we can write the gamma function in its integral representation, and the series representation of ζ_N converges absolutely. Note that the conditions $\Re(w) > 0$ and $\Re(a_j) > 0$ are required here to assure that the denominator doesn't vanish. We have

$$\begin{aligned} \zeta_N(s, w|a_1, \dots, a_N) \Gamma(s) &= \sum_{n_1, \dots, n_N} \frac{1}{(w + n_1 a_1 + \dots + n_N a_N)^s} \int_0^\infty x^{s-1} e^{-x} dx \\ &= \sum_{n_1, \dots, n_N} \int_0^\infty \frac{x^{s-1} e^{-x}}{(w + n_1 a_1 + \dots + n_N a_N)^s} dx \\ &= \sum_{n_1, \dots, n_N} \int_0^\infty u^{s-1} e^{-(w + n_1 a_1 + \dots + n_N a_N)u} du. \end{aligned}$$

The sum can be moved inside by the dominated convergence theorem, giving us

$$\zeta_N(s, w|a_1, \dots, a_N) \Gamma(s) = \int_0^\infty u^{s-1} e^{-wu} \sum_{n_1, \dots, n_N} e^{-(n_1 a_1 + \dots + n_N a_N)u} du,$$

and noting that $\Re(a_j) > 0$, we may apply the geometric series formula so that

$$\begin{aligned} \zeta_N(s, w|a_1, \dots, a_N) \Gamma(s) &= \int_0^\infty u^{s-1} e^{-wu} \frac{1}{1 - e^{-a_1 u}} \dots \frac{1}{1 - e^{-a_N u}} du \\ &= \int_0^\infty u^{s-1} e^{-wu} \prod_{j=1}^N (1 - e^{-a_j u})^{-1} du. \end{aligned}$$

The alternating version is proved similarly. □

A consequence of this general result is the following Berndt-type integral identity.

Corollary 1. Let $\Re(s) > 2$, $\Re(a) > 0$, and $-\Re(a) < \Im(b) < \Re(a)$. Then we have

$$\frac{1}{2\Gamma(s)} \int_0^\infty \frac{x^{s-1}}{\cosh(ax) - \cos(bx)} dx = \zeta_2(s, a|a - bi, a + bi),$$

and in particular,

$$\frac{1}{2\Gamma(s)} \int_0^\infty \frac{x^{s-1}}{\cosh(x) - \cos(x)} dx = \zeta_2(s, 1|1 - i, 1 + i).$$

The alternating case is given by

$$\frac{1}{2\Gamma(s)} \int_0^\infty \frac{x^{s-1}}{\cosh(ax) + \cos(bx)} dx = \tilde{\zeta}_2(s, a|a - bi, a + bi).$$

More generally, we may consider the Dirichlet-type multiple series

$$L_\chi(s, w|a_1, \dots, a_N) = \sum_{n_1, \dots, n_N \geq 0} \frac{\chi(n_1, \dots, n_N)}{(w + n_1 a_1 + \dots + n_N a_N)^s},$$

and hope for a similar integral representation. Formally, we have

$$\begin{aligned} L_\chi(s, w|a_1, \dots, a_N)\Gamma(s) &= \sum_{n_1, \dots, n_N \geq 0} \int_0^\infty \frac{x^{s-1} e^{-x} \chi(n_1, \dots, n_N)}{(w + n_1 a_1 + \dots + n_N a_N)^s} dx \\ &= \sum_{n_1, \dots, n_N \geq 0} \int_0^\infty \chi(n_1, \dots, n_N) x^{s-1} e^{-(w+n_1 a_1 + \dots + n_N a_N)x} dx \\ &= \int_0^\infty x^{s-1} \sum_{n_1, \dots, n_N \geq 0} \chi(n_1, \dots, n_N) e^{-(w+n_1 a_1 + \dots + n_N a_N)x} dx. \end{aligned}$$

The choice of $\chi = 1$ recovers Proposition 1. Meanwhile, other choices of χ result in similar identities. If χ is separable in the sense that $\chi(n_1, \dots, n_N) = \chi_1(n_1) \cdots \chi_N(n_N)$ for some χ_1, \dots, χ_N , then we observe that L_χ obtains the particularly simple form

$$L_\chi(s, w|a_1, \dots, a_N) = \frac{1}{\Gamma(s)} \int_0^\infty x^{s-1} e^{-wx} \prod_{k=1}^N \hat{\chi}_k(a_k x) dx,$$

with the Fourier transforms

$$\hat{\chi}_k(x) = \sum_{n \geq 0} \chi_k(n) e^{-nx},$$

and we list some examples of this type in Table 1.

Let us turn to the Berndt-type integrals (1) of arbitrary order and produce an evaluation using the Barnes zeta function and its alternating counterpart. In what follows, the notation $(a, b)^p$ indicates that the symbols a, b are repeated p times i.e. $(a, b)^3 = a, b, a, b, a, b$.

Proposition 2. Let $p > 0$ and $s > 2p$. Then

$$\int_0^\infty \frac{x^{s-1}}{(\cosh(x) - \cos(x))^p} dx = 2^p \Gamma(s) \zeta_{2p}(s, p|(1+i, 1-i)^p)$$

and

$$\int_0^\infty \frac{x^{s-1}}{(\cosh(x) + \cos(x))^p} dx = 2^p \Gamma(s) \tilde{\zeta}_{2p}(s, p|(1+i, 1-i)^p)$$

Proof. We will tackle the case of arbitrary products of sinh and cosh in the denominator and then specialize to the

| No. | $\chi(n_1, \dots, n_N)$ | $L_\chi(s, w a_1, \dots, a_N)$ |
|-----|--|--|
| 1 | 1 | $\frac{1}{\Gamma(s)} \int_0^\infty x^{s-1} e^{-wx} \prod_{j=1}^N (1 - e^{-a_j x})^{-1} dx$ |
| 2 | $(-1)^{n_1 + \dots + n_N}$ | $\frac{1}{\Gamma(s)} \int_0^\infty x^{s-1} e^{-wx} \prod_{j=1}^N (1 + e^{-a_j x})^{-1} dx$ |
| 3 | $n_1 \cdots n_N$ | $\frac{1}{\Gamma(s)} \int_0^\infty x^{s-1} e^{-wx} \prod_{j=1}^N (e^{a_j x/2} - e^{-a_j x/2})^{-2} dx$ |
| 4 | $\frac{c_1^{n_1} \cdots c_N^{n_N}}{n_1! \cdots n_N!}$ | $\frac{1}{\Gamma(s)} \int_0^\infty x^{s-1} e^{-wx} \prod_{j=1}^N e^{c_j e^{-a_j x}} dx$ |
| 5 | $\sin(n_1) \cdots \sin(n_N)$ | $\frac{1}{\Gamma(s)} \int_0^\infty x^{s-1} e^{-wx} \prod_{j=1}^N \left(\frac{-i(e^{2i} - 1)}{2} \right) (e^{a_j x + i} - e^{-(a_j x + i)})^{-1} dx$ |
| 6 | $\frac{1}{\Gamma(1 + \frac{n_1}{2}) \cdots \Gamma(1 + \frac{n_N}{2})}$ | $\frac{1}{\Gamma(s)} \int_0^\infty x^{s-1} e^{-wx} \prod_{j=1}^N e^{e^{-2a_j x}} (1 + \text{Erf}(e^{-a_j x})) dx$ |
| 7 | $H_{n_1} \cdots H_{n_N}$ | $\frac{1}{\Gamma(s)} \int_0^\infty x^{s-1} e^{-wx} (-1)^N \prod_{j=1}^N \log(1 - e^{-a_j x}) (1 - e^{-a_j x})^{-1} dx$ |
| 8 | $\frac{(-1)^{n_1 + \dots + n_N}}{(n_1 + 1) \cdots (n_N + 1)}$ | $\frac{1}{\Gamma(s)} \int_0^\infty x^{s-1} e^{-wx} \prod_{j=1}^N e^{a_j x} \log(1 + e^{-a_j x}) dx$ |
| 9 | $(-1)^{n_1 + \dots + n_N} n_1 \cdots n_N$ | $\frac{1}{\Gamma(s)} \int_0^\infty x^{s-1} e^{-wx} (-1)^N \prod_{j=1}^N (e^{a_j x/2} + e^{-a_j x/2})^{-2} dx$ |
| 10 | $\sqrt{n_1 \cdots n_N}$ | $\frac{1}{\Gamma(s)} \int_0^\infty x^{s-1} e^{-wx} \prod_{j=1}^N \text{polylog}(-\frac{1}{2}, e^{-a_j x}) dx$ |
| 11 | $(-1)^{n_1 + \dots + n_N} \sqrt{n_1 \cdots n_N}$ | $\frac{1}{\Gamma(s)} \int_0^\infty x^{s-1} e^{-wx} \prod_{j=1}^N \text{polylog}(-\frac{1}{2}, \sinh(a_j x) - \cosh(a_j x)) dx$ |

Table 1: List of Identities given by a choice of χ , where H_n denotes the n -th harmonic number. Here we assume that $\Re(s) > N$, $\Re(a_j) > 0$, and that $\Re(w) > 0$.

case of (1). Observe that

$$\begin{aligned}
I &:= \frac{1}{\Gamma(s)} \int_0^\infty \frac{x^{s-1} e^{-wx}}{\sinh(a_1 x) \cdots \sinh(a_M x) \cosh(b_1 x) \cdots \cosh(b_N x)} dx \\
&= \frac{1}{\Gamma(s)} \int_0^\infty \frac{x^{s-1} e^{-wx}}{\left(\prod_{i=1}^M \frac{e^{a_i x} - e^{-a_i x}}{2} \right) \left(\prod_{j=1}^N \frac{e^{b_j x} + e^{-b_j x}}{2} \right)} dx \\
&= \frac{2^{M+N}}{\Gamma(s)} \int_0^\infty \frac{x^{s-1} e^{-wx}}{\left(\prod_{i=1}^M (e^{a_i x} - e^{-a_i x}) \right) \left(\prod_{j=1}^N (e^{b_j x} + e^{-b_j x}) \right)} dx \\
&= \frac{2^{M+N}}{\Gamma(s)} \int_0^\infty \frac{x^{s-1} e^{-x(w + a_1 + \dots + a_M + b_1 + \dots + b_N)}}{\left(\prod_{i=1}^M (1 - e^{-2a_i x}) \right) \left(\prod_{j=1}^N (1 + e^{-2b_j x}) \right)} dx \\
&= \frac{2^{M+N}}{\Gamma(s)} \int_0^\infty x^{s-1} e^{-x(w + \sum_{i=1}^M a_i + \sum_{i=1}^N b_i)} \sum_{n_1, \dots, n_M, k_1, \dots, k_N \geq 0} (-1)^{k_1 + \dots + k_N} e^{-2x(\sum_{i=1}^M a_i n_i + \sum_{i=1}^N b_i k_i)} dx \\
&= \frac{2^{M+N}}{\Gamma(s)} \sum_{n_1, \dots, n_M, k_1, \dots, k_N \geq 0} \int_0^\infty (-1)^{k_1 + \dots + k_N} x^{s-1} e^{-x(w + \sum_{i=1}^M a_i + \sum_{i=1}^N b_i + 2(\sum_{i=1}^M a_i n_i + \sum_{i=1}^N b_i k_i))} dx \\
&= \frac{2^{M+N}}{\Gamma(s)} \sum_{n_1, \dots, n_M, k_1, \dots, k_N \geq 0} \int_0^\infty \frac{(-1)^{k_1 + \dots + k_N} x^{s-1} e^{-x} dx}{(w + \sum_{i=1}^M a_i + \sum_{i=1}^N b_i + 2(\sum_{i=1}^M a_i n_i + \sum_{i=1}^N b_i k_i))^s} \\
&= 2^{M+N} \sum_{n_1, \dots, n_M, k_1, \dots, k_N \geq 0} \frac{(-1)^{k_1 + \dots + k_N} dx}{(w + \sum_{i=1}^M a_i + \sum_{i=1}^N b_i + 2(\sum_{i=1}^M a_i n_i + \sum_{i=1}^N b_i k_i))^s}.
\end{aligned}$$

Now writing $\cosh(x) - \cos(x) = 2 \sinh(\frac{1-i}{2}x) \sinh(\frac{1+i}{2}x)$ and $\cosh(x) + \cos(x) = 2 \cosh(\frac{1-i}{2}x) \cosh(\frac{1+i}{2}x)$, the proposition follows. \square

Remark 1. In view of the previous proposition, it is worth pointing out that $\zeta_{2p}(s, p|(a, b)^p)$ can be simplified to a double sum for any value of $p \geq 2$, as a Dirichlet analog of ζ_2 . Indeed, noticing that

$$\sum_{n_1, \dots, n_p, m_1, \dots, m_p \geq 0} \frac{1}{(w + a(n_1 + \dots + n_p) + b(m_1 + \dots + m_p))^s} = \sum_{n, m \geq 0} \frac{P(n)P(m)}{(w + an + bm)^s},$$

where $P(n) = |\{n_1 \geq 0, \dots, n_p \geq 0 : n_1 + \dots + n_p = n\}| = \binom{n+p-1}{n}$ is a counting function, it follows that

$$\zeta_{2p}(s, w|(a, b)^p) = L_\chi(s, w|a, b) = \sum_{n, m \geq 0} \frac{\chi(n, m)}{(w + an + bm)^s}$$

with $\chi(n, m) = P(n)P(m) = \binom{n+p-1}{n} \binom{m+p-1}{m}$. This result extends to the more general case, assuming that $a_i, b_j \in \mathbb{Z}$

$$\sum_{n_1, \dots, n_p, m_1, \dots, m_p \geq 0} \frac{1}{(w + a_1 n_1 + \dots + a_p n_p + b_1 m_1 + \dots + b_p m_p)^s} = \sum_{n, m \geq 0} \frac{\chi_{a,b}(n, m)}{(w + an + bm)^s}$$

with $\chi_{a,b}(n, m) = \chi_a(n)\chi_b(m)$ and with the partition function $\chi_a(n) = |\{n_1 \geq 0, \dots, n_p \geq 0 : n_1 a_1 + \dots + n_p a_p = n\}|$.

In the work of Xu and Zhao, explicit evaluations of the quadratic Berndt-type integrals are given as rational linear combinations of products of $\Gamma(1/4)$ and π . For example, they show that

$$\int_0^\infty \frac{x^5}{(\cosh(x) - \cos(x))^2} dx = \frac{\Gamma^{16}(1/4)}{3 \cdot 2^{14} \pi^6} - \frac{\Gamma^8(1/4)}{2^8 \pi^2},$$

and so it follows from Proposition 2 that

$$\zeta(6, 2|(1+i, 1-i)^2) = \frac{1}{480} \left(\frac{\Gamma^{16}(1/4)}{3 \cdot 2^{14} \pi^6} - \frac{\Gamma^8(1/4)}{2^8 \pi^2} \right).$$

In this way, the results of Xu and Zhao, together with the above results, reveal explicit evaluations of the Barnes zeta function in terms of the gamma function. More generally, they show that for all integers $p \geq 1$ and $s \geq \lceil m/2 \rceil$, the Berndt-type integrals satisfy

$$I_+(4s+2, p) \in \mathbb{Q} [\Gamma^4(1/4), \pi^{-1}],$$

where $\mathbb{Q}[x, y]$ denotes all rational linear combinations of products of x and y (the polynomial ring generated by x, y with rational coefficients). Moreover, they show that the degrees of $\Gamma^4(1/4)$ have the same parity as p and are between $2s - p + 2$ and $2s + p$, inclusive, while the degrees of π^{-1} are between $2s - p + 2$ and $2s + 3p - 2$, inclusive. Similarly, they show that

$$I_-(4s, 2p+1) \in \mathbb{Q} [\Gamma^4(1/4), \pi^{-1}]$$

for all $s \geq k+1 \geq 1$ and that the degrees of $\Gamma^4(1/4)$ in each term are always even and between $2s - 2p$ and $2s + 2p$, while the degrees of π^{-1} are between $2s - 2p$ and $2s + 6p$. In the even order case, they show that

$$I_-(4s+1, 2p) \in \mathbb{Q} [\Gamma^4(1/4), \pi^{-1}]$$

for all integers $s \geq k \geq 1$ and the powers of $\Gamma^4(1/4)$ are always even and between $2s + 2 - 2p$ and $2s + 2p$, while the degrees of π^{-1} are between $2s + 2 - 2p$ and $2s + 6p - 2$. The following corollary therefore follows from Proposition 2.

Corollary 2. *Let $p \geq 1$ and $s \geq \lceil p/2 \rceil$ be integers. Then*

$$\tilde{\zeta}_{2p}(4s+2, p|(1+i, 1-i)^p) \in \mathbb{Q} [\Gamma^4(1/4), \pi^{-1}].$$

Similarly, if $s \geq p+1 \geq 1$, then

$$\zeta_{4p+2}(4s, 2p+1|(1+i, 1-i)^{2p+1}) \in \mathbb{Q} [\Gamma^4(1/4), \pi^{-1}],$$

and if $s \geq p \geq 1$, then

$$\zeta_{4p}(4s+1, 2p|(1+i, 1-i)^{2p}) \in \mathbb{Q} [\Gamma^4(1/4), \pi^{-1}].$$

We will close this section by pointing out two consequences of the connection between Berndt-type integrals and the Barnes zeta function.

Laplace transforms

The Barnes zeta representation lends itself to an interesting interpretation in terms of the Laplace transform. Indeed, at least formally, we have the following proposition showing that the integration of an analytic function f against the kernel $(\cosh ax - \cos bx)^{-1}$ performs a lattice summation of the Laplace transform of f .

Proposition 3. *Let $f(x) = \sum_{n \geq 2} \frac{a_n}{n!} x^n$ be an analytic function such that $f(0) = f'(0) = 0$, and let $F(p)$ denote its Laplace transform. Then*

$$\int_0^\infty \frac{f(x)}{\cosh ax - \cos bx} dx = 2 \sum_{p, q \geq 0} F(a + p(a - ib) + q(a + ib)). \quad (2)$$

Proof. The Laplace transform of f is given by

$$F(p) = \sum_{n \geq 2} \frac{a_n}{p^{n+1}}$$

so that formally

$$\begin{aligned} \int_0^\infty \frac{f(x)}{\cosh ax - \cos bx} dx &= \sum_{n \geq 2} \frac{a_n}{n!} \int_0^\infty \frac{x^n}{\cosh ax - \cos bx} dx \\ &= 2 \sum_{n \geq 2} \frac{a_n}{n!} \Gamma(n+1) \zeta_2(n+1, a, |a-ib, a+ib) \\ &= 2 \sum_{n \geq 2} a_n \sum_{p, q \geq 0} (a+p(a-ib) + q(a+ib))^{-n-1} \\ &= 2 \sum_{p, q \geq 0} F(a+p(a-ib) + q(a+ib)). \end{aligned}$$

□

This result extends to the case of the $(\cosh ax + \cos bx)^{-1}$ denominator as follows.

Proposition 4. Let $F(p)$ denote the Laplace transform of the analytic function $f(x) = \sum_{n \geq 0} \frac{a_n}{n!} x^n$. Then

$$\int_0^\infty \frac{f(x)}{\cosh ax + \cos bx} dx = 2 \sum_{p, q \geq 0} (-1)^{p+q} F(a+p(a-ib) + q(a+ib)). \quad (3)$$

Proof. The Laplace transform of f is given by

$$F(s) = \sum_{n \geq 0} \frac{a_n}{s^{n+1}},$$

from which we deduce

$$\begin{aligned} \int_0^\infty \frac{f(x)}{\cosh ax + \cos bx} dx &= \sum_{n \geq 0} \frac{a_n}{n!} \int_0^\infty \frac{x^n}{\cosh ax + \cos bx} dx \\ &= \sum_{n \geq 0} \frac{a_n}{n!} 2\Gamma(n+1) \tilde{\zeta}(n+1, a|a-ib, a+ib) \\ &= 2 \sum_{n \geq 0} a_n \sum_{p, q \geq 0} \frac{(-1)^{p+q}}{(a+p(a-ib) + q(a+ib))^{n+1}} \\ &= 2 \sum_{p, q \geq 0} (-1)^{p+q} F(a+p(a-ib) + q(a+ib)). \end{aligned}$$

□

Corollary 3. The special case $f(x) = \frac{\sin x}{x}$ produces the identity

$$\sum' (-1)^{p+q+1} \arctan\left(\frac{p+q+1}{p(p+1) + q(q+1)}\right) = \frac{\pi}{4}$$

where the \sum' sign indicates summation over the set of integers $\{p \geq 0, q \geq 0, (p, q) \neq (0, 0)\}$.

Proof. The Laplace transform of f is given by $F(s) = \arctan\left(\frac{1}{s}\right)$, from which it follows that

$$F(1+p+q+i(p-q)) = \arctan\left(\frac{1}{1+p+q+i(p-q)}\right).$$

We need only the real part of the Laplace transform. Denote

$$z = \frac{1}{1+p+q+i(p-q)} = x + iy.$$

Using [11, 4.23.36],

$$\Re(\arctan(x + iy)) = \frac{1}{2} \arctan \frac{2x}{1 - x^2 - y^2}.$$

Then with $x = \frac{1+p+q}{(1+p+q)^2+(p-q)^2}$ and $y = -\frac{p-q}{(1+p+q)^2+(p-q)^2}$, it follows that

$$\begin{aligned} \Re(F(1 + p(1+i) + q(1-i))) &= \Re\left(\arctan\left(\frac{1}{1+p+q+i(p-q)}\right)\right) \\ &= \frac{1}{2} \arctan\left(\frac{p+q+1}{p(p+1)+q(q+1)}\right) \end{aligned}$$

whereas, for $(p, q) = (0, 0)$,

$$F(1) = \arctan(1) = \frac{\pi}{4}.$$

We deduce

$$\begin{aligned} \frac{\pi}{4} &= \int_0^\infty \frac{\sin x}{x(\cosh x + \cos x)} dx \\ &= 2F(1) + 2 \sum' F(1 + p(1+i) + q(1-i)) \\ &= \frac{\pi}{2} + 2 \sum' (-1)^{p+q} \frac{1}{2} \arctan\left(\frac{p+q+1}{p(p+1)+q(q+1)}\right), \end{aligned}$$

and it follows that

$$\sum' (-1)^{p+q+1} \arctan \frac{p+q+1}{p(p+1)+q(q+1)} = \frac{\pi}{4}.$$

□

Analytic continuation

The second application of the Barnes zeta representation is an analytic continuation of a normalized version of a Berndt-type integral: since at $x = 0$,

$$\frac{x^{s-1}}{(\cosh x - \cos x)^p} \sim x^{s-1-2p}$$

the normalized integral

$$\hat{I}(s) = \frac{1}{\Gamma(s)} I(s) = \frac{1}{\Gamma(s)} \int_0^\infty \frac{x^{s-1}}{(\cosh x - \cos x)^p} dx$$

is convergent in the half-plane $\Re s > 2p$. Based on a result by Ruijsenaars', it can be analytically continued as follows.

Proposition 5. *The representation*

$$\hat{I}(s) = \frac{1}{\Gamma(s)} \int_0^\infty \frac{x^{s-1}}{(\cosh x - \cos x)^p} dx = 2^p \zeta_{2p}(s, p|(1+i, 1-i)^p)$$

produces an analytic continuation of $\hat{I}(s)$ to the whole complex plane except at the points $s = 0, 1, \dots, 2p$. Moreover, for positive $s > 2p$,

$$\hat{I}(-s) = \frac{(-2)^p}{(s+1)\dots(s+2p)} B_{2p+s}(p|(1+i, 1-i)^p)$$

with the Bernoulli-Barnes polynomials $B_n(z, \mathbf{a})$ defined by the generating function

$$\sum_{n \geq 0} \frac{B_n(z, (a_1, \dots, a_p))}{n!} u^n = e^{zu} \prod_{i=1}^p \frac{a_i u}{e^{a_i u} - 1}.$$

Proof. The multivariate integral representation in Ruijsenaars' article [23, (4.13)]

$$\zeta_{2p}(s, w|\mathbf{a}) = \frac{1}{(s-1)\dots(s-2p)} \int_{\mathbb{R}^n} \left(w + \sum_{i=1}^{2p} (ix_j - \frac{a_j}{2}) \right)^{2p-s} \prod_{i=1}^{2p} \phi\left(\frac{x_i}{a_i}\right) dx_i$$

with $\phi(x) = \frac{\pi}{2} \operatorname{sech}^2(\pi x)$ produces an analytic continuation (in the variable s) of the Barnes zeta function to the whole complex plane except at the points $s = 0, 1, \dots, 2p$. An integral representation for the Bernoulli Barnes polynomials, given by Ruijsenaars as

$$B_n(z, (a_1, \dots, a_p)) = \int_{\mathbb{R}^n} \left(w + \sum_{j=1}^{2p} (ix_j - \frac{a_j}{2}) \right)^n \prod_{i=1}^{2p} \phi\left(\frac{x_i}{a_i}\right) dx_i,$$

produces the desired result. \square

3 Kuznetsov's Approach

In [16], Kuznetsov gives a direct evaluation of a curious integral appearing in the work of Ismail and Valent [14] which takes the form

$$\int_{\mathbb{R}} \frac{dx}{\cos(\sqrt{x}K) + \cosh(\sqrt{x}K')} = 2,$$

where $K = K(k)$ denotes the complete elliptic integral of the first kind with elliptic modulus $k \in (0, 1)$, and $K' = K(k')$, where $k' = \sqrt{1 - k^2}$ is the complementary elliptic modulus.

In fact, Kuznetsov shows more generally that for $u \in \mathbb{C}$ satisfying $|\Re(u)| < K$ and $|\Im(u)| < K'$, we have the generating function

$$\int_{\mathbb{R}} \frac{\sin(\sqrt{x}u)}{\sqrt{x}} \frac{dx}{\cos(\sqrt{x}K) + \cosh(\sqrt{x}K')} = 2 \frac{\operatorname{sn}(u, k)}{\operatorname{cd}(u, k)} = 2\widetilde{\operatorname{nc}}(u, k), \quad (4)$$

where sn and cd are Jacobi elliptic functions and $\widetilde{\operatorname{nc}}(u, k) = \frac{d}{du} \log \operatorname{nc}(u, k)$. This identity has several implications which we will now discuss.

3.1 Connection to Berndt Integrals

Let us define the moment integrals

$$I_n^+ = \frac{1}{2} \int_{\mathbb{R}} \frac{x^n dx}{\cos(\sqrt{x}K) + \cosh(\sqrt{x}K')}, \quad (5)$$

which we note are the special cases $I_n^+ := I_+(n+1, 1)$ of the Berndt-type integrals defined in (1). By Taylor expanding both sides of identity (4) in u , it follows that

$$\frac{1}{2} \int_{\mathbb{R}} \frac{x^n dx}{\cos(\sqrt{x}K) + \cosh(\sqrt{x}K')} = (-1)^n \frac{d^{2n+1}}{du^{2n+1}} \widetilde{\operatorname{nc}}(u, k) \Big|_{u=0}. \quad (6)$$

Clearly the moment integrals I_n^+ are closely related to the Berndt-type integrals; indeed, a substitution $z = \sqrt{x}$ in (5) reveals that

$$\int_L \frac{z^{2n+1} dz}{\cos(zK) + \cosh(zK')} = (-1)^n \frac{d^{2n+1}}{du^{2n+1}} \widetilde{\operatorname{nc}}(u, k) \Big|_{u=0}, \quad (7)$$

where L is the contour which travels from infinity down the positive imaginary axis until reaching zero, where the contour then begins traveling along the orthogonal positive real axis to infinity. In the lemniscatic case, the modulus $\tilde{k} = \frac{1}{\sqrt{2}}$ is equal to the complementary modulus and the elliptic integral becomes $\tilde{K} = \frac{\pi^{\frac{3}{2}}}{2\Gamma^2(\frac{3}{4})}$. Thus, by making an additional substitution, we have

$$\int_L \frac{z^{2n+1} dz}{\cos(z) + \cosh(z)} = (-1)^n \tilde{K}^{2n+2} \frac{d^{2n+1}}{du^{2n+1}} \widetilde{\operatorname{nc}}(u, \tilde{k}) \Big|_{u=0},$$

and now substituting $x = -iz$ along the vertical portion of the contour finally produces

$$(1 + (-1)^n) \int_0^\infty \frac{x^{2n+1} dx}{\cos(x) + \cosh(x)} = \tilde{K}^{2n+2} \frac{d^{2n+1}}{du^{2n+1}} \widetilde{\operatorname{nc}}(u, \tilde{k}) \Big|_{u=0},$$

so that if $n = 2m$ is even, we obtain a Berndt-type integral representation in terms of Jacobi elliptic functions

$$\int_0^\infty \frac{x^{4m+1} dx}{\cos(x) + \cosh(x)} = \frac{1}{2} \left(\frac{\pi^{\frac{3}{2}}}{2\Gamma^2(\frac{3}{4})} \right)^{4m+2} \frac{d^{4m+1}}{du^{4m+1}} \widetilde{\operatorname{nc}}(u, 1/\sqrt{2}) \Big|_{u=0}.$$

The more general identity (7) is used in the next subsections to produce equivalent forms of Berndt's integrals in terms of special functions.

3.2 Moment Polynomials of Lomont and Brillhart

Evaluation of the first values of Kuznetsov's integral

$$I_0^+ = 1, I_1^+ = -2(1 - 2k^2), I_2^+ = 16(k^4 - k^2 + 1)$$

suggests that they can be expressed as a polynomial function of the elliptic modulus k . The work of Lomont and Brillhart in [20] allows us to identify these polynomials and some of their properties: for example, this connection can be leveraged to derive recurrence relations between Berndt-type integrals.

From [20, (5.35)], we have

$$\log \text{nc}(u, k) = \sum_{n \geq 0} 2^n P_n(1 - 2k^2, 4k^4 - 4k^2 + 4) \frac{u^{2n+2}}{(2n+2)!} \quad (8)$$

with $P_n(x, y)$ the moment polynomials defined in [20, Ch.4] by the recurrence

$$P_{n+2} = xP_{n+1} + (y - x^2) \sum_{j=0}^n \binom{2n+2}{2j} P_j \sum_{l=0}^{n-j} \binom{2n-2j+1}{2l} P_l P_{n-j-l}.$$

with initial values $P_0(x, y) = 1$, $P_1(x, y) = x$. These polynomials have integer coefficients and satisfy

$$\deg_x P_n(x, y) = \begin{cases} n & n \not\equiv 2 \pmod{3} \\ 3n & n \equiv 2 \pmod{3} \end{cases}$$

and

$$\deg_y P_n(x, y) = \left\lfloor \frac{n}{2} \right\rfloor.$$

First cases are [20, Table 4.1]

$$P_0(x, y) = 1, P_1(x, y) = x, P_2(x, y) = y, P_3(x, y) = -10x^3 + 11xy.$$

It follows from (8) that

$$\widetilde{\text{nc}}(u, k) = \sum_{n \geq 0} 2^n P_n(1 - 2k^2, 4k^4 - 4k^2 + 4) \frac{u^{2n+1}}{(2n+1)!},$$

producing the following evaluation of the integral considered by Kuznetsov.

Proposition 6. *Kuznetsov's integral satisfies*

$$I_n^+ = \frac{1}{2} \int_{\mathbb{R}} \frac{x^n dx}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} = (-1)^n 2^n P_n(1 - 2k^2, 4k^4 - 4k^2 + 4), \quad (9)$$

Identity (9) induces the following properties:

Corollary 4. *If k^2 is a rational number, then Kuznetsov's integral I_n is a rational number. Moreover, since*

$$I_1^+ = -2(1 - 2k^2), I_2^+ = 16(1 - k^2 + k^4),$$

the value of I_n^+ is a polynomial function of the initial values I_1 and I_2 given by

$$I_n^+ = (-2)^n P_n\left(-\frac{1}{2}I_1, \frac{1}{4}I_2\right).$$

It is instructive to list several of the first examples for the lemniscatic case $k = \frac{1}{\sqrt{2}}$ ¹:

$$I_0^+ = \frac{1}{2} \int_{\mathbb{R}} \frac{dx}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} = 1$$

$$I_1^+ = \frac{1}{2} \int_{\mathbb{R}} \frac{xdx}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} = 0$$

¹the sequence (1, 12, 3024, ...) appears as OEIS A104203 as it coincides with the Taylor coefficients of the sine lemniscate function

$$I_2^+ = \frac{1}{2} \int_{\mathbb{R}} \frac{x^2 dx}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} = 12$$

$$I_3^+ = \frac{1}{2} \int_{\mathbb{R}} \frac{x^3 dx}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} = 0$$

$$I_4^+ = \frac{1}{2} \int_{\mathbb{R}} \frac{x^4 dx}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} = 3024.$$

In this case, $P_{2n+1}(x, y) = P_{2n+1}(0, 3) = 0$ for all $n \geq 1$ as a consequence of the fact that x always factors in $P_{2n+1}(x, y)$, as can be seen from the recurrence relations satisfied by the polynomials P_n . Indeed, we have [20, (4.50)]

$$P_{n+2} = -n(2n+3)xP_{n+1} + \sum_{j=0}^n \left[2 \binom{2n+3}{2j+3} - \binom{2n+4}{2j+4} \right] P_{j+2} P_{n-j}$$

and

$$P_{n+2} = xP_{n+1} + (y-x^2) \sum_{j=0}^n \sum_{l=0}^{n-j} \binom{2n+2}{2j} \binom{2n-2j+1}{2l} P_j P_l P_{n-j-l},$$

from which we deduce the following proposition.

Proposition 7. *The integrals I_n^+ satisfy the recurrence relations*

$$I_{n+2}^+ = 2n(2n+3)(1-2k^2)I_{n+1}^+ + \sum_{j=0}^n \left[2 \binom{2n+3}{2j+3} - \binom{2n+4}{2j+4} \right] I_{j+2}^+ I_{n-j}^+$$

and

$$I_{n+2}^+ = -2(1-2k^2)I_{n+1}^+ + 12 \sum_{j=0}^n \binom{2n+2}{2j} I_j \sum_{l=0}^{n-j} \binom{2n-2j+1}{2l} I_l^+ I_{n-j-l}^+.$$

3.3 Symmetry Results

Kuznetsov's result (4) also allows us to recover some well-known symmetries of elliptic functions, as well as use these well-known symmetries to produce symmetries of Kuznetsov's integral. Jacobi's imaginary transformations

$$\operatorname{sn}(u, k) = -i \operatorname{sc}(iu, k')$$

and

$$\operatorname{cd}(u, k) = \operatorname{nd}(iu, k'),$$

imply that

$$\frac{\operatorname{sn}(u, k)}{\operatorname{cd}(u, k)} = -i \frac{\operatorname{sc}(iu, k')}{\operatorname{nd}(iu, k')} = -i \frac{\operatorname{sn}(iu, k')}{\operatorname{cn}(iu, k') \operatorname{nd}(iu, k')} = -i \frac{\operatorname{sn}(iu, k')}{\operatorname{cd}(iu, k')}$$

and we deduce

Proposition 8. *As a consequence of Jacobi's imaginary transformations, the ratio $\frac{\operatorname{sn}(u, k)}{\operatorname{cd}(u, k)}$ is invariant by the transformation $(k, u) \rightarrow (k', -iu)$.*

We give a proof of this result that relies solely on basic transformations of Kuznetsov's integral.

Proof. The change of variable $x \rightarrow -x$ in the integral yields

$$\frac{\operatorname{sn}(u, k)}{\operatorname{cd}(u, k)} = \frac{1}{2} \int_{\mathbb{R}} \frac{\sinh(u\sqrt{x})}{\sqrt{x}} \frac{dx}{\cos(K'\sqrt{x}) + \cosh(K\sqrt{x})}.$$

Replacing $k \rightarrow k'$ and $u \rightarrow \frac{u}{i}$ in the integral produces the result. \square

More basic invariances of the elliptic functions imply non trivial identities between Kuznetsov integrals as follows.

Proposition 9. *Kuznetsov's integrals satisfy the identities*

$$\left(\int_{\mathbb{R}} \frac{\sin(u\sqrt{x})}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} \frac{dx}{\sqrt{x}} \right) \left(\int_{\mathbb{R}} \frac{\sin((K-u)\sqrt{x})}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} \frac{dx}{\sqrt{x}} \right) = 4,$$

$$\left(\int_{\mathbb{R}} \frac{\sin(u\sqrt{x})}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} \frac{dx}{\sqrt{x}} \right) \left(\int_{\mathbb{R}} \frac{\sin((K+u)\sqrt{x})}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} \frac{dx}{\sqrt{x}} \right) = -4$$

and

$$\int_{\mathbb{R}} \frac{\sin((K-u)\sqrt{x})}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} \frac{dx}{\sqrt{x}} + \int_{\mathbb{R}} \frac{\sin((K+u)\sqrt{x})}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} \frac{dx}{\sqrt{x}} = 0.$$

Proof. The identity $\operatorname{sn}(u+K, k) = \operatorname{cd}(u, k)$ [27, pg. 500] and the fact that cd is an even function of u produces $\operatorname{sn}(K-u, k) = \operatorname{cd}(u, k)$, so that we have

$$\operatorname{dn}(u, k) = \frac{\operatorname{cn}(u, k)}{\operatorname{sn}(K-u, k)}.$$

As a consequence,

$$\frac{\operatorname{sn}(u, k)}{\operatorname{cd}(u, k)} = \frac{\operatorname{sn}(u, k)}{\operatorname{sn}(K-u, k)},$$

so that the change of variable $u \rightarrow K-u$ produces

$$\frac{\operatorname{sn}(K-u, k)}{\operatorname{cd}(K-u, k)} = \frac{\operatorname{sn}(K-u, k)}{\operatorname{sn}(u, k)} = \left(\frac{\operatorname{sn}(u, k)}{\operatorname{cd}(u, k)} \right)^{-1}$$

and, for $u > 0$, it follows that

$$\left(\int_{\mathbb{R}} \frac{\sin(u\sqrt{x})}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} \frac{dx}{\sqrt{x}} \right) \left(\int_{\mathbb{R}} \frac{\sin((K-u)\sqrt{x})}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} \frac{dx}{\sqrt{x}} \right) = 4.$$

In the same way, from $\operatorname{sn}(u+K, k) = \operatorname{cd}(u, k)$, we deduce that $\operatorname{dn}(u, k) = \frac{\operatorname{cn}(u, k)}{\operatorname{sn}(u+K, k)}$, from which it follows that $\frac{\operatorname{sn}(u, k)}{\operatorname{cd}(u, k)} = \frac{\operatorname{sn}(u, k)}{\operatorname{sn}(u+K, k)}$. Therefore, since $\operatorname{sn}(u+2K, k) = -\operatorname{sn}(u, k)$, we have

$$\frac{\operatorname{sn}(u+K, k)}{\operatorname{cd}(u+K, k)} = \frac{\operatorname{sn}(u+K, k)}{\operatorname{sn}(u+2K, k)} = -\frac{\operatorname{sn}(u+K, k)}{\operatorname{sn}(u, k)} = -\frac{\operatorname{cd}(u, k)}{\operatorname{sn}(u, k)},$$

and as a consequence, for all u , we deduce

$$\left(\int_{\mathbb{R}} \frac{\sin(u\sqrt{x})}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} \frac{dx}{\sqrt{x}} \right) \left(\int_{\mathbb{R}} \frac{\sin((K+u)\sqrt{x})}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} \frac{dx}{\sqrt{x}} \right) = -4$$

and

$$\int_{\mathbb{R}} \frac{\sin((K-u)\sqrt{x})}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} \frac{dx}{\sqrt{x}} + \int_{\mathbb{R}} \frac{\sin((K+u)\sqrt{x})}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} \frac{dx}{\sqrt{x}} = 0.$$

□

3.4 Lambert Series Representation

Kuznetsov's result also produces a Lambert series representation for Berndt-type integrals.

Proposition 10. *A Lambert series representation for Berndt-type integrals is*

$$\frac{1}{2} \int_{\mathbb{R}} \frac{x^{p-1} dx}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} = \left(\frac{\pi}{K} \right)^{2p} \left[-\frac{1}{2} E_{2p-1}(0) + 2 \sum_{n \geq 1} \frac{n^{2p} q^n}{n(1+(-q)^n)} \right].$$

Proof. Indeed, by [25, vol.3 p.15],

$$\log \operatorname{cn}(u, k) = \log \cos \left(\frac{\pi u}{2K} \right) - 4 \sum_{n \geq 1} q^n \frac{\sin^2 \left(\frac{n\pi u}{2K} \right)}{n(1+(-q)^n)}.$$

We expand, using [13, 1.518.2],

$$\log \cos \left(\frac{\pi u}{2K} \right) = - \sum_{k \geq 1} \frac{2^{2k-1} (2^{2k} - 1) |B_{2k}|}{k (2k)!} \left(\frac{\pi u}{2K} \right)^{2k}$$

with, for $k \geq 1$,

$$|B_{2k}| = (-1)^{k-1} B_{2k}$$

and

$$(2^{2k} - 1) B_{2k} = -k E_{2k-1}(0).$$

We deduce

$$\log \cos \left(\frac{\pi u}{2K} \right) = - \frac{1}{2} \sum_{k \geq 1} \frac{(-1)^k E_{2k-1}(0)}{(2k)!} \left(\frac{\pi u}{K} \right)^{2k}.$$

Observe that

$$\begin{aligned} \sum_{n \geq 1} q^n \frac{\sin^2 \left(\frac{n\pi u}{2K} \right)}{n (1 + (-q)^n)} &= \frac{1}{2} \sum_{n \geq 1} q^n \frac{1 - \cos \left(\frac{n\pi u}{K} \right)}{n (1 + (-q)^n)} \\ &= \frac{1}{2} \sum_{n \geq 1} q^n \frac{1}{n (1 + (-q)^n)} \left(1 - \sum_{p \geq 0} \frac{(-1)^p}{(2p)!} \left(\frac{n\pi u}{K} \right)^{2p} \right) \\ &= - \frac{1}{2} \sum_{n \geq 1} q^n \frac{1}{n (1 + (-q)^n)} \left(\sum_{p \geq 1} \frac{(-1)^p}{(2p)!} \left(\frac{n\pi u}{K} \right)^{2p} \right) \\ &= - \frac{1}{2} \sum_{p \geq 1} \frac{(-1)^p}{(2p)!} \left(\frac{\pi u}{K} \right)^{2p} \sum_{n \geq 1} \frac{n^{2p} q^n}{n (1 + (-q)^n)}, \end{aligned}$$

so that we deduce the Taylor series expansion

$$\begin{aligned} \log \operatorname{cn}(u, k) &= - \frac{1}{2} \sum_{p \geq 1} \frac{(-1)^p E_{2p-1}(0)}{(2p)!} \left(\frac{\pi u}{K} \right)^{2p} + 2 \sum_{p \geq 1} \frac{(-1)^p}{(2p)!} \left(\frac{\pi u}{K} \right)^{2p} \sum_{n \geq 1} \frac{n^{2p} q^n}{n (1 + (-q)^n)} \\ &= \sum_{p \geq 1} \frac{(-1)^p}{(2p)!} \left(\frac{\pi u}{K} \right)^{2p} \left[- \frac{1}{2} E_{2p-1}(0) + 2 \sum_{n \geq 1} \frac{n^{2p} q^n}{n (1 + (-q)^n)} \right]. \end{aligned}$$

Finally, noticing that

$$\begin{aligned} \frac{1}{2} \int_{\mathbb{R}} \frac{\sin(u\sqrt{x})}{\sqrt{x}} \frac{dx}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} &= - \frac{d}{du} \log \operatorname{cn}(u, k) \\ &= \sum_{p \geq 1} \frac{(-1)^{p-1}}{(2p-1)!} \left(\frac{\pi}{K} \right)^{2p} u^{2p-1} \left[- \frac{1}{2} E_{2p-1}(0) + 2 \sum_{n \geq 1} \frac{n^{2p} q^n}{n (1 + (-q)^n)} \right], \end{aligned}$$

and expanding the sine term in Kuznetsov's integral

$$\frac{1}{2} \int_{\mathbb{R}} \frac{\sin(u\sqrt{x})}{\sqrt{x}} \frac{dx}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} = \frac{1}{2} \sum_{p \geq 1} \frac{(-1)^{p-1} u^{2p-1}}{(2p-1)!} \int_{\mathbb{R}} \frac{x^{p-1} dx}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})},$$

we deduce the Lambert series representation

$$\frac{1}{2} \int_{\mathbb{R}} \frac{x^{p-1} dx}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} = \left(\frac{\pi}{K} \right)^{2p} \left[- \frac{1}{2} E_{2p-1}(0) + 2 \sum_{n \geq 1} \frac{n^{2p} q^n}{n (1 + (-q)^n)} \right].$$

□

3.5 Eisenstein Series Representation

Finally, we are able to produce an Eisenstein series representation for Berndt-type integrals.

Proposition 11. *The integral*

$$I_p^+ = \frac{1}{2} \int_{\mathbb{R}} \frac{x^p}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} dx$$

has Eisenstein series expansion

$$I_p^+ = (-1)^p (2p+1)! \sum'_{m,n} \left(\frac{1}{(2mK + i(2n+1)K')^{2p+2}} - \frac{1}{(iK')^{2p+2}} \right) - \left(\frac{1}{((2m+1)K + i2nK')^{2p+2}} - \frac{1}{K^{2p+2}} \right)$$

Proof. From [26, vol.5 p.11], with $c_{0,0} = 0$, $c_{m,n} = 1$ (m, n) \neq ($0, 0$)

$$\begin{aligned} & \frac{1}{2} \int_{\mathbb{R}} \frac{\sin(u\sqrt{x})}{\sqrt{x}} \frac{dx}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} = \widetilde{\text{nc}}(u, k) \\ & = -\eta_1 K + i\eta_2 K' - \sum_{\mathbb{Z}^2} \left[\frac{1}{2mK + i(2n+1)K' - u} - \frac{1}{(2m+1)K + i2nK' - u} + \frac{c_{m,n}(-K + iK')}{(2mK + i2nK')^2} \right] \\ & = -\eta_1 K + i\eta_2 K' - \left[\frac{1}{iK' - u} - \frac{1}{K - u} \right] \\ & \quad - \sum' \left[\frac{1}{2mK + i(2n+1)K' - u} - \frac{1}{(2m+1)K + i2nK' - u} + \frac{-K + iK'}{(2mK + i2nK')^2} \right] \end{aligned}$$

Expanding each term produces

$$\frac{1}{2mK + i(2n+1)K' - u} - \frac{1}{(2m+1)K + i2nK' - u} = \sum_{p \geq 0} \frac{u^p}{(2mK + i(2n+1)K')^{p+1}} - \frac{u^p}{((2m+1)K + i2nK')^{p+1}}$$

and

$$\frac{1}{iK' - u} - \frac{1}{K - u} = \sum_{p \geq 0} \frac{u^p}{(iK')^{p+1}} - \frac{u^p}{K^{p+1}},$$

from which we obtain

$$\widetilde{\text{nc}}(u, k) = \widetilde{\text{nc}}(0, k) + \sum_{p \geq 1} u^p \sum'_{m,n} \left(\frac{1}{(2mK + i(2n+1)K')^{p+1}} - \frac{1}{(iK')^{p+1}} \right) - \left(\frac{1}{((2m+1)K + i2nK')^{p+1}} - \frac{1}{K^{p+1}} \right)$$

with the constant term

$$\widetilde{\text{nc}}(0, k) = -\eta_1 K + i\eta_2 K' + \sum' \frac{-K + iK'}{(2mK + i2nK')^2} = 0$$

since $\widetilde{\text{nc}}(u, k)$ is an odd function. Identifying with

$$\widetilde{\text{nc}}(u, k) = \sum_{p \geq 0} \frac{(-1)^p}{(2p+1)!} I_p^+ u^{2p+1}$$

produces the Eisenstein series expansion for the integral

$$I_p^+ = (-1)^p (2p+1)! \sum'_{m,n} \left(\frac{1}{(2mK + i(2n+1)K')^{2p+2}} - \frac{1}{(iK')^{2p+2}} \right) - \left(\frac{1}{((2m+1)K + i2nK')^{2p+2}} - \frac{1}{K^{2p+2}} \right).$$

□

4 Extension of Kuznetsov's Result: The Difference Case

We will now endeavour to extend Kuznetsov's result to the case of the integrals

$$I_n^- = \int_{\mathbb{R}} \frac{x^{n+1}}{\cos(K\sqrt{x}) - \cosh(K'\sqrt{x})} dx$$

where the addition in the denominator is replaced by a difference.

4.1 Generating Function

We begin with a proof of the following theorem which gives a generating function for I_n^- .

Theorem 1. *Let $k \in (0, 1)$. Then for $u \in \mathbb{C}$ satisfying $|\Re(u)| < K/2$ and $|\Im(u)| < K'/2$, we have*

$$\int_{\mathbb{R}} \frac{\sqrt{x} \sin(u\sqrt{x})}{\cos(K\sqrt{x}) - \cosh(K'\sqrt{x})} dx = -8 \frac{\operatorname{sn}^2(u, k)}{\operatorname{cd}^2(u, k) \operatorname{sd}(2u, k)} = -2 \frac{d}{du} \widetilde{\operatorname{nc}}^2(u, k) \quad (10)$$

with the log-derivative $\widetilde{\operatorname{nc}}(u, k) = \frac{d}{du} \log \operatorname{nc}(u, k)$.

Proof. As in Kuznetsov's approach, we will establish (10) for $u = v(K + iK')/2$ with $v \in (-1, 1)$ and then extend by analytic continuation. Fix $v \in (-1, 1)$ and let

$$I := \int_{\mathbb{R}} \frac{f(\sqrt{x}v(K + iK')/2)}{\sqrt{x}(\cos(\sqrt{x}K) - \cosh(\sqrt{x}K'))} dx.$$

We will begin with the change of variables $z = K\sqrt{x}/2$, which maps the contour \mathbb{R} into the contour L considered earlier. The result of the change of variables is therefore

$$\begin{aligned} I &= \frac{4}{K} \int_L \frac{f(vz(1 + \tau))}{\cos(2z) - \cos(2z\tau)} dz \\ &= -\frac{2}{K} \int_L \frac{f(vz(1 + \tau))}{\sin(z(1 + \tau)) \sin(z(1 - \tau))} dz \end{aligned}$$

where we have defined $\tau := iK'/K$. Let $f(vz(1 + \tau)) = z^2 g(vz(1 + \tau))$ so that the singularity at $z = 0$ is removable. We care only about the simple poles appearing in the first quadrant which have the form $z_n = \pi n/(1 - \tau)$. Let us assume that g is analytic in the first quadrant. By closing the contour with a quarter circle and showing that this arc term decays exponentially, it follows from Cauchy's residue theorem that

$$\begin{aligned} I &= -\frac{4\pi i}{K} \sum_{n=1}^{\infty} \operatorname{Res}_{\pi n/(1-\tau)} \left(\frac{z^2 g(vz(1 + \tau))}{\sin(z(1 + \tau)) \sin(z(1 - \tau))} \right) \\ &= -\frac{4\pi^3 i}{K(1 - \tau)^3} \sum_{n=1}^{\infty} (-1)^n n^2 \frac{g(\pi n v t)}{\sin(\pi n t)}, \end{aligned}$$

where we have defined $t := \frac{1+\tau}{1-\tau}$. From [21, (2.16)], we have

$$\operatorname{sd}^2(vt\tilde{K}, \tilde{k}) = \frac{\tilde{E} - (\tilde{k}')^2 \tilde{K}}{\tilde{k}^2 (\tilde{k}')^2 \tilde{K}} + \frac{2\pi^2}{\tilde{k}^2 (\tilde{k}')^2 \tilde{K}^2} \sum_{n \geq 1} \frac{(-1)^n n q^n}{1 - q^{2n}} \cos(\pi n v t),$$

where $q = e^{i\pi t}$ is the nome associated to the lattice parameter t , $\tilde{K} := K(\tilde{k})$, and \tilde{k} is defined by $t = iK(\tilde{k}')/K(\tilde{k})$. In other words, we let t be the lattice parameter in [21, (2.16)] and distinguish the associated modulus and elliptic integrals from those related to the lattice parameter τ by including a tilde. Differentiating with respect to v produces

$$8\tilde{k}^2 (\tilde{k}')^2 \tilde{K}^3 \operatorname{sd}(vt\tilde{K}, \tilde{k}) \operatorname{cd}(vt\tilde{K}, \tilde{k}) \operatorname{nd}(vt\tilde{K}, \tilde{k}) = -4\pi^3 i \sum_{n \geq 1} (-1)^n n^2 \frac{\sin(\pi n v t)}{\sin(\pi n t)}.$$

By choosing $g(z) = \sin(z)$, it follows that

$$I = \frac{8\tilde{k}^2 (\tilde{k}')^2 \tilde{K}^3}{K(1 - \tau)^3} \operatorname{sd}(vt\tilde{K}, \tilde{k}) \operatorname{cd}(vt\tilde{K}, \tilde{k}) \operatorname{nd}(vt\tilde{K}, \tilde{k}).$$

With this choice of g , the integral becomes

$$I = \frac{K^2}{4} \int_{\mathbb{R}} \frac{\sqrt{x} \sin(u\sqrt{x})}{\cos(K\sqrt{x}) - \cosh(K'\sqrt{x})},$$

so that

$$J := \int_{\mathbb{R}} \frac{\sqrt{x} \sin(u\sqrt{x})}{\cos(K\sqrt{x}) - \cosh(K'\sqrt{x})} dx = \frac{32\tilde{k}^2 (\tilde{k}')^2 \tilde{K}^3}{K^3(1 - \tau)^3} \operatorname{sd}(vt\tilde{K}, \tilde{k}) \operatorname{cd}(vt\tilde{K}, \tilde{k}) \operatorname{nd}(vt\tilde{K}, \tilde{k}).$$

The identities [11, 22.2.6-22.2.8] relate the Jacobi elliptic functions seen here to theta functions, and the identities [11, 22.2.2] relate the modulus, the complementary modulus, and the elliptic integral to theta functions. Making use of these identities, we produce

$$J = \frac{4\pi^3}{K^3(1-\tau)^3} \theta_2^2(0, q) \theta_3^2(0, q) \theta_4^2(0, q) \frac{\theta_1(\pi vt/2, q) \theta_2(\pi vt/2, q) \theta_4(\pi vt/2, q)}{\theta_3^3(\pi vt/2, q)},$$

where the reader is reminded that the nome q is with lattice parameter t . We will now transform the theta functions which are currently in terms of t into theta functions which are in terms of τ . To distinguish the two cases we write $\theta_i(w|t) := \theta_i(w, q)$ when the nome is in terms of lattice parameter t ; that is, $q = e^{i\pi t}$. Now applying the theta function transformation identities [11, 20.7.26-20.7.29], we have

$$\begin{aligned} J &= \frac{4\pi^3}{K^3(1-\tau)^3} \theta_2^2(0|t) \theta_3^2(0|t) \theta_4^2(0|t) \frac{\theta_1(\pi vt/2|t) \theta_2(\pi vt/2|t) \theta_4(\pi vt/2|t)}{\theta_3^3(\pi vt/2|t)} \\ &= -\frac{4\pi^3}{K^3(1-\tau)^3} \theta_2^2(0|t+1) \theta_4^2(0|t+1) \theta_3^2(0|t+1) \frac{\theta_1(\pi vt/2|t+1) \theta_2(\pi vt/2|t+1) \theta_3(\pi vt/2|t+1)}{\theta_4^3(\pi vt/2|t+1)} \\ &= -\frac{4\pi^3}{K^3(1-\tau)^3} \theta_2^2\left(0 \middle| \frac{2}{1-\tau}\right) \theta_4^2\left(0 \middle| \frac{2}{1-\tau}\right) \theta_3^2\left(0 \middle| \frac{2}{1-\tau}\right) \frac{\theta_1\left(\pi vt/2 \middle| \frac{2}{1-\tau}\right) \theta_2\left(\pi vt/2 \middle| \frac{2}{1-\tau}\right) \theta_3\left(\pi vt/2 \middle| \frac{2}{1-\tau}\right)}{\theta_4^3\left(\pi vt/2 \middle| \frac{2}{1-\tau}\right)}. \end{aligned}$$

Next we apply the product reduction formulae [11, 20.7.11-20.7.12] to obtain

$$J = -\frac{\pi^3}{2K^3(1-\tau)^3} \theta_2^5\left(0 \middle| \frac{1}{1-\tau}\right) \theta_3^2\left(0 \middle| \frac{1}{1-\tau}\right) \theta_4^2\left(0 \middle| \frac{1}{1-\tau}\right) \frac{\theta_1\left(\pi vt/4 \middle| \frac{1}{1-\tau}\right) \theta_2\left(\pi vt/4 \middle| \frac{1}{1-\tau}\right) \theta_2\left(\pi vt/2 \middle| \frac{1}{1-\tau}\right)}{\theta_3^3\left(\pi vt/4 \middle| \frac{1}{1-\tau}\right) \theta_4^3\left(\pi vt/4 \middle| \frac{1}{1-\tau}\right)}.$$

Applying the lattice parameter transformations [11, 20.7.30-20.7.33] produces

$$J = \frac{\pi^3 \theta_4^5(0|\tau-1) \theta_3^2(0|\tau-1) \theta_2^2(0|\tau-1) \theta_1(-\pi v(\tau+1)/4|\tau-1) \theta_4(-\pi v(\tau+1)/4|\tau-1) \theta_4(-\pi v(\tau+1)/2|\tau-1)}{2K^3 \theta_3^3(-\pi v(\tau+1)/4|\tau-1) \theta_2^3(-\pi v(\tau+1)/4|\tau-1)}.$$

Using the lattice parameter transformations [11, 20.7.26-20.7.29] once more, we have

$$J = \frac{\pi^3 \theta_3^5(0|\tau) \theta_4^2(0|\tau) \theta_2^2(0|\tau) \theta_1(-\pi v(\tau+1)/4|\tau) \theta_3(-\pi v(\tau+1)/4|\tau) \theta_3(-\pi v(\tau+1)/2|\tau)}{2K^3 \theta_4^3(-\pi v(\tau+1)/4|\tau) \theta_2^3(-\pi v(\tau+1)/4|\tau)}.$$

We can write the elliptic integral K in terms of θ_3 by applying [11, 22.2.2], so that

$$J = \frac{4\theta_4^2(0|\tau) \theta_2^2(0|\tau) \theta_1(-\pi v(\tau+1)/4|\tau) \theta_3(-\pi v(\tau+1)/4|\tau) \theta_3(-\pi v(\tau+1)/2|\tau)}{\theta_3(0|\tau) \theta_4^3(-\pi v(\tau+1)/4|\tau) \theta_2^3(-\pi v(\tau+1)/4|\tau)}.$$

Now applying the duplication formula [11, 22.7.10] produces

$$J = \frac{8\theta_2(0|\tau) \theta_4(0|\tau) \theta_1^2(-\pi v(\tau+1)/4|\tau) \theta_3^2(-\pi v(\tau+1)/4|\tau) \theta_3(-\pi v(\tau+1)/2|\tau)}{\theta_3^2(0|\tau) \theta_4^2(-\pi v(\tau+1)/4|\tau) \theta_2^2(-\pi v(\tau+1)/4|\tau) \theta_1(-\pi v(\tau+1)/2|\tau)},$$

so that the definitions of the Jacobi elliptic functions [11, 22.2.4, 22.2.7, 22.2.8] finally give

$$\begin{aligned} J &= 8 \frac{\operatorname{sn}^2(-Kv(\tau+1)/2, k)}{\operatorname{cd}^2(-Kv(\tau+1)/2, k) \operatorname{sd}(-Kv(\tau+1), k)} \\ &= -8 \frac{\operatorname{sn}^2(Kv(\tau+1)/2, k)}{\operatorname{cd}^2(Kv(\tau+1)/2, k) \operatorname{sd}(Kv(\tau+1), k)}, \end{aligned}$$

where we have used the facts that sd and sn are odd and cd is even. Recall that $v = 2u/(K + iK') = 2u/(K(\tau + 1))$, so that we conclude

$$\int_{\mathbb{R}} \frac{\sqrt{x} \sin(u\sqrt{x})}{\cos(K\sqrt{x}) - \cosh(K'\sqrt{x})} dx = -8 \frac{\operatorname{sn}^2(u, k)}{\operatorname{cd}^2(u, k) \operatorname{sd}(2u, k)}$$

whenever $u = v(K + iK')/2$ with $v \in (-1, 1)$. Moreover, the integral converges absolutely and uniformly on compact subsets of the domain $D := \{u \in \mathbb{C} : |\Re(u)| < K/2 \text{ and } |\Im(u)| < K'/2\}$ and therefore defines an analytic function on D . Meanwhile, by locating the poles of sn and the zeros of cd and sd using [11, Table 22.4.1], we see that this quotient of Jacobi elliptic functions is analytic on D . Then by the identity theorem, the identity (10) holds on all of D .

The alternate expression of the generating function in terms of the log-derivative of the nc elliptic function is obtained by elementary transformations of the Jacobi elliptic functions as follows: observe that, by [25, vol 3, (839)]

$$\begin{aligned} \frac{\text{sn}(u, k)^2}{\text{cd}(u, k)^2} \frac{1}{\text{sd}(2u, k)} &= \frac{1 - \text{cn}(2u, k)}{1 + \text{dn}(2u, k)} \frac{1 + \text{dn}(2u, k)}{1 + \text{cn}(2u, k)} \frac{1}{\text{sd}(2u, k)} \\ &= \frac{1 - \text{cn}(2u, k)}{1 + \text{cn}(2u, k)} \frac{1}{\text{sd}(2u, k)} \\ &= \frac{1}{2} \frac{d}{du} \frac{1}{1 + \text{cn}(2u, k)}. \end{aligned}$$

Define $\bar{nc}(u, k) = \frac{d}{du} \log nc(u, k)$ so that, from [25, vol 3, (848)],

$$\frac{1}{1 + \text{cn}(2u, k)} = \frac{1}{2} (1 + \bar{nc}^2(u, k)),$$

and we deduce

$$\frac{\text{sn}(u, k)^2}{\text{cd}(u, k)^2} \frac{1}{\text{sd}(2u, k)} = \frac{1}{4} \frac{d}{du} \bar{nc}^2(u, k).$$

□

By Taylor expanding the sine function, we can now obtain an evaluation of the Berndt-type integrals I_n^- in terms of Jacobi elliptic functions.

Corollary 5. *Let $k \in (0, 1)$ and $n \geq 0$. Then*

$$\int_{\mathbb{R}} \frac{x^{n+1}}{\cos(K\sqrt{x}) - \cosh(K'\sqrt{x})} dx = (-1)^{n+1} 8 \frac{d^{2n+1}}{du^{2n+1}} \frac{\text{sn}^2(u, k)}{\text{cd}^2(u, k) \text{sd}(2u, k)} \Big|_{u=0}, \quad (11)$$

and if $n = 2m$ is even, it follows that

$$\int_0^\infty \frac{x^{4m+3}}{\cos(x) - \cosh(x)} dx = -2 \left(\frac{\pi^{\frac{3}{2}}}{2\Gamma^2(\frac{3}{4})} \right)^{4m+4} \frac{d^{4m+1}}{du^{4m+1}} \frac{\text{sn}^2(u, 1/\sqrt{2})}{\text{cd}^2(u, 1/\sqrt{2}) \text{sd}(2u, 1/\sqrt{2})} \Big|_{u=0}.$$

Proof. Expanding the sine in (10) produces

$$\sum_{m \geq 0} (-1)^m \frac{u^{2m+1}}{(2m+1)!} \int_{\mathbb{R}} \frac{x^{m+1}}{\cos(K\sqrt{x}) - \cosh(K'\sqrt{x})} dx = -8 \frac{\text{sn}^2(u, k)}{\text{cd}^2(u, k) \text{sd}(2u, k)},$$

so that differentiating $2n+1$ times recovers (11). In the lemniscatic case $k = 1/\sqrt{2}$, we have

$$\int_{\mathbb{R}} \frac{x^{n+1}}{\cos(K\sqrt{x}) - \cosh(K\sqrt{x})} dx = (-1)^{n+1} 8 \frac{d^{2n+1}}{du^{2n+1}} \frac{\text{sn}^2(u, 1/\sqrt{2})}{\text{cd}^2(u, 1/\sqrt{2}) \text{sd}(2u, 1/\sqrt{2})} \Big|_{u=0},$$

where $K = K(1/\sqrt{2}) = \frac{\pi^{\frac{3}{2}}}{2\Gamma^2(\frac{3}{4})}$. Substituting $K\sqrt{x} \rightarrow x$, we have

$$\frac{2}{K^{2n+4}} \int_{\mathbb{L}} \frac{x^{2n+3}}{\cos(x) - \cosh(x)} dx = (-1)^{n+1} 8 \frac{d^{2n+1}}{du^{2n+1}} \frac{\text{sn}^2(u, 1/\sqrt{2})}{\text{cd}^2(u, 1/\sqrt{2}) \text{sd}(2u, 1/\sqrt{2})} \Big|_{u=0},$$

and now substituting $ix \rightarrow x$ on the vertical part of the contour yields

$$\frac{2}{K^{2n+4}} ((-1)^n + 1) \int_0^\infty \frac{x^{2n+3}}{\cos(x) - \cosh(x)} dx = (-1)^{n+1} 8 \frac{d^{2n+1}}{du^{2n+1}} \frac{\text{sn}^2(u, 1/\sqrt{2})}{\text{cd}^2(u, 1/\sqrt{2}) \text{sd}(2u, 1/\sqrt{2})} \Big|_{u=0},$$

so that when n is even,

$$\begin{aligned} \int_0^\infty \frac{x^{2n+3}}{\cos(x) - \cosh(x)} dx &= -2K^{2n+4} \frac{d^{2n+1}}{du^{2n+1}} \frac{\operatorname{sn}^2(u, 1/\sqrt{2})}{\operatorname{cd}^2(u, 1/\sqrt{2})\operatorname{sd}(2u, 1/\sqrt{2})} \Big|_{u=0} \\ &= -2 \left(\frac{\pi^{\frac{3}{2}}}{2\Gamma^2(\frac{3}{4})} \right)^{2n+4} \frac{d^{2n+1}}{du^{2n+1}} \frac{\operatorname{sn}^2(u, 1/\sqrt{2})}{\operatorname{cd}^2(u, 1/\sqrt{2})\operatorname{sd}(2u, 1/\sqrt{2})} \Big|_{u=0}. \end{aligned}$$

Letting $n = 2m$ completes the proof. \square

4.2 More Symmetry Results

This generalization of Kuznetsov's result (10) allow us to establish additional identities between the two families of integrals I_n^- and I_n^+ .

Proposition 12. *From (10) and (4), we deduce the symmetry*

$$\int_{\mathbb{R}} \frac{\sqrt{x} \sin(u\sqrt{x})}{\cos(K\sqrt{x}) - \cosh(K'\sqrt{x})} dx = - \left(\int_{\mathbb{R}} \frac{\sin(u\sqrt{x})}{\sqrt{x}} \frac{dx}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} \right) \left(\int_{\mathbb{R}} \frac{\cos(u\sqrt{x})}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} dx \right).$$

At the level of the integrals

$$I_n^- = \int_{\mathbb{R}} \frac{x^{n+1}}{\cos K\sqrt{x} - \cosh K'\sqrt{x}} dx$$

and

$$I_n^+ = \frac{1}{2} \int_{\mathbb{R}} \frac{x^n}{\cos K\sqrt{x} + \cosh K'\sqrt{x}} dx,$$

these results translate into the binomial convolution identity

$$I_n^- = -4 \sum_{p=0}^n \binom{2n+1}{2p+1} I_p^+ I_{n-p}^+$$

and, with $c = K'(k)/K(k)$, $q = e^{-\pi c}$, into the identity

$$\begin{aligned} 2I_{n+1}^+ - I_n^- &= 8 \left(\frac{\pi}{K(k)} \right)^{2n+4} \sum_{m \geq 1} \frac{m^{2n+3} q^m}{1 - q^{2m}} \\ &= 8 \left(\frac{\pi}{K(k)} \right)^{2n+4} \left(-\frac{E_{2n+3}(0)}{4c^{2n+4}} + (-1)^n \frac{(2n+3)!}{\pi^{2n+4}} \sum_{p \geq 1, q \geq 1} \left(\frac{1}{(2p+ic(2q-1))^{2n+4}} + \frac{1}{(2p-ic(2q-1))^{2n+4}} \right) \right) \end{aligned} \quad (12)$$

Proof. From the two generating functions

$$J_+(u) = \int_{\mathbb{R}} \frac{\sin u\sqrt{x}}{\sqrt{x}} \frac{dx}{\cos K\sqrt{x} + \cosh K'\sqrt{x}} = 2\widetilde{\operatorname{nc}}(u, k) = \sum_{n \geq 0} \frac{(-1)^n}{(2n+1)!} u^{2n+1} I_n^+$$

and

$$J_-(u) = \int_{\mathbb{R}} \frac{\sqrt{x} \sin(u\sqrt{x})}{\cos K\sqrt{x} + \cosh K'\sqrt{x}} dx = -2 \frac{d}{du} \widetilde{\operatorname{nc}}^2(u, k) = \sum_{n \geq 0} \frac{(-1)^n}{(2n+1)!} u^{2n+1} I_n^-,$$

we deduce

$$\begin{aligned} J_-(u) &= -2\widetilde{\operatorname{nc}}(u, k) \frac{d}{du} 2\widetilde{\operatorname{nc}}(u, k) \\ &= - \sum_{p, q \geq 0} \frac{(-1)^p u^{2p+1}}{(2p+1)!} I_p^+ \frac{(-1)^q u^{2q}}{(2q)!} I_q^+ \\ &= - \sum_{p, n \geq 0} \frac{(-1)^n u^{2n+1}}{(2p+1)! (2n-2p)!} I_p^+ I_{n-p}^+ \end{aligned}$$

so that

$$I_n^- = - \sum_{p \geq 0} \binom{2n+1}{2p+1} I_p^+ I_{n-p}^+.$$

Moreover, from

$$I_{n+1}^+ = \frac{1}{2} \left(\frac{\pi}{K} \right)^{2n+4} \left(-E_{2n+3}(0) + 4 \sum_{p \geq 1} \frac{m^{2n+3} q^m}{1 + (-q)^m} \right)$$

and

$$I_n^- = - \left(\frac{\pi}{K} \right)^{2n+4} \left(E_{2n+3}(0) + 4 \sum_{p \geq 1} \frac{m^{2n+3} q^m}{1 - (-q)^m} \right),$$

we deduce the result by subtracting the second identity from twice the first one. The Lambert series is identified in [19] as the Weierstraß series

$$\sum_{m \geq 1} \frac{m^{2s-1}}{\sinh(\pi mc)} = \frac{2(2s-1)!}{\pi^{2s}} \left(\frac{U_{2s}}{c^{2s}} + (-1)^s \sum_{n \geq 1, m \geq 1} \left(\frac{1}{(2m + ic(2n-1))^{2s}} + \frac{1}{(2m - ic(2n-1))^{2s}} \right) \right)$$

with $U_{2s} = -\frac{\pi^{2s}}{4(2s-1)!} E_{2s-1}(0)$, so that

$$\sum_{m \geq 1} \frac{m^{2s-1}}{\sinh(\pi mc)} = 2 \left(-\frac{1}{4} E_{2s-1}(0) \frac{1}{c^{2s}} + (-1)^s \frac{(2s-1)!}{\pi^{2s}} \sum_{n \geq 1, m \geq 1} \left(\frac{1}{(2m + ic(2n-1))^{2s}} + \frac{1}{(2m - ic(2n-1))^{2s}} \right) \right)$$

and we have

$$2I_n^+ - I_n^- = 4 \left(\frac{\pi}{K(k)} \right)^{2n+4} \sum_{m \geq 1} \frac{m^{2n+3}}{\sinh(\pi mc)}$$

with $c = K'(k)/K(k)$, so that

$$\begin{aligned} 2I_n^+ - I_n^- &= 4 \left(\frac{\pi}{K(k)} \right)^{2n+4} \sum_{m \geq 1} \frac{m^{2n+3}}{\sinh(\pi mc)} \\ &= 8 \left(\frac{\pi}{K(k)} \right)^{2n+4} \left(-\frac{E_{2n+3}(0)}{4c^{2n+4}} + (-1)^n \frac{(2n+3)!}{\pi^{2n+4}} \sum_{p \geq 1, q \geq 1} \left(\frac{1}{(2p + ic(2q-1))^{2n+4}} + \frac{1}{(2p - ic(2q-1))^{2n+4}} \right) \right) \end{aligned}$$

□

4.3 The Polynomials of Lomont and Brillhart

Evaluating the first values of the integrals I_n^- produces

$$I_0^- = -4, \quad I_1^- = 32(1 - 2k^2), \quad I_2^- = -32(17 - 32k^2 + 32k^4),$$

suggesting that, as in the case of the integrals I_n^+ , they can be expressed as polynomials in the elliptic modulus k . This is confirmed by the following result.

Proposition 13. *Denote*

$$Q_n(x, y) = \sum_{p=0}^n \binom{2n+2}{2p+1} P_p(x, y) P_{n-p}(x, y) \quad (13)$$

the binomial convolution of the P_n elliptic polynomials with themselves. Then we have

$$\int_{\mathbb{R}} \frac{x^{n+1}}{\cos(K\sqrt{x}) - \cosh(K'\sqrt{x})} dx = (-2)^{n+1} Q_n(1 - 2k^2, 4k^4 - 4k^2 + 4). \quad (14)$$

Proof. Start from

$$\int_{\mathbb{R}} \frac{\sqrt{x} \sin(u\sqrt{x})}{\cos(K\sqrt{x}) - \cosh(K'\sqrt{x})} dx = -2 \frac{d}{du} \widetilde{\text{nc}}^2(u, k). \quad (15)$$

With $x = 1 - 2k^2$ and $y = 4k^4 - 4k^2 + 4$, we have

$$\widetilde{\text{nc}}(u, k) = \sum_{n \geq 0} 2^n P_n(x, y) \frac{u^{2n+1}}{(2n+1)!},$$

so that

$$\begin{aligned} \widetilde{\text{nc}}^2(u, k) &= \sum_{p, q \geq 0} 2^{p+q} P_p(x, y) P_q(x, y) \frac{u^{2p+1}}{(2p+1)!} \frac{u^{2q+1}}{(2q+1)!} \\ &= \sum_{n, p \geq 0} 2^n P_p(x, y) P_{n-p}(x, y) \frac{u^{2n+2}}{(2p+1)!(2n-2p+1)!} \\ &= \sum_{n \geq 0} 2^n Q_n(x, y) \frac{u^{2n+2}}{(2n+2)!}, \end{aligned}$$

where we have defined

$$Q_n(x, y) = \sum_{p=0}^n \binom{2n+2}{2p+1} P_p(x, y) P_{n-p}(x, y).$$

We deduce, still with $x = 1 - 2k^2$ and $y = 4k^4 - 4k^2 + 4$,

$$\begin{aligned} \int_{\mathbb{R}} \frac{\sqrt{x} \sin(u\sqrt{x})}{\cos(K\sqrt{x}) - \cosh(K'\sqrt{x})} dx &= -2 \frac{d}{du} \widetilde{\text{nc}}^2(u, k) \\ &= - \sum_{n \geq 0} 2^{n+1} Q_n(x, y) \frac{u^{2n+1}}{(2n+1)!}. \end{aligned}$$

The Taylor expansion

$$\int_{\mathbb{R}} \frac{\sqrt{x} \sin(u\sqrt{x})}{\cos(K\sqrt{x}) - \cosh(K'\sqrt{x})} dx = \sum_{n \geq 0} \frac{(-1)^n u^{2n+1}}{(2n+1)!} \int_{\mathbb{R}} \frac{x^{n+1}}{\cos(K\sqrt{x}) - \cosh(K'\sqrt{x})} dx$$

produces, upon identification with (15),

$$\int_{\mathbb{R}} \frac{x^{n+1}}{\cos(K\sqrt{x}) - \cosh(K'\sqrt{x})} dx = (-2)^{n+1} Q_n(x, y)$$

which is the desired result. \square

The first values of the polynomials $Q_n(x, y)$ are

$$Q_0(x, y) = 2, \quad Q_1(x, y) = 8x, \quad Q_2(x, y) = 20x^2 + 12y.$$

Notice that these polynomials appear in [20, Table 4.2] but they differ from the ones defined here by a factor 2.

Corollary 6. *If k^2 is a rational number, then the integral $I_n^- = \int_{\mathbb{R}} \frac{x^{n+1}}{\cos(K\sqrt{x}) - \cosh(K'\sqrt{x})} dx$ is a rational number.*

Moreover, since

$$I_1^- = 32(1 - 2k^2), \quad I_2^- = -32(17 - 32k^2 + 32k^4),$$

the value of I_n^- for $n \geq 3$ is a polynomial function of the initial values I_1 and I_2 given by

$$I_n^- = (-2)^{n+1} Q_n \left(\frac{1}{32} I_1^-, \frac{1}{8} \left(15 - \frac{I_2^-}{32} \right) \right).$$

Finally, with $x = 1 - 2k^2$ and $y = 4k^4 - 4k^2 + 4$, the integrals I_n^- satisfy the recurrence

$$I_{n+2}^- = -8x I_{n+1}^- - (y - x^2) \sum_{j=0}^n \binom{2n+4}{2j+2} I_j^- I_{n-j}^- \quad (16)$$

with initial values $I_0^- = -4$ and $I_1^- = 32(1 - 2k^2)$.

Proof. The integrality of the coefficients of $Q_n(x, y)$ is a consequence of the integrality of the coefficients of $P_n(x, y)$ and of their expression (13). The expression of I_n^- in terms of the initial values I_1^- and I_2^- is obtained from (14) by substituting x and y as functions of I_1^- and I_2^- respectively. Recurrence (16) is deduced from recurrence [20, (4.33)] where the Q_n that appear there are half those defined here. \square

Additional recursive identities between the two sets of polynomials $P_n(x, y)$ and $Q_n(x, y)$ can be found in [20, Chapter 4]; they induce identities between the two sets of integrals I_n^+ and I_n^- . For example, [20, (4.42)] induces the identity, for $n \geq 0$,

$$\sum_{j=0}^n (n-3j) \binom{2n+3}{2j+1} I_j^+ I_{n-j}^- = 0.$$

4.4 A Probabilistic Approach

The previous results can be given a probabilistic interpretation. Define the discrete random variable X by

$$\Pr \left\{ X = \pm \frac{(2n-1)\pi}{\chi} \right\} = \frac{\pi}{k'K'} \frac{q^{n-\frac{1}{2}}}{1+q^{2n-1}},$$

where $n \geq 1$, $\chi = \frac{K(k)}{K(k')}$, and $q = e^{-\pi \frac{K(k)}{K'(k)}}$. Then we can compute the moment generating function and the cumulants of X .

Proposition 14. *The moment generating function of X is*

$$\varphi_X(u) = \mathbb{E}e^{uX} = \text{nc}(u, k)$$

Proof. The proof is a straightforward computation. We have

$$\begin{aligned} \mathbb{E}e^{uX} &= \sum_{n \in \mathbb{Z}} p_n e^{ux_n} = \sum_{n \geq 1} \frac{\pi}{k'K'} \frac{q^{n-\frac{1}{2}}}{1+q^{2n-1}} e^{\frac{(2n-1)\pi}{\chi} u} + \sum_{n \geq 1} \frac{\pi}{k'K'} \frac{q^{n-\frac{1}{2}}}{1+q^{2n-1}} e^{-\frac{(2n-1)\pi}{\chi} u} \\ &= \frac{2\pi}{k'K'} \sum_{n \geq 1} \frac{q^{n-\frac{1}{2}}}{1+q^{2n-1}} \cosh\left(\frac{(2n-1)\pi}{\chi} u\right) = \text{nc}(u, k) \end{aligned}$$

by [24, p.19]. \square

Proposition 15. *For $n \geq 1$, the cumulants κ_{2n} of the discrete random variable X coincide with the moments I_{n-1}^+ of a continuous random variable Y with the probability density*

$$f_Y(z) = \frac{1}{2} \frac{1}{\cos(K\sqrt{z}) + \cosh(K'\sqrt{z})}.$$

Proof. The positivity of the function f_Y and the fact that $\int_{\mathbb{R}} f_Y(y) dy = 1$ make f_Y a probability density function. Moreover, the cumulants κ_X of X are defined by

$$\log(\text{nc}(iu, k)) = \sum_{n \geq 1} \kappa_n \frac{(iu)^n}{n!} = \sum_{n \geq 1} \kappa_{2n} (-1)^n \frac{u^{2n}}{(2n)!}$$

since $\log(\text{nc}(u, k))$ is an even function of u . Moreover, by [24, p.19],

$$\frac{d}{du} \log(\text{nc}(u, k)) = \sum_{n \geq 0} \frac{u^{2n+1}}{(2n+1)!} I_n^+$$

with

$$I_n^+ = \frac{1}{2} \int \frac{x^n}{\cos(K\sqrt{x}) + \cosh(K'\sqrt{x})} dx$$

so that, since $\text{nc}(0, k) = 1$, we have

$$\log(\text{nc}(iu, k)) = \sum_{n \geq 0} (-1)^n \frac{u^{2n+2}}{(2n+2)!} I_n^+,$$

and it follows that $\kappa_{2n} = I_{n-1}^+$ for $n \geq 1$. \square

4.5 Lambert Series Representation

As in Kuznetsov's case, our extension produces a Lambert series representation for Berndt-type integrals.

Proposition 16. *The I_n^- integrals have the following Lambert series representation*

$$\int_{\mathbb{R}} \frac{x^{n+1}}{\cos(K\sqrt{x}) - \cosh(K'\sqrt{x})} dx = - \left(\frac{\pi}{K}\right)^{2n+4} \left(E_{2n+3}(0) + 4 \sum_{m \geq 1} \frac{m^{2n+3} q^m}{1 - (-q)^m} \right)$$

with $E_n(x)$ the n -th Euler polynomial.

Proof. From [24, p. 134], the Lambert expansion for the generating function is

$$\frac{d}{du} \operatorname{nc}^2(u, k) = \frac{\pi^3}{4K^3} \frac{\tan\left(\frac{\pi u}{2K}\right)}{\cos^2\left(\frac{\pi u}{2K}\right)} + \frac{2\pi^3}{K^3} \sum_{n \geq 1} \frac{n^2 q^n \sin\left(\frac{n\pi u}{K}\right)}{1 - (-q)^n}.$$

The Taylor expansion

$$\tan z = \sum_{n \geq 1} (-1)^{n-1} \frac{2^{2n} (2^{2n} - 1) B_{2n}}{2n!} z^{2n-1}$$

together with the expression of the scaled Bernoulli numbers as Euler polynomials

$$(2^{2n} - 1) B_{2n} = -n E_{2n-1}(0)$$

produces

$$\tan z = \sum_{n \geq 1} (-1)^n \frac{2^{2n-1} E_{2n-1}(0)}{2n-1!} z^{2n-1}.$$

Since

$$\frac{\tan z}{\cos^2 z} = \frac{1}{2} \frac{d^2}{dz^2} \tan z,$$

we deduce

$$\frac{\tan z}{\cos^2 z} = \frac{1}{2} \frac{d^2}{dz^2} \sum_{n \geq 1} (-1)^n \frac{2^{2n-1} E_{2n-1}(0)}{(2n-1)!} z^{2n-1} = \sum_{n \geq 1} (-1)^{n-1} \frac{2^{2n} E_{2n+1}(0)}{(2n-1)!} z^{2n-1}.$$

Moreover, expanding

$$\sum_{n \geq 1} \frac{n^2 q^n \sin\left(\frac{n\pi u}{K}\right)}{1 - (-q)^n} = \sum_{n \geq 1} \frac{n^2 q^n}{1 - (-q)^n} \sum_{p \geq 0} \frac{(-1)^p}{(2p+1)!} \left(\frac{n\pi u}{K}\right)^{2p+1} = \sum_{p \geq 0} \frac{(-1)^p}{(2p+1)!} \left(\frac{\pi u}{K}\right)^{2p+1} \sum_{n \geq 1} \frac{n^{2p+3} q^n}{1 - (-q)^n}$$

produces

$$\begin{aligned} 2 \frac{\operatorname{sn}(u, k)^2}{\operatorname{cd}(u, k)^2} \frac{1}{\operatorname{sd}(2u, k)} &= \frac{\pi^3}{8K^3} \sum_{p \geq 0} (-1)^p \frac{2^{2p+2} E_{2p+3}(0)}{(2p+1)!} \left(\frac{\pi u}{2K}\right)^{2p+1} + \frac{\pi^3}{K^3} \sum_{p \geq 0} \frac{(-1)^p}{(2p+1)!} \left(\frac{\pi u}{K}\right)^{2p+1} \sum_{n \geq 1} \frac{n^{2p+3} q^n}{1 - (-q)^n} \\ &= \sum_{p \geq 0} (-1)^p \left(\frac{\pi}{K}\right)^{2p+4} \frac{u^{2p+1}}{(2p+1)!} \left[\frac{1}{4} E_{2p+3}(0) + \sum_{n \geq 1} \frac{n^{2p+3} q^n}{1 - (-q)^n} \right]. \end{aligned}$$

It follows that

$$2 \frac{d^{2n+1}}{du^{2n+1}} \frac{\operatorname{sn}(u, k)^2}{\operatorname{cd}(u, k)^2} \frac{1}{\operatorname{sd}(2u, k)} \Big|_{u=0} = (-1)^n \left(\frac{\pi}{K}\right)^{2n+4} \left[\frac{1}{4} E_{2n+3}(0) + \sum_{m \geq 1} \frac{m^{2n+3} q^m}{1 - (-q)^m} \right]$$

and

$$\int_{\mathbb{R}} \frac{x^{n+1}}{\cos(K\sqrt{x}) - \cosh(K'\sqrt{x})} dx = - \left(\frac{\pi}{K}\right)^{2n+4} \left[E_{2n+3}(0) + 4 \sum_{m \geq 1} \frac{m^{2n+3} q^m}{1 - (-q)^m} \right].$$

□

4.6 Eisenstein Series Representation

Proposition 17. *The I_n^- integrals have the Eisenstein series expansion*

$$I_n^- = \int_{\mathbb{R}} \frac{x^{n+1}}{\cos(K\sqrt{x}) - \cosh(K'\sqrt{x})} dx = (-1)^{n+1} 2(2n+3)! \sum_{(p,q) \in \mathbb{Z}^2} \frac{1}{((2q+p-1)K + ipK')^{2n+4}}.$$

Proof. The Weierstraß \wp_6 function has the double series representation [26, vol 5 p.10]

$$\wp_6(z, k) = \sum_{(p,q) \in \mathbb{Z}^2} \frac{1}{((2q+p-1)K + piK' - z)^2} - \frac{c_{p,q}}{((2q+p)K + piK')^2}$$

with

$$c_{p,q} = \begin{cases} 0, & p = q = 0 \\ 1, & \text{else,} \end{cases}$$

so that its derivative is

$$\wp_6'(u, k) = 2 \sum_{(p,q) \in \mathbb{Z}^2} \frac{1}{((2q+p-1)K + ipK' - u)^3}.$$

with Taylor expansion in u

$$\wp_6'(u, k) = 2 \sum_{n \geq 0} \binom{n+2}{2} u^n \sum_{(p,q) \in \mathbb{Z}^2} \frac{1}{((2q+p-1)K + ipK')^{n+3}}.$$

Comparing with the expansion

$$\int_{\mathbb{R}} \frac{\sqrt{x} \sin(u\sqrt{x})}{\cos(\sqrt{x}) - \cosh(K'\sqrt{x})} dx = \sum_{n \geq 0} \frac{(-1)^n}{(2n+1)!} u^{2n+1} I_n^- = -2\wp_6'(u, k)$$

where

$$I_n^- = \int_{\mathbb{R}} \frac{x^{n+1}}{\cos(K\sqrt{x}) - \cosh(K'\sqrt{x})} dx,$$

we obtain

$$\frac{(-1)^n}{(2n+1)!} I_n^- = -4 \binom{2n+3}{2} \sum_{(p,q) \in \mathbb{Z}^2} \frac{1}{((2q+p-1)K + ipK')^{2n+4}}$$

or

$$I_n^- = \int_{\mathbb{R}} \frac{x^{n+1}}{\cos(K\sqrt{x}) - \cosh(K'\sqrt{x})} dx = (-1)^{n+1} 2(2n+3)! \sum_{(p,q) \in \mathbb{Z}^2} \frac{1}{((2q+p-1)K + ipK')^{2n+4}}.$$

□

5 Conclusion and Perspectives

Berndt-type integrals are surprisingly rich with connections to various special functions. Central to our investigation was the utilization of the Barnes zeta function, which provided a powerful framework for evaluating Berndt-type integrals in terms of a multiple series representation. This approach not only extended the scope of known evaluations but also allowed us to exhibit the analytic continuation of Berndt-type integrals. Moreover, we have extended Kuznetsov's direct evaluation of the integral considered by Ismail and Valent related to the Nevanlinna parametrization of solutions to a certain indeterminate moment problem [5]. Kuznetsov's evaluations provide a direct link between the generating functions of Jacobi elliptic functions and integrals involving hyperbolic and trigonometric functions. By leveraging Kuznetsov's findings, we have demonstrated specific instances where Berndt-type integrals can be expressed in terms of Jacobi elliptic functions, thereby establishing a richer analytical understanding of their nature. This not only enhances our ability to compute these integrals but also opens avenues for exploring their broader implications within the realm of special functions and mathematical physics.

Through our investigations, we have demonstrated the versatility of a variety of approaches in handling Berndt-type integrals, from Lambert series representations to Barnes zeta function evaluations. Each method offers unique

insights and avenues for further exploration, revealing connections to broader classes of mathematical objects such as moment polynomials and Bernoulli-Barnes polynomials. In essence, the study of Berndt-type integrals exemplifies the enduring quest within mathematics to unify seemingly disparate concepts and reveal underlying symmetries. As we continue to delve deeper into their properties and connections—whether through the lens of Barnes zeta functions, Jacobi elliptic functions, or other mathematical frameworks—we anticipate further revelations and applications across disciplines. Thus, Berndt-type integrals not only present challenges in integration theory but also serve as gateways to new mathematical connections.

The fact that Lambert series can be expressed as polynomials in the elliptic modulus appears in Chapter 17 of Ramanujan’s work [7, Chapter 17] and more recently in S. Cooper’s work [9]. As we have shown, the study of these elliptic polynomials by Lomont and Brillhart gives detailed information about the Lambert series and, in turn, about Berndt’s integrals. A systematic way to link Lambert series to Eisenstein series is the object of several articles by Ling [17, 18]. In a future companion paper, our ultimate goal is to produce an integral representation for all elementary Lambert series $\sum_{n \geq 1} n^s \frac{(\pm 1)^n q^{a_n}}{1 \pm (\pm q)^{b_n}}$ with $(a_n, b_n) \in \{n, 2n\}$, their associated Eisenstein series and Barnes zeta series. A first example can in fact be deduced from (12) under the form

$$4 \left(\frac{\pi}{K(k)} \right)^{2n+4} \sum_{m \geq 1} \frac{m^{2n+3} q^m}{1 - q^{2m}} = - \int_{\mathbb{R}} \frac{x^{n+1} \cosh(K' \sqrt{x})}{\cos^2(K \sqrt{x}) - \cosh^2(K' \sqrt{x})} dx$$

As in the examples shown in this article, the integral representation contains symmetries that produce non elementary identities for these series.

References

- [1] Tewodros Amdeberhan, Olivier Espinosa, Ivan Gonzalez, Marshall Harrison, Victor Moll, and Armin Straub. Ramanujan’s master theorem. *The Ramanujan Journal*, 29, 12 2012. doi:10.1007/s11139-011-9333-y.
- [2] E. W. Barnes. On the theory of the multiple gamma function. *Trans. Cambridge Philos. Soc.*, 19:374–425, 1904.
- [3] Ernest William Barnes and Andrew Russell Forsyth. The theory of the double gamma function. *Philosophical Transactions of the Royal Society of London. Series A, Containing Papers of a Mathematical or Physical Character*, 196(274-286):265–387, 1901. doi:10.1098/rsta.1901.0006.
- [4] Abdelmejid Bayad and Matthias Beck. Relations for Bernoulli–Barnes numbers and Barnes zeta functions. *International Journal of Number Theory*, 10:1321–1335, 2013.
- [5] Christian Berg and Galliano Valent. The Nevanlinna parametrization for some indeterminate Stieltjes moment problems associated with birth and death processes. *Methods and applications of analysis*, 1, 1994.
- [6] Bruce Berndt. Integrals associated with Ramanujan and elliptic functions. *The Ramanujan Journal*, 41, 11 2016. doi:10.1007/s11139-016-9793-1.
- [7] Bruce C. Berndt. *Ramanujan’s Notebooks: Part III*. Springer, Germany, 1991. doi:10.1007/978-1-4612-0965-2.
- [8] Zachary P. Bradshaw and Christophe Vignat. An operational calculus generalization of Ramanujan’s master theorem. *Journal of Mathematical Analysis and Applications*, 523(2):127029, 2023. doi:10.1016/j.jmaa.2023.127029.
- [9] Shaun Cooper and H.Y. Lam. Sixteen Eisenstein series. *Ramanujan Journal*, 18:33–59, 10 2009. doi:10.1007/s11139-007-9039-3.
- [10] Atul Dixit, Victor Moll, and Christophe Vignat. The Zagier modification of Bernoulli numbers and a polynomial extension. Part I. *The Ramanujan Journal*, 33, 09 2012. doi:10.1007/s11139-013-9484-0.
- [11] *NIST Digital Library of Mathematical Functions*. <https://dlmf.nist.gov/>, Release 1.2.0 of 2024-03-15. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.
- [12] Eduardo Friedman and Simon Ruijsenaars. Shintani–Barnes zeta and gamma functions. *Advances in Mathematics*, 187(2):362–395, 2004. doi:10.1016/j.aim.2003.07.020.

- [13] I.S. Gradshteyn and I.M. Ryzhik. 1 - elementary functions. In Daniel Zwillinger, Victor Moll, I.S. Gradshteyn, and I.M. Ryzhik, editors, *Table of Integrals, Series, and Products (Eighth Edition)*, pages 25–62. Academic Press, Boston, eighth edition edition, 2014. doi:10.1016/B978-0-12-384933-5.00001-1.
- [14] Mourad E. H. Ismail and Galliano Valent. On a family of orthogonal polynomials related to elliptic functions. *Illinois Journal of Mathematics*, 42:294–312, 1998.
- [15] Lin Jiu, Victor H. Moll, and Christophe Vignat. A symbolic approach to some identities for Bernoulli–Barnes polynomials. *International Journal of Number Theory*, 12(03):649–662, March 2016. doi:10.1142/s1793042116500421.
- [16] Alexey Kuznetsov. A direct evaluation of an integral of Ismail and Valent. In *Frontiers in Orthogonal Polynomials and q -Series*, chapter 21, pages 417–424. World Scientific, 2017. doi:10.1142/9789813228887_0021.
- [17] Chih-Bing Ling. On summation of series of hyperbolic functions. *SIAM Journal on Mathematical Analysis*, 5(4):551–562, 1974. doi:10.1137/0505055.
- [18] Chih-Bing Ling. On summation of series of hyperbolic functions. II. *SIAM Journal on Mathematical Analysis*, 6(1):129–139, 1975. doi:10.1137/0506013.
- [19] Chih-Bing Ling. Generalization of certain summations due to Ramanujan. *SIAM Journal on Mathematical Analysis*, 9(1):34–48, 1978. doi:10.1137/0509003.
- [20] J.S. Lomont and J. Brillhart. *Elliptic Polynomials*. Taylor & Francis, 2000.
- [21] Stephen Milne. Infinite families of exact sums of squares formulas, Jacobi elliptic functions, continued fractions, and Schur functions. *Ramanujan Journal*, 6, 09 2000. doi:10.1023/A:1014865816981.
- [22] Srinivasa Ramanujan. On certain arithmetical functions. *Trans. Camb. Philos. Soc.*, 22:159–184, 1916.
- [23] S.N.M. Ruijsenaars. On Barnes’ multiple zeta and gamma functions. *Advances in Mathematics*, 156(1):107–132, 2000. doi:10.1006/aima.2000.1946.
- [24] F. Tölke. *Theta-Funktionen und spezielle Weierstraßsche Funktionen*, volume 2 of *Praktische Funktionenlehre*. Springer, 1966.
- [25] F. Tölke. *Jacobische elliptische Funktionen, Legendresche elliptische Normalintegrale und spezielle Weierstraßsche Zeta- und Sigma-Funktionen*, volume 3 of *Praktische Funktionenlehre*. Springer-Verlag, Berlin/Heidelberg/NewYork, 1967.
- [26] F. Tölke. *Allgemeine Weierstraßsche Funktionen und Ableitungen nach dem Parameter. Integrale der Theta-Funktionen und Bilinear-Entwicklungen*, volume 5 of *Praktische Funktionenlehre*. Springer, 1968.
- [27] E.T. Whittaker and G.N. Watson. *A Course of Modern Analysis*. A Course of Modern Analysis: An Introduction to the General Theory of Infinite Processes and of Analytic Functions, with an Account of the Principal Transcendental Functions. Cambridge University Press, 1996.
- [28] Ce Xu and Jianqiang Zhao. Berndt-type integrals and series associated with Ramanujan and Jacobi elliptic functions, 2023. arXiv:2301.08211.
- [29] Ce Xu and Jianqiang Zhao. General Berndt-type integrals and series associated with Jacobi elliptic functions, 2024. arXiv:2401.01385.

Appendix

An alternative to the approach in Section 2 is to make use of a symbolic technique involving the Bernoulli-Barnes and Euler-Barnes polynomials [4, 15]. With symbolic notations, the Bernoulli-Barnes polynomials are

$$B_p^{(2)}(x; a_1, a_2) = (x + a_1 B_1 + a_2 B_2)^p$$

and the Euler-Barnes polynomials are

$$E_p^{(2)}(x; a_1, a_2) = (x + a_1 E_1 + a_2 E_2)^p,$$

where B_i is the i -th Bernoulli number and E_i is the i -th Euler number. Their generating functions are

$$\sum_{n \geq 0} \frac{B_n^{(2)}(x; a_1, a_2)}{n!} z^n = e^{zx} \frac{a_1 z}{e^{a_1 z} - 1} \frac{a_2 z}{e^{a_2 z} - 1}$$

and

$$\sum_{n \geq 0} \frac{E_n^{(2)}(x; a_1, a_2)}{n!} z^n = e^{zx} \frac{2}{e^{a_1 z} + 1} \frac{2}{e^{a_2 z} + 1}.$$

We will make use of the following lemma.

Lemma 1. *The following expansions hold:*

$$\frac{x^2}{\cosh x - \cos x} = \frac{ix^2}{2} \csc\left(\sqrt{\frac{i}{2}}x\right) \operatorname{csch}\left(\sqrt{\frac{i}{2}}x\right) = \sum_{p \geq 0} \frac{x^p}{p!} g_1(p)$$

with

$$g_1(p) = (\sqrt{2i})^p \left(\left(B_1 + \frac{1}{2} \right) + i \left(B_2 + \frac{1}{2} \right) \right)^p = (\sqrt{2i})^p B_p^{(2)}\left(\frac{1+i}{2}; 1, i\right)$$

and

$$\frac{1}{\cosh x + \cos x} = \frac{1}{2} \operatorname{sech}\left(\sqrt{\frac{i}{2}}x\right) \sec\left(\sqrt{\frac{i}{2}}x\right) = \sum_{p \geq 0} (-1)^p \frac{x^{2p}}{2p!} g_2(2p)$$

with

$$g_2(2p) = \frac{1}{2} \left(\sqrt{\frac{i}{2}} \right)^{2p} E_{2p}^{(2)}(0; 1, i).$$

Ramanujan's master theorem [1] (see also [8] for a symbolic approach) allows us to compute the integrals I_- with $p = 1$ as

$$I_-(s+2, 1) = \int_0^\infty x^{s-1} \frac{x^2}{\cosh x - \cos x} dx = \Gamma(s) g_1(-s),$$

from which we deduce the result

Proposition 18. *With s a real number at least equal to 3, we have*

$$I_-(s, 1) = \int_0^\infty \frac{x^{s-1}}{\cosh x - \cos x} dx = 2\Gamma(s) \left(\frac{i}{2}\right)^{\frac{s}{2}} \zeta_2(s, 1; 1-i, 1+i).$$

More generally,

$$I_-(s, 1, a, b) := \int_0^\infty \frac{x^{s-1}}{\cosh ax - \cos bx} dx = 2\Gamma(s) \left(\frac{i}{2}\right)^{\frac{s}{2}} \zeta_2(s, a; a-ib, a+ib).$$

Proof. From Lemma 1, we have

$$\frac{x^2}{\cosh x - \cos x} = \sum_{p \geq 0} \frac{x^p}{p!} g_1(p)$$

with

$$\begin{aligned} g_1(p) &= (\sqrt{2i})^p \left(\left(B_1 + \frac{1}{2} \right) + i \left(B_2 + \frac{1}{2} \right) \right)^p \\ &= (\sqrt{2i})^p i^p \left(i \left(B_1 + \frac{1}{2} \right) + \left(B_2 + \frac{1}{2} \right) \right)^p \\ &= \left(\sqrt{\frac{2}{i}} \right)^p i^{2p} \left(i \left(B_1 + \frac{1}{2} \right) + \left(B_2 + \frac{1}{2} \right) \right)^p \\ &= (-1)^p \left(\sqrt{\frac{2}{i}} \right)^p \left(i \left(B_1 + \frac{1}{2} \right) + \left(B_2 + \frac{1}{2} \right) \right)^p \\ &= (-1)^p \left(\sqrt{\frac{2}{i}} \right)^p B_p^{(2)}\left(\frac{1+i}{2}; 1, i\right) \end{aligned}$$

so that

$$\frac{x^2}{\cosh x - \cos x} = \sum_{p \geq 0} (-1)^p \frac{x^p}{p!} g(p)$$

with

$$g(p) = \left(\sqrt{\frac{2}{i}} \right)^p B_p^{(2)} \left(\frac{1+i}{2}; 1, i \right).$$

The analytic continuation of the function g can be found using [4, Eq. (7)]

$$B_{k+2}^{(2)}(x; a_1, a_2) = a_1 a_2 (k+2)(k+1) \zeta_2(-k, x; a_1, a_2)$$

with the Barnes zeta function

$$\zeta_2(s, x; a_1, a_2) = \sum_{m_1, m_2 \geq 0} \frac{1}{(x + a_1 m_1 + a_2 m_2)^s}$$

so that

$$B_{-s}^{(2)}(x; a_1, a_2) = a_1 a_2 s(s+1) \zeta_2(s+2, x; a_1, a_2)$$

and

$$B_{-s}^{(2)}\left(\frac{1+i}{2}; 1, i\right) = is(s+1) \zeta_2\left(s+2, \frac{1+i}{2}; 1, i\right).$$

We therefore deduce

$$\begin{aligned} I_-(s+2, 1) &= \int_0^\infty x^{s-1} \frac{x^2}{\cosh x - \cos x} dx \\ &= \Gamma(s) \left(\sqrt{\frac{i}{2}} \right)^s is(s+1) \zeta_2\left(s+2, \frac{1+i}{2}; 1, i\right) \\ &= 2\Gamma(s+2) \left(\sqrt{\frac{i}{2}} \right)^{s+2} \zeta_2\left(s+2, \frac{1+i}{2}; 1, i\right), \end{aligned}$$

from which the result follows. □

We study now, for an integer $p \geq 1$,

$$I_-(s, p) = \int_0^\infty \frac{x^{s-1}}{(\cosh x - \cos x)^p} dx.$$

Proposition 19. *The integral $I_-(s, p)$ is equal to*

$$I_-(s, p) = 2^p \left(\sqrt{\frac{i}{2}} \right)^s \Gamma(s) \zeta_{2p}\left(s, p \frac{1+i}{2}; (1, i)^p\right) = 2^p \left(\sqrt{\frac{i}{2}} \right)^s \Gamma(s) \sum_{m, n \geq 0} \frac{\binom{p-1+m}{m} \binom{p-1+n}{n}}{\left(p \frac{1+i}{2} + m + in\right)^s}$$

Proof. Since

$$\frac{x^2}{\cosh x - \cos x} = \sum_{n \geq 0} \frac{x^n}{n!} g_1(n)$$

we have

$$f^p(x) = \left(\frac{x^2}{\cosh x - \cos x} \right)^p = \sum_{n \geq 0} \frac{x^n}{n!} g_1^{*p}(n)$$

with the convolution

$$g_1^{*p}(n) = (\sqrt{2i})^n \left(\left(B_1^{(1)} + \frac{1}{2} \right) + i \left(B_2^{(1)} + \frac{1}{2} \right) + \cdots + \left(B_1^{(p)} + \frac{1}{2} \right) + i \left(B_2^{(p)} + \frac{1}{2} \right) \right)^n$$

so that

$$\begin{aligned} g_1^{*p}(n) &= \left(\sqrt{\frac{2}{i}}\right)^n (-1)^n \left(\left(B_1^{(1)} + \frac{1}{2}\right) + i \left(B_2^{(1)} + \frac{1}{2}\right) + \cdots + \left(B_1^{(p)} + \frac{1}{2}\right) + i \left(B_2^{(p)} + \frac{1}{2}\right) \right)^n \\ &= \left(\sqrt{\frac{2}{i}}\right)^n (-1)^n B_n^{(2p)} \left(p \frac{1+i}{2}; (1, i)^p\right) \end{aligned}$$

with the notation $(1, i)^p = (1, i, 1, i, \dots, 1, i)$ so that

$$\left(\frac{x^2}{\cosh x - \cos x}\right)^p = \sum_{n \geq 0} (-1)^n \frac{x^n}{n!} g^{(p)}(n)$$

with

$$g^{(p)}(n) = \left(\sqrt{\frac{2}{i}}\right)^n B_n^{(2p)} \left(p \frac{1+i}{2}; (1, i)^p\right).$$

The Bernoulli-Barnes polynomial can be expressed as the value of the Barnes zeta function at negative index as

$$\zeta_{2p}(-k, x; \mathbf{a}) = (-1)^{2p} \frac{k!}{(2p+k)!} \frac{B_{2p+k}^{(2p)}(x; \mathbf{a})}{i^p}$$

so that

$$\begin{aligned} B_{-s}^{(2p)}(x; \mathbf{a}) &= (-1)^p (-s)(-s-1)\cdots(-s-2p+1) i^p \zeta_{2p}(s+p, x; \mathbf{a}) \\ &= i^p \frac{\Gamma(s+2p)}{\Gamma(s)} \zeta_{2p}(s+p, x; \mathbf{a}). \end{aligned}$$

We deduce

$$g^{(p)}(-s) = \left(\sqrt{\frac{i}{2}}\right)^s B_{-s}^{(2p)} \left(p \frac{1+i}{2}; (1, i)^p\right) = i^p \left(\sqrt{\frac{i}{2}}\right)^s \frac{\Gamma(s+2p)}{\Gamma(s)} \zeta_{2p}\left(s+2p, p \frac{1+i}{2}; (1, i)^p\right)$$

and

$$\begin{aligned} I_-(s+2p, p) &= \int_0^\infty x^{s-1} \frac{x^{2p}}{(\cosh x - \cos x)^p} dx = \Gamma(s) g(-s) \\ &= i^p \left(\sqrt{\frac{i}{2}}\right)^s \Gamma(s+2p) \zeta_{2p}\left(s+2p, p \frac{1+i}{2}; (1, i)^p\right) \end{aligned}$$

and finally

$$\begin{aligned} I_-(s, p) &= i^p \left(\sqrt{\frac{i}{2}}\right)^{s-2p} \Gamma(s) \zeta_{2p}\left(s, p \frac{1+i}{2}; (1, i)^p\right) \\ &= 2^p \left(\sqrt{\frac{i}{2}}\right)^s \Gamma(s) \zeta_{2p}\left(s, p \frac{1+i}{2}; (1, i)^p\right) \end{aligned}$$

Lastly, notice that the $2p$ -variate zeta function

$$\zeta_{2p}\left(s, p \frac{1+i}{2}; (1, i)^p\right) = \sum_{m_1, n_1, \dots, m_p, n_p \geq 0} \left(p \frac{1+i}{2} + m_1 + in_1 + \cdots + m_p + in_p\right)^{-s}$$

is in fact a two-variables Dirichlet series since

$$\zeta_{2p}\left(s, p \frac{1+i}{2}; (1, i)^p\right) = \sum_{m, n \geq 0} \frac{\binom{p-1+m}{m} \binom{p-1+n}{n}}{\left(p \frac{1+i}{2} + m + in\right)^s}$$

as a consequence of the counting function

$$\#\{(n_1, \dots, n_p) \in \mathbb{N}^p : n_1 + \cdots + n_p = n\} = \binom{p-1+n}{n}.$$

□