# ANALOGS OF THE DUAL CANONICAL BASES FOR CLUSTER ALGEBRAS FROM LIE THEORY

FAN QIN

ABSTRACT. For almost all the known (quantum) cluster algebras from Lie theory, we construct the common triangular bases. These bases provide analogs of the dual canonical bases which have long been anticipated in cluster theory. Moreover, when the generalized Cartan matrices are symmetric, these cluster algebras and bases are quasi-categorified.

We employ a unified approach based on the flexibility of cluster algebras, introducing and investigating various cluster operations for this purpose.

We obtain fruitful by products. First, we prove A = U for these quantum cluster algebras. Furthermore, we discover rich structures of the locally compactified quantum cluster algebras arising from double Bott-Samelson cells: T-systems, standard bases, Kazhdan-Lusztig type algorithms, and, in type ADE, monoidal categorifications. The monoidal categories appearing are associated with positive braids and consist of modules of quantum affine algebras. We also obtain monoidal categorifications of the quantized coordinate rings of algebraic groups in type ADE.

#### Contents

1. Introduction	1
Acknowledgments	5
2. Preliminaries	6
3. Freezing operators	11
4. Base changes	16
5. Based cluster algebras and monoidal categorifications	22
6. Cluster algebras associated with signed words	27
7. Applications: cluster algebras from Lie theory	35
8. Applications: cluster algebras from double Bott-Samelson cells	38
Appendix A. Principal coefficients	47
Appendix B. Freezing operators in additive categorifications	48
References	50

### 1. INTRODUCTION

1.1. **Background.** Cluster algebras were invented by Fomin and Zelevinsky [FZ02]. They are algebras with certain combinatorial structures. Particularly, they have distinguished elements called cluster monomials, which are recursively defined by combinatorial algorithms called mutations. They also have natural quantization [BZ05]. Originally, Fomin and Zelevinsky introduced them with the following ambitious expectations:

- (1) The (quantized) coordinate rings of many interesting varieties from Lie theory are isomorphic to appropriate cluster algebras.
- (2) These (quantized) coordinate rings possess distinguished bases, which are analogs of the dual canonical bases [Lus90, Lus91][Kas91] and contain the cluster monomials.

An ever-expanding list of varieties have been found to meet the first expectation. But the progress on the second expectation has been slow. Previously, it was shown that, for the quantized coordinate rings  $\mathcal{O}_q[N(w)]$  of unipotent subgroups N(w) viewed as cluster algebras, the dual canonical bases B<sup>\*</sup> contain the cluster monomials up to scalar multiples, see [Qin17, Qin20b] or [KKKO18][McN21].

We aim to show that the second expectation is met by almost all the known cluster algebras arising from Lie theory. Particularly, we will greatly expand known results from unipotent subgroups to other interesting varieties.

1.2. Main results. Let  $\mathbf{A}$  denote a classical or quantum cluster algebra of a chosen kind:  $\mathbf{A}$  might denote a cluster algebra  $\overline{\mathcal{A}}$ , an upper cluster algebra  $\overline{\mathcal{U}}$ , or their localization  $\mathcal{A}$ ,  $\mathcal{U}$  at the frozen variables. Moreover,  $\mathcal{A}$  and  $\mathcal{U}$  are  $\mathcal{R}$ -algebras over the frozen torus algebra  $\mathcal{R}$ , see Section 2.1. Recall that  $\mathbf{A}$  is constructed from a combinatorial data called a seed  $\mathbf{t}$ , denoted  $\mathbf{A} = \mathbf{A}(\mathbf{t})$ .

Assume that **A** has a basis  $\mathbb{B}$  subject to certain natural conditions, such as containing all cluster monomials (Definition 5.1). The pair (**A**,  $\mathbb{B}$ ) is called a based cluster algebra.

Let  $\mathcal{T}$  denote a monoidal category and K its (deformed) Grothendieck ring. We say  $(\mathbf{A}, \mathbb{B})$  is categorified by  $\mathcal{T}$  if there is an isomorphism  $\kappa : \mathbf{A} \simeq K$ , such that  $\kappa(\mathbb{B})$  consists of the isoclasses of the simple objects of  $\mathcal{T}$ . We say  $(\mathbf{A}, \mathbb{B})$  is quasi-categorified by  $\mathcal{T}$  if it is categorified up to quantization changes, base changes (changing the base  $\mathcal{R}$ ), and localization, see Definition 5.5. Note that, assuming mild conditions, we can keep track of the structures constants of  $\mathbb{B}$  under these changes using the correction technique (Theorem 4.6). Particularly, quasi-categorification implies that the structure constants of  $\mathbb{B}$  are positive.

**Theorem 1.1.** For almost all the quantum upper cluster algebras  $\mathcal{U}$  arising from Lie theory,  $\mathcal{U}$  possess the common triangular bases  $\mathbf{L}$  in the sense of [Qin17]. Moreover,  $(\mathcal{U}, \mathbf{L})$  are quasi-categorified by non-semisimple categories when the generalized Cartan matrices C are symmetric.

Note that **L** contains all cluster monomials by its definition. By [Qin17, Qin20b], any  $\mathcal{O}_q[N(w)]$  has **L** and it coincides with the dual canonical basis B<sup>\*</sup>. Therefore, **L** in Theorem 1.1 provides a natural analog of B<sup>\*</sup>, fulfilling the expectation (2). Theorem 1.1 applies to  $\mathcal{U}(\mathbf{t})$  where **t** belongs to the following families.

- (1) Algebraic group G: Assume C is of finite type. Let G denote the associated connected, simply connected, complex semisimple algebraic group G. We claim that the (quantized) coordinate rings  $\mathbb{C}[G]$  and  $\mathcal{O}_q[G]$  are cluster algebras  $\overline{\mathcal{U}}(\mathbf{t})$  for some  $\mathbf{t}$ , see Claim 7.6. The proofs will appear elsewhere ([FWZ20] treated  $\mathbb{C}[SL_n]$ ).
- (2) Subvarieties of G: Let  $\mathcal{O}_q[N(w)]$ ,  $\mathcal{O}_q[N^w]$ , and  $\mathcal{O}_q[G^{u,v}]$  denote quantum unipotent subgroups, quantum unipotent cells, and quantum double Bruhat cells. They are quantum cluster algebras associated with some seeds  $\mathbf{t}$ , see [GLS11, GLS13][GY16, GY20].

(3) Configurations of flags: The coordinate rings of Grassmannians, open Positroid varieties, open Richardson varieties, double Bott-Samelson cells, and braid varieties are classical cluster algebras for some t, see [Sco06] [SSBW19] [GL19] [SW21] [GLSBS22] [CGG<sup>+</sup>22].

The exotic cluster structures on  $\mathbb{C}[G]$  by [GSV23] and those on double Bott-Samelson cells by [EL21] are not examined in Theorem 1.1, see Remark 8.14.

Statements in Theorem 1.1 were previously known only for  $\mathcal{O}_q[N(w)]$ ,  $\mathcal{O}_q[N^w]$ [Qin17, Qin20b] [KKKO18] [McN21].

Remark 1.2. Family (3) includes cluster algebras arising from various monoidal categories: In type ADE, it includes those arising from representations of quantum affine algebras, see Section 7.6 and [HL10][Qin17]; In type  $A_1^{(1)}$ , it includes those arising from Satake categories, see the last quiver in [CW19, Figure 6.1].

Remark 1.3. In Theorem 1.1, **L** is positive when C is symmetric. In this case,  $(\mathcal{U}, \mathbf{L})$  is always categorified by a semisimple category, see Remark 5.6. We use modules of quiver Hecke algebras [KL09, KL10][Rou08] to provide a non-semisimple category for Theorem 1.1. When C is of type ADE, we can construct an alternative category using modules of quantum affine algebras by Theorem 5.17.

Remark 1.4. For varieties from Lie theory, the quantization of their coordinate rings are often not established in literature. But when the coordinate ring is a cluster algebra, we have natural quantization from cluster theory. And the quantum cluster algebra defined over  $\mathbb{Z}[q^{\pm \frac{1}{2}}]$  can be viewed as the integral form of the quantum cluster algebra defined over  $\mathbb{Q}(q^{\frac{1}{2}})$ . From this perspective, cluster theory provides quantized coordinate rings and their integral forms.

We have the following conjecture, see Example 7.8.

**Conjecture 1.5.** Up to  $q^{\frac{\mathbb{Z}}{2}}$  multiples, the common triangular basis **L** for  $\mathcal{O}_q[G]$  coincides with the global crystal basis of  $\mathcal{O}_q[G]$  in the sense of [Kas93].

We obtain fruitful byproducts in our approach to Theorem 1.1.

First, it is a fundamental yet largely open problem to determine if we have  $\mathcal{A} = \mathcal{U}$  and  $\overline{\mathcal{A}} = \overline{\mathcal{U}}$ . Most known cases concern classical A = U, see [Mul13] [CLS15] [SW21] [MW22] [IOS23]. At the quantum level, [GY20] proved  $\mathcal{A} = \mathcal{U}$  and  $\overline{\mathcal{A}} = \overline{\mathcal{U}}$  for quantum double Bruhat cells. In the following theorem, we make a significant progress on this problem at the quantum level.

**Theorem 1.6.** For the quantum upper cluster algebras in Theorem 1.1, the associated classical and quantum cluster algebras satisfy  $\mathcal{A} = \mathcal{U}$ . Moreover, we have  $\overline{\mathcal{A}}(\dot{\mathbf{t}}) = \overline{\mathcal{U}}(\dot{\mathbf{t}})$ , where  $\dot{\mathbf{t}}$  denote seeds associated with double Bott-Samelson cells as in [SW21] (Section 6).

Moreover, we discover rich structures on  $\overline{\mathcal{U}}(\mathbf{t})$  arising from double Bott-Samelson cells (dBS for short) with the help of the basis **L**. We introduce the interval variables  $W_{[j,k]}$  and the fundamental variables  $W_j$  for  $\overline{\mathcal{U}}(\mathbf{t})$  (Section 8). They are distinguished cluster variables in analogous to the Kirillov-Reshetikhin modules and the fundamental modules respectively.

**Theorem 1.7** (Proposition 8.6 Theorem 8.10).  $\overline{\mathcal{U}}(\dot{\mathbf{t}})$  has the following structures.

(1) The standard basis  $\mathbf{M}$ : it consists of normalized ordered products of fundamental variables.

(2) The T-systems: these are recursive equations relating the interval variables. Moreover, its common triangular basis L equals the Kazhdan-Lusztig basis associated with M.

In the subsequent paper [Qin24b], we will extend  $\overline{\mathcal{U}}(\dot{\mathbf{t}})$  to infinite ranks and compute the fundamental variables  $W_i$  via the braid group action from [JLO23].

**Conjecture 1.8.** If C is symmetric,  $(\overline{\mathcal{U}}(\dot{\mathbf{t}}), \mathbf{L})$  admits a monoidal categorification.

In Section 8.4, we prove Conjecture 1.8 for C of type ADE. For this purpose, we introduce a new monoidal category  $C_{\beta}$  consisting of finite dimensional modules of quantum affine algebras, where  $\beta$  is any positive braid. Then any  $(\overline{\mathcal{U}}(\dot{\mathbf{t}}), \mathbf{L})$  is categorified by  $C_{\beta}$  for the corresponding  $\beta$ .

Finally, assuming the claim that  $\mathcal{O}_q[G]$  is a cluster algebra (see Claim 7.6), we obtain the following result by using a monoidal subcategory of  $\mathcal{C}_{\beta}$ , where **L** denotes the common triangular basis of  $\mathcal{O}_q[G]$ .

**Theorem 1.9** (A special case of Theorem 8.22). In type ADE,  $(\mathcal{O}_q[G] \otimes \mathbb{Q}(q^{\frac{1}{2}}), \mathbf{L})$  admits a monoidal categorification.

1.3. **Strategy.** We employ a unified approach to various algebras in Theorem 1.1 based on the flexibility of cluster algebras. The following observation is crucial:

While varieties from Lie theory could be very different, their coordinate rings might possess closely-related seeds as cluster algebras.

Let  $\mathbf{A}$  denote  $\mathcal{A}$  or  $\mathcal{U}$ . Correspondingly, we introduce and investigate operations that relate cluster algebras  $\mathbf{A}$  associated with closely-related seeds, called cluster operations. Notably, they include the following:

- (1) Freezing  $\mathfrak{f}_F$  (Section 3): Recall that a seed  $\mathbf{t}$  is associated with a set of unfrozen vertices  $I_{uf}$  and a set of frozen vertices  $I_f$ . Choosing any  $F \subset I_{uf}$  and setting it to be frozen, we obtain a new seed  $\mathfrak{f}_F \mathbf{t}$ . We introduce freezing operators  $\mathfrak{f}_F$  sending basis elements of  $\mathbf{A}(\mathbf{t})$  to those of  $\mathbf{A}(\mathfrak{f}_F \mathbf{t})$ .
- (2) **Base change**  $\varphi$  (Section 4.2): Assume two seeds  $\mathbf{t}$ ,  $\mathbf{t}'$  are similar, i.e.,  $I_{uf} = I'_{uf}$  and their *B*-matrices satisfy  $\widetilde{B}_{I_{uf} \times I_{uf}} = \widetilde{B}'_{I_{uf} \times I_{uf}}$ . Under certain conditions, there is a homomorphism var :  $\mathcal{R} \to \mathcal{R}'$ , called a variation map  $[\mathrm{KQW23}][\mathrm{Fra16}]$ , such that we obtain an  $\mathcal{R}'$ -module isomorphism  $\mathcal{R}' \otimes_{\mathcal{R}}$  $\mathbf{A} \stackrel{\simeq}{\simeq} \mathbf{A}'$ , and  $\mathbb{B}'$  can be constructed from  $\mathbb{B}$  via  $\varphi$ .

Despite their elementary definitions, these operations demonstrate remarkable effectiveness and versatility. They propagate localized cluster monomials (Theorem 3.13), bases (Theorems 3.11 and 3.17, Lemmas 4.9 and 4.10, Proposition 4.18), quasi-categorifications or categorifications (Theorems 5.17 and 5.18), and the  $\mathcal{A} = \mathcal{U}$  property (Corollaries 3.16 and 4.12) from **t** to **t**'.

Note that Theorem 1.1 was known for quantum unipotent cells  $\mathcal{O}_q[N^w]$ , which are localization of  $\mathcal{O}_q[N(w)]$ . We hope to verify Theorem 1.1 for the (quantized) coordinate rings  $\Bbbk[\mathcal{V}]$  of other varieties  $\mathcal{V}$  using this known result, but  $\mathcal{V}$  might be more complicated than  $N^w$ . To circumvent this difficulty, we introduce and use the following extension and reduction technique:

Extend the generalized Cartan matrices C to  $\tilde{C}$ . Then study the cluster algebras  $\mathbb{k}[\mathcal{V}]$  by applying cluster operations to quantum unipotent cells  $\mathcal{O}_q[\tilde{N}^{\tilde{w}}]$  associated with  $\tilde{C}$ .

In practice, our approach to  $\mathbb{k}[\mathcal{V}]$  is the following (Sections 6 7):

- (1) Take  $\mathcal{O}_{q}[\widetilde{N}^{\widetilde{w}}]$  associated with an extended generalized Cartan matrix  $\widetilde{C}$ .
- (2) By freezing  $\mathcal{O}_q[\tilde{N}^{\tilde{w}}]$  and applying base changes, obtain  $\mathcal{U}(\dot{\mathbf{t}})$  associated with dBS and C.
- (3) Obtain  $\mathbb{k}[\mathcal{V}]$  from  $\mathcal{U}(\dot{\mathbf{t}})$ , possibly after applying cluster operations.

*Remark* 1.10. The extension and reduction technique is cluster theoretic. It would be interesting to understand it in Lie theory. Note that, when the author was preparing this paper, a paper [BH24] on totally positivity appeared, which also extended the Cartan matrix.

1.4. **Contents.** Section 2 contains basic notions and results for cluster algebras. New notions including dominant tropical points are introduced in Section 2.4.

In Section 3, we introduce and investigate the freezing operators. Base changes of cluster algebras are discussed in Section 4.

In Section 5, we introduced based cluster algebras. Then we discuss their categorifications and the monoidal subcategories associated with the freezing process.

In Section 6, we recall the seeds associated with double Bott-Samelson cells (dBS for short). Then we prove that the associated cluster algebras have common triangular bases and admit quasi-categorification, see Theorems 6.13 and 6.17.

In Section 7, applying the results on dBS in Section 6, we show Theorems 1.1 and 1.6 for almost all the known cluster algebras from Lie theory.

In Section 8, we explore rich structures of the cluster algebras associated with dBS. New monoidal categories  $C_{\beta}$  are introduced to categorify them.

In Section A, we recall principal coefficients seeds. In Section B, we interpret freezing operators in additive categorifications, which is of independent interests.

1.5. **Convention.** Unless otherwise specified, we work with a base ring  $\mathbb{k} = \mathbb{Z}$  for the classical case, and  $\mathbb{k} = \mathbb{Z}[q^{\pm \frac{1}{2}}]$  for the quantum case, where  $q^{\frac{1}{2}}$  is a formal parameter. The elements in  $\mathbb{k}_{\geq 0} := \mathbb{N}[q^{\pm \frac{1}{2}}]$  are called nonnegative. We say two elements  $z_1, z_2$  q-commute, denoted  $z_1 \sim z_2$ , if  $z_1 * z_2 = q^{\alpha} z_2 * z_1$  for some  $\alpha \in \mathbb{Q}$ .

We will use  $\cdot$  to denote the commutative multiplication, which is often omitted, and \* the q-twisted product, which is the default multiplication.

For any  $I \times J$  matrix B and  $I' \subset I$ ,  $J' \subset J$ ,  $B_{I' \times J'}$  denotes the  $I' \times J'$  submatrix of B. We denotes its (i, j)-th entry by  $B_{ij}$ ,  $B_{i,j}$ ,  $b_{ij}$ , or  $b_{i,j}$ . ()<sup>T</sup> denotes the matrix transpose. All vectors are column vectors unless otherwise specified.

Given any skew-symmetric  $\mathbb{Q}$ -matrix  $(b_{ij})_{i,j\in I}$ . We associate with it a quiver: the set of vertices is I, and there is an arrow from  $i \to j$  with weight  $b_{ij}$  if  $b_{ij} > 0$ . A collection of  $w_s$ -weighted arrows is equivalent to a  $\sum w_s$ -weighted arrow.

Given any  $I' \subset I$ , we take the natural decomposition  $\mathbb{Z}^I \simeq \mathbb{Z}^{I'} \oplus \mathbb{Z}^{I \setminus I'}$  and let  $\operatorname{pr}_{I'}$  denote the natural projection from  $\mathbb{Z}^I$  to  $\mathbb{Z}^{I'}$ . Let  $<_{\mathsf{lex}}$  denote the lexicographical order on  $\mathbb{Z}^{[1,l]}$ , and  $<_{\mathsf{rev}}$  the reverse lexicographical order, i.e.,  $(a_1, \ldots, a_l) <_{\mathsf{rev}} (b_1, \ldots, b_l)$  if  $(a_l, \ldots, a_1) <_{\mathsf{lex}} (b_l, \ldots, b_1)$ .

A word  $\underline{\eta} = (\eta_1, \eta_2, ...)$  has letters in  $\mathbb{Z}_{>0}$ , while a signed word  $\underline{\mathbf{i}} = (\mathbf{i}_1, \mathbf{i}_2, ...)$  has letters in  $\mathbb{Z} \setminus \{0\}$ .  $\forall j, k \in \mathbb{Z}, [j, k]$  denotes  $\{i \in \mathbb{Z} | j \leq i \leq k\}$  unless otherwise specified.

 $\mathsf{Span}_{\Bbbk}\mathcal{Z}$  denotes the set of finite  $\Bbbk$ -linear combinations of elements of  $\mathcal{Z}$ .

#### Acknowledgments

The author would like to thank Bernhard Keller, Melissa Sherman-Bennett, Peigen Cao, David Hernandez, Jianrong Li, Jianghua Lu, Linhui Shen, and Milen Yakimov for discussion. Particularly, he is grateful to Bernhard Keller for showing him the inspiring Figure 6.3 in MSRI, 2012.

### 2. Preliminaries

#### 2.1. Basics of cluster algebra.

Seeds. Let I denote a finite set of vertices and we choose its partition  $I = I_{\text{uf}} \sqcup I_{\text{f}}$ into the frozen vertices and unfrozen vertices. |I| is called the rank. Let  $d_i$ ,  $i \in I$ , denote a set of strictly positive integers, called symmetrizers. Let d be their least common multiplier. Denote the Langlands dual  $d_i^{\vee} = \frac{d}{d_i}$ .

Given  $b_{ij} \in \mathbb{Q}$ ,  $i, j \in I$ , such that  $d_i^{\vee} b_{ij} = -d_j^{\vee} b_{ji}$ . Define the *B*-matrix  $\widetilde{B} = (b_{ik})_{i \in I, k \in I_{uf}}$  and assume it is a  $\mathbb{Z}$ -matrix. Define its unfrozen part  $B := \widetilde{B}_{I_{uf} \times I_{uf}}$ .

Assumption 1. We assume that  $\widetilde{B}$  is of full rank unless otherwise specified.

Introduce lattices  $M^{\circ} = \mathbb{Z}^{I}$  and  $N_{\mathsf{uf}} = \mathbb{Z}^{I_{\mathsf{uf}}}$  with unit vectors  $f_i$  and  $e_k$  respectively. Define the linear map  $p^* : N_{\mathsf{uf}} \to M^{\circ}$  such that  $p^*(n) = \widetilde{B}n, \forall n \in N_{\mathsf{uf}}$ .

By [GSV03, GSV05], we can find a  $\mathbb{Z}$ -valued skew-symmetric bilinear form  $\lambda$ on  $M^{\circ}$  such that  $\lambda(f_i, p^*e_k) = \delta_{ik}\delta_k$  for strictly positive integers  $\delta_k$ .  $\lambda$  is called a compatible Poisson structure. Define the quantization matrix  $\Lambda = (\Lambda_{ij})_{i,j \in I} := (\lambda(f_i, f_j))_{i,j}$ . The pair  $(\tilde{B}, \Lambda)$  is called compatible by [BZ05].

**Definition 2.1.** A seed **t** is a collection  $(I, I_{uf}, (d_i)_{i \in I}, \tilde{B}, (x_i)_{i \in I})$ , where  $x_i$  are indeterminates. It is called a quantum seed if it is endowed with a compatible Poisson structure  $\lambda$ .

We shall often omit the symbols  $I, I_{uf}, (d_i)_{i \in I}$  when the context is clear.

Let k denote  $\mathbb{Z}$  or  $\mathbb{Z}[q^{\pm \frac{1}{2}}]$ . Define the Laurent polynomial  $\mathcal{LP} = \mathbb{k}[x_i^{\pm}]_{i \in I}$  using the usual commutative product  $\cdot$ . Denote the Laurent monomial  $x^g = \prod_{i \in I} x_i^{g_i}$ ,  $y_k = x^{p^* e_k}$ , and  $y^n = x^{p^* n}$  for  $g \in \mathbb{Z}^I$ ,  $k \in I_{uf}$ ,  $n \in \mathbb{Z}^{I_{uf}}$ . We further endow  $\mathcal{LP}$  with the following q-twisted product \*:

$$x^g * x^h = q^{\frac{1}{2}\lambda(g,h)} x^{g+h}, \ \forall g,h \in \mathbb{Z}^I.$$

By the multiplication of  $\mathcal{LP}$ , we mean the *q*-twisted product. Its subalgebra  $\overline{\mathcal{LP}} = \mathbb{k}[x_k^{\pm}]_{k \in I_{uf}}[x_j]_{j \in I_f}$  is called its partial compactification.<sup>1</sup> Let  $\mathcal{F}$  denote the skew-field of fractions of  $\mathcal{LP}$ .

Define the frozen monoid  $\overline{\mathcal{P}} := \{x^m | m \in \mathbb{N}^{I_f}\}$ , the frozen group  $\mathcal{P} := \{x^m | m \in \mathbb{Z}^{I_f}\}$ , and the frozen torus algebra  $\mathcal{R} := \Bbbk[x_j^{\pm}]_{j \in I_f}$ ,  $\overline{\mathcal{R}} := \Bbbk[x_j]_{j \in I_f}$ . We also define the bar involution  $\overline{(\ )}$  on  $\mathcal{LP}$  as the  $\mathbb{Z}$ -anti-automorphism such that  $\overline{q^{\alpha} x^g} = q^{-\alpha} x^g$ .

Mutations. Given a seed t. Choose any vertex  $k \in I$  and any sign  $\varepsilon \in \{+, -\}$ . Let  $[]_+$  denote max(, 0). Following [BFZ05], we have the following  $I \times I$  matrix  $\tilde{E}$  and the  $I_{uf} \times I_{uf}$ -matrix F:

$$(\widetilde{E}_{\varepsilon})_{ij} = \begin{cases} \delta_{ij} & k \notin \{i, j\} \\ -1 & i = j = k \\ [-\varepsilon b_{ik}]_+ & j = k, i \neq k \end{cases}, \quad (F_{\varepsilon})_{ij} = \begin{cases} \delta_{ij} & k \notin \{i, j\} \subset I_{uf} \\ -1 & i = j = k \\ [\varepsilon b_{kj}]_+ & i = k, j \neq k \in I_{uf} \end{cases}$$

<sup>&</sup>lt;sup>1</sup>The partial compactification  $\overline{\mathcal{LP}}$  should not be confused with the bar-involution () on  $\mathcal{LP}$ .

The mutation  $\mu_k$  produces a new seed  $\mathbf{t}' = \mu_k \mathbf{t} = (I, I_{uf}, (d_i)_{i \in I}, \tilde{B}', (x'_i)_{i \in I})$ such that  $\tilde{B}' = \tilde{E}_{\varepsilon} \tilde{B} F_{\varepsilon}$ . If  $\mathbf{t}$  is a quantum seed endowed with  $\lambda$ ,  $\mu_k$  also produces a new quantum seed  $\mathbf{t}'$  with a quantization matrix  $\Lambda'$ , such that  $\Lambda' = (\tilde{E}_{\varepsilon})^T \Lambda \tilde{E}_{\varepsilon}$ . Note that  $\mu_k$  is an involution and  $\tilde{B}'$  and  $\lambda'$  do not depend on  $\varepsilon$ , see [BZ05].

We further introduce a k-algebra isomorphism  $\mu_k^* : \mathcal{F}(t') \simeq \mathcal{F}(t)$ :

(2.1) 
$$\mu^* x'_i = \begin{cases} x_k^{-1} \cdot (x^{\sum_j [-b_{jk}]_+ f_j} + x^{\sum_i [b_{ik}]_+ f_i}) & i = k\\ x_i & i \neq k \end{cases}$$

We will often identify  $\mathcal{F}(\mathbf{t}')$  and  $\mathcal{F}(\mathbf{t})$  via  $\mu_k^*$  and omit the symbol  $\mu_k^*$ .

Let there be given an initial seed  $\mathbf{t}_0$ . For any sequence  $\underline{k} = (k_1, \ldots, k_l)$  in  $I_{uf}$ , we use  $\boldsymbol{\mu}_{\underline{k}}$  to denote the sequence of mutations  $\mu_{k_l} \cdots \mu_{k_2} \mu_{k_1}$  (read from right to left) and  $\boldsymbol{\mu}_{\underline{k}}^*$  the composition  $\mu_{k_1}^* \mu_{k_2}^* \cdots \mu_{k_l}^*$ . Denote  $\Delta^+ := \Delta_{\mathbf{t}_0}^+ := \{ \boldsymbol{\mu}_{\underline{k}} \mathbf{t}_0 | \forall \underline{k} \}$ . Take any seed  $\mathbf{t} \in \Delta^+$ . The associated x-variables  $x_i(\mathbf{t})$  are called the (quantum) cluster variables in  $\mathbf{t}$ . We call  $x_j := x_j(\mathbf{t}), j \in I_{\mathbf{f}}$ , the frozen variables. In  $\mathcal{LP}(\mathbf{t})$ , the monomials  $x^m, m \in \mathbb{N}^I$ , are called the cluster monomials in  $\mathbf{t}$ , and  $x^m, m \in$  $\mathbb{N}^{I_{uf}} \oplus \mathbb{Z}^{I_f}$ , the localized cluster monomials in  $\mathbf{t}$ .

**Definition 2.2.** The (partially compactified) cluster algebra  $\overline{\mathcal{A}}(\mathbf{t}_0)$  is the k-subalgebra of  $\mathcal{F}(\mathbf{t}_0)$  generated by the  $x_i(\mathbf{t}), \mathbf{t} \in \Delta^+$ . The (partially compactified) upper cluster algebra is  $\overline{\mathcal{U}}(\mathbf{t}_0) = \bigcap_{\mathbf{t} \in \Delta^+} \overline{\mathcal{LP}}(\mathbf{t})$ . Define the (localized) cluster algebra to be the localization  $\mathcal{A}(\mathbf{t}_0) := \overline{\mathcal{A}}(\mathbf{t}_0)[x_j^{-1}]_{j \in I_{\mathbf{f}}}$  and, similarly, the (localized) upper cluster algebra  $\mathcal{U}(\mathbf{t}_0) = \overline{\mathcal{U}}(\mathbf{t}_0)[x_j^{-1}]_{j \in I_{\mathbf{f}}} = \bigcap_{\mathbf{t} \in \Delta^+} \mathcal{LP}(\mathbf{t})$ .

By [FZ02][BZ05], we have  $\overline{\mathcal{A}}(\mathbf{t}_0) \subset \overline{\mathcal{U}}(\mathbf{t}_0)$  and  $\mathcal{A}(\mathbf{t}_0) \subset \mathcal{U}(\mathbf{t}_0)$ .

2.2. Dominance order, degrees, and support. Take any seed t. We define  $N_{\mathrm{uf}}^{\oplus}(\mathbf{t}) = \bigoplus_{k \in I_{\mathrm{uf}}} \mathbb{N}e_k(\mathbf{t})$  and  $N_{\mathrm{uf}}^+(\mathbf{t}) = N_{\mathrm{uf}}^{\oplus}(\mathbf{t}) \setminus \{0\}.$ 

**Definition 2.3** (Dominance order [Qin17][CILFS15, Proof of Proposition 4.3]).  $\forall g, h \in M^{\circ}(\mathbf{t})$ . We say g dominates h, denoted by  $h \preceq_{\mathbf{t}} g$ , if and only if  $h = g + p^* n$  for some  $n \in N_{\text{uf}}^{\oplus}(\mathbf{t})$ .

Omit the symbol **t** below for simplicity. Recall that we have Laurent monomials  $y^n := x^{p^*n} \in \mathcal{LP}$ . The monoid algebra  $\Bbbk[N_{\mathsf{uf}}^{\oplus}] := \Bbbk[y^n]_{n \in N_{\mathsf{uf}}^{\oplus}(t)}$  has a maximal ideal  $\Bbbk[N_{\mathsf{uf}}^+(t)]$ . Denote the corresponding completion by  $\widehat{\Bbbk[N_{\mathsf{uf}}^+(t)]}$ . We define the rings of formal Laurent series to be

$$\widehat{\mathcal{LP}} := \mathcal{LP} \otimes_{\Bbbk[N_{\mathsf{uf}}^{\oplus}]} \widehat{\Bbbk[N_{\mathsf{uf}}^{\oplus}]}, \ \widehat{\Bbbk[N_{\mathsf{uf}}]} := \Bbbk[N_{\mathsf{uf}}] \otimes_{\Bbbk[N_{\mathsf{uf}}^{\oplus}]} \widehat{\Bbbk[N_{\mathsf{uf}}^{\oplus}]}.$$

For  $m \in M$ , denote  $\widehat{\mathcal{LP}}_{\preceq m} := x^m \cdot \widehat{\Bbbk[N_{\mathsf{uf}}^{\oplus}]} \subset \widehat{\mathcal{LP}}$  and  $\mathcal{LP}_{\preceq m} := x^m \cdot \Bbbk[N_{\mathsf{uf}}^{\oplus}] \subset \mathcal{LP}$ .

**Definition 2.4.** Consider any formal sum  $z = \sum c_m x^m$ ,  $c_m \in k$ .

(1) If the set  $\{m | c_m \neq 0\}$  has a unique  $\prec_{\mathbf{t}}$ -maximal element g, we say z has degree g and denote  $\deg^{\mathbf{t}} z = g$ . We further say that z is g-pointed (or pointed at g) if  $c_g = 1$ . In this case,  $\forall \alpha \in \frac{1}{2}\mathbb{Z}$ , define the normalization of  $q^{\alpha}z$  to be  $[q^{\alpha}z]^{\mathbf{t}} := z$ .

(2) If the set  $\{m|c_m \neq 0\}$  has a unique  $\prec_t$ -minimal element h, we say z has codegree h and denote  $\operatorname{codeg}^t z = h$ . We further say that z is h-copointed (or copointed at h) if  $c_h = 1$ .

For example, the localized cluster monomials are pointed at distinct degrees and copointed at distinct codegrees in  $\mathcal{LP}(\mathbf{t})$ , see [FZ07][DWZ10][Tra11][GHKK18].

**Definition 2.5** (Support).  $\forall n = \sum n_k e_k \in N_{uf}$ , we define its support  $\text{supp} n := \{k \in I_{uf} | n_k \neq 0\}$ . For any  $z = x^g \cdot \sum_{n \in N_{uf}^{\oplus}} b_n y^n$  in  $\widehat{\mathcal{LP}}$  with  $b_n \in \Bbbk$ ,  $b_0 \neq 0$ . We define its support  $\text{supp} z := \bigcup_{n:b_n \neq 0} \text{supp} n$ . If further  $z \in \mathcal{LP}$ , we define its support dimension  $\text{suppDim} z \in N_{uf}^{\oplus}$  such that  $(\text{suppDim} z)_k = \max\{n_k | b_n \neq 0\}, \forall k \in I_{uf}$ .

Following [Qin17], we say  $\mathbf{t}_0$  injective-reachable if there exists a seed  $\mathbf{t}_0[1] = \Sigma \mathbf{t}_0$  for some mutation sequence  $\Sigma$  and some permutation  $\sigma$  of  $I_{uf}$ , such that  $\mathsf{deg}^{\mathbf{t}_0} x_{\sigma k}(\mathbf{t}_0[1]) \in -f_k + \mathbb{Z}^{I_f}$ , for any  $k \in I_{uf}$ . In this case,  $\Sigma$  is called a green to red sequence following [Kell1]. Note that, if  $\mathbf{t}_0$  is injective-reachable, then so are all seeds  $\mathbf{t} \in \Delta^+$  [Qin17][Mul16].

**Definition 2.6** ([Qin17]). Let  $z_s, s \in \mathbb{N}$ , denote a collection of pointed elements in  $\widehat{\mathcal{LP}}$ . A (possibly infinite) decomposition in  $\widehat{\mathcal{LP}}$  is called  $\prec_{\mathbf{t}}$ -unitriangular if it takes the form  $z = z_0 + \sum_{\deg z_s \prec_{\mathbf{t}} \deg z_0} b_s z_s, b_s \in \mathbb{k}$ . It is further called  $(\prec_{\mathbf{t}}, \mathbf{m})$ unitriangular, where  $\mathbf{m} := q^{-\frac{1}{2}} \mathbb{Z}[q^{-\frac{1}{2}}]$ , if  $b_s \in \mathbf{m}$ .

**Lemma 2.7.** Take any  $\Theta \subset M^{\circ}(\mathbf{t})$ . Let  $\mathbb{B} = \{b_m | m \in \Theta\}$  denote a subset of  $\mathcal{LP}(\mathbf{t})$ , where  $b_m$  are m-pointed. Denote  $V = \operatorname{Span}_{\Bbbk} \mathbb{B}$ . Let  $Z = \{z_m | m \in \Theta\}$  denote any subset of V where  $z_m$  are m-pointed. Then Z is also a k-basis of V if  $\prec_{\mathbf{t}}$  is bounded from below on  $\Theta$ .

Proof. We have a decomposition  $z_m = b_m + \sum_{m' \prec_t m \in \Theta} c_{m'} b_{m'}, c_{m'} \in \mathbb{k}, \forall m \in \Theta$ . Take the inverse transition  $b_m = z_m + \sum_{m' \prec_t m \in \Theta} c'_{m'} z_{m'}, c'_{m'} \in \mathbb{k}$ , see [Qin17]. This is a finite sum by the boundedness of  $\prec_t$  on  $\Theta$ . The desired claim follows.

2.3. Tropical transformations. Take seeds  $\mathbf{t}' = \mu_k \mathbf{t}$  for some  $k \in I_{uf}$ . The tropical transformation  $\phi_{\mathbf{t}',\mathbf{t}} : M^{\circ}(\mathbf{t}) \to M^{\circ}(\mathbf{t}')$  is piecewise linear such that, for any  $m = \sum m_i f_i(\mathbf{t}) \in \oplus \mathbb{Z} f_i(\mathbf{t}) = M^{\circ}(\mathbf{t})$ , its image  $m' = \sum m'_i f_i(\mathbf{t}') \in M^{\circ}(\mathbf{t}')$  is given by  $m'_i = \begin{cases} -m_k & i = k \\ m_i + b_{ik}(t)[m_k]_+ & b_{ik}(t) \ge 0. \\ m_i + b_{ik}(t)[-m_k]_+ & b_{ik}(t) \le 0 \end{cases}$ 

For any seeds  $\mathbf{t}' = \boldsymbol{\mu}_{\underline{k}} \mathbf{t}$ , we define  $\phi_{\mathbf{t}',\mathbf{t}}$  to be the composition of the tropical transformations along  $\boldsymbol{\mu}_{\underline{k}}$ . It is independent of the choice of  $\boldsymbol{\mu}_{\underline{k}}$ , see [GHK15].

**Definition 2.8** ([Qin24c]).  $\forall m \in M^{\circ}(\mathbf{t})$ , the tropical point represented by m is defined to be the equivalence class [m] in  $\sqcup_{\mathbf{t}'\in\Delta^+} M^{\circ}(\mathbf{t}')$  under the identifications  $\phi_{\mathbf{t}',\mathbf{t}}, \forall \mathbf{t}'$ . Define the set of tropical points  $\mathcal{M}^{\circ}$  to be  $\{[m]|\forall m \in M^{\circ}(\mathbf{t})\}$ .

**Definition 2.9** ([Qin24c]). Given an *m*-pointed element  $z \in \mathcal{LP}(\mathbf{t})$ . It is said to be compatibly pointed at  $\mathbf{t}, \mathbf{t}'$  if z is  $\phi_{t',t}m$ -pointed in  $\mathcal{LP}(t')$ . We say z is [m] pointed if it is compatibly pointed at  $\mathbf{t}, \mathbf{t}'$ , for all  $\mathbf{t}' \in \Delta^+$ .

For example, by [DWZ10][Tra11][GHKK18], the localized cluster monomials are pointed at distinct tropical points.

**Definition 2.10.** Let  $\Theta$  denote a subset of  $M^{\circ}(\mathbf{t})$  or  $\mathcal{M}^{\circ}$  and  $\mathcal{S} = \{s_g | g \in \Theta\}$  be a subset of  $\widehat{\mathcal{LP}}(\mathbf{t})$ . If  $s_g$  are g-pointed,  $\mathcal{S}$  is said to be  $\Theta$ -pointed.

**Lemma 2.11** ([Qin24c, Lemma 3.4.12]). Assume that t is injective-reachable. Take any  $u \in \mathcal{U}(t)$ . If there exists a localized cluster monomial z such that u is [deg z]pointed, then u = z. 2.4. **Partial compactification and optimized seeds.** Take any seed **t** without assuming  $\widetilde{B}(\mathbf{t})$  to be of full rank. For any frozen vertex j, let  $\nu_j$  denote the order of vanishing at  $x_j = 0$  on  $\mathcal{F}(\mathbf{t})$ , i.e.,  $\forall Z \in \mathcal{F}(t)$ , we have the reduced form  $Z = x_j^{\nu_j(Z)} * P * Q^{-1}$ , where  $P, Q \in \mathbb{k}[x_i]_{i \in I}$  are not divisible by  $x_j$ . Note that  $\nu_j$  is a valuation. We say  $Z \in \mathcal{F}(t)$  is regular at  $x_j = 0$  if  $\nu_j(Z) \ge 0$ .

**Lemma 2.12.** Take any  $\mathbf{t}' = \mu_k \mathbf{t}$ ,  $k \in I_{uf}$ . Let  $\nu_j$  and  $\nu'_j$  denote the orders of vanishing on  $\mathcal{F}(\mathbf{t})$  and  $\mathcal{F}(\mathbf{t}')$  respectively. Then for any  $Z \in \mathcal{F}(\mathbf{t}')$ , we have  $\nu'_j(Z) = \nu_j(\mu_k^*Z)$ .

Proof. Take any polynomial  $P' \in \mathbb{k}[x_i(\mathbf{t}')]_{i\in I}$  not divisible by  $x_j$ . View it as a polynomial in  $x_j$  and  $x_k$ , we can write  $P' = \sum_{r,s\geq 0} c_{r,s} * (x'_k)^r * x^s_j$  with  $0 \neq \sum_r c_{r,0} * (x'_k)^r$ . Note that  $\mu_k^* c_{r,s} = c_{r,s}$  and  $\mu_k^* x'_k|_{x_j\mapsto 0} = M_r * x_k^{-r} \in \mathbb{k}[x_i]_{i\in I}[x_k^{-1}]$  for  $M_r$  a monomial in  $x_i, i \neq j, k$ . We deduce that  $\mu_k^* P'|_{x_j\mapsto 0} = \sum_r c_{r,s} * M_r * x_k^{-r} \neq 0$ . Therefore,  $\mu_k^* P'$  is not divisible by  $x_j$ . The desired statement follows by applying this result to the reduced form of Z.

Observe that  $\overline{\mathcal{U}} = \{ z \in \mathcal{U} | \nu_j(z) \ge 0, \forall j \in I_f \}.$ 

**Lemma 2.13.** For any m-pointed  $z \in \mathcal{U}(\mathbf{t}) \subset \mathcal{LP}(\mathbf{t})$ , if some  $k \in I_{uf}$  is not contained in its support (Definition 2.5), we have  $\nu_k(z) \geq 0$ .

*Proof.* We can assume that **t** is the seed  $\mathbf{t}^{\text{prin}}$  of principal coefficients (Section A). Otherwise, we replace the pointed element **t** by a similar element z' in  $\mathcal{LP}(\mathbf{t}^{\text{prin}})$  as in Definition 4.4, for which we will have supp z = supp z',  $\nu_k(z) = \nu_k(z')$ , and  $z' \in \mathcal{U}(\mathbf{t}^{\text{prin}})$  by Lemma 4.7.

When  $\mathbf{t} = \mathbf{t}^{\text{prin}}$ , one can prove the claim from the condition  $z \in \mathcal{LP}(\mathbf{t}) \cap \mathcal{LP}(\mu_k \mathbf{t})$ . Let us recall the following proof from [CKQ24].

Denote  $I_{uf} := I_{uf}(\mathbf{t})$ . Since  $\mathbf{t} = \mathbf{t}^{\text{prin}}$ , we have  $I_f := I_f(\mathbf{t}) = \{k'|k \in I_{uf}\}$ . Denote  $U_{\widehat{k}} := \mathbb{k}[x_i^{\pm}, x_i^{\pm}]_{i \in I_{uf}, i \neq k}$  and  $V_{\widehat{k}} := U_{\widehat{k}}[x_{k'}^{\pm}]$ . Since  $z \in \mathcal{LP}(\mathbf{t})$ , we have  $z = \sum_{s \geq 0} L_s * x_k^s + \sum_{s \geq 0} x_k^{-s} * L_{-s}$  in  $\mathcal{LP}(\mathbf{t}) = V_{\widehat{k}}[x_k^{\pm}]$ , where  $L_{\pm s} \in V_{\widehat{k}}$ .

Denote  $\mathbf{t}' = \mu_k \mathbf{t}$  and  $x_k = x_k(\mathbf{t}')$ . When  $s \in \mathbb{N}$ , the Laurent expansion of  $x_k^s$  in  $\mathcal{LP}(\mathbf{t}')$  takes the form  $x_k^s = \widetilde{P}_s * (x_k')^{-s}$ , where  $\widetilde{P} \in U_{\widehat{k}}[x_{k'}]$  is a degree s polynomial in  $x_{k'}$  with non-vanishing constant term.

Then we obtain  $z = \sum_{s\geq 0}^{\circ} L_s * \widetilde{P}_s * (x'_k)^{-s} + \sum_{s>0}^{\circ} (x'_k)^s * \widetilde{P}_s^{-1} * L_{-s}$  in  $\mathcal{LP}(\mathbf{t}') = V_{\widehat{k}}[(x'_k)^{\pm}]$ . So we must have  $\widetilde{P}_s^{-1} * L_{-s} \in V_{\widehat{k}} = U_{\widehat{k}}[x_{k'}^{\pm}]$ . Denote  $Q_s = \widetilde{P}_s^{-1} * L_{-s} \in U_{\widehat{k}}[x_{k'}^{\pm}]$ , we have  $\widetilde{P}_s * Q_s = L_{-s} \in U_{\widehat{k}}$ . Recall that  $\widetilde{P}_s$  is a polynomial in  $x_{k'}$  of degree s with non-vanishing constant term. By analyzing the degree of  $x_{k'}$ , we deduce that  $Q_s = L_{-s} = 0, \forall s > 0$ . Equivalently,  $\nu_k(z) \geq 0$ .

Let j denote any frozen vertex of a seed  $\mathbf{t}$ . We say j is optimized in  $\mathbf{t}$  if  $b_{jk}(\mathbf{t}) \geq 0$ for any  $k \in I_{uf}$ , or, equivalently,  $\nu_j(y_k(\mathbf{t}_j)) \geq 0$ . We say j can be optimized if it is optimized in some seed  $\mathbf{t}' \in \Delta_{\mathbf{t}}^+$ , and  $\mathbf{t}$  can be optimized if its frozen vertices can be. Assume  $\widetilde{B}(\mathbf{t})$  is of full rank from now on.

**Lemma 2.14.** Take any  $j \in I_f$  with an optimized seed  $\mathbf{t}_j$ . If  $z \in \mathcal{LP}(\mathbf{t}_j)$  has degree m, then  $\nu_j(z) = \nu_j(m)$ .

The following result is analogous to [GHKK18, Proposition 9.7].

**Proposition 2.15.** Let S denote any basis of U such that it is  $\mathcal{M}^{\circ}$ -pointed. Assume that each  $j \in I_{f}$  has an optimized seed  $\mathbf{t}_{j}$ . Then  $S \cap \overline{U}$  is a basis of  $\overline{U}$ .

*Proof.* For any element  $Z \in \overline{\mathcal{U}}$ , we have a finite sum  $Z = \sum_{i=1}^{r} c_i S_i$  for basis elements  $S_i \in \mathcal{S}$  and  $0 \neq c_i \in \mathbb{k}$ . It suffices to show that, for any  $j \in I_{\mathsf{f}}, \nu_j(S_i) \geq 0$ .

Working inside  $\mathcal{LP}(\mathbf{t}_j)$  and denote  $m_i = \deg^{\mathbf{t}_j} S_i$ . Let  $m_k$  denote a  $\prec_{\mathbf{t}_j}$ -maximal element of  $\{m_i | i \in [1, r]\}$ . Then  $x^{m_k}$  must appear in the Laurent expansion of Z in  $\mathcal{LP}(\mathbf{t}_j)$ . So it is regular at  $x_j = 0$ . By Lemma 2.14,  $\nu_j(S_k) \ge 0$ . Replace Z by  $Z - S_k$  and consider  $Z - S_k = \sum_{i \neq k} c_i S_i$ . Repeating this process, we obtain that all  $S_i$  appearing are regular at  $x_j = 0$ , i.e.,  $\nu_j(S_i) \ge 0$ .

**Definition 2.16.** Choose any subset  $\mathcal{Z} = \{z_{[m]} | [m] \in \mathcal{M}^{\circ}\}$  of  $\mathcal{U}$  such that  $z_{[m]}$  are [m]-pointed. We define the associated set of dominant tropical points to be  $\overline{\mathcal{M}}^{\circ} := \{[m] \in \mathcal{M}^{\circ} | z_{[m]} \in \overline{\mathcal{U}}\}$ . Denote  $\overline{\mathcal{M}}^{\circ}(\mathbf{t}) := \{m \in \mathcal{M}^{\circ}(\mathbf{t}) | [m] \in \overline{\mathcal{M}}^{\circ}\}$ .

By Lemma 2.14,  $\overline{\mathcal{M}}^{\circ}$  can be calculated as below.

**Lemma 2.17.** If all  $j \in I_f$  have optimized seeds  $\mathbf{t}_j$ ,  $\overline{M}^{\circ}(\mathbf{t})$  equals  $\{m | (\phi_{\mathbf{t}_j, \mathbf{t}} m)_j \geq 0, \forall j \in I_f\}$ .

In general, we will choose  $\mathcal{Z}$  to be the set of theta functions [GHKK18], because their existence has been verified in vast generalities.

*Remark* 2.18. Z can be taken as the theta basis, the generic basis, or triangular basis, when they exist. We conjecture that the resulting  $\overline{\mathcal{M}}^{\circ}$  are equivalent for many (if not all) cluster algebras.

**Definition 2.19.** A frozen vertex j is called non-essential if  $\widetilde{B}_{\{j\}\times I_{uf}} = 0$ , and essential otherwise.

If j is non-essential, let t' denote the (quantum) seed naturally obtained from t by removing j. Then we have  $\overline{\mathcal{U}} = \Bbbk[x_j]\overline{\mathcal{U}}$  and  $\overline{\mathcal{A}} = \Bbbk[x_j]\overline{\mathcal{A}}$ .

2.5. Triangular bases. Take  $\mathbb{k} = \mathbb{Z}[q^{\pm \frac{1}{2}}]$ . Let A denote  $\mathcal{A}$  or  $\mathcal{U}$ . Assume t is injective-reachable.

**Definition 2.20** ([Qin17]). Let  $\mathbf{L} = {\mathbf{L}_m | m \in M^\circ(\mathbf{t})}$  be a k-basis of  $\mathbf{A}(\mathbf{t})$ , such that  $\mathbf{L}_m$  are *m*-pointed.  $\mathbf{L}$  is called the triangular basis with respect to the seed  $\mathbf{t}$  if we have the following:

- $\overline{\mathbf{L}_m} = \mathbf{L}_m$ .
- **L** contains the cluster monomials in  $\mathbf{t}$  and  $\mathbf{t}[1]$ .
- $\forall x_i(\mathbf{t}), \mathbf{L}_m$ , the decomposition of the normalization  $[x_i(\mathbf{t}) * \mathbf{L}_m]^{\mathbf{t}}$  is  $(\prec_t, \mathbf{m})$ unitriangular:

(2.2) 
$$[x_i(\mathbf{t}) * \mathbf{L}_g]^{\mathbf{t}} = \mathbf{L}_{g+f_i} + \sum_{g' \prec_{t_0} g+f_i} b_{g'} \mathbf{L}_{g'}, \text{ for some } b_{g'} \in \mathbf{m} := q^{-\frac{1}{2}} \mathbb{Z}[q^{-\frac{1}{2}}].$$

The decomposition on RHS of (2.2) is related to Leclerc's conjecture [Lec03] [Qin20a]. Note that  $[x_j^{\pm} * \mathbf{L}_m]^{\mathbf{t}} = \mathbf{L}_{m \pm f_j}, \forall j \in I_{\mathbf{f}}$ . The triangular basis with respect to **t** is determined by a Kazhdan-Lusztig type algorithm, see [Qin20b, Section 6.1]. In particular, it is unique if it exists.

**Definition 2.21** ([Qin21, Definition 5.5, Remark 5.6][Qin17]). A k-basis L of A is called the common triangular basis if it is the triangular basis with respect to any seed  $\mathbf{t} \in \Delta^+$ .

By [Qin20b, Proposition 6.4.3], if **L** is the triangular basis with respect to the seed **t**, then it is the common triangular basis if and only if it contains the cluster monomials of any seeds  $\mathbf{t}' \in \Delta^+$ . In this case, the basis elements  $\mathbf{L}_m$  are [m]-pointed by [Qin20b, Proposition 6.4.3]. Then [Qin24c, Theorem 4.3.1] implies that **L** is a basis for  $\mathcal{U}$  and, consequently,  $\mathbf{A} = \mathcal{U}$ .

Finally, if **L** is the common triangular basis (resp. the triangular basis with respect to **t**) of **A** such that  $\mathbf{L} \cap \overline{\mathbf{A}}$  is a basis of  $\overline{\mathbf{A}}$ , we call  $\mathbf{L} \cap \overline{\mathbf{A}}$  the common triangular basis (resp. the triangular basis with respect to **t**) of  $\overline{\mathbf{A}}$ .

#### 3. Freezing operators

3.1. Sub seeds. Take two seeds  $\mathbf{t} = (I, I_{uf}, (d_i), \widetilde{B}, (x_i)_{i \in I}), \mathbf{t}' = (I', I'_{uf}, (d'_i), \widetilde{B}', (x'_i)_{i \in I'}).$ 

**Definition 3.1.** A cluster embedding  $\iota$  from **t** to **t**' is an inclusion  $\iota: I' \hookrightarrow I$  such that  $\iota I'_{\mathsf{uf}} \subset I_{\mathsf{uf}}, d'_i = d_{\iota(i)}, \text{ and } b'_{i,k} = b_{\iota(i),\iota(k)}, \forall i \in I', k \in I'_{\mathsf{uf}}.$  If **t** and **t**' are quantum seeds, we further require  $\Lambda(\mathbf{t}')_{i,j} = \Lambda(\mathbf{t})_{\iota(i),\iota(j)}, \forall i, j \in I'.$ 

Assume a cluster embedding  $\iota : I' \hookrightarrow I$  is given.  $\forall$  mutation sequence  $\boldsymbol{\mu} = \mu_{k_r} \cdots \mu_{k_1}$  on  $I'_{uf}$ , let  $\iota(\boldsymbol{\mu})$  denote  $\mu_{\iota(k_r)} \cdots \mu_{\iota(k_1)}$ . By the mutation rules,  $\iota$  is also a cluster embedding from  $\boldsymbol{\mu}\mathbf{t}'$  to  $\iota(\boldsymbol{\mu})\mathbf{t}$ .

We obtain the inclusion  $\iota : \mathcal{F}(\mathbf{t}') \subset \mathcal{F}(\mathbf{t})$  sending  $x'_i$  to  $x_{\iota(i)}$ . In this case, we say  $\mathbf{t}'$  is a sub seed of  $\mathbf{t}$  via  $\iota$ , denoted  $\iota : \mathbf{t}' \subset \mathbf{t}$ . If we further have  $b_{i,k} = 0, \forall k \in \iota(I'_{uf})$  and  $i \in I \setminus \iota(I'), \mathbf{t}'$  is called a good sub seed of  $\mathbf{t}$ . We often omit  $\iota$  for simplicity.

By the mutation rules, we have the following results.

**Proposition 3.2.** Assume that  $\mathbf{t}'$  is a good sub seed of  $\mathbf{t}$  via  $\iota$ . Then  $\boldsymbol{\mu}\mathbf{t}'$  is a good sub seed of  $\iota(\boldsymbol{\mu})\mathbf{t}$  via  $\iota$ ,  $\iota(x_i(\boldsymbol{\mu}\mathbf{t}')) = x_i(\boldsymbol{\mu}\mathbf{t})$ , and  $\iota(\overline{\mathcal{LP}}(\boldsymbol{\mu}\mathbf{t}')) \subset \overline{\mathcal{LP}}(\iota(\boldsymbol{\mu})\mathbf{t})$  for any  $i \in I'$  and any mutation sequence  $\boldsymbol{\mu}$  on  $I'_{uf}$ . Particularly, we have  $\iota(\overline{\mathcal{A}}(\mathbf{t}')) \subset \overline{\mathcal{A}}(\mathbf{t})$ .

**Proposition 3.3.** When  $\mathbf{t}'$  is a good sub seed of  $\mathbf{t}$  via  $\iota$ , we have  $\overline{\mathcal{U}}' \subset \overline{\mathcal{U}}$  under the assumption that  $\mathbf{t}$  is of full rank.

Proof. We omit the symbol  $\iota$  for simplicity. Since  $\mathbf{t}$  is of full rank, we have  $\mathcal{U} = \mathcal{LP} \cap (\bigcap_{k \in I_{uf}} \mathcal{LP}(\mu_k \mathbf{t}))$  by [BFZ05, Corollary 1.9][BZ05, Theorem 5.1]. It follows that  $\overline{\mathcal{U}} = \overline{\mathcal{LP}} \cap (\bigcap_{k \in I_{uf}} \overline{\mathcal{LP}}(\mu_k \mathbf{t}))$ . Take any  $z \in \overline{\mathcal{U}}'$ . It belongs to  $\overline{\mathcal{LP}}' \cap (\bigcap_{k' \in I'_{uf}} \overline{\mathcal{LP}}(\mu_{k'} \mathbf{t}')) \subset \overline{\mathcal{LP}} \cap (\bigcap_{k' \in I'_{uf}} \overline{\mathcal{LP}}(\mu_{k'} \mathbf{t}))$ . Moreover,  $\forall k \in I_{uf} \setminus I'_{uf}, z \in \overline{\mathcal{LP}}' \subset \overline{\mathcal{LP}}$  is also contained in  $\overline{\mathcal{LP}}(\mu_k \mathbf{t})$  since  $\nu_k(z) \geq 0$ . Therefore,  $z \in \overline{\mathcal{U}}$ .

3.2. Freezing. Choose any subset of unfrozen vertices  $F \subset I_{uf}$ . By freezing F, we obtain a good sub seed of  $\mathbf{t}$ , denoted  $\mathfrak{f}_F \mathbf{t} := (I, I_{uf} \setminus F, (d_i), \widetilde{B}_{I \times (I_{uf} \setminus F)}, (x_i)_{i \in I})$ . If  $\mathbf{t}$  is a quantum seed,  $\mathfrak{f}_F \mathbf{t}$  is also quantized where we choose  $\Lambda(\mathfrak{f}_F \mathbf{t}) := \Lambda(\mathbf{t})$ .

Remark 3.4. Note that  $M^{\circ}(\mathfrak{f}_{F}\mathbf{t}) = M^{\circ}(\mathbf{t}), m' \preceq_{\mathfrak{f}_{F}\mathbf{t}} m$  implies  $m' \preceq_{\mathbf{t}} m, x_{i}(\mathfrak{f}_{F}\mathbf{t}) = x_{i}(\mathbf{t}), \text{ and } \mathcal{F}(\mathfrak{f}_{F}\mathbf{t}) = \mathcal{F}(\mathbf{t})$ . The identification  $M^{\circ}(\mathfrak{f}_{F}\mathbf{t}) = M^{\circ}(\mathbf{t})$  induces a natural identification between the set of tropical points for  $\mathcal{U}(\mathfrak{f}_{F}\mathbf{t})$  and  $\mathcal{U}(\mathbf{t})$ , which we both denote by  $\mathcal{M}^{\circ}$ .

**Definition 3.5** (Freezing operator). For any degree  $m \in M^{\circ}(\mathbf{t})$  and  $F \subset I_{uf}$ , we define the freezing operator  $\mathfrak{f}_{F,m}^{\mathbf{t}}$  to be the k-linear map from  $\widehat{\mathcal{LP}}_{\preceq m}(\mathbf{t})$  to  $\widehat{\mathcal{LP}}_{\prec m}(\mathfrak{f}_F \mathbf{t})$  such that

$$\mathbf{f}^{\mathbf{t}}_{F,m}(\sum_{n\in N_{\mathrm{uf}}^{\oplus}}c_nx^m\cdot y^n)=\sum_{n:\mathrm{supp}\,n\cap F=\emptyset}c_nx^m\cdot y^n,\;\forall c_n\in \Bbbk.$$

In general, by freezing  $F' \subset I$ , we mean freezing  $F := F' \cap I_{\text{uf}}$ , i.e.,  $\mathfrak{f}_{F'} \mathbf{t} := \mathfrak{f}_F \mathbf{t}$ ,  $\mathfrak{f}_{F',m}^{\mathbf{t}} := \mathfrak{f}_{F,m}^{\mathbf{t}}$ . When the context is clear, we omit the scripts  $\mathbf{t}$ , F and/or m for simplicity. In particular, we often denote  $\mathfrak{f}_F^{\mathbf{t}}(z) = \mathfrak{f}_{F,m}^{\mathbf{t}}(z)$  when z has a unique  $\prec_{\mathbf{t}}$ -maximal degree m.

Lemma 3.6. (1)  $\forall m' \preceq_{\mathbf{t}} m \text{ such that } m' = m + \widetilde{B}n \text{ and } z \in \widehat{\mathcal{LP}}_{\preceq m'}(\mathbf{t}) \subset \widehat{\mathcal{LP}}_{\preceq m}(\mathbf{t}),$ we have  $\mathfrak{f}_m(z) = \begin{cases} \mathfrak{f}_{m'}(z) & \operatorname{supp} n \cap F = \emptyset \\ 0 & \operatorname{supp} n \cap F \neq \emptyset \end{cases}$ (2)  $\forall m_1, m_2 \text{ and } z_1 \in \widehat{\mathcal{LP}}_{\preceq m_1}(\mathbf{t}), z_2 \in \widehat{\mathcal{LP}}_{\preceq m_2}(\mathbf{t}), we have$  $\mathfrak{f}_{m_1}(z_1) * \mathfrak{f}_{m_2}(z_2) = \mathfrak{f}_{m_1+m_2}(z_1 * z_2).$ 

*Proof.* The claims follow from definition of the freezing operators.

**Example 3.7.** Let **t** denote a seed such that  $I = \{1, 2\}, I_{uf} = \{1\}, \widetilde{B} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ . The upper cluster algebra  $\mathcal{U}(t)$  equals  $\mathbb{k}[x_2^{\pm}][x_1, x_1']$  where  $x_1' := x_1^{-1} \cdot (1+y_1)$ . Its localized cluster monomials are  $s_m = x_1^{m_1} \cdot x_2^{m_2}$  when  $m_1 \ge 0$ , and  $s_m = (x_1')^{-m_1} \cdot x_2^{m_2}$  when  $m_1 < 0$ . Choose  $F = \{1\}$  and construct the seed  $\mathfrak{f}_F \mathfrak{t}$  by freezing F. We have  $\mathcal{U}(\mathfrak{f}_F t) = \mathbb{k}[x_1^{\pm}, x_2^{\pm}]$ . Its localized cluster monomials  $s'_m := x^m$  satisfy  $s'_m = \mathfrak{f}_{F,m} s_m$ .

**Proposition 3.8.** Take any  $z \in \mathcal{LP}_{\preceq m}(\mathbf{t})$ . If  $z \in \mathcal{U}(\mathbf{t})$ , then  $\mathfrak{f}_m z \in \mathcal{U}(\mathfrak{f}_F \mathbf{t})$ .

Proof. Take any sequence  $\underline{k}$  on  $I_{uf} \setminus F$  and denote  $\mathbf{t}' = \boldsymbol{\mu}_{\underline{k}} \mathbf{t}$ . Since  $z \in \mathcal{LP}(\mathbf{t}')$ , we have  $x(\mathbf{t}')^{d'} * z = \sum_{g'} b_{g'} x(\mathbf{t}')^{g'}$  for some  $d', g' \in \mathbb{N}^I$ ,  $b_{g'} \in \mathbb{k}$ . Now work in  $\mathcal{LP}(\mathbf{t})$  and denote  $d = \mathsf{deg}^{\mathbf{t}} x(\mathbf{t}')^{d'}$ ,  $g = \mathsf{deg}^{\mathbf{t}} x(\mathbf{t}')^{g'}$ . Since  $z \in \mathcal{LP}_{\preceq m}(\mathbf{t})$ , we have  $g \preceq_{\mathbf{t}} d + m$  whenever  $b_{g'} \neq 0$  (see [Qin17, Lemma 3.1.10(iii)]). Applying  $\mathfrak{f}_{F,d+m}$ , we obtain  $\mathfrak{f}_d x(\mathbf{t}')^{d'} * \mathfrak{f}_m z = \sum_{g'} b_{g'} \mathfrak{f}_{d+m} x(\mathbf{t}')^{g'}$ . Since  $\underline{k}$  is a sequence on  $I_{uf} \setminus F$ ,  $\forall i$ ,  $x_i(\boldsymbol{\mu}_{\underline{k}}\mathbf{t}) = x_i(\boldsymbol{\mu}_{\underline{k}}\mathfrak{f}_F\mathbf{t})$  has support in  $I_{uf} \setminus F$ . Thus  $\mathfrak{f}_d x(\mathbf{t}')^{d'}$  equals  $x(\boldsymbol{\mu}_{\underline{k}}\mathfrak{f}_F\mathbf{t})^{d'}$ , and  $\mathfrak{f}_{d+m} x(\mathbf{t}')^{g'}$  is either  $x(\boldsymbol{\mu}_{\underline{k}}\mathfrak{f}_F\mathbf{t})^{g'}$  or 0. We deduce that  $\mathfrak{f}_m z \in \mathcal{LP}(\boldsymbol{\mu}_{\underline{k}}\mathfrak{f}_F\mathbf{t})$ .

3.3. **Properties and applications.** We will prove some key properties of the freezing operators. More results are in [Qin24a]. The following lemma is crucial.

**Lemma 3.9** (Mutation compatibility). Take seeds  $\mathbf{t} = \mu_k \mathbf{t}'$  for some  $k \in I_{uf} \setminus F$ and  $z \in \mathcal{LP}(\mathbf{t}) \cap (\mu_k^*)^{-1} \mathcal{LP}(\mathbf{t}')$ . Assume that z has a unique  $\prec_{\mathbf{t}}$ -maximal degree min  $\mathcal{LP}(\mathbf{t})$  and  $\mu_k^* z$  has a unique  $\prec_{\mathbf{t}'}$ -maximal degree m' in  $\mathcal{LP}(\mathbf{t}')$ . Then we have  $f_{F,m'}^{\mathbf{t}}(\mu_k^* z) = \mu_k^* f_{F,m}^*(z)$ .

Proof. For any  $n = \sum n_i e_i(\mathbf{t}) \in N_{uf}^{\oplus}(\mathbf{t})$  with  $n_i \in \mathbb{N}$ , denote  $n_F := (n_j)_{j \in F} \in \mathbb{N}^F$ . Decompose z as  $z = \sum_{n \in N_1} c_n x^m \cdot y^n + \sum_{n \in N_2} c_n x^m \cdot y^n$ , where  $c_n \in \mathbb{k}$ ,  $N_1 = \{n | c_n \neq 0, n_F = 0\}$ , and  $N_2 = \{n | c_n \neq 0, n_F > 0\}$ . Then  $\mathfrak{f}_{F,m}^{\mathbf{t}} z = \sum_{n \in N_1} c_n x^m \cdot y^n$ . Consider the pointed element  $\mu_k^*(y^n)$  of  $\widehat{\mathcal{LP}}(\mathbf{t}')$ . We have  $\deg^{\mathbf{t}'}\mu_k^*(y^n) = \deg^{\mathbf{t}'}(y')^{h(n)}$ , where  $h(n) \in N_{uf}(\mathbf{t}')$ . Note that h(n) = h(n') if and only if n = n'. Moreover,  $n_i = h(n)_i, \forall i \neq k$ , see [Qin24c, (2.4)], which implies  $h(n)_F = n_F$ .

Note that we have a finite decomposition  $\mu_k^* z = \sum_{n:c_n \neq 0} \mu_k^* (c_n x^m \cdot y^n)$  in  $\widehat{\mathcal{LP}}(\mathbf{t}')$ , see [Qin24c, Lemma 3.3.7(2)]. Since m' is the unique  $\prec_{\mathbf{t}'}$ -maximal degree of z, it equals  $\deg^{\mathbf{t}'} x^m + \deg^{\mathbf{t}'} y^{h(\widetilde{n})}$  for some  $c_{\widetilde{n}} \neq 0$  such that  $h(n) > h(\widetilde{n})$ ,  $\forall n \neq \widetilde{n}$ ,  $c_n \neq 0$ . Denote  $N = N'_1 \sqcup N'_2$  for  $N'_1 = \{n|c_n \neq 0, h(n)_F = h(\widetilde{n})_F\}$  and  $N'_2 = \{n|c_n \neq 0, h(n)_F > h(\widetilde{n})_F\}$ . We have  $f_{F,m'}^{\mathbf{t}'}\mu_k^* z = \sum_{n \in N'_1} \mu_k^* (c_n x^m \cdot y^n)$ . Moreover,  $0 \in N_1$  implies  $h(\widetilde{n})_F = h(0)_F = 0$ . Consequently,  $N'_1 = \{n|c_n \neq 0, h(n)_F = 0\} = \{n|c_n \neq 0, n_F = 0\} = N_1$ . The desired claim follows.

The following result follows from Lemma 3.9.

reachable, then  $\mathfrak{fS}$  is a  $\mathcal{M}^{\circ}$ -pointed k-basis of  $\mathcal{U}(\mathfrak{f}_{F}\mathbf{t})$ .

**Theorem 3.10.** If an element  $z \in \mathcal{U}(t_0)$  is compatibly pointed at all seeds in  $\Delta_{t_0}^+$ , then  $\mathfrak{f}_F z \in \mathcal{U}(\mathfrak{f}_F t_0)$  is compatibly pointed at all seeds in  $\Delta_{\mathfrak{f}_F t_0}^+$ .

Theorem 3.10, [Qin24c, Theorem 4.3.1], and Remark 3.4 imply the result below. **Theorem 3.11.** Assume that S is a  $\mathcal{M}^{\circ}$ -pointed  $\Bbbk$ -basis of  $\mathcal{U}(\mathbf{t})$ . If  $\mathfrak{f}_{F}\mathbf{t}$  is injective-

In [Mul16, Theorem 1.4.1], Muller showed that, if **t** is injective-reachable, then  $\mathfrak{f}_F \mathbf{t}$  is injective-reachable too. His proof also yields the following result. Details could be found in [Qin24a].

**Lemma 3.12.** If  $m \in M^{\circ}(\mathbf{t})$  is the degree of a localized cluster monomial for  $\mathcal{U}(\mathbf{t})$ , then m is also the degree of a localized cluster monomial for  $\mathcal{U}(\mathfrak{f}_{F}\mathbf{t})$ .

**Theorem 3.13.** Assume that  $\mathfrak{f}_F \mathbf{t}$  is injective-reachable. Let z be any localized (quantum) cluster monomial of  $\mathcal{U}(\mathbf{t})$ . Then  $\mathfrak{f}_F z$  is a localized (quantum) cluster monomial of  $\mathcal{U}(\mathfrak{f}_F \mathbf{t})$ .

*Proof.* By Lemma 3.12,  $\deg^{\mathfrak{f}_F \mathbf{t}} \mathfrak{f}_F z$  equals  $\deg^{\mathfrak{f}_F \mathbf{t}} u$  for some localized cluster monomial of  $\mathcal{U}(\mathfrak{f}_F \mathbf{t})$ . By Theorem 3.10,  $\mathfrak{f}_F z$  is compatibly pointed at  $\Delta^+_{\mathfrak{f}_F \mathbf{t}}$ . Then Lemma 2.11 implies  $\mathfrak{f}_F z = u$ .

In [Qin24a], we will prove Theorem 3.13 without the assumption on  $\mathfrak{f}_F \mathfrak{t}$ . From now on, we will use it in this generality.

**Definition 3.14.** Take any  $z \in \mathcal{A}(\mathbf{t})$ . Assume there is a finite decomposition  $z = \sum_i b_i z_i, b_i \in \mathbb{k}$ , such that  $z_i$  have distinct degrees in  $\mathcal{LP}(\mathbf{t})$  and are products of cluster variables and  $x_j^{-1}, j \in I_{\mathrm{f}}$ . The decomposition is called a nice cluster decomposition in  $\mathbf{t}$ .

**Lemma 3.15.** Take any  $z \in \mathcal{LP}_{\leq m}(\mathbf{t})$ . If it has a nice cluster decomposition in  $\mathbf{t}$ ,  $\mathfrak{f}_m z$  belongs to  $\mathcal{A}(\mathfrak{f}_F \mathbf{t})$ .

*Proof.* Denote the nice cluster decomposition by  $z = \sum_i b_i z_i$ ,  $b_i \in \mathbb{k}$ . Since  $z_i$  have distinct degrees,  $z_i \in \mathcal{LP}_{\preceq m}(\mathbf{t})$ , see [Qin24c, Lemma 4.1.1]. Denote  $z_i = s_1 * s_2 * \cdots * s_l$ , where  $s_j$  are cluster variables or  $x_j^{-1}$ ,  $j \in I_f$ . By Lemma 3.6,  $\mathfrak{f}_{F,m}(z_i)$  equals  $\mathfrak{f}_{F,\deg s_1}s_1 * \cdots * \mathfrak{f}_{F,\deg s_l}s_l$  or 0. By Theorem 3.13,  $\mathfrak{f}_{F,\deg s_i}s_i$  are localized cluster monomials. Therefore,  $z_i \in \mathcal{A}(\mathfrak{f}_F \mathbf{t})$  and the claim follows.

Combining Theorem 3.11 and Lemma 3.15, we obtain the following useful result.

**Corollary 3.16.** Assume that  $\mathcal{U}(\mathbf{t})$  has a  $\mathcal{M}^{\circ}$ -pointed k-basis  $\mathcal{S} = \{s_m | m \in M^{\circ}(\mathbf{t})\}$ such that  $s_m$  are [m]-pointed. If  $\mathcal{A}(\mathbf{t}) = \mathcal{U}(\mathbf{t})$ ,  $\mathfrak{f}_F \mathbf{t}$  is injective-reachable, and each  $s_m$  has a nice cluster decomposition, then  $\mathcal{A}(\mathfrak{f}_F \mathbf{t}) = \mathcal{U}(\mathfrak{f}_F \mathbf{t}) = \mathsf{Span}_{\Bbbk}\{\mathfrak{f}_{F,m}s_m | m \in M^{\circ}(\mathbf{t})\}$ . Moreover, each  $\mathfrak{f}_{F,m}s_m$  has a nice cluster decomposition.

**Theorem 3.17.** Take  $\mathbb{k} = \mathbb{Z}[q^{\pm \frac{1}{2}}]$ . Assume that  $\mathcal{U}(\mathbf{t})$  possesses the common triangular basis  $\mathbf{L} = {\mathbf{L}_m | m \in M^\circ(\mathbf{t})}$ , where  $\mathbf{L}_m$  are [m]-pointed. Then  $\mathfrak{f}_F \mathbf{L} := {\mathfrak{f}_{F,m} \mathbf{L}_m | m \in M^\circ(\mathbf{t})}$  is the common triangular basis for  $\mathcal{U}(\mathfrak{f}_F \mathbf{t})$ .

*Proof.* Since **L** is the common triangular basis, it contains all cluster monomials of  $\mathcal{U}(\mathbf{t})$ . Note that any cluster monomials of  $\mathcal{U}(\mathfrak{f}_F \mathbf{t})$  must be a cluster monomial of  $\mathcal{U}(\mathbf{t})$  with support in  $I_{uf} \setminus F$ , see Remark 3.4. So  $\mathfrak{f}\mathbf{L}$  contains all cluster monomials of  $\mathcal{U}(\mathfrak{f}_F \mathbf{t})$ . We claim that  $\mathfrak{f}\mathbf{L}$  is the triangular basis for  $\mathcal{U}(\mathfrak{f}_F \mathbf{t})$  with respect to the seed  $\mathfrak{f}_F \mathbf{t}$ . If so, it follows from [Qin20b, Proposition 6.4.3, Theorem 6.5.3] that  $\mathfrak{f}_F \mathbf{L}$  is the common triangular basis for  $\mathcal{U}(\mathfrak{f}_F \mathbf{t})$ .

By Theorem 3.11,  $\mathfrak{f}_F \mathbf{L}$  is a  $M^\circ(\mathbf{t})$ -pointed basis for  $\mathcal{U}(\mathfrak{f}_F \mathbf{t})$ . Its bar-invariance is clear. We have also seen that it contains the cluster monomials of  $(\mathfrak{f}_F \mathbf{t})[1]$ . It remains to check the triangularity. Applying  $\mathfrak{f}_{F,m+f_i}$  to (2.2) and using Lemma 3.6, we get  $[x_i * \mathfrak{f}_{F,m} \mathbf{L}_m]^{\mathbf{t}} = \mathbf{L}_{m+f_k} + \sum_{m' \prec \mathbf{t}_m + f_k} b_{m'} \mathfrak{f}_{F,m+f_k} \mathbf{L}_{m'}, b'_m \in \mathbf{m}, \forall i$ , where  $\mathfrak{f}_{F,m+f_i} \mathbf{L}_{m'}$  is 0 or  $\mathfrak{f}_{F,m'} \mathbf{L}_{m'}$ . So  $\mathfrak{f}_F \mathbf{L}$  is the triangular basis with respect to  $\mathbf{t}$ .

**Theorem 3.18.** Let  $\mathbf{L} = {\mathbf{L}_m | m \in M^\circ(\mathbf{t})}$  denote the common triangular basis. For any  $\mathbf{L}_m$  and  $k \in F$ , when  $d_k \in \mathbb{N}$  are large enough, we have

$$\mathfrak{f}_{F,m}\mathbf{L}_m = x^{-\sum_{k\in F} d_k f_k} \cdot \mathbf{L}_{m+\sum_{k\in F} d_k f_k},$$

and  $x_j \cdot \mathbf{L}_{m + \sum_{k \in F} d_k f_k} \in \mathbf{L}$  for any  $j \in F$ .

We will prove Theorem 3.18 in the end of this subsection.

Corollary 3.19. The following claims are true.

(1)  $\mathfrak{f}_F \mathbf{L} = \{ x^{-p} \cdot b | b \in \mathbf{L} \cap \overline{\mathcal{U}}, \operatorname{supp} b \cap F = \emptyset, p \in \mathbb{N}^{F \sqcup I_f} \}.$ 

(2)  $\mathfrak{f}_F \mathbf{L} = \{ x^{-p} \cdot b | b \in \mathbf{L} \cap \overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t}), p \in \mathbb{N}^{F \sqcup I_{\mathfrak{f}}} \}.$ 

(3)  $\mathfrak{f}_F \mathbf{L} \cap \overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t}) = \mathbf{L} \cap \overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t}).$ 

(4)  $\mathbf{L} \cap \overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t}) = \{ b \in \mathbf{L} \cap \overline{\mathcal{U}}, \operatorname{supp} b \cap F = \emptyset \}.$ 

*Proof.* (1) Take any  $x^{-p} \cdot b$  from the set on the right hand side. We have  $x^{-p} \cdot b = x^{-p} \cdot \mathfrak{f}_F b \in x^{-p} \cdot \mathfrak{f}_F \mathbf{L}$ . Since  $\mathfrak{f}_F \mathbf{L}$  is the common triangular basis of  $\mathcal{U}(\mathfrak{f}_F \mathbf{t})$  by Theorem 3.17, we have  $x^{-p} \cdot \mathfrak{f}_F \mathbf{L} = \mathfrak{f}_F \mathbf{L}$ . Thus  $x^{-p} \cdot b \in \mathfrak{f}_F \mathbf{L}$ .

Conversely, take any  $\mathbf{L}_m \in \mathbf{L}$ . By Theorem 3.18, when d large enough,  $\mathfrak{f}_F \mathbf{L}_m = x^{-d\sum_{k\in F} f_k} \cdot \mathbf{L}_{m+d\sum_{k\in F} f_k}$  with  $\operatorname{supp} \mathbf{L}_{m+d\sum_{k\in F} f_k} \cap F = \emptyset$ . Rewrite  $\mathfrak{f}_F \mathbf{L}_m = x^{-d\sum_{i\in F\sqcup I_f} f_i} \cdot \mathbf{L}_{m+d\sum_{i\in F\sqcup I_f} f_i}$ . We have  $\mathbf{L}_{m+d\sum_{i\in F\sqcup I_f} f_i} = \mathfrak{f}_F \mathbf{L}_{m+d\sum_{i\in F\sqcup I_f} f_i} \in \mathfrak{f}_F \mathbf{L}$ . For d large enough,  $\nu_i(\mathbf{L}_{m+d\sum_{i\in F\sqcup I_f} f_i}) \geq 0$ ,  $\forall i \in F \sqcup I_f$ . So we have  $\mathbf{L}_{m+d\sum_{i\in F\sqcup I_f} f_i} \in \overline{\mathcal{U}}$  and  $\mathbf{L}_{m+d\sum_{i\in F\sqcup I_f} f_i} = \mathfrak{f}_F \mathbf{L}_{m+d\sum_{i\in F\sqcup I_f} f_i} \in \overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t})$ . Particularly,  $\mathfrak{f}_F \mathbf{L}_m$  belongs to the set on the right.

(2) In the proof of (1), we have shown  $\mathfrak{f}_F \mathbf{L} \subset \{x^{-p} \cdot b | b \in \mathbf{L} \cap \overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t}), \operatorname{supp} b \cap F = \emptyset, p \in \mathbb{N}^{F \sqcup I_{\mathfrak{f}}}\}$ . Therefore,  $\mathcal{U}(\mathfrak{f}_F \mathbf{t}) = \operatorname{Span} \mathfrak{f}_F \mathbf{L} \subset \operatorname{Span} \{x^{-p} \cdot b | b \in \mathbf{L} \cap \overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t}), p \in \mathbb{N}^{F \sqcup I_{\mathfrak{f}}}\} \subset \mathcal{U}(\mathfrak{f}_F \mathbf{t})$ . The claim follows.

(3) The claim follows from (2).

(4) By (2), any  $b \in \mathbf{L} \cap \overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t})$  is contained in  $\mathfrak{f}_F \mathbf{L}$ . It follows that  $\mathfrak{supp} b \cap F = \emptyset$ . Note that  $\overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t}) \subset \overline{\mathcal{U}}(\mathbf{t})$  by Proposition 3.3. So  $\mathbf{L} \cap \overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t})$  is a subset of the set on the right. Conversely, for any b from the set on the right, we have  $\mathfrak{f}_F b = b$ , thus  $b \in \mathcal{U}(\mathfrak{f}_F \mathbf{t})$ . Note that  $\nu_i(\mathfrak{f}_F b) = \nu_i(b) \ge 0$  for any  $i \in F \sqcup I_{\mathfrak{f}}$  by Lemma 2.13. We deduce that  $b = \mathfrak{f}_F b \in \overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t})$ . So we obtain the inverse inclusion.

Let us consider some general properties. Take any  $k \in I$ . Let  $\mathcal{Z}$  denote a klinearly independent subset of  $\mathcal{LP}(\mathbf{t})$ . Assume that, for any  $z \in \mathcal{Z}$ , there is a finite decomposition

(3.1) 
$$x_k * z = \sum_{z_j \in \mathcal{Z}} b_j z_j, \ b_j \in \mathbb{Z}.$$

**Definition 3.20.** We say that  $\mathcal{Z}$  has the stabilization property with respect to  $x_k$ , if the following holds: Take any  $z \in \mathcal{Z}$  and  $d \in \mathbb{N}$ , consider the decomposition

(3.2) 
$$x_k^d * z = \sum_{z_j^{(d)} \in \mathcal{Z}} b_j^{(d)} z_j^{(d)}, \ b_j^{(d)} \in \Bbbk.$$

such that, when d is large enough,  $x_k^{-d} * z_j^{(d)}$  stabilize, and  $x_k * z_j^{(d)} \in q^{\frac{1}{2}\mathbb{Z}}\mathcal{Z}$  for any  $z_i^{(d)}$  appearing.

**Lemma 3.21.** If the Laurent coefficients of the elements of Z and the decomposition coefficients in (3.1) always belong to  $\mathbb{k}_{\geq 0}$ , Z has the stabilization property with respect to  $x_k$ .

*Proof.* Consider the decomposition (3.2) in  $\mathbb{Z}[q^{\pm \frac{1}{2}}, (x_i^{\pm})_{i \in I}]$ . As d grows, the sum  $(\sum b_j^{(d)})|_{q \mapsto 1}$  weakly increases. Since the total number of Laurent monomials  $q^{\alpha}x^m$  appearing in  $x_k^d * z_j$  is constant for varying d, the sum must stabilizes when d is large enough. In this case, for any  $z_j^{(d)}$  appearing, we have  $x_k * z_j^{(d)} = z'_j$  for some  $z'_j = q^{\alpha_j} z_j^{(d+1)}$ , where  $\alpha_j \in q^{\frac{\mathbb{Z}}{2}}, z_j^{(d+1)} \in \mathbb{Z}$ . The desired claim follows.

We work at the quantum level by choosing  $\mathbb{k} = \mathbb{Z}[q^{\pm \frac{1}{2}}]$  from now on.

**Definition 3.22.** Assume that the elements of Z are pointed at distinct degrees. We say Z satisfies the triangularity condition with respect to  $x_k$  if,  $\forall z \in Z$ , we have a finite decomposition

$$(3.3) [x_k * z] = z_0 + \sum_{\deg z_j \prec_t \deg z_0} b_j z_j$$

such that  $b_j \in q^{-\frac{1}{2}}\mathbb{Z}[q^{-\frac{1}{2}}]$  for  $j \neq 0$ .

**Lemma 3.23.** Assume the elements of  $\mathcal{Z}$  are bar-invariant and pointed at distinct degrees. If it has the stabilization property and satisfies triangularity condition with respect to  $x_k$ , then we have  $\mathfrak{f}_{\{k\}}z = [x_k^{-d} * z_0^{(d)}]$  for  $d \in \mathbb{N}$  large enough, where  $[x_k^d * z] = z_0^{(d)} + \sum_{\deg z_j^{(d)} \prec_{\mathfrak{t}} \deg z_0^{(d)}} b_j^{(d)} z_j^{(d)}$ .

*Proof.* The case  $k \in I_{\mathsf{f}}$  is trivial. Now assume  $k \in I_{\mathsf{uf}}$ . By the stabilization property, we can take d large enough such that  $[x_k * z_0^{(d)}] = z_0^{(d+1)}$ . Then the bar-invariance of  $z_0^{(d+1)}$  implies that  $z_0^{(d)} q$ -commutes with  $x_k$ , i.e.,  $\mathfrak{f}_{\{k\}, \deg z_0^{(d)}} z_0^{(d)} = z_0^{(d)}$ . Take

any  $z_i^{(d)}$  such that  $\deg z_i^{(d)}$  is  $\prec_t$ -maximal in  $\{\deg z_j^{(d)} | j \neq 0, b_j \neq 0\}$ . Then the Laurent coefficient of  $x^{\deg z_i^{(d)}}$  in  $[x_k^d * z]$ , contributed from  $z_0^{(d)}$  and  $b_i^{(d)} z_i^{(d)}$ , is not bar-invariant since  $b_i^{(d)} \in q^{-\frac{1}{2}} \mathbb{Z}[q^{-\frac{1}{2}}]$ . So we must have  $\deg z_i^{(d)} = \deg z_0^{(d)} + p^* n^{(i)}$  with  $n_k^{(i)} > 0$ . We deduce that, for any  $1 \leq j \leq s$ ,  $\deg z_j^{(d)} = \deg z_0^{(d)} + \tilde{B}n^{(j)}$  with  $n_k^{(j)} > 0$ . Applying  $\mathfrak{f}_{\deg z_0^{(d)}} = \mathfrak{f}_{\deg z+df_k}$  to both sides of the decomposition of  $[x_k^d * z]$ , we obtain  $\mathfrak{f}_{\deg z_0^{(d)}} z_j^{(d)} = 0$ ,  $\forall j \neq 0$ , see Lemma 3.6. Thus  $\mathfrak{f}_{\deg z_0^{(d)}}[x_k^d * z] = z_0^{(d)}$  and the claim follows.

**Lemma 3.24.** Assume that Z satisfies triangularity condition with respect to  $x_k$  and the elements of Z are pointed at distinct degrees and copointed at distinct codegrees. Then it has the stabilization property with respect to  $x_k$ .

*Proof.* The case  $k \in I_{\rm f}$  is trivial. Now assume  $k \in I_{\rm uf}$ . In the finite decomposition (3.3), we have  $\operatorname{codeg} z + f_k \preceq_{\rm t} \operatorname{codeg} z_j$  and the equality holds for a unique  $z_s$ , see [Qin17, Lemma 3.1.10(iii)].

First, assume  $z_s = z_0$ . Then the Laurent monomial  $x^{\operatorname{codeg} z+f_k}$  of  $[x_k * z]$  is 1, i.e.,  $\lambda(f_k, \deg z) = \lambda(f_k, \operatorname{codeg} z)$ . It follows that  $(\operatorname{suppDim} z)_k = 0$ . Thus  $x_k$  and z q-commute. Applying the bar-involution to (3.3), we deduce that  $[x_k * z] = z_0$ .

Otherwise, we always have  $\operatorname{codeg} z + f_k \preceq_{\mathbf{t}} \operatorname{codeg} z_j$  and  $\operatorname{deg} z_j \preceq_{\mathbf{t}} \operatorname{deg} z_0 + f_k$  and the equalities do not hold at the same time. So all  $z_j$  appearing satisfy  $\operatorname{suppDim} z_j < \operatorname{suppDim} z$ . By induction on the support dimension, we can assume that  $z_j$  satisfy the stabilization property. Then we deduce that z satisfies this property as well.

*Proof of Theorem 3.18.* Note that the common triangular basis elements are copointed at distinct codegrees [Qin20a]. Theorem 3.18 follows from Lemmas 3.24 and 3.23.

#### 4. Base changes

We interpret changes in the frozen part of cluster algebras as base changes in schemes. While the majority of our claims will be evident in practical use, we provide a general treatment for the sake of completeness.

4.1. Similarity and the correction technique. Let t denote any seed. Recall that, in the quantum case, we have  $\lambda(f_i, p^*e_k) = -\delta_{ik}\delta_k$  for some  $\delta_k \in \mathbb{N}_{>0}$ .

**Definition 4.1** ([Qin14][Qin17]). Let there be given two seeds  $\mathbf{t}, \mathbf{t}'$  not necessarily related by a mutation sequence,  $I'_{uf} = I_{uf}$ , and  $\sigma$  a permutation on  $I_{uf}$ . We say  $\mathbf{t}$  and  $\mathbf{t}'$  are similar (up to  $\sigma$ ) if  $b'_{\sigma i,\sigma j} = b_{ij}$  and  $d'_{\sigma i} = d_i, \forall i, j \in I_{uf}$ . In the quantum case, we further require that there exists some  $\rho \in \mathbb{Q}_{>0}$  such that  $\delta'_{\sigma k} = \rho \delta_k, \forall k \in I_{uf}$ .

Let  $\sigma$  denote any permutation of I such that  $\sigma I_{uf} = I_{uf}$ . If we are given a permutation  $\sigma$  on  $I_{uf}$ , extend  $\sigma$  by letting it acts on  $I_f$  trivially. We define the permuted seed  $\sigma \mathbf{t}$  by relabeling the vertices using  $\sigma$ , such that  $b_{\sigma i,\sigma j}(\sigma \mathbf{t}) = b_{ij}$ ,  $x_{\sigma i}(\sigma \mathbf{t}) = x_i$ ,  $d_{\sigma i}(\sigma \mathbf{t}) = d_i$ . In the quantum case, we endow  $\sigma \mathbf{t}$  with the bilinear form  $\sigma \lambda$  such that  $\sigma \lambda(\sigma m, \sigma g) := \lambda(m, g)$ , where  $(\sigma m)_{\sigma i} := m_i \forall m = (m_i) \in \mathbb{Z}^I$ . Apparently,  $\mathbf{t}$  and  $\sigma \mathbf{t}$  are similar.

From now on, let **t** and **t'** denote two seeds similar up to  $\sigma$ . Modifying **t'** by permuting its vertices if necessary, we can assume  $\sigma$  is trivial. Note that  $\mu$ **t** and  $\mu$ **t'** are similar,  $\forall \mu$ , see [Qin17].

At the quantum level, it is often sufficient to consider the case  $\rho = 1$ . For completeness, let us treat the general case  $\rho = \frac{d'_{\rho}}{d_{\rho}}$  for coprime  $d_{\rho}, d'_{\rho} \in \mathbb{N}_{>0}$ . Recall  $\mathbb{k} = \mathbb{Z}[q^{\pm \frac{1}{2}}]$ . Define  $\mathbb{k}' := \mathbb{Z}[q^{\pm \frac{1}{2d_{\rho}}}]$  and  $\mathbb{k}'' := \mathbb{Z}[q^{\pm \frac{1}{2d'_{\rho}}}]$ . Then  $\mathbb{k}' = \bigoplus_{j=0}^{d_{\rho}-1} \mathbb{k} \cdot v_j$ is a free k-module, where  $v_j := q^{\frac{j}{2d_{\rho}}}$ . Define  $\mathcal{LP}'_{\mathbb{k}'}, \ \widehat{\mathcal{LP}}'_{\mathbb{k}'}, \ \mathcal{A}'_{\mathbb{k}'}$ , and  $\mathcal{U}'_{\mathbb{k}'}$  associated to  $\mathbf{t}'$  using  $\mathbb{k}'$  instead of  $\mathbb{k}$  in their constructions. We have  $\mathcal{LP}'_{\mathbb{k}'} = \mathcal{LP}' \otimes \mathbb{k}',$  $\widehat{\mathcal{LP}}'_{\mathbb{k}'} = \widehat{\mathcal{LP}}' \otimes \mathbb{k}'$ , and  $\mathcal{A}'_{\mathbb{k}'} = \mathcal{A}' \otimes \mathbb{k}'$ .

**Proposition 4.2.** We have  $\mathcal{U}'_{\Bbbk'} = \mathcal{U}' \otimes \Bbbk'$ .

*Proof.* The skew-field of fractions  $\mathcal{F}(\mathbf{t}')_{\Bbbk'}$  for  $\mathcal{LP}(\mathbf{t}')_{\Bbbk'}$  contains  $\mathcal{F}(\mathbf{t}') \otimes \Bbbk'$ . Take any  $\mathbf{s} = \boldsymbol{\mu} \mathbf{t}' \in \Delta_{\mathbf{t}'}^+$ . We have  $\boldsymbol{\mu}^* : \mathcal{F}(\mathbf{s})_{\Bbbk'} \simeq \mathcal{F}(\mathbf{t}')_{\Bbbk'}$ . Since  $\boldsymbol{\mu}^*(v_j) = v_j$ ,  $(\boldsymbol{\mu}^* \mathcal{LP}(\mathbf{s})) \otimes \Bbbk'$  equals  $\boldsymbol{\mu}^* \mathcal{LP}(\mathbf{s})_{\Bbbk'}$  in  $\mathcal{F}(\mathbf{t}')_{\Bbbk'}$ .

For any  $\mathbf{s}' = \boldsymbol{\mu}' \mathbf{t}' \in \Delta_{\mathbf{t}'}^+$ , we have  $(\boldsymbol{\mu}^* \mathcal{LP}(\mathbf{s}) \cdot v_j) \cap ((\boldsymbol{\mu}')^* \mathcal{LP}(\mathbf{s}') \cdot v_{j'}) \subset (\mathcal{F}(\mathbf{t}') \cdot v_j) \cap (\mathcal{F}(\mathbf{t}') \cdot v_{j'}) = 0$  whenever  $j \neq j'$ . Therefore,  $\mathcal{U}'_{\mathbf{k}'} = \cap_{\mathbf{s} = \boldsymbol{\mu} \mathbf{t}' \in \Delta_{\mathbf{t}'}^+} (\boldsymbol{\mu}^* \mathcal{LP}(\mathbf{s}) \otimes \mathbf{k}') = \oplus_j (\cap_{\mathbf{s}} (\boldsymbol{\mu}^* \mathcal{LP}(\mathbf{s}) \cdot v_j)) = \oplus_j (\cap_{\mathbf{s}} \boldsymbol{\mu}^* \mathcal{LP}(\mathbf{s})) \cdot v_j = \mathcal{U}' \otimes \mathbf{k}'.$ 

**Lemma 4.3.** For any subset  $\mathcal{Z}$  of  $\widehat{\mathcal{LP}}$ ,  $\widehat{\mathcal{LP}} \cap \operatorname{Span}_{\Bbbk'} \mathcal{Z}$  equals  $\operatorname{Span}_{\Bbbk} \mathcal{Z}$  in  $\widehat{\mathcal{LP}} \otimes \Bbbk'$ . Proof. Recall  $\widehat{\mathcal{LP}} \otimes \Bbbk' = \bigoplus_{j=0}^{d_{\rho}-1} \widehat{\mathcal{LP}} \cdot v_j$ ,  $v_j = q^{\frac{j}{2d_{\rho}}}$ . Obviously,  $\widehat{\mathcal{LP}} \cap \operatorname{Span}_{\Bbbk'} \mathcal{Z}$ contains  $\operatorname{Span}_{\Bbbk} \mathcal{Z}$ . For any  $w \in \operatorname{Span}_{\Bbbk'} \mathcal{Z}$ , we have  $w = \sum_i (\sum_{j=0}^{d_{\rho}-1} c_{ij}v_j)z_i$  for  $z_i \in \mathcal{Z}$ ,  $c_{ij} \in \Bbbk$ . Then  $w = \sum_j w_j$  for  $w_j := \sum_i c_{ij}v_jz_i \in \widehat{\mathcal{LP}} \cdot v_j$ . If  $w \in \widehat{\mathcal{LP}}$ , then  $w = w_0$ . Therefore,  $\operatorname{Span}_{\Bbbk} \mathcal{Z}$  contains  $\widehat{\mathcal{LP}} \cap \operatorname{Span}_{\Bbbk'} \mathcal{Z}$ .

**Definition 4.4** ([Qin14]). Take any  $z = x^m \cdot (1 + \sum_{n \in \mathbb{N}^{I_{uf}}} c_n y^n) \in \widehat{\mathcal{LP}}, c_n \in \mathbb{Z}[q^{\pm \frac{1}{2}}].$ The elements  $z' = (x')^{m'} \cdot (1 + \sum_{n \in \mathbb{N}^{I_{uf}}} c'_n(y')^n) \in \widehat{\mathcal{LP}}'_{\Bbbk'}, c'_n \in \Bbbk'$ , are said to be similar to z if  $\operatorname{pr}_{I_{uf}} m' = \operatorname{pr}_{I_{uf}} m$  and  $c'_n = c_n|_{q^{\frac{1}{2}} \mapsto q^{\frac{1}{2}\rho}}$ . If  $q^{\frac{1}{2}} = 1$ , we require  $c'_n = c_n$ .

Note that, when z is given, any z' similar to z is determined by  $\deg^{t'} z'$ .

Let us extend the construction of similar elements.  $\forall m \in M^{\circ}(\mathbf{t})$ , define a  $\widehat{\Bbbk[N_{uf}]}$ -bimodule and a  $\Bbbk[N_{uf}]$ -bimodule:

(4.1) 
$$\widehat{V}_m := x^m \cdot \widehat{\Bbbk[N_{\mathsf{uf}}]}, \ V_m := x^m \cdot \Bbbk[N_{\mathsf{uf}}].$$

Then  $\widehat{V}_m \cap \widehat{V}_h \neq 0$  if and only if  $h \in m + p^* N_{uf}$ , which implies  $\widehat{V}_m = \widehat{V}_h$ . Therefore,  $\widehat{\mathcal{LP}} = \bigoplus_{m_i} \widehat{V}_{m_i}$ , where  $m_i$  are any chosen representatives of  $M^{\circ}/p^* N_{uf}$ . Similarly, we have  $\mathcal{LP} = \bigoplus_{m_i} V_{m_i}$ .

Choose  $m' \in M^{\circ}(\mathbf{t}')$  such that  $\operatorname{pr}_{I_{\operatorname{uf}}} m = \operatorname{pr}_{I_{\operatorname{uf}}} m'$ . Similarly define  $\widehat{V}'_{m'} \subset \widehat{\mathcal{LP}}'$  for  $\mathbf{t}'$ . We have the  $\mathbb{Z}$ -algebra isomorphism  $\operatorname{var} : \overline{\Bbbk''[N_{\operatorname{uf}}]} \to \widehat{\Bbbk'[N'_{\operatorname{uf}}]}$  such that  $\operatorname{var}(y^n) = (y')^n$ ,  $\operatorname{var}(q^{\frac{1}{2d'_{\rho}}}) = q^{\frac{1}{2d'_{\rho}}\rho} = q^{\frac{1}{2d_{\rho}}}$ . It induces a bijective  $\mathbb{Z}$ -linear map  $\operatorname{var} := \operatorname{var}_{m',m}$  from  $\widehat{V}_m \otimes \Bbbk''$  to  $(\widehat{V}'_{m'}) \otimes \Bbbk'$  such that  $\operatorname{var}(q^{\frac{1}{2d'_{\rho}}}) = q^{\frac{1}{2d_{\rho}}}$ ,  $\operatorname{var}(x^m \cdot y^n) := (x')^{m'} \cdot (y')^n$ . We have  $\operatorname{var}(a * z * b) = \operatorname{var}(a) * \operatorname{var}(z) * \operatorname{var}(b)$ ,  $\forall a, b \in \widehat{\Bbbk''[N_{\operatorname{uf}}]}, z \in V_m$ .

We observe that, for any  $(m + \widetilde{B}n)$ -pointed element  $z \in \widehat{\mathcal{LP}}$ ,  $n \in N_{uf}$ , the image  $\operatorname{var}_{m',m}(z)$  is  $(m' + \widetilde{B'}n)$ -pointed and similar to z in the sense of Definition 4.4.

In general, any  $z \in \widehat{\mathcal{LP}}$  belongs to some  $\bigoplus_{m_i} \widehat{V}_{m_i}$ . If z is further contained in  $\mathcal{LP}$ , then  $z \in \bigoplus_{m_i} V_{m_i}$ . Denote  $z = \sum_i z_i$  with  $z_i \in \widehat{V}_{m_i}$ . By choosing  $m'_i$  such that  $\operatorname{pr}_{I_{\mathrm{uf}}} m'_i = \operatorname{pr}_{I_{\mathrm{uf}}} m_i$ , we obtain an element  $\operatorname{var}_{\underline{m}',\underline{m}}(z) := \sum_i \operatorname{var}_{m'_i,m_i}(z_i)$ , where  $\underline{m} = (m_i)_i$  and  $\underline{m}' = (m'_i)_i$ . We could say  $\operatorname{var}_{\underline{m}',\underline{m}}(z)$  is similar to z.

**Lemma 4.5.** Assume  $z = \sum_i z_i, z_i \in V_{m_i}, V_{m_i} \cap V_{m_j} = 0$  when  $i \neq j$ . If  $z \in U$ , then  $z_i \in U$ .

Proof. It suffices to show  $z_i \in \mathcal{LP}(\mathbf{s})$  for any seed  $\mathbf{s} = \boldsymbol{\mu}^{-1}\mathbf{t}$ . Following [Qin24c, Section 3.3], define the bijective linear map  $\psi := \psi_{\mathbf{s},\mathbf{t}} : M^{\circ} \to M^{\circ}(\mathbf{s})$  such that  $\psi(\mathsf{deg}^{\mathbf{t}} x_i) = \mathsf{deg}^{\mathbf{s}} \boldsymbol{\mu}^* x_i, \forall i$ . Then,  $\forall m \in M^{\circ}, \boldsymbol{\mu}^* x^m$  is a  $\psi m$ -pointed element in  $\widehat{\mathcal{LP}}(\mathbf{s})$ . Moreover, we can deduce from the mutation rule of y-variables ([Qin24c, (2.4)]) that  $\psi$  restricts to a bijection from  $p^* N_{\mathsf{uf}}$  to  $p^* N_{\mathsf{uf}}(\mathbf{s})$ .

We have  $\boldsymbol{\mu}^* z = \sum_i \boldsymbol{\mu}^* z_i$  in  $\widehat{\mathcal{LP}}(\mathbf{s})$ , where  $\boldsymbol{\mu}^* z_i \in \widehat{V}(\mathbf{s})_{h_i}$ ,  $h_i = \psi m_i$ . Moreover, if  $h_i \in h_j + p^* N_{uf}(\mathbf{s})$ , we will have  $m_i \in m_j + p^* N_{uf}$  by applying  $\psi^{-1}$ . In this case, we obtain  $V_{m_i} = V_{m_j}$  and thus i = j. Therefore,  $\widehat{V}(\mathbf{s})_{h_i} \cap \widehat{V}(\mathbf{s})_{h_j} = 0$  if  $i \neq j$ . Combining with  $\boldsymbol{\mu}^* z \in \mathcal{LP}(\mathbf{s})$ , we obtain  $\boldsymbol{\mu}^* z_i \in V(\mathbf{s})_{h_i} \subset \mathcal{LP}(\mathbf{s})$ .

We slightly generalize the correction technique in [Qin14, Qin17] as below.

**Theorem 4.6.** Given  $m_s$ -pointed elements  $z_s \in \widehat{\mathcal{LP}}$  and  $m'_s$ -pointed elements  $z'_s \in \widehat{\mathcal{LP}}'_{\Bbbk'}$ , for  $r \in \mathbb{N}, 1 \leq s \leq r$ , such that  $z'_s$  are similar to  $z_s$ . We do not assume  $m_s$  to be distinct. Take any  $z \in V_m$  and  $z' \in (V'_{m'}) \otimes \Bbbk'$  such that  $\mathsf{pr}_{I_{uf}} m = \mathsf{pr}_{I_{uf}} m'$ ,  $z' = \mathsf{var}_{m',m}(z)$ .

(1) If  $z = [z_1 * \cdots * z_r]^{\mathbf{t}}$ , then  $z' = p' \cdot [z'_1 * \cdots * z'_r]^{\mathbf{t}'}$  for  $p' \in \mathcal{P}'$ , where  $\deg^{\mathbf{t}'} p' = m' - \sum_{r=1}^r m'_r$ .

 $\begin{aligned} & \deg^{\mathbf{t}'}p' = m' - \sum_{s=1}^{r} m'_{s}. \\ & (2) \ If \ z \ = \ \sum_{s=1}^{r} b_{s}z_{s}, \ b_{s} \ \in \ \Bbbk, \ we \ have \ z \ = \ \sum_{s\in J} b_{s}z_{s} \ for \ J \ = \ \{s|\exists n_{s} \ \in \ \mathbb{Z}^{I_{\mathrm{uf}}}, \ m_{s} \ = \ m + \widetilde{B}n_{s}\}. \end{aligned}$ 

*Proof.* (1) follows from direct calculation.

(2) The first claim follows from the decomposition of z in  $\oplus_g \widehat{V}_g$ . Using the  $\mathbb{Z}$ -linear map  $\operatorname{var} := \operatorname{var}_{m',m} : V_m \to V'_{m'} \otimes \Bbbk'$ , we obtain  $z' = \sum_{s \in J} \operatorname{var}(b_s z_s) = \sum_{s \in J} b'_s \operatorname{var}(z_s)$ , where  $\operatorname{var}(z_s)$  is similar to  $z_s$  with  $\operatorname{deg} \operatorname{var}(z_s) = m' + \widetilde{B}' n_s$ . The claim follows by rewriting  $\operatorname{var}(z_s) = p'_s \cdot z'_s$ .

By [FZ07][Tra11][GHKK18], for any mutation sequence  $\boldsymbol{\mu}$  and  $i \in I$ ,  $x_i(\boldsymbol{\mu}\mathbf{t}) \in \mathcal{LP}$  and  $x_i(\boldsymbol{\mu}\mathbf{t}') \in \mathcal{LP}' \subset \mathcal{LP}'_{\Bbbk'}$  are similar.

**Lemma 4.7.** Let  $z \in \mathcal{LP}$  and  $z' \in \mathcal{LP}'_{\Bbbk'}$  be similar pointed elements. (1) If  $z \in \mathcal{A}$ , then  $z' \in \mathcal{A}'_{\Bbbk'}$ . (2) If  $z \in \mathcal{U}$ , then  $z' \in \mathcal{U}'_{\Bbbk'}$ .

*Proof.* (1) z is a polynomial of cluster variables of  $\mathcal{A}$  and  $x_j^{-1}$ ,  $j \in I_{\rm f}$ , with coefficients in k. Then Theorem 4.6 implies that z' is a polynomial of the similar cluster variables of  $\mathcal{A}'$  and  $(x'_i)^{-1}$ ,  $j \in I'_{\rm f}$ , with coefficients in k'. The claim follows.

(2) For any mutation sequence  $\boldsymbol{\mu}$ , denote  $\mathbf{s} = \boldsymbol{\mu}\mathbf{t}$  and  $\mathbf{s}' = \boldsymbol{\mu}\mathbf{t}'$ . We want to show  $z' \in \mathcal{LP}(\mathbf{s}')_{\Bbbk'}$ . Since  $z \in \mathcal{LP}(\mathbf{s})$ , we have a finite decomposition  $z = \sum_m b_m [x(\mathbf{s})^{-d} * x(\mathbf{s})^m]^{\mathbf{t}}$  in  $\widehat{\mathcal{LP}}(\mathbf{t})$ , where  $d, m \in \mathbb{N}^I$  and  $b_m \in \mathbb{k}$ . Note that  $[x(\mathbf{s})^{-d} * x(\mathbf{s})^m]^{\mathbf{t}} \in \widehat{\mathcal{LP}}(\mathbf{t})$  and  $[x(\mathbf{s}')^{-d} * x(\mathbf{s}')^m]^{\mathbf{t}'} \in \widehat{\mathcal{LP}}(\mathbf{t}')$  are similar. By Theorem 4.6, we have a finite decomposition  $z' = \sum_m b'_m p'_m \cdot [x(\mathbf{s}')^{-d} * x(\mathbf{s}')^m]^{\mathbf{t}'}$  in  $\widehat{\mathcal{LP}}(\mathbf{t}')$  where  $b'_m \in \mathbb{k}'$  and  $p'_m \in \mathcal{P}'$ . Therefore,  $z' \in \mathcal{LP}(\mathbf{s}')_{\mathbb{k}'}$ .

4.2. **Base changes.** Let **t** and **t**' denote two similar seeds. Given any subset  $\mathcal{Z}$  of  $\mathcal{LP}(\mathbf{t})$  such that its elements are pointed and it is closed under the  $\mathcal{P}$  commutative multiplication, i.e.,  $x_j^{\pm} \cdot \mathcal{Z} \subset \mathcal{Z}, \forall j \in I_{\mathsf{f}}$ . In practice,  $\mathcal{Z}$  will be  $M^\circ$ -pointed.

Let  $\mathcal{Z}'$  denote the set of elements in  $\mathcal{LP}(\mathbf{t}')_{\Bbbk'}$  similar to the elements of  $\mathcal{Z}'$ . Then  $\mathcal{Z}'$  has pointed elements and is closed under the  $\mathcal{P}'$  commutative multiplication.

**Lemma 4.8.** (1) Let  $\mathcal{Z}''$  denote the set of the elements in  $\mathcal{LP}(\mathbf{t})_{\Bbbk''}$  that are similar to those of  $\mathcal{Z}'$ . Then  $\mathcal{Z}''$  coincides with  $\mathcal{Z}$ .

(2)  $\mathcal{Z}$  is k-linearly independent in  $\mathcal{LP}(\mathbf{t})$  if and only if it is k"-linearly independent in  $\mathcal{LP}(\mathbf{t})_{k''}$ .

(3) Z is k''-linearly independent if and only if Z' is k'-linearly independent.

*Proof.* For any m''-pointed  $z'' \in \mathcal{Z}''$ , there exists some m'-pointed  $z' \in \mathcal{Z}'$  similar to it. Similarly, there exists some m-pointed  $z \in \mathcal{Z}$  similar to z'. Note that  $\Pr_{I_{uf}} m'' = \Pr_{I_{uf}} m' = \Pr_{I_{uf}} m$ . Denote  $p = x^{m''-m}$ . Then  $p \cdot z \in \mathcal{Z}$  and it coincides with z'', since they are both similar to z' and have the same degree.

Claim (2) can be proved using similar arguments as those in the proof of Lemma 4.3. Finally, Claim (3) follows from Theorem 4.6(2) and Claim (1).

**Lemma 4.9.** If  $\mathcal{Z}$  is a k-basis of  $\mathcal{A}$ , then  $\mathcal{Z}'$  is a k'-basis of  $\mathcal{A}'_{k'}$ .

*Proof.* Lemma 4.7 and Lemma 4.8 imply  $\mathcal{Z}' \subset \mathcal{A}'_{\Bbbk'}$  and  $\mathcal{Z}'$  is linearly independent.

Let z' denote any product of cluster variables and  $(x')_j^{-1}$ ,  $j \in I'_{\mathsf{f}}$  of  $\mathcal{A}'$ . It is similar to a product z of the similar cluster variables of  $\mathcal{A}$  and  $x_j^{-1}$ ,  $j \in I_{\mathsf{f}}$ , which belongs to  $\mathsf{Span}_{\Bbbk} \mathcal{Z}$ . By Theorem 4.6, z' belongs to  $\mathsf{Span}_{\Bbbk'} \mathcal{Z}'$ . We deduce that  $\mathsf{Span}_{\Bbbk'} \mathcal{Z}' = \mathcal{A}'_{\Bbbk'}$ .

# **Lemma 4.10.** If $\mathcal{Z}$ is a k-basis of $\mathcal{U}$ , then $\mathcal{Z}'$ is a k'-basis of $\mathcal{U}'_{k'}$ .

*Proof.* Lemma 4.7 implies  $\mathcal{Z}' \subset \mathcal{U}'_{\Bbbk'}$ . Lemma 4.8 implies that  $\mathcal{Z}'$  is linearly independent. By Lemma 4.5, it remains to show that,  $\forall m' \in M^{\circ}(\mathbf{t}')$ , any  $u' \in \mathcal{U}'_{\Bbbk'} \cap (V'_{m'} \otimes \Bbbk')$  is contained in  $\mathsf{Span}_{\Bbbk'} \mathcal{Z}'$ .

Choose any  $m \in M^{\circ}(\mathbf{t})$  such that  $\operatorname{pr}_{I_{\mathrm{uf}}} m = \operatorname{pr}_{I_{\mathrm{uf}}} m'$ . Recall that we have bijective  $\mathbb{Z}$ -linear maps  $\operatorname{var}_{m,m'}: V'_{m'} \otimes \Bbbk' \to V_m \otimes \Bbbk''$  and  $\operatorname{var}_{m',m}: V_m \otimes \Bbbk'' \to V'_{m'} \otimes \Bbbk'$  as before, such that  $\operatorname{var}_{m,m'}$  and  $\operatorname{var}_{m',m}$  are inverse to each other. Denote  $u := \operatorname{var}_{m,m'}(u') \in V_m \otimes \Bbbk''$ . Since  $u' \in \mathcal{U}'_{\Bbbk'}$ , we have  $u \in \mathcal{U}_{\Bbbk''}$  by Lemma 4.7. In addition,  $\mathcal{U}_{\Bbbk''}$  equals  $\mathcal{U} \otimes \Bbbk''$  by Proposition 4.2, which has the  $\Bbbk''$ -basis  $\mathcal{Z}$ . So  $u \in \operatorname{Span}_{\Bbbk''} \mathcal{Z}$ . Therefore,  $u' = \operatorname{var}_{m',m}(u)$  is contained in  $\operatorname{Span}_{\Bbbk'} \mathcal{Z}'$  by Theorem 4.6.

**Lemma 4.11.** Let  $\mathbf{A}$  denote  $\mathcal{A}$  or  $\mathcal{U}$ . If  $\mathcal{Z}'$  is a  $\Bbbk'$ -basis of  $\mathbf{A}'_{\Bbbk'}$ , then  $\mathcal{Z}$  is a  $\Bbbk$ -basis of  $\mathbf{A}$ .

*Proof.* Let  $\mathcal{Z}''$  be the set of the elements similar to those of  $\mathcal{Z}'$  in  $\mathcal{LP}_{\Bbbk''}$ . It is a  $\Bbbk''$ -basis of  $\mathbf{A}_{\Bbbk''}$  by Lemmas 4.9 and 4.10. Note that  $\mathbf{A}_{\Bbbk''} = \mathbf{A} \otimes \Bbbk''$  (Proposition 4.2). By Lemmas 4.8 and 4.3,  $\mathcal{Z}$  coincides with  $\mathcal{Z}''$  and it is a  $\Bbbk$ -basis of  $\mathbf{A}$ .

Lemma 4.9, Lemma 4.10, and Theorem 4.6 imply the following.

**Corollary 4.12.** (1) If  $\mathcal{A} = \mathcal{U} = \operatorname{Span}_{\Bbbk} \mathcal{Z}$ , then  $\mathcal{A}'_{\Bbbk'} = \mathcal{U}'_{\Bbbk'} = \operatorname{Span}_{\Bbbk'} \mathcal{Z}'$ .

(2) If the elements of Z have nice cluster decompositions in  $\mathbf{t}$ , so do those of Z' in  $\mathbf{t}'$ .

From now on, assume there is a  $\mathbb{Z}$ -algebra homomorphism  $\operatorname{var} : \mathcal{LP}_{\Bbbk''} \to \mathcal{LP}'_{\Bbbk'}$ such that  $\operatorname{var}(x_k) = x'_k \cdot p'_k$ ,  $\operatorname{var}(y_k) = y'_k$ ,  $\operatorname{var}(q^{\frac{1}{2d_{\rho}}}) = q^{\frac{1}{2d_{\rho}}}$ , for  $k \in I_{\operatorname{uf}}, p'_k \in \mathcal{P}'$ , and  $\operatorname{var}(\mathcal{P}) \subset \mathcal{P}'$ , called a variation map following [KQW23] (see [Fra16] for  $\Bbbk = \mathbb{Z}$ ).

Note that  $\mathcal{LP}_{\Bbbk''}$  is a free  $\mathcal{R}_{\Bbbk''}$ -bimodule via multiplication. Similarly,  $\mathcal{R}'_{\Bbbk'}$  is a  $\mathcal{R}'_{\Bbbk'}$ -bimodule, and thus an  $\mathcal{R}_{\Bbbk''}$ -bimodule via var. Introduce the left  $\mathcal{R}'_{\Bbbk'}$ -module homomorphism

(4.2) 
$$\varphi: \mathcal{R}'_{\Bbbk'} \otimes_{\mathcal{R}_{\Bbbk''}} \mathcal{LP}_{\Bbbk''} \simeq \mathcal{LP}'_{\Bbbk'},$$

such that  $\varphi(p' \otimes x^m) := p' * \operatorname{var}(x^m), \forall m \in \mathbb{Z}^I, p' \in \mathcal{P}'$ . Then  $\varphi$  is an isomorphism.

**Definition 4.13.**  $\varphi$  is called the base change from  $\mathcal{LP}_{\Bbbk''}$  to  $\mathcal{LP}'_{\Bbbk'}$ .

The multiplication on  $\mathcal{LP}'_{\Bbbk'}$  induces that on  $\mathcal{R}'_{\Bbbk'} \otimes_{\mathcal{R}_{\Bbbk''}} \mathcal{LP}_{\Bbbk''}$  such that,  $\forall h \in \mathbb{Z}^I$ ,  $r' \in \mathcal{P}', \ (p' \otimes x^m) * (r' \otimes x^h) = q^{\lambda'(\deg^{\mathbf{t}'}(\operatorname{var} x^m), \deg^{\mathbf{t}'} r')}(p' * r') \otimes (x^m * x^h)$ . Then  $\varphi$  is a  $\mathbb{Z}$ -algebra isomorphism. It is a  $\mathbb{K}$ -algebra isomorphism when  $\rho = 1$ .

Note that  $\mathcal{Z}' = \mathcal{P}' \cdot \operatorname{var}(\mathcal{Z}) = \{ [p' * \operatorname{var}(z)]^{\mathbf{t}'} \mid \forall z \in \mathcal{Z}, p' \in \mathcal{P}' \}$ , also denoted as  $\mathcal{Z}' = [\varphi(\mathcal{P}' \otimes_{\mathcal{P}} \mathcal{Z})]^{\mathbf{t}'}$ . We say that  $\mathcal{Z}'$  is obtained from  $\mathcal{Z}$  by the base change  $\varphi$ .

**Theorem 4.14.**  $\varphi$  restricts to a  $\mathbb{Z}$ -algebra isomorphism  $\mathcal{R}'_{\Bbbk'} \otimes_{\mathcal{R}_{\Bbbk''}} \mathcal{A}_{\Bbbk''} \simeq \mathcal{A}'_{\Bbbk'}$ .

*Proof.* Let  $p * z_1 * z_2 * \cdots * z_l$  denote any product of unfrozen cluster variables of  $\mathcal{A}$  and  $p \in \mathcal{P}$ . Take any  $p' \in \mathcal{P}'$ . Then  $\mathsf{var}(p' \otimes p * z_1 * \cdots * z_l) = p' * \mathsf{var}(p) * \mathsf{var}(z_1) * \cdots * \mathsf{var}(z_l)$ . It is contained in  $\mathcal{A}'$  since  $\mathsf{var}(z_i)$  are localized cluster monomials. Therefore,  $\varphi(\mathcal{R}'_{\Bbbk'} \otimes_{\mathcal{R}_{\Bbbk''}} \mathcal{A}_{\Bbbk''}) \subset \mathcal{A}'_{\Bbbk'}$ .

Conversely, take any  $p' \in \mathcal{P}'$  and any product  $z'_1 * \cdots * z'_l$  of unfrozen cluster variables in  $\mathcal{A}'$ . Let  $z_i$  denote the cluster variables of  $\mathcal{A}$  similar to  $z'_i$ . Then Theorem 4.6 implies that  $p' * z'_1 * \cdots * z'_l = q^{\alpha}r' * \varphi(1 \otimes z_1 * \cdots * z_l) = \varphi(q^{\alpha}r' \otimes z_1 * \cdots * z_l)$  for some  $r' \in \mathcal{P}', \alpha \in \frac{\mathbb{Z}}{2d_{\rho}}$ . Therefore,  $\mathcal{A}'_{\Bbbk'} \subset \varphi(\mathcal{R}'_{\Bbbk'} \otimes_{\mathcal{R}_{\Bbbk''}} \mathcal{A}_{\Bbbk''})$ .

**Theorem 4.15.** If  $\mathcal{Z}$  is a  $\Bbbk''$ -basis of  $\mathcal{U}_{\Bbbk''}$  or  $\mathcal{Z}'$  is a  $\Bbbk'$ -basis of  $\mathcal{U}'_{\Bbbk'}$ , then  $\varphi$  restricts to a  $\mathbb{Z}$ -algebra isomorphism  $\mathcal{R}'_{\Bbbk'} \otimes_{\mathcal{R}_{\Bbbk''}} \mathcal{U}_{\Bbbk''} \simeq \mathcal{U}'_{\Bbbk'}$ .

*Proof.* Note that  $\mathcal{Z}$  is a k-basis of  $\mathcal{U}$  and  $\mathcal{Z}'$  is a k'-basis of  $\mathcal{U}_{k'}$  by Proposition 4.2 and Lemmas 4.3, 4.10 and 4.11. The claim follows from the equality  $\mathcal{Z}' = \mathcal{P}' \cdot \mathsf{var}(\mathcal{Z})$ .

If  $\Bbbk = \mathbb{Z}$ , we obtain a base change of schemes  $\mathsf{Spec}\,\mathcal{U}' = \mathsf{Spec}\,\mathcal{R}' \times_{\mathsf{Spec}\,\mathcal{R}} \mathsf{Spec}\,\mathcal{U}$ .

Remark 4.16. Let **A** denote  $\mathcal{A}$  or  $\mathcal{U}$ . We claim that the construction of  $\varphi$ :  $\mathcal{R}'_{\Bbbk'} \otimes_{\mathcal{R}_{\Bbbk''}} \mathbf{A}_{\Bbbk''} \simeq \mathbf{A}'_{\Bbbk'}$  is independent of the choice of **t**. To see this, take any  $k \in I_{uf}$ , denote  $\mathbf{s} = \mu_k \mathbf{t}, \ \mathbf{s}' = \mu_k \mathbf{t}'$ . Then var induces a new variation map  $\begin{aligned} \mathsf{var}_{\mathbf{s}} &: \mathcal{LP}(\mathbf{s})_{\Bbbk''} \to \mathcal{LP}(\mathbf{s}')_{\Bbbk'} \text{ such that } \mathsf{var}_{\mathbf{s}} |_{\mathbf{A}(\mathbf{s})_{\Bbbk''}} \text{ coincides with the composition} \\ \mathbf{A}(\mathbf{s})_{\Bbbk''} \xrightarrow{\mu_k^*} \mathbf{A}(\mathbf{t})_{\Bbbk''} \xrightarrow{\mathsf{var}} \mathbf{A}(\mathbf{t}')_{\Bbbk'} \xrightarrow{(\mu_k^*)^{-1}} \mathbf{A}(\mathbf{s}')_{\Bbbk'}. \text{ For more details, see [KQW23, Section 3.3 and Proposition 3.1.4], for which we replace <math>\mathcal{U}$  by  $\mathbf{A}$  and drop the condition  $\mathbf{t}' \in \Delta_{\mathbf{t}}^+$  without loss of generality. We have  $\mu_k^* : \mathcal{R}'_{\Bbbk'} \otimes_{\mathcal{R}_{\Bbbk''}} \mathbf{A}(\mathbf{s})_{\Bbbk''} \xrightarrow{\sim} \mathcal{R}'_{\Bbbk'} \otimes_{\mathcal{R}_{\Bbbk''}} \mathbf{A}(\mathbf{t})_{\Bbbk''} \text{ such that } \mu_k^*(p' \otimes z) := \mu_k^*(p') \otimes \mu_k^*(z) = p' \otimes \mu_k^*(z). \end{aligned}$ 

Introduce  $\varphi_{\mathbf{s}} : \mathcal{R}'_{\Bbbk'} \otimes_{\mathcal{R}_{\Bbbk''}} \mathbf{A}(\mathbf{s})_{\Bbbk''} \simeq \mathbf{A}(\mathbf{s}')_{\Bbbk'}$  associated with  $\mathsf{var}_{\mathbf{s}}$ . Then we have  $\varphi_{\mathbf{s}} = (\mu_k^*)^{-1} \varphi \mu_k^*$ . Therefore,  $\varphi_{\mathbf{s}}$  is identified with  $\varphi$  via the mutation maps  $\mu_k^*$ .

Let **A** denote either  $\mathcal{A}$  or  $\mathcal{U}$ . Let  $\mathbf{t}^{\text{prin}}$  denote the seed of principal coefficients. For some choice  $\lambda_1$  of  $\lambda(\mathbf{t}^{\text{prin}})$  with  $\delta_k(\mathbf{t}^{\text{prin}}) = \delta_k(\mathbf{t})$ , the variation map  $\mathsf{var}_1 : \mathcal{LP}(\mathbf{t}^{\text{prin}}) \to \mathcal{LP}(\mathbf{t})$  exists. See Section **A** for details. Denote the associated  $\mathbf{A}(\mathbf{t}^{\text{prin}})$  by  $\mathbf{A}(\mathbf{t}^{\text{prin}}, \lambda_1)$ . Similarly, the variation map  $\mathsf{var}_2 : \mathcal{LP}(\mathbf{t}^{\text{prin}}) \to \mathcal{LP}(\mathbf{t}')$  exists for some choice  $\lambda_2$  of  $\lambda(\mathbf{t}^{\text{prin}})$  with with  $\delta_k(\mathbf{t}^{\text{prin}}) = \delta_k(\mathbf{t}')$ .

If  $\mathbf{A} = \mathcal{A}$ ,  $\mathbf{A}$  and  $\mathbf{A}'$  are related by base changes  $\varphi_1, \varphi_2$  and a quantization change, where the squares associated with base changes are commutative:

$$(4.3) \qquad \begin{array}{ccc} \mathbf{A} & \stackrel{\varphi_1}{\longleftarrow} & \mathbf{A}(\mathbf{t}^{\mathrm{prin}}, \lambda_1) & \stackrel{\mathrm{quantization}}{\sim} & \mathbf{A}(\mathbf{t}^{\mathrm{prin}}, \lambda_2) & \stackrel{\varphi_2}{\longrightarrow} & \mathbf{A}' \\ \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \\ \mathcal{R} & \stackrel{\varphi_1}{\longleftarrow} & \mathcal{R}(\mathbf{t}^{\mathrm{prin}}, \lambda_1) & \stackrel{\mathrm{quantization}}{\underset{\mathrm{change}}{\sim}} & \mathcal{R}(\mathbf{t}^{\mathrm{prin}}, \lambda_2) & \stackrel{\varphi_2}{\longrightarrow} & \mathcal{R}' \end{array}$$

Note that the quantum seeds  $(\mathbf{t}^{\text{prin}}, \lambda_1)$  and  $(\mathbf{t}^{\text{prin}}, \lambda_2)$  are similar with  $\rho \in \mathbb{Q}$ . If we want to replace the middle square by a base change, we need to extend the algebras over  $\Bbbk''$  and  $\Bbbk'$  respectively. More precisely, we have the  $\mathbb{Z}$ -algebra isomorphism var :  $\mathcal{LP}(\mathbf{t}^{\text{prin}}, \lambda_1)_{\Bbbk''} \simeq \mathcal{LP}(\mathbf{t}^{\text{prin}}, \lambda_2)_{\Bbbk'}$ , sending  $q^{\alpha}x^m$  to  $q^{\alpha\rho}x^m$  such that var  $\mathcal{R}(\mathbf{t}^{\text{prin}}, \lambda_1)_{\Bbbk''} = \mathcal{R}(\mathbf{t}^{\text{prin}}, \lambda_2)_{\Bbbk'}$ . It induces the base change  $\varphi : \mathcal{R}'(\mathbf{t}^{\text{prin}}, \lambda_2)_{\Bbbk'} \otimes_{\mathcal{R}(\mathbf{t}^{\text{prin}}, \lambda_1)_{\Bbbk''}} \mathbf{A}(\mathbf{t}^{\text{prin}}, \lambda_1)_{\Bbbk''} \simeq \mathbf{A}(\mathbf{t}^{\text{prin}}, \lambda_2)_{\Bbbk'}$  between  $\mathbb{Z}$ -algebras. When  $\mathbf{A} = \mathcal{U}$ , we still have (4.3) under the following assumptions:

- (1) A has a k-basis  $\mathbb{B}$ , or  $\mathbf{A}(\mathbf{t}^{\text{prin}}, \lambda_1)$  has a k-basis  $\mathbb{B}_1$ , such that it consists of pointed elements and is closed under the  $\mathcal{P}$  commutative multiplication.
- (2) The similar assumption for  $\mathbf{A}'$  and  $\mathbf{A}(\mathbf{t}^{\text{prin}}, \lambda_2)$ .

Under the above assumptions, **A** and  $\mathbf{A}(\mathbf{t}^{\text{prin}}, \lambda_1)$  have bases  $\mathbb{B}$ ,  $\mathbb{B}_1$  respectively, which are related by a base change:  $\mathbb{B} = \mathcal{P} \cdot \text{var}_1(\mathbb{B}_1) = [\varphi_1(\mathcal{P} \otimes_{\mathcal{P}(\mathbf{t}^{\text{prin}}, \lambda_1)} \mathbb{B}_1)]$ . Similar, **A'** and  $\mathbf{A}(\mathbf{t}^{\text{prin}}, \lambda_2)$  have bases  $\mathbb{B}'$ ,  $\mathbb{B}_2$  related by a base change. Sometimes, we could choose  $\mathbb{B}_1$  and  $\mathbb{B}_2$  related by a quantization change:  $\mathbb{B}_1|_{q^{\frac{1}{2}} \mapsto q^{\frac{1}{2d_{\rho}}}} = \mathbb{B}_2$ , see Section 4.4.

4.3. Localization. Let  $\mathbf{A}$  denote either  $\overline{\mathcal{A}}$  or  $\overline{\mathcal{U}}$ . Assume it has a k-basis  $\mathcal{Z}$  whose elements are pointed in  $\mathcal{LP}$  and closed under the  $\overline{\mathcal{P}}$  commutative multiplication: we have  $p \cdot z \in \mathcal{Z}, \forall p \in \overline{\mathcal{P}}, z \in \mathcal{Z}$ . Let  $\widetilde{\mathbf{A}}$  denote the localization  $\mathbf{A}[x_j^{-1}]_{j \in I_{\mathbf{f}}}$ .

Similar to the previous discussion, we have an isomorphism  $\varphi : \mathcal{R} \otimes_{\overline{\mathcal{R}}} \overline{\mathcal{LP}} \xrightarrow{\sim} \mathcal{LP}$ , still called a base change. Note that  $\widetilde{\mathcal{Z}} := \mathcal{P} \cdot \mathcal{Z} = \{p \cdot z | p \in \mathcal{P}, z \in \mathcal{Z}\}$  is a basis of  $\widetilde{\mathbf{A}}$ , denoted  $\widetilde{\mathcal{Z}} = [\varphi(\mathcal{P} \otimes_{\overline{\mathcal{P}}} \mathcal{Z})]$ . We deduce that the base change  $\varphi$  restricts to an isomorphism  $\mathcal{R} \otimes_{\overline{\mathcal{R}}} \mathbf{A} \simeq \widetilde{\mathbf{A}}$ .

4.4. **Triangular bases.** As before, given  $\mathcal{Z} \subset \mathcal{LP}$  whose elements are pointed. Let  $\mathcal{Z}'$  be the subset of  $\mathcal{LP}'_{k'}$  consisting of the elements similar to those of  $\mathcal{Z}$ . **Lemma 4.17.** Let  $\mathbf{A}$  denote  $\mathcal{A}$  or  $\mathcal{U}$ . If  $\mathcal{Z}$  is the triangular basis of  $\mathbf{A}$  with respect to the seed  $\mathbf{s} := \mu \mathbf{t}$  for some  $\mu$ , then  $\mathbf{A}'$  has the triangular basis with respect to  $\mathbf{s}' := \mu \mathbf{t}'$ , and it equals  $\mathcal{Z}'$ .

*Proof.* By Lemma 4.10,  $\mathcal{Z}'$  is a  $\Bbbk'$ -basis of  $\mathcal{U}'_{\Bbbk'}$ .  $\mathcal{Z}'$  is bar-invariant and contains the cluster monomials in  $\mathbf{s}' = \boldsymbol{\mu} \mathbf{t}', \, \mathbf{s}'[1] = (\boldsymbol{\mu} \mathbf{t}')[1]$ . Let us check the triangularity condition (last condition).

Take any *m*-pointed  $z_m \in \mathcal{Z}$  and similar *m'*-pointed  $z'_{m'} \in \mathcal{Z}'$ . We can write  $z_m = q^{\alpha}x^m * F_m$  and  $z'_{m'} = q^{\alpha'}(x')^{m'} * F'_{m'}$ , where  $q^{\alpha}, q^{\alpha'}$  are determined by the bar-invariance,  $\operatorname{pr}_{I_{uf}} m = \operatorname{pr}_{I_{uf}} m'$ ,  $F_m \in \mathbb{k}[N_{uf}^{\oplus}]$ , and  $\operatorname{var}(F_m) = F'_{m'}$  under the map  $\operatorname{var} : \mathbb{k}[N_{uf}^{\oplus}] \to \mathbb{k}'[(N'_{uf})^{\oplus}]$  sending  $y_k$  to  $y'_k$ ,  $q^{\frac{1}{2}}$  to  $q^{\frac{\rho}{2}}$ . By [Qin17, Lemma 4.2.2],  $\mu \mathbf{t}$  and  $\mu \mathbf{t}'$  are similar. Moreover, as elements in  $\widehat{\mathcal{LP}}(\mathbf{s})$  and  $\widehat{\mathcal{LP}}(\mathbf{s}')_{\mathbb{k}'}$  respectively,  $x^m$  and  $(x')^{m'}$  are similar, and  $F_m$  and  $F'_{m'}$  are similar. Then Theorem 4.6 implies that  $z_m$  and  $z'_{m'}$  are similar as elements in  $\mathcal{LP}(\mathbf{s})$  and  $\mathcal{LP}(\mathbf{s'})_{\mathbb{k}'}$  respectively.

Since  $[x_i(\mathbf{s}) * z_m]^{\mathbf{s}}$  has a  $(\prec_{\mathbf{s}}, q^{-\frac{1}{2}} \mathbb{Z}[q^{-\frac{1}{2}}])$ -decomposition into  $\mathcal{Z}$  in  $\mathcal{LP}(\mathbf{s}), [x_i(\mathbf{s}') * z'_{m'}]^{\mathbf{s}'}$  has a  $(\prec_{\mathbf{s}'}, q^{-\frac{\rho}{2}} \mathbb{Z}[q^{-\frac{\rho}{2}}])$ -decomposition into  $\mathcal{Z}'$  in  $\mathcal{LP}(\mathbf{s}')_{\mathbb{k}'}$  by Theorem 4.6.

Therefore,  $\mathcal{Z}'$  is the triangular basis for  $\mathbf{A}'_{\Bbbk'}$  with respect to  $\mathbf{s}'$  over  $\Bbbk'$ . We claim that  $\mathcal{Z}' \subset \mathcal{LP}(\mathbf{s}')$ . If so,  $\mathcal{Z}'$  is a  $\Bbbk$ -basis of  $\mathbf{A}$  by Lemma 4.3.

Note that any  $z' \in \mathcal{Z}'$  has a  $(\prec_{\mathbf{s}}, q^{-\frac{\rho}{2}}\mathbb{Z}[q^{-\frac{\rho}{2}}])$ -decomposition into  $[x(\mathbf{s}')^{g_{+}} * x(\mathbf{s}'[1])^{g_{-}}]^{\mathbf{s}'}$ , for  $g_{+} \in \mathbb{N}^{I}$ ,  $g_{-} \in \mathbb{N}^{I_{\mathrm{uf}}}$ , see [Qin17, Lemmas 6.2.1(iii) and 3.1.11]. Since  $[x(\mathbf{s}')^{g_{+}} * x(\mathbf{s}'[1])^{g_{-}}]^{\mathbf{s}'}$  is contained in  $\mathcal{LP}(\mathbf{s}')$  and  $\mathcal{Z}'$  is bar-invariant, we obtain  $z' \in \widehat{\mathcal{LP}}(\mathbf{s}')$ . The desired claim follows.

Lemma 4.17 implies the following result.

**Proposition 4.18.** If Z is the common triangular basis for U, then the common triangular basis for U' exists and it equals Z'.

#### 5. Based cluster algebras and monoidal categorifications

**Definition 5.1.** Let A denote  $\overline{\mathcal{A}}$ ,  $\mathcal{A}$ ,  $\overline{\mathcal{U}}$ , or  $\mathcal{U}$ , and  $\mathbb{B}$  its k-basis satisfying:

- (1)  $\mathbb{B}$  contains all the cluster monomials.
- (2)  $\forall b \in \mathbb{B}, j \in I_{f}, x_{j} \cdot b \in \mathbb{B}.$
- (3) At the quantum level, the elements of  $\mathbb{B}$  are invariant under the bar involution  $q^{\frac{1}{2}} \mapsto q^{-\frac{1}{2}}$ .
- (4) (Homogeneity) Each  $b \in \mathbb{B}$  is contained in some  $x^m \cdot \Bbbk[N_{uf}]$ .

The pair  $(\mathbf{A}, \mathbb{B})$  is called a based cluster algebra.

We call  $(\mathbf{A}, \mathbb{B})$   $\Theta$ -pointed if  $\mathbb{B}$  is, where  $\Theta$  is a subset of  $M^{\circ}(\mathbf{t})$  or  $\mathcal{M}^{\circ}$ .

Localization of  $(\mathbf{A}, \mathbb{B})$  produces a based cluster algebra  $(\mathbf{A}[x_j^{-1}]_{j \in I_f}, \mathbb{B}')$  where  $\mathbb{B}' = \{p \cdot b | p \in \mathcal{P}, b \in \mathbb{B}\}.$ 

5.1. Monoidal categories. Let  $\mathbb{F}$  be a field. Let  $\mathcal{T}$  denote a  $\mathbb{F}$ -linear tensor category in the sense of [KKK18, Section A.1]. In particular,  $\mathcal{T}$  is endowed with an exact bifunctor () $\otimes$ () called the tensor functor, its objects have finite lengths, and its Grothendieck ring  $K_0(\mathcal{T})$  is unital and associative. Let [X] denote the isoclass of  $X \in \mathcal{T}$ . We have  $[X \otimes Y] = \sum_{S \in \mathsf{Simp}} c_{XY}^S[S]$  for  $X, Y \in \mathcal{T}$  and  $c_{XY}^S \in \mathbb{N}$ . Recall that the multiplication of  $K_0(\mathcal{T})$  is induced from the tensor product  $\otimes$ . Its unit is

[1] for some unit object 1. Let Simp denote the set of simple objects of  $\mathcal{T}$ . Simple objects are often referred to as simples.

When  $\mathbb{k} = \mathbb{Z}$ , we assume  $K_0(\mathcal{T})$  is commutative. Denote  $K := K_0(\mathcal{T})$ . When  $\mathbb{k} = \mathbb{Z}[q^{\pm \frac{1}{2}}]$ , we make one of the following assumptions.

- (1) We assume that  $\mathcal{T}$  has an object Q satisfying [KKKO18, (6.2)(ii)(iii)]. In particular,  $\mathbf{q} := Q \otimes ()$  is an auto-equivalence functor, such that  $[\mathbf{q}(X \otimes Y)] = [(\mathbf{q}X) \otimes Y] = [X \otimes (\mathbf{q}Y)]$ . We denote  $X \sim Y$  if  $X \simeq \mathbf{q}^d(Y)$  for some  $d \in \mathbb{Z}$ . Then  $K_0(\mathcal{T})$  is a  $\mathbb{Z}[q^{\pm}]$ -algebra such that  $q^{\pm}[X] := [\mathbf{q}^{\pm}X]$ , where q is a formal parameter. Denote  $K := K_0(\mathcal{T}) \otimes \mathbf{k}$ .
- (2) Or we assume that the k-space  $K := K_0(\mathcal{T}) \otimes_{\mathbb{Z}} \mathbb{k}$  has a q-twisted product \* such that, for  $X, Y \in \mathcal{T}$ , its multiplication satisfies  $[X] * [Y] = \sum_{S \in \mathsf{Simp}} c(q^{\frac{1}{2}})_{XY}^S[S]$ , where  $c(q^{\frac{1}{2}})_{XY}^S \in \mathbb{N}[q^{\pm \frac{1}{2}}]$  and  $c(1)_{XY}^S = c_{XY}^S$ .

The k-algebra K will be called the (deformed) Grothendieck ring of  $\mathcal{T}$ .

Remark 5.2. Assume that there is an order 2 anti-automorphism  $\overline{()}$  on K such that  $\overline{[S]} \sim [S]$  for any simple S. Take another simple V such that  $V \otimes S$  is simple. We claim that [S] and [V] q-commute,  $S \otimes V$  is simple, and  $S \otimes V \sim V \otimes S$ .

To see this, denote  $[V \otimes S] = [V] * [S] = [W]$  for some simple W. Applying  $\overline{()}$ , we obtain  $[S \otimes V] = [S] * [V] \sim [W]$ . The claim follows.

5.2. Categorifications of based cluster algebras. We will use the following notion of monoidal categorification, which is weaker than the one in [HL10].

**Definition 5.3.** We say  $\mathcal{T}$  admits a cluster structure **A** if the following holds.

- (1) There exists a k-algebra embedding  $\kappa : \mathbf{A} \hookrightarrow K$ .
- (2) For any cluster monomial z, there exists a simple S such that  $\kappa(z) \sim [S]$ .

We further say  $\mathcal{T}$  categorifies **A** if  $\kappa$  is isomorphic. We say  $\mathcal{T}$  categorifies **A** after localization if  $\kappa$  extends to  $\kappa : \mathbf{A}[x_j^{-1}]_{j \in I_{\mathbf{f}}} \simeq K[\kappa(x_j)^{-1}]_{j \in I_{\mathbf{f}}}$ . We can omit the symbol  $\kappa$  in this case.

**Definition 5.4.** We say  $\mathcal{T}$  categorifies  $(\mathbf{A}, \mathbb{B})$  if the following holds.

- (1)  $\mathcal{T}$  categorifies **A**.
- (2) For any simple object  $S, q^{\alpha}[S] \in \kappa \mathbb{B}$  for some  $\alpha$ .

Categorification after localization is defined in a similar manner. In addition, the notion of categorification for cluster algebras defined over the any extension of k can also be similarly established.

**Definition 5.5.** We say  $\mathcal{T}$  quasi-categorifies a based cluster algebra  $(\mathbf{A}', \mathbb{B}')$  if there exists  $(\mathbf{A}, \mathbb{B})$ , which is related to  $(\mathbf{A}', \mathbb{B}')$  by a finite sequence of base changes and quantization changes (see (4.3), section 4.3), such that  $\mathcal{T}$  categorifies  $(\mathbf{A}, \mathbb{B})$ after localization.

Note that, in Definition 5.5, when the elements of  $\mathbb{B}$  are pointed elements in some  $\mathcal{LP}(\mathbf{t})$  up to  $q^{\frac{\mathbb{Z}}{2}}$ -multiples, the structure constants of  $\mathbb{B}'$  are determined from those for  $\mathbb{B}$ , see Theorem 4.6.

Now assume that  $\mathcal{T}$  categories  $(\mathbf{A}, \mathbb{B})$ . For any cluster variables  $x_i(\mathbf{t})$ , choose a simple object  $S_i(\mathbf{t})$  such that  $q^{\alpha}[S_i(\mathbf{t})] = \kappa(x_i(\mathbf{t}))$ . We will often omit the symbols  $\kappa$  and  $\mathbf{t}$  for simplicity. Let us present some immediate consequences.

Apply the bar-involution  $\overline{(\ )}$  in **A** to  $[S], S \in \mathsf{Simp}$ , which is the identity map at the classical level. We obtain  $\overline{[S]} \sim [S]$ .

The simple objects  $S_j$ ,  $j \in I_f$ , are said to be frozen. Since  $q^{\frac{Z}{2}}\mathbb{B}$  is closed under multiplication by the frozen variables  $x_j$ , for any simple object S,  $S_j \otimes S$  and  $S \otimes S_j$  remain simple.

Finally, the structure constants of  $\mathbb{B}$  are non-negative.

Remark 5.6. If  $\mathbb{B}$  has positive structure constants, we can construct a semisimple monoidal category  $\mathcal{T}$  as below. The set of simple objects of  $\mathcal{T}$  is  $q^{\mathbb{Z}}\mathbb{B}$ . For any simple objects  $S, S' \in \mathcal{T}$ , we define their tensor product by multiplication:  $S \otimes S' := \bigoplus_j S_j^{c_j}$ whenever  $S * S' = \sum_{S_j \in q^{\mathbb{Z}} \mathbb{B}} c_j S_j, c_j \in \mathbb{N}$ , as elements in **A**. Then  $\mathcal{T}$  categorifies (**A**,  $\mathbb{B}$ ). Usually, we desire categories with richer structures than  $\mathcal{T}$ .

5.3. Triangular bases. Work at the quantum case  $\mathbb{k} = \mathbb{Z}[q^{\pm \frac{1}{2}}]$ .

**Condition (C)** The category  $\mathcal{T}$  is said to satisfy the commuting condition, or Condition (C), with respect to a simple X if, for any simple S,  $X \otimes S$  and  $S \otimes X$  are simple whenever [X] and [S] q-commute.

Now, assume we have an inclusion  $K \subset \mathcal{LP}(\mathbf{t})$  such that, for any simple S,  $\overline{[S]} \sim [S]$  (see also Remark 5.2). Further assume that for each simple  $S \in \mathcal{T}$ ,  $q^{\beta}S$  is pointed in  $\mathcal{LP}(\mathbf{t})$  for some  $\beta \in \frac{1}{2}\mathbb{Z}$ . Moreover,  $\deg S \neq \deg S'$  whenever  $S \nsim S'$  for simples S, S'.

**Condition (T)** The category  $\mathcal{T}$  is said to satisfy the  $\prec_t$ -triangularity condition, or Condition (T), with respect to a simple X if we have the following decomposition in K for any simple S:

(5.1) 
$$[X] * [S] = \sum_{j=0}^{s} c_j q^{\alpha_j} q^{\beta_j} [S_j],$$

where  $S_j$  are simple,  $\overline{q^{\beta_j}[S_j]} = q^{\beta_j}[S_j]$ ,  $c_0 = 1$ ,  $c_j \in \mathbb{N}$ , and  $\alpha_j < \alpha_0$  for j > 0.

Remark 5.7. We claim that Condition (T) implies Condition (C) with respect to X. To see this, denote  $[X] * [S] = q^{\gamma}[S] * [X]$ . Then  $[X] * [S] = q^{\gamma}[\overline{X}] * [S]$ . Decomposing [X] \* [S] as in (5.1), we deduce that  $q^{\gamma}\overline{c_j} = c_j$ . So  $\gamma = 2\alpha_0$  and  $c_j = 0$  for j > 0. Therefore,  $X \otimes S$  is simple. By Remark 5.2,  $S \otimes X$  is simple too.

**Theorem 5.8.** Take  $\mathbb{k} = \mathbb{Z}[q^{\pm \frac{1}{2}}]$  and  $\mathbf{A} = \mathcal{A}$  or  $\mathcal{U}$ . Assume that  $\mathbf{A}$  is injectivereachable,  $\mathcal{T}$  categorifies  $(\mathbf{A}, \mathbb{B})$  after localization, and the elements of  $\mathbb{B}$  are pointed at distinct degrees in  $\mathcal{LP}(\mathbf{t})$  for some seed  $\mathbf{t}$ . If  $\mathcal{T}$  satisfies the triangularity condition with respect to  $S_k(\mathbf{t})$ ,  $k \in I_{uf}$ , then  $\mathbb{B}$  is the common triangular basis of  $\mathbf{A}$ .

*Proof.* Note that  $\mathcal{T}$  always satisfy the Condition (T) with respect to the frozen simples  $S_j, j \in I_f$ .

By definition of based cluster algebras,  $\mathbb{B}$  contains all the cluster monomials, including those in **t** and **t**[1]. Then the Condition (T) with respect to  $S_i(\mathbf{t})$ ,  $i \in I$ , implies that  $\mathbb{B}$  is the triangular basis with respect to **t**. Since this triangular basis contains all the cluster monomials, it is the common triangular basis by [Qin20b, Proposition 6.4.3, Theorem 6.5.3].

Remark 5.9. Condition (T) holds for finite dimensional module categories of KLR algebras for symmetric C with respect to the real simple modules, see [KKKO18,

Corollary 3.8]. In type ADE, it also holds for finite dimensional modules categories of quantum affine algebras with respect to the cluster variables, see [Qin17].

Conversely, if  $\mathbb{B}$  is the triangular basis with respect to any seed  $\mathbf{t}$ , then Condition (T) holds with respect to the simples corresponding to the cluster monomials in  $\mathbf{t}$ .

5.4. Monoidal subcategories. Work at the quantum case  $\mathbb{k} = \mathbb{Z}[q^{\pm \frac{1}{2}}]$ . Let **t** denote a seed of  $\mathcal{A}$ . Assume we have an inclusion of  $\mathbb{k}$ -algebras  $K \hookrightarrow \mathcal{LP}(\mathbf{t})$  such that  $q^{\alpha_k}[S_i] = x_i$  for some simples  $S_i = S_i(\mathbf{t})$ , where  $i \in I$ , and  $\alpha_i \in \frac{1}{2}\mathbb{Z}$ . Assume  $\overline{|S|} \sim |S|, \forall S \in \text{Simp.}$ 

**Proposition 5.10** ([HL10, Proposition 2.2.(i)]). For any object V of  $\mathcal{T}$ , the Laurent expansion of [V] in  $\mathcal{LP}(\mathbf{t})$  have non-negative coefficients.

Choose any  $F \subset I_{uf}$  and introduce the following set:

(5.2)  $\operatorname{Simp}' := \{ S \in \operatorname{Simp} | [S_k] \text{ } q \text{-commutes with } [S], \forall k \in F \}.$ 

If  $\mathcal{T}$  satisfies Condition (C) with respect to  $S_k, k \in F$ , we have

 $\operatorname{Simp}' = \{ S \in \operatorname{Simp} | S_k \otimes S \text{ is simple}, \forall k \in F \}.$ 

Let  $\mathcal{T}'$  denote the full-subcategory of  $\mathcal{T}$  whose objects have composition factors in Simp'. Then  $\mathcal{T}'$  is closed under taking biproducts, extensions, kernels, and quotients. In particular,  $\mathcal{T}'$  is an abelian subcategory. If  $\mathcal{T}$  is equipped with the auto-equivalent functor  $Q \otimes ()$  as in Section 5.1,  $\mathcal{T}'$  is closed under this autoequivalence. The (deformed) Grothendieck group of  $\mathcal{T}'$ , denoted K', is defined to be the submodule of K spanned by  $\{[S]|S \in \mathsf{Simp}'\}$ .

**Lemma 5.11.** For any  $k \in F$  and object  $V \in \mathcal{T}$ , if [V] q-commutes with  $[S_k]$ , then [W] q-commutes with  $S_k$  for every subquotient W of V.

*Proof.* We have  $[S_k] * [V] = q^{\alpha}[V] * [S_k]$  for some  $\alpha \in \mathbb{Z}$ . Consider the corresponding Laurent polynomials in  $\mathcal{LP}(\mathbf{t})$  and note that  $[S_k] \sim x_k$ . Then all Laurent monomials  $x^m$  appearing in [V] satisfies  $x^m * x_k = q^{\alpha} x_k * x^m$ .

We have  $[V] = \sum b_i[V_i]$  for  $b_i \in \mathbb{N}$ ,  $V_i$  composition factors of V. By Proposition 5.10,  $[V_i]$  have positive Laurent expansions. So the Laurent monomials appearing in  $[V_i]$  already appear in [V]. Therefore,  $x_k * [V_i] = q^{\alpha}[V_i] * x_k$ . The claim follows.

# **Lemma 5.12.** The subcategory $\mathcal{T}'$ is closed under the tensor product.

Proof. For any two simples V, W in  $\mathcal{T}'$ ,  $[V \otimes W] = [V] * [W]$  q-commutes with  $S_k, k \in F$ . By Lemma 5.11, all composition factors of  $V \otimes W$  q-commute with  $S_k$ . Therefore,  $V \otimes W$  remains in  $\mathcal{T}'$ . In general, for any object V' and W' in  $\mathcal{T}'$ , we have  $[V'] = \sum b_i[V_i]$  and  $[W'] = \sum c_j[W_j], b_i, c_j \in \mathbb{N}$ , for composition factors  $V_i, W_j$ . Note that  $[V' \otimes W'] = [V'] * [W'] = \sum b_i c_j [V_i] * [W_j] = \sum b_i c_j [V_i \otimes W_j]$ . Any composition factor of  $V' \otimes W'$  must be a composition factor of some  $V_i \otimes W_j$ , which we have seen is contained in  $\mathcal{T}'$ . So  $V' \otimes W'$  belongs to  $\mathcal{T}'$ .

Remark 5.13. Any cluster monomial z of  $\overline{\mathcal{A}}(\mathfrak{f}_F \mathbf{t})$  q-commutes with  $x_k, \forall k \in F$ . So any simple  $S \in \mathcal{T}$  such that  $[S] \sim z$  belongs to  $\mathcal{T}'$ . Therefore, if  $\mathcal{T}$  admits a cluster structure  $\overline{\mathcal{A}}, \mathcal{T}'$  admits a cluster structure  $\overline{\mathcal{A}}(\mathfrak{f}_F \mathbf{t})$ .

By Proposition 5.10, the set of simples is a positive  $\mathbb{N}$ -basis of  $K \subset \mathcal{LP}(\mathbf{t})$  that has positive Laurent expansions. Therefore, it has the following stabilization property, see Lemma 3.21.

**Proposition 5.14.** Take any object V in  $\mathcal{T}$  and  $i \in I$ . Consider the decomposition  $[S_i^{\otimes d} \otimes V] = \sum b_j^{(d)}[V_j^{(d)}]$  where  $V_j^{(d)}$  are composition factors and  $b_j^{(d)} \in \mathbb{N}$ . When d is large enough,  $x_i^{-d} * [V_j^{(d)}]$  and  $b_j^{(d)}$  stabilize, and  $V_j^{(d)} \in \mathcal{T}'$ .

5.5. Categorifications via monoidal subcategories. For the moment, assume  $\mathcal{T}$  categorifies  $(\mathbf{A}, \mathbb{B})$  after localization and it has the triangularity property with respect to  $S_k = S_k(\mathbf{t}), \ k \in F$ . Assume  $\mathbb{B} = \{b_m | m \in M^\circ(\mathbf{t})\}$  with *m*-pointed  $b_m$  in  $\mathcal{LP}(\mathbf{t})$ . If  $\kappa b_m \sim [S]$  for some simple S, we choose a simple  $S_m$  such that  $\kappa b_m \sim [S_m]$ . Omit  $\kappa$  for simplicity.

**Lemma 5.15.** When  $d \in \mathbb{N}$  is large enough, we have  $[S_{m+df_i}] * [S_k] \sim [S_{m+(d+1)f_i}]$ .

*Proof.* Note that  $[S_{m+df_i}]$  is a term appearing in  $[S_k]^d * [S_m]$ . So, when d is large enough, the claim follows from the stabilization property, see Proposition 5.14.

 $\Box$ 

**Theorem 5.16.** For any simple  $S_m \in \mathcal{T}$ , the freezing operator acts on  $[S_m]$  by

$$\mathfrak{f}_F[S_m] \sim x^{-d\sum_{k \in F} f_k} * [S_{m+d\sum_{k \in F} f_k}]$$

for  $d \in \mathbb{N}$  large enough. In this case,  $S_{m+d\sum_{k\in F} f_k} \in \mathcal{T}'$ .

*Proof.* By Lemma 5.15,  $S_{m+d\sum_{k\in F} f_k} \in \mathcal{T}'$ . The claim follows from Lemma 3.23.

**Theorem 5.17.** If  $\mathbb{B}$  is  $\mathcal{M}^{\circ}$ -pointed and  $\mathfrak{f}_{F}\mathbf{t}$  is injective-reachable, then  $\mathcal{T}'$  categorifies  $(\mathcal{U}(\mathfrak{f}_{F}\mathbf{t}), \mathfrak{f}_{F}\mathbb{B})$  after localization.

*Proof.* For any  $S_m \in \text{Simp}' \subset \mathcal{T}'$ , we have  $b_m = \mathfrak{f}_F b_m$ , which is contained in  $\mathcal{U}(\mathfrak{f}_F \mathbf{t})$ , see Proposition 3.8. Therefore,  $K' \subset \mathcal{U}(\mathfrak{f}_F \mathbf{t})$ .

Since  $\mathbb{B}$  is  $\mathcal{M}^{\circ}$ -pointed,  $\mathfrak{f}_{F}\mathbb{B}$  is a  $\mathcal{M}^{\circ}$ -pointed basis of  $\mathcal{U}(\mathfrak{f}_{F}\mathbf{t})$  by Theorem 3.11. Take any  $b_{m}$  such that  $b_{m} \sim [S_{m}]$ . By Theorem 5.16,  $\mathfrak{f}_{F}b_{m} = x^{-d\sum_{k\in F}f_{k}} \cdot q^{\alpha}[S_{m+d\sum_{k\in F}f_{k}}]$  for  $d\in\mathbb{N}$  large enough,  $\alpha\in\mathbb{Z}_{2}$ , and  $S_{m+d\sum_{k\in F}f_{k}}\in\mathsf{Simp}'\subset\mathcal{T}'$ . We deduce that  $\mathcal{U}(\mathfrak{f}_{F}\mathbf{t})\subset K'[[S_{i}]^{-1}]_{i\in I_{\mathsf{f}}\sqcup F}$ .

From now on, we drop the assumptions in the beginning of Section 5.5.

**Theorem 5.18.** Let  $(\overline{\mathcal{U}}, \mathbb{B})$  be a based cluster algebra such that  $\mathbb{B}$  is  $M^{\circ}(\mathbf{t})$ -pointed in  $\mathcal{LP}(\mathbf{t})$ . Assume that  $\mathbb{B}' := \mathbb{B} \cap \overline{\mathcal{U}}(\mathfrak{f}_{F}\mathbf{t})$  is a basis for  $\overline{\mathcal{U}}(\mathfrak{f}_{F}\mathbf{t})$ . If  $\mathcal{T}$  categorifies  $(\overline{\mathcal{U}}, \mathbb{B})$ , then  $\mathcal{T}'$  categorifies  $(\overline{\mathcal{U}}(\mathfrak{f}_{F}\mathbf{t}), \mathbb{B}')$ .

*Proof.* Since  $\mathbb{B}'$  contains all cluster monomials of  $\overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t})$ , it remains to show  $K' = \overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t})$ .

 $\forall S \in \text{Simp}' \subset \mathcal{T}'$ , since [S] q-commute with  $x_k, k \in F$ , we have  $\text{supp}[S] \cap F = \emptyset$ . Then Lemma 2.13 implies  $\nu_k([S]) \geq 0$ ,  $\forall k \in F$ . Note that  $\nu_j([S]) \geq 0$  for  $j \in I_{\mathsf{f}}$ since  $[S] \in \overline{\mathcal{U}}$ . Combining with  $[S] = \mathfrak{f}_F[S] \in \mathcal{U}(\mathfrak{f}_F \mathbf{t})$ , we obtain  $[S] \in \overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t})$ . We deduce that  $K' \subset \overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t})$ .

We have  $\text{Span}_{\Bbbk}\{[S]|S \in \text{Simp}, [S] \in \overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t})\} = \text{Span}_{\Bbbk}(\mathbb{B} \cap \overline{\mathcal{U}})$ , which coincides with  $\overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t})$  by our assumption. Take any  $S \in \text{Simp}$  such that  $[S] \in \overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t})$ . Then [S] q-commutes with  $x_k, k \in F$ , since  $x_k$  are frozen variables in  $\overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t})$  and [S] is pointed in  $\mathcal{LP}(\mathfrak{f}_F \mathbf{t})$ . So we obtain  $S \in \text{Simp}' \subset \mathcal{T}'$ . We deduce that  $\overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t})$  is contained in K'.

Now assume  $\mathcal{U}$  has the common triangular basis  $\mathbf{L}$ . Then  $\mathcal{U}(\mathfrak{f}_F \mathbf{t})$  has the common triangular basis  $\mathbf{L}' = \mathfrak{f}_F \mathbf{L}$  (Theorem 3.17). Assume that  $\mathbf{t}$  and  $\mathfrak{f}_F \mathbf{t}$  can be optimized. Then  $\overline{\mathbf{L}} := \mathbf{L} \cap \overline{\mathcal{U}}$  is a basis of  $\overline{\mathcal{U}}$  and  $\overline{\mathbf{L}}' = \mathbf{L}' \cap \overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t})$  is a basis of  $\overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t})$ .

**Corollary 5.19.** If  $\mathcal{T}$  categorifies  $(\overline{\mathcal{U}}, \overline{\mathbf{L}})$ , then  $\mathcal{T}'$  categorifies  $(\overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t}), \overline{\mathbf{L}}')$ .

*Proof.* Proposition 3.3 implies  $\overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t}) \subset \overline{\mathcal{U}}(\mathbf{t})$ . Corollary 3.19(2) implies that the basis  $\overline{\mathbf{L}}' = \mathfrak{f}_F \mathbf{L} \cap \overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t})$  equals  $\mathbf{L} \cap \overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t})$ , which can be written as  $\mathbf{L} \cap (\overline{\mathcal{U}} \cap \overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t})) = (\mathbf{L} \cap \overline{\mathcal{U}}) \cap \overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t}) = \overline{\mathbf{L}} \cap \overline{\mathcal{U}}(\mathfrak{f}_F \mathbf{t})$ . The desired claim follows from Theorem 5.18.

#### 6. Cluster Algebras Associated with signed words

#### 6.1. Seeds from trapezoids.

Signed words. A signed word  $\underline{\mathbf{i}} = (\mathbf{i}_1, \mathbf{i}_2, \dots, \mathbf{i}_l)$  is a sequence of the elements of  $\mathbb{Z} \setminus \{0\}$ . For  $1 \leq j \leq k \leq l$ , denote  $\underline{\mathbf{i}}_{[j,k]} = (\mathbf{i}_j, \mathbf{i}_{j+1}, \dots, \mathbf{i}_k)$ . For any  $k \in [1, l]$ , define its successor and predecessor by

$$k[1] := \min(\{k' \in [k+1, l], |\mathbf{i}_{k'}| = |\mathbf{i}_k|\} \cup \{+\infty\}), k[-1] := \min(\{k' \in [1, k-1], |\mathbf{i}_{k'}| = |\mathbf{i}_k|\} \cup \{-\infty\}).$$

Inductively, define  $k[d \pm 1] := k[d][\pm 1]$  when  $k[d] \in \mathbb{Z}, d \in \mathbb{Z}$ .  $\forall a \in \mathbb{N}_{>0}$ , define

$$\begin{split} O^{\underline{\mathbf{i}}}([j,k];a) &= |\{s \in [j,k], \ |\mathbf{i}_s| = a\}|, \ O^{\underline{\mathbf{i}}}(a) := O^{\underline{\mathbf{i}}}([1,l];a), \\ o^{\underline{\mathbf{i}}}_{-}(k) &:= O^{\underline{\mathbf{i}}}([1,k-1];|\mathbf{i}_k|), \ o^{\underline{\mathbf{i}}}_{+}(k) := O^{\underline{\mathbf{i}}}([k+1,l];|\mathbf{i}_k|). \end{split}$$

Denote  $k^{\min} = k[o_{-}^{\underline{\mathbf{i}}}(k)], \ k^{\max} = k[o_{+}^{\underline{\mathbf{i}}}(k)].$ 

Define the length  $l(\underline{\mathbf{i}}) := l$ ,  $-\underline{\mathbf{i}} := (-\mathbf{i}_1, \ldots, -\mathbf{i}_l)$ , and the opposite word  $\underline{\mathbf{i}}^{\mathsf{op}} = (\mathbf{i}_l, \cdots, \mathbf{i}_2, \mathbf{i}_1)$ . Define the support  $\mathsf{supp} \, \underline{\mathbf{i}} := \{ |\mathbf{i}_k| \mid k \in [1, l] \}$ . For any  $s \in \mathbb{N}$ , we also denote s-copies of  $\underline{\mathbf{i}}$  by  $\underline{\mathbf{i}}^s := (\underline{\mathbf{i}}, \ldots, \underline{\mathbf{i}})$ .

Introduce  $I(\underline{\mathbf{i}}) := \{ \begin{pmatrix} a \\ d \end{pmatrix}^{\underline{\mathbf{i}}} | a \in \mathsf{supp} \underline{\mathbf{i}}, 0 \leq d < O^{\underline{\mathbf{i}}}(a) \}$ . Then we have an isomorphism  $[1, l] \simeq I(\underline{\mathbf{i}})$ , such that k is identified  $\binom{|\mathbf{i}_k|}{o_{\underline{\mathbf{i}}}^{\underline{\mathbf{i}}}(k)}^{\underline{\mathbf{i}}}$ . The order < on [1, l] induces the order  $<_{\mathbf{i}}$  on  $I(\underline{\mathbf{i}})$ . We also introduce  $\ddot{I}(\underline{\mathbf{i}}) := \{ \begin{pmatrix} a \\ -1 \end{pmatrix} | a \in J \} \sqcup I(\underline{\mathbf{i}}).$ 

We often omit the symbol  $\underline{\mathbf{i}}$  for simplicity. We will identify [1, l] and  $I(\underline{\mathbf{i}})$  and use their symbols interchangeably.

If  $\underline{\mathbf{i}}'$  is another signed word such that  $O^{\underline{\mathbf{i}}}(a) = O^{\underline{\mathbf{i}}'}(a), \forall a \in J$ , we will also identify  $I(\underline{\mathbf{i}})$  with  $I(\underline{\mathbf{i}}')$  such that  $\binom{a}{d}^{\underline{\mathbf{i}}} = \binom{a}{d}^{\underline{\mathbf{i}}'}$ . But  $<_{\underline{\mathbf{i}}}$  and  $<_{\mathbf{i}'}$  are different.

**Example 6.1.** Take  $\underline{\mathbf{i}} = (1, 2, 1, -2, -1, -2)$ . Then the ordered set  $I(\underline{\mathbf{i}}) = \{ \begin{pmatrix} 1 \\ 0 \end{pmatrix} < \begin{pmatrix} 2 \\ 0 \end{pmatrix} < \begin{pmatrix} 1 \\ 1 \end{pmatrix} < \begin{pmatrix} 2 \\ 1 \end{pmatrix} < \begin{pmatrix} 2 \\ 2 \end{pmatrix} \\ (\underline{\mathbf{i}}) < \begin{pmatrix} 2 \\ 2 \end{pmatrix} \}$  is isomorphic to [1, 6]. For  $\underline{\mathbf{i}}' = (2, 1, -2, -2, -1, -1)$ , we have  $I(\underline{\mathbf{i}}) = I(\underline{\mathbf{i}}')$  but their isomorphisms with [1, 6] are different.

Positive Braids. Choose and fix a finite subset  $J \subset \mathbb{Z}_{>0}$ . Let  $C = (C_{ab})_{a,b \in J}$  denote a generalized Cartan matrix with symmetrizers  $\mathsf{D}_a \in \mathbb{Z}_{>0}$ :  $C_{aa} = 2, C_{ab} \in \mathbb{Z}_{\leq 0}$  $\forall a \neq b$ , and  $\mathsf{D}_b C_{ab} = \mathsf{D}_a C_{ba}$ .

The monoid  $Br^+$  of positive braids is generated by  $\sigma_a, \forall a \in J$ , such that

$$\sigma_a \sigma_b = \sigma_b \sigma_a \text{ if } C_{ab} C_{ba} = 0, \qquad \sigma_a \sigma_b \sigma_a = \sigma_b \sigma_a \sigma_b \text{ if } C_{ab} C_{ba} = 1,$$
  
$$(\sigma_a \sigma_b)^2 = (\sigma_b \sigma_a)^2 \text{ if } C_{ab} C_{ba} = 2, \qquad (\sigma_a \sigma_b)^3 = (\sigma_b \sigma_a)^3 \text{ if } C_{ab} C_{ba} = 3.$$

Note that the generalized braid group Br is generated by these  $\sigma_a$  and their inverses.

A word  $\eta = (\eta_1, \eta_2, \dots, \eta_l)$  is a sequence formed from the letters of J. We associate with it the positive braid  $\beta_{\underline{\eta}} := \sigma_{\eta_1} \cdots \sigma_{\eta_l} \in \mathsf{Br}^+$ . The Weyl group W is the quotient of  $\mathsf{Br}$  by the relations  $\sigma_a^2 = \mathrm{Id}$ ,  $\forall a$ . Denote the image of  $\beta \in \mathsf{Br}$  in W by  $[\beta]$ . Denote  $w_{\underline{\eta}} := [\beta_{\underline{\eta}}]$  and  $s_a := [\sigma_a]$ . The length of an element  $w \in W$ , denoted l(w), is defined to be the minimum of the length of  $\underline{\eta}$  such that  $w = w_{\eta}$  in W, in which case we say  $\eta$  is a reduced word.

Let  $(\zeta, \eta)$  denote any pair of words. We will introduce the seed  $\mathbf{t}(\zeta, \eta, \Delta)$  associated to any triangulation  $\Delta$  of a trapezoid defined in [SW21]. We will also denote it by  $\mathbf{\ddot{t}}(\mathbf{\underline{i}})$  for the corresponding signed word  $\mathbf{\underline{i}}$ . See Example 6.2.

Triangulated trapezoids. Let  $\Sigma_{\underline{\eta}}^{\underline{\zeta}}$  denote a trapezoid in  $\mathbb{R}^2$  with horizontal bases, whose top base is subdivided into intervals labeled by  $-\zeta$  from left to right (i.e., along the (1,0)-direction), and its bottom base is analogously subdivided and labeled by  $\eta$ . The marked points are the endpoints of these intervals. A diagonal is an interior line segment in  $\Sigma_{\eta}^{\underline{s}}$  connecting a top marked point with a bottom one. A triangulation  $\Delta$  is a maximal collection of non-intersecting distinct diagonals.

Note that  $\Delta$  produces a partition of  $\Sigma_{\eta}^{\underline{\zeta}}$  into  $(l(\underline{\eta}) + l(\underline{\zeta}))$ -many triangles. For any triangle T in the partition, denoted  $T \in \Delta$ , we define  $\mathbf{i}_T \in \pm J$  to be the label of its horizontal edge. Define  $sign(T) := sign(i_T)$ .

Seeds. Draw distinct a-labeled horizontal lines  $L_a$  between the bases of  $\Sigma_{\eta}^{\underline{\zeta}}$  for  $a \in J$ , called layers. We insert one relative interior point  $p_T$  in  $L_{|\mathbf{i}_T|} \cap T$  for each  $T \in \Delta$ . The collection of  $L_a$  and  $p_T$  is called a string diagram, denoted  $D(\zeta, \eta, \Delta)$ .

Each layer  $L_a$  is subdivided by the nodes into intervals, which are labeled by the vectors  $\binom{a}{-1}, \binom{a}{0}, \ldots, \binom{a}{O(a)-1}$  from left to right. Let  $\ddot{I}$  consist of these intervals,  $I_{\mathsf{uf}} := \ddot{I}_{\mathsf{uf}}$  the bounded ones,  $\ddot{I}_{\mathsf{f}}$  the unbounded ones,  $I_{\mathsf{f}} = \{ \begin{pmatrix} a \\ O(a)-1 \end{pmatrix}, \forall a \}$  the rightunbounded ones, and  $I_{\rm f}^- = \{ \begin{pmatrix} a \\ -1 \end{pmatrix}, \forall a \}$  the left-unbounded ones. Any triangle  $T_k$ with  $|\mathbf{i}_k| = a$  intersects one interval  $\ell_b = \begin{pmatrix} b \\ O([1,k-1];b)-1 \end{pmatrix} \subset L_b$  for each  $b \neq a$ , an interval  $\ell_a^- = \begin{pmatrix} a \\ o_-(k) - 1 \end{pmatrix} \subset L_a$  on the left of  $p_T$ , and an interval  $\ell_a^+ = \begin{pmatrix} a \\ o_-(k) \end{pmatrix} \subset L_a$ on the right of  $p_T$ . It contributes a skew-symmetric  $\ddot{I} \times \ddot{I} \xrightarrow{\mathbb{Z}}$ -matrix  $B_{\Gamma}^{(T)}$  whose non-zero entries are:

- $\begin{array}{l} \bullet \ (B_{\Gamma}^{(T)})_{\ell_{a}^{-},\ell_{a}^{+}} = -(B_{\Gamma}^{(T)})_{\ell_{a}^{+},\ell_{a}^{-}} = \operatorname{sign}(T). \\ \bullet \ \forall b \neq a, (B_{\Gamma}^{(T)})_{\ell_{b},\ell_{a}^{-}} = -(B_{\Gamma}^{(T)})_{\ell_{b},\ell_{a}^{+}} = \frac{1}{2}\operatorname{sign}(T), (B_{\Gamma}^{(T)})_{\ell_{a}^{+},\ell_{b}} = -(B_{\Gamma}^{(T)})_{\ell_{a}^{-},\ell_{b}} = -(B_{\Gamma}^{(T)})_{\ell_{b}^{-},\ell_{b}} = -(B_{\Gamma}^{(T)})_{\ell_{b}^{$  $\frac{1}{2}$  sign(T).

Denote  $B_{\Gamma} := \sum_{T \in \Delta} B_{\Gamma}^{(T)}$  and its quiver by  $\Gamma$ . Define  $\ddot{B} = (b_{\ell,\ell'})_{\ell,\ell' \in \ddot{I}}$  by

- $b_{\ell,\ell'} = (B_{\Gamma})_{\ell,\ell'}$  if  $\ell, \ell' \subset L_a$  for some  $a \in J$ .
- $b_{\ell,\ell'} = (B_{\Gamma})_{\ell,\ell'}(-C_{ab})$  if  $\ell \subset L_a, \ell' \subset L_b$ , for some  $a \neq b \in J$ .

Then  $\ddot{B}$  is skew-symmetrizable with symmetrizers  $d_l$ , such that  $d_l^{\vee} = \mathsf{D}_a$  for  $\ell \subset L_a$ . In addition, the submatrix  $\ddot{B}_{\ddot{I} \times I_{uf}}$  is a  $\mathbb{Z}$ -matrix (see (6.1) or [SW21, Remark 3.6]).

The above data gives rise to a seed  $\mathbf{t}(\underline{\zeta}, \underline{\eta}, \Delta)$  whose *B*-matrix is  $B_{I \times I_{uf}}$ . Removing the frozen vertices in  $I_f^-$ , we further obtain the reduced seed  $\mathbf{t}(\underline{\zeta}, \underline{\eta}, \Delta)$  whose vertex set is  $I := I_{uf} \sqcup I_f$ .

**Example 6.2.** Choose the Cartan matrix  $C = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$ ,  $\underline{\zeta} = \underline{\eta} = (1, 2, 1)$ , and a triangulation for  $\Sigma_{\underline{\eta}}^{\underline{\zeta}}$ . Figure 6.1(A) shows the non-zero entries of  $B_{\Gamma}^{(T)}$  for single triangles T, Figure 6.1(B) shows the string diagram, and Figures 6.1(C)(D) show the quiver associated with  $\ddot{B}$ , where we identify  $\ddot{I}$  with  $[-2, -1] \sqcup [1, 6]$ . Circular nodes denote unfrozen vertices, and rectangular nodes denote frozen vertices.




FIGURE 6.1. The quiver for  $\ddot{B}(\ddot{t}(1,2,1,-1,-2,-1))$ .

6.2. Quantum seeds associated to signed words. Successively label the triangles in the triangulation  $\Delta$  of  $\Sigma_{\underline{\eta}}^{\underline{\zeta}}$  by  $T_1, T_2, \ldots$  from left to right, we obtain  $\underline{\mathbf{i}} = (\mathbf{i}_1, \mathbf{i}_2, \ldots) := (\mathbf{i}_{T_1}, \mathbf{i}_{T_2}, \ldots)$ . Then  $\underline{\mathbf{i}}$  is a (riffle) shuffle of the signed words  $-\underline{\zeta}$ and  $\underline{\eta}$ . Conversely, any signed word  $\underline{\mathbf{i}}$  determines a triangulation  $\Delta$ . For example,  $\underline{\mathbf{i}} = (1, 2, 1, -1, -2, -1)$  in Figure 6.1. Since the string diagram  $D(\underline{\zeta}, \underline{\eta}, \Delta)$  is determined by  $\underline{\mathbf{i}}$ , we can denote  $\ddot{\mathbf{t}}(\underline{\zeta}, \underline{\eta}, \Delta) = \ddot{\mathbf{t}}(\underline{\mathbf{i}})$  and  $\dot{\mathbf{t}}(\underline{\zeta}, \underline{\eta}, \Delta) = \dot{\mathbf{t}}(\underline{\mathbf{i}})$ .

We identify the set  $\ddot{I}$  of intervals with  $(-J) \sqcup I(\underline{\mathbf{i}})$ , sending  $\begin{pmatrix} a \\ -1 \end{pmatrix}$  to -a and  $\begin{pmatrix} a \\ d \end{pmatrix}$  to  $\begin{pmatrix} a \\ d \end{pmatrix}^{\underline{\mathbf{i}}}$ ,  $\forall a \in J, d \in [0, O^{\underline{\mathbf{i}}}(a) - 1]$ . Then we have  $I = I(\underline{\mathbf{i}}) \simeq [1, l]$ , such that the interval  $\ell^+_{|\mathbf{i}_k|}$  associated with  $T_k$  is identified with  $\begin{pmatrix} |\mathbf{i}_k| \\ o_{-(k)} \end{pmatrix}^{\underline{\mathbf{i}}}$  and k. We will denote the elements in I by  $\begin{pmatrix} a \\ d \end{pmatrix}^{\underline{\mathbf{i}}}$  and  $k \in [1, l]$  interchangeably. We also denote  $\begin{pmatrix} a \\ d \end{pmatrix}^{\underline{\mathbf{i}}}$  by  $\begin{pmatrix} a \\ d \end{pmatrix}^{\underline{\mathbf{i}}(\underline{\mathbf{i}})}$ ,  $\begin{pmatrix} a \\ d \end{pmatrix}^{\underline{\mathbf{i}}(\underline{\mathbf{i}})}$ , or simply  $\begin{pmatrix} a \\ d \end{pmatrix}$ . Denote  $\mathbf{i}_{-a} = -a$ .

Define  $\varepsilon_k := \operatorname{sign}(T_k), \forall k \in [1, l]$ . Extend  $[1] : I \to I \sqcup \{+\infty\}$  to a map from  $\ddot{I}$  to  $I \sqcup \{+\infty\}$  such that  $\binom{a}{d}[1] := \binom{a}{d+1}, \forall d \in [-1, O(a) - 1]$ , where we understand  $\binom{a}{O(a)}$  as  $+\infty$ .

**Lemma 6.3.** The matrix entries  $b_{jk}$ ,  $(j,k) \in \ddot{I} \times I_{uf}$ , of  $\ddot{B}$  are given as below:

$$(6.1) b_{jk} = \begin{cases} \varepsilon_k & k = j[1] \\ -\varepsilon_j & j = k[1] \\ \varepsilon_k C_{|\mathbf{i}_j|, |\mathbf{i}_k|} & \varepsilon_{j[1]} = \varepsilon_k, \ j < k < j[1] < k[1] \\ \varepsilon_k C_{|\mathbf{i}_j|, |\mathbf{i}_k|} & \varepsilon_k = -\varepsilon_{k[1]}, \ j < k < k[1] < j[1] \\ -\varepsilon_j C_{|\mathbf{i}_j|, |\mathbf{i}_k|} & \varepsilon_{k[1]} = \varepsilon_j, \ k < j < k[1] < j[1] \\ -\varepsilon_j C_{|\mathbf{i}_j|, |\mathbf{i}_k|} & \varepsilon_j = -\varepsilon_{j[1]}, \ k < j < j[1] < k[1] \\ 0 & otherwise \end{cases}$$

*Proof.* The case  $|\mathbf{i}_j| = |\mathbf{i}_k|$  is trivial. When  $|\mathbf{i}_j| \neq |\mathbf{i}_k|$ , the following triangles contribute to  $b_{jk}$ .

(i)  $T_k$  such that j < k < j[1]: We have  $\ell_{|\mathbf{i}_k|}^+ = k$  and it contributes  $\varepsilon_k \frac{1}{2} C_{|\mathbf{i}_j|, |\mathbf{i}_k|}$ . (ii)  $T_{k[1]}$  such that j < k[1] < j[1]: We have  $\ell_{|\mathbf{i}_k|}^- = k$  and contributes  $-\varepsilon_{k[1]} \frac{1}{2} C_{|\mathbf{i}_j|, |\mathbf{i}_k|}$ . (iii)  $T_j$  such that k < j < k[1]: We have  $\ell_{|\mathbf{i}_j|}^+ = j$  and it contributes  $-\varepsilon_j \frac{1}{2} C_{|\mathbf{i}_j|, |\mathbf{i}_k|}$ . (iv)  $T_{j[1]}$  such that k < j[1] < k[1]: We have  $\ell_{|\mathbf{i}_j|}^- = j$  and it contributes  $\varepsilon_{j[1]} \frac{1}{2} C_{|\mathbf{i}_j|, |\mathbf{i}_k|}$ .

Note that (i) and (iii) cannot appear simultaneously, and similar for (ii) and (iv). Then (i) and (iv) sum to the 3rd identity, (i) and (ii) sum to the 4th identity, (iii) and (ii) sum to the 5th identity, and (iii) and (iv) sum to the 6th identity.

Let  $\underline{\mathbf{j}}$  denote a word in which each -a appears exactly once, where  $a \in J$ . By Lemma 6.3, we have  $\ddot{B}_{I \times I_{uf}} = -\tilde{B}$  for the matrix  $\tilde{B}$  associated to  $(\underline{\mathbf{j}}, \underline{\mathbf{i}})$  in [BZ05, (8.7)]. Then [BZ05, (8.5)(8.9)], after a sign change, provides us a quantization matrix  $\ddot{\Lambda}$  for  $\mathbf{t}(\underline{\mathbf{i}})$ .

**Lemma 6.4.** The matrix  $\widetilde{B}(\dot{\mathbf{t}}(\underline{\mathbf{i}}))$  is of full rank.


Proof. (6.1) implies  $\deg^{\mathbf{t}} y_k = -\varepsilon_{k[1]} f_{k[1]} + \sum_{i < k[1]} b_{ik} f_i$ ,  $\forall k \in I_{uf}$ . Moreover, [1] is injective on  $I_{uf}$ . We deduce that  $\deg^{\mathbf{t}} y_k$ ,  $k \in I_{uf}$ , are linearly independent. The desired claim follows. Alternatively, the claim is implied by [CGG<sup>+</sup>22, Section 8.1].

Lemma 6.4 guarantees that we can quantize the seed  $\dot{t}(\underline{i})$ .

## 6.3. Seeds operations.

Mutations and reflections. Let  $\varepsilon$  denote any sign. By [SW21, Proposition 3.7], the following operations on  $\mathbf{t} = \ddot{\mathbf{t}}(\underline{\zeta}, \underline{\eta}, \Delta) = \ddot{\mathbf{t}}(\underline{\mathbf{i}})$  will generate new seeds  $\mathbf{t}' = \ddot{\mathbf{t}}(\zeta, \eta, \Delta') = \ddot{\mathbf{t}}(\underline{\mathbf{i}}')$ :

• (Flips) Change  $\mathbf{i} = (\dots, \varepsilon a, -\varepsilon b, \dots)$  to  $\mathbf{i}' = (\dots, -\varepsilon b, \varepsilon a, \dots)$ , where  $\varepsilon a$  is the k-th signed letter of  $\mathbf{i}$ . Then  $\mathbf{t}' = \mu_{\ell_{a,T_k}^+} \mathbf{t}$  when a = b, and  $\mathbf{t}' = \mathbf{t}$  otherwise. Note that the associated string diagrams are related by the diagonal flip of the quadrilateral  $T_k \cup T_{k+1}$ :



• (Braid moves) Assume that  $\beta_{\underline{\eta}} = \beta_{\underline{\eta}'} \in \mathsf{Br}^+$  such that  $\underline{\eta}$  take the form  $\underline{\mathbf{i}}_{[j,k]}$ . Replacing  $\underline{\eta}$  by  $\underline{\eta}'$  in  $\underline{\mathbf{i}}$ , we obtain  $\underline{\mathbf{i}}'$ . Then  $\mathbf{t}'$  and  $\mathbf{t}$  are related by a sequence of mutations  $\boldsymbol{\mu}_{\underline{\mathbf{i}}',\underline{\mathbf{i}}}$  at vertices  $r \in [j,k]$  such that  $r[1] \leq k$  and a permutation  $\sigma_{\underline{\mathbf{i}}',\underline{\mathbf{i}}}$  on [j,k]. We use  $\boldsymbol{\mu}_{\mathbf{i}',\mathbf{i}}^{\sigma}$  to denote this permuted mutation sequence.

Therefore, for any given words  $\underline{\eta}$ ,  $\underline{\zeta}$ ,  $\underline{\eta}'$ ,  $\underline{\zeta}'$ , such that  $\beta_{\underline{\eta}} = \beta_{\underline{\eta}'}$ ,  $\beta_{\underline{\zeta}} = \beta_{\underline{\zeta}'}$ , and any shuffles  $\underline{\mathbf{i}}$  of  $(-\underline{\zeta}, \underline{\eta})$  and  $\underline{\mathbf{i}}'$  of  $(-\underline{\zeta}', \underline{\eta}')$ , the seeds  $\mathbf{\ddot{t}}(\underline{\mathbf{i}})$  and  $\mathbf{\ddot{t}}(\underline{\mathbf{i}}')$  are connected by (permuted) mutations.

Following [SW21, Section 2.3], we introduce the reflection operations which will generate new seeds  $\mathbf{\ddot{t}}(\mathbf{\underline{i}}') = \mathbf{\ddot{t}}(\zeta', \eta', \Delta')$  similar to  $\mathbf{\ddot{t}}(\mathbf{\underline{i}}) = \mathbf{\ddot{t}}(\zeta, \eta, \Delta)$ .

- Left reflection: change  $\underline{\mathbf{i}} = (\varepsilon a, \ldots)$  to  $\underline{\mathbf{i}}' = (-\varepsilon a, \ldots)$ .
- Right reflection: change  $\underline{\mathbf{i}} = (\dots, \varepsilon a)$  to  $\underline{\mathbf{i}}' = (\dots, -\varepsilon a)$ .

Note that  $\Sigma_{\underline{\eta}'}^{\underline{\zeta}'} \neq \Sigma_{\underline{\eta}}^{\underline{\zeta}}$  and  $\ddot{\mathbf{t}}(\underline{\mathbf{i}}') \neq \ddot{\mathbf{t}}(\underline{\mathbf{i}})$ , but we have  $\dot{\mathbf{t}}(\underline{\mathbf{i}}') = \dot{\mathbf{t}}(\underline{\mathbf{i}})$  under left reflections. Using flips and left reflects, we obtain  $\overline{\mathcal{A}}(\dot{\mathbf{t}}(\underline{\mathbf{i}})) = \overline{\mathcal{A}}(\dot{\mathbf{t}}(\underline{\zeta}^{\mathsf{op}}\underline{\eta}))$ , which only depends on  $\beta_{\underline{\zeta}^{\mathsf{op}}\underline{\eta}} \in \mathsf{Br}^+$  by braid moves.

It is worth noting that the seeds connected by these operations provide the same cluster algebra. More precisely, let there be given signed words  $\underline{\mathbf{i}}^{(i)}$ , i = 1, 2, 3. Let  $\mathbf{t}^{(i)}$  denote either  $\mathbf{\dot{t}}(\underline{\mathbf{i}}^{(i)})$  or  $\mathbf{\ddot{t}}(\underline{\mathbf{i}}^{(i)})$ . In the former case, we assume  $\underline{\mathbf{i}}^{(i)}$  and  $\underline{\mathbf{i}}^{(j)}$ ,  $\forall i, j \in [1, 3]$ , are connected by flips, braid moves, and left reflections; in the latter case, we assume they are connected by flips and braid moves. Let us choose permutation mutation sequences  $\boldsymbol{\mu}^{\sigma}_{\underline{\mathbf{i}}^{(j)},\underline{\mathbf{i}}^{(i)}}$  associated to these operations such that  $\mathbf{t}^{(j)} = \boldsymbol{\mu}^{\sigma}_{\mathbf{i}^{(j)},\mathbf{i}^{(i)}} \mathbf{t}^{(i)}$ . Consider the following diagram for  $\mathcal{A}^{(i)} := \mathcal{A}(\mathbf{t}^{(i)})$ .

(6.2) 
$$\begin{array}{ccc} & (\boldsymbol{\mu}_{\underline{i}^{(3)},\underline{i}^{(1)}})^{*} \\ \mathcal{A}^{(3)} & \stackrel{\sim}{\longrightarrow} & \mathcal{A}^{(1)} \\ \downarrow^{\wr} (\boldsymbol{\mu}_{\underline{i}^{(3)},\underline{i}^{(2)}})^{*} & & \parallel & . \\ & (\boldsymbol{\mu}_{\underline{i}^{(2)},\underline{i}^{(1)}})^{*} \\ \mathcal{A}^{(2)} & \stackrel{\sim}{\longrightarrow} & \mathcal{A}^{(1)} \end{array}$$

Diagram (6.2) is commutative at the classical level by [SW21, Theorem 1.1]. By tracking the degrees of the quantum cluster variables, which are the same as those of their classical counterparts, we could easily deduce that the diagram is commutative at the quantum level as well (details could be found in [Qin24b, Lemma 3.2]).

Subwords. Take any  $1 \leq j \leq k \leq l$ . Define  $\mathbf{i}' = (\mathbf{i}'_1, \dots, \mathbf{i}'_{k-j+1})$  to be  $\mathbf{i}_{[j,k]} = (\mathbf{i}_j, \mathbf{i}_{j+1}, \dots, \mathbf{i}_k), \ \mathbf{\ddot{t}}' := \mathbf{\ddot{t}}(\mathbf{\underline{i}}'), \ \mathbf{\ddot{I}}' := I(\mathbf{\ddot{t}}'), \ \mathbf{I}'_{\mathsf{uf}} := I_{\mathsf{uf}}(\mathbf{\ddot{t}}'), \text{ and } I' := I(\mathbf{\dot{t}}(\mathbf{\underline{i}}')).$ 

**Definition 6.5.** Define the embedding  $\iota_{\underline{\mathbf{i}},\underline{\mathbf{i}}'}: \ddot{I}' \hookrightarrow \ddot{I}$ , such that  $\iota\binom{a}{d}^{\underline{\mathbf{i}}'} = \binom{a}{d+O([1,j-1];a)}^{\underline{\mathbf{i}}}$ .

Abbreviate  $\iota_{\underline{\mathbf{i}},\underline{\mathbf{i}}'}$  by  $\iota$ . Note that  $\iota(I'_{\mathsf{uf}}) \subset I_{\mathsf{uf}}, \iota(I') \subset I$ . Under the identification  $I' \simeq [1, k - j + 1]$  and  $I \simeq [1, l]$ , we have  $\iota(s) = s + j - 1, \forall s \in [1, k - j + 1]$ . In particular, s < s' if and only if  $\iota(s) < \iota(s')$  for  $s, s' \in I'$ . We have the following.


**Lemma 6.6.** The embedding  $\iota$  is a cluster embedding from  $\ddot{\mathbf{t}}(\underline{\mathbf{i}}')$  to  $\ddot{\mathbf{t}}(\underline{\mathbf{i}})$ , such that  $\ddot{\mathbf{t}}(\underline{\mathbf{i}}')$  is a good sub seed of  $\ddot{\mathbf{t}}(\underline{\mathbf{i}})$ . It restricts to a cluster embedding  $\iota$  from  $\dot{\mathbf{t}}(\underline{\mathbf{i}}')$  to  $\dot{\mathbf{t}}(\underline{\mathbf{i}})$ . When  $\underline{\mathbf{i}}' = \underline{\mathbf{i}}_{[1,k]}$ ,  $\dot{\mathbf{t}}(\underline{\mathbf{i}}')$  is a good sub seed of  $\dot{\mathbf{t}}(\underline{\mathbf{i}})$ .

*Proof.* Let  $\Sigma(\mathbf{i})$  denote the triangulated trapezoid associated with  $\mathbf{i}$ , whose triangles are  $T_1, \ldots, T_{l(\mathbf{i})}$ . Then  $T_j, \ldots, T_k$  form a triangulated trapezoid  $\Sigma(\mathbf{i}')$  associated

with  $\mathbf{i}'$ , such that  $T_r$ ,  $j \leq r \leq k$ , are identified with the triangle  $T'_{r-j+1}$  of  $\Sigma(\mathbf{i}')$ . Under this identification, the interval  $\binom{a}{d}^{\mathbf{i}'}$  only overlaps  $\iota\binom{a}{d}^{\mathbf{i}'}$ . Summing up the contributions from  $T_1, \ldots, T_{l(\mathbf{i})}$ , we deduce  $b'_{jk} = b_{\iota(j),\iota(k)}, \forall j \in \ddot{I}', k \in I'_{uf}$  and, moreover,  $b_{j',\iota(k)} = 0 \forall j' \in \ddot{I} \setminus \iota(\ddot{I}')$ . Therefore,  $\iota$  is a cluster embedding from  $\ddot{\mathbf{t}}(\mathbf{i}')$  to  $\ddot{\mathbf{t}}(\mathbf{i})$  such that  $\ddot{\mathbf{t}}(\mathbf{i}')$  is a good sub seed, and it restricts to a cluster embedding from  $\dot{\mathbf{t}}(\mathbf{i}')$  to  $\dot{\mathbf{t}}(\mathbf{i})$ .

If 
$$\underline{\mathbf{i}}' = \underline{\mathbf{i}}_{[1,k]}, I \setminus \iota(I')$$
 equals  $\{j'|j' > k\} = I \setminus \iota(I')$ . So  $\mathbf{t}(\underline{\mathbf{i}}')$  is a good sub seed.

**Example 6.7.** Take  $\underline{\zeta} = \underline{\eta} = (1, 2, 1)$ . The quiver associated to  $\mathbf{\ddot{t}}(\underline{\mathbf{i}}), \underline{\mathbf{i}} := (\underline{\eta}, -\underline{\zeta})$  is drawn in Figure 6.1. The quiver associated to  $\mathbf{\ddot{t}}(\underline{\mathbf{i}}'), \underline{\mathbf{i}}' := (\underline{\eta}_{[2,3]}, -\underline{\zeta})$  is drawn in Figure 6.2. Moreover,  $\mathbf{\ddot{t}}(\underline{\mathbf{i}}')$  is a good seed of  $\mathbf{\ddot{t}}(\underline{\mathbf{i}})$  under the cluster embedding  $\iota_{\underline{\mathbf{i}},\underline{\mathbf{i}}'}$ , such that  $\iota_{\underline{\mathbf{i}},\underline{\mathbf{i}}'}(-1) = 1$ ,  $\iota_{\underline{\mathbf{i}},\underline{\mathbf{i}}'}(-2) = 2$ , and  $\iota_{\underline{\mathbf{i}},\underline{\mathbf{i}}'}(s) = s + 1$ ,  $\forall s \in [1,5]$ .



*Extensions.* Let  $\widetilde{J}$  be any subset of  $\mathbb{Z}_{>0}$  containing J and  $\widetilde{C}$  any  $\widetilde{J} \times \widetilde{J}$  generalized Cartan matrix,  $\underline{\widetilde{i}}$  any signed word in the letters from  $\pm \widetilde{J}$ .

**Definition 6.8** (Extension). If  $\tilde{C}_{J \times J} = C$ ,  $\tilde{C}$  is called an extension of C.  $\underline{\tilde{i}}$  is called a letter extension of a signed word  $\underline{i}$ , if  $\underline{i}$  can be obtained from  $\underline{\tilde{i}}$  by removing the letters from  $\pm \tilde{J} \setminus J$ .

Assume  $\widetilde{C}$  is an extension of C and  $\underline{\widetilde{i}}$  a letter extension of  $\underline{i}$  from now on. Denote  $\mathbf{t}(\underline{\widetilde{i}}) = \mathbf{t}(\underline{\widetilde{\zeta}}, \underline{\widetilde{\eta}}, \underline{\widetilde{\Delta}})$ . Then the string diagram  $D(\underline{\zeta}, \underline{\eta}, \underline{\Delta})$  is contained in  $D(\underline{\widetilde{\zeta}}, \underline{\widetilde{\eta}}, \underline{\widetilde{\Delta}})$ . Note that a vertex of  $\mathbf{t}(\underline{\widetilde{i}})$  is either a vertex of  $\mathbf{t}(\underline{i})$ , or an interval on  $L_c$  for  $c \in \widetilde{J} \setminus J$ . We have the following observation.

**Lemma 6.9.** The matrix  $\ddot{B}(\ddot{\mathbf{t}}(\underline{\mathbf{i}}))$  equals  $\ddot{B}(\ddot{\mathbf{t}}(\underline{\mathbf{i}}))_{\ddot{I}\times\ddot{I}}$ .

The string diagram (or an associated quiver) of  $\underline{\tilde{i}}$  is called a layer extension of that of  $\underline{i}$ , see Figure 6.3. Let  $\widetilde{W}$  denote the Weyl group associated to  $\widetilde{C}$ . Then W is a subgroup of  $\widetilde{W}$ .


**Lemma 6.10** (Extension of words). For any word  $\underline{\eta}$  in J, there exist an extended generalized Cartan matrix  $\widetilde{C}$  and a reduced word  $\underline{\widetilde{\eta}}$  for  $\widetilde{W}$ , such that  $\underline{\widetilde{\eta}}$  is a letter extension of  $\eta$ .

*Proof.* Choose any  $c > \max J$ . Extend C to a  $(J \sqcup \{c\}) \times (J \sqcup \{c\})$ -matrix Cartan  $\widetilde{C}$  such that  $C_{a,c}C_{a,c} \ge 4$ ,  $\forall a \in J$ .  $\underline{\widetilde{\eta}} := (\eta_1, c, \eta_2, c, \dots, c, \eta_l)$  meets our expectation.

**Example 6.11.** Take  $\underline{\eta} = (1, 1, 1, 1)$  where  $C = (2)_{\{1\} \times \{1\}}$ . The quiver for the seed  $\ddot{\mathbf{t}}(\eta)$  is drawn in Figure 6.3(A). Choose the letter extension  $\tilde{\eta} = (1, 2, 1, 2, 1, 2, 1, 2)$ 

for  $\widetilde{C} = \begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}_{[1,2]\times[1,2]}$ . The quiver for the seed  $\mathbf{\ddot{t}}(\underline{\tilde{\eta}})$  is drawn in Figure 6.3(B). By removing its vertices 5, 6, 7, 8 (they correspond to the intervals on  $L_2$ ), we recover Figure 6.3(A).

FIGURE 6.3. A quiver and an extension



From now on, let  $\underline{\mathbf{i}}$  be a shuffle of  $(-\underline{\zeta}, \underline{\eta})$  and denote  $\underline{\xi} := (\underline{\zeta}^{\mathsf{op}}, \underline{\eta})$ .

**Proposition 6.12.** There exists a mutation sequence  $\boldsymbol{\mu}$  such that  $\dot{\mathbf{t}}(\underline{\mathbf{i}}) = \boldsymbol{\mu}\dot{\mathbf{t}}(\underline{\xi})$ . Moreover, we can find an extended Cartan matrix  $\widetilde{C}$ , a letter extension  $\underline{\widetilde{\xi}}$  of  $\underline{\xi}$ , and  $F \subset I_{uf}(\dot{\mathbf{t}}(\underline{\widetilde{\xi}}))$ , such that  $\underline{\widetilde{\xi}}$  is reduced, and the classical seed  $\dot{\mathbf{t}}(\underline{\xi})$  can be obtain from  $\mathfrak{f}_F \dot{\mathbf{t}}(\underline{\widetilde{\xi}})$  by removing frozen vertices.

*Proof.* Note that  $\dot{\mathbf{t}}(\underline{\mathbf{i}})$  can be obtained from  $\dot{\mathbf{t}}(\underline{\xi})$  by iterating flips (mutations) and left reflections (who act trivially on the seeds), see [Cao22, Proposition 4.3].

By Lemma 6.10, we can find an extended  $\widetilde{J} \times \widetilde{J}$  generalized Cartan matrix  $\widetilde{C}$ and a letter extension  $\underline{\widetilde{\xi}}$  of the word  $\underline{\xi}$  which is reduced for the Weyl group  $\widetilde{W}$ . Note that  $\widetilde{B}(\dot{\mathbf{t}}(\underline{\xi}))$  equals  $\widetilde{B}(\dot{\mathbf{t}}(\underline{\widetilde{\xi}}))_{I \times I_{uf}}$  by Lemma 6.9. Therefore,  $\dot{\mathbf{t}}(\underline{\xi})$  can be obtained from  $\dot{\mathbf{t}}(\underline{\widetilde{\xi}})$  by freezing and then removing the vertices on the lines  $L_c, c \in \widetilde{J} \setminus J$ .

**Theorem 6.13.** Let  $\mathbf{t}$  denote the seed  $\dot{\mathbf{t}}(\underline{\mathbf{i}})$  or the seed  $\ddot{\mathbf{t}}(\underline{\mathbf{i}})$  arising from (decorated) double Bott-Samelson cells. The following statements hold for  $\mathcal{U}(\mathbf{t})$ :

(1) It possesses the common triangular basis, denoted L.

(2) When C is symmetric,  $(\mathcal{U}(\mathbf{t}), \mathbf{L})$  admits quasi-categorification.

*Proof.* Since  $\dot{\mathbf{t}}(\underline{\mathbf{i}})$  are  $\ddot{\mathbf{t}}(\underline{\mathbf{i}})$  are similar, it suffices to check the claim for  $\mathbf{t} = \dot{\mathbf{t}}(\underline{\mathbf{i}})$ .

Let  $\underline{\mathbf{i}}$  denote a shuffle of  $(-\underline{\zeta},\underline{\eta})$  and denote  $\underline{\xi} = (\underline{\zeta}^{\mathsf{op}},\underline{\eta})$ . Then  $\mathbf{\dot{t}}(\underline{\xi}) = \mu \mathbf{\dot{t}}(\underline{\mathbf{i}})$  for a mutation sequence  $\mu$ , see Proposition 6.12. Let  $\underline{\widetilde{\xi}}$  denote a letter extension of  $\underline{\xi}$ as in Proposition 6.12 and denote  $w = w_{\underline{\widetilde{\xi}}} \in \widetilde{W}$ . Note that, with an appropriate quantization,  $\mathbf{\dot{t}}(\underline{\widetilde{\xi}})$  is a quantum seed for the quantum cluster algebra  $\mathcal{O}_q[N_w]$ . The dual canonical basis of  $\mathcal{O}_q[N_w]$  is the common triangular basis [Qin17], denoted  $\mathbf{L}_w$ .

By Proposition 6.12, for some  $F \subset I_{uf}(\dot{\mathbf{t}}(\underline{\boldsymbol{\xi}}))$ , the classical seed  $\mathbf{t}$  is obtained from  $\mathbf{t}' := \mathfrak{f}_F \dot{\mathbf{t}}(\underline{\tilde{\boldsymbol{\xi}}})$  by removing frozen vertices. Note that  $\mathbf{t}$  and  $\mathbf{t}'$  are similar as classical seeds. In addition,  $\mathcal{U}(\mathbf{t}')$  has the common triangular basis  $\mathfrak{f}_F \mathbf{L}_w$  (Theorem 3.17). We deduce that the  $\mathcal{U}(\mathbf{t})$  and  $\mathcal{U}(\mathbf{t}')$  are related by base changes and quantization changes, see Section 4.2, (4.3):

(6.3) 
$$\mathcal{U}(\dot{\mathbf{t}}(\underline{\widetilde{\xi}})) \xrightarrow{\mathfrak{f}_{F}} \mathcal{U}(\mathbf{t}') \xleftarrow{\varphi_{1}} \mathcal{U}(\mathbf{t}^{\mathrm{prin}},\lambda_{1}) \xrightarrow[\mathrm{change}]{\mathrm{quantization}} \mathcal{U}(\mathbf{t}^{\mathrm{prin}},\lambda_{2}) \xrightarrow{\varphi_{2}} \mathcal{U}(\mathbf{t}),$$

where  $(\mathbf{t}^{\text{prin}}, \lambda_1)$  and  $(\mathbf{t}^{\text{prin}}, \lambda_2)$  are two principal coefficients seeds with different quantization. By Proposition 4.18, the common triangular basis  $\mathbf{L}$  for  $\mathcal{U}(\mathbf{t})$  exists and consists of the elements similar to  $\mathfrak{f}_F \mathbf{L}_w$ .

If C is symmetric,  $(\mathcal{U}', \mathfrak{f}_F \mathbf{L}_w)$  is categorified by a monoidal subcategory  $\mathcal{C}'$  of  $\widetilde{\mathcal{C}}_w$  after localization (Theorem 5.17). Therefore,  $(\mathcal{U}(\mathbf{t}), \mathbf{L})$  is quasi-categorified by  $\mathcal{C}'$ .

**Theorem 6.14** (Classical  $\mathcal{A} = \mathcal{U}$  by [SW21]). For  $\mathbf{t} = \ddot{\mathbf{t}}(\underline{\mathbf{i}})$  or  $\dot{\mathbf{t}}(\underline{\mathbf{i}})$ , we have  $\mathcal{A}(\mathbf{t}) = \mathcal{U}(\mathbf{t})$ , and its common triangular basis elements have nice cluster decomposition in  $\mu \mathbf{t}$ , where  $\mu$  satisfies  $\mu(\dot{\mathbf{t}}(\underline{\mathbf{i}})) = \dot{\mathbf{t}}((\underline{\zeta}^{\mathsf{op}}, \underline{\eta}))$ .

Proof. Recall that  $\underline{\xi} = (\underline{\zeta}^{\text{op}}, \underline{\eta})$  and  $\overline{\mathcal{A}}(\mathbf{t}(\underline{\xi})) = \overline{\mathcal{U}}(\mathbf{t}(\underline{\xi})) = \mathcal{O}_q[N(w)]$  (see [GY16]). So  $\mathcal{A}(\mathbf{t}(\underline{\xi})) = \mathcal{U}(\mathbf{t}(\underline{\xi}))$  and the elements of  $\mathbf{L}_w$  have nice cluster decompositions in  $\mathbf{t}(\underline{\xi})$ , since  $\mathcal{O}_q[N(w)]$  has the dual PBW basis. Note that all algebras in (6.3) have the common triangular bases related by freezing or similarity. Applying Corollaries 3.16 and 4.12 to  $(\mathcal{A}(\mathbf{t}(\underline{\xi})), \mathbf{L}_w)$  in (6.3), we deduce that  $\mathcal{A}(\mathbf{t}(\underline{\xi})) = \mathcal{U}(\mathbf{t}(\underline{\xi}))$  and its common triangular basis elements have nice cluster decompositions in  $\mathbf{t}(\xi)$ .

The claim for  $\mu \dot{\mathbf{t}}(\underline{\mathbf{i}})$  follows from its similarity with  $\dot{\mathbf{t}}(\xi)$  and Corollary 4.12.

## 6.4. Optimized seeds.

**Lemma 6.15.** Denote  $\mathbf{t} = \dot{\mathbf{t}}(\underline{\eta})$  for any word  $\underline{\eta}$ . Then,  $\forall k \in I$ ,  $k^{\max}$  is optimized in  $\Sigma_{k^{\min}} \mathbf{t}$ .

*Proof.* Let us give a short proof using [Cao22]. Extend  $\underline{\eta}$  to  $\underline{\eta}' = (\underline{\eta}, \eta_k)$ . Then  $I(\mathbf{t}') = I \sqcup \{l(\underline{\eta}) + 1\}, k^{\max}$  becomes unfrozen in  $\mathbf{t}' := \dot{\mathbf{t}}(\underline{\eta}')$ , and  $\widetilde{B}(\mathbf{t}) = \widetilde{B}(\mathbf{t}')_{I \times I_{\text{uf}}}$ . By [Cao22, Lemma 4.6], we have  $b_{j,k^{\max}}(\Sigma_{k^{\min}}\mathbf{t}') \leq 0$  for all  $j \in I_{\text{uf}}$ .

### **Proposition 6.16.** For any signed word $\underline{\mathbf{i}}$ , $\mathbf{\ddot{t}}(\underline{\mathbf{i}})$ can be optimized.

*Proof.* (1) Let us prove the claim for any frozen vertex  $k^{\max}$ ,  $k \in I$ . For this purpose, it suffices to show  $k^{\max}$  can be optimized in the seed  $\dot{\mathbf{t}}(\underline{\mathbf{i}})$ , which has less frozen vertices. We can mutate  $\dot{\mathbf{t}}(\underline{\mathbf{i}})$  to a seed of the form  $\dot{\mathbf{t}}(-\underline{\zeta},\underline{\eta})$  for words  $\underline{\zeta},\underline{\eta}$  in J. It is connected to  $\dot{\mathbf{t}}(\underline{\zeta}^{\text{op}},\underline{\eta})$  by mutations [Cao22, Proposition 4.3]. Then  $k^{\max}$  can be optimized by Lemma 6.15.

(2) Let us prove the claim for any frozen vertex  $-\eta_k$ ,  $k \in I$ . Observe that  $\mathbf{\ddot{t}}(\underline{\mathbf{i}})$  is identified with  $\mathbf{\ddot{t}}(-\underline{\mathbf{i}}^{\mathsf{op}})$  after we identify the vertices  $\binom{a}{-1+d}$  of  $\mathbf{\ddot{t}}(\underline{\mathbf{i}})$  with  $\binom{a}{O(a)-1-d}$  of  $\mathbf{\ddot{t}}(-\underline{\mathbf{i}}^{\mathsf{op}})$ ,  $\forall d \in [0, O(a)]$ . Specifically, the frozen vertex  $-\eta_k = \binom{\eta_k}{-1}$  of  $\mathbf{\ddot{t}}(\underline{\mathbf{i}})$  becomes identified with the frozen vertex  $k^{\max} = \binom{\eta_k}{O(\eta_k)-1}$  of  $\mathbf{\ddot{t}}(-\underline{\mathbf{i}}^{\mathsf{op}})$ , which can be optimized by (1).

**Theorem 6.17.** Let t denote  $\dot{t}(\underline{i})$  or  $\ddot{t}(\underline{i})$ . Then the following statements hold.

- (1)  $\overline{\mathcal{U}}(\mathbf{t})$  possesses the common triangular basis, denoted **L**.
- (2) When C is symmetric,  $(\overline{\mathcal{U}}(\mathbf{t}), \mathbf{L})$  admits quasi-categorification.

<sup>&</sup>lt;sup>2</sup>Our quiver is opposite to that of [Cao 22] but the *B*-matrices are the same.

*Proof.* (1) Since all frozen vertices of  $\dot{\mathbf{t}}(\underline{\mathbf{i}})$  and  $\ddot{\mathbf{t}}(\underline{\mathbf{i}})$  can be optimized, combining Theorem 6.13 and Proposition 2.15, we obtain the desired claim.

(2) Note that we allow localization in quasi-categorification. The claim follows from Theorem 6.13(2).

#### 

## 7. Applications: cluster algebras from Lie theory

7.1. Double Bott-Samelson cells. As in Section 6, let C denote a  $J \times J$  generalized Cartan matrix. For any words  $\underline{\zeta}$ ,  $\underline{\eta}$ , let  $\ddot{X}_{\beta\underline{\eta}}^{\underline{\beta}\underline{\zeta}}$  denote the decorated double Bott-Samelson cell in [SW21, Section 2]. Choose any shuffle  $\underline{\mathbf{i}}$  of  $-\underline{\zeta}$  and  $\underline{\eta}$ . Then the coordinate ring  $\mathbb{C}[\ddot{X}_{\beta\underline{\eta}}^{\underline{\beta}\underline{\zeta}}]$  is isomorphic to  $\mathcal{U}(\ddot{\mathbf{t}}(\underline{\mathbf{i}}))$  for  $\mathbb{k} = \mathbb{C}$  by [SW21, Theorem 1.1]. Similarly, let  $\dot{X}_{\beta\underline{\eta}}^{\underline{\beta}\underline{\zeta}}$  denote the associated (half-decorated) double Bott-Samelson cell (dBS for short). Then  $\mathbb{C}[\dot{X}_{\beta\underline{\eta}}^{\underline{\beta}\underline{\zeta}}]$  is isomorphic to  $\mathcal{U}(\dot{\mathbf{t}}(\underline{\mathbf{i}}))$  for  $\mathbb{k} = \mathbb{C}$ , see [SW21, Section 2.4]. Theorems 6.13 and 6.14 imply the following.

**Theorem 7.1.** Theorems 1.1 and 1.6 hold for the coordinate rings of the decorated (or half-decorated) double Bott-Samelson cells.

We will mainly consider  $\mathcal{U}(\dot{\mathfrak{t}}(\underline{\zeta}))$  arising from  $\dot{X}_{\beta_{\underline{\zeta}}}^{e}$ , where e denotes the identity. For the reader's convenience, let us briefly recall  $\dot{X}_{\beta_{\underline{\zeta}}}^{e}$  for C of finite type following [CGG<sup>+</sup>22]. Let G be a connected, simply connected, complex semisimple algebraic group. We choose  $B_{\pm}$  its Borel subgroups, unipotent subgroup  $U_{\pm} = [B_{\pm}, B_{\pm}]$ , and maximal torus  $T = B_{+} \cap B_{-}$ . We fix pinning of the group G, i.e., for each  $a \in J$ , we select an isomorphism  $\varphi_{a} : \mathrm{SL}_{2}(\mathbb{C}) \to U_{a}$ , where  $U_{a}$  is the corresponding root subgroup. Define  $s_{a} := \varphi_{a} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ . For  $z \in \mathbb{C}$ , define  $B_{a}(z) := s_{a}\varphi_{a} \begin{pmatrix} 1 & z \\ 0 & 1 \end{pmatrix} = \varphi_{a} \begin{pmatrix} z & -1 \\ 1 & 0 \end{pmatrix}$ . For  $\underline{z} = (z_{k})_{k \in [1,l]} \in \mathbb{C}^{l}$ , define  $B_{\beta_{\underline{\zeta}}}(z) := B_{\zeta_{1}}(z_{1}) \cdots B_{\zeta_{l}}(z_{l})$ . Then we define the double Bott-Samelson variety as  $\dot{X}_{\beta_{c}}^{e} = \{\underline{z} \in \mathbb{C}^{l} | B_{\beta_{\zeta}}(z) \in B_{-}B_{+}\}$ .

7.2. Braid varieties. Assume C is of finite type. We briefly recall the braid varieties following [CGG<sup>+</sup>22, Section 3.3, Corollary 3.7]. Let  $\delta$  denote the Demazure product map from Br<sup>+</sup> to W, satisfying

$$\delta(\sigma_a) = s_a, \text{ and } \delta(\beta\sigma_a) = \begin{cases} \delta(\beta)s_a & \text{if } l(\delta(\beta)s_a) = l(\delta(\beta)) + 1\\ \delta(\beta) & \text{if } l(\delta(\beta)s_a) = l(\delta(\beta)) - 1 \end{cases}$$

Take any word  $\underline{\eta} = (\eta_1, \ldots, \eta_l)$ . Let  $\overrightarrow{w}_0$  denote any reduced word for the longest element  $w_0$  of W. We define the braid variety  $X(\beta)$  as  $\{\underline{z} \in \mathbb{C}^l | \delta(\beta_{\overrightarrow{w}_0})^{-1} B_{\beta_{\underline{\zeta}}}(\underline{z}) \in B\}$ . It is a smooth, irreducible affine variety of dimension  $l(\beta) - l(\delta(\beta))$ . Moreover, we have  $X(\beta) \simeq X(\beta\sigma_a)$  if  $\delta(\beta\sigma_a) = \delta(\beta)s_a$  ([CGG<sup>+</sup>22, Lemma 3.4(1)]). [CGG<sup>+</sup>22, Theorem 1.1] shows that  $\mathbb{C}[X(\beta)]$  is isomorphic to the cluster algebra  $\mathcal{U}(\mathbf{t}_{\overline{\mathfrak{m}}}(\underline{\eta}))$  for a specific seed  $\mathbf{t}_{\overline{\mathfrak{m}}}(\eta)$ .

By [CGG<sup>+</sup>22, Lemma 3.16], we can naturally identify  $X(\beta_{\vec{w}_0}\beta_{\underline{\eta}})$  with the dBS  $\dot{X}_{\beta_{\underline{\eta}}}$ . By [CGG<sup>+</sup>22, Corollary 4.42, Proposition 5.7], we further have an isomorphism  $\mathbb{C}[X(\beta_{\vec{w}_0}\beta_{\eta})] \simeq \mathbb{C}[\dot{X}_{\beta_{\eta}}]$  identifying  $\mathbf{t}_{\widehat{\mathbf{tm}}}(\vec{w}_0,\underline{\eta})$  with  $\dot{\mathbf{t}}(\underline{\eta})$ .

**Lemma 7.2.** Choose  $\vec{w}_0 = (\vec{w}'', \vec{w}')$ , such that  $\vec{w}'$  is a reduced word for  $w_0 \delta(\beta)^{-1}$ . Then the seed  $\mathbf{t}_{\underline{\mathfrak{m}}}(\underline{\eta})$  could be obtained from  $\mathbf{t}_{\underline{\mathfrak{m}}}(\vec{w}_0, \underline{\eta})$  by repeating the process: freezing and then removing a non-essential frozen vertex.

*Proof.* By [CGG<sup>+</sup>22, Lemma 4.44], for any word  $\underline{\zeta}$ , if  $\delta(\sigma_a\beta_{\underline{\zeta}}) = \delta(\beta_{\underline{\zeta}})$ ,  $\mathbf{t}_{\overline{\mathfrak{m}}}(a,\underline{\zeta})$  has an optimized frozen vertex j, such that  $\mathbf{t}_{\overline{\mathfrak{m}}}(\underline{\zeta})$  can be obtained from  $\mathbf{t}_{\overline{\mathfrak{m}}}(a,\underline{\zeta})$  by freezing all vertices incident to j and then removing the non-essential frozen vertex j; otherwise,  $\delta(\sigma_a\beta_{\zeta}) = s_a\delta(\beta_{\zeta})$  and  $\mathbf{t}_{\overline{\mathfrak{m}}}(a,\zeta) = \mathbf{t}_{\overline{\mathfrak{m}}}(\zeta)$ .

Since  $\delta(\beta_{\overrightarrow{w}},\beta) = w_0$ , we can obtain  $\mathbf{t}_{\overleftarrow{\mathfrak{m}}}(\overrightarrow{w}',\underline{\eta})$  by repeatedly applying this process to  $\mathbf{t}_{\overleftarrow{\mathfrak{m}}}(\overrightarrow{w}_0,\underline{\eta})$ . Since  $\overrightarrow{w}'$  is a reduced word for  $w_0\delta(\beta)^{-1}$ , we can identify  $\mathbf{t}_{\overleftarrow{\mathfrak{m}}}(\overrightarrow{w}',\underline{\eta})$ with  $\mathbf{t}_{\overleftarrow{\mathfrak{m}}}(\eta)$ . The claim follows.

Applying operations to  $\mathcal{U}(\mathbf{t}_{\overline{\mathfrak{m}}}(\overline{w}_0, \underline{\eta})) = \mathcal{U}(\mathbf{t}(\underline{\eta}))$  following Lemma 7.2. Theorems 6.13 6.14 and Corollaries 3.16 4.12 imply the following.

**Theorem 7.3.** Theorems 1.1 and 1.6 hold for the (quantum) cluster algebra  $\mathcal{U}(\mathbf{t}_{\mathrm{in}}(\eta))$ .

Note that Theorem 7.3 apply to open Richardson varieties as well, because they are special cases of braid varieties  $[CGG^+22, Section 3.6]$ .

7.3. Double Bruhat cells. Assume that C is of finite type. Take any pair of reduced words  $\underline{\zeta}, \underline{\eta}$  and denote  $u := w_{\underline{\zeta}}, v := w_{\underline{\eta}}$ . Define the double Bruhat cell  $G^{u,v} = B_+ u B_+ \cap B_- v B_-$ . The quantum double Bruhat cell  $\mathcal{O}_q[G^{u,v}]$  is isomorphic to the quantum cluster algebra  $\mathcal{U}(\mathbf{t}_{\mathrm{BZ}}(\underline{\mathbf{i}}))$ , where  $\underline{\mathbf{i}} = (\underline{\eta}, -\underline{\zeta})$  and  $\mathbf{t}_{\mathrm{BZ}}(\underline{\mathbf{i}})$  denote the Berenstein-Zelevinsky quantum seed in [BZ05], see [GY20, Section 3.3 Theorem 9.5] for details. Recall that  $\widetilde{B}(\mathbf{t}_{\mathrm{BZ}}(\underline{\mathbf{i}})) = -\widetilde{B}(\mathbf{\ddot{t}}(\underline{\mathbf{i}}))$  by Lemma 6.3. So we can identify  $\mathcal{U}(\mathbf{t}_{\mathrm{BZ}}(\underline{\mathbf{i}}))$  with  $\mathcal{U}(\mathbf{\ddot{t}}(\underline{\mathbf{i}}))$ . Then Theorems 6.13 and 6.14 imply the following.

**Theorem 7.4.** Theorems 1.1 and 1.6 hold for the quantum double Bruhat cells.

**Example 7.5.** Take  $G = SL_3$ ,  $\underline{\eta} = \underline{\zeta} = (1, 2, 1)$ ,  $\underline{\mathbf{i}} = (\underline{\eta}, -\underline{\zeta})$ . Then  $\mathcal{O}_q[G^{w_0, w_0}]$  is isomorphic to  $\mathcal{U}(\mathbf{t}_{\mathrm{BZ}}(\underline{\mathbf{i}}))$ . See Figure 6.1 for the quiver of  $-\widetilde{B}(\mathbf{t}_{\mathrm{BZ}}(\underline{\mathbf{i}})) = \widetilde{B}(\mathbf{\ddot{t}}(\underline{\mathbf{i}}))$ .

7.4. Algebraic groups. Assume that C is of finite type. Then G coincides with the closure  $\overline{G^{w_0,w_0}}$ , where  $w_0$  denote the longest element in W. Let  $\underline{\eta}, \underline{\zeta}$  denote any two chosen reduced words for  $w_0$ . Recall that the quantum double Bruhat cell  $\mathcal{O}_q[G^{w_0,w_0}]$  is isomorphic to  $\mathcal{U}(\mathbf{\ddot{t}}(\mathbf{\underline{i}}))$ , where  $\mathbf{\underline{i}} = (\underline{\eta}, -\underline{\zeta})$ . We claim the following natural statements are true.

Claim 7.6. (1)  $\mathbb{C}[G]$  is isomorphic to  $\overline{\mathcal{U}}(\mathbf{\dot{t}}(\mathbf{\underline{i}})) \otimes \mathbb{C}$  for  $\mathbb{k} = \mathbb{Z}$ . (2)  $\mathcal{O}_q[G] \otimes \mathbb{Q}(q^{\frac{1}{2}})$  is isomorphic to  $\overline{\mathcal{U}}(\mathbf{\ddot{t}}(\mathbf{\underline{i}})) \otimes \mathbb{Q}(q^{\frac{1}{2}})$  for  $\mathbb{k} = \mathbb{Z}[q^{\pm \frac{1}{2}}]$ .

Proofs will appear elsewhere. Here, the quantized coordinate ring  $\mathcal{O}_q[G]$  is defined as the quantum function algebra spanned by the matrix coefficients, see [Kas93]. Assuming these claims, Theorem 6.17 implies the following result.

**Theorem 7.7.** Theorem 1.1 holds for the (quantized) coordinate rings of G.

**Example 7.8.** [Kas93] provides an explicit calculation for  $\mathcal{O}_q[SL_2]$ , which has four generators u, y, v, x such that  $y * x = 1 + q^{-1}uv$ . It is isomorphic to a quantum cluster algebra  $\overline{\mathcal{U}}(\mathbf{t})$  for  $\mathbf{t} = \ddot{\mathbf{t}}((1,-1))$ , where  $\ddot{I} = \{-1,1,2\}$ , such that we send u, y, v, x to  $x_{-1}(\mathbf{t}), x_1(\mathbf{t}), x_2(\mathbf{t})$ , and  $x_1(\mu_1 \mathbf{t})$ . Its global crystal basis consists of the cluster monomials up to  $q^{\frac{Z}{2}}$  multiples, while the common triangular basis consists of the cluster monomials.

7.5. Other coordinate rings. Recall that open positroid varieties are special cases of double Bott-Samelson cells. So Theorems 7.1 and 1.6 apply to them.

Recall that the coordinate ring of a Grassmannian is a cluster algebra with a distinguished seed  $\mathbf{t}_{\mathsf{Gr}}$  [Sco06]. Note that  $\mathbf{t}_{\mathsf{Gr}}$  is similar to  $\dot{\mathbf{t}}(\underline{\gamma})$  for some adapted word  $\underline{\gamma}$  in type A. It is already known that  $\dot{\mathbf{t}}(\underline{\gamma})$  has the common triangular basis **L** and admits categorification by [Qin17], see Theorem 7.10. Combining with Theorem 6.14 and Corollary 4.12, we deduce that Theorems 1.1 and 1.6 hold for  $\mathcal{U}(\mathbf{t}_{\mathsf{Gr}})$ .

7.6. Cluster structures from quantum affine algebras. Choose a Coxeter word  $\underline{c} = (c_1, \ldots, c_{|J|})$ , i.e., each  $a \in J$  appears in it exactly once. Note that  $\underline{c}$  determines a skew-symmetric  $J \times J$ -matrix  $B^{\Delta}$  such that  $B^{\Delta}_{c_j,c_k} := \operatorname{sign}(C_{c_j,c_k}) \in \{0,-1\}$  for j < k. A letter  $c_k$  is called a sink of  $\underline{c}$  if  $C_{c_k,c_j} \ge 0 \forall j < k$ .

We introduce  $\mathbb{W} := \{ \langle_d^a \rangle | a \in J, d \in \mathbb{Z} \}$  as in [KQ14]. Endow it with the order < such that  $\langle_d^{c_j} \rangle < \langle_r^{c_k} \rangle$  if d < r or d = r but j < k. Let  $e_{\langle_d^a \rangle}$  denote the  $\langle_d^a \rangle$ -th unit vector of  $\mathbb{N}^{\mathbb{W}}$ .


For any sink letter  $a \in [1, |J|]$ , we define the mutation  $\mu_{a\underline{c}} := (\underline{c} \setminus \{a\}, a)$ . Choose any <u>c</u>-adapted word  $\underline{\gamma} = (\gamma_i)$ , i.e.,  $\gamma_i$  is a sink of  $\mu_{\gamma_{i-1}} \cdots \mu_{\gamma_1} \underline{c}$ ,  $\forall i$ . Define  $\beta_k := \beta_k(\underline{\gamma}) := e_{\langle a \ \gamma_k \ (\gamma_k) \rangle}$ .

Assume that the quiver associated with  $B^{\Delta}$  is a tree. Then we can choose a  $\mathbb{Z}$ -valued function  $\xi$  on J, such that  $\xi_{c_k} = \xi_{c_1} - 1$  when  $C_{c_j,c_k} < 0$  and k > 1. Denote  $J_{\mathbb{Z}}(\xi) := \{(a,p) | a \in J, p \in \xi_a + 2\mathbb{Z}\}$ . In this case, we identify  $\mathbb{W}$  with  $J_{\mathbb{Z}}(\xi)$  such that  $\langle a \\ \rangle$  corresponds to  $(a, \xi_a - 2d)$ . Then we can denote  $e_{(a,\xi_a-2d)}$  by  $e_{\langle a \\ \langle d \rangle}$ .

**Example 7.9.** Assume  $C = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$ . We choose  $\underline{c} = (1, 2), \underline{\gamma} = (1, 2, 1, 2, 1, 2)$ . The quiver for  $\dot{\mathbf{t}}(\underline{\gamma})$  is drawn in Figure 7.1, where we denote elements of  $I(\dot{\mathbf{t}}(\underline{\gamma}))$  by  $k \in [1, 6], \langle_a^a \rangle \in \mathbb{W}$ , or  $(a, \xi_a - 2d) \in J_{\mathbb{Z}}(\xi)$ .

In this case, the letter 1 is a sink,  $\xi_2 = \xi_1 - 1$  and  $\mu_1 \underline{c} = (2, 1)$ .

FIGURE 7.1. The quiver for  $\dot{t}(1, 2, 1, 2, 1, 2)$ 



From now on, assume C to be of type ADE. Let  $U_{\varepsilon}(\widehat{\mathfrak{g}})$  denote the associated quantum affine algebra, where  $\varepsilon \in \mathbb{C}^{\times}$  is not a root of unity. It has distinguished finite dimensional simple modules  $L_{a,\varepsilon'}, \forall a \in J, \varepsilon' \in \mathbb{C}^{\times}$ , called fundamental modules. Denote  $L(e_{\langle a \\ -d \rangle}) := L_{a,\varepsilon^{\langle a+2d \rangle}}$ . The composition factors of the tensor products of  $L(e_{\langle a \\ -d \rangle})$  could be parameterized by L(w), for  $w \in \bigoplus \mathbb{N}e_{\langle a \\ -d \rangle}$ . We refer the reader to  $[\operatorname{Nak04}][\operatorname{Nak04}][\operatorname{HL10}]$  for details.

Following [HL10], let  $\mathcal{C}_{\mathbb{Z}}(\xi)$  denote the monoidal category whose objects have these composition factors L(w). Let  $\mathcal{C}_{\underline{\gamma}}(\xi)$  denote the monoidal subcategory of  $\mathcal{C}_{\mathbb{Z}}(\xi)$  whose objects have the composition factors L(w), for  $w \in \bigoplus_k \mathbb{N}\beta_k$ .  $\forall j \leq k$ ,  $\gamma_j = \gamma_k, L(\beta_{[j,k]})$  are called the Kirillov-Reshetikhin modules, where  $\beta_{[j,k]} := \beta_j + \beta_{j[1]} + \cdots + \beta_k$ . Its deformed Grothendieck ring, denoted by K, is constructed from quiver varieties [Nak04][VV03][Qin17, Sections 7, 8.4].

By [Qin17, Theorem 8.4.3], we have an isomorphism  $\kappa : \overline{\mathcal{A}}(\mathbf{t}(\underline{\gamma})) \simeq K$ , such that  $\kappa(x_i(\mathbf{t}(\underline{\gamma}))) = [L(\beta_{[i^{\min},i]})]$ , where the quantization of K and  $\mathbf{t}(\underline{\gamma})$  are chosen as in [Qin17, Section 7.3].

**Theorem 7.10** ([Qin17, Theorem 1.2.1(II)]).  $\overline{\mathcal{A}}(\dot{\mathbf{t}}(\underline{\gamma}))$  has the common triangular basis **L**. And  $(\overline{\mathcal{A}}(\dot{\mathbf{t}}(\underline{\gamma})), \mathbf{L})$  is categorified by  $C_{\underline{\gamma}}(\xi)$ , such that  $\kappa \mathbf{L}_m = [L(w)]$ , where  $m = (m_i)_{i \in [1,l]}$  satisfies  $\sum_i m_i \beta_{[i^{\min},i]} = \sum_i w_i \beta_i$ .

Now we have a new proof for Theorem 7.10: The first statement is a special case of Theorem 6.17. Combining with [Qin17, Corollary 9.1.9], it implies the second statement.

Denote  $\underline{c}^l := (\underline{c}, \underline{c}, \dots, \underline{c})$  for *l*-copies of  $\underline{c}$ . Particularly, Theorem 7.10 implies the conjecture of Hernandez of Leclerc that  $C_{\underline{c}^l}(\xi)$  categorifies  $\overline{\mathcal{A}}(\dot{\mathbf{t}}(\underline{c}^l))$  (Definition 5.3).

# 8. Applications: cluster algebras from double Bott-Samelson cells

Take any word  $\underline{\eta}$ . Denote  $l = l(\underline{\eta})$ ,  $\mathbf{\dot{t}} = \mathbf{\dot{t}}(\underline{\eta})$ . We will explore rich structures of the quantum cluster algebra  $\overline{\mathcal{U}}(\mathbf{\dot{t}})$  arising from the double Bott-Samelson cell  $\dot{X}_{\beta_n}^e$ .

8.1. Mutation sequence  $\Sigma$ .  $\forall j \leq k \in [1, l] \simeq I(\underline{\eta}) = I$ ,  $\eta_j = \eta_k$ , define  $\boldsymbol{\mu}_{[j,k]} := \mu_k \cdots \mu_{j[1]} \mu_j$ ,  $\Sigma_k := \Sigma_k(\underline{\eta}) := \boldsymbol{\mu}_{[k^{\min},k^{\min}[o+(k)-1]]} = \boldsymbol{\mu}_{[k^{\min},k^{\max}[-1-o-(k)]]}$ ,  $\Sigma := \Sigma(\underline{\eta}) := \Sigma_l \cdots \Sigma_2 \Sigma_1$ , where the mutation sequences  $\Sigma_{k^{\max}}$  are the identity.

Recall that  $k \in [1, l]$  is identified with  $\binom{\eta_k}{o_{-}(k)}^{\underline{\eta}}$ , abbreviated as  $\binom{\eta_k}{o_{-}(k)}$ . Denote  $\binom{a}{-1-r} = -\infty$ ,  $\binom{a}{O(a)+r} = +\infty$  for  $r \in \mathbb{N}$ . The order < on [1, l] induces the order  $<_{\eta}$  on I, abbreviated as <.

Denote  $\dot{\mathbf{t}} := \dot{\mathbf{t}}(\underline{\eta})$ . When we apply  $\Sigma$  to  $\dot{\mathbf{t}}$ , the seeds appearing can be described as  $\dot{\mathbf{t}}\{r\}_s$ : For any  $r \in [0, l]$ , denote  $\dot{\mathbf{t}}\{0\} := \dot{\mathbf{t}}, \dot{\mathbf{t}}\{r\} := \Sigma_r \cdots \Sigma_2 \Sigma_1 \dot{\mathbf{t}}$ . Further denote  $\dot{\mathbf{t}}\{r\}_0 := \dot{\mathbf{t}}\{r\}, r' := r + 1$ , and  $\dot{\mathbf{t}}\{r\}_s := \boldsymbol{\mu}_{[r',r'[s-1]]}\dot{\mathbf{t}}\{r\}$  for  $s \in [0, o_+(r')]$ . Note that  $\dot{\mathbf{t}}\{r\}_{o_+(r')} = \dot{\mathbf{t}}\{r'\}_0$ .

 $\begin{array}{l} \text{Lemma 8.1. } (1) \ \dot{\mathbf{t}}\{r\} = \dot{\mathbf{t}}(\underline{\eta}_{[r+1,l]}, -\underline{\eta}_{[1,r]}^{\mathsf{op}}). \\ (2) \ \dot{\mathbf{t}}\{r\}_s = \dot{\mathbf{t}}(\underline{\eta}_{[r'+1,r'[s]]}, -\eta_{r'}, \underline{\eta}_{[r'[s]+1,l]}, -\underline{\eta}_{[1,r]}^{\mathsf{op}}). \\ (3) \ Take \ any \ \dot{\mathbf{t}}\{r\}_s \neq \Sigma \dot{\mathbf{t}}. \ Denote \ a := \eta_{r'}. \ Then \ we \ have \ b_{\binom{a}{s-1},\binom{a}{s}}(\dot{\mathbf{t}}\{r\}_s) = \\ b_{\binom{a}{s+1},\binom{a}{s}}(\dot{\mathbf{t}}\{r\}_s) = -1, \ b_{\binom{b}{d},\binom{a}{s}}(\dot{\mathbf{t}}\{r\}_s) = -C_{ba} \ if \ b \neq a \ and \ \binom{b}{d+O([1,r];b)} < \underline{\eta} \\ \binom{b}{s+O([1,r];a)+1} < \underline{\eta} \ \binom{b}{d+O([1,r];b)+1}, \ and \ b_{j,\binom{a}{s}}(\dot{\mathbf{t}}\{r\}_s) = 0 \ for \ other \ j \in I. \end{array}$ 

*Proof.* (1)(2) Consider the seeds appearing when we apply  $\Sigma_1$  to  $\dot{\mathbf{t}} = \dot{\mathbf{t}}\{0\}$ . In this case, r' = 1. We rewrite  $\dot{\mathbf{t}}(\eta_1, \eta_{[2,l]}) = \dot{\mathbf{t}}(-\eta_1, \eta_{[2,l]})$  by the left reflection. Applying  $\Sigma_1$  to  $\dot{\mathbf{t}}(-\eta_1, \eta_{[2,l]})$  via flips (Section 6.3), we get  $\dot{\mathbf{t}}_s = \dot{\mathbf{t}}(\underline{\eta}')$  for  $\underline{\eta}' = (\underline{\eta}_{[2,1[s]]}, -\eta_1, \underline{\eta}_{[1[s]+1,l]})$ . Next, use the reflection  $\dot{\mathbf{t}}(\eta_2, \eta_{[3,l]}, -\eta_1) = \dot{\mathbf{t}}(-\eta_2, \eta_{[3,l]}, -\eta_1)$  and apply  $\Sigma_2$  to  $\dot{\mathbf{t}}(-\eta_2, \eta_{[3,l]}, -\eta_1)$  via flips. Repeat the previous arguments, we obtained the desired claims.

(3) We prove the claim by induction on l. The case l = 1 is trivial. Assume they have been proved for lengths  $\leq l - 1$ .

Take r = 0. Consider  $\dot{\mathbf{t}}_s = \dot{\mathbf{t}}(\underline{\eta}')$  where  $\underline{\eta}' = (\underline{\eta}_{[2,1[s]]}, -\eta_1, \underline{\eta}_{[1[s]+1,l]}), 1[s] = \binom{\eta_1}{s}$ . So the  $\binom{\eta_1}{s}$ -th triangle  $T'_{\binom{\eta_1}{s}}$  for  $\underline{\eta}'$  satisfies  $\mathbf{i}_{T'_{\binom{\eta_1}{s}}} = -\eta_1$ , and  $\binom{\eta_1}{s}$  is the interval  $\ell_{\eta_1}^+$  incident to its interior point. (6.1) implies  $b_{\binom{\eta_1}{s-1},\binom{\eta_1}{s}}(\dot{\mathbf{t}}_s) = b_{\binom{\eta_1}{s+1},\binom{\eta_1}{s}}(\dot{\mathbf{t}}_s) = -1$ ,  $b_{\binom{b}{d},\binom{\eta_1}{s}}(\dot{\mathbf{t}}_s) = -C_{b\eta_1}$  if  $b \neq \eta_1$ ,  $\binom{b}{d} < \binom{\eta_1}{s} < \binom{\eta_1}{s+1} < \binom{b}{d+1}$  or  $\binom{\eta_1}{s} < \binom{b}{d} < \binom{\eta_1}{s+1} < \binom{b}{d} < \binom{\eta_1}{s+1} < \binom{b}{d+1}$ , and  $b_{j,\binom{\eta_1}{s}} = 0$  for other  $j \in I$ .


Therefore the claim is true for r = 0. Particularly, it holds for  $\dot{\mathbf{t}}' := \dot{\mathbf{t}}\{1\} = \dot{\mathbf{t}}(\underline{\eta}_{[2,l]}, -\eta_1)$ , and  $b_{1^{\max},k}(\dot{\mathbf{t}}') = 0$  for  $k \in I_{\mathrm{uf}} \setminus \{1^{\max}[-1]\}$ . Denote  $\mathbf{u} := \dot{\mathbf{t}}(\underline{\eta}_{[2,l]})$ . By Lemma 6.6, we have the cluster embedding  $\iota$  from  $\mathbf{u}$  to  $\dot{\mathbf{t}}$  such that  $\iota\binom{b}{d}^{\mathbf{u}} = \begin{pmatrix} b \\ d+\delta_{b,\eta_1} \end{pmatrix}$  and the cluster embedding  $\iota'$  from  $\mathbf{u}$  to  $\dot{\mathbf{t}}'$  such that  $\iota\binom{b}{d}^{\mathbf{u}} = \begin{pmatrix} b \\ d \end{pmatrix}$ . Recall that  $\binom{b}{d}^{\mathbf{u}} \leq \frac{\eta}{d} \leq \binom{c}{h}$ .

Note that the mutation sequence  $\Sigma' := \Sigma_l \cdots \Sigma_2$  does not act on  $1^{\max}[-1]$ . So we have  $b_{1^{\max},k}(\mathbf{t}\{r\}_s) = 0$ ,  $\forall \mathbf{t}\{r\}_s$ ,  $k \in I_{\mathsf{uf}} \setminus \{1^{\max}[-1]\}$ . Moreover,  $\Sigma'$  is identified with the mutation sequence  $\Sigma(\underline{\eta}_{[2,l]})$  via  $\iota'$ . Let  $\boldsymbol{\mu}$  denote any proper sub sequence of  $\Sigma(\underline{\eta}_{[2,l]})$ , such that  $\Sigma(\underline{\eta}_{[2,l]}) = \cdots = \mu_{k_{h+1}} \mu_{k_h} \cdots = \mu_{k_{h+1}} \mu$ . Then  $\boldsymbol{\mu}\mathbf{u}$  takes the form  $\mathbf{u}\{r\}_s$ , and  $\iota'$  is a cluster embedding from  $\boldsymbol{\mu}\mathbf{u}$  to  $(\iota'\boldsymbol{\mu})\mathbf{t}' = \mathbf{t}\{r+1\}_s$ .

Note that  $O^{\underline{n}_{[2,l]}}([1,r];b) = O([2,r+1];b)$ . By the induction hypothesis on  $\mathbf{u}$  and using  $\iota'$ , we obtain the following claim for  $\dot{\mathbf{t}}\{r+1\}_s$ :  $b_{\binom{a}{s-1},\binom{a}{s}}(\dot{\mathbf{t}}\{r+1\}_s) = b_{\binom{a}{s+1},\binom{a}{s}}(\dot{\mathbf{t}}\{r+1\}_s) = -1, b_{\binom{b}{d},\binom{a}{s}}(\dot{\mathbf{t}}\{r+1\}_s) = -C_{ba}$  if  $b \neq a, \binom{b}{d+O([2,r+1];b)} < \underline{n}_{[2,l]}$  $\binom{a}{(s+O([2,r+1];a)+1)} < \underline{n}_{[2,l]} \binom{b}{(d+O([2,r+1];b)+1)}$ , and  $b_{j,\binom{a}{s}}(\dot{\mathbf{t}}\{r+1\}_s) = 0$  for other  $j \in \iota'I(\mathbf{u}) = I \setminus \{1^{\max}\}$ . In addition, we already know  $b_{1^{\max},\binom{a}{s}}(\dot{\mathbf{t}}\{r+1\}_s) = 0$ . Since  $\iota, \iota'$  are cluster embeddings, we have  $b(\dot{\mathbf{t}}')_{j,k} = b_{(\iota')^{-1}j,(\iota')^{-1}k}(\mathbf{u}) = b_{\iota(\iota')^{-1}j,(\iota')^{-1}k}(\dot{\mathbf{t}})$  for  $j, k \in (\iota')I(\mathbf{s})$ . Then we obtain the desired claim by applying  $\iota(\iota')^{-1}$  and using  $\iota(\iota')^{-1}\binom{b}{d} = \binom{b}{d+\delta_{b,\eta_1}}$ .

**Example 8.2.** Take  $C = \begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}$ ,  $\underline{\eta} = (1, 2, 1, 1, 2, 2, 1)$ ,  $\dot{\mathbf{t}} = \dot{\mathbf{t}}(\underline{\eta})$ . We have  $\Sigma = \Sigma_5 \Sigma_4 \Sigma_3 \Sigma_2 \Sigma_1 = \mu_2 \mu_1 \mu_{[1,3]} \mu_{[2,5]} \mu_{[1,4]} = \mu_2 \mu_1 (\mu_3 \mu_1) (\mu_5 \mu_2) (\mu_4 \mu_3 \mu_1)$  (read from right to left). We can check Lemma 8.1 directly. See Figure 8.1 for some  $\ddot{B}(\dot{\mathbf{t}}\{r\})$ .

FIGURE 8.1. Quivers with weights associated with  $\ddot{B}$ .



8.2. Interval variables and fundamental variables.

**Definition 8.3.** Consider the seeds  $\dot{\mathbf{t}}\{r\}$ ,  $r \in [0, l]$ .  $\forall i \in I, a \in J$ , denote  $r_a = O([1, r]; a)$  and  $r_i := r_{\eta_i}$ . Then,  $\forall i = i^{\min}$ ,  $d < O(\eta_i) - r_i$ ,  $W_{[i[r_i], i[r_i+d]]}(\dot{\mathbf{t}}) := x_{i[d]}(\dot{\mathbf{t}}\{r\})$  are called the interval variables of  $\overline{\mathcal{A}}(\dot{\mathbf{t}}(\underline{\eta}))$ .  $W_j(\dot{\mathbf{t}}\{r\}) := W_{[j,j]}(\dot{\mathbf{t}}\{r\})$  are called the fundamental variables.<sup>3</sup>

Equivalently,  $W_{[\binom{a}{r_a},\binom{a}{r_a+d}]}(\dot{\mathbf{t}}) = x_{\binom{a}{d}}(\dot{\mathbf{t}}\{r\})$  for  $d < O(a) - r_a$ . We often omit  $\dot{\mathbf{t}}$  for simplicity. The degrees  $\beta_{[j,k]} := \deg^{\mathbf{t}} W_{[j,k]}, \beta_j := \beta_{[j,j]}$ , have the following surprisingly nice description.

**Lemma 8.4.** We have  $\beta_{[j,k]} = f_k - f_{j[-1]}$ , where we denote  $f_{\pm \infty} = 0$ .

An example is given as Example 8.9. Before proving Lemma 8.4, let us make some preparation.

Given any  $1 \leq j \leq k \leq l$ . By Lemma 6.6,  $\mathbf{u} := \dot{\mathbf{t}}(\underline{\eta}_{[j,k]})$  is a good sub seed of  $\dot{\mathbf{t}}' := \dot{\mathbf{t}}(\underline{\eta}_{[j,l]}, -\underline{\eta}_{[1,j-1]}^{\mathsf{op}})$  via the cluster embedding  $\iota'$  such that  $\iota'\binom{a}{d}^{\mathbf{u}} = \binom{a}{d}$ , where  $\binom{a}{d}^{\mathbf{u}} \in I(\mathbf{u})$  and  $\binom{a}{d} \in I(\dot{\mathbf{t}}') = I(\dot{\mathbf{t}})$ . Then we obtain the following inclusion of algebras:

(8.1) 
$$\iota_{[j,k]}: \overline{\mathcal{A}}(\dot{\mathbf{t}}(\underline{\eta}_{[j,k]})) \stackrel{\iota'}{\hookrightarrow} \overline{\mathcal{A}}(\dot{\mathbf{t}}(\underline{\eta}_{[j,l]}, -\underline{\eta}_{[1,j-1]}^{\mathsf{op}}) \stackrel{(\Sigma_{j-1}\cdots\Sigma_1)^*}{\simeq} \overline{\mathcal{A}}(\dot{\mathbf{t}}(\underline{\eta}))$$

such that  $\iota_{[j,k]}x_h(\mathbf{u}) = x_{\iota'(h)}(\dot{\mathbf{t}}'), \forall h \in I(\mathbf{u})$ , where we omit the symbol  $(\Sigma_{j-1}\cdots\Sigma_1)^*$ as usual. For any mutation sequence  $\boldsymbol{\mu}$  on  $I_{uf}(\mathbf{u})$ , let  $\iota'\boldsymbol{\mu}$  denote the corresponding sequence on  $I_{uf}(\dot{\mathbf{t}}')$ . Then we obtain  $\iota_{[j,k]}x_h(\boldsymbol{\mu}\mathbf{u}) = x_{\iota'(h)}((\iota'\boldsymbol{\mu})\dot{\mathbf{t}}')$ . Particularly, by taking  $\boldsymbol{\mu}$  such that  $\Sigma_l\cdots\Sigma_{k+1}=\cdots\mu_{k_{d+1}}\mu_{k_d}\cdots\mu_{k_1}=\cdots\mu_{k_{d+1}}\boldsymbol{\mu}$ , we deduce the following result by definitions of the interval variables.

**Lemma 8.5.** The inclusion  $\iota_{[j,k]} : \overline{\mathcal{A}}(\dot{\mathbf{t}}(\underline{\eta}_{[j,k]})) \hookrightarrow \overline{\mathcal{A}}(\dot{\mathbf{t}}(\underline{\eta}_{[j,l]}, -\underline{\eta}_{[1,j-1]}^{\mathsf{op}})) = \overline{\mathcal{A}}(\dot{\mathbf{t}}(\underline{\eta}))$ sends the fundamental variables  $W_{[\binom{a}{d}^{\underline{\eta}_{[j,k]}},\binom{a}{d'}^{\underline{\eta}_{[j,k]}}]}(\dot{\mathbf{t}}(\underline{\eta}_{[j,k]}))$  to the fundamental variables  $W_{[\binom{a}{d+O([1,j-1];a]}),\binom{a}{d'+O([1,j-1];a])}]}(\dot{\mathbf{t}}(\underline{\eta})).$ 

*Proof of Lemma 8.4.* We prove the claim by induction on l. The case l = 1 is trivial. Assume it has been proved for lengths  $\leq l - 1$ .

Denote  $\mathbf{u} = \dot{\mathbf{t}}(\underline{\eta}_{[2,l]})$ . It is a good sub seed of  $\dot{\mathbf{t}}' := \dot{\mathbf{t}}\{1\} = \dot{\mathbf{t}}(\underline{\eta}_{[2,l]}, -\eta_1)$ . Then we have the cluster embedding from  $\mathbf{u}$  to  $\dot{\mathbf{t}}'$  sending  $\binom{a}{r}^{\mathbf{u}} \in I(\mathbf{u})$  to  $\binom{a}{r} \in I(\dot{\mathbf{t}}') = I(\dot{\mathbf{t}}')$ . By Lemma 8.5, the inclusion  $\iota_{[2,l]} : \overline{\mathcal{A}}(\mathbf{u}) \stackrel{\iota'}{\hookrightarrow} \overline{\mathcal{A}}(\dot{\mathbf{t}}') \simeq \overline{\mathcal{A}}(\dot{\mathbf{t}})$  sends  $W_{[\binom{a}{r})^{\mathbf{u}}, \binom{a}{r+d}^{\mathbf{u}}]}(\mathbf{u})$  to  $W_{[\binom{a}{r+d}, \eta_1+d)]}$ . The induction hypothesis implies  $\deg^{\mathbf{u}}(W_{[\binom{a}{r})^{\mathbf{u}}, \binom{a}{r+d}^{\mathbf{u}}]}(\mathbf{u})) = f_{\binom{a}{r+d}^{\mathbf{u}}}(\mathbf{u}) - f_{\binom{a}{r-1}^{\mathbf{u}}}(\mathbf{u})$ . Using the cluster embedding  $\iota'$ , we deduce

$$\deg^{\mathbf{t}'}(W_{[j,k]}) = \begin{cases} f_k(\mathbf{t}') - f_{j[-1]}(\mathbf{t}') & a \neq \eta_1, j = \binom{a}{r}, k = \binom{a}{r+d} \in I \\ f_{k[-1]}(\mathbf{t}') - f_{j[-2]}(\mathbf{t}') & r > 0, j = \binom{\eta_1}{r}, k = \binom{\eta_1}{r+d} \in I \end{cases}$$

Recall that  $\dot{\mathbf{t}}_s = \boldsymbol{\mu}_{[1,1[s-1]]} \dot{\mathbf{t}}$  and  $\dot{\mathbf{t}}_{o_+(1)} = \dot{\mathbf{t}}\{1\} = \dot{\mathbf{t}}'$ . For the composition of tropical mutations  $\phi_{\dot{\mathbf{t}}_0,\dot{\mathbf{t}}_{o_+(1)}} = \phi_{\dot{\mathbf{t}}_0,\dot{\mathbf{t}}_1} \circ \phi_{\dot{\mathbf{t}}_1,\dot{\mathbf{t}}_2} \circ \cdots \circ \phi_{\dot{\mathbf{t}}_{o_+(1)-1},\dot{\mathbf{t}}_{o_+(1)}}$ , we have

<sup>&</sup>lt;sup>3</sup>When C is of type ADE and  $\underline{\eta}$  is adapted,  $W_{[j,k]}$  and  $W_{[j,j]}$  correspond to the Kirillov-Reshetikhin modules and the fundamental modules of the quantum affine algebras respectively, whence the names. The fundamental variables might also be called root variables in analogous to the root vectors for PBW bases.

 $\deg^{\mathbf{t}_0} W_{[j,k]} = \phi_{\mathbf{t}_0, \mathbf{t}_{o_+}(1)} \deg^{\mathbf{t}_{o_+}(1)} W_{[j,k]}$ . If  $\eta_j = \eta_k \neq \eta_1$ , the vector  $f_k - f_{j[-1]}$  remains unchanged and the desired claim for  $W_{[j,k]}$  follows.

Next, compute  $\deg^{\mathbf{t}}(W_{[1[d],1[s]]}), \forall d \geq 1$ . Note that  $\dot{\mathbf{t}}_{s+1} = \mu_{1[s]}\dot{\mathbf{t}}_{s}, \forall s \in [0, o_{+}(1) - 1]$ . Lemma 8.1 implies  $b_{1[s-1],1[s]}(\dot{\mathbf{t}}_{s+1}) = b_{1[s+1],1[s]}(\dot{\mathbf{t}}_{s+1}) = 1$  and  $b_{i,1[s]}(\dot{\mathbf{t}}_{s+1}) \leq 0$  for  $\eta_{i} \neq \eta_{1}$ . So we deduce

$$\begin{aligned} \phi_{\dot{\mathbf{t}}_{s},\dot{\mathbf{t}}_{s+1}}(f_{1[s]}(\dot{\mathbf{t}}_{s+1})) &= f_{1[s+1]}(\dot{\mathbf{t}}_{s}) - f_{1[s]}(\dot{\mathbf{t}}_{s}) + f_{1[s-1]}(\dot{\mathbf{t}}_{s-1}) \\ \phi_{\dot{\mathbf{t}}_{s},\dot{\mathbf{t}}_{s+1}}(\pm f_{1[d]}(\dot{\mathbf{t}}_{s+1})) &= \pm f_{1[d]}(\dot{\mathbf{t}}_{s}), \ \forall d \neq s. \end{aligned}$$

Then direct computation shows the following results for  $0 < d \le s \le o_+(1)$ :

$$\begin{aligned} \deg^{\mathbf{t}_{s}} W_{[1[d],1[s]]} &= \phi_{\mathbf{t}_{s},\mathbf{t}_{s+1}} \cdots \phi_{\mathbf{t}_{o_{+}(1)-1},\mathbf{t}_{o_{+}(1)}} \left( \deg^{\mathbf{t}_{o_{+}(1)}} W_{[1[d],1[s]]} \right) \\ &= f_{1[s-1]}(\mathbf{t}_{s}) - f_{1[d-2]}(\mathbf{t}_{s}) \text{ (coordinates are unchanged)} \\ \\ \deg^{\mathbf{t}_{d-1}} W_{[1[d],1[s]]} &= \phi_{\mathbf{t}_{d-1},\mathbf{t}_{d}} \cdots \phi_{\mathbf{t}_{s-1},\mathbf{t}_{s}} \left( \deg^{\mathbf{t}_{s}} W_{[1[d],1[s]]} \right) \\ &= f_{1[s]}(\mathbf{t}_{d-1}) - f_{1[d-1]}(\mathbf{t}_{s-1}) \\ \\ \\ \deg^{\mathbf{t}_{0}} W_{[1[d],1[s]]} &= \phi_{\mathbf{t}_{0},\mathbf{t}_{1}} \cdots \phi_{\mathbf{t}_{d-2},\mathbf{t}_{d-1}} \left( \deg^{\mathbf{t}_{d-1}} W_{[1[d],1[s]]} \right) \\ &= f_{1[s]}(\mathbf{t}_{0}) - f_{1[d-1]}(\mathbf{t}_{0}) \text{ (coordinates are unchanged)} \end{aligned}$$

Finally, note that  $W_{[1,1[s]]}$  are cluster variable of  $\dot{\mathbf{t}}$ . Thus  $\deg^{\dot{\mathbf{t}}} W_{[1,1[s]]} = f_{1[s]} = f_{1[s]} = f_{1[s]} - f_{-\infty}$ . Therefore, we have verified the desired statement for all  $W_{[j,k]}$ .

Recall that we use  $[]^{\dot{\mathbf{t}}}$  to denote normalization in  $\mathcal{LP}(\dot{\mathbf{t}})$ , abbreviated as [].

**Proposition 8.6.**  $\forall r \in [0, l]$ , denote  $a = \eta_{r+1}$  and  $r_b = O([1, r]; b)$  for  $b \in J$ . The interval variables satisfy the following equations, called the *T*-systems:

$$(8.2) W_{[\binom{a}{r_a},\binom{a}{r_a+s}]} * W_{[\binom{a}{r_a+1},\binom{a}{r_a+s+1}]} = q^{\alpha} [W_{[\binom{a}{r_a+1},\binom{a}{r_a+s}]} * W_{[\binom{a}{r_a},\binom{a}{r_a+s+1}]} + q^{\alpha'} [\prod_{\binom{b}{r_b+d}} W_{[\binom{b}{r_b},\binom{b}{r_b+d}]}^{-C_{ba}}],$$

where  $\binom{b}{r_b+d}$  appearing satisfy  $b \neq a$ ,  $\binom{b}{r_b+d} < \underline{n} \binom{a}{r_a+s+1} < \underline{n} \binom{b}{r_b+d+1}$ , and  $\alpha = \frac{1}{2}\lambda(\beta_{[\binom{a}{r_a},\binom{a}{r_a+s}]}, \beta_{[\binom{a}{r_a+1},\binom{a}{r_a+s+1}]}), \alpha' = \frac{1}{2}\lambda(\beta_{[\binom{a}{r_a},\binom{a}{r_a+s}]}, -\sum_{\binom{b}{r_b+d}}C_{ba}\beta_{[\binom{b}{r_b},\binom{b}{r_b+d}]}).$ We have  $\alpha > \alpha'$ .

Equivalently,  $\forall 1 \leq j \leq j [s] \leq l$ , we can rewrite the *T*-systems as

(8.3) 
$$W_{[j,j[s]]} * W_{[j[1],j[s+1]]} = q^{\alpha} [W_{[j[1],j[s]]} * W_{[j,j[s+1]]}] + q^{\alpha'} [\prod W_{[i,i[d]]}^{-C_{\eta_i,\eta_j}}],$$
  
where  $[i, i[d]]$  appearing satisfy  $i = i^{\min}[O([1, j - 1]; \eta_i)], i[d] < j[s+1] < i[d+1].$ 

*Proof.* Denote  $\dot{\mathbf{t}}' = \dot{\mathbf{t}}\{r\}$  and  $\dot{\mathbf{t}}'' = \dot{\mathbf{t}}\{r+1\}$ . We use ' and " denote their associated data respectively. Recall that  $\mu_{\binom{a}{s}}\dot{\mathbf{t}}\{r\}_s = \dot{\mathbf{t}}\{r\}_{s+1}$  for r' = r+1,  $s < o_+(r')$ ,  $a = \eta_{r'}$ , and  $x'_{\binom{b}{d}} = W_{[\binom{b}{r_b},\binom{b}{r_b+d}]}$  when  $b \neq a$ ,  $\binom{b}{r_b+d} \in I$ . Mutate  $\dot{\mathbf{t}}\{r\}_s$  at  $\binom{a}{s}$ . Then Lemma 8.1 implies

(8.4) 
$$x'_{\binom{a}{s}} * x''_{\binom{a}{s}} = q^{\alpha} [x''_{\binom{a}{s-1}} * x'_{\binom{a}{s+1}}]^{\mathbf{\dot{t}}'} + q^{\alpha'} [\prod (x'_{\binom{b}{d}})^{-C_{ba}}]^{\mathbf{\dot{t}}'},$$

such that  $\binom{b}{d}$  appearing satisfy  $\binom{b}{d+r_b} < \frac{\alpha}{n} \binom{a}{(s+r_a+1)} < \frac{\alpha}{n} \binom{b}{(d+r_b+1)}$ , and  $\alpha, \alpha'$  satisfy  $\alpha > \alpha', \alpha = \Lambda'(\deg' x'_{\binom{a}{s}}), \deg' x''_{\binom{a}{s}}), \alpha' = \Lambda'(\deg x'_{\binom{a}{s}}), -\sum C_{ba} \deg' x'_{\binom{b}{d}}).$ Since  $\binom{a}{s+r_a} \in I$ , we have  $\binom{b}{d+r_b} \in I$ . So the cluster variables in (8.4) are the

Since  $\binom{a}{s+r_a} \in I$ , we have  $\binom{b}{d+r_b} \in I$ . So the cluster variables in (8.4) are the interval variables in (8.2). Note that normalization and the coefficients  $\alpha, \alpha'$  could be equally computed in the seed  $\dot{\mathbf{t}}$ , since the cluster variables involved belong to the same cluster ([Qin17]). Then we obtain (8.2) from (8.4).

Let  $\sigma$  denote the permutation on  $I_{\text{uf}}$  such that  $\sigma k^{\min}[r] = k^{\max}[-1-r]$ . Note that we have  $x_{\sigma k}(\Sigma \dot{\mathbf{t}}) = W_{k[1],k^{\max}}(\dot{\mathbf{t}})$  for  $k \in I_{\text{uf}}$  by construction. Lemma 8.4 implies the following result.

**Proposition 8.7.** We have deg<sup>t</sup>  $x_{\sigma k}(\Sigma \dot{\mathbf{t}}) = f_{k^{\max}} - f_k$  for  $k \in I_{uf}$ .

Particularly,  $\Sigma$  is a green to red sequence. We will denote  $\dot{\mathbf{t}}[1] := \Sigma \dot{\mathbf{t}} = \dot{\mathbf{t}}(-\underline{\eta}^{\mathsf{op}})$ . Note that  $x_{j^{\max}}(\dot{\mathbf{t}}[1]) = W_{j^{\min},j^{\max}}$  for any frozen  $j^{\max}$ .

**Lemma 8.8.**  $\forall k \in I_{uf}$ , we have

$$\deg y_k(\dot{\mathbf{t}}) = -\beta_k - \beta_{k[1]} - \sum_{j,j[d]} C_{\eta_j \eta_k} \beta_{[j,j[d]]}$$

where j, j[d] appearing satisfy  $j[-1] < k < j \le j[d] < k[1] < j[d+1]$ .

*Proof.* We have  $\deg y_k(\dot{\mathbf{t}}) = f_{k[-1]} - f_{k[1]} + \sum_{j:\eta_j \neq \eta_k} b_{jk}f_j$ , and, by Lemma 8.4,  $f_{k[-1]} - f_{k[1]} = -\beta_k - \beta_{k[1]}$ . By (6.1), the other entries  $b_{jk}$  are given by  $b_{j_1k} = C_{a,\eta_k}$  for  $j_1 < k < j_1[1] < k[1]$  or  $b_{j_2k} = -C_{a,\eta_k}$  for  $k < j_2 < k[1] < j_2[1]$ , where  $a = \eta_{j_1} = \eta_{j_2} \neq \eta_k$ . If  $j_1, j_2$  both exist, we have  $j_1 < k < j_1[1] < \cdots < j_2 < k[1] < j_2[1]$ .

Note that  $k[1] < +\infty$ . If  $j_2$  does not exist, then  $j_1$  does not exist. Otherwise, define j such that  $j[-1] < k < j \leq j_2$ , where we denote  $j[-1] = -\infty$  if  $j = j^{\min}$ . Denote  $j_2 = j[d]$ . Then  $j[-1] < k < j \leq j[d] < k[1] < j[d+1]$ . Moreover, j[-1] and  $j_2$  contribute to  $-C_{a,\eta_k}(f_{j[d]} - f_{j[-1]})$ , which equals  $-C_{a,\eta_k}\beta_{[j,j[d]]}$  by Lemma 8.4. The desired claim follows.

**Example 8.9.** Continue Example 8.2. Then  $x_i(\mathbf{t}) = W_{[i^{\min},i]}, \beta_i = \mathsf{deg}^{\mathbf{t}} W_{i,i}$ . We can verify Lemma 8.8. For example,  $\mathsf{deg} y_4(\mathbf{t}) = f_3 - f_7 + 2(f_6 - f_2) = -\beta_{[4,7]} + 2\beta_{[5,6]}$ . Applying the mutation sequence  $\Sigma = \boldsymbol{\mu}_{(1,3,4,2,5,1,3,1,2)}$  to  $\mathbf{t}$ , we obtain the seed

 $\mathbf{t}$ [1]. We obtain the following new cluster variables along  $\Sigma$  with distinct degrees:

1		$W_{[3,3]}$	$W_{[3,4]}$	$W_{[3,7]}$	$W_{[5,5]}$	$W_{[5,6]}$
	$deg^{\mathbf{t}}$	$-f_1 + f_3$	$-f_1 + f_4$	$-f_1 + f_7$	$-f_2 + f_5$	$-f_2 + f_6$
		$W_{[4,4]}$	$W_{[4,7]}$	$W_{[7,7]}$	$W_{[6,6]}$	
	$deg^{\mathbf{t}}$	$-f_3 + f_4$	$-f_3 + f_7$	$-f_4 + f_7$	$-f_5 + f_6$	

We have  $x_{\sigma k}(\dot{\mathbf{t}}[1]) = W_{[k[1],k^{\max}]}, \ \operatorname{\mathsf{deg}}^{\dot{\mathbf{t}}} x_{\sigma k}(\dot{\mathbf{t}}[1]) = -f_k + f_{k^{\max}}, \ \forall k \in I_{uf}, \ \text{where} \ \sigma = (1,4)(2,5).$ 

8.3. Standard bases. For any  $w = \sum w_i \beta_i \in \bigoplus_{i \in I} \mathbb{Z}\beta_i$ , define the *w*-pointed standard monomial  $\mathbf{M}(w) := [W_1^{w_1} * \cdots * W_l^{w_l}]^{\mathbf{t}}$ . Note that  $\{\beta_i | i \in I\}$  is a  $\mathbb{Z}$ -basis of  $M^{\circ}(\mathbf{t}) = \bigoplus_{i=1}^l \mathbb{Z}\beta_i$ . Let  $<_{\mathsf{lex}}$  denote lexicographical order and  $<_{\mathsf{rev}}$  the

reverse lexicographical order on  $\mathbb{Z}^{[1,l]} \simeq \bigoplus_{i=1}^{l} \mathbb{Z}\beta_i$ . Working at the quantum level  $\mathbb{k} = \mathbb{Z}[q^{\pm \frac{1}{2}}]$ , we will prove the follow result.

**Theorem 8.10.** (1) The set  $\mathbf{M} := {\mathbf{M}(w) | w \in \mathbb{N}^{[1,l]}}$  is a k-basis of  $\overline{\mathcal{U}}(\dot{\mathbf{t}}(\underline{\eta}))$ . (2) It satisfies the analog of the Levendorskii-Soibelman straightening law:

(8.5) 
$$W_k W_j - q^{\lambda(\deg W_k, \deg W_j)} W_j W_k \in \sum_{w \in \mathbb{N}^{[j+1,k-1]}} \Bbbk W(w), \ \forall j \le k.$$

(3) We have  $\overline{\mathcal{A}}(\dot{\mathbf{t}}(\eta)) = \overline{\mathcal{U}}(\dot{\mathbf{t}}(\eta)).$ 

(4) The common triangular basis element  $\mathbf{L}_{\sum w_i\beta_i}$  for  $\overline{\mathcal{U}}(\mathbf{t}(\underline{\eta}))$  equals the Kazhdan-Lusztig basis element  $\mathbf{L}(w)$  associated with  $\mathbf{M}$  sorted by  $\langle_{\mathsf{rev}}$ , i.e.,  $\mathbf{L}(w)$  is barinvariant and satisfies

$$\mathbf{L}(w) = M(w) + \sum_{w' <_{\mathsf{rev}} w} b_{w'} M(w'), \ b_{w'} \in q^{-\frac{1}{2}} \mathbb{Z}[q^{\frac{1}{2}}].$$

This statement still holds if when we replace  $<_{rev}$  by  $<_{lex}$ .

The k-basis **M** will be called the standard basis. Since  $\dot{\mathbf{t}}(\underline{\eta})$  can be optimized by Lemma 6.15, the set of dominant degrees  $\overline{M}^{\circ}(\dot{\mathbf{t}}(\eta))$  is given by Lemma 2.17.

**Lemma 8.11.** The set of dominant degrees  $\overline{M}^{\circ}(\dot{\mathbf{t}}(\underline{\eta}))$  (Definition 2.16) coincides with  $\bigoplus_{i \in [1,l]} \mathbb{N}\beta_i$ .

*Proof.* We give an indirect proof based on freezing operators. Note that  $\beta_i$  are degrees of  $W_{i,i} \in \overline{\mathcal{U}}$ . It remains to check the claim  $\overline{M}^{\circ}(\dot{\mathbf{t}}(\underline{\eta})) \subset \bigoplus_{i \in [1,l]} \mathbb{N}\beta_i$ .

(i) The claim is true when  $\underline{\eta} = (a, a, \dots, a)$ . In this case,  $\overline{\mathcal{U}}(\mathbf{\dot{t}})$  is isomorphic to the deformed Grothendieck ring of the level l-1 monoidal category  $\mathcal{C}_{l-1}$  consisting of the  $U_q(\widehat{\mathfrak{sl}}_2)$ -modules, whose properties are well-known, see [HL10][Qin17].

(ii) Take any  $g \in \overline{M}^{\circ}(\mathbf{t})$ . Let  $\mathbf{L}_g(\mathbf{t}) \in \overline{\mathcal{U}}(\mathbf{t})$  denote the common triangular basis element pointed at the tropical point [g].<sup>4</sup> Denote  $g = \sum c_i \beta_i$  for  $i \in \mathbb{Z}$ . We need to show that, for any  $k = k^{\min}$ , we have  $c_{k'} \geq 0$  whenever  $\eta_{k'} = \eta_k$ .

to show that, for any  $k = k^{\min}$ , we have  $c_{k'} \ge 0$  whenever  $\eta_{k'} = \eta_k$ . Denote  $I' := \{k, k[1], \ldots, k^{\max}\}$ . Then  $I \setminus I' = \{i | \eta_i \neq \eta_k\}$ . Choose  $F := (I \setminus I') \cap I_{uf}$ . We have  $g' = \operatorname{pr}_{I'} g = \sum_{I \setminus I'} c_i \beta_i$  and  $p = \operatorname{pr}_{\bigoplus_{i \in I \setminus I'} \mathbb{Z}f_i} g = \sum_{i \in I \setminus I'} c_i \beta_i$ . Moreover, we have  $\mathfrak{f}_F \mathbf{L}_g(\dot{\mathbf{t}}) = \mathbf{L}_{g'}(\mathfrak{f}_F \dot{\mathbf{t}}) \cdot x^p$ , see Theorems 3.17 and 3.18. Take  $\mathbb{k} = \mathbb{Z}$  from now on for simplifying arguments.

We obtain the seed  $\mathbf{\dot{t}}' := \mathbf{\dot{t}}((\eta_k, \eta_{k[1]}, \dots, \eta_{k^{\max}}))$  by deleting the frozen vertices  $I \cap I'$  from  $\mathfrak{f}_F \mathbf{\dot{t}}$ . The g'-pointed triangular basis element  $\mathbf{L}_{g'}(\mathbf{\dot{t}}')$  of  $\mathcal{U}(\mathbf{\dot{t}}')$  is obtained from  $\mathbf{L}_{g'}(\mathfrak{f}_F \mathbf{\dot{t}})$  by evaluating  $x_i$  to 1 for  $\eta_i \neq \eta_k$ . Since  $\mathbf{L}_g \in \overline{\mathcal{U}}(\mathbf{\dot{t}})$ ,  $\mathbf{L}_{g'}(\mathfrak{f}_F \mathbf{\dot{t}})$  and  $\mathbf{L}_{g'}(\mathbf{\dot{t}}')$  are still regular at  $x_{k^{\max}} = 0$ . Then (i) implies that  $g' \in \bigoplus_{r \in [0, o_+(1)]} \mathbb{N}\beta_{k[r]}$ .

By Lemma 8.8,  $\prec_{\mathbf{t}}$  implies  $<_{\mathsf{rev}}$  and  $<_{\mathsf{lex}}$ . Then  $\prec_{\mathbf{t}}$  is bounded on  $\overline{M}^{\circ}(\mathbf{t}(\underline{\eta}))$  by Lemma 8.11.

Lemma 8.12. M is a k-basis of  $\overline{\mathcal{U}}(\dot{\mathbf{t}})$ .

*Proof.* By Lemma 6.15 and Proposition 2.15,  $\{\mathbf{L}_m | m \in \overline{M}^{\circ}(\mathbf{\dot{t}})\}$  is a  $\overline{M}^{\circ}(\mathbf{\dot{t}})$ -pointed basis for  $\overline{\mathcal{U}}(\mathbf{\dot{t}})$ . Then the desired statement follows from the boundedness of  $\prec_{\mathbf{\dot{t}}}$ , see Lemmas 8.11 and 2.7.

 $<sup>^{4}</sup>$ Alternatively, we can work with the theta function instead of a triangular basis element. By [Qin24a], the freezing operator sends it to a theta function.

For any  $i \leq j \in [1, l]$ , we define  $\overline{\mathcal{A}}_{[i,j]}$  to be the k-subalgebra of  $\overline{\mathcal{A}}(t(\underline{\eta}))$  generated by  $W_i, W_{i+1}, \ldots, W_j$ . We set  $\overline{\mathcal{A}}_{[i,j]} = \mathbb{k}$  if i > j.

**Lemma 8.13.** We have 
$$W_i * W_j - q^{\lambda(\deg W_j, \deg W_i)} W_j * W_i \in \overline{\mathcal{A}}_{[i+1,j-1]}, \forall i \leq j$$
.

*Proof.* We give a proof by induction on l. The case l = 1 is trivial. Let us assume the desired statement has been verified for lengths  $\leq l - 1$ .

Use the natural inclusion  $\overline{\mathcal{A}}(\dot{\mathbf{t}}(\underline{\eta}_{[1,l-1]})) \subset \overline{\mathcal{A}}(\dot{\mathbf{t}}(\underline{\eta}))$ . Then,  $\forall i \in [1, l-1]$ , the fundamental variables  $W_i$  of  $\overline{\mathcal{A}}(\dot{\mathbf{t}}(\underline{\eta}_{[1,l-1]}))$  are identified with the fundamental variables  $W_i$  of  $\overline{\mathcal{A}}(\dot{\mathbf{t}}(\underline{\eta}))$ . Therefore, the induction hypothesis implies the desired statement for  $i, j \in [1, l-1]$ .

Next, take the seed  $\dot{\mathbf{t}}\{1\} := \Sigma_1 \dot{\mathbf{t}} = \dot{\mathbf{t}}(\eta_2, \dots, \eta_l, -\eta_1)$ . Use the natural inclusion  $\overline{\mathcal{A}}(\dot{\mathbf{t}}(\underline{\eta}_{[2,l]})) \subset \overline{\mathcal{A}}(\dot{\mathbf{t}}\{1\})$ . It follows from the construction of fundamental variables that,  $\forall i \in [1, l-1]$ , the fundamental variables  $W'_i$  of  $\overline{\mathcal{A}}(\dot{\mathbf{t}}(\underline{\eta}_{[2,l]}))$  are identified with  $W_{i+1}$  of  $\overline{\mathcal{A}}(\dot{\mathbf{t}})$ . Therefore, the induction hypothesis implies the desired statement for  $i, j \in [2, l]$ .

It remains to show  $z := W_1 * W_l - q^{\lambda(\deg W_l, \deg W_1)} W_l * W_1$  belongs to  $\overline{\mathcal{A}}_{[2,l-1]}$ . By Lemma 8.12, we have a finite decomposition  $z = \sum_{w \in \mathbb{N}^{[1,l]}} c_w M(w), c_w \in \mathbb{k}$ . Since M(w) are pointed at distinct degrees, we have  $c_w \neq 0$  only if  $w \prec_{\mathbf{i}} \beta_1 + \beta_l$  (see [Qin17, Lemma 3.1.10(iii)]). By Lemma 8.8,  $w \prec_{\mathbf{i}} \beta_1 + \beta_l$  implies  $w <_{\mathsf{rev}} \beta_1 + \beta_l$ and  $w <_{\mathsf{lex}} \beta_1 + \beta_l$ . Moreover, any element of  $\bigoplus_{k \in I_{uf}} \mathbb{N} \deg^{\mathbf{i}} y_k$  takes the form  $-b_{k_1}\beta_{k_1} + \sum_{k_1 < j < k_2} b_j\beta_j - b_{k_2}\beta_{k_2}$  for  $b_{k_1}, b_{k_2} \in \mathbb{N}, b_j \in \mathbb{Z}$ . Denote  $w = \sum_i w_i\beta_i, w_i \in \mathbb{N}$ . Then  $w <_{\mathsf{rev}} \beta_1 + \beta_l$  implies  $w = \beta_l$  or  $w_l = 0$ . The

Denote  $w = \sum_{i} w_i \beta_i$ ,  $w_i \in \mathbb{N}$ . Then  $w <_{\mathsf{rev}} \beta_1 + \beta_l$  implies  $w = \beta_l$  or  $w_l = 0$ . The former case is impossible since  $w - (\beta_1 + \beta_l) = -\beta_1$  is not contained in  $\bigoplus_k \mathbb{N} \deg^{\mathbf{t}} y_k$ , i.e.  $w \not\prec_{\mathbf{t}} \beta_1 + \beta_l$ . Similarly,  $w <_{\mathsf{lex}} \beta_1 + \beta_l$  implies  $w = \beta_1$  or  $w_1 = 0$ . The former case is impossible since  $w - (\beta_1 + \beta_l) = -\beta_l$  is not contained in  $\bigoplus_k \mathbb{N} \deg^{\mathbf{t}} y_k(\mathbf{t})$ . We conclude  $w \in \mathbb{N}^{[2,l-1]}$  as desired.

Proof of Theorem 8.10. (1) and (2) were proved in Lemmas 8.12 and 8.13. Since  $\mathbf{M} \subset \overline{\mathcal{A}}(\dot{\mathbf{t}})$ , Lemma 8.12 implies (3). Finally, we claim that the Kazhdan-Lusztig basis associated to the standard basis  $\mathbf{M}$ , sorted by  $<_{\mathsf{rev}}$ , is the triangular basis with respect to  $\dot{\mathbf{t}}$ . This will imply (4).

Let us give a proof of this claim based on induction of l. The case l = 1 is trivial. Assume the claim has been proved for l - 1.

Use the natural inclusion  $\overline{\mathcal{A}}(\dot{\mathbf{t}}') \subset \overline{\mathcal{A}}(\dot{\mathbf{t}})$  where  $\dot{\mathbf{t}}' = \dot{\mathbf{t}}(\underline{\eta}_{[1,l-1]})$ . Let  $\mathbf{L}$  and  $\mathbf{L}'$ denote the (common) triangular basis for  $\overline{\mathcal{A}}(\dot{\mathbf{t}})$  and  $\overline{\mathcal{A}}(\dot{\mathbf{t}}')$  respectively. Note that  $\dot{\mathbf{t}}'$ can be obtained from  $\dot{\mathbf{t}}$  by freezing  $l^{\max}[-1]$  and deleting  $l^{\max}$ . Take the freezing operator  $\mathbf{f} := \mathbf{f}_{\{l^{\max}[-1]\}}$ . By Theorem 3.17, for any  $w \in \bigoplus_i \mathbb{N}\beta_i$ ,  $\mathbf{fL}(w)$  is the *w*pointed triangular basis element of  $\mathcal{U}(\mathbf{f}\dot{\mathbf{t}})$ . By the similarity between  $\dot{\mathbf{t}}'$  and  $\mathbf{f}\dot{\mathbf{t}}$ , we have  $\mathbf{fL}(w) = \mathbf{L}'(u) \cdot x_l^{w_l}$ , where  $\mathbf{L}'(u)$  is the *u*-pointed triangular basis for  $\mathcal{U}(\dot{\mathbf{t}}')$ ,  $u \in \bigoplus_{i=1}^{l-1} \mathbb{Z}\beta_i$ , and  $u + w_l \beta_{[l^{\min},l]} = w$ .

We aim to show that  $\forall w = \sum w_i \beta_i$  with  $w_i \in \mathbb{N}$ ,  $w_l = 0$ ,  $\mathbf{M}(w)$  is  $(\prec_{\mathbf{t}}, \mathbf{m})$ unitriangular to **L**. If so, an arbitrary standard monomial  $[\mathbf{M}(w) * W_l^{w_l}]$  must be  $(\prec_{\mathbf{t}}, \mathbf{m})$ -unitriangular to **L** as well, since  $W_l^{w_l} = x_{l^{\min}}(\mathbf{t}[1])$ , see [Qin17].

Denote the decomposition  $\mathbf{M}(w) = \mathbf{L}(w) + \sum_{w' \prec_{\mathbf{t}} w} c_{w'} \mathbf{L}(w')$  in  $\overline{\mathcal{U}}$ . Then w' appearing satisfy  $w' <_{\mathsf{rev}} w$ . It follows that  $w' \in \bigoplus_{i \in [1,l-1]} \mathbb{N}\beta_i$ . Applying the freezing operator  $\mathfrak{f} := \mathfrak{f}_w := \mathfrak{f}_{l^{\max}[-1],w}$ , we obtain  $\mathfrak{f}\mathbf{M}(w) = \mathfrak{f}\mathbf{L}(w) + \sum_{w' <_{\mathsf{rev}} w} c_{w'} \mathfrak{f}_w \mathbf{L}(w')$ .

Note that w' appearing satisfy  $w' - w = \sum_{k \in I_{uf}} n_k \deg y_k$ ,  $n_k \in \mathbb{N}$ . In addition,  $\deg y_k$  are given by Lemma 8.8. Since  $w_l = 0$  and  $w'_l \ge 0$ , we must have  $n_{l^{\max}[-1]} = 0$  and  $w'_l = 0$ . Therefore,  $\mathbf{fL}(w) = \mathbf{L}'(w)$  and  $\mathbf{f}_w \mathbf{L}(w') = \mathbf{L}'(w')$  by the above discussion of freezing triangular basis elements. In addition, for  $i \in [1, l - 1]$ ,  $\mathbf{f}_{\beta_i} W_i$  is the  $\beta_i$ -pointed localized cluster monomial of  $\overline{\mathcal{A}}(\mathbf{t}')$ , which must be the  $\beta_i$  pointed fundamental variables  $W'_i$  for  $\overline{\mathcal{A}}(\mathbf{t}')$  by Lemma 8.4. So  $\mathbf{fM}(w)$  is a standard monomial of  $\overline{\mathcal{A}}(\mathbf{t}')$ . Then the induction hypothesis implies that  $c_{w'} \in \mathbf{m}$  as desired.

*Remark* 8.14. When the Cartan matrix is of finite type, Elek and Lu [EL21] showed that the coordinate rings of Bott-Samelson cells are symmetric Poisson CGL extensions. Their work can be lifted to the quantum level by [Mi18]. Then these are quantum cluster algebras by [GY16, GY18]. We don't know if the cluster algebras obtained in this way are the same as ours.

8.4. Categories associated to positive braids. Assume C is of finite type. We will prove that  $C(\underline{\eta})$  admits categorification in this case (Conjecture 1.8). Consider the following partial orders in analogous to the weak Bruhat orders.

**Definition 8.15.** For any positive braids  $\beta, \beta', \beta'' \in Br^+$ , denote  $\beta' \leq_R \beta, \beta'' \leq_L \beta$  if  $\beta = \beta' \beta''$ .

Let  $\vec{w}_0$  denote any reduced word for  $w_0$ . The corresponding positive braid  $\Delta := \beta_{\vec{w}_0} \in \mathsf{Br}^+$  is known to be independent of the choice of  $\vec{w}_0$ . There exists an involution  $\nu$  on J such that  $\sigma_a \Delta = \Delta \sigma_{\nu(a)}, \forall a$ . We also have  $\sigma_a \leq_R \Delta$ .

**Lemma 8.16.**  $\forall \beta \in Br^+$ , there exists  $d \in \mathbb{N}$  such that  $\beta \leq_R \Delta^d$ ,  $\beta \leq_L \Delta^d$ .

*Proof.* We only prove  $\beta \leq_R \Delta^d$  by induction on  $l(\beta)$ , and  $\beta \leq_L \Delta^d$  can be proved similarly. The case  $l(\beta) = 1$  is trivial. Decompose  $\beta = \sigma_a \beta'$  for some  $\beta' \in \mathsf{Br}^+$ . By the induction hypothesis, we have  $\Delta^{d'} = \beta' \zeta'$  for some  $d' \in \mathbb{N}$ . It follows that  $\beta \leq_R \sigma_a \Delta^{d'} = \Delta^{d'} \sigma_{\nu^{d'}(a)} \leq_R \Delta^{d'+1}$ .

Choose  $\beta := \beta_{\underline{\eta}}$  associated with  $\underline{\eta}$ . Denote  $l = l(\beta)$ . By Lemma 8.16, we can choose a word  $\underline{\zeta}$  such that  $\Delta^d = \beta_{\underline{\zeta}}, \ \underline{\eta} = \underline{\zeta}_{[1,l]}$ . Choose any Coxeter word  $\underline{c}$  and any height function  $\xi$  as in Section 7.6. Then we can find an adapted word  $\underline{\gamma}$  such that  $\Delta^d = \beta_{\underline{\gamma}}$ , see the knitting algorithm [Kel08]. Note that we have  $\dot{\mathbf{t}}(\underline{\zeta}) = \boldsymbol{\mu}^{\sigma} \dot{\mathbf{t}}(\underline{\gamma})$  where  $\boldsymbol{\mu}^{\sigma}$  is given in Section 6.3.

Recall that  $\overline{\mathcal{U}}(\dot{\mathbf{t}}(\underline{\gamma}))$  is categorified by a monoidal category  $\mathcal{C}_{\underline{\gamma}}(\xi)$ . We have  $\kappa$ :  $\overline{\mathcal{U}}(\dot{\mathbf{t}}(\gamma)) \simeq K$ , where K is the deformed Grothendieck ring. See Section 7.6.

Then we have  $\overline{\mathcal{A}}(\dot{\mathbf{t}}(\underline{\zeta})) = \overline{\mathcal{A}}(\dot{\mathbf{t}}(\underline{\gamma})) \stackrel{\kappa}{\simeq} K$ . Let  $S_i$  denote the simple module such that  $[S_i] = \kappa(x_i(\dot{\mathbf{t}}(\underline{\zeta})))$ , and  $L_{[j,k]}$  denote the simple module such that  $[L_{[j,k]}] = \kappa(W_{[j,k]}(\dot{\mathbf{t}}(\underline{\zeta})))$ , called the interval modules. Denote  $L_k := L_{[k,k]} = \kappa(W_k(\dot{\mathbf{t}}(\underline{\zeta})))$ .  $w \in \mathbb{N}^{[1,dl(\overline{\Delta})]}$ , define the generalized standard modules  $M(w) := L_1^{w_1} \otimes \cdots \otimes L_{dl(\overline{\Delta})}^{w_{dl}(\Delta)}$ . **Definition 8.17.** Let  $\text{Simp}_{\beta}$  denote the set of the composition factors of M(w),  $\forall w \in \mathbb{N}^{[1,l]}$ . Define  $\mathcal{C}_{\beta}(\xi)$  as the full subcategory of  $\mathcal{C}_{\underline{\gamma}}(\xi)$  whose objects have the composition factors in  $\text{Simp}_{\beta}$ .

Denote  $\mathbf{\dot{t}} = \mathbf{\dot{t}}(\underline{\zeta}), \ \overline{\mathcal{A}} = \overline{\mathcal{U}} = \overline{\mathcal{U}}(\mathbf{\dot{t}}), \ \mathbf{\dot{t}}' = \mathbf{\dot{t}}(\underline{\eta}), \ \overline{\mathcal{A}}' = \overline{\mathcal{U}}' = \overline{\mathcal{U}}(\mathbf{\dot{t}}').$  Let **M** and **L** (resp. **M**' and **L**') denote the standard basis and the common triangular basis for  $\overline{\mathcal{U}}$  (resp.  $\overline{\mathcal{U}}'$ ). Since  $\underline{\eta} = \underline{\zeta}_{[1,l]}, \mathbf{\dot{t}}$  is a good sub seed of  $\mathbf{\dot{t}}(\underline{\zeta})$  via the cluster embedding  $\iota := \iota_{\underline{\zeta},\underline{\eta}}, \$  which sends  $j \in [1,l]$  to j (Lemma 6.6). So we have the inclusion  $\overline{\mathcal{A}}' \subset \overline{\mathcal{A}}, \$  such that  $W_{[j,k]}(\mathbf{\dot{t}}')$  is identified with  $W_{[j,k]}(\mathbf{\dot{t}})$  by Lemma 8.5. It follows that  $\mathbf{M}' = \{\mathbf{M}(w)|w \in \bigoplus_{j \in [1,l]} \mathbb{N}\beta_j\}.$  Then  $\mathbf{L}'$  equals  $\{\mathbf{L}(w)|w \in \bigoplus_{j \in [1,l]} \mathbb{N}\beta_j\}$  since it is the Kazhdan-Lusztig basis associated to  $\mathbf{M}'$  (Theorem 8.10).

**Lemma 8.18.** (1) We have  $\text{Simp}_{\beta} = \{S \in \text{Simp} | [S] \in \kappa \mathbf{L}'\}$ . (2)  $\mathcal{C}_{\beta}(\xi)$  is a monoidal category.

Proof. (1) Any  $S \in \text{Simp}_{\beta}$  is a composition factor of some M(w),  $w \in \bigoplus_{j=1}^{l} \mathbb{N}\beta_{j}$ . Then  $\kappa^{-1}[S]$  is some common triangular basis element  $\mathbf{L}(w')$  that appears in the decomposition of  $\mathbf{M}(w)$  into  $\mathbf{L}$ . Since  $w \in \bigoplus_{i=1}^{l} \mathbb{N}\beta_{j}$ ,  $\mathbf{M}(w)$  lies in  $\text{Span}_{\Bbbk} \mathbf{L}'$ . Therefore,  $\mathbf{L}(w') \in \mathbf{L}'$ .

Conversely, any  $\mathbf{L}(w) \in \mathbf{L}'$  appears in the decomposition of M(w) into  $\mathbf{L}$ . So, when we denote  $\kappa \mathbf{L}(w) = [S]$ , S is a composition factor of  $M(w) \in \mathbf{M}'$ . Thus  $S \in \text{Simp}_{\beta}$ .

(2) Note that  $\operatorname{Span}_{\mathbb{k}} \mathbf{L}'$  is closed under multiplication since it equals  $\overline{\mathcal{U}}'$ . We deduce that  $\mathcal{C}_{\beta}$  is closed under the tensor product.

Lemma 8.18 has the following consequence.

**Theorem 8.19.** When C is of type ADE,  $(\overline{\mathcal{U}}(\dot{\mathbf{t}}(\eta)), \mathbf{L}')$  is categorified by  $\mathcal{C}_{\beta}$ .

Let us give an alternative description of  $\operatorname{Simp}_{\beta}$ .  $\forall$  modules  $V_1, V_2$ , we say  $V_2|V_1$ if there exists a module  $V_3$  such that  $V_2 \otimes V_3 \simeq V_1$ . Associated with  $\underline{\zeta}$ , we define  $F := \{j \in [1, l(\underline{\zeta})] | j[1] > l\}, F' = [l + 1, l(\underline{\zeta})]$ . Then  $\dot{\mathbf{t}}(\underline{\eta})$  can be obtained from  $f_F \dot{\mathbf{t}}(\boldsymbol{\zeta})$  by removing the non-essential frozen vertices  $j \in F'$ .

Proposition 8.20. We have the following description

 $\mathsf{Simp}_{\beta} = \{ S \in \mathsf{Simp} \, | S \otimes S_j \text{ is simple } \forall j \in F, \text{ and } S_j \nmid S \, \forall j \in F' \}.$ 

*Proof.* Denote the right hand side by Simp'. Then  $\kappa \mathbf{L}' \subset \{[S] | S \in \mathsf{Simp'}\}$ .

Conversely, if  $S(w) \in \text{Simp}'$ , then  $\mathbf{L}(w) := \kappa^{-1}[S(w)]$  q-commutes with  $x_j$ ,  $\forall j \in F$ , and  $\mathbf{L}(w) \neq x_j \cdot \mathbf{L}(w')$ ,  $\forall j \in F'$ . The first condition implies that  $\mathbf{L}(w)$  is a common triangular basis element for  $\mathcal{U}(\mathfrak{f}_F \dot{\mathbf{t}})$ , see Theorem 5.17. By the similarity between  $\mathfrak{f}_F \dot{\mathbf{t}}$  and  $\dot{\mathbf{t}}'$ , we have  $\mathbf{L}(w) = x^m \cdot \mathbf{L}(w')$  for some  $\mathbf{L}(w') \in \mathbf{L}'$  such that  $m \in \bigoplus_{j \in F'} \mathbb{Z}f_j$ . Moreover, since the vertices of F' are non essential in  $\mathfrak{f}_F \dot{\mathbf{t}}$ , the order of vanishing  $\nu_j(\mathbf{L}(w'))$  equals  $0, \forall j \in F'$  (Section 6.4). Then  $m_j = \nu_j(\mathbf{L}(w)) \ge 0$ ,  $\forall j \in F'$ , implies  $m \ge 0$ . By the second condition, we further have m = 0. So  $\mathbf{L}(w) = \mathbf{L}(w') \in \mathbf{L}'$  and thus  $S(w) \in \text{Simp}_{\beta}$ .

**Proposition 8.21.**  $C_{\beta}(\xi)$  does not depend on  $\underline{\zeta}$  where  $\beta_{\underline{\zeta}} = \Delta^d$ ,  $\beta_{\underline{\zeta}_{1,\eta}} = \beta$ .

The subsequent paper [Qin24b] will offer an alternative proof using braid groups.

*Proof.* Let  $\underline{\alpha}$  be another choice. Since  $\beta_{\underline{\alpha}_{[1,l]}} = \beta_{\underline{\zeta}_{[1,l]}} = \beta$  and  $\beta_{\underline{\alpha}} = \beta_{\underline{\zeta}} = \Delta^d$ , we have  $\beta_{\underline{\alpha}_{[l+1,l(\underline{\zeta})]}} = \beta_{\underline{\zeta}_{[l+1,l(\underline{\zeta})]}}$ . Denote  $\underline{\alpha}' := \underline{\alpha}_{[1,l]}, \underline{\alpha}'' := \underline{\alpha}_{[l+1,l(\underline{\zeta})]}, \underline{\zeta}' = \underline{\zeta}_{[1,l]} = \underline{\eta}, \underline{\zeta}'' = \underline{\zeta}_{[l+1,l(\underline{\zeta})]}$ . Using the permuted mutation sequences associated to braid moves and the cluster embeddings in Section 6.3, we obtain commutative diagrams:

$$\begin{array}{cccc} (\mu^{\sigma}_{\underline{\alpha}'',\underline{\zeta}''})^{*} & \\ \overline{\mathcal{U}}(\dot{\mathbf{t}}(\underline{\alpha}')) & \subset & \overline{\mathcal{U}}(\dot{\mathbf{t}}(\underline{\alpha}',\underline{\zeta}'')) & \stackrel{\sim}{\leftarrow} & \overline{\mathcal{U}}(\dot{\mathbf{t}}(\underline{\alpha}',\underline{\alpha}'')) \\ \downarrow^{\wr} (\mu^{\sigma}_{\underline{\alpha}',\underline{\zeta}'})^{*} & & \downarrow^{\wr} (\mu^{\sigma}_{\underline{\alpha}',\underline{\zeta}'})^{*} & & \downarrow^{\wr} (\mu^{\sigma}_{\underline{\alpha}',\underline{\zeta}'})^{*} \\ \overline{\mathcal{U}}(\dot{\mathbf{t}}(\underline{\zeta}')) & \subset & \overline{\mathcal{U}}(\dot{\mathbf{t}}(\underline{\zeta}',\underline{\zeta}'')) & = & \overline{\mathcal{U}}(\dot{\mathbf{t}}(\underline{\zeta}',\underline{\zeta}'')) \end{array}$$

Then the common triangular basis  $\mathbf{L}'$  for  $\overline{\mathcal{U}}(\mathbf{t}(\underline{\alpha}'))$  and  $\overline{\mathcal{U}}(\mathbf{t}(\underline{\zeta}'))$  are identified in  $\overline{\mathcal{U}}(\mathbf{t}(\underline{\zeta}',\underline{\zeta}''))$ . The claim follows from Lemma 8.18(1).

Finally, let  $\underline{c}'$  denote any Coxeter word. Denote  $\underline{\tilde{\eta}} = (\underline{c}', \underline{\eta})$  and  $\beta = \beta_{\underline{\tilde{\eta}}}$ . Note that  $\mathbf{\ddot{t}}(\underline{\eta}) = \mathfrak{f}_F \mathbf{\dot{t}}(\underline{\tilde{\eta}})$ , where  $F = \{ \begin{pmatrix} a \\ 0 \end{pmatrix}^{\underline{\tilde{\eta}}} | a \in J \}$ . By Theorem 8.19,  $\overline{\mathcal{U}}(\mathbf{\dot{t}}(\underline{\tilde{\eta}}))$  is categorified by  $\mathcal{C}_{\beta}$ . Choose simples  $S_k$  of  $\mathcal{C}_{\beta}$  such that  $[S_k] \sim x_k$  for  $k \in F$ . Let  $\mathcal{C}'_{\beta}$  denote the monoidal subcategory of  $\mathcal{C}_{\beta}$  with the simples objects  $\{S \in \mathcal{C}_{\beta} | S \otimes S_k \text{ is simple}, \forall k \in F \}$ , see Section 5.4. By Theorem 6.17,  $\overline{\mathcal{U}}(\mathbf{\ddot{t}}(\underline{\eta}))$  has the common triangular basis, denoted **L**. Combining Theorem 8.19, Proposition 6.16 and Corollary 5.19, we deduce the following result.

**Theorem 8.22.**  $(\overline{\mathcal{U}}(\mathbf{\check{t}}(\underline{\eta})), \mathbf{L})$  is categorified by  $\mathcal{C}'_{\widetilde{\beta}}$  when C is of type ADE.

Recall that, for any reduced word  $\underline{\eta}$  of  $w_0$ , we have  $\overline{\mathcal{U}}(\mathbf{\dot{t}}(\underline{\eta})) \otimes \mathbb{Q}(q^{\frac{1}{2}}) \simeq \mathcal{O}_q[G] \otimes \mathbb{Q}(q^{\frac{1}{2}})$  by Claim 7.6. Therefore, Theorem 8.22 applies to  $\mathcal{O}_q[G] \otimes \mathbb{Q}(q^{\frac{1}{2}})$ , where we extend the deformed Grothendieck ring K to  $K \otimes Q(q^{\frac{1}{2}})$ .

# Appendix A. Principal coefficients

Given any seed **t**. Let  $I_{\rm f}^{\rm prin} := \{k' | k \in I_{\rm uf}\}$  denote a copy of  $I_{\rm uf}$ . We recall the seed of principal coefficients  $t^{\rm prin}$  [FZ07]. Its set of vertices is  $I^{\rm prin} := I_{\rm uf} \sqcup I_{\rm f}^{\rm prin}$  with  $I_{\rm f}^{\rm prin}$  being frozen. The symmetrizers are  $d_k^{\rm prin} = d_{k'}^{\rm prin} = d_k$  for  $k \in I_{\rm uf}$ .

 $\text{Introduce the matrix entries } b_{ij}^{\text{prin}} := \begin{cases} b_{ij} & i, j \in I_{\text{uf}} \\ \delta_{kj} & j, k \in I_{\text{uf}}, i = k' \\ -\delta_{ik} & i, k \in I_{\text{uf}}, j = k' \\ 0 & i, j \in I_{\text{f}}^{\text{prin}} \end{cases} \text{. Then its $B$-matrix is } \end{cases}$ 

 $\begin{array}{ccc} \mathbf{l} 0 & i,j \in I_{\mathbf{f}}^{\mathrm{prin}} \\ \widetilde{B}^{\mathrm{prin}} \coloneqq (b_{ij}^{\mathrm{prin}})_{i \in I_{\mathrm{uf}}, j \in I^{\mathrm{prin}}}. \text{ Let var denote the linear map from } \mathbb{Z}^{I^{\mathrm{prin}}} = \oplus_{i \in I^{\mathrm{prin}}} \mathbb{Z}f_i \\ \mathrm{to } \ \mathbb{Z}^I = \oplus_{i \in I} \mathbb{Z}f_i \text{ such that } \mathsf{var}(f_i) = \begin{cases} f_i & i \in I_{\mathrm{uf}} \\ \sum_{j \in I_{\mathbf{f}}} b_{jk}f_j & i = k', k \in I_{\mathrm{uf}} \end{cases}. \\ \mathsf{var}(\sum_{j \in I^{\mathrm{prin}}} b_{jk}^{\mathrm{prin}}f_j) = \sum_{i \in I} b_{ik}f_i, \forall k \in I_{\mathrm{uf}}. \\ \mathrm{When } \mathbf{t} \text{ is a quantum seed endowed with a compatible skew-symmetric bilinear} \end{cases}$ 

When **t** is a quantum seed endowed with a compatible skew-symmetric bilinear form  $\lambda$ , we define the skew-symmetric bilinear form  $\lambda^{\text{prin}}$  on  $\mathbb{Z}^{I^{\text{prin}}}$  by  $\lambda^{\text{prin}}(m, m') = \lambda(\text{var } m, \text{var } m')$ . We deduce the following result.

**Lemma A.1.** The bilinear form  $\lambda(t^{\text{prin}})$  is compatible with  $\widetilde{B}(t^{\text{prin}})$ . Moreover, the quantum seed  $t^{\text{prin}}$  is similar to t with the scalar  $\rho = 1$ .

The linear map var induces a monomial k-algebra homomorphism var :  $\mathcal{LP}(\mathbf{t}^{\text{prin}}) \rightarrow \mathcal{LP}(\mathbf{t})$ , sending  $x^m(\mathbf{t}^{\text{prin}})$  to  $x^{\text{var}(m)}(\mathbf{t})$ . It sends an *m*-pointed element in  $\mathcal{LP}(\mathbf{t}^{\text{prin}})$  to a var(*m*)-pointed similar element in  $\mathcal{LP}(\mathbf{t})$ . It is a (quantum) variation map in the sense of [KQW23].

### APPENDIX B. FREEZING OPERATORS IN ADDITIVE CATEGORIFICATIONS

We refer the reader to [Qin12, Section 2.3] for necessary notions of additive categorification of cluster algebras following [Pla11].

Assume  $\mathbb{k} = \mathbb{Z}$  and B is skew-symmetric. We associate with the initial seed  $\mathbf{t}_0$  the generalized cluster category  $\mathcal{C}$ . For simplicity, we assume  $\mathcal{C}$  is Hom-finite to avoid discussing subcategories of  $\mathcal{C}$  as in [Pla11]. Let  $T = \bigoplus_{i \in I} T_i$  denote the cluster tilting object associated to  $\mathbf{t}_0$ . Let [1] denote the shift functor of  $\mathcal{C}$ . Note that  $\operatorname{Hom}(T, T[1]) = 0$ . Denote  $T^m = \oplus T_i^{m_i}$  for  $m = (m_i)_{i \in I} \in \mathbb{N}^I$ . The corresponding completed Jacobian algebra J equals  $\operatorname{End}_{\mathcal{C}}(T)^{\operatorname{op}}$ .

For any object V of C, let  $\underline{V}$  denote the (left) J-module  $\operatorname{Hom}_{\mathcal{C}}(T, V[1])$ . Further assume  $(\dim \underline{V})_j = 0$  for  $j \in I_f$ . There exists a triangle  $V \to T^{m_-}[1] \to T^{m_+}[1] \to V[1]$ . Define the index  $\operatorname{ind}_T V = m_+ - m_-$  (it equals the coindex  $-\operatorname{coind} V$  associated with T[-1] in [Pal08]). Then the cluster character of V is given by

$$\operatorname{CC}(V) = x^{\operatorname{ind}_T V} \cdot (\sum_{n \in \mathbb{N}^{I_{\operatorname{uf}}}} \chi(\operatorname{Gr}_n \underline{V}) \cdot y^n),$$

where  $\chi$  denote the Euler characteristics, and  $\operatorname{Gr}_{\underline{n}} \underline{V}$  the quiver Grassmannian consisting of the *n*-dimensional submodules of  $\underline{V}$ .

Choose any  $F \subset I_{uf}$  and denote  $T_F = \bigoplus_{k \in F} T_k$ . Choose any  $d \ge \dim \operatorname{Hom}(V, T_F[1])$ and let V'[1] be the cone of a generic map ev in  $\operatorname{Hom}(V, T_F^{\oplus d}[1])$ . We then have a triangle  $T_F^{\oplus d} \to V' \to V \xrightarrow{\operatorname{ev}} T_F^d[1]$  and view V' as the universal extension of V and  $T_F^{\oplus d}$ . Denote  $\underline{V}' = \operatorname{Hom}(T, V'[1])$ .

**Proposition B.1.**  $\underline{V}'$  is the maximal J-submodule of  $\underline{V}$  such that  $(\dim \underline{V}')_k = 0$  for  $k \in F$ .


*Proof.* (1) We first show that  $\underline{V}'$  is a submodule of  $\underline{V}$  satisfying  $(\dim \underline{V}')_k = 0$  for  $k \in F$ .

Applying  $\operatorname{Hom}(T, ()[1])$  to the triangle  $T_F^{\oplus d} \to V' \to V \xrightarrow{\operatorname{ev}} T_F^{\oplus d}[1]$ , we obtain an exact sequence  $0 \to \operatorname{Hom}(T, V'[1]) \to \operatorname{Hom}(T, V[1])$ . It follows that  $\underline{V}'$  is a submodule of  $\underline{V}$ .

Applying Hom(,  $T_F[1]$ ) to the triangle  $T_F^{\oplus d} \to V' \to V \xrightarrow{\text{ev}} T_F^{\oplus d}[1]$ , we obtain an exact sequence Hom $(T_F^{\oplus d}[1], T_F[1]) \xrightarrow{\text{ev}^*} \text{Hom}(V, T_F[1]) \to \text{Hom}(V', T_F[1]) \to 0$ . Since ev is generic and  $d \ge \dim \text{Hom}(V, T_F[1])$ , ev<sup>\*</sup> is surjective. It follows that  $\text{Hom}(V', T_F[1]) = 0$ . Since  $\mathcal{C}$  is 2-Calabi-Yau [Ami09], we obtain  $\text{Hom}(T_F, V'[1]) = 0$ .

(2) Let  $\underline{X}$  denote any submodule of  $\underline{V}$  satisfying  $(\dim \underline{X})_k = 0$  for  $k \in F$ . Then there exists a triangle  $Y \xrightarrow{-r[-1]} X \to V \xrightarrow{u} Y[1]$  such that  $\operatorname{Hom}(T, X[1]) = \underline{X}$ and -r[-1] factors through add T ([Pal08, Lemma 3.1 Proposition 1.1]). Applying  $\operatorname{Hom}(T_F^{\oplus d}[1])$  to it, we obtain  $\operatorname{Hom}(Y[1], T_F^{\oplus d}[1]) \xrightarrow{u^*} \operatorname{Hom}(V, T_F^{\oplus d}[1]) \to$ 

# FIGURE B.1. Octahedral axiom



 $\mathsf{Hom}(X, T_F^{\oplus d}[1]) = 0$ . Then  $ev \in \mathsf{Hom}(V, T_F^{\oplus d}[1])$  equals the composition  $v \circ u$  for some  $v \in \mathsf{Hom}(Y[1], T_F^{\oplus d}[1])$ .

Applying the octahedral axiom to the triangles  $V \stackrel{\text{ev}}{\to} T_F^{\oplus d}[1] \to V'[1] \to V[1]$ ,  $V \stackrel{u}{\to} Y[1] \stackrel{r}{\to} X[1] \to V[1]$ , and  $Y[1] \stackrel{v}{\to} T_F^{\oplus d}[1] \to Z \stackrel{s}{\to} Y[2]$ , we obtain a triangle  $X[1] \to V'[1] \to Z \stackrel{h}{\to} X[2]$  such that  $h = r[1] \circ s$ , see **B.1**(A). Since r[-1] factors through add T, r[1] factors through add T[2]. Thus Hom(T[1], h) vanishes. Applying Hom(T[1], ) to  $Z \stackrel{h}{\to} X[2] \to V'[2] \to Z[1]$ , we obtain that  $\underline{X}$  is a submodule of  $\underline{V'}$ .

**Lemma B.2.** We have ind  $V = \text{ind } V' - d \sum_{k \in F} f_k$ .

*Proof.* Recall that we have a triangle  $T_F^d \xrightarrow{u} V' \to V \to T_F^d[1]$ . Choose triangles  $V' \xrightarrow{v} T^{m'_-}[1] \to T^{m'_+}[1] \to V'[1]$  and  $T_F^d \xrightarrow{0} T^{m'_-}[1] \to T^{m'_-}[1] \oplus T_F^d[1] \to T_F^d[1]$ . Note that  $vu \in \operatorname{Hom}(T, T[1]) = 0$ . Using the octahedral axiom, we obtain a triangle  $V \to T^{m'_-}[1] \oplus T_F^d[1] \to T^{m'_+}[1] \to V[1]$ , see Figure B.1(B). The desired claim follows.

Proposition B.1 and Lemma B.2 imply the following.

**Theorem B.3.** The freezing operator  $\mathfrak{f}_F$  acts on  $\mathrm{CC}(V)$  by

(B.1) 
$$f_F CC(V) = x^{-d\sum_{k \in F} f_k} CC(V')$$

for  $d \in \mathbb{N}$  large enough. In this case,  $\underline{V}'$  is the maximal J-submodule of  $\underline{V}$  such that  $(\dim \underline{V}')_k = 0$  for  $k \in F$ .

Let  $\mathcal{C}'$  denote the subcategory of  $\mathcal{C}$  such that it consists of objects X satisfying  $\mathsf{Hom}(T_F, X[1]) = 0$ . Then  $V' \in \mathcal{C}'$  by Proposition B.1. This category provides additive categorification for  $\mathcal{U}(\mathfrak{f}_F \mathbf{t})$ .

#### References

- [Ami09] Claire Amiot, Cluster categories for algebras of global dimension 2 and quivers with potential, Annales de l'institut Fourier 59 (2009), no. 6, 2525–2590, arXiv:0805.1035.
- [BFZ05] Arkady Berenstein, Sergey Fomin, and Andrei Zelevinsky, Cluster algebras. III. Upper bounds and double Bruhat cells, Duke Math. J. 126 (2005), no. 1, 1–52.
- [BH24] Huanchen Bao and Xuhua He, Product structure and regularity theorem for totally nonnegative flag varieties, Inventiones mathematicae (2024), 1–47.
- [BZ05] Arkady Berenstein and Andrei Zelevinsky, Quantum cluster algebras, Adv. Math. 195 (2005), no. 2, 405–455, arXiv:math/0404446v2.
- [Cao22] Peigen Cao, On exchange matrices from string diagrams, arXiv:2203.07822.
- [CGG<sup>+</sup>22] Roger Casals, Eugene Gorsky, Mikhail Gorsky, Ian Le, Linhui Shen, and José Simental, Cluster structures on braid varieties, arXiv:2207.11607.
- [CILFS15] Giovanni Cerulli Irelli, Daniel Labardini-Fragoso, and Jan Schröer, Caldero-chapoton algebras, Transactions of the American Mathematical Society 367 (2015), no. 4, 2787– 2822, arXiv:1208.3310.
- [CKQ24] Peigen Cao, Bernhard Keller, and Fan Qin, 2024, in preparation.
- [CLS15] Ilke Canakci, Kyungyong Lee, and Ralf Schiffler, On cluster algebras from unpunctured surfaces with one marked point, Proceedings of the American Mathematical Society, Series B 2 (2015), no. 3, 35–49.
- [CW19] Sabin Cautis and Harold Williams, Cluster theory of the coherent Satake category, Journal of the American Mathematical Society 32 (2019), no. 3, 709–778, arXiv:1801.08111.
- [DWZ10] Harm Derksen, Jerzy Weyman, and Andrei Zelevinsky, Quivers with potentials and their representations II: Applications to cluster algebras, J. Amer. Math. Soc. 23 (2010), no. 3, 749–790, arXiv:0904.0676.
- [EL21] Balázs Elek and Jiang-Hua Lu, Bott-Samelson varieties and Poisson ore extensions, International Mathematics Research Notices 2021 (2021), no. 14, 10745–10797.
- [Fra16] Chris Fraser, Quasi-homomorphisms of cluster algebras, Advances in Applied Mathematics 81 (2016), 40–77, arXiv:1509.05385.
- [FWZ20] Sergey Fomin, Lauren Williams, and Andrei Zelevinsky, Introduction to Cluster Algebras. Chapter 6, arXiv:2008.09189.
- [FZ02] Sergey Fomin and Andrei Zelevinsky, Cluster algebras. I: Foundations, Journal of the American Mathematical Society 15 (2002), no. 2, 497–529, arXiv:math/0104151.
- [FZ07] \_\_\_\_\_, Cluster algebras IV: Coefficients, Compositio Mathematica 143 (2007), 112– 164, arXiv:math/0602259.
- [GHK15] Mark Gross, Paul Hacking, and Sean Keel, Birational geometry of cluster algebras, Algebraic Geometry 2 (2015), no. 2, 137–175.
- [GHKK18] Mark Gross, Paul Hacking, Sean Keel, and Maxim Kontsevich, Canonical bases for cluster algebras, Journal of the American Mathematical Society 31 (2018), no. 2, 497–608, arXiv:1411.1394.
- [GL19] Pavel Galashin and Thomas Lam, Positroid varieties and cluster algebras, arXiv:1906.03501.
- [GLS11] Christof Gei
  ß, Bernard Leclerc, and Jan Schröer, Kac-Moody groups and cluster algebras, Advances in Mathematics 228 (2011), no. 1, 329–433, arXiv:1001.3545.
- [GLS13] \_\_\_\_\_, Cluster structures on quantum coordinate rings, Selecta Mathematica 19 (2013), no. 2, 337–397, arXiv:1104.0531.
- [GLSBS22] Pavel Galashin, Thomas Lam, Melissa Sherman-Bennett, and David Speyer, Braid variety cluster structures, I: 3D plabic graphs, arXiv:2210.04778.
- [GSV03] Michael Gekhtman, Michael Shapiro, and Alek Vainshtein, Cluster algebras and poisson geometry, Mosc. Math. J 3 (2003), no. 3, 899–934, arXiv:arXiv:math/0208033.
- [GSV05] \_\_\_\_\_, Cluster algebras and Weil-Petersson forms, Duke Math. J. **127** (2005), no. 2, 291–311.
- [GSV23] Misha Gekhtman, Michael Shapiro, and Alek Vainshtein, A unified approach to exotic cluster structures on simple Lie groups, arXiv:2308.16701.
- [GY16] K.R. Goodearl and M.T. Yakimov, Quantum cluster algebra structures on quantum nilpotent algebras, Memoirs of the American Mathematical Society 247 (2016), no. 1169, arXiv:1309.7869.

- [GY18] KR Goodearl and MT Yakimov, Cluster algebra structures on poisson nilpotent algebras, arXiv:1801.01963.
- [GY20] K.R. Goodearl and M.T. Yakimov, The Berenstein-Zelevinsky quantum cluster algebra conjecture, Journal of the European Mathematical Society 22 (2020), no. 8, 2453-2509, arXiv:1602.00498.
- [HL10] David Hernandez and Bernard Leclerc, Cluster algebras and quantum affine algebras, Duke Math. J. 154 (2010), no. 2, 265–341, arXiv:0903.1452.
- [IOS23] Tsukasa Ishibashi, Hironori Oya, and Linhui Shen, A = U for cluster algebras from moduli spaces of G-local systems, Advances in Mathematics **431** (2023), 109256.
- [JLO23] Il-Seung Jang, Kyu-Hwan Lee, and Se-jin Oh, Braid group action on quantum virtual Grothendieck ring through constructing presentations, arXiv:2305.19471.
- [Kas91] M. Kashiwara, On crystal bases of the q-analogue of universal enveloping algebras, Duke Math. J. 63 (1991), no. 2, 465–516.
- [Kas93] \_\_\_\_\_, Global crystal bases of quantum groups, Duke Math. J. 69 (1993), no. 2, 455–485, Available from: http://dx.doi.org/10.1215/S0012-7094-93-06920-7, doi:10.1215/S0012-7094-93-06920-7. MR 1203234 (94b:17024)
- [Kel08] Bernhard Keller, Cluster algebras, quiver representations and triangulated categories, 2008, arXiv:0807.1960v11.
- [Kel11] \_\_\_\_\_, On cluster theory and quantum dilogarithm identities, Representations of Algebras and Related Topics, Editors A. Skowronski and K. Yamagata, EMS Series of Congress Reports, European Mathematical Society, 2011, pp. 85–11.
- [KKK18] Seok-Jin Kang, Masaki Kashiwara, and Myungho Kim, Symmetric quiver Hecke algebras and R-matrices of quantum affine algebras, Inventiones mathematicae 211 (2018), no. 2, 591–685.
- [KKK018] S.-J. Kang, M. Kashiwara, M. Kim, and S.-j. Oh, Monoidal categorification of cluster algebras, J. Amer. Math. Soc. 31 (2018), 349-426, arXiv:1412.8106, doi:https://doi.org/10.1090/jams/895.
- [KL09] Mikhail Khovanov and Aaron D. Lauda, A diagrammatic approach to categorification of quantum groups I, Represent. Theory 13 (2009), 309–347, arXiv:0803.4121.
- [KL10] \_\_\_\_\_, A diagrammatic approach to categorification of quantum groups III, Quantum Topology 1 (2010), no. 1, 1–92, arXiv:0807.3250.
- [KQ14] Yoshiyuki Kimura and Fan Qin, Graded quiver varieties, quantum cluster algebras and dual canonical basis, Advances in Mathematics 262 (2014), 261–312, arXiv:1205.2066.
- [KQW23] Yoshiyuki Kimura, Fan Qin, and Qiaoling Wei, Twist automorphisms and Poisson structures, SIGMA. Symmetry, Integrability and Geometry: Methods and Applications 19 (2023), 105, arXiv:2201.10284.
- [Lec03] B. Leclerc, Imaginary vectors in the dual canonical basis of  $U_q(\mathfrak{n})$ , Transform. Groups 8 (2003), no. 1, 95–104.
- [Lus90] G. Lusztig, Canonical bases arising from quantized enveloping algebras, J. Amer. Math. Soc. 3 (1990), no. 2, 447–498.
- [Lus91] \_\_\_\_\_, Quivers, perverse sheaves, and quantized enveloping algebras, J. Amer. Math. Soc. 4 (1991), no. 2, 365–421.
- [McN21] Peter J McNamara, Cluster monomials are dual canonical, arXiv:2112.04109.
- [Mi18] Yipeng Mi, Quantization of Poisson CGL extensions, arXiv:1808.09854.
- [Mul13] Greg Muller, Locally acyclic cluster algebras, Advances in Mathematics 233 (2013), no. 1, 207–247.
- [Mul16] \_\_\_\_\_, The existence of a maximal green sequence is not invariant under quiver mutation, Electron. J. Combin. 23 (2016), no. 2, arXiv:arXiv:1503.04675.
- [MW22] HB Moon and HELEN Wong, Compatibility of skein algebra and cluster algebra on surfaces and applications, arXiv:2201.08833.
- [Nak01] Hiraku Nakajima, Quiver varieties and finite-dimensional representations of quantum affine algebras, J. Amer. Math. Soc. 14 (2001), no. 1, 145–238 (electronic), arXiv:math/9912158.
- [Nak04] \_\_\_\_\_, Quiver varieties and t-analogs of q-characters of quantum affine algebras, Ann. of Math. (2) 160 (2004), no. 3, 1057–1097, arXiv:math/0105173v2.
- [Pal08] Yann Palu, Cluster characters for 2-Calabi-Yau triangulated categories, Ann. Inst. Fourier (Grenoble) 58 (2008), no. 6, 2221–2248, arXiv:math/0703540v2.

- [Pla11] Pierre-Guy Plamondon, Cluster algebras via cluster categories with infinite-dimensional morphism spaces, Compos. Math. 147 (2011), no. 6, 1921–1934, arXiv:1004.0830v1.
- [Qin12] Fan Qin, Quantum cluster variables via Serre polynomials, J. Reine Angew. Math. 2012 (2012), no. 668, 149–190, with an appendix by Bernhard Keller, arXiv:1004.4171, doi:10.1515/CRELLE.2011.129.
- [Qin14] \_\_\_\_\_, t-analog of q-characters, bases of quantum cluster algebras, and a correction technique, International Mathematics Research Notices 2014 (2014), no. 22, 6175– 6232, arXiv:1207.6604, doi:10.1093/imrn/rnt115.
- [Qin17] \_\_\_\_\_, Triangular bases in quantum cluster algebras and monoidal categorification conjectures, Duke Mathematical Journal 166 (2017), no. 12, 2337-2442, arXiv:1501.04085.
- [Qin20a] \_\_\_\_\_, An analog of Leclerc's conjecture for bases of quantum cluster algebras, SIGMA. Symmetry, Integrability and Geometry: Methods and Applications 16 (2020), 122, arXiv:2004.12466.
- [Qin20b] \_\_\_\_\_, Dual canonical bases and quantum cluster algebras, arXiv:2003.13674.
- [Qin21] \_\_\_\_\_, Cluster algebras and their bases, proceedings of ICRA2020, to appear (2021), arXiv:2108.09279.
- [Qin24a] \_\_\_\_\_, Applications of the freezing operators on cluster algebras, to appear in the special issue "Cluster Algebras and Related Topics" of Journal of Algebra and its Applications (2024).
- [Qin24b] \_\_\_\_\_, Based cluster algebras of infinite ranks, arXiv:2409.02881.
- [Qin24c] \_\_\_\_\_, Bases for upper cluster algebras and tropical points, Journal of the European Mathematical Society **26** (2024), no. 4, 1255–1312, arXiv:1902.09507.
- [Rou08] Raphaël Rouquier, 2-Kac-Moody algebras, 2008, arXiv:0812.5023.
- [Sco06] Joshua S Scott, Grassmannians and cluster algebras, Proceedings of the London Mathematical Society 92 (2006), no. 02, 345–380, arXiv:arXiv:math/0311148.
- [SSBW19] Khrystyna Serhiyenko, Melissa Sherman-Bennett, and Lauren Williams, Cluster structures in Schubert varieties in the Grassmannian, Proceedings of the London Mathematical Society 119 (2019), no. 6, 1694–1744.
- [SW21] Linhui Shen and Daping Weng, Cluster structures on double Bott-Samelson cells, Forum of Mathematics, Sigma, vol. 9, Cambridge University Press, 2021, arXiv:1904.07992.
- [Tra11] Thao Tran, F-polynomials in quantum cluster algebras, Algebr. Represent. Theory 14 (2011), no. 6, 1025–1061, arXiv:0904.3291v1.
- [VV03] M. Varagnolo and E. Vasserot, Perverse sheaves and quantum Grothendieck rings, Studies in memory of Issai Schur (Chevaleret/Rehovot, 2000), Progr. Math., vol. 210, Birkhäuser Boston, Boston, MA, 2003, pp. 345-365, arXiv:math/0103182v3. MR MR1985732 (2004d:17023)

Email address: qin.fan.math@gmail.com