Graphon Particle Systems, Part II: Dynamics of Distributed Stochastic Continuum Optimization

Yan Chen and Tao Li, Senior Member, IEEE

Abstract

We study the distributed optimization problem over a graphon with a continuum of nodes, which is regarded as the limit of the distributed networked optimization as the number of nodes goes to infinity. Each node has a private local cost function. The global cost function, which all nodes cooperatively minimize, is the integral of the local cost functions on the node set. We propose stochastic gradient descent and gradient tracking algorithms over the graphon. We establish a general lemma for the upper bound estimation related to a class of time-varying differential inequalities with negative linear terms, based upon which, we prove that for both kinds of algorithms, the second moments of the nodes' states are uniformly bounded. Especially, for the stochastic gradient tracking algorithm, we transform the convergence analysis into the asymptotic property of coupled nonlinear differential inequalities with time-varying coefficients and develop a decoupling method. For both kinds of algorithms, we show that by choosing the time-varying algorithm gains properly, all nodes' states achieve \mathscr{L}^{∞} -consensus for a connected graphon. Furthermore, if the local cost functions are strongly convex, then all nodes' states converge to the minimizer of the global cost function and the auxiliary states in the stochastic gradient tracking algorithm converge to the gradient value of the global cost function at the minimizer uniformly in mean square.

Index Terms

This work was funded by the National Natural Science Foundation of China under Grant No. 62261136550. This paper was presented in part at the 62nd IEEE Conference on Decision and Control, Marina Bay Sands, Singapore, Dec. 13-15, 2023. Corresponding author: Tao Li.

Yan Chen is with the School of Mathematical Sciences, East China Normal University, Shanghai 200241, China (e-mail: YanChen@stu.ecnu.edu.cn).

Tao Li is with the Shanghai Key Laboratory of Pure Mathematics and Mathematical Practice, School of Mathematical Sciences, East China Normal University, Shanghai 200241, China, and he is also with Institute of Mathematical Sciences at New York University Shanghai, Shanghai, 200062, China (email: tli@math.ecnu.edu.cn). Graphon mean field theory, graphon particle system, stochastic gradient descent algorithm, stochastic gradient tracking algorithm.

I. INTRODUCTION

In a distributed optimization problem over a network, all nodes cooperatively optimize a global cost function which is the sum of local cost functions, and each node only knows its own local cost function. Distributed optimization involving information exchange among nodes over a large-scale network can be found applications in distributed machine learning ([1]), multiagent target tracking ([2]), distributed resource allocation ([3]), and so on. The dimensions of these algorithms explode as the number of nodes increases, and it is of interest to investigate the limiting case as the number of nodes tends to infinity. In fact, games and optimal control problems with a continuum of individuals have been studied intensively in the field called mean field games, which was pioneered independently by Huang, Caines and Malhamé ([4]) and Lasry and Lions ([5]), respectively. They attempt to understand the behaviors of the limiting systems of the dynamic games with a large number of individuals. In the past decades, there has been an increasing intention in mean field games and their applications ([6]-[13]). Motivated by the distributed optimization over large-scale networks and the developing theory of mean-field control and games, we investigate the limiting model of the distributed optimization problem as the number of nodes tends to infinity, that is, the distributed optimization problem over a graphon with a continuum of nodes.

Let [0,1] be the set of a continuum of nodes, each element of which corresponds to a node. The connecting structure among nodes is given by the graphon A, which is a symmetric measurable function from $[0,1] \times [0,1]$ to [0,1] ([14]). Any node $p \in [0,1]$ has a private local cost function $V(p,x) : [0,1] \times \mathbb{R}^n \to \mathbb{R}$, which is strongly convex and continuously differentiable w.r.t. $x \in \mathbb{R}^n$ and is integrable w.r.t. $p \in [0,1]$. The objective of all nodes is to cooperatively solve the optimization problem

$$\min_{x \in \mathbb{R}^n} V(x) \triangleq \int_{[0,1]} V(p,x) dp,$$
(1)

for which there is a unique minimizer of V(x) denoted by x^* .

In the distributed optimization over the network with finite nodes, all nodes interact through the underlying network. The interactions among nodes depend on their labels and so are heterogeneous. In the graphon mean field theory, the concept of graph limit is introduced into the

3

mean field theory, which provides a powerful tool for modeling the heterogeneous interactions among a large number of individuals ([15]-[24]). Representing the heterogeneous interactions among nodes in terms of the coupled mean field terms based on the graphon, we propose the following distributed stochastic gradient descent algorithm for the problem (1). For any node $p \in [0, 1]$,

$$dx_p(t) = \alpha_1(t) \int_{\mathbb{R}^n \times [0,1]} A(p,q)(x-x_p(t)) \mu_t(dx,dq) dt - \alpha_2(t) \nabla_x V(p,x_p(t)) dt$$
$$- \alpha_2(t) \Sigma_1 dw_p(t), \tag{2}$$

where $x_p(t) \in \mathbb{R}^n$ is the state of node p at time t, representing its local estimate of x^* ; $\nabla_x V(p, x_p(t)) \in \mathbb{R}^n$ is the gradient value of the local cost function at $x_p(t)$; $\int_{\mathbb{R}^n \times [0,1]} A(p,q)(x-x_p(t)) \mu_t(dx,dq)$ is the coupled mean field term based on the graphon A. Let $(\Omega, \mathscr{F}, \mathscr{P})$ be a complete probability space with a family of non-decreasing σ -algebras $\{\mathscr{F}_t, t \ge 0\} \subseteq \mathscr{F}$. For any $t \ge 0$, $\mu_t(dx,dq)$ is the distribution on $\mathbb{R}^n \times [0,1]$ and satisfies the following conditions. (i) The marginal distribution $\mu_t(dq)$ is always the uniform distribution on [0,1], that is, $\mu_t(dq) = dq$, $\forall t \ge 0$. (ii) Given $q \in [0,1]$, the conditional distribution $\mu_t(dx|q)$ is the distribution of $x_q(t)$. Here, $\{(w_p(t),\mathscr{F}_t), t\ge 0, p \in [0,1]\}$ is a family of independent *n*-dimensional standard Brownian motions and the initial states $\{x_p(0), p \in [0,1]\}$ are adapted to \mathscr{F}_0 and independent of $\{w_p(t), t \ge 0, p \in [0,1]\}$. The terms $\alpha_1(t)$ and $\alpha_2(t)$ are time-varying algorithm gains and $\Sigma_1 \in \mathbb{R}^{n \times n}$. We also propose the following stochastic gradient tracking algorithm

$$dz_{p}(t) = \beta_{3}(t) \int_{[0,1]\times\mathbb{R}^{n}} A(p,q)(z-z_{p}(t)) \mu_{t,q}(dz) dq dt - \beta_{1}(t) y_{p}(t) dt -\beta_{1}(t) \beta_{3}(t) \int_{[0,1]\times\mathbb{R}^{n}} A(p,q)(y-y_{p}(t)) v_{t,q}(dy) dq dt, dy_{p}(t) = \beta_{3}(t) \int_{[0,1]\times\mathbb{R}^{n}} A(p,q)(y-y_{p}(t)) v_{t,q}(dy) dq dt + \beta_{2}(t) H(V(p,z_{p}(t))) dz_{p}(t) +\beta_{2}(t) \eta_{p}(t) dt + \beta_{2}'(t) \nabla_{x} V(p,z_{p}(t)) dt,$$

$$(3)$$

 $\forall p \in [0,1]$, where $z_p(t) \in \mathbb{R}^n$ is the state of node p at time t, representing its local estimate of x^* ; $y_p(t) \in \mathbb{R}^n$ is the auxiliary state of node p at time t, tracking the average $\nabla_x \left(\int_{[0,1]} V(p, z_p(t)) dp \right)$ and satisfying that $E[y_p(0)] = E[V(p, z_p(0))]; \nabla_x V(p, z_p(t)) \in \mathbb{R}^n$ is the gradient value of the local cost function at $z_p(t); H(V(p, z_p(t)))$ is the Hessian matrix of the local cost function at $z_p(t);$ $\mu_{t,q}(dz)$ and $v_{t,q}(dy)$ are the distributions of $z_q(t)$ and $y_q(t); \int_{[0,1] \times \mathbb{R}^n} A(p,q)(z-z_p(t))\mu_{t,q}(dz)dq$ and $\int_{[0,1] \times \mathbb{R}^n} A(p,q)(y-y_p(t))v_{t,q}(dy)dq$ are the coupled mean field terms of the states and the auxiliary states based on the graphon A. Here, $\{(\eta_p(t), \mathscr{F}_t), t \ge 0, p \in [0,1]\}$ is a family of independent n-dimensional stochastic processes, the processes $\{w_p(t), t \ge 0, p \in [0,1]\}$ and $\{\eta_p(t), t \ge 0, p \in [0,1]\}$ are mutually independent, and the initial states $\{z_p(0), p \in [0,1]\}$ and auxiliary states $\{y_p(0), p \in [0,1]\}$ are adapted to \mathscr{F}_0 and independent of $\{\eta_p(t), t \ge 0, p \in [0,1]\}$ and $\{w_p(t), t \ge 0, p \in [0,1]\}$. The terms $\beta_1(t)$, $\beta_2(t)$ and $\beta_3(t)$ are time-varying algorithm gains and $\beta'_2(t)$ is the derivative of $\beta_2(t)$ w.r.t. t.

Both the systems (2) and (3) belong to the following graphon particle system

$$dz_{p}(t) = \left[c_{1}(t)\int_{[0,1]\times\mathbb{R}^{m}}A(p,q)(z-z_{p}(t))\mu_{t,q}(dz)dq + c_{2}(t)\int_{[0,1]\times\mathbb{R}^{m}}A(p,q)(f(q,z,t) - f(p,z_{p}(t),t))\mu_{t,q}(dz)dq + c_{3}(t)g(p,z_{p}(t),t) + c_{4}(t)\xi_{p}(t)\right]dt + c_{5}(t)\Sigma dw_{p}(t), \quad (4)$$

 $\forall p \in [0,1]$, where $f(p,z,t) : [0,1] \times \mathbb{R}^m \times [0,\infty) \to \mathbb{R}^m$ and $g(p,z,t) : [0,1] \times \mathbb{R}^m \times [0,\infty) \to \mathbb{R}^m$ are the functions satisfying appropriate conditions; $\{(\xi_p(t),\mathscr{F}_t), t \ge 0, p \in [0,1]\}$ is a family of independent *m*-dimensional stochastic processes; the processes $\{w_p(t), t \ge 0, p \in [0,1]\}$ and $\{\xi_p(t), t \ge 0, p \in [0,1]\}$ are mutually independent; the initial states $\{z_p(0), p \in [0,1]\}$ are adapted to \mathscr{F}_0 and independent of $\{\xi_p(t), t \ge 0, p \in [0,1]\}$ and $\{w_p(t), t \ge 0, p \in [0,1]\}$; $c_i(t), i =$ $1, \ldots, 5$ are the time-varying coefficients, $\Sigma \in \mathbb{R}^{m \times m}$ and *m* is a positive integer.

Up to now, most of existing works ([15]-[17]) focused on the existence and uniqueness of the solutions for different graphon particle systems and the convergence of finite particle systems to graphon particle systems. Only few works ([18]-[19]) are concerned with the asymptotic properties of the graphon particle systems. Bayraktar and Wu ([18]) showed that the distribution of each node's state converges to a limiting distribution as time goes to infinity. They also provided an exponential concentration bound for the Wasserstein distance between the empirical distribution and the integral of the limiting distributions on the node set in [19]. Note that all aforementioned works on the graphon particle systems only prove the existence of limiting distributions but do not characterize what these limiting distributions specifically are, particularly, they do not reveal the relation between the limiting distributions and system dynamics. However, for many practical problems, people are more interested in how the limiting distribution is related to the system dynamics. In particular, for the problem (1) and the algorithms (2) and (3), people expect to figure out whether the states $\{x_p(t), t \ge 0, p \in [0, 1]\}$ and $\{z_p(t), t \ge 0, p \in [0, 1]\}$ converge to the minimizer of the global cost function under some proper assumptions.

Motivated by the above, we investigate the asymptotic properties of the graphon particle systems (2) and (3). We prove that if the graphon is connected and the local cost functions are strongly convex, then by properly choosing algorithm gains, both the states $\{x_p(t), t \ge 0, p \in$

[0,1] in (2) and $\{z_p(t), t \ge 0, p \in [0,1]\}$ in (3) converge to the minimizer of the global cost function in mean square. Different from Bayraktar and Wu ([18]), we weaken the assumptions on the local cost functions and obtain stronger results. Bayraktar and Wu ([18]) assumed that the dissipativity of the drift term is strictly twice greater than the Lipschitz constant of the interaction term. For the systems (2) and (3), this assumption is equivalent to the strong convexity constant of the local cost functions being greater than two, which is not reasonable for distributed optimization problems. In this paper, the local cost functions are only assumed to be strongly convex and there is no further requirement on the strong convexity constant. Bayraktar and Wu ([18]) proved the existence of the limiting distributions but also reveal that the limiting distribution is right the Dirac measure at the minimizer of the global cost function. Besides, Bayraktar and Wu ([18]) proved that all nodes' states converge in distribution, while we prove the convergence in mean square, which is stronger than convergence in distribution.

The introducing of time-varying algorithm gains removes the requirement on the strong convexity constant of the local cost functions in (2) and (3), which is introduced in [18] for time-invariant graphon particle systems. This leads to a time-varying general system (4) and it poses difficulties in establishing the relationship between the \mathscr{L}^2 -consensus $\lim_{t\to\infty} \int_{[0,1]} ||E[z_p(t)]| - \int_{[0,1]} E[z_q(t)] dq ||^2 dp = 0$ and \mathscr{L}^∞ -consensus $\lim_{t\to\infty} \sup_{p\in[0,1]} ||E[z_p(t)]] - \int_{[0,1]} E[z_q(t)] dq ||^2 = 0$. To this end, we give a key lemma to estimate the upper bounds of a class of functions satisfying time-varying differential inequalities with negative linear terms, based upon which, by coordinating the decaying rates of the time-varying coefficients and combining the connectivity of the graphon, we prove that the \mathscr{L}^2 -consensus implies \mathscr{L}^∞ -consensus for the system (4) if the integral of the second moments of all nodes' states on the node set is uniformly bounded.

For the stochastic gradient decent algorithm (2), by choosing the algorithm gains properly, we obtain that if the graphon is connected, then the \mathscr{L}^2 -consensus is achieved. By the key lemma established, we prove that if the local cost functions are strongly convex, then $\int_{[0,1]} E[||x_p(t)||^2] dp$ is uniformly bounded, which together with the \mathscr{L}^2 -consensus implies that the \mathscr{L}^∞ -consensus is also achieved. This in turn derives that all nodes' states converge to the minimizer of the global cost function uniformly in mean square. Besides, we qualify how the convergence rate of \mathscr{L}^2 -consensus relates to the parameters of the system dynamics (2), especially the algebraic connectivity of the graphon.

For the stochastic gradient tracking algorithm (3), which is indeed a double-variable system, the

convergence analysis is more challenging. Since the states and the auxiliary states are coupled by the time-varying algorithm gains, the analysis method for the system (2) is no longer applicable. At first, we use a transformation of the auxiliary states and transform the convergence analysis into the asymptotic properties of a class of coupled nonlinear differential inequalities with timevarying coefficients. Then we give a general lemma, in which we develop a decoupling method for the asymptotic properties of this kinds of coupled differential inequalities. We prove that if the graphon is connected and the local cost functions are strongly convex, then the \mathcal{L}^2 -consensus of the states and the transformed auxiliary states is achieved. In combination with the comparison theorem, we prove that the integral of the expectations of the nodes' states on the node set tends to the minimizer of the global cost function. Furthermore, by the relationship between the \mathcal{L}^2 -consensus and \mathcal{L}^∞ -consensus for the general system (4), we prove that if the local cost functions are strongly convex, then the nodes' states converge to the minimizer of the global cost function and the auxiliary states converge to the gradient value of the global cost function at the minimizer uniformly in mean square, respectively.

This work is the companion paper of [35], in which we have proved the existence and uniqueness of the solution to the system (4) and the law of large numbers.

The rest of the paper is organized as follows. In Section II, the relationship between the \mathcal{L}^2 consensus and \mathcal{L}^{∞} -consensus for the system (4) is established. In Section III, the convergence
of the stochastic gradient descent algorithm (2) is proved. In Section IV, the convergence of the
stochastic gradient tracking algorithm (3) is proved. In Section IV, the conclusions and future
works are given.

The following notations will be used throughout this paper. Denote the set of all real numbers by \mathbb{R} . Denote the *n*-dimensional Euclidean space by \mathbb{R}^n and the Euclidean norm by $\|\cdot\|$. For a given matrix $A \in \mathbb{R}^{n \times n}$, $\operatorname{Tr}(A)$ denotes its trace. For a given vector $x \in \mathbb{R}^n$, x^{T} denotes its transpose. For a given random vector $X \in \mathbb{R}^n$, denote its mathematical expectation and distribution by E[X]and $\mathscr{L}(X)$, respectively. Denote the set of probability measures on \mathbb{R}^n by $\mathscr{P}(\mathbb{R}^n)$. For a given measurable space (F, \mathscr{G}) and $x \in F$, where \mathscr{G} is a σ -algebra on F, the Dirac measure δ_x at x is the measure defined by $\delta_x(A) := \begin{cases} 1, & x \in A \\ 0, & x \notin A \end{cases}$, $\forall A \in \mathscr{G}$.

II. Relationship Between \mathscr{L}^2 -Consensus and \mathscr{L}^{∞} -Consensus

In this section, we prove that the \mathscr{L}^2 -consensus implies \mathscr{L}^{∞} -consensus for the system (4) under some conditions. See Appendix B for the proofs.

We make the following assumptions on the graphon, the local cost functions in (1) and the system (4).

Assumption 2.1:

- (i) Graphon A is connected.
- (ii) There exists a constant $\kappa > 0$, such that $\|\nabla_x V(p, x_1) \nabla_x V(p, x_2)\| \leq \kappa \|x_1 x_2\|, \forall x_1, x_2 \in \mathbb{R}^n, p \in [0, 1]$. There exist constants $\sigma_v > 0$ and $C_v > 0$, such that $\|\nabla_x V(p, x)\| \leq \sigma_v \|x\| + C_v$, $\forall x \in \mathbb{R}^n, p \in [0, 1]$.
- (iii) The local cost function V(p,x) is uniformly strongly convex w.r.t. x, that is, there exists $\kappa_2 > 0$, such that $(x_1 x_2)^{\mathsf{T}}(\nabla_x V(p, x_1) \nabla_x V(p, x_2)) \ge \kappa_2 ||x_1 x_2||^2, \forall x_1, x_2 \in \mathbb{R}^n, p \in [0,1].$

Assumption 2.2: There exists a nonnegative constant λ_1 , such that $||f(p,z_1,t) - f(p,z_2,t)|| + ||g(p,z_1,t) - g(p,z_2,t)|| \leq \lambda_1 ||z_1 - z_2||$, $\forall z_1, z_2 \in \mathbb{R}^m$, $t \ge 0$, $p \in [0,1]$; there exist nonnegative constants λ_{11} and λ_{12} such that $||f(p,z,t)|| + ||g(p,z,t)|| \leq \lambda_{11} ||z|| + \lambda_{12}$, $\forall z \in \mathbb{R}^m$, $t \ge 0$, $p \in [0,1]$; there exist nonnegative constants λ_3 and λ_4 , such that for any $\varepsilon > 0$, there exists $\delta > 0$, such that if $||t_1 - t_2|| < \delta$, then $||f(p,z,t_1) - f(p,z,t_2)||^2 + ||g(p,z,t_1) - g(p,z,t_2)||^2 < \varepsilon (\lambda_3 ||z||^2 + \lambda_4)$, $\forall t_1, t_2 \ge 0$, $z \in \mathbb{R}^m$, $p \in [0,1]$; f(p,z,t) and g(p,z,t) are measurable w.r.t. p, $\forall z \in \mathbb{R}^m$, $t \ge 0$; the map $[0,1] \ni p \mapsto \mu_{0,p} = \mathscr{L}(z_p(0)) \in \mathscr{P}(\mathbb{R}^m)$ is measurable and there exists a constant $\varsigma \ge 0$ such that $\sup_{p \in [0,1]} E[||z_p(0)||^2] \le \varsigma$.

Assumption 2.3: The map $[0,1] \ni p \mapsto \mathscr{L}(\xi_p(0)) \in \mathscr{P}(\mathbb{R}^m)$ is measurable; $E[\xi_p(t)] = 0, \forall p \in [0,1], t \ge 0$; there exists $r_1 \ge 0$, such that $\sup_{t\ge 0, p\in [0,1]} E[||\xi_p(t)||^2] \le r_1; \xi_p(\cdot)$ satisfies that, for any $\varepsilon > 0$, there exists $\delta > 0$, such that if $|t_1 - t_2| < \delta$, then $E[||\xi_p(t_1) - \xi_p(t_2)||^2] < \varepsilon, \forall t_1, t_2 \in [0,\infty), p \in [0,1].$

Assumption 2.4: The time-varying coefficients satisfy that $c_1(t) > 0$, $c_2(t) \ge 0$, $c_3(t) \ge 0$, $c_4(t) \ge 0$, $c_5(t) \ge 0$, $\forall t \ge 0$, $c_i(t)$, i = 1, ..., 5 are continuous w.r.t. t, $\lim_{t\to\infty} \frac{c_i(t)}{c_1(t)} = 0$, i = 2, ..., 5, $\int_0^{\infty} c_5^2(t) dt < \infty$, $\int_0^{\infty} c_1(t) = +\infty$ and $\lim_{t\to\infty} c_1(t) = 0$.

The following lemma illustrates that the variances of the nodes' states tend to zero.

Lemma 2.1: For the graphon particle system (4), if Assumption 2.1 (i) and Assumptions 2.2-2.4 hold, then

$$\lim_{t \to \infty} \sup_{p \in [0,1]} E\left[\|z_p(t) - E[z_p(t)]\|^2 \right] = 0.$$
(5)

To give the relation between the \mathscr{L}^2 -consensus and \mathscr{L}^∞ -consensus, we also need the following lemma to show the time-varying upper bounds of a class of functions satisfying time-varying differential inequalities with negative linear terms.

Lemma 2.2: If $y(\cdot): [0,\infty) \to [0,\infty)$ satisfy

$$y'(t) \leq -a_1(t)y(t) + a_2(t)\sqrt{y(t)} + a_3(t), \quad \forall t \geq 0,$$
 (6)

where $a_1(\cdot): [0,\infty) \to (0,\infty)$, $a_2(\cdot)$, $a_3(\cdot): [0,\infty) \to [0,\infty)$, and $a_i(\cdot)$, i = 1,2,3 are continuous, then

$$y(t) \leq \left(\frac{a_2(t)}{2a_1(t)} + \sqrt{\frac{1}{4}\frac{a_2^2(t)}{a_1^2(t)} + \frac{a_3(t)}{a_1(t)}}\right)^2, \quad \forall \ t \ge 0$$
(7)

and

$$y(t) \leq \max\left\{y(0), \left(\sup_{s \in [0,t]} \frac{a_2(s)}{2a_1(s)} + \left(\sup_{s \in [0,t]} \frac{1}{4} \frac{a_2^2(s)}{a_1^2(s)} + \sup_{s \in [0,t]} \frac{a_3(s)}{a_1(s)}\right)^{\frac{1}{2}}\right)^2\right\}, \quad \forall \ t \ge 0.$$
(8)

By Lemmas 2.1-2.2, we give the following theorem which shows that the \mathscr{L}^2 -consensus implies \mathscr{L}^{∞} -consensus for the system (4) if the integral of the second moments of all nodes' states on the node set is uniformly bounded.

Theorem 2.1: For the graphon particle system (4), if Assumption 2.1 (i) and Assumptions 2.2-2.4 hold, $\lim_{t\to\infty} \int_{[0,1]} ||E[z_p(t)] - \int_{[0,1]} E[z_q(t)] dq ||^2 dp = 0$ and $\sup_{t\geq 0} \int_{[0,1]} E[||z_p(t)||^2] dp < \infty$, then

$$\lim_{t \to \infty} \sup_{p \in [0,1]} \left\| E[z_p(t)] - \int_{[0,1]} E[z_q(t)] dq \right\|^2 = 0.$$
(9)

III. CONVERGENCE OF STOCHASTIC GRADIENT DESCENT ALGORITHM

In this section, we prove the convergence of the stochastic gradient descent algorithm (2). Denote $\mu_t(dx|q) = \mu_{t,q}(dx)$. Then $\mu_t(dx,dq) = \mu_{t,q}(dx)dq$. Therefore, (2) can be written as

$$dx_p(t) = \alpha_1(t) \int_{[0,1]} A(p,q) \left(\int_{\mathbb{R}^n} (x - x_p(t)) \mu_{t,q}(dx) \right) dq dt$$
$$- \alpha_2(t) \nabla_x V(p, x_p(t)) dt - \alpha_2(t) \Sigma_1 dw_p(t).$$
(10)

We give the following assumptions on the algorithm (2).

Assumption 3.1: The map $[0,1] \ni p \mapsto \mu_{0,p} = \mathscr{L}(z_p(0)) \in \mathscr{P}(\mathbb{R}^n)$ is measurable and there exists $\zeta_2 > 0$ such that $\sup_{p \in [0,1]} E[\|x_p(0)\|^2] \leq \zeta_2$.

Assumption 3.2: The time-varying algorithm gains satisfy that $\alpha_1(t) > 0$, $\alpha_2(t) > 0$, $\forall t \ge 0$, $\alpha_1(t)$ and $\alpha_2(t)$ are continuous w.r.t. t, $\int_0^{\infty} \alpha_2(t) dt = \infty$, $\int_0^{\infty} \alpha_2^2(t) dt < \infty$, $\lim_{t\to\infty} \frac{\alpha_2(t)}{\alpha_1(t)} = 0$ and $\lim_{t\to\infty} \alpha_1(t) = 0$.

The following lemma illustrates that all nodes' states achieve \mathscr{L}^{∞} -consensus.

Lemma 3.1: For the problem (1) and the algorithm (10), if Assumption 2.1 and Assumptions 3.1-3.2 hold, then there exists $K_0 \ge 0$, such that

$$\sup_{t \ge 0, p \in [0,1]} E\left[\|x_p(t)\|^2 \right] \le K_0,$$

$$\int_{[0,1]} \left\| E[x_p(t)] - \int_{[0,1]} E[x_q(t)] dq \right\|^2 dp$$
(11)

$$\leq \Psi_0(0,t)\zeta + \int_0^t 8\left(\sigma_v K_0 + C_v K_0^{\frac{1}{2}}\right) \alpha_2(s)\Psi_0(s,t)ds,\tag{12}$$

$$\lim_{t \to \infty} \sup_{p \in [0,1]} \left\| E[x_p(t)] - \int_{[0,1]} E[x_q(t)] dq \right\|^2 = 0,$$
(13)

where $\Psi_0(s,t) = e^{-2\lambda_2(\mathbb{L}_A)\int_s^t \alpha_1(s')ds'}$ and $\lambda_2(\mathbb{L}_A)$ is the algebraic connectivity of the graphon *A* defined by (A.1).

Proof 1: See Appendix C for the proof.

Then we prove that the integral of the expectations of the states on the node set converges to the minimizer of the global cost function. By Assumption 2.1 (iii), we know that V(x) is strongly convex w.r.t. x and $\nabla_x V(p,x)$ is continuous w.r.t. x. Then, $\nabla_x V(x^*) = \int_{[0,1]} \nabla_x V(p,x^*) dp = 0$.

Lemma 3.2: For the problem (1) and the algorithm (10), if Assumption 2.1 and Assumptions 3.1-3.2 hold, then $\lim_{t\to\infty} \|\int_{[0,1]} E[x_p(t)]dp - x^*\|^2 = 0.$

Proof 2: See Appendix C for the proof.

Finally, we show that the state of each node converges to the minimizer of the global cost function in mean quare.

Theorem 3.1: For the problem (1) and the algorithm (2), if Assumption 2.1 and Assumptions 3.1-3.2 hold, then

$$\lim_{t \to \infty} \sup_{p \in [0,1]} E\left[\|x_p(t) - x^*\|^2 \right] = 0.$$
(14)

Proof 3: By C_r inequality, we have $\sup_{p \in [0,1]} E[\|x_p(t) - x^*\|^2] \leq 3 \sup_{p \in [0,1]} E[\|x_p(t) - E[x_p(t)]\|^2] + 3 \sup_{p \in [0,1]} \|E[x_p(t)] - \int_{[0,1]} E[x_q(t)] dq \|^2 + 3 \|\int_{[0,1]} E[x_q(t)] dq - x^* \|^2$. This together with Lemmas 3.1-3.2 and Lemma C.1 gives (14).

Remark 3.1: The graphon particle system (2) is equivalent to the following system in the sense of weak solution. Given the initial state $x(0) = x_P(0)$,

$$dx(t) = \alpha_1(t) \int_{\mathbb{R}^n \times [0,1]} A(P,q)(x - x(t)) \mu_t(dx, dq) dt - \alpha_2(t) \nabla_x V(P, x(t)) dt - \alpha_2(t) \Sigma_1 dw(t), \quad (15)$$

where *P* is uniformly distributed on [0,1] and for any $t \ge 0$, $\mu_t(dx, dq)$ is the distribution on $\mathbb{R}^n \times [0,1]$ and satisfies the following conditions. (i) The marginal distribution $\mu_t(dq)$ is always the uniform distribution on [0,1], that is, $\mu_t(dq) = dq$, $\forall t \ge 0$. (ii) The marginal distribution $\mu_t(dx) = \int_{[0,1]} \mu_t(dx|q) dq$ is the distribution of x(t). Here, $\{w(t), t \ge 0\}$ is an *n*-dimensional standard Brownian motion. From Theorem 3.1, we know that $\mu_t(dx|q)$ in (15) converges to $\delta_{x^*}(dx)$ uniformly. Then, the distribution $\mu_t(dx)$ converges to $\delta_{x^*}(dx)$.

IV. CONVERGENCE OF STOCHASTIC GRADIENT TRACKING ALGORITHM

In this section, we prove the convergence of the stochastic gradient tracking algorithm (3).

We give some assumptions on the system (3).

Assumption 4.1: The time-varying algorithm gains satisfy that $\beta_1(t) > 0$, $\beta_2(t) > 0$, $\beta_3(t) > 0$, $\forall t \ge 0$, $\beta_2(0) = 1$, $\beta_1(t)$ and $\beta_3(t)$ are continuous w.r.t. t, $\beta_2(t)$ is differentiable w.r.t. t, $\int_0^{\infty} \beta_3(t) dt = \infty$, $\int_0^{\infty} \beta_1(t) \beta_2(t) dt = \infty$, $\lim_{t \to \infty} \frac{\beta_1(t)}{\beta_3(t)} = 0$, $\lim_{t \to \infty} \frac{\beta_2(t)}{\beta_3(t)} = 0$ and $\lim_{t \to \infty} \beta_3(t) = 0$.

Assumption 4.2: The map $[0,1] \ni p \mapsto \mathscr{L}(z_p(0), y_p(0)) \in \mathscr{P}(\mathbb{R}^{2n})$ is measurable and there exist ζ and $\zeta_0 > 0$ such that $\sup_{p \in [0,1]} E[\|z_p(0)\|^2] \leqslant \zeta$ and $\sup_{p \in [0,1]} E[\|y_p(0)\|^2] \leqslant \zeta_0$.

Assumption 4.3: The map $[0,1] \ni p \mapsto \mathscr{L}(\eta_p(0)) \in \mathscr{P}(\mathbb{R}^n)$ is measurable; $E[\eta_p(t)] = 0, \forall p \in [0,1], t \ge 0$; there exists $b_1 \ge 0$ such that $\sup_{t\ge 0, p\in[0,1]} E[\|\eta_p(t)\|^2] \le b_1$; for any $\varepsilon > 0$, there exists $\delta > 0$, such that if $|t_1 - t_2| < \delta$, then $E[\|\eta_p(t_1) - \eta_p(t_2)\|^2] < \varepsilon, \forall t_1, t_2 \in [0,\infty), p \in [0,1]$.

Inspired by [25], by the transformation $\tilde{y}_p(t) = y_p(t) - \beta_2(t) \nabla_x V(p, z_p(t))$, we have the following transformed graphon particle system

$$\begin{cases} dz_{p}(t) = \left(-\beta_{1}(t)\widetilde{y}_{p}(t) - \beta_{1}(t)\beta_{2}(t)\nabla_{x}V(p,z_{p}(t))\right)dt \\ +\beta_{3}(t)\int_{[0,1]\times\mathbb{R}^{n}}A(p,q)(z-z_{p}(t))\mu_{t,q}(dz)dqdt \\ -\beta_{1}(t)\beta_{2}(t)\beta_{3}(t)\int_{[0,1]\times\mathbb{R}^{n}}A(p,q)\left(\nabla_{x}V(q,z) - \nabla_{x}V(p,z_{p}(t))\right)\mu_{t,q}(dz)dqdt \\ -\beta_{1}(t)\beta_{3}(t)\int_{[0,1]\times\mathbb{R}^{n}}A(p,q)(y-\widetilde{y}_{p}(t))\widetilde{v}_{t,q}(dy)dqdt, \\ d\widetilde{y}_{p}(t) = \beta_{3}(t)\int_{[0,1]\times\mathbb{R}^{n}}A(p,q)(y-\widetilde{y}_{p}(t))\widetilde{v}_{t,q}(dy)dqdt + \beta_{2}(t)\eta_{p}(t)dt \\ +\beta_{2}(t)\beta_{3}(t)\int_{[0,1]\times\mathbb{R}^{n}}A(p,q)\left(\nabla_{x}V(q,z) - \nabla_{x}V(p,z_{p}(t))\right)\mu_{t,q}(dz)dqdt, \end{cases}$$
(16)

where $\mu_{t,q}(dz)$ and $\tilde{v}_{t,q}(dy)$ are the distributions of $z_p(t)$ and $\tilde{y}_p(t)$. Here, $\tilde{y}_p(t)$ is called the transformed auxiliary state.

We transform the convergence analysis of the algorithm (16) into the asymptotic properties of a class of coupled differential inequalities with time-varying coefficients and develop a decoupling method in the following lemma.

Lemma 4.1: If
$$Y_1(\cdot), Y_2(\cdot) : [0, \infty) \to [0, \infty)$$
 are differentiable and

$$\frac{dY_1(t)}{dt} \leq (-a_1(t) + a_2(t))Y_1(t) + a_3(t)Y_2(t) + a_4(t), \qquad (17)$$

$$\frac{dY_2(t)}{dt} \leq (-a_1(t) + a_2(t))Y_1(t) + a_3(t)Y_2(t) + a_4(t), \qquad (17)$$

$$\frac{dY_2(t)}{dt} \leqslant -b_1(t)Y_2(t) + b_2(t)Y_2^{\frac{1}{2}}(t)\left(Y_1^{\frac{1}{2}}(t) + Y_3(t)\right)$$
(18)

hold, where the time-varying coefficients satisfy that $a_1(t) > 0$, $a_i(t) \ge 0$, i = 2, 3, 4, $\lim_{t\to\infty} \frac{a_2(t)}{a_1(t)} = 0$, $\lim_{t\to\infty} \frac{a_3(t)}{a_1(t)} = 0$, $\lim_{t\to\infty} \frac{a_4(t)}{a_1(t)} = 0$, $\int_0^\infty a_1(t)dt = \infty$, $b_1(t) > 0$, $b_2(t) \ge 0$, $b_1(t)$ and $b_2(t)$ are continuous w.r.t. t, $\int_0^\infty b_1(t) = \infty$, $\limsup_{t\to\infty} \frac{b_2(t)}{b_1(t)} < \infty$ and $\lim_{t\to\infty} Y_3(t) = 0$, then

$$\lim_{t \to \infty} Y_1(t) = 0, \tag{19}$$

$$\lim_{t \to \infty} Y_2(t) = 0. \tag{20}$$

Proof 4: See Appendix D for the proof.

Remark 4.1: The main idea of decoupling inequalities in the above lemma lies in that the time-varying coefficients of (18) have same orders, which together with Lemma 2.2 shows that $Y_2(t)$ can be bounded by $Y_1(t)$. Replacing the upper bound of $Y_2(t)$ into the inequality of $Y_1(t)$ and using the comparison theorem, we can show (19) and then (20) follows.

By the above lemma and Theorem 2.1, we show that the states and transformed auxiliary states achieve \mathscr{L}^{∞} -consensus and the integral of the expectations of the states on the node set tends to the minimizer of the global cost function.

Lemma 4.2: For the problem (1) and the stochastic gradient tracking algorithm (16), if Assumption 2.1 and Assumptions 4.1-4.3 hold, then

$$\sup_{t \ge 0, p \in [0,1]} E\left[\|z_p(t)\|^2 \right] < \infty,$$
(21)

$$\sup_{t \ge 0, p \in [0,1]} E\left[\|\widetilde{y}_p(t)\|^2 \right] < \infty,$$
(22)

$$\lim_{t \to \infty} \sup_{p \in [0,1]} \left\| E[z_p(t)] - \int_{[0,1]} E[z_q(t)] dq \right\|^2 = 0,$$
(23)

$$\lim_{t \to \infty} \sup_{p \in [0,1]} \left\| E\left[\widetilde{y}_p(t)\right] \right\|^2 = 0,$$
(24)

$$\lim_{t \to \infty} \left\| \int_{[0,1]} E[z_p(t)] dp - x^* \right\|^2 = 0.$$
(25)

Proof 5: See Appendix D for the proof.

The following theorem shows that all nodes' states and auxiliary states converge to the minimizer of the global cost function and the gradient value of the global cost function at the minimizer uniformly in mean square, respectively.

Theorem 4.1: For the problem (1) and the stochastic gradient tracking algorithm (3), if Assumption 2.1 and Assumptions 4.1-4.3 hold, then

$$\lim_{t \to \infty} \sup_{p \in [0,1]} E\left[\|z_p(t) - x^*\|^2 \right] = 0,$$
(26)

$$\lim_{t \to \infty} \sup_{p \in [0,1]} E\left[\left\| y_p(t) - \nabla_x \left(\int_{[0,1]} V(q, x^*) dq \right) \right\|^2 \right] = 0.$$
(27)

Proof 6: By Lemma D.1, Lemma 4.2 and C_r inequality, we have (26). Similar to the proof of (D.11) in Lemma 4.2 and by Assumption 2.1 (iii), we have $\int_{[0,1]} E[\tilde{y}_q(t)]dq = 0$ and $\nabla_x (\int_{[0,1]} V(q, x^*)dq) = 0$. This together with Assumption 2.1 (ii) and C_r inequality gives

$$E\left[\left\|y_{p}(t) - \nabla_{x}\left(\int_{[0,1]} V(q,x^{*})dq\right)\right\|^{2}\right]$$

$$\leq 3E\left[\left\|\widetilde{y}_{p}(t) - E[\widetilde{y}_{p}(t)]\right\|^{2}\right] + 3\beta_{2}^{2}(t)E\left[\left\|\nabla_{x}V(z_{p}(t),p)\right\|^{2}\right] + 3\left\|E[\widetilde{y}_{p}(t)]\right\|^{2}$$

$$\leq 3E\left[\left\|\widetilde{y}_{p}(t) - E[\widetilde{y}_{p}(t)]\right\|^{2}\right] + 6C_{v}^{2}\beta_{2}^{2}(t) + 3\left\|E[\widetilde{y}_{p}(t)]\right\|^{2} + 6\sigma_{v}^{2}\beta_{2}^{2}(t)\sup_{p\in[0,1],\ t\geq0} E\left[\left\|z_{p}(t)\right\|^{2}\right]. \quad (28)$$

By (26) and C_r inequality, we have $\sup_{p \in [0,1], t \ge 0} E[||z_p(t)||^2] < \infty$. Then, by Assumption 4.1, Lemma D.1, Lemma 4.2 and (28), we have (27).

We have proposed the stochastic gradient descent and gradient tracking algorithms over the graphon for solving the distributed optimization problem with a continuum of nodes. By establishing the lemma for the upper bound estimation related to a class of time-varying differential inequalities with negative linear terms, we have proved the uniform boundness of the second moments of the nodes' states in both kinds of algorithms. Besides, we have proved that if the graphon is connected and the time-varying algorithm gains are chosen properly, then the states in both kinds of algorithms achieve \mathscr{L}^{∞} -consensus. Moreover, if the local cost functions are strongly convex, then the states in both kinds of algorithms converge to the minimizer of the global cost function and the auxiliary states in the stochastic gradient tracking algorithm converge to the global cost function at the minimizer uniformly in mean square.

Note that the analysis of the asymptotic properties of the graphon particle systems relies on the special linear interactions among nodes in the proposed two kinds of algorithms, while for many graphon particle systems, such as Kuramoto oscillator ([26]), neural mean-field ([27]), SIS epidemics ([28]) and so on, the interactions are nonlinear. This results in that the methods in this paper are inapplicable. Besides, the graphon considered in this paper is static. In many practical scenarios, networks among nodes receive the feedback from nearby individuals and then make changes to better adapt to the world, such as adaptive Kuramoto-type network models in [29], which leads to a dynamic graphon. The asymptotic properties of the graphon particle systems with dynamic graphons are still open so far, which is of major importance from an applied perspective but highly mathematically challenging.

APPENDIX A

PROPERTIES OF GRAPHONS

The following notations will be used in the appendix. Denote $L^2([0,1],\mathbb{R}^n) = \{f : [0,1] \to \mathbb{R}^n, f \text{ is measurable}, \int_{[0,1]} ||f(x)||^2 dx < \infty\}$. Denote the set of all bounded linear operators from $L^2([0,1],\mathbb{R}^n)$ to $L^2([0,1],\mathbb{R}^n)$ by $\mathscr{L}(L^2([0,1],\mathbb{R}^n))$. Denote the inner product on $L^2([0,1],\mathbb{R}^n)$ by $\langle \cdot, \cdot \rangle_{L^2([0,1],\mathbb{R}^n)}$, that is, for any given $f, g \in L^2([0,1],\mathbb{R}^n), \langle f, g \rangle_{L^2([0,1],\mathbb{R}^n)} \triangleq \int_{[0,1]} f^{\mathsf{T}}(x)g(x)dx$. For a given function $f : F \to \mathbb{R}$, $\operatorname{supp}(f) = \{x \in F : f(x) \neq 0\}$ denotes the support set of f.

For a given graphon W, the Graphon-Laplacian $\mathbb{L}_W \in \mathscr{L}(L^2([0,1], \mathbb{R}^n))$ generated by W is given by, for any $z \in L^2([0,1], \mathbb{R}^n)$, $(\mathbb{L}_W z)(p) = \int_{[0,1]} W(p,q)(z(p) - z(q))dq$, $\forall p \in [0,1]$. For

a graphon W, the algebraic connectivity of W is defined by

$$\lambda_2(\mathbb{L}_W) = \inf_{z \in \mathscr{C}^\perp} \frac{\langle \mathbb{L}_W z, z \rangle_{L^2([0,1], \mathbb{R}^n)}}{\langle z, z \rangle_{L^2([0,1], \mathbb{R}^n)}^2} \ge 0, \tag{A.1}$$

where $\mathscr{C}^{\perp} = \{z \in L^2([0,1], \mathbb{R}^n) : \int_{[0,1]} z(p) dp = 0\}$. By Proposition 4.9 in [30], for the graphon *W*, the algebraic connectivity can also be written as

$$\lambda_2(\mathbb{L}_W) = \inf_{z \in \mathscr{C}^\perp} \frac{\int_{[0,1] \times [0,1]} W(p,q) z^{\mathsf{T}}(p)(z(p) - z(q)) dq dp}{\int_{[0,1]} \|z(p) - \int_{[0,1]} z(q) dq \|^2 dp}.$$
 (A.2)

Definition A.1: ([30]) For a graphon W, if the following conditions hold, then the graphon W is said to be connected.

- (i) For any $p \in [0,1]$ and $q \in [0,1] \setminus \{p\}$, there exists an integer $m \ge 1$ and a finite sequence $(l_k)_{1 \le k \le m} \subset [0,1]$ satisfying that $p = l_1$, $q = l_m$ and $l_{k+1} \in \text{supp}(W(l_k, \cdot)), \forall k \in \{1, \dots, m-1\}$.
- (ii) $\inf_{p \in [0,1]} \int_{[0,1]} W(p,q) \mathrm{d}q > 0.$

The following lemma shows the connection between the algebraic connectivity and the connectivity of a graphon.

Lemma A.1: ([30]) The graphon W is connected in the sense of Definition A.1 if and only if $\lambda_2(\mathbb{L}_W) > 0$.

APPENDIX B

Proof of Lemma 2.1: Noting that $\mu_{t,p}$ is the distribution of $z_p(t)$ in (4), the system (4) can be written as

$$dz_{p}(t) = \left[c_{2}(t)\int_{[0,1]}A(p,q)\left(E\left[f(q,z_{q}(t),t)\right] - f(p,z_{p}(t),t)\right)dq + c_{1}(t)\int_{[0,1]}A(p,q)\left(E\left[z_{q}(t)\right] - z_{p}(t)\right)dq + c_{4}(t)\xi_{p}(t) + c_{3}(t)g\left(p,z_{p}(t)\right)\right]dt + c_{5}(t)\Sigma dw_{p}(t).$$
(B.1)

By Assumption 2.4 and Theorem 2.3.1 in [31], we have $E\left[\int_0^t c_5(s)\Sigma dw_p(s)\right] = 0$. Then, by Assumption 2.3 and (B.1), we have

$$dE[z_{p}(t)] = c_{1}(t) \int_{[0,1]} A(p,q) \left(E[z_{q}(t)] - E[z_{p}(t)] \right) dq dt + c_{3}(t) E[g(p,z_{p}(t))] dt + c_{2}(t) \int_{[0,1]} A(p,q) \left(E[f(q,z_{q}(t),t)] - E[f(p,z_{p}(t),t)] \right) dq dt.$$
(B.2)

Denote $S_p(t) = ||z_p(t) - E[z_p(t)]||^2$. By Theorem 2.1 in [35] and Assumption 2.4, we have $E\left[\int_0^t ||c_5(s)(z_p(s) - E[z_p(s)])^{\mathsf{T}}\Sigma||^2 ds\right] \leq E\left[\sup_{0 \leq s \leq t} ||z_p(s)||^2\right] \int_0^t c_5^2(s) ds ||\Sigma||^2 < \infty$. Then, by Theorem 2.3.1 in [31], we have

$$\begin{split} E\left[\int_{0}^{t}c_{5}(s)(z_{p}(s)-E[z_{p}(s)])^{\mathsf{T}}\Sigma dw_{p}(s)\right] &= 0.\\ \text{By (B.2), } E\left[(z_{p}(t)-E[z_{p}(t)])^{\mathsf{T}}g(p,E[z_{p}(t)])\right] &= 0, \ E\left[(z_{p}(t)-E[z_{p}(t)])^{\mathsf{T}}(E[f(p,z_{p}(t),t)]\right] \\ &-f(p,E[z_{p}(t)],t))\right] &= 0, \ \text{Assumptions 2.2-2.3 and Itô formula, we have}\\ \frac{dE\left[S_{p}(t)\right]}{dt} \\ &= \int_{[0,1]} A(p,q)dq \left(-2c_{1}(t)E\left[S_{p}(t)\right]+2c_{2}(t)E\left[(z_{p}(t)-E[z_{p}(t)])^{\mathsf{T}}\left(E[f(p,z_{p}(t),t)]\right] \\ &-f(p,z_{p}(t),t)\right)\right] \right) + 2c_{3}(t)E\left[(z_{p}(t)-E[z_{p}(t)])^{\mathsf{T}}g(p,z_{p}(t))\right] + 2c_{4}(t)E\left[(z_{p}(t) - E[z_{p}(t)])^{\mathsf{T}}\xi_{p}(t)\right] + \operatorname{Tr}\left(\Sigma^{\mathsf{T}}\Sigma\right)c_{5}^{2}(t) \\ &\leqslant \left(-2c_{1}(t)\inf_{p\in[0,1]}\int_{[0,1]}A(p,q)dq + c_{4}(t)\right)E\left[S_{p}(t)\right] \\ &+ 2c_{2}(t)\int_{[0,1]}A(p,q)dqE\left[||z_{p}(t)-E[z_{p}(t)]||||f(p,z_{p}(t),t) - f(p,E[z_{p}(t)],t)||\right] \\ &+ 2c_{3}(t)E\left[||z_{p}(t)-E[z_{p}(t)]||||g(p,z_{p}(t)) - g(p,E[z_{p}(t)])||\right] + c_{4}(t)r_{1} + \operatorname{Tr}\left(\Sigma^{\mathsf{T}}\Sigma\right)c_{5}^{2}(t) \\ &\leqslant \phi(t)E\left[S_{p}(t)\right] + c_{4}(t)r_{1} + \operatorname{Tr}(\Sigma^{\mathsf{T}}\Sigma)c_{5}^{2}(t), \end{split}$$

where $\phi(t) = -2c_1(t) \inf_{p \in [0,1]} \int_{[0,1]} A(p,q) dq + 2\lambda_1(c_2(t) + c_3(t)) + c_4(t)$. This together with the comparison theorem ([32]) gives

$$E\left[S_p(t)\right] \leqslant \int_0^t \left(c_4(s)r_1 + \operatorname{Tr}(\Sigma^{\mathsf{T}}\Sigma)c_5^2(s)\right) e^{\int_s^t \phi(s')ds'} ds + e^{\int_0^t \phi(s)ds} E[S_p(0)].$$

By Assumption 2.2, we have $\sup_{p \in [0,1]} E[||S_p(0)||^2] \leq \sup_{p \in [0,1]} E[||z_p(0)||^2] \leq \varsigma$. Then, by the above inequality, we have

$$\sup_{p\in[0,1]} E\left[S_p(t)\right] \leqslant e^{\int_0^t \phi(s)ds} \varsigma + \int_0^t \left(c_4(s)r_1 + \operatorname{Tr}(\Sigma^{\mathsf{T}}\Sigma)c_5^2(s)\right) e^{\int_s^t \phi(s')ds'} ds.$$
(B.3)

By Assumption 2.4, we know that there exists $T \ge 0$, such that if $t \ge T$, then

$$2\lambda_1 \frac{(c_2(t) + c_3(t))}{c_1(t)} + \frac{c_4(t)}{c_1(t)} \leqslant \inf_{p \in [0,1]} \int_{[0,1]} A(p,q) dq$$

which together with $\int_0^{\infty} c_1(t) dt = \infty$, Assumption 2.1 (i) and Definition A.1 gives

$$\int_0^\infty \phi(s)ds = -\infty. \tag{B.4}$$

For the first term on the r.h.s. of (B.3), by the above equality, we have

$$\lim_{t \to \infty} e^{\int_0^t \phi(s) ds} \varsigma = 0. \tag{B.5}$$

For the second term on the r.h.s. of (B.3), by Assumption 2.1 (i), Assumption 2.4, (B.4) and L'Hospital's rule, we have

$$\lim_{t\to\infty}\int_0^t (c_4(s)r_1 + \operatorname{Tr}(\Sigma^{\mathsf{T}}\Sigma)c_5^2(s))e^{\int_s^t \phi(s')ds'}ds = 0,$$

which together with (B.3) and (B.5) gives (5).

Proof of Lemma 2.2: By (6) and $a_1(t) > 0$, $\forall t \ge 0$, we have

$$y'(t) \leq -a_1(t) \left(\sqrt{y(t)} - \frac{a_2(t)}{2a_1(t)}\right)^2 + \frac{a_2^2(t)}{4a_1(t)} + a_3(t).$$

Therefore, we know that if $\sqrt{y(t)} > \frac{a_2(t)}{2a_1(t)} + \left(\frac{1}{4}\frac{a_2^2(t)}{a_1^2(t)} + \frac{a_3(t)}{a_1(t)}\right)^{\frac{1}{2}}$, then y'(t) < 0 and $y(t) \le \left(\frac{a_2(t)}{2a_1(t)} + \sqrt{\frac{1}{4}\frac{a_2^2(t)}{a_1^2(t)} + \frac{a_3(t)}{a_1(t)}}\right)^2$, $\forall t \ge 0$, which leads to (8).

Proof of Theorem 2.1: Denote $B_p(t) = E\left[\|z_p(t)\|^2\right]$ and $R_p(t) = \left\|E[z_p(t)] - \int_{[0,1]} E[z_q(t)]dq\right\|^2$. By $\sup_{t\geq 0} \int_{[0,1]} E\left[\|z_p(t)\|^2\right] dp < \infty$, we know that there exists $K_1 \geq 0$, such that

$$\sup_{t \ge 0} \int_{[0,1]} E\left[\|z_p(t)\|^2 \right] dp \leqslant K_1.$$
(B.6)

By Theorem 2.1 in [35] and Assumption 2.4, we have $E\left[\int_0^t ||2c_5(s)z_p^{\mathsf{T}}(s)\Sigma||^2 ds\right] \leq 4||\Sigma||^2 \sup_{s \in [0,t]} E\left[||z_p(s)||^2\right] \int_0^t c_5^2(s) ds < \infty$. Then, by Theorem 2.3.1 in [31], we have

$$E\left[\int_0^t 2c_5(s) z_p^{\mathsf{T}}(s) \Sigma dw_p(s)\right] = 0,$$

which together with Assumptions 2.2-2.3, Itô formula, (4), (B.6), Hölder inequality and Jensen inequality gives

$$\begin{aligned} \frac{dB_{p}(t)}{dt} \\ =& 2 \int_{[0,1]} A(p,q) \left(E[z_{p}^{\mathsf{T}}(t)] E[z_{q}(t)] - E[z_{p}^{\mathsf{T}}(t)z_{p}(t)] \right) dqc_{1}(t) \\ &+ 2c_{2}(t) \int_{[0,1]} A(p,q) \left(E[z_{p}^{\mathsf{T}}(t)] E[f(q,z_{q}(t),t)] - E[z_{p}^{\mathsf{T}}(t)f(p,z_{p}(t),t)] \right) dq \\ &+ 2c_{4}(t) E[z_{p}^{\mathsf{T}}(t)\xi_{p}(t)] + 2c_{3}(t) E[z_{p}^{\mathsf{T}}(t)g(p,z_{p}(t))] + \operatorname{Tr}(\Sigma^{\mathsf{T}}\Sigma)c_{5}^{2}(t) \\ \leqslant \left(-2 \int_{[0,1]} A(p,q) dqc_{1}(t) + 2c_{2}(t) + c_{3}(t) + c_{4}(t) \right) B_{p}(t) \\ &+ 2c_{1}(t) \int_{[0,1]} \|E[z_{q}(t)]\| dq \|E[z_{p}(t)]\| + c_{2}(t) \left(\int_{[0,1]} \|E[f(q,z_{q}(t),t)]\|^{2} dq \end{aligned}$$

July 4, 2024

$$+ \left[\|f(p, z_{p}(t), t)\|^{2} \right] + c_{3}(t)E\left[\|g(p, z_{p}(t))\|^{2} \right]$$

$$+ \operatorname{Tr}(\Sigma^{\mathsf{T}}\Sigma)c_{5}^{2}(t) + c_{4}(t)E\left[\|\xi_{p}(t)\|^{2} \right]$$

$$\leq \left(-2\inf_{p\in[0,1]} \int_{[0,1]} A(p,q)dqc_{1}(t) + 2c_{2}(t) + c_{3}(t) + c_{4}(t) \right)B_{p}(t)$$

$$+ 2c_{1}(t)\left(\int_{[0,1]} \|E[z_{q}(t)]\|^{2}dq \right)^{\frac{1}{2}}E\left[\|z_{p}(t)\| \right] + c_{2}(t)\int_{[0,1]} \|E[f(q, z_{q}(t), t)]\|^{2}dq$$

$$+ c_{2}(t)E\left[\|f(p, z_{p}(t), t)\|^{2} \right] + c_{4}(t)r_{1} + c_{3}(t)E\left[\|g(p, z_{p}(t))\|^{2} \right] + \operatorname{Tr}(\Sigma^{\mathsf{T}}\Sigma)c_{5}^{2}(t)$$

$$\leq h_{1}(t)B_{p}(t) + 2c_{1}(t)K_{1}^{\frac{1}{2}}B_{p}^{\frac{1}{2}}(t) + 2\lambda_{12}^{2}c_{2}(t) + 2\lambda_{11}^{2}c_{3}(t) + c_{4}(t)r_{1} + \operatorname{Tr}(\Sigma^{\mathsf{T}}\Sigma)c_{5}^{2}(t)$$

$$\leq h_{1}(t)B_{p}(t) + 2c_{1}(t)K_{1}^{\frac{1}{2}}B_{p}^{\frac{1}{2}}(t) + 2\lambda_{12}^{2}c_{3}(t) + c_{4}(t)r_{1}$$

$$+ \operatorname{Tr}(\Sigma^{\mathsf{T}}\Sigma)c_{5}^{2}(t) + 2c_{2}(t)\left(2\lambda_{12}^{2} + \lambda_{11}^{2}\int_{[0,1]}E\left[\|z_{q}(t)\|^{2} \right]dq \right)$$

$$\leq h_{1}(t)B_{p}(t) + h_{2}(t)B_{p}^{\frac{1}{2}}(t) + h_{3}(t),$$
(B.7)

where $h_1(t) = -2\inf_{p \in [0,1]} \int_{[0,1]} A(p,q) dq c_1(t) + (2+2\lambda_{11}^2)c_2(t) + (1+2\lambda_{11}^2)c_3(t) + c_4(t), h_2(t) = 2c_1(t)K_1^{\frac{1}{2}}$ and $h_3(t) = (2\lambda_{11}^2K_1 + 4\lambda_{12}^2)c_2(t) + 2\lambda_{12}^2c_3(t) + c_4(t)r_1 + \operatorname{Tr}(\Sigma^{\mathsf{T}}\Sigma)c_5^2(t))$. By Assumption 2.4, there exists $T_2 \ge 0$, such that

$$\left((2+2\lambda_{11}^2)c_2(t)+(1+2\lambda_{11}^2)c_3(t)+c_4(t)\right)\frac{1}{c_1(t)}<2\inf_{p\in[0,1]}\int_{[0,1]}A(p,q)dq,\;\forall\;t\geqslant T_2,$$

that is, $h_1(t) < 0$, $\forall t \ge T_2$. By Theorem 2.1 in [35], there exists $L_0 \ge 0$, such that $\sup_{p \in [0,1], t \in [0,T_2]} B_p(t) \le L_0$. For any $t \ge T_2$, by (B.7) and Lemma 2.2, we have

$$B_p(t) \le \max\left\{B_p(T_2), \left(\sup_{s \in [T_2,t]} \frac{h_2(s)}{-2h_1(s)} + \left(\sup_{s \in [T_2,t]} \frac{1}{4} \frac{h_2^2(s)}{h_1^2(s)} + \sup_{s \in [T_2,t]} \frac{h_3(s)}{-h_1(s)}\right)^{\frac{1}{2}}\right)^2\right\}.$$
 (B.8)

By Assumption 2.4, we have

$$\lim_{t \to \infty} \frac{h_2(t)}{-2h_1(t)} = \left(\inf_{p \in [0,1]} \int_{[0,1]} A(p,q) dq\right)^{-1} \frac{K_1^{\frac{1}{2}}}{2}$$

 $\lim_{t \to \infty} \frac{h_3(t)}{-h_1(t)} = 0 \text{ and } \lim_{t \to \infty} \frac{1}{4} \frac{2h_2^2(t)}{h_1^2(t)} = \frac{K_1}{4} \left(\inf_{p \in [0,1]} \int_{[0,1]} A(p,q) dq \right)^{-2}. \text{ Then, there exist non-negative constants } L_1, L_2 \text{ and } L_3, \text{ such that } \sup_{t \ge 0} \frac{h_2(t)}{-2h_1(t)} \le L_1, \sup_{t \ge 0} \frac{1}{4} \frac{h_2^2(t)}{h_1^2(t)} \le L_2 \text{ and } \sup_{t \ge 0} \frac{h_3(t)}{-h_1(t)} \le L_3. \text{ Denote } L = \max\{L_0, (\sqrt{L_2 + L_3} + L_1)^2\}. \text{ Then, by (B.8), we have } \sup_{t \ge 0} B_p(t) \le L. \text{ Noting that } L \text{ is independent of } p, \text{ we have}$

$$\sup_{p \in [0,1], \ t \ge 0} E\left[\|z_p(t)\|^2 \right] \le L.$$
(B.9)

By $\lim_{t\to\infty} \int_{[0,1]} R_p(t) dp = 0$, we know that, for any $\varepsilon > 0$, there exists $T_{\varepsilon} > 0$, such that if $t \ge T_{\varepsilon}$, then $\int_{[0,1]} R_p(t) dp < \varepsilon^2$. For any $t \ge T_{\varepsilon}$, denote $S_{\varepsilon}^t = \{p \in [0,1] : R_p(t) > \varepsilon\}$. By Theorem 2.1 in [35] and (5.3.1) in [33], we know that S_{ε}^t is a measurable set of [0,1]. For any $t \ge T_{\varepsilon}$, we have

$$\varepsilon m(S^t_{\varepsilon}) < \int_{S^t_{\varepsilon}} R_p(t) dp \leqslant \int_{[0,1]} R_p(t) dp < \varepsilon^2,$$

that is, $m(S_{\varepsilon}^t) < \varepsilon$. Taking the derivative of $R_p(t)$ on $t \ge T_{\varepsilon}$ and combining (B.2) in Lemma 2.1 with the symmetry of the graphon A give

$$\frac{dR_{p}(t)}{dt} = -2c_{1}(t)\int_{[0,1]}A(p,q)dqR_{p}(t) + 2c_{1}(t)\left(E[z_{p}(t)] - \int_{[0,1]}E[z_{q}(t)]dq\right)^{\mathsf{T}}\left(\int_{[0,1]}A(p,q)\left(E[z_{q}(t)]\right) \\
-\int_{[0,1]}E[z_{q'}(t)]dq'\right)dq\right) + 2c_{2}(t)\left(E[z_{p}(t)] - \int_{[0,1]}E[z_{q}(t)]dq\right)^{\mathsf{T}}\left(\int_{[0,1]}A(p,q)\right) \\
\times \left(E[f(q,z_{q}(t),t)] - E[f(p,z_{p}(t),t)]\right)dq\right) + 2c_{3}(t)\left(E[z_{p}(t)] - \int_{[0,1]}E[z_{q}(t)]dq\right)^{\mathsf{T}} \\
\times \left(E[g(p,z_{p}(t))] - \int_{[0,1]}E[g(q,z_{q}(t))]dq\right).$$
(B.10)

For the second term on the r.h.s. of the above equality, by C_r inequality, Hölder inequality, Jensen inequality and (B.9), we have

$$2c_{1}(t)\left(E[z_{p}(t)] - \int_{[0,1]} E[z_{q}(t)]dq\right)^{\mathsf{T}}\left(\int_{[0,1]} A(p,q)\left(E[z_{q}(t)] - \int_{[0,1]} E[z_{q'}(t)]dq'\right)dq\right)$$

$$\leq 2c_{1}(t)R_{p}^{\frac{1}{2}}(t)\left(\int_{[0,1]} R_{q}(t)dq\right)^{\frac{1}{2}}$$

$$\leq 2c_{1}(t)R_{p}^{\frac{1}{2}}(t)\varepsilon^{\frac{1}{2}}\left(\sup_{q\in[0,1]} R_{q}(t) + 1\right)^{\frac{1}{2}}$$

$$\leq 2c_{1}(t)R_{p}^{\frac{1}{2}}(t)\varepsilon^{\frac{1}{2}}\left(2\sup_{q\in[0,1]} \left\|E[z_{q}(t)]\right\|^{2} + 2\left\|\int_{[0,1]} E[z_{q'}(t)]dq'\right\|^{2} + 1\right)^{\frac{1}{2}}$$

$$\leq 2c_{1}(t)R_{p}^{\frac{1}{2}}(t)\varepsilon^{\frac{1}{2}}\left(2L + 2\int_{[0,1]} E\left[\left\|z_{q'}(t)\right\|^{2}\right]dq' + 1\right)^{\frac{1}{2}}$$

$$\leq 2c_{1}(t)R_{p}^{\frac{1}{2}}(t)\varepsilon^{\frac{1}{2}}(4L + 1)^{\frac{1}{2}}.$$
(B.11)

For the third term on the r.h.s. of (B.10), by C_r inequality, Hölder inequality, Jensen inequality, Assumption 2.2 and (B.9), we have

$$-2c_{2}(t)\left(E[z_{p}(t)]-\int_{[0,1]}E[z_{q}(t)]dq\right)^{\mathsf{T}}\left(\int_{[0,1]}A(p,q)\left(E\left[f\left(q,z_{q}(t),t\right)\right]\right)$$

$$\times -E\left[f(p,z_{p}(t),t)\right] dq$$

$$\leq c_{2}(t) \left(R_{p}(t) + \left\| \int_{[0,1]} A(p,q) \left(E\left[f\left(q,z_{q}(t),t\right)\right] - E[f(p,z_{p}(t),t)]\right] dq \right\|^{2} \right)$$

$$\leq c_{2}(t) \left(R_{p}(t) + 2\left\| \int_{[0,1]} A(p,q) E\left[f\left(q,z_{q}(t),t\right)\right] dq \right\|^{2} + 2\left\|E\left[f(p,z_{p}(t),t)\right]\right\|^{2} \right)$$

$$\leq c_{2}(t) \left(R_{p}(t) + 2\int_{[0,1]} \left\|E\left[f\left(q,z_{q}(t),t\right)\right]\right\|^{2} dq + 2E\left[\left\|f(p,z_{p}(t),t)\right\|^{2}\right] \right)$$

$$\leq c_{2}(t) R_{p}(t) + 4c_{2}(t) \left(\lambda_{11}^{2} \int_{[0,1]} E\left[\left\|z_{q}(t)\right\|^{2}\right] dq + 2\lambda_{12}^{2} + \lambda_{11}^{2} E\left[\left\|z_{p}(t)\right\|^{2}\right] \right)$$

$$\leq c_{2}(t) \left(R_{p}(t) + 8\lambda_{11}^{2}L + 8\lambda_{12}^{2}\right).$$
(B.12)

For the fourth term on the r.h.s. of (B.10), by C_r inequality, Hölder inequality, Jensen inequality, Assumption 2.2 and (B.9), we have

$$2c_{3}(t)\left(E\left[z_{p}(t)\right] - \int_{\left[0,1\right]} E\left[z_{q}(t)\right] dq\right)^{\mathsf{T}} \left(\int_{\left[0,1\right]} E\left[g(q,z_{q}(t))\right] dq - E\left[g(p,z_{p}(t))\right]\right)$$

$$\leq 2c_{3}(t)\left\|E\left[z_{p}(t)\right] - \int_{\left[0,1\right]} E\left[z_{q}(t)\right] dq\right\|\left\|\int_{\left[0,1\right]} E\left[g(q,z_{q}(t))\right] dq - E\left[g(p,z_{p}(t))\right]\right\|\right\|$$

$$\leq 2\sqrt{2}c_{3}(t)R_{p}^{\frac{1}{2}}(t)\left(\left\|\int_{\left[0,1\right]} E\left[g(q,z_{q}(t))\right] dq\right\|^{2} + \left\|E\left[g(p,z_{p}(t))\right]\right\|^{2}\right)^{\frac{1}{2}}$$

$$\leq 2\sqrt{2}c_{3}(t)R_{p}^{\frac{1}{2}}(t)\left(\int_{\left[0,1\right]} E\left[g(q,z_{q}(t))\right]\right\|^{2} dq + E\left[\left\|g(p,z_{p}(t))\right\|^{2}\right]\right)^{\frac{1}{2}}$$

$$\leq 2\sqrt{2}c_{3}(t)R_{p}^{\frac{1}{2}}(t)\left(\sup_{q\in\left[0,1\right]} E\left[\left\|g(q,z_{q}(t))\right\|^{2}\right] + E\left[\left\|g(p,z_{p}(t))\right\|^{2}\right]\right)^{\frac{1}{2}}$$

$$\leq 4\sqrt{2}c_{3}(t)R_{p}^{\frac{1}{2}}(t)\left(\lambda_{11}^{2}\sup_{q\in\left[0,1\right]} E\left[\left\|z_{q}(t)\right\|^{2}\right] + \lambda_{12}^{2}\right)^{\frac{1}{2}}$$

$$\leq 4\sqrt{2}\left(\lambda_{11}^{2}L + \lambda_{12}^{2}\right)^{\frac{1}{2}}c_{3}(t)R_{p}^{\frac{1}{2}}(t).$$
(B.13)

By (B.9), Jensen inequality and Hölder inequality, we have $R_p(t) \leq 4L$, which gives $R_p^{\frac{1}{2}}(t) \leq 2L^{\frac{1}{2}}, \forall t > 0, p \in [0,1]$. Then, by (B.10)-(B.13), we have

$$\begin{split} &\frac{dR_{p}(t)}{dt} \\ \leqslant &-2\inf_{p\in[0,1]}\int_{[0,1]}A(p,q)dqc_{1}(t)R_{p}(t)+4L^{\frac{1}{2}}\varepsilon^{\frac{1}{2}}(4L+1)^{\frac{1}{2}}c_{1}(t)+4\left(L+2\lambda_{11}^{2}L+2\lambda_{12}^{2}\right)c_{2}(t) \\ &+8\sqrt{2}L^{\frac{1}{2}}\left(\lambda_{11}^{2}L+\lambda_{12}^{2}\right)^{\frac{1}{2}}c_{3}(t), \ \forall \ t \geqslant T_{\varepsilon}. \end{split}$$

This together with the comparison theorem ([32]) gives

$$\sup_{p \in [0,1]} R_p(t)$$

$$\leq \Psi_1(T_{\varepsilon},t) \sup_{p \in [0,1]} R_p(T_{\varepsilon}) + \int_{T_{\varepsilon}}^t 4L^{\frac{1}{2}} \varepsilon^{\frac{1}{2}} (4L+1)^{\frac{1}{2}} c_1(s) \Psi_1(s,t) ds + \int_{T_{\varepsilon}}^t \left(4\left(L+2\lambda_{11}^2L+2\lambda_{12}^2\right) c_2(s) + 8\sqrt{2}L^{\frac{1}{2}} \left(\lambda_{11}^2L+\lambda_{12}^2\right)^{\frac{1}{2}} c_3(s) \right) \Psi_1(s,t) ds, \qquad (B.14)$$

where $\psi_1(s,t) = e^{-\int_s^t 2 \inf_{p \in [0,1]} \int_{[0,1]} A(p,q) dq c_1(s') ds'}$. By Assumption 2.1, Assumption 2.4 and L'Hospital's rule, we have

$$\lim_{t \to \infty} \int_0^t c_1(s) \psi_1(s,t) ds = \frac{1}{2} \left(\inf_{p \in [0,1]} \int_{[0,1]} A(p,q) dq \right)^{-1}$$

Then, we know that there exists $K_3 \ge 0$, such that $\sup_{t\ge 0} \int_0^t \psi_1(s,t)c_1(s)ds \le K_3$. Therefore, for the second term on the r.h.s. of (B.14), we have

$$4L^{\frac{1}{2}}(4L+1)^{\frac{1}{2}}\varepsilon^{\frac{1}{2}}\int_{T_{\varepsilon}}^{t}c_{1}(s)\psi_{1}(s,t)ds \leq 4L^{\frac{1}{2}}\varepsilon^{\frac{1}{2}}(4L+1)^{\frac{1}{2}}K_{3}.$$

By the arbitrariness of ε , there exists $\widetilde{\varepsilon} > 0$, such that $4L^{\frac{1}{2}}\widetilde{\varepsilon}^{\frac{1}{2}}(4L+1)^{\frac{1}{2}}K_3 < \frac{\delta}{3}$. By $\lim_{t\to\infty} \int_{[0,1]} R_p(t) dp = 0$, we know that there exists $T_{\widetilde{\varepsilon}} > 0$, such that if $t \ge T_{\widetilde{\varepsilon}}$, then $\int_{[0,1]} R_p(t) dp < \widetilde{\varepsilon}^2$. Then, for any $t \ge T_{\widetilde{\varepsilon}}$, (B.14) can be written as

$$\sup_{p \in [0,1]} R_p(t) \leqslant \psi_1(T_{\tilde{\varepsilon}}, t) \sup_{p \in [0,1]} R_p(T_{\varepsilon}) + \frac{\delta}{3} + \int_{T_{\tilde{\varepsilon}}}^t \psi_1(s, t) \Big(4 \Big(L + 2\lambda_{12}^2 + 2\lambda_{11}^2 L \Big) c_2(s) + 8\sqrt{2}L^{\frac{1}{2}} \Big(\lambda_{11}^2 L + \lambda_{12}^2 \Big)^{\frac{1}{2}} c_3(s) \Big) ds.$$
(B.15)

By Assumption 2.1 (i), Assumption 2.4 and $\sup_{p \in [0,1], t \ge 0} R_p^{\frac{1}{2}}(t) \le 2L^{\frac{1}{2}}$, we have $\lim_{t \to \infty} \psi_1(T_{\tilde{\varepsilon}}, t)$ $\sup_{p \in [0,1]} R_p(T_{\tilde{\varepsilon}}) = 0$. Therefore, for the first term on the r.h.s. of (B.15), there exists $T_1 > 0$, such that if $t > T_1$, then

$$\Psi_1(T_{\widetilde{\varepsilon}},t) \sup_{p \in [0,1]} R_p(T_{\widetilde{\varepsilon}}) < \frac{\delta}{3}.$$
(B.16)

By Assumption 2.1 (i), Assumption 2.4 and L'Hospital's rule, we have $\lim_{t\to\infty} \int_{T_{\tilde{\varepsilon}}}^{t} (4(L+2\lambda_{11}^2L+2\lambda_{12}^2)c_2(s)+8\sqrt{2}L^{\frac{1}{2}}(\lambda_{11}^2L+\lambda_{12}^2)^{\frac{1}{2}}c_3(s))\psi_1(s,t)ds = 0$. Therefore, for the third term on the r.h.s. of (B.15), there exists $T_{11} > 0$ such that if $t > T_{11}$, then

$$\int_{T_{\tilde{\varepsilon}}}^{t} \left(4 \left(L + 2\lambda_{11}^{2}L + 2\lambda_{12}^{2} \right) c_{2}(s) + 8\sqrt{2}L^{\frac{1}{2}} \left(\lambda_{11}^{2}L + \lambda_{12}^{2} \right)^{\frac{1}{2}} c_{3}(s) \right) \psi_{1}(s,t) ds < \frac{\delta}{3}.$$

Therefore, for any $\delta > 0$, taking $T = \max\{T_1, T_{11}\}$ and by (B.15)-(B.16) and the above inequality, we know that, if $t \ge T$, then $\sup_{p \in [0,1]} R_p(t) < \delta$, that is, (9) holds.

APPENDIX C

Proof of Lemma 3.1: By Theorem 2.1, it's sufficient to prove $\lim_{t\to\infty} \int_{[0,1]} \|\int_{[0,1]} E[x_q(t)] dq - E[x_p(t)]\|^2 dp = 0$ and (11) for (13). By $\mu_{t,q} = \mathcal{L}(x_q(t))$ in (10), the system (10) can be written as

$$dx_{p}(t) = \alpha_{1}(t) \int_{[0,1]} A(p,q) \left(E[x_{q}(t)] - x_{p}(t) \right) dq dt - \alpha_{2}(t) \nabla_{x} V(p, x_{p}(t)) dt - \alpha_{2}(t) \Sigma_{1} dw_{p}(t).$$
(C.1)

By Assumption 3.2 and Theorem 2.3.1 in [31], we have $E\left[\int_0^t \alpha_2(s)\Sigma_1 dw_p(s)\right] = 0$. This together with (C.1) gives

$$dE[x_p(t)] = \alpha_1(t) \int_{[0,1]} A(p,q) \left(E[x_q(t)] - E[x_p(t)] \right) dq dt - \alpha_2(t) E\left[\nabla_x V(p, x_p(t)) \right] dt.$$
(C.2)

Denote $Y(t) = \int_{[0,1]} E[||x_p(t)||^2] dp$ and $R(t) = \int_{[0,1]} ||E[x_p(t)] - \int_{[0,1]} E[x_q(t)] dq ||^2 dp$. By Assumption 3.2, Corollary 3.1 in [35] and Theorem 2.3.1 in [31], we have $E[\int_0^t 2\alpha_2(s)x_p^{\mathsf{T}}(s)\Sigma_1 dw_p(s)] = 0$. Then, by Itô formula and (C.1), we have

$$dY(t) = 2\alpha_{1}(t) \int_{[0,1]\times[0,1]} A(p,q) \left(E[x_{p}^{\mathsf{T}}(t)]E[x_{q}(t)] - E[\|x_{p}(t)\|^{2}] \right) dqdpdt + \alpha_{2}^{2}(t) \operatorname{Tr}(\Sigma_{1}^{\mathsf{T}}\Sigma_{1}) dt - 2\alpha_{2}(t) \int_{[0,1]} E\left[x_{p}^{\mathsf{T}}(t)\nabla_{x}V(p,x_{p}(t))\right] dpdt.$$
(C.3)

By Assumption 2.1 (iii), we know $(x_p(t) - x_p(0))^{\mathsf{T}}(\nabla_x V(p, x_p(t)) - \nabla_x V(p, x_p(0))) \ge \kappa_2 ||x_p(t) - x_p(0)||^2$. Then, by Cauchy-Schwartz inequality and Hölder inequality, we have

$$-2\alpha_{2}(t)E\left[x_{p}^{\mathsf{T}}(t)\nabla_{x}V(p,x_{p}(t))\right]$$

$$\leq 2\alpha_{2}(t)\left(-\kappa_{2}E\left[\|x_{p}(t)\|^{2}\right]+E\left[\|x_{p}(t)\|^{2}\right]^{\frac{1}{2}}E\left[\|\nabla_{x}V(p,x_{p}(0))\|^{2}\right]^{\frac{1}{2}}+E\left[\|x_{p}(0)\|^{2}\right]^{\frac{1}{2}}$$

$$\times E\left[\|\nabla_{x}V(p,x_{p}(t))\|^{2}\right]^{\frac{1}{2}}+E\left[\|x_{p}(0)\|^{2}\right]^{\frac{1}{2}}E\left[\|\nabla_{x}V(p,x_{p}(0))\|^{2}\right]^{\frac{1}{2}}+2\kappa_{2}E\left[\|x_{p}(t)\|^{2}\right]^{\frac{1}{2}}$$

$$\times E\left[\|x_{p}(0)\|^{2}\right]^{\frac{1}{2}}\right).$$

By Assumption 2.1 (ii) and C_r inequality, we have $\|\nabla_x V(p, x_p(t))\|^2 \leq 3(\kappa^2 \|x_p(t)\|^2 + \kappa^2 \|x_p(0)\|^2 + \|\nabla_x V(p, x_p(0))\|^2)$. This together with the above inequality, C_r inequality, Assumption 2.1 and Assumption 3.1 gives

$$-2\alpha_{2}(t)E\left[x_{p}^{\mathsf{T}}(t)\nabla_{x}V(p,x_{p}(t))\right]$$

$$\leq -2\kappa_{2}\alpha_{2}(t)E\left[\|x_{p}(t)\|^{2}\right]+2\alpha_{2}(t)E\left[\|x_{p}(t)\|^{2}\right]^{\frac{1}{2}}E\left[\|\nabla_{x}V(p,x_{p}(0))\|^{2}\right]^{\frac{1}{2}}$$

$$+2\alpha_{2}(t)E\left[3\kappa^{2}\|x_{p}(t)\|^{2}+3\|\nabla_{x}V(p,x_{p}(0))\|^{2}+3\kappa^{2}\|x_{p}(0)\|^{2}\right]^{\frac{1}{2}}E[\|x_{p}(0)\|^{2}]^{\frac{1}{2}}$$

$$+2\alpha_{2}(t)E[\|x_{p}(0)\|^{2}]^{\frac{1}{2}}E[\|\nabla_{x}V(p,x_{p}(0))\|^{2}]^{\frac{1}{2}}+4\alpha_{2}(t)\kappa_{2}E[\|x_{p}(t)\|^{2}]^{\frac{1}{2}}E[\|x_{p}(0)\|^{2}]^{\frac{1}{2}}$$

$$\leq -2\kappa_{2}\alpha_{2}(t)E[\|x_{p}(t)\|^{2}]+\left(2\left(2\sigma_{\nu}^{2}\zeta_{2}+2C_{\nu}^{2}\right)^{\frac{1}{2}}+2\sqrt{3}\kappa\zeta_{2}^{\frac{1}{2}}+4\kappa_{2}\zeta_{2}^{\frac{1}{2}}\right)\alpha_{2}(t)E[\|x_{p}(t)\|^{2}]^{\frac{1}{2}}$$

$$+\left(2(\sqrt{3}+1)\zeta_{2}^{\frac{1}{2}}\left(2\sigma_{\nu}^{2}\zeta_{2}+2C_{\nu}^{2}\right)^{\frac{1}{2}}+2\sqrt{3}\zeta_{2}\kappa\right)\alpha_{2}(t).$$
(C.4)

By Assumption 2.1 (i) and (A.2), we have

$$2\alpha_{1}(t)\int_{[0,1]\times[0,1]}A(p,q)\left(E[x_{p}^{\mathsf{T}}(t)]E[x_{q}(t)]-E\left[\|x_{p}(t)\|^{2}\right]\right)dqdp \leq -2\alpha_{1}(t)\lambda_{2}(\mathbb{L}_{A})R(t) \leq 0.$$
(C.5)

This together with (C.3)-(C.4) gives

$$\frac{dY(t)}{dt} \leqslant -l_1(t)Y(t) + l_2(t)Y^{\frac{1}{2}}(t) + l_3(t),$$

where $l_1(t) = 2\kappa_2\alpha_2(t)$, $l_2(t) = 2((2\sigma_v^2\zeta_2 + 2C_v)^{\frac{1}{2}} + \sqrt{3}\kappa\zeta_2^{\frac{1}{2}} + 2\kappa_2\zeta_2^{\frac{1}{2}})\alpha_2(t)$ and $l_3(t) = 2((\sqrt{3} + 1) \times \zeta_2^{\frac{1}{2}}(2\sigma_v^2\zeta_2 + 2C_v^2)^{\frac{1}{2}} + \sqrt{3}\zeta_2\kappa)\alpha_2(t) + \alpha_2^2(t)\operatorname{Tr}(\Sigma_1^{\mathsf{T}}\Sigma_1)$. By Assumption 2.1 (iii), Assumption 3.2 and Lemma 2.2, we have

$$Y(t) \leq \max\left\{Y(0), \left(\sup_{0 \leq s \leq t} \frac{l_2(s)}{2l_1(s)} + \left(\sup_{0 \leq s \leq t} \frac{1}{4} \frac{l_2^2(s)}{l_1^2(s)} + \sup_{0 \leq s \leq t} \frac{l_3(s)}{l_1(s)}\right)^{\frac{1}{2}}\right)^2\right\}.$$
 (C.6)

By Assumption 3.1, we get $Y(0) = \int_{[0,1]} E[\|x_p(0)\|^2] \leq \sup_{p \in [0,1]} E[\|x_p(0)\|^2] \leq \zeta_2$. By Assumption 3.2, we have

$$\lim_{t \to \infty} \frac{l_2(t)}{2l_1(t)} = \frac{1}{2\kappa_2} \left(\left(2\sigma_v^2 \zeta_2 + C_v^2 \right)^{\frac{1}{2}} + \left(\sqrt{3}\kappa + 2\kappa_2 \right) \zeta_2^{\frac{1}{2}} \right),$$
$$\lim_{t \to \infty} \frac{1}{4} \frac{l_2^2(t)}{l_1^2(t)} = \frac{1}{4\kappa_2^2} \left(\left(2\sigma_v^2 \zeta_2 + 2C_v^2 \right)^{\frac{1}{2}} + \left(\sqrt{3}\kappa + 2\kappa_2 \right) \zeta_2^{\frac{1}{2}} \right)^2$$

and

$$\lim_{t \to \infty} \frac{l_3(t)}{l_1(t)} = \frac{\zeta_2^{\frac{1}{2}}}{\kappa_2} \left((\sqrt{3}+1) \left(2\sigma_v^2 \zeta_2 + 2C_v^2 \right)^{\frac{1}{2}} + \sqrt{3}\zeta_2^{\frac{1}{2}} \right).$$

By the above three equalities, there exist non-negative constants M_1 , M_2 and M_3 , such that $\sup_{t\geq 0} \frac{l_2(t)}{2l_1(t)} \leq M_1$, $\sup_{t\geq 0} \frac{1}{4} \frac{l_2^2(t)}{l_1^2(t)} \leq M_2$ and $\sup_{t\geq 0} \frac{l_3(t)}{l_1(t)} \leq M_3$. Denote $K = \max \{\zeta_2, (\sqrt{M_2 + M_3} + M_1)^2\}$. Then, by (C.6), we have $\sup_{t\geq 0} Y(t) \leq K$. Then, similar to the proof of (B.9) in Theorem 2.1 and by Assumption 2.1 (i), we have (11).

Combining (C.2) and the symmetry of the graphon A gives

$$\frac{d\left(\int_{[0,1]} E[x_p(t)]dp\right)}{dt} = -\alpha_2(t) \int_{[0,1]} E\left[\nabla_x V\left(p, x_p(t)\right)\right] dp.$$

This together with (C.2) gives

$$\frac{dR(t)}{dt} = 2\alpha_{1}(t) \int_{[0,1]} E[x_{p}^{\mathsf{T}}(t)] \left(\int_{[0,1]} A(p,q) \left(E[x_{q}(t)] - E[x_{p}(t)] \right) dq \right) dp + 2\alpha_{2}(t) \\
\times \int_{[0,1]} \left(E[x_{p}(t)] - \int_{[0,1]} E[x_{q}(t)] dq \right)^{\mathsf{T}} \left(\int_{[0,1]} E[\nabla_{x}V(q,x_{q}(t))] dq - E[\nabla_{x}V(p,x_{p}(t))] \right) dp.$$
(C.7)

Combining Assumption 2.1 (ii), C_r inequality, Jensen inequality, Hölder inequality and (11) gives

$$\begin{split} &2\alpha_{2}(t)\int_{[0,1]}\left(E[x_{p}(t)]-\int_{[0,1]}E[x_{q}(t)]dq\right)^{\mathsf{T}}\left(\int_{[0,1]}E\left[\nabla_{x}V(q,x_{q}(t))\right]dq\\ &-E\left[\nabla_{x}V(p,x_{p}(t))\right]\right)dp\\ &\leqslant &2\alpha_{2}(t)\int_{[0,1]}\left\|E[x_{p}(t)]-\int_{[0,1]}E[x_{q}(t)]dq\right\|\\ &\times\left\|\int_{[0,1]}E\left[\nabla_{x}V(q,x_{q}(t))\right]dq-E\left[\nabla_{x}V(p,x_{p}(t))\right]\right\|dp\\ &\leqslant &4\sqrt{2}\alpha_{2}(t)R^{\frac{1}{2}}(t)\left(\int_{[0,1]}\left\|E\left[\nabla_{x}V(p,x_{p}(t))\right]\right\|^{2}dp\right)^{\frac{1}{2}}\\ &\leqslant &4\sqrt{2}\alpha_{2}(t)R^{\frac{1}{2}}(t)\left(\int_{[0,1]}E\left[\|\nabla_{x}V(p,x_{p}(t))\|^{2}\right]dp\right)^{\frac{1}{2}}\\ &\leqslant &8\alpha_{2}(t)R^{\frac{1}{2}}(t)\left(\sigma_{v}^{2}\int_{[0,1]}E\left[\|x_{p}(t)\|^{2}\right]dp+C_{v}^{2}\right)^{\frac{1}{2}}\\ &\leqslant &8\alpha_{2}(t)R^{\frac{1}{2}}(t)\left(\sigma_{v}K_{0}^{\frac{1}{2}}+C_{v}\right). \end{split}$$

Then, by (C.5), (C.7) and the above inequality, we have

$$\frac{dR(t)}{dt} \leqslant -2\alpha_1(t)\lambda_2(\mathbb{L}_A)R(t) + 8\alpha_2(t)R^{\frac{1}{2}}(t)(\sigma_v K_0^{\frac{1}{2}} + C_v).$$
(C.8)

By (11) and Jensen inequality, we get $R(t) \leq \int_{[0,1]} ||E[x_p(t)]||^2 dp \leq \int_{[0,1]} E[||x_p(t)||^2] dp \leq K_0$, and then $R^{\frac{1}{2}}(t) \leq K_0^{\frac{1}{2}}$. This together with (C.8) gives

$$\frac{dR(t)}{dt} \leqslant -2\alpha_1(t)\lambda_2(\mathbb{L}_A)R(t) + 8\alpha_2(t)(\sigma_v K_0 + C_v K_0^{\frac{1}{2}}),$$

which together with the comparison theorem ([32]) gives

$$R(t) \leqslant e^{-2\lambda_{2}(\mathbb{L}_{A})\int_{0}^{t}\alpha_{1}(s)ds}R(0) + \int_{0}^{t} \left(8\left(\sigma_{\nu}K_{0} + C_{\nu}K_{0}^{\frac{1}{2}}\right)\alpha_{2}(s)e^{-2\lambda_{2}(\mathbb{L}_{A})\int_{s}^{t}\alpha_{1}(s')ds'}\right)ds.$$
(C.9)

This together with $R(0) \leq \zeta_2 < \infty$ gives (12). Then, by (C.9), Assumption 3.2 and L'Hospital's rule, we have

$$\lim_{t\to\infty}\left[e^{-2\lambda_2(\mathbb{L}_A)\int_0^t\alpha_1(s)ds}R(0)+\int_0^t\left(\left(8\sigma_v K_0+8C_v K_0^{\frac{1}{2}}\right)\alpha_2(s)e^{-2\lambda_2(\mathbb{L}_A)\int_s^t\alpha_1(s')ds'}\right)ds\right]=0.$$

This together with (11) and Theorem 2.1 leads to (13).

To prove Lemma 3.2, we need the following lemma, the proof of which is directly from Lemma 2.1.

Lemma C.1: For the problem (1) and the algorithm (10), if Assumption 2.1 and Assumptions 3.1-3.2 hold, then $\lim_{t\to\infty} \sup_{p\in[0,1]} E[\|x_p(t) - E[x_p(t)]\|^2] = 0.$

Proof of Lemma 3.2: Denote $L(t) = \|\int_{[0,1]} E[x_p(t)] dp - x^*\|^2$, $L_1(t) = \int_{[0,1]} E[x_p(t)] dp$, $L_2(t) = \sup_{p \in [0,1]} E[\|x_p(t) - E[x_p(t)]\|^2]$ and $L_3(t) = \sup_{p \in [0,1]} \|E[x_p(t)] - L_1(t)\|^2$. By the symmetry of the graphon A and (C.2) in Lemma 2.2, we have

$$\begin{aligned} \frac{dL(t)}{dt} =& 2\alpha_2(t) \left(x^* - L_1(t) \right)^\mathsf{T} \left(\int_{[0,1]} \left(E\left[\nabla_x V(p, x_p(t)) \right] - \nabla_x V(p, L_1(t)) \right. \\ &+ \left. \nabla_x V\left(p, L_1(t)\right) - \nabla_x V(p, x^*) \right) dp \right). \end{aligned}$$

This together with Assumption 2.1, Hölder inequality and Jensen inequality gives

$$\begin{aligned} \frac{dL(t)}{dt} \\ \leqslant 2\alpha_{2}(t) \left(L^{\frac{1}{2}}(t) \right\| \int_{[0,1]} \left(E[\nabla_{x}V(p,x_{p}(t))] - \nabla_{x}V(p,L_{1}(t))\right) dp \left\| - \kappa_{2}L(t) \right) \\ \leqslant 2\alpha_{2}(t) \left(- \kappa_{2}L(t) + \left(\int_{[0,1]} \left\| E[\nabla_{x}V(p,x_{p}(t))] - \nabla_{x}V(p,L_{1}(t))\right\|^{2} dp \right)^{\frac{1}{2}} L^{\frac{1}{2}}(t) \right) \\ \leqslant 2\alpha_{2}(t) \left(\sqrt{2}L^{\frac{1}{2}}(t) \left(\int_{[0,1]} \left\| E[\nabla_{x}V(p,x_{p}(t))] - \nabla_{x}V(p,E[x_{p}(t)])\right\|^{2} dp \right)^{\frac{1}{2}} \\ - \kappa_{2}L(t) + \sqrt{2}L^{\frac{1}{2}}(t) \left(\int_{[0,1]} \left\| \nabla_{x}V(p,E[x_{p}(t)]) - \nabla_{x}V(p,L_{1}(t))\right\|^{2} dp \right)^{\frac{1}{2}} \right) \\ \leqslant 2\alpha_{2}(t) \left(- \kappa_{2}L(t) + \sqrt{2}\kappa L^{\frac{1}{2}}(t) \left(\left(\int_{[0,1]} E\left[\left\| x_{p}(t) - E[x_{p}(t)] \right\|^{2} \right] dp \right)^{\frac{1}{2}} \\ + \left(\int_{[0,1]} \left\| E[x_{p}(t)] - L_{1}(t)\right\|^{2} dp \right)^{\frac{1}{2}} \right) \right) \\ \leqslant 2\alpha_{2}(t) \left(- \kappa_{2}L(t) + \sqrt{2}\kappa L^{\frac{1}{2}}(t) \left(L^{\frac{1}{2}}(t) + L^{\frac{1}{3}}(t) \right) \right). \end{aligned}$$
(C.10)

By Lemma 3.1, C_r inequality, Hölder inequality and Jensen inequality, we have $L(t) \leq 2||x^*||^2 + 2\int_{[0,1]} E[||x_p(t)||^2] dp \leq 2K + 2||x^*||^2$ and $L^{\frac{1}{2}}(t) \leq (2K_0 + 2||x^*||^2)^{\frac{1}{2}}$, which together with (C.10) gives

$$\frac{dL(t)}{dt} \leqslant -2\kappa_2\alpha_2(t)L(t) + 4\kappa\alpha_2(t)\left(K_0 + \|x^*\|^2\right)^{\frac{1}{2}}\left(L_2^{\frac{1}{2}}(t) + L_3^{\frac{1}{2}}(t)\right).$$

This together with the comparison theorem ([32]) leads to

$$L(t) \leq \Psi_{2}(0,t)L(0) + 4\kappa \left(K_{0} + \|x^{*}\|^{2}\right)^{\frac{1}{2}} \int_{0}^{t} \alpha_{2}(s)L_{2}^{\frac{1}{2}}(s)\Psi_{2}(s,t)ds + 4\kappa \left(K_{0} + \|x^{*}\|^{2}\right)^{\frac{1}{2}} \int_{0}^{t} \alpha_{2}(s)L_{3}^{\frac{1}{2}}(s)\Psi_{2}(s,t)ds,$$
(C.11)

where $\psi_2(s,t) = e^{-2\kappa_2 \int_s^t \alpha_2(s')ds'}$. For the first term on the r.h.s. of (C.11), by Assumption 3.1, C_r inequality and Hölder inequality, we have $L(0) = \left\| \int_{[0,1]} E[x_p(0)] dp - x^* \right\|^2 \leq 2 \|x^*\|^2 + 2 \sup_{p \in [0,1]} \|E[x_p(0)]\|^2 \leq 2\zeta_2 + 2 \|x^*\|^2$, which together with Assumption 2.1 (iii) and Assumption 3.2 gives

$$\lim_{t \to \infty} \psi_2(0, t) L(0) = 0.$$
 (C.12)

For the second term on the r.h.s. of (C.11), by Assumption 2.1 (iii), Assumption 3.2, Lemma C.1 and L'Hospital's rule, we have

$$\lim_{t \to \infty} 2\sqrt{2\kappa} \left(2K_0 + 2\|x^*\|^2 \right)^{\frac{1}{2}} \int_0^t \alpha_2(s) \psi_2(s,t) L_2^{\frac{1}{2}}(s) ds = 0.$$
(C.13)

For the third term on the r.h.s. of (C.11), by Assumption 2.1 (iii), Assumption 3.2, Lemma 3.1 and L'Hospital's rule, we have

$$\lim_{t\to\infty} 2\sqrt{2}\kappa (2K_0 + 2\|x^*\|^2)^{\frac{1}{2}} \int_0^t \alpha_2(s) \psi_2(s,t) L_3^{\frac{1}{2}}(s) ds = 0.$$

This together with (C.11)-(C.13) gives $\lim_{t\to\infty} L(t) = 0$.

APPENDIX D

Proof of Lemma 4.1: By $\lim_{t\to\infty} Y_3(t) = 0$, we know that there exists $N_1 \ge 0$, such that

$$\sup_{t \ge 0} Y_3(t) \leqslant N_1. \tag{D.1}$$

Then, by (18), we have

$$\frac{dY_2(t)}{dt} \leqslant -b_1(t)Y_2(t) + b_2(t)Y_2^{\frac{1}{2}}(t)\left(Y_1^{\frac{1}{2}}(t) + N_1^{\frac{1}{2}}\right).$$
(D.2)

By $\limsup_{t\to\infty} \frac{b_2(t)}{b_1(t)} < \infty$, we know that there exists $N_2 \ge 0$, such that $\sup_{t\ge \frac{b_2(t)}{b_1(t)}} \le N_2$. Then, by $b_1(t) > 0$, (D.2), Lemma 2.2 and C_r inequality, we have

$$Y_2(t) \leqslant \left(\frac{b_2(t)}{b_1(t)} \left(Y_1^{\frac{1}{2}}(t) + N_2^{\frac{1}{2}}\right)\right)^2 \leqslant 2N_2^2(Y_1(t) + N_1), \forall t \ge 0.$$

This together with (17) gives

$$\frac{dY_1(t)}{dt} \leqslant \left(-a_1(t) + a_2(t) + 2N_2^2 a_3(t)\right) Y_1(t) + 2N_1 N_2^2 a_3(t) + a_4(t).$$
(D.3)

By $a_1(t) > 0$, $a_2(t), a_3(t) \ge 0$, $\lim_{t\to\infty} \frac{a_2(t)}{a_1(t)} = 0$ and $\lim_{t\to\infty} \frac{a_3(t)}{a_1(t)} = 0$, we know that there exists $T \ge 0$, such that if $t \ge T$, then $-a_1(t) + a_2(t) + 2N_2^2a_3(t) < 0$. Then, by (D.3) and comparison theorem ([32]), we have

$$Y_1(t) \leqslant \int_T^t \psi_3(s,t) (2N_1 N_2^2 a_3(s) + a_4(s)) ds + \psi_3(T,t) Y_1(T), \forall t \ge T,$$
(D.4)

where $\psi_3(s,t) = e^{\int_s^t (-a_1(s') + a_2(s') + 2N_2^2 a_3(s'))ds'}$. By $\int_0^\infty a_1(t)dt = \infty$, $\lim_{t \to \infty} \frac{a_2(t)}{a_1(t)} = 0$ and $\lim_{t \to \infty} \frac{a_3(t)}{a_1(t)} = 0$, we have

$$\int_{T}^{\infty} (-a_1(s) + a_2(s) + 2N_2^2 a_3(s)) ds = -\infty.$$
 (D.5)

For the first term on the r.h.s. of (D.4), by (D.5), $\lim_{t\to\infty} \frac{a_2(t)}{a_1(t)} = 0$, $\lim_{t\to\infty} \frac{a_3(t)}{a_1(t)} = 0$, $\lim_{t\to\infty} \frac{a_4(t)}{a_1(t)} = 0$ and L'Hospital's rule, we have

$$\lim_{t \to \infty} \int_T^t \psi_3(s,t) (2N_1 N_2^2 a_3(s) + a_4(s)) ds = 0.$$
 (D.6)

Noting that $Y_1(t)$ is continuous w.r.t. t, then we have $Y_1(T) < \infty$. Then, for the second term on the r.h.s. of (D.4), by (D.5), we have

$$\lim_{t\to\infty}\psi_3(T,t)Y_1(T)=0.$$

By the above equality, (D.4) and (D.6), we have (19). By (18) and comparison theorem ([32]), we have

$$Y_2(t) \leqslant e^{-\int_0^t b_1(s)ds} Y_2(0) + \int_0^t e^{-\int_s^t b_1(s')ds'} b_2(s) \left(Y^{\frac{1}{2}}(s) + Y_3(s)\right) ds.$$
(D.7)

For the first term on the r.h.s. of (D.7), by $\int_0^{\infty} b_1(t) dt = \infty$, we have

$$\lim_{t \to \infty} e^{-\int_0^t b_1(s) ds} Y_2(0) = 0.$$
 (D.8)

By (19), we have $\lim_{t\to\infty} Y_1^{\frac{1}{2}}(t) = 0$. Then, for the second term on the r.h.s. of (D.7), by $\lim_{t\to\infty} Y_3(t) = 0$, $\limsup_{t\to\infty} \frac{b_2(t)}{b_1(t)} < \infty$ and L'Hospital's rule, we have

$$\lim_{t\to\infty}\int_0^t e^{-\int_s^t b_1(s')ds'}b_2(s)\left(Y^{\frac{1}{2}}(s)+Y_3(s)\right)ds=0.$$

By the above equality, (D.7) and (D.8), we have (20).

Before we prove Lemma 4.2, we need the following lemma whose proof is directly from Lemma 2.1.

Lemma D.1: For the problem (1) and the stochastic gradient tracking algorithm (16), if Assumption 2.1 and Assumptions 4.1-4.3 hold, then

$$\lim_{t \to \infty} \sup_{p \in [0,1]} E\left[\|z_p(t) - E[z_p(t)]\|^2 \right] = 0,$$

$$\lim_{t \to \infty} \sup_{p \in [0,1]} E\left[\|\widetilde{y}_p(t) - E[\widetilde{y}_p(t)]\|^2 \right] = 0.$$

Proof of Lemma 4.2: Noting that $\tilde{\mu}_{t,q}$ and $\tilde{v}_{t,q}$ are the distributions of $z_p(t)$ and $\tilde{y}_p(t)$, respectively and by Assumption 4.3, we have

$$\frac{dE[z_{p}(t)]}{dt} = \beta_{3}(t) \int_{[0,1]} A(p,q) \left(E[z_{q}(t)] - E[z_{p}(t)] \right) dq - \beta_{1}(t) \beta_{3}(t) \int_{[0,1]} A(p,q) \left(E[\widetilde{y}_{q}(t)] - E[\widetilde{y}_{p}(t)] + \beta_{2}(t) \left(E[\nabla_{x}V(q,z_{q}(t))] - E[\nabla_{x}V(p,z_{p}(t))] \right) \right) dq - \beta_{1}(t) \left(E[\widetilde{y}_{p}(t)] + \beta_{2}(t) E[\nabla_{x}V(p,z_{p}(t))] \right) \tag{D.9}$$

and

$$\frac{dE\left[\widetilde{y}_{p}(t)\right]}{dt} = \beta_{3}(t) \int_{[0,1]} A(p,q) \left(E\left[\widetilde{y}_{q}(t)\right] - E\left[\widetilde{y}_{p}(t)\right] \right) dq + \beta_{2}(t) \beta_{3}(t) \int_{[0,1]} A(p,q) \left(E\left[\nabla_{x}V(q,z_{q}(t))\right] - E\left[\nabla_{x}V(p,z_{p}(t))\right] \right) dq.$$
(D.10)

By (3) and Assumption 4.1, we have $E[\tilde{y}_p(0)] = E[y_p(0)] - \beta_2(0)E[\nabla_x V(p, z_p(0))] = 0$, which together with the above equality and the symmetry of the graphon A gives

$$\int_{[0,1]} E[\tilde{y}_p(t)] dp = 0.$$
 (D.11)

Then, by (D.9), we have

$$\frac{d\mathscr{R}(t)}{dt} = -2\beta_{1}(t)\int_{[0,1]} \left(E[z_{p}(t)] - R_{1}(t)\right)^{\mathsf{T}}E[\widetilde{y}_{p}(t)]dp - 2\beta_{1}(t)\beta_{2}(t)\int_{[0,1]} \left(E[z_{p}(t)] - R_{1}(t)\right)^{\mathsf{T}} \times E[\nabla_{x}V(p,z_{p}(t))]dp + 2\beta_{3}(t)\int_{[0,1]}E[z_{p}^{\mathsf{T}}(t)]\left(\int_{[0,1]}A(p,q)(E[z_{q}(t)] - E[z_{p}(t)])dq\right)dp$$

$$-2\beta_{1}(t)\beta_{3}(t)\int_{[0,1]}E[z_{p}^{\mathsf{T}}(t)]\left(\int_{[0,1]}A(p,q)\left(E\left[\widetilde{y}_{q}(t)\right]-E\left[\widetilde{y}_{p}(t)\right]\right)dq\right)dp$$

$$-2\beta_{1}(t)\beta_{2}(t)\beta_{3}(t)\int_{[0,1]}E[z_{p}^{\mathsf{T}}(t)]\left(\int_{[0,1]}A(p,q)\left(E\left[\nabla_{x}V\left(q,z_{q}(t)\right)\right]\right)dq\right)dp.$$

$$-E[\nabla_{x}V\left(p,z_{p}(t)\right)]dq\right)dp.$$
 (D.12)

For the first term on the r.h.s. of the above equality, by C_r inequality, we have

$$-2\beta_{1}(t)\int_{[0,1]} \left(E[z_{p}(t)]-R_{1}(t)\right)^{\mathsf{T}}E[\widetilde{y}_{p}(t)]dp \leqslant \beta_{1}(t)(\mathscr{R}(t)+\widetilde{\mathscr{R}}(t)).$$
(D.13)

For the second term on the r.h.s. of (D.12), by Assumption 2.1 (ii), C_r inequality and Jensen inequality, we have

$$\begin{aligned} &-2\beta_{1}(t)\beta_{2}(t)\int_{[0,1]}\left(E\left[z_{p}(t)\right]-R_{1}(t)\right)^{\mathsf{T}}E\left[\nabla_{x}V\left(p,z_{p}(t)\right)\right]dp\\ &=-2\beta_{1}(t)\beta_{2}(t)\int_{[0,1]}\left(E\left[z_{p}(t)\right]-R_{1}(t)\right)^{\mathsf{T}}\left(\nabla_{x}V(p,x^{*})-\nabla_{x}V(p,x^{*})+E\left[\nabla_{x}V\left(p,z_{p}(t)\right)\right]\right)dp\\ &\leqslant\beta_{1}(t)\beta_{2}(t)\left(2\mathscr{R}(t)+\int_{[0,1]}\left\|E\left[\nabla_{x}V\left(p,z_{p}(t)\right)\right]-\nabla_{x}V\left(p,x^{*}\right)\right\|^{2}dp+\int_{[0,1]}\left\|\nabla_{x}V(p,x^{*})\right\|^{2}dp\right)\\ &\leqslant\beta_{1}(t)\beta_{2}(t)\left(2\mathscr{R}(t)+2\int_{[0,1]}\left\|E\left[\nabla_{x}V\left(p,z_{p}(t)\right)\right]-\nabla_{x}V\left(p,E\left[z_{p}(t)\right]\right)\right\|^{2}dp\\ &+2\int_{[0,1]}\left\|\nabla_{x}V\left(p,E\left[z_{p}(t)\right]\right)-\nabla_{x}V(p,x^{*})\right\|^{2}dp+\int_{[0,1]}\left\|\nabla_{x}V(p,x^{*})\right\|^{2}dp\right)\\ &\leqslant\beta_{1}(t)\beta_{2}(t)\left(2\mathscr{R}(t)+2\int_{[0,1]}E\left[\left\|\nabla_{x}V\left(p,z_{p}(t)\right)-\nabla_{x}V\left(p,E\left[z_{p}(t)\right]\right)\right\|^{2}\right]dp\\ &+2\kappa^{2}\int_{[0,1]}\left\|E\left[z_{p}(t)\right]-x^{*}\right\|^{2}dp+\int_{[0,1]}\left\|\nabla_{x}V\left(p,x^{*}\right)\right\|^{2}dp\right)\\ &\leqslant\beta_{1}(t)\beta_{2}(t)\left(2\mathscr{R}(t)+2\kappa^{2}B(t)+2\kappa^{2}\int_{[0,1]}\left\|E\left[z_{p}(t)\right]-x^{*}\right\|^{2}dp+\int_{[0,1]}\left\|\nabla_{x}V(p,x^{*})\right\|^{2}dp\right)\\ &\leqslant2\beta_{1}(t)\beta_{2}(t)\left((1+2\kappa^{2})\mathscr{R}(t)+\kappa^{2}(2\mathscr{S}(t)+B(t))+\sigma_{\nu}^{2}\|x^{*}\|^{2}+C_{\nu}^{2}\right).\end{aligned}$$

For the third term on the r.h.s. of (D.12), by (A.2), we have

$$2\beta_{3}(t)\int_{[0,1]} E\left[z_{p}^{\mathsf{T}}(t)\right]\left(\int_{[0,1]} A(p,q)\left(E\left[z_{q}(t)\right]-E\left[z_{p}(t)\right]\right)dq\right)dp \leqslant -2\beta_{3}(t)\lambda_{2}\left(\mathbb{L}_{A}\right)\mathscr{R}(t).$$
(D.15)

For the fourth term on the r.h.s. of (D.12), by C_r inequality, we have

$$-2\beta_{1}(t)\beta_{3}(t)\int_{[0,1]}E\left[z_{p}^{\mathsf{T}}(t)\right]\left(\int_{[0,1]}A(p,q)\left(E\left[\widetilde{y}_{q}(t)\right]-E\left[\widetilde{y}_{p}(t)\right]\right)dq\right)dp$$
$$\leq 2\beta_{1}(t)\beta_{3}(t)\int_{[0,1]\times[0,1]}\left(\left\|E\left[z_{p}(t)\right]\right\|\left\|E\left[\widetilde{y}_{q}(t)\right]\right\|+\left\|E\left[z_{p}(t)\right]\right\|\left\|E\left[\widetilde{y}_{p}(t)\right]\right\|\right)dqdp$$

$$\leq 2\beta_{1}(t)\beta_{3}(t)\int_{[0,1]} \left(\left\| E\left[z_{p}(t)\right] \right\|^{2} + \left\| E\left[\widetilde{y}_{p}(t)\right] \right\|^{2} \right) dp$$

$$\leq 2\beta_{1}(t)\beta_{3}(t) \left(3\mathscr{R}(t) + 3\mathscr{S}(t) + 3\left\|x^{*}\right\|^{2} + \widetilde{\mathscr{R}}(t) \right).$$
(D.16)

For the fifth term on the r.h.s. of (D.12), by Assumption 2.1 (ii), C_r inequality and Jensen inequality, we have

$$-2\beta_{1}(t)\beta_{2}(t)\beta_{3}(t)\int_{[0,1]} E\left[z_{p}^{\mathsf{T}}(t)\right] \left(\int_{[0,1]} A(p,q)\left(E\left[\nabla_{x}V\left(q,z_{q}(t)\right)\right]\right) dq\right) dp$$

$$\leq 2\beta_{1}(t)\beta_{2}(t)\beta_{3}(t)\int_{[0,1]\times[0,1]} \left(\left\|E\left[\nabla_{x}V\left(q,z_{q}(t)\right)\right]\right\|\left\|E\left[z_{p}(t)\right]\right\|$$

$$+\left\|E\left[\nabla_{x}V\left(p,z_{p}(t)\right)\right]\right\|\left\|E\left[z_{p}(t)\right]\right\|\right) dq dp$$

$$\leq 2\beta_{1}(t)\beta_{2}(t)\beta_{3}(t)\left(\int_{[0,1]} \left\|E\left[z_{p}(t)\right]\right\|^{2} dp + \int_{[0,1]} \left\|E\left[\nabla_{x}V\left(p,z_{p}(t)\right)\right]\right\|^{2} dp\right)$$

$$\leq 2\beta_{1}(t)\beta_{2}(t)\beta_{3}(t)\left(2\int_{[0,1]} \left\|\nabla_{x}V\left(p,E\left[z_{p}(t)\right]\right)\right\|^{2} dp + \int_{[0,1]} \left\|E\left[z_{p}(t)\right]\right\|^{2} dp$$

$$+2\int_{[0,1]} E\left[\left\|\nabla_{x}V\left(p,z_{p}(t)\right) - \nabla_{x}V\left(p,E\left[z_{p}(t)\right]\right)\right|^{2}\right] dp\right)$$

$$\leq 2\beta_{1}(t)\beta_{2}(t)\beta_{3}(t)\left(\int_{[0,1]} \left\|E\left[z_{p}(t)\right]\right\|^{2} dp + 2\kappa^{2}B(t) + 2\int_{[0,1]} \left\|\nabla_{x}V\left(p,E\left[z_{p}(t)\right]\right)\right\|^{2} dp\right)$$

$$\leq 2\beta_{1}(t)\beta_{2}(t)\beta_{3}(t)\left(\left(1+4\sigma_{v}^{2}\right)\int_{[0,1]} \left\|E\left[z_{p}(t)\right]\right\|^{2} dp + 2\kappa^{2}B(t) + 4C_{v}^{2}\right)$$

$$\leq 2\beta_{1}(t)\beta_{2}(t)\beta_{3}(t)\left(3\left(1+4\sigma_{v}^{2}\right)\left(\mathscr{R}(t)+\mathscr{K}(t)+\left\|x^{*}\right\|^{2}\right) + 4C_{v}^{2}+2\kappa^{2}B(t)\right).$$
(D.17)

Combining (D.12)-(D.17) gives

$$\frac{d\mathscr{R}(t)}{dt} \leq \left(\beta_{1}(t) + (2 + 4\kappa^{2})\beta_{1}(t)\beta_{2}(t) - 2\lambda_{2}(\mathbb{L}_{A})\beta_{3}(t) + 6\beta_{1}(t)\beta_{3}(t) + 3(2 + 8\sigma_{\nu}^{2})\beta_{1}(t)\beta_{2}(t)\beta_{3}(t)\right) \\
\times \mathscr{R}(t) + \left(\beta_{1}(t) + 2\beta_{1}(t)\beta_{3}(t)\right)\widetilde{\mathscr{R}}(t) + \left(4\kappa^{2}\beta_{2}(t) + 6\beta_{3}(t) + 3(2 + 8\sigma_{\nu}^{2})\beta_{2}(t)\beta_{3}(t)\right)\beta_{1}(t) \\
\times \mathscr{S}(t) + 2\kappa^{2}\beta_{1}(t)\beta_{2}(t)\left(1 + 2\beta_{3}(t)\right)B(t) + \left(2\sigma_{\nu}^{2}\beta_{2}(t) + 3(2 + 8\sigma_{\nu}^{2})\beta_{2}(t)\beta_{3}(t) + 6\beta_{3}(t)\right) \\
\times \beta_{1}(t)\|x^{*}\|^{2} + 2C_{\nu}^{2}\beta_{1}(t)\beta_{2}(t)(1 + 4\beta_{3}(t)).$$
(D.18)

By (D.10), we have

$$\frac{d\widetilde{\mathscr{R}}(t)}{dt} = 2\beta_3(t) \int_{0,1]\times[0,1]} A(p,q) \left(E\left[\widetilde{y}_p^{\mathsf{T}}(t)\right] E\left[\widetilde{y}_q(t)\right] - E\left[\widetilde{y}_p^{\mathsf{T}}(t)\right] E\left[\widetilde{y}_p(t)\right] \right) dqdp$$

$$+2\beta_{2}(t)\beta_{3}(t)\int_{0,1]\times[0,1]}A(p,q)\left(E\left[\widetilde{y}_{p}^{\mathsf{T}}(t)\right]E\left[\nabla_{x}V\left(q,z_{q}(t)\right)\right]-E\left[\widetilde{y}_{p}^{\mathsf{T}}(t)\right]\right)$$
$$\times E\left[\nabla_{x}V\left(p,z_{p}(t)\right)\right]dqdp.$$
(D.19)

For the first term on the r.h.s. of the above equality, combining (A.2) and (D.11) gives

$$2\beta_{3}(t)\int_{0,1]\times[0,1]}A(p,q)\left(E[\widetilde{y}_{p}^{\mathsf{T}}(t)]E\left[\widetilde{y}_{q}(t)\right]-E\left[\widetilde{y}_{p}^{\mathsf{T}}(t)\right]E[\widetilde{y}_{p}(t)]\right)dqdp \leqslant -2\beta_{3}(t)\lambda_{2}(\mathbb{L}_{A})\widetilde{\mathscr{R}}(t).$$
(D.20)

For the second term on the r.h.s. of (D.19), similar to the proof of (D.17) and by Assumption 2.1 (ii), C_r inequality and Jensen inequality, we have

$$2\beta_{2}(t)\beta_{3}(t)\int_{0,1]\times[0,1]}A(p,q)\left(E\left[\widetilde{y}_{p}^{\mathsf{T}}(t)\right]E\left[\nabla_{x}V\left(q,z_{q}(t)\right)\right]\right)$$
$$-E\left[\widetilde{y}_{p}^{\mathsf{T}}(t)\right]E\left[\nabla_{x}V\left(p,z_{p}(t)\right)\right]\right)dqdp$$
$$\leq 2\beta_{2}(t)\beta_{3}(t)\left(\int_{[0,1]}\left\|E\left[\widetilde{y}_{p}(t)\right]\right\|^{2}dp+2\kappa^{2}B(t)+4\int_{[0,1]}\left(\sigma_{v}^{2}\left\|E\left[z_{p}(t)\right]\right\|^{2}+C_{v}^{2}\right)dp\right)$$
$$\leq 2\beta_{2}(t)\beta_{3}(t)\left(\widetilde{\mathscr{R}}(t)+2\kappa^{2}B(t)+12\sigma_{v}^{2}\left(\mathscr{R}(t)+\mathscr{S}(t)+\|x^{*}\|^{2}\right)+4C_{v}^{2}\right).$$

This together with (D.19)-(D.20) gives

$$\frac{d\widetilde{\mathscr{R}}(t)}{dt} \leq \left(-2\lambda_2(\mathbb{L}_A) + 2\beta_2(t)\right)\beta_3(t)\widetilde{\mathscr{R}}(t) + 4\beta_2(t)\beta_3(t)\left(6\sigma_v^2\mathscr{R}(t) + 6\sigma_v^2\mathscr{S}(t) + \kappa^2 B(t) + 6\sigma_v^2\|x^*\|^2 + 2C_v^2\right).$$
(D.21)

Then, by (D.18), (D.21) and $\max\{z, y\} = \frac{z+y+|z-y|}{2}, \forall z, y \in \mathbb{R}$, we have

$$\frac{d\widetilde{\mathscr{Y}(t)}}{dt} \leq \max\left\{\beta_{1}(t) + (2+4\kappa^{2})\beta_{1}(t)\beta_{2}(t) - 2\lambda_{2}(\mathbb{L}_{A})\beta_{3}(t) + 6\beta_{1}(t)\beta_{3}(t) + 3\left(2+8\sigma_{\nu}^{2}\right)\beta_{1}(t)\beta_{2}(t)\beta_{3}(t) + 24\sigma_{\nu}^{2}\beta_{2}(t)\beta_{3}(t), \ \beta_{1}(t) + 2\beta_{1}(t)\beta_{3}(t) - 2\beta_{3}(t)\lambda_{2}(\mathbb{L}_{A}) + 2\beta_{2}(t)\beta_{3}(t)\right\}\widetilde{\mathscr{Y}(t)} + m_{2}(t)\mathscr{S}(t) + m_{3}(t) \leq m_{1}(t)\widetilde{\mathscr{Y}(t)} + m_{2}(t)\mathscr{S}(t) + m_{3}(t),$$
(D.22)

where $m_1(t) = -2\beta_3(t)\lambda_2(\mathbb{L}_A) + \beta_1(t) + (2+4\kappa^2)\beta_1(t)\beta_2(t) + 6\beta_1(t)\beta_3(t) + 3(2+8\sigma_v^2)\beta_1(t)\beta_2(t)$ $\times \beta_3(t) + 24\sigma_v^2\beta_2(t)\beta_3(t), m_2(t) = 4\kappa^2\beta_1(t)\beta_2(t) + 6\beta_1(t)\beta_3(t) + 3(2+8\sigma_v^2)\beta_1(t)\beta_2(t)\beta_3(t) + 24\sigma_v^2\beta_2(t)\beta_3(t)$ and $m_3(t) = 2\beta_2(t)B(t)(\kappa^2\beta_1(t) + 2\kappa^2\beta_1(t)\beta_3(t) + 2\kappa^2\beta_3(t)) + (2\sigma_v^2\beta_1(t)\beta_2(t) + 3(2+8\sigma_v^2)\beta_1(t)\beta_2(t)\beta_3(t) + 6\beta_1(t)\beta_3(t) + 24\sigma_v^2\beta_2(t)\beta_3(t)) \|x^*\|^2 + 2\beta_2(t)(C_v^2\beta_1(t) + 4C_v^2\beta_1(t))$ $\times \beta_3(t) + 4C_{\nu}^2\beta_3(t)$). By (D.9)-(D.11), Assumption 2.1 (ii), C_r inequality, Hölder inequality and Jensen inequality, we have

$$\begin{aligned} \frac{d\mathscr{P}(t)}{dt} \\ &= -2\beta_{1}(t)\beta_{2}(t)(R_{1}(t) - x^{*})^{\mathsf{T}}\int_{[0,1]} \left(E\left[\nabla_{x}V\left(p,z_{p}(t)\right)\right] - \nabla_{x}V\left(p,R_{1}(t)\right)\right)dp \\ &- 2\beta_{1}(t)\beta_{2}(t)(R_{1}(t) - x^{*})^{\mathsf{T}}\int_{[0,1]} \left(\nabla_{x}V\left(p,R_{1}(t)\right) - \nabla_{x}V\left(p,x^{*}\right)\right)dp \\ &\leqslant -2\kappa_{2}\beta_{1}(t)\beta_{2}(t)\mathscr{P}(t) + 2\beta_{1}(t)\beta_{2}(t)\mathscr{P}^{\frac{1}{2}}(t)\left(\int_{[0,1]} E\left[\|\nabla_{x}V\left(p,z_{p}(t)\right)\right. \\ &- \nabla_{x}V(p,R_{1}(t))\|^{2}\right]dp\right)^{\frac{1}{2}} \\ &\leqslant -2\kappa_{2}\beta_{1}(t)\beta_{2}(t)\mathscr{P}(t) + 2\beta_{1}(t)\beta_{2}(t)\mathscr{P}^{\frac{1}{2}}(t)\left(\int_{[0,1]} \kappa^{2}E\left[\|z_{p}(t) - R_{1}(t)\|^{2}\right]dp\right)^{\frac{1}{2}} \\ &\leqslant 2\beta_{1}(t)\beta_{2}(t)\left(-\kappa_{2}\mathscr{P}(t) + \mathscr{P}^{\frac{1}{2}}(t)\left(2\kappa^{2}B(t) + 2\kappa^{2}\int_{[0,1]} \|E[z_{p}(t)] - R_{1}(t)\|^{2}dp\right)^{\frac{1}{2}}\right) \\ &\leqslant 2\beta_{1}(t)\beta_{2}(t)\left(-\kappa_{2}\mathscr{P}(t) + \sqrt{2}\kappa\mathscr{P}^{\frac{1}{2}}(t)\left(B^{\frac{1}{2}}(t) + \mathscr{P}^{\frac{1}{2}}(t)\right)\right) \\ &\leqslant 2\beta_{1}(t)\beta_{2}(t)\left(-\kappa_{2}\mathscr{P}(t) + \sqrt{2}\kappa\mathscr{P}^{\frac{1}{2}}(t)\left(B^{\frac{1}{2}}(t) + \mathscr{P}^{\frac{1}{2}}(t)\right)\right). \end{aligned}$$
(D.23)

By $B(t) \leq \sup_{p \in [0,1]} E[||z_p(t) - E[z_p(t)]||^2]$ and Lemma D.1, we have

$$\lim_{t \to \infty} B(t) = 0. \tag{D.24}$$

Let $Y_1(t) = \mathscr{Y}(t), Y_2(t) = \mathscr{S}(t), Y_3(t) = B(t), a_1(t) = 2\beta_3(t)\lambda_2(\mathbb{L}_A), a_2(t) = m_1(t) - a_1(t), a_3(t) = m_2(t), a_4(t) = m_3(t), b_1(t) = 2\kappa_2\beta_1(t)\beta_2(t)$ and $b_2(t) = 2\sqrt{2}\kappa\beta_1(t)\beta_2(t)$ in Lemma 4.1. Then, by (D.22), (D.23), Lemma 4.1 and Assumption 4.1, we have

$$\lim_{t \to \infty} \mathscr{R}(t) = 0, \tag{D.25}$$

$$\lim_{t \to \infty} \widetilde{\mathscr{R}}(t) = 0 \tag{D.26}$$

and (25). Then, by the above equalities and Lemma D.1, we have $\sup_{t \ge 0, p \in [0,1]} E[\|z_p(t) - E[z_p(t)]\|^2] < \infty$, $\sup_{t \ge 0} \int_{[0,1]} \|E[z_p(t)] - \int_{[0,1]} E[z_q(t)] dq \|^2 dp < \infty$, $\sup_{t \ge 0} \|\int_{[0,1]} E[z_q(t)] dq - x^* \|^2 < \infty$, $\sup_{t \ge 0, p \in [0,1]} E[\|\widetilde{y}_p(t) - E[\widetilde{y}_p(t)]\|^2] < \infty$ and $\sup_{t \ge 0} \int_{[0,1]} \|E[\widetilde{y}_p(t)]\|^2 dp < \infty$. This together with C_r inequality gives

$$\sup_{t \ge 0} \int_{[0,1]} E\big[\|z_p(t)\|^2 \big] dp$$

$$\leq 4 \sup_{t \geq 0} \int_{[0,1]} E\left[\|z_p(t) - E[z_p(t)]\|^2 \right] dp + 4 \sup_{t \geq 0} \int_{[0,1]} \left\| E\left[z_p(t)\right] - \int_{[0,1]} E\left[z_q(t)\right] dq \right\|^2 dp + 4 \sup_{t \geq 0} \left\| \int_{[0,1]} E\left[z_q(t)\right] dq - x^* \right\|^2 + 4 \|x^*\|^2 \leq 4 \sup_{t \geq 0, p \in [0,1]} E\left[\|z_p(t) - E[z_p(t)]\|^2 \right] + 4 \|x^*\|^2 + 4 \sup_{t \geq 0} \int_{[0,1]} \left\| E\left[z_p(t)\right] - \int_{[0,1]} E\left[z_q(t)\right] dq \right\|^2 dp + 4 \sup_{t \geq 0} \left\| \int_{[0,1]} E\left[z_q(t)\right] dq - x^* \right\|^2 < \infty$$
 (D.27)

and

$$\sup_{t \ge 0} \int_{[0,1]} E\left[\|\widetilde{y}_{p}(t)\|^{2} \right] dp
\leqslant \sup_{t \ge 0} \int_{[0,1]} E\left[\|\widetilde{y}_{p}(t) - E[\widetilde{y}_{p}(t)]\|^{2} \right] dp + \sup_{t \ge 0} \int_{[0,1]} \|E[\widetilde{y}_{p}(t)]\|^{2} dp
\leqslant \sup_{t \ge 0, p \in [0,1]} E\left[\|\widetilde{y}_{p}(t) - E[\widetilde{y}_{p}(t)]\|^{2} \right] + \sup_{t \ge 0} \int_{[0,1]} \|E[\widetilde{y}_{p}(t)]\|^{2} dp < \infty.$$
(D.28)

Then, similar to the proof of (B.9) in Theorem 2.1 and by Assumption 2.1 (i), we have (21) and (22). By (21), (22), (D.25), (D.26) and Theorem 2.1, we have (23) and (24).

REFERENCES

- [1] B. McMahan, E. Moore, D. Ramage, S. Hampson and B. A. Arcas, "Communicationefficient learning of deep networks from decentralized data," in *Proc. 20th Int. Conf. Artif. Intell. Stat.*, Fort Lauderdale, USA, Apr. 20-22, 2017, pp. 1273-1282.
- [2] D. Li, K. D. Wong, Y. H. Hu and A. M. Sayeed, "Detection, classification and tracking of targets," *IEEE Signal Process. Mag.*, vol. 19, no. 2, pp. 17-29, 2002.
- [3] R. Madan and S. Lall, "Distributed algorithms for maximum lifetime routing in wireless sensor networks," *IEEE Trans. Wirel. Commun.*, vol. 5, no. 8, pp. 2185-2193, 2006.
- [4] M. Huang, P. E. Caines and R. P. Malhamé, "Individual and mass behaviour in large population stochastic wireless power control problems: Centralized and Nash equilibrium solutions," in *Proc. 42nd IEEE Conf. Decision Control*, Hawaii, USA, Dec. 9-12, 2003, pp. 98-103.
- [5] J. M. Lasry and P. L. Lions, "Jeux à champ moyen. I. Le cas stationnaire," C. R. Math., vol. 343, no. 9, pp. 619-625, 2006.
- [6] A. Lachapelle, J. Salomon and G. Turinici, "Computation of mean field equilibria in economics," *Math. Models Meth. Appl. Sci.*, vol. 20, no. 4, pp. 567-588, 2010.

July 4, 2024

- [7] D. Gomes, L. Lafleche and L. Nurbekyan, "A mean-field game economic growth model," in *Proc. 2016 Amer. Control Conf.*, Boston, USA, Jul. 6-8, 2016, pp. 4693-4698.
- [8] A. Bensoussan, K. C. J. Sung, S. C. P. Yam and S. P. Yung, "Linear-quadratic mean field games," J. Optim. Theory Appl., vol. 169, pp. 496-529, 2016.
- [9] M. Bardi, "Explicit solutions of some linear-quadratic mean field games," *Netw. Heterog. Media*, vol. 7, no. 2, pp. 243-261, 2012.
- [10] M. Bardi and F. S. Priuli, "Linear-quadratic N-person and mean-field games with ergodic cost," SIAM J. Control Optim., vol. 52, no. 5, pp. 3022-3052, 2014.
- [11] J. Moon and T. Başar, "Linear quadratic risk-sensitive and robust mean field games," *IEEE Trans. Autom. Control*, vol. 62, no. 3, pp. 1062-1077, 2016.
- [12] M. Nourian and P. E. Caines, "ε-Nash mean field game theory for nonlinear stochastic dynamical systems with major and minor agents," *SIAM J. Control Optim.*, vol. 51, no. 4, pp. 3302-3331, 2013.
- [13] P. Cardaliaguet and C. A. Lehalle, "Mean field game of controls and an application to trade crowding," *Math. Financ. Econ.*, vol. 12, pp. 335-363, 2018.
- [14] L. Lovász, Large Networks and Graph Limits. Rhode Island: American Mathematical Society, 2012.
- [15] E. Bayraktar, S. Chakraborty and R. Wu. "Graphon mean field systems," *Ann. Appl. Probab.*, vol. 33, no. 5, pp. 3587-3619, 2023.
- [16] E. Bayraktar, R. Wu and X. Zhang, "Propagation of chaos of forward-backward stochastic differential equations with graphon interactions," *Appl. Math. Optim.*, vol. 88, no. 25, 2023.
- [17] G. Bet, F. Coppini and F. R. Nardi, "Weakly interacting oscillators on dense random graphs," *J. Appl. Probab.*, vol. 61, no. 1, pp. 255-278, 2024.
- [18] E. Bayraktar and R. Wu, "Stationarity and uniform in time convergence for the graphon particle system," *Stoch. Process. Their Appl.*, vol. 150, pp. 532-568, 2022.
- [19] E. Bayraktar and R. Wu, "Graphon particle system: Uniform-in-time concentration bounds," *Stoch. Process. Their Appl.*, vol. 156, pp. 196-225, 2023.
- [20] P. E. Caines and M. Huang, "Graphon mean field games and their equations," SIAM J. Control Optim., vol. 59, no. 6, pp. 4373-4399, 2021.
- [21] S. Gao, P. E. Caines and M. Huang, "LQG graphon mean field games: Graphon invariant subspaces," in *Proc. 60th IEEE Conf. Decision Control*, Austin, USA, Dec. 13-17, 2021, pp. 5253-5260.

- [22] C. Fabian, K. Cui and H. Koeppl, "Mean field games on weighted and directed graphs via colored digraphons," *IEEE Control Syst. Lett.*, vol. 7, pp. 877-882, 2022.
- [23] R. Foguen-Tchuendom, S. Gao, P. E. Caines and M. Huang, "Infinite horizon LQG graphon mean field games: Explicit Nash values and local minima," *Syst. Control Lett.*, vol. 187, pp. 105780, 2024.
- [24] S. Gao, P. E. Caines and M. Huang, "LQG graphon mean field games: Analysis via graphoninvariant subspaces," *IEEE Trans. Autom. Control*, vol. 68, no. 12, pp. 7482-7497, 2023.
- [25] M. Bin, I. Notarnicola, L. Marconi and G. Notarstefano, "A system theoretical perspective to gradient-tracking algorithms for distributed quadratic optimization," in *Proc. 58th IEEE Conf. Decision Control*, Nice, France, Dec. 11-13, 2019, pp. 2994-2999.
- [26] M. A. Gkogkas and C. Kuehn, "Graphop mean-field limits for Kuramoto-type models," SIAM J. Appl. Dyn. Syst., vol. 21, no. 1, pp. 248-283, 2022.
- [27] J. Baladron, D. Fasoli, O. Faugeras and J. Touboul, "Mean-field description and propagation of chaos in networks of Hodgkin-Huxley and FitzHugh-Nagumo neurons," J. Math. Neurosc., vol. 2, no. 10, pp. 1-50, 2012.
- [28] R. Vizuete, P. Frasca and F. Garin, "Graphon-based sensitivity analysis of SIS epidemics," *IEEE Control Syst. Lett.*, vol. 4, no. 3, pp. 542-547, 2020.
- [29] M. A. Gkogkas, C. Kuehn and C. Xu, "Mean field limits of co-evolutionary heterogeneous networks," arXiv:2202.01742, 2022.
- [30] B. Bonnet, N. P. Duteil and M. Sigalotti, "Consensus formation in first-order graphon models with time-varying topologies," *Math. Models Meth. Appl. Sci.*, vol. 32, no. 11, pp. 2121-2188, 2022.
- [31] B. Øksendal, Stochastic Differential Equations: an Introduction With Applications, 6th ed. Berlin: Springer, 2003.
- [32] A. N. Michel and R. K. Miller, *Qualitative Analysis of Large Scale Dynamical Systems*. New York: Academic Press, 1977.
- [33] L. Ambrosio, N. Gigli and G. Savaré, Gradient Flows: in Metric Spaces and in the Space of Probability Measures. Berlin: Springer Science & Business Media, 2005.
- [34] Y. Chen and T. Li, "A large-scale stochastic gradient descent algorithm over a graphon," in *Proc. 62nd IEEE Conf. Decision Control*, Marina Bay Sands, Singapore, Dec. 13-15, 2023, pp. 4806-4811.
- [35] Y. Chen and T. Li, "Graphon particle systems, part I: Spatio-temporal approximation and

July 4, 2024

law of large numbers." arXiv:2405.16760, 2024.