

# THE STRUCTURE OF DEFORMED DOUBLE COMPLEXES ON THE IWASAWA MANIFOLD

YAN HU, WEI XIA

ABSTRACT. The Kuranishi family of the Iwasawa manifold give rise naturally to a family of (deformed) double complexes. By using the structure theorem of double complexes due to Stelzig and Qi-Khovanov, we show there are exactly 3 isomorphism types in this family and determine explicitly structures of these 3 types. As an application, we computed the Frölicher spectral sequence for each fiber in the Kuranishi family of the Iwasawa manifold.

## 1. INTRODUCTION

Given a complex manifold  $X$ , the space of (complex) differential forms  $A^\bullet(X)$  admits a natural double complex structure with differential given by  $d = \partial + \bar{\partial}$ . From this double complex, denoted by

$$(1.1) \quad (A^\bullet(X), d = \partial + \bar{\partial}),$$

the Dolbeault/Bott-Chern/Aeppli cohomology and the Frölicher spectral sequence can be derived. These objects has been extensively studied in recent years (see e.g. [AK17, ZR15, RYY19, AT13, Kas13, PSU21, CFG91, CFUG97]). In order to attain a deeper comprehension of these complex structure invariants, it is necessary to study the structure of  $(A^\bullet(X), d = \partial + \bar{\partial})$ . In fact, according to the structure theorem of double complexes due to Stelzig [Ste21] and Qi-Khovanov [KQ20], any double complex can be decomposed into a direct sum of indecomposable ones (i.e. squares and zigzags) and the Dolbeault/Bott-Chern/Aeppli cohomology, the Frölicher spectral sequence can be determined by counting zigzags of specified types.

Instead of a single complex manifold  $X$ , we are more interested in how the double complex  $(A^\bullet(X), d = \partial + \bar{\partial})$  vary as the complex structure on  $X$  deforms.

Let  $\pi : (\mathcal{X}, X) \rightarrow (B, 0)$  be a small deformation of the compact complex manifold  $X$  such that for each  $t \in B$  the complex structure on  $X_t$  is represented by Beltrami differential  $\phi(t) \in A^{0,1}(X, T_X^{1,0})$ . For any  $t \in B$ , there is a double complex

$$(1.2) \quad (A^\bullet(X), d_{\phi(t)} = \partial + \bar{\partial}_{\phi(t)}),$$

where  $\bar{\partial}_{\phi(t)} = \bar{\partial} - \mathcal{L}_{\phi(t)}^{1,0}$  and  $\mathcal{L}_{\phi(t)}^{1,0} = i_{\phi(t)}\partial - \partial i_{\phi(t)}$ . (1.2) may be regarded as an analytic family of deformations of the usual double complex (1.1) because it is reduced to (1.1)

---

*Date:* July 4, 2024.

This work is supported by the National Natural Science Foundation of China No. 11901590, the Natural Science Foundation of Chongqing (China) No. CSTB2022NSCQ-MSX0876 and Scientific Research Foundation of Chongqing University of Technology.

when  $t = 0$ . The double complex (1.2) is naturally related to  $(A^\bullet(X_t), d = \partial_t + \bar{\partial}_t)$ : there is an isomorphism of filtered complex (see Subsection 2.2)

$$e^{i\phi(t)} = \sum_{k=0}^{\infty} i_{\phi(t)}^k : (F^\bullet A^\bullet(X), d_{\phi(t)}) \rightarrow (F^\bullet A^\bullet(X_t), d).$$

In particular, the Frölicher spectral sequence of the double complex (1.2) is isomorphic to the Frölicher spectral sequence of  $X_t$ .


The Iwasawa manifold (or more generally, complex parallelizable solvmanifolds) and its Kuranishi family has often been adopted as a test for many propositions about cohomology and the Frölicher spectral sequence, see e.g. [Nak75, Ang13, RZ18, Mas20, Xia22]. For the Kuranishi family  $\{X_t\}_{t \in \mathcal{B}}$  of the Iwasawa manifold  $X = X_0$ , there is a natural double sub-complex

$$(\wedge^\bullet, d_{\phi(t)} = \partial + \bar{\partial}_{\phi(t)})$$


of (1.2) for any  $t \in \mathcal{B}$  which is constituted of left invariant forms on  $X$  and the inclusion map is an  $E_1$ -isomorphism in the sense of Stelzig [Ste21].

By using the structure theorem of double complexes, Flavi [Fla20] recently computed the structure of the double sub-complex  $(\wedge^\bullet(X_t), d = \partial_t + \bar{\partial}_t)$  of  $(A^\bullet(X_t), d = \partial_t + \bar{\partial}_t)$  for some small deformations of the Iwasawa manifold. Based on this, Flavi determined the Frölicher spectral sequence  $E_r^{\bullet, \bullet}$  correspondingly. We are curious about the whole picture about deformation stability of degeneration of the Frölicher spectral sequence [WX23]. Motivated by these, in this paper, we explicitly decompose the double complex  $(\wedge^\bullet, d_{\phi(t)} = \partial + \bar{\partial}_{\phi(t)})$  as indecomposable ones (i.e. squares and zigzags) for the Kuranishi family of the Iwasawa manifold. The main result of this paper may be summarized as follows:

**Theorem 1.1.** *Let  $\{X_t\}_{t \in \mathcal{B}}$  be the Kuranishi family of the Iwasawa manifold  $X = X_0$  which is defined by Beltrami differentials  $\{\phi(t) \in A^{0,1}(X, T_X^{1,0})\}_{t \in \mathcal{B}}$ . Denoted by  $\wedge^{\bullet, \bullet} \subset A^{\bullet, \bullet}(X)$  the space of left invariant forms. Then for any  $t \in \mathcal{B}$ , the structure of double complex  $(\wedge^\bullet, d_{\phi(t)} = \partial + \bar{\partial}_{\phi(t)})$  is described as follows:*



(i)  $(t_{11}, t_{12}, t_{21}, t_{22}) = 0, D(t) = 0$



where  $D(t) = t_{11}t_{22} - t_{21}t_{12}$ .

**Remark 1.2.** Compared to the result of Flavi [Fla20], we see that the structure of double complex  $(\wedge^\bullet, d_{\phi(t)} = \partial + \bar{\partial}_{\phi(t)})$  seems less complicated than that of  $(\wedge^\bullet(X_t), d = \partial_t + \bar{\partial}_t)$ : the former has exactly 3 isomorphism types while the latter has at least 4 isomorphism types. According to the computation of the Bott-Chern cohomology by Angella [Ang13], there should be at least 5 isomorphism types of  $(\wedge^\bullet(X_t), d = \partial_t + \bar{\partial}_t)$ .

As a consequence, we can easily compute the deformed Dolbeault/Bott-Chern/Aeppli cohomology [Xia22, Xia21b, HX24] and the Frölicher spectral sequence of  $X_t$ . In view of these (see Table 1 and Table 2 in Section 5), we get the following

**Corollary 1.3.** *The following statements hold:*

- (1) *Any small deformation of the Iwasawa manifold satisfies:  $E_2 = E_\infty$ . There are small deformations (i.e. case (iii)) of the Iwasawa manifold for which  $E_1 = E_\infty$  holds;*
- (2) *For the Kuranishi family of the Iwasawa manifold, both  $\dim \operatorname{im} d_1^{p,q}$  and  $\dim \frac{E_1^{p,q}}{E_2^{p,q}} = \dim \operatorname{im} d_1^{p,q} + \dim \operatorname{im} d_1^{p-1,q}$  are upper semi-continuous function of  $t \in \mathcal{B}$  (under the analytic Zariski topology);*
- (3) *The varying of  $\sum_{i=0}^q (-1)^{q-i} \dim E_2^{p,i}$  may be neither upper nor lower semi-continuous.*

It is known [Kas15] that for complex parallelizable solvmanifolds,  $E_2 = E_\infty$  always holds, that is, the Frölicher spectral sequence degenerates at the second page. Moreover, it is conjectured by Kasuya-Stelzig [KS23] that this holds for all complex parallelisable manifold. We note that for case (ii) and (iii), the deformation  $X_t$  of the Iwasawa manifold is not complex parallelisable since  $\dim H_{\bar{\partial}}^{2,0}(X_t) = 2 \neq 3$ .

A special case of a result due to Flenner [Fle81] says that the alternating sum of Hodge numbers:  $\sum_{i=0}^q (-1)^{q-i} \dim E_1^{p,i}$ , is upper semi-continuous for any  $p$  and  $q$ . From (3) of Corollary 1.3, we see that this statement does not hold for  $\sum_{i=0}^q (-1)^{q-i} \dim E_2^{p,i}$ . It is perhaps worth noting that the upper or lower semi-continuous property of

$\sum_{i=0}^q (-1)^{q-i} \dim E_2^{p,i}$  for all  $p, q$  implies that of  $\dim E_2^{p,q}$  but not conversely, compare [Mas20, Fla20].

**Question 1.4.** Let  $\{X_t\}_{t \in B}$  be a complex analytic family and  $(E_r^{\bullet, \bullet}, d_r^{\bullet, \bullet})$  the Frölicher spectral sequence on  $X_t$ . Is  $\dim \operatorname{im} d_1^{p,q}$  or  $\dim \operatorname{im} \frac{E_1^{p,q}}{E_2^{p,q}} = \dim \operatorname{im} d_1^{p,q} + \dim \operatorname{im} d_1^{p-1,q}$  upper semi-continuous function of  $t \in B$ ?

## 2. PRELIMINARY

**2.1. The double complex  $(A^\bullet(X), d = \partial + \bar{\partial})$ .** Let  $X$  be a complex manifold and  $A^\bullet(X)$  the space of (complex valued) smooth forms on  $X$ . The de Rham complex  $(A^\bullet(X), d)$  naturally has a double complex structure given by

$$A^\bullet(X) = \bigoplus_{p \in \mathbb{N}} A^{p, \bullet-p}(X), \quad d = \partial + \bar{\partial},$$

where  $A^{p, \bullet-p}(X)$  is the space of  $(p, \bullet-p)$ -forms on  $X$ . This double complex  $(A^\bullet(X), d = \partial + \bar{\partial})$  induce a filtered complex

$$(F^\bullet A^\bullet(X), d), \quad \text{with } F^p A^\bullet(X) = \bigoplus_{\lambda \geq p} A^{\lambda, \bullet-\lambda}(X),$$

whose associated spectral sequence  $(E_r^{p,q}(X), d_r(X))$  is called the *Frölicher spectral sequence* of  $X$ .

**2.2. The deformed double complex  $(A^\bullet(X), d_{\phi(t)} = \partial + \bar{\partial}_{\phi(t)})$ .** Now, let

$$\pi : (\mathcal{X}, X) \rightarrow (B, 0)$$

be a complex analytic family (i.e. a surjective, proper and homomorphic submersion between complex manifolds). We make the following conventions which will be used throughout this paper:

- $X = \pi^{-1}(0)$  and  $X_t = \pi^{-1}(t)$  for any  $t \in B$ ;
- since only small deformations will be considered, we may assume  $B$  is a small polydisc in some Euclidean space;
- for any  $t \in B$ , the complex structure on  $X_t = \pi^{-1}(t)$  is represented by (the Beltrami differential)  $\phi(t) \in A^{0,1}(X, T^{1,0})$ .

By using the Beltrami differential  $\phi(t)$ , we can relate smooth forms on  $X$  to those on  $X_t$ . In fact, the exponential operator [Tod89, Cle05, LSY09, LRY15]

$$e^{i\phi(t)} := \sum_{k=0}^{\infty} \frac{i^k \phi(t)^k}{k!},$$

where  $i_{\phi(t)}$  is the contraction operator, satisfy (see [FM06, FM09, WZ24])

$$(2.1) \quad e^{i\phi(t)} : F^p A^k(X) \longrightarrow F^p A^k(X_t).$$

Set (see e.g. [LRY15, FM06, Xia19, Man04])

$$(2.2) \quad d_{\phi(t)} := e^{-i\phi(t)} d e^{i\phi(t)} = \partial + \bar{\partial}_{\phi(t)} = \partial + \bar{\partial} - \mathcal{L}_{\phi(t)}^{1,0},$$

where  $\bar{\partial}_{\phi(t)} := \bar{\partial} - \mathcal{L}_{\phi(t)}^{1,0}$  and  $\mathcal{L}_{\phi(t)}^{1,0} := i_{\phi(t)}\partial - \partial i_{\phi(t)}$  is the Lie derivative. Then  $(A^\bullet(X), d_{\phi(t)} = \partial + \bar{\partial}_{\phi(t)})$  constitutes a family of *deformed double complex*: when  $t = 0$ , it is reduced to  $(A^\bullet(X), d = \partial + \bar{\partial})$ .

There is an intimate relation between  $(A^\bullet(X), d_{\phi(t)} = \partial + \bar{\partial}_{\phi(t)})$  and  $(A^\bullet(X_t), d = \partial_t + \bar{\partial}_t)$ : indeed, as observed in [WX23],

$$(F^\bullet A^\bullet(X), d_{\phi(t)}) \xrightarrow{e^{i\phi(t)}} (F^\bullet A^\bullet(X_t), d)$$

is an isomorphism of filtered complexes which induces the following isomorphism

$$E_r^{\bullet,\bullet}(X, d_{\phi(t)}) \cong E_r^{\bullet,\bullet}(X_t),$$

where  $E_r^{\bullet,\bullet}(X, d_{\phi(t)})$  is the spectral sequence associated to  $(F^\bullet A^\bullet(X), d_{\phi(t)})$ .

**2.3. The structure theorem for double complexes.** In this subsection, we briefly recall the structure theorem for double complexes. The readers is referred to [Ste21, KQ20] for more information.

A *double complex* (over  $\mathbb{C}$ ) is a bigraded  $\mathbb{C}$ -vector space

$$A = \bigoplus_{p,q \in \mathbb{Z}} A^{p,q}$$

together with linear maps

$$\partial_1 : A^{p,q} \longrightarrow A^{p+1,q}, \quad \partial_2 : A^{p,q} \longrightarrow A^{p,q+1}, \quad p, q \in \mathbb{Z},$$

such that  $d = \partial_1 + \partial_2$  is a differential, i.e.  $(\partial_1 + \partial_2)^2 = 0$ . A nonzero double complex  $A$  is called *indecomposable* if there exist no nontrivial decomposition  $A = A_1 \oplus A_2$ . It is possible to write down all indecomposable double complexes. To describe these, we make the following conventions:

- (1) All drawn components are one-dimensional;
- (2) All drawn maps are isomorphisms;
- (3) All components and maps not drawn are zero.

With these in minds, indecomposable double complexes may be listed as following

- Squares:

$$\begin{array}{ccc} A^{p-1,q} & \xrightarrow{\partial_1} & A^{p,q} \\ \uparrow \partial_2 & & \uparrow \partial_2 \\ A^{p-1,q-1} & \xrightarrow{\partial_1} & A^{p,q-1}, \end{array}$$

- Zigzags:

$$\begin{array}{ccccccc}
A^{p,q}(\text{dot}), & A^{p,q+1}, & A^{p,q} \xrightarrow{\partial_1} & A^{p+1,q}, & A^{p,q+1} & & , \\
& \uparrow \partial_2 & & & \uparrow \partial_2 & & \\
& A^{p,q} & & & A^{p,q} \xrightarrow{\partial_1} & A^{p+1,q} & \\
A^{p-1,q} \xrightarrow{\partial_1} & A^{p,q}, & A^{p-1,q} \xrightarrow{\partial_1} & A^{p,q} & & & , \dots \\
\uparrow \partial_2 & & & \uparrow \partial_2 & & & \\
A^{p-1,q-1} & & & A^{p,q-1} \xrightarrow{\partial_1} & A^{p+1,q-1} & & 
\end{array}$$

For a square or a zigzag  $A$ , the *shape* is defined to be

$$S(A) = \{(p, q) \in \mathbb{Z}^2 \mid A^{p,q} \neq 0\}.$$

For any bounded  $S \subset \mathbb{Z}^2$ , there is an (essentially unique) double complex  $A$  with

$$A^{p,q} = \begin{cases} 0, & (p, q) \in S, \\ \mathbb{C}, & (p, q) \notin S, \end{cases}$$

and  $(\partial_1)^{p,q} = (-1)^q \text{Id}$ ,  $(\partial_2)^{p,q} = \text{Id}$ . This double complex will be denoted by  $C(S)$ . Now we are ready to state the structure theorem of double complexes due to Stelzig [Ste21, Thm. 3.4] and Qi-Khovanov [KQ20]:

**Theorem 2.1** (Structure theorem of double complexes). *For every bounded double complex  $A$ , there exist unique cardinal numbers  $\text{mult}_S(A)$  and a (non-unique) isomorphism*

$$A \cong \bigoplus_S C(S)^{\oplus \text{mult}_S(A)},$$

where  $S \subset \mathbb{Z}^2$  runs over the set of all shapes of squares and zigzags.

We will apply this theorem to the double complex  $(\wedge^\bullet, d_{\phi(t)} = \partial + \bar{\partial}_{\phi(t)})$  for the Kuranishi family of the Iwasawa manifold. In this case, each component  $\wedge^{p,q}$  of the double complex is of finite dimensional so that the decomposition process terminates at finite steps.

Since the (Dolbeault, Bott-Chern, Aeppli) cohomology groups and the Frölicher spectral sequence are most easy to compute for squares and zigzags, as a consequence of Theorem 2.1, we have the following (see [RZ20] for some related discussion)

**Theorem 2.2** (Proposition 6 and Lemma 8 in [Ste21]). *Let  $(A = \bigoplus A^{p,q}, d = \partial_1 + \partial_2)$  be a bounded double complex with a decomposition into elementary complexes with pairwise distinct support  $\varphi : \bigoplus A_i \xrightarrow{\cong} A$ .*

- (1) For every  $(p, q) \in \mathbb{Z}^2$ , the maps induced by  $\varphi$

$$\bigoplus_{\substack{A_i \text{ zigzag} \\ (p,q) \in S(A_i) \\ (p-1,q), (p+1,q) \notin S(A_i)}} H_{\partial_1}^{p,q}(A_i) \longrightarrow H_{\partial_1}^{p,q}(A)$$

$$\bigoplus_{\substack{A_i \text{ zigzag} \\ (p,q) \in S(A_i) \\ (p,q-1), (p,q+1) \notin S(A_i)}} H_{\partial_2}^{p,q}(A_i) \longrightarrow H_{\partial_2}^{p,q}(A)$$

are isomorphisms.

(2) For every  $(p, q) \in \mathbb{Z}^2$ , the maps induced by  $\varphi$

$$\bigoplus_{\substack{A_i \text{ zigzag} \\ (p,q) \in S(A_i) \\ (p+1,q), (p,q+1) \notin S(A_i)}} H_{BC}^{p,q}(A_i) \longrightarrow H_{BC}^{p,q}(A)$$

$$\bigoplus_{\substack{A_i \text{ zigzag} \\ (p,q) \in S(A_i) \\ (p-1,q), (p,q-1) \notin S(A_i)}} H_A^{p,q}(A_i) \longrightarrow H_A^{p,q}(A)$$

are isomorphisms.

(3) For any  $i = 1, 2$ , let  $({}_i E_r^{p,q}, {}_i d_r^{p,q})$  be the spectral sequence associated to the filtered complex

$$({}_i F_\bullet A^k, d), \text{ with } F_1^\bullet A^k = \bigoplus_{p \geq \bullet} A^{p,k-p}, \quad F_2^\bullet A^k = \bigoplus_{q \geq \bullet} A^{k-q,q}.$$

Then there is an equality

$$\text{mult}_{S_{i,r}^{p,q}}(A) = \dim \text{im } {}_i d_r^{p,q},$$

where  $\text{mult}_{S_{i,r}^{p,q}}(A)$  is the number of zigzags (with length  $2r$ ) in  $\bigoplus_i A_i$  which has one and only one outgoing arrow at  $(p, q)$ ,  $i$  is 1 (if this arrow is  $\partial_1$ ) or 2 (if this arrow is  $\partial_2$ ).

**Definition 2.3.** Let  $r \in \mathbb{N}$ . A morphism  $f : A \rightarrow B$  of double complexes is called an  $E_r$ -isomorphism [Ste21, Def. 10] if it induces an isomorphism on the  $r$ -th page of both Frölicher spectral sequences (where the 0-th page is defined to be the complex itself).

### 3. THE KURANISHI FAMILY OF THE IWASAWA MANIFOLD AND THE DOUBLE COMPLEX $(\wedge^\bullet, d_{\phi(t)} = \partial + \bar{\partial}_{\phi(t)})$

**3.1. Iwasawa manifold and its Kuranishi family.** Let  $G$  be the complex Lie group defined by

$$G := \left\{ \begin{pmatrix} 1 & z^1 & z^3 \\ 0 & 1 & z^2 \\ 0 & 0 & 1 \end{pmatrix} \in \text{GL}(3; \mathbb{C}) \mid z^1, z^2, z^3 \in \mathbb{C} \right\} \cong \mathbb{C}^3,$$

where the product is the one induced by matrix multiplication. This is usually called the *Heisenberg group*. Consider the discrete subgroup  $\Gamma$  defined by

$$\Gamma := \left\{ \begin{pmatrix} 1 & \omega^1 & \omega^3 \\ 0 & 1 & \omega^2 \\ 0 & 0 & 1 \end{pmatrix} \in G \mid \omega^1, \omega^2, \omega^3 \in \mathbb{Z}[\sqrt{-1}] \right\},$$

The quotient  $X = \Gamma \backslash G$  is called the *Iwasawa manifold*. A basis of  $H^0(X, \Omega^1)$  is given by

$$\varphi^1 = dz^1, \quad \varphi^2 = dz^2, \quad \varphi^3 = dz^3 - z^1 dz^2,$$

and a dual basis  $\theta^1, \theta^2, \theta^3 \in H^0(X, T_X^{1,0})$  is given by

$$\theta^1 = \frac{\partial}{\partial z^1}, \quad \theta^2 = \frac{\partial}{\partial z^2} + z^1 \frac{\partial}{\partial z^3}, \quad \theta^3 = \frac{\partial}{\partial z^3}.$$

Notice that  $\varphi^1, \varphi^2, \varphi^3$  are left invariant forms and  $\theta^1, \theta^2, \theta^3$  are left invariant vector fields on  $G$ . Denote by  $(\wedge^\bullet, d = \partial + \bar{\partial})$  the double sub-complex of  $(A^\bullet(X), d = \partial + \bar{\partial})$  consisting of left invariant forms, then

$$\wedge^\bullet = \bigoplus_{p \geq 0} \wedge^{p, \bullet - p} \quad \text{and} \quad \wedge^{1,0} = \mathbb{C}\{\varphi^1, \varphi^2, \varphi^3\},$$

where  $\wedge^{p, \bullet - p}$  is the vector space of left invariant  $(p, \bullet - p)$ -forms. The structure equation in terms of the basis  $\{\varphi^1, \varphi^2, \varphi^3\}$  is given by

$$(3.1) \quad d\varphi^1 = 0, \quad d\varphi^2 = 0, \quad d\varphi^3 = -\varphi^{12}.$$

$X$  is equipped with the Hermitian metric  $\sum_{i=1}^3 \varphi^i \otimes \bar{\varphi}^i$ . The Beltrami differential of the Kuranishi family of  $X$  is

$$\phi(t) = \sum_{i=1}^3 \sum_{\lambda=1}^2 t_{i\lambda} \theta^i \bar{\varphi}^\lambda - D(t) \theta^3 \bar{\varphi}^3, \quad \text{with } D(t) = t_{11}t_{22} - t_{21}t_{12},$$

and the Kuranishi space of  $X$  is

$$\mathcal{B} = \{t = (t_{11}, t_{12}, t_{21}, t_{22}, t_{31}, t_{32}) \in \mathbb{C}^6 \mid |t_{i\lambda}| < \epsilon, i = 1, 2, 3, \lambda = 1, 2\},$$

where  $\epsilon > 0$  is sufficiently small. Since  $\phi(t)$  is also left invariant, we see that for any  $t \in \mathcal{B}$ ,

$$(\wedge^\bullet, d_{\phi(t)} = \partial + \bar{\partial}_{\phi(t)}) \xrightarrow{\iota} (A^\bullet(X), d_{\phi(t)} = \partial + \bar{\partial}_{\phi(t)}),$$

is a double sub-complex. It is known that the Dolbeault cohomology of  $(A^\bullet(X), d_{\phi(t)} = \partial + \bar{\partial}_{\phi(t)})$  can be computed by using only left forms [Xia21a]. As a consequence,  $\iota$  is an  $E_1$ -isomorphism for each  $t \in \mathcal{B}$ . To determine the decomposition of  $(\wedge^\bullet, d = \partial + \bar{\partial}_{\phi(t)})$  indicated in Theorem 2.1, we need first to compute the differentials  $\partial$  and  $\bar{\partial}_{\phi(t)}$  so as to find all possible shapes of squares and zigzags. The former is easy to get by using the structure equation (3.1). So it is left to compute  $\bar{\partial}_{\phi(t)}$ .

**3.2. Compute  $\bar{\partial}_{\phi(t)}$  on the subcomplex of invariant forms.** Set

$$\phi_1 = \sum_{i=1}^3 \sum_{\lambda=1}^2 t_{i\lambda} \theta^i \bar{\varphi}^\lambda, \quad \phi_2 = -D(t) \theta^3 \bar{\varphi}^3,$$

then by direct computations, we have

$$\mathcal{L}_{\phi_1}^{1,0} \varphi^1 = \mathcal{L}_{\phi_1}^{1,0} \varphi^2 = \mathcal{L}_{\phi_1}^{1,0} \bar{\varphi}^1 = \mathcal{L}_{\phi_1}^{1,0} \bar{\varphi}^2 = \mathcal{L}_{\phi_1}^{1,0} \bar{\varphi}^3 = 0,$$

$$\mathcal{L}_{\phi_1}^{1,0} \varphi^3 = \sum_{\lambda=1}^2 (t_{1\lambda} \varphi^2 - t_{2\lambda} \varphi^1) \wedge \bar{\varphi}^\lambda, \quad \mathcal{L}_{\phi_2}^{1,0} \varphi^i = \mathcal{L}_{\phi_2}^{1,0} \bar{\varphi}^i = 0, \quad i = 1, 2, 3.$$



In particular,

$$(3.2) \quad \bar{\partial}_{\phi(t)} = \bar{\partial} - \mathcal{L}_{\phi_1}^{1,0}.$$

3.2.1.  $\bar{\partial}_{\phi(t)} : \wedge^{1,0} \rightarrow \wedge^{1,1} = \mathbb{C}\{\varphi^{i\bar{j}} \mid 1 \leq i, j \leq 3\}$ . Recall that  $\wedge^{1,0} = \mathbb{C}\{\varphi^1, \varphi^2, \varphi^3\}$  and  $\wedge^{1,1} = \mathbb{C}\{\varphi^{i\bar{j}} \mid 1 \leq i, j \leq 3\}$ .

$$\begin{aligned} \bar{\partial}_{\phi(t)}\varphi^1 &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^1 = 0, \\ \bar{\partial}_{\phi(t)}\varphi^2 &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^2 = 0, \\ \bar{\partial}_{\phi(t)}\varphi^3 &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^3 = -\mathcal{L}_{\phi_1}^{1,0}\varphi^3 \\ &= -\sum_{\lambda=1}^2 (t_{1\lambda}\varphi^2 - t_{2\lambda}\varphi^1) \wedge \varphi^{\bar{\lambda}} \\ &= t_{21}\varphi^{1\bar{1}} - t_{11}\varphi^{2\bar{1}} + t_{22}\varphi^{1\bar{2}} - t_{12}\varphi^{2\bar{2}}. \end{aligned}$$

We see that

$$\dim \operatorname{im} \bar{\partial}_{\phi(t)} \cap \wedge^{1,1} = \begin{cases} 0, & (t_{11}, t_{12}, t_{21}, t_{22}) = 0 \\ 1, & (t_{11}, t_{12}, t_{21}, t_{22}) \neq 0 \end{cases}.$$

3.2.2.  $\bar{\partial}_{\phi(t)} : \wedge^{0,1} \rightarrow \wedge^{0,2}$ . Recall that  $\wedge^{0,1} = \mathbb{C}\{\varphi^{\bar{1}}, \varphi^{\bar{2}}, \varphi^{\bar{3}}\}$  and  $\wedge^{0,2} = \mathbb{C}\{\varphi^{\bar{1}\bar{2}}, \varphi^{\bar{1}\bar{3}}, \varphi^{\bar{2}\bar{3}}\}$ .

$$\begin{aligned} \bar{\partial}_{\phi(t)}\varphi^{\bar{1}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{\bar{1}} = 0, \\ \bar{\partial}_{\phi(t)}\varphi^{\bar{2}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{\bar{2}} = 0, \\ \bar{\partial}_{\phi(t)}\varphi^{\bar{3}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{\bar{3}} = -\varphi^{\bar{1}\bar{2}} \neq 0. \end{aligned}$$

We see that  $\dim \operatorname{im} \bar{\partial}_{\phi(t)} \cap \wedge^{0,2} = 1$ .

3.2.3.  $\bar{\partial}_{\phi(t)} : \wedge^{2,0} \rightarrow \wedge^{2,1}$ . Recall that  $\wedge^{2,0} = \mathbb{C}\{\varphi^{12}, \varphi^{13}, \varphi^{23}\}$  and  $\wedge^{2,1} = \mathbb{C}\{\varphi^{ij\bar{k}} \mid 1 \leq i, j, k \leq 3, i < j\}$ .

$$\begin{aligned} \bar{\partial}_{\phi(t)}\varphi^{12} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{12} = 0, \\ \bar{\partial}_{\phi(t)}\varphi^{13} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{13} = t_{11}\varphi^{12\bar{1}} + t_{12}\varphi^{12\bar{2}} \\ \bar{\partial}_{\phi(t)}\varphi^{23} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{23} = t_{21}\varphi^{12\bar{1}} + t_{22}\varphi^{12\bar{2}}. \end{aligned}$$

Let  $\sigma_0 = a_1\varphi^{13} + a_2\varphi^{23}$ , then

$$\begin{aligned} \bar{\partial}_{\phi(t)}\sigma_0 &= a_1(t_{11}\varphi^{12\bar{1}} + t_{12}\varphi^{12\bar{2}}) + a_2(t_{21}\varphi^{12\bar{1}} + t_{22}\varphi^{12\bar{2}}) \\ &= (a_1t_{11} + a_2t_{21})\varphi^{12\bar{1}} + (a_1t_{12} + a_2t_{22})\varphi^{12\bar{2}}. \end{aligned}$$

Set

$$T = \begin{pmatrix} t_{11} & t_{21} \\ t_{12} & t_{22} \end{pmatrix},$$

then

(1) When  $(t_{11}, t_{12}, t_{21}, t_{22}) = 0$ , we have  $\operatorname{rank}(T) = 0$  and  $\dim \operatorname{im} \bar{\partial}_{\phi(t)} \cap \wedge^{2,1} = 0$ .

- (2) When  $(t_{11}, t_{12}, t_{21}, t_{22}) \neq 0, D(t) = 0$ , we have  $\text{rank}(T) = 1$  and  $\dim \text{im } \bar{\partial}_{\phi(t)} \cap \wedge^{2,1} = 1$ .
- (3) When  $(t_{11}, t_{12}, t_{21}, t_{22}) \neq 0, D(t) \neq 0$ , we have  $\text{rank}(T) = 2$  and  $\dim \text{im } \bar{\partial}_{\phi(t)} \cap \wedge^{2,1} = 2$ .

3.2.4.  $\bar{\partial}_{\phi(t)} : \wedge^{1,1} \rightarrow \wedge^{1,2}$ . Recall that  $\wedge^{1,1} = \mathbb{C}\{\varphi^{1\bar{1}}, \varphi^{1\bar{2}}, \varphi^{1\bar{3}}, \varphi^{2\bar{1}}, \varphi^{2\bar{2}}, \varphi^{2\bar{3}}, \varphi^{3\bar{1}}, \varphi^{3\bar{2}}, \varphi^{3\bar{3}}\}$  and  $\wedge^{1,2} = \mathbb{C}\{\varphi^{i\bar{j}\bar{k}} \mid 1 \leq i, j, k \leq 3, j < k\}$ .

$$\begin{aligned}
\bar{\partial}_{\phi(t)}\varphi^{1\bar{1}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{1\bar{1}} = 0, \\
\bar{\partial}_{\phi(t)}\varphi^{1\bar{2}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{1\bar{2}} = 0, \\
\bar{\partial}_{\phi(t)}\varphi^{2\bar{1}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{2\bar{1}} = 0, \\
\bar{\partial}_{\phi(t)}\varphi^{2\bar{2}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{2\bar{2}} = 0, \\
\bar{\partial}_{\phi(t)}\varphi^{1\bar{3}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{1\bar{3}} = \varphi^{1\bar{1}\bar{2}}, \\
\bar{\partial}_{\phi(t)}\varphi^{2\bar{3}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{2\bar{3}} = \varphi^{2\bar{1}\bar{2}}, \\
\bar{\partial}_{\phi(t)}\varphi^{3\bar{1}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{3\bar{1}} = -\mathcal{L}_{\phi_1}^{1,0}\varphi^{3\bar{1}} \\
&= -\sum_{\lambda=1}^2 (t_{1\lambda}\varphi^2 - t_{2\lambda}\varphi^1) \wedge \varphi^{\bar{\lambda}} \wedge \varphi^{\bar{1}} \\
&= -t_{22}\varphi^{1\bar{1}\bar{2}} + t_{12}\varphi^{2\bar{1}\bar{2}}, \\
\bar{\partial}_{\phi(t)}\varphi^{3\bar{2}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{3\bar{2}} = -\mathcal{L}_{\phi_1}^{1,0}\varphi^{3\bar{2}} \\
&= -\sum_{\lambda=1}^2 (t_{1\lambda}\varphi^2 - t_{2\lambda}\varphi^1) \wedge \varphi^{\bar{\lambda}} \wedge \varphi^{\bar{2}} \\
&= t_{21}\varphi^{1\bar{1}\bar{2}} - t_{11}\varphi^{2\bar{1}\bar{2}}.
\end{aligned}$$

Let  $\sigma_0 = a_1\varphi^{1\bar{3}} + a_2\varphi^{2\bar{3}} + a_3\varphi^{3\bar{1}} + a_4\varphi^{3\bar{2}}$ , then

$$\begin{aligned}
\bar{\partial}_{\phi(t)}\sigma_0 &= a_1\varphi^{1\bar{1}\bar{2}} + a_2\varphi^{2\bar{1}\bar{2}} + a_3(-t_{22}\varphi^{1\bar{1}\bar{2}} + t_{12}\varphi^{2\bar{1}\bar{2}}) + a_4(t_{21}\varphi^{1\bar{1}\bar{2}} - t_{11}\varphi^{2\bar{1}\bar{2}}) \\
&= (a_1 - a_3t_{22} + a_4t_{21})\varphi^{1\bar{1}\bar{2}} + (a_2 + a_3t_{12} - a_4t_{11})\varphi^{2\bar{1}\bar{2}}.
\end{aligned}$$

Set

$$T = \begin{pmatrix} 1 & 0 & -t_{22} & t_{21} \\ 0 & 1 & t_{12} & -t_{11} \end{pmatrix},$$

we see that the rank of

$$\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{1\bar{3}}, \varphi^{2\bar{3}}, \varphi^{3\bar{1}}, \varphi^{3\bar{2}}\} \longrightarrow \mathbb{C}\{\varphi^{1\bar{1}\bar{2}}, \varphi^{2\bar{1}\bar{2}}\},$$

is equal to  $\text{rank}(T) = 2$ . Similarly, from

$$\begin{aligned}\bar{\partial}_{\phi(t)}\varphi^{3\bar{3}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{3\bar{3}} = -\varphi^3 \wedge \bar{\partial}\varphi^{\bar{3}} - \mathcal{L}_{\phi_1}^{1,0}\varphi^3 \wedge \varphi^{\bar{3}} \\ &= \varphi^{3\bar{1}\bar{2}} - \sum_{\lambda=1}^2 (t_{1\lambda}\varphi^2 - t_{2\lambda}\varphi^1) \wedge \varphi^{\bar{\lambda}} \wedge \varphi^{\bar{3}} \\ &= \varphi^{3\bar{1}\bar{2}} - t_{11}\varphi^{2\bar{1}\bar{3}} + t_{21}\varphi^{1\bar{1}\bar{3}} - t_{12}\varphi^{2\bar{2}\bar{3}} + t_{22}\varphi^{1\bar{2}\bar{3}},\end{aligned}$$

we see that there the rank of

$$\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{3\bar{3}}\} \longrightarrow \mathbb{C}\{\varphi^{1\bar{1}\bar{3}}, \varphi^{1\bar{2}\bar{3}}, \varphi^{2\bar{1}\bar{3}}, \varphi^{2\bar{2}\bar{3}}, \varphi^{3\bar{1}\bar{2}}\},$$

is equal to 1.

3.2.5.  $\bar{\partial}_{\phi(t)} : \wedge^{0,2} \rightarrow \wedge^{0,3}$ . Recall that  $\wedge^{0,2} = \mathbb{C}\{\varphi^{\bar{1}\bar{2}}, \varphi^{\bar{1}\bar{3}}, \varphi^{\bar{2}\bar{3}}\}$  and  $\wedge^{0,3} = \mathbb{C}\{\varphi^{\bar{1}\bar{2}\bar{3}}\}$ .

$$\begin{aligned}\bar{\partial}_{\phi(t)}\varphi^{\bar{1}\bar{2}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{\bar{1}\bar{2}} = 0, \\ \bar{\partial}_{\phi(t)}\varphi^{\bar{1}\bar{3}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{\bar{1}\bar{3}} = 0, \\ \bar{\partial}_{\phi(t)}\varphi^{\bar{2}\bar{3}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{\bar{2}\bar{3}} = 0.\end{aligned}$$

We see that  $\dim \text{im } \bar{\partial}_{\phi(t)} \cap \wedge^{0,3} = 0$ .

3.2.6.  $\bar{\partial}_{\phi(t)} : \wedge^{3,0} \rightarrow \wedge^{3,1}$ . Recall that  $\wedge^{3,0} = \mathbb{C}\{\varphi^{123}\}$  and  $\wedge^{3,1} = \mathbb{C}\{\varphi^{123\bar{1}}, \varphi^{123\bar{2}}, \varphi^{123\bar{3}}\}$ .

$$\bar{\partial}_{\phi(t)}\varphi^{123} = (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{123} = -\mathcal{L}_{\phi_1}^{1,0}\varphi^{123} = 0.$$

We see that  $\dim \text{im } \bar{\partial}_{\phi(t)} \cap \wedge^{3,1} = 0$ .

3.2.7.  $\bar{\partial}_{\phi(t)} : \wedge^{2,1} \rightarrow \wedge^{2,2}$ . Recall that  $\wedge^{2,1} = \mathbb{C}\{\varphi^{ijk} \mid 1 \leq i, j, k \leq 3, i < j\}$  and  $\wedge^{2,2} = \mathbb{C}\{\varphi^{ijk\bar{l}} \mid 1 \leq i, j, k, l \leq 3, k < l\}$ .

$$\begin{aligned}\bar{\partial}_{\phi(t)}\varphi^{12\bar{1}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{12\bar{1}} = 0, \\ \bar{\partial}_{\phi(t)}\varphi^{12\bar{2}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{12\bar{2}} = 0, \\ \bar{\partial}_{\phi(t)}\varphi^{12\bar{3}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{12\bar{3}} = -\varphi^{12\bar{1}\bar{2}},\end{aligned}$$

$$\bar{\partial}_{\phi(t)}\varphi^{13\bar{1}} = (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{13\bar{1}} = -\mathcal{L}_{\phi_1}^{1,0}\varphi^{13\bar{1}} = -t_{12}\varphi^{12\bar{1}\bar{2}},$$

$$\bar{\partial}_{\phi(t)}\varphi^{13\bar{2}} = (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{13\bar{2}} = -\mathcal{L}_{\phi_1}^{1,0}\varphi^{13\bar{2}} = t_{11}\varphi^{12\bar{1}\bar{2}},$$

$$\bar{\partial}_{\phi(t)}\varphi^{23\bar{1}} = (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{23\bar{1}} = -\mathcal{L}_{\phi_1}^{1,0}\varphi^{23\bar{1}} = -t_{22}\varphi^{12\bar{1}\bar{2}},$$

$$\bar{\partial}_{\phi(t)}\varphi^{23\bar{2}} = (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{23\bar{2}} = -\mathcal{L}_{\phi_1}^{1,0}\varphi^{23\bar{2}} = t_{21}\varphi^{12\bar{1}\bar{2}}.$$

Let  $\sigma_0 = a_1\varphi^{12\bar{3}} + a_2\varphi^{13\bar{1}} + a_3\varphi^{13\bar{2}} + a_4\varphi^{23\bar{1}} + a_5\varphi^{23\bar{2}}$ , then

$$\bar{\partial}_{\phi(t)}\sigma_0 = (-a_1 - a_2t_{12} + a_3t_{11} - a_4t_{22} + a_5t_{21})\varphi^{12\bar{1}\bar{2}}.$$

Set

$$T = \begin{pmatrix} -1 & -t_{21} & t_{11} & -t_{22} & t_{21} \end{pmatrix}.$$

We see that the rank of

$$\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{12\bar{3}}, \varphi^{13\bar{1}}, \varphi^{13\bar{2}}, \varphi^{23\bar{1}}, \varphi^{23\bar{2}}\} \longrightarrow \mathbb{C}\{\varphi^{12\bar{1}\bar{2}}\}$$

is equal to  $\text{rank}(T) = 1$ . Similarly, from

$$\begin{aligned} \bar{\partial}_{\phi(t)}\varphi^{13\bar{3}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{13\bar{3}} \\ &= -\varphi^{13\bar{1}\bar{2}} + \varphi^1 \wedge [(t_{11}\varphi^2 - t_{21}\varphi^1) \wedge \varphi^{\bar{1}\bar{3}} + (t_{12}\varphi^2 - t_{22}\varphi^1) \wedge \varphi^{\bar{2}\bar{3}}] \\ &= -\varphi^{13\bar{1}\bar{2}} + t_{11}\varphi^{12\bar{1}\bar{3}} + t_{12}\varphi^{12\bar{2}\bar{3}}, \end{aligned}$$

we see that the rank of

$$\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{13\bar{3}}\} \longrightarrow \mathbb{C}\{\varphi^{13\bar{1}\bar{2}}, \varphi^{12\bar{1}\bar{3}}, \varphi^{12\bar{2}\bar{3}}\},$$

is equal to 1.

From

$$\begin{aligned} \bar{\partial}_{\phi(t)}\varphi^{23\bar{3}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{23\bar{3}} \\ &= -\varphi^{23\bar{1}\bar{2}} + \varphi^2 \wedge \left[ \sum_{\lambda=1}^2 (t_{1\lambda}\varphi^2 - t_{2\lambda}\varphi^1) \wedge \varphi^{\bar{\lambda}} \right] \wedge \varphi^{\bar{3}} \\ &= -\varphi^{23\bar{1}\bar{2}} + t_{21}\varphi^{12\bar{1}\bar{3}} + t_{22}\varphi^{12\bar{2}\bar{3}}, \end{aligned}$$

the rank of

$$\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{23\bar{3}}\} \longrightarrow \mathbb{C}\{\varphi^{23\bar{1}\bar{2}}, \varphi^{12\bar{1}\bar{3}}, \varphi^{12\bar{2}\bar{3}}\},$$

is equal to 1.

3.2.8.  $\bar{\partial}_{\phi(t)} : \wedge^{1,2} \rightarrow \wedge^{1,3}$ . Recall that  $\wedge^{1,2} = \mathbb{C}\{\varphi^{i\bar{j}k} \mid 1 \leq i, j, k \leq 3, j < k\}$  and  $\wedge^{1,3} = \mathbb{C}\{\varphi^{1\bar{1}\bar{2}\bar{3}}, \varphi^{2\bar{1}\bar{2}\bar{3}}, \varphi^{3\bar{1}\bar{2}\bar{3}}, \}$ .

$$\begin{aligned} \bar{\partial}_{\phi(t)}\varphi^{1\bar{1}\bar{2}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{1\bar{1}\bar{2}} = 0, \\ \bar{\partial}_{\phi(t)}\varphi^{1\bar{1}\bar{3}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{1\bar{1}\bar{3}} = 0, \\ \bar{\partial}_{\phi(t)}\varphi^{1\bar{2}\bar{3}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{1\bar{2}\bar{3}} = 0, \\ \bar{\partial}_{\phi(t)}\varphi^{2\bar{1}\bar{2}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{2\bar{1}\bar{2}} = 0, \\ \bar{\partial}_{\phi(t)}\varphi^{2\bar{1}\bar{3}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{2\bar{1}\bar{3}} = 0, \\ \bar{\partial}_{\phi(t)}\varphi^{2\bar{2}\bar{3}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{2\bar{2}\bar{3}} = 0, \end{aligned}$$

$$\begin{aligned}
\bar{\partial}_{\phi(t)}\varphi^{3\bar{1}\bar{2}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{3\bar{1}\bar{2}} = -\mathcal{L}_{\phi_1}^{1,0}\varphi^{3\bar{1}\bar{2}} \\
&= -\sum_{\lambda=1}^2(t_{1\lambda}\varphi^2 - t_{2\lambda}\varphi^1) \wedge \varphi^{\bar{\lambda}} \wedge \varphi^{\bar{1}\bar{2}} = 0, \\
\bar{\partial}_{\phi(t)}\varphi^{3\bar{1}\bar{3}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{3\bar{1}\bar{3}} \\
&= -\sum_{\lambda=1}^2(t_{1\lambda}\varphi^2 - t_{2\lambda}\varphi^1) \wedge \varphi^{\bar{\lambda}} \wedge \varphi^{\bar{1}\bar{3}} \\
&= t_{12}\varphi^{2\bar{1}\bar{2}\bar{3}} - t_{22}\varphi^{1\bar{1}\bar{2}\bar{3}}, \\
\bar{\partial}_{\phi(t)}\varphi^{3\bar{2}\bar{3}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{3\bar{2}\bar{3}} \\
&= -\sum_{\lambda=1}^2(t_{1\lambda}\varphi^2 - t_{2\lambda}\varphi^1) \wedge \varphi^{\bar{\lambda}} \wedge \varphi^{\bar{2}\bar{3}} \\
&= t_{21}\varphi^{1\bar{1}\bar{2}\bar{3}} - t_{11}\varphi^{2\bar{1}\bar{2}\bar{3}}.
\end{aligned}$$

Let  $\sigma_0 = a_1\varphi^{3\bar{1}\bar{3}} + a_2\varphi^{3\bar{2}\bar{3}}$ , then

$$\begin{aligned}
\bar{\partial}_{\phi(t)}\sigma_0 &= a_1(t_{12}\varphi^{2\bar{1}\bar{2}\bar{3}} - t_{22}\varphi^{1\bar{1}\bar{2}\bar{3}}) + a_2(t_{21}\varphi^{1\bar{1}\bar{2}\bar{3}} - t_{11}\varphi^{2\bar{1}\bar{2}\bar{3}}) \\
&= (-a_1t_{22} - a_2t_{21})\varphi^{1\bar{1}\bar{2}\bar{3}} + (a_1t_{12} - a_2t_{11})\varphi^{2\bar{1}\bar{2}\bar{3}}.
\end{aligned}$$

Set

$$T = \begin{pmatrix} -t_{22} & t_{21} \\ t_{12} & -t_{11} \end{pmatrix},$$

then

- (1) When  $(t_{11}, t_{12}, t_{21}, t_{22}) = 0$ , we have  $\text{rank}(T) = 0$  and the rank of  $\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{3\bar{1}\bar{3}}, \varphi^{3\bar{2}\bar{3}}\} \rightarrow \mathbb{C}\{\varphi^{1\bar{1}\bar{2}\bar{3}}, \varphi^{2\bar{1}\bar{2}\bar{3}}\}$  is 0.
- (2) When  $(t_{11}, t_{12}, t_{21}, t_{22}) \neq 0, D(t) = 0$ , we have  $\text{rank}(T) = 1$ , and the rank of  $\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{3\bar{1}\bar{3}}, \varphi^{3\bar{2}\bar{3}}\} \rightarrow \mathbb{C}\{\varphi^{1\bar{1}\bar{2}\bar{3}}, \varphi^{2\bar{1}\bar{2}\bar{3}}\}$  is 1.
- (3) When  $(t_{11}, t_{12}, t_{21}, t_{22}) \neq 0, D(t) \neq 0$ , we have  $\text{rank}(T) = 2$ , and the rank of  $\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{3\bar{1}\bar{3}}, \varphi^{3\bar{2}\bar{3}}\} \rightarrow \mathbb{C}\{\varphi^{1\bar{1}\bar{2}\bar{3}}, \varphi^{2\bar{1}\bar{2}\bar{3}}\}$  is 2.

3.2.9.  $\bar{\partial}_{\phi(t)} : \wedge^{3,1} \rightarrow \wedge^{3,2}$ . Recall that  $\wedge^{3,1} = \mathbb{C}\{\varphi^{123\bar{1}}, \varphi^{123\bar{2}}, \varphi^{123\bar{3}}\}$  and  $\wedge^{3,2} = \mathbb{C}\{\varphi^{123\bar{1}\bar{2}}, \varphi^{123\bar{1}\bar{3}}, \varphi^{123\bar{2}\bar{3}}\}$ .

$$\begin{aligned}
\bar{\partial}_{\phi(t)}\varphi^{123\bar{1}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{123\bar{1}} = -\mathcal{L}_{\phi_1}^{1,0}\varphi^{123\bar{1}} \\
&= -\varphi^{12} \wedge \left[ \sum_{\lambda=1}^2(t_{1\lambda}\varphi^2 - t_{2\lambda}\varphi^1) \wedge \varphi^{\bar{\lambda}} \right] \wedge \varphi^{\bar{1}} = 0,
\end{aligned}$$

$$\begin{aligned}
\bar{\partial}_{\phi(t)}\varphi^{123\bar{2}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{123\bar{2}} = -\mathcal{L}_{\phi_1}^{1,0}\varphi^{123\bar{2}} \\
&= -\varphi^{12} \wedge \left[ \sum_{\lambda=1}^2 (t_{1\lambda}\varphi^2 - t_{2\lambda}\varphi^1) \wedge \varphi^{\bar{\lambda}} \right] \wedge \varphi^{\bar{2}} = 0, \\
\bar{\partial}_{\phi(t)}\varphi^{123\bar{3}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{123\bar{3}} \\
&= \varphi^{123\bar{1}\bar{2}} - \varphi^{12} \wedge \left[ \sum_{\lambda=1}^2 (t_{1\lambda}\varphi^2 - t_{2\lambda}\varphi^1) \wedge \varphi^{\bar{\lambda}} \right] \wedge \varphi^{\bar{3}} \\
&= \varphi^{123\bar{1}\bar{2}}.
\end{aligned}$$

We see that the rank of  $\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{123\bar{3}}\} \rightarrow \mathbb{C}\{\varphi^{123\bar{1}\bar{2}}\}$  is 1.

3.2.10.  $\bar{\partial}_{\phi(t)} : \wedge^{2,2} \rightarrow \wedge^{2,3}$ . Recall that  $\wedge^{2,2} = \mathbb{C}\{\varphi^{ijk\bar{l}} \mid 1 \leq i, j, k \leq 3, i < j, k < l\}$  and  $\wedge^{2,3} = \mathbb{C}\{\varphi^{12\bar{1}\bar{2}\bar{3}}, \varphi^{13\bar{1}\bar{2}\bar{3}}, \varphi^{23\bar{1}\bar{2}\bar{3}}\}$ .

$$\begin{aligned}
\bar{\partial}_{\phi(t)}\varphi^{12\bar{1}\bar{2}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{12\bar{1}\bar{2}} = 0, \\
\bar{\partial}_{\phi(t)}\varphi^{12\bar{1}\bar{3}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{12\bar{1}\bar{3}} = 0, \\
\bar{\partial}_{\phi(t)}\varphi^{12\bar{2}\bar{3}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{12\bar{2}\bar{3}} = 0, \\
\bar{\partial}_{\phi(t)}\varphi^{13\bar{1}\bar{2}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{13\bar{1}\bar{2}} = -\mathcal{L}_{\phi_1}^{1,0}\varphi^{13\bar{1}\bar{2}} \\
&= \varphi^1 \wedge \left[ \sum_{\lambda=1}^2 (t_{1\lambda}\varphi^2 - t_{2\lambda}\varphi^1) \wedge \varphi^{\bar{\lambda}} \right] \wedge \varphi^{\bar{1}\bar{2}} = 0, \\
\bar{\partial}_{\phi(t)}\varphi^{23\bar{1}\bar{2}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{23\bar{1}\bar{2}} = -\mathcal{L}_{\phi_1}^{1,0}\varphi^{23\bar{1}\bar{2}} \\
&= \varphi^2 \wedge \left[ \sum_{\lambda=1}^2 (t_{1\lambda}\varphi^2 - t_{2\lambda}\varphi^1) \wedge \varphi^{\bar{\lambda}} \right] \wedge \varphi^{\bar{1}\bar{2}} = 0,
\end{aligned}$$

$$\begin{aligned}
\bar{\partial}_{\phi(t)}\varphi^{13\bar{1}\bar{3}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{13\bar{1}\bar{3}} \\
&= \varphi^1 \wedge \left[ \sum_{\lambda=1}^2 (t_{1\lambda}\varphi^2 - t_{2\lambda}\varphi^1 \wedge \varphi^{\bar{\lambda}}) \wedge \varphi^{\bar{1}\bar{3}} \right] \\
&= -t_{12}\varphi^{12\bar{1}\bar{2}\bar{3}}, \\
\bar{\partial}_{\phi(t)}\varphi^{13\bar{2}\bar{3}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{13\bar{2}\bar{3}} \\
&= \varphi^1 \wedge \left[ \sum_{\lambda=1}^2 (t_{1\lambda}\varphi^2 - t_{2\lambda}\varphi^1 \wedge \varphi^{\bar{\lambda}}) \wedge \varphi^{\bar{2}\bar{3}} \right] \\
&= t_{11}\varphi^{12\bar{1}\bar{2}\bar{3}},
\end{aligned}$$

$$\begin{aligned}
\bar{\partial}_{\phi(t)}\varphi^{23\bar{1}\bar{3}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{23\bar{1}\bar{3}} = -\mathcal{L}_{\phi_1}^{1,0}\varphi^{23\bar{1}\bar{3}} \\
&= \varphi^2 \wedge \left[ \sum_{\lambda=1}^2 (t_{1\lambda}\varphi^2 - t_{2\lambda}\varphi^1) \wedge \varphi^{\bar{\lambda}} \right] \wedge \varphi^{\bar{1}\bar{3}} \\
&= -t_{22}\varphi^{12\bar{1}\bar{2}\bar{3}}, \\
\bar{\partial}_{\phi(t)}\varphi^{23\bar{2}\bar{3}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{23\bar{2}\bar{3}} = -\mathcal{L}_{\phi_1}^{1,0}\varphi^{23\bar{2}\bar{3}} \\
&= \varphi^2 \wedge \left[ \sum_{\lambda=1}^2 (t_{1\lambda}\varphi^2 - t_{2\lambda}\varphi^1) \wedge \varphi^{\bar{\lambda}} \right] \wedge \varphi^{\bar{2}\bar{3}} \\
&= t_{21}\varphi^{12\bar{1}\bar{2}\bar{3}}.
\end{aligned}$$

Let  $\sigma_0 = a_1\varphi^{13\bar{1}\bar{3}} + a_2\varphi^{13\bar{2}\bar{3}} + a_3\varphi^{23\bar{1}\bar{3}} + a_4\varphi^{23\bar{2}\bar{3}}$ , then

$$\begin{aligned}
\bar{\partial}_{\phi(t)}\sigma_0 &= -a_1t_{12}\varphi^{12\bar{1}\bar{2}\bar{3}} + a_2t_{11}\varphi^{12\bar{1}\bar{2}\bar{3}} - a_3t_{22}\varphi^{12\bar{1}\bar{2}\bar{3}} + a_4a_1t_{21}\varphi^{12\bar{1}\bar{2}\bar{3}} \\
&= (-a_1t_{12} + a_2t_{11} - a_3t_{22} + a_4t_{21})\varphi^{12\bar{1}\bar{2}\bar{3}}.
\end{aligned}$$

Set

$$T = \begin{pmatrix} -t_{12} & t_{11} & -t_{22} & t_{21} \end{pmatrix},$$

then

- (1) When  $(t_{11}, t_{12}, t_{21}, t_{22}) = 0$ , we have  $\text{rank}(T) = 0$  and the rank of  $\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{13\bar{1}\bar{3}}, \varphi^{13\bar{2}\bar{3}}, \varphi^{23\bar{1}\bar{3}}, \varphi^{23\bar{2}\bar{3}}\} \rightarrow \mathbb{C}\{\varphi^{12\bar{1}\bar{2}\bar{3}}\}$  is 0.
- (2) When  $(t_{11}, t_{12}, t_{21}, t_{22}) \neq 0, D(t) = 0$ , we have  $\text{rank}(T) = 1$  and the rank of  $\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{13\bar{1}\bar{3}}, \varphi^{13\bar{2}\bar{3}}, \varphi^{23\bar{1}\bar{3}}, \varphi^{23\bar{2}\bar{3}}\} \rightarrow \mathbb{C}\{\varphi^{12\bar{1}\bar{2}\bar{3}}\}$  is 1.
- (3) When  $(t_{11}, t_{12}, t_{21}, t_{22}) \neq 0, D(t) \neq 0$ , we have  $\text{rank}(T) = 1$  and the rank of  $\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{13\bar{1}\bar{3}}, \varphi^{13\bar{2}\bar{3}}, \varphi^{23\bar{1}\bar{3}}, \varphi^{23\bar{2}\bar{3}}\} \rightarrow \mathbb{C}\{\varphi^{12\bar{1}\bar{2}\bar{3}}\}$  is 1.


3.2.11.  $\bar{\partial}_{\phi(t)} : \wedge^{3,2} \rightarrow \wedge^{3,3}$ . Recall that  $\wedge^{3,2} = \mathbb{C}\{\varphi^{123\bar{1}\bar{2}}, \varphi^{123\bar{1}\bar{3}}, \varphi^{123\bar{2}\bar{3}}\}$  and  $\wedge^{3,3} = \mathbb{C}\{\varphi^{123\bar{1}\bar{2}\bar{3}}\}$ .

$$\begin{aligned}
\bar{\partial}_{\phi(t)}\varphi^{123\bar{1}\bar{2}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{123\bar{1}\bar{2}} = -\mathcal{L}_{\phi_1}^{1,0}\varphi^{123\bar{1}\bar{2}} \\
&= -\varphi^{12} \wedge \left[ \sum_{\lambda=1}^2 (t_{1\lambda}\varphi^2 - t_{2\lambda}\varphi^1) \wedge \varphi^{\bar{\lambda}} \right] \wedge \varphi^{\bar{1}\bar{2}} = 0, \\
\bar{\partial}_{\phi(t)}\varphi^{123\bar{1}\bar{3}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{123\bar{1}\bar{3}} \\
&= -\varphi^{12} \wedge \left[ \sum_{\lambda=1}^2 (t_{1\lambda}\varphi^2 - t_{2\lambda}\varphi^1) \wedge \varphi^{\bar{\lambda}} \right] \wedge \varphi^{\bar{1}\bar{3}} = 0, \\
\bar{\partial}_{\phi(t)}\varphi^{123\bar{2}\bar{3}} &= (\bar{\partial} - \mathcal{L}_{\phi_1}^{1,0})\varphi^{123\bar{2}\bar{3}} \\
&= -\varphi^{12} \wedge \left[ \sum_{\lambda=1}^2 (t_{1\lambda}\varphi^2 - t_{2\lambda}\varphi^1) \wedge \varphi^{\bar{\lambda}} \right] \wedge \varphi^{\bar{2}\bar{3}} = 0.
\end{aligned}$$

We see that  $\dim \text{im } \bar{\partial}_{\phi(t)} \cap \wedge^{3,3} = 0$ .

#### 4. THE DECOMPOSITION OF $(\wedge^\bullet, d_{\phi(t)} = \partial + \bar{\partial}_{\phi(t)})$

According to Theorem 2.1, we may decompose the double complex  $(\wedge^\bullet, d_{\phi(t)} = \partial + \bar{\partial}_{\phi(t)})$  as direct sums of squares and zigzags. To realize this decomposition, we need to choose suitable basis of each components  $\wedge^{p,q}$  so that they are compatible with the actions of differentials  $\partial, \bar{\partial}_{\phi(t)}$ . This means that if  $\varphi_1, \dots, \varphi_k$  is chosen as a basis for  $\wedge^{p,q}$ , then  $\bar{\partial}_{\phi(t)}\varphi_1, \dots, \bar{\partial}_{\phi(t)}\varphi_k$  should be contained in the basis we have chosen for  $\wedge^{p,q+1}$  and  $\partial\varphi_1, \dots, \partial\varphi_k$  should also be contained in the basis we have chosen for  $\wedge^{p+1,q}$ . For  $t = 0$ , this has already done by Angella in [Ang13] and the result is pictured as follows:



where signs in several places are omitted (e.g.  $\varphi^{1\bar{2}}$ ) because we are only interested in the double complex structure. The same convention applies in latter subsections so long as this does not cause any trouble.

#### 4.1. The case that $(t_{11}, t_{12}, t_{21}, t_{22}) \neq 0, D(t) = 0$ .

**Proposition 4.1.** *There exist the following basis*

- $\wedge^{1,0} = \mathbb{C}\{\underbrace{\varphi^1, \varphi^2, \varphi^3}_{\ker \bar{\partial}_{\phi(t)}}\};$
- $\wedge^{2,0} = \mathbb{C}\{\underbrace{\xi_1^{2,0}, \varphi^{1\bar{3}}, \xi_2^{2,0}}_{\ker \bar{\partial}_{\phi(t)}}\};$
- $\wedge^{1,1} = \mathbb{C}\{\underbrace{\bar{\partial}_{\phi(t)}\varphi^3, \xi_1^{1,1}, \xi_2^{1,1}, \xi_3^{1,1}, \xi_4^{1,1}, \xi_5^{1,1}}_{\ker \bar{\partial}_{\phi(t)}}, \varphi^{1\bar{3}}, \varphi^{2\bar{3}}, \varphi^{3\bar{3}}\};$
- $\wedge^{2,1} = \mathbb{C}\{\underbrace{\partial\xi_4^{1,1}, \partial\xi_5^{1,1}, \xi_1^{2,1}, \xi_2^{2,1}, \xi_3^{2,1}, \xi_4^{2,1}}_{\ker \bar{\partial}_{\phi(t)}}, \varphi^{1\bar{2}\bar{3}}, \varphi^{1\bar{3}\bar{3}}, \varphi^{2\bar{3}\bar{3}}\};$



- $\Lambda^{1,2} = \mathbb{C}\{\underbrace{\bar{\partial}_{\phi(t)}\varphi^{3\bar{3}}, \varphi^{1\bar{1}\bar{2}}, \varphi^{2\bar{1}\bar{2}}, \xi_1^{1,2}, \xi_2^{1,2}, \xi_3^{1,2}, \xi_4^{1,2}, \xi_5^{1,2}, \xi_6^{1,2}}_{\ker \bar{\partial}_{\phi(t)}}\};$
- $\Lambda^{2,2} = \mathbb{C}\{\underbrace{\bar{\partial}_{\phi(t)}\varphi^{13\bar{3}}, \bar{\partial}_{\phi(t)}\varphi^{23\bar{3}}, \varphi^{12\bar{1}\bar{2}}, \partial\xi_5^{1,2}, \partial\xi_6^{1,2}, \xi_1^{2,2}, \xi_2^{2,2}, \xi_3^{2,2}, \xi_4^{2,2}}_{\ker \bar{\partial}_{\phi(t)}}\};$
- $\Lambda^{1,3} = \mathbb{C}\{\bar{\partial}_{\phi(t)}\xi_6^{1,2}, \xi_1^{1,3}, \varphi^{3\bar{1}\bar{2}\bar{3}}\},$

such that

- $\mathbb{C}\{\xi_1^{2,0}, \xi_2^{2,0}\} = \mathbb{C}\{\varphi^{13}, \varphi^{23}\};$
- $\mathbb{C}\{\xi_1^{1,1}, \xi_2^{1,1}, \xi_3^{1,1}, \bar{\partial}_{\phi(t)}\varphi^3\} = \mathbb{C}\{\varphi^{1\bar{1}}, \varphi^{1\bar{2}}, \varphi^{2\bar{1}}, \varphi^{2\bar{2}}\}$  and
 
$$\mathbb{C}\{\xi_4^{1,1}, \xi_5^{1,1}\} = \mathbb{C}\{\varphi^{3\bar{1}}, \varphi^{3\bar{2}}\}, \quad \partial\xi_4^{1,1} = \bar{\partial}_{\phi(t)}\xi_2^{2,0};$$
- $\mathbb{C}\{\varphi^{12\bar{3}}, \xi_1^{2,1}, \xi_2^{2,1}, \xi_3^{2,1}, \xi_4^{2,1}\} = \mathbb{C}\{\varphi^{12\bar{3}}, \varphi^{13\bar{1}}, \varphi^{13\bar{2}}, \varphi^{23\bar{1}}, \varphi^{23\bar{2}}\};$
- $\mathbb{C}\{\xi_1^{1,2}, \xi_2^{1,2}, \xi_3^{1,2}, \xi_4^{1,2}, \bar{\partial}_{\phi(t)}\varphi^{3\bar{3}}\} = \mathbb{C}\{\varphi^{1\bar{1}\bar{3}}, \varphi^{1\bar{2}\bar{3}}, \varphi^{2\bar{1}\bar{3}}, \varphi^{2\bar{2}\bar{3}}, \varphi^{3\bar{1}\bar{2}}\}$  and
 
$$\mathbb{C}\{\xi_5^{1,2}, \xi_6^{1,2}\} = \mathbb{C}\{\varphi^{3\bar{1}\bar{3}}, \varphi^{3\bar{2}\bar{3}}\}, \quad \text{with } \bar{\partial}_{\phi(t)}\xi_6^{1,2} \neq 0;$$
- $\mathbb{C}\{\partial\xi_5^{1,2}, \partial\xi_6^{1,2}, \bar{\partial}_{\phi(t)}\varphi^{13\bar{3}}, \bar{\partial}_{\phi(t)}\varphi^{23\bar{3}}\} = \mathbb{C}\{\varphi^{12\bar{1}\bar{3}}, \varphi^{12\bar{2}\bar{3}}, \varphi^{13\bar{1}\bar{2}}, \varphi^{23\bar{1}\bar{2}}\};$
- $\mathbb{C}\{\xi_1^{1,3}, \bar{\partial}_{\phi(t)}\xi_6^{1,2}\} = \mathbb{C}\{\varphi^{1\bar{1}\bar{2}\bar{3}}, \varphi^{2\bar{1}\bar{2}\bar{3}}\}.$

*Proof.* From Subsection 3.2.1, we know that  $\ker \bar{\partial}_{\phi(t)} \cap \Lambda^{1,0} = \mathbb{C}\{\varphi^1, \varphi^2\}$  and the rank of  $\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^3\} \rightarrow \mathbb{C}\{\varphi^{1\bar{1}}, \varphi^{2\bar{1}}, \varphi^{1\bar{2}}, \varphi^{2\bar{2}}\}$  is 1.

From Subsection 3.2.3, we know that  $\dim \ker \bar{\partial}_{\phi(t)} \cap \Lambda^{2,0} = 2$  and  $\bar{\partial}_{\phi(t)}\varphi^{12} = 0$ . On the other hand, the rank of  $\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{13}, \varphi^{23}\} \rightarrow \mathbb{C}\{\varphi^{12\bar{1}}, \varphi^{12\bar{2}}\}$  is 1. So we may choose  $\xi_1^{2,0}, \xi_2^{2,0} \in \Lambda^{2,0}$  such that  $\mathbb{C}\{\xi_1^{2,0}, \xi_2^{2,0}\} = \mathbb{C}\{\varphi^{13}, \varphi^{23}\}$  and

$$\bar{\partial}_{\phi(t)}\xi_1^{2,0} = 0, \quad \bar{\partial}_{\phi(t)}\xi_2^{2,0} \neq 0.$$

Assume  $\xi_2^{2,0} = a\varphi^{13} + b\varphi^{23}$ , then

$$\begin{aligned} \bar{\partial}_{\phi(t)}\xi_2^{2,0} &= (at_{11} + bt_{21})\varphi^{12\bar{1}} + (at_{12} + bt_{22})\varphi^{12\bar{2}} \\ &= \partial[-(at_{11} + bt_{21})\varphi^{3\bar{1}} - (at_{12} + bt_{22})\varphi^{3\bar{2}}]. \end{aligned}$$

Set  $\xi_4^{1,1} = -(at_{11} + bt_{21})\varphi^{3\bar{1}} - (at_{12} + bt_{22})\varphi^{3\bar{2}}$ , then  $\partial\xi_4^{1,1} = \bar{\partial}_{\phi(t)}\xi_2^{2,0}$ . We may choose  $\xi_5^{1,1} \in \Lambda^{1,1}$  so that  $\mathbb{C}\{\xi_4^{1,1}, \xi_5^{1,1}\} = \mathbb{C}\{\varphi^{3\bar{1}}, \varphi^{3\bar{2}}\}$ . In particular,  $\mathbb{C}\{\partial\xi_4^{1,1}, \partial\xi_5^{1,1}\} = \mathbb{C}\{\varphi^{12\bar{1}}, \varphi^{12\bar{2}}\}$ .

From Subsection 3.2.4, we know that  $\dim \ker \bar{\partial}_{\phi(t)} \cap \Lambda^{1,1} = 6$  and

$$\ker \bar{\partial}_{\phi(t)} = \mathbb{C}\{\varphi^{1\bar{1}}, \varphi^{1\bar{2}}, \varphi^{2\bar{1}}, \varphi^{2\bar{2}}, \xi_4^{1,1}, \xi_5^{1,1}\}.$$

On the other hand,  $\bar{\partial}_{\phi(t)}\varphi^{13} = \varphi^{1\bar{1}\bar{2}}, \bar{\partial}_{\phi(t)}\varphi^{23} = \varphi^{2\bar{1}\bar{2}}$  and the rank of  $\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{3\bar{3}}\} \rightarrow \mathbb{C}\{\varphi^{1\bar{1}\bar{3}}, \varphi^{1\bar{2}\bar{3}}, \varphi^{2\bar{1}\bar{3}}, \varphi^{2\bar{2}\bar{3}}, \varphi^{3\bar{1}\bar{2}}\}$  is 1. Since  $0 \neq \bar{\partial}_{\phi(t)}\varphi^3 \in \mathbb{C}\{\varphi^{1\bar{1}}, \varphi^{1\bar{2}}, \varphi^{2\bar{1}}, \varphi^{2\bar{2}}\}$ , we may choose  $\xi_1^{1,1}, \xi_2^{1,1}, \xi_3^{1,1}$  such that  $\mathbb{C}\{\xi_1^{1,1}, \xi_2^{1,1}, \xi_3^{1,1}, \bar{\partial}_{\phi(t)}\varphi^3\} = \mathbb{C}\{\varphi^{1\bar{1}}, \varphi^{1\bar{2}}, \varphi^{2\bar{1}}, \varphi^{2\bar{2}}\}$ . Notice that

$$\partial\bar{\partial}_{\phi(t)}\varphi^{3\bar{3}} = -\varphi^{12\bar{1}\bar{2}} = -\bar{\partial}_{\phi(t)}\varphi^{12\bar{3}} = \bar{\partial}_{\phi(t)}\partial\varphi^{3\bar{3}},$$

From Subsection 3.2.7, we know that  $\mathbb{C}\{\partial\xi_4^{1,1}, \partial\xi_5^{1,1}\} = \mathbb{C}\{\varphi^{12\bar{1}}, \varphi^{12\bar{2}}\} \subset \ker \bar{\partial}_{\phi(t)}$ , the rank of  $\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{12\bar{3}}, \varphi^{13\bar{1}}, \varphi^{13\bar{2}}, \varphi^{23\bar{1}}, \varphi^{23\bar{2}}\} \rightarrow \mathbb{C}\{\varphi^{12\bar{1}\bar{2}}\}$  is 1,  $\bar{\partial}_{\phi(t)}\varphi^{13\bar{3}}$  and  $\bar{\partial}_{\phi(t)}\varphi^{23\bar{3}}$  are linearly independent. We may choose  $\xi_1^{2,1}, \xi_2^{2,1}, \xi_3^{2,1}, \xi_4^{2,1} \in \ker \bar{\partial}_{\phi(t)}$  so that

$$\mathbb{C}\{\varphi^{12\bar{3}}, \xi_1^{2,1}, \xi_2^{2,1}, \xi_3^{2,1}, \xi_4^{2,1}\} = \mathbb{C}\{\varphi^{12\bar{3}}, \varphi^{13\bar{1}}, \varphi^{13\bar{2}}, \varphi^{23\bar{1}}, \varphi^{23\bar{2}}\}.$$

For  $\wedge^{1,2}$ , we already see that  $\varphi^{1\bar{1}\bar{2}} = \bar{\partial}_{\phi(t)}\varphi^{1\bar{3}}, \varphi^{2\bar{1}\bar{2}} = \bar{\partial}_{\phi(t)}\varphi^{2\bar{3}}$  and the rank of  $\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{3\bar{3}}\} \rightarrow \mathbb{C}\{\varphi^{1\bar{1}\bar{3}}, \varphi^{1\bar{2}\bar{3}}, \varphi^{2\bar{1}\bar{3}}, \varphi^{2\bar{2}\bar{3}}, \varphi^{3\bar{1}\bar{2}}\}$  is 1. From Subsection 3.2.8, we know that  $\varphi^{1\bar{1}\bar{2}}, \varphi^{1\bar{1}\bar{3}}, \varphi^{1\bar{2}\bar{3}}, \varphi^{2\bar{1}\bar{2}}, \varphi^{2\bar{1}\bar{3}}, \varphi^{2\bar{2}\bar{3}}, \varphi^{3\bar{1}\bar{2}}$  are in  $\ker \bar{\partial}_{\phi(t)}$  and the rank of  $\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{3\bar{1}\bar{3}}, \varphi^{3\bar{2}\bar{3}}\} \rightarrow \mathbb{C}\{\varphi^{1\bar{1}\bar{2}\bar{3}}, \varphi^{2\bar{1}\bar{2}\bar{3}}\}$  is 1. So we may choose  $\xi_1^{1,2}, \dots, \xi_6^{1,2}$  so that  $\mathbb{C}\{\xi_1^{1,2}, \xi_2^{1,2}, \xi_3^{1,2}, \xi_4^{1,2}, \bar{\partial}_{\phi(t)}\varphi^{3\bar{3}}\} = \mathbb{C}\{\varphi^{1\bar{1}\bar{3}}, \varphi^{1\bar{2}\bar{3}}, \varphi^{2\bar{1}\bar{3}}, \varphi^{2\bar{2}\bar{3}}, \varphi^{3\bar{1}\bar{2}}\}$  and

$$\mathbb{C}\{\xi_5^{1,2}, \xi_6^{1,2}\} = \mathbb{C}\{\varphi^{3\bar{1}\bar{3}}, \varphi^{3\bar{2}\bar{3}}\}, \text{ with } \bar{\partial}_{\phi(t)}\xi_5^{1,2} = 0, \bar{\partial}_{\phi(t)}\xi_6^{1,2} \neq 0.$$

For  $\wedge^{2,2}$ , we already see that

$$\bar{\partial}_{\phi(t)}\varphi^{13\bar{3}}, \bar{\partial}_{\phi(t)}\varphi^{23\bar{3}} \in \mathbb{C}\{\varphi^{13\bar{1}\bar{2}}, \varphi^{12\bar{1}\bar{3}}, \varphi^{12\bar{2}\bar{3}}, \varphi^{23\bar{1}\bar{2}}\},$$

are linearly independent. From Subsection 3.2.10, we know that  $\varphi^{12\bar{1}\bar{2}}, \varphi^{12\bar{1}\bar{3}}, \varphi^{12\bar{2}\bar{3}}, \varphi^{13\bar{1}\bar{2}}, \varphi^{23\bar{1}\bar{2}} \in \ker \bar{\partial}_{\phi(t)}$  and the rank of  $\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{13\bar{1}\bar{3}}, \varphi^{13\bar{2}\bar{3}}, \varphi^{23\bar{1}\bar{3}}, \varphi^{23\bar{2}\bar{3}}\} \rightarrow \mathbb{C}\{\varphi^{12\bar{1}\bar{2}\bar{3}}\}$  is 1. Notice that  $\mathbb{C}\{\varphi^{12\bar{1}\bar{3}}, \varphi^{12\bar{2}\bar{3}}\} = \mathbb{C}\{\partial\xi_5^{1,2}, \partial\xi_6^{1,2}\}$  and

$$\mathbb{C}\{\bar{\partial}_{\phi(t)}\varphi^{13\bar{3}}, \bar{\partial}_{\phi(t)}\varphi^{23\bar{3}}, \varphi^{12\bar{1}\bar{2}}, \varphi^{12\bar{1}\bar{3}}, \varphi^{12\bar{2}\bar{3}}\} = \mathbb{C}\{\varphi^{12\bar{1}\bar{2}}, \varphi^{12\bar{1}\bar{3}}, \varphi^{12\bar{2}\bar{3}}, \varphi^{13\bar{1}\bar{2}}, \varphi^{23\bar{1}\bar{2}}\} \subset \ker \bar{\partial}_{\phi(t)}.$$

So

$$\mathbb{C}\{\bar{\partial}_{\phi(t)}\varphi^{13\bar{3}}, \bar{\partial}_{\phi(t)}\varphi^{23\bar{3}}, \varphi^{12\bar{1}\bar{2}}, \partial\xi_5^{1,2}, \partial\xi_6^{1,2}\} = \mathbb{C}\{\varphi^{12\bar{1}\bar{2}}, \varphi^{12\bar{1}\bar{3}}, \varphi^{12\bar{2}\bar{3}}, \varphi^{13\bar{1}\bar{2}}, \varphi^{23\bar{1}\bar{2}}\} \subset \ker \bar{\partial}_{\phi(t)}.$$

Meanwhile, because the rank of  $\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{13\bar{1}\bar{3}}, \varphi^{13\bar{2}\bar{3}}, \varphi^{23\bar{1}\bar{3}}, \varphi^{23\bar{2}\bar{3}}\} \rightarrow \mathbb{C}\{\varphi^{12\bar{1}\bar{2}\bar{3}}\}$  is 1, we may choose  $\xi_1^{2,2}, \xi_2^{2,2}, \xi_3^{2,2}, \xi_4^{2,2} \in \ker \bar{\partial}_{\phi(t)}$  so that


$$\mathbb{C}\{\xi_1^{2,2}, \xi_2^{2,2}, \xi_3^{2,2}, \xi_4^{2,2}\} = \mathbb{C}\{\varphi^{13\bar{1}\bar{3}}, \varphi^{13\bar{2}\bar{3}}, \varphi^{23\bar{1}\bar{3}}, \varphi^{23\bar{2}\bar{3}}\} \text{ and } \bar{\partial}_{\phi(t)}\xi_4^{2,2} = \varphi^{12\bar{1}\bar{2}\bar{3}}.$$

For  $\wedge^{1,3}$ , we already see that  $\bar{\partial}_{\phi(t)}\xi_6^{1,2} \in \mathbb{C}\{\varphi^{1\bar{1}\bar{2}\bar{3}}, \varphi^{2\bar{1}\bar{2}\bar{3}}\}$  is nonzero, therefore we may choose  $\xi_1^{1,3}$  so that

$$\mathbb{C}\{\bar{\partial}_{\phi(t)}\xi_6^{1,2}, \xi_1^{1,3}\} = \mathbb{C}\{\varphi^{1\bar{1}\bar{2}\bar{3}}, \varphi^{2\bar{1}\bar{2}\bar{3}}\}.$$

□

By using Proposition 4.1, we immediately get the following diagrammatic view of the structure of  $(\wedge^\bullet, d_{\phi(t)} = \partial + \bar{\partial}_{\phi(t)})$  for  $(t_{11}, t_{12}, t_{21}, t_{22}) \neq 0, D(t) = 0$ :



4.2. The case that  $(t_{11}, t_{12}, t_{21}, t_{22}) \neq 0, D(t) \neq 0$ .

**Proposition 4.2.** *There exist the following basis*

- $\wedge^{1,0} = \mathbb{C}\{\underbrace{\varphi^1, \varphi^2, \varphi^3}_{\ker \bar{\partial}_{\phi(t)}}\};$
- $\wedge^{2,0} = \mathbb{C}\{\underbrace{\varphi^{12}, \varphi^{13}, \varphi^{23}}_{\ker \bar{\partial}_{\phi(t)}}\};$
- $\wedge^{1,1} = \mathbb{C}\{\underbrace{\bar{\partial}_{\phi(t)}\varphi^3, \xi_1^{1,1}, \xi_2^{1,1}, \xi_3^{1,1}, \eta_1^{1,1}, \eta_2^{1,1}}_{\ker \bar{\partial}_{\phi(t)}}, \varphi^{1\overline{3}}, \varphi^{2\overline{3}}, \varphi^{3\overline{3}}\};$

- $\wedge^{2,1} = \mathbb{C}\{\underbrace{\partial\eta_1^{1,1} = \bar{\partial}_{\phi(t)}\varphi^{23}, \partial\eta_2^{1,1} = \bar{\partial}_{\phi(t)}\varphi^{13}, \xi_2^{2,1}, \xi_3^{2,1}, \xi_4^{2,1}}_{\ker \bar{\partial}_{\phi(t)}}, \varphi^{12\bar{3}}, \varphi^{13\bar{3}}, \varphi^{23\bar{3}}\};$
- $\wedge^{1,2} = \mathbb{C}\{\underbrace{\bar{\partial}_{\phi(t)}\varphi^{3\bar{3}}, \varphi^{1\bar{1}\bar{2}}, \varphi^{2\bar{1}\bar{2}}, \xi_1^{1,2}, \xi_2^{1,2}, \xi_3^{1,2}, \xi_4^{1,2}}_{\ker \bar{\partial}_{\phi(t)}}, \varphi^{3\bar{1}\bar{3}}, \varphi^{3\bar{2}\bar{3}}\};$
- $\wedge^{2,2} = \mathbb{C}\{\underbrace{\bar{\partial}_{\phi(t)}\varphi^{13\bar{3}}, \bar{\partial}_{\phi(t)}\varphi^{23\bar{3}}, \varphi^{12\bar{1}\bar{2}}, \varphi^{12\bar{1}\bar{3}}, \varphi^{12\bar{2}\bar{3}}, \xi_1^{2,2}, \xi_2^{2,2}, \xi_3^{2,2}, \xi_4^{2,2}}_{\ker \bar{\partial}_{\phi(t)}}\};$
- $\wedge^{1,3} = \mathbb{C}\{\underbrace{\bar{\partial}_{\phi(t)}\varphi^{3\bar{1}\bar{3}}, \bar{\partial}_{\phi(t)}\varphi^{3\bar{2}\bar{3}}}_{\in \ker \bar{\partial}_{\phi(t)}}, \varphi^{3\bar{1}\bar{2}\bar{3}}\},$

such that

- $\mathbb{C}\{\xi_1^{1,1}, \xi_2^{1,1}, \xi_3^{1,1}, \bar{\partial}_{\phi(t)}\varphi^3\} = \mathbb{C}\{\varphi^{1\bar{1}}, \varphi^{1\bar{2}}, \varphi^{2\bar{1}}, \varphi^{2\bar{2}}\}$  and  
 $\mathbb{C}\{\eta_1^{1,1}, \eta_2^{1,1}\} = \mathbb{C}\{\varphi^{3\bar{1}}, \varphi^{3\bar{2}}\}, \quad \partial\eta_1^{1,1} = \bar{\partial}_{\phi(t)}\varphi^{23}, \quad \partial\eta_2^{1,1} = \bar{\partial}_{\phi(t)}\varphi^{13},$   
 $\partial\varphi^{3\bar{3}} = -\varphi^{12\bar{3}}, \quad \bar{\partial}_{\phi(t)}\varphi^{3\bar{3}} = \varphi^{1\bar{1}\bar{2}}, \quad \bar{\partial}_{\phi(t)}\varphi^{2\bar{3}} = \varphi^{2\bar{1}\bar{2}};$
- $\mathbb{C}\{\varphi^{12\bar{3}}, \xi_1^{2,1}, \xi_2^{2,1}, \xi_3^{2,1}, \xi_4^{2,1}\} = \mathbb{C}\{\varphi^{12\bar{3}}, \varphi^{13\bar{1}}, \varphi^{13\bar{2}}, \varphi^{23\bar{1}}, \varphi^{23\bar{2}}\}$  and  
 $\bar{\partial}_{\phi(t)}\varphi^{12\bar{3}} = -\varphi^{12\bar{1}\bar{2}}, \quad \mathbb{C}\{\bar{\partial}_{\phi(t)}\varphi^{13\bar{3}}, \bar{\partial}_{\phi(t)}\varphi^{23\bar{3}}\} \subset \mathbb{C}\{\varphi^{13\bar{1}\bar{2}}, \varphi^{23\bar{1}\bar{2}}, \varphi^{12\bar{1}\bar{3}}, \varphi^{12\bar{2}\bar{3}}\}$  is of dimension 2;
- $\mathbb{C}\{\xi_1^{1,2}, \xi_2^{1,2}, \xi_3^{1,2}, \xi_4^{1,2}, \bar{\partial}_{\phi(t)}\varphi^{3\bar{3}}\} = \mathbb{C}\{\varphi^{1\bar{1}\bar{3}}, \varphi^{1\bar{2}\bar{3}}, \varphi^{2\bar{1}\bar{3}}, \varphi^{2\bar{2}\bar{3}}, \varphi^{3\bar{1}\bar{2}}\}$  and  
 $\mathbb{C}\{\bar{\partial}_{\phi(t)}\varphi^{3\bar{1}\bar{3}}, \bar{\partial}_{\phi(t)}\varphi^{3\bar{2}\bar{3}}\} = \mathbb{C}\{\varphi^{1\bar{1}\bar{2}\bar{3}}, \varphi^{2\bar{1}\bar{2}\bar{3}}\}, \quad \text{with } \partial\varphi^{3\bar{1}\bar{3}} = -\varphi^{12\bar{1}\bar{3}}, \quad \partial\varphi^{3\bar{2}\bar{3}} = -\varphi^{12\bar{2}\bar{3}};$
- $\mathbb{C}\{\xi_1^{2,2}, \xi_2^{2,2}, \xi_3^{2,2}, \xi_4^{2,2}\} = \mathbb{C}\{\varphi^{13\bar{1}\bar{3}}, \varphi^{13\bar{2}\bar{3}}, \varphi^{23\bar{1}\bar{3}}, \varphi^{23\bar{2}\bar{3}}\}$  with  
 $\xi_1^{2,2}, \xi_2^{2,2}, \xi_3^{2,2} \in \ker \bar{\partial}_{\phi(t)}$  and  $\bar{\partial}_{\phi(t)}\xi_4^{2,2} = \varphi^{12\bar{1}\bar{2}\bar{3}};$
- $\mathbb{C}\{\bar{\partial}_{\phi(t)}\varphi^{3\bar{1}\bar{3}}, \bar{\partial}_{\phi(t)}\varphi^{3\bar{2}\bar{3}}\} = \mathbb{C}\{\varphi^{1\bar{1}\bar{2}\bar{3}}, \varphi^{2\bar{1}\bar{2}\bar{3}}\}.$

*Proof.* From Subsection 3.2.1, we know that  $\ker \bar{\partial}_{\phi(t)} \cap \wedge^{1,0} = \mathbb{C}\{\varphi^1, \varphi^2\}$  and the rank of  $\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^3\} \rightarrow \mathbb{C}\{\varphi^{1\bar{1}}, \varphi^{2\bar{1}}, \varphi^{1\bar{2}}, \varphi^{2\bar{2}}\}$  is 1.

From Subsection 3.2.3, we know that  $\dim \ker \bar{\partial}_{\phi(t)} \cap \wedge^{2,0} = 1$  and  $\bar{\partial}_{\phi(t)}\varphi^{12} = 0$ . On the other hand, the rank of  $\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{13}, \varphi^{23}\} \rightarrow \mathbb{C}\{\varphi^{12\bar{1}}, \varphi^{12\bar{2}}\}$  is 2. So we have

$$\mathbb{C}\{\bar{\partial}_{\phi(t)}\varphi^{12\bar{1}}, \bar{\partial}_{\phi(t)}\varphi^{12\bar{2}}\} = \mathbb{C}\{\varphi^{12\bar{1}}, \varphi^{12\bar{2}}\}.$$

From Subsection 3.2.4, we know that  $\dim \ker \bar{\partial}_{\phi(t)} \cap \wedge^{1,1} = 6$  and the rank of

$$\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{1\bar{3}}, \varphi^{2\bar{3}}, \varphi^{3\bar{1}}, \varphi^{3\bar{2}}\} \longrightarrow \mathbb{C}\{\varphi^{1\bar{1}\bar{2}}, \varphi^{2\bar{1}\bar{2}}\},$$

is equal to  $\text{rank}(T) = 2$ . So we may choose  $\xi_1^{1,1}, \xi_2^{1,1}, \xi_3^{1,1}$  so that  $\mathbb{C}\{\xi_1^{1,1}, \xi_2^{1,1}, \xi_3^{1,1}, \bar{\partial}_{\phi(t)}\varphi^3\} = \mathbb{C}\{\varphi^{1\bar{1}}, \varphi^{1\bar{2}}, \varphi^{2\bar{1}}, \varphi^{2\bar{2}}\}$ , and  $\eta_1^{1,1}, \eta_2^{1,1}$  so that (indeed, we may set  $\eta_1^{1,1} = -t_{11}\varphi^{3\bar{1}} - t_{12}\varphi^{3\bar{2}}$  and  $\eta_2^{1,1} = -t_{21}\varphi^{3\bar{1}} - t_{22}\varphi^{3\bar{2}}$ )

$$\mathbb{C}\{\eta_1^{1,1}, \eta_2^{1,1}\} = \mathbb{C}\{\varphi^{3\bar{1}}, \varphi^{3\bar{2}}\}, \quad \partial\eta_1^{1,1} = \bar{\partial}_{\phi(t)}\varphi^{23}, \quad \partial\eta_2^{1,1} = \bar{\partial}_{\phi(t)}\varphi^{13}.$$

$$\ker \bar{\partial}_{\phi(t)} = \mathbb{C}\{\varphi^{1\bar{1}}, \varphi^{1\bar{2}}, \varphi^{2\bar{1}}, \varphi^{2\bar{2}}, \xi_4^{1,1}, \xi_5^{1,1}\}.$$

On the other hand,  $\bar{\partial}_{\phi(t)}\varphi^{1\bar{3}} = \varphi^{1\bar{1}\bar{2}}$ ,  $\bar{\partial}_{\phi(t)}\varphi^{2\bar{3}} = \varphi^{2\bar{1}\bar{2}}$  and the rank of  $\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{3\bar{3}}\} \rightarrow \mathbb{C}\{\varphi^{1\bar{1}\bar{3}}, \varphi^{1\bar{2}\bar{3}}, \varphi^{2\bar{1}\bar{3}}, \varphi^{2\bar{2}\bar{3}}, \varphi^{3\bar{1}\bar{2}}\}$  is 1. Notice that

$$\partial\bar{\partial}_{\phi(t)}\varphi^{3\bar{3}} = -\varphi^{1\bar{2}\bar{1}\bar{2}} = -\bar{\partial}_{\phi(t)}\varphi^{1\bar{2}\bar{3}} = \bar{\partial}_{\phi(t)}\partial\varphi^{3\bar{3}},$$

From Subsection 3.2.7, we know that  $\mathbb{C}\{\partial\xi_4^{1,1}, \partial\xi_5^{1,1}\} = \mathbb{C}\{\varphi^{1\bar{2}\bar{1}}, \varphi^{1\bar{2}\bar{2}}\} \subset \ker \bar{\partial}_{\phi(t)}$ , the rank of  $\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{1\bar{2}\bar{3}}, \varphi^{1\bar{3}\bar{1}}, \varphi^{1\bar{3}\bar{2}}, \varphi^{2\bar{3}\bar{1}}, \varphi^{2\bar{3}\bar{2}}\} \rightarrow \mathbb{C}\{\varphi^{1\bar{2}\bar{1}\bar{2}}\}$  is 1,  $\bar{\partial}_{\phi(t)}\varphi^{1\bar{3}\bar{3}}$  and  $\bar{\partial}_{\phi(t)}\varphi^{2\bar{3}\bar{3}}$  are linearly independent. We may choose  $\xi_1^{2,1}, \xi_2^{2,1}, \xi_3^{2,1}, \xi_4^{2,1} \in \ker \bar{\partial}_{\phi(t)}$  so that

$$\mathbb{C}\{\varphi^{1\bar{2}\bar{3}}, \xi_1^{2,1}, \xi_2^{2,1}, \xi_3^{2,1}, \xi_4^{2,1}\} = \mathbb{C}\{\varphi^{1\bar{2}\bar{3}}, \varphi^{1\bar{3}\bar{1}}, \varphi^{1\bar{3}\bar{2}}, \varphi^{2\bar{3}\bar{1}}, \varphi^{2\bar{3}\bar{2}}\}.$$

For  $\wedge^{1,2}$ , we already see that  $\varphi^{1\bar{1}\bar{2}} = \bar{\partial}_{\phi(t)}\varphi^{1\bar{3}}, \varphi^{2\bar{1}\bar{2}} = \bar{\partial}_{\phi(t)}\varphi^{2\bar{3}}$  and the rank of  $\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{3\bar{3}}\} \rightarrow \mathbb{C}\{\varphi^{1\bar{1}\bar{3}}, \varphi^{1\bar{2}\bar{3}}, \varphi^{2\bar{1}\bar{3}}, \varphi^{2\bar{2}\bar{3}}, \varphi^{3\bar{1}\bar{2}}\}$  is 1. From Subsection 3.2.8, we know that  $\varphi^{1\bar{1}\bar{2}}, \varphi^{1\bar{1}\bar{3}}, \varphi^{1\bar{2}\bar{3}}, \varphi^{2\bar{1}\bar{2}}, \varphi^{2\bar{1}\bar{3}}, \varphi^{2\bar{2}\bar{3}}, \varphi^{3\bar{1}\bar{2}}$  are in  $\ker \bar{\partial}_{\phi(t)}$  and the rank of  $\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{3\bar{1}\bar{3}}, \varphi^{3\bar{2}\bar{3}}\} \rightarrow \mathbb{C}\{\varphi^{1\bar{1}\bar{2}\bar{3}}, \varphi^{2\bar{1}\bar{2}\bar{3}}\}$  is 1. So we may choose  $\xi_1^{1,2}, \dots, \xi_6^{1,2}$  so that  $\mathbb{C}\{\xi_1^{1,2}, \xi_2^{1,2}, \xi_3^{1,2}, \xi_4^{1,2}, \bar{\partial}_{\phi(t)}\varphi^{3\bar{3}}\} = \mathbb{C}\{\varphi^{1\bar{1}\bar{3}}, \varphi^{1\bar{2}\bar{3}}, \varphi^{2\bar{1}\bar{3}}, \varphi^{2\bar{2}\bar{3}}, \varphi^{3\bar{1}\bar{2}}\}$  and

$$\mathbb{C}\{\xi_5^{1,2}, \xi_6^{1,2}\} = \mathbb{C}\{\varphi^{3\bar{1}\bar{3}}, \varphi^{3\bar{2}\bar{3}}\}, \text{ with } \bar{\partial}_{\phi(t)}\xi_5^{1,2} = 0, \bar{\partial}_{\phi(t)}\xi_6^{1,2} \neq 0.$$

For  $\wedge^{2,2}$ , we already see that

$$\bar{\partial}_{\phi(t)}\varphi^{1\bar{3}\bar{3}}, \bar{\partial}_{\phi(t)}\varphi^{2\bar{3}\bar{3}} \in \mathbb{C}\{\varphi^{1\bar{3}\bar{1}\bar{2}}, \varphi^{1\bar{2}\bar{1}\bar{3}}, \varphi^{1\bar{2}\bar{2}\bar{3}}, \varphi^{2\bar{3}\bar{1}\bar{2}}\},$$

are linearly independent. From Subsection 3.2.10, we know that

$$\varphi^{1\bar{2}\bar{1}\bar{2}}, \varphi^{1\bar{2}\bar{1}\bar{3}}, \varphi^{1\bar{2}\bar{2}\bar{3}}, \varphi^{1\bar{3}\bar{1}\bar{2}}, \varphi^{2\bar{3}\bar{1}\bar{2}} \in \ker \bar{\partial}_{\phi(t)}$$

and the rank of  $\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{1\bar{3}\bar{1}\bar{3}}, \varphi^{1\bar{3}\bar{2}\bar{3}}, \varphi^{2\bar{3}\bar{1}\bar{3}}, \varphi^{2\bar{3}\bar{2}\bar{3}}\} \rightarrow \mathbb{C}\{\varphi^{1\bar{2}\bar{1}\bar{2}\bar{3}}\}$  is 1. Notice that  $\mathbb{C}\{\varphi^{1\bar{2}\bar{1}\bar{3}}, \varphi^{1\bar{2}\bar{2}\bar{3}}\} = \mathbb{C}\{\partial\xi_5^{1,2}, \partial\xi_6^{1,2}\}$  and

$$\mathbb{C}\{\bar{\partial}_{\phi(t)}\varphi^{1\bar{3}\bar{3}}, \bar{\partial}_{\phi(t)}\varphi^{2\bar{3}\bar{3}}, \varphi^{1\bar{2}\bar{1}\bar{2}}, \varphi^{1\bar{2}\bar{1}\bar{3}}, \varphi^{1\bar{2}\bar{2}\bar{3}}\} = \mathbb{C}\{\varphi^{1\bar{2}\bar{1}\bar{2}}, \varphi^{1\bar{2}\bar{1}\bar{3}}, \varphi^{1\bar{2}\bar{2}\bar{3}}, \varphi^{1\bar{3}\bar{1}\bar{2}}, \varphi^{2\bar{3}\bar{1}\bar{2}}\} \subset \ker \bar{\partial}_{\phi(t)}.$$

So


$$\mathbb{C}\{\bar{\partial}_{\phi(t)}\varphi^{1\bar{3}\bar{3}}, \bar{\partial}_{\phi(t)}\varphi^{2\bar{3}\bar{3}}, \varphi^{1\bar{2}\bar{1}\bar{2}}, \partial\xi_5^{1,2}, \partial\xi_6^{1,2}\} = \mathbb{C}\{\varphi^{1\bar{2}\bar{1}\bar{2}}, \varphi^{1\bar{2}\bar{1}\bar{3}}, \varphi^{1\bar{2}\bar{2}\bar{3}}, \varphi^{1\bar{3}\bar{1}\bar{2}}, \varphi^{2\bar{3}\bar{1}\bar{2}}\} \subset \ker \bar{\partial}_{\phi(t)}.$$

Meanwhile, because the rank of  $\bar{\partial}_{\phi(t)} : \mathbb{C}\{\varphi^{1\bar{3}\bar{1}\bar{3}}, \varphi^{1\bar{3}\bar{2}\bar{3}}, \varphi^{2\bar{3}\bar{1}\bar{3}}, \varphi^{2\bar{3}\bar{2}\bar{3}}\} \rightarrow \mathbb{C}\{\varphi^{1\bar{2}\bar{1}\bar{2}\bar{3}}\}$  is 1, we may choose  $\xi_1^{2,2}, \xi_2^{2,2}, \xi_3^{2,2}, \xi_4^{2,2} \in \ker \bar{\partial}_{\phi(t)}$  so that

$$\mathbb{C}\{\xi_1^{2,2}, \xi_2^{2,2}, \xi_3^{2,2}, \xi_4^{2,2}\} = \mathbb{C}\{\varphi^{1\bar{3}\bar{1}\bar{3}}, \varphi^{1\bar{3}\bar{2}\bar{3}}, \varphi^{2\bar{3}\bar{1}\bar{3}}, \varphi^{2\bar{3}\bar{2}\bar{3}}\} \text{ and } \bar{\partial}_{\phi(t)}\xi_4^{2,2} = \varphi^{1\bar{2}\bar{1}\bar{2}\bar{3}}.$$

For  $\wedge^{1,3}$ , we already see that  $\mathbb{C}\{\bar{\partial}_{\phi(t)}\varphi^{3\bar{1}\bar{3}}, \bar{\partial}_{\phi(t)}\varphi^{3\bar{2}\bar{3}}\} = \mathbb{C}\{\varphi^{1\bar{1}\bar{2}\bar{3}}, \varphi^{2\bar{1}\bar{2}\bar{3}}\}$  and it is obvious that  $\bar{\partial}_{\phi(t)}\varphi^{3\bar{1}\bar{2}\bar{3}} = 0$ .  $\square$

By using Proposition 4.2, we immediately get the following diagrammatic view of the structure of  $(\wedge^\bullet, d_{\phi(t)} = \partial + \bar{\partial}_{\phi(t)})$  for  $(t_{11}, t_{12}, t_{21}, t_{22}) \neq 0, D(t) \neq 0$ :



### 4.3. Proof of Theorem 1.1.

*Proof of Theorem 1.1.* The conclusions of Theorem 1.1 follows immediately from the two diagrams in Subsection 4.1 and 4.2.  $\square$

## 5. THE DIMENSION OF COHOMOLOGIES AND $E_r^{p,q}(X_t)$

In this section, we compute the Dolbeault/Bott-Chern cohomology and the Frölicher spectral sequence  $(E_r^{\bullet,\bullet}, d_r^{\bullet,\bullet})$  of the double complex  $(\wedge^\bullet, d_{\phi(t)} = \partial + \bar{\partial}_{\phi(t)})$  by counting zigzags in diagrams obtained in Subsection 4.1 and 4.2.

According to Theorem 2.2 and the fact that  $(\wedge^\bullet, d_{\phi(t)} = \partial + \bar{\partial}_{\phi(t)}) \hookrightarrow (A^\bullet(X), d_{\phi(t)} = \partial + \bar{\partial}_{\phi(t)})$  is an  $E_1$ -isomorphism, we immediately get the following

TABLE 1. Dimensions of  $H_{\bar{\partial}_{\phi(t)}}^{\bullet, \bullet}(X)$ ,  $H_{BC, \phi(t)}^{\bullet, \bullet}(X)$  and  $\text{im } d_1^{\bullet, \bullet}(X_t)$

	(i)	(ii)	(iii)		(i)	(ii)	(iii)		(i)	(ii)	(iii)
$h_{\bar{\partial}_{\phi(t)}}^{1,0}$	3	2	2	$h_{BC, \phi(t)}^{1,0}$	2	2	2	$\text{im } d_1^{1,0}$	1	0	0
$h_{\bar{\partial}_{\phi(t)}}^{0,1}$	2	2	2	$h_{BC, \phi(t)}^{0,1}$	2	2	2	$\text{im } d_1^{0,1}$	0	0	0
$h_{\bar{\partial}_{\phi(t)}}^{2,0}$	3	2	1	$h_{BC, \phi(t)}^{2,0}$	3	2	1	$\text{im } d_1^{2,0}$	0	0	0
$h_{\bar{\partial}_{\phi(t)}}^{1,1}$	6	5	5	$h_{BC, \phi(t)}^{1,1}$	4	4	4	$\text{im } d_1^{1,1}$	2	1	0
$h_{\bar{\partial}_{\phi(t)}}^{0,2}$	2	2	2	$h_{BC, \phi(t)}^{0,2}$	3	3	3	$\text{im } d_1^{0,2}$	0	0	0
$h_{\bar{\partial}_{\phi(t)}}^{3,0}$	1	1	1	$h_{BC, \phi(t)}^{3,0}$	1	1	1	$\text{im } d_1^{3,0}$	0	0	0
$h_{\bar{\partial}_{\phi(t)}}^{2,1}$	6	5	4	$h_{BC, \phi(t)}^{2,1}$	6	6	6	$\text{im } d_1^{2,1}$	0	0	0
$h_{\bar{\partial}_{\phi(t)}}^{1,2}$	6	5	4	$h_{BC, \phi(t)}^{1,2}$	6	6	6	$\text{im } d_1^{1,2}$	2	1	0
$h_{\bar{\partial}_{\phi(t)}}^{0,3}$	1	1	1	$h_{BC, \phi(t)}^{0,3}$	1	1	1	$\text{im } d_1^{0,3}$	0	0	0
$h_{\bar{\partial}_{\phi(t)}}^{3,1}$	2	2	2	$h_{BC, \phi(t)}^{3,1}$	2	2	2	$\text{im } d_1^{3,1}$	0	0	0
$h_{\bar{\partial}_{\phi(t)}}^{2,2}$	6	5	5	$h_{BC, \phi(t)}^{2,2}$	8	7	7	$\text{im } d_1^{2,2}$	0	0	0
$h_{\bar{\partial}_{\phi(t)}}^{1,3}$	3	2	1	$h_{BC, \phi(t)}^{1,3}$	2	2	2	$\text{im } d_1^{1,3}$	1	0	0
$h_{\bar{\partial}_{\phi(t)}}^{3,2}$	2	2	2	$h_{BC, \phi(t)}^{3,2}$	3	3	3	$\text{im } d_1^{3,2}$	0	0	0
$h_{\bar{\partial}_{\phi(t)}}^{2,3}$	3	2	2	$h_{BC, \phi(t)}^{2,3}$	3	3	3	$\text{im } d_1^{2,3}$	0	0	0

where  $\text{im } d_1^{p,q}$  is short for  $\dim \text{im } d_1^{p,q}(X_t) = \dim \text{im } d_1^{p,q}(X, d_{\phi(t)})$  and we know that  $H_{\bar{\partial}_{\phi(t)}}^{\bullet, \bullet}(X)$  is isomorphic to  $H_{\bar{\partial}_t}^{\bullet, \bullet}(X_t)$  [Xia22]. It is clear that for  $r > 1$ ,  $d_r^{p,q} = 0$  for any  $(p, q)$  because all zigzags appeared have length less than 4. Furthermore, because

$$E_{r+1}^{p,q} \cong \frac{\ker d_r^{p,q}}{\text{im } d_r^{p-r, q+r-1}} \quad \text{and} \quad E_r^{p,q} / \ker d_r^{p,q} \cong \text{im } d_r^{p,q}, \quad \forall r \geq 1,$$

we have

$$(5.1) \quad \begin{cases} \dim E_2^{p,q} &= \dim E_1^{p,q} - \dim \text{im } d_1^{p,q} - \dim \text{im } d_1^{p-1, q}, \\ \dim E_3^{p,q} &= \dim E_2^{p,q} - \dim \text{im } d_2^{p,q} - \dim \text{im } d_2^{p-2, q+1}, \\ &\vdots \\ \dim E_{r+1}^{p,q} &= \dim E_r^{p,q} - \dim \text{im } d_r^{p,q} - \dim \text{im } d_r^{p-r, q+r-1}. \end{cases}$$

It follows that

TABLE 2.  $e_r^{\bullet,\bullet} = \dim E_r^{\bullet,\bullet}(X_t)$  and  $\chi_q^p(E_2) = \sum_{i=0}^q (-1)^{q-i} e_2^{p,i}$

	(i)	(ii)	(iii)		(i)	(ii)	(iii)		(i)	(ii)	(iii)
$e_1^{1,0}$	3	2	2	$e_2^{1,0}$	2	2	2	$\chi_1^0(E_2)$	1	1	1
$e_1^{0,1}$	2	2	2	$e_2^{0,1}$	2	2	2	$\chi_2^0(E_2)$	1	1	1
$e_1^{2,0}$	3	2	1	$e_2^{2,0}$	2	2	1	$\chi_3^0(E_2)$	0	0	0
$e_1^{1,1}$	6	5	5	$e_2^{1,1}$	4	4	5	$\chi_1^1(E_2)$	2	2	3
$e_1^{0,2}$	2	2	2	$e_2^{0,2}$	2	2	2	$\chi_2^1(E_2)$	2	2	1
$e_1^{3,0}$	1	1	1	$e_2^{3,0}$	1	1	1	$\chi_3^1(E_2)$	0	0	0
$e_1^{2,1}$	6	5	4	$e_2^{2,1}$	4	4	4	$\chi_1^2(E_2)$	2	2	3
$e_1^{1,2}$	6	5	4	$e_2^{1,2}$	4	4	4	$\chi_2^2(E_2)$	2	2	2
$e_1^{0,3}$	1	1	1	$e_2^{0,3}$	1	1	1	$\chi_3^2(E_2)$	0	0	0
$e_1^{3,1}$	2	2	2	$e_2^{3,1}$	2	2	2	$\chi_1^3(E_2)$	1	1	1
$e_1^{2,2}$	6	5	5	$e_2^{2,2}$	4	4	5	$\chi_2^3(E_2)$	1	1	1
$e_1^{1,3}$	3	2	1	$e_2^{1,3}$	2	2	1	$\chi_3^3(E_2)$	0	0	0
$e_1^{3,2}$	2	2	2	$e_2^{3,2}$	2	2	2				
$e_1^{2,3}$	3	2	2	$e_2^{2,3}$	2	2	2				

Since  $d_r^{p,q} = 0$  for any  $(p, q)$  and  $r > 1$ , we have  $E_2^{p,q} = E_\infty^{p,q}$ . Furthermore, we see from Table 2 that  $\chi_1^1(E_2)$  is lower semi-continuous while  $\chi_2^1(E_2)$  is upper semi-continuous.

#### REFERENCES

- [AK17] D. Angella and H. Kasuya. Bott-Chern cohomology of solvmanifolds. *Ann. Global Anal. Geom.*, 52(4):363–411, 2017.
- [Ang13] D. Angella. The cohomologies of the Iwasawa manifold and of its small deformations. *J. Geom. Anal.*, 23(3):1355–1378, 2013.
- [AT13] D. Angella and A. Tomassini. On the  $\partial\bar{\partial}$ -lemma and Bott-Chern cohomology. *Invent. Math.*, 192(1):71–81, 2013.
- [CFG91] L. A. Cordero, M. Fernández, and A. Gray. The Frölicher spectral sequence for compact nilmanifolds. *Illinois J. Math.*, 35(1):56–67, 1991.
- [CFUG97] L. Cordero, M. Fernández, L. Ugarte, and A. Gray. A general description of the terms in the Frölicher spectral sequence. *Differential Geom. Appl.*, 7(1):75–84, 1997.
- [Cle05] H. Clemens. Geometry of formal Kuranishi theory. *Adv. Math.*, 198(1):311–365, 2005.
- [Fla20] C. Flavi. On the Frölicher spectral sequence of the Iwasawa manifold and its small deformations. *Riv. Math. Univ. Parma (N.S.)*, 11(2):283–292, 2020.



- [Fle81] H. Flenner. Ein Kriterium für die Offenheit der Versalität. *Math. Z.*, 178(4):449–473, 1981.
- [FM06] D. Fiorenza and M. Manetti. L-infinity algebras, Cartan homotopies and period maps. arXiv:math/0605297, 2006.
- [FM09] D. Fiorenza and M. Manetti. A period map for generalized deformations. *J. Noncommut. Geom.*, 3(4):579–597, 2009.
- [HX24] Y. Hu and W. Xia. Deformed Aeppli cohomology: canonical deformations and jumping formulas. preprint, 2024.
- [Kas13] H. Kasuya. Techniques of computations of Dolbeault cohomology of solvmanifolds. *Math. Z.*, 273(1-2):437–447, 2013.
- [Kas15] H. Kasuya. The Frölicher spectral sequence of certain solvmanifolds. *J. Geom. Anal.*, 25(1):317–328, 2015.
- [KQ20] M. Khovanov and Y. Qi. A faithful braid group action on the stable category of tri-complexes. *SIGMA Symmetry Integrability Geom. Methods Appl.*, 16:Paper No. 019, 32, 2020.
- [KS23] H. Kasuya and J. Stelzig. Frölicher spectral sequence and Hodge structures on the cohomology of complex parallelisable manifolds. *Transform. Groups*, 28(4):1595–1606, 2023.
- [LRY15] K. Liu, S. Rao, and X. Yang. Quasi-isometry and deformations of Calabi-Yau manifolds. *Invent. Math.*, 199(2):423–453, 2015.
- [LSY09] K. Liu, X. Sun, and S.-T. Yau. Recent development on the geometry of the Teichmüller and moduli spaces of Riemann surfaces. In *Geometry of Riemann surfaces and their moduli spaces*, volume XIV of *Surveys in differential geometry*, pages 221–259. 2009.
- [Man04] M. Manetti. Lectures on deformations of complex manifolds (deformations from differential graded viewpoint). *Rend. Mat. Appl. (7)*, 24(1):1–183, 2004.
- [Mas20] M. Maschio. On the degeneration of the Frölicher spectral sequence and small deformations. *Complex Manifolds*, 7(1):62–72, 2020.
- [Nak75] I. Nakamura. Complex parallelisable manifolds and their small deformations. *J. Differential Geom.*, 10(1):85–112, 1975.
- [PSU21] D. Popovici, J. Stelzig, and L. Ugarte. Higher-page Bott–Chern and Aeppli cohomologies and applications. *J. Reine Angew. Math.*, 777:157–194, 2021.
- [RYY19] S. Rao, S. Yang, and X.-D. Yang. Dolbeault cohomologies of blowing up complex manifolds. *J. Math. Pures Appl.*, 130:68–92, 2019.
- [RZ18] S. Rao and Q. Zhao. Several special complex structures and their deformation properties. *J. Geom. Anal.*, 28(4):2984–3047, 2018.
- [RZ20] S. Rao and Y. Zou.  $\partial\bar{\partial}$ -lemma and double complex. to appear in *Commun. Math. Stat.*, 2020.
- [Ste21] J. Stelzig. On the structure of double complexes. *J. Lond. Math. Soc. (2)*, 104(2):956–988, 2021.
- [Tod89] A. N. Todorov. The Weil-Petersson geometry of the moduli space of  $SU(\geq 3)$  (Calabi-Yau) manifolds i. *Comm. Math. Phys.*, 126(2):325–346, 1989.
- [WX23] X. Wan and W. Xia. Deformations of  $(p, q)$ -forms and degenerations of the Frölicher spectral sequence. arXiv:2312.13692, 2023.
- [WZ24] D. Wei and S. Zhu. Two Applications of the  $\partial\bar{\partial}$ -Hodge Theory. *Chinese Ann. Math. Ser. B*, 45(1):137–150, 2024.
- [Xia19] W. Xia. Derivations on almost complex manifolds. *Proc. Amer. Math. Soc.*, 147:559–566, 2019. Errata in arXiv:1809.07443v3.
- [Xia21a] W. Xia. Deformations of Dolbeault cohomology classes for Lie algebra with complex structures. *Ann. Global Anal. Geom.*, 60(3):709–734, 2021.
- [Xia21b] W. Xia. On the deformed Bott-Chern cohomology. *J. Geom. Phys.*, 166:Paper No. 104250, 19, 2021.
- [Xia22] W. Xia. Deformations of Dolbeault cohomology classes. *Math. Z.*, 300:2931–2973, 2022.
- [ZR15] Q. Zhao and S. Rao. Extension formulas and deformation invariance of Hodge numbers. *C. R. Math. Acad. Sci. Paris*, 353(11):979–984, 2015.

YAN HU, MATHEMATICAL SCIENCE RESEARCH CENTER, CHONGQING UNIVERSITY OF TECHNOLOGY, CHONGQING 400054, CHINA

*Email address:* `huyan1128@126.com`

WEI XIA, MATHEMATICAL SCIENCE RESEARCH CENTER, CHONGQING UNIVERSITY OF TECHNOLOGY, CHONGQING 400054, CHINA

*Email address:* `xiaweiwei3@126.com`, `xiawei@cqut.edu.cn`