

ON BANDLIMITED MULTIPLIERS ON MATRIX-WEIGHTED L^p -SPACES

MORTEN NIELSEN

ABSTRACT. We extend a classical result by Triebel on boundedness of bandlimited multipliers on $L^p(\mathbb{R}^n)$, $0 < p \leq 1$, to a vector-valued and matrix-weighted setting with boundedness of the bandlimited multipliers obtained on $L^p(W)$, $0 < p \leq 1$, for matrix-weights $W : \mathbb{R}^n \rightarrow \mathbb{C}^{N \times N}$ that satisfies a matrix Muckenhoupt A_p -condition.

1. INTRODUCTION

An $N \times N$ matrix weight on \mathbb{R}^n is a locally integrable and almost everywhere positive definite matrix function $W : \mathbb{R}^n \rightarrow \mathbb{C}^{N \times N}$. The matrix-weighted L^p -space $L^p(W)$, $0 < p < \infty$, is defined for any matrix-weight $W : \mathbb{R}^n \rightarrow \mathbb{C}^{N \times N}$ as the family of measurable functions $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{C}^N$ satisfying

$$(1.1) \quad \|\mathbf{f}\|_{L^p(W)} := \left(\int_{\mathbb{R}^n} |W^{1/p}(x)\mathbf{f}(x)|^p dx \right)^{1/p} < \infty.$$

Using the standard identification of (vector-)functions that differ on a set of measure zero, one can verify that $L^p(W)$ is a (quasi-)Banach space.

The matrix-weighted L^p -spaces have attracted a great deal of attention recently (see, e.g., [1, 10, 11, 13]) due to the fact that the setup generates a number interesting mathematical questions for vector valued functions that naturally connect to classical results on Muckenhoupt weights in harmonic analysis. A highlight in the matrix-weighted case is the formulation of a suitable matrix A_p condition by Nazarov, Treil and Volberg that completely characterizes boundedness of the Riesz-transform(s) on $L^p(W)$ for $1 < p < \infty$, see [14, 16], see also [8].

In the present paper we study Fourier multiplier on $L^p(W)$ with a focus on the case $0 < p \leq 1$. As is well-known, a scalar Fourier multiplier is a function $\varphi \in L^\infty(\mathbb{R}^n)$ that induces a corresponding bounded multiplier operator

$$\varphi(D)f := \mathcal{F}^{-1}(\varphi \mathcal{F}f), \quad f \in L^2(\mathbb{R}^n),$$

where \mathcal{F} denotes the Fourier transform on $L^2(\mathbb{R}^n)$, where we used the normalisation specified in Eq. (2.4) below.

In case, $\mathcal{F}^{-1}\varphi \in L^1(\mathbb{R}^n)$, Young's inequality provides an easy extension of the Fourier multiplier to $L^p(\mathbb{R}^n)$, $1 \leq p \leq \infty$, through the estimate

$$(1.2) \quad \|\varphi(D)f\|_{L^p(\mathbb{R}^n)} \leq C \|\mathcal{F}^{-1}\varphi\|_{L^1(\mathbb{R}^n)} \|f\|_{L^p(\mathbb{R}^n)}.$$

Let us now lift the multiplier to the vector-setting. Given a suitably nice $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{C}^N$, we let $\varphi(D)$ act coordinate-wise on $\mathbf{f} = (f_1, \dots, f_N)^T$, i.e., $\varphi(D)\mathbf{f} := (\varphi(D)f_1, \dots, \varphi(D)f_N)^T$. A much more challenging question is then whether $\varphi(D)$ extends to a bounded operator on $L^p(W)$ for a given matrix-weight $W : \mathbb{R}^n \rightarrow \mathbb{C}^{N \times N}$? This problem has been studied in detail in the case $1 \leq p < \infty$ by various authors. In case $1 < p < \infty$, the results on singular integrals obtain by Goldberg in [8] provides boundedness $\varphi(D) : L^p(W) \rightarrow L^p(W)$ provided

2010 *Mathematics Subject Classification.* Primary 42A45, 47B37; Secondary 47B38.

Key words and phrases. Matrix weight, Muckenhoupt condition, band-limited functions, sampling theorem, Fourier multiplier operator.

that $\mathcal{F}^{-1}\varphi$ satisfies a mild decay condition, and, more importantly, also provided that the matrix-weight W satisfies the so-called matrix Muckenhoupt A_p -condition, see Definition 2.1 below. Frazier and Roudenko extended this result in [7] to obtain boundedness in the end-point case $\varphi(D) : L^1(W) \rightarrow L^1(W)$, provided W satisfies a matrix Muckenhoupt A_1 -condition. This leaves the range $0 < p < 1$, which we will study in the present article.

Even in the scalar case, it is known that there are no unrestricted extensions of Young's estimate (1.2) to the range $0 < p < 1$, so let us discuss a framework that makes the case $0 < p < 1$ manageable. Here the notion of bandlimited functions will be central. Triebel showed [15, Theorem 1.5.1] that in case $0 < p \leq 1$, $\varphi : B(0,1) \rightarrow \mathbb{C}$ is compactly supported, and f is a bandlimited tempered distribution in the sense that the frequency support of f satisfies $\text{supp}(\hat{f}) \subseteq \{x \in \mathbb{R}^n : |x| < 1\}$, there exists C independent of f such that

$$\|\varphi(D)f\|_{L^p(\mathbb{R}^n)} \leq C\|f\|_{L^p(\mathbb{R}^n)},$$

provided $\mathcal{F}^{-1}\varphi$ satisfies a mild decay condition.

We will extend Triebel's result on bandlimited multipliers to the matrix-weighted case in Section 2 for weights W that satisfies a matrix A_p -condition. Finally, in Section 3, we provide a specific motivation for studying vector-valued Fourier multipliers in the case $0 < p \leq 1$ by considering an application of the multiplier result to the study of matrix-weighted smoothness spaces.

2. MUCKENHOUPHT WEIGHTS AND FOURIER MULTIPLIERS

The matrix A_p -conditions will be of fundamental importance in order to derive our main multiplier result, so let us first define these conditions. We let \mathcal{Q} denote the collection of all cubes $\{Q(z,r)\}_{z \in \mathbb{R}^n, r > 0}$ in \mathbb{R}^n , where $Q(z,r) := z + r[-1/2, 1/2]^n$.

Definition 2.1. Let $W : \mathbb{R}^n \rightarrow \mathbb{C}^{N \times N}$ be a matrix weight. We say that W satisfies the matrix Muckenhoupt A_p condition for $1 < p < \infty$ provided

$$(2.1) \quad [W]_{\mathbf{A}_p(\mathbb{R}^n)} := \sup_{Q \in \mathcal{Q}} \int_Q \left(\int_Q \|W^{1/p}(x)W^{-1/p}(t)\|^{p'} \frac{dt}{|Q|} \right)^{p/p'} \frac{dx}{|Q|} < +\infty.$$

In case $0 < p \leq 1$, W is said to satisfy the matrix Muckenhoupt A_p condition provided

$$(2.2) \quad [W]_{\mathbf{A}_p(\mathbb{R}^n)} := \sup_{Q \in \mathcal{Q}} \text{ess sup}_{y \in Q} \frac{1}{|Q|} \int_Q \|W(t)^{1/p}W^{-1/p}(y)\|^p dt < +\infty.$$

The norm $\|\cdot\|$ appearing in the integrals is any matrix norm on the $N \times N$ matrices. In case either (2.1) or (2.2) applies, we write $W \in \mathbf{A}_p(\mathbb{R}^n)$.

Remark 2.2. The matrix A_p -condition for $1 < p < \infty$ was introduced in [12, 14, 16] using the notion of dual norms. The condition given in (2.1) was first studied by Roudenko in [13], where the condition is also proven to be equivalent to the original matrix A_p -condition. For $0 < p \leq 1$, Frazier and Roudenko introduced the condition (2.2) in [6]. Also, in the scalar case $N = 1$, one can verify for $1 \leq p < \infty$ that the conditions in Definition 2.1 are equivalent to the well known scalar A_p conditions.

Remark 2.3. Since the family of cubes \mathcal{Q} in \mathbb{R}^n is clearly invariant under dilations $\mathbf{x} \rightarrow R\mathbf{x}$ for $R > 0$, it easily follows from (2.1) and (2.2) that $\mathbf{A}_p(\mathbb{R}^n)$ is invariant under such dilations. In fact, for any $W \in \mathbf{A}_p(\mathbb{R}^n)$, we have

$$[W(R\cdot)]_{\mathbf{A}_p(\mathbb{R}^n)} = [W]_{\mathbf{A}_p(\mathbb{R}^n)}.$$

Let us suppose that $W \in \mathbf{A}_p(\mathbb{R}^n)$. For a fixed vector $\mathbf{x} \in \mathbb{C}^N$, consider the scalar weight $w_{\mathbf{x}}(t) := \|W^{1/p}(t)x\|^p$. For $1 < p < \infty$, it is known that $w_{\mathbf{x}}$ is a scalar A_p -weight with A_p constant depending only on $[W]_{\mathbf{A}_p(\mathbb{R}^n)}$, see [8, Corollary 2.2], and for $0 < p \leq 1$, it was shown in [6, Lemma 2.1] that $w_{\mathbf{x}}$ is in scalar A_1 with an A_1 constant that only depends

on $[W]_{\mathbf{A}_p(\mathbb{R}^n)}$. In both cases, we may conclude, see, e.g., [9, Proposition 9.1.5.], that $w_{\mathbf{x}}$ induces a doubling measure in the sense that there exists a constant C independent of \mathbf{x} such that for any $Q(z, r) \in \mathcal{Q}$,

$$(2.3) \quad \int_{Q(z, 2r)} w_{\mathbf{x}}(t) dt \leq C \int_{Q(z, r)} w_{\mathbf{x}}(t) dt.$$

The doubling exponent β is defined by letting $2^\beta = C$, with C the smallest value of C satisfying (2.3). We observe that for the unit cubes $Q_k := Q(k, 1)$, $k \in \mathbb{Z}^n$, the doubling condition ensures that there exists a constant $c > 0$, independent of \mathbf{x} , such that for $k, \ell \in \mathbb{Z}^n$,

$$\int_{Q_k} w_{\mathbf{x}}(t) dt \leq c(1 + |k - \ell|)^\beta \int_{Q_\ell} w_{\mathbf{x}}(t) dt,$$

with β the doubling exponent of $w_{\mathbf{x}}$, due to the fact that $Q(k, 1) \subseteq Q(\ell, |k - \ell| + 1)$.

Let us specify our chosen normalisation of the Fourier transform. For $f \in L_1(\mathbb{R}^n)$, we let

$$(2.4) \quad \mathcal{F}(f)(\xi) := (2\pi)^{-n/2} \int_{\mathbb{R}^n} f(x) e^{-ix \cdot \xi} dx, \quad \xi \in \mathbb{R}^n,$$

denote the Fourier transform, and we use the standard notation $\hat{f}(\xi) = \mathcal{F}(f)(\xi)$. With this normalisation, the Fourier transform extends to a unitary transform on $L^2(\mathbb{R}^n)$ and we denote the inverse Fourier transform by \mathcal{F}^{-1} .

For $R > 0$, we define the following class of vector-valued functions with band-limited coordinate functions, where $B(a, r)$ denotes the (open) Euclidean ball in \mathbb{R}^n centered at a with radius r , and $\mathcal{S}'(\mathbb{R}^n)$ denotes the tempered distributions defined on \mathbb{R}^n ,

$$(2.5) \quad E_R := \{\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{C}^N : f_i \in \mathcal{S}'(\mathbb{R}^n) \text{ and } \text{supp}(\hat{f}_i) \subseteq B(0, R), i = 1, \dots, N\}.$$

We are now ready to state and prove our main result.

Proposition 2.4. *Let $W \in \mathbf{A}_p(\mathbb{R}^n)$ for some $0 < p \leq 1$ and let $\beta > 0$ be the doubling exponent from Eq. (2.3) associated with W . Suppose there is a constant K such that the compactly supported function $\varphi : B(0, R) \rightarrow \mathbb{C}$ satisfies*

$$|\mathcal{F}^{-1}(\varphi)(x)| \leq KR^n(1 + R|x|)^{-M}, \quad x \in \mathbb{R}^n,$$

for some $M > (n + \beta)/p$. Then there exists a finite constant $C := C([W]_{\mathbf{A}_p(\mathbb{R}^n)}, K, p)$ such that the Fourier multiplier

$$\varphi(D)\mathbf{f} := \mathcal{F}^{-1}[\varphi \cdot \mathcal{F}(\mathbf{f})],$$

defined for $\mathbf{f} \in \mathcal{S}'(\mathbb{R}^n)$ with $\text{supp}(\hat{\mathbf{f}}) \subseteq B(0, R)$, satisfies

$$\|\varphi(D)\mathbf{f}\|_{L^p(W)} \leq C\|\mathbf{f}\|_{L^p(W)}$$

for all $\mathbf{f} \in E_R \cap L^p(W)$.

Proof. Let us first consider the special case $R = 1$. For any $\mathbf{f} \in E_1$, we notice that $\varphi(D)\mathbf{f} = (\mathcal{F}^{-1}\varphi) * \mathbf{f}$, and for $t, u \in \mathbb{R}^n$, we have the sampling representation (see, e.g., [6, Section 3] or [5, Section 6]),

$$(\mathcal{F}^{-1}\varphi * \mathbf{f})(t) = \sum_{\ell \in \mathbb{Z}^n} \mathbf{f}(\ell + u) [\mathcal{F}^{-1}\varphi](t - u - \ell).$$

Hence,

$$W^{1/p}(t)(\mathcal{F}^{-1}\varphi * \mathbf{f})(t) = \sum_{\ell \in \mathbb{Z}^n} W^{1/p}(t)\mathbf{f}(\ell + u) [\mathcal{F}^{-1}\varphi](t - u - \ell),$$

which implies that,

$$|W^{1/p}(t)(\mathcal{F}^{-1}\varphi * \mathbf{f})(t)| \leq \sum_{\ell \in \mathbb{Z}^n} |W^{1/p}(t)\mathbf{f}(\ell + u) [\mathcal{F}^{-1}\varphi](t - u - \ell)|.$$

As before, we let $Q_k := Q(k, 1)$ for $k \in \mathbb{Z}^n$. Using $0 < p \leq 1$, we arrive at the following estimate valid for $u \in Q_0$, $t \in \mathbb{R}^n$,

$$\begin{aligned}
|W^{1/p}(t)(\mathcal{F}^{-1}\varphi * \mathbf{f})(t)|^p &\leq \sum_{\ell \in \mathbb{Z}^n} |W^{1/p}(t)\mathbf{f}(\ell + u)|^p |[\mathcal{F}^{-1}\varphi](t - u - \ell)|^p \\
&\leq K^p \sum_{\ell \in \mathbb{Z}^n} |W^{1/p}(t)\mathbf{f}(\ell + u)|^p (1 + |t - u - \ell|)^{-Mp} \\
(2.6) \qquad \qquad \qquad &\leq c_M K^p \sum_{\ell \in \mathbb{Z}^n} |W^{1/p}(t)\mathbf{f}(\ell + u)|^p (1 + |t - \ell|)^{-Mp}.
\end{aligned}$$

We now average the estimate (2.6) over $u \in Q_0$, noticing that $|Q_0| = 1$,

$$\begin{aligned}
|W^{1/p}(t)(\mathcal{F}^{-1}\varphi * \mathbf{f})(t)|^p &\leq c_M K^p \int_{Q_0} \sum_{\ell \in \mathbb{Z}^n} |W^{1/p}(t)\mathbf{f}(\ell + u)|^p (1 + |t - \ell|)^{-Mp} du \\
&= c_M K^p \sum_{\ell \in \mathbb{Z}^n} (1 + |t - \ell|)^{-Mp} \int_{Q_0} |W^{1/p}(t)\mathbf{f}(\ell + u)|^p du \\
(2.7) \qquad \qquad \qquad &= c_M K^p \sum_{\ell \in \mathbb{Z}^n} (1 + |t - \ell|)^{-Mp} \int_{Q_\ell} |W^{1/p}(t)\mathbf{f}(y)|^p dy,
\end{aligned}$$

where we have used Tonelli's theorem. By the doubling condition satisfied by the scalar weight $w_{\mathbf{w}}(t) := |W^{1/t}(t)\mathbf{w}|^p$, with doubling exponent β independent of $\mathbf{w} \in \mathbb{C}^N$, there exists a finite constant c_w such that for any $k, \ell \in \mathbb{Z}^n$, and $y \in \mathbb{R}^n$,

$$(2.8) \qquad \int_{Q_k} |W^{1/p}(t)\mathbf{f}(y)|^p dt \leq c_w (1 + |k - \ell|)^\beta \int_{Q_\ell} |W^{1/p}(t)\mathbf{f}(y)|^p dt.$$

For $k \in \mathbb{Z}^n$, we integrate inequality (2.7) over $t \in Q_k$, using Tonelli's theorem once more together with the estimate (2.8),

$$\begin{aligned}
\int_{Q_k} |W^{1/p}(t)(\varphi(D)\mathbf{f})(t)|^p dt &\leq c_M K^p \int_{Q_k} \sum_{\ell \in \mathbb{Z}^n} (1 + |k - \ell|)^{-Mp} \int_{Q_\ell} |W^{1/p}(t)\mathbf{f}(y)|^p dy dt \\
&\leq c'_M \sum_{\ell \in \mathbb{Z}^n} (1 + |k - \ell|)^{-Mp+\beta} \int_{Q_\ell} \int_{Q_\ell} |W^{1/p}(t)\mathbf{f}(y)|^p dt dy,
\end{aligned}$$

with $c'_M := c_w c_M K^p$. In the following concluding estimate, the matrix A_p condition (2.2) for W will be essential. We first notice that $\{Q_k\}_k$ forms a partition of \mathbb{R}^n with $|Q_k| = 1$, so we have, using the assumption that $Mp - \beta > n$ and putting $L := \sum_k (1 + |k|)^{-Mp+\beta} < \infty$,

$$\begin{aligned}
\|\varphi(D)\mathbf{f}\|_{L^p(W)}^p &= \sum_{k \in \mathbb{Z}^n} \int_{Q_k} |W^{1/p}(t)(\varphi(D)\mathbf{f})(t)|^p dt \\
&\leq c'_M \sum_{k \in \mathbb{Z}^n} \sum_{\ell \in \mathbb{Z}^n} (1 + |k - \ell|)^{-Mp+\beta} \int_{Q_\ell} \int_{Q_\ell} |W^{1/p}(t)\mathbf{f}(y)|^p dt dy \\
&= Lc'_M \sum_{\ell \in \mathbb{Z}^n} \int_{Q_\ell} \int_{Q_\ell} |W^{1/p}(t)\mathbf{f}(y)|^p dt dy \\
&= Lc'_M \sum_{\ell \in \mathbb{Z}^n} \int_{Q_\ell} \int_{Q_\ell} |W^{1/p}(t)W^{-1/p}(y)W^{1/p}(y)\mathbf{f}(y)|^p dt dy \\
&\leq Lc'_M \sum_{\ell \in \mathbb{Z}^n} \int_{Q_\ell} \left(\int_{Q_\ell} \|W^{1/p}(t)W^{-1/p}(y)\|^p dt \right) |W^{1/p}(y)\mathbf{f}(y)|^p dy \\
&= Lc'_M \sum_{\ell \in \mathbb{Z}^n} \int_{Q_\ell} \left(\frac{1}{|Q_\ell|} \int_{Q_\ell} \|W^{1/p}(t)W^{-1/p}(y)\|^p dt \right) |W^{1/p}(y)\mathbf{f}(y)|^p dy
\end{aligned}$$

$$\begin{aligned}
&\leq Lc'_M[W]_{\mathbf{A}_p(\mathbb{R}^d)} \sum_{\ell \in \mathbb{Z}^n} \int_{Q_\ell} |W^{1/p}(y)\mathbf{f}(y)|^p dy \\
&= Lc'_M[W]_{\mathbf{A}_p(\mathbb{R}^d)} \|\mathbf{f}\|_{L^p(W)}^p \\
&= C^p \|\mathbf{f}\|_{L^p(W)}^p,
\end{aligned}$$

with $C^p := Lc'_M[W]_{\mathbf{A}_p(\mathbb{R}^d)}$, which completes the proof in the case $R = 1$. For general $0 < R < \infty$, we consider the multiplier $\psi : B(0, 1) \rightarrow \mathbb{C}$ defined by $\psi(\cdot) = \varphi(R\cdot)$. We clearly have

$$\mathcal{F}^{-1}(\psi)(x) = R^{-n} \mathcal{F}^{-1}(\varphi)(R^{-1}x) \implies |\mathcal{F}^{-1}(\psi)(x)| \leq K(1 + |x|)^{-M}.$$

Now, notice that for any $\mathbf{f} \in E_R$, we have $\mathbf{g} := R^{-n}\mathbf{f}(R^{-1}\cdot) \in E_1$. Hence, using the result from the first part of the proof, we have for $\mathbf{f} \in E_R$,

$$\begin{aligned}
\|\varphi(D)\mathbf{f}\|_{L^p(W)}^p &= \int_{\mathbb{R}^n} |W^{1/p}(t)\varphi(D)\mathbf{f}(t)|^p dt \\
&= R^{-n} \int_{\mathbb{R}^n} |W^{1/p}(R^{-1}u)(\varphi(D)\mathbf{f})(R^{-1}u)|^p du \\
&= R^{-n+np} \int_{\mathbb{R}^n} |W^{1/p}(R^{-1}u)(\psi(D)\mathbf{g})(u)|^p du \\
&\leq CR^{-n+np} \int_{\mathbb{R}^n} |W^{1/p}(R^{-1}u)\mathbf{g}(u)|^p du \\
&= CR^{-n} \int_{\mathbb{R}^n} |W^{1/p}(R^{-1}u)\mathbf{f}(R^{-1}u)|^p du \\
&= C \int_{\mathbb{R}^n} |W^{1/p}(t)\mathbf{f}(t)|^p dt,
\end{aligned}$$

where $C := C([W(R^{-1}\cdot)]_{\mathbf{A}_p(\mathbb{R}^n)}, K, p)$. However, as mentioned in Remark 2.3, the matrix \mathbf{A}_p condition is dilation invariant in the sense that $[W(R^{-1}\cdot)]_{\mathbf{A}_p(\mathbb{R}^n)} = [W]_{\mathbf{A}_p(\mathbb{R}^n)}$ for any $R > 0$, making C independent of R . This completes the proof. \square

Remark 2.5. As mentioned in the introduction, in case $1 \leq p < \infty$, a corresponding multiplier result holds without any assumptions on the spectrum of the multiplier nor on the spectrum of \mathbf{f} , see [7, Lemma 4.4]: Suppose $W \in \mathbf{A}_p(\mathbb{R}^n)$ and assume there is a constant K such that $\varphi : \mathbb{R}^n \rightarrow \mathbb{C}$ satisfies

$$|\mathcal{F}^{-1}(\varphi)(x)| \leq KR^n(1 + R|x|)^{-n-1}, \quad x \in \mathbb{R}^n,$$

then there exists a finite constant $C := C([W]_{\mathbf{A}_p(\mathbb{R}^n)}, K, p)$ such that

$$\|\varphi(D)\mathbf{f}\|_{L^p(W)} \leq C\|\mathbf{f}\|_{L^p(W)}, \quad \mathbf{f} \in L^p(W).$$

3. AN APPLICATION: MATRIX-WEIGHTED SMOOTHNESS SPACES

An important application of matrix-weighted L^p -spaces is to the construction of various matrix-weighted smoothness spaces obtained by imposing suitable weighted L^p -restrictions on local components of a vector-function. The local components are often defined with an aim to capture local frequency content of the (vector-)function and can naturally be obtained by applying a family of suitable bandlimited Fourier multipliers (frequency filters) compatible with a desired decomposition of the frequency space.

Roudenko was the first to apply such an approach in the matrix weighted setup, see [13], where she introduced a very natural notion of matrix-weighted Besov spaces $B_{p,q}^s(W)$ based on dyadic decompositions of vector-functions. This work was later extended by Frazier and Roudenko [6, 7] to matrix-weighted Triebel-Lizorkin spaces and Besov spaces for the full range $0 < p < \infty$. A very extensive recent study by Bu et al. of various properties of matrix-weighted Besov and Triebel-Lizorkin type spaces can be found in [2–4].

We shall only touch upon one aspect of this rather involved theory here, namely the issue of such smoothness spaces being well-defined.

Let $W \in \mathbf{A}_p(\mathbb{R}^n)$ for some $0 < p < \infty$, and let $\beta > 0$ be the doubling exponent from Eq. (2.3) associated with W .

We say that $\Psi := \{\psi_j\}_{j \in \mathbb{Z}}$ is a bounded admissible partition of unity if it is a smooth resolution of the identity on $\mathbb{R}^n \setminus \{0\}$, i.e., $\sum_j \psi_j(x) \equiv 1$ on $\mathbb{R}^n \setminus \{0\}$, and there exist $c_1 := c_1(\Psi), c_2 := c_2(\Psi) > 0$ such that $\text{supp}(\psi_j) \subseteq \{x \in \mathbb{R}^n : c_1 2^j \leq |x| < c_2 2^j\}$, $j \in \mathbb{Z}$. Also suppose there exists $C := C(\Psi)$ such that

$$(3.1) \quad |\mathcal{F}^{-1}\psi_j(x)| \leq C 2^j (1 + 2^j |x|)^{-M}, \quad j \in \mathbb{Z},$$

for some $M > (n + \beta) / \min\{1, p\}$. Then we follow Frazier and Roudenko and define the homogeneous matrix-weighted Besov space $\dot{B}_{p,q}^s(W) := \dot{B}_{p,q}^s(W, \Psi)$ for $0 < q \leq \infty$ as the collection of $\mathbf{f} = (f_1, \dots, f_N)$ with $f_i \in \mathcal{S}'(\mathbb{R}^n) / \mathcal{P}$ (the tempered distributions modulo polynomials), $i = 1, \dots, N$, satisfying

$$\|\mathbf{f}\|_{\dot{B}_{p,q}^s(W)} := \left(\sum_{j=-\infty}^{\infty} 2^{jsq} \|\psi_j(D)\mathbf{f}\|_{L^p(W)}^q \right)^{1/q} < +\infty,$$

with the sum replaced by \sup_j in the case $q = \infty$. Now, let us take another bounded admissible partition of unity $\Phi := \{\varphi_j\}_{j \in \mathbb{Z}}$. Clearly, it is desirable that the construction of $\dot{B}_{p,q}^s(W)$ is well-defined in the sense that $\dot{B}_{p,q}^s(W, \Psi) = \dot{B}_{p,q}^s(W, \Phi)$ up to equivalence of norms, and this important fact is indeed proved in [6, 13], but the reader may verify that the proofs presented in [6, 13] of this fact are somewhat involved relying on a theory of almost diagonal matrices in the matrix-weighted setting developed by the same authors. We now give an alternative, more transparent, proof of the mentioned equivalence relying *solely* on Proposition 2.4.

We first notice that for $j \in \mathbb{Z}$,

$$(3.2) \quad \psi_j(D)\mathbf{f} = \psi_j(D) \sum_{k \in A_j} \varphi_k(D)\mathbf{f},$$

with $A_j = \{k \in \mathbb{Z} : \text{supp}(\psi_j) \cap \text{supp}(\varphi_k) \neq \emptyset\}$, where it is easy to verify that $\#A_j$ is bounded by a constant n_0 independent of j due to the dyadic nature of the support sets in Ψ and Φ . Hence, using the decay assumption (3.1), we may call on Proposition 2.4 (if $1 < p < \infty$, we simply use the result mentioned in Remark 2.5),

$$\|\psi_j(D)\mathbf{f}\|_{L^p(W)} \leq C \sum_{k \in A_j} \|\varphi_k(D)\mathbf{f}\|_{L^p(W)}.$$

Also observe that for $k \in A_j$, we have $2^{ks} \asymp 2^{js}$ uniformly in j . It follows from this observation that

$$2^{js} \|\psi_j(D)\mathbf{f}\|_{L^p(W)} \leq C \sum_{k \in A_j} 2^{ks} \|\varphi_k(D)\mathbf{f}\|_{L^p(W)}.$$

Using the uniform bounds on the cardinality of the sets A_j , it is then straightforward to verify that

$$\begin{aligned} \|\mathbf{f}\|_{\dot{B}_{p,q}^s(W, \Psi)} &:= \left(\sum_{j=-\infty}^{\infty} 2^{jsq} \|\psi_j(D)\mathbf{f}\|_{L^p(W)}^q \right)^{1/q} \\ &\leq C \left(\sum_{j=-\infty}^{\infty} \left(\sum_{k \in A_j} 2^{ks} \|\varphi_k(D)\mathbf{f}\|_{L^p(W)} \right)^q \right)^{1/q} \\ &\leq C' \left(\sum_{j=-\infty}^{\infty} \sum_{k \in A_j} 2^{ksq} \|\varphi_k(D)\mathbf{f}\|_{L^p(W)}^q \right)^{1/q} \end{aligned}$$

$$\begin{aligned} &\leq C'' \left(\sum_{k=-\infty}^{\infty} 2^{ksq} \|\varphi_k(D)\mathbf{f}\|_{L^p(W)}^q \right)^{1/q} \\ &=: C'' \|\mathbf{f}\|_{\dot{B}_{p,q}^s(W,\Phi)}. \end{aligned}$$

Interchanging the roles of Ψ and Φ provides the reverse estimate, yielding the wanted norm equivalence between $\dot{B}_{p,q}^s(W,\Psi)$ and $\dot{B}_{p,q}^s(W,\Phi)$.

REFERENCES

- [1] M. Bownik. Inverse volume inequalities for matrix weights. *Indiana Univ. Math. J.*, 50(1):383–410, 2001.
- [2] F. Bu, T. Hytönen, D. Yang, and W. Yuan. Matrix-weighted besov-type and triebel–lizorkin-type spaces ii: Sharp boundedness of almost diagonal operators. *arXiv preprint arXiv:2312.13548*, 2023.
- [3] F. Bu, T. Hytönen, D. Yang, and W. Yuan. Matrix-weighted besov-type and triebel–lizorkin-type spaces iii: Characterizations of molecules and wavelets, trace theorems, and boundedness of pseudo-differential operators and calderón–zygmund operators. *arXiv preprint arXiv:2312.13549*, 2023.
- [4] F. Bu, T. P. Hytönen, D. Yang, and W. Yuan. Matrix-weighted besov-type and triebel–lizorkin-type spaces i: a_p -dimensions of matrix weights and φ -transform characterizations. *arXiv preprint arXiv:2304.00292*, 2023.
- [5] M. Frazier, B. Jawerth, and G. Weiss. *Littlewood-Paley theory and the study of function spaces*, volume 79 of *CBMS Regional Conference Series in Mathematics*. Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1991.
- [6] M. Frazier and S. Roudenko. Matrix-weighted Besov spaces and conditions of A_p type for $0 < p \leq 1$. *Indiana Univ. Math. J.*, 53(5):1225–1254, 2004.
- [7] M. Frazier and S. Roudenko. Littlewood-Paley theory for matrix-weighted function spaces. *Math. Ann.*, 380(1-2):487–537, 2021.
- [8] M. Goldberg. Matrix A_p weights via maximal functions. *Pacific J. Math.*, 211(2):201–220, 2003.
- [9] L. Grafakos. *Modern Fourier analysis*, volume 250 of *Graduate Texts in Mathematics*. Springer, New York, third edition, 2014.
- [10] J. Isralowitz, H.-K. Kwon, and S. Pott. Matrix weighted norm inequalities for commutators and paraproducts with matrix symbols. *J. Lond. Math. Soc. (2)*, 96(1):243–270, 2017.
- [11] J. Isralowitz, S. Pott, and S. Treil. Commutators in the two scalar and matrix weighted setting. *J. Lond. Math. Soc. (2)*, 106(1):1–26, 2022.
- [12] F. L. Nazarov and S. R. Treil'. The hunt for a Bellman function: applications to estimates for singular integral operators and to other classical problems of harmonic analysis. *Algebra i Analiz*, 8(5):32–162, 1996.
- [13] S. Roudenko. Matrix-weighted Besov spaces. *Trans. Amer. Math. Soc.*, 355(1):273–314, 2003.
- [14] S. Treil and A. Volberg. Wavelets and the angle between past and future. *J. Funct. Anal.*, 143(2):269–308, 1997.
- [15] H. Triebel. *Theory of function spaces*, volume 78 of *Monographs in Mathematics*. Birkhäuser Verlag, Basel, 1983.
- [16] A. Volberg. Matrix A_p weights via S -functions. *J. Amer. Math. Soc.*, 10(2):445–466, 1997.

DEPARTMENT OF MATHEMATICAL SCIENCES, AALBORG UNIVERSITY, SKJERNVEJ 4A, DK-9220 AALBORG EAST, DENMARK
Email address: mnielsen@math.aau.dk