

CONJUGATION GROUPS AND STRUCTURE GROUPS OF QUANDLES

VICTORIA LEBED

ABSTRACT. Quandles are certain algebraic structures showing up in different mathematical contexts. A group G with the conjugation operation forms a quandle, $\text{Conj}(G)$. In the opposite direction, one can construct a group $\text{As}(Q)$ starting from any quandle Q . These groups are useful in practice, but hard to compute. We explore the group $\text{As}(\text{Conj}(G))$ for so-called \overline{C} -groups G . These are groups admitting a presentation with only conjugation and power relations. Symmetric groups S_n are typical examples. We show that for \overline{C} -groups, $\text{As}(\text{Conj}(G))$ injects into $G \times \mathbb{Z}^m$, where m is the number of conjugacy classes of G . From this we deduce information about the torsion, center, and derived group of $\text{As}(\text{Conj}(G))$. As an application, we compute the second quandle homology group of $\text{Conj}(S_n)$ for all n , and unveil rich torsion therein.

1. INTRODUCTION

A *quandle* is a set Q endowed with an idempotent binary operation \triangleleft for which all right translations are quandle automorphisms. In other words, it satisfies the following axioms for all $a, b, c \in Q$:

1. idempotence: $a \triangleleft a = a$;
2. bijectivity of the right translations $R_c: Q \rightarrow Q, a \mapsto a \triangleleft c$;
3. self-distributivity: $(a \triangleleft b) \triangleleft c = (a \triangleleft c) \triangleleft (b \triangleleft c)$.

Quandles and generalisations thereof have remarkably wide ranging applications: construction of invariants of knots and knotted surfaces (the three quandle axioms encoding the three Reidemeister moves), classification of Hopf algebras, construction and study of set-theoretic solutions to the Yang–Baxter equation, integration of Leibniz algebras (non-symmetric generalisations of Lie algebras). See the surveys [AG03, Kin07, Car12, EN15, Kam16, Leb18] for more detail. Recently, the relevance of quandles in theoretical physics was brought to light as well [Fri24, Doi24].

Quandles can be constructed starting from mathematical structures of very different nature. The two fundamental examples are:

1. *Alexander quandles* $\text{Al}(M)$, i.e., $\mathbb{Z}[t^{\pm 1}]$ -modules M with

$$a \triangleleft b = ta + (1 - t)b,$$

2. *conjugation quandles* $\text{Conj}(G)$, i.e., groups G with the conjugation operation

$$a \triangleleft b = b^{-1}ab.$$

2020 *Mathematics Subject Classification.* 57K12, 20B30, 20F05, 16S15, 16T25.

Key words and phrases. Conjugation quandle, structure group, quandle cohomology, symmetric group, conjugation group, Yang–Baxter equation.

The *structure group* (also called the *associated*, *enveloping*, or *adjoint group*) of a quandle (Q, \triangleleft) is defined by the following quadratic presentation:

$$(1) \quad \text{As}(Q) = \langle e_a, a \in Q \mid e_a e_b = e_b e_{a \triangleleft b}, a, b \in Q \rangle.$$

It can be considered from different viewpoints.

First, it is a deformation of a free abelian group, and inherits nice properties thereof. In particular, for finite Q , $\text{As}(Q)$ is virtually free abelian: carefully chosen powers of its generators $e_a^{d_a}$ form a central free abelian subgroup with a *Coxeter-like finite quotient* $\overline{\text{As}}(Q)$ [LV18]. This gives a rich source of nicely behaving groups.

Second, the structure group construction yields a functor from the category of quandles to that of groups, which is left adjoint to the conjugation quandle functor.

Third, $\text{As}(Q)$ is an invariant of the quandle Q , which opens quandle theory to group-theoretic tools. In particular, a sufficient understanding of $\text{As}(Q)$ allows one to compute the second *quandle homology* group $H_2(Q)$, which is another quandle invariant crucial for applications and in general difficult to compute (see the above referenced surveys for detail). This can be done via Eisermann's formula [Eis14]

$$(2) \quad H_2(Q) \cong \bigoplus_{\mathcal{O} \in \text{Orb}(Q)} (\text{Stab}_{\text{As}(Q)}(a_{\mathcal{O}}) \cap \text{Ker}(\varepsilon))_{Ab}.$$

Here $\text{Orb}(Q)$ is the set of the *orbits* of Q with respect to the action of the *inner group* $\text{Inn}(Q)$, which is the group of permutations of Q generated by the right translations R_c ; $a_{\mathcal{O}} \in Q$ is any representative of the orbit \mathcal{O} ; $\text{As}(Q)$ acts on Q via $a \cdot e_b = a \triangleleft b$; and ε is the *degree map*, which is the group morphism defined by

$$(3) \quad \begin{aligned} \varepsilon: \text{As}(Q) &\rightarrow \mathbb{Z}, \\ e_a &\mapsto 1. \end{aligned}$$

The free part of $H_2(Q)$ is known to be $\mathbb{Z}^{c(Q)(c(Q)-1)}$ [EG03, LN03]. Here $c(Q)$ is the number of the $\text{Inn}(Q)$ -orbits of Q ; for $Q = \text{Conj}(G)$, this is precisely the number of conjugacy classes of G . It is the torsion part that is used in applications, and it can be highly non-trivial.

Structure groups are very difficult to compute in practice. We know of several notable examples only: an explicit description of the groups $\text{As}(\text{Al}(M))$ for connected (with respect to the $\text{Inn}(\text{Al}(M))$ action) Alexander quandles [Cla10], resulting in a computation of H_2 for some of them [GIV16]; a classification of all quandles with free abelian structure groups, together with partial results on the structure groups and homology of abelian quandles (having commuting right translations) [LM21]; and full or partial results for certain more involved classes of quandles in [Nos17]. Note that in all these cases, H_2 does have torsion in general.

The original aim of this work was to compute the structure group and the second homology group of the conjugation quandle of a symmetric group S_n . The tools we developed turned out to apply to a much wider class of groups, which we call \overline{C} -groups. These are groups admitting a presentation with only conjugation relations ($b^{-1}ab = c$) and power relations ($a^k = 1$, $k \geq 2$). Without losing generality, we allow only one power relation per conjugacy class. If only conjugation relations appear, we talk about *conjugation groups*, or briefly *C-groups*. Among \overline{C} -groups one finds free and free abelian groups; Artin and Coxeter groups, and various generalisations thereof (virtual, welded, homotopy, etc.) [BPT23, Dar24]; cactus and twin groups, and various generalisations thereof [NNS24, Mos23, CN24];

Thompson's group F [Cho23, Szy24]; knot groups; structure groups and their finite quotients; and many more. Indeed, the commutation relation, for instance, can be interpreted as $b^{-1}ab = a$, the braid relation $aba = bab$ as $b^{-1}ab = c$ and $a^{-1}ca = b$ (c being an extra generator), etc.

C -groups appeared in [Kul93, Kul94], where it was shown that, for any $n \geq 2$, being a C -group is the same as being the fundamental group of the complement of an n -dimensional compact orientable manifold without boundary in \mathbb{S}^{n+2} . We borrow this term, even if it has an alternative usage in group theory. Note that these groups independently appeared in [GH95] in a different context and under a different name, that of *LOG-groups* (LOG = labelled oriented graph).

There are two easy ways to extract some partial information from an element of the structure group

$$A(G) := \text{As}(\text{Conj}(G)).$$

The first one is to abelianise. Relation (1) becomes then $e_a = e_{a \triangleleft b}$, thus all generators from the same orbit become undistinguishable. Hence a map

$$\text{Ab}: A(G) \rightarrow \mathbb{Z}^{\mathfrak{C}(G)},$$

where $\mathfrak{C}(G)$ is the set of *conjugacy classes* of G . This map works for all quandles. The second one is proper to conjugation quandles: one can return to the original group G via the group morphism defined by

$$\begin{aligned} \pi: A(G) &\rightarrow G, \\ e_a &\mapsto a. \end{aligned}$$

It turns out that for a \overline{C} -group G , the combination of these two simple maps suffices to understand the entire group $A(G)$:

Theorem A. For a \overline{C} -group G , the following map is injective:

$$\pi \times \text{Ab}: A(G) \rightarrow G \times \mathbb{Z}^{\mathfrak{C}(G)}.$$

The image of this map is described in terms of a pullback in Theorem 3.2.

As a result, various properties of the group $A(G)$ can be deduced from those of the \overline{C} -group G , as long as one understands the conjugacy classes of G reasonably well. Here are only some useful corollaries. To announce some of them, we will need to fix a \overline{C} -type presentation Π of G . These properties will be made explicit for our running example $G = S_n$, with its classical presentation.

1. The kernel of the map π is a subgroup of $\mathbb{Z}^{\mathfrak{C}(G)}$, hence free abelian. A more refined analysis (Proposition 5.3) yields an isomorphism

$$K(G) := \text{Ker}(\pi) \cong \mathbb{Z}^{\mathfrak{C}'(G)},$$

where $\mathfrak{C}'(G)$ is the set of conjugacy classes of G containing no infinite-order generators from Π . Thus $A(G)$ is a central extension of G . We study the properties of the associated group 2-cocycle from $H^2(G, \mathbb{Z}^{\mathfrak{C}'(G)})$ (Propositions 5.1 and 5.2). In particular, this yields a short exact sequence

$$0 \rightarrow \mathbb{Z}^{P(n)} \rightarrow A(S_n) \xrightarrow{\pi} S_n \rightarrow 0,$$

$P(n)$ being the *partition number* of n , which is the number of conjugacy classes of S_n .

2. The map π restricts to an isomorphism of the derived groups:

$$A(G)' \cong G'.$$

3. The torsion of $A(G)$ is $\text{Ker}(\text{Ab}) \cap \pi^{-1}(\text{Tor}(G))$, which is isomorphic to a subgroup of $\text{Tor}(G)$. For a **finite** \overline{C} -group G , $\text{Tor}(G) = G$, hence

$$\text{Tor}(A(G)) = \text{Ker}(\text{Ab}) = A(G)' \cong G'.$$

For S_n , one obtains the alternating group:

$$\text{Tor}(A(S_n)) = A(S_n)' \cong S_n' = A_n.$$

In particular, contrary to C -groups, S_n is not a subgroup of $A(S_n)$.

4. The center of $A(G)$ is

$$Z(A(G)) = \pi^{-1}(Z(G)).$$

For a centerless group, this is a subgroup of $\mathbb{Z}^{\mathfrak{e}(G)}$, hence free abelian. In particular, for $n > 2$, one gets

$$Z(A(S_n)) = K(S_n) \cong \mathbb{Z}^{P(n)}.$$

5. An element $g \in A(G)$ acts trivially on $a \in G$ if and only if $\pi(g)$ and a commute in G . Thus,

$$\text{Stab}_{A(G)}(a) = \pi^{-1}(\text{Stab}_G(a)).$$

We will develop this idea further in Proposition 6.1. Since commutation in S_n is well understood, we will use this to compute in Theorem 6.2 the torsion of $H_2(\text{Conj}(S_n))$, which turns out to be very rich.

While the above subgroup interpretation of $A(G)$ is instrumental in studying its properties, an alternative direct product type description is better adapted for other purposes. To get to that, we first recall a somewhat surprising characterisation of C -groups and certain \overline{C} -groups:

- Theorem B.* 1. [DRS23] A group G is a C -group if and only if $G \cong \text{As}(Q)$ for some quandle Q .
 2. [AHT22] A finite centerless group G is a \overline{C} -group if and only if $G \cong \overline{\text{As}}(Q)$ for some finite quandle Q .

We give short proofs of these results with our methods.

Note that the isomorphism $G \cong \text{As}(Q)$ or $\overline{\text{As}}(Q)$ yields a new presentation of G with C -type relations for **all** instead of **some** pairs of generators only.

In [SV24] the theorem was extended to *pq-generated groups* G , which allow conjugation and generalised power relations ($a^k = b^m$, $k, m \in \mathbb{Z}$) in their presentation, and to appropriate quotients of the structure group $\text{As}(\text{Conj}(G))$. Those quotients were studied in detail there.

A quandle Q satisfying the theorem is constructed explicitly, starting from a \overline{C} -type presentation Π of G . Indeed, one takes the union Γ^G of G -conjugacy classes of the generator set Γ of Π . This is reminiscent of dual braid monoids [BKL98, Bes03], generated by all “positive” conjugates of the classical generators of braid groups. The quandle Γ^G is a particular case of *Dehn quandles* from [DRS23]. When G is Thompson’s group F , we can reformulate [Szy24, Prop. 6.4] to say that Thompson’s quandle, described by generators and relations in [Szy24], is isomorphic to our Γ^F .

For our favourite groups S_n with their classical presentation, Γ^{S_n} is the set T_n of all transpositions from S_n , endowed with the conjugation operation. We thus recover the following isomorphism from [GIV16]:

$$S_n \cong \overline{\text{As}}(T_n).$$

Our next result shows that $A(G)$ is a trivial extension of the structure group $G^{\mathcal{D}} := \text{As}(\Gamma^G)$ of the much smaller quandle Γ^G :

Theorem C. For any \overline{C} -group G , one has a group isomorphism

$$A(G) \cong G^{\mathcal{D}} \times \mathbb{Z}^{\mathfrak{C}_{\text{ng}}(G)},$$

where $\mathfrak{C}_{\text{ng}}(G)$ is the set of conjugacy classes of G containing no generators from Π .

We propose several alternative descriptions of $G^{\mathcal{D}}$, adapted to different purposes.

First, it is an intermediate object between G and the *Artin-type lift* $G^{\mathcal{A}}$ obtained from a presentation Π of G by forgetting all power relations:

Theorem D. For a group G with a \overline{C} -type presentation Π , one has a sequence

$$G^{\mathcal{A}} \twoheadrightarrow G^{\mathcal{D}} \twoheadrightarrow G$$

of group surjections, where the first map imposes the element a^k to be central whenever Π contains a relation $a^k = 1$; and the second map collapses all these central elements, presenting $G^{\mathcal{D}}$ as a central extension of G :

$$(4) \quad 0 \rightarrow \mathbb{Z}^{\mathfrak{C}_{\text{fg}}(G)} \rightarrow G^{\mathcal{D}} \rightarrow G \rightarrow 0.$$

Here $\mathfrak{C}_{\text{fg}}(G)$ is the set of all conjugacy classes of G containing finite order generators.

In particular,

- for a C -group G , the three groups above coincide, and we recover the isomorphism $A(G) \cong G \times \mathbb{Z}^{\mathfrak{C}_{\text{ng}}(G)}$ from [Ryd93], where it was conjectured to characterise C -groups;
- for Coxeter groups, these surjections were studied in [Aki20] (with the notations $W = G$, $A_W = G^{\mathcal{A}}$, $Q_W = \Gamma^G$), generalising the S_n case from [AFGV11].

Second, we inject $G^{\mathcal{D}}$ into $G \times \mathbb{Z}^{\mathfrak{C}_{\text{fg}}(G)}$ in Theorem 4.3, and interpret this as a pullback in Proposition 4.4.

Finally, when all generators of G are from the same conjugacy class, Proposition 4.12 presents $G^{\mathcal{D}}$ as a semi-direct product

$$G^{\mathcal{D}} \cong G' \rtimes \mathbb{Z}.$$

For S_n , one has

$$S_n^{\mathcal{A}} \cong B_n \twoheadrightarrow S_n^{\mathcal{D}} \cong B_n \times \langle t \rangle / \sigma_1^2 = t \twoheadrightarrow S_n \cong B_n / \sigma_1^2 = 1,$$

and Theorem C yields

$$A(S_n) \cong S_n^{\mathcal{D}} \times \mathbb{Z}^{P(n)-1}.$$

Further, $S_n^{\mathcal{D}}$ is a central extension of S_n ,

$$S_n^{\mathcal{D}} \cong S_n \times_{\varphi} \mathbb{Z},$$

for the 2-cocycle

$$\begin{aligned} \varphi: \quad S_n^2 &\rightarrow \mathbb{Z}, \\ (\alpha, \beta) &\mapsto (-\text{lg}(\alpha\beta) + \text{lg}(\alpha) + \text{lg}(\beta))/2, \end{aligned}$$

where lg is the minimal number of transpositions needed to write a permutation. Finally, we recover the following description from [Eis14, Aki20]:

$$S_n^{\mathcal{D}} \cong A_n \rtimes \mathbb{Z},$$

where $1 \in \mathbb{Z}$ acts on A_n by conjugation by, say, the generator σ_1 .

2. STRUCTURE GROUPS AS SUBGROUPS

The aim of this section is to prove Theorem A, that is, to present the structure group $A(G)$ as a subgroup of the much more manageable direct product $G \times \mathbb{Z}^{\mathfrak{C}(G)}$.

Fix a group G with a \overline{C} -type presentation Π . Denote its set of generators by Γ . We will need several notations, relevant for all subsequent sections.

First, let us split the set of the conjugacy classes of G as

$$\mathfrak{C}(G) = \mathfrak{C}_g(G) \sqcup \mathfrak{C}_{ng}(G), \quad \mathfrak{C}_g(G) = \mathfrak{C}_{fg}(G) \sqcup \mathfrak{C}_{ig}(G),$$

where the classes from $\mathfrak{C}_g(G)$ contain some generator $a \in \Gamma$, while those from $\mathfrak{C}_{ng}(G)$ do not; and in a class $\mathcal{O} \in \mathfrak{C}_{fg}(G)$ all elements are of finite order, denoted by $k(\mathcal{O})$, while in $\mathfrak{C}_{ig}(G)$ everyone is of infinite order. Recall that we allow only one power relation per conjugacy class, so $k(\mathcal{O})$ is exactly the exponent in the only power relation $a^k = 1$ with $a \in \mathcal{O}$.

From the presentation Π , one directly deduces the abelianisation of G :

$$(5) \quad \text{Ab}(G) \cong \mathbb{Z}^{\mathfrak{C}_{ig}(G)} \oplus G_f, \quad \text{where } G_f := \bigoplus_{\mathcal{O} \in \mathfrak{C}_{fg}(G)} \mathbb{Z}_{k(\mathcal{O})}.$$

Further notations are needed to make the abelianisation map for the structure group, $\text{Ab}: A(G) \rightarrow \mathbb{Z}^{\mathfrak{C}(G)}$, more explicit. Denote by $c_{\mathcal{O}}$ the generator of $\mathbb{Z}^{\mathfrak{C}(G)}$ corresponding to $\mathcal{O} \in \mathfrak{C}(G)$; thus $\text{Ab}(e_a) = c_{\mathcal{O}}$ for $a \in \mathcal{O}$. Also, denote by $\varepsilon_{\mathcal{O}}: A(G) \rightarrow \mathbb{Z}$ the composition of Ab with the projection onto the corresponding factor of $\mathbb{Z}^{\mathfrak{C}(G)}$; thus $\varepsilon_{\mathcal{O}}(e_a) = 1$ if $a \in \mathcal{O}$, and 0 if $a \notin \mathcal{O}$. Observe that $\text{Ab} = \bigoplus_{\mathcal{O} \in \mathfrak{C}(G)} \varepsilon_{\mathcal{O}}$, and the degree map is just the sum $\varepsilon = \sum_{\mathcal{O} \in \mathfrak{C}(G)} \varepsilon_{\mathcal{O}}$.

The group $A(G)$ has a very big center. This will be discussed in more detail later; now we will need only central elements of the form e_a^k . To study those, let us recall a very general lemma, rediscovered by a remarkable number of authors:

Lemma 2.1. *If a power a^k is central in a (not necessarily \overline{C} -) group G , then the power e_a^k is central in $A(G)$, and one has $e_a^k = e_b^k$ for any b from the same conjugacy class. Taking at most one such power per conjugacy class, one generates a central free abelian subgroup of $A(G)$.*

Note that such central powers exist in the structure group of any finite quandle Q , and are used to construct the finite quotients $\overline{\text{As}}(Q)$.

Proof. The centrality statement follows from the computation

$$e_c \triangleleft (e_a^k) = e_{c \triangleleft (a^k)} = e_c$$

for all $c \in G$. Here and below the symbol \triangleleft in a group always stands for the conjugation operation. This implies

$$e_a^k = e_a^k \triangleleft e_c = (e_a \triangleleft e_c)^k = e_{a \triangleleft c}^k,$$

hence $e_a^k = e_b^k$ for any b from the same conjugacy class as a . Finally, the freeness follows from the freeness of the image in the abelianisation, since

$$\text{Ab}(e_a^k) = kc_{\mathcal{O}} \in \bigoplus_{\mathcal{O} \in \mathfrak{C}(G)} \mathbb{Z}c_{\mathcal{O}} = \mathbb{Z}^{\mathfrak{C}(G)}. \quad \square$$

For a \overline{C} -group G , to any $\mathcal{O} \in \mathfrak{C}_{fg}$ one then associates the central element

$$(6) \quad t_{\mathcal{O}} = e_a^{k(\mathcal{O})}, \quad a \in \mathcal{O},$$

which is independent of $a \in \mathcal{O}$. Indeed, the element $a^{k(\mathcal{O})} = 1$ is central in G .

Example 2.2. To help the reader better digest these notations, we write them out explicitly for the symmetric group S_n with its classical presentation, enhanced by extra generators to get to the \overline{C} -type:

$$(7) \quad S_n = \left\langle \begin{array}{ll} \sigma_i, & 1 \leq i < n, \\ \sigma_{i,i+1}, & 1 \leq i < n-1, \end{array} \left| \begin{array}{l} \sigma_i \triangleleft \sigma_j = \sigma_i, \quad i < j-1, \\ \sigma_i \triangleleft \sigma_{i+1} = \sigma_{i,i+1}, \\ \sigma_{i,i+1} \triangleleft \sigma_i = \sigma_{i+1}, \\ \sigma_i^2 = 1 \end{array} \right. \right\rangle.$$

The set of conjugacy classes $\mathfrak{C}(S_n)$ is in bijection with the set of partitions of n , which encode the cycle decomposition of permutations. All generators fall into the class $\mathcal{O}_{2,1,\dots,1}$, with $k(\mathcal{O}_{2,1,\dots,1}) = 2$. The central element $t_{\mathcal{O}_{2,1,\dots,1}}$ can be written as $e_{\sigma_i}^2$ for any i . Thus $\#\mathfrak{C}(S_n) = P(n)$ (the partition number of n), $\mathfrak{C}_{\text{ig}}(S_n)$ is empty, $\#\mathfrak{C}_{\text{fg}}(S_n) = 1$, and $(S_n)_{\text{f}} = \mathbb{Z}_2$.

Example 2.3. For the closely related braid groups B_n , a C -type presentation is obtained by removing the power relations $\sigma_i^2 = 1$ from the above presentation for S_n . The set $\mathfrak{C}(B_n)$ is now infinite, $\mathfrak{C}_{\text{fg}}(B_n)$ is empty, and $\#\mathfrak{C}_{\text{ig}}(B_n) = 1$.

Proof of Theorem A. Take an element $g \in A(G)$ satisfying $\pi(g) = 1_G$ and $\varepsilon_{\mathcal{O}}(g) = 0$ for all $\mathcal{O} \in \mathfrak{C}(G)$. We need to explain why g is trivial in $A(G)$. Our argument is based on rewriting.

Fix a representative $a_{\mathcal{O}} \in \mathcal{O}$ for any $\mathcal{O} \in \mathfrak{C}(G)$; if $\mathcal{O} \in \mathfrak{C}_{\text{g}}(G)$, we require $a_{\mathcal{O}} \in \Gamma$. Any other b from the same conjugacy class can be written as $b = a_{\mathcal{O}} \triangleleft a$ for some $a \in G$. Express a in terms of the generators of Π : $a = a_1^{\alpha_1} \cdots a_s^{\alpha_s}$, with $a_i \in \Gamma$, $\alpha_i = \pm 1$. Then

$$b = a_{\mathcal{O}} \triangleleft (a_1^{\alpha_1} \cdots a_s^{\alpha_s}) = (\cdots (a_{\mathcal{O}} \triangleleft^{\alpha_1} a_1) \cdots \triangleleft^{\alpha_s} a_s),$$

where $\triangleleft^{-1} = \triangleleft$, and the operation \triangleleft^{-1} is defined by $a \triangleleft b = c \Leftrightarrow c \triangleleft^{-1} b = a$. Then the defining relation (1) of the structure group yields

$$(8) \quad e_b = (\cdots (e_{a_{\mathcal{O}}} \triangleleft^{\alpha_1} e_{a_1}) \cdots \triangleleft^{\alpha_s} e_{a_s}).$$

Now, express our $g \in A(G)$ in terms of the generators e_a of $A(G)$. Using (1), these generators can be rearranged according to their conjugacy classes, which we order in an arbitrary way (the indices a in e_a might change along the way, but not their conjugacy class). Apply procedure (8) to all the e_a with $a \in \sqcup_{\mathcal{O} \in \mathfrak{C}_{\text{ng}}(G)} \mathcal{O}$. This presents g in terms of the $e_{a_{\mathcal{O}}}$, $\mathcal{O} \in \mathfrak{C}_{\text{ng}}(G)$, and the e_a , $a \in \sqcup_{\mathcal{O} \in \mathfrak{C}_{\text{g}}(G)} \mathcal{O}$. Using (1), the latter can be pushed through the $e_{a_{\mathcal{O}}}$ to the right of the expression, receiving \triangleleft -actions on the way. Now, apply procedure (8) once again, this time to all the e_a with $a \in \sqcup_{\mathcal{O} \in \mathfrak{C}_{\text{g}}(G)} \mathcal{O}$. The result can be re-organised as

$$(9) \quad g = e_{a_{\mathcal{O}_1}}^{d_1} e_{a_{\mathcal{O}_2}}^{d_2} \cdots e_{b_1}^{\beta_1} \cdots e_{b_t}^{\beta_t}, \quad \mathcal{O}_i \in \mathfrak{C}_{\text{ng}}(G), \quad d_i \in \mathbb{Z}, \quad b_j \in \Gamma, \quad \beta_j = \pm 1,$$

where only a finite number of the d_i 's are non-zero.

On the one hand, for any $\mathcal{O}_i \in \mathfrak{C}_{\text{ng}}(G)$, one has $\varepsilon_{\mathcal{O}_i}(g) = 0$. On the other hand, (9) yields $\varepsilon_{\mathcal{O}_i}(g) = d_i$. Hence all the d_i vanish. Further, since $1_G = \pi(g) = b_1^{\beta_1} \cdots b_t^{\beta_t}$, the word $b_1^{\beta_1} \cdots b_t^{\beta_t}$ can be made trivial in G using only relations from the presentation Π . But this procedure can be imitated in $A(G)$ for the element g , since a relation $a \triangleleft b = c$ from Π has an analogue $e_a \triangleleft e_b = e_{a \triangleleft b} = e_c$ in $A(G)$, and a relation $a^k = 1$ has an analogue $e_a^k = t_{\mathcal{O}}$, where $a \in \mathcal{O} \in \mathfrak{C}_{\text{g}}(G)$, and $t_{\mathcal{O}}$ is

defined in (6). As the $t_{\mathcal{O}}$'s are central, one can keep them all at the beginning of the word, and the rewriting procedure eventually leaves us with an expression

$$g = t_{\mathcal{O}_1}^{p_1} t_{\mathcal{O}_2}^{p_2} \cdots, \quad \mathcal{O}_i \in \mathfrak{C}_{\text{fg}}(G), \quad p_i \in \mathbb{Z}.$$

As usual, only a finite number of the p_i 's are non-zero. Now, for any \mathcal{O}_i , which is from $\mathfrak{C}_{\text{fg}}(G)$ this time, one has $0 = \varepsilon_{\mathcal{O}_i}(g) = p_i k(\mathcal{O}_i)$. Hence all the p_i 's vanish, and $g = 1_{A(G)}$ as desired. \square

Remark 2.4. We actually proved a stronger (but less elegant) group inclusion

$$A(G) \hookrightarrow G \times \mathbb{Z}^{\mathfrak{C}_{\text{fg}}(G)} \times \mathbb{Z}^{\mathfrak{C}_{\text{ng}}(G)},$$

obtained from the one in the theorem by ignoring the $\mathbb{Z}^{\mathfrak{C}_{\text{fg}}(G)}$ part of $\mathbb{Z}^{\mathfrak{C}(G)}$.

The theorem allows us to reduce many computations in $A(G)$ to G , especially those related to commutativity. The most straightforward example is

Corollary 2.5. *Let G be a \overline{C} -group. Then elements f and g commute in $A(G)$ if and only if $\pi(f)$ and $\pi(g)$ commute in G .*

Proof. Denoting the commutator as

$$[f, g] = fgf^{-1}g^{-1},$$

one has

$$(\pi \times \text{Ab})([f, g]) = ([\pi(f), \pi(g)], 0) \in G \times \mathbb{Z}^{\mathfrak{C}(G)},$$

so, by the injectivity of $\pi \times \text{Ab}$, we get $[f, g] = 1_{A(G)} \Leftrightarrow [\pi(f), \pi(g)] = 1_G$. \square

Similarly, one proves

Corollary 2.6. *Let G be a \overline{C} -group. Then the center of $A(G)$ can be computed as*

$$Z(A(G)) = \pi^{-1}(Z(G)).$$

In particular, for a centerless group, one has $Z(A(G)) = \text{Ker}(\pi)$.

Corollary 2.7. *Let G be a \overline{C} -group. The map π restricts to an isomorphism of the derived subgroups:*

$$A(G)' \cong G'.$$

Proof. Recall that the derived group is generated by the commutators. Since $\pi([f, g]) = [\pi(f), \pi(g)]$, the map π restricts to a map $A(G)' \rightarrow G'$. Since $\text{Ab}([f, g]) = 0$, the theorem guarantees that this restriction is injective. Since $\pi([e_a, e_b]) = [a, b]$ for all $a, b \in G$, it is also surjective. \square

Corollary 2.8. *Let G be a \overline{C} -group. The torsion of $A(G)$ is*

$$\text{Tor}(A(G)) = \text{Ker}(\text{Ab}) \cap \pi^{-1}(\text{Tor}(G)),$$

which is isomorphic to a subgroup of $\text{Tor}(G)$.

Proof. By the injectivity of $\pi \times \text{Ab}$, g is a torsion element in $A(G)$ if and only if $\pi(g)$ is a torsion element in G , and $\text{Ab}(g)$ is a torsion element in a free abelian group, which means $\text{Ab}(g) = 0$. The injectivity of $\pi \times \text{Ab}$ is then used once more to show the injectivity of the restriction of π to $\text{Ker}(\text{Ab}) \cap \pi^{-1}(\text{Tor}(G)) \rightarrow \text{Tor}(G)$. \square

This is particularly useful in the two extreme cases:

1. If G is finite (e.g., $G = S_n$), then $\text{Tor}(G) = G$, hence

$$\text{Tor}(A(G)) = \text{Ker}(\text{Ab}) = A(G)' \cong G'.$$

2. If, on the contrary, G has no torsion (e.g., $G = B_n$), then neither has $A(G)$:

$$\begin{aligned} \text{Tor}(A(G)) &= \text{Ker}(\text{Ab}) \cap \pi^{-1}(\text{Tor}(G)) = \text{Ker}(\text{Ab}) \cap \text{Ker}(\pi) \\ &= \text{Ker}(\pi \times \text{Ab}) = \{1_{A(G)}\}. \end{aligned}$$

Example 2.9. Our favourite \overline{C} -group S_n is finite and centerless for $n > 2$. The above corollaries then yield:

$$\begin{aligned} \text{Tor}(A(S_n)) &= \text{Ker}(\text{Ab}) = A(S_n)' \cong S_n' = A_n, \\ Z(A(S_n)) &= \text{Ker}(\pi). \end{aligned}$$

Here A_n is the alternating group.

Example 2.10. Our favourite C -group B_n has no torsion, hence neither has $A(B_n)$. The center of B_n is cyclic, $Z(B_n) = \langle \Delta_n^2 \rangle$ for $n > 2$, thus $Z(A(B_n))$ is generated by $\text{Ker}(\pi)$ and $e_{\Delta_n^2} \in \pi^{-1}(\Delta_n^2)$. This yields an isomorphism $Z(A(B_n)) \cong \text{Ker}(\pi) \times \mathbb{Z}$.

3. STRUCTURE GROUPS AS PULLBACKS

Let us now describe the image of the group inclusion $\pi \times \text{Ab}: A(G) \hookrightarrow G \times \mathbb{Z}^{\mathfrak{e}(G)}$ for a \overline{C} -group G .

Lemma 3.1. *There exists a unique group morphism $\overline{\pi}$ completing the following commutative square:*

$$\begin{array}{ccc} A(G) & \xrightarrow{\text{Ab}} & \mathbb{Z}^{\mathfrak{e}(G)} \\ \pi \downarrow & & \downarrow \overline{\pi} \\ G & \xrightarrow{\text{Ab}} & \mathbb{Z}^{\mathfrak{e}_{\text{ig}}(G)} \oplus G_{\text{f}} = \text{Ab}(G) \end{array}$$

(where we use the computation (5)).

Proof. Given a generator $c_{\mathcal{O}}$ of $\mathbb{Z}^{\mathfrak{e}(G)}$, take any of its Ab -preimages $\widehat{c}_{\mathcal{O}} \in A(G)$, and declare $\overline{\pi}(c_{\mathcal{O}})$ to be $\text{Ab}(\pi(\widehat{c}_{\mathcal{O}}))$. Different choices of the lift $\widehat{c}_{\mathcal{O}}$ differ by commutator factors $ghg^{-1}h^{-1}$, with $g, h \in A(G)$. But these are sent by π to commutators in G , which are killed by Ab :

$$ghg^{-1}h^{-1} \xrightarrow{\pi} \pi(g)\pi(h)\pi(g)^{-1}\pi(h)^{-1} \xrightarrow{\text{Ab}} 0.$$

Thus our $\overline{\pi}(c_{\mathcal{O}})$ does not depend on the choice of $\widehat{c}_{\mathcal{O}}$. Since the target group is abelian, we get a well defined group morphism $\overline{\pi}$. It is surjective since the $c_{\mathcal{O}}$'s with $\mathcal{O} \in \mathfrak{e}_{\text{g}}(G)$ are sent to the corresponding generators of $\text{Ab}(G)$. The square obtained is commutative by the construction of $\overline{\pi}$, and this completion of the square is unique since the upper map Ab is surjective. \square

Theorem 3.2. *For a \overline{C} -group G , the structure group $A(G)$ is the pullback*

$$A(G) \cong G \times_{\text{Ab}, \text{Ab}(G), \overline{\pi}} \mathbb{Z}^{\mathfrak{e}(G)}.$$

The above pullback will be denoted as $G \times_{\text{Ab}(G)} \mathbb{Z}^{\mathfrak{e}(G)}$ for brevity.

Example 3.3. The pullback looks far less scary in our favourite example $G = S_n$. Here $\mathbb{Z}^{\mathfrak{e}(S_n)} \cong \mathbb{Z}^{P(n)}$, $\text{Ab}(S_n) \cong \mathbb{Z}_2$, and the maps $\text{Ab}: S_n \rightarrow \mathbb{Z}_2$ (resp., $\bar{\pi}: \mathbb{Z}^{P(n)} \rightarrow \mathbb{Z}_2$) are the signature maps: they send even permutations (resp., generators corresponding to even partitions) to 0 and the remaining ones to 1. Thus $A(S_n)$ is really close to being a direct product:

$$A(S_n) \cong S_n \times_{\mathbb{Z}_2} \mathbb{Z}^{P(n)}.$$

Example 3.4. In the case of braid groups, the group $\mathbb{Z}^{\mathfrak{e}(B_n)}$ is infinitely generated; $\text{Ab}(B_n) \cong \mathbb{Z}$; the map $\text{Ab}: B_n \rightarrow \mathbb{Z}$ computes the length lg of a word written in the standard (equivalently, extended standard, in the sense of (7)) generators; and the map $\bar{\pi}: \mathbb{Z}^{\mathfrak{e}(B_n)} \rightarrow \mathbb{Z}$ sends the generator $c_{\mathcal{O}}$ to $\text{lg}(a)$ for any $a \in \mathcal{O} \subset B_n$. Hence

$$A(B_n) \cong B_n \times_{\mathbb{Z}} \mathbb{Z}^{\mathfrak{e}(B_n)}.$$

Proof of Theorem 3.2. Recall the definition of the pullback:

$$G \times_{\text{Ab}(G)} \mathbb{Z}^{\mathfrak{e}(G)} = \{ (a, x) \mid a \in G, x \in \mathbb{Z}^{\mathfrak{e}(G)}, \text{Ab}(a) = \bar{\pi}(x). \}$$

Theorem A together with Lemma 3.1 guarantee that the injection $\pi \times \text{Ab}$ actually takes values in $G \times_{\text{Ab}(G)} \mathbb{Z}^{\mathfrak{e}(G)}$. We need to show the surjectivity, that is, given

$a \in G$ and $x \in \mathbb{Z}^{\mathfrak{e}(G)}$ with $\text{Ab}(a) = \bar{\pi}(x)$, we are looking for a $g \in A(G)$ satisfying $\pi(g) = a$ and $\text{Ab}(g) = x$.

Assume first that $x \in \mathbb{Z}^{\mathfrak{e}_{\mathbb{S}}(G)}$, that is, its components indexed by $\mathcal{O} \in \mathfrak{C}_{\text{ng}}(G)$ are trivial. Write a as $a_1^{\alpha_1} \cdots a_s^{\alpha_s}$, with $a_i \in \Gamma$, $\alpha_i = \pm 1$. Put $g' = e_{a_1}^{\alpha_1} \cdots e_{a_s}^{\alpha_s}$. It is almost what we need: $\pi(g') = a$, $p_{\mathcal{O}}(\text{Ab}(g')) = 0$ for $\mathcal{O} \in \mathfrak{C}_{\text{ng}}(G)$, and

$$(10) \quad p_{\mathcal{O}}(\text{Ab}(g')) = p_{\mathcal{O}}(\text{Ab}(a)) = p_{\mathcal{O}}(\bar{\pi}(x)) = p_{\mathcal{O}}(x)$$

for $\mathcal{O} \in \mathfrak{C}_{\text{ig}}(G)$, where $p_{\mathcal{O}}$ projects $\mathbb{Z}^{\mathfrak{e}(G)}$ or $\mathbb{Z}^{\mathfrak{e}_{\text{ig}}(G)} \oplus G_{\text{f}}$ onto the component indexed by \mathcal{O} . The case of conjugacy classes from $\mathfrak{C}_{\text{fg}}(G)$ is more subtle, since in this case $p_{\mathcal{O}}(\text{Ab}(a)) = p_{\mathcal{O}}(\bar{\pi}(x)) \in \mathbb{Z}_{k(\mathcal{O})}$, whereas $p_{\mathcal{O}}(\text{Ab}(g'))$ and $p_{\mathcal{O}}(x)$ live in \mathbb{Z} . Equalities (10) then hold modulo $k(\mathcal{O})$ only. But then g' can be replaced with

$$g = g' t_{\mathcal{O}_1}^{p_1} t_{\mathcal{O}_2}^{p_2} \cdots, \quad \mathcal{O}_i \in \mathfrak{C}_{\text{fg}}(G), p_i \in \mathbb{Z},$$

with the exponents p_i chosen to get the equalities $p_{\mathcal{O}}(\text{Ab}(g)) = p_{\mathcal{O}}(x)$ in \mathbb{Z} for all $\mathcal{O} \in \mathfrak{C}_{\text{fg}}(G)$. Since $p_{\mathcal{O}}(\text{Ab}(g)) = p_{\mathcal{O}}(\text{Ab}(g')) = p_{\mathcal{O}}(x)$ for other \mathcal{O} 's, we get $\text{Ab}(g) = x$. Finally, recalling that $\pi(t_{\mathcal{O}}) = \pi(e_{a_{\mathcal{O}}}^{k(\mathcal{O})}) = a_{\mathcal{O}}^{k(\mathcal{O})} = 1_G$, we get $\pi(g) = \pi(g') = a$, and we are done.

In the general case, consider $x + y \in \mathbb{Z}^{\mathfrak{e}(G)}$, with $x \in \mathbb{Z}^{\mathfrak{e}_{\mathbb{S}}(G)}$ and $y \in \mathbb{Z}^{\mathfrak{e}_{\text{ng}}(G)}$, and $b \in G$ such that $\text{Ab}(b) = \bar{\pi}(x + y)$. Since the map Ab is surjective, there is some $h \in A(G)$ with $\text{Ab}(h) = y$. One can decompose b as $(b\pi(h)^{-1})\pi(h)$. Since

$$\text{Ab}(b\pi(h)^{-1}) = \text{Ab}(b) - \text{Ab}(\pi(h)) = \bar{\pi}(x + y) - \bar{\pi}(\text{Ab}(h)) = \bar{\pi}(x) + \bar{\pi}(y) - \bar{\pi}(y) = \bar{\pi}(x),$$

the above argument yields a $g \in A(G)$ satisfying $\pi(g) = b\pi(h)^{-1}$ and $\text{Ab}(g) = x$. But then $gh \in A(G)$ satisfies

$$\pi(gh) = \pi(g)\pi(h) = (b\pi(h)^{-1})\pi(h) = b$$

and $\text{Ab}(gh) = \text{Ab}(g) + \text{Ab}(h) = x + y$, as requested. \square

Remark 3.5. The lemma and the theorem above are in fact a particular case of Goursat's lemma, valid for any subdirect product of groups.

4. STRUCTURE GROUPS AS DIRECT PRODUCTS

In the proof of Theorem A, we have seen that generators e_a of $A(G)$ with $a \in \mathfrak{C}_{\text{ng}}(G)$ and those with $a \in \mathfrak{C}_{\text{g}}(G)$ play very different roles. We will now show how to split $A(G)$ into two parts, a trivial one corresponding to $\mathfrak{C}_{\text{ng}}(G)$, and an interesting one corresponding to $\mathfrak{C}_{\text{g}}(G)$.

To do this, we will use a \overline{C} -type presentation Π of G , with generator set Γ , to produce many elements in $\text{Ker}(\pi) \subseteq Z(A(G))$. Concretely, for any $\mathcal{O} \in \mathfrak{C}_{\text{ng}}(G)$, take an $a \in \mathcal{O}$, write it as $a = a_1^{\alpha_1} \cdots a_s^{\alpha_s}$, with $a_i \in \Gamma$, $\alpha_i = \pm 1$. Put

$$(11) \quad t_{\mathcal{O}} := e_a e_{a_s}^{-\alpha_s} \cdots e_{a_1}^{-\alpha_1} \in \text{Ker}(\pi), \quad \mathcal{O} \in \mathfrak{C}_{\text{ng}}(G).$$

Recall also the elements $t_{\mathcal{O}} \in \text{Ker}(\pi)$ defined in (6) for all $\mathcal{O} \in \mathfrak{C}_{\text{fg}}(G)$.

Proposition 4.1. *The above elements $t_{\mathcal{O}}$, with $\mathcal{O} \in \mathfrak{C}_{\text{ng}}(G) \sqcup \mathfrak{C}_{\text{fg}}(G)$, are free abelian generators of a subgroup of $\text{Ker}(\pi)$ isomorphic to $\mathbb{Z}^{\mathfrak{C}_{\text{ng}}(G)} \times \mathbb{Z}^{\mathfrak{C}_{\text{fg}}(G)}$.*

We will later prove that this recovers the whole $\text{Ker}(\pi)$.

Proof. The freeness can be seen in the abelianisation. Indeed, for $\mathcal{O} \in \mathfrak{C}_{\text{ng}}(G)$,

$$\varepsilon_{\mathcal{O}}(t_{\mathcal{O}'}) = \begin{cases} 1 & \text{if } \mathcal{O}' = \mathcal{O}, \\ 0 & \text{otherwise.} \end{cases}$$

Further for $\mathcal{O}, \mathcal{O}' \in \mathfrak{C}_{\text{fg}}(G)$,

$$\varepsilon_{\mathcal{O}}(t_{\mathcal{O}'}) = \begin{cases} k_{\mathcal{O}} & \text{if } \mathcal{O}' = \mathcal{O}, \\ 0 & \text{otherwise.} \end{cases} \quad \square$$

This is the trivial part of $A(G)$. Let us now describe the interesting part.

Definition 4.2. We call the *Dehn quandle* of (G, Π) the union Γ^G of the G -conjugacy classes of the generator set Γ of Π . Equivalently, it is the subquandle of $\text{Conj}(G)$ generated by Γ .

The structure group of this quandle will be denoted by

$$G^{\mathcal{D}} := \text{As}(\Gamma^G) = \langle e_a, a \in \Gamma^G \mid e_a e_b = e_b e_{a \triangleleft b}, a, b \in \Gamma^G \rangle.$$

The arguments from the proofs of Theorems A and 3.2 can be applied verbatim to $G^{\mathcal{D}}$, simplified significantly since one no longer needs to take the conjugacy classes from $\mathfrak{C}_{\text{ng}}(G)$ into account. This yields:

Theorem 4.3. *For a \overline{C} -group G , the map*

$$\pi \times \text{Ab}_{\text{f}}: G^{\mathcal{D}} \rightarrow G \times \mathbb{Z}^{\mathfrak{C}_{\text{fg}}(G)}$$

is injective.

Here we abusively keep the notation π for the map $G^{\mathcal{D}} \rightarrow G$, $e_a \mapsto a$; and Ab_{f} is the composition of the abelianisation map $G^{\mathcal{D}} \rightarrow \mathbb{Z}^{\mathfrak{C}_{\text{g}}(G)} = \mathbb{Z}^{\mathfrak{C}_{\text{fg}}(G)} \oplus \mathbb{Z}^{\mathfrak{C}_{\text{ig}}(G)}$ with the projection onto its part $\mathbb{Z}^{\mathfrak{C}_{\text{fg}}(G)}$ corresponding to finite-order generators.

Proposition 4.4. *For a \overline{C} -group G , the structure group $G^{\mathcal{D}}$ is the pullback*

$$G^{\mathcal{D}} \cong G \times_{\text{Ab}_{\text{f}}, G_{\text{f}}, \overline{\pi}} \mathbb{Z}^{\mathfrak{C}_{\text{fg}}(G)},$$

where :

- $\bar{\pi}: \mathbb{Z}^{\mathfrak{C}_{\text{fg}}(G)} \twoheadrightarrow G_{\text{f}} = \bigoplus_{\mathcal{O} \in \mathfrak{C}_{\text{fg}}(G)} \mathbb{Z}_{k(\mathcal{O})}$ is the obvious projection of each copy of \mathbb{Z} in $\mathbb{Z}^{\mathfrak{C}_{\text{fg}}(G)}$ indexed by $\mathcal{O} \in \mathfrak{C}_{\text{fg}}(G)$ onto $\mathbb{Z}_{k(\mathcal{O})}$;
- $\text{Ab}_{\text{f}}: G \rightarrow G_{\text{f}}$ is the composition of the abelianisation map $G \rightarrow \mathbb{Z}^{\mathfrak{C}_{\text{fg}}(G)} \oplus G_{\text{f}}$ with the projection onto G_{f} .

In other words, one has a pullback

$$\begin{array}{ccc} G^{\mathcal{D}} & \xrightarrow{\text{Ab}_{\text{f}}} & \mathbb{Z}^{\mathfrak{C}_{\text{fg}}(G)} \\ \pi \downarrow & & \downarrow \bar{\pi} \\ G & \xrightarrow{\text{Ab}_{\text{f}}} & G_{\text{f}}. \end{array}$$

Example 4.5. For the presentation (7) of S_n , $\Gamma^{S_n} \subset S_n$ is the subset T_n of all transpositions, endowed with the conjugation operation. Our pullback becomes

$$\begin{array}{ccc} S_n^{\mathcal{D}} = \text{As}(T_n) & \xrightarrow{\varepsilon} & \mathbb{Z} \\ \pi \downarrow & & \downarrow \\ S_n & \xrightarrow{\text{sign}} & \mathbb{Z}_2. \end{array}$$

The horizontal maps are abelianisations: the degree map (3), and the signature map for permutations. The rightmost map is the usual quotient $\mathbb{Z} \rightarrow \mathbb{Z}_2$.

The group $G^{\mathcal{D}}$ can be described in a more accessible way:

Theorem 4.6. Fix a $\bar{\mathcal{C}}$ -type presentation Π of a group G , with generator set Γ . A presentation of the group $G^{\mathcal{D}}$ can then be obtained from Π as follows:

- generators: e_a , $a \in \Gamma$;
- relations of two types:
 1. $e_a \triangleleft e_b = e_c$ for each conjugation relation $a \triangleleft b = c$ in Π ;
 2. $e_a^k e_b = e_b e_a^k$ for all $b \in \Gamma$, and each power relation $a^k = 1$ in Π .

Thus $G^{\mathcal{D}}$ is a relaxed version of G : the powers e_a^k are required to be central rather than trivial. We will call it the *Dehn-type lift* of G , and denote the presentation from the statement of the theorem as $\Pi^{\mathcal{D}}$.

Proof. Relations $e_a \triangleleft e_b = e_c$ hold in $G^{\mathcal{D}}$ by its definition, and the powers e_a^k are central due to Lemma 2.1. The e_a with $a \in \Gamma$ generate $G^{\mathcal{D}}$ using the rewriting mechanism from (8). It remains to check that if a word

$$w = e_{b_1}^{\beta_1} \cdots e_{b_s}^{\beta_s}, \quad b_i \in \Gamma, \beta_i = \pm 1,$$

represents a trivial element $g \in G^{\mathcal{D}}$, then it can be made trivial using relations from the presentation $\Pi^{\mathcal{D}}$ only. We will use the same rewriting strategy as in the proof of Theorem A. Since $\pi(g) = 1_G$, one can imitate a rewriting sequence transforming $b_1^{\beta_1} \cdots b_s^{\beta_s}$ into 1_G , to transform w into a finite product of terms $(e_{a_i}^{k_i})^{d_i}$, at most one for each power relation $e_{a_i}^{k_i} = 1$ in Π , using the presentation $\Pi^{\mathcal{D}}$ only. But since $\text{Ab}_{\text{f}}(g) = 0 \in \mathbb{Z}^{\mathfrak{C}_{\text{fg}}(G)}$, all the exponents p_i must be trivial. \square

One can relax the power relations $a^k = 1$ even further, by simply forgetting them. This yields an *Artin-type lift* of G :

Definition 4.7. Given a group G with a $\bar{\mathcal{C}}$ -type presentation Π , the group $G^{\mathcal{A}}$ is defined by keeping all the generators and only conjugation relations from Π .

Theorem 4.6 and Lemma 2.1 now immediately imply Theorem D. Theorem B requires slightly more work:

Proof of Theorem B. For a C -group G , the three groups above have the same presentation, hence

$$G \cong G^{\mathcal{D}} = \text{As}(\Gamma^G),$$

which proves the first part of the theorem.

Further, for a finite \overline{C} -group G , the finite quotient $\overline{\text{As}}(\Gamma^G)$ is the quotient of $\text{As}(\Gamma^G)$ by the powers e_a^k of all generators chosen to have trivial action on G . But this is equivalent to k being the minimal positive integer rendering a^k central in G . If G has trivial center, this translates as $a^k = 1$, which means that Π contains a conjugate of the power relation $a^k = 1$ (the minimality follows by passing to the abelianisation). By Theorem D, the group G can be regarded as a quotient of $G^{\mathcal{D}} = \text{As}(\Gamma^G)$ by the same elements. Therefore $G \cong \overline{\text{As}}(\Gamma^G)$. Hence the second part of the theorem. \square

Example 4.8. For S_n , the Artin-type lift gives the presentation for the braid group B_n recalled in Example 2.3. The group $S_n^{\mathcal{D}}$ is then obtained from B_n by imposing the centrality of the square of a generator, say, σ_1^2 . This is summarised as follows:

$$\begin{aligned} S_n^{\mathcal{A}} &\cong \text{As}(\Gamma^{B_n}) \cong B_n \\ &\downarrow \\ S_n^{\mathcal{D}} &= \text{As}(T_n) \cong B_n \times \langle t \rangle / \sigma_1^2 = t \\ &\downarrow \\ S_n &\cong \overline{\text{As}}(T_n) \cong B_n / \sigma_1^2 = 1 \end{aligned}$$

Here Γ^{B_n} is the set of all conjugates of the generator σ_1 in B_n .

Remark 4.9. Theorem B implies in particular that the image of the structure group functor is the category of C -groups.

We are now ready to prove the promised splitting of $A(G)$ into two parts.

Theorem 4.10. *For any \overline{C} -group G , one has a group isomorphism*

$$\begin{aligned} \phi: G^{\mathcal{D}} \times \mathbb{Z}^{\mathfrak{C}_{\text{ng}}(G)} &\xrightarrow{\sim} A(G), \\ e_a &\mapsto e_a, & a \in \Gamma^G, \\ c_{\mathcal{O}} &\mapsto t_{\mathcal{O}}, & \mathcal{O} \in \mathfrak{C}_{\text{ng}}(G), \end{aligned}$$

where $c_{\mathcal{O}}$ is the generator of $\mathbb{Z}^{\mathfrak{C}_{\text{ng}}(G)}$ corresponding to \mathcal{O} , and $t_{\mathcal{O}}$ is defined in (11).

This yields Theorem C from the Introduction.

Proof. The map ϕ is well defined because:

- the defining relations for the generators e_a in $G^{\mathcal{D}}$ are the same as for the corresponding generators in $A(G)$;
- the $t_{\mathcal{O}}$'s all lie in $\text{Ker}(\pi)$, and are thus central in $A(G)$

It is surjective since its image contains all the generators of $A(G)$:

- the e_a 's with $a \in \Gamma$ as the images of the corresponding e_a 's from $G^{\mathcal{D}}$;

- e_a for a representative a of each $\mathcal{O} \in \mathfrak{C}_{\text{ng}}$, constructed from $\phi(c_{\mathcal{O}}) = t_{\mathcal{O}}$ and from the e_{a_i} 's with $a_i \in \Gamma$;
- all other e_a 's as conjugates of the ones describes above, via (8).

To show the injectivity, pick some $(g, x) \in \text{Ker}(\phi)$. Applying to $\phi(g, x)$ the $\varepsilon_{\mathcal{O}}$'s with $\mathcal{O} \in \mathfrak{C}_{\text{ng}}(G)$, one sees that the corresponding components of $x \in \mathbb{Z}^{\mathfrak{C}_{\text{ng}}(G)}$ must be trivial, hence $x = 0$. But then, composing ϕ with $\pi \times \text{Ab}$, we see that $(\pi \times \text{Ab}_f)(g)$ is trivial. By Theorem 4.3, this yields the triviality of g . \square

An immediate consequence of the theorem is the (non-obvious!) group inclusion

$$(12) \quad \xi: \text{As}(\Gamma^G) = G^{\mathcal{D}} \hookrightarrow A(G),$$

$$e_a \mapsto e_a.$$

Since the factor $\mathbb{Z}^{\mathfrak{C}_{\text{ng}}(G)}$ is abelian, this implies

Corollary 4.11. *Let G be a \overline{C} -group. The map ξ restricts to an isomorphism of the derived subgroups:*

$$(G^{\mathcal{D}})' \cong A(G)' \cong G'.$$

We used Corollary 2.7 for the second isomorphism. Note that the groups G^A and G have different derived groups in general: for instance, $(S_n)' = A_n$ is finite, while $S_n^A \cong B_n$ has no torsion.

This property is particularly useful when all generators in Π are from the same conjugacy class:

Proposition 4.12. *Let Π be a \overline{C} -presentation of a group G such that all its generators fall into the same conjugacy class. Then $G^{\mathcal{D}}$ decomposes as a semi-direct product*

$$G^{\mathcal{D}} \cong G' \rtimes \mathbb{Z},$$

where $1 \in \mathbb{Z}$ acts on G' by conjugation by a chosen generator a of Π .

Proof. In the short exact sequence

$$0 \rightarrow \text{Ker}(\text{Ab}) \rightarrow G^{\mathcal{D}} \xrightarrow{\text{Ab}} \mathbb{Z} \rightarrow 0,$$

one can replace $\text{Ker}(\text{Ab})$ by $(G^{\mathcal{D}})'$, and then use the section

$$\mathbb{Z} \rightarrow G^{\mathcal{D}},$$

$$k \mapsto e_a^k.$$

This yields $G^{\mathcal{D}} \cong (G^{\mathcal{D}})' \rtimes \mathbb{Z}$, with $k \in \mathbb{Z}$ acting on $(G^{\mathcal{D}})'$ by conjugation by e_a^k . Under the isomorphism $(G^{\mathcal{D}})' \cong G'$, this becomes conjugation by a^k . \square

Example 4.13. For S_n , we recover the decomposition

$$S_n^{\mathcal{D}} \cong A_n \rtimes \mathbb{Z},$$

where $1 \in \mathbb{Z}$ acts on A_n by conjugation by, say, the generator σ_1 .

Finally, recall that for a C -group G , the group $G^{\mathcal{D}}$ is simply G itself. This gives

Corollary 4.14. *For any C -group G , the structure group $A(G)$ splits as*

$$A(G) \cong G \times \mathbb{Z}^{\mathfrak{C}_{\text{ng}}(G)}.$$

In other words, $\text{As}(\text{Conj}((G)))$ is only a trivial extension of G .

5. STRUCTURE GROUP $A(G)$ AS AN EXTENSION OF G

Let us now focus on the map π connecting the structure group $A(G)$ to the original group G , which we suppose to be arbitrary (not necessarily of \overline{C} -type) for the moment. The difference between the two groups is measured by the central (hence abelian) subgroup $K(G) := \text{Ker}(\pi)$ of $A(G)$, which is hard to compute in general.

The map π has an obvious *set-theoretic section*

$$\begin{aligned} s: G &\rightarrow A(G), \\ a &\mapsto e_a. \end{aligned}$$

It is not a group morphism: $e_a e_b$ and e_{ab} always have different degrees. It is classical to study the “*morphicity defect*” of such a section, which is a map

$$\begin{aligned} \varphi: G \times G &\rightarrow K(G), \\ (a, b) &\mapsto e_{ab}^{-1} e_a e_b. \end{aligned}$$

Such maps govern extensions of G by $K(G)$. This is written as

$$A(G) \cong G \times_{\varphi} K(G).$$

Proposition 5.1. *The above map φ is a symmetric equivariant group 2-cocycle. That is, it satisfies the following properties for all $a, b, c \in G$:*

- (13) $\varphi(b, c) - \varphi(ab, c) + \varphi(a, bc) - \varphi(a, b) = 0,$
- (14) $\varphi(a, 1) = e_1,$
- (15) $\varphi(a \triangleleft c, b \triangleleft c) = \varphi(a, b),$
- (16) $\varphi(a, b) = \varphi(b, a).$

Thus in principle the knowledge of the second group cohomology of G could shed light on the structure group $A(G)$.

Proof. The proof of the 2-cocycle condition (13) is classical:

$$\begin{aligned} \varphi(a, bc) + \varphi(b, c) &= e_{a(bc)}^{-1} e_a e_{bc} e_{bc}^{-1} e_b e_c = e_{abc}^{-1} e_a e_b e_c, \\ \varphi(ab, c) + \varphi(a, b) &= e_{(ab)c}^{-1} e_{ab} e_c e_{ab}^{-1} e_a e_b = e_{abc}^{-1} e_{ab} e_{ab}^{-1} e_a e_b e_c = e_{abc}^{-1} e_a e_b e_c, \end{aligned}$$

where we used the centrality of $e_{ab}^{-1} e_a e_b = \varphi(a, b)$. Note that we alternate between the multiplicative notation in the whole group $A(G)$, and the additive notation in the abelian subgroup $K(G)$.

The normalisation condition (14) is straightforward.

The equivariance condition (15) follows from

$$\begin{aligned} \varphi(a \triangleleft c, b \triangleleft c) &= e_{(a \triangleleft c)(b \triangleleft c)}^{-1} e_{a \triangleleft c} e_{b \triangleleft c} = e_{(ab) \triangleleft c}^{-1} e_{a \triangleleft c} e_{b \triangleleft c} = (e_{ab}^{-1} \triangleleft e_c)(e_a \triangleleft e_c)(e_b \triangleleft e_c) \\ &= (e_{ab}^{-1} e_a e_b) \triangleleft e_c = e_{ab}^{-1} e_a e_b, \end{aligned}$$

since the central element $e_{ab}^{-1} e_a e_b$ commutes with e_c .

The symmetry condition (16) follows from

$$\begin{aligned} \varphi(b, a) &= e_{ba}^{-1} e_b e_a = e_{a(b \triangleleft a)}^{-1} e_a (e_b \triangleleft e_a) = e_{(a \triangleleft a)(b \triangleleft a)}^{-1} e_{a \triangleleft a} e_{b \triangleleft a} \\ &= \varphi(a \triangleleft a, b \triangleleft a) = \varphi(a, b). \end{aligned}$$

We used the equivariance condition (15). □

It is remarkable that, in some sense, the 2-cocycle φ cannot be too trivial:

Proposition 5.2. *The image of φ generates the whole abelian group $K(G)$.*

Proof. Take a word w in the letters e_a , $a \in G$, representing some $g \in K(G)$. Using the relation

$$\varphi(a, b)e_{ab} = e_a e_b,$$

one can replace two neighbouring letters by one, at the cost of introducing a central factor $\varphi(a, b)^{\pm 1}$. Repeating this procedure, one gets one letter $e_d^{\pm 1}$ times some φ factors. Since $\pi(g) = 1$ and $\pi(\varphi(a, b)) = 1$, one gets $d^{\pm 1} = \pi(e_d^{\pm 1}) = 1$, hence $d = 1$. But $e_1 = \varphi(1, 1)$. In total, we have presented g as a product of the values of φ and their inverses. \square

Let us now return to \overline{C} -groups. Our understanding of the structure groups $A(G)$ in this case is sufficient to determine $K(G)$ completely.

Proposition 5.3. *A \overline{C} -group G fits into the exact sequence*

$$0 \rightarrow \mathbb{Z}^{\mathfrak{C}_{\text{ng}}(G)} \times \mathbb{Z}^{\mathfrak{C}_{\text{fg}}(G)} \xrightarrow{\iota} A(G) \xrightarrow{\pi} G \rightarrow 0,$$

where ι is the map constructed in Proposition 4.1.

Hence $K(G)$ is the free abelian group $\mathbb{Z}^{\mathfrak{C}_{\text{ng}}(G)} \times \mathbb{Z}^{\mathfrak{C}_{\text{fg}}(G)}$.

Proof. Proposition 4.1 guarantees that ι is injective, and its image lies in $\text{Ker}(\pi)$. It remains to explain why $\text{Im}(\iota)$ is the whole $\text{Ker}(\pi)$. By Theorem 4.10, $A(G)$ splits as $G^{\mathcal{D}} \times \mathbb{Z}^{\mathfrak{C}_{\text{ng}}(G)}$, where $\mathbb{Z}^{\mathfrak{C}_{\text{ng}}(G)}$ is the subgroup generated by the $t_{\mathcal{O}}$'s with $\mathcal{O} \in \mathfrak{C}_{\text{ng}}(G)$. But this is precisely $\iota(\mathbb{Z}^{\mathfrak{C}_{\text{ng}}(G)})$. Further, $\iota(\mathbb{Z}^{\mathfrak{C}_{\text{fg}}(G)})$ becomes the subgroup of $G^{\mathcal{D}}$ generated by appropriate powers of its generators, which is precisely the kernel of the map $G^{\mathcal{D}} \rightarrow G$ as described in Theorem D. \square

Example 5.4. For $G = S_n$, we get a short exact sequence

$$0 \rightarrow \mathbb{Z}^{P(n)} = \mathbb{Z}^{P(n)-1} \times \mathbb{Z} \xrightarrow{\iota} A(S_n) \xrightarrow{\pi} S_n \rightarrow 0,$$

and the 2-cocycle φ can be computed as follows. Given two permutations $\alpha, \beta \in S_n$, the $\mathbb{Z}^{P(n)-1}$ part of $\varphi(\alpha, \beta)$ is $-c_{\mathcal{O}(\alpha\beta)} + c_{\mathcal{O}(\alpha)} + c_{\mathcal{O}(\beta)}$, where $\mathcal{O}(\gamma)$ is the conjugacy class of $\gamma \in S_n$, and $c_{\mathcal{O}}$ is the generator of $\mathbb{Z}^{P(n)-1} = \mathbb{Z}^{\mathfrak{C}_{\text{ng}}(G)}$ corresponding to the class \mathcal{O} if $\mathcal{O} \in \mathfrak{C}_{\text{ng}}(G)$, and 0 otherwise. The \mathbb{Z} -part is given by the map

$$\begin{aligned} \varphi' : S_n^2 &\rightarrow \mathbb{Z}, \\ (\alpha, \beta) &\mapsto (-\text{lg}(\alpha\beta) + \text{lg}(\alpha) + \text{lg}(\beta))/2, \end{aligned}$$

where lg is the minimal number of transpositions needed to write a permutation. For instance, for two transpositions σ, τ , one has

$$\varphi'(\sigma, \tau) = \begin{cases} 0 & \text{if } \sigma \neq \tau, \\ 1 & \text{if } \sigma = \tau. \end{cases}$$

For a C -group, the map π also admits a group-theoretic section σ , which sends any generator $a \in \Gamma$ to e_a , and a general $a \in G$, written in terms of the generators a_i as $a_1^{\alpha_1} \cdots a_s^{\alpha_s}$, to $e_{a_1}^{\alpha_1} \cdots e_{a_s}^{\alpha_s}$. This is a well defined group morphism. Also, $\mathfrak{C}_{\text{fg}}(G)$ is empty for such G . We are thus left with the split sequence

$$0 \rightarrow \mathbb{Z}^{\mathfrak{C}_{\text{ng}}(G)} \xrightarrow{\iota} A(G) \xrightleftharpoons[\sigma]{\pi} G \rightarrow 0,$$

which recovers the splitting $A(G) \cong G \times \mathbb{Z}^{\mathfrak{e}_{\text{ng}}(G)}$ from Corollary 4.14. In this case the 2-cocycle φ must be a coboundary. Indeed:

Proposition 5.5. *For a C -group, one has $\varphi = df$, for the map*

$$\begin{aligned} f: G &\rightarrow K(G), \\ a &\mapsto \sigma(a)^{-1}e_a. \end{aligned}$$

Proof. The map f indeed takes values in $K(G)$, since

$$\pi(f(a)) = \pi(\sigma(a)^{-1}e_a) = \pi(\sigma(a))^{-1}\pi(e_a) = a^{-1}a = 1.$$

Further,

$$\begin{aligned} df(a, b) &= f(a) - f(ab) + f(b) = \sigma(a)^{-1}e_a e_{ab}^{-1} \sigma(ab) \sigma(b)^{-1} e_b = \sigma(a)^{-1}e_a e_{ab}^{-1} \sigma(a) e_b \\ &= e_{ab}^{-1} \sigma(a) \sigma(a)^{-1} e_a e_b = e_{ab}^{-1} e_a e_b = \varphi(a, b). \end{aligned}$$

We used the fact that σ is a group morphism, and the centrality of $\sigma(a)^{-1}e_a$. \square

6. HOMOLOGY

In this section, we will explain how our understanding of the group $A(G)$ can be used, via Eisermann's formula, to compute the second homology group of the conjugation quandle $\text{Conj}(G)$ of a \overline{C} -group G . As an illustration, we provide an explicit computation of $H_2(\text{Conj}(S_n))$.

Proposition 6.1. *Let G be a \overline{C} -group, \mathcal{O} its conjugacy class, $a_{\mathcal{O}} \in \mathcal{O}$ any representative of this class, and $\mathcal{O}_1 = \{1_G\}$ the class of 1_G . The corresponding component of formula (2) then splits as follows:*

$$(17) \quad (\text{Stab}_{A(G)}(a_{\mathcal{O}}) \cap \text{Ker}(\varepsilon))_{Ab} \cong (\text{Stab}_{G^{\mathcal{D}}}(a_{\mathcal{O}}))_{Ab} \times \mathbb{Z}^{\mathfrak{e}_{\text{ng}}(G) \setminus \{\mathcal{O}_1\}}.$$

Further, the map $\pi: G^{\mathcal{D}} \rightarrow G$ induces the short exact sequence

$$0 \rightarrow \mathbb{Z}^{\mathfrak{e}_{\text{fg}}(G)} \rightarrow (\text{Stab}_{G^{\mathcal{D}}}(a_{\mathcal{O}}))_{Ab} \rightarrow (\text{Stab}_G(a_{\mathcal{O}}))_{Ab} \rightarrow 0.$$

Thus, to determine $H_2(\text{Conj}(G))$, one can compute stabilisers in the initial group G , lift them to $G^{\mathcal{D}}$ via the set-theoretic section $a \mapsto e_a$, and establish all relations among these lifts and the free part $\mathbb{Z}^{\mathfrak{e}_{\text{fg}}(G)}$ in the abelian group $(\text{Stab}_{G^{\mathcal{D}}}(a_{\mathcal{O}}))_{Ab}$.

Proof. Recall first that both $G(A)$ and $G^{\mathcal{D}}$ act on G on the right via $a \cdot e_b = a \triangleleft b$, while G acts by conjugation: $a \cdot b = a \triangleleft b = b^{-1}ab$. In particular, for $g \in A(G)$ or $G^{\mathcal{D}}$, we have

$$(18) \quad a \cdot g = a \cdot \pi(g) = a \triangleleft \pi(g).$$

As a result, the group inclusion $G^{\mathcal{D}} \hookrightarrow A(G)$ from (12) and the map $\pi: A(G) \rightarrow G$ intertwine these actions on G .

Now, in the splitting $A(G) \cong G^{\mathcal{D}} \times \mathbb{Z}^{\mathfrak{e}_{\text{ng}}(G)}$ from Theorem C, the part $\mathbb{Z}^{\mathfrak{e}_{\text{ng}}(G)}$ lies in $\text{Ker}(\pi)$, hence acts trivially on G . Consequently,

$$\text{Stab}_{A(G)}(a_{\mathcal{O}}) \cong \text{Stab}_{G^{\mathcal{D}}}(a_{\mathcal{O}}) \times \mathbb{Z}^{\mathfrak{e}_{\text{ng}}(G)}.$$

Next, $\mathbb{Z}^{\mathfrak{e}_{\text{ng}}(G)}$ splits as $\mathbb{Z}^{\mathfrak{e}_{\text{ng}}(G) \setminus \{\mathcal{O}_1\}} \times \mathbb{Z}^{\{\mathcal{O}_1\}}$, and the last factor is generated by the central element e_1 . Since $\varepsilon(e_1) = 1$, one can mod e_1 out and get the isomorphism

$$\text{Stab}_{As(Q)}(a_{\mathcal{O}}) \cap \text{Ker}(\varepsilon) \cong \text{Stab}_{G^{\mathcal{D}}}(a_{\mathcal{O}}) \times \mathbb{Z}^{\mathfrak{e}_{\text{ng}}(G) \setminus \{\mathcal{O}_1\}}.$$

The second factor is central, hence survives in the abelianisation, yielding (17).

We now focus on the interesting factor $(\text{Stab}_{G^{\mathcal{D}}}(a_{\mathcal{O}}))_{Ab}$. According to (18), an element $g \in G^{\mathcal{D}}$ acts trivially on $a \in G$ if and only if $\pi(g)$ and a commute in G . In particular, $\text{Ker}(\pi)$ acts trivially on any a . Thus the short exact sequence (4) from Theorem D induces the short exact sequence

$$0 \rightarrow \mathbb{Z}^{\mathfrak{e}_{\text{fg}}(G)} \rightarrow \text{Stab}_{G^{\mathcal{D}}}(a_{\mathcal{O}}) \rightarrow \text{Stab}_G(a_{\mathcal{O}}) \rightarrow 0.$$

Since the group $\mathbb{Z}^{\mathfrak{e}_{\text{fg}}(G)}$ is abelian, one goes to the abelianisation and deduces the exact sequence

$$\mathbb{Z}^{\mathfrak{e}_{\text{fg}}(G)} \rightarrow (\text{Stab}_{G^{\mathcal{D}}}(a_{\mathcal{O}}))_{Ab} \rightarrow (\text{Stab}_G(a_{\mathcal{O}}))_{Ab} \rightarrow 0.$$

Finally, the leftmost map is injective, since when composed with the map

$$(\text{Stab}_{G^{\mathcal{D}}}(a_{\mathcal{O}}))_{Ab} \rightarrow (G^{\mathcal{D}})_{Ab} \cong \mathbb{Z}^{\mathfrak{e}_{\text{fg}}(G)} \times \mathbb{Z}^{\mathfrak{e}_{\text{ig}}(G)}$$

induced by the inclusion $\text{Stab}_{G^{\mathcal{D}}}(a_{\mathcal{O}}) \hookrightarrow G^{\mathcal{D}}$ it yields multiplication by $k(\mathcal{O})$ on the factor corresponding to $\mathcal{O} \in \mathfrak{e}_{\text{fg}}(G)$, which is injective. \square

Finally, we apply these techniques to our favourite example $G = S_n$. Some notations are needed to announce our result. For a partition λ of n (written as $\lambda \vdash n$), we call its *support* the set $\text{Supp}(\lambda)$ of sizes of its parts, and *repetition support* the set $\text{RSupp}(\lambda)$ of part sizes repeating in λ at least twice. Put

$$R(\lambda) = \begin{cases} \#\text{RSupp}(\lambda) - 1 & \text{if RSupp}(\lambda) \text{ contains an odd element,} \\ \#\text{RSupp}(\lambda) & \text{otherwise.} \end{cases}$$

Given an even $u \in \mathbb{N}$, let $s(n, u)$ be the number of partitions $\lambda \vdash n$ satisfying the following conditions:

- there is no odd $v \in \text{RSupp}(\lambda)$;
- $u \in \text{Supp}(\lambda)$;
- a power of 2 dividing u necessarily divides all other even $u' \in \text{Supp}(\lambda)$;
- u is minimal among even numbers satisfying the two preceding properties.

Recall also that $P(n)$ denotes the partition number of n .

Theorem 6.2. *The second homology group of the conjugation quandle of the symmetric group can be described as follows:*

$$\begin{aligned} H_2(\text{Conj}(S_n)) &\cong \mathbb{Z}^{P(n)(P(n)-1)} \times \mathbb{Z}_2^{\sum_{\lambda \vdash n} R(\lambda)} \\ &\times \prod_{\text{odd } u \leq n} \mathbb{Z}_u^{P(n-u)} \times \prod_{\text{even } u \leq n} \left(\mathbb{Z}_u^{P(n-u)-s(n,u)} \times \mathbb{Z}_{\frac{u}{2}}^{s(n,u)} \right). \end{aligned}$$

Note the presence of torsion which grows very fast with n . In particular, adding a part 1 to any partition $\lambda \vdash n$, one sees that $H_2(\text{Conj}(S_n))$ is a subgroup of $H_2(\text{Conj}(S_{n+1}))$ for any n . We get a much more interesting picture than the group homology case, where $H_2(S_n)$ is simply 0 for $n < 4$ and \mathbb{Z}_2 for $n \geq 4$.

For small n , our formulas yield

$$\begin{aligned} H_2(\text{Conj}(S_3)) &\cong \mathbb{Z}^6 \times \mathbb{Z}_3, \\ H_2(\text{Conj}(S_4)) &\cong \mathbb{Z}^{20} \times \mathbb{Z}_2^3 \times \mathbb{Z}_3, \\ H_2(\text{Conj}(S_5)) &\cong \mathbb{Z}^{42} \times \mathbb{Z}_2^3 \times \mathbb{Z}_3^2 \times \mathbb{Z}_5, \\ H_2(\text{Conj}(S_6)) &\cong \mathbb{Z}^{110} \times \mathbb{Z}_2^4 \times \mathbb{Z}_3^4 \times \mathbb{Z}_4^2 \times \mathbb{Z}_5, \\ H_2(\text{Conj}(S_7)) &\cong \mathbb{Z}^{210} \times \mathbb{Z}_2^7 \times \mathbb{Z}_3^6 \times \mathbb{Z}_4^2 \times \mathbb{Z}_5^2 \times \mathbb{Z}_7. \end{aligned}$$

Proof. Take the conjugacy class \mathcal{O}_λ of S_n indexed by a partition $\lambda = (\lambda_1, \dots, \lambda_k) \vdash n$, and its representative $a_\lambda = c_1 \cdots c_k$ written as a product of cycles c_i of length λ_i with disjoint support. The proposition above yields the isomorphism,

$$(\text{Stab}_{A(S_n)}(a_\lambda) \cap \text{Ker}(\varepsilon))_{Ab} \cong (\text{Stab}_{S_n^{\mathcal{D}}}(a_\lambda))_{Ab} \times \mathbb{Z}^{P(n)-2},$$

and the short exact sequence

$$(19) \quad 0 \rightarrow \mathbb{Z} \xrightarrow{\iota} (\text{Stab}_{S_n^{\mathcal{D}}}(a_\lambda))_{Ab} \xrightarrow{\tilde{\pi}} (\text{Stab}_{S_n}(a_\lambda))_{Ab} \rightarrow 0.$$

The subgroup $\text{Stab}_{S_n}(a_\lambda)$ of S_n is generated by the cycles c_i , and, for each $v \in \text{RSupp}(\lambda)$ appearing in λ $k_v \geq 2$ times, by a subgroup isomorphic to S_{k_v} , which permutes the supports of the v -cycles. In the abelianisation, the cycle c_i with $\lambda_i \notin \text{RSupp}(\lambda)$ yields a factor \mathbb{Z}_{λ_i} , and the $\mathbb{Z}_v^{k_v} \rtimes S_{k_v}$ component of $\text{Stab}_{S_n}(a_\lambda)$ corresponding to a $v \in \text{RSupp}(\lambda)$ yields $\mathbb{Z}_v \times \mathbb{Z}_2$. Hence

$$(20) \quad (\text{Stab}_{S_n}(a_\lambda))_{Ab} \cong \prod_{u \in \text{Supp}(\lambda)} \mathbb{Z}_u \times \prod_{v \in \text{RSupp}(\lambda)} \mathbb{Z}_2.$$

For each $u \in \text{Supp}(\lambda)$, $u \neq 1$, take a generator $c_i = c_{i;1} \cdots c_{i;u-1}$ of the corresponding component \mathbb{Z}_u written as a product of $u-1$ transpositions, and put $e_u := e_{c_{i;1}} \cdots e_{c_{i;u-1}}$ (we abusively use the same notations in a group and in its abelianisation). Finally, for $v \in \text{RSupp}(\lambda)$, denote by d_v any element of S_n permuting two cycles of length v in a_λ . It generates the copy of \mathbb{Z}_2 in $\mathbb{Z}_v \times \mathbb{Z}_2$. It can be written as a composition of v transpositions with disjoint support, $d_v = d_{v;1} \cdots d_{v;v}$. Let f_v be its lift $e_{d_{v;1}} \cdots e_{d_{v;v}}$ in $S_n^{\mathcal{D}}$. By (19), the abelian group $(\text{Stab}_{S_n^{\mathcal{D}}}(a_\lambda))_{Ab}$ is generated by the e_u , the f_v , and a generator $t = e_\tau^2$ of $\text{Im}(\iota)$, independent of the choice of transposition τ . We need to determine all relations between these generators. The most obvious ones concern the powers of these generators:

$$(21) \quad e_u^u = t^{\frac{u(u-1)}{2}}, \quad u \in \text{Supp}(\lambda), u \neq 1$$

$$(22) \quad f_v^2 = t^v, \quad v \in \text{RSupp}(\lambda).$$

Indeed, we chose (the minimal) exponents such that these powers of the generators e_u and f_v lie in $\text{Ker}(\tilde{\pi})$ and thus coincide with some powers of t , which are determined by computing degrees. Further, consider a relation

$$(23) \quad \prod_{u \in \text{Supp}(\lambda)} e_u^{\alpha_u} \prod_{v \in \text{RSupp}(\lambda)} f_v^{\beta_v} = t^\gamma, \quad \alpha_u, \beta_v, \gamma \in \mathbb{Z},$$

in $(\text{Stab}_{S_n^{\mathcal{D}}}(a_\lambda))_{Ab}$. Applying $\tilde{\pi}$ and comparing the image in (20), one sees that each α_u should be a multiple of u , and each β_v a multiple of 2. Thus relations (21)-(22) can be used to turn (23) into an equality between two powers of t , which can only be trivial since t generates a copy of \mathbb{Z} .

Let us now try to simplify relations (21)-(22) by variable substitution.

- For odd u , put $E_u := e_u t^{-\frac{(u-1)}{2}}$. Relation (21) then becomes $E_u^u = 1$.
- For even u , put $E_u := e_u t^{-\frac{(u-2)}{2}}$. Relation (21) then becomes $E_u^u = t^{\frac{u}{2}}$.
- For odd v , put $F_v := f_v t^{-\frac{(v-1)}{2}}$. Relation (22) then becomes $F_v^2 = t$.
- For even v , put $F_v := f_v t^{-\frac{v}{2}}$. Relation (22) then becomes $F_v^2 = 1$.

Case 1: There exists an odd $v_0 \in \text{RSupp}(\lambda)$. Put $\tau := F_{v_0}$. The variable $t = \tau^2$ can then be omitted. For even u , put $\overline{E}_u := E_u \tau^{-1}$. Relation (21) then becomes $\overline{E}_u^u = 1$. Also, for odd $v \neq v_0$, put $\overline{F}_v := F_v \tau^{-1}$. Relation (22)

then becomes $\overline{F}_v^2 = 1$. All relations are thus transformed into the triviality of certain powers of generators. Put together, they yield

$$(\text{Stab}_{S_n^{\mathcal{P}}}(a_\lambda))_{Ab} \cong \left(\prod_{u \in \text{Supp}(\lambda)} \mathbb{Z}_u \right) \times \mathbb{Z}_2^{\#\text{RSupp}(\lambda)-1} \times \mathbb{Z}.$$

Case 2: There is no odd $v \in \text{RSupp}(\lambda)$. Then one has a factor \mathbb{Z}_u for each odd $u \in \text{Supp}(\lambda)$, and a factor \mathbb{Z}_2 for each (necessarily even) $v \in \text{RSupp}(\lambda)$. The remaining part G_1 has as generators the E_u for even $u \in \text{Supp}(\lambda)$, together with the generator t ; they are subject to the relations $E_u^u = t^{\frac{u}{2}}$ only. Denote the values of u concerned as $2h_1, \dots, 2h_s$, $h_i \in \mathbb{N}$, arranged so that h_1 is the minimal among the h_i having the minimal $v_2(h_i)$ (where $v_2(h_i)$ is the maximal power of 2 dividing h_i). It will be denoted by $m(\lambda) := h_1$. Let us show that the abelian group G_1 isomorphic to

$$G_2 := \mathbb{Z}_{h_1} \times \mathbb{Z}_{2h_2} \times \dots \times \mathbb{Z}_{2h_s} \times \mathbb{Z}.$$

Our defining relations for G_1 can be arranged into the following matrix:

$$M_1 = \begin{bmatrix} 2h_1 & & & & -h_1 \\ & 2h_2 & & & -h_2 \\ & & \dots & & \dots \\ & & & 2h_s & -h_s \end{bmatrix}.$$

For G_2 , one gets

$$M_2 = \begin{bmatrix} h_1 & & & & 0 \\ & 2h_2 & & & 0 \\ & & \dots & & 0 \\ & & & 2h_s & 0 \end{bmatrix}.$$

By the existence and uniqueness of the Smith normal form, our two abelian groups are isomorphic if and only if, for all i , the matrices above share the same greatest common divisors (gcd) of the determinants of all their $i \times i$ minors. For M_1 , such a determinant is a product of factors $2h_q$ for distinct q , possibly times a factor h_r . So the gcd of such determinants for a fixed i is 2^{i-1} times the gcd of the products $h_{q_1} \dots h_{q_i}$ for all $1 \leq q_1 < \dots < q_i \leq n$. For M_2 , such a determinant is a product of factors $2h_q$ with distinct $q > 1$, possibly times the factor h_1 . For any prime $p > 2$, their gcd will contain the same power of p that the gcd for M_1 . For $p = 2$, one of the products containing the minimal power of 2 for both matrices is $h_1 * 2h_2 * \dots * 2h_i$. Hence the two gcd coincide. Summarising, one gets

$$(24) \quad (\text{Stab}_{S_n^{\mathcal{P}}}(a_\lambda))_{Ab} \cong \mathbb{Z}_{\frac{m(\lambda)}{2}} \times \left(\prod_{\substack{u \in \text{Supp}(\lambda) \\ u \neq m(\lambda)}} \mathbb{Z}_u \right) \times \mathbb{Z}_2^{\#\text{RSupp}(\lambda)} \times \mathbb{Z},$$

where the first term is omitted if λ contains no even parts.

By (2), the terms we have just computed need to be summed for all $\lambda \vdash n$.

- The free part of each of these $P(n)$ terms is $\mathbb{Z} \times \mathbb{Z}^{P(n)-2} \cong \mathbb{Z}^{P(n)-1}$. Hence the free part of the sum is $\mathbb{Z}^{P(n)(P(n)-1)}$, as predicted by [EG03, LN03].

- Summing the \mathbb{Z}_2 factors corresponding to the generators F_v , one gets

$$\prod_{\lambda \vdash n} \mathbb{Z}_2^{R(\lambda)} \cong \mathbb{Z}_2^{\sum_{\lambda \vdash n} R(\lambda)}.$$

- Each odd $u \in \text{Supp}(\lambda)$ brings a factor \mathbb{Z}_u . Since there are $P(n - u)$ partitions $\lambda \vdash n$ having $u \in \text{Supp}(\lambda)$, the overall contribution to H_2 is $\mathbb{Z}_u^{P(n-u)}$.
- The most delicate case is that of even $u \in \text{Supp}(\lambda)$, since they bring either a factor \mathbb{Z}_u , or, when λ has no repeating odd parts and $u = m(\lambda)$, a factor $\mathbb{Z}_{\frac{u}{2}}$. The sum of these factors is

$$\mathbb{Z}_u^{P(n-u)-s(n,u)} \times \mathbb{Z}_{\frac{u}{2}}^{s(n,u)}.$$

The sum of these four parts yields the announced result. \square

Remark 6.3. The proof easily adapts to a computation of $H_2(T_n)$ for the 1-orbit quandle T_n of all transpositions in S_n , endowed with the conjugation operation. Since $\text{As}(T_n) = S_n^{\mathcal{D}}$, one gets

$$H_2(T_n) \cong (\text{Stab}_{S_n^{\mathcal{D}}}(a_{(2,1,\dots,1)}) \cap \text{Ker}(\varepsilon))_{Ab}.$$

This is generated by elements t , e_2 , and, if $n \geq 4$, f_1 , subject to relations $e_2^2 = f_1^2 = t$. Since $\varepsilon(e_2) = \varepsilon(f_1) = 1$, intersection with $\text{Ker}(\varepsilon)$ is zero for $n \leq 3$, and is generated by the element $e_2 f_1^{-1}$ of order 2 otherwise. One obtains

$$H_2(T_n) \cong \begin{cases} 0, & n \leq 3, \\ \mathbb{Z}_2, & n \geq 4, \end{cases}$$

which recovers a computation from [GIV16].

Remark 6.4. Note that in all the cases from the proof, the torsion of $\text{Stab}_{S_n^{\mathcal{D}}}(a_\lambda)$ differs from its finite quotient $\text{Stab}_{S_n}(a_\lambda)$, which behaves more regularly, by at most one factor (one factor \mathbb{Z}_2 disappears, or one factor \mathbb{Z}_{2h} becomes \mathbb{Z}_h). This suggests that an alternative organisation of terms in the most difficult case (24) could probably lead to a more compact final formula, which we were unable to find.

Acknowledgments. The author is grateful to Carsten Dietzel, Eddy Godelle, Neha Nanda, and Markus Szymik for helpful remarks and pointers to the literature.

REFERENCES

- [AFGV11] N. Andruskiewitsch, F. Fantino, G. A. García, and L. Vendramin. On Nichols algebras associated to simple racks. In *Groups, algebras and applications*, volume 537 of *Contemp. Math.*, pages 31–56. Amer. Math. Soc., Providence, RI, 2011.
- [AG03] Nicolás Andruskiewitsch and Matías Graña. From racks to pointed Hopf algebras. *Adv. Math.*, 178(2):177–243, 2003.
- [AHT22] Toshiyuki Akita, Aoi Hasegawa, and Masayoshi Tanno. Structure of the associated groups of quandles. *Kodai Math. J.*, 45(2):270–281, 2022.
- [Aki20] Toshiyuki Akita. The adjoint group of a Coxeter quandle. *Kyoto J. Math.*, 60(4):1245–1260, 2020.
- [Bes03] David Bessis. The dual braid monoid. *Ann. Sci. École Norm. Sup. (4)*, 36(5):647–683, 2003.
- [BKL98] Joan Birman, Ki Hyoung Ko, and Sang Jin Lee. A new approach to the word and conjugacy problems in the braid groups. *Adv. Math.*, 139(2):322–353, 1998.
- [BPT23] Paolo Bellingeri, Luis Paris, and Anne-Laure Thiel. Virtual Artin groups. *Proc. Lond. Math. Soc. (3)*, 126(1):192–215, 2023.

- [Car12] J. Scott Carter. A survey of quandle ideas. In *Introductory lectures on knot theory*, volume 46 of *Ser. Knots Everything*, pages 22–53. World Sci. Publ., Hackensack, NJ, 2012.
- [Cho23] Fabienne Chouraqui. The Yang-Baxter equation and Thompson’s group F . *Internat. J. Algebra Comput.*, 33(3):547–584, 2023.
- [Cla10] F. J. B. J. Clauwens. The adjoint group of an Alexander quandle. *Preprint: arXiv:1011.1587*, 2010.
- [CN24] Hugo Chemin and Neha Nanda. Minimal presentation, finite quotients and lower central series of cactus groups. *Preprint: arXiv:2401.07924*, 2024.
- [Dar24] Jacques Darné. Milnor invariants of braids and welded braids up to homotopy. *Algebr. Geom. Topol.*, 24(3):1277–1320, 2024.
- [Doi24] Anastasia Doikou. Parametric set-theoretic yang-baxter equation: p-racks, solutions & quantum algebras. *Preprint: arXiv:2405.04088*, 2024.
- [DRS23] Neeraj K. Dhanwani, Hitesh R. Raundal, and Mahender Singh. Dehn quandles of groups and orientable surfaces. *Fund. Math.*, 263(2):167–201, 2023.
- [EG03] P. Etingof and M. Graña. On rack cohomology. *J. Pure Appl. Algebra*, 177(1):49–59, 2003.
- [Eis14] Michael Eisermann. Quandle coverings and their Galois correspondence. *Fund. Math.*, 225:103–168, 2014.
- [EN15] Mohamed Elhamdadi and Sam Nelson. *Quandles—an introduction to the algebra of knots*, volume 74 of *Student Mathematical Library*. American Mathematical Society, Providence, RI, 2015.
- [Fri24] Tobias Fritz. Self-distributive structures in physics. *Preprint: arXiv:2403.14458*, 2024.
- [GH95] N. D. Gilbert and James Howie. LOG groups and cyclically presented groups. *J. Algebra*, 174(1):118–131, 1995.
- [GIV16] Agustín García Iglesias and Leandro Vendramin. An explicit description of the second cohomology group of a quandle. *Mathematische Zeitschrift*, pages 1–23, 2016.
- [Kam16] Seiichi Kamada. Quandles and knot invariants. *Sugaku Expositions*, 29(1):17–39, 2016.
- [Kin07] Michael K. Kinyon. Leibniz algebras, Lie racks, and digroups. *J. Lie Theory*, 17(1):99–114, 2007.
- [Kul93] Vik. S. Kulikov. Alexander polynomials of plane algebraic curve. *Izv. Ross. Akad. Nauk Ser. Mat.*, 57(1):76–101, 1993.
- [Kul94] V. S. Kulikov. Geometric realization of C -groups. *Izv. Ross. Akad. Nauk Ser. Mat.*, 58(4):194–203, 1994.
- [Leb18] Victoria Lebed. Applications of self-distributivity to Yang-Baxter operators and their cohomology. *J. Knot Theory Ramifications*, 27(11):1843012, 20, 2018.
- [LM21] Victoria Lebed and Arnaud Mortier. Abelian quandles and quandles with abelian structure group. *J. Pure Appl. Algebra*, 225(1):Paper No. 106474, 22, 2021.
- [LN03] R. A. Litherland and Sam Nelson. The Betti numbers of some finite racks. *J. Pure Appl. Algebra*, 178(2):187–202, 2003.
- [LV18] Victoria Lebed and Leandro Vendramin. On structure groups of set-theoretic solutions to the Yang–Baxter equation. *Proc. Edimb. Math. Soc.*, 2018.
- [Mos23] Jacob Mostovoy. Round twin groups on few strands. *Preprint: arXiv:2303.10737*, 2023.
- [NNS24] Tushar Kanta Naik, Neha Nanda, and Mahender Singh. Virtual planar braid groups and permutations. *J. Group Theory*, 27(3):443–483, 2024.
- [Nos17] Takefumi Nosaka. Central extensions of groups and adjoint groups of quandles. In *Geometry and analysis of discrete groups and hyperbolic spaces*, volume B66 of *RIMS Kôkyûroku Bessatsu*, pages 167–184. 2017.
- [Ryd93] Hayley Ryder. The structure of racks, 1993. PhD thesis, University of Warwick.
- [SV24] Markus Szymik and Torstein Vik. Groups, conjugation and powers. *To appear in Involve*, 2024.
- [Szy24] Markus Szymik. Thompson’s quandle. *Preprint: arXiv:2407.06058*, 2024.

NORMANDIE UNIV, UNICAEN, CNRS, LMNO, 14000 CAEN, FRANCE

Email address: victoria.lebed@unicaen.fr