

COUNTING RATIONAL POINTS IN NON-ISOTROPIC NEIGHBORHOODS OF MANIFOLDS

RAJULA SRIVASTAVA

ABSTRACT. In this manuscript, we initiate the study of the number of rational points with bounded denominators, contained in a *non-isotropic* $\delta_1 \times \dots \times \delta_R$ neighborhood of a compact submanifold \mathcal{M} of codimension R in \mathbb{R}^M . We establish an upper bound for this counting function which holds when \mathcal{M} satisfies a strong curvature condition, first introduced by Schindler-Yamagishi in [21]. Further, even in the isotropic case when $\delta_1 = \dots = \delta_R = \delta$, we obtain an asymptotic formula which holds beyond the range of distance to \mathcal{M} established in [21]. Our result is also a generalization of the work of J.J. Huang [9] for hypersurfaces.

As an application, we establish for the first time an upper bound for the Hausdorff dimension of the set of weighted simultaneously well approximable points on a manifold \mathcal{M} satisfying the strong curvature condition, which agrees with the lower bound obtained by Allen-Wang in [2]. Moreover, for $R > 1$, we obtain a new upper bound for the number of rational points *on* \mathcal{M} , which goes beyond the bound in an analogue of Serre's dimension growth conjecture for submanifolds of \mathbb{R}^M .

1. INTRODUCTION

The aim of this manuscript is to establish an asymptotic formula for the number of rational points close to smooth manifolds satisfying a certain geometric condition first studied in [21]. In the case when the codimension of such a manifold is greater than one, we establish, for the first time, upper bounds on the number of rational points in a nonisotropic neighborhood of the manifold. Further, even in the isotropic case, we extend the main result of [21] (see Theorem 1.3), by obtaining an asymptotic formula which counts rational points beyond the range of distance to the manifold established in [21].

Let \mathcal{M} be a bounded immersed submanifold of \mathbb{R}^M with boundary, of dimension n and codimension R . For $Q \in \mathbb{Z}_{\geq 1}$ and $\delta \in (0, 1/2)$, we define the counting function

$$N_{\mathcal{M}}(Q, \delta) := \#\{(\mathbf{p}, q) \in \mathbb{Z}^{M+1} : 1 \leq q \leq Q, \text{dist}(\mathcal{M}, \mathbf{p}/q) \leq \delta/q\}.$$

Here dist denotes the distance with respect to the L^∞ norm on \mathbb{R}^M , that is,

$$\text{dist}(\mathcal{M}, \mathbf{p}/q) := \inf_{\mathbf{x} \in \mathcal{M}} \|\mathbf{x} - \mathbf{p}/q\|_\infty.$$

The study of rational points near manifolds has seen rapid development in the recent years. While the problem of obtaining precise asymptotics and upper bounds for $N_{\mathcal{M}}(Q, \delta)$ is interesting in its own right, it is also closely related to questions in Diophantine approximation and the dimension growth problem for submanifolds of \mathbb{R}^M (see §1.2 and Conjecture 1.7 further below).

We have the trivial upper bound

$$N_{\mathcal{M}}(Q, \delta) \leq c_{\mathcal{M}} Q^{n+1},$$

2020 *Mathematics Subject Classification*. 11D75; 11J13; 11J83; 11K55; 11J25; 11K60; 42B20.

Key words and phrases. Rational points near manifolds, Diophantine approximation on manifolds, oscillatory integrals, Hausdorff dimension.

with $c_{\mathcal{M}} > 0$ depending only on \mathcal{M} . Indeed, if \mathcal{M} is a (compact piece) of a rational hyperplane in \mathbb{R}^M , then the above estimate is the best we can hope for, as there exist constants $c'_{\mathcal{M}}, c_{\mathcal{M}}$ depending only on \mathcal{M} so that

$$c'_{\mathcal{M}}Q^{n+1} \leq N_{\mathcal{M}}(Q, \delta) \leq c_{\mathcal{M}}Q^{n+1}.$$

However, if \mathcal{M} is curved in some sense, a probabilistic heuristic suggests that

$$c'_{\mathcal{M}}\delta^R Q^{n+1} \leq N_{\mathcal{M}}(Q, \delta) \leq c_{\mathcal{M}}\delta^R Q^{n+1}. \quad (1.1)$$

We are still far from understanding the precise curvature conditions that would be sufficient for the heuristic above to be true. However, the class of nondegenerate manifolds, which arises quite frequently in Diophantine approximation, is a reasonable one to consider. Broadly speaking, a smooth (i.e., C^∞) connected submanifold of \mathbb{R}^M is nondegenerate if it is not contained in a proper affine subspace of \mathbb{R}^M .

Definition 1.1. Let $\mathcal{U}, \mathcal{U}'$ be bounded open subsets of \mathbb{R}^n with $\overline{\mathcal{U}} \subseteq \mathcal{U}'$. We say that an l -times continuously differentiable map $\Phi : \mathcal{U}' \rightarrow \mathbb{R}^M$ is l -nondegenerate at a point $\mathbf{x} \in \mathcal{U}'$ if the partial derivatives of Φ of order up to l at the point \mathbf{x} span \mathbb{R}^M . The map Φ is said to be l -nondegenerate if it is l -nondegenerate almost everywhere on \mathcal{U}' with respect to the n -dimensional Lebesgue measure. We say that an immersed manifold $\mathcal{M} := \Phi(\overline{\mathcal{U}})$ is l -nondegenerate if Φ is l -nondegenerate.

In the celebrated work [3], Beresnevich established the lower bound in (1.1) for *analytic*, nondegenerate manifolds, in the range

$$\delta > Q^{-\frac{1}{R}}.$$

In the recent work [20], Schindler, Technau and the author proved the indicated lower bound for *smooth*, nondegenerate manifolds in the range

$$\delta > Q^{-\frac{3}{2M-1}}.$$

Huxley, in [12], was the first to prove a near-optimal upper bound for sufficiently regular planar curves with non-vanishing curvature. This was followed by the remarkable work [25] of Vaughan and Velani, in which they established the sharp result for such curves under a slightly stronger regularity assumption. A recent breakthrough came in [9], where J.J. Huang proved an asymptotic for $N_{\mathcal{M}}(Q, \delta)$ when \mathcal{M} is a sufficiently smooth hypersurface with *non-vanishing Gaussian curvature*, in the optimal range

$$\delta > Q^{-1+\epsilon}. \quad (1.2)$$

Further, in [10], he made the following conjecture for submanifolds \mathcal{M} of \mathbb{R}^M of arbitrary dimension, satisfying the aforementioned nondegeneracy condition.

Conjecture 1.2 ([10], Conjecture 3.1). *Let \mathcal{M} be a bounded immersed submanifold of \mathbb{R}^M with boundary, of dimension n and codimension R . Suppose that \mathcal{M} is l -nondegenerate everywhere with $l \leq R + 1$. Then there exists a constant $c_{\mathcal{M}} > 0$ depending only on \mathcal{M} such that*

$$N_{\mathcal{M}}(Q, \delta) \leq c_{\mathcal{M}}\delta^R Q^{n+1},$$

when $\delta \geq Q^{-\frac{1}{R}+\epsilon}$ for some $\epsilon > 0$ and $Q \rightarrow \infty$.

The main theorems in [9, 25] demonstrate that non-vanishing Gaussian curvature is sufficient to establish Conjecture 1.2 for hypersurfaces. However, in [23], Technau and the author showed that the conjecture is also true for certain hypersurfaces with Gaussian curvature vanishing at a single point, provided the “degree of flatness” is below a critical value depending only on the dimension of the hypersurface. Further, when the degree of flatness is large,

[23] establishes a new asymptotic for $N_{\mathcal{M}}(Q, \delta)$ incorporating the contribution due to the “local flatness”.

For smooth, nondegenerate manifolds \mathcal{M} of arbitrary dimension, the current best upper bounds and asymptotics for a smoothed version of $N_{\mathcal{M}}(Q, \delta)$, in terms of the range of δ , are contained in [20, Theorem 1.4 and 1.6]. We also refer the reader to [5] for a previous result on upper bounds.

However, all of these results on upper bounds and asymptotics remain valid only within the range of δ prescribed by Conjecture 1.2. It therefore came as a surprise when in [21], Schindler and Yamagishi established an asymptotic for $N_{\mathcal{M}}(Q, \delta)$ for manifolds \mathcal{M} satisfying a strong curvature condition, in a range of δ which goes beyond Conjecture 1.2 when the codimension is bigger than one! To state their result (and later ours) precisely, we first need some basic set-up.

Recall that \mathcal{M} is a bounded immersed submanifold of \mathbb{R}^M with boundary, of dimension n and codimension R . Since \mathcal{M} is compact, we can work locally. Using the implicit function theorem, we may assume without loss of generality that \mathcal{M} has the parametrization

$$\mathcal{M} := \{(\mathbf{x}, f_1(\mathbf{x}), \dots, f_R(\mathbf{x})) \in \mathbb{R}^{n+R} : \mathbf{x} \in \overline{B_{\varepsilon_0}(\mathbf{x}_0)}\}. \quad (1.3)$$

Here $\mathbf{x}_0 \in \mathbb{R}^n$, $f_r : \mathbb{R}^n \rightarrow \mathbb{R}$ are C^∞ functions for $1 \leq r \leq R$ and $\overline{B_{\varepsilon_0}(\mathbf{x}_0)}$ denotes the closed ball in \mathbb{R}^n centred at \mathbf{x}_0 and of small enough radius ε_0 .

Further, we assume that \mathcal{M} satisfies the following

Curvature Condition: Given any $\mathbf{t} = (t_1, \dots, t_R) \in \mathbb{R}^R \setminus \{\mathbf{0}\}$, there exists a constant $C_{\mathbf{t}} > 0$ such that

$$\min_{\mathbf{x} \in \overline{B_{2\varepsilon_0}(\mathbf{x}_0)}} \left| \det H_{\sum_{i=1}^R t_i f_i}(\mathbf{x}) \right| > C_{\mathbf{t}}. \quad (\text{CC})$$

Note that when $R = 1$, condition (CC) reduces to $\det H_{f_1}(\mathbf{x}_0) \neq 0$, which in turn is equivalent to non-vanishing Gaussian curvature for hypersurfaces. The main result of [21] is the following.

Theorem 1.3 ([21], Corollary 1.3). *Let \mathcal{M} be as in (1.3) and let $n \geq 2$. Suppose Condition (CC) holds and that $\varepsilon_0 > 0$ is sufficiently small. Then there exists a constant $c_{\mathcal{M}} > 0$ depending only on \mathcal{M} such that*

$$N_{\mathcal{M}}(Q, \delta) \sim c_{\mathcal{M}} \delta^R Q^{n+1}$$

when

$$\delta \geq Q^{-\frac{n}{n+2(R-1)} + \epsilon} \quad (1.4)$$

for any $\epsilon > 0$ sufficiently small and $Q \rightarrow \infty$. In particular, Conjecture 1.2 holds in this case.

Note that

$$Q^{-\frac{n}{n+2(R-1)}} < Q^{-\frac{1}{R}},$$

whenever $R > 1$. Consequently, Theorem 1.3 goes beyond the range of δ hypothesized in Conjecture 1.2 for manifolds satisfying (CC) with codimension strictly bigger than one. We refer the reader to [21, Section 7] for examples of such manifolds. For a generalization of Theorem 1.3 for manifolds satisfying a less restrictive curvature condition, see [16].

1.1. Main Results. Following [9, 21, 23], we establish our results for a smoothed version of the counting function $N_{\mathcal{M}}(Q, \delta)$. The same asymptotic and bounds for $N_{\mathcal{M}}(Q, \delta)$ then follow by approximating the characteristic function of the ball $\overline{B_{\varepsilon_0}(\mathbf{x}_0)}$ by smooth weight functions as in [9, Section 7]. Let $w : \mathbb{R}^n \rightarrow [0, 1]$ be a smooth function with

$$\text{supp } w \subseteq B_{\varepsilon_0}(\mathbf{x}_0).$$

For $\delta \in (0, \frac{1}{2})$, we define

$$\mathfrak{N}_{w, \mathcal{M}}(Q, \delta) := \sum_{\substack{\mathbf{a} \in \mathbb{Z}^n \\ 1 \leq q \leq Q \\ \|qf_1(\mathbf{a}/q)\| \leq \delta \\ \vdots \\ \|qf_R(\mathbf{a}/q)\| \leq \delta}} w \left(\frac{\mathbf{a}}{q} \right). \quad (1.5)$$

We also introduce the exponent

$$\begin{aligned} \Theta &:= \max \left(\frac{n(n+R+1)}{n+2R}, n+1 - \frac{nR}{n+2(R-1) - \frac{4}{n}} \right) \\ &= \begin{cases} \frac{n(n+R+1)}{n+2R}, & 1 \leq R \leq 2. \\ n+1 - \frac{nR}{n+2(R-1) - \frac{4}{n}}, & R \geq 3. \end{cases} \end{aligned} \quad (1.6)$$

Our first result establishes an asymptotic for $\mathfrak{N}_{w, \mathcal{M}}(Q, \delta)$.

Theorem 1.4. *For $n \geq 2$ and a sufficiently small $\varepsilon_0 > 0$, let \mathcal{M} be as in (1.3). Suppose condition (CC) holds. Let Θ be as defined in (1.6). Then there exists a constant $C_{w, \mathcal{M}} > 0$ (depending only on w and \mathcal{M}) such that for all $Q \geq 1$ and $\delta \in (0, 1/2)$, we have*

$$\mathfrak{N}_{w, \mathcal{M}}(Q, \delta) = \frac{2\hat{w}(\mathbf{0})}{n} \delta^R Q^{n+1} + C_{w, \mathcal{M}} \left(\delta^{R-1} Q^{n+1 + \frac{\Theta - (n+1)}{R}} \tilde{\mathcal{E}}_n(Q)^{\frac{1}{R}} + Q^\Theta \mathcal{E}_n(Q) \right), \quad (1.7)$$

where

$$\mathcal{E}_n(Q) = \begin{cases} \exp(\mathfrak{c}_1 \sqrt{\log 4Q}), & \text{if } n = 2, R = 1 \\ (\log 4Q)^{\mathfrak{c}_2}, & \text{if } n \geq 3, R = 1 \\ \exp(\mathfrak{c}_2 (\log \log 4Q)^2) & \text{if } n \geq 3, R \geq 2 \end{cases}, \quad (1.8)$$

and

$$\tilde{\mathcal{E}}_n(Q) = \begin{cases} \exp(\mathfrak{c}_1 \sqrt{\log 4Q}), & \text{if } n = 2 \\ (\log 4Q)^{\mathfrak{c}_2} & \text{if } n \geq 3 \end{cases}, \quad (1.9)$$

for large enough constants $\mathfrak{c}_1, \mathfrak{c}_2 > 0$ depending only on w and \mathcal{M} which can be calculated from the proof.

By approximating the characteristic function of the ball $\overline{B_{\varepsilon_0}(\mathbf{x}_0)}$ by smooth weight functions from above and below using standard arguments, we obtain the following asymptotic for $N_{\mathcal{M}}(Q, \delta)$. For $R > 1$, this is valid in a range of δ beyond Theorem 1.3, and in particular, the one in Conjecture 1.2.

Corollary 1.5. *For $n \geq 2$ and a sufficiently small $\varepsilon_0 > 0$, let \mathcal{M} be as in (1.3). Suppose condition (CC) holds. Let Θ be as defined in (1.6). Then there exists a constant $c_{\mathcal{M}} > 0$ (depending only on \mathcal{M}) such that*

$$N_{\mathcal{M}}(Q, \delta) \sim c_{\mathcal{M}} \delta^R Q^{n+1} \quad (1.10)$$

whenever

$$\delta \geq Q^{\frac{\Theta - (n+1)}{R} + \epsilon} = \max \left(Q^{-\frac{n+2}{n+2R} + \epsilon}, Q^{-\frac{n}{n+2(R-1) - \frac{4}{n}} + \epsilon} \right) \quad (1.11)$$

for any sufficiently small $\epsilon > 0$ and $Q \rightarrow \infty$.

When $\delta = 0$, the term $\mathfrak{N}_{w, \mathcal{M}}(Q, 0)$ counts the weighted number of rational points with denominator bounded by Q lying on the manifold \mathcal{M} . Conjecture 1.2 would imply

$$N_{\mathcal{M}}(Q, 0) \ll N_{\mathcal{M}}(Q, Q^{-\frac{1}{R} + \epsilon}) \ll Q^{n+\epsilon R} \quad (1.12)$$

for any $\epsilon > 0$ sufficiently small, whenever \mathcal{M} is an n -dimensional bounded immersed submanifold of \mathbb{R}^M of codimension R , which is l -nondegenerate with $l \leq R+1$. One can consider the above as an analogue of Serre's dimension growth conjecture, but for submanifolds of \mathbb{R}^M of dimension n . The original formulation for irreducible projective varieties is stated below.

Conjecture 1.6 (Dimension Growth Conjecture). *Let $X \subseteq \mathbb{P}_{\mathbb{Q}}^{M-1}$ be an irreducible projective variety of degree at least two defined over \mathbb{Q} . Let $N_X(B)$ be the number of rational points on X of naive height bounded by B . Then*

$$N_X(B) \ll B^{\dim X} (\log B)^c$$

for some constant $c > 0$.

In [19], Salberger established a version of the above conjecture with B^ϵ in place of $(\log B)^c$. We refer the reader to [7, 21] and the references therein for an introduction to the topic, and for further refinements. In general, the upper bound in Conjecture 1.6 is sharp; for example, if X contains a rational linear divisor. However, by excluding divisors of small degree and imposing stronger conditions (say on the degree), it is possible to obtain a better upper bound.

In view of Conjecture 1.2 and the previous discussion, it is reasonable to formulate the following analogue of the dimension growth conjecture for the class of nondegenerate submanifolds of \mathbb{R}^M (see Definition 1.1).

Conjecture 1.7 (A Dimension Growth Conjecture for Nondegenerate Manifolds). *Let \mathcal{M} be a bounded immersed submanifold of \mathbb{R}^M with boundary, of dimension n and codimension R . Further, suppose that \mathcal{M} is l -nondegenerate for $l \leq R+1$. Then there exists a constant $c_{\mathcal{M}} > 0$ depending only on \mathcal{M} such that*

$$N_{\mathcal{M}}(Q, 0) \leq c_{\mathcal{M}} Q^{n+\epsilon},$$

for some $\epsilon > 0$ and all $Q \geq 1$.

The nondegeneracy condition, which implies that \mathcal{M} is not contained in a proper affine subspace of \mathbb{R}^M , can be considered to be a replacement for the requirement in Conjecture 1.6 that the projective variety be irreducible and of degree at least two.

In [21], as an immediate consequence of their main theorem, Schindler and Yamagishi obtained that

$$N_{\mathcal{M}}(Q, 0) \ll Q^{n - \frac{(n-2)(R-1)}{n+2(R-1)}} (\log Q)^c$$

for some constant $c > 0$, whenever the compact manifold \mathcal{M} satisfies condition (CC). In particular, for submanifolds of \mathbb{R}^M satisfying this much stronger curvature condition, their estimate broke the Q^n barrier in Conjecture 1.7.

In this paper, as a corollary of Theorem 1.4, we obtain the following improvement over [21, Corollary 1.4], which goes even further in pushing through the barrier in Conjecture 1.7 for submanifolds of \mathbb{R}^M satisfying the curvature condition (CC).

Corollary 1.8. *For $n \geq 3$ and a sufficiently small $\varepsilon_0 > 0$, let \mathcal{M} be as in (1.3). Suppose condition (CC) holds. Let Θ be as defined in (1.6), and let $\mathcal{E}_n(Q)$ be as in (1.8). Then*

$$N_{\mathcal{M}}(Q, 0) \ll Q^{\Theta} \mathcal{E}_n(Q)$$

for all $Q \geq 1$, with the implicit depending only on \mathcal{M} .

We now come to our main estimate, which is an upper bound for the number of rational points with bounded denominators contained in a non-isotropic neighborhood of a smooth manifold \mathcal{M} satisfying (CC). It specializes to the upper bound in Theorem 1.4 for isotropic

neighborhoods of \mathcal{M} . For $Q \in \mathbb{Z}_{\geq 1}$ and $\boldsymbol{\delta} = (\delta_1, \dots, \delta_R) \in (0, 1/2)^R$, we define the counting function

$$N_{\mathcal{M}}(Q, \boldsymbol{\delta}) := \#\{(\mathbf{a}, q) \in \mathbb{Z}^{n+1} : 1 \leq q \leq Q, \|qf_r(\mathbf{a}/q)\| \leq \delta_r/q \text{ for } 1 \leq r \leq R\},$$

and its smoothed version

$$\mathfrak{N}_{w, \mathcal{M}}(Q, \boldsymbol{\delta}) := \sum_{\substack{\mathbf{a} \in \mathbb{Z}^n \\ 1 \leq q \leq Q \\ \|qf_1(\mathbf{a}/q)\| \leq \delta_1 \\ \vdots \\ \|qf_R(\mathbf{a}/q)\| \leq \delta_R}} w\left(\frac{\mathbf{a}}{q}\right). \quad (1.13)$$

When $\delta_1 = \delta_2 = \dots = \delta_R = \delta$, we shall simply refer the above as $N_{\mathcal{M}}(Q, \delta)$ and $\mathfrak{N}_{w, \mathcal{M}}(Q, \delta)$ respectively, so as to be consistent with the notations for the corresponding isotropic counting functions.

We again have the trivial upper bound

$$N_{\mathcal{M}}(Q, \boldsymbol{\delta}) \leq c_{\mathcal{M}} Q^{n+1}.$$

However, using a probabilistic heuristic, we expect that

$$c'_{\mathcal{M}} \left(\prod_{r=1}^R \delta_r \right) Q^{n+1} \leq N_{\mathcal{M}}(Q, \boldsymbol{\delta}) \leq c_{\mathcal{M}} \left(\prod_{r=1}^R \delta_r \right) Q^{n+1}.$$

Let

$$\boldsymbol{\delta}^{\times} := \prod_{r=1}^R \delta_r; \quad (1.14)$$

and for $1 \leq r \leq R$, set

$$\delta_r^{\times} := \prod_{\substack{1 \leq s \leq R \\ s \neq r}} \delta_s = \frac{\boldsymbol{\delta}^{\times}}{\delta_r}. \quad (1.15)$$

Our main theorem is an upper bound for $\mathfrak{N}_{w, \mathcal{M}}(Q, \boldsymbol{\delta})$, which is the first ever non-trivial estimate for rational point count in a non-isotropic neighborhood of a submanifold of \mathbb{R}^M .

Theorem 1.9 (Main Theorem). *For $n \geq 2$ and a sufficiently small $\varepsilon_0 > 0$, let \mathcal{M} be as in (1.3). Suppose condition (CC) holds. Let Θ be as defined in (1.6). Then for all $Q \geq 1$ and $\boldsymbol{\delta} \in (0, 1/2)^R$, we have*

$$\mathfrak{N}_{w, \mathcal{M}}(Q, \boldsymbol{\delta}) \ll \boldsymbol{\delta}^{\times} Q^{n+1} + \sum_{r=1}^R \delta_r^{\times} Q^{n+1 + \frac{\Theta - (n+1)}{R}} \tilde{\mathcal{E}}_n(Q)^{\frac{1}{R}} + Q^{\Theta} \mathcal{E}_n(Q), \quad (1.16)$$

where $\mathcal{E}_n(Q)$ and $\tilde{\mathcal{E}}_n(Q)$ are as defined in (1.8) and (1.9) respectively. The implicit constant depends only on w and \mathcal{M} .

The following upper bound for $N_{\mathcal{M}}(Q, \boldsymbol{\delta})$ is a direct corollary of the above result.

Corollary 1.10. *For $n \geq 2$ and a sufficiently small $\varepsilon_0 > 0$, let \mathcal{M} be as in (1.3). Suppose condition (CC) holds. Let Θ be as defined in (1.6). Then there exists a constant $c_{\mathcal{M}} > 0$ (depending only on \mathcal{M}) such that*

$$N_{\mathcal{M}}(Q, \boldsymbol{\delta}) \leq c_{\mathcal{M}} \boldsymbol{\delta}^{\times} Q^{n+1} \quad (1.17)$$

whenever

$$\min_{1 \leq r \leq R} \delta_r \geq Q^{\frac{\Theta - (n+1)}{R} + \epsilon} = \max\left(Q^{-\frac{n+2}{n+2R} + \epsilon}, Q^{-\frac{n}{n+2(R-1) - \frac{1}{n}} + \epsilon}\right), \quad (1.18)$$

for any sufficiently small $\epsilon > 0$ and $Q \rightarrow \infty$.

1.2. Applications to Diophantine approximation. Next, we discuss applications of Theorem 1.9 to Diophantine approximation on the manifold \mathcal{M} . To do so, we need to introduce a definition and some notations.

Definition 1.11. Given a family $\boldsymbol{\psi} = (\psi_0, \psi_1, \dots, \psi_R)$ of monotonic functions $\psi_r : (0, +\infty) \rightarrow (0, 1)$ with $0 \leq r \leq R$, we call a point $\mathbf{y} \in \mathbb{R}^{n+R}$ $\boldsymbol{\psi}$ -approximable if the conditions

$$\left| y_i - \frac{a_i}{q} \right| < \frac{\psi_0(q)}{q}, \quad 1 \leq i \leq n, \quad (1.19)$$

$$\left| y_i - \frac{a_i}{q} \right| < \frac{\psi_{i-n}(q)}{q}, \quad n+1 \leq i \leq n+R, \quad (1.20)$$

hold for infinitely many $(q, \mathbf{a}) = (q, a_1, \dots, a_{n+R}) \in \mathbb{Z}_{\geq 0} \times \mathbb{Z}^M$.

We shall denote the set of $\boldsymbol{\psi}$ -approximable points in $\mathbb{R}^{n+R} = \mathbb{R}^M$ by $\mathcal{S}_{n+R}(\boldsymbol{\psi})$. For $\boldsymbol{\tau} \in \mathbb{R}_{>0}^{1+R}$, given the approximation function family $\boldsymbol{\psi}_{\boldsymbol{\tau}} = (q^{-\tau_0}, q^{-\tau_1}, \dots, q^{-\tau_R})$, we shall abbreviate notation and just write $\mathcal{S}_{n+R}(\boldsymbol{\tau}) := \mathcal{S}_{n+R}(\boldsymbol{\psi}_{\boldsymbol{\tau}})$. We shall also call this the set of $\boldsymbol{\tau}$ -approximable points, or the set of $\boldsymbol{\tau}$ -weighted simultaneously well approximable points. By Dirichlet's theorem [22], $\mathcal{S}_{n+R}(1/n, 1/n, \dots, 1/n) = \mathbb{R}^M$.

The weighted simultaneous approximation result below deals with the convergence case of Khintchine's theorem [14]. In the case when $\psi_0 = \psi_1 = \dots = \psi_R$, it complements the divergence case for *analytic* nondegenerate submanifolds of \mathbb{R}^M in [3, Theorem 2.5].

Theorem 1.12. For $n \geq 2$ and a sufficiently small $\epsilon_0 > 0$, let \mathcal{M} be as in (1.3). Suppose condition (CC) holds. Let $s > \frac{nR}{R+1}$ and let $\boldsymbol{\psi} = (\psi_0, \psi_1, \dots, \psi_R)$ be a family of monotonic approximation functions $\psi_r : (0, +\infty) \rightarrow (0, 1)$ for $0 \leq r \leq R$, with

$$\psi_0 \leq \min\{\psi_1, \dots, \psi_R\}, \quad (1.21)$$

and

$$\sum_{q=1}^{\infty} q^n \left(\frac{\psi_0(q)}{q} \right)^s \prod_{r=1}^R \psi_r(q) < \infty. \quad (1.22)$$

Then

$$\mathcal{H}^s(\mathcal{S}_{n+R}(\boldsymbol{\psi}) \cap \mathcal{M}) = 0. \quad (1.23)$$

Recall that the n -dimensional Hausdorff measure \mathcal{H}^n is a constant multiple of the Lebesgue measure μ_n . Thus, setting $s = n$ in the above theorem establishes the convergence case of Khintchine's theorem for \mathcal{M} . This is a weighted extension of [5, Theorem 1.2] (also see [20]) in the specific setting of smooth manifolds satisfying the strong condition (CC). Further, Theorem 1.12 generalizes the convergence case of [6, Theorem 2] for weighted simultaneous Diophantine approximation on planar nondegenerate curves when $\psi_0 \leq \psi_1$. It can be checked that for smooth planar curves, nondegeneracy (see Definition 1.1) is equivalent to the curvature condition (CC). We refer to [5, Proposition 2.13] for a proof of this fact.

As a corollary of Theorem 1.12, we can establish for the first time an upper bound for the set of weighted simultaneously well approximable points on \mathcal{M} corresponding to $\boldsymbol{\tau} \in \left(\frac{1}{n}, \frac{1}{R}\right)^{R+1}$. In fact, since the complementary lower bound for more general manifolds has already been established in [2, Theorem 1.1] (also see [4, Theorem 8] for an earlier result), we obtain the exact Hausdorff dimension of the set of such points.

Corollary 1.13. *For $n \geq 2$ and a sufficiently small $\varepsilon_0 > 0$, let \mathcal{M} be as in (1.3). Suppose condition (CC) holds. For $\boldsymbol{\tau} = (\tau_0, \tau_1, \dots, \tau_R) \in [\frac{1}{n}, \frac{1}{R}]^{R+1}$ with $\tau_0 \geq \max\{\tau_1, \dots, \tau_R\}$, we have*

$$\dim(\mathcal{M} \cap \mathcal{S}_{n+R}(\boldsymbol{\tau})) = \frac{n + R + 1 + \sum_{r=1}^R (\tau_0 - \tau_r)}{\tau_0 + 1} - R. \quad (1.24)$$

Note that for the much more general class of l -nondegenerate manifolds of dimension n and codimension R , in the case when $\tau_0 = \tau_1 = \dots = \tau_R$, the currently best known result is [20, Corollary 1.13], where the range of τ is given by

$$\tau < \frac{3\alpha + 1}{(2(n + R) - 1)\alpha + n + R}, \quad \text{with } \alpha := \frac{1}{n(2l - 1)(n + R + 1)}.$$

In other words, this result covers τ close to $1/n$. Corollary 1.13 is valid for a bigger range of τ , but requires the manifold \mathcal{M} to satisfy the strong curvature assumption (CC).

Proof of Corollary 1.13. Let

$$s > \frac{n + R + 1 + \sum_{r=1}^R (\tau_0 - \tau_r)}{\tau_0 + 1} - R = \frac{n + R + 1}{\tau_0 + 1} - \sum_{r=1}^R \frac{(\tau_r + 1)}{\tau_0 + 1}$$

and $\boldsymbol{\psi}(q) = (q^{-\tau_0}, q^{-\tau_1}, \dots, q^{-\tau_R})$. A straightforward calculation using the lower bound on s shows that (1.22) is convergent. We can thus apply Theorem 1.12 to conclude that $\mathcal{H}^s(\mathcal{S}_{n+R}(\boldsymbol{\tau}) \cap \mathcal{M}) = 0$, and consequently $\dim(\mathcal{S}_{n+R}(\boldsymbol{\tau}) \cap \mathcal{M}) \leq s$. Since $s > \frac{n+R+1}{\tau_0+1} - \sum_{r=1}^R \frac{(\tau_r+1)}{\tau_0+1}$ is arbitrary we conclude that

$$\dim(\mathcal{M} \cap \mathcal{S}_{n+R}(\boldsymbol{\tau})) \leq \frac{n + R + 1 + \sum_{r=1}^R (\tau_0 - \tau_r)}{\tau_0 + 1} - R. \quad (1.25)$$

To upgrade the above inequality to an equality, we can use the lower bound provided by [2, Theorem 1.1] which is true for any C^2 submanifold of \mathbb{R}^M of dimension n , whenever $\sum_{r=1}^R \tau_r < 1$. Applied to our setting, it says that

$$\dim(\mathcal{M} \cap \mathcal{S}_n(\boldsymbol{\tau})) \geq \frac{n + R + 1 + \sum_{r=1}^R (\tau_0 - \tau_r)}{\tau_0 + 1} - R. \quad (1.26)$$

Now (1.24) follows by combining (1.25) and (1.26). \square

1.3. Novelties and comparison with previous work. We compare our methods and results with the previous works of J.J. Huang (for hypersurfaces) and Schindler-Yamagishi (for manifolds satisfying the condition (CC)).

In [9], Huang used a novel combination of projective duality, stationary phase and induction on scales to develop a bootstrapping argument. Starting with the trivial estimate $N_{\mathcal{M}}(Q, \delta) \ll Q^{n+1}$, a repeated iteration of this process yielded the conjectured error term (of order Q^{n-1}) for the asymptotic expansion of $N_{\mathcal{M}}(Q, \delta)$ in the case when \mathcal{M} is a hypersurface with non-vanishing Gaussian curvature. The bedrock of this argument was a self-improving estimate relying on the fact that the Legendre dual of a hypersurface with non-vanishing Gaussian curvature is also a hypersurface with the same property. Furthermore, the Legendre transform is an involution. Thus after every two steps of this iteration, one returns to the original counting problem one started with, albeit with better estimates owing to repeated applications of stationary phase and induction on scales. It was not clear, however, whether such an argument could be adapted to manifolds of arbitrary dimension, or what duality would even mean in such a setting.

In [21], a deep insight of Schindler-Yamagishi was generalizing the notion of Legendre duality to manifolds of arbitrary dimension n and codimension R , but satisfying the geometric condition (CC). In essence, they exploit the curvature condition (CC) to “freeze” all but one codimension variables, followed by stationary phase for the family of hypersurfaces thus obtained (one for each discrete choice of the $(R - 1)$ frozen variables). This paves the way for the application of van der Corput’s B process for each such hypersurface, thus linking the manifold \mathcal{M} immersed in $\mathbb{R}^{n+R} = \mathbb{R}^M$ to a dual family of hypersurfaces in \mathbb{R}^M . Some major work is then involved in showing that these hypersurfaces also possess non-vanishing curvature. Once this is established, the authors use another application of stationary phase and duality (van der Corput’s B process) to return to the counting function for the original manifold \mathcal{M} .

However, after these two steps, the argument proceeds by using exactly one of the codimensions to project to a lower dimensional counting problem associated to a hypersurface in \mathbb{R}^{n+1} and summing trivially in the remaining in $R - 1$ codimension variables. This allows for the use of the sharp estimate for the rational point count close to hypersurfaces from [9] as a blackbox to deduce estimates for the rational point counting function associated to this family of projected hypersurfaces. The authors thus apply a similar bootstrapping procedure as in [9], but with only two steps. This is already sufficient to establish improved estimates for $\mathfrak{N}_{w,\mathcal{M}}(Q, \delta)$, in a range of δ beyond Conjecture 1.2! However, the involutive nature of Legendre duality is not fully exploited. Further, the process of summing up trivially in all but one codimension variables in the second step, does not fully utilize the information in these directions.

In this paper, we establish an inductive argument in the vein of [9] which exploits the duality between the manifold \mathcal{M} and a dual family of hypersurfaces, as formulated in [21].

- However, instead of using the main result from [9] for hypersurfaces as a blackbox, we use the involutive nature of the Legendre transform to return to the original counting problem associated to \mathcal{M} after every two steps (see Proposition 3.1). This allows us to develop a bootstrapping argument which takes as input a trivial estimate for $\mathfrak{N}_{w,\mathcal{M}}(Q, \delta)$, and eventually yields the upper bound contained in Theorem 1.9, which is valid in a bigger range of δ than in both Conjecture 1.2 and Theorem 1.3.
- Moreover, in a major departure from [9], the two inductive substeps (see Propositions 3.2 and 3.3) develop a connection between counting functions associated with two entirely different geometric objects: the manifold \mathcal{M} of codimension R immersed in \mathbb{R}^M on one hand, and a dual family of hypersurfaces in \mathbb{R}^M on the other. As mentioned previously, in [9], both counting functions were associated to hypersurfaces with non-vanishing curvature, whereas [21] only utilized this connection in one direction. In [23], Technau and the author developed such a duality argument for locally flat and rough geometric objects, but they were both required to be *hypersurfaces*.
- An important new ingredient in our argument is Proposition 3.5, which allows for passage from the sum of the dual weights (associated to hypersurfaces in \mathbb{R}^M) to the rational point count in a neighborhood of \mathcal{M} . This should be compared to [21, Proposition 5.3 and §6.3], where the original counting problem is projected to a lower dimensional one associated to a family of hypersurfaces in \mathbb{R}^{n+1} , with a trivial summing up in the remaining $R - 1$ directions. In contrast, we utilize a dyadic version of $\mathfrak{N}_{w,\mathcal{M}}(Q, \delta)$ and retain the counting problem in all the codimensions.
- In order to utilize the information from all codimensions independently, our argument *necessarily requires* estimates for a non-isotropic counting function (or else it is not possible to sum the dyadic terms (4.2) in the proof of Proposition 3.2). This inspired

us to develop our argument entirely in a non-isotropic setting, leading to Theorem 1.9.

- The class of manifolds satisfying the condition (CC) provides a fertile ground for the above ideas to be developed. However, we expect them to be useful for the problem of counting rational points near more general manifolds like curves satisfying a weaker curvature condition. We aim to address this in a future work.

1.4. Sharpness. The main result from [21] (see Theorem 1.3) broke through the $\delta > Q^{-\frac{1}{R}}$ threshold for manifolds \mathcal{M} satisfying (CC), while our results (Corollaries 1.5 and 1.10) go even beyond. A natural question is: what is the biggest range of δ in which the asymptotic/upper bound in Corollary 1.5/Corollary 1.10 holds true? The answer naturally depends on the order of the error term in the asymptotic expansion (1.7), or more generally, in the upper bound in (1.17).

In this paper, we establish that this error term is of the order of $Q^\Theta \mathcal{E}_n(Q)$, where $\mathcal{E}_n(Q)$ is as in (1.8) and

$$\Theta := \begin{cases} \frac{n(n+R+1)}{n+2R}, & 1 \leq R \leq 2. \\ n+1 - \frac{nR}{n+2(R-1) - \frac{4}{n}}, & R \geq 3. \end{cases}.$$

Proposition 3.1 states the combined effect of the two inductive substeps connecting the counting function $\mathfrak{N}_{w,\mathcal{M}}(Q, \delta)$ to its dual and vice versa (via the van der Corput B process). Neglecting logarithmic terms for the course of this discussion, a single application of Proposition 3.1 brings down the order of the error term from Q^β to $Q^{\tilde{\beta}}$, with

$$\tilde{\beta} = n+1 - \frac{nR}{n+2\left(R - \frac{n}{2\beta-n}\right)}.$$

Note that we have the trivial estimate $\mathfrak{N}_{w,\mathcal{M}}(Q, \delta) \ll \delta^\times Q^{n+1}$. Starting with $\beta_0 = n+1$, the above recursive relation yields a decreasing sequence $\{\beta_i\}_{i \geq 0}$ which converges to

$$\frac{n(n+R+1)}{n+2R}$$

after roughly $\log \log Q$ many steps, *irrespective of* whether $R \geq 2$ or $R \leq 2$. Indeed, when $R = 2$, and \mathcal{M} satisfies (CC), we conjecture that the up to logarithmic losses, the error term is of the order of

$$Q^{\frac{n(n+R+1)}{n+2R}}.$$

When $R = 1$ and \mathcal{M} is a hypersurface with non-vanishing curvature, this conjecture is true (as established in [9]); while our Theorem 1.9 establishes that same for $R = 2$.

On the other hand, for $R \geq 3$, the induction process stops once the error terms is of the order of Q^Θ with

$$\Theta = n+1 - \frac{nR}{n+2(R-1) - \frac{4}{n}}. \tag{1.27}$$

This is because below this critical value, it is the so called ‘‘error term’’ from the stationary phase expansion for the oscillatory integrals associated to the dual hypersurfaces (the last term on the right in (3.38)), which begins to dominate over the main terms (the first two terms on the right in (3.38)). In other words, for larger values of R , the order of the error term obtained from a single term stationary phase expansion determines the critical value Θ .

It is interesting to ask whether it is possible to go below even this threshold when $R \geq 3$. For hypersurfaces with non-vanishing Gaussian curvature (when $R = 1$), it is the order of the error term in the first order stationary phase expansion which determines the sharp order of

the error term in the asymptotic expansion for $N_{\mathcal{M}}(Q, \delta)$. It is thus plausible that for $R \geq 3$, the lowest possible order of the error term in the upper bound for $\mathfrak{N}_{w, \mathcal{M}}(Q, \delta)$ is indeed

$$Q^{n+1 - \frac{nR}{n+2(R-1) - \frac{4}{n}}}.$$

. At any rate, due to the obstruction just discussed, this exponent appears to be the optimal one possible using the method of stationary phase.

1.5. Outline of the paper.

- Section 2 collates technical results used throughout the paper.
- Section 3 contains an outline of the proof of Theorem 1.9.
 - We first deduce it as a consequence of Proposition 3.1.
 - Next, this proposition is broken down into two further sub steps: Propositions 3.2 and 3.3. The former uses an upper bound on the counting function associated with \mathcal{M} to obtain a better bound on a dual counting function associated to a family of n dimensional hypersurfaces in \mathbb{R}^{n+1} . Proposition 3.3 does the reverse: it bounds the counting function associated to \mathcal{M} using the upper bound on the dual counting function.
 - The proofs of Propositions 3.2 and 3.3 rely on a combination of projective duality and van der Corput's B-process to pass from \mathcal{M} to the family of dual hypersurfaces, and back. This connection is made precise in Propositions 3.4 and 3.5. The former enables passage from the sum of the dual weights to rational point count in the neighborhood of \mathcal{M} , while the latter achieves the same in the reverse direction.
- Section 4 contains the proof of Proposition 3.2 modulo Proposition 3.4 which is itself proven in Section 6. Similarly, Section 5 contains the proof of Proposition 3.3 modulo Proposition 3.5 which is proven in Section 7.
- Theorem 1.4 is proven in Section 8 using Theorem 1.9.
- Theorem 1.12 is deduced as a corollary of Theorem 1.9 in Section 9.

Remark 1.14. The condition (CC) imposes strong restrictions on how large the codimension R of \mathcal{M} can be with respect to its dimension n . Indeed, as mentioned in [21], the problem of finding n dimensional matrices satisfying (CC) is connected to question of determining the number of linearly independent vector fields on spheres. An R -tuple of symmetric matrices satisfying the condition gives rise to a system of $(R-1)$ linearly independent vector fields on the $n-1$ dimensional sphere in \mathbb{R}^n . We refer the interested reader to [1] for further reading. In particular,

$$R \leq \rho_{\text{RH}}(n), \tag{1.28}$$

where $\rho_{\text{RH}}(n)$ are Radon-Hurwitz numbers (see [11, 17]) defined as follows: if $n = (2n_1 + 1)2^{4n_2 + n_3}$ with $n_3 \in \{0, 1, 2, 3\}$ and for some $n_1 \in \{0, 1, 2, 3, \dots\}$, then $\rho_{\text{RH}}(n) = 8n_2 + 2^{n_3}$. Notice that for odd n , we have

$$R \leq \rho_{\text{RH}}(n) = 1. \tag{1.29}$$

For the interested reader, we also point out that the works of Radon and Hurwitz had originally been in the setting of Heisenberg type groups, where an analogous connection dictates how large the dimension of the centre of the Lie algebra can be as compared to the dimension of the group. We refer to [13] and [18] for further reading.

1.6. Notation. All vectors shall be denoted by boldface letters, e.g., $\mathbf{x}, \mathbf{y}, \mathbf{z}, \boldsymbol{\delta}, \mathbf{k}$. We shall use $|\cdot|$ to denote the ℓ^∞ norm of the vector under consideration. In other words, for $\mathbf{z} \in \mathbb{R}^k$ (where $k \in \mathbb{N}$),

$$|\mathbf{z}| := \max_{1 \leq i \leq k} |z_i|.$$

Given $\varepsilon > 0$ and $\mathbf{x} \in \mathbb{R}^n$, $B_\varepsilon(\mathbf{x})$ shall denote the open ball defined with respect to the metric induced by this norm, centred at \mathbf{x} and of radius ε ; i.e.,

$$B_\varepsilon(\mathbf{x}) := \{\mathbf{y} : |\mathbf{x} - \mathbf{y}| < \varepsilon\}.$$

For an open set $\mathcal{X} \subset \mathbb{R}^n$ and $\ell \in \mathbb{N}$, we shall denote the set of all ℓ -times continuously differentiable functions defined on \mathcal{X} by $C^\ell(\mathcal{X})$, the set of all smooth functions defined on \mathcal{X} by $C^\infty(\mathcal{X})$ and the set of all smooth functions defined on \mathcal{X} with compact support by $C_0^\infty(\mathcal{X})$. The gradient of a function $f \in C^1(\mathcal{X})$ shall be denoted by

$$\nabla f := \left(\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n} \right).$$

For $t \in \mathbb{R}$, we define

$$e(t) := e^{2\pi it}.$$

In the following, given a positive integer k and functions $A, B : \mathcal{X} \subset \mathbb{R}^k \rightarrow \mathbb{C}$, we shall use the notation $A \ll B$ to denote the fact that $|A(\mathbf{x})| < C|B(\mathbf{x})|$ for all $\mathbf{x} \in \mathcal{X}$, where $C > 0$ is a constant which is allowed to depend on w and \mathcal{M} , and therefore also on

- the dimension n and codimension R ,
- ε_0 and \mathbf{x}_0 ,
- the constant \mathfrak{C}_0 in condition (3.25),
- upper bounds for the functions f_r ($1 \leq r \leq R$) and w , and for their finitely many derivatives, on the domain $B_{4\varepsilon_0}(\mathbf{x}_0)$.

1.7. Acknowledgement. The author is supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy - EXC-2047/1 - 390685813 as well as SFB 1060. She is grateful to Lillian Pierce, Andreas Seeger and Niclas Technau for enlightening discussions at different stages of this project. The author also thanks Lars Becker for reading Section 1 of the paper and providing useful comments, and Jing-Jing Huang for directing her to a precise version of Conjecture 1.2.

2. PRELIMINARIES

In this section, we collect various technical results used throughout the paper.

2.1. Oscillatory Integral Estimates.

Lemma 2.1 (Non-Stationary Phase). *Let $d, K \in \mathbb{Z}_{>0}$ and $\mathcal{U} \subseteq \mathbb{R}^d$ be a bounded open set. Let $\omega \in C^\infty(\mathbb{R}^d)$, with $\text{supp}(\omega) \subseteq \mathcal{U}$ and $\varphi \in C^\infty(\mathcal{U})$ with $\nabla\varphi(\mathbf{x}) \neq \mathbf{0}$ for all $\mathbf{x} \in \text{supp}(\omega)$. Then for any $\lambda > 0$, we have*

$$\left| \int_{\mathbb{R}^d} \omega(\mathbf{x}) e(\lambda\varphi(\mathbf{x})) \, d\mathbf{x} \right| \ll \lambda^{-K+1},$$

where the implied constant depends only on K, d , upper bounds for the absolute values of finitely many derivatives of ω and φ on \mathcal{U} , and the lower bound for $|\nabla\varphi|$ on $\text{supp} \omega$.

Proof. See [8, Theorem 7.7.1] □

Lemma 2.2 (Stationary Phase). *Let $\mathcal{U}, \mathcal{U}_1 \subseteq \mathbb{R}^d$ be bounded open sets, with $\overline{\mathcal{U}} \subset \mathcal{U}_1$. Let $\omega \in C^\infty(\mathbb{R}^d)$, with $\text{supp}(\omega) \subseteq \mathcal{U}$ and $\varphi \in C^\infty(\mathcal{U}_1)$. Suppose there exists $\mathbf{v}_0 \in \mathcal{U}$ with $\nabla\varphi(\mathbf{v}_0) = 0$, and $\nabla\varphi(\mathbf{x}) \neq 0$ for all $\mathbf{x} \in \overline{\mathcal{U}} \setminus \{\mathbf{v}_0\}$. Further, let*

$$\Delta := |\det H_\varphi(\mathbf{v}_0)| \neq 0$$

and σ be the signature of $H_\varphi(\mathbf{v}_0)$. Then for any $\lambda > 0$, we have

$$\int_{\mathbb{R}^d} e(\lambda\varphi(\mathbf{x})) \omega(\mathbf{x}) d\mathbf{x} = \lambda^{-\frac{d}{2}} \Delta^{-\frac{1}{2}} e\left(\lambda\varphi(\mathbf{v}_0) + \frac{\sigma}{8}\right) (\omega(\mathbf{v}_0) + O(\lambda^{-1})), \quad (2.1)$$

where the implicit constant depends only on d , upper bounds for (the absolute values of) finitely many derivatives of ω and φ on \mathcal{U}_1 , an upper bound for

$$\sup_{\mathbf{x} \in \mathcal{U}_1 \setminus \{\mathbf{v}_0\}} \frac{|\mathbf{x} - \mathbf{v}_0|}{|\nabla\varphi(\mathbf{x})|}$$

and a lower bound for Δ .

Proof. See [24, Chapter VIII, Proposition 6] or [8, Theorem 7.7.5]. \square

2.2. Selberg Magic Functions. Let $I = (\alpha, \beta)$ be an arc of \mathbb{R}/\mathbb{Z} with $0 < \beta - \alpha < 1$, and let $\mathbb{1}_I$ denote its characteristic function. Then given $J \in \mathbb{Z}_{>0}$, there exist finite trigonometric polynomials $S_{J\pm} : [0, 1] \rightarrow \mathbb{R}$ of degree at most J

$$S_{J\pm}(x) := \sum_{j=-J}^J \hat{S}_{J\pm}(j) e(jx),$$

satisfying the following properties:

$$S_{J-}(\theta) \leq \mathbb{1}_I(\theta) \leq S_{J+}(\theta), \quad \theta \in \mathbb{R}/\mathbb{Z}; \quad (2.2)$$

$$\hat{S}_{J\pm}(0) = \beta - \alpha \pm \frac{1}{J+1}; \quad \text{and} \quad (2.3)$$

$$|\hat{S}_{J\pm}(j)| \leq \frac{1}{J+1} + \min\left(\beta - \alpha, \frac{1}{\pi|j|}\right). \quad (2.4)$$

We refer to [15, Chapter 1] for details about the construction of these functions.

2.3. The Fejér Kernel. For $D \in \mathbb{Z}_{>0}$, let $\mathcal{F}_D : [0, 1] \rightarrow \mathbb{R}$ be the Fejér kernel of degree D given by

$$\mathcal{F}_D(\theta) = \sum_{d=-D}^D \frac{D-|d|}{D^2} e(d\theta) = \left(\frac{\sin(\pi D\theta)}{D \sin(\pi\theta)}\right)^2. \quad (2.5)$$

Let $\delta^* \in (0, 1/2)$ be such that $D = \lfloor \frac{1}{2\delta^*} \rfloor$. Since $|\sin(\pi x)| \geq 2x$ for $x \in [0, 1/2]$, we have

$$\left(\frac{\sin(\pi D\theta)}{D \sin(\pi\theta)}\right)^2 \geq \left(\frac{2D\|\theta\|}{D\pi\|\theta\|}\right)^2 = \frac{4}{\pi^2}$$

whenever $\|\theta\| \in (0, \delta^*)$. In other words, letting $\mathbb{1}_{\delta^*}$ denote the characteristic function of the set $\{x \in \mathbb{R} : \|x\| \leq \delta^*\}$, we have

$$\mathbb{1}_{\delta^*}(\theta) \leq \frac{4}{\pi^2} \mathcal{F}_D(\theta). \quad (2.6)$$

3. OVERVIEW OF THE PROOF OF THEOREM 1.9

We now fix $n \geq 2$ and a sufficiently small $\varepsilon_0 > 0$. Further, define

$$\beta_{\text{st}} := \max \left(\frac{n(n+R+1)}{n+2R}, \frac{n(n+1)}{n+2} \right). \quad (3.1)$$

Observe that $\beta_{\text{st}} = \Theta$ for $R \in \{1, 2\}$, while $\beta_{\text{st}} > \Theta$ for $R \geq 3$.

Theorem 1.9 will be a consequence of the following self-improving estimate for the manifold \mathcal{M} , assumed to satisfy condition (CC).

Proposition 3.1. *Suppose there exists $\beta \in [\beta_{\text{st}}, n+1]$, $A \geq 1$ and $a \geq 0$ such that for all $Q \geq 1$ and for all $\delta \in (0, 1/2)^R$, it is true that*

$$\mathfrak{N}_{w, \mathcal{M}}(Q, \delta) \leq A \left(\delta^\times Q^{n+1} + \sum_{r=1}^R \delta_r^\times Q^{\frac{(n+1)(R-1)}{R}} Q^{\frac{\beta}{R}} + Q^\beta (\log 4Q)^{a(R-1)+R} \right). \quad (3.2)$$

Then there exists a positive constants C_0 depending only on w and \mathcal{M} , such that for all $Q \geq 1$ and for all $\delta \in (0, 1/2)^R$, we also have

$$\mathfrak{N}_{w, \mathcal{M}}(Q, \delta) \leq C_0 \delta^\times Q^{n+1} + C_0 A \left(\sum_{r=1}^R \delta_r^\times Q^{\frac{(n+1)(R-1)}{R}} Q^{\frac{\tilde{\beta}}{R}} + Q^{\tilde{\beta}} (\log 4Q)^{(a+1)(R-1)+R} \right); \quad (3.3)$$

with

$$\tilde{\beta} = n+1 - \frac{nR}{n+2 \left(R - \frac{n}{2\beta-n} \right)}. \quad (3.4)$$

Proof of Theorem 1.9, assuming Proposition 3.1. Using the trivial estimate

$$\mathfrak{N}_{w, \mathcal{M}}(Q, \delta) \leq \delta^\times Q^{n+1} + Q^{n+1} \leq \delta^\times Q^{n+1} + \sum_{r=1}^R \delta_r^\times Q^{\frac{(n+1)(R-1)}{R}} Q^{\frac{n+1}{R}} + Q^{n+1}$$

as the starting point, we apply Proposition 3.1 repeatedly. We first consider the case when $\beta_{\text{st}} = \frac{n(n+R+1)}{n+2R}$, which arises when $R \in \{1, 2\}$.

Applying Proposition 3.1 N many times, we get

$$\mathfrak{N}_{w, \mathcal{M}}(Q, \delta) \leq C_0 \delta^\times Q^{n+1} + C^N \left(\sum_{r=1}^R \delta_r^\times Q^{\frac{(n+1)(R-1)}{R}} Q^{\frac{\beta_N}{R}} + Q^{\beta_N} (\log 4Q)^{N(R-1)+R} \right), \quad (3.5)$$

where $C_0, C \geq 1$ are constants depending on w and \mathcal{M} ; and β_N is obtained from the recursive relation

$$\beta_i = n+1 - \frac{nR}{2R+n \left(1 - \frac{2}{2\beta_{i-1}-n} \right)}, \quad \beta_0 := n+1. \quad (3.6)$$

To determine how rapidly this sequence converges to $\frac{n(n+R+1)}{n+2R}$, we calculate

$$\begin{aligned} \beta_i - \frac{n(n+R+1)}{n+2R} &= n+1 - \frac{nR}{2R+n \left(1 - \frac{2}{2\beta_{i-1}-n} \right)} - \frac{n(n+R+1)}{n+2R} \\ &= \frac{(n+2)R}{n+2R} - \frac{nR}{2R+n \left(1 - \frac{2}{2\beta_{i-1}-n} \right)}. \end{aligned} \quad (3.7)$$

We can rewrite the last expression above as

$$\begin{aligned} & \frac{(n+2)R}{n+2R} - \frac{nR(2\beta_{i-1}-n)}{(n+2R)(2\beta_{i-1}-n)-2n} \\ &= \frac{R}{(n+2R)(2\beta_{i-1}-n)-2n} \left[\frac{(n+2)((n+2R)(2\beta_{i-1}-n)-2n) - n(n+2R)(2\beta_{i-1}-n)}{n+2R} \right]. \end{aligned} \quad (3.8)$$

The expression in square brackets equals

$$\begin{aligned} & \frac{-2n^2 + 2((n+2R)(2\beta_{i-1}-n)-2n)}{n+2R} = 4\beta_{i-1} - \frac{2n^2 + 2n(n+2R) + 4n}{n+2R} \\ &= 4\beta_{i-1} - \frac{4n(n+R+1)}{n+2R}. \end{aligned} \quad (3.9)$$

From (3.7), (3.8) and (3.9), we conclude that

$$\beta_i - \frac{n(n+R+1)}{n+2R} = \frac{4R}{(2\beta_{i-1}-n)(n+2R)-2n} \left(\beta_{i-1} - \frac{n(n+R+1)}{n+2R} \right). \quad (3.10)$$

Now since $\beta_{i-1} \geq \frac{n(n+R+1)}{n+2R}$, we have

$$(2\beta_{i-1}-n)(n+2R)-2n \geq n(n+2)-2n = n^2.$$

Plugging this into (3.10), we get

$$\beta_i - \frac{n(n+R+1)}{n+2R} \leq \frac{4R}{n^2} \left(\beta_{i-1} - \frac{n(n+R+1)}{n+2R} \right). \quad (3.11)$$

We start with the estimate

$$\beta_0 - \frac{n(n+R+1)}{n+2R} = n+1 - \frac{n(n+R+1)}{n+2R} = \frac{(n+2)R}{n+2R} \leq R.$$

After N steps, using (3.11), we get

$$\beta_N - \frac{n(n+R+1)}{n+2R} \leq \left(\frac{4R}{n^2} \right)^N R. \quad (3.12)$$

We now consider two subcases. When $n \geq 3$, it follows from Remark 1.14 and the definition of the Radon-Hurwitz numbers that $R < n$. Further, recall (1.29) which says that $R = 1$ whenever n is odd. Thus,

$$\frac{4R}{n^2} \leq \begin{cases} \frac{4}{5}, & \text{if } n \geq 5 \\ \frac{3}{4}, & \text{if } n = 4 \\ \frac{4}{9}, & \text{if } n = 3. \end{cases} \quad (3.13)$$

Combining the above with (3.12), we get

$$\beta_N - \frac{n(n+R+1)}{n+2R} \leq \left(\frac{4}{5} \right)^N R.$$

Thus, after

$$N = \left\lceil \frac{\log R + \log \log 4Q}{\log(5/4)} \right\rceil$$

many steps, we conclude from (3.5) that

$$\mathfrak{N}_{w,\mathcal{M}}(Q, \delta) \ll \delta^\times Q^{n+1} + \sum_{r=1}^R \delta_r^\times Q^{\frac{(n+1)(R-1)}{R}} Q^{\frac{n(n+R+1)}{R(n+2R)}} (\log 4Q)^{c_2} + Q^{\frac{n(n+R+1)}{n+2R}} \exp(c_2 (\log \log 4Q)^2),$$

for $R = 2$; and

$$\mathfrak{N}_{w,\mathcal{M}}(Q, \delta) \ll \delta^\times Q^{n+1} + \sum_{r=1}^R \delta_r^\times Q^{\frac{(n+1)(R-1)}{R}} Q^{\frac{n(n+R+1)}{R(n+2R)}} (\log 4Q)^{c_2} + Q^{\frac{n(n+R+1)}{n+2R}} (\log 4Q)^{c_2},$$

for $R = 1$. Here $c_2 > 0$ and the implied constant depend only on w and \mathcal{M} . This finishes the proof for $n \geq 3$ and $R \in \{1, 2\}$.

In the subcase when $n = 2$ and $R = 1$, the relation (3.10) reduces to

$$\beta_i - 2 = \frac{\beta_{i-1} - 2}{2\beta_{i-1} - 3}.$$

or equivalently, the identity

$$\frac{1}{\beta_i - 2} = 2 + \frac{1}{\beta_{i-1} - 2},$$

with $\beta_0 = 3$. This lets us conclude that

$$\beta_i = 2 + \frac{1}{2i + 1}.$$

This sequence also converges to the desired value of 2, albeit at a much slower rate. Indeed, after

$$N = \lfloor \sqrt{\log 4Q} \rfloor$$

many steps, we obtain using (3.5) that

$$\mathfrak{N}_{w,\mathcal{M}}(Q, \delta_1) \ll \delta_1 Q^3 + Q^2 \exp(c_1 \sqrt{\log 4Q}),$$

for a large enough constant $c_1 > 0$ depending only on w and \mathcal{M} . This establishes our result also in the case when $n = 2$.

We now come to the case when $\beta_{\text{st}} = \frac{n(n+1)}{2}$, which arises when $R \geq 3$. The argument proceeds in almost the same way as the previous case, except for one key difference. The sequence $\{\beta_i\}$ is still defined by the recursive relation (3.6) and therefore converges to $\frac{n(n+R+1)}{n+2R}$. However, Proposition 3.1 can be applied only until

$$\beta_i \geq \frac{n(n+1)}{2} > \frac{n(n+R+1)}{n+2R}.$$

Consequently, let β_{k-1}, β_k be such that

$$\beta_k < \frac{n(n+1)}{2} \leq \beta_{k-1}.$$

Applying Proposition 3.1 k many times, we obtain

$$\begin{aligned} \mathfrak{N}_{w,\mathcal{M}}(Q, \delta) &\leq C_0 \delta^\times Q^{n+1} + C^k \left(\sum_{r=1}^R \delta_r^\times Q^{\frac{(n+1)(R-1)}{R}} Q^{\frac{\beta_k}{R}} + Q^{\beta_k} (\log 4Q)^{(a+1)(R-1)+R} \right), \\ &\leq C_0 \delta^\times Q^{n+1} + C^k \left(\sum_{r=1}^R \delta_r^\times Q^{\frac{(n+1)(R-1)}{R}} Q^{\frac{n(n+1)}{2R}} + Q^{\frac{n(n+1)}{2}} (\log 4Q)^{(a+1)(R-1)+R} \right), \end{aligned}$$

where C_0, C_1 are positive constants depending only on w and \mathcal{M} . We apply Proposition 3.1 one more time, with $\beta = \frac{n(n+1)}{2}$, which gives

$$\mathfrak{N}_{w, \mathcal{M}}(Q, \delta) \leq C_0 \delta^\times Q^{n+1} + C^{k+1} \left(\sum_{r=1}^R \delta_r^\times Q^{\frac{(n+1)(R-1)}{R}} Q^{\frac{\Theta}{R}} + Q^\Theta (\log 4Q)^{(a+1)(R-1)+R} \right), \quad (3.14)$$

with

$$\Theta = n + 1 - \frac{nR}{n + 2 \left(R - \frac{n}{\frac{2n(n+1)}{2} - n} \right)} = n + 1 - \frac{nR}{n + 2(R-1) - \frac{4}{n}}.$$

Since $\Theta \geq \frac{n(n+R+1)}{n+2R}$, we conclude using (3.12) and (3.13) that

$$\beta_k - \Theta \leq \beta_k - \frac{n(n+R+1)}{n+2R} \leq \left(\frac{4}{5} \right)^k R.$$

Therefore we can again bound the number of steps required by

$$\left\lfloor \frac{\log R + \log \log 4Q}{\log(5/4)} \right\rfloor + 1.$$

Plugging the above into (3.14), we get

$$\mathfrak{N}_{w, \mathcal{M}}(Q, \delta) \ll \delta^\times Q^{n+1} + \sum_{r=1}^R \delta_r^\times Q^{\frac{(n+1)(R-1)}{R}} Q^{\frac{\Theta}{R}} (\log 4Q)^{c_2} + Q^\Theta \exp(c_2 (\log \log 4Q)^2),$$

for $R > 1$; and

$$\mathfrak{N}_{w, \mathcal{M}}(Q, \delta) \ll \delta^\times Q^{n+1} + \sum_{r=1}^R \delta_r^\times Q^{\frac{(n+1)(R-1)}{R}} Q^{\frac{n(n+R+1)}{R(n+2R)}} (\log 4Q)^{c_2} + Q^\Theta (\log 4Q)^{c_2},$$

for $R = 1$. Here $c_2 > 0$ and the implied constant depend only on w and \mathcal{M} .

This establishes (1.16) also in the case when $R \geq 3$ (note that the condition (1.28) forces n to be at least 3), and finishes the proof. \square

Proposition 3.1 will be a direct consequence of two inductive sub steps, built on the connection between the rational point count near the manifold \mathcal{M} and the sum of certain ‘‘dual’’ weights associated to a *family* of compact hypersurfaces in \mathbb{R}^{n+1} . We need some technical preparation first.

Let $\mathbb{1}_\delta$ denote the characteristic function of the set $\{\theta : \|\theta\| \leq \delta\}$, and for $1 \leq r \leq R$, set

$$J_r := \left\lfloor \frac{1}{2\delta_r} \right\rfloor. \quad (3.15)$$

Using the Selberg magic functions of degree J_r to estimate $\mathbb{1}_\delta$, as in (2.2), we can bound

$$\mathfrak{N}_{w, \mathcal{M}}(Q, \delta) = \sum_{\substack{\mathbf{a} \in \mathbb{Z}^n \\ 1 \leq q \leq Q}} w \left(\frac{\mathbf{a}}{q} \right) \prod_{r=1}^R \mathbb{1}_\delta \left(qf_r \left(\frac{\mathbf{a}}{q} \right) \right) \leq \sum_{\substack{\mathbf{a} \in \mathbb{Z}^n \\ 1 \leq q \leq Q}} w \left(\frac{\mathbf{a}}{q} \right) \prod_{r=1}^R S_{J_r^+} \left(qf_r \left(\frac{\mathbf{a}}{q} \right) \right).$$

Expanding the Selberg magic functions into their fourier series and multiplying, we get

$$\prod_{r=1}^R S_{J_r^+} \left(qf_r \left(\frac{\mathbf{a}}{q} \right) \right) = \sum_{\substack{\mathbf{j} \in \mathbb{Z}^R \\ |j_r| \leq J_r}} \left(\prod_{r=1}^R \widehat{S_{J_r^+}}(j_r) \right) \exp \left(\sum_{r=1}^R j_r qf_r \left(\frac{\mathbf{a}}{q} \right) \right).$$

The upshot is that

$$\mathfrak{N}_{w,\mathcal{M}}(Q, \delta) \leq \sum_{\substack{\mathbf{j} \in \mathbb{Z}^R: \\ |j_r| \leq J_r \\ 1 \leq r \leq R}} \left(\prod_{r=1}^R \widehat{S}_{J_r^+}(j_r) \right) \sum_{\substack{\mathbf{a} \in \mathbb{Z}^n \\ 1 \leq q \leq Q}} w \left(\frac{\mathbf{a}}{q} \right) \exp \left(\sum_{r=1}^R j_r q f_r \left(\frac{\mathbf{a}}{q} \right) \right). \quad (3.16)$$

Using (2.3) with $\beta - \alpha = 2\delta_r$ and $J = J_r$ for each $1 \leq r \leq R$, we conclude that the contribution from the regime when $\mathbf{j} = \mathbf{0}$ is

$$\prod_{r=1}^R \left(2\delta_r + \frac{1}{J_r + 1} \right) \sum_{\substack{\mathbf{a} \in \mathbb{Z}^n \\ 1 \leq q \leq Q}} w \left(\frac{\mathbf{a}}{q} \right) \ll_w \delta^\times Q^{n+1}. \quad (3.17)$$

Combining (3.16), (3.17) and the upper bound

$$|\widehat{S}_{J_r^+}(j_r)| \leq \frac{1}{J_r + 1} + \min \left(2\delta_r, \frac{1}{\pi |j_r|} \right) \leq \frac{1}{J_r + 1},$$

we get

$$\mathfrak{N}_{w,\mathcal{M}}(Q, \delta) \leq C_w \delta^\times Q^{n+1} + \sum_{\substack{\mathbf{j} \in \mathbb{Z}^R \setminus \{\mathbf{0}\}: \\ |j_r| \leq J_r \\ 1 \leq r \leq R}} \left(\prod_{r=1}^R \frac{1}{J_r + 1} \right) \left| \sum_{\substack{\mathbf{a} \in \mathbb{Z}^n \\ 1 \leq q \leq Q}} w \left(\frac{\mathbf{a}}{q} \right) \exp \left(\sum_{r=1}^R j_r q f_r \left(\frac{\mathbf{a}}{q} \right) \right) \right|. \quad (3.18)$$

For $1 \leq s \leq R$ and $X \in \mathbb{Z}_{>0}$, we define the ‘‘pencil set’’ of indices corresponding to the codimension indexed by s to be

$$\mathcal{J}^s(X) := \{\mathbf{j} \in \mathbb{Z}_{\geq 0}^R : 0 < \|\mathbf{j}\|_\infty = j_s \leq X\}. \quad (3.19)$$

For later use, we also introduce the slightly larger set

$$\mathcal{J}_b^s(X) := \{\mathbf{j} \in \mathbb{Z}^R : 0 < \|\mathbf{j}\|_\infty \leq 2j_s \leq 2X\}. \quad (3.20)$$

Further, for each $1 \leq s \leq R$, $\mathbf{j} \in \mathcal{J}^s(X)$ and $\gamma \in \{-1, 1\}^R$, let

$$F_{s,\mathbf{j},\gamma} := \gamma_s f_s + \sum_{\substack{1 \leq r \leq R \\ r \neq s}} \gamma_r \frac{j_r}{j_s} f_r \quad (3.21)$$

and

$$\begin{aligned} \mathfrak{N}_{w,\mathcal{M}}^{s,\gamma}(Q, \delta) &:= \delta^\times Q^{n+1} + \sum_{\mathbf{j} \in \mathcal{J}^s(J_s)} \left(\prod_{r=1}^R b_{j_r} \right) \left| \sum_{\substack{\mathbf{a} \in \mathbb{Z}^n \\ 1 \leq q \leq Q}} w \left(\frac{\mathbf{a}}{q} \right) \exp \left(\sum_{r=1}^R \gamma_r j_r q f_r \left(\frac{\mathbf{a}}{q} \right) \right) \right| \\ &= \delta^\times Q^{n+1} + \sum_{\mathbf{j} \in \mathcal{J}^s(J_s)} \left(\prod_{r=1}^R \frac{1}{j_r + 1} \right) \left| \sum_{\substack{\mathbf{a} \in \mathbb{Z}^n \\ 1 \leq q \leq Q}} w \left(\frac{\mathbf{a}}{q} \right) \exp \left(q j_s F_{s,\mathbf{j},\gamma} \left(\frac{\mathbf{a}}{q} \right) \right) \right|. \end{aligned} \quad (3.22)$$

Then it follows from the triangle inequality that

$$\mathfrak{N}_{w,\mathcal{M}}(Q, \delta) \leq \sum_{\gamma \in \{-1, 1\}^R} \sum_{s=1}^R \mathfrak{N}_{w,\mathcal{M}}^{s,\gamma}(Q, \delta). \quad (3.23)$$

Our argument shall be independent of the signs of the coefficients j_r . Thus by conjugation, if need be, we can always reduce matters to the case when $\gamma = (1, 1, \dots, 1)$. Henceforth,

we shall specialize to this choice of γ and suppress it from notation. In principle, the same argument can be made for the choice of a distinguished codimension s , but for clarity, we shall state the subsequent propositions for a general $s \in \{1, \dots, R\}$.

Let

$$\mathcal{D} := B_{2\varepsilon_0}(\mathbf{x}_0), \quad U := \text{supp } w \subseteq B_{\varepsilon_0}(\mathbf{x}_0), \quad (3.24)$$

with ε_0 chosen small enough so that there exists a constant $\mathfrak{C}_0 > 1$ such that for all $1 \leq s \leq R$ and $\mathbf{t} \in [-2, 2]^{R-1}$, we have

$$\frac{1}{\mathfrak{C}_0} \leq \left| \det H_{f_s + \sum_{\substack{r=1 \\ r \neq s}}^R t_r f_r} \right| \leq \mathfrak{C}_0; \quad (3.25)$$

and the maps

$$\mathbf{x} \rightarrow \nabla_{\mathbf{x}} \left(f_s + \sum_{\substack{1 \leq r \leq R \\ r \neq s}} t_r f_r \right)$$

are smooth diffeomorphisms on \mathcal{D} .

Then $\nabla F_{s,\mathbf{j}}$ is a diffeomorphism on \mathcal{D} and U for all $s \in \{1, \dots, R\}$ and $\mathbf{j} \in \mathcal{J}_b^s(J_s)$, and therefore also for all $\mathbf{j} \in \mathcal{J}^s(J_s)$. Let

$$\mathcal{R}_{s,\mathbf{j}} := \nabla F_{s,\mathbf{j}}(\mathcal{D}), \quad V_{s,\mathbf{j}} := \nabla F_{s,\mathbf{j}}(U). \quad (3.26)$$

Note that there exists a compact set \mathcal{L} , independent of s and J_1, \dots, J_R , such that $V_{s,\mathbf{j}} \subseteq \mathcal{R}_{s,\mathbf{j}} \subseteq \mathcal{L}$ for all $\mathbf{j} \in \mathcal{J}_b^s(J_s)$. For these \mathbf{j} tuples, we can now define the (Legendre) dual family of functions $F_{s,\mathbf{j}}^* : \mathcal{R}_{s,\mathbf{j}} \rightarrow \mathbb{R}$ by

$$F_{s,\mathbf{j}}^*(\mathbf{y}) := \mathbf{y} \cdot (\nabla F_{s,\mathbf{j}})^{-1}(\mathbf{y}) - (F_{s,\mathbf{j}} \circ (\nabla F_{s,\mathbf{j}})^{-1})(\mathbf{y}). \quad (3.27)$$

For each dual function $F_{s,\mathbf{j}}^*$, its gradient $\nabla F_{s,\mathbf{j}}^*$ is a smooth diffeomorphism mapping $\mathcal{R}_{s,\mathbf{j}}$ onto \mathcal{D} . For $\mathbf{y} = \nabla F_{s,\mathbf{j}}(\mathbf{x})$, we have

$$\nabla F_{s,\mathbf{j}}^*(\mathbf{y}) = \mathbf{x}, \quad H_{F_{s,\mathbf{j}}^*}(\mathbf{y}) = H_{F_{s,\mathbf{j}}}(\mathbf{x})^{-1}. \quad (3.28)$$

It can also be verified that the Legendre transform is an involution, i.e.

$$(F_{s,\mathbf{j}})^{**} = F_{s,\mathbf{j}}.$$

For $s \in \{1, \dots, R\}$ and $\mathbf{j} \in \mathcal{J}_b^s(J_s)$, we define the dual family of weights $w_{s,\mathbf{j}}^* : V_{s,\mathbf{j}} \rightarrow \mathbb{R}$ given by

$$w_{s,\mathbf{j}}^*(\mathbf{y}) := w \circ (\nabla F_{s,\mathbf{j}})^{-1}. \quad (3.29)$$

Finally, for $Q^* \geq 1$, $\delta^* \in (0, 1/2)$ and $s \in \{1, \dots, R\}$, define

$$\mathfrak{N}_{w,\mathcal{M}}^{*,s}(Q^*, \delta^*) := \sum_{\mathbf{j} \in \mathcal{J}^s(Q^*)} \sum_{\substack{\mathbf{a} \in \mathbb{Z}^n \\ \|j_s F_{s,\mathbf{j}}^*(\mathbf{a}/j_s)\| < \delta^*}} \frac{w_{s,\mathbf{j}}^*\left(\frac{\mathbf{a}}{j_s}\right)}{\sqrt{\left| \det H_{F_{s,\mathbf{j}}} \left((\nabla F_{s,\mathbf{j}})^{-1} \left(\frac{\mathbf{a}}{j_s} \right) \right) \right|}} \quad (3.30)$$

We are now ready to state the first sub step in the proof of Proposition 3.1, which converts an upper bound for $\mathfrak{N}_{w,\mathcal{M}}^s(Q, \delta)$ into an improved upper bound for the sum of the dual family of weights $\mathfrak{N}_{w,\mathcal{M}}^{*,s}(Q^*, \delta^*)$. In analogy with (3.1), we set

$$\alpha_{st} := \max \left(\frac{n(n+R+1)}{n+2}, n+R-1-\frac{2}{n} \right). \quad (3.31)$$

Proposition 3.2. *Suppose there exist $\beta \in [\beta_{st}, n+1]$, $A_1 \geq 1$ and $a_1 \geq 0$ such that for all $Q \geq 1$ and for all $\delta \in (0, 1/2)^R$, it is true that*

$$\mathfrak{N}_{w, \mathcal{M}}(Q, \delta) \leq A_1 \left(\delta^\times Q^{n+1} + \sum_{r=1}^R \delta_r^\times Q^{\frac{(n+1)(R-1)}{R}} Q^{\frac{\beta}{R}} + Q^\beta (\log 4Q)^{a_1(R-1)+R} \right). \quad (3.32)$$

Then there exists a positive constant C_1 depending only on w and \mathcal{M} such that for all $s \in \{1, \dots, R\}$, $Q^ \geq 1$ and $\delta^* \in (0, 1/2)$, we have*

$$\mathfrak{N}_{w, \mathcal{M}}^{*,s}(Q^*, \delta^*) \leq C_1 \delta^* (Q^*)^{n+R} + C_1 A_1 (Q^*)^\alpha \left((\log 4Q^*) \left(\frac{n+2R}{n+2} \right) \right)^{a_1(R-1)+R}, \quad (3.33)$$

with

$$\alpha = \max \left(n+R - \frac{n}{2\beta-n}, n+R-1 - \frac{2}{n} \right) \in [\alpha_{st}, n+R]. \quad (3.34)$$

The second sub step works in the reverse direction: it converts the upper bound for the sum of the above dual family of weights into an improved upper bound for $\mathfrak{N}_{w, \mathcal{M}}^s(Q, \delta)$.

Proposition 3.3. *Suppose there exists $\alpha \in \left[\frac{n(n+R+1)}{n+2}, n+R \right]$, $A_2 \geq 1$ and $a_2 \geq 0$ such that for all $s \in \{1, \dots, R\}$, $Q^* \geq 1$ and $\delta^* \in (0, 1/2)$, it is true that*

$$\mathfrak{N}_{w, \mathcal{M}}^{*,s}(Q^*, \delta^*) \leq A_2 \left(\delta^* (Q^*)^{n+R} + (Q^*)^\alpha \left((\log 4Q^*) \left(\frac{n+2R}{n+2} \right) \right)^{a_2(R-1)+R} \right). \quad (3.35)$$

Then there exists a positive constant C_2 depending only on w and \mathcal{M} such that for all $s \in \{1, \dots, R\}$, $Q \geq 1$ and $\delta \in (0, 1/2)^R$, we have

$$\mathfrak{N}_{w, \mathcal{M}}^s(Q, \delta) \leq C_2 \delta^\times Q^{n+1} + C_2 A_2 \left(\delta_s^\times Q^{\frac{(n+1)(R-1)}{R}} Q^{\frac{\tilde{\beta}}{R}} + Q^{\tilde{\beta}} (\log 4Q)^{(a_2+1)(R-1)+R} \right); \quad (3.36)$$

with

$$\tilde{\beta} = n+1 - \frac{nR}{2\alpha-n} \in \left[\frac{n(n+R+1)}{n+2R}, n+1 \right]. \quad (3.37)$$

Proof of Proposition 3.1, given Propositions 3.2 and 3.3. Using the hypothesis (3.2), we apply Proposition 3.2 to obtain (3.33) with $A_1 = A$, $a_1 = a$ and

$$\alpha = \max \left(n+R - \frac{n}{2\beta-n}, n+R-1 - \frac{2}{n} \right) = n+R - \frac{n}{2\beta-n}.$$

Indeed, since

$$\beta \geq \beta_{st} \geq \frac{n(n+1)}{n+2},$$

we have

$$n+R - \frac{n}{2\beta-n} \geq n+R - \frac{n}{\frac{2n(n+1)}{n+2} - n} = n+R-1 - \frac{2}{n}.$$

This in turn implies that the hypothesis of Proposition 3.3 is true with the same α , and with $a_2 = a$ and $A_2 = \max\{1, \mathfrak{C}_0 C_1 A\}$. Here \mathfrak{C}_0 is the constant from condition (3.25). Applying Proposition 3.3 next, we conclude that for all $s \in \{1, \dots, R\}$, $Q \geq 1$ and $\delta \in (0, 1/2)^R$,

$$\mathfrak{N}_{w, \mathcal{M}}^s(Q, \delta) \leq C'_0 \delta^\times Q^{n+1} + C'_0 A_2 \left(\sum_{r=1}^R \delta_r^\times Q^{\frac{(n+1)(R-1)}{R}} Q^{\frac{\tilde{\beta}}{R}} + Q^{\tilde{\beta}} (\log 4Q)^{(a+1)R} \right)$$

where C'_0 depends only on w and \mathcal{M} and

$$\tilde{\beta} = n + 1 - \frac{nR}{2\alpha - n}n + 1 - \frac{nR}{2\left(n + R - \frac{n}{2\beta - n}\right) - n} = n + 1 - \frac{nR}{2R + n\left(1 - \frac{2}{2\beta - n}\right)}.$$

Summing up in s and using (3.23), we obtain (3.3) with $C_0 = R2^R C'_0$. \square

Both Propositions 3.2 and 3.3 rely on a combination of projective duality and van der Corput's B-process to pass from \mathcal{M} to the family of dual hypersurfaces, and back. The following propositions make this connection precise.

We state the dual version first: for passage from the sum of the dual weights to rational point count in the neighborhood of \mathcal{M} . This should be compared to [21, Proposition 5.3 and §6.3], where the original counting problem is projected to a lower dimensional one associated to a family of hypersurfaces in \mathbb{R}^{n+1} , with a trivial summing up in the remaining $R - 1$ directions.

Proposition 3.4 (Dual van der Corput B-Process). *Let $s \in \{1, \dots, R\}$. For all $Q^* \geq 1$ and $\delta^* \in (0, 1/2)$ with $D := \lfloor \frac{1}{2\delta^*} \rfloor$, we have*

$$\begin{aligned} \mathfrak{N}_{w, \mathcal{M}}^{*,s}(Q^*, \delta^*) &\ll \delta^*(Q^*)^{n+R} + \frac{(Q^*)^{\frac{n}{2}}}{D} \sum_{d=1}^D d^{-\frac{n}{2}} \sum_{\mathbf{k} \in \mathbb{Z}^n} w\left(\frac{\mathbf{k}}{d}\right) \left(\prod_{r=1}^R \min(\|df_r(\mathbf{k}/d)\|^{-1}, Q^*) \right) \\ &\quad + (Q^*)^{\frac{n}{2}+R-1} D^{\frac{n}{2}-1}, \end{aligned} \quad (3.38)$$

with the implicit constant depending only on w and \mathcal{M} .

The final proposition gives an upper bound for $\mathfrak{N}_{w, \mathcal{M}}^s(Q, \boldsymbol{\delta})$ in terms of rational points around the dual family of hypersurfaces.

Proposition 3.5 (van der Corput B-Process for \mathcal{M}). *Let $s \in \{1, \dots, R\}$. For $Q \geq 1$ and $\boldsymbol{\delta} \in (0, 1/2)^R$, let $\mathfrak{N}_{w, \mathcal{M}}^s(Q, \boldsymbol{\delta})$ be as defined in (3.22) with $J_s := \lfloor \frac{1}{2\delta_s} \rfloor$. We have*

$$\begin{aligned} \mathfrak{N}_{w, \mathcal{M}}^s(Q, \boldsymbol{\delta}) &\ll \boldsymbol{\delta}^\times Q^{n+1} \\ &\quad + Q^{\frac{n}{2}} \sum_{\mathbf{j} \in \mathcal{J}^s(J_s)} \left(\prod_{r=1}^R \frac{1}{j_r + 1} \right) \sum_{\mathbf{k} \in \mathbb{Z}^n} \frac{w_{s, \mathbf{j}}^*\left(\frac{\mathbf{k}}{j_s}\right)}{\sqrt{|\det H_{F_{s, \mathbf{j}}}((\nabla F_{s, \mathbf{j}})^{-1}(\mathbf{k}/j_s))|}} j_s^{-\frac{n}{2}} \min(\|j_s F_{s, \mathbf{j}}^*(\mathbf{k}/j_s)\|^{-1}, Q) \\ &\quad + J_s^{\frac{n}{2}-1} Q^{\frac{n}{2}} (\log 4J_1)^R. \end{aligned} \quad (3.39)$$

The implicit constant depends only on w and \mathcal{M} .

We shall present the proofs of Propositions 3.2-3.5 for $s = 1$. The other cases can be reduced to this one by a relabelling of the variables j_1, \dots, j_r . Consequently, in the subsequent sections, we shall suppress notation and omit the parameter s . In other words, \mathcal{J}^s shall be denoted by \mathcal{J} , \mathcal{J}_b^s by \mathcal{J}_b , $F_{s, \mathbf{j}}$ by $F_{\mathbf{j}}$, $w_{s, \mathbf{j}}$ by $w_{\mathbf{j}}$, $\mathcal{R}_{s, \mathbf{j}}$ by $\mathcal{R}_{\mathbf{j}}$, $\mathcal{V}_{s, \mathbf{j}}$ by $\mathcal{V}_{\mathbf{j}}$, and so on.

4. PROOF OF PROPOSITION 3.2 USING PROPOSITION 3.4

Let $D := \lfloor \frac{1}{2\delta^*} \rfloor$. By Proposition 3.4, we have

$$\begin{aligned} \mathfrak{N}_{w, \mathcal{M}}^{*,1}(Q^*, \delta^*) &\ll \delta^*(Q^*)^{n+R} + \frac{(Q^*)^{\frac{n}{2}}}{D} \sum_{d=1}^D d^{-\frac{n}{2}} \sum_{\mathbf{k} \in \mathbb{Z}^n} w\left(\frac{\mathbf{k}}{d}\right) \left(\prod_{r=1}^R \min(\|df_r(\mathbf{k}/d)\|^{-1}, Q^*) \right) \\ &\quad + (Q^*)^{\frac{n}{2}+R-1} D^{\frac{n}{2}-1} \end{aligned} \quad (4.1)$$

with the implicit constant depending only on w and \mathcal{M} . To deal with the second term on the right, we employ dyadic decomposition based on the size of $\|df_r(\mathbf{k}/d)\|$ with respect to Q^* to obtain

$$\sum_{\mathbf{k} \in \mathbb{Z}^n} w\left(\frac{\mathbf{k}}{d}\right) \left(\prod_{r=1}^R \min(\|df_r(\mathbf{k}/d)\|^{-1}, Q^*) \right) \leq (Q^*)^R \sum_{\substack{\mathbf{i} \in \mathbb{Z}_{\geq 0}^R \\ \|\mathbf{i}\|_{\infty} \leq \frac{\log 4Q^*}{\log 2}}} 2^{-\sum_{r=1}^R i_r} \sum_{\substack{\mathbf{k} \in \mathbb{Z}^n \\ \|df_r(\frac{\mathbf{k}}{d})\| \leq \frac{2^{i_r+1}}{Q^*} \\ 1 \leq r \leq R}} w\left(\frac{\mathbf{k}}{d}\right). \quad (4.2)$$

Our induction hypothesis (3.32) implies that

$$\begin{aligned} \sum_{d=1}^D \sum_{\substack{\mathbf{k} \in \mathbb{Z}^n \\ \|df_r(\frac{\mathbf{k}}{d})\| \leq \frac{2^{i_r+1}}{Q^*}: 1 \leq r \leq R}} w\left(\frac{\mathbf{k}}{d}\right) &= \mathfrak{N}_{w, \mathcal{M}}\left(D, \frac{2^{i_1}}{Q^*}, \dots, \frac{2^{i_R}}{Q^*}\right) \\ &\leq A_1 \left(2^{\sum_{r=1}^R i_r} (Q^*)^{-R} D^{n+1} + \sum_{s=1}^R \frac{2^{\sum_{r=1}^R i_r}}{2^{i_s}} (Q^*)^{-R+1} D^{\frac{(n+1)(R-1)}{R}} D^{\frac{\beta}{R}} + D^{\beta} (\log 4D)^{a_1(R-1)+R} \right). \end{aligned}$$

Partial summation in the d variable then lets us conclude that

$$\begin{aligned} \sum_{d=1}^D d^{-\frac{n}{2}} \sum_{\substack{\mathbf{k} \in \mathbb{Z}^n \\ \|df_r(\frac{\mathbf{k}}{d})\| \leq \frac{2^{i_r+1}}{Q^*}: 1 \leq r \leq R}} w\left(\frac{\mathbf{k}}{d}\right) &\leq \\ A_1 D^{-\frac{n}{2}} \left(2^{\sum_{r=1}^R i_r} (Q^*)^{-R} D^{n+1} + \sum_{s=1}^R \frac{2^{\sum_{r=1}^R i_r}}{2^{i_s}} (Q^*)^{-R+1} D^{\frac{(n+1)(R-1)+\beta}{R}} + D^{\beta} (\log 4D)^{a_1(R-1)+R} \right). \end{aligned}$$

We combine the above with (4.2) to get

$$\begin{aligned} \sum_{d=1}^D d^{-\frac{n}{2}} \sum_{\mathbf{k} \in \mathbb{Z}^n} w\left(\frac{\mathbf{k}}{d}\right) \left(\prod_{r=1}^R \min(\|df_r(\mathbf{k}/d)\|^{-1}, Q^*) \right) &\leq A_1 D^{-\frac{n}{2}} (Q^*)^R \sum_{\substack{\mathbf{i} \in \mathbb{Z}_{\geq 0}^R \\ \|\mathbf{i}\|_{\infty} \leq \frac{\log 4Q^*}{\log 2}}} 2^{-\sum_{i=1}^R i_r} \left(2^{\sum_{r=1}^R i_r} (Q^*)^{-R} D^{n+1} + \sum_{s=1}^R \frac{2^{\sum_{r=1}^R i_r}}{2^{i_s}} (Q^*)^{-R+1} D^{\frac{(n+1)(R-1)+\beta}{R}} \right. \\ &\quad \left. + D^{\beta} (\log 4D)^{a_1(R-1)+R} \right) \\ &\leq A_1 \left(\left(\frac{\log 4Q^*}{\log 2} \right)^R D^{\frac{n}{2}+1} + RQ^* \left(\frac{\log 4Q^*}{\log 2} \right)^{R-1} D^{\frac{(n+1)(R-1)+\beta}{R} - \frac{n}{2}} + D^{\beta - \frac{n}{2}} (\log 4D)^{a_1(R-1)+R} (Q^*)^R \right). \quad (4.3) \end{aligned}$$

Plugging the above in (4.1) lets us conclude that $\mathfrak{N}_{w, \mathcal{M}}^*(Q^*, \delta^*)$ is bounded from above by a positive constant times

$$\begin{aligned} \delta^* (Q^*)^{n+R} + A_1 \frac{(Q^*)^{\frac{n}{2}}}{D} \left((\log 4Q^*)^R D^{\frac{n}{2}+1} + Q^* (\log 4Q^*)^{R-1} D^{\frac{(n+1)(R-1)+\beta}{R} - \frac{n}{2}} \right. \\ \left. + (Q^*)^R D^{\beta - \frac{n}{2}} (\log 4D)^{a_1(R-1)+R} \right) + (Q^*)^{\frac{n}{2}+R-1} D^{\frac{n}{2}-1}. \end{aligned}$$

Recalling that $D = \lfloor \frac{1}{2\delta^*} \rfloor$, we get

$$\begin{aligned} \mathfrak{N}_{w, \mathcal{M}}^{\ast, 1}(Q^*, \delta^*) &\leq C\delta^*(Q^*)^{n+R} + CA_1 \left((\log 4Q^*)^R (Q^*)^{\frac{n}{2}} (\delta^*)^{-\frac{n}{2}} + (Q^*)^{\frac{n}{2}+R} (\delta^*)^{\frac{n}{2}+1-\beta} (\log(4/\delta^*))^{a_1(R-1)+R} \right. \\ &\quad \left. + (\log 4Q^*)^{R-1} (Q^*)^{\frac{n}{2}+1} (\delta^*)^{\frac{n}{2}+1-\frac{(n+1)(R-1)+\beta}{R}} \right) + C(Q^*)^{\frac{n}{2}+R-1} (\delta^*)^{1-\frac{n}{2}}. \end{aligned} \quad (4.4)$$

Observe that all terms except the first on the right hand side of (4.4) involve a negative power of δ^* . To see this for the penultimate term, we estimate, first for $R \geq 2$,

$$\frac{n}{2} + 1 - \frac{(n+1)(R-1) + \beta}{R} = \frac{n+1-\beta}{R} - \frac{n}{2} \leq \frac{1-\beta}{2} \leq 0.$$

For $R = 1$, the power of δ^* on the penultimate term reduces to

$$\frac{n}{2} + 1 - \beta \leq 0,$$

as $\beta \geq n$ in this case.

We record a few small calculations which shall be of use later. Since $\beta \geq \frac{n(n+R+1)}{n+2R}$, we have

$$\frac{n}{2\beta - n} \leq \frac{n}{\frac{2n(n+R+1)}{n+2R} - n} \leq \frac{n+2R}{n+2}. \quad (4.5)$$

It is also straightforward to verify that with the aforementioned lower bound on β ,

$$\frac{n}{2} \left(1 + \frac{n}{2\beta - n} \right) \leq n + R - \frac{n}{2\beta - n}. \quad (4.6)$$

Further, since $\beta \geq \frac{n(n+1)}{n+2}$, we have

$$\left(\frac{n}{2\beta - n} \right) \left(\frac{n}{2} - 1 \right) \leq \frac{n \left(\frac{n}{2} - 1 \right)}{\frac{2n(n+1)}{n+2} - n} = \frac{n+2}{n} \left(\frac{n}{2} - 1 \right) = \frac{n}{2} - \frac{2}{n}. \quad (4.7)$$

We now return to (4.4), and consider two cases based on the size of δ^* with respect to Q^* . If $\delta^* \geq (Q^*)^{-\frac{n}{2\beta-n}}$, then we can estimate

$$(Q^*)^{\frac{n}{2}} (\delta^*)^{-\frac{n}{2}} \leq (Q^*)^{\frac{n}{2} \left(1 + \frac{n}{2\beta-n} \right)} \stackrel{(4.6)}{\leq} (Q^*)^{n+R-\frac{n}{2\beta-n}}. \quad (4.8)$$

Next, we deal with the last term on the right hand side in (4.4) as follows

$$\begin{aligned} (Q^*)^{\frac{n}{2}+R-1} (\delta^*)^{1-\frac{n}{2}} &\leq (Q^*)^{\frac{n}{2}+R-1+\left(\frac{n}{2\beta-n}\right)\left(\frac{n}{2}-1\right)} \stackrel{(4.7)}{\leq} (Q^*)^{\frac{n}{2}+R-1+\frac{n}{2}-\frac{2}{n}} \\ &= (Q^*)^{n+R-1-\frac{2}{n}}. \end{aligned} \quad (4.9)$$

We now estimate the two remaining middle terms which determine the relation of α with respect to β . The first one is easy

$$\begin{aligned} (Q^*)^{\frac{n}{2}+R} (\delta^*)^{\frac{n}{2}+1-\beta} (\log 4(1/\delta^*))^{a_1(R-1)+R} &\leq (Q^*)^{\frac{n}{2}+R+\left(\frac{n}{2\beta-n}\right)\left(\beta-\frac{n}{2}-1\right)} \left((\log 4Q^*) \left(\frac{n}{2\beta-n} \right) \right)^{a_1(R-1)+R} \\ &\stackrel{(4.5)}{\leq} (Q^*)^{n+R-\frac{n}{2\beta-n}} \left((\log 4Q^*) \left(\frac{n+2R}{n+2} \right) \right)^{a_1(R-1)+R}. \end{aligned} \quad (4.10)$$

We also have

$$\begin{aligned} (Q^*)^{\frac{n}{2}+1} (\delta^*)^{\frac{n}{2}+1-\frac{(n+1)(R-1)+\beta}{R}} &= \left((Q^*)^{\frac{n}{2}} (\delta^*)^{-\frac{n}{2}} \right)^{\frac{R-1}{R}} \left((Q^*)^{\frac{n}{2}+R} (\delta^*)^{\frac{n}{2}+1-\beta} \right)^{\frac{1}{R}} \\ &\stackrel{(4.8)+(4.10)}{\leq} \left((Q^*)^{\frac{(R-1)}{R}+\frac{1}{R}} \right)^{n+R-\frac{n}{2\beta-n}} = (Q^*)^{n+R-\frac{n}{2\beta-n}}. \end{aligned} \quad (4.11)$$

Combining (4.4), (4.8), (4.9), (4.10) and (4.11), yields

$$\mathfrak{N}_{w,\mathcal{M}}^{*,1}(Q^*, \delta^*) \leq C' \delta^* (Q^*)^{n+R} + C' A_1 (Q^*)^\alpha \left((\log 4Q^*) \left(\frac{n+2R}{n+2} \right) \right)^{a_1(R-1)+R}$$

for $\delta^* \geq (Q^*)^{-\frac{n}{2\beta-n}}$.

In the complementary case when $\delta^* \leq (Q^*)^{-\frac{n}{2\beta-n}}$, we use monotonicity of the function $\mathfrak{N}_{w,\mathcal{M}}^{*,1}(Q^*, \delta^*)$ to deduce that

$$\mathfrak{N}_{w,\mathcal{M}}^{*,1}(Q^*, \delta^*) \leq \mathfrak{N}_{w,\mathcal{M}}^{*,1}\left(Q^*, (Q^*)^{-\frac{n}{2\beta-n}}\right). \quad (4.12)$$

To estimate the right hand side, we use (4.4) with $\delta^* = (Q^*)^{-\frac{n}{2\beta-n}}$. Applying the same arguments as in (4.8)-(4.11), and evaluating the first term directly, gives

$$\begin{aligned} \mathfrak{N}_{w,\mathcal{M}}^{*,1}\left(Q^*, (Q^*)^{-\frac{n}{2\beta-n}}\right) &\leq C(Q^*)^{-\frac{n}{2\beta-n}+n+R} + C A_1 (Q^*)^\alpha \left((\log 4Q^*) \left(\frac{n+2R}{n+2} \right) \right)^{a_1(R-1)+R} \\ &\leq C'' A_1 (Q^*)^\alpha \left((\log 4Q^*) \left(\frac{n+2R}{n+2} \right) \right)^{a_1(R-1)+R}. \end{aligned} \quad (4.13)$$

This establishes (3.33) also in the case when $\delta^* \leq (Q^*)^{-\frac{n}{2\beta-n}}$, and thus finishes the proof of Proposition 3.2. The constant C_1 in (3.33) can be taken to be $\max(C', C'')$ and depends only on w and \mathcal{M} .

5. PROOF OF PROPOSITION 3.3 USING PROPOSITION 3.5

Recall that $J_1 := \left\lfloor \frac{1}{2\delta_1} \right\rfloor$. We begin by applying Proposition 3.5 to obtain

$$\begin{aligned} \mathfrak{N}_{w,\mathcal{M}}^1(Q, \delta) &\ll \delta^\times Q^{n+1} + J_1^{\frac{n}{2}-1} Q^{\frac{n}{2}} (\log 4J_1)^R \\ &+ Q^{\frac{n}{2}} \sum_{\mathbf{j} \in \mathcal{J}^1(J_1)} \left(\prod_{r=1}^R \frac{1}{j_r+1} \right) \sum_{\mathbf{k} \in \mathbb{Z}^n} \frac{w_{\mathbf{j}}^*\left(\frac{\mathbf{k}}{j_1}\right)}{\sqrt{|\det H_{F_{\mathbf{j}}}}((\nabla F_{\mathbf{j}})^{-1}(\mathbf{k}/j_1))}} j_1^{-\frac{n}{2}} \min(\|j_1 F_{\mathbf{j}}^*(\mathbf{k}/j_1)\|^{-1}, Q). \end{aligned} \quad (5.1)$$

To deal with the last term on the right, we use dyadic decomposition based on the size of $\|j_1 F_{\mathbf{j}}^*(\mathbf{k}/j_1)\|$ with respect to Q to split

$$\begin{aligned} &\sum_{\mathbf{k} \in \mathbb{Z}^n} \frac{w_{\mathbf{j}}^*\left(\frac{\mathbf{k}}{j_1}\right)}{\sqrt{|\det H_{F_{\mathbf{j}}}}((\nabla F_{\mathbf{j}})^{-1}(\mathbf{k}/j_1))}} \min(\|j_1 F_{\mathbf{j}}^*(\mathbf{k}/j_1)\|^{-1}, Q) \\ &\leq Q \sum_{\substack{\mathbf{k} \in \mathbb{Z}^n \\ \|j_1 F_{\mathbf{j}}^*(\mathbf{k}/j_1)\| < Q^{-1}}} \frac{w_{\mathbf{j}}^*\left(\frac{\mathbf{k}}{j_1}\right)}{\sqrt{|\det H_{F_{\mathbf{j}}}}((\nabla F_{\mathbf{j}})^{-1}(\mathbf{k}/j_1))}} \\ &+ \sum_{0 \leq i \leq \frac{\log 4Q}{\log 2}} 2^{-i} Q \sum_{\substack{\mathbf{k} \in \mathbb{Z}^n \\ \frac{2^i}{Q} < \|j_1 F_{\mathbf{j}}^*(\mathbf{k}/j_1)\| \leq \frac{2^{i+1}}{Q}}} \frac{w_{\mathbf{j}}^*\left(\frac{\mathbf{k}}{j_1}\right)}{\sqrt{|\det H_{F_{\mathbf{j}}}}((\nabla F_{\mathbf{j}})^{-1}(\mathbf{k}/j_1))}}. \end{aligned} \quad (5.2)$$

For $-1 \leq i \leq \frac{\log 4Q}{\log 2}$, our induction hypothesis (3.35) implies that

$$\begin{aligned} & \sum_{\mathbf{j} \in \mathcal{J}^1(J_1)} \sum_{\substack{\mathbf{k} \in \mathbb{Z}^n \\ \|j_1 F_{\mathbf{j}}^*(\mathbf{k}/j_1)\| \leq \frac{2^{i+1}}{Q}}} \frac{w_{\mathbf{j}}^*\left(\frac{\mathbf{k}}{j_1}\right)}{\sqrt{|\det H_{F_{\mathbf{j}}}((\nabla F_{\mathbf{j}})^{-1}(\mathbf{k}/j_1))|}} \\ &= \mathfrak{N}_{w, \mathcal{M}}^{*,1}\left(J_1, \frac{2^i}{Q}\right) \leq A_2 \left(2^{i+1}Q^{-1}J_1^{n+R} + J_1^\alpha \left((\log 4J_1) \left(\frac{n+2R}{n+2}\right)\right)^{a_2(R-1)+R}\right). \end{aligned} \quad (5.3)$$

Partial summation in the j_r variables, keeping in mind that $b_{j_r} \leq \frac{1}{j_r}$ and $j_r \leq j_1 \leq J_1$, then yields

$$\begin{aligned} & \sum_{\mathbf{j} \in \mathcal{J}^1(J_1)} \left(\prod_{r=1}^R \frac{1}{j_r + 1}\right) \sum_{\substack{\mathbf{k} \in \mathbb{Z}^n \\ \|j_1 F_{\mathbf{j}}^*(\mathbf{k}/j_1)\| \leq \frac{2^{i+1}}{Q}}} \frac{w_{\mathbf{j}}^*\left(\frac{\mathbf{k}}{j_1}\right) j_1^{-\frac{n}{2}}}{\sqrt{|\det H_{F_{\mathbf{j}}}((\nabla F_{\mathbf{j}})^{-1}(\mathbf{k}/j_1))|}} \\ & \leq \sum_{0 \leq s_2, \dots, s_R \leq \frac{\log J_1}{\log 2} + 1} \sum_{\substack{1 \leq j_1 \leq J_1 \\ j_r \in [2^{s_r-1}, 2^{s_r}] \\ (2 \leq r \leq R)}} \left(\prod_{r=2}^R 2^{-s_r}\right) j_1^{-\frac{n}{2}-1} \sum_{\substack{\mathbf{k} \in \mathbb{Z}^n \\ \|j_1 F_{\mathbf{j}}^*(\mathbf{k}/j_1)\| \leq \frac{2^{i+1}}{Q}}} \frac{w_{\mathbf{j}}^*\left(\frac{\mathbf{k}}{j_1}\right)}{\sqrt{|\det H_{F_{\mathbf{j}}}((\nabla F_{\mathbf{j}})^{-1}(\mathbf{k}/j_1))|}} \\ & \leq CA_2 (\log 4J_1)^R J_1^{-\frac{n}{2}} \left(2^{i+1}Q^{-1}J_1^n + J_1^{\alpha-R} \left((\log 4J_1) \left(\frac{n+2R}{n+2}\right)\right)^{a_2(R-1)+R}\right). \end{aligned}$$

Combining the decomposition in (5.2) with the above estimate, we get

$$\begin{aligned} & Q^{\frac{n}{2}} \sum_{\mathbf{j} \in \mathcal{J}^1(J_1)} \left(\prod_{r=1}^R \frac{1}{j_r + 1}\right) \sum_{\mathbf{k} \in \mathbb{Z}^n} \frac{w_{\mathbf{j}}^*\left(\frac{\mathbf{k}}{j_1}\right) j_1^{-\frac{n}{2}}}{\sqrt{|\det H_{F_{\mathbf{j}}}((\nabla F_{\mathbf{j}})^{-1}(\mathbf{k}/j_1))|}} \min(\|j_1 F_{\mathbf{j}}^*(\mathbf{k}/j_1)\|^{-1}, Q) \\ & \leq CA_2 Q^{\frac{n}{2}+1} (\log 4J_1)^R J_1^{-\frac{n}{2}} \sum_{0 \leq i \leq \frac{\log 4Q}{\log 2}} 2^{-i} \left(2^{i+1}Q^{-1}J_1^n + J_1^{\alpha-R} \left((\log 4J_1) \left(\frac{n+2R}{n+2}\right)\right)^{a_2(R-1)+R}\right) \\ & \leq CA_2 \left((\log 4J_1) \left(\frac{n+2R}{n+2}\right)\right)^b \left((\log 4Q) Q^{\frac{n}{2}} J_1^{\frac{n}{2}} + Q^{\frac{n}{2}+1} J_1^{\alpha-\frac{n}{2}-R}\right), \end{aligned}$$

where

$$b := (a_2 + 1)(R - 1) + R.$$

Plugging the above in (5.1) lets us conclude that

$$\mathfrak{N}_{w, \mathcal{M}}^1(Q, \delta) \leq C\delta^\times Q^{n+1} + CA_2 \left((\log 4J_1) \left(\frac{n+2R}{n+2}\right)\right)^b \left((\log 4Q) Q^{\frac{n}{2}} J_1^{\frac{n}{2}} + Q^{\frac{n}{2}+1} J_1^{\alpha-\frac{n}{2}-R}\right).$$

Recalling that $J_1 = \left\lfloor \frac{1}{2\delta_1} \right\rfloor$, we get

$$\mathfrak{N}_{w, \mathcal{M}}^1(Q, \delta) \leq C\delta^\times Q^{n+1} + CA_2 \left((\log 4\delta_1^{-1}) \left(\frac{n+2R}{n+2}\right)\right)^b \left((\log 4Q) Q^{\frac{n}{2}} \delta_1^{-\frac{n}{2}} + Q^{\frac{n}{2}+1} \delta_1^{\frac{n}{2}+R-\alpha}\right). \quad (5.4)$$

Observe that, since $\alpha \geq \frac{n(n+R+1)}{n+2}$, we have

$$\frac{n}{2\alpha - n} \left(\frac{n}{2} + R - \alpha \right) \leq \frac{n}{\frac{2n(n+R+1)}{n+2} - n} = \frac{n+2}{n+2R}. \quad (5.5)$$

It is also straightforward to verify that with the aforementioned lower bound on α , we have

$$\frac{n}{2} \left(1 + \frac{n}{2\alpha - n} \right) \leq n + 1 - \frac{nR}{2\alpha - n}. \quad (5.6)$$

We now return to (5.4) and consider two cases, depending on the size of δ_1 with respect to $Q^{-\frac{n}{2\alpha-n}}$. In the case when $\delta_1 \geq Q^{-\frac{n}{2\alpha-n}}$, we can bound

$$\begin{aligned} & \mathfrak{N}_{w, \mathcal{M}}^1(Q, \boldsymbol{\delta}) \\ & \leq C \boldsymbol{\delta}^\times Q^{n+1} + CA_2 \left((\log 4Q^{\frac{n}{2\alpha-n}}) \left(\frac{n+2R}{n+2} \right) \right)^b \left((\log 4Q) Q^{\frac{n}{2}(1+\frac{n}{2\alpha-n})} + Q^{\frac{n}{2}+1-\frac{n}{2\alpha-n}(\frac{n}{2}+R-\alpha)} \right) \\ & \stackrel{(5.5)+(5.6)}{\leq} C \boldsymbol{\delta}^\times Q^{n+1} + CA_2 (\log 4Q)^b \left(\frac{n+2R}{n+2} \right)^{-b} \left(\frac{n+2R}{n+2} \right)^b Q^{n+1-\frac{nR}{2\alpha-n}} \\ & \leq C' \boldsymbol{\delta}^\times Q^{n+1} + C' A_2 (\log 4Q)^{(a_2+1)(R-1)+R} Q^{\tilde{\beta}}. \end{aligned}$$

This establishes (3.36) when $\delta_1 \geq Q^{-\frac{n}{2\alpha-n}}$. In the complementary case, we use monotonicity of $\mathfrak{N}_{w, \mathcal{M}}^1(Q, \boldsymbol{\delta})$ as a function of δ_1 to deduce that

$$\begin{aligned} & \mathfrak{N}_{w, \mathcal{M}}^1(Q, \boldsymbol{\delta}) \\ & \leq \mathfrak{N}_{w, \mathcal{M}}^1 \left(Q, Q^{-\frac{n}{2\alpha-n}}, \delta_2, \dots, \delta_R \right) \\ & \stackrel{(5.4)}{\leq} C \boldsymbol{\delta}_1^\times Q^{n+1-\frac{n}{2\alpha-n}} + CA_2 (\log 4Q^{\frac{n}{2\alpha-n}})^b \left(\frac{n+2R}{n+2} \right)^b \left((\log 4Q) Q^{\frac{n}{2}(1+\frac{n}{2\alpha-n})} \right. \\ & \quad \left. + Q^{\frac{n}{2}+1-\frac{n}{2\alpha-n}(\frac{n}{2}+R-\alpha)} \right) \\ & \stackrel{(5.5)+(5.6)}{\leq} C'' \boldsymbol{\delta}_1^\times Q^{\frac{(R-1)(n+1)}{R}} Q^{\frac{1}{R}(n+1-\frac{nR}{2\alpha-n})} + C''' A_2 (\log 4Q)^b Q^{n+1-\frac{nR}{2\alpha-n}} \\ & = C'' \boldsymbol{\delta}_1^\times Q^{\frac{(R-1)(n+1)}{R}} Q^{\frac{\tilde{\beta}}{R}} + C''' A_2 (\log 4Q)^{(a_2+1)(R-1)+R} Q^{\tilde{\beta}}. \end{aligned}$$

This establishes (3.36) also in the case when $\delta_1 \leq Q^{-\frac{n}{2\alpha-n}}$, and finishes the proof.

6. PROOF OF PROPOSITION 3.4

For convenience, we recall the definition

$$\mathfrak{N}_{w, \mathcal{M}}^{*,1}(Q^*, \boldsymbol{\delta}^*) := \sum_{\mathbf{j} \in \mathcal{J}(Q^*)} \sum_{\substack{\mathbf{a} \in \mathbb{Z}^n \\ \|\mathbf{j}_1 F_{\mathbf{j}}^*(\mathbf{a}/\mathbf{j}_1)\| < \delta^*}} \frac{w_{\mathbf{j}}^* \left(\frac{\mathbf{a}}{\mathbf{j}_1} \right)}{\sqrt{|\det H_{F_{\mathbf{j}}}((\nabla F_{\mathbf{j}})^{-1}(\mathbf{a}/\mathbf{j}_1))|}}. \quad (6.1)$$

For technical reasons, it will be helpful to work with a dyadic version of the above counting function. For $\ell \in \mathbb{Z}_{>0}$, let

$$\mathcal{J}_\ell^1 := \mathcal{J}_\ell = \{ \mathbf{j} \in \mathbb{Z}_{\geq 0}^R : 2^{\ell-1} \leq j_1 < 2^\ell, \max_{r \neq 1} j_r \leq 2^\ell \}. \quad (6.2)$$

The counting function in (6.1) can be dominated by its dyadic version given by

$$N_{w,\mathcal{M}}^{*,1}(Q^*, \delta^*) := \sum_{\ell=1}^{\lceil \frac{\log 4Q^*}{\log 2} \rceil} \sum_{\mathbf{j} \in \mathcal{J}_\ell} \sum_{\substack{\mathbf{a} \in \mathbb{Z}^n \\ \|\mathbf{j}_1 F_{\mathbf{j}}^*(\mathbf{a}/j_1)\| < \delta^*}} \frac{w_{\mathbf{j}}^*\left(\frac{\mathbf{a}}{j_1}\right)}{\sqrt{|\det H_{F_{\mathbf{j}}}((\nabla F_{\mathbf{j}})^{-1}(\mathbf{a}/j_1))|}}. \quad (6.3)$$

For each $\ell \in \mathbb{Z}_{>0}$, \mathcal{J}_ℓ is a subset of \mathcal{J}_b (as defined in (3.20) with $s = 1$). Consequently, the condition (3.25) continues to hold and $\nabla F_{\mathbf{j}}$ is still a diffeomorphism on \mathcal{D} and U for all $\mathbf{j} \in \mathcal{J}_\ell$. The dyadic counting function above is therefore well defined.

We shall establish (3.38) with $\mathfrak{N}_{w,\mathcal{M}}^{*,1}(Q^*, \delta^*)$ replaced by $N_{w,\mathcal{M}}^{*,1}(Q^*, \delta^*)$.

Lemma 6.1. *Let $D := \lfloor \frac{1}{2\delta^*} \rfloor$. Then*

$$N_{w,\mathcal{M}}^{*,1}(Q^*, \delta^*) \ll (\delta^*)(Q^*)^{n+R} + \left| \sum_{\ell=1}^{\lceil \frac{\log 4Q^*}{\log 2} \rceil} \sum_{\mathbf{j} \in \mathcal{J}_\ell} \sum_{d=1}^D \frac{D-d}{D^2} \sum_{\mathbf{k} \in \mathbb{Z}^n} j_1^n I^*(d, \mathbf{j}, \mathbf{k}) \right|, \quad (6.4)$$

with

$$I^*(d, \mathbf{j}, \mathbf{k}) := \int_{\mathbb{R}^n} \frac{w_{\mathbf{j}}^*(\mathbf{z}) e(dj_1 (F_{\mathbf{j}}^*(\mathbf{z}) - j_1^{-1} \mathbf{k} \cdot \mathbf{z}))}{\sqrt{|\det H_{F_{\mathbf{j}}}((\nabla F_{\mathbf{j}})^{-1}(\mathbf{z}))|}} d\mathbf{z}. \quad (6.5)$$

Proof. Let $\mathcal{F}_D : \mathbb{R} \rightarrow [0, 1]$ be the Fejér kernel of degree D given by

$$\mathcal{F}_D(\theta) := \sum_{d=-D}^D \frac{D-|d|}{D^2} e(d\theta) = \left(\frac{\sin(\pi D\theta)}{D \sin(\pi\theta)} \right)^2. \quad (6.6)$$

Recall from (2.6) that this function has the property

$$\mathbf{1}_{\delta^*}(\theta) \leq \frac{\pi^2}{4} \mathcal{F}_D(\theta) \quad \text{for all } \theta \in \mathbb{R}. \quad (6.7)$$

Therefore $N_{w,\mathcal{M}}^{*,1}(Q^*, \delta^*)$ can be dominated by a positive constant times

$$\left| \sum_{\ell=1}^{\lceil \frac{\log 4Q^*}{\log 2} \rceil} \sum_{\mathbf{j} \in \mathcal{J}_\ell} \sum_{\mathbf{a} \in \mathbb{Z}^n} \frac{w_{\mathbf{j}}^*\left(\frac{\mathbf{a}}{j_1}\right)}{\sqrt{|\det H_{F_{\mathbf{j}}}((\nabla F_{\mathbf{j}})^{-1}(\mathbf{a}/j_1))|}} \sum_{d=-D}^D \frac{D-|d|}{D^2} e(dj_1 F_{\mathbf{j}}^*(\mathbf{a}/j_1)) \right|. \quad (6.8)$$

For all $\mathbf{j} \in \mathcal{J}_b$ (see (3.20)), the sets

$$\text{supp } w_{\mathbf{j}}^* := V_{\mathbf{j}}$$

are contained in a compact set $\mathcal{L} \subseteq \mathbb{R}^n$. Thus the contribution from the term corresponding to $d = 0$ in (6.8) is given by

$$\frac{1}{D} \sum_{\ell=1}^{\lceil \frac{\log 4Q^*}{\log 2} \rceil} \sum_{\mathbf{j} \in \mathcal{J}_\ell} \sum_{\mathbf{a} \in \mathbb{Z}^n} \frac{w_{\mathbf{j}}^*\left(\frac{\mathbf{a}}{j_1}\right)}{\sqrt{|\det H_{F_{\mathbf{j}}}((\nabla F_{\mathbf{j}})^{-1}(\mathbf{a}/j_1))|}} \ll \frac{1}{D} \left(\sum_{\ell=1}^{\lceil \frac{\log 4Q^*}{\log 2} \rceil} 2^{\ell(R-1)} \sum_{j_1=2^{\ell-1}}^{2^\ell} j_1^n \right) \ll (\delta^*)(Q^*)^{n+R}, \quad (6.9)$$

where the implicit constants depends only on w, \mathcal{M} and \mathcal{L} . Next, to handle the terms with $d \neq 0$, we apply the n -dimensional Poisson summation formula and a change of variables to

get

$$\begin{aligned} & \sum_{\mathbf{a} \in \mathbb{Z}^n} \frac{w_{\mathbf{j}}^* \left(\frac{\mathbf{a}}{j_1} \right) e \left(dj_1 F_{\mathbf{j}}^* \left(\frac{\mathbf{a}}{j_1} \right) \right)}{\sqrt{|\det H_{F_{\mathbf{j}}}((\nabla F_{\mathbf{j}})^{-1}(\mathbf{a}/j_1))|}} \\ &= \sum_{\mathbf{k} \in \mathbb{Z}^n} \int_{\mathbb{R}^n} \frac{w_{\mathbf{j}}^* \left(\frac{\mathbf{z}}{j_1} \right)}{\sqrt{|\det H_{F_{\mathbf{j}}}((\nabla F_{\mathbf{j}})^{-1}(\mathbf{a}/j_1))|}} e \left(dj_1 F_{\mathbf{j}}^* \left(\frac{\mathbf{z}}{j_1} \right) - \mathbf{k} \cdot \mathbf{z} \right) d\mathbf{z} = j_1^n \sum_{\mathbf{k} \in \mathbb{Z}^n} I^*(d, \mathbf{j}, \mathbf{k}), \end{aligned} \quad (6.10)$$

with $I^*(d, \mathbf{j}, \mathbf{k})$ as in (6.5). From (6.8), (6.9) and (6.10), we conclude

$$\begin{aligned} \mathfrak{N}_{w, \mathcal{M}}^{*,1}(Q^*, \delta^*) &\ll (\delta^*)(Q^*)^{n+R} + \left| \sum_{\ell=1}^{\lceil \frac{\log 4Q^*}{\log 2} \rceil} \sum_{\mathbf{j} \in \mathcal{J}_\ell} \sum_{|d|=1}^D \frac{D-d}{D^2} \sum_{\mathbf{k} \in \mathbb{Z}^n} j_1^n I^*(d, \mathbf{j}, \mathbf{k}) \right| \\ &\ll (\delta^*)(Q^*)^{n+R} + 2 \left| \sum_{\ell=1}^{\lceil \frac{\log 4Q^*}{\log 2} \rceil} \sum_{\mathbf{j} \in \mathcal{J}_\ell} \sum_{d=1}^D \frac{D-d}{D^2} \sum_{\mathbf{k} \in \mathbb{Z}^n} j_1^n I^*(d, \mathbf{j}, \mathbf{k}) \right|, \end{aligned}$$

where the last inequality follows from complex conjugation. \square

Let

$$\varphi^{(d, \mathbf{j}, \mathbf{k})}(\mathbf{x}) := F_{\mathbf{j}}^*(\mathbf{x}) - \frac{\mathbf{k}}{d} \cdot \mathbf{x}, \quad \varphi_1^{(d, \mathbf{j}, \mathbf{k})}(\mathbf{x}) := \frac{F_{\mathbf{j}}^*(\mathbf{x}) - \mathbf{k} \cdot \mathbf{x}}{\text{dist}(\mathbf{k}, dU)}. \quad (6.11)$$

Recall the index set \mathcal{J} (defined in (3.19)), and the fact that $\nabla F_{\mathbf{j}}$ is a diffeomorphism on the closure of \mathcal{D} for all $\mathbf{j} \in \mathcal{J}$. Further, recall the definitions of the sets $\mathcal{R}_{\mathbf{j}}$ and $V_{\mathbf{j}}$ from (3.26).

To analyze the oscillatory integrals $I^*(d, \mathbf{j}, \mathbf{k})$, we need a few preliminary estimates for the phase functions $\varphi^{(d, \mathbf{j}, \mathbf{k})}$ (and $\varphi_1^{(d, \mathbf{j}, \mathbf{k})}$) and the corresponding amplitudes. These are very similar to those proven in [21, §5 and §6]. The main work is needed to show that these are independent of the parameters d, \mathbf{j} and \mathbf{k} . We mention these estimates in the next lemma and briefly sketch their proofs. The interested reader is encouraged to consult [21] for the details.

Lemma 6.2. *Let $\mathcal{J}(J_1)$ be as defined in (3.19). Let $\alpha \in \mathbb{Z}_{\geq 0}^n$ be an n -dimensional multi-index, $d \leq D$ be a positive integer, $\mathbf{j} \in \mathcal{J}(J_1)$ and $\mathbf{k} \in \mathbb{Z}^n$. Then*

(i)

$$\sup_{\mathbf{x} \in V_{\mathbf{j}}} \left| \frac{\partial^\alpha \varphi_1^{d, \mathbf{j}, \mathbf{k}}}{\partial \mathbf{x}^\alpha}(\mathbf{x}) \right| \ll 1. \quad (6.12)$$

(ii)

$$\sup_{\mathbf{x} \in V_{\mathbf{j}}} \left| \frac{\partial^\alpha \varphi^{d, \mathbf{j}, \mathbf{k}}}{\partial \mathbf{x}^\alpha}(\mathbf{x}) \right| \ll 1. \quad (6.13)$$

(iii)

$$\sup_{\mathbf{x} \in V_{\mathbf{j}}} \left| \frac{\partial^\alpha}{\partial \mathbf{x}^\alpha} \left(w_{\mathbf{j}}^* (|\det H_{F_{\mathbf{j}}}|)^{-\frac{1}{2}} \right) (\mathbf{x}) \right| \ll 1. \quad (6.14)$$

(iv) *Under the additional assumption that $\mathbf{k} \in d\mathcal{D}$, we have*

$$\sup_{\mathbf{x} \in \mathcal{R}_{\mathbf{j}} \setminus \{x_{d, \mathbf{j}, \mathbf{k}}\}} \frac{|\mathbf{x} - \mathbf{x}_{d, \mathbf{j}, \mathbf{k}}|}{|\nabla \varphi^{d, \mathbf{j}, \mathbf{k}}(\mathbf{x})|} \ll 1, \quad (6.15)$$

where

$$\mathbf{x}_{d,\mathbf{j},\mathbf{k}} := (\nabla F_{\mathbf{j}}^*)^{-1}(\mathbf{k}/d) \in \mathcal{R}_{\mathbf{j}}. \quad (6.16)$$

The implicit constants in the estimates above depend on $\boldsymbol{\alpha}$ (for the first three inequalities), upper bounds for finitely many derivatives of f_r ($1 \leq r \leq R$) as well as w on U , and the constant \mathfrak{C}_0 in the condition (3.25), but are independent of d, \mathbf{j} and \mathbf{k} .

Proof. For the proof of (i), we refer the reader to Lemma 6.1 in [21]. The proof of (ii) also proceeds in the same way.

For part (iii), we also refer the reader to [21]. There is a slight difference though. In [21], the authors use Lemma 6.1 (or its proof) to deduce uniform upper bounds for the derivatives of the weight $w_{\mathbf{j}}^*$, whereas we need to establish (6.14) for $w_{\mathbf{j}}^* |\det H_{F_{\mathbf{j}}}|^{-\frac{1}{2}}$. However, this change is harmless owing to the condition (3.25) on the determinant of $H_{F_{\mathbf{j}}}$. This is easily seen for the zeroth derivative. Using the Leibniz rule, we know that the higher order derivatives of $w_{\mathbf{j}}^* |\det H_{F_{\mathbf{j}}}|^{-\frac{1}{2}}$ are sums of products of derivatives of $w_{\mathbf{j}}^*$ and $|\det H_{F_{\mathbf{j}}}|^{-\frac{1}{2}}$. We note that any partial derivative of $|\det H_{F_{\mathbf{j}}}|^{-\frac{1}{2}}$ is a real polynomial expression with uniformly bounded coefficients (independent of \mathbf{j} and \mathbf{k}), in terms of the powers of $|\det H_{F_{\mathbf{j}}}|^{-\frac{1}{2}}$ and the derivatives of $\nabla F_{\mathbf{j}}$ (which are again bounded by constants independent of \mathbf{j} and \mathbf{k}). Combining these observations with the proof of Lemma 6.1 in [21] establishes (6.14).

Finally, part (iv) is exactly the same as Lemma 6.2 in [21]. \square

As is standard for these types of problems, we divide our consideration into three regimes, based on whether we need to apply the method of stationary phase, non-stationary phase or a hybrid argument. Let

$$\rho^* := \frac{1}{2} \text{dist}(\partial \mathcal{D}, \partial U). \quad (6.17)$$

We split $\mathbb{Z}^n = \mathcal{K}_1 \cup \mathcal{K}_2 \cup \mathcal{K}_3$ where

$$\begin{aligned} \mathcal{K}_1 &:= \left\{ \mathbf{k} \in \mathbb{Z}^n : \frac{\mathbf{k}}{d} \in U \right\}, \\ \mathcal{K}_2 &:= \left\{ \mathbf{k} \in \mathbb{Z}^n : \text{dist} \left(\frac{\mathbf{k}}{d}, U \right) \geq \rho^* \right\}, \text{ and} \\ \mathcal{K}_3 &:= \mathbb{Z}^n \setminus (\mathcal{K}_1 \cup \mathcal{K}_2). \end{aligned}$$

For $i \in \{1, 2, 3\}$, we define the respective contributions

$$M_i^*(Q^*, \delta^*) := \left| \sum_{\ell=1}^{\lfloor \frac{\log 4Q^*}{\log 2} \rfloor} \sum_{\mathbf{j} \in \mathcal{J}_{\ell}} \sum_{d=1}^D \frac{D-d}{D^2} \sum_{\mathbf{k} \in \mathcal{K}_i} j_1^n I^*(d, \mathbf{j}, \mathbf{k}) \right|. \quad (6.18)$$

By (6.4), we have

$$N_{w, \mathcal{M}}^{*,1}(Q^*, \delta^*) \ll (\delta^*)(Q^*)^{n+R} + M_1^*(Q^*, \delta^*) + M_2^*(Q^*, \delta^*) + M_3^*(Q^*, \delta^*). \quad (6.19)$$

We first estimate the contribution from the non-stationary regime \mathcal{K}_2 .

Lemma 6.3.

$$M_2(Q^*, \delta^*) \ll (Q^*)^{R-1} \log(4Q^*),$$

with implicit constants depending only on \mathcal{M} and w .

Proof. Recall $\varphi^{d,\mathbf{j},\mathbf{k}}$ from (6.11), and let

$$\lambda_1 = j_1 \cdot \text{dist}(k, dU)$$

For each $\mathbf{k} \in \mathcal{K}_2$, we have the lower bound

$$\inf_{\mathbf{x} \in V_{\mathbf{j}}} |\nabla \varphi_1^{d, \mathbf{j}, \mathbf{k}}(\mathbf{x})| \geq 1.$$

Further, by parts (i) and (iii) of Lemma 6.2, we know that the derivatives of $\varphi_1^{d, \mathbf{j}, \mathbf{k}}$ and $w_{\mathbf{j}}^*(\mathbf{x}) |\det H_{F_{\mathbf{j}}}|^{-\frac{1}{2}}$ are bounded on $V_{\mathbf{j}}$ independently of d, \mathbf{j} and \mathbf{k} . Thus we can apply Lemma 2.1 (integration by parts), with phase $\varphi_1^{d, \mathbf{j}, \mathbf{k}}$ and $\lambda = \lambda_1$, to conclude that

$$I^*(d, \mathbf{j}, \mathbf{k}) \ll_m \lambda_1^{-m+1} = (j_1 \cdot \text{dist}(\mathbf{k}, dU))^{-m+1},$$

for $m \in \mathbb{Z}_{\geq 0}$, with implicit constants independent of d, \mathbf{j} and \mathbf{k} . In particular, taking $m \geq n+2$, we get

$$\begin{aligned} \sum_{\mathbf{k} \in \mathcal{K}_2} I^*(d, \mathbf{j}, \mathbf{k}) &\ll j_1^{-m+1} \sum_{\mathbf{k} \in \mathcal{K}_2} \text{dist}(\mathbf{k}, dU) \\ &\leq j_1^{-n-1} \sum_{i=0}^{\infty} \sum_{\substack{\mathbf{k} \in \mathbb{Z}^n \\ 2^i \rho^* \leq \text{dist}(\mathbf{k}, dU) < 2^{(i+1)} \rho^*}} 2^{-(i+1)(n+1)} (\rho^*)^{-(n+1)} \\ &\ll j_1^{-n-1} \sum_{i=0}^{\infty} 2^{(i+1)(-n-1+n)} \leq j_1^{-n-1}. \end{aligned}$$

Thus

$$\begin{aligned} M_2^*(Q^*, \delta^*) &\ll \sum_{\ell=1}^{\lceil \frac{\log 4Q^*}{\log 2} \rceil} \sum_{\mathbf{j} \in \mathcal{J}_{\ell}} \sum_{d=1}^D \frac{D-d}{D^2} j_1^{n-n-1} \ll \left(\sum_{\ell=1}^{\lceil \frac{\log 4Q^*}{\log 2} \rceil} 2^{\ell(R-1)} \sum_{j_1=2^{\ell-1}}^{2^{\ell}} j_1^{-1} \right) \\ &\ll (Q^*)^{R-1}. \end{aligned}$$

□

Recall $\varphi^{d, \mathbf{j}, \mathbf{k}}$ from (6.11). Since $\nabla F_{\mathbf{j}}^*$ is a diffeomorphism on $\mathcal{R}_{\mathbf{j}} = (\nabla F_{\mathbf{j}})^{-1}(\mathcal{D})$, each $\mathbf{k} \in d\mathcal{D}$ determines a unique critical point of $\varphi^{d, \mathbf{j}, \mathbf{k}}$ given by

$$x_{d, \mathbf{j}, \mathbf{k}} := (\nabla F_{\mathbf{j}}^*)^{-1}(\mathbf{k}/d) = \nabla F_{\mathbf{j}}(\mathbf{k}/d) \in \mathcal{R}_{\mathbf{j}}.$$

For the intermediate regime, we have the following estimate.

Lemma 6.4.

$$M_3(Q^*, \delta^*) \ll (Q^*)^{\frac{n}{2}+R-1} D^{\frac{n}{2}-1},$$

with implicit constants depending only on \mathcal{M} and w .

Proof. For $\mathbf{k} \in \mathcal{K}_3 \subseteq \mathcal{D} \setminus dU$, we know that $x_{d, \mathbf{j}, \mathbf{k}} \notin \nabla F_{\mathbf{j}}(U) = \text{supp } w_{\mathbf{j}}^*$. Recall that the phase $\varphi^{d, \mathbf{j}, \mathbf{k}}$ and the amplitude $\left(w_{\mathbf{j}}^*(\mathbf{x}) |\det H_{F_{\mathbf{j}}}|^{-\frac{1}{2}}\right)$ satisfy the estimates (6.13) and (6.14) respectively. Further,

$$H_{\varphi^{d, \mathbf{j}, \mathbf{k}}}(x_{d, \mathbf{j}, \mathbf{k}}) = H_{F_{\mathbf{j}}^*}((\nabla F_{\mathbf{j}}^*)^{-1}(\mathbf{k}/d)) = H_{F_{\mathbf{j}}}(\mathbf{k}/d) \gg \mathfrak{C}_0^{-1},$$

by (3.28) and (3.25). Thus, we can apply Lemma 2.2 (stationary phase principle), with phase $\varphi^{d, \mathbf{j}, \mathbf{k}}$ and $\lambda := dj_1$, to conclude that

$$I^*(d, \mathbf{j}, \mathbf{k}) \ll (dj_1)^{-\frac{n}{2}-1}$$

for each $\mathbf{k} \in \mathcal{K}_3$. Since $\#\mathcal{K}_3 \ll_{U,\rho^*} d^n$, we can estimate

$$\begin{aligned} M_3(Q^*, \delta^*) &\ll \sum_{\ell=1}^{\lceil \frac{\log 4Q^*}{\log 2} \rceil} \sum_{\mathbf{j} \in \mathcal{J}_\ell} \sum_{d=1}^D \frac{D-d}{D^2} \sum_{\mathbf{k} \in \mathcal{K}_3} j_1^n |I^*(d, \mathbf{j}, \mathbf{k})| \\ &\ll D^{-1} \sum_{\ell=1}^{\lceil \frac{\log 4Q^*}{\log 2} \rceil} 2^{\ell(R-1)} \sum_{j_1=2^{\ell-1}}^{2^\ell} \sum_{d=1}^D (j_1 d)^{n-\frac{n}{2}-1} \\ &\ll (Q^*)^{\frac{n}{2}+R-1} D^{\frac{n}{2}-1}. \end{aligned}$$

□

Finally, we estimate the contribution from the critical stationary phase regime.

Lemma 6.5.

$$M_1(Q^*, \delta^*) \ll \frac{(Q^*)^{\frac{n}{2}}}{D} \sum_{d=1}^D d^{-\frac{n}{2}} \sum_{\mathbf{k} \in \mathbb{Z}^n} w\left(\frac{\mathbf{k}}{d}\right) \left(\prod_{r=1}^R \min(\|df_r(\mathbf{k}/d)\|^{-1}, Q^*) \right) + (Q^*)^{\frac{n}{2}+R-1} D^{\frac{n}{2}-1},$$

with implicit constants depending only on \mathcal{M} and w .

Proof. We again intend to apply the stationary phase principle to evaluate the integrals $I^*(d, \mathbf{k}, \mathbf{j})$ for $\mathbf{k} \in dU$. As in the proof of Lemma 6.4, we note that the phase $\varphi^{d,\mathbf{j},\mathbf{k}}$ and the amplitude $\left(w_{\mathbf{j}}^*(\mathbf{x}) |\det H_{F_{\mathbf{j}}}|^{-\frac{1}{2}}\right)$ satisfy the estimates (6.13) and (6.14) respectively. Further, by (3.28) and (3.25), we know that

$$H_{\varphi^{d,\mathbf{j},\mathbf{k}}}(x_{d,\mathbf{j},\mathbf{k}}) = H_{F_{\mathbf{j}}}^*((\nabla F_{\mathbf{j}}^*)^{-1}(\mathbf{k}/d)) = H_{F_{\mathbf{j}}}(\mathbf{k}/d) \gg \mathfrak{C}_0^{-1}. \quad (6.20)$$

An application of Lemma 2.2 (stationary phase principle), with $\lambda = dj_1$, phase $\varphi^{d,\mathbf{j},\mathbf{k}}$ and amplitude function $\frac{w_{\mathbf{j}}^*}{\sqrt{|\det H_{F_{\mathbf{j}}}^*|}}$, gives

$$\begin{aligned} I^*(d, \mathbf{j}, \mathbf{k}) &= \frac{w_{\mathbf{j}}^*(\mathbf{x}_{d,\mathbf{j},\mathbf{k}})}{\sqrt{|\det H_{F_{\mathbf{j}}}^*(\mathbf{x}_{d,\mathbf{j},\mathbf{k}})|}} \frac{1}{\sqrt{|\det H_{F_{\mathbf{j}}}(\mathbf{k}/d)|}} (j_1 d)^{-\frac{n}{2}} e(-j_1 d \varphi^{d,\mathbf{j},\mathbf{k}}(\mathbf{x}_{d,\mathbf{j},\mathbf{k}}) + \sigma_{\mathbf{j}}/8) \\ &\quad + O((j_1 d)^{-\frac{n}{2}-1}). \end{aligned}$$

For $\mathbf{k} \in \mathcal{K}_1$, we have

$$\frac{w_{\mathbf{j}}^*(\mathbf{x}_{d,\mathbf{j},\mathbf{k}})}{\sqrt{|\det H_{F_{\mathbf{j}}}^*(\mathbf{x}_{d,\mathbf{j},\mathbf{k}})|}} = w\left(\frac{\mathbf{k}}{d}\right) \sqrt{|\det H_{F_{\mathbf{j}}}(\mathbf{k}/d)|}. \quad (6.21)$$

We can also simplify

$$\varphi^{d,\mathbf{j},\mathbf{k}}(\mathbf{x}_{d,\mathbf{j},\mathbf{k}}) = \left(F_{\mathbf{j}}^* \circ (\nabla F_{\mathbf{j}}^*)^{-1}\right)\left(\frac{\mathbf{k}}{d}\right) - \frac{\mathbf{k}}{d} \cdot (\nabla F_{\mathbf{j}}^*)^{-1}\left(\frac{\mathbf{k}}{d}\right) = -F_{\mathbf{j}}\left(\frac{\mathbf{k}}{d}\right). \quad (6.22)$$

Plugging (6.21) and (6.22) into the stationary phase expansion for $I^*(d, \mathbf{j}, \mathbf{k})$, we get

$$\begin{aligned} I^*(d, \mathbf{j}, \mathbf{k}) &= w\left(\frac{\mathbf{k}}{d}\right) (j_1 d)^{-\frac{n}{2}} e\left(-j_1 d F_{\mathbf{j}}\left(\frac{\mathbf{k}}{d}\right) + \frac{\sigma_{\mathbf{j}}}{8}\right) + O((j_1 d)^{-\frac{n}{2}-1}) \\ &= w\left(\frac{\mathbf{k}}{d}\right) (j_1 d)^{-\frac{n}{2}} e\left(-d \sum_{r=1}^R j_r f_r\left(\frac{\mathbf{k}}{d}\right) + \frac{\sigma_{\mathbf{j}}}{8}\right) + O((j_1 d)^{-\frac{n}{2}-1}). \end{aligned} \quad (6.23)$$

Now

$$\begin{aligned} & \left| \sum_{\ell=1}^{\lceil \frac{\log 4Q^*}{\log 2} \rceil} \sum_{\mathbf{j} \in \mathcal{J}_\ell} j_1^{n-\frac{n}{2}} e \left(-d \sum_{r=1}^R j_r f_r \left(\frac{\mathbf{k}}{d} \right) \right) \right| \\ & \leq \sum_{\ell=1}^{\lceil \frac{\log 4Q^*}{\log 2} \rceil} \left(\left| \sum_{j_1 \in \mathbb{Z} \cap [2^{\ell-1}, 2^\ell]} j_1^{\frac{n}{2}} e \left(-dj_1 f_1 \left(\frac{\mathbf{k}}{d} \right) \right) \right| \right) \left(\prod_{r=2}^R \left| \sum_{j_r \in \mathbb{Z} \cap [0, 2^\ell]} e \left(-dj_r f_r \left(\frac{\mathbf{k}}{d} \right) \right) \right| \right). \end{aligned}$$

By geometric summation, for $2 \leq r \leq R$, we have

$$\left| \sum_{j_r \in \mathbb{Z} \cap [0, 2^\ell]} e \left(-dj_r f_r \left(\frac{\mathbf{k}}{d} \right) \right) \right| \leq \min(2^\ell, \|df_r(\mathbf{k}/d)\|^{-1}),$$

while for $r = 1$, we use geometric and partial summation to conclude that

$$\left| \sum_{j_1 \in \mathbb{Z} \cap [2^{\ell-1}, 2^\ell]} j_1^{\frac{n}{2}} e \left(-dj_1 f_1 \left(\frac{\mathbf{k}}{d} \right) \right) \right| \leq 2^{\ell \frac{n}{2}} \min(2^\ell, \|df_1(\mathbf{k}/d)\|^{-1}).$$

Putting the above estimates together, we conclude that

$$\begin{aligned} \left| \sum_{\ell=1}^{\lceil \frac{\log 4Q^*}{\log 2} \rceil} \sum_{\mathbf{j} \in \mathcal{J}_\ell} j_1^{\frac{n}{2}} e \left(-d \sum_{r=1}^R j_r f_r \left(\frac{\mathbf{k}}{d} \right) \right) \right| & \leq \sum_{\ell=1}^{\lceil \frac{\log 4Q^*}{\log 2} \rceil} 2^{\ell \frac{n}{2}} \prod_{r=1}^R \min(2^\ell, \|df_r(\mathbf{k}/d)\|^{-1}) \\ & \leq \left(\sum_{\ell=1}^{\lceil \frac{\log 4Q^*}{\log 2} \rceil} 2^{\ell \frac{n}{2}} \right) \prod_{r=1}^R \min(2Q^*, \|df_r(\mathbf{k}/d)\|^{-1}) \\ & \leq 4(Q^*)^{\frac{n}{2}} \prod_{r=1}^R \min(Q^*, \|df_r(\mathbf{k}/d)\|^{-1}). \end{aligned} \quad (6.24)$$

We now have all ingredients in place to estimate $M_3^*(Q^*, \delta^*)$. We first use (6.23) to conclude that $M_3^*(Q^*, \delta^*)$ is dominated by

$$\sum_{d=1}^D \frac{D-d}{D^2} d^{-\frac{n}{2}} \sum_{\mathbf{k} \in \mathcal{K}_3} w \left(\frac{\mathbf{k}}{d} \right) \left| \sum_{\ell=1}^{\lceil \frac{\log 4Q^*}{\log 2} \rceil} \sum_{\mathbf{j} \in \mathcal{J}_\ell} j_1^{n-\frac{n}{2}} e \left(-d \sum_{r=1}^R j_r f_r \left(\frac{\mathbf{k}}{d} \right) \right) \right| \quad (6.25)$$

$$+ \sum_{\ell=1}^{\lceil \frac{\log 4Q^*}{\log 2} \rceil} \sum_{\mathbf{j} \in \mathcal{J}_\ell} \sum_{d=1}^D \frac{D-d}{D^2} \sum_{\mathbf{k} \in \mathcal{K}_3} j_1^{n-\frac{n}{2}-1} d^{-\frac{n}{2}-1}. \quad (6.26)$$

Using (6.24), we deduce that (6.25) can be bounded from above by a positive constant times

$$\frac{(Q^*)^{\frac{n}{2}}}{D} \sum_{d=1}^D d^{-\frac{n}{2}} \sum_{\mathbf{k} \in \mathbb{Z}^n} w \left(\frac{\mathbf{k}}{d} \right) \left(\prod_{r=1}^R \min(\|df_r(\mathbf{k}/d)\|^{-1}, Q^*) \right).$$

On the other hand, using the fact that $(\#\mathcal{K}_3) \ll d^{-n}$, (6.26) can be estimated as follows

$$\begin{aligned} \sum_{\ell=1}^{\lceil \frac{\log 4Q^*}{\log 2} \rceil} \sum_{\mathbf{j} \in \mathcal{J}_\ell} \sum_{d=1}^D \frac{D-d}{D^2} \sum_{\mathbf{k} \in \mathcal{K}_3} j_1^{n-\frac{n}{2}-1} d^{-\frac{n}{2}-1} &\leq \frac{1}{D} \left(\sum_{\ell=1}^{\lceil \frac{\log 4Q^*}{\log 2} \rceil} 2^{\ell(R-1)} \sum_{j_1=2^{\ell-1}}^{2^\ell} j_1^{\left(\frac{n}{2}-1\right)} \right) \left(\sum_{d=1}^D (\#\mathcal{K}_3) d^{-\frac{n}{2}-1} \right) \\ &\ll \left(\sum_{\ell=1}^{\lceil \frac{\log 4Q^*}{\log 2} \rceil} 2^{\ell\left(\frac{n}{2}+R-1\right)} \right) \left(D^{-1} \sum_{d=1}^D d^{n-\frac{n}{2}-1} \right) \leq (Q^*)^{\frac{n}{2}+R-1} D^{\frac{n}{2}-1}. \end{aligned}$$

Adding the two estimates finishes the proof. \square

Concluding the proof of Proposition 3.4. Combining (6.19) with Lemmas 6.3-6.5, we conclude that

$$\begin{aligned} \mathfrak{N}_{w,\mathcal{M}}^{*,1}(Q^*, \delta^*) &\leq N_{w,\mathcal{M}}^{*,1}(Q^*, \delta^*) \ll (\delta^*)(Q^*)^{n+R} + (Q^*)^{R-1} + (Q^*)^{\frac{n}{2}+R-1} D^{\frac{n}{2}-1} \\ &\quad + \frac{(Q^*)^{\frac{n}{2}}}{D} \sum_{d=1}^D d^{-\frac{n}{2}} \sum_{\mathbf{k} \in \mathbb{Z}^n} w\left(\frac{\mathbf{k}}{d}\right) \left(\prod_{r=1}^R \min(\|df_r(\mathbf{k}/d)\|^{-1}, Q^*) \right), \end{aligned}$$

which implies (3.38).

7. PROOF OF PROPOSITION 3.5

Recall that

$$\mathfrak{N}_{w,\mathcal{M}}^1(Q, \delta) = C_w \delta^\times Q^{n+1} + \sum_{\mathbf{j} \in \mathcal{J}(J_1)} \left(\prod_{r=1}^R \frac{1}{1+j_r} \right) \left| \sum_{\substack{\mathbf{a} \in \mathbb{Z}^n \\ 1 \leq q \leq Q}} w\left(\frac{\mathbf{a}}{q}\right) \exp\left(qj_1 F_{\mathbf{j}}\left(\frac{\mathbf{a}}{q}\right)\right) \right|,$$

where \mathcal{J} is as defined in (3.19) with $s = 1$. Using the n -dimensional Poisson summation formula (and a change of variables) for the sum inside absolute values, we can write

$$\sum_{\substack{\mathbf{a} \in \mathbb{Z}^n \\ 1 \leq q \leq Q}} w\left(\frac{\mathbf{a}}{q}\right) \exp\left(qj_1 F_{\mathbf{j}}\left(\frac{\mathbf{a}}{q}\right)\right) = \sum_{\substack{\mathbf{k} \in \mathbb{Z}^n \\ 1 \leq q \leq Q}} q^n I(q, \mathbf{j}, \mathbf{k}),$$

with

$$I(q, \mathbf{j}, \mathbf{k}) = \int_{\mathbb{R}^n} \exp\left(qj_1 \left(F_{\mathbf{j}}(\mathbf{x}) - \frac{\mathbf{k} \cdot \mathbf{x}}{j_1}\right)\right) w(\mathbf{x}) d\mathbf{x}. \quad (7.1)$$

We thus have

$$\mathfrak{N}_{w,\mathcal{M}}^1(Q, \delta) = C_w \delta^\times Q^{n+1} + \sum_{\mathbf{j} \in \mathcal{J}(J_1)} \left(\prod_{r=1}^R \frac{1}{1+j_r} \right) \left| \sum_{\substack{\mathbf{k} \in \mathbb{Z}^n \\ 1 \leq q \leq Q}} q^n I(q, \mathbf{j}, \mathbf{k}) \right|. \quad (7.2)$$

Recall the set \mathcal{D} from (3.24), and that $\nabla F_{\mathbf{j}}$ is a diffeomorphism on \mathcal{D} for all $\mathbf{j} \in \mathcal{J}$. Further, recall the definitions of the sets $\mathcal{R}_{\mathbf{j}}$ and $V_{\mathbf{j}}$ from (3.26) (for $s = 1$).

Let

$$\varphi^{(\mathbf{j}, \mathbf{k})}(\mathbf{x}) := F_{\mathbf{j}}(\mathbf{x}) - \frac{\mathbf{k}}{j_1} \cdot \mathbf{x}, \quad \varphi_1^{(\mathbf{j}, \mathbf{k})}(\mathbf{x}) := \frac{F_{\mathbf{j}}(\mathbf{x}) - \mathbf{k} \cdot \mathbf{x}}{\text{dist}(\mathbf{k}, j_1 V_{\mathbf{j}})}. \quad (7.3)$$

As in the last section, to analyze the oscillatory integrals $I(d, \mathbf{j}, \mathbf{k})$, we need a few preliminary estimates for the phase functions $\varphi^{(\mathbf{j}, \mathbf{k})}$ (and $\varphi_1^{(\mathbf{j}, \mathbf{k})}$) and the corresponding amplitude functions. These estimates are again very similar to those proven in [21, §5], with the main

work spent in showing that they are independent of the parameters \mathbf{j} and \mathbf{k} . The following lemma is analogous to Lemma 6.2 from the previous section.

Lemma 7.1. *Let $\mathcal{J}(J_1)$ be as defined in (3.19). Let $\boldsymbol{\alpha} \in \mathbb{Z}_{\geq 0}^n$ be an n -dimensional multi-index, $\mathbf{j} \in \mathcal{J}(J_1)$ and $\mathbf{k} \in \mathbb{Z}^n$. Then*

(i)

$$\sup_{\mathbf{x} \in U} \left| \frac{\partial^{\boldsymbol{\alpha}} \varphi_1^{\mathbf{j}, \mathbf{k}}}{\partial \mathbf{x}^{\boldsymbol{\alpha}}}(\mathbf{x}) \right| \ll 1. \quad (7.4)$$

(ii)

$$\sup_{\mathbf{x} \in U} \left| \frac{\partial^{\boldsymbol{\alpha}} \varphi^{\mathbf{j}, \mathbf{k}}}{\partial \mathbf{x}^{\boldsymbol{\alpha}}}(\mathbf{x}) \right| \ll 1. \quad (7.5)$$

(iii) *Under the additional assumption that $\mathbf{k} \in j_1 \mathcal{D}$, we have*

$$\sup_{\mathbf{x} \in \mathcal{D} \setminus \{x_{\mathbf{j}, \mathbf{k}}\}} \frac{|\mathbf{x} - \mathbf{x}_{\mathbf{j}, \mathbf{k}}|}{|\nabla \varphi^{\mathbf{j}, \mathbf{k}}(\mathbf{x})|} \ll 1, \quad (7.6)$$

where

$$\mathbf{x}_{\mathbf{j}, \mathbf{k}} := (\nabla F_{\mathbf{j}})^{-1}(\mathbf{k}/j_1) \in \mathcal{D}. \quad (7.7)$$

The implicit constants in the estimates above depend on $\boldsymbol{\alpha}$ (for the first three inequalities), upper bounds for finitely many derivatives of f_r ($1 \leq r \leq R$) as well as w on U , and the constant \mathfrak{C}_0 in the condition (3.25), but are independent of \mathbf{j} and \mathbf{k} .

Proof. See Lemmas 5.1 and 5.2 in [21]. □

We divide our consideration into the stationary, non-stationary and intermediate regimes. Recall the sets $\mathcal{D} = B_{2\varepsilon_0}(\mathbf{x}_0)$ and $U := \text{supp } w \subseteq B_{\varepsilon_0}(\mathbf{x}_0)$ from (3.24). Let

$$\rho := \frac{1}{2} \inf_{\mathbf{t} \in [0, 2]^{R-1}} \text{dist} \left(\partial \left(\nabla_{\mathbf{x}}(f_1 + \sum_{r=2}^R t_r f_r)(\mathcal{D}) \right), \partial \left(\nabla(f_1 + \sum_{r=2}^R t_r f_r)(B_{\varepsilon_0}(\mathbf{x}_0)) \right) \right) > 0. \quad (7.8)$$

For each $\mathbf{j} \in \mathcal{J}(J_1)_1$, we split $\mathbb{Z}^n = \mathcal{K}_{\mathbf{j}, 1} \cup \mathcal{K}_{\mathbf{j}, 2} \cup \mathcal{K}_{\mathbf{j}, 3}$ where

$$\begin{aligned} \mathcal{K}_{\mathbf{j}, 1} &:= \left\{ \mathbf{k} \in \mathbb{Z}^n : \frac{\mathbf{k}}{j_1} \in V_{\mathbf{j}} \right\}, \\ \mathcal{K}_{\mathbf{j}, 2} &:= \left\{ \mathbf{k} \in \mathbb{Z}^n : \text{dist} \left(\frac{\mathbf{k}}{j_1}, V_{\mathbf{j}} \right) \geq \rho \right\}, \text{ and} \\ \mathcal{K}_{\mathbf{j}, 3} &:= \mathbb{Z}^n \setminus (\mathcal{K}_{\mathbf{j}, 1} \cup \mathcal{K}_{\mathbf{j}, 2}). \end{aligned}$$

For $i \in \{1, 2, 3\}$, we define the respective contributions

$$M_i(Q, \boldsymbol{\delta}) := \sum_{\mathbf{j} \in \mathcal{J}(J_1)} \left(\prod_{r=1}^R \frac{1}{1 + j_r} \right) \left| \sum_{\substack{\mathbf{k} \in \mathcal{K}_{\mathbf{j}, i} \\ 1 \leq q \leq Q}} q^n I(q, \mathbf{j}, \mathbf{k}) \right|. \quad (7.9)$$

By (7.2), we then have

$$\mathfrak{N}_{w, \mathcal{M}}^1(Q, \boldsymbol{\delta}) \leq C_w \boldsymbol{\delta}^{\times Q} Q^{n+1} + M_1(Q, \boldsymbol{\delta}) + M_2(Q, \boldsymbol{\delta}) + M_3(Q, \boldsymbol{\delta}). \quad (7.10)$$

We first estimate the contribution from the non-stationary regime $\mathcal{K}_{\mathbf{j}, 2}$.

Lemma 7.2.

$$M_2(Q, \boldsymbol{\delta}) \ll (\log 4Q)(\log 4J_1)^R,$$

with the implicit constant depending only on \mathcal{M} and w .

Proof. Recall $\varphi^{\mathbf{j}, \mathbf{k}}$ from (7.3), and let

$$\lambda_1 = q \cdot \text{dist}(k, j_1 V_{\mathbf{j}})$$

For each $\mathbf{k} \in \mathcal{K}_{\mathbf{j}, 2}$, we have the lower bound

$$\inf_{\mathbf{x} \in U} |\nabla \varphi_1^{\mathbf{j}, \mathbf{k}}(\mathbf{x})| \geq 1.$$

Further, by part (i) of Lemma 7.1, we know that the derivatives of $\varphi_1^{\mathbf{j}, \mathbf{k}}$ are bounded on U independently of \mathbf{j} and \mathbf{k} . Thus we can apply Lemma 2.1 (integration by parts), with phase $\varphi_1^{\mathbf{j}, \mathbf{k}}$ and $\lambda = \lambda_1$, to conclude that

$$I(q, \mathbf{j}, \mathbf{k}) \ll_m \lambda_1^{-m+1} = (q \cdot \text{dist}(\mathbf{k}, j_1 V_{\mathbf{j}}))^{-m+1},$$

for $m \in \mathbb{Z}_{\geq 0}$, with implicit constants independent of q, \mathbf{j} and \mathbf{k} . In particular, taking $m \geq n+2$ and arguing like in the proof of Lemma 6.3, we get

$$\sum_{\mathbf{k} \in \mathcal{K}_{\mathbf{j}, 2}} I(q, \mathbf{j}, \mathbf{k}) \ll q^{-n-1} \sum_{i=0}^{\infty} 2^{(i+1)(-n-1+n)} \leq q^{-n-1}.$$

Thus

$$\begin{aligned} M_2(Q, \boldsymbol{\delta}) &\ll \sum_{\mathbf{j} \in \mathcal{J}(J_1)} \left(\prod_{r=1}^R \frac{1}{1+j_r} \right) \left| \sum_{\substack{\mathbf{k} \in \mathcal{K}_{\mathbf{j}, 2} \\ 1 \leq q \leq Q}} q^n I(q, \mathbf{j}, \mathbf{k}) \right| \ll \sum_{j_1=1}^{J_1} \sum_{j_2, \dots, j_R=0}^{j_1} \left(\prod_{r=1}^R \frac{1}{1+j_r} \right) \sum_{q=1}^Q q^{n-n-1} \\ &\ll (\log 4Q)(\log 4J_1)^R. \end{aligned}$$

□

Recall $\varphi^{\mathbf{j}, \mathbf{k}}$ from (7.3). Since $\nabla F_{\mathbf{j}}$ is a diffeomorphism mapping \mathcal{D} onto $\mathcal{R}_{\mathbf{j}}$, each $\mathbf{k} \in j_1 \mathcal{R}_{\mathbf{j}}$ determines a unique critical point of $\varphi^{\mathbf{j}, \mathbf{k}}$ given by

$$x_{\mathbf{j}, \mathbf{k}} := (\nabla F_{\mathbf{j}})^{-1}(\mathbf{k}/j_1) \in \mathcal{D}.$$

For the intermediate regime, we have the following estimate.

Lemma 7.3.

$$M_3(Q, \boldsymbol{\delta}) \ll Q^{\frac{n}{2}} J_1^{\frac{n}{2}-1} (\log 4J_1)^R,$$

with the implicit constant depending only on \mathcal{M} and w .

Proof. For $\mathbf{k} \in \mathcal{K}_{\mathbf{j}, 3} \subseteq \mathcal{R}_{\mathbf{j}} \setminus j_1 V_{\mathbf{j}}$, we know that $x_{\mathbf{j}, \mathbf{k}} \notin (\nabla F_{\mathbf{j}})^{-1}(V_{\mathbf{j}}) = U \supset \text{supp } w$. Recall that the phase $\varphi^{\mathbf{j}, \mathbf{k}}$ satisfies the estimates (7.5). Further,

$$H_{\varphi^{\mathbf{j}, \mathbf{k}}}(x_{\mathbf{j}, \mathbf{k}}) = H_{F_{\mathbf{j}}}((\nabla F_{\mathbf{j}})^{-1}(\mathbf{k}/j_1)) \gg \mathfrak{C}_0^{-1},$$

by (3.28) and (3.25). Thus, we can apply Lemma 2.1 (stationary phase principle) with phase $\varphi^{\mathbf{j}, \mathbf{k}}$ and $\lambda := qj_1$, to conclude that

$$I(q, \mathbf{j}, \mathbf{k}) \ll (qj_1)^{-\frac{n}{2}-1}$$

for each $\mathbf{k} \in \mathcal{K}_{j,3}$. Since $\#\mathcal{K}_3 \ll_{U,\rho} j_1^n$, we can estimate

$$\begin{aligned} M_3(Q, \delta) &\ll \sum_{\mathbf{j} \in \mathcal{J}(J_1)} \left(\prod_{r=1}^R \frac{1}{1+j_r} \right) j_1^{n-\frac{n}{2}-1} \sum_{1 \leq q \leq Q} q^{n-\frac{n}{2}-1} \leq \left(\sum_{\mathbf{j} \in \mathcal{J}(J_1)} \prod_{r=1}^R \frac{1}{1+j_r} \right) Q^{\frac{n}{2}} j_1^{\frac{n}{2}-1} \\ &\leq Q^{\frac{n}{2}} j_1^{\frac{n}{2}-1} (\log 4J_1)^R. \end{aligned}$$

□

Finally, we estimate the contribution from the critical stationary phase regime.

Lemma 7.4.

$$\begin{aligned} M_1(Q, \delta) &\ll Q^{\frac{n}{2}} \sum_{\mathbf{j} \in \mathcal{J}^1(J_1)} \left(\prod_{r=1}^R b_{j_r} \right) \sum_{\mathbf{k} \in \mathbb{Z}^n} \frac{w_{\mathbf{j}}^* \left(\frac{\mathbf{k}}{j_1} \right)}{\sqrt{|\det H_{F_{\mathbf{j}}}((\nabla F_{\mathbf{j}})^{-1}(\mathbf{k}/j_1))|}} j_1^{-\frac{n}{2}} \min(\|j_1 F_{\mathbf{j}}^*(\mathbf{k}/j_1)\|^{-1}, Q) \\ &\quad + Q^{\frac{n}{2}} j_1^{\frac{n}{2}-1} (\log 4J_1)^R, \end{aligned}$$

with implicit constants depending only on \mathcal{M} and w .

Proof. We intend to apply the stationary phase principle to evaluate the integrals $I(q, \mathbf{k}, \mathbf{j})$ for $\mathbf{k} \in j_1 V_{\mathbf{j}}$. As in the proof of Lemma 7.3, we note that the phase $\varphi^{\mathbf{j}, \mathbf{k}}$ satisfies the estimates (7.5). Further, by (3.28) and (3.25), we know that

$$H_{\varphi^{\mathbf{j}, \mathbf{k}}}(x_{\mathbf{j}, \mathbf{k}}) = H_{F_{\mathbf{j}}}((\nabla F_{\mathbf{j}})^{-1}(\mathbf{k}/j_1)) \gg \mathfrak{C}_0^{-1}. \quad (7.11)$$

An application of the stationary phase principle (Lemma 2.2) with $\lambda = dj_1$, phase $\varphi^{\mathbf{j}, \mathbf{k}}$ and amplitude function w , gives

$$I(q, \mathbf{j}, \mathbf{k}) = \frac{w(\mathbf{x}_{\mathbf{j}, \mathbf{k}})}{\sqrt{|\det H_{F_{\mathbf{j}}}((\nabla F_{\mathbf{j}})^{-1}(\mathbf{k}/j_1))|}} (qj_1)^{-\frac{n}{2}} e(-qj_1 \varphi^{\mathbf{j}, \mathbf{k}}(\mathbf{x}_{\mathbf{j}, \mathbf{k}}) + \sigma/8) + O((qj_1)^{-\frac{n}{2}-1}).$$

For $\mathbf{k} \in \mathcal{K}_{j,1}$, we have

$$w(\mathbf{x}_{\mathbf{j}, \mathbf{k}}) = w((\nabla F_{\mathbf{j}})^{-1}(\mathbf{k}/j_1)) = w_{\mathbf{j}}^* \left(\frac{\mathbf{k}}{j_1} \right). \quad (7.12)$$

We can also simplify

$$\varphi^{\mathbf{j}, \mathbf{k}}(\mathbf{x}_{\mathbf{j}, \mathbf{k}}) = (F_{\mathbf{j}} \circ (\nabla F_{\mathbf{j}})^{-1}) \left(\frac{\mathbf{k}}{j_1} \right) - \frac{\mathbf{k}}{j_1} \cdot (\nabla F_{\mathbf{j}})^{-1} \left(\frac{\mathbf{k}}{j_1} \right) = -F_{\mathbf{j}}^* \left(\frac{\mathbf{k}}{j_1} \right). \quad (7.13)$$

Plugging (7.12) and (7.13) into the stationary phase expansion for $I(q, \mathbf{j}, \mathbf{k})$, we get

$$I(q, \mathbf{j}, \mathbf{k}) = \frac{w^* \left(\frac{\mathbf{k}}{j_1} \right)}{\sqrt{|\det H_{F_{\mathbf{j}}}((\nabla F_{\mathbf{j}})^{-1}(\mathbf{k}/j_1))|}} (qj_1)^{-\frac{n}{2}} e \left(-qj_1 F_{\mathbf{j}}^* \left(\frac{\mathbf{k}}{j_1} \right) + \frac{\sigma}{8} \right) + O((qj_1)^{-\frac{n}{2}-1}). \quad (7.14)$$

By geometric and partial summation, we have

$$\sum_{q=1}^Q q^n I(q, \mathbf{j}, \mathbf{k}) \ll Q^{\frac{n}{2}} \frac{w^* \left(\frac{\mathbf{k}}{j_1} \right)}{\sqrt{|\det H_{F_{\mathbf{j}}}((\nabla F_{\mathbf{j}})^{-1}(\mathbf{k}/j_1))|}} j_1^{-\frac{n}{2}} \min(\|j_1 F_{\mathbf{j}}^*(\mathbf{k}/j_1)\|^{-1}, Q) + Q^{\frac{n}{2}} j_1^{-\frac{n}{2}-1}.$$

Therefore

$$\begin{aligned} M_1(Q, \delta) &\ll Q^{\frac{n}{2}} \sum_{\mathbf{j} \in \mathcal{J}(J_1)} \left(\prod_{r=1}^R \frac{1}{1+j_r} \right) \sum_{\mathbf{k} \in \mathbb{Z}^n} \frac{w_{\mathbf{j}}^* \left(\frac{\mathbf{k}}{j_1} \right)}{\sqrt{|\det H_{F_{\mathbf{j}}}((\nabla F_{\mathbf{j}})^{-1}(\mathbf{k}/j_1))|}} j_1^{-\frac{n}{2}} \min(\|j_1 F_{\mathbf{j}}^*(\mathbf{k}/j_1)\|^{-1}, Q) \\ &+ \sum_{\mathbf{j} \in \mathcal{J}(J_1)} \left(\prod_{r=1}^R \frac{1}{1+j_r} \right) \sum_{\mathbf{k} \in \mathcal{K}_{\mathbf{j},1}} Q^{\frac{n}{2}} j_1^{-\frac{n}{2}-1}. \end{aligned} \quad (7.15)$$

Arguing as in the proof of Lemma 7.3, we can deduce that $\#\mathcal{K}_{\mathbf{j},1} \ll j_1^n$. Thus the second term in (7.15) can be estimated as below

$$\sum_{\mathbf{j} \in \mathcal{J}(J_1)} \left(\prod_{r=1}^R \frac{1}{1+j_r} \right) \sum_{\mathbf{k} \in \mathcal{K}_{\mathbf{j},1}} Q^{\frac{n}{2}} j_1^{-\frac{n}{2}-1} \ll Q^{\frac{n}{2}} \sum_{j_1=1}^{J_1} \sum_{j_2, \dots, j_R=0}^{j_1} \left(\prod_{r=1}^R \frac{1}{1+j_r} \right) j_1^{\frac{n}{2}-1} \leq Q^{\frac{n}{2}} J_1^{\frac{n}{2}-1} (\log 4J_1)^R.$$

This establishes Lemma 7.4. \square

Concluding the proof of Proposition 3.5. The required estimate follows by combining (7.10) with Lemmas 7.2-7.4.

8. PROOF OF THEOREM 1.4

Let

$$\mathfrak{N}_0(Q) := \sum_{\substack{\mathbf{a} \in \mathbb{Z}^n \\ 1 \leq q \leq Q}} w \left(\frac{\mathbf{a}}{q} \right). \quad (8.1)$$

Applying the Poisson summation formula and using the rapid decay of \hat{w} , we get

$$\mathfrak{N}_0 = \sum_{1 \leq q \leq Q} q^n \sum_{\mathbf{k} \in \mathbb{Z}^n} \hat{w}(q\mathbf{k}) = \frac{\hat{w}(\mathbf{0})}{n} Q^{n+1} + O(Q^n). \quad (8.2)$$

Next, we use an R -fold product of the Selberg magic functions of degree X to estimate the characteristic function of the set $(-\delta, \delta)$. Let

$$b_{j_r} := \frac{1}{X+1} + \min \left\{ 2\delta, \frac{1}{j_r} \right\},$$

and

$$E_{s,\gamma}(Q, X) := \sum_{\substack{\mathbf{j} \in \mathbb{Z}_{\geq 0}^R \\ \|\mathbf{j}\|_{\infty} = j_s \in [1, X]}} \left(\prod_{r=1}^R b_{j_r} \right) \left| \sum_{\substack{\mathbf{a} \in \mathbb{Z}^n \\ 1 \leq q \leq Q}} w \left(\frac{\mathbf{a}}{q} \right) \exp \left(\sum_{r=1}^R \gamma_r j_r q f_r \left(\frac{\mathbf{a}}{q} \right) \right) \right|. \quad (8.3)$$

Separating the term corresponding to $j_1 = \dots = j_R = 0$ and using (2.3), we can write

$$|\mathfrak{N}_{w,\mathcal{M}}(Q, \delta) - 2\delta^R \mathfrak{N}_0| \ll \delta^{R-1} \frac{1}{X} Q^{n+1} + \frac{1}{X^R} Q^{n+1} + \sum_{s=1}^R \sum_{\gamma \in \{-1,1\}^R} E_{s,\gamma}(Q, X), \quad (8.4)$$

For ease of exposition, let us focus on the case when $s = 1$ and $\gamma = (1, 1, \dots, 1) \in \mathbb{Z}^R$; the other cases can be dealt with using the same argument. We shall suppress notation and refer

to $E_{1,(1,\dots,1)}(Q, X)$ simply as $E(Q, X)$. Arguing the same way as in the proof of Proposition 3.5, we can conclude that

$$E(Q, X) \ll Q^{\frac{n}{2}} \sum_{\substack{1 \leq j_1 \leq X \\ 0 \leq j_{r'} \leq j_1 \\ 2 \leq r' \leq R}} \left(\prod_{r=1}^R b_{j_r} \right) \sum_{\mathbf{k} \in \mathbb{Z}^n} \frac{w_{\mathbf{j}}^* \left(\frac{\mathbf{k}}{j_1} \right)}{\sqrt{|\det H_{F_{\mathbf{j}}}((\nabla F_{\mathbf{j}})^{-1}(\mathbf{k}/j_1))|}} j_1^{-\frac{n}{2}} \min(\|j_1 F_{\mathbf{j}}^*(\mathbf{k}/j_1)\|^{-1}, Q) \\ + X^{\frac{n}{2}-1} Q^{\frac{n}{2}} (\log 4X)^R. \quad (8.5)$$

with the implicit constant depending only on w and \mathcal{M} . We recall that the function $F_{\mathbf{j}}$ is given by

$$F_{\mathbf{j}}(\mathbf{x}) := f_1(\mathbf{x}) + \sum_{r=2}^R \frac{j_r}{j_1} f_r(\mathbf{x}).$$

We will need two preliminary lemmas. The first one is a consequence of Theorem 1.9.

Lemma 8.1. *Let α_{st} be as defined in (3.31). There exists a positive constant C_3 depending only on w and \mathcal{M} such that for all $Q^* \geq 1$ and for all $\delta^* \in (0, 1/2)$, we have*

$$\sum_{\substack{1 \leq j_1 \leq Q^* \\ 0 \leq j_{r'} \leq j_1 \\ 2 \leq r' \leq R}} \sum_{\substack{\mathbf{k} \in \mathbb{Z}^n \\ \|j_1 F_{\mathbf{j}}^*(\mathbf{k}/j_1)\| < \delta^*}} \frac{w_{\mathbf{j}}^* \left(\frac{\mathbf{k}}{j_1} \right)}{\sqrt{|\det H_{F_{\mathbf{j}}}((\nabla F_{\mathbf{j}})^{-1}(\mathbf{k}/j_1))|}} \leq C_3 \delta^* (Q^*)^{n+R} + C_3 (Q^*)^{\alpha_{st}} \mathcal{E}_n(Q^*), \quad (8.6)$$

where $\mathcal{E}_n(Q)$ is as defined in (1.8).

Proof. The proof proceeds in exactly the same way as that of Proposition 3.2, except that we use the conclusion of Theorem 1.9 instead of (3.32) as our input. Indeed, by Theorem 1.9, we know that for all $Q \geq 1$ and for all $\delta \in (0, 1/2)^R$,

$$\mathfrak{N}_{w, \mathcal{M}}(Q, \delta) \ll \mathcal{E}_n(Q) \left(\delta^R Q^{n+1} + \sum_{r=1}^R \delta^{R-1} Q^{n+1 + \frac{\Theta - (n+1)}{R}} + Q^{\Theta} \right),$$

with Θ is as defined in (1.6). Then by following the proof of Proposition 3.2, we can conclude that there exists a positive constant C_3 depending only on w and \mathcal{M} , such that for all $Q^* \geq 1$ and for all $\delta^* \in (0, 1/2)$, we have

$$\mathfrak{N}_{w, \mathcal{M}}^{*,1}(Q^*, \delta^*) \leq C_3 \delta^* (Q^*)^{n+R} + C_3 (Q^*)^{\alpha_{st}} \mathcal{E}_n(Q^*),$$

with

$$\alpha_{st} = n + R - \frac{n}{2\Theta - n} = \max \left(\frac{n(n+R+1)}{n+2}, n + R - 1 - \frac{2}{n} \right).$$

Here $\mathfrak{N}_{w, \mathcal{M}}^{*,1}(Q^*, \delta^*)$ is as defined in (3.30) and is exactly the left hand side of (8.6). Thus we are done. \square

The second lemma takes the conclusion of the previous one as input to derive an estimate for $E(Q, X)$, using (8.5). The proof is exactly the same as the first half of the proof of Proposition 3.3, using Proposition 3.5.

Lemma 8.2. *There exists a positive constant C_4 depending only on w and \mathcal{M} such that for all $Q \geq 1$ and for all $X \in (2, \infty)$, we have*

$$E(Q, X) \leq C_4 \mathcal{E}_n(X) (\log 4X)^R \left((\log 4Q) Q^{\frac{n}{2}} X^{\frac{n}{2}} + Q^{\frac{n}{2}+1} X^{\alpha_{st} - \frac{n}{2} - R} \right).$$

Proof. We proceed as in the proof of Proposition 3.3. We make a dyadic decomposition as in (5.2) based on the size of $\|j_1 F_j^*(\mathbf{k}/j_1)\|$ and then use Lemma 8.1 and partial summation on the j_r variables, to conclude that for each $0 \leq i \leq \frac{\log 4Q}{\log 2}$, we have

$$\begin{aligned} & \sum_{\substack{1 \leq j_1 \leq X \\ 0 \leq j_r \leq X \\ 2 \leq r \leq R}} \left(\prod_{r=1}^R b_{j_r} \right) \sum_{\substack{\mathbf{k} \in \mathbb{Z}^n \\ \|j_1 F_j^*(\mathbf{k}/j_1)\| \leq \frac{2^{i+1}}{Q}}} \frac{w_j^*\left(\frac{\mathbf{k}}{j_1}\right) j_1^{-\frac{n}{2}}}{\sqrt{|\det H_{F_j}((\nabla F_j)^{-1}(\mathbf{k}/j_1))|}} \\ & \ll \mathcal{E}_n(X) (\log 4X)^R X^{-\frac{n}{2}} (2^{i+1} Q^{-1} X^n + X^{\alpha_{\text{st}} - R}). \end{aligned}$$

Using (8.5) and summing up in the dyadic scales i , we get

$$\begin{aligned} E(Q, X) & \ll Q^{\frac{n}{2}} \sum_{\substack{1 \leq j_1 \leq J_1 \\ 0 \leq j_r \leq X \\ 2 \leq r \leq R}} \left(\prod_{r=1}^R b_{j_r} \right) \sum_{\mathbf{k} \in \mathbb{Z}^n} \frac{w_j^*\left(\frac{\mathbf{k}}{j_1}\right)}{\sqrt{|\det H_{F_j}((\nabla F_j)^{-1}(\mathbf{k}/j_1))|}} j_1^{-\frac{n}{2}} \min(\|j_1 F_j^*(\mathbf{k}/j_1)\|^{-1}, Q) \\ & + X^{\frac{n}{2}-1} Q^{\frac{n}{2}} \\ & \ll \mathcal{E}_n(X) Q^{\frac{n}{2}+1} (\log 4X)^R X^{-\frac{n}{2}} \sum_{0 \leq i \leq \frac{\log 4Q}{\log 2}} 2^{-i} (2^{i+1} Q^{-1} X^n + X^{\alpha_{\text{st}} - R}) \\ & \ll \mathcal{E}_n(X) (\log 4X)^R ((\log 4Q) Q^{\frac{n}{2}} X^{\frac{n}{2}} + Q^{\frac{n}{2}+1} X^{\alpha_{\text{st}} - \frac{n}{2} - R}). \end{aligned}$$

□

We now return to the proof of Theorem 1.4. The estimate in Lemma 8.2 for each $E^{r,\gamma}(X, Q)$, independently of Q and X . Therefore, (8.4) yields

$$\begin{aligned} & |\mathfrak{N}_{w, \mathcal{M}}(Q, \delta) - 2\delta^R \mathfrak{N}_0| \\ & \ll \delta^{R-1} \frac{1}{X} Q^{n+1} + \frac{1}{X^R} Q^{n+1} + \mathcal{E}_n(X) (\log 4X)^R ((\log 4Q) Q^{\frac{n}{2}} X^{\frac{n}{2}} + Q^{\frac{n}{2}+1} X^{\alpha_{\text{st}} - \frac{n}{2} - R}). \end{aligned}$$

Plugging in (8.2), we get

$$\begin{aligned} & \left| \mathfrak{N}_{w, \mathcal{M}}(Q, \delta) - \frac{2\hat{w}(\mathbf{0})}{n} \delta^R Q^{n+1} \right| \\ & \ll \delta^{R-1} \frac{1}{X} Q^{n+1} + \frac{1}{X^R} Q^{n+1} + \mathcal{E}_n(X) (\log 4X)^R ((\log 4Q) Q^{\frac{n}{2}} X^{\frac{n}{2}} + Q^{\frac{n}{2}+1} X^{\alpha_{\text{st}} - \frac{n}{2} - R}) + O(Q^n). \end{aligned}$$

We now choose our parameter X and set it to be $Q^{\frac{n+2}{n+2R}}$. This yields

$$\begin{aligned} & \left| \mathfrak{N}_{w, \mathcal{M}}(Q, \delta) - \frac{2\hat{w}(\mathbf{0})}{n} \delta^R Q^{n+1} \right| \\ & \ll \delta^{R-1} Q^{n+1 - \frac{n+2}{n+2R}} + Q^{n+1 - \frac{(n+2)R}{n+2R}} + \mathcal{E}_n(Q) (\log 4Q)^R \left((\log 4Q) Q^{\frac{n(n+R+1)}{n+2R}} + Q^{\frac{n}{2}+1} Q^{\left(\frac{n+2}{n+2R}\right)\left(\frac{n(n+R+1)}{n+2} - \frac{n}{2} - R\right)} \right) \\ & + O(Q^n) \\ & \ll \delta^{R-1} Q^{n+1 - \frac{n+2}{n+2R}} + \mathcal{E}_n(Q) Q^{\frac{n(n+R+1)}{n+2R}} = \delta^{R-1} Q^{\frac{(n+1)(R-1)}{R} + \frac{n(n+R+1)}{R(n+2R)}} + \mathcal{E}_n(Q) Q^{\frac{n(n+R+1)}{n+2R}}, \end{aligned}$$

with the constants \mathfrak{c}_2 (or \mathfrak{c}_1) in the definition of $\mathcal{E}_n(Q)$ chosen large enough. This establishes (1.7), and hence Theorem 1.4.

9. PROOF OF THEOREM 1.12

Since ψ is monotonic, by a slight generalisation of Cauchy's condensation test, (1.22) being convergent is equivalent to

$$\sum_{i=1}^{\infty} 2^{i(n+1)} \left(\frac{\psi_0(2^i)}{2^i} \right)^s \prod_{r=1}^R \psi_r(2^i) < \infty. \quad (9.1)$$

Let $\boldsymbol{\psi} := (\psi_0, \psi_1, \dots, \psi_R)$ and $\mathcal{P}_{\mathcal{M}}(\boldsymbol{\psi})$ denote the projection of $\mathcal{M} \cap \mathcal{S}_{n+R}(\boldsymbol{\psi})$ onto $\mathcal{D} \subseteq \mathbb{R}^n$. Since the functions f_1, \dots, f_R are smooth, this projection is bi-Lipschitz. Consequently,

$$\mathcal{H}^s(\mathcal{M} \cap \mathcal{S}_{n+R}(\boldsymbol{\psi})) = 0 \iff \mathcal{H}^s(\mathcal{P}_{\mathcal{M}}(\boldsymbol{\psi})) = 0,$$

and it suffices to show the latter equality. As $\mathcal{H}^s(\mathcal{P}_{\mathcal{M}}(\boldsymbol{\psi})) = 0$ for $s > n$, we may assume without loss of generality that $s \leq n$. To be able to use Theorem 1.4, by replacing $\psi_r(q)$ and with

$$\max \left(\psi_r(q), q^{-\frac{n+2}{n+2R}+\eta}, q^{-\frac{n}{n+2(R-1)-\frac{4}{n}+\eta}} \right),$$

if need be, we may also assume that

$$\psi_r(q) \geq \max \left(q^{-\frac{n+2}{n+2R}+\eta}, q^{-\frac{n}{n+2(R-1)-\frac{4}{n}+\eta}} \right), \quad \text{for all } q \in \mathbb{Z}_{>0} \text{ and } 0 \leq r \leq R. \quad (9.2)$$

For $(q, \mathbf{p}) = (q, \mathbf{a}, \mathbf{b}) \in \mathbb{Z}_{>0} \times \mathbb{Z}^n \times \mathbb{Z}^R$, we define

$$\sigma(\mathbf{p}/q) := \left\{ \mathbf{x} \in \mathcal{P}_{\mathcal{M}}(\boldsymbol{\psi}) : \left\| \mathbf{x} - \frac{\mathbf{a}}{q} \right\|_2 \leq \frac{\psi_0(q)}{q}, \left| f_r(\mathbf{x}) - \frac{b_r}{q} \right| \leq \frac{\psi_r(q)}{q} \text{ for } 1 \leq r \leq R \right\}. \quad (9.3)$$

Then $\sigma(\mathbf{p}/q)$ is contained in a set of radius $\frac{\psi_0(q)}{q}$. Moreover, if $\sigma(\mathbf{p}/q) \neq \emptyset$, then by the triangle inequality and using the Lipschitz property of f_r , we can conclude that

$$\left| f_r \left(\frac{\mathbf{a}}{q} \right) - \frac{b_r}{q} \right| \ll \frac{\psi_0(q)}{q} + \frac{\psi_r(q)}{q} \ll \frac{\psi_r(q)}{q},$$

with the implicit constant depending only on f_r . For each $i \in \mathbb{Z}_{\geq 0}$, we define

$$B_i := \{(q, \mathbf{p}) \in \mathbb{Z}_{\geq 0} \times \mathbb{Z}^{n+R} : 2^i \leq q < 2^{i+1}, \sigma(\mathbf{p}/q) \neq \emptyset\}. \quad (9.4)$$

Then we have

$$\begin{aligned} \#B_i &\leq \#\{\mathbf{p}/q = (\mathbf{a}/q, \mathbf{b}/q) \in \mathbb{Q}^{n+R} : 1 \leq q < 2^{i+1}, \mathbf{a}/q \in \mathcal{D}, |f_r(\mathbf{a}/q) - b_r/q| \ll \psi_r(q)/q \text{ for } 1 \leq r \leq R\} \\ &\leq \#\{\mathbf{a}/q \in \mathbb{Q}^n : 1 \leq q \leq 2^i, \mathbf{a}/q \in \mathcal{D}, \|f_r(\mathbf{a}/q)\| \ll \psi_r(q)/q \text{ for } 1 \leq r \leq R\} \\ &\ll \mathfrak{N}_{w, \mathcal{M}}(2^{i+1}, \psi_1(2^i), \dots, \psi_R(2^i)) \ll 2^{i(n+1)} \prod_{r=1}^R \psi_r(2^i) \end{aligned}$$

where we used (9.2) and Corollary 1.10 to obtain the last estimate. Defining

$$\Sigma_i := \bigcup_{\mathbf{p}/q \in B_i} \sigma(\mathbf{p}/q),$$

we observe that

$$\mathcal{P}_{\mathcal{M}}(\boldsymbol{\psi}) = \bigcap_{T \geq 0} \bigcup_{i \geq T} \Sigma_i.$$

Now,

$$\mathcal{H}^s \left(\bigcup_{i \geq T} \Sigma_i \right) \leq \sum_{i \geq T} \sum_{(q, \mathbf{p}) \in B_i} \left(\frac{\psi_0(q)}{q} \right)^s \ll \sum_{i \geq T} 2^{i(n+1)} \left(\frac{\psi_0(2^i)}{2^i} \right)^s \prod_{r=1}^R \psi_r(2^i),$$

which converges to 0 as $T \rightarrow \infty$, due to (9.1). Therefore, we conclude that

$$\mathcal{H}^s(\mathcal{P}_{\mathcal{M}}(\psi)) = 0.$$

REFERENCES

- [1] J F Adams, *Vector fields on spheres*, Annals of Mathematics (1962), 603–632.
- [2] D Allen and B Wang, *A note on weighted simultaneous diophantine approximation on manifolds*, Journal de Theorie des Nombres de Bordeaux (2022).
- [3] Victor Beresnevich, *Rational points near manifolds and metric diophantine approximation*, Annals of Mathematics (2012), 187–235.
- [4] Victor Beresnevich, Jason Levesley, and Ben Ward, *A lower bound for the Hausdorff dimension of the set of weighted simultaneously approximable points over manifolds*, International Journal of Number Theory **17** (2021), no. 08, 1795–1814.
- [5] Victor Beresnevich and Lei Yang, *Khintchine’s theorem and diophantine approximation on manifolds*, Acta Mathematica (2023).
- [6] Victor V Beresnevich and Sanju L Velani, *A note on simultaneous diophantine approximation on planar curves*, Mathematische Annalen **337** (2007), no. 4, 769–796.
- [7] Wouter Castryck, Raf Cluckers, Philip Dittmann, and Kien Huu Nguyen, *The dimension growth conjecture, polynomial in the degree and without logarithmic factors*, Algebra & Number Theory **14** (2020), no. 8, 2261–2294.
- [8] Lars Hörmander, *The analysis of linear partial differential operators I. Distribution theory and Fourier analysis*, Grundlehren **256** (1990).
- [9] Jing-Jing Huang, *The density of rational points near hypersurfaces*, Duke Mathematical Journal **169** (2020), no. 11, 2045–2077.
- [10] ———, *Extremal affine subspaces and Khintchine-Jarník type theorems*, Geometric and Functional Analysis **34** (2024), no. 1, 113–163.
- [11] Adolf Hurwitz, *Über die Komposition der quadratischen Formen*, Math. Ann. **88** (1922), no. 1-2, 1–25. MR1512117
- [12] MN Huxley, *The rational points close to a curve*, Annali della Scuola Normale Superiore di Pisa-Classe di Scienze **21** (1994), no. 3, 357–375.
- [13] Aroldo Kaplan, *Fundamental solutions for a class of hypoelliptic PDE generated by composition of quadratic forms*, Trans. Amer. Math. Soc. **258** (1980), no. 1, 147–153. MR554324
- [14] Aleksandr Khintchine, *Zur metrischen theorie der diophantischen approximationen*, Mathematische Zeitschrift **24** (1926), no. 1, 706–714.
- [15] Hugh L Montgomery, *Ten lectures on the interface between analytic number theory and harmonic analysis*, American Mathematical Soc., 1994.
- [16] Florian Munkelt, *On the number of rational points close to a compact manifold under a less restrictive curvature condition*, arXiv preprint arXiv:2205.06183 (2022).
- [17] Johann Radon, *Lineare Scharen orthogonaler Matrizen*, Abh. Math. Sem. Univ. Hamburg **1** (1922), no. 1, 1–14. MR3069384
- [18] Joris Roos, Andreas Seeger, and Rajula Srivastava, *Lebesgue space estimates for spherical maximal functions on Heisenberg groups*, International Mathematics Research Notices **2022** (2022), no. 24, 19222–19257.
- [19] Per Salberger, *Counting rational points on projective varieties*, Proceedings of the London Mathematical Society **126** (2023), no. 4, 1092–1133.
- [20] Damaris Schindler, Rajula Srivastava, and Niclas Technau, *Rational points near manifolds, homogeneous dynamics, and oscillatory integrals*, arXiv preprint arXiv:2310.03867 (2023).
- [21] Damaris Schindler and Shuntaro Yamagishi, *Density of rational points near/on compact manifolds with certain curvature conditions*, Advances in Mathematics **403** (2022), 108358.
- [22] Wolfgang M Schmidt, *Diophantine approximation*, Vol. 785, Springer, 2009.
- [23] Rajula Srivastava and Niclas Technau, *Density of rational points near flat/rough hypersurfaces*, arXiv preprint arXiv:2305.01047 (2023).
- [24] Elias M Stein and Timothy S Murphy, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Vol. 3, Princeton University Press, 1993.
- [25] Robert C Vaughan and Sanju Velani, *Diophantine approximation on planar curves: the convergence theory*, Inventiones mathematicae **166** (2006), no. 1, 103–124.

MATHEMATICAL INSTITUTE, UNIVERSITY OF BONN, ENDENICHER ALLEE 60, 53115, BONN, GERMANY,
AND
MAX PLANCK INSTITUTE FOR MATHEMATICS, VIVATSGASSE 7, 53111, BONN, GERMANY.
Email address: `rajulas@math.uni-bonn.de`