Flows of SU(2)-structures

Udhav Fowdar^{1,2} and Henrique N. Sá Earp²

¹Università di Torino (UniTo) ²Universidade Estadual de Campinas (Unicamp)

31st July 2024

Abstract

This paper initiates a classification programme of flows of SU(2)-structures on 4-manifolds which have short-time existence and uniqueness. Our approach adapts a representation-theoretic method originally due to Bryant in the context of G_2 geometry. We show how this strategy can also be used to deduce the number of geometric flows of a given H-structure; we illustrate this in the G_2 , $Spin(7)$ and $SU(3)$ cases. Our investigation also leads us to derive explicit expressions for the Ricci and self-dual Weyl curvature in terms of the intrinsic torsion of the underlying SU(2)-structure. We compute the first variation formulae of all the quadratic functionals in the torsion; these provide natural building blocks for SU(2) gradient flows. In particular, our results demonstrate that both the negative gradient flow of the Dirichlet energy of the intrinsic torsion and the harmonic Ricci flow are parabolic after a modified DeTurck's trick.

Contents

1 Introduction

This article is a thorough study of variations of $SU(2)$ -structures on a connected and oriented 4-manifold M, as an instance of the abstract theory of flows of geometric H -structures, i.e., flows of tensor fields defining H -reductions of the frame bundle, for closed and connected subgroups $H \subset SO(n)$, developed in [\[LSE23,](#page-39-0) [FLMSE22\]](#page-39-1). The subject currently attracts substantial interest, most notably in the cases of $H = G_2$, $Spin(7)$, see references below and others therein.

In particular, the recent work by Dwivedi-Gianniotis-Karigiannis [\[DGK23\]](#page-39-2) brings forth a systematic study of the secondorder differential invariants of a G₂-structure, as tentative building blocks for *all possible* such geometric flows, up to lowerorder terms, which could be 'heat-like' [ibid.], i.e. parabolic. By sorting out those invariants which correspond to symmetric (2-tensors) and skew-symmetric (vector) objects, the authors are able to identify essentially all the *independent* such degrees of freedom for the construction of short-time flows. This in turn is a concrete expression of the representation-theoretic perspective on H-structure torsion introduced by Bryant in [\[Bry05\]](#page-39-3), which allows one to determine beforehand *how many* such independent invariants one ought to find, beyond which additional invariants will necessarily be determined by relations. In this context, our approach explicitly combines both narratives as a coherent programme, exploiting from the fact that SU(2) is 'sufficiently small' that its representation theory is still completely tractable, and yet 'sufficiently nontrivial' to exhibit several interesting phenomena tying in geometric torsion and Riemannian geometry. It is worth pointing out that the smaller the group H , the greater the number of invariants, and hence the more possible ways of evolving the underlying H -structure; thereby making a full classification of parabolic flows the more difficult.

1.1 $SU(2)$ -flows

Let M be a 4-manifold endowed with an SU(2)-structure determined by the triple of 2-forms $\omega_1, \omega_2, \omega_3$. Using the induced Riemannian metric q , we have the well-known decomposition^{[1](#page-1-2)}

$$
\text{End}(TM) \cong T^*M \otimes T^*M \cong \langle g \rangle \oplus \Sigma_0^2(M) \oplus \Lambda_+^2(M) \oplus \Lambda_-^2(M).
$$

While the latter decomposition is irreducible as a sum of $SO(4)$ -modules, it can be further split into $SU(2)$ -modules:

$$
\text{End}(TM) \cong \langle g \rangle \oplus \bigoplus_{i=1}^3 \Sigma^2_{0,i}(M) \oplus \bigoplus_{i=1}^3 \langle \omega_i \rangle \oplus \Lambda^2_-(M),
$$

where the irreducible spaces $\Sigma_{0,i}^2(M)$ are defined by [\(13\)](#page-4-0). Thus, given a general 2-tensor A, we can write $A = B + C \in$ $\Sigma^2(M)\oplus \Lambda^2(M)$ as

$$
B = f_0 g + B_{0,1} + B_{0,2} + B_{0,3},
$$

\n
$$
C = f_1 \omega_1 + f_2 \omega_2 + f_3 \omega_3 + C^2,
$$

where f_i are functions, each $B_{0,i}$ corresponds to the projection of the symmetric 2-tensor B onto $\Sigma_{0,i}^2(M)$, and C_{-}^2 is the anti-self-dual component of the 2-form C . In the above notation, following the general theory developed in [\[FLMSE22\]](#page-39-1), any variation of the SU(2)-triple $\omega := (\omega_1, \omega_2, \omega_3)$ can be expressed as

$$
\frac{d}{dt}\omega = A \diamond \omega,\tag{1}
$$

where \diamond denotes the endomorphism action, see [\(5\)](#page-3-3) for the precise definition. More explicitly, we can express [\(1\)](#page-1-3) as

$$
\frac{d}{dt}\omega_1 = +f_0\omega_1 + f_3\omega_2 - f_2\omega_3 + B_{0,1}\diamond\omega_1,\tag{2}
$$

$$
\frac{d}{dt}\omega_2 = -f_3\omega_1 + f_0\omega_2 + f_1\omega_3 + B_{0,2}\diamond \omega_2,\tag{3}
$$

$$
\frac{d}{dt}\omega_3 = +f_2\omega_1 - f_1\omega_2 + f_0\omega_3 + B_{0,3}\diamond\omega_3,\tag{4}
$$

where each $B_{0,i} \otimes \omega_i$ corresponds to an anti-self-dual 2-form on M. Since the term C^2_- lies in the kernel of $\otimes \omega$ it does not occur in [\(2\)](#page-1-4)-[\(4\)](#page-1-5) and hence, without loss of generality, we shall henceforth set it to zero. It follows that the infinitesimal deformations of $SU(2)$ -structures are given by sections of a rank 13 vector bundle; flows of $SU(2)$ -structures are therefore determined by particular choices of such sections.

This abstract framework for $SU(2)$ -flows bears relations with several well-known geometric flows in the literature:

• The SU(2)-flow [\(2\)](#page-1-4)-[\(4\)](#page-1-5) can be interpreted as a U(2)-flow, say for the multi-tensor (g, J_1, ω_1) , simply by setting $f_1 = 0$ and viewing ω_2 and ω_3 as being locally defined; or equivalently, by substituting the pair [\(3\)](#page-1-6)-[\(4\)](#page-1-5) by [\(16\)](#page-5-0), for (i, j, k) = $(1, 2, 3)$. In particular, the *Kähler-Ricci flow* is obtained when the almost complex structure J_1 is integrable and $d\omega_1 = 0$, so that $f_2 = f_3 = 0 = B_{0,2} = B_{0,3}$ and $B = f_0g + B_{0,1} = -2Ric(g)$. By contrast, the *Ricci flow* is obtained when $f_1 = f_2 = f_3 = 0$ and $B = -2Ric(g)$; this is called the *Ricci H-flow* in [\[FLMSE22,](#page-39-1) Example 1.26].

¹We denote the space of symmetric 2-tensors by $\Sigma^2(M)$ and reserve the notation S^2 for projection map of 2-tensors in $\Sigma^2(M)$

- • In $[LSE23]$ Loubeau and the second-named author introduced the *harmonic* H-flow. When $H = SU(2)$, the harmonic flow is obtained when $B = 0$ and $f_i = \text{div}(a_i)$, where a_i are the torsion 1-forms defined in Proposition [2.9.](#page-7-0) Stationary points of this flow correspond to H-structures which are in certain sense in Coulomb gauge, cf. [\[Gri17\]](#page-39-4) and also Corollary [2.13](#page-9-2) below.
- It is worth pointing out that the geometric flow considered here is different from the *hypersymplectic flow* introduced by Fine and Yao in [\[FY18\]](#page-39-5). In the latter framework, one starts with a triple of closed 2-forms and evolve them within their cohomology classes; the closed triple in [\[FY18\]](#page-39-5) do *not* constitute to an SU(2)-triple unless they are critical points. By contrast, in our setup we already start with an $SU(2)$ -triple and instead evolve them without a priori preserving any differential or cohomological conditions.

1.2 Main results

In this article, we classify all the second-order invariants of an $SU(2)$ -structure. By choosing the deformation A among suitable linear combinations of these invariants, we are able to classify all 'heat-like' flows. We then establish short-time existence and uniqueness for a large class of these SU(2)-flows, using a modified version of DeTurck's trick akin to the Ricci flow case in [\[DeT83\]](#page-39-6). While the theory is formally similar to the G_2 and $Spin(7)$ cases recently investigated in [\[DGK23,](#page-39-2) [Dwi24\]](#page-39-7), a few important differences emerge. First, our classification of these second-order invariants is entirely based on the representationtheoretic method of $[By05]$; indeed in many ways this paper is an adaptation of Bryant's ideas to the $SU(2)$ framework. Second, unlike with those groups, certain SU(2)-modules occur with multiplicities, and this gives rise to many more ways of using the same second-order invariants to define different flows. This article is an attempt to unify the ideas of $\frac{Bry05}{}$, [\[DGK23\]](#page-39-2) and [\[FLMSE22\]](#page-39-1), by proposing a systematic classification procedure for second-order flows of H-structures and their parabolicity regimes, which we carry out in detail for $H = SU(2)$. We highlight the following results, by relevance:

- We show in Theorem [4.2](#page-20-1) that there are 22 (independent) second-order SU(2)-invariants. With the exception of the anti-self-dual Weyl curvature, all the remaining 21 invariants can be used to evolve the underlying SU(2)-structure. In full generality this leads to a 72-parameter family of $SU(2)$ -flows.
- Our method of derivation provides a systematic way of classifying geometric flows of H-structures which evolve by its second-order invariants. We explain in $\S6.1$ how one can already deduce the number of possible second-order H flows only from the knowledge of its representation. We illustrate this concretely in the $SU(3)$, G_2 and $Spin(7)$ cases recovering some recent results of [\[DGK23,](#page-39-2)[Dwi24\]](#page-39-7).
- Using the second-order invariants, we derive an explicit expression for the Ricci curvature in terms of the intrinsic torsion of the $SU(2)$ -structure, see Proposition [4.4.](#page-21-1) This is analogous to Bryant's derivation of the Ricci curvature in terms of the intrinsic torsion of the G₂-structure in $Bry05$, §4.5.3]. Moreover, we also derive a corresponding expression for the self-dual Weyl curvature in Proposition [4.7.](#page-23-1)
- There are 15 SU(2)-invariant functionals which are quadratic in the intrinsic torsion. Arguably the most natural one is the L^2 -norm of the intrinsic torsion. In Theorem [5.4](#page-28-1) we prove short time existence and uniqueness of a 2-parameter family of $SU(2)$ -flow which includes its negative gradient flow and the coupled Ricci harmonic $SU(2)$ -flow. These provide natural $SU(2)$ -flows to study to find Einstein metrics, see also the recent article [\[BK24\]](#page-39-8) for another such functional.
- In [§5.4](#page-31-0) we show how one can extend Theorem [5.4](#page-28-1) to more general SU(2)-flows. In particular, we show that one can incorporate self-dual Weyl curvature terms in flows of SU(2)-structures which is rather different from flows of merely O(4)-structures, i.e. Riemannian metrics, which by contrast can only involve the Ricci curvature. We also demonstrate short time existence for the negative gradient flow of the weighted energy functional of the intrinsic torsion for suitable parameters.
- In [§3.3](#page-14-0) we compute the first variational formulae of the aforementioned 15 quadratic functionals (which by symmetry it suffices to consider only 5). Their negative gradient flows provide natural candidates of geometric flows to study, for instance, these can be used to evolve an almost complex structure by the negative gradient flow of the L^2 -norm of its Nijenhuis tensor. Thus, these provide building blocks of natural SU(2) functionals and their corresponding gradient flows. We expect these to be useful for constructing geometric flows beyond those considered in this article.

Acknowledgements: The authors are grateful to Shubham Dwivedi, Daniel Fadel, Spiro Karigiannis, Kirill Krasnov, and Andrés Moreno for several interesting discussions. The authors would also like to thank the organisers and participants of the *Special Holonomy and Geometric Structures on Complex Manifolds* workshop held at IMPA in March 2024, for their feedback on this work.

UF was funded by the São Paulo Research Foundation (Fapesp) [2021/07249-0] from November 2021 to January 2024, in a postdoctoral grant subordinate to the thematic project *Gauge theory and algebraic geometry* [2018/21391-1], led by Marcos Jardim. HSE was supported by the Fapesp [2021/04065-6] *BRIDGES collaboration* and a Brazilian National Council for Scientific and Technological Development (CNPq) [311128/2020-3] Productivity Grant level 1D.

1.3 Notation and conventions

We define the *diamond operator* \diamond , induced by the infinitesimal action of an endomorphism $v \otimes \alpha \in \mathfrak{gl}(4,\mathbb{R}) = \mathbb{R}^4 \otimes (\mathbb{R}^4)^*$ on k -tensors, by

$$
(v \otimes \alpha) \diamond (\beta_1 \otimes \ldots \otimes \beta_k) := \frac{1}{2} (\beta_1(v) \alpha \otimes \beta_2 \otimes \ldots \otimes \beta_k + \cdots + \beta_k(v) \beta_1 \otimes \ldots \otimes \beta_{k-1} \otimes \alpha).
$$
 (5)

This differs from the convention in [\[FLMSE22\]](#page-39-1), see Remark [2.1](#page-6-1) below. Given k-forms α, β , we define their inner product by

$$
g(\alpha,\beta) := \frac{1}{k!} g^{i_1 j_1} \dots g^{i_k j_k} \alpha_{i_1 \dots i_k} \beta_{j_1 \dots j_k}.
$$

With this convention, $|dx_{i_1...i_k}|^2 = 1$ with respect to the Euclidean metric; this contrasts with the conventions of [\[FLMSE22\]](#page-39-1), where the factor of $k!$ above is absent. On the other hand, for tensors we shall use the standard inner product. As a consequence of these differences in convention, the reader should be careful when comparing formulae in this article with other related papers in the literature, such as [\[Kar09,](#page-39-9) [FLMSE22,](#page-39-1) [DGK23,](#page-39-2) [DLSE24\]](#page-39-10).

Denoting cyclic permutation by ∼, we shall often use the Levi-Civita symbols

$$
\varepsilon_{ijk} := \begin{cases}\n+1 & \text{if } (i,j,k) \sim (1,2,3), \\
-1 & \text{if } (i,j,k) \sim (1,3,2), \\
0 & \text{otherwise.} \n\end{cases}
$$

For a given 1-form α and a vector field X, we define $J(\alpha)(X) := \alpha(JX)$, so that on 1-forms we have the relation

$$
J_i\circ J_j=-\varepsilon_{ijk}J_k.
$$

Also, given a smooth function f, we define $d^{c_i} f := J_i(df)$. These conventions might differ from others in the literature by a sign.

For simplicity, we shall often identify the SU(2)-modules $\otimes^k \mathbb{R}^4 \otimes^l (\mathbb{R}^4)^*$ (as well as their symmetrisations and skewsymmetrisations) with their associated vector bundles $\otimes^k TM \otimes^l T^*M$ and also with their space of sections; this should not cause any confusion for our purposes. We shall write l.o.t. (x) to mean lower-order differential terms in x.

2 Background on SU(2)-structures

2.1 Preliminaries

The main purpose of this section is to gather some algebraic facts about $SU(2)$ -structures that we shall use throughout this article and to fix the notation. We begin by recalling the basic definition.

Definition 2.1. An SU(2)-structure on a 4-manifold M is determined by a triple of 2-forms $\boldsymbol{\omega} = (\omega_1, \omega_2, \omega_3)$ which at each point can be expressed as

$$
\omega_1 = dx_{12} + dx_{34}, \n\omega_2 = dx_{13} + dx_{42}, \n\omega_3 = dx_{14} + dx_{23},
$$

for some local coordinates $\{x_i\}.$

The triple $\omega_1, \omega_2, \omega_3$ in turn defines a Riemannian metric g and a triple of almost complex structures J_1, J_2, J_3 on M^4 as follows:

$$
J_i X \lrcorner \omega_j := X \lrcorner \omega_k, \quad \text{for} \quad (i, j, k) \sim (1, 2, 3) \quad \text{and} \quad X \in \mathcal{X}(M). \tag{6}
$$

The metric q is then given by

$$
\omega_i(\cdot,\cdot) = g(J_i \cdot, \cdot). \tag{7}
$$

We also define a volume form on M by vol := $\frac{1}{2}\omega_1 \wedge \omega_1$ and the Hodge star operator $*$ by

$$
\alpha \wedge * \beta = g(\alpha, \beta) \text{vol}, \quad \text{for} \quad \alpha, \beta \in \Lambda^k(M)
$$
 (8)

In particular, note that $*$ and J_i commute on differential forms. It is also worth pointing out there are other equivalent definitions of $SU(2)$ -structures in the literature cf. $[CS07, FY18]$ $[CS07, FY18]$. Probably the most common one is:

Definition 2.2 (alternative). An SU(2)-structure on a 4-manifold M is given by a triple of non-degenerate 2-forms $\omega_1, \omega_2, \omega_3$ satisfying the conditions

$$
\omega_i\wedge\omega_j=\delta_{ij}\omega_1\wedge\omega_1,
$$

and that $\omega_3(X, Y) \geq 0$, for every pair of vector fields X, Y such that $X \perp \omega_1 = Y \perp \omega_2$. Here *non-degenerate* means that $\omega_i \wedge \omega_i$ is nowhere-vanishing and the second condition corresponds to an orientation of the triple.

Note that one could equivalently work with the data (q, J_1, J_2, J_3) instead of $(\omega_1, \omega_2, \omega_3)$, but the latter description fits better with the approach in the G_2 and $Spin(7)$ cases, as we shall see later on. We shall use the following frequently in computations:

Lemma 2.3. *Given arbitrary* 1*-forms* α , β *and* 2*-form* γ *, the following identities hold:*

- *1.* $\alpha \wedge \omega_1 = -J_3(\alpha) \wedge \omega_2 = J_2(\alpha) \wedge \omega_3$.
- 2. $*(\alpha \wedge \omega_i) = -J_i(\alpha)$.
- *3.* $\gamma \wedge \omega_i = q(\gamma, \omega_i)$ vol.
- *4.* $q(\alpha, J_i \beta) = q(\alpha \land \beta, \omega_i)$.

Given an oriented Riemannian 4-manifold (M^4, g, vol) , there is a natural orthogonal splitting of 2-forms into Hodge self-dual and anti-self-dual forms:

$$
\Lambda^{2}(M) = \Lambda^{2}_{+}(M) \oplus \Lambda^{2}_{-}(M) \cong \mathfrak{su}(2)^{+} \oplus \mathfrak{su}(2)^{-}.
$$
 (9)

Geometrically the data (q, vol) correspond to a structure group reduction of the frame bundle from $GL(4, \mathbb{R})$ to $SO(4)$. The existence of a global orthonormal basis of $\Lambda^2_+(M)$ allows one to further reduce the structure group from SO(4) to SU(2) := SU(2)[−]. Thus, as SU(2)-modules, we have the irreducible splitting

$$
\Lambda^2(M) = \langle \omega_1, \omega_2, \omega_3 \rangle \oplus \Lambda^2(M), \tag{10}
$$

where SU(2) acts trivially on each ω_i and $\Lambda^2(M) \cong \mathbb{R}^3$ corresponds to its adjoint representation. We define the projection map π^2 : $\Lambda^2(M) \to \Lambda^2_-(M)$ by

$$
\pi_-^2(\alpha) := \frac{1}{2}(\alpha - * \alpha). \tag{11}
$$

As SU(2)-modules, the spaces $\Lambda^1(M)$ and $\Lambda^3(M)$ are both irreducible and correspond to the standard representation of $\text{Sp}(1) \cong \text{SU}(2)$ on $\mathbb{H} \cong \mathbb{C}^2$.

Next we consider the space of symmetric 2-tensors $\Sigma^2(M)$. As an SO(4)-module, we have that

$$
\Sigma^2(M) = \langle g \rangle \oplus \Sigma_0^2(M),\tag{12}
$$

where $\Sigma_0^2(M)$ denotes the space of traceless symmetric 2-tensors. Under the action of SU(2), we can further decompose $\Sigma_0^2(M)$ into three irreducible representations $\Sigma_{0,1}^2(M)$, $\Sigma_{0,2}^2(M)$ and $\Sigma_{0,3}^2(M)$ – all isomorphic to the adjoint representation $\mathfrak{su}(2) \cong \mathbb{R}^3$. Explicitly, in terms of our conventions in [§1.3,](#page-3-0)

$$
\Sigma_{0,i}^{2}(M) = \{ \alpha \in \Sigma^{2}(M) \mid J_{i}(\alpha) = \alpha, J_{j}(\alpha) = J_{k}(\alpha) = -\alpha \}, \text{ for } (i, j, k) \sim (1, 2, 3). \tag{13}
$$

Thus a symmetric 2-tensor $\alpha \odot \beta := \alpha \otimes \beta + \beta \otimes \alpha \in \Sigma^2(M) = \langle g \rangle \oplus \Sigma^2_{0,1}(M) \oplus \Sigma^2_{0,2}(M) \oplus \Sigma^2_{0,3}(M)$ decomposes into four irreducible components:

$$
\alpha \odot \beta = +\frac{1}{4} (\alpha \odot \beta + J_1 \alpha \odot J_1 \beta + J_2 \alpha \odot J_2 \beta + J_3 \alpha \odot J_3 \beta)
$$

+
$$
\frac{1}{4} (\alpha \odot \beta + J_1 \alpha \odot J_1 \beta - J_2 \alpha \odot J_2 \beta - J_3 \alpha \odot J_3 \beta)
$$

+
$$
\frac{1}{4} (\alpha \odot \beta - J_1 \alpha \odot J_1 \beta + J_2 \alpha \odot J_2 \beta - J_3 \alpha \odot J_3 \beta)
$$

+
$$
\frac{1}{4} (\alpha \odot \beta - J_1 \alpha \odot J_1 \beta - J_2 \alpha \odot J_2 \beta + J_3 \alpha \odot J_3 \beta).
$$

Let us define the projection maps $S^2_{0,i}$, given on a simple 2-tensor $\alpha \otimes \beta$ by

$$
S_{0,1}^2: T^*(M) \otimes T^*(M) \to \Sigma_{0,1}^2(M)
$$

$$
S_{0,1}^2(\alpha \otimes \beta) := \frac{1}{8} (\alpha \odot \beta + J_1 \alpha \odot J_1 \beta - J_2 \alpha \odot J_2 \beta - J_3 \alpha \odot J_3 \beta),
$$

and likewise for $S^2_{0,2}(\alpha \otimes \beta)$ and $S^2_{0,3}(\alpha \otimes \beta)$.

As mentioned above, the spaces $\Lambda^2_-(M)$ and $\Sigma^2_{0,i}(M)$ are all isomorphic as SU(2)-modules. In what follows we will need this identification explicitly, and to this end we define linear maps P_i , given on a simple 2-form $\alpha = \alpha_1 \wedge \alpha_2$ by

$$
P_i: \Lambda^2_-(M) \to \Sigma^2_{0,i}(M),
$$

$$
P_i(\alpha) := \alpha_1 \otimes (\alpha_2^{\sharp} \cup \omega_i) - \alpha_2 \otimes (\alpha_1^{\sharp} \cup \omega_i).
$$

The above maps are manifestly $SU(2)$ -equivariant, hence by Schur's lemma each P_i is either an isomorphism or the zero map. One easily checks on the Euclidean model that $P_1(dx_{12} - dx_{34}) = -dx_1^2 - dx_2^2 + dx_3^2 + dx_4^2$, and thus we deduce that P_1 is indeed an isomorphism; similarly for P_2 and P_3 .

Lemma 2.4. *Given* $\alpha \in \Lambda^2_{-}(M)$ *and* $\beta \in \Sigma^2_{0,i}(M)$ *, the following identities hold:*

- *1.* $P_i(\alpha)(\cdot, \cdot) = \alpha(J_i \cdot, \cdot).$
- 2. $P_i(\alpha) \diamond \omega_i = -\delta_{ij}\alpha$.
- 3. $P_i(\beta \diamond \omega_i) = -\beta$.

Proof. Identity 1. follows from the fact that $X \perp \omega_i = g(J_i X, \cdot) = (J_i X)^{\flat}$. The proofs of 2. and 3. are simple applications of Schur's lemma. Since P_i and ∞_i are both SU(2)-equivariant maps, we know that $P_i(\alpha) \circ \omega_j = \lambda \alpha$ for some constant λ , hence we can find λ in each case by checking on a simple example. hence we can find λ in each case by checking on a simple example.

Since the triple $(\omega_1, \omega_2, \omega_3)$ completely determines g, vol, J_1, J_2, J_3 , it follows that the evolutions of latter tensors are determined by $(2)-(4)$ $(2)-(4)$. Indeed, explicitly:

Corollary 2.5. *Under a variation of the* SU(2)*-structure given by [\(2\)](#page-1-4)-[\(4\)](#page-1-5), the underlying metric, volume form and almost complex structures evolve by:*

$$
\frac{d}{dt}g = f_0g + B_{0,1} + B_{0,2} + B_{0,3},\tag{14}
$$

$$
\frac{d}{dt}\text{vol} = 2f_0\text{vol},\tag{15}
$$

$$
\frac{d}{dt}(J_i) = f_k J_j - f_j J_k - B^i_{0,j} - B^i_{0,k},\tag{16}
$$

 \Box

where $(i, j, k) \sim (1, 2, 3)$ *and the endomorphisms* $B_{0,i}^j$ *are defined by* $B_{0,i}(J_j \cdot, \cdot) =: g(B_{0,i}^j \cdot, \cdot)$ *.*

Proof. As [\(14\)](#page-5-1) and [\(15\)](#page-5-2) are straightforward, we only prove [\(16\)](#page-5-0). It suffices to consider the case $(i, j, k) = (1, 2, 3)$, since the same argument holds for the other cyclic permutations of indices. Differentiating $g(J_1 \cdot, \cdot) = \omega_1(\cdot, \cdot)$ and using [\(2\)](#page-1-4), we get

$$
g(\frac{d}{dt}(J_1)\cdot, \cdot) = f_0\omega_1 + B_{0,1}\diamond \omega_1 + f_3\omega_2 - f_2\omega_3 - B(J_1\cdot, \cdot).
$$

Observing that

$$
B(J_1 \cdot, \cdot) = f_0 \omega_1 + \sum_{i=1}^3 B_{0,i}(J_1 \cdot, \cdot),
$$

we know moreover from Lemma [2.4](#page-5-3) that $(B_{0,1} \diamond \omega_1)(\cdot, \cdot) = B_{0,1}(J_1 \cdot, \cdot)$, and the result follows.

It is clear from [\(16\)](#page-5-0) that the almost complex structures are invariant under conformal rescalings, which correspond to the function f_0 . We also see that the variations f_i and $B_{0,i}$ leave the J_i unchanged.

Let ∇ denote the Levi-Civita connection of g. Given an arbitrary 1-form α , we can decompose the 2-tensor $\nabla \alpha$ in terms of its SU(2)-irreducible components as

$$
\nabla \alpha = -\frac{1}{4} (\delta \alpha) g + \sum_{i=1}^{3} S_{0,i}^{2} (\nabla \alpha) + \frac{1}{4} \sum_{i=1}^{3} g(d\alpha, \omega_{i}) \omega_{i} + \frac{1}{2} \pi^{2} (d\alpha), \tag{17}
$$

where $\delta \alpha = - * d * \alpha$ denotes the codifferential. Appealing to Lemma [2.4](#page-5-3) and using the definition of $\Sigma_{0,i}^2(M)$, we have the following identifications:

$$
S_{0,1}^2(\nabla\alpha)(J_1,\cdot) = S_{0,1}^2(\nabla\alpha) \diamond \omega_1 = S_{0,1}^2(\nabla\alpha)(J_3,J_2),\tag{18}
$$

$$
S_{0,1}^2(\nabla\alpha)(J_2,\cdot) = -P_3(S_{0,1}^2(\nabla\alpha)\diamond\omega_1) = S_{0,1}^2(\nabla\alpha)(J_3,J_1),\tag{19}
$$

$$
S_{0,1}^2(\nabla\alpha)(J_3,\cdot) = +P_2(S_{0,1}^2(\nabla\alpha)\diamond\omega_1) = -S_{0,1}^2(\nabla\alpha)(J_1,J_2). \tag{20}
$$

The first expression identifies $S_{0,1}^2(\nabla \alpha) \in \Sigma_{0,1}^2(M)$ with a 2-form in $\Lambda^2(M)$, the second with a symmetric 2-tensor in $\Sigma_{0,3}^2(M)$ and the third with a symmetric 2-tensor in $\Sigma_{0,2}^2(M)$; analogous statements HOLD for the other $S_{0,i}^2$, by cyclic permutation. For instance, from (18) we also deduce that

$$
S_{0,2}^{2}(\nabla \alpha)(J_{2},) = S_{0,2}^{2}(\nabla \alpha) \diamond \omega_{2} = S_{0,2}^{2}(\nabla \alpha)(J_{1}, J_{3}) \in \Lambda_{-}^{2}(M).
$$

The identifications [\(18\)](#page-5-4)-[\(20\)](#page-5-5) will be particularly useful in \S 3 and \S 4. Lastly we recall the following simple lemma, certainly known to most readers, that we shall use frequently in \S 5 and as such we also give a proof:

Lemma 2.6. *Given a vector field* X *and a 2-form* ω *on* M, we have that

$$
\mathcal{L}_X \omega = \nabla_X \omega + (\mathcal{L}_X g - dX^{\flat}) \diamond \omega.
$$
 (21)

Proof. Given vector fields $X, Y, Z \in \mathcal{X}(M)$, from the relation $\mathcal{L}_X(\omega(Y, Z)) = \nabla_X(\omega(Y, Z))$ we derive

$$
(\mathcal{L}_X\omega)(Y,Z)=(\nabla_X\omega)(Y,Z)+\omega(\nabla_YX,Z)+\omega(Y,\nabla_ZX).
$$

i.e. $(\mathcal{L}_X \omega)_{ij} = X^k \nabla_k \omega_{ij} + \nabla_i X^k \omega_{ki} + \nabla_j X^k \omega_{ik}$. On the other hand, we know that

$$
(\mathcal{L}_X g)_{ij} = \nabla_i X^k g_{kj} + \nabla_j X^k g_{ki}.
$$

Acting with the latter on a 2-form ω under \diamond , as defined in [§1.3,](#page-3-0) we get

$$
((\mathcal{L}_X g) \diamond \omega)_{ij} = \frac{1}{2} (\nabla_a X^k g_{ki} g^{ab} \omega_{bj} + \nabla_i X^b \omega_{bj} + \nabla_a X^k g_{kj} g^{ab} \omega_{ib} + \nabla_j X^b \omega_{ib}),
$$

and the result follows from the fact that $\nabla X^{\flat} = \frac{1}{2}(\mathcal{L}_X g + dX^{\flat}).$

One way to interpret the above formula, say when $\omega = \omega_1$, is that to highest order in X, the anti-self-dual part of $\mathcal{L}_X \omega_1$ and the ω_1 -component are determined by the deformation of the metric, i.e. by $\mathcal{L}_{X}g$, while the isometric deformation is determined by dX^{\flat} . Observe that the lower-order term $\nabla_X \omega_1$ in X lies in the linear span $\langle \omega_2, \omega_3 \rangle$. This interpretation of [\(21\)](#page-5-6) will be useful when considering DeTurck's trick to prove parabolicity, in [§5.](#page-24-0) Another immediate application of the above lemma is the following:

Corollary 2.7. *If* $X \in \mathcal{X}(M)$ *is a Killing vector field, then*

 $\mathcal{L}_X \omega_i \in \langle \omega_j, \omega_k \rangle$, *for* $(i, j, k) \sim (1, 2, 3)$.

Conversely, if $\mathcal{L}_X \omega_1 = \mathcal{L}_X \omega_2 = \mathcal{L}_X \omega_3$ *, then* X *is a Killing vector field.*

Having covered the algebraic preliminaries of $SU(2)$ -structures, we next consider their first-order invariants, namely, the intrinsic torsion.

2.2 Intrinsic torsion of $SU(2)$ -structures

We begin by considering the more general set up, as described in [\[Sal89,](#page-39-12) Chapter 2]. Given an H-structure on the manifold M^n , such that $H \subset SO(n)$, we have a natural inclusion $\mathfrak{h} \subset \mathfrak{so}(n) \cong \Lambda^2(M)$ and hence an orthogonal splitting $\Lambda^2(M) = \mathfrak{h} \oplus \mathfrak{h}^{\perp}$. The intrinsic torsion of the H-structure is determined by a tensor $T \in \Lambda^1(M) \otimes \mathfrak{h}^{\perp}$ which measures the failure of the holonomy algebra of the Levi-Civita connection ∇ to reduce from $\mathfrak{so}(n)$ to h. If H can be defined as the stabiliser of a (multi-)tensor ξ , then T can be identified with $\nabla \xi$.

In our setup, we have $H = \text{SU}(2)$, with $\mathfrak{h} = \mathfrak{su}(2)^-$ and $\mathfrak{h}^{\perp} = \mathfrak{su}(2)^+$, and the stabilised multi-tensor is $\xi = (\omega_1, \omega_2, \omega_3)$. Note that \mathfrak{h}^{\perp} is not in general a Lie subalgebra, this occurs in dimension 4 because $\text{SO}(4) \cong (\text{SU}(2) \times \text{SU}(2))/\mathbb{Z}_2$. Thus, the intrinsic torsion of an $SU(2)$ -structure is given by a tensor

$$
T \in \Lambda^1(M) \otimes \mathfrak{h}^\perp \cong \Lambda^1(M) \otimes \Lambda^2_+(M). \tag{22}
$$

To describe the latter explicitly, we first note that \circ induces an action of $\Lambda^2 \cong \mathfrak{so}(4)$ on itself. In particular:

Lemma 2.8. *The operator* \diamond *acts on* $\Lambda^2_+(M) = \langle \omega_1, \omega_2, \omega_3 \rangle$ *by*

$$
\omega_i \diamond \omega_j = -\omega_j \diamond \omega_i = \omega_k, \quad \text{for} \quad (i, j, k) \sim (1, 2, 3),
$$

and $\omega_i \diamond \omega_i = 0$.

The above lemma is not really surprising, since the operator \diamond corresponds (up to a constant factor) to the Lie bracket on $\mathfrak{su}(2)^+ \subset \Lambda^2$, and hence Lemma [2.8](#page-6-2) identifies $\langle \omega_1, \omega_2, \omega_3 \rangle$ with the algebra of imaginary quaternions $\langle i, j, k \rangle$. More generally, we have an action of $\mathfrak{gl}(4,\mathbb{R}) = \mathbb{R}^4 \otimes (\mathbb{R}^4)^*$ on tensors given by [\(5\)](#page-3-3). In local coordinates, given $A = A_j^i \partial_{x_i} \otimes dx_j$ and a 2-tensor $\alpha = \alpha_{ij} dx_i \otimes dx_j$, we have

$$
(A \diamond \alpha)_{ij} = \frac{1}{2} (A_i^b \alpha_{bj} + A_j^b \alpha_{ib}).
$$

Using the metric, we have the identification $\mathfrak{gl}(4,\mathbb{R}) \cong (\mathbb{R}^4)^* \otimes (\mathbb{R}^4)^* = \langle g \rangle \oplus \Sigma_0^2(M) \oplus \Lambda^2(M)$, and thus shall also view the \diamond operator as an action of 2-tensors, i.e for $A = A_{ij} dx_i \otimes dx_j$,

$$
(A \diamond \alpha)_{ij} = \frac{1}{2} (A_{ai} g^{ab} \alpha_{bj} + A_{aj} g^{ab} \alpha_{ib}).
$$

With this convention, notice that $g \diamond \omega_i = \omega_i$.

Remark 2.1. The infinitesimal action \diamond defined here differs from the one given in [\[FLMSE22\]](#page-39-1) by a factor of one half; furthermore, the latter authors view $\mathfrak{gl}(4,\mathbb{R})$ as $(\mathbb{R}^4)^* \otimes \mathbb{R}^4$, and as a result there is a $-1/2$ factor difference for the diamond action of 2-forms, and only a factor of $+1/2$ for action of symmetric 2-tensors. In local coordinates, $A_j^a g_{ai} = A_{ji}$ in the convention of [\[FLMSE22\]](#page-39-1), whereas here $A_j^a g_{ai} = A_{ij}$; this does not matter if A is a symmetric 2-tensor, but there is a sign difference for 2-forms. So the reader should bear these differences in mind if comparing results; fortunately, we shall not need to rely directly on formulae computed therein for this article.

Denoting the SU(2)-triple by the row vector $\omega = (\omega_1, \omega_2, \omega_3)$, from [\[LSE23,](#page-39-0) Lemma 1] we know that

$$
\nabla_X \omega = T(X) \diamond \omega, \quad \text{for} \quad X \in \mathcal{X}(M). \tag{23}
$$

Under the identification [\(22\)](#page-6-3), we can view $T(X)$ as a self-dual 2-form

$$
T = a_1 \otimes \omega_1 + a_2 \otimes \omega_2 + a_3 \otimes \omega_3, \tag{24}
$$

where a_1, a_2, a_3 are *torsion* 1*-forms*, which can be explicitly described as follows:

Proposition 2.9. *The torsion tensor* T *can be expressed as*

$$
T = \frac{1}{2} \big(g(\nabla \omega_2, \omega_3) \otimes \omega_1 + g(\nabla \omega_3, \omega_1) \otimes \omega_2 + g(\nabla \omega_1, \omega_2) \otimes \omega_3 \big).
$$
 (25)

Equivalently, the torsion forms [\(24\)](#page-7-1) *are given by*

$$
a_1 = -\frac{1}{2}(*d\omega_1 + J_2(*d\omega_3) - J_3(*d\omega_2)),
$$
\n(26)

$$
a_2 = -\frac{1}{2}(*d\omega_2 - J_1(*d\omega_3) + J_3(*d\omega_1)),
$$
\n(27)

$$
a_3 = -\frac{1}{2}(*d\omega_3 + J_1(*d\omega_2) - J_2(*d\omega_1)).
$$
\n(28)

In particular,

 $\nabla \omega_i = 0 \Leftrightarrow a_j = a_k = 0, \quad \text{for} \quad (i, j, k) \sim (1, 2, 3).$

Moreover, $T = 0 \Leftrightarrow \nabla \omega = 0 \Leftrightarrow d\omega = 0$.

Proof. Expanding [\(23\)](#page-7-2) with Lemma [2.8,](#page-6-2) we get

$$
\nabla_X \omega^t = (-a_2(X)\omega_3 + a_3(X)\omega_2, a_1(X)\omega_3 - a_3(X)\omega_1, -a_1(X)\omega_2 + a_2(X)\omega_1)^t
$$

=
$$
\begin{pmatrix} 0 & +a_3(X) & -a_2(X) \\ -a_3(X) & 0 & +a_1(X) \\ +a_2(X) & -a_1(X) & 0 \end{pmatrix} \begin{pmatrix} \omega_1 \\ \omega_2 \\ \omega_3 \end{pmatrix} .
$$
 (29)

It follows that

$$
a_i(X) = \frac{1}{2}g(\nabla_X \omega_j, \omega_k) = -\frac{1}{2}g(\nabla_X \omega_k, \omega_j), \text{ for } (i, j, k) \sim (1, 2, 3),
$$
 (30)

and this proves [\(25\)](#page-7-3).

From [\(29\)](#page-7-4), we also have

$$
d\omega_i = a_k \wedge \omega_j - a_j \wedge \omega_k. \tag{31}
$$

Using the above together with Lemma [2.3,](#page-4-1) a short calculation yields [\(26\)](#page-7-5)-[\(28\)](#page-7-6).

Among SU(2)-structures with non-vanishing torsion, one important subclass is that of *hyper-Hermitian* structures i.e when the almost complex structures J_1, J_2, J_3 are all integrable. In our setup, we can characterise this condition as follows:

Proposition 2.10. *For each* $(i, j, k) \sim (1, 2, 3)$, the almost complex structure J_i is integrable if, and only if,

$$
J_i(a_k) = a_j.
$$

In particular, $\{J_1, J_2, J_3\}$ *defines a hypercomplex structure if, and only if,* $a_2 = -J_3(a_1)$ *and* $a_3 = J_2(a_1)$ *.*

Proof. The almost complex structure J_1 is integrable if, and only if, $d(\omega_2 + i\omega_3) \in \Lambda^{3,0} \oplus \Lambda^{2,1}$, in terms of the J_1 -type decomposition. It is easy to see this is equivalent to $a_3 - ia_2$ being a (1,0)-form (J₁-type) i.e. $J_1(a_3) = a_2$. The same argument holds for J₂ and J₃. argument holds for J_2 and J_3 .

Next we want to relate the derivative of the intrinsic torsion with the Riemann curvature tensor. To this end we recall the alternative characterisation of the intrinsic torsion, found eg. in [\[Bry05,](#page-39-3) §4.2] :

$$
g(\nabla_X Y - \nabla_X^{\mathfrak{h}} Y, Z) = \frac{1}{2} T(X)(Y, Z)
$$
\n(32)

where ∇ denotes the Levi-Civita connection and $\nabla^{\mathfrak{h}}$ denotes the natural SU(2)-connection obtained by projecting ∇ onto $\mathfrak{h} = \mathfrak{su}(2)^-$. In particular, $\nabla^{\mathfrak{h}} \omega_i = 0$, for $i = 1, 2, 3$, and $T = 0$ precisely when $\nabla = \nabla^{\mathfrak{h}}$. The connection $\nabla^{\mathfrak{h}}$ is in general not torsion-free, and instead its torsion can be identified with T .

 \Box

Denoting the curvature forms of ∇ and $\nabla^{\mathfrak{h}}$ by $F_{\nabla} = \text{Rm}$ and $F_{\nabla^{\mathfrak{h}}}$, respectively, and writing $d_{\nabla^{\mathfrak{h}}}$ for the exterior covariant derivative, we have

$$
F_{\nabla} = F_{\nabla^{\mathfrak{h}}} + \frac{1}{2} d_{\nabla^{\mathfrak{h}}} T + \frac{1}{8} [T \wedge T], \tag{33}
$$

where T is viewed as an $\mathfrak{h}^{\perp} = \mathfrak{sp}(1)^+$ -valued 1-form and $[\cdot \wedge \cdot]$ corresponds to simultaneously wedging the differential forms and taking the Lie bracket, cf. $[DK90, §2.1.2]$ $[DK90, §2.1.2]$. Note that the relations [\(32\)](#page-7-7) and [\(33\)](#page-8-0) hold more generally for any Hstructure with $H \subset SO(n)$, and $\nabla^{\mathfrak{h}}$ is again understood to be the projection of the Levi-Civita connection onto $\mathfrak{h} \subset \mathfrak{so}(n)$ cf. [\[Bry05,](#page-39-3)[Sal89\]](#page-39-12).

A unique feature of our setup, due to the splitting $\mathfrak{so}(4) = \mathfrak{sp}(1)^+ \oplus \mathfrak{sp}(1)^-$, is that $\nabla^{\mathfrak{h}}$ induces the flat connection on $\Lambda^2_+(M)$, and hence $d_{\nabla^b}T = dT$. Moreover, just like the diffeomorphism-invariance of the Riemann curvature yields the Bianchi identity, cf. $[CK04, §2.2]$, the diffeomorphism-invariance of the intrinsic torsion of an H-structure also yields a Bianchi-type identity; see for instance $[Kar09,$ Theorem 4.2] for the G_2 case, $[DLSE24,$ Theorem 3.8] for the $Spin(7)$ case, [\[FSE24,](#page-39-15) Proposition 2.5] for the Sp(2)Sp(1) case and [\[FLMSE22,](#page-39-1) Corollary 1.38] for the general case. This identity expresses the $\Lambda^2(M) \otimes \mathfrak{h}^{\perp} \subset \Lambda^2(M) \otimes \Lambda^2(M)$ component of the Riemann curvature in terms of derivatives of the intrinsic torsion, and it is obtained by the endomorphism action of [\(33\)](#page-8-0) on the tensor ξ stabilised by H, see also [§4.2](#page-19-2) below.

Corollary 2.11 (Bianchi-type identity). *The diffeomorphism-invariance of the intrinsic torsion* T *is expressed by:*

$$
2F_{\nabla} \diamond \omega = ((da_1 + a_2 \wedge a_3) \otimes \omega_1 + (da_2 + a_3 \wedge a_1) \otimes \omega_2 + (da_3 + a_1 \wedge a_2) \otimes \omega_3) \diamond \omega.
$$

Proof. The claim follows directly from a simple computation using [\(33\)](#page-8-0), along with $F_{\nabla^b} \diamond \omega = 0$ and $d_{\nabla^b} T = dT$, as discussed above. discussed above.

Remark 2.2. Corollary [2.11](#page-8-1) implies that the $\mathfrak{su}(2)^\perp$ component of the curvature form F_∇ is determined, up to lower-order terms, by da_i . Based on this observation, we can already deduce that, to highest order, the Ricci and self-dual Weyl curvature tensors can be expressed in terms of these second-order quantities – we use on this in [§4](#page-19-0) to derive explicit formulae.

Next we characterise *harmonic* SU(2)-structures, in the sense of [\[LSE23\]](#page-39-0). We recall that an SU(2)-structure defined by the triple ω is called harmonic if

$$
\operatorname{div}(T) := g^{ij} \nabla_i T_{jk} = 0
$$

(this holds true for more general H-structures with $H \subset SO(n)$, cf. [\[LSE23,](#page-39-0) Lemma 2]). In our setup we can state the condition more explicitly as follows:

Proposition 2.12. *The divergence of the intrinsic torsion* T *can be expressed as*

$$
\operatorname{div}(T) = \sum_{i=1}^{3} \operatorname{div}(a_i) \omega_i.
$$
 (34)

Equivalently, in terms of ∇ *and the triple* ω *alone,*

$$
\operatorname{div}(T) = \frac{1}{2} \Big(\big(g(\nabla^* \nabla \omega_2, \omega_3) + g(\nabla \omega_2, \nabla \omega_3) \big) \omega_1 + \big(g(\nabla^* \nabla \omega_3, \omega_1) + g(\nabla \omega_3, \nabla \omega_1) \big) \omega_2 + (g(\nabla^* \nabla \omega_1, \omega_2) + g(\nabla \omega_1, \nabla \omega_2) \big) \omega_3 \Big),
$$

 $where \nabla^* \nabla \omega_i := \text{tr}(\nabla^2 \omega_i)$. Thus, $\text{div}(T) = 0$ if, and only if, the 1*-forms* a_i are all coclosed.

Proof. Let $\{e_i\}$ denote a local SU(2)-frame, then using [\(25\)](#page-7-3) we compute

$$
2\text{div}(T) = g(\nabla^*\nabla\omega_1, \omega_2)\omega_3 + g(\nabla^*\nabla\omega_3, \omega_1)\omega_2 + g(\nabla^*\nabla\omega_2, \omega_3)\omega_1
$$

+
$$
\sum_{i=1}^4 g(\nabla_{e_i}\omega_1, \nabla_{e_i}\omega_2)\omega_3 + g(\nabla_{e_i}\omega_3, \nabla_{e_i}\omega_1)\omega_2 + g(\nabla_{e_i}\omega_2, \nabla_{e_i}\omega_3)\omega_1
$$

+
$$
\sum_{i=1}^4 g(\nabla_{e_i}\omega_1, \omega_2)\nabla_{e_i}\omega_3 + g(\nabla_{e_i}\omega_3, \omega_1)\nabla_{e_i}\omega_2 + g(\nabla_{e_i}\omega_2, \omega_3)\nabla_{e_i}\omega_1
$$

=
$$
g(\nabla^*\nabla\omega_1, \omega_2)\omega_3 + g(\nabla^*\nabla\omega_3, \omega_1)\omega_2 + g(\nabla^*\nabla\omega_2, \omega_3)\omega_1
$$

-
$$
2g(a_1, a_2)\omega_3 - 2g(a_3, a_1)\omega_2 - 2g(a_2, a_3)\omega_1
$$

=
$$
(g(\nabla^*\nabla\omega_1, \omega_2) + g(\nabla\omega_1, \nabla\omega_2))\omega_3
$$

+
$$
(g(\nabla^*\nabla\omega_2, \omega_3) + g(\nabla\omega_2, \nabla\omega_3))\omega_1
$$

+
$$
(g(\nabla^*\nabla\omega_3, \omega_1) + g(\nabla\omega_3, \nabla\omega_1))\omega_2.
$$

The equivalent expression [\(34\)](#page-8-2) can be deduced easily from the proof of Proposition [3.5](#page-12-0) below.

 \Box

In $[Gri17]$ Grigorian showed that harmonic G_2 -structures can be interpreted as G_2 -structures in 'octonionic Coulomb gauge', where the gauge group corresponds to the octonions O and hence is non-associative. This interpretation is also valid in the SU(2) case, see the discussion in [§6.2,](#page-37-0) but since the quaternions $\mathbb H$ are associative we instead get the usual notion of Coulomb gauge:

Corollary 2.13. The SU(2)-structure determined by $\langle \omega_1, \omega_2, \omega_3 \rangle$ is harmonic if, and only if, the induced Levi-Civita connec*tion* ∇ *on* $\Lambda^2_+(M)$ *is in Coulomb gauge for the frame* $\langle \omega_1, \omega_2, \omega_3 \rangle \subset \Lambda^2_+(M)$ *.*

Proof. The $\mathfrak{so}(3)$ matrix of a 1-form $\mathbf{a}^t = -\mathbf{a}$, as defined in [\(29\)](#page-7-4), simply corresponds to the induced Levi-Civita connection on Λ^2_+ , viewed here as a matrix of 1-forms with respect to the frame $\langle \omega_1, \omega_2, \omega_3 \rangle$. Thus, the harmonic condition $\text{div}(T) = 0$ is equivalent to $\delta \mathbf{a} = 0$.

Remark 2.3. From the general theory of harmonic structures [\[LSE23\]](#page-39-0), we already know that harmonic SU(2)-structures are critical points of the Dirichlet energy functional

$$
E(\omega) := \int_M |T(\omega)|^2 \text{vol} = 2 \int_M |\mathbf{a}|^2 \text{vol},\tag{35}
$$

whereby we only consider isometric variations of the SU(2)-structure i.e. q is kept fixed while the triple ω are allowed to vary as g-compatible 2-forms. Thus, Corollary [2.13](#page-9-2) asserts that Coulomb gauge minimises (or rather is the critical gauge for) the L^2 -norm of the connection 1-form a; this was already suggested in [\[DK90,](#page-39-13) p. 56] motivated by the theory of Abelian connections. In [§3](#page-9-0) we shall consider more general functionals, for which the metric is also allowed to vary.

Since throughout this article we shall express everything in terms of the torsion 1-forms a_i , we conclude this section by identifying several well-known classes of $SU(2)$ -structures in terms of intrinsic torsion:

Definition 2.14. In the above framework, $(M^4, \omega_1, \omega_2, \omega_3)$ is

- 1. hyperKähler, if $a_1 = a_2 = a_3 = 0$.
- 2. Kähler with respect to ω_i , if $a_j = a_k = 0$, for $(i, j, k) \sim (1, 2, 3)$.
- 3. Hermitian with respect to J_i , if $J_j(a_j) = J_k(a_k)$, for $(i, j, k) \sim (1, 2, 3)$.
- 4. hyperHermitian, if $J_1a_1 = J_2a_2 = J_3a_3$.
- 5. a harmonic SU(2)-structure, if $\delta(a_1) = \delta(a_2) = \delta(a_3) = 0$.
- 6. a harmonic U(2)-structure with respect to (J_i, ω_i) , if $\delta(a_j) = g(a_k, a_i)$ and $\delta(a_k) = -g(a_j, a_i)$, for $(i, j, k) \sim (1, 2, 3)$.

Note that the definition of harmonic $U(n)$ -structures in the literature is usually formulated in terms of the almost complex structure J, namely as $[\nabla^*\nabla J, J] = 0$, where $\nabla^*\nabla J := \text{tr}(\nabla^2 J)$ cf. [\[Woo93,](#page-39-16) [Woo95\]](#page-39-17) and also compare with Proposition [2.12](#page-8-3) above. In Definition [2.14](#page-9-3) (6) above we have instead expressed the harmonic condition in terms of the torsion 1-forms.

3 Variations and Functionals

We will now compute the evolution of the intrinsic torsion under [\(2\)](#page-1-4)-[\(4\)](#page-1-5), thence deriving the first variational formulae of all the quadratic functionals in the intrinsic torsion. The corresponding gradient flow provides natural candidates of $SU(2)$ -flows, among which we will have interest primary interest in the negative gradient flow of the Dirichlet energy [\(44\)](#page-11-1). Throughout this section we shall use the Einstein summation conventions i.e. we sum over repeated indices. We should mention that some of the computations here have already appeared in $[FLMSE22]$ but owing to the differences in conventions, see [§1.3](#page-3-0) and Remark [2.1,](#page-6-1) we find it safer to redo the computations.

3.1 Variation of intrinsic torsion

In this section we compute the evolution equation for the intrinsic torsion when the $SU(2)$ $SU(2)$ -structure evolves under (2)-[\(4\)](#page-1-5).

Lemma 3.1. *Under the variation of the* SU(2)*-structure given by [\(2\)](#page-1-4)-[\(4\)](#page-1-5),*

1. the Christoffel symbols Γ_{ij}^k *evolve by*

$$
\frac{d}{dt}(\Gamma_{ij}^k) = \frac{1}{2}g^{lk}(\nabla_j B_{il} + \nabla_i B_{jl} - \nabla_l B_{ij});
$$
\n(36)

2. the covariant derivative of an arbitrary 2*-tensor* ξ *evolves by*

$$
\frac{d}{dt}(\nabla_l \xi_{ij}) = \nabla_l \left(\frac{d}{dt}\xi_{ij}\right) - \frac{1}{2}\xi_{bj}g^{ab}(\nabla_i B_{la} + \nabla_l B_{ia} - \nabla_a B_{li}) - \frac{1}{2}\xi_{ib}g^{ab}(\nabla_j B_{la} + \nabla_l B_{ja} - \nabla_a B_{lj}).\tag{37}
$$

In particular, $\boldsymbol{\omega} = (\omega_1, \omega_2, \omega_3)$ *evolves by*

$$
\frac{d}{dt}(\nabla_l \boldsymbol{\omega}) = A \diamond \nabla_l \boldsymbol{\omega} + (\nabla_l C + \Lambda \nabla B_l) \diamond \boldsymbol{\omega},\tag{38}
$$

where $\Lambda \nabla B_l$ is the 2-form defined by $(\Lambda \nabla B_l)_{ij} := \nabla_i B_{lj} - \nabla_j B_{li}$.

Proof. The evolution of the Christoffel symbols is a standard result, cf. [\[Top06,](#page-39-18) Prop 2.3.1] or [\[CK04,](#page-39-14) Lemma 3.2]. Since $\nabla_l \xi_{ij} = \partial_l(\xi_{ij}) - \xi_{bj} \Gamma_{li}^b - \xi_{ib} \Gamma_{lj}^b$, we can easily deduce [\(37\)](#page-9-4) from [\(36\)](#page-9-5). So it only remains to prove [\(38\)](#page-10-0).

First, since $\frac{d}{dt}\omega = A \diamond \omega$ we have

$$
\nabla_l(\frac{d}{dt}\omega) = A \diamond \nabla_l \omega + \nabla_l A \diamond \omega
$$

= $A \diamond \nabla_l \omega + \nabla_l B \diamond \omega + \nabla_l C \diamond \omega$
= $A \diamond \nabla_l \omega + \frac{1}{2}g^{ab}(\nabla_l B_{ai}\omega_{bj} + \nabla_l B_{aj}\omega_{ib})dx_i \otimes dx_j + \nabla_l C \diamond \omega.$

Here we used the fact that the operator \diamond depends only on the metric and hence is preserved by ∇ . Setting $\xi = \omega$ in [\(37\)](#page-9-4) and using the above now yields

$$
\frac{d}{dt}(\nabla_l \omega) = \nabla_l (\frac{d}{dt} \omega) - \frac{1}{2} \omega_{bj} g^{ab} (\nabla_i B_{la} + \nabla_l B_{ai} - \nabla_a B_{li}) - \frac{1}{2} \omega_{ib} g^{ab} (\nabla_j B_{la} + \nabla_l B_{aj} - \nabla_a B_{lj})
$$
\n
$$
= A \diamond \nabla_l \omega + \nabla_l C \diamond \omega + \frac{1}{2} ((\nabla_a B_{li} - \nabla_i B_{la}) g^{ab} \omega_{bj} + (\nabla_a B_{lj} - \nabla_j B_{la}) g^{ab} \omega_{ib}) dx_i \otimes dx_j
$$

and this proves [\(38\)](#page-10-0).

Since the intrinsic torsion T is defined by (23) , we can now derive the evolution of T using (38) . In what follows it will also be important to distinguish the endomorphism-valued 1-form $T_{l,j}^i$ from the 3-tensor $T_{l,ij}$; to emphasise this subtlety, we shall denote the associated endomorphism as indeed a matrix by $[T_l],$ i.e

$$
[T_l]_j^i = T_{l,aj} g^{ai}
$$

.

and likewise we denote by $[P]$ the endomorphism associated to a 2-tensor P .

Proposition 3.2. Under the variation of the SU[\(2\)](#page-1-4)-structure given by (2)-[\(4\)](#page-1-5), the endomorphism $[T_l]$ evolves by

$$
\left(\frac{d}{dt}[T_l] - [\nabla_l C] - [\Lambda \nabla B_l] + \frac{1}{2}[[A],[T_l]]\right) \diamond \boldsymbol{\omega} = 0,\tag{39}
$$

where $[[A],[T_l]]^i_j=A^i_aT^a_{l,j}-T^i_{l,a}A^a_j$ denotes the usual Lie bracket on matrices. In particular,

$$
g\left(\frac{d}{dt}[T_l] - [\nabla_l C] - [\Lambda \nabla B_l], [T_l]\right) = 0,\tag{40}
$$

 \Box

where $[\nabla_l C]$ and $[\Lambda \nabla B_l]$ are viewed as endomorphisms.

Proof. Differentiating [\(23\)](#page-7-2),

$$
\frac{d}{dt}(\nabla\omega) = \frac{d}{dt}([T]) \diamond \omega + [T] \diamond \frac{d}{dt}\omega
$$

$$
= \frac{d}{dt}([T]) \diamond \omega + [T] \diamond (A \diamond \omega).
$$

Note that if we had considered T_l as a 2-tensor, instead of the endomorphism $[T_l]$, we would have had to differentiate \diamond as well, since it would depend on q . Using (38) , we can rewrite the above as

$$
\left(\frac{d}{dt}([T_l]) - [\nabla_l C] - [\Lambda \nabla B_l]\right) \diamond \omega + [T_l] \diamond (A \diamond \omega) - A \diamond ([T_l] \diamond \omega) = 0. \tag{41}
$$

Writing [A] for the endomorphism associated to the 2-tensor A, one can easily verify from the definition of \diamond that

$$
[T_l] \diamond (A \diamond \boldsymbol{\omega}) - A \diamond ([T_l] \diamond \boldsymbol{\omega}) = \frac{1}{2} [[A], [T_l]] \diamond \boldsymbol{\omega},
$$

which proves [\(39\)](#page-10-1). In particular, using the metric we see that $[[B],[T_1]]$ corresponds to a symmetric 2-tensor i.e. $[[B],[T_1]]_{ij} =$ $[[B],[T_l]]_{ji}$, while $[[C],[T_l]]$ corresponds to a 2-form. As a consequence,

$$
g([[A],[T_l]], [T_l]) = g([[C],[T_l]], [T_l])
$$

\n
$$
= [C]_k^a [T_l]_b^k [T_l]_j^i g_{ia} g^{bj} - [T_l]_k^a [C]_b^k [T_l]_j^i g_{ia} g^{bj}
$$

\n
$$
= [C]_k^a ([T_l]_b^k [T_l]_j^i g_{ia} g^{bj} - [T_l]_a^b [T_l]_j^i g_{ib} g^{kj})
$$

\n
$$
= [C]_k^a ([T_l]^{kj} [T_l])_{aj} - [T_l]_{ia} [T_l]^{ik})
$$

\n
$$
= 0
$$

Finally we derive the evolution of each irreducible component of the intrinsic torsion, namely the $\{a_i\}$.

Proposition 3.3. *Under the variation of the* SU[\(2\)](#page-1-4)*-structure given* (2)-[\(4\)](#page-1-5), *the torsion* 1*-forms* $\{a_i\}$ *evolve by*

$$
\frac{d}{dt}(a_i)_l = \nabla_l f_i + f_k(a_j)_l - f_j(a_k)_l + \frac{1}{2}g(\Lambda \nabla B_l, \omega_i), \quad \text{for} \quad (i, j, k) \sim (1, 2, 3), \tag{42}
$$

where $(a_i)_l := a_i(\partial_{x_l})$.

Proof. Differentiating $\nabla_l \omega_1 = (a_3)_l \omega_2 - (a_2)_l \omega_3$, we have

$$
\frac{d}{dt}(\nabla_l \omega_1) = \frac{d}{dt}((a_3)_l)\omega_2 - \frac{d}{dt}((a_2)_l)\omega_3 + (a_3)_l A \diamond \omega_2 - (a_2)_l A \diamond \omega_3.
$$

Now, from Lemma [3.1,](#page-9-6) we know that

$$
\frac{d}{dt}(\nabla_l \omega_1) = A \diamond \nabla_l \omega_1 + (\nabla_l C + \Lambda \nabla B_l) \diamond \omega_1.
$$

Combining both assertions we get

$$
\frac{d}{dt}((a_3)_l)\omega_2 - \frac{d}{dt}((a_2)_l)\omega_3 = (\nabla_l C + \Lambda \nabla B_l) \diamond \omega_1.
$$
\n(43)

Since $C = f_1 \omega_1 + f_2 \omega_2 + f_3 \omega_3$, we also have

$$
\nabla_l C = (\nabla_l f_1 - f_2(a_3)_l + f_3(a_2)_l)\omega_1 + (\nabla_l f_2 - f_3(a_1)_l + f_1(a_3)_l)\omega_2 + (\nabla_l f_3 - f_1(a_2)_l + f_2(a_1)_l)\omega_3,
$$

and, in view of Lemma [2.8,](#page-6-2) we get

$$
\frac{d}{dt}((a_3)_l)\omega_2 - \frac{d}{dt}((a_2)_l)\omega_3 = (\nabla_l f_3 - f_1(a_2)_l + f_2(a_1)_l)\omega_2 - (\nabla_l f_2 - f_3(a_1)_l + f_1(a_3)_l)\omega_3 + \Lambda \nabla B_l \diamond \omega_1.
$$

The fact that $\Lambda \nabla B_l \diamond \omega_1 = \frac{1}{2} g(\Lambda \nabla B_l, \omega_3) \omega_2 - \frac{1}{2} g(\Lambda \nabla B_l, \omega_2) \omega_3$ gives us the evolution of $(a_2)_l$ and $(a_3)_l$, and applying the same argument to $(a_1)_l$ yields the result. \Box

3.2 Energy of the intrinsic torsion

We are now in position to compute the first variation formula of the Dirichlet energy functional:

$$
E(\omega) := \int_M |T|^2 \text{vol} = 2 \int_M \sum_{i=1}^3 |a_i|^2 \text{vol.}
$$
 (44)

We should point out that (44) corresponds to a special choice among many other possible $SU(2)$ -functionals which are quadratic in the intrinsic torsion; these will be considered in [§3.3.](#page-14-0) Our particular interest stems from the fact that its negative gradient flow provides a natural tool to find torsion-free SU(2)-structures.

Proposition 3.4. *The first variation of the Dirichlet functional [\(44\)](#page-11-1) is given by*

$$
\frac{d}{dt}E(\omega) = -\int_M 2g(\text{div}(T), C)\text{vol} - \int_M g(\text{sym}(\text{div}(T^t)) + T \cdot T - \frac{1}{2}|T|^2 g, B)\text{vol}
$$
\n(45)

where

$$
\operatorname{div}(T)_{ij} := g^{ab} \nabla_a T_{b,ij}, \quad \operatorname{div}(T^t)_{ij} := g^{ab} \nabla_a T_{i,bj}, \quad \operatorname{sym}(A)_{ij} := A_{ij} + A_{ji},
$$

and

$$
(T * T)_{ab} := \frac{1}{2} g^{cd} g^{ef} T_{a,ce} T_{b,df}
$$

is a symmetric 2*-tensor.*

Proof. Recall our convention to take the inner product of 2-forms on the last two indices of T, so that

$$
|T|^2 = \frac{1}{2}g^{ab}g^{cd}g^{ef}T_{a,ce}T_{b,df}.
$$

With this in mind,

$$
\frac{d}{dt}|T|^2 = g^{ab}g^{cd}g^{ef}\frac{d}{dt}(T_{a,ce})T_{b,df} - \frac{1}{2}B^{ab}g^{cd}g^{ef}T_{a,ce}T_{b,df} - g^{ab}B^{cd}g^{ef}T_{a,ce}T_{b,df}
$$

$$
=g^{ab}g^{cd}g^{ef}(\frac{d}{dt}([T_a]_e^k)g_{ck}+[T_a]_e^kB_{ck})T_{b,df}-\frac{1}{2}B^{ab}g^{cd}g^{ef}T_{a,ce}T_{b,df}-g^{ab}B^{cd}g^{ef}T_{a,ce}T_{b,df}
$$

\n
$$
=g^{ab}g^{cd}g^{ef}\frac{d}{dt}([T_a]_e^k)g_{ck}T_{b,df}-\frac{1}{2}B^{ab}g^{cd}g^{ef}T_{a,ce}T_{b,df}-g^{ab}B^{cd}g^{ef}T_{a,ce}T_{b,df}+g^{ab}g^{cd}g^{ef}[T_a]_e^kB_{ck}T_{b,df}
$$

\n
$$
=g^{ab}g^{cd}g^{ef}\frac{d}{dt}([T_a]_e^k)g_{ck}T_{b,df}-\frac{1}{2}B^{ab}g^{cd}g^{ef}T_{a,ce}T_{b,df}
$$

where, for the last equality, we used

$$
g^{ab}g^{cd}g^{ef}[T_a]_e^k B_{ck}T_{b,df} = g^{ab}g^{cd}g^{ef}T_{a,ie}g^{ik}B_{ck}T_{b,df} = g^{ab}B^{di}g^{ef}T_{a,ie}T_{b,df}.
$$

Using [\(40\)](#page-10-2), and integrating by parts,

$$
\begin{split} \int_{M} (\frac{d}{dt}|T|^2) \mathrm{vol} &= \int_{M} (g^{ab}g^{cd}g^{ef} \frac{d}{dt} ([T_a]_e^k) g_{ck} T_{b,df} - \frac{1}{2} B^{ab}g^{cd}g^{ef} T_{a,ce} T_{b,df}) \mathrm{vol} \\ &= \int_{M} (g^{ab}g^{cd}g^{ef} (\nabla_a C_{ce} + (\Lambda \nabla B_a)_{ce}) T_{b,df} - \frac{1}{2} B^{ab}g^{cd}g^{ef} T_{a,ce} T_{b,df}) \mathrm{vol} \\ &= - \int_{M} 2g(\mathrm{div}(T), C) \mathrm{vol} + \int_{M} (g^{ab}g^{cd}g^{ef} (2\nabla_c B_{ae}) T_{b,df} - \frac{1}{2} B^{ab}g^{cd}g^{ef} T_{a,ce} T_{b,df}) \mathrm{vol} \\ &= - \int_{M} 2g(\mathrm{div}(T), C) \mathrm{vol} - \int_{M} (g^{ab}g^{cd}g^{ef} (2B_{ae}) \nabla_c T_{b,df} + \frac{1}{2} B^{ab}g^{cd}g^{ef} T_{a,ce} T_{b,df}) \mathrm{vol} \\ &= - \int_{M} 2g(\mathrm{div}(T), C) \mathrm{vol} - \int_{M} g(2\mathrm{div}(T^t) + T * T, B) \mathrm{vol}. \end{split}
$$

Note again that the factor '2' in front of the $div(T)$ term is due to our inner product convention on differential forms. The result now follows from the fact that $\frac{d}{dt}$ vol = $\frac{1}{2}g(g, B)$ vol, from [\(15\)](#page-5-2). \Box

Thus, the critical points of the energy functional $E(\omega)$ are given by:

$$
\operatorname{div}(T) = 0,\tag{46}
$$

$$
sym(\text{div}(T^t)) + T * T - \frac{1}{2}|T|^2 g = 0.
$$
\n(47)

We now describe these critical points in terms of the 1-forms $\{a_i\}$.

Proposition 3.5. *In terms of the torsion* 1*-forms* $\{a_i\}$ *, as defined by [\(24\)](#page-7-1),*

1. div(T) = $\sum_{i=1}$ div(a_i) ω_i = $-\sum_{i=1}$ $\delta(a_i)\omega_i$, 2. div(T^t) = $-\sum_{i,j} (\nabla_{e_i} a_j) \otimes J_j(e^i) - \sum_{i,j,k} \varepsilon_{ijk} a_i \otimes J_j a_k$, *3.* $T * T = 2(a_1 \otimes a_1 + a_2 \otimes a_2 + a_3 \otimes a_3)$, 4. $|T|^2 = 2(|a_1|^2 + |a_2|^2 + |a_3|^2)$.

Proof. As usual, let e_i (resp. e^i) denote a local SU(2) (co-)frame, so that $\omega_1 = e^{12} + e^{34}$, $\omega_2 = e^{13} - e^{24}$ and $\omega_3 = e^{14} + e^{23}$. Using [\(24\)](#page-7-1) and [\(29\)](#page-7-4), a direct computation yields

$$
\nabla_{e_i} T = \sum_j (\nabla_{e_i} a_j) \otimes \omega_j + a_1 \otimes (a_3(e_i)\omega_2 - a_2(e_i)\omega_3) + a_2 \otimes (-a_3(e_i)\omega_1 + a_1(e_i)\omega_3) + a_3 \otimes (a_2(e_i)\omega_1 - a_1(e_i)\omega_2).
$$

We deduce the first expression easily from the above since $\text{div}(T) = \sum_i e_i \cdot \nabla_{e_i} T$. On the other hand, using the identity $X \perp \omega_i = -J_i(X^{\flat}),$

$$
\operatorname{div}(T^t) = -\sum_{i,j} (\nabla_{e_i} a_j) \otimes J_j(e^i) + a_1 \otimes (-J_2 a_3 + J_3 a_2) + a_2 \otimes (J_1 a_3 - J_3 a_1)
$$

+ $a_3 \otimes (-J_1 a_2 + J_2 a_1)$
= $-\sum_{i,j} (\nabla_{e_i} a_j) \otimes J_j(e^i) - \sum_{i,j,k} \varepsilon_{ijk} a_i \otimes J_j a_k.$

Finally, the terms $T * T$ and $|T|^2$ are easily computed from [\(24\)](#page-7-1).

Note that the expressions in Proposition [3.5](#page-12-0) are not split into SU(2)-irreducible components. However, in the sequel it will be important to have an explicit description of such components of the highest-order term of sym $(\text{div}(T^t))$. To this end, we make the following observation:

 \Box

Remark 3.1. Since $\nabla_X a_i = (\nabla a_i)(X, \cdot) = \sum_{j=1}^4 e^j(X) \otimes \nabla_{e_j} a_i$, it follows that

$$
\nabla_{JX} a_i = \sum_{j=1}^4 J(e^j)(X) \otimes \nabla_{e_j} a_i.
$$

Hence, in view of Proposition [3.5–](#page-12-0)[\(2\)](#page-12-1), up to a factor of -1 , the highest-order term of [\(47\)](#page-12-2) is

$$
\mathrm{sym}(\sum_{i,j}(\nabla_{e_i}a_j)\otimes J_j(e^i))=\sum_{i,j}(\nabla_{e_i}a_j)\odot J_j(e^i)=\sum_{j=1}^3\mathrm{sym}(\nabla_{J_j}a_j),
$$

where sym $(\nabla_{J_i} a_j)(X, Y) := (\nabla_{J_i X} a_j)(Y) + (\nabla_{J_i Y} a_j)(X)$. Using [\(17\)](#page-5-7) and [\(18\)](#page-5-4)-[\(20\)](#page-5-5), it follows that

$$
sym(\nabla_{J_1} a_1) = -\frac{1}{2}g(da_1, \omega_1)g + P_1(\pi_-^2(da_1)) + 2P_3(S_{0,2}^2(\nabla a_1) \diamond \omega_2) - 2P_2(S_{0,3}^2(\nabla a_1) \diamond \omega_3),\tag{48}
$$

and analogous expressions can be deduced for sym($\nabla_{J_2} a_2$) and sym($\nabla_{J_2} a_2$), by permuting indices. This shows that sym($\nabla_{J_1} a_1$) does not depend on the second-order terms δa_1 , $g(da_1, \omega_2)$, $g(da_1, \omega_3)$ and $S_{0,1}^2(\nabla a_1)$. Note that such a formula was to be expected, since $\text{sym}(\nabla_{J_1}a_1)$ is determined by the SU(2)-invariant components of ∇a ; but we have $\text{sym}(\nabla_{J_1}a_1) \in S^2(M) \cong \mathbb{R} \oplus 3\mathbb{R}^3$, while $\nabla a_1 \in T^* \otimes T^* \cong 4\mathbb{R} \oplus 4\mathbb{R}^3$, so the former term had to be independent of the terms generating a copy of $3\mathbb{R} \oplus \mathbb{R}^3$ which is indeed consistent with the above expression (see also [§4.2](#page-19-2) below).

Note that we can also express $T * T$ in terms of its SU(2)-irreducible components as

$$
T * T = \frac{1}{2} \sum_{i} |a_i|^2 g + S_{0,1}^2 (T * T) + S_{0,2}^2 (T * T) + S_{0,3}^2 (T * T). \tag{49}
$$

Together with Proposition [3.5](#page-12-0) it follows that the 1-dimensional component of (47) is given by

$$
g(\operatorname{sym}(\operatorname{div}(T^t)), g) = 2\sum_{i} |a_i|^2. \tag{50}
$$

Applying the above to the critical condition [\(47\)](#page-12-2), we deduce:

Corollary 3.6. If ω is a critical point of $E(\omega)$ and $\text{div}(T^t)$ is traceless, then ω is an absolute minimum.

Observe that, since $SU(2) \subset U(2)$, we are implicitly also considering the almost Hermitian case. In fact, the tensors

$$
T_i := a_j \otimes \omega_j + a_k \otimes \omega_k \in \Lambda^1 \otimes (\langle \omega_i \rangle \oplus \Lambda^2_-)^{\perp}, \quad \text{for} \quad (i, j, k) \sim (1, 2, 3), \tag{51}
$$

correspond to the intrinsic torsion of the U(2)-structure defined by (g, ω_i, J_i) i.e.

$$
T_i \diamond \omega_i = \nabla \omega_i. \tag{52}
$$

Hence it is natural to consider the 'partial' energy functionals defined by

$$
E_i(\boldsymbol{\omega}) := \int_M |T_i|^2 \text{vol}, \quad \text{for} \quad i = 1, 2, 3,
$$
 (53)

each corresponding to the Dirichlet energy of the intrinsic torsion of the associated $U(2)$ -structure. In particular, it is not hard to see that the computations in the proofs of Proposition [3.2](#page-10-3) and [3.4](#page-11-2) also apply to each torsion tensor T_i , and we immediately deduce the following:

Proposition 3.7. *The first variation of [\(53\)](#page-13-0) is given by*

$$
\frac{d}{dt}E_i(\omega) = -\int_M g(\text{div}(T_i), C)\text{vol} - \int_M g(\text{sym}(\text{div}(T_i^t)) + T_i * T_i - \frac{1}{2}|T_i|^2 g, B)\text{vol.}
$$
\n(54)

It is also worth pointing out that

$$
E(\boldsymbol{\omega}) = \frac{1}{2} \Big(E_1(\boldsymbol{\omega}) + E_2(\boldsymbol{\omega}) + E_3(\boldsymbol{\omega}) \Big), \tag{55}
$$

and moreover this holds for any $Sp(n)$ -structure, not just $Sp(1) = SU(2)$. Next we consider more general $SU(2)$ functionals which are quadratic in the torsion.

3.3 General quadratic functionals of the intrinsic torsion

In the previous section we considered energy functionals involving the intrinsic torsion and computed their first variation formulae. The negative gradient flows of these functionals provide natural candidates for flows of $U(2)$ - and $SU(2)$ -structures, since the absolute minima correspond precisely to torsion-free structures. In our context, however, there are also other interesting functionals that one can consider; for instance, the functional $\int_M |J_1a_2+a_3|^2$ vol can be identified with the L^2 -norm of the Nijenhuis tensor of the almost complex structure J_1 , see Definition [2.14–](#page-9-3)(3), and hence its negative gradient flow provides a natural way to deform J_1 (optimistically) towards a complex structure. Thus, we are led to consider the first variation of functionals of the form

$$
\int_M g(a_i, a_j) \text{vol} \quad \text{and} \quad \int_M g(a_i, J_j(a_k)) \text{vol}.
$$

From [\(66\)](#page-21-2) below, it follows that there are in total 15 such distinct functionals. These can be viewed as building blocks of general functionals of $SU(2)$ -structures which are quadratic in the intrinsic torsion T. Note that it suffices to consider the 5 functionals

$$
\int_M g(a_1, a_2) \text{vol}, \quad \int_M g(a_1, a_1) \text{vol}, \quad \int_M g(a_1, J_3 a_2) \text{vol}, \quad \int_M g(a_1, J_1 a_2) \text{vol}, \quad \int_M g(a_1, J_1 a_3) \text{vol},
$$

since the remaining 10 terms can be obtained by cyclically permuting indices. Next we compute their first-variation formulae.

Proposition 3.8. *Under the variation of the* SU(2)*-structure given by [\(2\)](#page-1-4)-[\(4\)](#page-1-5),*

$$
\frac{d}{dt} \int_{M} g(a_1, a_2) \text{vol} = \frac{1}{2} \int_{M} g(B, \nabla_{J_1} a_2 + \nabla_{J_2} a_1 + g(a_1, a_2)g - 2a_1 \otimes a_2 + a_1 \otimes J_3 a_1 - a_1 \otimes J_1 a_3
$$

+ $a_2 \otimes J_2 a_3 - a_2 \otimes J_3 a_2$)vol

$$
-\frac{1}{2} \int_{M} g(C, (\text{div}(a_2) - g(a_1, a_3))\omega_1 + (\text{div}(a_1) + g(a_2, a_3))\omega_2 + (|a_1|^2 - |a_2|^2)\omega_3)
$$
vol. (56)

Proof. Using $\partial_t g^{ij} = -B^{ij}$ and $\partial_t \text{vol} = 2f_0 \text{vol}$,

$$
\frac{d}{dt} \int_M g(a_1, a_2) \text{vol} = \frac{d}{dt} \int_M g^{ij}(a_1)_i (a_2)_j \text{vol}
$$
\n
$$
= \int_M \left(-B^{ij}(a_1)_i (a_2)_j + 2f_0 g(a_1, a_2) + g^{ij} \frac{d}{dt} (a_1)_i (a_2)_j + g^{ij} (a_1)_i \frac{d}{dt} (a_2)_j \right) \text{vol}
$$

Using the evolution equations for the torsion forms in Proposition [3.3,](#page-11-3) and integrating by parts, we can rewrite the above as

$$
\frac{d}{dt} \int_{M} g(a_1, a_2) \text{vol} = \int_{M} \left(-B^{ij}(a_1)_i (a_2)_j + 2f_0 g(a_1, a_2) - g^{ij} \nabla_i (a_2)_j f_1 + g^{ij} f_3(a_2)_i (a_2)_j \right. \\
\left. - g^{ij} f_2(a_2)_j (a_3)_i - g^{ij} f_2 \nabla_j (a_1)_i + g^{ij} f_1(a_1)_i (a_3)_j - g^{ij} f_3(a_1)_i (a_1)_j \right. \\
\left. + \frac{1}{4} g^{ij} g^{ab} g^{cd} (a_2)_j (\omega_1)_{bd} (\nabla_a B_{ic} - \nabla_c B_{ia}) + \frac{1}{4} g^{ij} g^{ab} g^{cd} (a_1)_i (\omega_2)_{bd} (\nabla_a B_{jc} - \nabla_c B_{ja}) \right) \text{vol}.
$$

Recall from Proposition [2.9](#page-7-0) that $\nabla \omega_1 = -a_2 \otimes \omega_3 + a_3 \otimes \omega_2$, and similarly for $\nabla \omega_2$, $\nabla \omega_3$; on the other hand, for general 1-form α , we have $(J_i\alpha)_a = -(\omega_i)_{k\alpha} g^{bk}\alpha_b$. Using these facts, and integrating by parts again, we get

$$
\frac{d}{dt} \int_{M} g(a_1, a_2) \text{vol} = \int_{M} \left(-B^{ij}(a_1)_i (a_2)_j + 2f_{0}g(a_1, a_2) - g^{ij} \nabla_i(a_2)_j f_1 + g^{ij} f_3(a_2)_i (a_2)_j \right. \\
\left. - g^{ij} f_2(a_2)_j (a_3)_i - g^{ij} f_2 \nabla_j(a_1)_i + g^{ij} f_1(a_1)_i (a_3)_j - g^{ij} f_3(a_1)_i (a_1)_j \right. \\
\left. - \frac{1}{2} g^{ij} g^{ab} g^{cd} B_{ic}(-a_2)_a (\omega_3)_{bd}(a_2)_j + (a_3)_a (\omega_2)_{bd}(a_2)_j + (\omega_1)_{bd} \nabla_a(a_2)_j \right) \\
\left. - \frac{1}{2} g^{ij} g^{ab} g^{cd} B_{jc}(-a_3)_a (\omega_1)_{bd}(a_1)_i + (a_1)_a (\omega_3)_{bd}(a_1)_i + (\omega_2)_{bd} \nabla_a(a_1)_i \right) \right) \text{vol}, \\
= \int_{M} \left(-g(B, a_1 \otimes a_2) + 2f_{0}g(a_1, a_2) - f_1 \operatorname{div}(a_2) - f_2 \operatorname{div}(a_1) + f_3 |a_2|^2 - f_3 |a_1|^2 \right. \\
\left. - f_2 g(a_2, a_3) + f_1 g(a_1, a_3) - \frac{1}{2} g^{ij} g^{ab} g^{cd} B_{ic}(\omega_1)_{bd} \nabla_a(a_2)_j - \frac{1}{2} g^{ij} g^{ab} g^{cd} B_{jc}(\omega_2)_{bd} \nabla_a(a_1)_i \right. \\
\left. + \frac{1}{2} g(B, a_1 \otimes J_3 a_1 - a_1 \otimes J_1 a_3 + a_2 \otimes J_2 a_3 - a_2 \otimes J_3 a_2) \right) \text{vol}.
$$

Observe that the highest-order symmetric part in [\(56\)](#page-14-1) is given by the symmetric components of $\nabla a_2(J_1, \cdot) + \nabla a_1(J_2, \cdot)$, while the highest-order skew-symmetric part is given by $\text{div}(a_2)\omega_1 + \text{div}(a_1)\omega_2$. While the latter term lies in an irreducible $SU(2)$ -module, the former terms can be further decomposed as follows. Using (17) and $(18)-(20)$ $(18)-(20)$ $(18)-(20)$ we can express these as

$$
\nabla a_2(J_1,) = -\frac{1}{4}g(da_2,\omega_1)g + \frac{1}{2}P_1(\pi_-^2(da_2)) - P_2(S_{0,3}^2(\nabla a_2)\diamond\omega_3) + P_3(S_{0,2}^2(\nabla a_2)\diamond\omega_2)
$$

$$
+\frac{1}{4}(-(\delta a_2)\omega_1+g(da_2,\omega_3)\omega_2-g(da_2,\omega_2)\omega_3)
$$

and

$$
\nabla a_1(J_2,) = -\frac{1}{4}g(da_1, \omega_2)g + \frac{1}{2}P_2(\pi^2_-(da_1)) - P_3(S^2_{0,1}(\nabla a_1) \diamond \omega_1) + P_1(S^2_{0,3}(\nabla a_1) \diamond \omega_3) + \frac{1}{4}(-(\delta a_1)\omega_2 + g(da_1, \omega_1)\omega_3 - g(da_1, \omega_3)\omega_1).
$$

The key point here is that we are considering the inner product with the symmetric endomorphism B , so we may disregard the skew-symmetric terms in the above two expressions. By an analogous computation, one can show:

Proposition 3.9. *Under the variation of the* SU(2)*-structure given by [\(2\)](#page-1-4)-[\(4\)](#page-1-5),*

$$
\frac{d}{dt} \int_M g(a_1, a_1) \text{vol} = \int g(B, \nabla_{J_1} a_1 + \frac{1}{2}|a_1|^2 g - a_1 \otimes a_1 - a_1 \otimes J_3 a_2 + a_1 \otimes J_2 a_3) \text{vol} \n- \int g(C, \text{div}(a_1)\omega_1 + g(a_1, a_3)\omega_2 - g(a_1, a_2)\omega_3) \text{vol}.
$$

From Proposition [3.9](#page-15-0) we can also deduce the analogous expressions when a_1 is replaced by a_2 and a_3 , simply by permuting indices. In particular, the reader can easily verify that the first variation of the Dirichlet functional, as given in Proposition [3.4,](#page-11-2) is just a consequence of the latter, in view of [\(44\)](#page-11-1). More generally, we can deduce the critical points of the weighted energy functionals

$$
E_{\lambda}(\omega) := 2 \int_{M} \sum_{i=1}^{3} \lambda_i |a_i|^2 \text{vol}, \qquad (57)
$$

where $\lambda = (\lambda_1, \lambda_2, \lambda_3) \in \mathbb{R}^3$; the cases $\lambda = (1, 1, 0), (1, 0, 1), (0, 1, 1)$ correspond to the U(2) energy functionals [\(53\)](#page-13-0), and $\lambda = (1, 1, 1)$ to the Dirichlet energy [\(44\)](#page-11-1).

Proposition 3.10. *Under the variation of the* SU(2)*-structure given by [\(2\)](#page-1-4)-[\(4\)](#page-1-5),*

$$
\frac{d}{dt} \int_{M} g(a_1, J_3 a_2) \text{vol} = \int \frac{1}{2} g(B, (\nabla a_2)(J_1, J_3) - (\nabla a_1)(J_2, J_3)) \text{vol} \n+ \int \frac{1}{2} g(B, J_3 a_1 \otimes J_1 a_3 - J_3 a_1 \otimes J_3 a_1 + J_2 a_1 \otimes J_2 a_1 - J_2 a_2 \otimes J_1 a_1) \text{vol} \n+ \int \frac{1}{2} g(B, J_2 a_3 \otimes J_3 a_2 - J_3 a_2 \otimes J_3 a_2 + J_1 a_2 \otimes J_1 a_2 - J_1 a_1 \otimes J_2 a_2) \text{vol} \n+ \int g(B, a_2 \otimes J_3 a_1 - a_1 \otimes J_3 a_2 + \frac{3}{4} g(a_1, J_3 a_2) g + S_{0,3}^2 (J_3 a_2 \otimes a_1)) \text{vol} \n- \int \frac{1}{2} g(C, (g(da_2, \omega_3) - g(a_1, J_3 a_3)) \omega_1 - (g(da_1, \omega_3) + g(a_3, J_3 a_2)) \omega_2) \text{vol}.
$$
\n(S8)

Proof. From the outset, we have

$$
\frac{d}{dt} \int_{M} g(a_1, J_3 a_2) \text{vol} = -\frac{d}{dt} \int_{M} g^{ij}(a_1)_i (\omega_3)_{kj} g^{bk}(a_2)_b \text{vol}
$$
\n
$$
= -\int_{M} g(B, a_1 \otimes J_3 a_2) \text{vol} + \int_{M} g(a_1, J_3(\frac{d}{dt} a_2)) \text{vol} + \int_{M} g(\frac{d}{dt} a_1, J_3 a_2) \text{vol}
$$
\n
$$
+ \int_{M} 2f_0 g(a_1, J_3 a_2) \text{vol} + \int_{M} g^{ij} B^{bk}(a_1)_i (\omega_3)_{kj}(a_2)_b \text{vol}
$$
\n
$$
+ \int_{M} g^{ij} g^{bk}(a_1)_i \frac{d}{dt} (\omega_3)_{kj}(a_2)_b \text{vol}.
$$

Using [\(4\)](#page-1-5) and Proposition [3.3,](#page-11-3) we further have

$$
\frac{d}{dt} \int_{M} g(a_1, J_3 a_2) \text{vol} = -\int_{M} g(B, a_1 \otimes J_3 a_2) \text{vol} + \int_{M} 2f_0 g(a_1, J_3 a_2) \text{vol} \n- \int_{M} g^{ij} g^{bk} (\nabla_i f_1 + f_3(a_2)_i - f_2(a_3)_i + \frac{1}{2} g(\Lambda \nabla B_i, \omega_1)) (\omega_3)_{kj}(a_2)_b \text{vol} \n- \int_{M} g^{ij} g^{bk}(a_1)_i (\omega_3)_{kj} (\nabla_b f_2 + f_1(a_3)_b - f_3(a_1)_b + \frac{1}{2} g(\Lambda \nabla B_b, \omega_2)) \text{vol} \n+ \int_{M} g^{ij} B^{bk}(a_1)_i (\omega_3)_{kj}(a_2)_b \text{vol}
$$

$$
-\int_{M} g^{ij}g^{bk}(a_{1})_{i}(f_{0}(\omega_{3})_{kj} + f_{2}(\omega_{1})_{kj} - f_{1}(\omega_{2})_{kj} + (B_{0,3} \circ \omega_{3})_{kj})(a_{2})_{b}vol= -\int_{M} g(B, a_{1} \otimes J_{3}a_{2})vol + \int_{M} 2f_{0}g(a_{1}, J_{3}a_{2})vol+ \int_{M} g^{ij}g^{bk}f_{1}(\nabla_{i}(\omega_{3})_{kj}(a_{2})_{b} + (\omega_{3})_{kj}\nabla_{i}(a_{2})_{b})vol- \int_{M} g^{ij}g^{bk}(f_{3}(a_{2})_{i} - f_{2}(a_{3})_{i})(\omega_{3})_{kj}(a_{2})_{b}vol- \int_{M} \frac{1}{2}g^{ij}g^{bk}g^{\alpha\beta}g^{cd}\nabla_{\alpha}B_{ic}(\omega_{1})_{\beta d}(\omega_{3})_{kj}(a_{2})_{b}vol+ \int_{M} g^{ij}g^{bk}(\nabla_{b}(a_{1})_{i}(\omega_{3})_{kj} + (a_{1})_{i}\nabla_{b}(\omega_{3})_{kj})f_{2}vol- \int_{M} g^{ij}g^{bk}(a_{1})_{i}(\omega_{3})_{kj}(f_{1}(a_{3})_{b} - f_{3}(a_{1})_{b})vol- \int_{M} \frac{1}{2}g^{ij}g^{bk}(a_{1})_{i}(\omega_{3})_{kj}g^{\alpha\beta}g^{cd}\nabla_{\alpha}B_{bc}(\omega_{2})_{\beta d}vol+ \int_{M} g^{ij}B^{bk}(a_{1})_{i}(\omega_{3})_{kj}(a_{2})_{b}vol- \int_{M} g^{ij}g^{bk}(a_{1})_{i}(f_{0}(\omega_{3})_{kj} + f_{2}(\omega_{1})_{kj} - f_{1}(\omega_{2})_{kj} + (B_{0,3} \circ \omega_{3})_{kj})(a_{2})_{b}vol.
$$

Since $(J_i\alpha)_a = -(\omega_i)_{ka} g^{bk}\alpha_b$ for any 1-form α , we then use Proposition [2.9](#page-7-0) to compute $\nabla \omega_i$ and integrate by parts:

$$
\begin{split} \frac{d}{dt} \int_{M} g(a_{1}, J_{3}a_{2}) \mathrm{vol} & = -\int_{M} g(B, a_{1} \otimes J_{3}a_{2}) \mathrm{vol} + \int_{M} 2f_{0}g(a_{1}, J_{3}a_{2}) \mathrm{vol} + \int_{M} g^{ij}g^{bk}f_{1}(\omega_{3})_{kj} \nabla_{i}(a_{2})_{b} \mathrm{vol} \\ & + \int_{M} g^{ij}g^{bk}f_{1}(\nabla_{i}(\omega_{3})_{kj}(a_{2})_{b}) \mathrm{vol} \\ & = \int_{M} g^{ij}g^{bk}(f_{3}(a_{2})_{i} - f_{2}(a_{3})_{i})(\omega_{3})_{kj}(a_{2})_{b} \mathrm{vol} \\ & + \int_{M} \frac{1}{2}g^{ij}g^{bk}g^{3}g^{cd}B_{ic} \nabla_{\alpha}((\omega_{1})_{\beta}g(\omega_{3})_{kj}(a_{2})_{b}) \mathrm{vol} \\ & + \int_{M} g^{ij}g^{bk}(\nabla_{b}(a_{1})_{i}(\omega_{3})_{kj} + (a_{1})_{i} \nabla_{b}(\omega_{3})_{kj} f_{2} \mathrm{vol} \\ & - \int_{M} g^{ij}g^{bk}(a_{1})_{i}(\omega_{3})_{kj} (f_{1}(a_{3})_{b} - f_{3}(a_{1})_{b}) \mathrm{vol} \\ & + \int_{M} \frac{1}{2}g^{ij}g^{bk}(a_{1})_{i}(\omega_{3})_{kj}(a_{2})_{b} \mathrm{vol} \\ & + \int_{M} g^{ij}B^{bk}(a_{1})_{i}(\omega_{3})_{kj}(a_{2})_{b} \mathrm{vol} \\ & = - \int_{M} g^{ij}g^{bk}(a_{1})_{i} (f_{0}(\omega_{3})_{kj} + f_{2}(\omega_{1})_{kj} - f_{1}(\omega_{2})_{kj} + (B_{0,3} \circ \omega_{3})_{kj})(a_{2})_{b} \mathrm{vol} \\ & = - \int_{M} g^{ij}g^{bk}(a_{1})_{i} (f_{0}(\omega_{3})_{kj} + f_{2}(\omega_{1})_{kj} - f_{1}(\omega_{2})_{kj} + (B_{0,3} \circ \omega_{3})_{kj})(a_{2})_{b} \mathrm{vol} \\ & + \int_{M} g^{ij}g^{bk}(a_{1})_{i} (\omega_{2})_{
$$

+
$$
\int_M \frac{1}{2} g^{ij} g^{bk} g^{\alpha\beta} g^{cd} B_{bc}(a_1)_i (\nabla_\alpha(\omega_3)_{kj}(\omega_2)_{\beta d} + (\omega_3)_{kj} \nabla_\alpha(\omega_2)_{\beta d}) \text{vol}
$$

+
$$
\int_M g^{ij} B^{bk}(a_1)_i (\omega_3)_{kj}(a_2)_b \text{vol}
$$

$$
- \int_M g^{ij} g^{bk}(a_1)_i (f_0(\omega_3)_{kj} + f_2(\omega_1)_{kj} - f_1(\omega_2)_{kj} + (B_{0,3} \diamond \omega_3)_{kj})(a_2)_b \text{vol}.
$$

The assertion follows by simplifying the above and applying $g(B_{0,3}, J_3a_2 \otimes a_1) = -g(B_{0,3} \otimes \omega_3, a_2 \otimes a_1)$ which follows from a simple computation using the results in §2.1. from a simple computation using the results in [§2.1.](#page-3-2)

The highest-order symmetric part in [\(58\)](#page-15-1) is given by the symmetric components of $(\nabla a_2)(J_1, J_3)$ and $(\nabla a_1)(J_2, J_3)$. Using (17) and $(18)-(20)$ $(18)-(20)$ $(18)-(20)$, we can express these as

$$
(\nabla a_2)(J_1, J_3) = -\frac{1}{4}g(da_2, \omega_2)g - P_1(S_{0,3}^2(\nabla a_2) \diamond \omega_3) - \frac{1}{2}P_2(\pi_-^2(da_2)) - P_3(S_{0,1}^2(\nabla a_2) \diamond \omega_1) + \frac{1}{4}(g(da_2, \omega_3)\omega_1 + (\delta a_2)\omega_2 + g(da_2, \omega_1)\omega_3) + S_{0,2}^2(\nabla a_2) \diamond \omega_2
$$

and

$$
(\nabla a_1)(J_2, J_3) = +\frac{1}{4}g(da_1, \omega_1)g + \frac{1}{2}P_1(\pi_-^2(da_1)) - P_2(S_{0,3}^2(\nabla a_1) \diamond \omega_3) - P_3(S_{0,2}^2(\nabla a_1) \diamond \omega_2) + \frac{1}{4}(-(\delta a_1)\omega_1 + g(da_1, \omega_3)\omega_2 + g(da_1, \omega_2)\omega_3) - S_{0,1}^2(\nabla a_1) \diamond \omega_1,
$$

where as before the skew-symmetric terms do not play a role in the variational formula.

Proposition 3.11. *Under the variation of the* SU(2)*-structure given by [\(2\)](#page-1-4)-[\(4\)](#page-1-5),*

$$
\frac{d}{dt} \int_{M} g(a_1, J_1 a_2) \text{vol} = \int_{M} \frac{1}{2} g(B, (\nabla a_2)(J_1, J_1) - (\nabla a_1)(J_2, J_1)) \text{vol} \n+ \int_{M} \frac{1}{2} g(B, J_1 a_2 \otimes J_2 a_3 + J_2 a_2 \otimes J_1 a_3 - J_1 a_2 \otimes J_3 a_2 - J_3 a_2 \otimes J_1 a_2) \text{vol} \n+ \int_{M} \frac{1}{2} g(B, J_3 a_1 \otimes J_2 a_2 + J_1 a_1 \otimes J_1 a_3 - J_1 a_1 \otimes J_3 a_1 - J_2 a_1 \otimes J_2 a_3) \text{vol} \n+ \int_{M} g(B, a_2 \otimes J_1 a_1 - a_1 \otimes J_1 a_2 + \frac{1}{2} g(a_1, J_1 a_2) g - \frac{1}{4} g(a_2, J_1 a_1) g) \text{vol} \n- \int_{M} g(B, S_{0,1}^2 (J_1 a_1 \otimes a_2)) \text{vol} \n+ \int_{M} \frac{1}{2} g(C, -g(da_2, \omega_1)\omega_1 + g(da_1, \omega_1)\omega_2) \text{vol} \n+ \int_{M} \frac{1}{2} g(C, (g(a_2, J_2 a_3) - g(a_3, J_1 a_1))\omega_1 + (g(a_2, J_3 a_1) + g(a_3, J_2 a_1) - g(a_3, J_1 a_2) - g(a_2, J_3 a_1)) \omega_2 - g(a_2, J_2 a_1) \omega_3) \text{vol}.
$$
\n(59)

Proof. By direct computation,

$$
\frac{d}{dt} \int_{M} g(a_1, J_1 a_2) \text{vol} = -\frac{d}{dt} \int_{M} g^{ij}(a_1)_i (\omega_1)_{kj} g^{bk}(a_2)_b \text{vol}
$$
\n
$$
= -\int_{M} g(B, a_1 \otimes J_1 a_2) \text{vol} + \int_{M} g(a_1, J_1(\frac{d}{dt} a_2)) \text{vol} + \int_{M} g(\frac{d}{dt} a_1, J_1 a_2) \text{vol} + \int_{M} 2f_0 g(a_1, J_1 a_2) \text{vol} + \int_{M} g^{ij} B^{bk}(a_1)_i (\omega_1)_{kj}(a_2)_b \text{vol}
$$
\n
$$
+ \int_{M} g^{ij} g^{bk}(a_1)_i \frac{d}{dt} (\omega_1)_{kj}(a_2)_b \text{vol}.
$$

A long computation as before shows

$$
\frac{d}{dt} \int_{M} g(a_1, J_1 a_2) \text{vol} = -\int_{M} g(B, a_1 \otimes J_1 a_2) \text{vol} + \int_{M} 2f_0 g(a_1, J_1 a_2) \text{vol} + \int_{M} g^{ij} g^{bk} f_1(\omega_1)_{kj} \nabla_i(a_2)_b \text{vol} \n+ \int_{M} g^{ij} g^{bk} f_1 \nabla_i(\omega_1)_{kj}(a_2)_b \text{vol} \n- \int_{M} g^{ij} g^{bk} (f_3(a_2)_i - f_2(a_3)_i)(\omega_1)_{kj}(a_2)_b \text{vol}
$$

+
$$
\int_M \frac{1}{2} g^{ij} g^{bk} g^{\alpha\beta} g^{cd} B_{ic} \nabla_{\alpha} ((\omega_1)_{\beta d} (\omega_1)_{kj} (a_2)_b)
$$
vol
+ $\int_M g^{ij} g^{bk} (\nabla_b (a_1)_i (\omega_1)_{kj} + (a_1)_i \nabla_b (\omega_1)_{kj}) f_2$ vol
- $\int_M g^{ij} g^{bk} (a_1)_i (\omega_1)_{kj} (f_1(a_3)_b - f_3(a_1)_b)$ vol
+ $\int_M \frac{1}{2} g^{ij} g^{bk} g^{\alpha\beta} g^{cd} B_{bc} \nabla_{\alpha} ((a_1)_i (\omega_1)_{kj} (\omega_2)_{\beta d})$ vol
+ $\int_M g^{ij} B^{bk} (a_1)_i (\omega_1)_{kj} (a_2)_b$ vol
- $\int_M g^{ij} g^{bk} (a_1)_i (f_0(\omega_1)_{kj} + f_3(\omega_2)_{kj} - f_2(\omega_3)_{kj} + (B_{0,1} \diamond \omega_1)_{kj}) (a_2)_b$ vol

 \Box

and the result follows by simplifying the above.

Note that the highest-order symmetric terms in [\(59\)](#page-17-0) can be expressed as:

$$
\int_M g(B,(\nabla a_2)(J_1,J_1))\text{vol} = \int_M g(B, -\frac{1}{4}(\delta a_2)g + S_{0,1}^2(\nabla a_2) - S_{0,2}^2(\nabla a_2) - S_{0,3}^2(\nabla a_2))\text{vol}
$$

and

$$
\int_M g(B,(\nabla a_1)(J_2,J_1))\text{vol} = -\int_M g(B,\frac{1}{4}g(da_1,\omega_3)g + P_1(S_{0,2}^2(\nabla a_1)\diamond\omega_2) + P_2(S_{0,1}^2(\nabla a_1)\diamond\omega_1) + \frac{1}{2}P_3(\pi_-(da_1)))\text{vol}.
$$

Finally, by similar computations, we have:

Proposition 3.12. *Under the variation of the* SU(2)*-structure given by [\(2\)](#page-1-4)-[\(4\)](#page-1-5),*

$$
\frac{d}{dt} \int_{M} g(a_{1}, J_{1}a_{3}) \text{vol} = \int_{M} \frac{1}{2} g(B, (\nabla a_{3})(J_{1}, J_{1}) - (\nabla a_{1})(J_{3}, J_{1})) \text{vol} \n+ \int_{M} \frac{1}{2} g(B, J_{1}a_{3} \otimes J_{2}a_{3} + J_{3}a_{1} \otimes J_{3}a_{2} - J_{1}a_{3} \otimes J_{3}a_{2} - J_{3}a_{3} \otimes J_{1}a_{2}) \text{vol} \n+ \int_{M} \frac{1}{2} g(B, J_{1}a_{1} \otimes J_{2}a_{1} + J_{1}a_{3} \otimes J_{2}a_{3} - J_{1}a_{1} \otimes J_{1}a_{2} - J_{2}a_{1} \otimes J_{3}a_{3}) \text{vol} \n+ \int_{M} g(B, a_{3} \otimes J_{1}a_{1} - a_{1} \otimes J_{1}a_{3} + \frac{1}{2} g(a_{1}, J_{1}a_{3})g + \frac{1}{4} g(a_{1}, J_{1}a_{3})g) \text{vol} \n- \int_{M} g(B, S_{0,1}^{2}(J_{1}a_{1} \otimes a_{3})) \text{vol} \n+ \int_{M} \frac{1}{2} g(C, -g(da_{3}, \omega_{1})\omega_{1} + g(da_{1}, \omega_{1})\omega_{3}) \text{vol} \n+ \int_{M} \frac{1}{2} g(C, (g(a_{2}, J_{1}a_{1}) - g(a_{3}, J_{3}a_{2}))\omega_{1} + (g(a_{2}, J_{1}a_{3}) + g(a_{3}, J_{2}a_{1}) + g(a_{1}, J_{2}a_{3}) - g(a_{2}, J_{3}a_{1}))\omega_{3} - g(a_{1}, J_{3}a_{3})\omega_{2}) \text{vol}.
$$
\n(60)

Note that the highest order symmetric terms in [\(60\)](#page-18-0) can be expressed as:

$$
\int_M g(B,(\nabla a_3)(J_1,J_1))\text{vol} = \int_M g(B, -\frac{1}{4}(\delta a_3)g + S_{0,1}^2(\nabla a_3) - S_{0,2}^2(\nabla a_3) - S_{0,3}^2(\nabla a_3))\text{vol}
$$

and

$$
\int_M g(B, (\nabla a_1)(J_3, J_1)) \text{vol} = \int_M g(B, \frac{1}{4}g(da_1, \omega_2)g - P_1(S_{0,3}^2(\nabla a_1) \diamond \omega_3) + \frac{1}{2}P_2(\pi_+^2(da_1)) - P_3(S_{0,1}^2(\nabla a_1) \diamond \omega_1)) \text{vol}.
$$

Having computed the first variation formulae for all the quadratic functionals in the intrinsic torsion we next classify the second-order invariants of SU(2)-structures.

4 Second-order invariants

It is a classical fact that all the second-order differential invariants of a Riemannian manifold, i.e. tensors defined by second derivatives of the metric, are determined by the Riemann curvature tensor. This is however not the case for general Hstructures with $H \subset SO(n)$, see for instance [\[Bry05\]](#page-39-3) for a detailed description of the G_2 case. Furthermore, the first-order invariants given by the components of the intrinsic torsion determine part of the curvature tensor, see eg. [§2.2;](#page-6-0) this is to say that there are relations between ∇T and Rm, see Corollary [2.11.](#page-8-1) In the case $H = G_2$ considered in [\[DGK23\]](#page-39-2), the authors derive these relations by direct computations from the G_2 Bianchi identity, then use these to extract a basis of second-order invariants, in a process which is computationally quite involved. There is however an alternative approach using representation theory due to Bryant [ibid.], which effectively allows us to deduce the number of second-order invariants by purely algebraic methods, then it is just a matter of choosing a basis of such invariants. In this section, we give an explicit description of this phenomenon in the $H = SU(2)$ case. Later on, in [§6.1,](#page-35-1) we systematise the corresponding results in the SU(3), G_2 and Spin(7) cases.

4.1 The Riemann curvature tensor

On an oriented Riemannian manifold $(M⁴, g)$, the curvature tensor Rm decomposes into 4 independent components: the scalar curvature s, the traceless Ricci tensor Ric₀, and the self-dual and anti-self-dual parts of the Weyl curvature tensor W^{\pm} . More concretely, we can describe these irreducible SO(4)-modules as follows:

$$
\operatorname{Rm} = \operatorname{Scal} + \operatorname{Ric}_0 + W^+ + W^- \in \mathbb{R} \oplus \Lambda^2_+ \otimes \Lambda^2_- \oplus S_0^2(\Lambda^2_+) \oplus S_0^2(\Lambda^2_-) \cong \ker(S^2(\Lambda^2) \xrightarrow{\wedge} \Lambda^4).
$$

If M admits an SU(2)-structure, say determined by the triple $\{\omega_1, \omega_2, \omega_3\}$, then we can further decompose Rm as follows:

- 1. The scalar curvature Scal is 1-dimensional, so it clearly does not further split.
- 2. Ric₀ splits into 3 components, according to the splitting of symmetric tensors $\Sigma_0^2 = \Sigma_{0,1}^2 \oplus \Sigma_{0,2}^2 \oplus \Sigma_{0,3}^2$.
- 3. W^+ can be expressed as a sum of the 5 terms

$$
2\omega_1 \otimes \omega_1 - \omega_2 \otimes \omega_2 - \omega_3 \otimes \omega_3, \quad \omega_2 \otimes \omega_2 - \omega_3 \otimes \omega_3, \quad \omega_1 \odot \omega_2, \quad \omega_1 \odot \omega_3 \quad \text{and} \quad \omega_2 \odot \omega_3,
$$

with suitable coefficient functions. In other words, the latter 5 terms provide an orthogonal basis for $S_0^2(\Lambda_+^2)$.

4. The anti-self-dual Weyl tensor W^- remains irreducible as a section of $S_0^2(\Lambda^2_-)$.

As already seen in Corollary [2.11,](#page-8-1) the derivatives of a_i already determine part of the curvature tensor. Furthermore, from the above decomposition, it is not hard to see that these include all the curvature components aside from W^- , since the latter lies in the kernel of ∞ . Before deriving explicit expressions for the Ricci curvature and W^+ in terms of a_i , we shall next describe all the second-order invariants according to Bryant's perspective.

4.2 Some representation theory

Following $\left[\text{Bry05}\right]$, we describe the space of first- and second-order invariants of an SU(2)-structure abstractly as irreducible SU(2)-modules. We shall give an explicit basis of these second-order invariants in Theorem [4.2.](#page-20-1)

We begin by considering the case of general H-structures, with $H \subset SO(n)$. In [\[Bry05,](#page-39-3) §4.2], all the k-th order invariants of an H-structure are obtained as sections of the associated bundle to the H-module $V_k(\mathfrak{h})$, which is implicitly defined by

$$
V_k(\mathfrak{h}) \oplus (\Lambda^1 \otimes S^{k+1}(\Lambda^1)) \cong (S^2(\Lambda^1) \oplus \mathfrak{h}^\perp) \otimes S^k(\Lambda^1). \tag{61}
$$

In this article we are only be concerned with $k = 1, 2$. Since Bryant does not include the proof of [\(61\)](#page-19-3) therein, we sketch the proof for $k = 2$; the case $k = 1$ is just the space where intrinsic torsion lies, which is already well-known, cf. [\[Bry87,](#page-39-19) [Sal89\]](#page-39-12).

Proof (sketch). Recall from the beginning of $\S2.2$ that the intrinsic torsion T of a H-structure is given by

$$
\nabla \xi = T \diamond \xi,
$$

where ξ is the (multi-)tensor defining the H-structure. It follows, by differentiation, that

$$
\nabla^2 \xi = \nabla T \diamond \xi + T \diamond (T \diamond \xi).
$$

Hence, skew-symmetrising on the first two indices and using [\(32\)](#page-7-7), we have

$$
F_{\nabla} \diamond \xi = alt(\nabla^{\mathfrak{h}} T) \diamond \xi + \text{l.o.t}(\xi).
$$

Since $\nabla^{\mathfrak{h}}T \in \Lambda^1 \otimes \Lambda^1 \otimes \mathfrak{h}^{\perp}$, it follows that $\text{alt}(\nabla^{\mathfrak{h}}T) \diamond \xi \in \Lambda^2 \otimes \mathfrak{h}^{\perp}$. The second-order invariants are obtained from ∇T and Rm, but these overlap precisely in the space $\Lambda^2 \otimes \mathfrak{h}^{\perp}$, so $V_2(\mathfrak{h})$ is given by $S^2(\Lambda^1) \otimes \mathfrak{h}^{\perp}$ (consisting of those invariants not coming from the curvature) together with the space of curvature tensors ($S^2(\Lambda^1) \otimes S^2(\Lambda^1)$ modulo the terms in $\Lambda^1 \otimes S^3(\Lambda^1)$), cf. [\[BBG83,](#page-39-20) §5]. This concludes the proof. For the cases $k > 2$, the proof is similar but one must also use the differential Bianchi identity. \Box

We now specialise to the case $H = SU(2)$. We already know from [§2.2](#page-6-0) that

$$
V_1(\mathfrak{su}(2)) \cong 3\Lambda^1 \cong 3\mathbb{R}^4
$$

is precisely the space of intrinsic torsion. On the other hand $V_2(\mathfrak{su}(2))$ is implicitly defined by

$$
V_2(\mathfrak{su}(2)) \oplus (\Lambda^1 \otimes S^3(\Lambda^1)) \cong (S^2(\Lambda^1) \oplus 3\mathbb{R}) \otimes S^2(\Lambda^1). \tag{62}
$$

A straightforward calculation using the Clebsch–Gordan formula, cf. [\[Sal89\]](#page-39-12), shows that

$$
(S^2(\Lambda^1) \oplus 3\mathbb{R}) \otimes S^2(\Lambda^1) \cong 9\mathbb{R}^5 \oplus 24\mathbb{R}^3 \oplus 13\mathbb{R},
$$

$$
\Lambda^1 \otimes S^3(\Lambda^1) \cong 8\mathbb{R}^5 \oplus 12\mathbb{R}^3 \oplus 4\mathbb{R}.
$$

Since $\mathbb{R}^3 \cong \mathfrak{su}(2)$ and $\mathbb{R}^5 \cong S_0^2(\mathbb{R}^3)$, as SU(2)-representations, we deduce that

$$
V_2(\mathfrak{su}(2)) \cong \mathbb{R}^5 \oplus 12\mathbb{R}^3 \oplus 9\mathbb{R}.\tag{63}
$$

On the other hand, as an $SO(4)$ -module, we have

$$
V_2(\mathfrak{so}(4)) \cong \mathbb{R}^5_+ \oplus \mathbb{R}^5_- \oplus (\mathbb{R}^3_+ \otimes \mathbb{R}^3_-) \oplus \mathbb{R},
$$

which is of course just the space of the curvature tensor for an oriented Riemannian 4-manifold. Refining the latter as an $SU(2)$ -module gives

$$
V_2(\mathfrak{so}(4)) \cong \mathbb{R}^5 \oplus 3\mathbb{R}^3 \oplus 6\mathbb{R}.\tag{64}
$$

The latter is exactly the decomposition given by (1)-(4) in [§4.1.](#page-19-1) Comparing we see that $V_2(\mathfrak{su}(2))$ has an extra $9\mathbb{R}^3 \oplus 3\mathbb{R}$ modules, which correspond precisely to the second-order invariants which do not arise from curvature terms. In the next section we shall find an explicit generator for each component of $V_2(\mathfrak{su}(2))$.

4.3 Second-order invariants explicitly

Recall that $T \in \Lambda^1 \otimes \Lambda^2_+ \cong 3\Lambda^1$, since $\Lambda^2_+ = \langle \omega_1, \omega_2, \omega_3 \rangle$, and hence T is identified with the triple of 1-forms (a_1, a_2, a_3) . It follows that ∇T is determined by ∇a_i :

$$
\nabla a_i \in 4\mathbb{R} \oplus \Lambda^2_- \oplus \Sigma^2_{0,1} \oplus \Sigma^2_{0,2} \oplus \Sigma^2_{0,3},\tag{65}
$$

where for each i the four copies of R are determined by δa_i and $g(da_i, \omega_j)$, for $j = 1, 2, 3$; the Λ^2 -component is just the anti-self-dual part of da_i ; and the $S^2_{0,j}$ components are the type-decomposition of the symmetric part of ∇a_i . We should point out that these second-order terms are not all independent. We summarise their dependency relations in the next proposition.

Proposition 4.1. *The following identities hold:*

1.
$$
g(da_2, \omega_3) = g(da_3, \omega_2) + g(a_1, J_2(a_2)) + g(a_1, J_3(a_3))
$$

\n2. $g(da_3, \omega_1) = g(da_1, \omega_3) + g(a_2, J_3(a_3)) + g(a_2, J_1(a_1))$
\n3. $g(da_1, \omega_2) = g(da_2, \omega_1) + g(a_3, J_1(a_1)) + g(a_3, J_2(a_2))$

Proof. The result follows by taking the exterior derivative of the relations [\(31\)](#page-7-8) and simplifying. For instance, differentiating $d\omega_1 = -a_2 \wedge \omega_3 + a_3 \wedge \omega_2$ gives $0 = -da_2 \wedge \omega_3 + a_1 \wedge a_2 \wedge \omega_2 + da_3 \wedge \omega_2 + a_1 \wedge a_3 \wedge \omega_3$, which yields the first identity using Lemma 2.3, and likewise for the remaining ones. using Lemma [2.3,](#page-4-1) and likewise for the remaining ones.

This shows that the exterior derivatives of the torsion 1-forms a_i are not completely independent: the 9 second-order terms $g(da_i, \omega_j)$ are related by the above 3 relations, up to addition of lower-order terms which are quadratic in T. Recall from the representation theory above that there are 9 copies of $\mathbb R$ in $V_2(\mathfrak{su}(2))$. On the other hand, we also saw that there are 12 R-components which can be obtained from ∇a_i , namely, δa_i and $g(d\dot{a_i}, \omega_j)$. The above proposition shows that there are 3 relations between the latter terms and hence a total of $12 - 3 = 9$ independent R-terms arising from ∇a_i . Thus, all the \mathbb{R} -components of $V_2(\mathfrak{su}(2))$ are precisely given by them, and there are no further relations; this would not have been obvious, had one not computed the space $V_2(\mathfrak{su}(2))$ in [\(63\)](#page-20-2) beforehand. In the G₂ case considered in [\[DGK23\]](#page-39-2), the authors derive all the relations coming from ∇T and Rm, which is as impressive as it is laborious; our approach circumvents this effort. The conclusion of Proposition [4.1,](#page-20-3) (62) and (65) can be summarised into:

Theorem 4.2. *All second-order invariants of an* SU(2)*-structure can be expressed from these linearly independent terms:*

- *1. The* 9 functions $\delta a_1, \delta a_2, \delta a_3$ and $g(da_i, \omega_j)$, where $1 \leq i \leq j \leq 3$.
- 2. The 12 anti-self-dual forms $\pi_{-}^2(da_i)$, where $i = 1, 2, 3$, and $S_{0,j}^2(\nabla a_i) \diamond \omega_j$, where $i, j = 1, 2, 3$.
- *3. The anti-self-dual Weyl curvature* $W^- \in S_0^2(\Lambda_-^2)$ *.*

As we already saw in [§3.3,](#page-14-0) all the above second order operators (with the exception of W^-) arise naturally when considering the first variational formulae of $SU(2)$ functionals, see Propositions [3.8](#page-14-2) - [3.12.](#page-18-1)

Remark 4.1. As a point of comparison, the results in $\left[\text{Bry05}, \frac{84.3}{84.3}\right]$ show that, in the G₂ case, there are in fact 11 second-order invariants; 6 of which were described explicitly in [\[DGK23,](#page-39-2) §5], as these can be used to flow the underlying G_2 -structure. In [§6.1](#page-35-1) below, we provide a systematic method for describing the full generality of second-order flows for arbitrary H-structures.

Finally, we derive formulae to compute certain components of $d(J_i a_j)$ from those of $d a_i$, similar to those we found in Proposition [4.1.](#page-20-3)

Proposition 4.3. *Let* α *be an arbitrary* 1*-form on* M. Then the 12 *second-order terms given by* $g(d(J_i(\alpha)), \omega_i)$ *and* $\delta(J_i\alpha))$ *are in fact completely determined by the* 4 *second-order terms* $g(d\alpha, \omega_i)$ *and* $\delta\alpha$ *, up to addition of lower-order terms. More precisely,*

$$
\delta \alpha = g(d(J_1 \alpha), \omega_1) - g(\alpha, J_2 \alpha_2) - g(\alpha, J_3 \alpha_3)
$$

= $g(d(J_2 \alpha), \omega_2) - g(\alpha, J_3 \alpha_3) - g(\alpha, J_1 \alpha_1)$
= $g(d(J_3 \alpha), \omega_3) - g(\alpha, J_1 \alpha_1) - g(\alpha, J_2 \alpha_2)$

and

$$
g(d(J_i\alpha), \omega_j) = +g(d\alpha, \omega_k) - g(\alpha, J_i(a_j) + a_k),
$$

$$
g(d(J_i\alpha), \omega_k) = -g(d\alpha, \omega_j) - g(\alpha, J_i(a_k) - a_j),
$$

for $(i, j, k) \sim (1, 2, 3)$ *.*

Proof. From Lemma [2.3–](#page-4-1)(1), we know that $\alpha \wedge \omega_1 = -J_3(\alpha) \wedge \omega_2 = J_2(\alpha) \wedge \omega_3$. The claim follows by applying the exterior derivative to the latter relation. For instance,

$$
d(J_3(\alpha)) \wedge \omega_2 = -d\alpha \wedge \omega_1 - \alpha \wedge a_2 \wedge \omega_3 - a_1 \wedge J_3(\alpha) \wedge \omega_3,
$$

and hence, using Lemma [2.3,](#page-4-1)

$$
g(d(J_3\alpha), \omega_2) = -g(d\alpha, \omega_1) - g(\alpha, J_3a_2) + g(\alpha, a_1).
$$

Likewise, since

$$
(\delta \alpha) \text{vol} = d((J_1 \alpha) \wedge \omega_1) = d((J_2 \alpha) \wedge \omega_2) = d((J_3 \alpha) \wedge \omega_3),
$$

we have

$$
\delta \alpha = g(d(J_1\alpha), \omega_1) - g(\alpha, J_2\alpha_2) - g(\alpha, J_3\alpha_3).
$$

The other expressions are obtained analogously.

In the next section we express the Ricci curvature and the self-dual Weyl tensor in terms of the second-order invariants of Theorem [4.2](#page-20-1) and quadratic terms in the first-order invariants which lie in

$$
S^2(V_1(\mathfrak{su}(2))) \cong 15\mathbb{R} \oplus 21\mathbb{R}^3. \tag{66}
$$

Note that the 15 functions of [\(66\)](#page-21-2) correspond to $g(a_i, a_j)$ and $g(a_i, J_j a_k)$, whose first variation of the L^2 -norm we computed in [§3,](#page-9-0) and the \mathbb{R}^3 components are spanned by terms of the form $\pi^2 \cdot (J_i a_j \wedge a_k)$ and $S^2_{0,i} (J_j a_k \otimes a_l)$.

4.3.1 Ricci tensor in terms of intrinsic torsion

Proposition 4.4. *The traceless Ricci tensor of* g *can be computed explicitly in terms of the intrinsic torsion, by*

$$
Ric_0(g) = P_1(\pi^2_-(\Phi_1)) + P_2(\pi^2_-(\Phi_2)) + P_3(\pi^2_-(\Phi_3)) \in \Sigma^2_{0,1}(M) \oplus \Sigma^2_{0,2}(M) \oplus \Sigma^2_{0,3}(M),
$$

where

$$
\Phi_1 := da_1 + \frac{1}{4}(J_2(a_2) \wedge J_2(a_3) - J_1(a_1) \wedge J_2(a_3) + J_2(a_2) \wedge a_1)
$$

$$
-\frac{1}{4}(J_1(a_1) \wedge J_3(a_2) + J_3(a_3) \wedge a_1) - \frac{3}{4}(J_3(a_3) \wedge J_3(a_2)),
$$

and Φ_2 , Φ_3 *are obtained by cyclically permuting the indices of* Φ_1 *. The scalar curvature is given by*

$$
Scal(g) = -2\Big(\sum_{i=1}^{3} g(da_i, \omega_i) + g(J_1(a_1), J_2(a_2)) + g(J_1(a_1), J_3(a_3)) + g(J_2(a_2), J_3(a_3))\Big). \tag{67}
$$

 \Box

Proof. From Theorem [4.2](#page-20-1) we know that there are 12 possible second-order terms to compute Ric₀, whereas from Corollary [2.11](#page-8-1) we know that one can restrict to just the $\pi^2(a_i)$, so there are only 3 possible terms. On the other hand, the lower-order quadratic terms belong in [\(66\)](#page-21-2), of which there are 21 possibilities. It is now just a matter of identifying the coefficient constants for each of these terms, which can be done by considering suitable examples. \Box

Remark 4.2. The G_2 analogue of Proposition [4.4](#page-21-1) was derived in [\[Bry05,](#page-39-3) §4.5.3], and the Spin(7) analogue in [\[Fow20,](#page-39-21) Proposition 4.1].

To the best of our knowledge, the above expressions for the irreducible components of the Ricci tensor in terms of the SU(2) torsion forms do not seem to have appeared in the literature; except under further assumptions on the underlying structure, such as Hermitian or Kähler. In particular, we make the following observations:

- 1. *q* is an Einstein metric if, and only if, Φ_i are all self-dual 2-forms.
- 2. if ω_1 is a Kähler form, i.e. $a_2 = a_3 = 0$, then $\Phi_1 = -da_1$ and $\Phi_2 = \Phi_3 = 0$. Thus, $\text{Ric}(g) \in \langle g \rangle \oplus \Sigma_{0,1}^2$ i.e. it is of type $(1, 1)$, which is of course just the well-known fact that the closed 2-form Φ_1 is the Ricci form, up to a constant factor, cf. [\[KN63,](#page-39-22) Chapter IX. 4]. It is also easy to see from [\(16\)](#page-5-0) that the Ricci flow preserves the complex structure J_1 , since in this case $C = 0$ and $B = f_0 g + B_{0,1} = -2Ric \in \langle g \rangle \oplus \Sigma_{0,1}^2$.

The following result appears to be originally due to Vaisman [\[Vai82,](#page-39-23) Theorem 3.1], see also [\[Boy86\]](#page-39-24).

Corollary 4.5. *A compact hyper-Hermitian* 4*-manifold* M *has non-negative total scalar curvature. Moreover, the total scalar curvature is zero if, and only if, the hyper-Hermitian structure is hyperKähler.*

Proof. Using Proposition [4.3,](#page-21-3) we can equivalently express the scalar curvature [\(67\)](#page-21-4) as

Scal(g) =
$$
2\left(\sum_{i=1}^{3} \delta(J_i(a_i)) + g(J_1(a_1), J_2(a_2)) + g(J_1(a_1), J_3(a_3)) + g(J_2(a_2), J_3(a_3))\right)
$$
.

From Definition [2.14](#page-9-3) it follows that, for a hyper-Hermitian 4-manifold,

$$
Scal(g) = 6(\delta(J_1(a_1)) + |a_1|^2).
$$

Assuming $M⁴$ is compact, and integrating the latter expression, Stokes' theorem gives the result.

We conclude this section by relating the Ricci tensor to the symmetrisation of $div(T^t)$, which is the highest-order term in [\(47\)](#page-12-2). This will be important when we investigate parabolic flows of $SU(2)$ -structures in [§5.](#page-24-0)

Proposition 4.6. *Denoting by* T *the torsion of an* $SU(2)$ *-structure* ω *, the following holds*

$$
\text{sym}(\text{div}(T^t)) = -2\text{Ric}(g) - \sum_{i=1}^3 \mathcal{L}_{(J_i a_i)^{\sharp}} g + \text{l.o.t.}(\boldsymbol{\omega}),\tag{68}
$$

where the lower-order terms are quadratic in the torsion forms a_i .

Proof. From Proposition [4.4,](#page-21-1) it follows that

$$
Ric(g) = -\frac{1}{2} \sum_{i=1}^{3} g(da_i, \omega_i)g + P_1(\pi_-^2(da_1)) + P_2(\pi_-^2(da_2)) + P_3(\pi_-^2(da_3)) + \text{l.o.t.}(\omega).
$$
 (69)

On the other hand from Proposition [3.5](#page-12-0) and Remark [3.1](#page-13-1) we see that

$$
sym(\text{div}(T^t)) = +\frac{1}{2} \sum_{i=1}^3 g(da_i, \omega_i)g - P_1(\pi_-^2(da_1)) - P_2(\pi_-^2(da_2)) - P_3(\pi_-^2(da_3))
$$

$$
- 2P_3(S_{0,2}^2(\nabla a_1) \diamond \omega_2) + 2P_2(S_{0,3}^2(\nabla a_1) \diamond \omega_3)
$$

$$
- 2P_1(S_{0,3}^2(\nabla a_2) \diamond \omega_3) + 2P_3(S_{0,1}^2(\nabla a_2) \diamond \omega_1)
$$

$$
- 2P_2(S_{0,1}^2(\nabla a_3) \diamond \omega_1) + 2P_1(S_{0,2}^2(\nabla a_3) \diamond \omega_2) + \text{l.o.t.}(\boldsymbol{\omega}).
$$

Observe that, for any 1-form α and vector field X, we have

 $\nabla_X (J\alpha) = (\nabla_X J)\alpha + J(\nabla_X \alpha) = (\nabla_X J)\alpha + (\nabla \alpha)(X, J \cdot)$

i.e. $\nabla_X (J\alpha) = (\nabla \alpha)(X, J\cdot) + 1 \cdot \text{o.t.}(\alpha)$. Using [\(17\)](#page-5-7) and [\(18\)](#page-5-4)-[\(20\)](#page-5-5), we can express the latter in terms of its irreducible components as

$$
(\nabla \alpha)(\cdot, J_1 \cdot) = +\frac{1}{4}g(d\alpha, \omega_1)g - \frac{1}{2}P_1(\pi^2 - (d\alpha)) - P_2(S^2_{0,3}(\nabla \alpha) \diamond \omega_3) + P_3(S^2_{0,2}(\nabla \alpha) \diamond \omega_2)
$$

 \Box

$$
+\frac{1}{4}(\delta\alpha)\omega_1+\frac{1}{4}g(d\alpha,\omega_3)\omega_2-\frac{1}{4}g(d\alpha,\omega_2)\omega_3-S_{0,1}^2(\nabla\alpha)\diamond\omega_1,
$$

and similarly for $J = J_2, J_3$. Applying the above to compute $\sum_{i=1}^3 \mathcal{L}_{(J_ia_i)} \not\equiv \sum_{i=1}^3 \text{sym}(\nabla (J_ia_i))$, we then have

$$
\sum_{i=1}^{3} \mathcal{L}_{(J_ia_i)^{\sharp}} g = + \frac{1}{2} g(da_1, \omega_1) g - P_1(\pi_-^2(da_1)) - 2P_2(S_{0,3}^2(\nabla a_1) \diamond \omega_3) + 2P_3(S_{0,2}^2(\nabla a_1) \diamond \omega_2)
$$

+
$$
\frac{1}{2} g(da_2, \omega_2) g - P_2(\pi_-^2(da_2)) - 2P_3(S_{0,1}^2(\nabla a_2) \diamond \omega_1) + 2P_1(S_{0,3}^2(\nabla a_2) \diamond \omega_3)
$$

+
$$
\frac{1}{2} g(da_3, \omega_3) g - P_3(\pi_-^2(da_3)) - 2P_1(S_{0,2}^2(\nabla a_3) \diamond \omega_2) + 2P_2(S_{0,1}^2(\nabla a_3) \diamond \omega_1)
$$

+
$$
\text{l.o.t.}(\omega)
$$

=
$$
+ \frac{1}{2} \sum_{i=1}^{3} g(da_i, \omega_i) g - P_1(\pi_-^2(da_1)) - P_2(\pi_-^2(da_2)) - P_3(\pi_-^2(da_3))
$$

+
$$
+ 2P_3(S_{0,2}^2(\nabla a_1) \diamond \omega_2) - 2P_2(S_{0,3}^2(\nabla a_1) \diamond \omega_3) +
$$

+
$$
+ 2P_1(S_{0,3}^2(\nabla a_2) \diamond \omega_3) - 2P_3(S_{0,1}^2(\nabla a_2) \diamond \omega_1) +
$$

+
$$
+ 2P_2(S_{0,1}^2(\nabla a_3) \diamond \omega_1) - 2P_1(S_{0,2}^2(\nabla a_3) \diamond \omega_2) + \text{l.o.t.}(\omega).
$$

This proposition shows that the symmetric part of the negative gradient flow of (44) is not the Ricci flow to highest order, but rather there emerges another second-order term coming from the Lie derivative of g, by a certain combination of the torsion forms. As such, the short-time existence of negative gradient flow of [\(44\)](#page-11-1) is not immediate; we shall investigate this in [§5.](#page-24-0)

4.3.2 Self-dual Weyl in terms of intrinsic torsion

Proposition 4.7. *The self-dual Weyl curvature is explicitly given in terms of the intrinsic torsion by*

$$
W^{+} = -\frac{1}{24} \Big(2g(da_{1}, \omega_{1}) - g(da_{2}, \omega_{2}) - g(da_{3}, \omega_{3})
$$

+ $g(J_{3}a_{1}, a_{2}) - g(J_{2}a_{1}, a_{3}) - 2g(J_{1}a_{2}, a_{3}) \Big) (2\omega_{1} \otimes \omega_{1} - \omega_{2} \otimes \omega_{2} - \omega_{3} \otimes \omega_{3})$
- $\frac{1}{8} \Big(g(da_{2}, \omega_{2}) - g(da_{3}, \omega_{3}) + g(J_{3}a_{1}, a_{2}) + g(J_{2}a_{1}, a_{3}) \Big) (\omega_{2} \otimes \omega_{2} - \omega_{3} \otimes \omega_{3})$
- $\frac{1}{4} (g(da_{2}, \omega_{1}) + g(J_{1}a_{1}, a_{3})) (\omega_{1} \odot \omega_{2})$
- $\frac{1}{4} (g(da_{3}, \omega_{2}) + g(J_{2}a_{2}, a_{1})) (\omega_{2} \odot \omega_{3})$
- $\frac{1}{4} (g(da_{1}, \omega_{3}) + g(J_{3}a_{3}, a_{2})) (\omega_{3} \odot \omega_{1}),$

where $\omega_i \odot \omega_j := \omega_i \otimes \omega_j + \omega_j \otimes \omega_i \in S_0^2(\Lambda_+^2)$ for $i \neq j$.

Proof. The proof is again similar to that of the Ricci case. First, from Theorem [4.2](#page-20-1) and Corollary [2.11,](#page-8-1) we know that there are only 6 possible second-order terms, namely $g(da_i, \omega_j)$, with $i \leq j$. For the quadratic terms, we have 15 possibilities in view of [\(66\)](#page-21-2). So it is again a simple matter of determining the coefficient constants, which are obtained by verifying in suitable examples. \Box

Corollary 4.8. If J_1 is an integrable almost complex structure, then W^+ belong to a 3-dimensional subspace of $S_0^2(\Lambda_+^2)$, and *it is explicitly given by*

$$
W^{+} = -\frac{1}{24} \Big(2g(da_{1}, \omega_{1}) - g(da_{2}, \omega_{2}) - g(da_{3}, \omega_{3})
$$

+ $g(J_{3}a_{1}, a_{2}) - g(J_{2}a_{1}, a_{3}) - 2g(J_{1}a_{2}, a_{3}) \Big) (2\omega_{1} \otimes \omega_{1} - \omega_{2} \otimes \omega_{2} - \omega_{3} \otimes \omega_{3})$
- $\frac{1}{4} (g(da_{2}, \omega_{1}) + g(J_{1}a_{1}, a_{3})) (\omega_{1} \odot \omega_{2})$
- $\frac{1}{4} (g(da_{1}, \omega_{3}) + g(J_{3}a_{3}, a_{2})) (\omega_{3} \odot \omega_{1}).$

In particular, if M *is hyper-Hermitian, then* $W^+ = 0$ *. On the other hand, if* $\nabla J_1 = 0$ *i.e.* (M, g, J_1, ω_1) *is Kähler, then* W^+ belongs to a 1 -dimensional subspace of $S_0^2(\Lambda_+^2)$, and it is explicitly given by

$$
W^{+} = \frac{\text{Scal(g)}}{24} \Big(2\omega_1 \otimes \omega_1 - \omega_2 \otimes \omega_2 - \omega_3 \otimes \omega_3 \Big). \tag{70}
$$

Proof. Recall from Definition [2.14](#page-9-3) that $N_{J_1} = 0$ means $J_2 a_2 = J_3 a_3$, i.e. $a_2 = J_1 a_3$. To prove the first result, we take the exterior derivative of the relation $a_2 \wedge \omega_2 = J_1 a_3 \wedge \omega_2 = a_3 \wedge \omega_3$ to get $da_2 \wedge \omega_2 = da_3 \wedge \omega_3$. The second identity we need is obtained by differentiating $d\omega_1 = -2a_2 \wedge \omega_3 = 2a_3 \wedge \omega_2$. Together with Proposition [4.7,](#page-23-1) this proves the first expression. The last assertion follows by the same argument, replacing J_1 with J_2 and J_3 . In the Kähler situation we have $a_2 = a_3 = 0$, and hence $da_1 \wedge \omega_2 = da_1 \wedge \omega_3 = 0$ from [\(31\)](#page-7-8). Finally, from Proposition [4.4,](#page-21-1) $Scal(g) = 2\delta(J_1a_1) = -2g(da_1, \omega_1)$, and this concludes the proof. concludes the proof.

In the Kähler case, the expression [\(70\)](#page-23-2) for W^+ appears to be originally due to Derdzinski in [[Der83,](#page-39-25) Proposition 2], and the fact that hyper-Hermitian 4-manifolds have $W^+ = 0$ appears to be originally due to Boyer in [\[Boy88,](#page-39-26) Theorem 2].

Remark 4.3. While the anti-self-dual Weyl curvature W[−] cannot be computed from the SU(2) torsion forms, we can nonetheless express it as follows. If we choose a local SU(2) coframing $\{e^1, e^2, e^3, e^4\}$ on M^4 , so that Λ^2 can be trivialised by

$$
\omega_1 = e^{12} + e^{34}
$$
, $\omega_2 = e^{13} + e^{42}$, $\omega_3 = e^{14} + e^{23}$,
\n $\sigma_1 = e^{12} - e^{34}$, $\sigma_2 = e^{13} - e^{42}$, $\sigma_3 = e^{14} - e^{23}$,

then as in Proposition [2.9](#page-7-0) we can define "anti-self-dual torsion 1-forms" b_i by

$$
d\sigma_1 = -b_2 \wedge \sigma_3 + b_3 \wedge \sigma_2, d\sigma_2 = -b_3 \wedge \sigma_1 + b_1 \wedge \sigma_3, d\sigma_3 = -b_1 \wedge \sigma_2 + b_2 \wedge \sigma_1.
$$

The anti-self-dual Weyl curvature W^- can then be expressed as in Proposition [4.7,](#page-23-1) by replacing (a_i, ω_i) with (b_i, σ_i) .

We now proceed to our main section, where we establish short-time existence for a large class of $SU(2)$ -flows.

5 Parabolicity of $SU(2)$ -flows

As we showed in Theorem [4.2,](#page-20-1) all the second-order invariants of an $SU(2)$ -structure are determined by the suitable irreducible components of ∇a_i and W^- . Since W^- lies the 5-dimensional module $S_0^2(\mathbb{R}^3)$ and the latter does not occur in End(\mathbb{R}^4)/ $\mathfrak{su}(2) \cong 4\mathbb{R} \oplus 3\mathbb{R}^3$, it follows that we cannot evolve ω_i using W^- via a second order geometric flow, see also [§6.1.](#page-35-1) On the other hand, all the components of ∇a_i lie in the SU(2)-modules R and R³, hence all diffeomorphism-invariant second order flows of SU(2)-structures, to highest order, can be obtained by choosing f_0, f_1, f_2, f_3 among $\delta(a_i)$ and $g(da_i, \omega_j)$ $(i \leq j)$, and $B_{0,1}, B_{0,2}, B_{0,3}$ among $\pi^2 \cdot (da_i)$ and $S_{0,i}^2(\nabla a_j)$ (each suitably interpreted as a tensor of the right type via the identifications defined in $\S2.1$). In this section we shall establish short-time existence for a large class of such flows.

5.1 Linearisations and principal symbols

We begin by computing the principal symbols of the differential operators in Theorem [4.2](#page-20-1) that lead to SU(2)-flows. For convenience we have summarised the result in Appendix [A.](#page-38-0) First we recall the basic definitions following [\[Top06\]](#page-39-18).

Definition 5.1. Let E and F denote smooth vector bundles over M and let $L : \Gamma(E) \to \Gamma(F)$ be a linear second order partial differential operator. The *principal symbol* of L is the vector bundle homomorphism $\sigma(L) : \pi^*(E) \to \pi^*(F)$, where $\pi: T^*M \to M$, defined as follows: given $(x,\xi) \in T^*M$, $s \in \Gamma(E)$ such that $s(x) = v$ and $f \in C^{\infty}(M)$ with $df(x) = \xi$, then 2

$$
\sigma(L)(x,\xi)v:=\lim_{t\to\infty}2t^{-2}e^{-tf}L(e^{tf}s)(x).
$$

When, moreover, $E = F$ and $\langle \cdot, \cdot \rangle$ denotes any fibre metric on E, we say that the flow

$$
\partial_t u_t = L(u_t)
$$

is *parabolic* if there exists $\lambda > 0$ such that

$$
\langle \sigma(L)(x,\xi)v,v\rangle \geq \lambda |\xi|^2 |v|^2, \quad \forall (x,\xi) \in T^*M \quad \text{and} \quad v \in \Gamma(E).
$$

We extend the above notions to a non-linear partial differential operator P by requiring that its linearisation L_p , at any given $v \in \Gamma(E)$, satisfies the above parabolicity condition.

Here we are interested in non-linear differential operators of the form

$$
\mathcal{P}: \mathcal{M} \to \Omega^2(M), \n\omega \mapsto \mathcal{P}(\omega)
$$

where $\mathcal{M} \subset \Omega^2(M) \times \Omega^2(M) \times \Omega^2(M)$ is a bundle whose fibres are isomorphic to the 13-dimensional space GL(4, R)/SU(2), and P is a linear combination of the invariants given in 1. and 2. of Theorem [4.2,](#page-20-1) up to $SU(2)$ isomorphism. More precisely, we are viewing the invariants of Theorem [4.2](#page-20-1) as second order differential operators acting on ω . The reason we emphasise 'up to SU(2) isomorphism' above is because, for instance, the term $\pi^2_-(da_1)$ can be viewed a tensor in $\Lambda^2_-, \Sigma^2_{0,1}, \Sigma^2_{0,2}$ or $\Sigma^2_{0,3}$ (which are all equivalent as SU(2)-modules) but depending on what type of tensor we identify it with, this leads to a different geometric flow, see also the last paragraph of [§5.4](#page-31-0) below.

 2 The extra factor of '2' in the definition of the principal symbol is merely for convenience.

Proposition 5.2. The principal symbols of the non-linear second order differential operators ∇a_i at the point $(x,\xi) \in T^*M$ *are given by*

$$
\sigma(L_{\nabla a_1})(x,\xi)(A) = -\xi \otimes *(\xi \wedge \tilde{B}_{0,1}) - \xi \otimes * (J_2 \xi \wedge \tilde{B}_{0,3}) + \xi \otimes * (J_3 \xi \wedge \tilde{B}_{0,2}) + 2f_1 \xi \otimes \xi - f_0 \xi \otimes J_1 \xi,
$$

$$
\sigma(L_{\nabla a_2})(x,\xi)(A) = -\xi \otimes *(\xi \wedge \tilde{B}_{0,2}) - \xi \otimes * (J_3 \xi \wedge \tilde{B}_{0,1}) + \xi \otimes * (J_1 \xi \wedge \tilde{B}_{0,3}) + 2f_2 \xi \otimes \xi - f_0 \xi \otimes J_2 \xi,
$$

$$
\sigma(L_{\nabla a_3})(x,\xi)(A) = -\xi \otimes *(\xi \wedge \tilde{B}_{0,3}) - \xi \otimes * (J_1 \xi \wedge \tilde{B}_{0,2}) + \xi \otimes * (J_2 \xi \wedge \tilde{B}_{0,1}) + 2f_3 \xi \otimes \xi - f_0 \xi \otimes J_3 \xi,
$$

where $L_{\nabla a_i}$ denotes the linearisation of ∇a_i , $A = f_0 g + \sum_{i=1}^3 B_{0,1} + \sum_{i=1}^3 f_i \omega_i$ is the variation of ω as given in [\(2\)](#page-1-4)-[\(4\)](#page-1-5) and $\tilde{B}_{0,i} := B_{0,i} \diamond \omega_i \in \Lambda^2_-(M).$

Proof. From Proposition [2.9,](#page-7-0) we have

$$
\nabla a_1 = -\frac{1}{2}\nabla(*d\omega_1 + J_2(*d\omega_3) - J_3(*d\omega_2)).
$$

It follows that the linearisation of the second order operator ∇a_1 , at the point $\omega = (\omega_1, \omega_2, \omega_3)$, in the direction of A, as given by $(2)-(4)$ $(2)-(4)$ is

$$
L_{\nabla a_1}(A) = -\frac{1}{2} \Big(\nabla * d(f_0 \omega_1 - f_2 \omega_3 + f_3 \omega_2 + \tilde{B}_{0,1}) + \nabla J_2 * d(f_0 \omega_3 - f_1 \omega_2 + f_2 \omega_1 + \tilde{B}_{0,3}) -
$$

$$
\nabla J_3 * d(f_0 \omega_2 - f_3 \omega_1 + f_1 \omega_3 + \tilde{B}_{0,2}) \Big) + \text{l.o.t.}(A).
$$

Given an arbitrary function f ,

$$
-2L_{\nabla a_1}(e^{tf}A) = t^2 e^{tf} df \otimes \left((-f_0 d^{c_1} f + f_2 d^{c_3} f - f_3 d^{c_2} f + *(df \wedge \tilde{B}_{0,1})) +
$$

\n
$$
J_2(-f_0 d^{c_3} f + f_1 d^{c_2} f - f_2 d^{c_1} f + *(df \wedge \tilde{B}_{0,3})) -
$$

\n
$$
J_3(-f_0 d^{c_2} f + f_3 d^{c_1} f - f_1 d^{c_3} f + *(df \wedge \tilde{B}_{0,2})) + l.o.t.(t)
$$

\n
$$
= t^2 e^{tf} df \otimes \left((-f_0 d^{c_1} f + f_2 d^{c_3} f - f_3 d^{c_2} f + *(df \wedge \tilde{B}_{0,1})) +
$$

\n
$$
(+f_0 d^{c_1} f - f_1 df - f_2 d^{c_3} f + *(d^{c_2} f \wedge \tilde{B}_{0,3})) +
$$

\n
$$
(+f_0 d^{c_1} f + f_3 d^{c_2} f - f_1 df - *(d^{c_3} f \wedge \tilde{B}_{0,2})) \right) + l.o.t.(t)
$$

\n
$$
= t^2 e^{tf} df \otimes \left(f_0 d^{c_1} f - 2f_1 df + *(df \wedge \tilde{B}_{0,1}) + *(d^{c_2} f \wedge \tilde{B}_{0,3}) - *(d^{c_3} f \wedge \tilde{B}_{0,2}) \right) + l.o.t.(t)
$$

where we used Lemma [2.3](#page-4-1) and the fact that $J_j(\tilde{B}_{0,i}) = \tilde{B}_{0,i}$, since $\tilde{B}_{0,i} \in \Omega^2(M)$. The principal symbol of $L_{\nabla a_1}$ now follows immediately from the above, and an analogous computation applies to $L_{\nabla a_1}$ and $L_{\nabla a_2}$. \Box

We can now perform a similar computation to derive the principal symbols of each SU(2)-invariant second-order differential operator δa_i , $g(da_i, \omega_j)$, $S^2_{0,i}(\nabla a_j)$ and $\pi^2(d a_i)$. However, there is a simpler way to do this, since the principal symbols of the latter terms can be deduced from those of ∇a_i , by an appropriate projection map. For instance, for any 1-form α , we have $d\alpha(X, Y) = (\nabla_X \alpha)(Y) - (\nabla_Y \alpha)(X)$ and thus, from the above Proposition, we immediately deduce

$$
\sigma(L_{da_1})(x,\xi)(A) = -\xi \wedge *(\xi \wedge \tilde{B}_{0,1}) - \xi \wedge * (J_2 \xi \wedge \tilde{B}_{0,3}) + \xi \wedge * (J_3 \xi \wedge \tilde{B}_{0,2}) - f_0 \xi \wedge J_1 \xi, \tag{71}
$$

and similarly for $\sigma(L_{da_2})$ and $\sigma(L_{da_3})$. Unlike the skew-symmetrisation operator, the SU(2) projection maps do depend on ω . However, since these maps are only of zeroth order in ω , these do not affect the principal symbols when we consider their linearisations. Thus, their components in ω_i and $\Lambda^2(M)$ can be computed using Lemmas [2.3](#page-4-1) and [2.4.](#page-5-3) For instance, one can express [\(71\)](#page-25-0) explicitly as

$$
\sigma(L_{da_1})(x,\xi)(A) = +\frac{1}{2}(f_0|\xi|^2 - g(\xi \wedge J_1\xi, \tilde{B}_{0,1}) + g(\xi \wedge J_2\xi, \tilde{B}_{0,2}) + g(\xi \wedge J_3\xi, \tilde{B}_{0,3})) \omega_1 \n- \frac{1}{2}(g(\xi \wedge J_2\xi, \tilde{B}_{0,1}) + g(\xi \wedge J_1\xi, \tilde{B}_{0,2})) \omega_2 \n- \frac{1}{2}(g(\xi \wedge J_3\xi, \tilde{B}_{0,1}) + g(\xi \wedge J_1\xi, \tilde{B}_{0,3})) \omega_3 \n+ \frac{1}{2}|\xi|^2 \tilde{B}_{0,1} - \frac{1}{2}f_0 \sum_i g(\xi \wedge J_1\xi, \sigma_i)\sigma_i \n- \sum_{i,j,k} \varepsilon_{ijk} \frac{1}{4}(g(\xi \wedge J_3\xi, \sigma_i)g(\tilde{B}_{0,2}, \sigma_j))\sigma_k + \sum_{i,j,k} \varepsilon_{ijk} \frac{1}{4}(g(\xi \wedge J_2\xi, \sigma_i)g(\tilde{B}_{0,3}, \sigma_j))\sigma_k
$$

and similarly for $\sigma(L_{da_2})$ and $\sigma(L_{da_3})$. In the same vein, one can use Lemma [2.3](#page-4-1) to show that

$$
\sigma(L_{\delta a_1})(x,\xi)(A) = -2f_1|\xi|^2 - *(\xi \wedge J_2\xi \wedge \tilde{B}_{0,3}) + *(\xi \wedge J_3\xi \wedge \tilde{B}_{0,2}),
$$
\n(72)

and similarly for $\sigma(L_{\delta a_1})$ and $\sigma(L_{\delta a_1})$. It now remains to compute the principal symbols of the terms $S^2_{0,i}(\nabla a_j)$. Thus, we need to apply the projection maps $S_{0,i}^2$ defined in [§2.1](#page-3-2) to the principal symbols computed in Proposition [5.2,](#page-25-1) to do so we appeal to the following lemma:

Lemma 5.3. Let α, β be 1-forms, $B_{0,i}$ be arbitrary symmetric 2-tensors in $S_{0,i}^2$, and $\tilde{B}_{0,i} := B_{0,i} \circ \omega_i \in \Lambda^2_-,$ then the *following relations hold:*

$$
S_{0,i}^{2}(\alpha \otimes J_{i}\beta) \diamond \omega_{i} = \frac{1}{2}\pi_{-}^{2}(\alpha \wedge \beta),
$$

\n
$$
S_{0,i}^{2}(\alpha \otimes B_{0,i}(\alpha^{\sharp},\cdot)) = \frac{1}{4}|\alpha|^{2}B_{0,i},
$$

\n
$$
\pi_{-}^{2}(\alpha \wedge *(\alpha \wedge \tilde{B}_{0,i})) = -\frac{1}{2}|\alpha|^{2}\tilde{B}_{0,i},
$$

\n
$$
*(\alpha \wedge \tilde{B}_{0,i}) = -B_{0,i}(J_{i}(\alpha^{\sharp}),\cdot) = -\alpha^{\sharp}\Box\tilde{B}_{0,i},
$$

\n
$$
8S_{0,i}^{2}(\alpha \otimes B_{0,j}(\beta^{\sharp},\cdot)) \diamond \omega_{i} = -\sum_{a,b,c} \varepsilon_{abcd}g(\tilde{B}_{0,j},\sigma_{a})g(\alpha \wedge J_{k}\beta,\sigma_{b})\sigma_{c} + 2g(\alpha,J_{k}\beta)B_{0,j},
$$

\n
$$
8S_{0,i}^{2}(\alpha \otimes B_{0,k}(\beta^{\sharp},\cdot)) \diamond \omega_{i} = +\sum_{a,b,c} \varepsilon_{abc}g(\tilde{B}_{0,k},\sigma_{a})g(\alpha \wedge J_{j}\beta,\sigma_{b})\sigma_{c} - 2g(\alpha,J_{j}\beta)B_{0,k},
$$

where $(i, j, k) \sim (1, 2, 3)$ *in the last two expressions. In particular,*

$$
8S_{0,i}^2(\alpha \otimes B_{0,j}(\alpha^{\sharp},\cdot)) \diamond \omega_i = -\sum_{a,b,c} \varepsilon_{abc} g(\tilde{B}_{0,j},\sigma_a) g(\alpha \wedge J_k \alpha, \sigma_b) \sigma_c,
$$

$$
8S_{0,i}^2(\alpha \otimes B_{0,k}(\alpha^{\sharp},\cdot)) \diamond \omega_i = +\sum_{a,b,c} \varepsilon_{abc} g(\tilde{B}_{0,k},\sigma_a) g(\alpha \wedge J_j \alpha, \sigma_b) \sigma_c.
$$

Proof. The maps involved are all SU(2)-invariant, and hence by Schur's lemma it suffices to verify the above expressions for suitable choices of α , β and $B_{0,i}$. This is a long but straightforward calculation, using the above local expressions for ω_i and σ_i . \Box

Since $S^2_{0,i}(M) \cong \Lambda^2_{-}(M)$, describing the latter principal symbols as anti-self-dual forms will facilitate comparisons. Using Proposition [5.2](#page-25-1) and Lemma [5.3,](#page-26-0)

$$
\sigma(L_{S_{0,1}^2(\nabla a_1)})(x,\xi)(A) \diamond \omega_1 = \frac{1}{2}\pi^2 \left(\xi \wedge J_1(\xi^{\sharp}) \Box \tilde{B}_{0,1} - \xi \wedge J_3(\xi^{\sharp}) \Box \tilde{B}_{0,3} - \xi \wedge J_2(\xi^{\sharp}) \Box \tilde{B}_{0,2} - 2f_1\xi \wedge J_1\xi\right)
$$

\n
$$
= -\frac{1}{2}f_1 \sum_i g(\xi \wedge J_1\xi, \sigma_i)\sigma_i - \frac{1}{8} \sum_{i,j,k} \varepsilon_{ijk}g(\xi \wedge J_2\xi, \sigma_j)g(\tilde{B}_{0,2}, \sigma_i)\sigma_k
$$

\n
$$
+ \frac{1}{8} \sum_{i,j,k} \varepsilon_{ijk}g(\xi \wedge J_1\xi, \sigma_j)g(\tilde{B}_{0,1}, \sigma_i)\sigma_k - \frac{1}{8} \sum_{i,j,k} \varepsilon_{ijk}g(\xi \wedge J_3\xi, \sigma_j)g(\tilde{B}_{0,3}, \sigma_i)\sigma_k,
$$

$$
\sigma(L_{S_{0,2}^{2}(\nabla a_{1})})(x,\xi)(A) \diamond \omega_{2} = \frac{1}{2}\pi_{-}^{2}\left(\xi \wedge J_{2}(\xi^{\sharp})\Box \tilde{B}_{0,1} + \xi \wedge \xi^{\sharp}\Box \tilde{B}_{0,3} + \xi \wedge J_{1}(\xi^{\sharp})\Box \tilde{B}_{0,2} - 2f_{1}\xi \wedge J_{2}\xi + f_{0}\xi \wedge J_{3}\xi\right)
$$

\n
$$
= +\frac{1}{4}|\xi|^{2}\tilde{B}_{0,3} + \frac{1}{4}f_{0}\sum_{i}g(\xi \wedge J_{3}\xi,\sigma_{i})\sigma_{i} - \frac{1}{2}f_{1}\sum_{i}g(\xi \wedge J_{2}\xi,\sigma_{i})\sigma_{i}
$$

\n
$$
+ \frac{1}{8}\sum_{i,j,k}\varepsilon_{ijk}g(\xi \wedge J_{2}\xi,\sigma_{j})g(\tilde{B}_{0,1},\sigma_{i})\sigma_{k} + \frac{1}{8}\sum_{i,j,k}\varepsilon_{ijk}g(\xi \wedge J_{1}\xi,\sigma_{j})g(\tilde{B}_{0,2},\sigma_{i})\sigma_{k},
$$

$$
\sigma(L_{S_{0,3}^{2}(\nabla a_{1})})(x,\xi)(A) \diamond \omega_{3} = \frac{1}{2}\pi^{2}\left(\xi \wedge J_{3}(\xi^{\sharp})\varphi_{0,1} + \xi \wedge (J_{1}\xi^{\sharp})\varphi_{0,3} - \xi \wedge \xi^{\sharp}\varphi_{0,2} - 2f_{1}\xi \wedge J_{3}\xi - f_{0}\xi \wedge J_{2}\xi\right)
$$

\n
$$
= -\frac{1}{4}|\xi|^{2}\tilde{B}_{0,2} - \frac{1}{4}f_{0}\sum_{i}g(\xi \wedge J_{2}\xi,\sigma_{i})\sigma_{i} - \frac{1}{2}f_{1}\sum_{i}g(\xi \wedge J_{3}\xi,\sigma_{i})\sigma_{i}
$$

\n
$$
+ \frac{1}{8}\sum_{i,j,k}\varepsilon_{ijk}g(\xi \wedge J_{3}\xi,\sigma_{j})g(\tilde{B}_{0,1},\sigma_{i})\sigma_{k} + \frac{1}{8}\sum_{i,j,k}\varepsilon_{ijk}g(\xi \wedge J_{1}\xi,\sigma_{j})g(\tilde{B}_{0,3},\sigma_{i})\sigma_{k}.
$$

To obtain the remaining 9 terms, it suffices to take cyclic permutation of the indices (123) above; this is summarised in Appendix [A.](#page-38-0) Having computed the principal symbols, we next investigate which $SU(2)$ -flows are parabolic.

5.2 The Ricci flow and DeTurck's trick

Let us briefly recall the proof of short-time existence of the Ricci flow, since we shall apply a very similar strategy. There are several excellent references on the topic, cf. [\[CK04,](#page-39-14) [CLN06\]](#page-39-27), but we shall follow the exposition by Topping [\[Top06\]](#page-39-18). From now on the manifold $M⁴$ will be assumed to be compact, in order to appeal to the standard result that parabolic flows exist for short time and are unique, given regular enough initial data.

From [\[Top06,](#page-39-18) Chapter 5], the principal symbol of the linearisation of the Ricci map, $g \mapsto \text{Ric}(g)$, is

$$
-\sigma(L_{\mathrm{Ric}(g)})(x,\xi)(A) = |\xi|^2 B - \xi \otimes B(\xi^{\sharp},\cdot) - B(\xi^{\sharp},\cdot) \otimes \xi + 4f_0(\xi \otimes \xi),\tag{73}
$$

where we recall that the variation of g is given by (14) . In particular, we see that the Ricci map is not elliptic, since its principal symbol is not invertible. From the work of Hamilton and DeTurck, we know that the kernel of the principal symbol is precisely generated by the diffeomorphism group. Using the results from Appendix [A](#page-38-0) and Lemma [5.3,](#page-26-0) it is not hard to verify directly that

$$
S^2_{0,i}(\sigma(L_{\mathrm{Ric}(g)})(x,\xi)(A)) \diamond \omega_i = -\sigma(L_{\pi^2_{-}(da_i)})(x,\xi)(A)
$$

and

$$
\text{tr}(\sigma(L_{\text{Ric}(g)})(x,\xi)(A)) = -\frac{1}{2}\sigma(L_{g(da_1,\omega_1)+g(da_2,\omega_2)+g(da_3,\omega_3)}(x,\xi)(A).
$$

In view of Proposition [4.4](#page-21-1) this was indeed to be expected, so this confirms that everything is consistent. We shall now recall DeTurck's trick to prove short-time existence of the Ricci flow.

We begin by considering the *gravitation map* $G : S^2(M) \to S^2(M)$, defined by

$$
G(H) = H - \frac{1}{2}(\operatorname{tr} H)g,
$$

where H is an arbitrary positive definite symmetric tensor, and its divergence by $\delta G(H) = -\text{tr}(\nabla H) + \frac{1}{2}d(\text{tr }H)$. Observe that $\delta G(H)$ is a first-order differential operator acting on g. It is shown in [\[Top06,](#page-39-18) Chapter 5] that the linearisation of $\mathcal{L}_{(H^{-1}\delta G(H))}$ ‡g is given by

$$
\partial_t(\mathcal{L}_{(H^{-1}\delta G(H))^{\sharp}}g) = -\mathcal{L}_{(\delta G(B))^{\sharp}}g + \text{l.o.t}(g)
$$

and hence its principal symbol is

$$
\sigma(\mathcal{L}_{(\delta G(B))^{\sharp}}g)(x,\xi)(A)=-2\xi\otimes B(\xi^{\sharp},\cdot)-2B(\xi^{\sharp},\cdot)\otimes \xi+8f_0(\xi\otimes \xi).
$$

It follows that

$$
-2\sigma(L_{\text{Ric}(g)})(x,\xi)(A) - \sigma(\mathcal{L}_{(\delta G(B))^{\sharp}}g)(x,\xi)(A) = 2|\xi|^{2}B,
$$
\n(74)

and we deduce that the Ricci-DeTurck flow

$$
\partial_t(g(t)) = -2\text{Ric}(g(t)) + \mathcal{L}_{(H^{-1}\delta G(H))^{\sharp}}g(t)
$$
\n(75)

is parabolic, admitting as such a unique solution for short time, given smooth initial data $g(0) = g_0$. We now let $X_t =$ $-\tilde{H}^{-1}\delta G(H)$ [‡] denote a time-dependent vector field (associated to the metric solving [\(75\)](#page-27-1)) and $\{\varphi_t\}$ the associated 1parameter family of diffeomorphisms defined by

$$
X_t \circ \varphi_t(p) = \frac{d}{dt} \varphi_t(p), \quad \text{with} \quad \varphi_0 = \text{Id}.
$$
 (76)

Setting $\hat{g}(t) := \varphi_t^*(g(t))$, we then have

$$
\partial_t(\hat{g}(t)) = \varphi_t^*(-2\mathrm{Ric}(g) + \mathcal{L}_{(H^{-1}\delta G(H))^{\sharp}}g + \mathcal{L}_{X_t}g) = -2\mathrm{Ric}(\varphi_t^*g(t)),
$$

and this gives the desired solution to the Ricci flow starting at $\hat{g}(0) = g_0$.

Since our main goal is to classify parabolic flows of $SU(2)$ -structures, it is natural to consider those which induce the Ricci flow on the metric, to highest order. From Proposition [4.4,](#page-21-1) it follows that such a flow is necessarily of the form:

$$
\frac{d}{dt}\omega_1 = +\sum_{i=1}^3 g(da_i, \omega_i)\omega_1 + f_3\omega_2 - f_2\omega_3 + 2\pi^2_-(da_1) + \text{l.o.t.}(\omega)
$$
\n(77)

$$
\frac{d}{dt}\omega_2 = -f_3\omega_1 + \sum_{i=1}^3 g(da_i, \omega_i)\omega_2 + f_1\omega_3 + 2\pi^2 (da_2) + \text{l.o.t.}(\omega)
$$
\n(78)

$$
\frac{d}{dt}\omega_3 = +f_2\omega_1 - f_1\omega_2 + \sum_{i=1}^3 g(da_i, \omega_i)\omega_3 + 2\pi^2(da_3) + \text{l.o.t.}(\omega)
$$
\n(79)

or, more compactly,

$$
\partial_t(\boldsymbol{\omega}) = \left(\sum_{i=1}^3 g(da_i, \omega_i)g - 2\sum_{i=1}^3 P_i(\pi_{-}^2(da_i)) + \sum_{i=1}^3 f_i\omega_i\right) \diamond \boldsymbol{\omega} + \text{l.o.t.}(\boldsymbol{\omega})
$$

$$
= \left(-2\text{Ric}(g) + \sum_{i=1}^3 f_i\omega_i\right) \diamond \boldsymbol{\omega} + \text{l.o.t.}(\boldsymbol{\omega})
$$

Observe that this SU(2)-flow is under-determined, since the functions f_1, f_2, f_3 are not prescribed. Motivated by the results in $[LSE23]$ (see also Proposition [5.5](#page-28-2) below), it is natural to ask if the Ricci flow, when coupled with the harmonic flow, is parabolic. We investigate more general classes of such flows in the next two sections.

5.3 Harmonic Ricci-like flows

The main goal of this section is to establish the following result:

Theorem 5.4. *Consider the flow of* SU(2)*-structures given by*

$$
\partial_t(\omega) = \left(-2\text{Ric}(g) - \mu \sum_{i=1}^3 \mathcal{L}_{(J_i a_i)^{\sharp}} g + \lambda \sum_{i=1}^3 \text{div}(a_i) \omega_i\right) \diamond \omega + \text{l.o.t.}(\omega),
$$

$$
\omega(0) = \omega_0,
$$
 (80)

where $\mu, \lambda \in \mathbb{R}$ and ω_0 *is a smooth* SU(2)-triple. Then there exists a smooth solution $\omega(t)$ to [\(80\)](#page-28-3) provided that $9 - 4\sqrt{5}$ < $\lambda + \frac{\mu}{2} < 9 + 4\sqrt{5}$. Moreover, this solution is unique.

Before proving the above theorem, we first make a few observations. If we consider the flow given by

$$
\partial_t(\boldsymbol{\omega}) = \left(-2\varepsilon \text{Ric}(g) - \mu \sum_{i=1}^3 \mathcal{L}_{(J_i a_i)^{\sharp}} g + \lambda \sum_{i=1}^3 \text{div}(a_i) \omega_i\right) \diamond \boldsymbol{\omega},\tag{81}
$$

then by rescaling we can always assume that $\varepsilon = -1$, 0 or 1. We disregard the case $\varepsilon = -1$, since backwards heat-type flows are never parabolic. Henceforth, we will ignore lower-order terms and omit writing $l.o.t.(\omega)$ altogether, as these do not play any role in the proof of Theorem [5.4.](#page-28-1) The family of flows given by [\(81\)](#page-28-4) contains three notable cases:

- 1. If $\varepsilon = \mu = 0$ and $\lambda = 1$, then the flow reduces to the *harmonic flow* of SU(2)-structures cf. [\[LSE23\]](#page-39-0). This is the negative gradient flow of [\(44\)](#page-11-1) whereby the metric is fixed; it is sometimes also called the *isometric flow* in the literature.
- 2. If $\varepsilon = \lambda = 1$ and $\mu = 0$, then it corresponds to the *harmonic Ricci flow*. We should point out that, since the metric is also evolving, it is not immediately clear that this flow exists.
- 3. If $\varepsilon = \mu = 1$ and $\lambda = 1/2$, then this corresponds to the negative *Dirichlet gradient flow* of [\(44\)](#page-11-1); this follows from Proposition [3.4,](#page-11-2) [3.5](#page-12-0) and [4.6.](#page-22-0)

The authors of $[LSE23]$ establish short-time existence and uniqueness of any harmonic flow of H-structures by showing that it can be interpreted as a harmonic map heat flow for certain equivariant maps. In the case of harmonic G_2 -structures considered in [\[DGK21\]](#page-39-28), this was proven directly by showing that the flow is parabolic. In our setup here, we can also give a direct proof, as an immediate consequence of the results in [§5.1:](#page-24-1)

Proposition 5.5. *The harmonic* SU(2)-flow exists for short time and is unique, for given smooth initial data ω_0 .

Proof. If we restrict to only isometric variations of ω , i.e. we set $B = 0$ and $C = \sum_{i=1}^{3} f_i \omega_i$ in [\(2\)](#page-1-4)-[\(4\)](#page-1-5), then from the principal symbols listed in [A](#page-38-0)ppendix A we see that $\sigma(L_{div(a_i)})(x,\xi)(C) = 2f_i|\xi|^2$, for $i = 1, 2, 3$. It follows that

$$
\langle \sigma(L_{\sum_{i=1}^3 \text{div}(a_i)\omega_i})(x,\xi)(C), C \rangle = 4|\xi|^2 \sum_{i=1}^3 f_i^2 = 2|\xi|^2|C|^2.
$$

Thus, the harmonic flow is immediately seen to be parabolic and the result follows.

It worth pointing out that we did not have to rely on DeTurck's trick in the above proof, as the flow was already parabolic. With these observations out of the way, we now begin the proof of Theorem [5.4.](#page-28-1)

Consider the modified flow of (80) given by

$$
\partial_t(\boldsymbol{\omega}) = (-2\mathrm{Ric}(g) - \mu \sum_{i=1}^3 \mathcal{L}_{(J_i a_i)^{\sharp}} g + \lambda \sum_{i=1}^3 \mathrm{div}(a_i) \omega_i) \diamond \boldsymbol{\omega} + \mathcal{L}_{(H^{-1} \delta G(H))^{\sharp} + \mu \sum_{i=1}^3 (J_i a_i)^{\sharp}} \boldsymbol{\omega}.
$$
(82)

 \Box

From Lemma [2.6,](#page-5-8) we can express the latter equivalently as

$$
\partial_t(\omega) = (-2\mathrm{Ric}(g) + \mathcal{L}_{(H^{-1}\delta G(H))^{\sharp}}g + \lambda \sum_{i=1}^3 \mathrm{div}(a_i)\omega_i - d(H^{-1}\delta G(H) + \mu \sum_{i=1}^3 J_i a_i)) \diamond \omega. \tag{83}
$$

For convenience we shall write

$$
\mathcal{P}(\omega) = \mathcal{P}_{sym}(\omega) + \mathcal{P}_{alt}(\omega)
$$

 := $\left(-2\text{Ric}(g) + \mathcal{L}_{(H^{-1}\delta G(H))^{\sharp}}g\right) + \left(\lambda \sum_{i=1}^{3} \text{div}(a_{i})\omega_{i} - d(H^{-1}\delta G(H) + \mu \sum_{i=1}^{3} (J_{i}a_{i}))\right),$

where P_{sym} and P_{alt} denote the symmetric and skew-symmetric parts respectively. In particular, observe that P_{sym} induces precisely the Ricci-DeTurck flow [\(75\)](#page-27-1), hence it follows from [\(74\)](#page-27-2) that

$$
\langle \sigma(L_{\mathcal{P}})(x,\xi)(A),B\rangle = 2|\xi|^2|B|^2.
$$
\n(84)

The choice of the modified flow [\(82\)](#page-28-5) was precisely intended to obtain [\(84\)](#page-29-0). Thus, it only remains to consider the skewsymmetric part.

First note that, to highest order, $d(J_i a_i) = 2\pi^2 (\nabla a_i(\cdot, J_i \cdot))$, where π^2 denotes the projection of a 2-tensor on $\Lambda^2(M)$. On the other hand, using [\(17\)](#page-5-7),

$$
\nabla a_1(\cdot, J_1 \cdot) = \frac{1}{4} (\delta a_1) \omega_1 + \frac{1}{4} g(da_1, \omega_3) \omega_2 - \frac{1}{4} g(da_1, \omega_2) \omega_3 + \cdots,
$$

where \cdots consists of the components of $\nabla a_1(\cdot, J_1)$ in the complement of $\langle \omega_1, \omega_2, \omega_3 \rangle$ in $T^* \otimes T^*$; likewise we get analogous expressions for $\nabla a_2(\cdot, J_2 \cdot)$ and $\nabla a_3(\cdot, J_3 \cdot)$. Using the above together with Appendix [A,](#page-38-0) and the fact that $\Lambda^2_-(M) \subset \text{ker}(\infty)$, we deduce that

$$
2\sigma(L_{\sum_{i=1}^{3}d(J_{i}a_{i})})(x,\xi)(A) \diamond \omega = \left(\sigma(L_{\delta a_{1}-g(da_{2},\omega_{3})+g(da_{3},\omega_{2})}(x,\xi)(A)\omega_{1} + \sigma(L_{\delta a_{2}-g(da_{3},\omega_{1})+g(da_{1},\omega_{3})}(x,\xi)(A)\omega_{2} + \sigma(L_{\delta a_{3}-g(da_{1},\omega_{2})+g(da_{2},\omega_{1})}(x,\xi)(A)\omega_{3}\right) \diamond \omega,
$$

$$
= \left(\sum_{i=1}^{3}\sigma(L_{\delta a_{i}})(x,\xi)(A)\omega_{i}\right) \diamond \omega.
$$

Next we need to compute the self-dual components of the principal symbol of $d(H^{-1}\delta G(H))$, whose linearisation, we recall, is given by $-\delta G(B) = \text{tr}(\nabla B) - \frac{1}{2}d(\text{tr }B)$. Thus, we have that

$$
\sigma(d(\delta G(B)))(x,\xi) = -2\xi \wedge \left(f_0 g(\xi^{\sharp},\cdot) + \sum_{i=1}^3 \text{tr}(\xi \otimes B_{0,i})\right)
$$

$$
= -2\xi \wedge (\sum_{i=1}^3 B_{0,i}(\xi^{\sharp},\cdot))
$$

$$
= +*(\xi \wedge J_3\xi \wedge \tilde{B}_{0,2} - \xi \wedge J_2\xi \wedge \tilde{B}_{0,3}) \omega_1
$$

$$
+*(\xi \wedge J_1\xi \wedge \tilde{B}_{0,3} - \xi \wedge J_3\xi \wedge \tilde{B}_{0,1}) \omega_2
$$

$$
+*(\xi \wedge J_2\xi \wedge \tilde{B}_{0,1} - \xi \wedge J_1\xi \wedge \tilde{B}_{0,2}) \omega_3 + \cdots,
$$

where \cdots denotes the anti-self-dual terms, and for the last equality we used

$$
\xi \wedge B_{0,i}(\xi^{\sharp},\cdot) \wedge \omega_j = \xi \wedge J_j(J_i\xi) \wedge \tilde{B}_{0,i}, \text{ for } i,j = 1,2,3.
$$

The latter identity is obtained as follows: since $\tilde{B}_{0,i} \wedge \omega_j = 0$, it follows that $(X \Box \tilde{B}_{0,i}) \wedge \omega_j = -\tilde{B}_{0,i} \wedge (X \Box \omega_j) =$ $\tilde{B}_{0,i} \wedge J_j(X^{\flat})$, for any vector field X. The result follows from the fact that $B_{0,i}(\cdot, \cdot) = -\tilde{B}_{0,i}(J_i, \cdot)$, cf. Lemma [2.4.](#page-5-3) Using the principal symbols in Appendix [A,](#page-38-0) we can summarise the result of the above computations into

$$
\sigma(L_{\mathcal{P}_{alt}})(x,\xi)(A) = (2\lambda + \mu)f_1|\xi|^2 + (\lambda + \frac{\mu}{2} - 1)(*(\xi \wedge J_2 \xi \wedge \tilde{B}_{0,3} - \xi \wedge J_3 \xi \wedge \tilde{B}_{0,2}))\omega_1 +
$$

$$
(2\lambda + \mu)f_2|\xi|^2 + (\lambda + \frac{\mu}{2} - 1)(*(\xi \wedge J_3 \xi \wedge \tilde{B}_{0,1} - \xi \wedge J_1 \xi \wedge \tilde{B}_{0,3}))\omega_2 +
$$

$$
(2\lambda + \mu)f_3|\xi|^2 + (\lambda + \frac{\mu}{2} - 1)(*(\xi \wedge J_1 \xi \wedge \tilde{B}_{0,2} - \xi \wedge J_2 \xi \wedge \tilde{B}_{0,1}))\omega_3 + \cdots,
$$

where \cdots consists of the terms in $\Lambda^2_-(M)$. Together with [\(84\)](#page-29-0) we now have that

$$
\langle \sigma(L_{\mathcal{P}})(x,\xi)(A), A \rangle = 2|\xi|^2|B|^2 + 2|\xi|^2(2\lambda + \mu)(f_1^2 + f_2^2 + f_3^2) +
$$

\n
$$
2f_1(\lambda + \frac{\mu}{2} - 1)(*(\xi \wedge J_2\xi \wedge \tilde{B}_{0,3} - \xi \wedge J_3\xi \wedge \tilde{B}_{0,2})) +
$$

\n
$$
2f_2(\lambda + \frac{\mu}{2} - 1)(*(\xi \wedge J_3\xi \wedge \tilde{B}_{0,1} - \xi \wedge J_1\xi \wedge \tilde{B}_{0,3})) +
$$

\n
$$
2f_3(\lambda + \frac{\mu}{2} - 1)(*(\xi \wedge J_1\xi \wedge \tilde{B}_{0,2} - \xi \wedge J_2\xi \wedge \tilde{B}_{0,1}))
$$

\n
$$
= 2|\xi|^2|B|^2 + 2|\xi|^2(2\lambda + \mu)(f_1^2 + f_2^2 + f_3^2) +
$$

\n
$$
2(\lambda + \frac{\mu}{2} - 1)(*(\tilde{B}_{0,1} \wedge (f_2\xi \wedge J_3\xi - f_3\xi \wedge J_2\xi))) +
$$

\n
$$
2(\lambda + \frac{\mu}{2} - 1)(*(\tilde{B}_{0,2} \wedge (f_3\xi \wedge J_1\xi - f_1\xi \wedge J_3\xi))) +
$$

\n
$$
2(\lambda + \frac{\mu}{2} - 1)(*(\tilde{B}_{0,3} \wedge (f_1\xi \wedge J_2\xi - f_2\xi \wedge J_1\xi))).
$$

We see already that if $\lambda + \frac{\mu}{2} = 1$ then the flow is parabolic (in particular, this applies to the harmonic Ricci flow). Our goal is to find an optimal range of values for λ , μ so that the flow [\(83\)](#page-29-1) stays parabolic. Since SU(2) acts transitively on $S^3 \subset \mathbb{R}^4$, we can identify pointwise $(\omega_1, \omega_2, \omega_3)$ with the standard triple on \mathbb{R}^4 and ξ with $|\xi| dx_1$. Writing $\tilde{B}_{0,1} = \sum_i b_i \sigma_i$, where σ_i denote the standard anti-self-dual 2-forms on \mathbb{R}^4 , we have

$$
\begin{aligned}\n *(\tilde{B}_{0,1} \wedge (f_2 \xi \wedge J_3 \xi - f_3 \xi \wedge J_2 \xi)) &= *(|\xi|^2 (b_3 f_2 - b_2 f_3) dx_{1234}) \\
 &= |\xi|^2 (b_3 f_2 - b_2 f_3) \\
&\ge -|\xi|^2 \sqrt{b_2^2 + b_3^2} \sqrt{f_2^2 + f_3^2} \\
&\ge -|\xi|^2 \frac{|\tilde{B}_{0,1}|}{\sqrt{2}} \sqrt{f_2^2 + f_3^2} \\
&\ge -\frac{1}{2}|\xi|^2 \Big(\frac{|\tilde{B}_{0,1}|^2}{2\varepsilon} + \varepsilon (f_2^2 + f_3^2)\Big)\n\end{aligned}
$$

where we used Cauchy-Schwarz for the first inequality and Young's for the last one: $ab \leq (a^2/\varepsilon + \varepsilon b^2)/2$, for any $\varepsilon > 0$. An analogous argument holds for the other two terms, hence, using $|B_{0,i}|^2 = 2|\tilde{B}_{0,i}|^2$,

$$
\langle \sigma(L_{\mathcal{P}})(x,\xi)(A), A \rangle \ge 2|\xi|^2|B|^2 + 2|\xi|^2(2\lambda + \mu)(f_1^2 + f_2^2 + f_3^2)
$$

$$
-|\xi|^2|\lambda + \frac{\mu}{2} - 1|\left(\frac{|\tilde{B}_{0,1}|^2 + |\tilde{B}_{0,2}|^2 + |\tilde{B}_{0,3}|^2}{2\varepsilon} + 2\varepsilon(f_1^2 + f_2^2 + f_3^2)\right)
$$

$$
= 8|\xi|^2f_0^2 + \frac{1}{2}|\xi|^2(8 - \frac{1}{\varepsilon}|\lambda + \frac{\mu}{2} - 1|)\sum_{i=1}^3|\tilde{B}_{0,i}|^2 + 2|\xi|^2(2\lambda + \mu - \varepsilon|\lambda + \frac{\mu}{2} - 1|)\sum_{i=1}^3 f_i^2.
$$
 (85)

We see that the modified flow [\(83\)](#page-29-1) is parabolic for λ , μ such that $|x-1| < 8\varepsilon$ and $|x-1| < 2x/\varepsilon$, where $x := \lambda + \frac{\mu}{2}$ $\frac{\mu}{2}$. Observe that we still have the freedom to choose $\varepsilon > 0$ so as to maximise the range of values of x satisfying these inequalities. It is not hard to see that the minimum value of x is attained when $2x/\varepsilon = 8\varepsilon = 1 - x \Leftrightarrow \varepsilon = -1 + \sqrt{5}/2$, and the maximum when $2x/\varepsilon = 8\varepsilon = x - 1 \Leftrightarrow \varepsilon = 1 + \sqrt{5}/2$. The corresponding range of values of x is precisely as given in Theorem [5.4.](#page-28-1)

To conclude existence of the flow [\(80\)](#page-28-3), we argue as in DeTurck's trick: we define a 1-parameter family of diffeomorphisms $\{\varphi_t\}$ by [\(76\)](#page-27-3), where

$$
X_t = -(H^{-1}\delta G(H))^{\sharp} - \mu \sum_{i=1}^3 (J_i a_i)^{\sharp}
$$

and $\varphi_0 = \text{Id}$. Setting $\hat{\omega}(t) := \varphi_t^*(\omega(t))$ then gives a short-time solution to [\(80\)](#page-28-3), ie. the existence part of Theorem [5.4.](#page-28-1)

The proof of uniqueness is also similar to the Ricci case, but we need to proceed in two steps (see Remark [5.1](#page-31-1) below). The strategy is similar to the one employed in [\[DGK23\]](#page-39-2) for the G₂ case. Suppose that there are two solutions $\omega_1(t), \omega_2(t)$ to [\(80\)](#page-28-3), with $\omega_1(0) = \omega_2(0) = \omega(0)$. We define vector fields

$$
Y_u(t) := \mu \sum_{i=1}^3 ((J_i)_u (a_i)_u)^\sharp
$$
\n(86)

associated to $\omega_u(t)$, for $u = 1, 2$. It should be understood that $(a_i)_u$ correspond to the torsion forms associated to $\omega_u(t)$, and similarly for the almost complex structures $(J_i)_u$. We denote by $\{f_u(t)\}\$ the 1-parameter family of diffeomorphisms associated to $\{Y_u(t)\}\$, with $f_u(0) = \text{Id}$. Let $\bar{\omega}_u(t) := f_u(t)^* \omega_u(t)$, then we have

$$
\partial_t(\bar{\boldsymbol{\omega}}_u(t)) = f_u(t)^* \left((-2\mathrm{Ric}(g_u) - \mu \sum_{i=1}^3 \mathcal{L}_{((J_i)_u(a_i)_u)^{\sharp}} g_u + \lambda \sum_{i=1}^3 \mathrm{div}((a_i)_u)(\omega_i)_u) \diamond \boldsymbol{\omega}_u + \mathcal{L}_{Y_u(t)} \boldsymbol{\omega}_u \right)
$$

$$
= f_u(t)^* \left((-2\mathrm{Ric}(g_u) + \lambda \sum_{i=1}^3 \mathrm{div}((a_i)_u)(\omega_i)_u - d(\mu \sum_{i=1}^3 (J_i)_u(a_i)_u)) \diamond \omega_u \right)
$$

$$
= \left(-2\mathrm{Ric}(\bar{g}_u) + \lambda \sum_{i=1}^3 \mathrm{div}((\bar{a}_i)_u)(\bar{\omega}_i)_u - d(\mu \sum_{i=1}^3 (\bar{J}_i)_u(\bar{a}_i)_u) \right) \diamond \bar{\omega}_u,
$$

where we used Lemma [2.6](#page-5-8) for the second equality. Observe that the symmetric part of the latter flow is precisely the Ricci flow. Thus, following [\[CK04,](#page-39-14) §4.1], we can define another 1-parameter of diffeomorphisms $\{\phi_u(t)\}\$ by the harmonic map heat flow

$$
\partial_t \phi_u(t) = \Delta_{\bar{g}_u(t), H} \phi_u(t)
$$
, with $\phi_1(0) = \phi_2(0) = \text{Id}$,

where we recall that H denotes a fixed background metric. Letting $\hat{\omega}_u(t) := (\phi_u(t)^{-1})^* \bar{\omega}_u(t)$, and computing as before,

$$
\partial_t(\hat{\omega}_u(t)) = (\phi_u(t)^{-1})^* \Big(\left(-2\mathrm{Ric}(\bar{g}_u) + \lambda \sum_{i=1}^3 \mathrm{div}((\bar{a}_i)_u)(\bar{\omega}_i)_u - d \left(\mu \sum_{i=1}^3 (\bar{J}_i)_u(\bar{a}_i)_u \right) \right) \diamond \bar{\omega}_u \Big) + \mathcal{L}_{(H^{-1}\hat{\delta}G(H))^{\sharp}} \hat{\omega}_u
$$

\n
$$
= \left(-2\mathrm{Ric}(\hat{g}_u) + \lambda \sum_{i=1}^3 \mathrm{div}((\hat{a}_i)_u)(\hat{\omega}_i)_u - d \left(\mu \sum_{i=1}^3 (\hat{J}_i)_u(\hat{a}_i)_u \right) \right) \diamond \hat{\omega}_u + \mathcal{L}_{(H^{-1}\hat{\delta}G(H))^{\sharp}} \hat{\omega}_u
$$

\n
$$
= \left(-2\mathrm{Ric}(\hat{g}_u) + \mathcal{L}_{(H^{-1}\hat{\delta}G(H))^{\sharp}} \hat{g}_u + \lambda \sum_{i=1}^3 \mathrm{div}((\hat{a}_i)_u)(\hat{\omega}_i)_u - d \left(H^{-1}\hat{\delta}G(H) + \mu \sum_{i=1}^3 (\hat{J}_i)_u(\hat{a}_i)_u \right) \right) \diamond \hat{\omega}_u,
$$

where the first equality follows from the relation between the harmonic map flow and the Ricci flow, see [\[CK04,](#page-39-14) Lemma 3.27], and the last one is from Lemma [21.](#page-5-6) Thus, we see that $\hat{\omega}_1(t)$ and $\hat{\omega}_2(t)$ are both solutions to [\(83\)](#page-29-1) with the same initial data. But we already established above that [\(83\)](#page-29-1) is parabolic, hence by uniqueness $\hat{\omega}_1(t) = \hat{\omega}_2(t)$. Since $\hat{g}_1(t) = \hat{g}_2(t)$, it also follows that $\phi_1(t) = \phi_2(t)$ from the relation between the harmonic map heat flow and the Ricci flow, see [\[CK04,](#page-39-14) §4.4], and so $\bar{\omega}_1(t) = \bar{\omega}_2(t)$. Thus, to conclude, we only need to show that $f_1(t) = f_2(t)$. Since

$$
0 = \partial_t(f_u(t) \circ f_u(t)^{-1}(p))
$$

= $\partial_t(f_u(t)) \circ f_u(t)^{-1}(p) + (f_u(t))_* \partial_t(f_u(t)^{-1}(p))$
= $Y(t)(p) + (f_u(t))_* \partial_t(f_u(t)^{-1}(p)),$

we have

$$
\partial_t(f_u(t)^{-1}(p)) = -(f_u(t)^{-1})_* Y(t)(p) = -\mu \sum_{i=1}^3 ((\bar{J}_i)_u(\bar{a}_i)_u)^{\sharp} \circ f_u(t)^{-1}(p)
$$

i.e. $f_u(t)^{-1}$ is the diffeomorphism generated by the vector field $-\mu((\bar{J}_i)_u(\bar{a}_i)_u)^{\sharp}$. As $\bar{\omega}_1(t) = \bar{\omega}_2(t)$ and $f_1(0) = f_2(0) = \text{Id}$, the result follows by uniqueness. This concludes the proof of Theorem [5.4.](#page-28-1)

Remark 5.1. The point of defining two diffeomorphisms $f_u(t)$ and $\phi_u(t)$ in succession, rather than just one, is that we relied on the parabolicity of the harmonic map heat flow, cf. [\[ES64\]](#page-39-29), and also its relation to Ricci flow, cf. [\[CK04,](#page-39-14) Lemma 3.27]. Otherwise one would have to demonstrate parabolicity of a different flow of maps and establish a corresponding relation. This parallels Remark 6.82 of [\[DGK23\]](#page-39-2).

5.4 More general SU(2)-flows

We are interested in $SU(2)$ -flows which evolve, to second order (highest), by the invariants given in Theorem [4.2.](#page-20-1) This means that each f_i in [\(2\)](#page-1-4)-[\(4\)](#page-1-5) can be chosen as a linear combination of the 9 functions from Theorem [4.2–](#page-20-1)1., and each $B_{0,i}$ as a linear combination of the 12 terms in the space (isomorphic to) $\mathfrak{su}(2) \cong \mathbb{R}^3$ from Theorem [4.2–](#page-20-1)2.. This gives a $(4 \times 9 + 3 \times 12) = 72$ parameter family of SU(2)-flows. Deriving precise conditions for parabolicity in such generality is much too difficult; in this section we explain how one can nonetheless extend our results to a wider class of flows, by suitable modifications.

Modification by Lie derivative terms. A proof along the lines of $\S 5.3$ applies more generally to SU(2)-flows of the form

$$
\partial_t(\boldsymbol{\omega}) = (-2\mathrm{Ric}(g) - \sum_{i=0}^3 \sum_{j=1}^3 \mu_{ij} \mathcal{L}_{(J_i a_j)^{\sharp}} g + \sum_{j=1}^3 \lambda_j \operatorname{div}(a_j) \omega_j) \diamond \boldsymbol{\omega},\tag{87}
$$

where $\mu_{ij}, \lambda_j \in \mathbb{R}$, and we set $J_0 := \text{Id}$ for summation convenience. Denoting, for clarity, the vector field

$$
W := (H^{-1} \delta G(H))^{\sharp} + \sum_{i=0}^{3} \sum_{j=1}^{3} \mu_{ij} \mathcal{L}_{(J_i a_j)^{\sharp}},
$$

and again using Lemma [2.6](#page-5-8) for the second equality, we can consider the modified flow given by

$$
\partial_t(\omega) = \left(-2\mathrm{Ric}(g) - \sum_{i=0}^3 \sum_{j=1}^3 \mu_{ij} \mathcal{L}_{(J_i a_j)^{\sharp}} g + \sum_{j=1}^3 \lambda_j \operatorname{div}(a_j) \omega_j\right) \diamond \omega + \mathcal{L}_W \omega
$$

= $\left(-2\mathrm{Ric}(g) + \mathcal{L}_{(H^{-1}\delta G(H))^{\sharp}} g + \sum_{j=1}^3 \lambda_j \operatorname{div}(a_j) \omega_j - d(H^{-1}\delta G(H) + \sum_{i=0}^3 \sum_{j=1}^3 \mu_{ij} J_i a_j)\right) \diamond \omega,$

whose symmetric part is precisely given by the Ricci DeTurck flow. Thus, it suffices to compute the principal symbol of the skew-symmetric part and extract a parabolicity range for the parameters. The calculation at this level generality is still rather cumbersome, but the reader might have specific cases of interest which could be easier to work out. It should be clear that if the μ_{ij} and λ_i are sufficiently close to the flow of Theorem [5.4,](#page-28-1) then the modified flow of [\(87\)](#page-31-2) will also exist, as invertibility of the principal symbol is an open condition. The solution to the original flow is obtained by pulling back by the corresponding diffeomorphism as we did before. The proof of uniqueness also extends to this situation.

Negative gradient flow of the weighted Dirichlet energy functional. In \S 5.4, we established the existence and uniqueness of the negative gradient flow of the functional [\(44\)](#page-11-1). We shall now show how one can extend this to the weighted functional $E_{\lambda}(\omega)$ defined by [\(57\)](#page-15-2). From Proposition [3.9](#page-15-0) and Remark [3.1,](#page-13-1) it is easy to see that its first variation formula is

$$
\frac{d}{dt}E_{\lambda}(\omega) = \int_{M} g\Big(B, \lambda_{1}\left(-\frac{1}{2}g(da_{1}, \omega_{1})g + P_{1}(\pi_{-}^{2}(da_{1})) - 2P_{2}(S_{0,3}^{2}(\nabla a_{1}) \diamond \omega_{3}) + 2P_{3}(S_{0,2}^{2}(\nabla a_{1}) \diamond \omega_{2})\right) \n+ \lambda_{2}\left(-\frac{1}{2}g(da_{2}, \omega_{2})g + P_{2}(\pi_{-}^{2}(da_{2})) - 2P_{3}(S_{0,1}^{2}(\nabla a_{2}) \diamond \omega_{1}) + 2P_{1}(S_{0,3}^{2}(\nabla a_{2}) \diamond \omega_{3})\right) \n+ \lambda_{3}\left(-\frac{1}{2}g(da_{3}, \omega_{3})g + P_{3}(\pi_{-}^{2}(da_{3})) - 2P_{1}(S_{0,2}^{2}(\nabla a_{3}) \diamond \omega_{2}) + 2P_{2}(S_{0,1}^{2}(\nabla a_{3}) \diamond \omega_{1})\right) \text{vol} \n- 2\sum_{i=1}^{3} \int_{M} g\Big(C, \lambda_{i} \operatorname{div}(a_{i})\omega_{i}\Big) \text{vol} + \text{l.o.t}(\omega).
$$

Writing $c_i = \lambda_i^{-1}$, the corresponding negative gradient flow is, to highest order, given by

$$
\partial_t(\omega) = \left(-2\text{Ric}(g) - \mu \sum_{i=1}^3 \mathcal{L}_{(J_ia_i)^{\sharp}} g + \lambda \sum_{i=1}^3 \text{div}(a_i) \omega_i + \sum_{i=1}^3 \left(\frac{c_i}{2} + \frac{\mu}{2} - 1\right) g(da_i, \omega_i)g
$$

+ $P_1((2 - c_1 - \mu)\pi^2 (da_1) + 2(c_3 - \mu) S_{0,2}^2(\nabla a_3) \diamond \omega_2 - 2(c_2 - \mu) S_{0,3}^2(\nabla a_2) \diamond \omega_3)$
+ $P_2((2 - c_2 - \mu)\pi^2 (da_2) + 2(c_1 - \mu) S_{0,3}^2(\nabla a_1) \diamond \omega_3 - 2(c_3 - \mu) S_{0,1}^2(\nabla a_3) \diamond \omega_1)$
+ $P_3((2 - c_3 - \mu)\pi^2 (da_3) + 2(c_2 - \mu) S_{0,1}^2(\nabla a_2) \diamond \omega_1 - 2(c_1 - \mu) S_{0,k}^2(\nabla a_1) \diamond \omega_2)$
+ $\sum_{i=1}^3 \left(\frac{c_i}{2} - \lambda\right) \text{div}(a_i) \omega_i\right) \diamond \omega =: \mathcal{P}(\omega).$ (88)

Observe that, when $\mu = c_1 = c_2 = c_3 = 1$ and $\lambda = 1/2$, we indeed recover the negative gradient flow of [\(44\)](#page-11-1), as discussed in [§5.3.](#page-28-0) Our goal here is to identify a range of values for the constants c_i such that this flow remains parabolic. To this end we set

$$
L_1 := \sup_i \{|2 - \mu - c_i|\}, \quad L_2 := \sup_i \{|\mu - c_i|\}, \quad \text{and} \quad L_3 := \sup_i \{|\frac{c_i}{2} - \lambda|\}.
$$

Using the principal symbols given in Appendix [A,](#page-38-0) alongside the Cauchy-Schwarz and Young inequalities, one easily obtains: Lemma 5.6. *The following bounds hold*

$$
|\langle \sigma(L_{g(da_1,\omega_1)g})(x,\xi)(A),A\rangle| \leq |\xi|^2(10f_0^2 + 2\sum_{i=1}^3 |\tilde{B}_{0,i}|^2),
$$

$$
|\langle \sigma(L_{P_1(\pi^2_{-}(da_1))})(x,\xi)(A),A\rangle| \leq |\xi|^2(f_0^2 + 4|\tilde{B}_{0,1}|^2 + |\tilde{B}_{0,2}|^2 + |\tilde{B}_{0,3}|^2),
$$

$$
|\langle \sigma(L_{P_1(S^2_{0,2}(\nabla a_3)\circ\omega_2)})(x,\xi)(A),A\rangle| \leq |\xi|^2(\frac{1}{2}f_0^2 + f_3^2 + 3|\tilde{B}_{0,1}|^2 + \frac{1}{2}(|\tilde{B}_{0,2}|^2 + |\tilde{B}_{0,3}|^2))
$$

$$
|\langle \sigma(L_{P_1(S^2_{0,3}(\nabla a_2)\circ\omega_3)})(x,\xi)(A),A\rangle| \leq |\xi|^2(\frac{1}{2}f_0^2 + f_2^2 + 3|\tilde{B}_{0,1}|^2 + \frac{1}{2}(|\tilde{B}_{0,2}|^2 + |\tilde{B}_{0,3}|^2))
$$

$$
|\langle \sigma(L_{\text{div}(a_1)\omega_1})(x,\xi)(A),A\rangle| \leq |\xi|^2(6f_1^2 + |\tilde{B}_{0,2}|^2 + |\tilde{B}_{0,3}|^2),
$$

and similar expressions hold for cyclic permutation of the indices.

Proof. We prove the first one, to establish the procedure; the rest follow by analogous computations.

$$
\begin{split} |\langle \sigma(L_{g(da_1,\omega_1)g})(x,\xi)(A),A\rangle| &= 4|f_0^2|\xi|^2 - f_0g(\xi \wedge J_1\xi, \tilde{B}_{0,1}) + f_0g(\xi \wedge J_2\xi, \tilde{B}_{0,2}) + f_0g(\xi \wedge J_3\xi, \tilde{B}_{0,3})| \\ &\leq 4f_0^2|\xi|^2 + 4|\xi|^2(|f_0||\tilde{B}_{0,1}| + |f_0||\tilde{B}_{0,2}| + |f_0||\tilde{B}_{0,3}|) \\ &\leq 4f_0^2|\xi|^2 + 2|\xi|^2(3f_0^2 + |\tilde{B}_{0,1}|^2 + |\tilde{B}_{0,2}|^2 + |\tilde{B}_{0,3}|^2) \\ &= |\xi|^2(10f_0^2 + 2\sum_{i=1}^3 |\tilde{B}_{0,i}|^2). \end{split}
$$

As in the proof of Theorem [5.4,](#page-28-1) we can apply the modified DeTurck trick to [\(88\)](#page-32-0), using the same vector field. Computing as in \S 5.3, using the above estimates and (85) , we find

$$
\langle \sigma(L_{\mathcal{P}})(x,\xi)(A), A \rangle \ge |\xi|^2 f_0^2 (8 - 18L_1 - 6L_2) +
$$

$$
|\xi|^2 (4 - \frac{1}{2\varepsilon}|\lambda + \frac{\mu}{2} - 1| - 9L_1 - 16L_2 - 2L_3) \sum_{i=1}^3 |\tilde{B}_{0,i}|^2 +
$$

$$
|\xi|^2 (2(2\lambda + \mu - \varepsilon|\lambda + \frac{\mu}{2} - 1|) - 4L_2 - 6L_3) \sum_{i=1}^3 f_i^2.
$$

In particular, setting $\mu = 1$ and $\lambda = 1/2$, we see that if c_i are such $9L_1 + 3L_2 < 4$, $9L_1 + 16L_2 + 2L_3 < 4$ and $2L_2 + 3L_3 < 2$, then the flow [\(88\)](#page-32-0) exists for short time. Note that the range of values here are not necessarily sharp.

Weyl curvature terms decrease parabolicity. Consider the flow given by

$$
\partial_t(\boldsymbol{\omega}) = (\nu g(da_1, \omega_2)g - 2\mathrm{Ric}(g) - \mu \sum_{i=1}^3 \mathcal{L}_{(J_i a_i)^{\sharp}} g + \lambda \sum_{i=1}^3 \mathrm{div}(a_i) \omega_i) \diamond \boldsymbol{\omega},\tag{89}
$$

with $\nu, \mu, \lambda \in \mathbb{R}$. This is clearly a modified version of [\(80\)](#page-28-3) by a 'self-dual Weyl-type' term $g(da_1, \omega_2)$, in the terminology of Proposition [4.7.](#page-23-1)

From Appendix [A,](#page-38-0) we have

$$
\langle \sigma(L_{g(da_1,\omega_2)})(x,\xi)(A)g,A\rangle = -4f_0(g(\xi \wedge J_2\xi,\tilde{B}_{0,1}) + g(\xi \wedge J_1\xi,\tilde{B}_{0,2})).
$$

Observe that this term does not come with a 'sign', ie. the flow [\(89\)](#page-33-0) will exist for a smaller range of values than that in Theorem [5.4.](#page-28-1) To illustrate this more explicitly, we again start by using Cauchy-Schwarz and Young's inequalities:

$$
\begin{aligned} |\langle \sigma(L_{g(da_1,\omega_2)})(x,\xi)(A)g,A\rangle| &= 4|f_0||g(\xi \wedge J_2\xi, \tilde{B}_{0,1}) + g(\xi \wedge J_1\xi, \tilde{B}_{0,2})| \\ &\le 4|\xi|^2|f_0|(|\tilde{B}_{0,1}| + |\tilde{B}_{0,2}|) \\ &\le 2|\xi|^2(2f_0^2 + (|\tilde{B}_{0,1}|^2 + |\tilde{B}_{0,2}|^2)). \end{aligned}
$$

Repeating the calculations in the proof of Theorem [5.4](#page-28-1) for the flow [\(89\)](#page-33-0), we obtain, instead of [\(85\)](#page-30-0),

$$
\langle \sigma(L_{\mathcal{P}})(x,\xi)(A), A \rangle \ge 2|\xi|^2|B|^2 + 2|\xi|^2(2\lambda + \mu)(f_1^2 + f_2^2 + f_3^2)
$$

\n
$$
-|\xi|^2|\lambda + \frac{\mu}{2} - 1|\left(\frac{|\tilde{B}_{0,1}|^2 + |\tilde{B}_{0,2}|^2 + |\tilde{B}_{0,3}|^2}{2\varepsilon} + 2\varepsilon(f_1^2 + f_2^2 + f_3^2)\right)
$$

\n
$$
-2|\nu||\xi|^2(2f_0^2 + (|\tilde{B}_{0,1}|^2 + |\tilde{B}_{0,2}|^2))
$$

\n
$$
= (8 - 4|\nu|)|\xi|^2 f_0^2 + \frac{1}{2}|\xi|^2(8 - \frac{1}{\varepsilon}|\lambda + \frac{\mu}{2} - 1| - 4|\nu|)\sum_{i=1}^2 |\tilde{B}_{0,i}|^2
$$

\n
$$
+ \frac{1}{2}|\xi|^2(8 - \frac{1}{\varepsilon}|\lambda + \frac{\mu}{2} - 1|)|\tilde{B}_{0,3}|^2 + 2|\xi|^2(2\lambda + \mu - \varepsilon|\lambda + \frac{\mu}{2} - 1|)\sum_{i=1}^3 f_i^2.
$$

In order to get parabolicity, we now need $|\nu| < 2$, $|x-1| < (8-4|\nu|)\varepsilon$ and $|x-1| < 2x/\varepsilon$, where $x := \lambda + \frac{\mu}{2}$ $\frac{\mu}{2}$. Comparing with the proof of Theorem [5.4,](#page-28-1) we see that the range of values for $\lambda + \frac{\mu}{2}$ will be smaller than that of Theorem [5.4.](#page-28-1) This is in fact a general feature: the Weyl curvature terms reduce parabolicity, since unlike the Ricci tensor they lack a Laplacian term, cf. their principal symbols given in Appendix [A.](#page-38-0) Thus, addition of the SU(2)-invariants of the form $g(da_i, \omega_j)$, for $i < j$, always makes the family of weighted flows 'less parabolic', in this precise sense. The crucial conclusion is that we need the Ricci curvature term to dominate the Weyl terms, in order to ensure short-time existence.

By comparison to the G_2 case, the reader can observe in $[DGK23,$ Theorem 6.76] that the Weyl curvature term, denoted by F therein, reduces the parabolicity range; this is precisely the feature we described above. Unlike the G_2 case, however, there is here also an issue of multiplicity. To explain this, consider a flow given by

$$
\partial_t(\boldsymbol{\omega}) = (\nu g(da_1, \omega_2)\omega_1 - 2\text{Ric}(g) - \mu \sum_{i=1}^3 \mathcal{L}_{(J_ia_i)^{\sharp}}g + \lambda \sum_{i=1}^3 \text{div}(a_i)\omega_i) \diamond \boldsymbol{\omega}.
$$
\n(90)

Both in [\(89\)](#page-33-0) and [\(90\)](#page-34-1), we are modifying the flow [\(80\)](#page-28-3) by the same term $g(da_1, \omega_2)$, but the resulting flows are very different. Similarly, adding either $S_{0,1}^2(\nabla a_1)$ or $P_2(S_{0,1}^2(\nabla a_1)\diamond\omega_1)$ results in different flows, however much these terms are equivalent as SU(2)-invariants. This type of phenomenon occurs because $S^2(M) \oplus \mathfrak{su}(2)^\perp \cong 4\mathbb{R} \oplus 3\mathbb{R}^3$ contains multiple copies of some SU(2)-modules; this also occurs in the SU(n) and Sp(n) cases, for $n > 1$, but not in the G₂ and Spin(7) cases, see [§6.1](#page-35-1) below. The point here is that flow families of the forms [\(89\)](#page-33-0) and [\(90\)](#page-34-1), although built from the same second-order invariants, will have different parabolicity ranges.

5.5 Solitons

In this section we define the notion of solitons for the $SU(2)$ -we considered in [§5.3,](#page-28-0) while leaving a more detailed study of those solutions as a suggestion for future investigations. We begin by considering a general $SU(2)$ -flow of the form

$$
\partial_t(\boldsymbol{\omega}) = \mathcal{P}(\boldsymbol{\omega}),\tag{91}
$$

where P denotes a *diffeomorphism-invariant* partial differential operator, ie.

$$
\varphi^* \mathcal{P}(\boldsymbol{\omega}) = \mathcal{P}(\varphi^* \boldsymbol{\omega}), \quad \forall \varphi \in \text{Diff}(M),
$$

and which is also *homothetic scaling-invariant*, ie. $\mathcal{P}(c^2\omega) = \mathcal{P}(\omega)$, for $c \in \mathbb{R}$. Note that in particular the Ricci map $g \mapsto \text{Ric}(g)$ satisfies these two properties. Suppose now that ω_0 is a solution to

$$
\mathcal{P}(\omega_0) = \mathcal{L}_Y \omega_0 - \nu \omega_0, \tag{92}
$$

for some vector field Y and $\nu \in \mathbb{R}$. Let $\omega(t) := (1 - \nu t) \varphi_t^* \omega_0$, $X(t) := (1 - \nu t)^{-1} Y$ and denote by φ_t the 1-parameter family of diffeomorphisms generated by $X(t)$ as in [\(76\)](#page-27-3), then we have

$$
\partial_t \omega(t) = -\nu \varphi_t^* \omega_0 + (1 - \nu t) \varphi_t^* (\mathcal{L}_{X(t)} \omega_0)
$$

= $\varphi^* (\mathcal{P}(\omega_0))$
= $\mathcal{P}(\omega(t)).$

Thus, $\omega(t)$ is a solution to [\(91\)](#page-34-2) which evolves only by the symmetries of the flow. Mimicking the definition of Ricci soliton cf. [\[Top06,](#page-39-18) Definition 1.2.2], we define:

Definition 5.7. The data (ω_0, Y, ν) satisfying [\(92\)](#page-34-3) is called a *shrinking*, *steady* or *expanding* soliton for the flow [\(91\)](#page-34-2) if $\nu < 0$, $\nu = 0$ or $\nu > 0$ respectively.

From Proposition [2.9,](#page-7-0) it is not hard to see that the torsion forms remain unchanged under the homothetic scaling $\tilde{\omega} := c^2 \omega$, i.e. $\tilde{a}_i = a_i$. Furthermore, the $a_i = a_i(\omega)$ are diffeomorphism-invariant; indeed the 'Bianchi-type identity' in Corollary [2.11](#page-8-1) is a consequence of this invariance, cf. [\[Kar09,](#page-39-9) §4.1]. Since $\tilde{J} \tilde{a}_i = J a_i$, we also have $(\tilde{J} \tilde{a}_i)^{\tilde{\sharp}} = c^{-2} (J a_i)^{\sharp}$, and similarly one can check that $\tilde{\delta} \tilde{a}_i = c^{-2} \delta a_i$. We deduce that the flow given by

$$
\partial_t(\boldsymbol{\omega}) = \left(-2\mathrm{Ric}(g) - \mu \sum_{i=1}^3 \mathcal{L}_{(J_i a_i)^{\sharp}} g + \lambda \sum_{i=1}^3 \mathrm{div}(a_i) \omega_i\right) \diamond \boldsymbol{\omega} + Q(\boldsymbol{a}),\tag{93}
$$

where $Q(a)$ denotes a function in a_i , satisfies the properties of [\(91\)](#page-34-2). This also applies to other flows, such as [\(87\)](#page-31-2). Note that in practice we will only require $Q(a)$ to be a quadratic polynomial in the torsion. Thus, a soliton of [\(93\)](#page-34-4) is a solution to [\(92\)](#page-34-3), where $\mathcal{P}(\omega)$ is the RHS of [\(93\)](#page-34-4). In particular, it would interesting to find solitons when

$$
\mathcal{P}(\boldsymbol{\omega}) = \left(\text{sym}(\text{div}(T^t)) + T \ast T - \frac{1}{2}|T|^2 g + \text{div}(T)\right) \diamond \boldsymbol{\omega}
$$

=
$$
\sum_{i} \left(-\text{sym}(\nabla_{J_i} a_i) - \sum_{j,k} \varepsilon_{ijk} a_i \odot J_j a_k + 2a_i \otimes a_i - |a_i|^2 g + \text{div}(a_i) \omega_i\right) \diamond \boldsymbol{\omega},
$$

corresponding to the negative gradient flow of [\(44\)](#page-11-1) and where the second equality follows from Proposition [3.5.](#page-12-0)

6 Discussion of results

6.1 $SU(3)$, G_2 and $Spin(7)$ second-order invariants revisited

We now provide several illustrations for our reading of Bryant's remarks, and we explain how this approach applies to general H-structures. We give an alternative proof of [\[DGK23,](#page-39-2) Theorem 6.2], which states that there are exactly 6 second order G_2 invariants that lead to geometric flows of the underlying G_2 -structure 3-form. Our approach differs from the one given in $[DGK23, §5]$ $[DGK23, §5]$, in that it uses the representation-theoretic method of [§4.2,](#page-19-2) following $[Bry05]$. In the Spin(7) case, we show that there are exactly 4 such terms; this was also recently determined in $Dw124$. In the case of SU(3)-structures, we use results in $[BVO7]$ to show that there are exactly 16 such terms, leading to a 19-parameter family of possible $SU(3)$ -flows, to second order; this doesn't seem to have been mentioned heretofore in the literature. In particular, we shall see that both the G_2 and $Spin(7)$ cases admit fewer geometric flows, compared to the $SU(2)$ case; this explains why a full classification of parabolic flows of $SU(2)$ -structures is so much more involved.

We first recall from [\[FLMSE22\]](#page-39-1) that, in order to define a geometric flow of an H-structure, with $H \subset SO(n)$, say determined by a (multi-)tensor ξ , we need to choose an endomorphism $A = A(\xi)$ and evolve the underlying tensor by

$$
\partial_t \xi = A \diamond \xi. \tag{94}
$$

Denoting the space of second-order invariants by $V_2(h)$, as in [§4.2,](#page-19-2) let us now restrict to cases in which A is at most second order in ξ, ie. $A \in V_2(\mathfrak{h})$, to highest order. Note that we also need the endomorphism A to lie in the complement of ker($\diamond \xi$), since H is the stabiliser of ξ by definition; we denote this space by

$$
\mathrm{End}(\mathbb{R}^n)_{\mathfrak{h}}:=\mathrm{End}(\mathbb{R}^n)/\mathfrak{h}\cong \Sigma^2(\mathbb{R}^n)\oplus \mathfrak{h}^\perp,
$$

so that the assumption is $A \in \text{End}(\mathbb{R}^n)_{\mathfrak{h}} \cap V_2(\mathfrak{h})$. To be precise, $V_2(\mathfrak{h}) \cap \text{End}(\mathbb{R}^n)$ refers to the subspace of $V_2(\mathfrak{h})$ consisting of those H-modules which also occur in $\text{End}(\mathbb{R}^n)_{\mathfrak{h}}$, *counted with their multiplicities*. This is precisely the approach we considered in this article, and we shall next apply it to the cases $H = G_2$ and Spin(7).

The G₂ case. The space $V_2(\mathfrak{g}_2)$ was computed in [\[Bry05,](#page-39-3) (4.7)]. From this we easily see that

$$
V_2(\mathfrak{g}_2) \cap \text{End}(\mathbb{R}^7)_{\mathfrak{g}_2} \cong \mathbb{R} \oplus 2\Lambda^1(\mathbb{R}^7) \oplus 3\Sigma_0^2(\mathbb{R}^7),\tag{95}
$$

where, in the notation of [\[Bry05\]](#page-39-3), $V_{0,0} \cong \mathbb{R}$, $V_{1,0} \cong \Lambda^1(\mathbb{R}^7)$ and $V_{2,0} \cong \Sigma_0^2(\mathbb{R}^7)$. Observe that there are exactly 6 irreducible G_2 -modules as expected from [\[DGK23,](#page-39-2) Theorem 6.2]. Moreover, as G_2 -modules, we also know from [\[Bry05,](#page-39-3) (4.8)] that

$$
V_2(\mathfrak{so}(7)) \cap \text{End}(\mathbb{R}^7)_{\mathfrak{g}_2} \cong \mathbb{R} \oplus 2\Sigma_0^2(\mathbb{R}^7). \tag{96}
$$

Hence the $\mathbb R$ component in $V_2(\mathfrak{g}_2) \cap (\text{End}(\mathbb{R}^7))_{\mathfrak{g}_2}$ is given by the scalar curvature, and two of the three components of $\Sigma_0^2(\mathbb{R}^7)$ by the traceless Ricci tensor and a Weyl curvature term W_{27} . Clearly the third term in $\Sigma_0^2(\mathbb{R}^7)$ has to be given by $\mathcal{L}_{\tau_1^{\sharp}}g$, since it does not lie in $V_2(\mathfrak{so}(7))$; here τ_1 denotes the G_2 torsion 1-form, as defined in [\[Bry05,](#page-39-3) (3.7)]. Among the two terms spanning $\Lambda^1(\mathbb{R}^7)$, one has to be given by the harmonic part div(T); compare with Proposition [2.12,](#page-8-3) see also [\[LSE23\]](#page-39-0). We should point out that, strictly speaking, the harmonic part lies in $\mathfrak{h}^{\perp} \subset \Lambda^2(\mathbb{R}^n)$, and in the G_2 case it so happens that

$$
\mathfrak{g}_2^{\perp} = \Lambda_7^2(\mathbb{R}^7) \cong \Lambda^1(\mathbb{R}^7),
$$

see eg. [\[Bry05,](#page-39-3) (2.14)]. The second term in $\Lambda^1(\mathbb{R}^7)$, denoted by $\text{div}(T^t)$ in [\[DGK23\]](#page-39-2), corresponds to other $\Lambda^1(\mathbb{R}^7)$ -component of ∇T , where T denotes the full intrinsic torsion. We now explain the appearance of this term. As described below for general H, the term $div(T^t)$ in fact belongs to space

$$
\Sigma^2(\mathbb{R}^7)\otimes\mathfrak{g}_2^\perp\cong2\Lambda^1(\mathbb{R}^7)\oplus\Sigma^2_0(\mathbb{R}^7)\oplus V_{1,1}\oplus V_{0,1}\oplus\oplus V_{3,0}.
$$

It follows that $div(T^t)$ is obtained by suitably tracing ∇T : whereas the trace on the first two indices yields the usual $div(T)$, the trace of the first and third indices gives precisely $div(T^t)$. Note that the occurrence of $\Sigma_0^2(\mathbb{R}^7)$ in the decomposition corresponds to the aforementioned $\mathcal{L}_{\tau_1^{\sharp}}g$ term. Since $\text{End}(\mathbb{R}^7)_{\mathfrak{g}_2} \cong \mathbb{R} \oplus \Lambda^1(\mathbb{R}^7) \oplus \Sigma_0^2(\mathbb{R}^7)$, it follows that there is a 6-parameter family of G₂-flows, to second order. Thus, specialising [\(94\)](#page-35-2) to flows of the 3-form $\xi = \varphi$, we get

$$
\partial_t(\varphi) = \left(\lambda_1 \text{Scal}(g)g + \lambda_2 \text{Ric}(g) + \lambda_3 W_{27} + \lambda_4 \mathcal{L}_{\tau_1^{\sharp}} g + \lambda_5 \operatorname{div}(T) + \lambda_6 \operatorname{div}(T^{\dagger})\right) \diamond \varphi. \tag{97}
$$

as the general G_2 -flow, to second order, up to lower order terms; this was indeed shown in [\[DGK23\]](#page-39-2).

Owing to the fact that Bryant had already computed the spaces $V_1(\mathfrak{g}_2)$ and $V_2(\mathfrak{g}_2)$, and introduced the Laplacian flow in $Bry05$, we believe that he was already aware of the existence of a 6-parameter family of G_2 -flows, albeit not explicitly mentioning it therein. The reason for singling out the Laplacian flow of closed G_2 -structures in this family is the fact that the underlying metric evolves by the Ricci flow (to highest order), see $[Bry05, (6.15)]$ $[Bry05, (6.15)]$, and the stationary points are precisely the torsion-free G_2 -structures. More generally, Bryant's results show that, for any flow of closed G_2 -structures, the metric can only evolve by the Ricci curvature.

The Spin(7) **case.** The space $V_2(\text{spin}(7))$ was computed in [\[Fow20,](#page-39-21) §4]. From this we easily see that

$$
V_2(\mathfrak{spin}(7)) \cap \text{End}(\mathbb{R}^8)_{\mathfrak{spin}(7)} \cong \mathbb{R} \oplus \mathfrak{spin}(7)^\perp \oplus 2\Sigma_0^2(\mathbb{R}^8),\tag{98}
$$

where, in their notation, $V_{0,0,0} \cong \mathbb{R}$, $V_{1,0,0} \cong \mathfrak{spin}(7)^{\perp} \cong \mathbb{R}^7$ and $V_{0,0,2} \cong \Sigma_0^2(\mathbb{R}^8)$. There are exactly 4 irreducible $Spin(7)$ -modules, which is consistent with $Dwi24$. On the other hand, it is also easy to show that, as $Spin(7)$ -modules,

$$
V_2(\mathfrak{so}(8)) \cap \text{End}(\mathbb{R}^8)_{\mathfrak{spin}(7)} \cong \mathbb{R} \oplus \Sigma_0^2(\mathbb{R}^8). \tag{99}
$$

This implies that the $\mathbb R$ component in $V_2(\text{spin}(7)) \cap (\text{End}(\mathbb{R}^8))_{\text{spin}(7)}$ is again given by the scalar curvature, and one of the two $\Sigma_0^2(\mathbb{R}^7)$ components is given by the traceless Ricci tensor. Clearly the second term in $\Sigma_0^2(\mathbb{R}^8)$ has to be given by $\mathcal{L}_{T_s^1}$ #g, since it does not lie in $V_2(\mathfrak{so}(8))$; here T_8^1 denotes the $Spin(7)$ torsion 1-form, as defined in [\[Fow20,](#page-39-21) (2.5)]. The remaining term in $\text{spin}(7)^{\perp}$ is again just the harmonic term $\text{div}(T)$, where T denotes the $\text{Spin}(7)$ intrinsic torsion. This exhausts all the secondorder invariants which can be used to evolve the Spin(7)-structure 4-form. Since $\text{End}(\mathbb{R}^8)_{\text{spin}(7)} \cong \mathbb{R} \oplus \text{spin}(7)^\perp \oplus \Sigma_0^2(\mathbb{R}^7)$, we conclude that there is a 4-parameter family of $Spin(7)$ flows, to second order. Explicitly, specialising [\(94\)](#page-35-2) to flows of the 4-form $\xi = \Phi$, we get

$$
\partial_t(\Phi) = (\lambda_1 \text{Scal}(g)g + \lambda_2 \text{Ric}(g) + \lambda_3 \mathcal{L}_{T_8^{\perp \sharp}} g + \lambda_4 \operatorname{div}(T)) \diamond \Phi.
$$
\n(100)

We observe that, while there is exactly one Weyl curvature term that can occur in a G_2 -flow, none at all can occur in the Spin(7) case. By further contrast, we saw in the $SU(2)$ case that all of the 5 components of W^+ can occur.

The general H case. Given $H \subset SO(n)$, as in Berger's list [\[Ber55\]](#page-39-31), we always have

$$
\mathbb{R} \oplus \mathfrak{h}^{\perp} \oplus 2\Sigma_0^2(\mathbb{R}^n) \subset V_2(\mathfrak{h}) \cap \text{End}(\mathbb{R}^n)_{\mathfrak{h}},\tag{101}
$$

where, as before, the $\mathbb R$ component corresponds to the scalar curvature, and one of the two $\Sigma_0^2(\mathbb R^n)$ components is given by the traceless Ricci tensor. The remaining term in $\Sigma_0^2(\mathbb{R}^n)$ is of the form $\mathcal{L}_{\tau^{\sharp}}g$, where τ is a 1-form component in the space of intrinsic torsion $V_1(\mathfrak{h})$, sometimes called the *Lee form* in the literature, eg. $\tau = \sum_{i=1}^3 J_i a_i$, for $H = SU(2)$. The \mathfrak{h}^{\perp} component corresponds as before to the harmonic term, cf. [\[LSE23\]](#page-39-0). Note that we are not assuming that these are irreducible as H-modules, which will in general depend on the group H . In particular, we see that the $Spin(7)$ setup actually realises the minimal instance of [\(101\)](#page-36-0). Moreover, observe that the second-order invariants that arise from the curvature tensor lie in the sub-module

$$
V_2(\mathfrak{so}(n)) \cong \mathbb{R} \oplus \Sigma_0^2(\mathbb{R}^n) \oplus W \subset V_2(\mathfrak{h}),
$$

where W denotes the space of Weyl curvature terms, and furthermore, from the proof of (61) , we know that the remaining ones lie in

$$
\Sigma^2(\mathbb{R}^n) \otimes \mathfrak{h}^\perp \cong \Sigma^2_0(\mathbb{R}^n) \otimes \mathfrak{h}^\perp \oplus \mathfrak{h}^\perp,
$$

where of course the harmonic term $div(T)$ lies in the last summand. This provides an efficient method to extract the secondorder invariants, seeing as only these arise independently from the curvature and sym(∇T) $\diamond \xi$. The number of possible second-order flow to highest order can then be deduced from the multiplicity of each irreducible H-module in $V_2(\mathfrak{h})$ which also appears in $\text{End}(\mathbb{R}^n)_{\mathfrak{h}}$.

In the cases $H = SU(n)$ and $Sp(n)$, there are many other terms aside from those in [\(101\)](#page-36-0), as already illustrated by our results for $H = SU(2) = Sp(1)$. Another distinctive feature of the SU(2) case, relative to G_2 or Spin(7), is the fact that there occur H-modules with repeated multiplicities in $\text{End}(\mathbb{R}^n)_{\mathfrak{h}}$, which is why we had to carefully distinguish between the spaces $\Sigma_{0,1}^2$, $\Sigma_{0,2}^2$, $\Sigma_{0,3}^2$ and Λ_{-}^2 , and likewise for g , ω_1 , ω_2 and ω_3 . This indeed explains why there are so many possible flows in this context, which also occurs in the $SU(n)$ and $Sp(n)$ cases. For instance, we illustrate the $SU(3)$ setup next as a novel application of our approach.

The SU(3) case. The space $V_2(\mathfrak{su}(3))$ was computed in [\[BV07,](#page-39-30) §2.6]. From this we easily see that

$$
V_2(\mathfrak{su}(3)) \cap \text{End}(\mathbb{R}^6)_{\mathfrak{su}(3)} \cong 3\mathbb{R} \oplus 4\Lambda^1(\mathbb{R}^6) \oplus 5S_+^2(\mathbb{R}^6) \oplus 4S_-^2(\mathbb{R}^6),\tag{102}
$$

where $\Sigma_0^2(\mathbb{R}^6) = S_+^2(\mathbb{R}^6) \oplus S_-^2(\mathbb{R}^6)$, as irreducible SU(3)-modules of dimensions 8 and 12, respectively [\[BV07,](#page-39-30) §2.3]. In the above decomposition we used the identifications

$$
\mathfrak{su}(3)^{\perp} \cong \mathbb{R} \oplus \Lambda^{1}(\mathbb{R}^{6}) \quad \text{and} \quad \mathfrak{su}(3) \cong S^{2}_{+}(\mathbb{R}^{6}).
$$

Thus, there are a total 16 possible second-order $SU(3)$ -invariants which can be used to evolve an $SU(3)$ -structure. On the other hand, as $SO(6)$ -modules, it is not hard to compute that

$$
V_2(\mathfrak{so}(6)) \cong \mathbb{R} \oplus \Sigma_0^2(\mathbb{R}^6) \oplus V_{0,2,2},
$$
\n(103)

where $V_{0,2,2}$ is the irreducible SO(6)-module of dimension 84, corresponding to the Weyl curvature. One still has to check whether any components of the Weyl term occur in $V_2(\mathfrak{su}(3)) \cap \text{End}(\mathbb{R}^6)$, and indeed a calculation shows that, as SU(3)modules,

$$
V_{0,2,2} \cap \text{End}(\mathbb{R}^6)_{\mathfrak{su}(3)} \cong \mathbb{R} \oplus \Lambda^1(\mathbb{R}^6) \oplus S^2_+(\mathbb{R}^6) \oplus S^2_-(\mathbb{R}^6).
$$

Thus, among the 16 second-order SU(3)-invariants, precisely 7 can be computed from the Riemann curvature tensor. Since

$$
\mathrm{End}(\mathbb{R}^6)_{\mathfrak{su}(3)}\cong 2\mathbb{R}\oplus \Lambda^1(\mathbb{R}^6)\oplus S^2_+(\mathbb{R}^6)\oplus S^2_-(\mathbb{R}^6),
$$

note that there is a $(2 \times 3 + 4 + 5 + 4) = 19$ -parameter family of SU(3)-flows, to second order. Geometric SU(3)-flows arising from this framework will be investigated in a future work.

6.2 Quaternionic bundle

In [\[Gri17\]](#page-39-4), Grigorian derived the notion of harmonic G_2 -structures independently of [\[DGK21,](#page-39-28) [LSE23\]](#page-39-0)), by interpreting the intrinsic torsion as a connection on a rank 8 octonionic vector bundle $\mathbb{O}M$, whose natural 'Coulomb gauge' condition is equivalent to gweometric harmonincity. In our $SU(2)$ case, we can define bya analogy a *quaternionic bundle* over $M⁴$ as follows.

We consider the rank 4 vector bundle $\mathbb{H}M := \Lambda^0 \oplus \Lambda^2_+$, and define a quaternionic product on sections by

$$
(a, \alpha) \cdot (b, \beta) := (ab - \frac{1}{2}g(\alpha, \beta), a\beta + b\alpha + \alpha \diamond \beta),
$$
\n(104)

corresponding to the usual multiplication on \mathbb{H} , whereby $\text{Im}(\mathbb{H}) \simeq \Lambda^2_+(M)$, see Lemma [2.8.](#page-6-2) We define an inner product on $\mathbb{H}M$ by

$$
\langle (a,\alpha), (b,\beta) \rangle := ab + \frac{1}{2}g(\alpha,\beta),\tag{105}
$$

where the half factor stems from our convention $g(\omega_i, \omega_j) = 2\delta_{ij}$. Since the operator \diamond only depends on the metric, it is covariantly constant and hence:

Proposition 6.1. *The induced covariant derivative on* HM *is compatible with the quaternionic product, ie.*

$$
\nabla_X((a,\alpha)\cdot(b,\beta)) = \nabla_X((a,\alpha))\cdot(b,\beta) + (a,\alpha)\cdot\nabla_X((b,\beta)),\tag{106}
$$

where $\nabla_X((c,\gamma)) := (\nabla_X c, \nabla_X \gamma)$ *.*

Comparing with $[Gri17,$ Proposition 6.1], we see that in the G_2 case the above compatibility does not hold for the octonionic bundle $\mathbb{O}M$, and in fact a term involving the intrinsic torsion arises. This seems to be due to the fact that, exceptionally in dimension 4, the Levi-Civita connection preserves the decomposition $\Lambda^2_+ \oplus \Lambda^2_-$ without any holonomy constraint. In [\[Gri17\]](#page-39-4), the octonionic bundle $\mathbb{O}M$ is defined as $\Lambda^0 \oplus \Lambda_7^1$, and the appearance of Λ_7^1 stems from the G₂-equivariant isomorphism identifying it with $\Lambda^2_7 \cong \mathfrak{g}_2^{\perp}$. Such an identification does not hold in general (certainly not in the SU(2) case), so for general H-structures one should really consider \mathfrak{h}^{\perp} . In particular, it would be interesting to investigate this perspective on harmonic H-structures for the quaternionic structures $\text{Sp}(n)$ and $\text{Sp}(n)\text{Sp}(1)$, $n > 1$.

6.3 Ongoing and future work

We briefly discuss some natural open questions related to our results, which might be object of investigation by the authors and the wider community in the immediate future.

- 1. One important issue which we have overlooked in this paper is the choice of lower-order terms in the flows. While for short-time existence one can restrict attention to the highest-order terms, this says nothing about long-time existence, let alone convergence. Neither did we address what the stationary points of the flows actually are expected to be; one should be able to find suitable expressions involving terms in $V_2(\mathfrak{su}(2))$ and $S^2(V_1(\mathfrak{su}(2)))$ such that the stationary points have special torsion – most optimistically no torsion at all.
- 2. Our results suggest that the most interesting flows to consider are the harmonic Ricci flow and the negative gradient flow the Dirichlet energy functional [\(44\)](#page-11-1). In both cases, it is still unclear whether there are more critical points, aside from the torsion-free ones. Moreover, one still needs to develop a long-time theory for them, including Shi-type estimates, monotone quantities etc. In particular, what can one say in terms of singularity formation, solitons etc.?
- 3. We are currently investigating the general flows of $SU(n)$ and $Sp(n)$ -structures, for $n \geq 2$, together with the other authors of [\[FLMSE22\]](#page-39-1). The present paper should be considered as a first step in the broader programme of analytical classification of such flows, according to their special stationarity and parabolicity.
- 4. It would be interesting to study the flows introduced here in explicit settings, such as homogeneous spaces, cohomogeneity one etc.
- 5. It was recently shown in $[BK24, Theorem 1.3]$ that there is an $SU(2)$ functional, different from [\(44\)](#page-11-1), whose critical points are Einstein metrics. Using the $SU(2)$ functionals defined here in [§3.3,](#page-14-0) can one find other such functionals capable of detecting special metrics?

A Principal symbols

We summarise the principal symbols of all the second order diffeomorphism-invariants belonging to 3-dimensional SU(2)module (which we identify with $\Lambda^2_-(M)$):

$$
\sigma(L_{S_{0,1}^{2}(\nabla a_{1})})(x,\xi)(A) \diamond \omega_{1} = \frac{1}{2}\pi^{2}(\xi \wedge * (J_{1}\xi \wedge \tilde{B}_{0,1}) - \xi \wedge * (J_{3}\xi \wedge \tilde{B}_{0,3}) - \xi \wedge * (J_{2}\xi \wedge \tilde{B}_{0,2}) - 2f_{1}\xi \wedge J_{1}\xi)
$$

$$
\sigma(L_{S_{0,2}^{2}(\nabla a_{2})})(x,\xi)(A) \diamond \omega_{2} = \frac{1}{2}\pi^{2}(\xi \wedge * (J_{2}\xi \wedge \tilde{B}_{0,2}) - \xi \wedge * (J_{1}\xi \wedge \tilde{B}_{0,1}) - \xi \wedge * (J_{3}\xi \wedge \tilde{B}_{0,3}) - 2f_{2}\xi \wedge J_{2}\xi)
$$

$$
\sigma(L_{S_{0,3}^{2}(\nabla a_{3})})(x,\xi)(A) \diamond \omega_{3} = \frac{1}{2}\pi^{2}(\xi \wedge * (J_{3}\xi \wedge \tilde{B}_{0,3}) - \xi \wedge * (J_{2}\xi \wedge \tilde{B}_{0,2}) - \xi \wedge * (J_{1}\xi \wedge \tilde{B}_{0,1}) - 2f_{3}\xi \wedge J_{3}\xi)
$$

$$
\sigma(L_{S_{0,3}^{2}(\nabla a_{1})})(x,\xi)(A) \diamond \omega_{2} = \frac{1}{2}\pi^{2}(\xi \wedge * (J_{2}\xi \wedge \tilde{B}_{0,1}) + \frac{1}{2}|\xi|^{2}\tilde{B}_{0,3} + \xi \wedge * (J_{1}\xi \wedge \tilde{B}_{0,2}) - 2f_{1}\xi \wedge J_{2}\xi + f_{0}\xi \wedge J_{3}\xi)
$$

$$
\sigma(L_{S_{0,3}^{2}(\nabla a_{1})})(x,\xi)(A) \diamond \omega_{3} = \frac{1}{2}\pi^{2}(\xi \wedge * (J_{3}\xi \wedge \tilde{B}_{0,2}) + \frac{1}{2}|\xi|^{2}\tilde{B}_{0,1} + \xi \wedge * (J_{2}\xi \wedge \tilde{B}_{0,3}) - 2f_{2}\xi \wedge J_{3}\xi + f_{0}\xi \wedge J_{1}\xi)
$$

Observe that the above comprises of 12 independent terms which is indeed consistent with Theorem [4.2.](#page-20-1) Next we summarise the principal symbols of all the second-order invariants belonging to the trivial representation of SU(2):

$$
\sigma(L_{\delta a_{1}})(x,\xi)(A) = -2f_{1}|\xi|^{2} - *(\xi \wedge J_{2}\xi \wedge \tilde{B}_{0,3}) + *(\xi \wedge J_{3}\xi \wedge \tilde{B}_{0,2}).
$$

$$
\sigma(L_{\delta a_{2}})(x,\xi)(A) = -2f_{2}|\xi|^{2} - *(\xi \wedge J_{3}\xi \wedge \tilde{B}_{0,1}) + *(\xi \wedge J_{1}\xi \wedge \tilde{B}_{0,3}).
$$

$$
\sigma(L_{\delta a_{3}})(x,\xi)(A) = -2f_{3}|\xi|^{2} - *(\xi \wedge J_{1}\xi \wedge \tilde{B}_{0,2}) + *(\xi \wedge J_{2}\xi \wedge \tilde{B}_{0,1}).
$$

$$
\sigma(L_{g(da_{1},\omega_{1})})(x,\xi)(A) = f_{0}|\xi|^{2} - g(\xi \wedge J_{1}\xi, \tilde{B}_{0,1}) + g(\xi \wedge J_{2}\xi, \tilde{B}_{0,2}) + g(\xi \wedge J_{3}\xi, \tilde{B}_{0,3})
$$

$$
\sigma(L_{g(da_{2},\omega_{2})})(x,\xi)(A) = f_{0}|\xi|^{2} - g(\xi \wedge J_{2}\xi, \tilde{B}_{0,2}) + g(\xi \wedge J_{3}\xi, \tilde{B}_{0,3}) + g(\xi \wedge J_{1}\xi, \tilde{B}_{0,1})
$$

$$
\sigma(L_{g(da_{3},\omega_{3})})(x,\xi)(A) = f_{0}|\xi|^{2} - g(\xi \wedge J_{3}\xi, \tilde{B}_{0,3}) + g(\xi \wedge J_{1}\xi, \tilde{B}_{0,1}) + g(\xi \wedge J_{2}\xi, \tilde{B}_{0,2})
$$

$$
\sigma(L_{g(da_{3},\omega_{3})})(x,\xi)(A) = -g(\xi \wedge J_{2}\xi, \tilde{B}_{0,1}) - g(\xi \wedge J_{1}\xi, \tilde{B}_{0,2})
$$

$$
\sigma(L_{g(da_{3},\omega_{1})})(x,\xi)(A) = -g(\xi \wedge J_{3}\xi, \tilde{B}_{0,2}) - g(\xi \wedge J_{2}\xi, \tilde{B}_{0,3})
$$

$$
\sigma(L_{g(da_{3},
$$

Observe that the above comprises of $9 = 12 - 3$ independent terms since $\sigma(L_{g(da_i, \omega_i)}) = \sigma(L_{g(da_j, \omega_i)})$ which is again consistent with Theorem [4.2.](#page-20-1)

References

- [BBG83] Eric Berger, Robert Bryant, and Phillip Griffiths, *The Gauss equations and rigidity of isometric embeddings*, Duke Mathematical Journal 50 (1983), no. 3, 803–892. ↑[20](#page-19-4)
- [Ber55] Marcel Berger, *Sur les groupes d'holonomie homogènes de variétés à connexion affine et des variétés Riemanniennes*, Bulletin de la Société Mathématique de France 83 (1955), 279–330 (fr). MR18,149a ↑[37](#page-36-1)
- [BK24] Niren Bhoja and Kirill Krasnov, *SU(2) structures in four dimensions and Plebanski formalism for GR*, arXiv:2405.15408 (2024). ↑[3,](#page-2-1) [38](#page-37-2)
- [Boy86] Charles Boyer, *Conformal duality and compact complex surfaces*, Mathematische Annalen 274 (1986), 517–526. ↑[23](#page-22-1)
- [Boy88] , *A note on hyperhermitian four-manifolds*, Proceedings of the American Mathematical Society 102 (1988), no. 1, 157–164. ↑[25](#page-24-3)
- [Bry05] Robert Bryant, *Some remarks on* G2*-structures*, Proceedings of 12th Gökova Geometry-Topology Conference, 2005, pp. 75–109. ↑[2,](#page-1-7) [3,](#page-2-1) [8,](#page-7-9) [9,](#page-8-4) [20,](#page-19-4) [22,](#page-21-5) [23,](#page-22-1) [36](#page-35-3)
- [Bry87] , *Metrics with exceptional holonomy*, Annals of Mathematics 126 (1987), no. 3, 525–576. ↑[20](#page-19-4)
- [BV07] Lucio Bedulli and Luigi Vezzoni, *The Ricci tensor of SU(3)-manifolds*, Journal of Geometry and Physics 57 (2007), no. 4, 1125–1146. ↑[36,](#page-35-3) [37](#page-36-1)
- [CK04] Bennett Chow and Dan Knopf, *The Ricci Flow: An Introduction*, Mathematical surveys and monographs, American Mathematical Society, 2004. ↑[9,](#page-8-4) [11,](#page-10-4) [28,](#page-27-4) [32](#page-31-3)
- [CLN06] Bennett Chow, Peng Lu, and Lei Ni, *Hamilton's Ricci Flow*, Graduate studies in mathematics, American Mathematical Society/Science Press, 2006. ↑[28](#page-27-4)
- [CS07] Diego Conti and Simon Salamon, *Generalized killing spinors in dimension 5*, Transactions of the American Mathematical Society 359 (2007), no. 11, 5319–5343. ↑[4](#page-3-4)
- [Der83] Andrzej Derdziński, Self-dual Kähler manifolds and Einstein manifolds of dimension four, Compositio Mathematica 49 (1983), no. 3, 405-433 (en). MR707181 ↑[25](#page-24-3)
- [DeT83] Dennis M. DeTurck, *Deforming metrics in the direction of their Ricci tensors*, Journal of Differential Geometry 18 (1983), no. 1, 157–162. ↑[3](#page-2-1)
- [DGK21] Shubham Dwivedi, Panagiotis Gianniotis, and Spiro Karigiannis, *A gradient flow of isometric* G₂-structures, The Journal of Geometric Analysis 31 (2021), no. 2, 1855–1933. †[29,](#page-28-6) [38](#page-37-2)
- [DGK23] Shubham Dwivedi, Panagiotis Gianniotis, and Spiro Karigiannis, *Flows of* G₂-structures, II: Curvature, torsion, symbols, and functionals, arXiv:2311.05516 (2023). ↑[2,](#page-1-7) [3,](#page-2-1) [4,](#page-3-4) [20,](#page-19-4) [21,](#page-20-6) [22,](#page-21-5) [31,](#page-30-1) [32,](#page-31-3) [35,](#page-34-5) [36](#page-35-3)
- [DK90] Simon Donaldson and Peter Kronheimer, *The geometry of four-manifolds*, Oxford University Press, 1990. ↑[9,](#page-8-4) [10](#page-9-7)
- [DLSE24] Shubham Dwivedi, Eric Loubeau, and Henrique N. Sá Earp, *Harmonic flow of* Spin(7)*-structures*, Ann. Sc. Norm. Super. Pisa, Cl. Sci. (5) 25 (2024), no. 1, 151–215 (English). ↑[4,](#page-3-4) [9](#page-8-4)
- [Dwi24] Shubham Dwivedi, *A gradient flow of Spin(7)-structures*, arXiv:2404.00870 (2024). ↑[3,](#page-2-1) [36,](#page-35-3) [37](#page-36-1)
- [ES64] James Eells and Joseph H. Sampson, *Harmonic Mappings of Riemannian Manifolds*, American Journal of Mathematics 86 (1964), no. 1, 109– 160. ↑[32](#page-31-3)
- [FLMSE22] Daniel Fadel, Eric Loubeau, Andrés J. Moreno, and Henrique N. Sá Earp, *Flows of geometric structures*, arXiv:2211.05197 (2022). ↑[2,](#page-1-7) [3,](#page-2-1) [4,](#page-3-4) [7,](#page-6-4) [9,](#page-8-4) [10,](#page-9-7) [36,](#page-35-3) [38](#page-37-2)
	- [Fow20] Udhav Fowdar, S 1 *-quotient of* Spin(7)*-structures*, Annals of Global Analysis and Geometry 57 (2020), no. 4, 489–517. ↑[23,](#page-22-1) [37](#page-36-1)
	- [FSE24] Udhav Fowdar and Henrique N. Sá Earp, *Harmonic flow of quaternion-Kähler structures*, The Journal of Geometric Analysis 34 (2024), no. 183. ↑[9](#page-8-4)
	- [FY18] Joel Fine and Chengjian Yao, *Hypersymplectic 4-manifolds, the* G2*-Laplacian flow, and extension assuming bounded scalar curvature*, Duke Math. J. 167 (2018), no. 18, 3533–3589. ↑[3,](#page-2-1) [4](#page-3-4)
	- [Gri17] Sergey Grigorian, G2*-structures and octonion bundles*, Advances in Mathematics 308 (2017), 142–207. ↑[3,](#page-2-1) [10,](#page-9-7) [38](#page-37-2)
	- [Kar09] Spiro Karigiannis, *Flows of* G₂-structures, I., The Quarterly Journal of Mathematics 60 (2009), no. [4,](#page-3-4) 487–522. ↑4, [9,](#page-8-4) [35](#page-34-5)
	- [KN63] Shoshichi Kobayashi and Katsumi Nomizu, *Foundations of Differential Geometry II*, Volume 61 of Wiley Classics Library, Wiley, 1963. ↑[23](#page-22-1)
	- [LSE23] Eric Loubeau and Henrique N. Sá Earp, *Harmonic flow of geometric structures*, Annals of Global Analysis and Geometry 64 (2023), no. 4, 23. ↑[2,](#page-1-7) [3,](#page-2-1) [8,](#page-7-9) [9,](#page-8-4) [10,](#page-9-7) [29,](#page-28-6) [36,](#page-35-3) [37,](#page-36-1) [38](#page-37-2)
	- [Sal89] Simon Salamon, *Riemannian geometry and holonomy groups*, Longman Scientific & Technical, 1989. ↑[7,](#page-6-4) [9,](#page-8-4) [20,](#page-19-4) [21](#page-20-6)
	- [Top06] Peter Topping, *Lectures on the Ricci Flow*, London Mathematical Society Lecture Note Series, Cambridge University Press, 2006. ↑[11,](#page-10-4) [25,](#page-24-3) [28,](#page-27-4) [35](#page-34-5)
	- [Vai82] Izu Vaisman, *Some curvature properties of complex surfaces*, Annali di Matematica Pura ed Applicata 132 (1982), no. 1, 1–18. ↑[23](#page-22-1)
	- [Woo93] Chris M. Wood, *Instability of the nearly-Kähler six-sphere*, Journal für die reine und angewandte Mathematik 439 (1993), 205–212. ↑[10](#page-9-7)
	- [Woo95] , *Harmonic almost-complex structures*, Compositio Mathematica 99 (1995), no. 2, 183–212. ↑[10](#page-9-7)