# Matrices with Hyperbolical Krein Space Numerical Range

N. Bebiano<sup>1</sup>, R. Lemos<sup>2</sup> and G. Soares<sup>3</sup>

**Abstract.** This paper is devoted to matrices with hyperbolical Krein space numerical range. This shape characterizes the 2-by-2 case and persists for certain classes of matrices, independently of their size. Necessary and sufficient conditions for low dimensional tridiagonal matrices to have this shape are obtained only involving the matrix entries.

AMS subject Classification: Primary 47A12; Secondary 15A60

**Keywords;** numerical range, Krein space, centrosymmetric matrices, tridiagonal matrices, Hyperbolical Range Theorem

#### 1 Introduction

The investigation of the numerical range and its generalizations has attracted the interest of researchers from different areas of pure and applied science [4, 10], since the concept was born with Toeplitz [26] and Hausdorff [19] in the second decade of the last century. The numerical range of linear operators in an indefinite inner product space is a topic which has deserved the study by some authors; we refer the interested reader to [3, 5, 6, 11, 13, 14, 22, 23, 24, 25]. Ilya Spitkovsky and collaborators published several inspiring papers on the ellipticity of the classical numerical range (see e.g. [12, 15, 16, 20, 21]). Recently, the numerical range of some structured tridiagonal matrices was studied [7, 8]. In this note, we investigate classes of matrices with hyperbolical Krein space numerical range, which can be seen as counterparts of these results for the indefinite inner product space context.

Let  $M_n$  stand for the associative algebra of  $n \times n$  complex matrices and  $I_n$  be the identity matrix. Consider the complex vector space  $\mathbb{C}^n$  equipped with the Krein space structure induced by the *indefinite inner product*  $[x,y]_J := y^*Jx$ , for  $x,y \in \mathbb{C}^n$ , with  $J = P^T(I_r \oplus -I_{n-r})P$ , for some permutation matrix  $P \in M_n$  and  $0 \le r \le n$  (for references on Krein spaces see e.g. [1, 17]).

The Krein space numerical range (KNR), also called indefinite numerical range, of  $A \in M_n$  is denoted and defined as

$$W^{J}(A) := -W_{-}^{J}(A) \cup W_{+}^{J}(A)$$

<sup>&</sup>lt;sup>1</sup>CMUC and Mathematics Department, University of Coimbra, 3001-501 Coimbra (bebiano@mat.uc.pt)

<sup>&</sup>lt;sup>2</sup>CIDMA, Department of Mathematics, University of Aveiro, Aveiro, Portugal (rute@ua.pt)

<sup>&</sup>lt;sup>3</sup>CM-UTAD, Department of Mathematics, University of Trás-os-Montes e Alto Douro, Vila Real, Portugal (gsoares@utad.pt)

with

$$W_{+}^{J}(A) := \{ [Ax, x]_{J} : x \in \mathbb{C}^{n}, [x, x]_{J} = \pm 1 \}.$$

It is clear that  $W_{-}^{J}(A) = W_{+}^{-J}(-A)$  and  $W^{J}(A) = \{1\}$  if and only if  $A = I_n$ . If  $J = I_n$ , then  $W^{J}(A) = W_{+}^{J}(A)$  reduces to the well-known classical numerical range W(A), a compact and convex set as asserted by the Toeplitz-Hausdorff Theorem [19, 26].

These sets share some properties, but they also have big differences, such as the possible degeneracy of  $W^J(A)$  to the complex plane, which never occurs for W(A) when  $A \in M_n$ . Although each set  $W_{\pm}^J(A)$  is connected, it may not be closed or bounded [24]. Both sets  $W_{+}^J(A)$ ,  $W_{-}^J(A)$  are convex, and  $W^J(A)$  is pseudo-convex, that is, for any pair of distinct points  $z_1, z_2 \in W^J(A)$ , either  $W^J(A)$  contains the closed line segment  $\{tz_1 + (1-t)z_2 : 0 \le t \le 1\}$  or  $W^J(A)$  contains the half-lines  $\{tz_1 + (1-t)z_2 : t \le 0 \text{ or } t \ge 1\}$ .

A matrix  $A \in M_n$  is called *J-Hermitian* if it equals its *J-adjoint* which is given by  $A^{\#} := JA^*J$  and it is *J-unitary* if  $AA^{\#} = I_n$ . Consider the following representation

$$A = \Re^J(A) + i\,\Im^J(A),$$

where

$$\Re^{J}(A) = \frac{A + A^{\#}}{2}$$
 and  $\Im^{J}(A) = \frac{A - A^{\#}}{2i}$ 

are J-Hermitian matrices. For each angle  $\theta \in \mathbb{R}$ , let

$$H_{\theta}(A) = \Re^{J}(A)\cos\theta + \Im^{J}(A)\sin\theta, \qquad \theta \in \mathbb{R},$$
 (1)

which is also J-Hermitian, and so its eigenvalues are real or occur in complex conjugate pairs [17]. It is known that if  $H_{\theta}(A)$  has non-real eigenvalues, then  $W^{J}(H_{\theta}(A))$  is the whole real line [5, Proposition 2.1]. To avoid this trivial case, we assume throughout that all its eigenvalues are real. Let us define

$$\sigma_{\pm}^{J}(H_{\theta}(A)) := \{ \lambda \in \mathbb{R} : \exists x \in \mathbb{C}^{n}, [x, x]_{J} = \pm 1, H_{\theta}(A) x = \lambda x \},$$

where

$$\sigma_{+}^{J}(H_{\theta}(A)) = \{\lambda_{1}(\theta), \dots, \lambda_{r}(\theta)\}, \quad \lambda_{1}(\theta) \ge \dots \ge \lambda_{r}(\theta),$$
  
$$\sigma_{-}^{J}(H_{\theta}(A)) = \{\lambda_{r+1}(\theta), \dots, \lambda_{n}(\theta)\}, \quad \lambda_{r+1}(\theta) \ge \dots \ge \lambda_{n}(\theta).$$

We will be mainly concerned with the class  $\mathcal{J}$  of matrices  $H_{\theta}(A)$  with real eigenvalues, such that  $\sigma_{+}^{J}(H_{\theta}(A))$  and  $\sigma_{-}^{J}(H_{\theta}(A))$  do not interlace, that is, either  $\lambda_{r}(\theta) > \lambda_{r+1}(\theta)$  or  $\lambda_{n}(\theta) > \lambda_{1}(\theta)$ . Notice that if the eigenvalues of  $H_{\theta}(A)$  interlace, then  $W^{J}(H_{\theta}(A))$  is the whole real line too.

We recall that a support line of a convex set  $S \subset \mathbb{C}$  is a line containing a boundary point of S and defining two half-planes, such that one of them does not contain S. The support lines of  $W^J(A)$  are by definition the support lines of the convex sets  $-W^J_-(A)$  and  $W^J_+(A)$ .

Let  $\Omega_A = (\eta, \xi)$  be an interval of angles  $\theta$ , with greatest possible diameter, such that  $H_{\theta}(A) \in \mathcal{J}$ . For  $\theta \in \Omega_A$ , if  $\lambda_r(\theta) > \lambda_{r+1}(\theta)$  holds, then the maximal eigenvalue in  $\sigma_-^J(H_{\theta}(A))$  is denoted by  $\lambda_L(H_{\theta}(A))$  and the minimal eigenvalue in  $\sigma_+^J(H_{\theta}(A))$  by  $\lambda_R(H_{\theta}(A))$ . It is known that  $x = \lambda_L(H_{\theta}(A))$  and  $x = \lambda_R(H_{\theta}(A))$  are supports lines of  $-W_-^J(A)$  and  $W_+^J(A)$ , respectively. Analogously, if  $\lambda_n(\theta) > \lambda_1(\theta)$  holds, the maximal eigenvalue in  $\sigma_+^J(H_{\theta}(A))$  is denoted by  $\lambda_L(H_{\theta}(A))$  and the minimal eigenvalue in  $\sigma_-^J(H_{\theta}(A))$  by  $\lambda_R(H_{\theta}(A))$ . Due to the convexity of  $W_\pm^J(A)$ , these families of eigenvalues, when  $\theta$  ranges over  $\Omega_A$ , provide the characterization of these sets as described below.

The KNR generating polynomial of  $W^{J}(A)$  is defined as

$$p_A(z,\theta) = \det \left( \Re^{J}(A) \cos \theta + \Im^{J}(A) \sin \theta - z I_n \right)$$

and plays a central role in the study of this set. Indeed, to a matrix  $A \in M_n$ , through the equation

$$\det (u \Re^{J}(A) + v \Im^{J}(A) + w I_n) = 0,$$

is associated an algebraic curve of class n in homogeneous line coordinates, called the boundary generating curve of  $W^J(A)$ . Its real part, denoted by  $C^J(A)$ , generates  $W^J(A)$  as its pseudoconvex hull, which is obtained as follows: for any two points  $z_1 = [Ax, x]_J$ ,  $z_2 = [Ay, y]_J$  in the boundary generating curve, take the line segment joining them if  $[x, x]_J[y, y]_J > 0$  and the two rays  $\{tz_1 + (1-t)z_2 : t \leq 0 \text{ or } t \geq 1\}$  if  $[x, x]_J[y, y]_J < 0$  (see e.g. [3] for more details).

All the possible shapes of the numerical range  $W^J(A)$  of matrices of size 3 were classified in [13, 14] based on the factorability of the KNR generating polynomial  $p_A$ . If this polynomial factors into three linear factors, then  $C^J(A)$  consists of three points, which are the eigenvalues of A, and  $W^J(A)$  is the pseudo-convex hull of the eigenvalues. Another possibility is that  $p_A$  factors into a linear factor and a quadratic one. Thus  $C^J(A)$  consists of a point, the eigenvalue p corresponding to the linear factor, and a hyperbola or ellipse  $\mathcal{H}$ , corresponding to the quadratic factor, according to the signature of the involved submatrix of J. If p lies outside the hyperbola (resp. ellipse), then p is a nondifferentiable boundary point and there are flat portions on the boundary of  $W^J(A)$ . If p lies in the interior of  $\mathcal{H}$ , then  $W^J(A)$  is the hyperbolic (resp. elliptic) disc bounded by  $\mathcal{H}$  and the matrix A may, or may not, be J-unitarily reducible. We recall that a matrix A is J-unitarily reducible if there exists a J-unitary matrix U, such that

$$U^{\#}AU = A_1 \oplus A_2,$$

where the sum of the sizes of the blocks  $A_1$ ,  $A_2$  is the size of A.

In general, not too much is known about the boundary of  $W^J(A)$ , denoted by  $\partial W^J(A)$ . Here, we restrict our attention to structured matrices with lower dimensional order. The investigation of the hyperbolicity of the Krein space numerical range of tridiagonal matrices of order 3 and of centrosymmetric tridiagonal matrices of order 5 with biperiodic main diagonal is investigated and the obtained results are given in terms of the matrices entries. We also consider centrosymmetric tridiagonal matrices of order 4 with quasi-biperiodic main diagonal, and we show that  $C^J(A)$  may be formed by two hyperbolas. In this case, flat portions may eventually appear on  $\partial W^J(A)$  in conjunction with hyperbolic arcs, or  $W^J(A)$  may degenerate into the whole complex plane.

The remaining of this note is organized as follows. Section 2 contains auxiliary results useful in subsequent discussions. In Section 3, counterparts of classical numerical range results, some recently given in [8] and others mainly due to Spitkovsky and collaborators (e.g. [12, 15, 21]), are presented in the set up of indefinite inner product spaces. Illustrative examples of the obtained results are also given. In Section 4, some observations on the case of  $6 \times 6$  matrices are performed.

# 2 Pre-requisites

For  $A \in M_n$  the following basic properties of  $W^J(A)$  will be used throughout.

- (i)  $W^{J}(\alpha A + \beta I_n) = \alpha W^{J}(A) + \beta$  for any  $\alpha, \beta \in \mathbb{C}$ .
- (ii)  $W^J(A)$  is J-unitarily invariant:  $W^J(U^\#AU) = W^J(A)$  for any J-unitary  $U \in M_n$ .
- (iii)  $W^{J[\kappa]}(A[\kappa]) \subseteq W^J(A)$ , where  $A[\kappa]$  (resp.  $J[\kappa]$ ) denotes the principal submatrix of A (resp. J) that lies in rows and columns indexed by the set  $\kappa \subset \{1, \ldots, n\}$  [18, Lemma 2.3].
- (iv) For  $A = (a_{ij})$  tridiagonal,  $W^J(A)$  is invariant under interchange of the entries  $a_{j,j+1}$ ,  $a_{j+1,j}$ ,  $j = 1, \ldots, n-1$  [11, Lemma 2.1].
- (v) Hyperbolical Range Theorem [3, 6]: For J = diag(1, -1) and  $A \in M_2$ , the set  $W^J(A)$  is bounded by a non-degenerate hyperbola with foci at the eigenvalues  $\lambda_1, \lambda_2$  of A, transverse and non-transverse axes of length

$$\left(\operatorname{Tr}(A^{\#}A) - 2\Re(\lambda_1\bar{\lambda}_2)\right)^{\frac{1}{2}}$$
 and  $\left(|\lambda_1|^2 + |\lambda_2|^2 - \operatorname{Tr}(A^{\#}A)\right)^{\frac{1}{2}}$ ,

respectively, if and only if

$$2\Re(\bar{\lambda}_1\lambda_2) < \operatorname{Tr}(A^{\#}A) < |\lambda_1|^2 + |\lambda_2|^2.$$

If  $A \in M_2$  is J-Hermitian non-scalar, then  $W^J(A)$  degenerates into either two half-rays with endpoints at the eigenvalues  $\lambda_1, \lambda_2$  of A, when the eigenvalues are real, or the whole real line,

when they are complex; and it reduces to a singleton if and only if A is a scalar matrix. Moreover,  $W^{J}(A)$  may be the whole complex plane, eventually except a line (see [2] for a detailed description).

(v) If z is a corner of  $W^J(A)$ , that is, if z lies on more than one support line of  $W^J(A)$ , then z is an eigenvalue of A [22].

The following criterion of non-degenerate hyperbolicity of  $W^{J}(A)$  was obtained in [6].

**Theorem 2.1** Let  $\tilde{a}, \tilde{b} > 0$ ,  $J = I_r \oplus -I_{n-r}$ , 0 < r < n, and  $A \in M_n$ . The set  $W^J(A)$  is bounded by the non-degenerate hyperbola centered at the origin, with horizontal transverse and vertical non-transverse semi-axes of lengths, respectively,  $\tilde{a}$  and  $\tilde{b}$  if and only if

$$\lambda_R(H_{\theta}(A)) = (\widetilde{a}^2 - \widetilde{c}^2 \sin^2 \theta)^{\frac{1}{2}} \text{ and } \lambda_L(H_{\theta}(A)) = -(\widetilde{a}^2 - \widetilde{c}^2 \sin^2 \theta)^{\frac{1}{2}}$$

where  $\widetilde{c}^2 = \widetilde{a}^2 + \widetilde{b}^2$  for all  $\theta \in \Omega_A = (-\theta_0, \theta_0)$  and  $\theta_0 = \arctan(\widetilde{a}/\widetilde{b})$ .

The criterion in Theorem 2.1 can be modified as follows.

Corollary 2.1 Let  $p, q, t \in \mathbb{R}$ , s = q + it,  $\gamma = \frac{1}{2} \arg(s)$ ,  $p^2 < q^2 + t^2$  and let  $\mathcal{H}$  be the hyperbola centered at the origin, transverse semi-axis parallel to  $e^{i\gamma}$  and non-transverse semi-axis of lengths  $\sqrt{|s|+p}$  and  $\sqrt{|s|-p}$ , respectively. If  $J = I_r \oplus -I_{n-r}$ , 0 < r < n and  $A \in M_n$ ,  $W^J(A)$  is bounded by the non-degenerate hyperbola  $\mathcal{H}$  if and only if  $z = \lambda_R(H_\theta(A))$  and  $z = \lambda_L(H_\theta(A))$  satisfy the quadratic equation

$$z^2 = p + q\cos(2\theta) + t\sin(2\theta) \tag{2}$$

for all  $\theta \in \Omega_A = (\gamma - \theta_0, \gamma + \theta_0)$  and

$$\theta_0 = \arctan\left(\frac{|s|+p}{|s|-p}\right)^{\frac{1}{2}}.$$

**Proof.** Under the hypothesis, we have |s| > |p|. Remark that  $\partial W^J(A) = \mathcal{H}$  if and only if  $\partial W^J(\mathrm{e}^{-i\gamma}A) = \mathrm{e}^{-i\gamma}\mathcal{H}$  is the rotated hyperbola centered at the origin, with horizontal transverse and vertical non-transverse semi-axes, respectively, of lengths  $\widetilde{a} = \sqrt{|s| + p} > 0$  and  $\widetilde{b} = \sqrt{|s| - p} > 0$ . By Theorem 2.1, this is equivalent to

$$\lambda_R(H_{\eta}(e^{-i\gamma}A)) = (p+|s|-2|s|\sin^2\eta)^{\frac{1}{2}} = -\lambda_L(H_{\eta}(e^{-i\gamma}A))$$

for  $\eta \in (-\theta_0, \theta_0)$  and  $\theta_0 = \arctan(\tilde{a}/\tilde{b})$ , observing that  $\tilde{a}^2 + \tilde{b}^2 = 2|s|$ . Since  $H_{\eta}(e^{-i\gamma}A) = H_{\eta+\gamma}(A)$ , considering  $\theta = \eta + \gamma$ , the previous hyperbolicity condition holds if and only if

$$\lambda_R(H_{\theta}(A)) = \lambda_R(H_{\theta-\gamma}(e^{-i\gamma}A)) = (p+|s|-2|s|\sin^2(\theta-\gamma))^{\frac{1}{2}},$$

$$\lambda_L(H_{\theta}(A)) = -\lambda_R(H_{\theta}(A)), \quad \theta \in (\gamma - \theta_0, \gamma + \theta_0).$$

By trivial trigonometric transformations, we find

$$|s| - 2|s| \sin^2(\theta - \gamma) = |s| \cos^2(\theta - \gamma) - |s| \sin^2(\theta - \gamma)$$
$$= |s| \cos(2\theta - 2\gamma)$$
$$= |s| \cos(2\gamma) \cos(2\theta) + |s| \sin(2\gamma) \sin(2\theta)$$
$$= q \cos(2\theta) + t \sin(2\theta).$$

Then the previous hiperbolicity condition occurs if and only if  $z = \lambda_R(H_{\theta}(A))$  and  $z = \lambda_L(H_{\theta}(A))$  satisfy condition (2), having in mind that the RHS of the quadratic equation in (2) is positive for all angles  $\theta$  in the interval  $\Omega_A = (\gamma - \theta_0, \gamma + \theta_0)$ .

In Theorem 2.1 and Corollary 2.1, we may consider, for 0 < r < n,  $J = P^T(I_r \oplus -I_{n-r}) P$ , with  $P \in M_n$  a permutation matrix.

**Remark 2.1** In parallel to the classical numerical range (see, e.g. [20, Proposition 1]), if  $C^J(A)$  contains a hyperbola centered at a point (c,d), then the KNR generating polynomial of  $W^J(A)$  is divisible by a factor of the form

$$(z - c\cos\theta - d\sin\theta)^2 - (p + q\cos(2\theta) + t\sin(2\theta)),$$

where p, q, t are defined as in Corollary 2.1.

#### 3 Results

### 3.1 Hyperbolicity of $W^{J}(A)$ for orders 3 and 5

In this subsection, we will be concerned with the Krein space numerical range of tridiagonal matrices of odd order with biperiodic main diagonal, that is, matrices  $A = (a_{ij}) \in M_{2n+1}$ , such that  $a_{ij} = 0$ , |i - j| > 1, with  $a_{ii} = a_1$  if i is odd and  $a_{ii} = a_2$  if i is even, considering

$$J = diag(1, -1, 1, \dots, -1, 1) \in M_{2n+1}$$

with biperiodic main diagonal too. Without loss of generality, we may focus on tridiagonal matrices with real biperiodic main diagonal, denoted by  $T_{2n+1}(\mathbf{c}, \mathbf{a}, \mathbf{b})$ , where

$$\mathbf{a} = (a, -a, a, \dots, -a, a), \quad \mathbf{b} = (b_1, b_2, \dots, b_{2n}), \quad \mathbf{c} = (c_1, c_2, \dots, c_{2n})$$

are the main diagonal, the first upper and the first lower subdiagonals, respectively. In fact, for m = 2n + 1, we easily see that

$$A = e^{i\tau} T_m(\mathbf{c}, \mathbf{a}, \mathbf{b}) + \delta I_m, \tag{3}$$

considering  $a = \frac{1}{m}|{\rm Tr}(JA)|,\; \delta = \frac{1}{m}{\rm Tr}(A),\; \tau = \arg{\rm Tr}(JA)$  and

$$b_j = a_{j,j+1} e^{-i\tau}, \qquad c_j = a_{j+1,j} e^{-i\tau}, \qquad j = 1, \dots, m-1.$$
 (4)

By property (i),  $W^{J}(A)$  is readily obtained from  $W^{J}(T_{m}(\mathbf{c}, \mathbf{a}, \mathbf{b}))$ .

We are interested in deriving conditions ensuring nondegenerate hyperbolicity of the KNR. We start with the  $3 \times 3$  tridiagonal case.

**Theorem 3.1** Let J = diag(1, -1, 1),  $A = T_3(\mathbf{c}, \mathbf{a}, \mathbf{b})$  with  $\mathbf{a} = (a, -a, a)$ ,  $\mathbf{b} = (b_1, b_2)$ ,  $\mathbf{c} = (c_1, c_2)$  and  $\Delta = a^2 + b_1c_1 + b_2c_2$ . The set  $W^J(A)$  is a nondegenerate hyperbolic disc bounded by  $\mathcal{H}$ , with foci at  $\pm \Delta^{\frac{1}{2}}$  and non-transverse axis of length

$$(2|\Delta| - 2a^2 + |b_1|^2 + |b_2|^2 + |c_1|^2 + |c_2|^2)^{\frac{1}{2}},$$
(5)

if and only if

$$a^2 - 2|\Delta| < \text{Tr}(A^\# A) < a^2 + 2|\Delta|.$$
 (6)

In this case,  $C^{J}(A)$  is the union of the hyperbola  $\mathcal{H}$  and the point a.

**Proof.** Since  $\det(A - \lambda I_3) = -(\lambda - a)(\lambda^2 - \Delta)$ , the eigenvalues of A are easily obtained:  $a, -\Delta^{\frac{1}{2}}, \Delta^{\frac{1}{2}}$ , and we find that

$$H_{\theta}(A) = \frac{1}{2} \begin{bmatrix} 2a & b_1 - \overline{c_1} & 0 \\ c_1 - \overline{b_1} & -2a & c_2 - \overline{b_2} \\ 0 & b_2 - \overline{c_2} & 2a \end{bmatrix} \cos \theta + \frac{1}{2i} \begin{bmatrix} 0 & b_1 + \overline{c_1} & 0 \\ c_1 + \overline{b_1} & 0 & c_2 + \overline{b_2} \\ 0 & b_2 + \overline{c_2} & 0 \end{bmatrix} \sin \theta.$$

The KNR generating polynomial of  $W^{J}(A)$  is

$$p_A(z,\theta) = -(z - a\cos\theta) \left(z^2 - p - q\cos(2\theta) - t\sin(2\theta)\right)$$

with

$$p = \frac{a^2}{2} - \frac{1}{4} (|b_1|^2 + |b_2|^2 + |c_1|^2 + |c_2|^2) = \frac{1}{4} (\operatorname{Tr}(A^{\#}A) - a^2),$$

$$q = \frac{a^2}{2} + \frac{1}{2} \Re (b_1 c_1 + b_2 c_2) = \frac{1}{2} \Re (\Delta),$$

$$t = \frac{1}{2} \Im (b_1 c_1 + b_2 c_2) = \frac{1}{2} \Im (\Delta).$$

Since  $\cos(2\theta) = 2\cos^2\theta - 1$ , we can see that

$$q\cos(2\theta) + t\sin(2\theta) = a^2\cos^2\theta - \frac{a^2}{2} + \frac{1}{2}\Re(e^{-2i\theta}(b_1c_1 + b_2c_2))$$

and using  $\Re(e^{-2i\theta}(b_1c_1 + b_2c_2) \le |b_1c_1 + b_2c_2|$ , we get

$$p + q\cos(2\theta) + t\sin(2\theta) \le a^2\cos^2\theta. \tag{7}$$

We have  $q + it = \frac{1}{2}\Delta$  and  $p^2 < q^2 + t^2$  holds if and only if

$$\frac{1}{2} \left| \operatorname{Tr}(A^{\#}A) - a^2 \right| < |\Delta|,$$

which is equivalent to (6) as shown below.

Firstly, suppose that (6) holds. In this case,  $\Delta \neq 0$ , the LHS of (7) is positive and  $p_A(z,\theta) = 0$  if and only if  $z = a \cos \theta$  or

$$z = \pm \left(p + q\cos(2\theta) + t\sin(2\theta)\right)^{\frac{1}{2}} \tag{8}$$

for  $\theta \in (\Gamma - \theta_0, \Gamma + \theta_0)$ , where  $\Gamma = \frac{1}{2} \arg(\Delta)$  and

$$\theta_0 = \arctan\left(\frac{|\Delta| + 2p}{|\Delta| - 2p}\right)^{\frac{1}{2}}.$$

Let  $\lambda_{\pm}(\theta)$  be the eigenvalues of  $H_{\theta}(A)$  given by the RHS of (8). We find  $w_{\theta} = (b_2 - \overline{c_2} e^{2i\theta}, 0, \overline{b_1} e^{2i\theta} - c_1)$  as an eigenvector of  $H_{\theta}(A)$  associated to the eigenvalue  $a \cos \theta$ , such that  $[w_{\theta}, w_{\theta}]_J > 0$ . Further,

$$u_{\theta,\pm} = \left(\overline{c_1} e^{i\theta} - b_1 e^{-i\theta}, 2(a\cos\theta - \lambda_{\pm}(\theta)), \overline{b_2} e^{i\theta} - c_2 e^{-i\theta}\right)$$

is an eigenvector of  $H_{\theta}(A)$  associated with  $\lambda_{\pm}(\theta)$ , satisfying

$$[u_{\theta,\pm}, u_{\theta,\pm}]_J = 8\lambda_{\pm}(\theta)(a\cos\theta - \lambda_{\pm}(\theta))$$

and so  $[u_{\theta,+}u_{\theta,+}]_J[u_{\theta,-},u_{\theta,-}]_J < 0$ , because  $H_{\theta}(A)$  is not diagonal, then (7) does not hold as equality and

$$(\lambda_{+}(\theta) - a\cos\theta)(\lambda_{-}(\theta) - a\cos\theta) > 0.$$
(9)

Therefore, one of the eigenvalues  $\lambda_{\pm}(\theta)$  belongs to  $\sigma_{-}^{J}(H_{\theta}(A))$ , the other and  $a\cos\theta$  belong to  $\sigma_{+}^{J}(H_{\theta}(A))$ . We conclude that  $H_{\theta}(A) \in \mathcal{J}$ , with

$$\lambda_R(H_{\theta}(A)) = \lambda_+(\theta)$$
 and  $\lambda_L(H_{\theta}(A)) = \lambda_-(\theta)$ 

for all  $\theta \in \Omega_A = (\Gamma - \theta_0, \Gamma + \theta_0)$ .

By Corollary 2.1,  $W^J(A)$  is a non-degenerate hyperbolic disc centered at the origin, with transverse axis parallel to  $e^{i\Gamma}$ , that is, on the line containing the eigenvalues  $\pm \Delta^{\frac{1}{2}}$ , of length  $(2|\Delta| + 4p)^{\frac{1}{2}}$ 

and with non-transverse axis of length  $(2|\Delta|-4p)^{\frac{1}{2}}$  equal to (5). The foci of the hyperbola are  $\pm \Delta^{\frac{1}{2}}$  and the remaining eigenvalue a of A is in the interior, or on the boundary, of this hyperbolic disc.

Conversely, if  $W^{J}(A)$  is a non-degenerate hyperbolic disc with the foci and non-transverse axis given, we easily see that

$$(2|\Delta| - a^2 + \text{Tr}(A^{\#}A))^{\frac{1}{2}}$$
 and  $(2|\Delta| + a^2 - \text{Tr}(A^{\#}A))^{\frac{1}{2}}$ 

are the axes lengths, this implying that (6) must be satisfied.

Remark 3.1 The hyperbola describing  $\partial W^J(A)$  for A in Theorem 3.1 under the condition (6) can be characterized using the matrix invariants, namely, the eigenvalues  $\lambda_1, \lambda_2, \lambda_3$  of A and the trace of  $A^\#A$ . In fact, the non-transverse axis has length

$$(|\lambda_1|^2 + |\lambda_2|^2 + |\lambda_3|^2 - \text{Tr}(A^{\#}A))^{\frac{1}{2}},$$

at least one of the eigenvalues equals the trace of A and the remaining two are the foci of the hyperbola.

**Example 3.1** Let J = diag(1, -1, 1) and  $A = T_3(\mathbf{c}, \mathbf{a}, \mathbf{d})$  with  $\mathbf{a} = (4, -4, 4)$ ,  $\mathbf{b} = (3 + i, 1)$ ,  $\mathbf{c} = (-4 + i, -i)$ . The eigenvalues of A, given by  $4, -\sqrt{3 - 2i}$  and  $\sqrt{3 - 2i}$ , are represented by green dots in Figure 1. The boundary generating curve  $C^J(A)$  consists of the point (4, 0) and the hyperbola of Figure 1, being  $W^J(A)$  bounded by this hyperbola.




Figure 1:  $C^{J}(A)$  for the matrices in Example 3.1

Now, we apply the previous result to study the case of centrosymmetric tridiagonal matrices of order 5.

**Theorem 3.2** Consider J = diag(1, -1, 1, -1, 1) and  $A = T_5(\mathbf{c}, \mathbf{a}, \mathbf{b})$  with  $\mathbf{a} = (a, -a, a, -a, a)$ ,  $\mathbf{b} = (b_1, b_2, b_3, b_4)$ ,  $\mathbf{c} = (b_4, b_3, b_2, b_1)$ . Let

$$\Delta_1 = a^2 + b_1 b_4, \qquad M_1 = 2a^2 - |b_1|^2 - |b_4|^2,$$

$$\Delta_2 = \Delta_1 + 2b_2b_3, \qquad M_2 = M_1 - 2|b_2|^2 - 2|b_3|^2$$

and  $\mathcal{H}_k$  be the hyperbola with foci  $\pm \sqrt{\Delta_k}$  and non-transverse axis of length

$$(2|\Delta_k| - M_k)^{\frac{1}{2}}$$
.

The set  $W_J(A)$  is bounded by the nondegenerate hyperbola  $\mathcal{H}_2$  if and only if  $|M_2| < 2|\Delta_2|$ . In this case,  $C_J(A)$  consists of the point a and the hyperbolas  $\mathcal{H}_1$ ,  $\mathcal{H}_2$ , with  $\mathcal{H}_1$  eventually degenerating into the points  $\pm \sqrt{\Delta_1}$ .

**Proof.** Let  $E_2$  denote the exchange matrix of order 2, obtained from  $I_2$  by reversing the order of its columns. Then

$$Q = \frac{\sqrt{2}}{2} \begin{bmatrix} I_2 & 0 & E_2 \\ 0 & \sqrt{2} & 0 \\ -E_2 & 0 & I_2 \end{bmatrix}$$

is such that  $Q^{\#}Q = I_5$ . Via this *J*-orthogonal matrix Q, the matrix A is *J*-orthogonaly similar to the block diagonal matrix

$$Q^{\#}AQ = \begin{bmatrix} a & b_1 & 0 & 0 & 0 \\ b_4 & -a & 0 & 0 & 0 \\ 0 & 0 & a & \sqrt{2}b_3 & 0 \\ 0 & 0 & \sqrt{2}b_2 & -a & b_4 \\ 0 & 0 & 0 & b_1 & a \end{bmatrix}.$$

Let  $J_1 = \text{diag}(1, -1)$  and  $J_2 = \text{diag}(1, -1, 1)$ . The boundary generating curves of  $W^J(A)$  are those of  $W^{J_1}(R)$  and  $W^{J_2}(S)$ , where

$$R = \begin{bmatrix} a & b_1 \\ b_4 & -a \end{bmatrix} \quad \text{and} \quad S = \begin{bmatrix} a & \sqrt{2}b_3 & 0 \\ \sqrt{2}b_2 & -a & b_4 \\ 0 & b_1 & a \end{bmatrix}.$$

Using properties (ii)-(iv), considering  $\nu = \{2, 3\}$ , we easily find

$$W^{J_1}(R) = W^{E_2J_1E_2}(E_2RE_2) = W^{J_2[\nu]}(S[\nu]) \subseteq W^{J_2}(S),$$

because  $E_2RE_2 = S[\nu]$ . The eigenvalues  $\lambda_1$ ,  $\lambda_2$  of R are  $\pm\sqrt{\Delta_1}$  and

$$|\lambda_1|^2 + |\lambda_2|^2 - \text{Tr}(J_1 R^* J_1 R) = 2|\Delta_1| - 2a^2 + |b_1|^2 + |b_4|^2.$$

By the Hyperbolical Range Theorem,  $W^{J_1}(R)$  is the hyperbolic disc bounded by  $\mathcal{H}_1$ .

Now, applying Theorem 3.1 to the tridiagonal matrix S with the eigenvalues  $a, \pm \sqrt{\Delta_2}$ , we conclude that  $W^{J_2}(S)$  is a nondegenerate hyperbolic disc bounded by  $\mathcal{H}_2$  if and only if

$$a^2 - 2|\Delta_2| < \text{Tr}(J_2 S^* J_2 S) < a^2 + 2|\Delta_2|.$$

Note that

$$Tr(J_2 S^* J_2 S) = 3a^2 - |b_1|^2 - 2|b_2|^2 - 2|b_3|^2 - |b_4|^2,$$

and the previous double inequality may be witten as  $|M_2| < 2|\Delta_2|$ . In this case,  $a \neq 0$  and  $C^{J_1}(R) = \mathcal{H}_1$  may eventually degenerate into the pair of points  $\pm \sqrt{\Delta_1}$ . Then  $C^J(A) = \mathcal{H}_1 \cup C^{J_2}(S)$  contains also the point a and the non-degenerate hyperbola  $\mathcal{H}_2$ .

Remark 3.2 Under the hypothesis of Theorem 3.2, we have

$$Tr(JA^*JA) = 5a^2 - 2|b_1|^2 - 2|b_2|^2 - 2|b_3|^2 - 2|b_4|^2.$$

Considering the eigenvalues  $\lambda_k$  of A, k = 1, ..., 5, we easily find that

$$\sum_{k=1}^{5} |\lambda_k|^2 - \text{Tr}(JA^*JA) = \sum_{k=1}^{2} (2|\Delta_k| - M_k)$$

is the sum of the squares of the non-transverse axes of the hyperbolas  $\mathcal{H}_1$  and  $\mathcal{H}_2$  belonging to  $C^J(A)$ .

**Example 3.2** Consider J = diag(1, -1, 1, -1, 1) and  $A = T_5(\mathbf{c}, \mathbf{a}, \mathbf{d})$  with  $\mathbf{a} = (6, -6, 6, -6, 6)$ ,  $\mathbf{b} = (-3, 5, i, 2)$ ,  $\mathbf{c} = (2, i, 5, -3)$ . The eigenvalues of A given by 6,  $\pm \sqrt{30}$  and  $\pm \sqrt{30 + 10i}$  are represented by rose dots in Figure 2. The boundary generating curve  $C^J(A)$  consists of the point (6,0) and two nested hyperbolas, displayed in red and blue in Figure 2, being  $W^J(A)$  bounded by the outer hyperbola.




Figure 2:  $C^{J}(A)$  for the matrices in Example 3.2

## 3.2 Bihyperbolic $C^{J}(A)$ for matrices of order 4

In this subsection, we will consider tridiagonal matrices of even order  $A = (a_{ij}) \in M_{2n}$  with main diagonal, satisfying  $a_{ii} = a_{2n-i+1,2n-i+1}$  and  $a_{ii} = a_1$  if  $i \le n$  is odd,  $a_{ii} = a_2$  if  $i \le n$  is even. Let

$$J = diag(1, -1, \dots, -1, 1) \in M_{2n}, \tag{10}$$

that is, with the main diagonal entries  $j_1, \ldots, j_n$  of J, satisfying  $j_i = j_{n-i+1}$ ,  $i = 1, \ldots, n$ . Without loss of generality, by property (i), we may focus on tridiagonal matrices with real *quasi*-biperiodic main diagonal, denoted by  $T_{2n}(\mathbf{c}, \mathbf{a}, \mathbf{b})$ , where

$$\mathbf{a} = (a, -a, \dots, -a, a), \quad \mathbf{b} = (b_1, b_2, \dots, b_{2n-1}), \quad \mathbf{c} = (c_1, c_2, \dots, c_{2n-1})$$

are the main diagonal, the first upper and lower subdiagonals, respectively. In fact, let m = 2n. Now, (3) holds too, considering  $a, \delta, \tau$  defined as before,  $b_j, c_j$  written as (4), but with J given by (10).

Now, the KNR of centrosymmetric tridiagonal matrices of order 4 is studied. As shown below,  $C^{J}(A)$  may be bihyperbolic.

**Theorem 3.3** Let J = diag(1, -1, -1, 1),  $A = T_4(\mathbf{c}, \mathbf{a}, \mathbf{b})$ ,  $\mathbf{a} = (a, -a, -a, a)$ ,  $\mathbf{b} = (b_1, b_2, b_3)$  and  $\mathbf{c} = (b_3, b_2, b_1)$ . Consider

$$\Delta = a^2 + b_1 b_3, \qquad \Delta_{\pm} = \Delta \mp a b_2 + \frac{b_2^2}{4},$$

$$M = 2a^2 - |b_1|^2 - |b_3|^2, \qquad M_{\pm} = M + \frac{1}{2} |b_2|^2 \pm 2 a \Re(b_2).$$

(a)  $C^{J}(A)$  is the union of two nondegenerate and nonconcentric hyperbolas  $\mathcal{H}_{+}$  and  $\mathcal{H}_{-}$  if and only if

$$b_2 \neq 0$$
,  $|M_+| < 2|\Delta_+|$  and  $|M_-| < 2|\Delta_-|$ .

In this case,  $W^{J}(A)$  is the pseudo-convex hull of  $\mathcal{H}_{+}$  and  $\mathcal{H}_{-}$ , where  $\mathcal{H}_{+}$  (resp.  $\mathcal{H}_{-}$ ) has foci  $\frac{1}{2}b_{2} \pm \sqrt{\Delta_{+}}$  (resp.  $-\frac{1}{2}b_{2} \pm \sqrt{\Delta_{-}}$ ) and non-transverse axis of length

$$(2|\Delta_{+}|-M_{+})^{\frac{1}{2}}$$
  $(resp. (2|\Delta_{-}|-M_{-})^{\frac{1}{2}}).$ 

Flat portions on the boundary may eventually exist, or  $W_J(A)$  may degenerate into the whole complex plane.

(b) In particular, if  $b_2 = 0$  and  $|M| < 2|\Delta|$ , then  $W^J(A)$  is a nondegenerate hyperbolic disc with foci at  $\pm \sqrt{\Delta}$ , non-transverse axis of length  $(2|\Delta| - M)^{\frac{1}{2}}$ .

**Proof.** It is easy to see that A is J-orthogonally similar to the block diagonal matrix

$$Q^{\#}AQ = \begin{bmatrix} a & b_1 & 0 & 0 \\ b_3 & -b_2 - a & 0 & 0 \\ 0 & 0 & b_2 - a & b_3 \\ 0 & 0 & b_1 & a \end{bmatrix}$$

via the real J-orthogonal matrix

$$Q = \frac{\sqrt{2}}{2} \begin{bmatrix} I_2 & E_2 \\ -E_2 & I_2 \end{bmatrix}$$

with  $E_2$  the exchange matrix of order 2. Then  $C^J(A)$  is the union of the boundary generating curves of  $W^{J_1}(S_-)$  and  $W^{-J_1}(R)$ , with  $J_1 = \text{diag}(1, -1)$  and

$$S_{\pm} = \begin{bmatrix} a & b_1 \\ b_3 & \pm b_2 - a \end{bmatrix}, \qquad R = \begin{bmatrix} b_2 - a & b_3 \\ b_1 & a \end{bmatrix}.$$

Concerning the last set, it is clear that

$$W^{-J_1}(R) = W^{E_2J_1E_2}(E_2S_+E_2) = W^{J_1}(S_+).$$

By the Hyperbolical Range Theorem, each set  $W^{J_1}(S_{\pm})$  is bounded by a non-degenerate hyperbola  $\mathcal{H}_{\pm}$  with foci at the eigenvalues of  $S_{\pm}$ ,

$$\lambda_{1,\pm} = \pm \frac{1}{2} b_2 + \sqrt{\Delta_{\pm}}, \qquad \lambda_{2,\pm} = \pm \frac{1}{2} b_2 - \sqrt{\Delta_{\pm}},$$

and non-transverse axes of length given by the square root of

$$|\lambda_{1,\pm}|^2 + |\lambda_{2,\pm}|^2 - \text{Tr}(J_1 S_{\pm}^* J_1 S_{\pm}) = 2|\Delta_{\pm}| - M_{\pm}$$

if and only if  $-2|\Delta_{\pm}| < M_{\pm} < 2|\Delta_{\pm}|$ . The hyperbolas  $\mathcal{H}_{+}$  and  $\mathcal{H}_{-}$  are not concentric, except when  $b_2 = 0$ , in which case they coincide and (b) is obtained. Otherwise,  $W^{J}(A)$  has a nondegenerate bihyperbolic boundary generating curve as described in (a). Henceforth, we may obtain:

- i. two nested hyperbolas as  $C^{J}(A)$  and  $W^{J}(A)$  is a hyperbolic disc bounded by the outer hyperbola;
- ii. hyperbolic arcs from  $\mathcal{H}_+$  and  $\mathcal{H}_-$  in  $\partial W_J(A)$ , when these hyperbolas are not nested, and eventually some flat portions on  $\partial W^J(A)$ , generated when taking the pseudo-convex hull of  $\mathcal{H}_+$  and  $\mathcal{H}_-$ ;
- iii. the set  $W^J(A)$  as the whole complex plane, when the eigenvalues of  $H_{\theta}(A)$  interlace for some angles  $\theta \in \Omega_A$ .

All these three cases may occur as illustrated in Examples 3.3 - 3.5 presented below.

As the next example shows, the converse of Theorem 3.3 (b) does not hold, since  $W^{J}(A)$  may be a hyperbolic disc even when  $b_2 \neq 0$ .

**Example 3.3** Let J = diag(1, -1, -1, 1),  $A = T_4(\mathbf{c}, \mathbf{a}, \mathbf{d})$ ,  $\mathbf{a} = (5, -5, -5, 5)$ ,  $\mathbf{b} = (1, 2, 7)$  and  $\mathbf{c} = (7, 2, 1)$ . The eigenvalues of A, which are given by  $-1 \pm \sqrt{43}$ ,  $1 \pm \sqrt{23}$ , are represented by green dots in Figure 3. We have  $C^J(A) = C^J(S_+) \cup C^J(S_-)$ , where

$$S_{+} = \begin{bmatrix} 5 & 1 \\ 7 & -3 \end{bmatrix}$$
 and  $S_{-} = \begin{bmatrix} 5 & 1 \\ 7 & -7 \end{bmatrix}$ ,

using the notation in the proof of Theorem 3.3. Then  $C^J(A)$  consists of two nonconcentric nested hyperbolas displayed in Figure 3, where the blue (red) curves represent the branches of the hyperbolas that bound  $W_+^{J_1}(S_\pm)$  (resp.,  $-W_-^{J_1}(S_\pm)$ ). We conclude that  $W^J(A) = W^J(S_+)$  is bounded by the outer hyperbola, centered at the point (1,0) with focal distance equal to  $2\sqrt{23}$  and vertical nontransverse axis of length  $\sqrt{58}$ . This example illustrates the case i. presented in the proof above.




Figure 3:  $C^{J}(A)$  for the matrices in Example 3.3

**Example 3.4** Let J = diag(1, -1, -1, 1),  $A = T_4(\mathbf{c}, \mathbf{a}, \mathbf{d})$ ,  $\mathbf{a} = (6, -6, -6, 6)$ ,  $\mathbf{b} = (i, 2 + i, 2)$  and  $\mathbf{c} = (2, 2 + i, i)$ . The eigenvalues of A are

$$\frac{1}{2}\left((-2-i)\pm\sqrt{195+36i}\right),\qquad \frac{1}{2}\left((2+i)\pm\sqrt{99-12i}\right),$$

represented by green dots and  $C_J(A)$  consists of a pair of displaced hyperbolas displayed in Figure 4. As in the previous example, the blue (red) curves represent the branches of the hyperbolas associated to points  $z = [Ax, x]_J$  in  $C^J(A)$  generated by vectors x with  $[x, x]_J > 0$  (resp.,  $[x, x]_J < 0$ ). In this case, from the pseudo-convex hull of these two hyperbolas, flat portions will appear on  $\partial W^J(A)$ . This example illustrates the case ii. in the proof of Theorem 3.3.




Figure 4:  $C^{J}(A)$  for the matrices in Example 3.4

**Example 3.5** Let J = diag(1, -1, -1, 1),  $A = T_4(\mathbf{c}, \mathbf{a}, \mathbf{d})$ ,  $\mathbf{a} = (1, -1, -1, 1)$ ,  $\mathbf{b} = (3, 5, 1/2)$  and  $\mathbf{c} = (1/2, 5, 3)$ . The eigenvalues of A are  $\frac{1}{2}(-5 \pm \sqrt{55})$ ,  $\frac{1}{2}(5 \pm \sqrt{15})$ , represented by violet dots and  $C_J(A)$  consists of the pair of nonconcentric hyperbolas displayed in Figure 5. The blue (red) curves represent again the branches of the hyperbolas obtained from points  $z = [Ax, x]_J$  in  $C^J(A)$  with generating vectors x such that  $[x, x]_J > 0$  (resp.,  $[x, x]_J < 0$ ). In this case, we can easily conclude that  $W^J(A)$  is the whole complex plane.




Figure 5:  $C^{J}(A)$  for the matrices in Example 3.5

### 4 Final remarks

In the case of matrices of order 6, the previous cenario changes. Now, let J = diag(1, -1, 1, 1, -1, 1) and  $A = T_6(\mathbf{c}, \mathbf{a}, \mathbf{b})$  be the centrosymmetric matrix with  $\mathbf{a} = (a, -a, a, a, -a, a)$ ,  $\mathbf{b} = (b_1, b_2, b_3, b_4, b_5)$ ,  $\mathbf{c} = (b_5, b_4, b_3, b_2, b_1)$ . It may be seen that A is J-orthogonally similar to the block diagonal matrix:

$$Q^{\#}AQ = \begin{bmatrix} a & b_1 & 0 & 0 & 0 & 0 \\ b_5 & -a & b_2 & 0 & 0 & 0 \\ 0 & b_4 & a - b_3 & 0 & 0 & 0 \\ 0 & 0 & 0 & a + b_3 & b_4 & 0 \\ 0 & 0 & 0 & b_2 & -a & b_5 \\ 0 & 0 & 0 & 0 & b_1 & a \end{bmatrix},$$

via the real J-orthogonal matrix

$$Q = \frac{\sqrt{2}}{2} \left[ \begin{array}{cc} I_3 & E_3 \\ -E_3 & I_3 \end{array} \right].$$

where  $E_3$  is the exchange matrix of order 3.

If  $b_3 = 0$ , then  $C^J(A) = C^{J_2}(S) \cup C^{J_2}(R')$ , where  $J_2 = \text{diag}(1, -1, 1)$ ,

$$S = \begin{bmatrix} a & b_1 & 0 \\ b_5 & -a & b_2 \\ 0 & b_4 & a \end{bmatrix} = E_2 R' E_2, \qquad R' = \begin{bmatrix} a & b_4 & 0 \\ b_2 & -a & b_5 \\ 0 & b_1 & a \end{bmatrix}.$$

As  $J_2 = E_3 J_2 E_3$ ,  $W^{J_2}(R') = W^{J_2}(E_2 R' E_2) = W^{J_2}(S)$  and  $C^J(A)$  reduces to  $C^{J_2}(S)$ . By Theorem 3.1, the set  $W^{J_2}(S)$  is a nondegenerate hyperbolic disc, bounded by the hyperbola  $\mathcal{K}$  with foci at

the points

$$\pm (a^2 + b_1b_5 + b_2b_4)^{\frac{1}{2}}$$

and non-transverse axis of length

$$(2|a^2 + b_1b_5 + b_2b_4| - 2a^2 + |b_1|^2 + |b_2|^2 + |b_4|^2 + |b_5|^2)^{\frac{1}{2}},$$

if and only if

$$\left| \text{Tr}(J_2 S^* J_2 S) - a^2 \right| < 2|a^2 + b_1 b_5 + b_2 b_4|.$$

In this last case,  $C^{J}(A) = \{a\} \cup \mathcal{K}$  with the point a belonging to the interior, or the boundary, of the hyperbolic disc bounded by  $\mathcal{K}$ .

As it is well-known, tridiagonal matrices are too sensitive to perturbations, and even small perturbation provokes dramatic changes in their behaviour. In particular, the necessary and sufficient condition for hyperbolicity is violated for small changes in the (3,3) entry of S.

If  $b_3 \neq 0$ , then the boundary generating curves for both tridiagonal blocks or order 3 in  $Q^{\#}AQ$  may not be hyperbolas. As illustrated in the next example, their KNR generating polynomials are irreducible cubics and the corresponding boundary generating curves are algebraic curves of 6th degree, with hyperbolic-like shapes and deltoids in their interior.

**Example 4.1** Consider J = diag(1, -1, 1, 1, -1, 1) and  $A = T_6(\mathbf{c}, \mathbf{a}, \mathbf{b})$  with  $\mathbf{a} = (4, -4, 4, 4, -4, 4)$ ,  $\mathbf{b} = (1, 2, -2, 4, 5)$ ,  $\mathbf{c} = (5, 4, -2, 2, 1)$ . Moreover, let  $A_0 = T_6(\mathbf{c}, \mathbf{a}, \mathbf{b}_0)$ ,  $\mathbf{b}_0 = (1, 2, 0, 4, 5)$ , which only differs from the matrix A on the entry  $b_3$ .

From the previous discussion, it is clear that  $W^J(A_0)$  is bounded by the hyperbola centered at the origin, with foci at the points  $\pm \sqrt{29}$ , horizontal tranverse axis and vertical non-transverse axis of length  $6\sqrt{2}$ . This hyperbola is the dashed curve displayed in Figure 6, which forms the boundary generating curve of  $W^J(A_0)$  together with the point (5,0) represented by a rose dot in its interior; the other two rose dots correspond to the remaining eigenvalues of  $A_0$ , which are the foci given above.

Concerning the matrix A, the KNR generating polynomial of  $W^{J}(A)$  is the product of two irreducible polynomials given by

$$p_1(u, v, w) = w^3 - 68u^3 - 11u^2w + 3u(30v^2 + 2w^2) + 18v^2w,$$
  
$$p_2(u, v, w) = w^3 - 20u^3 - 11u^2w + 2u(27v^2 + w^2) + 18v^2w.$$

We can easily see that  $p_1(u, v, w)$  is an irreducible polynomial. In fact, supposing that it is a product p(u, v, w)q(u, v, w) of some real linear and quadratic homogeneous polynomials of type

$$p(u, v, w) = au + bv + w$$
  

$$q(u, v, w) = gu^2 + huv + juw + kv^2 + lvw + w^2,$$




Figure 6:  $C^{J}(A_0)$  and  $C^{J}(A)$  for the matrices in Example 4.1

after performing this product and using Mathematica 13.0, from the equality  $p_1(u, v, w) = p(u, v, w)q(u, v, w)$ , we obtain a contradiction. Is is also clear that  $p_1(u, v, w)$  is not the product of three real linear homogeneous polynomials. Analogousgly, we can prove that  $p_2(u, v, w)$  is irreducible. As displayed in Figure 6,  $C^J(A)$  contains two hyperbolic-like curves, which are not hyperbolas, and two deltoids in its interior. The left-most blue curve and right-most red curve represent the boundaries of the convex components  $W_+^J(A)$  and  $-W_-^J(A)$ , respectively, that is,  $W^J(A)$  is bounded that these two curves. Now, the eigenvaluess of A are marked as the green dots.

After several experimental tests, we found that this type of shape is kept for the Krein space numerical range of matrices of type  $A' = T_6(\mathbf{c}, \mathbf{a}, \mathbf{b}')$  with  $\mathbf{b}' = (1, 2, b_3, 4, 5)$ , considering different values of  $b_3$  in a neighbourhood of 0, and that the deltoids are getting smaller and smaller as we consider values of  $b_3$  approaching 0, indicating that we are approaching hyperbolicty of the set, which only occurs for  $b_3 = 0$ .

#### Acknowledgment

The first author was partially supported by the Centre for Mathematics of the University of Coimbra (funded by the Portuguese Government through FCT/MCTES, DOI 10.54499/UIDB/00324/2020). The second author was supported by the Center for Research and Development in Mathematics and Applications (CIDMA) through the Portuguese Foundation for Science and Technology (FCT - Fundação para a Ciência e a Tecnologia), project UIDB/04106/2020 (https://doi.org/10.54499/UIDB/04106/2020).

The third author was financed by CMAT-UTAD through the Portuguese Foundation for Science and Technology (FCT - Fundação para a Ciência e a Tecnologia), reference UIDB/00013/2020, (https://doi.org/10.54499/UIDB/00013/2020).

#### References

- T. Ya. Azizov and I. S. Iokhvidov, Linear Operators in Spaces with an Indefinite Metric, Nauka, Moscow, 1986, English Translation: Wiley, New York, 1989.
- [2] N. Bebiano, R. Lemos, J. da Providência and G. Soares, Remarks on generalized numerical ranges of operators on indefinite inner product spaces *Online Proceedings of SIAM Applied Linear Algebra Conference*, College of William and Mary, Williamsburg (2003) 15–19.
- [3] N. Bebiano, R. Lemos, J. da Providência and G. Soares, On generalized numerical ranges of operators on a indefinite inner product space, *Linear Multilin. Algebra* **52** (2004), 203–233.
- [4] N. Bebiano, R. Lemos and J. da Providência, Numerical ranges of unbounded operators arising in quantum physics, *Linear Algebra Appl.* **381** (2004), 259–279.
- [5] N. Bebiano, R. Lemos, J. da Providência and G. Soares, On the geometry of numerical ranges in spaces with an indefinite inner product, *Linear Algebra Appl.* **399** (2005), 17–34.
- [6] N. Bebiano, R. Lemos and G. Soares, On the hyperbolicity of the Krein space numerical range, Linear Multilin. Algebra 52 (2023), 1–21.
- [7] N. Bebiano, R. Lemos and G. Soares, On the numerical range of Kac-Sylvester matrices, *Electron. J. Linear Algebra* **39** (2023), 241–259.
- [8] N. Bebiano, R. Lemos and G. Soares, Algebraic curves associated with centrosymmetric matrices of orders up to 6. Adv. Oper. Theory 9, 56 (2024).
- [9] N. Bebiano, H. Nakazato, R. Lemos, J. da Providência and G. Soares, Inequalities for J-Hermitian matrices, Linear Algebra Appl. 407 (2005), 67–84.
- [10] N. Bebiano and J. da Providência, Numerical ranges in physics, Linear and Multilin. Algebra 43 (4) (1998), 327–337.
- [11] N. Bebiano, J. da Providência, A. Nata and G. Soares, Krein spaces numerical ranges and their computer generation, *Electr. J. Linear Algebra* 17 (2008), 192–208.

- [12] N. Bebiano, J. da Providência, I. Spitkovsky and V. Kenya, Kippenhahn curves of some tridiagonal matrices, Filomat 35 (2021), 3047–3061.
- [13] N. Bebiano, J. da Providência and R. Teixeira, Indefinite numerical range of 3 × 3 matrices. Czech Math. J. 59 (2009), 221—239.
- [14] N. Bebiano, J. da Providência and R. Teixeira, Flat portions on the boundary of the indefinite numerical range of 3 × 3 matrices, *Linear Algebra Appl.* **428** (2008), 2863–2879.
- [15] E. Brown and I. M. Spitkovsky, On matrices with elliptical numerical ranges, *Linear Multilin*. Algebra 52 (2004), 177–193.
- [16] T. Geryba and I. M. Spitkovsky, On the numerical range of some block matrices with scalar diagonal blocks, *Linear Multilin. Algebra* 69: 5 (2021), 772–785.
- [17] I. Gohberg, P. Lancaster and L. Rodman, Matrices and Indefinite Scalar Products, OT 8, Birkäuser Verlag, Basel, 1983.
- [18] A. Guterman, R. Lemos and G. Soares, More on geometry of Krein space C-numerical range, *Appl. Math. Comput.* **352** (2019), 258–269.
- [19] F. Hausdorff, Der Wertvorrat einer Bilinearform. Math. Z. 3 (1919), 314–316.
- [20] M. Jiang and I. M. Spitkovsky, On some reciprocal matrices with elliptical components of their Kippenhahn curves, Special Matrices 10 (2022), 117–130.
- [21] D. S. Keeler, L. Rodman and I. M. Spitkovsky, The numerical range of 3 × 3 matrices, *Linear Algebra Appl.* 252 (1997), 115–139.
- [22] C.-K. Li and L. Rodman, Remarks of numerical ranges of operators in spaces with an indefinite inner product, *Proc. Amer. Math. Soc.* **126** (1998), 973–982.
- [23] C.-K. Li and L. Rodman, Shapes and computer generation of numerical ranges of Krein space operators, *Electr. J. Linear Algebra*, **3** (1998), 31–47.
- [24] C.-K. Li, N.-K. Tsing and F. Uhlig, Numerical ranges of an operator on an indefinite inner product space, *Electr. J. Linear Algebra* 1 (1996), 1–17.
- [25] F. Philipp and C. Trunk, The numerical range of non-negative operators in Krein spaces, *Linear Algebra Appl.* 438 (2013), 2542–2556.
- [26] O. Toeplitz, Das algebraische Analogon zu einem Satze von Fejér. Math. Z. 2 (1918), 187–197.