Rings of simple range 2 B.V. Zabavsky, O.M. Romaniv, A.V. Sagan

Corresponding author: O.M. Romaniv

Address: Ivan Franko National University of Lviv, Ukraine (O.M. Romaniv, A.V. Sagan)

Abstract: We introduce the concept of rings of simple range 2. Based on this concept, we build a theory diagonal reduction of matrices over Bezout domain. In particular we show that invariant Bezout domain is an elementary divisor ring if and only if it is a rings of simple range 2.

Keywords and phrases: Bezout ring, Hermite ring, stable range, diagonal reduction, simple ring, simple range

Mathematics Subject Classification: 06F20, 13F99, 13E15

1 Introduction

The study of linear groups over fields goes back to the mid-19th century. However, practically nothing was known about the ring except for semilocal and some types of arithmetic rings until the mid-1960s. A real commenting revolution was initiated by work H. Bass [2]. In particular, H. Bass introduced a new concept of dimension of rings, a stable range and found that the main results on the structure of complete linear group GL_n over the field with the necessary changes are transferred to rings, whose stable range is less than n. Note that a stable range is also closely related to the issues of stable structure of projective modules. In modern algebraic research there is a close connection between the concept of a stable range of ring and the open questions of diagonalization of matrices over these rings. The problem of diagonalization of matrices is a classic one. An overview can be found in [10]. It is known [10, Theorem 1.2.40] that the stable range of an elementary divisor ring did not exceed 2. I. Kaplansky noted that a commutative Bezout ring of stable range 1 is an elementary divisor ring [6]. M. Henriksen showed that a unit regular ring is a ring over which the matrices are diagonalized [5]. Even

in the case of commutative rings, the bound of stable range is found to be cramped. In particular [9, 11, 12] introduced a generalization of stable range of rings (rings of Gelfand range 1, rings of dyadic range 1) which allowed to reduce tha problem of complete description of the elementary divisor rings to the question of existence of a nontrivial Gelfand element [11]. In this paper we introduce the concept of a ring of simple range 2, which allows us to describe the new classes of noncommutative elementary divisor rings.

2 Notations and preliminary results

Let R be an associative ring with non-zero unit. We say that matrices A and B over a ring R are equivalent if there exist invertible matrices P and Q over R such that B = PAQ. If a matrix A over R is equivalent to a diagonal matrix $D = (d_{ii})$ with the property that d_{ii} is a total divisor of $d_{i+1,i+1}$ (i.e. $Rd_{i+1,i+1}R \subset d_{i,i}R \cap Rd_{i,i}$), then we say that A admits a canonical diagonal reduction. A ring R over which every matrix admits a canonical diagonal reduction is called an *elementary divisor ring*. A ring R is called *right (left)* Hermite if each matrix $A \in R^{1\times 2}$ ($A \in R^{2\times 1}$) admits a diagonal reduction. A ring which is right and left Hermite is called a *Hermite ring*. Obviously a commutative left (right) Hermite ring is a Hermite ring. Moreover, each elementary divisor ring is Hermite, and a right (left) Hermite ring is a right (left) *Bézout ring*, i.e. a ring in which any finitely generated right (left) ideal of R is principal [10].

A row $(a_1, \ldots, a_n) \in \mathbb{R}^n$ is called *unimodular* if $a_1R + a_2R + \cdots + a_nR = R$. An unimodular *n*-row $(a_1, \ldots, a_n) \in \mathbb{R}^n$ over a ring *R* is called *reducible* if there exist a (n-1)-row $(b_1, \ldots, b_{n-1}) \in \mathbb{R}^{n-1}$ such that

$$(a_1 + a_n b_1, a_2 + a_n b_2, \dots, a_{n-1} + a_n b_{n-1}) \in \mathbb{R}^{n-1}$$

is unimodular. If $n \in \mathbb{N}$ is the smallest number such that any unimodular (n+1)-row is reducible, then R has stable range n, where $n \geq 2$. In particular, a ring R has stable range 1 if aR + bR = R implies that (a + bt)R = R for some $t \in R$. A ring R is a ring of stable range 2 if aR + bR + cR = R implies that (a + cx)R + (b + cy)R = R for some $x, y \in R$. A right (left) Hermite ring is a ring of stable range 2 [10, Theorem 1.2.40]. In the case of commutative Bezout we have a results.

Theorem 1. [10, Theorem 2.1.2] A commutative Bezout ring is a Hermite ring if and only if it is a ring of stable range 2. This is an open question: is right (left) Bezout ring of stable range 2 a right (left) Hermite ring?

By [1] we have that Bezout domain is a Hermite ring, i.e. Bezout domain is a ring of stable range 2.

In the future we will consider rings in which for any $a \in R$ we have $RaR = a_*R = Ra_*$ for some $a_* \in R$. Examples of such ring are simple rings [10, Section 4.2], quasi-duo elementary divisor rings, semi-local semi-prime elementary divisor rings [3, Theore 1], principal ideal domains. Condition $RaR = a_*R = Ra_*$ for any $a \in R$ is called *Dubrovin condition* [10]. Recall that nonzero element $a \in R$ is right (left) invariant if aR (Ra) is two-sided ideal, if both conditions hold that is, if aR = Ra then a is called invariant element. If in a domain R every factor of an invariant element is invariant we say that R is a domain with Komarnytsky condition. Obvious example of domain with Komarnytsky condition is a simple domain and domain $\mathbb{H}[x]$, where \mathbb{H} is quaternion divisor ring, we have invariant element $x^2 + 1$ but not any factor of it is not invariant. D-K elementary divisor rings are closely related with rings with Dubrovin and Komarnytsky conditions (hence forth, rings with Dubrovin-Komarnytsky (D-K) condition).

Definition 1. Ring R is called an D-K elementary divisor ring if an arbitrary matrix A over R is the equivalent matrix $\operatorname{diag}(\varepsilon_1, \varepsilon_2, \ldots, \varepsilon_r, 0, \ldots, 0)$ where $R\varepsilon_{i+1}R \subset \varepsilon_i R \cap R\varepsilon_i$ for all $i \in 1, r-1$ and $\varepsilon_1, \varepsilon_2, \ldots, \varepsilon_{r-1}$ are invariant elements.

A simple elementary divisor domain is an example of D-K elementary divisor ring. By [10, Theorem 4.1.1] a simple Bezout domain is an elementary divisor ring if and only if it is 2-simple domain. i.e. for any $a \in R \setminus \{0\}$ we have that $u_1 a v_1 + u_2 a v_2 = 1$ for some $u_1, u_2, v_1, v_2 \in R$. Consequently we have a next result.

Corollary 1. Simple Bezout domain is D-K elementary divisor ring if and only if it is 2-simple domain.

Let R be a simple ring. Clearly $RaR = \{\sum_{i=1}^{n} u_i a v_i \mid u_i, v_i \in R\} = R$ for each $a \in R \setminus \{0\}$. Then we have $n \in \mathbb{N}$ such that

$$u_1 a v_1 + u_2 a v_2 + \dots + u_n a v_n = 1.$$
(1)

If for all $a \in R \setminus \{0\}$ there exists a minimal n which satisfies (1), then R is called *n*-simple.

Description D-K elementary divisor ring is quite an interesting and extremes challenging task for future usearch. Just note the next open task: is there *n*-simple Bezout domain where $n \ge 3$?

3 Rings of simple range 2

Definition 2. We say a ring R is a ring of simple range 2 if RaR + RbR + RcR = R where $c \neq 0$ implied R(pa + qb)R + RpcR = R for some $p, q \in R$.

Proposition 1. Let R be Bezout ring. The following statements are equivalent:

- **1)** R is a ring of simple range 2;
- 2) $RaR+RbR+RcR, c \neq 0$, implied (pa+qb)R+pcR = dR where RdR = Rfor some $p, q \in R$.

Proof. Let RaR + RbR + RcR = R, $c \neq 0$, implied (pa + qb)R + pcR = dRwhere RdR = R for some $p, q \in R$. Since $pcR \subseteq RpcR$ and $(pa + qb)R \subseteq R(pa + qb)R$ we have $(pa + qb)R + pcR \subseteq R(pa + qb)R + RpcR$. So on RdR = R and $d \in R(pa + qb)R + RpcR$ we have R(pa + qb)R + RpcR = R.

Let condition RaR + RbR + RcR = R, $c \neq 0$, implied R(pa + qb)R + RpcR = R for some $p, q \in R$. Since R is a Bezout ring then (pa + qb)R + pcR = dR for some $d \in R$. Since $(pa + qb)R \subset dR$ and $pcR \subseteq RdR$ and $R(pa + qb)R \subseteq RdR$. So on R(pa + qb)R + RpcR = R we have that RdR = R.

Obviously a simple ring is ring of simple range 2.

A ring R is called *purely infinite* if $R \cong R^2$ as R-modules. For example for any ring $S = End_R(R^{\infty})$ is purely infinite, where R^{∞} denotes an infinite direct product of R. This ring is simple ring, i.e. a ring of simple range 2 which have infinite stable range. Nevertheless we have a next result.

Theorem 2. A Bezout ring of stable range 1 is a ring of simple range 2.

Proof. Let RaR + RbR + RcR = R, $c \neq 0$. Since R is a Bezout ring we have aR + bR = dR, $a = da_1$, $b = db_1$ and au + bv = d for some $d, a_1, b_1, u, v \in R$. Then $d(1 - a_1u - b_1v) = 0$. Let $c = 1 - a_1u - b_1v$ we have $a_1R + b_1R + cR = R$ and dc = 0. Since any ring of stable range 1 is a ring of stable range 2 we have that $(a_1 + c\lambda)R + (b_1 + c\mu)R = R$ for some $\lambda, \mu \in R$. Let $a_0 = a_1 + c\lambda$, $b_0 = b_1 + c\mu$, then $a_oR + b_0R = R$ and $a = da_0$, $b = db_0$. Since R is a ring of stable range 1 we have $(b_0\lambda + a_0)R = R$. i.e. $b_0\lambda + a_0 = u$ is invertible element R. Since RaR + RbR + RcR = R and $b\lambda + a = du$ we have $(b\lambda + a)R = duR = dR$ and $(b\lambda + a)R \subseteq R(b\lambda + a)R$, $dR \subseteq RdR$ and RaR + RbR = RdR. Then $R(b\lambda + a)R + RcR = RaR + RbR + RcR = R$. i.e. R is a ring of simple range 2.

Theorem 3. D-K elementary divisor domain is a ring of simple range 2.

Proof. Let R be a D-K elementary divisor domain and RaR + RbR + RcR = R, $c \neq 0$. Consider a matrix $A = \begin{pmatrix} a & c \\ b & 0 \end{pmatrix}$. Let $b \neq 0$, then

$$PAQ = \begin{pmatrix} z & 0\\ 0 & \alpha \end{pmatrix} \tag{2}$$

where $R\alpha R \subset zR \cap Rz$, for some invertible matrices P and Q. Since $b \neq 0$ we have that $\alpha \neq 0$. Since RaR + RbR + RcR = R and $R\alpha R \subset aR \cap Rz$ according to (2) we have R = RaR + RbR + RcR = zR = Rz, i.e. z is an invertible element R. We denote $P = \binom{p \ q}{**}$. According to (2) we have (pa + qb)R + pcR = R. By Proposition 1 we have R(pa + qb)R + RpcR = R. If b = 0 we have R(a + b)R + RcR = R. We proved that R is a ring of simple range 2.

We will note that if R is a Bezout domain and RaR = R and if $aR \subset bR$ or $Ra \subset Rc$ we have that RbR = R and RcR = R. We have a next result.

Proposition 2. Let R be a Bezout domain with D-K condition. If RaR = R and RbR = R then RabR = R.

Proof. In R there exists $a, b \in R$ such that RaR = R and RbR = R such that $RabR = \alpha R = R\alpha$ where α is not invertible element of R. Since R is a Bezout domain we have $bR + \alpha R = \beta R$ for some $\beta \in R$. Since $\alpha R \subset \beta R$ we have that β is an invariant element. Since $bR \subset \beta R$ and RbR = R we have that β is invertible element R.

Then we have that $bu + \alpha v = 1$ for some $u, v \in R$. Since $abu + a\alpha = a$ and $RabR = \alpha R = R\alpha$ we have that $ab = s\alpha$ for some $s \in R$. Then $su'\alpha + a\alpha = a$ where $\alpha u = u'\alpha$. So on we have $Ra \subset R\alpha$ and $R = RaR \subset R\alpha R$ we have that α is an invertible element. The contradiction obtained proves Proposition 2.

Let R be a Bezout domain of simple range 2, i.e. RaR + RbR + RcR = R, $c \neq 0$, imply R(pa + qb)R + RpcR = R.

Remark 1. According to Proposition 1, Proposition 2 and the fact then R is Bezout domain we can assume that it is equality R(pa+qb)R+RpcR=R elements p, q such that pR+qR=R. If pR+qR=tR, then $p=tp_0, q=tq_0, p_0R+q_0R=R$ for some $t, p_0, q_0 \in R$. Since R(pa+qb)R+RpcR=R we have RtR=R and $R(p_0a+q_0b)R+Rp_0cR=R$.

We will need the next ancient result in the future.

Proposition 3. [7, Lemma 1] Let R be a Bezout domain and pR + qR = R, Ru + Rv = R for some $p, q, u, v \in R$. Then there exist invertible matrices P and Q of the form

$$P = \begin{pmatrix} p & q \\ * & * \end{pmatrix}, \quad Q = \begin{pmatrix} u & * \\ v & * \end{pmatrix}.$$

As a result we get the main result of this paper in terms of constructing the theorem of matrix diagonalization over noncommutative rings.

Lemma 1. Let R be a Bezout domain of simple range 2. Then any matrix $A = \begin{pmatrix} a & c \\ b & 0 \end{pmatrix}$ where $c \neq 0$ and RaR + RbR + RcR = R is equivalent to matrix $X = \begin{pmatrix} x & z \\ y & 0 \end{pmatrix}$ such that RxR = R.

Proof. Since RaR + RbR + RcR = R and R is a ring of simple range 2, then (pa + qb)R + pcR = R where RxR = R. We have

$$\begin{pmatrix} p & q \end{pmatrix} \begin{pmatrix} a & c \\ b & 0 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} = x$$

where (pa+qb)u+pcv = x for some $u, v \in R$. By Remark 1 and Proposition 3 we have

$$PAQ = \begin{pmatrix} x & * \\ * & * \end{pmatrix}$$

for some invertible matrices

$$P = \begin{pmatrix} p & q \\ * & * \end{pmatrix}, \quad Q = \begin{pmatrix} u & * \\ v & * \end{pmatrix}.$$

According to Theorem 3 D-K elementary divisor domain is a ring of simple range 2. In what follows we shall prove that for invariant Bezout domain the condition of a simple range 2 is sufficient, that is, the following theorem holds.

Theorem 4. Let R be invariant Bezout domain of simple range 2. Then R is D-K elementary divisor ring.

Proof. Invariant Bezout domain is obvious example of a domain with D-K conditions. According to [4, Proposition 1] in order to prove Theorem 4 we need to prove that matrix $A = \begin{pmatrix} a & c \\ b & 0 \end{pmatrix}$ where $c \neq 0$ and RaR + RbR + RcR = R hold canonical diagonal reduction. By Lemma 1, matrix A is equivalent to matrix $X = \begin{pmatrix} x & y \\ 0 & z \end{pmatrix}$ where RxR = R. Since R is an invariant domain we have that x is an invertible element. Elementary transformations of row and columns matrix X will be reduced to appearance $D = \begin{pmatrix} 1 & 0 \\ 0 & \Delta \end{pmatrix}$ for some $d \in R$, i.e. we proved that a matrix A hold a canonical diagonal reduction.

References

- Amitsur, S. A. (1963). Remarks on principal ideal rings. Osaka Math. J. 15: 59–69.
- [2] Bass, H. (1964). K-theory and stable algebra. I.H.E.S. 22: 5–60.
- [3] Dubrovin, N. I. (1986). On rings with elementary divisors. Soviet Math. 30(11): 16-24.
- [4] Gatalevich, A. I., Zabavsky, B. V. (1999). Noncommutative elementary divisor rings. J. Math. Sci. 96: 3013–3016.
- [5] Henriksen, M. (1973). On a class of regular rings that are elementary divisor rings. Arch. Math. 24: 133–141.
- [6] Kaplansky, I. (1954). Infinite abelian groups. Michgan Univ. of Michigan Press 96p.
- [7] Williams, R. E. (1968). A note on weak Bezout rings. Proc. Amer. Math Soc. 19: 951–952.
- [8] Zabavsky, B. V., Komarnytsky, M. Ya. (1990). Distributive elementary divisor domains. Ukr. Math. J. 42(7): 890–892.

- Zabavsky, B. V., Romaniv, O. M. (2019). A Bezout ring of stable range 2 which has square stable range 1. *Commun. Algebra* 47(12): 5392–5397; doi: 10.1080/00927872.2019.1623239
- [10] Zabavsky, B. V. (2012). Diagonal reduction of matrices over rings. Mathematical Studies Monograph Series, Vol. XVI. Ukraine: VNTL Publisher.
- [11] Zabavsky, B. V. (2017). Conditions for stable range of an elementary divisor rings. Commun. Algebra 45(9): 4062–4066; doi: 10.1080/00927872.2016.1259418
- [12] Zabavsky, B. V. (2019). Rings of dyadic rings 1, J. Alg. Appl. 18(11): 1950206; doi: 10.1142/S0219498819502062