

# QUADRATIC DIFFERENTIAL EQUATIONS IN TWO VARIABLES FREE OF MOVABLE CRITICAL POINTS

ADOLFO GUILLOT

ABSTRACT. We study ordinary differential equations in the complex domain, and investigate the absence of movable critical points (the Painlevé property) for systems of non-autonomous polynomial quadratic differential equations in two variables. We classify the systems having this property by studying, when it is possible, some obstructions described by Malmquist, and by reducing them to second-order equations of the first degree in one variable, for which the analysis of Painlevé and Gambier can be applied. We find that, in many cases, overcoming Malmquist's obstructions guarantees the absence of movable critical points.

## 1. INTRODUCTION

Within the study of ordinary differential equations in the complex domain, the following problem is attributed to Goursat: *to classify the systems of equations*

$$(1) \quad \begin{aligned} x' &= P(x, y, t), \\ y' &= Q(x, y, t), \end{aligned}$$

with  $P$  and  $Q$  rational in  $x$  and  $y$ , analytic in the independent variable  $t$ , which are free of movable critical points, with two systems differing by a time-dependent birational transformation in the dependent variables considered equal [Gar60]. Following Painlevé, a *critical point* of a function is a singular point around which at least two of its branches are permuted (see the first of his *Leçons de Stockholm* [Pai73]). The above is a natural and broad generalization of the problem, solved by Painlevé and Gambier, concerning second-order equations in one variable. Garnier wondered if such general systems could define new transcendents; he estimated this to be a difficult problem, and gave a first step towards its solution, by addressing it in the case where (1) is autonomous, and homogeneous in the dependent variables [Gar60].

One may restrict this problem by considering systems (1) with  $P$  and  $Q$  polynomial in  $x$  and  $y$  (up to either affine or polynomial automorphisms). These polynomial equations form a rich class, which, as shown by Okamoto, already contains all six Painlevé transcendents [Oka79]. Goffar-Lombet investigated the question for polynomial systems having some particular homogeneous polynomials, of degrees four and five, as highest-degree terms, obtaining precise and complete results for such systems [GL74]. In 1980, Kimura and Matuda announced a classification of polynomial systems of the form (1), free of movable critical points, of degree up to five, and reported that there were no novel ways of integrating these beyond

---

*Key words and phrases.* Absence of movable critical points, Painlevé Property, Painlevé transcendent.

2020 *Mathematics Subject Classification:* 34M04, 34M55.

elementary ones, and those given by the Painlevé transcendents [KM80]. To the best of our knowledge, these results were never published.

In this article, we classify the nonlinear systems of equations of the form (1), with  $P$  and  $Q$  polynomials in  $x$  and  $y$  of degree at most two, that are free of movable branch points, up to time-dependent affine transformations. We describe the integration of these equations: they either reduce to a succession of two Riccati (or linear) equations, or are birationally equivalent to rational second-order equations in one variable, falling within Painlevé's analysis. We thus bring to completion Kimura and Matuda's announcement for the quadratic case, and confirm their observation that only the Painlevé I, II and IV transcendents are needed to integrate quadratic systems.

Our tools are those used by Painlevé, Goffar-Lombet, and Kimura and Matuda, and will be reviewed in Section 2. The systems may be grouped following the linear equivalence class of their quadratic parts, which may be supposed to be constant. The admissible classes of quadratic parts, which we will call *types*, constitute eight equations plus two countable families, and appear in Table 1. This homogeneous part has some special solutions, to some of which we may associate the necessary condition for the absence of movable critical points studied by Malmquist; this condition is of an algebro-differential nature, and may be explicitly calculated (see Section 2.2). After imposing these conditions, we may either solve the resulting equation by integrating one variable separately, or, in most cases, transform the system birationally into a second-order equation, where one can use the Painlevé-Gambier analysis to establish necessary and sufficient conditions for the absence of movable critical points. The calculation of Malmquist's conditions is probably not necessary in all cases, but allows us to join the Painlevé-Gambier analysis at a more advanced stage, and to give a concrete, intrinsic, geometric, and localized interpretation to some of the obstructions revealed by this analysis.

Equations of Type IV, which Goursat saw as natural generalizations of Riccati equations, are always free of movable critical points. It results from our analysis that, for systems of Types  $I_{3,3,3}$ ,  $I_{2,3,6}$ ,  $I_{2,4,4}$ ,  $I_{2,2,\infty}$ ,  $I_\infty$ ,  $II_1$ , and  $I_1$ ,  $I_2$ ,  $I_3$ , the Malmquist conditions, necessary for the absence of movable critical points, are also sufficient ones (we do not know if this is still the case for systems of Types  $I_n$  for  $n \geq 4$ ).

## 2. THE TOOLKIT

We now describe some of the tools we will use to establish the absence of movable critical points for the equations under consideration. In Section 2.1, we describe the obstructions found in the quadratic part of the equations. The Malmquist condition for the absence of movable critical points will be presented in Section 2.2. The background on second-order equations in one variable that we will need will be discussed in Section 2.3. Finally, in Section 2.4, we describe how to reduce the number of parameters involved in the equations.

**2.1. Quadratic homogeneous equations and univalence.** Consider a system of equations of the form (1), that we will write as

$$(2) \quad z' = \sum_{i=0}^2 X_i(z, t),$$

with  $z \in \mathbf{C}^2$ , and  $X_i$  homogeneous of degree  $i$  in  $z$  and holomorphic in  $t$ . For  $\alpha \neq 0$ , for  $\zeta = \alpha^{-1}z$ , and  $\tau = t_0 + \alpha t$ ,

$$\frac{d\zeta}{d\tau} = \sum_{i=0}^2 \alpha^{2-i} X_i(\zeta, t_0 + \alpha\tau).$$

The limit as  $\alpha \rightarrow 0$  is the autonomous, homogeneous equation  $d\zeta/d\tau = X_2(\zeta, t_0)$ . If the original equation is free of movable critical points, the latter will have only single-valued solutions [Pai00]. The first problem is thus to determine the quadratic homogeneous systems in two variables with single-valued solutions.

For a system of the form (1), with  $P$  and  $Q$  rational and homogeneous of the same degree,  $s = x/y$  is either constant, or a solution to an equation of the form

$$(3) \quad -\frac{s''}{(s')^2} = F(s),$$

and its univalence can be studied through the work of Briot and Bouquet [BB55]: the above expression has only simple poles, whose residues are either  $-1$ , or of the form  $1/m - 1$  for  $m \in \mathbf{Z}$  (see also Painlevé [Pai00, §10]; see [Gui06, Section 2.2] for a geometric interpretation). Based on this, Garnier [Gar60] established a birational classification of these equations. Finer classifications of polynomial systems, up to linear transformations, were given, on the one hand, by Goffar-Lombet in [GL74, Ch. I], and by Ghys and Rebelo in [GR97, Thm. C] (with a correction in [GR00]).

Let us discuss part of this classification through the notion of *Kowalevski exponent*, very much related to the obstructions that lead to Briot and Bouquet's classification. Let  $X$  be a quadratic homogeneous vector field on  $\mathbf{C}^n$  (we will sometimes consider the differential equations as vector fields). Let  $E$  denote Euler's vector field  $\sum z_i \partial_{z_i}$ . A *radial orbit* for  $X$  is a line through the origin of  $\mathbf{C}^n$  (a point in  $\mathbf{P}^{n-1}$ ) where  $E$  and  $X$  are collinear. If  $X$  has a radial orbit in the direction of the vector  $v \in \mathbf{C}^n$ , which is *non-degenerate* in the sense that  $X(v) \neq 0$ , and  $v$  is normalized so that  $X(v) = -v$ , a solution along this line is given by  $t \mapsto t^{-1}v$ . If  $X = \sum X_i \partial_{z_i}$ , let  $M$  be the constant matrix  $(\partial X_i / \partial z_j)|_v$ . The first equation of variations along this solution reads

$$(4) \quad t\zeta' = M\zeta.$$

The *Kowalevski exponents* of  $X$  at  $[v]$  are the eigenvalues of  $\mathbf{I} - M$  in the normal direction to  $v$  (which is an eigenvector of  $M$  associated to the eigenvalue  $-2$ ). These receive many other names: *resonances*, *Fuchs indices*, etc., and can be defined in many ways; see [Gor00].

If  $u$  is an eigenvector of  $M$  associated to the eigenvalue  $\lambda$ , a solution of (4) is given by  $t^\lambda u$ . Hence, *for a quadratic homogeneous vector field to be univalent, it is necessary that the Kowalevski exponents of its non-degenerate radial orbits are integers*. For example, for the vector field  $x_1(x_1 \partial_{x_1} + \sum_{i>1} m_i x_i \partial_{x_i})$ , for the radial orbit along the normalized vector  $v = (-1, 0, \dots, 0)$ , the Kowalevski exponents are  $1 - m_2, \dots, 1 - m_n$ . In dimension two, a Kowalevski exponent  $\nu$  corresponds to a residue  $1/\nu - 1$  in the associated simple pole of (3).

On  $\mathbf{C}^2$ , for a quadratic homogeneous vector field  $X$ , if not every line through the origin is a radial orbit (if  $X$  and  $E$  are not collinear), there are three radial orbits (counted with multiplicity). Let us convene to say that a simple radial orbit along which  $X$  vanishes identically has Kowalevski exponent  $\infty$ . If a quadratic

Type	equation	rad. orbs.	exponents
I <sub>3,3,3</sub>	$x(x-2y)\partial_x + y(y-2x)\partial_y$	3	3, 3, 3
I <sub>2,3,6</sub>	$x(2x-5y)\partial_x + y(y-4x)\partial_y$	3	2, 3, 6
I <sub>2,4,4</sub>	$x(x-3y)\partial_x + y(y-3x)\partial_y$	3	2, 4, 4
I <sub>2,2,\infty</sub>	$(x-y)(x\partial_x - y\partial_y)$	3	2, 2, $\infty$
I <sub>n</sub>	$x^2\partial_x + y(y-(n-1)x)\partial_y, \quad n \geq 1$	3	1, $n, -n$
I <sub><math>\infty</math></sub>	$x(x-y)\partial_x$	3	1, $\infty, \infty$
II <sub>n</sub>	$x(x\partial_x - (n-1)y\partial_y), \quad n \neq 0$	2	$n$
II <sub><math>\infty</math></sub>	$xy\partial_x$	2	$\infty$
III	$y^2\partial_x$	1	—
IV	$x(x\partial_x + y\partial_y)$	$\infty$	—

TABLE 1. The quadratic homogeneous equations in  $\mathbf{C}^2$  along with the Kowalevski exponents of its simple radial orbits ( $n \in \mathbf{Z}$ ).

homogeneous vector field  $X$  has three radial orbits, with Kowalevski exponents  $\nu_i$ ,

$$(5) \quad \frac{1}{\nu_1} + \frac{1}{\nu_2} + \frac{1}{\nu_3} = 1$$

(this follows, for instance, from the residue theorem).

A generic quadratic homogeneous vector field on  $\mathbf{C}^2$  is determined, up to linear equivalence, by its Kowalevski exponents. (These vector fields form a six-dimensional vector space, on which the four-dimensional group of linear transformations of  $\mathbf{C}^2$  acts linearly; the symmetric functions of the Kowalevski exponents are functions on the associated quotient).

The linear equivalence classes the univalent quadratic homogeneous vector fields on  $\mathbf{C}^2$  appears in Table 1. The classification of the univalent vector fields with three different radial orbits starts by solving (5) as a Diophantine equation (admitting that  $\nu_i$  can take the value  $\infty$ ), from which the first six lines of Table 1 are obtained (the corresponding vector fields turn out to be univalent). The other lines of this table follow from organizing quadratic homogeneous vector fields according to the multiplicities of their radial orbits and to the Kowalevski exponents of the simple ones, and from calculating the expression (3).

**2.2. Malmquist's conditions.** Consider the system of equations:

$$(6) \quad \begin{aligned} y \frac{dx}{dt} &= na(t)x + b(t)y + \sum_{i+j \geq 2} f_{ij}(t)x^i y^j, \\ \frac{dy}{dt} &= a(t) + \sum_{i+j \geq 1} g_{ij}(t)x^i y^j \end{aligned}$$

with holomorphic right-hand side, with  $n \in \mathbf{Z}$ ,  $n \geq 1$ , and  $a$  a non-vanishing function. We have the following result:

**Proposition 1** (Malmquist). *In the case  $n = 1$ , for the system (6) to be free of movable branch points, it is necessary that  $b \equiv 0$ .*

We refer the reader to Malmquist's work for its proof [Mal21, §11]. The idea is, first, to establish, via a theorem of Darboux, that, for every  $t_0$ , there exists an invariant surface in the extended phase space of the form  $t = F(x, y)$  with  $F(0, 0) = t_0$ , and that, in restriction to this surface,  $x$ , as a function of  $y$ , satisfies

an equation of Briot-Bouquet type; the solution of the latter will have logarithmic terms unless  $b(t_0) = 0$ .

If  $n > 1$ , we may, in the extended phase space, blow up the curve  $(x, y) = (0, 0)$ : for

$$u = \frac{b}{(n-1)a},$$

and for  $\xi$  defined by  $x = (\xi - u)y$ , the system becomes

$$\begin{aligned} y \frac{d\xi}{dt} &= (n-1)a\xi + yh(\xi, y), \\ \frac{dy}{dt} &= a + \sum_{i+j \geq 1} g_{ij}(\xi - u)^i y^{i+j}, \end{aligned}$$

for

$$h(\xi, y) = u' + \sum_{i+j \geq 2} f_{ij}(\xi - u)^i y^{i+j-2} - \sum_{i+j \geq 1} g_{ij}(\xi - u)^i y^{i+j-1}.$$

The transformed system has the same form as the original, but, in it,  $n$  has been replaced by  $n - 1$ . By repeatedly applying this procedure, we arrive to the case  $n = 1$ , where we can then apply the criterion of Proposition 1.

The context in which we will apply this criterion is the following one. For an equation of the form (2) whose quadratic part at  $t = t_0$  has a radial orbit in the direction  $[a : b]$  with positive Kowalevski exponent  $n$ , by extending  $\mathbf{C}^2 \times U$  into  $\mathbf{P}^2 \times U$  by  $((x, y), t) \mapsto ([x : y : 1], t)$  the resulting system will have poles along the line at infinity for all values of  $t$ , with a line of singularities at  $\{[a : b : 0]\} \times U$ , along which the system will have the form (6). Malmquist's condition may be calculated after some successive blow-ups are performed.

**2.3. The Painlevé-Gambier classification.** We will need to state conditions for the absence of movable critical points in some equations of the form

$$(7) \quad u'' = \left(1 - \frac{1}{m}\right) \frac{(u')^2}{u} + \left(au + b + \frac{c}{u}\right) u' + du^3 + eu^2 + fu + g + \frac{h}{u},$$

with  $m \in \mathbf{Z}^*$ , in which the coefficients are analytic functions of the independent variable  $t$ . This falls within the problem solved by Painlevé and Gambier's titanic enterprise, leading to the discovery of the Painlevé transcendents [Pai02, Gam10].

It is natural to consider these equations up to changes of coordinates of the form

$$(8) \quad u(t) = \lambda(t)w(\tau), \quad \frac{d\tau}{dt} = \rho(t).$$

Such a change of coordinates transforms (7) into an equation

$$\frac{d^2w}{d\tau^2} = \left(1 - \frac{1}{m}\right) \frac{1}{w} \left(\frac{dw}{d\tau}\right)^2 + \left(a_1w + b_1 + \frac{c_1}{w}\right) \frac{dw}{d\tau} + d_1w^3 + \dots + \frac{h_1}{w},$$

of the same form, in which

$$(9) \quad \begin{aligned} a_1 &= \frac{\lambda}{\rho} a, \quad b_1 = \frac{1}{\rho} \left( b - \left(\frac{2}{m}\right) \frac{\lambda'}{\lambda} - \frac{\rho'}{\rho} \right), \quad c_1 = \frac{1}{\lambda\rho} c, \quad d_1 = \frac{\lambda^2}{\rho^2} d, \\ e_1 &= \frac{\lambda}{\rho^2} \left( e + a \frac{\lambda'}{\lambda} \right), \quad f_1 = \frac{1}{\rho^2} \left( f + b \frac{\lambda'}{\lambda} - \frac{1}{m} \left(\frac{\lambda'}{\lambda}\right)^2 - \left(\frac{\lambda'}{\lambda}\right)' \right), \\ g_1 &= \frac{1}{\lambda\rho^2} \left( g + c \frac{\lambda'}{\lambda} \right), \quad h_1 = \frac{1}{\lambda^2\rho^2} h \end{aligned}$$

(this corrects a slight misprint in table (F) in [Gam10, p. 26]). Observe that, in equation (7), the vanishing of either  $a$ ,  $c$ ,  $d$ , or  $h$  is an intrinsic condition, and that the ratios  $a^2 : d$  and  $c^2 : h$  are also intrinsic. The condition  $b \equiv 0$  is preserved by the changes of coordinates for which  $\lambda^2 \rho^m$  is constant.

When  $m = 1$ ,  $c$  and  $h$  must vanish if the equation is free from movable branch points [Pai00, §11], and, in this case, the changes of coordinates of the form  $u \mapsto u + v(t)$  should also be taken into account.

We will sometimes use Ince's systematic presentation of Painlevé and Gambier's results as a flowchart for deciding if a given equation is free of movable branch points [Inc44, Ch. XIV].

**2.4. Normalizations.** In order to alleviate our calculations, it will be useful to consider equations with few parameters.

**Proposition 2.** *Consider an equation of the form (2). If the linear equivalence class of  $X_2(z, t)$  is independent of  $t$  in neighborhood of  $t_0$  then, after a suitable time-dependent affine map, we may assume that, up to restricting the equation to a neighborhood of  $t_0$ , the following conditions hold:*

- (1)  $X_0$  is identically zero;
- (2)  $X_2$  is independent of  $t$ ;
- (3) if  $X_1 = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ , and  $\alpha, \beta, \gamma$  and  $\delta$  are constants with  $\alpha \neq \delta$ ,  $\alpha A + \beta B + \gamma C + \delta D = 0$ ; furthermore,  $X_1$  can be multiplied by any nonzero constant.

*Proof.* (1) For any function  $u(t)$ ,  $\zeta = z - u$  satisfies the equation

$$\zeta' = X_2(z + u, t) + X_1(z + u, t) + X_0(t) + u'.$$

Thus, if  $u$  is a solution to the equation

$$u' + X_2(u, t) + X_1(u, t) + X_0(t) = 0,$$

there are no terms of degree 0 in the equation satisfied by  $\zeta$ . We will henceforth suppose that  $X_0 \equiv 0$ .

(2) Let  $M(t)$  be a linear transformation such that  $MX_2(M^{-1}\zeta, t) = X_2(\zeta, t_0)$ . Then, for  $\zeta = M(t)z$ ,

$$\frac{d\zeta}{dt} = MX_2(M^{-1}\zeta, t) + MX_1(M^{-1}\zeta, t) + M'M^{-1}\zeta,$$

whose quadratic term is constant. From now on,  $X_2$  will be supposed to be constant.

(3) For a non-vanishing function  $\rho(t)$ , for  $\zeta(\tau) = \rho(t)z(t)$  with  $\tau'(t) = \rho(t)$ , the equation gets transformed into

$$\frac{d\zeta}{d\tau} = X_2(\zeta) + \rho X_1(\zeta, t) + \rho'\zeta.$$

Thus, if  $A_1, \dots, D_1$ , are the coefficients of the linear part of the new equation,

$$\alpha A_1 + \beta B_1 + \gamma C_1 + \delta D_1 = (\alpha A + \beta B + \gamma C + \delta D)\rho + (\alpha + \delta)\rho',$$

and, if  $\alpha + \delta \neq 0$ , the function  $\rho$  can be chosen as to make this expression vanish. The changes of coordinates with a constant  $\rho$  multiply  $X_1$  by a constant, preserving the previous normalizations.  $\square$

## 3. THE EQUATIONS

For each quadratic homogeneous system appearing in Table 1, we consider the polynomial systems having them as homogeneous component of top degree, with the simplifications of Section 2.4. We then follow the strategy outlined at the beginning of Section 2.

**3.1. Type  $I_{3,3,3}$ .** Following Proposition 2, it is sufficient to consider systems of equations of the form

$$\begin{aligned}x' &= x(x - 2y) + Ax + By, \\y' &= y(y - 2x) + Cx - Ay.\end{aligned}$$

Let us indicate how to calculate the Malmquist condition for the radial orbit  $[0 : 1]$  of the quadratic part. Let  $(X, Y) = (x/y, 1/y)$ , so that  $[x : y : 1] = [X : 1 : Y]$ . These satisfy the system

$$\begin{aligned}YX' &= -3X + BY + 3X^2 + 2AXY - 2CX^2Y, \\Y' &= -1 + 2X + AY - CXY,\end{aligned}$$

which has the form (6) for  $n = 3$ . One can then follow the procedure in Section 2.2 to find that the Malmquist condition for the radial orbit  $[0 : 1]$  of the quadratic part is  $B'' = (\frac{1}{2}B^2 + AB)'$ . By symmetry, the condition for the radial orbit  $[1 : 0]$  is  $C'' = (\frac{1}{2}C^2 - AC)'$ .

The linear involution  $(x, y) \mapsto (-x, -x + y)$  preserves the quadratic part of the system and the particular form of its linear part, and acts upon the coefficients of the latter by

$$(A, B, C) \mapsto (A + B, -B, 2A + B - C).$$

Through this symmetry, and with the simplifications brought by the first two Malmquist conditions, the Malmquist condition for the radial orbit  $[1 : 1]$  is found to be

$$2A'' + B'' - C'' = (\frac{1}{2}C^2 - \frac{1}{2}B^2 - AC - AB)',$$

equivalent to  $(A + B - C)'' = 0$ . We will suppose that  $B' = \frac{1}{2}B^2 + AB + 2\beta$ , and that  $C' = \frac{1}{2}C^2 - AC + 2\gamma$ , for some constants  $\beta$  and  $\gamma$ ; we will also suppose that  $A = 2L - B + C$ , for  $L$  a polynomial of degree at most one.

The function  $u = x - \frac{1}{2}B$  solves the second-order equation

$$(10) \quad u'' = \frac{(u')^2}{2u} + \frac{3}{2}u^3 + eu^2 + fu + \frac{h}{u},$$

for

$$e = 4L, \quad f = 2L^2 + 2L' - \beta + 2\gamma, \quad h = -\frac{1}{2}\beta^2.$$

Through the expression of  $x'$ ,  $y$  is a rational function of  $u$  and  $u'$  whose coefficients are analytic in  $t$ , and the Painlevé property will be present in Eq. (10) if and only if it is present in the original system.

Equation (10) is free of movable critical points. Let us integrate it, following [Inc44, §14.331, case 6°, p. 339]. If  $L$  is constant, then so are  $e$  and  $f$ ; Eq. (10) has the first integral  $K$  given by

$$(u')^2 = u^4 + eu^3 + 2fu^2 + 4Ku + \beta^2,$$

and  $u$  is thus an elliptic function. If  $L$  is not constant then, according to Proposition 2, we may suppose that  $L' = 1$ , and, up to a translation in  $t$ , that  $L = t$ . Eq. (10) reads

$$u'' = \frac{(u')^2}{2u} + \frac{3}{2}u^3 + 4tu^2 + 2(t^2 - \delta)u - \frac{1}{2}\beta^2,$$

for  $\delta = \frac{1}{2}\beta - \gamma - 1$ . This is the general form of the Painlevé IV equation. For each one of these equations, there is a two-dimensional family of inequivalent systems of Type  $I_{3,3,3}$  transformed into it, parametrized by the values of  $B$  and  $C$  at a given point.

For the equations this type, satisfying the Malmquist conditions at the three radial orbits of its quadratic part is sufficient for the system to be free of movable critical points.

**3.2. Type  $I_{2,3,6}$ .** It is sufficient to consider equations of the form

$$(11) \quad \begin{aligned} x' &= x(x - y) + Ax + By, \\ y' &= y(y - 2x) + 2Cx - \left(\frac{3}{2}A + B + \frac{1}{2}C\right)y. \end{aligned}$$

The Malmquist condition for the radial orbit  $[0 : 1]$  is  $B' = B^2 + AB$ . Assuming it, the one for the radial orbit  $[1 : 0]$  is found to be

$$C'' = AC^2 - \frac{1}{4}C^3 + BC^2 - \frac{3}{4}A^2C - 3ABC - 2B^2C - 2AC' - 2BC' + \frac{3}{2}CC' - \frac{3}{2}CA'.$$

Assuming these two, we find that, for  $u = -x - \frac{1}{2}A + \frac{1}{2}C$ ,

$$(12) \quad u'' = -uu' + u^3 - 12qu + 12q',$$

for

$$q = \frac{1}{48}(A^2 - 8AC - 8BC + 3C^2 - 2A' - 2C').$$

The Malmquist condition for the radial orbit  $[2 : 3]$  is  $(q'' - 6q^2)'' = 0$ .

Under this last condition, Eq. (12) is free of movable critical points; its integration is carried out in [Pai02, p. 23, eq. 6]. If  $z$  is a solution to  $z'' - 6z^2 = q'' - 6q^2$  different from  $q$ ,

$$u = \frac{z' - q'}{z - q}$$

is a solution to (12). If  $q'' - 6q^2$  is constant, so is  $z'' - 6z^2$ , and both  $q$  and  $z$  are elliptic functions. If  $q'' - 6q^2$  is not constant, it is a polynomial of degree one, and both  $q$  and  $z$  satisfy the Painlevé I equation (up to an inessential scaling of the independent variable).

We have that  $(x, y)$  is birationally equivalent to  $(u, u')$ , and the system (11) is free of movable critical points as soon as the three Malmquist conditions are satisfied.

**3.3. Type  $I_{2,4,4}$ .** We will consider the equations in the form

$$\begin{aligned} x' &= x(x - y) + Ax + By, \\ y' &= y(y - 3x) + 3Cx - (2A + B)y. \end{aligned}$$

The Malmquist condition for the radial orbit  $[0 : 1]$  reads  $B' = B^2 + AB$ .

The function  $u = x + \frac{1}{2}A - \frac{1}{2}C$  solves the equation

$$(13) \quad u'' = 2u^3 + fu + g,$$



for

$$\begin{aligned} f &= \frac{1}{2}(6AC - A^2 - 3C^2 + 6BC + 2A'), \\ g &= \frac{1}{2}(3AC^2 - 2A^2C - 3ABC + 3BC^2 - C^3 - AA' + CA' + A'' - C''). \end{aligned}$$

The Malmquist condition for the radial orbit  $[1 : 0]$  is  $f'' = 2g'$ .

The linear involution  $(x, y) \mapsto (-x, y - 2x)$  preserves the quadratic part of the system, fixing the radial orbit  $[0 : 1]$  and exchanging the radial orbits  $[1 : 0]$  and  $[1 : 0]$ ; it preserves the chosen form of the linear part, on which it acts by

$$(A, B, C) \mapsto (A + 2B, -B, 2A + 2B - C).$$

Under this involution,  $f \mapsto f$  and  $g \mapsto -g$ ; the Malmquist condition for the radial orbit  $[1 : 1]$  is thus  $f'' = -2g'$ . Together, these conditions are equivalent to  $f'' = 0$  and  $g' = 0$ . Under them, equation (13) is free of movable branch points. It is integrated in [Inc44, §14.315, p. 333]. If  $f$  is constant, (13) has the first integral

$$u^4 + fu^2 + 2gu - (u')^2,$$

and  $u$  is an elliptic function; if  $f$  is linear, the equation is the most general Painlevé II equation (up to scaling  $f$  and translating the independent variable).

Again, since  $(x, y)$  is birationally equivalent to  $(u, u')$ , the original system is free of movable branch points if the Malmquist conditions at the three radial orbits of its quadratic part are satisfied.

**3.4. Types  $I_{n+1}$  ( $n \geq 0$ ).** Consider, in a first time, the equations in the form

$$(14) \quad \begin{aligned} x' &= x^2 + Ax + By, \\ y' &= y(y - nx) + Cx + Dy. \end{aligned}$$

The Malmquist condition for the radial orbit  $[1 : 0]$  is  $B \equiv 0$ , which will be assumed from now on. Thus,  $x$  satisfies a Riccati equation with holomorphic coefficients, and  $y$  solves a Riccati equation, having, however, meromorphic ones. The poles of  $x$  might produce movable critical points for  $y$ .

For  $n = 0$ , the Malmquist condition for the radial orbit  $[0 : 1]$  reads  $C \equiv 0$ , and the equation separates into two independent Riccati equations with holomorphic coefficients.

We will henceforth suppose that  $n \geq 1$ , and consider the equations in the form

$$\begin{aligned} x' &= x \left( x + \frac{2}{n}A \right), \\ y' &= y(y - nx) + nCx - (A + 2C)y. \end{aligned}$$

The function  $u = -y + C$  solves the equation

$$(15) \quad u'' = \left( 1 - \frac{1}{n} \right) \frac{(u')^2}{u} + \left( au + \frac{c}{u} \right) u' + du^3 + fu + g + \frac{h}{u},$$

for

$$\begin{aligned} a &= -\frac{n+2}{n}, \quad c = -\left( \frac{n-2}{n} \right) E, \quad d = -\frac{1}{n}, \\ f &= \frac{2}{n}E + \frac{1}{n}A^2 - A', \quad g = E', \quad h = -\frac{1}{n}E^2, \end{aligned}$$

with  $E = AC + C^2 + C'$ .

3.4.1. *When  $E \equiv 0$ .* We will see that, under this condition, the systems are free of movable critical points.

For  $n = 1$ , this condition is exactly Malmquist's one for the radial orbit  $[1 : 0]$ , and the equation reduces to

$$u'' = -3uu' - u^3 + fu,$$

for  $f = A^2 - A'$ . The solutions are of the form  $u = v'/v$  for  $v''' = fv'$  (see [Inc44, §14.313, p. 331]). These equations are thus free from movable critical points. For greater values of  $n$ , the Eqs. (15) are particular cases of an equation studied by Gambier in [Gam10, p. 28, §6], where explicit formulas giving their integration, exhibiting the absence of moving critical points, may be found.

The sufficiency of the condition  $E \equiv 0$  for the absence of movable critical points can be interpreted geometrically, in close connection to the ideas in [GR97, Section 3.4], which are somewhat hinted at by Gambier's formulas. Let us, for a moment, not assume that  $E \equiv 0$ . In the coordinates  $(z, u) = (x, y - c)$ , the time-dependent vector field in  $\mathbf{C}^2 \times U$  associated to the system (14) with  $B \equiv 0$  is

$$z \left( z + \frac{2}{n}A \right) \partial_z + (u^2 - nzu - Au - E) \partial_u.$$

Gluing another copy of  $\mathbf{C}^2 \times U$ , with coordinates  $(w, v)$  of  $\mathbf{C}^2$ , through  $(w, v) = (1/z, z^n u)$ , we find the vector field

$$-(1 + \frac{2}{n}Aw) \partial_w + (v^2 w^n + Av - v^{-n}E) \partial_v.$$

This gluing produces, at each  $t$ , the total space of the line bundle  $O(-n)$  over  $\mathbf{P}^1$ . The original vector field gives a meromorphic vector field on the second chart, which is holomorphic whenever  $E \equiv 0$ . When compactifying  $O(-n)$  into the  $n$ -th Hirzebruch surface  $F_n$  (by  $u \mapsto 1/u$  in the first chart; by  $v \mapsto 1/v$  in the second), the vector field extends, still under the condition  $E \equiv 0$ , holomorphically. Hence, if  $E \equiv 0$ , the original extended phase space  $\mathbf{C}^2 \times U \rightarrow U$  can be extended to a space with compact fibers  $F_n \times U \rightarrow U$ , in which the vector is holomorphic, and hence transverse to the fibration. This guarantees the absence of movable critical points.

3.4.2. *When  $E \not\equiv 0$ .* For  $n = 2$ , the Malmquist condition at  $[1 : 0]$  is  $E' \equiv 0$ . Let us show that this condition is sufficient for the system to be free of movable critical points. Assuming, with Proposition 2, that  $E \equiv 1$ , reduces the equation to

$$(16) \quad u'' = \frac{(u')^2}{2u} - 2uu' - \frac{u^3}{2} + fu - \frac{1}{2u},$$

for  $f = 1 + \frac{1}{2}A^2 - A'$ . For its integration, we follow [Inc44, §14.331, case 4°, p. 338]. For  $u = v'/v$ ,

$$2v'''v' = (v'')^2 + 2f(v')^2 - v^2.$$

which, by differentiation, yields the fourth-order linear equation

$$v^{IV} = 2fv'' + f'v' - v.$$

(This seems to challenge Kimura and Matuda's claim in [KM80], that no linear equations of order greater than three are needed to integrate the equations under consideration. Is there a way to integrate (16) without resorting to fourth-order equations?)

For greater values of  $n$ , a change of coordinates of the form (7), with  $\lambda \equiv 1$  and  $\rho = E$ , transforms the equation into one whose coefficients are

$$a_1 = -\frac{n+2}{nE}, \quad b_1 = -\frac{E'}{E^2}, \quad c_1 = -\frac{n-2}{n},$$

$$d_1 = -\frac{1}{nE^2}, \quad f_1 = \frac{f}{E^2}, \quad g_1 = \frac{E'}{E^2}, \quad h_1 = -\frac{1}{n}.$$

(These are functions of  $t$ , which is, on its turn, a function of the new independent variable  $\tau$ .) Since  $dE/d\tau = E'/E$ , we have that  $da_1/d\tau = a_1b_1$ . These equations are particular cases of the family studied by Gambier in [Gam10, Ch. VII, Eq. 5, p. 53], and have a rather intricate scheme of integration. For them, either the absence of movable critical points can be established directly through the vanishing of a particular explicit expression, or the equation may be transformed into one of the same family, in which  $n$  is replaced by  $n-2$ . We refer to Gambier's work for details.

We ignore the extent to which the Malmquist conditions ensure the absence of movable critical points, beyond what has already been said.

**3.5. Type  $I_{2,2,\infty}$ .** Consider the equations

$$x' = x(x-y) + Ax + By,$$

$$y' = y(y-x) + Cx - (A+B+C)y.$$

Malmquist's conditions at the radial orbits  $[0 : 1]$  and  $[1 : 0]$  are, respectively,  $B' = (A+B)B$ , and  $C' = -(A+B)C$ . With these,  $u = -x + B$  satisfies the equation

$$u'' = -2uu' + qu' + q'u,$$

for  $q = A + 2B - C$ . The equation is integrated in [Inc44, §14.312, p. 330]: it has the first integral  $u' + u^2 - qu$ , and  $u$  is the solution of a Riccati equation.

**3.6. Type  $I_\infty$ .** These are the equations of the form

$$x' = x(x-y) + Ax + By,$$

$$y' = Cx + Dy.$$

Malmquist's condition for the radial orbit  $[1 : 0]$  reads  $C \equiv 0$ :  $y$  can be integrated independently, as a holomorphic function, after which,  $x$  satisfies a Riccati equation with holomorphic coefficients.

**3.7. Types  $II_{n+1}$ ,  $n \neq -1$ .** We will consider the equations in the form

$$x' = x^2 + By,$$

$$y' = -nxy + Cx + Dy.$$

After a rescaling of the form  $y \mapsto \rho(t)y$ , we will suppose that  $D \equiv 0$ . The direction  $[1 : 0]$  gives a radial orbit for the quadratic part, with Kowalevski exponent  $n+1$  (positive, and hence subject to Malmquist's test, if  $n \geq 0$ ).

For  $n = 0$ , Malmquist's condition for the radial orbit  $[1 : 0]$  reads  $C \equiv 0$ :  $y$  is a first integral, and  $x$  solves a Riccati equation with holomorphic coefficients.

We will henceforth suppose that  $n \neq 0$ . For  $u = -ny + C$ ,

$$(17) \quad u'' = \left(1 - \frac{1}{n}\right) \frac{(u')^2}{u} + \frac{cu'}{u} + eu^2 + fu + g + \frac{h}{u},$$

for

$$c = -\frac{n-2}{n}C', \quad e = B, \quad f = -BC, \quad g = C'', \quad h = -\frac{1}{n}(C')^2.$$

If  $B \equiv 0$ , following [Gam10, p. 28, Eq. 1] (where the equation solved by  $1/u$  is integrated), the solutions are of the form

$$u(t) = L^n(t) \int^t \frac{C'(s)}{L^n(s)} ds,$$

with  $L$  a polynomial of degree at most one. For the solutions to be free of movable critical points, these integrals must be free of logarithmic terms for all choices of  $L$ , this is,  $d^n C/dt^n \equiv 0$  (this is the Malmquist condition at  $[1 : 0]$  when  $B \equiv 0$ ).

We will henceforth assume that  $B \neq 0$ . We begin by showing that, for equation (17) to be free of movable critical points, it is necessary that  $n$  divides 4 and that  $n \neq -2$ . Let  $t_0$  be such that  $e(t_0) \neq 0$ . By considering

$$w(t) = \alpha^2 u(t_0 + \alpha t),$$

and taking the limit of the equation it satisfies as  $\alpha \rightarrow 0$ , we obtain the autonomous equation

$$w'' = \left(1 - \frac{1}{n}\right) \frac{(w')^2}{w} + \epsilon w^2,$$

with  $\epsilon = e(t_0)$ . If (17) is free from movable branch points, the solutions to this equation are univalent. For  $s = w^3/(w')^2$ ,

$$(18) \quad -\frac{s''}{(s')^2} = -\frac{3n\epsilon s - n - 1}{s(2n\epsilon s - n - 2)}.$$

Briot and Bouquet have shown that, for  $s$  to be univalent, this expression must have only simple poles, with residues equal either to  $-1$  or to  $1/m - 1$  for some  $m \in \mathbf{Z}^*$  [BB55]. The expression (18) has a double pole at  $s = 0$  when  $n = -2$ , and this case may be discarded. The residue of (18) at the pole  $s = (n+2)/(2n\epsilon)$  is

$$\frac{n}{2(n+2)} - 1.$$

If  $s$  is univalent,  $2(n+2)/n = 2 + 4/n$  is an integer, and  $n$  divides 4.

Let us now study the special cases where  $B \neq 0$  and  $n \in \{-4, 1, 2, 4\}$ .

3.7.1. *Case  $n = -4$ .* For  $v = 1/u$ , we have

$$(19) \quad v'' = \frac{3(v')^2}{4v} + avv' + dv^3 + ev^2 + fv + g,$$

for

$$a = -\frac{3}{2}C', \quad d = -\frac{1}{4}(C')^2, \quad e = -C'', \quad f = BC, \quad g = -B,$$

which has the form (7) for  $m = 4$ . We will see that the condition for the absence of movable critical points is that  $B$  is constant.

If  $C' \equiv 0$ , the coefficients  $a$ ,  $d$  and  $e$  vanish identically. A suitable change of coordinates in equation (19) of the form (8), with  $\lambda\rho^2 \equiv 1$ , preserves the form of the equation, makes  $f$  vanish, and does not modify  $g$  (but for the change of independent variable), reducing the equation to

$$v'' = \frac{3(v')^2}{4v} + g.$$

Following [Inc44, §14.331, case 1°, Eq. XXII, p. 337], for this equation to be free of movable critical points, it is necessary that  $g$  is constant, in which case the equation is solved by  $v = (\alpha t^2 + \beta t + \gamma)^2$ , with  $\beta^2 = 4\alpha\gamma - g$ .

If  $C' \neq 0$ , by a change of coordinates of the form (8), with  $\lambda \equiv 1$  and  $\rho = C'$ , we obtain a new equation of the form (7), with  $m = 4$ , whose nonzero coefficients are

$$a_1 = -\frac{3}{2}, \quad b_1 = -\frac{C''}{(C')^2}, \quad d_1 = -\frac{1}{4}, \quad e_1 = b_1, \quad f_1 = \frac{BC}{(C')^2}, \quad g_1 = -\frac{B}{(C')^2}.$$

(Again, these are functions of  $t$ , which is, on its turn, function of the new independent variable  $\tau$ .) Following [Inc44, §14.331, case 2°, p. 338], a necessary and sufficient condition for the equation to be free of movable critical points is that

$$b_1 = \frac{1}{2g_1} \frac{dg_1}{d\tau},$$

this is, that  $B$  is constant (this is an intrinsic condition). The resulting equation appears as Eq. XXV in [Inc44, §14.331, p. 338], where the reader may find the explicit form of its integration ( $u$  is expressed in terms of the logarithmic derivative of the solution of a linear equation of the third order).

3.7.2. *Case  $n = 1$ .* The Malmquist condition at  $[1 : 0]$  is  $C' \equiv 0$ . Eq. (17) reduces to  $u'' = eu^2 + fu$ , for  $e = B$ ,  $f = -BC$ . A suitable change of coordinates of the form (8) with  $\lambda^2\rho \equiv 1$  transforms this equation into one of the form

$$\frac{d^2w}{d\tau^2} = 6w^2 + f_1w.$$

For  $v = w + \frac{1}{12}f_1$ ,

$$\frac{d^2v}{d\tau^2} = 6v^2 + \frac{1}{24} \left( 2\frac{d^2f_1}{d\tau^2} - f_1^2 \right).$$

The condition for the absence of movable critical points is that the expression in parenthesis is a polynomial in  $\tau$  of degree at most one. If the latter is constant, the equation is solved by elliptic functions; if it has degree one, we have the first Painlevé equation (see [Inc44, §14.311, p. 328–329]).

3.7.3. *Case  $n = 2$ .* The Malmquist condition at  $[1 : 0]$  is  $C'' \equiv 0$ . Eq. (17) reduces to

$$u'' = \frac{(u')^2}{2u} + eu^2 + fu + \frac{h}{u},$$

for

$$e = B, \quad f = -BC, \quad h = -\frac{1}{2}(C')^2.$$

A suitable change of coordinates of the form (8) with  $\lambda\rho \equiv 1$  preserves the form of the equation, makes  $e_1$  an arbitrary (nonzero) constant, and preserves  $h$ , which remains constant. The conditions for the resulting equations to be free of movable branch points are as follows.

Suppose, in a first time, that  $C$  is constant. According to [Inc44, §14.331, case 1°, p. 337], for the equation

$$u'' = \frac{(u')^2}{2u} + 4u^2 + fu$$

to be free of movable critical points, it is necessary that  $f'' \equiv 0$ . If  $f$  is constant, the equation has the first integral  $K$  defined by

$$(u')^2 = 4u(u^2 + \frac{1}{2}fu + K),$$

and  $u$  is an elliptic function (this corresponds to equations XVIII and XI in [Inc44, p. 337]). If  $f$  is not constant then, up to a scaling and a translation in the independent variable, we may suppose that  $f = 2t$ . In this case, for  $w^2 = u$ , we have that  $w'' = 2w^3 + tw$ , a particular case of the Painlevé II equation (this is Eq. XX in [Inc44, p. 337]).

If  $C$  is not constant, it is sufficient, according to Proposition 2, to consider the case where  $C' \equiv 1$ . According to [Inc44, §14.331, case 3°, p. 338], for the equation

$$u'' = \frac{(u')^2}{2u} + 4\alpha u^2 + fu - \frac{1}{2u},$$

with  $\alpha \in \mathbf{C}^*$ , to be free of movable critical points, it is necessary and sufficient that  $f'' \equiv 0$ . For  $\beta \in \mathbf{C}$ ,

$$u'' = \frac{(u')^2}{2u} + 4\alpha u^2 + \beta u - \frac{1}{2u}$$

has the first integral  $(u')^2 = 4\alpha u^3 + 2\beta u^2 + 4Ku + 1$ . If

$$u'' = \frac{(u')^2}{2u} + 4\alpha u^2 - tu - \frac{1}{2u},$$

with  $\alpha \in \mathbf{C}^*$ ,  $v = \frac{1}{2}(u' + 1)/u$  is a solution to the Painlevé II equation

$$v'' = 2v^3 + tv - 2\alpha - \frac{1}{2}.$$

(These are equations XXXIII and XXXIV in [Inc44, p. 339].)

3.7.4. *Case  $n = 4$ .* The Malmquist condition at the radial orbit  $[1 : 0]$  reads

$$(20) \quad 6BCC'' + 4B(C')^2 + 3B'CC' + 6C^{IV} = 0,$$

but we will not assume it for the time being.

We are dealing with the equation

$$(21) \quad u'' = \frac{3(u')^2}{4u} + c\frac{u'}{u} + eu^2 + fu + g + \frac{h}{u},$$

for

$$c = -\frac{1}{2}C', \quad e = B, \quad f = -BC, \quad g = C'', \quad h = -\frac{1}{4}(C')^2,$$

with  $B \neq 0$ . When such an equation is free of movable critical points, it can be transformed, through a change of coordinates of the form (8), into one of the equations in cases 1° and 3° in §14.331 in Ince's treatise [Inc44, p. 337–339] (cases 2° and 4° cannot arrive when the coefficient of  $uu'$  in the right-hand side of (21) is zero while the coefficient of  $u^2$  is not).

Case 1° can only arrive if  $C' \equiv 0$  (in which Malmquist's condition (20) is satisfied), giving an equation of the form (21) in which both  $c$ ,  $g$  and  $h$  vanish identically. The absence of movable critical point requires that, when the equation is transformed as to make  $e \equiv 3$ , that  $f$  is constant, leading to Ince's Eq. XXI when  $f = 0$ , and to a particular case of Ince's Eq. XXIII when  $f \neq 0$ . In both cases,  $u^2$  satisfies an equation with an elliptic first integral. These cases arrive, for instance, when  $B$  is a nonzero constant.

Case 3° corresponds to the canonical form

$$(22) \quad u'' = \frac{3(u')^2}{4u} + 6q'\frac{u'}{u} + 3u^2 + 12qu - 12q'' - 36\frac{(q')^2}{u},$$

with  $(q'' - 6q^2)'' = 0$ , appearing both in [Gam10, p. 29], and, as Eq. XXVI, in [Inc44, §14.331, p. 338], whose solutions are given as follows. Let  $z$  be a solution to  $z'' + z^2 = q'' + 6q^2$ ,  $z \neq q$ , and let

$$v = \frac{z' - q'}{z - q}.$$

Then,  $u = \frac{1}{3}(-2v' + v^2 - 12q)$  is a solution to (22) (this corrects a misprint in Gambier's article, repeated later in Ince's treatise).

In equation (21), after a suitable change of coordinates of the form (8) in which  $\lambda\rho^2$  is constant, we can make the coefficient of  $u^2$  in the right-hand side of (21) equal to 3 while keeping the coefficient of  $u'$  equal to 0, obtaining an equation of the form

$$u'' = \frac{3(u')^2}{4u} + 6p\frac{u'}{u} + 3u^2 + 12qu - 12p' - 36\frac{p^2}{u},$$

for  $p = \frac{1}{6}c_1$ ,  $q = \frac{1}{12}f_1$ , with  $c_1$  and  $f_1$  given by (9). The first condition for the latter to fit within the canonical form (22) is that  $p = dq/d\tau$ , which reads

$$64B^4C' + 48CB^3B' - 48B^2B''' + 180BB'B'' - 135(B')^3 = 0.$$

Given this, Malmquist's condition (20) is equivalent to

$$\frac{d^2}{d\tau^2} \left( \frac{d^2q}{d\tau^2} - 6q^2 \right) = 0,$$

the second necessary condition for redressing the equation into its canonical form.

**3.8. Type  $II_\infty$ .** Consider the equations

$$\begin{aligned} x' &= xy + Ax + By, \\ y' &= Cx. \end{aligned}$$

Up to a suitable rescaling for the form  $x \mapsto \rho(t)x$ , we may suppose that  $A \equiv 0$ , while preserving the above form. If  $C \equiv 0$ ,  $y$  is constant, and  $x$  satisfies a linear equation. We will henceforth suppose that this is not the case.

The function  $u = -\frac{1}{2}y$  solves the equation

$$u'' = -2uu' + bu' + fu,$$

for  $b = C'/C$  and  $f = BC$ . This equation is studied in [Inc44, §14.312, p. 331]. For it to be free of movable branch points, it is necessary that the intrinsic condition  $f' = b$  is satisfied, in which case we have the first integral  $u' + u^2 - bu$ .

**3.9. Type III.** Consider the equations

$$(23) \quad \begin{aligned} x' &= 6y^2 + Ax + By, \\ y' &= Cx + Dy. \end{aligned}$$

If  $C \equiv 0$ , the system separates into a linear equation and a Riccati equation with holomorphic coefficients, and we will henceforth suppose that this is not the case. For the new variables  $(X, Y)$  defined by

$$\begin{aligned} x(t) &= \alpha(t)X(\tau) + \beta(t)Y(\tau), \\ y(t) &= \gamma(t)Y(\tau), \end{aligned}$$

with  $d\tau/dt = \rho(t)$ ,

$$\begin{aligned}\frac{dX}{d\tau} &= \frac{6\gamma^2}{\alpha\rho}Y^2 + \frac{1}{\rho}\left(A - \frac{\beta}{\gamma}C - \frac{\alpha'}{\alpha}\right)X + (\dots)Y, \\ \frac{dY}{d\tau} &= \frac{\alpha C}{\gamma\rho}X + \frac{1}{\rho}\left(\frac{\beta}{\gamma}C + D - \frac{\gamma'}{\gamma}\right)Y.\end{aligned}$$

By conveniently choosing  $\alpha, \beta, \gamma$  and  $\rho$ , we may suppose (after relabeling the variables), that, in the system (23),  $A \equiv 0$ ,  $D \equiv 0$ , and  $C \equiv 1$ , reducing it to  $y'' = 6y^2 + By$  (with  $x = y'$ ). This equation was discussed in Section 3.7.2; it either reduces to the Painlevé I equation or is solved by elliptic functions.

**3.10. Type IV.** These are the equations

$$\begin{aligned}x' &= x^2 + Ax + By, \\ y' &= xy + Cx + Dy.\end{aligned}$$

After a suitable change of coordinates of the form  $(x, y) \mapsto (x, \rho(t)y + \mu(t)x)$ , we may suppose that both  $C$  and  $D$  vanish identically. Then,  $u = 1/y$  satisfies the linear equation  $u'' = Au' - B$ .

Goursat observed that these are legitimate generalizations of the Riccati equation, and that they are free of movable critical points [Gou29, Ch. XXI]: when  $\mathbf{C}^2 \times U$  is compactified in the direction of the dependent variables into  $\mathbf{P}^2 \times U$  by  $((x, y), t) \mapsto ([x : y : 1], t)$ , the system extends holomorphically.

#### REFERENCES

- [BB55] C. BRIOT and J.-C. BOUQUET, Recherches sur les fonctions doublement périodiques, *C. R. Acad. Sci. Paris* **40** no. 7 (1855), 342–344.
- [Gam10] B. GAMBIER, Sur les équations différentielles du second ordre et du premier degré dont l'intégrale générale est à points critiques fixes, *Acta Math.* **33** no. 1 (1910), 1–55. <https://doi.org/10.1007/BF02393211>.
- [Gar60] R. GARNIER, Sur des systèmes différentiels du second ordre dont l'intégrale générale est uniforme, *Ann. Sci. École Norm. Sup. (3)* **77** (1960), 123–144. <https://doi.org/10.24033/asens.1088>.
- [GR97] É. GHYS and J.-C. REBELO, Singularités des flots holomorphes. II, *Ann. Inst. Fourier (Grenoble)* **47** no. 4 (1997), 1117–1174. <https://doi.org/10.5802/aif.1594>.
- [GR00] É. GHYS and J.-C. REBELO, Erratum: “Singularités des flots holomorphes. II”, *Ann. Inst. Fourier (Grenoble)* **50** no. 3 (2000), 1019–1020. <https://doi.org/10.5802/aif.1783>.
- [GL74] J. GOFFAR-LOMBET, Sur des systèmes polynomiaux d'équations différentielles dont l'intégrale générale est à points critiques fixes, *Acad. Roy. Belg. Cl. Sci. Mém. (2)* **XLI** no. 3 (1974). Available at [https://academieroyale.be/Academie/documents/XLI3\\_GoffarLombet\\_Surdessystemespolynomiaux\\_197413525.pdf](https://academieroyale.be/Academie/documents/XLI3_GoffarLombet_Surdessystemespolynomiaux_197413525.pdf).
- [Gou29] E. GOURSAT, *Cours d'analyse mathématique. Tome II*, cinquième ed., Gauthier-Villars et Cie., éditeurs, 1929.
- [Gor00] A. GORIELY, A brief history of Kovalevskaya exponents and modern developments, *Regul. Chaotic Dyn.* **5** no. 1 (2000), 3–15. <https://doi.org/10.1070/rd2000v005n01ABEH000120>.
- [Gui06] A. GUILLOT, Semicompleteness of homogeneous quadratic vector fields, *Ann. Inst. Fourier (Grenoble)* **56** no. 5 (2006), 1583–1615. <https://doi.org/10.5802/aif.2221>.
- [Inc44] E. L. INCE, *Ordinary Differential Equations*, Dover Publications, New York, 1944.
- [KM80] T. KIMURA and T. MATUDA, On systems of differential equations of order two with fixed branch points, *Proc. Japan Acad. Ser. A Math. Sci.* **56** no. 10 (1980), 445–449. <https://doi.org/10.3792/pjaa.56.445>.



- [Mal21] J. MALMQUIST, Sur les points singuliers des équations différentielles, 2ème mémoire, *Ark. Mat. Astr. Fys.* **15** no. 27 (1921), 1–78.
- [Oka79] K. OKAMOTO, Sur les feuilletages associés aux équations du second ordre à points critiques fixes de P. Painlevé, *Japan. J. Math. (N.S.)* **5** no. 1 (1979), 1–79. <https://doi.org/10.4099/math1924.5.1>.
- [Pai00] P. PAINLEVÉ, Mémoire sur les équations différentielles dont l'intégrale générale est uniforme, *Bull. Soc. Math. France* **28** (1900), 201–261. <https://doi.org/10.24033/bsmf.633>.
- [Pai02] P. PAINLEVÉ, Sur les équations différentielles du second ordre et d'ordre supérieur dont l'intégrale générale est uniforme, *Acta Math.* **25** no. 1 (1902), 1–85. <https://doi.org/10.1007/BF02419020>.
- [Pai73] P. PAINLEVÉ, *Oeuvres de Paul Painlevé. Tome I*, Éditions du Centre National de la Recherche Scientifique, Paris, 1973, compiled by R. Gérard, G. Reeb, and A. Sec.

INSTITUTO DE MATEMÁTICAS, UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO, CIUDAD UNIVERSITARIA 04510, CIUDAD DE MÉXICO, MEXICO

*Email address:* `adolfo.guillot@im.unam.mx`