INTEGRABILITY OF THE INOZEMTSEV SPIN CHAIN

OLEG CHALYKH

ABSTRACT. We show that the Inozemtsev spin chain is integrable. The conserved quantities (commuting Hamiltonians) are constructed using elliptic Dunkl operators. We also suggest a generalisation.

1. INTRODUCTION

The Inozemtsev quantum spin chain [I1] is described by the Hamiltonian

$$\mathscr{H} = \sum_{i < j}^{n} \wp\left(\frac{i - j}{n}\right) P_{ij} \,. \tag{1.1}$$

Here $\wp(z) = \wp(z|1,\tau)$ is the Weierstrass \wp -function with periods $1,\tau$ and P_{ij} acts on the tensor power $U^{\otimes n}$ of $U = \mathbb{C}^m$ by permuting the factors. This is viewed as a spin chain on n sites, with each copy of U representing local spin states, and with $\wp\left(\frac{i-j}{n}\right)$ expressing the strength of interaction between the *i*th and *j*th sites. To show that it is integrable, one would like to find elements of $\operatorname{End}_{\mathbb{C}}(U^{\otimes n})$ commuting with \mathscr{H} . This is an old problem, see [I2, DI1, DI2, SS, FGL] for some of its history and related context. The trigonometric case – with $\sin^{-2}(\pi z)$ in place of $\wp(z)$ – describes the celebrated Haldane–Shastry chain [H, S] which is integrable and has the Yangian symmetry [BGHP]. Nothing of that sort in known in the elliptic case so, as it stands, (1.1) falls outside the class of problems solved by Quantum Inverse Scattering or similar methods.

Nevertheless, as will be demonstrated below, one can construct commuting Hamiltonians from those of the quantum elliptic spin Calogero–Moser system, by "freezing". This builds on (and elucidates) the original idea by Polychronakos [P2], see also related works [FM, MP, TH]. Our main ingredients are the elliptic Dunkl operators and Cherednik algebra. Along the way, we find a generalisation of the Inozemtsev spin chain.

One would also like to calculate the eigenvalues and eigenvectors of \mathscr{H} , see [KL] for the state of the art and references. Hopefully, the link to the elliptic Dunkl operators can shed some light onto this. Our methods apply, almost verbatim, to elliptic Calogero–Moser systems (and spin chains) for arbitrary root systems. They can also be extended to Calogero–Moser systems of *R*-matrix type and the corresponding spin chains from [LOZ, SZ], which will be done separately [ChL].

Acknowledgement. I would like to thank Andrei Zotov for useful conversations. I am also grateful to Jules Lamers and Rob Klabbers for stimulating discussions and for waiting patiently for this paper to be written up.

2. RATIONAL CALOGERO-MOSER SYSTEM AND DUNKL OPERATORS

Here we recall the well-known relationship between the rational Calogero–Moser system, Dunkl operators and Cherednik algebra, see [E] and references therein.

2.1. Let $V = \mathbb{C}^n$ be the *n*-dimensional complex Euclidean vector space with the standard basis (e_i) and coordinates (x_i) . The symmetric group S_n acts on V by permuting the basis vectors. This induces an S_n -action on the algebra $\mathbb{C}(V)$ of meromorphic functions in *n* variables by $(w.f)(x) = f(w^{-1}.x)$ for $x \in V$, $w \in S_n$. The crossed product $\mathbb{C}(V) * S_n$ is formed by taking the vector space $\mathbb{C}(V) \otimes \mathbb{C}S_n$ with the multiplication $(f \otimes w)(f' \otimes w') = f(w.f') \otimes ww'$. Let $\mathscr{D}(V)$ denote the ring of differential operators on V with meromorphic coefficients. It is generated by the partial derivatives $\partial_i = \frac{\partial}{\partial x_i}$ and operators of multiplication by $g \in \mathbb{C}(V)$. We have a natural action of S_n on $\mathscr{D}(V)$, hence the crossed product $\mathscr{D}(V) * S_n$. As an algebra, $\mathscr{D}(V) * S_n$ is generated by its two subalgebras, $1 \otimes \mathbb{C}S_n$ and $\mathscr{D}(V) \otimes 1$ which can be identified with $\mathbb{C}W$ and $\mathscr{D}(V)$, respectively. Using these identifications, we replace $a \otimes w$ by aw, so each element of $\mathscr{D}(V) * S_n$ is written uniquely as $a = \sum_{w \in S_n} a_w w$ with $a_w \in \mathscr{D}(V)$.

2.2. The *quantum rational* Calogero–Moser (CM) system is described by the Hamiltonian

$$H^{(r)} = \sum_{i=1}^{n} \hbar^2 \partial_i^2 - \sum_{i< j}^{n} \frac{2\kappa(\kappa - \hbar)}{(x_i - x_j)^2}.$$
 (2.1)

Here $\kappa \in \mathbb{C}$ is the coupling constant and $\hbar \neq 0$ is a quantum parameter ¹. This is a completely integrable system, in the sense that it admits *n* commuting Hamiltonians. These are constructed using the *Dunkl operators*:

$$y_i^{(r)} = \hbar \partial_i - \kappa \sum_{j \neq i}^n \frac{1}{x_i - x_j} s_{ij}, \qquad i = 1, \dots, n.$$
 (2.2)

The key property of the these operators is their commutativity, $[y_i^{(r)}, y_j^{(r)}] = 0$, and equivariance,

$$s_{ij}y_i^{(r)} = y_j^{(r)}s_{ij}, \qquad s_{ij}y_k^{(r)} = y_k^{(r)}s_{ij} \quad (k \neq i, j)$$

As a result, the assignment $e_i \mapsto y_i^{(r)}$ extends to a S_n -equivariant algebra map

$$\mathbb{C}[V^*] \to \mathscr{D}(V) * S_n \,, \quad q \mapsto q(y^{(r)}) \,. \tag{2.3}$$

Introduce the linear map

$$\operatorname{Res}: \mathscr{D}(V) * S_n \to \mathscr{D}(V), \qquad \sum_{w \in S_n} a_w w \mapsto \sum_{w \in S_n} a_w . \tag{2.4}$$

Combining this map with (2.3), define

$$L_q^{(r)} = \operatorname{Res} q(y^{(r)}), \qquad q \in \mathbb{C}[V^*]^{S_n}.$$
 (2.5)

The S_n -invariance of q and the commutativity of the Dunkl operators imply that (1) each $L_q^{(r)}$ is S_n -invariant, (2) $L_q^{(r)} L_{q'}^{(r)} = L_{qq'}^{(r)}$ for any $q, q' \in \mathbb{C}[V^*]^{S_n}$, (3) the family $\{L_q^{(r)}, q \in \mathbb{C}[V^*]^{S_n}\}$ is commutative.

¹To follow the standard conventions of quantum mechanics, \hbar should be replaced $-i\hbar$ and κ should be purely imaginary.

Taking $q = e_1^2 + \cdots + e_n^2 \in S^2 V \subset \mathbb{C}[V^*]$, a direct calculation shows that

$$q(y^{(r)}) = (y_1^{(r)})^2 + \dots + (y_n^{(r)})^2 = \sum_{i=1}^n \hbar^2 \partial_i^2 - \sum_{i(2.6)$$

Hence, $L_q^{(r)} = \operatorname{Res} q(y^{(r)})$ in this case is precisely the CM Hamiltonian (2.1). Other symmetric combinations of $y_1^{(r)}, \ldots, y_n^{(r)}$ produce higher commuting Hamiltonians.

2.3. The classical limit corresponds to taking $\hbar \to 0$. More precisely, we view the Dunkl operators as elements of the algebra

$$A_{\hbar} = \mathbb{C}[[\hbar]] \otimes \mathbb{C}(V)[\hat{p}_1, \dots, \hat{p}_n], \qquad \hat{p}_i = \hbar \partial_i.$$

The quantum momenta \hat{p}_i satisfy the relations $[\hat{p}_i, f] = \hbar \partial_i f$ for $f \in \mathbb{C}(V)$. We have an algebra isomorphism

$$\eta_0: (A_\hbar * S_n) / \hbar (A_\hbar * S_n) \to A_0 * S_n, \qquad f \mapsto f, \ \hat{p}_i \mapsto p_i, \ w \mapsto w, \qquad (2.7)$$

where

$$A_0 = \mathbb{C}(V) \otimes \mathbb{C}[V^*] = \mathbb{C}(V)[p_1, \dots, p_n]$$

is the classical version of A_{\hbar} . Therefore, A_{\hbar} and $A_{\hbar} * S_n$ are formal deformations of A_0 and $A_0 * S_n$, respectively. Note that A_0 is commutative, with Poisson bracket defined by $\{\eta_0(a), \eta_0(b)\} = \eta_0(\hbar^{-1}[a, b])$ for $a, b \in A_{\hbar}$. For any $a \in A_{\hbar} * S_n$, we call $\eta_0(a)$ the classical limit of a. For instance, the classical limit of (2.2) are the classical Dunkl operators

$$y_{i,c}^{(r)} = p_i - \kappa \sum_{j \neq i}^n \frac{1}{x_i - x_j} s_{ij}, \qquad i = 1, \dots, n.$$
(2.8)

These are commuting, equivaraint elements of $A_0 * S_n$, and we have a classical variant of (2.3):

$$\mathbb{C}[V^*] \to A_0 * S_n , \quad q \mapsto q(y_c^{(r)}) .$$
(2.9)

2.4. The classical limit of (2.1) gives the Hamiltonian

$$H_c^{(r)} = \sum_{i=1}^n p_i^2 - \sum_{i< j}^n \frac{2\kappa^2}{(x_i - x_j)^2}.$$
 (2.10)

Its complete integrability can be established similarly to the quantum case. Namely, using the classical variant of the map (2.4), we set

$$L_{q,c}^{(r)} = \operatorname{Res} q(y_c^{(r)}), \qquad q \in \mathbb{C}[V^*]^{S_n}.$$
(2.11)

By construction, $L_{q,c}^{(r)}$ is the classical limit of $L_q^{(r)}$. Hence, the family $\{L_{q,c}^{(r)}, q \in \mathbb{C}[V^*]^{S_n}\}$ is Poisson commutative, i.e. it defines a classical integrable system. Passing to the classical limit in (2.6), we get

$$(y_{1,c}^{(r)})^2 + \dots + (y_{n,c}^{(r)})^2 = \sum_{i=1}^n p_i^2 - \sum_{i< j}^n \frac{2\kappa^2}{(x_i - x_j)^2},$$
(2.12)

which is the Hamiltonian (2.10). Note that in this case there is no need to apply Res. This is true for the higher Hamiltonians as well:

Lemma 2.1 (Lemma 2.2 [EFMV]). For any $q \in \mathbb{C}[V^*]^{S_n}$, we have $q(y_c^{(r)}) \in A_0$. Hence, $L_{q,c}^{(r)} = q(y_c^{(r)})$, i.e. the application of Res is not necessary.

An analogous statement in the elliptic case will be important below.

2.5. By definition, the rational Cherednik algebra $\mathbb{H}_{\hbar,\kappa}$ of type S_n is the subalgebra of $\mathscr{D}(V) * S_n$ generated by $w \in S_n$, x_i (i = 1, ..., n), and the Dunkl operators (2.2). The spherical subalgebra of $\mathbb{H}_{\hbar,\kappa}$ is defined as $e \mathbb{H}_{\hbar,\kappa} e$, where

$$e = \frac{1}{n!} \sum_{w \in S_n} w \,. \tag{2.13}$$

Restricting (2.4) onto the spherical subalgebra, one obtains an algebra map

$$\operatorname{Res}: e_{\operatorname{H}_{\hbar,\kappa}e} \hookrightarrow \mathscr{D}(V)^{S_n}.$$

$$(2.14)$$

whose image is denoted as $B_{\hbar,\kappa}$. Elements of $B_{\hbar,\kappa}$ can be obtained by applying **Res** to symmetric combinations of x_i and $y_i^{(r)}$. Inside $B_{0,\kappa}$ we have the commutative algebra of quantum rational CM Hamiltonians $\{L_q^{(r)}, q \in \mathbb{C}[V^*]^{S_n}\}$.

The classical limits are defined in a similar way. That is, $\mathbb{H}_{0,\kappa}$ is the subalgebra of $A_0 * S_n$ generated by $w \in S_n$, x_i (i = 1, ..., n), and the classical Dunkl operators (2.8). The spherical subalgebra is defined as $e \mathbb{H}_{0,\kappa} e$, and we have an algebra map

$$\operatorname{Res}: e \operatorname{H}_{0,\kappa} e \hookrightarrow A_0^{S_n}.$$

$$(2.15)$$

Its image, $B_{0,\kappa}$, is a Poisson subalgebra of $A_0^{S_n}$, containing a Poisson commutative algebra, $\{L_{q,c}^{(r)}, q \in \mathbb{C}[V^*]^{S_n}\}$, of classical rational CM Hamiltonians.

3. Elliptic Dunkl operators and Calogero-Moser system

3.1. The quantum elliptic CM system is described by the Hamiltonian

$$H = \hbar^2 \sum_{i=1}^n \partial_i^2 - 2\kappa(\kappa - \hbar) \sum_{i < j}^n \wp(x_i - x_j).$$
(3.1)

Here $\wp(z) = \wp(z|1,\tau)$ is the Weierstrass \wp -function. The classical Hamiltonian is

$$H_c = \sum_{i=1}^{n} p_i^2 - 2\kappa^2 \sum_{i< j}^{n} \wp(x_i - x_j).$$
(3.2)

3.2. As in the rational case, both quantum and classical systems are completely integrable [Ca, Pe, OP1, OP2, OS]. Following [EFMV], this can be shown using *elliptic Dunkl operators* [BFV] which, in the quantum case, are

$$y_i = \hbar \partial_i - \kappa \sum_{j \neq i}^n \phi(\lambda_i - \lambda_j, x_i - x_j) s_{ij}, \qquad \phi(\mu, z) = \frac{\sigma(z - \mu)}{\sigma(z)\sigma(-\mu)}.$$
(3.3)

Here $\sigma(z) = \sigma(z|1,\tau)$ is the Weierstrass σ -function, and $\lambda_1, \ldots, \lambda_n$ are auxiliary variables referred to as the *spectral variables*. Hence, $y_i = y_i(\lambda)$ are λ -dependent elements of $\mathscr{D}(V) * S_n$.

For $\xi \in V$, we write $y_{\xi} = \sum_{i=1}^{n} \xi_i y_i$. Two main properties of the operators y_{ξ} are their commutativity and equivariance: for all $\xi, \eta \in V$ and $w \in S_n$,

$$y_{\xi} y_{\eta} = y_{\eta} y_{\xi}, \qquad w y_{\xi}(\lambda) = y_{w\xi}(w\lambda) w.$$
 (3.4)

Note that in the second relation the group action affects both ξ and λ . As before, the assignment $\xi \mapsto y_{\xi}$ extends to an algebra map

$$\mathbb{C}[V^*] \to \mathscr{D}(V) * S_n , \quad q \mapsto q(y) . \tag{3.5}$$

However, unlike in the rational case, this map is not S_n -equivariant and constructing commuting quantum Hamiltonians requires certain regularization.

3.3. To explain the regularization procedure [EFMV], we extend the algebra map (3.5) by allowing polynomials with λ -dependent coefficients:

$$\mathbb{C}(V) \otimes \mathbb{C}[V^*] \to \mathscr{D}(V) * S_n , \quad f \mapsto f(\lambda, y) .$$
 (3.6)

Next, recall the rational classical Hamiltonians $L_{q,c}^{(r)} = q(y_c^{(r)}), q \in \mathbb{C}[V^*]^{S_n}$. These are S_n -invariant elements of $A_0 = \mathbb{C}(V) \otimes \mathbb{C}[V^*]$.

Theorem 3.1 ([EFMV]). For $q \in \mathbb{C}[V^*]^{S_n}$, consider $L_{q,c}^{(r)} \in A_0^{S_n}$. Identify A_0 with $\mathbb{C}(V) \otimes \mathbb{C}[V^*]$ in (3.6) and obtain $L_{q,c}^{(r)}(\lambda, y) \in \mathscr{D}(V) * W$ by applying (3.6), i.e. by replacing classical momenta in $L_{q,c}^{(r)}$ with elliptic Dunkl operators.

(1) The elements $L_{q,c}^{(r)}(\lambda, y)$ are regular near $\lambda = 0$ and so have a well-defined limit at $\lambda = 0$.

(2) Setting $L_q := \operatorname{Res} \lim_{\lambda \to 0} L_{q,c}^{(r)}(\lambda, y)$ defines an algebra embedding $\mathbb{C}[V^*]^{S_n} \to \mathscr{D}(V)^{S_n}, q \mapsto L_q$.

3.4. To illustrate the theorem, let us follow a calculation in [BFV]. Choose $L_{q,c}^{(r)}$ to be the Hamiltonian $H_c^{(r)}$ (2.10). In this case,

$$H_c^{(r)}(\lambda, y) = y_1^2 + \dots + y_n^2 - \sum_{i < j} \frac{2\kappa^2}{(\lambda_i - \lambda_j)^2}.$$

By a direct calculation, with the help of the identity $\phi(\mu, z)\phi(\mu, -z) = \wp(\mu) - \wp(z)$,

$$y_1^2 + \dots + y_n^2 = \hbar^2 \sum_{i=1}^n \partial_i^2 - 2\hbar\kappa \sum_{i$$

Here $\phi'(\mu, z) = \frac{d}{dz}\phi(\mu, z)$. Using that

$$\wp(z) = \frac{1}{z^2} + o(z)$$
 as $z \to 0$, $\lim_{\mu \to 0} \phi'(\mu, z) = -\wp(z)$,

one finds that

$$\lim_{\lambda \to 0} H_c^{(r)}(\lambda, y) = \hbar^2 \sum_{i=1}^n \partial_i^2 - \sum_{i< j}^n 2\kappa(\kappa - \hbar s_{ij})\wp(x_i - x_j).$$
(3.8)

Applying the map (2.4) gives the Hamiltonian (3.1):

$$\operatorname{Res}\lim_{\lambda \to 0} H_c^{(r)}(\lambda, y) = H.$$
(3.9)

Remark 3.2. For the Calogero–Moser particles, their total momentum

$$P = p_1 + \dots + p_n$$

is conserved. Replacing p_i with the Dunkl operators y_i , one finds that

$$y_1 + \dots + y_n = \hbar \partial_1 + \dots + \hbar \partial_n$$
, (3.10)

reflecting conservation of the total momentum in the quantum system.

3.5. By passing to the classical limit in the above constructions, one obtains Hamiltonians for the classical elliptic CM system. Namely, we think of the operators (3.11) as elements of $A_{\hbar} * S_n$, so their classical limit are the following commuting elements of $A_0 * S_n$:

$$y_{i,c} = p_i - \kappa \sum_{j \neq i}^n \phi(\lambda_i - \lambda_j, x_i - x_j) s_{ij}.$$
(3.11)

The classical limit of (3.6) is the map

$$\mathbb{C}(V) \otimes \mathbb{C}[V^*] \to A_0 * S_n , \quad f \mapsto f(\lambda, y_c) . \tag{3.12}$$

Theorem 3.3 ([EFMV]). For $q \in \mathbb{C}[V^*]^{S_n}$, consider $L_{q,c}^{(r)} \in A_0^{S_n}$. Identify A_0 with $\mathbb{C}(V) \otimes \mathbb{C}[V^*]$ in (3.12) and obtain $L_{q,c}^{(r)}(\lambda, y_c) \in A_0 * W$ by applying (3.12), i.e. by replacing classical momenta in $L_{q,c}^{(r)}$ with classical elliptic Dunkl operators.

(1) The elements $L_{q,c}^{(r)}(\lambda, y_c)$ are regular near $\lambda = 0$ and so have a well-defined limit at $\lambda = 0$.

(2) Setting $L_{q,c} := \operatorname{Res} \lim_{\lambda \to 0} L_{q,c}^{(r)}(\lambda, y_c)$ defines a Poisson-commutative subalgebra $\{L_{q,c}, q \in \mathbb{C}[V^*]^{S_n}\}$ in $A_0^{S_n}$.

(3) In particular, similarly to (3.8),

$$\operatorname{Res}\lim_{\lambda \to 0} H_c^{(r)}(\lambda, y_c) = \sum_{i=1}^n p_i^2 - 2\kappa^2 \sum_{i < j}^n \wp(x_i - x_j) \,,$$

which is the Hamiltonian (3.2).

3.6. For later use, let us consider what happens when we substitute elliptic Dunkl operators into *elliptic* Hamiltonians (rather than rational ones).

Proposition 3.4 (Proposition 5.1 [Ch]). For $q \in \mathbb{C}[V^*]^{S_n}$, let $L_{q,c} \in A_0^{S_n}$ be the classcal Hamiltonian from Theorem 3.3. Then:

(1) $L_{q,c}(\lambda, y)$ is regular near $\lambda = 0$,

(2) $L_{q,c}(\lambda, y_c) \in A_0 * S_n$ does not depend on λ . Furthermore, expanding it into $\sum_{w \in S_n} a_w w$, we have $a_w = 0$ for $w \neq id$, i.e. $L_{q,c}(\lambda, y_c) \in A_0 \subset A_0 * S_n$.

Note that part (2) of the proposition is an elliptic analogue of Lemma 2.1. Now, viewing $L_{q,c}(\lambda, y)$ as elements of $A_{\hbar} * S_n$, denote

$$\mathsf{H}_q := \operatorname{Res} \lim_{\lambda \to 0} L_{q,c}(\lambda, y) \,, \qquad q \in \mathbb{C}[V^*]^{S_n}. \tag{3.13}$$

Clearly, H_q belongs to $A_h^{S_n}$ and commutes with all higher CM Hamiltonians².

Corollary 3.5. For $q \in \mathbb{C}[V^*]^{S_n}$, we have

$$\lim_{\lambda \to 0} L_{q,c}(\lambda, y) = \mathsf{H}_q + \hbar \mathsf{A}_q, \qquad \text{for some } \mathsf{A}_q \in (A_\hbar * S_n)^{S_n}.$$
(3.14)

Indeed, the classical limit of the l.h.s. is $\lim_{\lambda\to 0} L_{q,c}(\lambda, y_c) = \operatorname{Res} \lim_{\lambda\to 0} L_{q,c}(\lambda, y_c)$ (the last equality is by part (2) of Prop. 3.4). Hence, $\lim_{\lambda\to 0} L_{q,c}(\lambda, y)$ and H_q have the same classical limit.

4. Spin Calogero-Moser system

The Dunkl operators can be used to construct a spin generalisation of the elliptic Calogero–Moser system. For the trigonometric case, see [P1, BGHP, Che].

²In fact, one can show that $\mathsf{H}_q = L_{q'}$ for some $q' \in \mathbb{C}[V^*]^{S_n}$, but we will not need this.

4.1. Consider the Hamiltonian

$$\widehat{H} = \sum_{i=1}^{n} \hbar^2 \partial_i^2 - 2 \sum_{i< j}^{n} \wp(x_i - x_j) \kappa(\kappa - \hbar \widehat{s}_{ij}).$$
(4.1)

This is viewed as an element of the algebra $\mathscr{D}(V) \otimes \mathbb{C}S_n$, with the tensor-product sign omitted and with the "hat" symbol used to distinguish this from the crossed product $\mathscr{D}(V) * S_n$. Choosing an S_n -module \mathscr{U} makes \widehat{H} a matrix-valued differential operator, i.e. an element of $\mathscr{D}(V) \otimes \operatorname{End}_{\mathbb{C}} \mathscr{U}$, with \widehat{s}_{ij} acting on \mathscr{U} (but not on $\mathbb{C}(V)$). For example, we may choose $\mathscr{U} = U^{\otimes n}$, $U = \mathbb{C}^m$, with \widehat{s}_{ij} acting as the permutation P_{ij} of the tensor factors. Our goal is to construct a commutative subalgebra of $\mathscr{D}(V) \otimes \mathbb{C}S_n$ containing \widehat{H} .

4.2. We begin with considerations as in [P1, BGHP, Che], rephrased for convenience. Let us enlarge the algebra $\mathscr{D}(V) * S_n$ by adding another copy of the group algebra:

$$\mathscr{A} := (\mathscr{D}(V) * S_n) \otimes \mathbb{C}S_n \, .$$

Elements of \mathscr{A} can be uniquely written as

$$a = \sum_{w,w' \in S_n} a_{ww'} w \otimes \widehat{w'} \quad \text{with } a_{ww'} \in \mathscr{D}(V) \,.$$

Define a linear map

$$\widehat{\operatorname{Res}}: \mathscr{A} \to \mathscr{D}(V) \otimes \mathbb{C}S_n, \qquad \sum_{w,w' \in S_n} a_{ww'} w \otimes \widehat{w'} \mapsto \sum_{w,w' \in S_n} a_{ww'} \otimes \widehat{w'w^{-1}}.$$
(4.2)

Equivalently, $\widehat{\operatorname{Res}}(a)$ is the unique element $L_a \in \mathscr{D}(V) \otimes \mathbb{C}S_n$ such that

$$a\widehat{e} = L_a\widehat{e}, \qquad \widehat{e} := \frac{1}{n!} \sum_{w \in S_n} w \otimes \widehat{w}.$$
 (4.3)

We have an S_n -action on \mathscr{A} and $\mathscr{D}(V) \otimes \mathbb{C}S_n$ by conjugation,

$$a \mapsto (w \otimes \widehat{w})a(w^{-1} \otimes \widehat{w^{-1}}) \qquad \forall \ w \in S_n \,.$$

$$(4.4)$$

It is easy to check that the map $\widehat{\text{Res}}$ is S_n -equivariant with respect to this action.

Lemma 4.1. The restriction $\widehat{\text{Res}}$: $\mathscr{A}^{S_n} \to (\mathscr{D}(V) \otimes \mathbb{C}S_n)^{S_n}$ of the map (4.2) is an algebra homomorphism.

Indeed, if $a \in \mathscr{A}^{S_n}$ and $L_a = \widehat{\operatorname{Res}}(a)$, then $a\widehat{e} = \widehat{e}a$ and $L_a\widehat{e} = \widehat{e}L_a$. Hence, for $a, b \in \mathscr{A}^{S_n}$ we have $L_{ab}\widehat{e} = ab\widehat{e} = (a\widehat{e})(b\widehat{e}) = (L_a\widehat{e})(L_b\widehat{e}) = L_aL_b\widehat{e}$.

4.3. On $\mathscr{D}(V) * S_n \subset \mathscr{A}$, the action (4.4) is simply $a \mapsto waw^{-1}$. Thus, restricting (4.2) further gives an algebra map

$$\widehat{\operatorname{Res}} : (\mathscr{D}(V) * S_n)^{S_n} \to (\mathscr{D}(V) \otimes \mathbb{C}S_n)^{S_n}, \qquad \sum_{w \in S_n} a_w w \mapsto \sum_{w \in S_n} a_w \otimes \widehat{w^{-1}}, \quad (4.5)$$

converting reflection-differential operators to $\mathbb{C}S_n$ -valued differential operators. Recall that, by Theorem 3.1, we have well-defined elements $\lim_{\lambda\to 0} L_{q,c}^{(r)}(\lambda, y)$, for every $q \in \mathbb{C}[V^*]^{S_n}$. Since $wL_{q,c}^{(r)}(\lambda, y)w^{-1} = L_{q,c}^{(r)}(w\lambda, y)$, the limit at $\lambda = 0$ is S_n -invariant. Also, since these elements are constructed from commuting Dunkl operators, they pairwise commute. Therefore, we have the following result.

Proposition 4.2. For $q \in \mathbb{C}[V^*]^{S_n}$, define $\widehat{\mathsf{L}}_q = \widehat{\operatorname{Res}} \lim_{\lambda \to 0} L_{q,c}^{(r)}(\lambda, y)$. The family $\{\widehat{\mathsf{L}}_q, q \in \mathbb{C}[V^*]^{S_n}\}$ forms a commutative subalgebra in $(\mathscr{D}(V) \otimes \mathbb{C}S_n)^{S_n}$.

Using our calculations in $\S3.4$, we find

$$\mathbf{L}_{q} = \hbar\partial_{1} + \dots + \hbar\partial_{n} \qquad \qquad \text{for } q = e_{1} + \dots + e_{n}, \quad (4.6)$$
$$\widehat{\mathbf{L}}_{q} = \sum_{n=1}^{n} \hbar^{2} \partial^{2} - 2\sum_{n=1}^{n} e_{n}(\mathbf{x}_{n} - \mathbf{x}_{n}) \mathbf{u}(\mathbf{x}_{n} - \hbar\widehat{\mathbf{x}}_{n}) \qquad \qquad \text{for } q = e_{1}^{2} + \dots + e_{n}^{2}, \quad (4.7)$$

$$\widehat{\mathsf{L}}_q = \sum_{i=1} \hbar^2 \partial_i^2 - 2 \sum_{i < j} \wp(x_i - x_j) \kappa(\kappa - \hbar \widehat{s}_{ij}) \qquad \text{for } q = e_1^2 + \dots + e_n^2 \,. \tag{4.7}$$

Hence, the operator (4.1) belongs to the constructed commutative family.

Remark 4.3. The algebra $\mathscr{D}(V) * S_n$ has a nontrivial centre, spanned by the central idempotents $c_{\pi} \in \mathbb{C}S_n$, $\pi \in \mathrm{Irr}S_n$. The elements $\widehat{c_{\pi}}$ commute with any of the spin Hamiltonians.

4.4. Due to extra degrees of freedom in the spin system, one expects more commuting Hamiltonians. Recall the classical spherical subalgebra $B_{0,\kappa}$, a Poisson subalgebra of $A_0^{S_n}$, see §2.5. Also, recall the map $A_0 \to \mathscr{D}(V) * S_n$, $f \mapsto f(\lambda, y)$, see (3.6). We have the following result strengthening Proposition 4.2.

Theorem 4.4. For any $f \in B_{0,\kappa}$, the element $f(\lambda, y) \in \mathscr{D}(V) * S_n$ is regular near $\lambda = 0$. Define $\widehat{\mathsf{L}}_f := \widehat{\mathsf{Res}} \lim_{\lambda \to 0} f(\lambda, y)$. Then the family $\{\widehat{\mathsf{L}}_f, f \in \mathsf{B}_{0,\kappa}\}$ forms a commutative subalgebra in $(\mathscr{D}(V) \otimes \mathbb{C}S_n)^{S_n}$.

The fact that $f(\lambda, y)$ are regular near $\lambda = 0$ for any $f \in B_{0,\kappa}$ follows from the second proof of [EFMV, Theorem 3.1]. To be precise, in *loc. cit.* this is shown for $f = L_{q,c}^{(r)}$, cf. Theorem 3.1. However, the proof applies verbatim to any $f \in B_{0,\kappa}$, see [EFMV, Sec. 5.3] for the details. The remaining statements are clear.

4.5. We can also substitute Dunkl operators into elliptic Hamiltonians (cf. Prop. 3.4) and obtain

$$\widehat{\mathsf{H}}_q := \widehat{\operatorname{Res}} \lim_{\lambda \to 0} L_{q,c}(\lambda, y) \,, \qquad q \in \mathbb{C}[V^*]^{S_n} \,. \tag{4.8}$$

These obviously commute between themselves and with any of the Hamiltonians $\widehat{\mathsf{L}}_f$, $f \in \mathsf{B}_{0,\kappa}$.³ We will refer to the elements $\widehat{\mathsf{L}}_f$, $f \in \mathsf{B}_{0,\kappa}$ as higher spin CM Hamiltonians, and to $\widehat{\mathsf{H}}_q$, $q \in \mathbb{C}[V^*]^{S_n}$ as principal spin CM Hamiltonians.

5. INOZEMTSEV QUANTUM SPIN CHAIN

We are now in a position to construct commuting Hamiltonians for the Inozemtsev spin chain (1.1). They will be obtained from the spin CM Hamiltonians by taking classical limits and evaluating at an equilibrium. In particular, this will provide a full justification for the "freezing" procedure [P2].

5.1. Consider the classical Hamiltonian H (3.2), assuming $x_i, p_i \in \mathbb{R}$. Suppose $\tau \in i\mathbb{R}_+$, so $\wp(z)$ is real for real z. The following fact is well known (it follows easily from the convexity of $\wp(z)$ for 0 < z < 1).

Lemma 5.1. In the region $x_1 < \cdots < x_n < x_1 + 1$, all possible equilibria (x, p) of H are of the form $x_i = c + \frac{i}{n}$ (with arbitrary $c \in \mathbb{R}$), $p_i = 0$ (i = 1, ..., n).

³One can show that, in fact, $H_q = \widehat{L}_f$ for some $f \in B_{0,\kappa}$, but we will not need this result.

Write (x^*, p^*) for an equilibrium of H of the form $x_i = \frac{i}{n}$, $p_i = 0$.

Corollary 5.2. If a function F = F(x, p) satisfies $\{F, H\} = \{F, x_1 + \dots + x_n\} = 0$ then (x^*, p^*) is an equilibrium for F.

Indeed, the Hamiltonian flow defined by F should map an equilibrium for H to an equilibrium, at the same time preserving $x_1 + \cdots + x_n$.

Remark 5.3. Assuming that F depends analytically on τ , the above corollary remains valid for arbitrary, not necessarily real, τ . This further implies that for any primitive period $\omega \in \mathbb{Z} + \mathbb{Z}\tau$, the point (x, p) with $x_i = \frac{i}{n}\omega$, $p_i = 0$ is an equilibrium for F.

5.2. Recall CM Hamiltonians H_q , see § 3.6. Let us restrict the choice of q to

$$\mathbb{C}[V^*]_0^{S_n} := \mathbb{C}[V^*]^{S_n} / \langle e_1 + \dots + e_n \rangle$$

In other words, we assume that q is a symmetric polynomial of

$$\xi_i = e_i - \frac{1}{n}(e_1 + \dots + e_n), \qquad i = 1, \dots, n$$

Let $y_i^0 = y_{\xi_i}$ be the corresponding elliptic Dunkl operators. It is easy to check that $[y_i^0, x_1 + \cdots + x_n] = 0$. Hence, the quantum CM Hamiltonians H_q built from y_i^0 would commute with $x_1 + \cdots + x_n$, and their classical limit, $\mathsf{H}_{q,c}$, would Poisson-commute with $x_1 + \cdots + x_n$, so we can apply the Corollary 5.2.

Proposition 5.4. The classical CM Hamiltonians $\mathsf{H}_{q,c} = \eta_0(\mathsf{H}_q)$ with $q \in \mathbb{C}[V^*]_0^{S_n}$ have a joint equilibrium at (x^*, p^*) . Therefore,

$$\{\mathsf{H}_{q,c}, G\} = 0 \quad at \ (x, p) = (x^*, p^*) \qquad \forall \ G = G(x, p). \tag{5.1}$$

5.3. Recall the commuting spin CM Hamiltonians \widehat{L}_f , \widehat{H}_q , see § 4.4, § 4.5. Let us view them as elements of $A_\hbar \otimes \mathbb{C}S_n$, so we can take classical limits using a map, similar to (2.7):

$$\eta_0 : A_{\hbar} \otimes \mathbb{C}S_n \to A_0 \otimes \mathbb{C}S_n, \qquad f \mapsto f, \quad \hat{p}_i \mapsto p_i, \quad w \mapsto w.$$

By applying $\widehat{\text{Res}}$ to (3.14), we have

$$\widehat{\mathsf{H}}_q = \mathsf{H}_q + \hbar \widehat{\mathsf{A}}_q \qquad \text{with } \widehat{\mathsf{A}}_q \in A_\hbar \otimes \mathbb{C}S_n \,.$$
 (5.2)

We define *principal Inozemtsev Hamiltonians* by taking the classical limit of the spin part \widehat{A}_q and evaluating it at equilibrium, i.e.

$$\mathscr{H}_q := \eta_0(\widehat{\mathsf{A}}_q)|_{(x,p)=(x^*,p^*)}, \quad q \in \mathbb{C}[V^*]^{S_n}.$$
(5.3)

For example, for $q = e_1^2 + \cdots + e_n^2$ one finds from (4.7):

$$\mathscr{H}_q = 2\kappa \sum_{i < j}^n \wp\left(\frac{i-j}{n}\right) \left(\widehat{s}_{ij} - \widehat{\mathrm{id}}\right).$$

The multiple of the identity can be removed so this is essentially the same as (1.1).

Theorem 5.5. The elements $\mathscr{H}_q \in \mathbb{C}S_n$ pairwise commute.

Proof. First, note that for $q = e_1 + \cdots + e_n$, $\mathscr{H}_q = 0$. Hence, it is sufficient to prove commutativity for $q \in \mathbb{C}[V^*]_0^{S_n}$. We need the following elementary lemma.

Lemma 5.6. Let $B = B_0 + \hbar B_1$, $C = C_0 + \hbar C_1$ be elements of $A_{\hbar} \otimes \mathbb{C}S_n$ with $B_0, C_0 \in A_{\hbar} \subset A_{\hbar} \otimes \mathbb{C}S_n$. Denote by b_0, b_1, c_0, c_1 the classical limits of B_0, B_1, C_0, C_1 . Then, assuming $[B_0, C_0] = 0$, one has $[B, C] \in \hbar^2 A_{\hbar} \otimes \mathbb{C}S_n$, and

$$\eta_0(\hbar^{-2}[B,C]) = \{b_0,c_1\} - \{c_0,b_1\} + [b_1,c_1]$$

Here the Poisson bracket in the r.h.s. is understood as $\{a, b \otimes \widehat{w}\} := \{a, b\} \otimes \widehat{w}$ for $a, b \in A_0$.

Now, given $q, q' \in \mathbb{C}[V^*]_0^{S_n}$, we apply the lemma to $B = \widehat{H}_q$, $C = \widehat{H}_{q'}$ and use the decomposition (5.2). Since [B, C] = 0 in this case, we get $0 = \{b_0, c_1\} - \{c_0, b_1\} + [b_1, c_1]$. The first two terms in the r.h.s. vanish at $(x, p) = (x^*, p^*)$, by Prop. 5.4. As a result, $[b_1, c_1]$ vanishes at (x^*, p^*) , i.e. $[\mathscr{H}_q, \mathscr{H}_{q'}] = 0.4$

5.4. We proceed to define higher Inozemtsev Hamiltonians by

$$\mathscr{H}_f := \eta_0(\mathbf{L}_f)|_{(x,p)=(x^*,p^*)}, \quad f \in \mathsf{B}_{0,\kappa}.$$
(5.4)

Theorem 5.7. The elements \mathcal{H}_f pairwise commute, as well as commuting with the principal Hamiltonians \mathcal{H}_q .

The fact that $[\mathscr{H}_f, \mathscr{H}_{f'}] = 0$ for $f, f' \in \mathsf{B}_{0,\kappa}$ is immediate from $[\widehat{\mathsf{L}}_f, \widehat{\mathsf{L}}_{f'}] = 0$ (evaluation at (x^*, p^*) plays no role here). The fact that $[\mathscr{H}_q, \mathscr{H}_f] = 0$ follows from the following lemma (obtained by setting $C_0 = 0$ in Lemma 5.6).

Lemma 5.8. Let $B = B_0 + \hbar B_1$, C be elements of $A_{\hbar} \otimes \mathbb{C}S_n$ with $B_0 \in A_{\hbar} \subset A_{\hbar} \otimes \mathbb{C}S_n$. Denote by b_0, b_1, c the classical limits of B_0, B_1, C . Then one has $[B, C] \in \hbar A_{\hbar} \otimes \mathbb{C}S_n$, and

$$\eta_0(\hbar^{-1}[B,C]) = \{b_0,c\} + [b_1,c].$$

Remark 5.9. The above results and constructions remain valid for elliptic Calogero-Moser systems associated for any root system (including the BC_n version) - all the necessary ingredients can already be found in the cited sources. In general, equilibria for these systems will not have such a simple form as above, which makes the corresponding spin chains less natural. In some cases, however, simple equilibria configurations are possible, see, e.g., [IS, BPS].

6. Explicit examples

Here we calculate two of the Hamiltonians explicitly, as an illustration and to compare with known results.

6.1. We use the following shorthand notation: $\phi_{ij}^{ab} = \phi(\lambda_a - \lambda_b, x_i - x_j), \phi_{ij} := \phi_{ij}^{ij},$ $h_{ij}^{ab} = -\phi'(\lambda_a - \lambda_b, x_i - x_j), h_{ij} := h_{ij}^{ij}, u_{ij} = \wp(x_i - x_j), u^{ab} = \wp(\lambda_a - \lambda_b), \theta_i = \sum_{j \neq i} \phi_{ij} s_{ij}.$ With these notations, the Dunkl operators are $y_i = \hat{p}_i - \kappa \theta_i$. Let us substitute them into the cubic classical Hamiltonian $L = L_q$ with $q = \sum_{i < j < k} e_i e_j e_k$:

$$L = \sum_{i < j < k} \left(p_i p_j p_k + \kappa^2 u^{ij} p_k + \kappa^2 u^{ik} p_j + \kappa^2 u^{jk} p_i \right) \,.$$

⁴When finalising this paper, we learned about the work [LRS] where a similar idea was used for the Haldane–Shastry chain.

Thus, we need to evaluate

$$L(\lambda, y) = \frac{1}{6} \sum_{i \neq j \neq k} y_i y_j y_k + \kappa^2 \sum_{i \neq j \neq k} u^{ij} y_k.$$
(6.1)

First, consider

 $y_i y_j y_k = \hat{p}_i \hat{p}_j \hat{p}_k - \kappa (\hat{p}_i \hat{p}_j \theta_k + \hat{p}_i \theta_j \hat{p}_k + \theta_i \hat{p}_j \hat{p}_k) + \kappa^2 (\theta_i \theta_j \hat{p}_k + \theta_i \hat{p}_j \theta_k + \hat{p}_i \theta_j \theta_k) - \kappa^3 \theta_i \theta_j \theta_k$. When expanding each θ_a in terms of $\phi_{ab} s_{ab}$ and taking the sum of these over all $i \neq j \neq k$, some terms cancel due to $\phi_{ia} = -\phi_{ai}$. For example, the term $\hat{p}_i \hat{p}_j \phi_{ak} s_{ka}$ (with $a \neq i, j, k$) coming from $y_i y_j y_k$ cancels with a similar term $\hat{p}_i \hat{p}_j \phi_{ak} s_{ak}$ coming from $y_i y_j y_a$. In addition, some terms linear in momenta cancel due to Fay's identity,

$$\phi_{ij}^{ik}\phi_{jk} = \phi_{ik}^{jk}\phi_{ij} + \phi_{jk}^{ji}\phi_{ik} \qquad (i \neq j \neq k)$$

which implies that for $a \neq i, j$

$$\phi_{ia}s_{ia}\phi_{ja}s_{ja} + (\text{cyclic permutations of } i, j, a) = 0$$

Similarly, when expanding $\theta_i \theta_j \theta_k$, we may neglect terms with indices outside $\{i, j, k\}$ because of the previous identity and its higher analogue, valid for $a \neq i, j, k$:

 $\phi_{ia}s_{ia}\phi_{ja}s_{ja}\phi_{ka}s_{ka} + (\text{cyclic permutations of } i, j, k, a) = 0.$

As a result, when calculating $\sum_{i \neq j \neq k} y_i y_j y_k$, it is sufficient to keep in each $y_i y_j y_k$ only the terms with indices from $\{i, j, k\}$ set (i.e. calculating $y_i y_j y_k$ as if it was just for three particles at x_i, x_j, x_k .) By the same token, in the second sum in (6.1), we have a cancellation of, e.g., $u^{ij} \phi_{ka} s_{ka}$ with $u^{ij} \phi_{ak} s_{ak}$ (if $a \neq i, j, k$). As a result, we may replace

$$u^{ij}y_k \mapsto u^{ij}(\hat{p}_k - \kappa \phi_{ki}s_{ki} - \kappa \phi_{kj}s_{kj}).$$

Altogether, (6.1) can be replaced with $\sum_{i < j < k} H_{ijk}$ where

$$H_{ijk} = (\hat{p}_i - \kappa \phi_{ij} s_{ij} - \kappa \phi_{ik} s_{ik})(\hat{p}_j - \kappa \phi_{ji} s_{ji} - \kappa \phi_{jk} s_{jk})(\hat{p}_k - \kappa \phi_{ki} s_{ki} - \kappa \phi_{kj} s_{kj}) + \kappa^2 \left\{ u^{ij}(\hat{p}_k - \kappa \phi_{ki} s_{ki} - \kappa \phi_{kj} s_{kj}) + \text{ (cyclic permutations of } i, j, k) \right\}.$$

With the help of Fay's identity and $\phi_{ij}^{ij}\phi_{ij}^{ji} = u_{ij} - u^{ij}$, this becomes

$$\begin{split} H_{ijk} = & \hat{p}_i \hat{p}_j \hat{p}_k + \kappa^2 u_{ij} \hat{p}_k + \kappa^2 u_{ik} \hat{p}_j + \kappa^2 u_{jk} \hat{p}_i \\ & - \hbar \kappa (h_{ij} \hat{p}_k s_{ij} + h_{jk} \hat{p}_i s_{jk} + h_{ki} \hat{p}_j s_{ki}) \\ & + \hbar \kappa^2 \left\{ (h_{ij} \phi_{ki}^{kj} + h_{ij}^{ik} \phi_{jk}) + \text{ (cyclic permutations of } i, j, k) \right\} (ijk) \\ & + \hbar \kappa^2 \left\{ (h_{ji} \phi_{kj}^{ki} + h_{ji}^{jk} \phi_{ik}) + \text{ (cyclic permutations of } i, j, k) \right\} (kji) \,. \end{split}$$

Taking $\lambda \to 0$ gives

$$\begin{aligned} H_{ijk} = \hat{p}_i \hat{p}_j \hat{p}_k + \kappa^2 u_{ij} \hat{p}_k + \kappa^2 u_{ik} \hat{p}_j + \kappa^2 u_{jk} \hat{p}_i \\ &- \hbar \kappa (u_{ij} \hat{p}_k s_{ij} + u_{jk} \hat{p}_i s_{jk} + u_{ki} \hat{p}_j s_{ki}) \\ &+ \hbar \kappa^2 \left\{ (u_{ij} + u_{jk} + u_{ki}) (\zeta_{ij} + \zeta_{jk} + \zeta_{ki}) + \frac{1}{2} (u'_{ij} + u'_{jk} + u'_{ki}) \right\} (ijk) \\ &+ \hbar \kappa^2 \left\{ (u_{kj} + u_{ji} + u_{ik}) (\zeta_{kj} + \zeta_{ji} + \zeta_{ik}) + \frac{1}{2} (u'_{kj} + u'_{ji} + u'_{ik}) \right\} (kji) \end{aligned}$$

Applying $\widehat{\text{Res}}$ gives a spin Hamiltonian that matches the expression for I_2 in [DI2]. Furthermore, picking terms of order \hbar and evaluating at an equilibrium in the classical limit, we get a principal Inozemtsev Hamiltonian

$$\mathscr{H}_{3} = \sum_{i < j < k} \left\{ (\wp_{ij} + \wp_{jk} + \wp_{ki})(\zeta_{ij} + \zeta_{jk} + \zeta_{ki}) + \frac{1}{2}(\wp_{ij}' + \wp_{jk}' + \wp_{ki}') \right\} (\widehat{(ijk)} - \widehat{(kji)}) .$$

Here $\wp_{ij} = \wp(\frac{i-j}{n}), \ \zeta_{ij} = \zeta(\frac{i-j}{n}), \ \wp'_{ij} = \wp'(\frac{i-j}{n}).$ Note that by the addition formula for $\wp(z), \ \wp_{ij} + \wp_{jk} + \wp_{ki} = (\zeta_{ij} + \zeta_{jk} + \zeta_{ki})^2$, so \mathscr{H}_3 matches \hat{I}_2 from [I1].

6.2. Let us also consider higher Hamiltonians obtained from $f \in B_{0,\kappa}$. First, let us try

$$f = \operatorname{Res} \sum_{i=1}^n x_i y_{i,c}^{(r)} = \sum_{i=1}^n x_i p_i$$

Replacing the classical momenta with elliptic Dunkl operators, we get

$$f(\lambda, y) = \sum_{i=1}^{n} \lambda_i y_i = \sum_{i=1}^{n} \lambda_i (\hat{p}_i + \kappa \sum_{j \neq i} \phi_{ij} s_{ij}).$$

We have

$$\phi_{ij} = -\frac{1}{\lambda_i - \lambda_j} + \zeta(x_i - x_j) + o(1) \quad \text{near } \lambda = 0.$$
(6.2)

Using this in the previous formula, we find that

$$\lim_{\lambda \to 0} f(\lambda, y) = -\sum_{i < j} s_{ij}$$

which commutes with other spin Hamiltonians for trivial reasons. A more interesting choice is

$$f = \sum_{i \neq j \neq k} x_i \left(p_j p_k + \frac{\kappa^2}{x_{jk}^2} \right) \,.$$

Such f can be obtained by applying **Res** to $\sum_{i \neq j \neq k} x_i y_{j,c}^{(r)} y_{k,c}^{(r)}$ (and subtracting a multiple of $p_1 + \cdots + p_n$). We then need to consider

$$f(\lambda, y) = \sum_{i \neq j \neq k} \lambda_i \left(y_j y_k + \frac{\kappa^2}{\lambda_{jk}^2} \right) \,.$$

Since we are interested in the $\lambda \to 0$ limit, we may expand $y_j y_k$ using (6.2) and keep track only of the terms singular in λ . After a straightforward rearrangement, the limiting form of $f(\lambda, y)$ is found as

$$-\kappa \sum_{i \neq j \neq k} \hat{p}_i s_{jk} - \kappa^2 \sum_{i < j < k} \left\{ \zeta(x_i - x_j) + \zeta(x_j - x_k) + \zeta(x_k - x_i) \right\} \left((ijk) - (kji) \right).$$

Applying Res gives an additional spin CM Hamiltonian

$$\widehat{\mathsf{H}}_{f} = -\kappa \sum_{i \neq j \neq k} \widehat{p}_{i} \widehat{s_{jk}} + \kappa^{2} \sum_{i < j < k} \left\{ \zeta(x_{i} - x_{j}) + \zeta(x_{j} - x_{k}) + \zeta(x_{k} - x_{i}) \right\} (\widehat{(ijk)} - \widehat{(kji)}).$$

This matches the quantum spin Hamiltonian I_1 found in [DI2]. Taking classical limit and evaluating at equilibrium, we obtain (setting $\kappa = 1$)

$$\mathscr{H}_{f} = \sum_{i < j < k} \left\{ \zeta_{ij} + \zeta_{jk} + \zeta_{ki} \right\} \left(\widehat{(ijk)} - \widehat{(kji)} \right), \qquad \zeta_{ij} = \zeta \left(\frac{i-j}{n} \right) \,.$$

This coincides with \hat{I}_1 from [I1]. Note that it is surprisingly difficult to check that $[\hat{I}_1, \hat{I}_2] = 0$ by a direct calculation, see [DI1].

Remark 6.1. For calculating \mathscr{H}_f with $f(x, p) \in B_{0,\kappa}$, we may take classical limit and evaluate at equilibrium immediately, i.e. substitute $x_i \mapsto \lambda$, $p_i \mapsto \theta_i = \sum_{j \neq i} \phi_{ij} s_{ij}$. However, finding the limiting value at $\lambda \to 0$ seems a lengthy process in general.

7. A GENERALISATION

7.1. Consider the following modification of the Hamiltonian (1.1):

$$\mathscr{H}^{\vee} = \sum_{i < j}^{n} \phi'(\lambda_i - \lambda_j, \frac{i - j}{n}) P_{ij}.$$
(7.1)

Here $\phi'(\mu, z) = \frac{d}{dz}\phi(\mu, z)$ is considered as a formal series in μ ,

$$\phi'(\mu, z) = \sum_{i=0}^{\infty} c_i \mu^i, \qquad c_0 = -\wp(z), \ c_1 = \wp'(z) + 2\zeta(z)\wp(z), \dots,$$

making $\phi'(\lambda_i - \lambda_j, \frac{i-j}{n})$ a formal series in λ_i, λ_j . The operator (7.1) is viewed as acting on the tensor product $U^{\otimes n}$ of $U = \mathbb{C}^m \otimes \mathbb{C}[[s]]$, with P_{ij} permuting the factors, and with the *i*th spectral variable λ_i acting as $s^{(i)}$, i.e. as multiplication by *s* on the *i*th factor and identity on the others. Note that the action of \mathscr{H} naturally descends onto $(U/s^{l+1}U)^{\otimes n}$, for any $l \geq 0$. In particular, taking l = 0 recovers the original Hamiltonian (1.1).

7.2. To demonstrate the integrability of the Hamiltonian (7.1), we first construct a dynamical spin model, modifying the approach in §4 to incorporate dependence on the spectral variables λ . It will be convenient to introduce an auxiliary space $V^{\vee} \cong V$ with $\lambda \in V^{\vee}$, and denote by S_n^{\vee} a copy of the symmetric group acting on V^{\vee} . Consider the algebra $\mathscr{D}^{\vee}(V)$ of differential operators on V whose coefficients also depend on λ ; it is generated by (operators of multiplication by) functions $f \in \mathbb{C}(V \times V^{\vee})$ and the derivations $\partial_i = \partial/\partial x_i$. The group $S_n \times S_n^{\vee}$ acts on $V \times V^{\vee}$ and on $\mathscr{D}^{\vee}(V)$, so we can form a crossed product

$$\mathscr{A}^{\vee} := \mathscr{D}^{\vee}(V) * (S_n \times S_n^{\vee}).$$

Elements of \mathscr{A}^{\vee} can be uniquely written as

$$a = \sum_{w_1, w_2 \in S_n} a_{w_1 w_2} w_1 \otimes w_2^{\vee} \quad \text{with } a_{w_1 w_2} \in \mathscr{D}^{\vee}(V) \,.$$

Define a linear map

$$\operatorname{Res}^{\vee}: \mathscr{A}^{\vee} \to \mathscr{D}^{\vee}(V) * S_n^{\vee},$$
$$\sum_{w_1, w_2 \in S_n} a_{w_1 w_2} w_1 \otimes w_2^{\vee} \mapsto \sum_{w_1, w_2 \in S_n} a_{w_1 w_2} (w_2 w_1^{-1})^{\vee}.$$
(7.2)

Equivalently, $\operatorname{Res}^{\vee}(a)$ is the unique element $L_a \in \mathscr{D}^{\vee}(V) * S_n^{\vee}$ such that

$$ae^{\vee} = L_a e^{\vee}, \qquad e^{\vee} := \frac{1}{n!} \sum_{w \in S_n} w \otimes w^{\vee}.$$
 (7.3)

We have an S_n -action on \mathscr{A}^{\vee} and $\mathscr{D}^{\vee}(V) * S_n^{\vee}$ by conjugation,

$$a \mapsto (w \otimes w^{\vee})a(w \otimes w^{\vee})^{-1} \qquad \forall \ w \in S_n \,.$$

$$(7.4)$$

It is easy to check that the map $\operatorname{Res}^{\vee}$ is S_n -equivariant with respect to this action.

Lemma 7.1. The restriction $\operatorname{Res}^{\vee}$: $(\mathscr{A}^{\vee})^{S_n} \to (\mathscr{D}^{\vee}(V) * S_n^{\vee})^{S_n}$ of the map (4.2) is an algebra homomorphism.

Proof is identical to that of Lemma 4.1.

7.3. Restricting (7.2) further gives an algebra map

$$\begin{split} \operatorname{Res}^{\vee} &: (\mathscr{D}^{\vee}(V) \ast S_n)^{S_n} \to (\mathscr{D}^{\vee}(V) \ast S_n^{\vee})^{S_n} \,, \\ & \sum_{w \in S_n} a_w w \mapsto \sum_{w \in S_n} a_w (w^{-1})^{\vee} \,. \end{split}$$

Comparing with a similar map in §4, it takes elements invariant under "diagonal" S_n -action (on both x, λ). Another difference is the crossed-product structure in the resulting operator so it acts on the spectral variables.

As for examples of elements in $(\mathscr{D}^{\vee}(V) * S_n)^{S_n}$, we can substitute elliptic Dunkl operators into any S_n -invariant function on $V^{\vee} \otimes V^*$. However, we would like the result to be regular near $\lambda = 0$ so we can expand the coefficients into a formal series on λ as we did for (7.1). This motivates the following definitions:

$$\mathsf{L}_{f}^{\vee} := \operatorname{\mathsf{Res}}^{\vee} f(\lambda, y) \,, \quad \mathsf{H}_{q}^{\vee} := \operatorname{\mathsf{Res}}^{\vee} L_{q,c}(\lambda, y) \,, \qquad \text{with } f \in \mathsf{B}_{0,\kappa} \,, q \in \mathbb{C}[V^{*}]^{S_{n}} \,. \tag{7.5}$$

These are similar to the definitions in §4, but one important difference is that we do not take limit $\lambda \to 0$.

Theorem 7.2. The elements L_{f}^{\vee} and H_{q}^{\vee} are pairwise commuting, S_{n} -equivariant elements of $\mathscr{D}^{\vee}(V) * S_{n}^{\vee}$, regular near $\lambda = 0$.

Proofs are the same as in $\S 4$.

As an example, using the calculations in §3.4, we find that for $q = e_1^2 + \cdots + e_n^2$,

$$\mathsf{H}_{q}^{\vee} = \sum_{i=1}^{n} \hbar^{2} \partial_{i}^{2} - 2\kappa^{2} \sum_{i< j}^{n} \wp(x_{i} - x_{j}) - 2\hbar\kappa \sum_{i< j}^{n} \phi'(\lambda_{i} - \lambda_{j}, x_{i} - x_{j}) s_{ij}^{\vee}.$$
(7.6)

This is a generalisation of the spin Calogero–Moser Hamiltonian (4.1). Choosing any $\mathbb{C}[[V^{\vee}]] * S_n^{\vee}$ -module \mathscr{U} , and expanding H_q^{\vee} into a formal series in λ makes it a differential operator acting on \mathscr{U} -valued functions $\varphi : V \to \mathscr{U}$. For example, we can choose $\mathscr{U} = U^{\otimes n}$ with $U = \mathbb{C}^m \otimes \mathbb{C}[[s]]$ and with λ_i acting by $s^{(i)}$, in the same way as it was done for the Hamiltonian (7.1).

7.4. Now the commuting Hamiltonians for the spin chain (7.1) can be constructed in the same way as before. We omit the proofs as they are identical to those in § 5 and use evaluation at an equilibrium (x^*, p^*) .

Let us view the Hamiltonians (7.5) as elements of $A_{\hbar} * S_n^{\vee}$, so we can take classical limits. We have a counterpart of (5.2):

$$\mathsf{H}_{q}^{\vee} = \mathsf{H}_{q} + \hbar \mathsf{A}_{q}^{\vee} \qquad \text{with } \mathsf{A}_{q}^{\vee} \in A_{\hbar} * S_{n}^{\vee} \,. \tag{7.7}$$

We define *principal Hamiltonians* by

$$\mathscr{H}_{q}^{\vee} := \eta_{0}(\mathsf{A}_{q}^{\vee})|_{(x,p)=(x^{*},p^{*})}, \quad q \in \mathbb{C}[V^{*}]^{S_{n}}.$$
(7.8)

For example, for $q = e_1^2 + \cdots + e_n^2$ one finds from (7.6):

$$\mathscr{H}_{q}^{\vee} = -2\kappa \sum_{i< j}^{n} \phi'\left(\lambda_{i} - \lambda_{j}, \frac{i-j}{n}\right) s_{ij}^{\vee} - 2\kappa \sum_{i< j}^{n} \wp\left(\frac{i-j}{n}\right) \mathrm{id}^{\vee}.$$
(7.9)

The multiple of the identity can be removed, giving

$$\mathscr{H}^{\vee} = \sum_{i < j}^{n} \phi'\left(\lambda_i - \lambda_j, \frac{i - j}{n}\right) s_{ij}^{\vee}.$$
(7.10)

We also define *higher Hamiltonians* by

$$\mathscr{H}_f^{\vee} := \eta_0(\mathsf{L}_f^{\vee})|_{(x,p)=(x^*,p^*)}, \quad f \in \mathsf{B}_{0,\kappa}.$$

$$(7.11)$$

Theorem 7.3. The elements \mathscr{H}_q^{\vee} , \mathscr{H}_f^{\vee} with $q \in \mathbb{C}[V^*]^{S_n}$, $f \in \mathsf{B}_{0,\kappa}$ form a commutative family in $\mathbb{C}[[V^{\vee}]] * S_n^{\vee}$.

Choosing $\mathscr{U} = U^{\otimes n}$ with $U = \mathbb{C}^m \otimes \mathbb{C}[[s]]$ as a representation of $\mathbb{C}[[V^{\vee}]] * S_n^{\vee}$ identifies (7.10) with the Hamiltonian (7.1), hence, it is integrable.

References

- [BGHP] D. Bernard, M. Gaudin, F. D. M. Haldane, V. Pasquier: Yang-Baxter equation in spin chains with long range interactions. J. Phys. A: Math. Gen. 26, 5219–523 (1993)
- [BPS] D. Bernard, V. Pasquier, D. Serban: Exact solution of long-range interacting spin chains with boundaries. Europhys. Lett. 30 301–306 (1995)
- [BFV] V. Buchstaber, G. Felder, A. Veselov: Elliptic Dunkl operators, root systems, and functional equations. Duke Math. J. 76 (3) (1994), 885–911.
- [Ca] F. Calogero: Exactly solvable one-dimensional many-body systems. Lett. Nuovo Cimento 13, 411–415 (1975)
- [Ch] O. Chalykh: Quantum Lax pairs via Dunkl and Cherednik operators. Commun. Math. Phys. 369(1), 261–316 (2019)
- [ChL] O. Chalykh, J. Lamers: R-matrix Calogero-Moser systems and spin chains. In preparation.
- [Che] I. Cherednik: Integration of quantum many-body problems by affine Knizhnik-Zamolodchikov equations. Adv. Math. 106(1), 65–95 (1994)
- [DI1] J. Dittrich, V. Inozemtsev: The commutativity of integrals of motion for quantum spin chains and elliptic functions identities. Regul. Chaotic Dyn. 13, 19–26 (2008)
- [DI2] J. Dittrich, V. I. Inozemtsev: Towards the proof of complete integrability of quantum elliptic many-body systems with spin degrees of freedom. Regul. Chaotic Dyn. 14, 218–222 (2009)
- [E] P. Etingof: Calogero-Moser systems and representation theory. Zürich Lectures in Advanced Mathematics, Eur. Math. Soc., Zürich 2007.
- [EFMV] P. Etingof, G. Felder, X. Ma, A. Veselov: On elliptic Calogero-Moser systems for complex crystallographic reflection groups. J. Algebra 329, 107–129 (2011)
- [FGL] F. Finkel, A. González-López: A new perspective on the integrability of Inozemtsev's elliptic spin chain. Ann. Phys. 351, 797–827 (2014)

- [FM] M. Fowler, J. A. Minahan: Invariants of the Haldane-Shastry SU(N) chain. Phys. Rev. Lett. 70(15), 2325–2328 (1993)
- [H] F. D. M. Haldane: Exact Jastrow-Gutzwiller resonating-valence-bond ground state of the spin-1/2 antiferromagnetic Heisenberg chain with 1/r² exchange. Phys. Rev. Lett. **60**, 635–638 (1988)
- [I1] V. I. Inozemtsev: On the connection between the one-dimensional S = 1/2 Heisenberg chain and Haldane–Shastry model. J. Stat. Phys. **59**, 1143–1155 (1990)
- [I2] V. Inozemtsev: Integrable Heisenberg-van Vleck chains with variable range exchange. Phys. Part. Nucl. 34, 166–193 (2003)
- [IS] V. I. Inozemtsev, R. Sasaki: Universal Lax pairs for spin Calogero-Moser models and spin exchange models. J. Phys. A: Math. Gen. 34(37), 7621–7632 (2001)
- [KL] R. Klabbers, J. Lamers: How coordinate Bethe ansatz works for Inozemtsev model. Comm. Math. Phys. 390(2), 827–905 (2022)
- [LOZ] A. Levin, M. Olshanetsky, A. Zotov, A.: Planck constant as spectral parameter in integrable systems and KZB equations. JHEP 10, 1–29 (2014)
- [LRS] A. Liashyk, N. Reshetikhin, I. Sechin: Quantum integrable systems on a classical integrable background. arXiv:2405.17865 [math-ph] (2024)
- [MP] J. A. Minahan, A. P. Polychronakos: Integrable systems for particles with internal degrees of freedom. Phys. Lett. B 302(2-3), 265–270 (1993)
- [OS] T. Oshima, H. Sekiguchi; Commuting families of differential operators invariant under the action of a Weyl group. J. Math. Sci. Univ. Tokyo 2(1), 1–75 (1995)
- [OP1] M. A. Olshanetsky, A. M. Perelomov: Classical integrable systems related to Lie algebras. Phys. Rep. 71 (5), 313–400 (1981)
- [OP2] M. A. Olshanetsky, A. M. Perelomov: Quantum integrable systems related to Lie algebras. Phys. Rep. 94(6), 313–404 (1983)
- [Pe] A. M. Perelomov: Completely integrable classical systems connected with semisimple Lie algebras. III. Lett. Math. Phys. 1(6), 531–534 (1977)
- [P1] A. P. Polychronakos: Exchange operator formalism for integrable systems of particles. Phys. Rev. Lett. 69(5), 703–705 (1992)
- [P2] A. P. Polychronakos: Lattice integrable systems of Haldane-Shastry type. Phys. Rev. Lett. 70(15), 2329–2331 (1993)
- [SZ] I. Sechin, A. Zotov; *R-matrix-valued Lax pairs and long-range spin chains*. Phys. Lett. B 781, 1–7 (2018)
- [SS] D. Serban, M. Staudacher: Planar N = 4 gauge theory and the Inozemtsev long range spin chain. JHEP **06** 001 (2004)
- [S] B. S. Shastry: Exact solution of an S = 1/2 Heisenberg antiferromagnetic chain with longranged interactions. Phys. Rev. Lett. **60**, 639–642 (1988)
- [TH] J. C. Talstra, F. D. M. Haldane: Integrals of motion of the Haldane–Shastry model. J. Phys. A: Math. Gen. 28, 2369–2377 (1995)

SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LEEDS LS2 9JT, UK *Email address*: o.chalykh@leeds.ac.uk