

July 4, 2024

**DERIVATIVE OF THE RIEMANN-HILBERT MAP**

VLADIMIR MARKOVIĆ AND OGNJEN TOŠIĆ

ABSTRACT. Given a pair  $(X, \nabla)$ , consisting of a closed Riemann surface  $X$  and a holomorphic connection  $\nabla$  on the trivial principal bundle  $X \times \mathrm{SL}_2(\mathbb{C}) \rightarrow X$ , the Riemann–Hilbert map sends  $(X, \nabla)$  to its monodromy representation. We compute the derivative of this map, and provide a simple description of the locus where it is injective, recovering in the process several previously obtained results.

## 1. INTRODUCTION

By  $\Sigma_g$  we denote a closed topological surface of genus  $g$ , and by  $\mathcal{T}_g$  the Teichmüller space of marked closed Riemann surfaces of genus  $g$ . Given  $X \in \mathcal{T}_g$ , we let  $\Omega^1(X)$  denote the space of Abelian differentials on  $X$ . By  $\mathfrak{sl}_2(\Omega^1(X))$  we denote the space of traceless  $2 \times 2$  matrices of holomorphic 1-forms on  $X$ .

Fix  $A \in \mathfrak{sl}_2(\Omega^1(X))$ , and consider the trivial holomorphic principal bundle  $\underline{\mathrm{SL}}_2(\mathbb{C}) := X \times \mathrm{SL}_2(\mathbb{C}) \rightarrow X$  equipped with the connection  $\nabla = d + A$ . This is a flat holomorphic connection over a marked Riemann surface  $X$  inducing the monodromy representation

$$\rho_A : \pi_1(\Sigma_g) \rightarrow \mathrm{SL}_2(\mathbb{C}).$$

The space of  $\mathfrak{sl}_2$ -systems over  $\Sigma_g$  is defined as

$$\mathrm{Syst}_g = \{(X, A) : X \in \mathcal{T}_g, A \in \mathfrak{sl}_2(\Omega^1(X)), \rho_A \text{ is irreducible}\} // \mathrm{SL}_2(\mathbb{C})$$

where we identify pairs  $(X, A)$  and  $(X, B)$  if  $A$  and  $B$  are conjugated by an element of  $\mathrm{SL}_2(\mathbb{C})$ . The Riemann–Hilbert map

$$\mathbf{M} : \mathrm{Syst}_g \rightarrow \mathrm{Rep}_g$$

is defined by  $\mathbf{M}(X, A) = \rho_A$ . Here

$$\mathrm{Rep}_g = \{\rho \in \mathrm{Hom}(\pi_1(\Sigma_g) \rightarrow \mathrm{SL}_2(\mathbb{C})) : \rho \text{ is irreducible}\} // \mathrm{SL}_2(\mathbb{C})$$

is the character variety of irreducible representations of the surface group  $\pi_1(\Sigma_g)$ .

It is well known that  $\mathbf{M}$  is a holomorphic map between connected complex manifolds of complex dimension  $6g - 6$  (see [1] and [2]). Our main result is Theorem 2 below is the explicit computation of the derivative  $D\mathbf{M}$ . But first, we state its main corollary characterising the pairs  $(X, A)$  where the

---

This work was supported by the *Simons Investigator Award* 409745 from the Simons Foundation

2010 *Mathematics Subject Classification*. Primary 34M03.

derivative of  $\mathbf{M}$  is injective (and hence bijective), answering a well known question (see [2], and also [4] for the broader picture).

Denote by  $\mathrm{QD}(X)$  the space of holomorphic quadratic differentials on  $X$ . Given  $\alpha, \beta, \gamma \in \Omega^1(X)$ , we define the subspace  $\mathrm{QD}(\alpha, \beta, \gamma) \leq \mathrm{QD}(X)$  by

$$\mathrm{QD}(\alpha, \beta, \gamma) = \{\alpha\varphi_1 + \beta\varphi_2 + \gamma\varphi_3 : \varphi_1, \varphi_2, \varphi_3 \in \Omega^1(X)\}.$$

**Theorem 1.** *Let  $(X, A) \in \mathrm{Syst}_g$ , and write*

$$A = \begin{pmatrix} \alpha & \beta \\ \gamma & -\alpha \end{pmatrix}$$

for  $\alpha, \beta, \gamma \in \Omega^1(X)$ . *The derivative  $D_{(X,A)}\mathbf{M}$  is injective if and only if*

$$(1) \quad \mathrm{QD}(\alpha, \beta, \gamma) = \mathrm{QD}(X).$$

**Remark 1.** This theorem has a number of applications which we prove in the last section. In particular, we show how it can be used to recover theorems by Biswas-Dumitrescu [1], Calsamiglia-Deroin-Heu-Loray [2]. In Proposition 3.1 in [1], Biswas-Dumitrescu give a sufficient condition on the pair  $(X, A)$  for the derivative  $D_{(X,A)}\mathbf{M}$  to be injective. This condition can be viewed as a cohomological version of the equality (1).

Before stating Theorem 2 we describe our parameterisation of the tangent spaces to  $\mathrm{Syst}_g$  and  $\mathrm{Rep}_g$ .

**1.1. Beltrami differentials and tamed matrix forms.** Let  $\mathrm{Belt}(X)$  denote the vector space of smooth Beltrami differentials on  $X$ . Two differentials  $\mu, \nu \in \mathrm{Belt}(X)$  are equivalent if

$$\int_X \mu\phi = \int_X \nu\phi$$

for every holomorphic quadratic differential  $\phi \in \mathrm{QD}(X)$ . This is a linear equivalence relation, that we denote  $\sim$ , and the quotient vector space  $\mathrm{Belt}(X)/\sim$  is naturally isomorphic to the tangent space of the Teichmüller space  $T_X\mathcal{T}_g$ .

Next, we define the notion of a tame matrix valued form.

**Definition 1.** Let  $A \in \mathfrak{sl}_2(\Omega^1(X))$ , and  $\mu \in \mathrm{Belt}(X)$ . We say that  $\dot{A}$  is  $(\mu, A)$ -tamed form if  $\dot{A}$  a closed  $\mathfrak{sl}_2(\mathbb{C})$ -valued 1-form whose  $(0, 1)$  part satisfies the equality  $\dot{A}^{0,1} = \mu A$ .

The following lemma is elementary and its proof is left to the reader.

**Lemma 1.** *Fix  $A \in \mathfrak{sl}_2(\Omega^1(X))$ . Let  $\mu, \nu \in \mathrm{Belt}(X)$ , and suppose  $\dot{A}$  and  $\dot{B}$  are  $(\mu, A)$  and  $(\nu, A)$  tamed respectively. If  $\mu \sim \nu$  then there exists a smooth function  $T : X \rightarrow \mathfrak{sl}_2(\mathbb{C})$  such that*

$$\dot{A} - \dot{B} - dT \in \mathfrak{sl}_2(\Omega^1(X)).$$

**1.2. Parametrisation of the tangent space to  $\text{Syst}_g$ .** Fix  $(X, A) \in \text{Syst}_g$ , and define the vector space

$$Z(X, A) = \{(\mu, \dot{A}) : \mu \in \text{Belt}(X), \dot{A} \text{ is } (\mu, A)\text{-tamed}\},$$

That is,  $Z(X, A)$  is a set of pairs  $(\mu, \dot{A})$  where  $\mu$  is a smooth Beltrami form on  $X$ , and  $\dot{A}$  a closed  $\mathfrak{sl}_2(\mathbb{C})$ -valued 1-form satisfying the equality  $\dot{A}^{0,1} = \mu A$ . We define the linear equivalence relation on  $Z(X, A)$  by letting  $(\mu, \dot{A}) \sim (\nu, \dot{B})$  if the following two conditions are satisfied

- (1)  $\mu \sim \nu$ ,
- (2)  $\dot{A}^{1,0} - \dot{B}^{1,0} - \partial T \in [A, \mathfrak{sl}_2(\mathbb{C})]$ , where  $T : X \rightarrow \mathfrak{sl}_2(\mathbb{C})$  is the function from Lemma 1.

**Remark 2.** Whenever  $\mu \sim \nu$ , we have  $\dot{A}^{1,0} - \dot{B}^{1,0} - \partial T \in \mathfrak{sl}_2(\Omega^1(X))$ . For the pairs  $(\mu, \dot{A})$  and  $(\nu, \dot{B})$  to be equivalent we require this element to live in  $[A, \mathfrak{sl}_2(\mathbb{C})] \leq \mathfrak{sl}_2(\Omega^1(X))$ .

It will be shown in Lemma 5 that the vector space  $Z(X, A)/\sim$  is naturally isomorphic to  $T_{(X,A)}\text{Syst}_g$ .

**1.3. Derivative of the monodromy map  $\mathbf{M}$ .** The tangent space  $T_\rho \text{Rep}_g$  is well-known to coincide with  $H^1(\text{Ad}_\rho)$ , where  $\text{Ad}_\rho$  denotes the  $\pi_1(\Sigma_g)$ -module with underlying vector space  $\mathfrak{sl}_2(\mathbb{C})$  and the action  $(\gamma, T) \rightarrow \rho(\gamma)T\rho(\gamma)^{-1}$ . Since  $\text{Ad}_\rho$  is precisely the monodromy of the flat connection

$$d_A := d + \text{ad}(A)$$

on the bundle  $\underline{\mathfrak{sl}_2(\mathbb{C})} := X \times \mathfrak{sl}_2(\mathbb{C}) \rightarrow X$ , there is a de Rham isomorphism

$$(2) \quad \iota : H^1(X, \mathcal{E}) \rightarrow H^1(\text{Ad}_\rho).$$

Here  $\mathcal{E}$  denotes the sheaf of  $d_A$ -flat sections of the bundle  $\underline{\mathfrak{sl}_2(\mathbb{C})}$ . We are now ready to state our main result.

**Theorem 2.** *Let  $(X, A) \in \text{Syst}_g$  and  $(\mu, \dot{A}) \in Z(X, A)$ . Then  $\dot{A}$  is a  $d_A$ -closed  $\mathfrak{sl}_2(\mathbb{C})$ -valued 1-form, and hence defines a cohomology class  $\chi \in H^1(X, \mathcal{E})$ . Moreover,  $\text{DM}([\mu, \dot{A}]) = -\iota(\chi)$ , where  $[\mu, \dot{A}] \in Z(X, A)/\sim$  denotes the corresponding tangent vector under the identification  $T_{(X,A)}\text{Syst}_g \approx Z(X, A)/\sim$ .*

**1.4. Organisation.** Theorem 1 is proved in Section 2. In Section 3 we describe the tangent space to  $\text{Syst}_g$  and prove that  $T_{(X,A)}\text{Syst}_g$  is isomorphic to  $Z(X, A)/\sim$ . In Section 4 we recall the description of the tangent space of  $\text{Rep}_g$  and construct the de Rham isomorphism (2). In Section 5 we prove Theorem 2. Applications of Theorem 1 are derived in Section 6.

2. THE KERNEL OF  $DM$ 

Fix  $(X, A) \in \text{Syst}_g$ , and consider a tangent vector  $(\mu, \dot{A}) \in Z(X, A)$ . The following three lemmas easily follow from Theorem 2, and the isomorphism  $T_{(X,A)}\text{Syst}_g \approx Z(X, A)/\sim$ . We then use these lemmas to prove Theorem 1.

**Lemma 2.** *Let  $(\mu, \dot{A}) \in Z(X, A)$ . The derivative  $DM(\mu, \dot{A})$  vanishes if and only if there exists a smooth function  $\dot{F} : X \rightarrow \mathfrak{sl}_2(\mathbb{C})$  such that*

$$(3) \quad \mu A = \bar{\partial}\dot{F} \quad \text{and} \quad \dot{A} = \partial\dot{F} + [A, \dot{F}] + \mu A.$$

*Proof.* By Theorem 2 the derivative  $DM(\mu, \dot{A})$  vanishes if and only if  $\chi = [\dot{A}] \in H^1(X, \mathcal{E})$  vanishes, that is, if  $\dot{A}$  is  $d_A$ -exact. This means that  $\dot{A} = d_A\dot{F} = d\dot{F} + [A, \dot{F}]$ , for some smooth function  $\dot{F} : X \rightarrow \mathfrak{sl}_2(\mathbb{C})$ . Splitting this equality into  $(0, 1)$  and  $(1, 0)$  parts, and using that  $\dot{A}^{0,1} = \mu A$ , we derive (3). □

**Lemma 3.** *Suppose  $\mu$  is a Beltrami differential on  $X$ . Then there exists  $\dot{A}$  such that  $DM(\mu, \dot{A}) = 0$  if and only if the 1-forms  $\mu\alpha, \mu\beta, \mu\gamma$  are all  $\bar{\partial}$ -exact, where*

$$A = \begin{pmatrix} \alpha & \beta \\ \gamma & -\alpha \end{pmatrix}$$

*Proof.* If such  $\dot{A}$  exists then by Lemma 2 there exists  $\dot{F} : X \rightarrow \mathfrak{sl}_2(\mathbb{C})$  satisfying the system (3). In particular, the equality  $\mu A = \bar{\partial}\dot{F}$  holds implying that the 1-forms  $\mu\alpha, \mu\beta, \mu\gamma$  are all  $\bar{\partial}$ -exact.

Suppose now that  $\mu\alpha, \mu\beta, \mu\gamma$  are all  $\bar{\partial}$ -exact. Then we can construct  $\dot{F} : X \rightarrow \mathfrak{sl}_2(\mathbb{C})$  such that  $\mu A = \bar{\partial}\dot{F}$ . Set  $\dot{A} = \mu A + \partial\dot{F} + [A, \dot{F}]$ . One can now verify that  $(\mu, \dot{A})$  solves the system (3). □

**Lemma 4.** *Suppose  $\mu$  is zero in  $T_X\mathcal{T}_g$ . Then  $DM(\mu, \dot{A}) = 0$  if and only if the pair  $(\mu, \dot{A})$  represents the zero vector in  $T_{(X,A)}\text{Syst}_g$ .*

*Proof.* Suppose that  $DM(\mu, \dot{A}) = 0$ . We need to prove that  $(\mu, \dot{A})$  represents the zero vector in  $T_{(X,A)}\text{Syst}_g$ . Since  $\mu$  is zero in  $T_X\mathcal{T}_g$ , we may assume  $\mu = 0$ . Let  $\dot{F} : X \rightarrow \mathfrak{sl}_2(\mathbb{C})$  be the function from Lemma 2. Then  $\bar{\partial}\dot{F} = 0$ , and hence  $\dot{F} : X \rightarrow \mathfrak{sl}_2(\mathbb{C})$  is holomorphic. Therefore  $\dot{F}$  is constant, and by the second equality in (3) we obtain  $\dot{A} = \dot{A}^{1,0} = [A, \dot{F}]$ . But then  $\dot{A} \in [A, \mathfrak{sl}_2(\mathbb{C})]$ , and so  $(0, \dot{A})$  vanishes in  $T_{(X,A)}\text{Syst}_g$ . □

**2.1. Proof of Theorem 1.** We are ready to complete the proof of Theorem 1. Let  $\mu$  denote a Beltrami differential on  $X$ . Then by Lemma 3 and Lemma 4 there exists  $\dot{A}$  such that  $(\mu, \dot{A})$  is a non-zero vector with  $DM(\mu, \dot{A}) = 0$  if and only if  $\mu$  is not equal to zero in  $T_X\mathcal{T}_g$ , and the 1-forms  $\mu\alpha, \mu\beta, \mu\gamma$  are all  $\bar{\partial}$ -exact. By Serre duality, the last condition is equivalent to

$$\int_X \mu\phi = 0$$

for every  $\phi \in \alpha\Omega^1(X) + \beta\Omega^1(X) + \gamma\Omega^1(X) \leq \text{QD}(X)$ . Thus,  $DM$  is injective if and only if the equality (1) holds.

### 3. THE SPACE OF $\mathfrak{sl}_2$ -SYSTEMS

Recall that

$$\text{Syst}_g = \{(X, A) : X \in \mathcal{T}_g, A \in \mathfrak{sl}_2(\Omega^1(X)), \rho_A \text{ is irreducible}\} // \text{SL}_2(\mathbb{C}).$$

Note that the quotient here is by the conjugation action of  $\text{SL}_2(\mathbb{C})$ , in the sense of geometric invariant theory. Since we have restricted to the irreducible locus, this coincides with the quotient in the sense of topology. It is standard that  $\text{Syst}_g$  is a smooth complex manifold.

Our aim in this section is to describe the tangent space  $T_{(X,A)}\text{Syst}_g = Z(X, A) / \sim$ , as described in Section 1. Given a smooth Beltrami form  $\mu \in \text{Belt}(X)$  with  $\|\mu\|_{L^\infty(X)} < 1$ , denote by  $f^\mu : X \rightarrow X^\mu$  the smooth map to a Riemann surface  $X^\mu$  such that

$$\bar{\partial}f^\mu = \mu\partial f^\mu.$$

**Lemma 5.** *Let  $(X, A) \in \text{Syst}_g$  and  $(\mu, \dot{A}) \in Z(X, A)$ . Then there exists a path of pairs  $(X^{t\mu}, A_t)$  based at  $(X, A)$ , such that*

$$(4) \quad \left. \frac{d}{dt} \right|_{t=0} \left( (f^{t\mu})^* A_t \right) = \dot{A}.$$

Moreover, the map  $\tau : Z(X, A) \rightarrow T_{(X,A)}\text{Syst}_g$ , defined by taking the tangent vector to this path at  $t = 0$ , is surjective and has kernel  $B(X, A)$  consisting of pairs  $(\mu, \dot{A}) \in Z(X, A)$  such that

- $\mu$  vanishes in  $T_X\mathcal{T}_g$ , and
- $\dot{A} - dT \in [A, \mathfrak{sl}_2(\mathbb{C})]$  for some function  $T : X \rightarrow \mathfrak{sl}_2(\mathbb{C})$ .

We split the proof of Lemma 5 into two parts. We construct the path with property (4) in §3.1, and then we describe the kernel of  $\tau$  in §3.2.

**3.1. Constructing the path.** The path we construct will in fact be in

$$\{(X, A) : X \in \mathcal{T}_g, A \in \mathfrak{sl}_2(\Omega^1(X))\}$$

rather than  $\text{Syst}_g$ . Note that this space is isomorphic to the total space of  $\mathcal{H}_g \otimes \mathfrak{sl}_2(\mathbb{C})$ , where  $\mathcal{H}_g \rightarrow \mathcal{T}_g$  is the Hodge bundle of Abelian differentials over a varying Riemann surface, i.e. the fibre of  $\mathcal{H}_g$  over  $X \in \mathcal{T}_g$  is  $\Omega^1(X)$ .

Let  $A_t$  be a path of  $\mathfrak{sl}_2(\mathbb{C})$  valued holomorphic 1-forms on  $X^{t\mu}$ , chosen so that  $[\text{Re}(A_t)] \in H^1(\Sigma_g, \mathfrak{sl}_2(\mathbb{R}))$  depends smoothly on  $t$ , and has  $\left. \frac{d}{dt} \right|_{t=0} [\text{Re}(A_t)] = [\text{Re}(\dot{A})]$ . By the smoothness of the Gauss–Manin connection on  $\mathcal{H}_g$ , it follows that  $(f^{t\mu})^* A_t$  depend smoothly on  $t$ , and hence we can define

$$B = \left. \frac{d}{dt} \right|_{t=0} \left( (f^{t\mu})^* A_t \right).$$

Since the 1-forms  $A_t$  are holomorphic, they are also closed, and hence so is  $B$ . Moreover,  $\operatorname{Re}(\dot{A} - B)$  is exact by construction.

**Claim 1.** We have  $B^{0,1} = \mu A$ .

*Proof.* We work in a local coordinate  $z$  on  $X$ , and abuse notation slightly to denote the Beltrami form by  $\mu \frac{d\bar{z}}{dz}$  in this coordinate. Then  $\tilde{A}_t := (f^{t\mu})^* A_t$  is proportional to  $f_z^{t\mu} (dz + t\mu d\bar{z})$ . In particular,

$$\tilde{A}_t^{0,1} = t\mu \tilde{A}_t^{1,0}.$$

Differentiating at  $t = 0$ , we see that  $B^{0,1} = \mu A$ , as desired.  $\square$

In particular, it follows that  $B - \dot{A}$  is a closed form of type  $(1,0)$ , and is therefore holomorphic. Since  $B - \dot{A}$  has exact real part, it follows by Hodge theory that  $B - \dot{A} = 0$ . We have thus constructed a path with (4).

**3.2. Surjectivity and the kernel of  $\tau$ .** We first show the surjectivity of  $\tau$ . Given any tangent vector  $V \in T_{(X,A)}(\mathcal{H}_g \otimes \mathfrak{sl}_2(\mathbb{C}))$ , let  $(X^{t\mu}, A_t)$  be a path in  $\mathcal{H}_g \otimes \mathfrak{sl}_2(\mathbb{C})$  tangent to  $V$  at  $(X, A)$ . Note that each  $A_t$  is closed, so is  $(f^{t\mu})^* A_t$ . Thus  $\dot{A} := \left. \frac{d}{dt} \right|_{t=0} (f^{t\mu})^* A_t$  is closed, and by Claim 1 has  $(0,1)$ -part equal to  $\mu A$ . Thus  $\dot{A}$  is  $(\mu, A)$ -tamed and hence  $V = \tau(\mu, \dot{A})$ . Thus  $\tau$  is surjective.

We now show that  $\ker \tau = B(X, A)$ . Since the paths constructed in §3.1 lie in  $\mathcal{H}_g \otimes \mathfrak{sl}_2(\mathbb{C})$ ,  $\tau$  factors as

$$Z(X, A) \xrightarrow{\tilde{\tau}} T_{(X,A)}(\mathcal{H}_g \otimes \mathfrak{sl}_2(\mathbb{C})) \rightarrow T_{(X,A)}\operatorname{Syst}_g.$$

Since  $\rho_A$  is irreducible, the point  $(X, A)$  is stable for the  $\operatorname{SL}_2(\mathbb{C})$ -action on  $\mathcal{H}_g \otimes \mathfrak{sl}_2(\mathbb{C})$ . Thus  $\ker \tau = \tilde{\tau}^{-1}((0, [A, \mathfrak{sl}_2(\mathbb{C})])) = (0, [A, \mathfrak{sl}_2(\mathbb{C})]) + \ker \tilde{\tau}$  by standard geometric invariant theory. In the rest of this subsection, we show that

$$\ker \tilde{\tau} = \{(\mu, \dot{A}) \in Z(X, A) : \mu \text{ vanishes in } T_X \mathcal{T}_g \text{ and } \dot{A} \text{ is exact}\},$$

which implies  $B(X, A) = \ker \tau$  immediately.

Suppose first that  $(\mu, \dot{A}) \in \ker \tilde{\tau}$ . Then  $\mu$  corresponds to the zero tangent vector in  $T_X \mathcal{T}_g$ . By the construction of the path in §3.1, it follows that  $\operatorname{Re}(\dot{A})$  is exact. Since the period matrix of  $X^{t\mu} = X$  remains constant up to terms of order  $o(t)$  (for  $\mu$  is zero in the  $T\mathcal{T}_g$ ), we see that  $\operatorname{Im}(\dot{A})$  is exact as well.

Conversely, if  $\mu$  vanishes in  $T_X \mathcal{T}_g$  and  $\dot{A} = dT$ , then the cohomology class  $[\dot{A}] \in H^1(\Sigma_g, \mathfrak{sl}_2(\mathbb{C}))$  vanishes. Thus the path in §3.1 represents a path of holomorphic 1-forms on a fixed Riemann surface, with the same cohomology. Thus the path is constant, and  $(\mu, \dot{A}) \in \ker \tilde{\tau}$ .

#### 4. THE REPRESENTATION VARIETY AND DE RHAM ISOMORPHISM

In this section, we explain the tangent space to the representation variety  $\operatorname{Rep}_g$  and the de Rham isomorphism for local systems.

**4.1. Tangent space to  $\text{Rep}_g$ .** The description of  $T\text{Rep}_g$  is much more standard than  $T\text{Syst}_g$ , and can be found for instance in [6]. We summarise the results here for the sake of completeness.

Given an irreducible representation  $\rho : \pi_1(\Sigma_g) \rightarrow \text{SL}_2(\mathbb{C})$ , the tangent space  $T_\rho\text{Rep}_g$  can be described as  $H^1(\text{Ad}_\rho) := Z(\text{Ad}_\rho)/B(\text{Ad}_\rho)$ , where

$$Z(\text{Ad}_\rho) = \{\eta : \pi_1(\Sigma_g) \rightarrow \mathfrak{sl}_2(\mathbb{C}) \text{ with } \eta(\gamma_1\gamma_2) = \eta(\gamma_1) + \rho(\gamma_1)\eta(\gamma_2)\rho(\gamma_1)^{-1}\},$$

and

$$B(\text{Ad}_\rho) = \{\rho T \rho^{-1} - T \text{ for } T \in \mathfrak{sl}_2(\mathbb{C})\}.$$

Given an element  $\eta \in Z(\text{Ad}_\rho)$ , there exists a smooth path  $\rho_t : \pi_1(\Sigma_g) \rightarrow \text{SL}_2(\mathbb{C})$  such that

$$\left. \frac{d}{dt} \right|_{t=0} \rho_t(\gamma) = \eta(\gamma)\rho(\gamma),$$

and this path is tangent to  $\eta + B(\text{Ad}_\rho) \in H^1(\text{Ad}_\rho)$ .

**4.2. De Rham isomorphism for local systems.** There is a standard correspondence between flat vector bundles  $E$  of rank  $n$  over a manifold  $M$ , and representations  $\rho : \pi_1(M) \rightarrow \text{GL}_n(\mathbb{C})$  of its fundamental group. Taking the derivative of this correspondence gives an isomorphism  $H^1(M, \mathcal{E}) \cong H^1(\pi_1(M), \text{Ad}_\rho)$ , where  $\mathcal{E}$  is the sheaf of parallel sections of  $E$ . In this section, we describe this isomorphism explicitly. We omit the proofs as they are standard application of homological algebra. A much more general result can be found in Proposition 6.3 in [7].

We first describe how  $H^1(M, \mathcal{E})$  and  $H^1(\text{Ad}_\rho)$  can be understood explicitly. Consider the de Rham resolution of  $\mathcal{E}$

$$\mathcal{E} \rightarrow E \xrightarrow{d^\nabla} E \otimes T^*M \xrightarrow{d^\nabla} E \otimes \bigwedge^2 T^*M.$$

This is a soft resolution, and its sections can hence be used to compute the cohomology

$$H^1(\Sigma_g, \mathcal{E}) = \frac{\{A \in C^\infty(E \otimes T^*\Sigma_g) : d^\nabla A = 0\}}{\{d^\nabla B : B \in C^\infty(E)\}}.$$

Also recall that  $H^1(\text{Ad}_\rho) = Z(\text{Ad}_\rho)/B(\text{Ad}_\rho)$ , where

$$Z(\text{Ad}_\rho) = \{\eta : \pi_1(\Sigma_g) \rightarrow \mathbb{C}^n \text{ such that } \eta(\gamma_1\gamma_2) = \eta(\gamma_1) + \rho(\gamma_1)\eta(\gamma_2)\rho(\gamma_1)^{-1}\},$$

and

$$B(\text{Ad}_\rho) = \{\rho(\gamma)\eta\rho(\gamma)^{-1} - \eta : \eta \in \mathbb{C}^n\}.$$

We now describe the de Rham isomorphism  $\iota : H^1(M, \mathcal{E}) \rightarrow H^1(\text{Ad}_\rho)$ . Let  $p : \tilde{M} \rightarrow M$  be the universal cover, and  $\tilde{E} = p^*E$  be the pullback of  $E$ . Note that the deck group action of  $\pi_1(M)$  on  $\tilde{M}$  naturally extends to an action on  $\tilde{E}$ . Fix an arbitrary basepoint  $x \in \tilde{M}$ .

Given a  $d^\nabla$ -closed  $E$ -valued 1-form  $A$ , the class  $\iota([A])$  can be computed as follows. Since  $\tilde{E}$  is the trivial flat bundle, it admits a section  $T$  such that

$dT = p^*A$ . It is easily seen that  $\gamma T - T$  is parallel for any  $\gamma \in \pi_1(M)$ , and hence

$$\gamma \longrightarrow (\gamma T - T)(x)$$

defines a cycle in  $Z(\text{Ad}_\rho)$ . The image of this cycle in  $H^1(\text{Ad}_\rho)$  is  $\iota([\chi])$ .

## 5. PROOF OF THEOREM 2

**5.1. Trivializing the flat  $\text{SL}_2(\mathbb{C})$ -bundle.** We now describe the setup in which we will prove Theorem 2.

Let  $X$  be a Riemann surface and  $A \in \mathfrak{sl}_2(\Omega^1(X))$ . Let  $\rho : \pi_1(X) \rightarrow \text{SL}_2(\mathbb{C})$  be the monodromy of  $d + A$ . The following lemma shows how to trivialize  $(\text{SL}_2(\mathbb{C}), d + A)$ . Denote by  $\tilde{X}$  the universal cover of  $X$ . This lemma is entirely contained in the literature (see e.g. the introduction to [2]), but we include a proof for completeness.

**Lemma 6.** *There exists a map  $F : \tilde{X} \rightarrow \text{SL}_2(\mathbb{C})$  such that*

$$dF + AF = 0 \text{ and } F(\gamma x) = F(x)\rho(\gamma)^{-1}.$$

*Then the map  $\Phi : \tilde{X} \times \text{SL}_2(\mathbb{C}) \rightarrow \tilde{X} \times \text{SL}_2(\mathbb{C})$  defined by  $\Phi(x, T) = (x, F(x)T)$  is an isomorphism of principal bundles with the following two properties*

- (1)  $\Phi$  conjugates the  $\pi_1(X)$ -action  $\gamma \cdot (x, T) = (\gamma x, \rho(\gamma)T)$  to the action  $\gamma \cdot (x, T) = (\gamma x, T)$ , and
- (2)  $\Phi^*(d + A) = d$ .

*Proof.* Fix an arbitrary basepoint  $\tilde{x}_0 \in \tilde{X}$ , and let  $F_0 \in \text{SL}_2(\mathbb{C})$  be such that  $\rho$  is the monodromy of  $d + A$  with basepoint  $\tilde{x}_0$  relative to the basis that consists of the columns of  $F_0$ . Define  $F(x)$  as follows: for  $i = 1, 2$ , its  $i$ -th column is the  $d_A$ -parallel transport of the  $i$ -th column of  $F_0$  along an arbitrary path from  $\tilde{x}_0$  to  $x$ . It is immediate that  $dF + AF = 0$ , since  $F$  is obtained through parallel transport. Since the monodromy of  $d + A$  is precisely  $\rho$ , it follows that  $F(x) = F(\gamma x)\rho(\gamma)$ .

Moreover, if we let  $\sigma : \tilde{X} \rightarrow \text{SL}_2(\mathbb{C})$  be a section of the bundle  $(\text{SL}_2(\mathbb{C}), d_A)$ , then

$$\begin{aligned} \nabla(F^{-1}\sigma) &= \sigma^{-1}Fd(F^{-1}\sigma) \\ &= \sigma^{-1}F\left(-F^{-1}(dF)F^{-1}\sigma + F^{-1}d\sigma\right) \\ &= \sigma^{-1}d\sigma + \sigma^{-1}A\sigma = d_A\sigma, \end{aligned}$$

and the result is shown.  $\square$

**5.2. Computing the derivative.** We now let  $(X^{t\mu}, A_t)$  be the path at  $(X, A) \in \text{Syst}_g$  tangent to the vector  $(\mu, \dot{A})$  with  $\dot{A}^{0,1} = \mu A$ , as in Lemma 5. Let  $F_t : \tilde{X}^{t\mu} \rightarrow \text{SL}_2(\mathbb{C})$  be as in Lemma 6, and suppose that the monodromy of  $d + A_t$  is  $\rho_t : \pi_1(X^{t\mu}) \rightarrow \text{SL}_2(\mathbb{C})$ .



We use  $f^{t\mu}$  to transport  $A_t, F_t$  and  $\rho_t$  to  $X$ . Recall that  $\frac{d}{dt}\Big|_{t=0} A_t = \dot{A}$ . We have

$$dF_t + A_t F_t = 0 \text{ and } F_t(\gamma x) = F_t(x) \cdot \rho_t(\gamma)^{-1}.$$

Let  $F_t = (\text{id} + t\dot{F} + o(t)) \cdot F$  for some map  $\dot{F} : \tilde{X} \rightarrow \mathfrak{sl}_2(\mathbb{C})$ . Then

$$d_A \dot{F} + \dot{A} = d\dot{F} + \dot{A} + [A, \dot{F}] = 0.$$

We thus have  $d_A \dot{F} = -\dot{A}$ . Similarly, if we set  $\rho_t(\gamma) = (\text{id} + t\dot{\rho}(\gamma) + o(t)) \cdot \rho(\gamma)$ , we have

$$\dot{\rho}(\gamma) = F(\gamma x)^{-1} \left( \dot{F}(x) - \dot{F}(\gamma x) \right) F(\gamma x).$$

Set  $B = F^{-1} \dot{F} F$ . Then  $\dot{\rho}(\gamma) = \rho(\gamma) B(x) \rho(\gamma)^{-1} - B(\gamma x)$ . It is easy to see that  $\rho(\gamma) B(x) \rho(\gamma)^{-1} - B(\gamma x)$  does not depend on  $x \in \tilde{X}$ , so we have

$$\dot{\rho}(\gamma) = \rho(\gamma) B(\gamma^{-1} x) \rho(\gamma)^{-1} - B(x).$$

From the description of the de Rham isomorphism  $H^1(X, \mathcal{E}) \rightarrow H^1(\text{Ad}_\rho)$  in §4.2, we see that  $[\dot{\rho}] = -\iota[\dot{A}]$ .

## 6. APPLICATIONS

Recall that Theorem 1 states that the derivative  $DM$  is injective on the tangent space  $T_{(X,A)} \text{Syst}_g$  if and only if  $\text{QD}(\alpha, \beta, \gamma) = \text{QD}(X)$ , where

$$A = \begin{pmatrix} \alpha & \beta \\ \gamma & -\alpha. \end{pmatrix}$$

In this section we derive several applications of this theorem.

**6.1. Preliminaries.** Let  $X$  be a Riemann surface of genus at least two, and let  $\xi_i \in \Omega^1(X)$ ,  $i = 1, \dots, k$ . Let  $\text{QD}(\xi_1, \dots, \xi_k) \leq \text{QD}(X)$  be the vector subspace generated by the products  $\xi_i \varphi$ ,  $i = 1, \dots, k$ , and  $\varphi \in \Omega^1(X)$ . The following was proved by Rauch (see statement C in [10]).

**Theorem 3.** *Let  $X$  be a hyperelliptic Riemann surface of genus at least two, and suppose  $\xi_1, \dots, \xi_g$  is a basis of  $\Omega^1(X)$ . Then  $\dim(\text{QD}(\xi_1, \dots, \xi_g)) = 2g - 1$ .*

The next theorem is a classical result.

**Theorem 4.** *Suppose  $g \geq 3$ . There exists a dense open subset  $U \subset \mathcal{H}_g^{\oplus 3}$  such that the equality  $\text{QD}(\alpha, \beta, \gamma) = \text{QD}(X)$  holds for each  $(\alpha, \beta, \gamma) \in U$ .*

**Remark 3.** Let  $p : \mathcal{H}_g^{\oplus 3} \rightarrow \mathcal{T}_g$  be the natural projection. According to the result of [5], the set  $U$  from Theorem 4 can in fact be taken such that  $U \cap p^{-1}(X)$  is dense in  $p^{-1}(X)$  for any non-hyperelliptic  $X \in \mathcal{T}_g$ . The same result can be recovered from the well known Max Noether theorem (see III.11.20 in [3]).

*Proof.* Let  $X$  be a non-hyperelliptic Riemann surface. By Theorem 1.1 in [5], the set of triples  $(\alpha, \beta, \gamma) \in (\Omega^1(X))^3$  such that  $\text{QD}(\alpha, \beta, \gamma) = \text{QD}(X)$  is an open and dense subset of  $(\Omega^1(X))^3$ . Taking the union of these dense open subsets over all non-hyperelliptic Riemann surface in  $\mathcal{T}_g$  yields the required open and dense subset of  $\mathcal{H}_g^{\oplus 3}$ .  $\square$

**6.2. Two corollaries.** The following is a corollary of Theorem 1 and Theorem 3.

**Theorem 5.** *Suppose  $X \in \mathcal{T}_g$  is hyperelliptic and let  $(X, A) \in \text{Syst}_g$ . Then  $DM$  is injective on the tangent space  $T_{(X,A)}\text{Syst}_g$  if and only if  $g = 2$ .*

**Remark 4.** That  $DM$  is injective on the tangent space  $T_{(X,A)}\text{Syst}_g$  if  $g = 2$  is a well known theorem by Calsamiglia-Derooin-Heu-Loray [2].

*Proof.* Assume at first that  $g > 2$ . Then by Theorem 3 the following inequality holds

$$\dim(\text{QD}(\alpha, \beta, \gamma)) \leq 2g - 1 < 3g - 3 = \dim(\text{QD}(X)).$$

Thus, the equality (1) can never hold on such  $X$  regardless of the choice of Abelian differentials  $\alpha, \beta, \gamma$ .

We now consider the case  $g = 2$ . Since  $(X, A) \in \text{Syst}_g$ , we know that the monodromy  $\rho_A : \pi_1(\Sigma_g) \rightarrow \text{SL}_2(\mathbb{C})$  is irreducible. Then the three Abelian differentials  $\alpha, \beta$ , and  $\gamma$ , span  $\Omega^1(X)$ . This was observed by Biswas-Dumitrescu (see the proof of Proposition 4.1 in [1]). Since  $\dim(\text{QD}(X)) = 2$ , without loss of generality we may assume that  $\alpha$  and  $\beta$  span  $\Omega^1(X)$ . But then by Theorem 3 we know that the three quadratic differentials  $\alpha^2, \alpha\beta, \beta^2$  represent a basis for  $\text{QD}(X)$ . By Theorem 1 the derivative  $DM$  is injective on  $T_{(X,A)}\text{Syst}_g$ .  $\square$

In the remainder of this section we let

$$I : \mathcal{H}_g^{\oplus 3} \rightarrow \mathcal{H}_g \otimes \mathfrak{sl}_2(\mathbb{C})$$

denote the identification

$$(\alpha, \beta, \gamma) \mapsto A = \begin{pmatrix} \alpha & \beta \\ \gamma & -\alpha \end{pmatrix}.$$

**Theorem 6.** *The derivative  $DM$  is locally injective on an open and dense subset of  $\text{Syst}_g$ .*

**Remark 5.** This is a recent theorem by Biswas-Dumitrescu [1].

*Proof.* Let  $U \subset \mathcal{H}_g^{\oplus 3}$  be the open dense set from Theorem 4. Moreover, let  $V$  be the open dense subset of  $\mathcal{H}_g \otimes \mathfrak{sl}_2(\mathbb{C})$  that corresponds to irreducible flat connections. Then  $I(U) \cap V$  is a dense open subset of  $\mathcal{H}_g \otimes \mathfrak{sl}_2(\mathbb{C})$ . Since the quotient map  $q : V \rightarrow V // \text{SL}_2(\mathbb{C})$  is open, it follows that  $q(I(U) \cap V)$  is the desired dense open set by Theorem 1.  $\square$

## REFERENCES

- [1] I. Biswas and S. Dumitrescu, *The monodromy map from differential systems to character variety is generically immersive*. Publ. Res. Inst. Math. Sci. 59 (2023), 821-842.
- [2] G. Calsamiglia, B. Deroin, V. Heu and F. Loray, *The Riemann-Hilbert mapping for  $\mathfrak{sl}_2$  systems over genus two curves*. Bull. Soc. Math. France 147 (2019), 159-195.
- [3] H. Farkas, I. Kra, *Riemann surfaces*. Graduate Texts in Mathematics, 71. Springer-Verlag, New York-Berlin (1980).
- [4] E. Ghys. *Déformations des structures complexes sur les espaces homogènes de  $SL(2, \mathbb{C})$* . J. Reine Angew. Math. 468 (1995) 113-138.
- [5] D. Gieseker, *A lattice version of the KP equation*. Acta Math. 168 (1992), 219-248.
- [6] W. Goldman, *The symplectic nature of fundamental groups of surfaces*. Advances in Mathematics 54 (1984), 200-225.
- [7] W. Goldman, J. Millson, *The deformation theory of representations of fundamental groups of compact Kähler manifolds*. Publications Mathématiques de l'IHÉS 67 (1988), 43-96.
- [8] V. Marković, *Unramified correspondences and virtual properties of mapping class groups*. Bull. Lond. Math. Soc. 54, no. 6 (2022) 2324-2337.
- [9] V. Marković, O. Tošić, *The second variation of the Hodge norm and higher Prym representations*. Journal of Topology 17 (2024).
- [10] E. Rauch, *On the transcendental moduli of algebraic Riemann surfaces*. Proc. Nat. Acad. Sci. U.S.A. 41 (1955), 42-49.

MATHEMATICAL INSTITUTE  
UNIVERSITY OF OXFORD  
UNITED KINGDOM

*Email address:* ognjen.tosic@gmail.com

ALL SOULS COLLEGE  
UNIVERSITY OF OXFORD  
UNITED KINGDOM

*Email address:* markovic@maths.ox.ac.uk