

# Fair Division of Indivisible Chores via Earning Restricted Equilibria\*

Jugal Garg<sup>†</sup>  
jugal@illinois.edu

Aniket Murhekar<sup>‡</sup>  
aniket2@illinois.edu

John Qin<sup>§</sup>  
johnqin2@illinois.edu

## Abstract

We study fair division of  $m$  indivisible chores among  $n$  agents with additive preferences. We consider the desirable fairness notions of envy-freeness up to any chore (EFX) and envy-freeness up to  $k$  chores (EF $k$ ), alongside the efficiency notion of Pareto optimality (PO). We present the first constant approximations of these notions, showing the existence of:

- 4-EFX allocations, which improves the best-known factor of  $O(n^2)$ -EFX.
- 2-EF2 and PO allocations, which improves the best-known factor of EF $m$  and PO. In particular, we show the existence of an allocation that is PO and for every agent, either EF2 or 2-EF1.
- 3-EFX and PO allocations for the special case of bivalued instances, which improves the best-known factor of  $O(n)$ -EFX without any efficiency guarantees.

A notable contribution of our work is the introduction of the novel concept of *earning-restricted (ER) competitive equilibrium* for fractional allocations, which limits agents' earnings from each chore. Technically, our work addresses two main challenges: proving the existence of an ER equilibrium and designing algorithms that leverage ER equilibria to achieve the above results. To tackle the first challenge, we formulate a linear complementarity problem (LCP) formulation that captures all ER equilibria and show that the classic complementary pivot algorithm on the LCP must terminate at an ER equilibrium. For the second challenge, we carefully set the earning limits and use properties of ER equilibria to design sophisticated procedures that involve swapping and merging bundles to meet the desired fairness and efficiency criteria. We expect that the concept of ER equilibrium will be instrumental in deriving further results on related problems.

## 1 Introduction

Allocation problems frequently arise in various contexts, such as task allocation, partnership dissolution, or the division of inheritance. The question of fairly allocating indivisible items has been studied across multiple disciplines, including computer science, economics, social choice theory, and multi-agent systems. Although earlier work primarily focused on *goods* (items that provide value to agents), there has been a growing interest in the fair division of *chores* (items that impose a cost to agents) in recent years. In this paper, we study the fundamental problem of *fairly* allocating a set of  $m$  *indivisible chores* among  $n$  agents with additive disutility functions.

\*Work supported by NSF Grants CCF-1942321 and CCF-2334461.

<sup>†</sup>University of Illinois at Urbana-Champaign, USA

<sup>‡</sup>University of Illinois at Urbana-Champaign, USA

<sup>§</sup>University of Illinois at Urbana-Champaign, USA

Among various choices, *envy-freeness* (EF) [22] is the most natural fairness concept, where every agent (weakly-)prefers the items allocated to her over those assigned to other agents. However, with indivisible items, EF allocations need not exist, e.g., consider allocating one task between two agents, suggesting relaxations of EF to suit the discrete case. The two most popular and compelling relaxations are envy-freeness up to any item (EFX) and envy-freeness up to one item (EF1). For goods, an allocation is said to be EFX (resp. EF1) if no agent  $i$  is envious of another agent  $h$  *after* the counter-factual removal of *any* (resp. some) good assigned to  $h$ . Conversely, for chores, which impose a cost on the agents receiving them, an allocation of chores is said to be EFX (resp. EF1) if no agent  $i$  is envious of another agent  $h$  *after* the counter-factual removal of *any* (resp. some) chore assigned to  $i$ . Thus, EFX is the closest discrete analog to EF, with the relationship being  $\text{EF} \implies \text{EFX} \implies \text{EF1}$ . In contrast to EF, EF1 allocations always exist and can be computed in polynomial time for both goods [32] and chores [13]. However, the existence of EFX allocations remains a major open problem in discrete fair division for more than three agents in the case of goods and for more than two agents in the case of chores.

**EFX for Chores.** The existence of EFX allocations is regarded as the ‘*biggest open problem in discrete fair division*’ [37]. For chores, the existence of EFX allocations is known only under certain additional assumptions about the instance. For example, they are known to exist when the number of chores is at most twice the number of agents [29], or when there are two types of chores [8] (see Section 1.3 for an expanded discussion). Alternatively, one may inquire about the existence of approximate-EFX allocations for all instances. In an  $\alpha$ -EFX allocation of chores, the disutility of an agent after the removal of any chore assigned to her is at most  $\alpha$  times her disutility for the chores assigned to any other agent, for some  $\alpha \geq 1$ . Currently, the best-known result for chores is the existence of  $O(n^2)$ -EFX allocations [38]. In contrast, for goods, the existence of 0.618-EFX allocations is known [1], and is improved to  $2/3$  [3] for special cases. This disparity motivates the first main question of this paper:

*Do  $\alpha$ -EFX allocations of chores exist for some constant  $\alpha \geq 1$ ?*

Since a fair allocation can be subpar in terms of overall efficiency, seeking allocations that are both fair and efficient is an ideal goal. The standard notion of economic efficiency is Pareto-optimality (PO): an allocation is PO if no redistribution makes at least one agent better off without making any other agent worse off. Therefore, we seek allocations that are simultaneously PO and satisfy some relaxation of envy-freeness.

**EFX and PO for Chores.** For goods, it is known that EFX and PO allocations need not exist, and verifying their existence is NP-hard [24]. However, in the case of *bivalued* instances, where the value of each agent for the items is one of two given numbers  $\{a, b\}$ , an EFX and PO allocation exists [1] and can be computed in polynomial time [24] for goods. Bivalued instances model soft and hard preferences and are extensively studied in discrete fair division [1, 24, 21, 25, 27, 38, 4]. For bivalued instances of chores, the existence of EFX and PO allocations is known for  $n = 3$  agents [27], and  $(n - 1)$ -EFX when  $n \geq 4$ , without efficiency guarantees [38]. This prompts the question:

*Do  $\alpha$ -EFX and PO allocations exist for any  $\alpha \geq 1$ , even for structured classes such as bivalued instances?*

**EF1 and PO for Chores.** The existence of an allocation that is both EF1 and PO is another important open question in discrete fair division. While EF1 allocations always exist, whether they can be achieved with PO is unclear: verifying PO itself is coNP-hard [20].

For goods, Caragiannis et al. [14] proved that an allocation with the highest Nash welfare – the product of agents’ utilities – is both EF1 and PO. While this shows existence, it does not lead to fast computation, given that computing an allocation with maximum Nash welfare is APX-hard [30, 26]. An alternative approach to obtain PO is through *competitive equilibria* (CE). In CE allocations, agents are endowed with a fictitious amount of money, goods are assigned prices, and each agent is allocated goods that give them ‘maximum value-for-money’. The latter property ensures that the allocation is fractionally PO (fPO), a stronger efficiency property than PO. Barman et al. [10] and later Garg and Murhekar [23] utilized this approach to develop pseudo-polynomial time algorithms for computing an EF1 and PO allocation for goods.

The case of chores turns out to be significantly harder than that of goods. In fact, the existence of an EF1 and PO allocation of chores is an open question, let alone its computation. Moreover, it is unclear whether allocations that are PO and satisfy multiplicative approximations to envy-freeness up to the removal of  $k$  chores ( $\alpha$ -EF $k$ ) exist for any  $\alpha \geq 1$  and constant  $k$ ! In an  $\alpha$ -EF $k$  allocation of chores, for any two agents  $i$  and  $\ell$ , the disutility of  $i$  after the removal of  $k$  chores assigned to her is at most  $\alpha$  times her disutility for the chores assigned to  $\ell$ . This raises the following fundamental question addressed in this paper:

*What are the best values of  $\alpha$  and  $k$  for which an  $\alpha$ -EF $k$  and PO allocation of chores is guaranteed to exist?*

It is important to note that an arbitrary PO allocation is EF $m$  given  $m$  chores. However, for any constant  $k$ , it may not satisfy  $\alpha$ -EF $k$  for any  $\alpha \geq 1$ . A natural approach to address this question is to adapt techniques from goods to chores. However, the settings are only superficially similar, and all approaches fall short. Firstly, it is unclear whether there is a welfare function analogous to Nash, which guarantees EF1 and PO, for the chores. The second approach of using CE for chores has shown promise, but is currently limited to specific settings. The existence of an EF1 and PO allocation is known only for structured instances like  $n = 3$  agents [27], bivalued instances [25, 21], or two types of chores [8]. These algorithms use the CE approach to move in the space of PO allocations towards an EF1 allocation, with termination proofs heavily relying on the specialized structure of the instance. For generic instances, it remains unclear whether the CE approach will terminate in a finite time. Similarly, the existing results for EFX heavily leverage the specific structure of the instance. This prompts the following question:

*Can novel techniques be developed for the fair and efficient allocation of chores across all instances without relying on their special structure?*

## 1.1 Our Contributions

In this paper, we address the aforementioned questions regarding the existence of fair (and efficient) allocations of chores. We establish the following results:

- For *all* instances, there exists an allocation that is 4-EFX (Theorem 9).
- For *all* instances, there exists an allocation that is 2-EF2 and PO. In fact, we show a stronger guarantee: the allocation is PO, and for each agent it is 2-EF1 or EF2 (Theorem 3).
- For *bivalued* instances, there exists an allocation that is 3-EFX and PO (Theorem 12).

To prove these results, we introduce a novel concept of *earning-restricted* competitive equilibrium. In a (unrestricted) CE for chores, each agent  $i$  aims to earn an amount  $e_i > 0$  by performing chores in exchange for payment. Each chore  $j$  pays an amount  $p_j > 0$  for the completion of

the chore; if agent  $i$  performs a fraction  $x_{ij} \in [0, 1]$  of chore  $j$ , she earns  $p_j \cdot x_{ij}$ . An allocation  $\mathbf{x} = \{x_{ij}\}_{i \in [n], j \in [m]}$  and a set of chore payments  $\mathbf{p} > 0$  is a CE if all chores are fully allocated, and each agent receives her earning requirement  $e_i$ , subject to performing chores of the least disutility.

We define the concept of earning restriction by imposing a limit  $c_j > 0$  on collective earnings, agents can derive from each chore  $j$ . Consequently, agents earn  $\min\{p_j, c_j\}$  in total from chore  $j$ , and only a  $\min\{p_j, c_j\}/p_j$  fraction of chore  $j$  is allocated. This model has a natural economic interpretation: an enterprise intends to execute a project  $j$  that costs  $p_j$ , but due to financial constraints or contractual agreements, it can only allocate up to  $c_j$  towards the completion of the project. The analogous concept in the case of goods is spending restricted (SR) equilibrium, where there is a limit on the amount agents can spend on a good. Cole and Gkatzelis [17] showed the existence of an SR equilibrium. Moreover, an SR equilibrium can also be computed efficiently: it is captured by a convex program formulation, and there are polynomial-time flow-based algorithms for its computation [18, 17]. However, it is unclear whether we can design such a program for ER equilibria, and, in fact, it is not clear if an ER equilibrium even exists. Indeed, if agents demand more aggregate earning than can be disbursed by chores, an ER equilibrium cannot exist. Our next main result shows existence under a natural condition:

- An earning-restricted competitive equilibrium exists whenever  $\sum_{i \in [n]} e_i \leq \sum_{j \in [m]} c_j$  (Theorem 2).

Although for our fair division results, we only use  $e_i = 1$  for each agent  $i$  and all earning limits  $c_j$  to the same value, we establish the existence of ER competitive equilibria for arbitrary agent earning requirements  $e_i > 0$  and chore earning limits  $c_j > 0$  provided that the aforementioned condition is met.

**ER competitive equilibria for fair chore division.** We utilize earning-restricted competitive equilibria for fair and efficient chore allocation. All our algorithms begin by rounding a fractional ER equilibrium to an integral PO allocation. We ensure the resulting allocation is fair by choosing agent earning requirements and chore earning limits carefully. Intuitively, the advantage of an earning-restricted CE over an unrestricted CE for constructing a fair allocation lies in the following insight. The earning restriction on a lucrative chore forces agents to seek out less lucrative chores in the equilibrium. In doing so, the ER equilibrium reveals more information about agent preferences than an unrestricted equilibrium (see Example 2 for an illustration). In fact, without earning limits, no rounding of a CE with equal agent earnings is  $\alpha$ -EF $k$ , for any  $\alpha, k \geq 1$  (see Example 3).

Additionally, we obtain the following results.

- Existence of an allocation that is  $(n - 1)$ -EF1 and PO for all chore instances (Theorem 6).
- A polynomial time algorithm for computing a PO allocation that is *balanced*, i.e., the number of chores assigned to each agent differs by at most one (Theorem 5).
- A polynomial time algorithm for computing an EFX allocation when the number of chores is at most twice the number of agents (Theorem 10).
- A polynomial time algorithm for computing an EFX and PO allocation for bivalued instances where the number of chores is at most twice the number of agents (Theorem 14).

Our work advances the state-of-the-art by improving the best-known approximation factor  $\alpha$  for the existence of  $\alpha$ -EFX for chores from  $O(n^2)$  [38] to a constant. Moreover, we provide the first results showing the existence of  $\alpha$ -EF $k$  and PO allocations for *all* chore allocation instances for

constant  $\alpha$  and  $k$ . We note that the (non-)existence of an EF1 and PO allocation has remained elusive. In fact, an EF1 and PO allocation may not even exist in general, and 2-EF2 and PO might represent the best achievable outcome in all instances. For bivalued instances, our result showing the existence of a 3-EFX and PO allocation improves the best previously known result of  $(n - 1)$ -EFX (without PO) [38] on both fairness and efficiency fronts. Finally, we believe the concept of earning-restricted equilibrium and its existence are noteworthy and anticipate its application in other fair chore allocation problems.

## 1.2 Technical Overview

In a CE allocation, the earning of an agent serves as a proxy for her disutility in the allocation. With this, roughly speaking, it suffices to balance agents' earnings to obtain a fair allocation, while the allocation being a CE ensures that it is PO.

**Algorithm 1: 2-EF2 and PO allocation.** We uniformly set agent earning requirements to 1, and chore earning limits to  $\beta = 1/2$ ; we assume  $m \geq 2n$  to ensure an ER equilibrium exists. Given an ER equilibrium of such an instance, we design a polynomial time rounding algorithm — Algorithm 1 — that carefully rounds the ER equilibrium to ensure that the earning of every agent is at least  $\frac{1}{2}$ , and is at most 1 up to the removal of at most two chores. This gives us a 2-EF2 and PO allocation.

To obtain an approximate-EF1 guarantee, we set  $\beta = 1$  and assume  $m \geq n$ . We design another polynomial time algorithm (Algorithm 6) which rounds the ER equilibrium of such an instance and returns a  $2(n - 1)$ -EF1 and PO allocation. By carefully choosing the ‘root’ agents in the rounding, we present an improved polynomial time algorithm that computes an  $(n - 1)$ -EF1 and PO allocation given an ER equilibrium.

**Algorithm 2: Balanced PO allocation.** To address the case of  $m \leq 2n$  left out by the above approach, we design Algorithm 2: a polynomial time algorithm that gives a balanced PO allocation for any number of chores, i.e., every agent gets roughly  $m/n$  chores. When  $m \leq 2n$ , a balanced allocation is EF2, since every agent gets at most two chores. Similarly, when  $m \leq n$ , a balanced allocation is EF1, since every agent gets at most one chore.

**Algorithm 3: EFX allocation for  $m \leq 2n$ .** We design an algorithm that computes an EFX allocation for instances with  $m \leq 2n$ . Our algorithm fixes an order of the agents, say agent 1 to agent  $n$ , and first allocates chores following a picking sequence. Following the sequence, each agent picks their least-disutility chore among the remaining chores in their turn. When  $m \leq n$ , the picking sequence is simply  $1, \dots, n$ . In this case, the resulting allocation is EFX since each agent gets at most one chore. When  $m = n + r$  for  $r \in [n]$ , the picking sequence is  $r, r - 1, \dots, 1$ , followed by  $1, 2, \dots, n$ . The resulting allocation may not be EFX as the agents in  $[r]$  get two chores. To fix their EFX-envy, we perform *swaps* between the bundles of an agent  $i \in [r]$  and the agent  $\ell$  who  $i$  most envies. In such a swap,  $i$  receives the entire bundle of  $\ell$ , and  $\ell$  receives the higher-disutility chore of  $i$ . We argue that after such a swap, agents  $i$  and  $\ell$  are both EFX. We carefully argue that each agent in  $[r]$  undergoes a swap at most once, and becomes EFX after the swap. Thus, Algorithm 3 terminates with an EFX allocation after at most  $r \leq n$  swaps.

We note that although the existence of EFX allocations for  $m \leq 2n$  is known from prior work [29], our algorithm is faster and arguably simpler.

**Algorithm 4: 4-EFX allocation.** Algorithm 4 is our most technically involved algorithm, and relies on several novel ideas. Since Algorithm 3 computes an EFX allocation for instances with  $m \leq 2n$ , we focus on instances where  $m \geq 2n$ . Given an ER equilibrium of such an instance with  $e_i = 1$  and  $\beta = 1/2$ , we compute a 2-EF2 and PO allocation using Algorithm 1. Based on their payment, we partition the chores into two sets: the set  $L$  of low paying chores with payment at most  $1/2$ , and the set  $H$  of high paying chores with payment more than  $1/2$ . The 2-EF2 allocation may not be  $O(1)$ -EFX due to chores in  $H$ . We show that  $|H| \leq 2n$  and compute an EFX re-allocation of  $H$  using Algorithm 3. Using the structure of the initial rounded allocation and the EFX property of the allocation of  $H$ , we show the counter-intuitive fact that agents with at least two  $H$  chores are 4-EFX. This leaves us to address the EFX-envy of agents with a single  $H$  chore. For such an agent  $i$ , we *swap* the  $H$  chore of  $i$  with all chores belonging to the agent  $\ell$  that  $i$  most envies, termed an  $(i, \ell)$  swap. We carefully argue that there can only be polynomially many swap steps before the algorithm terminates with a 4-EFX allocation. To do this, we show that immediately after an  $(i, \ell)$  swap, agents  $i$  and  $\ell$  are both 4-EFX. Moreover, by carrying out swaps in a carefully chosen order, we can argue that an agent  $i$  who participated in an  $(i, \ell)$  swap remains 4-EFX throughout the execution of the algorithm. This shows that Algorithm 4 terminates with a 4-EFX allocation after at most  $n$  swaps.

**Algorithm 5: 3-EFX and PO for bivalued instances.** We next design Algorithm 5, which returns a 3-EFX and PO allocation for bivalued instances given an ER equilibrium with  $\beta = 1/2$ ; for this we assume  $m > 2n$ . Algorithm 5 uses the ideas of chore swaps used in Algorithm 4, but the additional structure offered by the bivalued nature of the instance allows us to improve the approximation guarantee to 3-EFX while also maintaining PO.

**Algorithm 7: EFX and PO for bivalued instances with  $m \leq 2n$ .** For bivalued instances with  $m \leq 2n$ , we design Algorithm 7 which computes an EFX and PO allocation in polynomial time. Algorithm 7 begins with a balanced allocation computed using Algorithm 2, and then essentially runs Algorithm 5. Since the number of chores is limited, a careful analysis shows that the guarantee of the resulting allocation can be improved to EFX and PO.

**Existence of ER equilibrium.** Proving the existence of an ER equilibrium turns out to be quite non-trivial. Existing flow-based algorithms designed for computing a CE or a spending-restricted equilibrium in the context of goods do not apply straightforwardly to chores. While computing a (unrestricted) CE for chores is more challenging compared to goods, several combinatorial algorithms exist that compute approximate CEs. However, these algorithms also do not seem to extend to show the existence and computation of ER equilibria for chores.

To prove existence, we devise a linear complementarity problem (LCP) formulation that captures all ER equilibria of an instance. We then show that Lemke’s scheme – a simplex-like complementary pivot algorithm that traverses the vertices of a polyhedron associated with the LCP – must terminate at an ER equilibrium. This requires a careful design of the LCP to show that Lemke’s scheme does not end up on vertices that do not correspond to ER equilibria or on infinite edges of the polyhedron.

### 1.3 Further Related Work

We discuss other related literature that is most relevant to EF1/EFX and PO allocations. For other related work, we refer the reader to excellent surveys [7, 2].

**EFX.** For goods, EFX allocations are known to exist for two agents (via cut-or-choose), identically ordered (IDO) instances [36], three agents [15] and two types of agents [33]. Recent works [12, 3] improve the approximation guarantee beyond 0.618 in special cases. For chores, existence is known for two agents, IDO instances [31] and two types of chores [8]. Using matching-based approaches, [29] showed existence for instances where (i) the number of chores is at most twice the number of agents, (ii) all but one agent have IDO disutility functions, and (iii) there are three agents with 2-ary disutilities. For chores, [16] showed the existence of 2-EFX allocations for  $n = 3$  agents.

**EF1/EFX and PO.** For goods, an EF1 and PO allocation can be computed in polynomial time for binary instances [11], for  $k$ -ary instances where  $k$  is a constant, and for a constant number of agents [23]. For chores, existence and polynomial time computation is also known for (i) identical agents (folklore), (ii) two agents [5], and (iii) three types of agents [27, 28]. [24] showed that checking the existence of EFX and PO allocations is NP-hard for goods. For chores, [27] showed that EFX and fPO allocations need not exist even for two agents with 2-ary disutilities. Later, [27] proved an EFX and PO allocation can be computed in polynomial time for  $n = 3$  agents with bivalued disutility functions.

**Spending restricted equilibrium and CE rounding.** Cole and Gkatzelis [17] introduced the concept of spending-restricted equilibria for goods and gave a polynomial time algorithm for computing such an equilibrium. SR equilibria have been used to develop approximation algorithms for Nash welfare [17, 18]. [35] used SR equilibria to compute approximately Nash optimal allocations that are PO and satisfy Prop1: a discrete relaxation of the share-based fairness notion of proportionality. [9] showed that a CE can be rounded to give a Prop1 and PO allocation. A similar result was obtained by [6]. Example 1 shows that such an allocation need not be  $\alpha$ -EF $k$  for any  $\alpha, k \geq 1$ .

**Organization of the remainder of the paper.** Section 2 defines the problem formally and introduces the earning-restricted model. Section 3 presents our results on the existence of allocations that are (i) 2-EF2 and PO (one lemma in Appendix B) and (ii)  $(n - 1)$ -EF1 and PO (deferred to Appendix C). Section 4 presents our algorithms proving the existence 4-EFX allocations. The case of  $m \leq 2n$  is discussed in Section 4.1, while the general case is discussed in Section 4.2 and Section 4.3. Section 5 presents the existence of 3-EFX and PO allocations for bivalued instances. The existence of EFX and PO allocations of bivalued chores when  $m \leq 2n$  is discussed in Appendix E. Finally, we show the existence of ER equilibria in Section 6. Appendix A contains illustrative examples.

## 2 Preliminaries

**Problem instance.** An instance  $(N, M, D)$  of the chore allocation problem consists of a set  $N = [n]$  of  $n$  agents, a set  $M = [m]$  of  $m$  indivisible chores, and a list  $D = \{d_i\}_{i \in N}$ , where  $d_i : 2^M \rightarrow \mathbb{R}_{\geq 0}$  is agent  $i$ 's *disutility* function over the chores. Let  $d_{ij} > 0$  denote the disutility of chore  $j$  for agent  $i$ . We assume that the disutility functions are additive; thus for every  $i \in N$  and  $S \subseteq M$ ,  $d_i(S) = \sum_{j \in S} d_{ij}$ . An instance is said to be *bivalued* if there exist  $a, b \in \mathbb{R}_{>0}$  such that  $d_{ij} \in \{a, b\}$  for all  $i \in N, j \in M$ .

**Allocation.** An *integral allocation*  $\mathbf{x} = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)$  is an  $n$ -partition of the chores; here  $\mathbf{x}_i \subseteq M$  is the set of chores assigned to agent  $i$ , who receives disutility  $d_i(\mathbf{x}_i)$ . In a *fractional allocation*  $\mathbf{x} \in [0, 1]^{n \times m}$ , chores are divisible and  $x_{ij} \in [0, 1]$  denotes the fraction of chore  $j$  assigned

to agent  $i$ , who receives disutility  $d_i(\mathbf{x}_i) = \sum_{j \in M} d_{ij} \cdot x_{ij}$ . We assume allocations are integral unless specified.

**Fairness and efficiency notions.** An allocation  $\mathbf{x}$  is said to be:

1.  $\alpha$ -Envy-free up to  $k$  chores ( $\alpha$ -EF $k$ ) if for all  $i, h \in N$ , there exists  $S \subseteq \mathbf{x}_i$  with  $|S| \leq k$  such that  $d_i(\mathbf{x}_i \setminus S) \leq \alpha \cdot d_i(\mathbf{x}_h)$ .
2.  $\alpha$ -Envy-free up to any chore ( $\alpha$ -EFX) if for all  $i, h \in N$  and  $j \in \mathbf{x}_i$ ,  $d_i(\mathbf{x}_i \setminus \{j\}) \leq \alpha \cdot d_i(\mathbf{x}_h)$ .
3. Pareto optimal (PO) if there is no allocation  $\mathbf{y}$  that dominates  $\mathbf{x}$ . An allocation  $\mathbf{y}$  dominates allocation  $\mathbf{x}$  if for all  $i \in N$ ,  $d_i(\mathbf{y}_i) \leq d_i(\mathbf{x}_i)$ , and there exists  $h \in N$  such that  $d_h(\mathbf{y}_h) < d_h(\mathbf{x}_h)$ .
4. Fractionally Pareto-optimal (fPO) if there is no fractional allocation that dominates  $\mathbf{x}$ . An fPO allocation is clearly PO, but not vice-versa.

## 2.1 Competitive equilibrium

An instance  $(N, M, D, e)$  of a Fisher model for chores consists of a set  $N$  of agents, set  $M$  of chores, list  $D = \{d_i\}_{i \in N}$  specifying the disutility functions of the agents, and an *earning requirement*  $e_i > 0$  for each agent  $i \in N$ . We associate payments  $\mathbf{p} = (p_1, \dots, p_m) \in \mathbb{R}_{>0}^m$  with the chores, i.e. chore  $j$  pays  $p_j$ . Each agent  $i$  aims to earn at least  $e_i$  by performing chores in exchange for payment. In a (fractional) allocation  $\mathbf{x}$  with payments  $\mathbf{p}$ , the *earning* of agent  $i$  is  $\mathbf{p}(\mathbf{x}_i) = \sum_{j \in M} p_j \cdot x_{ij}$ . An allocation  $(\mathbf{x}, \mathbf{p})$  is said to be a competitive equilibrium if all chores are allocated and all agents earn their earning requirement subject to performing chores of least possible disutility. For additive disutilities, the latter condition can be expressed in terms of disutility-to-payment ratios as follows.

**Definition 1.** (*MPB allocation*) For each agent  $i$ , the *pain-per-buck* ratio  $\alpha_{ij}$  of chore  $j$  is defined as  $\alpha_{ij} = d_{ij}/p_j$ , and the *minimum-pain-per-buck* (MPB) ratio of agent  $i$  is then given by  $\alpha_i = \min_{j \in M} \alpha_{ij}$ . Let  $MPB_i = \{j \in M \mid d_{ij}/p_j = \alpha_i\}$  denote the set of chores which are MPB for agent  $i$  for payments  $\mathbf{p}$ . An allocation  $(\mathbf{x}, \mathbf{p})$  is called an *MPB allocation* if for all  $i \in N$  and  $j \in M$ ,  $x_{ij} > 0$  implies  $j \in MPB_i$ , i.e., agents are only assigned chores which are MPB for them.

**Definition 2.** (*Competitive equilibrium*) We say that  $(\mathbf{x}, \mathbf{p})$  is a *competitive equilibrium* (CE) for the instance  $(N, M, D, e)$  if (i) for all  $j \in M$ ,  $\sum_{i \in N} x_{ij} = 1$ , i.e., all chores are completely allocated, (ii) for all  $i \in N$ ,  $\mathbf{p}(\mathbf{x}_i) = e_i$ , i.e., each agent receives her earning requirement, and (iii)  $(\mathbf{x}, \mathbf{p})$  is an MPB allocation.

The First Welfare Theorem [34] shows that for a competitive equilibrium  $(\mathbf{x}, \mathbf{p})$  of instance  $(N, M, D, e)$ , the allocation  $\mathbf{x}$  is fPO. Using this fact, we can argue:

**Proposition 1.** *Let  $(\mathbf{x}, \mathbf{p})$  be an MPB allocation. Then  $\mathbf{x}$  is fPO.*

*Proof.* We create an associated Fisher market instance  $I = (N, M, D, e)$  by defining  $e_i = \mathbf{p}(\mathbf{x}_i)$  for each  $i \in N$ . It is easy to see that  $(\mathbf{x}, \mathbf{p})$  is a CE for  $I$ . By the First Welfare Theorem,  $\mathbf{x}$  is fPO.  $\square$

The above proposition shows that MPB allocations are useful in ensuring efficiency. We now discuss how such allocations can be utilized for fairness. For an MPB allocation  $(\mathbf{x}, \mathbf{p})$  where  $\mathbf{x}$  is integral, we let  $\mathbf{p}_{-k}(\mathbf{x}_i) := \min_{S \subseteq \mathbf{x}_i, |S| \leq k} \mathbf{p}(\mathbf{x}_i \setminus S)$  denote the payment agent  $i$  receives from  $\mathbf{x}_i$  excluding her  $k$  highest paying chores. Likewise, we let  $\mathbf{p}_{-X}(\mathbf{x}_i) := \max_{j \in \mathbf{x}_i} \mathbf{p}(\mathbf{x}_i \setminus \{j\})$  denote the payment  $i$  receives from  $\mathbf{x}_i$  excluding her lowest paying chore.



**Definition 3** (Payment EFk and Payment EFX). An allocation  $(\mathbf{x}, \mathbf{p})$  is said to be  $\alpha$ -payment envy-free up to  $k$  chores ( $\alpha$ -pEFk) if for all  $i, h \in N$  we have  $\mathbf{p}_{-k}(\mathbf{x}_i) \leq \alpha \cdot \mathbf{p}(\mathbf{x}_h)$ . Agent  $i$   $\alpha$ -pEFk-envies  $h$  if  $\mathbf{p}_{-k}(\mathbf{x}_i) > \alpha \cdot \mathbf{p}(\mathbf{x}_h)$ .

An allocation  $(\mathbf{x}, \mathbf{p})$  is said to be  $\alpha$ -payment envy-free up any chore ( $\alpha$ -pEFX) if for all  $i, h \in N$  we have  $\mathbf{p}_{-X}(\mathbf{x}_i) \leq \alpha \cdot \mathbf{p}(\mathbf{x}_h)$ . Agent  $i$   $\alpha$ -pEFX-envies  $h$  if  $\mathbf{p}_{-X}(\mathbf{x}_i) > \alpha \cdot \mathbf{p}(\mathbf{x}_h)$ .

We derive a sufficient condition for computing an  $\alpha$ -EFk/ $\alpha$ -EFX and PO allocation.

**Lemma 1.** *Let  $(\mathbf{x}, \mathbf{p})$  be an MPB allocation where  $\mathbf{x}$  is integral.*

(i) *If  $(\mathbf{x}, \mathbf{p})$  is  $\alpha$ -pEFk, then  $\mathbf{x}$  is  $\alpha$ -EFk and fPO.*

(ii) *If  $(\mathbf{x}, \mathbf{p})$  is  $\alpha$ -pEFX, then  $\mathbf{x}$  is  $\alpha$ -EFX and fPO.*

*Proof.* Since  $(\mathbf{x}, \mathbf{p})$  is an MPB allocation, Proposition 1 shows  $\mathbf{x}$  is fPO. Let  $\alpha_i$  be the MPB ratio of agent  $i$  in  $(\mathbf{x}, \mathbf{p})$ . Consider any pair of agents  $i, h \in N$ .

(i) If  $(\mathbf{x}, \mathbf{p})$  is  $\alpha$ -pEFk, then:

$$\min_{S \subseteq \mathbf{x}_i, |S| \leq k} d_i(\mathbf{x}_i \setminus S) = \alpha_i \cdot \mathbf{p}_{-k}(\mathbf{x}_i) \leq \alpha_i \cdot \alpha \cdot \mathbf{p}(\mathbf{x}_h) \leq \alpha \cdot d_i(\mathbf{x}_h),$$

where the first and last transitions use that  $(\mathbf{x}, \mathbf{p})$  is on MPB, and the middle inequality uses that  $(\mathbf{x}, \mathbf{p})$  is  $\alpha$ -pEFk (Definition 3). Thus,  $\mathbf{x}$  is  $\alpha$ -EFk.

(ii) If  $(\mathbf{x}, \mathbf{p})$  is  $\alpha$ -pEFX, then:

$$\max_{j \in \mathbf{x}_i} d_i(\mathbf{x}_i \setminus \{j\}) = \alpha_i \cdot \mathbf{p}_{-X}(\mathbf{x}_i) \leq \alpha_i \cdot \alpha \cdot \mathbf{p}(\mathbf{x}_h) \leq \alpha \cdot d_i(\mathbf{x}_h),$$

where the first and last transitions use that  $(\mathbf{x}, \mathbf{p})$  is on MPB, and the middle inequality uses that  $(\mathbf{x}, \mathbf{p})$  is  $\alpha$ -pEFX (Definition 3). Thus,  $\mathbf{x}$  is  $\alpha$ -EFX.  $\square$

## 2.2 Earning-restricted equilibrium

We introduce a new concept of *earning-restricted (ER) competitive equilibrium* for chores. An instance  $(N, M, D, e, c)$  of the ER competitive equilibrium problem consists of a set  $N = [n]$  of  $n$  agents, a set  $M = [m]$  of  $m$  chores, a list  $D = \{d_i\}_{i \in N}$  of additive agent disutility functions, a list  $e = \{e_i\}_{i \in N}$  of agent earning requirements, and a list  $c = \{c_j\}_{j \in M}$  of chore earning-restrictions. Each agent  $i \in N$  aims to earn at least  $e_i > 0$  by performing chores in exchange for payment from the chores. Further, the earning limit  $c_j \geq 0$  specifies a limit on the money that agents can collectively earn from chore  $j \in M$ .

An equilibrium  $(\mathbf{x}, \mathbf{p})$  of an ER instance consists of a partial fractional allocation  $\mathbf{x}$  and a set of chore payments  $\mathbf{p}$  such that each agent  $i$  earns her earning requirement  $e_i$  while performing chores of least possible disutility, with the restriction that the earning from each chore  $j$  is at most  $c_j$ . Once a chore has paid  $c_j$  to the agents, the rest of the chore is not assigned. Define the *earning vector*  $\mathbf{q} \in \mathbb{R}^{n \times m}$  associated with  $(\mathbf{x}, \mathbf{p})$  given by  $q_{ij} := p_j x_{ij}$  which denotes the amount agent  $i$  earns from chore  $j$ . Let  $q_j = \sum_i q_{ij} = \sum_i p_j x_{ij}$  denote the total earning from chore  $j$ . We now formally define an ER equilibrium  $(\mathbf{x}, \mathbf{p})$ .

**Definition 4** (Earning-restricted equilibrium). *Let  $\mathbf{q}$  be the earning vector associated with an allocation  $(\mathbf{x}, \mathbf{p})$ . Then  $(\mathbf{x}, \mathbf{p})$  is an earning-restricted equilibrium of an ER instance  $(N, M, D, e, c)$  if*

- (i) (Agents)  $(\mathbf{x}, \mathbf{p})$  is an MPB allocation, i.e., for  $i \in N, j \in M, x_{ij} > 0$  implies  $j \in MPB_i$ . Moreover, for each  $i \in N, \sum_j q_{ij} = e_i$ .
- (ii) (Chores) For each  $j \in M$ , either  $\sum_i x_{ij} = 1$  and  $q_j = p_j \leq c_j$ , or  $\sum_i x_{ij} < 1$  and  $q_j = c_j < p_j$ . In other words, for each  $j, q_j = \min\{p_j, c_j\}$ .

For notational convenience, we often use both  $(\mathbf{x}, \mathbf{p})$  and  $(\mathbf{x}, \mathbf{p}, \mathbf{q})$  to denote an ER equilibrium. See Example 2 for an illustration of an ER equilibrium. Clearly, an ER equilibrium can exist only if  $\sum_i e_i \leq \sum_j c_j$ , i.e., the chores must collectively pay enough so all agents can earn their earning requirements. In Section 6, we prove that this condition is in fact sufficient for existence.

**Theorem 2.** *Every ER instance  $(N, M, D, e, c)$  satisfying  $\sum_i e_i \leq \sum_j c_j$  admits an ER equilibrium.*

### 3 Existence of 2-EF2 and PO Allocations

In this section, we prove our main result concerning the existence of 2-EF2 and fPO allocations for all chore allocation instances.

**Theorem 3.** *Any chore allocation instance admits a 2-EF2 and fPO allocation. More precisely, for every agent the allocation is either 2-EF1 or EF2.*

The main idea behind Theorem 3 is to use ER equilibria to compute a fair and efficient allocation. Given a chore allocation instance, we set uniform agent earning requirements  $e_i = 1$  and impose a uniform earning limit of  $\beta \in [1/2, 1)$  on all chores. When  $m \cdot \beta \geq n$ , the feasible earning condition is satisfied and an ER equilibrium  $(\mathbf{z}, \mathbf{p})$  exists by Theorem 2. We design a polynomial time algorithm Algorithm 1 which carefully rounds the fractional allocation  $\mathbf{z}$  to an integral allocation  $\mathbf{x}$  that is approximately-EF $k$  and fPO. With different choices of  $\beta$ , the rounded integral allocation satisfies different fairness guarantees. In particular, setting  $\beta = 1/2$  gives a 2-EF2 and fPO allocation.

To handle the case of  $m \leq 2n$ , we design a polynomial time algorithm (Algorithm 2), which computes an EF2 and fPO allocation. Specifically, for any number of chores, Algorithm 2 produces an fPO allocation in which the number of chores in agent bundles differ by at most one, i.e., is balanced. Thus, for  $m \leq 2n$ , each agent gets at most two chores and hence the allocation is EF2. The algorithm starts with an imbalanced allocation and transfers chores from agents with a higher number of chores to agents with a lower number chores until the allocation is balanced, while preserving fPO.

#### 3.1 Earning-restricted rounding

We now describe Algorithm 1, which rounds a fractional ER equilibrium  $(\mathbf{y}, \mathbf{p})$  of an instance with uniform chore earning limit  $\beta$  to an approximately-EF $k$  and fPO allocation.

**Definition 5** (Payment graph). *The payment graph  $G = (N, M, E)$  associated with an allocation  $(\mathbf{x}, \mathbf{p})$  is a weighted bipartite graph with vertex set  $V(G) = N \sqcup M$ , and edge set  $E(G) = \{(i, j) : i \in N, j \in M, x_{ij} > 0\}$ . The weight of edge  $(i, j)$  is  $p_j \cdot x_{ij}$ , which is the earning of agent  $i$  from chore  $j$ .*

Algorithm 1 first transforms the given equilibrium into one whose payment graph is acyclic, i.e., is a collection of trees. This is due to the following lemma, proved in Appendix B.

---

**Algorithm 1** Earning-Restricted Rounding

---

**Input:** Instance  $(N, M, D)$  with  $m\beta \geq n$ ; ER equilibrium  $(\mathbf{y}, \mathbf{p})$  with earning limit  $\beta \in [1/2, 1)$

**Output:** An integral allocation  $\mathbf{x}$

```
1:  $(\mathbf{z}, \mathbf{p}) \leftarrow \text{MakeAcyclic}(\mathbf{y}, \mathbf{p})$ 
2: Let  $G = (N, M, E)$  be the payment graph associated with  $(\mathbf{z}, \mathbf{p})$ 
3: Root each tree of  $G$  at some agent and orient edges
4:  $\mathbf{x}_i \leftarrow \emptyset$  for all  $i \in N$  ▷ Initialize empty allocation
5:  $L = \{j \in M : p_j \leq \beta\}$ ,  $H = \{j \in M : p_j > \beta\}$  ▷ Low, High paying chores
   — Phase 1: Round leaf chores —
6: for all leaf chores  $j$  do
7:    $\mathbf{x}_i \leftarrow \mathbf{x}_i \cup \{j\}$  for  $i = \text{parent}(j)$ ; delete  $j$  from  $G$ 
   — Phase 2: Allocate  $L$  —
8: for every tree  $T$  of  $G$  do
9:   for every agent  $i$  of  $T$  in BFS order do
10:    if  $\mathbf{p}(\mathbf{x}_i) > 1$  then
11:      for every  $j \in \text{child}(i) \cap H$  do
12:        Assign  $j$  to agent  $h \in \text{child}(j)$  earning most from  $j$  among  $\text{child}(j)$ ; delete  $j$ 
13:      while  $\exists j \in \text{child}(i) \cap L$  s.t.  $\mathbf{p}(\mathbf{x}_i \cup \{j\}) \leq 1$  do
14:         $\mathbf{x}_i \leftarrow \mathbf{x}_i \cup \{j\}$ ; delete  $j$  from  $G$ 
15:      for every  $j \in \text{child}(i) \cap L$  do
16:        Assign  $j$  to arbitrary agent  $h \in \text{child}(j)$ ; delete  $j$  from  $G$ 
   — Phase 3: Pruning trees —
17: for chore  $j \in V(G) \cap M$  do
18:   if agent  $i \in \text{child}(j)$  does not earn the most from  $j$  among agents in  $\text{child}(j)$  then
19:     Delete edge  $(j, i)$  from  $G$ 
   — Phase 4: Matching to allocate  $H$  —
20: for every tree  $T = (N(T) \cup M(T), E(T))$  of  $G$  do
21:    $h \leftarrow \arg \max_{i \in N(T)} \mathbf{p}(\mathbf{x}_i)$ 
22:   Compute a matching  $\sigma$  of  $i \in N(T) \setminus \{h\}$  to  $M(T)$ 
23:   for  $i \in N(T) \setminus \{h\}$  do
24:      $\mathbf{x}_i \leftarrow \mathbf{x}_i \cup \{\sigma(i)\}$ 
25: return  $\mathbf{x}$ 
```

---

**Lemma 2.** *There is a polynomial time algorithm `MakeAcyclic` which takes as input an ER equilibrium of instance  $I$  and returns an another ER equilibrium of  $I$  whose payment graph is acyclic.*

Given the ER equilibrium  $(\mathbf{z}, \mathbf{p})$  whose payment graph  $G$  is acyclic, Algorithm 1 roots each tree of  $G$  at some agent and orients its edges. For a node  $v \in V(G)$ , let  $\text{child}(v)$  denote the children nodes of  $v$  and  $\text{parent}(v)$  denote the parent node of  $v$ . Note that the root nodes of trees in  $G$  are agents and the leaf nodes are chores. We let  $\mathbf{x}$  denote the integral allocation of chores to agents made by Algorithm 1, which is initially empty. We classify chores into two sets:  $L = \{j \in M : p_j \leq \beta\}$  comprising of low cost chores, and  $H = \{j \in M : p_j > \beta\}$  comprising of high cost or overpaying chores. Algorithm 1 proceeds in four phases.

Phase 1 rounds every leaf chore  $j$  to their parent agent  $\text{parent}(j)$  and then deletes  $j$  from  $G$ . After this, all chores in  $G$  have edges to at least two agents, i.e., are *shared* chores. Note that there can be at most  $(n - 1)$  shared chores, since  $G$  is acyclic.

Phase 2 assigns chores in  $L$ . In each tree  $T$  of  $G$ , we visit agents in breadth-first order starting from the root. At agent  $i$ , we first check if  $\mathbf{p}(\mathbf{x}_i) > 1$ . Note that this can happen only if  $i$  received the  $\mathbf{parent}(i)$  chore. If so, we assign every chore  $j \in \mathbf{child}(i)$  to a child agent of  $j$ . A chore  $j \in \mathbf{child}(i) \cap L$  is assigned to an arbitrary child of  $j$ , while  $j \in \mathbf{child}(i) \cap H$  is assigned to an agent who earns the most from  $j$  among children of  $j$ . After this, such an agent  $i$  is not assigned any further chores in the algorithm. Otherwise, if  $\mathbf{p}(\mathbf{x}_i) \leq 1$  when visiting  $i$ , we iteratively assign the child chores of  $i$  in  $L$  as long as  $\mathbf{p}(\mathbf{x}_i) \leq 1$ . Any remaining child chore  $j \in L$  is assigned to an arbitrary child agent of  $j$ . Thus at the end of phase 2, all chores in  $L$  have been allocated, and the graph  $G$  is a collection of ‘Phase 2 trees’ whose vertices are agents and chores from  $H$ .

Phase 3 prunes Phase 2 trees by deleting certain edges. For every shared chore  $j \in H$ , we delete the edge  $(j, i)$  for  $i \in \mathbf{child}(j)$  if  $i$  does not earn the most from  $j$  among the child agents of  $j$ . As a result, we obtain ‘Phase 3’ trees in which each chore  $j \in H$  is adjacent to exactly two agents.

Phase 4 assigns the remaining shared chores in  $H$ . Due to the pruning phase, each Phase 3 tree  $T = (N(T) \cup M(T), E(T))$  with  $|N(T)| = r$  agents has exactly  $|M(T)| = r - 1$  shared chores from  $H$ . We identify an agent  $h \in N(T)$  with the highest earning  $\mathbf{p}(\mathbf{x}_h)$ , and then assign the  $(r - 1)$  chores of  $M(T)$  to the  $(r - 1)$  agents of  $N(T) \setminus \{h\}$  via a matching. Such a matching is possible because in a Phase 3 tree, each shared chore is adjacent to exactly two agents. Thus during Phase 4, every agent gets at most one chore from  $H$ , and all chores are allocated.

**Lemma 3.** *Given an ER equilibrium  $(\mathbf{y}, \mathbf{p})$  of an instance  $(N, M, D)$ , Algorithm 1 returns an integral allocation  $\mathbf{x}$  in  $\text{poly}(n, m)$  time.*

*Proof.* Lemma 2 shows that the procedure `MakeAcyclic` results in an allocation with an acyclic payment graph in polynomial time. In the following phases, Algorithm 1 assigns all chores to agents. Each phase takes polynomial time since they involve polynomial time operations such as BFS in the payment graph or computing a matching in a tree.  $\square$

We now analyze the properties of the allocation  $\mathbf{x}$  returned by Algorithm 1. We first show:

**Lemma 4.** *The allocation  $\mathbf{x}$  returned by Algorithm 1 is fPO.*

*Proof.* Since  $(\mathbf{z}, \mathbf{p})$  is an ER equilibrium,  $(\mathbf{z}, \mathbf{p})$  is an MPB allocation. Let  $Z_i = \{j : z_{ij} > 0\}$ . Note that throughout Algorithm 1,  $\mathbf{x}_i \subseteq Z_i$ . Hence,  $(\mathbf{x}, \mathbf{p})$  is also an MPB allocation. Consequently, Proposition 1 implies that  $\mathbf{x}$  is fPO.  $\square$

To analyze fairness properties of  $\mathbf{x}$ , we first prove upper bounds on agent earnings. Essentially, the following lemma states that the earning up to one chore of each agent is at most 1, except when the agent has two chores from  $H$ ; in the latter case the agent earns at most  $1 - \beta$  from other chores.

**Lemma 5.** *Let  $(\mathbf{x}, \mathbf{p})$  be the allocation returned by Algorithm 1 with earning restriction  $\beta \in [1/2, 1)$ . Then for each  $i \in N$ , either  $\mathbf{p}_{-1}(\mathbf{x}_i) \leq 1$ , or  $|\mathbf{x}_i \cap H| = 2$  and  $\mathbf{p}_{-2}(\mathbf{x}_i) \leq 1 - \beta$ .*

*Proof.* Let  $\mathbf{x}^t$  denote the allocation after Phase  $t$ , for  $t \in [4]$ ; note that  $\mathbf{x}^4 = \mathbf{x}$ . Consider an agent  $i \in N$ . Let  $\hat{\mathbf{x}}_i$  be the allocation when Algorithm 1 visits  $i$  in Phase 2. Suppose  $\mathbf{p}(\hat{\mathbf{x}}_i) \leq 1$ . Then we have  $\mathbf{p}(\mathbf{x}_i^2) \leq 1$  at the end of Phase 2 after  $i$  is assigned a subset of  $\mathbf{child}(i) \cap L$ . Subsequently,  $i$  can be assigned one more chore in Phase 4. Hence we have  $\mathbf{p}_{-1}(\mathbf{x}_i) \leq 1$  in this case.

On the contrary, suppose  $\mathbf{p}(\hat{\mathbf{x}}_i) > 1$ . Then Algorithm 1 will not allocate any chore to  $i$  in Phase 4, and hence  $\mathbf{x}_i = \mathbf{x}_i^2 = \hat{\mathbf{x}}_i$ . Note that either  $\hat{\mathbf{x}}_i = \mathbf{x}_i^1$  or  $\hat{\mathbf{x}}_i = \mathbf{x}_i^1 \cup \{j\}$ , where  $j = \mathbf{parent}(i)$ . That is,  $\hat{\mathbf{x}}_i$  includes the chores  $\mathbf{x}_i^1$  allocated to  $i$  in Phase 1, and may include  $i$ ’s parent chore  $j$ . Recall that Phase 1 rounds leaf chores to their parent agents, hence  $\mathbf{x}_i^1$  comprises of the leaf chores that

are also child chores of  $i$ . Due to the earning restriction of  $\beta$ , agent  $i$  earns exactly  $\beta$  from any chore in  $\mathbf{x}_i^1 \cap H$ . Since  $\beta \geq 1/2$  and  $e_i = 1$ , we have  $|\mathbf{x}_i^1 \cap H| \leq 2$ . We consider three cases:

- $|\mathbf{x}_i^1 \cap H| = 0$ . Then we have  $\mathbf{p}(\mathbf{x}_i^1) \leq 1$ . Hence  $\mathbf{p}_{-1}(\hat{\mathbf{x}}_i) \leq \mathbf{p}(\hat{\mathbf{x}}_i \setminus \{j\}) \leq \mathbf{p}(\mathbf{x}_i^1) \leq 1$ .
- $|\mathbf{x}_i^1 \cap H| = 1$ , and let  $\mathbf{x}_i^1 \cap H = \{j_1\}$ . Then  $\mathbf{p}(\mathbf{x}_i^1 \setminus \{j_1\}) \leq 1 - \beta$ , since the earning of  $i$  from the  $j_1$  is exactly  $\beta$ . We consider cases depending on the payment of  $i$ 's parent chore  $j$ .
  - If  $j \notin H$ , then  $p_j \leq \beta$ . Hence  $\mathbf{p}_{-1}(\hat{\mathbf{x}}_i) = \mathbf{p}(\hat{\mathbf{x}}_i \setminus \{j_1\}) \leq \mathbf{p}((\mathbf{x}_i^1 \setminus \{j_1\}) \cup \{j\}) \leq (1 - \beta) + \beta = 1$ .
  - If  $j \in H$  and  $j \notin \hat{\mathbf{x}}_i$ , then  $\mathbf{p}_{-1}(\hat{\mathbf{x}}_i) = \mathbf{p}(\mathbf{x}_i^1 \setminus \{j_1\}) \leq 1 - \beta < 1$ .
  - If  $j \in H$  and  $j \in \hat{\mathbf{x}}_i$ , then  $|\mathbf{x}_i \cap H| = 2$ , and  $\mathbf{p}_{-2}(\hat{\mathbf{x}}_i \setminus \{j, j_1\}) \leq \mathbf{p}(\mathbf{x}_i^1 \setminus \{j_1\}) \leq 1 - \beta$ .
- $|\mathbf{x}_i^1 \cap H| = 2$ . In this case,  $\mathbf{p}(\mathbf{x}_i^1) \geq 2\beta$ , since  $i$  earns exactly  $\beta$  from each chore in  $\mathbf{x}_i^1 \cap H$ . However since  $\beta \geq 1/2$  and  $e_i = 1$ , this case can only arise if  $\beta = 1/2$ , in which case  $i$  can only be earning from the two chores in  $\mathbf{x}_i^1$ . Thus  $i$  has no parent chore, and  $\mathbf{x}_i = \mathbf{x}_i^1$ . Hence,  $|\mathbf{x}_i \cap H| = 2$  and  $\mathbf{p}_{-2}(\mathbf{x}_i) = 0$ .

This proves the lemma. □

We next establish lower bounds on agent earnings. These bounds are derived by investigating the allocation computed by the matching phase (Phase 4). We say agent  $i$  loses a chore  $j$  if  $i$  is earning from  $j$  in the fractional solution  $\mathbf{z}$  but not in the integral allocation  $\mathbf{x}$ , i.e.,  $z_{ij} > 0$  but  $j \notin \mathbf{x}_i$ .

**Lemma 6.** *Let  $(\mathbf{x}, \mathbf{p})$  be the allocation returned by Algorithm 1. Then for each agent  $i \in N$ ,  $\mathbf{p}(\mathbf{x}_i) \geq \min\{\beta, 1 - \beta\}$ .*

*Proof.* Let  $(\mathbf{z}, \mathbf{p})$  be the ER equilibrium whose payment graph is acyclic, which is computed before Phase 1 begins. Let  $\mathbf{x}^t$  denote the allocation after Phase  $t$  of Algorithm 1, for  $t \in [4]$ . Note that  $\mathbf{x}^2 = \mathbf{x}^3$  since Phase 3 does not assign any chores and only deletes edges in  $G$ . Also note  $\mathbf{x}^4 = \mathbf{x}$ .

Let  $T = (N(T) \cup M(T), E(T))$  be a Phase 3 tree rooted at agent  $i_0$ . Since  $T$  is a Phase 3 tree,  $T$  has exactly  $|N(T)| - 1$  chores, all of which belong to  $H$ . Phase 4 identifies the agent  $h \in \arg \max_{i \in N(T)} \mathbf{p}(\mathbf{x}_i^3)$ , and assigns a chore  $\sigma(i) \in H$  to every agent  $i \in N(T) \setminus \{h\}$  by computing a matching of  $M(T)$  to  $N(T) \setminus \{h\}$ . Since  $p_j > \beta$  for  $j \in H$ , we have  $\mathbf{p}(\mathbf{x}_i) \geq p_{\sigma(i)} > \beta$  for all  $i \in N(T) \setminus \{h\}$ . Hence we only need to prove lower bounds on the earning  $\mathbf{p}(\mathbf{x}_h)$  of the agent  $h$ . Note that  $\mathbf{x}_h = \mathbf{x}_h^3 = \mathbf{x}_h^2$ , since  $h$  is not allocated any chores in Phase 3 or 4. By choice of  $h$ , we also have that  $\mathbf{p}(\mathbf{x}_h) \geq \mathbf{p}(\mathbf{x}_i^3) = \mathbf{p}(\mathbf{x}_i^2)$  for all  $i \in N(T)$ . We now analyze three scenarios.

- (i) Some agent  $i \in N(T)$  lost a child chore  $j \in \text{child}(i)$ . Suppose  $i$  lost  $j$  in Phase 2. If  $j \in H$ , then it must be that  $\mathbf{p}(\mathbf{x}_i^2) > 1$ . If  $j \in L$ , then it must be that  $\mathbf{p}(\mathbf{x}_i^2) \geq 1 - \beta$ ; otherwise we could have assigned  $j$  to  $i$  in Phase 2. In either case, we have  $\mathbf{p}(\mathbf{x}_i^2) \geq 1 - \beta$ , and hence  $\mathbf{p}(\mathbf{x}_h) \geq \mathbf{p}(\mathbf{x}_i^2) \geq 1 - \beta$  by choice of  $h$ . Note that  $i$  cannot lose  $j \in \text{child}(i)$  in Phase 3 since Phase 3 only deletes edges from a chore to some of its child agents. Thus,  $\mathbf{p}(\mathbf{x}_h) \geq 1 - \beta$  in this case.
- (ii) No agent in  $N(T)$  lost a child chore. In this case, no agent in  $N(T) \setminus \{i_0\}$  has lost any chore they were earning from in  $(\mathbf{z}, \mathbf{p})$ ; the root agent  $i_0$  could have potentially lost its parent chore  $j_0 = \text{parent}(i_0)$ . We evaluate the amount of earning  $i_0$  loses due to losing  $j_0$ . Suppose  $j_0 \in H$ . Then  $i_0$  must have lost  $j_0$  in either Phase 2 or 3 to some agent  $i' \in \text{child}(j_0)$  since  $i_0$  was not earning the most from  $j_0$  among agents in  $\text{child}(i_0)$ . Due to the earning limit, agents can earn at most  $\beta$  from  $j_0$ . Hence the earning from  $i_0$  from  $j_0$  is at most  $\beta/2$ . On the

other hand, if  $j_0 \in L$ , then  $i_0$  earns at most  $p_{j_0} \leq \beta$  from  $j_0$ . In either case, we find that  $i_0$  has only lost  $\beta$  in earning. Hence the total earning of agents in  $N(T)$  is at least  $|N(T)| - \beta$ , while that from the chores in  $M(T)$  is at most  $\beta \cdot (|N(T)| - 1)$ . Hence there is at least one agent  $i \in N(T)$  whose earning  $\mathbf{p}(\mathbf{x}_i^2)$  satisfies:

$$\mathbf{p}(\mathbf{x}_i^2) \geq \frac{|N(T)| - \beta - \beta \cdot (|N(T)| - 1)}{|N(T)|} = 1 - \beta.$$

Since  $\mathbf{p}(\mathbf{x}_h) \geq \mathbf{p}(\mathbf{x}_i^2)$  by choice of  $h$ , this implies  $\mathbf{p}(\mathbf{x}_h) \geq 1 - \beta$ .  $\square$

To conclude, we established that  $\mathbf{p}(\mathbf{x}_i) \geq \beta$  for the agents  $i$  that are matched to a chore in Phase 4, or  $\mathbf{p}(\mathbf{x}_h) \geq 1 - \beta$  for the agents  $h$  that are not matched. Thus, for all  $i \in N$ ,  $\mathbf{p}(\mathbf{x}_i) \geq \min\{\beta, 1 - \beta\}$ .

Note that the maximum value of the lower bound on agent earnings is given by Lemma 6 is obtained at  $\beta = 1/2$ . We now prove the main theorem of this section.

**Theorem 4.** *Given an ER equilibrium of an instance  $(N, M, D)$  where  $m \geq 2n$ , Algorithm 1 returns a 2-EF2 and fPO allocation in polynomial time. More precisely, for every agent the allocation is either 2-EF1 or EF2.*

*Proof.* Let  $(\mathbf{x}, \mathbf{p})$  be the allocation returned by running Algorithm 1 with an ER equilibrium of the instance with  $\beta = 1/2$ . Lemma 6 then implies that for every agent  $i \in N$ ,  $\mathbf{p}(\mathbf{x}_i) \geq \frac{1}{2}$ .

Lemma 5 implies that for every agent  $h \in N$ , either  $\mathbf{p}_{-1}(\mathbf{x}_h) \leq 1$ , or  $\mathbf{p}_{-2}(\mathbf{x}_h) \leq 1 - \beta = \frac{1}{2}$ . Thus for any agent  $h$ :

- If  $\mathbf{p}_{-1}(\mathbf{x}_h) \leq 1$ , then  $\mathbf{x}$  is 2-EF1 for agent  $h$ , as  $\mathbf{p}_{-1}(\mathbf{x}_h) \leq 1 \leq 2 \cdot \mathbf{p}(\mathbf{x}_i)$ .
- If  $\mathbf{p}_{-2}(\mathbf{x}_h) \leq \frac{1}{2}$ , then  $\mathbf{x}$  is EF2 for agent  $h$ , as  $\mathbf{p}_{-2}(\mathbf{x}_h) \leq \frac{1}{2} \leq \mathbf{p}(\mathbf{x}_i)$ .

Using Lemma 1, this shows that every agent  $h$  is either 2-EF1 or EF2 towards any another agent  $i$ . Overall, the allocation is 2-EF2. Lemma 4 implies  $\mathbf{x}$  is fPO and Lemma 3 shows Algorithm 1 runs in polynomial time.  $\square$

Example 5 shows the tightness of our approach with  $\beta = 1/2$ .

### 3.2 Balanced chore allocation

In this section, we show the existence of an EF2 and fPO allocation when  $m \leq 2n$ . For this, we design Algorithm 2, which computes a balanced fPO allocation for any given instance. Formally, an allocation  $\mathbf{x}$  is balanced iff  $||\mathbf{x}_i| - |\mathbf{x}_h|| \leq 1$  for all agents  $i, h \in N$ , i.e., the sizes of agent bundles differ by at most one. In particular, when  $m \leq 2n$ , every agent in a balanced allocation has at most two chores, so the allocation returned by Algorithm 2 is EF2 and fPO. Similarly, when  $m \leq n$ , Algorithm 2 returns an EF1 and fPO allocation since every agent has at most one chore.

**Definition 6** (MPB graph). *The MPB graph  $G = (N, M, E)$  associated with an integral MPB allocation  $(\mathbf{x}, \mathbf{p})$  is a directed bipartite graph with vertex set  $V(G) = N \sqcup M$ , and edge set  $E(G) = \{(i, j) : i \in N, j \in M, j \in \mathbf{x}_i\} \cup \{(j, i) : i \in N, j \in M, j \in MPB_i \setminus \mathbf{x}_i\}$ .*

Algorithm 2 first allocates the entire set of chores  $M$  to an arbitrary agent  $h$ , with the payment of each chore  $j$  set as  $p_j = d_{hj}$  to ensure the initial allocation is MPB. We make progress towards achieving a balanced allocation by reducing the number of chores assigned to  $h$ , the agent with the most chores, and increasing the number of chores assigned to  $\ell$ , the agent with the fewest chores.

---

**Algorithm 2** Balanced PO allocation

---

**Input:** Chore allocation instance  $(N, M, D)$ **Output:** An balanced PO allocation  $\mathbf{x}$ 

```
1: For some agent  $h \in N$ , set  $\mathbf{x}_h \leftarrow M$ ,  $\mathbf{x}_i \leftarrow \emptyset$  for  $i \neq h$ 
2: For each  $j \in M$ , set  $p_j \leftarrow d_{hj}$ 
3:  $\ell \leftarrow \arg \min_{i \in N} |\mathbf{x}_i|$  ▷ Agent with fewest number of chores
4: while  $|\mathbf{x}_h| > |\mathbf{x}_\ell| + 1$  do
5:    $C_h \leftarrow$  Vertices reachable from  $h$  in the MPB graph of  $(\mathbf{x}, \mathbf{p})$ 
6:   if  $\ell \in C_h$  then ▷ Transfer chores along a path in the MPB graph
7:      $P \leftarrow (h = i_0, j_1, i_1, j_2, \dots, i_{k-1}, j_k, i_k = \ell)$  ▷ Path from  $h$  to  $\ell$ 
8:     for  $1 \leq r \leq k$  do
9:        $\mathbf{x}_{i_{r-1}} \leftarrow \mathbf{x}_{i_{r-1}} \setminus \{j_r\}$ ,  $\mathbf{x}_{i_r} \leftarrow \mathbf{x}_{i_r} \cup \{j_r\}$  ▷ Chore transfers along path  $P$ 
10:     $\ell \leftarrow \arg \min_{i \in N} |\mathbf{x}_i|$ 
11:   else ▷ Raise payments of chores in  $C_h$ 
12:      $\gamma \leftarrow \min_{i \in N \setminus C_h, j \in C_h \cap M} \frac{\alpha_i}{d_{ij}/p_j}$ 
13:     for  $j \in C_h$  do
14:        $p_j \leftarrow \gamma \cdot p_j$ 
15: return  $\mathbf{x}$ 
```

---

Let  $C_h$  denote the set of vertices in the MPB graph reachable from  $h$ . If  $\ell \in C_h$ , i.e.,  $\ell$  is reachable from  $h$  in the MPB graph via a path  $P = (h = i_0, j_1, i_1, j_2, \dots, j_{k-1}, i_{k-1}, j_k, i_k = \ell)$ , then chore  $j_r$  is transferred from  $i_{r-1}$  to  $i_r$ , for all  $r \in [k]$ . The result of such transfers is that  $h$  has one fewer chore,  $\ell$  has one more chore, and all other agents maintain the same number of chores. Observe that since  $j_r \in MPB_{i_r}$ , the allocation after the transfers is also MPB. If  $\ell \notin C_h$ , then we uniformly raise the payments of all chores in  $C_h$ , until a chore  $j \in C_h$  becomes MPB for an agent  $i \notin C_h$ . Such a payment raise respects the MPB condition for all agents, hence the resulting allocation remains MPB. We repeat payment raises until  $\ell$  is reachable from  $h$ . After this, we transfer a chore from  $h$  to  $\ell$  along a path in the MPB graph, and repeat this process until the allocation is balanced. Algorithm 2 runs in polynomial time, as there are at most  $m$  transfers, and there are at most  $n$  payment raises between two transfers.

**Theorem 5.** *For any instance  $(N, M, D)$ , Algorithm 2 returns a balanced fPO allocation  $\mathbf{x}$  in polynomial time. In particular,  $\mathbf{x}$  is EF2 and fPO when  $m \leq 2n$ , and EF1 and fPO when  $m \leq n$ .*

*Proof.* Let  $(\mathbf{x}, \mathbf{p})$  be an allocation in the run of Algorithm 2 before termination. We claim that the allocation excluding  $\mathbf{x}_h$  is always a balanced allocation, i.e., for  $i_1, i_2 \in N \setminus h$ ,  $\|\mathbf{x}_{i_1}\| - \|\mathbf{x}_{i_2}\| \leq 1$ . We prove this by induction. Indeed, the initial allocation excluding  $h$  is trivially balanced. A chore transfer step from  $h$  to  $\ell$  along a path  $P$  only changes the number of chores assigned to  $h$  and  $\ell$  and no other agents in  $P$ . Since  $\ell$  has the fewest chores and gains only one additional chore, the allocation excluding  $h$  remains balanced. Since the payment raise step does not change the allocation, the claim holds by induction. We next show that while the allocation  $\mathbf{x}$  is not balanced,  $h$  is the unique agent with the highest number of chores. Since Algorithm 2 has not terminated,  $|\mathbf{x}_h| > |\mathbf{x}_\ell| + 1$ . Since the allocation excluding  $h$  is balanced, we have for  $i \in N \setminus \{h, \ell\}$  that  $|\mathbf{x}_i| - |\mathbf{x}_\ell| \leq 1$ , implying that  $|\mathbf{x}_i| \leq |\mathbf{x}_\ell| + 1$ . This shows  $|\mathbf{x}_i| < |\mathbf{x}_h|$ .

We can now complete the proof for the termination of Algorithm 2. In any chore transfer step, agent  $h$  loses exactly one chore, so  $|\mathbf{x}_h|$  decreases by one. Additionally,  $|\mathbf{x}_h|$  does not change in any payment raise iteration. Algorithm 2 therefore terminates after at most  $m$  chore transfer steps.

We claim that there can be at most  $n$  payment raise iterations between chore transfer iterations. A payment raise step results in an agent  $i \notin C_h$  getting added to  $C_h$ , and does not remove any agents from  $C_h$ . Thus, after at most  $n$  payment raise steps it must be that  $\ell$  is reachable from  $h$ , and Algorithm 2 performs a chore transfer step. Thus, Algorithm 2 terminates after at most  $mn + m$  iterations. On termination with  $\mathbf{x}$ , it must be that  $|\mathbf{x}_h| - |\mathbf{x}_\ell| \leq 1$ , implying that the final allocation is balanced. In particular, if  $m \leq 2n$ , each agent has at most two chores and  $\mathbf{x}$  is EF2. Likewise, if  $m \leq n$ , each agent has at most one chore and  $\mathbf{x}$  is EF1.

Since the initial allocation is MPB, and every transfer step and payment raise preserves the MPB condition, the resulting allocation is MPB as well. Thus, Proposition 1 shows that the final allocation is fPO.  $\square$

### 3.3 Approximate-EF1 and PO

We now turn to the existence of approximately-EF1 and PO allocations. The main result of this section is:

**Theorem 6.** *Any chore allocation instance with  $n$  agents admits an  $(n-1)$ -EF1 and fPO allocation.*

First, we observe from Theorem 5 that if  $m \leq n$ , an EF1 and fPO allocation can be computed in polynomial time using Algorithm 2. Hence we assume  $m \geq n$  in the remainder of the section.

Following the ideas developed in Section 3.1 which rounds a fractional solution of an ER equilibrium, the natural approach towards obtaining an approximate-EF1 guarantee is to ensure that every agent gets at most one overpaying chore in the rounded solution. Clearly, this requires the number of overpaying chores to be at most  $n$ , which cannot be guaranteed for earning limit  $\beta < 1$ . However, Lemma 6 does not show good lower bounds on the agent earnings when  $\beta = 1$ .

To fix this, we design Algorithm 6 (pseudocode in Appendix C) by modifying the rounding procedure of Algorithm 1. Since  $m \geq n$ , an ER equilibrium with  $\beta = 1$  exists. Given such an equilibrium  $(\mathbf{y}, \mathbf{p})$ , Algorithm 6 defines  $L$  to be the set of chores with payment at most  $\beta/2 = 1/2$ , and  $H$  to be the set of chores with payment exceeding  $1/2$ . We prove that by using the same rounding procedure but with  $L$  and  $H$  defined this way, we obtain an integral MPB allocation  $(\mathbf{x}, \mathbf{p})$  where  $\mathbf{p}_{-1}(\mathbf{x}_i) \leq 1$  and  $\mathbf{p}(\mathbf{x}_i) \geq \frac{1}{2(n-1)}$  for all agents  $i \in N$ . Using Lemma 1, this implies that  $\mathbf{x}$  is  $2(n-1)$ -EF1 and fPO.

**Theorem 7.** *Given an ER equilibrium for an instance  $(N, M, D)$  where  $m \geq n$ , Algorithm 6 returns a  $2(n-1)$ -EF1 and fPO allocation in polynomial time.*

Algorithm 6 and its analysis is presented in Appendix C. In Example 4, we show a lower bound on our approach with  $\beta = 1$  by presenting an instance for which no rounding of the ER equilibrium is  $(n-1-\delta)$ -EF1, for any  $\delta > 0$ . In Appendix C, we close this gap in the approximation factor by designing an improved algorithm.

**Theorem 8.** *Given an ER equilibrium of an instance with  $m \geq n$ , an  $(n-1)$ -EF1 and fPO allocation can be found in polynomial time.*

## 4 Existence of 4-EFX Allocations

In this section, we prove our main result on approximately-EFX allocations of chores.

**Theorem 9.** *Any chore allocation instance admits a 4-EFX allocation.*



---

**Algorithm 3** Computes an EFX allocation for instances with  $m \leq 2n$

---

**Input:** Instance  $(N, M, D)$  with  $m \leq 2n$

**Output:** An integral allocation  $\mathbf{x}$

```

1:  $r \leftarrow \max\{0, m - n\}$ 
2:  $M' \leftarrow M, \mathbf{x}_i \leftarrow \emptyset$  for all  $i \in [n]$ 
   — Phase 1: Agents in  $N_2$  pick chores in order from  $r$  to 1 —
3: for  $i = r$  down to 1 do
4:    $e_i \leftarrow \arg \min_{j \in M'} d_i(j)$ 
5:    $\mathbf{x}_i \leftarrow \mathbf{x}_i \cup \{e_i\}, M' \leftarrow M' \setminus \{e_i\}$ 
   — Phase 2: Agents pick chores in order from 1 to  $n$  —
6: for  $i = 1$  to  $n$  do
7:    $j_i \leftarrow \arg \min_{j \in M'} d_i(j)$ 
8:    $\mathbf{x}_i \leftarrow \mathbf{x}_i \cup \{j_i\}, M' \leftarrow M' \setminus \{j_i\}$ 
   — Phase 3: Chore swaps —
9: while  $\mathbf{x}$  is not EFX do
10:   $i \leftarrow \arg \min\{i' \in N : i' \text{ is not EFX}\}$ 
11:   $\ell \leftarrow \arg \min\{d_i(\mathbf{x}_h) : h \in N\}$ 
12:  Perform  $(i, \ell)$  swap:  $\mathbf{x}_i \leftarrow \mathbf{x}_i \cup \mathbf{x}_\ell \setminus \{j_i\}, \mathbf{x}_\ell \leftarrow \{j_i\}$ 
13: return  $\mathbf{x}$ 

```

---

We first show that when  $m \leq 2n$ , an EFX allocation exists and can be computed in polynomial time. Note that an  $O(n^3)$ -time algorithm that uses matching based techniques to compute an EFX allocation is known from prior work [29]. We design an alternative algorithm (Algorithm 3) which is faster (runs in  $O(n^2)$  time), and arguably simpler. Moreover, our algorithm returns an EFX allocation with certain special properties and introduces the idea of chore swaps, both of which are important in the general case.

**Theorem 10.** *For a chore allocation instance with  $n$  agents ordered 1 through  $n$ , and  $m \leq 2n$  chores, Algorithm 3 returns in  $O(n^2)$  time an allocation  $\mathbf{x}$  s.t.*

(i)  $\mathbf{x}$  is EFX.

(ii) If  $m > n$ , then  $|\mathbf{x}_i| = 1$  for all  $i > m - n$ .

With the above result, it only remains to establish Theorem 11 for instances with  $m \geq 2n$ . For this setting, we design a polynomial time algorithm Algorithm 4, which computes a 4-EFX allocation for a chore allocation instance with  $m \geq 2n$ , given its ER equilibrium with earning limit  $\beta = 1/2$ . Algorithm 4 uses Algorithm 3 as a subroutine and crucially relies on the properties outlined in Theorem 10.

**Theorem 11.** *Given an ER equilibrium of a chore allocation instance with  $m \geq 2n$  and earning limit  $\beta = 1/2$ , Algorithm 4 returns a 4-EFX allocation in polynomial time.*

Theorems 10 and 11 together prove Theorem 9, and are proved in the following sections.

#### 4.1 EFX for $m \leq 2n$

This section proves Theorem 10 by describing and analyzing Algorithm 3, which computes an EFX allocation for instances with  $m \leq 2n$ . Algorithm 3 iteratively allocates chores to agents in three

phases. We initialize  $M'$  to  $M$  and update this set as chores are allocated. For simplicity, we first assume  $m > n$ , letting  $r = m - n$ .

In Phase 1, proceeding in the order  $r, r - 1, \dots, 1$ , each agent  $i$  iteratively picks their least disutility chore  $e_i$  among the set of remaining items  $M'$ . Let  $L = \{e_1, \dots, e_r\}$ . Then in Phase 2, proceeding in the order 1 to  $n$ , each agent  $i$  picks their least disutility chore  $j_i$  in  $M'$ . Let  $H = \{j_1, \dots, j_n\}$ . Let  $\mathbf{x}^0$  be the allocation at the end of Phase 2. Let  $N_2 = [r]$  be the set of agents with two chores in  $\mathbf{x}^0$ , and let  $N_1 = [n] \setminus [r]$  be the set of agents with one chore in  $\mathbf{x}^0$ . Clearly, agents in  $N_1$  are EFX, and hence only agents in  $N_2$  may be EFX-envious.

Starting with  $\mathbf{x}^0$ , Phase 3 of Algorithm 3 performs *chore swaps* between an agent  $i \in N_2$  who is not EFX, and the agent  $\ell$  who  $i$  envies the most. We refer to such a swap as an  $(i, \ell)$ -swap. In an  $(i, \ell)$  swap in an allocation  $\mathbf{x}$ , the bundle  $\mathbf{x}_\ell$  is transferred to  $i$ , and the higher disutility chore  $j_i$  is transferred from  $i$  to  $\ell$ . When there are multiple envious agents  $i$ , we break ties following the agent ordering. We argue that after an  $(i, \ell)$  swap, the agent  $i$  becomes EFX and remains EFX throughout the subsequent execution of the algorithm. Thus, every agent  $i \in N_2$  undergoes an  $(i, \ell)$  swap at most once, and these swaps happen in the order of agents 1 through  $r$ . This implies that Algorithm 3 terminates in at most  $r$  steps with an EFX allocation.

Finally, we note that when  $m \leq n$ , i.e.,  $r = 0$ , Algorithm 3 only executes Phase 2 and returns an allocation in which each agent gets a single chore, and hence is EFX.

**Analysis of Algorithm 3.** Since it is clear that Algorithm 3 returns an EFX allocation when  $m \leq n$ , we assume  $m > n$  in the following analysis. Purely for the purpose of analysis, we implement Phase 3 as follows:

---

```

9: for  $i = 1$  to  $r$  do
10:   if  $i$  is not EFX then
11:      $\ell \leftarrow \arg \min \{d_i(\mathbf{x}_h) : h \in N\}$ 
12:     Perform  $(i, \ell)$  swap:  $\mathbf{x}_i \leftarrow \mathbf{x}_i \cup \mathbf{x}_\ell \setminus \{j_i\}$ ,  $\mathbf{x}_\ell \leftarrow \{j_i\}$ 

```

---

Lemma 7 below refers to the above implementation of Phase 3 of Algorithm 3. Let  $\mathbf{x}^i$  denote the allocation after iteration  $i$  of Phase 3.

**Lemma 7.** *For each  $i \in [r]$ ,*

- (i) *Before iteration  $i$ , agents  $N \setminus [i - 1]$  do not participate in any swap.*
- (ii) *In iteration  $i$ , if agent  $i$  participates in an  $(i, \ell)$  swap, then  $i$  is EFX after the swap. Moreover,  $\max_{j \in \mathbf{x}_i^i} d_i(\mathbf{x}_i^i \setminus \{j\}) < d_i(j_i)$  immediately after the swap.*
- (iii) *After iteration  $i$ , agents in  $N_2 \cap [i]$  are EFX. Agents in  $N_1$  have a single chore, and are EFX.*

*Proof.* We prove the invariants inductively, beginning with  $i = 1$ . Consider the allocation  $\mathbf{x}^0$  before Phase 3 begins. We show that the invariants hold for  $i = 1$  as follows.

- (i) Invariant (i) holds trivially, since no agent has participated in any swap before iteration 1.
- (ii) Note that  $\mathbf{x}_1^0 = \{e_1, j_1\}$  and  $j_h \in \mathbf{x}_h^0$  for any  $h \geq 2$  in the allocation  $\mathbf{x}^0$  before iteration 1. By the order in which agent 1 picks chores, we have  $d_1(e_1) \leq d_1(j_1) \leq d_1(j_h)$  for any  $h \geq 2$ . Hence,  $\max_{j \in \mathbf{x}_1^0} d_1(\mathbf{x}_1^0 \setminus \{j\}) = d_1(j_1) \leq d_1(j_h) \leq d_1(\mathbf{x}_h^0)$ . This shows that agent 1 is EFX in the allocation  $\mathbf{x}^0$ , and hence no swap takes place in iteration 1. Invariant (ii) thus holds vacuously.

- (iii) As argued above, agent 1 is EFX in  $\mathbf{x}^0$ , and since no swap takes place in iteration 1, we have  $\mathbf{x}^1 = \mathbf{x}^0$ . In  $\mathbf{x}^1$ , agent 1 is EFX, and agents in  $N_1$  have a single chore, and hence are EFX. Thus, invariant (iii) holds.

Assume that the invariants (i)-(iii) holds for some  $i \in [r-1]$ . We will prove that the invariants hold for  $i+1$  as well.

- (i) Let  $i$  and  $\ell$  be the agents participating in an  $(i, \ell)$  swap in iteration  $i$ . By invariant (i) of the inductive hypothesis, agents  $\{i, i+1, \dots, n\}$  have not undergone a swap before iteration  $i$ . Hence the allocation before iteration  $i$  satisfies  $\mathbf{x}_h^{i-1} = \mathbf{x}^0$  for any  $h \in N_1 \cup N_2 \setminus [i-1]$ . By the order in which agent  $i$  picks chores, we have:

$$\max_{j \in \mathbf{x}_i^{i-1}} d_i(\mathbf{x}_1^{i-1} \setminus \{j\}) = d_i(j_i) \leq d_i(j_h) \leq d_i(\mathbf{x}_h^{i-1}),$$

which shows that agent  $i$  does not EFX-envy any agent in  $\{i+1, \dots, n\}$ . Since  $i$  is EFX-envious before iteration  $i$ , we must have  $\ell \in [i-1]$ . Thus, after iteration  $i$ , only agents in  $[i]$  have participated in swaps, establishing invariant (i).

- (ii) Suppose agent  $(i+1)$  participates in an  $(i+1, k)$  swap with agent  $k$  in iteration  $(i+1)$  resulting in the allocation  $\mathbf{x}^{i+1}$ . Before iteration  $(i+1)$ , we know from invariant (i) that agents in  $N \setminus [i]$  have not participated in any swap. Thus,  $\mathbf{x}_{i+1}^i = \{e_{i+1}, j_{i+1}\}$ . Using the fact that agent  $(i+1)$  is not EFX in the allocation  $\mathbf{x}^i$ , and the choice of agent  $k$ , we have  $d_{i+1}(j_{i+1}) > d_{i+1}(\mathbf{x}_k^i)$ . Once again, using the order in which agent  $(i+1)$  picked chores, we see that  $k \in [i]$ .

Next, we claim that  $d_{i+1}(e_i) \leq d_{i+1}(j)$  for any  $j \in \mathbf{x}_k^i$ . To see this, note that by the order in which agent  $(i+1)$  picks chores, the only chores that have disutility less than  $d_{i+1}(e_i)$  for agent  $(i+1)$  could be the chores  $e_{i+2}, \dots, e_r$ . However, since agents  $(i+1), \dots, r$  have not undergone any swap step, these chores cannot belong to  $\mathbf{x}_k^i$ . The claim thus holds.

We now prove invariant (ii). After iteration  $(i+1)$ , we have  $\mathbf{x}_{i+1}^{i+1} = \{e_{i+1}\} \cup \mathbf{x}_k^i$ , and  $\mathbf{x}_k^{i+1} = \{j_{i+1}\}$ . Observe that:

$$\max_{j \in \mathbf{x}_{i+1}^{i+1}} d_{i+1}(\mathbf{x}_{i+1}^{i+1} \setminus \{j\}) = d_{i+1}(\mathbf{x}_k^i) < d_{i+1}(j_{i+1}),$$

where we use the claim that  $d_{i+1}(e_i) \leq d_{i+1}(j)$  for any  $j \in \mathbf{x}_k^i$  in the first equality, and the fact that agent  $(i+1)$  EFX-envies agent  $k$  in the allocation  $\mathbf{x}^i$  in the second inequality. This proves the second claim of invariant (ii) and also shows that agent  $(i+1)$  does EFX-envy agent  $k$  immediately after the swap. Consider some other agent  $h \notin \{i+1, k\}$ . By the choice of agent  $k$ , we have  $d_{i+1}(\mathbf{x}_k^i) \leq d_{i+1}(\mathbf{x}_h^i)$ . Thus,  $\max_{j \in \mathbf{x}_{i+1}^{i+1}} d_{i+1}(\mathbf{x}_{i+1}^{i+1} \setminus \{j\}) = d_{i+1}(\mathbf{x}_k^i) \leq d_{i+1}(\mathbf{x}_h^i)$ . Thus, agent  $(i+1)$  is EFX after the swap in iteration  $(i+1)$ .

- (iii) Consider the allocation  $\mathbf{x}^{i+1}$  after iteration  $(i+1)$ . Invariant (ii) shows that agent  $(i+1)$  is EFX in  $\mathbf{x}^{i+1}$ . Moreover, agent  $k$  is EFX in  $\mathbf{x}^{i+1}$  since she has a single chore. Since we argued above that  $k \in [i]$ , the agents in  $N_1$  continue to have a single chore and are EFX.

Thus, it only remains to be shown that an agent  $h \in [i] \setminus \{k\}$  who was EFX in the allocation  $\mathbf{x}^i$  remains EFX in the allocation  $\mathbf{x}^{i+1}$  after the  $(i+1, k)$  swap. First note that  $h$  is EFX towards  $\mathbf{x}_{h'}^{i+1}$  for any  $h' \notin \{i+1, k\}$ , since  $\mathbf{x}_{h'}^i = \mathbf{x}_{h'}^{i+1}$  and  $h$  is EFX in  $\mathbf{x}^i$ .

Next, observe that agent  $h$  is EFX towards the bundle  $\mathbf{x}_k^i$ . Since  $\mathbf{x}_k^i \subset \mathbf{x}_{i+1}^{i+1}$ , agent  $h$  is EFX towards  $\mathbf{x}_{i+1}^{i+1}$  as well.

Finally, we show that  $h$  is EFX towards the bundle  $\mathbf{x}_k^{i+1} = \{j_{i+1}\}$ . If agent  $h$  underwent a swap of the form  $(i', h)$  during iteration  $i' \in [h+1, i+1]$ , then agent  $h$  has a single chore and will be EFX in  $\mathbf{x}^{i+1}$ . Hence we assume agent  $h$  did not undergo any swap during iterations  $[h+1, i+1]$ , and hence  $\mathbf{x}_h^{i+1} = \mathbf{x}_h^h$ . Now, observe that:

$$\max_{j \in \mathbf{x}_h^h} d_h(\mathbf{x}_h^h \setminus \{j\}) \leq d_h(j_h).$$

This is true, because if  $h$  does not undergo a swap in iteration  $h$ , we have  $\mathbf{x}_h^h = \{e_h, j_h\}$ , and  $d_h(e_h) \leq d_h(j_h)$ . If  $h$  does undergo a swap in iteration  $h$ , invariant (ii) implies the same observation. Finally, by the order in which  $h$  picked chores, we have  $d_h(j_h) \leq d_h(j_{i+1})$ . Using  $\mathbf{x}_h^{i+1} = \mathbf{x}_h^h$ , we obtain:

$$\max_{j \in \mathbf{x}_h^{i+1}} d_h(\mathbf{x}_h^{i+1} \setminus \{j\}) = \max_{j \in \mathbf{x}_h^h} d_h(\mathbf{x}_h^h \setminus \{j\}) \leq d_h(j_h) \leq d_h(j_{i+1}) = d_h(\mathbf{x}_k^{i+1}),$$

which shows that  $h$  is EFX towards the bundle  $\mathbf{x}_k^{i+1}$ . In conclusion, invariant (iii) holds.

By induction, the invariants hold for all  $i \in [r]$ . □

With Lemma 7 in hand, Theorem 10 follows easily.

*Proof of Theorem 10.* For  $m \leq n$ , Theorem 10 only executes Phase 2 which assigns a single chore to each agent, thus returning an EFX. For  $m > n$ , Invariant (iii) of Lemma 7 for  $i = r$  implies that in the allocation returned by Algorithm 3, agents in  $N_2$  are EFX. Moreover, agents in  $N_1 = [n] \setminus [r]$  have a single chore and hence are EFX. This proves both properties (1) and (2) claimed by Theorem 10.

Finally, note that Algorithm 3 runs in  $O(n^2)$ -time: Phases 1 and 2 involve  $m \leq 2n$  steps of identifying an agent's favorite chore ( $O(m)$  time each), and Phase 3 involves at most  $r \leq n$  swap steps. □

## 4.2 4-EFX allocation: Algorithm overview

We now prove the existence of 4-EFX allocations for chore allocation instances with  $m \geq 2n$ . We design a polynomial time algorithm, Algorithm 4, which returns a 4-EFX allocation for an instance when given an ER equilibrium of the instance with earning limit  $\beta = 1/2$  as input.

Algorithm 4 first runs Algorithm 1 on the given ER equilibrium  $(\mathbf{y}, \mathbf{p})$  with earning limit  $\beta = 1/2$  to obtain a 2-EF2 and fPO allocation  $(\mathbf{x}, \mathbf{p})$ . As in Section 3, we classify chores based on their payments as  $L = \{j \in M : p_j \leq 1/2\}$  and  $H = \{j \in M : p_j > 1/2\}$ . Thus  $L$  is the set of low cost chores, and  $H$  is the set of high cost or *overpaying* chores, whose payment exceeds the earning limit  $\beta = 1/2$ .

We partition the bundle of each agent  $i$  as  $\mathbf{x}_i = S_i \cup H_i$ , where  $S_i \subseteq L$  and  $H_i \subseteq H$ . Let  $N_H$  denote the set of agents who are assigned overpaying chores, and let  $N_0 = N \setminus N_H$ . The following lemma records properties of the allocation  $\mathbf{x}$ .

**Lemma 8.** *The allocation  $(\mathbf{x}, \mathbf{p})$  satisfies:*

(i) For any  $i \in N$ :  $\mathbf{p}(\mathbf{x}_i) = \mathbf{p}(S_i \cup H_i) \geq 1/2$ .

(ii) For  $i \in N_H$ :  $\mathbf{p}(S_i) \leq 1$ .

(iii) For  $i \in N_0$ :  $\mathbf{p}(\mathbf{x}_i) = \mathbf{p}(S_i) \leq 3/2$ .

---

**Algorithm 4** Computes a 4-EFX allocation

---

**Input:** Instance  $(N, M, D)$  with  $m \geq 2n$  and its ER equilibrium  $(\mathbf{y}, \mathbf{p})$  with  $\beta = 1/2$

**Output:** An integral allocation  $\mathbf{x}$

- 1:  $(\mathbf{x}, \mathbf{p}) \leftarrow$  Algorithm 1( $\mathbf{y}, \mathbf{p}$ )
  - 2:  $L = \{j \in M : p_j \leq 1/2\}$ ,  $H = \{j \in M : p_j > 1/2\}$  ▷ Low, High paying chores
  - 3: Partition each  $\mathbf{x}_i = S_i \sqcup H_i$ , where  $S_i \subseteq L$  and  $H_i \subseteq H$
  - 4:  $N_H \leftarrow \{i \in N : H_i \neq \emptyset\}$ , and  $N_0 \leftarrow \{i \in N : H_i = \emptyset\}$ .
  - 5: Re-order agents s.t. agents in  $N_H$  are ordered before agents in  $N_0$
  - 6:  $\mathbf{z}' \leftarrow$  EFX allocation of  $H$  to  $N$  using Algorithm 3, with agents ordered as above
  - 7: Partition agents into  $N_L, N_H^1, N_H^2$  using Definition 8
  - 8:  $H' \leftarrow \cup_{i \in N_H^1} \mathbf{z}'_i$
  - 9:  $(\mathbf{z}, \mathbf{q}) \leftarrow$  Min cost matching of  $H'$  to  $N_H^1$ , and associated dual variables
  - 10: For each  $i \in N_H^1$ ,  $\mathbf{x}_i \leftarrow S_i \cup \mathbf{z}_i$
  - 11: For each  $i \notin N_H^1$ ,  $\mathbf{x}_i \leftarrow S_i \cup \mathbf{z}'_i$
  - 12: **while**  $\exists i \in N_H^1$  not 4-EFX **do**
  - 13:      $i \leftarrow \arg \min\{\mathbf{q}(\mathbf{z}_h) : h \in N_H^1 \text{ not 4-EFX}\}$
  - 14:      $\ell \leftarrow \arg \min\{d_i(\mathbf{x}_h) : h \in N\}$
  - 15:     Perform  $(i, \ell)$  swap:  $\mathbf{x}_i \leftarrow \mathbf{x}_i \cup \mathbf{x}_\ell \setminus \mathbf{z}_i$ ,  $\mathbf{x}_\ell \leftarrow \mathbf{z}_i$
  - 16: **return**  $\mathbf{x}$
- 

*Proof.* For  $\beta = 1/2$ , Lemmas 5 and 6 imply that for all agents  $i \in N$ ,  $(\mathbf{x}, \mathbf{p})$  satisfies  $\mathbf{p}(\mathbf{x}_i) \geq 1/2$ , and either  $\mathbf{p}_{-1}(\mathbf{x}_i) \leq 1$ , or  $|\mathbf{x}_i \cap H| = 2$  and  $\mathbf{p}_{-2}(\mathbf{x}_i) \leq 1/2$ . The lemma then follows from the definitions of the partition of agents  $N = N_H \sqcup N_0$  and the chores bundles  $\mathbf{x}_i = S_i \sqcup H_i$ .  $\square$

Notice that the above lemma implies that agents in  $N_0$  are 3-EFX. Thus,  $\mathbf{x}$  may not be  $O(1)$ -EFX only because of agents in  $N_H$ , who are assigned one or more overpaying chores. Therefore, our algorithm must address the  $O(1)$ -EFX envy of agents in  $N_H$ .

**Chore swaps.** For simplicity, let us assume that all agents in  $N_H$  are assigned a single overpaying chore. Consider an agent  $i$  with  $H_i = \{j_i\}$ , who is not 4-EFX in  $\mathbf{x}$ . To fix the 4-EFX envy that  $i$  has towards other agents, we re-introduce the idea of a ‘chore swap’:

**Definition 7.** Consider an allocation  $\mathbf{x}$  in which an agent  $i \in N_H$  is not 4-EFX. Let  $j_i$  be the overpaying chore in  $\mathbf{x}_i$ . Let  $\ell$  be the agent who  $i$  envies the most, i.e.  $\ell = \arg \min\{h \in N : d_i(\mathbf{x}_h)\}$ . An  $(i, \ell)$  swap on the allocation  $\mathbf{x}$  results in an allocation  $\mathbf{x}'$  obtained by transferring all the chores of  $\ell$  to  $i$ , and transferring the chore  $j_i$  from  $i$  to  $\ell$ . That is,  $\mathbf{x}'_i = \mathbf{x}_i \cup \mathbf{x}_\ell \setminus \{j_i\}$ ,  $\mathbf{x}_\ell = \{j_i\}$ , and  $\mathbf{x}'_h = \mathbf{x}_h$  for all  $h \neq \{i, \ell\}$ .

Note the similarity to the definition of chore swaps involved in Algorithm 3. Similar to the analysis in Lemma 40, we claim that immediately after the  $(i, \ell)$  swap,  $i$  is 4-EFX towards all agents. To see this, let us scale the disutility function of each agent so that every agent has MPB ratio 1. This allows us to measure payments and disutilities on the same scale. Let  $\mathbf{x}'$  be the allocation resulting from an  $(i, \ell)$  swap on  $\mathbf{x}$ . Note that  $\mathbf{x}'_i = \mathbf{x}_i \cup \mathbf{x}_\ell \setminus \{j_i\} = S_i \cup \mathbf{x}_\ell$ , and  $\mathbf{x}'_\ell = \{j_i\}$ , while  $\mathbf{x}'_h = \mathbf{x}_h$  for all  $h \neq \{i, \ell\}$ . Since  $\mathbf{p}(S_i) \leq 1$  and  $\mathbf{p}(\mathbf{x}_\ell) \geq 1/2$ , we have that  $d_i(\mathbf{x}'_i) \leq 3 \cdot d_i(\mathbf{x}_\ell)$ . By the choice of  $\ell$ , for any  $h \neq \{i, \ell\}$ ,  $d_i(\mathbf{x}_\ell) \leq d_i(\mathbf{x}_h) = d_i(\mathbf{x}'_h)$ . Thus  $d_i(\mathbf{x}'_i) \leq 3 \cdot d_i(\mathbf{x}'_h)$ , showing that  $i$  does not 3-EFX envy (and hence 4-EFX envy) agent  $h$  after the swap. Similarly, the fact that  $i$  is 4-EFX envious of the bundle  $\mathbf{x}_\ell$  establishes a lower bound on the disutility of  $j_i$  for  $i$ ,

which we can use to prove that  $i$  will not 4-EFX envy  $\mathbf{x}'_\ell = \{j_i\}$  after the swap. Moreover, agent  $\ell$  is EFX after the swap since she has a single chore.

In conclusion, after an  $(i, \ell)$  chore swap, both agents  $i$  and  $\ell$  are 4-EFX, i.e., the  $O(1)$ -EFX envy of agent  $i$  is temporarily resolved. The above idea suggests repeatedly performing chore swaps until the allocation is  $O(1)$ -EFX. However, two things remain unclear: (i) how to address agents in  $N_H$  with two overpaying chores, and (ii) whether an agent  $i$  who underwent a swap develops EFX-envy subsequently in the run of the algorithm. Our algorithm addresses both these issues by separately treating overpaying chores and the agents to whom they are assigned, and using clever design choices.

**Re-allocating overpaying chores.** Observe that  $\mathbf{p}(S_i) \leq O(1) \cdot p_j$  for any agent  $i$  and overpaying chore  $j \in H$ . This means that for any agent  $i$ , the chores in  $S_i$  have cumulatively less than *any* single overpaying chore, up to a constant factor. Thus we should ‘balance’ out the envy created among the agents due to an imbalanced allocation of the overpaying chores. To do this, we compute an EFX allocation  $\mathbf{z}'$  of the overpaying chores  $H$  using Algorithm 3. This is possible since there are at most  $2n$  overpaying chores, i.e.,  $|H| \leq 2n$  as each agent has at most two overpaying chores in the rounded allocation. In our invocation of Algorithm 3, we order the agents in  $N_H$  to appear before the agents in  $N_0$ .

We then classify the agents based on the EFX allocation  $\mathbf{z}'$  as follows:

**Definition 8.** (*Classification of Agents in Algorithm 4*) Agents are classified as:

- $N_L = \{i \in N : |\mathbf{z}'_i| = 0\}$ , i.e. agents with no overpaying chores.
- $N_H^1 = \{i \in N : |\mathbf{z}'_i| = 1\}$ , i.e. agents with a single overpaying chore.
- $N_H^2 = \{i \in N : |\mathbf{z}'_i| \geq 2\}$ , i.e. agents with at least two overpaying chores.

Note that agents in  $N_H^2$  can have more than two overpaying chores since we re-allocated the overpaying chores via the EFX allocation  $\mathbf{z}'$ . However, since  $\mathbf{z}'$  is EFX, it cannot be that both  $N_H^2 \neq \emptyset$  and  $N_L \neq \emptyset$ .

Additionally, by ordering the agents in  $N_H$  before agents in  $N_0$ , we can leverage Theorem 10 to show that  $N_H^2 \subseteq N_H$ . That is, if an agent  $i$  obtains two or more overpaying chores after re-allocating the overpaying chores, then  $i$  must have had an overpaying chore to begin with. Recall from Lemma 8 that the earning of such agents is at most 1 from chores in  $L$ , i.e., that  $\mathbf{p}(S_i) \leq 1$  for each such agent  $i \in N_H$ . This property is useful to bound the total EFX-envy of agents in  $N_H^2$ .

Having re-allocated the overpaying chores  $H$ , we add back the chores from  $L$  to obtain the allocation  $\mathbf{x}'$  given by  $\mathbf{x}'_i = S_i \cup \mathbf{z}'_i$  for all agents  $i$ . We note that each agent  $i \in N_H^2$  is actually  $O(1)$ -EFX in  $\mathbf{x}'$ : since  $\mathbf{p}(S_i) \leq O(1) \cdot \mathbf{p}(\mathbf{z}'_i)$ , we have that  $d_i(\mathbf{x}'_i) \leq O(1) \cdot d_i(\mathbf{z}'_i)$ . Since  $\mathbf{z}'$  is EFX and  $i$  has at least two overpaying chores, we have that  $d_i(\mathbf{z}'_i) \leq 2 \cdot d_i(\mathbf{z}'_k)$  for all  $k$ . Thus  $d_i(\mathbf{x}'_i) \leq O(1) \cdot d_i(\mathbf{x}'_k)$ . This proves a surprising property of the allocation  $\mathbf{x}'$ : the agents in  $N_H^2$  who have two or more overpaying chores are actually  $O(1)$ -EFX! The EFX re-allocation of the  $H$  chores thus leaves us to tackle the agents in  $N_H^1$  with exactly one overpaying chore. For these agents, we use chore swaps as described earlier.

**Performing swaps involving  $N_H^1$  agents.** Consider an  $(i, \ell)$  swap between an agent  $i \in N_H^1$  who was not 4-EFX and the agent  $\ell$  who  $i$  envied the most. We argued that after the swap  $D_i \leq 4 \cdot d_i(j_i)$ , where  $D_i$  is the disutility of  $i$  after the swap, and  $j_i \in H$  is the overpaying chore of  $i$  that was transferred to  $\ell$ . Consider a subsequent swap  $(h, k)$  between  $h \in N_H^1$  and  $k \in N$ , after

which the overpaying chore  $j_h \in H$  of agent  $h$  is (the only chore) assigned to  $k$ . Roughly speaking, since  $i$  was 4-EFX after the  $(i, \ell)$  swap,  $i$  does not 4-EFX envy  $k$ 's bundle before the swap. Hence,  $i$  will not envy  $h$  after the  $(h, k)$  swap. However, it could happen that  $i$  develops  $O(1)$ -EFX envy towards  $k$  after the  $(h, k)$  swap, if the swaps are made arbitrarily. But observe that if  $i$ 's disutility for  $j_h$  is at least that of  $j_i$ , then we will have that  $D_i \leq 4 \cdot d_i(j_i) \leq 4 \cdot d_i(j_h)$ , showing that  $i$  will not 4-EFX envy  $k$ . This observation suggests that we can avoid agents who participate in a swap from becoming envious again *by performing swaps in a carefully chosen order*. This order depends on the disutilities of  $N_H^1$  agents for the set  $H'$  of overpaying chores assigned to them, i.e.,  $H' = \cup_{i \in N_H^1} z'_i$ .

To determine this order, we re-allocate  $H'$  to the  $N_H^1$  agents by computing an fPO allocation  $(\mathbf{z}, \mathbf{q})$ , where every agent in  $N_H^1$  gets exactly one chore of  $H'$  in  $\mathbf{z}$ , and  $\mathbf{q}$  is the set of supporting payments. We show in Lemma 10 that such an allocation  $\mathbf{z}$  can be found by solving a linear program for minimum cost matching, and the payments  $\mathbf{q}$  can be computed from the dual variables of this program. Algorithm 4 then performs chore swaps in the following order: at each time step  $t$ , among all the agents in  $N_H^1$  who are not 4-EFX, we pick the agent  $i$  with the overpaying chore with the minimum payment  $\mathbf{q}(z_i)$ , and perform an  $(i, \ell)$  swap. An involved analysis shows that this design choice ensures Algorithm 4 does not cause an agent in  $N_H^1$  to re-develop 4-EFX envy. Algorithm 4 thus terminates in at most  $n$  steps.

With the above ideas, we argue that the resulting allocation  $\mathbf{x}$  is 4-EFX: (i) agents in  $N_H^1$  are addressed via swaps, (ii) agents in  $N_H^2$  remain 4-EFX during swaps, and (iii) agents in  $N_L$  are 3-EFX since their earning is at most  $3/2$  and every agent has an earning of at least  $1/2$ ; the latter property is maintained during every swap as well. Thus, Algorithm 4 returns a 4-EFX allocation.

### 4.3 4-EFX allocation: Algorithm analysis

We begin by providing a recap of Algorithm 4. In Phase 1, we compute a 2-EF2 and fPO allocation  $(\mathbf{x}', \mathbf{p})$  using the Algorithm 1 algorithm. We normalize the disutilities so that the MPB ratio of each agent is 1 for the payment vector  $\mathbf{p}$ . We then partition the chores of each agent  $i$  as  $\mathbf{x}'_i = S_i \cup H$ , where  $S_i \subseteq L$  contains low cost chores and  $H_i \subseteq H$  contains overpaying chores. This partitions the set of agents as  $N = N_H \sqcup N_L$ , where agents in  $N_H$  receive one or two overpaying chores, and agents in  $N_0$  receive none. In Phase 2, we re-allocate  $H$  by computing an EFX allocation  $\mathbf{z}'$  using Algorithm 3 with agents ordered as  $N_H$  first followed by  $N_0$ . We then categorize agents into  $N_H^1, N_H^2$  and  $N_L$  depending on the number of  $H$  chores they are assigned (see Definition 8) in  $\mathbf{z}'$ . The following is a useful property of  $\mathbf{z}'$ .

**Lemma 9.** *With agents ordered as  $N_H$  first, followed by  $N_0$ , Algorithm 3 computes an EFX allocation  $\mathbf{z}'$  of the overpaying chores  $H$  s.t. for each agent  $i$  with  $|z'_i| \geq 2$ , we have  $\mathbf{p}(S_i) \leq 1$ . In other words,  $N_H^2 \subseteq N_H$ .*

*Proof.* From Lemma 8, we know that for  $i \in N_H$ , we have  $\mathbf{p}(S_i) \leq 1$ . Hence it suffices to argue that if  $|z'_i| \geq 2$ , then  $i \in N_H$ . We will prove the contrapositive statement: if  $i \in N_0$ , then  $|z'_i| \leq 1$ .

First note that if  $|H| \leq n$ , then  $|z'_i| \leq 1$  for all  $i \in N$ . If  $|H| > n$ , then Theorem 10 shows that Algorithm 3 returns an EFX allocation in which agents with index greater than  $r := |H| - n$  have a single chore. These are the last  $2n - |H|$  agents in the order. Thus, it suffices to prove that  $|N_0| \leq 2n - |H|$ , since the agents in  $N_0$  appear last in our order.

Clearly,  $n = |N_H| + |N_0|$ . Since agents in  $N_H$  have exactly one or two overpaying chores, we have  $|H| \leq 2 \cdot |N_H|$ . This gives  $|N_0| + |H| \leq |N_0| + 2 \cdot |N_H| \leq 2n$ . This implies  $|N_0| \leq 2n - |H|$ , which is what we aimed to show.

In conclusion, any agent  $i \in N_0$  has  $|z'_i| \leq 1$ , thus proving the lemma.  $\square$

Next, we re-compute a matching  $\mathbf{z}$  of the chores  $H' = \cup_{i \in N_H^1} \mathbf{z}'_i$  to agents in  $N_H^1$ , and let  $\mathbf{q}$  be a set of payments of chores in  $H'$  such that  $(\mathbf{z}, \mathbf{q})$  is on MPB. The following lemma (proved in Appendix D) shows that such an allocation  $(\mathbf{z}, \mathbf{q})$  is computable in polynomial time.

**Lemma 10.** *Given a chore allocation instance with  $m = n$  chores, an MPB allocation  $(\mathbf{z}, \mathbf{q})$  where  $|\mathbf{z}_i| = 1$  for each  $i \in N$  can be computed in polynomial time.*

In Phase 3, we perform *swaps* (Definition 7) involving agents in  $N_H^1$  which are not 4-EFX. In each step, we pick the 4-EFX envious agent  $i \in N_H^1$  with the minimum  $\mathbf{q}(\mathbf{z}_i)$ .

Let  $\mathbf{x}$  be the allocation computed at the end of Phase 2, before any Phase 3 swaps are performed. For  $t \in \mathbb{Z}_{\geq 0}$ , we use the phrase ‘at time step  $t$ ’ to refer to the  $t^{\text{th}}$  iteration of the while loop of Algorithm 4, and use it interchangeably with ‘just before the swap at time step  $t$ ’. Let  $\mathbf{x}^t$  denote the allocation at time step  $t$ , with  $\mathbf{x} = \mathbf{x}^0$ . We first prove a few basic invariants maintained by Algorithm 4.

**Lemma 11.** *At any time step  $t$  in the run of Algorithm 4, we have:*

- (i) *For any agent  $i \in N$ ,  $\mathbf{p}(\mathbf{x}_i^t) \geq 1/2$ .*
- (ii) *If an agent  $i \in N_H^2 \cup N_L$  has participated in a swap at time  $t' < t$ , then  $i$  is EFX at  $t$ .*
- (iii) *Any agent  $i \in N_H^2$  is 4-EFX.*
- (iv) *Any agent  $i \in N_L$  is 3-EFX.*

*Proof.* We first prove claim (i) by an inductive argument. For the allocation  $(\mathbf{x}', \mathbf{p})$  obtained by rounding the ER equilibrium, we have that  $\mathbf{p}(\mathbf{x}'_i) \geq 1/2$ . This remains true at time step  $t = 0$  after the overpaying chores are re-allocated in Phase 2, since each overpaying chore pays at least  $1/2$ . Suppose claim (i) holds at time step  $t$  before an  $(i, \ell)$  swap takes place. After the swap,  $\ell$  is assigned an overpaying chore, hence  $\mathbf{p}(\mathbf{x}_\ell^{t+1}) > 1/2$ . Moreover, since  $i$  receives chores earlier assigned to  $\ell$ , we have  $\mathbf{p}(\mathbf{x}_i^{t+1}) \geq \mathbf{p}(\mathbf{x}_\ell^t) \geq 1/2$  using the inductive hypothesis at time  $t$ . Thus, claim (i) holds at every time step in the run of the algorithm.

For claim (ii), observe that an agent  $i \in N_H^2 \cup N_L$  can only participate in a swap of the form  $(h, i)$ , where  $h \in N_H^1$ . Then  $i$  is EFX immediately after the swap since  $i$  is assigned a single chore. This remains true even after subsequent swaps that  $i$  participates in, and hence  $i$  remains EFX at any time step of the algorithm.

Since the allocation of overpaying chores  $\mathbf{z}'$  is EFX, we have that  $N_H^2 \neq \emptyset$  and  $N_L \neq \emptyset$  cannot both be true. We first assume  $N_H^2 \neq \emptyset$  and prove claim (iii). In this case,  $N_L = \emptyset$ . Consider an agent  $i \in N_H^2$ . If  $i$  participated in an  $(h, i)$  swap before  $t$ , then  $i$  is EFX at  $t$  due to claim (ii). Hence we assume that  $i \in N_H^2$  did not participate in any swap, and thus  $\mathbf{x}_i^t = S_i \cup \mathbf{z}'_i$ . Let  $j_0 = \arg \min_{j \in \mathbf{z}'_i} d_i(j)$ . Since  $i \in N_H^2$ ,  $\mathbf{z}'_i \setminus \{j\} \neq \emptyset$ . Consider any other agent  $h \in N$  at time step  $t$ . We have that  $\mathbf{x}_h^t \supseteq \mathbf{z}'_k$  for some  $k \in N$  since an agent participating in a chore swap always swaps all of her overpaying chores. We now show that  $i$  is 4-EFX towards  $h$  as follows.

$$\begin{aligned}
d_i(\mathbf{x}_i^t) &= d_i(S_i) + d_i(\mathbf{z}'_i) && \text{(since } i \text{ did not undergo any swap)} \\
&\leq \mathbf{p}(S_i) + d_i(\mathbf{z}'_i \setminus \{j_0\}) + d_i(j_0) \\
&\leq \mathbf{p}(S_i) + 2 \cdot d_i(\mathbf{z}'_i \setminus \{j_0\}) && \text{(since } \mathbf{z}'_i \setminus \{j_0\} \neq \emptyset \text{ and } j_0 = \arg \min_{j \in \mathbf{z}'_i} d_i(j)) \\
&\leq \mathbf{p}(S_i) + 2 \cdot d_i(\mathbf{z}'_k) && \text{(since } \mathbf{z}' \text{ is EFX)} \\
&\leq 1 + 2 \cdot d_i(\mathbf{z}'_k) && \text{(since } \mathbf{p}(S_i) \leq 1 \text{ using Lemma 9)} \\
&\leq 4 \cdot d_i(\mathbf{z}'_k) && \text{(using } \mathbf{p}(\mathbf{z}'_k) > 1/2 \text{ since } \mathbf{z}'_k \subseteq H) \\
&\leq 4 \cdot d_i(\mathbf{x}_h^t) && \text{(since } \mathbf{x}_h^t \supseteq \mathbf{z}'_k \text{).}
\end{aligned}$$



Finally, we assume  $N_L \neq \emptyset$  and prove claim (iv). In this case,  $N_H^2 = \emptyset$ . Consider an agent  $i \in N_L$ . As before, if  $i$  participated in a swap at a time before  $t$ , then  $i$  is EFX at  $t$  due to claim (ii). Hence we assume that  $i \in N_L$  did not participate in any swap, and thus  $\mathbf{x}_i^t = S_i$ . Consider any other agent  $h \in N$  at time step  $t$ . We show that  $i$  is 3-EFX towards  $h$  as follows.

$$\begin{aligned} d_i(\mathbf{x}_i^t) &= \mathbf{p}(S_i) && \text{(since } \mathbf{x}_i^t = \mathbf{x}'_i = S_i \text{ is on MPB)} \\ &\leq \frac{3}{2} < 3 \cdot \mathbf{p}(\mathbf{x}_h^t) && \text{(since claim (ii) shows } \mathbf{p}(\mathbf{x}_h^t) > 1/2) \\ &\leq 3 \cdot d_i(\mathbf{x}_h^t). && \text{(using the MPB condition)} \end{aligned}$$

This proves the lemma.  $\square$

The above lemma shows that  $N_H^2$  and  $N_L$  agents are 4-EFX and we need to address the  $N_H^1$  agents. Let us examine the change in disutility of agent  $i \in N_H^1$  after an  $(i, \ell)$  swap.

**Lemma 12.** *Suppose  $i \in N_H^1$  participates in an  $(i, \ell)$  swap for the first time at time step  $t$ . Then,  $d_i(\mathbf{x}_i^{t+1}) < 4 \cdot d_i(\mathbf{z}_i)$ , and  $i$  is 4-EFX immediately after the swap.*

*Proof.* Since  $i$  has not undergone a swap until time step  $t$ , we have  $\mathbf{x}_i^t = S_i \cup \mathbf{z}_i$ . By the definition of a swap, we have  $\mathbf{x}_i^{t+1} = S_i \cup \mathbf{x}_\ell^t$ ,  $\mathbf{x}_\ell^{t+1} = \mathbf{z}_i$ , and  $\mathbf{x}_h^{t+1} = \mathbf{x}_h^t$  for all  $h \notin \{i, \ell\}$ . Since  $i$  is not 4-EFX towards  $\ell$  at time  $t$ , we know  $d_i(\mathbf{x}_i^t) > 4 \cdot d_i(\mathbf{x}_\ell^t)$ . We prove the first part of the lemma using the above observations.

$$\begin{aligned} d_i(\mathbf{x}_i^{t+1}) &= d_i(S_i) + d_i(\mathbf{x}_\ell^t) \\ &< d_i(S_i) + \frac{d_i(\mathbf{x}_i^t)}{4} && \text{(using } d_i(\mathbf{x}_i^t) > 4 \cdot d_i(\mathbf{x}_\ell^t)) \\ &= d_i(S_i) + \frac{d_i(S_i) + d_i(\mathbf{z}_i)}{4} && \text{(since } \mathbf{x}_i^t = S_i \cup \mathbf{z}_i) \\ &= \frac{5}{4} \cdot d_i(S_i) + \frac{1}{4} \cdot d_i(\mathbf{z}_i) \\ &= \frac{5}{4} \cdot \mathbf{p}(S_i) + \frac{1}{4} \cdot d_i(\mathbf{z}_i) && \text{(since } d_i(S_i) = \mathbf{p}(S_i) \text{ using the MPB condition)} \\ &< \frac{15}{4} \cdot d_i(\mathbf{z}_i) + \frac{1}{4} \cdot d_i(\mathbf{z}_i) && \text{(using } \mathbf{p}(S_i) \leq 3/2 \text{ and } d_i(\mathbf{z}_i) > 1/2) \\ &= 4 \cdot d_i(\mathbf{z}_i). \end{aligned}$$

By the choice of  $\ell$ , it holds that  $d_i(\mathbf{x}_h^t) \geq d_i(\mathbf{x}_\ell^t)$  for  $h \in N \setminus \{\ell\}$ . We next prove that  $i$  is 4-EFX in the allocation  $\mathbf{x}^{t+1}$  after the swap.

- $i$  is 4-EFX towards  $\ell$ , since  $d_i(\mathbf{x}_i^{t+1}) < 4 \cdot d_i(\mathbf{z}_i)$  as argued previously and  $\mathbf{x}_\ell^{t+1} = \mathbf{z}_i$ .
- $i$  is 4-EFX towards an agent  $h \in N \setminus \{\ell\}$ , since:

$$\begin{aligned} d_i(\mathbf{x}_i^{t+1}) &= d_i(S_i) + d_i(\mathbf{x}_\ell^t) && \text{(since } \mathbf{x}_i^{t+1} = S_i \cup \mathbf{x}_\ell^t) \\ &= \mathbf{p}(S_i) + d_i(\mathbf{x}_\ell^t) && \text{(since } d_i(S_i) = \mathbf{p}(S_i) \text{ using the MPB condition)} \\ &\leq \frac{3}{2} + d_i(\mathbf{x}_\ell^t) && \text{(using } \mathbf{p}(S_i) \leq \frac{3}{2}) \\ &\leq 3 \cdot \mathbf{p}(\mathbf{x}_\ell^t) + d_i(\mathbf{x}_\ell^t) && \text{(since } \mathbf{p}(\mathbf{x}_\ell^t) \geq 1/2 \text{ by Lemma 11)} \\ &\leq 4 \cdot d_i(\mathbf{x}_\ell^t) && \text{(since the MPB condition implies } d_i(\mathbf{x}_\ell^t) \geq \mathbf{p}(\mathbf{x}_\ell^t)) \\ &\leq 4 \cdot d_i(\mathbf{x}_h^t) = 4 \cdot d_i(\mathbf{x}_h^{t+1}). && \text{(by choice of } \ell) \end{aligned}$$

$\square$

The above lemma shows that an agent  $i \in N_H^1$  is 4-EFX immediately after the first  $(i, \ell)$  swap she participates in. Next, we argue that such an agent cannot develop 4-EFX envy again. The key idea is to choose among all  $N_H^1$  agents who are not 4-EFX, the agent  $i$  with minimum  $\mathbf{q}(\mathbf{z}_i)$ . Let  $\alpha_i$  denote the MPB ratio of  $i$  in  $(\mathbf{z}, \mathbf{q})$ . Note that  $|\mathbf{z}_i| = 1$  for all  $i \in N_H^1$ . For an  $(i, \ell)$  swap at time step  $t$ , we let  $q_t = \mathbf{q}(\mathbf{z}_i)$  denote the payment of the overpaying chore  $\mathbf{z}_i$  transferred from  $i$  to  $\ell$ . We now prove the following set of invariants of Algorithm 4.

**Lemma 13.** *At any time step  $t$  in the while loop of Algorithm 4, the following hold:*

- (i) *Every agent  $i$  has participated in at most one  $(i, \ell)$  swap until time  $t$ .*
- (ii) *If an agent  $i$  has participated in an  $(i, \ell)$  swap at time  $t' < t$ , then  $i$  is 4-EFX at  $t$ .*
- (iii) *If an agent  $i \in N_H^1$  is not 4-EFX at  $t$ , then  $q(\mathbf{z}_i) \geq q_{t-1}$ .*
- (iv)  $q_t \geq q_{t-1} \geq \dots \geq q_1 \geq q_0 := 0$ .

*Proof.* We prove this by induction on  $t$ . Since no agent has participated in a swap before  $t = 1$  and  $q_0 = 0$ , claims (i)-(iv) are vacuously true at  $t = 1$ . Suppose claims (i)-(iv) hold true at some time step  $t \geq 1$ . Consider a swap  $(i, \ell)$  taking place at  $t$ . We prove that claims (i)-(iv) hold at time  $(t + 1)$  after the swap has taken place.

- (i) Suppose  $i$  has already participated in a swap at time  $t' < t$ , then claim (ii) of the induction hypothesis implies that  $i$  is 4-EFX at time  $t$ , contradicting the fact that an  $(i, \ell)$  swap takes place at  $t$ . Thus  $i$  participates in her first swap at  $t$ . Together with claim (i) of the induction hypothesis for an agent  $h \neq i$ , this establishes claim (i) at time  $(t + 1)$ .
- (ii) We will prove that every agent  $h$  who has participated in a swap at time  $t' < (t + 1)$  is 4-EFX at  $(t + 1)$ . We first consider the case of  $h = i$ . Note that Lemma 12 implies that  $i$  is 4-EFX after the swap at  $t$ , i.e.,  $i$  is 4-EFX at time  $(t + 1)$ .

Next we prove the claim for agents  $h \neq i$ . Suppose an agent  $h \neq i$  participated in a swap of the form  $(h, k)$  at time  $t' < (t + 1)$ . Since  $h \neq i$ , and the  $(i, \ell)$  swap takes place at time  $t$ , we know  $t' < t$ . We therefore apply claim (ii) of the induction hypothesis to obtain that  $h$  is 4-EFX at time  $t$ . In particular, this shows that at time  $(t + 1)$ , agent  $h$  remains 4-EFX towards all agents  $h' \neq \{i, \ell\}$  who don't participate in the  $(i, \ell)$  swap at time  $t$ . Moreover, since  $h$  is 4-EFX towards the bundle  $\mathbf{x}_\ell^t$  and  $\mathbf{x}_{i+1}^t \supseteq \mathbf{x}_\ell^t$ ,  $h$  remains 4-EFX towards agent  $i$  after the swap at  $(t + 1)$ .

It remains to be shown that  $h$  is 4-EFX towards agent  $\ell$  at time  $(t + 1)$ . By claim (i) of the induction hypothesis,  $h$  does not participate in swaps during times  $[t' + 1, t]$ . Thus  $\mathbf{x}_h^{t+1} = \mathbf{x}_h^{t'+1}$ . Then, Lemma 12 implies that  $d_h(\mathbf{x}_h^{t'+1}) \leq 4 \cdot d_h(\mathbf{z}_h)$ . Using the MPB condition for the fPO allocation  $(\mathbf{z}, \mathbf{q})$ , we get that  $d_h(\mathbf{z}_h) = \alpha_h \mathbf{q}(\mathbf{z}_h)$ . Using  $\mathbf{q}(\mathbf{z}_h) = q_{t'}$ , we conclude that  $d_h(\mathbf{x}_h^{t+1}) \leq 4 \cdot \alpha_h q_{t'}$ . With this, the following chain of inequalities shows that  $h$  remains 4-EFX towards  $\ell$  at  $(t + 1)$ .

$$\begin{aligned}
d_h(\mathbf{x}_h^{t+1}) &\leq 4 \cdot \alpha_h q_{t'} && \text{(as argued above)} \\
&\leq 4 \cdot \alpha_h q_t && \text{(using claim (iv) of the induction hypothesis)} \\
&= 4 \cdot \alpha_h \mathbf{q}(\mathbf{z}_i), && \text{(using } \mathbf{q}(\mathbf{z}_i) = q_t) \\
&\leq 4 \cdot d_h(\mathbf{z}_i), && \text{(using the MPB condition)} \\
&\leq 4 \cdot d_h(\mathbf{x}_\ell^{t+1}). && \text{(since } \mathbf{x}_\ell^{t+1} = \mathbf{z}_i)
\end{aligned}$$

This proves that claim (ii) holds at time  $(t + 1)$ .

- (iii) We prove that for an agent  $h \in N_H^1$  who is not 4-EFX at time  $(t+1)$ , it holds that  $\mathbf{q}(\mathbf{z}_h) \geq q_t$ . Clearly,  $h \neq i$ . Moreover, claim (ii) proved above shows that  $h$  has not participated in a swap at any time step  $t' < t+1$ . Thus,  $\mathbf{x}_h^{t+1} = S_h \cup \mathbf{z}_h$ . Thus we have:

$$\begin{aligned}
d_h(\mathbf{x}_h^{t+1}) &= d_h(S_h) + d_h(\mathbf{z}_h) && \text{(using } \mathbf{x}_h^{t+1} = S_h \cup \mathbf{z}_h) \\
&= \mathbf{p}(S_h) + d_h(\mathbf{z}_h) && \text{(since } d_h(S_h) = \mathbf{p}(S_h)) \\
&\leq \frac{3}{2} + d_h(\mathbf{z}_h) && \text{(using } \mathbf{p}(S_h) \leq \frac{3}{2}) \\
&\leq 4 \cdot d_h(\mathbf{z}_h) && \text{(using } d_h(\mathbf{z}_h) > \frac{1}{2}) \\
&= 4 \cdot \alpha_h \mathbf{q}(\mathbf{z}_h). && \text{(using the MPB condition for } (\mathbf{z}, \mathbf{q}))
\end{aligned} \tag{1}$$

If  $h$  was not 4-EFX at  $t$ , then  $\mathbf{q}(\mathbf{z}_h) \geq \mathbf{q}(\mathbf{z}_i)$ , since Algorithm 4 chose to perform a swap involving  $i$  instead of  $h$ . Thus  $\mathbf{q}(\mathbf{z}_h) \geq \mathbf{q}(\mathbf{z}_i) = q_t$  in this case.

On the other hand, suppose  $h$  was 4-EFX at  $t$ . Since  $h$  is not 4-EFX at  $(t+1)$ ,  $h$  became envious at  $(t+1)$  due to the  $(i, \ell)$  swap at  $t$ . Since  $\mathbf{x}_i^{t+1} \supseteq \mathbf{x}_\ell^t$ ,  $h$  does not 4-EFX envy  $i$  at  $(t+1)$ . This implies that  $h$  4-EFX envies  $\ell$  at time  $(t+1)$ .

Thus,

$$\begin{aligned}
4 \cdot \alpha_h \mathbf{q}(\mathbf{z}_h) &\geq d_h(\mathbf{x}_h^{t+1}) && \text{(using Eq. (1))} \\
&> 4 \cdot d_h(\mathbf{x}_\ell^{t+1}) && \text{(since } h \text{ 4-EFX envies } \ell) \\
&= 4 \cdot d_h(\mathbf{z}_i) && \text{(using claim (i))} \\
&= 4 \cdot \alpha_h q_t. && \text{(using the MPB condition of } (\mathbf{z}, \mathbf{q}))
\end{aligned} \tag{2}$$

Dividing each side by  $4 \cdot \alpha_h$ , we obtain  $\mathbf{q}(\mathbf{z}_h) \geq q_t$ , as claimed.

- (iv) Consider a swap  $(h, k)$  taking place at  $(t+1)$ . Since  $h$  is not 4-EFX at  $(t+1)$ , we have that  $\mathbf{q}(\mathbf{z}_h) \geq q_t$  by claim (iii) proved above. Thus  $q_{t+1} := \mathbf{q}(\mathbf{z}_h) \geq q_t$ . With claim (iv) of the induction hypothesis at  $t$ , we obtain  $q_{t+1} \geq q_t \geq \dots \geq q_1 \geq q_0$ , as desired.  $\square$

We can now prove the main result of this section.

**Theorem 11.** *Given an ER equilibrium of a chore allocation instance with  $m \geq 2n$  and earning limit  $\beta = 1/2$ , Algorithm 4 returns a 4-EFX allocation in polynomial time.*

*Proof.* Algorithm 4 runs as long as there is an agent  $i \in N_H^1$  who is not 4-EFX. Lemma 21 shows that once an agent  $i \in N_H^1$  who participates in an  $(i, \ell)$  swap, she remains 4-EFX in the subsequent run of the algorithm. Thus, there can only be  $n$  swap steps before the algorithm terminates.

We argue that the resulting allocation is 4-EFX. If an agent  $i \in N_H^1$  is not 4-EFX, then the algorithm would not have terminated. Thus, all agents in  $N_H^1$  are 4-EFX upon termination of the algorithm. Finally, Lemma 11 shows that the agents in  $N_H^2$  are 4-EFX and agents in  $N_L$  are 3-EFX throughout the run of the algorithm. In conclusion, given an ER equilibrium for a chore allocation instance with  $m \geq 2n$ , Algorithm 4 returns a 4-EFX allocation in polynomial time.  $\square$

## 5 Approximate EFX and PO for Bivalued Instances

We now turn to the problem of computing (approximately-)EFX and PO allocations for *bivalued* instances. Recall that in a bivalued instance  $(N, M, D)$  there exist  $a, b \in \mathbb{R}_{>0}$  s.t.  $d_{ij} \in \{a, b\}$ . Note that we can re-scale the disutilities so that  $d_{ij} \in \{1, k\}$ , where  $k > 1$ . We refer to such an instance as a  $\{1, k\}$ -bivalued instance. The main result of this section is that:

**Theorem 12.** *Any bivalued instance admits a 3-EFX and fPO allocation.*

To prove Theorem 12, we design and analyze two algorithms: Algorithm 5, which computes a 3-EFX and PO allocation when  $m > 2n$ ; and Algorithm 7 which computes an EFX and PO allocation when  $m \leq 2n$ . Both algorithms begin with initial allocations with certain desirable properties and perform subsequent chore transfers to achieve (approximate-)EFX and PO. Algorithm 5 begins with the 2-EF2 and PO allocation obtained by rounding an ER equilibria using Algorithm 1, while Algorithm 7 begins with the balanced allocation obtained using Algorithm 2. Before discussing our algorithms, we note that the bivalued nature of the instance allows us to prove some important properties of any competitive equilibrium  $(\mathbf{x}, \mathbf{p})$ .

**Lemma 14.** *Let  $(\mathbf{x}, \mathbf{p})$  be a CE of a  $\{1, k\}$ -bivalued instance with  $\rho = \min_j p_j$ . Then:*

- (i) *For every  $j \in M$ ,  $\rho \leq p_j \leq \rho k$ .*
- (ii) *Let  $j \in \mathbf{x}_i$  be s.t.  $p_j \in (\rho, \rho k)$ . Then for all  $j' \in \mathbf{x}_i$ ,  $p_{j'} = p_j$ .*

*Proof.* For any  $j \in M$ ,  $p_j \geq \rho$  follows from the definition of  $\rho$ . Suppose for some  $j_1 \in M$ ,  $p_{j_1} > \rho k$ . Let  $j_0 \in \mathbf{x}_\ell$  be such that  $p_{j_0} = \rho$ . Then the MPB condition for agent  $\ell$  implies that  $\frac{d_{\ell j_0}}{p_{j_0}} \leq \frac{d_{\ell j_1}}{p_{j_1}}$ . This implies  $\frac{d_{\ell j_0}}{d_{\ell j_1}} \leq \frac{p_{j_0}}{p_{j_1}} < \frac{1}{k}$ . However this is a contradiction since  $d_{ij} \in \{1, k\}$  for all  $i \in N, j \in M$ . This proves (i).

For (ii), suppose  $\exists j, j' \in \mathbf{x}_i$  s.t.  $p_j \in (\rho, \rho k)$  and  $p_{j'} \neq p_j$ . Then the MPB condition for  $i$  implies that  $\frac{d_{ij}}{p_j} = \frac{d_{ij'}}{p_{j'}}$ , implying that  $\frac{p_j}{p_{j'}} = \frac{d_{ij}}{d_{ij'}}$ . Since  $d_{ij}, d_{ij'} \in \{1, k\}$ , we know  $\frac{p_j}{p_{j'}} \in \{1/k, 1, k\}$ . Since  $p_j \neq p_{j'}$ , we have  $\frac{p_j}{p_{j'}} \in \{1/k, k\}$ . Thus  $p_{j'} = kp_j$  or  $p_{j'} = p_j/k$ . Since  $p_j \in (\rho, \rho k)$  and  $k > 1$ , this implies either  $p_{j'} > \rho k$  or  $p_{j'} < \rho$ , both of which contradict (i).  $\square$

### 5.1 3-EFX and PO for $m > 2n$

We first present Algorithm 5: a polynomial-time algorithm that computes a 3-EFX and fPO allocation for a bivalued instance with  $m > 2n$ , given its ER equilibrium  $(\mathbf{y}, \mathbf{p})$  as input. Algorithm 5 first rounds  $(\mathbf{y}, \mathbf{p})$  using Algorithm 1 with the chore earning limit set as  $\beta = 1/2$ . The resulting allocation  $\mathbf{x}^0$  is already a good starting point: it is fPO, and Lemmas 5 and 6 with  $\beta = 1/2$  show its fairness properties.

**Lemma 15.** *The allocation  $(\mathbf{x}^0, \mathbf{p})$  is fPO and satisfies:*

- (i) *For all  $i \in N$ ,  $\mathbf{p}(\mathbf{x}_i^0) \geq 1/2$ .*
- (ii) *For all  $h \in N$ , either  $\mathbf{p}_{-1}(\mathbf{x}_h^0) \leq 1$ , or  $|\mathbf{x}_h^0 \cap \{j : p_j > 1/2\}| = 2$  and  $\mathbf{p}_{-2}(\mathbf{x}_h^0) \leq 1/2$ .*

Let  $\rho = \min_j p_j$  be the minimum chore payment.

**Lemma 16.** *The minimum chore payment satisfies  $\rho < 1/2$ .*

*Proof.* Let  $q_j = \min\{p_j, 1/2\}$  be the earning from chore  $j \in M$ . If  $\rho \geq 1/2$ , then  $p_j \geq 1/2$  for all  $j$ , implying that  $q_j = 1/2$ . Hence, the total earning from chores is  $\sum_j q_j = \frac{m}{2}$ . Since  $\sum_j q_j = \sum_i e_i = n$ , we obtain that  $n = \frac{m}{2}$ , which contradicts our assumption that  $m > 2n$ .  $\square$

The next lemma shows that if the largest chore payment is small,  $\mathbf{x}^0$  is already fair.

**Lemma 17.** *If  $\rho k \leq 1/2$ , then  $\mathbf{x}^0$  is 3-EF.*

*Proof.* If  $\rho k \leq 1$ , then  $p_j \leq 1/2$  for all  $j \in M$  by Lemma 14. Lemma 15 then implies that:

- (i) For all  $i \in N$ ,  $\mathbf{p}(\mathbf{x}_i^0) \geq 1/2$ .
- (ii) For all  $h \in N$ ,  $\mathbf{p}_{-1}(\mathbf{x}_h^0) \leq 1$ , or  $\mathbf{p}_{-2}(\mathbf{x}_h^0) \leq 1/2$ . Thus  $\mathbf{p}(\mathbf{x}_h^0) \leq \max\{1 + \frac{1}{2}, \frac{1}{2} + 2 \cdot \frac{1}{2}\} = \frac{3}{2}$ .

We therefore have that  $(\mathbf{x}^0, \mathbf{p})$  is 3-EF, since for any  $i, h \in N$ , we have  $\mathbf{p}(\mathbf{x}_h^0) \leq \frac{3}{2} \leq 3 \cdot \mathbf{p}(\mathbf{x}_i^0)$ .  $\square$

Thus, Algorithm 5 simply returns  $(\mathbf{x}^0, \mathbf{p})$  if  $\rho k \leq 1/2$ . Hence, we assume  $\rho k > 1/2$  in the subsequent discussion. Note that  $\rho < 1/2$  by Lemma 16.

**Definition 9.** (*Classification of Chores and Agents in Algorithm 5*) Chores are categorized as:

- $L = \{j \in M : p_j = \rho\}$ , i.e., low paying or  $L$ -chores. Note  $p_j = \rho < 1/2$  for all  $j \in L$ .
- $H = \{j \in M : p_j = \rho k\}$ , i.e., high paying or  $H$ -chores. Note  $p_j = \rho k > 1/2$  for all  $j \in H$ .
- $M' = M \setminus (L \cup H) = \{j \in M : p_j \in (\rho, \rho k)\}$ , or  $M'$ -chores.

Lemma 14 (ii) shows that agents can either be assigned chores from  $M'$  or from  $L$  and  $H$ , but not both. With this observation, we classify agents into four categories given an allocation  $(\mathbf{x}, \mathbf{p})$ :

- $N_L = \{i \in N : \mathbf{x}_i \subseteq L\}$ , i.e., agents who are only assigned  $L$ -chores.
- $N_H^1 = \{i \in N : |\mathbf{x}_i \cap H| = 1\}$ , i.e., agents who are assigned exactly one  $H$ -chore.
- $N_H^2 = \{i \in N : |\mathbf{x}_i \cap H| = 2\}$ , i.e., agents who are assigned exactly two  $H$ -chores.
- $N_0 = \{i \in N : \mathbf{x}_i \subseteq M'\}$ , i.e., agents who are only assigned  $M'$ -chores.

Let  $N_H = N_H^1 \cup N_H^2$ . We begin by exploring the source of EFX-envy in  $\mathbf{x}^0$ . We prove a general lemma concerning the EFX-envy of agents in  $N_0$ .

**Lemma 18.** *Consider an allocation  $(\mathbf{x}, \mathbf{p})$  s.t.  $\mathbf{x}_i = \mathbf{x}_i^0$  for all  $i \in N_0$  and  $\mathbf{p}(\mathbf{x}_h) \geq 1/2$  for all  $h \in N$ . Then  $\mathbf{x}$  is 2-EFX for any  $i \in N_0$ .*

*Proof.* Consider an agent  $i \in N_0$ . As per Lemma 15, we consider two cases regarding  $\mathbf{x}_i^0$ :

- (i)  $\mathbf{p}_{-1}(\mathbf{x}_i^0) \leq 1$ . Since Lemma 21 implies that  $\mathbf{p}(\mathbf{x}_h) \geq 1/2$  for any  $h \in N$ , we obtain that  $\mathbf{p}_{-1}(\mathbf{x}_i^0) \leq 2 \cdot \mathbf{p}(\mathbf{x}_h)$ . Thus  $i$  is 2-pEFX and hence 2-EFX towards any  $h \in N$  by Lemma 1.
- (ii)  $|\mathbf{x}_i^0 \cap \{j : p_j > 1/2\}| = 2$  and  $\mathbf{p}_{-2}(\mathbf{x}_i^0) \leq 1/2$ . Let  $\mathbf{x}_i^0 \cap \{j : p_j > 1/2\} = \{j_1, j_2\}$ . By Lemma 14, all chores in  $\mathbf{x}_i^0$  have the same payment  $\rho' \in (\rho, \rho k)$ . Hence  $p_{j_1} = p_{j_2} = \rho' > 1/2$ . Thus  $\mathbf{p}_{-2}(\mathbf{x}_i^0) \leq 1/2$  implies that  $\mathbf{x}_i^0 \setminus \{j : p_j > 1/2\} = \emptyset$ , i.e.,  $\mathbf{x}_i^0 = \{j_1, j_2\}$ .

Suppose  $i$  is not EFX towards an agent  $h \in N$ . If  $\exists j \in \mathbf{x}_h$  s.t.  $d_{ij} = k$ , then  $\max_{j' \in \mathbf{x}_h} d_i(\mathbf{x}_i \setminus \{j'\}) \leq k = d_{ij} \leq d_i(\mathbf{x}_h)$ , implying that  $i$  is EFX towards  $h$ . On the contrary, if  $\forall j \in \mathbf{x}_h$ ,  $d_{ij} = 1$ , then the MPB condition for agent  $i$  implies  $\frac{d_{ij_1}}{p_{j_1}} \leq \frac{d_{ij}}{p_j}$  for  $j \in \mathbf{x}_h$ . This gives:  $k \cdot p_j \leq p_{j_1} < \rho k$ , implying  $p_j < \rho$ . This contradicts the definition of  $\rho$ .  $\square$

Since  $\mathbf{p}(\mathbf{x}_h^0) \geq 1/2$  for all  $h \in N$  by Lemma 15, the above lemma shows that  $\mathbf{x}^0$  is 2-EFX for agents in  $N_0$ . Next, we show that  $\mathbf{x}^0$  is also 2-EFX for agents in  $N_L$ .

**Lemma 19.**  $\mathbf{x}^0$  is 2-EFX for agents in  $N_L$ .

---

**Algorithm 5** 3-EFX + PO for bivalued instances with  $m > 2n$ 

---

**Input:**  $\{1, k\}$ -bivalued instance with  $m > 2n$ , its ER equilibrium  $(\mathbf{y}, \mathbf{p})$  with  $\beta = 1/2$

**Output:** An integral allocation  $\mathbf{x}$

- 1:  $(\mathbf{x}, \mathbf{p}) \leftarrow$  Run Algorithm 1 with  $(\mathbf{y}, \mathbf{p})$
  - 2:  $\rho \leftarrow \min_j p_j$
  - 3: **if**  $\rho k \leq 1/2$  **then return**  $\mathbf{x}$   $\triangleright \mathbf{x}$  is 3-EFX by Lemma 17
  - 4:  $L = \{j \in M : p_j = \rho\}$ ,  $H = \{j \in M : p_j = \rho k\}$   $\triangleright$  Low, High paying chores
  - 5: Classify agents as  $N_L, N_H^1, N_H^2, N_0$  (See Definition 9)  
— Phase 1: Address  $N_H^2$  agents —
  - 6: **while**  $\exists i \in N_H^2$  not 3-EFX **do**  $\triangleright$  Lemma 22 shows  $\ell \in N_L$
  - 7:  $\ell \leftarrow$  agent 3-EFX-envied by  $i$
  - 8: **if**  $\mathbf{p}(\mathbf{x}_\ell) > 1$  **then**  $S \leftarrow j_1$  for some  $j_1 \in \mathbf{x}_\ell$
  - 9: **else**  $S \leftarrow \emptyset$
  - 10:  $j \in \mathbf{x}_i \cap H$
  - 11:  $\mathbf{x}_\ell \leftarrow \mathbf{x}_\ell \setminus S \cup \{j\}$
  - 12:  $\mathbf{x}_i \leftarrow \mathbf{x}_i \cup S \setminus \{j\}$
  - 13:  $N_H^1 \leftarrow N_H^1 \cup \{i, \ell\}$ ,  $N_H^2 \leftarrow N_H^2 \setminus \{i\}$ ,  $N_L \leftarrow N_L \setminus \{\ell\}$   
— Phase 2: Address  $N_H^1$  agents —
  - 14: **while**  $\exists i \in N_H^1$  not 3-EFX **do**  $\triangleright$  Lemma 22 shows  $\ell \in N_L$
  - 15:  $\ell \leftarrow \arg \min\{\mathbf{p}(\mathbf{x}_h) : h \in N \text{ s.t. } i \text{ 3-EFX envies } h\}$
  - 16:  $j \in \mathbf{x}_i \cap H$
  - 17:  $\mathbf{x}_i \leftarrow \mathbf{x}_i \cup \mathbf{x}_\ell \setminus \{j\}$
  - 18:  $\mathbf{x}_\ell \leftarrow \{j\}$
  - 19:  $N_H^1 \leftarrow N_H^1 \cup \{\ell\} \setminus \{i\}$ ,  $N_L \leftarrow N_L \cup \{i\} \setminus \{\ell\}$
  - 20: **return**  $\mathbf{x}$
- 

*Proof.* Consider an agent  $i \in N_L$ . Since  $\mathbf{x}_i^0 \subseteq L$ ,  $\mathbf{p}_{-X}(\mathbf{x}_i^0) = \mathbf{p}_{-1}(\mathbf{x}_i^0)$ . Lemma 15 implies that  $\mathbf{p}_{-1}(\mathbf{x}_i^0) \leq 1 \leq 2 \cdot \mathbf{p}(\mathbf{x}_h^0)$  for any  $h \in N$ . This shows  $i$  is 2-EFX towards any  $h \in N$ .  $\square$

If  $H = \emptyset$ ,  $N = N_0 \cup N_L$ . Thus  $\mathbf{x}^0$  is 2-EFX, and Algorithm 5 will simply return  $\mathbf{x}^0$ . Hence, we assume  $H \neq \emptyset$  in the subsequent discussion. Lemmas 18 and 19 show that  $\mathbf{x}^0$  is 2-EFX for agents in  $N_0 \cup N_L$ . Hence if  $\mathbf{x}^0$  is not 3-EFX, some agent in  $N_H$  must 3-EFX-envy another agent. Intuitively, an agent  $i \in N_H$  2-EFX-envies another agent  $\ell$  since  $i$  has one or two high-paying  $H$ -chores in addition to some low-paying  $L$ -chores. Algorithm 5 addresses the EFX-envy of these agents by swapping some chores between agents  $i$  and  $\ell$ , and does so in two phases.

In Phase 1, Algorithm 5 addresses agents in  $N_H^2$ . An agent  $i \in N_H^2$  has two  $H$ -chores, and earns at most  $1/2$  from  $L$ -chores. We show in Lemma 22 that if  $i$  3-EFX-envies an agent  $\ell$ , we must have  $\ell \in N_L$ . We then transfer one  $H$ -chore from  $i$  to  $\ell$ , and if needed, transfer a single  $L$ -chore from  $\ell$  to  $i$  so that both agents earn at most 1 from their  $L$ -chores. We show that such a swap preserves that the allocation is MPB. After the swap, both  $i$  and  $\ell$  are added to  $N_H^1$  and removed from  $N_H^2$  and  $N_L$  respectively. This implies that Phase 1 terminates after at most  $n/2$  swaps, after which the allocation is 3-EFX for all agents in  $N_H^2$ .

In Phase 2, Algorithm 5 addresses agents in  $N_H^1$ . An agent  $i \in N_H^1$  has one  $H$ -chore, and earns at most 1 from  $L$ -chores. Once again, Lemma 22 shows that if  $i$  3-EFX-envies an agent  $\ell$ , then  $\ell \in N_L$ . We then transfer the  $H$ -chore from  $i$  to  $\ell$ , and transfer all the chores of  $\ell$  to  $i$ . As before, we argue that such a swap preserves that the allocation is MPB. After the swap,  $i$  gets added to

$N_L$  and removed from  $N_H^1$ , while  $\ell$  is added to  $N_H^1$  and removed from  $N_L$ . Since  $\ell$  is now assigned a single  $H$ -chore,  $\ell$  does not EFX-envy any agent. This implies that Phase 2 terminates after at most  $n$  swaps since the number of agents in  $N_H^1$  who are not 2-EFX strictly decreases. The resulting allocation is 3-EFX for all agents in  $N_H^1$ .

Lastly, we show that throughout the algorithm, agents in  $N_0$  are 3-EFX towards all other agents (Lemma 18), agents in  $N_L$  are 3-EFX towards all other agents (Lemma 25), and those in  $N_H$  are 3-EFX towards agents in  $N_H \cup N_0$  (Lemma 22). Since the algorithm addresses 3-EFX-envy from agents in  $N_H$  towards those in  $N_L$  in at most  $3n/2$  swaps, it terminates with a 3-EFX and fPO allocation.

We now formally prove the above claims. We begin by recording a lemma regarding the MPB ratio  $\alpha_i$  of an agent  $i \in N_L \cup N_H$ .

**Lemma 20.** *Assume  $H \neq \emptyset$ . Then:*

- (i) *For all  $i \in N_L$ ,  $\alpha_i = 1/\rho$ . Moreover for every  $j \in H$ ,  $d_{ij} = k$  and  $j \in MPB_i$ .*
- (ii) *For all  $i \in N_H$ ,  $\alpha_i \in \{\frac{1}{\rho}, \frac{1}{\rho k}\}$ .*
- (iii) *For all  $i \in N_H$ , if  $\mathbf{x}_i \setminus H \neq \emptyset$  then  $\alpha_i = 1/\rho$ .*

*Proof.* Let  $j_0 \in H$  with  $p_j = \rho k$ . For (i), consider  $i \in N_L$ , and let  $j \in \mathbf{x}_i$ . Since  $j \in L$ ,  $p_j = \rho$ . The MPB condition for  $i$  implies  $\frac{d_{ij}}{p_j} \leq \frac{d_{ij_0}}{p_{j_0}}$ . This gives  $kd_{ij} \leq d_{ij_0}$ . Since  $d_{ij}, d_{ij_0} \in \{1, k\}$ , the above inequality must be an equality and  $d_{ij} = 1$  and  $d_{ij_0} = k$ . Thus  $\alpha_i = 1/\rho$  for  $i \in N_L$ . Now consider any  $j' \in H$ . The MPB condition for  $i$  implies  $\alpha_i \leq \frac{d_{ij'}}{p_{j'}}$ . This implies  $d_{ij'} \geq k$ . Since  $d_{ij'} \in \{1, k\}$ , we have  $d_{ij'} = k$  and  $j' \in MPB_i$ .

For (ii), let  $i \in N_H$  and  $j \in \mathbf{x}_i \cap H$ . Then  $\alpha_i = \frac{d_{ij}}{p_j} \in \{\frac{1}{\rho}, \frac{1}{\rho k}\}$ , since  $d_{ij} \in \{1, k\}$  and  $p_j = \rho k$ .

For (iii), consider  $i \in N_H$  with  $j_1 \in \mathbf{x}_i \setminus H$  and  $j_2 \in \mathbf{x}_i \cap H$ . The MPB condition for  $i$  implies  $\frac{d_{ij_1}}{p_{j_1}} = \frac{d_{ij_2}}{p_{j_2}}$ , which gives  $d_{ij_1} = 1$  and  $d_{ij_2} = k$ . Thus  $\alpha_i = 1/\rho$ .  $\square$

We next show that Algorithm 5 maintains the following invariants.

**Lemma 21.** *(Invariants of Alg.5) Let  $(\mathbf{x}, \mathbf{p})$  be an allocation in the run of Algorithm 5. Then:*

- (i)  *$(\mathbf{x}, \mathbf{p})$  is an MPB allocation.*
- (ii) *For all  $i \in N$ ,  $\mathbf{p}(\mathbf{x}_i) \geq \frac{1}{2}$ .*
- (iii) *For all  $i \in N_L$ ,  $\mathbf{p}_{-1}(\mathbf{x}_i) \leq 1$  during Phase 1.*
- (iv) *For all  $i \in N_L$ ,  $\mathbf{p}(\mathbf{x}_i) \leq \frac{4}{3} + \frac{\rho k}{3}$ .*
- (v) *For all  $i \in N_H^1$ ,  $\mathbf{p}_{-1}(\mathbf{x}_i) \leq 1$ .*
- (vi) *For all  $i \in N_H^2$ ,  $\mathbf{p}_{-2}(\mathbf{x}_i) \leq \frac{1}{2}$ .*

*Proof.* We prove this using Lemmas 22, 23 and 24 below.  $\square$

We first prove that some conditions must hold if an agent in  $N_H$  3-EFX-envies another agent.

**Lemma 22.** *Consider an allocation  $(\mathbf{x}, \mathbf{p})$  satisfying the invariants of Lemma 21. Then if  $i \in N_H$  3-EFX-envies  $\ell$ , then  $\alpha_i = 1/\rho$ ,  $\ell \in N_L$ , and  $\mathbf{x}_\ell \subseteq MPB_i$ .*

*Proof.* Consider  $i \in N_H$  who 3-EFX-envies  $\ell \in N$ . We know from Lemma 20 that  $\alpha_i \in \{\frac{1}{\rho}, \frac{1}{\rho k}\}$ . Suppose  $\alpha_i = \frac{1}{\rho k}$ . Then  $d_{ij} = 1$  for all  $j \in \mathbf{x}_i$ . By the contrapositive of Lemma 20 (iii), we get  $\mathbf{x}_i \subseteq H$ . Since  $|\mathbf{x}_i \cap H| \leq 2$ , we get  $|\mathbf{x}_i| \leq 2$ . Thus,  $\max_{j \in \mathbf{x}_i} d_i(\mathbf{x}_i \setminus \{j\}) \leq 1 \leq d_i(\mathbf{x}_\ell)$ , since  $\mathbf{x}_\ell \neq \emptyset$  and the instance is bivalued. Thus,  $i$  is EFX towards  $\ell$  if  $\alpha_i = \frac{1}{\rho k}$ , which implies that  $\alpha_i = 1/\rho$ .

If  $\exists j \in \mathbf{x}_\ell$  such that  $d_{ij} = k$ , then using the constraints on  $\mathbf{p}(\mathbf{x}_i)$  implied by invariants (v) and (vi), we have:

$$d_i(\mathbf{x}_i) = \alpha_i \cdot \mathbf{p}(\mathbf{x}_i) \leq \frac{1}{\rho} \cdot \max\{1 + \rho k, \frac{1}{2} + 2\rho k\} = \max\left\{\frac{1}{\rho} + k, \frac{1}{2\rho} + 2k\right\} < 3k \leq 3d_i(\mathbf{x}_h),$$

where we used  $\rho k > 1/2$  in the penultimate inequality and  $d_i(\mathbf{x}_h) \geq k$  in the final inequality. This shows that  $i$  is 3-EFX towards  $\ell$ . Thus it must be that for all  $j \in \mathbf{x}_\ell$ ,  $d_{ij} = 1$ . The MPB condition for  $i$  implies that  $\alpha_i \leq d_{ij}/p_j$ , showing that  $p_j \leq \rho$ . Lemma 14 implies that  $p_j = \rho$  for all  $j \in \mathbf{x}_\ell$ . Thus  $\ell \in N_L$ . Moreover, for any  $j \in \mathbf{x}_\ell$ ,  $\alpha_i = d_{ij}/p_j$ , and hence  $\mathbf{x}_\ell \subseteq MPB_i$ .  $\square$

The next two lemmas establish the invariants claimed by Lemma 21.

**Lemma 23.** *The invariants of Lemma 21 are maintained during Phase 1 of Algorithm 5.*

*Proof.* We prove the statement inductively. We first show that the invariants hold at  $(\mathbf{x}^0, \mathbf{p})$ . Invariants (i), (ii), (iii), (v) and (vi) follow from Lemma 15. For (iv), note that for any  $i \in N_L$ , we have  $\mathbf{p}_{-1}(\mathbf{x}_i^0) \leq 1$ . Thus  $\mathbf{p}(\mathbf{x}_i^0) \leq 1 + \rho < 1 + \rho k$ .

Suppose the invariants hold at an allocation  $(\mathbf{x}, \mathbf{p})$  during Phase 1. Consider a Phase 1 swap involving agents  $i \in N_H^2$  and  $\ell \in N$ . Given that Algorithm 5 performed the swap,  $i$  must 3-EFX-envy  $\ell$ . Lemma 22 implies that  $\ell \in N_L$  and hence  $\mathbf{x}_\ell \in L$ . As per Algorithm 5, if  $\mathbf{p}(\mathbf{x}_\ell) > 1$ , then  $S = \{j_1\}$  for some  $j_1 \in \mathbf{x}_\ell$ , otherwise  $S = \emptyset$ . Let  $j \in \mathbf{x}_i \cap H$ .

Let  $\mathbf{x}'$  be the resulting allocation. Thus  $\mathbf{x}'_i = \mathbf{x}_i \setminus \{j\} \cup S$ ,  $\mathbf{x}'_\ell = \mathbf{x}_\ell \setminus S \cup \{j\}$ , and  $\mathbf{x}'_h = \mathbf{x}_h$  for all  $h \notin \{i, \ell\}$ . We show that the invariants hold at  $(\mathbf{x}', \mathbf{p})$ . Since a Phase 1 step removes agents  $i$  and  $\ell$  from  $N_H^2$  and  $N_L$  respectively, invariants (iii), (iv), (vi) continue to hold. For the rest, observe:

- (i)  $(\mathbf{x}', \mathbf{p})$  is on MPB. This is because Lemma 22 implies  $S \subseteq \mathbf{x}_\ell \subseteq MPB_i$ , showing  $\mathbf{x}'_i \subseteq MPB_i$ . Since  $\ell \in N_L$  at  $(\mathbf{x}, \mathbf{p})$  and  $j \in H$ , Lemma 20 shows  $j \in MPB_\ell$  and hence  $\mathbf{x}'_\ell \subseteq MPB_\ell$ .
- (ii) Since  $|\mathbf{x}'_i \cap H| = |\mathbf{x}_i \cap H| = 1$ , we have  $\mathbf{p}(\mathbf{x}'_i) \geq \rho k$  and  $\mathbf{p}(\mathbf{x}'_\ell) \geq \rho k$ . Invariant (ii) follows by noting that  $\rho k > 1/2$ .
- (v) For agent  $i$ , note that  $\mathbf{p}_{-1}(\mathbf{x}'_i) \leq \mathbf{p}(\mathbf{x}_i \setminus \{j\} \cup S) = \mathbf{p}_{-1}(\mathbf{x}_i \setminus \{j\}) + \mathbf{p}(S)$ . Invariant (iii) implies  $\mathbf{p}_{-1}(\mathbf{x}_i \setminus \{j\}) = \mathbf{p}_{-2}(\mathbf{x}_i) \leq 1/2$ , and  $\mathbf{p}(S) \leq 1/2$  by construction. Hence  $\mathbf{p}_{-1}(\mathbf{x}'_i) \leq 1$ .  
For agent  $\ell$ , note that  $\mathbf{p}_{-1}(\mathbf{x}'_\ell) = \mathbf{p}(\mathbf{x}_\ell \setminus S) = \mathbf{p}(\mathbf{x}_\ell) - \mathbf{p}(S)$ . If  $\mathbf{p}(\mathbf{x}_\ell) \leq 1$ , then  $S = \emptyset$ , implying that  $\mathbf{p}_{-1}(\mathbf{x}'_\ell) \leq 1$ . On the other hand suppose  $\mathbf{p}(\mathbf{x}_\ell) > 1$ . Since invariant (v) holds at  $(\mathbf{x}, \mathbf{p})$ , we have  $\mathbf{p}_{-1}(\mathbf{x}_\ell) \leq 1$ , which gives  $\mathbf{p}(\mathbf{x}_\ell) \leq 1 + \rho$ . With  $\mathbf{p}(S) = p_{j_1} = \rho$ , we obtain  $\mathbf{p}_{-1}(\mathbf{x}'_\ell) = \mathbf{p}(\mathbf{x}_\ell) - \mathbf{p}(S) \leq 1$ .

Since the swap does not affect any  $h \notin \{i, \ell\}$ , the invariants continue to hold for  $h$  after the swap. By induction, we have shown that the invariants of Lemma 21 hold after any Phase 1 swap.  $\square$

**Lemma 24.** *The invariants of Lemma 21 are maintained during Phase 2 of Algorithm 5. Moreover, agents in  $N_H^2$  remain 3-EFX towards other agents.*



*Proof.* We prove the statement inductively. Lemma 23 shows the invariants hold at the end of Phase 1. Suppose the invariants hold at an allocation  $(\mathbf{x}, \mathbf{p})$  during Phase 2. Consider a Phase 2 swap involving agents  $i \in N_H^1$  and  $\ell \in N$ . Given that Algorithm 5 performed the swap,  $i$  must 3-EFX-envy  $\ell$ . Lemma 22 implies that  $\ell \in N_L$  and hence  $\mathbf{x}_\ell \in L$ . Let  $j \in \mathbf{x}_i \cap H$ .

Let  $\mathbf{x}'$  be the resulting allocation. Thus  $\mathbf{x}'_i = \mathbf{x}_i \setminus \{j\} \cup \mathbf{x}_\ell$ ,  $\mathbf{x}'_\ell = \{j\}$ , and  $\mathbf{x}'_h = \mathbf{x}_h$  for all  $h \notin \{i, \ell\}$ . We now show that the invariants hold at  $(\mathbf{x}', \mathbf{p})$ . Since we are in Phase 2, invariant (iii) does not apply, and since Phase 2 swaps do not alter the allocation of agents in  $N_H^2$ , invariant (vi) continues to hold. For the rest, observe:

- (i)  $(\mathbf{x}', \mathbf{p})$  is on MPB. This is because Lemma 22 implies  $\mathbf{x}_\ell \subseteq MPB_i$ , showing  $\mathbf{x}'_i \subseteq MPB_i$ . Since  $\ell \in N_L$  at  $(\mathbf{x}, \mathbf{p})$  and  $j \in H$ , Lemma 20 shows  $j \in MPB_\ell$  and hence  $\mathbf{x}'_\ell \subseteq MPB_\ell$ .
- (ii) For agent  $i$ ,  $\mathbf{p}(\mathbf{x}'_i) \geq \mathbf{p}(\mathbf{x}_\ell) \geq 1/2$ , since invariant (ii) holds in  $(\mathbf{x}, \mathbf{p})$ . For agent  $\ell$ , note that  $\mathbf{p}(\mathbf{x}'_\ell) = p_j = \rho k > 1/2$ .
- (iv) We want to show  $\mathbf{p}(\mathbf{x}'_i) \leq 1 + \rho k$ . To see this note that since  $i$  3-EFX-envies  $\ell$  in  $\mathbf{x}$ ,  $i$  must 3-pEF-envy  $\ell$  in  $(\mathbf{x}, \mathbf{p})$ . Thus  $\mathbf{p}(\mathbf{x}_i) > 3 \cdot \mathbf{p}(\mathbf{x}_\ell)$ . Now  $\mathbf{p}_{-1}(\mathbf{x}_i) \leq 1$  due to invariant (v), which shows  $\mathbf{p}(\mathbf{x}_i) \leq 1 + \rho k$ . We therefore obtain  $\mathbf{p}(\mathbf{x}_\ell) < \frac{\mathbf{p}(\mathbf{x}_i)}{3} = \frac{1+\rho k}{3}$ .  
Now  $\mathbf{p}(\mathbf{x}'_i) = \mathbf{p}(\mathbf{x}_i \setminus \{j\}) + \mathbf{p}(\mathbf{x}_\ell) \leq 1 + \frac{1+\rho k}{3}$ , where we once again used  $\mathbf{p}_{-1}(\mathbf{x}_i) = \mathbf{p}(\mathbf{x}_i \setminus \{j\}) = 1$ . The invariant thus follows.
- (v) Note that  $\ell \in N_H^1$  in  $(\mathbf{x}', \mathbf{p})$ , and  $\mathbf{p}_{-1}(\mathbf{x}_\ell) = 0 < 1$ .

The swap does not affect an agent  $h \notin \{i, \ell\}$  and hence the invariants continue to hold for  $h$  after the swap. By induction, we conclude that the invariants of Lemma 21 hold after any Phase 2 swap.

We now show that  $i \in N_H^2$  cannot 3-EFX-envy an agent  $\ell \in N$ . Lemma 22 implies that  $\ell \in N_L$  and hence  $\mathbf{x}_\ell \subseteq L$ . Let  $\mathbf{x}^1$  be the allocation at the end of Phase 1. Note that the bundle  $\mathbf{x}_\ell \subseteq L$  is obtained via a series of Phase 2 swaps initiated with some agent  $\ell_1$  in  $(\mathbf{x}^1, \mathbf{p})$ . Here,  $\ell_1 \in N_L$  at  $(\mathbf{x}^1, \mathbf{p})$ . Thus  $\mathbf{x}_\ell \supseteq \mathbf{x}_{\ell_1}^1$ . Agent  $i \in N_H^2$  did not 3-EFX-envy  $\ell_1$  in  $\mathbf{x}^1$ , otherwise Algorithm 5 would have performed a Phase 1 swap between agent  $i$  and  $\ell_1$ . Since  $\mathbf{x}_i = \mathbf{x}_i^1$  as Algorithm 5 does not alter allocation of agents in  $N_H^2$  and  $\mathbf{x}_\ell \supseteq \mathbf{x}_{\ell_1}^1$ ,  $i$  will not 3-EFX-envy  $\ell$  in  $\mathbf{x}$  either. Thus, all agents in  $N_H^2$  continue to remain 3-EFX during Phase 2.  $\square$

We require one final lemma showing that  $N_L$  agents do not 3-EFX-envy any other agent.

**Lemma 25.** *At any allocation  $(\mathbf{x}, \mathbf{p})$  in the run of Algorithm 5,  $\mathbf{x}$  is 3-EFX for every agent in  $N_L$ .*

*Proof.* Lemma 19 shows that the initial allocation  $\mathbf{x}^0$  is 3-EFX for agents in  $N_L$ . Let  $\mathbf{x}$  be the earliest allocation in the run of Algorithm 5 in which an agent  $i \in N_L$  3-EFX-envies another agent  $h \in N$ . Using  $\alpha_i = 1/\rho$  from Lemma 20, the bound on  $\mathbf{p}(\mathbf{x}_i)$  from Lemma 21 (iv), and  $\rho k > 1/2$ , we note:

$$d_i(\mathbf{x}_i) = \alpha_i \cdot \mathbf{p}(\mathbf{x}_i) \leq \frac{1}{\rho} \cdot \left( \frac{4}{3} + \frac{\rho k}{3} \right) < 3k. \quad (3)$$

Thus if  $\exists j \in \mathbf{x}_h$  s.t.  $d_{ij} = k$ , then by (3),  $d_i(\mathbf{x}_i) < 3k \leq d_i(\mathbf{x}_h)$ , showing that  $i$  does not 3-EFX-envy  $h$  in  $\mathbf{x}$ . Hence it must be that for all  $j \in \mathbf{x}_h$ ,  $d_{ij} = 1$ . This also implies  $\mathbf{x}_h \subseteq MPB_i$ , since  $\alpha_i = 1/\rho = d_{ij}/p_j$  for any  $j \in \mathbf{x}_h$ . We now consider three cases based on the category of  $h$ .

- $h \in N_0$ . For  $j \in \mathbf{x}_h$ , the MPB condition of  $i$  implies  $\alpha_i \leq d_{ij}/p_j$ , implying  $d_{ij} > p_j/\rho$ . Since  $h \in N_0$ , we have  $j \in M'$  and  $p_j > \rho$ . Thus  $d_{ij} = k$  for  $j \in \mathbf{x}_h$ , which is a contradiction.

- $h \in N_H$ . By definition of  $N_H$ ,  $\exists j \in \mathbf{x}_h$  s.t.  $j \in H$ . Since  $i \in N_L$ , by Lemma 20 (i) we get  $d_{ij} = k$ , which is a contradiction.
- $h \in N_L$ . Since  $\mathbf{x}^0$  is 3-EFX for agents in  $N_L$ , and Phase 1 swaps only remove agents from  $N_L$ , it cannot be that  $i$  starts 3-EFX-envying  $h \in N_L$  during Phase 1. Let  $\mathbf{x}'$  be the preceding allocation, at which Algorithm 5 performed a Phase 2 swap. Since  $\mathbf{x}$  is the earliest allocation in which  $i$  3-EFX-envies  $\ell$ , it must be that in  $\mathbf{x}'$ , agent  $i$  was in  $N_H^1$  and was involved with a Phase 2 swap with another agent  $\ell \in N_L$ . Since  $\mathbf{x}_h \subseteq MPB_i$  and Algorithm 5 did *not* perform a swap between agents  $i$  and  $h$  in the allocation  $\mathbf{x}'$ , we must have  $\mathbf{p}(\mathbf{x}'_\ell) \leq \mathbf{p}(\mathbf{x}'_h)$  by the choice of  $\ell$  at  $(\mathbf{x}', \mathbf{p})$ .

Note that  $\mathbf{x}_i = (\mathbf{x}'_i \setminus H) \cup \mathbf{x}'_\ell$ . By Lemma 21 (i), we know  $\mathbf{p}_{-1}(\mathbf{x}'_i) = \mathbf{p}(\mathbf{x}'_i \setminus H) \leq 1$ . Thus:

$$\mathbf{p}(\mathbf{x}_i) = \mathbf{p}(\mathbf{x}'_i \setminus H) + \mathbf{p}(\mathbf{x}'_\ell) \leq 1 + \mathbf{p}(\mathbf{x}'_\ell) \leq 3\mathbf{p}(\mathbf{x}_h),$$

where the last inequality uses  $\mathbf{x}_h = \mathbf{x}'_h$  and  $\mathbf{p}(\mathbf{x}_h) \geq 1/2$ . Thus  $i$  is actually 3-EFX towards  $h$ .

Since these cases are exhaustive, we conclude that it is not possible for an agent  $i \in N_L$  to 3-EFX-envy any other agent during the course of Algorithm 5.  $\square$

We are now in a position to summarize and conclude our analysis of Algorithm 5.

**Theorem 13.** *Given an ER equilibrium of a bivalued instance with  $m > 2n$  and chore earning limit  $\beta = 1/2$ , Algorithm 5 returns a 3-EFX and fPO allocation in polynomial time.*

*Proof.* Let  $(\mathbf{x}^0, \mathbf{p})$  be the initial allocation obtained by using Algorithm 1 on the ER equilibrium, and let  $\rho = \min_j p_j$  be the minimum payment. Lemma 17 shows that  $\mathbf{x}^0$  is 3-EFX if  $\rho k \leq 1/2$ , hence we assume otherwise. Since any allocation  $(\mathbf{x}, \mathbf{p})$  during the course of Algorithm 5 satisfies invariant (ii) of Lemma 21, Lemma 18 implies that  $\mathbf{x}$  is 3-EFX for agents in  $N_0$ .

Lemma 25 shows that any allocation  $\mathbf{x}$  in the course of Algorithm 5 is 3-EFX for agents in  $N_L$ . Any potential EFX-envy is therefore from some agent  $i \in N_H$ . Lemma 22 shows that if  $i \in N_H$  is not 3-EFX towards  $\ell$ , then  $\ell \in N_L$ . If  $i \in N_H^2$ ,  $i$  participates in a Phase 1 swap with agent  $\ell$ , after which  $i$  and  $\ell$  get removed from  $N_H^2$  and  $N_L$  respectively. This implies that Phase 1 terminates after at most  $n/2$  swaps, and the resulting allocation is 3-EFX for all agents in  $N_H^2$ . If  $i \in N_H^1$ ,  $i$  participates in a Phase 2 swap with agent  $\ell$ , after which  $\ell$  is added to  $N_H^1$  and is assigned a single chore and  $\ell$  does not have EFX-envy. This implies that Phase 2 terminates after at most  $n$  swaps, since the number of agents in  $N_H^1$  who are not 3-EFX strictly decreases. The resulting allocation is 3-EFX for all agents in  $N_H^1$ . Lemma 24 also shows that Phase 2 swaps do not cause  $N_H^2$  agents to start 3-EFX-envying any agent in  $N_L$ . Thus the allocation on termination of Algorithm 5 is 3-EFX. By invariant (i) of Lemma 21,  $\mathbf{x}$  is also fPO. Since there are at most  $3n/2$  swaps, Algorithm 5 terminates in polynomial time.  $\square$

## 5.2 EFX and PO for $m \leq 2n$

We design Algorithm 7 for bivalued instances with  $m \leq 2n$ . Algorithm 7 begins with a balanced allocation computed using Algorithm 2, and essentially runs Algorithm 5. Since the number of chores is limited, a careful analysis shows that the guarantee of the resulting allocation can be improved to EFX and fPO.

**Theorem 14.** *Given a bivalued instance with  $m \leq 2n$ , Algorithm 7 returns an EFX and fPO allocation in polynomial time.*

Since the main ideas of the analysis are similar to those presented in Section 5.1, we defer this section in a self-contained Appendix E. Surprisingly, Example 6 shows that if we slightly generalize the class to 2-ary instances, an  $\alpha$ -EFX and fPO allocation need not exist.

## 6 Existence of Earning-Restricted Equilibria

We prove Theorem 2 in this section.

**Theorem 2.** *Every ER instance  $(N, M, D, e, c)$  satisfying  $\sum_i e_i \leq \sum_j c_j$  admits an ER equilibrium.*

In what follows, we assume ER instances satisfy the feasible earning condition  $\sum_i e_i \leq \sum_j c_j$ . We prove Theorem 2 by designing a linear complementarity problem (LCP) whose solution corresponds to an ER equilibrium. We begin with some background on LCPs.

### 6.1 Linear complementarity problems and Lemke's scheme

A Linear Complementary Problem (LCP) is a generalization of linear programming (LP) with complementary slackness conditions: given a matrix  $A \in \mathbb{R}^{n \times n}$  and a  $\mathbf{b} \in \mathbb{R}^n$ , the problem is:

$$\text{LCP}(A, \mathbf{b}): \text{Find } \mathbf{y} \geq \mathbf{0} \text{ such that } A\mathbf{y} \leq \mathbf{b}, \text{ and } y_i \cdot (A\mathbf{y} - \mathbf{b})_i = 0 \text{ for all } i \in [n]. \quad (4)$$

We use the shorthand notation  $(A\mathbf{y})_i \leq b_i \perp y_i$  to represent the constraints  $A\mathbf{y} \leq \mathbf{b}$ ,  $y_i \geq 0$ , and  $y_i \cdot (A\mathbf{y} - \mathbf{b})_i$ . If  $\mathbf{b} \geq \mathbf{0}$ , then  $\mathbf{y} = \mathbf{0}$  is a trivial solution to the LCP. If  $\mathbf{b} \not\geq \mathbf{0}$ , then the LCP may not have a solution; indeed, LCPs are general enough to capture NP-hard problems [19].

**Lemke's scheme.** Let  $\mathcal{P} = \{\mathbf{y} \in \mathbb{R}^n : A\mathbf{y} \leq \mathbf{b}, \mathbf{y} \geq \mathbf{0}\}$ . We assume that the polyhedron  $\mathcal{P}$  is non-degenerate, i.e., exactly  $n - d$  constraints hold with equality on any  $d$ -dimensional face of  $\mathcal{P}$ . With this assumption, each solution to (4) corresponds to a vertex of  $\mathcal{P}$  since exactly  $n$  equalities must be satisfied. Lemke's scheme finds such a vertex solution by working with an *augmented* LCP which adds a scalar variable  $z$  to  $\text{LCP}(A, \mathbf{b})$ , resulting in the following program:

$$\text{Augmented LCP}(A, \mathbf{b}): z \geq 0; \text{ and } A\mathbf{y} - z \cdot \mathbf{1} \leq \mathbf{b}, \text{ and } y_i \cdot ((A\mathbf{y} - \mathbf{b})_i - z) = 0 \text{ for all } i \in [n]. \quad (5)$$

Note that  $(\mathbf{y}, z = 0)$  is a solution to (5) iff  $\mathbf{y}$  is solution to (4). Let  $\mathcal{P}' = \{(\mathbf{y}, z) \in \mathbb{R}^{n+1} : A\mathbf{y} - z \cdot \mathbf{1} \leq \mathbf{b}, \mathbf{y} \geq \mathbf{0}, z \geq 0\}$ . Assuming  $\mathcal{P}'$  is non-degenerate, a solution to (5) still satisfies  $n$  constraints of  $\mathcal{P}'$  with equality. Since  $\mathcal{P}'$  is  $(n + 1)$ -dimensional, this means the set of solutions  $S$  is a subset of the 1-skeleton of  $\mathcal{P}'$ , i.e., edges (1-dimensional faces) and vertices (0-dimensional faces). Moreover,  $\mathbf{y}$  is a solution of (4) iff  $(\mathbf{y}, 0)$  is a vertex of  $\mathcal{P}'$ .

The set  $S$  of solutions to the augmented LCP has some important structural properties. We say that the label  $i$  is present at  $(\mathbf{y}, z) \in \mathcal{P}'$  if  $y_i = 0$  or  $(A\mathbf{y})_i - z = b_i$ . Every solution in  $S$  is *fully labeled* where all the labels are present. A solution  $s \in S$  contains *double label*  $i$  if  $(A\mathbf{y})_i - z = b_i$  for  $i \in [n]$ . Since there are only two ways to relax the double label while keeping all other labels, there are two edges of  $S$  incident to  $s$ . The above observations imply that  $S$  consists of paths and cycles. Clearly, any solution  $s$  to (5) with  $z = 0$  contains no double labels. Relaxing  $z = 0$  gives the unique edge incident to  $s$  at this vertex. We note that some of the edges in  $S$  are unbounded. An unbounded edge of  $S$  incident to vertex  $(\mathbf{y}^*, z^*)$  with  $z^* > 0$  is called a ray. Formally, a ray  $R$  has the form  $R = \{(\mathbf{y}^*, z^*) + \alpha \cdot (\mathbf{y}', z') : \alpha \geq 0\}$ , where  $(\mathbf{y}', z') \neq \mathbf{0}$  solves (5) with  $\mathbf{b} = \mathbf{0}$ . The *primary ray* is the ray  $\{(\mathbf{0}, z) : z \geq |\min_i b_i|\}$ , which contains solutions with  $\mathbf{y} = \mathbf{0}$  and  $z$  sufficiently large to satisfy (5). All other rays are called *secondary rays*.

Starting from the primary ray, Lemke's scheme follows a path on the 1-skeleton of  $\mathcal{P}'$  with a guarantee that it never revisits a vertex. If a vertex  $s$  is non-degenerate, i.e., has a *unique* double label, then Lemke's scheme *pivots* by relaxing one of the two constraints and travelling along the edge of  $\mathcal{P}'$  to the next vertex solution. Therefore, if the vertices are non-degenerate, Lemke's scheme eventually either reaches a vertex with  $z = 0$  (which is a solution of the original LCP (4)), or it ends up on a secondary ray. In the latter case, the algorithm fails to find a solution; in fact the problem may not have a solution. Note that it suffices to introduce  $z$  in the  $(A\mathbf{y})_i \leq b_i$  constraint only if  $b_i < 0$ , without changing the role of  $z$ .

## 6.2 Basic LCP for ER equilibrium

We first capture ER equilibria in an instance  $(N, M, D, e, c)$  via the following LCP with variables  $\mathbf{p} = \{p_j\}_{j \in M}$ ,  $\mathbf{q} = \{q_{ij}\}_{i \in N, j \in M}$ ,  $\mathbf{r} = \{r_i\}_{i \in N}$ , and  $\boldsymbol{\beta} = \{\beta_j\}_{j \in M}$ .

$$\forall i \in N : \quad e_i \leq \sum_j q_{ij} \quad \perp \quad r_i \quad (6a)$$

$$\forall j \in M : \quad \sum_i q_{ij} \leq p_j - \beta_j \quad \perp \quad p_j \quad (6b)$$

$$\forall i \in N, j \in M : \quad p_j \leq d_{ij} r_i \quad \perp \quad q_{ij} \quad (6c)$$

$$\forall j \in M : \quad p_j - \beta_j \leq c_j \quad \perp \quad \beta_j \quad (6d)$$

*Notation.* We follow the following notational convention. For a constraint labelled  $L$ , we represent its complementarity constraint expressing the non-negativity of a variable by  $L'$ . For example, (6a) is the constraint  $e_i \leq \sum_j q_{ij}$  for agent  $i$ , and (6a)' is the constraint  $r_i \geq 0$ .

*Interpretation of the LCP.* In the above LCP,  $p_j$  denotes the payment of chore  $j$ ,  $q_{ij}$  denotes the earning of agent  $i$  from chore  $j$ ,  $r_i$  denotes the reciprocal of the MPB ratio of agent  $i$ , and  $\beta_j$  denotes the excess payment of chore  $j$ , i.e.,  $q_j := p_j - \beta_j$  is the total earning from chore  $j$ .

Constraint (6a) imposes that each agent  $i$  earns at least their earning requirement of  $e_i$ . Constraint (6b) imposes that the earning  $\sum_i q_{ij}$  from each chore  $j$  is at most  $q_j = p_j - \beta_j$ . Constraint (6c) enforces the MPB condition. Constraint (6d) enforces the earning restriction on each chore. Constraints (6a)', (6b)', (6c)', (6d)' enforce non-negativity of the LCP variables. The constraints (6a)-(6d) and (6a)'-(6d)' together define LCP(6).

We now demonstrate the correspondence between the ER equilibria and certain solutions to LCP(6).

**Lemma 26.** *Any ER equilibrium can be used to construct a solution to LCP(6).*

*Proof.* Let  $(\mathbf{x}, \mathbf{p})$  be an ER equilibrium. Let  $\alpha_i$  be the MPB ratio of agent  $i$  in  $(\mathbf{x}, \mathbf{p})$ . Since all  $d_{ij} > 0$ , we have  $\alpha_i > 0$  for all  $i \in N$ . Define  $r_i = \alpha_i^{-1}$ . Since  $\mathbf{p} > 0$  in any ER equilibrium, we have that  $r_i > 0$  for all  $i \in N$ . Let  $q_{ij} = p_j x_{ij}$  be the earning of agent  $i$  from chore  $j$ , and let  $q_j = \sum_i q_{ij}$ . Finally, define  $\beta_j = \max\{0, p_j - c_j\}$  for each  $j \in M$ . We show that  $(\mathbf{p}, \mathbf{q}, \mathbf{r}, \boldsymbol{\beta})$  is a solution to LCP(6) as follows:

(Constraint (6a)) For all  $i \in N$ ,  $e_i = \sum_j q_{ij}$  since  $(\mathbf{x}, \mathbf{p})$  is an ER equilibrium (Def. 4 (i)). Also for all  $i \in N$ ,  $r_i > 0$ .

(Constraints (6c)) Since  $\mathbf{x}$  is an MPB allocation, for all  $i \in N, j \in M$  we have  $d_{ij}/p_j \geq \alpha_i$ , with equality if  $x_{ij} > 0$ . We then note that  $\alpha_i^{-1} = r_i$  and  $q_{ij} = p_j x_{ij}$ .

(Constraints (6b), (6d)) We consider two cases for each chore  $j \in M$ . If  $p_j \leq c_j$ , then  $q_j = p_j$  and  $\beta_j = 0$ . Otherwise,  $p_j > c_j$ , and we have  $q_j = c_j$  and  $\beta_j = p_j - c_j$ . In both cases, the complementarity constraints (6b) and (6d) hold for each  $j \in M$ .  $\square$

**Lemma 27.** Any solution  $(\mathbf{p}, \mathbf{q}, \mathbf{r}, \boldsymbol{\beta})$  to LCP(6) with  $\mathbf{p} > 0$  can be used to construct an ER equilibrium  $(\mathbf{x}, \mathbf{p}, \mathbf{q})$ .

*Proof.* We first argue that  $\forall i \in N : r_i > 0$ . Suppose  $r_i = 0$  for some  $i \in N$ . Then constraint (6c) implies  $p_j = 0$  for all  $j \in M$ . In turn, with constraint (6b) this implies that  $q_{ij} = 0$  for all  $i, j$ . Then constraint (6a) cannot be satisfied for agent  $i$  since  $e_i > 0$ , leading to the contradiction that  $(\mathbf{p}, \mathbf{q}, \mathbf{r}, \boldsymbol{\beta})$  is a solution to LCP(6). Therefore  $\forall i \in N : r_i > 0$ .

Define an allocation  $\mathbf{x}$  as  $x_{ij} = q_{ij}/p_j$  and let  $q_j = \sum_i q_{ij}$ . We show  $(\mathbf{x}, \mathbf{p}, \mathbf{q})$  is an ER equilibrium by showing that it satisfies the conditions of Definition 4 as follows.

(Agents) The complementarity constraint (6c) implies that  $(\mathbf{x}, \mathbf{p})$  is an MPB allocation with  $1/r_i$  being the MPB ratio of agent  $i$ . Moreover, constraint (6a) implies  $e_i = \sum_j q_{ij}$  for all  $i \in N$ , since  $r_i > 0$  for all  $i \in N$ .

(Chores) Since  $p_j > 0$  for all  $j \in M$  by assumption, we have  $q_j = p_j - \beta_j$ . We consider two cases for each chore  $j \in M$ . If  $\beta_j = 0$ , then  $q_j = p_j$  (from (6b)), and also  $q_j \leq c_j$  (from (6d)). Thus  $q_j = \min\{p_j, c_j\}$ . Otherwise,  $\beta_j > 0$  and  $q_j = p_j - \beta_j < p_j$  (from (6b)) and  $q_j = c_j$  (from (6d)). Thus  $q_j = \min\{p_j, c_j\}$  in this case as well.  $\square$

### 6.3 Main LCP for ER equilibrium

However, Lemke's scheme for LCP(6) may not converge to a solution with  $\mathbf{p} > 0$ . We address this issue by performing a change of variables. First, we show that that chore payments can be assumed to be upper bounded by some constant  $P$ .

**Lemma 28.** For every ER instance  $(N, M, D, e, c)$ , there exists a constant  $P$  such that for every ER equilibrium  $(\mathbf{x}, \mathbf{p}, \mathbf{q})$  there exists a scaled ER equilibrium  $(\mathbf{x}', \mathbf{p}', \mathbf{q})$  s.t.  $\forall j \in M, p'_j \leq P$ .

*Proof.* We set the constant  $P$  to  $\frac{d_{max}}{d_{min}} \cdot c_{max}$ , where  $d_{max} = \max_{i,j} d_{ij}$ ,  $d_{min} = \min_{i,j} d_{ij} > 0$  and  $c_{max} = \max_j c_j$ . If there exists some chore  $j \in \mathbf{x}_i$  with  $p_j \leq c_j$ , then the MPB condition implies that for every chore  $k \in M$ ,  $\frac{d_{ij}}{p_j} \leq \frac{d_{ik}}{p_k}$ , showing that  $p_k \leq \frac{d_{max}}{d_{min}} \cdot p_j \leq P$ .

Therefore, suppose  $p_j > c_j$  for all  $j \in M$ . Then we uniformly decrease the payments as  $p'_j = \frac{p_j}{\min_k p_k / c_k}$ . We therefore have  $p'_j \geq c_j$  for every  $j$ , but  $p'_k = c_k$  for some  $k$ . We update the allocation  $\mathbf{x}'$  s.t.  $x'_{ij} \cdot p'_j = x_{ij} \cdot p_j$  for all  $i, j$ , ensuring that the earning vector of  $(\mathbf{x}', \mathbf{p}')$  stays the same as  $\mathbf{q}$ . Since payments are decreased uniformly and  $x'_{ij} > 0$  iff  $x_{ij} > 0$ ,  $(\mathbf{x}', \mathbf{p}')$  is MPB. Thus  $(\mathbf{x}', \mathbf{p}', \mathbf{q})$  is an ER equilibrium. The MPB condition for an agent  $i$  s.t.  $x_{ik} > 0$  implies that  $\frac{d_{ik}}{p_k} \leq \frac{d_{ij}}{p_j}$  for any  $j \in M$ . Thus, for any  $j \in M$ ,  $p'_j \leq \frac{d_{ij}}{d_{ik}} \cdot p'_k \leq \frac{d_{max}}{d_{min}} \cdot c_{max} = P$ , as desired.  $\square$

The upper bound  $P$  on payments implies an upper bound  $R$  on the reciprocal of the MPB ratios. Let  $R$  be chosen so that  $R \cdot \min_{i,j} d_{ij} > P$ . We replace variable  $p_j$  with  $P - p_j$  and  $r_i$  with  $R - r_i$ , while keeping the complementary constraints the same. Finally, we uniformly scale the input parameters  $e$  and  $c$  and obtain the following LCP.

$$\forall i \in N : \quad e_i \cdot \frac{\sum_j (P - p_j - \beta_j)}{\sum_h e_h} \leq \sum_j q_{ij} \quad \perp \quad r_i \quad (7a)$$

$$\forall j \in M : \quad \sum_i q_{ij} \leq (P - p_j - \beta_j) \quad \perp \quad p_j \quad (7b)$$

$$\forall i \in N, j \in M : \quad P - p_j \leq d_{ij}(R - r_i) \quad \perp \quad q_{ij} \quad (7c)$$

$$\forall j \in M : \quad P - p_j - \beta_j \leq c_j \cdot \frac{\sum_k (P - p_k - \beta_k)}{\sum_i e_i} \quad \perp \quad \beta_j \quad (7d)$$

Similar to Lemma 27, we first establish a correspondence between certain solutions to LCP(7) and ER equilibria. We call a solution  $(\mathbf{p}, \mathbf{q}, \mathbf{r}, \boldsymbol{\beta})$  'good' if  $\forall j \in M : p_j + \beta_j < P$  and  $\forall i \in N : r_i < R$ .

**Lemma 29.** *Any good solution of LCP(7) can be used to construct an ER equilibrium with all payments at most  $P$ , and vice versa.*

*Proof.* Let  $\mathbf{v} = (\mathbf{p}, \mathbf{q}, \mathbf{r}, \boldsymbol{\beta})$  be a good solution of LCP(7). Let  $Q := \frac{\sum_j (P - p_j - \beta_j)}{\sum_i e_i}$ . Since  $\mathbf{v}$  is good,  $Q > 0$ . We first show that for all  $i \in N, r_i > 0$ . For the sake of contradiction, let  $r_i = 0$  for some  $i \in N$ . Since  $d_{ij}R > P$ , constraint (7c) is not tight, which implies due to complementarity that  $q_{ij} = 0$  for all  $j \in M$ . Hence  $\sum_j q_{ij} = 0$ , which implies that (7a) cannot hold, as  $e_i \cdot Q > 0$ . Thus,  $\forall i : r_i > 0$ . By complementarity, this means constraints (7a) must be tight:  $\forall i \in N : e_i = \sum_j q_{ij}/Q$ .

We next show that constraints (7b) must be tight. Suppose for some  $j \in M, \sum_i q_{ij} < P - p_j - \beta_j$ . Using inequalities (7a) and (7b), we have:

$$\sum_i e_i \cdot Q = \sum_i \sum_j q_{ij} = \sum_j \sum_i q_{ij} < \sum_j (P - p_j - \beta_j) = \sum_i e_i \cdot Q,$$

which is a contradiction. Thus for all  $j \in M, \sum_i q_{ij} = P - p_j - \beta_j$ .

For  $i \in N$  and  $j \in M$ , define:

$$\hat{p}_j := \frac{P - p_j}{Q}, \quad \hat{q}_{ij} := \frac{q_{ij}}{Q}, \quad \hat{q}_j := \frac{P - p_j - \beta_j}{Q}, \quad \hat{x}_{ij} = \frac{\hat{q}_{ij}}{e_i}$$

We show  $(\hat{\mathbf{x}}, \hat{\mathbf{p}}, \hat{\mathbf{q}})$  is an ER equilibrium by showing it satisfies the conditions of Definition 4.

(Agents) Since  $e_i = \sum_j q_{ij}/Q$ , we have  $e_i = \sum_j \hat{q}_{ij}$  for all  $i \in N$ . Moreover the complementarity constraint (7c) implies  $(\hat{\mathbf{x}}, \hat{\mathbf{p}})$  is an MPB allocation and  $Q/(R - r_i)$  is the MPB ratio of agent  $i$ .

(Chores) Since  $\sum_i q_{ij} = P - p_j - \beta_j$ , we have  $\hat{q}_j = \sum_i \hat{q}_{ij}$  for each  $j \in M$ . We consider two cases for each  $j \in M$ . If  $\beta_j = 0$ , then  $\hat{q}_j = \hat{p}_j$  (by definition), and also  $\hat{q}_j \leq c_j$  (from (7d)). Thus  $\hat{q}_j = \min\{\hat{p}_j, c_j\}$ . Otherwise  $\beta_j > 0$ , and  $\hat{q}_j = \hat{p}_j - \beta_j/Q < \hat{p}_j$  (by definition) and  $\hat{q}_j = c_j$  (from (7d)). Thus  $\hat{q}_j = \min\{\hat{p}_j, c_j\}$  in this case as well.

Thus a good solution of LCP(7) can be used to construct an ER equilibrium. In the other direction, the argument of Lemma 26 with the appropriate change of variables shows that an ER equilibrium with payments at most  $P$  can be used to construct a good solution to LCP(7).  $\square$

We now give the augmented LCP for LCP(7) so we can apply Lemke's scheme as discussed in Section 6.1. As is standard practice, we add the variable  $z$  in the constraints whose right hand side is negative. We thus obtain LCP(8).

$$\forall i \in N : \quad e_i \cdot \frac{\sum_j (-p_j - \beta_j)}{\sum_h e_h} - \sum_j q_{ij} - z \leq -e_i \cdot \frac{mP}{\sum_h e_h} \quad \perp \quad r_i \quad (8a)$$

$$\forall j \in M : \quad p_j + \beta_j + \sum_i q_{ij} \leq P \quad \perp \quad p_j \quad (8b)$$

$$\forall i \in N, j \in M : \quad d_{ij}r_i - p_j \leq d_{ij}R - P \quad \perp \quad q_{ij} \quad (8c)$$

$$\forall j \in M : \quad -p_j - \beta_j + c_j \cdot \frac{\sum_k (p_k + \beta_k)}{\sum_i e_i} - z \leq -P + c_j \cdot \frac{mP}{\sum_i e_i} \quad \perp \quad \beta_j \quad (8d)$$

$$z \geq 0 \quad (8e)$$

Let  $\mathcal{P}$  be the polyhedron defined by the constraints of LCP(8). The primary ray in Lemke's algorithm will set  $\mathbf{p}, \mathbf{q}, \mathbf{r}, \boldsymbol{\beta}$  to zero and  $z = \max\{\max_i \frac{e_i \cdot mP}{\sum_h e_h}, \max_j (c_j \cdot \frac{mP}{\sum_i e_i} - P)\}$  as the initial vertex solution. Lemke's scheme involves pivoting from *non-degenerate* vertices of  $\mathcal{P}$ , i.e., those with a unique double label (see Section 6.1).

**Definition 10** (Non-degenerate instance). *An instance  $(N, M, D, e, c)$  is non-degenerate if there is no polynomial relation between the input parameters, i.e., there is no polynomial  $\phi$  s.t.  $\phi(D, e, c) = 0$ .*

We can assume our instance is non-degenerate without loss of generality, as there are standard ways of handling degeneracy in the input parameters like the lexico-minimum test [19].

**Lemma 30.** *For a non-degenerate instance  $(N, M, D, e, c)$ , every vertex  $\mathbf{v} = (\mathbf{p}, \mathbf{q}, \mathbf{r}, \boldsymbol{\beta}, z)$  encountered in Lemke's scheme which is good and satisfies  $z > 0$  is non-degenerate.*

*Proof.* For the sake of contradiction, suppose  $\mathbf{v} = (\mathbf{p}, \mathbf{q}, \mathbf{r}, \boldsymbol{\beta}, z)$  is a vertex encountered in Lemke's scheme which is good, where  $z > 0$ , and which is degenerate. We show that the parameters of the instance have a polynomial relation, contradicting the instance being non-degenerate.

Let  $N$  be the number of variables in LCP(7). Hence the augmented LCP has  $N + 1$  variables, with  $z$  being the additional variable. Let the augmented LCP polyhedron be given by  $\mathcal{P} = \{\mathbf{y} \in \mathbb{R}^N, z \in \mathbb{R} : A\mathbf{y} \leq \mathbf{b}, \mathbf{y} \geq 0, z \geq 0\}$ , where  $A \in \mathbb{R}^{N \times (N+1)}$ , and  $\mathbf{b} \in \mathbb{R}^N$ . Thus, there are  $N$  linear constraints given by the rows of  $A$ , and each such constraint may involve  $(N + 1)$  variables, including  $z$ .

Since  $\mathbf{v}$  is a vertex of  $\mathcal{P}$ , exactly  $(N + 1)$  inequalities (out of the  $2N + 1$  inequalities, including the non-negativity constraints) must be tight at  $\mathbf{v}$ . Let  $I$  be the set of non-zero variables of  $\mathbf{v}$  excluding  $z$ , and let  $|I| = N'$ . By complementarity, the constraint  $(A\mathbf{v} - \mathbf{b})_i = 0$  for each such variable  $i \in I$ . Consider the subsystem of  $A\mathbf{y} \leq \mathbf{b}$  corresponding to the variables in  $I$  and  $z$ . This can be represented as a collection of equalities given by  $A' \cdot \mathbf{v}' = \mathbf{b}'$ , where  $A' \in \mathbb{R}^{N' \times (N'+1)}$ ,  $\mathbf{b}' \in \mathbb{R}^{N'}$  and  $\mathbf{v}' \in \mathbb{R}^{N'+1}$ . Note that  $\mathbf{v}' = (\mathbf{v}'', z)$  is simply the subvector of  $\mathbf{v}$  with non-zero entries. By separating out the terms involving  $z$ , we can transform the above system into an equation of the form  $z \cdot \boldsymbol{\gamma} + A'' \cdot \mathbf{v}'' = \mathbf{b}'$ , where  $A'' \in \mathbb{R}^{N' \times N'}$  and  $\boldsymbol{\gamma} \in \mathbb{R}^{N'}$ . This gives  $\mathbf{v}'' = (A'')^{-1} \mathbf{b}' - z \cdot (A'')^{-1} \boldsymbol{\gamma}$ , which expresses each non-zero variable in the set  $I$  as a linear term in  $z$  with coefficients that are polynomials in the input parameters.

Now observe that the degeneracy of  $\mathbf{v}$  implies that  $\mathbf{v}$  has at least two double labels (see Section 6.1). That is, there are two variables  $i, j \notin I$  s.t.  $y_i = y_j = (A\mathbf{v} - \mathbf{b})_i = (A\mathbf{v} - \mathbf{b})_j = 0$ . We use one of these equalities to solve for  $z$  by replacing each non-zero variable with its linear expression in  $z$  obtained earlier. We then substitute this value of  $z$  into the second equality to obtain a polynomial relation in the input parameters. This contradicts the fact that the instance is non-degenerate.  $\square$

## 6.4 Convergence of Lemke's scheme

In this section, we show that Lemke's scheme converges to a good solution  $(\mathbf{p}, \mathbf{q}, \mathbf{r}, \boldsymbol{\beta}, z)$  with  $z = 0$  for LCP(8) for non-degenerate instances. A solution to LCP(8) with  $z = 0$  is a solution to LCP(7). With Lemma 29, this implies the existence of ER equilibria and proves Theorem 2. Further, it provides an algorithm for computing an ER equilibrium.

To show convergence, we need to show that starting from the primary ray, Lemke's scheme only reaches good vertex solutions and does not reach a secondary ray. We prove the former using Lemmas 31, 32, and 33, and the latter using Lemma 34. Recall that a solution  $(\mathbf{p}, \mathbf{q}, \mathbf{r}, \boldsymbol{\beta}, z)$  is good if  $\forall j : p_j + \beta_j < P$  and  $\forall i : r_i < R$ .

**Lemma 31.** *Let  $\mathbf{v} = (\mathbf{p}, \mathbf{q}, \mathbf{r}, \boldsymbol{\beta}, z)$  be a vertex encountered by Lemke's scheme starting from the primary ray of LCP(8). If  $\forall j : p_j + \beta_j < P$  holds, then  $\forall i : r_i < R$  also holds.*

*Proof.* Suppose at a vertex  $p_j + \beta_j < P$  for all  $j$ . Then for every  $i$ , (8c) implies  $d_{ij}(r_i - R) \leq p_j - P$ . Since  $p_j < P$  and  $d_{ij} > 0$ , we have  $r_i < R$  for all  $i$ .  $\square$

**Lemma 32.** *Let  $\mathbf{v} = (\mathbf{p}, \mathbf{q}, \mathbf{r}, \boldsymbol{\beta}, z)$  be a vertex encountered by Lemke's scheme starting from the primary ray of LCP(8), and let  $\mathbf{v}' = (\mathbf{p}', \mathbf{q}', \mathbf{r}', \boldsymbol{\beta}', z')$  be the next vertex after a pivoting step. If  $\forall j : p_j + \beta_j < P$  holds, then it cannot happen that  $p'_j + \beta'_j = P$  holds for a strict non-empty subset of  $M$ .*

*Proof.* Let  $M_0 \subseteq M$  be the set of chores in  $\mathbf{v}'$  for which  $p'_j + \beta'_j = P$ . Assume  $\emptyset \neq M_0 \subsetneq M$ . Hence there are chores  $g$  and  $k$  s.t.  $g \in M_0$  and  $k \in M \setminus M_0$ . Thus,  $p'_g + \beta'_g = P$  and  $p'_k + \beta'_k < P$ . Let  $E$  be the edge from  $\mathbf{v}$  to  $\mathbf{v}'$ . Along  $E$ , at least one of (8b) and (8b)' and at least one of (8d) and (8d)' have to remain tight for chore  $g$ , due to complementarity. We now consider three cases:

- The constraint (8d) remains tight along  $E$  for  $g$ . Thus,  $P - p_g - \beta_g = c_g \frac{\sum_j (P - p_j - \beta_j)}{\sum_i e_i} + z$  holds along  $E$ . Since  $P - p_g - \beta_g$  goes to 0 along  $E$ , we must have that  $c_g \frac{\sum_j (P - p_j - \beta_j)}{\sum_i e_i} + z$  goes to 0 along  $E$ . Since  $p_j + \beta_j \leq P$  for all  $j \in M$  due to constraint (8b) and  $z \geq 0$ , it must be that  $P - p'_j - \beta'_j = 0$  for all  $j \in M$  at  $\mathbf{v}'$ , contradicting that  $p'_k + \beta'_k < P$ .
- The constraints (8b) and (8d)' remain tight along  $E$  for  $g$ . Thus  $\sum_i q_{ig} = P - p_g - \beta_g$  and  $\beta_g = 0$  along  $E$ . Thus  $p_g < P$  pivots to  $p'_g = P$  along  $E$  from  $\mathbf{v}$  to  $\mathbf{v}'$ . Since  $\sum_i q_{ij} = P - p_g > 0$  at  $\mathbf{v}$  and along  $E$ , there is some agent  $i$  s.t.  $q_{ij} > 0$ . By complementarity for (8c), we have  $d_{ig}r_i - p_g = d_{ig}R - P$ . Moreover  $d_{ik}r_i - p_k \leq d_{ik}R - P$ . This implies:

$$R - r_i = \frac{P - p_g}{d_{ig}} \geq \frac{P - p_k}{d_{ik}}.$$

Since  $p_g$  pivots to  $P$  along  $E$ , the above inequality implies  $R - r_i$  also pivots to 0. Hence  $p_k$  must pivot to  $P$  along  $E$ , i.e.,  $p'_k = P$ . This contradicts the fact that at  $\mathbf{v}'$ ,  $p'_k + \beta'_k < P$ .

- The constraints (8b)' and (8d)' remain tight along  $E$  for  $g$ . Thus,  $p_g = \beta_g = 0$  along  $E$ . In particular at  $\mathbf{v}'$ ,  $p'_g + \beta'_g = 0$ . This contradicts  $p'_g + \beta'_g = P$ . Therefore it cannot happen that both (8b)' and (8d)' remain tight along  $E$ .

Thus, it cannot happen that at  $\mathbf{v}'$ ,  $p'_j + \beta'_j = P$  holds for a strict non-empty subset of chores.  $\square$

**Lemma 33.** *Let  $\mathbf{v} = (\mathbf{p}, \mathbf{q}, \mathbf{r}, \boldsymbol{\beta}, z)$  be a vertex encountered by Lemke's scheme starting from the primary ray of LCP(8), and let  $\mathbf{v}' = (\mathbf{p}', \mathbf{q}', \mathbf{r}', \boldsymbol{\beta}', z')$  be the next vertex after a pivoting step. If  $\forall j : p_j + \beta_j < P$  holds, then  $\forall j : p'_j + \beta'_j = P$  cannot hold.*

*Proof.* For sake of contradiction, suppose  $\forall j \in M, p'_j + \beta'_j = P$  at  $\mathbf{v}'$ . Let  $E$  be the edge from  $\mathbf{v}$  to  $\mathbf{v}'$ . Then for each chore  $j$ , either  $p_j$  or  $\beta_j$  increases along  $E$ . We consider three cases.

- Suppose  $p_j$  increases along  $E$  for each  $j \in M$ . By complementarity, constraint (8b) must be tight at  $\mathbf{v}$ . Thus  $\forall j : \sum_i q_{ij} = P - p_j - \beta_j$ , which implies:

$$\sum_j \sum_i q_{ij} = \sum_j (P - p_j - \beta_j). \quad (9)$$

Let  $N_1 = \{i \in N : r_i = 0\}$  and  $N_2 = N \setminus N_1$ . For  $i \in N_1$ , since  $r_i = 0$ ,  $p_j \geq 0$  and  $d_{ij}R - P > 0$ , (8c) is not tight. By complementarity,  $q_{ij} = 0$  for all  $j \in M$ . Thus (8a) becomes



$e_i \cdot \frac{\sum_j (P - p_j - \beta_j)}{\sum_h e_h} \leq z$  for  $i \in N_1$ . For  $i \in N_2$ , since  $r_i > 0$ , complementarity implies (8a) must be tight. Thus  $e_i \cdot \frac{\sum_j (P - p_j - \beta_j)}{\sum_h e_h} - z = \sum_j q_{ij}$  for  $i \in N_2$ . Using these observations, we have:

$$\begin{aligned} \sum_i \sum_j q_{ij} &= \sum_{i \in N_2} \sum_j q_{ij} = \sum_{i \in N_2} \left( e_i \cdot \frac{\sum_j (P - p_j - \beta_j)}{\sum_h e_h} - z \right) \\ &= \frac{\sum_{i \in N_2} e_i}{\sum_h e_h} \cdot \sum_j (P - p_j - \beta_j) - |N_2|z. \end{aligned} \quad (10)$$

Putting (9) and (10) together and rearranging, we obtain:

$$\frac{\sum_{i \in N_1} e_i}{\sum_h e_h} \cdot \sum_j (P - p_j - \beta_j) = -|N_2|z.$$

Since  $p_j + \beta_j < P$  for all  $j \in M$ , we have  $\sum_j (P - p_j - \beta_j) > 0$ . Moreover  $z > 0$  at  $\mathbf{v}$ . Since either  $N_1 \neq \emptyset$  or  $N_2 \neq \emptyset$ , the above equality cannot hold.

- Suppose  $\beta_j$  increases along  $E$  for each  $j \in M$ . By complementarity, constraint (8d) must be tight. Hence  $P - p_j - \beta_j - z = c_j \cdot \frac{\sum_k (P - p_k - \beta_k)}{\sum_i e_i}$  for all  $j \in M$ . Summing over all  $j$  and rearranging gives:

$$\sum_j (P - p_j - \beta_j) \cdot \left( \frac{\sum_j c_j}{\sum_i e_i} - 1 \right) = -mz.$$

Since  $p_j + \beta_j < P$  for all  $j \in M$ , we have  $\sum_j (P - p_j - \beta_j) > 0$ . Since  $\sum_i e_i \leq \sum_j c_j$ , hence the left side of the above equation is non-negative. Since  $z > 0$  at  $\mathbf{v}$ , the right side is negative, and hence the above equality cannot hold.

- There are two chores  $j$  and  $k$  such that  $p_j = 0$  and  $\beta_k = 0$  hold all along the edge  $E$ . It cannot be that  $j = k$ , since  $p'_j + \beta'_j = P$ . Since  $p_k + \beta_k$  increases to  $P$  while  $\beta_k = 0$  along  $E$ , it must mean that  $p_k$  increases to  $P$  along  $E$ . By complementarity, (8b) is tight along  $E$ . Thus,  $\sum_i q_{ik} = P - p_k > 0$  along  $E$ , since  $p_k < P$  along  $E$ . Thus, there is some agent  $i \in N$  for which  $q_{ik} > 0$  along  $E$ . By complementarity, (8c) implies that  $d_{ik}r_i - p_k = d_{ik}R - P$  holds along  $E$ . Moreover the constraint (8c) also implies  $d_{ij}r_i - p_j \leq d_{ij}R - P$  holds along  $E$ . This implies that:

$$R - r_i = \frac{P - p_k}{d_{ik}} \geq \frac{P - p_j}{d_{ij}}$$

holds along the edge  $E$ . However, since  $p_j = 0$  along  $E$ , we have  $p_k \leq P \cdot (1 - d_{ik}/d_{ij}) < P$ , where  $d_{ik} \neq d_{ij}$  follows from the non-degeneracy of the instance. Thus,  $p_k$  always remains strictly below  $P$  along the edge  $E$ . Therefore, it cannot happen that  $p'_k = P$  at  $\mathbf{v}'$ , which is a contradiction.

Since these cases are exhaustive, the lemma holds.  $\square$

**Lemma 34.** *Lemke's scheme starting from the primary ray of LCP(8) does not reach a secondary ray.*

*Proof.* Suppose Lemke's scheme starting from the primary ray of LCP(8) reaches a vertex  $\mathbf{v}^0 = (\mathbf{p}^0, \mathbf{q}^0, \mathbf{r}^0, \beta^0, z^0)$  and then pivots to a secondary ray given by  $\mathcal{R} = \{\mathbf{v}^0 + \alpha \cdot \mathbf{v}' : \alpha \geq 0\}$ , where

$\mathbf{v}' = (\mathbf{p}', \mathbf{q}', \mathbf{r}', \boldsymbol{\beta}', z')$  with  $z' > 0$ . We first show that  $\mathbf{v}' = \mathbf{0}$  by arguing that if this is not the case then some constraint of LCP(8) will be violated at some point on the secondary ray.

If  $p'_j < 0$  for some  $j \in M$  then eventually the non-negativity constraint (8b)' will be violated. On the other hand if  $p'_j > 0$  then eventually the constraint (8b) will be violated. Thus  $\mathbf{p}' = \mathbf{0}$ . By similar arguments considering constraints (8c)' and (8b) for  $\mathbf{q}'$ , (8a)' and (8c) for  $\mathbf{r}'$ , and (8d)' and (8b) for  $\boldsymbol{\beta}'$ , we can conclude that  $\mathbf{q}' = \mathbf{0}$ ,  $\mathbf{r}' = \mathbf{0}$  and  $\boldsymbol{\beta}' = \mathbf{0}$ . If  $z' < 0$  then the  $z \geq 0$  constraint will be violated eventually. Suppose  $z' > 0$ . Then (8a) becomes strict, implying that  $r_i = r_i^0 = 0$  for each  $i \in N$ . Since  $d_{ij}R - P > 0$  for all  $i, j$ , (8c) is strict. By complementarity,  $q_{ij} = q_{ij}^0 = 0$  for all  $i \in N, j \in M$ . Similarly, since  $z' > 0$  (8d) eventually becomes strict and by complementarity  $\beta_j = \beta_j^0 = 0$  for all  $j \in M$ . Since  $\mathbf{v}^0$  is a vertex encountered in Lemke's scheme starting from the primary ray, Lemma 32 and Lemma 33 imply that  $\forall j : p_j^0 + \beta_j^0 < P$ . This means that (8b) is strict, and hence by complementarity  $p_j = p_j^0 = 0$  for all  $j \in M$ . Thus,  $\mathbf{p}^0 = \mathbf{0}$ ,  $\mathbf{q}^0 = \mathbf{0}$ ,  $\mathbf{r}^0 = \mathbf{0}$  and  $\boldsymbol{\beta}^0 = \mathbf{0}$ . We therefore have  $\mathcal{R} = \{(0, 0, 0, 0, z^0) + \alpha \cdot (0, 0, 0, 0, z') : \alpha \geq 0\}$  where  $z^0 > 0$  and  $z' > 0$ . However this is the same as the primary ray, thus showing  $\mathcal{R}$  cannot be a secondary ray.  $\square$

We conclude the above discussion in the following lemma.

**Lemma 35.** *Lemke's scheme starting from the primary ray of LCP(8) converges to a good solution  $(\mathbf{p}, \mathbf{q}, \mathbf{r}, \boldsymbol{\beta}, z)$  where  $z = 0$ .*

*Proof.* Lemma 34 shows that no secondary rays are encountered. Lemma 32 and Lemma 33 together show that if at a vertex  $\mathbf{v}$  it holds that  $\forall j : p_j + \beta_j < P$ , then the same holds at the next vertex  $\mathbf{v}'$  after pivoting. Since the primary ray sets all  $p_j = \beta_j = 0$ , this is true initially. Hence  $\forall j : p_j + \beta_j < P$  at every vertex encountered by Lemke's scheme. Finally Lemma 31 shows that at every such vertex  $\forall i : r_i < R$  also holds, hence such a vertex is good. Lemma 30 shows that every good vertex with  $z > 0$  is non-degenerate. Hence pivoting to the next step is always possible and Lemke's scheme eventually reaches a good solution with  $z = 0$ .  $\square$

Lemma 29 and Lemma 35 thus prove Theorem 2: the existence of an ER equilibrium under the feasible earning condition, and also show that Lemke's scheme can be used to compute it.

## A Examples

**Example 1.** A Prop and PO allocation need not be  $\alpha$ -EFk for any  $\alpha, k \geq 1$ .

Consider an instance with three agents  $a, b$ , and  $c$  and three types of chores, each with  $s > k$  many copies. The disutility of each agent for each chore type is given below.

	Type 1	Type 2	Type 3
$a$	1	$t$	$3t$
$b$	1	$t$	$3t$
$c$	$t$	$t$	1

We claim that for  $t > \frac{\alpha \cdot s}{s-k}$ , the allocation  $\mathbf{x}$  in which agent  $a$  receives all type 1 chores,  $b$  receives all type 2 chores, and  $c$  receives all type 3 chores is Prop + PO but fails to be  $\alpha$ -EFk. We first note that as  $\alpha \geq 1$  and  $\frac{s}{s-k} > 1$ , we have  $t > 1$ . It is then easily verified that  $\mathbf{x}$  is Prop. Additionally, since  $\mathbf{x}$  is social welfare maximizing, it is necessarily fPO. We now show that agent  $b$   $\alpha$ -EFk-envies agent  $a$ . We have that

$$\min_{S \subseteq \mathbf{x}_b, |S| \leq k} d_b(\mathbf{x}_b \setminus S) = t(s-k) > \frac{\alpha \cdot s}{s-k} \cdot (s-k) = \alpha \cdot s = \alpha \cdot d_b(\mathbf{x}_a),$$

showing the result.

**Example 2** (ER equilibrium). Consider an instance with two agents  $\{a_1, a_2\}$  and three chores  $\{j_1, j_2, j_3\}$ , with disutilities given in the following table.

	$j_1$	$j_2$	$j_3$
$a_1$	1	3	6
$a_2$	6	3	1

Suppose each agent has an earning requirement of 1. Let  $\mathbf{x} = (\mathbf{x}_1, \mathbf{x}_2)$  be the allocation given by  $\mathbf{x}_1 = \{j_1\}$  and  $\mathbf{x}_2 = \{j_2, j_3\}$ . Let  $\mathbf{p} = (p_{j_1}, p_{j_2}, p_{j_3}) = (1, \frac{3}{4}, \frac{1}{4})$  be a payment vector. It is easily verified that  $(\mathbf{x}, \mathbf{p})$  satisfies Definition 2 and is a (unrestricted) CE.

We now impose a uniform earning limit  $\beta = \frac{2}{3}$  on each chore, giving us an earning-restricted instance. As  $q_{j_1} = 1 > \frac{2}{3}$  and  $q_{j_2} = \frac{3}{4} > \frac{2}{3}$  in  $(\mathbf{x}, \mathbf{p})$ , it is not an ER equilibrium. Consider the allocation  $(\mathbf{x}', \mathbf{p}')$  given by  $\mathbf{x}'_1 = \{j_1, \frac{1}{6}j_2\}$ ,  $\mathbf{x}'_2 = \{\frac{1}{6}j_2, j_3\}$ , and  $\mathbf{p}' = (p_{j_1}, p_{j_2}, p_{j_3}) = (\frac{2}{3}, 2, \frac{2}{3})$ . Since it satisfies Definition 4,  $(\mathbf{x}', \mathbf{p}')$  is an ER equilibrium. We highlight that here  $j_2$  is not completely allocated but agents collectively earn the earning limit  $\beta = \frac{2}{3}$  from  $j_2$ . Moreover, we learn more about agent preferences from  $(\mathbf{x}', \mathbf{p}')$  than  $(\mathbf{x}, \mathbf{p})$ : in addition to their favorite chore, each agent is partially allocated their second favorite chore in  $\mathbf{x}'$ , but not in  $\mathbf{x}$ .

**Example 3.** A competitive equilibrium from equal earning (CEEE) need not admit a rounding which is  $\alpha$ -EFk.

We construct a CEEE and show that it admits no rounded  $\alpha$ -EFk allocation. We consider three agents  $a, b$ , and  $c$  with identical disutility functions. There exists one shared chore  $j$  among the agents such that  $a, b$ , and  $c$  earn  $\frac{1}{2}$ ,  $1 - \frac{1}{5\alpha}$ , and  $1 - \frac{1}{5\alpha}$  from  $j$ , respectively. Each agent  $i$  is integrally allocated a set of chores  $S_i$  such that:

- agent  $a$  earns  $\frac{1}{2}$  from  $S_a$ .
- agent  $b$  earns  $\frac{1}{5\alpha}$  from  $S_b$ .

- agent  $c$  earns  $\frac{1}{5\alpha}$  from  $S_c$ .

Specifically, we note that  $S_a$  consists of  $2k$  identical chores which each pay  $\frac{1}{4k}$ . In conjunction with their earning from  $j$ , we see that each agent earns 1, showing equal earnings. We now show that  $\alpha$ -EF $k$ -envy between agents persists regardless of whom the single shared chore  $j$  is rounded to in the integral allocation  $\mathbf{x}$ . We have that:

$$\min_{S \subseteq S_a, |S| \leq k} d_a(S_a \setminus S) = \alpha_a \cdot \mathbf{p}_{-k}(S_a) = \alpha_a \cdot \frac{1}{4} > \alpha_a \cdot \frac{1}{5} = \alpha_a \cdot \alpha \cdot \mathbf{p}(S_b) = \alpha \cdot d_a(S_b).$$

Note that for  $i \in \{a, b, c\}$ ,  $S_i \subseteq \mathbf{x}_i$ . Thus, we have that:

$$\min_{S \subseteq \mathbf{x}_a, |S| \leq k} d_a(\mathbf{x}_a \setminus S) \geq \min_{S \subseteq S_a, |S| \leq k} d_a(S_a \setminus S) > \alpha \cdot d_a(S_b).$$

It follows then that if  $\mathbf{x}_b = S_b$ , agent  $a$  will  $\alpha$ -EF $k$ -envy agent  $b$ . An analogous argument shows that if  $\mathbf{x}_c = S_c$ , agent  $a$  will  $\alpha$ -EF $k$ -envy agent  $c$ . Since  $j$  can only be rounded to one agent, it must be that either  $\mathbf{x}_b = S_b$  or  $\mathbf{x}_c = S_c$ , so agent  $a$  must  $\alpha$ -EF $k$ -envy some agent.

**Example 4.** *There exists an ER equilibrium for which no rounding is  $(n - 1 - \delta)$ -EF1.*

We construct an ER equilibrium with  $n = 2k + 1$  agents  $i_1, \dots, i_{2k+1}$ ,  $2k - 1$  shared chores  $j_1, \dots, j_{2k-1}$ , and uniform chore earning limit  $\beta = 1$ . Note that agents may have other chores which are not shared with other agents.

We describe the structure of the payment graph  $G$ .  $G$  is a forest with two trees. The first tree  $T_1$  consists of the lone agent  $i_{2k+1}$  and  $\frac{1}{\varepsilon}$  many small,  $\varepsilon$ -paying chores which are integrally allocated to  $i_{2k+1}$ . The second tree  $T_2$  contains agents  $i_1$  to  $i_{2k}$  and all of the  $2k - 1$  shared chores. We note that each agent  $i \in T_2$  earns  $\frac{1}{2k}$  from a set of chores  $S_i$  which is integrally allocated to  $i$ , so we focus our attention on the edges that are incident to the  $2k - 1$  shared chores.

Let  $T_2$  be rooted at chore  $j_{2k-1}$  so that  $j_{2k-1}$  has two agents  $i_{2k-1}$  and  $i_{2k}$  as children, each of whom earn  $\frac{1}{2}$  from  $j_{2k-1}$ . Then,  $i_{2k-1}$  and  $i_{2k}$  each have  $k - 1$  children chores, with  $i_{2k-1}$  having children  $j_1, \dots, j_{k-1}$  and  $i_{2k}$  having children  $j_k, \dots, j_{2k-2}$ . Both  $i_{2k-1}$  and  $i_{2k}$  earn  $\frac{1}{2k}$  from each of their children. Finally, for  $r \in \{1, \dots, 2k - 2\}$ , chore  $j_r$  has one child agent  $i_r$  who earns  $1 - \frac{1}{2k}$  from  $j_r$ . We verify that each agent meets their earning requirement in the following:

- For  $1 \leq r \leq 2k - 2$ ,  $e_{i_r} = \frac{1}{2k} + \left(1 - \frac{1}{2k}\right) = 1$ .
- For  $r \in \{2k - 1, 2k\}$ ,  $e_{i_r} = \frac{1}{2k} + \frac{1}{2} + (k - 1) \cdot \frac{1}{2k} = 1$ .
- For  $r = 2k + 1$ ,  $e_{i_r} = \frac{1}{\varepsilon} \cdot \varepsilon = 1$ .

Additionally, each shared chore meets the earning limit:

- For  $1 \leq r \leq 2k - 2$ ,  $q_r = \frac{1}{2k} + \left(1 - \frac{1}{2k}\right) = 1$ .
- For  $r = 2k - 1$ ,  $q_r = \frac{1}{2} + \frac{1}{2} = 1$ .

It is trivial to have the chores in a set  $S_i$  satisfy the earning limit by increasing the number of chores in  $S_i$  and thus decreasing the individual earning from each chore. We now show that there is no rounding of the ER equilibrium that is better than  $(n - 1)$ -EF1. We may assume that the payment of any chore  $j$  is equal to its payout to the agents, i.e., for all  $j \in M$ ,  $p_j = q_j$ . In any rounding there must be some agent  $h$  in  $T_2$  who receives no shared chore and thus earns only  $\frac{1}{2k}$ : this is because there are  $2k$  agents in  $T_2$  but only  $2k - 1$  shared chores. Suppose for the sake of contradiction that

a rounded allocation  $\mathbf{x}$  is  $(n - 1 - \delta)$ -EF1 for some  $\delta > 0$ . Note that the bundle of agent  $i_{2k+1}$  is the same in any rounding, as  $i_{2k+1}$  does not share chores with any agent. Then, letting  $\varepsilon < \frac{\delta}{n-1}$  and letting the disutility function of  $i_{2k+1}$  be such that for all  $S \subseteq M$ ,  $d_{i_{2k+1}}(S) = \mathbf{p}(S)$  (so  $\alpha_{i_{2k+1}} = 1$  and all chores are MPB for  $i_{2k+1}$ ), we have that

$$\mathbf{p}_{-1}(\mathbf{x}_{i_{2k+1}}) = \min_{j \in \mathbf{x}_{i_{2k+1}}} d_{i_{2k+1}}(\mathbf{x}_{i_{2k+1}} \setminus \{j\}) \leq (n - 1 - \delta) \cdot d_{i_{2k+1}}(\mathbf{x}_h) = (n - 1 - \delta) \cdot \mathbf{p}(\mathbf{x}_h),$$

where the first and last equalities stem from our definition of  $d_{i_{2k+1}}(\cdot)$  and the inequality stems from the fact that  $\mathbf{x}$  is  $(n - 1 - \delta)$ -EF1. Using  $\mathbf{p}_{-1}(\mathbf{x}_{i_{2k+1}}) \leq (n - 1 - \delta) \cdot \mathbf{p}(\mathbf{x}_h)$ , we have that  $1 - \varepsilon \leq (n - 1 - \delta) \cdot \frac{1}{2k} = (n - 1 - \delta) \cdot \frac{1}{n-1}$  and equivalently that  $\varepsilon \geq \frac{\delta}{n-1}$ , a contradiction. Thus, such an ER equilibrium has no rounding which is  $(n - 1 - \delta)$ -EF1 for any  $\delta > 0$ .

**Example 5.** *There exists an ER equilibrium for which no rounding is  $(2 - \delta)$ -EF2.*

We modify the ER equilibrium from Example 4 with  $n = 2k + 1$  agents. Setting  $\beta = \frac{1}{2}$ , we aim to change the agent earnings for each chore so that each agent still receives their earning requirement 1 but the total earning from any chore is at most  $\frac{1}{2}$ . The changes are as follows:

- each agent  $i \in T_2$  earns  $\frac{1}{2} + \frac{1}{4k}$  from their set of integrally allocated chores  $S_i$ ,
- agents  $i_{2k-1}$  and  $i_{2k}$  each earn  $\frac{1}{4}$  from their parent chore  $j_{2k-1}$ ,
- agents  $i_{2k-1}$  and  $i_{2k}$  earn  $\frac{1}{4k}$  from each of their  $k - 1$  children,
- for  $r \in \{1, \dots, 2k - 2\}$ , agent  $i_r$  earns  $\frac{1}{2} - \frac{1}{4k}$  from their parent chore.

As in Example 4, it can be verified that each agent meets her earning requirement and each chore satisfies the earning limit. Then, also as in Example 4, it must be that some agent in  $T_2$  receives only their integrally allocated chores. That is, there exists some  $i \in T_2$  such that  $\mathbf{x}_i = S_i$  and  $d_{i_{2k+1}}(\mathbf{x}_i) = \mathbf{p}(\mathbf{x}_i) = \frac{1}{2} + \frac{1}{4k}$ . Recall that agent  $i_{2k+1}$  is only allocated  $\frac{1}{\varepsilon}$  many small,  $\varepsilon$ -paying chores and  $\min_{S \subseteq \mathbf{x}_{i_{2k+1}}, |S| \leq 2} d_{i_{2k+1}}(\mathbf{x}_{i_{2k+1}} \setminus S) = p_{-2}(\mathbf{x}_{i_{2k+1}})$ . Then, for any  $\delta > 0$ , we may choose sufficiently large  $k$  and sufficiently small  $\varepsilon$  so that

$$p_{-2}(\mathbf{x}_{i_{2k+1}}) \geq \varepsilon \cdot \left(\frac{1}{\varepsilon} - 2\right) = 1 - 2\varepsilon > (2 - \delta) \cdot \left(\frac{1}{2} + \frac{1}{4k}\right) = (2 - \delta) \cdot \mathbf{p}(\mathbf{x}_i) = (2 - \delta) \cdot d_{i_{2k+1}}(\mathbf{x}_i)$$

and thus agent  $i_{2k+1}$   $(2 - \delta)$ -EF2-envies agent  $i$ .

**Example 6.** *An  $\alpha$ -EFX + fPO allocation need not exist for 2-ary instances.*

Consider the following 2-ary instance with two agents  $\{a, b\}$ , four chores  $\{j_1, j_2, j_3, j_4\}$  with disutilities given by:

	$j_1$	$j_2$	$j_3$	$j_4$
$a$	1	1	$3\alpha$	$3\alpha$
$b$	1	1	$3\alpha^2 + 3\alpha$	$3\alpha^2 + 3\alpha$

We show that this instance does not admit an allocation which is both  $\alpha$ -EFX and fPO. Suppose that  $\mathbf{x}$  is an  $\alpha$ -EFX allocation. Given that  $\alpha \geq 1$ , we see that neither agent may receive both  $j_3$  and  $j_4$  under  $\mathbf{x}$ , as for  $i \in \{a, b\}$  we have

$$\max_{j \in \{j_3, j_4\}} d_i(\{j_3, j_4\} \setminus \{j\}) \geq 3\alpha > 2\alpha = \alpha \cdot d_i(\{j_1, j_2\}).$$

Thus, we assume without loss of generality that  $j_3 \in \mathbf{x}_a$  and  $j_4 \in \mathbf{x}_b$ . We next argue that  $j_1, j_2 \in \mathbf{x}_a$ . Suppose for sake of contradiction (again w.l.o.g.) that  $j_2 \in \mathbf{x}_b$ . We have

$$\max_{j \in \mathbf{x}_b} d_b(\mathbf{x}_b \setminus \{j\}) \geq d_b(\{j_2, j_4\} \setminus \{j_2\}) = 3\alpha^2 + 3\alpha > 3\alpha^2 + \alpha = \alpha \cdot (3\alpha + 1) = \alpha \cdot d_b(\{j_1, j_3\}) \geq \alpha \cdot d_b(\mathbf{x}_a),$$

so agent  $b$  would  $\alpha$ -EFX-envy agent  $a$ . Thus, it must be that  $\mathbf{x}_a = \{j_1, j_2, j_3\}$  and  $\mathbf{x}_b = \{j_4\}$ . We now show however that  $\mathbf{x}$  is not fPO as it is dominated by the fractional allocation  $\mathbf{y}$  where  $\mathbf{y}_a = \{j_1, j_3, \frac{1}{3\alpha}j_4\}$  and  $\mathbf{y}_b = \{j_2, \frac{3\alpha-1}{3\alpha}j_4\}$ . Indeed, we have that  $d_a(\mathbf{y}_a) = 3\alpha + 2 = d_a(\mathbf{x}_a)$  and  $d_b(\mathbf{y}_b) = 3\alpha^2 + 2\alpha < 3\alpha^2 + 3\alpha = d_b(\mathbf{x}_b)$ . Agent  $a$ 's disutility remains the same under  $\mathbf{y}$  while agent  $b$ 's disutility strictly decreases under  $\mathbf{y}$ , so  $\mathbf{y}$  dominates  $\mathbf{x}$  and  $\mathbf{x}$  not fPO. Thus, the given instance admits no  $\alpha$ -EFX and fPO allocation.

## B Appendix to Section 3

**Lemma 2.** *There is a polynomial time algorithm `MakeAcyclic` which takes as input an ER equilibrium of instance  $I$  and returns another ER equilibrium of  $I$  whose payment graph is acyclic.*

*Proof.* `MakeAcyclic` begins with the payment graph  $G = (N \sqcup M, E)$  of the ER equilibrium  $(\mathbf{y}, \mathbf{p})$ . If  $G$  is acyclic, it returns  $(\mathbf{y}, \mathbf{p})$ . Otherwise, suppose an intermediate allocation  $(\mathbf{z}, \mathbf{p})$  has a cycle  $C = (i_1, j_1, i_2, j_2, \dots, i_k, j_k, i_1)$ , where  $i_\ell \in N$  are agents and  $j_\ell \in M$  are chores, and  $C$  contains the edges  $(i_\ell, j_\ell) \in E$  and  $(j_\ell, i_{\ell+1}) \in E$  for  $1 \leq \ell \leq k$ , with the notation that  $i_{k+1} = i_1$ . The earning of an agent  $i$  from chore  $j$  is  $q_{ij} = p_j \cdot y_{ij}$ . Without loss of generality, assume  $(i_1, j_1)$  is the edge with minimum  $q_{ij}$  among the edges  $(i, j)$  in  $C$ . Let  $s = q_{i_1 j_1}$ .

Now consider the allocation  $(\mathbf{z}, \mathbf{p}, \mathbf{q}')$ , where for all  $\ell \in [k]$ ,  $q'_{i_\ell j_\ell} = q_{i_\ell j_\ell} - s$ , and  $q'_{i_\ell j_{\ell-1}} = q_{i_\ell j_{\ell-1}} + s$ , and  $q'_{ij} = q_{ij}$  for all  $(i, j) \notin C$ . This has the effect of circulating agent earnings around the cycle  $C$  and the edge  $(i_1, j_1)$  is no longer present in the payment graph of  $(\mathbf{z}, \mathbf{p})$ . `MakeAcyclic` updates the allocation to  $(\mathbf{z}, \mathbf{p})$  and continues deleting cycles until the payment graph becomes acyclic. Since each step strictly decreases the number of edges in the payment graph and cycles can be found efficiently, `MakeAcyclic` terminates in polynomial time.

We prove using induction that the resulting allocation is an ER equilibrium. The initial allocation  $(\mathbf{y}, \mathbf{p})$  is an ER equilibrium and suppose the claim holds at some iteration with an updated allocation  $(\mathbf{y}, \mathbf{p})$ . Let  $(\mathbf{z}, \mathbf{p}, \mathbf{q}')$  be the next allocation. Notice for each agent  $i$ ,  $\sum_j q'_{ij} = \sum_j q_{ij} = e_i$ . Next for each chore, we have  $\sum_i q'_{ij} = \sum_i q_{ij} = \min\{p_j, c_j\}$ . Lastly if  $z_{ij} > 0$  then  $y_{ij} > 0$  as well. Thus the conditions of Definition 4 is satisfied, implying that  $(\mathbf{z}, \mathbf{p}, \mathbf{q}')$  is an ER equilibrium.  $\square$

## C Appendix to Section 3.3: $(n - 1)$ -EF1 and PO

In this section, we first present Algorithm 6, which takes as input an ER equilibrium  $(\mathbf{y}, \mathbf{p})$  of an instance with  $m \geq n$  and earning limit  $\beta = 1$ . Algorithm 6 performs essentially the same rounding algorithm as in Algorithm 1, except that the chore sets  $L$  and  $H$  are defined differently as  $L = \{j \in M : p_j \leq 1/2\}$  and  $H = \{j \in M : p_j > 1/2\}$ . We note that Lemma 3 (polynomial run-time) and Lemma 4 (allocation is always fPO) are still applicable to Algorithm 6.

Analogous to Lemmas 5 and 6, we prove upper and lower bounds on the earning of agents in the allocation returned by Algorithm 6.

**Lemma 36.** *Let  $(\mathbf{x}, \mathbf{p})$  be the allocation returned by Algorithm 6 with earning restriction  $\beta = 1$ . Then for each  $i \in N$ ,  $\mathbf{p}_{-1}(\mathbf{x}_i) \leq 1$ .*

---

**Algorithm 6** ER Rounding for  $(n - 1)$ -EF1 and PO

---

**Input:** Instance  $(N, M, D)$ , with  $m \geq n$  for earning limit  $\beta = 1$ ; an ER equilibrium  $(\mathbf{y}, \mathbf{p})$

**Output:** An integral allocation  $\mathbf{x}$

```
1:  $(\mathbf{z}, \mathbf{p}) \leftarrow \text{MakeAcyclic}(\mathbf{y}, \mathbf{p})$ 
2: Let  $G = (N, M, E)$  be the payment graph associated with  $(\mathbf{z}, \mathbf{p})$ 
3: Root each tree of  $G$  at some agent and orient edges
4:  $\mathbf{x}_i \leftarrow \emptyset$  for all  $i \in N$  ▷ Initialize empty allocation
5:  $L = \{j \in M : p_j \leq 1/2\}$ ,  $H = \{j \in M : p_j > 1/2\}$  ▷ Low, High paying chores
   — Phase 1: Round leaf chores —
6: for all leaf chores  $j$  do
7:    $\mathbf{x}_i \leftarrow \mathbf{x}_i \cup \{j\}$  for  $i = \text{parent}(j)$ ; delete  $j$  from  $G$ 
   — Phase 2: Allocate  $L$  —
8: for every tree  $T$  of  $G$  do
9:   for every agent  $i$  of  $T$  in BFS order do
10:    if  $\mathbf{p}(\mathbf{x}_i) > 1$  then
11:      for every  $j \in \text{child}(i) \cap H$  do
12:        Assign  $j$  to agent  $h \in \text{child}(j)$  earning most from  $j$  among  $\text{child}(j)$ ; delete  $j$ 
13:      while  $\exists j \in \text{child}(i) \cap L$  s.t.  $\mathbf{p}(\mathbf{x}_i \cup \{j\}) \leq 1$  do
14:         $\mathbf{x}_i \leftarrow \mathbf{x}_i \cup \{j\}$ ; delete  $j$  from  $G$ 
15:      for every  $j \in \text{child}(i) \cap L$  do
16:        Assign  $j$  to arbitrary agent  $h \in \text{child}(j)$ ; delete  $j$  from  $G$ 
   — Phase 3: Pruning trees —
17: for chore  $j \in V(G) \cap M$  do
18:   if a  $i \in \text{child}(j)$  does not earn the most from  $j$  among agents in  $\text{child}(j)$  then
19:     Delete edge  $(j, i)$  from  $G$ 
   — Phase 4: Matching to allocate  $H$  —
20: for every tree  $T = (N(T) \cup M(T), E(T))$  of  $G$  do
21:    $h \leftarrow \arg \max_{i \in N(T)} \mathbf{p}(\mathbf{x}_i)$ 
22:   Compute a matching  $\sigma$  of  $i \in N(T) \setminus \{h\}$  to  $M(T)$ 
23:   for  $i \in N(T) \setminus \{h\}$  do
24:      $\mathbf{x}_i \leftarrow \mathbf{x}_i \cup \{\sigma(i)\}$ 
25: return  $\mathbf{x}$ 
```

---

*Proof.* Let  $\mathbf{x}^t$  denote the allocation after Phase  $t$ , for  $t \in [4]$ ; note that  $\mathbf{x}^4 = \mathbf{x}$ . Consider an agent  $i \in N$ . Let  $\hat{\mathbf{x}}_i$  be the allocation when Algorithm 1 visits  $i$  in Phase 2. Suppose  $\mathbf{p}(\hat{\mathbf{x}}_i) \leq 1$ . Then we have  $\mathbf{p}(\mathbf{x}_i^2) \leq 1$  at the end of Phase 2 after  $i$  is assigned a subset of  $\text{child}(i) \cap L$ . Subsequently,  $i$  can be assigned one more chore in Phase 4. Hence we have  $\mathbf{p}_{-1}(\mathbf{x}_i) \leq 1$  in this case.

On the contrary, suppose  $\mathbf{p}(\hat{\mathbf{x}}_i) > 1$ . Then Algorithm 1 will not allocate any chore to  $i$  in Phase 4, and hence  $\mathbf{x}_i = \mathbf{x}_i^2 = \hat{\mathbf{x}}_i$ . Note that either  $\hat{\mathbf{x}}_i = \mathbf{x}_i^1$  or  $\hat{\mathbf{x}}_i = \mathbf{x}_i^1 \cup \{j\}$ , where  $j = \text{parent}(i)$ . That is,  $\hat{\mathbf{x}}_i$  includes the chores  $\mathbf{x}_i^1$  allocated to  $i$  in Phase 1, and may include  $i$ 's parent chore  $j$ . Recall that Phase 1 rounds leaf chores to their parent agents, hence  $\mathbf{x}_i^1$  comprises of the leaf chores that are child chores of  $i$ .

Suppose there exists a chore  $j_1 \in \mathbf{x}_i^1$  such that  $p_{j_1} > 1$ , i.e., there is a leaf chore  $j_1$  rounded to  $i$  whose payment exceeds the earning limit  $\beta = 1$ . Then agent  $i$  earns  $e_i = 1$  from  $j_1$  and no other chore, implying that  $\mathbf{x}_i = \hat{\mathbf{x}}_i = \{j_1\}$ . Then  $\mathbf{p}_{-1}(\mathbf{x}_i) = 0 \leq 1$ .

Otherwise,  $\mathbf{p}(\mathbf{x}_i^1) \leq 1$ . Then  $\mathbf{p}_{-1}(\mathbf{x}_i) = \mathbf{p}_{-1}(\hat{\mathbf{x}}_i) \leq \mathbf{p}(\hat{\mathbf{x}}_i \setminus \{j\}) = \mathbf{p}(\mathbf{x}_i^1) \leq 1$ , showing that the claim holds in this case too.  $\square$

**Lemma 37.** *Let  $(\mathbf{x}, \mathbf{p})$  be the allocation returned by Algorithm 6. Let  $T = (N(T) \cup M(T), E(T))$  be a Phase 3 tree rooted at agent  $i_0$ .*

- (i) *If some agent in  $N(T)$  lost a child chore, then for every  $i \in N(T)$ ,  $\mathbf{p}(\mathbf{x}_i) \geq \frac{1}{2}$ .*
- (ii) *If no agent in  $N(T)$  lost a child chore and  $i_0$  received  $\mathbf{parent}(i_0)$  chore, then for every  $i \in N(T)$ ,  $\mathbf{p}(\mathbf{x}_i) \geq \frac{1}{|N(T)|}$ .*
- (iii) *If no agent in  $N(T)$  lost a child chore and  $i_0$  lost  $\mathbf{parent}(i_0)$  chore, then for every  $i \in N(T)$ ,  $\mathbf{p}(\mathbf{x}_i) \geq \frac{1}{2|N(T)|}$ .*

*Proof.* Let  $(\mathbf{z}, \mathbf{p})$  be the acyclic ER equilibrium computed before Phase 1. Let  $\mathbf{x}^t$  denote the allocation after Phase  $t$  of Algorithm 6, for  $t \in [4]$ . Note that  $\mathbf{x}^2 = \mathbf{x}^3$  since Phase 3 does not assign any chores and only deletes edges in  $G$ . Also note  $\mathbf{x}^4 = \mathbf{x}$ .

Consider a Phase 3 tree  $T$  rooted at agent  $i_0$ . Since  $T$  is a Phase 3 tree,  $T$  has exactly  $|N(T)| - 1$  chores, all of which belong to  $H$ . Phase 4 identifies the agent  $h \in \arg \max_{i \in N(T)} \mathbf{p}(\mathbf{x}_i^3)$ , and assigns a chore  $\sigma(i) \in H$  to every agent  $i \in N(T) \setminus \{h\}$  by computing a matching of  $M(T)$  to  $N(T) \setminus \{h\}$ . Since  $p_j > 1/2$  for  $j \in H$ , we have  $\mathbf{p}(\mathbf{x}_i) \geq p_{\sigma(i)} > 1/2$  for all  $i \in N(T) \setminus \{h\}$ . Hence we only need to prove lower bounds on the earning  $\mathbf{p}(\mathbf{x}_h)$  of the agent  $h$ . Note that  $\mathbf{x}_h = \mathbf{x}_h^3 = \mathbf{x}_h^2$ , since  $h$  is not allocated any chores in Phase 3 or 4. By choice of  $h$ , we also have that  $\mathbf{p}(\mathbf{x}_h) \geq \mathbf{p}(\mathbf{x}_i^3) = \mathbf{p}(\mathbf{x}_i^2)$  for all  $i \in N(T)$ . We now analyze three scenarios.

- (i) Some agent  $i \in N(T)$  lost a child chore  $j \in \mathbf{child}(i)$ . Suppose  $i$  lost  $j$  in Phase 2. If  $j \in H$ , then it must be that  $\mathbf{p}(\mathbf{x}_i^2) > 1$ . If  $j \in L$ , then it must be that  $\mathbf{p}(\mathbf{x}_i^2) \geq 1/2$ ; otherwise we would have assigned  $j$  to  $i$  in Phase 2. In either case, we have  $\mathbf{p}(\mathbf{x}_i^2) \geq 1/2$ , and hence  $\mathbf{p}(\mathbf{x}_h) \geq \mathbf{p}(\mathbf{x}_i^2) \geq 1/2$  by choice of  $h$ . Note that  $i$  cannot lose  $j \in \mathbf{child}(i)$  in Phase 3 since Phase 3 only deletes edges from a chore to some of its child agents. This proves (i).
- (ii) No agent in  $N(T)$  lost a child chore and  $i_0$  received  $j_0 = \mathbf{parent}(i_0)$ ; it is possible that  $\mathbf{parent}(i_0) = \emptyset$ . This implies that no agent in  $N(T)$  has lost *any* chore they were earning from in  $(\mathbf{z}, \mathbf{p})$ . Since the earning of each agent in  $(\mathbf{z}, \mathbf{p})$  equals 1, the total earning of agents in  $N(T)$  is at least  $|N(T)|$ . The earning from the  $|N(T)| - 1$  chores in  $M(T)$  is at most  $(|N(T)| - 1)$  due to the earning restriction on each chore in  $M(T)$ . Hence there is at least one agent  $i \in N(T)$  whose earning  $\mathbf{p}(\mathbf{x}_i^2)$  satisfies:

$$\mathbf{p}(\mathbf{x}_i^2) \geq \frac{|N(T)| - (|N(T)| - 1)}{|N(T)|} = \frac{1}{|N(T)|}.$$

Since  $\mathbf{p}(\mathbf{x}_h) \geq \mathbf{p}(\mathbf{x}_i^2)$  by choice of  $h$ , this proves (ii).

- (iii) No agent in  $N(T)$  lost a child chore and  $i_0$  lost  $j_0 = \mathbf{parent}(i_0)$ . In this case, no agent in  $N(T)$  except  $i_0$  has lost any chore they were earning from in  $(\mathbf{z}, \mathbf{p})$ . We evaluate the amount of earning  $i_0$  loses due to losing  $j_0$ . Suppose  $j_0 \in H$ . Then  $i_0$  must have lost  $j_0$  in either Phase 2 or 3 to some agent  $i' \in \mathbf{child}(j_0)$  since  $i_0$  was not earning the most from  $j_0$  among agents in  $\mathbf{child}(i_0)$ . Due to the earning limit, agents can earn at most 1 from  $j_0$ . Hence the earning from  $i_0$  from  $j_0$  is at most  $1/2$ . On the other hand, if  $j_0 \in L$ , then  $i_0$  earns at most  $p_{j_0} \leq 1/2$  from  $j_0$ . In either case, we find that  $i_0$  has only lost  $1/2$  in earning. Hence the total



earning of agents in  $N(T)$  is at least  $|N(T)| - 1/2$ , while that from the chores in  $M(T)$  is at most  $(|N(T)| - 1)$ . Hence there is at least one agent  $i \in N(T)$  whose earning  $\mathbf{p}(\mathbf{x}_i^2)$  satisfies:

$$\mathbf{p}(\mathbf{x}_i^2) \geq \frac{|N(T)| - 1/2 - (|N(T)| - 1)}{|N(T)|} = \frac{1}{2|N(T)|}.$$

Since  $\mathbf{p}(\mathbf{x}_h) \geq \mathbf{p}(\mathbf{x}_i^2)$  by choice of  $h$ , this proves (iii).  $\square$

**Theorem 7.** *Given an ER equilibrium for an instance  $(N, M, D)$  where  $m \geq n$ , Algorithm 6 returns a  $2(n - 1)$ -EF1 and fPO allocation in polynomial time.*

*Proof.* Let  $(\mathbf{x}, \mathbf{p})$  be the allocation returned by Algorithm 6 with  $\beta = 1$ . Consider a Phase 3 tree  $T = (N(T) \cup M(T), E(T))$  rooted at agent  $i_0$ . Clearly,  $|N(T)| \leq n$ . If  $i_0$  lost the chore  $\mathbf{parent}(j_0)$  to another agent  $i_1$ , it must be that  $|N(T)| \leq n - 1$  since  $i_1 \notin N(T)$ . We use these facts together with Lemma 37 to obtain that for all  $i \in N$ :

$$\mathbf{p}(\mathbf{x}_i) \geq \min \left\{ \frac{1}{2}, \frac{1}{n}, \frac{1}{2(n-1)} \right\} = \frac{1}{2(n-1)},$$

since  $n \geq 2$ . Moreover Lemma 36 implies that  $\mathbf{p}_{-1}(\mathbf{x}_h) \leq 1$  for any  $h \in N$ . Thus, for any pair of agents  $i, h$ , we have:

$$\mathbf{p}_{-1}(\mathbf{x}_h) \leq 1 = 2(n-1) \cdot \frac{1}{2(n-1)} \leq 2(n-1) \cdot \mathbf{p}(\mathbf{x}_i),$$

thus showing that  $\mathbf{x}$  is  $2(n-1)$ -EF1 by Lemma 1. Lemma 4 implies  $\mathbf{x}$  is fPO and Lemma 3 shows Algorithm 1 runs in polynomial time.  $\square$

## C.1 An improved algorithm

Next, we improve our previous result by proving Theorem 8.

**Theorem 8.** *Given an ER equilibrium of an instance with  $m \geq n$ , an  $(n - 1)$ -EF1 and fPO allocation can be found in polynomial time.*

Let  $(\mathbf{z}, \mathbf{p})$  be the  $2(n - 1)$ -EF1 and PO allocation returned by Algorithm 6. We obtained this fairness guarantee by showing that  $\mathbf{p}_{-1}(\mathbf{z}_i) \leq 1$  and  $\mathbf{p}(\mathbf{z}_h) \geq \frac{1}{2(n-1)}$  for all agents  $i, h \in N$ . Improving the lower bound to  $\mathbf{p}(\mathbf{z}_h) \geq \frac{1}{n-1}$  for all  $h \in N$  would imply that  $\mathbf{z}$  is  $(n - 1)$ -EF1 and PO. Our algorithm aims to construct such an allocation in the event that  $\mathbf{z}$  is not already  $(n - 1)$ -EF1.

To do so, we revisit Lemma 37, which shows lower bounds on the earning of agents in the allocation resulting from the matching phase of Algorithm 6.

We call a Phase 3 tree  $T$  ‘problematic’ if after running Phase 4, some agent in  $T$  has an earning strictly less than  $\frac{1}{n-1}$  in the resulting allocation  $(\mathbf{z}, \mathbf{p})$ . By Lemma 37, if (i) some agent in  $T$  lost a child chore, or (ii)  $i_0$  received  $\mathbf{parent}(i_0)$  and  $|N(T)| \leq n - 1$ , or (iii) if  $|N(T)| \leq \frac{n-1}{2}$ , then  $\mathbf{p}(\mathbf{z}_i) \geq \frac{1}{n-1}$  for every  $i \in N(T)$ , and hence  $T$  is not-problematic. This leaves two possibilities for a problematic tree: (i)  $N(T) = [n]$ , or (ii)  $T$  is large, i.e.,  $|N(T)| > \frac{n-1}{2}$ , and no agent in  $T$  has lost a child chore, and its root  $i_1$  lost its parent chore  $j_1 = \mathbf{parent}(i_1)$ .

We eliminate case (i) by showing that a Phase 3 tree  $T$  with  $n$  agents and  $n - 1$  chores is not problematic. Phase 4 selects an agent  $h \in \arg \max_{i \in [n]} \mathbf{p}(\mathbf{x}_i)$ , where  $\mathbf{x}$  is the allocation at end of Phase 3. In the matching phase, each agent  $i \in [n] \setminus \{h\}$  is assigned a single chore  $j_i$ , while  $h$  is not assigned any chore. The resulting allocation  $\mathbf{z}$  is therefore given by  $\mathbf{z}_h = \mathbf{x}_h$  and  $\mathbf{z}_i = \mathbf{x}_i \cup \{j_i\}$  for all  $i \neq h$ . The following shows that  $\mathbf{z}$  is actually 2-EF1.

- (i)  $i \neq h$  does not EF1-envy  $h$ , as  $\mathbf{p}_{-1}(\mathbf{z}_i) \leq \mathbf{p}(\mathbf{z}_i \setminus \{j_i\}) = \mathbf{p}(\mathbf{x}_i) \leq \mathbf{p}(\mathbf{x}_h) = \mathbf{p}(\mathbf{z}_h)$ .
- (ii)  $i \in [n]$  does not 2-EF1-envy  $\ell \neq h$ , as  $\mathbf{p}_{-1}(\mathbf{z}_i) \leq 1 \leq 2 \cdot p_{j_\ell} \leq \mathbf{p}(\mathbf{z}_\ell)$ , since  $p_{j_\ell} \geq 1/2$  as  $j_\ell \in H$ .

Therefore, a tree  $T$  is problematic iff case (ii) holds. If the allocation returned by Algorithm 6 is not  $(n-1)$ -EF1, then there must exist a ‘problematic’ Phase 3 tree  $T_1$ . Then  $|N(T_1)| > \frac{n-1}{2}$ , no agent in  $T_1$  lost a child chore, and the root  $i_1$  lost the  $j_1 = \text{parent}(i_1)$  chore to another agent. We have two cases:

**Case 1.** We first handle the case of  $j_1 \in L$ . The  $(j_1, i_1)$  edge must have been deleted in Phase 2 when  $j_1$  was assigned another agent  $i_2$  (who is either  $\text{parent}(j_1)$  or a sibling of  $i_1$ ). Our algorithm ‘unrolls’ parts of Algorithm 6 in the ‘old run’ and re-visits the event in Phase 2 which deleted the edge  $(j_1, i_1)$ . This must have happened during a BFS call to agent  $i_0$  in Phase 2, which happened before the BFS call to agent  $i_1$ . Let  $T_0 \supseteq T_1$  be the Phase 1 tree containing  $T_1$ . At this point, we ‘re-run’ Phase 2 on  $T_0$  by starting with  $i_1$  as the root agent of  $T_0$ . The chore  $j_1$  now becomes the child of  $i_1$ . We visit all child chores of  $i_1$  before visiting  $j_1$ .

Let  $T'_1$  be the Phase 3 tree rooted at  $i_1$  in the new run. Since  $T_1$  is a problematic Phase 3 tree,  $i_1$  received all of her child  $L$ -chores during Phase 2 of the old run. Since these child chores are visited before  $j_1$ ,  $i_1$  receives all of them in the re-run as well. This shows that irrespective of whether  $i_1$  is assigned  $j_1$  or not, the Phase 3 tree  $T'_1$  produced in the new run is such that  $N(T'_1) = N(T_1)$ .

If  $i_1$  is assigned  $j_1$ , then  $T'_1$  is not problematic, as its root has not lost its parent:  $i_1$  has no parent chore in the re-run. On the other hand, suppose  $i_1$  loses  $j_1$  to another agent  $i_2$  who is a part of a Phase 3 tree  $T_2$ . Once again,  $T'_1$  is not problematic as an agent  $i_1$  has lost a child chore  $j_1$ . Suppose  $T_2$  is problematic. Then  $|N(T_2)| > \frac{n-1}{2}$ . Note that  $T_2$  is disjoint from  $T'_1$ , and since  $T_1$  is problematic we have  $|N(T'_1)| = |N(T_1)| > \frac{n-1}{2}$  as well. Since  $|N(T_1)| + |N(T_2)| \leq n$ , the above inequalities can only hold if  $n = 2n'$  for  $n' \in \mathbb{N}$  and  $|N(T_1)| = |N(T_2)| = n'$ . This implies that the Phase 1 tree  $T_0$  comprises of trees  $T_1$  and  $T_2$  rooted at  $i_1$  and  $i_2$  respectively, both of which have edges to the chore  $j_1$ . In this case, we simply round  $j_1$  to the agent in  $\{i_1, i_2\}$  who earns more from  $j_1$ . Without loss of generality, suppose this agent is  $i_2$ . Then  $T_2$  is not problematic since its root received its parent chore. In  $T_1$ , agents have lost an earning of at most  $1/4$ , since  $p_{j_1} \leq 1/2$  as  $j_1 \in L$ , and  $i_1$  earned at most as much as  $i_2$  did from  $j_1$ . Hence every agent in  $T_1$  earns at least  $\frac{1-1/4}{n'} = \frac{3}{2n}$ . For  $n \geq 3$ ,  $\frac{3}{2n} \geq \frac{1}{n-1}$ , showing that agents get the desired lower bound of  $\frac{1}{n-1}$  on their earning. For  $n = 2$ , it is easy to see that the resulting allocation is in fact EF1.

**Case 2.** We now handle the case of  $j_1 \in H$ . The  $(j_1, i_1)$  edge must have been deleted either in Phase 2 (during the BFS call to agent  $i_0 = \text{parent}(j_1)$ ) or Phase 3 (because  $i_1$  was not earning the most from  $j_1$  among  $\text{child}(j)$ ). In either case,  $j_1$  retains an edge to a sibling  $i_2$  of  $i_1$  in the Phase 3 tree  $T'$  containing  $i_2$ . Let  $T_2 \subseteq T'$  be the subtree rooted at  $i_2$ . Since  $T'$  is a Phase 3 tree,  $T_2$  is a Phase 3 tree as well, i.e., every chore in  $T_2$  is adjacent to exactly two agents.

Let  $s_1$  and  $s_2$  be the earning of agents  $i_1$  and  $i_2$  from  $j_1$  respectively. Following the proof of Lemma 37, we observe that agents in  $T_1$  have lost at most  $s_1$  total earning. Hence there must exist some agent in  $T_1$  who earns at least  $\frac{1-s_1}{|N(T_1)|}$  from the chores assigned integrally thus far. If  $\frac{1-s_1}{|N(T_1)|} \geq \frac{1}{n-1}$ , then the allocation  $\mathbf{z}$  must have already been  $(n-1)$ -EF1, hence we assume  $\frac{1-s_1}{|N(T_1)|} < \frac{1}{n-1}$ . This gives:  $s_1 > 1 - \frac{|N(T_1)|}{n-1}$ . Since the earning from each chore is at most 1, we have  $s_1 + s_2 \leq 1$ . Thus we obtain  $s_2 < \frac{|N(T_1)|}{n-1}$ .

Our algorithm now ‘unrolls’ parts of Algorithm 6 by re-visiting the event which deleted the edge  $(j_1, i_1)$ . Instead, the edge  $(j_1, i_1)$  is re-introduced and the edge  $(j_1, i_2)$  is deleted. This results

in a larger tree  $T''$  which contains  $T_1$ , and the Phase 3 tree  $T_2$ . After Phase 4, every agent in  $T_2$  earns at least:

$$\frac{1 - s_2}{|N(T_2)|} > \frac{1 - |N(T_1)|/(n-1)}{|N(T_2)|} \geq \frac{1}{n-1},$$

where we used the bound  $s_2 < \frac{|N(T_1)|}{n-1}$  derived earlier for the first inequality, and the second inequality used the fact that  $|N(T_1)| + |N(T_2)| \leq n-1$ , since  $i_0 = \mathbf{parent}(j_1) \notin T_1 \cup T_2$ . Thus  $T_2$  is not problematic. If  $T''$  is problematic, we recurse and repeat our algorithm with  $T''$  instead, i.e., set  $T_1 \leftarrow T''$ . Since  $N(T'') \supseteq N(T_1) \cup \{i_0\}$ , every recursive step increases the size of  $T_1$ . Eventually it must happen that  $T_1$  is non-problematic, or its root agent has no grand-parent agent. In the latter case, this tree  $T_1$  must be non-problematic, since its root has not lost its parent root, and  $|N(T_1)| \leq n-1$  since  $T_2 \subsetneq T_1$ . The algorithm therefore terminates with at most  $n$  recursive calls.

## D Appendix to Section 4

**Lemma 10.** *Given a chore allocation instance with  $m = n$  chores, an MPB allocation  $(\mathbf{z}, \mathbf{q})$  where  $|\mathbf{z}_i| = 1$  for each  $i \in N$  can be computed in polynomial time.*

*Proof.* We show that the required allocation  $\mathbf{z}$  can be computed via the following linear program for finding a minimum cost matching.

$$\begin{aligned} & \min \sum_{i \in N} \sum_{j \in M} \log d_{ij} \cdot x_{ij} \\ & \forall j \in M : \sum_{i \in N} x_{ij} = 1 \\ & \forall i \in N : \sum_{j \in M} x_{ij} = 1 \\ & \forall i \in N, j \in M : x_{ij} \geq 0. \end{aligned} \tag{11}$$

Note that the objective is well-defined since  $d_{ij} > 0$  for all  $i \in N, j \in M$ . Since the matching polytope is integral, there exists a integral optimal solution  $\mathbf{z}$  with  $|\mathbf{z}_i| = 1$  for all  $i \in N$ . We now show that we can compute chore payments  $\mathbf{q}$  such that  $(\mathbf{z}, \mathbf{q})$  is on MPB by using dual variables of the above program. Let  $\lambda_j$  and  $\alpha_i$  be the dual variables corresponding to the constraints corresponding to chore  $j$  and agent  $i$  respectively. The stationarity KKT condition corresponding to the variable  $x_{ij}$  implies:

$$\log d_{ij} + \lambda_j + \alpha_i \geq 0.$$

This implies that for all  $i \in N$  and  $j \in M$ ,  $\frac{d_{ij}}{e^{-\lambda_j}} \geq e^{-\alpha_i}$ . Moreover, the complementary slackness condition implies that the above inequality is an equality when  $x_{ij} > 0$ , i.e.,  $x_{ij} > 0 \Rightarrow \frac{d_{ij}}{e^{-\lambda_j}} = e^{-\alpha_i}$ . We set the chore payments  $\mathbf{q}$  as  $q_j = e^{-\lambda_j} > 0$ . The above observations then imply that  $(\mathbf{z}, \mathbf{q})$  is on MPB.  $\square$

## E Appendix to Section 5: EFX and PO for $m \leq 2n$

Algorithm 7 essentially follows the same template as Algorithm 5, except that it begins with a balanced allocation  $(\mathbf{x}^0, \mathbf{p})$  computed using Algorithm 2. When  $m \leq n$ ,  $\mathbf{x}^0$  is EFX and PO since every agent gets at most one chore. Hence we assume  $m > n$ . Since the allocation is balanced, we know  $1 \leq |\mathbf{x}_i^0| \leq 2$  for all  $i \in N$ .

For bivalued instances, we can scale the payments to ensure that for all  $j \in M$ ,  $p_j \in \{1, k\}$ .

**Lemma 38.** *Let  $\mathbf{p}$  be the payment vector at the end of Algorithm 2 on a  $\{1, k\}$ -bivalued instance. Then there exists  $r \in \mathbb{Z}_{\geq 0}$  such that  $p_j \in \{k^r, k^{r+1}\}$  for all  $j \in M$ .*

*Proof.* We show that all chore payments in the run of Algorithm 2 are powers of  $k$ . This is true initially since all chores are allocated to agent  $h$  and pay 1 or  $k$ . If all chore payments are powers of  $k$  and the two possible disutility values differ by a factor of  $k$ , then the payment raise coefficient  $\beta$  must also be a power of  $k$ . Thus, all chore payments continue to be powers of  $k$  after the payment raise. Thus, it must be that all chore payments remain a power of  $k$  throughout the run of Algorithm 2.

Now suppose for sake of contradiction that there exist chores  $j_1$  and  $j_2$  such that  $p_{j_1} = k^r$  and  $p_{j_2} = k^{r+s}$ , where  $s > 1$ . Since we have seen that Algorithm 2 maintains a CE, it must be that  $j_1$  is MPB for the agent  $i$  it is allocated to. However, we have that  $\alpha_{ij_2} = \frac{d_{ij_2}}{p_{j_2}} \leq \frac{k}{k^{r+s}} < \frac{1}{k^r} \leq \frac{d_{ij_1}}{p_{j_1}} = \alpha_{ij_1}$ . Thus,  $j_1$  cannot be MPB for agent  $i$ , and we have a contradiction. It must then be that in fact  $s = 1$ , showing the result.  $\square$

We classify the chores as low-paying,  $L = \{j : p_j = 1\}$ , and high paying  $H = \{j : p_j = k\}$ . As in Definition 9, we define classify agents into sets  $N_L$ ,  $N_H^1$ , and  $N_H^2$  depending on whether they have only  $L$ -chores, a single  $H$ -chore, or two  $H$ -chores. We first note that  $\mathbf{x}^0$  is EFX for agents in  $N_L$ , since  $\mathbf{p}_{-X}(\mathbf{x}_i) = \mathbf{p}_{-1}(\mathbf{x}_i) \leq 1 \leq \mathbf{p}(\mathbf{x}_h)$  for any  $i \in N_L$  and  $h \in N$ . Thus, if  $H = \emptyset$ ,  $N = N_L$  and Algorithm 7 simply returns  $\mathbf{x}^0$ . We therefore assume  $H \neq \emptyset$  subsequently. With this assumption, the following statement regarding the MPB ratios of agents analogous to Lemma 20 holds.

**Lemma 39.** *Assume  $H \neq \emptyset$ . Then:*

- (i) *For all  $i \in N_L$ ,  $\alpha_i = 1$ . Moreover for every  $j \in H$ ,  $d_{ij} = k$  and  $j \in \text{MPB}_i$ .*
- (ii) *For all  $i \in N_H$ ,  $\alpha_i \in \{1, 1/k\}$ .*
- (iii) *For all  $i \in N_H$ , if  $\mathbf{x}_i \setminus H \neq \emptyset$  then  $\alpha_i = 1$ .*

If  $\mathbf{x}^0$  is not EFX, some agent in  $N_H$  must EFX-envy another  $\ell$  agent. Algorithm 5 addresses the EFX-envy of agents in  $N_H$  by swapping some chores between agents  $i$  and  $\ell$  by performing the same swap steps defined in Algorithm 5. The only point of difference is that Algorithm 7 performs a swap if  $i$  EFX-envies  $\ell$ , whereas Algorithm 5 performs it if  $i$  3-EFX-envies  $\ell$ . Since there is a limited number of chores, Algorithm 7 can ensure agents in  $N_H$  do not have too much cost: agents in  $N_H^2$  have exactly two  $H$  chores and no other chores, while agents in  $N_H^1$  have exactly one  $H$  chore and at most one  $L$  chore.

We now prove the above claims formally.

**Lemma 40.** *(Invariants of Alg. 7) Let  $(\mathbf{x}, \mathbf{p})$  be an allocation in the run of Algorithm 7. Then:*

- (i)  *$(\mathbf{x}, \mathbf{p})$  is an MPB allocation.*
- (ii) *For all  $i \in N$ ,  $\mathbf{p}(\mathbf{x}_i) \geq 1$ .*
- (iii) *For all  $i \in N_L$ ,  $\mathbf{p}_{-1}(\mathbf{x}_i) \leq 1$  during Phase 1.*
- (iv) *For all  $i \in N_L$ ,  $\mathbf{p}(\mathbf{x}_i) < 1 + k$ .*
- (v) *For all  $i \in N_H^1$ ,  $|\mathbf{x}_i \setminus H| \leq |\mathbf{x}_i \cap H| = 1$ .*
- (vi) *For all  $i \in N_H^2$ ,  $|\mathbf{x}_i| = |\mathbf{x}_i \cap H| = 2$ .*

---

**Algorithm 7** EFX + PO for bivalued instances with  $m \leq 2n$ 

---

**Input:**  $\{1, k\}$ -bivalued instance  $(N, M, D)$  with  $m \leq 2n$ **Output:** An integral allocation  $\mathbf{x}$ 

- 1:  $(\mathbf{x}, \mathbf{p}) \leftarrow$  Algorithm 2 on  $(N, M, D)$  ▷ For all  $j \in M$ ,  $p_j \in \{1, k\}$
  - 2: **if**  $m \leq n$  **then return**  $\mathbf{x}$
  - 3:  $L = \{j \in M : p_j = 1\}$ ,  $H = \{j \in M : p_j = k\}$  ▷ Low, High paying chores
  - 4: Classify agents as  $N_L, N_H^1, N_H^2$  as before  
— Phase 1: Address  $N_H^2$  agents —
  - 5: **while**  $\exists i \in N_H^2$  not EFX **do** ▷ Lemma 22 shows  $\ell \in N_L$
  - 6:    $\ell \leftarrow$  agent EFX-envied by  $i$
  - 7:   **if**  $\mathbf{p}(\mathbf{x}_\ell) > 1$  **then**  $S \leftarrow j_1$  for some  $j_1 \in \mathbf{x}_\ell$
  - 8:   **else**  $S \leftarrow \emptyset$
  - 9:    $j \in \mathbf{x}_i \cap H$
  - 10:    $\mathbf{x}_\ell \leftarrow \mathbf{x}_\ell \setminus S \cup \{j\}$
  - 11:    $\mathbf{x}_i \leftarrow \mathbf{x}_i \cup S \setminus \{j\}$
  - 12:    $N_H^1 \leftarrow N_H^1 \cup \{i, \ell\}$ ,  $N_H^2 \leftarrow N_H^2 \setminus \{i\}$ ,  $N_L \leftarrow N_L \setminus \{\ell\}$   
— Phase 2: Address  $N_H^1$  agents —
  - 13: **while**  $\exists i \in N_H^1$  not EFX **do** ▷ Lemma 22 shows  $\ell \in N_L$
  - 14:    $\ell \leftarrow \arg \min\{\mathbf{p}(\mathbf{x}_h) : h \in N \text{ s.t. } i \text{ EFX envies } h\}$
  - 15:    $j \in \mathbf{x}_i \cap H$
  - 16:    $\mathbf{x}_i \leftarrow \mathbf{x}_i \cup \mathbf{x}_\ell \setminus \{j\}$
  - 17:    $\mathbf{x}_\ell \leftarrow \{j\}$
  - 18:    $N_H^1 \leftarrow N_H^1 \cup \{\ell\} \setminus \{i\}$ ,  $N_L \leftarrow N_L \cup \{i\} \setminus \{\ell\}$
  - 19: **return**  $\mathbf{x}$
- 

We prove the above lemma using Lemmas 42 and 43 below. Like Lemma 22, we show that an agent in  $N_H = N_H^1 \cup N_H^2$  can only EFX-envy another agent in  $N_L$ .

**Lemma 41.** *Consider an allocation  $(\mathbf{x}, \mathbf{p})$  satisfying the invariants of Lemma 40. If  $i \in N_H$  EFX-envies  $\ell$ , then  $\alpha_i = 1$ ,  $\ell \in N_L$ , and  $\mathbf{x}_\ell \subseteq MPB_i$ .*

*Proof.* Consider  $i \in N_H$  who EFX-envies  $\ell \in N$ . We know from Lemma 39 that  $\alpha_i \in \{1, 1/k\}$ . Suppose  $\alpha_i = 1/k$ . Then  $d_{ij} = 1$  for all  $j \in \mathbf{x}_i$ . By the contrapositive of Lemma 20 (iii), we get  $\mathbf{x}_i \subseteq H$ . Since  $|\mathbf{x}_i \cap H| \leq 2$ , we get  $|\mathbf{x}_i| \leq 2$ . Thus,  $\max_{j \in \mathbf{x}_i} d_i(\mathbf{x}_i \setminus \{j\}) \leq 1 \leq d_i(\mathbf{x}_\ell)$ , since  $\mathbf{x}_\ell \neq \emptyset$  and the instance is bivalued. Thus  $i$  is EFX towards  $\ell$  if  $\alpha_i = 1/k$ , which implies that  $\alpha_i = 1$ .

Suppose  $\exists j \in \mathbf{x}_\ell$  such that  $d_{ij} = k$ . Then invariants (v) and (vi) imply that  $\max_{j' \in \mathbf{x}_i} d_i(\mathbf{x}_i \setminus \{j'\}) \leq k \leq d_{ij} \leq d_i(\mathbf{x}_\ell)$ , showing that  $i$  is EFX towards  $\ell$ . Thus it must be that for all  $j \in \mathbf{x}_\ell$ ,  $d_{ij} = 1$ . The MPB condition for  $i$  implies that  $\alpha_i \leq d_{ij}/p_j$ , showing that  $p_j \leq 1$ , and hence  $p_j = 1$  for all  $j \in \mathbf{x}_\ell$ . Thus  $\ell \in N_L$ . Moreover, for any  $j \in \mathbf{x}_\ell$ ,  $\alpha_i = d_{ij}/p_j$ , and hence  $\mathbf{x}_\ell \subseteq MPB_i$ .  $\square$

The next two lemmas establish the invariants of Lemma 40.

**Lemma 42.** *The invariants of Lemma 40 are maintained during Phase 1 of Algorithm 7.*

*Proof.* We prove the statement inductively. We first show that the invariants hold at  $(\mathbf{x}^0, \mathbf{p})$ . Invariants (i), (ii), (iii), (v) and (vi) follow from the fact that  $\mathbf{x}^0$  is a balanced allocation. For (iv), note that for any  $i \in N_L$ , we have  $\mathbf{p}_{-1}(\mathbf{x}_i^0) \leq 1$ . Thus  $\mathbf{p}(\mathbf{x}_i^0) \leq 2 < 1 + k$ .

Suppose the invariants hold at an allocation  $(\mathbf{x}, \mathbf{p})$  during Phase 1. Consider a Phase 1 swap involving agents  $i \in N_H^2$  and  $\ell \in N$ . Given that Algorithm 5 performed the swap,  $i$  must EFX-envy

$\ell$ . Lemma 22 implies that  $\ell \in N_L$  and hence  $\mathbf{x}_\ell \in L$ . As per Algorithm 5, if  $\mathbf{p}(\mathbf{x}_\ell) > 1$ , then  $S = \{j_1\}$  for some  $j_1 \in \mathbf{x}_\ell$ , otherwise  $S = \emptyset$ . Let  $j \in \mathbf{x}_i \cap H$ .

Let  $\mathbf{x}'$  be the resulting allocation. Thus  $\mathbf{x}'_i = \mathbf{x}_i \setminus \{j\} \cup S$ ,  $\mathbf{x}'_\ell = \mathbf{x}_\ell \setminus S \cup \{j\}$ , and  $\mathbf{x}'_h = \mathbf{x}_h$  for all  $h \notin \{i, \ell\}$ . We show that the invariants hold at  $(\mathbf{x}', \mathbf{p})$ . Since a Phase 1 step removes agents  $i$  and  $\ell$  from  $N_H^2$  and  $N_L$  respectively, invariants (iii), (iv), (vi) continue to hold. For the rest, observe:

(i)  $(\mathbf{x}', \mathbf{p})$  is on MPB. This is because Lemma 41 implies  $S \subseteq \mathbf{x}_\ell \subseteq MPB_i$ , showing  $\mathbf{x}'_i \subseteq MPB_i$ . Since  $\ell \in N_L$  at  $(\mathbf{x}, \mathbf{p})$  and  $j \in H$ , Lemma 20 shows  $j \in MPB_\ell$  and hence  $\mathbf{x}'_\ell \subseteq MPB_\ell$ .

(ii) Follows from  $\mathbf{x}'_i \neq \emptyset$  and  $\mathbf{x}'_\ell \neq \emptyset$ .

(v) For agent  $i$ , note that  $\mathbf{x}'_i$  contains exactly one  $H$ -chore and perhaps one  $L$ -chore  $j_1$ . Hence  $|\mathbf{x}'_i \setminus H| \leq 1 = |\mathbf{x}'_i \cap H|$ , proving invariant (v).

For agent  $\ell$ , note that  $\mathbf{x}'_\ell$  contains exactly one  $H$ -chore  $j$ , hence  $|\mathbf{x}'_\ell \cap H| = 1$ . Since invariant (iii) implies  $|\mathbf{x}_\ell \setminus H| \leq 2$ ,  $|\mathbf{x}'_\ell \setminus H| \leq 1$  after the potential transfer of  $j_1$ , thus proving (v).

The swap does not affect an agent  $h \notin \{i, \ell\}$  and hence the invariants continue to hold for  $h$  after the swap. By induction, we have shown that the invariants of Lemma 40 hold after any Phase 1 swap.  $\square$

**Lemma 43.** *The invariants of Lemma 40 are maintained during Phase 2 of Algorithm 7. Moreover, agents in  $N_H^2$  remain EFX towards other agents.*

*Proof.* We prove the statement inductively. Lemma 42 shows the invariants hold at the end of Phase 1. Suppose the invariants hold at an allocation  $(\mathbf{x}, \mathbf{p})$  during Phase 2. Consider a Phase 2 swap involving agents  $i \in N_H^1$  and  $\ell \in N$ . Given that Algorithm 7 performed the swap,  $i$  must EFX-envy  $\ell$ . Lemma 41 implies that  $\ell \in N_L$  and hence  $\mathbf{x}_\ell \in L$ . Let  $j \in \mathbf{x}_i \cap H$ .

Let  $\mathbf{x}'$  be the resulting allocation. Thus  $\mathbf{x}'_i = \mathbf{x}_i \setminus \{j\} \cup \mathbf{x}_\ell$ ,  $\mathbf{x}'_\ell = \{j\}$ , and  $\mathbf{x}'_h = \mathbf{x}_h$  for all  $h \notin \{i, \ell\}$ . We now show that the invariants hold at  $(\mathbf{x}', \mathbf{p})$ . Since we are in Phase 2, invariant (iii) does not apply, and since Phase 2 swaps do not alter the allocation of agents in  $N_H^2$ , invariant (vi) continues to hold. For the rest, observe:

(i)  $(\mathbf{x}', \mathbf{p})$  is on MPB. This is because Lemma 41 implies  $\mathbf{x}_\ell \subseteq MPB_i$ , showing  $\mathbf{x}'_i \subseteq MPB_i$ . Since  $\ell \in N_L$  at  $(\mathbf{x}, \mathbf{p})$  and  $j \in H$ , Lemma 39 shows  $j \in MPB_\ell$  and hence  $\mathbf{x}'_\ell \subseteq MPB_\ell$ .

(ii) Follows from  $\mathbf{x}'_i \neq \emptyset$  and  $\mathbf{x}'_\ell \neq \emptyset$ .

(iv) We want to show  $\mathbf{p}(\mathbf{x}'_i) \leq 1 + k$ . To see this note that since  $i$  EFX-envies  $\ell$  in  $\mathbf{x}$ ,  $i$  must pEFX-envy  $\ell$  in  $(\mathbf{x}, \mathbf{p})$ . Using invariant (v), this means that  $\mathbf{p}_{-X}(\mathbf{x}_i) = k > \mathbf{p}(\mathbf{x}_\ell)$ .

Now  $\mathbf{p}(\mathbf{x}'_i) = \mathbf{p}(\mathbf{x}_i \setminus \{j\}) + \mathbf{p}(\mathbf{x}_\ell) < 1 + k$ , where we used  $\mathbf{p}(\mathbf{x}_i \setminus \{j\}) = 1$  since invariant (v) shows  $|\mathbf{x}_i \setminus H| \leq 1$ .

(v) Note that  $\ell \in N_H^1$  in  $(\mathbf{x}', \mathbf{p})$ , and  $|\mathbf{x}'_\ell \setminus H| = 0 < 1 = |\mathbf{x}'_\ell \cap H|$ .

The swap does not affect an agent  $h \notin \{i, \ell\}$  and hence the invariants continue to hold for  $h$  after the swap. By induction, we conclude that the invariants of Lemma 21 hold after any Phase 2 swap.

We now show that  $i \in N_H^2$  cannot EFX-envy an agent  $\ell \in N$ . Lemma 22 implies that  $\ell \in N_L$  and hence  $\mathbf{x}_\ell \subseteq L$ . Let  $\mathbf{x}^1$  be the allocation at the end of Phase 1. Note that the bundle  $\mathbf{x}_\ell \subseteq L$  is obtained via a series of Phase 2 swaps initiated with some agent  $\ell_1$  in  $(\mathbf{x}^1, \mathbf{p})$ . Here,  $\ell_1 \in N_L$  at  $(\mathbf{x}^1, \mathbf{p})$ . Thus  $\mathbf{x}_\ell \supseteq \mathbf{x}_{\ell_1}^1$ . Agent  $i \in N_H^2$  did not EFX-envy  $\ell_1$  in  $\mathbf{x}^1$ , otherwise Algorithm 7 would

have performed a Phase 1 swap between agent  $i$  and  $\ell_1$ . Since  $\mathbf{x}_i = \mathbf{x}_i^1$  as Algorithm 7 does not alter allocation of agents in  $N_H^2$  and  $\mathbf{x}_\ell \supseteq \mathbf{x}_{\ell_1}^1$ ,  $i$  will not EFX-envy  $\ell$  in  $\mathbf{x}$  either. Thus, all agents in  $N_H^2$  continue to remain EFX during Phase 2.  $\square$

We need one final lemma showing that  $N_L$  agents do not EFX-envy any other agent.

**Lemma 44.** *At any allocation  $(\mathbf{x}, \mathbf{p})$  in the run of Algorithm 7,  $\mathbf{x}$  is EFX for every agent in  $N_L$ .*

*Proof.* We know that the initial allocation  $\mathbf{x}^0$  is EFX for agents in  $N_L$ . Let  $\mathbf{x}$  be the earliest allocation in the run of Algorithm 7 in which an agent  $i \in N_L$  EFX-envies another agent  $h \in N$ . Using  $\alpha_i = 1$  from Lemma 39, the bound on  $\mathbf{p}(\mathbf{x}_i)$  from Lemma 40 (iv), we note:

$$\max_{j' \in \mathbf{x}_i} d_i(\mathbf{x}_i \setminus \{j'\}) = \alpha_i \cdot \mathbf{p}_{-X}(\mathbf{x}_i) < (1+k) - 1 = k. \quad (12)$$

Thus if  $\exists j \in \mathbf{x}_h$  s.t.  $d_{ij} = k$ , then by (12),  $d_i(\mathbf{x}_i) < k \leq d_i(\mathbf{x}_h)$ , showing that  $i$  does not EFX-envy  $h$  in  $\mathbf{x}$ . Hence it must be that for all  $j \in \mathbf{x}_h$ ,  $d_{ij} = 1$ . This also implies  $\mathbf{x}_h \subseteq MPB_i$ , since  $\alpha_i = 1 = d_{ij}/p_j$  for any  $j \in \mathbf{x}_h$ . We now consider two cases based on the category of  $h$ .

- $h \in N_H$ . By definition of  $N_H$ ,  $\exists j \in \mathbf{x}_h$  s.t.  $j \in H$ . Since  $i \in N_L$ , by Lemma 39 (i) we get  $d_{ij} = k$ , which is a contradiction.
- $h \in N_L$ . Since  $\mathbf{x}^0$  is EFX for agents in  $N_L$ , and Phase 1 swaps only remove agents from  $N_L$ , it cannot be that  $i$  starts EFX-envying  $h \in N_L$  during Phase 1. Let  $\mathbf{x}'$  be the preceding allocation, at which Algorithm 7 performed a Phase 2 swap. Since  $\mathbf{x}$  is the earliest allocation in which  $i$  EFX-envies  $h$ , it must be in  $\mathbf{x}'$ , agent  $i$  was in  $N_H^1$  and was involved with a Phase 2 swap with another agent  $\ell \in N_L$ . Since  $\mathbf{x}_h \subseteq MPB_i$  and Algorithm 7 did *not* perform a swap between agents  $i$  and  $h$  in the allocation  $\mathbf{x}'$ , we must have  $\mathbf{p}(\mathbf{x}'_\ell) \leq \mathbf{p}(\mathbf{x}'_h)$  by the choice of  $\ell$  at  $(\mathbf{x}', \mathbf{p})$ .

Note that  $\mathbf{x}_i = (\mathbf{x}'_i \setminus H) \cup \mathbf{x}'_\ell$ . By Lemma 21 (i), we know  $\mathbf{p}_{-1}(\mathbf{x}'_i) = \mathbf{p}(\mathbf{x}'_i \setminus H) \leq 1$ . Thus:

$$\mathbf{p}_{-X}(\mathbf{x}_i) = \mathbf{p}(\mathbf{x}'_i \setminus H) + \mathbf{p}(\mathbf{x}'_\ell) - 1 \leq \mathbf{p}(\mathbf{x}'_\ell) \leq \mathbf{p}(\mathbf{x}'_h),$$

where the last inequality used  $\mathbf{x}_h = \mathbf{x}'_h$ . Thus,  $i$  is EFX towards  $h$ .

We conclude that it is not possible for an agent  $i \in N_L$  to EFX-envy any other agent during the course of Algorithm 7.  $\square$

We are now in a position to summarize and conclude our analysis of Algorithm 7.

**Theorem 14.** *Given a bivalued instance with  $m \leq 2n$ , Algorithm 7 returns an EFX and fPO allocation in polynomial time.*

*Proof.* Let  $(\mathbf{x}^0, \mathbf{p})$  be the initial balanced allocation obtained by using Algorithm 2. If  $m \leq n$  or  $H = \emptyset$ , then  $\mathbf{x}^0$  is EFX, hence we assume otherwise.

Lemma 44 shows that any allocation  $\mathbf{x}$  in the course of Algorithm 7 is EFX for agents in  $N_L$ . Any potential EFX-envy is, therefore, from some agent  $i \in N_H$ . Lemma 41 shows that if  $i \in N_H$  is not EFX towards  $\ell$ , then  $\ell \in N_L$ . If  $i \in N_H^2$ ,  $i$  participates in a Phase 1 swap with agent  $\ell$ , after which  $i$  and  $\ell$  get removed from  $N_H^2$  and  $N_L$  respectively. This implies that Phase 1 terminates after at most  $n/2$  swaps, and the resulting allocation is EFX for all agents in  $N_H^2$ . If  $i \in N_H^1$ ,  $i$  participates in a Phase 2 swap with agent  $\ell$ , after which  $\ell$  is added to  $N_H^1$  and is assigned a single chore and  $\ell$  does not EFX-envy. This implies that Phase 2 terminates after at most  $n$  swaps since the number of agents in  $N_H^1$  who are not EFX strictly decreases. The resulting allocation is EFX

for all agents in  $N_H^1$ . Lemma 43 also shows that Phase 2 swaps do not cause  $N_H^2$  agents to start EFX-envying any agent in  $N_L$ . Thus the allocation on termination of Algorithm 7 is EFX. By invariant (i) of Lemma 40,  $\mathbf{x}$  is also fPO. Since there are at most  $3n/2$  swaps and Algorithm 2 takes polynomial time, Algorithm 7 terminates in polynomial time.  $\square$



## References

- [1] Georgios Amanatidis, Georgios Birmpas, Aris Filos-Ratsikas, Alexandros Hollender, and Alexandros A. Voudouris. Maximum Nash welfare and other stories about EFX. In *Proceedings of the Twenty-Ninth International Joint Conference on Artificial Intelligence, (IJCAI)*, pages 24–30, 2020.
- [2] Georgios Amanatidis, Haris Aziz, Georgios Birmpas, Aris Filos-Ratsikas, Bo Li, Hervé Moulin, Alexandros A. Voudouris, and Xiaowei Wu. Fair division of indivisible goods: Recent progress and open questions. *Artificial Intelligence*, 322:103965, 2023.
- [3] Georgios Amanatidis, Aris Filos-Ratsikas, and Alkmini Sgouritsa. Pushing the frontier on approximate EFX allocations. In *Conf. Economics and Computation (EC)*, 2024.
- [4] Haris Aziz, Péter Biró, Jérôme Lang, Julien Lesca, and Jérôme Monnot. Efficient reallocation under additive and responsive preferences. *Theoretical Computer Science*, 790:1 – 15, 2019.
- [5] Haris Aziz, Ioannis Caragiannis, Ayumi Igarashi, and Toby Walsh. Fair allocation of indivisible goods and chores. In *Proceedings of the 28th International Joint Conference on Artificial Intelligence (IJCAI)*, page 53–59, 2019.
- [6] Haris Aziz, Hervé Moulin, and Fedor Sandomirskiy. A polynomial-time algorithm for computing a Pareto optimal and almost proportional allocation. *Oper. Res. Lett.*, 48(5):573–578, 2020.
- [7] Haris Aziz, Bo Li, Herve Moulin, and Xiaowei Wu. Algorithmic fair allocation of indivisible items: A survey and new questions, 2022.
- [8] Haris Aziz, Jeremy Lindsay, Angus Ritossa, and Mashbat Suzuki. Fair allocation of two types of chores. In *Proceedings of the International Conference on Autonomous Agents and Multiagent Systems (AAMAS)*, page 143–151, 2023.
- [9] Siddharth Barman and Sanath Krishnamurthy. On the proximity of markets with integral equilibria. In *Proceedings of the 33rd AAAI Conference on Artificial Intelligence (AAAI)*, pages 1748–1755, 2019.
- [10] Siddharth Barman, Sanath Kumar Krishnamurthy, and Rohit Vaish. Finding fair and efficient allocations. In *Proceedings of the 19th ACM Conference on Economics and Computation (EC)*, pages 557–574, 2018.
- [11] Siddharth Barman, Sanath Kumar Krishnamurthy, and Rohit Vaish. Greedy algorithms for maximizing Nash social welfare. In *Proceedings of the 17th International Conference on Autonomous Agents and MultiAgent Systems (AAMAS)*, page 7–13, 2018.
- [12] Siddharth Barman, Debajyoti Kar, and Shraddha Pathak. Parameterized guarantees for almost envy-free allocations, 2023. URL <https://arxiv.org/abs/2312.13791>.
- [13] Umang Bhaskar, A. R. Sricharan, and Rohit Vaish. On approximate envy-freeness for indivisible chores and mixed resources. *CoRR*, abs/2012.06788, 2020. URL <https://arxiv.org/abs/2012.06788>.

- [14] Ioannis Caragiannis, David Kurokawa, Hervé Moulin, Ariel D. Procaccia, Nisarg Shah, and Junxing Wang. The unreasonable fairness of maximum Nash welfare. In *Proceedings of the 17th ACM Conference on Economics and Computation (EC)*, page 305–322, 2016.
- [15] Bhaskar Ray Chaudhury, Jugal Garg, and Kurt Mehlhorn. EFX exists for three agents. In *Proceedings of the 21st ACM Conference on Economics and Computation (EC)*, page 1–19, 2020.
- [16] Vasilis Christoforidis and Christodoulos Santorinaios. On the pursuit of efx for chores: Non-existence and approximations, 2024. URL <https://arxiv.org/abs/2406.10752>.
- [17] Richard Cole and Vasilis Gkatzelis. Approximating the Nash social welfare with indivisible items. In *Proceedings of the Forty-Seventh Annual ACM Symposium on Theory of Computing (STOC)*, page 371–380, 2015.
- [18] Richard Cole, Nikhil Devanur, Vasilis Gkatzelis, Kamal Jain, Tung Mai, Vijay V. Vazirani, and Sadra Yazdanbod. Convex program duality, Fisher markets, and Nash social welfare. In *Proceedings of the ACM Conference on Economics and Computation (EC)*, page 459–460, 2017.
- [19] Richard W. Cottle, Jong-Shi Pang, and Richard E. Stone. *The Linear Complementarity Problem*. Society for Industrial and Applied Mathematics, 2009.
- [20] Bart de Keijzer, Sylvain Bouveret, Tomas Klos, and Yingqian Zhang. On the complexity of efficiency and envy-freeness in fair division of indivisible goods with additive preferences. In Francesca Rossi and Alexis Tsoukias, editors, *Algorithmic Decision Theory*, pages 98–110, 2009.
- [21] Soroush Ebadian, Dominik Peters, and Nisarg Shah. How to fairly allocate easy and difficult chores. In *International Conference on Autonomous Agents and MultiAgent Systems (AAMAS)*, 2022.
- [22] D.K. Foley. Resource allocation and the public sector. *Yale Economic Essays*, 7(1):45–98, 1967.
- [23] Jugal Garg and Aniket Murhekar. On fair and efficient allocations of indivisible goods. In *Proceedings of the 35th AAAI Conference on Artificial Intelligence (AAAI)*, 2021.
- [24] Jugal Garg and Aniket Murhekar. Computing fair and efficient allocations with few utility values. *Theoretical Computer Science*, 962:113932, 2023.
- [25] Jugal Garg, Aniket Murhekar, and John Qin. Fair and efficient allocations of chores under bivalued preferences. *Proceedings of the 36th AAAI Conference on Artificial Intelligence (AAAI)*, pages 5043–5050, 2022.
- [26] Jugal Garg, Martin Hoefer, and Kurt Mehlhorn. Satiation in Fisher markets and approximation of Nash social welfare. *Mathematics of Operations Research*, 2023.
- [27] Jugal Garg, Aniket Murhekar, and John Qin. New algorithms for the fair and efficient allocation of indivisible chores. In *Proceedings of the Thirty-Second International Joint Conference on Artificial Intelligence (IJCAI)*, pages 2710–2718, 2023.

- [28] Jugal Garg, Aniket Murhekar, and John Qin. Weighted EF1 and PO allocations with few types of agents or chores. In *IJCAI*, 2024. URL <https://arxiv.org/abs/2402.17173>.
- [29] Yusuke Kobayashi, Ryoga Mahara, and Souta Sakamoto. EFX allocations for indivisible chores: Matching-based approach. In *Algorithmic Game Theory (SAGT)*, pages 257–270, 2023.
- [30] Euiwoong Lee. APX-hardness of maximizing Nash social welfare with indivisible items. *Information Processing Letters*, 122, 07 2015.
- [31] Bo Li, Yingkai Li, and Xiaowei Wu. Almost (weighted) proportional allocations for indivisible chores. In *Proceedings of the ACM Web Conference (WWW) 2022*, page 122–131, 2022.
- [32] Richard Lipton, Evangelos Markakis, Elchanan Mossel, and Amin Saberi. On approximately fair allocations of indivisible goods. In *In ACM Conference on Electronic Commerce (EC)*, pages 125–131, 2004.
- [33] Ryoga Mahara. Extension of additive valuations to general valuations on the existence of EFX. In *29th Annual European Symposium on Algorithms (ESA)*, 2021.
- [34] A. Mas-Colell, M.D. Whinston, and J.R. Green. *Microeconomic Theory*. Oxford University Press, 1995.
- [35] Peter McGlaughlin and Jugal Garg. Improving Nash social welfare approximations. *J. Artif. Intell. Res.*, 68:225–245, 2020.
- [36] Benjamin Plaut and Tim Roughgarden. Almost envy-freeness with general valuations. *SIAM Journal on Discrete Mathematics*, 34(2):1039–1068, 2020.
- [37] Ariel D. Procaccia. Technical perspective: An answer to fair division’s most enigmatic question. *Commun. ACM*, 63(4):118, 2020.
- [38] Shengwei Zhou and Xiaowei Wu. Approximately EFX allocations for indivisible chores. *Artif. Intell.*, 326:104037, 2024.