

When can weak latent factors be statistically inferred?

Jianqing Fan*

Yuling Yan†

Yuheng Zheng*

July 15, 2024

Abstract

This article establishes a new and comprehensive estimation and inference theory for principal component analysis (PCA) under the weak factor model that allow for cross-sectional dependent idiosyncratic components under nearly minimal the factor strength relative to the noise level or signal-to-noise ratio. Our theory is applicable regardless of the relative growth rate between the cross-sectional dimension N and temporal dimension T . This more realistic assumption and noticeable result requires completely new technical device, as the commonly-used leave-one-out trick is no longer applicable to the case with cross-sectional dependence. Another notable advancement of our theory is on PCA inference — for example, under the regime where $N \asymp T$, we show that the asymptotic normality for the PCA-based estimator holds as long as the signal-to-noise ratio (SNR) grows faster than a polynomial rate of $\log N$. This finding significantly surpasses prior work that required a polynomial rate of N . Our theory is entirely non-asymptotic, offering finite-sample characterizations for both the estimation error and the uncertainty level of statistical inference. A notable technical innovation is our closed-form first-order approximation of PCA-based estimator, which paves the way for various statistical tests. Furthermore, we apply our theories to design easy-to-implement statistics for validating whether given factors fall in the linear spans of unknown latent factors, testing structural breaks in the factor loadings for an individual unit, checking whether two units have the same risk exposures, and constructing confidence intervals for systematic risks. Our empirical studies uncover insightful correlations between our test results and economic cycles.

Keywords: factor model, principal component analysis, weak factors, cross-sectional correlation, inference, signal-to-noise ratio.

Contents

1	Introduction	2
2	Models, assumptions, and notation	4
2.1	Model setup	4
2.2	Basic assumptions	4
2.3	Notation	5
3	Main results	5
3.1	A first-order characterization of subspace perturbation errors	6
3.2	Implications: estimation guarantees and distributional characterizations	7
4	Comparison with previous work	9
5	Applications in econometrics	10
5.1	The factor specification tests	10
5.2	Test for structural breaks in betas	13
5.3	The two-sample test for betas	14
5.4	Statistical inference for the systematic risks	15

*Department of Operations Research and Financial Engineering, Princeton University, Princeton, NJ 08544, USA; Email: {jqfan, yuheng}@princeton.edu.

†Institute for Data, Systems, and Society, Massachusetts Institute of Technology, Cambridge, MA 02142, USA; Email: yulingy@mit.edu.

6	Numerical experiments	16
6.1	Monte Carlo simulations	16
6.2	Empirical studies	18
7	Other related works	21
8	Conclusions and discussions	22
A	Proof of Theorem 1: First-order approximations	23
A.1	Some useful lemmas	23
A.2	Proof of Theorem 1	36
B	Proof of Corollary 1	37
C	Inference for the factors and the factor loadings	38
C.1	Proof of Corollary 2	38
C.2	Proof of Corollary 3	41
D	Proof of Theorem 2: Factor test	41
D.1	Some useful lemmas	41
D.2	Proof of Theorem 2	49
D.3	Estimation of the idiosyncratic noise covariance matrix	52
E	Proof of Theorem 3: Test for structural breaks in betas	55
E.1	Some useful lemmas	55
E.2	Proof of Theorem 3	60
F	Proof of Theorem 4: Two-sample test for betas	62
F.1	A useful lemma	62
F.2	Proof of Theorem 4:	65
G	Proof of Theorem 5: Inference for the systematic risks	66
H	Technical lemmas	69

1 Introduction

The factor model, a pivotal tool for analyzing large panel data, has become a significant topic in finance and economics research (e.g., Chamberlain and Rothschild, 1983; Fama and French, 1993; Stock and Watson, 2002; Bai and Ng, 2002; Giglio and Xiu, 2021; Fan et al., 2021b). The estimation and inference for factor models are crucial in economic studies, particularly in areas like asset pricing and return forecasting. In the era of big data, the factor model has gained increased prominence in capturing the latent common structure for large panel data, where both the cross-sectional and temporal dimensions are ultra-high (see, e.g., recent surveys Bai and Wang, 2016; Fan et al., 2021a). Principal component analysis (PCA), known for its simplicity and effectiveness, is closely connected with the factor model and has long been a key research topic of interest in the econometric community (e.g., Stock and Watson, 2002; Bai and Ng, 2002; Bai, 2003; Onatski, 2012; Fan et al., 2013; Bai and Ng, 2013, 2023).

As pointed out by Giglio et al. (2023), most theoretical guarantees for the PCA approach to factor analysis rely on the pervasiveness assumption (e.g., Bai and Ng, 2002; Bai, 2003). This assumption requires the signal-to-noise ratio (SNR), which measures the factor strength relative to the noise level, to grow with the rate of \sqrt{N} – the square root of the cross-sectional dimension. However, many real datasets in economics do not exhibit sufficiently strong factors to meet this pervasiveness assumption. When the SNR grows slower than \sqrt{N} , the resulting model is often called the weak factor model (e.g., Onatski, 2009, 2010). Extensive research has been dedicated to the weak factor model (e.g., Onatski, 2012; Bailey et al., 2021; Freyaldenhoven, 2022; Uematsu and Yamagata, 2022a,b; Bai and Ng, 2023; Jiang et al., 2023; Choi and Yuan, 2024), among which

the PC estimators (the estimators via the PCA approach) have been a primary subject. Recently, [Bai and Ng \(2023\)](#); [Jiang et al. \(2023\)](#); [Choi and Yuan \(2024\)](#) studied the consistency and asymptotic normality of the PC estimators for factors and factor loadings in the weak factor model. In the extreme case (also called the super-weak factor model) where the SNR is $O(1)$, [Onatski \(2012\)](#) showed that the PC estimators are inconsistent.

This paper establishes a novel and comprehensive theory for PCA in the weak factor model. Our theory is non-asymptotic and can be easily translated to asymptotic results. The conditions we propose for the asymptotic normality of PC estimators are optimal in the sense that, surprisingly different from the existing literature, the required growth rate of the SNR for consistency aligns with that for asymptotic normality, differing only by a logarithmic factor. In particular, in the regime $N \asymp T$, where T is the temporal dimension, we prove that asymptotic normality holds as long as the SNR grows faster than a polynomial rate of $\log N$, and this result is a substantial advance compared with the existing results that require the SNR to grow with a polynomial rate of N (e.g., [Bai and Ng, 2023](#); [Jiang et al., 2023](#); [Choi and Yuan, 2024](#)).

The most innovative part of our theory lies in establishing a closed-form, first-order approximation of the PC estimator, that is, we decompose the PC estimator into three components — the ground truth, a first-order term, and the higher-order negligible term. We express the first-order term explicitly using the parameters in the factor model. This closed-form characterization paves the way for us to establish the asymptotic normality and design various test statistics for practical applications. Our theory is based on novel applications of the leave-one-out analysis and matrix concentration inequalities. Moreover, our findings provide valuable insights and practical implications for a range of econometric problems related with PCA, e.g., macroeconomic forecasting based on factor-augmented regressions (e.g., [Bai and Ng, 2006a](#)).

We demonstrate the practical applications of our theories using both synthetic and real datasets. First, we design an easy-to-implement statistic for factor specification test (e.g., [Bai and Ng, 2006b](#)) — whether an observed factor is in the linear space spanned by the latent common factors or not. A key innovation of our approach is the ability to conduct this test in any flexible subset of the whole period, owing to the row-wise error bound in our theory. We utilize the monthly return data of the S&P 500 constituents from 1995 to 2024, along with the data of the Fama-French three factors, then run the factor test in a rolling window manner (e.g., [Fan et al., 2015](#)). Our results uncover a notable decline in the importance and explanatory power of the size factor during the 2008 financial crisis, and of both the size and value factors during the COVID-19 pandemic around 2019. These findings are supported by our test results, which reject the null hypothesis that these factors are in the linear space of latent factors.

Then, we design a test statistic for the structural break of betas (e.g., [Stock and Watson, 2009](#); [Breitung and Eickmeier, 2011](#)), and apply it to the aforementioned S&P 500 constituents data. Our novelty is that our test statistic works in the weak factor model without the pervasiveness assumption, which is required in prior work. We test for each stock to determine if the beta has changed before and after the three recessions covered by our data — the Early 2000s Recession, the 2008 Great Recession, and the COVID-19 Recession. We find that in each recession, the sectors most affected by the shocks, where many stocks exhibited structural breaks in betas, correspond reasonably to the causes of these economic recessions. For example, during the 2008 Great Recession, marked by the subprime mortgage crisis, the financial sector experienced a strong impact, which aligns with its exposure to mortgage-backed securities and other related financial instruments. During the COVID-19 Recession, the Health Care and Real Estate sectors were significantly impacted, reflecting the uncertainties brought about by lockdowns and health crises due to the pandemic. Additionally, we develop statistical tests for the betas to evaluate the similarity in risk exposure between two stocks and construct valid confidence interval for the systematic risk of each stock.

The rest of the paper is organized as follows. Section 2 introduces the model setup, basic assumptions, and notation. In Section 3, we show our main results on the first-order approximations for the PC estimators, and present the asymptotic normality results as corollaries. Section 4 provides a detailed comparison of our results with related work. In Section 5, we showcase four applications in econometrics based on our main results. Section 6 collects the numerical results in both simulated data and real data. More related works are discussed in Section 7, and the paper concludes with a discussion on future directions in Section 8.

2 Models, assumptions, and notation

2.1 Model setup

Let N be the number of cross-sectional units and T be the number of observations. Consider the factor model for a panel of data $\{x_{i,t}\}$:

$$x_{i,t} = \mathbf{b}_i^\top \mathbf{f}_t + \varepsilon_{i,t}, \quad 1 \leq i \leq N, 1 \leq t \leq T, \quad (2.1)$$

where $\mathbf{f}_t = (f_{1,t}, f_{2,t}, \dots, f_{r,t})^\top$ is the latent factor, r is the number of factors, $\mathbf{b}_i = (b_{i,1}, b_{i,2}, \dots, b_{i,r})^\top$ is a vector of factor loadings, and $\varepsilon_{i,t}$ represents the idiosyncratic noise. Viewing $x_{i,t}$ as the excess return of the i -th asset at time t , the model (2.1) is intimately linked with the multi-factor pricing model. This model originates from the Arbitrage Pricing Theory (APT) developed by Ross (1976) and finds extensive applications in finance.

To compact the notation, we denote by $\mathbf{B} = (\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_N)^\top$ the $N \times r$ factor loading matrix. Let $\mathbf{x}_t = (x_{1,t}, x_{2,t}, \dots, x_{N,t})^\top$ and $\mathbf{e}_t = (\varepsilon_{1,t}, \varepsilon_{2,t}, \dots, \varepsilon_{N,t})^\top$. Then, the factor model (2.1) can be expressed as $\mathbf{x}_t = \mathbf{B}\mathbf{f}_t + \mathbf{e}_t$, or written in a matrix form as follows

$$\mathbf{X} = \mathbf{B}\mathbf{F}^\top + \mathbf{E}, \quad (2.2)$$

where $\mathbf{X} = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_T)$, $\mathbf{F} = (\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_T)^\top$, and $\mathbf{E} = (\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_T)$ are the $N \times T$ panel data, the $T \times r$ factor realizations, and the $N \times T$ idiosyncratic noise matrix, respectively.

In our setup, the only observable part is the panel data \mathbf{X} . We are interested in the estimation and inference for both the latent factors and factor loadings via the PCA approach.

2.2 Basic assumptions

We propose some basic assumptions as follows. Note that the factor model has the rotation (indeed affine transform) ambiguity (e.g., Bai and Ng, 2013), that is, the factor loadings and latent factors are not identifiable since $\mathbf{B}\mathbf{F}^\top = (\mathbf{B}\mathbf{H}^{-1})(\mathbf{H}\mathbf{F}^\top)^\top$ holds for any invertible matrix \mathbf{H} . Without loss of generality, we assume that the columns of \mathbf{B} are orthogonal and the covariance of \mathbf{f}_t is the identity matrix, as stated in Assumption 1 below.

Assumption 1. For $t = 1, 2, \dots, T$, the factor \mathbf{f}_t has mean zero and the identity covariance matrix. The factor loading matrix \mathbf{B} has orthogonal columns:

$$\mathbf{B}^\top \mathbf{B} = \mathbf{\Sigma}^2 \quad \text{with} \quad \mathbf{\Sigma} = \text{diag}(\sigma_1, \sigma_2, \dots, \sigma_r), \quad (2.3)$$

where $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_r > 0$.

Assumption 1 is a standard identification condition for the factor model (e.g., Fan et al., 2013). The singular values $\{\sigma_i\}$ of the factor loading matrix \mathbf{B} characterize the strengths of the latent factors.

Next, to accommodate the cross-sectional correlation in the noise and to facilitate leave-one-out analysis in our technical proof, we propose Assumption 2 for the noise matrix \mathbf{E} , specifying its structure and the distribution of its entries.

Assumption 2. The idiosyncratic noise matrix \mathbf{E} is given by

$$\mathbf{E} = \mathbf{\Sigma}_\varepsilon^{1/2} \mathbf{Z}. \quad (2.4)$$

Here, $\mathbf{\Sigma}_\varepsilon$ is a $N \times N$ positive definite matrix and $\mathbf{\Sigma}_\varepsilon^{1/2}$ is the symmetric square root of $\mathbf{\Sigma}_\varepsilon$; $\mathbf{Z} = (Z_{i,t})_{i=1}^N_{t=1}^T$ is a $N \times T$ matrix; The entries of \mathbf{Z} are independent sub-Gaussian random variables that satisfy

$$\mathbb{E}[Z_{i,t}] = 0, \quad \mathbb{E}[Z_{i,t}^2] = 1, \quad \|Z_{i,t}\|_{\psi_2} = O(1),$$

for $1 \leq i \leq N, 1 \leq t \leq T$, where $\|\cdot\|_{\psi_2}$ is the sub-Gaussian norm (see Definition 2.5.6 in Vershynin, 2017).

The nonzero off-diagonal entries of Σ_ε characterize the cross-sectional correlations in the idiosyncratic noise $\mathbf{e}_t = (\varepsilon_{1,t}, \varepsilon_{2,t}, \dots, \varepsilon_{N,t})^\top$. Though the noise terms $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_T$ are independent under Assumption 2, our theory could potentially be generalized to the cases where the temporal correlations are present in the noise matrix \mathbf{E} , and this generalization is an interesting future direction. The formulation (2.4) imposes additional structural constraints on the noise matrix. However, such assumptions are commonplace in the study of the weak factor model; see similar assumptions in Onatski (2010, 2012).

Then, to establish the non-asymptotic results via the concentration inequalities, we propose the following assumption on the distribution of the factors.

Assumption 3. *The factor realization $\mathbf{F} = (\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_T)^\top$ is independent with the noise matrix \mathbf{E} ; The factors $\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_T$ are independent and sub-Gaussian random vectors in \mathbb{R}^r satisfying that*

$$\|\mathbf{f}_t\|_{\psi_2} = O(1) \quad \text{for} \quad 1 \leq t \leq T.$$

The conditions in Assumption 3 are slightly stronger than those in the literature, e.g., Bai and Ng (2023), but remain standard and simplify our technical proofs.

2.3 Notation

We introduce some notation that will be used throughout the paper. For two sequences a_n and b_n , we write $a_n \lesssim b_n$ (or equivalently, $b_n \gtrsim a_n$) if $a_n = O(b_n)$, i.e., there exist a constant $C > 0$ and an integer $N > 0$ such that, $a_n \leq Cb_n$ holds for any $n > N$; write $a_n \asymp b_n$ if both $a_n \lesssim b_n$ and $a_n \gtrsim b_n$ hold and $a_n \ll b_n$ (or equivalently, $b_n \gg a_n$) if $a_n = o(b_n)$.

For a symmetric matrix \mathbf{M} , we denote by $\lambda_{\min}(\mathbf{M})$ and $\lambda_{\max}(\mathbf{M})$ its minimum and maximum eigenvalues. For a vector \mathbf{v} , we denote $\|\mathbf{v}\|_2$, $\|\mathbf{v}\|_1$, and $\|\mathbf{v}\|_\infty$ as the ℓ_2 -norm, ℓ_1 -norm, and supremum norm, respectively. Consider any matrix $\mathbf{A} \in \mathbb{R}^{m \times n}$. We denote by $\mathbf{A}_{i,\cdot}$ and $\mathbf{A}_{\cdot,j}$ the i -th row and the j -th column of \mathbf{A} . We let $\|\mathbf{A}\|_2$, $\|\mathbf{A}\|_F$, and $\|\mathbf{A}\|_{2,\infty}$ denote the spectral norm, the Frobenius norm, and the $\ell_{2,\infty}$ -norm (i.e., $\|\mathbf{A}\|_{2,\infty} := \sup_{1 \leq i \leq m} \|\mathbf{A}_{i,\cdot}\|_2$), respectively. For an index set $S \subseteq \{1, 2, \dots, m\}$ (resp. $S \subseteq \{1, 2, \dots, n\}$), we use $|S|$ to denote its cardinality, and use \mathbf{A}_S (resp. $\mathbf{A}_{\cdot,S}$) to denote a submatrix of \mathbf{A} whose rows (resp. columns) are indexed by S . We let $\text{col}(\mathbf{A})$, \mathbf{A}^+ , and $\mathbf{P}_\mathbf{A} := \mathbf{A}\mathbf{A}^+$ denote the column subspace of \mathbf{A} , the generalized inverse of \mathbf{A} , and the projection matrix onto the column space of \mathbf{A} .

We denote by $\text{diag}(a_1, a_2, \dots, a_r)$ the $r \times r$ diagonal matrix whose diagonal entries are given by a_1, a_2, \dots, a_r . Let \mathbf{I}_r be the $r \times r$ identity matrix. We denote by $\mathcal{O}^{r \times r}$ the set of all $r \times r$ orthonormal (or rotation) matrices. For a non-singular $n \times n$ matrix \mathbf{H} with SVD $\mathbf{U}_H \Sigma_H \mathbf{V}_H^\top$, we denote by $\text{sgn}(\mathbf{H})$ the following orthogonal matrix $\text{sgn}(\mathbf{H}) := \mathbf{U}_H \mathbf{V}_H^\top$. Then we have that, for any two matrices $\hat{\mathbf{U}}, \mathbf{U} \in \mathbb{R}^{n \times r}$ with $r \leq n$, among all rotation matrices, the one that best aligns $\hat{\mathbf{U}}$ and \mathbf{U} is precisely $\text{sgn}(\hat{\mathbf{U}}^\top \mathbf{U})$ (see, e.g., Appendix D.2.1 in Ma et al., 2020), namely,

$$\text{sgn}(\hat{\mathbf{U}}^\top \mathbf{U}) = \arg \min_{\mathbf{O} \in \mathcal{O}^{r \times r}} \|\hat{\mathbf{U}}\mathbf{O} - \mathbf{U}\|_F.$$

3 Main results

In this section, we demonstrate the desirable statistical performance of PCA in weak factor models. We will first present a master result (cf. Theorem 1) on subspace error decomposition. Based on this key result, we derive non-asymptotic distributional characterization for our estimators for the factors and factor loadings, paving the way to data-driven statistical inference.

First, we formally introduce the PC estimators of the factor and factor loadings in Algorithm 1.

Algorithm 1 The PCA-based method.

Input: panel data \mathbf{X} , rank r .

Compute the truncated rank- r SVD $\hat{\mathbf{U}}\hat{\Sigma}\hat{\mathbf{V}}^\top$ of $T^{-1/2}\mathbf{X}$, where $\hat{\mathbf{U}} \in \mathbb{R}^{N \times r}$ and $\hat{\mathbf{V}} \in \mathbb{R}^{T \times r}$ have orthonormal columns, and $\hat{\Sigma} = \text{diag}(\hat{\sigma}_1, \hat{\sigma}_2, \dots, \hat{\sigma}_r) \in \mathbb{R}^{r \times r}$ satisfies that $\hat{\sigma}_1 \geq \hat{\sigma}_2 \geq \dots \geq \hat{\sigma}_r$.

Output $\hat{\mathbf{F}} := T^{1/2}\hat{\mathbf{V}}$ as the estimator of factors \mathbf{F} , and $\hat{\mathbf{B}} := \hat{\mathbf{U}}\hat{\Sigma}$ as the estimator of factor loadings \mathbf{B} .

Our master result focuses on the subspace estimates $\widehat{\mathbf{U}}$ and $\widehat{\mathbf{V}}$. We define some relevant quantities first. Denote by $\mathbf{U}\mathbf{\Lambda}\mathbf{V}^\top$ the SVD of $T^{-1/2}\mathbf{B}\mathbf{F}^\top$, where both $\mathbf{U} \in \mathbb{R}^{N \times r}$ and $\mathbf{V} \in \mathbb{R}^{T \times r}$ have orthonormal columns, and $\mathbf{\Lambda} = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_r) \in \mathbb{R}^{r \times r}$ satisfies that $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_r \geq 0$. We define $n := \max(N, T)$.

3.1 A first-order characterization of subspace perturbation errors

Before presenting our master results, we list two key assumptions on the SNR. To characterize the SNR, we define

$$\theta := \sigma_r / \sqrt{\|\boldsymbol{\Sigma}_\varepsilon\|_2}, \quad \vartheta_k := \sigma_r / \|(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{k,\cdot}\|_1, \quad \text{and} \quad \vartheta := \sigma_r / \|\boldsymbol{\Sigma}_\varepsilon^{1/2}\|_1.$$

Here, the signal strength, which is the factor strength in our case, is effectively represented by σ_r , the smallest singular value of the factor loading matrix \mathbf{B} . The noise level is captured by the norm of the covariance matrix $\boldsymbol{\Sigma}_\varepsilon$ for the idiosyncratic noise $\mathbf{E} = \boldsymbol{\Sigma}_\varepsilon^{1/2}\mathbf{Z}$ (cf. (2.4)). The SNRs θ , ϑ_k , and ϑ are similar, though their denominators adopt different norms of $\boldsymbol{\Sigma}_\varepsilon$. The inclusion of ℓ_1 -norms $\|(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{k,\cdot}\|_1$ and $\|\boldsymbol{\Sigma}_\varepsilon^{1/2}\|_1 = \max_{1 \leq k \leq N} \|(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{k,\cdot}\|_1$, though less common than the spectral norm $\sqrt{\|\boldsymbol{\Sigma}_\varepsilon\|_2} = \|\boldsymbol{\Sigma}_\varepsilon^{1/2}\|_2$, is that our main results are on the row-wise error bound and we need these norms in the matrix inequalities for technical reasons. Then, two crucial assumptions on SNR are as follows.

Assumption 4. *There exists a sufficiently large constant $C_1 > 0$ such that*

$$\vartheta \equiv \frac{\sigma_r}{\|\boldsymbol{\Sigma}_\varepsilon^{1/2}\|_1} \geq C_1 \sqrt{\frac{n}{T} \log n}. \quad (3.1)$$

Assumption 5. *There exists a sufficiently large constant $C_2 > 0$ such that*

$$\theta \equiv \frac{\sigma_r}{\|\boldsymbol{\Sigma}_\varepsilon^{1/2}\|_2} \geq C_2 \sqrt{\frac{n}{T} \log n}. \quad (3.2)$$

Assumption 4 implies Assumption 5 due to the elementary inequality $\|\boldsymbol{\Sigma}_\varepsilon^{1/2}\|_2 \leq \|\boldsymbol{\Sigma}_\varepsilon^{1/2}\|_1$. On the other hand, $\|\boldsymbol{\Sigma}_\varepsilon^{1/2}\|_1 \leq s\|\boldsymbol{\Sigma}_\varepsilon^{1/2}\|_2$, where s is the maximum number of nonzero elements of $\boldsymbol{\Sigma}_\varepsilon^{1/2}$ in each row, which is small when $\boldsymbol{\Sigma}_\varepsilon$ is sparse. In particular, when the noise covariance matrix $\boldsymbol{\Sigma}_\varepsilon$ is diagonal (i.e., no cross-sectional correlations, also known as the strict factor model (Ross, 1976; Fan et al., 2008; Bai and Shi, 2011)), we have that $\|\boldsymbol{\Sigma}_\varepsilon^{1/2}\|_2 = \|\boldsymbol{\Sigma}_\varepsilon^{1/2}\|_1$. In this case, Assumption 4 is equivalent with Assumption 5. Throughout the paper, we denote by $C_0, C_1, C_2, C_U, C_V, c_0$, etc. the generic constants that may vary from place to place. We are now ready to present our main results as follows.

Theorem 1. *Assume that $T \geq C_0(r + \log n)$ and $n \geq C_0 r \log n$ for some sufficiently large constant $C_0 > 0$. Consider the first-order expansions*

$$\widehat{\mathbf{U}}\mathbf{R}_U - \mathbf{U} = \mathbf{G}_U + \boldsymbol{\Psi}_U \quad \text{with} \quad \mathbf{G}_U := T^{-1/2}\mathbf{E}\mathbf{V}\mathbf{\Lambda}^{-1}, \quad (3.3a)$$

$$\widehat{\mathbf{V}}\mathbf{R}_V - \mathbf{V} = \mathbf{G}_V + \boldsymbol{\Psi}_V \quad \text{with} \quad \mathbf{G}_V := T^{-1/2}\mathbf{E}^\top\mathbf{U}\mathbf{\Lambda}^{-1}, \quad (3.3b)$$

where $\mathbf{R}_U := \text{sgn}(\widehat{\mathbf{U}}^\top\mathbf{U})$ and $\mathbf{R}_V := \text{sgn}(\widehat{\mathbf{V}}^\top\mathbf{V})$ are two global rotation matrices. Under Assumptions 1, 2, and 3, we have, with probability at least $1 - O(n^{-2})$, the remainder terms $\boldsymbol{\Psi}_U$ and $\boldsymbol{\Psi}_V$ are higher-order negligible terms that satisfy the following bounds:

(i) Under Assumption 4, there exists some universal constant $C_U > 0$ such that, uniformly for all $k = 1, 2, \dots, N$,

$$\|(\boldsymbol{\Psi}_U)_{k,\cdot}\|_2 \leq C_U \frac{\sqrt{n}}{\vartheta_k \vartheta T} \sqrt{r} \log^{3/2} n + C_U \left(\frac{n}{\theta^2 T} + \frac{1}{\theta \sqrt{T}} \sqrt{r} \log n \right) \|\mathbf{U}_{k,\cdot}\|_2 + C_U \frac{n}{\vartheta_k \vartheta T} \log n \|\mathbf{U}\|_{2,\infty}.$$

(ii) Under Assumption 5, there exists some universal constant $C_V > 0$ such that, uniformly for all $l = 1, 2, \dots, T$,

$$\|(\boldsymbol{\Psi}_V)_{l,\cdot}\|_2 \leq C_V \frac{\sqrt{n}}{\theta^2 T} \sqrt{r} \log^{3/2} n + C_V \left(\frac{n}{\theta^2 T} + \frac{1}{\theta \sqrt{T}} \right) \sqrt{r} \log n \|\mathbf{V}_{l,\cdot}\|_2.$$

The perturbation bounds in Theorem 1 pave the way for the statistical inference of the factor loadings and factors. The reasons why the quantities presented in Theorem 1 relate to the factor loadings \mathbf{B} and factors \mathbf{F} are that, as demonstrated in Lemma 2, the column subspaces of \mathbf{U} and \mathbf{B} are identical, as are those of \mathbf{V} and \mathbf{F} .

To show how strong our SNR condition is, we examine scenarios where there is no correlation in the noise matrix \mathbf{E} , i.e., Σ_ε is diagonal as in the strict factor model. In this case, the SNRs satisfy $\vartheta = \theta = \sigma_r/\sigma_\varepsilon$, where $\sigma_\varepsilon = \|\Sigma_\varepsilon^{1/2}\|_2 = \|\Sigma_\varepsilon^{1/2}\|_1$. Then Assumptions 4 and 5 match the minimal condition $\theta \gg \sqrt{n/T}$ required for the consistent estimations of the left and right singular subspaces (e.g., Yan et al., 2024), up to log factors.

To explain that why Ψ_U and Ψ_V are higher-order negligible terms, we compare their bounds with the magnitude of the first-order terms \mathbf{G}_U and \mathbf{G}_V . For simplicity, we consider the setting where $r \asymp 1$, $N \asymp T$, Σ_ε is well-conditioned (i.e., $\kappa_\varepsilon := \lambda_{\max}(\Sigma_\varepsilon)/\lambda_{\min}(\Sigma_\varepsilon) \asymp 1$), and \mathbf{U} is incoherent (see, e.g., Definition 3.1 in Chen et al., 2021a); the matrix \mathbf{V} can be proven to be incoherent since $\|\mathbf{V}\|_{2,\infty} \lesssim T^{-1/2} \log^{1/2} n$. Then, the typical size of the k -th row of the first-order term \mathbf{G}_U can be measured by $\mathbb{E}^{1/2}[\|(\mathbf{G}_U)_{k,\cdot}\|_2^2]$. Standard computations yield $\mathbb{E}^{1/2}[\|(\mathbf{G}_U)_{k,\cdot}\|_2^2] \geq T^{-1/2} \sigma_r^{-1} \sqrt{\lambda_{\min}(\Sigma_\varepsilon)}$. Similarly, for the first-order term \mathbf{G}_V , we have that $\mathbb{E}^{1/2}[\|(\mathbf{G}_V)_{l,\cdot}\|_2^2] \geq T^{-1/2} \sigma_r^{-1} \sqrt{\lambda_{\min}(\Sigma_\varepsilon)}$. Then, using the row-wise bounds for Ψ_U and Ψ_V in Theorem 1, we obtain that

$$(\mathbb{E}^{1/2}[\|(\mathbf{G}_U)_{k,\cdot}\|_2^2])^{-1} \|(\Psi_U)_{k,\cdot}\|_2 \lesssim \left(\frac{\theta}{\vartheta_k \vartheta} + \frac{1}{\theta}\right) \log^{3/2} n + \frac{1}{\sqrt{N}} \log n \ll 1, \quad (3.4a)$$

and

$$(\mathbb{E}^{1/2}[\|(\mathbf{G}_V)_{l,\cdot}\|_2^2])^{-1} \|(\Psi_V)_{l,\cdot}\|_2 \lesssim \left(\frac{1}{\theta} + \frac{1}{\sqrt{T}}\right) \log^{3/2} n \ll 1. \quad (3.4b)$$

The last inequality in (3.4a) (resp. (3.4b)) implies that the first-order term $(\mathbf{G}_U)_{k,\cdot}$ (resp. $(\mathbf{G}_V)_{l,\cdot}$) dominates the higher-order term $(\Psi_U)_{k,\cdot}$ (resp. $(\Psi_V)_{l,\cdot}$), provided that the SNRs grow at polynomial rate of $\log n$: $\theta^{-1} \vartheta_k \vartheta \gg \log^{3/2} n$ and $\theta \gg \log^{3/2} n$ (resp. $\theta \gg \log^{3/2} n$). These conditions are less stringent than the assumptions in the prior work (e.g., Bai and Ng, 2023; Jiang et al., 2023; Choi and Yuan, 2024) that required the SNR to grow with a polynomial rate of n .

As we will show later, Theorem 1 is the foundation to design test statistics and conduct statistical inference for the factors, factor loadings, and other parameters of interest. To the best of our knowledge, the non-asymptotic first-order approximations (3.3a)–(3.3b) are new in the statistics and econometrics literature. While similar findings exist in the fields of low-rank matrix completion and spectral methods, the closest result to ours is Theorem 9 of Yan et al. (2024). However, they assumed that the entries of the noise matrix \mathbf{E} are independent, while we allow the presence of cross-sectional correlations in the noise matrix \mathbf{E} .

Our results in Theorem 1 are nontrivial generalizations of those in Yan et al. (2024). We note that their proof of row-wise error bounds via leave-one-out (LOO) technique requires constructing an auxiliary matrix that is independent with the target row. This construction is obtained by zeroing the row with the same position in noise matrix. However, this approach does not work in our case because it is still correlated with the target row due to the cross-sectional correlation among all rows. We overcome this challenge through applications of matrix concentration inequalities, and our results do not need any additional structural assumption (e.g., sparsity) on the noise covariance matrix Σ_ε . An additional advance of our theory compared with Yan et al. (2024) lies in that, our bounds for $\|(\Psi_U)_{k,\cdot}\|_2$ and $\|(\Psi_V)_{l,\cdot}\|_2$ are free of condition number $\kappa := \sigma_1/\sigma_r$, making our theory work even when the factor loading matrix \mathbf{B} is near-singular and ill-conditioned, i.e., κ is very large. In particular, our result accommodates the heterogeneous case of Bai and Ng (2023).

3.2 Implications: estimation guarantees and distributional characterizations

In this section, we present the immediate consequences of Theorem 1 — estimation error bounds and distributional characterization for the PC estimators. Later in Section 4, we will compare these results with the recent work (Bai and Ng (2023); Jiang et al. (2023); Choi and Yuan (2024)), and highlight the advantages of our theory.

Recall that our estimators for the factors \mathbf{F} and factor loadings \mathbf{B} in Algorithm 1 are given by $\widehat{\mathbf{F}} = \sqrt{T} \widehat{\mathbf{V}}$ and $\widehat{\mathbf{B}} = \widehat{\mathbf{U}} \widehat{\Sigma}$ respectively. Let us take a moment to see how to quantify the estimation error for $\widehat{\mathbf{F}}$ in the

face of rotational ambiguity. In view of Theorem 1, we know that $\widehat{\mathbf{F}}\mathbf{R}_V$ should be close to $\sqrt{T}\mathbf{V}$ in Euclidean distance. In addition, recall that $\mathbf{U}\mathbf{\Lambda}\mathbf{V}^\top$ is the SVD of the rank- r matrix $T^{-1/2}\mathbf{B}\mathbf{F}^\top$, which suggests the existence of an invertible, $\sigma(\mathbf{F})$ -measurable matrix $\mathbf{J} \in \mathbb{R}^{r \times r}$ such that $\mathbf{V} = T^{-1/2}\mathbf{F}\mathbf{J}$. Hence, by defining $\mathbf{R}_F := \mathbf{J}\mathbf{R}_V^\top$, we may use $\widehat{\mathbf{F}} - \mathbf{F}\mathbf{R}_F$ to evaluate the estimation error of $\widehat{\mathbf{F}}$. Similarly, we define $\mathbf{R}_B := (\mathbf{R}_F^{-1})^\top$ and use $\widehat{\mathbf{B}} - \mathbf{B}\mathbf{R}_B$ to measure the estimation error of $\widehat{\mathbf{B}}$.

We will show in Lemma 2 that \mathbf{R}_F and \mathbf{R}_B are close to rotation matrices. Indeed, in the study of PC estimators for factor model (e.g., Bai and Ng (2023); Jiang et al. (2023); Choi and Yuan (2024)), it is quite common to take $\mathbf{F}\mathbf{R}_F$ and $\mathbf{B}\mathbf{R}_B$, rather than \mathbf{F} and \mathbf{B} , as the groundtruth, though the specific choices of \mathbf{R}_F and \mathbf{R}_B vary across different works. Both the averaged and the row-wise estimation error bound can be deduced from Theorem 1. We record the results in the following corollary.

Corollary 1 (Estimation guarantees for factors and factor loadings). *Suppose that the assumptions in Theorem 1 hold. Then there exists a universal constant $C_0 > 0$ such that: with probability at least $1 - O(n^{-2})$, for factors, the averaged and the row-wise estimation error are given by*

$$\frac{1}{T} \|\widehat{\mathbf{F}} - \mathbf{F}\mathbf{R}_F\|_F^2 \leq C_0 \frac{1}{\theta^2} \frac{n}{T} r \quad \text{and} \quad \|(\widehat{\mathbf{F}} - \mathbf{F}\mathbf{R}_F)_{t,\cdot}\|_2 \leq C_0 \frac{1}{\theta} \sqrt{r} \log n, \forall t, \quad (3.5)$$

respectively; for factor loadings, we define $\bar{\mathbf{U}} := \mathbf{B}\mathbf{\Sigma}^{-1} \in \mathbb{R}^{N \times r}$, if there exists some universal constant $C_U > 0$ such that $\|\bar{\mathbf{U}}\|_{2,\infty} \leq C_U \theta / \vartheta$, $\|\bar{\mathbf{U}}\|_{2,\infty} \leq C_U \sqrt{T} \vartheta / n$ and $\|\bar{\mathbf{U}}\|_{2,\infty} \leq C_U \sqrt{T} \theta^2 / (\vartheta n)$, then the averaged estimation error bound is given by

$$\frac{1}{N} \|\widehat{\mathbf{B}} - \mathbf{B}\mathbf{R}_B\|_F^2 \leq C_0 \left(\frac{1}{T} \|\mathbf{\Sigma}_\varepsilon^{1/2}\|_1^2 + \left(\frac{n^2}{\theta^4 T^2} + \frac{1}{\theta^2 T} \right) \sigma_r^2 \right) r^3 \log^4 n, \quad (3.6a)$$

and the row-wise estimation error bound is given by

$$\|(\widehat{\mathbf{B}} - \mathbf{B}\mathbf{R}_B)_{i,\cdot}\|_2 \leq C_0 \left(\frac{1}{\sqrt{T}} \|(\mathbf{\Sigma}_\varepsilon^{1/2})_{i,\cdot}\|_1 + \left(\frac{n}{\theta^2 T} + \frac{1}{\theta \sqrt{T}} \right) \sigma_r \|\bar{\mathbf{U}}_{i,\cdot}\|_2 \right) r \log^2 n. \quad (3.6b)$$

Let us interpret the above bounds under some specific settings. For simplicity, we ignore the log factors $\log n$ and consider the setting discussed after Theorem 1, with the assumption that $\|\mathbf{\Sigma}_\varepsilon^{1/2}\|_1 \asymp \|\mathbf{\Sigma}_\varepsilon^{1/2}\|_2 \asymp 1$. Under this setting, to make the assumptions on $\|\bar{\mathbf{U}}\|_{2,\infty}$ hold, it suffices to assume that $\theta \gtrsim 1$, where $\theta \equiv \sigma_r / \|\mathbf{\Sigma}_\varepsilon^{1/2}\|_2$ is the SNR. For factors, the upper bounds for the averaged and row-wise estimation error rates in (3.5) are given by θ^{-2} and θ^{-1} , respectively; for factor loadings, the two bounds in (3.6a)–(3.6b) are given by $T^{-1} + N^{-1}\theta^{-2}$ and $T^{-1/2} + N^{-1/2}\theta^{-1}$, respectively. All these bounds go to zero under the condition that $\theta \gg 1$, which match the minimal condition required for the consistent estimations of the left and right singular subspaces if there is no correlation in the noise matrix (e.g., Yan et al., 2024).

Next, we present our results on the inference for factors and factor loadings in Corollaries 2 and 3, respectively.

Corollary 2 (Distributional theory for factors). *Suppose that Assumptions 1, 2, and 3 hold. For any given target error level $\delta > 0$, assume that $T \geq C_1(r + \log n)$, $n \geq C_1 r \log n$, $T \geq C_0 \kappa_\varepsilon \delta^{-2} r^2 \log^4 n$, $n \geq C_0 \delta^{-1/2}$,*

$$\theta \geq C_0 \delta^{-1} \sqrt{\kappa_\varepsilon} \frac{n}{T} r \log^{3/2} n, \quad \text{and} \quad \frac{\|\mathbf{\Sigma}_\varepsilon^{1/2} \bar{\mathbf{U}}\|_{2,\infty}}{\|\mathbf{\Sigma}_\varepsilon^{1/2}\|_2} \leq c_0 \delta \kappa_\varepsilon^{-1/2} r^{-5/4},$$

hold for some universal constant $c_0, C_0 > 0$ and sufficiently large constant $C_1 > 0$. Then, for any $t = 1, 2, \dots, T$, it holds that

$$\sup_{\mathcal{C} \in \mathcal{C}^r} \left| \mathbb{P}((\widehat{\mathbf{F}} - \mathbf{F}\mathbf{R}_F)_{t,\cdot} \in \mathcal{C}) - \mathbb{P}(\mathcal{N}(0, \mathbf{\Sigma}_{F,t}) \in \mathcal{C}) \right| \leq \delta, \quad (3.7)$$

where \mathcal{C}^r is the collection of all convex sets in \mathbb{R}^r , and the covariance matrix $\mathbf{\Sigma}_{F,t}$ is given by $\mathbf{\Sigma}_{F,t} = \mathbf{R}_V \mathbf{\Lambda}^{-1} \mathbf{U}^\top \mathbf{\Sigma}_\varepsilon \mathbf{U} \mathbf{\Lambda}^{-1} \mathbf{R}_V^\top$.

Remark 1. If all the entries of the matrix \mathbf{Z} are Gaussian, i.e., the noise is Gaussian, then the result (3.7) holds without the assumption that $\|\mathbf{\Sigma}_\varepsilon^{1/2} \bar{\mathbf{U}}\|_{2,\infty} / \|\mathbf{\Sigma}_\varepsilon^{1/2}\|_2 \leq c_0 \kappa_\varepsilon^{-1/2} r^{-5/4} \delta$. Indeed, in the scenario where $\|\mathbf{\Sigma}_\varepsilon^{1/2}\|_1 / \|\mathbf{\Sigma}_\varepsilon^{1/2}\|_2 \asymp 1$ and $\bar{\mathbf{U}}$ is μ -incoherent, fulfilling our assumption on $\|\mathbf{\Sigma}_\varepsilon^{1/2} \bar{\mathbf{U}}\|_{2,\infty} / \|\mathbf{\Sigma}_\varepsilon^{1/2}\|_2$ merely requires that $N \geq \mu \kappa_\varepsilon r^{5/2} c_0^{-2} \delta^{-2}$ on the growth rate of cross-sectional dimension N .

Corollary 3 (Distributional theory for factor loadings). *Suppose that Assumptions 1, 2, and 3 hold. Assume that there exists a constant $c_\varepsilon > 0$ such that $\lambda_{\min}(\mathbf{\Sigma}_\varepsilon) > c_\varepsilon$. For any given target error level $\delta > 0$, assume that $T \geq C_1(r + \log n)$, $n \geq C_1 r \log n$, $T \geq C_0 \delta^{-2}(r^2 \log n + \kappa_\varepsilon r \log^4 n)$,*

$$\vartheta_i \geq C_0 \delta^{-1} \frac{1}{\sqrt{c_\varepsilon}} \|\mathbf{\Sigma}_\varepsilon^{1/2}\|_1 \frac{n}{T} \sqrt{r} \log^{3/2} n, \quad \text{and} \quad \theta \geq C_0 \delta^{-1} \sqrt{\kappa_\varepsilon} \frac{n}{T} \sqrt{r} \log n,$$

for some universal constant $C_0 > 0$ and sufficiently large constant $C_1 > 0$. Then, for any $i = 1, 2, \dots, N$, it holds that

$$\sup_{\mathcal{C} \in \mathcal{C}^r} \left| \mathbb{P}((\widehat{\mathbf{B}} - \mathbf{B}\mathbf{R}_B)_{i,\cdot} \in \mathcal{C}) - \mathbb{P}(\mathcal{N}(0, \mathbf{\Sigma}_{B,i}) \in \mathcal{C}) \right| \leq \delta,$$

where \mathcal{C}^r is the collection of all convex sets in \mathbb{R}^r , and the covariance matrix $\mathbf{\Sigma}_{B,i}$ is given by $\mathbf{\Sigma}_{B,i} = T^{-1}(\mathbf{\Sigma}_\varepsilon)_{i,i} \mathbf{I}_r$.

Let us look at the assumptions about the SNRs θ and ϑ_i in Corollaries 2 and 3. Consider the setting where $r \asymp 1$, $N \asymp T$, $c_\varepsilon \asymp 1$, $\|\mathbf{\Sigma}_\varepsilon^{1/2}\|_1 \asymp \|\mathbf{\Sigma}_\varepsilon^{1/2}\|_2 \asymp 1$, $\|(\mathbf{\Sigma}_\varepsilon^{1/2})_{i,\cdot}\|_1 \asymp 1$, and $\mathbf{\Sigma}_\varepsilon$ is well-conditioned. Then, the SNR condition in Corollary 2 (resp. Corollary 3) is equivalent to $\theta \gg \log^{3/2} N$ (resp. $\theta \gg \log N$ and $\vartheta_i \gg \log^{3/2} N$), indicating that inference of factors and factor loadings is achievable as long as the SNR grows faster than a polynomial rate of $\log N$. Our SNR condition is less restrictive than the prior work (e.g., Bai and Ng, 2023; Jiang et al., 2023; Choi and Yuan, 2024) that required the SNR to grow with a polynomial rate of N . Regardless of log factors, our SNR condition for inference is equivalent to $\theta \gg 1$, which is optimal since it matches the same condition required for consistency as commented after Corollary 1. Also, in the special case where $\mathbf{\Sigma}_\varepsilon$ is diagonal, i.e., there is no correlation in the noise matrix, our SNR condition for inference matches the minimal condition required for the subspace inference (e.g., Yan et al., 2024).

Both Corollaries 2 and 3 are stated in a non-asymptotic sense, and can be easily translated to asymptotic normality results. In practice, the confidence regions for the factors and the factor loadings can be constructed by replacing the asymptotic covariance matrices $\mathbf{\Sigma}_{F,t}$ and $\mathbf{\Sigma}_{B,i}$ with their consistent estimators:

$$\widehat{\mathbf{\Sigma}}_{F,t} = \widehat{\mathbf{\Sigma}}^{-1} \widehat{\mathbf{U}}^\top \widehat{\mathbf{\Sigma}}_\varepsilon^\top \widehat{\mathbf{U}} \widehat{\mathbf{\Sigma}}^{-1} \quad \text{and} \quad \widehat{\mathbf{\Sigma}}_{B,i} = \frac{1}{T} (\widehat{\mathbf{\Sigma}}_\varepsilon^\top)_{i,i} \mathbf{I}_r,$$

respectively. Here, $\widehat{\mathbf{\Sigma}}_\varepsilon^\top$ is the estimator of the noise covariance matrix which we will introduce in detail in Section 5.1.1. The intuition behind the above consistent estimators is that, $\widehat{\mathbf{U}}$ and $\widehat{\mathbf{\Sigma}}$ are the consistent estimators of $\mathbf{U}\mathbf{R}_U$ and $\mathbf{R}_U^\top \mathbf{\Lambda} \mathbf{R}_V$, respectively, and we will show this fact in the proof of Theorem 1.

4 Comparison with previous work

In this section, we compare our main results with related ones established in prior works Bai and Ng (2023); Jiang et al. (2023); Choi and Yuan (2024). All of them studied the PC estimators in weak factor models, and established estimation error bounds and asymptotic normality under different assumptions. Distinct from these three papers, all our results are entirely non-asymptotic, providing finite-sample characterizations for both the estimation error and the uncertainty level of statistical inference. From a technical viewpoint, in the regime $N \asymp T$, a notable advancement of our theory is that, our assumptions for inference require the growth rate of SNR to be faster than a polynomial rate of $\log N$, unlike the polynomial rate of N required in these three papers. Also, all these three papers assume that the r singular values $\{\sigma_i\}_{i=1}^r$ of factor loading \mathbf{B} are distinct, while our theory does not require such eigengap condition.

Having provided an overview of the advantages of our theory, we now make the comparisons in detail.

- Bai and Ng (2023) studied both the homogeneous case where $\sigma_i^2 \asymp N^\alpha$ for $i = 1, 2, \dots, r$, and the heterogeneous case where $\sigma_i^2 \asymp N^{\alpha_i}$ with $1 \geq \alpha_1 \geq \alpha_2 \geq \dots \geq \alpha_r > 0$. They required $\alpha_r > 1/2$ to prove the asymptotic normality for inference. However, the case when $0 < \alpha_r \leq 1/2$ has not been covered by their inferential theory. Our theory fills this gap and establishes the inference results even when σ_r do not grow with a polynomial rate of N . Besides the inference, our theory provides row-wise estimation error bounds for factors and factor loadings as detailed in Section 3.2, while the bounds in Bai and Ng (2023) are Frobenius norm bounds measuring the average error over all rows. The consistency and asymptotic

normality results in [Jiang et al. \(2023\)](#) are similar to those in [Bai and Ng \(2023\)](#), while the focus of [Jiang et al. \(2023\)](#) is to identify the so-called pseudo-true parameter that is consistently estimated by the PCA estimator. We note that both their frameworks accommodate temporal correlation, an aspect not covered by our theory.

- [Choi and Yuan \(2024\)](#) adopted the leave-one-out analysis similar in spirit to those used in matrix completion to investigate the PCA estimators. The setup they studied is the homogeneous case in [Bai and Ng \(2023\)](#) because they assumed that $N^{-\alpha} \mathbf{B}^\top \mathbf{B} \rightarrow \boldsymbol{\Sigma}_B$, which requires $\sigma_i^2 \asymp N^\alpha$ for all the singular values $\{\sigma_i\}_{i=1,2,\dots,r}$ of factor loading \mathbf{B} . When temporal correlation does not exist, similar to our results, they also filled the gap to establish the inference results for $0 < \alpha \leq 1/2$. In comparison, our theory is fully non-asymptotic and does not assume any asymptotic growth assumptions on the singular values. Also, the homogeneous case they studied assumes that the condition number $\kappa = \sigma_1/\sigma_r$ satisfies that $\kappa \asymp 1$, while our theory does not need this assumption and allows any growth rate of the condition number κ .

To enhance the clarity of the comparison of the SNR assumption for inference, we detailed the results in [Table 1](#) under a specific setup: Assumptions [1](#), [2](#), [3](#) hold; The cross-sectional and temporal dimensions satisfy that $N \asymp T \asymp n$; the number of factors satisfies $r \asymp 1$; the noise covariance matrix satisfies $\kappa_\varepsilon \asymp 1$, $\|\boldsymbol{\Sigma}_\varepsilon\|_2 \asymp 1$, and $\|\boldsymbol{\Sigma}_\varepsilon^{1/2}\|_1 \asymp 1$. Since the scale of the noise is reflected by $\boldsymbol{\Sigma}_\varepsilon$ and in our current setup we have that $\|\boldsymbol{\Sigma}_\varepsilon\|_2 \asymp 1$ and $\|\boldsymbol{\Sigma}_\varepsilon^{1/2}\|_1 \asymp 1$, the assumptions on the SNRs $\theta = \sigma_r/\sqrt{\|\boldsymbol{\Sigma}_\varepsilon\|_2}$ and $\vartheta = \sigma_r/\|\boldsymbol{\Sigma}_\varepsilon^{1/2}\|_1$ defined in [Section 3](#) are fully represented by the growth rate of the smallest singular value σ_r of factor loading matrix \mathbf{B} .

Table 1: SNR assumptions for inference of factor and factor loading

	Factor	Factor loading
Bai and Ng (2023)	$\sigma_r \gg N^{1/4}$	$\sigma_r \gg N^{1/4}$
Jiang et al. (2023)	$\sigma_r \gg \max(N^{1/4}, \kappa^2)$	$\sigma_r \gg \kappa^{1/2} N^{1/4}$
Choi and Yuan (2024)	$\sigma_r \asymp N^{\alpha/2}$ with $\alpha > 0$	$\sigma_r \gg N^{1/6} \log^{\omega/6} N$
Our Theory	$\sigma_r \gg \log^{3/2} N$	$\sigma_r \gg \log^{3/2} N$

[Table 1](#) demonstrates the advancement of our theory. For the conditions of [Choi and Yuan \(2024\)](#) in [Table 1](#), the parameter $\omega > 0$ is defined in Assumption B'' (iv) therein on the noise structure and they need two additional growth rate assumptions on σ_r (see Assumption D'' (i) therein). For other regimes where $T \ll N$ or $N \ll T$, similar comparisons can be made, so we omit the details for the limit of space.

5 Applications in econometrics

Our first-order approximations do not only advance the existing theories on the PCA under the weak factor model, but also pave the path for various statistical tests that are useful in economics and finance. In this section, we show four applications of our results.

5.1 The factor specification tests

Recall that the factor model is given by $\mathbf{X} = \mathbf{B}\mathbf{F}^\top + \mathbf{E}$, where the matrix \mathbf{F} is the realization of latent factors. Suppose we have time series data of some observed factors, e.g., the Fama-French factors. Our focus is on testing if the observed factor is in the linear space spanned by the latent factors \mathbf{F} . In particular, we examine this linear dependence in a flexible range of the whole period $[1, T]$. Formally, we consider an index set $S = \{t_1, t_2, \dots, t_{|S|}\} \subseteq \{1, 2, \dots, T\}$ of interest, and we have the data $\mathbf{v} = (v_{t_1}, v_{t_2}, \dots, v_{t_{|S|}})^\top \in \mathbb{R}^{|S|}$ of

an observed factor recorded at the time index set S . We test the hypothesis as follows,

$$H_0 : \quad \text{There exists } \mathbf{w} \in \mathbb{R}^r \text{ such that } \mathbf{v} = \mathbf{F}_{S,\cdot} \mathbf{w}. \quad (5.1)$$

Under the null hypothesis H_0 , we have that $v_t = \mathbf{F}_{t,\cdot} \mathbf{w} = \mathbf{f}_t^\top \mathbf{w}$ for any $t \in S$.

Under the strong factor model where the pervasiveness assumption holds, [Bai and Ng \(2006b\)](#) studied this problem and designed test statistics for the whole set, i.e., $S = \{1, 2, \dots, T\}$. Our study extends to the case when S is a subset of the whole time span $\{1, 2, \dots, T\}$ under the weak factor model. The subset S can be any specific time window of interest. This scenario is economically meaningful because the relationship $v_t = \mathbf{f}_t^\top \mathbf{w}$ may only be valid for relatively short periods, not necessarily across the entire span; see [Bai and Ng \(2006b\)](#) for the connections between the CAPM analysis and this problem. In [Section 6](#), we show that our factor specification test results strikingly reconcile with the economic cycles and financial crisis.

Our test statistic relies on the estimation of the noise covariance matrix Σ_ε . We denote by $\widehat{\Sigma}_\varepsilon^\tau$ the estimator of Σ_ε and we will elaborate its construction in the next section. In [Theorem 2](#) below, we keep the error bound $\|\widehat{\Sigma}_\varepsilon^\tau - \Sigma_\varepsilon\|_2$ in the final result.

Theorem 2. *For any given target error level $\delta > 0$, assume there exist some universal constants $c_0, C_0 > 0$ and sufficiently large constant $C_1 > 0$ such that, $n \geq C_1 r \log n$, $|S| \geq C_1(r + \log n)$, $T \geq C_0 |S| \delta^{-2} \kappa_\varepsilon^2 r^2 \log^4 n$,*

$$\theta \geq C_0 \delta^{-1} \frac{n}{T} \sqrt{|S| \kappa_\varepsilon^4 r^2 \log^4 n}, \quad \text{and} \quad \frac{\|\Sigma_\varepsilon^{1/2} \bar{\mathbf{U}}\|_{2,\infty}}{\|\Sigma_\varepsilon^{1/2}\|_2} \leq c_0 \delta \kappa_\varepsilon^{-1} \sqrt{\frac{T}{n}} |S|^{-3/2}; \quad (5.2)$$

and that for the estimator $\widehat{\Sigma}_\varepsilon^\tau$ of the noise covariance matrix Σ_ε , it holds

$$\frac{1}{\|\Sigma_\varepsilon\|_2} \|\widehat{\Sigma}_\varepsilon^\tau - \Sigma_\varepsilon\|_2 \leq c_0 \delta |S|^{-1/2} \kappa_\varepsilon^{-4} r^{-1} \log^{-2} n.$$

Then, under [Assumptions 1, 2, 3](#), and the null hypothesis H_0 in [\(5.1\)](#), we have that

$$\left| \mathbb{P} \left(\widehat{\mathcal{T}}(S, \mathbf{v}) \leq \chi_{1-\alpha}^2(|S| - r) \right) - (1 - \alpha) \right| \leq \delta, \quad (5.3)$$

where the test statistic $\widehat{\mathcal{T}}(S, \mathbf{v})$ is defined by

$$\widehat{\mathcal{T}}(S, \mathbf{v}) := \frac{1}{\widehat{\phi}} \mathbf{v}^\top (\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}) \mathbf{v} \quad \text{with} \quad \widehat{\phi} := \frac{1}{T} ((\widehat{\mathbf{V}}_{S,\cdot})^+ \mathbf{v})^\top \widehat{\Sigma}^{-1} \widehat{\mathbf{U}}^\top \widehat{\Sigma}^\tau \widehat{\mathbf{U}} \widehat{\Sigma}^{-1} (\widehat{\mathbf{V}}_{S,\cdot})^+ \mathbf{v}.$$

Remark 2. If all the entries of the noise matrix \mathbf{Z} in [\(2.4\)](#) are Gaussian, then we do not need the second assumption in [\(5.2\)](#).

The idea of the test statistic $\widehat{\mathcal{T}}(S, \mathbf{v})$ is to utilize the fact that, under the null hypothesis H_0 , the residual vector of \mathbf{v} is zero after projection onto the column space of $\mathbf{F}_{S,\cdot}$. Note that $\widehat{\mathbf{V}}_{S,\cdot}$ estimates the column space of $\mathbf{F}_{S,\cdot}$. So, we construct $\widehat{\mathcal{T}}(S, \mathbf{v})$ by computing the ℓ_2 -norm of the projection residual vector $\|(\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}) \mathbf{v}\|_2^2 = \mathbf{v}^\top (\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}) \mathbf{v}$. The component $\widehat{\phi}$ estimates the variance of $(\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}) \mathbf{v}$, and needs the estimator $\widehat{\Sigma}_\varepsilon^\tau$ of noise covariance matrix as input.

Our focus is on the regime where the size of subset S is small. This is economically meaningful because the linear relationship between observed factors and latent factors usually holds only for a short horizon. Based on our distributional results in [\(5.3\)](#), the case when S is the whole period, i.e., $|S| = T$, can be handled straightforwardly by applying the Gaussian approximation to the Chi-square distribution. Our assumption on the size $|S|$ can be summarized as $r + \log n \ll |S| \ll T / (\kappa_\varepsilon^2 r^2 \log^4 n)$. On the one hand, we require that $|S|$ cannot be too small so that $\mathbf{F}_{S,\cdot}$ is of full column rank and its singular value is close to $\sqrt{|S|}$, which is crucial for our proof. On other hand, we require that $|S|$ cannot be too large, otherwise the null distribution $\chi^2(|S| - r)$ with mean $|S| - r$ would diverge in such a case, complicating the proof of proximity between $\chi^2(|S| - r)$ and the law of test statistic $\widehat{\mathcal{T}}(S, \mathbf{v})$.

The assumption on SNR is that $\theta \gg \frac{n}{T} \sqrt{|S| \kappa_\varepsilon^4 r^2 \log^4 n}$. Consider the setting where $r \asymp 1$, $N \asymp T$, and Σ_ε is well-conditioned (i.e., $\kappa_\varepsilon \asymp 1$). Then, the SNR condition becomes $\theta \gg \sqrt{|S|} \log^4 n$, illustrating the

effect of the subset size $|S|$ on the growth rate required for SNR θ . The $\sqrt{|S|}$ factor appears in the SNR because we use the Chi-square distribution as the null distribution to accommodate the general case when $|S|$ is small. When $|S| \asymp 1$, regardless of the log factors, the SNR condition is equivalent to $\theta \gg 1$, which match the minimal condition required for the consistent estimations of the left and right singular subspaces if there is no correlation in the noise matrix (e.g., [Yan et al., 2024](#)).

[Bai and Ng \(2006b\)](#) studied the factor specification test under the whole period case, i.e., $S = \{1, 2, \dots, T\}$. They derived the approximate cumulative distribution function (CDF) for their test statistic, but they did not provide a finite sample error bound for it and their results were under the strong factor model. In comparison, our results give a precise characterization of the error and adapt to the weak factor model. In our case, both the low SNR in the weak factor model and the subset S in the hypothesis H_0 make it challenging to design the test statistics. The subspace perturbation bounds in [Theorem 1](#) enable us to conquer these difficulties and establish non-asymptotic analysis of our test statistic for any arbitrary subset S .

5.1.1 Estimation of the noise covariance matrix Σ_ε

We accommodate the case where the cross-sectional dimension N is much larger than the temporal dimension T . In such case, estimating the high-dimensional covariance matrix Σ_ε becomes challenging yet is vital for numerous statistical tests, including the factor specification test and the two-sample test for betas discussed in the next section. Following the approaches in, e.g., [Bickel and Levina \(2008\)](#); [Fan et al. \(2011, 2013\)](#), we assume that the error covariance matrix Σ_ε is sparse in a suitable sense and use the adaptive thresholding method to estimate Σ_ε . It's important to note that the assumed sparsity of Σ_ε is solely for our statistical tests and is not a required condition to establish our main theories on the subspace perturbation bounds in [Theorem 1](#).

Assumption 6. *For some $q \in [0, 1)$, there exists a constant $s(\Sigma_\varepsilon) > 0$ such that*

$$\max_{1 \leq i \leq N} \sum_{j=1}^N ((\Sigma_\varepsilon)_{i,i} (\Sigma_\varepsilon)_{j,j})^{(1-q)/2} |(\Sigma_\varepsilon)_{i,j}|^q \leq s(\Sigma_\varepsilon).$$

The assumption is slightly weaker than those in, e.g., [Bickel and Levina \(2008\)](#); [Fan et al. \(2013\)](#), where they assumed that $\lambda_{\min}(\Sigma_\varepsilon) \asymp 1$, $\lambda_{\max}(\Sigma_\varepsilon) \asymp 1$, and $\max_{1 \leq i \leq N} \sum_{j=1}^N |(\Sigma_\varepsilon)_{i,j}|^q \leq s_0$ for some sparsity parameter s_0 . In particular, for $q = 0$, it constrains on the maximum number of nonzero elements of Σ_ε . This sparsity assumption for the noise covariance Σ_ε is also natural in economics and finance. As the latent common factors largely explain the co-movements in the panel data, the correlation among individual asset's idiosyncratic noises should be close to zero. A specific example of the sparse structure of Σ_ε arises from the remaining sector effects ([Gagliardini et al., 2016](#)).

We estimate the idiosyncratic noise matrix using $\widehat{\mathbf{E}} = \mathbf{X} - \widehat{\mathbf{B}}\widehat{\mathbf{F}}^\top$, where $\widehat{\mathbf{B}}$ and $\widehat{\mathbf{F}}$ are the PCA estimators of factor loadings and factors defined in [Section 3.2](#). Then, the pilot estimator of the covariance matrix Σ_ε is given by $\widehat{\Sigma}_\varepsilon = T^{-1}\widehat{\mathbf{E}}\widehat{\mathbf{E}}^\top$. We will show in [Lemma 1](#) later that $\max_{1 \leq i, j \leq N} |(\widehat{\Sigma}_\varepsilon - \Sigma_\varepsilon)_{i,j}| \lesssim \epsilon_{N,T} \sqrt{(\Sigma_\varepsilon)_{i,i} (\Sigma_\varepsilon)_{j,j}}$, where $\epsilon_{N,T} > 0$ is negligible as $N, T \rightarrow \infty$. Then we apply an adaptive thresholding method ([Bickel and Levina, 2008](#)) to obtain the sparse covariance matrix estimator $\widehat{\Sigma}_\varepsilon^\tau$ as follows,

$$(\widehat{\Sigma}_\varepsilon^\tau)_{i,j} := \begin{cases} (\widehat{\Sigma}_\varepsilon)_{i,j}, & \text{if } i = j, \\ h((\widehat{\Sigma}_\varepsilon)_{i,j}, \tau_{i,j}), & \text{if } i \neq j, \end{cases}$$

where $h(z, \tau)$ is a thresholding function with the threshold value τ . Here, the threshold value $\tau_{i,j}$ is set adaptively to $\tau_{i,j} = C\epsilon_{N,T} \sqrt{(\widehat{\Sigma}_\varepsilon)_{i,i} (\widehat{\Sigma}_\varepsilon)_{j,j}}$ with some large constant $C > 0$. The idea behind the adaptive thresholding is examining the sample correlation matrix and retaining entries exceeding $C\epsilon_{N,T}$ in magnitude. The specific examples of the function h include the hard-thresholding function $h(z, \tau) = z1_{\{|z| < \tau\}}$, among other common thresholding functions like the soft-thresholding function and SCAD ([Fan and Li, 2001](#)). In general, we require that the thresholding function $h(z, \tau)$ to satisfy: (i) $h(z, \tau) = 0$ if $|z| < \tau$; (ii) $|h(z, \tau) - z| < \tau$. The estimation error is given as follows.

Lemma 1. *Suppose that all the assumptions in Corollary 1 hold, and that there exists a sufficiently small constant $\epsilon_{N,T} > 0$ such that*

$$\max_{1 \leq i, j \leq N} \left(\frac{1}{\sqrt{T}} (1 + (1 + \gamma_{\epsilon, j})\gamma_i + (1 + \gamma_{\epsilon, i})\gamma_j) + (\beta_i \gamma_j + \beta_j \gamma_i) \left(\frac{n}{\theta^2 T} + \frac{1}{\theta \sqrt{T}} \right) \right) \cdot r \log^2 n \leq \epsilon_{N,T}, \quad (5.4)$$

where

$$\beta_i := \frac{\|\mathbf{b}_i\|_2}{\sqrt{(\boldsymbol{\Sigma}_\epsilon)_{i,i}}}, \quad \gamma_i := \frac{\sigma_r \|\bar{\mathbf{U}}_{i,\cdot}\|_2}{\sqrt{(\boldsymbol{\Sigma}_\epsilon)_{i,i}}} \quad \text{and} \quad \gamma_{\epsilon, i} := \frac{\|(\boldsymbol{\Sigma}_\epsilon^{1/2})_{i,\cdot}\|_1}{\|(\boldsymbol{\Sigma}_\epsilon^{1/2})_{i,\cdot}\|_2}.$$

Then, we have that, with probability at least $1 - O(n^{-10})$, it holds $\max_{1 \leq i, j \leq N} |(\boldsymbol{\Sigma}_\epsilon)_{i,i}(\boldsymbol{\Sigma}_\epsilon)_{j,j}|^{-1/2} |(\widehat{\boldsymbol{\Sigma}}_\epsilon - \boldsymbol{\Sigma}_\epsilon)_{i,j}| \leq C_0 \epsilon_{N,T}$, for a universal constant $C_0 > 0$. Further, under Assumption 6, the generalized thresholding estimator $\widehat{\boldsymbol{\Sigma}}_\epsilon^\tau$ satisfies that, there exists some universal constant $C_0 > 0$ such that, with probability at least $1 - O(n^{-10})$,

$$\|\widehat{\boldsymbol{\Sigma}}_\epsilon^\tau - \boldsymbol{\Sigma}_\epsilon\|_2 \leq C_0 (\epsilon_{N,T})^{1-q} s(\boldsymbol{\Sigma}_\epsilon).$$

The above error bounds for the estimators $\widehat{\boldsymbol{\Sigma}}_\epsilon$ and $\widehat{\boldsymbol{\Sigma}}_\epsilon^\tau$ differ from the existing literature owing to our row-wise subspace perturbation bounds in Theorem 1. Let us interpret the condition (5.4) under some specific settings. Consider the setting where $r \asymp 1$, $N \asymp T$, $\bar{\mathbf{U}}$ is μ -incoherent, $\|\mathbf{B}\|_{2,\infty} \lesssim 1$ and $\boldsymbol{\Sigma}_\epsilon$ is well-behaved such that $\gamma_{\epsilon, i} \asymp 1$, $(\boldsymbol{\Sigma}_\epsilon)_{i,i} \asymp 1$, and $\|\boldsymbol{\Sigma}_\epsilon\|_2 \asymp 1$. Then, the condition (5.4) is equivalent to $T \geq (\sigma_r \log^2 n) / \epsilon_{N,T}$, which requires the sample size T is sufficiently large. According to Lemma 1, fulfilling our assumption on the estimation error $\|\widehat{\boldsymbol{\Sigma}}_\epsilon^\tau - \boldsymbol{\Sigma}_\epsilon\|_2$ required for Theorem 2 merely requires that the estimation error $\epsilon_{N,T}$ satisfies

$$\epsilon_{N,T} \ll \left((\|\boldsymbol{\Sigma}_\epsilon\|_2 / s(\boldsymbol{\Sigma}_\epsilon)) |S|^{-1/2} \kappa_\epsilon^{-4} r^{-1} \log^{-2} n \right)^{1/(1-q)}.$$

In subsequent applications, we will directly use $\epsilon_{N,T}$, instead of $\widehat{\boldsymbol{\Sigma}}_\epsilon^\tau$, to express assumptions on the estimation error of the noise covariance matrix.

5.2 Test for structural breaks in betas

The discussion so far has studied the factor models with constant betas. However, in many cases, there is a possibility of time-variant betas, and it is important to test for structural changes in betas (e.g., Stock and Watson, 2009; Breitung and Eickmeier, 2011; Chen et al., 2014; Han and Inoue, 2015). We test for two time periods $\Gamma_1 = \{t_1 + 1, t_1 + 2, \dots, t_1 + T_1\}$ and $\Gamma_2 = \{t_2 + 1, t_2 + 2, \dots, t_2 + T_2\}$ that are possibly not consecutive, i.e., $t_1 + T_1 \leq t_2 + 1$. The factor models in these two periods are subject to a structural break in beta: for each given individual unit i ,

$$\begin{aligned} x_{i,t} &= \mathbf{f}_t^\top \mathbf{b}_i^1 + \varepsilon_{i,t} & \text{for } t \in \Gamma_1, \\ x_{i,t} &= \mathbf{f}_t^\top \mathbf{b}_i^2 + \varepsilon_{i,t} & \text{for } t \in \Gamma_2, \end{aligned}$$

where i is the index of the cross-sectional unit of interest. Here, we assume that the factors are the same, but the betas can be different. For example, we might ask if a external shock such as the 2008 financial crises has caused the changes in factor loadings before and after the shock with a transition period (T_1, T_2) .

The data consist of two panels, \mathbf{X}^1 and \mathbf{X}^2 , during Γ_1 and Γ_2 , respectively, where $\mathbf{X}^1 \in \mathbb{R}^{N \times T_1}$ and $\mathbf{X}^2 \in \mathbb{R}^{N \times T_2}$. We test the hypothesis as follows:

$$H_0 : \mathbf{b}_i^1 = \mathbf{b}_i^2 \leftrightarrow H_1 : \mathbf{b}_i^1 \neq \mathbf{b}_i^2.$$

Different from our approach, Chen et al. (2014); Han and Inoue (2015) tested for the entire factor loading, with their null hypothesis being $H_0 : \mathbf{b}_i^1 = \mathbf{b}_i^2$ for $\forall i$. Stock and Watson (2009); Breitung and Eickmeier (2011) also studied the test for a given i like ours, while Stock and Watson (2009) focused on empirical studies and Breitung and Eickmeier (2011) developed and established theories for the test statistics. Our work differs in that, as we will show below, the effectiveness of our test statistic does not require the pervasiveness assumption needed in Breitung and Eickmeier (2011). In addition, our formulation allows the

gap between the two periods Γ_1 and Γ_2 , which can model the special transition periods we purposely, e.g., financial crisis, enabling us to test for structural breaks before and after these special periods.

To construct the test statistic, we first merge the two panels \mathbf{X}^1 and \mathbf{X}^2 into a $N \times (T_1 + T_2)$ -dimensional data matrix $\mathbf{X} = (\mathbf{X}^1, \mathbf{X}^2)$. Next, we apply SVD as described in Algorithm 1 to obtain an estimator of the factors: $\widehat{\mathbf{F}} = \sqrt{T}\widehat{\mathbf{V}}$, where $T = T_1 + T_2$ and $\widehat{\mathbf{U}}\widehat{\mathbf{\Sigma}}\widehat{\mathbf{V}}^\top$ is the truncated rank- r SVD of $T^{-1/2}\mathbf{X}$ with $\widehat{\mathbf{U}} \in \mathbb{R}^{N \times r}$ and $\widehat{\mathbf{V}} \in \mathbb{R}^{T \times r}$. Then, we split the estimated factors into $\widehat{\mathbf{F}} = ((\widehat{\mathbf{F}}^1)^\top, (\widehat{\mathbf{F}}^2)^\top)^\top$ according to two time periods Γ_1 and Γ_2 , where $\widehat{\mathbf{F}}^j \in \mathbb{R}^{T_j \times r}$. Subsequently, we obtain the estimator for \mathbf{b}_i^j by regressing $\mathbf{X}_{i,\cdot}^j$ on $\widehat{\mathbf{F}}^j$:

$$\widehat{\mathbf{b}}_i^j = ((\widehat{\mathbf{F}}^j)^\top \widehat{\mathbf{F}}^j)^{-1} (\widehat{\mathbf{F}}^j)^\top (\mathbf{X}_{i,\cdot}^j)^\top \quad \text{for } j = 1, 2.$$

Finally, under null hypothesis H_0 , we show that $\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2$ is approximately Gaussian via the first-order approximation results in Theorem 1, and construct the test statistic using the plug-in estimator of the covariance matrix.

Theorem 3. *Suppose that the assumptions in Theorem 1 hold, and the covariance estimator $\widehat{\mathbf{\Sigma}}_\varepsilon^\tau$ satisfies the conditions as in Lemma 1. For any given target error level $\delta > 0$, assume there exists some universal constants $c_0, C_0, C_U > 0$ such that $\min(T_1, T_2) \geq C_0(r + \log n)$, $\|\mathbf{U}_{i,\cdot}\|_2 \leq C_U \sqrt{T/(T_1 T_2)}$, $T \geq C_0^2 C_U^2 \delta^{-2} \kappa_\varepsilon r \log^3 n$, $\vartheta_i \geq C_0 C_U^2 \delta^{-1} T^{-1/2} \kappa_\varepsilon^2 r \log n$,*

$$\theta \geq C_0 \left\{ C_U^2 \frac{1}{\delta} \left(\kappa_\varepsilon \frac{n}{T} + \kappa_\varepsilon^2 \sqrt{\frac{n}{T}} \|\mathbf{U}_{i,\cdot}\|_2 \right) + \sqrt{T_1 T_2 / T} \frac{1}{\delta} \kappa_\varepsilon \right\} r \log^{3/2} n,$$

and $C_0 r^{3/2} \|\mathbf{w}\|_\infty / \|\mathbf{w}\|_2 \leq \delta$, where \mathbf{w} is a vector defined as $\mathbf{w} := [\mathbf{\Sigma}_\varepsilon^{1/2}(\mathbf{I}_N + \mathbf{U}\mathbf{U}^\top)]_{\cdot, i}$; for the parameter $\epsilon_{N,T}$ (cf. Lemma 1) that captures the estimation error of the noise covariance matrix $\mathbf{\Sigma}_\varepsilon$, assume that $\epsilon_{N,T} \leq c_0 \delta / (\kappa_\varepsilon^2 r \log n)$ and $\epsilon_{N,T} \leq c_0 [\delta \|\mathbf{\Sigma}_\varepsilon\|_2 / (s(\mathbf{\Sigma}_\varepsilon) \|\mathbf{U}_{i,\cdot}\|_2 \kappa_\varepsilon^2 r \log n)]^{1/(1-q)}$. Then, under the null hypothesis H_0 : $\mathbf{b}_i^1 = \mathbf{b}_i^2$, we have that

$$\left| \mathbb{P} \left(\widehat{\mathcal{B}}_i \leq \chi_{1-\alpha}^2(r) \right) - (1 - \alpha) \right| \leq \delta,$$

where the test statistic $\widehat{\mathcal{B}}_i$ is given by

$$\widehat{\mathcal{B}}_i := \frac{1}{T \widehat{\varphi}_i} (\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2)^\top \left[\prod_{j=1}^2 (\widehat{\mathbf{F}}^j)^\top \widehat{\mathbf{F}}^j \right] (\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2),$$

with $\widehat{\varphi}_i := (\widehat{\mathbf{\Sigma}}_\varepsilon^\tau + \widehat{\mathbf{U}}\widehat{\mathbf{U}}^\top \widehat{\mathbf{\Sigma}}_\varepsilon^\tau + \widehat{\mathbf{\Sigma}}_\varepsilon^\tau \widehat{\mathbf{U}}\widehat{\mathbf{U}}^\top + \widehat{\mathbf{U}}\widehat{\mathbf{U}}^\top \widehat{\mathbf{\Sigma}}_\varepsilon^\tau \widehat{\mathbf{U}}\widehat{\mathbf{U}}^\top)_{i,i}$.

Remark 3. If all the entries of the noise matrix \mathbf{Z} in (2.4) are Gaussian, then we do not need the assumption that $C_0 r^{3/2} \|\mathbf{w}\|_\infty / \|\mathbf{w}\|_2 \leq \delta$, where $\mathbf{w} = [\mathbf{\Sigma}_\varepsilon^{1/2}(\mathbf{I}_N + \mathbf{U}\mathbf{U}^\top)]_{\cdot, i}$.

To illustrate our assumptions, we consider the following setting similar to that discussed after Theorem 1: $r \asymp 1$, $T_1 \asymp T \asymp N$, $T_1 \gg T_2$, $\mathbf{\Sigma}_\varepsilon$ is well-conditioned and its sparsity parameter satisfies $s(\mathbf{\Sigma}_\varepsilon) \lesssim \|\mathbf{\Sigma}_\varepsilon\|_2$, and the column subspace of \mathbf{B} is incoherent. Then, fulfilling our assumptions merely requires that, $\theta \gg \log^{3/2} n$ and $\vartheta_i \gg T^{-1/2} \log n$ for the SNR, $T \gg \log^3 n$ for the sample size, and $\epsilon_{N,T} \ll \min(\log^{-1} n, (N^{1/2} / \log n)^{1/(1-q)})$ for the estimation error of $\mathbf{\Sigma}_\varepsilon$. In particular, our SNR conditions are less restrictive than prior work (e.g., Breitung and Eickmeier, 2011) that needed the pervasiveness assumption, which required the SNR to grow as $\theta \gg N^{1/4}$. Our results adapt to the weak factor model with the cross-sectional correlations, and to the best of our knowledge, no prior work has developed the test statistics for the structural break test under this case.

5.3 The two-sample test for betas

In finance and economics, besides the latent common factors that drive the co-movements of asset returns, the factor loadings, also known as betas, are important as well, which measure the sensitivity of asset return to the movements of the factors. Consider the example that the panel data \mathbf{X} is the stock return, where the i -th row of \mathbf{X} represents the time series data of the i -th stock. Then, the factor loadings \mathbf{B} assess the risk exposure of these stocks to the latent common factors \mathbf{f}_t , with the i -th row \mathbf{b}_i being the i -th stock's beta.

For any distinct i and j , we aim to test the hypothesis that $\mathbf{b}_i = \mathbf{b}_j$, i.e., if the i -th and the j -th stocks have the same risk exposure on the common factors. It is a statistical approach to evaluate the similarity in risk structure between two stocks. Our test statistic is similar to the idea of two-sample test: we show that $\widehat{\mathbf{B}}_{i,\cdot} - \widehat{\mathbf{B}}_{j,\cdot}$ is approximately Gaussian and then derive a Chi-square test statistic. In Theorem 2 below, we construct the test statistic and show its validity.

Theorem 4. *Suppose that the assumptions in Theorem 1 hold. Assume that $s(\boldsymbol{\Sigma}_\varepsilon)/\lambda_{\min}(\boldsymbol{\Sigma}_\varepsilon) \leq C_\varepsilon$ for some universal constant C_ε , and the covariance estimator $\widehat{\boldsymbol{\Sigma}}_\varepsilon^T$ satisfies the conditions as in Lemma 1. For any given target error level $\delta > 0$, assume there exists some universal constants $c_0, C_0 > 0$ such that $T \geq C_0 r^2 \delta^{-2} \log n$,*

$$\max(\|\bar{\mathbf{U}}_{i,\cdot}\|_2, \|\bar{\mathbf{U}}_{j,\cdot}\|_2) \leq c_0 \delta \left(\frac{n}{\theta\sqrt{T}} + 1\right)^{-1} \kappa_\varepsilon^{-1/2} r^{1/2} \log^{-1} n, \quad (5.5a)$$

and

$$\min(\vartheta_i, \vartheta_j) \geq C_0 \delta^{-1} \sqrt{\kappa_\varepsilon} \frac{\theta n}{\vartheta\sqrt{T}} (\|\bar{\mathbf{U}}\|_{2,\infty} + \frac{1}{\sqrt{n}}) \sqrt{r} \log^{3/2} n; \quad (5.5b)$$

for the parameter $\epsilon_{N,T}$ (cf. Lemma 1) that captures the estimation error of the noise covariance matrix $\boldsymbol{\Sigma}_\varepsilon$, assume that $\epsilon_{N,T} \leq c_0 (\delta r^3 \log^2 n)^{1/(1-q)}$. Then, for any $i, j \in \{1, 2, \dots, N\}$ satisfying $i \neq j$, under the null hypothesis $H_0 : \mathbf{b}_i = \mathbf{b}_j$, we have that

$$\left| \mathbb{P} \left(\widehat{\mathcal{B}}_{ij} \leq \chi_{1-\alpha}^2(r) \right) - (1 - \alpha) \right| \leq \delta,$$

where the test statistic $\widehat{\mathcal{B}}_{ij}$ is given by

$$\widehat{\mathcal{B}}_{ij} := T((\widehat{\boldsymbol{\Sigma}}_\varepsilon^T)_{i,i} + (\widehat{\boldsymbol{\Sigma}}_\varepsilon^T)_{j,j} - 2(\widehat{\boldsymbol{\Sigma}}_\varepsilon^T)_{i,j})^{-1} \|\widehat{\mathbf{B}}_{i,\cdot} - \widehat{\mathbf{B}}_{j,\cdot}\|_2^2.$$

Remark 4. If all the entries of the noise matrix \mathbf{Z} in (2.4) are Gaussian, then we do not need the assumption that $T \geq C_0 r^2 \delta^{-2} \log n$ on the sample size T .

To illustrate the SNR conditions (5.5a)–(5.5b), we consider the setting discussed after Theorem 1, i.e., $r \asymp 1$, $N \asymp T$, $\boldsymbol{\Sigma}_\varepsilon$ is well-conditioned, and the column subspace of \mathbf{B} is incoherent. Then, fulfilling our assumptions (5.5a)–(5.5b) merely requires that $\theta \gg \log n$ and $\theta^{-1} \vartheta \min(\vartheta_i, \vartheta_j) \gg \log^{3/2} n$. These SNR conditions are less restrictive than the prior work (e.g., Bai and Ng, 2023; Jiang et al., 2023; Choi and Yuan, 2024) that required the SNR to grow with a polynomial rate of n .

To the best of our knowledge, no prior work has developed the test statistics for this two-sample test of betas. Our results are applicable to the weak factor model with the cross-sectional correlations. The subspace perturbation bounds in Theorem 1 pave the way for us to statistically assess the similarity for arbitrary two rows of factor loadings \mathbf{B} .

5.4 Statistical inference for the systematic risks

In the factor model $x_{i,t} = \mathbf{b}_i^\top \mathbf{f}_t + \varepsilon_{i,t}$, the risk associated with stock return $x_{i,t}$ is decomposed into two parts (e.g., Bai, 2003) — systematic risk from the common component $\mathbf{b}_i^\top \mathbf{f}_t$ and idiosyncratic risk from the noise $\varepsilon_{i,t}$. Systematic risk, often referred to as market risk, is integral in financial economics as it represents the inherent risk affecting the entire market or market segment. This risk, driven by broader economic forces such as inflation, political events, and changes in interest rates, comes from the risk factors \mathbf{f}_t that explain the systematic co-movements and impacts all the stocks.

A standard metric of the systematic risk is the variance of $\mathbf{b}_i^\top \mathbf{f}_t$, which is given by $\text{Var}(\mathbf{b}_i^\top \mathbf{f}_t) = \|\mathbf{b}_i\|_2^2$. Our focus is on constructing the confidence interval (CI) for systematic risk $\|\mathbf{b}_i\|_2^2$. Similar to the idea we conduct the inference for beta \mathbf{b}_i in Corollary 3 where the estimator of \mathbf{b}_i is $\widehat{\mathbf{B}}_{i,\cdot}$, we show that $\|\widehat{\mathbf{B}}_{i,\cdot}\|_2^2 - \|\mathbf{b}_i\|_2^2$ is approximately Gaussian and then construct the CI for $\|\mathbf{b}_i\|_2^2$. We present the CI and its validity in Theorem 5.

Theorem 5. *Suppose that the assumptions in Theorem 1 hold. Assume that covariance estimator $\widehat{\boldsymbol{\Sigma}}_\varepsilon^T$ satisfies the conditions as in Lemma 1. For any given target error level $\delta > 0$, assume there exist some*

universal constants $c_0, C_0, C_U > 0$ such that $T \geq C_0 \delta^{-2} \log n$, $\|\bar{\mathbf{U}}\|_{2,\infty} \leq C_U \theta / \vartheta$, $\|\bar{\mathbf{U}}\|_{2,\infty} \leq C_U \sqrt{T} \vartheta / n$, $\|\bar{\mathbf{U}}\|_{2,\infty} \leq C_U \sqrt{T} \theta^2 / (\vartheta n)$, $\|\mathbf{b}_i\|_2^2 / (\boldsymbol{\Sigma}_\varepsilon)_{i,i} \leq c_0 \delta^2 r^{-1} \log^{-1} n$,

$$\vartheta_i \geq C_0 \delta^{-1} \frac{\sqrt{\kappa_\varepsilon} \theta}{\vartheta_i \sqrt{T}} \frac{1}{\|\bar{\mathbf{U}}_{i,\cdot}\|_2} r \log^2 n, \quad \vartheta \geq C_0 \delta^{-1} \frac{\sqrt{\kappa_\varepsilon} \theta}{\vartheta_i \sqrt{T}} (\sqrt{n} + n \|\bar{\mathbf{U}}\|_{2,\infty}) r \log^2 n \quad (5.6a)$$

$$\vartheta_i \geq C_0 \delta^{-1} \sqrt{\kappa_\varepsilon} \left(\frac{1}{\sqrt{T}} + \frac{n}{\theta T} \right) \left(1 + \frac{1}{\|\bar{\mathbf{U}}_{i,\cdot}\|_2} \right) r \log n, \quad \text{and} \quad \|\bar{\mathbf{U}}_{i,\cdot}\|_2 \leq c_0 \delta \kappa_\varepsilon^{-1/2} \left(1 + \frac{n}{\theta \sqrt{T}} \right)^{-1}; \quad (5.6b)$$

for the parameter $\epsilon_{N,T}$ (cf. Lemma 1) that captures the estimation error of the noise covariance matrix $\boldsymbol{\Sigma}_\varepsilon$, assume that $\epsilon_{N,T} \leq c_0 \delta r^{-1} \log^{-1/2} n$. Then we have that, for any $1 \leq i \leq N$,

$$\left| \mathbb{P}(\|\mathbf{b}_i\|_2^2 \in CI_i^{\mathbf{B},(1-\alpha)}) - (1-\alpha) \right| \leq \delta,$$

where the confidence interval is constructed as

$$CI_i^{\mathbf{B},(1-\alpha)} := [\|\widehat{\mathbf{B}}_{i,\cdot}\|_2^2 - \widehat{\sigma}_{B,i} z_{1-\frac{1}{2}\alpha}, \|\widehat{\mathbf{B}}_{i,\cdot}\|_2^2 + \widehat{\sigma}_{B,i} z_{1-\frac{1}{2}\alpha}], \quad \text{with} \quad \widehat{\sigma}_{B,i} := \frac{2}{\sqrt{T}} \sqrt{(\widehat{\boldsymbol{\Sigma}}_\varepsilon^T)_{i,i}} \|\widehat{\mathbf{B}}_{i,\cdot}\|_2,$$

and z_p is the p -quantile of the standard Gaussian $\mathcal{N}(0,1)$.

Remark 5. If all the entries of the noise matrix \mathbf{Z} in (2.4) are Gaussian, then we do not need the assumption that $T \geq C_0 \delta^{-2} \log n$ on the sample size T .

We now explain the assumptions under the setting discussed after Theorem 4. In this case, to make (5.6a) hold, it suffices to assume that $\theta^{-1} \vartheta_i \vartheta \gg \log^2 n$ and $\theta^{-1} \vartheta_i^2 \gg \log^2 n$; to make (5.6b) hold, it suffices to assume that $\theta \gg 1$, $\vartheta_i \theta \gg \log n$, and $\vartheta_i \gg T^{-1/2} \log n$. All these conditions require only polynomial growth rates of $\log n$ for the SNRs ϑ_i , ϑ , and θ . The assumption on $\|\bar{\mathbf{U}}\|_{2,\infty}$ is the same with that we required in Corollary 1 to prove the estimation error for factor loading.

On one hand, as shown in Theorem 5, the CI width $\widehat{\sigma}_{B,i}$ is proportional to the square root of the product of noise level estimator $(\widehat{\boldsymbol{\Sigma}}_\varepsilon^T)_{i,i}$ and systematic risk estimator $\|\widehat{\mathbf{B}}_{i,\cdot}\|_2^2$. On the other hand, in the proof of Theorem 5, we show that the bias of systematic risk estimator $\|\widehat{\mathbf{B}}_{i,\cdot}\|_2^2$ is proportional to $\|\mathbf{b}_i\|_2^2$. Thus, the assumption that $\|\mathbf{b}_i\|_2^2 / (\boldsymbol{\Sigma}_\varepsilon)_{i,i} \leq c_0 \delta^2 r^{-1} \log^{-1} n$ is essential because the validity of inference hinges on the dominance of the CI width over the bias. Consequently, we assume that the ratio between systematic risk $\|\mathbf{b}_i\|_2^2$ and noise level $(\boldsymbol{\Sigma}_\varepsilon)_{i,i}$ cannot be too large. The row-wise subspace perturbation bounds in Theorem 1 facilitate us to conduct statistical inference for the systematic risk of any given stock in the panel data.

6 Numerical experiments

In this section, we conduct Monte Carlo simulations to demonstrate our inferential theories for the PC estimators in the weak factor models. Additionally, our empirical studies reveal that the testing results based on our test statistics surprisingly align with the economic cycles and financial crisis periods.

6.1 Monte Carlo simulations

To make our simulations similar to the real applications, we use the standard Fama-French three-factor model:

$$x_{i,t} = \mathbf{b}_i^\top \mathbf{f}_t + \varepsilon_{i,t}, \quad 1 \leq i \leq N, 1 \leq t \leq T,$$

where the dimension of the latent factor \mathbf{f}_t is set to $r = 3$. The idiosyncratic noise $\varepsilon_{i,t}$ exhibits cross-sectional correlations, and the noise covariance matrix $\boldsymbol{\Sigma}_\varepsilon$ is sparse.

We generate the factor loadings $\{\mathbf{b}_i\}_{i=1}^N$, the factors $\{\mathbf{f}_t\}_{t=1}^T$, and the noise terms $\{\varepsilon_t\}_{t=1}^T$ independently from $\mathcal{N}(0, \boldsymbol{\Sigma}_b)$, $\mathcal{N}(0, s_f \boldsymbol{\Sigma}_f)$, and $\mathcal{N}(0, \boldsymbol{\Sigma}_\varepsilon)$, respectively, where $\boldsymbol{\varepsilon}_t = (\varepsilon_{1,t}, \dots, \varepsilon_{N,t})^\top$. Both $\boldsymbol{\Sigma}_b$ and $\boldsymbol{\Sigma}_f$ are set to $r \times r$ identity matrices. We generate $\boldsymbol{\Sigma}_\varepsilon$ as a block-diagonal matrix $\boldsymbol{\Sigma}_\varepsilon = \text{diag}(\mathbf{A}_1, \mathbf{A}_2, \dots, \mathbf{A}_J)$, where the number of blocks is set to $J = 20$. We set $N = 300$ and $T = 200$. For each \mathbf{A}_i , we construct it as an equi-correlation matrix $\mathbf{A}_i = (1 - \rho_i) \mathbf{I}_m + \rho_i \mathbf{1}_m \mathbf{1}_m^\top$, where the block size m is set to $m = N/J = 15$, $\mathbf{1}_m$

is the m -dimensional vector of ones, and ρ_i is drawn from a uniform distribution on $[0, 0.5]$. To validate our test statistic for two-sample test of betas, we set \mathbf{b}_2 equal to \mathbf{b}_1 after generating the loading matrix \mathbf{B} . This slight modification allows us to examine our test statistics under the null hypothesis $\mathbf{b}_1 = \mathbf{b}_2$. In our simulation results, we report the values of $\theta = \sigma_r / \|\boldsymbol{\Sigma}_\varepsilon^{1/2}\|_2$ to reflect different levels of SNR.

First, we demonstrate the practical validity of the confidence regions constructed using Corollaries 2–3 and Theorem 5, for the factors, betas (i.e., factor loadings), and systematic risks, respectively.

- For factors (resp. betas), we construct 95% confidence regions by substituting the asymptotic covariance matrix $\boldsymbol{\Sigma}_{V,t}$ (resp. $\boldsymbol{\Sigma}_{B,i}$) with their consistent estimators as commented after Corollary 3. We define $\widehat{\text{Cov}}_F(t)$ (resp. $\widehat{\text{Cov}}_B(i)$) as the empirical probability that the constructed confidence region covers $\mathbf{f}_t^\top \mathbf{R}_F$ (resp. $\mathbf{b}_i^\top \mathbf{R}_B$) over 200 Monte Carlo trials, where \mathbf{R}_F (resp. \mathbf{R}_B) is the rotation matrix defined in Corollary 2 (resp. Corollary 3).
- For systematic risks, similarly, we define $\widehat{\text{Cov}}_{\ell_2}(i)$ as the empirical probability that the 95% confidence interval $\text{CI}_i^{\mathbf{b}, 0.95}$ constructed via Theorem 5 covers $\|\mathbf{b}_i\|_2^2$ over 200 Monte Carlo trials.

Finally, we compute the mean and standard deviation for $\{\widehat{\text{Cov}}_F(t)\}_{t=1}^T$, $\{\widehat{\text{Cov}}_B(i)\}_{i=1}^N$, and $\{\widehat{\text{Cov}}_{\ell_2}(i)\}_{i=1}^N$, then present these as $\text{Mean}(\widehat{\text{Cov}})$ and $\text{Std}(\widehat{\text{Cov}})$, with the results reported in Table 2. As indicated in Table 2, the coverage probabilities are close to 0.95 according to the mean value, and exhibit stability across different rows of \mathbf{F} and \mathbf{B} as evidenced by low standard deviation. As SNR θ goes down, the slight slippage of the coverage probabilities of the systematic risks reconciles with our comments after Theorem 5 that, $\|\mathbf{b}_i\|_2^2 / (\boldsymbol{\Sigma}_\varepsilon)_{i,i}$, which is close to SNR, reflects the ratio between the bias and the CI width and cannot be too large. These favorable numerical results persist even under low SNR, supporting our inferential theory under the weak factor model.

Table 2: Empirical coverage rates of 95%-CI

$(N, T) = (300, 200)$	Factor		Beta		Systematic Risk	
	Mean($\widehat{\text{Cov}}$)	Std($\widehat{\text{Cov}}$)	Mean($\widehat{\text{Cov}}$)	Std($\widehat{\text{Cov}}$)	Mean($\widehat{\text{Cov}}$)	Std($\widehat{\text{Cov}}$)
SNR θ						
4.5	0.9383	0.0172	0.9325	0.0171	0.9071	0.0400
3.5	0.9298	0.0190	0.9264	0.0184	0.9192	0.0323
2.5	0.9045	0.0210	0.9103	0.0215	0.9244	0.0292

Next, we demonstrate the effectiveness of our test statistics in Theorems 2 and 4 by showing their satisfactory size and power.

- For the null hypothesis $H_0 : \mathbf{v} \in \text{col}(\mathbf{F}_{S,\cdot})$ in the factor specification test of Theorem 2, the time index subset S is chosen as $S = \{\frac{1}{2}T + 1, \frac{1}{2}T + 2, \dots, \frac{1}{2}T + L\}$ with $|S| = L = 12$, and the observed factors \mathbf{v} is set as

$$\mathbf{v} = \mathbf{F}_{S,\cdot} \mathbf{w} + \delta \mathbf{g}.$$

The vector \mathbf{w} is set as $\mathbf{w} = (w_1, w_2, w_3)^\top = (1, 1, 0.5)^\top$, and the parameter δ controls the deviation of the alternatives from the null distribution. The vector \mathbf{g} is constructed as follows: a vector \mathbf{u} is drawn from the standard Gaussian $\mathcal{N}(0, \mathbf{I}_{|S|})$ and the projection residual $\mathbf{u}^\perp = (\mathbf{I}_{|S|} - \mathbf{P}_{\mathbf{F}_{S,\cdot}}) \mathbf{u}$ is computed, where $\mathbf{P}_{\mathbf{F}_{S,\cdot}}$ is the projection matrix on the column space of $\mathbf{F}_{S,\cdot}$. Then, we set $\mathbf{g} = 2\mathbf{u}^\perp / \|\mathbf{u}^\perp\|_2 \cdot \|\mathbf{F}_{S,\cdot}\|_F \|\mathbf{w}\|_2$. The formulation of \mathbf{g} ensures that the signal strength from the column space of $\mathbf{F}_{S,\cdot}$, which is captured by $\|\mathbf{F}_{S,\cdot}\|_F \|\mathbf{w}\|_2$, is balanced with that from the space orthogonal to $\mathbf{F}_{S,\cdot}$. The null hypothesis H_0 should not be rejected when $\delta = 0$, and should be rejected when $\delta > 0$.

- For the null hypothesis $H_0 : \mathbf{b}_i^1 = \mathbf{b}_i^2$ in the structural break test of betas in Theorem 3, the cross-sectional unit is set as $i = 1$, and the time subsets are set as $\Gamma_1 = \{1, 2, \dots, T_1\}$ and $\Gamma_2 = \{T_1 + 1, T_1 + 2, \dots, T_1 + T_2\}$ with $T_1 = T_2 = T/2 = 100$. The beta \mathbf{b}_i^1 on period Γ_1 is generated by the aforementioned procedure, while the beta \mathbf{b}_i^2 on period Γ_2 is set as

$$\mathbf{b}_i^2 = \mathbf{b}_i^1 + \Delta \|\mathbf{b}_i^1\|_2 \mathbf{1},$$

where $\mathbf{1} = (1, 1, 1)^\top$. The null hypothesis H_0 should not be rejected when $\Delta = 0$, and should be rejected when $\Delta > 0$.

- For the null hypothesis $H_0 : \mathbf{b}_i = \mathbf{b}_j$ in the two-sample test of betas in Theorem 4, we set $i = 1$ and study two cases for $j = 2$ and $j = 3$ respectively. According to our simulation setup, the null hypothesis H_0 should not be rejected when $j = 2$, and should be rejected when $j = 3$.

Tables 3–4 report the empirical rejections rates at 5% significance level over 200 Monte Carlo trials. Table 3 (resp. 4) shows that for the test statistics in Theorem 2 (resp. Theorems 3 and 4), the results are favorable, exhibiting appropriate size and power, even under a weak signal setup where the SNR θ is small.

Table 3: Empirical rejection rates at level 5% for the factor tests

$(N, T) = (300, 200)$	Factor specification test					
	SNR θ	$\delta = 0$	$\delta = 0.25$	$\delta = 0.5$	$\delta = 0.75$	$\delta = 1$
	5.5	0.000	0.000	0.745	0.990	1.000
	5.0	0.000	0.000	0.505	0.945	0.995
	4.5	0.000	0.000	0.250	0.895	0.980

Table 4: Empirical rejection rates at level 5% for the beta tests

$(N, T) = (300, 200)$	Structural break test					Two-sample test		
	SNR θ	$\Delta = 0$	$\Delta = 0.25$	$\Delta = 0.5$	$\Delta = 0.75$	$\Delta = 1$	$j = 2$	$j = 3$
	5.5	0.030	0.510	0.990	1.000	1.000	0.045	1.000
	5.0	0.030	0.445	0.970	1.000	1.000	0.050	1.000
	4.5	0.035	0.355	0.945	1.000	1.000	0.050	1.000

6.2 Empirical studies

We analyze the monthly returns data of the S&P 500 constituents from the CRSP database for the period from January 1995 to March 2024. We apply the factor specification test in Theorem 2 and the structural break test for betas in Theorem 3 to the stock returns.

First, we consider the factor specification test. The observed factors we study are the Fama-French three factors: market (MKT), size (SMB), and value (HML), denoted as $\mathbf{v}^{(1)}$, $\mathbf{v}^{(2)}$, and $\mathbf{v}^{(3)}$, respectively. We obtain the time series data for these factors from Kenneth French’s website. We conduct our tests via a rolling window approach, moving a 60-month window $[t+1, t+T]$ ($T = 60$) through the dataset. To mitigate the survival bias, we keep the time series that have no more than 50% missing values in each window, i.e., the number of missing values is less than $T/2$, and fill the missing values by the median of each time series. For each window, the formed data matrix \mathbf{X}^t is an $N \times T$ matrix, where the cross-sectional dimension N varies with t . We assume that \mathbf{X}^t satisfies the factor model as in (2.2), i.e.,

$$\mathbf{X}^t = \mathbf{B}^t(\mathbf{F}^t)^\top + \mathbf{E}^t.$$

Here, the superscript t is added to each matrix to emphasize that we apply the PCA method across different time windows $[t+1, t+T]$. The number of factors is fixed as $r = 3$. We conduct the factor specification test in Theorem 2 to test the null hypothesis

$$\mathbf{v}_{S^t}^{(i)} = \mathbf{F}_{S^t}^t \mathbf{w}^{t,(i)},$$

for the three factors, corresponding to $i = 1, 2, 3$, respectively. The time index subset S^t is set as $S^t = [t+T-L+1, t+T]$ with $|S^t| = L = 12$.

The above procedure implies that, for each 12-month period $S^t = [t+T-L+1, t+T]$, to test if the observed factors are in the column space of the latent factors, we look back and utilize a broader historical data window $[t+1, t+T]$ ($T = 60$) to estimate the latent common factors \mathbf{F}^t . Then we test the null hypothesis for this specific 12-month period S^t , a subset at the end of the whole window $[t+1, t+T]$. Finally, we plot in Figure 1 the test statistics against the time index t for each factor, underscoring the 95% critical value. As highlighted by Fan et al. (2015), this rolling window manner not only utilizes the up-to-date information in the equity universe, but also alleviates the impacts of time-varying betas and sampling biases.




Figure 1: The evolution of the test statistics

In Figure 1, our findings indicate that during the financial crisis of 2007-2009, the null hypothesis that the size factor SMB lies in the latent factors' column space, is rejected at 95% confidence level. This suggests a diminished importance and reduced explanatory power of the size factor SMB for stock return data in this period. Note that the sizes of stocks can still be important or even more important than that in the normal period, but not necessarily captured by the variable SMB. Similar interpretations hold for the value factor HML around 2009. During the COVID period around 2019, both the size factor SMB and the value factor HML exhibit a loss in explanatory power. The spike for the size factor SMB during 2022 is probably due to the war between Russia and Ukraine that started in February 2022, which is an unexpected economic shock to the stock market. Notably, the market portfolio maintains its explanatory strength throughout, indicating its stability and resilience as an explanatory variable, even during distinct economic cycles.

Next, we consider the structural break test for betas for individual stocks. We test whether the betas have changed before and after the three economic recessions covered by the time horizon of our data – the Early 2000s Recession (Mar. 2001–Nov. 2001) due to the dot com bubble, the 2008 Great Recession due to housing bubble and financial crisis (Dec. 2007–Jun. 2009), and the COVID-19 Recession (Feb. 2020–Apr. 2020). Here, the start and the end of each recession are according to the NBER's Business Cycle Dating Committee.

For each recession period $[t_{\text{start}}, t_{\text{end}}]$, we first take the data \mathbf{X}^1 and \mathbf{X}^2 lying in the time window $[t_{\text{start}} - T_1, t_{\text{start}} - 1]$ and $[t_{\text{end}} + 1, t_{\text{end}} + T_2]$, respectively, where $T_1 = T_2 = 60$. Next, we merge the two panels to get $\mathbf{X} = (\mathbf{X}^1, \mathbf{X}^2)$, and fill the missing values in the same manner as in the factor specification test. We assume that \mathbf{X} satisfies the factor model as follows

$$\begin{aligned} x_{i,t} &= \mathbf{f}_t^\top \mathbf{b}_i^1 + \varepsilon_{i,t} & \text{for } t \in \{t_{\text{start}} - T_1, t_{\text{start}} - T_1 + 1, \dots, t_{\text{start}} - 1\}, \\ x_{i,t} &= \mathbf{f}_t^\top \mathbf{b}_i^2 + \varepsilon_{i,t} & \text{for } t \in \{t_{\text{end}} + 1, t_{\text{end}} + 2, \dots, t_{\text{end}} + T_2\}. \end{aligned}$$

The number of factors is chosen by a scree plot of the eigenvalues of \mathbf{X} , which is 4, 3, and 3 for the three recessions, respectively. We test the hypothesis $H_0 : \mathbf{b}_i^1 = \mathbf{b}_i^2 \leftrightarrow H_1 : \mathbf{b}_i^1 \neq \mathbf{b}_i^2$ for each cross-sectional unit i using the test statistic in Theorem 3. To report the test results, we group the stocks into 11 sectors by Global Industrial Classification Standard (GICS), and then count the numbers of stocks in each sector that

reject the null at the 95% confidence level. The test results for the three recessions are illustrated in Figures 2, 3, and 4, respectively.




Figure 2: Test for breaks in betas for Early 2000s Recession (total number of stocks is in bracket)

In Figure 2, we observe that the sectors with the highest number of stocks experiencing structural breaks in betas are Financials, Information Technology, Industrials, and Utilities. The impact on the Information Technology sector is likely due to the dot-com bubble burst, which was one of the triggers of the Early 2000s Recession. Another significant cause of this recession was the 9/11 attacks. Such severe economic shocks are possible reasons why typically stable sectors like Industrials and Utilities were also affected.




Figure 3: Test for breaks in betas for 2008 Great Recession (total number of stocks is in bracket)

In Figure 3, the 2007-2009 financial crisis, marked by the subprime mortgage crisis and the collapse of the United States housing bubble, affected many sectors. The financial sector experienced a strong impact due to direct exposure to mortgage-backed securities and other related financial instruments. The crisis

led to the failure or collapse of many of the United States’ largest financial institutions. Even though our analysis is inevitably influenced by survival bias, as we can only analyze stocks that existed before and after the crisis, we still observe that the financial sector had the most affected stocks. The impact in Real Estate reflects the direct consequences of the housing bubble burst. For sectors related to consumer spending, such as Consumer Discretionary, Consumer Staples, and Health Care, the shocks can be attributed to reduced consumer spending due to increased unemployment and economic uncertainty.




Figure 4: Test for breaks in betas for COVID-19 Recession (total number of stocks is in bracket)

In Figure 4, the economic effects of the pandemic can be seen in many affected sectors, such as Real Estate, Industrials, Health Care, and Financials. The significant changes in Real Estate likely reflect the uncertainties brought about by lockdowns and health crises, resulting in severe fluctuations in property values and rent payments. The impact on the Health Care sector may be tied to the heightened demand for medical services and supplies, alongside volatility in biotechnology investments. Notably, Utilities and Energy showed smaller changes, suggesting relative stability in these sectors despite overall market volatility. During the 2008 Great Recession, these sectors also demonstrated similar stability. This resilience could be attributed to the essential nature of services provided by these sectors, making them less susceptible to economic disruptions.

7 Other related works

The factor model is an important topic in finance and economics. The early econometric studies on factor model can be dated back to [Forni et al. \(2000\)](#); [Stock and Watson \(2002\)](#). Most previous works on factor model assumes that all factors are strong or pervasive, that is, the SNR grows at rate \sqrt{N} (e.g., [Bai and Ng, 2002](#); [Bai, 2003](#); [Fan et al., 2013](#)). When the SNR grows at a rate slower than \sqrt{N} , the model is often called the weak factor model, which has been a popular research topic in recent years. The method for determining the number of factors under the weak factor model has been studied by a few papers (e.g., [Onatski, 2009, 2010](#); [Freyaldenhoven, 2022](#)). Several recent works have pursued the estimation and inference in the weak factor model. To name a few examples, [Bai and Ng \(2023\)](#); [Jiang et al. \(2023\)](#); [Choi and Yuan \(2024\)](#) studied the consistency and asymptotic normality of PC estimators; [Uematsu and Yamagata \(2022a,b\)](#); [Wei and Zhang \(2023\)](#) studied the sparsity-induced weak factor models where the low SNR is due to the sparsity of the factor loading matrix; [Bailey et al. \(2021\)](#) proposed an estimator of factor strength and established its theoretical guarantee; [Onatski \(2012\)](#) showed that the PC estimators are inconsistent in the extreme case (also known as the super-weak factor model) where the SNR is $O(1)$.

The PCA is one of the most popular methods for the factor model, and has been studied in many papers mentioned previously (e.g., [Stock and Watson, 2002](#); [Bai and Ng, 2002](#); [Bai, 2003](#); [Onatski, 2012](#); [Fan et al., 2013](#); [Bai and Ng, 2023](#); [Jiang et al., 2023](#); [Choi and Yuan, 2024](#)). Among the enormous literature, [Bai and Ng \(2023\)](#); [Jiang et al. \(2023\)](#); [Choi and Yuan \(2024\)](#) are the most recent and closest to our paper, since we all focus on the PC estimators, especially on the inference side, under the weak factor model. Besides the theoretical analysis, the variants of PCA under the weak factor model have also been applied to empirical asset pricing, macroeconomic forecasting, and many other important problems in finance (e.g., [Giglio et al., 2021, 2023](#)). The estimation of factor model is closely related to the low-rank matrix denoising in the statistical machine learning community. In recent years, studying the factor model from the view of low-rank matrix denoising has provided lots of exciting findings and understandings (see [Fan et al. \(2021a\)](#); [Yan et al. \(2024\)](#) for comprehensive reviews), and our theory is partly inspired by this low-rank structure of factor model.

From a technical perspective, our estimation procedure is a spectral method, and is related to previous studies of spectral methods on PCA and subspace estimation ([Abbe et al., 2022](#); [Cai et al., 2021](#); [Chen et al., 2021b](#); [Yan et al., 2024](#); [Zhou and Chen, 2023](#)). Our analysis relies on the leave-one-out techniques that have found wide applications in analyzing spectral estimators and nonconvex optimization algorithms ([El Karoui, 2015](#); [Abbe et al., 2020](#); [Ma et al., 2020](#); [Chen et al., 2020, 2023](#)); see the recent monograph [Chen et al. \(2021a\)](#) for more details. In addition, the problem and analysis in this paper is also related to past works on inference for other low-rank models ([Chen et al., 2019](#); [Xia and Yuan, 2021](#); [Chernozhukov et al., 2023](#); [Choi et al., 2023, 2024](#); [Yan and Wainwright, 2024](#)).

8 Conclusions and discussions

In this paper, we establish a novel theory for PCA under the weak factor model, offering significant advancements on the inference of PCA over the existing literature. The weak factor model removes the pervasiveness assumption that requires of SNR growing at a \sqrt{N} rate, where N is the cross-sectional dimension. Our theory covers both estimation and inference of factors and factor loadings. Notably, in the regime $N \asymp T$, where T is the temporal dimension, we show that the asymptotic normality of PC estimators holds as long as the SNR grows faster than a polynomial rate of $\log N$, while the previous work required a polynomial rate of N . The optimality of our theory is in the sense that, the required growth rate of the SNR for consistency aligns with that for asymptotic normality, differing only by some logarithmic factors. Our theory paves the way to design easy-to-implement test statistics for practical applications, e.g., factor specification test, structural break test for betas, and build confidence regions for crucial model parameters, e.g., betas and systematic risks. We validate our statistical methods through extensive Monte Carlo simulations, and conduct empirical studies to find noteworthy correlations between our test results and specific economic cycles.

While the current scope of our theory is considerably wide-ranging, it is possible to further widen it and there are lots of topics that are worth pursuing. For instance, how to extend the theory to the setting where each observed variables have only fourth moment via robustification of the covariance input ([Fan et al., 2021c](#))? How to generalize our inferential theory to the case where the time-serial correlation also appears in the idiosyncratic noise? If the panel data is missing at random, are the PC estimators still valid under the weak factor model, and how to conduct statistical inference in this case? What benefit could our new theory bring to forecasting methods based on the factor-augmented regression? Among many directions, these topics can be investigated in future research.

Acknowledgements

J. Fan is supported in part by the NSF grants DMS-2053832 and DMS-2210833, and ONR grant N00014-22-1-2340. Y. Yan is supported in part by the Norbert Wiener Postdoctoral Fellowship from MIT.

A Proof of Theorem 1: First-order approximations

For ease of exposition, we introduce some additional notation that will be used throughout the proofs. For any $m \times n$ matrix $\mathbf{M} = (M_{i,j})_{1 \leq i \leq m, 1 \leq j \leq n}$, we let $\sigma_i(\mathbf{M})$, $\sigma_{\min}(\mathbf{M})$, and $\sigma_{\max}(\mathbf{M})$ denote the i -th largest, the minimum, and the maximum singular value of \mathbf{M} , respectively. For a symmetric matrix \mathbf{A} , we denote by $\text{tr}(\mathbf{A})$ its trace.

A.1 Some useful lemmas

To prove Theorem 1, we collect some useful lemmas as preparations. We start with a lemma to reveal the close relations between the SVD $\mathbf{U}\mathbf{\Lambda}\mathbf{V}^\top$ and the common component $\mathbf{B}\mathbf{F}^\top$ in the factor model (2.2), where $\mathbf{B} = \bar{\mathbf{U}}\mathbf{\Sigma}$.

Lemma 2. *Suppose that Assumptions 1 and 3 hold. Assuming that*

$$r + \log n \ll T, \quad (\text{A.1})$$

then we have that, there exists a $\sigma(\mathbf{F})$ -measurable event \mathcal{E}_0 with $\mathbb{P}(\mathcal{E}_0) > 1 - O(n^{-2})$, where $\sigma(\mathbf{F})$ is the σ -algebra generated by \mathbf{F} , such that, the following properties hold when \mathcal{E}_0 happens: (i) $\text{rank}(\mathbf{F}) = r$,

$$\left\| \frac{1}{T} \mathbf{F}^\top \mathbf{F} - \mathbf{I}_r \right\|_2 \lesssim \frac{\sqrt{r + \log n}}{\sqrt{T}}, \quad \text{and} \quad \|\mathbf{F}\|_{2,\infty} \lesssim \sqrt{\log n}.$$

(ii) There exists a $\sigma(\mathbf{F})$ -measurable matrix \mathbf{Q} satisfying that $\mathbf{U} = \bar{\mathbf{U}}\mathbf{Q}$ and $\mathbf{Q} \in \mathcal{O}^{r \times r}$ is a rotation matrix, i.e., $\mathbf{Q}\mathbf{Q}^\top = \mathbf{Q}^\top\mathbf{Q} = \mathbf{I}_r$.

(iii) $\lambda_i \asymp \sigma_i$ for $i = 1, 2, \dots, r$.

(iv) There exists a $\sigma(\mathbf{F})$ -measurable matrix \mathbf{J} satisfying that $\mathbf{V} = T^{-1/2}\mathbf{F}\mathbf{J}$ and $\mathbf{J} = \mathbf{\Sigma}\mathbf{Q}\mathbf{\Lambda}^{-1}$ is a $r \times r$ invertible matrix satisfying that $\sigma_i(\mathbf{J}) \asymp 1$ for $i = 1, 2, \dots, r$. Further, it holds

$$\|\mathbf{V}\|_{2,\infty} \lesssim \sqrt{\frac{\log n}{T}}.$$

Proof. Since $\mathbf{F} = (\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_T)^\top$, and $\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_T$ are independent sub-Gaussian random vectors under Assumption 3, we obtain by (4.22) in Vershynin (2017) that,

$$\left\| \frac{1}{T} \mathbf{F}^\top \mathbf{F} - \mathbf{I}_r \right\|_2 \lesssim \sqrt{\frac{r}{T}} + \sqrt{\frac{\log n}{T}} + \frac{r}{T} + \frac{\log n}{T} \lesssim \sqrt{\frac{r + \log n}{T}}, \quad (\text{A.2})$$

with probability at least $1 - O(n^{-2})$, where the last inequality is owing to (A.1) and the fact that $\kappa \geq 1$. Then using the sub-Gaussian property of \mathbf{f}_t and the standard concentration inequality, we have that

$$\|\mathbf{F}\|_{2,\infty} \lesssim \sqrt{\log n}, \quad (\text{A.3})$$

with probability at least $1 - O(n^{-2})$. In particular, we let \mathcal{E}_0 be the event that both (A.2) and (A.3) happen, then we have that $\mathbb{P}(\mathcal{E}_0) > 1 - O(n^{-2})$. In what follows, we show that \mathcal{E}_0 satisfies all the requirements.

When (A.2) happens, since (A.1) implies that $\sqrt{\frac{r + \log n}{T}} \ll 1$, we have that $\left\| \frac{1}{T} \mathbf{F}^\top \mathbf{F} - \mathbf{I}_r \right\|_2 \ll 1$, and thus $\vartheta_i(\frac{1}{T} \mathbf{F}^\top \mathbf{F}) \asymp 1$ for $i = 1, 2, \dots, r$, implying that $\sigma_i(\mathbf{F}) \asymp \sqrt{T}$ for $i = 1, 2, \dots, r$. Since $\sigma_i(\mathbf{F}) \asymp \sqrt{T}$ implies that $\sigma_{\min}(\mathbf{F}) > 0$, we obtain that $\text{rank}(\mathbf{F}) = r$, i.e., \mathbf{F} has full column rank.

By definition, the columns of \mathbf{U} (resp. $\bar{\mathbf{U}}$) are the left singular vectors of $\mathbf{B}\mathbf{F}^\top$ (resp. \mathbf{B}). When (A.2) happens, since $\text{rank}(\mathbf{F}_{S,\cdot}) = r$, the r -dimensional column spaces of \mathbf{U} and $\bar{\mathbf{U}}$ are the same with each other. Then since the columns of both \mathbf{U} and $\bar{\mathbf{U}}$ are orthonormal, we obtain that there exists a rotation matrix $\mathbf{Q} \in \mathcal{O}^{r \times r}$ such that $\mathbf{U} = \bar{\mathbf{U}}\mathbf{Q}$. By construction we have that \mathbf{Q} is $\sigma(\mathbf{F})$ -measurable.

Next, we prove the relations for eigenvalues. By definition, we have that $\{\lambda_i^2\}_{1 \leq i \leq r}$ are the eigenvalues of $(T^{-1/2}\mathbf{B}\mathbf{F}^\top)^\top T^{-1/2}\mathbf{B}\mathbf{F}^\top = T^{-1}\mathbf{F}\mathbf{B}^\top\mathbf{B}\mathbf{F}^\top$, and $\{\sigma_i^2\}_{1 \leq i \leq r}$ are the eigenvalues of $\mathbf{B}^\top\mathbf{B} = \mathbf{\Sigma}^2$. By Theorem A.2 in Braun (2006), we obtain that

$$|\lambda_i^2 - \sigma_i^2| = |\vartheta_i(T^{-1}\mathbf{F}\mathbf{B}^\top\mathbf{B}\mathbf{F}^\top) - \vartheta_i(\mathbf{B}^\top\mathbf{B})| \leq \sigma_i^2 \left\| T^{-1}\mathbf{F}^\top \mathbf{F} - \mathbf{I}_r \right\|_2 \lesssim \sigma_i^2 \sqrt{\frac{r + \log n}{T}} \ll \sigma_i^2,$$

where the last inequality is owing to (A.1). So we obtain that $\lambda_i \asymp \sigma_i$ for $i = 1, 2, \dots, r$.

By the property of SVD, we have that $\mathbf{V} = (T^{-1/2} \mathbf{B} \mathbf{F}^\top)^\top \mathbf{U} \mathbf{\Lambda}^{-1} = T^{-1/2} \mathbf{F} \mathbf{J}$, where \mathbf{J} is given by $\mathbf{J} := \mathbf{B}^\top \mathbf{U} \mathbf{\Lambda}^{-1} = \mathbf{\Sigma} \mathbf{U}^\top \mathbf{U} \mathbf{\Lambda}^{-1} = \mathbf{\Sigma} \mathbf{Q} \mathbf{\Lambda}^{-1}$ and \mathbf{J} is invertible. So we have that $T^{-1/2} \mathbf{F} = \mathbf{V} \mathbf{J}^{-1}$ and thus $T^{-1} \mathbf{F}^\top \mathbf{F} = (\mathbf{J}^{-1})^\top \mathbf{V}^\top \mathbf{V} \mathbf{J}^{-1} = (\mathbf{J}^{-1})^\top \mathbf{J}^{-1}$. Then, we obtain that $\|(\mathbf{J}^{-1})^\top \mathbf{J}^{-1} - \mathbf{I}_r\|_2 \leq \|T^{-1} \mathbf{F}^\top \mathbf{F} - \mathbf{I}_r\|_2 \lesssim T^{-1/2} \sqrt{r + \log n} \ll 1$, and thus we obtain that $\sigma_i(\mathbf{J}) \asymp 1$ for $i = 1, 2, \dots, r$. As a result, we have that

$$\|\mathbf{V}\|_{2,\infty} = \frac{1}{\sqrt{T}} \|\mathbf{F} \mathbf{J}\|_{2,\infty} \leq \frac{1}{\sqrt{T}} \|\mathbf{F}\|_{2,\infty} \|\mathbf{J}\|_2 \stackrel{(i)}{\lesssim} \frac{1}{\sqrt{T}} \|\mathbf{F}\|_{2,\infty} \stackrel{(ii)}{\lesssim} \sqrt{\frac{\log n}{T}},$$

where (i) and (ii) use $\sigma_i(\mathbf{J}) \asymp 1$ and $\|\mathbf{F}\|_{2,\infty} \lesssim \sqrt{\log n}$ in (A.3), respectively. \square

Lemma 2 shows how the SVD $\mathbf{U} \mathbf{\Lambda} \mathbf{V}^\top$ relates with $\mathbf{B} \mathbf{F}^\top$ and constructs a good event \mathcal{E}_0 on which we have nice properties of the quantities in our setup. Lemma 3 below establishes some useful results on the random matrix \mathbf{Z} which characterizes the noise matrix \mathbf{E} in (2.4).

Lemma 3. *Under Assumption 2, for any fixed matrices $\mathbf{A} \in \mathbb{R}^{T \times m}$ and $\mathbf{B} \in \mathbb{R}^{N \times m}$ with $m \leq n$, we have that, with probability at least $1 - O(n^{-2})$,*

$$\|\mathbf{Z}\|_2 \lesssim \sqrt{n}, \quad \|\mathbf{E}\|_2 \lesssim \sqrt{\|\mathbf{\Sigma}_\varepsilon\|_2} \sqrt{n}, \quad (\text{A.4})$$

and

$$\|\mathbf{Z} \mathbf{A}\|_{2,\infty} \lesssim \|\mathbf{A}\|_{\mathbb{F}} \log n, \quad \|\mathbf{Z}^\top \mathbf{B}\|_{2,\infty} \lesssim \|\mathbf{B}\|_{\mathbb{F}} \log n.$$

Proof. It suffices to prove that every inequality holds with probability at least $1 - O(n^{-2})$. Since the entries of \mathbf{Z} are independent and sub-Gaussian under Assumption 2, we obtain by the union bound argument that

$$\mathbb{P}(\mathcal{E}_Z) > 1 - O(n^{-2}) \text{ with } \mathcal{E}_Z := \left\{ \max_{1 \leq k \leq N, 1 \leq l \leq T} |\mathbf{Z}_{k,l}| \leq C \sqrt{\log n} \right\}, \quad (\text{A.5})$$

where $C > 0$ is a constant. Given \mathcal{E}_Z , it follows from (3.9) in Chen et al. (2021a) that, $\|\mathbf{Z}\|_2 \lesssim \sqrt{n} + \log n \lesssim \sqrt{n}$ with probability at least $1 - O(n^{-2})$, i.e., $\mathbb{P}(\mathcal{E}_1 | \mathcal{E}_Z) > 1 - O(n^{-2})$ where $\mathcal{E}_1 := \{\|\mathbf{Z}\|_2 \lesssim \sqrt{n}\}$. So, we obtain

$$\mathbb{P}(\mathcal{E}_1^c) = \mathbb{P}(\mathcal{E}_1^c | \mathcal{E}_Z) \mathbb{P}(\mathcal{E}_Z) + \mathbb{P}(\mathcal{E}_1^c | \mathcal{E}_Z^c) \mathbb{P}(\mathcal{E}_Z^c) \leq \mathbb{P}(\mathcal{E}_1^c | \mathcal{E}_Z^c) + \mathbb{P}(\mathcal{E}_Z^c) \leq O(n^{-2}), \quad (\text{A.6})$$

implying that our desired result $\mathbb{P}(\|\mathbf{Z}\|_2 \lesssim \sqrt{n}) = 1 - \mathbb{P}(\mathcal{E}_1^c) > 1 - O(n^{-2})$. Indeed, the argument for \mathcal{E}_1 in (A.6) adapts to all the events which we will prove to hold with probability at least $1 - O(n^{-2})$. In other words, for any event \mathcal{E} , to prove $\mathbb{P}(\mathcal{E}) > 1 - O(n^{-2})$, it suffices to prove that $\mathbb{P}(\mathcal{E} | \mathcal{E}_Z) > 1 - O(n^{-2})$. So, in what follows, we conduct the proofs conditioning on \mathcal{E}_Z .

It is easy to see that, $\|\mathbf{E}\|_2 \lesssim \sqrt{\|\mathbf{\Sigma}_\varepsilon\|_2} \sqrt{n}$ follows from

$$\|\mathbf{E}\|_2 = \|\mathbf{\Sigma}_\varepsilon^{1/2} \mathbf{Z}\|_2 \leq \|\mathbf{\Sigma}_\varepsilon^{1/2}\|_2 \|\mathbf{Z}\|_2 = \sqrt{\|\mathbf{\Sigma}_\varepsilon\|_2} \|\mathbf{Z}\|_2 \lesssim \sqrt{\|\mathbf{\Sigma}_\varepsilon\|_2} \sqrt{n}.$$

Then, we prove that

$$\mathbb{P}(\mathcal{E}_A | \mathcal{E}_Z) > 1 - O(n^{-2}) \text{ with } \mathcal{E}_A := \left\{ \|\mathbf{Z} \mathbf{A}\|_{2,\infty} \lesssim \|\mathbf{A}\|_{\mathbb{F}} \log n \right\}. \quad (\text{A.7})$$

To prove (A.7), we note that, for $k = 1, 2, \dots, N$, $\mathbf{Z}_{k,\cdot} \mathbf{A} = \sum_{l=1}^T \mathbf{Z}_{k,l} \mathbf{A}_{l,\cdot}$ is a sum of independent mean zero random vectors. Given \mathcal{E}_Z , it holds $\|\mathbf{Z}_{k,l} \mathbf{A}_{l,\cdot}\|_2 \leq C \sqrt{\log n} \|\mathbf{A}_{l,\cdot}\|_2$ for $l = 1, 2, \dots, T$. Then, using the matrix Hoeffding inequality (Tropp, 2012, Theorem 1.3), we obtain that

$$\mathbb{P}(\|\mathbf{Z}_{k,\cdot} \mathbf{A}\|_2 \geq \delta | \mathcal{E}_Z) \leq 2m \exp\left(-\frac{\delta^2}{8\sigma^2}\right),$$

where $\sigma = [\sum_{l=1}^T (C \sqrt{\log n} \|\mathbf{A}_{l,\cdot}\|_2)^2]^{1/2} = C \sqrt{\log n} \|\mathbf{A}\|_{\mathbb{F}}$. Then we obtain by the union bound argument that $\mathbb{P}(\|\mathbf{Z} \mathbf{A}\|_{2,\infty} \leq \delta | \mathcal{E}_Z) > 1 - 2Nm \exp(-\frac{\delta^2}{8\sigma^2})$. Letting $\delta = 100\sigma \sqrt{\log n}$, we obtain that $\mathbb{P}(\|\mathbf{Z} \mathbf{A}\|_{2,\infty} \leq \delta | \mathcal{E}_Z) > 1 - O(n^{-2})$, which implying (A.7). The proof for $\|\mathbf{Z}^\top \mathbf{B}\|_{2,\infty} \lesssim \|\mathbf{B}\|_{\mathbb{F}} \log n$ follows from the same manner, so we omit the details for the sake of brevity. \square

Next, we consider the truncated rank- r SVD $\widehat{\mathbf{U}}\widehat{\boldsymbol{\Sigma}}\widehat{\mathbf{V}}^\top$ of $\frac{1}{\sqrt{T}}\mathbf{X} = \frac{1}{\sqrt{T}}(\mathbf{B}\mathbf{F}^\top + \mathbf{E})$. We define

$$\mathbf{H}_U := \widehat{\mathbf{U}}^\top \mathbf{U} \text{ and } \mathbf{H}_V := \widehat{\mathbf{V}}^\top \mathbf{V}.$$

Lemma 4 below give the perturbation bounds for the singular spaces, i.e., the column spaces of \mathbf{U} and \mathbf{V} , under the spectral norm, and some basic facts on $\widehat{\boldsymbol{\Sigma}}$, \mathbf{H}_U , and \mathbf{H}_V .

Lemma 4. *Suppose that Assumptions 1, 2, and 3 hold. For simplicity of notations, we define*

$$\rho := \frac{\sqrt{\|\boldsymbol{\Sigma}_\varepsilon\|_2} \sqrt{n}}{\sigma_r \sqrt{T}} = \frac{\sqrt{n}}{\theta \sqrt{T}}. \quad (\text{A.8})$$

Assuming that

$$\rho \ll 1, \quad (\text{A.9})$$

then we have that, with probability at least $1 - O(n^{-2})$,

$$\max \left\{ \|\widehat{\mathbf{U}}\mathbf{R}_U - \mathbf{U}\|_2, \|\widehat{\mathbf{V}}\mathbf{R}_V - \mathbf{V}\|_2, \sqrt{\|\mathbf{H}_U - \mathbf{R}_U\|_2}, \sqrt{\|\mathbf{H}_V - \mathbf{R}_V\|_2} \right\} \lesssim \rho,$$

and $\max\{\|\widehat{\mathbf{U}}\mathbf{R}_U - \mathbf{U}\|_F, \|\widehat{\mathbf{V}}\mathbf{R}_V - \mathbf{V}\|_F\} \lesssim \rho\sqrt{r}$, as well as

$$\widehat{\sigma}_i \asymp \sigma_i, \quad \sigma_i(\mathbf{H}_U) \asymp 1, \quad \sigma_i(\mathbf{H}_V) \asymp 1, \quad \text{for } i = 1, 2, \dots, r.$$

Also, we have that

$$\max \left\{ \|\widehat{\boldsymbol{\Sigma}}(\mathbf{H}_U - \mathbf{R}_U)\|_2, \|\widehat{\boldsymbol{\Sigma}}(\mathbf{H}_V - \mathbf{R}_V)\|_2 \right\} \lesssim \sigma_r \rho^2.$$

Proof. Using the argument as discussed after (A.6), for any event \mathcal{E} , to prove $\mathbb{P}(\mathcal{E}) > 1 - O(n^{-2})$, it suffices to prove that $\mathbb{P}(\mathcal{E}|\mathcal{E}_Z \cap \mathcal{E}_0) > 1 - O(n^{-2})$. Here, the event \mathcal{E}_Z is defined in (A.5) and the event \mathcal{E}_0 is defined in Lemma 2. To see this, we note that, as long as $\mathbb{P}(\mathcal{E}|\mathcal{E}_Z \cap \mathcal{E}_0) > 1 - O(n^{-2})$, it holds

$$\begin{aligned} \mathbb{P}(\mathcal{E}^c) &= \mathbb{P}(\mathcal{E}^c|\mathcal{E}_Z \cap \mathcal{E}_0) \mathbb{P}(\mathcal{E}_Z \cap \mathcal{E}_0) + \mathbb{P}(\mathcal{E}^c|(\mathcal{E}_Z \cap \mathcal{E}_0)^c) \mathbb{P}((\mathcal{E}_Z \cap \mathcal{E}_0)^c) \\ &\leq \mathbb{P}(\mathcal{E}^c|\mathcal{E}_Z \cap \mathcal{E}_0) + \mathbb{P}((\mathcal{E}_Z \cap \mathcal{E}_0)^c) \leq O(n^{-2}) + \mathbb{P}(\mathcal{E}_Z^c \cup \mathcal{E}_0^c) \\ &\leq O(n^{-2}) + \mathbb{P}(\mathcal{E}_Z^c) + \mathbb{P}(\mathcal{E}_0^c) \leq O(n^{-2}). \end{aligned} \quad (\text{A.10})$$

So, in what follows, we conduct the proofs conditioning on $\mathcal{E}_Z \cap \mathcal{E}_0$.

Recall that $\widehat{\mathbf{U}}\widehat{\boldsymbol{\Sigma}}\widehat{\mathbf{V}}^\top$ is the truncated rank- r SVD of $\frac{1}{\sqrt{T}}\mathbf{X}$, and $\mathbf{U}\boldsymbol{\Lambda}\mathbf{V}^\top$ is the SVD of $\frac{1}{\sqrt{T}}\mathbf{B}\mathbf{F}^\top$. Note that $\frac{1}{\sqrt{T}}\mathbf{X} - \frac{1}{\sqrt{T}}\mathbf{B}\mathbf{F}^\top = \frac{1}{\sqrt{T}}\mathbf{E}$, and it follows from (A.4) in Lemma 3 that

$$\frac{1}{\sqrt{T}} \|\mathbf{E}\|_2 \lesssim \frac{1}{\sqrt{T}} \sqrt{\|\boldsymbol{\Sigma}_\varepsilon\|_2} \sqrt{n} \ll \sigma_r,$$

where the last inequality is owing to (A.9). So we obtain $\frac{1}{\sqrt{T}} \|\mathbf{E}\|_2 \ll \lambda_r$, since $\lambda_i \asymp \sigma_i$ for $i = 1, 2, \dots, r$ according to Lemma 2. We write the full SVD of $T^{-1/2}\mathbf{X}$ as

$$\frac{1}{\sqrt{T}}\mathbf{X} = \begin{bmatrix} \widehat{\mathbf{U}} & \widehat{\mathbf{U}}_\perp \end{bmatrix} \begin{bmatrix} \widehat{\boldsymbol{\Sigma}} & 0 \\ 0 & \widehat{\boldsymbol{\Sigma}}_\perp \end{bmatrix} \begin{bmatrix} \widehat{\mathbf{V}}^\top \\ (\widehat{\mathbf{V}}_\perp)^\top \end{bmatrix},$$

where $\widehat{\mathbf{U}}_\perp \in \mathbb{R}^{N \times (N-r)}$ and $\widehat{\mathbf{V}}_\perp \in \mathbb{R}^{T \times (T-r)}$ are the orthogonal complements of $\widehat{\mathbf{U}}$ and $\widehat{\mathbf{V}}$, and $\widehat{\boldsymbol{\Sigma}}_\perp \in \mathbb{R}^{(N-r) \times (T-r)}$ contains the smaller singular values and $\widehat{\boldsymbol{\Sigma}}_\perp$ is not necessarily a diagonal matrix. Then by Wedin's $\sin \Theta$ theorem (Chen et al., 2021a, (2.26a)–(2.26b)), we have that

$$\|\widehat{\mathbf{U}}\mathbf{R}_U - \mathbf{U}\|_2 \lesssim \|\widehat{\mathbf{U}}\widehat{\mathbf{U}}^\top - \mathbf{U}\mathbf{U}^\top\|_2 \lesssim \|\mathbf{U}^\top \widehat{\mathbf{U}}_\perp\|_2 \lesssim \frac{\frac{1}{\sqrt{T}} \|\mathbf{E}\|_2}{\lambda_r - \lambda_{r+1}} \lesssim \frac{\frac{1}{\sqrt{T}} \sqrt{\|\boldsymbol{\Sigma}_\varepsilon\|_2} \sqrt{n}}{\sigma_r} = \rho, \quad (\text{A.11})$$

and $\|\widehat{\mathbf{V}}\mathbf{R}_V - \mathbf{V}\|_2 \lesssim \|\mathbf{V}^\top \widehat{\mathbf{V}}_\perp\|_2 \lesssim \rho$, and similarly $\max\{\|\widehat{\mathbf{U}}\mathbf{R}_U - \mathbf{U}\|_F, \|\widehat{\mathbf{V}}\mathbf{R}_V - \mathbf{V}\|_F\} \lesssim \rho\sqrt{r}$, where we use the fact that $\lambda_r \asymp \sigma_r$ and $\lambda_{r+1} = 0$, since λ_i is the i -th largest singular value of $\frac{1}{\sqrt{T}}\mathbf{B}\mathbf{F}^\top$ and $\text{rank}(\mathbf{B}\mathbf{F}^\top) = r$. Next, for \mathbf{H}_U , using the arguments in Section C.3.1 of Yan et al. (2024) based on Wedin's $\sin \Theta$ theorem, we obtain that $\|\mathbf{H}_U - \mathbf{R}_U\|_2 \lesssim \|\widehat{\mathbf{U}}\mathbf{R}_U - \mathbf{U}\|_2^2 \lesssim \rho^2 \ll 1$. Then, since $\mathbf{R}_U \in \mathcal{O}^{r \times r}$ implies that $\sigma_i(\mathbf{R}_U) = 1$ for $i = 1, 2, \dots, r$, we obtain $\sigma_i(\mathbf{H}_U) \asymp 1$ for $i = 1, 2, \dots, r$. The results for \mathbf{H}_V can be proven in the same manner. Also, we obtain by Weyl's inequality (Chen et al., 2021a, Lemma 2.2) that, for $i = 1, 2, \dots, r$, $|\widehat{\sigma}_i - \lambda_i| \leq \frac{1}{\sqrt{T}} \|\mathbf{E}\|_2 \ll \sigma_r$, implying that $\widehat{\sigma}_i \asymp \sigma_i$ for $i = 1, 2, \dots, r$.

We now prove the results for $\|\widehat{\Sigma}(\mathbf{H}_U - \mathbf{R}_U)\|_2$ and $\|\widehat{\Sigma}(\mathbf{H}_V - \mathbf{R}_V)\|_2$. Similar to the arguments for proving (F.33) in Yan and Wainwright (2024), we have that

$$\|\widehat{\Sigma}(\mathbf{H}_U - \mathbf{R}_U)\|_2 \lesssim \rho \|\widehat{\Sigma}\widehat{\mathbf{U}}\mathbf{U}_\perp\|_2,$$

and $\widehat{\Sigma}\widehat{\mathbf{U}}\mathbf{U}_\perp = T^{-1/2}\widehat{\mathbf{V}}^\top \mathbf{E}^\top \mathbf{U}_\perp$, where $\mathbf{U}_\perp \in \mathbb{R}^{N \times (N-r)}$ is the orthogonal complement of \mathbf{U} . Next, we have that

$$\begin{aligned} \|\widehat{\Sigma}\widehat{\mathbf{U}}\mathbf{U}_\perp\|_2 &= \frac{1}{\sqrt{T}} \|(\mathbf{U}_\perp)^\top \mathbf{E} \widehat{\mathbf{V}}\|_2 \leq \frac{1}{\sqrt{T}} \|(\mathbf{U}_\perp)^\top \Sigma_\varepsilon^{1/2} \mathbf{Z} \mathbf{V}\|_2 + \frac{1}{\sqrt{T}} \|(\mathbf{U}_\perp)^\top \mathbf{E} (\widehat{\mathbf{V}}\mathbf{R}_V - \mathbf{V})\|_2 \\ &\stackrel{(i)}{\lesssim} \frac{1}{\sqrt{T}} \|\widehat{\mathbf{V}}^\top \Sigma_\varepsilon^{1/2}\|_2 \|\mathbf{U}_\perp\|_2 \sqrt{r + (N-r) + \log n} + \frac{1}{\sqrt{T}} \|\mathbf{E}\|_2 \|(\widehat{\mathbf{V}}\mathbf{R}_V - \mathbf{V})\|_2 \\ &\stackrel{(ii)}{\lesssim} \frac{1}{\sqrt{T}} \sqrt{\|\Sigma_\varepsilon\|_2} \sqrt{N + \log n} + \frac{1}{\sqrt{T}} \sqrt{\|\Sigma_\varepsilon\|_2} \sqrt{n} \rho \stackrel{(iii)}{\lesssim} \frac{1}{\sqrt{T}} \sqrt{\|\Sigma_\varepsilon\|_2} \sqrt{n}, \end{aligned}$$

where (i) uses (G.3) in Lemma 19 of Yan and Wainwright (2024) and the fact that $\text{rank}(\mathbf{U}_\perp) = N - r$, (ii) follows from (A.4) and $\|\widehat{\mathbf{V}}\mathbf{R}_V - \mathbf{V}\|_2 \lesssim \rho$, and (iii) is because $\rho \ll 1$ as we assumed. So we conclude that

$$\|\widehat{\Sigma}(\mathbf{H}_U - \mathbf{R}_U)\|_2 \lesssim \rho \|\widehat{\Sigma}\widehat{\mathbf{U}}\mathbf{U}_\perp\|_2 \lesssim \rho \frac{1}{\sqrt{T}} \sqrt{\|\Sigma_\varepsilon\|_2} \sqrt{n} = \sigma_r \rho^2.$$

The results for $\|\widehat{\Sigma}(\mathbf{H}_V - \mathbf{R}_V)\|_2$ can be proven in the same manner. \square

Then we establish the close relation between $\widehat{\Sigma}$ and $\mathbf{\Lambda}$. In particular, when we prove all of the following results, we will use the results that have been proven to hold with probabilities at least $1 - O(n^{-2})$ in the previous lemmas. When we use these previous results, it is equivalent to add a new event in $\mathcal{E}_Z \cap \mathcal{E}_0$ and then repeat the argument in (A.10). Since all of our results are to be proven to hold with probabilities at least $1 - O(n^{-2})$, the argument in (A.10) always works and in what follows we will use it many times without mentioning it.

Lemma 5. *Suppose that (A.9) and Assumptions 1, 2, and 3 hold. We have that, with probability at least $1 - O(n^{-2})$,*

$$\begin{aligned} \|(\mathbf{H}_U)^\top \widehat{\Sigma} \mathbf{H}_V - \mathbf{\Lambda}\|_2 &\lesssim \sigma_r \rho^3 + \sigma_r \rho \sqrt{\frac{1}{n}(r + \log n)}, \\ \|(\mathbf{R}_U)^\top \widehat{\Sigma} \mathbf{R}_V - \mathbf{\Lambda}\|_2 &\lesssim \sigma_r \rho^2 + \sigma_r \rho \sqrt{\frac{1}{n}(r + \log n)}. \end{aligned}$$

Proof. Note that

$$\|(\mathbf{H}_U)^\top \widehat{\Sigma} \mathbf{H}_V - \mathbf{\Lambda}\|_2 \leq \underbrace{\|(\mathbf{H}_U)^\top \widehat{\Sigma} \mathbf{H}_V - \mathbf{U}^\top (T^{-1/2} \mathbf{X}) \mathbf{V}\|_2}_{=:\varepsilon_1} + \underbrace{\|\mathbf{U}^\top (T^{-1/2} \mathbf{X}) \mathbf{V} - \mathbf{\Lambda}\|_2}_{=:\varepsilon_2}.$$

For ε_1 , using the full SVD of $T^{-1/2} \mathbf{X}$ in the proof of Lemma 4, we have that

$$(\mathbf{H}_U)^\top \widehat{\Sigma} \mathbf{H}_V - \mathbf{U}^\top (T^{-1/2} \mathbf{X}) \mathbf{V} = \mathbf{U}^\top \widehat{\mathbf{U}} \widehat{\Sigma} \widehat{\mathbf{V}}^\top \mathbf{V} - \mathbf{U}^\top (T^{-1/2} \mathbf{X}) \mathbf{V} = -\mathbf{U}^\top \widehat{\mathbf{U}}_\perp \widehat{\Sigma}_\perp (\widehat{\mathbf{V}}_\perp)^\top \mathbf{V}.$$

By (A.4) in Lemma 3 and Weyl's inequality (Chen et al., 2021a, Lemma 2.2), we have

$$\|\widehat{\Sigma}_\perp\|_2 \leq \sigma_{r+1} \left(\frac{1}{\sqrt{T}} \mathbf{B} \mathbf{F}^\top \right) + \frac{1}{\sqrt{T}} \|\mathbf{E}\|_2 = \frac{1}{\sqrt{T}} \|\mathbf{E}\|_2 \lesssim \frac{1}{\sqrt{T}} \sqrt{\|\Sigma_\varepsilon\|_2} \sqrt{n}.$$

By Lemmas 2.5-2.6 in Chen et al. (2021a) and Lemma 4, we have that $\|\mathbf{U}^\top \widehat{\mathbf{U}}_\perp\|_2 \lesssim \|\widehat{\mathbf{U}} \mathbf{R}_U - \mathbf{U}\|_2 \lesssim \rho$ and $\|\mathbf{V}^\top \widehat{\mathbf{V}}_\perp\|_2 \lesssim \|\widehat{\mathbf{V}} \mathbf{R}_V - \mathbf{V}\|_2 \lesssim \rho$. So we obtain

$$\varepsilon_1 = \|\mathbf{U}^\top \widehat{\mathbf{U}}_\perp \widehat{\Sigma}_\perp (\widehat{\mathbf{V}}_\perp)^\top \mathbf{V}\|_2 \leq \|\mathbf{U}^\top \widehat{\mathbf{U}}_\perp\|_2 \|\widehat{\Sigma}_\perp\|_2 \|(\widehat{\mathbf{V}}_\perp)^\top \mathbf{V}\|_2 \lesssim \sigma_r \rho^3.$$

For ε_2 , we start by writing

$$\mathbf{U}^\top (T^{-1/2} \mathbf{X}) \mathbf{V} - \mathbf{A} = \mathbf{U}^\top (T^{-1/2} \mathbf{X}) \mathbf{V} - \mathbf{U}^\top (T^{-1/2} \mathbf{B} \mathbf{F}^\top) \mathbf{V} = \mathbf{U}^\top (T^{-1/2} \mathbf{E}) \mathbf{V} = T^{-1/2} \mathbf{U}^\top \Sigma_\varepsilon^{1/2} \mathbf{Z} \mathbf{V}.$$

Using (G.3) in Lemma 19 of Yan and Wainwright (2024), we obtain that

$$\varepsilon_2 \lesssim \frac{1}{\sqrt{T}} \|\mathbf{U}^\top \Sigma_\varepsilon^{1/2}\|_2 \|\mathbf{V}\|_2 \sqrt{r + r + \log n} \lesssim \frac{1}{\sqrt{T}} \sqrt{\|\Sigma_\varepsilon\|_2} \sqrt{r + \log n} = \sigma_r \rho \sqrt{\frac{1}{n} (r + \log n)}.$$

Then, we conclude that

$$\|(\mathbf{H}_U)^\top \widehat{\Sigma} \mathbf{H}_V - \mathbf{A}\|_2 \leq \varepsilon_1 + \varepsilon_2 \lesssim \sigma_r \rho^3 + \sigma_r \rho \sqrt{\frac{1}{n} (r + \log n)}.$$

For $(\mathbf{R}_U)^\top \widehat{\Sigma} \mathbf{R}_V - \mathbf{A}$, we note that

$$\begin{aligned} \|(\mathbf{R}_U)^\top \widehat{\Sigma} \mathbf{R}_V - (\mathbf{H}_U)^\top \widehat{\Sigma} \mathbf{H}_V\|_2 &\leq \|(\mathbf{R}_U)^\top \widehat{\Sigma} (\mathbf{R}_V - \mathbf{H}_V)\|_2 + \|(\mathbf{R}_U - \mathbf{H}_U)^\top \widehat{\Sigma} \mathbf{H}_V\|_2 \\ &\leq \|\mathbf{R}_U\|_2 \|\widehat{\Sigma} (\mathbf{H}_V - \mathbf{R}_V)\|_2 + \|\widehat{\Sigma} (\mathbf{H}_U - \mathbf{R}_U)\|_2 \|\mathbf{H}_V\|_2 \\ &\lesssim \sigma_r \rho^2, \end{aligned}$$

where the last inequality is owing to Lemma 4. So we obtain

$$\begin{aligned} \|(\mathbf{R}_U)^\top \widehat{\Sigma} \mathbf{R}_V - \mathbf{A}\|_2 &\leq \|(\mathbf{R}_U)^\top \widehat{\Sigma} \mathbf{R}_V - (\mathbf{H}_U)^\top \widehat{\Sigma} \mathbf{H}_V\|_2 + \|(\mathbf{H}_U)^\top \widehat{\Sigma} \mathbf{H}_V - \mathbf{A}\|_2 \\ &\lesssim \sigma_r \rho^2 + \sigma_r \rho^3 + \sigma_r \rho \sqrt{\frac{1}{n} (r + \log n)} \\ &\lesssim \sigma_r \rho^2 + \sigma_r \rho \sqrt{\frac{1}{n} (r + \log n)}, \end{aligned}$$

where the last inequality is because $\rho \ll 1$ in (A.9). □

We now prove an important lemma where the leave-one-out (LOO) technique plays a key role.

Lemma 6. *Suppose that Assumptions 1, 2, and 3 hold. For simplicity of notations, we define*

$$\omega_k := \frac{\|(\Sigma_\varepsilon^{1/2})_{k,\cdot}\|_1 \sqrt{n}}{\sigma_r \sqrt{T}} = \frac{\sqrt{n}}{\vartheta_k \sqrt{T}} \quad \text{for } k = 1, 2, \dots, N, \quad (\text{A.12})$$

as well as

$$\omega := \max_{1 \leq k \leq N} \omega_k = \frac{\|\Sigma_\varepsilon^{1/2}\|_1 \sqrt{n}}{\sigma_r \sqrt{T}} = \frac{\sqrt{n}}{\vartheta \sqrt{T}},$$

(i) Assuming that (3.2) holds, i.e.,

$$\rho \sqrt{\log n} = \frac{\sqrt{\|\Sigma_\varepsilon\|_2} \sqrt{n \log n}}{\sigma_r \sqrt{T}} \ll 1,$$

then we have that, for $l = 1, 2, \dots, T$, with probability at least $1 - O(n^{-2})$,

$$\begin{aligned} & \left\| (\widehat{\mathbf{V}} \widehat{\boldsymbol{\Sigma}} \mathbf{H}_U - (T^{-1/2} \mathbf{X})^\top \mathbf{U})_{l,\cdot} \right\|_2 \\ & \lesssim \sigma_r \rho^2 \left(\sqrt{\frac{r}{n}} \log^{3/2} n + \log n \|\mathbf{V}_{l,\cdot}\|_2 + \log n \left\| (\widehat{\mathbf{V}} \mathbf{H}_V - \mathbf{V})_{l,\cdot} \right\|_2 \right). \end{aligned}$$

(ii) Assuming that (3.1) holds, i.e.,

$$\omega \sqrt{\log n} = \frac{\left\| \boldsymbol{\Sigma}_\varepsilon^{1/2} \right\|_1 \sqrt{n \log n}}{\sigma_r \sqrt{T}} \ll 1,$$

then we have that, for $k = 1, 2, \dots, N$, with probability at least $1 - O(n^{-2})$,

$$\begin{aligned} & \left\| (\widehat{\mathbf{U}} \widehat{\boldsymbol{\Sigma}} \mathbf{H}_V - (T^{-1/2} \mathbf{X}) \mathbf{V})_{k,\cdot} \right\|_2 \\ & \lesssim \sigma_r \left(\omega_k \rho \sqrt{\frac{r}{n}} \log^{3/2} n + \rho^2 \|\mathbf{U}_{k,\cdot}\|_2 + \omega_k \omega \log n \left(\|\mathbf{U}\|_{2,\infty} + \|\widehat{\mathbf{U}} \mathbf{H}_U - \mathbf{U}\|_{2,\infty} \right) \right), \end{aligned}$$

where $\omega_k = \left\| (\boldsymbol{\Sigma}_\varepsilon^{1/2})_{k,\cdot} \right\|_1 \sqrt{n} / (\sigma_r \sqrt{T})$ is defined in (A.12).

Proof. The proof consists of four steps. In the first (resp, third) step, we prove the upper bound for $\left\| (\widehat{\mathbf{V}} \widehat{\boldsymbol{\Sigma}} \mathbf{H}_U - (T^{-1/2} \mathbf{X})^\top \mathbf{U})_{l,\cdot} \right\|_2$ (resp. $\left\| (\widehat{\mathbf{U}} \widehat{\boldsymbol{\Sigma}} \mathbf{H}_V - (T^{-1/2} \mathbf{X}) \mathbf{V})_{k,\cdot} \right\|_2$), and in the second (resp. fourth) step, we add the details of the proof for an important inequality used in the proof for $\left\| (\widehat{\mathbf{V}} \widehat{\boldsymbol{\Sigma}} \mathbf{H}_U - (T^{-1/2} \mathbf{X})^\top \mathbf{U})_{l,\cdot} \right\|_2$ (resp. $\left\| (\widehat{\mathbf{U}} \widehat{\boldsymbol{\Sigma}} \mathbf{H}_V - (T^{-1/2} \mathbf{X}) \mathbf{V})_{k,\cdot} \right\|_2$).

Step 1 – By calculations, we have that, for $l = 1, 2, \dots, T$,

$$\left(\widehat{\mathbf{V}} \widehat{\boldsymbol{\Sigma}} \mathbf{H}_U - (T^{-1/2} \mathbf{X})^\top \mathbf{U} \right)_{l,\cdot} = \underbrace{\mathbf{V}_{l,\cdot} \boldsymbol{\Lambda} \mathbf{U}^\top (\widehat{\mathbf{U}} \mathbf{H}_U - \mathbf{U})}_{=:\alpha_1} + \underbrace{T^{-1/2} (\mathbf{Z}^\top)_{l,\cdot} \boldsymbol{\Sigma}_\varepsilon^{1/2} (\widehat{\mathbf{U}} \mathbf{H}_U - \mathbf{U})}_{=:\alpha_2}.$$

For α_1 , using the full SVD of $T^{-1/2} \mathbf{X}$ in the proof of Lemma 4, we have that, similar to Step 1 in Section F.2.3 of Yan and Wainwright (2024), it holds

$$\boldsymbol{\Lambda} \mathbf{U}^\top (\widehat{\mathbf{U}} \mathbf{H}_U - \mathbf{U}) = \mathbf{V}^\top \widehat{\mathbf{V}}_\perp (\widehat{\boldsymbol{\Sigma}}_\perp)^\top (\widehat{\mathbf{U}}_\perp)^\top \mathbf{U} - T^{-1/2} \mathbf{V}^\top \mathbf{E}^\top (\widehat{\mathbf{U}} \widehat{\mathbf{U}}^\top - \mathbf{I}_N) \mathbf{U}.$$

Using (A.11), the spectral norm of the first matrix can be bounded by

$$\left\| \mathbf{V}^\top \widehat{\mathbf{V}}_\perp (\widehat{\boldsymbol{\Sigma}}_\perp)^\top (\widehat{\mathbf{U}}_\perp)^\top \mathbf{U} \right\|_2 \leq \left\| \mathbf{V}^\top \widehat{\mathbf{V}}_\perp \right\|_2 \|\widehat{\boldsymbol{\Sigma}}_\perp\|_2 \|\mathbf{U}^\top \widehat{\mathbf{U}}_\perp\|_2 \lesssim \sigma_r \rho^3,$$

where we use the fact that $\|\widehat{\boldsymbol{\Sigma}}_\perp\|_2 = \hat{\sigma}_r \lesssim T^{-1/2} \|\mathbf{E}\|_2 \lesssim \sigma_r \rho$. The spectral norm of the second matrix can be bounded by

$$\begin{aligned} \left\| T^{-1/2} \mathbf{V}^\top \mathbf{E}^\top (\widehat{\mathbf{U}} \widehat{\mathbf{U}}^\top - \mathbf{I}_N) \mathbf{U} \right\|_2 & \leq \frac{1}{\sqrt{T}} \|\mathbf{E} \mathbf{V}\|_2 \left\| (\widehat{\mathbf{U}} \widehat{\mathbf{U}}^\top - \mathbf{I}_N) \mathbf{U} \right\|_2 = \frac{1}{\sqrt{T}} \|\boldsymbol{\Sigma}_\varepsilon^{1/2} \mathbf{Z} \mathbf{V}\|_2 \|\widehat{\mathbf{U}} \mathbf{H}_U - \mathbf{U}\|_2 \\ & \stackrel{(i)}{\lesssim} \frac{1}{\sqrt{T}} \sqrt{\|\boldsymbol{\Sigma}_\varepsilon\|_2} \sqrt{N + \log n} \cdot \rho \lesssim \sigma_r \rho^2, \end{aligned}$$

where (i) uses (G.3) in Lemma 19 of Yan and Wainwright (2024) and the fact that $\widehat{\mathbf{U}} \mathbf{H}_U - \mathbf{U} = \widehat{\mathbf{U}} \widehat{\mathbf{U}}^\top \mathbf{U} - \mathbf{U} = (\widehat{\mathbf{U}} \widehat{\mathbf{U}}^\top - \mathbf{U} \mathbf{U}^\top) \mathbf{U}$ implying $\|\widehat{\mathbf{U}} \mathbf{H}_U - \mathbf{U}\|_2 \leq \|\widehat{\mathbf{U}} \widehat{\mathbf{U}}^\top - \mathbf{U} \mathbf{U}^\top\|_2 \lesssim \rho$ according to (A.11). So, we obtain that

$$\|\alpha_1\|_2 \leq \|\mathbf{V}_{l,\cdot}\|_2 (\sigma_r \rho^3 + \sigma_r \rho^2) \lesssim \sigma_r \rho^2 \|\mathbf{V}_{l,\cdot}\|_2. \quad (\text{A.13})$$

For α_2 , we use the leave-one-out (LOO) technique. We define $\mathbf{Z}^{\{l\}}$ as the matrix obtained by replacing the l th column of \mathbf{Z} with a zero vector, and define $\mathbf{X}^{\{l\}} := \mathbf{B} \mathbf{F}^\top + \boldsymbol{\Sigma}_\varepsilon^{1/2} \mathbf{Z}^{\{l\}}$. Then, we define $\widehat{\mathbf{U}}^{\{l\}}, \widehat{\mathbf{V}}^{\{l\}}$,

$\mathbf{H}_U^{\{l\}}$, and $\mathbf{H}_V^{\{l\}}$ w.r.t. $T^{-1/2}\mathbf{X}^{\{l\}}$, in the same way as how we define $\widehat{\mathbf{U}}$, $\widehat{\mathbf{V}}$, \mathbf{H}_U , and \mathbf{H}_V w.r.t. $T^{-1/2}\mathbf{X}$. We have that

$$\alpha_2 = \underbrace{T^{-1/2}(\mathbf{Z}^\top)_{l,\cdot}\boldsymbol{\Sigma}_\varepsilon^{1/2}(\widehat{\mathbf{U}}^{\{l\}}\mathbf{H}_U^{\{l\}} - \mathbf{U})}_{=:\gamma_1} + \underbrace{T^{-1/2}(\mathbf{Z}^\top)_{l,\cdot}\boldsymbol{\Sigma}_\varepsilon^{1/2}(\widehat{\mathbf{U}}\mathbf{H}_U - \widehat{\mathbf{U}}^{\{l\}}\mathbf{H}_U^{\{l\}})}_{=:\gamma_2}. \quad (\text{A.14})$$

For γ_1 , since $(\widehat{\mathbf{U}}^{\{l\}}\mathbf{H}_U^{\{l\}} - \mathbf{U})$ is independent with $(\mathbf{Z}^\top)_{l,\cdot}$, we apply Lemma 3 to obtain that $\|\gamma_1\|_2 \lesssim \frac{1}{\sqrt{T}}\|\boldsymbol{\Sigma}_\varepsilon^{1/2}(\widehat{\mathbf{U}}^{\{l\}}\mathbf{H}_U^{\{l\}} - \mathbf{U})\|_{\mathbb{F}} \log n$. Note that

$$\begin{aligned} \|\boldsymbol{\Sigma}_\varepsilon^{1/2}(\widehat{\mathbf{U}}^{\{l\}}\mathbf{H}_U^{\{l\}} - \mathbf{U})\|_{\mathbb{F}} &\leq \|\boldsymbol{\Sigma}_\varepsilon^{1/2}\|_2 \|\widehat{\mathbf{U}}^{\{l\}}\mathbf{H}_U^{\{l\}} - \mathbf{U}\|_{\mathbb{F}} \\ &\leq \sqrt{\|\boldsymbol{\Sigma}_\varepsilon\|_2} \left(\|\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U}\|_{\mathbb{F}} + \|\widehat{\mathbf{U}}^{\{l\}}\mathbf{H}_U^{\{l\}} - \widehat{\mathbf{U}}\mathbf{H}_U\|_{\mathbb{F}} \right). \end{aligned}$$

Then since $\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U} = \widehat{\mathbf{U}}\widehat{\mathbf{U}}^\top\mathbf{U} - \mathbf{U} = (\widehat{\mathbf{U}}\widehat{\mathbf{U}}^\top - \mathbf{U}\mathbf{U}^\top)\mathbf{U}$, we obtain that

$$\|\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U}\|_{\mathbb{F}} \leq \|\widehat{\mathbf{U}}\widehat{\mathbf{U}}^\top - \mathbf{U}\mathbf{U}^\top\|_{\mathbb{F}} \lesssim \|\widehat{\mathbf{U}}\mathbf{R}_U - \mathbf{U}\|_{\mathbb{F}} \lesssim \rho\sqrt{r}, \quad (\text{A.15})$$

where the last inequality is from Lemma 4. So, we conclude that

$$\|\gamma_1\|_2 \lesssim \frac{1}{\sqrt{T}}\sqrt{\|\boldsymbol{\Sigma}_\varepsilon\|_2}(\rho\sqrt{r} + \|\mu_{\{l\}}\|_{\mathbb{F}}) \log n \quad \text{with} \quad \mu_{\{l\}} := \widehat{\mathbf{U}}^{\{l\}}\mathbf{H}_U^{\{l\}} - \widehat{\mathbf{U}}\mathbf{H}_U. \quad (\text{A.16})$$

For γ_2 , using the event \mathcal{E}_Z defined in (A.5), we have that

$$\|\gamma_2\|_2 \lesssim \frac{1}{\sqrt{T}}\sqrt{\|\boldsymbol{\Sigma}_\varepsilon\|_2}\sqrt{n \log n} \|\mu_{\{l\}}\|_2 \leq \frac{1}{\sqrt{T}}\sqrt{\|\boldsymbol{\Sigma}_\varepsilon\|_2 n \log n} \|\mu_{\{l\}}\|_{\mathbb{F}}, \quad (\text{A.17})$$

with probability at least $1 - O(n^{-2})$. Combining the bounds for γ_1 and γ_2 , we obtain that

$$\|\alpha_2\|_2 \leq \|\gamma_1\|_2 + \|\gamma_2\|_2 \lesssim \frac{1}{\sqrt{T}}\sqrt{\|\boldsymbol{\Sigma}_\varepsilon\|_2}\rho\sqrt{r} \log n + \frac{1}{\sqrt{T}}\sqrt{\|\boldsymbol{\Sigma}_\varepsilon\|_2 n \log n} \|\mu_{\{l\}}\|_{\mathbb{F}}. \quad (\text{A.18})$$

We will use self-bounding technique to prove in *Step 2* that

$$\|\mu_{\{l\}}\|_{\mathbb{F}} \lesssim \rho\sqrt{\frac{r}{n}} \log n + \rho\sqrt{\log n} \left(\|\mathbf{V}_{l,\cdot}\|_2 + \|(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})_{l,\cdot}\|_2 \right).$$

As long as we have this, plugging the above upper bound of $\|\mu_{\{l\}}\|_{\mathbb{F}}$ into (A.18), and combining (A.13) and (A.18), we obtain the desired bound for $(\widehat{\mathbf{V}}\widehat{\boldsymbol{\Sigma}}\mathbf{H}_U - (T^{-1/2}\mathbf{X})^\top\mathbf{U})_{l,\cdot}$ as below,

$$\begin{aligned} &\|(\widehat{\mathbf{V}}\widehat{\boldsymbol{\Sigma}}\mathbf{H}_U - (T^{-1/2}\mathbf{X})^\top\mathbf{U})_{l,\cdot}\|_2 \\ &\leq \|\alpha_1\|_2 + \|\alpha_2\|_2 \\ &\lesssim \sigma_r\rho^2\|\mathbf{V}_{l,\cdot}\|_2 + \frac{1}{\sqrt{T}}\sqrt{\|\boldsymbol{\Sigma}_\varepsilon\|_2}\rho\sqrt{r} \log n \\ &\quad + \frac{1}{\sqrt{T}}\sqrt{\|\boldsymbol{\Sigma}_\varepsilon\|_2 n \log n} \left[\rho\sqrt{\frac{r}{n}} \log n + \rho\sqrt{\log n} \left(\|\mathbf{V}_{l,\cdot}\|_2 + \|(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})_{l,\cdot}\|_2 \right) \right] \\ &\lesssim \sigma_r\rho^2 \left(\sqrt{\frac{r}{n}} \log^{3/2} n + \log n \|\mathbf{V}_{l,\cdot}\|_2 + \log n \|(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})_{l,\cdot}\|_2 \right). \end{aligned}$$

Step 2 – Recall that $\mu_{\{l\}} = \widehat{\mathbf{U}}^{\{l\}}\mathbf{H}_U^{\{l\}} - \widehat{\mathbf{U}}\mathbf{H}_U$ as defined in (A.16). We define

$$\nu_{\{l\}} := \widehat{\mathbf{V}}^{\{l\}}\mathbf{H}_V^{\{l\}} - \widehat{\mathbf{V}}\mathbf{H}_V.$$

By definition, we have that $\mu_{\{l\}} = (\widehat{\mathbf{U}}^{\{l\}}(\widehat{\mathbf{U}}^{\{l\}})^\top - \widehat{\mathbf{U}}\widehat{\mathbf{U}}^\top)\mathbf{U}$, so $\|\mu_{\{l\}}\|_{\mathbb{F}} \leq \|\widehat{\mathbf{U}}^{\{l\}}(\widehat{\mathbf{U}}^{\{l\}})^\top - \widehat{\mathbf{U}}\widehat{\mathbf{U}}^\top\|_{\mathbb{F}}$, and similarly we have $\|\nu_{\{l\}}\|_{\mathbb{F}} \leq \|\widehat{\mathbf{V}}^{\{l\}}(\widehat{\mathbf{V}}^{\{l\}})^\top - \widehat{\mathbf{V}}\widehat{\mathbf{V}}^\top\|_{\mathbb{F}}$. Denote by $E_l := \frac{1}{\sqrt{T}}\mathbf{X}^{\{l\}} - \frac{1}{\sqrt{T}}\mathbf{X} = \frac{1}{\sqrt{T}}\boldsymbol{\Sigma}_\varepsilon^{1/2}(\mathbf{Z}^{\{l\}} -$

\mathbf{Z}). Then using (A.9) and our argument in proof of Lemma 4, we have that $\|E_l\| \ll \sigma_r$, and $\sigma_i(\frac{1}{\sqrt{T}}\mathbf{X}^{\{l\}}) \asymp \sigma_i$ for $i = 1, 2, \dots, r$, and $\sigma_{r+1}(\frac{1}{\sqrt{T}}\mathbf{X}^{\{l\}}) \ll \sigma_r$. By Wedin's sin Θ theorem (Chen et al., 2021a, Theorem 2.9), we obtain that

$$\begin{aligned} & \max \left\{ \|\mu_{\{l\}}\|_{\mathbb{F}}, \|\nu_{\{l\}}\|_{\mathbb{F}} \right\} \\ & \leq \max \left\{ \|\widehat{\mathbf{U}}^{\{l\}}(\widehat{\mathbf{U}}^{\{l\}})^{\top} - \widehat{\mathbf{U}}\widehat{\mathbf{U}}^{\top}\|_{\mathbb{F}}, \|\widehat{\mathbf{V}}^{\{l\}}(\widehat{\mathbf{V}}^{\{l\}})^{\top} - \widehat{\mathbf{V}}\widehat{\mathbf{V}}^{\top}\|_{\mathbb{F}} \right\} \\ & \lesssim \frac{1}{\sigma_r(\frac{1}{\sqrt{T}}\mathbf{X}^{\{l\}}) - \sigma_{r+1}(\frac{1}{\sqrt{T}}\mathbf{X}^{\{l\}}) - \|E_l\|_2} \max \left\{ \|E_l\widehat{\mathbf{V}}^{\{l\}}\|_{\mathbb{F}}, \|(E_l)^{\top}\widehat{\mathbf{U}}^{\{l\}}\|_{\mathbb{F}} \right\} \\ & \lesssim \frac{1}{\sigma_r} \max \left\{ \|E_l\widehat{\mathbf{V}}^{\{l\}}\|_{\mathbb{F}}, \|(E_l)^{\top}\widehat{\mathbf{U}}^{\{l\}}\|_{\mathbb{F}} \right\}. \end{aligned}$$

Next, by calculations we have that

$$\begin{aligned} \|(E_l)^{\top}\widehat{\mathbf{U}}^{\{l\}}\|_{\mathbb{F}} &= \left\| \frac{1}{\sqrt{T}}(\mathbf{Z}^{\{l\}} - \mathbf{Z})^{\top}\boldsymbol{\Sigma}_{\varepsilon}^{1/2}\widehat{\mathbf{U}}^{\{l\}} \right\|_{\mathbb{F}} \\ &= \frac{1}{\sqrt{T}} \left\| (\mathbf{Z}_{\cdot, l})^{\top}\boldsymbol{\Sigma}_{\varepsilon}^{1/2}\widehat{\mathbf{U}}^{\{l\}} \right\|_2 \stackrel{(i)}{\lesssim} \frac{1}{\sqrt{T}} \left\| (\mathbf{Z}_{\cdot, l})^{\top}\boldsymbol{\Sigma}_{\varepsilon}^{1/2}\widehat{\mathbf{U}}^{\{l\}}\mathbf{H}_U^{\{l\}} \right\|_2 \\ &\leq \frac{1}{\sqrt{T}} \left\| (\mathbf{Z}_{\cdot, l})^{\top}\boldsymbol{\Sigma}_{\varepsilon}^{1/2}\mathbf{U} \right\|_2 + \frac{1}{\sqrt{T}} \left\| (\mathbf{Z}_{\cdot, l})^{\top}\boldsymbol{\Sigma}_{\varepsilon}^{1/2}(\widehat{\mathbf{U}}^{\{l\}}\mathbf{H}_U^{\{l\}} - \mathbf{U}) \right\|_2 \\ &\lesssim \frac{1}{\sqrt{T}}\sqrt{\|\boldsymbol{\Sigma}_{\varepsilon}\|_2 r \log n} + \frac{1}{\sqrt{T}}\sqrt{\|\boldsymbol{\Sigma}_{\varepsilon}\|_2}(\rho\sqrt{r} + \|\mu_{\{l\}}\|_{\mathbb{F}})\log n. \end{aligned}$$

Here, (i) is because $\sigma_{\min}(\mathbf{H}_U^{\{l\}}) \asymp 1$, which can be proven in the same manner as we prove $\sigma_i(\mathbf{H}_U) \asymp 1$ in Lemma 4; the last inequality uses the bound (A.16) for γ_1 defined in (A.14) and the fact that, by Lemma 3,

$$\left\| (\mathbf{Z}_{\cdot, l})^{\top}\boldsymbol{\Sigma}_{\varepsilon}^{1/2}\mathbf{U} \right\|_2 \lesssim \left\| \boldsymbol{\Sigma}_{\varepsilon}^{1/2}\mathbf{U} \right\|_{\mathbb{F}} \log n \leq \|\boldsymbol{\Sigma}_{\varepsilon}^{1/2}\|_2 \|\mathbf{U}\|_{\mathbb{F}} \log n = \sqrt{\|\boldsymbol{\Sigma}_{\varepsilon}\|_2} r \log n.$$

Also, we have that

$$\begin{aligned} \|E_l\widehat{\mathbf{V}}^{\{l\}}\|_{\mathbb{F}} &= \left\| \frac{1}{\sqrt{T}}\boldsymbol{\Sigma}_{\varepsilon}^{1/2}(\mathbf{Z}^{\{l\}} - \mathbf{Z})\widehat{\mathbf{V}}^{\{l\}} \right\|_{\mathbb{F}} \leq \frac{1}{\sqrt{T}} \|\boldsymbol{\Sigma}_{\varepsilon}^{1/2}\|_2 \left\| (\mathbf{Z}^{\{l\}} - \mathbf{Z})\widehat{\mathbf{V}}^{\{l\}} \right\|_{\mathbb{F}} \\ &= \frac{1}{\sqrt{T}}\sqrt{\|\boldsymbol{\Sigma}_{\varepsilon}\|_2} \|\mathbf{Z}_{\cdot, l}\|_2 \left\| (\widehat{\mathbf{V}}^{\{l\}})_{l, \cdot} \right\|_2 \stackrel{(i)}{\lesssim} \frac{1}{\sqrt{T}}\sqrt{\|\boldsymbol{\Sigma}_{\varepsilon}\|_2} \|\mathbf{Z}_{\cdot, l}\|_2 \left\| (\widehat{\mathbf{V}}^{\{l\}})_{l, \cdot}\mathbf{H}_V^{\{l\}} \right\|_2 \\ &\leq \frac{1}{\sqrt{T}}\sqrt{\|\boldsymbol{\Sigma}_{\varepsilon}\|_2} \|\mathbf{Z}_{\cdot, l}\|_2 \left(\|\mathbf{V}_{l, \cdot}\|_2 + \|(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})_{l, \cdot}\|_2 + \|(\widehat{\mathbf{V}}^{\{l\}}\mathbf{H}_V^{\{l\}} - \widehat{\mathbf{V}}\mathbf{H}_V)_{l, \cdot}\|_2 \right) \\ &\leq \frac{1}{\sqrt{T}}\sqrt{\|\boldsymbol{\Sigma}_{\varepsilon}\|_2} n \log n \left(\|\mathbf{V}_{l, \cdot}\|_2 + \|(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})_{l, \cdot}\|_2 + \|\nu_{\{l\}}\|_{\mathbb{F}} \right). \end{aligned}$$

Here (i) is because $\sigma_{\min}(\mathbf{H}_V^{\{l\}}) \asymp 1$, which can be proven in the same manner as we prove $\sigma_i(\mathbf{H}_V) \asymp 1$ in Lemma 4; the last inequality uses the event \mathcal{E}_Z defined in (A.5).

Combining the above bounds for $\|(E_l)^{\top}\widehat{\mathbf{U}}^{\{l\}}\|_{\mathbb{F}}$ and $\|E_l\widehat{\mathbf{V}}^{\{l\}}\|_{\mathbb{F}}$, and using $\rho \ll 1$, we obtain

$$\begin{aligned} & \max \left\{ \|\mu_{\{l\}}\|_{\mathbb{F}}, \|\nu_{\{l\}}\|_{\mathbb{F}} \right\} \\ & \lesssim \frac{1}{\sigma_r\sqrt{T}}\sqrt{\|\boldsymbol{\Sigma}_{\varepsilon}\|_2 r \log n} + \frac{1}{\sigma_r\sqrt{T}}\sqrt{\|\boldsymbol{\Sigma}_{\varepsilon}\|_2}(\rho\sqrt{r} + \|\mu_{\{l\}}\|_{\mathbb{F}})\log n \\ & + \frac{1}{\sigma_r\sqrt{T}}\sqrt{\|\boldsymbol{\Sigma}_{\varepsilon}\|_2} n \log n \left(\|\mathbf{V}_{l, \cdot}\|_2 + \|(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})_{l, \cdot}\|_2 + \|\nu_{\{l\}}\|_{\mathbb{F}} \right) \\ & \lesssim \rho\sqrt{\frac{r}{n}} \log n + \rho\sqrt{\log n} \left(\|\mathbf{V}_{l, \cdot}\|_2 + \|(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})_{l, \cdot}\|_2 \right) + \rho\sqrt{\log n} \max \left\{ \|\mu_{\{l\}}\|_{\mathbb{F}}, \|\nu_{\{l\}}\|_{\mathbb{F}} \right\}. \end{aligned}$$

Note that both sides of the above inequality include the term $\max\{\|\mu_{\{l\}}\|_{\mathbb{F}}, \|\nu_{\{l\}}\|_{\mathbb{F}}\}$. Then, since $\rho\sqrt{\log n} \ll 1$ according to (3.2), we obtain that

$$\max\{\|\mu_{\{l\}}\|_{\mathbb{F}}, \|\nu_{\{l\}}\|_{\mathbb{F}}\} \lesssim \rho\sqrt{\frac{r}{n}}\log n + \rho\sqrt{\log n} \left(\|\mathbf{V}_{l,\cdot}\|_2 + \|(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})_{l,\cdot}\|_2 \right).$$

Step 3 – For β_1 , in the same manner as we did for α_1 in (A.13), we have that $\|\beta_1\|_2 \lesssim \sigma_r \rho^2 \|\mathbf{U}_{k,\cdot}\|_2$. For β_2 , our starting point is the following inequality

$$\|\beta_2\|_2 \leq \frac{1}{\sqrt{T}} \|(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{k,\cdot}\|_1 \|\mathbf{Z}(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})\|_{2,\infty}. \quad (\text{A.19})$$

Then, the problem boils down to deriving an upper bound for $\|\mathbf{Z}(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})\|_{2,\infty} = \max_{1 \leq m \leq N} \|\mathbf{Z}_{m,\cdot}(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})\|_2$. We use the LOO technique to handle $\mathbf{Z}_{m,\cdot}(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})$ for any $m = 1, 2, \dots, N$. We define $\mathbf{Z}^{(m)}$ as the matrix obtained by replacing the m th row of \mathbf{Z} with a zero vector, and define $\mathbf{X}^{(m)} := \mathbf{B}\mathbf{F}^\top + \boldsymbol{\Sigma}_\varepsilon^{1/2}\mathbf{Z}^{(m)}$. Then, we define $\widehat{\mathbf{U}}^{(m)}$, $\widehat{\mathbf{V}}^{(m)}$, $\mathbf{H}_U^{(m)}$, and $\mathbf{H}_V^{(m)}$ w.r.t. $\frac{1}{\sqrt{T}}\mathbf{X}^{(m)}$, in the same way as how we define $\widehat{\mathbf{U}}$, $\widehat{\mathbf{V}}$, \mathbf{H}_U , and \mathbf{H}_V w.r.t. $\frac{1}{\sqrt{T}}\mathbf{X}$. We have that

$$\mathbf{Z}_{m,\cdot}(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V}) = \underbrace{\mathbf{Z}_{m,\cdot}(\widehat{\mathbf{V}}^{(m)}\mathbf{H}_V^{(m)} - \mathbf{V})}_{=:\zeta_1} + \underbrace{\mathbf{Z}_{m,\cdot}(\widehat{\mathbf{V}}\mathbf{H}_V - \widehat{\mathbf{V}}^{(m)}\mathbf{H}_V^{(m)})}_{=:\zeta_2}. \quad (\text{A.20})$$

For ζ_1 , since $(\widehat{\mathbf{V}}^{(m)}\mathbf{H}_V^{(m)} - \mathbf{V})$ is independent with $\mathbf{Z}_{m,\cdot}$, we apply Lemma 3 to obtain that $\|\zeta_1\|_2 \lesssim \|\widehat{\mathbf{V}}^{(m)}\mathbf{H}_V^{(m)} - \mathbf{V}\|_{\mathbb{F}} \log n$. Note that

$$\|\widehat{\mathbf{V}}^{(m)}\mathbf{H}_V^{(m)} - \mathbf{V}\|_{\mathbb{F}} \leq \|\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V}\|_{\mathbb{F}} + \|\widehat{\mathbf{V}}^{(m)}\mathbf{H}_V^{(m)} - \widehat{\mathbf{V}}\mathbf{H}_V\|_{\mathbb{F}} \lesssim \rho\sqrt{r} + \|\widehat{\mathbf{V}}^{(m)}\mathbf{H}_V^{(m)} - \widehat{\mathbf{V}}\mathbf{H}_V\|_{\mathbb{F}}.$$

Here, the last inequality is because $\|\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V}\|_{\mathbb{F}} \lesssim \rho\sqrt{r}$, which can be proven in the same manner as (A.15). So, we conclude that

$$\|\zeta_1\|_2 \lesssim (\rho\sqrt{r} + \|\nu_{(m)}\|_{\mathbb{F}}) \log n \quad \text{with} \quad \nu_{(m)} := \widehat{\mathbf{V}}^{(m)}\mathbf{H}_V^{(m)} - \widehat{\mathbf{V}}\mathbf{H}_V. \quad (\text{A.21})$$

For ζ_2 , using the event \mathcal{E}_Z defined in (A.5), we have that

$$\|\zeta_2\|_2 \lesssim \sqrt{n \log n} \|\nu_{(m)}\|_2,$$

with probability at least $1 - O(n^{-2})$. Combining the bounds for ζ_1 and ζ_2 , we obtain that

$$\|\mathbf{Z}_{m,\cdot}(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})\|_2 \leq \|\zeta_1\|_2 + \|\zeta_2\|_2 \lesssim \rho\sqrt{r} \log n + \sqrt{n \log n} \|\nu_{(m)}\|_{\mathbb{F}}. \quad (\text{A.22})$$

We will use self-bounding technique to prove in *Step 4* that

$$\|\nu_{(m)}\|_{\mathbb{F}} \lesssim \rho\sqrt{\frac{r}{n}} \log n + \omega\sqrt{\log n} \left(\|\mathbf{U}\|_{2,\infty} + \|\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U}\|_{2,\infty} \right).$$

As long as we have this, plugging the above upper bound of $\|\nu_{(m)}\|_{\mathbb{F}}$ into (A.22), we obtain the upper bound for $\|\mathbf{Z}(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})\|_{2,\infty} = \max_{1 \leq m \leq N} \|\mathbf{Z}_{m,\cdot}(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})\|_2$, which leads to an upper bound for $\|\beta_2\|_2$ in (A.19). Finally, combining the bound for β_1 and the bound for β_2 in (A.19), we obtain the desired bound for $(\widehat{\mathbf{V}}\widehat{\boldsymbol{\Sigma}}\mathbf{H}_U - (T^{-1/2}\mathbf{X})^\top\mathbf{U})_{k,\cdot}$ as below,

$$\begin{aligned} & \|(\widehat{\mathbf{V}}\widehat{\boldsymbol{\Sigma}}\mathbf{H}_U - (T^{-1/2}\mathbf{X})^\top\mathbf{U})_{k,\cdot}\|_2 \\ & \leq \|\beta_1\|_2 + \|\beta_2\|_2 \\ & \lesssim \sigma_r \rho^2 \|\mathbf{U}_{k,\cdot}\|_2 + \frac{1}{\sqrt{T}} \|(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{k,\cdot}\|_1 \rho\sqrt{r} \log n \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{\sqrt{T}} \left\| (\Sigma_\varepsilon^{1/2})_{k,\cdot} \right\|_1 \sqrt{n \log n} \left[\rho \sqrt{\frac{r}{n}} \log n + \omega \sqrt{\log n} \left(\|U\|_{2,\infty} + \|\widehat{U}H_U - U\|_{2,\infty} \right) \right] \\
& \lesssim \sigma_r \omega_k \rho \sqrt{\frac{r}{n}} \log^{3/2} n + \sigma_r \rho^2 \|U_{k,\cdot}\|_2 + \sigma_r \omega_k \omega \log n \left(\|U\|_{2,\infty} + \|\widehat{U}H_U - U\|_{2,\infty} \right).
\end{aligned}$$

Step 4 – Recall that $\nu_{(m)} := \widehat{V}^{(m)} H_V^{(m)} - \widehat{V} H_V$ as defined in (A.21). We define

$$\mu_{(m)} := \widehat{U}^{(m)} H_U^{(m)} - \widehat{U} H_U.$$

By definition, we have that $\mu_{(m)} = (\widehat{U}^{(m)} (\widehat{U}^{(m)})^\top - \widehat{U} \widehat{U}^\top) U$, so $\|\mu_{(m)}\|_F \leq \|(\widehat{U}^{(m)} (\widehat{U}^{(m)})^\top - \widehat{U} \widehat{U}^\top)\|_F$, and similarly we have $\|\nu_{(m)}\|_F \leq \|(\widehat{V}^{(m)} (\widehat{V}^{(m)})^\top - \widehat{V} \widehat{V}^\top)\|_F$. Denote by $E_m := \frac{1}{\sqrt{T}} \mathbf{X}^{(m)} - \frac{1}{\sqrt{T}} \mathbf{X} = \frac{1}{\sqrt{T}} \Sigma_\varepsilon^{1/2} (\mathbf{Z}^{(m)} - \mathbf{Z})$. Then using (A.9) and our argument in proof of Lemma 4, we have that $\|E_m\| \ll \sigma_r$, and $\sigma_i(\frac{1}{\sqrt{T}} \mathbf{X}^{(m)}) \asymp \sigma_i$ for $i = 1, 2, \dots, r$, and $\sigma_{r+1}(\frac{1}{\sqrt{T}} \mathbf{X}^{(m)}) \ll \sigma_r$. By Wedin's sin Θ theorem (Chen et al., 2021a, Theorem 2.9), we obtain that

$$\begin{aligned}
& \max \left\{ \|\mu_{(m)}\|_F, \|\nu_{(m)}\|_F \right\} \\
& \leq \max \left\{ \|(\widehat{U}^{(m)} (\widehat{U}^{(m)})^\top - \widehat{U} \widehat{U}^\top)\|_F, \|(\widehat{V}^{(m)} (\widehat{V}^{(m)})^\top - \widehat{V} \widehat{V}^\top)\|_F \right\} \\
& \lesssim \frac{1}{\sigma_r (\frac{1}{\sqrt{T}} \mathbf{X}^{(m)}) - \sigma_{r+1} (\frac{1}{\sqrt{T}} \mathbf{X}^{(m)}) - \|E_m\|_2} \max \left\{ \|E_m \widehat{V}^{(m)}\|_F, \|(E_m)^\top \widehat{U}^{(m)}\|_F \right\} \\
& \lesssim \frac{1}{\sigma_r} \max \left\{ \|E_m \widehat{V}^{(m)}\|_F, \|(E_m)^\top \widehat{U}^{(m)}\|_F \right\}.
\end{aligned}$$

Next, by calculations we have that

$$\begin{aligned}
\|(E_m)^\top \widehat{U}^{(m)}\|_F & = \left\| \frac{1}{\sqrt{T}} (\mathbf{Z}^{(m)} - \mathbf{Z})^\top \Sigma_\varepsilon^{1/2} \widehat{U}^{(m)} \right\|_F = \frac{1}{\sqrt{T}} \left\| (\mathbf{Z}_{m,\cdot})^\top (\Sigma_\varepsilon^{1/2} \widehat{U}^{(m)})_{m,\cdot} \right\|_F \\
& = \frac{1}{\sqrt{T}} \|\mathbf{Z}_{m,\cdot}\|_2 \left\| (\Sigma_\varepsilon^{1/2} \widehat{U}^{(m)})_{m,\cdot} \right\|_2 \stackrel{(i)}{\lesssim} \frac{1}{\sqrt{T}} \|\mathbf{Z}_{m,\cdot}\|_2 \left\| (\Sigma_\varepsilon^{1/2} \widehat{U}^{(m)})_{m,\cdot} H_U^{(m)} \right\|_2 \\
& \leq \frac{1}{\sqrt{T}} \|\mathbf{Z}_{m,\cdot}\|_2 \left\| \Sigma_\varepsilon^{1/2} \widehat{U}^{(m)} H_U^{(m)} \right\|_{2,\infty} \leq \frac{1}{\sqrt{T}} \|\mathbf{Z}_{m,\cdot}\|_2 \left\| \Sigma_\varepsilon^{1/2} \right\|_1 \left\| \widehat{U}^{(m)} H_U^{(m)} \right\|_{2,\infty} \\
& \leq \frac{1}{\sqrt{T}} \left\| \Sigma_\varepsilon^{1/2} \right\|_1 \|\mathbf{Z}_{m,\cdot}\|_2 \left(\|U\|_{2,\infty} + \|\widehat{U}H_U - U\|_{2,\infty} + \|\widehat{U}^{(m)} H_U^{(m)} - \widehat{U} H_U\|_{2,\infty} \right) \\
& \lesssim \frac{1}{\sqrt{T}} \left\| \Sigma_\varepsilon^{1/2} \right\|_1 \sqrt{n \log n} \left(\|U\|_{2,\infty} + \|\widehat{U}H_U - U\|_{2,\infty} + \|\mu_{(m)}\|_F \right).
\end{aligned}$$

Here (i) is because $\sigma_{\min}(H_U^{(m)}) \asymp 1$, which can be proven in the same manner as we prove $\sigma_i(H_U) \asymp 1$ in Lemma 4, and the last inequality uses the event \mathcal{E}_Z defined in (A.5).

Also, we have that

$$\begin{aligned}
\|E_m \widehat{V}^{(m)}\|_F & = \left\| \frac{1}{\sqrt{T}} \Sigma_\varepsilon^{1/2} (\mathbf{Z}^{(m)} - \mathbf{Z}) \widehat{V}^{(m)} \right\|_F \leq \frac{1}{\sqrt{T}} \left\| \Sigma_\varepsilon^{1/2} \right\|_2 \left\| (\mathbf{Z}^{(m)} - \mathbf{Z}) \widehat{V}^{(m)} \right\|_F \\
& = \frac{1}{\sqrt{T}} \sqrt{\|\Sigma_\varepsilon\|_2} \left\| \mathbf{Z}_{m,\cdot} \widehat{V}^{(m)} \right\|_2 \stackrel{(i)}{\lesssim} \frac{1}{\sqrt{T}} \sqrt{\|\Sigma_\varepsilon\|_2} \left\| \mathbf{Z}_{m,\cdot} \widehat{V}^{(m)} H_V^{(m)} \right\|_2 \\
& \leq \frac{1}{\sqrt{T}} \sqrt{\|\Sigma_\varepsilon\|_2} \left(\|\mathbf{Z}_{m,\cdot} \mathbf{V}\|_2 + \left\| \mathbf{Z}_{m,\cdot} (\widehat{V}^{(m)} H_V^{(m)} - \mathbf{V}) \right\|_2 \right) \\
& \lesssim \frac{1}{\sqrt{T}} \sqrt{\|\Sigma_\varepsilon\|_2} r \log n + \frac{1}{\sqrt{T}} \sqrt{\|\Sigma_\varepsilon\|_2} (\rho \sqrt{r} + \|\nu_{(m)}\|_F) \log n.
\end{aligned}$$

Here, (i) is because $\sigma_{\min}(H_V^{(m)}) \asymp 1$, which can be proven in the same manner as we prove $\sigma_i(H_V) \asymp 1$ in Lemma 4; the last inequality uses the bound (A.21) for ζ_1 defined in (A.20) and the fact that, by Lemma 3, $\|\mathbf{Z}_{m,\cdot} \mathbf{V}\|_2 \lesssim \|\mathbf{V}\|_F \log n = \sqrt{r} \log n$.

Combining the above bounds for $\|(E_m)^\top \widehat{\mathbf{U}}^{(m)}\|_{\mathbb{F}}$ and $\|E_m \widehat{\mathbf{V}}^{(m)}\|_{\mathbb{F}}$, and using $\rho \ll 1$, we obtain

$$\begin{aligned} & \max \{ \|\mu_{(m)}\|_{\mathbb{F}}, \|\nu_{(m)}\|_{\mathbb{F}} \} \\ & \lesssim \frac{1}{\sigma_r \sqrt{T}} \sqrt{\|\Sigma_\varepsilon\|_2 r \log n} + \frac{1}{\sqrt{T}} \sqrt{\|\Sigma_\varepsilon\|_2} (\rho \sqrt{r} + \|\nu_{(m)}\|_{\mathbb{F}}) \log n \\ & + \frac{1}{\sigma_r \sqrt{T}} \|\Sigma_\varepsilon^{1/2}\|_1 \sqrt{n \log n} \left(\|\mathbf{U}\|_{2,\infty} + \|\widehat{\mathbf{U}} \mathbf{H}_U - \mathbf{U}\|_{2,\infty} + \|\mu_{(m)}\|_{\mathbb{F}} \right) \\ & \lesssim \rho \sqrt{\frac{r}{n}} \log n + \omega \sqrt{\log n} \left(\|\mathbf{U}\|_{2,\infty} + \|\widehat{\mathbf{U}} \mathbf{H}_U - \mathbf{U}\|_{2,\infty} \right) + \omega \sqrt{\log n} \max \{ \|\mu_{(m)}\|_{\mathbb{F}}, \|\nu_{(m)}\|_{\mathbb{F}} \}. \end{aligned}$$

Note that, both sides of the above inequality include the term $\max \{ \|\mu_{(m)}\|_{\mathbb{F}}, \|\nu_{(m)}\|_{\mathbb{F}} \}$. Then, since $\omega \sqrt{\log n} \ll 1$ according to (3.1), we obtain that

$$\max \{ \|\mu_{(m)}\|_{\mathbb{F}}, \|\nu_{(m)}\|_{\mathbb{F}} \} \lesssim \rho \sqrt{\frac{r}{n}} \log n + \omega \sqrt{\log n} \left(\|\mathbf{U}\|_{2,\infty} + \|\widehat{\mathbf{U}} \mathbf{H}_U - \mathbf{U}\|_{2,\infty} \right).$$

□

Then we handle $\widehat{\mathbf{U}} \mathbf{H}_U - \mathbf{U}$ and $\widehat{\mathbf{V}} \mathbf{H}_V - \mathbf{V}$.

Lemma 7. *Suppose that Assumptions 1, 2, and 3 hold, and*

$$\rho \ll \sqrt{\log n} \text{ and } r \log n \ll n. \quad (\text{A.23})$$

(i) *Assuming (3.2), i.e., $\rho \sqrt{\log n} \ll 1$, then we have that, for $l = 1, 2, \dots, T$, with probability at least $1 - O(n^{-2})$,*

$$\|(\widehat{\mathbf{V}} \mathbf{H}_V - \mathbf{V})_{l,\cdot}\|_2 \lesssim \rho \sqrt{\frac{r}{n}} \log n + (\rho^2 + \rho \sqrt{\frac{r}{n}}) \log n \|\mathbf{V}_{l,\cdot}\|_2.$$

(ii) *Assuming (3.1), i.e., $\omega \sqrt{\log n} \ll 1$, then we have that, with probability at least $1 - O(n^{-2})$,*

$$\|\widehat{\mathbf{U}} \mathbf{H}_U - \mathbf{U}\|_{2,\infty} \lesssim \omega \sqrt{\frac{r}{n}} \log n + (\rho \sqrt{\frac{r}{n}} + \omega^2) \log n \|\mathbf{U}\|_{2,\infty},$$

and for $k = 1, 2, \dots, N$,

$$\|(\widehat{\mathbf{U}} \mathbf{H}_U - \mathbf{U})_{k,\cdot}\|_2 \lesssim \omega_k \sqrt{\frac{r}{n}} \log n + (\rho \sqrt{\frac{r}{n}} + \omega^2) \log n \|\mathbf{U}_{k,\cdot}\|_2 + \omega_k \omega \log n \|\mathbf{U}\|_{2,\infty},$$

where $\omega_k = \|((\Sigma_\varepsilon)^{1/2})_{k,\cdot}\|_1 \sqrt{n}/(\sigma_r \sqrt{T})$ is defined in (A.12).

Proof. We will prove (i) and (ii) in *Step 1* and *Step 2*, respectively.

Step 1 – Since $\lambda_i \asymp \sigma_i$ according to Lemma 2, we obtain that $\|(\widehat{\mathbf{V}} \mathbf{H}_V - \mathbf{V})_{l,\cdot}\|_2 \lesssim \frac{1}{\sigma_r} \|(\widehat{\mathbf{V}} \mathbf{H}_V \Lambda - \mathbf{V} \Lambda)_{l,\cdot}\|_2$. Next, by calculations, we have that

$$\widehat{\mathbf{V}} \mathbf{H}_V \Lambda - \mathbf{V} \Lambda = \widehat{\mathbf{V}} \widehat{\Sigma} \mathbf{H}_U - (T^{-1/2} \mathbf{X})^\top \mathbf{U} + T^{-1/2} \mathbf{Z}^\top \Sigma_\varepsilon^{1/2} \mathbf{U} + \widehat{\mathbf{V}} \mathbf{H}_V \Lambda - \widehat{\mathbf{V}} \widehat{\Sigma} \mathbf{H}_U.$$

So we obtain that

$$\begin{aligned} & \|(\widehat{\mathbf{V}} \mathbf{H}_V - \mathbf{V})_{l,\cdot}\|_2 \\ & \lesssim \underbrace{\frac{1}{\sigma_r} \|(\widehat{\mathbf{V}} \widehat{\Sigma} \mathbf{H}_U - (T^{-1/2} \mathbf{X})^\top \mathbf{U})_{l,\cdot}\|_2}_{=: \mathbf{a}_1} + \underbrace{\frac{1}{\sigma_r} \|T^{-1/2} (\mathbf{Z}^\top)_{l,\cdot} \Sigma_\varepsilon^{1/2} \mathbf{U}\|_2}_{=: \mathbf{a}_2} + \underbrace{\frac{1}{\sigma_r} \|\widehat{\mathbf{V}}_{l,\cdot} (\mathbf{H}_V \Lambda - \widehat{\Sigma} \mathbf{H}_U)\|_2}_{=: \mathbf{a}_3}. \end{aligned}$$

The term \mathbf{a}_1 can be bounded using Lemma 6. For \mathbf{a}_2 , we obtain by Lemma 3 that

$$\mathbf{a}_2 \leq \frac{1}{\sigma_r \sqrt{T}} \|\Sigma_\varepsilon^{1/2} \mathbf{U}\|_{\mathbb{F}} \log n \leq \frac{1}{\sigma_r \sqrt{T}} \|\Sigma_\varepsilon^{1/2}\|_2 \|\mathbf{U}\|_{\mathbb{F}} \log n = \rho \sqrt{\frac{r}{n}} \log n.$$

For \mathbf{a}_3 , we have that $\|\widehat{\mathbf{V}}_{l,\cdot}(\mathbf{H}_V\boldsymbol{\Lambda} - \widehat{\boldsymbol{\Sigma}}\mathbf{H}_U)\|_2 \leq \|\widehat{\mathbf{V}}_{l,\cdot}\|_2\|\mathbf{H}_V\boldsymbol{\Lambda} - \widehat{\boldsymbol{\Sigma}}\mathbf{H}_U\|_2$ and we will handle the two terms in the right hand side. For $\|\widehat{\mathbf{V}}_{l,\cdot}\|_2$, since $\sigma_i(\mathbf{H}_V) \asymp 1$ in Lemma 4, we have that

$$\|\widehat{\mathbf{V}}_{l,\cdot}\|_2 \lesssim \|\widehat{\mathbf{V}}_{l,\cdot}\mathbf{H}_V\|_2 = \|(\widehat{\mathbf{V}}\mathbf{H}_V)_{l,\cdot}\|_2 \leq \|\mathbf{V}_{l,\cdot}\|_2 + \|(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})_{l,\cdot}\|_2.$$

For $\|\mathbf{H}_V\boldsymbol{\Lambda} - \widehat{\boldsymbol{\Sigma}}\mathbf{H}_U\|_2$, using the full SVD of $T^{-1/2}\mathbf{X}$ in the proof of Lemma 4, we have that

$$\begin{aligned} \mathbf{H}_V\boldsymbol{\Lambda} - \widehat{\boldsymbol{\Sigma}}\mathbf{H}_U &= \widehat{\mathbf{V}}^\top \mathbf{V}\boldsymbol{\Lambda} - \widehat{\boldsymbol{\Sigma}}\widehat{\mathbf{U}}^\top \mathbf{U} = \widehat{\mathbf{V}}^\top (\mathbf{V}\boldsymbol{\Lambda}\mathbf{U}^\top - \widehat{\mathbf{V}}\widehat{\boldsymbol{\Sigma}}\widehat{\mathbf{U}}^\top)\mathbf{U} \\ &= \widehat{\mathbf{V}}^\top (T^{-1/2}\mathbf{E}^\top - \widehat{\mathbf{V}}_\perp(\widehat{\boldsymbol{\Sigma}}_\perp)^\top(\widehat{\mathbf{U}}_\perp)^\top)\mathbf{U} = T^{-1/2}\widehat{\mathbf{V}}^\top \mathbf{E}^\top \mathbf{U}. \end{aligned}$$

Thus, we obtain that

$$\begin{aligned} \|\mathbf{H}_V\boldsymbol{\Lambda} - \widehat{\boldsymbol{\Sigma}}\mathbf{H}_U\|_2 &\stackrel{(i)}{\lesssim} T^{-1/2}\|\mathbf{U}^\top \mathbf{E}\widehat{\mathbf{V}}\mathbf{H}_V\|_2 \leq T^{-1/2}\|\mathbf{U}^\top \mathbf{E}\mathbf{V}\|_2 + T^{-1/2}\|\mathbf{U}^\top \mathbf{E}(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})\|_2 \\ &\lesssim T^{-1/2}\|\mathbf{U}^\top \boldsymbol{\Sigma}_\varepsilon^{1/2}\mathbf{Z}\mathbf{V}\|_2 + T^{-1/2}\|\mathbf{U}^\top \boldsymbol{\Sigma}_\varepsilon^{1/2}\mathbf{Z}\|_2\|\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V}\|_2 \\ &\stackrel{(ii)}{\lesssim} \sigma_r\rho\sqrt{\frac{1}{n}(r + \log n)} + \frac{1}{\sqrt{T}}\sqrt{\|\boldsymbol{\Sigma}_\varepsilon\|_2}\sqrt{r + T + \log n} \cdot \rho \\ &\lesssim \sigma_r\rho\sqrt{\frac{1}{n}(r + \log n)} + \sigma_r\rho^2, \end{aligned}$$

where (i) is because $\sigma_i(\mathbf{H}_V) \asymp 1$ in Lemma 4, (ii) uses (G.3) in Lemma 19 of Yan and Wainwright (2024) and the upper bound for $\|\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V}\|_2$ is similar to that in (A.11). Combing the bounds for $\|\widehat{\mathbf{V}}_{l,\cdot}\|_2$ and $\|\mathbf{H}_V\boldsymbol{\Lambda} - \widehat{\boldsymbol{\Sigma}}\mathbf{H}_U\|_2$, we obtain that

$$\begin{aligned} \mathbf{a}_3 &\lesssim \frac{1}{\sigma_r} \left(\|\mathbf{V}_{l,\cdot}\|_2 + \|(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})_{l,\cdot}\|_2 \right) \left(\sigma_r\rho\sqrt{\frac{1}{n}(r + \log n)} + \sigma_r\rho^2 \right) \\ &\lesssim \left(\|\mathbf{V}_{l,\cdot}\|_2 + \|(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})_{l,\cdot}\|_2 \right) \left(\sqrt{\frac{1}{n}(r + \log n)} + \rho \right) \rho. \end{aligned}$$

Finally, we combine the bounds for \mathbf{a}_1 , \mathbf{a}_2 , and \mathbf{a}_3 to obtain that, for $l = 1, 2, \dots, T$,

$$\begin{aligned} &\|(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})_{l,\cdot}\|_2 \\ &\lesssim \rho^2 \left(\sqrt{\frac{r}{n}} \log^{3/2} n + \log n \|\mathbf{V}_{l,\cdot}\|_2 + \log n \|(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})_{l,\cdot}\|_2 \right) \\ &\quad + \rho\sqrt{\frac{r}{n}} \log n + \left(\|\mathbf{V}_{l,\cdot}\|_2 + \|(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})_{l,\cdot}\|_2 \right) \left(\sqrt{\frac{1}{n}(r + \log n)} + \rho \right) \rho \\ &\lesssim \rho\sqrt{\frac{r}{n}} \log n + (\rho^2 + \rho\sqrt{\frac{r}{n}}) \log n \|\mathbf{V}_{l,\cdot}\|_2 \\ &\quad + (\rho^2 + \rho\sqrt{\frac{r}{n}}) \log n \|(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})_{l,\cdot}\|_2, \end{aligned}$$

where we use $\rho\sqrt{\log n} \ll 1$ in (3.2). Then using the self-bounding technique, since $(\rho^2 + \rho\sqrt{\frac{r}{n}}) \log n \ll 1$ owing to (3.2) and (A.23), we conclude that

$$\|(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})_{l,\cdot}\|_2 \lesssim \rho\sqrt{\frac{r}{n}} \log n + (\rho^2 + \rho\sqrt{\frac{r}{n}}) \log n \|\mathbf{V}_{l,\cdot}\|_2.$$

Step 2 – Since $\lambda_i \asymp \sigma_i$ according to Lemma 2, we obtain that $\|(\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U})_{k,\cdot}\|_2 \lesssim \frac{1}{\sigma_r} \|(\widehat{\mathbf{U}}\mathbf{H}_U\boldsymbol{\Lambda} - \mathbf{U}\boldsymbol{\Lambda})_{k,\cdot}\|_2$. Next, by calculations, we have that

$$\widehat{\mathbf{U}}\mathbf{H}_U\boldsymbol{\Lambda} - \mathbf{U}\boldsymbol{\Lambda} = \widehat{\mathbf{U}}\widehat{\boldsymbol{\Sigma}}\mathbf{H}_V - (T^{-1/2}\mathbf{X})\mathbf{V} + T^{-1/2}\boldsymbol{\Sigma}_\varepsilon^{1/2}\mathbf{Z}\mathbf{V} + \widehat{\mathbf{U}}\mathbf{H}_U\boldsymbol{\Lambda} - \widehat{\mathbf{U}}\widehat{\boldsymbol{\Sigma}}\mathbf{H}_V.$$

So we obtain that

$$\begin{aligned} & \|(\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U})_{k,\cdot}\|_2 \\ & \lesssim \underbrace{\frac{1}{\sigma_r} \|(\widehat{\mathbf{U}}\widehat{\Sigma}\mathbf{H}_V - (T^{-1/2}\mathbf{X})\mathbf{V})_{k,\cdot}\|_2}_{=: \mathbf{b}_1} + \underbrace{\frac{1}{\sigma_r} \|T^{-1/2}(\Sigma_\varepsilon^{1/2})_{k,\cdot}\mathbf{Z}\mathbf{V}\|_2}_{=: \mathbf{b}_2} + \underbrace{\frac{1}{\sigma_r} \|\widehat{\mathbf{U}}_{k,\cdot}(\mathbf{H}_U\mathbf{\Lambda} - \widehat{\Sigma}\mathbf{H}_V)\|_2}_{=: \mathbf{b}_3}. \end{aligned}$$

The term \mathbf{b}_1 can be bounded using Lemma 6. For \mathbf{b}_2 , we obtain by Lemma 3 that

$$\begin{aligned} \mathbf{b}_2 & \leq \frac{1}{\sigma_r\sqrt{T}} \|(\Sigma_\varepsilon^{1/2})_{k,\cdot}\|_1 \|\mathbf{Z}\mathbf{V}\|_{2,\infty} \lesssim \frac{1}{\sigma_r\sqrt{T}} \|(\Sigma_\varepsilon^{1/2})_{k,\cdot}\|_1 \|\mathbf{V}\|_{\mathbb{F}} \log n \\ & = \frac{1}{\sigma_r\sqrt{T}} \|(\Sigma_\varepsilon^{1/2})_{k,\cdot}\|_1 \sqrt{r} \log n = \omega_k \sqrt{\frac{r}{n}} \log n. \end{aligned}$$

For \mathbf{b}_3 , we have that $\|\widehat{\mathbf{U}}_{k,\cdot}(\mathbf{H}_U\mathbf{\Lambda} - \widehat{\Sigma}\mathbf{H}_V)\|_2 \leq \|\widehat{\mathbf{U}}_{k,\cdot}\|_2 \|\mathbf{H}_U\mathbf{\Lambda} - \widehat{\Sigma}\mathbf{H}_V\|_2$ and we will handle the two terms in the right hand side. For $\|\widehat{\mathbf{U}}_{k,\cdot}\|_2$, since $\sigma_i(\mathbf{H}_U) \asymp 1$ in Lemma 4, we have that

$$\|\widehat{\mathbf{U}}_{k,\cdot}\|_2 \lesssim \|\widehat{\mathbf{U}}_{k,\cdot}\mathbf{H}_U\|_2 = \|(\widehat{\mathbf{U}}\mathbf{H}_U)_{k,\cdot}\|_2 \leq \|\mathbf{U}_{k,\cdot}\|_2 + \|(\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U})_{k,\cdot}\|_2.$$

For $\|\mathbf{H}_U\mathbf{\Lambda} - \widehat{\Sigma}\mathbf{H}_V\|_2$, similar to the upper bound for $\|\mathbf{H}_V\mathbf{\Lambda} - \widehat{\Sigma}\mathbf{H}_U\|_2$ in Step 1, we have that

$$\|\mathbf{H}_U\mathbf{\Lambda} - \widehat{\Sigma}\mathbf{H}_V\|_2 \lesssim \sigma_r \rho \sqrt{\frac{1}{n}(r + \log n)} + \sigma_r \rho^2.$$

Combining the bounds for $\|\widehat{\mathbf{U}}_{k,\cdot}\|_2$ and $\|\mathbf{H}_U\mathbf{\Lambda} - \widehat{\Sigma}\mathbf{H}_V\|_2$, we obtain that

$$\mathbf{b}_3 \lesssim \left(\|\mathbf{U}_{k,\cdot}\|_2 + \|(\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U})_{k,\cdot}\|_2 \right) \left(\rho \sqrt{\frac{1}{n}(r + \log n)} + \rho^2 \right).$$

Finally, we combine the bounds for \mathbf{b}_1 , \mathbf{b}_2 , and \mathbf{b}_3 to obtain that, for $k = 1, 2, \dots, N$,

$$\begin{aligned} & \|(\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U})_{k,\cdot}\|_2 \\ & \lesssim \left(\omega_k \rho \sqrt{\frac{r}{n}} \log^{3/2} n + \rho^2 \|\mathbf{U}_{k,\cdot}\|_2 + \omega_k \omega \log n \left(\|\mathbf{U}\|_{2,\infty} + \|\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U}\|_{2,\infty} \right) \right) \\ & + \omega_k \sqrt{\frac{r}{n}} \log n + \left(\|\mathbf{U}_{k,\cdot}\|_2 + \|(\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U})_{k,\cdot}\|_2 \right) \left(\rho \sqrt{\frac{1}{n}(r + \log n)} + \rho^2 \right) \\ & \lesssim \omega_k \sqrt{\frac{r}{n}} \log n + \left(\rho \sqrt{\frac{1}{n}(r + \log n)} + \rho^2 \right) \left(\|\mathbf{U}_{k,\cdot}\|_2 + \|(\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U})_{k,\cdot}\|_2 \right) \\ & + \omega_k \omega \log n \left(\|\mathbf{U}\|_{2,\infty} + \|\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U}\|_{2,\infty} \right), \end{aligned}$$

where we use the fact that $\rho\sqrt{\log n} \leq \omega\sqrt{\log n} \ll 1$. Taking supremum w.r.t. k , we obtain that

$$\|\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U}\|_{2,\infty} \lesssim \omega \sqrt{\frac{r}{n}} \log n + \left(\rho \sqrt{\frac{1}{n}(r + \log n)} + \omega^2 \log n \right) \left(\|\mathbf{U}\|_{2,\infty} + \|\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U}\|_{2,\infty} \right).$$

Then using the self-bounding technique for $\|\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U}\|_{2,\infty}$, since $\omega\sqrt{\log n} \ll 1$ and $\rho\sqrt{\frac{1}{n}(r + \log n)} \ll 1$ and $\rho \leq \omega$, we conclude that

$$\|\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U}\|_{2,\infty} \lesssim \omega \sqrt{\frac{r}{n}} \log n + \left(\rho \sqrt{\frac{r}{n}} \log n + \omega^2 \log n \right) \|\mathbf{U}\|_{2,\infty}.$$

Plugging the above bound of $\|\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U}\|_{2,\infty}$ into the previous inequality for $\|(\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U})_{k,\cdot}\|_2$, we obtain that

$$\begin{aligned} \|(\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U})_{k,\cdot}\|_2 &\lesssim \omega_k \sqrt{\frac{r}{n}} \log n + \left(\rho \sqrt{\frac{1}{n}(r + \log n)} + \rho^2 \right) \left(\|\mathbf{U}_{k,\cdot}\|_2 + \|(\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U})_{k,\cdot}\|_2 \right) \\ &\quad + \omega_k \omega \log n \left(\|\mathbf{U}\|_{2,\infty} + \omega \sqrt{\frac{r}{n}} \log n + \left(\rho \sqrt{\frac{r}{n}} + \omega^2 \right) \log n \|\mathbf{U}\|_{2,\infty} \right). \end{aligned}$$

Then using the self-bounding technique for $\|(\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U})_{k,\cdot}\|_2$, since $\rho\sqrt{\log n} \leq \omega\sqrt{\log n} \ll 1$ in (3.1), we conclude that

$$\|(\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U})_{k,\cdot}\|_2 \lesssim \omega_k \sqrt{\frac{r}{n}} \log n + \left(\rho \sqrt{\frac{r}{n}} \log n + \omega^2 \log n \right) \|\mathbf{U}_{k,\cdot}\|_2 + \omega_k \omega \log n \|\mathbf{U}\|_{2,\infty}.$$

□

A.2 Proof of Theorem 1

We are now ready to prove Theorem 1. We will prove (i) and (ii) in *Step 1* and *Step 2*, respectively.

Step 1 – By calculations, we have that

$$\boldsymbol{\Psi}_U = (\widehat{\mathbf{U}}\widehat{\boldsymbol{\Sigma}}\mathbf{H}_V - (T^{-1/2}\mathbf{X})\mathbf{V})\boldsymbol{\Lambda}^{-1} + \widehat{\mathbf{U}}\widehat{\boldsymbol{\Sigma}}(\mathbf{R}_V - \mathbf{H}_V)\boldsymbol{\Lambda}^{-1} + (\widehat{\mathbf{U}}\mathbf{R}_U)(\boldsymbol{\Lambda} - (\mathbf{R}_U)^\top \widehat{\boldsymbol{\Sigma}}\mathbf{R}_V)\boldsymbol{\Lambda}^{-1}.$$

Since $\|\widehat{\mathbf{U}}_{k,\cdot}\mathbf{R}_U\|_2 = \|\widehat{\mathbf{U}}_{k,\cdot}\|_2$, we obtain that

$$\begin{aligned} &\|(\boldsymbol{\Psi}_U)_{k,\cdot}\|_2 \\ &\lesssim \|(\widehat{\mathbf{U}}\widehat{\boldsymbol{\Sigma}}\mathbf{H}_V - (T^{-1/2}\mathbf{X})\mathbf{V})_{k,\cdot}\|_2 \|\boldsymbol{\Lambda}^{-1}\|_2 + \|\widehat{\mathbf{U}}_{k,\cdot}\|_2 \left(\|\widehat{\boldsymbol{\Sigma}}(\mathbf{R}_V - \mathbf{H}_V)\boldsymbol{\Lambda}^{-1}\|_2 + \|(\boldsymbol{\Lambda} - (\mathbf{R}_U)^\top \widehat{\boldsymbol{\Sigma}}\mathbf{R}_V)\boldsymbol{\Lambda}^{-1}\|_2 \right) \\ &\stackrel{(i)}{\lesssim} \|(\widehat{\mathbf{U}}\widehat{\boldsymbol{\Sigma}}\mathbf{H}_V - (T^{-1/2}\mathbf{X})\mathbf{V})_{k,\cdot}\|_2 \frac{1}{\sigma_r} + \|\widehat{\mathbf{U}}_{k,\cdot}\|_2 \left(\sigma_r \rho^2 \frac{1}{\sigma_r} + \frac{1}{\sigma_r} \left(\sigma_r \rho^2 + \sigma_r \rho \sqrt{\frac{1}{n}(r + \log n)} \right) \right) \\ &\lesssim \|(\widehat{\mathbf{U}}\widehat{\boldsymbol{\Sigma}}\mathbf{H}_V - (T^{-1/2}\mathbf{X})\mathbf{V})_{k,\cdot}\|_2 \frac{1}{\sigma_r} + \|\widehat{\mathbf{U}}_{k,\cdot}\|_2 \left(\rho^2 + \rho \sqrt{\frac{r}{n}} \sqrt{\log n} \right), \end{aligned}$$

where (i) is owing to the inequalities in Lemmas 4 and 5. Using $\sigma_i(\mathbf{H}_U) \asymp 1$ in Lemma 4, we have that $\|\widehat{\mathbf{U}}_{k,\cdot}\|_2 \lesssim \|(\widehat{\mathbf{U}}\mathbf{H}_U)_{k,\cdot}\|_2 \leq \|\mathbf{U}_{k,\cdot}\|_2 + \|(\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U})_{k,\cdot}\|_2$. Then, we obtain by Lemma 6 that,

$$\begin{aligned} \|(\boldsymbol{\Psi}_U)_{k,\cdot}\|_2 &\lesssim \left(\omega_k \rho \sqrt{\frac{r}{n}} \log^{3/2} n + \rho^2 \|\mathbf{U}_{k,\cdot}\|_2 + \omega_k \omega \log n \left(\|\mathbf{U}\|_{2,\infty} + \|(\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U})_{k,\cdot}\|_2 \right) \right) \\ &\quad + \left(\|\mathbf{U}_{k,\cdot}\|_2 + \|(\widehat{\mathbf{U}}\mathbf{H}_U - \mathbf{U})_{k,\cdot}\|_2 \right) \left(\rho^2 + \rho \sqrt{\frac{r}{n}} \sqrt{\log n} \right). \end{aligned}$$

Finally, we obtain by Lemma 7 that,

$$\begin{aligned} &\|(\boldsymbol{\Psi}_U)_{k,\cdot}\|_2 \\ &\lesssim \left(\omega_k \rho \sqrt{\frac{r}{n}} \log^{3/2} n + \rho^2 \|\mathbf{U}_{k,\cdot}\|_2 + \omega_k \omega \log n \left(\|\mathbf{U}\|_{2,\infty} + \omega \sqrt{\frac{r}{n}} \log n \right. \right. \\ &\quad \left. \left. + (\rho \sqrt{\frac{r}{n}} + \omega^2) \log n \|\mathbf{U}\|_{2,\infty} \right) \right) \\ &\quad + \left(\|\mathbf{U}_{k,\cdot}\|_2 + \omega_k \sqrt{\frac{r}{n}} \log n \right. \\ &\quad \left. + (\rho \sqrt{\frac{r}{n}} + \omega^2) \log n \|\mathbf{U}_{k,\cdot}\|_2 + \omega_k \omega \log n \|\mathbf{U}\|_{2,\infty} \right) \left(\rho^2 + \rho \sqrt{\frac{r}{n}} \sqrt{\log n} \right) \\ &\lesssim \omega_k \omega \sqrt{\frac{r}{n}} \log^{3/2} n + \left(\rho^2 + \rho \sqrt{\frac{r}{n}} \log n \right) \|\mathbf{U}_{k,\cdot}\|_2 + \omega_k \omega \log n \|\mathbf{U}\|_{2,\infty}, \end{aligned}$$

where, in the last inequality, we use the assumptions that $\omega\sqrt{\log n} \ll 1$, $\rho\sqrt{\log n} \ll 1$, and $r \log n \ll n$ owing to (3.2) and (3.1), and $\omega_k \leq \omega$ as well as $\rho \leq \omega$.

Step 2 – By calculations, we have that

$$\Psi_V = (\widehat{\mathbf{V}}\widehat{\Sigma}\mathbf{H}_U - (T^{-1/2}\mathbf{X})^\top \mathbf{U})\mathbf{\Lambda}^{-1} + \widehat{\mathbf{V}}\widehat{\Sigma}(\mathbf{R}_U - \mathbf{H}_U)\mathbf{\Lambda}^{-1} + (\widehat{\mathbf{V}}\mathbf{R}_V)(\mathbf{\Lambda} - (\mathbf{R}_V)^\top \widehat{\Sigma}\mathbf{R}_U)\mathbf{\Lambda}^{-1}.$$

Since $\|\widehat{\mathbf{V}}_{l,\cdot}\mathbf{R}_V\|_2 = \|\widehat{\mathbf{V}}_{l,\cdot}\|_2$, we obtain that

$$\begin{aligned} & \|(\Psi_V)_{l,\cdot}\|_2 \\ & \lesssim \|(\widehat{\mathbf{V}}\widehat{\Sigma}\mathbf{H}_U - (T^{-1/2}\mathbf{X})^\top \mathbf{U})_{l,\cdot}\|_2 \|\mathbf{\Lambda}^{-1}\|_2 + \|\widehat{\mathbf{V}}_{l,\cdot}\|_2 \left(\|\widehat{\Sigma}(\mathbf{R}_U - \mathbf{H}_U)\mathbf{\Lambda}^{-1}\|_2 + \|(\mathbf{\Lambda} - (\mathbf{R}_V)^\top \widehat{\Sigma}\mathbf{R}_U)\mathbf{\Lambda}^{-1}\|_2 \right) \\ & \stackrel{(i)}{\lesssim} \|(\widehat{\mathbf{V}}\widehat{\Sigma}\mathbf{H}_U - (T^{-1/2}\mathbf{X})^\top \mathbf{U})_{l,\cdot}\|_2 \frac{1}{\sigma_r} + \|\widehat{\mathbf{V}}_{l,\cdot}\|_2 \left(\sigma_r \rho^2 \frac{1}{\sigma_r} + \frac{1}{\sigma_r} \left(\sigma_r \rho^2 + \sigma_r \rho \sqrt{\frac{r}{n}} \log n \right) \right) \\ & \lesssim \|(\widehat{\mathbf{V}}\widehat{\Sigma}\mathbf{H}_U - (T^{-1/2}\mathbf{X})^\top \mathbf{U})_{l,\cdot}\|_2 \frac{1}{\sigma_r} + \|\widehat{\mathbf{V}}_{l,\cdot}\|_2 \left(\rho^2 + \rho \sqrt{\frac{r}{n}} \log n \right), \end{aligned}$$

where (i) is owing to the inequalities in Lemmas 4 and 5, and the fact that $\|\mathbf{\Lambda} - (\mathbf{R}_V)^\top \widehat{\Sigma}\mathbf{R}_U\|_2 = \|(\mathbf{\Lambda} - (\mathbf{R}_V)^\top \widehat{\Sigma}\mathbf{R}_U)^\top\|_2 = \|\mathbf{\Lambda} - (\mathbf{R}_U)^\top \widehat{\Sigma}\mathbf{R}_V\|_2$. Using $\sigma_i(\mathbf{H}_V) \asymp 1$ in Lemma 4, we have that $\|\widehat{\mathbf{V}}_{l,\cdot}\|_2 \lesssim \|(\widehat{\mathbf{V}}\mathbf{H}_V)_{l,\cdot}\|_2 \leq \|\mathbf{V}_{l,\cdot}\|_2 + \|(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})_{l,\cdot}\|_2$. Then, we obtain by Lemma 6 that,

$$\begin{aligned} \|(\Psi_V)_{l,\cdot}\|_2 & \lesssim \sigma_r \rho^2 \left(\sqrt{\frac{r}{n}} \log^{3/2} n + \log n \|\mathbf{V}_{l,\cdot}\|_2 + \log n \|(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})_{l,\cdot}\|_2 \right) \\ & \quad + \left(\|\mathbf{V}_{l,\cdot}\|_2 + \|(\widehat{\mathbf{V}}\mathbf{H}_V - \mathbf{V})_{l,\cdot}\|_2 \right) \left(\rho^2 + \rho \sqrt{\frac{r}{n}} \log n \right). \end{aligned}$$

Finally, we obtain by Lemma 7 that,

$$\begin{aligned} & \|(\Psi_V)_{l,\cdot}\|_2 \\ & \lesssim \rho^2 \left(\sqrt{\frac{r}{n}} \log^{3/2} n + \log n \|\mathbf{V}_{l,\cdot}\|_2 + \log n \left(\rho \sqrt{\frac{r}{n}} \log n + (\rho^2 + \rho \sqrt{\frac{r}{n}} \log n) \|\mathbf{V}_{l,\cdot}\|_2 \right) \right) \\ & \quad + \left(\|\mathbf{V}_{l,\cdot}\|_2 + \rho \sqrt{\frac{r}{n}} \log n \right) \left(\rho^2 + \rho \sqrt{\frac{r}{n}} \log n \right) \\ & \lesssim \rho^2 \sqrt{\frac{r}{n}} \log^{3/2} n + \left(\rho^2 \log n + \rho \sqrt{\frac{r}{n}} \log n \right) \|\mathbf{V}_{l,\cdot}\|_2, \end{aligned}$$

where, in the last inequality, we use the assumptions that, $\rho \sqrt{\log n} \ll 1$ and $r \log n \ll n$ owing to (3.2) and (A.23).

B Proof of Corollary 1

For the first-order terms \mathbf{G}_U and \mathbf{G}_V in Theorem 1, we obtain by Lemma 3 that, with probability at least $1 - O(n^{-2})$,

$$\begin{aligned} \|(\mathbf{G}_V)_{l,\cdot}\|_2 & = \|(T^{-1/2}\mathbf{E}^\top \mathbf{U}\mathbf{\Lambda}^{-1})_{l,\cdot}\|_2 = \|(T^{-1/2}\mathbf{Z}^\top \Sigma_\varepsilon^{1/2} \mathbf{U}\mathbf{\Lambda}^{-1})_{l,\cdot}\|_2 \\ & = \|T^{-1/2}(\mathbf{Z}^\top)_{l,\cdot} \Sigma_\varepsilon^{1/2} \mathbf{U}\mathbf{\Lambda}^{-1}\|_2 \lesssim \frac{1}{\sqrt{T}} \|\Sigma_\varepsilon^{1/2} \mathbf{U}\mathbf{\Lambda}^{-1}\|_{\mathbb{F}} \log n \\ & \lesssim \frac{1}{\sqrt{T}} \|\Sigma_\varepsilon^{1/2}\|_2 \|\mathbf{U}\|_{\mathbb{F}} \frac{1}{\sigma_r} \log n = \rho \sqrt{\frac{r}{n}} \log n, \end{aligned}$$

and similarly we obtain that

$$\|(\mathbf{G}_U)_{k,\cdot}\|_2 \lesssim \frac{1}{\vartheta_k \sqrt{T}} \sqrt{r} \log n.$$

Step 1 – Derive the error bounds for factors.

Using the upper bounds for $\|\widehat{\mathbf{U}}\mathbf{R}_U - \mathbf{U}\|_{\mathbb{F}}$ and $\|\widehat{\mathbf{V}}\mathbf{R}_V - \mathbf{V}\|_{\mathbb{F}}$ in Lemma 4 and the upper bound for $\|(\mathbf{R}_U)^\top \widehat{\boldsymbol{\Sigma}}\mathbf{R}_V - \boldsymbol{\Lambda}\|_2$ in Lemma 5, we have that

$$\frac{1}{T}\|\widehat{\mathbf{F}} - \mathbf{F}\mathbf{J}(\mathbf{R}_V)^\top\|_{\mathbb{F}}^2 = \|\widehat{\mathbf{V}}\mathbf{R}_V(\mathbf{R}_V)^\top - \mathbf{V}(\mathbf{R}_V)^\top\|_{\mathbb{F}}^2 \lesssim \frac{1}{\theta^2} \frac{n}{T} r.$$

For $\widehat{\mathbf{V}}\mathbf{R}_V - \mathbf{V} = \mathbf{G}_V + \boldsymbol{\Psi}_V$, using the fact that $\|\mathbf{V}\|_{2,\infty} \lesssim \sqrt{\frac{\log n}{T}}$ in Lemma 2, we obtain that,

$$\begin{aligned} \|\boldsymbol{\Psi}_V\|_{2,\infty} &= \sup_{1 \leq l \leq T} \|(\boldsymbol{\Psi}_V)_{l,\cdot}\|_2 \lesssim \rho^2 \sqrt{\frac{r}{n}} \log^{3/2} n + \left(\rho^2 + \rho \sqrt{\frac{r}{n}}\right) \log n \|\mathbf{V}\|_{2,\infty} \\ &\lesssim \rho^2 \sqrt{\frac{r}{n}} \log^{3/2} n + \left(\rho^2 + \rho \sqrt{\frac{r}{n}} \log n\right) \sqrt{\frac{\log n}{T}} \lesssim \rho \sqrt{\frac{r}{n}} \log n, \end{aligned}$$

where the last inequality is because $\rho\sqrt{\log n} \ll 1$ and $\log n \ll T$. Since $\|\mathbf{G}_V\|_{2,\infty} \lesssim \rho\sqrt{\frac{r}{n}} \log n$, we obtain that $\|\widehat{\mathbf{V}}\mathbf{R}_V - \mathbf{V}\|_{2,\infty} \leq \|\mathbf{G}_V\|_{2,\infty} + \|\boldsymbol{\Psi}_V\|_{2,\infty} \lesssim \rho\sqrt{\frac{r}{n}} \log n$, which implies the row-wise error bound for $\|(\widehat{\mathbf{F}} - \mathbf{F}\mathbf{R}_F)_{t,\cdot}\|_2$.

Step 2 - Derive the error bounds for factor loadings.

For $\widehat{\mathbf{U}}\mathbf{R}_U - \mathbf{U} = \mathbf{G}_U + \boldsymbol{\Psi}_U$, the problem boils down to comparing the upper bounds of $\|(\mathbf{G}_U)_{k,\cdot}\|_2$ and $\|(\boldsymbol{\Psi}_U)_{k,\cdot}\|_2$. For the first term in the upper bound of $\|(\boldsymbol{\Psi}_U)_{k,\cdot}\|_2$, we obtain by $\omega\sqrt{\log n} \ll 1$ that $\omega_k \omega \sqrt{\frac{r}{n}} \log^{3/2} n = \omega_k \sqrt{\frac{r}{n}} \log n (\omega\sqrt{\log n}) \ll \omega_k \sqrt{\frac{r}{n}} \log n$. Next, for the second and the third terms in the upper bound of $\|(\boldsymbol{\Psi}_U)_{k,\cdot}\|_2$, we have that

$$\frac{(\rho^2 + \rho\sqrt{\frac{r}{n}} \log n) \|\mathbf{U}_{k,\cdot}\|_2}{\omega_k \sqrt{\frac{r}{n}} \log n} = \left(\frac{\rho}{\sqrt{\frac{r}{n}} \log n} + 1\right) \frac{\rho}{\omega_k} \|\mathbf{U}_{k,\cdot}\|_2 \leq (\rho\sqrt{n} + 1) \frac{\rho}{\omega_k} \|\bar{\mathbf{U}}_{k,\cdot}\|_2 \lesssim 1,$$

and

$$\frac{\omega_k \omega \log n \|\mathbf{U}\|_{2,\infty}}{\omega_k \sqrt{\frac{r}{n}} \log n} = \omega \sqrt{\frac{n}{r}} \|\mathbf{U}\|_{2,\infty} \leq \omega \sqrt{n} \|\bar{\mathbf{U}}\|_{2,\infty} \lesssim 1.$$

So we obtain that

$$\|(\widehat{\mathbf{U}}\mathbf{R}_U - \mathbf{U})_{k,\cdot}\|_2 \lesssim \omega_k \sqrt{\frac{r}{n}} \log n \quad \text{and} \quad \|\widehat{\mathbf{U}}\mathbf{R}_U - \mathbf{U}\|_{2,\infty} \lesssim \omega \sqrt{\frac{r}{n}} \log n.$$

The row-wise error bound for $\widehat{\mathbf{B}} - \mathbf{B}\mathbf{R}_B$ follows from combining the error bounds of $\widehat{\mathbf{U}}\mathbf{R}_U - \mathbf{U}$ and $(\mathbf{R}_U)^\top \widehat{\boldsymbol{\Sigma}}\mathbf{R}_V - \boldsymbol{\Lambda}$. Then the averaged estimation error bound follows from averaging the row-wise error bounds.

C Inference for the factors and the factor loadings

C.1 Proof of Corollary 2

Step 1 - bounding $\mathbb{P}((\widehat{\mathbf{V}}\mathbf{R}_V\mathbf{J}^{-1} - \frac{1}{\sqrt{T}}\mathbf{F})_{t,\cdot} \in \mathcal{C})$.

It follows from Lemma 2 that, there exists a $\sigma(\mathbf{F})$ -measurable matrix \mathbf{J} satisfying that $\mathbf{V} = \frac{1}{\sqrt{T}}\mathbf{F}\mathbf{J}$ and \mathbf{J} is a $r \times r$ invertible matrix satisfying that $\sigma_i(\mathbf{J}) \asymp 1$ for $i = 1, 2, \dots, r$. We define

$$\mathbf{R}_F := \mathbf{J}\mathbf{R}_V^\top,$$

then we obtain by Theorem 1 that

$$\widehat{\mathbf{F}} - \mathbf{F}\mathbf{R}_F = \sqrt{T}(\widehat{\mathbf{V}}\mathbf{R}_V\mathbf{J}^{-1} - \mathbf{V}\mathbf{J}^{-1})\mathbf{R}_F = \sqrt{T}(\mathbf{G}_V + \boldsymbol{\Psi}_V)\mathbf{J}^{-1}\mathbf{R}_F = (\sqrt{T}\mathbf{G}_V\mathbf{J}^{-1} + \sqrt{T}\boldsymbol{\Psi}_V\mathbf{J}^{-1})\mathbf{R}_F.$$

We also note that $\Sigma_{F,t} = \mathbf{R}_V \mathbf{\Lambda}^{-1} \mathbf{U}^\top \Sigma_\varepsilon \mathbf{U} \mathbf{\Lambda}^{-1} \mathbf{R}_V^\top = \mathbf{R}_V \mathbf{J}^\top \Sigma^{-1} \bar{\mathbf{U}}^\top \Sigma_\varepsilon \bar{\mathbf{U}} \Sigma^{-1} \mathbf{J} \mathbf{R}_V^\top = \mathbf{R}_F^\top (\Sigma^{-1} \bar{\mathbf{U}}^\top \Sigma_\varepsilon \bar{\mathbf{U}} \Sigma^{-1}) \mathbf{R}_F$. Denote by $(\bar{\mathbf{U}}^\top \Sigma_\varepsilon \bar{\mathbf{U}})^{-1/2}$ the matrix such that $(\bar{\mathbf{U}}^\top \Sigma_\varepsilon \bar{\mathbf{U}})^{-1/2} (\bar{\mathbf{U}}^\top \Sigma_\varepsilon \bar{\mathbf{U}}) (\bar{\mathbf{U}}^\top \Sigma_\varepsilon \bar{\mathbf{U}})^{-1/2} = \mathbf{I}_r$. So, we have that

$$\begin{aligned} & \sup_{\mathcal{C} \in \mathcal{C}^r} \left| \mathbb{P}((\hat{\mathbf{F}} - \mathbf{F} \mathbf{R}_F)_{t,\cdot} \in \mathcal{C}) - \mathbb{P}(\mathcal{N}(0, \Sigma_{F,t}) \in \mathcal{C}) \right| \\ &= \sup_{\mathcal{C} \in \mathcal{C}^r} \left| \mathbb{P}(\sqrt{T}(\mathbf{G}_V \mathbf{J}^{-1} + \Psi_V \mathbf{J}^{-1})_{t,\cdot} \in \mathcal{C}) - \mathbb{P}(\mathcal{N}(0, \Sigma^{-1} \bar{\mathbf{U}}^\top \Sigma_\varepsilon \bar{\mathbf{U}} \Sigma^{-1}) \in \mathcal{C}) \right| \\ &= \sup_{\mathcal{C} \in \mathcal{C}^r} \left| \mathbb{P}(\sqrt{T}(\bar{\mathbf{U}}^\top \Sigma_\varepsilon \bar{\mathbf{U}})^{-1/2} \Sigma((\mathbf{G}_V \mathbf{J}^{-1} + \Psi_V \mathbf{J}^{-1})_{t,\cdot})^\top \in \mathcal{C}) - \mathbb{P}(\mathcal{N}(0, \mathbf{I}_r) \in \mathcal{C}) \right|, \end{aligned}$$

where the last equality is because \mathcal{C}^r is invariant under nondegenerate linear transformation and the matrix $(\bar{\mathbf{U}}^\top \Sigma_\varepsilon \bar{\mathbf{U}})^{-1/2} \Sigma$ is not degenerate.

To study $\sqrt{T}(\bar{\mathbf{U}}^\top \Sigma_\varepsilon \bar{\mathbf{U}})^{-1/2} \Sigma((\mathbf{G}_V \mathbf{J}^{-1} + \Psi_V \mathbf{J}^{-1})_{t,\cdot})^\top$, we define the vector $\mathbf{K}_{V,t}$ and the scalar ε_1 , respectively, as below:

$$\mathbf{K}_{V,t} := \sqrt{T}(\bar{\mathbf{U}}^\top \Sigma_\varepsilon \bar{\mathbf{U}})^{-1/2} \Sigma((\mathbf{G}_V \mathbf{J}^{-1})_{t,\cdot})^\top \quad \text{and} \quad \varepsilon_1 := \left\| \sqrt{T}(\bar{\mathbf{U}}^\top \Sigma_\varepsilon \bar{\mathbf{U}})^{-1/2} \Sigma((\Psi_V \mathbf{J}^{-1})_{t,\cdot})^\top \right\|_2.$$

Then we have the following inequality

$$\mathbb{P}(\mathbf{K}_{V,t} \in \mathcal{C}^{-\varepsilon_1}) \leq \mathbb{P}(\sqrt{T}(\bar{\mathbf{U}}^\top \Sigma_\varepsilon \bar{\mathbf{U}})^{-1/2} \Sigma((\mathbf{G}_V \mathbf{J}^{-1} + \Psi_V \mathbf{J}^{-1})_{t,\cdot})^\top \in \mathcal{C}) \leq \mathbb{P}(\mathbf{K}_{V,t} \in \mathcal{C}^{\varepsilon_1}).$$

For any non-empty convex set $\mathcal{C} \in \mathcal{C}^r$, we define $\mathcal{C}^\varepsilon := \{x \in \mathbb{R}^r : \delta_{\mathcal{C}}(x) \leq \varepsilon\}$ where $\delta_{\mathcal{C}}(x)$ is the signed distance of the point x to the set \mathcal{C} . We will show later that ε_1 is negligible.

Step 2 – Proving the approximate Gaussianity of $\mathbf{K}_{V,t}$.

In this step, we establish the upper bound for the constant τ_V defined below,

$$\tau_V := \sup_{\mathcal{C}_1 \in \mathcal{C}^r} |\mathbb{P}(\mathbf{K}_{V,t} \in \mathcal{C}_1) - \mathbb{P}(\mathcal{N}(0, \mathbf{I}_r) \in \mathcal{C}_1)|.$$

We obtain by Theorem 1 that $\mathbf{G}_V := T^{-1/2} \mathbf{E}^\top \mathbf{U} \mathbf{\Lambda}^{-1}$. It follows from Lemma 2 that $\mathbf{J} = \Sigma \mathbf{Q} \mathbf{\Lambda}^{-1}$ and $\mathbf{U} = \bar{\mathbf{U}} \mathbf{Q}$, where $\mathbf{Q} \in \mathcal{O}^{r \times r}$ is a rotation matrix, i.e., $\mathbf{Q} \mathbf{Q}^\top = \mathbf{Q}^\top \mathbf{Q} = \mathbf{I}_r$. So, we obtain that $\mathbf{U} \mathbf{Q}^{-1} = \bar{\mathbf{U}}$ and

$$\begin{aligned} \sqrt{T} \mathbf{G}_V \mathbf{J}^{-1} &= \mathbf{E}^\top \mathbf{U} \mathbf{\Lambda}^{-1} \mathbf{\Lambda} \mathbf{Q}^{-1} \Sigma^{-1} = \mathbf{E}^\top \mathbf{U} \mathbf{Q}^{-1} \Sigma^{-1} \\ &= \mathbf{E}^\top \bar{\mathbf{U}} \Sigma^{-1} = \mathbf{Z}^\top \Sigma_\varepsilon^{1/2} \bar{\mathbf{U}} \Sigma^{-1} = \mathbf{Z}^\top \mathbf{H} \quad \text{with} \quad \mathbf{H} := \Sigma_\varepsilon^{1/2} \bar{\mathbf{U}} \Sigma^{-1}. \end{aligned}$$

Going back to our target vector, we have that

$$\begin{aligned} \mathbf{K}_{V,t} &= (\bar{\mathbf{U}}^\top \Sigma_\varepsilon \bar{\mathbf{U}})^{-1/2} \Sigma(\sqrt{T}(\mathbf{G}_V \mathbf{J}^{-1})_{t,\cdot})^\top = (\bar{\mathbf{U}}^\top \Sigma_\varepsilon \bar{\mathbf{U}})^{-1/2} \Sigma((\mathbf{Z}^\top \mathbf{H})_{t,\cdot})^\top \\ &= (\bar{\mathbf{U}}^\top \Sigma_\varepsilon \bar{\mathbf{U}})^{-1/2} \Sigma \mathbf{H}^\top \mathbf{Z}_{\cdot,t} = (\bar{\mathbf{U}}^\top \Sigma_\varepsilon \bar{\mathbf{U}})^{-1/2} \Sigma \Sigma^{-1} (\Sigma_\varepsilon^{1/2} \bar{\mathbf{U}})^\top \mathbf{Z}_{\cdot,t} \\ &= \sum_{k=1}^N \mathbf{Z}_{k,t} (\bar{\mathbf{U}}^\top \Sigma_\varepsilon \bar{\mathbf{U}})^{-1/2} ((\Sigma_\varepsilon^{1/2} \bar{\mathbf{U}})_{k,\cdot})^\top. \end{aligned}$$

Note that both $\bar{\mathbf{U}}$ and Σ_ε are deterministic. So, we obtain that the vector $\mathbf{K}_{V,t}$ is the summation of N independent and mean zero vectors, and the covariance matrix of $\mathbf{K}_{V,t}$ is given by

$$\begin{aligned} \text{cov}(\mathbf{K}_{V,t}) &= \sum_{k=1}^N (\bar{\mathbf{U}}^\top \Sigma_\varepsilon \bar{\mathbf{U}})^{-1/2} ((\Sigma_\varepsilon^{1/2} \bar{\mathbf{U}})_{k,\cdot})^\top (\Sigma_\varepsilon^{1/2} \bar{\mathbf{U}})_{k,\cdot} (\bar{\mathbf{U}}^\top \Sigma_\varepsilon \bar{\mathbf{U}})^{-1/2} \\ &= (\bar{\mathbf{U}}^\top \Sigma_\varepsilon \bar{\mathbf{U}})^{-1/2} (\bar{\mathbf{U}}^\top \Sigma_\varepsilon \bar{\mathbf{U}}) (\bar{\mathbf{U}}^\top \Sigma_\varepsilon \bar{\mathbf{U}})^{-1/2} = \mathbf{I}_r. \end{aligned}$$

Given the above analysis of $\mathbf{K}_{V,t}$, we are now ready to bound the term τ_V :

(i) When all entries of \mathbf{Z} are Gaussian, we have that, the vector $\mathbf{K}_{V,t}$ is also a Gaussian vector because it is a linear combination of Gaussian vectors. Since $\text{cov}(\mathbf{K}_{V,t}) = \mathbf{I}_r$, we obtain that the law of $\mathbf{K}_{V,t}$ is

the r -dimensional standard Gaussian law $\mathcal{N}(0, \mathbf{I}_r)$. Thus, we have that $\tau_V = 0$ when all entries of \mathbf{Z} are Gaussian.

(ii) When the entries of \mathbf{Z} are sub-Gaussian, we obtain by the Berry-Esseen theorem (Theorem 1.1 in Raič (2019)) that

$$\tau_V \lesssim r^{1/4} \tau_V^{(0)},$$

and

$$\begin{aligned} \tau_V^{(0)} &= \sum_{k=1}^N \mathbb{E} \left[\left\| \mathbf{Z}_{k,t} (\bar{\mathbf{U}}^\top \boldsymbol{\Sigma}_\varepsilon \bar{\mathbf{U}})^{-1/2} ((\boldsymbol{\Sigma}_\varepsilon^{1/2} \bar{\mathbf{U}})_{k,\cdot})^\top \right\|_2^3 \right] = \sum_{k=1}^N \left\| (\boldsymbol{\Sigma}_{V,t})^{-1/2} (\mathbf{H}_{k,\cdot})^\top \right\|_2^3 \mathbb{E} \left[\left\| \mathbf{Z}_{k,t} \right\|_2^3 \right] \\ &\stackrel{(i)}{\lesssim} \sum_{k=1}^N \left\| (\bar{\mathbf{U}}^\top \boldsymbol{\Sigma}_\varepsilon \bar{\mathbf{U}})^{-1/2} ((\boldsymbol{\Sigma}_\varepsilon^{1/2} \bar{\mathbf{U}})_{k,\cdot})^\top \right\|_2^3 \mathbb{E} \left[\left\| \mathbf{Z}_{k,t} \right\|_2^3 \right] \\ &\leq \max_{1 \leq k \leq N} \left\| (\bar{\mathbf{U}}^\top \boldsymbol{\Sigma}_\varepsilon \bar{\mathbf{U}})^{-1/2} ((\boldsymbol{\Sigma}_\varepsilon^{1/2} \bar{\mathbf{U}})_{k,\cdot})^\top \right\|_2 \cdot \sum_{k=1}^N \left\| (\bar{\mathbf{U}}^\top \boldsymbol{\Sigma}_\varepsilon \bar{\mathbf{U}})^{-1/2} ((\boldsymbol{\Sigma}_\varepsilon^{1/2} \bar{\mathbf{U}})_{k,\cdot})^\top \right\|_2^2 \\ &\leq (\lambda_{\min}(\bar{\mathbf{U}}^\top \boldsymbol{\Sigma}_\varepsilon \bar{\mathbf{U}}))^{-1/2} \max_{1 \leq k \leq N} \left\| (\boldsymbol{\Sigma}_\varepsilon^{1/2} \bar{\mathbf{U}})_{k,\cdot} \right\|_2 \cdot \text{tr} \left[(\bar{\mathbf{U}}^\top \boldsymbol{\Sigma}_\varepsilon \bar{\mathbf{U}})^{-1/2} \bar{\mathbf{U}}^\top \boldsymbol{\Sigma}_\varepsilon \bar{\mathbf{U}} (\bar{\mathbf{U}}^\top \boldsymbol{\Sigma}_\varepsilon \bar{\mathbf{U}})^{-1/2} \right] \\ &= (\lambda_{\min}(\bar{\mathbf{U}}^\top \boldsymbol{\Sigma}_\varepsilon \bar{\mathbf{U}}))^{-1/2} \left\| \boldsymbol{\Sigma}_\varepsilon^{1/2} \bar{\mathbf{U}} \right\|_{2,\infty} r, \end{aligned}$$

where (i) is because the entries of \mathbf{Z} are sub-Gaussian and their sub-Gaussian norms satisfy $\|Z_{i,t}\|_{\psi_2} = O(1)$. Then, since $\bar{\mathbf{U}}$ has full column rank, we have that

$$\lambda_{\min}(\bar{\mathbf{U}}^\top \boldsymbol{\Sigma}_\varepsilon \bar{\mathbf{U}}) = \inf_{\|\mathbf{x}\|_2=1} \mathbf{x}^\top \bar{\mathbf{U}}^\top \boldsymbol{\Sigma}_\varepsilon \bar{\mathbf{U}} \mathbf{x} = \inf_{\|\mathbf{x}\|_2=1} (\bar{\mathbf{U}} \mathbf{x})^\top \boldsymbol{\Sigma}_\varepsilon \bar{\mathbf{U}} \mathbf{x} = \inf_{\|\mathbf{x}\|_2=1} \mathbf{x}^\top \boldsymbol{\Sigma}_\varepsilon \mathbf{x} = \lambda_{\min}(\boldsymbol{\Sigma}_\varepsilon).$$

Recall that $\rho := \sqrt{\|\boldsymbol{\Sigma}_\varepsilon\|_2} \sqrt{n} / (\sigma_r \sqrt{T})$. So, we conclude that

$$\tau_V^{(0)} \lesssim \frac{1}{\lambda_{\min}(\boldsymbol{\Sigma}_\varepsilon)} \left\| \boldsymbol{\Sigma}_\varepsilon^{1/2} \bar{\mathbf{U}} \right\|_{2,\infty} r = \sqrt{\kappa_\varepsilon} \frac{\left\| \boldsymbol{\Sigma}_\varepsilon^{1/2} \bar{\mathbf{U}} \right\|_{2,\infty}}{\left\| \boldsymbol{\Sigma}_\varepsilon^{1/2} \right\|_2} r, \quad \text{and} \quad \tau_V \lesssim r^{1/4} \tau_V^{(0)} \lesssim \sqrt{\kappa_\varepsilon} r^{5/4} \frac{\left\| \boldsymbol{\Sigma}_\varepsilon^{1/2} \bar{\mathbf{U}} \right\|_{2,\infty}}{\left\| \boldsymbol{\Sigma}_\varepsilon^{1/2} \right\|_2},$$

where we use the fact that $\|\boldsymbol{\Sigma}_\varepsilon\|_2 = \lambda_{\max}(\boldsymbol{\Sigma}_\varepsilon)$ and $\kappa_\varepsilon = \lambda_{\max}(\boldsymbol{\Sigma}_\varepsilon) / \lambda_{\min}(\boldsymbol{\Sigma}_\varepsilon)$ is the condition number of $\boldsymbol{\Sigma}_\varepsilon$.

Step 3 – Establishing the validity of confidence regions.

Since we obtained in Step 2 that $|\mathbb{P}(\mathbf{K}_{V,t} \in \mathcal{C}_1) - \mathbb{P}(\mathcal{N}(0, \mathbf{I}_r) \in \mathcal{C}_1)| \leq \tau_V$ holds for any convex set \mathcal{C}_1 , we let \mathcal{C}_1 be $\mathcal{C}^{-\varepsilon_1}$ and $\mathcal{C}^{\varepsilon_1}$, respectively to obtain the probabilities bounds as below

$$\mathbb{P}(\mathcal{N}(0, \mathbf{I}_r) \in \mathcal{C}^{-\varepsilon_1}) - \tau_V \leq \mathbb{P}(\sqrt{T}(\bar{\mathbf{U}}^\top \boldsymbol{\Sigma}_\varepsilon \bar{\mathbf{U}})^{-1/2} \boldsymbol{\Sigma}((\mathbf{G}_V \mathbf{J}^{-1} + \boldsymbol{\Psi}_V \mathbf{J}^{-1})_{t,\cdot})^\top \in \mathcal{C}) \leq \mathbb{P}(\mathcal{N}(0, \mathbf{I}_r) \in \mathcal{C}^{\varepsilon_1}) + \tau_V.$$

Next, by Theorem 1.2 in Raič (2019) for multi-dimensional standard Gaussian vector, we obtain

$$\begin{aligned} &\mathbb{P}(\mathcal{N}(0, \mathbf{I}_r) \in \mathcal{C}) - (K_1 r^{1/4} + K_0) \varepsilon_1 - \tau_V \\ &\leq \mathbb{P}(\sqrt{T}(\bar{\mathbf{U}}^\top \boldsymbol{\Sigma}_\varepsilon \bar{\mathbf{U}})^{-1/2} \boldsymbol{\Sigma}((\mathbf{G}_V \mathbf{J}^{-1} + \boldsymbol{\Psi}_V \mathbf{J}^{-1})_{t,\cdot})^\top \in \mathcal{C}) \\ &\leq \mathbb{P}(\mathcal{N}(0, \mathbf{I}_r) \in \mathcal{C}) + (K_1 r^{1/4} + K_0) \varepsilon_1 + \tau_V. \end{aligned}$$

Since the above inequality holds for any convex set $\mathcal{C} \in \mathcal{C}^r$, we obtain that

$$\begin{aligned} &\sup_{\mathcal{C} \in \mathcal{C}^r} \left| \mathbb{P}((\hat{\mathbf{F}} - \mathbf{F} \mathbf{R}_F)_{t,\cdot} \in \mathcal{C}) - \mathbb{P}(\mathcal{N}(0, \boldsymbol{\Sigma}_{V,t}) \in \mathcal{C}) \right| \\ &= \sup_{\mathcal{C} \in \mathcal{C}^r} \left| \mathbb{P}(\sqrt{T}(\bar{\mathbf{U}}^\top \boldsymbol{\Sigma}_\varepsilon \bar{\mathbf{U}})^{-1/2} \boldsymbol{\Sigma}((\mathbf{G}_V \mathbf{J}^{-1} + \boldsymbol{\Psi}_V \mathbf{J}^{-1})_{t,\cdot})^\top \in \mathcal{C}) - \mathbb{P}(\mathcal{N}(0, \mathbf{I}_r) \in \mathcal{C}) \right| \\ &\leq (K_1 r^{1/4} + K_0) \varepsilon_1 + \tau_V \lesssim r^{1/4} \varepsilon_1 + \tau_V. \end{aligned}$$

Finally, we derive the upper bound for $\varepsilon_1 := \frac{1}{\sqrt{T}} \left\| (\boldsymbol{\Sigma}_{V,t})^{-1/2} ((\boldsymbol{\Psi}_V \mathbf{J}^{-1})_{t,\cdot})^\top \right\|_2$.

Using the fact that $\sigma_i(\mathbf{J}) \asymp 1$ for $i = 1, 2, \dots, r$ in Lemma 2, we obtain that, with probability at least $1 - O(n^{-2})$, it holds

$$\varepsilon_1 = \left\| \sqrt{T} (\bar{\mathbf{U}}^\top \boldsymbol{\Sigma}_\varepsilon \bar{\mathbf{U}})^{-1/2} \boldsymbol{\Sigma} ((\boldsymbol{\Psi}_V \mathbf{J}^{-1})_{t,\cdot})^\top \right\|_2 \leq \frac{\sqrt{T} \|(\boldsymbol{\Psi}_V \mathbf{J}^{-1})_{t,\cdot} \boldsymbol{\Sigma}\|_2}{(\lambda_{\min}(\boldsymbol{\Sigma}_\varepsilon))^{1/2}} = \frac{\sqrt{\kappa_\varepsilon} \sqrt{T}}{\|\boldsymbol{\Sigma}_\varepsilon^{1/2}\|_2} \|(\boldsymbol{\Psi}_V \mathbf{J}^{-1})_{t,\cdot} \boldsymbol{\Sigma}\|_2.$$

Then, using the upper bound $\|\mathbf{V}\|_{2,\infty} \lesssim \sqrt{\frac{\log n}{T}}$ obtained in Lemma 2 and the formula of $\boldsymbol{\Psi}_V$ in the proof of Theorem 1, we obtain that

$$\begin{aligned} \|(\boldsymbol{\Psi}_V \mathbf{J}^{-1})_{t,\cdot} \boldsymbol{\Sigma}\|_2 &\lesssim \sigma_r \rho^2 \sqrt{\frac{r}{n}} \log^{3/2} n + \sigma_r \left(\rho^2 + \rho \sqrt{\frac{r}{n}} \log n \right) \|\mathbf{V}_{l,\cdot}\|_2 \\ &\lesssim \sigma_r \rho^2 \sqrt{\frac{r}{n}} \log^{3/2} n + \sigma_r \left(\rho^2 + \rho \sqrt{\frac{r}{n}} \log n \right) \sqrt{\frac{\log n}{T}}. \end{aligned}$$

So, we obtain that, with probability at least $1 - O(n^{-2})$, it holds

$$\begin{aligned} r^{1/4} \varepsilon_1 &\lesssim r^{1/4} \frac{\sqrt{\kappa_\varepsilon} \sqrt{T} \sigma_r}{\|\boldsymbol{\Sigma}_\varepsilon^{1/2}\|_2} \rho \cdot \left(\rho \sqrt{\frac{r}{n}} \log^{3/2} n + \left(\rho + \sqrt{\frac{r}{n}} \log n \right) \sqrt{\frac{\log n}{T}} \right) \\ &= r^{1/4} \sqrt{\kappa_\varepsilon} \sqrt{n} \left(\rho \sqrt{\frac{r}{n}} \log^{3/2} n + \left(\rho + \sqrt{\frac{r}{n}} \log n \right) \sqrt{\frac{\log n}{T}} \right) \\ &= \sqrt{\kappa_\varepsilon} \left(\rho + (\rho \sqrt{n} + 1) \frac{1}{\sqrt{T}} \right) r \log^{3/2} n \\ &\lesssim \sqrt{\kappa_\varepsilon} \left(\rho \sqrt{\frac{n}{T}} + \frac{1}{\sqrt{T}} \right) r \log^{3/2} n. \end{aligned}$$

Thus, by a standard conditioning argument, we conclude that

$$\begin{aligned} &\sup_{\mathcal{C} \in \mathcal{C}^r} \left| \mathbb{P}((\widehat{\mathbf{V}} \mathbf{R}_F - \mathbf{F})_{t,\cdot} \in \mathcal{C}) - \mathbb{P}(\mathcal{N}(0, \boldsymbol{\Sigma}_{V,t}) \in \mathcal{C}) \right| \\ &\lesssim \sqrt{\kappa_\varepsilon} \left(\rho \sqrt{\frac{n}{T}} + \frac{1}{\sqrt{T}} \right) r \log^{3/2} n + \tau_V + O(n^{-2}), \end{aligned}$$

which is our desired result. The sufficient conditions for $\sqrt{\kappa_\varepsilon} \left(\rho \sqrt{\frac{n}{T}} + \frac{1}{\sqrt{T}} \right) r \log^{3/2} n \lesssim \delta$ to hold can be rewritten as two inequalities as follows: $C_0 n^{-2} \leq \delta$, $\sqrt{\kappa_\varepsilon} r^{5/4} \|\boldsymbol{\Sigma}_\varepsilon^{1/2} \bar{\mathbf{U}}\|_{2,\infty} \leq \delta \|\boldsymbol{\Sigma}_\varepsilon^{1/2}\|_2$,

$$C_0 \sqrt{\kappa_\varepsilon} \frac{\sqrt{\|\boldsymbol{\Sigma}_\varepsilon\|_2}}{\sigma_r} \frac{n}{T} r \log^{3/2} n \leq \delta, \quad \text{and} \quad C_0 \kappa_\varepsilon r^2 \log^3 n \ll \delta^2 T.$$

C.2 Proof of Corollary 3

The proof is similar to that for the factors. So we omit the proof due to the limit of space.

D Proof of Theorem 2: Factor test

D.1 Some useful lemmas

To prove Theorem 2, we collect some useful lemmas as preparations. Recall that, we already showed in Lemma 2 that $\mathbf{V} = \frac{1}{\sqrt{T}} \mathbf{F} \mathbf{J}$, implying that $\mathbf{V}_{S,\cdot} = \frac{1}{\sqrt{T}} \mathbf{F}_{S,\cdot} \mathbf{J}$ and thus the column space of $\mathbf{F}_{S,\cdot}$ is closely related to that of $\mathbf{V}_{S,\cdot}$. Then, we have the following lemma for some properties of $\mathbf{V}_{S,\cdot}$.

Lemma 8. Suppose that Assumptions 1 and 3 hold. Assume that (A.1), i.e., $r + \log n \ll T$, and

$$r + \log n \ll |S|, \quad (\text{D.1})$$

then we have that, there exists a $\sigma(\mathbf{F})$ -measurable event \mathcal{E}_S with $\mathbb{P}(\mathcal{E}_S) > 1 - O(n^{-2})$, such that, the following properties hold when $\mathcal{E}_0 \cap \mathcal{E}_S$ happens:

- (i) $\sigma_i(\mathbf{V}_{S,\cdot}) \asymp \sqrt{\frac{|S|}{T}}$ for $i = 1, 2, \dots, r$, and $\text{rank}(\mathbf{V}_{S,\cdot}) = r$, i.e., $\mathbf{V}_{S,\cdot}$ has full column rank.
- (ii) $\sigma_i((\mathbf{V}_{S,\cdot})^+) \asymp \sqrt{\frac{T}{|S|}}$ for $i = 1, 2, \dots, r$, where $(\mathbf{V}_{S,\cdot})^+$ is the generalized inverse of $\mathbf{V}_{S,\cdot}$.

Proof. Since $\mathbf{F} = (\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_T)^\top$, and $\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_T$ are independent sub-Gaussian random vectors under Assumption 3, we obtain by (4.22) in Vershynin (2017) that,

$$\left\| \frac{1}{|S|} (\mathbf{F}_{S,\cdot})^\top \mathbf{F}_{S,\cdot} - \mathbf{I}_r \right\|_2 = \left\| \frac{1}{|S|} \sum_{i \in S} \mathbf{f}_i (\mathbf{f}_i)^\top - \mathbf{I}_r \right\|_2 \lesssim \sqrt{\frac{r}{|S|}} + \sqrt{\frac{\log n}{|S|}} + \frac{r}{|S|} + \frac{\log n}{|S|} \lesssim \sqrt{\frac{r + \log n}{|S|}}, \quad (\text{D.2})$$

with probability at least $1 - O(n^{-2})$, where the last inequality is owing to $(r + \log n) \ll |S|$ in (D.1). In particular, we let \mathcal{E}_0 be the event that (D.2) happens, then we show that \mathcal{E}_S satisfies all the requirements.

When (D.2) happens, since $(r + \log n) \ll |S|$, we obtain that $\left\| \frac{1}{|S|} (\mathbf{F}_{S,\cdot})^\top \mathbf{F}_{S,\cdot} - \mathbf{I}_r \right\|_2 \ll 1$, and thus $\lambda_i(\frac{1}{|S|} (\mathbf{F}_{S,\cdot})^\top \mathbf{F}_{S,\cdot}) \asymp 1$ for $i = 1, 2, \dots, r$, implying that $\sigma_i(\mathbf{F}_{S,\cdot}) \asymp \sqrt{|S|}$ for $i = 1, 2, \dots, r$. Since $\sigma_r(\mathbf{F}_{S,\cdot}) \asymp \sqrt{|S|}$ implies that $\sigma_{\min}(\mathbf{F}_{S,\cdot}) > 0$, we obtain that $\text{rank}(\mathbf{F}_{S,\cdot}) = r$, i.e., $\mathbf{F}_{S,\cdot}$ has full column rank.

Then, recall that, given \mathcal{E}_0 in Lemma 2, it holds $\mathbf{V} = \frac{1}{\sqrt{T}} \mathbf{F} \mathbf{J}$ and \mathbf{J} is a $r \times r$ invertible matrix satisfying that $\|\mathbf{J} - \mathbf{Q}\|_2 \ll 1$ and $\sigma_i(\mathbf{J}) \asymp 1$ for $i = 1, 2, \dots, r$, where $\mathbf{Q} \in \mathcal{O}^{r \times r}$ is the rotation matrix given in Lemma 2. When $\mathcal{E}_0 \cap \mathcal{E}_S$ happens, since $\|\mathbf{J} - \mathbf{Q}\|_2 \ll 1$ and $\left\| \frac{1}{|S|} (\mathbf{F}_{S,\cdot})^\top \mathbf{F}_{S,\cdot} - \mathbf{I}_r \right\|_2 \ll 1$, we have that

$$\begin{aligned} \left\| \frac{T}{|S|} (\mathbf{V}_{S,\cdot})^\top \mathbf{V}_{S,\cdot} - \mathbf{I}_r \right\|_2 &= \left\| \mathbf{J}^\top \left(\frac{1}{|S|} (\mathbf{F}_{S,\cdot})^\top \mathbf{F}_{S,\cdot} \right) \mathbf{J} - \mathbf{Q}^\top \mathbf{I}_r \mathbf{Q} \right\|_2 \\ &\leq \left\| \mathbf{J}^\top \left(\frac{1}{|S|} (\mathbf{F}_{S,\cdot})^\top \mathbf{F}_{S,\cdot} - \mathbf{I}_r \right) \mathbf{J} \right\|_2 + \|\mathbf{J}^\top \mathbf{J} - \mathbf{Q}^\top \mathbf{Q}\|_2 \\ &\leq \left\| \frac{1}{|S|} (\mathbf{F}_{S,\cdot})^\top \mathbf{F}_{S,\cdot} - \mathbf{I}_r \right\|_2 + \|(\mathbf{J}^\top - \mathbf{Q}^\top) \mathbf{Q}\|_2 + \|\mathbf{J}^\top (\mathbf{J} - \mathbf{Q})\|_2 \\ &\ll 1. \end{aligned}$$

So, we obtain that $\lambda_i(\frac{T}{|S|} (\mathbf{V}_{S,\cdot})^\top \mathbf{V}_{S,\cdot}) \asymp 1$ for $i = 1, 2, \dots, r$, and thus $\sigma_i(\mathbf{V}_{S,\cdot}) \asymp \sqrt{\frac{|S|}{T}}$ for $i = 1, 2, \dots, r$, implying that $\text{rank}(\mathbf{V}_{S,\cdot}) = r$.

Given $\mathcal{E}_0 \cap \mathcal{E}_S$, since $\mathbf{V}_{S,\cdot}$ has full column rank, we have that $(\mathbf{V}_{S,\cdot})^+ = [(\mathbf{V}_{S,\cdot})^\top \mathbf{V}_{S,\cdot}]^{-1} (\mathbf{V}_{S,\cdot})^\top$. So, on the one hand, we have that

$$\sigma_{\max}((\mathbf{V}_{S,\cdot})^+) \leq \|[(\mathbf{V}_{S,\cdot})^\top \mathbf{V}_{S,\cdot}]^{-1}\|_2 \|\mathbf{V}_{S,\cdot}\|_2 \leq \frac{\sigma_{\max}(\mathbf{V}_{S,\cdot})}{(\sigma_{\min}(\mathbf{V}_{S,\cdot}))^2} \lesssim \left(\sqrt{\frac{|S|}{T}} \right)^{-2} \sqrt{\frac{|S|}{T}} = \sqrt{\frac{T}{|S|}}.$$

On the other hand, similarly we have that

$$\sigma_{\min}((\mathbf{V}_{S,\cdot})^+) \geq \frac{\sigma_{\min}(\mathbf{V}_{S,\cdot})}{(\sigma_{\max}(\mathbf{V}_{S,\cdot}))^2} \gtrsim \left(\sqrt{\frac{|S|}{T}} \right)^{-2} \sqrt{\frac{|S|}{T}} = \sqrt{\frac{T}{|S|}}.$$

So we conclude that $\sigma_i((\mathbf{V}_{S,\cdot})^+) \asymp \sqrt{\frac{T}{|S|}}$ for $i = 1, 2, \dots, r$. \square

Using the properties of $\mathbf{V}_{S,\cdot}$ obtained in Lemma 8, we are able to establish the following properties on the projection matrices.

Lemma 9. *Suppose that Assumptions 1, 2, and 3 hold. Assume that $r + \log n \ll T$, $r \log n \ll n$, and $\rho\sqrt{\log n} \ll 1$ as in Theorem 2, and $r + \log n \ll |S|$ in (D.1). Then we have that*

(i) *The projection matrices have the perturbation bound as below: with probability at least $1 - O(n^{-2})$, it holds*

$$\|\mathbf{P}_{\mathbf{V}_{S,\cdot}} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}\|_2 \lesssim \rho\sqrt{\frac{T}{n}}\sqrt{r}\log^{3/2}n.$$

(ii) *Under the null hypothesis H_0 in (5.1), we have that, with probability at least $1 - O(n^{-2})$,*

$$\|(\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}})\mathbf{v}\|_2 \lesssim \sqrt{|S|}\rho\sqrt{\frac{r}{n}}\log n\|(\mathbf{V}_{S,\cdot})^+\mathbf{v}\|_2,$$

and the following first-order approximation

$$(\mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}} - \mathbf{I}_{|S|})\mathbf{v} = \mathbf{G}_P + \Upsilon_P, \quad \text{with} \quad \mathbf{G}_P := (\mathbf{I}_{|S|} - \mathbf{P}_{\mathbf{V}_{S,\cdot}})(\mathbf{G}_V)_{S,\cdot}(\mathbf{V}_{S,\cdot})^+\mathbf{v},$$

where \mathbf{G}_V is the first-order term in (3.3b) and is given by $\mathbf{G}_V = T^{-1/2}\mathbf{E}^\top\mathbf{U}\mathbf{\Lambda}^{-1}$, and with probability at least $1 - O(n^{-2})$, it holds

$$\|\Upsilon_P\|_2 \lesssim \sqrt{|S|}\left(\rho^2\frac{1}{\sqrt{T}} + \rho\frac{1}{\sqrt{nT}}\right)r\log^2n\|(\mathbf{V}_{S,\cdot})^+\mathbf{v}\|_2.$$

Proof. The proof consists of three steps.

Step 1 – Projection matrix perturbation bound.

Since the rotation matrix $\mathbf{R}_V \in \mathcal{O}^{r \times r}$ is invertible, we have that $\text{col}(\widehat{\mathbf{V}}_{S,\cdot}) = \text{col}(\widehat{\mathbf{V}}_{S,\cdot}\mathbf{R}_V)$, implying that $\mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}} = \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}\mathbf{R}_V}$. By Theorem 1.2 in Chen et al. (2016) on the perturbation bound of projection matrix, we have that

$$\begin{aligned} \|\mathbf{P}_{\mathbf{V}_{S,\cdot}} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}\|_2 &= \|\mathbf{P}_{\mathbf{V}_{S,\cdot}} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}\mathbf{R}_V}\|_2 \leq \min\left\{\|(\mathbf{V}_{S,\cdot})^+\|_2, \|(\widehat{\mathbf{V}}_{S,\cdot}\mathbf{R}_V)^+\|_2\right\}\|\widehat{\mathbf{V}}_{S,\cdot}\mathbf{R}_V - \mathbf{V}_{S,\cdot}\|_2 \\ &\leq \|(\mathbf{V}_{S,\cdot})^+\|_2\|\widehat{\mathbf{V}}_{S,\cdot}\mathbf{R}_V - \mathbf{V}_{S,\cdot}\|_2. \end{aligned}$$

The term $\|(\mathbf{V}_{S,\cdot})^+\|_2$ can be bounded using Lemma 8. For the term $\|\widehat{\mathbf{V}}_{S,\cdot}\mathbf{R}_V - \mathbf{V}_{S,\cdot}\|_2$, we have that

$$\|\widehat{\mathbf{V}}_{S,\cdot}\mathbf{R}_V - \mathbf{V}_{S,\cdot}\|_2 \leq \sqrt{|S|}\|\widehat{\mathbf{V}}_{S,\cdot}\mathbf{R}_V - \mathbf{V}_{S,\cdot}\|_{2,\infty} \leq \sqrt{|S|}\|\widehat{\mathbf{V}}\mathbf{R}_V - \mathbf{V}\|_{2,\infty} \lesssim \sqrt{|S|}\rho\sqrt{\frac{r}{n}}\log n,$$

where the last inequality uses the upper bound of $\|\widehat{\mathbf{V}}\mathbf{R}_V - \mathbf{V}\|_{2,\infty}$ obtained in the proof of Corollary 1. So, we obtain

$$\|\mathbf{P}_{\mathbf{V}_{S,\cdot}} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}\|_2 \leq \|(\mathbf{V}_{S,\cdot})^+\|_2\|\widehat{\mathbf{V}}_{S,\cdot}\mathbf{R}_V - \mathbf{V}_{S,\cdot}\|_2 \lesssim \sqrt{\frac{T}{|S|}}\sqrt{|S|}\rho\sqrt{\frac{r}{n}}\log n = \rho\sqrt{\frac{T}{n}}\sqrt{r}\log n.$$

Step 2 – First-order approximation expression.

For ease of exposition, we denote

$$\mathbf{G} := \widehat{\mathbf{V}}_{S,\cdot}\mathbf{R}_V - \mathbf{V}_{S,\cdot}.$$

We obtain by the identity after (4.6) in Stewart (1977) that

$$\begin{aligned} \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}} - \mathbf{P}_{\mathbf{V}_{S,\cdot}} &= \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}\mathbf{R}_V} - \mathbf{P}_{\mathbf{V}_{S,\cdot}} \\ &= (\widehat{\mathbf{V}}_{S,\cdot}\mathbf{R}_V)^+\mathbf{P}_{(\widehat{\mathbf{V}}_{S,\cdot}\mathbf{R}_V)^\top}G^\top(\mathbf{I}_{|S|} - \mathbf{P}_{\mathbf{V}_{S,\cdot}}) + (\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}})G\mathbf{P}_{(\mathbf{V}_{S,\cdot})^\top}(\mathbf{V}_{S,\cdot})^+. \end{aligned}$$

Next, we obtain by Lemma 2 that, there exists a $\sigma(\mathbf{F})$ -measurable and invertible matrix \mathbf{J} such that $\mathbf{V} = \frac{1}{\sqrt{T}}\mathbf{F}\mathbf{J}$, implying that $\mathbf{V}_{S,\cdot} = \frac{1}{\sqrt{T}}\mathbf{F}_{S,\cdot}\mathbf{J}$. So, under the null hypothesis H_0 in (5.1), since $\mathbf{v} \in \text{col}(\mathbf{F}_{S,\cdot})$ implies that $\mathbf{v} \in \text{col}(\mathbf{V}_{S,\cdot})$, we have that $(\mathbf{I}_{|S|} - \mathbf{P}_{\mathbf{V}_{S,\cdot}})\mathbf{v} = 0$. Then, we obtain that

$$(\mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}} - \mathbf{I}_{|S|})\mathbf{v} = (\mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}\mathbf{R}_V} - \mathbf{P}_{\mathbf{V}_{S,\cdot}})\mathbf{v} = (\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}})G\mathbf{P}_{(\mathbf{V}_{S,\cdot})^\top}(\mathbf{V}_{S,\cdot})^+\mathbf{v}$$

$$= (\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}})G(\mathbf{V}_{S,\cdot})^+ \mathbf{v}.$$

Here, the last equality is because $\mathbf{V}_{S,\cdot}$ has full column rank, implying that $\mathbf{P}_{(\mathbf{V}_{S,\cdot})^\top} = \mathbf{I}_r$.

Then, since the projection matrix $\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}$ satisfies that $\|\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}\|_2 = 1$, we obtain that

$$\|(\mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}} - \mathbf{I}_{|S|})\mathbf{v}\|_2 \leq \|G\|_2 \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2 \lesssim \sqrt{|S|} \rho \sqrt{\frac{r}{n}} \log n \cdot \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2,$$

where the last inequality uses the upper bound of $\|G\|_2 = \|\widehat{\mathbf{V}}_{S,\cdot} \mathbf{R}_V - \mathbf{V}_{S,\cdot}\|_2$ obtained in Step 1.

Step 3 – Decomposition and upper bounds for the higher-order terms.

To establish the first-order approximation, we do the decomposition as follows:

$$\begin{aligned} & (\mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}} - \mathbf{I}_{|S|})\mathbf{v} \\ &= (\mathbf{I}_{|S|} - \mathbf{P}_{\mathbf{V}_{S,\cdot}})G(\mathbf{V}_{S,\cdot})^+ \mathbf{v} + (\mathbf{P}_{\mathbf{V}_{S,\cdot}} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}})G(\mathbf{V}_{S,\cdot})^+ \mathbf{v} \\ &= (\mathbf{I}_{|S|} - \mathbf{P}_{\mathbf{V}_{S,\cdot}})(\mathbf{G}_V)_{S,\cdot}(\mathbf{V}_{S,\cdot})^+ \mathbf{v} \\ & \quad + \underbrace{(\mathbf{I}_{|S|} - \mathbf{P}_{\mathbf{V}_{S,\cdot}})(\boldsymbol{\Psi}_V)_{S,\cdot}(\mathbf{V}_{S,\cdot})^+ \mathbf{v}}_{=:\psi_1} + \underbrace{(\mathbf{P}_{\mathbf{V}_{S,\cdot}} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}})G(\mathbf{V}_{S,\cdot})^+ \mathbf{v}}_{=:\psi_2}. \end{aligned}$$

Here, the last equation is because $G = \widehat{\mathbf{V}}_{S,\cdot} \mathbf{R}_V - \mathbf{V}_{S,\cdot} = (\widehat{\mathbf{V}} \mathbf{R}_V - \mathbf{V})_{S,\cdot} = (\mathbf{G}_V)_{S,\cdot} + (\boldsymbol{\Psi}_V)_{S,\cdot}$. So, we get that $\boldsymbol{\Upsilon}_P = \psi_1 + \psi_2$, and thus the problems boil down to deriving the upper bound of the norms of ψ_1 and ψ_2 .

For ψ_1 , since $\|(\mathbf{I}_{|S|} - \mathbf{P}_{\mathbf{V}_{S,\cdot}})\|_2 = 1$, we have that

$$\begin{aligned} \|\psi_1\|_2 &\leq \|(\boldsymbol{\Psi}_V)_{S,\cdot}\|_2 \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2 \leq \sqrt{|S|} \|\boldsymbol{\Psi}_V\|_{2,\infty} \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2 \\ &\stackrel{(i)}{\lesssim} \sqrt{|S|} \left(\rho^2 \sqrt{\frac{r}{n}} \log^{3/2} n + \left(\rho^2 + \rho \sqrt{\frac{r}{n}} \log n \right) \|\mathbf{V}\|_{2,\infty} \right) \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2 \\ &\stackrel{(ii)}{\lesssim} \sqrt{|S|} \left(\rho^2 \sqrt{\frac{r}{n}} \log^{3/2} n + \left(\rho^2 + \rho \sqrt{\frac{r}{n}} \right) \log n \sqrt{\frac{\log n}{T}} \right) \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2. \end{aligned}$$

where (i) uses the upper bound for $\|(\boldsymbol{\Psi}_V)_{S,\cdot}\|_2$ in Theorem 1, and (ii) uses the upper bound $\|\mathbf{V}\|_{2,\infty} \lesssim \sqrt{\frac{\log n}{T}}$ obtained in Lemma 2.

For ψ_2 , using the bounds of $\|\mathbf{P}_{\mathbf{V}_{S,\cdot}} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}\|_2$ and $\|G\|_2 = \|\widehat{\mathbf{V}}_{S,\cdot} \mathbf{R}_V - \mathbf{V}_{S,\cdot}\|_2$ obtained in Step 1, we have that

$$\begin{aligned} \|\psi_2\|_2 &\leq \|\mathbf{P}_{\mathbf{V}_{S,\cdot}} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}\|_2 \|G\|_2 \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2 \\ &\lesssim \rho \sqrt{\frac{T}{n}} \sqrt{r} \log n \cdot \sqrt{|S|} \rho \sqrt{\frac{r}{n}} \log n \cdot \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2 = \sqrt{|S|} \rho^2 \frac{\sqrt{T}}{n} r \log^2 n \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2. \end{aligned}$$

Then, we combine the above results for $\|\psi_1\|_2$ and $\|\psi_2\|_2$ to obtain that

$$\begin{aligned} \|\boldsymbol{\Upsilon}_P\|_2 &\leq \|\psi_1\|_2 + \|\psi_2\|_2 \\ &\lesssim \sqrt{|S|} \left(\rho^2 \sqrt{\frac{r}{n}} \log^{3/2} n + \left(\rho^2 + \rho \sqrt{\frac{r}{n}} \right) \log n \sqrt{\frac{\log n}{T}} + \frac{\sqrt{T}}{n} \rho^2 r \log^2 n \right) \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2, \\ &\lesssim \sqrt{|S|} \left(\rho^2 \frac{1}{\sqrt{T}} + \rho \frac{1}{\sqrt{nT}} \right) r \log^2 n \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2 \\ &\lesssim \sqrt{|S|} \left(\rho^2 \frac{1}{\sqrt{T}} + \rho \frac{1}{\sqrt{nT}} \right) r \log^2 n \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2, \end{aligned}$$

which is our desired result. \square

Then, we establish Gaussian approximation for the first-order term. For simplicity of notations, we denote $\chi^2(n)$ as a Chi-square random variable with degree of freedom equal to n , and denote by $\chi_\phi^2(n)$ its ϕ -quantile.

Lemma 10. *Suppose that the assumptions in Lemma 9 hold. Then, under the null hypothesis H_0 in (5.1), we have that, for any random variable ζ satisfying $|\zeta| \ll |S|$, it holds*

$$|\mathbb{P}(\mathcal{T}(S, \mathbf{v}) + \zeta \leq \chi_{1-\alpha}^2(|S| - r)) - (1 - \alpha)| \lesssim \sqrt{|S|} \left(\rho \sqrt{\frac{n}{T}} + \frac{1}{\sqrt{T}} \right) \kappa_\varepsilon r \log^2 n + \frac{1}{\sqrt{|S|}} |\zeta| + s_3, \quad (\text{D.3})$$

where

$$\mathcal{T}(S, \mathbf{v}) := \frac{1}{\phi} \mathbf{v}^\top (\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}) (\mathbf{I}_{|S|} - \mathbf{P}_{\mathbf{V}_{S,\cdot}}) (\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}) \mathbf{v},$$

with

$$\phi := \frac{1}{T} ((\mathbf{V}_{S,\cdot})^+ \mathbf{v})^\top \boldsymbol{\Lambda}^{-1} \mathbf{U}^\top \boldsymbol{\Sigma}_\varepsilon \mathbf{U} \boldsymbol{\Lambda}^{-1} (\mathbf{V}_{S,\cdot})^+ \mathbf{v},$$

and $s_3 \geq 0$ is a constant bounded by

$$s_3 \lesssim \kappa_\varepsilon \sqrt{\frac{n}{T}} |S|^{3/2} \frac{\|\boldsymbol{\Sigma}_\varepsilon^{1/2} \bar{\mathbf{U}}\|_{2,\infty}}{\|\boldsymbol{\Sigma}_\varepsilon^{1/2}\|_2}.$$

Further, if all the entries of the matrix \mathbf{Z} are Gaussian, i.e., the noise is Gaussian, then s_3 in the inequality (D.3) is equal to zero, i.e., the inequality (D.3) holds when $s_3 = 0$.

In addition, we have that

$$\phi \gtrsim \frac{1}{\kappa_\varepsilon^2 n} \rho^2 \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2^2.$$

Proof. Our starting point is the following inequality

$$\mathbb{P}(\mathcal{T}(S, \mathbf{v}) \leq \chi_{1-\alpha}^2(|S| - r) - |\zeta|) \leq \mathbb{P}(\mathcal{T}(S, \mathbf{v}) + \zeta \leq \chi_{1-\alpha}^2(|S| - r)) \leq \mathbb{P}(\mathcal{T}(S, \mathbf{v}) \leq \chi_{1-\alpha}^2(|S| - r) + |\zeta|). \quad (\text{D.4})$$

Then, we will prove both the upper bound and the lower bound are close to $(1 - \alpha)$.

We will use three steps. We conduct the proof conditioning on $\mathcal{E}_0 \cap \mathcal{E}_S$.

Step 1 - Bounding $\mathbb{P}(\mathcal{T}(S, \mathbf{v}) + \zeta \leq \chi_{1-\alpha}^2(|S| - r))$.

We already obtained the first-order approximation of $(\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}) \mathbf{v} = \mathbf{G}_P + \boldsymbol{\Upsilon}_P$ in Lemma 9. Indeed, the first-order term \mathbf{G}_P can be rewritten as

$$\mathbf{G}_P = (\mathbf{I}_{|S|} - \mathbf{P}_{\mathbf{V}_{S,\cdot}}) \mathbf{u} \quad \text{with} \quad \mathbf{u} := (\mathbf{G}_V)_{S,\cdot} (\mathbf{V}_{S,\cdot})^+ \mathbf{v} = (T^{-1/2} \mathbf{E}^\top \mathbf{U} \boldsymbol{\Lambda}^{-1})_{S,\cdot} (\mathbf{V}_{S,\cdot})^+ \mathbf{v}. \quad (\text{D.5})$$

Later we will show that the $|S|$ -dimensional random vector \mathbf{u} is close to a Gaussian vector, so that the desired Chi-square law results can be proven. But now let us focus on how this term appear at the term $\mathcal{T}(S, \mathbf{v})$. To do this, we need to decompose $(\mathbf{I}_{|S|} - \mathbf{P}_{\mathbf{V}_{S,\cdot}})$ as follows.

By Lemma 8, given the event $\mathcal{E}_0 \cap \mathcal{E}_S$, $\mathbf{V}_{S,\cdot}$ has rank r , and the $|S| \times |S|$ projection matrix $(\mathbf{I}_{|S|} - \mathbf{P}_{\mathbf{V}_{S,\cdot}})$ is idempotent with rank $(|S| - r)$. So the eigen-decomposition of $\mathbf{I}_{|S|} - \mathbf{P}_{\mathbf{V}_{S,\cdot}}$ is given by

$$\mathbf{I}_{|S|} - \mathbf{P}_{\mathbf{V}_{S,\cdot}} = \mathbf{H} \mathbf{I}_{|S|-r} \mathbf{H}^\top = \mathbf{H} \mathbf{H}^\top, \quad (\text{D.6})$$

where \mathbf{H} is a $\sigma(\mathbf{F})$ -measurable $|S| \times (|S| - r)$ matrix, and \mathbf{H} has orthonormal columns, i.e., $\mathbf{H}^\top \mathbf{H} = \mathbf{I}_{|S|-r}$. So we obtain that

$$\mathbf{G}_P = (\mathbf{I}_{|S|} - \mathbf{P}_{\mathbf{V}_{S,\cdot}}) \mathbf{u} = \mathbf{H} \mathbf{H}^\top \mathbf{u},$$

and thus $\mathcal{T}(S, \mathbf{v})$ can be rewritten as

$$\begin{aligned} \mathcal{T}(S, \mathbf{v}) &= \frac{1}{\|\mathbf{m}\|_2^2} (\mathbf{G}_P + \boldsymbol{\Upsilon}_P)^\top \mathbf{H} \mathbf{H}^\top (\mathbf{G}_P + \boldsymbol{\Upsilon}_P) = \frac{1}{\|\mathbf{m}\|_2^2} (\mathbf{H} \mathbf{H}^\top \mathbf{u} + \boldsymbol{\Upsilon}_P)^\top \mathbf{H} \mathbf{H}^\top (\mathbf{H} \mathbf{H}^\top \mathbf{u} + \boldsymbol{\Upsilon}_P) \\ &= \frac{1}{\|\mathbf{m}\|_2^2} \|\mathbf{H}^\top (\mathbf{H} \mathbf{H}^\top \mathbf{u} + \boldsymbol{\Upsilon}_P)\|_2^2 = \frac{1}{\|\mathbf{m}\|_2^2} \|\mathbf{H}^\top \mathbf{u} + \mathbf{H}^\top \boldsymbol{\Upsilon}_P\|_2^2 = \left\| \frac{1}{\|\mathbf{m}\|_2} \mathbf{H}^\top \mathbf{u} + \frac{1}{\|\mathbf{m}\|_2} \mathbf{H}^\top \boldsymbol{\Upsilon}_P \right\|_2^2. \end{aligned} \quad (\text{D.7})$$

The expression (D.7) shows how $\frac{1}{\|\mathbf{m}\|_2} \mathbf{H}^\top \mathbf{u}$ plays a role in $\mathcal{T}(S, \mathbf{v})$.

To further use the probabilities about $\frac{1}{\|\mathbf{m}\|_2} \|\mathbf{H}^\top \mathbf{u}\|_2$ to bound the probabilities about $\mathcal{T}(S, \mathbf{v})$, we define

$$\tau := \sqrt{\mathcal{T}(S, \mathbf{v})} - \frac{1}{\|\mathbf{m}\|_2} \|\mathbf{H}^\top \mathbf{u}\|_2. \quad (\text{D.8})$$

Then, we obtain that, for the upper bound in (D.4)

$$\begin{aligned} \mathbb{P}(\mathcal{T}(S, \mathbf{v}) \leq \chi_{1-\alpha}^2(|S| - r) + |\zeta|) &= \mathbb{P}\left(\frac{1}{\|\mathbf{m}\|_2} \|\mathbf{H}^\top \mathbf{u}\|_2 + \tau \leq \sqrt{\chi_{1-\alpha}^2(|S| - r) + |\zeta|}\right) \\ &\leq \mathbb{P}\left(\frac{1}{\|\mathbf{m}\|_2} \|\mathbf{H}^\top \mathbf{u}\|_2 \leq \sqrt{\chi_{1-\alpha}^2(|S| - r) + |\zeta|} + |\tau|\right), \end{aligned}$$

and similarly for the lower bound in (D.4) we have that $\mathbb{P}(\mathcal{T}(S, \mathbf{v}) \leq \chi_{1-\alpha}^2(|S| - r) - |\zeta|) \geq \mathbb{P}\left(\frac{1}{\|\mathbf{m}\|_2} \|\mathbf{H}^\top \mathbf{u}\|_2 \leq \sqrt{\chi_{1-\alpha}^2(|S| - r) - |\zeta|} - |\tau|\right)$. Thus, we obtain by the inequality (D.4) that

$$\begin{aligned} &\mathbb{P}\left(\frac{1}{\|\mathbf{m}\|_2} \|\mathbf{H}^\top \mathbf{u}\|_2 \leq \sqrt{\chi_{1-\alpha}^2(|S| - r) - |\zeta|} - |\tau|\right) \\ &\leq \mathbb{P}(\mathcal{T}(S, \mathbf{v}) + \zeta \leq \chi_{1-\alpha}^2(|S| - r)) \\ &\leq \mathbb{P}\left(\frac{1}{\|\mathbf{m}\|_2} \|\mathbf{H}^\top \mathbf{u}\|_2 \leq \sqrt{\chi_{1-\alpha}^2(|S| - r) + |\zeta|} + |\tau|\right). \end{aligned} \quad (\text{D.9})$$

Step 2 - Expressing $\frac{1}{\|\mathbf{m}\|_2} \mathbf{H}^\top \mathbf{u}$ as a sum to show its proximity to a Gaussian vector.

We define

$$s_3 := \sup_{R \geq 0} \left| \mathbb{P}\left(\frac{1}{\|\mathbf{m}\|_2} \|\mathbf{H}^\top \mathbf{u}\|_2 \leq R\right) - \mathbb{P}(\chi^2(|S| - r) \leq R^2) \right|. \quad (\text{D.10})$$

We look at the random vector $\mathbf{u} \in \mathbb{R}^{|S|}$ first. Conditioning on \mathbf{F} , the random vector \mathbf{u} defined in (D.5) can be written as a sum of independent and mean zero $|S|$ -dimensional vectors as follows:

$$\begin{aligned} \mathbf{u} &= (T^{-1/2} \mathbf{E}^\top \mathbf{U} \mathbf{\Lambda}^{-1})_{S, \cdot} (\mathbf{V}_{S, \cdot})^+ \mathbf{v} = (T^{-1/2} \mathbf{Z}^\top \mathbf{\Sigma}_\varepsilon^{1/2} \mathbf{U} \mathbf{\Lambda}^{-1})_{S, \cdot} (\mathbf{V}_{S, \cdot})^+ \mathbf{v} \\ &= T^{-1/2} (\mathbf{Z}^\top)_{S, \cdot} \mathbf{\Sigma}_\varepsilon^{1/2} \mathbf{U} \mathbf{\Lambda}^{-1} (\mathbf{V}_{S, \cdot})^+ \mathbf{v} \\ &= (\mathbf{Z}_{\cdot, S})^\top \mathbf{m} = \sum_{k=1}^N m_k (\mathbf{Z}_{k, S})^\top \quad \text{with} \quad \mathbf{m} := (m_1, m_2, \dots, m_N)^\top = T^{-1/2} \mathbf{\Sigma}_\varepsilon^{1/2} \mathbf{U} \mathbf{\Lambda}^{-1} (\mathbf{V}_{S, \cdot})^+ \mathbf{v}. \end{aligned} \quad (\text{D.11})$$

So we obtain that $\frac{1}{\|\mathbf{m}\|_2} \mathbf{H}^\top \mathbf{u}$ can be expressed as a summation as follows,

$$\frac{1}{\|\mathbf{m}\|_2} \mathbf{H}^\top \mathbf{u} = \sum_{k=1}^N \frac{1}{\|\mathbf{m}\|_2} m_k \mathbf{H}^\top (\mathbf{Z}_{k, S})^\top. \quad (\text{D.12})$$

Here, the matrix \mathbf{H} defined in (D.6) is $\sigma(\mathbf{F})$ -measurable; the vector \mathbf{m} defined in (D.11) is also $\sigma(\mathbf{F})$ -measurable under the Null because \mathbf{v} is in the column space of $\mathbf{F}_{S, \cdot}$. By the independence between \mathbf{F} and \mathbf{Z} in Assumption 2, we obtain that, the vectors $\frac{1}{\|\mathbf{m}\|_2} m_k \mathbf{H}^\top (\mathbf{Z}_{k, S})^\top \in \mathbb{R}^{|S|-r}$ for $k = 1, 2, \dots, N$ are independent and mean zero, and the covariance matrix is given by

$$\text{cov}\left(\frac{1}{\|\mathbf{m}\|_2} \mathbf{H}^\top \mathbf{u} | \mathbf{F}\right) = \frac{1}{\|\mathbf{m}\|_2^2} \sum_{k=1}^N m_k^2 \mathbf{H}^\top \mathbf{H} = \frac{1}{\|\mathbf{m}\|_2^2} \|\mathbf{m}\|_2^2 \mathbf{I}_{|S|-r} = \mathbf{I}_{|S|-r}.$$

Here, we use the fact that $\text{cov}((\mathbf{Z}_{k, S})^\top | \mathbf{F}) = \mathbf{I}_{|S|-r}$, and $\mathbf{H}^\top \mathbf{H} = \mathbf{I}_{|S|-r}$.

Given the above analysis of $\frac{1}{\|\mathbf{m}\|_2} \mathbf{H}^\top \mathbf{u}$, we are now ready to bound the term s_3 defined in (D.10):

(i) When all entries of \mathbf{Z} are Gaussian, we have that, conditioning on \mathbf{F} , the vector $\frac{1}{\|\mathbf{m}\|_2} \mathbf{H}^\top \mathbf{u}$ is also a Gaussian vector because it is a linear combination of Gaussian vectors. Since $\text{cov}\left(\frac{1}{\|\mathbf{m}\|_2} \mathbf{H}^\top \mathbf{u} | \mathbf{F}\right) =$

$\mathbf{I}_{|S|-r}$, and $\mathbf{I}_{|S|-r}$ is independent with \mathbf{F} , we obtain that the unconditional law of $\frac{1}{\|\mathbf{m}\|_2} \mathbf{H}^\top \mathbf{u}$ the $(|S| - r)$ -dimensional standard Gaussian law $\mathcal{N}(0, \mathbf{I}_{|S|-r})$, implying that the conditional law of $\frac{1}{\|\mathbf{m}\|_2} \mathbf{u}$ given \mathbf{F} is the $|S|$ -dimensional standard Gaussian law $\mathcal{N}(0, \mathbf{I}_{|S|})$. Note that $\chi^2(|S| - r)$ has the same distribution with the 2-norm of $\mathcal{N}(0, \mathbf{I}_{|S|-r})$. Thus, we have that $s_3 = 0$ when all entries of \mathbf{Z} are Gaussian.

(ii) When the entries of \mathbf{Z} are sub-Gaussian, we obtain by Lemma 16 that

$$s_3 \lesssim \sum_{k=1}^N \mathbb{E} \left[\left\| \frac{1}{\|\mathbf{m}\|_2} \mathbf{H}^\top m_k (\mathbf{Z}_{k,S})^\top \right\|_2^3 | \mathbf{F} \right] = \sum_{k=1}^N \frac{1}{\|\mathbf{m}\|_2^3} \mathbb{E} \left[\left\| \mathbf{H}^\top m_k (\mathbf{Z}_{k,S})^\top \right\|_2^3 | \mathbf{F} \right].$$

Then, since $\|\mathbf{H}\|_2 = 1$ owing to $\mathbf{H}^\top \mathbf{H} = \mathbf{I}_{|S|-r}$, we have that

$$\begin{aligned} \mathbb{E} \left[\left\| \mathbf{H}^\top m_k (\mathbf{Z}_{k,S})^\top \right\|_2^3 | \mathbf{F} \right] &\leq \mathbb{E} \left[\left\| m_k (\mathbf{Z}_{k,S})^\top \right\|_2^3 | \mathbf{F} \right] = \mathbb{E} \left[|m_k|^3 \|\mathbf{Z}_{k,S}\|_2^3 | \mathbf{F} \right] \\ &\stackrel{(i)}{=} |m_k|^3 \mathbb{E} \left[\|\mathbf{Z}_{k,S}\|_2^3 | \mathbf{F} \right] \stackrel{(ii)}{\leq} |m_k|^2 \mathbb{E} \left[\|\mathbf{Z}_{k,S}\|_2^3 \right] \max_{1 \leq i \leq N} |m_i|, \end{aligned}$$

where (i) is because m_k is $\sigma(\mathbf{F})$ -measurable, (ii) is because \mathbf{Z} is independent with \mathbf{F} . On the one hand, we have that

$$\mathbb{E} \left[\|\mathbf{Z}_{k,S}\|_2^3 \right] = \mathbb{E} \left[\left(\sum_{i=1}^{|S|} |Z_{k,i}|^2 \right)^{3/2} \right] \stackrel{(i)}{\leq} \mathbb{E} \left[\left(\sum_{i=1}^{|S|} |Z_{k,i}|^3 \right) \left(\sum_{i=1}^{|S|} 1 \right)^{1/2} \right] = |S|^{3/2} \mathbb{E} \left[|Z_{k,i}|^3 \right] \stackrel{(ii)}{\lesssim} |S|^{3/2},$$

where (i) uses Holder's inequality, and (ii) uses the fact that the entries of \mathbf{Z} are sub-Gaussian and their sub-Gaussian norms satisfy $\|Z_{k,i}\|_{\psi_2} = O(1)$ under Assumption 2. Then, On the other hand, for $\max_{1 \leq i \leq N} |m_i|$, by definition of the vector \mathbf{m} in (D.11), we have that

$$\begin{aligned} \max_{1 \leq i \leq N} |m_i| &= \|T^{-1/2} \Sigma_\varepsilon^{1/2} \mathbf{U} \Lambda^{-1} (\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_{2,\infty} \leq \frac{1}{\sqrt{T}} \|\Sigma_\varepsilon^{1/2} \mathbf{U}\|_{2,\infty} \|\Lambda^{-1} (\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2 \\ &\leq \frac{1}{\sqrt{T}} \|\Sigma_\varepsilon^{1/2} \mathbf{U}\|_{2,\infty} \|\Lambda^{-1}\|_2 \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2 \stackrel{(i)}{\leq} \frac{1}{\sqrt{T}} \|\Sigma_\varepsilon^{1/2} \bar{\mathbf{U}}\|_{2,\infty} \|\mathbf{Q}\|_2 \frac{1}{\lambda_r} \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2 \\ &\stackrel{(ii)}{\lesssim} \frac{1}{\sqrt{T}} \|\Sigma_\varepsilon^{1/2} \bar{\mathbf{U}}\|_{2,\infty} \frac{1}{\sigma_r} \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2, \end{aligned}$$

where (i) uses $\mathbf{U} = \bar{\mathbf{U}} \mathbf{Q}$ in Lemma 2, (ii) uses the facts that $\lambda_i \asymp \sigma_i$, and $\mathbf{Q} \in \mathcal{O}^{r \times r}$ is a rotation matrix so that $\|\mathbf{Q}\|_2 = 1$. Combining the above upper bounds of $\mathbb{E}[\|\mathbf{Z}_{k,S}\|_2^3]$ and $\max_{1 \leq i \leq N} |m_i|$, we obtain that

$$s_3 \leq \mathbb{E} \left[\|\mathbf{Z}_{k,S}\|_2^3 \right] \max_{1 \leq i \leq N} |m_i| \sum_{k=1}^N \frac{1}{\|\mathbf{m}\|_2^3} |m_k|^2 \lesssim |S|^{3/2} \frac{1}{\sqrt{T}} \|\Sigma_\varepsilon^{1/2} \mathbf{U}\|_{2,\infty} \frac{1}{\sigma_r} \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2 \frac{1}{\|\mathbf{m}\|_2}.$$

The lower bound of $\|\mathbf{m}\|_2$ is derived as follows. Recall that $\mathbf{m} = T^{-1/2} \Sigma_\varepsilon^{1/2} \mathbf{U} \Lambda^{-1} (\mathbf{V}_{S,\cdot})^+ \mathbf{v}$ as defined in (D.11). So, we obtain that

$$\begin{aligned} \|\mathbf{m}\|_2 &= \sqrt{\mathbf{m}^\top \mathbf{m}} = \frac{1}{\sqrt{T}} \left[((\mathbf{V}_{S,\cdot})^+ \mathbf{v})^\top \Lambda^{-1} \mathbf{U} \Sigma_\varepsilon \mathbf{U} \Lambda^{-1} (\mathbf{V}_{S,\cdot})^+ \mathbf{v} \right]^{1/2} \\ &\geq \frac{1}{\sqrt{T}} \sqrt{\lambda_{\min}(\Lambda^{-1} \mathbf{U}^\top \Sigma_\varepsilon \mathbf{U} \Lambda^{-1})} \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2. \end{aligned}$$

Next, since $\mathbf{U}^\top \mathbf{U} = \mathbf{I}_r$ and $\Lambda = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_r)$ satisfies $\lambda_i \asymp \sigma_i$ as shown in Lemma 2, we obtain that

$$\begin{aligned} \lambda_{\min}(\Lambda^{-1} \mathbf{U}^\top \Sigma_\varepsilon \mathbf{U} \Lambda^{-1}) &= \inf_{\|\mathbf{x}\|_2=1} \mathbf{x}^\top \Lambda^{-1} \mathbf{U}^\top \Sigma_\varepsilon \mathbf{U} \Lambda^{-1} \mathbf{x} \geq \frac{1}{\lambda_r^2} \inf_{\|\mathbf{x}\|_2=1} \mathbf{x}^\top \mathbf{U}^\top \Sigma_\varepsilon \mathbf{U} \mathbf{x} \\ &\gtrsim \frac{1}{\sigma_r^2} \inf_{\|\mathbf{x}\|_2=1} (\mathbf{U} \mathbf{x})^\top \Sigma_\varepsilon \mathbf{U} \mathbf{x} \geq \frac{1}{\sigma_r^2} \lambda_{\min}(\Sigma_\varepsilon). \end{aligned}$$

Recall that $\rho := \sqrt{\|\boldsymbol{\Sigma}_\varepsilon\|_2} \sqrt{n}/(\sigma_r \sqrt{T})$. So, we conclude that

$$\|\mathbf{m}\|_2 \gtrsim \frac{1}{\sqrt{T}} \frac{1}{\sigma_r} \sqrt{\lambda_{\min}(\boldsymbol{\Sigma}_\varepsilon)} \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2 = \frac{1}{\sqrt{n}} \frac{1}{\kappa_\varepsilon} \rho \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2, \quad (\text{D.13})$$

where we use the fact that $\|\boldsymbol{\Sigma}_\varepsilon\|_2 = \lambda_{\max}(\boldsymbol{\Sigma}_\varepsilon)$ and $\kappa_\varepsilon = \lambda_{\max}(\boldsymbol{\Sigma}_\varepsilon)/\lambda_{\min}(\boldsymbol{\Sigma}_\varepsilon)$ is the condition number of $\boldsymbol{\Sigma}_\varepsilon$. So, when the entries of \mathbf{Z} are sub-Gaussian, we obtain that

$$s_3 \lesssim |S|^{3/2} \frac{1}{\sqrt{T}} \|\boldsymbol{\Sigma}_\varepsilon^{1/2} \mathbf{U}\|_{2,\infty} \frac{1}{\sigma_r} \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2 \frac{1}{\frac{1}{\sqrt{n}} \frac{1}{\kappa_\varepsilon} \rho \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2} = \kappa_\varepsilon \sqrt{\frac{n}{T}} |S|^{3/2} \frac{\|\boldsymbol{\Sigma}_\varepsilon^{1/2} \bar{\mathbf{U}}\|_{2,\infty}}{\|\boldsymbol{\Sigma}_\varepsilon^{1/2}\|_2}.$$

By definition of s_3 , we obtain by the inequality (D.9) in Step 1 for $\mathbb{P}(\mathcal{T}(S, \mathbf{v}) + \zeta \leq \chi_{1-\alpha}^2(|S| - r))$ that

$$\begin{aligned} & \mathbb{P}\left(\sqrt{\chi^2(|S| - r)} \leq \sqrt{\chi_{1-\alpha}^2(|S| - r) - |\zeta| - |\tau|}\right) - s_3 \\ & \leq \mathbb{P}(\mathcal{T}(S, \mathbf{v}) + \zeta \leq \chi_{1-\alpha}^2(|S| - r)) \\ & \leq \mathbb{P}\left(\sqrt{\chi^2(|S| - r)} \leq \sqrt{\chi_{1-\alpha}^2(|S| - r) + |\zeta| + |\tau|}\right) + s_3. \end{aligned} \quad (\text{D.14})$$

Step 3 - Establishing chi-squared distributional characterization.

By Lemma 14, we have that $\chi_{1-\alpha}^2(|S| - r) \asymp (|S| - r) \asymp |S|$, where we use the fact that $r \ll |S|$ and thus $(|S| - r) \asymp |S|$. Since we assume that $|\zeta| \ll |S|$, we have that $|\zeta| \lesssim \chi_{1-\alpha}^2(|S| - r)$. For $a > 0$ and $b \in \mathbb{R}$ satisfying $|b| \ll a$, we have that $|\sqrt{a+b} - \sqrt{a}| = \frac{|b|}{\sqrt{a+b} + \sqrt{a}} \lesssim \frac{|b|}{\sqrt{a}}$. So we obtain that $\sqrt{\chi_{1-\alpha}^2(|S| - r) - |\zeta|} \geq \sqrt{\chi_{1-\alpha}^2(|S| - r) - g}$ and $\sqrt{\chi_{1-\alpha}^2(|S| - r) + |\zeta|} \leq \sqrt{\chi_{1-\alpha}^2(|S| - r) + g}$, where $g > 0$ and g satisfies that $g \lesssim \frac{1}{\sqrt{\chi_{1-\alpha}^2(|S| - r)}} |\zeta| \lesssim \frac{1}{\sqrt{|S|}} |\zeta|$. Then, using the inequality (D.14), we conclude that, for $h = g + |\tau|$, it holds

$$\begin{aligned} & \mathbb{P}\left(\sqrt{\chi^2(|S| - r)} \leq \sqrt{\chi_{1-\alpha}^2(|S| - r) - h}\right) - s_3 \\ & \leq \mathbb{P}(\mathcal{T}(S, \mathbf{v}) + \zeta \leq \chi_{1-\alpha}^2(|S| - r)) \\ & \leq \mathbb{P}\left(\sqrt{\chi^2(|S| - r)} \leq \sqrt{\chi_{1-\alpha}^2(|S| - r) + h}\right) + s_3. \end{aligned} \quad (\text{D.15})$$

Here, $h > 0$ satisfies that $h \lesssim |\tau| + \frac{1}{\sqrt{|S|}} |\zeta|$.

Note that, for any $R > 0$, we have that $\mathbb{P}(\sqrt{\chi^2(n)} \leq R) = \mathbb{P}(\mathcal{N}(0, \mathbf{I}_n) \in B(R))$ where $B(R)$ is the ball. So, we obtain by Lemma 15 that

$$\begin{aligned} & \left| \mathbb{P}\left(\sqrt{\chi^2(|S| - r)} \leq \sqrt{\chi_{1-\alpha}^2(|S| - r) \pm h}\right) - (1 - \alpha) \right| \\ & = \left| \mathbb{P}\left(\sqrt{\chi^2(|S| - r)} \leq \sqrt{\chi_{1-\alpha}^2(|S| - r) \pm h}\right) - \mathbb{P}\left(\sqrt{\chi^2(|S| - r)} \leq \sqrt{\chi_{1-\alpha}^2(|S| - r)}\right) \right| \lesssim h. \end{aligned}$$

Thus, using the two-side bounds for $\mathbb{P}(\mathcal{T}(S, \mathbf{v}) + \zeta \leq \chi_{1-\alpha}^2(|S| - r))$ in the inequality (D.15), we obtain that

$$\left| \mathbb{P}(\mathcal{T}(S, \mathbf{v}) + \zeta \leq \chi_{1-\alpha}^2(|S| - r)) - (1 - \alpha) \right| \lesssim h + s_3 \lesssim |\tau| + \frac{1}{\sqrt{|S|}} |\zeta| + s_3.$$

So, to prove the desired result in the lemma, it suffices to establish the upper bound for $|\tau|$, which is defined in (D.8).

Step 4 - Deriving an upper bound for $|\tau|$.

For τ defined in (D.8), using (D.7) and the triangle inequality, we obtain that

$$|\tau| = \frac{1}{\|\mathbf{m}\|_2} \left| \|\mathbf{H}^\top \mathbf{u} + \mathbf{H}^\top \boldsymbol{\Upsilon}_P\|_2 - \|\mathbf{H}^\top \mathbf{u}\|_2 \right| \leq \frac{1}{\|\mathbf{m}\|_2} \|\mathbf{H}^\top \boldsymbol{\Upsilon}_P\|_2 \leq \frac{1}{\|\mathbf{m}\|_2} \|\mathbf{H}\|_2 \|\boldsymbol{\Upsilon}_P\|_2 = \frac{1}{\|\mathbf{m}\|_2} \|\boldsymbol{\Upsilon}_P\|_2,$$

where the last equality uses the fact that $\|\mathbf{H}\|_2 = 1$ since $\mathbf{H}^\top \mathbf{H} = \mathbf{I}_{|S|-r}$. Combining the lower bound of $\|\mathbf{m}\|_2$ in (D.13) and the upper bound of $\|\Upsilon_P\|_2$ in Lemma 9, we obtain that

$$\begin{aligned} |\tau| &\leq \frac{1}{\|\mathbf{m}\|_2} \|\Upsilon_P\|_2 \\ &\lesssim \frac{\kappa_\varepsilon \sqrt{n}}{\rho \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2} \sqrt{|S|} \left(\rho^2 \frac{1}{\sqrt{n}} \sqrt{\frac{n}{T}} + \rho \frac{1}{\sqrt{nT}} \right) r \log^2 n \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2 \\ &= \lesssim \kappa_\varepsilon \sqrt{|S|} \left(\rho \sqrt{\frac{n}{T}} + \frac{1}{\sqrt{T}} \right) r \log^2 n. \end{aligned}$$

Finally, the lower bound of ϕ can be proven using the lower bound of $\|\mathbf{m}\|_2$ in (D.13) as follows,

$$\phi = \|\mathbf{m}\|_2^2 \gtrsim \left(\frac{1}{\sqrt{n}} \frac{1}{\kappa_\varepsilon} \rho \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2 \right)^2 = \frac{1}{\kappa_\varepsilon^2} \frac{1}{n} \rho^2 \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2^2,$$

which is our desired result. \square

D.2 Proof of Theorem 2

To prove the desired results, we use Lemma 10 with the small perturbation $\zeta = \widehat{\mathcal{T}}(S, \mathbf{v}) - \mathcal{T}(S, \mathbf{v})$. Then, the problem boils down to deriving the upper bound for $\frac{1}{\sqrt{|S|}} |\zeta|$. We do the following decomposition:

$$\zeta = \widehat{\mathcal{T}}(S, \mathbf{v}) - \mathcal{T}(S, \mathbf{v}) = r_1 + r_2,$$

where r_1 and r_2 are respectively defined by

$$r_1 = \frac{1}{\phi} \mathbf{v}^\top (\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}) \mathbf{v} - \mathcal{T}(S, \mathbf{v}) \quad \text{and} \quad r_2 = \widehat{\mathcal{T}}(S, \mathbf{v}) - \frac{1}{\phi} \mathbf{v}^\top (\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}) \mathbf{v}.$$

Step 1 – Upper bound for $\frac{1}{\sqrt{|S|}} |r_1|$.

Since $(\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}})$ is idempotent, we can rewrite r_1 as

$$\begin{aligned} r_1 &= \frac{1}{\phi} \mathbf{v}^\top (\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}) (\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}) (\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}) \mathbf{v} - \frac{1}{\phi} \mathbf{v}^\top (\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}) (\mathbf{I}_{|S|} - \mathbf{P}_{\mathbf{V}_{S,\cdot}}) (\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}) \mathbf{v} \\ &= \frac{1}{\phi} \mathbf{v}^\top (\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}) (\mathbf{P}_{\mathbf{V}_{S,\cdot}} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}) (\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}) \mathbf{v}. \end{aligned}$$

Then we get

$$\begin{aligned} |r_1| &\leq \frac{1}{\phi} \|\mathbf{P}_{\mathbf{V}_{S,\cdot}} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}\|_2 \|(\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}) \mathbf{v}\|_2^2 \\ &\stackrel{(i)}{\lesssim} \frac{\kappa_\varepsilon^2 n}{\rho^2 \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2^2} \rho \sqrt{\frac{T}{n}} \sqrt{r} \log^{3/2} n \left(\sqrt{|S|} \rho \sqrt{\frac{r}{n}} \log n \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2 \right)^2 \\ &\leq \kappa_\varepsilon^2 |S| \rho \sqrt{\frac{T}{n}} r^{3/2} \log^{7/2} n, \end{aligned}$$

where (i) uses the results in Lemmas 9 and 10. So we obtain

$$\frac{|r_1|}{\sqrt{|S|}} \lesssim \kappa_\varepsilon^2 \rho \sqrt{|S|} \sqrt{\frac{T}{n}} r^{3/2} \log^{7/2} n \ll 1.$$

Step 2 – Upper bound for $\frac{1}{\sqrt{|S|}} |r_2|$.

Since $(\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}})$ is idempotent, we can rewrite r_2 as

$$\begin{aligned} r_2 &= \frac{1}{\widehat{\phi}} \mathbf{v}^\top (\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}) (\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}) \mathbf{v} - \frac{1}{\phi} \mathbf{v}^\top (\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}) (\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}) \mathbf{v} \\ &= \left(\frac{1}{\widehat{\phi}} - \frac{1}{\phi} \right) \| (\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}) \mathbf{v} \|_2^2 = \frac{\phi - \widehat{\phi}}{\widehat{\phi}} \frac{\| (\mathbf{I}_{|S|} - \mathbf{P}_{\widehat{\mathbf{V}}_{S,\cdot}}) \mathbf{v} \|_2^2}{\phi}. \end{aligned}$$

Then, using the results in Lemmas 9 and 10, we obtain that

$$|r_2| \lesssim \frac{1}{\widehat{\phi}} |\widehat{\phi} - \phi| \cdot \frac{\left(\sqrt{|S|} \rho \sqrt{\frac{\varepsilon}{n}} \log n \| (\mathbf{V}_{S,\cdot})^+ \mathbf{v} \|_2 \right)^2}{\frac{1}{\kappa_\varepsilon^2} \frac{1}{n} \rho^2 \| (\mathbf{V}_{S,\cdot})^+ \mathbf{v} \|_2^2} \leq \frac{1}{\widehat{\phi}} |\widehat{\phi} - \phi| \cdot |S| \kappa_\varepsilon^2 r \log^2 n. \quad (\text{D.16})$$

So, to derive the upper bound for $|r_2|$, we need to derive the upper bound of $\frac{1}{\widehat{\phi}} |\widehat{\phi} - \phi|$.

Note that $\widehat{\phi}$ and ϕ are nonnegative owing the property of positive semidefinite matrix. We will derive the upper bound for $|\widehat{\phi} - \phi|$ first, and then use it and the lower bound of ϕ obtained in Lemma 10 to get the lower bound of $\widehat{\phi}$, because $\widehat{\phi} \geq \phi - |\widehat{\phi} - \phi|$. Our starting point is to rewrite ϕ as the following form:

$$\widehat{\phi} = \frac{1}{T} ((\widehat{\mathbf{V}}_{S,\cdot} \mathbf{R}_V)^+ \mathbf{v})^\top ((\mathbf{R}_U)^\top \widehat{\boldsymbol{\Sigma}}^{-1} \mathbf{R}_V)^\top (\widehat{\mathbf{U}} \mathbf{R}_U)^\top \widehat{\boldsymbol{\Sigma}}_\varepsilon^\top (\widehat{\mathbf{U}} \mathbf{R}_U) ((\mathbf{R}_U)^\top \widehat{\boldsymbol{\Sigma}}^{-1} \mathbf{R}_V) (\widehat{\mathbf{V}}_{S,\cdot} \mathbf{R}_V)^+ \mathbf{v},$$

where we use the fact that $(\widehat{\mathbf{V}}_{S,\cdot} \mathbf{R}_V)^+ = (\mathbf{R}_V)^\top (\widehat{\mathbf{V}}_{S,\cdot})^+$ since $\widehat{\mathbf{V}}_{S,\cdot}$ has full column rank and $(\mathbf{V}_{S,\cdot})^+ = [(\mathbf{V}_{S,\cdot})^\top \mathbf{V}_{S,\cdot}]^{-1} (\mathbf{V}_{S,\cdot})^\top$. Recall that

$$\phi = \frac{1}{T} ((\mathbf{V}_{S,\cdot})^+ \mathbf{v})^\top \boldsymbol{\Lambda}^{-1} \mathbf{U}^\top \boldsymbol{\Sigma}_\varepsilon \mathbf{U} \boldsymbol{\Lambda}^{-1} (\mathbf{V}_{S,\cdot})^+ \mathbf{v}.$$

Next, for simplicity of notations, we denote

$$A_0 = (\mathbf{V}_{S,\cdot})^+, \quad B_0 = \boldsymbol{\Lambda}^{-1}, \quad C_0 = \mathbf{U}, \quad D_0 = \boldsymbol{\Sigma}_\varepsilon,$$

and

$$A_1 = (\widehat{\mathbf{V}}_{S,\cdot} \mathbf{R}_V)^+, \quad B_1 = (\mathbf{R}_U)^\top \widehat{\boldsymbol{\Sigma}}^{-1} \mathbf{R}_V, \quad C_1 = \widehat{\mathbf{U}} \mathbf{R}_U, \quad D_1 = \widehat{\boldsymbol{\Sigma}}_\varepsilon^\top,$$

so that $\widehat{\phi}$ and ϕ can be written as

$$\widehat{\phi} = \frac{1}{T} \mathbf{v}^\top A_1^\top B_1^\top C_1^\top D_1 C_1 B_1 A_1 \mathbf{v} \quad \text{and} \quad \phi = \frac{1}{T} \mathbf{v}^\top A_0^\top B_0^\top C_0^\top D_0 C_0 B_0 A_0 \mathbf{v},$$

respectively.

To derive the upper bound for $|\widehat{\phi} - \phi|$, we will establish the upper bounds for $\|A_1 - A_0\|_2$, $\|B_1 - B_0\|_2$, $\|C_1 - C_0\|_2$, and $\|D_1 - D_0\|_2$ respectively.

Step 2.1. For $\|A_1 - A_0\|_2$, using Theorem 3.3 in Stewart (1977), we obtain that

$$\begin{aligned} \|A_1 - A_0\|_2 &= \|(\widehat{\mathbf{V}}_{S,\cdot} \mathbf{R}_V)^+ - (\mathbf{V}_{S,\cdot})^+\|_2 \\ &\lesssim \max(\|(\mathbf{V}_{S,\cdot})^+\|_2^2, \|(\widehat{\mathbf{V}}_{S,\cdot} \mathbf{R}_V)^+\|_2^2) \|\widehat{\mathbf{V}}_{S,\cdot} \mathbf{R}_V - \mathbf{V}_{S,\cdot}\|_2. \end{aligned}$$

For the term $\|\widehat{\mathbf{V}}_{S,\cdot} \mathbf{R}_V - \mathbf{V}_{S,\cdot}\|_2$, we have that

$$\|\widehat{\mathbf{V}}_{S,\cdot} \mathbf{R}_V - \mathbf{V}_{S,\cdot}\|_2 \leq \sqrt{|S|} \|\widehat{\mathbf{V}}_{S,\cdot} \mathbf{R}_V - \mathbf{V}_{S,\cdot}\|_{2,\infty} \leq \sqrt{|S|} \|\widehat{\mathbf{V}} \mathbf{R}_V - \mathbf{V}\|_{2,\infty} \lesssim \sqrt{|S|} \rho \sqrt{\frac{r}{n}} \log n,$$

where the last inequality uses the upper bound of $\|\widehat{\mathbf{V}} \mathbf{R}_V - \mathbf{V}\|_{2,\infty}$ obtained in the proof of Corollary 1. We obtain by Lemma 8 that $\sigma_i(\mathbf{V}_{S,\cdot}) \asymp \sqrt{\frac{|S|}{T}}$ for $i = 1, 2, \dots, r$. Next, since (5.2) implies that $\rho \sqrt{\frac{T}{n}} \sqrt{r} \log n \ll$

1, we obtain that $\|\widehat{\mathbf{V}}_{S,\cdot}\mathbf{R}_V - \mathbf{V}_{S,\cdot}\|_2 \lesssim \sqrt{|S|}\rho\sqrt{\frac{T}{n}}\log n \ll \sqrt{\frac{|S|}{T}}$, and thus $\sigma_i(\widehat{\mathbf{V}}_{S,\cdot}\mathbf{R}_V) \asymp \sqrt{\frac{|S|}{T}}$ for $i = 1, 2, \dots, r$. Then, similar to the proof of $\sigma_i((\mathbf{V}_{S,\cdot})^+) \asymp \sqrt{\frac{T}{|S|}}$ for $i = 1, 2, \dots, r$ in Lemma 8, we obtain that $\sigma_i((\widehat{\mathbf{V}}_{S,\cdot}\mathbf{R}_V)^+) \asymp \sqrt{\frac{T}{|S|}}$ for $i = 1, 2, \dots, r$. So we obtain

$$\begin{aligned} \|A_1 - A_0\|_2 &\lesssim \max(\|(\mathbf{V}_{S,\cdot})^+\|_2^2, \|(\widehat{\mathbf{V}}_{S,\cdot}\mathbf{R}_V)^+\|_2^2) \|\widehat{\mathbf{V}}_{S,\cdot}\mathbf{R}_V - \mathbf{V}_{S,\cdot}\|_2 \\ &\lesssim \frac{T}{|S|} \cdot \sqrt{|S|}\rho\sqrt{\frac{r}{n}}\log n = \frac{T}{\sqrt{n|S|}}\rho\sqrt{r}\log n. \end{aligned}$$

Also, we obtain by Lemma 8 that $\|A_0\|_2 = \|(\mathbf{V}_{S,\cdot})^+\|_2 \lesssim \sqrt{\frac{T}{|S|}}$. Since $\sigma_i((\widehat{\mathbf{V}}_{S,\cdot}\mathbf{R}_V)^+) \asymp \sqrt{\frac{T}{|S|}}$ for $i = 1, 2, \dots, r$, we obtain that $\|A_1\|_2 = \|(\widehat{\mathbf{V}}_{S,\cdot}\mathbf{R}_V)^+\|_2 \lesssim \sqrt{\frac{T}{|S|}}$.

For $\|B_1 - B_0\|_2$, since \mathbf{R}_U and \mathbf{R}_V are rotation matrices, we have that

$$\begin{aligned} \|B_1 - B_0\|_2 &= \|(\mathbf{R}_U)^\top \widehat{\Sigma}^{-1} \mathbf{R}_V - \Lambda^{-1}\|_2 \\ &= \|[(\mathbf{R}_U)^\top \widehat{\Sigma}^{-1} - \Lambda^{-1}(\mathbf{R}_V)^\top] \mathbf{R}_V\|_2 \\ &\leq \|\Lambda^{-1}[\Lambda(\mathbf{R}_U)^\top - (\mathbf{R}_V)^\top \widehat{\Sigma}] \widehat{\Sigma}^{-1}\|_2 \leq \frac{1}{\lambda_r} \|\Lambda(\mathbf{R}_U)^\top - (\mathbf{R}_V)^\top \widehat{\Sigma}\|_2 \frac{1}{\widehat{\sigma}_r} \\ &\stackrel{(i)}{\lesssim} \frac{1}{\widehat{\sigma}_r^2} \|[\Lambda - (\mathbf{R}_V)^\top \widehat{\Sigma} \mathbf{R}_U] (\mathbf{R}_U)^\top\|_2 \leq \frac{1}{\widehat{\sigma}_r^2} \|(\mathbf{R}_U)^\top \widehat{\Sigma} \mathbf{R}_V - \Lambda\|_2 \\ &\stackrel{(ii)}{\lesssim} \frac{1}{\widehat{\sigma}_r^2} \left(\sigma_r \rho^2 + \sigma_r \rho \sqrt{\frac{r}{n}} \sqrt{\log n} \right) = \frac{1}{\sigma_r} \left(\rho^2 + \rho \sqrt{\frac{r}{n}} \sqrt{\log n} \right), \end{aligned}$$

where (i) uses $\widehat{\sigma}_i \asymp \sigma_i$ and $\lambda_i \asymp \sigma_i$ in Lemmas 2 and 4, respectively, and (ii) uses Lemma 5. Also, since $\|B_0\|_2 = \|\Lambda^{-1}\|_2 \lesssim \frac{1}{\sigma_r}$ and the above result implies that $\|B_1 - B_0\|_2 \ll \frac{1}{\sigma_r}$ owing to $\rho + \sqrt{\frac{r}{n}} \sqrt{\log n} \ll 1$ as assumed, so we obtain $\|B_1\|_2 \leq \|B_1 - B_0\|_2 + \|B_0\|_2 \lesssim \frac{1}{\sigma_r}$.

For $\|C_1 - C_0\|_2 = \|\widehat{\mathbf{U}} \mathbf{R}_U - \mathbf{U}\|_2$, we obtain by Lemma 4 that $\|C_1 - C_0\|_2 \lesssim \rho$. Also, since $\widehat{\mathbf{U}}^\top \widehat{\mathbf{U}} = \mathbf{U}^\top \mathbf{U} = \mathbf{I}_r$ and $\mathbf{R}_U \in \mathcal{O}^{r \times r}$ is a rotation matrix, we have that $\|C_1\|_2 = \|\widehat{\mathbf{U}} \mathbf{R}_U\|_2 \leq 1$ and $\|C_0\|_2 = \|\mathbf{U}\|_2 \leq 1$.

For $\|D_1 - D_0\|_2 = \|\widehat{\Sigma}_\varepsilon^r - \Sigma_\varepsilon\|_2$, by assumption we have that $\|D_1 - D_0\|_2 \sqrt{|S|} \cdot \kappa_\varepsilon^4 r \log^2 n \ll \|\Sigma_\varepsilon\|_2$.

In particular, the above results imply that, $\|A_i\|_2 \lesssim K_A$, $\|B_i\|_2 \lesssim K_B$, $\|C_i\|_2 \lesssim K_C$, and $\|D_i\|_2 \lesssim K_D$ for $i = 0, 1$, where $K_A = \sqrt{\frac{T}{|S|}}$, $K_B = \frac{1}{\sigma_r}$, $K_C = 1$, and $K_D = \|\Sigma_\varepsilon\|_2$.

Step 2.2. We obtain by the representations of $\widehat{\phi}$ and ϕ that,

$$|\widehat{\phi} - \phi| \lesssim \frac{1}{T} \left(\frac{\|A_1 - A_0\|_2}{K_A} + \frac{\|B_1 - B_0\|_2}{K_B} + \frac{\|C_1 - C_0\|_2}{K_C} + \frac{\|D_1 - D_0\|_2}{K_D} \right) K_A^2 K_B^2 K_C^2 K_D \|\mathbf{v}\|_2^2.$$

Then using the results in Step 2.1, we obtain that

$$\begin{aligned} &|\widehat{\phi} - \phi| \\ &\lesssim \frac{1}{T} \left| \frac{1}{\|\Sigma_\varepsilon\|_2} \|\widehat{\Sigma}_\varepsilon^r - \Sigma_\varepsilon\|_2 + \frac{\rho}{1} + \frac{\frac{1}{\sigma_r} \rho \left(\rho + \sqrt{\frac{r}{n}} \sqrt{\log n} \right)}{\frac{1}{\sigma_r}} + \frac{\frac{T}{\sqrt{n|S|}} \rho \sqrt{r} \|\mathbf{v}\|_2}{\sqrt{\frac{T}{|S|}} \|\mathbf{v}\|_2} \right| \cdot \frac{T}{|S|} \cdot \frac{1}{\sigma_r^2} \cdot \|\Sigma_\varepsilon\|_2 \cdot \|\mathbf{v}\|_2^2 \\ &\lesssim \left| \frac{1}{\|\Sigma_\varepsilon\|_2} \|\widehat{\Sigma}_\varepsilon^r - \Sigma_\varepsilon\|_2 + \rho + \rho \left(\rho + \sqrt{\frac{r}{n}} \sqrt{\log n} \right) + \rho \sqrt{\frac{T}{n}} \sqrt{r} \right| \cdot \frac{1}{|S|} \rho^2 \frac{T}{n} \|\mathbf{v}\|_2^2 \\ &\stackrel{(i)}{\lesssim} \left| \frac{1}{\|\Sigma_\varepsilon\|_2} \|\widehat{\Sigma}_\varepsilon^r - \Sigma_\varepsilon\|_2 + \rho + \rho \sqrt{\frac{T}{n}} \sqrt{r} \right| \cdot \frac{1}{|S|} \rho^2 \frac{T}{n} \|\mathbf{v}\|_2^2. \end{aligned}$$

where (i) uses in (5.2). We obtain by the bound of ϕ in Lemma 10 and the bound of $\sigma_{\min}((\mathbf{V}_{S,\cdot})^+)$ in Lemma 8 that

$$\phi \gtrsim \frac{1}{\kappa_\varepsilon^2 n} \rho^2 \|(\mathbf{V}_{S,\cdot})^+ \mathbf{v}\|_2^2 \geq \frac{1}{\kappa_\varepsilon^2 n} \rho^2 [\sigma_{\min}((\mathbf{V}_{S,\cdot})^+)]^2 \|\mathbf{v}\|_2^2 \gtrsim \frac{1}{\kappa_\varepsilon^2 n} \rho^2 \frac{T}{|S|} \|\mathbf{v}\|_2^2.$$

Then, we conclude that $|\widehat{\phi} - \phi| \ll \phi$ because the ratio between the above upper bound of $|\widehat{\phi} - \phi|$ and the above lower bound of ϕ is given by

$$\frac{\left| \frac{1}{\|\widehat{\Sigma}_\varepsilon\|_2} \|\widehat{\Sigma}_\varepsilon^\tau - \Sigma_\varepsilon\|_2 + \rho + \rho \sqrt{\frac{T}{n}} \sqrt{r} \right| \cdot \frac{1}{|S|} \rho^2 \frac{T}{n} \|\mathbf{v}\|_2^2}{\frac{1}{\kappa_\varepsilon^2 n} \rho^2 \frac{T}{|S|} \|\mathbf{v}\|_2^2} = \kappa_\varepsilon^2 \left| \frac{1}{\|\Sigma_\varepsilon\|_2} \|\widehat{\Sigma}_\varepsilon^\tau - \Sigma_\varepsilon\|_2 + \rho + \rho \sqrt{\frac{T}{n}} \sqrt{r} \right| \ll 1,$$

where the last inequality is owing to the assumptions. So $\widehat{\phi} \geq \phi - |\widehat{\phi} - \phi| \gtrsim \phi \gtrsim \frac{1}{\kappa_\varepsilon^2 n} \rho^2 \frac{T}{|S|} \|\mathbf{v}\|_2^2$. Using this lower bound of $\widehat{\phi}$ and the above upper bound of $|\widehat{\phi} - \phi|$, we obtain that

$$\frac{1}{\widehat{\phi}} |\widehat{\phi} - \phi| \lesssim \frac{\left| R_\varepsilon + \rho + \rho \sqrt{\frac{T}{n}} \sqrt{r} \right| \cdot \frac{1}{|S|} \rho^2 \frac{T}{n} \|\mathbf{v}\|_2^2}{\frac{1}{\kappa_\varepsilon^2 n} \rho^2 \frac{T}{|S|} \|\mathbf{v}\|_2^2} = \kappa_\varepsilon^2 \left| \frac{1}{\|\Sigma_\varepsilon\|_2} \|\widehat{\Sigma}_\varepsilon^\tau - \Sigma_\varepsilon\|_2 + \rho + \rho \sqrt{\frac{T}{n}} \sqrt{r} \right|. \quad (\text{D.17})$$

Using (D.16), (D.17), and the fact that $\kappa_\varepsilon \geq 1$, we obtain that

$$\begin{aligned} \frac{|r_2|}{\sqrt{|S|}} &\lesssim \frac{1}{\widehat{\phi}} |\widehat{\phi} - \phi| \sqrt{|S|} \kappa_\varepsilon^2 r \log^2 n \lesssim \left| \frac{1}{\|\Sigma_\varepsilon\|_2} \|\widehat{\Sigma}_\varepsilon^\tau - \Sigma_\varepsilon\|_2 + \rho + \sqrt{\frac{T}{n}} \rho \sqrt{r} \right| \sqrt{|S|} \kappa_\varepsilon^4 r \log^2 n. \\ &\lesssim \frac{1}{\|\Sigma_\varepsilon\|_2} \|\widehat{\Sigma}_\varepsilon^\tau - \Sigma_\varepsilon\|_2 \sqrt{|S|} \kappa_\varepsilon^4 r \log^2 n + \rho \sqrt{|S|} \kappa_\varepsilon^4 r^2 \log^2 n. \end{aligned}$$

Step 3. We now assemble all the above upper bounds. We obtain by Lemma 10 that

$$\begin{aligned} &\left| \mathbb{P} \left(\widehat{\mathcal{T}}(S, \mathbf{v}) \leq \chi_{1-\alpha}^2(|S| - r) \right) - (1 - \alpha) \right| \\ &\lesssim \sqrt{|S|} \left(\rho \sqrt{\frac{n}{T}} + \frac{1}{\sqrt{T}} \right) \kappa_\varepsilon r \log^2 n + \kappa_\varepsilon^2 \rho \sqrt{|S|} \sqrt{\frac{T}{n}} r^{3/2} \log^{7/2} n \\ &\quad + \frac{1}{\|\Sigma_\varepsilon\|_2} \|\widehat{\Sigma}_\varepsilon^\tau - \Sigma_\varepsilon\|_2 \sqrt{|S|} \kappa_\varepsilon^4 r \log^2 n + \rho \sqrt{|S|} \kappa_\varepsilon^4 r^2 \log^2 n + s_3 \\ &\lesssim \frac{1}{\|\Sigma_\varepsilon\|_2} \|\widehat{\Sigma}_\varepsilon^\tau - \Sigma_\varepsilon\|_2 \sqrt{|S|} \kappa_\varepsilon^4 r \log^2 n + \sqrt{|S|} \rho \sqrt{\frac{n}{T}} \kappa_\varepsilon^4 r^2 \log^4 n + \frac{\sqrt{|S|}}{\sqrt{T}} \kappa_\varepsilon r \log^2 n + s_3 \end{aligned}$$

where, as defined in Lemma 10,

$$s_3 \lesssim \kappa_\varepsilon \sqrt{\frac{n}{T}} |S|^{3/2} \frac{\|\Sigma_\varepsilon^{1/2} \widehat{\mathbf{U}}\|_{2,\infty}}{\|\Sigma_\varepsilon^{1/2}\|_2},$$

and, if all the entries of the matrix \mathbf{Z} are Gaussian, i.e., the noise is Gaussian, then the above inequality for $|\mathbb{P}(\widehat{\mathcal{T}}(S, \mathbf{v}) \leq \chi_{1-\alpha}^2(|S| - r)) - (1 - \alpha)|$ holds when $s_3 = 0$. Then the desired non-asymptotic results follow from the above upper bound of $|\mathbb{P}(\widehat{\mathcal{T}}(S, \mathbf{v}) \leq \chi_{1-\alpha}^2(|S| - r)) - (1 - \alpha)|$.

D.3 Estimation of the idiosyncratic noise covariance matrix

Step 1 – Estimation error for the sample covariance matrix $\widehat{\Sigma}_\varepsilon = \frac{1}{T} \widehat{\mathbf{E}} \widehat{\mathbf{E}}^\top$.

We obtain by $x_{i,t} = \mathbf{b}_i^\top \mathbf{f}_t + \varepsilon_{i,t}$ that $\widehat{\varepsilon}_{i,t} - \varepsilon_{i,t} = \mathbf{b}_i^\top \mathbf{f}_t - \widehat{\mathbf{b}}_i^\top \widehat{\mathbf{f}}_t$. So, the sample covariance estimator $\widehat{\Sigma}_\varepsilon$ is given by

$$\begin{aligned} \widehat{\Sigma}_\varepsilon &= \frac{1}{T} \widehat{\mathbf{E}} \widehat{\mathbf{E}}^\top = \frac{1}{T} (\mathbf{X} - \widehat{\mathbf{B}} \widehat{\mathbf{F}}^\top) (\mathbf{X} - \widehat{\mathbf{B}} \widehat{\mathbf{F}}^\top)^\top \\ &= \frac{1}{T} (\mathbf{X} \mathbf{X}^\top - \mathbf{X} \widehat{\mathbf{B}} \widehat{\mathbf{F}}^\top - \widehat{\mathbf{B}} \widehat{\mathbf{F}}^\top \mathbf{X} + \widehat{\mathbf{B}} \widehat{\mathbf{F}}^\top \widehat{\mathbf{F}} \widehat{\mathbf{B}}^\top) = \frac{1}{T} (\mathbf{X} \mathbf{X}^\top - T \widehat{\mathbf{B}} \widehat{\mathbf{B}}^\top), \end{aligned}$$

where we use the fact that $\frac{1}{T} \mathbf{X} \widehat{\mathbf{F}} = \frac{1}{T} \mathbf{X} \widehat{\mathbf{V}} = \widehat{\mathbf{U}} \widehat{\Sigma} = \widehat{\mathbf{B}}$. Then, we use the identity $\mathbf{X} = \mathbf{B} \mathbf{F}^\top + \mathbf{E}$ to obtain that

$$\widehat{\Sigma}_\varepsilon = \frac{1}{T} ((\mathbf{B} \mathbf{F}^\top + \mathbf{E})(\mathbf{B} \mathbf{F}^\top + \mathbf{E})^\top - T \widehat{\mathbf{B}} \widehat{\mathbf{B}}^\top)$$

$$= \frac{1}{T} \mathbf{E} \mathbf{E}^\top + \left(\frac{1}{T} \mathbf{E} \mathbf{F} \mathbf{B}^\top + \frac{1}{T} \mathbf{B} \mathbf{F}^\top \mathbf{E}^\top \right) + \left(\frac{1}{T} \mathbf{B} \mathbf{F}^\top \mathbf{F} \mathbf{B}^\top - \widehat{\mathbf{B}} \widehat{\mathbf{B}}^\top \right).$$

Then, using the fact that $\frac{1}{T} \mathbf{B} \mathbf{F}^\top \mathbf{F} \mathbf{B}^\top = \mathbf{U} \mathbf{\Lambda}^2 \mathbf{U}^\top$ and $\widehat{\mathbf{B}} \widehat{\mathbf{B}}^\top = \widehat{\mathbf{U}} \widehat{\mathbf{\Sigma}}^2 \widehat{\mathbf{U}}$, we obtain that

$$\widehat{\mathbf{\Sigma}}_\varepsilon - \mathbf{\Sigma}_\varepsilon = \left(\frac{1}{T} \mathbf{E} \mathbf{E}^\top - \mathbb{E} \left[\frac{1}{T} \mathbf{E} \mathbf{E}^\top \right] \right) + \left(\frac{1}{T} \mathbf{E} \mathbf{F} \mathbf{B}^\top + \frac{1}{T} \mathbf{B} \mathbf{F}^\top \mathbf{E}^\top \right) + \left(\mathbf{U} \mathbf{\Lambda}^2 \mathbf{U}^\top - \widehat{\mathbf{U}} \widehat{\mathbf{\Sigma}}^2 \widehat{\mathbf{U}} \right).$$

We now look at the estimator error $|(\widehat{\mathbf{\Sigma}}_\varepsilon - \mathbf{\Sigma}_\varepsilon)_{i,j}|$ of the (i, j) -th entry by analyzing the terms in the above three brackets.

Step 1.1 – Bound for $(\frac{1}{T} \mathbf{E} \mathbf{E}^\top - \mathbb{E}[\frac{1}{T} \mathbf{E} \mathbf{E}^\top])$.

First, we have that

$$\left(\frac{1}{T} \mathbf{E} \mathbf{E}^\top - \mathbb{E} \left[\frac{1}{T} \mathbf{E} \mathbf{E}^\top \right] \right)_{i,j} = \frac{1}{T} \sum_{t=1}^T \mathbf{E}_{i,t} \mathbf{E}_{j,t} - \mathbb{E}[\mathbf{E}_{i,t} \mathbf{E}_{j,t}]$$

is the sum of T independent zero-mean sub-Gaussian random variables. Since $\mathbf{E}_{i,t} = \sum_{k=1}^N (\mathbf{\Sigma}_\varepsilon^{1/2})_{i,k} \mathbf{Z}_{k,t}$, we obtain that $\|\mathbf{E}_{i,t}\|_{\psi_2} \lesssim (\sum_{k=1}^N ((\mathbf{\Sigma}_\varepsilon^{1/2})_{i,k})^2)^{1/2} = \sqrt{(\mathbf{\Sigma}_\varepsilon)_{i,i}}$. By Lemma 2.7.7 in [Wainwright \(2019\)](#), we obtain that $\mathbf{E}_{i,t} \mathbf{E}_{j,t}$ is sub-exponential and $\|\mathbf{E}_{i,t} \mathbf{E}_{j,t}\|_{\psi_1} \leq \|\mathbf{E}_{i,t}\|_{\psi_2} \|\mathbf{E}_{j,t}\|_{\psi_2} \lesssim \sqrt{(\mathbf{\Sigma}_\varepsilon)_{i,i} (\mathbf{\Sigma}_\varepsilon)_{j,j}}$, and the centering $\mathbf{E}_{i,t} \mathbf{E}_{j,t} - \mathbb{E}[\mathbf{E}_{i,t} \mathbf{E}_{j,t}]$ does not hurt the sub-exponential properties by Exercise 2.7.8 in [Wainwright \(2019\)](#). Then we obtain by Theorem 2.8.1 in [Wainwright \(2019\)](#) that, with probability at least $1 - O(n^{-10})$,

$$\left| \left(\frac{1}{T} \mathbf{E} \mathbf{E}^\top - \mathbb{E} \left[\frac{1}{T} \mathbf{E} \mathbf{E}^\top \right] \right)_{i,j} \right| \lesssim \frac{1}{\sqrt{T}} \sqrt{(\mathbf{\Sigma}_\varepsilon)_{i,i} (\mathbf{\Sigma}_\varepsilon)_{j,j}} \log n.$$

Step 1.2 – Bound for $(\frac{1}{T} \mathbf{E} \mathbf{F} \mathbf{B}^\top + \frac{1}{T} \mathbf{B} \mathbf{F}^\top \mathbf{E}^\top)$.

Next, we have that, conditioning on \mathbf{F} ,

$$\left(\frac{1}{T} \mathbf{E} \mathbf{F} \mathbf{B}^\top \right)_{i,j} = \frac{1}{T} (\mathbf{\Sigma}_\varepsilon^{1/2})_{i,\cdot} \mathbf{Z} \mathbf{F} \mathbf{b}_j = \frac{1}{T} \sum_{t=1}^T \sum_{k=1}^N (\mathbf{\Sigma}_\varepsilon^{1/2})_{i,k} \mathbf{Z}_{k,t} \mathbf{f}_t^\top \mathbf{b}_j$$

is the sum of NT independent zero-mean sub-Gaussian random variables. We obtain by Theorem 2.6.3 in [Wainwright \(2019\)](#) that, with probability at least $1 - O(n^{-10})$,

$$\begin{aligned} \left| \left(\frac{1}{T} \mathbf{E} \mathbf{F} \mathbf{B}^\top \right)_{i,j} \right| &\lesssim \frac{1}{T} \sqrt{\sum_{t=1}^T \sum_{k=1}^N ((\mathbf{\Sigma}_\varepsilon^{1/2})_{i,k})^2 (\mathbf{f}_t^\top \mathbf{b}_j)^2 \log n} = \frac{1}{T} \sqrt{(\mathbf{\Sigma}_\varepsilon)_{i,i} \mathbf{b}_j^\top \mathbf{F}^\top \mathbf{F} \mathbf{b}_j \log n} \\ &= \frac{1}{\sqrt{T}} \sqrt{(\mathbf{\Sigma}_\varepsilon)_{i,i} \mathbf{b}_j^\top \left(\frac{1}{T} \mathbf{F}^\top \mathbf{F} - \mathbf{I}_r + \mathbf{I}_r \right) \mathbf{b}_j \log n} \\ &\lesssim \frac{1}{\sqrt{T}} \sqrt{(\mathbf{\Sigma}_\varepsilon)_{i,i} \mathbf{b}_j^\top \mathbf{I}_r \mathbf{b}_j \log n} = \frac{1}{\sqrt{T}} \|\mathbf{b}_j\|_2 \sqrt{(\mathbf{\Sigma}_\varepsilon)_{i,i} \log n} \\ &\lesssim \frac{1}{\sqrt{T}} \sigma_1 \|\bar{\mathbf{U}}_{j,\cdot}\|_2 \sqrt{(\mathbf{\Sigma}_\varepsilon)_{i,i} \log n}, \end{aligned}$$

where the last line is owing to the fact that $\|\mathbf{b}_i\|_2 \lesssim \sigma_1 \|\bar{\mathbf{U}}_{i,\cdot}\|_2$ for all $1 \leq i \leq N$. Similarly we have $|(\frac{1}{T} \mathbf{B} \mathbf{F}^\top \mathbf{E}^\top)_{i,j}| = |(\frac{1}{T} \mathbf{E} \mathbf{F} \mathbf{B}^\top)_{j,i}| \lesssim \frac{1}{\sqrt{T}} \sigma_1 \|\bar{\mathbf{U}}_{i,\cdot}\|_2 \sqrt{(\mathbf{\Sigma}_\varepsilon)_{j,j} \log n}$.

Step 1.3 – Bound for $(\mathbf{U} \mathbf{\Lambda}^2 \mathbf{U}^\top - \widehat{\mathbf{U}} \widehat{\mathbf{\Sigma}}^2 \widehat{\mathbf{U}})$.

Finally, we have that $(\mathbf{U} \mathbf{\Lambda}^2 \mathbf{U}^\top - \widehat{\mathbf{U}} \widehat{\mathbf{\Sigma}}^2 \widehat{\mathbf{U}})_{i,j} = \tilde{\mathbf{B}}_{i,\cdot} \tilde{\mathbf{B}}_{j,\cdot}^\top - \widehat{\mathbf{B}}_{i,\cdot} \widehat{\mathbf{B}}_{j,\cdot}^\top$. We obtain by Lemma 2 that $\mathbf{B} \mathbf{R}_B = \mathbf{B} (\mathbf{J}^{-1})^\top \mathbf{R}_V^\top = \tilde{\mathbf{B}} \mathbf{R}_V^\top$. So, using the error bounds for $\|(\widehat{\mathbf{B}} - \mathbf{B} \mathbf{R}_B)_{i,\cdot}\|_2$ in Corollary 1, we obtain that, with probability at least $1 - O(n^{-10})$

$$\left| (\mathbf{U} \mathbf{\Lambda}^2 \mathbf{U}^\top - \widehat{\mathbf{U}} \widehat{\mathbf{\Sigma}}^2 \widehat{\mathbf{U}})_{i,j} \right| = \left| (\mathbf{B} \mathbf{R}_B)_{i,\cdot} (\mathbf{B} \mathbf{R}_B)_{j,\cdot}^\top - \widehat{\mathbf{B}}_{i,\cdot} \widehat{\mathbf{B}}_{j,\cdot}^\top \right|$$

$$\begin{aligned}
&\lesssim \|(\mathbf{BR}_B)_{i,\cdot} - \widehat{\mathbf{B}}_{i,\cdot}\|_2 \|(\mathbf{BR}_B)_{j,\cdot}\|_2 + \|(\mathbf{BR}_B)_{j,\cdot} - \widehat{\mathbf{B}}_{j,\cdot}\|_2 \|\widehat{\mathbf{B}}_{i,\cdot}\|_2 \\
&\stackrel{(i)}{\lesssim} \left(\frac{1}{\sqrt{T}} \|(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{i,\cdot}\|_1 + \left(\frac{n}{\theta^2 T} + \frac{1}{\theta\sqrt{T}}\right) \sigma_r \|\bar{\mathbf{U}}_{i,\cdot}\|_2\right) r \log^2 n \|\mathbf{b}_j\|_2 \\
&+ \left(\frac{1}{\sqrt{T}} \|(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{j,\cdot}\|_1 + \left(\frac{n}{\theta^2 T} + \frac{1}{\theta\sqrt{T}}\right) \sigma_r \|\bar{\mathbf{U}}_{j,\cdot}\|_2\right) r \log^2 n \|\mathbf{b}_i\|_2
\end{aligned}$$

where (i) is because $\|\mathbf{R}_B\|_2 \leq \|\mathbf{J}^{-1}\|_2 \|\mathbf{R}_V\|_2 \lesssim 1$ by Lemma 2, and $\|\widehat{\mathbf{B}}_{i,\cdot}\|_2 \leq \|(\mathbf{BR}_B)_{i,\cdot}\|_2 + \|\widehat{\mathbf{B}}_{i,\cdot} - (\mathbf{BR}_B)_{i,\cdot}\|_2 \lesssim \|\mathbf{B}_{i,\cdot}\|_2$ since $\|\widehat{\mathbf{B}}_{i,\cdot} - (\mathbf{BR}_B)_{i,\cdot}\|_2$ is negligible by Corollary 1.

In summary, using a standard union bound argument, we obtain that, with probability at least $1 - O(n^{-8})$, the following bound holds simultaneously for all $1 \leq i, j \leq N$:

$$\begin{aligned}
|(\widehat{\boldsymbol{\Sigma}}_\varepsilon - \boldsymbol{\Sigma}_\varepsilon)_{i,j}| &\lesssim \frac{1}{\sqrt{T}} \sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i} (\boldsymbol{\Sigma}_\varepsilon)_{j,j} \log n} + \frac{1}{\sqrt{T}} \|\mathbf{b}_i\|_2 \sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{j,j} \sqrt{\log n}} + \frac{1}{\sqrt{T}} \|\mathbf{b}_j\|_2 \sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i} \sqrt{\log n}} \\
&+ \left(\frac{1}{\sqrt{T}} (\|(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{i,\cdot}\|_1 \|\mathbf{b}_j\|_2 + \|(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{j,\cdot}\|_1 \|\mathbf{b}_i\|_2) + \left(\frac{n}{\theta^2 T} + \frac{1}{\theta\sqrt{T}}\right) \sigma_r (\|\bar{\mathbf{U}}_{i,\cdot}\|_2 \|\mathbf{b}_j\|_2 + \|\bar{\mathbf{U}}_{j,\cdot}\|_2 \|\mathbf{b}_i\|_2)\right) r \log^2 n.
\end{aligned}$$

The desired bound for $|(\widehat{\boldsymbol{\Sigma}}_\varepsilon - \boldsymbol{\Sigma}_\varepsilon)_{i,j}|$ follows from simplifying the above bound using β_i , γ_i , and $\gamma_{\varepsilon,i}$.

Step 2 – Estimation error for the generalized thresholding estimator $\widehat{\boldsymbol{\Sigma}}_\varepsilon^\tau$.

Since the spectral norm is not larger than the maximum L_1 -norm of rows, we obtain that

$$\|\widehat{\boldsymbol{\Sigma}}_\varepsilon^\tau - \boldsymbol{\Sigma}_\varepsilon\|_2 \leq \max_{1 \leq i \leq N} \sum_{j=1}^N |(\widehat{\boldsymbol{\Sigma}}_\varepsilon^\tau)_{i,j} - (\boldsymbol{\Sigma}_\varepsilon)_{i,j}| = \max_{1 \leq i \leq N} \sum_{j=1}^N |h((\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{i,j}, \tau_{i,j}) - (\boldsymbol{\Sigma}_\varepsilon)_{i,j}|.$$

For the summand $|h((\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{i,j}, \tau_{i,j}) - (\boldsymbol{\Sigma}_\varepsilon)_{i,j}|$, we have that

$$\begin{aligned}
|h((\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{i,j}, \tau_{i,j}) - (\boldsymbol{\Sigma}_\varepsilon)_{i,j}| &= |h((\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{i,j}, \tau_{i,j}) \mathbf{1}_{\{ |(\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{i,j}| \geq \tau_{i,j} \}} - (\boldsymbol{\Sigma}_\varepsilon)_{i,j} \mathbf{1}_{\{ |(\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{i,j}| \geq \tau_{i,j} \}} - (\boldsymbol{\Sigma}_\varepsilon)_{i,j} \mathbf{1}_{\{ |(\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{i,j}| < \tau_{i,j} \}}| \\
&= |h((\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{i,j}, \tau_{i,j}) - (\boldsymbol{\Sigma}_\varepsilon)_{i,j} \mathbf{1}_{\{ |(\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{i,j}| \geq \tau_{i,j} \}} + |(\boldsymbol{\Sigma}_\varepsilon)_{i,j} \mathbf{1}_{\{ |(\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{i,j}| < \tau_{i,j} \}}|.
\end{aligned}$$

Then we derive the upper bounds for the above two terms.

For the first term, without loss of generality, we take for instance the hard thresholding function $h(z, \tau) = z \mathbf{1}_{\{|z| \geq \tau\}}$, and the proof for other thresholding functions follows from the similar ways. We have that

$$\begin{aligned}
|h((\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{i,j}, \tau_{i,j}) - (\boldsymbol{\Sigma}_\varepsilon)_{i,j} \mathbf{1}_{\{ |(\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{i,j}| \geq \tau_{i,j} \}} &= |(\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{i,j} - (\boldsymbol{\Sigma}_\varepsilon)_{i,j} \mathbf{1}_{\{ |(\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{i,j}| \geq \tau_{i,j} \}} \\
&\leq |(\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{i,j} - (\boldsymbol{\Sigma}_\varepsilon)_{i,j}| \lesssim \epsilon_{N,T} \sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i} (\boldsymbol{\Sigma}_\varepsilon)_{j,j}} \\
&\lesssim (\epsilon_{N,T} \sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i} (\boldsymbol{\Sigma}_\varepsilon)_{j,j}})^{1-q} |(\boldsymbol{\Sigma}_\varepsilon)_{i,j}|^q,
\end{aligned}$$

where the last second inequality is because $\epsilon_{N,T} \sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i} (\boldsymbol{\Sigma}_\varepsilon)_{j,j}} \lesssim |(\boldsymbol{\Sigma}_\varepsilon)_{i,j}|$.

Next, for the second term, we have that

$$\begin{aligned}
|(\boldsymbol{\Sigma}_\varepsilon)_{i,j} \mathbf{1}_{\{ |(\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{i,j}| < \tau_{i,j} \}} &= |(\boldsymbol{\Sigma}_\varepsilon)_{i,j}|^q \cdot |(\boldsymbol{\Sigma}_\varepsilon)_{i,j}|^{1-q} \cdot \mathbf{1}_{\{ |(\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{i,j}| < \tau_{i,j} \}} \\
&\lesssim |(\boldsymbol{\Sigma}_\varepsilon)_{i,j}|^q \cdot (|(\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{i,j}| + \epsilon_{N,T} \sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i} (\boldsymbol{\Sigma}_\varepsilon)_{j,j}})^{1-q} \cdot \mathbf{1}_{\{ |(\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{i,j}| < \tau_{i,j} \}} \\
&\leq |(\boldsymbol{\Sigma}_\varepsilon)_{i,j}|^q \cdot (\tau_{i,j} + \epsilon_{N,T} \sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i} (\boldsymbol{\Sigma}_\varepsilon)_{j,j}})^{1-q}.
\end{aligned}$$

For $\tau_{i,j}$, since $\max_{1 \leq i, j \leq N} |(\widehat{\boldsymbol{\Sigma}}_\varepsilon - \boldsymbol{\Sigma}_\varepsilon)_{i,j}| \lesssim \epsilon_{N,T} \sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i} (\boldsymbol{\Sigma}_\varepsilon)_{j,j}}$ and $\epsilon_{N,T} \ll 1$, we obtain that $\max_{1 \leq i \leq N} |(\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{i,i} - (\boldsymbol{\Sigma}_\varepsilon)_{i,i}| \lesssim \epsilon_{N,T} (\boldsymbol{\Sigma}_\varepsilon)_{i,i}$ and thus $(\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{i,i} \asymp (\boldsymbol{\Sigma}_\varepsilon)_{i,i}$, implying $\tau_{i,j} = C \epsilon_{N,T} \sqrt{(\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{i,i} (\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{j,j}} \lesssim \epsilon_{N,T} \sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i} (\boldsymbol{\Sigma}_\varepsilon)_{j,j}}$. So, we obtain that

$$|(\boldsymbol{\Sigma}_\varepsilon)_{i,j} \mathbf{1}_{\{ |(\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{i,j}| < \tau_{i,j} \}} \lesssim |(\boldsymbol{\Sigma}_\varepsilon)_{i,j}|^q (\epsilon_{N,T} \sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i} (\boldsymbol{\Sigma}_\varepsilon)_{j,j}} + \epsilon_{N,T} \sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i} (\boldsymbol{\Sigma}_\varepsilon)_{j,j}})^{1-q}$$

$$\lesssim (\epsilon_{N,T})^{1-q} ((\boldsymbol{\Sigma}_\varepsilon)_{i,i} (\boldsymbol{\Sigma}_\varepsilon)_{j,j})^{(1-q)/2} |(\boldsymbol{\Sigma}_\varepsilon)_{i,j}|^q.$$

Combining the bounds for the above two terms, we conclude that, using the assumption on the sparsity of $\boldsymbol{\Sigma}_\varepsilon$,

$$\begin{aligned} \|\widehat{\boldsymbol{\Sigma}}_\varepsilon^\tau - \boldsymbol{\Sigma}_\varepsilon\|_2 &\leq \max_{1 \leq i \leq N} \sum_{j=1}^N |(\widehat{\boldsymbol{\Sigma}}_\varepsilon^\tau)_{i,j} - (\boldsymbol{\Sigma}_\varepsilon)_{i,j}| \\ &\lesssim (\epsilon_{N,T})^{1-q} \max_{1 \leq i \leq N} \sum_{j=1}^N ((\boldsymbol{\Sigma}_\varepsilon)_{i,i} (\boldsymbol{\Sigma}_\varepsilon)_{j,j})^{(1-q)/2} |(\boldsymbol{\Sigma}_\varepsilon)_{i,j}|^q \\ &\leq (\epsilon_{N,T})^{1-q} s(\boldsymbol{\Sigma}_\varepsilon). \end{aligned}$$

E Proof of Theorem 3: Test for structural breaks in betas

E.1 Some useful lemmas

To prove Theorem 2, we collect some useful lemmas as preparations. Recall that, we already showed in Lemma 8 for some properties of \mathbf{V}_S , for any subset S . Using these properties of \mathbf{V}_S , we are able to establish the following lemma.

Lemma 11. *Suppose that $r + \log n \ll \min(T_1, T_2)$ and the assumptions in Theorem 1 hold. Then we have that, with probability at least $1 - O(n^{-2})$:*

$$(i) \text{ It holds } \sigma_i(\widehat{\mathbf{V}}^j) \asymp \sigma_i(\mathbf{V}^j) \asymp (T_j/T)^{1/2}, \quad \sigma_i((\widehat{\mathbf{V}}^j)^+) \asymp \sigma_i((\mathbf{V}^j)^+) \asymp (T/T_j)^{1/2},$$

$$\|(\widehat{\mathbf{V}}^j)^+ - (\mathbf{V}^j \mathbf{R}_V^\top)^+\|_2 \lesssim (T/T_j)^{1/2} \frac{1}{\theta} \sqrt{r} \log n,$$

and

$$\left\| \prod_{j=1}^2 (\widehat{\mathbf{V}}^j)^\top \widehat{\mathbf{V}}^j - \prod_{j=1}^2 (\mathbf{V}^j \mathbf{R}_V^\top)^\top \mathbf{V}^j \mathbf{R}_V^\top \right\|_2 \lesssim (T_1 T_2 / T^2) \frac{1}{\theta} \sqrt{r} \log n.$$

(ii) Under the null hypothesis $H_0 : \mathbf{b}_i^1 = \mathbf{b}_i^2$, it holds

$$\|\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2\|_2 \lesssim \frac{1}{\theta} \sqrt{r} \log n \|\mathbf{b}_i^j\|_2 + \frac{1}{\theta} \sqrt{r} \log^{3/2} n \|(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{i,\cdot}\|_2 + \sum_{j=1}^2 \frac{1}{\sqrt{T_j}} \|(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{i,\cdot}\|_2 \sqrt{r \log n}$$

and $\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2$ admits the following first-order approximation

$$\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2 = \mathbf{G}_{b_i} + \boldsymbol{\Upsilon}_{b_i}, \quad \text{with} \quad \mathbf{G}_{b_i} := \frac{1}{\sqrt{T}} \left[\sum_{j=1}^2 (-1)^{j-1} (\mathbf{V}^j \mathbf{R}_V^\top)^+ (\mathbf{Z}^j)^\top \right] \boldsymbol{\Sigma}_\varepsilon^{1/2} [\mathbf{I}_N + \mathbf{U} \mathbf{U}^\top]_{\cdot, i},$$

where the term $\boldsymbol{\Upsilon}_{b_i}$ satisfies that

$$\|\boldsymbol{\Upsilon}_{b_i}\|_2 \lesssim \frac{1}{\theta} \sqrt{r} \log^{3/2} n \cdot \sqrt{\varphi_i} + \left(\frac{n}{\theta^2 T} + \frac{1}{\theta \sqrt{T}} \right) \sqrt{r} \log^{3/2} n \cdot \sigma_r \|\mathbf{U}_{i,\cdot}\|_2,$$

with $\varphi_i := [\boldsymbol{\Sigma}_\varepsilon + \mathbf{U} \mathbf{U}^\top \boldsymbol{\Sigma}_\varepsilon + \boldsymbol{\Sigma}_\varepsilon \mathbf{U} \mathbf{U}^\top + \mathbf{U} \mathbf{U}^\top \boldsymbol{\Sigma}_\varepsilon \mathbf{U} \mathbf{U}^\top]_{i,i}$.

Proof. For the first-order approximation of $\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2$, we start by rewriting the regression coefficients $\widehat{\mathbf{b}}_i^j$ as $\widehat{\mathbf{b}}_i^j = T^{-1/2} (\widehat{\mathbf{V}}^j)^+ (\mathbf{X}_{i,\cdot}^j)^\top$, where $(\widehat{\mathbf{V}}^j)^+ = ((\widehat{\mathbf{V}}^j)^\top \widehat{\mathbf{V}}^j)^{-1} (\widehat{\mathbf{V}}^j)^\top$ is the pseudo inverse of $\widehat{\mathbf{F}}^j$. We obtain by our master theorem that $\widehat{\mathbf{V}} - \mathbf{V} \mathbf{R}_V^\top = \mathbf{G}_V \mathbf{R}_V^\top + \boldsymbol{\Psi}_V \mathbf{R}_V^\top$. We partition the matrices as $\widehat{\mathbf{V}} = ((\widehat{\mathbf{V}}^1)^\top, (\widehat{\mathbf{V}}^2)^\top)^\top$, $\mathbf{V} = ((\mathbf{V}^1)^\top, (\mathbf{V}^2)^\top)^\top$, $\mathbf{E} = (\mathbf{E}^1, \mathbf{E}^2) = (\boldsymbol{\Sigma}_\varepsilon^{1/2} \mathbf{Z}^1, \boldsymbol{\Sigma}_\varepsilon^{1/2} \mathbf{Z}^2)$, where $\mathbf{V}^j, \widehat{\mathbf{V}}^j \in \mathbb{R}^{T_j \times r}$, $\mathbf{E}^j, \mathbf{Z}^j \in \mathbb{R}^{N \times T_j}$, and denote $\mathbf{D}^j := \widehat{\mathbf{V}}^j - \mathbf{V}^j \mathbf{R}_V^\top$.

Proof of (i). Similar to Step 1 in proof of Lemma 11, we have that

$$\|\widehat{\mathbf{V}}^j - \mathbf{V}^j \mathbf{R}_V^\top\|_2 \lesssim \sqrt{T_j} \rho \sqrt{\frac{r}{n}} \log n = \sqrt{T_j/T} \frac{1}{\theta} \sqrt{r} \log n.$$

Then using the above bounds we can prove that, with probability at least $1 - O(n^{-2})$, the ranks of $\widehat{\mathbf{V}}^j$ and \mathbf{V}^j are both equal to r . Similar to the proof of Lemma 8, we have that $\sigma_i(\widehat{\mathbf{V}}^j) \asymp \sigma_i(\mathbf{V}^j) \asymp (T_j/T)^{1/2}$. We obtain by Theorem 3.4 in Stewart (1977) that

$$\begin{aligned} \|(\widehat{\mathbf{V}}^j)^+ - (\mathbf{V}^j \mathbf{R}_V^\top)^+\|_2 &\lesssim \|(\widehat{\mathbf{V}}^j)^+\|_2 \|(\mathbf{V}^j \mathbf{R}_V^\top)^+\|_2 \|\widehat{\mathbf{V}}^j - \mathbf{V}^j \mathbf{R}_V^\top\|_2 \\ &\lesssim (\|(\widehat{\mathbf{V}}^j)^+ - (\mathbf{V}^j \mathbf{R}_V^\top)^+\|_2 + \|(\mathbf{V}^j \mathbf{R}_V^\top)^+\|_2) \|(\mathbf{V}^j \mathbf{R}_V^\top)^+\|_2 \|\widehat{\mathbf{V}}^j - \mathbf{V}^j \mathbf{R}_V^\top\|_2 \\ &\lesssim \|(\widehat{\mathbf{V}}^j)^+ - (\mathbf{V}^j \mathbf{R}_V^\top)^+\|_2 \times \|(\mathbf{V}^j \mathbf{R}_V^\top)^+\|_2 \|\widehat{\mathbf{V}}^j - \mathbf{V}^j \mathbf{R}_V^\top\|_2 \\ &\quad + \|(\mathbf{V}^j \mathbf{R}_V^\top)^+\|_2^2 \|\widehat{\mathbf{V}}^j - \mathbf{V}^j \mathbf{R}_V^\top\|_2. \end{aligned}$$

Note that the coefficient for $\|(\widehat{\mathbf{V}}^j)^+ - (\mathbf{V}^j \mathbf{R}_V^\top)^+\|_2$ in the right hand side of the above inequality satisfies that $\|(\mathbf{V}^j \mathbf{R}_V^\top)^+\|_2 \|\widehat{\mathbf{V}}^j - \mathbf{V}^j \mathbf{R}_V^\top\|_2 \lesssim (T/T_j)^{1/2} \sqrt{T_j} \rho \sqrt{\frac{r}{n}} \log n \ll 1$. Thus, by the self-bounding method, we obtain that

$$\|(\widehat{\mathbf{V}}^j)^+ - (\mathbf{V}^j \mathbf{R}_V^\top)^+\|_2 \lesssim \|(\mathbf{V}^j \mathbf{R}_V^\top)^+\|_2^2 \|\widehat{\mathbf{V}}^j - \mathbf{V}^j \mathbf{R}_V^\top\|_2 \lesssim \frac{T}{T_j} \sqrt{T_j} \rho \sqrt{\frac{r}{n}} \log n \lesssim (T/T_j)^{1/2} \frac{1}{\theta} \sqrt{r} \log n.$$

Then we obtain that $\sigma_i((\widehat{\mathbf{V}}^j)^+) \asymp \sigma_i((\mathbf{V}^j)^+) \asymp (T/T_j)^{1/2}$.

We have that

$$\begin{aligned} \|(\widehat{\mathbf{V}}^j)^\top \widehat{\mathbf{V}}^j - (\mathbf{V}^j \mathbf{R}_V^\top)^\top \mathbf{V}^j \mathbf{R}_V^\top\|_2 &\lesssim (\|\widehat{\mathbf{V}}^j\|_2 + \|\mathbf{V}^j\|_2) \|\widehat{\mathbf{V}}^j - \mathbf{V}^j \mathbf{R}_V^\top\|_2 \\ &\lesssim (T_j/T)^{1/2} \sqrt{T_j} \rho \sqrt{\frac{r}{n}} \log n \lesssim T_j \rho \sqrt{\frac{r}{nT}} \log n = (T_j/T) \frac{1}{\theta} \sqrt{r} \log n. \end{aligned}$$

Then, we obtain that

$$\begin{aligned} \left\| \prod_{j=1}^2 (\widehat{\mathbf{V}}^j)^\top \widehat{\mathbf{V}}^j - \prod_{j=1}^2 (\mathbf{V}^j \mathbf{R}_V^\top)^\top \mathbf{V}^j \mathbf{R}_V^\top \right\|_2 &\lesssim ((T_1/T)^{1/2})^2 T_2 \rho \sqrt{\frac{r}{nT}} \log n + ((T_2/T)^{1/2})^2 T_1 \rho \sqrt{\frac{r}{nT}} \log n \\ &\lesssim \frac{1}{T} T_1 T_2 \rho \sqrt{\frac{r}{nT}} \log n \lesssim (T_1 T_2 / T^2) \frac{1}{\theta} \sqrt{r} \log n. \end{aligned}$$

Proof of (ii). For the upper bound of $\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2$, we have that

$$\begin{aligned} \widehat{\mathbf{b}}_i^j &= T^{-1/2} (\widehat{\mathbf{V}}^j)^+ (\mathbf{X}_{i,\cdot}^j)^\top = T^{-1/2} (\widehat{\mathbf{V}}^j)^+ (\mathbf{F}^j \mathbf{b}_i^j + (\mathbf{E}_{i,\cdot}^j)^\top) \\ &= (\widehat{\mathbf{V}}^j)^+ \mathbf{V}^j \mathbf{R}_V^\top (\mathbf{R}_V \mathbf{J}^{-1} \mathbf{b}_i^j) + T^{-1/2} (\widehat{\mathbf{V}}^j)^+ (\mathbf{E}_{i,\cdot}^j)^\top \\ &= [\mathbf{I}_r - (\widehat{\mathbf{V}}^j)^+ (\widehat{\mathbf{V}}^j - \mathbf{V}^j \mathbf{R}_V^\top)] (\mathbf{R}_V \mathbf{J}^{-1} \mathbf{b}_i^j) + T^{-1/2} (\widehat{\mathbf{V}}^j)^+ (\mathbf{E}_{i,\cdot}^j)^\top. \end{aligned}$$

So we obtain that, under the null hypothesis that $\mathbf{b}_i^1 = \mathbf{b}_i^2$, it holds

$$\begin{aligned} \widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2 &= \sum_{j=1}^2 (-1)^j [(\widehat{\mathbf{V}}^j)^+ (\widehat{\mathbf{V}}^j - \mathbf{V}^j \mathbf{R}_V^\top) (\mathbf{R}_V \mathbf{J}^{-1} \mathbf{b}_i^j) - T^{-1/2} (\widehat{\mathbf{V}}^j)^+ (\mathbf{E}_{i,\cdot}^j)^\top] \\ &= \sum_{j=1}^2 (-1)^j [(\widehat{\mathbf{V}}^j)^+ (\widehat{\mathbf{V}}^j - \mathbf{V}^j \mathbf{R}_V^\top) (\mathbf{R}_V \mathbf{J}^{-1} \mathbf{b}_i^j)] \\ &\quad + \sum_{j=1}^2 (-1)^{j-1} T^{-1/2} \{[(\widehat{\mathbf{V}}^j)^+ - (\mathbf{V}^j \mathbf{R}_V^\top)^+] (\mathbf{Z}^j)^\top (\boldsymbol{\Sigma}_\varepsilon^{1/2})_{\cdot,i} + (\mathbf{V}^j \mathbf{R}_V^\top)^+ (\mathbf{Z}^j)^\top (\boldsymbol{\Sigma}_\varepsilon^{1/2})_{\cdot,i}\}. \end{aligned}$$

Then we obtain by (G.3) in Lemma 19 of [Yan and Wainwright \(2024\)](#) that

$$\begin{aligned} \|\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2\|_2 &\lesssim \sum_{j=1}^2 (T/T_j)^{1/2} \sqrt{T_j/T} \frac{1}{\theta} \sqrt{r} \log n \|\mathbf{b}_i^j\|_2 \\ &+ \sum_{j=1}^2 T^{-1/2} \cdot (T/T_j)^{1/2} \frac{1}{\theta} \sqrt{r} \log n \cdot \|(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{i,\cdot}\|_2 \sqrt{T_j \log n} + T^{-1/2} \cdot \sqrt{T/T_j} \|(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{i,\cdot}\|_2 \sqrt{r \log n} \\ &\lesssim \frac{1}{\theta} \sqrt{r} \log n \|\mathbf{b}_i^j\|_2 + \frac{1}{\theta} \sqrt{r} \log^{3/2} n \|(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{i,\cdot}\|_2 + \sum_{j=1}^2 (1/T_j)^{1/2} \|(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{i,\cdot}\|_2 \sqrt{r \log n}. \end{aligned}$$

Then, we establish the first-order approximation for $\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2$.

Step 1 – compute the first-order approximation.

We obtain by Theorem 3.2 in [Stewart \(1977\)](#) that

$$(\widehat{\mathbf{V}}^j)^+ - (\mathbf{V}^j \mathbf{R}_V^\top)^+ = (\widehat{\mathbf{V}}^j)^+ \mathbf{D}^j (\mathbf{V}^j \mathbf{R}_V^\top)^+ + ((\widehat{\mathbf{V}}^j)^\top \widehat{\mathbf{V}}^j)^{-1} (\mathbf{D}^j)^\top (\mathbf{I}_{T_j} - \mathbf{P}_{V^j}),$$

for $j = 1, 2$. Then, we obtain that

$$\begin{aligned} \widehat{\mathbf{b}}_i^j - T^{-1/2} (\mathbf{V}^j \mathbf{R}_V^\top)^+ (\mathbf{X}_{i,\cdot}^j)^\top &= T^{-1/2} ((\widehat{\mathbf{V}}^j)^+ - (\mathbf{V}^j \mathbf{R}_V^\top)^+) (\mathbf{X}_{i,\cdot}^j)^\top \\ &= T^{-1/2} ((\widehat{\mathbf{V}}^j)^+ - (\mathbf{V}^j \mathbf{R}_V^\top)^+) \mathbf{F}^j \mathbf{b}_i^j + T^{-1/2} ((\widehat{\mathbf{V}}^j)^+ - (\mathbf{V}^j \mathbf{R}_V^\top)^+) (\mathbf{E}_{i,\cdot}^j)^\top \\ &= T^{-1/2} (\widehat{\mathbf{V}}^j)^+ \mathbf{D}^j (\mathbf{V}^j \mathbf{R}_V^\top)^+ \mathbf{F}^j \mathbf{b}_i^j + T^{-1/2} ((\widehat{\mathbf{V}}^j)^+ - (\mathbf{V}^j \mathbf{R}_V^\top)^+) (\mathbf{E}_{i,\cdot}^j)^\top \\ &= (\widehat{\mathbf{V}}^j)^+ \mathbf{D}^j \mathbf{R}_V \mathbf{J}^{-1} \mathbf{b}_i^j + T^{-1/2} ((\widehat{\mathbf{V}}^j)^+ - (\mathbf{V}^j \mathbf{R}_V^\top)^+) (\mathbf{E}_{i,\cdot}^j)^\top. \end{aligned}$$

Note that the term in the left hand side can be decomposed into

$$T^{-1/2} (\mathbf{V}^j \mathbf{R}_V^\top)^+ (\mathbf{X}_{i,\cdot}^j)^\top = T^{-1/2} (\mathbf{V}^j \mathbf{R}_V^\top)^+ (\mathbf{F}^j \mathbf{b}_i^j + (\mathbf{E}_{i,\cdot}^j)^\top) = \mathbf{R}_V \mathbf{J}^{-1} \mathbf{b}_i^j + T^{-1/2} (\mathbf{V}^j \mathbf{R}_V^\top)^+ (\mathbf{E}_{i,\cdot}^j)^\top,$$

and the first term in the right hand side can be decomposed into

$$\begin{aligned} (\widehat{\mathbf{V}}^j)^+ \mathbf{D}^j \mathbf{R}_V \mathbf{J}^{-1} \mathbf{b}_i^j &= (\widehat{\mathbf{V}}^j)^+ \mathbf{D}^j \mathbf{R}_V \mathbf{J}^{-1} \mathbf{b}_i^j = (\widehat{\mathbf{V}}^j)^+ (\mathbf{G}_V^j \mathbf{R}_V^\top + \boldsymbol{\Psi}_V^j \mathbf{R}_V^\top) \mathbf{R}_V \mathbf{J}^{-1} \mathbf{b}_i^j \\ &= (\widehat{\mathbf{V}}^j)^+ \mathbf{G}_V^j \mathbf{J}^{-1} \mathbf{b}_i^j + (\widehat{\mathbf{V}}^j)^+ \boldsymbol{\Psi}_V^j \mathbf{J}^{-1} \mathbf{b}_i^j \\ &= (\mathbf{V}^j \mathbf{R}_V^\top)^+ \mathbf{G}_V^j \mathbf{J}^{-1} \mathbf{b}_i^j + [((\widehat{\mathbf{V}}^j)^+ - (\mathbf{V}^j \mathbf{R}_V^\top)^+) \mathbf{G}_V^j + (\widehat{\mathbf{V}}^j)^+ \boldsymbol{\Psi}_V^j] \mathbf{J}^{-1} \mathbf{b}_i^j. \end{aligned}$$

So, we obtain by $\mathbf{G}_V^j \mathbf{J}^{-1} \mathbf{b}_i^j = T^{-1/2} (\mathbf{E}^j)^\top \mathbf{U} (\mathbf{U}_{i,\cdot})^\top$ that

$$\begin{aligned} \widehat{\mathbf{b}}_i^j - \mathbf{R}_V \mathbf{J}^{-1} \mathbf{b}_i^j &= T^{-1/2} (\mathbf{V}^j \mathbf{R}_V^\top)^+ (\mathbf{E}_{i,\cdot}^j)^\top + (\mathbf{V}^j \mathbf{R}_V^\top)^+ \mathbf{G}_V^j \mathbf{J}^{-1} \mathbf{b}_i^j + \mathbf{r}_0^j \\ &= T^{-1/2} (\mathbf{V}^j \mathbf{R}_V^\top)^+ [(\mathbf{E}_{i,\cdot}^j)^\top + (\mathbf{E}^j)^\top \mathbf{U} (\mathbf{U}_{i,\cdot})^\top] + \mathbf{r}_0^j \\ &= T^{-1/2} (\mathbf{V}^j \mathbf{R}_V^\top)^+ (\mathbf{E}^j)^\top [\mathbf{I}_N + \mathbf{U} \mathbf{U}^\top]_{\cdot,i} + \mathbf{r}_0^j, \end{aligned}$$

where the remainder term \mathbf{r}_0^j is given by

$$\begin{aligned} \mathbf{r}_0^j &= T^{-1/2} ((\widehat{\mathbf{V}}^j)^+ - (\mathbf{V}^j \mathbf{R}_V^\top)^+) (\mathbf{E}_{i,\cdot}^j)^\top + [((\widehat{\mathbf{V}}^j)^+ - (\mathbf{V}^j \mathbf{R}_V^\top)^+) \mathbf{G}_V^j + (\widehat{\mathbf{V}}^j)^+ \boldsymbol{\Psi}_V^j] \mathbf{J}^{-1} \mathbf{b}_i^j \\ &= T^{-1/2} ((\widehat{\mathbf{V}}^j)^+ - (\mathbf{V}^j \mathbf{R}_V^\top)^+) (\mathbf{E}^j)^\top [\mathbf{I}_N + \mathbf{U} \mathbf{U}^\top]_{\cdot,i} + (\widehat{\mathbf{V}}^j)^+ \boldsymbol{\Psi}_V^j \mathbf{J}^{-1} \mathbf{b}_i^j. \end{aligned}$$

Under the null hypothesis $H_0 : \mathbf{b}_i^1 = \mathbf{b}_i^2$, we have that $\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2 = \mathbf{G}_{b_i} + \boldsymbol{\Upsilon}_{b_i}$ where the first-order term \mathbf{G}_{b_i} is given by

$$\begin{aligned} \mathbf{G}_{b_i} &= T^{-1/2} [(\mathbf{V}^1 \mathbf{R}_V^\top)^+ (\mathbf{E}^1)^\top - (\mathbf{V}^2 \mathbf{R}_V^\top)^+ (\mathbf{E}^2)^\top] [\mathbf{I}_N + \mathbf{U} \mathbf{U}^\top]_{\cdot,i} \\ &= T^{-1/2} [(\mathbf{V}^1 \mathbf{R}_V^\top)^+ (\mathbf{Z}^1)^\top - (\mathbf{V}^2 \mathbf{R}_V^\top)^+ (\mathbf{Z}^2)^\top] \boldsymbol{\Sigma}_\varepsilon^{1/2} [\mathbf{I}_N + \mathbf{U} \mathbf{U}^\top]_{\cdot,i}, \end{aligned}$$

and the remainder term $\boldsymbol{\Upsilon}_{b_i}$ is given by $\boldsymbol{\Upsilon}_{b_i} = \mathbf{r}_0^1 - \mathbf{r}_0^2$.

Step 2 – derive the upper bound for Υ_{b_i} .

Then we have that, by the expression of Ψ_V^j and (G.3) in Lemma 19 of Yan and Wainwright (2024)

$$\begin{aligned}
\|\mathbf{r}_0^j\|_2 &\lesssim T^{-1/2} \|(\widehat{\mathbf{V}}^j)^+ - (\mathbf{V}^j \mathbf{R}_V^\top)^+\|_2 \|(\mathbf{Z}^j)^\top \Sigma_\varepsilon^{1/2} [\mathbf{I}_N + \mathbf{U}\mathbf{U}^\top]_{\cdot,i}\|_2 \\
&\quad + \|(\widehat{\mathbf{V}}^j)^+\|_2 \cdot \sqrt{T_j} \left(\frac{\sqrt{n}}{\theta^2 T} \sqrt{r} \log^{3/2} n + \left(\frac{n}{\theta^2 T} + \frac{1}{\theta \sqrt{T}} \right) \sqrt{r} \log n \sqrt{\frac{\log n}{T}} \right) \cdot \sigma_r \|\mathbf{U}_{i,\cdot}\|_2 \\
&\lesssim T^{-1/2} \cdot (T/T_j)^{1/2} \frac{1}{\theta} \sqrt{r} \log n \cdot \sqrt{T_j \log n} \sqrt{\varphi_i} + \sqrt{T} \left(\frac{\sqrt{n}}{\theta^2 T} + \left(\frac{n}{\theta^2 T} + \frac{1}{\theta \sqrt{T}} \right) \frac{1}{\sqrt{T}} \right) \sqrt{r} \log^{3/2} n \cdot \sigma_r \|\mathbf{U}_{i,\cdot}\|_2 \\
&\lesssim \frac{1}{\theta} \sqrt{r} \log^{3/2} n \cdot \sqrt{\varphi_i} + \left(\frac{n}{\theta^2 T} + \frac{1}{\theta \sqrt{T}} \right) \sqrt{r} \log^{3/2} n \cdot \sigma_r \|\mathbf{U}_{i,\cdot}\|_2.
\end{aligned}$$

Then, the upper bound for $\|\Upsilon_{b_i}\|_2$ follows from $\|\Upsilon_{b_i}\|_2 \leq \|\mathbf{r}_0^1\|_2 + \|\mathbf{r}_0^2\|_2$. \square

Then, we establish Gaussian approximation for the first-order term. For simplicity of notations, we denote $\chi^2(n)$ as a Chi-square random variable with degree of freedom equal to n , and denote by $\chi_\phi^2(n)$ its ϕ -quantile.

Lemma 12. *Suppose that the assumptions in Lemma 11 hold. Then, under the null hypothesis $H_0 : \mathbf{b}_i^1 = \mathbf{b}_i^2$, we have that, for any random variable ζ satisfying $|\zeta| \ll r$, it holds*

$$|\mathbb{P}(\mathcal{B}_i + \zeta \leq \chi_{1-\alpha}^2(r)) - (1-\alpha)| \lesssim |\tau| + r^{-1/2} |\zeta| + s_3, \quad (\text{E.1})$$

where $\varphi_i = [\Sigma_\varepsilon + \mathbf{U}\mathbf{U}^\top \Sigma_\varepsilon + \Sigma_\varepsilon \mathbf{U}\mathbf{U}^\top + \mathbf{U}\mathbf{U}^\top \Sigma_\varepsilon \mathbf{U}\mathbf{U}^\top]_{i,i}$ as defined in Lemma 11, \mathcal{B}_i is defined by

$$\mathcal{B}_i := \frac{T}{\varphi_i} (\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2)^\top \left[\prod_{j=1}^2 (\mathbf{V}^j \mathbf{R}_V^\top)^\top \mathbf{V}^j \mathbf{R}_V^\top \right] (\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2),$$

and $|\tau|$ satisfies that

$$|\tau| \lesssim \left[\frac{1}{\theta} + \sqrt{\kappa_\varepsilon} \|\mathbf{U}_{i,\cdot}\|_2 \left(\frac{1}{\theta} \frac{n}{T} + \frac{1}{\sqrt{T}} \right) \right] \sqrt{T_1 T_2 / T} \sqrt{r} \log^{3/2} n.$$

Here, $s_3 \geq 0$ is a constant bounded by

$$s_3 \lesssim \frac{r^{3/2} \max_{1 \leq k \leq N} \left| [\Sigma_\varepsilon^{1/2} (\mathbf{I}_N + \mathbf{U}\mathbf{U}^\top)]_{k,i} \right|}{\|[\Sigma_\varepsilon^{1/2} (\mathbf{I}_N + \mathbf{U}\mathbf{U}^\top)]_{\cdot,i}\|_2}.$$

Further, if all the entries of the matrix \mathbf{Z} are Gaussian, i.e., the noise is Gaussian, then s_3 in the inequality (E.1) is equal to zero, i.e., the inequality (E.1) holds when $s_3 = 0$.

Proof. Our starting point is the following inequality

$$\mathbb{P}(\mathcal{B}_i \leq \chi_{1-\alpha}^2(r) - |\zeta|) \leq \mathbb{P}(\mathcal{B}_i + \zeta \leq \chi_{1-\alpha}^2(r)) \leq \mathbb{P}(\mathcal{B}_i \leq \chi_{1-\alpha}^2(r) + |\zeta|).$$

Then, we will prove both the upper bound and the lower bound are close to $(1-\alpha)$. We will use three steps.

Step 1 – Bounding $\mathbb{P}(\mathcal{B}_i + \zeta \leq \chi_{1-\alpha}^2(r))$.

We already obtained the first-order approximation of in Lemma 11. Indeed, the first-order term \mathbf{G}_P can be rewritten as

$$\mathbf{G}_{b_i} = T^{-1/2} ((\mathbf{V}^1 \mathbf{R}_V^\top)^+ (\mathbf{Z}^1)^\top - (\mathbf{V}^2 \mathbf{R}_V^\top)^+ (\mathbf{Z}^2)^\top) \Sigma_\varepsilon^{1/2} (\mathbf{I}_N + \mathbf{U}\mathbf{U}^\top)_{\cdot,i}.$$

Denote $\mathbf{K} \in \mathbb{R}^{r \times r}$ the matrix such that $\prod_{j=1}^2 (\mathbf{V}^j \mathbf{R}_V^\top)^\top \mathbf{V}^j \mathbf{R}_V^\top = \mathbf{K}^\top \mathbf{K}$. Then \mathcal{B}_i can be rewritten as

$$\mathcal{B}_i = \frac{T}{\varphi_i} (\mathbf{G}_{b_i} + \Upsilon_{b_i})^\top \mathbf{K}^\top \mathbf{K} (\mathbf{G}_{b_i} + \Upsilon_{b_i}) = \|(T/\varphi_i)^{1/2} \mathbf{K} (\mathbf{G}_{b_i} + \Upsilon_{b_i})\|_2^2.$$

We denote

$$\tau := \sqrt{\mathcal{B}_i} - \left\| \sqrt{\frac{T}{\varphi_i}} \mathbf{K} \mathbf{G}_{b_i} \right\|_2 = \left\| \sqrt{\frac{T}{\varphi_i}} \mathbf{K} (\mathbf{G}_{b_i} + \Upsilon_{b_i}) \right\|_2 - \left\| \sqrt{\frac{T}{\varphi_i}} \mathbf{K} \mathbf{G}_{b_i} \right\|_2.$$

Then we get

$$\mathbb{P}(\mathcal{B}_i \leq \chi_{1-\alpha}^2(r) + |\zeta|) = \mathbb{P}\left(\left\| \sqrt{\frac{T}{\varphi_i}} \mathbf{K} \mathbf{G}_{b_i} \right\|_2 + \tau \leq \sqrt{\chi_{1-\alpha}^2(r) + |\zeta|}\right) \leq \mathbb{P}\left(\left\| \sqrt{\frac{T}{\varphi_i}} \mathbf{K} \mathbf{G}_{b_i} \right\|_2 \leq \sqrt{\chi_{1-\alpha}^2(r) + |\zeta|} + |\tau|\right),$$

and thus

$$\mathbb{P}\left(\left\| \sqrt{\frac{T}{\varphi_i}} \mathbf{K} \mathbf{G}_{b_i} \right\|_2 \leq \sqrt{\chi_{1-\alpha}^2(r) - |\zeta|} - |\tau|\right) \leq \mathbb{P}(\mathcal{B}_i + \zeta \leq \chi_{1-\alpha}^2(r)) \leq \mathbb{P}\left(\left\| \sqrt{\frac{T}{\varphi_i}} \mathbf{K} \mathbf{G}_{b_i} \right\|_2 \leq \sqrt{\chi_{1-\alpha}^2(r) + |\zeta|} + |\tau|\right).$$

Step 2 – Expressing $\mathbf{K} \mathbf{G}_{b_i}$ as a sum to show its proximity to a Gaussian vector.

We define

$$s_3 := \sup_{R \geq 0} \left| \mathbb{P}\left(\left\| \sqrt{\frac{T}{\varphi_i}} \mathbf{K} \mathbf{G}_{b_i} \right\|_2 \leq R\right) - \mathbb{P}(\chi^2(r) \leq R^2) \right|.$$

Conditioning on \mathbf{F} , the random vector \mathbf{G}_{b_i} can be written as a sum of N independent and mean zero $|S|$ -dimensional vectors as follows:

$$\mathbf{G}_{b_i} = \sum_{k=1}^N \mathbf{g}_k \quad \text{with} \quad \mathbf{g}_k = T^{-1/2} \left(\sum_{j=1}^2 (-1)^{j-1} (\mathbf{V}^j \mathbf{R}_V^\top)^\top + (\mathbf{Z}_{k,\cdot}^j)^\top \right) [\boldsymbol{\Sigma}_\varepsilon^{1/2} (\mathbf{I}_N + \mathbf{U} \mathbf{U}^\top)]_{k,i}.$$

By calculations, we have that $\mathbf{K}^\top \mathbf{K} = \prod_{j=1}^2 (\mathbf{V}^j \mathbf{R}_V^\top)^\top \mathbf{V}^j \mathbf{R}_V^\top = [\sum_{j=1}^2 ((\mathbf{V}^j \mathbf{R}_V^\top)^\top \mathbf{V}^j \mathbf{R}_V^\top)^{-1}]^{-1}$, the covariance matrix is given by $\text{cov}(\mathbf{G}_{b_i} | \mathbf{F}) = \sum_{k=1}^N \text{cov}(\mathbf{g}_k | \mathbf{F}) = (\varphi_i/T) \sum_{j=1}^2 ((\mathbf{V}^j \mathbf{R}_V^\top)^\top \mathbf{V}^j \mathbf{R}_V^\top)^{-1} = (\varphi_i/T) (\mathbf{K}^\top \mathbf{K})^{-1} = (\varphi_i/T) \mathbf{K}^{-1} (\mathbf{K}^\top)^{-1}$, and thus

$$\text{cov}((T/\varphi_i)^{1/2} \mathbf{K} \mathbf{G}_{b_i} | \mathbf{F}) = (T/\varphi_i) \mathbf{K} \text{cov}(\mathbf{G}_{b_i} | \mathbf{F}) \mathbf{K}^\top = \mathbf{K} \mathbf{K}^{-1} (\mathbf{K}^\top)^{-1} \mathbf{K}^\top = \mathbf{I}_r.$$

Then we consider two cases.

- (i) When all entries of \mathbf{Z} are Gaussian, we have that $s_3 = 0$.
- (ii) When the entries of \mathbf{Z} are sub-Gaussian, we obtain by Lemma 16 that

$$\begin{aligned} s_3 &\lesssim \sum_{k=1}^N \mathbb{E} \left[\|(T/\varphi_i)^{1/2} \mathbf{K} \mathbf{g}_k\|_2^3 | \mathbf{F} \right] \leq (1/\varphi_i)^{3/2} \mathbb{E} \left[\left\| \mathbf{K} [(\mathbf{V}^1 \mathbf{R}_V^\top)^\top + (\mathbf{Z}_{k,\cdot}^1)^\top - (\mathbf{V}^2 \mathbf{R}_V^\top)^\top + (\mathbf{Z}_{k,\cdot}^2)^\top] \right\|_2^3 | \mathbf{F} \right] \\ &\quad \times \max_{1 \leq k \leq N} \left| [\boldsymbol{\Sigma}_\varepsilon^{1/2} (\mathbf{I}_N + \mathbf{U} \mathbf{U}^\top)]_{k,i} \right| \cdot \sum_{k=1}^N ([\boldsymbol{\Sigma}_\varepsilon^{1/2} (\mathbf{I}_N + \mathbf{U} \mathbf{U}^\top)]_{k,i})^2 \\ &\lesssim \varphi_i^{-1/2} \max_{1 \leq k \leq N} \left| [\boldsymbol{\Sigma}_\varepsilon^{1/2} (\mathbf{I}_N + \mathbf{U} \mathbf{U}^\top)]_{k,i} \right| \mathbb{E} \left[\left\| \mathbf{K} [(\mathbf{V}^1 \mathbf{R}_V^\top)^\top + (\mathbf{Z}_{k,\cdot}^1)^\top - (\mathbf{V}^2 \mathbf{R}_V^\top)^\top + (\mathbf{Z}_{k,\cdot}^2)^\top] \right\|_2^3 | \mathbf{F} \right]. \end{aligned}$$

Recall that $\prod_{j=1}^2 (\mathbf{V}^j \mathbf{R}_V^\top)^\top \mathbf{V}^j \mathbf{R}_V^\top = \mathbf{K}^\top \mathbf{K}$, so we obtain that $\|\mathbf{K}\|_2 \lesssim \prod_{j=1}^2 (T_j/T)^{1/2} \leq \sqrt{T_1 T_2}/T$. since $\mathbf{K} [(\mathbf{V}^1 \mathbf{R}_V^\top)^\top + (\mathbf{Z}_{k,\cdot}^1)^\top - (\mathbf{V}^2 \mathbf{R}_V^\top)^\top + (\mathbf{Z}_{k,\cdot}^2)^\top]$ is a r -dimensional sub-Gaussian vector whose covariance matrix is equal to $\mathbf{K} [\sum_{j=1}^2 ((\mathbf{V}^j \mathbf{R}_V^\top)^\top \mathbf{V}^j \mathbf{R}_V^\top)^{-1}] \mathbf{K}^\top = \mathbf{K} (\mathbf{K}^\top \mathbf{K})^{-1} \mathbf{K}^\top = \mathbf{I}_r$. Then we have that, similar to Step 2 in the proof of Lemma 10, it holds $\mathbb{E} \left[\left\| \mathbf{K} [(\mathbf{V}^1 \mathbf{R}_V^\top)^\top + (\mathbf{Z}_{k,\cdot}^1)^\top - (\mathbf{V}^2 \mathbf{R}_V^\top)^\top + (\mathbf{Z}_{k,\cdot}^2)^\top] \right\|_2^3 | \mathbf{F} \right] \lesssim r^{3/2}$. So we obtain

$$s_3 \lesssim r^{3/2} \varphi_i^{-1/2} \max_{1 \leq k \leq N} \left| [\boldsymbol{\Sigma}_\varepsilon^{1/2} (\mathbf{I}_N + \mathbf{U} \mathbf{U}^\top)]_{k,i} \right|.$$

Step 3 – Establishing chi-squared distributional characterization.

By definition of s_3 , we obtain that

$$\mathbb{P}\left(\sqrt{\chi^2(r)} \leq \sqrt{\chi_{1-\alpha}^2(r) - |\zeta| - |\tau|}\right) - s_3 \leq \mathbb{P}(\mathcal{B}_i + \zeta \leq \chi_{1-\alpha}^2(r)) \leq \mathbb{P}\left(\sqrt{\chi^2(r)} \leq \sqrt{\chi_{1-\alpha}^2(r) + |\zeta| + |\tau|}\right) + s_3.$$

Similar to Step 3 in the proof of Lemma 10, we obtain that

$$|\mathbb{P}(\mathcal{B}_i + \zeta \leq \chi_{1-\alpha}^2(r)) - (1 - \alpha)| \lesssim |\tau| + r^{-1/2} |\zeta| + s_3.$$

So, to prove the desired result in the lemma, it suffices to establish the upper bound for $|\tau|$, which is defined in Step 1. Using (D.7) and the triangle inequality, we obtain that

$$|\tau| = \left| \left\| \sqrt{\frac{T}{\varphi_i}} \mathbf{K}(\mathbf{G}_{b_i} + \mathbf{Y}_{b_i}) \right\|_2 - \left\| \sqrt{\frac{T}{\varphi_i}} \mathbf{K} \mathbf{G}_{b_i} \right\|_2 \right| \leq \left\| \sqrt{\frac{T}{\varphi_i}} \mathbf{K} \mathbf{Y}_{b_i} \right\|_2.$$

Finally, using the upper bound of $\|\mathbf{Y}_{b_i}\|_2$, we obtain the upper bound for $|\tau|$ as

$$\begin{aligned} |\tau| &\lesssim (T/\varphi_i)^{1/2} \cdot \frac{1}{T} \sqrt{T_1 T_2} \cdot \|\mathbf{Y}_{b_i}\|_2 = (1/\varphi_i)^{1/2} \sqrt{T_1 T_2 / T} \cdot \|\mathbf{Y}_{b_i}\|_2 \\ &\lesssim \frac{1}{\theta} \sqrt{T_1 T_2 / T} \sqrt{r} \log^{3/2} n + \sqrt{\kappa_\varepsilon} \|\mathbf{U}_{i,\cdot}\|_2 \left(\frac{n}{\theta T} + \frac{1}{\sqrt{T}} \right) \sqrt{T_1 T_2 / T} \sqrt{r} \log^{3/2} n. \end{aligned}$$

□

E.2 Proof of Theorem 3

To prove the desired results, we use Lemma 12 with the small perturbation $\zeta = \widehat{\mathcal{B}}_i - \mathcal{B}_i$. Then, the problem boils down to deriving the upper bound for $r^{-1/2} |\zeta|$. Rewrite the test statistic as $\widehat{\mathcal{B}}_i = (T/\varphi_i)(\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2)^\top [\prod_{j=1}^2 (\widehat{\mathbf{V}}^j)^\top \widehat{\mathbf{V}}^j] (\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2)$. We do the decomposition $\zeta = \widehat{\mathcal{B}}_i - \mathcal{B}_i = r_1 + r_2$, where r_1 and r_2 are respectively defined by

$$r_1 := \frac{T}{\varphi_i} (\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2)^\top \left[\prod_{j=1}^2 (\widehat{\mathbf{V}}^j)^\top \widehat{\mathbf{V}}^j \right] (\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2) - \mathcal{B}_i,$$

and

$$r_2 := \widehat{\mathcal{B}}_i - \frac{T}{\varphi_i} (\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2)^\top \left[\prod_{j=1}^2 (\widehat{\mathbf{V}}^j)^\top \widehat{\mathbf{V}}^j \right] (\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2).$$

Step 1 – Upper bound for $|r_1|$.

We have that

$$r_1 = \frac{T}{\varphi_i} (\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2)^\top \left[\prod_{j=1}^2 (\widehat{\mathbf{V}}^j)^\top \widehat{\mathbf{V}}^j - \prod_{j=1}^2 (\mathbf{V}^j \mathbf{R}_V^\top)^\top \mathbf{V}^j \mathbf{R}_V^\top \right] (\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2).$$

Then using the results in Lemma 11, we obtain

$$\begin{aligned} |r_1| &\leq \frac{T}{\varphi_i} \left\| \prod_{j=1}^2 (\widehat{\mathbf{V}}^j)^\top \widehat{\mathbf{V}}^j - \prod_{j=1}^2 (\mathbf{V}^j \mathbf{R}_V^\top)^\top \mathbf{V}^j \mathbf{R}_V^\top \right\|_2 \|\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2\|_2^2 \\ &\lesssim \frac{T}{\varphi_i} \cdot (T_1 T_2 / T^2) \frac{1}{\theta} \sqrt{r} \log n \cdot \|\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2\|_2^2 = \frac{1}{\theta} \sqrt{r} \log n \cdot \frac{1}{\varphi_i} \frac{T_1 T_2}{T} \|\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2\|_2^2. \end{aligned}$$

Step 2 – Upper bound for $|r_2|$.

By definition, we have that

$$r_2 = T \left(\frac{1}{\widehat{\varphi}_i} - \frac{1}{\varphi_i} \right) \cdot (\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2)^\top \left[\prod_{j=1}^2 (\widehat{\mathbf{V}}^j)^\top \widehat{\mathbf{V}}^j \right] (\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2).$$

Then we have that

$$\begin{aligned} |r_2 / (\frac{1}{\widehat{\varphi}_i} - \frac{1}{\varphi_i})| &= T(\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2)^\top [\prod_{j=1}^2 (\widehat{\mathbf{V}}^j)^\top \widehat{\mathbf{V}}^j] (\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2) \\ &\lesssim T \prod_{j=1}^2 ((T_j/T)^{1/2})^2 \|\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2\|_2^2 \lesssim \frac{T_1 T_2}{T} \|\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2\|_2^2. \end{aligned}$$

So the problem boils down to bounding $|1/\widehat{\varphi}_i - 1/\varphi_i| = |\widehat{\varphi}_i - \varphi_i|/|\widehat{\varphi}_i \varphi_i|$. We obtain by Lemma 1 that $\|(\widehat{\boldsymbol{\Sigma}}_\varepsilon^T)_{i,\cdot} - (\boldsymbol{\Sigma}_\varepsilon)_{i,\cdot}\|_2 \leq \|\widehat{\boldsymbol{\Sigma}}_\varepsilon^T - \boldsymbol{\Sigma}_\varepsilon\|_2 \lesssim (\epsilon_{N,T})^{1-q} s(\boldsymbol{\Sigma}_\varepsilon)$ and $|(\widehat{\boldsymbol{\Sigma}}_\varepsilon^T)_{i,i} - (\boldsymbol{\Sigma}_\varepsilon)_{i,i}| = |(\widehat{\boldsymbol{\Sigma}}_\varepsilon)_{i,i} - (\boldsymbol{\Sigma}_\varepsilon)_{i,i}| \lesssim (\boldsymbol{\Sigma}_\varepsilon)_{i,i} \epsilon_{N,T}$. We obtain by Lemma 4 that $\|\widehat{\mathbf{U}}\mathbf{R}_U - \mathbf{U}\|_2 \lesssim \rho$, and we obtain by the proof of Corollary 1 that $\|(\widehat{\mathbf{U}}\mathbf{R}_U - \mathbf{U})_{i,\cdot}\|_2 \lesssim \omega_i \sqrt{\frac{r}{n}} \log n$. So we obtain that

$$\begin{aligned} |\widehat{\varphi}_i - \varphi_i| &\lesssim \epsilon_{N,T} (\boldsymbol{\Sigma}_\varepsilon)_{i,i} + (1 + \|\mathbf{U}_{i,\cdot}\|_2) \{\|\mathbf{U}_{i,\cdot}\|_2 (\epsilon_{N,T})^{1-q} s(\boldsymbol{\Sigma}_\varepsilon) \\ &\quad + \omega_i \sqrt{\frac{r}{n}} \log n \|(\boldsymbol{\Sigma}_\varepsilon)_{i,\cdot}\|_2 + \|\mathbf{U}_{i,\cdot}\|_2 \rho \|(\boldsymbol{\Sigma}_\varepsilon)_{i,\cdot}\|_2\}. \end{aligned}$$

Then we have that

$$\begin{aligned} \varphi_i &= (\mathbf{I}_N + \mathbf{U}\mathbf{U}^\top)_{i,\cdot} \boldsymbol{\Sigma}_\varepsilon (\mathbf{I}_N + \mathbf{U}\mathbf{U}^\top)_{i,\cdot} \geq \lambda_{\min}(\boldsymbol{\Sigma}_\varepsilon) \|(\mathbf{I}_N + \mathbf{U}\mathbf{U}^\top)_{i,\cdot}\|_2^2 = \lambda_{\min}(\boldsymbol{\Sigma}_\varepsilon) (1 + 3\|\mathbf{U}_{i,\cdot}\|_2^2), \\ &\gtrsim \frac{1}{\kappa_\varepsilon} \|\boldsymbol{\Sigma}_\varepsilon\|_2 (1 + \|\mathbf{U}_{i,\cdot}\|_2^2). \end{aligned}$$

Finally, we obtain that

$$\begin{aligned} |1/\widehat{\varphi}_i - 1/\varphi_i| &= \frac{|\widehat{\varphi}_i - \varphi_i|}{|\widehat{\varphi}_i| |\varphi_i|} \leq \frac{1}{\varphi_i \frac{1}{\kappa_\varepsilon} \|\boldsymbol{\Sigma}_\varepsilon\|_2 (1 + \|\mathbf{U}_{i,\cdot}\|_2^2)} \\ &\quad \times \{\epsilon_{N,T} (\boldsymbol{\Sigma}_\varepsilon)_{i,i} + (1 + \|\mathbf{U}_{i,\cdot}\|_2) \|\mathbf{U}_{i,\cdot}\|_2 (\epsilon_{N,T})^{1-q} s(\boldsymbol{\Sigma}_\varepsilon) + \omega_i \sqrt{\frac{r}{n}} \log n \|(\boldsymbol{\Sigma}_\varepsilon)_{i,\cdot}\|_2 + \|\mathbf{U}_{i,\cdot}\|_2 \rho \|(\boldsymbol{\Sigma}_\varepsilon)_{i,\cdot}\|_2\} \end{aligned}$$

We assemble all the above error bounds to obtain

$$\begin{aligned} |r_2| &\lesssim \frac{1}{\varphi_i} \frac{T_1 T_2}{T} \|\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2\|_2^2 \cdot \frac{\kappa_\varepsilon}{\|\boldsymbol{\Sigma}_\varepsilon\|_2} \\ &\quad \times \{\epsilon_{N,T} (\boldsymbol{\Sigma}_\varepsilon)_{i,i} + \sqrt{r} \|\mathbf{U}_{i,\cdot}\|_2 (\epsilon_{N,T})^{1-q} s(\boldsymbol{\Sigma}_\varepsilon) + \omega_i \sqrt{\frac{r}{n}} \log n \|(\boldsymbol{\Sigma}_\varepsilon)_{i,\cdot}\|_2 + \|\mathbf{U}_{i,\cdot}\|_2 \rho \|(\boldsymbol{\Sigma}_\varepsilon)_{i,\cdot}\|_2\} \\ &\lesssim \frac{1}{\varphi_i} \frac{T_1 T_2}{T} \|\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2\|_2^2 \cdot \kappa_\varepsilon \{\epsilon_{N,T} + (\epsilon_{N,T})^{1-q} \sqrt{r} \|\mathbf{U}_{i,\cdot}\|_2 \frac{s(\boldsymbol{\Sigma}_\varepsilon)}{\|\boldsymbol{\Sigma}_\varepsilon\|_2} + \frac{1}{\vartheta_i \sqrt{T}} r \log n + \sqrt{r} \|\mathbf{U}_{i,\cdot}\|_2 \frac{1}{\theta} \sqrt{\frac{n}{T}}\} \end{aligned}$$

where we use the fact that $(\boldsymbol{\Sigma}_\varepsilon)_{i,i} \leq \|\boldsymbol{\Sigma}_\varepsilon\|_2$ and $\|(\boldsymbol{\Sigma}_\varepsilon)_{i,\cdot}\|_2 \leq \|\boldsymbol{\Sigma}_\varepsilon\|_2$.

Step 3 – Combining all the bounds.

We obtain that

$$\begin{aligned} |\zeta| &\leq |r_1| + |r_2| \lesssim \frac{1}{\varphi_i} \frac{T_1 T_2}{T} \|\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2\|_2^2 \\ &\quad \times \left\{ \frac{1}{\theta} \sqrt{r} \log n + \kappa_\varepsilon [\epsilon_{N,T} + (\epsilon_{N,T})^{1-q} \sqrt{r} \|\mathbf{U}_{i,\cdot}\|_2 \frac{s(\boldsymbol{\Sigma}_\varepsilon)}{\|\boldsymbol{\Sigma}_\varepsilon\|_2} + \frac{1}{\vartheta_i \sqrt{T}} r \log n + \sqrt{r} \|\mathbf{U}_{i,\cdot}\|_2 \frac{1}{\theta} \sqrt{\frac{n}{T}}] \right\}. \end{aligned}$$

Then, the desired results follow from the inequality (E.1) in Lemma 12:

$$\begin{aligned} \left| \mathbb{P} \left(\widehat{\mathcal{B}}_i \leq \chi_{1-\alpha}^2(r) \right) - (1 - \alpha) \right| &\lesssim |\tau| + r^{-1/2} |\zeta| + s_3 \\ &\lesssim \left[\frac{1}{\theta} + \sqrt{\kappa_\varepsilon} \|\mathbf{U}_{i,\cdot}\|_2 \left(\frac{1}{\theta} \frac{n}{T} + \frac{1}{\sqrt{T}} \right) \right] \sqrt{T_1 T_2 / T} \sqrt{r} \log^{3/2} n + \frac{1}{\varphi_i} \frac{T_1 T_2}{T} \|\widehat{\mathbf{b}}_i^1 - \widehat{\mathbf{b}}_i^2\|_2^2 \end{aligned}$$

$$\begin{aligned}
& \times \left\{ \frac{1}{\theta} \sqrt{r} \log n + \kappa_\varepsilon \epsilon_{N,T} + \kappa_\varepsilon (\epsilon_{N,T})^{1-q} \sqrt{r} \|\mathbf{U}_{i,\cdot}\|_2 \frac{s(\boldsymbol{\Sigma}_\varepsilon)}{\|\boldsymbol{\Sigma}_\varepsilon\|_2} + \kappa_\varepsilon \frac{1}{\vartheta_i \sqrt{T}} r \log n + \kappa_\varepsilon \sqrt{r} \|\mathbf{U}_{i,\cdot}\|_2 \frac{1}{\theta} \sqrt{\frac{n}{T}} \right\} \\
& \lesssim \frac{1}{\theta} \sqrt{T_1 T_2 / T} \sqrt{r} \log^{3/2} n + \sqrt{\kappa_\varepsilon} \|\mathbf{U}_{i,\cdot}\|_2 \left(\frac{1}{\theta} \frac{n}{T} + \frac{1}{\sqrt{T}} \right) \sqrt{T_1 T_2 / T} \sqrt{r} \log^{3/2} n \\
& + \{ \kappa_\varepsilon \|\mathbf{U}_{i,\cdot}\|_2^2 \frac{T_1 T_2}{T} r \log n + \kappa_\varepsilon \frac{1}{\theta^2} \frac{T_1 T_2}{T} r \log n + \kappa_\varepsilon r \log n \} \\
& \times \left\{ \frac{1}{\theta} + \kappa_\varepsilon \epsilon_{N,T} + \kappa_\varepsilon (\epsilon_{N,T})^{1-q} \|\mathbf{U}_{i,\cdot}\|_2 \frac{s(\boldsymbol{\Sigma}_\varepsilon)}{\|\boldsymbol{\Sigma}_\varepsilon\|_2} + \kappa_\varepsilon \frac{1}{\vartheta_i \sqrt{T}} + \kappa_\varepsilon \|\mathbf{U}_{i,\cdot}\|_2 \frac{1}{\theta} \sqrt{\frac{n}{T}} \right\}.
\end{aligned}$$

F Proof of Theorem 4: Two-sample test for betas

F.1 A useful lemma

We prove a lemma that will be useful when establishing Theorem 4.

Lemma 13. *Assume that the assumptions in Theorem 4 hold. For any $i \neq j$ and $i, j \in \{1, 2, \dots, N\}$, under the null hypothesis $\mathbf{b}_i = \mathbf{b}_j$, we have that, for any random variable ζ satisfying $|\zeta| \ll r$, it holds*

$$\begin{aligned}
|\mathbb{P}(\mathcal{B}_{ij} + \zeta \leq \chi_{1-\alpha}^2(r)) - (1 - \alpha)| & \lesssim \frac{|\zeta|}{\sqrt{r}} + \tau_U^{(i,j)} \\
& + \frac{\sqrt{T} \sigma_r (\omega_i + \omega_j) \omega \sqrt{\frac{T}{n}} \log^{3/2} n + \sqrt{T} \sigma_r (\omega_i + \omega_j) \omega \log n \|\bar{\mathbf{U}}\|_{2,\infty}}{\sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i} + (\boldsymbol{\Sigma}_\varepsilon)_{j,j} - 2(\boldsymbol{\Sigma}_\varepsilon)_{i,j}}} \\
& + \frac{\sqrt{T} \sigma_r (\rho^2 + \rho \sqrt{\frac{T}{n}} \log n) (\|\bar{\mathbf{U}}_{i,\cdot}\|_2 + \|\bar{\mathbf{U}}_{j,\cdot}\|_2)}{\sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i} + (\boldsymbol{\Sigma}_\varepsilon)_{j,j} - 2(\boldsymbol{\Sigma}_\varepsilon)_{i,j}}},
\end{aligned}$$

where \mathcal{B}_{ij} is defined as

$$\mathcal{B}_{ij} := T((\boldsymbol{\Sigma}_\varepsilon)_{i,i} + (\boldsymbol{\Sigma}_\varepsilon)_{j,j} - 2(\boldsymbol{\Sigma}_\varepsilon)_{i,j})^{-1} \left\| (\hat{\mathbf{U}}_{i,\cdot} - \hat{\mathbf{U}}_{j,\cdot}) \hat{\boldsymbol{\Sigma}} \right\|_2^2,$$

and $\tau_U^{(i,j)}$ is a constant bounded by $\tau_U^{(i,j)} \lesssim r \sqrt{\frac{\log n}{T}}$. Further, if all the entries of the matrix \mathbf{Z} are Gaussian, i.e., the noise is Gaussian, then result holds even when $\tau_U^{(i,j)} = 0$.

Proof. Step 1 – Expand $(\hat{\mathbf{U}}_{i,\cdot} - \hat{\mathbf{U}}_{j,\cdot}) \hat{\boldsymbol{\Sigma}}$ and develop the whole roadmap for the proof.

We already have the expansion of $\hat{\mathbf{U}} \mathbf{R}_U - \mathbf{U}$ and the error bound for $\varepsilon_\Sigma := (\mathbf{R}_U)^\top \hat{\boldsymbol{\Sigma}} \mathbf{R}_V - \boldsymbol{\Lambda}$, so we obtain that

$$\begin{aligned}
\hat{\mathbf{U}} \hat{\boldsymbol{\Sigma}} \mathbf{R}_V & = \hat{\mathbf{U}} \mathbf{R}_U (\mathbf{R}_U)^\top \hat{\boldsymbol{\Sigma}} \mathbf{R}_V = (\mathbf{U} + \mathbf{G}_U + \boldsymbol{\Psi}_U) (\boldsymbol{\Lambda} + \varepsilon_\Sigma) = (\mathbf{U} + \mathbf{G}_U + \boldsymbol{\Psi}_U) \boldsymbol{\Lambda} + \hat{\mathbf{U}} \mathbf{R}_U \varepsilon_\Sigma \\
& = \mathbf{U} \boldsymbol{\Lambda} + T^{-1/2} \boldsymbol{\Sigma}_\varepsilon^{1/2} \mathbf{Z} \mathbf{V} + (\hat{\mathbf{U}} \mathbf{R}_U \varepsilon_\Sigma + \boldsymbol{\Psi}_U \boldsymbol{\Lambda}).
\end{aligned}$$

Note that, by Lemma 2, conditioning on the good event \mathcal{E}_0 , we have that $\mathbf{U}_{i,\cdot} - \mathbf{U}_{j,\cdot} = 0$ under the null $\mathbf{b}_i = \mathbf{b}_j$. The reason is that, there exists a rotation matrix \mathbf{Q} such that $\mathbf{U} = \bar{\mathbf{U}} \mathbf{Q}$, and by definition we have that $\bar{\mathbf{U}}_{i,\cdot} = \bar{\mathbf{U}}_{j,\cdot}$, since $\bar{\mathbf{U}} = \mathbf{B} \boldsymbol{\Sigma}^{-1}$ and $\mathbf{b}_i = \mathbf{b}_j$, i.e., $\mathbf{B}_{i,\cdot} = \mathbf{B}_{j,\cdot}$.

The following derivations are conditioning on \mathcal{E}_0 and we will take the effect of \mathcal{E}_0 into consideration in the final non-asymptotic probability bound. Then we have that

$$\begin{aligned}
& (\hat{\mathbf{U}}_{i,\cdot} - \hat{\mathbf{U}}_{j,\cdot}) \hat{\boldsymbol{\Sigma}} \mathbf{R}_V \\
& = (\mathbf{U}_{i,\cdot} - \mathbf{U}_{j,\cdot}) \boldsymbol{\Lambda} + T^{-1/2} ((\boldsymbol{\Sigma}_\varepsilon^{1/2})_{i,\cdot} - (\boldsymbol{\Sigma}_\varepsilon^{1/2})_{j,\cdot}) \mathbf{Z} \mathbf{V} + (\hat{\mathbf{U}}_{i,\cdot} - \hat{\mathbf{U}}_{j,\cdot}) \mathbf{R}_U \varepsilon_\Sigma + ((\boldsymbol{\Psi}_U)_i - (\boldsymbol{\Psi}_U)_j) \boldsymbol{\Lambda} \\
& = T^{-1/2} \mathbf{a} \mathbf{Z} \mathbf{V} + ((\boldsymbol{\Psi}_U)_i - (\boldsymbol{\Psi}_U)_j) \boldsymbol{\Lambda} = T^{-1/2} \sum_{t=1}^T (\mathbf{a} \mathbf{Z}_{\cdot,t}) \mathbf{V}_{t,\cdot} + ((\boldsymbol{\Psi}_U)_i - (\boldsymbol{\Psi}_U)_j) \boldsymbol{\Lambda},
\end{aligned}$$

where, for simplicity of notations, we denote $a_k := T^{-1/2}(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{i,k} - (\boldsymbol{\Sigma}_\varepsilon^{1/2})_{j,k}$ and denote $\mathbf{a} = (a_1, a_2, \dots, a_N)$ as a row vector satisfying that

$$\|\mathbf{a}\|_2^2 = \|\mathbf{a}\|_F^2 = \sum_{k=1}^N a_k^2 = \frac{1}{T}((\boldsymbol{\Sigma}_\varepsilon)_{i,i} + (\boldsymbol{\Sigma}_\varepsilon)_{j,j} - 2(\boldsymbol{\Sigma}_\varepsilon)_{i,j}).$$

We will show proximity between the law of $\sum_{t=1}^T (\mathbf{a}\mathbf{Z}_{\cdot,t})\mathbf{V}_{t\cdot}$ and Gaussian, and prove that the term

$$\varepsilon_2 := \frac{1}{\|\mathbf{a}\|_2} \|((\boldsymbol{\Psi}_U)_i, -(\boldsymbol{\Psi}_U)_j)\boldsymbol{\Lambda}\|_2$$

is negligible.

We will establish the upper bound for the parameter

$$\tau_U^{(i,j)} := \sup_{R \geq 0} \left| \mathbb{P} \left(\frac{1}{\|\mathbf{a}\|_2} \left\| \sum_{t=1}^T (\mathbf{a}\mathbf{Z}_{\cdot,t})\mathbf{V}_{t\cdot} \right\|_2 \leq R \right) - \mathbb{P}(\chi^2(r) \leq R^2) \right|.$$

The motivations for establishing the upper bound for $\tau_U^{(i,j)}$ are as follows: since $\|\mathbf{a}\|_2^2 = \frac{1}{T}((\boldsymbol{\Sigma}_\varepsilon)_{i,i} + (\boldsymbol{\Sigma}_\varepsilon)_{j,j} - 2(\boldsymbol{\Sigma}_\varepsilon)_{i,j})$, we can rewrite \mathcal{B}_{ij} as

$$\mathcal{B}_{ij} := \frac{1}{\|\mathbf{a}\|_2} \left\| \sum_{t=1}^T (\mathbf{a}\mathbf{Z}_{\cdot,t})\mathbf{V}_{t\cdot} + ((\boldsymbol{\Psi}_U)_i, -(\boldsymbol{\Psi}_U)_j)\boldsymbol{\Lambda} \right\|_2^2.$$

Similar to the techniques we used for projection matrix of factors in Lemma 10, we have that, for any random variable ζ satisfying $|\zeta| \ll r$,

$$\begin{aligned} & \mathbb{P} \left(\frac{1}{\|\mathbf{a}\|_2} \left\| \sum_{t=1}^T (\mathbf{a}\mathbf{Z}_{\cdot,t})\mathbf{V}_{t\cdot} \right\|_2 \leq \sqrt{\chi_{1-\alpha}^2(r) - |\zeta| - \varepsilon_2} \right) \\ & \leq \mathbb{P}(\mathcal{B}_{ij} + \zeta \leq \chi_{1-\alpha}^2(r)) \\ & \leq \mathbb{P} \left(\frac{1}{\|\mathbf{a}\|_2} \left\| \sum_{t=1}^T (\mathbf{a}\mathbf{Z}_{\cdot,t})\mathbf{V}_{t\cdot} \right\|_2 \leq \sqrt{\chi_{1-\alpha}^2(r) + |\zeta| + \varepsilon_2} \right). \end{aligned}$$

Then by definition of $\tau_U^{(i,j)}$, we have that

$$\mathbb{P} \left(\sqrt{\chi^2(r)} \leq \sqrt{\chi_{1-\alpha}^2(r) - |\zeta| - \varepsilon_2} \right) - \tau_U^{(i,j)} \leq \mathbb{P}(\mathcal{B}_{ij} + \zeta \leq \chi_{1-\alpha}^2(r)) \leq \mathbb{P} \left(\sqrt{\chi^2(r)} \leq \sqrt{\chi_{1-\alpha}^2(r) + |\zeta| + \varepsilon_2} \right) + \tau_U^{(i,j)}.$$

Similar to Step 3 of the proof of Lemma 10, since $|\zeta| \ll r \asymp \chi_{1-\alpha}^2(r)$, we have that the left bound of the above inequality satisfies that

$$\mathbb{P}(\chi^2(r) \leq \chi_{1-\alpha}^2(r)) - h \lesssim \mathbb{P} \left(\sqrt{\chi^2(r)} \leq \sqrt{\chi_{1-\alpha}^2(r) - h} \right) \leq \mathbb{P} \left(\sqrt{\chi^2(r)} \leq \sqrt{\chi_{1-\alpha}^2(r) - |\zeta| - \varepsilon_2} \right),$$

where $h > 0$ satisfies that $h \lesssim |\varepsilon_2| + \frac{1}{\sqrt{r}}|\zeta|$. Similar inequality holds for the right bound $\mathbb{P}(\sqrt{\chi^2(r)} \leq \sqrt{\chi_{1-\alpha}^2(r) + |\zeta| + \varepsilon_2})$ of $\mathbb{P}(\mathcal{B}_{ij} + \zeta \leq \chi_{1-\alpha}^2(r))$, so we obtain that

$$|\mathbb{P}(\mathcal{B}_{ij} + \zeta \leq \chi_{1-\alpha}^2(r)) - (1 - \alpha)| \lesssim \varepsilon_2 + \frac{1}{\sqrt{r}}|\zeta| + \tau_U^{(i,j)}.$$

Thus, the problems boil down to proving the upper bounds for $\tau_U^{(i,j)}$ and ε_2 .

Step 2 – Prove the upper bound for $\tau_U^{(i,j)}$ to show the proximity of $\sum_{t=1}^T (\mathbf{a}\mathbf{Z}_{\cdot,t})\mathbf{V}_{t\cdot}$ with a Gaussian vector.

Conditioning on \mathbf{F} , the variance of $\sum_{t=1}^T (\mathbf{a}\mathbf{Z}_{\cdot,t})\mathbf{V}_{t,\cdot}$ is given by

$$\sum_{t=1}^T \mathbb{E}[(\mathbf{a}\mathbf{Z}_{\cdot,t})^2](\mathbf{V}_{t,\cdot})^\top \mathbf{V}_{t,\cdot} = \|\mathbf{a}\|_2^2 \mathbf{V}^\top \mathbf{V} = \|\mathbf{a}\|_2^2 \mathbf{I}_r,$$

where we use the fact that \mathbf{V} is $\sigma(\mathbf{F})$ -measurable and \mathbf{Z} is independent with \mathbf{F} . Then we have that

(i) When all entries of \mathbf{Z} are Gaussian, we have that, the law of $\frac{1}{\|\mathbf{a}\|_2} \sum_{t=1}^T (\mathbf{a}\mathbf{Z}_{\cdot,t})\mathbf{V}_{t,\cdot}$ is r -dimensional standard Gaussian $\mathcal{N}(0, \mathbf{I}_r)$, because it is a linear combination of Gaussian vectors. Thus, we have that $\tau_U^{(i,j)} = 0$ when all entries of \mathbf{Z} are Gaussian.

(ii) When the entries of \mathbf{Z} are sub-Gaussian, we obtain by Lemma 16 that

$$\begin{aligned} \tau_U^{(i,j)} &\lesssim \frac{1}{\|\mathbf{a}\|_2^3} \sum_{t=1}^T \mathbb{E} \left[\left\| (\mathbf{a}\mathbf{Z}_{\cdot,t})\mathbf{V}_{t,\cdot} \right\|_2^3 \right] = \frac{1}{\|\mathbf{a}\|_2^3} \mathbb{E} \left[\left\| (\mathbf{a}\mathbf{Z}_{\cdot,t}) \right\|_2^3 \right] \max_{1 \leq t \leq T} \|\mathbf{V}_{t,\cdot}\|_2 \sum_{t=1}^T \|\mathbf{V}_{t,\cdot}\|_2^2 \\ &\lesssim \frac{1}{\|\mathbf{a}\|_2^3} \|\mathbf{a}\|_2^3 \cdot \max_{1 \leq t \leq T} \|\mathbf{V}_{t,\cdot}\|_2 \cdot r \lesssim r \sqrt{\frac{\log n}{T}}, \end{aligned}$$

where we used the fact that $\mathbb{E} \left[\left\| (\mathbf{a}\mathbf{Z}_{\cdot,t}) \right\|_2^3 \right] \lesssim \|\mathbf{a}\|_2^3$. We now prove it as follows. By Exercise 6.3.5 in Vershynin (2017), we have that, since $\|\mathbf{a}\|_2 = \|\mathbf{a}\|_{\mathbb{F}}$, there exist constants $C_1 > 0, C_2 > 0$ such that, for any $t \geq 0$, it holds

$$\mathbb{P}(|\mathbf{a}\mathbf{Z}_{\cdot,t}| \geq C_1 \|\mathbf{a}\|_2 + t) \leq \exp(-C_2 \frac{t^2}{\|\mathbf{a}\|_2^2}), \text{ i.e., } \mathbb{P}(|\mathbf{a}\mathbf{Z}_{\cdot,t}| \geq C_1 \|\mathbf{a}\|_2(1+t)) \leq \exp(-C_2 t^2)$$

So similarly we have

$$\begin{aligned} \mathbb{E} [|\mathbf{a}\mathbf{Z}_{\cdot,t}|^3] &= \int_0^{+\infty} 3s^2 \mathbb{P}(|\mathbf{a}\mathbf{Z}_{\cdot,t}| \geq s) ds \\ &= \int_0^{C_1 \|\mathbf{a}\|_2} 3s^2 \mathbb{P}(|\mathbf{a}\mathbf{Z}_{\cdot,t}| \geq s) ds \\ &\quad + \int_0^{+\infty} 3(C_1 \|\mathbf{a}\|_2(1+t))^2 \mathbb{P}(|\mathbf{a}\mathbf{Z}_{\cdot,t}| \geq C_1 \|\mathbf{a}\|_2(1+t)) dt \\ &\leq \int_0^{C_1 \|\mathbf{a}\|_2} 3s^2 ds + \int_0^{+\infty} 3(C_1 \|\mathbf{a}\|_2(1+t))^2 \exp(-C_2 t^2) dt \\ &\lesssim \|\mathbf{a}\|_2^3. \end{aligned}$$

Step 3 – Prove the upper bound for ε_2 .

Recall that

$$\varepsilon_2 = \frac{1}{\|\mathbf{a}\|_2} \left\| ((\Psi_U)_i, -(\Psi_U)_j) \mathbf{\Lambda} \right\|_2 = \frac{\sqrt{T}}{\sqrt{(\Sigma_\varepsilon)_{i,i} + (\Sigma_\varepsilon)_{j,j} - 2(\Sigma_\varepsilon)_{i,j}}} \left\| ((\Psi_U)_i, -(\Psi_U)_j) \mathbf{\Lambda} \right\|_2.$$

Then using the decomposition of Ψ_U in the proof of Theorem 1, we obtain that

$$\begin{aligned} \left\| ((\Psi_U)_i, -(\Psi_U)_j) \mathbf{\Lambda} \right\|_2 &\lesssim \sigma_r(\omega_i + \omega_j) \omega \sqrt{\frac{r}{n}} \log^{3/2} n + \sigma_r(\omega_i + \omega_j) \omega \log n \|\bar{\mathbf{U}}\|_{2,\infty} \\ &\quad + \sigma_r \left(\rho^2 + \rho \sqrt{\frac{r}{n}} \log n \right) (\|\bar{\mathbf{U}}_{i,\cdot}\|_2 + \|\bar{\mathbf{U}}_{j,\cdot}\|_2), \end{aligned}$$

Finally, the desired result follows from assembling the upper bounds of $\tau_U^{(i,j)}$ and ε_2 . \square

F.2 Proof of Theorem 4:

We are now ready to prove the main theorem. Define $\mathbf{e}_{ij} \in \mathbb{R}^N$ as the vector where the i -th (resp. j -th) entry is 1 (resp. -1) and the other entries are all zero. Note that $T\|\mathbf{a}\|_2^2 = (\boldsymbol{\Sigma}_\varepsilon)_{i,i} + (\boldsymbol{\Sigma}_\varepsilon)_{j,j} - 2(\boldsymbol{\Sigma}_\varepsilon)_{i,j} = \mathbf{e}_{ij}^\top \boldsymbol{\Sigma}_\varepsilon \mathbf{e}_{ij} \geq \lambda_{\min}(\boldsymbol{\Sigma}_\varepsilon) \|\mathbf{e}_{ij}\|_2^2 = 2\lambda_{\min}(\boldsymbol{\Sigma}_\varepsilon)$, $|\mathbf{e}_{ij}^\top (\widehat{\boldsymbol{\Sigma}}_\varepsilon^\tau - \boldsymbol{\Sigma}_\varepsilon) \mathbf{e}_{ij}| \lesssim (\epsilon_{N,T})^{1-q} s(\boldsymbol{\Sigma}_\varepsilon) \ll 1$, and thus $(\widehat{\boldsymbol{\Sigma}}_\varepsilon^\tau)_{i,i} + (\widehat{\boldsymbol{\Sigma}}_\varepsilon^\tau)_{j,j} - 2(\widehat{\boldsymbol{\Sigma}}_\varepsilon^\tau)_{i,j} = \mathbf{e}_{ij}^\top \boldsymbol{\Sigma}_\varepsilon \mathbf{e}_{ij} + \mathbf{e}_{ij}^\top (\widehat{\boldsymbol{\Sigma}}_\varepsilon^\tau - \boldsymbol{\Sigma}_\varepsilon) \mathbf{e}_{ij} \gtrsim \lambda_{\min}(\boldsymbol{\Sigma}_\varepsilon)$. So we obtain that

$$|1/\mathbf{e}_{ij}^\top \widehat{\boldsymbol{\Sigma}}_\varepsilon^\tau \mathbf{e}_{ij} - 1/\mathbf{e}_{ij}^\top \boldsymbol{\Sigma}_\varepsilon \mathbf{e}_{ij}| = \frac{|\mathbf{e}_{ij}^\top (\widehat{\boldsymbol{\Sigma}}_\varepsilon^\tau - \boldsymbol{\Sigma}_\varepsilon) \mathbf{e}_{ij}|}{(\mathbf{e}_{ij}^\top \widehat{\boldsymbol{\Sigma}}_\varepsilon^\tau \mathbf{e}_{ij})(\mathbf{e}_{ij}^\top \boldsymbol{\Sigma}_\varepsilon \mathbf{e}_{ij})} \lesssim (\epsilon_{N,T})^{1-q} s(\boldsymbol{\Sigma}_\varepsilon) \frac{1}{\lambda_{\min}(\boldsymbol{\Sigma}_\varepsilon) T \|\mathbf{a}\|_2^2}.$$

We let

$$\zeta := \widehat{\mathcal{B}}_{ij} - \mathcal{B}_{ij} = T(1/\mathbf{e}_{ij}^\top \widehat{\boldsymbol{\Sigma}}_\varepsilon^\tau \mathbf{e}_{ij} - 1/\mathbf{e}_{ij}^\top \boldsymbol{\Sigma}_\varepsilon \mathbf{e}_{ij}) \|(\widehat{\mathbf{U}}_{i,\cdot} - \widehat{\mathbf{U}}_{j,\cdot}) \widehat{\boldsymbol{\Sigma}}\|_2^2.$$

We derive the upper bound for $\|(\widehat{\mathbf{U}}_{i,\cdot} - \widehat{\mathbf{U}}_{j,\cdot}) \widehat{\boldsymbol{\Sigma}}\|_2^2$ first. Recall that

$$(\widehat{\mathbf{U}}_{i,\cdot} - \widehat{\mathbf{U}}_{j,\cdot}) \widehat{\boldsymbol{\Sigma}} \mathbf{R}_V = \sum_{t=1}^T (\mathbf{a} \mathbf{Z}_{\cdot,t}) \mathbf{V}_{t,\cdot} + ((\boldsymbol{\Psi}_U)_i, -(\boldsymbol{\Psi}_U)_j) \boldsymbol{\Lambda}.$$

Note that $\sum_{t=1}^T (\mathbf{a} \mathbf{Z}_{\cdot,t}) \mathbf{V}_{t,\cdot}$ is the sum of independent mean zero random vectors. Given the event \mathcal{E}_Z defined in Lemma 3, it holds $\|(\mathbf{a} \mathbf{Z}_{\cdot,t}) \mathbf{V}_{t,\cdot}\|_2 \leq C\sqrt{\log n} \|\mathbf{a}\|_2 \|\mathbf{V}_{t,\cdot}\|_2$ for $t = 1, 2, \dots, T$. Then, we obtain by the matrix Hoeffding inequality (Tropp, 2012, Theorem 1.3) that $\|\sum_{t=1}^T (\mathbf{a} \mathbf{Z}_{\cdot,t}) \mathbf{V}_{t,\cdot}\|_2 \lesssim \sigma\sqrt{\log n}$, where $\sigma = [\sum_{t=1}^T (C\sqrt{\log n} \|\mathbf{a}\|_2 \|\mathbf{V}_{t,\cdot}\|_2)^2]^{1/2} = C\sqrt{\log n} \|\mathbf{a}\|_2 r$. So, we obtain that, with probability at least $1 - O(n^{-2})$, it holds

$$\left\| \sum_{t=1}^T (\mathbf{a} \mathbf{Z}_{\cdot,t}) \mathbf{V}_{t,\cdot} \right\|_2 \lesssim \|\mathbf{a}\|_2 r \log n.$$

Then we obtain that

$$\begin{aligned} \|(\widehat{\mathbf{U}}_{i,\cdot} - \widehat{\mathbf{U}}_{j,\cdot}) \widehat{\boldsymbol{\Sigma}}\|_2^2 &\lesssim \left\| \sum_{t=1}^T (\mathbf{a} \mathbf{Z}_{\cdot,t}) \mathbf{V}_{t,\cdot} \right\|_2^2 + \|((\boldsymbol{\Psi}_U)_i, -(\boldsymbol{\Psi}_U)_j) \boldsymbol{\Lambda}\|_2^2 \\ &\lesssim \|\mathbf{a}\|_2^2 r^2 \log^2 n + \|\mathbf{a}\|_2^2 \varepsilon_2^2, \end{aligned}$$

where ε_2 is defined in the proof of Lemma 13. Thus, we obtain that

$$\begin{aligned} |\zeta| &= T(1/\mathbf{e}_{ij}^\top \widehat{\boldsymbol{\Sigma}}_\varepsilon^\tau \mathbf{e}_{ij} - 1/\mathbf{e}_{ij}^\top \boldsymbol{\Sigma}_\varepsilon \mathbf{e}_{ij}) \|(\widehat{\mathbf{U}}_{i,\cdot} - \widehat{\mathbf{U}}_{j,\cdot}) \widehat{\boldsymbol{\Sigma}}\|_2^2 \\ &\lesssim T(\epsilon_{N,T})^{1-q} s(\boldsymbol{\Sigma}_\varepsilon) \frac{1}{\lambda_{\min}(\boldsymbol{\Sigma}_\varepsilon) T \|\mathbf{a}\|_2^2} (\|\mathbf{a}\|_2^2 r^2 \log^2 n + \|\mathbf{a}\|_2^2 \varepsilon_2^2) \\ &\lesssim (\epsilon_{N,T})^{1-q} (r^2 \log^2 n + \varepsilon_2^2), \end{aligned}$$

where we used the assumption that $s(\boldsymbol{\Sigma}_\varepsilon)/\lambda_{\min}(\boldsymbol{\Sigma}_\varepsilon) \leq C_\varepsilon$ for some universal constant C_ε . Plugging the above bound of $|\zeta|$ into Lemma 13, we obtain that

$$\begin{aligned} \left| \mathbb{P}\left(\widehat{\mathcal{B}}_{ij} \leq \chi_{1-\alpha}^2(r)\right) - (1-\alpha) \right| &\lesssim (\epsilon_{N,T})^{1-q} r^{3/2} \log^2 n + \tau_U^{(i,j)} + \frac{1}{\sqrt{r}} (\epsilon_{N,T})^{1-q} \varepsilon_2^2 + \varepsilon_2 \\ &\lesssim (\epsilon_{N,T})^{1-q} r^{3/2} \log^2 n + \tau_U^{(i,j)} + \varepsilon_2, \end{aligned}$$

where we use the assumption that $\epsilon_{N,T} \ll 1$ and $\varepsilon_2 \ll 1$. Using the upper bound of ε_2 obtained in the proof of Lemma 13, we have that

$$\varepsilon_2 \lesssim \frac{\sqrt{T} \sigma_r(\omega_i + \omega_j) \omega \sqrt{\frac{T}{n}} \log^{3/2} n + \sqrt{T} \sigma_r(\omega_i + \omega_j) \omega \log n \|\bar{\mathbf{U}}\|_{2,\infty}}{\sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i} + (\boldsymbol{\Sigma}_\varepsilon)_{j,j} - 2(\boldsymbol{\Sigma}_\varepsilon)_{i,j}}}$$

$$\begin{aligned}
& + \frac{\sqrt{T}\sigma_r (\rho^2 + \rho\sqrt{\frac{r}{n}}\log n) (\|\bar{U}_{i,\cdot}\|_2 + \|\bar{U}_{j,\cdot}\|_2)}{\sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i} + (\boldsymbol{\Sigma}_\varepsilon)_{j,j} - 2(\boldsymbol{\Sigma}_\varepsilon)_{i,j}}} \\
& \lesssim \frac{\sqrt{\|\boldsymbol{\Sigma}_\varepsilon\|_2}}{\sqrt{\lambda_{\min}(\boldsymbol{\Sigma}_\varepsilon)}} \sqrt{T}\theta(\omega_i + \omega_j)\omega(\sqrt{\frac{r}{n}}\log^{3/2}n + \log n\|\bar{U}\|_{2,\infty}) \\
& + \frac{\sqrt{\|\boldsymbol{\Sigma}_\varepsilon\|_2}}{\sqrt{\lambda_{\min}(\boldsymbol{\Sigma}_\varepsilon)}} \sqrt{T}\theta \left(\rho^2 + \rho\sqrt{\frac{r}{n}}\log n \right) (\|\bar{U}_{i,\cdot}\|_2 + \|\bar{U}_{j,\cdot}\|_2) \\
& \lesssim \sqrt{\kappa_\varepsilon} \frac{\theta n}{\vartheta\sqrt{T}} \left(\frac{1}{\vartheta_i} + \frac{1}{\vartheta_j} \right) (\|\bar{U}\|_{2,\infty} + \frac{1}{\sqrt{n}}) \sqrt{r}\log^{3/2}n + \sqrt{\kappa_\varepsilon} \left(\frac{n}{\theta\sqrt{T}} + 1 \right) (\|\bar{U}_{i,\cdot}\|_2 + \|\bar{U}_{j,\cdot}\|_2) \sqrt{r}\log n.
\end{aligned}$$

The final result follows from the above upper bound.

G Proof of Theorem 5: Inference for the systematic risks

We start by rewriting our target error as follows,

$$\left| \mathbb{P} \left(\frac{1}{\hat{\sigma}_{B,i}} (\|\tilde{\mathbf{B}}_{i,\cdot}\|_2^2 - \|(\hat{U}\hat{\Sigma})_{i,\cdot}\|_2^2) \in [-z_{1-\frac{1}{2}\alpha}, z_{1-\frac{1}{2}\alpha}] \right) - (1-\alpha) \right|.$$

Step 1 – Rewrite the first-order approximation to show the Gaussian term.

Our starting point is to prove that, for any random variable ζ satisfying $|\zeta| \ll 1$, it holds

$$\left| \mathbb{P} \left(\frac{1}{\sigma_{B,i}} (\|\tilde{\mathbf{B}}_{i,\cdot}\|_2^2 - \|(\hat{U}\hat{\Sigma})_{i,\cdot}\|_2^2) + \zeta \leq z_{1-\frac{1}{2}\alpha} - (1-\alpha) \right) \right| \lesssim |\varepsilon_{B,i}| + |\zeta| + \tau_{B,i}^i,$$

where $\sigma_{B,i} = \frac{2}{\sqrt{T}} \sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i}} \|\tilde{\mathbf{B}}_{i,\cdot}\|_2$. Then we let

$$\zeta = \left(\frac{1}{\hat{\sigma}_{B,i}} - \frac{1}{\sigma_{B,i}} \right) (\|\tilde{\mathbf{B}}_{i,\cdot}\|_2^2 - \|(\hat{U}\hat{\Sigma})_{i,\cdot}\|_2^2) = \left(\frac{1}{\sqrt{(\hat{\Sigma}_\varepsilon^T)_{i,i}}} - \frac{1}{\sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i}}} \right) \frac{\sqrt{T}}{2\|\tilde{\mathbf{B}}_{i,\cdot}\|_2} (\|\tilde{\mathbf{B}}_{i,\cdot}\|_2^2 - \|(\hat{U}\hat{\Sigma})_{i,\cdot}\|_2^2),$$

and later we will derive an upper bound for $|\zeta|$. We already have the expansion of $\hat{U}\mathbf{R}_U - U$ and the error bound for $\varepsilon_\Sigma := (\mathbf{R}_U)^\top \hat{\Sigma} \mathbf{R}_V - \boldsymbol{\Lambda}$, so we obtain that

$$\begin{aligned}
\hat{U}\hat{\Sigma}\mathbf{R}_V &= \hat{U}\mathbf{R}_U \cdot (\mathbf{R}_U)^\top \hat{\Sigma} \mathbf{R}_V = (\mathbf{U} + \mathbf{G}_U + \boldsymbol{\Psi}_U)(\boldsymbol{\Lambda} + \varepsilon_\Sigma) \\
&= \tilde{\mathbf{B}} + \mathbf{G}_U \boldsymbol{\Lambda} + \hat{U}\mathbf{R}_U \varepsilon_\Sigma + \boldsymbol{\Psi}_U \boldsymbol{\Lambda}.
\end{aligned}$$

Since \mathbf{R}_V is a rotation matrix, we obtain

$$\begin{aligned}
\|(\hat{U}\hat{\Sigma})_{i,\cdot}\|_2^2 &= \|(\hat{U}\hat{\Sigma})_{i,\cdot} \mathbf{R}_V\|_2^2 = (\hat{U}\hat{\Sigma}\mathbf{R}_V)_{i,\cdot} ((\hat{U}\hat{\Sigma}\mathbf{R}_V)_{i,\cdot})^\top \\
&= (\hat{U}\mathbf{R}_U)_{i,\cdot} (\boldsymbol{\Lambda}^2 + \varepsilon_\Sigma \boldsymbol{\Lambda} + \boldsymbol{\Lambda}(\varepsilon_\Sigma)^\top + \varepsilon_\Sigma(\varepsilon_\Sigma)^\top) ((\hat{U}\mathbf{R}_U)_{i,\cdot})^\top \\
&= (\hat{U}\mathbf{R}_U)_{i,\cdot} (\boldsymbol{\Lambda}^2) ((\hat{U}\mathbf{R}_U)_{i,\cdot})^\top + \varepsilon_1 = (\mathbf{U} + \mathbf{G}_U + \boldsymbol{\Psi}_U)_{i,\cdot} (\boldsymbol{\Lambda}^2) ((\mathbf{U} + \mathbf{G}_U + \boldsymbol{\Psi}_U)_{i,\cdot})^\top + \varepsilon_1 \\
&= \|\tilde{\mathbf{B}}_{i,\cdot}\|_2^2 + (\mathbf{U}_{i,\cdot} \boldsymbol{\Lambda}^2 ((\mathbf{G}_U)_{i,\cdot})^\top + (\mathbf{G}_U)_{i,\cdot} \boldsymbol{\Lambda}^2 (\mathbf{U}_{i,\cdot})^\top) + \varepsilon_1 + \varepsilon_2 \\
&= \|\mathbf{B}_{i,\cdot}\|_2^2 + (\mathbf{U}_{i,\cdot} \boldsymbol{\Lambda}^2 ((\mathbf{G}_U)_{i,\cdot})^\top + (\mathbf{G}_U)_{i,\cdot} \boldsymbol{\Lambda}^2 (\mathbf{U}_{i,\cdot})^\top) + \varepsilon_1 + \varepsilon_2 + \varepsilon_J,
\end{aligned}$$

where ε_1 , ε_2 , and ε_J are scalars defined by

$$\begin{aligned}
\varepsilon_1 &:= (\hat{U}\mathbf{R}_U)_{i,\cdot} (\varepsilon_\Sigma \boldsymbol{\Lambda} + \boldsymbol{\Lambda}(\varepsilon_\Sigma)^\top + \varepsilon_\Sigma(\varepsilon_\Sigma)^\top) ((\hat{U}\mathbf{R}_U)_{i,\cdot})^\top, \\
\varepsilon_2 &:= \mathbf{U}_{i,\cdot} \boldsymbol{\Lambda}^2 ((\boldsymbol{\Psi}_U)_{i,\cdot})^\top + (\boldsymbol{\Psi}_U)_{i,\cdot} \boldsymbol{\Lambda}^2 (\mathbf{U}_{i,\cdot})^\top + (\mathbf{G}_U + \boldsymbol{\Psi}_U)_{i,\cdot} (\boldsymbol{\Lambda}^2) ((\mathbf{G}_U + \boldsymbol{\Psi}_U)_{i,\cdot})^\top, \\
\varepsilon_J &:= \|\tilde{\mathbf{B}}_{i,\cdot}\|_2^2 - \|\mathbf{B}_{i,\cdot}\|_2^2 = \mathbf{B}_{i,\cdot} ((\mathbf{J}^{-1})^\top \mathbf{J}^{-1} - \mathbf{I}_r) (\mathbf{B}_{i,\cdot})^\top.
\end{aligned}$$

The expression of ε_J uses the fact that $\mathbf{B} = \tilde{\mathbf{B}}\mathbf{J}^\top$. Note that we have

$$\begin{aligned} \mathbf{U}_{i,\cdot}\boldsymbol{\Lambda}^2((\mathbf{G}_U)_{i,\cdot})^\top &= (\mathbf{G}_U)_{i,\cdot}\boldsymbol{\Lambda}^2(\mathbf{U}_{i,\cdot})^\top = T^{-1/2}(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{i,\cdot}\mathbf{Z}\mathbf{V}\boldsymbol{\Lambda}(\mathbf{U}_{i,\cdot})^\top \\ &= T^{-1/2}(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{i,\cdot}\mathbf{Z}\mathbf{V}(\tilde{\mathbf{B}}_{i,\cdot})^\top \\ &= T^{-1/2}\sum_{t=1}^T(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{i,\cdot}\mathbf{Z}_{\cdot,t}\mathbf{V}_{t,\cdot}(\tilde{\mathbf{B}}_{i,\cdot})^\top \end{aligned}$$

is a scalar. So we obtain

$$\frac{1}{\sigma_{B,i}}(\|(\widehat{\mathbf{U}}\widehat{\boldsymbol{\Sigma}})_{i,\cdot}\|_2^2 - \|\tilde{\mathbf{B}}_{i,\cdot}\|_2^2) = \mathbf{K}_{B,i} + \varepsilon_{B,i}, \quad \text{with} \quad \mathbf{K}_{B,i} := \frac{1}{\sigma_{B,i}}2T^{-1/2}\sum_{t=1}^T(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{i,\cdot}\mathbf{Z}_{\cdot,t}\mathbf{V}_{t,\cdot}(\tilde{\mathbf{B}}_{i,\cdot})^\top,$$

and

$$|\varepsilon_{B,i}| \leq \frac{1}{\sigma_{B,i}}(|\varepsilon_1| + |\varepsilon_2| + |\varepsilon_J|).$$

Step 2 – Show the proximity of $\mathbf{K}_{B,i}$ to a Gaussian vector and simplify the inequality.

Note that, the variance of $\mathbf{K}_{B,i}$ is equal to

$$\frac{4}{\sigma_{B,i}^2 T} \sum_{t=1}^T \text{Var}[(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{i,\cdot}\mathbf{Z}_{\cdot,t}\mathbf{V}_{t,\cdot}(\tilde{\mathbf{B}}_{i,\cdot})^\top] = \frac{4}{\sigma_{B,i}^2 T}(\boldsymbol{\Sigma}_\varepsilon)_{i,i}\tilde{\mathbf{B}}_{i,\cdot}(\tilde{\mathbf{B}}_{i,\cdot})^\top = 1,$$

since $\sigma_{B,i} = \frac{2}{\sqrt{T}}\sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i}}\|\tilde{\mathbf{B}}_{i,\cdot}\|_2$. We will establish the upper bound for the parameter

$$\tau_B^i := \sup_{R \geq 0} |\mathbb{P}(|\mathbf{K}_{B,i}| \leq R) - \mathbb{P}(|\mathcal{N}(0,1)| \leq R)|.$$

Then by the Berry-Esseen theorem, we obtain that

$$\begin{aligned} \tau_B^i &\leq \frac{2\sqrt{2}}{\sigma_{B,i}^3 T^{3/2}} \sum_{t=1}^T \mathbb{E}[\|(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{i,\cdot}\mathbf{Z}_{\cdot,t}\mathbf{V}_{t,\cdot}(\tilde{\mathbf{B}}_{i,\cdot})^\top\|_2^3] \\ &\leq \frac{2\sqrt{2}}{\sigma_{B,i}^3 T^{3/2}} \mathbb{E}[\|(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{i,\cdot}\mathbf{Z}_{\cdot,t}\|_2^3] \max_{1 \leq t \leq T} |\mathbf{V}_{t,\cdot}(\tilde{\mathbf{B}}_{i,\cdot})^\top| \sum_{t=1}^T |\mathbf{V}_{t,\cdot}(\tilde{\mathbf{B}}_{i,\cdot})^\top|_2^2 \\ &= \frac{2\sqrt{2}}{\sigma_{B,i}^3 T^{3/2}} ((\boldsymbol{\Sigma}_\varepsilon)_{i,i})^{3/2} \max_{1 \leq t \leq T} |\mathbf{V}_{t,\cdot}(\tilde{\mathbf{B}}_{i,\cdot})^\top| \sum_{t=1}^T |\mathbf{V}_{t,\cdot}(\tilde{\mathbf{B}}_{i,\cdot})^\top|_2^2 \\ &\lesssim \sqrt{\frac{\log n}{T}}, \end{aligned}$$

where we used the fact that

$$\sum_{t=1}^T |\mathbf{V}_{t,\cdot}(\tilde{\mathbf{B}}_{i,\cdot})^\top|_2^2 = \tilde{\mathbf{B}}_{i,\cdot} \left[\sum_{t=1}^T (\mathbf{V}_{t,\cdot})^\top \mathbf{V}_{t,\cdot} \right] (\tilde{\mathbf{B}}_{i,\cdot})^\top = \tilde{\mathbf{B}}_{i,\cdot} \mathbf{I}_r (\tilde{\mathbf{B}}_{i,\cdot})^\top = \tilde{\mathbf{B}}_{i,\cdot} (\tilde{\mathbf{B}}_{i,\cdot})^\top = \|\tilde{\mathbf{B}}_{i,\cdot}\|_2^2,$$

and

$$\mathbb{E}[\|(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{i,\cdot}\mathbf{Z}_{\cdot,t}\|_2^3] \lesssim \|(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{i,\cdot}\|_2^3 = ((\boldsymbol{\Sigma}_\varepsilon)_{i,i})^{3/2},$$

which follows from the same argument as we proved $\mathbb{E}[\|(\mathbf{a}\mathbf{Z}_{\cdot,t})\|_2^3] \lesssim \|\mathbf{a}\|_2^3$ in the proof of Lemma 13.

Recall that our target is to prove the upper bound of $|\varepsilon_{B,i}|$ and the inequality

$$\left| \mathbb{P}(|\mathbf{K}_{B,i} + \varepsilon_{B,i} + \zeta| \leq z_{1-\frac{1}{2}\alpha}) - (1 - \alpha) \right| \lesssim |\varepsilon_{B,i}| + |\zeta| + \tau_B^i.$$

Note that

$$\mathbb{P}(|\mathbf{K}_{B,i}| \leq z_{1-\frac{1}{2}\alpha} - |\varepsilon_{B,i}| - |\zeta|) < \mathbb{P}(|\mathbf{K}_{B,i} + \varepsilon_{B,i} + \zeta| \leq z_{1-\frac{1}{2}\alpha}) \leq \mathbb{P}(|\mathbf{K}_{B,i}| \leq z_{1-\frac{1}{2}\alpha} + |\varepsilon_{B,i}| + |\zeta|).$$

Then by definition of τ_B^i , we obtain

$$\mathbb{P}(|\mathcal{N}(0, 1)| \leq z_{1-\frac{1}{2}\alpha} - |\varepsilon_{B,i}| - |\zeta|) - \tau_B^i < \mathbb{P}(|\mathbf{K}_{B,i} + \varepsilon_{B,i} + \zeta| \leq z_{1-\frac{1}{2}\alpha}) \leq \mathbb{P}(|\mathcal{N}(0, 1)| \leq z_{1-\frac{1}{2}\alpha} + |\varepsilon_{B,i}| + |\zeta|) + \tau_B^i.$$

Then using the property of Gaussian density, or by Lemma 15, we obtain that, there exists a constant $c_G > 0$ such that

$$\mathbb{P}(|\mathcal{N}(0, 1)| \leq z_{1-\frac{1}{2}\alpha}) - c_G(|\varepsilon_{B,i}| + |\zeta|) - \tau_B^i \leq \mathbb{P}(|\mathbf{K}_{B,i} + \varepsilon_{B,i} + \zeta| \leq z_{1-\frac{1}{2}\alpha}) \leq \mathbb{P}(|\mathcal{N}(0, 1)| \leq z_{1-\frac{1}{2}\alpha}) + c_G(|\varepsilon_{B,i}| + |\zeta|) + \tau_B^i.$$

Then since $\mathbb{P}(|\mathcal{N}(0, 1)| \leq z_{1-\frac{1}{2}\alpha}) = 1 - \alpha$, we obtain that

$$\left| \mathbb{P}(|\mathbf{K}_{B,i} + \varepsilon_{B,i} + \zeta| < \tau_{1-\frac{1}{2}\alpha}) - (1 - \alpha) \right| \lesssim c_G(|\varepsilon_{B,i}| + |\zeta|) + \tau_B^i \lesssim |\varepsilon_{B,i}| + |\zeta| + \tau_B^i.$$

Step 3 – Prove the upper bound for $|\varepsilon_{B,i}|$.

By definition, we have that $|\varepsilon_{B,i}| \leq \frac{1}{\sigma_{B,i}}(|\varepsilon_1| + |\varepsilon_2| + |\varepsilon_J|)$.

For $\varepsilon_1 = (\widehat{\mathbf{U}}\mathbf{R}_U)_{i,\cdot}(\varepsilon_\Sigma\mathbf{\Lambda} + \mathbf{\Lambda}(\varepsilon_\Sigma)^\top + \varepsilon_\Sigma(\varepsilon_\Sigma)^\top)((\widehat{\mathbf{U}}\mathbf{R}_U)_{i,\cdot})^\top$, since $\|\varepsilon_\Sigma\|_2 \lesssim \sigma_r(\rho^2 + \rho\sqrt{\frac{r}{n}}\log n) \lesssim \sigma_r \lesssim \|\mathbf{\Lambda}\|_2$, we have that

$$\begin{aligned} \frac{1}{\sigma_{B,i}}|\varepsilon_1| &\lesssim \frac{\sqrt{T}}{\sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i}}\|\widetilde{\mathbf{B}}_{i,\cdot}\|_2} \cdot \|(\widehat{\mathbf{U}}_{i,\cdot}\mathbf{R}_U - \mathbf{U}_{i,\cdot})\mathbf{\Lambda} + \mathbf{U}_{i,\cdot}\mathbf{\Lambda}\|_2 \cdot \|\varepsilon_\Sigma\|_2 \cdot \|(\widehat{\mathbf{U}}\mathbf{R}_U)_{i,\cdot}\|_2 \\ &\stackrel{(i)}{\lesssim} \frac{\sqrt{T}}{\sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i}}\|\widetilde{\mathbf{U}}_{i,\cdot}\|_2\sigma_r} \cdot (\omega_i\sqrt{\frac{r}{n}}\log n \cdot \sigma_r) \cdot \sigma_r(\rho^2 + \rho\sqrt{\frac{r}{n}}\log n) \cdot \sqrt{r} \\ &\quad + \frac{\sqrt{T}}{\sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i}}\|\widetilde{\mathbf{B}}_{i,\cdot}\|_2} \|\mathbf{U}_{i,\cdot}\mathbf{\Lambda}\|_2\sigma_r(\rho^2 + \rho\sqrt{\frac{r}{n}}\log n) \cdot \|(\widehat{\mathbf{U}}_{i,\cdot}\mathbf{R}_U - \mathbf{U}_{i,\cdot}) + \mathbf{U}_{i,\cdot}\|_2 \\ &\stackrel{(ii)}{\lesssim} \frac{\sqrt{T}}{\sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i}}\|\widetilde{\mathbf{U}}_{i,\cdot}\|_2\sigma_r} \cdot (\omega_i\sqrt{\frac{r}{n}}\log n \cdot \sigma_r) \cdot \sigma_r(\rho^2 + \rho\sqrt{\frac{r}{n}}\log n)\sqrt{r} \\ &\quad + \frac{\sqrt{T}}{\sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i}}} \sigma_r(\rho^2 + \rho\sqrt{\frac{r}{n}}\log n) \cdot (\omega_i\sqrt{\frac{r}{n}}\log n + \|\widetilde{\mathbf{U}}_{i,\cdot}\|_2), \end{aligned}$$

where (i) uses the fact that $\|\widetilde{\mathbf{B}}_{i,\cdot}\|_2 = \|\mathbf{U}_{i,\cdot}\mathbf{\Lambda}\|_2 \gtrsim \|\mathbf{U}_{i,\cdot}\|_2\sigma_r$ and the bound $\|(\widehat{\mathbf{U}}_{i,\cdot}\mathbf{R}_U - \mathbf{U}_{i,\cdot})\mathbf{\Lambda}\|_2 \lesssim \omega_i\sqrt{\frac{r}{n}}\log n \cdot \sigma_r$ obtained by the decomposition of $(\widehat{\mathbf{U}}_{i,\cdot}\mathbf{R}_U - \mathbf{U}_{i,\cdot})$ as we did in proof of Corollary 1, and (ii) uses the fact that $\|\mathbf{U}_{i,\cdot}\|_2 = \|\widetilde{\mathbf{U}}_{i,\cdot}\|_2$.

For $\varepsilon_2 = \mathbf{U}_{i,\cdot}\mathbf{\Lambda}^2((\boldsymbol{\Psi}_U)_{i,\cdot})^\top + (\boldsymbol{\Psi}_U)_{i,\cdot}\mathbf{\Lambda}^2(\mathbf{U}_{i,\cdot})^\top + (\mathbf{G}_U + \boldsymbol{\Psi}_U)_{i,\cdot}(\mathbf{\Lambda}^2)((\mathbf{G}_U + \boldsymbol{\Psi}_U)_{i,\cdot})^\top$, we have that

$$\begin{aligned} \frac{1}{\sigma_{B,i}}|\varepsilon_2| &\lesssim \frac{\sqrt{T}}{2\sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i}}\|\widetilde{\mathbf{B}}_{i,\cdot}\|_2} (\|\widetilde{\mathbf{B}}_{i,\cdot}\|_2\|(\boldsymbol{\Psi}_U)_{i,\cdot}\mathbf{\Lambda}\|_2 + \|(\widehat{\mathbf{U}}_{i,\cdot}\mathbf{R}_U - \mathbf{U}_{i,\cdot})\mathbf{\Lambda}\|_2^2) \\ &\lesssim \frac{\sqrt{T}\sigma_r}{\sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i}}} \left(\omega_i\omega\sqrt{\frac{r}{n}}\log^{3/2}n + \left(\rho^2 + \rho\sqrt{\frac{r}{n}}\log n \right) \|\widetilde{\mathbf{U}}_{i,\cdot}\|_2 + \omega_i\omega\log n\|\bar{\mathbf{U}}\|_{2,\infty} \right) \\ &\quad + \frac{\sqrt{T}(\omega_i\sqrt{\frac{r}{n}}\log n)^2\sigma_r}{\sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i}}\|\widetilde{\mathbf{U}}_{i,\cdot}\|_2}, \end{aligned}$$

where we use the decomposition and the upper bound for $(\boldsymbol{\Psi}_U)_{i,\cdot}$ in the proof of Theorem 1, and the bound $\|(\widehat{\mathbf{U}}_{i,\cdot}\mathbf{R}_U - \mathbf{U}_{i,\cdot})\mathbf{\Lambda}\|_2 \lesssim \omega_i\sqrt{\frac{r}{n}}\log n \cdot \sigma_r$ obtained by the decomposition of $(\widehat{\mathbf{U}}_{i,\cdot}\mathbf{R}_U - \mathbf{U}_{i,\cdot})$ as we did in proof of Corollary 1.

For $\varepsilon_J = \mathbf{B}_{i,\cdot}((\mathbf{J}^{-1})^\top\mathbf{J}^{-1} - \mathbf{I}_r)(\mathbf{B}_{i,\cdot})^\top$, we have that

$$\frac{1}{\sigma_{B,i}}|\varepsilon_J| \lesssim \frac{\sqrt{T}}{2\sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i}}\|\mathbf{B}_{i,\cdot}\|_2} \cdot \|\mathbf{B}_{i,\cdot}\|_2^2 \sqrt{\frac{r + \log n}{T}} \lesssim \frac{\|\mathbf{b}_i\|_2}{\sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i}}} \sqrt{r + \log n}.$$

Step 4 – Prove the upper bound for $|\zeta|$ and assemble all the bounds to get the final result.

We have that

$$\zeta = (((\widehat{\boldsymbol{\Sigma}}_\varepsilon^\tau)_{i,i})^{-1/2} - ((\boldsymbol{\Sigma}_\varepsilon)_{i,i})^{-1/2}) \frac{\sqrt{T}}{2\|\widetilde{\mathbf{B}}_{i,\cdot}\|_2} (\|\widetilde{\mathbf{B}}_{i,\cdot}\|_2^2 - \|(\widehat{\mathbf{U}}\widehat{\boldsymbol{\Sigma}})_{i,\cdot}\|_2^2).$$

Note that $|((\widehat{\boldsymbol{\Sigma}}_\varepsilon^\tau)_{i,i} - (\boldsymbol{\Sigma}_\varepsilon)_{i,i})| \lesssim \epsilon_{N,T}(\boldsymbol{\Sigma}_\varepsilon)_{i,i}$ and $\epsilon_{N,T} \ll 1$, so we obtain by the mean value theorem that

$$|((\widehat{\boldsymbol{\Sigma}}_\varepsilon^\tau)_{i,i})^{-1/2} - ((\boldsymbol{\Sigma}_\varepsilon)_{i,i})^{-1/2}| \lesssim ((\boldsymbol{\Sigma}_\varepsilon)_{i,i})^{-3/2} \cdot \epsilon_{N,T}(\boldsymbol{\Sigma}_\varepsilon)_{i,i} = ((\boldsymbol{\Sigma}_\varepsilon)_{i,i})^{-1/2} \cdot \epsilon_{N,T}.$$

So we obtain that

$$\begin{aligned} |\zeta| &\lesssim \epsilon_{N,T} \cdot \frac{\sqrt{T}}{\sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i}}\|\widetilde{\mathbf{B}}_{i,\cdot}\|_2} \|\widetilde{\mathbf{B}}_{i,\cdot}\|_2^2 - \|(\widehat{\mathbf{U}}\widehat{\boldsymbol{\Sigma}})_{i,\cdot}\|_2^2 \\ &\lesssim \epsilon_{N,T} \cdot \frac{1}{\sigma_{B,i}} \mathbf{K}_{B,i} + \epsilon_{N,T} \cdot \frac{1}{\sigma_{B,i}} (|\varepsilon_1| + |\varepsilon_2| + |\varepsilon_J|). \end{aligned}$$

Using (G.3) in Lemma 19 of [Yan and Wainwright \(2024\)](#), the leading order term is bounded by

$$\begin{aligned} \epsilon_{N,T} \cdot \frac{1}{\sigma_{B,i}} \mathbf{K}_{B,i} &\lesssim \epsilon_{N,T} \cdot \frac{\sqrt{T}}{\sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i}}\|\widetilde{\mathbf{B}}_{i,\cdot}\|_2} \frac{1}{\sqrt{T}} \|(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{i,\cdot}\|_2 \|\widetilde{\mathbf{B}}_{i,\cdot}\|_2 \sqrt{r} \sqrt{1+r+\log n} \\ &\lesssim \epsilon_{N,T} \cdot r \sqrt{\log n}, \end{aligned}$$

where we use the fact that $\|(\boldsymbol{\Sigma}_\varepsilon^{1/2})_{i,\cdot}\|_2^2 = (\boldsymbol{\Sigma}_\varepsilon)_{i,i}$.

Combining the above bounds, we obtain that

$$\left| \mathbb{P}\left(\frac{1}{\widehat{\sigma}_{B,i}} (\|\widetilde{\mathbf{B}}_{i,\cdot}\|_2^2 - \|(\widehat{\mathbf{U}}\widehat{\boldsymbol{\Sigma}})_{i,\cdot}\|_2^2) \in [-z_{1-\frac{1}{2}\alpha}, z_{1-\frac{1}{2}\alpha}] \right) - (1-\alpha) \right| \lesssim |\varepsilon_{B,i}| + |\zeta| + \tau_B^i,$$

where $\tau_B^i \lesssim \sqrt{\frac{\log n}{T}}$, and since $\epsilon_{N,T} \ll 1$, we have that

$$\begin{aligned} |\varepsilon_{B,i}| + |\zeta| &\lesssim \frac{1}{\sigma_{B,i}} (|\varepsilon_1| + |\varepsilon_2| + |\varepsilon_J|) + \epsilon_{N,T} \cdot \frac{1}{\sigma_{B,i}} \mathbf{K}_{B,i} + \epsilon_{N,T} \cdot \frac{1}{\sigma_{B,i}} (|\varepsilon_1| + |\varepsilon_2| + |\varepsilon_J|) \\ &\lesssim \frac{1}{\sigma_{B,i}} (|\varepsilon_1| + |\varepsilon_2| + |\varepsilon_J|) + \epsilon_{N,T} \cdot \frac{1}{\sigma_{B,i}} \mathbf{K}_{B,i} \\ &\lesssim \frac{\sqrt{T}\sigma_r}{\sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i}}} \left(\omega_i \omega \sqrt{\frac{r}{n}} \log^{3/2} n + \left(\rho^2 + \rho \sqrt{\frac{r}{n}} \log n \right) \|\bar{\mathbf{U}}_{i,\cdot}\|_2 + \omega_i \omega \log n \|\bar{\mathbf{U}}\|_{2,\infty} \right) \\ &\quad + \frac{\sqrt{T}(\omega_i \sqrt{\frac{r}{n}} \log n)^2 \sigma_r}{\sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i}} \|\bar{\mathbf{U}}_{i,\cdot}\|_2} + \frac{\sqrt{T}}{\sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i}} \|\bar{\mathbf{U}}_{i,\cdot}\|_2 \sigma_r} \cdot \omega_i \sqrt{\frac{r}{n}} \log n \cdot \sigma_r \cdot \sigma_r (\rho^2 + \rho \sqrt{\frac{r}{n}} \log n) \sqrt{r} \\ &\quad + \frac{\sqrt{T}}{\sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i}}} \sigma_r (\rho^2 + \rho \sqrt{\frac{r}{n}} \log n) \cdot (\omega_i \sqrt{\frac{r}{n}} \log n + \|\bar{\mathbf{U}}_{i,\cdot}\|_2) + \frac{\|\mathbf{b}_i\|_2}{\sqrt{(\boldsymbol{\Sigma}_\varepsilon)_{i,i}}} \sqrt{r + \log n} + \epsilon_{N,T} \cdot r \sqrt{\log n}. \end{aligned}$$

The desired result follows from the above bound.

H Technical lemmas

We collect some technical lemmas in this section. First, we establish the results on the quantile of Chi-square distribution. Recall that, we denote $\chi^2(n)$ as a random variable follows Chi-square distribution with degree of freedom n , and denote by $\chi_q^2(n)$ its q -quantile.

Lemma 14. *For any fixed $q \in (0, 1)$, the q -quantile of Chi-square distribution satisfies that $\chi_q^2(n) \asymp n$.*

Proof. Consider n i.i.d. standard Gaussian $\mathcal{N}(0, 1)$ random variables X_1, X_2, \dots, X_n . Define $S_n := \sum_{i=1}^n X_i^2$, which follows $\chi^2(n)$ distribution. Using the Berry-Esseen theorem (Vershynin, 2017, Theorem 2.1.3), we have that

$$\sup_{t \in \mathbb{R}} \left| \mathbb{P}\left(\frac{S_n - n}{\sqrt{2n}} \leq t\right) - \int_{-\infty}^t \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x^2} dx \right| < c \frac{1}{\sqrt{n}},$$

where c is a constant independent with n . We denote by z_q the q -quantile of standard Gaussian $\mathcal{N}(0, 1)$. Letting $t = z_q$ in the above inequality, we obtain that

$$q - c \frac{1}{\sqrt{n}} < \mathbb{P}(S_n \leq n + z_q \sqrt{2n}) < q + c \frac{1}{\sqrt{n}}.$$

Note that $\chi_q^2(n)$ is the q -quantile of S_n . By the monotonicity of cdf, we obtain that

$$n + z_{q-c\frac{1}{\sqrt{n}}} \sqrt{2n} \leq \chi_q^2(n) \leq n + z_{q+c\frac{1}{\sqrt{n}}} \sqrt{2n}.$$

Then, we prove that $z_{q-c\frac{1}{\sqrt{n}}}$ and $z_{q+c\frac{1}{\sqrt{n}}}$ are close to z_q .

On the one hand, when $c\frac{1}{\sqrt{n}} < \frac{1-q}{2}$, i.e., $n > \frac{4c^2}{(1-q)^2}$, we have that

$$\begin{aligned} c \frac{1}{\sqrt{n}} &= \left(q + c \frac{1}{\sqrt{n}}\right) - q = \int_{z_q}^{z_{q+c\frac{1}{\sqrt{n}}}} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x^2} dx \\ &\stackrel{(i)}{=} (z_{q+c\frac{1}{\sqrt{n}}} - z_q) \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x^2} \Big|_{x=\vartheta z_{q+c\frac{1}{\sqrt{n}}} + (1-\vartheta)z_q} \\ &\geq (z_{q+c\frac{1}{\sqrt{n}}} - z_q) \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x^2} \Big|_{x=z_{(1+q)/2}}, \end{aligned}$$

where (i) uses the mean value theorem for integral. On the other hand, similarly we have that

$$c \frac{1}{\sqrt{n}} \geq (z_q - z_{q-c\frac{1}{\sqrt{n}}}) \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x^2} \Big|_{x=z_q}.$$

So, there exists a constant c_1 which does not depend on n , such that $\max(z_{q+c\frac{1}{\sqrt{n}}} - z_q, z_q - z_{q-c\frac{1}{\sqrt{n}}}) \leq c_1 \frac{1}{\sqrt{n}}$ holds for sufficiently large n . As a result, we obtain

$$n + z_q \sqrt{2n} - \sqrt{2}c_1 = n + (z_q - c_1 \frac{1}{\sqrt{n}}) \sqrt{2n} \leq \chi_q^2(n) \leq n + (z_q + c_1 \frac{1}{\sqrt{n}}) \sqrt{2n} \leq n + z_q \sqrt{2n} + \sqrt{2}c_1.$$

For simplicity of notations, we write the above results for $\chi_q^2(n)$ as follows: for sufficiently large n , it holds

$$|\chi_q^2(n) - (n + C_1 \sqrt{n})| < C_0,$$

where $C_1, C_0 > 0$ are constants independent with n . So we obtain that $\chi_q^2(n) \asymp n$. \square

Next, we consider another results on Chi-square distribution, which can be regarded as the probability that a Gaussian vector lies in a spherical shell.

Lemma 15. *There exists a constant c_B which does not depend on the dimension m , such that, for any ε satisfying $|\varepsilon| < R$, it holds*

$$\sup_{R \geq 0} |\mathbb{P}(\chi^2(m) \leq (R + \varepsilon)^2) - \mathbb{P}(\chi^2(m) \leq R^2)| \leq c_B |\varepsilon|.$$

Proof. Denote by $B_m(R)$ the ball in \mathbb{R}^m centered at the origin with the radius R . By (2.3) in Sazonov (1972) and the comments after it, we obtain that $|\mathbb{P}(\mathcal{N}(0, \mathbf{I}_m) \in B_m(R + \varepsilon)) - \mathbb{P}(\mathcal{N}(0, \mathbf{I}_m) \in B_m(R))| \leq c_B |\varepsilon|$, where $c_B > 0$ is a constant and c_B does not depend on m , R , and ε . So the desired result follows from the fact that $\mathbb{P}(\chi^2(m) \leq R^2) = \mathbb{P}(\mathcal{N}(0, \mathbf{I}_m) \in B(R))$. \square

Then we consider a special case of multi-dimensional Berry-Esseen theorem.

Lemma 16. Let $\xi_1, \xi_2, \dots, \xi_n$ be independent d -dimensional random vectors with zero means, and let $W = \sum_{k=1}^n \xi_k$. Suppose that $\text{cov}(W) = \mathbf{I}_d$. Then we have that

$$\sup_{R \geq 0} |\mathbb{P}(\|W\|_2 \leq R) - \mathbb{P}(\chi^2(d) \leq R^2)| \leq c_G \sum_{k=1}^n \mathbb{E} \left[\|\xi_k\|_2^3 \right],$$

where c_G is a constant and c_G does not depend on n and d .

Proof. The proof is based on the results in Raič (2019) and we will directly adopt the notations therein to conduct the proof. Note that $\mathbb{P}(\chi^2(d) \leq R^2) = \mathbb{P}(\mathcal{N}(0, \mathbf{I}_d) \in B_d(R))$, where $B_d(R)$ is the ball in \mathbb{R}^d centered at the origin with the radius R . According to (1.4) in Theorem 1.3 and Example 1.2, we have that

$$\begin{aligned} \sup_{R \geq 0} |\mathbb{P}(\|W\|_2 \leq R) - \mathbb{P}(\chi^2(d) \leq R^2)| &= \sup_{R \geq 0} |\mathbb{P}(W \in B_d(R)) - \mathbb{P}(\mathcal{N}(0, \mathbf{I}_d) \in B_d(R))| \\ &\lesssim (1 + \gamma^*(\mathcal{A}|\rho)) \sum_{k=1}^n \mathbb{E} \left[\|\xi_k\|_2^3 \right]. \end{aligned} \quad (\text{H.1})$$

Here, for $\gamma^*(\mathcal{A}|\rho)$ defined in Raič (2019), following Example 1.2 therein for the class of all balls, \mathcal{A} is the set of all balls in \mathbb{R}^d , and $\rho_A = \delta_A$. Then we obtain $A^{t|\rho} = \{x; \rho_A(x) \leq t\} = \{x; \delta_A(x) \leq t\} = A^t$, and thus by definition we get $\gamma^*(\mathcal{A}|\rho) = \gamma^*(\mathcal{A})$. Since \mathcal{A} is the set of all balls in \mathbb{R}^d , we obtain by definition of $\gamma^*(\mathcal{A})$ that

$$\gamma^*(\mathcal{A}) = \sup_{v \in \mathbb{R}^d, R \geq 0, \varepsilon \neq 0} \frac{1}{|\varepsilon|} |\mathbb{P}(\mathcal{N}(0, \mathbf{I}_d) \in B_d(R + \varepsilon; v)) - \mathbb{P}(\mathcal{N}(0, \mathbf{I}_d) \in B_d(R; v))|,$$

where $B_d(R; v)$ is the ball in \mathbb{R}^d centered at v with the radius R . By (2.3) in Sazonov (1972) and the comments after it, we obtain that, there exists a constant c_B which does not depend on the dimension d , such that $\gamma^*(\mathcal{A}) \leq c_B$. Then the desired result follows from (H.1) and the fact that $\gamma^*(\mathcal{A}|\rho) = \gamma^*(\mathcal{A}) \leq c_B$. \square

References

- Abbe, E., Fan, J., and Wang, K. (2022). An ℓ_p theory of PCA and spectral clustering. *The Annals of Statistics*, 50(4):2359 – 2385.
- Abbe, E., Fan, J., Wang, K., Zhong, Y., et al. (2020). Entrywise eigenvector analysis of random matrices with low expected rank. *Annals of Statistics*, 48(3):1452–1474.
- Bai, J. (2003). Inferential theory for factor models of large dimensions. *Econometrica*, 71(1):135–171.
- Bai, J. and Ng, S. (2002). Determining the number of factors in approximate factor models. *Econometrica*, 70(1):191–221.
- Bai, J. and Ng, S. (2006a). Confidence intervals for diffusion index forecasts and inference for factor-augmented regressions. *Econometrica*, 74(4):1133–1150.
- Bai, J. and Ng, S. (2006b). Evaluating latent and observed factors in macroeconomics and finance. *Journal of Econometrics*, 131(1-2):507–537.
- Bai, J. and Ng, S. (2013). Principal components estimation and identification of static factors. *Journal of econometrics*, 176(1):18–29.
- Bai, J. and Ng, S. (2023). Approximate factor models with weaker loadings. *Journal of Econometrics*.
- Bai, J. and Shi, S. (2011). Estimating high dimensional covariance matrices and its applications. *Annals of Economics and Finance*, 12(2):199–215.
- Bai, J. and Wang, P. (2016). Econometric analysis of large factor models. *Annual Review of Economics*, 8:53–80.

- Bailey, N., Kapetanios, G., and Pesaran, M. H. (2021). Measurement of factor strength: Theory and practice. *Journal of Applied Econometrics*, 36(5):587–613.
- Bickel, P. J. and Levina, E. (2008). Covariance regularization by thresholding. *The Annals of Statistics*, 36(6):2577–2604.
- Braun, M. L. (2006). Accurate error bounds for the eigenvalues of the kernel matrix. *The Journal of Machine Learning Research*, 7:2303–2328.
- Breitung, J. and Eickmeier, S. (2011). Testing for structural breaks in dynamic factor models. *Journal of Econometrics*, 163(1):71–84.
- Cai, C., Li, G., Chi, Y., Poor, H. V., and Chen, Y. (2021). Subspace estimation from unbalanced and incomplete data matrices: $\ell_{2,\infty}$ statistical guarantees. *The Annals of Statistics*, 49(2):944–967.
- Chamberlain, G. and Rothschild, M. (1983). Arbitrage, factor structure, and mean-variance analysis on large asset markets. *Econometrica*, pages 1281–1304.
- Chen, L., Dolado, J. J., and Gonzalo, J. (2014). Detecting big structural breaks in large factor models. *Journal of Econometrics*, 180(1):30–48.
- Chen, Y., Chi, Y., Fan, J., Ma, C., et al. (2021a). Spectral methods for data science: A statistical perspective. *Foundations and Trends® in Machine Learning*, 14(5):566–806.
- Chen, Y., Chi, Y., Fan, J., Ma, C., and Yan, Y. (2020). Noisy matrix completion: Understanding statistical guarantees for convex relaxation via nonconvex optimization. *SIAM Journal on Optimization*, 30(4):3098–3121.
- Chen, Y., Fan, J., Ma, C., and Yan, Y. (2019). Inference and uncertainty quantification for noisy matrix completion. *Proceedings of the National Academy of Sciences*, 116(46):22931–22937.
- Chen, Y., Fan, J., Ma, C., and Yan, Y. (2021b). Bridging convex and nonconvex optimization in robust pca: Noise, outliers and missing data. *The Annals of Statistics*, 49(5):2948–2971.
- Chen, Y., Fan, J., Wang, B., and Yan, Y. (2023). Convex and nonconvex optimization are both minimax-optimal for noisy blind deconvolution under random designs. *Journal of the American Statistical Association*, 118(542):858–868.
- Chen, Y. M., Chen, X. S., and Li, W. (2016). On perturbation bounds for orthogonal projections. *Numerical Algorithms*, 73(2):433–444.
- Chernozhukov, V., Hansen, C., Liao, Y., and Zhu, Y. (2023). Inference for low-rank models. *The Annals of statistics*, 51(3):1309–1330.
- Choi, J., Kwon, H., and Liao, Y. (2023). Inference for low-rank models without estimating the rank. *arXiv preprint arXiv:2311.16440*.
- Choi, J., Kwon, H., and Liao, Y. (2024). Inference for low-rank completion without sample splitting with application to treatment effect estimation. *Journal of Econometrics*, 240(1):105682.
- Choi, J. and Yuan, M. (2024). High dimensional factor analysis with weak factors. *arXiv preprint arXiv:2402.05789*.
- El Karoui, N. (2015). On the impact of predictor geometry on the performance on high-dimensional ridge-regularized generalized robust regression estimators. *Probability Theory and Related Fields*, pages 1–81.
- Fama, E. F. and French, K. R. (1993). Common risk factors in the returns on stocks and bonds. *Journal of financial economics*, 33(1):3–56.
- Fan, J., Fan, Y., and Lv, J. (2008). High dimensional covariance matrix estimation using a factor model. *Journal of Econometrics*, 147(1):186–197.

- Fan, J., Li, K., and Liao, Y. (2021a). Recent developments in factor models and applications in econometric learning. *Annual Review of Financial Economics*, 13:401–430.
- Fan, J., Li, K., and Liao, Y. (2021b). Recent developments on factor models and its applications in econometric learning. *Annual Review of Financial Economics*, 13:to appear.
- Fan, J. and Li, R. (2001). Variable selection via nonconcave penalized likelihood and its oracle properties. *Journal of the American statistical Association*, 96(456):1348–1360.
- Fan, J., Liao, Y., and Mincheva, M. (2011). High dimensional covariance matrix estimation in approximate factor models. *Annals of statistics*, 39(6):3320.
- Fan, J., Liao, Y., and Mincheva, M. (2013). Large covariance estimation by thresholding principal orthogonal complements. *Journal of the Royal Statistical Society. Series B, Statistical methodology*, 75(4).
- Fan, J., Liao, Y., and Yao, J. (2015). Power enhancement in high-dimensional cross-sectional tests. *Econometrica*, 83(4):1497–1541.
- Fan, J., Wang, K., Zhong, Y., and Zhu, Z. (2021c). Robust high-dimensional factor models with applications to statistical machine learning. *Statistical Science*, 36(2):303–327.
- Forni, M., Hallin, M., Lippi, M., and Reichlin, L. (2000). The generalized dynamic-factor model: Identification and estimation. *Review of Economics and statistics*, 82(4):540–554.
- Freyaldenhoven, S. (2022). Factor models with local factors — determining the number of relevant factors. *Journal of Econometrics*, 229(1):80–102.
- Gagliardini, P., Ossola, E., and Scaillet, O. (2016). Time-varying risk premium in large cross-sectional equity data sets. *Econometrica*, 84(3):985–1046.
- Giglio, S. and Xiu, D. (2021). Asset pricing with omitted factors. *Journal of Political Economy*, 129(7):1947–1990.
- Giglio, S., Xiu, D., and Zhang, D. (2021). Test assets and weak factors. Technical report, National Bureau of Economic Research.
- Giglio, S., Xiu, D., and Zhang, D. (2023). Prediction when factors are weak. *University of Chicago, Becker Friedman Institute for Economics Working Paper*, (2023-47).
- Han, X. and Inoue, A. (2015). Tests for parameter instability in dynamic factor models. *Econometric Theory*, 31(5):1117–1152.
- Jiang, P., Uematsu, Y., and Yamagata, T. (2023). Revisiting asymptotic theory for principal component estimators of approximate factor models. *arXiv preprint arXiv:2311.00625*.
- Ma, C., Wang, K., Chi, Y., and Chen, Y. (2020). Implicit regularization in nonconvex statistical estimation: Gradient descent converges linearly for phase retrieval, matrix completion, and blind deconvolution. *Foundations of Computational Mathematics*, 20(3):451–632.
- Onatski, A. (2009). Testing hypotheses about the number of factors in large factor models. *Econometrica*, 77(5):1447–1479.
- Onatski, A. (2010). Determining the number of factors from empirical distribution of eigenvalues. *The Review of Economics and Statistics*, 92(4):1004–1016.
- Onatski, A. (2012). Asymptotics of the principal components estimator of large factor models with weakly influential factors. *Journal of Econometrics*, 168(2):244–258.
- Raič, M. (2019). A multivariate Berry-Esseen theorem with explicit constants. *Bernoulli*, 25(4A):2824–2853.
- Ross, S. A. (1976). The arbitrage theory of capital asset pricing. *Journal of Economic Theory*, 13(3):341–360.

- Sazonov, V. (1972). On a bound for the rate of convergence in the multidimensional central limit theorem. In *Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability, Volume 2: Probability Theory*, volume 6, pages 563–582. University of California Press.
- Stewart, G. W. (1977). On the perturbation of pseudo-inverses, projections and linear least squares problems. *SIAM review*, 19(4):634–662.
- Stock, J. H. and Watson, M. (2009). Forecasting in dynamic factor models subject to structural instability. *The Methodology and Practice of Econometrics. A Festschrift in Honour of David F. Hendry*, 173:205.
- Stock, J. H. and Watson, M. W. (2002). Forecasting using principal components from a large number of predictors. *Journal of the American statistical association*, 97(460):1167–1179.
- Tropp, J. A. (2012). User-friendly tail bounds for sums of random matrices. *Found. Comput. Math.*, 12(4):389–434.
- Uematsu, Y. and Yamagata, T. (2022a). Estimation of sparsity-induced weak factor models. *Journal of Business & Economic Statistics*, 41(1):213–227.
- Uematsu, Y. and Yamagata, T. (2022b). Inference in sparsity-induced weak factor models. *Journal of Business & Economic Statistics*, 41(1):126–139.
- Vershynin, R. (2017). *High Dimensional Probability*. Cambridge University Press.
- Wainwright, M. (2019). *High-Dimensional Statistics: A Non-Asymptotic Viewpoint*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press.
- Wei, J. and Zhang, Y. (2023). Does principal component analysis preserve the sparsity in sparse weak factor models? *arXiv preprint arXiv:2305.05934*.
- Xia, D. and Yuan, M. (2021). Statistical inferences of linear forms for noisy matrix completion. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 83(1):58–77.
- Yan, Y., Chen, Y., and Fan, J. (2024). Inference for heteroskedastic pca with missing data. *The Annals of Statistics*, 52(2):729–756.
- Yan, Y. and Wainwright, M. J. (2024). Entrywise inference for causal panel data: A simple and instance-optimal approach. *arXiv preprint arXiv:2401.13665*.
- Zhou, Y. and Chen, Y. (2023). Deflated heteropca: Overcoming the curse of ill-conditioning in heteroskedastic pca. *arXiv preprint arXiv:2303.06198*.