

UNIVERSAL PIECEWISE POLYNOMIALITY FOR COUNTING CURVES IN TORIC SURFACES

MARVIN ANAS HAHN AND VINCENZO REDA

ABSTRACT. Inspired by piecewise polynomiality results of double Hurwitz numbers, Ardila and Brugallé introduced an enumerative problem which they call *double Gromov–Witten invariants of Hirzebruch surfaces* in [AB17]. These invariants serve as a two-dimensional analogue and satisfy a similar piecewise polynomial structure. More precisely, they introduced the enumeration of curves in Hirzebruch surfaces satisfying point conditions and tangency conditions on the two parallel toric boundaries. These conditions are stored in four partitions and the resulting invariants are piecewise polynomial in their entries. Moreover, they found that these expressions also behave polynomially with respect to the parameter determining the underlying Hirzebruch surfaces. Based on work of Ardila and Block [AB13], they proposed that such a polynomiality could also hold while changing between more general toric surfaces corresponding to h -transverse polygons. In this work, we answer this question affirmatively. Moreover, we express the resulting invariants for h -transverse polygons as matrix elements in the two-dimensional bosonic Fock space.

CONTENTS

1. Introduction	1
1.1. Double Gromov–Witten invariants for h -transverse polygons	2
1.2. Results	4
2. Preliminaries	5
2.1. Floor diagrams	5
2.2. Ehrhart theory	10
3. Proofs on Theorems 1.7, 1.8 and 1.10	12
3.1. Proof of Theorems 1.7 and 1.10	12
3.2. Proof of Theorem 1.8	15
3.3. Example	15
4. Double Gromov–Witten invariants via the bosonic Fock space	20
References	23

1. INTRODUCTION

Double Gromov–Witten invariants of Hirzebruch surfaces were introduced by Ardila and Brugallé in [AB17] as a two-dimensional generalisation of so-called *double Hurwitz numbers*. While double Hurwitz numbers count branched morphisms to the Riemann sphere with two relative conditions describing the ramification data, this two-dimensional analogue enumerates curves in a Hirzebruch surface passing through a certain number of points and having prescribed tangency order at two distinguished toric boundary divisors .

2020 *Mathematics Subject Classification.* 14N10,14T90,14N35.

Key words and phrases. Floor diagrams, Gromov–Witten invariants, bosonic Fock space, tropical geometry.

Acknowledgements. We thank Hannah Markwig for many valuable discussions.

Double Hurwitz numbers admit a piecewise polynomial structure and tropical geometry has proved to be a powerful framework in the study of the polynomiality of Hurwitz numbers. [CJM10; CJM11; HL22; HL20; HM22; FHK23] A key idea in these tropical approaches is to express Hurwitz numbers as a weighted enumeration of abstract graphs. For the enumeration of curves in Hirzebruch surfaces, **Floor diagrams** provide a two-dimensional analogue. Motivated by this and by storing the tangency orders in partitions, Ardila and Brugallé used tropical geometry to show that double Gromov–Witten invariants of Hirzebruch surfaces admit a piecewise polynomial behaviour in the entries of those partitions that closely mirrors the piecewise polynomiality of double Hurwitz numbers in the entries of the ramification profiles. Moreover, they observed that the piecewise polynomiality extends when moving between different Hirzebruch surfaces. To be precise, recall that Hirzebruch surfaces \mathbb{F}_k live in a one-dimensional family indexed by natural numbers $k \geq 0$. The polynomials describing double Gromov–Witten invariants of Hirzebruch surfaces are also polynomial in the parameter k . Based on this observation, Ardila and Brugallé asked in [AB17, Section 7] whether this polynomial interpolation between Hirzebruch surfaces extends to enumerations of curves in more complicated toric surfaces – an analogous statement for Severi degrees of toric surfaces was derived in [AB13; LO18]. In this work, we answer this question affirmatively for the important family of toric surfaces corresponding to *h-transverse polygons*.

Definition 1.1. A polygon P is said to be *h-transverse* if every vertex has integer coordinates and every edge has slope 0 , ∞ or $\frac{1}{k}$, with $k \in \mathbb{Z}$.

An important feature of *h-transverse polygons* is their enumerative geometry may be studied via so-called *floor diagram* in tropical geometry. Floor diagrams are an important tool in tropical geometry that were introduced in [BM09; BM16]. For curves in toric surfaces corresponding to *h-transverse polygons*, they allow to translate the enumerative problem in terms of a weighted count of abstract decorated graphs. This perspective was also taken in [AB17] to study double Gromov–Witten invariants of Hirzebruch surfaces. In this work, we take the same approach. A detailed discussion of floor diagrams may be found in Section 2.1.



1.1. Double Gromov–Witten invariants for *h-transverse polygons*. We now set-up our counting problem of *double Gromov–Witten invariants for h-transverse polygons*. To begin with, we parametrise *h-transverse polygons*.

Let P be an *h-transverse polygon*. We will always assume that P has two edges of slope 0, a top edge and a bottom edge. We denote by $d^t > 0$ the lattice length of the top edge and $d^b > 0$ the lattice length of the bottom edge. Moreover, we record the direction of the edges on the left of P by $(c_1^l, -1), \dots, (c_m^l, -1)$ in counterclockwise order. Similarly, we record the direction of the edges on the right of P by $(c_1^r, -1), \dots, (c_n^r, -1)$ in a clockwise manner. Note that c_i^l and c_j^r are integers. Moreover, we have $c_1^l < \dots < c_m^l$ and $c_1^r > \dots > c_n^r$. In addition, we denote the lattice length of c_i^l by d_i^l and the lattice length of c_j^r by d_j^r . We define partitions $\mathbf{c}^l = (c_1^l, \dots, c_m^l)$, $\mathbf{c}^r = (c_1^r, \dots, c_n^r)$, $\mathbf{d}^l = (d_1^l, \dots, d_m^l)$ and $\mathbf{d}^r = (d_1^r, \dots, d_n^r)$. Denoting $\mathbf{c} = (\mathbf{c}^r, \mathbf{c}^l)$ and $\mathbf{d} = (\mathbf{d}^r, \mathbf{d}^l)$, we see that \mathbf{c} and \mathbf{d} completely determine the *h-transverse polygon* P and we will denote $P = P(\mathbf{c}, \mathbf{d})$. Moreover, we denote the toric surface corresponding to $P(\mathbf{c}, \mathbf{d})$ by $S(\mathbf{c})$.

Remark 1.2. Since the normal fan of P is balanced, we obtain the following relations:

$$d^t + \sum_{i=1}^n c_i^r d_i^r - d^b - \sum_{j=1}^m c_j^l d_j^l = 0$$

$$\sum_{i=1}^n d_i^r = \sum_{j=1}^m d_j^l.$$


 FIGURE 1. The polygon corresponding to the Hirzebruch surface \mathbb{F}_2 .

 FIGURE 2. The h -transverse polygon associated to the data in Example 1.4.

Furthermore, it follows from the definition of h -transverse polygons that the vector \mathbf{c} has integer coordinates.

Example 1.3. An important class of h -transverse polygons are the polygons corresponding to Hirzebruch surfaces, that are obtained by this construction by setting $n = m = 1$, $c^r = k \geq 0$ and $c^l = 0$, see Figure 1

Example 1.4. For $\mathbf{c} = (\mathbf{c}^r; \mathbf{c}^l) = (3, 1, -3; -1, 0)$ and $\mathbf{d} = (d^t; \mathbf{d}^r; \mathbf{d}^l) = (2; 1, 2, 1; 2, 2)$, one obtains the h -transverse polygon in Figure 2.

Let $P(\mathbf{c}, \mathbf{d})$ an h -transverse polygon as above. Furthermore, consider $g \geq 0$ and four sequences $\alpha, \beta, \tilde{\alpha}, \tilde{\beta}$ with

$$\sum_i i(\alpha_i + \beta_i) = d^b \quad \text{and} \quad \sum_i i(\tilde{\alpha}_i + \tilde{\beta}_i) = d^t.$$

Let D_b and D_t the toric divisors corresponding to the bottom and top edge of $P(\mathbf{c}, \mathbf{d})$ respectively. The enumerative problem, we study in this article counts genus g curves in $S(\mathbf{c})$ passing through an appropriate number of points and satisfying tangency conditions at the toric divisors D_b and D_t . The tangency conditions at D_b are determined by α and β , while $\tilde{\alpha}$ and $\tilde{\beta}$ determine the conditions at D_t .

We define $a = \sum_{i=1}^n d_i^r = \sum_{j=1}^m d_j^l$ and $l = 2a + g + \sum_{i \geq 1} (\beta_i + \tilde{\beta}_i) - 1$. Finally, we say that a curve in $S(\mathbf{c})$ with Newton polygon $P(\mathbf{c}, \mathbf{d})$ has multidegree \mathbf{d} .

Now, we are ready to define double Gromov–Witten invariants for h -transverse polygons.

Definition 1.5. In the set-up above, we consider a generic configuration

$$\omega = (p_1, \dots, p_l, q_1^1, \dots, q_{\alpha_1}^1, \dots, q_1^i, \dots, q_{\alpha_i}^i, \dots, \tilde{q}_1^1, \dots, \tilde{q}_{\tilde{\alpha}_1}^1, \dots, \tilde{q}_1^i, \dots, \tilde{q}_{\tilde{\alpha}_i}^i, \dots),$$

where $q_j^i \in D_b$, $\tilde{q}_j^i \in D_t$ and $p_i \in S(\mathbf{c}) \setminus (D_t \cup D_b)$.

We denote by $N_{\mathbf{c},g}^{\alpha,\beta,\tilde{\alpha},\tilde{\beta}}(\mathbf{d})$ the number of irreducible complex algebraic curves C of multidegree \mathbf{d} in $S(\mathbf{c})$ of genus g such that:

- C passes through all the points q_j^i , \tilde{q}_j^i and p_i ;
- C has order of contact i with D_b at q_j^i , and has β_i other non-prescribed points with order of contact i with D_b ;
- C has order of contact i with D_t at \tilde{q}_j^i , and has $\tilde{\beta}_i$ other non-prescribed points with order of contact i with D_t .

This number is finite and does not depend on the chosen generic configuration of points. We call this number a *double Gromov-Witten invariant* of $S(\mathbf{c})$. When we allow reducible curves, we denote the resulting invariant by $N_{\mathbf{c},g}^{\alpha,\beta,\tilde{\alpha},\tilde{\beta},\bullet}(\mathbf{d})$.

Remark 1.6. The number of prescribed points in the interior of $S(\mathbf{c})$ in the configuration ω is given by

$$\begin{aligned} & |\partial P(\mathbf{c}, \mathbf{d}) \cap \mathbb{Z}^2| + g - 1 - \sum_{i \geq 1} (i-1)(\alpha_i + \tilde{\alpha}_i + \beta_i + \tilde{\beta}_i) = \\ &= \sum_{i=1}^n d_i^r + \sum_{j=1}^m d_j^l + d^t + d^b + g - 1 - \sum_{i \geq 1} (i-1)(\alpha_i + \tilde{\alpha}_i + \beta_i + \tilde{\beta}_i) \\ &= 2a + g + \sum_{i \geq 1} (\alpha_i + \tilde{\alpha}_i + \beta_i + \tilde{\beta}_i) - 1. \end{aligned}$$

Note that we subtract the quantity $\sum_{i \geq 1} (i-1)(\alpha_i + \tilde{\alpha}_i + \beta_i + \tilde{\beta}_i)$ in order to have just the number of points without their multiplicities. Therefore, the number l we defined above is just the number of points in the configuration without the prescribed points belonging to the toric divisors D_t and D_b .

1.2. Results. Our main objective is to study the polynomiality properties of the invariant $N_{\mathbf{c},g}^{\alpha,\beta,\tilde{\alpha},\tilde{\beta}}(\mathbf{d})$. For this, we first reparametrise the invariant and interpret it as a function.

Let $\mathbf{d}^l, \mathbf{d}^r$ as above and $n_1, n_2 \geq 0$. Then, we define

$$\Lambda = \left\{ (x_1, \dots, x_{n_1}, y_1, \dots, y_{n_2}) \in \mathbb{Z}^{n_1} \times \mathbb{Z}^{n_2} \mid \sum_{i=1}^{n_1} x_i + \sum_{j=1}^{n_2} y_j + \sum_{i=1}^n c_i^r d_i^r - \sum_{j=1}^m c_j^l d_j^l = 0 \right\}.$$

For $(\mathbf{x}, \mathbf{y}) \in \Lambda$, we associate a tuple $(\alpha, \beta, \tilde{\alpha}, \tilde{\beta})$ as follows: α_i is the number of elements $x_j = -i$, β_i is the number of elements $y_j = -i$, $\tilde{\alpha}_i$ is the number of elements $x_j = i$ and $\tilde{\beta}_i$ is the number elements $y_j = i$. Obviously (\mathbf{x}, \mathbf{y}) and $(\alpha, \beta, \tilde{\alpha}, \tilde{\beta})$ determine each other.

We consider the function

$$\begin{aligned} F_{(\mathbf{d}^r, \mathbf{d}^l), \mathbf{c}, g}^{n_1, n_2} : \Lambda &\longrightarrow \mathbb{Z} \\ (\mathbf{x}, \mathbf{y}) &\longmapsto N_{\mathbf{c}, g}^{\alpha, \beta, \tilde{\alpha}, \tilde{\beta}}(\mathbf{d}). \end{aligned}$$

In order to study the polynomiality properties of $F_{(\mathbf{d}^r, \mathbf{d}^l), \mathbf{c}, g}^{n_1, n_2}$, consider the hyperplane arrangement in Λ consisting of all hyperplanes

$$\begin{aligned} \sum_{i \in S} x_i + \sum_{j \in T} y_j + \sum_{i=1}^n c_i^r k_i - \sum_{j=1}^m c_j^l t_j &= 0 \\ y_i - y_j &= 0 \quad 1 \leq i < j \leq n_2 \end{aligned}$$

where $S \subseteq [n_1]$, $T \subseteq [n_2]$, $0 \leq k_i \leq d_i^r$ for $i = 1, \dots, n$ and $0 \leq t_j \leq d_j^l$ for $j = 1, \dots, m$. We denote this hyperplane arrangement by $\mathcal{H}^{n_1, n_2}(\mathbf{c})$.

We call the elements of $\mathcal{H}^{n_1, n_2}(\mathbf{c})$ walls and the connected components of $\Lambda \setminus \mathcal{H}^{n_1, n_2}(\mathbf{c})$ chambers.

The following is the first main result of this work.

Theorem 1.7. *Let $(\mathbf{d}^r, \mathbf{d}^l) > 0$ a vector with integer coordinates and $g \geq 0$, $n_1, n_2 > 0$ fixed integers and $\mathbf{c} = (\mathbf{c}^r, \mathbf{c}^l) \in \mathbb{Z}^{n+m}$ such that $c_1^r > \dots > c_n^r$ and $c_1^l < \dots < c_m^l$. The function $F_{(\mathbf{d}^r, \mathbf{d}^l), \mathbf{c}, g}^{n_1, n_2}(\mathbf{x}, \mathbf{y})$ of double Gromov-Witten invariants of the toric surface $S(\mathbf{c})$ is polynomial in each chamber of $\Lambda \setminus \mathcal{H}^{n_1, n_2}(\mathbf{c})$.*

In a next step we further study the polynomials appearing in Theorem 1.7 and prove the following result.

Theorem 1.8. *Each polynomial piece of $F_{(\mathbf{d}^r, \mathbf{d}^l), \mathbf{c}, g}^{n_1, n_2}(\mathbf{x}, \mathbf{y})$ has degree $n_2 + 3g + 2a - 2$, and is either even or odd.*

Remark 1.9. For the special case of $S(\mathbf{c})$ a Hirzebruch surface, Theorem 1.7 was proved in [AB17, Theorem 1.3] and Theorem 1.8 was proved in [AB17, Theorem 1.4].

The results in Theorems 1.7 and 1.8 establish the piecewise polynomiality of $N_{\mathbf{c}, g}^{\alpha, \beta, \tilde{\alpha}, \tilde{\beta}}(\mathbf{d})$ and determine the parity of the involved polynomials for a fixed toric surface. The following result shows that $N_{\mathbf{c}, g}^{\alpha, \beta, \tilde{\alpha}, \tilde{\beta}}(\mathbf{d})$ also behaves polynomially while changing the underlying toric surface, answering the aforementioned question posed in [AB17] affirmatively.

Theorem 1.10. *Let $(\mathbf{d}^r, \mathbf{d}^l) > 0$ a vector with integer coordinates and $g, n_1, n_2, n, m \geq 0$ fixed integers. The function $F_{(\mathbf{d}^r, \mathbf{d}^l), g}^{n_1, n_2}(\mathbf{x}, \mathbf{y}, \mathbf{c})$ of double Gromov-Witten invariants depending also on the toric surface $S(\mathbf{c})$ is piecewise polynomial relative to the chambers of the hyperplane arrangement*

$$\begin{aligned} \sum_{i \in S} x_i + \sum_{j \in T} y_j + \sum_{i=1}^n c_i^r k_i - \sum_{j=1}^m c_j^l t_j &= 0 \\ y_i - y_j &= 0 \quad 1 \leq i < j \leq n_2 \\ c_1^r > \dots > c_n^r \quad \text{and} \quad c_1^l < \dots < c_m^l \end{aligned}$$

where $S \subseteq [n_1]$, $T \subseteq [n_2]$, $0 \leq k_i \leq d_i^r$ for $i = 1, \dots, n$ and $0 \leq t_j \leq d_j^l$ for $j = 1, \dots, m$, inside $\Lambda \times \mathbb{Z}^{n+m}$.

Finally, in Section 4, we express our invariants as matrix elements in the bosonic Fock space. For this purpose, we introduce a certain operator whose vacuum expectations are equal to our invariants up to normalisation. The precise statement may be found in Theorem 4.5.

2. PRELIMINARIES

2.1. Floor diagrams. We introduce floor diagrams following the notation in [AB13] and [AB17]. Let $P(\mathbf{c}, \mathbf{d})$ be a h -transverse polygon and $S(\mathbf{c})$ the corresponding toric surface. We denote by D_r and D_l two multisets containing the directions of the right and left sides respectively:

$$\begin{aligned} D_r &= \underbrace{\{c_1^r, \dots, c_1^r\}}_{d_1^r \text{-times}}, \dots, \underbrace{\{c_n^r, \dots, c_n^r\}}_{d_n^r \text{-times}} \\ D_l &= \underbrace{\{c_1^l, \dots, c_1^l\}}_{d_1^l \text{-times}}, \dots, \underbrace{\{c_m^l, \dots, c_m^l\}}_{d_m^l \text{-times}}. \end{aligned}$$

Note that $|D_r| = |D_l| = a$. Let $r = (r_1, \dots, r_a)$ and $l = (l_1, \dots, l_a)$ be permutations of D_r and D_l respectively.

Definition 2.1. A **marked floor diagram** \mathcal{D} for $S(\mathbf{c})$ is a tuple (V, E, w) , such that

- (1) The vertex set V is decomposed as $V = L \cup C \cup R$. Moreover, we have that C is totally ordered from left to right, while $L = \{\tilde{q}_1, \dots, \tilde{q}_l\}$ is unordered and to the left of C , and $R = \{q_1, \dots, q_r\}$ is unordered and to the right of C .
- (2) The vertices in V are coloured black, white and grey and every vertex in L and R is white.
- (3) The set of edges E is directed from left to right, such that:
 - the resulting graph is connected;
 - every white vertex has valency one and is connected to precisely one black vertex;
 - every grey vertex has valency two with one incoming and one outgoing edge, each connecting it to a black vertex.
- (4) We have a map

$$w : E \rightarrow \mathbb{Z}_{>0}$$

such that if we define the divergence of v to be

$$\text{div}(v) = \sum_{e:v \rightarrow v'} w(e) - \sum_{e:v' \rightarrow v} w(e)$$

then

- $\text{div}(B_i) = r_i - l_i$, where B_i is the i -th black vertex in \mathcal{D} , for all $i = 1, \dots, a$;
- $\text{div}(v) = 0$ for every grey vertex v .

Moreover, we define the following notions associated to a marked floor diagram \mathcal{D} :

- We call (n_1, n_2) the **type** of \mathcal{D} , where \mathcal{D} has n_1 white vertices in $L \cup R$ and n_2 white vertices in C .
- Let
 - $\mathbf{x} = (\text{div}(\tilde{q}_1), \dots, \text{div}(\tilde{q}_l), \text{div}(q_1), \dots, \text{div}(q_r))$ the sequence of divergences of white vertices in L and R ;
 - \mathbf{y} the sequence of divergences of white vertices in C , listed from left to right.

We call the vector $(\mathbf{x}, \mathbf{y}) \in \mathbb{Z}^{n_1} \times \mathbb{Z}^{n_2}$ of length $n_1 + n_2$ the **divergence sequence** of \mathcal{D} . Since the divergences for a fixed marked floor diagram sum to zero, we have:

$$\sum_{i=1}^{n_1} x_i + \sum_{j=1}^{n_2} y_j = \sum_{j=1}^m c_j^l d_j^l - \sum_{i=1}^n c_i^r d_i^r = d^t - d^b.$$

- To each marked floor diagram \mathcal{D} , we associate the **divergence multiplicity vector** $(\alpha(\mathbf{x}), \beta(\mathbf{y}), \tilde{\alpha}(\mathbf{x}), \tilde{\beta}(\mathbf{y}))$ where
 - α_i is the number of vertices $v \in R$ such that $\text{div}(v) = -i$;
 - $\tilde{\alpha}_i$ is the number of vertices $v \in L$ such that $\text{div}(v) = i$;
 - β_i is the number of white vertices $v \in C$ such that $\text{div}(v) = -i$;
 - $\tilde{\beta}_i$ is the number of white vertices $v \in C$ such that $\text{div}(v) = i$;
- We define the **multidegree** of \mathcal{D} as the vector $\mathbf{d} = (d^t, \mathbf{d}^r, \mathbf{d}^l)$ with

$$\sum_i i(\alpha_i + \beta_i) = d^t + \sum_{i=1}^n d_i^r c_i^r - \sum_{j=1}^m d_j^l c_j^l \quad \text{and} \quad \sum_i i(\tilde{\alpha}_i + \tilde{\beta}_i) = d^t.$$

- We define the **genus** $g(\mathcal{D})$ of \mathcal{D} to be the first Betti number of the underlying graph, i. e. $g(\mathcal{D}) = 1 - |V| + |E|$.
- An edge is **internal** if it connects two vertices of C . We define the **multiplicity** $\mu(\mathcal{D})$ of \mathcal{D} as

$$\mu(\mathcal{D}) = \prod w(e)$$

where the product runs over all internal edges.

We now collect the two combinatorial properties of floor diagrams, see [AB17, Lemma 5.1 and Lemma 5.2] for the special case of Hirzebruch surfaces.

Proposition 2.2. *Let \mathcal{D} be a floor diagram. The genus of \mathcal{D} is $g(\mathcal{D}) = 1 - v_b + v_g$, where v_b and v_g is the number of black and grey vertices respectively.*

Proof. The number of vertices $|V|$ in \mathcal{D} is the sum $v_w + v_b + v_g$, while the number of edges $|E|$ is $v_w + 2v_g$, where v_w is the number of white vertices in \mathcal{D} . Therefore

$$g(\mathcal{D}) = 1 - |V| + |E| = 1 - (v_w + v_b + v_g) + (v_w + 2v_g) = 1 - v_b + v_g.$$

□

Proposition 2.3. *A floor diagram for $S(\mathbf{c})$ of multidegree \mathbf{d} and type (n_1, n_2) has:*

- $a = |D_r| = |D_l|$ black vertices, $g + a - 1$ grey vertices and $n_1 + n_2$ white vertices;
- $2(g + a - 1)$ black-grey edges and $n_1 + n_2$ black-white edges.

Proof. • From point (4) in Definition 2.1, it follows immediately that the number of black vertices is equal to a . By Proposition 2.2 $g = 1 - v_b + v_g$, therefore $v_g = g + v_b - 1 = g + a - 1$. Since a floor diagram of type (n_1, n_2) has n_1 white vertices in R and L and n_2 white vertices in C , the total number of white vertices is $n_1 + n_2$.

- From point (3) in Definition 2.1, it follows that for each grey vertex there are two edges connecting it to two different black vertices and each white vertex is connected to precisely one black vertex, hence the number of black-grey edges is $2v_g = 2(g + a - 1)$ and the number of black-white edges is $n_1 + n_2$.

□

Theorem 2.4. *Let $\mathbf{d} = (d^t, \mathbf{d}^r, \mathbf{d}^l) > 0$ be a vector of integer numbers, $g \geq 0$ an integer and \mathbf{x} a vector with coordinates in $\mathbb{Z} \setminus \{0\}$. We write $\alpha(\mathbf{x}) = \alpha$ and $\tilde{\alpha}(\mathbf{x}) = \tilde{\alpha}$. Then, for any two sequences of non-negative integer numbers $\beta = (\beta_i)_{i \geq 1}$ and $\tilde{\beta} = (\tilde{\beta}_i)_{i \geq 1}$ such that*

$$\sum_i i(\alpha_i + \beta_i) = d^t + \sum_{i=1}^n c_i^r d_i^r - \sum_{j=1}^m c_j^l d_j^l \quad \text{and} \quad \sum_i i(\tilde{\alpha}_i + \tilde{\beta}_i) = d^t,$$

one has

$$N_{\mathbf{c}, g}^{\alpha, \beta, \tilde{\alpha}, \tilde{\beta}}(\mathbf{d}) = \sum_{\mathcal{D}} \mu(\mathcal{D})$$

where the sum runs over all floor diagrams \mathcal{D} of multidegree \mathbf{d} , genus g , left-right sequence \mathbf{x} , and divergence multiplicity vector $(\alpha, \beta, \tilde{\alpha}, \tilde{\beta})$ for $S(\mathbf{c})$.

Proof. In [BM09] a proof of this result is given for

$$\alpha = \tilde{\alpha} = 0, \quad \beta = (d^b, 0, \dots, 0), \quad \tilde{\beta} = (d^t, 0, \dots, 0).$$

employing Mikhalkin's correspondence Theorem [Miko05] that expresses enumerations of curves in toric surfaces via *tropical curves* which are piecewise linear graphs in the plane. In [Shu12, Theorem 2] a generalisation of Mikhalkin's correspondence Theorem which covers the case of curves satisfying tangency conditions with toric divisors was proved. For the sake of convenience, we provide a sketch of the proof for a tropically inclined reader. Shustin's strategy was to dissipate each point p with multiplicity $k > 1$ in k points in a neighbourhood of p . Considering the curves passing through the new configuration of points and then specialising back to the original one, Shustin proved that each of these curves converges to a curve satisfying the prescribed tangency. This yields a correspondence theorem between enumeration of classical and tropical curves with point and tangency conditions. In order to obtain our result, we need to construct a bijection between tropical curves and floor diagrams.

Let $P = P(\mathbf{c}, \mathbf{d})$ be an h -transverse polygon and $S(\mathbf{c})$ be the corresponding toric surface. There is a correspondence between lattice subdivisions of P and the tropical curves of multidegree \mathbf{d} having P as a Newton polygon:

- the vertices of the tropical curve correspond to polygons in the lattice subdivision;
- the edges of the tropical curve emanated from a vertex v correspond to the normal lines to the sides of the polygon corresponding to v ;
- the faces determined by two edges emanated from a vertex v correspond to vertices of the polygon in the lattice subdivision corresponding to v .

Fixing a preferred direction in \mathbb{R}^2 allows to distinguish between edges of the tropical curve which are parallel to this direction and the others. The direction we fix is given by the vector $(0, 1)$. We call the unbounded edges of the tropical curve leaves and the edges in direction $(0, 1)$ elevators. A connected component of the tropical curve after removing bounded and unbounded elevators is called a floor.

We want to associate a decorated graph $\mathcal{D}(T)$ to the tropical curve T : black vertices of $\mathcal{D}(T)$ correspond to floors of T , grey vertices of $\mathcal{D}(T)$ correspond to bounded elevators of T and white vertices of $\mathcal{D}(T)$ correspond to unbounded elevators of T . More precisely, each vertex of the graph corresponds to a point in a fixed generic configuration. The weights of black-grey edges are the weights of the corresponding elevators, while the weights of the black-white edges are the weights of the corresponding leaves. We divide the graph in three blocks by adding vertical dashed lines, the white vertices in the central block correspond to non-prescribed points, while the white vertices in the left and right blocks correspond to points in the generic configuration belonging to D_t and D_b respectively. Furthermore, to each vertex we associate a number called divergence:

- to each black vertex we associate the difference between the slopes of the right and left leaves of the corresponding floor;
- to each grey vertex we associate 0;
- to each white vertex we associate the weight of the corresponding leaf if the leaf points to $+\infty$ or minus the weight of the corresponding leaf if the leaf points to $-\infty$.

The construction explained above yields a bijection between tropical curves and floor diagrams. \square

Example 2.5. In Figure 3, we give an example of a floor diagram constructed by using the technique explained in the proof of Theorem 2.4 in the case $P = P(\mathbf{c}, \mathbf{d})$, where $\mathbf{c} = (2; -1, 0)$, $\mathbf{d} = (1; 2; 1, 1)$ and $x_1 = 1$, $x_2 = -5$ and $y_1 = -1$.

Example 2.6. Let us consider $n = 2$, $m = 3$, $n_1 = 2$, $n_2 = 1$, $g = 0$, $\mathbf{c}^r = (c_1^r, c_2^r)$, $\mathbf{c}^l = (c_1^l, c_2^l, c_3^l)$, $\mathbf{d}^r = (2, 2)$ and $\mathbf{d}^l = (1, 1, 2)$. Then, $a = d_1^r + d_2^r = d_1^l + d_2^l + d_3^l = 4$ and the multisets are

$$D_r = \{c_1^r, c_1^r, c_2^r, c_2^r\} \quad D_l = \{c_1^l, c_2^l, c_3^l, c_3^l\}.$$

Let $r = (c_1^r, c_2^r, c_1^r, c_2^r)$ and $l = (c_3^l, c_1^l, c_3^l, c_2^l)$, so $r - l = (c_1^r - c_3^l, c_2^r - c_1^l, c_1^r - c_3^l, c_2^r - c_2^l)$. The floor diagram in Figure 4 is of type $(2, 1)$, with divergence sequence $(3, -4, -2)$, divergence multiplicity vector $(\alpha, \beta, \tilde{\alpha}, \tilde{\beta}) = (0001, 01, 001, 0)$, multidegree $\mathbf{d} = (3; 2, 2; 1, 1, 2)$ and multiplicity

$$\mu(\mathcal{D}) = 2(c_1^r - c_3^l + 3)^2 (c_2^r - c_1^l + c_1^r - c_3^l + 3)^2 (6 - c_2^r + c_2^l)^2.$$

This floor diagram contributes non-zero to the Gromov-Witten invariant

$$N_{\mathbf{c}, 0}^{0001, 01, 001, 0}(3; 2, 2; 1, 1, 2) = F_{(2, 2; 1, 1, 2), \mathbf{c}, 0}^{2, 1}(3, -4, -2)$$




FIGURE 3.




FIGURE 4.

of the surface $S(\mathbf{c})$ in the case all weights of the edges are positive, namely

$$\begin{cases} c_1^r - c_3^l + 3 > 0 \\ c_2^r - c_1^l + c_1^r - c_3^l + 3 > 0 \\ 6 - c_2^r + c_2^l > 0 \end{cases} \implies \begin{cases} c_1^r > c_3^l - 3 \\ c_1^l < c_2^r < c_2^l + 6 \end{cases}$$

Since $c_1^r > c_2^r$ we must have $c_3^l \geq c_2^l + 9$. Hence, the conditions on the slopes are

$$\begin{cases} c_1^r > c_3^l - 3 \\ c_1^l < c_2^r < c_2^l + 6 \\ c_3^l \geq c_2^l + 9 \end{cases}$$

Example 2.7. Fix $n = 2$, $m = 3$, $n_1 = 3$, $n_2 = 1$, $g = 1$, $\mathbf{c} = (\mathbf{c}^r; \mathbf{c}^l) = (c_1^r, c_2^r; c_1^l, c_2^l, c_3^l)$ and $(\mathbf{d}^r, \mathbf{d}^l) = (2, 1; 1, 1, 1)$. Furthermore, let $r = (c_1^r, c_2^r, c_1^l)$ and $l = (c_3^l, c_2^l, c_1^l)$, then $r - l = (c_1^r - c_3^l, c_2^r - c_2^l, c_1^r - c_1^l)$. Consider the floor diagram in Figure 5.

Note that the weights are uniquely determined by the variable w . Since every edge must have positive weight, we get the following inequalities:

$$x_1, x_2 > 0, \quad y_1, x_3 < 0,$$

$$w > 0, \quad c_2^l - c_2^r + w > 0, \quad -x_3 - w - c_1^r + c_1^l > 0.$$




FIGURE 5.

If we assume $c_1^l - c_1^r - x_3 > c_2^r - c_2^l$, the multiplicity of the floor diagram \mathcal{D} is the sum over the lattice points of the interval in \mathbb{R} described by the inequalities $\max\{0, c_2^r - c_2^l\} \leq w \leq c_1^l - c_1^r - x_3$, i. e.

$$\mu(\mathcal{D}) = \sum_{w=\max\{0, c_2^r - c_2^l\}}^{c_1^l - c_1^r - x_3} (-y_1)w^2(c_2^l - c_2^r + w)^2(-x_3 - w - c_1^r + c_1^l)^2.$$

Example 2.8. Let us consider $\mathbf{c}^r = 2$, $\mathbf{c}^l = (-1, 0)$, $g = 0$, $\alpha = 00001$, $\beta = 1$, $\tilde{\alpha} = 1$ and $\tilde{\beta} = 0$. In Figure 6, we see all the floor diagrams that contribute to $N_{(2; -1, 0), 0}^{00001, 1, 1, 0}(1; 2; 1, 1)$, so by Theorem 2.4 we get

$$N_{(2; -1, 0), 0}^{00001, 1, 1, 0}(1; 2; 1, 1) = 3\mu(\mathcal{D}_1) + \mu(\mathcal{D}_2) + 3\mu(\mathcal{D}_3) + \mu(\mathcal{D}_4) = 27 + 16 + 12 + 9 = 64.$$

2.2. Ehrhart theory. Here we review some results from [AB17] and examples on weighted partition functions and Ehrhart reciprocity.

Definition 2.9. Let $X = \{a_1, \dots, a_m\} \subseteq \mathbb{Z}^d$ be a finite multiset of lattice vectors of \mathbb{R}^d .

- The **rank** of X , denoted by $r(X)$, is the dimension of the real span of X .
- Writing X as an $m \times d$ matrix which columns are a_1, \dots, a_m , then X is **unimodular** if all the maximal minors are equal to $-1, 0$ and 1 .
- Define the **cone** of X as

$$\text{cone}(X) = \left\{ \sum_{i=1}^m t_i a_i \mid t_i \geq 0 \right\}.$$

We call X **pointed** if the unique linear subspace contained in $\text{cone}(X)$ is $\{0\}$, or, equivalently, if X lies in some open half-space of \mathbb{R}^d .

Definition 2.10. Let $X = \{a_1, \dots, a_m\} \subseteq \mathbb{Z}^d$ be a pointed vector configuration. Define the **partition function** $\mathcal{P}_X : \mathbb{Z}^d \rightarrow \mathbb{Z}$ as

$$\mathcal{P}_X(c) = \left| \left\{ (t_1, \dots, t_m) \in \mathbb{N}^m \mid c = \sum_{i=1}^m t_i a_i \right\} \right|$$

which is equivalent to the number of lattice points in the polytope

$$P_X(c) = \{t \in \mathbb{R}^m \mid Xt = c, t \geq 0\}.$$

Let $f \in \mathbb{R}[t_1, \dots, t_m]$ a polynomial, then we define the **weighted partition function** as

$$\mathcal{P}_{X,f}(c) = \sum_{t \in P_X(c) \cap \mathbb{Z}^m} f(t).$$




FIGURE 6.

The key idea in our investigation of the polynomiality of $F_{(d^r, d^l), c, g}^{n_1, n_2}$ is to express it as a weighted partition function, which exhibit polynomial behaviour. In order to be more precise, we need the following definitions.

Definition 2.11. The **chamber complex** $\text{Ch}(X)$ of X is a polyhedral complex supported on $\text{cone}(X)$. It is given by the common refinement of all cones spanned by subsets of X .

Definition 2.12. A function $f : \mathbb{Z}^d \rightarrow \mathbb{R}$ is **quasipolynomial** if there exists a sublattice $\Lambda \subseteq \mathbb{Z}^d$ of full rank and polynomials f_1, \dots, f_N corresponding to the different cosets $\Lambda_1, \dots, \Lambda_N$ of Λ such that $f(v) = f_i(v)$ for all $v \in \Lambda_i$. A function $f : \mathbb{Z}^d \rightarrow \mathbb{R}$ is **piecewise polynomial relative to** $\text{Ch}(X)$ if the restriction of f to any given face F of the chamber complex $\text{Ch}(X)$ is equal to a quasipolynomial function f^F depending on F .

Let $X = \{a_1, \dots, a_m\}$, $Y \subseteq X$ and $\pi_Y : \mathbb{R}^m \rightarrow \mathbb{R}$ be the function $\pi_Y(t_1, \dots, t_m) = \prod_{i \in Y} t_i$, where the components of the vector (t_1, \dots, t_m) correspond to the coefficients of the linear combination $t_1 a_1 + \dots + t_m a_m$. The following result was proved in [AB13, Theorem 4.2].

Theorem 2.13. *For any pointed vector configuration $X \subseteq \mathbb{Z}^d$ and for any subset $Y \subseteq X$, the weighted partition function \mathcal{P}_{X, π_Y} is piecewise quasipolynomial relative to the chamber complex $\text{Ch}(X)$. Furthermore, if X is unimodular, then \mathcal{P}_{X, π_Y} is piecewise polynomial. The polynomial pieces of \mathcal{P}_{X, π_Y} have degree $|X| + |Y| - r(X)$.*

The following example will be key for our proofs in Section 3.

Example 2.14 (Root system). Let e_1, \dots, e_d be the canonical basis of \mathbb{R}^d . The root system is the set

$$A_{d-1} = \{e_i - e_j \mid 1 \leq j < i \leq d\}.$$

It can be proved that A_{d-1} is unimodular. Moreover, A_{d-1} is contained in the hyperplane H_d contained in \mathbb{R}^d of equation $x_1 + \dots + x_d = 0$. Let us consider the discriminant arrangement S_d of H_d given by the hyperplanes

$$\sum_{i \in S} x_i = 0 \quad \emptyset \subset S \subset \{1, \dots, d\} = [d].$$

Note that $\sum_{i \in S} x_i = 0$ implies $\sum_{i \in [d] \setminus S} x_i = 0$. The elements in A_{d-1} span the hyperplanes in S_d . More precisely, any subset of A_{d-1} lies in one of the hyperplanes in S_d and any hyperplane of S_d contains a subset of A_{d-1} . Hence $\text{Ch}(A_{d-1})$ in H_d is the restriction of the discriminant arrangement S_d to $\text{cone}(A_{d-1})$, in other words the chamber complex of A_{d-1} in H_d is given by the hyperplanes in S_d intersected with the common refinement of $\text{cone}(A_{d-1})$.

We end this section with the following result which was proved in [AB13, Theorem 4.6].

Theorem 2.15 (Weighted Ehrhart reciprocity). *Let $P \subset \mathbb{R}^m$ be a rational polytope and $f : \mathbb{R}^m \rightarrow \mathbb{R}$ be a polynomial function. For each positive integer n , let*

$$L_{P,f}(n) = \sum_{z \in nP \cap \mathbb{Z}^m} f(z), \quad L_{P^\circ, f} = \sum_{z \in nP^\circ \cap \mathbb{Z}^m} f(z).$$

Then $L_{P,f}$ and $L_{P^\circ, f}$ extend to quasipolynomial functions which satisfy

$$L_{-P^\circ, f}(x) = (-1)^{\dim P} L_{P,f}(-x).$$

Furthermore, if P is a lattice polytope, then $L_{P,f}$ and $L_{P^\circ, f}$ are polynomial.

3. PROOFS ON THEOREMS 1.7, 1.8 AND 1.10

In this section, we prove Theorems 1.7, 1.8 and 1.10. In Section 3.1, we give the proofs of Theorems 1.7 and 1.10. We continue in Section 3.2 where we give the proof of Theorem 1.8. Finally, we illustrate our results in an example in Section 3.3.

3.1. Proof of Theorems 1.7 and 1.10. The root system presented in Example 2.14 will play a fundamental role. By Theorem 2.4 we have

$$F_{(\mathbf{d}^r, \mathbf{d}^l), \mathbf{c}, g}^{n_1, n_2}(\mathbf{x}, \mathbf{y}) = \sum_{\mathcal{D}} \mu(\mathcal{D})$$

where the sum runs over all floor diagrams \mathcal{D} for $S(\mathbf{c})$ having multidegree \mathbf{d} , genus g , divergence multiplicity vector $(\alpha, \beta, \tilde{\alpha}, \tilde{\beta})$ and left-right sequence \mathbf{x} . Let $\tilde{\mathcal{D}}$ be the floor diagram obtained by removing all weights of \mathcal{D} , but such that the underlying graph $\tilde{\mathcal{D}}$ inherit the partition $V = L \cup C \cup R$

of the vertices, the ordering of C and the coloring of the vertices. The collection \mathcal{G} of such graphs that contribute to $F_{(\mathbf{d}^r, \mathbf{d}^l), \mathbf{c}, g}^{n_1, n_2}(\mathbf{x}, \mathbf{y})$ is finite and depends only on g , $a = \sum_{i=1}^n d_i^r = \sum_{j=1}^m d_j^l$ and $n_1 + n_2$. Let us

denote by $Perm(D_r)$ and $Perm(D_l)$ the sets of permutations of the multisets D_r and D_l respectively and let $r \in Perm(D_r)$ and $l \in Perm(D_l)$. For each graph $G \in \mathcal{G}$, let $E(G)$ and $V(G)$ be the sets of edges and vertices of G respectively and define the set $W_{G, \mathbf{c}, r-l}(\mathbf{x}, \mathbf{y})$ of weights $w : E(G) \rightarrow \mathbb{N}$ for which the resulting weighted graph is a floor diagram for $S(\mathbf{c})$, i. e. such that the i -th black vertex has divergence $r_i - l_i$ and every grey vertex has divergence 0, with white divergence sequence (\mathbf{x}, \mathbf{y}) . Note that, by construction, the obtained floor diagram has genus g and multidegree \mathbf{d} . Our goal is to rewrite the sum $\sum_{\mathcal{D}} \mu(\mathcal{D})$ in a more convenient way, using the collection \mathcal{G} . To do that, call $\mathbb{R}^X = \{\mathbf{w} : X \rightarrow \mathbb{R}\}$

and let $\pi_{\text{int}} : \mathbb{R}^{E(G)} \rightarrow \mathbb{R}$ be the polynomial function defined by $\pi_{\text{int}}(\mathbf{w}) = \prod_{\substack{e \in E(G) \\ e \text{ internal}}} \mathbf{w}(e)$, which is the multiplicity of the floor diagram \mathcal{D} obtained from G adding the weights $\mathbf{w}(e)$ at every internal edge e . Hence

$$F_{G, \mathbf{c}, r-l}(\mathbf{x}, \mathbf{y}) = \sum_{\mathbf{w} \in W_{G, \mathbf{c}, r-l}} \pi_{\text{int}}(\mathbf{w}).$$

Note that $F_{G, \mathbf{c}, r-l}(\mathbf{x}, \mathbf{y})$ depends on the order of the entries of \mathbf{y} , while in $F_{(\mathbf{d}^r, \mathbf{d}^l), \mathbf{c}, g}^{n_1, n_2}(\mathbf{x}, \mathbf{y})$ we have to consider all the distinct orders for \mathbf{y} :

$$F_{(\mathbf{d}^r, \mathbf{d}^l), \mathbf{c}, g}^{n_1, n_2}(\mathbf{x}, \mathbf{y}) = \frac{1}{\beta_1! \beta_2! \cdots \tilde{\beta}_1! \tilde{\beta}_2! \cdots} \sum_{G \in \mathcal{G}} \sum_{(r, l)} \sum_{\sigma \in \mathcal{S}_{n_2}} F_{G, \mathbf{c}, r-l}(\mathbf{x}, \sigma(\mathbf{y}))$$

We want to have the setting of Theorem 2.13. Recall that the divergence of a vertex is defined as

$$\text{div}(v) = \sum_{e: v \rightarrow v'} w(e) - \sum_{e: v' \rightarrow v} w(e)$$

and that the adjacency matrix of the graph G is given by $A \in \mathbb{R}^{V(G) \times E(G)}$ which is, in our convention:

$$A(v, e) = \begin{cases} 1 & \text{when } e : v \rightarrow v' \text{ for some } v' \\ -1 & \text{when } e : v' \rightarrow v \text{ for some } v' \\ 0 & \text{otherwise} \end{cases}$$

Note that the columns of the matrix A are a subset of the root system $A_{|E(G)|-1}$, therefore the matrix A is unimodular. Now, take $\mathbf{k} \in \mathbb{R}^{V(G)}$ and define the flow polytope

$$\begin{aligned} \Phi_G(\mathbf{k}) &= \{\mathbf{w} \in \mathbb{R}^{E(G)} \mid \mathbf{w}(e) \geq 0 \text{ for all } e \in E(G), \text{div}(v) = \mathbf{k}(v) \text{ for all vertices } v\} \\ &= \{\mathbf{w} \in \mathbb{R}^{E(G)} \mid A\mathbf{w} = \mathbf{k}, \mathbf{w} \geq 0\}. \end{aligned}$$

If we take \mathbf{k} to be the vector which entries are given by (\mathbf{x}, \mathbf{y}) for the white vertices, $r - l$ for the black vertices and 0 for the gray vertices, then $W_{G, \mathbf{c}, r-l}(\mathbf{x}, \mathbf{y}) = \Phi_G(\mathbf{k})$ because A is unimodular and \mathbf{k} is a vector of integers, therefore the solutions of the linear system $A\mathbf{w} = \mathbf{k}$ must be vectors with integer entries.

By Theorem 2.13 and Example 2.14, the weighted partition function

$$\mathcal{P}_{G, \pi_{\text{int}}}(\mathbf{k}) = \sum_{\mathbf{w} \in \Phi_G(\mathbf{k})} \pi_{\text{int}}(\mathbf{w})$$

is piecewise polynomial relative to the chambers of the discriminant arrangement in $\{\mathbf{k} \in \mathbb{R}^{V(G)} \mid \sum \mathbf{k}(v) = 0\}$. Recall that this arrangement consists of the hyperplanes $\sum_{v' \in V'} \mathbf{k}(v') = 0$ for all subsets $V' \subseteq V$. Since our objects of interest are floor diagrams, we need the values of $\mathcal{P}_{G, \pi_{\text{int}}}(\mathbf{k})$ on the subspace H_{r-l} determined by the equations

$$\mathbf{k}(w_i) = x_i, \quad \mathbf{k}(w_j) = y_j, \quad \mathbf{k}(u) = 0 \text{ for all gray } u, \quad \mathbf{k}(b_i) = r_i - l_i \text{ for all black } b_i.$$

Since $\sum_{i=1}^a r_i - l_i = \sum_{i=1}^n c_i^r d_i^r - \sum_{j=1}^m c_j^l d_j^l$ and the sum of all divergences is 0, we have

$$\sum_{i=1}^{n_1} x_i + \sum_{j=1}^{n_2} y_j + \sum_{i=1}^n c_i^r d_i^r - \sum_{j=1}^m c_j^l d_j^l = 0$$

automatically, thus $H_{r-l} \subseteq \{\mathbf{k} \in \mathbb{R}^{V(G)} \mid \sum \mathbf{k}(v) = 0\}$. The restriction of $\mathcal{P}_{G, \pi_{\text{int}}}(\mathbf{k})$ to the subspace H_{r-l} is the function $F_{G, \mathbf{c}, r-l}(\mathbf{x}, \mathbf{y})$, which is piecewise polynomial in the chamber structure stated above. When we symmetrise, the result $\sum_{\sigma \in S_{n_2}} F_{G, \mathbf{c}, r-l}(\mathbf{x}, \sigma(\mathbf{y}))$ is still piecewise polynomial relative

to the same chambers, since the chamber structure is fixed under permutation of the n_2 \mathbf{y} variables. What remains to prove is that $\sum_{(r,l)} \sum_{\sigma \in S_{n_2}} F_{G, \mathbf{c}, r-l}(\mathbf{x}, \sigma(\mathbf{y}))$ is piecewise polynomial. Since every pair of

permutations (r, l) gives rise to a hyperplane arrangement S_{r-l} in H_{r-l} , then $\sum_{(r,l)} \sum_{\sigma \in S_{n_2}} F_{G, \mathbf{c}, r-l}(\mathbf{x}, \sigma(\mathbf{y}))$ is piecewise polynomial in the chambers of the hyperplane arrangement $\mathcal{H}^{n_1, n_2}(\mathbf{c}) = \bigcup_{(r,l)} S_{r-l}$. This

completes the proof. Finally, Theorem 1.10 follows from the same arguments.

Remark 3.1. In Example 2.7 we computed the polytope $\Phi_G(\mathbf{k}) = [\max\{0, c_2^r - c_2^l\}, c_1^l - c_1^r - x_3]$, where $\mathbf{k} = (x_1, x_2, c_1^r - c_3^l, y_1, 0, 0, c_2^r - c_2^l, 0, c_1^r - c_1^l, x_3)$.

Remark 3.2. In the case in which $m = 1$ something interesting happens to the hyperplane arrangements S_{r-l} . Indeed, in this case the only left permutation possible is the identity, since $D_l = \underbrace{\{c^l, \dots, c^l\}}_{d^l \text{-times}}$;

therefore $S_{r-id} = S_{\tilde{r}-id}$ with $r, \tilde{r} \in \text{Perm}(D_r)$. This means that the chamber structure for the piecewise polynomiality of the function $F_{(\mathbf{d}^r, \mathbf{d}^l), \mathbf{c}, g}^{n_1, n_2}(\mathbf{x}, \mathbf{y})$ does not depend on the permutation r .

Remark 3.3. In [CJM10] the authors presented a technique to get a wall crossing formula for genus 0 double Hurwitz numbers and further it was generalised to arbitrary genus in [CJM11]. We briefly sketch the technique in the case of genus 0 double Hurwitz numbers: let C_1 and C_2 be two adjacent chambers of polynomiality for genus 0 double Hurwitz numbers and call $\delta = 0$ the equation of the wall dividing them. Let us consider a graph contributing to C_1 that presents an edge with weight δ . Once we pass through the wall what happens is that the orientation of the edge having weight δ will be inverted, which corresponds to cutting and regluing the edge in a suitable way to get the new graph and changing the sign of the edge weight. This operation divides the graph into two new graphs contributing to two genus 0 double Hurwitz numbers having new data and this provides the recursive formula. When adapting this to our case, each edge weight δ arises twice, thus we obtain a contribution of δ^2 . The squaring erases the sign change. Therefore, we expect that new techniques may be necessary to possibly derive wall-crossing formulae for our invariants.




FIGURE 7. The h -transverse polygon associated to the data above.

3.2. Proof of Theorem 1.8. In the notation of Theorem 1.7, it suffices to show these claim for the following piecewise polynomial function for each graph G and permutations $r \in \text{Perm}(D_r)$ and $l \in \text{Perm}(D_l)$:

$$F_{G,c,r-l}(\mathbf{x}, \mathbf{y}) = \sum_{\mathbf{w} \in W_{G,c,r-l}} \pi_{\text{int}}(\mathbf{w}).$$

The degree of the polynomial $\pi_{\text{int}}(\mathbf{w})$ is the number of interior edges, which is $n_2 + 2(g + a - 1)$ by Proposition 2.3. Moreover, the polytope $W_{G,c,r-l}$ has dimension g in each full dimensional chamber since g is the smallest number with the following property: if we fix the flow on g edges whose removal turns the graph into a tree, the whole flow vector will be uniquely determined. Therefore, summing over $W_{G,c,r-l}$ produce a polynomial piece of degree $g + [n_2 + 2(g + a - 1)] = n_2 + 3g + 2a - 2$. Let

$$L_{W_{G,c,r-l}, \pi_{\text{int}}}(t) = \sum_{\mathbf{w} \in tW_{G,c,r-l}} \pi_{\text{int}}(\mathbf{w}) = F_{G,c,r-l}(t\mathbf{x}, t\mathbf{y}).$$

By Theorem 2.15 we have

$$F_{G,c,r-l}(-t\mathbf{x}, -t\mathbf{y}) = L_{W_{G,c,r-l}, \pi_{\text{int}}}(-t) = (-1)^g L_{-W_{G,c,r-l}^\circ, \pi_{\text{int}}}(t).$$

Denote by $i = n_2 + 2(g + a - 1)$ the number of internal edges in G , then $\pi_{\text{int}}(-w) = (-1)^i \pi_{\text{int}}(w)$ for any weight w for G , therefore we get

$$F_{G,c,r-l}(-t\mathbf{x}, -t\mathbf{y}) = (-1)^g L_{-W_{G,c,r-l}^\circ, \pi_{\text{int}}}(t) = (-1)^{g+i} L_{W_{G,c,r-l}^\circ, \pi_{\text{int}}}(t) = (-1)^{n_2+3g+2a-2} F_{G,c,r-l}(t\mathbf{x}, t\mathbf{y}).$$

Hence, the function $F_{G,c,r-l}(t\mathbf{x}, t\mathbf{y})$ is either even or odd depending on the parity of $n_2 + 3g + 2a - 2$.

3.3. Example. The goal of this subsection is to explicitly compute the function $F_{(d^r, d^l), c, g}^{n_1, n_2}(\mathbf{x}, \mathbf{y})$ in the case $n_1 = 2, n_2 = 1, \mathbf{c}^r = (k, 0)$, where $k > 0$ is an integer, $\mathbf{c}^l = 0, \mathbf{d}^r = (1, 1), \mathbf{d}^l = 2$ and $g \geq 0$. The corresponding polytope is given in Figure 7.

In Tables 1 and 2 we list all the floor diagrams contributing to $F_{(1,1;2), (k,0;0), 0}^{2,1}(x_1, x_2, y_1)$ with divergence sequence $r_1 - l$ and $r_2 - l$ respectively. Note that, since $c_2^r = c^l = 0$ the domain will be

$$\Lambda = \{(\mathbf{x}, \mathbf{y}) \in \mathbb{Z}^2 \times \mathbb{Z} \mid x_1 + x_2 + y_1 + k = 0\}.$$

The following planes provide the subdivision of Λ in sixteen chambers, see Figure 8:

$$x_1 = 0, x_1 + k = 0, x_2 = 0, x_2 + k = 0, y_1 = 0, y_1 + k = 0.$$

Remark 3.4. We note that even though we do not work with a Hirzebruch surface, the chamber complex we obtain coincides with the one in [AB17, Section 6]. This is due to the fact that the polygon in Figure 7 only differs from the polygon of a Hirzebruch surface by a vertical edge.




FIGURE 8. The chamber complex for $F_{(1,1;2),(k,0;0),0}^{2,1}(x_1, x_2, y_1)$

As in [AB17], we label each chamber with a triple $s_{x_1}s_{x_2}s_{y_1}$ where each s_i is +, 0 or – according to whether the corresponding variable is > 0 , between $-k$ and 0, or $< -k$ respectively. For instance, the chamber $0 + -$ is given by the inequalities:

$$x_1 + k > 0 > x_1, \quad x_2 + k > x_2 > 0, \quad y_1 < y_1 + k < 0.$$

Since $F_{(1,1;2),(k,0;0),g}^{2,1}(x_1, x_2, y_1) = F_{(1,1;2),(k,0;0),0}^{2,1}(x_2, x_1, y_1)$, it is sufficient to compute this function for $x_1 \geq x_2$, therefore we consider just ten of the sixteen chambers and the corresponding polynomials are listed in Table 3. The polynomials in the remaining six chambers can be obtained by symmetry.

Let us discuss the case $g = 0$. The graphs listed in Tables 1 and 2 contribute to $F_{(1,1;2),(k,0;0),0}^{2,1}(x_1, x_2, y_1)$ and they are obtained by a careful analysis of the weights.

Example 3.5. Take the graphs $B2$ and $B'2$. The edge weights for the graph $B2$, from left to right, must be $x_1, -x_2, -x_2, -y_1, -x_2$, while for the graph $B'2$ they must be $x_1, -x_2 - k, -x_2 - k, -y_1, -x_2$. Therefore, the graph $B2$ contributes to $F_{(1,1;2),(k,0;0),0}^{2,1}(x_1, x_2, y_1)$ with weight $(-y_1)x_2^2$ as long as $x_1 > 0, x_2, y_1 < 0$; that is, in chambers $+0-, +--$ and $+ - 0$. On the other hand, the graph $B'2$ contributes to $F_{(1,1;2),(k,0;0),0}^{2,1}(x_1, x_2, y_1)$ with weight $(-y_1)(x_2 + k)^2$ as long as $x_1 > 0, x_2 < x_2 + k < 0, y_1 < 0$; that is, in chambers $+--$ and $+ - 0$.

Note that the graphs in $B2$ and $B'2$ are the same, but we have different divergence sequences and for this reason we have different weights.

In general, all the graphs in Tables 1 and 2 contribute to $F_{(1,1;2),(k,0;0),0}^{2,1}(x_1, x_2, y_1)/|y_1|$ with weight $(w + k)^2$, where $w \in \{0, x_1, x_2, x_1 - k, x_2 - k, y_1, y_1 - k\}$.

	A	B	C	D
1				
2				
3				
4				
5				
6				
7				
8				
9				
10				
11				
12				
13				

TABLE 1. The floor diagrams with divergence sequence $r_1 - l$ that contribute to $F_{(1,1;2),(k,0;0),0}^{2,1}(x_1, x_2, y_1)$

	A'	B'	C'	D'
1				
2				
3				
4				
5				
6				
7				
8				
9				

TABLE 2. The floor diagrams with divergence sequence $r_2 - l$ that contribute to $F_{(1,1;2),(k,0;0),0}^{2,1}(x_1, x_2, y_1)$

Remark 3.6. For each graph in rows from seven to thirteen in Table 1 and for each graph in rows from five to nine in Table 2 (i. e. when x_1 and x_2 have the same sign), there are a priori two different possibilities of labeling the vertices in L or R , respectively, with \tilde{q}_1 and \tilde{q}_2 or q_1 and q_2 . The two corresponding floor diagrams are the same for graphs in rows eight, ten, eleven and twelve in Table 1 and in rows six, seven and eight in Table 2, and they are different for graphs in rows seven, nine and thirteen in Table 1 and in rows five and nine in Table 2.

Let us discuss the case $g > 0$. In each graph, we need to replace the grey vertex and its two incident edges by $g + 1$ gray vertices and the corresponding $2(g + 1)$ edges. The position of an intermediate white vertex can be chosen among the $g + 1$ grey vertices; there are $g + 3$ choices. This gives rise to the factor $g + 3$ in Table 3. For example, in chamber $+ - -$ and genus $g = 0$ the graphs $A2, B2, C2$ are

Chamber	Graphs ($g=0$)	$F_{(1,1;2),(k,0;0),g}^{2,1}(x_1, x_2, y_1)/ y_1 $
++-	A'_5, A_5, A_7, A_7, A_8	$\Gamma(y_1 - k) + \Gamma(y_1) + \Gamma(x_1) +$ $+ \Gamma(x_1 - k) + \Gamma(x_2) + \Gamma(x_2 - k)$
+0-	$A_1, A'_1, A_2, B_2, C_2,$ A_3, A'_3, A_4, A_5	$\Gamma(x_1) + \Gamma(x_1 - k) + (g + 3)\Gamma(x_2 - k) +$ $+ \Gamma(y_1 - k) + \Gamma(y_1) + \Gamma(x_2) + \Gamma(0)$
+ -	$A_1, A'_1, A_2, A'_2, B_2,$ $B'_2, C_2, C'_2, A_3, A'_3,$ A_5	$\Gamma(x_1) + \Gamma(x_1 - k) + (g + 3)\Gamma(x_2 - k) +$ $+ (g + 3)\Gamma(x_2) + \Gamma(y_1 - k) + \Gamma(y_1) + \Gamma(0)$
00-	$A'_8, A_9, A_9, A_{10}, A_{12},$ $A_{13}, A_{13}, B_{13}, B_{13}, C_{13},$ C_{13}	$\Gamma(x_1) + \Gamma(x_2) + \Gamma(0) + \Gamma(y_1 - k) + \Gamma(y_1) +$ $+ (g + 3)\Gamma(x_2 - k) + (g + 3)\Gamma(x_1 - k)$
+ - 0	$A_1, A'_1, A_2, A'_2, B_2,$ $B'_2, C_2, C'_2, A_3, A_5,$ A_6, B_6, C_6	$\Gamma(x_1) + \Gamma(x_1 - k) + (g + 3)\Gamma(x_2 - k) +$ $+ (g + 3)\Gamma(x_2) + \Gamma(y_1 - k) + \Gamma(0) +$ $+ (g + 3)\Gamma(y_1)$
0 - 0	$A_9, A'_9, B'_9, C'_9, A_{10},$ $A_{11}, B_{11}, C_{11}, A_{12}, A_{13},$ $A_{13}, B_{13}, B_{13}, C_{13}, C_{13}$	$\Gamma(x_1) + \Gamma(0) + (g + 3)\Gamma(y_1) + \Gamma(y_1 - k) +$ $+ (g + 3)\Gamma(x_2 - k) + (g + 3)\Gamma(x_1 - k) +$ $+ (g + 3)\Gamma(x_2)$
000	$A_9, A_9, A_{10}, A_{11}, B_{11},$ $C_{11}, A_{12}, A_{13}, A_{13}, B_{13},$ B_{13}, C_{13}, C_{13}	$\Gamma(x_1) + \Gamma(x_2) + \Gamma(0) + (g + 3)\Gamma(y_1) +$ $+ \Gamma(y_1 - k) + (g + 3)\Gamma(0) + \Gamma(y_1) + \Gamma(y_1 - k)$
+ - +	$B_1, B'_1, C_1, C'_1, D_1,$ $D'_1, D_2, D'_2, D'_4, B_5,$ C_5, D_5, D_6	$(g + 3)\Gamma(x_1) + (g + 3)\Gamma(x_1 - k) + \Gamma(x_2 - k) +$ $+ \Gamma(x_2) + (g + 3)\Gamma(0) + \Gamma(y_1) + \Gamma(y_1 - k)$
0 - +	$D'_7, B_9, C_9, D_9, D'_9,$ $B_{10}, C_{10}, D_{10}, D_{11}, D_{13},$ D_{13}	$(g + 3)\Gamma(x_1) + (g + 3)\Gamma(0) + \Gamma(y_1) +$ $+ \Gamma(y_1 - k) + \Gamma(x_2) + \Gamma(x_2 - k) + \Gamma(x_1 - k)$
- +	$D'_7, D'_9, D'_9, B_{10}, C_{10},$ $D_{10}, D_{11}, D_{13}, D_{13}$	$(g + 3)\Gamma(0) + \Gamma(y_1) + \Gamma(y_1 - k) + \Gamma(x_2) +$ $+ \Gamma(x_2 - k) + \Gamma(x_1) + \Gamma(x_1 - k)$

 TABLE 3. The double Gromov-Witten invariants $F_{(1,1;2),(k,0;0),g}^{2,1}(x_1, x_2, y_1)$

isomorphic as unoriented graphs, and they account the three possible positions of the white vertex in C relative to the black and grey vertices.

Suppose w to be the weight of a black-grey edge in a genus 0 graph. In the genus g graph, that total weight w has to be distributed among $g + 1$ weights. Therefore the resulting contribution is

$$\Gamma_g(w) = \sum_{w_1 + \dots + w_{g+1} = w} \prod_{i=1}^{g+1} w_i^2$$

where w_1, \dots, w_{g+1} are positive integers. The function Γ in Table 3 is given by

$$\Gamma(w) = \Gamma_g(|w + k|).$$

4. DOUBLE GROMOV–WITTEN INVARIANTS VIA THE BOSONIC FOCK SPACE

In this section, we establish a connection between our invariants $N_{\mathbf{c},g}^{\alpha,\beta,\tilde{\alpha},\tilde{\beta},\bullet}(\mathbf{d})$ and the bosonic Fock space. More precisely, we express them as matrix elements of certain operators. For this, we first need to re-organise floor diagrams. In this, we follow the notion of floor diagram introduced in [CJMR21] for the study of curves in Hirzebruch surfaces. For this, we fix the same setting as for Definition 2.1.

Definition 4.1. Let \mathcal{F} a loop-free graph with vertex set V , edge set E . There are two types of edges: bounded edges which are composed of two half-edges adjacent to different vertices and unbound edges called *ends* with one flag. We call \mathcal{F} a **thickened floor diagram** of multidegree \mathbf{d} and relative to (\mathbf{x}, \mathbf{y}) if:

- Each vertex $v \in V$ carries a size s_v that is either 0 or 1.
- Each half-edge may be decorated with a thickening and for each bounded edge exactly one of its half-edges is thickened.
- At each vertex v , exactly $2 - 2s_v$ half-edges are thickened.
- We have a map

$$w : E \rightarrow \mathbb{Z}_{>0}$$

associating to each edge $e \in E$ an expansion factor $w(e)$.

- For each vertex v denote by E_v^- the set of incoming edges at v and by E_v^+ the set of outgoing edges at v . We define the divergence of v as

$$\text{div}(v) = \sum_{e \in E_v^+} w(e) - \sum_{e \in E_v^-} w(e).$$

Then, we require that

- if $s_v = 0$, then $\text{div}(v) = 0$
- if $s_v = 1$, then $\text{div}(v) = l_i - r_i$, where v is the i -th vertex with $s_v = 1$.
- The sequence of expansion factors of non-thick ends is given by \mathbf{x} , while the sequence of expansion factors of thick ends is given by \mathbf{y} , where negative entries of \mathbf{x} and \mathbf{y} correspond to ends pointing to the left and positive entries to ends pointing to the right.
- The ends of \mathcal{F} are labelled by the parts of \mathbf{x} and \mathbf{y} .

The genus of \mathcal{F} is the first Betti number of the underlying graph.

Finally, we denote by

$$\mu(\mathcal{F}) = \prod w(e),$$

where the product runs over all bounded edges of \mathcal{F} , the **weight** of \mathcal{F} .

Similarly to Theorem 2.4, thickened floor diagrams compute our invariants $N_{\mathbf{c},g}^{\alpha,\beta,\tilde{\alpha},\tilde{\beta},\bullet}(\mathbf{d})$. More precisely, we have the following theorem which follows from the same arguments as in [CJMR21, section 5].

Theorem 4.2. *Let $\mathbf{d} = (d^t, \mathbf{d}^r, \mathbf{d}^l) > 0$ be a vector of integer numbers, $g \geq 0$ an integer and \mathbf{x} a vector with coordinates in $\mathbb{Z} \setminus \{0\}$. We write $\alpha(\mathbf{x}) = \alpha$ and $\tilde{\alpha}(\mathbf{x}) = \tilde{\alpha}$. Then, for any two sequences of non-negative integer numbers $\beta = (\beta_i)_{i \geq 1}$ and $\tilde{\beta} = (\tilde{\beta}_i)_{i \geq 1}$ such that*

$$\sum_i i(\alpha_i + \beta_i) = d^t + \sum_{i=1}^n c_i^r d_i^r - \sum_{j=1}^m c_j^l d_j^l \quad \text{and} \quad \sum_i i(\tilde{\alpha}_i + \tilde{\beta}_i) = d^t,$$

one has

$$N_{\mathbf{c},g}^{\alpha,\beta,\tilde{\alpha},\tilde{\beta},\bullet}(\mathbf{d}) = \sum_{\mathcal{F}} \mu(\mathcal{F})$$

where the sum runs over all thickened floor diagrams \mathcal{F} of multidegree \mathbf{d} , genus g and relative to (\mathbf{x}, \mathbf{y}) for $S(\mathbf{c})$.

This interpretation of $N_{\mathbf{c},g}^{\alpha,\beta,\tilde{\alpha},\tilde{\beta},\bullet}(\mathbf{d})$ is the basis for the remainder of the section. Next, we introduce the bosonic Fock space and related notions.

Definition 4.3. We define the two-dimensional Heisenberg algebra \mathcal{H} as generated by $(a_n)_{n \in \mathbb{Z}}$ and $(b_n)_{n \in \mathbb{Z}}$ satisfying the commutator relations

$$[a_n, a_m] = 0, \quad [b_n, b_m] = 0 \quad \text{and} \quad [a_n, b_m] = n\delta_{n,-m}$$

where $\delta_{n,-m}$ is the Kronecker symbol. Moreover, we set $a_0 = b_0 = 0$.

We now consider the free action of \mathcal{H} on the so-called vacuum vector v_\emptyset where we set $a_n \cdot v_\emptyset = b_n \cdot v_\emptyset = 0$ for $n > 0$. For two partitions μ, ν , we define

$$v_{\mu,\nu} = \frac{1}{|\text{Aut}(\mu)||\text{Aut}(\nu)|} \prod_{i=1}^{\ell(\mu)} a_{-\mu_i} \prod_{j=1}^{\ell(\nu)} b_{-\nu_j} \cdot v_\emptyset.$$

The vectors $v_{\mu,\nu}$ form a basis of a vector space which we call the bosonic Fock space. We also define an inner product by declaring $\langle v_\emptyset | v_\emptyset \rangle = 1$, a_n as the adjoint of a_{-n} and b_n as the adjoint of b_{-n} . Thus, we obtain

$$\langle v_{\mu,\nu} | v_{\mu',\nu'} \rangle = \frac{1}{|\text{Aut}(\mu)|} \frac{1}{|\text{Aut}(\nu)|} \prod_i \mu_i \prod_{i'} \nu_{i'} \delta_{\mu,\nu'} \delta_{\mu',\nu}.$$

Moreover, for an operator $M \in \mathcal{H}$, we denote $\langle v | M | w \rangle = \langle v | Mw \rangle$. Finally, we define the **normal ordering** of a monomial of operators $:\prod \alpha_i \prod \beta_j:$ as the product, such that all α_i, β_j with $i, j < 0$ appear to left of the operators with positive index.

We now define the operator whose matrix elements will give our invariants.

Definition 4.4. Let $P(\mathbf{c}, \mathbf{d})$ an h -transverse polygon, $(\mathbf{x}, \mathbf{y}) \in \Lambda$ and $s \in \mathbb{Z} \setminus \{0\}$. Let

$$\mathbf{a}_n := \begin{cases} u a_n, & \text{if } n < 0 \\ a_n & \text{if } n > 0 \end{cases} \quad \text{and} \quad \mathbf{b}_n := \begin{cases} u b_n, & \text{if } n < 0 \\ b_n & \text{if } n > 0 \end{cases}$$

Then, we define

$$M_{\mathbf{c}} = u^{-1} \sum_{m \in \mathbb{Z}_{>0}} \sum_{\substack{\mathbf{z} \in (\mathbb{Z})^m \\ \sum z_i = c}} : \mathbf{a}_{z_1} \cdots \mathbf{a}_{z_m} :$$

and

$$M = u^{-1} \sum_{m \in \mathbb{Z}_{>0}} : \mathbf{b}_{-m} \mathbf{b}_m : .$$

We are now ready to state our main theorem of this section.

Theorem 4.5. *Consider the same data as in Theorem 1.7. Then, we have*

$$N_{c,g}^{\alpha,\beta,\tilde{\alpha},\tilde{\beta},\bullet}(\mathbf{d}) = \frac{|\text{Aut}(\mathbf{x})||\text{Aut}(\mathbf{y})|}{\prod |x_i| \prod |x_j|} \left\langle v_{x^-,y^-} \mid \left[u^{g-1+\ell(x^-)+\ell(y^-)} \right] \sum_{\sigma} \sum_{\substack{(r_1,\dots,r_a) \\ (l_1,\dots,l_a)}} \sigma \left(\prod_{i=1}^a M_{l_i-r_i} M^{a+g+\ell(\mathbf{y})-1} \right) \mid v_{x^+,y^+} \right\rangle,$$

where $[u^{g-1+\ell(x^-)+\ell(y^-)}]$ denotes the coefficient of the monomial $t^a u^{g-1}$ and the sum runs over all permutations of D_r and D_l . Moreover the first sum runs over all permutations $\sigma \in S_{3a+g+\ell(\mathbf{y})-1}$ that respect the ordering of $1, \dots, a$ and $\sigma \left(\prod_{i=1}^a M_{r_i-l_i} M^{a+g+\ell(\mathbf{y})-1} \right)$ denotes the permutations of the $2a + g + \ell(\mathbf{y}) - 1$ factors induced by σ .

The proof follows from a tropical formulation of Wick's theorem [Wic50] that may be found in [BG16, Proposition 5.2] and in [CJMR21, Proposition 6.6] for the version we employ. The key idea is to construct (coloured) Feynman diagrams corresponding to matrix elements of monomials in the bosonic Fock space. The idea, also outlined after [CJMR21, Definition 6.4] is as follows: Let $m_-, m_1, \dots, m_s, m_+$ be normally ordered monomials in the operators a_i, b_j , such that m_- contains only operators with negative index and m_+ only operators with positive index. Then, we associate to the monomial $P = m_+ m_1 \cdots m_s m_-$ a family of graphs, we call **Feynman diagrams** associated to P via the following procedure:

- (1) For each monomial m_l , we create a vertex v_l . For each a_i with $i < 0$ appearing in m_l , we create a half-edge e adjacent to v_l pointing to the left with expansion factor $w(e) = |i|$. For each a_i with $i > 0$ we do create a half-edge pointing to the right with expansion factor i . Moreover, for each b_j appearing in m_l , we create a thickened half-edge adjacent to v_l again pointing to the left for $j < 0$ and to the right for $j > 0$. For the monomial m_- and m_+ we create a set of unbounded half-edges of corresponding weight for each operator appearing in them. We thicken the half-edges coming from operators b_i .
- (2) We order all pieces by moving the half-edges corresponding to m_+ to the left and the half-edges corresponding to m_- to the right. All vertices m_l are ordered linearly ordered in between respecting the ordering of the indices l .
- (3) We connect half-edges to other according to the following rules:
 - A half-edge pointing to the left is connected to a half-edge pointing to the right and the connection respects the ordering of the vertices.
 - Connected half-edges must have the same expansion factor.
 - All resulting edges that adjacent to two vertices have exactly one thickened half-edge.

As such the operators M_c corresponds to vertices of size 1 and non-trivial divergence, whereas the operator M yield vertices of size 0 and trivial divergence. Note that by the construction of thickened floor diagrams in [CJMR21], all vertices of size 0 are 2-valent.

Proposition 4.6 ([BG16, Proposition 5.2],[CJMR21, Proposition 6.6]). *The vacuum expectation of an operator P as above is equal to the weighted sum of all Feynman diagrams associated to P , where each diagram is weighted the product of all edges (bounded and unbounded).*

We end this section with the proof of Theorem 4.5.

Proof of Theorem 4.5. Clearly all Feynman diagrams are thickend Floor diagrams and the operators M_c and M are built, such that all thickend floor diagrams contributing to $N_{c,g}^{\alpha,\beta,\tilde{\alpha},\tilde{\beta},\bullet}(\mathbf{d})$ appear. The weights only differ by the product of the expansion factors of the ends, which is why we divide by them. A simple Euler characteristic calculation shows that all Feynman diagrams are of the right genus. \square

REFERENCES

- [AB13] F. Ardila and F. Block. “Universal polynomials for Severi degrees of toric surfaces”. In: *Advances in Mathematics* 237 (2013), pp. 165–193.
- [AB17] F. Ardila and E. Brugallé. “The double Gromov–Witten invariants of Hirzebruch surfaces are piecewise polynomial”. In: *International Mathematics Research Notices* 2017.2 (2017), pp. 614–641.
- [BG16] F. Block and L. Göttsche. “Fock spaces and refined Severi degrees”. In: *International Mathematics Research Notices* 2016.21 (2016), pp. 6553–6580.
- [BM16] E. Brugallé and G. Mikhalkin. “Floor decompositions of tropical curves: the general case”. In: *preparation, preliminary version available at <http://www.math.jussieu.fr/brugalle/articles/FDn/FDGeneral.pdf>* (2016).
- [BM09] E. Brugallé and G. Mikhalkin. “Floor decompositions of tropical curves: the planar case”. English. In: *Proceedings of the 15th Gökova geometry-topology conference, Gökova, Turkey, May 26–31, 2008*. Cambridge, MA: International Press, 2009, pp. 64–90. ISBN: 978-1-57146-136-0.
- [CJM10] R. Cavalieri, P. Johnson, and H. Markwig. “Tropical Hurwitz numbers”. In: *Journal of Algebraic Combinatorics* 32.2 (2010), pp. 241–265.
- [CJM11] R. Cavalieri, P. Johnson, and H. Markwig. “Wall crossings for double Hurwitz numbers”. In: *Advances in Mathematics* 228.4 (2011), pp. 1894–1937.
- [CJMR21] R. Cavalieri, P. Johnson, H. Markwig, and D. Ranganathan. “Counting curves on Hirzebruch surfaces: tropical geometry and the Fock space”. In: *Mathematical Proceedings of the Cambridge Philosophical Society*. Vol. 171. 1. Cambridge University Press. 2021, pp. 165–205.
- [FHK23] S. G. Fitzgerald, M. A. Hahn, and S. Kelly. “Combinatorics of pruned Hurwitz numbers”. In: *arXiv preprint arXiv:2310.05225* (2023).
- [HL20] M. Hahn and D. Lewański. “Wall-crossing and recursion formulae for tropical Jucys covers”. In: *Transactions of the American Mathematical Society* 373.7 (2020), pp. 4685–4711.
- [HL22] M. A. Hahn and D. Lewanski. “Tropical Jucys covers”. In: *Mathematische Zeitschrift* 301.2 (2022), pp. 1719–1738.
- [HM22] M. A. Hahn and H. Markwig. “Twisted Hurwitz numbers: Tropical and polynomial structures”. In: *arXiv preprint arXiv:2210.00595* (2022).
- [LO18] F. Liu and B. Osserman. “Severi degrees on toric surfaces”. In: *Journal für die reine und angewandte Mathematik (Crelles Journal)* 2018.739 (2018), pp. 121–158.
- [Mik05] G. Mikhalkin. “Enumerative tropical algebraic geometry in \mathbb{R}^2 ”. In: *Journal of the American Mathematical Society* 18.2 (2005), pp. 313–377.
- [Shu12] E. Shustin. “Tropical and algebraic curves with multiple points”. In: *Perspectives in Analysis, Geometry, and Topology: On the Occasion of the 60th Birthday of Oleg Viro* (2012), pp. 431–464.
- [Wic50] G.-C. Wick. “The evaluation of the collision matrix”. In: *Physical review* 80.2 (1950), p. 268.

M. A. HAHN: SCHOOL OF MATHEMATICS, 17 WESTLAND ROW, TRINITY COLLEGE DUBLIN, DUBLIN 2, IRELAND
Email address: hahnma@maths.tcd.ie

V. REDA: SCHOOL OF MATHEMATICS, 17 WESTLAND ROW, TRINITY COLLEGE DUBLIN, DUBLIN 2, IRELAND
Email address: redav@tcd.ie