

A uniform-in-time nonlocal approximation of the standard Fokker-Planck equation

José A. Cañizo & Niccolò Tassi

Abstract

We study a nonlocal approximation of the Fokker-Planck equation in which we can estimate the speed of convergence to equilibrium in a way which respects the local limit of the equation. This uniform estimate cannot be easily obtained with standard inequalities or entropy methods, but can be obtained through the use of Harris’s theorem, finding interesting links to quantitative versions of the central limit theorem in probability. As a consequence one can prove that solutions of this nonlocal approximation converge to solutions of the usual Fokker-Planck equation uniformly in time—hence we show the approximation is asymptotic-preserving in this sense. The associated equilibrium has some interesting tail and regularity properties, which we also study.

Contents

1	Introduction	2
2	Well-posedness of the initial value problem	9
2.1	Existence and Uniqueness	9
2.2	Semigroup results	13
2.3	Representation of the solution	14
3	Asymptotic behavior	17
3.1	Harris’s Theorem	17
3.2	Lyapunov condition	18
3.3	An L^∞ version of Berry-Esseen’s central limit theorem	21
3.4	Positivity of the solution	27
4	Convergence: from nonlocal to local	31
5	Analysis of the equilibrium: regularity, moments, cumulants and tails	35
5.1	Regularity	35

5.2	Moments and cumulants	38
5.3	Tails	41
	References	42

1 Introduction

We consider the linear partial differential equation

$$\partial_t u = \frac{1}{\varepsilon^2} (J_\varepsilon * u - u) + \operatorname{div}(xu) =: L_\varepsilon u, \tag{NLFP}$$

where $u: [0, +\infty) \times \mathbb{R}^d \rightarrow \mathbb{R}$ is the unknown whose variables we usually name by $u \equiv u(t, x)$, and $\varepsilon > 0$ is a scaling parameter. This PDE involves a nonnegative kernel $J: \mathbb{R}^d \rightarrow [0, +\infty)$ which we always assume at least integrable, and such that

$$\int_{\mathbb{R}^d} J(x) dx = 1, \quad \int_{\mathbb{R}^d} J(x)x dx = 0 \quad \int_{\mathbb{R}^d} J(x)x_i x_j dx = 2\delta_{ij}, \tag{1}$$

for $i, j = 1, \dots, d$. Making straightforward scaling arguments one can also apply our results if J is not of mass 1.

A canonical example of J for which our results are meaningful is any smooth, compactly supported probability distribution satisfying the above, which is a helpful case to keep in mind. The scaled kernel J_ε is defined by

$$J_\varepsilon(x) := \varepsilon^{-d} J\left(\frac{x}{\varepsilon}\right), \quad x \in \mathbb{R}^d,$$

and the convolution in (NLFP) is understood to be in the x variable. Thus (NLFP) is, strictly speaking, a partial integro-differential equation since it involves an integral kernel. We choose this particular scaling in ε since it is well known that as $\varepsilon \rightarrow 0$ the PDE (NLFP) resembles the standard Fokker-Planck PDE

$$\partial_t u = \Delta u + \operatorname{div}(xu) =: L_0 u. \tag{FP}$$

In contrast with the local diffusion, the nonlocal PDE (NLFP) does not have instantaneous regularization properties. We refer to the book by [Andreu-Vaillio et al. \(2010\)](#) for a good explanation of this limit. Hence, equation (NLFP) can be understood as a nonlocal version of the standard Fokker-Planck equation. Both PDE (NLFP) and (FP) have a stochastic interpretation in terms of a Markov process; we refer to standard texts such as [Risken \(1996\)](#) for this. In this paper we are interested in understanding the asymptotic behavior of solutions to (NLFP) as $t \rightarrow +\infty$ by the use of a new method based on Harris's theorem, since techniques for nonlocal PDE still have many shortcomings and this equation is a good test case. We pay particular attention to whether the asymptotic behavior of (NLFP) matches that of the standard Fokker-Planck equation

as $\varepsilon \rightarrow 0$. The use of Harris's theorem is by no means new in the study of nonlocal PDE, but as far as we know it is the first time that it can be applied to give a result which is stable under the scaling in (NLFP). We consider this the main novelty of this paper.

Obviously, since our results will be valid also for a fixed ε , we may frame the problem in a slightly different way by studying the PDE

$$\partial_t u = c(J * u - u) + \operatorname{div}(xu),$$

for J a general probability distribution satisfying (1). We have chosen to write the PDE as in (NLFP) in order to emphasise that we are especially interested in its scaling limit to the Fokker-Planck equation.

We are not aware of any studies regarding the specific PDE (NLFP) except for Mischler and Tristani (2017), whose results are similar to ours but use very different techniques. There are plenty of results on related equations, of which we give a short summary now, but we highlight that nonlocal-to-local limits which preserve the asymptotic behavior in the limit are comparatively rare, and are a current source of interest in several contexts (this is also the point of view of Mischler and Tristani (2017), where other nonlocal PDE are investigated as well).

The approach from nonlocal diffusion-type operators of the form $J * u - u$ to local diffusion operators has been the subject of many previous works. The PDE $\partial_t u = J * u - u$ is studied in Chasseigne et al. (2006), and a review of related techniques (making strong use of explicit solutions via the Fourier transform) is given in Andreu-Vaillo et al. (2010). Generalizations of this result to similar nonlocal PDE with different integral kernels are given in Molino and Rossi (2019). The same PDE is studied in Rey and Toscani (2013), with a focus on its large-time behavior. In this case the motivation comes from both physics (since this PDE is thought to be a more appropriate diffusion model in some cases) and numerical analysis (see below for more on this). Some results on its large-time behavior can be obtained by using Fourier-type distances (Carrillo and Toscani, 2007), again by taking advantage of the fact that the PDE is explicitly solvable in Fourier variables. A different nonlocal and nonlinear version of the Fokker-Planck PDE (FP) is studied in Ignat and Rossi (2007). Its relationship to the local problem is also studied there, but the results on asymptotic behavior are not preserved when passing to the local limit. Still a different nonlocal version has been recently considered in Auricchio et al. (2023), motivated by a model in collective behavior.

We mentioned that equation (NLFP) is a good test case for ideas to study the asymptotic behavior of nonlocal PDE, and we would like to describe a few more involved situations where this is relevant. An equation close to (NLFP) was proposed in Cai et al. (2006) as a model for the probability of the concentration of a certain protein which is being expressed by a particular gene. This equation was later studied in Cañizo et al.

(2019) by means of an entropy inequality developed specifically for this PDE, which highlights that a general strategy is lacking even for such simple nonlocal equations.

Another important domain where these ideas are relevant is numerical analysis. In dimension $d = 1$, if we formally take $J = \frac{1}{2}(\delta_{-1} + \delta_1)$ then the equation $\partial_t u = \frac{1}{\varepsilon^2}(J_\varepsilon * u - u)$, when evaluated at a set $\{x_i\}_{i=0,\dots,N}$ of equally spaced nodes at distance ε , becomes a numerical scheme for solving the standard heat equation on \mathbb{R} . The question of whether a numerical method preserves the large-time behavior of its limiting equation is of practical importance, and it is a subject of recent research [Ayi et al. \(2022\)](#), [Dujardin et al. \(2020\)](#), [Cancès et al. \(2020\)](#) and [Bessemoulin-Chatard et al. \(2020\)](#). We sometimes refer to this as the property of the scheme being *time asymptotic-preserving* (just “asymptotic-preserving” is often used for schemes which are well-behaved under a certain scaling limit of their associated PDE, which is not our case). Numerical schemes for PDE are nonlocal (in the sense that the equation at one of the nodes depends on the value at other nodes, which are at a fixed positive distance) and the study of their asymptotic behavior often encounters similar difficulties as for nonlocal PDE. This was one of the motivations of [Rey and Toscani \(2013\)](#), and we have the same hope that developing different methods for nonlocal PDEs can also suggest ideas to understand the behavior of some numerical methods.

We finally mention that several methods to study the asymptotic behavior of nonlocal PDE have been developed in the context of kinetic theory. Some examples for linear PDE include semigroup tools and spectral methods ([Gualdani et al., 2018](#)), techniques for hypocoercive PDE ([Villani, 2009](#); [Dolbeault et al., 2009](#)), applications of Harris’s theorem ([Cañizo et al., 2020](#)), and entropy techniques, which are often useful also for nonlinear PDE ([Lods et al., 2008](#); [Carrillo and Toscani, 2007](#); [Bisi et al., 2015](#)). Of these methods, the first one based on spectral methods is the basis of [Mischler and Tristani \(2017\)](#); entropy techniques are the main tool of [Cañizo et al. \(2019\)](#) on a very similar PDE; and as far as we know our paper is the first time Harris-based techniques are used to get the correct behavior as $\varepsilon \rightarrow 0$. It does not seem straightforward to use entropy techniques to get a similar result, since an inequality along the lines of [Cañizo et al. \(2019\)](#) does not respect the scaling as $\varepsilon \rightarrow 0$, but finding a way to do so is in our opinion an interesting problem.

Let us describe more precisely the results we are able to show. We denote by $B_r(x_0)$, the ball in \mathbb{R}^d centered at x_0 with radius r . When the ball is centered at 0, we use B_r as a shorthand. We always denote by L^k_1 the usual weighted Lebesgue space with norm

$$\|\varphi\|_{L^k_1} = \int_{\mathbb{R}^d} (1 + |x|^k) |\varphi(x)| \, dx$$

or with the equivalent norm

$$\|\varphi\|_{L^k_{\langle x \rangle}} = \int_{\mathbb{R}^d} \langle x \rangle^k |\varphi(x)| \, dx,$$

where $\langle x \rangle^k := (1 + |x|^2)^{k/2}$, a “smoothed k -th power” which avoids regularity problems at $x = 0$. We denote by $\langle x \rangle$ the smoothed 1st power. Similarly we define $L_{\text{exp},a}^1$ as the Lebesgue space with weight $e^{a|x|}$ with natural norm

$$\|\varphi\|_{\text{exp},a} = \int_{\mathbb{R}^d} e^{a|x|} |\varphi(x)| \, dx$$

or with equivalent norm

$$\|\varphi\|_{\langle \text{exp} \rangle, a} = \int_{\mathbb{R}^d} e^{a\langle x \rangle} |\varphi(x)| \, dx.$$

An *equilibrium* of equation (NLFP) is a function $F_\varepsilon \in L^1$ which satisfies the stationary PDE

$$\frac{1}{\varepsilon^2} (J_\varepsilon * F - F) + \text{div}(xF) = 0$$

in the distributional sense. A *probability equilibrium* is a nonnegative equilibrium F_ε with $\int_{\mathbb{R}^d} F_\varepsilon(x) \, dx = 1$. It is easy to see that there exists at most one probability equilibrium (see Theorem (10)) and under mild additional conditions we also show existence (and additional estimates). Here is the main theorem we prove in this paper regarding asymptotic behavior of mild solutions.

Theorem 1.1. *Let $\varepsilon \in (0, 1]$ and let $J: \mathbb{R}^d \rightarrow [0, +\infty)$ satisfy (1) and, moreover, $J \in L^p$ for some $p \in (1, \infty]$. Then,*

- (i) *If $J \in L_{k'}^1$ for some $k' > 2$, then there exists a unique equilibrium F_ε of (NLFP) in $L_{k'}^1$. Additionally, for every $0 < k \leq k'$ there exist (explicit) constants $C \geq 1$ and $\gamma > 0$, independent of ε , such that any (mild) solution u of (NLFP) with a probability initial condition $u_0 \in L_k^1$ satisfies*

$$\|u(t, \cdot) - F_\varepsilon\|_{L_k^1} \leq C e^{-\gamma t} \|u_0 - F_\varepsilon\|_{L_k^1} \quad \text{for every } t \geq 0.$$

- (ii) *If $J \in L_{\text{exp},a'}^1$, for some $a' > 0$, then there exists a unique equilibrium F_ε of (NLFP) in $L_{\text{exp},a}^1$, for all $a < a'$. Additionally, there exist (explicit) constants $C \geq 1$ and $\gamma > 0$, independent of ε , such that any (mild) solution u of (NLFP) with a probability initial condition $u_0 \in L_{\text{exp},a}^1$ satisfies*

$$\|u(t, \cdot) - F_\varepsilon\|_{L_{\text{exp},a}^1} \leq C e^{-\gamma t} \|u_0 - F_\varepsilon\|_{L_{\text{exp},a}^1} \quad \text{for every } t \geq 0.$$

We remark that the unique equilibrium F_ε also belongs to the (smaller, but ε -dependent) space $L_{\text{exp},b}^1$ for any $b < a'\varepsilon^{-1}$.

In fact, from our proof of the theorem (or directly as a simple consequence) we get the same estimates substituting $u - F_\varepsilon$ by any solution w with initial mass 0 (i.e. $\int_{\mathbb{R}^d} w_0(x) \, dx = 0$). For example, we prove that under the conditions of point (i),

$$\|w(t, \cdot)\|_{L_k^1} \leq C e^{-\gamma t} \|w_0\|_{L_k^1} \quad \text{for every } t \geq 0.$$

This is indeed more general than our statement since one can take $w_0 = u_0 - F_\varepsilon$, but we choose to state the theorem as above for readability.

We remark that both points (i) and (ii) of the previous theorem, and all of our results concerning asymptotic behaviour, require in particular that there exists $s \in (0, 1]$ such that

$$\int_{\mathbb{R}^d} J(x)|x|^{2+s} dx =: \rho_{2+s} < \infty. \quad (2)$$

This quantity ρ_{2+s} will be fundamental for our estimates in Section 3.

Proving an estimate like the previous one has several consequences. One of these is the following uniform-in-time result on the closeness solutions to the local and nonlocal problems, which we prove in Section 4. For readability, we state the theorem with likely non-optimal hypotheses, since optimising it is not the main goal of this paper. It requires the following additional hypothesis on J :

$$\int_{\mathbb{R}^d} J(x)x^\alpha = 0 \quad \text{for all multiindices } \alpha \text{ of order } |\alpha| = 3. \quad (3)$$

(Which is satisfied for example if J is symmetric.)

Theorem 1.2 (Nonlocal-to-local approximation). *Take $k \geq 0$ and assume $J \in L^1_{k+4+d}$ satisfies (1) and (3), $J \in L^p$ for some $p \in (1, \infty]$. Let u be the mild solution to equation (NLFP) with a probability initial condition $u_0 \in L^1_k \cap \mathcal{C}^4$, and such that there exist $c > 0$ and $K > k + d$ with $|D^\alpha u_0(x)| \leq c \langle x \rangle^{-K}$ for all multiindices α with $|\alpha| = 4$.*

Let v the (unique) integrable solution to the (local) Fokker-Planck equation (FP) with the same initial condition. Then, there exists $C > 0$ such that

$$\|u(\cdot, t) - v(\cdot, t)\|_{L^1_k} \leq C\varepsilon^2 \quad \text{for every } t \geq 0 \text{ and } \varepsilon \in (0, 1]. \quad (4)$$

We emphasize that the constant C is independent of ε .

Remark 1.3. If we do not assume (3), then under slightly weaker assumptions ($u_0 \in L^1_k \cap \mathcal{C}^3$, $J \in L^1_{k+3+d}$) one can show that

$$\|u(\cdot, t) - v(\cdot, t)\|_{L^1_k} \leq C\varepsilon \quad \text{for every } t \geq 0 \text{ and } \varepsilon \in (0, 1], \quad (5)$$

for some (other) constant $C > 0$ depending only on d and u_0 . In order to obtain this it is enough to carry out the Taylor expansions in Section 4 to one order less.

Remark 1.4. With the same strategy one can prove the previous statement also in Lebesgue spaces with exponential weight. Assume $J \in L^1_{\text{exp}, a'}$ and $u_0 \in L^1_{\text{exp}, a'} \cap \mathcal{C}^3$. Then (5) holds also in $L^1_{\text{exp}, a}$ for every $a < a'/2$. Moreover if (3) holds and $u_0 \in \mathcal{C}^4$, then equation (4) is also valid in $L^1_{\text{exp}, a}$, for every $a < a'/2$. We don't give the proofs, since they are essentially the same as the ones in L^1_k spaces.

Throughout the paper (as was used in the theorem above), we will use multi-index notation for derivatives, so if $\alpha \in \mathbb{N}^d$ is a multi-index we write $|\alpha| = \sum_i \alpha_i$, $x^\alpha = x_1^{\alpha_1} x_2^{\alpha_2} \cdots x_d^{\alpha_d}$, $\alpha! = \alpha_1! \alpha_2! \cdots \alpha_d!$ and $D^\alpha \varphi(x) = \partial_{x_1}^{\alpha_1} \partial_{x_2}^{\alpha_2} \cdots \partial_{x_d}^{\alpha_d} \varphi(x)$.

We emphasise once more that, although being far from optimal, the constants we find are all explicit, since proofs are completely constructive.

We will also make a few remarks regarding the unique probability equilibrium F_ε associated to (NLFP). Theorems 1.1 and 1.2 imply in particular that $\|F_\varepsilon - G\|_{L_k^1} \leq C\varepsilon$, where G is the standard Gaussian probability distribution in \mathbb{R}^d . But the properties of F_ε are quite interesting in their own right. Theorem 1.1 includes the fact that F_ε has exponentially-decaying tails as long as J has the same property. However, when J is compactly supported we suspect that F_ε has tails which decay like the Poisson distribution, that is $F_\varepsilon \approx e^{-a|x| \log|x|}$ for large $|x|$. We are able to show an incomplete version of this statement in Section 5.3. We suspect that in general F_ε does *not* have Gaussian tails, but we haven't been able to prove this.

The regularity of F_ε is also interesting, as we explore in Section 5.1: independently of the regularity of J , F_ε appears to be smooth everywhere except at $x = 0$; and its regularity at $x = 0$ depends on ε , not on the regularity or tails of J . The equilibrium F_ε becomes more regular as $\varepsilon \rightarrow 0$. We think a more complete study of the properties of F_ε would be interesting for a future work.

Strategy of the proof The central point of our results is the proof of (i) and (ii) in Theorem 1.1. The proof of Theorem 1.2 can then be obtained as a consequence with fairly standard estimates. As mentioned, the proof of Theorem 1.1 uses Harris's theorem from probability, which requires us to show a certain positivity property of solutions of our nonlocal equation (NLFP); see Section 3.1, where we give a short introduction to this. Showing that solutions are positive for any *fixed* $\varepsilon > 0$ is not hard; and the fact that solutions to the (local) Fokker-Planck equation (FP) are positive is well known, and a consequence of its regularization properties and the parabolic Harnack inequality (or just a consequence of its explicit solution!). However, finding a positivity result which is valid for all $\varepsilon > 0$ and which does not degenerate in the limit $\varepsilon \rightarrow 0$ is not an easy task. We notice that it is linked to convergence results in the central limit theorem, since the solution u to (NLFP) can be written as (see Section 2.3)

$$u(t)(x) = \sum_{n=0}^{\infty} \left(\frac{1}{\varepsilon^2}\right)^n \int_0^t \int_0^{t_1} \cdots \int_0^{t_{n-1}} J_\varepsilon^{t_1, \dots, t_n} * u_0(e^t x) dt_n \cdots dt_1,$$

where it is understood that the term for $n = 0$ is just $u_0(e^t x)$, and

$$J_\varepsilon^{t_1, \dots, t_n} := J_{\varepsilon e^{t_1}} * \cdots * J_{\varepsilon e^{t_n}}(x).$$

Each of the terms represents the distribution of particles which have undergone n random jumps, if we see (NLFP) as the PDE for the distribution of a jump stochastic

process. Proving positivity of u can then be translated to the problem of understanding the behavior of repeated convolutions of scalings of the function J , which can in turn be understood in terms of the central limit theorem in probability. This can be made into a rigorous proof, as we show in this paper. We think it provides an intriguing link among these different tools, and it yields a result which is otherwise non-obvious to obtain.

Structure of the paper In Section 2, we present results related to the well-posedness of the equation (NLFP) in L_k^1 . For this purpose we define the *mild solutions* of (NLFP) by a standard use of Duhamel's formula, for which we prove existence, uniqueness and continuous dependence on the initial data via the well known fixed-point strategy. We point out that the same strategy can be used to prove well-posedness in measure spaces, and interpret these results also in terms of semigroups.

In the last part of Section 2 we give two explicit representations of the solution: a classical one via Fourier transform, and the Wild sum representation which we use in a crucial way.

Section 3 contains the proof of Theorem 1.1. We give a short introduction to Harris's Theorem, which requires two main conditions: a Lyapunov confinement condition which is fairly straightforward in our case, based on a Taylor's expansion; and a positivity condition which is the key part of our argument, as we discussed above. Proving this positivity condition requires a suitable version of the Berry-Esseen Theorem, a form of the Central Limit Theorem which gives a rate of convergence in the L^∞ norm. We need a result for independent, not necessarily identically distributed, random vectors. Afterwards, the proof of positivity is straightforward once a suitable scaling is carried out.

Section 4 deals with the speed of convergence of the Nonlocal Fokker Planck equation (NLFP) to the local one (FP) at all time t , and gives the proof of Theorem 1.2. The rate of convergence at finite times comes from a standard consistency and stability argument in the spirit of numerical methods. The latter result, combined with Theorem 1.1, allows us to extend the convergence for all times.

In the last section 5 we gather some independent results on the regularity of the equilibrium, showing that it improves as $\varepsilon \rightarrow 0$. Moreover, we give a representation of the solution of (NLFP) in terms of the cumulative generating function, which is nothing else than the logarithm of the moment generating function. This gives in particular an explicit representation for the moments of the equilibrium. We also investigate on the tails of the solution of (NLFP) and its equilibrium F_ε . In the local case, the equilibrium is a Gaussian and the solution has Gaussian tails if the initial data has. For the nonlocal equation (NLFP) we strongly suspect this is not the case, even for very nice initial data and kernel. We are able to prove that for compactly supported J the tails are at least Poisson-like.

2 Well-posedness of the initial value problem

We gather in this section some results on the well-posedness of equation (NLFP), that is,

$$\partial_t u = \frac{1}{\varepsilon^2} (J_\varepsilon * u - u) + \operatorname{div}(xu),$$

with the addition of an initial condition $u_0 \in L_k^1$.

$$u(0, x) = u_0(x), \quad x \in \mathbb{R}^d.$$

Though they are mostly standard results for linear PDE, it is useful to lay out the basic theory.

2.1 Existence and Uniqueness

Solutions to equation (NLFP) can be studied (for example) by using semigroup theory, which easily yields mild solutions $u \in \mathcal{C}([0, \infty), L_k^1(\mathbb{R}^d))$ whenever $u_0 \in L_k^1$, for some $k \geq 0$. Alternatively, we give a direct simple proof of existence of mild solutions. The first observation is that the equation

$$\partial_t u = \operatorname{div}(xu) - \frac{1}{\varepsilon^2} u, \quad u(0, x) = u_0(x)$$

has the explicit classical solution

$$u(t, x) = e^{(d-1/\varepsilon^2)t} u_0(e^t x), \quad t \geq 0, x \in \mathbb{R}^d,$$

at least for a differentiable u_0 . Hence, for any $u_0 \in L^1$ we define

$$T_t[u_0](x) \equiv T_t u_0(x) := e^{(d-1/\varepsilon^2)t} u_0(e^t x) \quad \text{for } t \geq 0, x \in \mathbb{R}^d. \quad (6)$$

Following a well known strategy, we use Duhamel's formula to define mild solutions of equation (NLFP) as follows:

Definition 2.1 (Mild solution). Let $k \geq 0$, and take J satisfying Hypothesis (1) on J and $J \in L_k^1$. Take $u_0 \in L_k^1$, and take $T^* \in (0, +\infty]$. A *mild solution* of equation (8) on $[0, T^*)$ with initial condition u_0 is a function $u \in \mathcal{C}([0, T^*), L_k^1)$ such that

$$u(t) = T_t u_0 + \frac{1}{\varepsilon^2} \int_0^t T_{t-s} [J_\varepsilon * u(s)] ds \quad \text{for all } t \in I,$$

where the integral is understood (equivalently) in the Riemann or Bochner sense.

2.1.1 Existence of L_k^1 solutions

We are going to show the following result:

Theorem 2.2. Take $k \geq 0$ and let J satisfy (1) and, in addition, $J \in L^k_1$. Then, there exists a unique mild solution of equation (NLFP) with initial condition $u_0 \in L^1_k$ (in the sense of Definition 2.1). In addition, the solution is continuously dependent on the initial data, in the following sense: for any two mild solutions u and v of (NLFP) with initial data u_0 and v_0

$$\|u(t) - v(t)\|_{L^1_k} \leq e^{\frac{t}{\varepsilon^2}(c_k - 1)} \|u_0 - v_0\|_{L^1_k}$$

The proof of this is fairly straightforward, and we give it after a couple of lemmas. We first notice that the convolution operator $u \mapsto J * u$ is a continuous operator under fairly mild conditions. For any measurable functions $u: \mathbb{R}^d \rightarrow \mathbb{R}$ and $V: \mathbb{R}^d \rightarrow [1, +\infty)$ we denote

$$\|u\|_V := \int_{\mathbb{R}^d} |u(x)|V(x) dx.$$

We need the following generalization of Young's inequality in weighted spaces

Lemma 2.3. Take a measurable $J: \mathbb{R}^d \rightarrow \mathbb{R}$ such that $\|J\|_V \|u\|_V < +\infty$. Then for every sub-additive function $V: \mathbb{R}^d \rightarrow [1, \infty)$ (that is, satisfying for all $x, y \in \mathbb{R}^d$ $V(x + y) \leq C(V(x) + V(y))$ for some $C > 0$) it holds that

$$\|J * u\|_V \leq 2C \|J\|_V \|u\|_V$$

for all measurable $u: \mathbb{R}^d \rightarrow \mathbb{R}$ such that $\|u\|_V < +\infty$. In particular, for $V(x) := 1 + |x|^k$ with $k \geq 0$ we have

$$\|J * u\|_{L^1_k} \leq 2^k \|J\|_{L^1_k} \|u\|_{L^1_k}. \quad (7)$$

Proof. We write a direct estimate as follows:

$$\begin{aligned} \|J * u\|_V &= \int_{\mathbb{R}^d} \left| \int_{\mathbb{R}^d} J(x - y)u(y) dx \right| V(x) dx \\ &\leq C \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} |J(x - y)|u(y)(V(x - y) + V(y)) dy dx \\ &= C(\|u\|_V \|J\|_1 + \|u\|_1 \|J\|_V) \leq 2C \|J\|_V \|u\|_V, \end{aligned}$$

where in the last inequality we used the assumption that $V \geq 1$ at all points. The last statement is a consequence of this, since $V(x) = 1 + |x|^k$ satisfies for all $x, y \in \mathbb{R}^d$ that

$$V(x + y) = 1 + |x + y|^k \leq 1 + 2^{k-1}(|x|^k + |y|^k) \leq 2^{k-1}(V(x) + V(y))$$

where we used that the convexity of $|\cdot|^k$ implies

$$\left| \frac{x + y}{2} \right|^k \leq \frac{|x|^k + |y|^k}{2} \implies |x + y|^k \leq 2^{k-1}(|x|^k + |y|^k).$$

□

We also notice that for any $t \geq 0$, the operator T_t is contractive in all L_k^1 for any $k \geq 0$:

Lemma 2.4 (Contractivity). *For all $k \geq 0$, $t > 0$ and $u \in L_k^1$, the operator T_t defined in (6) satisfies*

$$\|T_t u\|_{L_k^1} \leq e^{-t/\varepsilon^2} \|u\|_{L_k^1}.$$

Proof. We just write the following direct estimate:

$$\begin{aligned} \|T_t u\|_{L_k^1} &= \int (|x|^k + 1) |T_t u(x)| dx = e^{-t/\varepsilon^2} \int (|x|^k + 1) |u(xe^t)| e^{dt} dx \\ &= e^{-t/\varepsilon^2} \int (|e^{-t}y|^k + 1) |u(y)| dy < \int (|y|^k + 1) u(y) dy. \end{aligned}$$

□

Proof of Theorem 2.2. This is done via a standard contraction argument. Take $T^* > 0$ and define the Banach space $\mathcal{Y} := \mathcal{C}([0, T^*], L_k^1)$ equipped with the norm

$$\|u\|_{\mathcal{Y}} := \sup_{t \in [0, T^*]} \|u(t)\|_{L_k^1}.$$

Let us define $\Psi : \mathcal{Y} \rightarrow \mathcal{Y}$ by

$$\Psi[u](t) := T_t u_0 + \frac{1}{\varepsilon^2} \int_0^t T_{t-s} [J_\varepsilon * u(s)] ds$$

for any $u \in \mathcal{Y}$, so that u is a mild solution if and only if u is a fixed point for Ψ . In the following estimate u_s and v_s will be used as shorthand for $u(s)$ and $v(s)$. For every $u_t, v_t \in L_k^1$, use Lemmas 2.3 and 2.4 we have

$$\begin{aligned} \|\Psi[u](t) - \Psi[v](t)\|_{L_k^1} &= \left\| \frac{1}{\varepsilon^2} \int_0^t T_{t-s} [J_\varepsilon * (u_s - v_s)] ds \right\|_{L_k^1} \\ &\leq \frac{1}{\varepsilon^2} \int_0^t \|T_{t-s} [J_\varepsilon * (u_s - v_s)]\|_{L_k^1} ds \\ &\leq \frac{1}{\varepsilon^2} \int_0^t e^{-\frac{t-s}{\varepsilon^2}} \|J_\varepsilon * (u_s - v_s)\|_{L_k^1} ds \\ &\leq \frac{1}{\varepsilon^2} c_k \int_0^t \|u_s - v_s\|_{L_k^1} ds \\ &\leq \frac{1}{\varepsilon^2} c_k t \|u - v\|_{\mathcal{Y}}, \end{aligned}$$

where $c_k = 2^k \|J\|_{L_k^1} \geq 2^k \|J_\varepsilon\|_{L_k^1}$ from Lemma 2.3. Iterating the process, we obtain

$$\|\Psi^n[u](t) - \Psi^n[v](t)\|_{L_k^1} \leq \left(\frac{c_k t}{\varepsilon^2}\right)^n \frac{1}{n!} \|u - v\|_{\mathcal{Y}} \leq \left(\frac{c_k T^*}{\varepsilon^2}\right)^n \frac{1}{n!} \|u - v\|_{\mathcal{Y}}$$

and then, taking the supremum, leads to

$$\|\Psi^n[u](t) - \Psi^n[v](t)\|_{\mathcal{Y}} \leq \left(\frac{c_k T^*}{\varepsilon^2}\right)^n \frac{1}{n!} \|u - v\|_{\mathcal{Y}}$$

For every T^* , there exists $n \in \mathbb{N}$ such that $(\frac{c_k T^*}{\varepsilon^2})^n \frac{1}{n!} < 1$. There is a specific version of the contraction mapping theorem (see for example [Miklavcic \(1998\)](#), Thm. 1.1.3) which now implies Ψ has a fixed point which is a solution of (NLFP) in the sense of Definition 2.1. (Alternatively, one can consider the sequence $(\Psi^n(u_0))_{n \geq 0}$, where $u_0 \in \mathcal{Y}$ is the function constantly equal to u_0 on $[0, T^*]$; our last inequality easily shows that it is a Cauchy sequence, and hence must have a limit. This can be done for any $T^* > 0$, easily obtaining the result on $[0, +\infty)$.)

To prove continuity with respect to initial data we proceed as before: let v be another mild solution for some other initial data v_0 , then for every t

$$\begin{aligned} \|u(t) - v(t)\|_{L_k^1} &\leq \|T_t(u_0 - v_0)\|_{L_k^1} + \frac{1}{\varepsilon^2} \int_0^t \|T_{t-s}[J_\varepsilon * (u_s - v_s)]\|_{L_k^1} ds \\ &\leq e^{-t/\varepsilon^2} \|u_0 - v_0\|_{L_k^1} + \frac{c_k}{\varepsilon^2} \int_0^t e^{-\frac{t-s}{\varepsilon^2}} \|u(s) - v(s)\|_{L_k^1} ds. \end{aligned}$$

By the standard Gronwall inequality applied to $e^{\frac{t}{\varepsilon^2}} \|u(t) - v(t)\|_{L_k^1}$ we obtain the result. \square

2.1.2 Existence of measure solutions

We sketch similar arguments that give existence of solutions in a space of measures, since the Harris theorem we use later is usually stated for semigroups in a space of measures (or Markov semigroups in the context of probability). This is really not strictly necessary, since there are versions of the Harris theorem which hold similarly in weighted L^1 spaces (see [Cañizo and Mischler \(2021\)](#)), but since the proof is very similar we give the main steps.

Let us denote with $\mathcal{M}(\mathbb{R}^d)$ the space of signed finite Radon measures over \mathbb{R}^d and by $\|\cdot\|_{TV}$ and $\|\cdot\|_{\mathcal{M}_k}$ the total variation norm and the weighted total variation, respectively. Let us recall the definition of Bounded Lipschitz norm, i.e.

$$\|\mu\|_{BL} := \sup_{f \in \mathcal{F}} \left| \int_{\mathbb{R}^d} f d\mu \right|$$

where $\mathcal{F} := \{f \in W^{1,\infty}(\mathbb{R}^d) : \|f\|_{1,\infty} \leq 1\}$, with $\|f\|_{1,\infty} := \|f\|_\infty + \text{Lip}(f)$ and $\text{Lip}(f)$ denoting the Lipschitz constant of f . In the literature, the bounded Lipschitz norm is sometimes defined as the equivalent norm obtained by choosing in the previous definition $\mathcal{F} := \{f : \mathbb{R}^d \rightarrow \mathbb{R} : \|f\|_\infty \leq 1, \sup_{x \neq y} \frac{|f(x) - f(y)|}{|x - y|} \leq 1\}$.

We define the ball $\mathcal{X}_R := \{\mu \in \mathcal{M} : \|\mu\|_{\mathcal{M}_k} \leq R\}$ and the set $\mathcal{Z} = \mathcal{C}([0, T^*], \mathcal{X}_R)$ of functions which are continuous from $[0, T^*]$ to \mathcal{X}_R in the bounded Lipschitz norm. \mathcal{Z} is a complete metric space with the metric induced by the norm

$$\|\mu\|_{\mathcal{Z}} = \sup_{t \in [0, T^*]} \|\mu(t)\|_{BL}.$$

We stress that the same claim in the space $\mathcal{C}([0, \infty), \mathcal{M}_k)$ does not hold since in $(W^{1, \infty})^*$ there are elements that are not measures which can be approximated by Cauchy sequences in \mathcal{M}_k .

Following the path of the previous proof, we can prove existence and uniqueness of the mild solution of equation (NLFP) in \mathcal{Z} choosing a suitable R such that $\mu_0 \in \mathcal{X}_R$. One can prove a version of inequality (7) in weighted total variation norms:

$$\|J * \mu\|_{\mathcal{M}_k} \leq c_k \|\mu\|_{\mathcal{M}_k}$$

Moreover, the contractivity Lemma 2.4 still holds in the sense of measures in weighted total variation norm. The functional Ψ , defined as

$$\Psi[\mu](t) = T_t \mu_0 + \frac{1}{\varepsilon^2} \int_0^t T_{t-s} [J_\varepsilon * \mu_s] ds$$

is such that if $t \rightarrow \mu(t) \in \mathcal{Z}$, then so is $t \rightarrow \Psi[\mu](t)$ for small enough T^* . We briefly show a proof of this. Continuity comes straightforwardly from the fact that T_t is a continuous semigroup (in the bounded Lipschitz norm) and all the other operations are continuity-preserving.

We have left to show that $\Psi[\mu](t) \in \mathcal{X}_R$ for every $t \in [0, T^*]$ when $\mu(t) \in \mathcal{X}_R$. Set $R = 2 \|u_0\|_{\mathcal{M}_k}$

$$\begin{aligned} \|\Psi[\mu](t)\|_{\mathcal{M}_k} &\leq \|T_t \mu_0\|_{\mathcal{M}_k} + \frac{1}{\varepsilon^2} \int_0^t \|T_{t-s} [J_\varepsilon * \mu_s]\|_{\mathcal{M}_k} ds \\ &\leq \|\mu_0\|_{\mathcal{M}_k} + \frac{1}{\varepsilon^2} \int_0^t C_k \|\mu_s\|_{\mathcal{M}_k} ds \leq \|\mu_0\|_{\mathcal{M}_k} + \frac{1}{\varepsilon^2} C T^* \|\mu\|_{\mathcal{M}_k} \leq R \end{aligned}$$

for $T^* \leq \frac{\varepsilon^2}{2C_k}$. We then iterate the process for another time T_1^* until we extend the existence and uniqueness globally.

With the same approach, one can prove the previous results in the space $L_{\text{exp}, a}^1$. Since the procedure is the same *mutatis mutandis*, we have chosen to omit it.

2.2 Semigroup results

Our result on the well-posedness of the nonlocal Fokker-Planck equation (NLFP) can also be phrased in terms of semigroup theory. For an introduction and further details on the results we will be using below, we refer the reader to the books Engel and Nagel (2001) or Pazy (1983). We first point out that for any $k \geq 0$, the operators $(T_t)_{t \geq 0}$ given in (6) define a strongly continuous semigroup on the Banach space L_k^1 ; it is easy to check this directly. The generator of this semigroup is the operator B given by

$$B_\varepsilon u := \text{div}(xu) - \frac{1}{\varepsilon^2} u$$

with domain

$$\mathcal{D}(B_\varepsilon) := \{u \in L_k^1 \mid \text{div}(xu) \in L_k^1\},$$

which is a closed linear operator. Then, equation (NLFP) can be written as

$$\frac{d}{dt}u = L_\varepsilon u = A_\varepsilon u + B_\varepsilon u, \quad (8)$$

where L_ε and A_ε denote the operators

$$\begin{aligned} A_\varepsilon u &:= \frac{1}{\varepsilon^2} J_\varepsilon * u, \\ L_\varepsilon u &:= A_\varepsilon u + B_\varepsilon u = \frac{1}{\varepsilon^2} (J_\varepsilon * u - u) + \operatorname{div}(xu). \end{aligned}$$

Due to Lemma 2.3 we know that $A_\varepsilon: L_k^1 \rightarrow L_k^1$ is a bounded linear operator. Since we know B_ε is a generator in L_k^1 , standard results for bounded perturbations in semigroup theory (see for example Engel and Nagel (2001, chapter III.1)) ensure then that the operator $L_\varepsilon = A_\varepsilon + B_\varepsilon$ is also a generator with domain $\mathcal{D}(L_\varepsilon) = \mathcal{D}(B_\varepsilon)$. This is equivalent to the well-posedness of equation (8) in the sense of semigroups (Engel and Nagel, 2001, Chapter II.6). Solutions to (8) in the sense of semigroups must also be mild solutions in the sense of Definition 2.1 (Engel and Nagel, 2001, Section III.1.3, Corollary 1.4), so they coincide with the solutions given in Theorem 2.2.

2.3 Representation of the solution

For this kind of problems, we can express the solution via the classic Fourier transform or with the so-called Wild sum. We use the latter as an important tool in our later proofs, emphasizing that our main results do not seem easy to obtain by using the explicit representation in the terms of the Fourier transform. More importantly, we would like to explore strategies which would be valid even in cases in which Fourier transform methods are not available.

For the sake of completeness and to obtain regularity results in section 5.1, we will also briefly comment on the Fourier solution. From (1) we deduce that

$$\hat{J}(\xi) = 1 - \xi^2 + o(|\xi|^2) \quad \text{as } \xi \rightarrow 0.$$

Then if we take u_0 as an initial probability distribution in L^1 , we can prove that there exists at most one solution of (NLFP) in $\mathcal{C}([0, \infty), L^1)$. If it exists, it must satisfy

$$\hat{u}(t, \xi) = \hat{u}_0(e^{-t}\xi) e^{\frac{1}{\varepsilon^2} \int_0^t \zeta_\varepsilon(e^{-s}\xi) ds} \quad (9)$$

with $\zeta_\varepsilon(\xi) = \hat{J}_\varepsilon(\xi) - 1$. Moreover, there exists at most one equilibrium in L^1 of (NLFP), and it must satisfy

$$\hat{F}_\varepsilon(\xi) = e^{\frac{1}{\varepsilon^2} \int_0^\infty \zeta_\varepsilon(e^{-s}\xi) ds} \quad (10)$$

Indeed, passing to Fourier variables, equation (NLFP) becomes

$$\partial_t \hat{u} = \frac{1}{\varepsilon^2} (\hat{J}_\varepsilon \hat{u} - \hat{u}) - \xi \cdot \nabla_\xi \hat{u} = \frac{1}{\varepsilon^2} \zeta_\varepsilon(\xi) \hat{u} - \xi \cdot \nabla_\xi \hat{u}$$

which has equation (9) as its unique solution. We also notice that the transform of the equilibrium must satisfy

$$\hat{F}_\varepsilon(\xi) = \hat{u}(r, \xi) = \hat{F}_\varepsilon(e^{-r}\xi) e^{\frac{1}{\varepsilon^2} \int_0^r \zeta_\varepsilon(e^{-s}\xi) ds} \quad \text{for every } r \geq 0$$

with the initial condition $F_\varepsilon(0) = 1$, since $\|F_\varepsilon\|_{L^1} = 1$. In particular

$$\hat{F}_\varepsilon(\xi) = \lim_{r \rightarrow \infty} \hat{F}_\varepsilon(e^{-r}\xi) e^{\frac{1}{\varepsilon^2} \int_0^r \zeta_\varepsilon(e^{-s}\xi) ds} = e^{\frac{1}{\varepsilon^2} \int_0^\infty \zeta_\varepsilon(e^{-s}\xi) ds}.$$

Remark 2.5. With a more careful study of these arguments or other classical ones (like Tychonoff's Theorem) one could prove existence of a solution and of the equilibrium. We don't want to insist on these details since in the next section we will give a different proof for the existence and uniqueness of the equilibrium.

Wild Sum When dealing with linear convolution-type operators, Wild Sum—or Dyson-Phillips's Series (Engel and Nagel, 2001, p. 163)—give a particularly simple way to express the solution, based on a “Picard iteration” for PDEs.

We first present the following lemma that, for a fixed t , allows us to exchange the order of the operator T_t with the convolution operator and vice versa.

Lemma 2.6. *Let (T_t) be the semigroup associated to the solution of equation (6). Then*

$$J_\varepsilon * (T_t f) = T_t [J_{\varepsilon e^t} * f]$$

Proof. For every x

$$\begin{aligned} (J_\varepsilon * T_t f)(x) &= \int \varepsilon^{-d} J\left(\frac{x-y}{\varepsilon}\right) (T_t f)(y) dy = e^{(d-\frac{1}{\varepsilon^2})t} \int \varepsilon^{-d} J\left(\frac{x-y}{\varepsilon}\right) f(e^t y) dy \\ &= e^{-\frac{t}{\varepsilon^2}} \int \varepsilon^{-d} J\left(\frac{e^{-t}}{\varepsilon}(e^t x - z)\right) f(z) dz \\ &= \int e^{(d-\frac{1}{\varepsilon^2})t} (e^{-dt} \varepsilon^{-d}) J\left(\frac{e^t x - z}{e^t \varepsilon}\right) f(z) dz = e^{(d-\frac{1}{\varepsilon^2})t} (J_{\varepsilon e^t} * f)(e^t x) \end{aligned} \quad (11)$$

where we have used the change of variable $z = e^t y$. Hence,

$$J_\varepsilon * T_t f = T_t [J_{\varepsilon e^t} * f]. \quad \square$$

Theorem 2.7. *Let u be the unique solution to equation (NLFP) with initial condition $u_0 \in L^1$. Then, in the hypotheses of Theorem 2.2, we can write the solution of the equation via Wild sum, namely for every $t \geq 0$*

$$u(t)(x) = e^{(d-\frac{1}{\varepsilon^2})t} \left[u_0(e^t x) + \sum_{n=1}^{\infty} \left(\frac{1}{\varepsilon^2}\right)^n \int_0^t \int_0^{t_1} \dots \int_0^{t_{n-1}} J_\varepsilon^{t_1, \dots, t_n} * u_0(e^t x) dt_n \dots dt_1 \right]$$

where the convergence in the RHS is understood to be in L^1 for every t , and

$$J_\varepsilon^{t_1, \dots, t_n} := J_{\varepsilon e^{t_1}} * \dots * J_{\varepsilon e^{t_n}}(x).$$

for $0 \leq t_1 \leq \dots \leq t_n \leq t$.

Proof. Define

$$\begin{aligned} h(t) &:= T_{-t}u(t) = T_{-t}\left[T_t u_0 + \int_0^t T_{t-s}\left[\frac{1}{\varepsilon^2}J_\varepsilon * u(s)\right] ds\right] \\ &= u_0 + \frac{1}{\varepsilon^2} \int_0^t T_{-s}[J_\varepsilon * u(s)] \end{aligned}$$

Using Lemma 2.6

$$T_{-s}[J_\varepsilon * u_s] = T_{-s}[J_\varepsilon * T_s(T_{-s}u)] = T_{-s}[T_s J_{\varepsilon e^s} * h(s)] = J_{\varepsilon e^s} * h(s)$$

This means that u is a mild solution for Equation (NLFP) iff

$$h(t) = u_0 + \frac{1}{\varepsilon^2} \int_0^t J_{\varepsilon e^s} * T_{-s}u(s) ds = u_0 + \frac{1}{\varepsilon^2} \int_0^t J_{\varepsilon e^s} * h(s) ds,$$

that is if h is a fixed point for the functional Ψ

$$\Psi(g) := u_0 + \frac{1}{\varepsilon^2} \int_0^t J_{\varepsilon e^s} * g_s ds$$

To find an expression for this fixed point we perform a Picard Iteration defining $h_0 = u_0$ and

$$h_{n+1} = \Psi(h_n) \tag{12}$$

i.e.

$$h_n(t) = h_{n-1}(t) + \frac{1}{\varepsilon^{2n}} \int_0^t \int_0^{t_1} \dots \int_0^{t_{n-1}} J_\varepsilon^{t_1, \dots, t_n} * u_0 dt_n \dots dt_1,$$

where we use as a shorthand the following

$$J_\varepsilon^{t_1, \dots, t_n} := J_{\varepsilon e^{t_1}} * J_{\varepsilon e^{t_2}} * \dots * J_{\varepsilon e^{t_n}}$$

Clearly the sequence $(h_n)_n$ is pointwise monotone increasing. Since the functions h_n are all nonnegative and bounded above by h , the monotone convergence theorem shows that h_n converges in L^1 to some function $\tilde{h} = \tilde{h}(t) \geq 0$ when $n \rightarrow \infty$, namely

$$\tilde{h}(t) = u_0 + \sum_{n=1}^{\infty} \left(\frac{1}{\varepsilon^2}\right)^n \int_0^t \int_0^{t_1} \dots \int_0^{t_{n-1}} J_\varepsilon^{t_1, \dots, t_n} * u_0 dt_n \dots dt_1.$$

Passing to the limit in (12), one obtains that \tilde{h} is a fixed point for the functional Ψ , so in fact $\tilde{h} = h$. Therefore,

$$\begin{aligned} u(t)(x) &= (T_t h)(x) = e^{(d-1/\varepsilon^2)t} u_0(e^t x) + \\ &+ e^{(d-1/\varepsilon^2)t} \sum_{n=1}^{\infty} \varepsilon^{-2n} \int_0^t \int_0^{t_1} \dots \int_0^{t_{n-1}} \left(J_\varepsilon^{t_1, \dots, t_n} * u_0\right)(e^t x) dt_n \dots dt_1. \quad \square \end{aligned}$$

Remark 2.8. From the previous is immediate to see that if u_0 and J are radially symmetric functions then also the solution is radially symmetric. One can also say this of any property which is preserved by convolution with J . Analogously, the equilibrium will preserve the same properties.

3 Asymptotic behavior

In this section we study the asymptotic behavior of equation (NLFP). For this we use Harris’s theorem from probability, establishing exponential convergence to equilibrium of solutions in \mathcal{M}_k for $k > 0$. Our crucial advantage of our approach is that it yields a uniform bound on the convergence rate to equilibrium for all $\varepsilon \in (0, 1]$.

3.1 Harris’s Theorem

Let us first give a short overview of Harris’s theorem, in order to make this paper as self-contained as possible. We refer again to Hairer and Mattingly (2011); Cañizo and Mischler (2021) and the book by Meyn and Tweedie (2010) for a more complete exposition.

Let us denote by $\mathcal{M} \equiv \mathcal{M}(\Omega)$, the space of finite signed measures over a measurable space Ω , with $\|\mu\| = \|\mu\|_{TV} = \int |\mu|$, the total variation norm. \mathcal{P} will denote the set of probability measures in \mathcal{M} . If $V : \Omega \rightarrow [1, \infty)$ is a measurable function (which we call a *weight function*), we denote by \mathcal{M}_V the subspace of measures $\mu \in \mathcal{M}$ such that the norm $\|\mu\|_V = \int V|\mu| < \infty$. We will use \mathcal{P}_V to denote $\mathcal{M}_V \cap \mathcal{P}$.

A linear operator $S : \mathcal{M} \rightarrow \mathcal{M}$ is *stochastic* if it leaves \mathcal{P} invariant. A stochastic semigroup $(S_t)_{t \geq 0}$ is a family of stochastic operators S_t parameterised by t such that $S_0 = \text{Id}$ and $S_t \circ S_s = S_{t+s}$ for all $t, s \geq 0$.

We say that S is a stochastic operator on \mathcal{M}_V if it is a stochastic operator on \mathcal{M} and its restriction on \mathcal{M}_V is bounded in the $\|\cdot\|_V$ norm. Similarly, S_t is a stochastic semigroup on \mathcal{M}_V if it is a stochastic semigroup on \mathcal{M} and each S_t is a stochastic operator on \mathcal{M}_V .

A basic tool in the study of the asymptotic behavior of Markov Processes is Doeblin’s Theorem that under a (strong) positivity hypothesis gives existence of a unique stationary state, which is additionally exponentially stable. A well known generalization of it is Harris’s Theorem where Doeblin’s condition is replaced by a weaker local positivity condition and a Lyapunov confinement, which we state next.

Hypothesis 1 (Semigroup Lyapunov condition). *Let $V : \Omega \rightarrow [1, \infty)$ be a measurable (“weight”) function and let $(S_t)_{t \geq 0}$ be a stochastic semigroup on \mathcal{M}_V . We say that $(S_t)_{t \geq 0}$ satisfies a semigroup Lyapunov condition if there exist constants $\lambda, C > 0$ such that*

$$\|S_t \mu\|_V \leq e^{-\lambda t} \|\mu\|_V + \frac{C}{\lambda} (1 - e^{-\lambda t}) \|\mu\|. \quad (\text{SLC})$$

If S_t is a semigroup with generator L , a natural and better known version of this condition is to impose

$$L^* V \leq C - \lambda V \quad (13)$$

where L^* is the (formal) dual operator of L acting on “nice enough” functions. The latter condition has the technical burden of appropriately defining L^* (see for example

Hairer (2010)), where inequality (13) must be understood in a submartingale sense). Our approach is really the same as this, but we prefer to use Hypothesis 1, which does not have any technical issues. We actually show condition (13), in a suitable sense, for the generator associated to equation (NLFP) in Theorems 3.2 and 3.4. From this we will deduce Hypothesis 1.

The second condition needed is the *Harris condition*:

Hypothesis 2 (Harris condition). *A stochastic operator S on \mathcal{M} satisfies a Harris condition on a set $\mathcal{S} \subseteq \Omega$ if there exist $0 < \alpha < 1$ and $\rho \in \mathcal{P}$ such that*

$$S\mu \geq \alpha\rho \int_{\mathcal{S}} \mu \quad \text{for every } 0 \leq \mu \in \mathcal{M}. \quad (\text{HC})$$

We will use the following version of Harris's Theorem:

Theorem 3.1 (Harris's theorem for semigroups.). *Let $V : \mathbb{R}^d \rightarrow [1, \infty)$ be a measurable function and let $(S_t)_{t \geq 0}$ be a stochastic semigroup on \mathcal{M}_V . Assume that:*

1. *the semigroup $(S_t)_{t \geq 0}$ satisfies the semigroup Lyapunov condition (SLC), and*
2. *for some $T^* > 0$ and some $R > \frac{2C}{\lambda}$, the operator S_{T^*} satisfies the Harris condition (HC) on the set $\mathcal{S} := \{x : V(x) \leq R\}$.*

Then the semigroup has an invariant probability $\mu^ \in \mathcal{P}_V$ which is unique in \mathcal{P}_V , and there exist $C, \lambda > 0$ such that*

$$\|S_t\nu\|_V \leq Ce^{-\lambda t} \|\nu\|_V \quad \text{for all } \nu \in \mathcal{P}_V \text{ with } \int_{\Omega} \nu = 0 \text{ and all } t \geq 0,$$

and in particular

$$\|S_t\mu - \mu^*\|_V \leq Ce^{-\lambda t} \|\mu - \mu^*\|_V \quad \text{for all } \mu \in \mathcal{P}_V \text{ and } t \geq 0.$$

Theorem 1.1 automatically follows once we prove the two conditions of Harris's Theorem with weight $V(x) = \langle x \rangle^k$. We notice that, since solutions to (NLFP) with L_k^1 initial data remain in L_k^1 for all times (and the \mathcal{M}_k norm in L_k^1 is just equal to the L_k^1 norm), Theorem 1.1 in L_k^1 really follows once we are able to apply Theorem 3.1 in \mathcal{M}_k . A similar consideration holds with exponential weights. Hence, the next two sections are devoted to proving the hypotheses required in Theorem 3.1 for the semigroup generated by equation (NLFP).

3.2 Lyapunov condition

We define L_ε^* , the formal dual operator of L_ε , acting on functions $\varphi \in \mathcal{C}^2$ (with appropriate growth as $|x| \rightarrow +\infty$) as

$$L_\varepsilon^*\varphi := A_\varepsilon^*\varphi + B_\varepsilon^*\varphi = \frac{1}{\varepsilon^2}[J_\varepsilon * \varphi - \varphi] - x\nabla\varphi.$$

This ‘‘appropriate growth’’ is just to ensure the convolution $J_\varepsilon * \varphi$ is properly defined. Let $(S_t)_{t \geq 0}$ be the semigroup associated to equation (NLFP).

Theorem 3.2. *Let J satisfy (1) and let $J \in L^1_{k'}$, for $k' \geq 2$. Then for every $0 < k \leq k'$ there exist constants $C, \lambda > 0$ depending only on k and J such that*

$$L_\varepsilon^*(\langle x \rangle^k) \leq C - \lambda \langle x \rangle^k$$

for any $0 < \varepsilon \leq 1$.

Proof. To lighten the notation in the following, we will use C_1, C_2 as general constants, whose value may change from line to line. Recall from (1), that for every α such that $|\alpha| = 1$,

$$\int J(z) z^\alpha dz = 0$$

Therefore, for every $\varphi \in \mathcal{C}^\infty$, expanding around x ,

$$\begin{aligned} \frac{1}{\varepsilon^2} [J_\varepsilon * \varphi - \varphi] &= \frac{1}{\varepsilon^2} \int_{\mathbb{R}^d} J(z) (\varphi(x - \varepsilon z) - \varphi(x)) dz \\ &= \frac{1}{\varepsilon^2} \int_{\mathbb{R}^d} J(z) \left(\varphi(x) + \frac{1}{2} \sum_{|\alpha|=2} R_\alpha(\xi) \varepsilon^2 z^\alpha - \varphi(x) \right) dz \\ &= \frac{1}{2} \int_{\mathbb{R}^d} J(z) \sum_{i=1}^d \partial_i^2 \varphi(\xi) z_i^2 dz \end{aligned}$$

where $R_\alpha(\xi)$ is the Lagrange remainder for some $\xi(x, z) \in [x, x - \varepsilon z]$, the segment that joins x and $x - \varepsilon z$. Set $\varphi(x) = \langle x \rangle^k$.

With a simple computation

$$\partial_i^2 \varphi(x) = k(\langle x \rangle^{k-2} + (k-2)x_i^2 \langle x \rangle^{k-4}) \quad (14)$$

For $k \in (0, 2)$ $\partial_i^2 \varphi(\xi) \leq k$, for every i, ξ . Hence

$$\frac{1}{\varepsilon^2} [J_\varepsilon * \langle x \rangle^k - \langle x \rangle^k] \leq \frac{k}{2} \int_{\mathbb{R}^d} J(z) |z|^2 dz = k$$

For $k \geq 2$, since $\langle \cdot \rangle^j$ is a radially nondecreasing *subadditive* function for every $j \geq 0$, we can bound (14), by

$$\partial_i^2 \varphi(\xi) \leq k(k-1) \langle \xi \rangle^{k-2} \leq k(k-1) \langle |x| + \varepsilon |z| \rangle_{k-2} \leq C_1 (\langle x \rangle^{k-2} + \langle z \rangle^{k-2})$$

It follows that

$$\begin{aligned} \frac{1}{\varepsilon^2} [J_\varepsilon * \langle x \rangle^k - \langle x \rangle^k] &\leq \frac{1}{2} \int_{\mathbb{R}^d} J(z) \sum_{i=1}^d z_i^2 C_1 (\langle x \rangle^{k-2} + \langle z \rangle^{k-2}) dz \\ &\leq C_1 \langle x \rangle^{k-2} \frac{1}{2} \int_{\mathbb{R}^d} J(z) |z|^2 dz + C_1 \int_{\mathbb{R}^d} J(z) \langle z \rangle^k dz \leq C_2 \langle x \rangle^{k-2} \end{aligned}$$

Consequently, for every $k \geq 0$

$$\begin{aligned} L_\varepsilon^* \varphi &= \frac{1}{\varepsilon^2} [J_\varepsilon * \varphi - \varphi] - x \nabla \varphi \leq C_2 \langle x \rangle^{k-2} - k|x|^2 \langle x \rangle^{k-2} \\ &= (C_2 + k) \langle x \rangle^{k-2} - k(1 + |x|^2) \langle x \rangle^{k-2} \leq C - \lambda \langle x \rangle^k \end{aligned} \quad (15)$$

for some $C, \lambda > 0$.

□

Remark 3.3. We point out that as k approaches 0, the RHS of (15) goes to 0 as well, since we don't have Lyapunov confinement in Lebesgue spaces without weight.

With almost the same procedure, we can prove Lyapunov condition in $L_{\text{exp},a}^1$.

Theorem 3.4. *Assume $J \in L_{\text{exp},a'}^1$ for some $a' > 0$. Then for every $0 < a < a'$ there exist $C, \lambda > 0$ (only depending on a, d and J), such that*

$$L_\varepsilon^*(e^{a\langle x \rangle}) \leq C - \lambda e^{a\langle x \rangle}.$$

Proof. To lighten the notation in the following, we will use C_1, C_2 as general constants, whose value may change from line to line. Again we notice that for every α such that $|\alpha| = 1$

$$\int J(z) z^\alpha = 0$$

As before, for every $\varphi \in \mathcal{C}^\infty$, we Taylor expand around x , obtaining

$$\frac{1}{\varepsilon^2} [J_\varepsilon * \varphi - \varphi] = \frac{1}{2\varepsilon^2} \int_{\mathbb{R}^d} J(z) \sum_{i=1}^d \partial_i^2 \varphi(\xi) \varepsilon^2 z_i^2 dz$$

where the Lagrange remainder is evaluated at some $\xi(x, z) \in [x, x - \varepsilon z]$.

Set now $\varphi = e^{a\langle x \rangle}$.

One can see that there exists $C_1 > 0$ a dimension dependent constant, such that

$$\partial_i^2 \varphi(\xi) = \frac{ae^{a\langle \xi \rangle}}{\langle \xi \rangle_2} \left(a\xi_i^2 + \langle \xi \rangle - \frac{\xi_i^2}{\langle \xi \rangle} \right) \leq C_1 e^{a\langle \xi \rangle} \leq C_1 e^{a\langle \xi \rangle} \leq C_1 e^{a(|x| + \varepsilon|z|)} \leq C_1 e^{a(\langle x \rangle + \varepsilon\langle z \rangle)}$$

since φ is a symmetric, positive and increasing function in every component and,

$$\langle |x| + \varepsilon|y| \rangle \leq \langle x \rangle + \varepsilon\langle y \rangle.$$

Thus

$$\begin{aligned} \frac{1}{\varepsilon^2} [J_\varepsilon * \varphi - \varphi] &\leq \frac{1}{2} \int_{\mathbb{R}^d} J(z) \sum_{i=1}^d C_1 e^{\langle |x| + \varepsilon|z| \rangle} z_i^2 dz \\ &\leq \frac{C_1}{2} e^{a\langle x \rangle} \int_{\mathbb{R}^d} J(z) e^{a\varepsilon\langle z \rangle} |z|^2 dz \leq C_2 e^{a\langle x \rangle} \end{aligned}$$

since

$$\int_{\mathbb{R}^d} J(z) |z|^2 e^{a\varepsilon\langle z \rangle} dz \leq C_3 \int_{\mathbb{R}^d} J(z) e^{a'\langle z \rangle} dz < \infty.$$

(Let us point out that this is where the remark at the end of Theorem 1.1 comes from).
Then

$$L_\varepsilon^* e^{a\langle x \rangle} = \frac{1}{\varepsilon^2} [J_\varepsilon * \varphi - \varphi] - x \nabla \varphi \leq C_2 e^{a\langle x \rangle} - a \frac{|x|^2}{\langle x \rangle} e^{a\langle x \rangle} \leq C - \lambda e^{a\langle x \rangle}$$

for some $C, \lambda > 0$. □

We thus have proved Lyapunov condition on the generator for $V = \langle \cdot \rangle^k$ and $V = e^{a\langle \cdot \rangle}$. For these weight functions (and in generally for \mathcal{C}^2 functions), we see that actually Assumption (13) implies Assumption (SLC), which is the statement of the next Theorem.

Theorem 3.5 (Semigroup Lyapunov condition). *Assume u_0 and J satisfy the hypothesis of Theorem 3.2 (respectively 3.4). For $\varphi = \langle x \rangle^k$ (respectively $\varphi = e^{a\langle x \rangle}$), the semigroup associated to equation (NLFP) satisfies a Lyapunov condition (SLC) with $V \equiv \varphi$ and the same C and λ of Theorem 3.2 (respectively 3.4).*

Proof. We just prove it for L_k^1 , being analogous in the other case.

Let u_0 be in $D(L)$ by standard semigroup theory $u(t) \in \mathcal{C}^1([0, \infty); L_k^1)$, and consequently we compute

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}^d} u(t, \cdot) \varphi(x) dx &= \int_{\mathbb{R}^d} \frac{d}{dt} u(t, \cdot) \varphi(x) dx \\ &= \int_{\mathbb{R}^d} (Lu) \varphi(x) dx \stackrel{*}{=} \int_{\mathbb{R}^d} u(L^* \varphi) \leq \int_{\mathbb{R}^d} u(-\lambda \varphi + C) \end{aligned}$$

where to prove $\stackrel{*}{=}$ we use classic cutoff arguments with bump functions.

We then use Gronwall Lemma with the \mathcal{C}^1 function $f(t) := \int_{\mathbb{R}^d} u(t) \varphi$: That is,

$$f'(t) \leq -\lambda f(t) + C \implies f(t) \leq e^{-\lambda t} f(0) + \frac{C}{\lambda} (1 - e^{-\lambda t})$$

Since the $D(L)$ is dense in L_k^1 , the claim is true also for any $u_0 \in L_k^1$ by classic density argument. □

3.3 An L^∞ version of Berry-Esseen's central limit theorem

Berry-Esseen-type theorems are quantitative versions of the central limit theorem which provide an estimate on the error between the normal distribution and the scaled law. A result of this kind will be crucial for our proof in Section 3.4 of positivity for the nonlocal Fokker-Planck equation. The first version of this theorem was proved independently by Berry (1941) and Esseen (1942) for real, centred i.i.d. random variables X_1, \dots, X_n with $\text{Var}(X_1) = \sigma^2$ and $\mathbb{E}(|X_1|^3) = \rho$, obtaining a result of the type

$$|F_n(x) - \Phi(x)| \leq \frac{C\rho}{\sqrt{n}\sigma^3}$$

where Φ is the cumulative distribution function (c.d.f.) of the normal distribution, F_n is the c.d.f. of $\bar{S}_n = \frac{1}{\sigma\sqrt{n}} \sum_{i=1}^n X_i$, and C is a universal constant depending on the dimension d . Several generalisations have been developed, proving the same kind of results for not i.i.d. random vectors in \mathbb{R}^d , as well as improving the precision of the value of C . Recent versions can be found in [Bentkus \(2005\)](#) and [Raič \(2019\)](#).

The majority of this kind of results are “weak” results in the sense that they deal with the cumulative distribution function. For our results we would like to work with the density instead. An important result in this direction is [Lions and Toscani \(1995\)](#) in which the authors provide a stronger version of the CLT on the densities, and later in [Goudon et al. \(2002\)](#), [Carlen et al. \(2010\)](#), [Hauray and Mischler \(2014\)](#). For this paper we need an estimate on the densities of independent but not identically distributed random vectors in \mathbb{R}^d , so we will extend the proofs of the previous cited papers to our case. Following the ideas in [Goudon et al. \(2002\)](#), we weaken the hypothesis of finiteness of the third moment. We require only a “bit more” than finite variance, paying a price on the speed of convergence to zero of the error.

Let $f : \mathbb{R}^d \rightarrow [0, \infty)$ be a probability density function such that there exists $s \in (0, 1]$ with

$$\int_{\mathbb{R}^d} x f(x) dx = 0, \quad \int_{\mathbb{R}^d} x_i x_j f(x) dx = \delta_{ij}, \quad \int_{\mathbb{R}^d} |x|^{2+s} f(x) dx := \rho_{2+s} < \infty. \quad (16)$$

Notice that this normalisation of moments of order 2 is different from (1) by a factor of 2. We prefer the present normalisation in this section, which is a natural one in a probabilistic setting, and is the same one used in most statements of Berry-Esseen’s theorem. By a simple scaling it is straightforward to apply the result if one assumes the normalisation (1) instead.

The main result of this section is the following L^∞ version of Berry-Esseen’s Theorem for independent, non-identically distributed random vectors:

Theorem 3.6. *Let $p \in (1, \infty)$ and $f \in L^1(\mathbb{R}) \cap L^p(\mathbb{R})$ satisfying (16). Let $(\sigma_i)_{i \geq 1}$ be a sequence of positive numbers, and define for $n \geq 1$,*

$$\begin{aligned} f^{\sigma_1, \dots, \sigma_n}(x) &:= f_{\sigma_1} * f_{\sigma_2} * \dots * f_{\sigma_n}(x) \\ \mathfrak{f}_n(x) &:= (\bar{\sigma}^2 n)^{d/2} f^{\sigma_1, \dots, \sigma_n}(\sqrt{n} \bar{\sigma} x), \end{aligned}$$

with $\bar{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n \sigma_i^2$. Suppose, in addition, that there exist $l, L > 0$ such that

$$0 < l \leq \sigma_i \leq L \quad i = 1, \dots, n$$

Then there exist an integer $N = N(p)$ and a constant C_{BE} depending on $\frac{L}{l}, p, d, \rho_{2+s}, \|f\|_{L^p}$ such that $\mathfrak{f}_n \in L^\infty$ for all $n \geq N$ and

$$\|\mathfrak{f}_n - G\|_{L^\infty} \leq \frac{C_{BE}}{n^{s/2}} \quad \text{for every } n \geq N,$$

where G is the density of the standard normalised Gaussian.

In order to prove this we first present some preliminary results. The first one is a generalisation of the identity $a^n - b^n = \sum_{i=1}^n (a - b)a^{i-1}b^{n-i}$, that can be easily proven by induction:

Lemma 3.7. *For real a_1, \dots, a_n and b_1, \dots, b_n , for any $n \geq 1$*

$$\prod_{i=1}^n a_i - \prod_{i=1}^n b_i = \sum_{i=1}^n \left[(a_i - b_i) \left(\prod_{j=1}^{i-1} b_j \right) \left(\prod_{j=i+1}^n a_j \right) \right]$$

The second one is a Hölder regularity result for the Fourier transform.

Lemma 3.8. *Let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be a probability distribution such that for some $s \in (0, 1)$,*

$$\int_{\mathbb{R}^d} f(x) |x|^{2+s} dx = \rho_{2+s} < +\infty.$$

Then $\hat{f} \in \mathcal{C}^{2,s}$; that is, $\hat{f} \in \mathcal{C}^2$ and for every $x, y \in \mathbb{R}^d$

$$\max_{|\alpha|=2} |D^\alpha \hat{f}(x) - D^\alpha \hat{f}(y)| \leq L|x + y|^s,$$

with $L := 4\rho_{2+s}$.

Proof. The \mathcal{C}^2 -regularity is immediate from Fourier transform properties. We are left to show the Hölder-continuity of the derivatives of order 2, namely, for every multi-index α such that $|\alpha| = 2$,

$$\begin{aligned} \left| D^\alpha \hat{f}(\xi) - D^\alpha \hat{f}(\eta) \right| &= \left| -\widehat{x^\alpha f}(\xi) + \widehat{x^\alpha f}(\eta) \right| = \left| \int_{\mathbb{R}^d} (e^{-ix\xi} - e^{-ix\eta}) x^\alpha f(x) dx \right| \\ &\leq \int_{|x| \leq R} |e^{-ix\xi} - e^{-ix\eta}| |x|^2 |f(x)| dx + \int_{|x| > R} |e^{-ix\xi} - e^{-ix\eta}| |x|^2 |f(x)| dx \\ &\leq 2|\xi - \eta| \int_{|x| \leq R} |x|^{2-s} |x|^{1-s} |f(x)| dx + 2R^{-s} \int_{|x| > R} |x|^{2+s} |f(x)| dx \\ &\leq 2\rho_{2+s} (|\xi - \eta| R^{1-s} + R^{-s}). \end{aligned}$$

Choosing $R = |\xi - \eta|^{-1}$ we conclude the proof. \square

The last ingredient we need is the following technical Lemma, that is basically (Hauray and Mischler, 2014, lemma 4.8) with a weaker assumption on moments.

Lemma 3.9. *Let f be a probability distribution on \mathbb{R}^d satisfying (16). Then*

(i) *There exist $\delta \in (0, 1)$ depending on ρ_{2+s} such that*

$$\forall \xi \in B_\delta(0) \quad |\hat{f}(\xi)| \leq e^{-\frac{|\xi|^2}{4}}$$

(ii) *Assume additionally that $f \in L^p$, with $p \in (1, \infty]$. For any $\delta > 0$ there exists $\kappa = \kappa(\rho_{2+s}, \delta, \|f\|_{L^p}) \in (0, 1)$ such that*

$$\sup_{|\xi| \geq \delta} |\hat{f}(\xi)| \leq \kappa(\delta).$$

Proof. (i) For $\hat{f} : \mathbb{R}^d \rightarrow \mathbb{R}$ and for ξ define $h : [0, |\xi|] \rightarrow \mathbb{R}$ such that $h(t) := \hat{f}\left(t \frac{\xi}{|\xi|}\right)$. Applying twice the fundamental theorem of calculus to h we write

$$\hat{f}(\xi) = h(|\xi|) = 1 - \frac{|\xi|^2}{2} + \int_0^{|\xi|} (h''(z) - h''(0))(z - |\xi|) dz$$

where

$$h''(z) = \left(\frac{\xi}{|\xi|}\right)^T H_{\hat{f}}\left(z \frac{\xi}{|\xi|}\right) \left(\frac{\xi}{|\xi|}\right).$$

Hence, applying Lemma 3.8, we can bound the reminder

$$\begin{aligned} & \left| \int_0^{|\xi|} \left(\frac{\xi}{|\xi|}\right)^T \left(H_{\hat{f}}\left(z \frac{\xi}{|\xi|}\right) - H_{\hat{f}}(0)\right) \left(\frac{\xi}{|\xi|}\right) (z - |\xi|) dz \right| \\ & \leq \int_0^{|\xi|} \left\| \left(H_{\hat{f}}\left(z \frac{\xi}{|\xi|}\right) - H_{\hat{f}}(0)\right) \right\| |z - |\xi|| dz \\ & \leq \int_0^{|\xi|} L |z|^s |z - |\xi|| dz = \frac{L |\xi|^{2+s}}{(s+1)(s+2)}, \end{aligned}$$

where $H_f(x)$ is the Hessian matrix of f applied at point x and $\|\cdot\|$ is the spectral norm which is always less or equal than the max- norm used in Lemma 3.8.

Thus, for small $|\xi| < \delta$ which explicitly depends only on s and ρ_{2+s} we have

$$\left| \hat{f}(\xi) \right| \leq \left| 1 - \frac{3}{8} |\xi|^2 \right| \leq e^{-\frac{|\xi|^2}{4}}.$$

- (ii) This statement is a direct consequence of (Carlen et al., 2010, Prop. 26 (iii)), where the assumption of finiteness of the entropy is ensured by our assumption that $f \in L^1 \cap L^p$ and has some finite moment. \square

We can now present the proof of our version of Berry-Esseen theorem, that follows the ideas presented in (Goudon et al., 2002, Thm. 1) and (Hauray and Mischler, 2014, Thm 4.6).

Proof of Theorem 3.6. First, we notice that \mathfrak{f}_n satisfies the hypotheses of Lemma 3.9 for every n , and, in particular

$$\int_{\mathbb{R}^d} \mathfrak{f}_n(x) x_i x_k = \delta_{ij}.$$

We can rewrite

$$\begin{aligned} \hat{\mathfrak{f}}_n(\xi) &= \int_{\mathbb{R}^d} e^{-ix\xi} (\bar{\sigma}^2 n)^{d/2} f^{\sigma_1, \dots, \sigma_n}(\sqrt{n}\bar{\sigma}x) dx = \int_{\mathbb{R}^d} e^{-iy \frac{\xi}{\sqrt{n}\bar{\sigma}}} f_{\sigma_1} * \dots * f_{\sigma_n}(y) dy \\ &= \mathcal{F}(f_{\sigma_1} * \dots * f_{\sigma_n})\left(\frac{\xi}{\sqrt{n}\bar{\sigma}}\right) = \prod_{i=1}^n \hat{f}_{\sigma_i}\left(\frac{\xi}{\sqrt{n}\bar{\sigma}}\right) = \prod_{i=1}^n \int_{\mathbb{R}^d} e^{-ix \frac{\xi}{\sqrt{n}\bar{\sigma}}} \sigma_i^{-d} f\left(\frac{x}{\sigma_i}\right) dx \\ &= \prod_{i=1}^n \int_{\mathbb{R}^d} e^{-iy \frac{\xi \sigma_i}{\sqrt{n}\bar{\sigma}}} f(y) dy = \prod_{i=1}^n \hat{f}\left(\frac{\xi \sigma_i}{\sqrt{n}\bar{\sigma}}\right), \end{aligned}$$

and

$$\hat{G}(\xi) = \prod_{i=1}^n \hat{G}\left(\frac{\xi \sigma_i}{\sqrt{n\bar{\sigma}}}\right).$$

Without loss of generality, we can assume $p \leq 2$. Hausdorff-Young's inequality then implies that \hat{f} is in $L^{p'} \cap L^\infty$ with $p' \in [2, \infty)$. Using Hölder's inequality,

$$\left\| \prod_{i=1}^n \hat{f}\left(\frac{\xi \sigma_i}{\sqrt{n\bar{\sigma}}}\right) \right\|_{L^1} \leq \prod_{i=1}^n \left\| \hat{f}\left(\frac{\xi \sigma_i}{\sqrt{n\bar{\sigma}}}\right) \right\|_{L^n} \leq \prod_i (\bar{\sigma} \sqrt{n})^{d/n} \|\hat{f}\|_{L^n} \leq n^{d/2} \left(\frac{L}{l}\right)^d \|\hat{f}\|_{L^n}^n,$$

which is finite for $n \geq p'$, i.e. \hat{f}_n is in L^1 for any $n \geq p'$. We can thus control the L^∞ norm with L^1 norm of the Fourier transform, since

$$|\hat{f}_n(x) - G(x)| = (2\pi)^d \left| \int_{\mathbb{R}^d} (\hat{f}_n(\xi) - \hat{G}(\xi)) e^{i\xi x} d\xi \right| \leq (2\pi)^d \int_{\mathbb{R}^d} |\hat{f}_n - \hat{G}| d\xi$$

Let δ be given by Lemma 3.9 (i) for \hat{f} and define $B_\delta = B_\delta(n, l, L) := \{\xi : |\xi| < \sqrt{n} \frac{L}{l} \delta\}$. Following known techniques, we split the integral into high and low frequencies

$$\|\hat{f}_n - G\|_{L^\infty} \leq \int_{\xi \in B_{\delta, n}^c} |\hat{f}_n| d\xi + \int_{\xi \in B_{\delta, n}^c} |\hat{G}| d\xi + \int_{\xi \in B_{\delta, n}} |\hat{f}_n - \hat{G}| d\xi := T_1 + T_2 + T_3$$

The first two terms can be bounded in a similar way. For T_1 , and without loss of generality, assume p' is an integer (otherwise we can work with $\lceil p' \rceil$, the smallest integer larger or equal than p' , since \hat{f} is also in $L^{\lceil p' \rceil}$). Then

$$\begin{aligned} \int_{B_\delta^c} \prod_{i=1}^n \left| \hat{f}\left(\frac{\xi \sigma_i}{\sqrt{n\bar{\sigma}}}\right) \right| d\xi &= n^{d/2} \int_{|\eta| \geq \frac{L}{l} \delta} \prod_{i=1}^n \left| \hat{f}\left(\frac{\sigma_i}{\bar{\sigma}} \eta\right) \right| d\eta \\ &\leq n^{d/2} \sup_{|\eta| \geq \frac{L}{l} \delta} \left| \hat{f}\left(\frac{\sigma_{p'+1}}{\bar{\sigma}} \eta\right) \right| \cdots \sup_{|\eta| \geq \frac{L}{l} \delta} \left| \hat{f}\left(\frac{\sigma_n}{\bar{\sigma}} \eta\right) \right| \int_{|\eta| \geq \frac{L}{l} \delta} \left| \hat{f}\left(\frac{\sigma_1}{\bar{\sigma}} \eta\right) \right| \cdots \left| \hat{f}\left(\frac{\sigma_{p'}}{\bar{\sigma}} \eta\right) \right| d\eta. \end{aligned} \quad (17)$$

For each $i = p' + 1, \dots, n$

$$\sup_{|\eta| \geq \frac{L}{l} \delta} \left| \hat{f}\left(\frac{\sigma_i}{\bar{\sigma}} \eta\right) \right| \leq \sup_{|\eta| \geq \frac{\bar{\sigma}}{\sigma_i} \delta} \left| \hat{f}\left(\frac{\sigma_i}{\bar{\sigma}} \eta\right) \right| = \sup_{|\eta'| \geq \delta} |\hat{f}(\eta')| \leq \kappa(\delta) \quad (18)$$

with $\kappa(\delta) \in (0, 1)$, coming from Lemma 3.9 (i). The factors inside the integral in Equation (17) can be bounded by using Hölder's inequality: we rewrite

$$\begin{aligned} &\int_{|\eta| \geq \frac{L}{l} \delta} \left| \hat{f}\left(\frac{\sigma_1}{\bar{\sigma}} \eta\right) \right| \cdots \left| \hat{f}\left(\frac{\sigma_{p'}}{\bar{\sigma}} \eta\right) \right| d\eta \\ &\leq \left(\int_{\mathbb{R}^d} \left| \hat{f}\left(\frac{\sigma_1}{\bar{\sigma}} \eta\right) \right|^{p'} d\eta \right)^{1/p'} \cdots \left(\int_{\mathbb{R}^d} \left| \hat{f}\left(\frac{\sigma_{p'}}{\bar{\sigma}} \eta\right) \right|^{p'} d\eta \right)^{1/p'} \\ &= \left(\frac{\sigma_1}{\bar{\sigma}}\right)^{d/p'} \|\hat{f}\|_{p'} \cdots \left(\frac{\sigma_{p'}}{\bar{\sigma}}\right)^{d/p'} \|\hat{f}\|_{p'} \leq \left(\frac{L}{l}\right)^d \|\hat{f}\|_{p'}^{p'} \leq \left(\frac{L}{l}\right)^d C_p \|f\|_p^{p'}, \end{aligned} \quad (19)$$

after first applying Hölder's inequality and then Hausdorff-Young's inequality (C_p is the Hausdorff-Young constant). Hence, plugging (18) and (19) into equation (17) we obtain

$$T_1 \leq C_p \left(\sqrt{n} \frac{L}{l} \right)^d \kappa(\delta)^{n-p'} \|f\|_p^{p'}. \quad (20)$$

Proceed similarly for T_2 to get that there exists C_1 such that

$$T_1 + T_2 \leq \frac{C_1}{n^{s/2}} \quad \text{for } n > p'.$$

Now we focus our attention on T_3 . Using the factorisation in Lemma 3.7 we obtain

$$\begin{aligned} \left| \hat{\mathbf{f}}_n(\xi) - \hat{G} \right| &= \prod_{i=1}^n \hat{f} \left(\frac{\xi \sigma_i}{\sqrt{n\bar{\sigma}}} \right) - \prod_{i=1}^n \hat{G} \left(\frac{\xi \sigma_i}{\sqrt{n\bar{\sigma}}} \right) \\ &= \sum_{i=1}^n \left| \hat{f} \left(\frac{\xi \sigma_i}{\sqrt{n\bar{\sigma}}} \right) - \hat{G} \left(\frac{\xi \sigma_i}{\sqrt{n\bar{\sigma}}} \right) \right| \left| \prod_{j=1}^{i-1} \hat{f} \left(\frac{\xi \sigma_j}{\sqrt{n\bar{\sigma}}} \right) \right| \left| \prod_{j=i+1}^n \hat{G} \left(\frac{\xi \sigma_j}{\sqrt{n\bar{\sigma}}} \right) \right|. \end{aligned}$$

From Lemma 3.9 (i) it follows that for $\xi \in B_\delta$

$$\left| \hat{f} \left(\frac{\xi \sigma_j}{\sqrt{n\bar{\sigma}}} \right) \right| \leq e^{-\frac{|\xi|^2 \sigma_j^2}{4n \bar{\sigma}^2}} \leq e^{-\frac{|\xi|^2}{4n} \left(\frac{l}{L} \right)^2}.$$

and

$$\left| \hat{G} \left(\frac{\xi \sigma_j}{\sqrt{n\bar{\sigma}}} \right) \right| = (2\pi)^{-d/2} e^{-\frac{|\xi|^2 \sigma_j^2}{2n \bar{\sigma}^2}} \leq e^{-\frac{|\xi|^2}{2n} \left(\frac{l}{L} \right)^2}.$$

Then

$$\begin{aligned} \frac{\left| \hat{\mathbf{f}}_n(\xi) - \hat{G} \right|}{|\xi|^{2+s}} &= \frac{1}{|\xi|^{2+s}} \left[\sum_{i=1}^n \left| \hat{f} \left(\frac{\xi \sigma_i}{\sqrt{n\bar{\sigma}}} \right) - \hat{G} \left(\frac{\xi \sigma_i}{\sqrt{n\bar{\sigma}}} \right) \right| \left| \prod_{j=1}^{i-1} \hat{f} \left(\frac{\xi \sigma_j}{\sqrt{n\bar{\sigma}}} \right) \right| \left| \prod_{j=i+1}^n \hat{G} \left(\frac{\xi \sigma_j}{\sqrt{n\bar{\sigma}}} \right) \right| \right] \\ &\leq \frac{1}{|\xi|^{2+s}} \sum_{i=1}^n \left| \hat{f} \left(\frac{\xi \sigma_i}{\sqrt{n\bar{\sigma}}} \right) - \hat{G} \left(\frac{\xi \sigma_i}{\sqrt{n\bar{\sigma}}} \right) \right| e^{-\frac{|\xi|^2}{4n} (i-1) \left(\frac{l}{L} \right)^2} e^{-\frac{|\xi|^2}{2n} (n-1) \left(\frac{l}{L} \right)^2} \\ &\leq \frac{1}{|\xi|^{2+s}} \sum_{i=1}^n \left| \frac{\xi \sigma_i}{\sqrt{n\bar{\sigma}}} \right|^{2+s} \frac{\left| \hat{f} \left(\frac{\xi \sigma_i}{\sqrt{n\bar{\sigma}}} \right) - \hat{G} \left(\frac{\xi \sigma_i}{\sqrt{n\bar{\sigma}}} \right) \right|}{\left| \frac{\xi \sigma_i}{\sqrt{n\bar{\sigma}}} \right|^{2+s}} e^{-\frac{|\xi|^2}{4} \frac{n-1}{n} \left(\frac{l}{L} \right)^2} \\ &\leq \frac{1}{n^{\frac{2+s}{2}}} \left(\frac{L}{l} \right)^{2+s} n \sup_{\eta'} \frac{|f(\eta') - \hat{G}(\eta')|}{|\eta'|^{2+s}} e^{-\frac{|\xi|^2}{8} \left(\frac{l}{L} \right)^2}. \end{aligned}$$

Since f and G are both centred and have the same covariance matrix, in the same spirit of the proof of Lemma 3.9 (i), we obtain

$$\sup_{\eta'} \frac{|f(\eta') - \hat{G}(\eta')|}{|\eta'|^{2+s}} \leq \sup_{\eta'} \left(\frac{|f(\eta') - f(0)|}{|\eta'|^{2+s}} + \frac{|\hat{G}(\eta') - \hat{G}(0)|}{|\eta'|^{2+s}} \right) \leq C_1 (\rho_{2+s} + M_{2+s}(G)).$$

where $M_{2+s}(G) = \int |x|^{2+s} G(x) dx$. This leads to the final estimate on the third term (with the ‘‘correct’’ rate of decay), that is

$$\begin{aligned} T_3 &\leq \frac{C_1}{n^{s/2}} \left(\frac{L}{l} \right)^{2+s} (\rho_{2+s} + M_{2+s}(G)) \int_{\mathbb{R}^d} e^{-\frac{|\xi|^2}{8} \left(\frac{l}{L} \right)^2} d\xi \\ &:= \frac{1}{n^{s/2}} \left(\frac{L}{l} \right)^{2+s} (\rho_{2+s} + M_{2+s}(G)) C_{\frac{L}{l}, d}. \end{aligned}$$

Adding up T_1, T_2 and T_3 , the proof is completed. \square

3.4 Positivity of the solution

In order to show Harris's positivity condition (Hypothesis (HC)) we are in fact able to prove a stronger result, namely a local positivity condition at any positive time $t > 0$:

Theorem 3.10 (Lower bound). *Let $J: \mathbb{R}^d \rightarrow [0, \infty)$ satisfy (1), (2), and $J \in L^p$ for some $p \in (1, \infty]$. Then, the solution u of equation (NLFP) with a probability initial condition u_0 is uniformly positive on compact sets for all $0 < \varepsilon \leq 1$. That is: for every $0 < \varepsilon \leq 1$, $t > 0$, $R_1 > 0$ and $R_2 > 0$, there exists an ε -independent constant α , such that the solution of (NLFP) with initial data $u_0 \in L^1_k$ is such that*

$$u(t, x) \geq \alpha \int_{B_{R_2}} u_0 \quad \text{for } x \in B_{R_1}.$$

This is a nontrivial estimate since we seek the independence of α from ε ; in particular, we seek positivity that holds uniformly in the limit $\varepsilon \rightarrow 0$.

Our proof of this estimate is based on the Wild sum representation of the solution from Section 2.3 and on the version of Berry-Esseen's theorem we proved in Section 3.3. We distinguish two cases which are described by the two following Lemmas:

Lemma 3.11. *Let J satisfy (1) and (2). Then, for any $t, \eta > 0$, there exists $\varepsilon_0 > 0$ and a (ε -independent) constant $A > 0$ such that*

$$J_\varepsilon^{t_1, \dots, t_n}(x) \geq A \quad \text{for all } x \in B_\eta$$

for all $\varepsilon < \varepsilon_0$ and for any t_1, \dots, t_n with $t \geq t_1 \geq \dots \geq t_n \geq 0$ and n such that

$$\frac{t}{\varepsilon^2} \leq n \leq 2 \frac{t}{\varepsilon^2}.$$

Proof. Define the independent random vectors $X_i \sim J_{\varepsilon e^{t_i}}$ for which

$$\int_{\mathbb{R}^d} x J_{\varepsilon e^{t_i}} dx = 0 \quad \int_{\mathbb{R}^d} x_j x_k J_{\varepsilon e^{t_i}}(x) dx = 2\varepsilon^2 e^{2t_i} \delta_{jk}.$$

This implies that $S_n = \sum_{i=1}^n X_i \sim J_\varepsilon^{t_1, \dots, t_n}$ and

$$\int_{\mathbb{R}^d} x J_\varepsilon^{t_1, \dots, t_n}(x) dx = 0 \quad \int_{\mathbb{R}^d} x_j x_k J_\varepsilon^{t_1, \dots, t_n}(x) dx = 2\varepsilon^2 n \bar{\tau}^2 \delta_{jk},$$

where $\bar{\tau}^2 := \frac{1}{n} \sum_{i=1}^n e^{2t_i}$ and $1 \leq \bar{\tau}^2 \leq e^{2t}$. Notice that

$$l := \varepsilon \leq \varepsilon e^{t_n} \leq \dots \leq \varepsilon e^{t_1} \leq \varepsilon e^t := L$$

and thus $\frac{L}{l} = e^t$. Let $N > 0$ be the threshold for which the Berry-Esseen estimate 3.6 holds, and define $\varepsilon_1 = \sqrt{\frac{t}{N}}$. Let $\varepsilon_2 > 0$ be such that

$$\left[G\left(\frac{\eta}{\sqrt{t}}\right) - \frac{C_{BE}}{\sqrt{t}} \varepsilon_2^s \right] \geq \frac{1}{2} G\left(\frac{\eta}{\sqrt{t}}\right).$$

Take $\varepsilon_0 := \min\{\varepsilon_1, \varepsilon_2\}$. For all $\varepsilon < \varepsilon_0$, we can apply Theorem 3.6, since, by hypothesis, $n \geq \frac{t}{\varepsilon^2} \geq \frac{t}{\varepsilon_1^2} = N$. Hence, since G is radially decreasing,

$$\begin{aligned} J_\varepsilon^{t_1, \dots, t_n}(x) &\geq (2\varepsilon^2 n \bar{\tau}^2)^{-d/2} \left[G\left(\frac{x}{\sqrt{n\varepsilon\bar{\tau}}}\right) - \frac{C_{BE}}{n^{s/2}} \right] \\ &\geq (2\varepsilon^2 n \bar{\tau}^2)^{-d/2} \left[G\left(\frac{\eta}{\sqrt{t\bar{\tau}}}\right) - \frac{C_{BE}}{\sqrt{t}} \varepsilon^s \right] \\ &\geq (2\varepsilon^2 n \bar{\tau}^2)^{-d/2} \left[G\left(\frac{\eta}{\sqrt{t\bar{\tau}}}\right) - \frac{C_{BE}}{\sqrt{t}} \varepsilon_2^s \right] \\ &\geq (4te^{2t})^{-d/2} \frac{1}{2} G\left(\frac{\eta}{\sqrt{t}}\right) := A. \quad \square \end{aligned}$$

Notice that the BE constant depends on $\frac{L}{t} = e^t$ which is ε -independent.

If ε has a positive lower bound ε_0 , then we prove the following Lemma in the same spirit:

Lemma 3.12. *Let J satisfies (1) and (2). Take $\varepsilon_0 \in (0, 1]$, $t > 0$ and $\eta > 0$. Then, there exist an explicit $N \in \mathbb{N}$, $B > 0$ such that for all $n \geq N$ and $\varepsilon \in [\varepsilon_0, 1]$,*

$$J_\varepsilon^{t_1, \dots, t_n}(x) \geq \frac{B}{n^{\frac{d}{2}}} =: B_n \quad \text{for all } x \in B_n$$

where B depends on ε_0 , t , d , η and the Berry-Esseen constant.

Proof. Let $N = \max\{N_1, N_2\}$, where N_1 is the threshold for which estimate in Theorem 3.6 holds, and N_2 is such that e.g.

$$G\left(\frac{\eta}{\varepsilon_0}\right) - \frac{C_{BE}}{N_2^{s/2}} \geq \frac{1}{2} G\left(\frac{\eta}{\varepsilon_0}\right).$$

Then for $n \geq N$, applying Berry - Esseen CLT, we obtain

$$\begin{aligned} J_\varepsilon^{t_1, \dots, t_n}(x) &\geq (2\varepsilon^2 n \bar{\tau}^2)^{-d/2} \left[G\left(\frac{x}{\sqrt{n\varepsilon\bar{\tau}}}\right) - \frac{C_{BE}}{n^{s/2}} \right] \\ &\geq \frac{1}{n^{\frac{d}{2}}} \left(\frac{1}{\sqrt{2e^t}}\right)^d \left[G\left(\frac{\eta}{\varepsilon_0}\right) - \frac{C_{BE}}{N_2^{s/2}} \right] \\ &\geq \frac{1}{n^{\frac{d}{2}}} \left(\frac{1}{\sqrt{2e^t}}\right)^d \frac{1}{2} G\left(\frac{\eta}{\varepsilon_0}\right) =: \frac{B}{n^{\frac{d}{2}}}. \quad \square \end{aligned}$$

The last ingredient we need to prove Theorem 3.10 is the following lemma on the partial sums of the densities of Poisson-distributed random variables.

Lemma 3.13. *There exists an explicit constant C_L such that*

$$s_m := e^{-m} \sum_{n=m}^{2m} \frac{m^n}{n!} \geq C_L \quad \text{for all } m \geq 1,$$

and $\lim_{m \rightarrow \infty} s_m = \frac{1}{2}$.

Proof. Let X_m be a Poisson random variable of parameter m . For a fixed m , using a known result on bounds for the tail probabilities of a Poisson random variable, we have that

$$s_m = 1 - \mathbb{P}(X_m < m) - \mathbb{P}(X_m \leq 2m + 1) \geq \frac{1}{2} - e^{m+1} \left(\frac{m}{2m+1} \right)^{2m+1} := b_m \geq b_1 =: C_L,$$

since b_m is increasing for $m \geq 1$. To prove that $s_m \rightarrow 1/2$, let us now take a random variable X_m , Poisson-distributed with parameter m . Let $Y_i, i = 1, \dots, m$ be Poisson-distributed random variables with parameter 1 so that $X_m = \sum_{i=1}^m Y_i$. We have

$$\begin{aligned} e^{-m} \sum_{n=m}^m \frac{m^n}{n!} &= e^{-m} \sum_{n=0}^{2m} \frac{m^n}{n!} - e^{-m} \sum_{n=0}^{m-1} \frac{m^n}{n!} \\ &= \mathbb{P}\left(\sum_{i=1}^m Y_i - m \leq m\right) - \mathbb{P}\left(\frac{1}{\sqrt{m}}\left(\sum_{i=1}^m Y_i - m\right) \leq 0\right) \rightarrow 1 - \frac{1}{2} = \frac{1}{2}, \end{aligned}$$

thanks to the law of large numbers and the central limit theorem. \square

Remark 3.14. For our proofs we will use the above lemma as stated. However, with a very similar proof one can show that for any $r > 1$, the lemma still holds with a partial sum from m to $\lceil rm \rceil$ (instead of m to $2m$), with the same limit of $1/2$.

Proving Theorem 3.10 is now straightforward:

Proof of Theorem 3.10. Expressing the solution via Wild sums we have.

$$u(t, x) \geq e^{(d-\frac{1}{\varepsilon^2})t} \sum_{n=1}^{\infty} \varepsilon^{-2n} \int_0^t \dots \int_0^{t_{n-1}} J_{\varepsilon}^{t_1, \dots, t_n} * u_0(e^t x) dt_n \dots dt_1$$

Set $\eta = e^t R_1 + R_2$ and take ε_0 from Lemma 3.11. If $\varepsilon \leq \tilde{\varepsilon}_0 := \min\{\varepsilon_0, \sqrt{t}\}$, applying Lemma 3.11 we have

$$\begin{aligned} u(t, x) &\geq e^{(d-\frac{1}{\varepsilon^2})t} \sum_{n=1}^{\infty} \varepsilon^{-2n} \int_0^t \dots \int_0^{t_{n-1}} \int_{\mathbb{R}^d} J_{\varepsilon}^{t_1, \dots, t_n}(e^t x - y) u_0(y) dy dt_n \dots dt_1 \\ &\geq e^{(d-\frac{1}{\varepsilon^2})t} \sum_{n=\lfloor \frac{t}{\varepsilon^2} \rfloor}^{2\lfloor \frac{t}{\varepsilon^2} \rfloor} \varepsilon^{-2n} \int_0^t \dots \int_0^{t_{n-1}} \int_{B_{R_2}} J_{\varepsilon}^{t_1, \dots, t_n}(e^t x - y) u_0(y) dy dt_n \dots dt_1 \\ &\geq A e^{dt} e^{-\frac{t}{\varepsilon^2}} \sum_{n=\lfloor \frac{t}{\varepsilon^2} \rfloor}^{2\lfloor \frac{t}{\varepsilon^2} \rfloor} \left(\frac{t}{\varepsilon^2}\right)^n \frac{1}{n!} \int_{B_{R_2}} u_0(y) dy \\ &\geq \left(A e^{dt} e^{-(\frac{t}{\varepsilon^2} - \lfloor \frac{t}{\varepsilon^2} \rfloor)} \int_{B_{R_2}} u_0(y) dy\right) e^{-\lfloor \frac{t}{\varepsilon^2} \rfloor} \sum_{n=\lfloor \frac{t}{\varepsilon^2} \rfloor}^{2\lfloor \frac{t}{\varepsilon^2} \rfloor} \left[\frac{t}{\varepsilon^2}\right]^n \frac{1}{n!} \\ &\geq \left(A e^{dt-1} \int_{B_{R_2}} u_0(y) dy\right) e^{-m} \sum_{n=m}^{2n} \frac{m^n}{n!} \geq AC_L e^{dt-1} \int_{B_{R_2}} u_0(y) dy \end{aligned}$$

with $m = \lfloor \frac{t}{\varepsilon^2} \rfloor \geq 1$ and C_L the constant found in Lemma 3.13. Notice that we have used $\tilde{\varepsilon}_0 \leq \varepsilon_0$ in order to apply Lemma 3.11, and also $\tilde{\varepsilon}_0 \leq \sqrt{t}$ to make sure that the bound we obtain is not trivial (that is, to avoid $m = 0$).

If $\varepsilon \geq \tilde{\varepsilon}_0$ then, applying Lemma 3.12 to $\tilde{\varepsilon}_0$, for all $x \in B_\eta$ we obtain

$$\begin{aligned} u(t, x) &\geq e^{(d-\frac{1}{\varepsilon^2})t} \sum_{n=1}^{\infty} \varepsilon^{-2n} \int_0^t \dots \int_0^{t_{n-1}} \int_{\mathbb{R}^d} J_\varepsilon^{t_1, \dots, t_n}(e^t x - y) u_0(y) \, dy \, dt_n \dots \, dt_1 \\ &= e^{(d-\frac{1}{\varepsilon^2})t} \sum_{n=1}^{\infty} \varepsilon^{-2n} \int_0^t \dots \int_0^{t_{n-1}} \int_{B_{R_2}} J_\varepsilon^{t_1, \dots, t_n}(e^t x - y) u_0(y) \, dy \, dt_n \dots \, dt_1 \\ &\geq e^{(d-\frac{1}{\varepsilon^2})t} \sum_{n=N}^{\infty} \left(\frac{t}{\varepsilon^2}\right)^n \frac{1}{n!} B_n \int_{B_{R_2}} u_0(y) \, dy \geq B e^{(d-\frac{1}{\varepsilon_0^2})t} \left(\frac{t}{\varepsilon_0^2}\right)^N \frac{1}{N^{\frac{d}{2}}} \int_{B_{R_2}} u_0(y) \, dy. \end{aligned}$$

We finally take

$$\alpha := \min\{AC_L e^{dt-1}, B e^{(d-\frac{1}{\varepsilon_0^2})t} \left(\frac{t}{\varepsilon_0^2}\right)^N \frac{1}{N^{\frac{d}{2}}}\}.$$

□

Remark 3.15. The previous proof of positivity works almost without changes for the equation

$$\partial_t u = \frac{1}{\varepsilon^2} (J_\varepsilon * u - u) + \operatorname{div}(\nabla V u)$$

for any, even *time-dependent* potential $V : \mathbb{R}^d \times \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\frac{d}{dt} X(t) = \nabla V(X(t), t)$$

is a linear ODE. It seems harder to extend the proof to potentials with different expressions.

Proof of the main result

We have now all the ingredients to prove Theorem 1.1.

Proof. The proof comes straightforwardly thanks to Harris's Theorem 3.1. From Theorem 3.10, at every time $t > 0$ and for every $R_1, R_2 > 0$, the operator S_t satisfies Hypothesis (HC)

$$S_t \delta_{x_0} \geq \alpha \mathbf{1}_{B_{R_1}} \int_{\mathcal{S}} u_0(y) \, dy$$

on the "small set" $\mathcal{S} = B_{R_2}$.

Theorem 3.2 gives the confining condition on the generator L_ε in $L^1_{(k)}$ and consequently in L^1_k . Let u_0 be in $D(L_\varepsilon)$. By standard semigroup theory $u(t) \in \mathcal{C}^1([0, \infty); L^1_k)$, and consequently we compute with $\varphi = \langle x \rangle^k$,

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}^d} u(t, \cdot) \varphi(x) \, dx &= \int_{\mathbb{R}^d} \frac{d}{dt} u(t, \cdot) \varphi(x) \, dx \\ &= \int_{\mathbb{R}^d} (L_\varepsilon u) \varphi(x) \, dx = \int_{\mathbb{R}^d} u (L_\varepsilon^* \varphi) \, dx \leq \int_{\mathbb{R}^d} u (-\lambda \varphi + C) \, dx \end{aligned}$$

where to prove $\stackrel{*}{=}$ we use a classical cutoff argument with a bump function. We then use Gronwall's lemma with the C^1 function $f(t) := \int_{\mathbb{R}^d} u(t)\varphi$: That is,

$$f'(t) \leq -\lambda f(t) + C \implies f(t) \leq e^{-\lambda t} f(0) + \frac{C}{\lambda}(1 - e^{-\lambda t})$$

Since $D(L)$ is dense in L_k^1 , the claim is true also for any $u_0 \in L_k^1$ by standard density arguments, hence proving the Lyapunov condition on semigroups. Similarly, part (ii) of Theorem 1.1 is an immediate consequence of Theorem 3.10 and Theorem 3.4, proving a semigroup Lyapunov condition in a similar manner with $\varphi = e^{a\langle x \rangle}$. \square

4 Convergence: from nonlocal to local

As a consequence of Theorem 1.1, proved in Section 3, will now prove results on convergence of the nonlocal Fokker-Planck equation (NLFP) to the local one (FP). We present proofs which give a speed of convergence of order ε^2 , assuming the ‘‘symmetry’’ condition (3) on the kernel J . As we mentioned in Remark 1.3, one may make minor modifications in the proofs below in order to avoid condition (3), at the price of obtaining a slower speed of convergence of order ε . Since the proofs are very similar, for the sake of brevity we present only the ε^2 case. We will thus assume (3) for the whole section.

We follow a strategy of proof which is common in numerical analysis. We begin by proving suitable consistency and stability results in order to prove convergence for finite times, assuming nice regularity and decay hypotheses.

Proposition 4.1 (Consistency). *Take $k \geq 0$. Assume J satisfies (1), (3), and suppose additionally $J \in L_{k+d+4}^1$. Let $v \in C^4 \cap L_k^1$ be a function such that there exist $c > 0$ and $K > k + d$ with $|D^\alpha v(x)| \leq c \langle x \rangle^{-K}$ for all multiindices α with $|\alpha| = 4$.*

Then, there exists a constant $C = C(K, d, k, \|J\|_{L_{4+k+d}^1}, c) > 0$ such that

$$\|(L_\varepsilon - L_0)v\|_{L_k^1} \leq C\varepsilon^2$$

Proof. To lighten the notation we denote by $\varphi = \langle \cdot \rangle^{-K}$. We write (understanding that all operators are evaluated at $x \in \mathbb{R}^d$)

$$L_\varepsilon v - L_0 v = \frac{1}{\varepsilon^2} [J_\varepsilon * v - v] - \Delta v = \int_{\mathbb{R}^d} J(z) \left[\frac{1}{\varepsilon^2} (v(x - \varepsilon z) - v(x)) - \Delta_x v \right] dz,$$

and we Taylor-expand v around x up to the fourth order:

$$v(x - \varepsilon z) = \sum_{|\alpha| \leq 3} D^\alpha v(x) \frac{(-\varepsilon z)^\alpha}{\alpha!} + \sum_{|\alpha|=4} D^\alpha v(\xi_{x,\varepsilon z}) \frac{(-\varepsilon z)^\alpha}{\alpha!}$$

with $\xi_{x,\varepsilon z} \in [x, x - \varepsilon z]$. From assumptions (1) and (3) when integrating in z , for α with odd entry α_i , all terms in the first sum disappear. This leads to

$$\begin{aligned} |L_\varepsilon v - L_0 v| &= \left| \int_{\mathbb{R}^d} J(z) \left[\frac{1}{\varepsilon^2} (v(x - \varepsilon z) - v(x)) - \Delta_x v \right] dz \right| \\ &\leq \varepsilon^2 \int_{\mathbb{R}^d} J(z) \sum_{|\alpha|=4} \frac{1}{\alpha!} |D^\alpha v|(\xi_{x,\varepsilon z}) |z|^4 dz \leq c\varepsilon^2 \int_{\mathbb{R}^d} J(z) \varphi(\xi_{x,\varepsilon z}) |z|^4 dz. \end{aligned}$$

Thus, we are left with

$$\begin{aligned} \|(L_\varepsilon - L_0)v\|_{L_k^1} &\leq c\varepsilon^2 \int_{\mathbb{R}^d} (1 + |x|^k) \int_{\mathbb{R}^d} J(z) \varphi(\xi_{x,\varepsilon z}) |z|^4 dz dx \\ &= c\varepsilon^2 \int_{\mathbb{R}^d} J(z) |z|^4 \int_{\mathbb{R}^d} \varphi(\xi_{x,\varepsilon z}) (1 + |x|^k) dx dz. \end{aligned}$$

We split the inner integral:

$$\int_{\mathbb{R}^d} \varphi(\xi_{x,\varepsilon z}) (1 + |x|^k) dx = \int_{|x| \leq 2\varepsilon|z|} \varphi(\xi_{x,\varepsilon z}) (1 + |x|^k) dx + \int_{|x| > 2\varepsilon|z|} \varphi(\xi_{x,\varepsilon z}) (1 + |x|^k) dx.$$

The first term can be bounded by

$$\int_{|x| \leq 2\varepsilon|z|} \varphi(\xi_{x,\varepsilon z}) (1 + |x|^k) dx \leq \varphi(0) (1 + |2\varepsilon|z|^k) |B_{2\varepsilon|z}| \leq C_1 \varepsilon^d |z|^d + C_2 \varepsilon^{k+d} |z|^{k+d}$$

Since in the second integral $\varepsilon|z| < \frac{|x|}{2}$ and φ is radial, we can bound

$$\begin{aligned} \int_{|x| > 2\varepsilon|z|} \varphi(\xi_{x,\varepsilon z}) (1 + |x|^k) dx &\leq \int_{|x| > 2\varepsilon|z|} \varphi(|x| - \varepsilon|z|) (1 + |x|^k) dx \\ &\leq \int_{|x| > 2\varepsilon|z|} \varphi(x/2) (1 + |x|^k) dx \leq 2^{k+d} \|\varphi\|_{L_k^1} \end{aligned}$$

Then

$$\begin{aligned} \|(L_\varepsilon - L_0)v\|_{L_k^1} &\leq c\varepsilon^2 \int_{\mathbb{R}^d} J(z) |z|^4 [2^{k+d} \|\varphi\|_{L_k^1} + C_2 \varepsilon^d |z|^d + C_3 \varepsilon^{k+d} |z|^{k+d}] dz \\ &\leq c\varepsilon^2 \left[C_1 \int_{\mathbb{R}^d} J(z) |z|^4 dz + C_2 \varepsilon^d \int_{\mathbb{R}^d} J(z) |z|^{4+d} dz + C_3 \varepsilon^{k+d} \int_{\mathbb{R}^d} J(z) |z|^{4+k+d} dz \right] \\ &\leq c\varepsilon^2 \left(C_1 \|J\|_{L_4^1} + C_2 \varepsilon^d \|J\|_{L_{4+d}^1} + C_3 \varepsilon^{k+d} \|J\|_{L_{4+k+d}^1} \right) \leq C\varepsilon^2 \end{aligned}$$

where C only depends on $c, k, K, d,$, and $\|J\|_{L_{4+k+d}^1}$. □

Remark 4.2. Notice that for any v such that $D^\alpha v = 0$ with $|\alpha| \leq 4$, like polynomials of order 3 or lower, the local and the nonlocal operators agree.

We now study the stability of our nonlocal PDE under small perturbations:

Proposition 4.3 (Stability). *Let $u \in \mathcal{C}([0, \infty); L_k^1)$ be the solution of*

$$\partial_t u = L_\varepsilon u$$

Let $h, v \in \mathcal{C}([0, \infty); L_k^1)$ satisfy in the mild sense

$$\partial_t v = L_\varepsilon v + h(t)$$

with the same initial data $u(0) = v(0) = u_0$. Then, there exists constants $\omega > 0, M \geq 1$ depending only on k and d such that

$$\|u(t) - v(t)\|_{L_k^1} \leq M \int_0^t e^{\omega(t-s)} \|h(s)\|_{L_k^1} ds \quad \text{for all } t \geq 0.$$

Proof. $\psi := v - u$ solves

$$\begin{cases} \partial_t \psi = L_\varepsilon \psi + h(t) \\ \psi(0) = 0 \end{cases}$$

Using Duhamel's formula we have

$$\|\psi(t)\|_{L_k^1} = \left\| \int_0^t e^{(t-s)L_\varepsilon} h(s) ds \right\|_{L_k^1} \leq M \int_0^t e^{\omega(t-s)} \|h(s)\|_{L_k^1} ds,$$

since for a semigroup there exist $\omega > 0, M \geq 1$, such that

$$\|e^{tL_\varepsilon} f\|_{L_k^1} \leq M e^{\omega t} \|f\|_{L_k^1}. \quad \square$$

Remark 4.4. One could improve the growth rate in the above result by a more careful estimate of $e^{(t-s)L_\varepsilon} h(s)$. However, since the point of this is to later obtain a local-in-time bound, we give only the above rough estimate.

We can therefore prove the following:

Proposition 4.5 (Local-in-time convergence). *Let J satisfy assumptions (1), (3), and assume $J \in L_{k+d+4}^1$ $k \geq 0$. Let u be the solution of the nonlocal Fokker-Planck equation (NLFP) and v the solution of the local Fokker-Planck equation (FP) with the same initial data $u_0 \in L_k^1 \cap \mathcal{C}^4$. Suppose that there exists $c > 0$ and $K > k + d$ with $|D^\alpha u_0(x)| \leq c \langle x \rangle^{-K}$ for all multiindices α with $|\alpha| = 4$. Then for all $T^* > 0$ there exist $C, \omega > 0$ such that,*

$$\|u(t) - v(t)\|_{L_k^1} \leq C e^{\omega T^*} \varepsilon^2 \quad \text{for } t \in [0, T^*].$$

Proof. By standard theory of Fokker-Planck equation, the solution $v = v(t, x)$ of (FP) is in \mathcal{C}^4 , for every $t \in [0, T^*]$ and $|D^\alpha v(t, x)| \leq c \langle x \rangle^{-K}$ uniformly in $[0, T^*]$, due to our assumptions on the derivatives of u_0 . Hence, v satisfies the hypotheses of Proposition 4.1, uniformly for all $t \in [0, T^*]$. We rewrite

$$Lv = L_\varepsilon v + (L_0 v - L_\varepsilon v) := L_\varepsilon v + h(t).$$

Then

$$\|u(t) - v(t)\|_{L_k^1} \leq M e^{\omega t} \int_0^t e^{-\omega s} \|L_0 v - L_\varepsilon v\|_{L_k^1} ds \leq M \frac{e^{\omega t} - 1}{\omega} C_1 \varepsilon^2 \leq C e^{\omega T^*} \varepsilon^2$$

where we used our results on stability of Proposition 4.3 in the first inequality and the consistency with constant C_1 of Proposition 4.1 in the second inequality. \square

Once we have studied the convergence for finite times, we use Theorem 1.1 to gather information on the equilibrium and the behaviour of the solution at large times. As a first consequence of the previous results, we can estimate the speed convergence of the equilibrium F_ε of (NLFP) to the equilibrium G of (FP) as $\varepsilon \rightarrow 0$. Indeed, having a spectral gap for an operator (semigroup) provides information of the norm of the resolvent of the latter, assuring that the norm of the resolvent remains bounded by a constant.

Proposition 4.6 (Convergence of the equilibria). *Let $k > 0$ and suppose J satisfies (1) and (3). Additionally assume $J \in L^1_{k+d+4}$. Then*

$$\|F_\varepsilon - G\|_{L^1_k} \leq C\varepsilon^2$$

Proof. Feller-Miyadera-Philips's general case of Hille-Yosida Theorem for semigroups—see (Engel and Nagel, 2001, Part 2, Thm. 3.8)—states that having an estimate for a strongly continuous semigroup (S_t) in a Banach space \mathcal{Y} of the type

$$\|S_t\| \leq Me^{-\gamma t} \quad \text{for } \gamma > 0, M > 1, t > 0$$

(where $\|\cdot\|$ denotes the operator norm) is equivalent to the property that for every $\lambda > -\gamma$, λ is in the resolvent set $\rho(L_\varepsilon)$ of its infinitesimal generator $(L_\varepsilon, D(L_\varepsilon))$, and

$$\|R(\lambda, L_\varepsilon)^n\| \leq \frac{M}{(\lambda + \gamma)^n} \quad \text{for every } n \in \mathbb{N}, \quad (21)$$

where R denotes the resolvent operator. Choosing $n = 1$ and $\lambda = 0 > -\gamma$ in (21), from Theorem 1.1 it follows that L_ε has an inverse with $\|L_\varepsilon^{-1}\| \leq M/\gamma$. Then,

$$\begin{aligned} \|F_\varepsilon - G\|_{L^1_k} &= \|L_\varepsilon^{-1}L_\varepsilon(F_\varepsilon - G)\| \leq \|L_\varepsilon^{-1}\| \cdot \|L_\varepsilon F_\varepsilon - L_\varepsilon G\|_{L^1_k} \\ &\leq \frac{M}{\gamma} \|L_\varepsilon G\|_{L^1_k} = \frac{M}{\gamma} \|(L_\varepsilon - L_0)G\|_{L^1_k}. \end{aligned}$$

Given the regularity and decay properties of the Gaussian, by consistency 4.1 we obtain

$$\|(L_\varepsilon - L_0)G\|_{L^1_k} \leq C\varepsilon^2,$$

concluding the proof □

As a consequence of the previous proposition and, again, Theorem 1.1 we can estimate the speed of convergence at large times.

Proposition 4.7 (Convergence for large times). *Let J satisfy the hypotheses of Theorem 1.1 (i), Assumption (3) and, additionally $J \in L^1_{k+d+4}$. Let u be the solution of (NLFP) and v be the solution of (FP) with the same initial data $u_0 \in L^1_k$. Then, there exist $T^* > 0$ and $C > 0$ (depending on d, k, p , and u_0) such that*

$$\|u(t, \cdot) - v(t, \cdot)\|_{L^1_k} \leq C\varepsilon^2 \quad \text{for all } t > T^*.$$

Proof. From Theorem 1.1

$$\|u(t, \cdot) - F_\varepsilon\|_{L_k^1} \leq C_1 e^{-\lambda_1 t} \|u_0 - F_\varepsilon\|_{L_k^1}.$$

Moreover, we have a similar result for the local case— see e.g. [Gualdani et al. \(2018\)](#).

$$\|v(t, \cdot) - G\|_{L_k^1} \leq C_2 e^{-\lambda_2 t} \|u_0 - G\|_{L_k^1}$$

where v is the solution of the local Fokker Planck equation with the same initial data. This means

$$\|u(t, \cdot) - v(t, \cdot)\|_{L_k^1} \leq \|F_\varepsilon - G\|_{L_k^1} + C e^{-\lambda t} \left(\|u_0 - F_\varepsilon\|_{L_k^1} - \|u_0 - G\|_{L_k^1} \right).$$

From Theorem 4.6,

$$\|F_\varepsilon - G\|_{L_k^1} \leq \tilde{C} \varepsilon^2.$$

Let us define T^* such that

$$e^{-\lambda T^*} \left(\|u_0 - F_\varepsilon\|_{L_k^1} - \|u_0 - G\|_{L_k^1} \right) \leq \varepsilon^2.$$

Then for all $t > T^*$

$$\|u(t, \cdot) - v(t, \cdot)\|_{L_k^1} \leq C \varepsilon^2. \quad \square$$

We finally combine the previous results to give the proof of Theorem 1.2:

Proof of Theorem 1.2. The proof comes directly from the previous estimates in this section. Indeed, take T^* as in Proposition 4.7. If $t \in [0, T^*]$, then apply Proposition 4.5. If $t > T^*$, then Proposition 4.7 gives the result. \square

5 Analysis of the equilibrium: regularity, moments, cumulants and tails

Thanks to Harris's Theorem we have proved the existence and uniqueness of an exponentially stable equilibrium for the equation (NLFP) at each F_ε . We can infer some information about its regularity and shape, especially on the cumulants, moments and tails.

5.1 Regularity

Thanks to the the explicit form (10) of the equilibrium in the frequency domain, one can study the regularity by standard results on the Fourier transform.

Theorem 5.1. *The equilibrium F_ε of equation (NLFP) has the following regularity properties. For all $\varepsilon \in (0, 1]$*

(i) $F_\varepsilon \in \mathcal{C}^{k-d}$ for all $k < \frac{1}{\varepsilon^2}$ such that $k \geq d$.

(ii) $F_\varepsilon \notin \mathcal{C}^k$ for all $k > \frac{1}{\varepsilon^2}$.

Proof. Recall that the Fourier transform of the equilibrium is

$$\hat{F}_\varepsilon(\xi) = \exp\left(\frac{1}{\varepsilon^2} \int_0^\infty \zeta_\varepsilon(e^{-s}\xi) ds\right)$$

with $\zeta_\varepsilon(\xi) = \hat{J}_\varepsilon(\xi) - 1$. Let us assume ζ_ε radially symmetric. If not, define $\underline{\zeta}_\varepsilon$ and $\bar{\zeta}_\varepsilon$ by

$$\underline{\zeta}_\varepsilon(z) = \min_{|y|=|z|} \zeta_\varepsilon(y) \quad \bar{\zeta}_\varepsilon(z) = \max_{|y|=|z|} \zeta_\varepsilon(y).$$

These functions are then radially symmetric and satisfy

$$\begin{aligned} \underline{\zeta}_\varepsilon(\xi) &\leq \zeta_\varepsilon(\xi) \leq \bar{\zeta}_\varepsilon(\xi), \\ \underline{\zeta}_\varepsilon(\xi) &= -|\xi|^2 + o(|\xi|^3) \quad \text{and} \quad \bar{\zeta}_\varepsilon(\xi) = -|\xi|^2 + o(|\xi|^3) \quad \text{as } \xi \rightarrow 0. \end{aligned}$$

After a change of variable $y = e^{-s}|\xi|$, we obtain

$$\int_0^\infty \zeta_\varepsilon(e^{-s}|\xi|) ds = \int_0^{|\xi|} \frac{\zeta_\varepsilon(y)}{y} dy$$

Since $\zeta_\varepsilon(y) = -\varepsilon^2 y^2 + o(y^2)$, the singularity at zero is cancelled and, moreover, close to $\xi = 0$ we have

$$\hat{F}_\varepsilon(\xi) \approx e^{-\frac{|\xi|^2}{2}}.$$

To study the regularity, we are interested in the decay properties of the Fourier transform

(i) To prove the first claim, it suffices to show that

$$|\hat{F}_\varepsilon(\xi)| \leq \frac{c}{1 + |\xi|^{k+\eta}} \quad \text{for } k < \frac{1}{\varepsilon^2} \quad (22)$$

for some (small) positive η . Indeed, by the Riemann-Lebesgue lemma, $\hat{J}(y) \rightarrow 0$ as $|y| \rightarrow \infty$. Let $y_\delta > 0$ be such that

$$|\hat{J}(y)| \leq \delta := \frac{\varepsilon^2}{2} \left(\frac{1}{\varepsilon^2} - k \right) \quad \text{for all } |y| > y_\delta$$

Without loss of generality, we can assume $y_\delta < |\xi|$, otherwise there is nothing to prove. Then

$$\int_0^{|\xi|} \frac{\zeta_\varepsilon(y)}{y} dy = \int_0^{y_\delta} \frac{\zeta_\varepsilon(y)}{y} dy + \int_{y_\delta}^{|\xi|} \frac{\zeta_\varepsilon(y)}{y} dy \leq C_1 - (1 - \delta)(\log |\xi| - \log |y_\delta|)$$

which implies that

$$|\hat{F}_\varepsilon(\xi)| \leq C \exp\left(-\frac{1-\delta}{\varepsilon^2} \log |\xi|\right) = \frac{C}{|\xi|^{\frac{1-\delta}{\varepsilon^2}}} \leq \frac{C}{|\xi|^{k+(\frac{1}{\varepsilon^2}-k)/2}}.$$

For $|\xi|$ large, the previous implies condition (22) with $\eta < (\frac{1}{\varepsilon^2} - k)/2$ and by standard argument on Fourier transform $F_\varepsilon \in \mathcal{C}^{k-d}$ for all $k < \frac{1}{\varepsilon^2}$.

(ii) To prove the second claim, we will show that

$$|\xi|^k |\hat{F}_\varepsilon(\xi)| \notin L^\infty \quad \text{for } k > \frac{1}{\varepsilon^2}$$

proceeding in a similar way as before. Notice that since $\xi \mapsto \hat{J}(\xi)$ is continuous and decays to zero as $|\xi| \rightarrow +\infty$, there exists y_δ such that we can bound \hat{J} from below by

$$\hat{J}(x) \geq -\delta := \frac{\varepsilon^2}{2} \left(\frac{1}{\varepsilon^2} - k \right) \quad \text{for all } y \geq y_\delta.$$

As a consequence we see that

$$|\xi|^k \hat{F}_\varepsilon(\xi) \geq |\xi|^k C \exp \left(-\frac{1+\delta}{\varepsilon^2} \log |x| \right) = |\xi|^{k-\frac{1+\delta}{\varepsilon^2}} \notin L^\infty$$

and thus by standard properties of the Fourier transform, $F_\varepsilon \notin C^k$.

□

Remark 5.2. We just pointed out that the regularity keeps improving when $\varepsilon \rightarrow 0$ although the result is not optimal. It would be interesting to study the regularity of the equilibrium which is, as far as we know, an open question.

Hence we have proved that the regularity of the equilibrium keeps improving as $\varepsilon \rightarrow 0$. In fact, we suspect that under quite general conditions, the only point at which F_ε is not C^∞ is the origin. We are able to show this in dimension 1, and for radially symmetric J :

Theorem 5.3. *Let $\Omega \subset\subset \mathbb{R}^d$ such that $0 \notin \Omega$. Suppose (at least) one of the two following condition is verified*

(a) $d = 1$

(b) J is radially symmetric.

then $F_\varepsilon \in C^\infty(\mathbb{R}^d \setminus \{0\})$.

Proof. We proceed via a bootstrap argument: the equilibrium satisfies the stationary equation

$$\operatorname{div}(xF_\varepsilon) = -\frac{1}{\varepsilon^2}(J_\varepsilon * F_\varepsilon - F_\varepsilon) \quad (23)$$

Since $F_\varepsilon \in L^1(\Omega)$, the RHS is also in $L^1(\Omega)$, and thus $\operatorname{div}(xF_\varepsilon)$ must be in $L^1(\Omega)$ as well.

(a) If $d = 1$ then $\operatorname{div}(xF_\varepsilon) \in L^1(\Omega)$ and $F_\varepsilon \in L^1(\Omega)$ implies that $xF' \in L^1(\Omega)$ and, being $\operatorname{dist}(0, \Omega) \geq \delta$, also F' is integrable. This, by the fundamental theorem of calculus, leads to continuity for F and consequently for the RHS of (23). Iterating the process one gets the result.

(b) If $d \neq 1$, the fact that $\operatorname{div}(xF_\varepsilon)$ is integrable in Ω does not assure that $\nabla F \in L^1(\Omega)$, even when we are far from 0. Therefore, we need an additional hypothesis. If J is radially symmetric, F_ε inherits this property (see Remark 2.8), and for such functions the following holds:

$$F_\varepsilon \in W^{j,1}(\Omega), \quad \operatorname{div}(xF_\varepsilon) \in W^{j,1}(\Omega) \implies F_\varepsilon \in W^{j+1,1}(\Omega) \quad \text{for all } j \geq 0$$

Indeed, since $\operatorname{div}(xF_\varepsilon) = dF_\varepsilon + x \cdot \nabla F_\varepsilon$ and, by hypotheses $x \cdot F_\varepsilon \in W^{1,1}(\Omega)$, due to radial symmetry we can write

$$x \cdot \nabla F_\varepsilon = x \cdot F'_\varepsilon(|x|) \frac{x}{|x|} = F'_\varepsilon(|x|)|x|.$$

Since $0 \notin \Omega$, (i.e. $|x| \geq \delta$), it follows that $F'(|x|) \in W^{1,1}(\Omega)$. By definition of ∇F_ε and again the fact that $0 \notin \Omega$, we conclude that $\nabla F_\varepsilon \in W^{j,1}(\Omega)$, which implies that $F_\varepsilon \in W^{j+1,1}(\Omega)$.

We then iterate the process as in step (a) using (23), improving the regularity up to $F_\varepsilon \in W^{d,1}(\Omega)$, where d is the dimension. By the Sobolev embedding theorem, this implies that $F_\varepsilon \in \mathcal{C}(\Omega)$.

If we keep repeating the argument $(k+1)d$ times, we have that $F_\varepsilon \in \mathcal{C}^k(\Omega)$ for every k and thus the \mathcal{C}^∞ regularity follows.

□

5.2 Moments and cumulants

Let $\alpha = (\alpha_1, \dots, \alpha_d)$ be a multi-index with order $|\alpha| = n$. For a nonnegative Borel measure f on \mathbb{R}^d We define the *absolute moment of order n* by

$$M_n(f) := \int_{\mathbb{R}^d} |x|^n f(x) dx,$$

(always well-defined but possibly $+\infty$) and the α -moment by

$$m_\alpha(f) := \int_{\mathbb{R}^d} x^\alpha f(x) dx = \int_{\mathbb{R}^d} x_1^{\alpha_1} \cdots x_d^{\alpha_d} f(x) dx,$$

which is well-defined whenever $M_n[f] < +\infty$. The *moment generating function* $\mathfrak{M}(\xi)$ is given by

$$\mathfrak{M}[f](\xi) := \int_{\mathbb{R}^d} e^{x\xi} u(t, x) dx$$

whenever $x \mapsto e^{x\xi} u(t, x)$ is integrable. Moments can then also defined via the moment generating function by

$$m_\alpha(f) = D^\alpha \mathfrak{M}[f](\xi) \Big|_{\xi=0}.$$

Cumulants are an alternative way to describe a distribution, similar to the moments. They can be defined through the *cumulative generating function* (c.g.f.) $C[f]$, which is the logarithm of the moment generating function $\mathfrak{M}(\xi)$:

$$C[f](\xi) := \log \int_{\mathbb{R}^d} e^{x\xi} f(x) dx.$$

The α -cross cumulant is then defined by

$$\kappa_\alpha(f) = D^\alpha C[f](\xi) \Big|_{\xi=0}.$$

If two distributions have the same moments than they have the same cumulants and vice versa. However, moments always depend on the moments of lower order while the cumulants don't.

Using Fáa di Bruno's formula, a generalization of the chain rule for repetitive derivatives, one can prove that cumulants and moments are linked via Bell's polynomials, defined in [Comtet \(1974\)](#). These are well known combinatorial objects in \mathbb{R} , defined through their exponential generating function, that is

$$\exp \left(\sum_{i=1}^{\infty} \frac{x_i}{i!} y^i \right) = \sum_{n=0}^{\infty} \frac{B_n(x_1, \dots, x_n)}{n!} y^n. \quad (24)$$

In the rest of this section, u denotes a solution to [\(NLFP\)](#), and $C = C(t, \xi)$ is its cumulative generating function:

$$C(t, \xi) = \log \int_{\mathbb{R}^d} e^{x\xi} u(t, x) dx.$$

This can be explicitly computed in a simple way:

Theorem 5.4. *Let J satisfy [\(1\)](#) and let u be the solution of [\(NLFP\)](#) with initial data u_0 . Assume that $\int_{\mathbb{R}^d} e^{|x|} u_0 dx < \infty$ and $\int_{\mathbb{R}^d} e^{|x|} J dx < \infty$. Then*

$$C(t, \xi) = C_0(e^{-t}\xi) + \frac{1}{\varepsilon^2} \int_0^t \zeta(e^{-s}\xi) ds$$

with $C_0(\xi) = \log \int_{\mathbb{R}^d} e^{x\xi} u_0(x) dx$ and $\zeta(\xi) = \mathfrak{M}(J) - 1 = \int_{\mathbb{R}^d} e^{x\xi} J(x) dx - 1$.

Proof. We proceed similarly to the work of [Alfaro and Veruete \(2020\)](#) and [Gil et al. \(2017\)](#). The partial derivatives of the c.g.f. are

$$\partial_{\xi_i} C(t, \xi) = \frac{\int_{\mathbb{R}^d} x_i e^{x\xi} dx}{\int_{\mathbb{R}^d} e^{x\xi} dx}$$

and

$$\partial_t C(t, \xi) = \frac{\int_{\mathbb{R}^d} e^{x\xi} \partial_t u dx}{\int_{\mathbb{R}^d} e^{x\xi} u dx} = \frac{1}{\varepsilon^2} \frac{\int_{\mathbb{R}^d} e^{x\xi} (J_\varepsilon * u - u) dx}{\int_{\mathbb{R}^d} e^{x\xi} u dx} + \frac{\int_{\mathbb{R}^d} e^{x\xi} \operatorname{div}(xu) dx}{\int_{\mathbb{R}^d} e^{x\xi} u dx}$$

Analysing the first term we get

$$\begin{aligned} \int_{\mathbb{R}^d} e^{x\xi} (J_\varepsilon * u) \, dx &= \int_{\mathbb{R}^d} e^{x\xi} \left(\int_{\mathbb{R}^d} J_\varepsilon(x-y) u(t, y) \, dy \right) \, dx \\ &= \int_{\mathbb{R}^d} e^{y\xi} u(t, y) \, dy \int_{\mathbb{R}^d} J_\varepsilon(x-y) e^{(x-y)\xi} \, dx = \left(\int_{\mathbb{R}^d} e^{y\xi} u(t, y) \, dy \right) \mathfrak{M}(J_\varepsilon). \end{aligned}$$

For the second term, integrating by parts,

$$\int_{\mathbb{R}^d} e^{x\xi} \operatorname{div}_x(xu) \, dx = - \int_{\mathbb{R}^d} \nabla_x(e^{x\xi}) \cdot (xu) = -\xi \cdot \int_{\mathbb{R}^d} e^{x\xi} xu = -\xi \cdot \nabla_\xi C(t, \xi) \int_{\mathbb{R}^d} e^{x\xi} u \, dx$$

Therefore

$$\partial_t C(t, \xi) = \frac{1}{\varepsilon^2} \zeta_\varepsilon(\xi) - \xi \cdot \nabla_\xi C(t, \xi).$$

One can check that the only solution to this above first order PDE is the function C in the statement (of course, the PDE can be solved, for example by the method of characteristics, to obtain the given formula). Hence the proof is complete. \square

Corollary 5.5. *In the hypotheses of the previous theorem, let α be a multi-index with $|\alpha| = n$. Then,*

(i) *The α -cross cumulant is*

$$\kappa_\alpha(u(t)) = e^{-|\alpha|t} \kappa_\alpha(u_0) + (1 - e^{-|\alpha|t}) \frac{1}{|\alpha|\varepsilon^2} m_\alpha(J_\varepsilon)$$

As a consequence

$$\kappa_\alpha(F_\varepsilon) = \frac{1}{|\alpha|\varepsilon^2} m_\alpha(J_\varepsilon)$$

(ii) *If $d = 1$, then, the n -th moment is*

$$m_n(u) = B_n(\kappa_1, \dots, \kappa_n) = \sum_{k=1}^n B_{n,k}(\kappa_1, \dots, \kappa_{n-k+1}) \quad (25)$$

where B_n are the complete (exponential) Bell's Polynomials defined in (24) and $B_{n,k}$ are called partial Bell's Polynomials¹.

Proof. From Theorem 5.4 the α -th derivative of the c.g.f. is

$$D^\alpha C(t, \xi) = e^{-nt} (D^\alpha C_0)(e^{-t\xi}) + \frac{1 - e^{-nt}}{n\varepsilon^2} (D^\alpha \beta)(e^{-t\xi})$$

and thus, evaluating at $\xi = 0$ we get the α -mixed cumulant

$$\kappa_\alpha(u(t)) = e^{-nt} \kappa_\alpha(u_0) + (1 - e^{-nt}) \frac{1}{n\varepsilon^2} m_\alpha(J_\varepsilon)$$

Expression (25) comes directly from the definition of Bell's polynomials (24). \square

¹From a combinatorial point of view $B_{n,k}(x_1, \dots, x_{n-k+1})$ enumerate the possible ways to partition a set of n elements into k non-empty subsets. Each term of the polynomials express specific way to distribute the n elements in the k subsets. For example, $B_{5,2} = 10x_2x_3 + 5x_1x_4$ means that there are 10 ways to partition a set of 5 elements into two subsets of cardinality respectively 2 and 3 and 5 ways of doing so in two subsets of cardinality 1 and 4.

Remark 5.6. One can generalise the last result to higher dimension but it appears that the multidimensional version of Bell’s polynomials is not very well known. We refer to [Withers and Nadarajah \(2010\)](#), [Schumann \(2019\)](#) and references therein for further information.

Remark 5.7. With a direct computation one can obtain a recursive formula for the moments of the solution (and consequently for the equilibrium)

$$m_\alpha(u(t)) = \gamma_\alpha(1 - e^{-nt}) + e^{-nt}m_\alpha(u_0)$$

where

$$\gamma_\alpha = \frac{1}{n\varepsilon^2} \sum_{\substack{\beta \leq \alpha \\ \beta \neq 0}}^n \binom{\alpha}{\beta} m_\beta(J) m_{\alpha-\beta}(u).$$

This can be obtained by looking at the time derivative of the α moment, given by:

$$\frac{d}{dt} m_\alpha(u(t)) = \frac{1}{\varepsilon^2} (m_\alpha(J_\varepsilon * u) - m_\alpha(u)) - m_\alpha(\operatorname{div}(xu)).$$

5.3 Tails

For the usual (local) Fokker-Planck equation ([FP](#)), the the is a Gaussian. In the nonlocal approximation we have proposed the tails of the equilibrium seem to decay slower, no matter how fast the kernel J is decaying. Indeed, as a consequence of the Lyapunov condition ([Theorem 3.4](#)), for an exponentially decaying kernel $J \in L_{\exp, a'}$, the unique probability equilibrium has exponentially decaying tails; that is, for every $a < a'\varepsilon^{-1}$

$$\int F_\varepsilon(x) e^{a(x)} dx < \infty.$$

We conjecture that for a kernel J decaying fast enough, the tails of the equilibrium have a “Poisson-like” behaviour: they decay roughly like $e^{-|x|\log(|x|+1)}$. This implies that

$$\int F_\varepsilon(x) e^{a(x)^\gamma} dx = \infty \quad \text{for every } \gamma > 1.$$

We have not been able to prove the previous claim in general, but we can give a good approximation when J has compact support:

Proposition 5.8. *Let $J : \mathbb{R}^d \rightarrow \mathbb{R}$ be compactly supported and its support is in the ball B_R for some $R > 0$. Then*

$$\int F_\varepsilon(x) e^{a|x|\log(|x|+1)} dx < \infty$$

for any $a \leq \frac{1}{\varepsilon R}$.

Proof. Let $\varphi(x) = \exp\{a|x|\log(1+|x|)\} \in C^2$ with a to be determined later. It follows

$$\begin{aligned} J_\varepsilon * \varphi(x) &= \int_{\mathbb{R}^d} J_\varepsilon(y)\varphi(x-y) \, dy = \int_{B_R} J(z)\varphi(x-\varepsilon z) \, dz \\ &\leq \varphi(|x| + \varepsilon R) \int_{B_R} J(z) \, dz = \varphi(|x| + R) \\ &= e^{a(|x|+\varepsilon R)\log(|x|+\varepsilon R+1)} = e^{a|x|\log(|x|+\varepsilon R+1)} e^{a\varepsilon R\log(|x|+\varepsilon R+1)} \\ &= (|x| + \varepsilon R)^{\varepsilon a R} e^{a|x|\log(|x|+\varepsilon R+1)} \end{aligned}$$

with a slight abuse of notation, having denoted with φ also the function $\varphi : \mathbb{R} \rightarrow \mathbb{R}$. Moreover,

$$x \cdot \nabla \varphi = x \cdot a \left(\frac{x}{|x|} \log(|x|+1) + \frac{x}{|x|+1} \right) \varphi = a|x|\varphi \left(\log(|x|+1) + \frac{|x|}{|x|+1} \right)$$

Hence,

$$\begin{aligned} L_\varepsilon^* \varphi &= J_\varepsilon * \varphi - \varphi - x \cdot \nabla \varphi \leq \varphi(|x| + \varepsilon R) - \varphi(x) - x \cdot \nabla \varphi(x) \\ &= (|x| + \varepsilon R)^{a\varepsilon R} e^{a|x|\log(|x|+\varepsilon R+1)} - a\varphi(x)|x| \left[\log(|x|+1) + \frac{|x|}{|x|+1} \right] \\ &\leq \varphi(x) \left[C_2(|x| + \varepsilon R)^{a\varepsilon R} - a|x|(\log(|x|+1)) \right] \leq C - \lambda\varphi \end{aligned}$$

for some $C, \lambda > 0$ and $a \leq 1/\varepsilon R$. □

Acknowledgments

We would like to thank Jo Evans for discussions which gave rise to the early ideas of this paper. We would also like to thank Alejandro Gárriz for several discussions and suggestions.

The project that gave rise to these results received the support of a fellowship from the “la Caixa” Foundation (ID 100010434). The fellowship code is LCF/BQ/DI22/11940032. Both authors acknowledge support from the “Maria de Maeztu” Excellence Unit IMAG, reference CEX2020-001105-M, funded by MCIN/AEI/10.13039/501100011033/. They were also supported by grant PID2020-117846GB-I00 and the research network RED2022-134784-T from MCIN/AEI/10.13039/501100011033/.

References

Alfaro, Matthieu and Veruete, Mario. [Density dependent replicator-mutator models in directed evolution](#). *Discrete and Continuous Dynamical Systems - B*, 25(6):2203–2221, May 2020. ISSN 1531-3492. Publisher: Discrete and Continuous Dynamical Systems - B.

- Andreu-Vaillo, Fuensanta; Mazón, José M.; Rossi, Julio D., and Toledo-Melero, Julián J. *Nonlocal diffusion problems*. American Mathematical Society ; Real Sociedad Matemática Española, 2010. ISBN 9780821852309.
- Auricchio, Ferdinando; Toscani, Giuseppe, and Zanella, Mattia. [Trends to equilibrium for a nonlocal Fokker–Planck equation](#). *Applied Mathematics Letters*, 145:108746, 2023. ISSN 0893-9659.
- Ayi, Nathalie; Herda, Maxime; Hivert, Hélène, and Tristani, Isabelle. [On a structure-preserving numerical method for fractional Fokker-Planck equations](#), July 2022. arXiv:2107.13416 [cs, math].
- Bentkus, V. [A Lyapunov-type Bound in \$\mathbf{R}^d\$](#) . *Theory of Probability & Its Applications*, 49(2):311–323, 2005, <https://doi.org/10.1137/S0040585X97981123>.
- Berry, Andrew C. The Accuracy of the Gaussian Approximation to the Sum of Independent Variates. *Transactions of the american mathematical society*, 1941.
- Bessemoulin-Chatard, Marianne; Herda, Maxime, and Rey, Thomas. Hypocoercivity and diffusion limit of a finite volume scheme for linear kinetic equations. *Mathematics of Computation*, 89(323):1093–1133, 2020.
- Bisi, Marzia; Cañizo, José A., and Lods, Bertrand. Entropy dissipation estimates for the linear Boltzmann operator. *Journal of Functional Analysis*, 269(4):1028–1069, 2015.
- Cai, Long; Friedman, Nir, and Xie, X Sunney. Stochastic protein expression in individual cells at the single molecule level. *Nature*, 440(7082):358–362, 2006.
- Cancès, Clément; Chainais-Hillairet, Claire; Herda, Maxime, and Krell, Stella. Large time behavior of nonlinear finite volume schemes for convection-diffusion equations. *SIAM Journal on Numerical Analysis*, 58(5):2544–2571, 2020.
- Cañizo, José A.; Carrillo, José A., and Pájaro, Manuel. [Exponential equilibration of genetic circuits using entropy methods](#). *Journal of Mathematical Biology*, 78(1-2): 373–411, aug 2019.
- Cañizo, José A; Cao, Chuqi; Evans, Josephine, and Yoldaş, Havva. Hypocoercivity of linear kinetic equations via harris’s theorem. *Kinetic and Related Models*, 13(1): 97–128, 2020.
- Cañizo, José A. and Mischler, Stéphane. [Harris-type results on geometric and subgeometric convergence to equilibrium for stochastic semigroups](#), 2021, [arXiv:2110.09650v1](https://arxiv.org/abs/2110.09650v1).

- Carlen, E.A.; Carvalho, M.C.; Le Roux, J.; Loss, M., and Villani, C. [Entropy and chaos in the Kac model](#). *Kinetic and Related Models*, 3(1):85–122, 2010. ISSN 1937-5077.
- Carrillo, J. A. and Toscani, G. Contractive probability metrics and asymptotic behavior of dissipative kinetic equations. *Riv. Mat. Univ. Parma (7)*, 6:75–198, 2007.
- Chasseigne, Emmanuel; Chaves, Manuela, and Rossi, Julio. [Asymptotic behavior for nonlocal diffusion equations](#). *Journal de Mathématiques Pures et Appliquées*, 86: 271–291, 09 2006.
- Comtet, Louis. *Advanced combinatorics: Theory of finite and infinite expansions*. Reidel, Dordrecht, rev. and enlarged ed edition, 1974. ISBN 978-90-277-0441-2.
- Dolbeault, Jean; Mouhot, Clément, and Schmeiser, Christian. [Hypocoercivity for kinetic equations with linear relaxation terms](#). *Comptes Rendus Mathématique*, 347(9): 511–516, 2009. ISSN 1631-073X.
- Dujardin, Guillaume; Hérau, Frédéric, and Lafitte, Pauline. Coercivity, hypocoercivity, exponential time decay and simulations for discrete fokker–planck equations. *Numerische Mathematik*, 144(3):615–697, 2020.
- Engel, Klaus-Jochen and Nagel, Rainer. [One-parameter semigroups for linear evolution equations](#). *Semigroup Forum*, 63:278–280, 06 2001.
- Esseen, Carl-Gustav. On the liapunov limit error in the theory of probability. *Ark. Mat. Astr. Fys.*, 28:1–19, 1942.
- Gil, Marie-Eve; Hamel, Francois; Martin, Guillaume, and Roques, Lionel. [Mathematical Properties of a Class of Integro-differential Models from Population Genetics](#). *SIAM Journal on Applied Mathematics*, 77(4):1536–1561, January 2017. ISSN 0036-1399. Publisher: Society for Industrial and Applied Mathematics.
- Goudon, Thierry; Junca, Stéphane, and Toscani, Giuseppe. [Fourier-Based Distances and Berry-Esseen Like Inequalities for Smooth Densities](#). *Monatshefte für Mathematik*, 135(2):115–136, March 2002. ISSN 1436-5081.
- Gualdani, Maria P.; Mischler, Stéphane, and Mouhot, Clément. Factorization for non-symmetric operators and exponential H-theorem. *Mémoires de la Société Mathématique de France*, 2018, [arXiv:1006.5523](#). To appear.
- Hairer, Martin. [Convergence of Markov processes](#). Unpublished lecture notes, 2010.
- Hairer, Martin and Mattingly, Jonathan C. [Yet another look at Harris’ ergodic theorem for Markov chains](#). In Dalang, Robert; Dozzi, Marco, and Russo, Francesco, editors, *Seminar on Stochastic Analysis, Random Fields and Applications VI*, pages 109–117, Basel, October 2011. Springer Basel. ISBN 978-3-0348-0020-4, [arXiv:0810.2777](#).

- Hauray, Maxime and Mischler, Stéphane. [On Kac's chaos and related problems](#). *Journal of Functional Analysis*, 266(10):6055–6157, May 2014. ISSN 0022-1236.
- Ignat, Liviu I and Rossi, Julio D. A nonlocal convection–diffusion equation. *Journal of Functional Analysis*, 251(2):399–437, 2007.
- Lions, Pierre Louis and Toscani, Giuseppe. A strengthened central limit theorem for smooth densities. *Journal of Functional Analysis*, 129(1):148–167, 1995.
- Lods, Bertrand; Mouhot, Clément, and Toscani, Giuseppe. Relaxation rate, diffusion approximation and fick's law for inelastic scattering boltzmann models. *Kinetic and Related Models*, 1(2):223–248, 2008.
- Meyn, Sean P. and Tweedie, Richard L. [Markov chains and stochastic stability](#). Cambridge University Press, 2010. ISBN 9780521731829.
- Miklavcic, Milan. *Applied Functional Analysis And Partial Differential Equations*. World Scientific, August 1998. ISBN 978-981-4495-90-5.
- Mischler, Stéphane and Tristani, Isabelle. [Uniform semigroup spectral analysis of the discrete, fractional and classical Fokker-Planck equations](#). In *Journal de l'École polytechnique — Mathématiques*, volume 4, pages 389–433, March 2017. ISSN: 2270-518X.
- Molino, Alexis and Rossi, Julio D. Nonlocal approximations to Fokker-Planck equations. *Funkcialaj Ekvacioj*, 62(1):35–60, 2019.
- Pazy, A. [Semigroups of Linear Operators and Applications to Partial Differential Equations](#), volume 44 of *Applied Mathematical Sciences*. Springer, New York, NY, 1983. ISBN 978-1-4612-5563-5 978-1-4612-5561-1.
- Raič, Martin. [A Multivariate Berry-Esseen Theorem with explicit constants](#). *Bernoulli*, 25(4), 2019. ISSN 1350-7265, [1802.06475 \[math\]](#).
- Rey, Thomas and Toscani, Giuseppe. [Large-Time behavior of the solutions to Rosenau-Type approximations to the heat equation](#). *SIAM Journal on Applied Mathematics*, 73(4):1416–1438, January 2013. ISSN 0036-1399.
- Risken, Hannes. [The Fokker-Planck equation](#). Springer Berlin Heidelberg, 1996. ISBN 9783642615443.
- Schumann, Aidan. Multivariate bell polynomials and derivatives of composed functions. *arXiv preprint arXiv:1903.03899*, 2019.
- Villani, C. [Hypoocoercivity](#). *Memoirs of the American Mathematical Society*, 202:0–0, 2009.

Withers, C. S. and Nadarajah, S. [Multivariate bell polynomials](#). *International Journal of Computer Mathematics*, 87:2607–2611, 2010.

JOSÉ A. CAÑIZO. DEPARTAMENTO DE MATEMÁTICA APLICADA & IMAG, UNIVERSIDAD DE GRANADA. AVDA. FUENTENUEVA S/N, 18071 GRANADA, SPAIN. *E-mail address*: canizo@ugr.es

NICCOLÒ TASSI. DEPARTAMENTO DE MATEMÁTICA APLICADA & IMAG, UNIVERSIDAD DE GRANADA. AVDA. FUENTENUEVA S/N, 18071 GRANADA, SPAIN. *E-mail address*: tassi@ugr.es