

GENERIC NONEXPANSIVE HILBERT SPACE MAPPINGS

DAVIDE RAVASINI ¹ AND DAYLEN K. THIMM ²

ABSTRACT. We consider a closed convex set C in a separable, infinite-dimensional Hilbert space and endow the set $\mathcal{N}(C)$ of nonexpansive self-mappings on C with the topology of pointwise convergence. We introduce the notion of a somewhat bounded set and establish a strong connection between this property and the existence of fixed points for the generic $f \in \mathcal{N}(C)$, in the sense of Baire categories. Namely, if C is somewhat bounded, the generic nonexpansive mapping on C admits a fixed point, whereas if C is not somewhat bounded, the generic nonexpansive mapping on C does not have any fixed points. This results in a topological 0–1 law: the set of all $f \in \mathcal{N}(C)$ with a fixed point is either meager or residual. We further prove that, generically, there are no fixed points in the interior of C and, under additional geometric assumptions, we show the uniqueness of such fixed points for the generic $f \in \mathcal{N}(C)$ and the convergence of the iterates of f to its fixed point.

1. INTRODUCTION

The question whether nonexpansive mappings have fixed points when defined on closed, convex subsets of Hilbert spaces and more general spaces has been investigated from various viewpoints. Let us recall that on a metric space (X, ρ) a mapping $f: X \rightarrow X$ is nonexpansive if for all $x, y \in X$ the mapping satisfies

$$\rho(f(x), f(y)) \leq \rho(x, y).$$

A fundamental result is the classical theorem of Brouwer, which asserts that every continuous self-mapping of a bounded, closed, and convex subset of a Euclidean space has a fixed point. Notably, this theorem only works for the finite-dimensional case. Generalizing it to infinite dimensions necessitates to only consider nonexpansive mappings and additional geometric properties, as demonstrated in the fixed point theorem of Browder–Göhde–Kirk (see [7, 10, 11]). Finding a geometric restriction for a similar theorem to hold for all Lipschitz mappings is impossible: in every Banach space there is a Lipschitz retraction from the ball to the sphere [5], and the combination of this retraction with a suitable (also Lipschitz) reflection constitutes a Lipschitz mapping without a fixed point. Moreover, in some Banach spaces, e.g. $\mathcal{C}([0, 1])$, there are nonexpansive mappings on closed, convex, and bounded subsets without a fixed point.

This brings up the following question:

In infinite dimensions, how big is the set of nonexpansive mappings with a fixed point?

A pioneering result in this direction was found by de Blasi and Myjak [8, 9], who showed that on a bounded, closed, and convex subset of a Banach space, the generic

July 8, 2024

2020 Mathematics Subject Classification: 46C05, 54E52.

Keywords: nonexpansive mapping, fixed point, Hilbert space, somewhat bounded set.

nonexpansive mapping has a fixed point. Here generic is used to mean large in a topological sense. As the nonexpansive mappings with a fixed point form a large subset, one may wonder whether the strict contractions, i.e., Lipschitz mappings with Lipschitz constant $L < 1$, are responsible for this, as they all possess a fixed point. It turns out that this is not the case, as de Blasi and Myjak had already shown in [8] within the context of a Hilbert space that the set of strict contractions is negligible. Later it was shown that this result carries to Banach spaces in [1] and a large class of metric spaces in [2]. The type of mappings to be held responsible for the original result by de Blasi and Myjak have been found to be the Rakotch contractions, which all have fixed points. Recall that a Rakotch contraction is a mapping in which the distance of two images is not linearly bounded by the distance of the original points as in Lipschitz contractions, but by a factor that may depend on the distance. More precisely, a function f on a metric space (X, ρ) is a Rakotch contraction if there exists a decreasing function $\phi_f: [0, \infty) \rightarrow [0, 1]$ with $\phi_f(t) < 1$ for all $t > 0$ such that for all $x, y \in X$ we have that

$$\rho(f(x), f(y)) \leq \phi_f(\rho(x, y))\rho(x, y).$$

Due to the Rakotch fixed point theorem [13], all Rakotch contractions on a complete metric space have a fixed point. Furthermore, as shown by Reich and Zaslavski in [15, 16], almost all nonexpansive mappings on a bounded, closed, and convex set are Rakotch contractions.

Whereas the set of nonexpansive functions on *bounded* sets carries the topology of uniform convergence as a natural choice, in the *unbounded* case, one has multiple topologies that seem natural to choose from, such as the topology of uniform convergence on bounded sets and the topology of pointwise convergence. Moreover, there are multiple natural metrics generating these topologies. In contrast to the bounded case, where there are many Rakotch contractions, Strobin was able to prove that with respect to the topology of uniform convergence on bounded sets the opposite holds: the set of Rakotch contractions is negligible [17]. In fact, Strobin proved something more general. Recall that for $t \geq 0$ the modulus of continuity of f is defined by

$$\omega_f(t) := \sup\{\rho(f(x), f(y)) : x, y \in X, \rho(x, y) \leq t\}.$$

Given a concave and increasing function $\omega: [0, \infty) \rightarrow [0, \infty)$ with $\omega(0) = 0$, Strobin considered the space $C_\omega(X)$ of all continuous self-mappings f of X satisfying $\omega_f(t) \leq \omega(t)$ for all $t \geq 0$ with respect to the topology of uniform convergence on bounded sets. He showed that if X is an unbounded, closed, and convex subset of a Hilbert space, then the subset of all mappings satisfying $\omega_f(t) < \omega(t)$ for some $t > 0$ is meager. For the case $\omega(t) = t$ this coincides with the previous statement about Rakotch contractions. As an important tool in his proof, he uses the Grünbaum–Zarantonello extension theorem, which generalizes the Kirszbraun–Valentine theorem to uniformly continuous mappings. Both these theorems are limited to Hilbert spaces. Overcoming this limitation, Bargetz, Reich, and the second author were recently able to generalize Strobin’s result for nonexpansive mappings to hyperbolic spaces [3]. Shortly thereafter, the first author was able to extend this to arbitrary ω [14].

Furthermore, in [3], Bargetz, Reich, and the second author looked at the problem in the unbounded setting from the perspective of the topology of pointwise convergence. In an effort to prove the generic presence of fixed points, they showed that for the generic nonexpansive mapping the strict inequality $\rho(f(x), f(y)) < \rho(x, y)$ holds for residually many pairs of points $(x, y) \in X \times X$, thus hinting towards some kind of contractivity. However, we in fact disprove that the generic existence of fixed points is always guaranteed in the topology of pointwise convergence. This also provides an answer to the question asked at the end of [14].

We focus on separable Hilbert spaces and, for the first time, we observe that depending upon what type of unbounded set is given, the generic nonexpansive mapping may or may not have a fixed point. Indeed, we are able to precisely classify on what sets the generic nonexpansive mapping does or does not have a fixed point. Since the sets where the generic nonexpansive mapping does have a fixed point exhibit the features of both bounded and possibly unbounded sets, we name them *somewhat bounded* sets. We call all other sets totally unbounded. We are able to formulate this in two theorems that appear in the manuscript as Theorem 3.4 and Theorem 4.1, respectively.

Theorem. *Let C be a closed, convex, somewhat bounded set in a separable, infinite-dimensional Hilbert space H . Then there is a dense open set $\mathcal{R} \subset \mathcal{N}(C)$ such that every $f \in \mathcal{R}$ has a fixed point.*

Theorem. *Let C be a closed, convex set in a separable, infinite-dimensional Hilbert space H . If C is totally unbounded, then there is a residual set $\mathcal{R} \subset \mathcal{N}(C)$ such that every $f \in \mathcal{R}$ fails to have any fixed points.*

Note that these two statements do not logically imply each other, as there exist sets which are neither meager nor residual. As our main result we directly obtain Theorem 4.2, a topological 0–1 Law.

Theorem (0–1 Law). *Let C be a closed, convex set in a separable, infinite-dimensional Hilbert space H and let \mathcal{F} be the set of all $f \in \mathcal{N}(C)$ with a fixed point. Then either \mathcal{F} is meager or it contains a dense open set.*

As an additional result we show that the generic nonexpansive mapping on a somewhat bounded set has all fixed points on the boundary of the set. If the set is additionally locally uniformly rotund, we show that any sequence of iterates converges to the fixed point located on the boundary, as the set is, in particular, somewhat bounded. Since our proofs rely on the Kirszbraun–Valentine extension theorem, the fixed point theorem of Brouwer–Göhde–Kirk and the classical machinery of Hilbert spaces, our results are limited to subsets of a Hilbert space.

2. SETTING AND NOTATION

Given a metric space (X, ρ) , we denote by $\mathcal{N}(X)$ the space of nonexpansive mappings $f : X \rightarrow X$. If X is separable, we always assume $\mathcal{N}(X)$ to be endowed with the topology of pointwise convergence. This is the coarsest topology containing all open sets of the form $U_f(x, \varepsilon)$, where $f \in \mathcal{N}(X)$, $x \in X$, $\varepsilon > 0$ and

$$U_f(x, \varepsilon) = \{g \in \mathcal{N}(X) : \rho(g(x), f(x)) < \varepsilon\}.$$

If X is complete and separable, the topology of pointwise convergence can be metrized via a complete metric as follows. We consider a dense, countable set $\Theta = \{\vartheta_n\}_{n=1}^{\infty}$ and define $d_{\Theta} : \mathcal{N}(X) \times \mathcal{N}(X) \rightarrow \mathbb{R}$ by

$$d_{\Theta}(f, g) = \sum_{n=1}^{\infty} 2^{-n} \frac{\rho(f(\vartheta_n), g(\vartheta_n))}{1 + \rho(f(\vartheta_n), g(\vartheta_n))}.$$

The proof that d_{Θ} is indeed a complete metric on $\mathcal{N}(C)$ and that it induces the topology of pointwise convergence can be found in [3]. We stress that the existence of a complete metric for $\mathcal{N}(X)$ makes the use of Baire categories meaningful, as we shall recall at the end of this section.

The following two lemmas will be needed later.

Lemma 2.1. *Let (X, ρ) be a metric space. For every $f \in \mathcal{N}(X)$, $x \in X$, $\varepsilon > 0$ and every positive integer k the set*

$$V_f(x, \varepsilon, k) = \{g \in \mathcal{N}(X) : \rho(g^k(x), f^k(x)) < \varepsilon\}$$

is an open neighborhood of f in $\mathcal{N}(X)$.

Proof. We proceed by induction on k . For $k = 1$, note that $V_f(x, \varepsilon, 1) = U_f(x, \varepsilon)$ for every $f \in \mathcal{N}(X)$, $x \in X$ and $\varepsilon > 0$. Let now k be fixed and assume that $V_g(x, \delta, k)$ is open in $\mathcal{N}(X)$ for every $g \in \mathcal{N}(X)$, $x \in X$ and $\delta > 0$. Given now $f \in \mathcal{N}(X)$, $x \in X$ and $\varepsilon > 0$, we have to show that $V_f(x, \varepsilon, k + 1)$ is open. Let $g \in V_f(x, \varepsilon, k + 1)$. The goal is to find an open neighborhood W of g such that $\rho(h^{k+1}(x), f^{k+1}(x)) < \varepsilon$ for every $h \in W$. Pick $\delta > 0$ such that $2\delta + \rho(f^{k+1}(x), g^{k+1}(x)) < \varepsilon$ and set $W_1 = U_g(g^k(x), \delta)$, $W_2 = V_g(x, \delta, k)$. W_1 is open by the definition of the topology, whereas W_2 is open by the inductive assumption. Now set $W = W_1 \cap W_2$. If $h \in W$ we have

$$\begin{aligned} \rho(h^{k+1}(x), f^{k+1}(x)) &\leq \rho(h(h^k(x)), h(g^k(x))) + \rho(h(g^k(x)), g(g^k(x))) + \\ &\quad + \rho(g^{k+1}(x), f^{k+1}(x)) < \\ &< \rho(h^k(x), g^k(x)) + \delta + \rho(g^{k+1}(x), f^{k+1}(x)) < \\ &< 2\delta + \rho(f^{k+1}(x), g^{k+1}(x)) < \varepsilon, \end{aligned}$$

which is what we wanted. \square

Lemma 2.2. *Let (X, ρ) be a metric space, let f be a nonexpansive mapping on X and let $x \in X$ and $r > 0$ be such that $\rho(f(x), x) > 2r$. If $y \in X$ is a fixed point of f , then $\rho(x, y) > r$.*

Proof. We have

$$\rho(x, y) = \rho(x, f(y)) \geq \rho(x, f(x)) - \rho(f(x), f(y)) > 2r - \rho(x, y),$$

from which $\rho(x, y) > r$ follows. \square

Throughout the paper we will deal with a single real, separable Hilbert space H and its closed subspaces. We simply denote by $\langle x, y \rangle$ and $\|x\|$ the scalar product of two elements $x, y \in H$ and the norm of x respectively. B_H and S_H stand for the closed unit ball and the unit sphere of H respectively. Given a closed, convex set $C \subseteq H$, we denote by $\pi_C : H \rightarrow H$ the nearest point projection onto C . In the case of a closed subspace $F \subseteq H$, we use the notation π_F for the orthogonal projection onto F . This does not lead to any confusion, since the nearest point projection and

the orthogonal projection coincide. Finally, given a subset $C \subseteq H$ and $x \in H$ we denote by $\text{dist}(x, C)$ the distance of x to C , i.e.,

$$\text{dist}(x, C) = \inf\{\|x - y\| : y \in C\}.$$

We recall that Hilbert spaces are uniformly convex Banach spaces. A Banach space X is *uniformly convex* if for every $\varepsilon > 0$ there is $\delta > 0$ such that for every $x, y \in X$ with $\|x\| = \|y\| = 1$ and $\|x - y\| > \delta$ we have $\|x + y\| \leq 2 - \varepsilon$. In uniformly convex Banach spaces, the following theorem holds (see [7, 11, 10]).

Theorem 2.3 (Browder–Göhde–Kirk). *Let C be a closed, convex, bounded set in a uniformly convex Banach space X . Then every nonexpansive mapping $f : C \rightarrow C$ has a fixed point.*

Theorem 2.3 makes questions concerning the generic existence of fixed points rather trivial in the bounded case, which is the reason why our primary focus is on unbounded sets. Finally, we recall that in Hilbert spaces the Kirszbraun–Valentine Theorem on the extension of Lipschitz mappings is available (see, for instance, [4], Theorem 1.12). This will be needed in the proof of Theorem 3.4.

Theorem 2.4 (Kirszbraun–Valentine). *Let H_1 and H_2 be two Hilbert spaces, let $A \subseteq H_1$ be any subset and let $f_0 : A \rightarrow H_2$ be a Lipschitz mapping with Lipschitz constant L . Then there is a Lipschitz mapping $f_1 : H_1 \rightarrow H_2$ with Lipschitz constant L and such that $f_1(x) = f_0(x)$ for every $x \in A$.*

To conclude this preliminary section, we shall spend a few words on Baire categories, since they play a central role in the statement of all our results. A subset N of a topological space X is *nowhere dense* if the closure of N in X has empty interior or, equivalently, if $X \setminus N$ contains a dense open set. A subset of X is *meager* if it is a countable union of nowhere dense sets. The complement of a meager set is often called *comeager* or *residual* and, by definition, it contains a countable intersection of dense open sets. The importance of meager and residual sets lies in the Baire category theorem, which asserts that a completely metrizable space is never meager in itself, i.e., it cannot be a countable union of some of its nowhere dense subsets. Since countable unions of meager sets are also meager, the Baire category theorem implies that a countable family of meager sets will never be able to cover the whole space. This conveys the intuitive idea that meager sets are small, whereas residual sets almost fill the entire space. The points of a residual set are usually called the *generic* points of the space and a property which holds for every point in a residual set is usually said to be generic.

3. SOMEWHAT BOUNDED SETS

The central definition of our work is the following.

Definition 3.1. *A nonempty, non-singleton convex set C in an infinite-dimensional Hilbert space H is **somewhat bounded** if there are $x_0 \in C$, a finite-dimensional subspace $F \subset H$ and $\alpha > 0$ such that $\alpha B_F \subset C - x_0$ and $F^\perp \cap (C - x_0)$ is bounded. We say that C is **totally unbounded** if it is not somewhat bounded.*

For example, the set $\{x \in H : \text{dist}(x, F) \leq r\}$, where $F \subset H$ is a finite-dimensional subspace and $r \geq 0$, is unbounded but also somewhat bounded. We say that a convex set C in a Hilbert space H is *totally unbounded* if it is not somewhat bounded. H itself is totally unbounded, as well as all its infinite-dimensional subspaces.




FIGURE 1. Visualization of the proof of Lemma 3.2

Given a Hilbert space H , for every closed subspace $F \subset H$ and $\alpha, \beta > 0$ we define the set

$$D_F(\alpha, \beta) = \{x \in H : \|\pi_{F^\perp}(x)\| - \beta \leq \alpha^{-1}\beta\|\pi_F(x)\|\}.$$

Note that, by definition, $(x + F^\perp) \cap D_F(\alpha, \beta)$ is bounded for every $x \in H$.

Lemma 3.2. *Let C be a convex set in an infinite-dimensional Hilbert space H . Then the following are equivalent.*

- (1) C is somewhat bounded.
- (2) There are $z_0 \in C$, a finite-dimensional subspace $F \subset H$ and $\alpha, \beta > 0$ such that $\alpha B_F \subset C - z_0 \subset D_F(\alpha, \beta)$.

Proof. (1) \Rightarrow (2). Since C is somewhat bounded, we can find $z_0 \in C$, a finite-dimensional subspace $F \subset H$ and $\alpha > 0$ such that $\alpha B_F \subset C$ and $F^\perp \cap (C - z_0)$ is bounded. Since $F^\perp \cap (C - z_0)$ is bounded, there must be $\beta > 0$ such that $\beta S_{F^\perp} \cap (C - z_0) = \emptyset$. We want to show that $C - z_0 \subset D_F(\alpha, \beta)$. Assume that this is not the case and pick $y \in (C - z_0) \setminus D_F(\alpha, \beta)$. Set

$$\lambda = \frac{\beta}{\|\pi_{F^\perp}(y)\| - \beta}$$

and $w = -\lambda\pi_F(y)$. Note that

$$\|w\| = \lambda\|\pi_F(y)\| = \frac{\beta\|\pi_F(y)\|}{\|\pi_{F^\perp}(y)\| - \beta} < \beta\beta^{-1}\alpha = \alpha,$$




FIGURE 2. Visualization of the proof of Lemma 3.3

thus $w \in \alpha B_F \subset C - z_0$. Moreover, if we set

$$\mu = \frac{1}{1 + \lambda} = \frac{\|\pi_{F^\perp}(y)\| - \beta}{\|\pi_{F^\perp}(y)\|} \in (0, 1),$$

we obtain

$$\mu w + (1 - \mu)y = \frac{\beta \pi_{F^\perp}(y)}{\|\pi_{F^\perp}(y)\|} \in \beta S_{F^\perp} \cap (C - z_0).$$

This contradicts $\beta S_{F^\perp} \cap (C - z_0) = \emptyset$. Thus, we conclude that $C - z_0 \subset D_F(\alpha, \beta)$, as claimed.

(2) \Rightarrow (1). It is obvious, since $F^\perp \cap (C - z_0) \subseteq F^\perp \cap D_F(\alpha, \beta) = \beta B_{F^\perp}$ is a bounded set. \square

Lemma 3.3. *Assume that, for a closed, convex, unbounded set C in a Hilbert space H , there are $\alpha, \beta > 0$ and a subspace $F \subset H$ such that $\alpha B_F \subset C \subset D_F(\alpha, \beta)$. For every $\lambda, t > 0$ define*

$$E_F(\lambda, t) = \{x \in tS_F : \|\pi_C(x)\| \geq \lambda\}.$$

Then, for every $\lambda > 0$, there are $t \geq \lambda$ and $r, r' > 0$ such that $r' > r$ and

$$r' B_H \cap C \subseteq E(\lambda, t) + r B_H.$$

Proof. Let $\lambda > 0$ be given. We first want to show that there is $t \geq \lambda$ such that $E(\lambda, t)$ is nonempty. To this end, select t with

$$t - \frac{\beta}{\sqrt{\alpha^2 + \beta^2}}(\alpha + t) \geq \lambda.$$

Since C is unbounded and $C \subset D_F(\alpha, \beta)$, there must $p \in C$ with $\|\pi_F(p)\| \geq t$. Define

$$v = \frac{t\pi_F(p)}{\|\pi_F(p)\|} \in tS_F, \quad w = -\frac{\alpha\pi_F(p)}{\|\pi_F(p)\|} \in \alpha S_F.$$

The point on the line segment $\{w + \mu(p - w) : \mu \in [0, 1]\}$ which minimizes the distance to v is given by setting

$$\mu = \frac{\langle p - w, v - w \rangle}{\|p - w\|^2}.$$

If we call q such point, we have

$$\|q - v\|^2 = (\alpha + t)^2 - \frac{\langle p - w, v - w \rangle^2}{\|p - w\|^2}.$$

We now want to take a closer look at the right-hand side of the above equation. Using that $p \in D_F(\alpha, \beta)$ we get

$$\frac{\langle p - w, v - w \rangle}{\|p - w\|} = \|v - w\| \left\langle \frac{p - w}{\|p - w\|}, \frac{v - w}{\|v - w\|} \right\rangle \geq (\alpha + t) \frac{\alpha}{\sqrt{\alpha^2 + \beta^2}}.$$

This implies

$$\|q - v\|^2 \leq (\alpha + t)^2 - \frac{\alpha^2}{\alpha^2 + \beta^2} (\alpha + t)^2 = \frac{\beta^2}{\alpha^2 + \beta^2} (\alpha + t)^2.$$

Since $q \in C$, we now have

$$\|v\| - \|\pi_C(v)\| \leq \|v - \pi_C(v)\| \leq \|v - q\| \leq \frac{\beta}{\sqrt{\alpha^2 + \beta^2}} (\alpha + t),$$

therefore

$$\|\pi_C(v)\| \geq \|v\| - \frac{\beta}{\sqrt{\alpha^2 + \beta^2}} (\alpha + t) = t - \frac{\beta}{\sqrt{\alpha^2 + \beta^2}} (\alpha + t) \geq \lambda,$$

i.e., $v \in E(\lambda, t)$.

We now set

$$r = \sqrt{(2t)^2 + (\beta + \alpha^{-1}\beta t)^2}$$

and $r' = \sqrt{r^2 + t^2} > r$. Let $x \in C \cap r'B_H$. If $\|\pi_F(x)\| \leq t$, then $x \in \overline{B}(v, r)$, where $v \in E(\lambda, t)$ is the point we have determined before. We have indeed

$$\|x - v\|^2 = \|\pi_F(x) - v\|^2 + \|\pi_{F^\perp}(x)\|^2 \leq (2t)^2 + (\beta + \alpha^{-1}\beta t)^2.$$

If $\|\pi_F(x)\| > t$, then, as in the first part of the proof, we have

$$y = \frac{t\pi_F(x)}{\|\pi_F(x)\|} \in E(\lambda, t).$$

Moreover,

$$\begin{aligned} \|x - y\|^2 &= \|x\|^2 + \|y\|^2 - 2\langle x, y \rangle = \\ &= \|x\|^2 + t^2 - 2t\|\pi_F(x)\| \leq \\ &\leq r^2 + 2t^2 - 2t\|\pi_F(x)\| \leq \\ &\leq r^2 + 2t^2 - 2t^2 = \\ &= r^2. \end{aligned}$$

□

We are now ready to show the generic existence of fixed points for nonexpansive mappings on somewhat bounded sets.

Theorem 3.4. *Let C be a closed, convex, somewhat bounded set in a separable, infinite-dimensional Hilbert space H . Then there is a dense open set $\mathcal{R} \subset \mathcal{N}(C)$ such that every $f \in \mathcal{R}$ has a fixed point.*

Proof. First let us note that we may assume that C is unbounded, as every nonexpansive mapping has a fixed point on bounded sets due to Theorem 2.3. By translating C if needed, we may also assume that there are $\alpha, \beta > 0$ and a finite dimensional subspace $F \subset H$ with $\alpha B_F \subset C \subset D_F(\alpha, \beta)$. Let $\Theta = \{\vartheta_n\}_{n=1}^\infty$ be a dense set in C and let d_Θ be the corresponding metric on $\mathcal{N}(C)$. To prove the existence of \mathcal{R} , let us fix some $f \in \mathcal{N}(C)$ and show that arbitrarily close to f there is an open neighborhood of functions with a fixed point. Let us pick $\varepsilon > 0$ and find $n \in \mathbb{N}$ with $\sum_{j=n+1}^\infty 2^{-j} < \varepsilon$. Now set $M = \sup\{\|\vartheta_1\|, \dots, \|\vartheta_n\|, \|f(\vartheta_1)\|, \dots, \|f(\vartheta_n)\|\}$ and $\lambda = 2M$. By Lemma 3.3 we get $t \geq \lambda$ and $r', r > 0$ with $r' > r$ such that

$$r' B_H \cap C \subseteq E_F(\lambda, t) + r B_H,$$

where $E_F(\lambda, t)$ is defined as in the statement of Lemma 3.3. Let us choose $\delta > 0$ such that $r + \delta < r'$. Since F is finite-dimensional, $E_F(\lambda, t)$ is compact, therefore we can find a finite δ -net $N \subset E(\lambda, t)$. Note that this implies

$$C \cap r' B_H \subseteq N + (r + \delta) B_H.$$

Moreover, since the inequality $\|x - \pi_C(y)\| \leq \|x - y\|$ holds for every $x \in C$ and every $y \in H$, we also have $C \cap r' B_H \subseteq \pi_C(N) + (r + \delta) B_H$. Note further that $\|x\| \geq 2M$ for every $x \in \pi_C(N)$ by the definition of $E_F(\lambda, t)$. Let us define the function

$$g_0 : \{\vartheta_1, \dots, \vartheta_n\} \cup \pi_C(N) \rightarrow C$$

$$x \mapsto \begin{cases} f(x) & \text{if } x \in \{\vartheta_1, \dots, \vartheta_n\} \\ 0 & \text{if } x \in \pi_C(N) \end{cases}$$

and check that it is nonexpansive. Clearly, if $x, y \in \{\vartheta_1, \dots, \vartheta_n\}$ or $x, y \in \pi_C(N)$ we have that $\|g_0(x) - g_0(y)\| \leq \|x - y\|$, thus it only remains to show that the same holds for $x \in \{\vartheta_1, \dots, \vartheta_n\}$ and $y \in \pi_C(N)$. However, this is true as

$$\|x - y\| \geq \|y\| - \|x\| \geq 2M - M = M$$

and

$$\|g_0(x) - g_0(y)\| = \|f(x)\| \leq M \leq \|x - y\|.$$

Now, Theorem 2.4 yields a nonexpansive mapping $g_1 : C \rightarrow H$ which extends g_0 , i.e., such that $g_1(x) = g_0(x)$ for every $x \in \{\vartheta_1, \dots, \vartheta_n\} \cup \pi_C(N)$. Finally, let us define $g = \pi_C \circ g_1 \in \mathcal{N}(C)$ and note that g also extends g_0 . Furthermore, as g and f agree on $\{\vartheta_1, \dots, \vartheta_n\}$, we have that $d_\Theta(f, g) \leq \sum_{j=n+1}^\infty 2^{-j} < \varepsilon$. We now want to exhibit an open neighborhood U of g in $\mathcal{N}(C)$ such that every $h \in U$ has a fixed point. Choose $\eta > 0$ with $r + \delta + \eta < r'$. By the definition of the topology of pointwise convergence, the set

$$U = \{h \in \mathcal{N}(C) : \|h(x)\| < \eta \text{ for all } x \in \pi_C(N)\}$$

is an open neighborhood of g in $\mathcal{N}(C)$. For every $x \in r' B_H \cap C$ there is $y \in \pi_C(N)$ with $\|x - y\| \leq r + \delta$. If $h \in U$, we have that $\|h(y)\| < \eta$, hence

$$\|h(x)\| = \|h(x) - h(y)\| + \|h(y)\| < r + \delta + \eta < r'.$$

The above line shows that $h(r'B_H \cap C) \subseteq r'B_H \cap C$ for every $h \in U$. Since every $h \in U$ maps a closed, convex, bounded subset of C into itself, we can conclude by Theorem 2.3 that every $h \in U$ has a fixed point. Since f and ε were chosen arbitrarily, this concludes the proof. \square

Although Theorem 3.4 guarantees the generic existence of fixed points in the case of a somewhat bounded set C , we shall now see that, generically, none of them lie in the interior of C . We start with a Lemma, which we also need in Section 5.

Lemma 3.5. *Let C be a closed, convex, somewhat bounded set in a separable, infinite-dimensional Hilbert space H . Then there is a residual set $\mathcal{R} \subseteq \mathcal{N}(C)$ such that for every $f \in \mathcal{R}$ and every $x \in C$ we have $\text{dist}(f^k(x), \partial C) \rightarrow 0$ as $k \rightarrow \infty$.*

Proof. Let us assume without loss of generality that $0 \in C$ and that $F \subset H$ is a finite-dimensional subspace such that $C \cap (x + F^\perp)$ is bounded for every $x \in H$. Let $\Theta = \{\vartheta_n\}_{n=1}^\infty$ be a dense set in C and consider the corresponding metric d_Θ . Given a positive integer p and $r \in \mathbb{Q} \cap (0, 1)$ define

$$R(p, r) = \{f \in \mathcal{N}(C) : \text{there is } k \text{ with } \text{dist}(f^k(\vartheta_p), \partial C) < r\}.$$

We now fix p and r arbitrarily and show that $R(p, r)$ contains a dense open set. To this end, we pick $f \in \mathcal{N}(C)$, $\delta > 0$ and carry out the next part of the proof in two steps.

Step 1. We want to find $h \in \mathcal{N}(C)$ with $d_\Theta(h, f) < \delta$ and a positive integer k such that $h^k(\vartheta_p) \in \partial C$. Choose a positive integer n with $\sum_{j=n+1}^\infty 2^{-j} < \delta/2$ and set $m = \max\{n, p\}$. Define

$$\begin{aligned} G &= F + \text{span}\{\vartheta_1, \dots, \vartheta_m, f(\vartheta_1), \dots, f(\vartheta_m)\} \\ D &= \text{conv}\{\vartheta_1, \dots, \vartheta_m, f(\vartheta_1), \dots, f(\vartheta_m)\} \end{aligned}$$

Choose $y_0 \in S_{G^\perp}$. Since $G^\perp \subseteq F^\perp$, we have that $C \cap (x + \mathbb{R}y_0)$ is compact for every $x \in D$, hence we can define the function $\phi : D \rightarrow [0, \infty)$ by setting $\phi(x) = \max\{t \geq 0 : x + ty_0 \in C\}$ for every $x \in D$. The function ϕ is defined on a compact, convex subset of a finite-dimensional Hilbert space and it is easily seen to be concave, thus it must be continuous, which means that it attains a maximum value $M \geq 0$. Let $P : H \rightarrow F$ be the orthogonal projection onto F and let $Q : H \rightarrow \mathbb{R}y_0$ be the orthogonal projection onto $\mathbb{R}y_0$. Note that $\pi_D = \pi_D \circ P$. Define $g \in \mathcal{N}(C)$ as $g = \pi_D \circ f \circ \pi_D$ and $h : C \rightarrow C$ by setting

$$h(x) = \pi_C \left(g(x) + Q(x) + \frac{\delta}{2} y_0 \right)$$

for every $x \in C$. We have to check that h is nonexpansive. For every $x, y \in C$ we have

$$\begin{aligned} \|h(x) - h(y)\|^2 &\leq \|g(x) - g(y)\|^2 + \|Q(x) - Q(y)\|^2 = \\ &= \|(\pi_D \circ f \circ \pi_D \circ P)(x) - (\pi_D \circ f \circ \pi_D \circ P)(y)\|^2 + \\ &\quad + \|Q(x) - Q(y)\|^2 \leq \\ &\leq \|P(x) - P(y)\|^2 + \|Q(x) - Q(y)\|^2 = \|x - y\|^2, \end{aligned}$$

as we want. Next, we want $d_{\Theta}(h, f) < \delta$. Note that, for every $j \in \{1, \dots, m\}$, we have

$$\begin{aligned} \|h(\vartheta_j) - f(\vartheta_j)\| &= \left\| \pi_C \left(f(\vartheta_j) + \frac{\delta}{2} y_0 \right) - \pi_C(f(\vartheta_j)) \right\| \leq \\ &\leq \left\| f(\vartheta_j) + \frac{\delta}{2} y_0 - f(\vartheta_j) \right\| = \frac{\delta}{2}, \end{aligned}$$

hence

$$d_{\Theta}(f, h) < \sum_{j=1}^m 2^{-j} \|h(\vartheta_j) - f(\vartheta_j)\| + \frac{\delta}{2} < \frac{\delta}{2} + \frac{\delta}{2} = \delta.$$

Finally, we want a positive integer k with $h^k(\vartheta_p) \in \partial C$. Assume, by way of contradiction, that $h^k(\vartheta_p) \in \text{int}C$ for every positive integer k . Using that $\pi_C(x) \in \text{int}C$ only if $\pi_C(x) = x$, it is not hard to show by induction on k that the formula

$$h^k(\vartheta_p) = g^k(\vartheta_p) + \frac{k\delta}{2} y_0$$

holds for every $k \in \mathbb{N}$. Since $g^k(\vartheta_p) \in D$ for every k , we now have $k \leq 2\delta^{-1}M$ for every positive integer k , which is absurd.

Step 2. We want to show that there is an open neighborhood $U \subset \mathcal{N}(C)$ of h such that $\text{dist}(u^k(\vartheta_p), \partial C) < r$ for every $u \in U$, where k is the positive integer we have determined in Step 1. Using the notation of Lemma 2.1, this can be achieved by setting $U = V_h(\vartheta_p, r, k)$, which is indeed an open neighborhood of h such that every $u \in U$ has the desired property.

Step 1 and Step 2 prove that $R(p, q)$ contains a dense open set, as wished. Now define

$$\mathcal{R} = \bigcap_{r \in \mathbb{Q} \cap (0,1)} \bigcap_{p=1}^{\infty} R(p, r)$$

and observe that \mathcal{R} is residual. Let $f \in \mathcal{R}$, let $x \in C$ and let $\varepsilon > 0$. Find a positive integer n with $\|x - \vartheta_n\| < \varepsilon/2$. By our choice of f we can find a positive integer k with $\text{dist}(f^k(\vartheta_n), \partial C) < \varepsilon/2$. We now have

$$\text{dist}(f^k(x), \partial C) \leq \|f^k(x) - f^k(\vartheta_n)\| + \text{dist}(f^k(\vartheta_n), \partial C) \leq \|x - \vartheta_n\| + \frac{\varepsilon}{2} < \varepsilon.$$

Since x and ε are arbitrary, this shows that \mathcal{R} is the set we are looking for. \square

Theorem 3.6. *Let C be a closed, convex, somewhat bounded set in a separable, infinite-dimensional Hilbert space H . Then there is a residual set $\mathcal{R} \subseteq \mathcal{N}(C)$ such that, for every $f \in \mathcal{R}$, there are no fixed points of f in $\text{int}C$.*

Proof. Let \mathcal{R} be the residual set given by Lemma 3.5. Let $f \in \mathcal{R}$ and assume that x is a fixed point of f . Then

$$\text{dist}(x, \partial C) = \text{dist}(f^k(x), \partial C) \rightarrow 0$$

as $k \rightarrow \infty$. Thus, $\text{dist}(x, \partial C) = 0$. Since ∂C is closed, this is possible only if $x \in \partial C$. \square

Corollary 3.7. *Let C be a closed, strictly convex, somewhat bounded set in a separable, infinite-dimensional Hilbert space H . Then there exists a residual set $\mathcal{R} \subset \mathcal{N}(C)$ such that every $f \in \mathcal{R}$ has a unique fixed point.*

Proof. Theorem 3.4 yields a dense open set $\mathcal{R}_1 \subset \mathcal{N}(C)$ such that every $f \in \mathcal{R}_1$ has at least one fixed point, whereas Theorem 3.6 yields a residual set \mathcal{R}_2 such that, for every $f \in \mathcal{R}_2$, all fixed points of f lie in ∂C . Set $\mathcal{R} = \mathcal{R}_1 \cap \mathcal{R}_2$ and note that \mathcal{R} is residual. Note that, for every $f \in \mathcal{N}(C)$, the set of fixed points of f is convex because the norm of H is strictly convex. If $f \in \mathcal{R}$, then the set of fixed points of f is nonempty and contained in ∂C . Since ∂C does not admit any nontrivial convex subsets, we deduce that the fixed point of f must be unique. \square

4. TOTALLY UNBOUNDED SETS

In the present section we aim to show that the situation is completely opposite on totally unbounded sets, in that the generic nonexpansive mapping does not admit any fixed points. We can already formulate and prove the main theorem. The proof adapts and makes use of the same iteration technique we have already seen in Lemma 3.5.

Theorem 4.1. *Let C be a closed, convex set in a separable, infinite-dimensional Hilbert space H . If C is totally unbounded, then there is a residual set $\mathcal{R} \subset \mathcal{N}(C)$ such that every $f \in \mathcal{R}$ fails to have any fixed points.*

Proof. Without loss of generality, we can assume $0 \in C$. Let $\Theta = \{\vartheta_n\}_{n=1}^\infty$ be a dense subset in C and define the corresponding metric d_Θ . For every positive integer r , define $R(r)$ as the set of all $f \in \mathcal{N}(C)$ such that no fixed point of f lies in rB_H . We want to show that $R(r)$ contains a dense open set for every r . To this end, we consider $f \in \mathcal{N}(C)$ and $\varepsilon > 0$ and divide the next part of the proof in two steps.

Step 1. We want to find $h \in \mathcal{N}(C)$ and a positive integer k such that $d_\Theta(f, h) < \varepsilon$ and $\|h^k(0)\| \geq 3r$. Choose a positive integer n with $\sum_{j=n+1}^\infty 2^{-j} < \varepsilon/2$ and set

$$F = \text{span}\{\vartheta_1, \dots, \vartheta_n, f(\vartheta_1), \dots, f(\vartheta_n)\}.$$

Note that $C \cap F$ has nonempty relative interior in F , which implies that there are $x_0 \in C \cap F$ and $\alpha > 0$ with $\alpha B_F \subset C - x_0$. Set

$$\rho = \max\{\|\vartheta_1 - x_0\|, \dots, \|\vartheta_n - x_0\|, \|f(\vartheta_1) - x_0\|, \dots, \|f(\vartheta_n) - x_0\|\}$$

and pick any positive $\delta < \min\{1, (8\rho)^{-1}\varepsilon\}$. The assumption that C is totally unbounded implies that $C \cap (x_0 + F^\perp)$ is unbounded, hence we can find $y_0 \in S_{F^\perp}$ such that $x_0 + 3r\delta^{-1}y_0 \in C$. Define

$$D = \delta x_0 + (1 - \delta) \cdot \text{conv}\{\vartheta_1, \dots, \vartheta_n, f(\vartheta_1), \dots, f(\vartheta_n)\},$$

It follows from the convexity of C that

$$\delta x_0 + (1 - \delta)C + [0, 3r]y_0 \subseteq C.$$

Indeed, for every $x \in C$ and $t \in [0, 3r]$ one has

$$\begin{aligned} \delta x_0 + (1 - \delta)x + ty_0 &\in (1 - \delta)C + [\delta x_0, \delta x_0 + 3ry_0] = \\ &= (1 - \delta)C + \delta[x_0, x_0 + 3r\delta^{-1}y_0] \subseteq (1 - \delta)C + \delta C = C. \end{aligned}$$

In particular, one has $D + [0, 3r]y_0 \subseteq C$. Let $P : H \rightarrow F$ and $Q : H \rightarrow \mathbb{R}y_0$ be the orthogonal projections onto F and $\mathbb{R}y_0$ respectively. Pick a positive integer $k > 12r\varepsilon^{-1}$ and define $g \in \mathcal{N}(C)$ as $g = \pi_D \circ f \circ \pi_D$ and $h : C \rightarrow C$ by setting

$$h(x) = \pi_C(g(x) + Q(x) + 3rk^{-1}y_0)$$

for every $x \in C$. One checks that h is nonexpansive similarly to the proof of Lemma 3.5. To see that $d_{\Theta}(f, h) < \varepsilon$, first set

$$\rho = \max\{\|\vartheta_1 - x_0\|, \dots, \|\vartheta_n - x_0\|, \|f(\vartheta_1) - x_0\|, \dots, \|f(\vartheta_n) - x_0\|\}.$$

Then, note that, for every $j \in \{1, \dots, n\}$, one has

$$\begin{aligned} \|h(\vartheta_j) - f(\vartheta_j)\| &= \|\pi_D \circ f \circ \pi_D(\vartheta_j) + 3rk^{-1}y_0 - f(\vartheta_j)\| \leq \\ &\leq \|\pi_D \circ f \circ \pi_D(\vartheta_j) - \pi_D \circ f(\vartheta_j)\| + \\ &\quad + \|\pi_D \circ f(\vartheta_j) - f(\vartheta_j)\| + 3rk^{-1} \leq \\ &\leq \|\pi_D(\vartheta_j) - \vartheta_j\| + \|\pi_D \circ f(\vartheta_j) - f(\vartheta_j)\| + 3rk^{-1} \leq \\ &\leq \|\delta x_0 + (1 - \delta)\vartheta_j - \vartheta_j\| + \\ &\quad + \|\delta x_0 + (1 - \delta)f(\vartheta_j) - f(\vartheta_j)\| + 3rk^{-1} < \\ &< 2\delta\rho + \frac{\varepsilon}{4}. \end{aligned}$$

Thus, by our choice of δ we obtain $\|h(\vartheta_j) - f(\vartheta_j)\| < \varepsilon/2$ for every $j \in \{1, \dots, n\}$. It follows that

$$\begin{aligned} d_{\Theta}(h, f) &\leq \sum_{j=1}^n 2^{-j} \|h(\vartheta_j) - f(\vartheta_j)\| + \sum_{j=n+1}^{\infty} 2^{-j} < \\ &< \frac{\varepsilon}{2} \sum_{j=1}^{\infty} 2^{-j} + \sum_{j=n+1}^{\infty} 2^{-j} \leq \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon. \end{aligned}$$

Finally, using the inclusion $D + [0, 3r]y_0 \subseteq C$, it is not hard to check by induction that the formula $h^j(x) = g^j(x) + 3rk^{-1}jy_0$ holds for every $x \in C \cap F$ and every $j \in \{1, \dots, k\}$. In particular, we have $h^k(0) = g^k(0) + 3ry_0$. Since $g^k(0)$ and y_0 are orthogonal to each other, this implies

$$\|h^k(0)\|^2 = \|g^k(0)\|^2 + 9r^2 \geq 9r^2,$$

which is what we wanted.

Step 2. We want to find a neighborhood U of h in $\mathcal{N}(C)$ such that every $u \in U$ does not have any fixed points in rB_H . Using the notation of Lemma 2.1, set $U = V_h(0, r, k)$, where k is the positive integer we found in Step 1. If $x \in C$ is a fixed point of $u \in U$, then it is also a fixed point of u^k . Note that

$$\|u^k(0)\| \geq \|h^k(0)\| - \|u^k(0) - h^k(0)\| > 3r - r = 2r,$$

therefore it follows from Lemma 2.2 that $\|x\| > r$.

Step 1 and Step 2 show that $R(r)$ contains a dense open set for every r , as desired. Now set

$$\mathcal{R} = \bigcap_{r=1}^{\infty} R(r),$$

and note that \mathcal{R} is residual. If $f \in \mathcal{R}$ has a fixed point x , then $\|x\| > r$ for every positive integer r , which is impossible. Hence, f does not have any fixed points. \square

If we combine Theorem 3.4 and Theorem 4.1, we obtain the topological 0–1 law we mentioned in the introduction.

Theorem 4.2 (0–1 Law). *Let C be a closed, convex set in a separable, infinite-dimensional Hilbert space H and let \mathcal{F} be the set of all $f \in \mathcal{N}(C)$ with a fixed point. Then either \mathcal{F} is meager or it contains a dense open set.*

5. CONVERGENCE OF THE ITERATES

We dedicate this last section to the study of the iterates f^k , $k \in \mathbb{N}$, of the generic nonexpansive mapping $f \in \mathcal{N}(C)$, where C is a closed and convex set in a separable Hilbert space. Our convergence result will be obtained under a further geometric assumption on C , which is usually called *local uniform rotundity*. We say that C is *locally uniformly rotund*, or LUR, if $\partial C \neq \emptyset$ and for every $x \in C$ and $\varepsilon > 0$ there is $\delta > 0$ such that

$$\text{dist}\left(\frac{x+y}{2}, \partial C\right) > \varepsilon$$

whenever $y \in C$ and $\|x - y\| > \delta$. Note that LUR sets are always strictly convex and have therefore nonempty interior. Alternatively, one can define LUR sets via their *modulus of convexity*. For every $x \in C$, set $\Delta_C(x) = \sup\{\|y - x\| : y \in C\}$. Now, for every $x \in C$ and every $\varepsilon \in (0, \Delta_C(x))$, define

$$\delta_C(x, \varepsilon) = \inf\left\{\text{dist}\left(\frac{x+y}{2}, \partial C\right) : y \in C, \|y - x\| \geq \varepsilon\right\}.$$

C is locally uniformly rotund precisely when $\delta_C(x, \varepsilon) > 0$ for every $x \in C$ and every $\varepsilon \in (0, \Delta_C(x))$. LUR sets are a natural generalization of LUR norms and have been considered, for example, in [12] and in [6], where the authors investigate this property in connection with existence of extensions for quasiconvex functions. We will use the fact that the modulus of convexity $\delta_C(x, \varepsilon)$ is a continuous function of ε once x is fixed.

Lemma 5.1. *Let C be a LUR set in a Hilbert space H and let x, y, z be points in C with $y \in \partial C$. If $r, s > 0$ are such that $r \leq \|x - y\| \leq s$, then*

$$\left\langle \frac{z-y}{\|z-y\|}, \frac{x-y}{\|x-y\|} \right\rangle \geq -\frac{s}{\sqrt{s^2 + 4\delta_C(x, r)^2}}.$$

Proof. We may assume without loss of generality that $y = 0$. Define $F = \text{span}\{x, z\}$ and note that, since C is strictly convex, x and z cannot be linearly dependent, hence $\dim F = 2$. Let $w \in S_F$ be such that $\langle x, w \rangle = 0$ and, for every $t \in \mathbb{R}$, define

$$p_t = \frac{1}{2}x + tw.$$

Set $\gamma = \delta_C(x, r)$. Observe that, since $\text{dist}(x/2, \partial C) \geq \gamma$, we have that $p_t \in \text{int}C$ for every $t \in (-\gamma, \gamma)$. Thus, for every $t \in (-\gamma, \gamma)$, the half line $\{sp_t : s \leq 0\}$ does not contain any points of C besides 0. In particular, $C \cap F$ is contained in the cone

$$Q = \{\lambda x + \mu w : \lambda, \mu \in \mathbb{R}, 2\gamma\lambda \geq -|\mu|\}.$$

Let λ, μ be such that $z = \lambda x + \mu w$. If $\lambda \geq 0$ the statement is obvious. If $\lambda < 0$, we have $(\mu/\lambda)^2 \geq 4\gamma^2$. This implies

$$\begin{aligned} \left\langle \frac{z}{\|z\|}, \frac{x}{\|x\|} \right\rangle &= \frac{\lambda\|x\|}{\sqrt{\lambda^2\|x\|^2 + \mu^2}} = -\frac{\|x\|}{\sqrt{\|x\|^2 + (\mu/\lambda)^2}} \geq \\ &\geq -\frac{\|x\|}{\sqrt{\|x\|^2 + 4\gamma^2}} \geq -\frac{s}{\sqrt{s^2 + 4\gamma^2}}, \end{aligned}$$

as claimed. \square

Remark: on a geometric level, Lemma 5.1 asserts that the angle φ between the vectors $x - y$ and $z - y$ must be bounded away from π by a quantity which depends on r and s and on the modulus of convexity of C at x . More precisely, we have

$$\varphi \leq \arccos \left(-\frac{s}{\sqrt{s^2 + 4\delta_C(x, r)^2}} \right) = \pi - \arccos \left(\frac{s}{\sqrt{s^2 + 4\delta_C(x, r)^2}} \right).$$

Lemma 5.2. *Let C be a LUR set in a Hilbert space H and let $f : C \rightarrow C$ be a nonexpansive mapping such that $\text{dist}(f^k(x), \partial C) \rightarrow 0$ for every $x \in C$ as $k \rightarrow \infty$. Moreover, let $x_0 \in C$ be a fixed point of f . Then there is a continuous function $\alpha : (0, \Delta_C(x_0)) \rightarrow (0, 1)$ such that for every $r \in (0, \Delta_C(x_0))$ and for every $x \in C$ with $\|x - x_0\| = r$ there is a positive integer k with $\|f^k(x) - x_0\| \leq \alpha(r)r$.*

Proof. Let $\delta : [0, \Delta_C(x_0)) \rightarrow [0, \infty)$ be the modulus of convexity of C at the point p . We introduce the function $\beta : (0, \Delta_C(x_0)) \rightarrow (0, 1)$ given by

$$\beta(r) = \frac{5r}{\sqrt{25r^2 + 256\delta(r/8)^2}}$$

and observe that

$$\sqrt{\frac{1 + \beta(r)}{2}} < 1$$

holds for every $r \in (0, \Delta_C(x_0))$. Therefore, we can find a continuous function $\varepsilon : (0, \Delta_C(x_0)) \rightarrow (0, 1/4)$ with

$$\sqrt{\frac{1 + \beta(r)}{2}} \cdot (1 + \varepsilon(r)) < 1.$$

We claim that the function α we are looking for can be defined by

$$\alpha(r) = \max \left\{ \frac{3}{4}, \sqrt{\frac{1 + \beta(r)}{2}} \cdot (1 + \varepsilon(r)) \right\}.$$

Fix $r \in (0, \Delta_C(x_0))$, let $x \in C$ with $\|x - x_0\| = r$ and let y be the midpoint of the segment $[x_0, x]$. By our assumption on f there is a positive integer k such that $\text{dist}(f^k(y), \partial C) < r\varepsilon(r)/4$, which allows us to choose a point $z \in \partial C$ with $\|f^k(y) - z\| < r\varepsilon(r)/2$. Since f is nonexpansive we now get

$$\begin{aligned} \|x_0 - z\| &\leq \|x_0 - f^k(y)\| + \|f^k(y) - z\| \leq (1 + \varepsilon(r)) \frac{r}{2}, \\ \|f^k(x) - z\| &\leq \|f^k(x) - f^k(y)\| + \|f^k(y) - z\| \leq (1 + \varepsilon(r)) \frac{r}{2}. \end{aligned}$$

In particular, we have $\|x_0 - z\| \leq 5r/8$. If $\|f^k(y) - x_0\| < r/4$, we get

$$\|f^k(x) - x_0\| \leq \|f^k(x) - f^k(y)\| + \|f^k(y) - x_0\| < \frac{3}{4}r \leq \alpha(r)r$$

and the lemma holds. If $\|f^k(y) - x_0\| \geq r/4$, we have

$$\|x_0 - z\| \geq \|f^k(y) - x_0\| - \|f^k(y) - z\| \geq \frac{r}{4} - \frac{r\varepsilon(r)}{2} = \frac{r}{4}(1 - 2\varepsilon(r)) \geq \frac{r}{8}.$$

Now we want to apply Lemma 5.1 to the points x_0 , z and $f^k(x)$ and the bound $r/8 \leq \|x_0 - z\| \leq 5r/8$, thus obtaining

$$\left\langle \frac{f^k(x) - z}{\|f^k(x) - z\|}, \frac{x_0 - z}{\|x_0 - z\|} \right\rangle \geq -\frac{5r/8}{\sqrt{(5r/8)^2 + 4\delta(r/8)^2}} = -\beta(r).$$

Therefore,

$$\begin{aligned} \|f^k(x) - x_0\|^2 &= \|f^k(x) - z\|^2 + \|x_0 - z\|^2 \\ &\quad - 2\|f^k(x) - z\| \cdot \|x_0 - z\| \cdot \left\langle \frac{f^k(x) - z}{\|f^k(x) - z\|}, \frac{x_0 - z}{\|x_0 - z\|} \right\rangle \leq \\ &\leq (1 + \varepsilon(r))^2 \cdot \frac{r^2}{4} + (1 + \varepsilon(r))^2 \cdot \frac{r^2}{4} \\ &\quad + 2(1 + \varepsilon(r))^2 \cdot \frac{r^2}{4} \cdot \beta(r) = \\ &= \frac{1}{2}(1 + \beta(r)) \cdot (1 + \varepsilon(r))^2 r^2 \leq \alpha(r)^2 r^2, \end{aligned}$$

which again confirms our claim. \square

Theorem 5.3. *Let C be a closed, somewhat bounded set in a separable, infinite-dimensional Hilbert space. If C is a LUR set, then there exists a residual set $\mathcal{R} \subseteq \mathcal{N}(C)$ such that every $f \in \mathcal{R}$ has a unique fixed point and the sequence of iterates $(f^k(x))_{k \in \mathbb{N}}$ converges to the fixed point of f for every $x \in C$.*

Proof. Theorem 3.4 yields a dense open set $\mathcal{R}_1 \subseteq \mathcal{N}(C)$ such that every $f \in \mathcal{R}_1$ has a fixed point, whereas Lemma 3.5 gives us a residual set $\mathcal{R}_2 \subset \mathcal{N}(C)$ such that for every $f \in \mathcal{R}_2$ and every $x \in C$ we have $\text{dist}(f^k(x), \partial C) \rightarrow 0$ as $k \rightarrow \infty$. Set $\mathcal{R} = \mathcal{R}_1 \cap \mathcal{R}_2$. We have already concluded in Theorem 3.6 and the subsequent Corollary 3.7 that every $f \in \mathcal{R}$ has a unique fixed point, which in addition lies on ∂C . Let $f \in \mathcal{R}$ and let x_0 be its unique fixed point. Since f satisfies the assumptions of Lemma 5.2, we get a continuous function $\alpha : (0, \Delta_C(x_0)) \rightarrow [0, 1)$ such that for every $x \in C$ there is a positive integer k with $\|f^k(x) - x_0\| \leq \alpha(\|x - x_0\|)\|x - x_0\|$. Let $x \in C$, set $r_k = \|f^k(x) - x_0\|$ for every $k \in \mathbb{N}$ and note that the sequence $(r_k)_{k \in \mathbb{N}}$ is nonincreasing and has therefore a limit and infimum r . Assume that $r \neq 0$. Since α is continuous, we can find $\delta > 0$ such that $\alpha(t)t < r$ for every $t \in [r, r + \delta)$. Let k be such that $r_k < r + \delta$. Lemma 5.2 now yields $k' > k$ such that $r_{k'} \leq \alpha(r_k)r_k < r$. This contradicts the fact that r is the infimum, hence $r = 0$. \square

6. OPEN QUESTIONS

For a closed, convex, somewhat bounded set C in a separable Hilbert space H , the following questions are still open.

- (1) Does the generic mapping $f \in \mathcal{N}(C)$ have a unique fixed point even if C is not strictly convex?
- (2) Do the iterates $f^k(x)$ of the generic mapping $f \in \mathcal{N}(C)$ converge to a fixed point of f for every $x \in C$ even if C is not a LUR set?

However, the natural research direction concerns the possibility to extend our results to the Banach space setting. In what follows, C is a closed, convex set in a separable, infinite-dimensional Banach space.

- (3) Does it hold that the generic mapping $f \in \mathcal{N}(C)$ fails to have any fixed points in $\text{int}C$?
- (4) Let \mathcal{F} be the set of all $f \in \mathcal{N}(C)$ with a fixed point. Is \mathcal{F} either meager or residual?
- (5) Let \mathcal{F} be the set of all $f \in \mathcal{N}(C)$ which map a closed, convex, bounded subset of C to itself. Is \mathcal{F} either meager or residual?
- (6) What can be said about the uniqueness of fixed points and the convergence of the iterates for the generic $f \in \mathcal{N}(C)$ in case the set of all $f \in \mathcal{N}(C)$ with a fixed point is residual?

ACKNOWLEDGMENTS

The second Author was supported by the Doctoral Scholarship of the University of Innsbruck.

REFERENCES

- [1] C. Bargetz and M. Dymond. “ σ -porosity of the set of strict contractions in a space of non-expansive mappings”. *Israel Journal of Mathematics* 214 (2016), pp. 235–244.
- [2] C. Bargetz, M. Dymond, and S. Reich. “Porosity results for sets of strict contractions on geodesic metric spaces”. *Topol. Methods Nonlinear Anal.* 50.1 (2017), pp. 89–124.
- [3] C. Bargetz, S. Reich, and D. K. Thimm. “Generic properties of nonexpansive mappings on unbounded domains”. *Journal of Mathematical Analysis and Applications* 526.1 (Oct. 2023).
- [4] Y. Benyamini and J. Lindenstrauss. *Geometric Nonlinear Functional Analysis, Volume 1*. Vol. 48. American Mathematical Society, Colloquium Publications, 2000.
- [5] Y. Benyamini and Y. Sternfeld. “Spheres in infinite-dimensional normed spaces are Lipschitz contractible”. *Proceedings of the American Mathematical Society* 88.3 (1983), pp. 439–445.
- [6] C. A. de Bernardi and L. Veselý. “Rotundity properties, and non-extendability of Lipschitz quasiconvex functions”. *Journal of Convex Analysis* 30 (2023), pp. 329–342.
- [7] F. E. Browder. “Fixed-point theorems for noncompact mappings in Hilbert space”. *Proceedings of the National Academy of Sciences* 53.6 (1965), pp. 1272–1276.
- [8] F. S. De Blasi and J. Myjak. “Sur la convergence des approximations successives pour les contractions non linéaires dans un espace de Banach”. *Comptes Rendus de l’Académie des Sciences. Série I* 283 (1976), pp. 185–187.

- [9] F. S. De Blasi and J. Myjak. “Sur la porosité de l’ensemble des contractions sans point fixe”. *Comptes Rendus de l’Académie des Sciences. Série I* 308.2 (1989), pp. 51–54.
- [10] K. Goebel and W. A. Kirk. *Topics in metric fixed point theory*. Cambridge Studies in Advanced Mathematics 28. Cambridge university press, 1990.
- [11] K. Goebel and S. Reich. *Uniform convexity, hyperbolic geometry, and non-expansive mappings*. Monographs and Textbooks in Pure and Applied Mathematics 83. Marcel Dekker, Inc., New York, 1984.
- [12] V. L. Klee, L. Vesely, and C. Zanco. “Rotundity and smoothness of convex bodies in reflexive and nonreflexive spaces”. *Studia Mathematica* 120 (1996), pp. 191–204.
- [13] E. Rakotch. “A note on contractive mappings”. *Proceedings of the American Mathematical Society* 13.3 (1962), pp. 459–465.
- [14] D. Ravasini. “Generic uniformly continuous mappings on unbounded hyperbolic spaces”. *Journal of Mathematical Analysis and Applications* 538.1 (2024).
- [15] S. Reich and A. J. Zaslavski. “Almost all nonexpansive mappings are contractive”. *Mathematical Reports of the Academy of Sciences* 22.3 (2000), pp. 118–124.
- [16] S. Reich and A. J. Zaslavski. “The set of noncontractive mappings is σ -porous in the space of all nonexpansive mappings”. *Comptes Rendus de l’Académie des Sciences-Series I-Mathematics* 333.6 (2001), pp. 539–544.
- [17] F. Strobin. “Some porous and meager sets of continuous mappings”. *Journal of Nonlinear and Convex Analysis* 13.2 (2012), pp. 351–361.

(1) UNIVERSITÄT LEIPZIG, MATHEMATISCHES INSTITUT, AUGUSTUSPLATZ 10, 04109 LEIPZIG, GERMANY

Email address: `daive.ravasini@uni-leipzig.de`

(2) UNIVERSITÄT INNSBRUCK, DEPARTMENT OF MATHEMATICS, TECHNIKERSTRASSE 13, 6020 INNSBRUCK, AUSTRIA

Email address: `daylen.thimm@student.uibk.ac.at`