

Asymptotic integrability of nonlinear wave equations

A. M. Kamchatnov^{1,2}

¹*Institute of Spectroscopy, Russian Academy of Sciences, Troitsk, Moscow, 108840, Russia*

²*Skolkovo Institute of Science and Technology, Skolkovo, Moscow, 143026, Russia*

We introduce the notion of asymptotic integrability into the theory of nonlinear wave equations. It means that the Hamiltonian structure of equations describing propagation of high-frequency wave packets is preserved by hydrodynamic evolution of the large-scale background wave, so that these equations have an additional integral of motion. This condition is expressed mathematically as a system of equations for the carrier wave number as a function of the background variables. We show that a solution of this system for a given dispersion relation of linear waves is related with the quasiclassical limit of the Lax pair for the completely integrable equation having the corresponding dispersionless and linear dispersive behavior. We illustrate the theory by several examples.

PACS numbers: 02.30.Ik, 05.45.Yv, 47.35.Fg

At first sight it seems that the physical explanation of solitonic propagation as a manifestation of subtle balance of dispersive and nonlinear effects, on one side, and the mathematical idea of complete integrability of nonlinear wave equations, on the other side, are not related with each other. We show in this paper that a combined consideration of two limiting situations—the dispersionless evolution of large-scale waves and the propagation of high-frequency wave packets according to Hamilton’s optical-mechanical analogy—sheds new light on the complete integrability condition. As a result, we formulate a much weaker condition of asymptotic integrability which means that the Hamiltonian structure of equations for the packet’s propagation is preserved by the dispersionless evolution of the background wave providing thus an additional integral of motion. If this criterion is fulfilled, then we arrive at the quasiclassical limit of the Lax pair for some completely integrable equation. This theory is illustrated by several examples.

therein). At the same time, it seems that the original physical idea of interplay of nonlinear and dispersive effects has not played any essential role in this formal development. The aim of this article is to demonstrate that at least in some situations elaboration of this classical idea leads to interesting results useful for the nonlinear wave physics.

II. ASYMPTOTIC INTEGRABILITY

We suppose that our physical system is described by two variables which we will call for definiteness the “density” ρ and the “flow velocity” u . In one-dimensional geometry, they depend on the space x and time t variables: $\rho = \rho(x, t)$, $u = u(x, t)$. In the dispersionless limit of long wavelengths, the dynamics of our system is described by some hydrodynamic equations which can always be transformed (in case of two dependent variables) to a diagonal Riemann form (see, e.g.,¹³)

$$\frac{\partial r_+}{\partial t} + v_+ \frac{\partial r_+}{\partial x} = 0, \quad \frac{\partial r_-}{\partial t} + v_- \frac{\partial r_-}{\partial x} = 0 \quad (1)$$

for the so-called Riemann invariants $r_{\pm} = r_{\pm}(\rho, u)$. They are the most convenient variables in wave physics, since in many typical situations the ‘velocities’ v_{\pm} are given by the expressions

$$v_{\pm} = u \pm c, \quad (2)$$

where c is velocity of long sound waves propagating along a uniform background with constant density ρ and zero flow velocity u . This means that the velocities v_{\pm} correspond to the upstream or downstream propagation of sound waves. Naturally, both u and c as well as v_{\pm} can be expressed as functions of the Riemann invariants r_{\pm} . Equations (1) are nonlinear and they express the nonlinear properties of our system.

The dispersive properties of the system are described by the dispersion relations $\omega = \omega_{\pm}(k, r_+, r_-)$ (or $\omega = \omega_{\pm}(k, \rho, u)$) for harmonic linear waves $\propto \exp[i(kx - \omega t)]$ propagating along a uniform background with constant values of r_+, r_- (or ρ, u). It is important that in our formulation the dispersive and nonlinear properties are not independent of each other: the velocities (2) can be expressed as a long wavelength limit

I. INTRODUCTION

As was explained by classics of physics in the 19th century^{1–4}, the phenomenon of solitary wave (or solitonic) propagation is a result of subtle balance of nonlinear and dispersive effects of wave motion. Independently, the mathematicians of the 19th century developed the idea of complete integrability of equations of Newton dynamics (see, e.g.,⁵ and references therein), and this idea was based on some specific properties of Hamilton’s formulation of classical mechanics. Apparently, these two fundamental ideas had not been related with each other until the discovery of the inverse scattering transform method (IST) of integration of nonlinear wave equations^{6–8}. Soon after this discovery, it was realized^{9,10} that the nonlinear wave equations solvable by the IST method possess the property of complete integrability generalized to systems with infinite number of degrees of freedom. Since then, the Hamiltonian approach to nonlinear wave equations has become an important tool for their investigation and, as a result, the soliton physics has become a well-developed part of the modern mathematical physics (see, e.g.,^{11,12} and references

of the phase velocities of linear waves:

$$v_{\pm} = \lim_{k \rightarrow 0} \frac{\omega_{\pm}(k)}{k}, \quad (3)$$

so the two dispersion relations correspond to the two directions of propagation of linear waves.

Let now the background variables change little at distances of the order of the size of the short-wavelength wave packet built from the linear waves with wave numbers around the carrier wave number k . Then we can introduce the coordinate $x(t)$ of such a wave packet and consider its motion along a nonuniform time-dependent background wave whose evolution obeys Eqs. (1). In this case, the frequencies ω_{\pm} become slow functions of x and t via the background variables $r_{\pm} = r_{\pm}(x, t)$. Consequently, propagation of our wave packet obeys the Hamilton equations (see, e.g.,^{14,15})

$$\frac{dx}{dt} = \frac{\partial \omega}{\partial k}, \quad \frac{dk}{dt} = -\frac{\partial \omega}{\partial x}, \quad (4)$$

where ω is given by one of the branches ω_{\pm} of the dispersion relation. Thus, to find the path of the wave packet, we have to solve at first the hydrodynamics equations (1), then substitute the solution into the dispersion relation $\omega(k, r_+, r_-)$, and, at last, to solve the Hamilton equations (4).

The above general formulation of the packet's propagation problem refers to arbitrary dispersion relations and related with them according to Eq. (3) large-scale hydrodynamic equations. Now we want to specify this formulation in such a way, that it becomes 'integrable'. This means that the Hamilton equations (4) must have an integral of motion which allows one to solve these equations in quadratures. Thus, we demand that there exists such a function $k = k(r_+, r_-)$ that the Hamilton equations (4) are compatible with the background evolution according to the hydrodynamic equations (1). To express this condition in analytical form, let us consider two moments of time separated by a small interval dt and, consequently, by a small distance $dx = v_g dt$ between the corresponding locations of the wave packet, where $v_g = \partial \omega / \partial k$ is the group velocity of the wave packet (see the first Eq. (4)). Then the differences of the values of the background Riemann invariants at the locations of the wave packet are equal to

$$dr_{\pm} = \frac{\partial r_{\pm}}{\partial x} dx + \frac{\partial r_{\pm}}{\partial t} dt = \frac{\partial r_{\pm}}{\partial x} (v_g - v_{\pm}) dt,$$

where we have taken into account the hydrodynamic equations (1). This relation gives

$$\frac{\partial r_{\pm}}{\partial x} dt = -\frac{dr_{\pm}}{v_{\pm} - v_g}$$

and, consequently, the second Hamilton equation (4) yields the following expression for the small change of the wave number:

$$\begin{aligned} dk &= -\frac{\partial \omega}{\partial x} dt = -\left(\frac{\partial \omega}{\partial r_+} \frac{\partial r_+}{\partial x} + \frac{\partial \omega}{\partial r_-} \frac{\partial r_-}{\partial x} \right) dt \\ &= \frac{\partial \omega / \partial r_+}{v_+ - v_g} dr_+ + \frac{\partial \omega / \partial r_-}{v_- - v_g} dr_-. \end{aligned}$$

On the other hand, our assumption that the wave number k is only a function $k = k(r_+, r_-)$ of the background variables gives

$$dk = \frac{\partial k}{\partial r_+} dr_+ + \frac{\partial k}{\partial r_-} dr_-.$$

Since we consider an arbitrary general solution of Eqs. (1), the differentials dr_+ and dr_- can be considered as independent of each other, so comparison of the above expressions yields the system of equations

$$\frac{\partial k}{\partial r_+} = \frac{\partial \omega / \partial r_+}{v_+ - v_g}, \quad \frac{\partial k}{\partial r_-} = \frac{\partial \omega / \partial r_-}{v_- - v_g}, \quad (5)$$

which define the function $k = k(r_+, r_-)$ of two independent variables r_+, r_- , which is the desirable integral of motion. Naturally, for existence of such a function these derivatives must commute,

$$\frac{\partial}{\partial r_-} \left(\frac{\partial k}{\partial r_+} \right) = \frac{\partial}{\partial r_+} \left(\frac{\partial k}{\partial r_-} \right), \quad (6)$$

and this *asymptotic integrability condition* is only fulfilled for some special choices of the dispersion relation $\omega = \omega(k, r_+, r_-)$ and, consequently, of the hydrodynamic equations (1).

It is worth noticing that if we consider particular solutions of Eqs. (1) in the form of simple waves, then one of the Riemann invariants is constant and the above calculation leads to a single equation for the wave number $k = k(r)$ which becomes a function of the only varying Riemann invariant r ,

$$\frac{dk}{dr} = \frac{\partial \omega / \partial r}{V - v_g}, \quad (7)$$

where $\omega = \omega(k, r)$ and $V = \lim_{k \rightarrow 0} (\omega(k)/k)$, $v_g = \partial \omega(k) / \partial k$. This equation was obtained in Ref.¹⁶ in the theory of dispersive shock waves as a small-amplitude limit of the Whitham modulation equations and in Refs.¹⁷ from the Hamilton theory of wave packets propagation. The general equations (5) were considered in Refs.^{18,19} with applications to the problems of propagation of linear wave packets and asymptotic theory of solitons produced from an intensive initial pulse. Here we shall consider the relation of the asymptotic integrability condition to the complete integrability of nonlinear wave equations.

III. TRANSITION TO THE COMPLETE INTEGRABILITY

As was noticed in Ref.²⁰, the solution of Eqs. (5) is related with the quasiclassical limit of Lax pairs of nonlinear wave equations completely integrable in framework of the Ablowitz-Kaup-Newell-Segur (AKNS) scheme²¹ written in a scalar form²². Let the completely integrable equation can be written as a compatibility condition of two linear equations

$$\phi_{xx} = \mathcal{A} \phi, \quad \phi_t = -\frac{1}{2} \mathcal{B}_x \phi + \mathcal{B} \phi_x, \quad (8)$$

where the functions \mathcal{A} and \mathcal{B} depend on the wave variables ρ, u , their space derivatives of different orders, and on the spectral parameter λ . If we turn to the quasiclassical limit of these equations with fast oscillations of ϕ , so that we can neglect all the derivatives of the smooth functions ρ and u in the expressions for \mathcal{A} and \mathcal{B} , then we arrive at their quasiclassical limits $\overline{\mathcal{A}}$ and $\overline{\mathcal{B}}$ which only depend on the wave variables ρ and u and the spectral parameter.

The nonlinear equations under consideration reduce in the dispersionless limit to some hydrodynamic-like equations which can be transformed to the form (1) for the Riemann invariants r_+, r_- . On the other hand, linearization of our completely integrable equations yields the dispersion relation $\omega = \omega(k, r_+, r_-)$. Thus, we arrive at the system (5) which in case of completely integrable equations has a solution

$$k = k(r_+, r_-, q), \quad (9)$$

where q is an integration constant. As was shown in Ref.²⁰, the functions $\overline{\mathcal{A}}$ and $\overline{\mathcal{B}}$ are related with k and ω by the formulas

$$\begin{aligned} \overline{\mathcal{A}} &= -\frac{1}{4}k^2(r_+, r_-, q), \\ \overline{\mathcal{B}} &= -\frac{\omega(k(r_+, r_-, q), r_+, r_-)}{k(r_+, r_-, q)}, \end{aligned} \quad (10)$$

where q is related with the spectral parameter λ of the IST method. It is natural to ask, whether it is possible to go in the opposite direction and starting from the hydrodynamic equations and dispersion relation which satisfy the asymptotic integrability condition (6), so that the solution of Eqs. (5) is known, to arrive at the Lax pair (8) of some completely integrable nonlinear wave equation.

Actually, this procedure of generalization of quasiclassical formulas to their fully dispersive counterparts is analogous to the well-known quantization procedure of classical mechanical systems resulting in equations for their quantum-mechanical behavior. As is well known, such a quantization procedure is not trivial and unique, so we shall confine ourselves to the simplest methods widely used in physics.

In the first method, we can consider the functions $\overline{\mathcal{A}}, \overline{\mathcal{B}}$ given by Eqs. (10) as exact functions defining the Lax pair (8), so their compatibility condition $(\phi_{xx})_t = (\phi_t)_{xx}$, i.e.,

$$\mathcal{A}_t - 2\mathcal{B}_x \mathcal{A} - \mathcal{B} \mathcal{A}_x + \frac{1}{2}\mathcal{B}_{xxx} = 0 \quad (11)$$

yields the corresponding nonlinear wave system.

In the second method, we can transform Eqs. (10) to the variables ρ, u and combine these variables to a complex field

$$\psi = \sqrt{\rho} \exp\left(i \int^x u(x', t) dx'\right) \quad (12)$$

according to the well-known Madelung transformation²³ of the quantum-mechanical wave function. Then in the quasiclassical limit we have $u \approx -i\psi_x/\psi$ or $u \approx i\psi_x^*/\psi^*$, so we can try to return to the nonlinear dispersive equation for the field ψ by the replacements

$$\rho \rightarrow \psi\psi^*, \quad u \rightarrow -\frac{i\psi_x}{\psi} \quad \text{or} \quad u \rightarrow \frac{i\psi_x^*}{\psi^*} \quad (13)$$

in the quasiclassical expressions $\overline{\mathcal{A}}$ and $\overline{\mathcal{B}}$. Besides that, we might have to add to these functions some terms vanishing in the quasiclassical limit. The form of these term can be guessed by means of analogy with known results and correctness of their choice can be proved by checking the compatibility condition (11).

Let us illustrate this approach by several examples.

IV. SHALLOW WATER AND BOSE-EINSTEIN CONDENSATE

Let the system be described in the dispersionless limit by the shallow water equations

$$\rho_t + (\rho u)_x = 0, \quad u_t + uu_x + \rho_x = 0. \quad (14)$$

As a dispersion relation we choose the formula

$$\omega = k \left(u \pm \sqrt{\rho + \frac{k^2}{4}} \right) \quad (15)$$

which in the long wavelength limit gives Eqs. (2) with $c = \sqrt{\rho}$, that is

$$v_{\pm} = u \pm \sqrt{\rho}. \quad (16)$$

The Riemann invariants are defined by the formulas

$$r_{\pm} = u/2 \pm \sqrt{\rho}. \quad (17)$$

It is convenient to rewrite Eqs. (5) in terms of ρ and u , so we get the equations

$$\frac{\partial k^2}{\partial \rho} = -4, \quad \frac{\partial k^2}{\partial u} = -2\sqrt{k^2 + 4\rho}, \quad (18)$$

which define the function $k = k(\rho, u)$. The asymptotic integrability condition (6) is fulfilled and Eqs. (18) have the solution

$$\begin{aligned} k^2 &= (q - u)^2 - 4\rho \\ &= (q - 2r_+)(q - 2r_-). \end{aligned} \quad (19)$$

If we assume that $q > u$ and choose the upper sign in Eq. (15), then we obtain

$$\frac{\omega}{k} = \frac{1}{2}(q + u). \quad (20)$$

The first method of quantization yields the Lax pair (8) with

$$\mathcal{A} = \rho - \frac{1}{4}(q - u)^2, \quad \mathcal{B} = -\frac{1}{2}(q + u), \quad (21)$$

where q plays the role of the spectral parameter. This is the Lax pair for the Kaup-Boussinesq system²⁴

$$\rho_t + (\rho u)_x - \frac{1}{4}\rho_{xxx} = 0, \quad u_t + uu_x + \rho_x = 0. \quad (22)$$

It finds applications to dynamics of shallow water dispersive waves.

If we treat the system (14) as Landau equations²⁵ for the superfluid component and assume that Eq. (15) is the Bogoliubov dispersion relation²⁶ for linear waves propagating along a uniform condensate, then the second method of quantization seems appropriate for the condensate's wave function $\psi = \sqrt{\rho} \exp(i \int^x u(x', t) dx')$. We make replacements (13) in Eqs. (21) and after adding a higher order term for fulfillment of Eq. (11) we arrive at the Lax pair with

$$\begin{aligned} \mathcal{A} &= -\frac{1}{4} \left(q + \frac{i\psi_x}{\psi} \right)^2 + \psi\psi^* - \left(\frac{\psi_x}{2\psi} \right)_x, \\ \mathcal{B} &= -\frac{1}{2} \left(q - \frac{i\psi_x}{\psi} \right). \end{aligned} \quad (23)$$

This is the well-known scalar form of the Lax pair for the Gross-Pitaevskii^{27,28} (or nonlinear Schrödinger (NLS)²⁹) equation

$$i\psi_t + \frac{1}{2}\psi_{xx} - |\psi|^2\psi = 0. \quad (24)$$

V. DERIVATIVE NLS EQUATIONS

As we saw above, the shallow water equations (14) and the dispersion relation (15) lead to different completely integrable systems depending on the method of 'quantization' of the solution (19) of the asymptotic integrability equations. Consequently, if we have some completely integrable equation for a complex field $\psi = \sqrt{\rho} \exp(i \int^x u(x', t) dx')$, then the corresponding dispersionless system and dispersion relation for linear waves are also known and they satisfy the asymptotic integrability condition, so the solution (9) can be found together with the quasiclassical limit (10) of the functions \mathcal{A} and \mathcal{B} . Then they define another completely integrable system for the variables ρ and u which can be considered as a 'shallow water' counterpart of the equation for the complex field ψ . Here we shall consider such systems obtained from three forms of the derivative NLS (DNLS) equation.

A. Kaup-Newell DNLS equation

The Kaup-Newell DNLS equation³⁰ has the form

$$i\psi_t + \frac{1}{2}\psi_{xx} - i(|\psi|^2\psi)_x = 0. \quad (25)$$

The substitution $\psi = \sqrt{\rho} \exp(i \int^x u(x', t) dx')$ yields in the dispersionless limit the hydrodynamic system

$$\begin{aligned} \rho_t + \left[\rho \left(u - \frac{3}{2}\rho \right) \right]_x &= 0, \\ u_t + uu_x - (\rho u)_x &= 0. \end{aligned} \quad (26)$$

The characteristic velocities are given by the formulas

$$v_{\pm} = u - 2\rho \pm \sqrt{\rho(\rho - u)} \quad (27)$$

and Eqs. (26) can be cast to the diagonal form (1) for the Riemann invariants

$$r_{\pm} = \frac{u}{2} - \rho \pm \sqrt{\rho(\rho - u)}. \quad (28)$$

Linearization of Eq. (25) for small-amplitude waves propagating along a uniform background yields the dispersion relation

$$\omega = k \left(u - 2\rho \pm \sqrt{\rho(\rho - u) + \frac{k^2}{4}} \right). \quad (29)$$

Eqs. (5) can be transformed to the variables ρ, u , so they read

$$\begin{aligned} \frac{\partial k^2}{\partial \rho} &= 4 \left[\sqrt{k^2 + 4\rho(\rho - u)} - (2\rho - u) \right], \\ \frac{\partial k^2}{\partial u} &= -2 \left[\sqrt{k^2 + 4\rho(\rho - u)} - 2\rho \right] \end{aligned} \quad (30)$$

and have the solution

$$\begin{aligned} k^2 &= (q - u)^2 + 4q\rho \\ &= (q - 2r_+)(q - 2r_-), \end{aligned} \quad (31)$$

where q is an integration constant, so that

$$\frac{\omega}{k} = \frac{q}{2} + \frac{u}{2} - \rho. \quad (32)$$

We define the Lax pair according to Eqs. (10),

$$\mathcal{A} = -\frac{1}{4}(q - u)^2 - q\rho, \quad \mathcal{B} = -\frac{q}{2} - \frac{u}{2} + \rho, \quad (33)$$

where q plays the role of the spectral parameter. The compatibility condition yields the completely integrable system

$$\begin{aligned} \rho_t + \left[\rho \left(u - \frac{3}{2}\rho \right) \right]_x + \frac{1}{4u}(u - 2\rho)_{xxx} &= 0, \\ u_t + uu_x - (\rho u)_x + \frac{1}{2u}(u - 2\rho)_{xxx} &= 0. \end{aligned} \quad (34)$$

Obviously, this is a dispersive generalization of the hydrodynamic system (26) different from the Kaup-Newell equation transformed to the (ρ, u) -variables:

$$\begin{aligned} \rho_t + \left[\rho \left(u - \frac{3}{2}\rho \right) \right]_x &= 0, \\ u_t + uu_x - (\rho u)_x + \left(\frac{\rho_x^2}{8\rho^2} - \frac{\rho_{xx}}{4\rho} \right)_x &= 0. \end{aligned} \quad (35)$$

Although the substitutions

$$\bar{\rho} = \rho(\rho - u), \quad \bar{u} = u - 2\rho \quad (36)$$

transform Eqs. (34) to the Kaup-Boussinesq system (22) for the variables $(\bar{\rho}, \bar{u})$, the inverse transform is not single-valued, the system (34) is not genuinely nonlinear and describes nonlinear waves different from those for the Kaup-Boussinesq system.

B. Chen-Lee-Liu DNLS equation

The Chen-Lee-Liu equation³¹

$$i\psi_t + \frac{1}{2}\psi_{xx} - i|\psi|^2\psi_x = 0 \quad (37)$$

reduces after Madelung transformation (12) to the dispersionless equations

$$\rho_t + \left[\rho \left(u - \frac{\rho}{2} \right) \right]_x = 0, \quad u_t + uu_x - (\rho u)_x = 0 \quad (38)$$

and leads to the dispersion relation

$$\omega = k \left(u - \rho \pm \sqrt{-\rho u + \frac{k^2}{4}} \right), \quad (39)$$

so we get the characteristic velocities

$$v_{\pm} = u - \rho \pm \sqrt{-\rho u}. \quad (40)$$

The Riemann invariants can be defined by the formulas

$$r_{\pm} = \frac{1}{2}(u - \rho) \pm \sqrt{-\rho u}. \quad (41)$$

The asymptotic integrability equations

$$\begin{aligned} \frac{\partial k^2}{\partial \rho} &= 4u + 2\sqrt{k^2 - 4\rho u}, \\ \frac{\partial k^2}{\partial u} &= 4\rho - 2\sqrt{k^2 - 4\rho u} \end{aligned} \quad (42)$$

have the solution

$$\begin{aligned} k^2 &= (q - u + \rho)^2 + 4\rho u \\ &= (q - 2r_+)(q - 2r_-), \end{aligned} \quad (43)$$

so

$$\frac{\omega}{k} = \frac{1}{2}(q + u - \rho). \quad (44)$$

As a result we arrive at the Lax pair with

$$\mathcal{A} = -\frac{1}{4}(q - u + \rho)^2 - \rho u, \quad \mathcal{B} = -\frac{1}{2}(q + u - \rho), \quad (45)$$

and, consequently, obtain the completely integrable system

$$\begin{aligned} \rho_t + \left[\rho \left(u - \frac{\rho}{2} \right) \right]_x + \frac{1}{4(\rho + u)}(u - \rho)_{xxx} &= 0, \\ u_t + uu_x - (\rho u)_x + \frac{1}{4(\rho + u)}(u - \rho)_{xxx} &= 0. \end{aligned} \quad (46)$$

Again this system reduces to the Kaup-Boussinesq system for the variables

$$\bar{\rho} = -\rho u, \quad \bar{u} = u - \rho. \quad (47)$$

C. Gerdjikov-Ivanov DNLS equation

The Gerdjikov-Ivanov equation³²

$$i\psi_t + \psi_{xx} + i\psi^2\psi_x^* + \frac{1}{2}|\psi|^4\psi = 0 \quad (48)$$

yields the dispersionless equations

$$\begin{aligned} \rho_t + \left[\rho \left(2u + \frac{\rho}{2} \right) \right]_x &= 0, \\ u_t + 2uu_x - (\rho u)_x - \rho\rho_x &= 0, \end{aligned} \quad (49)$$

the dispersion relation

$$\omega = k \left(2u \pm \sqrt{-\rho(\rho + 2u) + k^2} \right), \quad (50)$$

and the characteristic velocities

$$v_{\pm} = 2u \pm \sqrt{-\rho(\rho + 2u)}. \quad (51)$$

The Riemann invariants are defined by the formulas

$$r_{\pm} = u \pm \sqrt{-\rho(\rho + 2u)}. \quad (52)$$

The asymptotic integrability equations

$$\begin{aligned} \frac{\partial k^2}{\partial \rho} &= 2(u + \rho), \\ \frac{\partial k^2}{\partial u} &= 2\rho - 2\sqrt{k^2 - 4\rho(\rho + 2u)} \end{aligned} \quad (53)$$

give the solution

$$\begin{aligned} k^2 &= (q - u)^2 + \rho(\rho + 2u) \\ &= (q - r_+)(q - r_-), \end{aligned} \quad (54)$$

so

$$\frac{\omega}{k} = q + u. \quad (55)$$

Consequently, we obtain the Lax pair with

$$\mathcal{A} = -\frac{1}{4}(q - u)^2 - \frac{1}{4}\rho(\rho + 2u), \quad \mathcal{B} = -q - u, \quad (56)$$

and the corresponding completely integrable system

$$\begin{aligned} \rho_t + \left[\rho \left(2u + \frac{\rho}{2} \right) \right]_x + \frac{1}{\rho + u}u_{xxx} &= 0, \\ u_t + 2uu_x - (\rho u)_x - \rho\rho_x &= 0. \end{aligned} \quad (57)$$

which can be transformed to the Kaup-Boussinesq system for the variables

$$\bar{\rho} = -\rho(\rho + 2u), \quad \bar{u} = 2u. \quad (58)$$

As is known, the above three DNLS equations are related with each other by gauge transformations³³⁻³⁵. As we see, their 'shallow water' counterparts are related with each other by simple algebraic formulas. At the same time, the physical meaning of these equations can be different, as it will be especially clearly seen in one more example considered in the next section.

VI. LANDAU-LIFSHITZ EQUATION AND TWO-LAYER 'SHALLOW WATER' MODEL

The Landau-Lifshitz equation

$$\mathbf{M}_t = (\mathbf{M}_{xx} - M_3 \mathbf{e}_3) \wedge \mathbf{M} \quad (59)$$

for the magnetization vector $\mathbf{M} = (M_1, M_2, M_3)$, $|\mathbf{M}| = 1$, with an easy-plane anisotropy, can be transformed to more convenient variables w, v by a Madelung-like substitution³⁶⁻³⁸

$$\begin{aligned} M_3 &= -w, \\ M_- &\equiv M_1 - iM_2 \\ &= \sqrt{1-w^2} \exp\left(-i \int^x v dx'\right). \end{aligned} \quad (60)$$

The resulting equations

$$\begin{aligned} w_- [v(1-w^2)]_x &= 0, \\ v_t - [w(1-v^2)]_x + \left[\frac{1}{\sqrt{1-w^2}} \left(\frac{w_x}{\sqrt{1-w^2}} \right) \right]_x &= 0 \end{aligned} \quad (61)$$

reduce to the dispersionless equations

$$\begin{aligned} w_t - [v(1-w^2)]_x &= 0, \\ v_t - [w(1-v^2)]_x &= 0 \end{aligned} \quad (62)$$

which can be transformed to the form (1) by introduction of the Riemann invariants

$$r_{\pm} = vw \pm \sqrt{(1-v^2)(1-w^2)}. \quad (63)$$

The characteristic velocities of Eqs. (62) are equal to

$$v_{\pm} = 2vw \pm \sqrt{(1-v^2)(1-w^2)}. \quad (64)$$

The whole system (61) after linearization with respect to small-amplitude waves propagating along a uniform background yields the dispersion relation

$$\omega = k \left(2vw \pm \sqrt{k^2 + (1-v^2)(1-w^2)} \right) \quad (65)$$

for harmonic waves. The asymptotic integrability equations (5) written in terms of the (v, w) -variables

$$\begin{aligned} \frac{\partial k^2}{\partial w} &= 2 \left[w(1-v^2) - v \sqrt{k^2 + (1-v^2)(1-w^2)} \right], \\ \frac{\partial k^2}{\partial v} &= 2 \left[v(1-w^2) - w \sqrt{k^2 + (1-v^2)(1-w^2)} \right], \end{aligned} \quad (66)$$

have the solution

$$\begin{aligned} k^2 &= (q-vw)^2 - (1-v^2)(1-w^2) \\ &= (q-r_+)(q-r_-) \end{aligned} \quad (67)$$

so (assuming $q-vw > 0$) we get

$$\frac{\omega}{k} = q + vw. \quad (68)$$

We define the Lax pair (8) with

$$\begin{aligned} \mathcal{A} &= \frac{1}{4} [(1-v^2)(1-w^2) - (q-vw)^2], \\ \mathcal{B} &= -q - vw \end{aligned} \quad (69)$$

and obtain the corresponding completely integrable equations

$$\begin{aligned} w_t - [v(1-w^2)]_x + \frac{v(vw)_{xxx}}{v^2-w^2} &= 0, \\ v_t - [w(1-v^2)]_x - \frac{w(vw)_{xxx}}{v^2-w^2} &= 0. \end{aligned} \quad (70)$$

This is a 'shallow water' counterpart of the Landau-Lifshitz equation (59).

It is remarkable that the dispersionless equations (62) were derived in Ref.³⁹ as a long wave shallow water approximation for dynamics of inner waves in the fluid composed of two thin layers with very small difference of densities of liquids in them (the lighter layer above the heavier one). The dispersion relation for linear inner waves in a two-layer system is described by a more complicated formula than Eq. (65) (see, e.g., Ref.⁴⁰). Nevertheless, it has similar general structure, so Eqs. (70) can be considered as a 'toy model' for dispersive dynamics of shallow inner waves. Although Eqs. (70) can be reduced to the Kaup-Boussinesq system for the variables

$$\bar{p} = (1-v^2)(1-w^2), \quad \bar{u} = 2vw, \quad (71)$$

the qualitative properties of these two systems are very different. For example, the Kaup-Boussinesq system is genuinely nonlinear and an initial discontinuity evolves according to it into one of six possible wave structures^{41,42}. The system (70) is not genuinely nonlinear and classification of possible wave structures includes various combined dispersive shock waves similar to the structures appearing in the theory of the Landau-Lifshitz equation³⁷. One might expect that such a classification of possible wave structures evolving from initial discontinuities in the theory of Eqs. (70) is qualitatively similar to such a classification for the actual two-layer dispersion relation.

As one can see, although the considered above equations belong to different physical situations, they all correspond to the same solution of the asymptotic integrability equations with the dispersion relation

$$\omega = k \left(r_+ + r_- \pm \sqrt{\frac{1}{4}(r_+ - r_-)^2 + k^2} \right), \quad (72)$$

so that the long wavelength characteristic velocities are given by

$$v_{\pm} = r_+ + r_- \pm \frac{1}{2}(r_+ - r_-), \quad (73)$$

and the solution can be written in the form

$$k^2 = \left[q - \frac{1}{2}(r_+ + r_-) \right]^2 - \frac{1}{4}(r_+ - r_-)^2. \quad (74)$$

Only the formulas connecting the Riemann invariants r_{\pm} with the physical variables depend on the choice of the physical system and the above list of equations is not exhaustive. In particular, one can easily check that the Jaulent-Miodek system^{45,46} is described by the same solution of the asymptotic integrability equations.

VII. ZAKHAROV-ITO SYSTEM

To demonstrate another type of solution of the asymptotic integrability equations, let us consider the hydrodynamic-like equations written in the Riemann form (1), where

$$v_+ = 6r_+ + 2r_-, \quad v_- = 2r_+ + 6r_- \quad (75)$$

One can check that the asymptotic integrability condition is fulfilled if we choose the dispersion relation

$$\omega = k \left\{ 4(r_+ + r_-) + k^2/2 \pm \sqrt{[2(r_+ + r_-) + k^2/2]^2 - 16r_+r_-} \right\}, \quad (76)$$

so that Eqs. (5) have the solution

$$k^2 = \frac{4}{q}(q - r_+)(q - r_-), \quad (77)$$

and then

$$\frac{\omega}{k} = 4q + 2(r_+ + r_-). \quad (78)$$

Let the Riemann invariants be related with the ‘physical’ variables ρ, u by the formulas

$$r_+ = \frac{1}{2}(u + \sqrt{u^2 - 4\rho}), \quad r_- = \frac{1}{2}(u - \sqrt{u^2 - 4\rho}), \quad (79)$$

so that

$$k^2 = 4 \left(q - u + \frac{\rho}{q} \right), \quad \frac{\omega}{k} = 4q + 2u. \quad (80)$$

Transition to the Lax pairs according to the rules (10) yields

$$\mathcal{A} = -q + u - \frac{\rho}{q}, \quad \mathcal{B} = -4q - 2u, \quad (81)$$

where q plays the role of the spectral parameter. The resulting completely integrable system is the well-known Zakharov-Ito system^{43,44}

$$\begin{aligned} \rho_t + 2u\rho_x + 4\rho u_x &= 0, \\ u_t + 6uu_x - 4\rho_x - u_{xxx} &= 0. \end{aligned} \quad (82)$$

It is worth noticing that although the dispersionless system with the velocities (75) written in (ρ, u) -variables

$$\begin{aligned} \rho_t + 2u\rho_x + 4\rho u_x &= 0, \\ u_t + 6uu_x - 4\rho_x &= 0 \end{aligned} \quad (83)$$

can be transformed to the shallow water system (14) for the variables

$$\bar{\rho} = 4u^2 - 16\rho, \quad \bar{u} = 4u, \quad (84)$$

these substitutions are not compatible with transformation of the dispersion relation (15). In fact, they cast the dispersion relation (76) to the form

$$\omega = k \left(4u + \frac{k^2}{2} \pm \sqrt{\left(2u + \frac{k^2}{2} \right)^2 - 16\rho} \right) \quad (85)$$

different from the relation (15) for the Kaup-Boussinesq system.

VIII. CONCLUSION

In this paper, we introduced the physically natural condition of asymptotic integrability of nonlinear wave equations which means that two asymptotic limits—dispersionless (hydrodynamic) evolution of a smooth background wave and propagation of high-frequency wave packets—are compatible with each other in the sense that the Hamiltonian structure of the packet’s propagation is preserved by evolution of the background wave and, consequently, the Hamilton equations have an integral of motion. In fact, this weaker integrability condition leads to the quasiclassical limit of complete integrability in framework of the AKNS scheme²⁰ and if the Lax pair of the equations under consideration does not depend on the space derivatives of the wave variables, then the asymptotic integrability condition reproduces the exact Lax pair. This observation sheds new light on the origin of the completely integrable nonlinear wave equations.

Besides that, it should be noticed that the relationship for the dependence of the packet’s wave number on the background variables can be converted to a similar relationship for the inverse half-width of narrow solitons propagating along smooth background, and this remark allows one to develop a Hamiltonian theory for propagation of narrow solitons along a non-uniform and time dependent background^{47–49}.

At last, Eqs. (5) of asymptotic integrability can also have approximate solutions correct in the limit of large wave numbers k even for not completely integrable equations. In these quite general situations, one can develop the theory of propagation of small-amplitude wave packets along a non-uniform and time-dependent background¹⁸ and to formulate a generalized Bohr-Sommerfeld quantization rule which determines the parameters of solitons produced from an intensive initial pulse¹⁹. Thus, the condition of asymptotic integrability turns out to be a useful tool for investigation of various problems in nonlinear physics.

ACKNOWLEDGMENTS

I thank E. A. Kuznetsov and M. V. Pavlov for useful discussions. This research is funded by the research project FFUU-2021-0003 of the Institute of Spectroscopy of the Russian

Academy of Sciences (Sections 1-3) and by the RSF grant number 19-72-30028 (Sections 4-7).

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the author upon reasonable request.

- ¹J. Boussinesq, Théorie des ondes et des remous qui se propagent le long d'un canal rectangulaire horizontal, en communiquant au liquide contenu dans ce canal des vitesses sensiblement pareilles de la surface au fond, *J. Math. Pures Appl.*, **17**, 55–108 (1872).
- ²Lord Rayleigh, On waves, *Phil. Mag.*, **1**, 257–279 (1876).
- ³J. J. Stokes, Note on the theory of the solitary waves, *Phil. Mag.* **31**, 314–316 (1891).
- ⁴D. J. Korteweg and G. de Vries, On the Change of Form of Long Waves advancing in a Rectangular Canal, and on a New Type of Long Stationary Waves, *Phil. Mag.*, **39**, 422–443 (1895).
- ⁵E. T. Whittaker, *Analytical Dynamics*, (Cambridge, CUP, 1927).
- ⁶S. C. Gardner, J. M. Greene, M. D. Kruskal, R. M. Miura, Method for solving the Korteweg-de Vries equation, *Phys. Rev. Lett.*, **19**, 1095–1097 (1967).
- ⁷P. D. Lax, Integrals of Nonlinear Equations of Evolution and Solitary Waves, *Comm. Pure Appl. Math.* **21**, 467–490 (1968).
- ⁸V. E. Zakharov, A. B. Shabat, Exact theory of two-dimensional self-focusing and one-dimensional self-modulation of waves in nonlinear media, *Zh. Eksp. Teor. Fiz.* **61**, 118–134 (1971) [*Sov. Phys. JETP*, **34**, 62–69 (1972)].
- ⁹V. E. Zakharov and L. D. Faddeev, Korteweg-de Vries equation: A completely integrable Hamiltonian system, *Funl. Analiz Prilozh.*, **5**, 18–27 (1971) [*Func. Anal. Appl.* **5**, 280–287 (1971)].
- ¹⁰S. C. Gardner, Korteweg-de Vries equation and generalizations. IV. Korteweg-de Vries as a Hamiltonian system, *J. Math. Phys.* **12**, 1548–1551 (1971).
- ¹¹L. A. Dickey, *Soliton Equations and Hamiltonian Systems*, (World Scientific, Singapore, 2003).
- ¹²L. D. Faddeev, L. A. Takhtajan, *Hamiltonian Methods in the Theory of Solitons*, (Springer, Berlin, 2007).
- ¹³B. L. Roždestvenskii and N. N. Yanenko, *Systems of Quasilinear Equations and Their Applications to Gas Dynamics*, (Providence, AMS, 1983).
- ¹⁴L. D. Landau and E. M. Lifshitz, *The Classical Theory of Fields*, (Pergamon, Oxford, 1971).
- ¹⁵Yu. A. Kravtsov, Yu. I. Orlov, *Geometrical Optics of Inhomogeneous Media*, (Springer, Berlin, 1990).
- ¹⁶G. A. El, Chaos, Resolution of a shock in hyperbolic systems modified by weak dispersion, **15**, 037103 (2005).
- ¹⁷A. M. Kamchatnov, Theory of quasi-simple dispersive shock waves and number of solitons evolved from a nonlinear pulse, *Chaos* **30**, 123148 (2020).
- ¹⁸D. V. Shaykin, and A. M. Kamchatnov, Propagation of wave packets along large-scale background waves, *Phys. Fluids*, **35**, 062108 (2023).
- ¹⁹A. M. Kamchatnov, Asymptotic theory of not completely integrable soliton equations, *Chaos* **33**, 093105 (2023).
- ²⁰A. M. Kamchatnov, D. V. Shaykin, Quasiclassical integrability condition in AKNS scheme, *Physica D*, **460**, 134085 (2024).
- ²¹M. J. Ablowitz, D. J. Kaup, A. C. Newell, H. Segur, The inverse scattering transform—Fourier analysis for nonlinear problems, *Stud. Appl. Math.*, **53**, 249–315 (1974).
- ²²A. M. Kamchatnov, R. A. Kraenkel, On the relationship between 2×2 matrix and second-order scalar spectral problems for integrable equations, *J. Phys. A: Math. Gen.* **35**, L13–L18 (2002).
- ²³E. Madelung, Quantentheorie in hydrodynamischer Form, *Z. Phys.* **40**, 322–326 (1927).
- ²⁴D. J. Kaup, A higher-order water-wave equation and the method for solving it, *Prog. Theor. Phys.* **54**, 396–408 (1975).
- ²⁵L. D. Landau, The theory of superfluidity of helium II, *J. Phys. (USSR)* **5**, 71–100 (1941).
- ²⁶N. N. Bogoliubov. On the theory of superfluidity, *J. Phys. (USSR)*, **11**, 23–32 (1947).
- ²⁷E. P. Gross, Structure of a Quantized Vertex in Boson Systems, *Nuovo Cimento*, **20**, 454–477 (1961).
- ²⁸L. P. Pitaevskii, Vortex Lines in an Imperfect Bose Gas, *Zh. Eksp. Teor. Fiz.* **40**, 646–651 (1961) [*Sov. Phys. JETP* **13**, 451–454 (1961)].
- ²⁹D. J. Benney and A. C. Newell, The propagation of nonlinear wave envelopes, *J. Math. & Phys.*, **46**, 133–139 (1967).
- ³⁰D. J. Kaup and A. C. Newell, An Exact Solution for a Derivative Nonlinear Schrödinger Equation, *J. Math. Phys.* **19**, 798–801 (1978).
- ³¹H. H. Chen, Y. C. Lee, C. S. Liu, Integrability of Nonlinear Hamiltonian Systems by Inverse Scattering Method, *Phys. Scr.* **20**, 490–492 (1979).
- ³²V. S. Gerdjikov, M. I. Ivanov, The quadratic bundle of general form and the nonlinear evolution equations, *Bulg. J. Phys.* **10**, 130–143 (1983).
- ³³M. Wadati and K. Sogo, Gauge Transformations in Soliton Theory, *J. Phys. Soc. Japan* **52**, 394–338 (1983).
- ³⁴A. Kundu, Landau-Lifshitz and higher-order nonlinear systems gauge generated from nonlinear Schrödinger type equations, *J. Math. Phys.* **25**, 3433–3438 (1984).
- ³⁵A. Kundu, Exact solutions to higher-order nonlinear equations through gauge transformations, *Physica D*, **25**, 399–406 (1987).
- ³⁶T. Congy, A. M. Kamchatnov, and N. Pavloff, Dispersive hydrodynamics of nonlinear polarization waves in two-component Bose-Einstein condensates, *SciPost Phys.*, **1**, 006 (2016).
- ³⁷S. K. Ivanov, A. M. Kamchatnov, T. Congy, and N. Pavloff, Solution of the Riemann problem for polarization waves in a two-component Bose-Einstein condensate, *Phys. Rev. E* **96**, 062202 (2017).
- ³⁸E. Iacocca, T. J. Silva, and M. A. Hoefer, Breaking of Galilean Invariance in the Hydrodynamic Formulation of Ferromagnetic Thin Films, *Phys. Rev. Lett.* **118**, 017203 (2017).
- ³⁹L. V. Ovsyannikov, Two-layer “shallow water” model, *J. Appl. Mech. Tech. Phys.* **20**, 127–135 (1979).
- ⁴⁰H. Lamb, *Hydrodynamics*, (Cambridge University Press, Cambridge, 1994).
- ⁴¹G. A. El, V. V. Geogjaev, A. V. Gurevich, and A. L. Krylov, Decay of an initial discontinuity in the defocusing NLS hydrodynamics, *Physica D* **87**, 186–192 (1995).
- ⁴²T. Congy, S. K. Ivanov, A. M. Kamchatnov, and N. Pavloff, Evolution of initial discontinuities in the Riemann problem for the Kaup-Boussinesq equation with positive dispersion, *Chaos*, **27**, 083107 (2017).
- ⁴³V. E. Zakharov, The inverse scattering method, *Solitons*, edited by R. K. Bullough and P. J. Caudrey, pp. 243–285 (Springer, Berlin, 1980).
- ⁴⁴M. Ito, Symmetries and conservation laws of a coupled nonlinear wave equation, *Phys. Lett. A* **91**, 335–338 (1982).
- ⁴⁵M. Jaulent, I. Miodek, Nonlinear evolution equations associated with ‘energy-dependent Schrödinger potentials’, *Lett. Math. Phys.*, **1**, 243–259 (1976).
- ⁴⁶R. Zhou, The finite-band solution of the Jaulent-Miodek equation, *J. Math. Phys.*, **38**, 2535–2546 (1997).
- ⁴⁷A. M. Kamchatnov, D. V. Shaykin, Propagation of generalized Korteweg-de Vries solitons along large-scale waves, *Phys. Rev. E* **108**, 054205 (2023).
- ⁴⁸A. M. Kamchatnov, Hamilton theory of dark soliton motion in the nonlinear Schrödinger equation theory, *Theor. Math. Phys.* **219**, 44–55 (2024).
- ⁴⁹A. M. Kamchatnov, D. V. Shaykin, Propagation of dark solitons of DNLS equations along a large-scale background, *Wave Motion*, **129**, 103349 (2024).