Delay differential equations with periodic coefficients: a numerical insight

Anatoli Ivanov and Sergiy Shelyag

Abstract Simple form scalar differential equation with delay and non-linear negative periodic feedback is considered. The existence of slowly oscillating periodic solutions with the same period as the feedback coefficient is shown numerically within the admissible range for the periods.

1 Introduction

This paper presents sample data obtained from the analysis of numerical solutions of scalar simple form delay differential equations (DDEs) with non-linear negative periodic feedback. The aim of this preliminary analysis is to study periodic solutions of these equations and their dynamics in general. A general form of the class of DDEs we aim our research efforts at is given by the following:

$$x'(t) = -\mu(t)x(t) + a(t)f(x(t-\tau)), \quad \tau > 0,$$
(1)

where the non-linearity f(x) is a continuous function and the coefficients a(t) and $\mu(t)$ are continuous *T*-periodic functions. This equation is a natural generalization to the periodic case of the well known autonomous equation

$$x'(t) = -\mu_0 x(t) + a_0 f(x(t-\tau)), \quad \mu_0, a_0 - \text{const},$$
(2)

Anatoli Ivanov

Sergiy Shelyag

Penn State Wilkes-Barre, 44 University Drive, Dallas, PA 18612, USA e-mail: afil@psu.edu

College of Science and Engineering, Flinders University at Tonsley, Adelaide, South Australia 5042, Australia

e-mail: sergiy.shelyag@flinders.edu.au

which is comprehensively studied in numerous publications (see e.g. [3, 8] and further references therein).

Equation (2) is used extensively in a broad range of applications to model various real world phenomena, most notably in biology, physiology, and population dynamics (see e.g. [4, 5, 6, 17, 16] and further references therein). Some of such models indeed achieve a better mathematical description and a more accurate prediction of the processes if certain intrinsic periodic factors in the environment, such as circadian rhythms and seasonal changes, are taken into account. This naturally leads from the initially autonomous DDEs given by Equation (2) to those with periodic coefficients of the form (1).

In this paper we present some numerical outcomes for DDEs (1) in the special case of $\mu(t) \equiv 0$, when it becomes of the form

$$x'(t) = a(t)f(x(t-\tau)), \tag{3}$$

and under the same assumptions of the continuity of f and a and the T-periodicity of the coefficient a(t). The corresponding autonomous equation (2) has the form

$$x'(t) = a_0 f(x(t-\tau)), \quad a_0 - \text{const},$$
 (4)

and is even simpler and well studied from the point of view of the existence of periodic solutions and their properties. The conditions for the existence of periodic solutions of the latter are well known, and can be found in multiple sources, see e.g. [3, 8]. To those general conditions mentioned above the standard additional ones are that the function *f* must satisfy the negative feedback condition $x \cdot f(x) < 0$, $\forall x \neq 0$, be one-sided bounded ($f(x) \le M > 0$, $\forall x \in \mathbf{R}$ (or $f(x) \ge -m < 0$, $\forall x \in \mathbf{R}$)), and the corresponding linearized DDE about the equilibrium x = 0 must be unstable.

The principal objective we are addressing in this work is to numerically identify and possibly find analytically conditions on the coefficient a(t) and the nonlinearity f which would yield the existence of periodic solutions with the same period T as the coefficient a(t). A natural approach to start is to consider the existing periodic solutions for the autonomous DDE (4) and follow their changes as the coefficient a_0 is subject to perturbations. We demonstrate numerically that small periodic perturbations in the coefficient of the form $a(t) = a_0 + \varepsilon a_1(t)$ can lead to induced periodic solutions to DDEs (1) of the same period.

Our subsequent and more extensive numerical simulations have lead to precise analytical results and calculations of explicit periodic solutions of the same period as that in the coefficient a(t); the corresponding results are contained in the recent papers [10, 11, 12].

2 Preliminaries

We consider a partial case of the periodic delay differential equation (1) when $\mu(t) = 0$ and $\tau > 0$, the coefficient $a(t) \ge 0$ is non-negative, and the nonlinearity *f*

Delay differential equations with periodic coefficients: a numerical insight

satisfies the negative feedback assumption $x \cdot f(x) < 0, x \neq 0$. In addition, the delay $\tau > 0$ can always be normalized to the standard one $\tau = 1$. Therefore, the equation (3) becomes

$$x'(t) = a(t)f(x(t-1)).$$
(5)

In the autonomous case of DDE (5), $a(t) = a_0 > 0$ unambiguously ensures the negative feedback condition. In this case, the conditions for the existence of periodic solutions are well known. They can be found in multiple sources, see e.g. monographs [3, 8] and further references therein; here we present a brief summary of associated results for the sake of readers' convenience. We distinguish between a general case of the nonlinearity f, and its partial special case when f is an odd function (symmetric).

2.1 Symmetric (odd) nonlinearity

In this subsection we assume that the nonlinearity f is odd, f(-x) = -f(x), $\forall x \neq 0$, which case we shall also call the "symmetric f". The existence of periodic solutions for the corresponding autonomous DDE (4) are well-known. They are primarily found as the so-called Kaplan-Yorke solutions first described in paper [15].

The following proposition provides conditions for the existence and uniqueness of the symmetric slowly oscillating periodic solution of period 4 ($\tau = 1$). One set of conditions guarantees that the solution is stable with the asymptotic phase; the other implies that the periodic solution is unstable (and therefore not detected numerically). The proposition is a partial case of the more general statements in papers [2, 15].

Proposition 1 (Existence and stability of symmetric periodic solutions).

Suppose that f(x) is odd, f(-x) = -f(x), satisfying the negative feedback condition $x \cdot f(x) < 0$, $\forall x \neq 0$, is strictly decreasing on **R** with $|f'(0)| > \pi/2$ and $\limsup_{x\to\infty} |f(x)/x| = \beta < \pi/2$. Then DDE (4) has at least one slowly oscillating periodic solution satisfying the following symmetry conditions: x(0) = 0, $x(-t) \equiv -x(t)$, $\forall t \in \mathbf{R}$. If, in addition, the derivative f'(x) is monotone on \mathbf{R}_+ , then:

- (i) In case when f'(x) is strictly increasing, delay differential equation (5) has a unique slowly oscillating periodic solution of period 4 which is asymptotically stable with asymptotic phase;
- (ii) In case when f'(x) is strictly decreasing on \mathbf{R}_+ , DDE (5) has a unique unstable slowly oscillating periodic solution with period 4.

2.2 Non-symmetric (general) nonlinearity

In this subsection we consider a general (non-symmetric) case of the nonlinearity f where only the negative feedback assumption is in place. The existence of periodic

solutions for the corresponding autonomous DDE (4) are also well-known. They go back to the classical work [13, 14]; they are also comprehensively described in monographs [3, 8].

Proposition 2 (*Existence of slowly oscillating periodic solutions*). Suppose that f satisfies the negative feedback assumption, $x \cdot f(x) < 0, x \neq 0$, differentiable at zero with $f'(0) > \pi/2$, and is one-sided bounded, $f(x) \le M > 0$, $\forall x \in \mathbf{R}$ (or $f(x) \ge -m < 0$, $\forall x \in \mathbf{R}$). Then delay differential equation (5) has a non-trivial slowly oscillating periodic solution $x_0(t)$, such that

$$x_0(t) > 0 \ \forall t \in (0, t_1), \ x_0(t) < 0 \ \forall t \in (t_1, t_2), \ x_0(t+T) \equiv x_0(t), \ T = t_1 + t_2.$$

Note that unlike as in Proposition 1 the uniqueness of the periodic solution cannot be derived in this more general case. In fact there are typical examples when DDE (4) has multiple periodic solutions [20, 1].

3 Results

The numerical solutions for the delay differential equation (5) have been calculated using the algorithm described in [1]. The algorithm has demonstrated fourth-order global convergence for non-linear systems of delay differential equations with discontinuous solutions. The periods of the solutions have been measured and their stability confirmed based on persistence of the distances between the consecutive zeros of the obtained solutions with the error less than 0.1%.

The numerical results in this section are given for the coefficient a(t) > 0 of the form $a(t) = 1 + \varepsilon b_0(t)$, where $b_0(t)$ is a fixed *T*-periodic function, and $\varepsilon \in (0, 1)$ is a constant parameter.

We have considered both symmetric and asymmetric functions f. For the symmetric case we have computed numerical solutions for the following two non-linearities:

$$f_1(x) = -C \tanh(cx), \ C > 0, c > 0 \tag{6}$$

and

$$f_2(x) = -\frac{px}{1+x^{2n}}, \ p > 0, n \ge 1, n \in \mathbb{N}.$$
(7)

The numerical data in Figure 1 is shown for the symmetric case with the first non-linearity f_1 when C = 1 and c = 5. According to Proposition 1 for these parametric values DDE (4) has a unique asymptotically stable slowly oscillating periodic solution with period T = 4. This solution clearly exists for small perturbation amplitudes. With increasing perturbation amplitude (left panel), the period of the solution grows until it becomes equal to the period of the perturbation. The effect of varying the perturbation period at a given amplitude of the perturbation is shown in the right panel of Figure 1. For a small amplitude a = 0.5, the range of perturbation periods, for which the period of the solution is equal to the perturbation period, is also nar-

row in agreement with the left panel of Figure 1. Notably, this range is asymmetric. Similar results have been obtained for the non-linearity given by Equation (7).



Fig. 1 Dependence of the detected period in solution of Equation (5) with the symmetric smooth f given by Equation (6) on the amplitude of the perturbation ε for three given perturbation periods (left panel) and on the perturbation period ω for a = 0.5. The base period of the autonomous equation solution is T = 4.

For the asymmetric case, the numerical simulations were run for the following two non-linearities:

$$f_3(x) = A[1 - \exp\{ax\}], \quad A > 0, \ a > 0,$$
(8)

and

$$f_4(x) = \begin{cases} -B \arctan(bx) & \text{if } x < 0, \\ -B \ln(1+bx) & \text{if } x \ge 0, \end{cases}$$
(9)

where B > 0 and b > 0. The latter f_4 , given by expression (9), was used to produce the data shown in Figure 2. The parametric values used are B = 2 and b = 30; the period of the corresponding autonomous periodic solution is T = 5.07. In the first case of the non-symmetric f_3 (Equation 8) analogous computations were performed for A = 2 and a = 1 with similar outcomes for the those in the case of f_4 (they are not depicted by graphs here). The existence of periodic solutions in both non-symmetric cases follows from Proposition 2.

Similar to the symmetric case, as shown in the left panel of Figure 2, there is an interval of perturbation amplitudes ε , for which the period of solution is equal to the period of the perturbation. As shown in the right panel of Figure 2, there is an interval of perturbation periods for a given perturbation amplitude, for which the period of solution is also equal to the period of perturbation.

4 Discussion

In this short paper, we provided a preliminary numerical analysis of a delay differential equation with periodic feedback coefficient. We have demonstrated that



Fig. 2 Dependence of the detected period in solution of Equation (5) with f given by Equation 9 on the amplitude of the perturbation ε for three given perturbation periods (left panel) and on the period ω of the perturbation with the constant amplitude a = 0.1 (right panel). The base period of the autonomous equation solution is T = 5.07.

stable periodic solutions with the period of the coefficient exist in a wide range of perturbation amplitudes and periods. The next steps will be to extend this study to a broader class of delay-differential equations (e.g. Equation 1) and to provide an analytical insight into the periodic behaviour of the studied equations.

Acknowledgements The authors thank the mathematical research institute MATRIX in Australia where part of this research was performed. Its final version resulted from collaborative activities of the authors during the workshop "Delay Differential Equations and Their Applications" (https://www.matrix-inst.org.au/events/delay-differential-equations-and-their-applications/) held in December 2023. The authors are also grateful for the financial support provided for these research activities by Simons Foundation (USA), Flinders University (Australia), and the Pennsylvania State University (USA). A.I.'s research was also supported in part by the Alexander von Humboldt Stiftung (Germany) during his visit to Justus-Liebig-Universität, Giessen, in June-August 2023.

References

- Angelova, M., Belyakov, G., Ivanov, A. and Shelyag, S. (2021): Global stability and periodicity in a glucose-insulin regulation model with a single delay. *Commun. Nonlin. Sci.* 95, 105659.
- 2. Chow, S.-N. and Walther, H.-O. (1988): Characteristic multipliers and stability of periodic solutions of $\dot{x} = g(x(t-1))$. *Trans. Amer. Math. Soc.* 307, 127–142.
- Diekmann, O., van Gils, S., Verdyn Lunel, S., Walther, H.-O.: Delay Equations: Complex, Functional, and Nonlinear Analysis. Springer-Verlag, Ser.: Applied Mathematical Sciences, Vol. 110 (1995).
- Faria, T. (2017): Periodic solutions for a non-monotone family of delayed differential equations with applications to Nicholson systems. J. Differential Equations, 263, no. 1, 509–533.
- Friedman, H.I. and Wu, J (1992): Periodic solutions of single-species models with periodic delay. SIAM J. Math. Anal. 23, no. 3, 689–701.
- Glass, L., Mackey, M.C.: From Clocks to Chaos (1988). The Rhythms of Life. Princeton University Press.
- Gurney, W.S.C., Blythe, S.P., Nisbet, R.M. (1980): Nicholson's blowflies revisited. *Nature*, 287, 17–21.

Delay differential equations with periodic coefficients: a numerical insight

- Hale, J.K., Verduyn Lunel, S.M. (1993): Introduction to Functional Differential Equations. Springer Applied Mathematical Sciences, vol. 99.
- Ivanov, A.F., Sharkovsky, A.N. (1991): Oscillations in singularly perturbed delay equations. Dynamics Reported (New Series), Springer Verlag, vol. 1, 165–224.
- Ivanov, A., Shelyag, S. (2023): Stable periodic solutions in scalar periodic differential delay equations. Archivum Mathematicum (Brno), 59 (1), 69–76.
- Ivanov, A. and Shelyag, S (2023): Explicit periodic solutions in a delay differential equation. Proceedings of 6th international conference AMMCS-2023. Preprint, arXiv 2402.08197, December 2023, 7 pp. (submitted)
- Ivanov, A. and Shelyag, S (2024): Periodic Solutions in a Simple Delay Differential Equation. Preprint, March 2024, 13 pp. (submitted)
- 13. Jones, G.S. (1962): The existence of periodic solutions of $f'(x) = -\alpha f(x-1)[1+f(x)]$. J. Math. Anal. Appl., 5, 435–450.
- 14. Jones, G.S. (1964): Periodic motions in Banach space and applications to functional differential equations. *Contributions to Differential Equations*, 3, 75–106.
- Kaplan, J. and Yorke, J. (1974):Ordinary differential equations which yield periodic solutions of functional differential equations. J. Math. Anal. Appl., 48, 317–324.
- Kuang, Y. (1993): Delay Differential Equations with Applications in Population Dynamics. Academic Press Inc. Series: Mathematics in Science and Engineering, Vol. 191, 398 pp.
- Li, Y. and Kuang, Y. (2001): Periodic solutions in periodic state-dependent delay equations and population models. *Proceedings of the American Mathematical Society*, 130, 1345–1353.
- Mackey, M.C., Glass, L. (1977): Oscillation and chaos in physiological control systems. *Science*, 197, 287–289.
- 19. Nussbaum, R.D. (1977): The range of periods of periodic solutions of $x'(t) = -\alpha f(x(t-1))$. *J. Math. Anal. Appl.*, 58, no. 2, 280–292.
- 20. Nussbaum, R.D. (1979): Uniqueness and nonuniqueness of periodic solutions of x'(t) = g(x(t-1)). *J. Differential Equations*, 34, 25–54.
- Perez, J.F., Malta, C.P., Coutinho, F.A.B. (1978): Qualitative analysis of oscillations in isolated populations of flies. J. Theoret. Biol., 71, 505–514.
- 22. Peters, H. (1980): Comportement chaotique d'une equation differentielle retardee. *C. R. Acad. Sci. Paris Sér. A-B*, 290, no. 24, A1119—A1122. (in French)