Study of the neutrino-oxygen cross sections of the charged-current reaction $^{16}{\rm O}(\bar{\nu}_e,e^+)^{16}{\rm N}(0$ $\text{MeV},\,2^{-})$ and the neutral-current reaction $^{16}{\rm O}(\nu, \nu')^{16}{\rm O}(12.97/12.53\,\,{\rm MeV},\,2^{-}),\,{\rm producing}$ high-energy γ rays

Makoto SAKUDA^{1,*}, Toshio SUZUKI^{2,3*}, Ken'ichiro NAKAZATO^{4,*}, and Hideyuki SUZUKI⁵

¹Physics Department, Okayama University, Okayama 700-8530, Japan 2 Department of Physics, College of Humanities and Sciences, Nihon University, Tokyo 156-8550, Japan

³NAT Research Center, NAT Corporation, 3129-45 Hibara Muramatsu, Tokai-mura, Naka-gun, Ibaraki 319-1112, Japan

⁴Faculty of Arts and Science, Kyushu University, Fukuoka 819-0395, Japan ⁵Department of Physics, Faculty of Science and Technology, Tokyo University of Science, Chiba 278-8510, Japan *

[∗]E-mail: sakuda-m@okayama-u.ac.jp, suzuki.toshio@nihon-u.ac.jp, nakazato@artsci.kyushu-u.ac.jp

> . In the previous work, we discussed the cross section and the detection of 4.4-MeV γ rays produced in the neutrino neutral-current (NC) reaction ${}^{16}O(\nu, \nu'){}^{16}O(12.97 \text{ MeV}$ and 12.53 MeV, 2[−]) in a water Cherenkov detector at the low energy below 100 MeV. In this report, we further investigated both the charged-current (CC) reaction ${}^{16}O(\bar{\nu}_e, e^+)^{16}N(0)$ MeV, 2⁻) and the NC reaction¹⁶ $O(\nu, \nu')$ ¹⁶ $O(12.97 \text{ MeV}$ and 12.53 MeV, 2⁻), producing high-energy γ rays, in which the more solid identification of the reactions can be applied via the coincidence method. .

Subject Index C43, D02, D03, D21, F22

1. Introduction

The 12.97-MeV and 12.53-MeV states are the first strong 2^- excited states of ¹⁶O just above the proton separation energy (12.1 MeV). The 12.97-MeV state, which is nearly an isospin $T=1$ state, is one of the dominant multipoles in the neutrino-oxygen interactions at low energy below 100 MeV. The electromagnetic form factors $F^2(q)$ of these states were measured in (e, e') reactions in 1960 [1-[4\]](#page-14-1). No new measurements of those states in (e, e') reactions have been performed since then. Donnelly and Walecka [\[5–](#page-14-2)[7\]](#page-14-3) calculated the neutrino-16O cross sections at $E_x=12-20$ MeV precisely with accuracy of 15-20% after they analysed the data of ¹⁶O(e, e')¹⁶O(E_x =12-20 MeV) scattering and semi-leptonic weak interactions (muon capture and β decay) and evaluated the reduction factors $(a/\xi=0.6-0.7)$ to the transition amplitudes

of their model. This reduction in transition amplitudes of a calculation model (or in the coupling constant) is sometimes called a quenching factor. Haxton [\[8\]](#page-14-4) calculated the cross sections of the charged-current (CC) neutrino-oxygen ¹⁶O(ν_e , e⁻)¹⁶F and ¹⁶O($\bar{\nu}_e$, e⁺)¹⁶N reactions, using the quenching factors for negative parity states, which were evaluated in Ref. [\[5–](#page-14-2)[7\]](#page-14-3). He further examined the CC cross sections to the bound states $(2^-, 0^-, 3^-, 0^+)$ 1^-) of ¹⁶N, which are followed by the β^- decay to the ground state or the excited state (6.13 MeV) of ¹⁶O. The total energy given by β^- and γ ray (6.13 MeV) was estimated to be about 8 MeV. He concluded that since the CC cross section to the bound states of ¹⁶N never exceeds 1% of the dominant inverse beta decay (IBD) reaction in supernova neutrino bursts, the extra delayed signal of 8 MeV scattered over the decay time $(T_{1/2}=7.13 \text{ sec})$ gives only negligible effect on the event timing, which is determined by the dominant IBD events.

At the time of these analysis, the isospin mixing of the two 2[−] states at 12.53 MeV and 12.97 MeV was not known and was not considered. There have been several reports on the isospin mixing between the 12.97 MeV ($T = 1$) and 12.53 MeV ($T = 0$) states previously [\[1,](#page-14-0) [9](#page-14-5)[–12\]](#page-15-0). These physical two 2⁻ states (the higher energy state $|U\rangle$ and the lower energy state $|D\rangle$ are written in terms of the pure isospin states as,

$$
|U\rangle = \sqrt{1 - \beta^2} |U, T = 1\rangle - \beta |U, T = 0\rangle,
$$

$$
|D\rangle = \sqrt{1 - \beta^2} |D, T = 0\rangle + \beta |D, T = 1\rangle,
$$
 (1)

where β is the isospin mixing parameter. A well-known example of the isospin mixing is that between the two excited states of ¹²C at 12.71 MeV $(1^+, T = 0)$ and 15.11 MeV $(1^+, T = 0)$ 1) [\[13–](#page-15-1)[15\]](#page-15-2).

In our previous work [\[16\]](#page-15-3), we followed the analysis by Donnelly and Walecka, evaluating both the quenching factor $f_s = g_s^{\text{eff}}/g_s$ of the spin g factor and the isospin-mixing parameter β of the two 2⁻ states to be $f_s = 0.65 \pm 0.05$ and $\beta = 0.25 \pm 0.05$, respectively, and also determining the quenching factor $f_A = g_A^{\text{eff}}/g_A$ of the axial-vector coupling constant to be $f_A = 0.68 \pm 0.05$. Then, we discussed the cross section of 4.4-MeV γ ray production in the neutrino neutral-current (NC) reaction ${}^{16}O(\nu, \nu'){}^{16}O(12.97 \text{ MeV}, 2^-)$ in a water Cherenkov detector at the low energy below 100 MeV.

The Super-Kamiokande (SK) experiment summarizes the following three detection channels from supernova (SN) neutrino bursts as described in Ref. [\[17,](#page-15-4) [18\]](#page-15-5): (1) the IBD reaction $p(\bar{\nu}_e, e^+)n$, (2) the neutrino-electron elastic scattering, and (3) the CC reactions, ¹⁶O(ν_e, e^{-})¹⁶F and ¹⁶O($\bar{\nu}_e, e^{+}$)¹⁶N. The first IBD reaction is the main interaction channel, responsible for about 90% of the reactions in water. The second one is a subdominant channel, useful for determining the direction of the SN. The third CC neutrino-oxygen interactions are also subdominant ones. Their cross sections were calculated initially by the authors of Ref. [\[8,](#page-14-4) [19\]](#page-15-6) and recently in Ref. [\[20\]](#page-15-7), and the electron spectra of the CC reactions were discussed by the authors of Ref. [\[17,](#page-15-4) [21,](#page-15-8) [22\]](#page-15-9).

The JUNO experiment [\[23\]](#page-15-10), a Liquid Scintillator detector of 20-kton fiducial mass, uses the 15.11-MeV γ -ray emission of the NC reaction ${}^{12}C(\nu, \nu'){}^{12}C(15.11 \text{ MeV}, 1^+),$ and the CC reactions, ¹²C(ν_e, e^{-})¹²N(g.s., 1⁺) and ¹²C($\bar{\nu}_e, e^{+}$)¹²B(g.s., 1⁺) [\[24\]](#page-15-11), as the main detection channels for the analysis of the SN neutrino bursts, in addition to the IBD reaction, elastic ν –p scattering and elastic ν –e scattering [\[25–](#page-15-12)[27\]](#page-15-13). We denote the ground state as g.s. hereafter.

This study on the CC/NC neutrino-oxygen reactions using 2^{-} (T = 1) states of ¹⁶N and ¹⁶O is motivated by the well-studied CC/NC neutrino-carbon reactions using 1^+ (T=1)

ground states of ¹²B, ¹²C and ¹²N, where both CC and NC reactions of neutrino-¹²C are already measured by the low-energy neutrino experiments [\[28–](#page-15-14)[36\]](#page-15-15). We investigate both the CC $^{16}O(\bar{\nu}_e, e^+)^{16}N(g.s., 2^-)$ and its β decay to ^{16}O . Furthermore, we investigate the NC reaction cross sections from the two 2^- states (12.97 MeV and 12.53 MeV) of ¹⁶O, producing high-energy γ rays. We discuss a possible coincidence method to identify these CC and NC reactions unambiguously, which can be used not only in the SK experiment but also in the future Hyper-K experiment [\[22\]](#page-15-9).

The three states, ¹⁶N(g.s., 2⁻), ¹⁶O(12.97 MeV, 2⁻) and ¹⁶F(0.42 MeV, 2⁻), form a $T = 1$ triplet $(T_z = -1, 0, 1)$. The energy levels of ¹⁶N, ¹⁶O and ¹⁶F near their ground states are shown in Fig. [1.](#page-3-0) Just above the ground state ¹⁶N(g.s., 2⁻), there are also T=1 bound states at 0.120 MeV (0^-) , 0.298 MeV (3^-) and 0.397 MeV (1^-) . They all decay electromagnetically to ${}^{16}N(g.s., 2^-)$, emitting a small γ ray. We call these bound states including the ground state (2^-) as the g.s. group of ¹⁶N in the present report. There are no bound states in ¹⁶F(T=1).

2. Charged-current reaction $^{16}{\rm O}(\bar{\nu}_e,e^+)^{16}{\rm N}$ (g.s. group)

The calculations of the electron/positron spectra from ¹⁶ $O(\nu_e, e^{-})^{16}F$ and ¹⁶ $O(\bar{\nu}_e, e^{+})^{16}N$ reactions were fully described in Ref. [\[21\]](#page-15-8) and the implications of the 18 O mixture in water on SN neutrino events were discussed in Ref [\[37\]](#page-15-16). In this section, we discuss on the CC reaction ${}^{16}O(\bar{\nu}_e, e^+){}^{16}N(g.s.$ group), where the g.s. group consists of the bound states at 0 MeV (2⁻), 0.120 MeV (0⁻), 0.298 MeV (3⁻) and 0.397 MeV (1⁻). The bound states (0⁻, 3^- , 1^-) decay electromagnetically to the ground state (2^-) , emitting a small γ ray. All of them are followed by the β decay from ¹⁶N(g.s.) to ¹⁶O. We discuss the g.s. group together, since the four bound states of the g.s. group in the CC reactions ${}^{16}O(\bar{\nu}_e, e^+)^{16}N(g.s.$ group) cannot be distinguished in a water-Cherenkov detector unless a small γ ray (0.120 MeV, 0.298 MeV, 0.397 MeV) can be identified. We describe some unique features of these CC reaction as compared to other CC reactions. The formula of the cross section calculation for the CC neutrino-oxygen reactions are given in Eq.(1) of Ref. [\[21\]](#page-15-8) and we calculated the cross section of ¹⁶O($\bar{\nu}_e$, e^+)¹⁶N(g.s. group) using the quenching factors $f_A=0.68\pm0.05$ and $f_s=0.65\pm0.05$, which were evaluated in the previous work [\[16,](#page-15-3) [37\]](#page-15-16).

First, the reaction $\bar{\nu}_e + {}^{16}O \rightarrow e^+ + {}^{16}N(g.s.$ group) can be uniquely identified by the coincidence between a prompt positron from the primary reaction and a 6.13-MeV γ ray (and partly 7.12-MeV and 8.87-MeV γ rays) from the subsequent β^- decay of $^{16}N(g.s.)$, $^{16}N(g.s.) \rightarrow ^{16}O(E_x > 0) + e^- + \bar{\nu}_e$, both of which are produced at the same interaction point during the time interval of the β decay. The detailed parameters of the β decay of $^{16}N(g.s.)$ to ^{16}O [\[38\]](#page-15-17) are summarized in Table 1 and the schematic diagram of the decay is also shown in dashed lines in Fig. [1.](#page-3-0) This coincidence method with the constraint on the same vertex position during on the decay interval will reduce the accidental background significantly. The identification of this reaction by applying the coincidence will reject other CC reactions ¹⁶ $O(\nu_e, e^{-})^{16}$ F and ¹⁶ $O(\bar{\nu}_e, e^{+})^{16}$ N $(E_x > 1$ MeV) [\[21\]](#page-15-8), which have larger cross sections than this reaction and have no delayed signals. Though some of these other CC reactions may be accompanied by the prompt γ rays above 5 MeV which are emitted from the strong-decay products as $15N^*$ or $15O^*$ within a microsecond, they can be further removed if the second signals from the first one microsecond in the coincidence are excluded out of from the long decay time $(T_{1/2}=7.13 \text{ sec}).$

Fig. 1 Energy levels of ¹⁶N, ¹⁶O and ¹⁶F near the ground state with isospin $T=1$ [\[38\]](#page-15-17).

Secondly, the visible energy T_{e^+} of the positron from the primary reaction of this channel (g.s. group) can be used to determine the incident neutrino energy as $E_{\bar{\nu}_e} = T_{e^+} + 11.44$ MeV above the threshold energy ($E_{th}=11.44$ MeV). A small γ ray (0.12 MeV, 0.298 MeV, 0.397 MeV) of the g.s. group is negligible as compared to 11.44 MeV. The electron or positron signal from other CC reactions cannot give the incident neutrino energy without knowing the excited states (E_x) of ¹⁶F and ¹⁶N. This CC reaction from ¹⁶O to ¹⁶N(g.s. group) has the lowest energy threshold among the neutrino-oxygen reaction, except for the CC ¹⁸O(ν_e, e^{-})¹⁸F reaction (E_{th} =1.66 MeV) [\[37\]](#page-15-16).

The cross section of ${}^{16}O(\bar{\nu}_e, e^+)^{16}N(g.s.$ group) is shown as a function of neutrino energy in Fig.2 and also given in Table 2. The cross section of the CC reaction from $^{16}O(g.s.)$ to $16N(g.s., 2⁻)$ is the largest among the CC reaction to the g.s. group, that to the 1⁻ state

is about 3/5 of that to the 2^- state below 50 MeV and that to the 0^- state is about 15% of that to the 2[−] state between 12 and 20 MeV. This feature is qualitatively explained by the strength of the transition $(S=1 \text{ and } L=1)$ proportional to $(2J+1)$. Above 50 MeV, the cross section is contributed to by the transition (S=1 and L=3) from ¹⁶O(g.s.) to the 2⁻ and 3 [−] states. The cross section to the 2[−] state becomes even larger and that to the 3[−] state becomes significant, about 10% of that to the 2[−] state, at higher energy than 100 MeV.

The cross section of the CC reaction ${}^{16}O(\bar{\nu}_e, e^+){}^{16}N(g.s.$ group) was first calculated by Haxton [\[8\]](#page-14-4). We find that our calculation of the CC cross section for ${}^{16}O(\bar{\nu}_e,e^+)^{16}N(g.s.$ group) is larger by about 40% than his calculation. We note that our evaluations of the quenching factors for these bound states, $f_s = 0.65 \pm 0.05$ and $f_A = 0.68 \pm 0.05$, were validated by the transverse form factor $F_T^2(q)$ of the (e, e') cross section near 13 MeV $(2^-, 1^-, 3^-)$ [\[1,](#page-14-0) [2,](#page-14-6) [4\]](#page-14-1), the rate of the partial muon capture (μ^-,ν_μ) from the 1s orbit on ¹⁶O(g.s., 0⁺) to the bound states $(2^-, 0^-, 3^-, 1^-)$ of ¹⁶N and the total muon capture rate from ¹⁶O to ¹⁶N(g.s., 2[−]) [\[16\]](#page-15-3).

Below 30 MeV, this cross section of ${}^{16}O(\bar{\nu}_e, e^+){}^{16}N(g.s.$ group) is dominant among all CC reactions. There are several excited states $(2^-, 1^-)$ of ${}^{16}N(E_x=3-25 \text{ MeV})$ with significant CC cross sections and they all decay hadronically to $n+$ ¹⁵N [\[21\]](#page-15-8). Above 30 MeV, the cross section to the g.s. group becomes smaller than the sum of other CC cross sections by an order of magnitude. However, we note again that the neutrino energy can be reconstructed, only when the electron/positron energy of other CC reactions can be measured and the excited states (E_x) of the nucleus in the reaction, either ¹⁶N or ¹⁶F, are measured for each event. In addition, those excited states (E_x) will decay hadronically and sometimes emit γ rays from 5 to 9 MeV, which will diffuse the primary electron/positron signal. On the other hand, the CC reaction ${}^{16}O(\bar{\nu}_e, e^+){}^{16}N(g.s.$ group) can be unambiguously identified and the neutrino energy can be reconstructed, if the delayed coincidence method can be used. This is a unique feature.

While the 6.13-MeV and 7.12-MeV states decay electromagnetically via $E3/E1$ transition to the ground state, producing a single 6.13-MeV and 7.12-MeV γ ray, the 8.87-MeV state (2⁻) decays mostly through cascade to the ground state, producing two γ rays $(2.74 \text{ MeV} + 6.13 \text{ MeV}, \text{ or } 1.75 \text{ MeV} + 7.12 \text{ MeV})$ and partly a single 8.87-MeV γ ray. After the β^- decay and the electromagnetic transitions, the branching ratios of emitting $γ$ rays becomes Br($E_γ=6.13$ MeV)=0.662 ± 0.006, Br(7.12 MeV)=0.048±0.004 and Br(8.87 MeV)=0.0106±0.0007 [\[38\]](#page-15-17) and the sum of them is $Br(E_{\gamma} \ge 6.13 \text{ MeV})=0.720 \pm 0.007$. The probability of no γ ray is 0.280 ± 0.004 . We note that the visible energy of the delayed signal is $E_d = E_{\gamma} + T_{\beta} = 10.42 \text{ MeV} - T_{\bar{\nu}_e}$ for all decay modes, including the decay of ¹⁶N(g.s) to ¹⁶O(g.s), where $T_{\bar{\nu}_e}$ is the neutrino energy from the β^- decay, ${}^{16}N(g.s.) \rightarrow {}^{16}O(E_x > 1)$ $0 + e^- + \bar{\nu}_e$. The delayed signal E_d is further contributed to by the kinetic energy T_β from the β decay by 1-2 MeV as shown in Table 1 and this will make the detection efficiency larger.

The β decay of ¹⁶N(g.s.) goes to ¹⁶O(g.s.) with the branching ratio 28.0%, producing an electron with $T_{\beta \text{ max}} = 10.42 \text{ MeV}$ and no γ ray. Even for this decay mode, the delayed coincidence between the primary positron and the delayed coincidence signal may be possible for the case of $T_{\beta} > 5$ MeV. The kinetic energy spectrum of this β decay mode is shown in Fig.3, where Q-value is equal to 10.42 MeV. About 49% of the electrons have $T_{\beta} > 5$ MeV and they can be detected. This will add a probability of about 14% (= 0.28 · 0.49) to that of the delayed coincidence signal producing γ rays with $E_{\gamma} \geq 6.13$ MeV (72%) and the total probability of the delayed coincidence signal with visible energy greater than 5 MeV, from the β decay of $^{16}N(g.s.)$, is estimated to be about 86%.

The SNO experiment and SK experiment use γ rays of 6.13 MeV and 7.12 MeV from ¹⁶N(g.s., 2⁻) β decay for a PMT calibration [\[39,](#page-15-18) [40\]](#page-15-19). The SK experiment also uses a 6.13-MeV γ ray to measure the NC neutrino-oxygen quasielastic interaction [\[41–](#page-15-20)[44\]](#page-15-21), which are consistent with the calculation [\[45\]](#page-15-22). Thus, it is clear that a 6.13-MeV γ ray can be observed in a water Cherenkov detector.

We comment on the unique feature of this CC reaction channel to $^{16}N(g.s.$ group). A detector such as SK-Gd [\[48–](#page-15-23)[50\]](#page-15-24) can measure the neutrino energy of the IBD reaction unambiguously by the neutron tagging and reconstruct the incident neutrino spectrum $F(E_{\bar{\nu}_e})$, using its well-known IBD cross section [\[25,](#page-15-12) [26\]](#page-15-25) and the relation $E_{\bar{\nu}_e} = T_{e^+} + 1.80$ MeV. Then, we can measure the CC cross section $\sigma(E_{\bar{\nu}_e})$, using the measured neutrino spectrum $F(E_{\bar{\nu}_e})$, since we can measure $F(E_{\bar{\nu}_e}) \cdot \sigma(E_{\bar{\nu}_e})$ from the measurement of the primary electron spectrum T_{e^+} , using the relation $E_{\bar{\nu}_e} = T_{e^+} + 11.44 \text{ MeV}$, where $\sigma(E_{\bar{\nu}_e})$ is the CC cross section to ${}^{16}N(g.s.$ group). We can compare the measured cross section with our calculated one and confirm/improve the calculation. The measurement of this CC reaction will also validate the calculation of the NC cross sections which we describe in the next section, since the calculations of these CC and NC reactions are related by the CVC hypothesis [\[16,](#page-15-3) [46,](#page-15-26) [47\]](#page-15-27). We also note that the delayed coincidence method to this reaction can be applied in the Hyper-K detector, even without the neutron tagging method.

Table 1 Parameters of the β^- decay of ¹⁶N(g.s.), ¹⁶N(g.s.) \rightarrow ¹⁶O(E_x) + e^- + $\bar{\nu}_e$ [\[38\]](#page-15-17). The half life of the decay is $T_{1/2} = 7.13 \pm 0.02$ sec. While the 6.13-MeV and 7.12-MeV states decay to the ground state, producing a single 6.13-MeV and 7.12-MeV γ ray, the 8.87-MeV state (2^-) decays through cascade to the ground state, producing mostly two γ rays (2.74) MeV + 6.13 MeV, or 1.75 MeV + 7.12 MeV) and partly a single 8.87-MeV γ ray.

β^- decay of ¹⁶ N(g.s.) to	Branching ratio	E_{γ}	Range of T_{β}
E_x of ¹⁶ O	$(\%)$	(MeV)	(MeV)
8.87 MeV	1.06 ± 0.07	$8.87\;\rm{MeV}$	$0 < T_{\beta} < 1.55 \text{ MeV}$
7.12 MeV	4.8 ± 0.4	$7.12~\mathrm{MeV}$	$0 < T_{\beta} < 3.30 \; \mathrm{MeV}$
6.13 MeV	66.2 ± 0.6	6.13 MeV	$0 < T_{\beta} < 4.29 \text{ MeV}$
0 MeV	28.0 ± 0.4	0.0	$0 < T_{\beta} < 10.42$ MeV

3. Neutral-current reaction $^{16}{\rm O}(\nu,\nu')^{16}{\rm O}(12.53\ {\rm MeV}$ and $12.97\ {\rm MeV},\ 2^-)$ and the branching ratios of the two 2^- states producing γ rays

We briefly review a feature of the NC reaction from the U and D states, which produces a 4.4-MeV γ ray from the α decay of these states [\[16\]](#page-15-3). Next, we discuss a high-energy γ ray emission from the electromagnetic decay of these two states. The latter cross sections for high-energy γ rays are small, but the signature of the events is so distinct with highenergy visible energy above 10 MeV that those events can be clearly identified. In these calculations, we use both the quenching factors of the spin g factor $(f_s = 0.65)$ and of the axial-vector coupling constant ($f_A = 0.68$), and the isospin-mixing parameter of the two 2[−]

Fig. 2 The cross sections of the CC reactions ${}^{16}O(\bar{\nu}_e, e^+)^{16}N(g.s., 2^-)$ (black dashed line), ¹⁶N(1⁻) (black dash-dotted line), and ¹⁶N(g.s. group) (black solid line); the NC cross sections of the 4.4-MeV γ ray from U and D states, $\sigma_{NC,\gamma}^U$ (red dashed line) and $\sigma_{NC,\gamma}^D$ (red dash-dotted line), and the sum of them $\sigma_{NC,\gamma}^{tot}$ (red solid line); the NC cross sections of the high-energy γ rays (12.97 MeV and 12.53 MeV) via electromagnetic decay of the U and D states, $\sigma_{\text{NC},2\gamma}^{U}$ (blue dashed line), $\sigma_{\text{NC},2\gamma}^{D}$ (blue dash-dotted line), and the sum of them $\sigma_{\text{NC},2\gamma}^{tot}$ (blue solid line) as a function of the neutrino energy. The IBD cross section is shown in black dotted lines for comparison.

states ($\beta = 0.25$). We summarize the decay properties of the two states in Table 3, which we use in the present paper as in the previous paper [\[16\]](#page-15-3). We also illustrate the NC reactions $^{16}O(\nu, \nu')$ ¹⁶O(12.53 MeV and 12.97 MeV, 2⁻) in Fig.4 which are relevant in this section.

First, we review the 4.4-MeV γ ray from the α decay of the two states. The α decay of the 2⁻ states to $\alpha +^{12}C(0 \text{ MeV}, 0^+)$ is forbidden by the angular momentum conservation and the α decay of the 2⁻ states to $\alpha + {}^{12}C(4.4 \text{ MeV}, 2^+)$ is allowed through the T=0 component. The three experiments reported the α -decay branching ratio Br($U \rightarrow \alpha +^{12}C(4.4 \text{ MeV})$)= Γ_{α_1}/Γ , which we denote as $Br^U(\alpha_1)$: Leavitt *et al.* [\[10\]](#page-14-7), Zijderhand and van der Leun [\[11\]](#page-15-28), and Charity et al. [\[12\]](#page-15-0) reported $Br^{U}(\alpha_1)$ to be 0.37 \pm 0.06, 0.22 \pm 0.04, and 0.46 \pm 0.08, respectively. We took a simple mean of the three values [\[10–](#page-14-7)[12\]](#page-15-0) and used this mean value, $Br^U(\alpha_1)=0.35$, to evaluate the 4.4-MeV γ -ray production cross section in the previous paper [\[16\]](#page-15-3) as well as in the present paper. We denote the NC cross section of the U and D states as σ_{NC}^{U} and $\sigma_{\rm NC}^D$, respectively, and the sum of them as $\sigma_{\rm NC}^{\rm tot} = \sigma_{\rm NC}^U + \sigma_{\rm NC}^D$. We also denote the 4.4-MeV γ-ray production cross section of the U and D states as $\sigma_{NC,\gamma}^U$ and $\sigma_{NC,\gamma}^D$, respectively, and the sum of them as $\sigma_{NC,\gamma}^{tot} = \sigma_{NC,\gamma}^U + \sigma_{NC,\gamma}^D$. We note that the NC cross sections σ_{NC}^U and $\sigma_{\rm NC}^D$ are calculated for an average of one neutrino flavor and its anti-neutrino flavor. In the previous work [\[16\]](#page-15-3), only the figures for the U state, σ_{NC}^U and $\sigma_{NC,\gamma}^U = \sigma_{NC}^U \cdot Br^U(\alpha_1)$, were

E_{ν} (MeV)	2^{-}	1^{-}	0^{-}	3^-	Sum $(2^-, 1^-, 0^-, 3^-)$
12	$1.59E - 04$	$4.04E - 05$	$2.51E - 05$	$0.0\,$	$2.24E - 04$
14	$2.10E - 03$	$1.46E - 03$	$3.03E - 04$	0.0	$3.86E - 03$
16	$7.79E - 03$	$5.74E - 03$	$8.69E - 04$	0.0	$1.44E - 02$
18	$1.98E - 02$	$1.44E - 02$	$1.73E - 03$	$1.20E - 06$	$3.60E - 02$
20	$4.13E - 02$	$2.92E - 02$	$2.88E - 03$	$4.24E - 06$	$7.34E - 02$
22	$7.57E - 02$	$5.20E - 02$	$4.32E - 03$	$1.23E - 05$	$1.32E - 01$
24	$1.27E - 01$	$8.46E - 02$	$6.07E - 03$	$3.11E - 05$	$2.18E - 01$
26	$2.00E - 01$	$1.29E - 01$	$8.13E - 03$	$7.07E - 05$	$3.37E - 01$
28	$2.97E - 01$	$1.87E - 01$	$1.05E - 02$	$1.48E - 04$	$4.95E - 01$
30	$4.25E - 01$	$2.60E - 01$	$1.32E - 02$	$2.88E - 04$	$6.99E - 01$
32	$5.86E - 01$	$3.51E - 01$	$1.62E - 02$	$5.29E - 04$	$9.54E - 01$
34	$7.86E - 01$	$4.60E - 01$	$1.95E - 02$	$9.27E - 04$	$1.27E + 00$
36	$1.03E + 00$	$5.88E - 01$	$2.32E - 02$	$1.56E - 03$	$1.64E + 00$
38	$1.31E + 00$	$7.36E - 01$	$2.72E - 02$	$2.52E - 03$	$2.08E + 00$
40	$1.65E + 00$	$9.06E - 01$	$3.16E - 02$	$3.95E - 03$	$2.59E + 00$
50	$4.11E + 00$	$2.06E + 00$	$5.90E - 02$	$2.57E - 02$	$6.25E + 00$
60	$7.88E + 00$	$3.64E + 00$	$9.63E - 02$	$1.07E - 01$	$1.17E + 01$
70	$1.27E + 01$	$5.37E + 00$	$1.44E - 01$	$3.31E - 01$	$1.85E + 01$
80	$1.80E + 01$	$6.96E + 00$	$2.02E - 01$	$8.22E - 01$	$2.60E + 01$
90	$2.32E + 01$	$8.18E + 00$	$2.69E - 01$	$1.73E + 00$	$3.34E + 01$
100	$2.78E + 01$	$8.93E + 00$	$3.43E - 01$	$3.18E + 00$	$4.02E + 01$

Table 2 Cross sections of the CC reaction ${}^{16}O(\bar{\nu}_e, e^+)^{16}N(g.s.$ group) as functions of the neutrino energy, E_{ν} (MeV). The unit of the cross section is 10^{-42} cm².

Table 3 The two 2^- states of ¹⁶O and their decay properties. The numbers are not yet established, but still uncertain [\[16\]](#page-15-3). We used $Br^U(\alpha_1)=0.35$ and $Br^D(\alpha_1)=0.83$.

Excited states (J^P, T)		Γ_{α_1}	Γ_p	Γ_{γ}	Reference
	(keV)	(keV)	(keV)	$\rm (eV)$	
$12.53 \text{ MeV} (2,0)$					
				3.4 ± 0.3	[51]
	$0.097 + 0.010$	$0.072 + 0.010$	$0.025 + 0.003$		[10]
	$0.108 + 0.010$	$0.092 + 0.010$	$0.016 + 0.003$	0.55 ± 0.06	$[11]$
Values we used	0.111 ± 0.010	0.092 ± 0.010	0.016 ± 0.003	3.4 ± 0.3	
12.97 MeV $(2^-, 1)$					
		0.69 ± 0.07		3.6 ± 0.6	$\left\lceil 51 \right\rceil$
	$1.59 + 0.14$	$0.60 + 0.08$	$0.99 + 0.12$		$\left[10\right]$
	1.34 ± 0.04	$0.30 + 0.06$	$1.04 + 0.07$	1.6 ± 0.3	$[11]$
Values we used	1.34 ± 0.04			3.6 ± 0.6	

shown. This time, $\sigma_{\text{NC},\gamma}^U$ (red dashed line) and $\sigma_{\text{NC},\gamma}^D = \sigma_{\text{NC}}^D \cdot \text{Br}^D(\alpha_1)$ (red dash-dotted line), with $Br^{U}(\alpha_1)=0.35$ and $Br^{D}(\alpha_1)=0.83$, respectively, are shown in Fig.2. The sum of them $\sigma_{\text{NC},\gamma}^{\text{tot}}$ (red solid line) is also shown and it is larger than $\sigma_{\text{NC},\gamma}^U$ by about 16%, since the ratio $\sigma_{\rm NC, \gamma}^D/\sigma_{\rm NC, \gamma}^U$ =0.16 at β =0.25.

Next, we discuss the high-energy γ -ray production via electromagnetic decay of the two states. The direct M2 electromagnetic transition from the two 2[−] states to the ground state 0^+ is suppressed and electromagnetic transitions to the ground state go through the cascade transitions producing more than two γ rays. Gorodetzky *et al.* [\[51\]](#page-15-29) measured both the electromagnetic cascade decay ${}^{15}N(p,\gamma\gamma){}^{16}O$ and the α decay ${}^{15}N(p,\alpha_1\gamma){}^{12}C(4.4 \text{ MeV})$ from the U and D states in the proton capture experiment. Zijderhand and van der Leun [\[11\]](#page-15-28) also measured both a single γ ray from the electromagnetic decay ${}^{15}N(p,\gamma){}^{16}O$ and the α

Fig. 3 The electron energy spectrum of the β decay, ¹⁶N(g.s.) \rightarrow ¹⁶O(g.s.)+ $e^- + \bar{\nu}_e$. The product of the nuclear shape factor $C(E)$, the phase space factor, and the Fermi function $F(E)$ is shown as a function of the electron kinetic energy T_β (MeV). Here, E and p_e are electron energy and momentum, respectively, and Q is the Q-value for the reaction.

Fig. 4 Illustrative figure for the NC reactions, $\nu + {}^{16}O \rightarrow \nu + {}^{16}O(12.97/12.53 \text{MeV}, 2^{-}).$

decay ${}^{15}N(p, \alpha_1 \gamma){}^{12}C(4.4 \text{ MeV})$ from the two states in the proton capture experiment. Their values are shown in Table 3. We use the radiative decay widths measured by Gorodetzky et al. [\[51\]](#page-15-29) in the present paper. The latest evaluation for the decay parameters of ^{16}O can be found in Ref. [\[38\]](#page-15-17) and the values for the radiative decay widths are the same as those we use in the present paper. As shown in Table 3, we use the total decay widths $\Gamma = 0.111 \pm 0.010$ (keV) and Γ =1.34 \pm 0.04 (keV) for the D and U states, respectively, to calculate the branching ratios of the two states producing γ rays, $Br(D \to \gamma \text{ rays})=\Gamma_{\gamma}/\Gamma=3.1\pm0.04\%$ and $Br(U \to$ γ rays)=0.27±0.02%. If we take Γ_{γ} =3.4±0.3 eV of Gorodetzky *et al.* [\[51\]](#page-15-29) for the D state,

Table 4 The branching ratios of the electromagnetic decay from the two states of ^{16}O . Gorodetzky *et al.* [\[51\]](#page-15-29) measured the widths of the cascade γ rays as in the first column and we calculated the branching ratios in the second column to compare them with the measurements by Zijderhand and van der Leun [\[11\]](#page-15-28). The sum of the branching ratios is normalized to 100% (the radiative width Γ_{γ}).

$12.53 \text{ MeV} (2^-)$	Γ_{γ} (eV)	Branching ratio $(\%)$	Branching ratio $(\%)$
Transition to the state			
$\rightarrow 0$ MeV (0^+)			6.0 ± 0.6
\rightarrow 8.87 MeV (2 ⁻)	0.86 ± 0.10	$25 + 3$	33 ± 2
\rightarrow 7.12 MeV (1 ⁻)	0.51 ± 0.10	15 ± 3	12.0 ± 0.7
\rightarrow 6.13 MeV (3 ⁻)	2.1 ± 0.2	60 ± 6	$49 + 2$
\rightarrow All states	3.4 ± 0.3	100	100 $(0.55 \pm 0.06 \text{ eV})$
Reference	Gorodetzky et al.[51]		Zijderhand and van der Leun[11]
$12.97 \text{ MeV} (2^-)$	Γ_{γ} (eV)	Branching ratio $(\%)$	Branching ratio $(\%)$
Transition to the state			
$\rightarrow 0$ MeV (0^+)			2.1 ± 0.4
\rightarrow 8.87 MeV (2 ⁻)	0.90 ± 0.10	25 ± 6	$42 + 2$
\rightarrow 7.12 MeV (1 ⁻)	0.44 ± 0.10	12 ± 2	6 ± 1
\rightarrow 6.13 MeV (3 ⁻)	2.3 ± 0.3	63 ± 6	50 ± 2
\rightarrow All states	3.6 ± 0.3	100	$100 (1.6 \pm 0.3 \text{ eV})$
Reference	Gorodetzky et al.[51]		Zijderhand and van der Leun[11]

we naturally obtain the total width Γ to be $\Gamma = \Gamma_{\alpha} + \Gamma_{p} + \Gamma_{\gamma} = 0.092 + 0.016 + 0.0034 = 0.1114$ (keV) , which agrees with the total width of the D state listed in Ref. [\[38\]](#page-15-17). Thus, the branching ratio of the D state producing γ rays is Γ_{γ}/Γ =3.1±0.04%. We note that Zijderhand and van der Leun [\[11\]](#page-15-28) adopted the total width Γ =0.108 keV by taking the average of the three previous measurements of the total width [\[10,](#page-14-7) [52,](#page-15-30) [53\]](#page-15-31), including the measurement by Leavitt et al. [\[10\]](#page-14-7). Thus, the measurements of the branching ratios for the D state are consistent with each other except for the γ -ray width. On the other hand, the measurements for the U states are not consistent with each other.

As shown in Table 4, the cascade electromagnetic decay of the U state produces a pair of two (sometimes three) γ rays, 6.84 MeV + 6.13 MeV (63%), 5.85 MeV + 7.12 MeV (12%), 4.10 MeV + 8.87 MeV (25%). As explained in the previous section, the 8.87-MeV state decays mostly through cascade to the ground state, producing two γ rays. Similarly, the cascade electromagnetic decay of the D state produces a pair of two (sometimes three) γ rays, 6.40 $\text{MeV} + 6.13 \text{ MeV} (60\%), 5.41 \text{ MeV} + 7.12 \text{ MeV} (15\%), 3.66 \text{ MeV} + 8.87 \text{ MeV} (25\%).$ We note that the electromagnetic decay will produce mostly two γ rays at the same time and at the same vertex position and that the sum of them is as high as 12.97 MeV and 12.53 MeV from the U and D states, respectively. This feature can be used to identify this reaction channel and the coincidence method can be also used if the multiple-ring reconstruction can be developed [\[54\]](#page-15-32).

We show the cross section of the high-energy γ -ray production via electromagnetic decay of the U and D states in Fig.2 and also in Table 5. We denote them as $\sigma_{\rm NC,2\gamma}^{U} = \sigma_{\rm NC}^{U} \cdot \rm{Br}(U \to \gamma)$ rays), $\sigma_{\text{NC},2\gamma}^D = \sigma_{\text{NC}}^D \cdot \text{Br}(D \to \gamma \text{ rays})$ and the sum of them $\sigma_{\text{NC},2\gamma}^{\text{tot}} = \sigma_{\text{NC},2\gamma}^U + \sigma_{\text{NC},2\gamma}^D$. Since the branching ratios for the U and D states are $Br(U \to \gamma \text{ rays}) = 0.27 \pm 0.02\%$ and $Br(D \to \gamma \text{ axis})$ γ rays) = 3.1 \pm 0.04%, respectively, each of their cross sections is smaller than the CC cross section by three orders of magnitude. However, it should be reminded that all flavors of

Table 5 NC cross sections of 4.4-MeV γ -ray production $(\sigma_{NC,\gamma}^U$ and $\sigma_{NC,\gamma}^D$), 12.97-MeV and 12.53-MeV γ -ray production $(\sigma_{NC,2\gamma}^U$ and $\sigma_{NC,2\gamma}^D$) from the U and D states as functions of the neutrino energy, E_{ν} (MeV). We note that each of the NC cross section is calculated for an average over one neutrino flavor and its anti-neutrino flavor. The unit of the cross section is 10^{-42} cm².

E_{ν} (MeV)	$\sigma_{\mathrm{NC},\gamma}$	$\sigma_{\rm NC, \gamma}$	$\sigma_{\rm NC,2\gamma}$	$\sigma_{\rm NC,2\gamma}$
12				
14	$4.61E - 05$	$1.55E - 05$	$3.56\mathrm{E}{-07}$	$5.81E - 07$
16	$5.41E - 04$	$1.17E - 04$	$4.17E - 06$	$4.37E - 06$
18	$1.95E - 03$	$3.82E - 04$	$1.51E - 05$	$1.43E - 05$
20	$4.88E - 03$	$9.10E - 04$	$3.76E - 05$	$3.40E - 05$
22	$1.00E - 02$	$1.82E - 03$	$7.74E - 05$	$6.81E - 05$
24	$1.83E - 02$	$3.26E - 03$	$1.41E - 04$	$1.22E - 04$
26	$3.05E - 02$	$5.39E - 03$	$2.36E - 04$	$2.01E - 04$
28	$4.79E - 02$	$8.37E - 03$	$3.69E - 04$	$3.13E - 04$
30	$7.14E - 02$	$1.24E - 02$	$5.51E - 04$	$4.63E - 04$
32	$1.02E - 01$	$1.76E - 02$	$7.89E - 04$	$6.59E - 04$
34	$1.42E - 01$	$2.43E - 02$	$1.09E - 03$	$9.08E - 04$
36	$1.91E - 01$	$3.26E - 02$	$1.47E - 03$	$1.22E - 03$
38	$2.50E - 01$	$4.26E - 02$	$1.93E - 03$	$1.59E - 03$
40	$3.22E - 01$	$5.47E - 02$	$2.48E - 03$	$2.04E - 03$
50	$8.90E - 01$	$1.49E - 01$	$6.87E - 03$	$5.58E - 03$
60	$1.86E + 00$	$3.10E - 01$	$1.43E - 02$	$1.16E - 02$
70	$3.21E + 00$	$5.33E - 01$	$2.48E - 02$	$1.99E - 02$
80	$4.85E + 00$	$8.02E - 01$	$3.74E - 02$	$3.00E - 02$
90	$6.59E + 00$	$1.09E + 00$	$5.08E - 02$	$4.06E - 02$
100	$8.24E + 00$	$1.36E + 00$	$6.35E - 02$	$5.07E - 02$

neutrinos and antineutrinos contribute to the number of NC events while the CC reaction depends on the flavor. Only new measurements will resolve the above inconsistencies in the branching ratios between Gorodetzky et al. [\[51\]](#page-15-29) and Zijderhand and van der Leun [\[11\]](#page-15-28).

4. Estimation of the number of events from $^{16}{\rm O}(\bar{\nu}_e,e^+)^{16}{\rm N}(0\,\,{\rm MeV},\,2^-)$ and $^{16}{\rm O}(\nu,\nu')^{16}{\rm O}(12.97/12.53\,\,{\rm MeV},\,2^-),$ induced by supernova neutrinos

We evaluate the number of these CC and NC events induced by neutrinos from SN explosion which can be observed by the SK, a 32 kton water Cherenkov detector [\[55\]](#page-15-33) in the Earth 1 1 . The four bound states of the g.s. group of ¹⁶N, 0 MeV (2⁻), 0.120 MeV (0⁻), 0.298 MeV (3⁻) and 0.397 MeV (1−), in the CC reactions cannot be distinguished in a water-Cherenkov detector unless a small γ ray can be identified. We thus estimate the number of ${}^{16}O(\bar{\nu}_e, e^+){}^{16}N(0)$ MeV, 2^-) and ¹⁶N(g.s. group).

We calculate the number of events using the following parametrization (called KRJ fit [\[56,](#page-15-34) 57) for the normalized SN neutrino spectra $f(E_{\nu})$ as we used in the previous work [\[16\]](#page-15-3):

$$
f(E_{\nu}) = \frac{(\alpha+1)^{\alpha+1}}{\Gamma(\alpha+1)\langle E_{\nu}\rangle^{\alpha+1}} E_{\nu}^{\alpha} \exp\left(-\frac{(\alpha+1)E_{\nu}}{\langle E_{\nu}\rangle}\right),\tag{2}
$$

¹ The evaluations in this section can be applied to the Hyper-K detector (187-kton fiducial vol-ume) [\[22\]](#page-15-9) if the detector threshold on the electron kinetic energy is taken into account. The SK detector can measure the kinetic energy of electron/positron larger than 3.5 MeV [\[55\]](#page-15-33) and the recent Hyper-K study on SN uses 5 MeV for the threshold on the kinetic energy.

where $\langle E_{\nu} \rangle$ is the average neutrino energy. In this expression, $\Gamma(\alpha + 1)$ is the Gamma function and α is the pinching parameter. As the value α becomes larger, the high-energy tail of the distribution is more suppressed for the same average energy. We only calculate the number of events using three typical values of the KRJ fit with $\alpha = 3$ and $\langle E_{\nu} \rangle = 10$, 12 and 14 MeV, and assume that the neutrino spectra are flavor independent in Table 6.

The time-integrated number spectrum of neutrinos emitted from a SN core, dN_{ν}/dE_{ν} , is related to the normalized neutrino spectra $f(E_{\nu})$ as

$$
\frac{dN_{\nu}}{dE_{\nu}} = \frac{E_{\nu}^{\text{tot}}}{\langle E_{\nu} \rangle} f(E_{\nu}),\tag{3}
$$

where E_{ν}^{tot} is the total energy emitted by one neutrino flavor. Hereafter, we set $E_{\nu}^{\text{tot}} =$ 5×10^{52} erg for each neutrino flavor. We calculate the number of events at various average energies using the neutrino flux $F(E_{\nu})$ at a detector on the Earth, which is given as

$$
F(E_{\nu}) = \frac{1}{4\pi d_{\rm SN}^2} \frac{E_{\nu}^{\rm tot}}{\langle E_{\nu} \rangle} f(E_{\nu}).
$$
\n(4)

We set the distance from a detector to the SN to $d_{SN} = 10$ kpc.

We calculate the number of events $N^{(i)}(E_{\nu})$ produced in the energy range from E_{ν} to $E_{\nu} + \Delta E_{\nu}$ for the IBD reaction and the CC reactions ¹⁶O($\bar{\nu}_e$, e^+)¹⁶N(g.s. group) by folding the neutrino flux and the cross sections as,

$$
N^{(i)}(E_{\nu}) = n_{\text{tar}}^{(i)} F(E_{\nu}) \sigma^{(i)}(E_{\nu}) \Delta E_{\nu}, \tag{5}
$$

where $\sigma^{(i)}(E_{\nu})$ stands for the cross section of either IBD or CC reaction and $n_{\text{tar}}^{(i)}$ is the number of either protons or 16 O targets for the case of a 32 kton fiducial volume of the SK detector [\[18\]](#page-15-5). For the IBD reaction (CC reactions), the relation $E_{\nu} = T_{e^+} + 1.80$ MeV (11.44 MeV) between the neutrino energy E_{ν} and the visible energy E_{ν} (= T_{e+}) of the positron holds. In Fig. [5,](#page-12-0) the number of events of the CC reactions are compared with that of the IBD reaction as a function of the visible energy E_{vis} with an energy width of $\Delta E_{\nu} = 2$ MeV, for the KRJ fit with $\alpha = 3$ and $\langle E_{\nu} \rangle = 12$ MeV. The numbers of events of the CC reactions integrated over the neutrino energy up to 100 MeV are summarized in Table 6, for the KRJ fit with $\alpha = 3$ and $\langle E_{\nu} \rangle = 10$, 12 and 14 MeV, where we show the effect of the requirement on the kinetic energy of a positron with $T_e > 5$ MeV or $T_e > 0$ MeV (Threshold). The primary positron spectrum from the CC reaction depends on the assumed SN flux and the cross section, and about 2% of the spectrum lies below 5 MeV for the KRJ fit ($\alpha = 3$) and $\langle E_{\nu} \rangle = 12$ MeV). We do not consider the efficiency of having the coincidence signal with the visible energy greater than 5 MeV, which is estimated to be about 86%. We point out again the importance of the low threshold energy of the CC reactions, since 98%, 54% and only 18% of the SN neutrino flux $F(E_\nu)$ remain after the requirement of $E_\nu > 1.80$ MeV (IBD reaction), 11.44 MeV (CC ¹⁶N(g.s.) reaction) and 18 MeV (typical E_{th} value for CC ¹⁶N($E_x > 3$ MeV) reactions) [\[21\]](#page-15-8), respectively, for the typical KRJ fit with $\alpha = 3$ and $\langle E_{\nu} \rangle = 12$ MeV, and the effect of the requirement on the positron kinetic energy T_e will be imposed additionally.

For the NC reactions, the γ ray of 4.4 MeV, 12.53 MeV or 12.97 MeV is produced independently of the incident neutrino energy E_{ν} , we can calculate only the total number of events $N^{(i)}$ integrated over E_{ν} as

$$
N^{(i)} = n_{\text{tar}} \int_0^{E_{\nu}^{\text{max}}} dE_{\nu} F(E_{\nu}) \sigma^{(i)}(E_{\nu}), \tag{6}
$$

where $\sigma^{(i)}(E_\nu)$ stands for the cross section of any type of NC reactions and n_{tar} is the number of ¹⁶O targets in a 32 kton fiducial volume of the SK detector and we set $E_{\nu}^{\max} = 100 \text{ MeV}$.

We show in Table 6 the total numbers of the NC events containing 4.4-MeV γ rays due to $\sigma_{NC,\gamma}^U$ and $\sigma_{NC,\gamma}^{tot}$ as well as those containing 12.97-MeV and 12.53-MeV γ rays due to $\sigma_{\text{NC},2\gamma}^U$ and $\sigma_{\text{NC},2\gamma}^{\text{tot}}$, for the KRJ fit with $\alpha = 3$ and $\langle E_{\nu} \rangle = 10$, 12 and 14 MeV. We also plot in Fig.5 the total numbers of those NC γ events due to $\sigma_{NC,\gamma}^{tot}$ at $E_{vis} = 4.4 \text{ MeV}$ and $\sigma_{NC,2\gamma}^{tot}$ at $E_{\text{vis}} = 12.97 \text{ MeV}$ in filled black squares, for the KRJ fit with $\alpha = 3$ and $\langle E_{\nu} \rangle = 12 \text{ MeV}$, assuming that the detection efficiency is 100% without considering the energy resolution.

The number of events due to the ${}^{16}O(\bar{\nu}_e, e^+){}^{16}N(g.s.$ group) reaction is smaller by two or three orders of magnitude than that of the IBD events. The total cross section of NC events are summed over 3 flavors of a neutrino and anti-neutrino, namely, 6 times the average NC cross section. That explains why the number of the NC events producing a 4.4-MeV γ ray due to $\sigma_{NC,\gamma}^U$ and $\sigma_{NC,\gamma}^{tot}$ is nearly the same as that of the ${}^{16}O(\bar{\nu}_e,e^+){}^{16}N(g.s.$ group) reaction. The numbers of NC events producing 12.97 and 12.97/12.53-MeV γ ray due to $\sigma_{\text{NC},2\gamma}^U$ and $\sigma_{\text{NC},2\gamma}^{\text{tot}}$, respectively, are smaller by two orders of magnitude than that of the ${}^{16}O(\bar{\nu}_e, e^+){}^{16}N(g.s.$ group) reaction.

Fig. 5 The visible energy spectrum of ${}^{16}O(\bar{\nu}_e, e^+)^{16}N(g.s.$ group) (open triangles) and that of the IBD event spectrum (open circles) are also plotted with an energy-bin width of 2 MeV as a function of the visible energy E_{vis} for the KRJ fit with $\alpha = 3$ and $\langle E_{\nu} \rangle = 12$ MeV. The total numbers of those NC γ events due to $\sigma_{NC,\gamma}^{tot}$ and $\sigma_{NC,2\gamma}^{tot}$ are plotted in filled black squares at $E_{\rm vis} = 4.4 \text{ MeV}$ and at $E_{\rm vis} = 12.97 \text{ MeV}$, for the same KRJ fit values.

5. Summary

In the previous work, we discussed the detection of a single 4.4-MeV γ ray produced in the neutrino NC reaction ¹⁶ $O(\nu, \nu')$ ¹⁶ $O(12.97 \text{ MeV}$ and 12.53 MeV, 2⁻), after evaluating both the isospin-mixing parameter β and the quenching factors, $f_s = g_s^{\text{eff}}/g_s$ of the spin g factor

	KRJ parameters $(\alpha, \langle E_{\nu} \rangle$ [MeV])	Condition	(3, 10)	(3, 12)	(3, 14)
IBD	$p(\bar{\nu}_e,e^+)n$	$T_e > 0$ MeV	4.83×10^{3}	4.88×10^{3}	6.88×10^{3}
		$T_e > 5.0$ MeV	4.71×10^3	4.81×10^{3}	6.84×10^{3}
CC	${}^{16}O(\bar{\nu}_e,e^+)^{16}N(g.s., 2^-)$	$T_e > 0$ MeV	$1.6\,$	4.6	10.6
		$T_e > 5.0$ MeV	1.5	4.5	10.5
	${}^{16}O(\bar{\nu}_e,e^+){}^{16}N(g.s.$ group)	$T_e > 0$ MeV	2.8	7.8	17.4
		$T_e > 5.0$ MeV	2.6	7.6	17.2
NC		$E_{\gamma} = 4.4 \text{ MeV}$	1.9	5.1	10.7
		$E_{\gamma} = 4.4 \text{ MeV}$	2.2	5.9	12.4
		$E_{2\gamma}$ =12.97 MeV	0.015	0.039	0.082
	$\sigma^U_{\mathrm{NC},\gamma} \ \sigma^{\mathrm{tot}}_{\mathrm{NC},\gamma} \ \sigma^U_{\mathrm{NC},2\gamma} \ \sigma^{\mathrm{tot}}_{\mathrm{NC},2\gamma}$	$E_{2\gamma}$ =12.97 or 12.53 MeV	0.027	0.072	0.151

Table 6 Expected number of neutrino events from a core-collapse SN at 10 kpc to be detected at SK (32-kton fiducial volume) for the models with KRJ fit.

and $f_A = g_A^{\text{eff}}/g_A$ of the axial-vector coupling constant g_A , of the two 2⁻ states [\[16\]](#page-15-3). In this report, we have further examined both the β decay of the CC reaction ${}^{16}O(\bar{\nu}_e, e^+){}^{16}N(g.s.$ group) and the electromagnetic decay of the NC reaction ${}^{16}O(\nu, \nu'){}^{16}O(12.97 \text{ MeV}$ and 12.53 MeV, 2^-) producing high-energy γ rays of 12.97 MeV and 12.53 MeV, using the same quenching factors for the two 2[−] states. We have evaluated the number of these CC and NC reactions induced by neutrinos from SN explosion which can be observed by the SK, a 32 kton water Cherenkov detector [\[55\]](#page-15-33) in the Earth.

Even though the cross sections of these CC/NC reactions are small, the application of the present work for the future SN neutrino events by water Cherenkov detectors has some unique features. Above all, the coincidence method can be applied for the more solid identification in the former CC reaction. In the latter NC reaction, the large visible energy of 12.97 MeV and 12.53 MeV can be produced and the coincidence method can be also used if the multiple-ring reconstruction can be developed [\[54\]](#page-15-32). Further, it is important for the study of SN physics to have the detection channels which have significant sensitivities at as low neutrino energy as 10 to 20 MeV, where the majority of the neutrino energy spectra from SN bursts lie. These neutrino-oxygen reactions have the lowest energy thresholds (11.44 MeV and 12.97 MeV) among the neutrino-oxygen reactions, which correspond to the energy levels of the $T=1$ ground states of ¹⁶N and ¹⁶O. They have the dominant cross sections from 12 to 30 MeV, above which the cross sections of other CC and NC reactions from the $T=1$ excited states dominate.

We also note that the delayed coincidence method to this reaction can be applied in the Hyper-K detector, even without the neutron tagging method using Gd. The recent study by the Hyper-K Collaboration on the detection of the SN neutrino events [\[22\]](#page-15-9) comments that they do not consider the γ -ray emission from the NC interactions on ¹⁶O nuclei, since a dominant channel ${}^{16}O(\nu, \nu'){}^{16}O(E_x > 16 \text{ MeV})$ mainly produces only γ rays with an energy of 5 MeV to 9 MeV [\[58\]](#page-15-36) and the visible energy from these events would typically be below 5 MeV (Hyper-K energy threshold) after Compton scattering on an electron or electronpositron pair production. Our study for the coincidence method including the high energy γ rays may turn out to be useful.

The JUNO experiment [\[23\]](#page-15-10) uses the NC reaction ${}^{12}C(\nu, \nu'){}^{12}C(15.11 \text{ MeV}, 1^+)$ and CC reactions ¹²C(ν_e , e^{-})¹²N(g.s., 1⁺) with the subsequent β decay and ¹²C($\bar{\nu}_e$, e^{+})¹²B(g.s., 1⁺)

with the subsequent β decay [\[24\]](#page-15-11), as the main detection channels for the analysis of the SN neutrino bursts, in addition to the IBD reaction, elastic ν -p scattering and elastic ν -e scattering [\[25–](#page-15-12)[27\]](#page-15-13). It is important to note some basic features of the CC/NC neutrinooxygen reactions which are different from those of the CC/NC neutrino-carbon reactions. The neutrino-carbon cross sections related to the 1^+ state are larger by two orders of magnitude than the neutrino-oxygen cross sections from the 12.97 and 12.53 states (2^-) since the former neutrino-¹²C reactions have a large matrix element causing the spin-flip transition from $1p_{3/2}$ to $1p_{1/2}$, while the latter neutrino-¹⁶O reactions go through the spin-dipole transition from fully occupied $1p$ shells to $2s - 1d$ shells, which is smaller by an order of magnitude than the former. In addition, the electromagnetic decay branching ratio $(\Gamma_{\gamma}/\Gamma=96\%)$ of the former state ¹²C(15.11 MeV, 1⁺) [\[59\]](#page-15-37) is larger by two order of magnitude than that (Γ_{γ}/Γ) of the 2^- states of ¹⁶O. This is because in the former state ¹²C(15.11 MeV), the electromagnetic decay is dominant and the hadronic decay to $p+^{11}B$ decay is suppressed due to the threshold $(E_{th} = 15.96 \text{ MeV})$, while in the latter 2⁻ states of ¹⁶O, the hadronic decays are allowed $(E_{\text{th}} = 12.13 \text{ MeV}$ for $p+15\text{N}$ decay) and the electromagnetic decay branching ratio becomes relatively very small.

We hope that new accurate measurements of the cross section of ${}^{16}O(e, e'){}^{16}O(12.53 \text{ MeV},$ 12.97 MeV, 2^-) and the branching ratios of ¹⁶O(12.53 MeV, 12.97 MeV, 2^-) decaying to the p, α and γ channels will be performed in the near future at the low-energy electron accelerators ($E_e = 30{\text{-}}100 \text{ MeV}$), at MESA accelerator [\[60\]](#page-15-38) or at the ULQ2 facility at the Research Center for Electron-Photon Science (Tohoku University) [\[61\]](#page-15-39), so that the prediction of both the CC/NC neutrino-oxygen cross sections for 12.53 MeV and 12.97 MeV (2−) can be accurate to a level of 10% or less.

Acknowledgment

This work was partially supported by JSPS Grant-in-Aid for Scientific Research (No. JP19K03855, JP20K03973, No. JP20K03989, and No. JP24K07021) and also by JSPS Grant-in-Aid for Scientific Research on Innovative Areas "Unraveling the History of the Universe and Matter Evolution with Underground Physics" (No. JP19H05802 and No. JP19H05811), Grant-in-Aid for Transformative Research Areas "The creation of multimessenger astrophysics" (No. JP24H01817), and Grant-in-Aid for Transformative Research Areas "Investigation on the Origin and Evolution of Matter in the Universe by Extremely Rare Events: Frontier of Creating a New Insight on the Matter in the Universe" (No. JP24H02245).

References

- [1] M. Stroetzel, Z. f. Physik **214**, 357 (1968).
- [2] J.C. Kim, R.P. Singhal and H.S. Caplan, Can. J. Phys.48, 83 (1970).
- [3] G.J. Vanpraet and W.C. Barber, Nucl.Phys. 79, 550 (1966).
- [4] I. Sick, E.B. Hughes, T.W. Donnely, J.D. Walecka and G.E. Walker, Phys. Rev.Lett. 23, 1117 (1969).
- [5] T.W. Donnelly and J.D. Walecka, Phys.Lett. B41, 275 (1972).
- [6] T.W. Donnelly and J.D. Walecka, Ann.Rev.Nucl.Sci.25, 329 (1975).
- [7] J. D. Walecka, in Muon Physics, eds V. H. Hughes and C. S. Wu (Academic, New York 1975), Section 4, Page 113-217.
- [8] W.C. Haxton, Phys.Rev.D36, 2283 (1987).
- [9] G.J. Wagner, K.T. Knopfle, G. Mairle, F. Doll, H. Hafner and J.L.C. Ford, Jr., Phys. Rev. C 16, 1271 (1977).
- [10] R. A. Leavitt, H. C. Evans, G. T. Ewan, H.-B. Mak, R. E. Azuma, C. Rolfs, and K. P. Jackson, Nucl.

Phys. A 410, 93 (1983).

- [11] F. Zijderhand and C. van der Leun, Nucl. Phys. A 460, 181 (1986).
- [12] R.J. Charity et al., Phys. Rev. C **99**, 044304 (2019).
- [13] E. Adelberger, R. Marrs, K. Snover and J. Bussoletti, Phys. Rev. C15, 484 (1977).
- [14] J.B. Flanz, R. Hicks, R. Lindgren, G. Peterson, J. Dubach and W. Haxton, Phys. Rev. Lett. 43, 1922 (1979).
- [15] P. von Neumann-Cosel, H.-D. Gräf, U. Krämer, A. Richter and E. Spamer, Nucl.Phys. A669, 3 (2000).
- [16] M. Sakuda, T. Suzuki, M.S. Reen, K. Nakazato and H. Suzuki, PTEP 2023, 013D02 (2023).
- [17] K. Abe et al. (SK Collaboration), Astropart. Phys. **81**, 39 (2016).
- [18] M. Ikeda et al. (SK Collaboration), Astrophys. J. **669**, 519 (2007).
- [19] E. Kolbe, K. Langanke, G. Martínez-Pinedo, and P. Vogel, J. Phys. G 29, 2569 (2003).
- [20] T. Suzuki, S. Chiba, T. Yoshida, K. Takahashi, and H. Umeda, Phys. Rev. C98, 034613 (2018).
- [21] K. Nakazato, T. Suzuki, M. Sakuda, PTEP 2018, 123E02 (2018).
- [22] K. Abe et al. (Hyper-K Collaboration), Astrophys. J. 916:15, 1-17 (2021).
- [23] F. An et al. (JUNO Collaboration), J. Phys. G 43, 030401 (2016).
- [24] M. Fukugita, Y. Kohyama and K. Kubodera, Phys. Lett. B 212, 139 (1988).
- [25] A. Strumia and S. Vissani, Phys. Lett. B **564**, 42-54 (2003) (Eq.25).
- [26] G. Ricciardi, N. Vignarolia, and F. Vissani, JHEP 08 **2022**, 212 (2022).
- [27] J. F. Beacom, W. M. Farr, and P. Vogel, Phys. Rev. D 66, 033001 (2002).
- [28] B. Bodmann et al. (KARMEN Collaboration), Phys. Lett. B 267, 321 (1991).
- [29] B. Armbruster et al.(KARMEN Collaboration), Phys. Lett. B 423, 15 (1998).
- [30] D.A. Krakauer et al.(E225 Collaboration), Phys. Rev. C 45, 2450 (1992).
- [31] B. Bodmann et al. (KARMEN Collaboration), Phys. Lett. B **332**, 251 (1994).
- [32] B. Bodmann et al. (KARMEN Collaboration), Phys. Lett. B 339, 215 (1994).
- [33] L.B. Auerbach et al., Phys. Rev. C 63, 112001 (2001).
- [34] L.B. Auerbach et al. (LSND Collaboration), Phys. Rev. C **64**, 065501 (2001).
- [35] C. Athanassopoulos et al. (LSND Collaboration), Phys. Rev. C 56, 2806 (1997).
- [36] L.B. Auerbach et al. (LSND Collaboration), Phys. Rev. C 66, 015501 (2002).
- [37] T. Suzuki, K. Nakazato, M. Sakuda, Nucl. Phys. A 1038, 122719 (2023).
- [38] D.R. Tilley, H.R. Weller and C.M. Cheves, Energy levels of light nuclei $A=16-17$, Nucl. Phys. A 564, 1 (1993).
- [39] M.Dragowsky et al. (SNO Collaboration), Nucl. Instrum. Meth. A 481, 284 (2002).
- [40] E. Blaufuss et al. (SK Collaboration), Nucl. Instrum. Meth. A 458, 638 (2001).
- [41] K. Abe et al. (T2K Collaboration), Phys. Rev. D90, 072012 (2014).
- [42] L Wan et al. (SK Collaboration), Phys. Rev. D99, 032005 (2019).
- [43] K. Abe et al. (T2K Collaboration), Phys. Rev. D100, 112009 (2019).
- [44] S. Sakai et al. (SK Collaboration), Phys. Rev. D109, L011101 (2024).
- [45] A.Ankowski, O.Benhar, T.Mori, R. Yamaguchi and M.Sakuda, Phys.Rev.Lett.108, 052505 (2012).
- [46] R.P Feynman and M. Gell-Mann, Phys. Rev 109, 193 (1958).
- [47] S.S. Gerstein and Y.B. Zeldovich, Sov.Phys.JETP 2, 576 (1956).
- [48] LI. Marti et al. (SK Collaboration), Nucl. Instrum. Meth. A **959**, 163549 (2020).
- [49] K. Abe et al. (SK Collaboration), Nucl. Instrum. Meth. A 1027, 166248 (2022).
- [50] J. F. Beacom and M. R. Vagins, Phys. Rev. Lett. 93, 171101 (2004).
- [51] S. Gorodetzky, J. C. Adloff, F. Brochard, P. Chevallier, D. Disdier, Ph. Gorodetzky, R. Modjtahed-Zadeh and F. Scheibling, Nucl. Phys. A 113, 221 (1968).
- [52] B. Maurel and G. Amsel, Nucl. Instr. Meth. 218, 159 (1983).
- [53] H. Damjantschitsch, M. Weiser, G. Heusser, S. Kalbitzer and H. Mannsperger, Nucl. Instr. Meth. 218, 129 (1983).
- [54] Z. Li et al. (SK Collaboration), Phys. Rev. D98, 052006 (2018).
- [55] K. Abe et al. (SK Collaboration), Phys. Rev. D94, 052010 (2016).
- [56] M. Th. Keil, G. Raffelt and H.-Th. Janka, Astrophys. J. 590, 971 (2003).
- [57] I. Tamborra, B. Müller, L. Hüdepohl, H.-Th. Janka and G. Raffelt, Phys. Rev. D86, 125031 (2012).
- [58] K. Langanke, P. Vogel and E. Kolbe, Phys. Rev. Lett. 76, 2629 (1996).
- [59] J.H. Kelley, J. Purcell and C. Shen, Nucl. Phys. A968, 71 (2017).
- [60] B.S. Schlimme et al. (MAGIX Collaboration), Nucl.Instrum.Meth. A 1013, 165668 (2021).
- [61] T. Suda, Low-energy electron scattering facilities in Japan, J. Phys. Conf. Ser. 2391, 012004 (2022).