

A Tunneling Method for Nonlinear Multi-objective Optimization Problems

Bikram Adhikary¹ and Md Abu Talhamainuddin Ansary^{2,*}

^{1, 2}Department of Mathematics, Indian Institute of Technology Jodhpur,
Jodhpur, 342030, India.

*Corresponding author(s). E-mail(s): md.abutalha2009@gmail.com;
Contributing authors: bikram.adhikary333@gmail.com;

Abstract

In this paper, a tunneling method is developed for nonlinear multi-objective optimization problems. The proposed method is free from any kind of priori chosen parameter or ordering information of objective functions. Using this method, global Pareto front can be generated for non-convex problems with more than one local front. An algorithm is developed using some ideas of single objective tunneling method. Convergence of this algorithm is justified under some mild assumptions. In addition to this, some numerical examples are included to justify the theoretical results.

Keywords: multi-objective optimization, global optimization, non-convex optimization; tunneling algorithm, efficient solution

MSC Classification: 49M37 , 65K05 , 90C29 , 90C30.

1 Introduction

In multi-objective optimization problem, several objective functions are minimized simultaneously. Applications of this type of problem can be found in environmental analysis [1, 2], space exploration[3, 4], portfolio optimization [5], management science [6, 7], medical science [8] etc.. Classical methods for solving multi-objective optimization problems are Scalarization methods [9–11] or heuristic methods [12, 13]. Scalarization techniques are user dependent and heuristic methods do not ensure a descending sequence and hence can not guarantee the converge to the optimal solution. In recent times, numerous researchers have developed descent methods for nonlinear

which are free from above limitations. These methods can be treated as extension of single objective descent methods to the multi-objective scenario. Major developments in this area are descent methods for smooth unconstrained [14–17], constrained multi-objective optimization problems [18–20], and unconstrained non-smooth multi-objective optimization problems [21–24]. Global/local convergence of each method is justified under certain assumptions. Solution of a multi-objective optimization problem is a set of efficient solutions. To generate well distributed approximate Pareto front, multi-start technique is used in most cases. Some different spreading techniques developed in [18, 20]. However in case of non-convex multi-objective optimization problems, all these methods can generate local efficient solution unless initial approximation is very close to global efficient solution. As global efficient solution dominate local efficient solutions, it is very difficult to generate global Pareto front for non-convex multi-objective optimization problems using line search techniques.

Recently many researchers have developed different algorithms to find global minima of non-convex single objective optimization problems (see [25–27]). In these methods, initially a local minimum solution is obtained by using some gradient descent method. Then another local minimum solution with lower objective value is obtained using some auxiliary function. This process is repeated to get a global minimum solution. In this paper, we have adopted some ideas of single objective tunneling method in [25] and developed a tunneling method for nonlinear multi-objective optimization problems.

Outline of this paper is as follows. Some prerequisites related to present contribution is discussed in Section 2. A multi-objective tunneling method is developed with the help of a multi-objective auxiliary function in Section 3. An algorithm is proposed and convergence of this algorithm is justified in this section. Finally, the proposed method is verified with some numerical examples in Section 4.

2 Preliminaries

Throughout the paper, we use the notations $\Lambda_p := \{1, 2, \dots, p\}$, $\mathbb{R}_+^p := \{x \in \mathbb{R}^p \mid x_i \geq 0, \forall i \in \Lambda_p\}$, $\mathbb{R}_{++}^p := \{x \in \mathbb{R}^p \mid x_i > 0, \forall i \in \Lambda_p\}$ where $p \in \mathbb{N}$. For $x, y \in \mathbb{R}^p$, vector inequalities are interpreted component wise.

A constrained nonlinear multi-objective optimization problem is of the form:

$$(MOP_C) : \begin{aligned} & \min_{x \in \mathbb{R}^n} (f_1(x), f_2(x), \dots, f_m(x)) \\ & \text{s.t.} \quad g_i(x) \leq 0, \quad i \in \Lambda_p. \end{aligned}$$

where $m \geq 2$ and $f_k, g_i : \mathbb{R}^n \rightarrow \mathbb{R}$, $\forall k \in \Lambda_m$ and $\forall i \in \Lambda_p$. Set of feasible solutions of (MOP_C) is $X = \{x \in \mathbb{R}^n \mid g_i(x) \leq 0, \forall i \in \Lambda_p\}$.

If there exists any $x^* \in X$, where all objective functions are minimized simultaneously, then x^* is said to be an ideal solution. However, in practical value of an objective function often increases when the value of another one decreases. Thus, the

idea of efficiency, rather than optimality, takes centre stage.

A feasible point x^* is deemed to be an *efficient solution* of (MOP_C) if there is no other $x \in X$, such that both $f(x) \leq f(x^*)$ and $f(x) \neq f(x^*)$ hold, where

$$f(x) = (f_1(x), f_2(x), \dots, f_m(x)).$$

Additionally, a feasible point x^* is characterized as a *weak efficient solution* of the (MOP_C) if there is no $x \in X$ for which $f(x) < f(x^*)$ is true. If X^* represents the collection of efficient solutions for (MOP_C) , then $f(X^*)$ is termed as the Pareto front of (MOP_C) .

The following theorem is a necessary condition for x^* to be a local weak efficient solution of (MOP_C) where $f_k, g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ are continuously differentiable for $k \in \Lambda_m$ and $i \in \Lambda_p$.

Theorem 2.1. Fritz John necessary condition (Theorem 3.1.1 in [9]) If any feasible point x^* is a weak efficient solution of (MOP_C) , then there exists $(\lambda, \mu) \in \mathbb{R}_+^m \times \mathbb{R}_+^p$, $(\lambda, \mu) \neq \mathbf{0}^{m+p}$, satisfying

$$\sum_{k \in \Lambda_m} \lambda_k \nabla f_k(x^*) + \sum_{i \in \Lambda_p} \mu_i \nabla g_i(x^*) = 0, \quad (2.1)$$

$$\mu_i g_i(x^*) = 0 \quad \forall i \in \Lambda_p. \quad (2.2)$$

The set of the vector $(\lambda, \mu) \in \mathbb{R}_+^m \times \mathbb{R}_+^p \setminus \{\mathbf{0}^{m+p}\}$, satisfying (2.1) and (2.2) are referred to as Fritz John multipliers corresponding to x^* . However, relying solely on the Fritz John condition does not guarantee the existence of $\lambda_k > 0$ for at least one $k \in \Lambda_m$. Hence, specific constraint qualifications or regularity conditions must be met to ensure this occurrence. In our analysis, we will consistently consider the Mangasarian–Fromovitz constraint qualification (MFCQ) in [28].

Definition 2.1. The MFCQ is considered to hold at $x \in X$, if there exists a $z \in \mathbb{R}^n$ such that $\nabla g_i(x)^T z < 0$ for $i \in I(x) = \{i \in \Lambda_p : g_i(x) = 0\}$, the set of active constraints.

By Gordan’s theorem of alternative, the MFCQ holds at a feasible point x if and only if $\sum_{i \in I(x)} \mu_i \nabla g_i(x) = 0$, with $\mu_i \geq 0$, has no non-zero solution (see [20]).

If MFCQ isn’t met at point x (not necessarily feasible), then there exists at least one non-zero multiplier $\mu_i > 0, i \in I(x)$ such that $\sum_{i \in I(x)} \mu_i \nabla g_i(x) = 0$. Then substituting $\lambda_k = 0, \forall k \in \Lambda_m$, we can show that x is a Fritz John point.

If MFCQ does not hold in any infeasible point, then there exists at least one non-zero multiplier $\mu_i > 0, i \in \bar{I}(x)$ where $\bar{I}(x) = \{i \in \Lambda_p : g_i(x) = \max\{g_i(x), i \in \Lambda_p\}\}$ such that $\sum_{i \in \bar{I}(x)} \mu_i \nabla g_i(x) = 0$. Again substituting $\lambda_k = 0, \forall k \in \Lambda_m$, we can show that x is an infeasible Fritz John point.

Definition 2.2. A feasible point x^* is said to be a critical point of (MOP_C) if (2.1) and (2.2) hold at x^* with $\lambda_k > 0$ for at least one $k \in \Lambda_m$.

3 Multi-objective tunneling method

In this section, we derive a multi-objective tunneling method for (MOP_C) that can help us to find a global weak efficient solution for non-convex multi-objective optimization problems. This algorithm is built of a series of cycles, each cycle consisting of two phases:

- (I) The minimization phase, which aims to reduce the current functional value until a local efficient solution is discovered.
- (II) The tunneling phase, which aims to find a point \bar{x} other than the previous local efficient solution found, such that when employed as the initial point for the subsequent minimization phase, the resulting critical point possesses a function value, which does not exceed the previous discovered local Pareto value.

Once finding out a local efficient solution, we construct a multi-objective variant of the auxiliary function, *tunneling function* as:

$$(TP) : \begin{aligned} & \min_{x \in \mathbb{R}^n} (T_1(x), T_2(x), \dots, T_m(x)) \\ & \text{s.t.} \quad T_k(x) \leq 0, \quad k \in \Lambda_m \\ & \quad \quad g_i(x) \leq 0, \quad i \in \Lambda_p \end{aligned}$$

where,

$$T_k(x) = \begin{cases} \frac{f_k(x) - f_k(x^*)}{[(x - x^*)^T(x - x^*)]^\eta}, & \text{for } x \neq x^*, \\ \infty, & \text{otherwise,} \end{cases}$$

for $k \in \Lambda_m$. Here $\eta > 0$ and x^* is a weak efficient solution of (MOP_C) . One can observe that if \bar{x} is local efficient solution of (TP) then $f_k(\bar{x}) \leq f_k(x)$ for all $k \in \Lambda_m$. In the following theorem, we have justified that any Fritz John point of (TP) is a critical point of (MOP_C) .

Theorem 3.1. If \bar{x} represents a Fritz John point of (TP) and MFCQ holds at \bar{x} , then \bar{x} also serves as a critical point for problem (MOP_C) .

Proof: Let us suppose, \bar{x} is a *Fritz John* point of the problem (TP) . Then there exists $(\lambda, \mu, \sigma) \in \mathbb{R}_+^m \times \mathbb{R}_+^m \times \mathbb{R}_+^p$, $(\lambda, \mu, \sigma) \neq \mathbf{0}^{m+m+p}$ such that

$$\sum_{k=1}^m (\lambda_k + \mu_k) \nabla T_k(\bar{x}) + \sum_{i=1}^p \sigma_i \nabla g_i(\bar{x}) = 0 \quad (3.1)$$

$$\mu_k T_k(\bar{x}) = 0, \quad \forall k \in \Lambda_m \quad (3.2)$$

$$\sigma_i g_i(\bar{x}) = 0, \quad \forall i \in \Lambda_p \quad (3.3)$$

hold.

Since $\lambda_k, \mu_k, \sigma_i > 0$ for at least one $k \in \Lambda_m$ or $i \in \Lambda_p$, $\lambda_k + \mu_k > 0$ or $\sigma_i > 0$ holds for at least one $k \in \Lambda_m$ or $i \in \Lambda_p$. Hence from (3.1), *MFCQ* does not hold at \bar{x} . Then

substituting $\lambda_k = 0, \forall k \in \Lambda_m$ in (3.1) we have,

$$\sum_{k=1}^m \mu_k \nabla T_k(\bar{x}) + \sum_{i=1}^p \sigma_i \nabla g_i(\bar{x}) = 0, \quad (3.4)$$

$$\mu_k T_k(\bar{x}) = 0, \quad \forall k \in \Lambda_m, \quad (3.5)$$

$$\sigma_i g_i(\bar{x}) = 0, \quad \forall i \in \Lambda_p. \quad (3.6)$$

Now for $k \in \Lambda_m$,

$$\nabla T_k(x) = \frac{\nabla f_k(x)}{\|x - x^*\|^{2\eta}} - \frac{2\eta \cdot (x - x^*)}{\|x - x^*\|^2} T_k(x).$$

Using (3.4), we have

$$\sum_{k=1}^m \mu_k \frac{\nabla f_k(\bar{x})}{\|\bar{x} - x^*\|^{2\eta}} - \sum_{k=1}^m \mu_k \frac{2\eta \cdot (\bar{x} - x^*)}{\|\bar{x} - x^*\|^2} T_k(\bar{x}) + \sum_{i=1}^p \sigma_i \nabla g_i(\bar{x}) = 0$$

This implies,

$$\sum_{k=1}^m \mu_k \frac{\nabla f_k(\bar{x})}{\|\bar{x} - x^*\|^{2\eta}} + \sum_{i=1}^p \sigma_i \nabla g_i(\bar{x}) = 0$$

Again this implies,

$$\sum_{k=1}^m \gamma_k \nabla f_k(\bar{x}) + \sum_{i=1}^p \sigma_i \nabla g_i(\bar{x}) = 0,$$

$$\text{where } \gamma_k = \frac{\mu_k}{\|\bar{x} - x^*\|^{2\eta}}, \quad \forall k \in \Lambda_m.$$

Hence, there exists $(\gamma, \sigma) \in \mathbb{R}_+^m \times \mathbb{R}_+^p, (\gamma, \sigma) \neq \mathbf{0}^{m+p}$ such that

$$\sum_{k=1}^m \gamma_k \nabla f_k(\bar{x}) + \sum_{i=1}^p \sigma_i \nabla g_i(\bar{x}) = 0, \quad (3.7)$$

$$\sigma_i g_i(\bar{x}) = 0, \quad \forall i \in \Lambda_p \quad (3.8)$$

hold and therefore, \bar{x} is a *Fritz John point* of (MOP_C) .

Next, we show that $\gamma_k > 0$, for at least one $k \in \Lambda_m$. Otherwise from (3.7), we have $\sum_{i=1}^p \sigma_i \nabla g_i(\bar{x}) = 0$, with $\sigma_i > 0$, for at least one $i \in \Lambda_p$. This implies, *MFCQ* does not hold at \bar{x} for (MOP_C) . This contradicts our assumption, that *MFCQ* holds

for every feasible point of (MOP_C) .

Hence,

$$\sum_{k=1}^m \gamma_k \nabla f_k(\bar{x}) + \sum_{i=1}^p \sigma_i \nabla g_i(\bar{x}) = 0, \quad (3.9)$$

$$\sigma_i g_i(\bar{x}) = 0, \quad \forall i \in \Lambda_p \quad (3.10)$$

hold at \bar{x} with $\gamma_k > 0$, for at least one $k \in \Lambda_m$ and therefore, \bar{x} is a critical point of (MOP_C) . \square

From this theorem one can observe that, using the tunneling phase, we can have a critical point with less objective values. Thus, multiple repetitions of the tunneling phase can generate a global weak efficient solution of (MOP_C) .

The above ideas are summarized in the following algorithm. In this algorithm, PF denotes the Pareto front of (MOP_C) prior to tunneling, while PFT signifies the Pareto front of (MOP_C) after tunneling. Furthermore, WPF represents the weak Pareto front of (MOP_C) before tunneling, and $WPFT$ refers to the weak Pareto front of (MOP_C) after tunneling.

Algorithm 1. (*Tunneling Algorithm for (MOP_C)*):

Sep 0: Supply $f, g, X^0 =$ a set of initial approximations.

Set $PF = PFT = WPF = WPFT = \emptyset$.

Sep 1: **for** each $x^0 \in X^0$, **do**

(a) Solve MOP_C using some descent method to find a local weak efficient solution x^* . Update $WPF = WPF \cup \{f(x^*)\}$.

(b) Construct tunneling function TP using x^* .

(c) Solve (TP) using some descent method to find \bar{x} , a Fritz John point of (TP) . Update $WPFT = WPFT \cup \{f(\bar{x})\}$.

Sep 2: Generate Pareto Front PF using WPF by removing dominated points.

Sep 3: Generate Pareto Front PFT using $WPFT$ by removing dominated points.

Output: PF and PFT .

4 Numerical illustration and discussion

In this section, Algorithm 1 is applied to several test problems. MATLAB (R2023b) code is developed to implement Algorithm 1. Both (MOP_C) and (TP) are solved using modified SQCQP method in [20].

In the following example from [18], we verify that a Fritz John point of (TP) will be a critical point of MOP_C :

$$(DTLZ1n2) : \min_{x \in \mathbb{R}^n} (f_1(x), f_2(x)) \\ \text{s.t. } 0 \leq x_1, x_2 \leq 1$$

where,

$$f_1(x) = 0.5(1 + gx)x_1, \\ f_2(x) = 0.5(1 + gx)(1 - x_1), \\ \text{and } gx = 100[1 + (x_2 - 0.5)^2 - \cos\{20\pi(x_2 - 0.5)\}].$$

In the process of applying the tunneling method and generating the Pareto front for the above problem, two cases arise.

Case 1: Let us consider the initial point $x^0 = (0.9956, 0.0018)^T$. Then, $x^* = (1, 0)^T$ and

$$\nabla f_1(x^*) = (13, -49.8569)^T, \quad \nabla f_2(x^*) = (-13, 0)^T, \\ \nabla g_1(x^*) = (-1, 0)^T, \quad \nabla g_2(x^*) = (0, -1)^T, \quad \nabla g_3(x^*) = (1, 0)^T, \quad \nabla g_4(x^*) = (0, 1)^T. \\ \text{Choosing } \lambda_1 = 0, \lambda_2 = 1, \mu_1 = 0, \mu_2 = 0, \mu_3 = 1, \mu_4 = 0 \text{ satisfies the equation (2.1)} \\ \text{showing that } x^* \text{ is a KKT point of the problem (DTLZ1n2).}$$

After tunneling, we get $\bar{x} = (0, 0.2997)^T$ with $T(\bar{x}) = (-11.7248, 2.2655)^T$, showing \bar{x} is infeasible for (TP) , which is constructed from $(DTLZ1n2)$ using x^* . But, \bar{x} is a KKT point of the problem $(DTLZ1n2)$, as:

$$\nabla f_1(\bar{x}) = (2.5119, 0)^T, \quad \nabla f_2(\bar{x}) = (-2.5119, -71.4928)^T,$$

$\nabla g_1(\bar{x}) = (-1, 0)^T, \quad \nabla g_2(\bar{x}) = (0, -1)^T, \quad \nabla g_3(\bar{x}) = (1, 0)^T, \quad \nabla g_4(\bar{x}) = [0, 1]^T$ and choosing $\lambda_1 = 1, \lambda_2 = 0, \mu_1 = 2.5119, \mu_2 = 0, \mu_3 = 0, \mu_4 = 0$ satisfies the equation (2.1).

Case 2: Let us consider the initial point $x^0 = (0.0048, 0.6076)^T$. Then, $x^* = (0, 0.5997)^T$ and

$$\nabla f_1(x^*) = (1.0080, 0)^T, \quad \nabla f_2(x^*) = (-1.0080, -57.1462)^T,$$

$\nabla g_1(x^*) = (-1, 0)^T, \quad \nabla g_2(x^*) = (0, -1)^T, \quad \nabla g_3(x^*) = (1, 0)^T, \quad \nabla g_4(x^*) = (0, 1)^T$. Choosing $\lambda_1 = 1, \lambda_2 = 0, \mu_1 = 1, \mu_2 = 0, \mu_3 = 0, \mu_4 = 0$ satisfies the equation (2.1) showing that x^* is a KKT point of the problem $(DTLZ1n2)$.

After tunneling, we get $\bar{x} = (0, 0.5014)^T$ with $T(\bar{x}) = (0, -80.0251)^T$ and $g_i(\bar{x}) \leq 0$, for $i = 1, 2, 3, 4$, showing \bar{x} is a feasible point of (TP) , which is constructed from $(DTLZ1n2)$ using x^* .

Also, \bar{x} is a KKT point of the problem $(DTLZ1n2)$, as:

$$\nabla f_1(\bar{x}) = (0.7028, 0)^T, \quad \nabla f_2(\bar{x}) = (-0.7028, 282.7663)^T,$$

$$\nabla g_1(\bar{x}) = (-1, 0)^T, \quad \nabla g_2(\bar{x}) = (0, -1)^T, \quad \nabla g_3(\bar{x}) = (1, 0)^T, \quad \nabla g_4(\bar{x}) = (0, 1)^T$$

and choosing $\lambda_1 = 1$, $\lambda_2 = 0$, $\mu_1 = 0.7028$, $\mu_2 = 0$, $\mu_3 = 0$, $\mu_4 = 0$ satisfies the equation (2.1).

In both cases, \bar{x} is a weak efficient solution of $(DTLZ1n2)$.

Using the aforementioned process with 200 initial points uniformly distributed from $(0,0)^T$ to $(1,1)^T$ for $(DTLZ1n2)$, and after eliminating the dominated points, the number of points in the Pareto front before tunneling is 8, while after tunneling it is 17.

Similarly, for the problem $(DTLZ3n2)$ with the same initial point distribution, the Pareto front contains 9 points before tunneling and 19 points after tunneling. The problem $(DTLZ3n2)$ is defined as follows:

$$(DTLZ3n2) : \min_{x \in \mathbb{R}^n} (f_1(x), f_2(x))$$

$$\text{s.t. } 0 \leq x_1, x_2 \leq 1$$

where,

$$f_1(x) = (1 + gx) \cos(0.5\pi x_1),$$

$$f_2(x) = (1 + gx) \sin(0.5\pi x_1),$$

$$\text{and } gx = 100[1 + (x_2 - 0.5)^2 - \cos\{20\pi(x_2 - 0.5)\}].$$

For both problems, the number of points in the Pareto front increases after tunneling,




Fig. 1: Weak efficient solutions of $(DTLZ1n2)$ for 200 random initial points

indicating a trend towards the global front. The weak efficient solutions of $(DTLZ1n2)$ and $(DTLZ3n2)$ are shown in Figures 1 and 2 respectively.

Similar to these test problems, Algorithm 1 is applied on a set of test problems sourced from different references. For details of these test problems, readers may see [18]. Each test problem is executed using $\eta = 1.2$ and a set of 200 uniformly distributed random initial points. Computational details are provided in Table 1. In this table, m , n , $PFBT$, and $PFAT$ represent the number of objective functions, the number of variables, number of approximate efficient solutions before tunneling, and number of




Fig. 2: Weak efficient solutions of $(DTLZ3n2)$ for 200 random initial points

approximate efficient solutions after tunneling respectively.

Form table 1, it is evident that the tunneling algorithm has a discernible impact on the set of efficient solutions. The comparison of the number of points in the pareto front (PF) before and after tunneling across various test problems reveals interesting insights into the algorithm's efficacy. One can observe that the tunneling algorithm demonstrates a notable increase in the number of points in the PF after its application, in most cases. This augmentation implies that the algorithm successfully enhances the diversity of the PF and potentially reveals previously undiscovered efficient solutions.

5 Conclusion

In this paper, we have developed a multi-objective tunneling method for nonlinear constrained multi-objective optimization problems. This method is free from any kind of priori chosen parameter or ordering information of objective functions. The tunneling algorithm showcases promising results in enriching the solution space and getting better efficient solutions for various nonlinear multi-objective optimization problems. Further investigation and experimentation could elucidate the algorithm's robustness and scalability across an even broader array of problem domains. Multi-start technique is used to generate approximate Pareto front. However, this technique fails in some cases. Some new spreading techniques similar to [18, 20] can be developed to generate well distributed approximate Pareto front. This is left as a slope of future research of this present contribution.

Acknowledment

Author Bikram Adhikary acknowledges funding from the *prime minister's research fellows (PMRF) scheme* (PMRF ID. 2202747), India.

Sl. No.	Test Problem	(m,n)	PFBT	PFAT
1	CEC09.3	(2,10)	4	6
2	Deb513	(2,2)	9	20
3	Deb521a.a	(2,2)	28	36
4	Deb521b	(2,2)	1	52
5	DTLZ1	(3,7)	51	58
6	DTLZ1n2	(2,2)	8	17
7	DTLZ2	(3,12)	27	31
8	DTLZ2n2	(2,2)	45	21
9	DTLZ3	(3,12)	53	33
10	DTLZ3n2	(2,2)	9	19
11	DTLZ4	(3,12)	24	25
12	DTLZ4n2	(2,2)	6	9
13	DTLZ5	(3,12)	1	2
14	DTLZ5n2	(2,2)	1	2
15	DTLZ6	(3,12)	6	7
16	DTLZ6n2	(2,2)	1	5
17	ex005	(2,2)	1	37
18	Far1	(2,2)	70	72
19	Fonseca	(2,2)	19	10
20	GE5	(3,3)	52	102
21	IKK1	(3,2)	1	17
22	IM1	(2,2)	3	59
23	Jin2.a	(2,10)	3	5
24	Jin3	(2,2)	4	17
25	Jin4.a	(2,2)	2	62
26	KW2	(2,2)	27	7
27	lovison2	(2,2)	87	65
28	lovison3	(2,2)	14	30
29	lovison4	(2,2)	52	78
30	lovison5	(3,3)	9	10
31	lovison6	(3,3)	12	15
32	LRS1	(2,2)	1	2
33	MHHM2	(3,2)	1	2
34	MLF2	(2,2)	34	47
35	MOP2	(2,2)	6	7
36	MOP3	(2,2)	25	15
37	MOP5	(3,2)	4	184
38	MOP6	(2,2)	8	7
39	MOP7	(3,2)	1	2
40	SK2	(2,4)	7	5
41	VFM1	(3,2)	1	25
42	ZDT1	(2,10)	5	6
43	ZDT2	(2,10)	6	9
44	ZDT3	(2,10)	10	15
45	ZDT6.a	(2,10)	7	16

Table 1: Comparison of (*PFBT*) and (*PFAT*) for various test problems.

References

- [1] T.M. Leschine, H. Wallenius, and W.A. Verdini. Interactive multiobjective analysis and assimilative capacity-based ocean disposal decisions. *European J. Oper. Res.*, 56(2):278–289, 1992.

- [2] J. Fliege. OLAF—a general modeling system to evaluate and optimize the location of an air polluting facility. *OR Spektrum*, 23:117–136, 2001.
- [3] G. Palermo, C. Silvano, S. Valsecchi, and V. Zaccaria. A system-level methodology for fast multi-objective design space exploration. In *Proceedings of the 13th ACM Great Lakes symposium on VLSI*, pages 92–95, 2003.
- [4] M. Tavana. A subjective assessment of alternative mission architectures for the human exploration of Mars at NASA using multicriteria decision making. *Comput. Oper. Res.*, 31(7):1147–1164, 2004.
- [5] S. Babaei, M.M. Sepehri, and E. Babaei. Multi-objective portfolio optimization considering the dependence structure of asset returns. *European J. Oper. Res.*, 244(2):525–539, 2015.
- [6] M. Gravel, J.M. Martel, R. Nadeau, W. Price, and R. Tremblay. A multicriterion view of optimal resource allocation in job-shop production. *European J. Oper. Res.*, 61(1-2):230–244, 1992.
- [7] P. De, J.B. Ghosh, and C.E. Wells. On the minimization of completion time variance with a bicriteria extension. *Oper. Res.*, 40(6):1148–1155, 1992.
- [8] A Hutterer and J Jahn. On the location of antennas for treatment planning in hyperthermia. *OR Spectrum*, 25:397–412, 2003.
- [9] K. Miettinen. *Nonlinear Multiobjective Optimization*. International series in operations research & management science. Springer, New York, Oct 2012.
- [10] M. Ehrgott. *Multicriteria Optimization*. Springer, Berlin, Germany, May 2005.
- [11] K. Deb. *Multi-Objective Optimization using Evolutionary Algorithms*. Wiley Interscience Series in Systems and Optimization. John Wiley & Sons, Chichester, England, May 2001.
- [12] M. Laumanns, L. Thiele, K. Deb, and E. Zitzler. Combining convergence and diversity in evolutionary multiobjective optimization. *Evol. Comput.*, 10(3):263–282, 2002.
- [13] S. Mostaghim, J. Branke, and H. Schmeck. Multi-objective particle swarm optimization on computer grids. In *Proceedings of the 9th annual conference on Genetic and evolutionary computation*, pages 869–875, 2007.
- [14] J. Fliege and B.F. Svaiter. Steepest descent methods for multicriteria optimization. *Math. Methods Oper. Res.*, 51:479–494, 2000.
- [15] J. Fliege, L.M.G. Drummond, and B.F. Svaiter. Newton’s method for multiobjective optimization. *SIAM J. Optim.*, 20(2):602–626, 2009.

- [16] M.A.T. Ansary and G. Panda. A modified quasi-Newton method for vector optimization problem. *Optimization*, 64(11):2289–2306, 2015.
- [17] S Qu, M Goh, and B Liang. Trust region methods for solving multiobjective optimisation. *Optim. Methods Softw.*, 28(4):796–811, 2012.
- [18] J. Fliege and A.I.F. Vaz. A method for constrained multiobjective optimization based on SQP techniques. *SIAM J. Optim.*, 26(4):2091–2119, 2016.
- [19] M.A.T. Ansary and G. Panda. A sequential quadratic programming method for constrained multi-objective optimization problems. *J. Appl. Math. Comput.*, 64(1):379–397, 2020.
- [20] M.A.T. Ansary and G. Panda. A globally convergent SQCQP method for multiobjective optimization problems. *SIAM J. Optim.*, 31(1):91–113, 2021.
- [21] G.C. Bento, J.X. Cruz Neto, and A. Soubeyran. A proximal point-type method for multicriteria optimization. *Set-Valued Var. Anal.*, 22:557–573, 2014.
- [22] H. Tanabe, E.H. Fukuda, and N. Yamashita. Proximal gradient methods for multiobjective optimization and their applications. *Comput. Optim. Appl.*, 72:339–361, 2019.
- [23] M.A.T. Ansary. A Newton-type proximal gradient method for nonlinear multi-objective optimization problems. *Optim. Methods Softw.*, 38(3):570–590, 2023.
- [24] M.A.T. Ansary and G. Panda. A sequential quadratically constrained quadratic programming technique for a multi-objective optimization problem. *Eng. Optim.*, 51(1):22–41, 2019.
- [25] A.V. Levy and A. Montalvo. The tunneling algorithm for the global minimization of functions. *SIAM J. Sci. Stat. Comput.*, 6(1):15–29, 1985.
- [26] C.K. Ng, D. Li, and L.S. Zhang. Global descent method for global optimization. *SIAM J. Optim.*, 20(6):3161–3184, 2010.
- [27] G.E. Rempu. A filled function method for finding a global minimizer of a function of several variables. *Math. Program.*, 46:191–204, 1990.
- [28] O.L. Mangasarian and S. Fromovitz. The Fritz John necessary optimality conditions in the presence of equality and inequality constraints. *J. Math. Anal. Appl.*, 17(1):37–47, 1967.