

Precision cosmology with exact inhomogeneous solutions of General Relativity: the Szekeres models.

Marie-Noëlle Célérier

*Laboratoire Univers et Théories, LUTH, Observatoire de Paris, Université PSL,
Université Paris Cité, CNRS, 5, Place Jules Janssen, 92190 Meudon, France**

(Dated: July 18 2024)

The exact Friedman solution to the field equations of General Relativity (GR) describing an homogeneous and isotropic universe together with its linear and higher-order perturbations are the theoretical roots of the current standard model of cosmology. However, despite its global successes, this standard model currently faces a number of tensions and anomalies, which occur mostly from a mismatch between early and late cosmic time region depictions. Actually, in the era of precision cosmology, the late inhomogeneities in the densities of the Universe components can no more be neglected. Moreover, since GR is fundamentally nonlinear, any linear perturbation theory is doomed to fail at reproducing precisely its features, while higher-order perturbed FLRW models cannot either claim the status of exact solution to Einstein's field equations. Fortunately, other GR solutions exist which are best suited for this purpose, e. g., exact inhomogeneous solutions able to represent a matter-cosmological constant dominated universe region. Among these, the Szekeres solution, which is devoid of any symmetry, appears as a proper tool for this purpose. Moreover, since these solutions possess the FLRW model as a homogeneous limit, they can be smoothly matched to the standard representation at the inhomogeneity-homogeneity transition. In this paper, the Szekeres solution and its main interesting properties, as well as the equations needed to use this solution in a cosmological context are presented. Then the prospects for a broader use of its abilities are sketched out. In particular, the use of neural networks is proposed to allow, in the future, the fitting of the huge amount of data becoming available to constrain the arbitrary functions and the constant parameters characterizing the Szekeres model, running for representing our late Universe with an increased precision.

I. INTRODUCTION

The exact Friedman solution to the field equations of General Relativity (GR) describing an homogeneous and isotropic universe equipped with its linear and higher-order perturbations are the theoretical roots of the current standard model of cosmology. However, besides its global successes and the progresses it has provided in our understanding of the Universe, this standard paradigm faces presently a number of tensions and misunderstandings [1] which hint at a need for a more accurate theoretical background.

Two main options are therefore possible. Either one dismisses GR and explores modified gravity theories, while taking care of reproducing all GR successful predictions such as dynamics in the solar system, gravitational waves, lensing, redshift, etc. Or one goes on within the GR framework and considers another exact solution able to satisfy the constraints implied by the full range of cosmological data. This latter option is adopted here.

Since the tensions and anomalies currently encountered seem to occur mostly from a mismatch between early and late cosmic time region depictions, we will consider a cosmological model where the inhomogeneities in the densities of the Universe components are intrinsic. The Szekeres solution which describes a dust dominated

spacetime devoid of symmetries appears as a widely general and accurate tool for our purpose. Moreover, this solution exhibits the FLRW model as a homogeneous limit. It seems therefore fairly well adapted to represent our Universe, since all the best early time predictions of the standard model can be preserved in its framework.

The Szekeres equations are, of course, more involved than the FLRW ones. Anyhow, the interesting physical properties of the model have lead a number of authors to try to overcome these complications to benefit from such properties. However, up to now, the method used has consisted mainly in assuming a priori simplified parameters and functions describing a given Szekeres model, designed in order to solve a particular cosmological problem. Then the observable quantities attached to this particular model were calculated and compared to those actually observed. Most of the works were numerically completed.

A few authors considered what they called finding “the metric of the Universe” [2, 3]. However, at the time such a proposal was done, the numerical calculation power needed to perform this plan was too low and these authors were limited in their legitimate ambition. Even the use of the simpler Lemaitre-Tolman-Bondi (LTB) models proved unable to reach the goal.

Now, we can benefit from the progresses realized in artificial intelligence and deep learning, to consider again such a program. In the present paper, we display a comprehensive set of observables calculated assuming a Szekeres-type Universe, i. e., using the Szekeres solution to represent the late Universe and its FLRW homoge-

*Electronic address: marie-noelle.celerier@obspm.fr

neous limit at larger scales. Then we show how these observables could be constrained by the cosmological data currently available and by those to be collected in a near future.

The paper is organised as follows. In Sec.II, we present the Szekeres solutions with their main physical properties and, in particular, their null geodesic, redshift and distance equations. The constraints from currently or soon available cosmological observations are discussed in Sec.III. Some effects observable in inhomogeneous models only are analyzed in Sec.IV. The use of deep learning for processing the data fitting is presented in Sec.V. Section VI is devoted to the conclusions. Finally, the Szekeres curvature symbols and tensor components needed for the calculations are displayed in Appendix A.

II. THE SZEKERES SOLUTIONS

An exact solution to Einstein's field equations for a dust gravitationally sourced spacetime with no symmetry has been found by Szekeres in 1975 [4]. Since then, this solution has been (too scarcely) considered in a cosmological context and part of its features has been analyzed. Here, we recall, precise, generalize, complete, and correct when needed, the main expressions and properties which will be used to build a consistent cosmological work program. Adopting a convenient coordinate system [5] the line element in synchronous and comoving coordinates, and in geometric units, can be written as

$$ds^2 = -dt^2 + \frac{(\Phi_{,r} - \Phi E_{,r}/E)^2}{\epsilon - k} dr^2 + \frac{\Phi^2}{E^2} (dp^2 + dq^2), \quad (1)$$

where

$$E(r, p, q) = \frac{S}{2} \left[\left(\frac{p-P}{S} \right)^2 + \left(\frac{q-Q}{S} \right)^2 + \epsilon \right], \quad (2)$$

with k , S , P , and Q being functions of the r coordinate alone. The parameter ϵ determines whether the (p, q) 2-surfaces of constant $\{r, t\}$ are unit 2-spheres ($\epsilon = +1$), unit 2-pseudo-spheres, i.e., hyperboloids ($\epsilon = -1$), or 2-planes ($\epsilon = 0$).

The quasi-spherical Szekeres (QSS) solutions ($\epsilon = +1$) can be thought of as a set of non-concentric evolving spheres, which exhibit a dipole distribution in the energy density variation around each sphere. The dipole arises from the $E_{,r}/E$ term. Actually, on each (p, q) 2-spheres, $E_{,r}/E$ vanishes on an "equator", is maximal at one "pole" and minimal at the other "pole", with $E_{,r}/E|_{min} = -E_{,r}/E|_{max}$.

The quasi-hyperbolic solutions ($\epsilon = -1$) can be described as a set of evolving right-hyperboloids piled "non-concentrically". In this case also, the hyperboloids display a "pseudo-spherical" dipole whose strength and orientation on each comoving shell is determined by $S(r)$,

$P(r)$, and $Q(r)$. However, half of the dipole is located in the hyperboloid sheet which is not free of shell crossings [6]. Therefore, if a shell-crossing should occur in a region spanned by the cosmological solution selected by the observational data, this might constitute a severe physical drawback. Hence, after fitting the data, a key verification will be to test whether the achieved model is actually free of shell crossings. The conditions for no shell-crossing have been displayed by Hellaby and Krasiński for spherical foliation ($\epsilon = +1$) [9] and for pseudospherical ($\epsilon = -1$) and planar ($\epsilon = 0$) foliations [6], and they will have to be checked.

The quasi-planar Szekeres solutions ($\epsilon = 0$) do not have any $E_{,r}/E$ extremum. Therefore, it does not exhibit any dipole-like structure.

Notice that, even though the quasi-spherical family has been mostly employed up to now in a cosmological context since it is the best understood case, the surfaces $r = \text{const}$ within a space $t = \text{const}$ might exhibit different geometries such as being quasi-spherical in regions of the space and quasi-hyperbolic elsewhere, with a zero curvature at the boundaries [6]. This planar boundary can either be a thin 3-surface or a 2-surface with finite width. If thin, its lack of dipole would not impair the dipole feature of spacetime as a whole, since it would only concern the instantaneous transition from quasi-spherical to quasi-hyperbolic dipole-like matter distributions.

More thorough descriptions of the mathematical and physical properties of these solutions can be found in textbooks [7, 8] and in a bunch of theoretical papers [5, 6, 9–12]. We give below a summary of the properties which will be needed for the present study.

The Einstein field equations written with a dust source and a cosmological constant reduce to the following two:

$$\Phi_{,t}^2 = \frac{2M}{\Phi} - k + \frac{\Lambda}{3}\Phi^2, \quad (3)$$

$$4\pi\rho(t, r, p, q) = \frac{M_{,r} - 3ME_{,r}/E}{\Phi^2(\Phi_{,r} - \Phi E_{,r}/E)}, \quad (4)$$

where ρ is the matter energy density, $M(r)$ is another function of the r coordinate which, in the quasi-spherical case where $\epsilon = +1$, represents the total active gravitational mass in a shell determined by $r = \text{const}$.

Eq. (3) can be integrated as

$$t - t_B(r) = \int_0^\Phi \frac{d\tilde{\Phi}}{\sqrt{\frac{2M}{\tilde{\Phi}} - k + \frac{\Lambda}{3}\tilde{\Phi}^2}}, \quad (5)$$

where t_B is an arbitrary function of r which represents the Big Bang or Crunch time.

A given Szekeres model is therefore a priori determined by six functions of r : $k(r)$, $S(r)$, $P(r)$, $Q(r)$, $M(r)$, and $t_B(r)$, one quasi-constant parameter ϵ and a cosmological

constant Λ ¹. However, the number of independent functions of r can be reduced to five by using the coordinate freedom to rescale r , e. g., through a choice of any one among the arbitrary functions.

To construct a Szekeres model able to reproduce a sufficiently large set of cosmological data, we need to express the observable quantities with the tools available to describe the Szekeres solutions. This is the main purpose of the present paper.

A. Null geodesic and redshift equations

The data obtained by cosmological surveys currently available and those which will be displayed in a near future are and will be mainly obtained by measuring electromagnetic signals from astrophysical objects located at some distance from the observer². To process them in view of constraining a cosmological model, the trajectories of the photons emitted by the source and measured by the observer in this model, as well as a method for calculating the redshift, need to be known. We give below

¹ The term quasi-constant employed here means that ϵ which can only have three different possible values can change this value according to the following mechanism: a quasi-spherical and a quasi-hyperbolic Szekeres region of spacetime whose constant r sphere radius, constant r hyperboloid radius respectively, diverges at some r value, can be joined at their planar boundaries, provided junctions conditions are verified [6]. Therefore, even

the main equations describing these features in the most general Szekeres model. The expressions for the geometrical quantities used for establishing these equations are given in Appendix A.

In the geometric optics approximation, based on the assumption that light wave-lengths are negligible with respect to the space curvature scales, which is a reasonable assumption in a cosmological framework, photons travel along null geodesics, and the light rays have no effect on the geometry, i.e., they are test objects. The tangent vector to the null geodesics, k^α , is given by the gradient of the wave phase and is identified with the photon momentum. It satisfies $k^\alpha k_{;\beta}^\alpha = 0$ and $k^\alpha k_\alpha = 0$. The first equation is the null geodesic equation which can be written, in an affine parametrization as

$$\frac{d^2 x^\alpha}{ds^2} + \Gamma_{\beta\gamma}^\alpha \frac{dx^\beta}{ds} \frac{dx^\gamma}{ds} = 0. \quad (6)$$

Inserting into (6) the non-zero values of the Christoffel symbols displayed in Appendix A, we obtain the four null geodesic equations as [13, 14]

though the values of ϵ are limited to a set of three numbers, its actually realized value depends on that of the r coordinate. The parameter ϵ can therefore be viewed as a step function of r

² Of course, gravitational waves are intended to become a key tool in cosmology, but their consideration is premature and therefore out of the scope of the present study.

$$\frac{d^2 t}{ds^2} + \left(\frac{\Phi_{,tr} - \Phi_{,t} E_{,r}/E}{\epsilon - k} \right) (\Phi_{,r} - \Phi E_{,r}/E) \left(\frac{dr}{ds} \right)^2 + \frac{\Phi \Phi_{,t}}{E^2} \left[\left(\frac{dp}{ds} \right)^2 + \left(\frac{dq}{ds} \right)^2 \right] = 0, \quad (7)$$

$$\begin{aligned} \frac{d^2 r}{ds^2} + 2 \left(\frac{\Phi_{,tr} - \Phi_{,t} E_{,r}/E}{\Phi_{,r} - \Phi E_{,r}/E} \right) \frac{dt}{ds} \frac{dr}{ds} + \left(\frac{\Phi_{,rr} - \Phi E_{,rr}/E}{\Phi_{,r} - \Phi E_{,r}/E} - \frac{E_{,r}}{E} + \frac{k_{,r}}{2(\epsilon - k)} \right) \left(\frac{dr}{ds} \right)^2 \\ + 2 \frac{\Phi}{E^2} \left(\frac{E_{,r} E_{,p} - E E_{,rp}}{\Phi_{,r} - \Phi E_{,r}/E} \right) \frac{dr}{ds} \frac{dp}{ds} + 2 \frac{\Phi}{E^2} \left(\frac{E_{,r} E_{,q} - E E_{,rq}}{\Phi_{,r} - \Phi E_{,r}/E} \right) \frac{dr}{ds} \frac{dq}{ds} \\ - \frac{\Phi}{E^2} \left(\frac{\epsilon - k}{\Phi_{,r} - \Phi E_{,r}/E} \right) \left[\left(\frac{dp}{ds} \right)^2 + \left(\frac{dq}{ds} \right)^2 \right] = 0, \end{aligned} \quad (8)$$

$$\begin{aligned} \frac{d^2 p}{ds^2} + 2 \frac{\Phi_{,t}}{\Phi} \frac{dt}{ds} \frac{dp}{ds} - \frac{\Phi_{,r} - \Phi E_{,r}/E}{\Phi(\epsilon - k)} (E_{,r} E_{,p} - E E_{,rp}) \left(\frac{dr}{ds} \right)^2 \\ + 2 \frac{\Phi_{,r} - \Phi E_{,r}/E}{\Phi} \frac{dr}{ds} \frac{dp}{ds} - 2 \frac{E_{,q}}{E} \frac{dp}{ds} \frac{dq}{ds} + \frac{E_{,p}}{E} \left[- \left(\frac{dp}{ds} \right)^2 + \left(\frac{dq}{ds} \right)^2 \right] = 0, \end{aligned} \quad (9)$$

$$\begin{aligned} \frac{d^2 q}{ds^2} + 2 \frac{\Phi_{,t}}{\Phi} \frac{dt}{ds} \frac{dq}{ds} - \frac{\Phi_{,r} - \Phi E_{,r}/E}{\Phi(\epsilon - k)} (E_{,r} E_{,q} - E E_{,rq}) \left(\frac{dr}{ds} \right)^2 \\ + 2 \frac{\Phi_{,r} - \Phi E_{,r}/E}{\Phi} \frac{dr}{ds} \frac{dq}{ds} - 2 \frac{E_{,p}}{E} \frac{dp}{ds} \frac{dq}{ds} + \frac{E_{,q}}{E} \left[\left(\frac{dp}{ds} \right)^2 - \left(\frac{dq}{ds} \right)^2 \right] = 0. \end{aligned} \quad (10)$$

In addition, we have the first integral of the geodesics, which reads

$$\left(\frac{dt}{ds} \right)^2 = \frac{(\Phi_{,r} - \Phi E_{,r}/E)^2}{\epsilon - k} \left(\frac{dr}{ds} \right)^2 + \frac{\Phi^2}{E^2} \left[\left(\frac{dp}{ds} \right)^2 + \left(\frac{dq}{ds} \right)^2 \right]. \quad (11)$$

Now, two methods can be used to calculate the redshift. First, we can solve the four second order or-

dinary differential null geodesic equations (7)-(10) for $\{t(s), r(s), p(s), q(s)\}$ where the coefficients are expressed with the metric functions which must be evaluated on the null cones while satisfying the field equations (3) and (4). After integration, we can calculate the tangent vector four components $\{k^t, k^r, k^p, k^q\}$, among which k^t will be used to obtain the redshift z , given by

$$1 + z = \frac{k_e^t}{k_o^t}, \quad (12)$$

where the subscripts e and o denote values taken at the emitter and at the observer, respectively. Since we have chosen to use comoving synchronous coordinates, the four-velocity of the matter fluid is $u_\alpha = (1, 0, 0, 0)$. The null geodesic tangent vector at the observer can be normalized so that $k_o^t = -1$, which implies

$$1 + z = -k_e^t. \quad (13)$$

The second method is inspired from that applied to the spherically symmetric Lemaître-Tolman-Bondi (LTB) model by H. Bondi [15] and is adapted here to the Szekeres model. A previous attempt restricted to the case of quasi-spherical symmetry can be consulted [16]. We give below the calculation adapted to the most general Szekeres model.

$$\frac{dt}{ds} \frac{dT}{ds} = T(s) \left\{ \left[\frac{\Phi_{,tr} \Phi_{,r} + \Phi \Phi_{,t} (E_{,r}/E)^2 - (\Phi_{,t} \Phi_{,r} + \Phi \Phi_{,tr}) (E_{,r}/E)}{\epsilon - k} \right] \left(\frac{dr}{ds} \right)^2 + \frac{\Phi \Phi_{,t}}{E^2} \left[\left(\frac{dp}{ds} \right)^2 + \left(\frac{dq}{ds} \right)^2 \right] \right\}. \quad (17)$$

Then, we write the redshift as

$$1 + z(s_e) = \frac{T(s_o)}{T(s_e)}. \quad (18)$$

Taking the derivative of (18) with respect to s_e , we get

$$\frac{dz}{ds_e} = -\frac{dT(s_e)}{ds_e} \frac{1 + z(s_e)}{T(s_e)}. \quad (19)$$

Then, we substitute (19) into (18) while dropping the subscripts for better readability and we obtain

$$\frac{d(\ln(1 + z))}{ds} = -\frac{1}{\frac{dT}{ds}} \left\{ \left[\frac{\Phi_{,tr} \Phi_{,r} + \Phi \Phi_{,t} (E_{,r}/E)^2 - (\Phi_{,t} \Phi_{,r} + \Phi \Phi_{,tr}) (E_{,r}/E)}{\epsilon - k} \right] \left(\frac{dr}{ds} \right)^2 + \frac{\Phi \Phi_{,t}}{E^2} \left[\left(\frac{dp}{ds} \right)^2 + \left(\frac{dq}{ds} \right)^2 \right] \right\}. \quad (20)$$

After solving the null geodesic equations, with the two field equations satisfied, we have all the results needed to obtain the redshift from (20).

B. Distance equations

Consider a congruence of light rays with surface area δS emitted by a source S and received by a point observer at O . Assume that the δS sheet at S is orthogonal to the central ray of the bundle emitted from S . This bundle encloses a solid angle $\delta\Omega$ at O . The image propagation

along the light rays is characterized by optical kinematic quantities defined in the observer's screen space, which is the two-dimensional space in the rest frame of the matter four-velocity u^α orthogonal to k^α . Defining the propagation vector k_α as $k_\alpha = \psi_{,\alpha}$, where ψ is the phase of the

$$\left(\frac{d(t+T)}{ds} \right)^2 = \frac{[\Phi_{,r}(t+T, r) - \Phi(t+T, r)E_{,r}/E]^2}{\epsilon - k} \left(\frac{dr}{ds} \right)^2 + \frac{\Phi^2(t+T, r)}{E^2} \left[\left(\frac{dp}{ds} \right)^2 + \left(\frac{dq}{ds} \right)^2 \right]. \quad (14)$$

Assuming that the period T of the light wave is negligible with respect to the time over which the light propagates - which is an assumption equivalent to the geometric optics approximation - and following Bondi [15], we Taylor expand Φ and $\Phi_{,r}$ to first order as

$$\Phi(t+T, r) = \Phi(t, r) + \Phi_{,t}(t, r)T, \quad (15)$$

$$\Phi_{,r}(t+T, r) = \Phi_{,r}(t, r) + \Phi_{,tr}(t, r)T. \quad (16)$$

By substituting (15) and (16) into (11) and (14), then subtracting (11) from (14), we obtain

light-wave, the light-rays have no vorticity. Furthermore, being geodesics, the light rays experience no acceleration. Only the null expansion and the null shear are non-zero. The null expansion θ , measures the area expansion rate of images. It is given, in terms of the affinely parametrized tangent vector by

$$\theta = k_{;\alpha}^{\alpha}. \quad (21)$$

The rate of change of the observer area distance, usually simply named area distance and denoted D_A , depends on the expansion rate of the light bundle as

$$\frac{d(\ln D_A)}{d\lambda} = \theta = k_{;\alpha}^{\alpha}. \quad (22)$$

Now, we insert into (22) the expression for $k_{;\alpha}^{\alpha}$ obtained with the use of the Christoffel symbols displayed in Appendix A. Then, we apply the total derivative expression to the result and integrate which gives

$$D_A = \frac{\Phi^2}{E^2} \left(\frac{\Phi_{,r} - \Phi E_{,r}/E}{\sqrt{\epsilon - k}} \right) \times \exp \left[\int_{s_o}^s (k_{,t}^t + k_{,r}^r + k_{,p}^p + k_{,q}^q) d\lambda \right], \quad (23)$$

where s_o is the value of the affine parameter at the observer, which can be set to zero.

Thus, the area distance is obtained from (23), where the k^α and their derivatives $k_{;\alpha}^{\alpha}$ emerge from (7)-(11).

Another means of obtaining an expression for the area distance is to use the Sachs optical equations for the null bundle expansion θ and its shear σ which read [17]

$$\frac{d\theta}{d\lambda} + \theta^2 + |\sigma|^2 = -\frac{1}{2} R_{\alpha\beta} k^\alpha k^\beta, \quad (24)$$

$$\frac{d\sigma_{\alpha\beta}}{d\lambda} + \theta\sigma_{\alpha\beta} = -C_{\alpha\beta\gamma\delta} k^\gamma k^\delta, \quad (25)$$

where $R_{\alpha\beta}$ is the Ricci tensor and $C_{\alpha\beta\gamma\delta}$ is the Weyl tensor.

By using (22) into (24), we obtain

$$\frac{d^2 D_A}{d\lambda^2} = - \left(|\sigma|^2 + \frac{1}{2} R_{\alpha\beta} k^\alpha k^\beta \right) D_A. \quad (26)$$

Thus, the area distance can be obtained from solving (26) together with (24), (25) and the null geodesic equations written for the k^α s, using the Ricci and Weyl tensor components displayed in Appendix A.

Finally, assuming that the photon number is conserved and that photons travel on unique null geodesics, and since we work in the framework of a metric theory of gravity, we can apply the reciprocity theorem to obtain the value of the luminosity distance of a given source at redshift z , D_L , as [18]

$$D_L = (1+z)^2 D_A. \quad (27)$$

III. CONSTRAINTS FROM CURRENTLY AVAILABLE COSMOLOGICAL OBSERVATIONS

A. Supernova data

Type Ia supernovae (SNIa) are used in cosmology as standard candles. The apparent bolometric magnitude, m , of a standard candle with absolute bolometric magnitude, M , at a given redshift, z , is a function of z and of the functions and parameters determining the cosmological model. It can be written as [19]

$$m = \mathcal{M} + 5 \log D_L(z; p_a), \quad (28)$$

where \mathcal{M} is the magnitude zero point, which can be measured from the apparent magnitude and redshift of low-redshift examples of the standard candles, and p_a represents the set of model parameters.

To constrain these parameters with SNIa data, one compares the apparent magnitudes of distant supernovae to local SNIa calibrators whose distance is evaluated through, e. g., measurements of Cepheid variables present in these SN Ia hosts. The set of apparent magnitude and redshift measurements for low-redshift SNIa is used to calibrate (28), and another set of such measurements at higher redshift is used to determine p_a , through (23), (26), and (27).

In the Szekeres case, the parameters of the model are included in the expression for D_L obtained through (27), where the redshift calculation is achieved by applying the method displayed in Sec.II A.

B. Galaxy number counts

Here, we adapt to the Szekeres models a reasoning formerly designed for LTB spacetimes [20]. Denoting $n(z)$ the number density of sources in redshift space per steradian and unit redshift interval, the number of sources observed in a given redshift interval dz and solid angle $d\Omega$ is $nd\Omega dz$ and the total rest mass between z and dz is

$$\mathcal{M} = 4\pi m n dz, \quad (29)$$

where $m(z)$ is the average mass per source, assumed to be known from astrophysical considerations.

The total rest mass in a volume element d^3V , defined as the proper volume on a constant time slice, evaluated on the null cone, can be written as

$$\mathcal{M} = \rho d^3V, \quad (30)$$

which reads, in a Szekeres framework,

$$\mathcal{M} = 4\pi\rho \frac{\Phi^2}{E^2} \left(\frac{\Phi_{,r} - \Phi E_{,r}/E}{\sqrt{\epsilon - k}} \right) dr. \quad (31)$$

In the Szekeres solutions, the energy density ρ is given by (4). Inserting the corresponding expression into (31),

we obtain

$$4\pi mn \frac{dz}{dr} = \frac{M_{,r} - 3ME_{,r}/E}{E^2\sqrt{\epsilon - k}}, \quad (32)$$

which gives the average mass density in redshift space $m(z)n(z)$. Since the redshift z can be obtained by the algorithm described in Sec.II A, the mass density in real space follows and can be compared to the measured data.

C. Cosmic microwave background (CMB)

1. CMB multipoles

The CMB temperature observed at each point in the sky can be calculated by generating, at the observer, a null geodesic reaching backwards in time the last scattering surface with an initial tangent vector pointing in the direction of the given point in the sky. This is done by integrating the geodesic equations (7)-(10). The redshift, which determines the last scattering location on the geodesics (defined by its redshift z_{ls}) hence, the CMB temperature seen along any line of sight, will then be obtained through (20) [21].

For an accurate calculation of the observed CMB multipoles over the whole sky, geodesics going in every direction will have to be integrated. Then the multipoles will be obtained by computing the coefficients of the spherical harmonic expansion

$$a_{lm} = \int_0^{2\pi} \int_0^\pi \frac{\Delta T}{T} Y_{lm}(\theta, \phi) \sin \theta d\theta d\phi. \quad (33)$$

In particular the CMB dipole will be obtained as

$$D = \sqrt{\sum_{m=-1}^1 |a_{1m}|^2}. \quad (34)$$

Now, in the quasi-spherical, as well as in the quasi-hyperbolic Szekeres models, the distribution of mass over each single sphere $\{t = \text{const}, r = \text{const}\}$ has the form of a mass-dipole superposed on a monopole [7, 22, 23]. This is not the case of the quasi-planar models, but since, in spacetimes where they might be present, such quasi-planar models can be assumed as infinitely thin, their absence of contribution to a dipole can be ignored. The Szekeres dipole-like contribution to matter density, is therefore given by [8]

$$\begin{aligned} \kappa \Delta \rho &= \frac{\chi_{,r} - \chi E_{,r}/E}{\Phi^2(\Phi_{,r} - \Phi E_{,r}/E)} \\ &\times \frac{6M\Phi_{,r} - 2M_{,r}\Phi}{\Phi^2(\Phi_{,r}\chi - \Phi\chi_{,r})}, \end{aligned} \quad (35)$$

where

$$\chi = \frac{1 + P^2 + Q^2}{2S} + \frac{S}{2}. \quad (36)$$

One could therefore assume that (part of) the somehow large dipole measured in the CMB data should be due to the Szekeres geometry of the intervening spacetime region, and use it to constrain the Szekeres parameter functions implied in (35).

Now, the dipole pertaining to the Szekeres model is an intrinsic dipole. To compare it with the CMB's, we might have to subtract the kinetic dipole due to our motion with respect to the CMB rest frame. However, the kinematic interpretation of the CMB dipole has been questioned by some survey analyses with high statistical confidence [24]. This might compromise, at least tentatively, the use of the CMB dipole as a proper tool for constraining the Szekeres model. However, an unbiased estimator for the kinematic dipole amplitude, proportional to the peculiar velocity of the observer, has been recently proposed [25], and might be used in a near future to estimate the kinetic part of the CMB dipole.

Now, imagine that, in the future, the intrinsic part of the CMB dipole should be properly identified, as well as the intrinsic values of the other low- l multipoles, e. g., from physics at last scattering, then the Szekeres determining functions and parameters would be only constrained from the CMB data through the geometry of the null geodesics followed by the microwave rays between last scattering and the observer.

However, in the meanwhile, it might become possible to constrain our cosmological model without taking the CMB data into account. Once the proper model is fixed by other data sets, we would be able to use (35) to deduce the intrinsic dipole and therefore to discriminate between the kinetic and intrinsic dipole, as suggested in Sec.III F. Finally, the amplitude and direction of the observer's tilt with respect to the Hubble bulk flow would be properly identified.

2. Sunyaev Zeldovitch effects

The Sunyaev Zeldovitch effect (SZE) is the added spectral distortion of the CMB through inverse Compton scattering by high-energy electrons in galaxy clusters, in which the low-energy photons receive an energy boost during their collision with the high-energy cluster electrons. [26] The SZE can be divided into three different types: the thermal, the kinetic, and the polarization effects.

The thermal Sunyaev Zeldovitch effect (TSZE) denotes the effect arising from the interaction of the CMB photons with electrons which owe their high-energy to their temperature. The TSZE spectral distortion of the CMB expressed as a temperature change ΔT_{TSZ} at dimensionless frequency $x = h\nu/(K_B T_{CMB})$ is given by

$$\frac{\Delta T_{TSZ}}{T_{CMB}} = f(x)y = f(x) \int n_e \frac{k_B T_e}{m_e c^2} \sigma_T dl, \quad (37)$$

where T_{CMB} is the CMB average temperature, y is the Compton y -parameter, which, for an isothermal clus-

ter, equals the optical depth times the fractional energy gain per scattering, σ_T is the Thomson scattering cross-section, n_e is the electron number density, T_e is the electron temperature, k_B is the Boltzmann constant, $m_e c^2$ is the electron rest mass energy, and the integration is along the line-of-sight. As can be seen in (37), $\Delta T_{TSZ}/T_{CMB}$ is independent of redshift. This unique feature of the TSZE makes it a powerful tool for investigating the high-redshift Universe.

Moreover, it has been shown [27] that the area diameter distance can be obtained from the normalizations N_{RJ} of the TSZE and N_X of the surface brightness distribution through

$$D_A = \frac{N_{RJ}^2}{N_X} \left(\frac{m_e c^2}{k_B T_{eo}} \right)^2 \frac{\Lambda_{eo}}{16\pi T_{CMB}^2 \sigma_T^2 (1+z)^3}, \quad (38)$$

where Λ_{eo} is the X-ray spectral emissivity and z is the redshift of the cluster gas. Equation (38) can be used to compare the expression of D_A in a Szekeres cosmology, obtained as described in Sec.II B, to its measured values as obtained through the right hand-side of (38).

The main features of the TSZE are: (i) it is a small distortion of the CMB, proportional to the cluster pressure integrated along the line-of-sight, see (37), (ii) it is redshift-independent, (iii) it possesses a unique spectral signature with a decrease in the CMB intensity at low frequencies and an increase at high frequencies, (iv) the integrated TSZE is proportional to the temperature-weighted mass (the total thermal energy) of the cluster, and mostly interesting for our purpose, (v) through (38) it allows us to determine the cluster area distance in a cosmology independent way.

The kinetic Sunyaev Zeldovitch effect (KSZE) denotes the Doppler effect on the scattered photons due to the cluster bulk velocity owing to its motion with respect to the CMB rest frame. If a component of the cluster velocity is projected along the line-of-sight from the observer to the cluster, then the Doppler effect yields an observable distortion of the CMB spectrum. It gives a method for calculating the peculiar velocity, valid in the non-relativistic limit. However, we have not found a proper way to use it in a Szekeres framework.

As regards the SZ polarisation effect on the CMB photons, it is predicted as too small for being measurable in a near future.

D. Baryon acoustic oscillations (BAO)

Before last scattering, the highly coupled photon-baryon plasma oscillates under the competitive effects of gravitational collapse and radiation pressure. The resultant acoustic waves travel in the plasma at the sound speed

$$c_s(z) = \frac{1}{\sqrt{3 \left(1 + \frac{3\rho_b(z)}{4\rho_\gamma(z)} \right)}}, \quad (39)$$

where ρ_b and ρ_γ are the baryon and radiation energy densities, respectively. At baryon decoupling, i. e., end of the drag epoch, the pressure on the baryons disappears and the baryon acoustic wave is frozen in, while the photons stream freely. An enhanced baryon over-density is thus formed at the distance travelled by the sound wave from the observer to the decoupling. This implies a baryon acoustic oscillation (BAO) peak in the galaxy correlation function and wiggles (oscillatory features) in their power spectrum [28, 29].

This feature being initiated in the early universe, it is not especially subject to initial modifications by the late time inhomogeneities. The only effect such inhomogeneities have on BAO measurements is a bending of the light rays used to establish the galaxy correlation function or power spectrum. The simplest model-independent BAO measurements use the angular 2-point correlation function which involves only the angular separation θ between galaxy pairs. Using thin enough redshift bins, one measures the angular BAO scale given by

$$\theta_{BAO}(z) = \frac{r_s}{(1+z)D_A(z)}, \quad (40)$$

where r_s is the sound horizon at the end of the drag epoch, which can be written as

$$r_s = \int_{z_d}^{\infty} c_s(1+z)dz. \quad (41)$$

Since it is large, around 130 Mpc comoving, the acoustic scale is usually assumed to be conserved along its travel through the late universe inhomogeneities. It is therefore considered as a proper standard ruler. We propose to assume temporarily that this property can be retained in a Szekeres framework.

The outcomes of a typical galaxy survey such as DESI [30] are displayed under the form of, either a couple of quantities $\{D_M/r_s, D_H/r_s\}$, where the comoving distances $D_M(z)$ and $D_H(z)$ are defined by $D_M(z) = r_s/\Delta\theta$ and $D_H(z) = r_s/\Delta z$, respectively, or by a single quantity D_V/r_s , where the angle-averaged distance D_V is defined by

$$D_V(z) = [zD_M^2(z)D_H(z)]^{1/3}, \quad (42)$$

which is measured in samples with low signal-to-noise, such as quasar surveys. We see that D_M is the comoving distance related to the area distance by the reciprocity theorem which yields: $D_M = (1+z)D_A$.

We have described in Sec.II A and IIB methods to compute the redshift and the area distance in a Szekeres framework. The value of r_s is independent of the redshift at which it is measured. Therefore, a set of measurements of the angular BAO scale at different redshifts will yield constraints on the Szekeres functions and parameters from the $(1+z)D_A(z)$ factor in the right hand side of (40).

Notice, however, that small acoustic scale changes of order some tenth of percent, have been identified from

second order effects in perturbation theory and N-body simulations [31]. These should be taken into account when addressing the most precise stages of cosmological modelling. The primary difficulty proceeds from the anisotropic distortions issuing from the galaxy peculiar velocity, see Sec.III F. The tackling of this issue in the BAO framework is however beyond the scope of the present paper and is left to future work.

E. Weak lensing

1. Generalities

In curved spacetimes, Maxwell's equations cannot be solved explicitly, save in cases of high symmetry. In a Szekeres model with no symmetry at all, plane waves do not exist.

However, in the short-wave (WKB) [32] and gravitational lens [33] approximations, we can consider a locally plane wave, propagating without interaction with matter, and associated with a hypersurface S , orthogonal to a congruence of null geodesics representing light-rays. We spot one fiducial (central) light-ray γ_o and consider the geodesic deviation of adjacent light-rays γ in the bundle relative to γ_o . This fiducial ray is described by the null tangent vector field $k^\mu = \partial x^\mu / \partial \lambda$, since it is assumed to be parametrized by an affine parameter λ . For a past-directed light cone, $\lambda = 0$ at the observer and increases while going back in time. We can write, at the observer,

$$(u^\mu u_\mu)_o = -1, \quad (43)$$

$$u_o^\mu k_\mu = 1, \quad (44)$$

where u^μ is the four-velocity of the matter fluid.

The light bundle is propagated from the observer along the fiducial ray and the deviation of nearby rays relative to this fiducial ray represent the lensing properties of any intervening mass. Adopting in the following the reasoning displayed by Troxel et al. [34], we define a deviation vector field or Jacobi field Y^μ as

$$Y^\mu(\vec{\vartheta}, \lambda) = \gamma^\mu(\vartheta, \lambda) - \gamma_o^\mu(\vartheta = 0, \lambda), \quad (45)$$

such that Y represents the changing separation between some ray γ in the bundle and the fiducial ray γ_o , from which the relative angular position is ϑ . The four-vector Y^μ is decomposed on a space-like basis $\{E_1, E_2\}$ which spans the plane orthogonal to both u^μ and k^μ in order to complete the tetrad along γ_o , u^μ resulting from parallelly propagating u_o^μ along γ_o . We can therefore rewrite Y as

$$Y^\mu = \xi_1 E_1^\mu + \xi_2 E_2^\mu + \xi_0 k^\mu, \quad (46)$$

where the screen components $\xi_i (i = 1, 2)$ change according to the deformation equation

$$\dot{\xi}_i = S_{ij} \xi_j, \quad S_{ij} = E_i^\alpha k_{\alpha;\beta} E_j^\beta, \quad (47)$$

where a dot denotes differentiation with respect to the affine parameter λ . In matrix form, (47) becomes

$$\dot{\xi} = \mathcal{S} \xi. \quad (48)$$

The optical deformation matrix \mathcal{S} is composed of the Sach's optical scalars of the beam [17], i. e., its expansion rate

$$\theta(\lambda) = k_{;\mu}^\mu, \quad (49)$$

and its complex shear rate

$$\sigma(\lambda) = \frac{1}{2} k_{\mu;\nu}(\lambda) \bar{\epsilon}^\mu(\lambda) \bar{\epsilon}^\nu(\lambda), \quad (50)$$

where $\epsilon^\mu = E_1^\mu + i E_2^\mu$ and an over-bar denotes complex conjugation. Then, the optical deformation matrix becomes

$$\mathcal{S}(\lambda) = \begin{pmatrix} \theta(\lambda) - \Re\sigma(\lambda) & \Im\sigma(\lambda) \\ \Im\sigma(\lambda) & \theta(\lambda) + \Re\sigma(\lambda) \end{pmatrix}. \quad (51)$$

Differentiating (48) with respect to λ gives

$$\ddot{\xi}(\lambda) = \mathcal{T}(\lambda) \xi(\lambda), \quad (52)$$

where the optical matrix \mathcal{T} is given by

$$\mathcal{T} = \dot{\mathcal{S}} + \mathcal{S}^2 = \begin{pmatrix} \mathcal{R}(\lambda) - \Re\mathcal{F}(\lambda) & \Im\mathcal{F}(\lambda) \\ \Im\mathcal{F}(\lambda) & \mathcal{R}(\lambda) + \Re\mathcal{F}(\lambda) \end{pmatrix}, \quad (53)$$

where, from Sach's transport equations, we have for θ and σ [17]

$$\dot{\theta} + \theta^2 + |\sigma|^2 = \mathcal{R}, \quad (54)$$

$$\dot{\sigma} + 2\theta\sigma = \mathcal{F}, \quad (55)$$

with

$$\mathcal{R} = -\frac{1}{2} R_{\alpha\beta} k^\alpha k^\beta, \quad (56)$$

$$\mathcal{F} = -\frac{1}{2} C_{\alpha\beta\gamma\delta} \bar{\epsilon}^\alpha k^\beta \bar{\epsilon}^\gamma k^\delta. \quad (57)$$

The source of convergence is written in terms of the Ricci curvature, \mathcal{R} , while the source of shear is written in terms of the Weyl curvature, \mathcal{F} .

For Szekeres spacetimes, the Ricci and Weyl curvatures can be calculated using the Ricci and Weyl tensor components displayed in Appendix A.

Equation (52) describes the deformation, or lensing, of an infinitesimal light bundle as it can be seen propagating as ξ_i on the screen corresponding to each event parametrized by λ along the fiducial ray. Due to the linearity of (52), we can use a Jacobi mapping between ξ_i

and the initial angle ϑ_i between the ray of interest and the fiducial ray, which reads

$$\xi(\lambda) = \mathcal{D}(\lambda)\vartheta. \quad (58)$$

The evolution of \mathcal{D} with respect to λ obeys

$$\dot{\mathcal{D}} = \mathcal{T}\mathcal{D}, \quad (59)$$

where, at $\lambda = 0$, we have $\mathcal{D} = 0$ and $\dot{\mathcal{D}} = 1$. The linearity of (52) implies that its solution $\xi(\lambda)$ is related to its initial value, $\xi(0) = \vec{\vartheta}$, by a λ -dependent linear transformation which can be written as (58). With such a choice of λ , $\vec{\vartheta}$ is the vectorial angle between γ_o and a neighbouring ray. The Jacobi map (58) takes infinitesimal changes of ray directions at the observer, back to a screen at an event of γ_o determined by the value of λ . If that event is taken on some source plane, $\mathcal{D}(\lambda)$ corresponds to the properly scaled magnification matrix \mathcal{A} of lens theory. Indeed, since we consider an inhomogeneous model with a limiting background FLRW spacetime, this relationship reads [35]

$$\mathcal{D} = D_A^{FLRW} \mathcal{A}, \quad (60)$$

where D_A^{FLRW} is the angular diameter distance in the background model and \mathcal{A} can be written as

$$\mathcal{A} = \begin{pmatrix} 1 - \kappa - \gamma_1 & \gamma_2 \\ \gamma_2 & 1 - \kappa + \gamma_1 \end{pmatrix}, \quad (61)$$

where the complex-valued lensing shear $\gamma = \gamma_1 + i\gamma_2$ describes the stretching of galaxy images due to lensing, and the convergence κ describes a change in size and brightness. It is common to write this in terms of the reduced shear:

$$g_i = \frac{\gamma_i}{1 - \kappa}, \quad (62)$$

which gives

$$\mathcal{A} = (1 - \kappa) \begin{pmatrix} 1 - g_1 & g_2 \\ g_2 & 1 + g_1 \end{pmatrix}. \quad (63)$$

Now, denoting a solution to (59) as

$$\mathcal{D} = \begin{pmatrix} D_{11} & D_{12} \\ D_{21} & D_{22} \end{pmatrix}. \quad (64)$$

We can write the convergence and shear due to the inhomogeneities in the Szekeres region of the Universe as

$$\kappa = 1 - \frac{D_{11} + D_{22}}{2D_A^{FLRW}}, \quad (65)$$

$$\gamma_1 = \frac{D_{22} - D_{11}}{2D_A^{FLRW}}, \quad (66)$$

$$\gamma_2 = \frac{D_{12}}{D_A} = \frac{D_{21}}{D_A^{FLRW}}, \quad (67)$$

$$\gamma = \sqrt{\gamma_1^2 + \gamma_2^2}. \quad (68)$$

Comparison of (48) and the derivative of (58) shows that

$$\dot{\mathcal{D}} = \mathcal{S}\mathcal{D}, \quad (69)$$

which implies that \mathcal{S} can be obtained from \mathcal{D} .

Now, from the definition (58) of the Jacobi map, it follows that

$$|\det\mathcal{D}(\lambda)| = \frac{\delta A(\lambda)}{\delta\Omega}. \quad (70)$$

This shows that $\det\mathcal{D}(\lambda)$ contains information about the area $\delta A(\lambda)$ as well as about the parity, i.e., the orientation of the beam at λ relative to that close to the vertex of the light cone.

2. Geodesic deviation in the Szekeres model

The key components of the optical tidal matrix are the Ricci and Weyl curvatures. Their components are given in Appendix A, after renaming the metric components as [34]

$$H^2 = \frac{(\Phi_{,r} - \Phi E_{,r}/E)^2}{\epsilon - k}, \quad F^2 = \frac{\Phi^2}{E^2}. \quad (71)$$

With this renaming, the null geodesic equations become

$$\dot{k}^t + HH_{,t}(k^r)^2 + FF_{,t}[(k^p)^2 + (k^q)^2] = 0, \quad (72)$$

$$\begin{aligned} & H^2 \dot{k}^r + 2H\dot{H}k^r - HH_{,r}(k^r)^2 \\ & - FF_{,r}[(k^p)^2 + (k^q)^2] = 0, \end{aligned} \quad (73)$$

$$\begin{aligned} & F^2 \dot{k}^p + 2F\dot{F}k^r - HH_{,p}(k^r)^2 \\ & - FF_{,p}[(k^p)^2 + (k^q)^2] = 0, \end{aligned} \quad (74)$$

$$\begin{aligned} & F^2 \dot{k}^q + 2F\dot{F}k^r - HH_{,q}(k^r)^2 \\ & - FF_{,q}[(k^p)^2 + (k^q)^2] = 0. \end{aligned} \quad (75)$$

To which we can add the null vector condition written under the form

$$(k^t)^2 - H^2(k^r)^2 - F^2[(k^p)^2 + (k^q)^2] = 0. \quad (76)$$

Inserting (76) into (72)-(75) and using the definition of the total derivative of H and F with respect to λ , we obtain

$$\dot{k}^t + \frac{F_{,t}}{F}(k^t)^2 + H^2 \left(\frac{H_{,t}}{H} - \frac{F_{,t}}{F} \right) (k^r)^2 = 0, \quad (77)$$

$$\dot{k}^r - \left(\frac{H_{,r}}{H} - \frac{F_{,r}}{F} \right) (k^r)^2 + \frac{\dot{H}}{H} k^r - \frac{F_{,r}}{H^2 F} (k^t)^2 = 0, \quad (78)$$

$$\dot{k}^p - \frac{H^2}{F^2} \left(\frac{H_{,p}}{H} - \frac{F_{,p}}{F} \right) (k^r)^2 + \frac{\dot{F}}{F} k^r - \frac{F_{,p}}{F^3} (k^t)^2 = 0, \quad (79)$$

$$\dot{k}^q - \frac{H^2}{F^2} \left(\frac{H_{,q}}{H} - \frac{F_{,q}}{F} \right) (k^r)^2 + \frac{\dot{F}}{F} k^r - \frac{F_{,q}}{F^3} (k^t)^2 = 0. \quad (80)$$

Finally, the screen basis vectors E_1^μ and E_2^μ are defined at the observer in terms of the null vector $k_o^\mu =$

$$(E_1^\mu)_o = \left(0, \frac{F}{H} \sqrt{(k_o^p)^2 + (k_o^q)^2}, -\frac{H}{F} \frac{k_o^r k_o^p}{\sqrt{(k_o^p)^2 + (k_o^q)^2}}, -\frac{H}{F} \frac{k_o^r k_o^q}{\sqrt{(k_o^p)^2 + (k_o^q)^2}} \right), \quad (82)$$

$$(E_2^\mu)_o = \left(0, 0, \frac{1}{F} \frac{k_o^p}{\sqrt{(k_o^p)^2 + (k_o^q)^2}}, -\frac{1}{F} \frac{k_o^q}{\sqrt{(k_o^p)^2 + (k_o^q)^2}} \right), \quad (83)$$

for non-zero k_o^p and k_o^q , which means that the null vectors are not initially radial at the observer.

Once E_1^μ and E_2^μ are determined at the observer, they are parallelly propagated together with u^μ and k^μ along the path of the fiducial light ray. Thus, at any position along this fiducial ray parametrized by λ , we have similarly the following differential equations to be solved

$$\dot{E}_a^\mu + \Gamma_{\alpha\beta}^\mu E_a^\alpha k^\beta = 0, \quad a = \{1, 2\}. \quad (84)$$

These equations can be written as

$$\dot{E}_a^t + H H_{,t} E_a^r k^r + F F_{,t} (E_a^p k^p + E_a^q k^q) = 0, \quad (85)$$

$$\begin{aligned} \dot{E}_a^r + \frac{\dot{H}}{H} E_a^r + \frac{k^r}{H} (H_{,t} E_a^t + H_{,p} E_a^p + H_{,q} E_a^q) \\ - \frac{F F_{,r}}{H^2} (E_a^p k^p + E_a^q k^q) = 0, \end{aligned} \quad (86)$$

$$\begin{aligned} \dot{E}_a^p + \frac{\dot{F}}{F} E_a^p + \frac{k^p}{F} (F_{,t} E_a^t + F_{,r} E_a^r + F_{,q} E_a^q) \\ - \frac{H H_{,p}}{F^2} E_a^r k^r - \frac{F_{,p}}{F} E_a^q k^q = 0, \end{aligned} \quad (87)$$

$$\begin{aligned} \dot{E}_a^q + \frac{\dot{F}}{F} E_a^q + \frac{k^q}{F} (F_{,t} E_a^t + F_{,r} E_a^r + F_{,p} E_a^p) \\ - \frac{H H_{,q}}{F^2} E_a^r k^r - \frac{F_{,q}}{F} E_a^p k^p = 0. \end{aligned} \quad (88)$$

In general, (59) has no analytic solution. Therefore, to evaluate the Jacobi matrix \mathcal{D} , and, through (60), the magnification matrix \mathcal{A} , we have to solve simultaneously the set of second order differential equations (59) for \mathcal{D} and the sets of first order differential equations, (77)-(80) for k^μ , and (85)-(88) for E_1^μ and E_2^μ .

Since weak lensing surveys usually yield, for each observed object: (i) the source galaxy redshift, (ii) the galaxy position on the sky, (iii) reduced shear estimates, the reconstruction of \mathcal{A} can be done from these data, and compared to the theoretical Szekeres lensing quantities.

$(-1, k_o^r, k_o^p, k_o^q)$, of the comoving velocity of the observer $u_o^\mu = (1, 0, 0, 0)$, and of the metric functions in terms of H and F . These screen basis vectors must satisfy the orthogonality conditions

$$k_\mu E_a^\mu = 0, \quad u_\mu E_a^\mu = 0, \quad E_{a\mu} E_b^\mu = 0, \quad (81)$$

where $a = \{1, 2\}$, and which imply the initial conditions

F. Redshift-space distortions

For observed galaxies, especially for faraway objects, the line-of-sight velocity is coupled with the line-of-sight distance. This phenomenon is referred to as the redshift-space distortion (RSD) effect and must be taken into account for cosmological analyses of galaxy position data.

A galaxy peculiar velocity can be divided into two parts: the peculiar velocity of the galaxy host halo, determined by the large-scale matter distribution, and the galaxy random velocity within the halo. Since the RSD effects at large scale are determined by the large scale density, and, at small scale, by the galaxy virial motion in the halo, one can try to model it separately at these two different scales.

The distance between a galaxy and the observer is measured by its redshift, z_o , which includes the cosmological redshift, z_{cos} , arising from the Hubble expansion, and the Doppler redshift, z_{pec} , caused by the galaxy peculiar velocity along the line-of-sight, v_{los} . This can be written as

$$\frac{\lambda_o}{\lambda_e} = 1 + z_o = (1 + z_{pec})(1 + z_{cos}). \quad (89)$$

On large scales, matter around overdensity regions is falling in and such a stream velocity makes the scatter of z_o for galaxies in this region smaller than that of their z_{cos} , resulting in a visual compression effect along the line-of-sight known as the Kaiser effect [36]. On small scales, some nearby galaxies around the target galaxy can have very large line-of-sight peculiar velocities due to virial motion, and thus will make the scatter of z_o along the line-of-sight for those galaxies much bigger than that of their z_{cos} , which makes these galaxies stretch along the line-of-sight. This is known as the finger-of-God effect [36, 37].

Therefore, accurate estimations of line-of-sight velocities is strongly needed. Eliminating the nonlinear effect

of RSD, e. g., in the BAO reconstruction, is indeed important. This implies an accurate determination of the redshift in the framework of the Szekeres model which yields proper equations for the null geodesics and the redshift, z_{cos} , see Sec.II A.

Since a galaxy peculiar velocity is determined by surrounding matter, which is normally traced by neighbouring galaxies, one needs to define variables which contain the necessary information to extract the galaxy line-of-sight velocity from survey data, which can be analysed and implemented with the tools of artificial neural networks. This has been done in a recent study completed with a Λ CDM background [38] which should be adapted to the Szekeres background in future works.

G. Redshift drift

The redshift drift is the redshift increase or decrease that a comoving observer can measure when looking at the same comoving source at two different instants. This drift proceeds from the fact that the source is measured on the observer's two different past light cones. It occurs in any expanding universe, but its magnitude depends on the geometry of the region travelled by the rays.

This effect has already been analyzed for quasi-spherical models, first, in the case where the geodesics are axially directed in an axially symmetric Szekeres spacetime [39], then, for randomly directed geodesics [40]. However, in the second analysis, a purely Szekeres effect, ensuring that two rays emitted subsequently by the same source are directed in two different directions, has been missed. This effect has been fully worked out by Krasiński and Bolejko [13] and their method will be used in Sec.IV B to study both effects, the redshift drift and the position drift, i. e., a new effect particular to general light rays travelling in an inhomogeneous spacetime.

Now, the cosmological redshift drift effect has been predicted a long time ago [41, 42]. It has not been much considered since then because of its tiny amplitude, i. e., of the order 10^{-18} s^{-1} . However, measuring it on a time-scale of a few years is now within reach. Some authors have indeed proposed to use strong-gravitational-lens time delays to complete such a measurement [43–45]. This could be done within a very short time, the change in time being supplied by the time delay instead of by waiting for years between observations. We can therefore look at future experiments able to measure it within a reasonable time lapse.

IV. EFFECTS OBSERVABLE IN INHOMOGENEOUS MODELS ONLY

We consider here some physical effects occurring only in inhomogeneous models and exhibiting a particular signature when considered in a Szekeres geometry. Should such effects be observed in our Universe, this might be

strong hints in favor of an inhomogeneous cosmology at scales corresponding with those of the effects.

A. Differential cosmic expansion and Hubble flow anisotropies

Inhomogeneous cosmological models such as LTB and Szekeres, exhibit differential cosmic expansion, i. e., the inhomogeneous counterpart of the Hubble parameter is no more a function of time only as in FLRW, but it becomes a function of space as well as of time.

In the geometric optics approximation, a formula for the redshift is [46]

$$\frac{1}{(1+z)^2} \frac{dz}{ds} = \frac{\Theta}{3} + \Sigma_{\alpha\beta} n^\alpha n^\beta + u_{;\beta}^\alpha u^\beta n_\alpha, \quad (90)$$

where u^α is the matter velocity field, n_α is the connecting covector field, locally orthogonal to u^α , $\Sigma_{\alpha\beta}$ is the shear of the velocity field, and $\Theta = u_{;\alpha}^\alpha$ is its expansion. In homogeneous and isotropic models, the shear vanishes and the expansion reduces to $\Theta = 3H(t)$, H being the Hubble parameter. Once cosmic structures are formed, the expansion field becomes non-uniform, and thus the shear and the acceleration, $\dot{u}^\alpha = u_{;\beta}^\alpha u^\beta$ vanishes no more which modifies formula (90) for the redshift.

If the redshift of an observed object can be described solely in terms of a homogeneous expansion and of a Doppler effect, such as

$$(1+z)_o = (1+z)_{FLRW} (1+z)_{Doppler}, \quad (91)$$

where the first term on the right hand side refers to the global homogeneous and isotropic expansion of a background FLRW model and the second term corresponds to a Doppler effect with respect to the FLRW frame which combines the motions of the observer (local boost) and of the observed object (peculiar motion), then the redshift anisotropy is called kinematic. The factor $(1+z)_{FLRW}$ must be defined with respect to the canonical choice of the local Lorentz frame corresponding to a local boost of the local group of galaxies.

If the redshift of an observed source (galaxy-type or CMB) cannot be explained entirely in terms of (91), then non-kinematic effects are present and can be a hint of inhomogeneous geometry.

Distances are also affected by the non vanishing of the acceleration and shear of the velocity field. Now, a modification of the expansion rate implies a modification of the distances. Therefore, a measurement of such effects, if it happens to occur, could be considered as a clear step toward inhomogeneous cosmology.

Indeed, for the study of the redshift dependence of the expansion anisotropy of the local Universe, at redshift $z \ll 1$, a Taylor expansion of the luminosity distance can be used. In a FLRW cosmology it reads

$$D_L(z) = \frac{c}{H_0} \left[z + \frac{1}{2}(1-q_0)z^2 - \frac{1}{6}(1-q_0) \right]$$

$$- \left[3q_0^2 + j_0 + \frac{kc^2}{H_0^2 a_0^2} \right] z^3 + O(z^4), \quad (92)$$

where q_0 is the deceleration parameter, j_0 is the jerk parameter, $a_0 = a(t_0)$ is the present cosmic scale factor and $k = \{-1, 0, 1\}$ is the spatial curvature.

If the Universe is described by an alternative space-time where the Taylor expansion still converges at small z , an expansion law written as a Taylor series in z can still apply [47]. However, in such cases, the second and higher order coefficients are different from FLRW ones. In particular, generic inhomogeneous cosmological models do not expand such as to keep their spatial curvature, k , constant.

A galaxy survey providing, for each galaxy, its angular position, its luminosity distance, and its redshift can be used to compute the redshift dependence of the anisotropy of the local Universe expansion [48]. For this purpose, each light ray is propagated, using the null geodesic equations (7)-(10), up until D_L equals the measured distance, and the corresponding redshift is obtained through (90). Then the dipole proceeds from the reconstruction of mock catalogues where the measured

galaxy redshifts are replaced by their Szekeres counterparts computed in the measured direction.

B. Redshift drift and position drift

As explained in Sec.III G an observer located in a Szekeres universe who measures the same light source at different instants, observes not only a shift in the object redshift, but also in its position on the sky. A method for calculating both effects can be inspired from that used by Krasiński and Bolejko [13]. The aim of these authors was to exhibit the new position drift effect and we will see that, as a bonus, we can obtain the redshift drift.

The null geodesic equations (7)-(10) can be written with respect to the r coordinate by implementing the following formula for the total derivative:

$$\frac{d^2 x^\alpha}{ds^2} = \left(\frac{dr}{ds} \right)^2 \frac{d^2 x^\alpha}{dr^2} + \frac{d^2 r}{ds^2} \frac{dx^\alpha}{dr}, \quad (93)$$

which gives

$$\frac{d^2 t}{dr^2} + \left(\frac{\Phi_{,tr} - \Phi_{,t} E_{,r}/E}{\epsilon - k} \right) (\Phi_{,r} - \Phi E_{,r}/E) + \frac{\Phi \Phi_{,t}}{E^2} \left[\left(\frac{dp}{dr} \right)^2 + \left(\frac{dq}{dr} \right)^2 \right] + U \frac{dt}{dr} = 0, \quad (94)$$

$$\begin{aligned} & \frac{d^2 p}{dr^2} + 2 \frac{\Phi_{,t}}{\Phi} \frac{dt}{dr} \frac{dp}{dr} - \frac{\Phi_{,r} - \Phi E_{,r}/E}{\Phi(\epsilon - k)} (E_{,r} E_{,p} - E E_{,rp}) \\ & + 2 \frac{\Phi_{,r} - \Phi E_{,r}/E}{\Phi} \frac{dp}{dr} - 2 \frac{E_{,q}}{E} \frac{dp}{dr} \frac{dq}{dr} + \frac{E_{,p}}{E} \left[- \left(\frac{dp}{dr} \right)^2 + \left(\frac{dq}{dr} \right)^2 \right] + U \frac{dp}{dr} = 0, \end{aligned} \quad (95)$$

$$\begin{aligned} & \frac{d^2 q}{dr^2} + 2 \frac{\Phi_{,t}}{\Phi} \frac{dt}{dr} \frac{dq}{dr} - \frac{\Phi_{,r} - \Phi E_{,r}/E}{\Phi(\epsilon - k)} (E_{,r} E_{,q} - E E_{,rq}) \\ & + 2 \frac{\Phi_{,r} - \Phi E_{,r}/E}{\Phi} \frac{dq}{dr} - 2 \frac{E_{,p}}{E} \frac{dp}{dr} \frac{dq}{dr} + \frac{E_{,q}}{E} \left[\left(\frac{dp}{dr} \right)^2 - \left(\frac{dq}{dr} \right)^2 \right] + U \frac{dq}{dr} = 0, \end{aligned} \quad (96)$$

where we have defined

$$\begin{aligned} U(t, r, p, q) = & -2 \frac{\Phi_{,tr} - \Phi_{,t} E_{,r}/E}{\Phi_{,r} - \Phi E_{,r}/E} \frac{dt}{dr} - \left[\frac{\Phi_{,rr} - \Phi E_{,rr}/E}{\Phi_{,r} - \Phi E_{,r}/E} - \frac{E_{,r}}{E} + \frac{k_{,r}}{2(\epsilon - k)} \right] - 2 \frac{\Phi}{E^2} \left(\frac{E_{,r} E_{,p} - E E_{,rp}}{\Phi_{,r} - \Phi E_{,r}/E} \right) \frac{dp}{dr} \\ & - 2 \frac{\Phi}{E^2} \left(\frac{E_{,r} E_{,q} - E E_{,rq}}{\Phi_{,r} - \Phi E_{,r}/E} \right) \frac{dq}{dr} + \frac{\Phi}{E^2} \left(\frac{\epsilon - k}{\Phi_{,r} - \Phi E_{,r}/E} \right) \left[\left(\frac{dp}{dr} \right)^2 + \left(\frac{dq}{dr} \right)^2 \right]. \end{aligned} \quad (97)$$

Consider two light rays directed from a unique source towards an observer with coordinates (r, p, q) and separated by a time-interval τ , short with respect to the time travelled by the rays between the source and the observer. The trajectory of the first ray is given by

$$(t, p, q) = (T(r), X(r), Y(r)). \quad (98)$$

The equation for the second signal is

$$(t, p, q) = (T(r) + \tau(r), X(r) + \zeta(r), Y(r) + \psi(r)). \quad (99)$$

This means that, while the first ray intersects some given hypersurface $r = \text{const}$ at $(t, p, q) = (T, X, Y)$, the second ray will intersect the same hypersurface, not only later on, but, in general, at a different comoving location. Therefore, in general, the two rays will intersect different sequences of intermediate matter worldlines.

Since we are using comoving coordinates, we have $(\zeta, \psi) = (0, 0)$ at the emitter and at the observer. Moreover, the directions of the first and second rays are determined by $(dp/dr, dq/dr)$ and $(dp/dr + \xi(r), dq/dr + \eta(r))$ respectively, with $\xi = d\zeta/dr$ and $\eta = d\psi/dr$. It is assumed that $d\tau/dr, \zeta, \psi, \xi$, and η are small, of the same order as τ , hence all terms nonlinear in any of them and terms involving their products will be neglected.

Since $\zeta = \psi = 0$ at the observer, they are not observable. Anyhow, they must be monitored along the rays because they enter the equation for τ which is connected to the redshift drift δz through

$$1 + z + \delta z = \frac{d(t + \tau)/dr|_{r=r_e}}{d(t + \tau)/dr|_{r=r_o}}. \quad (100)$$

Indeed, subtracting from (100) the redshift defining equation

$$\frac{d^2\tau}{dr^2} + \frac{\Phi_{01}\Delta\Phi_1 + \Phi_1\Delta\Phi_{01}}{\epsilon - k} + \frac{(\Phi_{,t}^2 + \Phi\Phi_{,tt})\Sigma\tau}{E^2} - 2\frac{\Phi\Phi_{,t}\Delta E\Sigma}{E^3} + \frac{\Phi\Phi_{,t}\Delta\Sigma}{E^2} + \Delta U\frac{dt}{dr} + U\frac{d\tau}{dr} = 0, \quad (103)$$

$$\begin{aligned} \frac{d^2\zeta}{dr^2} &+ 2\left(\frac{\Phi_{,tt}}{\Phi} - \frac{\Phi_{,t}^2}{\Phi^2}\right)\frac{dt}{dr}\frac{dp}{dr}\tau + 2\frac{\Phi_{,t}}{\Phi}\frac{dp}{dr}\frac{d\tau}{dr} + 2\frac{\Phi_{,t}}{\Phi}\frac{dt}{dr}\xi - \frac{\Delta\Phi_1 E_{12}}{(\epsilon - k)\Phi} + \frac{\Phi_{,t}\Phi_1 E_{12}\tau}{(\epsilon - k)\Phi^2} - \frac{\Phi_1\Delta E_{12}}{(\epsilon - k)\Phi} \\ &+ 2\left(\frac{\Delta\Phi_1}{\Phi} - \frac{\Phi_1\Phi_{,t}\tau}{\Phi^2}\right)\frac{dp}{dr} + 2\frac{\Phi_1}{\Phi}\xi - \left(\frac{\zeta}{SE} - \frac{E_{,p}\Delta E}{E^2}\right)\left(\frac{dp}{dr}\right)^2 - 2\frac{E_{,p}\xi}{E}\frac{dp}{dr} - 2\left(\frac{\psi}{SE} - \frac{E_{,q}\Delta E}{E^2}\right)\frac{dp}{dr}\frac{dq}{dr} \\ &- 2\frac{E_{,q}}{E}\left(\frac{dq}{dr}\xi + \frac{dp}{dr}\eta\right) + \left(\frac{\zeta}{SE} - \frac{E_{,p}\Delta E}{E^2}\right)\left(\frac{dq}{dr}\right)^2 + 2\frac{E_{,p}\eta}{E}\frac{dq}{dr} + \Delta U\frac{dp}{dr} + U\xi = 0, \end{aligned} \quad (104)$$

$$\begin{aligned} \frac{d^2\psi}{dr^2} &+ 2\left(\frac{\Phi_{,tt}}{\Phi} - \frac{\Phi_{,t}^2}{\Phi^2}\right)\frac{dt}{dr}\frac{dq}{dr}\tau + 2\frac{\Phi_{,t}}{\Phi}\frac{dq}{dr}\frac{d\tau}{dr} + 2\frac{\Phi_{,t}}{\Phi}\frac{dt}{dr}\eta - \frac{\Delta\Phi_1 E_{13}}{(\epsilon - k)\Phi} + \frac{\Phi_{,t}\Phi_1 E_{13}\tau}{(\epsilon - k)\Phi^2} - \frac{\Phi_1\Delta E_{13}}{(\epsilon - k)\Phi} \\ &+ 2\left(\frac{\Delta\Phi_1}{\Phi} - \frac{\Phi_1\Phi_{,t}\tau}{\Phi^2}\right)\frac{dq}{dr} + 2\frac{\Phi_1}{\Phi}\eta - \left(\frac{\psi}{SE} - \frac{E_{,q}\Delta E}{E^2}\right)\left(\frac{dp}{dr}\right)^2 + 2\frac{E_{,q}\xi}{E}\frac{dp}{dr} - 2\left(\frac{\zeta}{SE} - \frac{E_{,p}\Delta E}{E^2}\right)\frac{dp}{dr}\frac{dq}{dr} \\ &- 2\frac{E_{,p}}{E}\left(\frac{dq}{dr}\xi + \frac{dp}{dr}\eta\right) - \left(\frac{\psi}{SE} - \frac{E_{,q}\Delta E}{E^2}\right)\left(\frac{dq}{dr}\right)^2 - 2\frac{E_{,q}\eta}{E}\frac{dq}{dr} + \Delta U\frac{dq}{dr} + U\eta = 0, \end{aligned} \quad (105)$$

where

$$\Phi_1 \equiv \Phi_{,r} - \Phi\frac{E_{,r}}{E}, \quad \Phi_{01} \equiv \Phi_{,tr} - \Phi_{,t}\frac{E_{,r}}{E}, \quad \Phi_{11} \equiv \Phi_{,rr} - \Phi\frac{E_{,rr}}{E}, \quad (106)$$

$$E_{12} \equiv E_{,r}E_{,p} - EE_{,rp}, \quad E_{13} \equiv E_{,r}E_{,q} - EE_{,rq}, \quad \Sigma \equiv \left(\frac{dp}{dr}\right)^2 + \left(\frac{dq}{dr}\right)^2, \quad (107)$$

$$\Delta\Phi = \Phi_{,t}\tau, \quad \Delta(\Phi_{,t}) = \Phi_{,tt}\tau, \quad \Delta\frac{dt}{dr} = \frac{d\tau}{dr}, \quad \Delta\frac{dp}{dr} = \xi, \quad \Delta\frac{dq}{dr} = \eta, \quad (108)$$

$$\Delta E = E_{,p}\zeta + E_{,q}\psi, \quad \Delta E_{,p} = \frac{\zeta}{S}, \quad \Delta E_{,q} = \frac{\psi}{S}, \quad \Delta\Phi_1 = \Phi_{01}\tau + \frac{\Phi E_{12}}{E^2}\zeta + \frac{\Phi E_{13}}{E^2}\psi, \quad (109)$$

$$\Delta\Phi_{01} = (\Phi_{,ttr} - \Phi_{,tt}\frac{E_{,r}}{E})\tau + \frac{\Phi_{,t}E_{12}}{E^2}\zeta + \frac{\Phi_{,t}E_{13}}{E^2}\psi, \quad (110)$$

tion written as

$$1 + z = \frac{dt/dr|_{r=r_e}}{dt/dr|_{r=r_o}}, \quad (101)$$

and arranging, we obtain

$$\delta z = (1 + z) \left(\frac{\frac{d\tau/dr|_{r=r_e} - \frac{d\tau/dr|_{r=r_o}}{dt/dr|_{r=r_o}}}{1 + \frac{d\tau/dr|_{r=r_o}}{dt/dr|_{r=r_o}}} \right). \quad (102)$$

In the following, the Δ symbol applied to some expression will denote the difference between this expression taken at $(t + \tau, r, p + \zeta, q + \psi)$ and at (t, r, p, q) , linearized in (τ, ζ, ψ) . By applying Δ to (94)-(96), one obtains [13]

$$\Delta\Phi_{11} = (\Phi_{,ttr} - \Phi_{,t}\frac{E_{,rr}}{E})\tau + \frac{\Phi}{E^2} [(E_{,rr}E_{,p} - EE_{,rrp})\zeta + (E_{,rr}E_{,q} - EE_{,rrq})\psi], \quad (111)$$

$$\Delta E_{12} = (E_{,r}E_{,pp} - EE_{,rpp})\zeta + (E_{,rq}E_{,p} - E_{,q}E_{,rp})\psi, \quad \Delta E_{13} = (E_{,rp}E_{,q} - E_{,p}E_{,rq})\zeta + (E_{,r}E_{,qq} - EE_{,rqq})\psi, \quad (112)$$

$$\Delta\Sigma = 2\frac{dp}{dr}\xi + 2\frac{dq}{dr}\eta, \quad (113)$$

$$\begin{aligned} \Delta U &= 2\left(-\frac{\Delta\Phi_{01}}{\Phi_1} + \frac{\Phi_{01}\Delta\Phi_1}{\Phi_1^2}\right)\frac{dt}{dr} - 2\frac{\Phi_{01}}{\Phi_1}\frac{d\tau}{dr} - \frac{\Delta\Phi_{11}}{\Phi_1} + \frac{\Phi_{11}\Delta\Phi_1}{\Phi_1^2} + \frac{\Delta E_{,r}}{E} - \frac{E_{,r}\Delta E}{E^2} \\ &+ 2\left(-\frac{\Phi_{,t}E_{12}\tau}{E^2\Phi_1} + 2\frac{\Phi\Delta EE_{12}}{E^3\Phi_1} - \frac{\Phi\Delta E_{12}}{E^2\Phi_1} + \frac{\Phi E_{12}\Delta\Phi_1}{E^2\Phi_1^2}\right)\frac{dp}{dr} - 2\frac{\Phi E_{12}\xi}{E^2\Phi_1} \\ &+ 2\left(-\frac{\Phi_{,t}E_{13}\tau}{E^2\Phi_1} + 2\frac{\Phi\Delta EE_{13}}{E^3\Phi_1} - \frac{\Phi\Delta E_{13}}{E^2\Phi_1} + \frac{\Phi E_{13}\Delta\Phi_1}{E^2\Phi_1^2}\right)\frac{dq}{dr} - 2\frac{\Phi E_{13}\eta}{E^2\Phi_1} \\ &+ \frac{(\epsilon - k)\Phi\Sigma}{E^2\Phi_1} \left(\frac{\Phi_{,t}\tau}{\Phi} - 2\frac{\Delta E}{E} - \frac{\Delta\Phi_1}{\Phi_1} + \frac{\Delta\Sigma}{\Sigma}\right). \end{aligned} \quad (114)$$

Then, applying the Δ -operation to (11), written with r as a parameter on the null geodesics, one obtains

$$\frac{d\tau}{dr}\frac{dt}{dr} = \frac{\Phi_1\Delta\Phi_1}{\epsilon - k} + \left(\frac{\Phi\Phi_{,t}\tau}{E^2} - \frac{\Phi^2\Delta E}{E^3}\right)\Sigma + \frac{\Phi^2}{E^2}\left(\frac{dp}{dr}\xi + \frac{dq}{dr}\eta\right). \quad (115)$$

The redshift drift measured by the observer with coordinates (t, r, p, q) in a Szekeres spacetime is determined by $d\tau/dr$ and k^t through (102), but the null geodesic equations have to be solved together with (103)-(105), since ζ and ψ must be monitored along the ray because the equations which determine (τ, ζ, ψ) are coupled and the same applies to the determination of k^t , depending on the evolution of k^p , and k^q . Therefore, after solving this double set of equations, we have all the results needed to obtain the measurable redshift drift from (102).

Moreover, the second ray is emitted in a different direction and is received from a different direction by the observer who thus sees every light source slowly drifting across the sky [13, 49]. This effect is present in other inhomogeneous cosmological models, e. g., it affects non-radial rays in spherically symmetric LTB models. It has been predicted, in this framework, under the name of cosmic parallax [50, 51].

It has been demonstrated by Krasiński [52] that the only spacetimes in the Szekeres family where the position drift vanishes for every null geodesic pair are the Friedmann models. Therefore, an observation of this drift for any remote source would be an evidence for the Universe to be inhomogeneous at large scales. To obtain a rough estimate of the magnitude of this effect, Krasiński and Bolejko [13] have calculated the time-averaged rate of change of the source position in a LTB configuration where the null geodesics on which light travels are non-radial. They obtained a drift $\sim 10^{-6} - 10^{-7}$ arcsec/year. This effect is thus too tiny to be measured by current or coming soon experiments. However, since we experience regularly tremendous technical progresses, hope is left for a future probe of such a position drift.

C. Cosmological blueshifts in QSS spacetimes

For completeness, we stress that the possible existence of strongly blueshifted light sources in QSS spacetimes has been rather recently put forward [53]. It has been shown that, in an axially symmetric QSS model, infinite blueshift can appear, but only on axial rays, i. e., rays which intersect every space orthogonal to the dust flow on the symmetry axis. Moreover, using a particular axially symmetric QSS model, strong blueshifts for rays emitted shortly after the Big Bang and running close to the symmetry axis have been numerically obtained. Then, in a QSS toy model deprived of any symmetry, two null lines such that rays in their vicinity exhibit similar features to those in the vicinity of axial null geodesics in axially symmetric QSS models have been numerically found. This hints at the tentative possibility that rays observed with infinite or very strong blueshifts exist in general QSS spacetimes and are focussed around two special directions depending on the particular features of the model.

However, the Szekeres model validity region is bounded by the matter-radiation equality surface, since its gravitational fluid source, being dust, cannot account for any pressure component. Therefore, to move the initial frequencies of the relic radiation to an observable range, a particular form must be imposed to the $t_B(r)$ Bang function in a QSS region with no origin [54]. Hence, a possible future observation of such an effect would set additional stringent constraints on the Szekeres model to be fitted by other data, i. e., the quasi-spherical nature of the region of interest, implying there ($\epsilon = +1$), and the location and amplitude of the hump prescribed for

the $t_B(r)$ function.

V. PROCESSING THE DATA FITTING WITH DEEP LEARNING

The Szekeres equations displayed in this paper cannot be solved analytically. In previous studies involving these spacetimes, the method used was: assuming a set of given expressions for the Szekeres functions and parameters written with a finite number of constant parameters, fit these constant parameters to some cosmological data of interest and claim that the matched model solves the issue encompassed by the data [16, 21, 40, 48, 55–60]. In these works, the Szekeres equations are numerically integrated.

Now, with our more general model, besides one constant parameter, Λ , and one quasi-constant parameter, ϵ , we have also five functions of the r coordinate which are to be determined by the data. However, some simplifying constraints have to be imposed on these determining constants and functions. They are described below.

A. Basic physical constraints

To preserve the $(-+++)$ metric signature, we must have

$$\epsilon - k > 0. \quad (116)$$

This implies that, for quasi-hyperbolic regions ($\epsilon = -1$), only the hyperbolic evolution, obtained for $k < 0$, is possible, while quasi-planar regions can only experiment parabolic or hyperbolic evolution, where $k \leq 0$, and quasi-spherical regions can display the three types of evolution, elliptic ($k > 0$), parabolic ($k = 0$), and hyperbolic ($k < 0$). Such a crossed-constraint will have to be checked a posteriori for each cosmological model issuing from any data fit.

Shell-crossings are loci in spacetimes where the energy density diverges. They must therefore be avoided in a well-behaved cosmological model. The conditions for avoiding shell-crossing surfaces in quasi-spherical regions have been worked out by Szekeres [22] and enlarged to the other two classes of models by Hellaby and Krasiński [6, 9]. These requirements must be fulfilled by any quasi-spherical region in any proper Szekeres cosmological model and this will have to be checked also a posteriori.

For obvious physical reasons, the areal radius Φ has to be positive, which reads

$$\Phi(t, r) > 0, \quad \forall t \text{ and } r. \quad (117)$$

The metric should be nondegenerate and nonsingular, except at the Bang time $t_B(r)$. Since $(dp^2 + dq^2)/E^2$ maps to the unit sphere, plane, or pseudosphere, ,

$|S(r)| \neq 0$ is needed for a sensible mapping, and therefore, we can choose $S > 0$. In the cases where $\epsilon = 0$ or 1, E is bound to vanish at some (p, q) location. Thus the mapping is badly behaved there and the presence of such loci in a Szekeres region would rule out the corresponding model.

A well-behaved r coordinate implies

$$\infty > \frac{(\Phi_{,r} - \Phi E_{,r}/E)^2}{\epsilon - k} > 0. \quad (118)$$

In accordance with the weak energy condition, the energy density must be positive and the Kretschmann scalar must be finite which yields

$$\text{either } M_{,r} - 3ME_{,r}/E \geq 0 \quad \text{and} \quad \Phi_{,r} - \Phi E_{,r}/E \geq 0, \quad (119)$$

$$\text{or } M_{,r} - 3ME_{,r}/E \leq 0 \quad \text{and} \quad \Phi_{,r} - \Phi E_{,r}/E \leq 0. \quad (120)$$

Finally, the asymptotic recovery of an FLRW universe at the scales of inhomogeneity/homogeneity transition sets constraints on the forms of the functions to be fitted. Indeed, the Friedmann limit follows [11] when $\Phi(t, r) = rR(t)$, $k = k_0 r^2$, with $k_0 = \text{const}$, $B(r) = C(r) = 0$, and $D(r) = 4A(r) = \epsilon$, where

$$E(r, p, q) = A(p^2 + q^2) + 2Bp + 2Cq + D. \quad (121)$$

Using the E definition (2), this gives, for the most general Szekeres model where ϵ can be any among $\{-1, 0, +1\}$,

$$S = 2\epsilon, \quad P = Q = 0, \quad (122)$$

and still

$$\Phi(t, r) = rR(t), \quad k = k_0 r^2. \quad (123)$$

Therefore, we have five constraints to be fulfilled by any Szekeres cosmological model, which are summarized in (122) and (123) where $t \geq t_{trans}$ and $r \geq r_{trans}$ with the subscript 'trans' implying that the quantity is evaluated at the inhomogeneity/homogeneity transition. We are thus provided with two extra constant parameters for our model, t_{trans} and r_{trans} , which will have either to be determined through data fitting or, as a temporary simplification, fixed by an a priori but physically justified choice.

Now, although the choice of the way to determine the r coordinate is wide, it might be comfortable, for practical purpose, to choose a definition which includes readily the Friedmann limit, e. g., by setting $k(r) = k_0 r^2$ everywhere. In this case, only k_0 is left as a constant parameter to be determined by the observations. However, other choices can be convenient. For example, $t_B(r) = t_0$ implies that the Bang time is constant in the whole spacetime, which complies with Friedmann. One can, without loss of generality, equalize t_0 to zero, which amounts to a mere rescaling of the time coordinate. However, should the blueshift effect discussed in Sec.IV C be observed in the future, a more complicated Bang function

would arise. In the FLRW limit we have also, from (3), E/r^2 and M/r^3 constant [11]. Since, in the generic Szekeres solution, E is a function of r but also of p and q , a choice of the type $E = E_0 r^2$ might be too restrictive and might suppress some of the most interesting properties of the Szekeres solutions, while $M = M_0 r^3$ can be a better choice.

B. Fitting method

The problem is now to determine a cosmological model depending on five functions of r , chosen among S, P, Q, M, k , and t_B , and on two constant parameters, ϵ and Λ (or on four constant parameters if we decide to include t_{trans} and r_{trans}). For this, we have, in principle, an enormous set of data from surveys which are designed each to probe one or more particular cosmological effect. To deal with these data, we need a device very powerful and a method adapted to the fitting of functions. Both now exist. The device is one of those machines designed to implement deep learning and the method is symbolic regression.

Symbolic regression aims at determining among a huge set of data points the best-fit curve representing the data behavior. The results are obtained using machine-learning and they can be displayed either numerically or analytically depending on the problem at hand. This method can be employed to fit the Szekeres functions through a comparison of the model predictions to the actual cosmological data.

The ultimate goal will be actually to use the whole set of available cosmological data to thoroughly constrain the model. The aim of the present paper is therefore to give the most comprehensive set of tools needed to complete such a task in a Szekeres framework. However, we are aware that this will take a rather long time being carried out. But deep learning begins only to make a few steps in cosmology [61] and it should be appropriate that these steps should be done in the right direction.

We want also to stress that it is a current standard practice to automatically transform redshift data to the CMB rest frame before performing any analysis. The canonical CMB rest frame is defined by matching the 3.37 mK temperature of the CMB dipole to the dipole in a series expansion adapted to the Λ CDM paradigm. For a fully model independent study, this practice should be given up and the data used for our purpose should be the rawest as possible. This implies also to remove any Λ CDM inspired step from the data processing, such as bias or foreground removals.

VI. CONCLUSIONS

We have provided, in this paper, a thorough analysis of the Szekeres solution as a cosmological model. This solu-

tion benefits indeed from being devoid of any symmetry and, as an exact GR solution for dust plus a cosmological constant, it can be used as a substitute to the perturbed Λ CDM homogeneous model to describe the late Universe. At larger scales, it can be joined smoothly to a Friedmannian early time Universe which it has the nice property to include as a limit. We have here summarized, updated, corrected when needed, and completed with new results thirty years of work on the cosmological abilities of the Szekeres solutions. Actually, our aim has been to provide generalized, comprehensive, and consistent tools which can be readily used for future broad cosmological studies conducted with a well-adapted GR solution.

We have therefore presented the most general form of the equations needed to calculate the cosmological quantities which will have to be confronted to the different data surveys. We have displayed the light ray trajectories or null geodesics, the redshift, the different cosmological distances (area, luminosity, etc.), the intrinsic matter dipole, the BAO scale, the weak-lensing parameters (convergence and shear). Then we have proposed methods to use the current or soon-coming data surveys as probes to constraint the Szekeres independent parameters and functions. The effects which have been analyzed here are supernova dimming, galaxy number count evolution, CMB multipoles, SZ effects, BAO, weak lensing, redshift space distortions, redshift drift, differential cosmic expansion, position drift, and possible blueshift effect. The ultimate goal of finding the Szekeres model representing accurately the late Universe would imply that, at least, seven probes testing different physical effects should be used, one for each degree of freedom of the Szekeres parameters and functions. Given the heap of in principle available data, we have suggested to use machine learning to tackle more efficiently this issue.

The program described in this paper could appear rather ambitious. However, it seems to be one of the best ways to deal with the cosmological problem and to get rid of the current tensions and anomalies which plague the FLRW paradigm, as far as one wishes to keep GR as the proper gravitation theory, which is the main underlying physical paradigm of this endeavour. Moreover, the results displayed here can be of valuable use for any stepwise undertaking.

Acknowledgments

The author wishes to acknowledge enlightening discussions with Jean-Michel Alimi and with Andrzej Krasiński.

Appendix A: Szekeres geometrical quantities of interest

For completeness and to correct some errors which can be found in the literature [34], we provide here the expressions for the Christoffel symbols, the components of the Ricci and of the Weyl tensors pertaining to the Szekeres solutions and needed to complete the calculations described in this paper. These quantities have been obtained by running the SageMath software [62] with the simplifying notations of (71).

The non-zero Christoffel symbols are

$$\begin{aligned}\Gamma_{rr}^t &= HH_{,t} & \Gamma_{pp}^t &= \Gamma_{qq}^t = FF_{,t}, & \Gamma_{rt}^r &= \frac{H_{,t}}{H}, & \Gamma_{rr}^r &= \frac{H_{,r}}{H}, & \Gamma_{rp}^r &= \frac{H_{,p}}{H}, & \Gamma_{rq}^r &= \frac{H_{,q}}{H}, \\ \Gamma_{pp}^r &= \Gamma_{qq}^r = -\frac{FF_{,r}}{H^2}, & \Gamma_{pt}^p &= \Gamma_{qt}^q = \frac{F_{,t}}{F}, & \Gamma_{rp}^p &= \Gamma_{rq}^q = \frac{F_{,r}}{F}, & \Gamma_{rr}^p &= -\frac{HH_{,p}}{F^2}, & \Gamma_{rr}^q &= -\frac{HH_{,q}}{F^2}, \\ \Gamma_{pq}^p &= \Gamma_{qq}^q = -\Gamma_{pp}^q = \frac{F_{,q}}{F}, & \Gamma_{pp}^p &= -\Gamma_{qq}^p = \Gamma_{pq}^q = \frac{F_{,p}}{F}.\end{aligned}$$

The non-zero components of the Ricci tensor are

$$\begin{aligned}R_{tt} &= -\left(\frac{H_{,tt}}{H} + 2\frac{F_{,tt}}{F}\right), & R_{tr} &= 2\left(\frac{H_{,t}F_{,r}}{HF} - \frac{F_{,tr}}{F}\right), \\ R_{tp} &= -\frac{F_{,tp}}{F} - \frac{H_{,tp}}{H} + \frac{F_{,t}F_{,p}}{F^2} + \frac{H_{,p}F_{,t}}{HF}, & R_{tq} &= -\frac{F_{,tq}}{F} - \frac{H_{,tq}}{H} + \frac{F_{,t}F_{,q}}{F^2} + \frac{H_{,q}F_{,t}}{HF}, \\ R_{rr} &= 2\frac{H_{,t}F_{,t}}{F} + H_{,tt} - 2\frac{F_{,rr}}{F} + 2\frac{H_{,r}F_{,r}}{HF} - \frac{H}{F^2}(H_{,pp} + H_{,qq}), \\ R_{rp} &= -\frac{F_{,rp}}{F} + \frac{F_{,r}F_{,p}}{F^2} + \frac{H_{,p}F_{,r}}{HF}, & R_{rq} &= -\frac{F_{,rq}}{F} + \frac{F_{,r}F_{,q}}{F^2} + \frac{H_{,q}F_{,r}}{HF}, \\ R_{pp} &= (F_{,t})^2 + FF_{,tt} + \frac{FH_{,t}F_{,t}}{H} - \frac{(F_{,r})^2}{H^2} - \frac{FF_{,rr}}{H^2} + \frac{(F_{,p})^2}{F^2} \\ &\quad - \frac{F_{,pp}}{F} + \frac{(F_{,q})^2}{F^2} - \frac{F_{,qq}}{F} + \frac{FH_{,r}F_{,r}}{H^3} + \frac{H_{,p}F_{,p}}{HF} - \frac{H_{,pp}}{H} - \frac{H_{,q}F_{,q}}{HF}, \\ R_{pq} &= \frac{H_{,p}F_{,q}}{HF} - \frac{H_{,pq}}{H} + \frac{H_{,q}F_{,p}}{HF}, \\ R_{qq} &= (F_{,t})^2 + FF_{,tt} + \frac{FH_{,t}F_{,t}}{H} - \frac{(F_{,r})^2}{H^2} - \frac{FF_{,rr}}{H^2} + \frac{(F_{,p})^2}{F^2} \\ &\quad - \frac{F_{,pp}}{F} + \frac{(F_{,q})^2}{F^2} - \frac{F_{,qq}}{F} + \frac{FH_{,r}F_{,r}}{H^3} - \frac{H_{,p}F_{,p}}{HF} - \frac{H_{,qq}}{H} + \frac{H_{,q}F_{,q}}{HF}.\end{aligned}$$

The non-zero components of the Weyl tensor are

$$\begin{aligned}C_{trtr} &= \frac{H}{3F^2} \left\{ -H(F_{,t})^2 + HFF_{,tt} + FH_{,t}F_{,t} - F^2H_{,tt} + \frac{(F_{,r})^2}{H} - \frac{F}{H}F_{,rr} \right. \\ &\quad \left. - \frac{H}{F^2} [(F_{,p})^2 + (F_{,q})^2] + \frac{H}{F}(F_{,pp} + F_{,qq}) + \frac{FH_{,r}F_{,r}}{H^2} - \frac{1}{2}(H_{,pp} + H_{,qq}) \right\}, \\ C_{trtp} &= \frac{1}{2F} \left(-F_{,rp} + \frac{F_{,r}F_{,p}}{F} + \frac{H_{,p}F_{,r}}{H} \right), & C_{trtq} &= \frac{1}{2F} \left(-F_{,rq} + \frac{F_{,r}F_{,q}}{F} + \frac{H_{,q}F_{,r}}{H} \right),\end{aligned}$$

$$C_{tptp} = \frac{1}{6} \left\{ (F_{,t})^2 - FF_{,tt} - \frac{FH_{,t}F_{,t}}{H} + \frac{F^2H_{,tt}}{H} - \frac{(F_{,r})^2}{H^2} + \frac{FF_{,rr}}{H^2} + \frac{1}{F^2} [(F_{,p})^2 + (F_{,q})^2] \right. \\ \left. - \frac{1}{F} (F_{,pp} + F_{,qq}) - \frac{FH_{,r}F_{,r}}{H^3} + \frac{3}{HF} (H_{,p}F_{,p} - H_{,q}F_{,q}) - \frac{H_{,pp}}{H} + 2\frac{H_{,qq}}{H} \right\},$$

$$C_{tptq} = \frac{1}{2H} \left(-H_{,pq} + \frac{H_{,p}F_{,q}}{F} + \frac{H_{,q}F_{,p}}{F} \right),$$

$$C_{tqtq} = \frac{1}{6} \left\{ (F_{,t})^2 - FF_{,tt} - \frac{FH_{,t}F_{,t}}{H} + \frac{F^2H_{,tt}}{H} - \frac{(F_{,r})^2}{H^2} + \frac{FF_{,rr}}{H^2} + \frac{1}{F^2} [(F_{,p})^2 + (F_{,q})^2] \right. \\ \left. - \frac{1}{F} (F_{,pp} + F_{,qq}) - \frac{FH_{,r}F_{,r}}{H^3} + \frac{3}{HF} (-H_{,p}F_{,p} + H_{,q}F_{,q}) + 2\frac{H_{,pp}}{H} - \frac{H_{,qq}}{H} \right\},$$

$$C_{rttr} = \frac{H^2}{3F^2} \left\{ (F_{,t})^2 - FF_{,tt} - \frac{FH_{,t}F_{,t}}{H} + \frac{F^2H_{,tt}}{H} - \frac{(F_{,r})^2}{H^2} + \frac{FF_{,rr}}{H^2} + \frac{1}{F^2} [(F_{,p})^2 + (F_{,q})^2] \right. \\ \left. - \frac{1}{F} (F_{,pp} + F_{,qq}) - \frac{FH_{,r}F_{,r}}{H^3} + \frac{1}{2H} (H_{,pp} + H_{,qq}) \right\},$$

$$C_{trpr} = \frac{H^2}{2F} \left(F_{,tp} - \frac{F_{,t}F_{,p}}{F} - \frac{FH_{,tp}}{H} + \frac{H_{,p}F_{,t}}{H} \right), \quad C_{trqr} = \frac{H^2}{2F} \left(F_{,tq} - \frac{F_{,t}F_{,q}}{F} - \frac{FH_{,tq}}{H} + \frac{H_{,q}F_{,t}}{H} \right),$$

$$C_{rppr} = H^2 C_{tptp} - \frac{H}{F} (H_{,p}F_{,p} - H_{,q}F_{,q}) + \frac{H}{2} (H_{,pp} - H_{,qq}),$$

$$C_{rqqr} = H^2 C_{tqtq} + \frac{H}{F} (H_{,p}F_{,p} - H_{,q}F_{,q}) - \frac{H}{2} (H_{,pp} - H_{,qq}),$$

$$C_{rppq} = -F^2 C_{tptq}, \quad C_{rqpq} = F^2 C_{trtq}, \quad C_{rqqq} = F^2 C_{trtp}, \quad C_{qpqp} = -\frac{F^4}{H^2} C_{trtr}.$$

- [1] E. Abdalla, G. F. Abellán, A. Aboubrahim *et al.*, Cosmology intertwined: A review of the particle physics, astrophysics, and cosmology associated with the cosmological tensions and anomalies, *J. High En. Astrophys.* **34**, 49 (2022).
- [2] M. L. McClure and C. Hellaby, Determining the metric of the cosmos: stability, accuracy, and consistency, *Phys. Rev. D* **78**, 044005 (2008).
- [3] C. Hellaby and A. H. A. Alfedeel, Solving the observer metric, *Phys. Rev. D* **79**, 043501 (2009).
- [4] P. Szekeres, A class of inhomogeneous cosmological models, *Commun. Math. Phys.* **41**, 55 (1975a).
- [5] C. Hellaby, The null and KS limits of the Szekeres model, *Class. Quantum Grav.* **13**, 2537 (1996).
- [6] C. Hellaby and A. Krasinski, Physical and geometrical interpretation of the $\epsilon \leq 0$ Szekeres models, *Phys. Rev. D* **77**, 023529 (2008).
- [7] J. Plebański and A. Krasinski, *An Introduction to General Relativity and Cosmology* (Cambridge University Press, Cambridge, England, 2d edition, 2024).
- [8] K. Bolejko, A. Krasinski, C. Hellaby, and M.-N. Célérier, *Structures in the Universe by exact methods: formation, evolution, interactions* Cambridge Monographs on Mathematical Physics (Cambridge University Press, Cambridge, England, 2010).
- [9] C. Hellaby and A. Krasinski, You cannot get through Szekeres wormholes: Regularity, topology, and causality in quasispherical Szekeres models, *Phys. Rev. D* **66**, 084011 (2002).
- [10] A. Krasinski, Geometry and topology of the quasiplane Szekeres model, *Phys. Rev. D* **78**, 064038 (2008).
- [11] A. Krasinski and K. Bolejko, Geometry of the quasi-hyperbolic Szekeres models, *Phys. Rev. D* **86**, 104036 (2012).
- [12] I. Georg and C. Hellaby, Symmetry and equivalence in Szekeres models, *Phys. Rev. D* **95**, 124016 (2017).
- [13] A. Krasinski and K. Bolejko, Redshift propagation equations in the $\beta' \neq 0$ Szekeres models, *Phys. Rev. D* **83**,

- 083503 (2011).
- [14] A. Nwankwo, M. Ishak, and J. Thompson, Luminosity distance and redshift in the Szekeres inhomogeneous cosmological models, *JCAP* 05(2011)028.
- [15] H. Bondi, Spherically Symmetrical Models in General Relativity, *Mon. Not. R. Astron. Soc.* **107**, 410 (1947). Reprinted in *Gen. Relativ. Gravit.* **31**, 1783 (1999), with an editorial note by A. Krasinski, *Gen. Relativ. Gravit.* **31**, 1777 (1999).
- [16] M. Ishak, J. Richardson, D. Garred, D. Whittington, A. Nwankwo, and R. Sussman, Dark energy or apparent acceleration due to a relativistic cosmological model more complex than the Friedmann-Lemaître-Robertson-Walker model?, *Phys. Rev. D* **78**, 123531 (2008); **84** 089902(E) (2011).
- [17] R. K. Sachs, Gravitational waves in general relativity. 6 The outgoing radiation condition, *Proc. Roy. Soc. London A* **264** 309 (1961).
- [18] I. M. H. Etherington, LX. On the definition of distance in general relativity, *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science* **15**, 761 (1933). Reprinted in *Gen. Relativ. Gravit.* **39**, 1055 (2007), with an editorial note by G. F. R. Ellis, *Gen. Relativ. Gravit.* **39**, 1047 (2007).
- [19] S. Perlmutter, S. Gabi, G. Goldhaber, *et al.*, Measurements of the Cosmological Parameters Ω and Λ from the First Seven Supernovae at $z \geq 0.35$, *Astrophys. J.* **483**, 565 (1997).
- [20] M.-N. Célérier, K. Bolejko, and A. Krasinski, A (giant) void is not mandatory to explain away dark energy with a Lemaître-Tolman model, *Astron. Astrophys.* **518**, A21 (2010).
- [21] R. G. Buckley and E. M. Schlegel, CMB dipoles and other low-order multipoles in the quasispherical Szekeres model, *Phys. Rev. D* **87**, 023524 (2013).
- [22] P. Szekeres, Quasispherical gravitational collapse, *Phys. Rev. D* **12**, 2941 (1975b).
- [23] M. M. de Souza, Hidden symmetries of Szekeres quasispherical solutions, *Revista Brasileira de Física* **15**, 379 (1985).
- [24] N. J. Secrest, S. von Hausegger, M. Rameez, R. Mohayaee, S. Sarkar, and J. Colin, A Test of the Cosmological Principle with Quasars, *Astrophys. J.* **908**, L51 (2021).
- [25] C. Dalang, R. Durrer, and F. Lacasa, Statistical effects of the observer's peculiar velocity on source number counts, *Phys. Rev. D* **107**, 103514 (2023).
- [26] R. A. Sunyaev and Ya. B. Zel'dovich, Small-Scale Fluctuations of Relic Radiation, *Astrophys. and Space Science* **7**, 3 (1970).
- [27] M. Birkinshaw, J. P. Hugues, and K. A. Arnaud, A Measurement of the Value of the Hubble Constant from the X-Ray Properties and the Sunyaev-Zel'dovich Effect of Abell 665, *Astrophys. J.* **379**, 466 (1991).
- [28] W. Hu and N. Sugiyama, Small-scale cosmological perturbations: an analytic approach, *Astrophys. J.* **471**, 542 (1996).
- [29] D. J. Eisenstein and W. Hu, Baryonic features in the matter transfer function, *Astrophys. J.* **496**, 605 (1998).
- [30] DESI collaboration, DESI 2024 VI: Cosmological Constraints from the Measurements of Baryon Acoustic Oscillations, arXiv:2404.03002 [astro-ph.CO].
- [31] N. Padmanabhan and M. White, Constraining anisotropic baryon oscillations, *Phys. Rev. D* **77**, 123540 (2008).
- [32] P. Schneider, J. Ehlers, and E. E. Falco, *Gravitational Lenses*, (Springer -Verlag, New York, Berlin, Heidelberg, 1992).
- [33] S. Seitz, P. Schneider, and J. Ehlers, Light propagation in arbitrary spacetimes and the gravitational lens approximation, *Class. Quantum Grav.* **11**, 2345 (1994).
- [34] M. A. Troxel, M. Ishak, and A. Peel, The effects of structure anisotropy on lensing observables in an exact general relativistic setting for precision cosmology, *JCAP*03 (2014)040.
- [35] C. Clarkson, G. F. R. Ellis, A. Faltenbacher, R. Maartens, O. Umeh, and J.-P. Uzan, (Mis)interpreting supernovae observations in a lumpy universe, *Mon. Not. Roy. Astron. Soc.* **426**, 1121 (2012).
- [36] N. Kaiser, Clustering in real space and in redshift space, *Mon. Not. Roy. Astron. Soc.* **227**, 1 (1987).
- [37] J. C. Jackson, A Critique of Rees's Theory of Primordial Gravitational Radiation, *Mon. Not. Roy. Astron. Soc.* **156**, 1P (1972).
- [38] H. Chen, J. Wang, T. Mao, *et al.* Estimation of line-of-sight velocities of individual galaxies using neural network I. Modelling redshift-space distortions at large scales, arXiv: 23.03468 [astro-ph].
- [39] P. Mishra, M.-N. Célérier, and T. P. Singh, Redshift drift in axially symmetric quasispherical Szekeres models, *Phys. Rev. D* **86**, 083520 (2012).
- [40] P. Mishra, Redshift and redshift drift in $\Lambda = 0$ quasispherical Szekeres cosmological models and the effect of averaging, *Phys. Rev. D* **105**, 063520 (2022).
- [41] A. Sandage, The change of redshift and apparent luminosity of galaxies due to the deceleration of selected expanding universes, *Astrophys. J.* **136**, 319 (1962).
- [42] G. C. McVittie, Appendix to The change of redshift and apparent luminosity of galaxies due to the deceleration of selected expanding universes, *Astrophys. J.* **136**, 334 (1962).
- [43] O. Wucknitz, L. G. Spitler, and U.-L. Pen, Cosmology with gravitationally lensed repeating fast radio bursts, *Astron. Astrophys.* **645**, A44 (2021).
- [44] C. Wang, K. Bolejko, and G. F. Lewis, The Instantaneous Redshift Difference of Gravitationally Lensed Images: Theory and Observational Prospects, *Astrophys. J.* **940**, 16 (2022).
- [45] P. Helbig, Redshift drift and strong gravitational lensing, *Mon. Not. Roy. Astron. Soc.* **519**, 2769 (2023).
- [46] J. Ehlers, Beiträge zur relativistischen Mechanik kontinuierlicher Medien [Contributions to the relativistic mechanics of continuous media], *Abhandlungen der Mathematisch-Naturwissenschaftlichen Klasse der Akademie der Wissenschaften und Literatur Mainz*, No 11; English translation: *Gen. Relativ. Gravit.* **25**, 1225 (1993).
- [47] M.-N. Célérier, Do we really see a cosmological constant in the supernovae data, *Astron. Astrophys.* **353**, 63 (2000).
- [48] K. Bolejko, M. Ashan Nazer, and D. L. Wiltshire, Different cosmic expansion and the Hubble flow anisotropy, *JCAP*06(2016)035.
- [49] A. Krasinski and K. Bolejko, Exact Inhomogeneous Models and the Drift of Light Rays Induced by Nonsymmetric Flow of the Cosmic Medium, *Int. J. Mod. Phys. D* **22**, 1330013 (2013).
- [50] C. Quercellini, M. Quartin, and L. Amendola, Possibil-

- ity of Detecting Anisotropic Expansion of the Universe by Very Accurate Astrometry Measurements, *Phys. Rev. Lett.* **102**, 151302 (2009).
- [51] C. Quercellini, L. Amendola, A. Balbi, P. Cabella, and M. Quartin, Real-time cosmology, *Phys. Rep.* **521**, 95 (2012).
- [52] A. Krasinski, Spacetimes with no position drift, *Acta Physica Polonica B* **54**, 2-A1 (2023).
- [53] A. Krasinski, Existence of blueshifts in quasispherical Szekeres spacetimes, *Phys. Rev. D* **94**, 023515 (2016).
- [54] A. Krasinski, Gamma Radiation from Areal Radius Minima in a Quasi-Spherical Szekeres Metric, *Acta Physica Polonica B* **51**, 483 (2020).
- [55] K. Bolejko, The Szekeres Swiss Cheese model and the CMB observations, *Gen. Relativ. Gravit.* **41**, 1737 (2009).
- [56] K. Bolejko and M.-N. C el erier, Szekeres Swiss-cheese model and supernova observations, *Phys. Rev. D* **82**, 103510 (2010).
- [57] K. Bolejko, M.-N. C el erier, and A. Krasinski, Inhomogeneous cosmological models: exact solutions and their applications, *Class. Quantum Grav.* **28**, 164002 (2011).
- [58] M. A. Troxel, A. Peel, and M. Ishak, Effects of anisotropy on gravitational infall in galaxy clusters using an exact general relativistic model, *JCAP12* (2013)048.
- [59] M. Ishak, A. Peel, and M. A. Troxel, Stringent Restriction from the Growth of Large-Scale Structures on Apparent Acceleration in Inhomogeneous Cosmological Models, *Phys. Rev. Lett.* **111**, 251302 (2013).
- [60] A. Peel, M. A. Troxel, and M. Ishak, Effects of inhomogeneities on high precision measurements of cosmological distances, *Phys. Rev. D* **90**, 123536 (2014).
- [61] R. Luna, D. D. Doneva, J. A. Font, Jr-H. Lien, and S. S. Yazadjiev, Quasinormal modes in modified gravity using physics-informed neural networks, *Phys.Rev.D* **109**, 124064 (2024).
- [62] SageMath, the Sage Mathematics Software System, <https://www.sagemath.org>.