

β -DIMENSIONAL SHARP MAXIMAL FUNCTION AND ITS APPLICATIONS

YOU-WEI BENSON CHEN AND ALEJANDRO CLAROS

ABSTRACT. In this paper, we study β -dimensional sharp maximal operator defined as

$$\mathcal{M}_\beta^\# f(x) := \sup_Q \inf_{c \in \mathbb{R}} \chi_Q(x) \frac{1}{\ell(Q)^\beta} \int_Q |f - c| d\mathcal{H}_\infty^\beta,$$

where the supremum is taken over all cubes in \mathbb{R}^d with sides parallel to the coordinate axes, $\ell(Q)$ is the length side of Q and \mathcal{H}_∞^β is the Hausdorff content. In particular, we prove Fefferman-Stein inequality for $\mathcal{M}_\beta^\# f$ by giving a good lambda estimate for β -dimensional sharp maximal operator in the context of Hausdorff content. Additionally, we prove the Muckenhoupt-Wheeden inequality in this framework by establishing a good lambda inequality of independent interest.

1. INTRODUCTION

In the recent paper [8], a John-Nirenberg inequality is established within the framework of Hausdorff content. Specifically, for $\beta \in (0, d]$, the condition

$$(1.1) \quad \|u\|_{\text{BMO}^\beta(Q_0)} := \sup_{Q \subset Q_0} \inf_{c \in \mathbb{R}} \frac{1}{\ell(Q)^\beta} \int_Q |u - c| d\mathcal{H}_\infty^\beta < +\infty,$$

is shown to be equivalent to the following exponential decay estimate:

$$(1.2) \quad \mathcal{H}_\infty^\beta(\{x \in Q : |u(x) - c_Q| > t\}) \leq C_\beta \ell(Q)^\beta \exp(-c_\beta t),$$

for some constants $c_\beta, C_\beta > 0$, for all $t > 0$, for all subcubes $Q \subset Q_0$, and for a suitable choice of $c_Q \in \mathbb{R}$. In here, \mathcal{H}_∞^β denotes the Hausdorff content defined as

$$\mathcal{H}_\infty^\beta(E) := \inf \left\{ \sum_{i=1}^{\infty} \omega_\beta r_i^\beta : E \subset \bigcup_{i=1}^{\infty} B(x_i, r_i) \right\},$$

where $\omega_\beta := \pi^{\beta/2} / \Gamma(\frac{\beta}{2} + 1)$ is a normalized constant. When $\beta = d$, this result recovers the classical John-Nirenberg theorem given in [19]. For $\beta \in (0, d)$, this result demonstrates a self-improving property for functions whose bounded mean oscillation is controlled on subspaces of dimension β , quantified via the Hausdorff content. The motivation to define such a space arose from the consideration of certain, sharp forms of the Sobolev embedding in the critical exponent [1, 11, 13, 20, 28], as in this regime one often finds not only the well-known exponential integrability estimate but even estimates of this type along lower dimensional sets. The main contribution of [8] was the observation that the

maximal function $\mathcal{M}_{\mathcal{H}_\infty^{\beta,Q}}$ with respect to dyadic Hausdorff content $\mathcal{H}_\infty^{\beta,Q}$ (see Section 2 for definitions) exhibits the weak type (1, 1) estimate

$$\mathcal{H}_\infty^{\beta,Q} \left(\left\{ x \in \mathbb{R}^d : \mathcal{M}_{\mathcal{H}_\infty^{\beta,Q}} f > t \right\} \right) \leq \frac{C}{t} \int_{\mathbb{R}^d} |f| d\mathcal{H}_\infty^{\beta,Q}$$

despite the inherent nonlinearity of the Choquet integral with respect to $\mathcal{H}_\infty^{\beta,Q}$.

Introduced by Fefferman and Stein in [14], the sharp maximal function is defined as

$$\mathcal{M}^\# f(x) := \sup_Q \chi_Q(x) \frac{1}{|Q|} \int_Q |f - f_Q| dy,$$

where $f_Q = \frac{1}{|Q|} \int_Q f dy$. Moreover, the Fefferman-Stein inequality is established in [14] to study the intermediate spaces between BMO and $L^p(\mathbb{R}^d)$. Specifically, there exists a constant C , depending only on the dimension d and the exponent p , such that

$$(1.3) \quad \|\mathcal{M}f\|_{L^p(\mathbb{R}^d)} \leq C \|\mathcal{M}^\# f\|_{L^p(\mathbb{R}^d)} \quad \text{for } 1 < p < \infty.$$

Here, $\mathcal{M}f$ denotes the Hardy-Littlewood maximal function of f , defined as

$$\mathcal{M}f(x) := \sup_Q \frac{\chi_Q(x)}{|Q|} \int_Q |f| dy.$$

Additionally, the definition of BMO^β naturally leads to the introduction of the β -dimensional sharp maximal operator $\mathcal{M}_\beta^\#$ defined as follows.

Definition 1.1. *Let $0 < \beta \leq d \in \mathbb{N}$. The β -dimensional sharp maximal operator of a function f in \mathbb{R}^d is defined as*

$$(1.4) \quad \mathcal{M}_\beta^\# f(x) := \sup_Q \inf_{c \in \mathbb{R}} \chi_Q(x) \frac{1}{\ell(Q)^\beta} \int_Q |f - c| d\mathcal{H}_\infty^\beta$$

(see also [18, p. 16] for the β -dimensional sharp maximal function defined with respect to balls in Euclidean space).

It is easy to see that $\mathcal{M}_\beta^\# f \lesssim_{d,\beta} \mathcal{M}_{\mathcal{H}_\infty^\beta} f$, where $\mathcal{M}_{\mathcal{H}_\infty^\beta}$ denotes the β -dimensional maximal operator introduced in [9], defined as

$$\mathcal{M}_{\mathcal{H}_\infty^\beta} f(x) = \sup_{r>0} \frac{1}{\mathcal{H}_\infty^\beta(B(x,r))} \int_{B(x,r)} |f| d\mathcal{H}_\infty^\beta.$$

The first main result of this paper is an extension of (1.3) within the framework of $L^p(\mathcal{H}_\infty^\beta)$, we show the Fefferman-Stein inequality for the β -dimensional sharp maximal operator $\mathcal{M}_\beta^\# f$, which generalizes the result in (1.3). Our approach utilizes the good lambda inequality for the β -dimensional sharp maximal function $\mathcal{M}_\beta^\# f$ as stated in Theorem 3.1, combining the packing condition of Hausdorff content with the good lambda estimate in [14].

Theorem 1.2. *Let $0 < \beta \leq d \in \mathbb{N}$ and $0 < p_0 < \infty$. Let $f \in L_{loc}^1(\mathcal{H}_\infty^\beta)$ such that*

$$(1.5) \quad \sup_{0 < t \leq N} t^{p_0} \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : \mathcal{M}_{\mathcal{H}_\infty^\beta} f(x) > t\}) < \infty \quad \text{for all } N > 0.$$

Then there exists a constant $C = C(p, d, \beta) > 0$ such that

$$(1.6) \quad \left\| \mathcal{M}_{\mathcal{H}_\infty^\beta} f \right\|_{L^p(\mathcal{H}_\infty^\beta)} \leq C \left\| \mathcal{M}_\beta^\# f \right\|_{L^p(\mathcal{H}_\infty^\beta)}$$

for all $p_0 < p < \infty$, and also

$$(1.7) \quad \left\| \mathcal{M}_{\mathcal{H}_\infty^\beta} f \right\|_{L^{p,\infty}(\mathcal{H}_\infty^\beta)} \leq C \left\| \mathcal{M}_\beta^\# f \right\|_{L^{p,\infty}(\mathcal{H}_\infty^\beta)}$$

for all $p_0 \leq p < \infty$.

Remark 1.3. *The condition (1.5) is quite general, and it is satisfied, for example, for each function $f \in L^{p_0}(\mathcal{H}_\infty^\beta)$ if $p_0 \geq 1$.*

Let $0 < \alpha < d$. We recall that the Riesz potential of order α of a measure μ is defined by

$$I_\alpha \mu(x) := \frac{1}{\gamma(\alpha)} \int_{\mathbb{R}^d} \frac{d\mu(y)}{|x-y|^{d-\alpha}},$$

where $\gamma(\alpha) = \pi^{d/2} 2^\alpha \Gamma(\alpha/2) \Gamma((d-\alpha)/2)^{-1}$ is a normalization constant. We shall denote $I_\alpha f$ to represent $I_\alpha \mu$ where $d\mu(x) = f(x)dx$. It is well-known that B. Muckenhoupt and R. L. Wheeden proved in [21, Theorem 1] that for $1 < p < \infty$,

$$(1.8) \quad \|I_\alpha \mu\|_{L^p(\mathbb{R}^d)} \leq C \|\mathcal{M}_\alpha \mu\|_{L^p(\mathbb{R}^d)},$$

where the fractional maximal function $\mathcal{M}_\alpha \mu$ is defined as

$$\mathcal{M}_\alpha \mu(x) = \sup_{x \in Q} \frac{\mu(Q)}{\ell(Q)^{d-\alpha}}.$$

As before we shall denote $M_\alpha f$ to represent $M_\alpha \mu$ where $d\mu(x) = f(x)dx$. The proof of (1.8) employs the good lambda inequality for the fractional maximal function. Later, in [2], Adams provided an alternative proof of (1.8) with additional assumption $f \in L^r(\mathbb{R}^d)$ for some $r > 1$ based on the following pointwise estimate of independent interest:

$$(1.9) \quad \mathcal{M}^\#(I_\alpha f)(x) \cong \mathcal{M}_\alpha f(x),$$

for every nonnegative measurable function f in \mathbb{R}^d such that $I_\alpha f \in L^1_{loc}(\mathbb{R}^d)$.

Our second main result is a refinement of (1.9) in the context of the β -dimensional sharp maximal operator, formulated as follows:

Theorem 1.4. *Let f be a nonnegative measurable function in \mathbb{R}^d such that $I_\alpha f \in L^1_{loc}(\mathbb{R}^d)$. Then for $\beta \in (d-\alpha, d]$, it holds that*

$$\mathcal{M}_\beta^\#(I_\alpha f)(x) \cong \mathcal{M}_\alpha f(x).$$

Theorem 1.4 helps us to study the local behavior of functions of bounded β -dimensional mean oscillation, and it provides an analog of [10, Theorem 1.6] concerning the local Morrey space (see Section 2 for precise definition). More precisely, let Ω be an open set and $f \geq 0$ such that $I_\alpha f \in L^1_{loc}(\mathbb{R}^d)$. Then we have

$$\|I_\alpha f\|_{\text{BMO}^\beta(\Omega)} \cong \|f\|_{\mathcal{M}^\alpha(\Omega)} \quad \text{for } \beta \in (d-\alpha, d].$$

Theorem 1.2 and Theorem 1.4 naturally led one to inquire whether there is an analog of B. Muckenhoupt and R. L. Wheeden's result in (1.8) in the context of $L^p(\mathcal{H}_\infty^\beta)$ -boundedness. We now state the third result of this paper, which contains (1.8) as a particular case with $\beta = d$.

Theorem 1.5. *Let $0 < \alpha < d \in \mathbb{N}$, $\beta \in (d - \alpha, d]$, $1 \leq p < \infty$ and μ be a locally finite measure. Then there exists $C = C(d, \alpha, \beta) > 0$ such that*

$$(1.10) \quad \|I_\alpha \mu\|_{L^p(\mathcal{H}_\infty^\beta)} \leq C p \|\mathcal{M}_\alpha \mu\|_{L^p(\mathcal{H}_\infty^\beta)},$$

and

$$(1.11) \quad \|I_\alpha \mu\|_{L^{p,\infty}(\mathcal{H}_\infty^\beta)} \leq C p \|\mathcal{M}_\alpha \mu\|_{L^{p,\infty}(\mathcal{H}_\infty^\beta)}.$$

Remark 1.6. *The linear dependence on p is a consequence of the exponential decay in our good lambda inequality, stated in Theorem 5.3.*

As a consequence, we present the following corollary, which extends [2, Corollary (iii)] to the $L^p(\mathcal{H}_\infty^\beta)$ setting.

Corollary 1.7. *Let $0 < \alpha < d \in \mathbb{N}$, $r > 1$, $\beta \in (d - \alpha, d]$ and $f \in L^r(\mathbb{R}^d)$ be a non-negative function. Then the functions $I_\alpha f$, $\mathcal{M}_\alpha f$, and $\mathcal{M}_\beta^\#(I_\alpha f)$ have comparable norms in $L^p(\mathcal{H}_\infty^\beta)$ for $1 < p < \infty$.*

This paper is organized as follows: Section 2 reviews the necessary preliminaries for defining the Choquet integral with respect to Hausdorff content and discusses their basic properties, such as Hölder's inequality and the quasi-linearity of $L^1(\mathcal{H}_\infty^\beta)$. This section also covers the weak type $(1, 1)$ estimate for dyadic maximal function and a minor modification of the Calderón-Zygmund decomposition with respect to the dyadic Hausdorff content presented in [8, Theorem 3.1]. In Section 3, we give the good lambda estimates for the β -dimensional sharp maximal function $\mathcal{M}_\beta^\# f$, and we use it to prove Theorem 1.2. In section 4, we prove the pointwise estimate in Theorem 1.4 and use this result to provide an alternative proof of Theorem 1.5 with additional assumption, following the approach in [2]. Finally, in Section 5, we give the good lambda estimates for the fractional maximal function $\mathcal{M}_\alpha \mu$ and we prove Theorem 1.5.

2. PRELIMINARIES

We begin with recalling the definition of the Choquet integral with respect to an outer measure H . Let H be an outer measure, that is, H satisfies the following properties:

- (i) $H(\emptyset) = 0$;
- (ii) If $E \subseteq F$, then $H(E) \leq H(F)$;
- (iii) If $E \subseteq \bigcup_{i=1}^\infty E_i$, then

$$H(E) \leq \sum_{i=1}^\infty H(E_i).$$

The Choquet integral of a nonnegative function f over a set Ω with respect to outer measure H is then defined as:

$$(2.1) \quad \int_\Omega f dH := \int_0^\infty H(\{x \in \Omega : f(x) > t\}) dt.$$

A utilitarian way to deal with the Choquet integral with respect to the Hausdorff content \mathcal{H}_∞^β is to move between the dyadic Hausdorff content $\mathcal{H}_\infty^{\beta,Q}$ which is defined as follows:

Definition 2.1. Let $0 < \beta < d \in \mathbb{N}$ and Q be a cube in \mathbb{R}^d . Then for every $E \subseteq \mathbb{R}^d$, we define the dyadic Hausdorff content $\mathcal{H}_\infty^{\beta, Q}$ of E as

$$(2.2) \quad \mathcal{H}_\infty^{\beta, Q}(E) := \inf \left\{ \sum_{i=1}^{\infty} \ell(Q_i)^\beta : E \subset \bigcup_{i=1}^{\infty} Q_i, Q_i \in \mathcal{D}(Q) \right\}$$

where $\mathcal{D}(Q)$ denotes the collection of all dyadic cube generated by Q .

While the Hausdorff content \mathcal{H}_∞^β is not strongly subadditive if $\beta < d$, it can be shown that the dyadic Hausdorff content $\mathcal{H}_\infty^{\beta, Q}$ is strongly subadditive for any cube $Q \subseteq \mathbb{R}^d$ and thus satisfies

$$(2.3) \quad \int_{\mathbb{R}^d} \sum_{j=1}^{\infty} f_j d\mathcal{H}_\infty^{\beta, Q} \leq \sum_{j=1}^{\infty} \int f_j d\mathcal{H}_\infty^{\beta, Q}$$

(see [24, Proposition 3.5 and 3.6] for usual dyadic case and [8, Proposition 2.6 and Proposition 2.10] for general cube Q). Moreover, there exists a constant $C_\beta > 0$ such that for every cube $Q \subseteq \mathbb{R}^d$,

$$(2.4) \quad \frac{1}{C_\beta} \mathcal{H}_\infty^{\beta, Q}(E) \leq \mathcal{H}_\infty^\beta(E) \leq C_\beta \mathcal{H}_\infty^{\beta, Q}(E)$$

for all $E \subset \mathbb{R}^d$ (see Proposition 2.3 in [27] and Proposition 2.11 in [8]) and thus

$$(2.5) \quad \int_{\mathbb{R}^d} \sum_{j=1}^{\infty} f_j d\mathcal{H}_\infty^\beta \leq C_\beta^2 \sum_{j=1}^{\infty} \int_{\mathbb{R}^d} f_j d\mathcal{H}_\infty^\beta$$

by definition of Choquet integral (2.1).

Next, we outline the basic properties of the Choquet integral with respect to the Hausdorff content. The following lemma can be found in [15, p. 5], with its proof referred to in [4] and [5, Chapter 4].

Lemma 2.2. Let $0 < \beta \leq d \in \mathbb{R}^d$ and $\Omega \subseteq \mathbb{R}^d$. Then the following statements hold:

(i) For $a \geq 0$ and nonnegative functions f in \mathbb{R}^d , we have

$$\int_{\Omega} a f(x) d\mathcal{H}_\infty^\beta = a \int_{\Omega} f(x) d\mathcal{H}_\infty^\beta;$$

(ii) For nonnegative functions f_1 and f_2 in \mathbb{R}^d , we have

$$\int_{\Omega} f_1(x) + f_2(x) d\mathcal{H}_\infty^\beta \leq 2 \left(\int_{\Omega} f_1(x) d\mathcal{H}_\infty^\beta + \int_{\Omega} f_2(x) d\mathcal{H}_\infty^\beta \right);$$

(iii) Let $1 < p < \infty$, and p' defined by $\frac{1}{p} + \frac{1}{p'} = 1$. Then for nonnegative functions f_1 and f_2 in \mathbb{R}^d , we have

$$\int_{\Omega} f_1(x) f_2(x) d\mathcal{H}_\infty^\beta \leq 2 \left(\int_{\Omega} f_1(x)^p d\mathcal{H}_\infty^\beta \right)^{\frac{1}{p}} \left(\int_{\Omega} f_2(x)^{p'} d\mathcal{H}_\infty^\beta \right)^{\frac{1}{p'}}.$$

Furthermore, the following estimates of weak and strong types hold:

Lemma 2.3. Let $0 < \eta < d$.

(i) If $1 < p < \frac{d}{\eta}$ and $\beta = d - \eta p$, then there exists a constant A_1 depending on d, η and p such that

$$\left(\int_{\mathbb{R}^d} (\mathcal{M}_\eta f)^p d\mathcal{H}_\infty^\beta \right)^{\frac{1}{p}} \leq A_1 \left(\int_{\mathbb{R}^d} |f|^p dx \right)^{\frac{1}{p}};$$

(ii) If $p = \frac{\beta}{d-\eta}$ with $d - \eta \leq \beta \leq d$, then there exists a constant A_2 depending on d, η and p such that

$$\|\mathcal{M}_\eta f\|_{L^{p,\infty}(\mathbb{R}^d; \mathcal{H}_\infty^\beta)} \leq A_2 \int_{\mathbb{R}^d} |f| dx.$$

In here, $L^{p,\infty}(\mathbb{R}^d; \mathcal{H}_\infty^\beta)$ denotes the vector spaces of all the functions $f : \mathbb{R}^d \rightarrow \mathbb{R}$ satisfying

$$\|f\|_{L^{p,\infty}(\mathbb{R}^d; \mathcal{H}_\infty^\beta)} := \sup_{\lambda > 0} \lambda \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : |f(x)| > \lambda\})^{\frac{1}{p}} < \infty.$$

The demonstration of assertion (i) is based on the one weight inequality established in [25, Theorem B]. For assertion (ii), specifically when $\beta = n - \eta$, the derivation is given in [7, Lemma 3.2]. Moreover, both assertions (i) and (ii) are provided in [4, Theorem 7], which discusses these in the framework of Hausdorff content, as outlined in Lemma 2.3.

A function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is said to be in $L^p(\mathcal{H}_\infty^\beta; \Omega)$ provided f is \mathcal{H}_∞^β -quasi continuous (see [8, p. 2] for definition) and

$$\|f\|_{L^p(\mathcal{H}_\infty^\beta; \Omega)} := \left(\int_{\Omega} |f|^p d\mathcal{H}_\infty^\beta \right)^{\frac{1}{p}} < \infty.$$

In the rest of the paper, we adopt the convention $L^p(\mathcal{H}_\infty^\beta) = L^p(\mathcal{H}_\infty^\beta; \mathbb{R}^d)$. Moreover, it has been shown in [3] that

$$(2.6) \quad \|f\|_{L^1(\mathcal{H}_\infty^\beta; \Omega)} \asymp \sup_{\|\mu\|_{\mathcal{M}^\beta(\Omega)} \leq 1} \int_{\Omega} f d\mu.$$

In here, $\mathcal{M}^\beta(\Omega)$ denotes the vector space of all Radon measure in \mathbb{R}^d satisfying

$$\mathcal{M}^\beta(\Omega) := \left\{ \mu \in M_{loc}(\Omega) : \|\mu\|_{\mathcal{M}^\beta(\Omega)} := \sup_{x \in \Omega, r > 0} \frac{|\mu|(B(x, r) \cap \Omega)}{r^\beta} < \infty \right\},$$

where $M_{loc}(\Omega)$ is the set of locally finite Radon measures in Ω .

We now give the definition of the dyadic maximal function $\mathcal{M}_{\mathcal{H}_\infty^\beta, Q_0}$, which is introduced in [8].

Definition 2.4. Let $0 < \beta \leq d \in \mathbb{N}$ and Q_0 be a cube in \mathbb{R}^d . The dyadic maximal function associate to $\mathcal{H}_\infty^{\beta, Q_0}$ is defined as

$$\mathcal{M}_{\mathcal{H}_\infty^{\beta, Q_0}} f(x) := \sup_Q \frac{1}{\ell(Q)^\beta} \int_Q |f| d\mathcal{H}_\infty^{\beta, Q_0},$$

where the supremum is taken over all the dyadic cubes Q in $\mathcal{D}(Q_0)$ containing x .

In particular, a weak type (1, 1) estimate for dyadic maximal function proved in [8, Theorem 3.1] asserts that for any $t > 0$, the inequality

$$(2.7) \quad \mathcal{H}_\infty^{\beta, Q_0} \left(\left\{ x \in \mathbb{R}^d : \mathcal{M}_{\mathcal{H}_\infty^{\beta, Q_0}} f > t \right\} \right) \leq \frac{C}{t} \int_{\mathbb{R}^d} |f| d\mathcal{H}_\infty^{\beta, Q_0}$$

holds for all nonnegative function f .

Remark 2.5. Note that in [8, Theorem 3.1], f is assumed to be in $L^1(\mathcal{H}_\infty^\beta)$. However, one finds that the assumption

$$\int_{\mathbb{R}^d} |f| d\mathcal{H}_\infty^\beta < \infty$$

is sufficient for the weak type estimate (2.7) by just following the original proof. (That is, the assumption of f being quasi-continuous is not necessary for (2.7).)

The following lemma, concerning the comparison of the Hausdorff content of level sets of $\mathcal{M}_{\mathcal{H}_\infty^\beta}$ and $\mathcal{M}_{\mathcal{H}_\infty^{\beta, Q_0}}$, is given in [9, Lemma 2.4].

Lemma 2.6. Let Q_0 be a cube in \mathbb{R}^d . Then there exist constants $C = C(d, \beta)$ and $\tilde{C} = \tilde{C}(d, \beta)$ such that

$$\mathcal{H}_\infty^\beta \left(\left\{ x \in \mathbb{R}^d : \mathcal{M}_{\mathcal{H}_\infty^\beta} f(x) > t \right\} \right) \leq C \mathcal{H}_\infty^\beta \left(\left\{ x \in \mathbb{R}^d : \mathcal{M}_{\mathcal{H}_\infty^{\beta, Q_0}} f(x) > \tilde{C} 2^{-(\beta+d)} t \right\} \right)$$

holds for every nonnegative function f .

We next record a lemma concerning the packing condition of dyadic cubes, which is a small modification of [8, Proposition 2.11] and [22, Lemma 2].

Lemma 2.7. Suppose $\{Q_j\}$ is a family of non-overlapping dyadic cubes subordinate to some cube lattice $\mathcal{D}(Q)$. Then there exists a subfamily $\{Q_{j_k}\}$ and a family of non-overlapping ancestors \tilde{Q}_k such that

(1)

$$\bigcup_j Q_j \subset \bigcup_k Q_{j_k} \cup \bigcup_k \tilde{Q}_k$$

(2)

$$\sum_{Q_{j_k} \subset Q} \ell(Q_{j_k})^\beta \leq 2\ell(Q)^\beta, \text{ for each dyadic cube } Q.$$

(3)

$$\begin{aligned} \mathcal{H}_\infty^{\beta, Q_0}(\cup_j Q_j) &\leq \sum_{k, Q_{j_k} \not\subset \tilde{Q}_m} \ell(Q_{j_k})^\beta + \sum_k \ell(\tilde{Q}_k)^\beta \\ &\leq \sum_k \ell(Q_{j_k})^\beta \\ &\leq 2\mathcal{H}_\infty^{\beta, Q_0}(\cup_j Q_j). \end{aligned}$$

Proof. An application of Proposition 2.11 in [8] yields a subfamily $\{Q_{j_k}\}_k$ of $\{Q_j\}_j$ and a family of non-overlapping ancestors $\{\tilde{Q}_k\}_k$ which satisfy the properties (1), (2) and

$$\ell(\tilde{Q}_k)^\beta \leq \sum_{Q_{j_i} \subset \tilde{Q}_k} \ell(Q_{j_i})^\beta \text{ for each } \tilde{Q}_k.$$

Moreover, using the subadditive of $\mathcal{H}_\infty^{\beta, Q_0}$, we obtain

$$\begin{aligned} \mathcal{H}_\infty^{\beta, Q_0}(\cup_j Q_j) &\leq \sum_{k, Q_{j_k} \not\subseteq \tilde{Q}_m} \ell(Q_{j_k})^\beta + \sum_k \ell(\tilde{Q}_k)^\beta \\ &\leq \sum_{k, Q_{j_k} \not\subseteq \tilde{Q}_m} \ell(Q_{j_k})^\beta + \sum_{k, Q_{j_k} \subseteq \tilde{Q}_m} \ell(Q_{j_k})^\beta \\ (2.8) \quad &= \sum_k \ell(Q_{j_k})^\beta. \end{aligned}$$

Now, it is left to show that

$$(2.9) \quad \sum_k \ell(Q_{j_k})^\beta \leq 2\mathcal{H}_\infty^{\beta, Q_0}(\cup_j Q_j).$$

To this end, we let $\{Q_m\}_m \subseteq \mathcal{D}(Q_0)$ be a maximal collection of dyadic cubes such that

$$\bigcup_j Q_j \subseteq \bigcup_m Q_m.$$

The property (2) then yields that

$$\begin{aligned} \sum_k \ell(Q_{j_k})^\beta &= \sum_m \sum_{k, Q_{j_k} \subseteq Q_m} \ell(Q_{j_k})^\beta \\ &\leq \sum_m 2\ell(Q_m)^\beta \end{aligned}$$

and (2.9) follows by taking the infimum over all such families of dyadic cubes $\{Q_m\}_m$. \square

The condition (2) in Lemma 2.7 is called packing condition, and an important application is that for any $f \geq 0$, one has the estimate

$$(2.10) \quad \sum_k \int_{Q_{j_k}} f d\mathcal{H}_\infty^{\beta, Q_0} \leq 2 \int_{\cup_k Q_{j_k}} f \mathcal{H}_\infty^{\beta, Q_0}.$$

The following lemma restates the Calderón-Zygmund decomposition for dyadic Hausdorff content given in [8, Theorem 3.4]. This lemma will play an important role in the proof of Lemma 3.1, and we provide the proof here for the reader's convenience.

Lemma 2.8. *Let Q_0 be a cube and $\lambda > 0$. If*

$$\mathcal{H}_\infty^{\beta, Q_0}(\{x \in \mathbb{R}^d : \mathcal{M}_{\mathcal{H}_\infty^{\beta, Q_0}} f(x) > \lambda\}) < \infty,$$

then there exists a countable collection of non-overlapping dyadic cubes $\{Q_k\}$ subordinate to Q_0 such that

$$(i) \quad \bigcup_k Q_k = \{x \in \mathbb{R}^d : \mathcal{M}_{\mathcal{H}_\infty^{\beta, Q_0}} f(x) > \lambda\},$$

(ii) $\{Q_k\}$ is the maximal subcollection of all the dyadic cubes contained in

$$\{x \in \mathbb{R}^d : \mathcal{M}_{\mathcal{H}_\infty^{\beta, Q_0}} f(x) > \lambda\}.$$

(iii) $\lambda < \frac{1}{\ell(Q_k)^\beta} \int_{Q_k} |f| \mathcal{H}_\infty^{\beta, Q_0} \leq 2^\beta \lambda,$

(iv) $|f(x)| \leq \lambda$ for \mathcal{H}^β -a.e. $x \notin \bigcup_k Q_k,$

Proof. Let $\{Q_i\}_i$ be the collection of all the dyadic cubes contained in $\{x \in \mathbb{R}^d : \mathcal{M}_{\mathcal{H}_\infty^{\beta, Q_0}} f(x) > \lambda\}$. Since $\mathcal{H}_\infty^{\beta, Q_0}(\{x \in \mathbb{R}^d : \mathcal{M}_{\mathcal{H}_\infty^{\beta, Q_0}} f(x) > \lambda\}) < \infty$, the maximal subcollection of $\{Q_i\}_i$ exists and we denote it as $\{Q_k\}_k$. In particular, we have

$$\bigcup_k Q_k = \{x \in \mathbb{R}^d : \mathcal{M}_{\mathcal{H}_\infty^{\beta, Q_0}} f(x) > \lambda\}.$$

and thus

$$\lambda < \frac{1}{\ell(Q_k)^\beta} \int_{Q_k} |f| d\mathcal{H}_\infty^{\beta, Q_0}.$$

Now let \tilde{Q}_k denotes the parent of Q_k for each k . Using the fact \tilde{Q}_k is not contained in $x \in \mathbb{R}^d : \mathcal{M}_{\mathcal{H}_\infty^{\beta, Q_0}} f(x) > \lambda$, we have

$$\frac{1}{2^\beta \ell(Q_k)^\beta} \int_{Q_k} |f| d\mathcal{H}_\infty^{\beta, Q_0} \leq \frac{1}{\ell(\tilde{Q}_k)^\beta} \int_{\tilde{Q}_k} |f| d\mathcal{H}_\infty^{\beta, Q_0} \leq \lambda.$$

Finally, (iv) follows by the Lebesgue differential theorem with respect to dyadic Hausdorff content given in [8, Theorem 3.3].

□

3. GOOD LAMBDA ESTIMATES FOR THE β -DIMENSIONAL SHARP MAXIMAL FUNCTION AND PROOF OF THEOREM 1.2

In [14], Fefferman and Stein shows the $L^p(\mathbb{R}^d)$ boundeness of sharp maximal function $\mathcal{M}^\# f$ based on the good lambda estimate

$$(3.1) \quad |\{x \in \mathbb{R}^d : \mathcal{M}f > t\}| \leq |\{x \in \mathbb{R}^d : \mathcal{M}^\# f > \frac{t}{A}\}| + \frac{2}{A} |\{x \in \mathbb{R}^d : \mathcal{M}f > 2^{-n-1}t\}|$$

for all positive t and A . Here, we give an extension of (3.1) in the context of Hausdorff content whose proof is parallel to the one given in [14]. There are some difficulties due to the nonlinearity of the Choquet integral, which we overcome through the use of the packing condition in Lemma 2.7.

Theorem 3.1. *Let $0 < \beta \leq d \in \mathbb{N}$, Q_0 be a cube in \mathbb{R}^d , f be a function in \mathbb{R}^d and $\mu : (0, \infty) \rightarrow [0, \infty]$ be defined as*

$$(3.2) \quad \mu(t) := \mathcal{H}_\infty^{\beta, Q_0}(\{x \in \mathbb{R}^d : \mathcal{M}_{\mathcal{H}_\infty^{\beta, Q_0}} f(x) > t\}).$$

Then for any $t > 0$ and $A > 0$, we have

$$(3.3) \quad \mu(t) \leq \mathcal{H}_\infty^{\beta, Q_0}(\{x \in \mathbb{R}^d : \mathcal{M}_{\beta, Q_0}^\# f(x) > \frac{t}{A}\}) + \frac{8}{A} \mu(2^{-\beta-2}t),$$

where $\mathcal{M}_{\beta, Q_0}^\# f$ is the dyadic sharp maximal function defined by

$$\mathcal{M}_{\beta, Q_0}^\# f(x) := \sup_{Q \in \mathcal{D}(Q_0)} \inf_{c \in \mathbb{R}} \chi_Q(x) \frac{1}{\ell(Q)^\beta} \int_Q |f - c| d\mathcal{H}_\infty^{\beta, Q_0}.$$

Proof. Fix $t > 0$, we apply Lemma 2.8 and obtain countable collections of non-overlapping dyadic cubes $\{Q_k^t\}_k$ and $\{Q_k^{2^{-\beta-2}t}\}_k$ such that properties (i) to (iv) in Lemma 2.8 hold for $\lambda = t$ and $\lambda = 2^{-\beta-2}t$ respectively. That is, the collection of cubes $\{Q_k^t\}_k$ and $\{Q_k^{2^{-\beta-2}t}\}_k$ satisfy

- (A₁) $\bigcup_k Q_k^t = \{x \in \mathbb{R}^d : \mathcal{M}_{\mathcal{H}_\infty^{\beta, Q_0}} f(x) > t\}$,
- (A₂) $\{Q_k^t\}_k$ is the maximal subcollection of all the dyadic cubes contained in the set $\{x \in \mathbb{R}^d : \mathcal{M}_{\mathcal{H}_\infty^{\beta, Q_0}} f(x) > t\}$,
- (A₃) $t < \frac{1}{\ell(Q_k^t)^\beta} \int_{Q_k^t} |f| d\mathcal{H}_\infty^{\beta, Q_0} \leq 2^\beta t$,
- (A₄) $|f(x)| \leq t$ for \mathcal{H}^β -a.e. $x \notin \bigcup_k Q_k^t$

and

- (B₁) $\bigcup_k Q_k^{2^{-\beta-2}t} = \{x \in \mathbb{R}^d : \mathcal{M}_{\mathcal{H}_\infty^{\beta, Q_0}} f(x) > 2^{-\beta-2}t\}$,
- (B₂) $\{Q_k^{2^{-\beta-2}t}\}_k$ is the maximal subcollection of all the dyadic cubes contained in the set $\{x \in \mathbb{R}^d : \mathcal{M}_{\mathcal{H}_\infty^{\beta, Q_0}} f(x) > 2^{-\beta-2}t\}$,
- (B₃) $2^{-\beta-2}t < \frac{1}{\ell(Q_k^{2^{-\beta-2}t})^\beta} \int_{Q_k^{2^{-\beta-2}t}} |f| d\mathcal{H}_\infty^{\beta, Q_0} \leq \frac{t}{4}$,
- (B₄) $|f(x)| \leq 2^{-\beta-2}t$ for \mathcal{H}^β -a.e. $x \notin \bigcup_k Q_k^{2^{-\beta-2}t}$.

Since $t > 2^{-\beta-2}t$, we have the cubes in $\{Q_k^t\}_k$ are sub-cubes of the cubes in $\{Q_k^{2^{-\beta-2}t}\}_k$ by (A₂) and (B₂). Thus,

$$\{Q_k^t\}_k = \mathcal{A}_1 \cup \mathcal{A}_2,$$

where

$$\mathcal{A}_1 := \left\{ Q_k^t : Q_k^t \subseteq Q_{k_0}^{2^{-\beta-2}t} \text{ for some } k_0 \text{ and } Q_{k_0}^{2^{-\beta-2}t} \subseteq \{x \in \mathbb{R}^d : \mathcal{M}_{\beta, Q_0}^\# f(x) > \frac{t}{A}\} \right\},$$

and

$$\mathcal{A}_2 := \left\{ Q_k^t : Q_k^t \subseteq Q_{k_0}^{2^{-\beta-2}t} \text{ for some } k_0 \text{ and } Q_{k_0}^{2^{-\beta-2}t} \not\subseteq \{x \in \mathbb{R}^d : \mathcal{M}_{\beta, Q_0}^\# f(x) > \frac{t}{A}\} \right\}.$$

The subadditivity of $\mathcal{H}_\infty^{\beta, Q_0}$ then implies that

$$(3.4) \quad \mu(t) \leq \mathcal{H}_\infty^{\beta, Q_0} \left(\bigcup_{Q_k^t \in \mathcal{A}_1} Q_k^t \right) + \mathcal{H}_\infty^{\beta, Q_0} \left(\bigcup_{Q_k^t \in \mathcal{A}_2} Q_k^t \right).$$

It is easy to see that

$$(3.5) \quad \bigcup_{Q_k^t \in \mathcal{A}_1} Q_k^t \subseteq \left\{ x : \mathcal{M}_{\beta, Q_0}^\# f(x) > \frac{t}{A} \right\}$$

and thus

$$(3.6) \quad \mathcal{H}_\infty^{\beta, Q_0} \left(\bigcup_{Q_k^t \in \mathcal{A}_1} Q_k^t \right) \leq \mathcal{H}_\infty^{\beta, Q_0} \left(\left\{ x : \mathcal{M}_{\beta, Q_0}^\# f(x) > \frac{t}{A} \right\} \right).$$

Now the proof is complete if we show that

$$(3.7) \quad \mathcal{H}_\infty^{\beta, Q_0} \left(\bigcup_{Q_k^t \in \mathcal{A}_2} Q_k^t \right) \leq \frac{8}{A} \mu(2^{-\beta-2}t).$$

To this end, let $\{Q'_k\}_k$ be the subset of $\{Q_k^{2^{-\beta-2}t}\}_k$ which is defined as

$$\{Q'_k\}_k := \left\{ Q \in \{Q_k^{2^{-\beta-2}t}\}_k : Q \text{ contains some cube in } \mathcal{A}_2 \right\}.$$

Applying Lemma 2.7 to $\{Q'_k\}_k$, we then obtain a subfamily $\{Q'_{k_v}\}_v$ and a family of non-overlapping ancestors $\{\tilde{Q}_k\}_k$ such that

$$\bigcup_k Q'_k \subset \bigcup_v Q'_{k_v} \cup \bigcup_k \tilde{Q}_k,$$

$$\sum_{Q'_{k_v} \subset Q} \ell(Q'_{k_v})^\beta \leq 2\ell(Q)^\beta \text{ for each dyadic cube } Q \in \mathcal{D}(Q_0),$$

$$(3.8) \quad \begin{aligned} \mathcal{H}_\infty^{\beta, Q_0} \left(\bigcup_k Q'_k \right) &\leq \sum_{v, Q'_{k_v} \not\subseteq \tilde{Q}_k} \ell(Q'_{k_v})^\beta + \sum_k \ell(\tilde{Q}_k)^\beta \\ &\leq 2\mathcal{H}_\infty^{\beta, Q_0} \left(\bigcup_k Q'_k \right). \end{aligned}$$

In particular, let $Q' \in \{Q'_{k_v}\}_v \cup \{\tilde{Q}_k\}_k$, we have Q' contains a cube in $\mathcal{A}_2 \subseteq \{Q_k^{2^{-\beta-2}t}\}_k$ and thus

$$Q' \not\subseteq \left\{ x : \mathcal{M}_{\beta, Q_0}^\# f(x) > \frac{t}{A} \right\}.$$

It then follows from (B_2) , (B_3) and definition of $\mathcal{M}_{\beta, Q_0}^\# f$ that

$$(3.9) \quad \frac{1}{\ell(Q')^\beta} \int_{Q'} |f| d\mathcal{H}_\infty^{\beta, Q_0} \leq \frac{t}{4}$$

and

$$(3.10) \quad \frac{1}{\ell(Q')^\beta} \int_{Q'} |f - c_{Q'}| d\mathcal{H}_\infty^{\beta, Q_0} \leq \min \left(\frac{t}{A}, \frac{t}{4} \right),$$

where

$$c_{Q'} := \arg \min_{c \in \mathbb{R}} \frac{1}{\ell(Q')^\beta} \int_{Q'} |f - c| d\mathcal{H}_\infty^{\beta, Q_0}.$$

Moreover, the combination of (2.3), (3.9) and (3.10) gives

$$\begin{aligned}
|c_{Q'}| &= \frac{1}{\ell(Q')^\beta} \int_{Q'} |c_{Q'}| d\mathcal{H}_\infty^{\beta, Q_0} \\
&\leq \frac{1}{\ell(Q')^\beta} \int_{Q'} |c_{Q'} - f| d\mathcal{H}_\infty^{\beta, Q_0} + \frac{1}{\ell(Q')^\beta} \int_{Q'} |f| d\mathcal{H}_\infty^{\beta, Q_0} \\
&\leq \frac{t}{4} + \frac{t}{4} \\
&= \frac{t}{2}.
\end{aligned}$$

Thus, if $Q_k^t \in \mathcal{A}_2$ and $Q_k^t \subseteq Q'$, then

$$\begin{aligned}
(3.11) \quad \int_{Q_k^t} |f(x) - c_{Q'}| d\mathcal{H}_\infty^{\beta, Q_0} &\geq \int_{Q_k^t} |f| d\mathcal{H}_\infty^{\beta, Q_0} - c_{Q'} \ell(Q_k^t)^\beta \\
&\geq t \ell(Q_k^t)^\beta - \frac{t}{2} \ell(Q_k^t)^\beta \\
&= \frac{t}{2} \ell(Q_k^t)^\beta.
\end{aligned}$$

Now we apply Lemma 2.7 and (2.8) to \mathcal{A}_2 and obtain a non-overlapping subfamily $\{Q_{k_j}^t\}_j$ of \mathcal{A}_2 satisfying

$$(3.12) \quad \sum_{Q_{k_j}^t \subseteq Q} \ell(Q_{k_j}^t)^\beta \leq 2\ell(Q)^\beta \text{ for each dyadic cube } Q \in \mathcal{D}(Q_0)$$

and

$$(3.13) \quad \mathcal{H}_\infty^{\beta, Q_0} \left(\bigcup_{Q_k^t \in \mathcal{A}_2} Q_k^t \right) \leq \sum_v \ell(Q_{k_j}^t)^\beta.$$

Now combining estimates (3.11) and (3.13), we get

$$\begin{aligned}
(3.14) \quad \mathcal{H}_\infty^{\beta, Q_0} \left(\bigcup_{Q_k^t \in \mathcal{A}_2} Q_k^t \right) &\leq \sum_j \ell(Q_{k_j}^t)^\beta \\
&= \sum_v \sum_{Q_{k_j}^t \subseteq Q'_{k_v}} \ell(Q_{k_j}^t)^\beta + \sum_k \sum_{Q_{k_j}^t \subseteq \tilde{Q}_k} \ell(Q_{k_j}^t)^\beta \\
&\leq \frac{2}{t} \sum_v \sum_{Q_{k_j}^t \subseteq Q'_{k_v}} \int_{Q'_{k_v}} |f - c_{Q'_{k_v}}| d\mathcal{H}_\infty^{\beta, Q_0} \\
&\quad + \frac{2}{t} \sum_k \sum_{Q_{k_j}^t \subseteq \tilde{Q}_k} \int_{\tilde{Q}_k} |f - c_{\tilde{Q}_k}| d\mathcal{H}_\infty^{\beta, Q_0}, \\
&\leq \frac{4}{t} \sum_v \int_{Q'_{k_v}} |f - c_{Q'_{k_v}}| d\mathcal{H}_\infty^{\beta, Q_0} \\
&\quad + \frac{4}{t} \sum_k \int_{\tilde{Q}_k} |f - c_{\tilde{Q}_k}| d\mathcal{H}_\infty^{\beta, Q_0},
\end{aligned}$$

where the last inequality follows the packing condition (3.12) of $\{Q_{k_j}^t\}_j$ and (2.10). Finally, using the fact that $\{Q'_k\}_k$ is a subfamily of $\{Q_k^{2^{-\beta-2}t}\}_k$ and estimates (3.8), (3.10) and (3.14), we obtain

$$\begin{aligned} \mathcal{H}_\infty^{\beta, Q_0} \left(\bigcup_{Q_k^t \in \mathcal{A}_2} Q_k^t \right) &\leq \frac{4}{t} \sum_v \int_{Q'_{k_v}} |f - c_{Q'_{k_v}}| d\mathcal{H}_\infty^{\beta, Q_0} \\ &\quad + \frac{4}{t} \sum_k \int_{\tilde{Q}_k} |f - c_{\tilde{Q}_k}| d\mathcal{H}_\infty^{\beta, Q_0} \\ &\leq \frac{4}{t} \frac{t}{A} \left(\sum_v \ell(Q'_{k_v})^\beta + \sum_k \ell(\tilde{Q}_k)^\beta \right) \\ &\leq \frac{8}{A} \mathcal{H}_\infty^{\beta, Q_0} \left(\bigcup_k Q'_k \right) \\ &\leq \frac{8}{A} \mathcal{H}_\infty^{\beta, Q_0} \left(\bigcup_k Q_k^{2^{-\beta-2}t} \right) \\ &= \frac{8}{A} \mu(2^{-\beta-2}t), \end{aligned}$$

which completes the proof. \square

We next give the proof of Theorem 1.2.

Proof of Theorem 1.2. Let $Q_0 = [0, 1] \times \cdots \times [0, 1] \subset \mathbb{R}^d$. By Lemma 2.6, we deduce there exist $C_1, C_2, C_3 > 0$ such that

$$\begin{aligned} \int_{\mathbb{R}^d} (\mathcal{M}_{\mathcal{H}_\infty^\beta} f)^p d\mathcal{H}_\infty^\beta &= p \int_0^\infty t^{p-1} \mathcal{H}_\infty^\beta \left(\{x \in \mathbb{R}^d : \mathcal{M}_{\mathcal{H}_\infty^\beta} f(x) > t\} \right) dt \\ &\leq C_1 p \int_0^\infty t^{p-1} \mathcal{H}_\infty^\beta \left(\{x \in \mathbb{R}^d : \mathcal{M}_{\mathcal{H}_\infty^{\beta, Q_0}} f(x) > C_2 2^{-(\beta+d)} t\} \right) dt \\ &= C_3 \int_{\mathbb{R}^d} (\mathcal{M}_{\mathcal{H}_\infty^{\beta, Q_0}} f)^p d\mathcal{H}_\infty^{\beta, Q_0}. \end{aligned}$$

Moreover, it is easy to see that there exists $C > 0$ such that

$$\mathcal{M}_{\beta, Q_0}^\# f(x) \leq C \mathcal{M}_\beta^\# f(x).$$

Thus, the proof is complete once we show that there exists $C' = C'(p, d, \beta)$ such that

$$(3.15) \quad \int_{\mathbb{R}^d} (\mathcal{M}_\infty^{\beta, Q_0} f)^p d\mathcal{H}_\infty^{\beta, Q_0} \leq C' \int_{\mathbb{R}^d} (\mathcal{M}_{\beta, Q_0}^\# f)^p d\mathcal{H}_\infty^{\beta, Q_0}.$$

To this end, we apply Lemma 3.1 and obtain that for any $t > 0$ and $A > 0$,

$$(3.16) \quad \mu(t) \leq \mathcal{H}_\infty^{\beta, Q_0}(\{x \in \mathbb{R}^d : \mathcal{M}_{\beta, Q_0}^\# f(x) > \frac{t}{A}\}) + \frac{16}{A} \mu(2^{-\beta-2}t),$$

where

$$\mu(t) = \mathcal{H}_\infty^{\beta, Q_0} \left(\{x \in \mathbb{R}^d : \mathcal{M}_{\mathcal{H}_\infty^{\beta, Q_0}} f(x) > t\} \right).$$

In particular, for any $N > 0$, using (3.16) and change of variables, we obtain

$$\begin{aligned}
(3.17) \quad p \int_0^N t^{p-1} \mu(t) dt &\leq p \int_0^N t^{p-1} \mathcal{H}_\infty^{\beta, Q_0}(\{x \in \mathbb{R}^d : \mathcal{M}_{\beta, Q_0}^\# f(x) > \frac{t}{A}\}) dt \\
&\quad + \frac{8}{A} p \int_0^N t^{p-1} \mu(2^{-\beta-2}t) dt \\
&\leq A^p p \int_0^\infty t^{p-1} \mathcal{H}_\infty^{\beta, Q_0}(\{x \in \mathbb{R}^d : \mathcal{M}_{\beta, Q_0}^\# f(x) > t\}) dt \\
&\quad + 2^{(\beta+2)p} \frac{8}{A} p \int_0^N t^{p-1} \mu(t) dt \\
&= A^p \int_{\mathbb{R}^d} \left(\mathcal{M}_{\beta, Q_0}^\# f \right)^p d\mathcal{H}_\infty^{\beta, Q_0} + 2^{(\beta+2)p} \frac{8}{A} p \int_0^N t^{p-1} \mu(t) dt.
\end{aligned}$$

Moreover, we observe that

$$\int_0^N t^{p-1} \mu(t) dt \leq M_N \int_0^N t^{p-p_0-1} dt = M_N \frac{N^{p-p_0}}{p-p_0} < \infty,$$

where $M_N := \sup_{0 < t < N} t^{p_0} \mu(t) < \infty$ by (1.5). Therefore, we may subtract both sides of inequality (3.17) with $2^{(\beta+2)p} \frac{8}{A} p \int_0^N t^{p-1} \mu(t) dt$ and obtain

$$(3.18) \quad \left(1 - 2^{(\beta+2)p} \frac{8}{A} \right) p \int_0^N t^{p-1} \mu(t) dt \leq A^p \int_{\mathbb{R}^d} \left(\mathcal{M}_{\beta, Q_0}^\# f \right)^p d\mathcal{H}_\infty^{\beta, Q_0}.$$

By choosing $A = \frac{2^{(\beta+2)p}}{16}$ and letting $N \rightarrow \infty$ in inequality (3.18), we obtain

$$\begin{aligned}
\frac{1}{2} \int_{\mathbb{R}^d} \left(\mathcal{M}_{\mathcal{H}_\infty^{\beta, Q_0}} f \right)^p d\mathcal{H}_\infty^{\beta, Q_0} &= \lim_{N \rightarrow \infty} \left(1 - \frac{1}{2} \right) p \int_0^N t^{p-1} \mu(t) dt \\
&\leq \frac{2^{(\beta+2)p^2}}{16p} \int_{\mathbb{R}^d} \left(\mathcal{M}_{\beta, Q_0}^\# f \right)^p d\mathcal{H}_\infty^{\beta, Q_0}
\end{aligned}$$

by monotone converge theorem. This completes the proof of (3.15).

To prove (1.7), it is sufficient to show that there exist $C' = C'(p, d, \beta)$ such that

$$(3.19) \quad \left\| \mathcal{M}_\infty^{\beta, Q_0} f \right\|_{L^{p, \infty}(\mathcal{H}_\infty^\beta)} \leq C' \left\| \mathcal{M}_{\beta, Q_0}^\# f \right\|_{L^{p, \infty}(\mathcal{H}_\infty^\beta)}.$$

Let $N > 0$, using again (3.16), multiplying by t^p and taking the supremum over $0 < \lambda \leq N$ we obtain the following estimate,

$$\begin{aligned}
\sup_{0 < t \leq N} t^p \mu(t) &\leq \frac{16}{A} \sup_{0 < t \leq N} t^p \mu(2^{-\beta-2}t) + \sup_{0 < t \leq N} t^p \mathcal{H}_\infty^{\beta, Q_0}(\{x \in \mathbb{R}^d : \mathcal{M}_{\beta, Q_0}^\# f(x) > \frac{t}{A}\}) \\
&= \frac{2^{(\beta+2)p+4}}{A} \sup_{0 < t \leq 2^{-\beta-2}N} t^p \mu(t) + A^p \sup_{0 < t \leq \frac{N}{A}} t^p \mathcal{H}_\infty^{\beta, Q_0}(\{x \in \mathbb{R}^d : \mathcal{M}_{\beta, Q_0}^\# f(x) > t\}) \\
&\leq \frac{2^{(\beta+2)p+4}}{A} \sup_{0 < t \leq N} t^p \mu(t) + A^p \left\| \mathcal{M}_{\beta, Q_0}^\# f \right\|_{L^{p, \infty}(\mathcal{H}_\infty^\beta)},
\end{aligned}$$

for all $A > 0$. Observe that the first term is finite,

$$\sup_{0 < t \leq N} t^p \mu(t) \leq M_N N^{p-p_0} < \infty,$$

where $M_N := \sup_{0 < t < N} t^{p_0} \mu(t) < \infty$ by (1.5). We take $A > 0$ such that $\frac{2^{(\beta+2)p+4}}{A} = \frac{1}{2}$, that is $A = 2^{(\beta+2)p+5}$. Then, since the right-hand side is finite, we can absorb it into the left-hand side, and we obtain,

$$\sup_{0 < t \leq N} t^p \mu(t) \leq 2^{((\beta+2)p+5)p+1} \left\| \mathcal{M}_{\beta, Q_0}^{\#} f \right\|_{L^{p, \infty}(\mathcal{H}_{\infty}^{\beta})}.$$

We conclude the proof of (3.19) letting $N \rightarrow \infty$. \square

4. POINTWISE ESTIMATE OF FRACTIONAL MAXIMAL FUNCTION AND SHARP MAXIMAL FUNCTION OF $I_{\alpha} f$

In this section, we will first give the proof of Theorem 1.4 and then provide an analogous of Theorem 1.5 based on Adams's approach. We begin with a result which can be found in [9, Lemma 2.2] and [16, Proposition 2.3] and is an extension of [22, Lemma 3].

Lemma 4.1. *Let $f \geq 0$ and $0 < \alpha \leq \beta \leq d \in \mathbb{N}$. Then*

$$\int_{\mathbb{R}^d} f d\mathcal{H}_{\infty}^{\beta} \leq \frac{\beta}{\alpha} \left(\int_{\mathbb{R}^d} f^{\frac{\alpha}{\beta}} d\mathcal{H}_{\infty}^{\alpha} \right)^{\frac{\beta}{\alpha}}.$$

With Lemma 4.1, one can immediately establish

Lemma 4.2. *Let $0 < \alpha \leq \beta \leq d \in \mathbb{N}$. Then there exists a constant $C > 0$ depending only on α and β such that*

$$\mathcal{M}_{\beta}^{\#} f(x) \leq C \mathcal{M}_{\alpha}^{\#} f(x)$$

for all $x \in \mathbb{R}^d$.

Proof. Let $0 < \alpha \leq \beta \leq d$. Fix $x \in \mathbb{R}^d$ and Q be a cube containing x . An application of Lemma 4.1 and Hölder's inequality in Lemma 2.2 (iii) yields that

$$\begin{aligned} \frac{1}{\mathcal{H}_{\infty}^{\beta}(Q)} \int_Q |f - c_Q| d\mathcal{H}_{\infty}^{\beta} &\leq \frac{\beta}{\alpha} \frac{1}{\mathcal{H}_{\infty}^{\beta}(Q)} \left(\int_Q |f - c_Q|^{\frac{\alpha}{\beta}} d\mathcal{H}_{\infty}^{\alpha} \right)^{\frac{\beta}{\alpha}} \\ &\leq \frac{2\beta \mathcal{H}_{\infty}^{\alpha}(Q)^{\frac{\beta}{\alpha}}}{\alpha \mathcal{H}_{\infty}^{\beta}(Q)} \left(\frac{1}{\mathcal{H}_{\infty}^{\alpha}(Q)} \int_Q |f - c_Q| d\mathcal{H}_{\infty}^{\alpha} \right) \\ &\leq C \mathcal{M}_{\alpha}^{\#} f(x), \end{aligned}$$

where

$$c_Q = \arg \min_{c \in \mathbb{R}} \frac{1}{\ell(Q)^{\alpha}} \int_Q |f - c| d\mathcal{H}_{\infty}^{\alpha}$$

and

$$C = \frac{2\beta \mathcal{H}_{\infty}^{\alpha}(Q)^{\frac{\beta}{\alpha}}}{\alpha \mathcal{H}_{\infty}^{\beta}(Q)}.$$

Taking the supremum over all cubes Q such that $x \in Q$, we obtain

$$\mathcal{M}_{\beta}^{\#} f(x) \leq C \mathcal{M}_{\alpha}^{\#} f(x).$$

\square

We next record two results from [10, Lemma 3.5 and Lemma 3.8].

Lemma 4.3. *Let $0 < \alpha < d \in \mathbb{N}$, $0 < \varepsilon \leq \alpha$, Q be an open cube in \mathbb{R}^d with centre x_0 and f be a real-valued function in \mathbb{R}^d . If $\text{supp } f \subseteq 2Q$, then $I_\alpha f \in L^1(Q; \mathcal{H}_\infty^{d-\alpha+\varepsilon})$ and there exists a constant C depending only on d, α and ε such that*

$$(4.1) \quad \int_Q |I_\alpha f| d\mathcal{H}_\infty^{d-\alpha+\varepsilon} \leq C\ell(Q)^{d-\alpha+\varepsilon} \mathcal{M}_\alpha f(x_0).$$

Lemma 4.4. *Let $0 < \alpha < d \in \mathbb{N}$, $\beta \in (0, d]$, Q be an open cube in \mathbb{R}^d with centre x_0 and f be a real-valued function with $\text{supp } f \subseteq (2Q)^c$. If $I_\alpha f \in L^1_{loc}(\mathbb{R}^d)$, then $I_\alpha f$ is continuous in Q and there exists $c \in \mathbb{R}$ such that*

$$(4.2) \quad \int_Q |I_\alpha f - c| d\mathcal{H}_\infty^\beta \leq C\ell(Q)^\beta \mathcal{M}_\alpha f(x_0).$$

The combination of Lemma 4.3 and 4.4 readily yields the following pointwise estimate concerning β -dimensional sharp maximal function and fractional maximal function.

Lemma 4.5. *Let $0 < \alpha < d \in \mathbb{N}$, $0 < \varepsilon \leq \alpha$. If $I_\alpha f \in L^1_{loc}(\mathbb{R}^d)$, then $I_\alpha f \in L^1(Q; \mathcal{H}_\infty^{d-\alpha+\varepsilon})$ and for every $x \in \mathbb{R}^d$, there exists a constant C depending only on d, α, ε such that*

$$(4.3) \quad \mathcal{M}_\beta^\#(I_\alpha f)(x) \leq C\mathcal{M}_\alpha f(x).$$

For the proof of Lemma 4.5, we introduce β -dimensional centered sharp maximal operator $\mathcal{M}_\beta^{\#,c} f(x)$ of a function f which is defined as

$$\mathcal{M}_\beta^{\#,c} f(x) := \sup_Q \inf_{c \in \mathbb{R}} \frac{1}{\ell(Q)^\beta} \int_Q |f - c| d\mathcal{H}_\infty^\beta,$$

where the supremum is taken over all cubes Q with center x . It is easy to see that there exists a constant $C_\beta > 0$ such that

$$(4.4) \quad \mathcal{M}_\beta^{\#,c} f(x) \leq \mathcal{M}_\beta^\# f(x) \leq C_\beta \mathcal{M}_\beta^{\#,c} f(x).$$

Proof. Suppose that $I_\alpha f \in L^1_{loc}(\mathbb{R}^d)$. Let Q be a cube in \mathbb{R}^d with centre x_0 and $f_1 = f\chi_{2Q}$, $f_2 = f - f_1$. Then there exists a constant $c \in \mathbb{R}$ and C depending on d, α, ε such that

$$\begin{aligned} \int_Q |I_\alpha f - c| d\mathcal{H}_\infty^{d-\alpha+\varepsilon} &= \int_Q |I_\alpha f_1 + I_\alpha f_2 - c| d\mathcal{H}_\infty^{d-\alpha+\varepsilon} \\ &\leq 2 \left(\int_Q |I_\alpha f_1| d\mathcal{H}_\infty^{d-\alpha+\varepsilon} + \int_Q |I_\alpha f_2 - c| d\mathcal{H}_\infty^{d-\alpha+\varepsilon} \right) \\ &\leq 2C\ell(Q)^{d-\alpha+\varepsilon} \mathcal{M}_\alpha f(x_0), \end{aligned}$$

where we use Lemma 4.3 and Lemma 4.4 in the last inequality. Now dividing both sides with $\ell(Q)^{d-\alpha+\varepsilon}$ and taking the supremum over all cubes Q with centre x , we obtain

$$\mathcal{M}_\beta^{\#,c}(I_\alpha f)(x) \leq C\mathcal{M}_\alpha f(x)$$

and thus

$$\mathcal{M}_\beta^\#(I_\alpha f)(x) \leq C\mathcal{M}_\alpha f(x)$$

by (4.4). □

We now give the proof of Theorem 1.4

Proof. Let $x \in \mathbb{R}^d$. By Lemma 4.5, we have

$$\mathcal{M}_\beta^\#(I_\alpha f)(x) \lesssim \mathcal{M}_\alpha f(x).$$

The combination of (1.9) and Lemma 4.2 yields that

$$\mathcal{M}_\alpha f(x) \cong \mathcal{M}^\#(I_\alpha f)(x) \lesssim \mathcal{M}_\beta^\#(I_\alpha f)(x),$$

which completes the proof. □

Now, we use the previous result to provide an alternative proof of Theorem 1.5 with an additional assumption, following the approach in [2].

Theorem 4.6. *Let $0 < \alpha < d \in \mathbb{N}$, $0 < p < \infty$ and $\beta \in (d - \alpha, d]$. If f is a function in \mathbb{R}^d such that there exists $0 < p_0 < p$ such that*

$$(4.5) \quad \sup_{0 < t \leq N} t^{p_0} \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : \mathcal{M}_{\mathcal{H}_\infty^\beta} I_\alpha f(x) > t\}) < \infty \quad \text{for all } N > 0,$$

then there exists a constant $C = C(d, \alpha, \beta, p)$ such that

$$\|I_\alpha f\|_{L^p(\mathcal{H}_\infty^\beta)} \leq C \|\mathcal{M}_\alpha f\|_{L^p(\mathcal{H}_\infty^\beta)}.$$

Proof. Without loss of generality, we may assume $f \geq 0$ and

$$(4.6) \quad \|\mathcal{M}_\alpha f\|_{L^p(\mathcal{H}_\infty^\beta)} < \infty.$$

This yields that $I_\alpha f \in L_{loc}^1(\mathbb{R}^d)$. To see this, we observe that $\mathcal{M}_\alpha f \in L^p(\mathcal{H}_\infty^\beta)$ implies that $\mathcal{M}_\alpha f \in L^r(\mathbb{R}^d)$ for $r = \frac{pd}{\beta}$ by Lemma 4.1 and thus $I_\alpha f \in L^r(\mathbb{R}^d)$ by (1.8). It follows that $I_\alpha f \in L_{loc}^1(\mathbb{R}^d)$. Now we apply Theorem 1.2 and Theorem 1.4 to $I_\alpha f$ and deduce that there exists a constant $C = C(d, \alpha, \beta, p)$ such that

$$\begin{aligned} \|I_\alpha f\|_{L^p(\mathcal{H}_\infty^\beta)} &\leq \|\mathcal{M}_{\mathcal{H}_\infty^\beta}(I_\alpha f)\|_{L^p(\mathcal{H}_\infty^\beta)} \\ &\leq C \|\mathcal{M}_\beta^\#(I_\alpha f)\|_{L^p(\mathcal{H}_\infty^\beta)} \\ &\leq C \|\mathcal{M}_\alpha f\|_{L^p(\mathcal{H}_\infty^\beta)}, \end{aligned}$$

which completes the proof. □

Finally, we give the proof of Corollary 1.7.

Proof. Since $f \in L^r(\mathbb{R}^d)$, we have $I_\alpha f \in L_{loc}^1(\mathbb{R}^d)$. It then follows immediately from Theorem 1.4 that

$$\|\mathcal{M}_\beta^\#(I_\alpha f)\|_{L^p(\mathcal{H}_\infty^\beta)} \cong \|\mathcal{M}_\alpha f\|_{L^p(\mathcal{H}_\infty^\beta)}.$$

Moreover, since

$$\mathcal{M}_\alpha f(x) \leq C_{d,\alpha} I_\alpha f(x)$$

we have

$$(4.7) \quad \|\mathcal{M}_\alpha f\|_{L^p(\mathcal{H}_\infty^\beta)} \leq C_{d,\alpha} \|I_\alpha f\|_{L^p(\mathcal{H}_\infty^\beta)}.$$

The combination of Theorem 1.5 and inequality (4.7) gives

$$\|\mathcal{M}_\alpha f\|_{L^p(\mathcal{H}_\infty^\beta)} \cong \|I_\alpha f\|_{L^p(\mathcal{H}_\infty^\beta)},$$

which completes the proof. □

5. GOOD LAMBDA ESTIMATE FOR RIESZ POTENTIAL WITH RESPECT TO HAUSDORFF CONTENT AND APPLICATIONS

In this section, we present the proof of Theorem 1.5. To this end, we first prove a good lambda inequality with exponential decay, following the ideas of [17] and using the packing condition to avoid difficulties due to the nonlinearity of the Hausdorff content. We conclude the section with some consequences of the exponential decay on the good-lambda estimate.

First, we define a dyadic version of the Riesz potential, which we will use to approximate I_α . We begin with the definition of a dyadic grid. A dyadic grid \mathcal{D} is a countable family of cubes with the following properties

- (1) For each $Q \in \mathcal{D}$, $\ell(Q) = 2^k$ for some $k \in \mathbb{Z}$.
- (2) Given $Q, P \in \mathcal{D}$, then $Q \cap P \in \{\emptyset, P, Q\}$.
- (3) For each $k \in \mathbb{Z}$, the set $\mathcal{D}_k = \{Q \in \mathcal{D} : \ell(Q) = 2^k\}$ is a partition of \mathbb{R}^d .

The following dyadic operator will play a central role in the proof of the good lambda estimate.

Definition 5.1. *Let $0 < \alpha < d \in \mathbb{N}$ and let \mathcal{D} a dyadic grid. We define the dyadic Riesz potential with respect to \mathcal{D} of a locally finite measure μ , by*

$$I_\alpha^{\mathcal{D}} \mu(x) = \sum_{Q \in \mathcal{D}} \frac{\mu(Q)}{\ell(Q)^{d-\alpha}} \chi_Q(x).$$

Dyadic Riesz potentials were first introduced by E. Sawyer and R. L. Wheeden [26]. We will use the following result by D. Cruz-Uribe and K. Moen (see [12, Proposition 2.2]) to approximate the Riesz potential by a linear combination of dyadic Riesz potential.

Proposition 5.2. *Given $0 < \alpha < d \in \mathbb{N}$, and a locally finite measure μ , then for any dyadic grid \mathcal{D} ,*

$$(5.1) \quad I_\alpha^{\mathcal{D}} \mu(x) \leq c_{d,\alpha} I_\alpha \mu(x).$$

Conversely, we have that

$$I_\alpha \mu(x) \leq c_{d,\alpha} \max_{i \in \{0, \frac{1}{3}\}^d} I_\alpha^{\mathcal{D}^i} \mu(x),$$

where

$$\mathcal{D}^i = \{2^{-k}([0, 1)^d + m + (-1)^k i) : k \in \mathbb{Z}, m \in \mathbb{Z}^d\}, \quad i \in \{0, \frac{1}{3}\}^d.$$

Hence, the Riesz potential is pointwise equivalent to a linear combination of dyadic Riesz potentials, that is,

$$(5.2) \quad I_\alpha \mu(x) \cong_{d,\alpha} \sum_{i \in \{0, \frac{1}{2}\}^d} I_\alpha^{\mathcal{D}^i} \mu(x).$$

We should mention that the proof of the previous result in [12] is made for functions, but a detailed inspection of the proof shows that the same proof works for measures.

Now, we state the main result of this section. The result is an extension of the good lambda inequality proved in [17], which improves the classical result of B. Muckenhoupt and R. L. Wheeden [21].

Theorem 5.3. *Let $0 < \alpha < d$, $\beta \in (d - \alpha, d]$ and let \mathcal{D} be a dyadic grid. Then, there exist constants $C, c > 0$ depending on d, α, β such that*

$$(5.3) \quad \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : I_\alpha^{\mathcal{D}} \mu(x) > 2\lambda, \mathcal{M}_\alpha \mu(x) \leq \varepsilon \lambda\}) \leq C e^{-\frac{c}{\varepsilon}} \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : I_\alpha^{\mathcal{D}} \mu(x) > \lambda\})$$

for all $\lambda > 0$ and all $0 < \varepsilon < 1$ and all locally finite measure μ .

Proof. It is sufficient to prove (5.3) for the dyadic content $\mathcal{H}_\infty^{\beta, Q_0}$, where Q_0 is a cube in \mathcal{D} , since both contents are equivalent by (2.4).

Fix $\lambda > 0$ and $0 < \varepsilon < 1$. We denote $G_\lambda = \{x \in \mathbb{R}^d : I_\alpha^{\mathcal{D}} \mu(x) > \lambda\}$. We can assume without loss of generality $\mathcal{H}_\infty^{\beta, Q_0}(G_\lambda) < \infty$. Let $\{Q_j^\lambda\}_j$ the maximal collection of dyadic cubes subordinates to \mathcal{D} contained in G_λ with respect to the inclusion.

Applying Lemma 2.7 to the family $\{Q_j^\lambda\}_j$ we obtain a subfamily $\{Q_{j_k}^\lambda\}_k$ and a family of non-overlapping ancestors $\{\tilde{Q}_k\}_k$ such that

(1)

$$G_\lambda = \bigcup_k Q_{j_k}^\lambda \subset \bigcup_k Q_j^\lambda \cup \bigcup_k \tilde{Q}_k$$

(2)

$$\sum_{Q_{j_k}^\lambda \subset Q} \ell(Q_{j_k}^\lambda)^\beta \leq 2\ell(Q)^\beta, \text{ for each dyadic cube } Q.$$

(3)

$$\begin{aligned} \mathcal{H}_\infty^{\beta, Q_0}(\bigcup_j Q_j^\lambda) &\leq \sum_{k, Q_{j_k}^\lambda \not\subseteq \tilde{Q}_m} \ell(Q_{j_k}^\lambda)^\beta + \sum_k \ell(\tilde{Q}_k)^\beta \\ &\leq 2 \sum_k \ell(Q_{j_k}^\lambda)^\beta \\ &\leq 2\mathcal{H}_\infty^{\beta, Q_0}(\bigcup_j Q_j^\lambda). \end{aligned}$$

Define $S = \{x \in \mathbb{R}^d : I_\alpha^{\mathcal{D}} \mu(x) > 2\lambda, \mathcal{M}_\alpha \mu(x) \leq \varepsilon \lambda\}$. Since $G_{2\lambda} \subseteq G_\lambda$, we observe that

$$S \subseteq G_{2\lambda} \subseteq G_\lambda = \bigcup_k Q_{j_k}^\lambda \cup \bigcup_k \tilde{Q}_k,$$

and then we have

$$S \subseteq \bigcup_{k: Q_{j_k}^\lambda \not\subseteq \tilde{Q}_m} \{x \in Q_{j_k}^\lambda : I_\alpha^{\mathcal{D}} \mu(x) > 2\lambda, \mathcal{M}_\alpha \mu(x) \leq \varepsilon\lambda\} \\ \cup \bigcup_k \{x \in \tilde{Q}_k : I_\alpha^{\mathcal{D}} \mu(x) > 2\lambda, \mathcal{M}_\alpha \mu(x) \leq \varepsilon\lambda\}.$$

Using the subadditivity of $\mathcal{H}_\infty^{\beta, Q_0}$ we obtain

$$(5.4) \quad \mathcal{H}_\infty^{\beta, Q_0}(S) \leq \sum_{k: Q_{j_k}^\lambda \not\subseteq \tilde{Q}_m} \mathcal{H}_\infty^{\beta, Q_0}(\{x \in Q_{j_k}^\lambda : I_\alpha^{\mathcal{D}} \mu(x) > 2\lambda, \mathcal{M}_\alpha \mu(x) \leq \varepsilon\lambda\}) \\ + \sum_k \mathcal{H}_\infty^{\beta, Q_0}(\{x \in \tilde{Q}_k : I_\alpha^{\mathcal{D}} \mu(x) > 2\lambda, \mathcal{M}_\alpha \mu(x) \leq \varepsilon\lambda\}).$$

Fix $Q' \in \{Q_{j_k}^\lambda\}_k \cup \{\tilde{Q}_k\}_k$, and let us define $S_{Q'} = \{x \in Q' : I_\alpha^{\mathcal{D}} \mu(x) > 2\lambda, \mathcal{M}_\alpha \mu(x) \leq \varepsilon\lambda\}$. We can assume that $S_{Q'}$ is non-empty.

We have the following estimate on the tail of the potential

$$\sum_{\substack{Q \in \mathcal{D} \\ Q \supseteq Q'}} \frac{\mu(Q)}{\ell(Q)^{d-\alpha}} \leq \lambda,$$

by the definition of G_λ and observing that either Q' is a maximal cube of the family $\{Q_j^\lambda\}_j$, or it contains a maximal cube in $\{Q_j^\lambda\}_j$. Let $j_0 \in \mathbb{Z}$ such that $\ell(Q') = 2^{j_0}$ and $m \in \mathbb{N}$ to be chosen later. If $x \in S_{Q'}$, we have

$$I_\alpha^{\mathcal{D}} \mu(x) = \sum_{x \in Q \in \mathcal{D}} \frac{\mu(Q)}{\ell(Q)^{d-\alpha}} \\ = \sum_{\substack{x \in Q \in \mathcal{D} \\ Q \supseteq Q'}} \frac{\mu(Q)}{\ell(Q)^{d-\alpha}} + \sum_{\substack{x \in Q \in \mathcal{D} \\ 2^{j_0-m} < \ell(Q) \leq 2^{j_0}}} \frac{\mu(Q)}{\ell(Q)^{d-\alpha}} + \sum_{\substack{x \in Q \in \mathcal{D} \\ \ell(Q) \leq 2^{j_0-m}}} \frac{\mu(Q)}{\ell(Q)^{d-\alpha}} \\ \leq \lambda + m\mathcal{M}_\alpha \mu(x) + \sum_{\substack{x \in Q \in \mathcal{D} \\ \ell(Q) \leq 2^{j_0-m}}} \frac{\mu(Q)}{\ell(Q)^{d-\alpha}} \\ \leq \lambda + m\varepsilon\lambda + \sum_{k \leq j_0-m} g_k(x),$$

where $g_k(x)$ is defined as

$$g_k(x) = \sum_{x \in Q \in \mathcal{D}_k} \frac{\mu(Q)}{\ell(Q)^{d-\alpha}} \quad \text{for } k \in \mathbb{Z}$$

and $\mathcal{D}_k = \{Q \in \mathcal{D} : \ell(Q) = 2^k\}$. Observe that only one nonzero term exists for each x in the previous sum.

Let $\eta = \frac{\beta-(d-\alpha)}{2}$, we remark that since $\beta \in (d-\alpha, d]$ we have $\eta > 0$. We claim

$$(5.5) \quad S_{Q'} \subseteq \bigcup_{k \leq j_0-m} \{x \in Q' : g_k(x) > \lambda(1-m\varepsilon)2^{\eta(k-j_0+m)}(1-2^{-\eta})\}.$$

To this end, observe that if $x \in S_{Q'}$, we have

$$2\lambda < I_\alpha^D \mu(x) \leq \lambda + m\varepsilon\lambda + \sum_{k \leq j_0 - m} g_k(x)$$

and thus,

$$(5.6) \quad \lambda(1 - m\varepsilon) < \sum_{k \leq j_0 - m} g_k(x).$$

If we assume that

$$g_k(x) \leq \lambda(1 - m\varepsilon)2^{\eta(k-j_0+m)}(1 - 2^{-\eta})$$

for all $k \leq j_0 - m$, then

$$\sum_{k \leq j_0 - m} g_k(x) \leq \lambda(1 - m\varepsilon)(1 - 2^{-\eta}) \sum_{k \leq j_0 - m} 2^{\eta(k-j_0+m)} = \lambda(1 - m\varepsilon),$$

which contradicts to (5.6). Thus, there exists $k \leq j_0 - m$ such that

$$g_k(x) > \lambda(1 - m\varepsilon)2^{\eta(k-j_0+m)}(1 - 2^{-\eta}),$$

and we conclude the proof of the claim.

Observe that for any $\gamma > 0$ and any $k \leq j_0$, we have

$$\begin{aligned} \mathcal{H}_\infty^{\beta, Q_0}(\{x \in Q' : g_k(x) > \gamma\}) &\leq \frac{1}{\gamma} \int_{Q'} g_k d\mathcal{H}_\infty^{\beta, Q_0} \\ &= \frac{1}{\gamma} \sum_{\substack{x \in Q \in \mathcal{D}_k \\ Q \subseteq Q'}} \frac{\mu(Q)}{\ell(Q)^{d-\alpha}} \ell(Q)^\beta \\ &= \frac{1}{\gamma} \sum_{\substack{x \in Q \in \mathcal{D}_k \\ Q \subseteq Q'}} \ell(Q)^{\beta-(d-\alpha)} \mu(Q) \\ &\leq \frac{1}{\gamma} 2^{2\eta k} \mu(Q') \\ &= \frac{1}{\gamma} 2^{2\eta k} \frac{1}{\ell(Q')^{2\eta}} \ell(Q')^\beta \frac{\mu(Q')}{\ell(Q')^{d-\alpha}} \\ &\leq \frac{1}{\gamma} 2^{2\eta(k-j_0)} \ell(Q')^\beta \inf_{z \in Q'} \mathcal{M}_\alpha \mu(z) \\ &\leq \frac{1}{\gamma} 2^{2\eta(k-j_0)} \ell(Q')^\beta \mathcal{M}_\alpha \mu(x_{Q'}) \\ (5.7) \quad &\leq \frac{1}{\gamma} 2^{2\eta(k-j_0)} \ell(Q')^\beta \varepsilon \lambda, \end{aligned}$$

where $x_{Q'} \in Q'$ such that $\mathcal{M}_\alpha \mu(x_{Q'}) \leq \varepsilon \lambda$, that we know exists because otherwise, $S_{Q'}$ would be empty.

The combination of (5.5) and (5.7) then yields that

$$\begin{aligned}
\mathcal{H}_\infty^{\beta, Q_0}(S_{Q'}) &\leq \sum_{k \leq j_0 - m} \mathcal{H}_\infty^{\beta, Q_0}(\{x \in Q' : g_k(x) > \lambda(1 - m\varepsilon)2^{\eta(k - j_0 + m)}(1 - 2^{-\eta})\}) \\
&\leq \frac{1}{\lambda(1 - m\varepsilon)(1 - 2^{-\eta})} \ell(Q')^\beta \varepsilon \lambda \sum_{k \leq j_0 - m} \frac{2^{2\eta(k - j_0)}}{2^{\eta(k - j_0 + m)}} \\
&= \ell(Q')^\beta \frac{\varepsilon}{1 - m\varepsilon} \frac{2^{-2\eta m}}{1 - 2^{-\eta}} \sum_{k \leq j_0 - m} 2^{-\eta(j_0 - m - k)} \\
&= \ell(Q')^\beta \frac{\varepsilon}{1 - m\varepsilon} \frac{2^{-2\eta m}}{1 - 2^{-\eta}} \sum_{k=0}^{\infty} 2^{-\eta k} \\
&= C_\eta 2^{-2\eta m} \frac{\varepsilon}{1 - m\varepsilon} \ell(Q')^\beta.
\end{aligned}$$

We remark that the previous sum is finite because $\eta > 0$. Now we choose $m \in \mathbb{N} \cup \{0\}$ such that

$$\frac{1}{\varepsilon} - 2 \leq m < \frac{1}{\varepsilon} - 1,$$

then $\frac{\varepsilon}{1 - m\varepsilon} < 1$, and hence, there exists $c = c(d, \alpha, \beta)$ such that

$$\mathcal{H}_\infty^{\beta, Q_0}(S_{Q'}) \leq C_\eta 2^{-\frac{2\eta}{\varepsilon}} \ell(Q')^\beta = C_\eta e^{-\frac{c}{\varepsilon}} \ell(Q')^\beta.$$

Using the previous estimate and the packing condition of the family $\{Q_{j_k}^\lambda\}_k$, we have

$$\begin{aligned}
\mathcal{H}_\infty^{\beta, Q_0}(S) &\leq \sum_{k: Q_{j_k}^\lambda \not\subseteq \tilde{Q}_m} \mathcal{H}_\infty^{\beta, Q_0}(S_{Q_{j_k}^\lambda}) + \sum_k \mathcal{H}_\infty^{\beta, Q_0}(S_{\tilde{Q}_k}) \\
&\leq C_\eta e^{-\frac{c}{\varepsilon}} \left(\sum_{k: Q_{j_k}^\lambda \not\subseteq \tilde{Q}_m} \ell(Q_{j_k}^\lambda)^\beta + \sum_k \ell(\tilde{Q}_k)^\beta \right) \\
&\leq 2C_\eta e^{-\frac{c}{\varepsilon}} \mathcal{H}_\infty^{\beta, Q_0} \left(\bigcup_j Q_j^\lambda \right) \\
&= 2C_\eta e^{-\frac{c}{\varepsilon}} \mathcal{H}_\infty^{\beta, Q_0}(G_\lambda).
\end{aligned}$$

□

As a consequence of the previous good-lambda estimate, we give the proof of Theorem 1.5, which is an improved version of Theorem 4.6.

Proof of Theorem 1.5. We observe that it is enough to prove (1.10) and (1.11) for the dyadic Riesz potential $I_\alpha^\mathcal{D}$ with a constant independent of the dyadic grid \mathcal{D} . To show

this, we use (5.2), and we obtain the desired inequality,

$$\begin{aligned} \|I_\alpha \mu\|_{L^p(\mathcal{H}_\infty^\beta)} &\leq c_{d,\alpha} \left\| \sum_{i \in \{0, \frac{1}{3}\}^d} I_\alpha^{\mathcal{D}^i} \mu \right\|_{L^p(\mathcal{H}_\infty^\beta)} \\ &\leq c_\beta c_{d,\alpha} \sum_{i \in \{0, \frac{1}{3}\}^d} \|I_\alpha^{\mathcal{D}^i} \mu\|_{L^p(\mathcal{H}_\infty^\beta)} \\ &\leq Cp c_\beta c_{d,\alpha} \sum_{i \in \{0, \frac{1}{3}\}^d} \|\mathcal{M}_\alpha \mu\|_{L^p(\mathcal{H}_\infty^\beta)} \\ &= C_{\beta,\alpha,d} p \|\mathcal{M}_\alpha \mu\|_{L^p(\mathcal{H}_\infty^\beta)}. \end{aligned}$$

Similarly, we can obtain (1.11) from the dyadic estimate.

Let us first assume that μ has compact support. We may also assume that $\|\mathcal{M}_\alpha \mu\|_{L^p(\mathcal{H}_\infty^\beta)}$ is finite because otherwise, there is nothing to prove. Let \mathcal{D} a dyadic grid, using Theorem 5.3, and the subadditivity of \mathcal{H}_∞^β we obtain,

$$\begin{aligned} \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : I_\alpha^{\mathcal{D}} \mu(x) > 2\lambda\}) &\leq \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : I_\alpha^{\mathcal{D}} \mu(x) > 2\lambda, \mathcal{M}_\alpha \mu(x) \leq \varepsilon\lambda\}) \\ &\quad + \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : \mathcal{M}_\alpha \mu(x) > \varepsilon\lambda\}) \\ &\leq C e^{-\frac{\varepsilon}{2}} \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : I_\alpha^{\mathcal{D}} \mu(x) > \lambda\}) \\ &\quad + C \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : \mathcal{M}_\alpha \mu(x) > \varepsilon\lambda\}), \end{aligned}$$

for all $\lambda > 0$ and all $0 < \varepsilon < 1$. Let $N > 0$, we have

$$\begin{aligned} \int_0^N \lambda^{p-1} \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : I_\alpha^{\mathcal{D}} \mu(x) > \lambda\}) d\lambda &= 2^p \int_0^{\frac{N}{2}} \lambda^{p-1} \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : I_\alpha^{\mathcal{D}} \mu(x) > 2\lambda\}) d\lambda \\ &\leq 2^p C e^{-\frac{\varepsilon}{2}} \int_0^{\frac{N}{2}} \lambda^{p-1} \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : I_\alpha^{\mathcal{D}} \mu(x) > \lambda\}) d\lambda \\ &\quad + 2^p C \int_0^{\frac{N}{2}} \lambda^{p-1} \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : \mathcal{M}_\alpha \mu(x) > \varepsilon\lambda\}) d\lambda \\ &\leq 2^p C e^{-\frac{\varepsilon}{2}} \int_0^N \lambda^{p-1} \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : I_\alpha^{\mathcal{D}} \mu(x) > \lambda\}) d\lambda \\ &\quad + \frac{2^p C}{\varepsilon^p} \int_0^{\frac{N\varepsilon}{2}} \lambda^{p-1} \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : \mathcal{M}_\alpha \mu(x) > \lambda\}) d\lambda \end{aligned}$$

for all $0 < \varepsilon < 1$. Notice that the first term is finite because μ has compact support. Let B be a ball such that $\text{supp } \mu \subseteq B$. If $x \notin 2B$, we have $r(B) > |x - y|$ for all $y \in B$. Hence,

$$I_\alpha \mu(x) = \frac{1}{\gamma(\alpha)} \int_B \frac{d\mu(y)}{|x - y|^{d-\alpha}} \leq \frac{1}{\gamma(\alpha)} \frac{\mu(B)}{r(B)^{d-\alpha}} \leq \frac{1}{\gamma(\alpha)} \mathcal{M}_\alpha \mu(x),$$

for all $x \notin 2B$. Using the previous estimate, we obtain

$$\begin{aligned} \{x \in \mathbb{R}^d : I_\alpha \mu(x) > \lambda\} &= \{x \in 2B : I_\alpha \mu(x) > \lambda\} \cup \{x \in (2B)^c : I_\alpha \mu(x) > \lambda\} \\ &\subseteq \{x \in 2B : I_\alpha \mu(x) > \lambda\} \cup \{x \in (2B)^c : \mathcal{M}_\alpha \mu(x) > \gamma(\alpha)\lambda\}, \end{aligned}$$

for all $\lambda > 0$. Then, we have

$$\begin{aligned} \int_0^N \lambda^{p-1} \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : I_\alpha \mu(x) > \lambda\}) d\lambda &\leq \int_0^N \lambda^{p-1} \mathcal{H}_\infty^\beta(\{x \in 2B : I_\alpha \mu(x) > \lambda\}) d\lambda \\ &\quad + \int_0^N \lambda^{p-1} \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : \mathcal{M}_\alpha \mu(x) > \gamma(\alpha)\lambda\}) d\lambda \\ &\leq r(2B)^\beta \frac{N^p}{p} + \frac{1}{\gamma(\alpha)^p} \|\mathcal{M}_\alpha \mu\|_{L^p(\mathcal{H}_\infty^\beta)}^p \\ &< \infty, \end{aligned}$$

and using (5.1) we obtain that the dyadic counterpart is also finite.

We take ε such that $2^p C e^{-\frac{\varepsilon}{c}} = \frac{1}{2}$, that is $\varepsilon = \frac{c}{\log(2^{p+1}C)}$. Then, since the right-hand side is finite, we can absorb it into the left-hand side, and we obtain

$$\int_0^N \lambda^{p-1} \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : I_\alpha^D \mu(x) > \lambda\}) d\lambda \leq 2^{p+1} C \frac{\log(2^{p+1}C)^p}{c^p p} \int_{\mathbb{R}^d} |\mathcal{M}_\alpha \mu|^p d\mathcal{H}_\infty^\beta.$$

We obtain the desired inequality letting $N \rightarrow \infty$,

$$\begin{aligned} \left(\int_{\mathbb{R}^d} |I_\alpha^D \mu|^p d\mathcal{H}_\infty^\beta \right)^{\frac{1}{p}} &\leq c_d \frac{\log(2^{p+1}C)}{c} \left(\int_{\mathbb{R}^d} |\mathcal{M}_\alpha \mu|^p d\mathcal{H}_\infty^\beta \right)^{\frac{1}{p}} \\ &\leq \tilde{C} p \left(\int_{\mathbb{R}^d} |\mathcal{M}_\alpha \mu|^p d\mathcal{H}_\infty^\beta \right)^{\frac{1}{p}} \end{aligned}$$

where $\tilde{C} = \tilde{C}(d, \beta, \alpha) > 0$.

Let us consider now the general case. Given a measure μ , we define $\mu_R = \mu \chi_{B(0,R)}$. Since μ_R has compact support, we have

$$\|I_\alpha^D \mu_R\|_{L^p(\mathcal{H}_\infty^\beta)} \leq C p \|\mathcal{M}_\alpha \mu_R\|_{L^p(\mathcal{H}_\infty^\beta)} \leq C p \|\mathcal{M}_\alpha \mu\|_{L^p(\mathcal{H}_\infty^\beta)}.$$

Since \mathcal{H}_∞^β is continuous from below, we may apply the monotone convergence theorem (see [23] for instance), and we obtain the desired inequality.

In order to prove (1.11), let us first assume that μ has compact support. We may assume that $\|\mathcal{M}_\alpha \mu\|_{L^{p,\infty}(\mathcal{H}_\infty^\beta)} < \infty$, because otherwise there is nothing to prove. Let $N > 0$, using again Theorem 5.3, multiplying by λ^p and taking the supremum over $0 < \lambda \leq N$ we obtain the following estimate,

$$\begin{aligned} \sup_{0 < \lambda \leq N} \lambda^p \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : I_\alpha^D \mu(x) > 2\lambda\}) &\leq C e^{-\frac{\varepsilon}{c}} \sup_{0 < \lambda \leq N} \lambda^p \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : I_\alpha^D \mu(x) > \lambda\}) \\ &\quad + C \sup_{0 < \lambda \leq N} \lambda^p \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : \mathcal{M}_\alpha \mu(x) > \varepsilon \lambda\}). \end{aligned}$$

Therefore,

$$\begin{aligned} \frac{1}{2^p} \sup_{0 < \lambda \leq 2N} \lambda^p \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : I_\alpha^D \mu(x) > \lambda\}) &\leq C e^{-\frac{\varepsilon}{c}} \sup_{0 < \lambda \leq 2N} \lambda^p \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : I_\alpha^D \mu(x) > \lambda\}) \\ &\quad + \frac{C}{\varepsilon^p} \|\mathcal{M}_\alpha \mu\|_{L^{p,\infty}(\mathcal{H}_\infty^\beta)}^p. \end{aligned}$$

Observe that the first term is finite because μ has compact support. Let B be a ball such that $\text{supp } \mu \subseteq B$. Using the same argument that before, we have

$$\begin{aligned} \sup_{0 < \lambda \leq 2N} \lambda^p \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : I_\alpha \mu(x) > \lambda\}) &\leq \sup_{0 < \lambda \leq 2N} \lambda^p \mathcal{H}_\infty^\beta(\{x \in 2B : I_\alpha \mu(x) > \lambda\}) \\ &+ \sup_{0 < \lambda \leq 2N} \lambda^p \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : \mathcal{M}_\alpha \mu(x) > \gamma(\alpha)\lambda\}) \\ &\leq (2N)^p r(2B)^\beta + \frac{1}{\gamma(\alpha)^p} \|\mathcal{M}_\alpha \mu\|_{L^{p,\infty}(\mathcal{H}_\infty^\beta)}^p \\ &< \infty, \end{aligned}$$

and thus, using (5.1), we obtain that the dyadic counterpart is also finite.

We take again ε such that $2^p C e^{-\frac{\varepsilon}{c}} = \frac{1}{2}$, that is $\varepsilon = \frac{c}{\log(2^{p+1}C)}$. Then, since the right-hand side is finite, we can absorb it into the left-hand side, and we obtain

$$\begin{aligned} \sup_{0 < \lambda \leq 2N} \lambda \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : I_\alpha^D \mu(x) > \lambda\})^{\frac{1}{p}} &\leq c_d \frac{\log(2^{p+1}C)}{c} \|\mathcal{M}_\alpha \mu\|_{L^{p,\infty}(\mathcal{H}_\infty^\beta)} \\ &\leq C' p \|\mathcal{M}_\alpha \mu\|_{L^{p,\infty}(\mathcal{H}_\infty^\beta)}. \end{aligned}$$

Letting $N \rightarrow \infty$, we obtain the desired inequality. The case of a general measure μ is obtained in the same way as before. □

Using Proposition 5.2, it is easy to see that (5.3) implies the following good lambda inequality for the Riesz potential.

Corollary 5.4. *Let $0 < \alpha < d$, $\beta \in (d - \alpha, d]$ and let μ be a locally finite measure. Then, there exist constants $C, c > 0$ depending on d, α, β such that*

$$(5.8) \quad \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : I_\alpha \mu(x) > 2^{d+1} \gamma_1 \lambda, \mathcal{M}_\alpha \mu(x) \leq \varepsilon \lambda\}) \leq C e^{-\frac{\varepsilon}{c}} \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : I_\alpha \mu(x) > \gamma_2^{-1} \lambda\})$$

for all $\lambda > 0$ and all $0 < \varepsilon < 1$, where $\gamma_1, \gamma_2 > 1$ are the constants in (5.2) and (5.1), respectively.

As a consequence of the good lambda inequality with exponential decay (5.8), we can prove the following result about the exponential integrability of $I_\alpha \mu$ for measures in the local Morrey space $\mathcal{M}^{n-\alpha}(B)$. This is the natural endpoint $p = \infty$ in Theorem 1.5. We give an alternative proof of a result of Adams and Xiao [6] (see also [10, Theorem 1.7]).

Theorem 5.5. *Let $0 < \alpha < d$ and let $\beta \in (d - \alpha, d]$. Then, there exist $C = C(d, \alpha, \beta) > 0$ and $\gamma = \gamma(d, \alpha, \beta) > 0$ such that*

$$\frac{1}{r(2B)^\beta} \int_{2B} \exp\left(\frac{\gamma I_\alpha \mu(x)}{\|\mu\|_{\mathcal{M}^{d-\alpha}(B)}}\right) d\mathcal{H}_\infty^\beta \leq C,$$

for balls B such that $\mu \in \mathcal{M}^{d-\alpha}(B)$.

Proof. Let B be a ball in \mathbb{R}^d such that $\|\mu\|_{\mathcal{M}^{d-\alpha}(B)} < \infty$. We can assume $\|\mu\|_{\mathcal{M}^{d-\alpha}(B)} \leq 1$. Observe that $r(B) \leq |x - y|$ for any $y \in B$ and $x \notin 2B$, hence

$$I_\alpha \mu(x) = \frac{1}{\gamma(\alpha)} \int_B \frac{d\mu(y)}{|x - y|^{d-\alpha}} \leq \frac{1}{\gamma(\alpha)} \frac{\mu(B)}{r(B)^{d-\alpha}} \leq \frac{1}{\gamma(\alpha)} \|\mu\|_{\mathcal{M}^{d-\alpha}(B)} \leq \frac{1}{\gamma(\alpha)},$$

for each $x \notin 2B$.

Therefore, if $\lambda > \frac{1}{\gamma(\alpha)}$ we have $\{x \in \mathbb{R}^d : I_\alpha \mu(x) > \lambda\} \subseteq 2B$. For fixed $\lambda > \frac{\gamma_2}{\gamma(\alpha)} > \frac{1}{\gamma(\alpha)}$, applying (5.8) with $\varepsilon = \frac{1}{\lambda}$ we obtain

$$\begin{aligned} \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : I_\alpha \mu(x) > 2^{d+1} \gamma_1 \lambda\}) &\leq C e^{-c\lambda} \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : I_\alpha \mu(x) > \gamma_2^{-1} \lambda\}) \\ &\quad + \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : \mathcal{M}_\alpha \mu(x) > 1\}) \\ &= C e^{-c\lambda} \mathcal{H}_\infty^\beta(\{x \in \mathbb{R}^d : I_\alpha \mu(x) > \gamma_2^{-1} \lambda\}) \\ &\leq C e^{-c\lambda} r(2B)^\beta. \end{aligned}$$

Using the previous estimate, we can conclude the proof,

$$\begin{aligned} \int_{2B} \exp\left(\frac{c}{2^{d+2}\gamma_1} I_\alpha \mu(x)\right) d\mathcal{H}_\infty^\beta &= \frac{c}{2^{d+2}\gamma_1} \int_0^\infty e^{\frac{c}{2^{d+2}\gamma_1} \lambda} \mathcal{H}_\infty^\beta(\{x \in 2B : I_\alpha \mu(x) > \lambda\}) d\lambda \\ &= \frac{c}{2} \int_0^\infty e^{\frac{c}{2} \lambda} \mathcal{H}_\infty^\beta(\{x \in 2B : I_\alpha \mu(x) > 2^{d+1} \gamma_1 \lambda\}) d\lambda \\ &= \frac{c}{2} \int_0^{\frac{\gamma_2}{\gamma(\alpha)}} e^{\frac{c}{2} \lambda} \mathcal{H}_\infty^\beta(\{x \in 2B : I_\alpha \mu(x) > 2^{d+1} \gamma_1 \lambda\}) d\lambda \\ &\quad + \frac{c}{2} \int_{\frac{\gamma_2}{\gamma(\alpha)}}^\infty e^{\frac{c}{2} \lambda} \mathcal{H}_\infty^\beta(\{x \in 2B : I_\alpha \mu(x) > 2^{d+1} \gamma_1 \lambda\}) d\lambda \\ &\leq \frac{c e^{\frac{c}{2}}}{2} \frac{\gamma_2}{\gamma(\alpha)} r(2B)^\beta + C \frac{c}{2} r(2B)^\beta \int_{\frac{\gamma_2}{\gamma(\alpha)}}^\infty e^{-\frac{c}{2} \lambda} d\lambda \\ &= C' r(2B)^\beta. \end{aligned}$$

□

ACKNOWLEDGMENTS

A. Claros is supported by the Basque Government through the BERC 2022-2025 program, by the Ministry of Science and Innovation: Grant PRE2021-099091 funded by BCAM Severo Ochoa accreditation CEX2021-001142-S/MICIN/AEI/10.13039/501100011033 and by ESF+.

CONFLICTS OF INTEREST

The authors have no conflicts of interest to declare.

DATA AVAILABILITY STATEMENT

Data sharing not applicable to this article as no datasets were generated or analyzed during the current study.

REFERENCES

- [1] D. R. Adams, *Traces of potentials. II*, Indiana Univ. Math. J. **22** (1972/73), 907–918, DOI 10.1512/iumj.1973.22.22075. MR313783 \uparrow 1
- [2] ———, *A note on Riesz potentials*, Duke Math. J. **42** (1975), no. 4, 765–778. MR458158 \uparrow 3, 4, 17
- [3] ———, *A note on Choquet integrals with respect to Hausdorff capacity*, Function spaces and applications (Lund, 1986), Lecture Notes in Math., vol. 1302, Springer, Berlin, 1988, pp. 115–124, DOI 10.1007/BFb0078867. MR942261 \uparrow 6
- [4] ———, *Choquet integrals in potential theory*, Publ. Mat. **42** (1998), 3–66. \uparrow 5, 6
- [5] ———, *Morrey spaces*, Lecture Notes in Applied and Numerical Harmonic Analysis, Birkhäuser/Springer, Cham, 2015. MR3467116 \uparrow 5
- [6] D. R. Adams and J. Xiao, *Erratum to: Morrey potentials and harmonic maps [MR2851148]*, Comm. Math. Phys. **339** (2015), no. 2, 769–771, DOI 10.1007/s00220-015-2409-6. MR3370618 \uparrow 25
- [7] T. Bagby and W. P. Ziemer, *Pointwise differentiability and absolute continuity*, Trans. Amer. Math. Soc. **191** (1974), 129–148, DOI 10.2307/1996986. MR0344390 \uparrow 6
- [8] Y.-W. Chen and D. Spector, *On functions of bounded β -dimensional mean oscillation*, Adv. Calc. Var., DOI <https://doi.org/10.1515/acv-2022-0084>. \uparrow 1, 4, 5, 6, 7, 8, 9
- [9] You-Wei Benson Chen and Keng Hao Ooi and Daniel Spector, *Capacitary Maximal Inequalities and Applications* (2023), available at 2305.19046. \uparrow 2, 7, 15
- [10] Y.-W. B. Chen, *A self-improving property of Riesz potentials in BMO*, 2024. \uparrow 3, 16, 25
- [11] A. Cianchi, *Moser-Trudinger trace inequalities*, Adv. Math. **217** (2008), no. 5, 2005–2044, DOI 10.1016/j.aim.2007.09.007. MR2388084 \uparrow 1
- [12] D. Cruz-Uribe and K. Moen, *One and two weight norm inequalities for Riesz potentials*, Illinois J. Math. **57** (2013), no. 1, 295–323. MR3224572 \uparrow 18, 19
- [13] L. Fontana and C. Morpurgo, *Adams inequalities on measure spaces*, Adv. Math. **226** (2011), no. 6, 5066–5119, DOI 10.1016/j.aim.2011.01.003. MR2775895 \uparrow 1
- [14] C. Fefferman and E. M. Stein, *H^p spaces of several variables*, Acta Math. **129** (1972), no. 3-4, 137–193, DOI 10.1007/BF02392215. MR447953 \uparrow 2, 9
- [15] P. Harjulehto and R. Hurri-Syrjänen, *On Choquet integrals and Poincaré-Sobolev inequalities*, J. Funct. Anal. **284** (2023), no. 9, Paper No. 109862, 18, DOI 10.1016/j.jfa.2023.109862. MR4545158 \uparrow 5
- [16] P. Harjulehto and R. Hurri-Syrjänen, *On Sobolev inequalities with Choquet integrals*, 2023. \uparrow 15
- [17] P. Honzik and B. J. Jaye, *On the good- λ inequality for nonlinear potentials*, Proc. Amer. Math. Soc. **140** (2012), no. 12, 4167–4180, DOI 10.1090/S0002-9939-2012-11352-8. MR2957206 \uparrow 18, 19
- [18] P. Harjulehto and R. Hurri-Syrjänen, *On Hausdorff content maximal operator and Riesz potential for non-measurable functions* (2024), available at 2405.12113v1. \uparrow 2
- [19] F. John and L. Nirenberg, *On functions of bounded mean oscillation*, Comm. Pure Appl. Math. **14** (1961), 415–426, DOI 10.1002/cpa.3160140317. MR131498 \uparrow 1
- [20] Á. D. Martínez and D. Spector, *An improvement to the John-Nirenberg inequality for functions in critical Sobolev spaces*, Adv. Nonlinear Anal. **10** (2021), no. 1, 877–894, DOI 10.1515/anona-2020-0157. MR4191703 \uparrow 1
- [21] B. Muckenhoupt and R. Wheeden, *Weighted norm inequalities for fractional integrals*, Trans. Amer. Math. Soc. **192** (1974), 261–274, DOI 10.2307/1996833. MR0340523 \uparrow 3, 19
- [22] J. Oröbitg and J. Verdera, *Choquet integrals, Hausdorff content and the Hardy-Littlewood maximal operator*, Bull. London Math. Soc. **30** (1998), 145–150. \uparrow 7, 15
- [23] A. C. Ponce and D. Spector, *Some remarks on capacitary integrals and measure theory*, Potentials and partial differential equations—the legacy of David R. Adams, Adv. Anal. Geom., vol. 8, De Gruyter, Berlin, [2023] ©2023, pp. 235–263. MR4654520 \uparrow 24
- [24] H. Saito, H. Tanaka, and T. Watanabe, *Abstract dyadic cubes, maximal operators and Hausdorff content*, Bull. Sci. Math. **140** (2016), no. 6, 757–773, DOI 10.1016/j.bulsci.2016.02.001. MR3543752 \uparrow 5
- [25] E. T. Sawyer, *A characterization of a two-weight norm inequality for maximal operators*, Studia Math. **75** (1982), no. 1, 1–11, DOI 10.4064/sm-75-1-1-11. MR0676801 \uparrow 6
- [26] E. Sawyer and R. L. Wheeden, *Weighted inequalities for fractional integrals on Euclidean and homogeneous spaces*, Amer. J. Math. **114** (1992), no. 4, 813–874, DOI 10.2307/2374799. MR1175693 \uparrow 18

- [27] D. Yang and W. Yuan, *A note on dyadic Hausdorff capacities*, Bull. Sci. Math. **132** (2008), no. 6, 500–509, DOI 10.1016/j.bulsci.2007.06.005. MR2445577 ↑5
- [28] V. I. Judovič, *Some estimates connected with integral operators and with solutions of elliptic equations*, Dokl. Akad. Nauk SSSR **138** (1961), 805–808 (Russian). MR0140822 ↑1

(Y.-W. Chen) NATIONAL TAIWAN UNIVERSITY, DEPARTMENT OF MATHEMATICS, ASTRONOMY MATHEMATICS BUILDING 5F, No. 1, SEC. 4, ROOSEVELT RD., TAIPEI 10617, TAIWAN (R.O.C.)

Email address: `bensonchen.sc07@nycu.edu.tw`

(A. Claros) BCAM - BASQUE CENTER FOR APPLIED MATHEMATICS, BILBAO, SPAIN

Email address: `aclaros@bcamath.org`