

# ON CERTAIN DETERMINANTS AND THE SQUARE ROOT OF SOME DETERMINANTS INVOLVING LEGENDRE SYMBOLS

CHEN-KAI REN AND XIN-QI LUO

ABSTRACT. Let  $p > 3$  be a prime and  $\left(\frac{\cdot}{p}\right)$  be the Legendre symbol. For any integer  $d$  with  $p \nmid d$  and any positive integer  $m$ , Sun introduced the determinants

$$T_m(d, p) = \det \left[ (i^2 + dj^2)^m \left( \frac{i^2 + dj^2}{p} \right) \right]_{1 \leq i, j \leq (p-1)/2},$$

and

$$D_p^{(m)} = \det \left[ (i^2 - j^2)^m \left( \frac{i^2 - j^2}{p} \right) \right]_{1 \leq i, j \leq (p-1)/2}.$$

In this paper, we obtain some properties of  $T_m(d, p)$  and  $\sqrt{D_p^{(m)}}$  for some  $m$ . We also confirm some related conjectures posed by Zhi-Wei Sun.

## 1. INTRODUCTION

Determinants of certain special matrices are useful in many branches of mathematics. Readers may refer to Krattenthaler's survey paper [5] for recent progress and advanced techniques on this topic. In this paper, we study some determinants involving Legendre symbols.

Let  $p$  be an odd prime, and let  $\left(\frac{\cdot}{p}\right)$  be the Legendre symbol. Carlitz [1] determined the characteristic polynomial of the matrix

$$\left[ x + \left( \frac{i-j}{p} \right) \right]_{1 \leq i, j \leq p-1},$$

and Chapman [2] used quadratic Gauss sums to determine the values of the determinant

$$\det \left[ x + \left( \frac{i+j-1}{p} \right) \right]_{1 \leq i, j \leq (p-1)/2}.$$

Vsemirnov [13, 14] confirmed a challenging conjecture of Chapman by evaluating the determinant

$$\det \left[ \left( \frac{j-i}{p} \right) \right]_{1 \leq i, j \leq (p+1)/2}.$$

---

*Key words and phrases.* determinant, Legendre Symbol.

2020 *Mathematics Subject Classification.* Primary 11A25; Secondary 11N25.

Supported by the Natural Science Foundation of China (grant no. 12371004).

Let  $d$  be any integer. Sun [9] introduced the determinant

$$S(d, p) = \det \left[ \left( \frac{i^2 + dj^2}{p} \right) \right]_{1 \leq i, j \leq (p-1)/2}.$$

Sun [9, Theorem 1.2] proved that

$$\left( \frac{S(d, p)}{p} \right) = \begin{cases} \left( \frac{-1}{p} \right) & \text{if } \left( \frac{d}{p} \right) = 1, \\ 0 & \text{if } \left( \frac{d}{p} \right) = -1. \end{cases}$$

D. Grinberg, Sun and L. Zhao [3] showed that if  $p > 3$  then

$$\det \left[ (i^2 + dj^2) \left( \frac{i^2 + dj^2}{p} \right) \right]_{0 \leq i, j \leq (p-1)/2} \equiv 0 \pmod{p}.$$

Motivated by this, for any positive integer  $n$  with  $(p-1)/2 \leq n \leq p-1$  Sun introduced the determinant

$$S_n(d, p) = \det \left[ (i^2 + dj^2)^n \right]_{1 \leq i, j \leq (p-1)/2}.$$

In 2022, H.-L. Wu, Y.-F. She and L.-Y. Wang [15] proved the conjecture of Sun [9, Conjecture 4.5] that if  $p > 3$  and  $p \nmid d$  then

$$\left( \frac{S_{(p+1)/2}(d, p)}{p} \right) = \begin{cases} \left( \frac{d}{p} \right)^{(p-1)/4} & \text{if } p \equiv 1 \pmod{4}, \\ \left( \frac{d}{p} \right)^{(p+1)/4} (-1)^{h(-p)-1/2} & \text{if } p \equiv 3 \pmod{4}, \end{cases}$$

where  $h(-p)$  denotes the class number of the imaginary quadratic field  $\mathbb{Q}(\sqrt{-p})$ .

For any prime  $p \equiv 3 \pmod{4}$ , Sun [9, Remark 1.3] showed that

$$S_{p-2}(1, p) = \det \left[ \frac{1}{i^2 + j^2} \right]_{1 \leq i, j \leq (p-1)/2} \equiv \left( \frac{2}{p} \right) \pmod{p}.$$

Sun [12] proved that if  $p \equiv 3 \pmod{4}$ ,

$$\left( \frac{S_{p-3}(1, p)}{p} \right) = 1.$$

Ren and Sun [7] solved some conjectures of Sun concerning the Legendre symbols of  $S_{p-2}$ ,  $S_{p-3}$  and  $S_{p-4}$ . Sun posed the following conjecture [12, Remark 1.1].

**Conjecture 1.1.** *Let  $p > 3$  be a prime with  $p \neq 11$ . Let*

$$T_2(1, p) = \det \left[ (i^2 + j^2)^2 \left( \frac{i^2 + j^2}{p} \right) \right]_{1 \leq i, j \leq (p-1)/2},$$

where  $\left( \frac{\cdot}{p} \right)$  is the Legendre symbol. Then

$$\left( \frac{T_2(1, p)}{p} \right) = \begin{cases} \left( \frac{2}{p} \right) & \text{if } p \equiv 1 \pmod{4}, \\ \left( \frac{-6}{p} \right) & \text{if } p \equiv 3 \pmod{4}. \end{cases}$$

In this paper, we confirm this conjecture. Inspired by this, we start to study determinants on finite field. Let  $q = 2n + 1$  be an odd prime power and let  $\phi$  denote the unique quadratic multiplicative character of  $\mathbb{F}_q$ , which is the map  $\mathbb{F}_q \rightarrow \mathbb{C}$  that sends 0 to 0, each non-zero square to 1, and each non-square to  $-1$ . Let  $\mathbb{F}_q^\times := \mathbb{F}_q \setminus \{0\}$  and let

$$\mathbb{F}_q^{\times 2} := \{x^2 : x \in \mathbb{F}_q^\times\} = \{a_1, a_2, \dots, a_n\}.$$

We define

$$\tilde{T}_m(d, q) = \det[(a_i + da_j)^m \phi(a_i + da_j)]_{1 \leq i, j \leq n}.$$

We have the following generalized result.

**Theorem 1.1.** *Let  $q = 2n + 1$  be an odd prime power and let  $d \in \mathbb{F}_q^\times$ . Suppose  $\text{char}(\mathbb{F}_q) > 3$ . Let*

$$\tilde{T}_2(d, q) = \det[(a_i + da_j)^2 \phi(a_i + da_j)]_{1 \leq i, j \leq n}.$$

*If we view  $\tilde{T}_2(d, q)$  as a determinant over  $\mathbb{F}_q$ , then the following results hold:*

*(i) If  $q \equiv 1 \pmod{4}$ , then*

$$\tilde{T}_2(d, q) = d^{\frac{q-1}{4}} \times \frac{q-1}{2}! x_q(d)^2$$

*for some  $x_q(d) \in \mathbb{F}_q$ .*

*(ii) If  $q \equiv 3 \pmod{4}$ , then*

$$\tilde{T}_2(d, q) = (-1)^{\frac{q-3}{4}} 3d^{\frac{q+1}{4}} y_q(d)^2$$

*for some  $y_q(d) \in \mathbb{F}_q$ .*

Applying this result, we can obtain Conjecture 1.1.

**Corollary 1.1.** *Conjecture 1.1 holds.*

**Theorem 1.2.** *Let  $q = 2n + 1$  be an odd prime power and let  $d \in \mathbb{F}_q^\times$ . Suppose  $\text{char}(\mathbb{F}_q) > 7$ . Let*

$$\tilde{T}_{(q-11)/2}(d, q) = \det[(a_i + da_j)^{(q-11)/2} \phi(a_i + da_j)]_{1 \leq i, j \leq n}.$$

*If we view  $\tilde{T}_{(q-11)/2}(d, q)$  as a determinant over  $\mathbb{F}_q$ , then the following results hold:*

*(i) If  $q \equiv 1 \pmod{4}$ , then*

$$\tilde{T}_{(q-11)/2}(d, q) = d^{\frac{q-1}{4}} x_q(d)^2$$

*for some  $x_q(d) \in \mathbb{F}_q$ .*

*(ii) If  $q \equiv 3 \pmod{4}$ , then*

$$\tilde{T}_{(q-11)/2}(d, q) = 7(q-1)! d^{\frac{q+1}{4}} y_q(d)^2$$

*for some  $y_q(d) \in \mathbb{F}_q$ .*

Applying this result, we can get the following corollary.

**Corollary 1.2.** *Let  $p > 7$  be a prime and let  $d$  be any integer with  $\left(\frac{d}{p}\right) = 1$ . Let*

$$T_{(p-11)/2}(d, p) = \det \left[ (i^2 + dj^2)^{(p-11)/2} \left( \frac{i^2 + dj^2}{p} \right) \right]_{1 \leq i, j \leq (p-1)/2}.$$

*If  $\left(\frac{T_{(p-11)/2}(d, p)}{p}\right) = -1$ , then  $p \equiv 1$  or  $2$  or  $4 \pmod{7}$ .*

Let  $m, n \in \mathbb{Z}^+$  with  $n$  odd. Sun introduced the determinant

$$D_n^{(m)} = \det \left[ (i^2 - j^2)^m \left( \frac{i^2 - j^2}{n} \right) \right]_{1 \leq i, j \leq (n-1)/2}.$$

If  $2 \nmid m$  and  $4 \mid n - 1$ , then  $D_n^{(m)}$  is skew-symmetric and of even order. Hence, it is the square of a certain integer by Cayley's Theorem [8]. We also prove the following conjectures of Sun [12].

**Theorem 1.3.** *For any positive odd integer  $m$ , the set*

$$E(m) = \{p : p \text{ is a prime with } 4 \mid p - 1 \text{ and } p \mid D_p^{(m)}\}$$

*is finite. In particular,*

$$E(5) = \{29\}, \quad E(7) = \{13, 53\}, \quad E(9) = \{13, 17, 29\}$$

$$E(11) = \{17, 29\} \text{ and } E(13) = \{17, 109, 401\}.$$

**Theorem 1.4.** *For any prime  $p \equiv 1 \pmod{4}$ , we have*

$$\left( \frac{\sqrt{D_p^{(1)}}}{p} \right) = (-1)^{|\{0 < k < \frac{p}{4} : \left(\frac{k}{p}\right) = -1\}|} \left( \frac{p}{3} \right).$$

**Theorem 1.5.** *For any prime  $p \equiv 1 \pmod{4}$ , we have*

$$\left( \frac{\sqrt{D_p^{(3)}}}{p} \right) = (-1)^{|\{0 < k < \frac{p}{4} : \left(\frac{k}{p}\right) = -1\}|} \left( \frac{p}{4 + (-1)^{(p-1)/4}} \right).$$

## 2. SOME PREPARATIONS BEFORE THE PROOF

Let  $q = 2n + 1$  be an odd prime power and recall  $\phi$  denote the unique quadratic multiplicative character of  $\mathbb{F}_q$ . Let  $\mathbb{F}_q^\times := \mathbb{F}_q \setminus \{0\}$  and let

$$\mathbb{F}_q^{\times 2} := \{x^2 : x \in \mathbb{F}_q^\times\} = \{a_1, a_2, \dots, a_n\}.$$

For any  $x, y \in \mathbb{F}_q$ , we write  $x \equiv y \pmod{\mathbb{F}_q^{\times 2}}$  if there is an element  $z \in \mathbb{F}_q^\times$  such that  $x = yz^2$ .

We need the following lemma [5] on determinants.

**Lemma 2.1.** *Let  $R$  be a commutative ring with identity, and let  $P(x) = \sum_{i=0}^{n-1} b_i x^i \in R[x]$ . Then we have*

$$\det [P(X_i Y_j)]_{1 \leq i < j \leq n} = b_0 b_1 \cdots b_{n-1} \prod_{1 \leq i < j \leq n} (X_i - X_j)(Y_i - Y_j).$$

Let  $a$  be an integer with  $(a, n) = 1$ . We notice that multiplication by  $a$  introduces a permutation  $\tau_a$  of  $\mathbb{Z}/n\mathbb{Z}$ . Lerch [6] obtained the following result which determines the sign of  $\tau_a$ .

**Lemma 2.2.** *Let  $\text{sgn}(\tau_a^{(n)})$  denote the sign of  $\tau_a$ . Then*

$$\text{sgn}(\tau_a^{(n)}) = \begin{cases} \left(\frac{a}{n}\right) & \text{if } n \equiv 1 \pmod{2}, \\ 1 & \text{if } n \equiv 2 \pmod{4}, \\ (-1)^{\frac{a-1}{2}} & \text{if } n \equiv 0 \pmod{4}, \end{cases}$$

where  $\left(\frac{\cdot}{n}\right)$  denotes Jacobi symbol if  $n$  is odd.

Clearly  $\text{inv}_q : a_i \rightarrow a_i^{-1}$  is a permutation of  $a_1, \dots, a_n$ . Fix a generator  $g$  of the cyclic group  $\mathbb{F}_q^\times$ . If we reindex the elements  $g^0, g^1, \dots, g^{n-1}$  of  $\mathbb{F}_q^\times$  as  $0, 1, \dots, n-1$  of  $\mathbb{Z}/n\mathbb{Z}$ , then the permutation  $\text{inv}_q$  becomes the permutation  $\tau_{-1}$  that sends  $i$  to  $n-i$  for  $1 \leq i \leq n-1$  while leaving 0 fixed. This permutation has  $\frac{(n-1)(n-2)}{2}$  inversions. In view of the above we obtain the following result.

**Lemma 2.3.** *Let notations be as above. Then*

$$\text{sgn}(\text{inv}_q) = (-1)^{\frac{(n-1)(n-2)}{2}} = (-1)^{\frac{(q-3)(q-5)}{8}}.$$

### 3. PROOFS OF THEOREM 1.1 AND 1.2

**Proof of Theorem 1.1.** Recall that  $\phi$  denote the unique quadratic multiplicative character of  $\mathbb{F}_q$ . If we view  $\tilde{T}_2(d, q)$  as a determinant over  $\mathbb{F}_q$ , then we have

$$\tilde{T}_2(d, q) = \det \left[ (a_i + da_j)^{\frac{q+3}{2}} \right]_{1 \leq i, j \leq n}.$$

Thus

$$\tilde{T}_2(d, q) \equiv \det \left[ \left( \frac{a_i}{a_j} + d \right)^{\frac{q+3}{2}} \right]_{1 \leq i, j \leq n} \pmod{\mathbb{F}_q^{\times 2}}.$$

Noting that

$$\left( \frac{a_i}{a_j} \right)^{\frac{q-1}{2}} = 1 \text{ and } d^{\frac{q-1}{2}} = \phi(d),$$

we can verify that

$$\left( \frac{a_i}{a_j} + d \right)^{\frac{q+3}{2}}$$

$$\begin{aligned}
&= \sum_{k=0}^{\frac{q+3}{2}} \binom{\frac{q+3}{2}}{k} \left(\frac{a_i}{a_j}\right)^k d^{\frac{q+3}{2}-k} \\
&= \sum_{k=3}^{\frac{q-3}{2}} \binom{\frac{q+3}{2}}{k} \left(\frac{a_i}{a_j}\right)^k d^{\frac{q+3}{2}-k} + d^{\frac{q+3}{2}} + \frac{q+3}{2} \left(\frac{a_i}{a_j}\right) d^{\frac{q+1}{2}} + \frac{(q+3)(q+1)}{8} d^{\frac{q-1}{2}} \left(\frac{a_i}{a_j}\right)^2 \\
&\quad + \frac{(q+3)(q+1)}{8} d^2 \left(\frac{a_i}{a_j}\right)^{\frac{q-1}{2}} + \frac{q+3}{2} d \left(\frac{a_i}{a_j}\right)^{\frac{q+1}{2}} + \left(\frac{a_i}{a_j}\right)^{\frac{q+3}{2}} \\
&= \sum_{k=3}^{\frac{q-3}{2}} \binom{\frac{q+3}{2}}{k} \left(\frac{a_i}{a_j}\right)^k d^{\frac{q+3}{2}-k} + \phi(d)d^2 + \frac{3}{2} \left(\frac{a_i}{a_j}\right) d\phi(d) + \frac{3}{8}\phi(d) \left(\frac{a_i}{a_j}\right)^2 + \frac{3}{8}d^2 + \frac{3}{2}d \left(\frac{a_i}{a_j}\right) + \left(\frac{a_i}{a_j}\right)^2 \\
&= \sum_{k=3}^{\frac{q-3}{2}} \binom{\frac{q+3}{2}}{k} \left(\frac{a_i}{a_j}\right)^k d^{\frac{q+3}{2}-k} + \left(\frac{3}{8} + \phi(d)\right) d^2 + \frac{3}{2} \left(\frac{a_i}{a_j}\right) d(\phi(d) + 1) + \left(\frac{a_i}{a_j}\right)^2 \left(\frac{3}{8}\phi(d) + 1\right).
\end{aligned}$$

Hence

$$\left(\frac{a_i}{a_j} + d\right)^{\frac{q+3}{2}} = f\left(\frac{a_i}{a_j}\right)$$

where

$$\begin{aligned}
f(T) &= \left(\frac{3}{8} + \phi(d)\right) d^2 + \frac{3}{2}d(\phi(d) + 1)T + \left(\frac{3}{8}\phi(d) + 1\right) T^2 \\
&\quad + \sum_{k=3}^{\frac{q-3}{2}} \binom{\frac{q+3}{2}}{k} T^k d^{\frac{q+3}{2}-k}.
\end{aligned}$$

Observe that

$$\left(\frac{3}{8} + \phi(d)\right) \left(\frac{3}{8}\phi(d) + 1\right) \equiv 0 \text{ or } \phi(d) \pmod{\mathbb{F}_q^{\times 2}}$$

(just check both cases  $\phi(d) = 1$  or  $\phi(d) = -1$  and consider whether  $\frac{3}{8} + \phi(d) \equiv 0 \pmod{q}$ ).

Let  $C_f$  denote the product of coefficients of  $f(T)$ , then

$$\begin{aligned}
C_f &= \left(\frac{3}{8} + \phi(d)\right) \frac{3}{2} d^3 (\phi(d) + 1) \left(\frac{3}{8}\phi(d) + 1\right) \prod_3^{\frac{q-3}{2}} \binom{\frac{q+3}{2}}{k} d^{\frac{q+3}{2}-k} \\
&= \frac{3}{2} d^3 (\phi(d) + 1) \left(\frac{3}{8} + \phi(d)\right) \left(\frac{3}{8}\phi(d) + 1\right) \prod_3^{\frac{q-3}{2}} \binom{\frac{q+3}{2}}{k} \times d^{\frac{(q+3)(q-7)}{8}}
\end{aligned}$$

$$\equiv 6d^{\frac{(q-3)(q-1)}{8}}(\phi(d) + 1)\left(\frac{3}{8} + \phi(d)\right)\left(\frac{3}{8}\phi(d) + 1\right) \prod_3^{\frac{q-3}{2}} \binom{\frac{q+3}{2}}{k} \pmod{\mathbb{F}_q^{\times 2}}.$$

Thus applying Lemma 2.1 to  $P(T) = f(T)$ , we rewrite as

$$\begin{aligned} \tilde{T}_2(d, q) &\equiv \text{sgn}(\text{inv}_q) 6d^{\frac{(q-3)(q-1)}{8}}(\phi(d) + 1)\left(\frac{3}{8} + \phi(d)\right)\left(\frac{3}{8}\phi(d) + 1\right) \prod_3^{\frac{q-3}{2}} \binom{\frac{q+3}{2}}{k} \prod_{1 \leq i < j \leq n} (a_j - a_i)^2 \\ &\equiv \text{sgn}(\text{inv}_q) 6d^{\frac{(q-3)(q-1)}{8}}(\phi(d) + 1)\left(\frac{3}{8} + \phi(d)\right)\left(\frac{3}{8}\phi(d) + 1\right) \prod_3^{\frac{q-3}{2}} \binom{\frac{q+3}{2}}{k} \pmod{\mathbb{F}_q^{\times 2}}. \end{aligned}$$

Now we divide the remaining proof into two cases.

**Case 1.**  $q \equiv 1 \pmod{4}$ .

In this case, since  $\pm 1 \in \mathbb{F}_q^{\times 2}$  and  $\text{char}(\mathbb{F}_q) > 3$ , we can verify that

$$\text{sgn}(\text{inv}_q) 6d^{\frac{(q-3)(q-1)}{8}} \equiv 6d^{\frac{q-1}{4}} \pmod{\mathbb{F}_q^{\times 2}}$$

and that

$$\prod_{k=3}^{\frac{q-3}{2}} \binom{\frac{q+3}{2}}{k} \in \{0\} \cup \left(\frac{\frac{q+3}{2}}{\frac{q+3}{4}}\right) \mathbb{F}_q^{\times 2}$$

where  $\left(\frac{\frac{q+3}{2}}{\frac{q+3}{4}}\right) \equiv 3 \times \frac{q-1}{2}! \pmod{\mathbb{F}_q^{\times 2}}$ . Hence, we have

$$\begin{aligned} \tilde{T}_2(d, q) &\equiv 6d^{\frac{q-1}{4}}(\phi(d) + 1)\phi(d) \times 3\frac{q-1}{2}! \text{ or } 0 \\ &\equiv d^{\frac{q-1}{4}} \times \frac{q-1}{2}! \text{ or } 0 \pmod{\mathbb{F}_q^{\times 2}} \end{aligned}$$

Therefore, there is an element  $x_q(d) \in \mathbb{F}_q$  such that

$$\tilde{T}_2(d, q) = d^{\frac{q-1}{4}} \times \frac{q-1}{2}! x_q(d)^2.$$

**Case 2.**  $q \equiv 3 \pmod{4}$ .

In this case, we have

$$\prod_{k=3}^{\frac{q-3}{2}} \binom{\frac{q+3}{2}}{k} \in \{0\} \cup \mathbb{F}_q^{\times 2}.$$

By Lemma 2.3, we obtain that

$$\text{sgn}(\text{inv}_q) 6d^{\frac{(q-3)(q-1)}{8}} \equiv (-1)^{\frac{q-3}{4}} 6d^{\frac{q-3}{4}} \pmod{\mathbb{F}_q^{\times 2}}.$$

Noting that  $\phi(d) \equiv d \pmod{\mathbb{F}_q^{\times 2}}$ , we verify that

$$\begin{aligned}\tilde{T}_2(d, q) &\equiv (-1)^{\frac{q-3}{4}} 6d^{\frac{q-3}{4}} (\phi(d) + 1)\phi(d) \text{ or } 0 \\ &\equiv (-1)^{\frac{q-3}{4}} 3d^{\frac{q+1}{4}} \text{ or } 0 \pmod{\mathbb{F}_q^{\times 2}}.\end{aligned}$$

In view of the above there is an element  $y_q(d) \in \mathbb{F}_q$  such that

$$\tilde{T}_2(d, q) = (-1)^{\frac{q-3}{4}} 3d^{\frac{q+1}{4}} y_q(d)^2.$$

□

**Proof of Corollary 1.1.** If we view  $T_2(1, p)$  as the determinant over  $\mathbb{F}_p$ , we can take advantage of the proof above. Take  $d = 1$  and let  $q = p$  with  $p \geq 5$  and  $p \neq 11$ . We obtain that

$$\begin{aligned}T_2(1, p) &\equiv 3 \times \text{sgn}(\text{inv}_p) \left(\frac{3}{8} + 1\right)^2 \prod_3^{\frac{p-3}{2}} \binom{\frac{p+3}{2}}{k} \\ &\equiv 3 \times \text{sgn}(\text{inv}_p) \prod_3^{\frac{p-3}{2}} \binom{\frac{p+3}{2}}{k} \pmod{\mathbb{F}_p^{\times 2}}.\end{aligned}$$

Also, we divide the remaining proof into two cases.

**Case 1.**  $p \equiv 1 \pmod{4}$ . In this case, we have

$$\begin{aligned}T_2(1, p) &\equiv 3 \binom{\frac{p+3}{2}}{\frac{p+3}{4}} \prod_3^{\frac{p-1}{4}} \binom{\frac{p+3}{2}}{k}^2 \\ &\equiv \frac{p-1}{2}! \pmod{\mathbb{F}_p^{\times 2}}.\end{aligned}$$

Hence, by Lemma [9, Lemma 2.3], we have  $\left(\frac{T_2(1, p)}{p}\right) = \left(\frac{(\frac{p-1}{2})!}{p}\right) = \left(\frac{2}{p}\right)$ .

**Case 2.**  $p \equiv 3 \pmod{4}$ . In this case, by Lemma 2.3 we have

$$\begin{aligned}T_2(1, p) &\equiv 3 \times \text{sgn}(\text{inv}_q) \prod_3^{\frac{p+1}{4}} \binom{\frac{p+3}{2}}{k}^2 \\ &\equiv 3(-1)^{\frac{p-3}{4}} \pmod{\mathbb{F}_p^{\times 2}}.\end{aligned}$$



Therefore, for  $\left(\frac{(-1)^{\frac{p-3}{4}}}{p}\right) = (-1)^{\frac{p-3}{4}} = -1 \times (-1)^{\frac{(p-1)(p+1)}{8}} = \left(\frac{-2}{p}\right)$ , we obtain

$$\left(\frac{T_2(1, p)}{p}\right) = \left(\frac{3(-1)^{\frac{p-3}{4}}}{p}\right) = \left(\frac{-6}{p}\right).$$

□

**Proof of Theorem 1.2.** Recall that  $\phi$  denote the unique quadratic multiplicative character of  $\mathbb{F}_q$ . If we view  $\tilde{T}_{(q-11)/2}(d, q)$  as a determinant over  $\mathbb{F}_q$ , then we have

$$\tilde{T}_{(q-11)/2}(d, q) = \det [(a_i + da_j)^{q-6}]_{1 \leq i, j \leq n}.$$

Thus

$$\tilde{T}_{(q-11)/2}(d, q) \equiv \det \left[ \left( \frac{a_i}{a_j} + d \right)^{q-6} \right]_{1 \leq i, j \leq n} \pmod{\mathbb{F}_q^{\times 2}}.$$

Noting that

$$\left(\frac{a_i}{a_j}\right)^{\frac{q-1}{2}} = 1 \text{ and } d^{\frac{q-1}{2}} = \phi(d),$$

we can verify that

$$\begin{aligned} & \left(\frac{a_i}{a_j} + d\right)^{q-6} \\ &= \sum_{k=0}^{q-6} \binom{q-6}{k} \left(\frac{a_i}{a_j}\right)^k d^{q-6-k} \\ &= \sum_{k=0}^{\frac{q-11}{2}} \left( \binom{q-6}{k} d^{q-6-k} + \binom{\frac{q-6}{2} + k}{k} d^{q-6-k-(q-1)/2} \right) \left(\frac{a_i}{a_j}\right)^k \\ & \quad + \sum_{k=0}^3 \binom{q-6}{\frac{q-9}{2} + k} d^{(q-3)/2-k} \left(\frac{a_i}{a_j}\right)^{(q-9)/2+k} \\ &= \sum_{k=0}^{\frac{q-11}{2}} \left( \binom{q-6}{k} + \phi(d)^{-1} \binom{q-6}{\frac{q-11}{2} - k} \right) d^{q-6-k} \left(\frac{a_i}{a_j}\right)^k \\ & \quad + \sum_{k=0}^3 \binom{q-6}{\frac{q-9}{2} + k} d^{(q-3)/2-k} \left(\frac{a_i}{a_j}\right)^{(q-9)/2+k}. \end{aligned}$$

Hence

$$\left(\frac{a_i}{a_j} + d\right)^{q-6} = f \left(\frac{a_i}{a_j}\right)$$

where

$$f(T) = \sum_{k=0}^{\frac{q-11}{2}} \left( \binom{q-6}{k} + \phi(d)^{-1} \binom{q-6}{\frac{q-11}{2}-k} \right) d^{q-6-k} T^k \\ + \sum_{k=0}^3 \binom{q-6}{\frac{q-9}{2}+k} d^{(q-3)/2-k} T^{(q-9)/2+k}.$$

Observe that

$$\left( \binom{q-6}{k} + \phi(d)^{-1} \binom{q-6}{\frac{q-11}{2}-k} \right) \left( \binom{q-6}{\frac{q-11}{2}-k} + \phi(d)^{-1} \binom{q-6}{k} \right) \equiv 0 \text{ or } \phi(d) \pmod{\mathbb{F}_q^{\times 2}}$$

(just check both cases  $\phi(d) = 1$  or  $\phi(d) = -1$  and consider whether  $\binom{q-6}{k} + \phi(d)^{-1} \binom{q-6}{\frac{q-11}{2}-k} \equiv 0 \pmod{q}$ ). Let  $C_f$  denote the product of coefficients of  $f(T)$ , then

$$C_f = \prod_{k=0}^3 \binom{q-6}{\frac{q-9}{2}+k} \prod_{k=0}^3 d^{(q-3)/2-k} \prod_{k=0}^{\frac{q-11}{2}} \left( \binom{q-6}{k} + \phi(d)^{-1} \binom{q-6}{\frac{q-11}{2}-k} \right) \prod_{k=0}^{\frac{q-11}{2}} d^{q-6-k} \\ = \binom{q-6}{\frac{q-9}{2}}^2 \binom{q-6}{\frac{q-7}{2}}^2 \prod_{k=0}^{\frac{q-11}{2}} \left( \binom{q-6}{k} + \phi(d)^{-1} \binom{q-6}{\frac{q-11}{2}-k} \right) d^{\frac{(q-1)(3q-21)}{8}}.$$

Applying Lemma 2.1 to  $P(T) = f(T)$ , we rewrite as

$$\tilde{T}_{(q-11)/2}(d, q) \equiv \text{sgn}(\text{inv}_q) \prod_{k=0}^{\frac{q-11}{2}} \left( \binom{q-6}{k} + \phi(d)^{-1} \binom{q-6}{\frac{q-11}{2}-k} \right) d^{\frac{(q-1)(3q-21)}{8}} \\ \times \binom{q-6}{\frac{q-9}{2}}^2 \binom{q-6}{\frac{q-7}{2}}^2 \prod_{1 \leq i < j \leq n} (a_j - a_i)^2 \\ \equiv \text{sgn}(\text{inv}_q) \prod_{k=0}^{\frac{q-11}{2}} \left( \binom{q-6}{k} + \phi(d)^{-1} \binom{q-6}{\frac{q-11}{2}-k} \right) \\ \times \binom{q-6}{\frac{q-9}{2}}^2 \binom{q-6}{\frac{q-7}{2}}^2 d^{\frac{(q-1)(3q-21)}{8}} \pmod{\mathbb{F}_q^{\times 2}}.$$

Now we divide the remaining proof into two cases.

**Case 1.**  $q \equiv 1 \pmod{4}$ .

Since  $\pm 1 \in \mathbb{F}_q^{\times 2}$  and  $\text{char}(\mathbb{F}_q) > 7$ , we can verify that

$$\begin{aligned}
 \tilde{T}_{(q-11)/2}(d, q) &\equiv \text{sgn}(\text{inv}_q) \prod_{k=0}^{\frac{q-11}{2}} \left( \binom{q-6}{k} + \phi(d)^{-1} \binom{q-6}{\frac{q-11}{2}-k} \right) \left( \frac{q-6}{2} \right)^2 \left( \frac{q-6}{2} \right)^2 d^{\frac{(q-1)(3q-21)}{8}} \\
 &\equiv \prod_{k=0}^{\frac{q-13}{4}} \left( \binom{q-6}{k} + \phi(d)^{-1} \binom{q-6}{\frac{q-11}{2}-k} \right) \left( \binom{q-6}{\frac{q-11}{2}-k} + \phi(d)^{-1} \binom{q-6}{k} \right) \\
 &\quad \times \left( \frac{q-6}{2} \right)^2 \left( \frac{q-6}{2} \right)^2 d^{\frac{(q-1)(3q-21)}{8}} \\
 &\equiv d^{\frac{(q-1)(3q-21)}{8}} \phi(d)^{\frac{q-9}{4}} \text{ or } 0 \\
 &\equiv d^{\frac{q-1}{4}} \text{ or } 0 \pmod{\mathbb{F}_q^{\times 2}}.
 \end{aligned}$$

Hence there is an element  $x_q(d) \in \mathbb{F}_q$  such that

$$\tilde{T}_{(q-11)/2}(d, q) = d^{\frac{q-1}{4}} x_q(d)^2.$$

**Case 2.**  $q \equiv 3 \pmod{4}$ . We can verify that

$$\begin{aligned}
 \tilde{T}_{(q-11)/2}(d, q) &\equiv \text{sgn}(\text{inv}_q) \prod_{k=0}^{\frac{q-11}{2}} \left( \binom{q-6}{k} + \phi(d)^{-1} \binom{q-6}{\frac{q-11}{2}-k} \right) \left( \frac{q-6}{2} \right)^2 \left( \frac{q-6}{2} \right)^2 d^{\frac{(q-1)(3q-21)}{8}} \\
 &\equiv \text{sgn}(\text{inv}_q) \prod_{k=0}^{\frac{q-15}{4}} \left( \binom{q-6}{k} + \phi(d)^{-1} \binom{q-6}{\frac{q-11}{2}-k} \right) \left( \binom{q-6}{\frac{q-11}{2}-k} + \phi(d)^{-1} \binom{q-6}{k} \right) \\
 &\quad \times \left( \frac{q-6}{2} \right)^2 \left( \frac{q-6}{2} \right)^2 \left( \binom{q-6}{\frac{q-11}{4}} + \phi(d)^{-1} \binom{q-6}{\frac{q-11}{4}} \right) d^{\frac{(q-1)(3q-21)}{8}} \pmod{\mathbb{F}_q^{\times 2}}.
 \end{aligned}$$

If  $\phi(d) = -1$  or  $\left(\frac{q-6}{\frac{q-11}{4}}\right) \equiv 0 \pmod{q}$ ,  $\tilde{T}_{(q-11)/2}(d, q) \equiv 0 \pmod{q}$ . Otherwise,

$$\begin{aligned}
 &\left( \frac{q-6}{\frac{q-11}{4}} \right) + \phi(d)^{-1} \left( \frac{q-6}{\frac{q-11}{4}} \right) \\
 &= 2 \frac{(q-6)!}{\left(\frac{q-11}{4}\right)! \left(\frac{3q-13}{4}\right)!} \\
 &= 2 \frac{(q-6)! \times \frac{3q-9}{4} \frac{3q-5}{4} \dots \frac{3q+7}{4}}{\left(\left(\frac{q-11}{4}\right)!\right)^2 \times \frac{q-7}{4} \frac{q-3}{4} \dots \frac{3q+7}{4}} \\
 &\equiv \frac{-7(q-1)!}{(-1)^{\frac{q+9}{4}} \left(\frac{q-7}{4} \frac{q-3}{4} \dots \frac{q-1}{2}\right)^2}
 \end{aligned}$$

$$\equiv 7(-1)^{\frac{q+5}{4}}(q-1)! \pmod{\mathbb{F}_q^{\times 2}}.$$

In this case, by Lemma 2.3 and  $\phi(d) = 1$  we obtain that

$$\begin{aligned} \tilde{T}_{(q-11)/2}(d, q) &\equiv 7(-1)^{\frac{q+5}{4}}(q-1)!(-1)^{\frac{(q-3)(q-5)}{8}}d^{\frac{3q+3}{4}} \text{ or } 0 \\ &\equiv 7(q-1)!d^{\frac{q+1}{4}} \text{ or } 0 \pmod{\mathbb{F}_q^{\times 2}}. \end{aligned}$$

In view of the above, there is an element  $y_q(d) \in \mathbb{F}_q$  such that

$$\tilde{T}_{(q-11)/2}(d, q) = 7(q-1)!d^{\frac{q+1}{4}}y_q(d)^2.$$

□

Applying this result, we can obtain Corollary 1.2 easily.

**Proof of Corollary 1.2.** If  $p \equiv 1 \pmod{4}$  and  $\left(\frac{d}{p}\right) = 1$ , from the proof above, we obtain

$$\left(\frac{T_{(p-11)/2}(d, p)}{p}\right) = 1 \text{ or } 0.$$

In other words,  $\left(\frac{T_{(p-11)/2}(d, p)}{p}\right) \neq -1$ .

If  $p \equiv 3 \pmod{4}$  and  $\left(\frac{d}{p}\right) = 1$ , we obtain that  $\left(\frac{T_{(p-11)/2}(d, p)}{p}\right) = \left(\frac{-7}{p}\right)$  or 0. We get that if

$$\left(\frac{T_{(p-11)/2}(d, p)}{p}\right) = -1,$$

then  $p \equiv 1$  or  $2$  or  $4 \pmod{7}$  immediately. □

#### 4. PROOF OF THEOREM 1.3

**Proof** Let  $a = \left(\frac{p-1}{2}\right)!$ . For  $p \equiv 1 \pmod{4}$  we have  $a^2 = \left(\left(\frac{p-1}{2}\right)!\right)^2 \equiv -1 \pmod{p}$ . For each  $k = 1, \dots, (p-1)/2$ , let  $\pi_a^*(k)$  be the unique  $r \in \{1, \dots, (p-1)/2\}$  with  $ak$  congruent to  $r$  or  $-r$  modulo  $p$ . For the permutation  $\pi_a^*$  on  $\{1, \dots, (p-1)/2\}$ , Huang and Pan [4] showed that its sign is given by

$$\text{sgn}(\pi_a^*) = \left(\frac{a}{p}\right)^{(p+1)/2}.$$

If we view  $D_p^{(m)}$  as a determinant over  $\mathbb{F}_p$ , then we have

$$D_p^{(m)} = \det \left[ (i^2 + (aj)^2)^{m + \frac{p-1}{2}} \right]_{1 \leq i, j \leq (p-1)/2} = \text{sgn}(\pi_a^*) \det \left[ (a_i + a_j)^{m + \frac{p-1}{2}} \right]_{1 \leq i, j \leq (p-1)/2}.$$

Thus

$$D_p^{(m)} \equiv \text{sgn}(\pi_a^*) \det \left[ \left(\frac{a_i}{a_j} + 1\right)^{m + \frac{p-1}{2}} \right]_{1 \leq i, j \leq (p-1)/2} \pmod{\mathbb{F}_p^{\times 2}}.$$

Fix positive odd integer  $m$ . For integer  $k$  with  $0 \leq k \leq (m-1)/2$ , we define

$$F_m(k) = 2^{m-2k}(m-k)(m-k-1)\dots(k+1) + (2m-2k-1)(2m-2k-3)\dots(2k+1).$$

Let

$$M = \max_{0 \leq k \leq (m-1)/2} F_m(k).$$

We claim that  $p \nmid D_p^{(m)}$  when  $p > M$ . In other words, if  $p > M$ , then we have  $p \notin E(m)$ . Hence, the elements of  $E(m)$  is finite. Now, we prove the claim.

Noting that  $p > M \geq F_m(0) = 2^m m! + (2m-1)!! > 2m$  and  $\left(\frac{a_i}{a_j}\right)^{\frac{p-1}{2}} = 1$ , we can verify that

$$\begin{aligned} & \left(\frac{a_i}{a_j} + 1\right)^{m+\frac{p-1}{2}} \\ &= \sum_{k=0}^{m+\frac{p-1}{2}} \binom{m+\frac{p-1}{2}}{k} \left(\frac{a_i}{a_j}\right)^k \\ &= \sum_{k=0}^m \left( \binom{m+\frac{p-1}{2}}{k} + \binom{m+\frac{p-1}{2}}{\frac{p-1}{2}+k} \right) \left(\frac{a_i}{a_j}\right)^k + \sum_{k=m+1}^{\frac{p-3}{2}} \binom{m+\frac{p-1}{2}}{k} \left(\frac{a_i}{a_j}\right)^k. \end{aligned}$$

Hence

$$\left(\frac{a_i}{a_j} + 1\right)^{m+\frac{p-1}{2}} = f\left(\frac{a_i}{a_j}\right)$$

where

$$f(T) = \sum_{k=0}^m \left( \binom{m+\frac{p-1}{2}}{k} + \binom{m+\frac{p-1}{2}}{\frac{p-1}{2}+k} \right) T^k + \sum_{k=m+1}^{\frac{p-3}{2}} \binom{m+\frac{p-1}{2}}{k} T^k.$$

Let  $C_f$  denote the product of coefficients of  $f(T)$ , then

$$C_f = \prod_{k=0}^m \left( \binom{\frac{p-1}{2}+m}{k} + \binom{\frac{p-1}{2}+m}{m-k} \right) \prod_{k=m+1}^{\frac{p-3}{2}} \binom{\frac{p-1}{2}+m}{k}$$

Applying Lemma 2.1 to  $P(T) = f(T)$ , we have

$$\begin{aligned} D_p^{(m)} &\equiv \text{sgn}(\pi_a^*) \text{sgn}(\text{inv}_p) \prod_{k=0}^m \left( \binom{\frac{p-1}{2}+m}{k} + \binom{\frac{p-1}{2}+m}{m-k} \right) \prod_{k=m+1}^{\frac{p-3}{2}} \binom{\frac{p-1}{2}+m}{k} \prod_{1 \leq i < j \leq n} (a_j - a_i)^2 \\ &\equiv \text{sgn}(\pi_a^*) \text{sgn}(\text{inv}_p) \prod_{k=0}^m \left( \binom{\frac{p-1}{2}+m}{k} + \binom{\frac{p-1}{2}+m}{m-k} \right) \prod_{k=m+1}^{\frac{p-3}{2}} \binom{\frac{p-1}{2}+m}{k} \pmod{\mathbb{F}_p^{\times 2}}. \end{aligned}$$

Since  $p \nmid \prod_{m+1}^{\frac{p-3}{2}} \binom{\frac{p-1}{2}+m}{k}$  for  $p > 2m$ , we can deduce that if  $p \mid D_p^{(m)}$ , then

$$p \mid \prod_{k=0}^m \left( \binom{\frac{p-1}{2}+m}{k} + \binom{\frac{p-1}{2}+m}{m-k} \right).$$

For  $0 \leq k \leq m$ ,

$$\binom{\frac{p-1}{2}+m}{k} + \binom{\frac{p-1}{2}+m}{m-k} = \frac{(\frac{p-1}{2}+m)! ((m-k)! (\frac{p-1}{2}+k)! + k! (\frac{p-1}{2}+m-k)!)}{(m-k)! (\frac{p-1}{2}+k)! k! (\frac{p-1}{2}+m-k)!}.$$

Hence, if  $p \mid \binom{\frac{p-1}{2}+m}{k} + \binom{\frac{p-1}{2}+m}{m-k}$ , we have

$$(m-k)! \left( \frac{p-1}{2} + k \right)! + k! \left( \frac{p-1}{2} + m - k \right)! \equiv 0 \pmod{p}.$$

Without loss of generality, we assume  $k \leq (m-1)/2$  since  $m$  is an odd positive integer. Then we can deduce that

$$p \mid (m-k)(m-k-1) \dots (k+1) + \left( \frac{p-1}{2} + m - k \right) \left( \frac{p-1}{2} + m - k - 1 \right) \dots \left( \frac{p-1}{2} + k + 1 \right).$$

It means that

$$2^{m-2k} (m-k)(m-k-1) \dots (k+1) + (2m-2k-1)(2m-2k-3) \dots (2k+1) = F_m(k) \equiv 0 \pmod{p},$$

which is in contradiction to  $p > M$ . Therefore,  $p \nmid \binom{\frac{p-1}{2}+m}{k} + \binom{\frac{p-1}{2}+m}{m-k}$ . Hence the claim is true.

We can also compute  $E(m)$  for some specific  $m$ . For example, let  $m = 5$ . Firstly, we check all the prime number  $p$  with  $p \leq 2m + 3 = 13$ . We find that there is no prime  $p \in E(5)$ . For prime number  $p > 13$ , we get that if  $p \mid D_p^{(5)}$ , then  $p \mid \binom{\frac{p-1}{2}+5}{k} + \binom{\frac{p-1}{2}+5}{5-k}$  for some  $k$  from the illustration above. Moreover, we have  $p \equiv 1 \pmod{4}$  and  $p \mid F_5(k)$  for some  $k$ . We calculate  $F_5(0)$ ,  $F_5(1)$  and  $F_5(2)$  and only find that  $29 \mid F_5(0) = 4785$ . Therefore,  $E(5) = \{29\}$ .

In the same way, we obtain that

$$E(7) = \{13, 53\}, \quad E(9) = \{13, 17, 29\}$$

$$E(11) = \{17, 29\} \text{ and } E(13) = \{17, 109, 401\}.$$

□

## 5. PROOFS OF THEOREM 1.4 AND 1.5

**Proof of Theorem 1.4.** By Cayley's Theorem we have  $D_p^{(1)} = \text{pf}(D_p^{(1)})^2$ . For  $p \equiv 1 \pmod{4}$  we have  $((\frac{p-1}{2})!)^2 \equiv -1 \pmod{p}$ . Let  $a = (\frac{p-1}{2})!$  and define  $\pi_a^*(k)$  the same as before. By Lemma [9, Lemma 2.3], we have

$$\text{sgn}(\pi_a^*) = \left( \frac{a}{p} \right)^{(p+1)/2} = \left( \frac{2}{p} \right)^{(p+1)/2} = (-1)^{(p-1)/4}.$$

If we view  $D_p^{(1)}$  as a determinant over  $\mathbb{F}_p$ , we have

$$\begin{aligned}
 D_p^{(1)} &= \det \left[ (i^2 + (aj)^2)^{\frac{p+1}{2}} \right]_{1 \leq i, j \leq (p-1)/2} \\
 &= \text{sgn}(\pi_a^*) \det \left[ (a_i + a_j)^{\frac{p+1}{2}} \right]_{1 \leq i, j \leq (p-1)/2} \\
 &= (-1)^{(p-1)/4} \prod_{i=1}^{(p-1)/2} a_i \det \left[ \left( \frac{a_i}{a_j} + 1 \right)^{\frac{p+1}{2}} \right]_{1 \leq i, j \leq (p-1)/2} \\
 &= (-1)^{(p+3)/4} \det \left[ \left( \frac{a_i}{a_j} + 1 \right)^{\frac{p+1}{2}} \right]_{1 \leq i, j \leq (p-1)/2}.
 \end{aligned}$$

Noting that  $\left( \frac{a_i}{a_j} \right)^{\frac{p-1}{2}} = 1$ , we can verify that

$$\left( \frac{a_i}{a_j} + 1 \right)^{\frac{p+1}{2}} = \sum_{k=2}^{\frac{p-3}{2}} \binom{\frac{p+1}{2}}{k} \left( \frac{a_i}{a_j} \right)^k + \frac{p+3}{2} \left( \frac{a_i}{a_j} \right) + \frac{p+3}{2}.$$

Hence for

$$f(T) = \sum_{k=2}^{\frac{p-3}{2}} \binom{\frac{p+1}{2}}{k} T^k + \frac{p+3}{2} T + \frac{p+3}{2}$$

we have

$$\left( \frac{a_i}{a_j} + 1 \right)^{\frac{p+1}{2}} = f \left( \frac{a_i}{a_j} \right).$$

By Lemma 2.1 and Lemma 2.3, we obtain

$$\begin{aligned}
 D_p^{(1)} &= (-1)^{(p+3)/4} (-1)^{\frac{(p-3)(p-5)}{8}} \left( \frac{p+3}{2} \right)^2 \prod_{k=2}^{\frac{p-3}{2}} \binom{\frac{p+1}{2}}{k} \prod_{1 \leq i < j \leq \frac{p-1}{2}} (i^2 - j^2)^2 \\
 &= \left( \frac{p+3}{2} \right)^2 \prod_{k=2}^{\frac{p-1}{4}} \binom{\frac{p+1}{2}}{k}^2 \prod_{1 \leq i < j \leq \frac{p-1}{2}} (i^2 - j^2)^2.
 \end{aligned}$$

Let

$$t = \frac{p+3}{2} \prod_{k=2}^{\frac{p-1}{4}} \binom{\frac{p+1}{2}}{k} \prod_{1 \leq i < j \leq \frac{p-1}{2}} (j^2 - i^2).$$

We have

$$D_p^{(1)} = \text{pf}(D_p^{(1)})^2 \equiv t^2 \pmod{p}.$$

Therefore  $\sqrt{D_p^{(1)}} = \pm t \pmod{p}$ . For  $p \equiv 1 \pmod{4}$  and  $\left(\frac{-1}{p}\right) = 1$ , we obtain that

$$\left(\frac{\sqrt{D_p^{(1)}}}{p}\right) = \left(\frac{t}{p}\right).$$

Referring to [10], we have

$$\prod_{1 \leq i < j \leq \frac{p-1}{2}} (j^2 - i^2) = -\left(\frac{p-1}{2}\right)! \pmod{p}.$$

Observe that

$$\prod_{k=2}^{\frac{p-1}{4}} \binom{\frac{p+1}{2}}{k} = \left(\left(\frac{p+1}{2}\right)!\right)^{\frac{p-5}{4}} \prod_{k=2}^{\frac{p-3}{2}} k!.$$

By observation, we have

$$\prod_{k=2}^{\frac{p-3}{2}} k! \equiv \frac{p-3}{2} \left(\frac{p-5}{2}\right)^2 \left(\frac{p-7}{2}\right)^3 \dots 3^{\frac{p-3}{2}} 2^{\frac{p-5}{2}} \equiv \frac{p-3}{2} \frac{p-7}{2} \dots 3 \equiv \left(\frac{p-3}{2}\right)!! \pmod{\mathbb{F}_p^{\times 2}}.$$

We also have

$$\begin{aligned} \left(\frac{p-3}{2}\right)!! &\equiv \left(\frac{p-3}{2}\right)!! \left(\left(\frac{p-1}{2}\right)!!\right)^2 \\ &\equiv \left(\frac{p-1}{2}\right)! \left(\frac{p-1}{4}\right)! 2^{(p-1)/4} \pmod{\mathbb{F}_p^{\times 2}}. \end{aligned}$$

We conclude that

$$\begin{aligned} t &\equiv -\frac{3}{2} \left(\left(\frac{p+1}{2}\right)!\right)^{\frac{p-5}{4}} \left(\frac{p-1}{2}\right)! \left(\frac{p-1}{4}\right)! 2^{(p-1)/4} \left(\frac{p-1}{2}\right)! \\ &\equiv -3 \left(\left(\frac{p+1}{2}\right)!\right)^{\frac{p-5}{4}} \left(\frac{p-1}{4}\right)! 2^{(p-5)/4} \pmod{\mathbb{F}_p^{\times 2}}. \end{aligned}$$

For  $p \equiv 1 \pmod{4}$ , by Lemma [9, Lemma 2.3] we have

$$\left(\frac{\left(\frac{p+1}{2}\right)!}{p}\right) = \left(\frac{2^{-1}}{p}\right) \left(\frac{\left(\frac{p-1}{2}\right)!}{p}\right) = \left(\frac{2^{-1}}{p}\right) \left(\frac{2}{p}\right) = 1.$$



Notice that  $\left(\frac{2^{(p-5)/4}}{p}\right) = 1$  and  $\left(\frac{-1}{p}\right) = 1$ , we obtain

$$\left(\frac{\sqrt{D_p^{(1)}}}{p}\right) = \left(\frac{t}{p}\right) = \left(\frac{3}{p}\right) \left(\frac{\left(\frac{p-1}{4}\right)!}{p}\right) = (-1)^{|\{0 < k < \frac{p}{4} : \binom{k}{p} = -1\}|} \left(\frac{p}{3}\right). \quad \square$$

**Proof of Theorem 1.5.** For  $p = 5$ , we check that Theorem 1.5 is right. Now, we consider  $p \geq 13$  with  $p \equiv 1 \pmod{4}$ . By Cayley's Theorem we have  $D_p^{(3)} = \text{pf}(D_p^{(3)})^2$ .

For  $p \equiv 1 \pmod{4}$  we have  $\left(\left(\frac{p-1}{2}\right)!\right)^2 \equiv -1 \pmod{p}$ . Let  $a = \left(\frac{p-1}{2}\right)!$  and define  $\pi_a^*(k)$  the same as before. By Lemma [9, Lemma 2.3], we have

$$\text{sgn}(\pi_a^*) = \left(\frac{a}{p}\right)^{(p+1)/2} = \left(\frac{2}{p}\right)^{(p+1)/2} = (-1)^{(p-1)/4}.$$

If we view  $D_p^{(3)}$  as a determinant over  $\mathbb{F}_p$ , then we have

$$\begin{aligned} D_p^{(1)} &= \det \left[ (i^2 + (aj)^2)^{\frac{p+5}{2}} \right]_{1 \leq i, j \leq (p-1)/2} \\ &= \text{sgn}(\pi_a^*) \det \left[ (a_i + a_j)^{\frac{p+5}{2}} \right]_{1 \leq i, j \leq (p-1)/2} \\ &= (-1)^{(p-1)/4} \prod_{i=1}^{(p-1)/2} a_i \det \left[ \left( \frac{a_i}{a_j} + 1 \right)^{\frac{p+5}{2}} \right]_{1 \leq i, j \leq (p-1)/2} \\ &= (-1)^{(p+3)/4} \det \left[ \left( \frac{a_i}{a_j} + 1 \right)^{\frac{p+5}{2}} \right]_{1 \leq i, j \leq (p-1)/2}. \end{aligned}$$

Noting that  $\left(\frac{a_i}{a_j}\right)^{\frac{p-1}{2}} = 1$ , we can verify that

$$\begin{aligned} \left(\frac{a_i}{a_j} + 1\right)^{\frac{p+5}{2}} &= \sum_{k=4}^{\frac{p-3}{2}} \binom{\frac{p+5}{2}}{k} \left(\frac{a_i}{a_j}\right)^k + 1 + \binom{\frac{p+5}{2}}{1} \left(\frac{a_i}{a_j}\right) + \binom{\frac{p+5}{2}}{2} \left(\frac{a_i}{a_j}\right)^2 + \binom{\frac{p+5}{2}}{3} \left(\frac{a_i}{a_j}\right)^3 \\ &\quad + \binom{\frac{p+5}{2}}{\frac{p-1}{2}} \left(\frac{a_i}{a_j}\right)^{\frac{p-1}{2}} + \binom{\frac{p+5}{2}}{\frac{p+1}{2}} \left(\frac{a_i}{a_j}\right)^{\frac{p+1}{2}} + \binom{\frac{p+5}{2}}{\frac{p+3}{2}} \left(\frac{a_i}{a_j}\right)^{\frac{p+3}{2}} + \binom{\frac{p+5}{2}}{\frac{p+5}{2}} \left(\frac{a_i}{a_j}\right)^{\frac{p+5}{2}} \\ &= \sum_{k=4}^{\frac{p-3}{2}} \binom{\frac{p+5}{2}}{k} \left(\frac{a_i}{a_j}\right)^k + 1 + \binom{\frac{p+5}{2}}{\frac{p-1}{2}} + \left( \binom{\frac{p+5}{2}}{1} + \binom{\frac{p+5}{2}}{\frac{p+1}{2}} \right) \left(\frac{a_i}{a_j}\right) \\ &\quad + \left( \binom{\frac{p+5}{2}}{2} + \binom{\frac{p+5}{2}}{\frac{p+3}{2}} \right) \left(\frac{a_i}{a_j}\right)^2 + \left( \binom{\frac{p+5}{2}}{3} + \binom{\frac{p+5}{2}}{\frac{p+5}{2}} \right) \left(\frac{a_i}{a_j}\right)^3 \end{aligned}$$

$$= \sum_{k=4}^{\frac{p-3}{2}} \binom{\frac{p+5}{2}}{k} \left(\frac{a_i}{a_j}\right)^k + \frac{21}{16} + \frac{35}{8} \left(\frac{a_i}{a_j}\right) + \frac{35}{8} \left(\frac{a_i}{a_j}\right)^2 + \frac{21}{16} \left(\frac{a_i}{a_j}\right)^3.$$

Hence

$$\left(\frac{a_i}{a_j} + 1\right)^{\frac{p+1}{2}} = f\left(\frac{a_i}{a_j}\right)$$

where

$$f(T) = \sum_{k=4}^{\frac{p-3}{2}} \binom{\frac{p+5}{2}}{k} T^k + \frac{21}{16} + \frac{35}{8} T + \frac{35}{8} T^2 + \frac{21}{16} T^3.$$

By Lemma 2.1 and Lemma 2.3, we obtain

$$\begin{aligned} D_p^{(3)} &= (-1)^{(p+3)/4} (-1)^{\frac{(p-3)(p-5)}{8}} \left(\frac{21}{16}\right)^2 \left(\frac{35}{8}\right)^2 \prod_{k=4}^{\frac{p-3}{2}} \binom{\frac{p+5}{2}}{k} \prod_{1 \leq i < j \leq \frac{p-1}{2}} (i^2 - j^2)^2 \\ &= \left(\frac{21}{16}\right)^2 \left(\frac{35}{8}\right)^2 \prod_{k=4}^{\frac{p+3}{4}} \binom{\frac{p+5}{2}}{k}^2 \prod_{1 \leq i < j \leq \frac{p-1}{2}} (i^2 - j^2)^2. \end{aligned}$$

Let

$$s = \frac{21}{16} \times \frac{35}{8} \prod_{k=4}^{\frac{p+3}{4}} \binom{\frac{p+5}{2}}{k} \prod_{1 \leq i < j \leq \frac{p-1}{2}} (j^2 - i^2).$$

We have

$$D_p^{(3)} = \text{pf}(D_p^{(3)})^2 \equiv s^2 \pmod{p}.$$

Therefore  $\sqrt{D_p^{(3)}} = \pm s \pmod{p}$ . For  $p \equiv 1 \pmod{4}$  and  $\left(\frac{-1}{p}\right) = 1$ , we obtain that

$$\left(\frac{\sqrt{D_p^{(3)}}}{p}\right) = \left(\frac{s}{p}\right).$$

Referring to [10], we have

$$\prod_{1 \leq i < j \leq \frac{p-1}{2}} (j^2 - i^2) = -\left(\frac{p-1}{2}\right)! \pmod{p}.$$

Observe that

$$\prod_{k=4}^{\frac{p+3}{4}} \binom{\frac{p+5}{2}}{k} = \left(\left(\frac{p+5}{2}\right)!\right)^{\frac{p-9}{4}} \prod_{k=4}^{\frac{p-3}{2}} k!.$$

We notice that  $\frac{p-3}{2}$  appears once,  $\frac{p-5}{2}$  appears twice ... and 4 appears  $\frac{p-9}{2}$  times in  $\prod_{k=2}^{\frac{p-3}{2}} k!$ . Hence,

$$\prod_{k=2}^{\frac{p-3}{2}} k! \equiv \frac{p-3}{2} \frac{p-7}{2} \cdots 5 \equiv \frac{1}{3} \left(\frac{p-3}{2}\right)!! \pmod{\mathbb{F}_p^{\times 2}}.$$

We also have

$$\left(\frac{p-3}{2}\right)!! \equiv \left(\frac{p-1}{2}\right)! \left(\frac{p-1}{4}\right)! 2^{(p-1)/4} \pmod{\mathbb{F}_p^{\times 2}}.$$

We conclude that

$$\begin{aligned} s &\equiv -\frac{21}{16} \times \frac{35}{8} \left(\left(\frac{p+5}{2}\right)!\right)^{\frac{p-9}{4}} \frac{1}{3} \left(\frac{p-1}{2}\right)! \left(\frac{p-1}{4}\right)! 2^{(p-1)/4} \left(\frac{p-1}{2}\right)! \\ &\equiv -5 \left(\left(\frac{p+5}{2}\right)!\right)^{\frac{p-5}{4}} \left(\frac{p-1}{4}\right)! 2^{(p-5)/4} \pmod{\mathbb{F}_p^{\times 2}}. \end{aligned}$$

For  $p \equiv 1 \pmod{4}$ , by Lemma [9, Lemma 2.3] we have

$$\left(\frac{\left(\frac{p+5}{2}\right)!}{p}\right) = \left(\frac{\frac{15}{4}}{p}\right) \left(\frac{\left(\frac{p+1}{2}\right)!}{p}\right) = \left(\frac{3}{p}\right) \left(\frac{5}{p}\right).$$

Notice that  $\left(\frac{2^{(p-5)/4}}{p}\right) = 1$  and  $\left(\frac{-1}{p}\right) = 1$ , we obtain

$$\begin{aligned} \left(\frac{\sqrt{D_p^{(3)}}}{p}\right) &= \left(\frac{s}{p}\right) \\ &= \left(\frac{3}{p}\right)^{(p-9)/4} \left(\frac{5}{p}\right)^{(p-5)/4} \left(\frac{\left(\frac{p-1}{4}\right)!}{p}\right) \\ &= (-1)^{|\{0 < k < \frac{p}{4} : \binom{k}{p} = -1\}|} \left(\frac{p}{4 + (-1)^{(p-1)/4}}\right). \end{aligned}$$

□

**Acknowledgment.** The authors would like to thank Zhi-Wei Sun for the conjectures he provided in [12].

#### REFERENCES

- [1] L. Carlitz, *Some cyclotomic matrices*, Acta Arith., **5** (1959), 293–308.
- [2] R. Chapman, *Determinants of Legendre symbol matrices*, Acta Arith. **115** (2004), 231–244.
- [3] D. Grinberg, Z.-W. Sun and L. Zhao, *Proof of three conjectures on determinants related to quadratic residues*, Linear Multilinear Algebra **70** (2022), 3734–3746

- [4] C. Huang and H. Pan, *A remark on Zolotarev's theorem*, Colloq.Math., **171**, No.1 (2023), 159–166.
- [5] C. Krattenthaler, *Advanced determinant calculus: a complement*, Linear Algebra Appl., **411** (2005), 68–166
- [6] M. Lerch, *Sur un théorème arithmétique de Zolotarev*, Bull. Intern. de l'Acad. Francois Joseph **3** (1896), 34–37
- [7] Chen-kai Ren and Zhi-Wei Sun, *On some determinants arising from quadratic residues* arXiv: 2404.11547
- [8] R. P. Stanley, *Enumerative Combinatorics*, Vol. 1, Cambridge Univ. Press, Cambridge, 19
- [9] Z.-W. Sun, *On some determinants with Legendre symbol entries*, Finite Fields Appl., **56** (2019), 285–307.
- [10] Z.-W. Sun, *Quadratic residues and related permutations and identities*, Finite Fields Appl., **59** (2019), 246–283
- [11] Z.-W. Sun, *On some determinants and permanents*, Acta Math. Sinica Chin. Ser., **66** (2023), in press. See also arXiv:2207.13039.
- [12] Z.-W. Sun, *Some determinants involving quadratic residues module primes* arXiv:2401.14301v1
- [13] M. Vsemirnov, *On the evaluation of R. Chapman's "evil determinant"*, Linear Algebra Appl., **436** (2012), 4101–4106.
- [14] M. Vsemirnov, *On R. Chapman's "evil determinant": case  $p \equiv 1 \pmod{4}$* , Acta Arith., **159** (2013), 331–344.
- [15] H.-L. Wu, Y.-F. She and L.-Y. Wang, *Cyclotomic matrices and hypergeometric functions over finite fields*, Finite Fields Appl., **82** (2022), Article ID 102054

(CHEN-KAI REN) DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY, NANJING 210093, PEOPLE'S REPUBLIC OF CHINA

*Email address:* ckRen@smail.nju.edu.cn

(XIN-QI LUO ) DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY, NANJING 210093, PEOPLE'S REPUBLIC OF CHINA

*Email address:* lxq15995710087@163.com