# Determination Problems for Orbit Closures and Matrix Groups

Rida Ait El Manssour<sup>\*1</sup>, George Kenison<sup>†2</sup>, Mahsa Shirmohammadi<sup>‡1</sup>, and James Worrell<sup>§3</sup>

<sup>1</sup>IRIF, CNRS, Université Paris Cité, Paris, France <sup>2</sup>Liverpool John Moores University, Liverpool, UK <sup>3</sup>University of Oxford, Oxford, UK

#### Abstract

Computational problems concerning the orbit of a point under the action of a matrix group occur in numerous subfields of computer science, including complexity theory, program analysis, quantum computation, and automata theory. In many cases the focus extends beyond orbits proper to orbit closures under a suitable topology. Typically one starts from a group and several points and asks questions about the orbit closure of the points under the action of the group, e.g., whether two given orbit closures intersect.

In this paper we consider a collection of what we call determination problems concerning groups and orbit closures. These problems begin with a given variety and seek to understand whether and how it arises either as an algebraic group or as an orbit closure. The *how* question asks whether the underlying group is *s*-generated, meaning it is topologically generated by *s* matrices for a given number *s*. Among other applications, problems of this type have recently been studied in the context of synthesising loops subject to certain specified invariants on program variables.

Our main result is a polynomial-space procedure that inputs a variety V and a number s and determines whether V arises as an orbit closure of a point under an s-generated commutative matrix group. The main tools in our approach are rooted in structural properties of commutative algebraic matrix groups and lattice theory. We leave open the question of determining whether a variety is an orbit closure of a point under an algebraic matrix group (without the requirement of commutativity). In this regard, we note that a recent paper [Nos+22] gives an elementary procedure to compute the orbit closure of a point under finitely many matrices.

<sup>\*</sup>manssour@irif.fr

<sup>&</sup>lt;sup>†</sup>g.j.kenison@ljmu.ac.uk

<sup>&</sup>lt;sup>‡</sup>mahsa@irif.fr

<sup>§</sup>jbw@cs.ox.ac.uk

## 1 Introduction

**Orbit Problems.** The computational study of orbits of matrix groups stretches back many decades. One of the most fundamental problems in this area is determining, for a given field  $\mathbb{F}$ , whether a given pair of vectors  $\boldsymbol{u}, \boldsymbol{v} \in \mathbb{F}^d$  lie in the same orbit under the action of a finitely generated subgroup G of the general linear group  $\operatorname{GL}_d(\mathbb{F})$ ; that is, whether  $\boldsymbol{u} \in G \cdot \boldsymbol{v}$ . For cyclic matrix groups over  $\mathbb{Q}$ , this problem reduces to the Kannan–Lipton orbit problem. For such groups, there exists a polynomial-time algorithm for deciding whether a given pair of vectors lie in the same orbit [KL80; KL86]. The extension of the Kannan–Lipton orbit problem to a group with a finite generating set  $\{M_1, \ldots, M_s\}$  over  $\mathbb{Q}$  (and number fields) is known to be decidable when the generating set comprises commuting matrices. However, this problem becomes undecidable in the general case [Bab+96]. The exploration of this problem over finite fields was instrumental in leading Babai to introduce the concept of interactive proofs [Bab85].

**Orbit Closures.** For many applications, including in program analysis and geometric complexity theory, it makes sense to study *orbit closures* in lieu of orbits proper [Bür+11; Bür24; DJK05; Hru+23]. Given a field  $\mathbb{F}$  and a group  $G \in \operatorname{GL}_d(\mathbb{F})$  acting on  $\mathbb{F}^d$ , the orbit closure of  $\boldsymbol{v} \in \mathbb{F}^d$ , denoted by  $\overline{G \cdot \boldsymbol{v}}$ , is the closure of the orbit  $G \cdot \boldsymbol{v}$  in the Zariski topology. We note that over the field  $\mathbb{C}$ , if the group in question is Zariski closed then the Zariski closure of an orbit coincides with its closure in the Euclidean topology.

The orbit-closure containment problem asks, given vectors  $\boldsymbol{u}, \boldsymbol{v} \in \mathbb{F}^d$  and a group  $G \subseteq \operatorname{GL}_d(\mathbb{F})$ , to determine whether  $\boldsymbol{u}$  lies in  $\overline{G \cdot \boldsymbol{v}}$ , whereas the orbit-closure intersection problem asks whether  $\overline{G \cdot \boldsymbol{v}} \cap \overline{G \cdot \boldsymbol{u}} \neq \emptyset$ . Orbit closures feature in complexity theory, program analysis, quantum computation and automata theory. A striking application is geometric complexity theory, in which many recent studies centred around the formulation of the  $\mathbf{VP} = \mathbf{VNP}$  problem in terms of orbit-closure containment with respect to the action of  $\operatorname{GL}_d(\overline{\mathbb{Q}})$  on polynomial rings [Bür+11; Bür24].

In certain applications, such as non-convex optimisation problems, non-commutative rational identity testing, and graph isomorphism [Blä+21; Bür+21; DM20; FS13], one considers the orbit closure of a point  $\boldsymbol{v}$  under a linear algebraic group G that is explicitly presented via a given set of equations (that is, G is the zero set of a given polynomial ideal  $\langle P_1, \ldots, P_k \rangle$ ). In other applications, such as quantum computing and program analysis [DJK05; Hru+23], one wishes to compute the orbit closure of a group G that is implicitly presented via a finite set of topological generators (that is,  $G = \langle M_1, \ldots, M_s \rangle$ ). We refer to these settings as *explicit* and *implicit* presentations of the orbit-closure problems, respectively.

Given a polynomial ideal  $\langle P_1, \ldots, P_s \rangle \subseteq \mathbb{K}[x]$  defining an algebraic matrix group  $G \in \mathrm{GL}_d(\mathbb{F})$ with  $\mathbb{K}$  an effective subfield of  $\mathbb{F}$ , as highlighted in [Blä+21], orbit-closure- containment and intersection can easily be checked in existential fragment of the first order theory of the field  $\mathbb{F}$ . By Koiran's seminal results on Hilbert's Nullstellensatz over  $\overline{\mathbb{Q}}$  [Koi96], the explicit orbit-closure problems over  $\overline{\mathbb{Q}}$  are both in **AM** assuming the generalized Riemann hypothesis (GRH). Furthermore, answering a question posed by Wigderson, it is shown in [Blä+21] that the explicit orbit-closure containment problem is **NP**-hard, and shown in [Blä+21, Theorem 3.1.] that this problem over  $\mathbb{R}$  is polynomialtime equivalent to the existential theory over the reals (ETR)<sup>1</sup>. The primary application of the orbit problem in that work is the formulation of the slice rank of tensors in the union of orbit-closures. In [Bür+21], several other applications of the orbit-closure problems are identified in combinatorial optimisation and dynamical systems, specifically where the underlying group is assumed to be commutative. Notably the results in [Bür+21] fully resolve the problem for a subclass of commutative

<sup>&</sup>lt;sup>1</sup>In [Blä+21], the problem is simply called orbit-closure containment.

groups, namely tori. However, the complexity of explicit orbit-closure problems for commutative groups remains open.

In the implicit orbit-closure problems, the main challenge in computing an orbit closure lies in computing the Zariski closure of the group in question. Given a set  $\{M_1, \ldots, M_s\}$  of matrices over a number field K, the first algorithm to compute the polynomial ideal defining  $\langle M_1, \ldots, M_s \rangle$  was introduced in [DJK05]. The complexity bound of the algorithm therein is not known to be elementary, largely due to an iterative call to a subprocedure that takes the quotient of linear algebraic groups and the inherent difficulty of forming such quotients [Nos+22, Appendix C]. An elementary algorithm for implicit orbit-closure can be derived from a linearisation technique of [MS04a] together with a recent result in [Nos+22]. The key result in [Nos+22] is a quantitative structure lemma for algebraic matrix groups, providing an upper bound on the degree of the polynomials defining  $\langle M_1, \ldots, M_s \rangle$ . The above-mentioned complexity bounds for the implicit orbit-closure problem are in the order of seven fold exponential time when the generating matrices are over Q. It remains a challenging open problem to close the complexity gap, borrowing the lower bound from the explicit orbit-closure problems. Further algorithms to compute the Zariski closure of matrix groups and semigroups are presented in [Gra17; Hru+18; Hru+23].

Implicit orbit-closure problem in quantum computation, automata theory and program analysis have been the subject of extensive interest after the resolution of decade long open problems such as the equivalence problem for deterministic top-down tree-to-string transducers [SMK15], and the threshold problem for quantum automata [DJK05] (a natural version of the language-emptiness problem). Another application of closure problem in quantum setting is to test whether a finite set of quantum gates is universal [Nos+22]. Orbit closures feature in program analysis when one wants to automatically compute polynomial invariants of certain classes of loop programs [CK24; Hru+18; Kin+18; MS04b; SSM04]; the task of automatic invariant generation has been considered as the most important task in program verification [Bey+07].

#### 1.1 Determination Problems

In this paper, we investigate a series of determination problems related to groups and their orbit closures. These problems start with a given variety and examine whether it can be realized as an algebraic group or as an orbit closure, with the constraint that the underlying group is topologically s-generated. We define an algebraic group  $G \subseteq \operatorname{GL}_d(\overline{\mathbb{Q}})$  to be topologically s-generated if there is a set  $S \subseteq \operatorname{GL}_d(\overline{\mathbb{Q}})$  of matrices with cardinally s such that  $G = \overline{\langle S \rangle}$ . As noted in Proposition 5, algebraic groups are always topologically generated by a finite set.

In this context, determining whether a variety  $Z \subseteq \overline{\mathbb{Q}}^d$  arises as an orbit closure under the action of G is, in principle, straightforward. In such a case G is necessarily a subgroup of the group  $\operatorname{Sym}(Z) := \{A \in \operatorname{GL}_d(\overline{\mathbb{Q}}) : A(Z) = Z\}$  of matrices that fix Z, and so we may assume without loss of generality that G equals the latter group. But  $\operatorname{Sym}(Z)$  is definable in first-order logic over  $\overline{\mathbb{Q}}$  and hence the question of whether Z arises as the orbit closure of a point under  $\operatorname{Sym}(Z)$  reduces to the decision problem for this theory. It further holds by Proposition 5 that  $\operatorname{Sym}(Z)$  has a Zariski dense subgroup that is finitely generated (as a group). Whence Z is the orbit closure of a point under  $\operatorname{Sym}(Z)$ . In comparison, it is more challenging to determine whether a given variety is the orbit closure of a topologically *s*-generated group than simply determining whether it is an orbit closure *tout court*.

Our main determination problems are as follows. Let  $\boldsymbol{x} = \{x_{i,j}\}_{1 \leq i,j \leq d}$  be a tuple of variables. The **Group Determination Problem** asks, given  $s \in \mathbb{N}$  and a family of polynomials  $f_1, \ldots, f_m \in \mathbb{Q}[\boldsymbol{x}]$  of total degree at most b, to determine whether their zero locus  $Z \subseteq \operatorname{GL}_d(\overline{\mathbb{Q}})$  is an s-generated matrix group. The **Orbit-closure Determination Problem** asks, given  $s \in \mathbb{N}$  and a family of polynomials  $f_1, \ldots, f_m \in \mathbb{Q}[\mathbf{x}]$  of total degree at most b, determine whether their zero locus  $Z \subseteq \operatorname{GL}_d(\overline{\mathbb{Q}})$  is the orbit closure of some point  $\mathbf{v} \in \overline{\mathbb{Q}}^d$  under the action of an *s*-generated matrix group. In our complexity analysis we refer to the tuple (d, m, b) as the parameters of the problem instances, omitting the parameter s. This is without loss of generality as, by Proposition 7, Proposition 8 and Remark 10, the minimum number s of topological generators for the groups we study (commutative groups) is upper bounded by d.

This paper focuses on addressing the complexity of the determination problems for commutative groups. The extension of our results to the case of general matrix groups appears to be challenging. To approach the above version of the orbit determination problem, we rely on the observation that with respect to a convenient basis an orbit closure itself carries the structure of a matrix group. We then use basic structural results about semisimple and unipotent linear algebraic groups to identify when the above group is the closure of a commutative group.

**Orbit-Closure Determination Problem.** We reduce our determination problems to satisfiability problems of a fragment of the first-order theory of algebraically closed fields of characteristic zero. The following theorem gives a complexity bound on the decision problem for this theory.

**Theorem 1** ([CG84]). Consider a first-order sentence in the language of rings that mentions m polynomials in d variables, with total degree at most b, and with k quantifier alternations. The truth of such a sentence in  $\overline{\mathbb{Q}}$  can be decided in time  $(mb)^{d^{2k+2}}$ .

Following [BPR06, Remark 13.11], the truth of first-order sentences over  $\overline{\mathbb{Q}}$  with a fixed number of alternations can be decided in space  $(d \log b)^{O(1)}$ , by rewriting into the first-order theory of realclosed fields. Recall that the existential fragment is **NP**-hard and in **AM** assuming GRH [Koi96].

The following theorem is our main contribution:

**Theorem 2.** The orbit-closure determination problem for commutative matrices with the parameters (d, m, b) can be decided in time  $(mb)^{\text{poly}(d)}$ , and in space bounded by  $(d \log b)^{O(1)}$ .

Below, we give a brief and informal overview of the main ingredients of the proof of Theorem 2. Let G be a commutative algebraic group. It is known that the subset of semisimple matrices in G, denoted by  $G_s$ , forms an algebraic subgroup; likewise the set of unipotent matrices in G, denoted by  $G_u$ , forms an algebraic subgroup.

Let Z be the zero locus of the input polynomials. If Z is the orbit-closure of a point  $\boldsymbol{v}$  under the action of G, it can be written as  $\overline{G_u \cdot G_s \cdot \boldsymbol{v}}$ . Given that both  $G_s$  and  $G_u$  are s-generated, by Remark 10, G will be s-generated. By Proposition 8 and Lemma 15, it follows from the commutativity of G and the rational parameterisation of  $G_u$  that  $Z = G_u \cdot \overline{G_s \cdot \boldsymbol{v}}$ . The commutativity of  $G_s$  entails the existence of a matrix P and diagonal invertible matrices  $D_i$ ,  $1 \leq i \leq s$ , such that  $PG_sP^{-1} = \overline{\langle D_i : 1 \leq i \leq s \rangle}$ . By Lemma 12, we show that P can be chosen such that  $P\boldsymbol{v}$  is a zero-one vector.

The algorithm guesses the zero-one vector Pv. The semisimple group  $\overline{\langle D_i : 1 \leq i \leq s \rangle \cdot Pv}$  is a union of toric varieties. Denote by  $\Lambda$  the associated lattice of the defining ideal I of these toric varieties. By Proposition 7, since  $\overline{\langle D_i : 1 \leq i \leq s \rangle \cdot Pv}$  as a linear algebraic group has s topological generators, the torsion subgroup of  $\mathbb{Z}^d/\Lambda$  is s-generated. As a consequence of Proposition 6, the upper bound b on the degree of the defining polynomials of Z carries over to a generating set of I. Considering this degree bound b, the algorithm guesses a lattice  $\Lambda \subseteq \mathbb{Z}^k$  that is generated by vectors whose entries have absolute value at most b and has at most s elementary divisors not equal to one.

By Remark 10, there exist unipotent matrices  $U_i$ ,  $1 \leq i \leq s$ , that topologically generate the unipotent subgroup  $G_u$  of G. Furthermore, the equality  $G_u = \{\exp(\sum_{i=1}^s t_i \log U_i) : t_1, \ldots, t_s \in \overline{\mathbb{Q}}\}$  holds by Proposition 8.

Now we are ready to write an  $\exists^*\forall^*\exists$ -sentence in the theory of algebraically closed fields. The existential quantifiers encode the possible choices of the matrices P and  $U_1, \ldots, U_s$ , while the equality of Z and

$$\{\exp(\sum_{i=1}^{s} t_i \log U_i)h : t_1, \dots, t_s \in \overline{\mathbb{Q}}, h \in \overline{\langle D_i : 1 \le i \le s \rangle \cdot Pv}\}$$

is encoded by a  $\forall^*\exists$ -sentence with parameters P and  $U_1, \ldots, U_s$ . The algorithm returns "yes," meaning that Z is an orbit closure of a point v under the action of the group G, if the above sentence is satisfiable. By Theorem 1, the truth of such a sentence can be decided in time  $(mb)^{\text{poly}(d)}$ . Then the overall complexity bound follows from the fact that the number of choices of the lattice  $\Lambda$  and vector Pv is at most  $(2b)^{d^2+1}$ . So ends our informal overview of the proof of Theorem 2; the detailed proof can be found in Section 4.

**Example 3.** Let  $Z \subseteq \overline{\mathbb{Q}}^4$  be the zero set of the input ideal  $I := \langle F_1, F_2 \rangle$ , where

$$F_1 := x_2^2 - x_1 - x_4$$
 and  $F_2 := -2x_4x_2 - 2x_3^2 - \frac{1}{5}x_2x_3$ .

Our nondeterministic procedure in Theorem 2 shows that  $Z = \overline{\langle M \rangle \cdot v}$  where

$$M = \begin{pmatrix} 25 & 0 & -1 & 20\\ 0 & 5 & 0 & 0\\ 0 & -\frac{1}{2} & 5 & 0\\ 0 & 0 & 1 & 5 \end{pmatrix} \quad \text{and} \quad \boldsymbol{v} = \begin{pmatrix} 1\\ 1\\ 0\\ 0 \end{pmatrix}$$

An account of the procedural steps taken to produce M and v is given in Example 17.

**Orbit-Closure vs. Group Determination.** En route to proving Theorem 2 on orbit-closure determination, we consider a simpler variant-namely group determination. The two problems bear many similarities, especially in the case when the input polynomial ideal  $I \subseteq \mathbb{Q}[x_1, \ldots, x_d]$  defines a variety that is a union of toric varieties.

Let us first consider the group determination setting. In this setting, the sought group G can be topologically generated by diagonal matrices. By standard results, we associated a lattice  $\Lambda$  with the input ideal I. If the torsion subgroup of  $\mathbb{Z}^d/\Lambda$  is *s*-generated, then, by Proposition 7, the minimal number of topological generators of G is *s*. However, in the setting of orbit-closure determination, this lower bound on the number of generators may no longer hold. This phenomenon is witnessed by the following example.

**Example 4.** Let  $\Lambda$  be the lattice associated with the ideal  $I := \langle x_1^2 - x_3^2, x_2^2 - x_3^2 \rangle \subseteq \mathbb{Q}[x_1, x_2, x_3]$ . The torsion subgroup of  $\mathbb{Z}^3/\Lambda$  is 2-generated. By Proposition 7, the variety defined by the ideal I is 2-generated as an algebraic subgroup of diagonal matrices in  $\mathrm{GL}_3(\overline{\mathbb{Q}})$ , and in fact 2 is the minimal number of generators. An example of such generators are

$$\begin{pmatrix} 2 & 0 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & 2 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} -2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{pmatrix}.$$

By comparison, the variety defined by I arises as the orbit closure of v under the action of  $\overline{\langle M \rangle}$  where

$$M = \begin{pmatrix} 0 & -2 & 0 \\ 2 & 0 & 0 \\ 0 & 0 & 2 \end{pmatrix} \quad \text{and} \quad v = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}.$$

#### **1.2** Further Applications

**Orbit-Closure Determination and Loop Synthesis.** A related area of research to the orbitclosure determination problem is the synthesis of simple linear loops, which are branch-free loops characterized by linear update assignments. As explained above, while (implicit) orbit-closure problems are primarily used for invariant generation in program analysis, loop synthesis focuses on designing programs that conform to a given polynomial invariant.

Recent works [HBK20; Hit+24; Hum+22; KKV23] have focused on synthesising deterministic linear loops to ensure that a specified set of polynomial equalities holds among program variables at each loop iteration. This guarantees that the synthesised loops adhere to certain polynomial invariants. Geometrically, this amounts to finding an infinite orbit of a cyclic matrix group that lies inside a given variety<sup>2</sup>. Humenberger et al. [HBK20; Hum+22] give a method based on constraint solving that synthesises a loop satisfying a given polynomial invariant based on a user-supplied template. Recent work of Hitarth et al. [Hit+24] solves a version of the loop synthesis problem in which the polynomial invariant is given by a single quadratic equation. The authors of [KKV23] synthesise simple linear loops whose polynomial invariants are specified by ideals generated by pure difference binomials. The zero set of such an ideal is a union of toric varieties. The synthesis procedure in [KKV23] relies on a construction by Galuppi and Stanojkovski [GS21, Proposition 14] which demonstrates that for every toric variety V one can construct a diagonal rational matrix M such that  $\overline{\{M^n : n \in \mathbb{Z}\}} = V$ .

In terms of the results we present, the special case of Proposition 13 and Theorem 2 with s = 1 (i.e., when the sought for group is required to be cyclic), takes a variety  $Z \subseteq \overline{\mathbb{Q}}^d$  and determines whether Z is equal to the orbit-closure of the initial program variables under the action of a simple linear loop whose update assignments satisfy certain conditions. This is strictly stronger than the synthesis task in above references, which asks only that the infinite orbit closure of the synthesised loop be *contained in* Z. Furthermore, by lifting the restriction that the underling group is cyclic, we extend the scope from deterministic loops to nondeterministic loops.

**Group Determination and Matrix Completion.** The group determination problems for cyclic groups can also be seen as a type of matrix completion problem. Recall that the latter asks to determine whether a partially given matrix can be completed subject to some polynomial constraints on the entries, e.g. lower bounds on the rank [IKS10]. In the determination setting, we ask instead to complete the matrix subjected to polynomial constraints on *all powers* of the matrix.

Matrix completion has applications in areas such as combinatorial structures and perfect matching algorithms [BFS99; Edm67; IKS10; Lov79]. Arguably the most interesting application for matrix completion is that of polynomial identity testing [IKS10], arising from the fact that every arithmetic formula can be written as the determinant of a matrix of the linear forms [Val79]. Moreover, an arithmetic formula is non-zero if and only if the corresponding matrix can attain full rank.

## 2 Algebraic Background

Let  $\overline{\mathbb{Q}}$  denote the field of algebraic numbers and write  $\overline{\mathbb{Q}}[x_1, \ldots, x_d]$  for the ring of polynomials with coefficients in  $\overline{\mathbb{Q}}$  over the variables  $x_1, \ldots, x_d$ . A polynomial ideal I is an additive subgroup of  $\overline{\mathbb{Q}}[x_1, \ldots, x_d]$  that is closed under multiplication by polynomials in  $\overline{\mathbb{Q}}[x_1, \ldots, x_d]$ . Given a finite collection of polynomials  $S \subseteq \overline{\mathbb{Q}}[x_1, \ldots, x_d]$ , we denote by  $\langle S \rangle$  the *ideal generated by* S.

<sup>&</sup>lt;sup>2</sup>Notably, loop synthesis includes a non-triviality condition: asking that the orbit of the synthesised loop be infinite. For otherwise, the synthesis problem reduces to that of polynomial equation solving (see [Hit+24, Remark 2.8]). Such an assumption naturally aligns with that of a wandering point of an arithmetic dynamical system [Ben+19; Sil07].

An algebraic set (or variety) is the set of common zeroes of a finite collection of polynomials. By Hilbert's basis theorem every polynomial ideal  $I \subseteq \overline{\mathbb{Q}}[x_1, \ldots, x_d]$  is finitely generated. Thus the set

$$V(I) := \{ \boldsymbol{x} \in \overline{\mathbb{Q}}^d : f(\boldsymbol{x}) = 0 \text{ for all } f \in I \}$$

is a variety. The Zariski topology on  $\overline{\mathbb{Q}}^d$  has as its closed sets the varieties in  $\overline{\mathbb{Q}}^d$ . Given a set  $E \subseteq \overline{\mathbb{Q}}^d$ , we denote by  $\overline{E}$  the closure of E under the Zariski topology. Here  $\overline{E}$  is given by the smallest algebraic set that contains E.

Given an ideal  $I \subseteq \overline{\mathbb{Q}}[X]$ , where  $X = \{x_{i,j}\}_{1 \leq i,j \leq d}$ , and matrix  $M \in \overline{\mathbb{Q}}^{d \times d}$ , we write  $M \cdot I$  for the ideal  $\{f(MX) \in \overline{\mathbb{Q}}[X] : f \in I\}$ . Clearly  $V(M \cdot I) = \{A \in \overline{\mathbb{Q}}^{d \times d} : MA \in V(I)\}$ .

#### 2.1 Linear Algebraic Groups

Recall that a matrix  $M \in \overline{\mathbb{Q}}^{d \times d}$  is called *nilpotent* if there exists  $n \in \mathbb{N}$  such that  $M^n = 0_d$ ; M is *unipotent* if  $M - \mathrm{Id}_d$  is *nilpotent* (where  $\mathrm{Id}_d$  is the  $d \times d$  identity matrix), and M semisimple if it is diagonalisable over  $\overline{\mathbb{Q}}$ . We further say that M is *upper triangular* if all entries below the main diagonal are zero. We use the term *upper unitriangular* to refer to an upper triangular matrix whose entries along the main diagonal are all ones.

We write  $\operatorname{GL}_d(\overline{\mathbb{Q}})$  for the group of  $d \times d$  invertible matrices with entries in  $\overline{\mathbb{Q}}$ . We identify  $\operatorname{GL}_d(\overline{\mathbb{Q}})$  with the variety  $\{(M, y) \in \overline{\mathbb{Q}}^{d^2} \times \overline{\mathbb{Q}} : \det(M) \cdot y = 1\}$ . Under this identification, matrix multiplication is a polynomial map  $\operatorname{GL}_d(\overline{\mathbb{Q}}) \times \operatorname{GL}_d(\overline{\mathbb{Q}}) \to \operatorname{GL}_d(\overline{\mathbb{Q}})$ , and, by Cramer's rule, matrix inversion is also a polynomial map  $\operatorname{GL}_d(\overline{\mathbb{Q}}) \to \operatorname{GL}_d(\overline{\mathbb{Q}})$ . A *linear algebraic group* G is a Zariski-closed subgroup of  $\operatorname{GL}_d(\overline{\mathbb{Q}})$ . Recall that the identity component of G, denoted by  $G^\circ$ , is its irreducible component containing the identity.

We say that G is topologically generated by  $S \subseteq \operatorname{GL}_d(\overline{\mathbb{Q}})$  if G is the smallest Zariski closed subgroup of  $\operatorname{GL}_d(\overline{\mathbb{Q}})$  that contains S. In this case we write  $G = \overline{\langle S \rangle}$ . In case S is finite with s elements we say that G is s-generated. If G is a commutative algebraic group then the subset of semisimple matrices in G, denoted by  $G_s$ , forms an algebraic subgroup; likewise the set of unipotent matrices in G, denoted by  $G_u$ , forms an algebraic subgroup.

#### **Proposition 5.** Let G be an algebraic group, then G is topologically generated by a finite set.

*Proof.* Below, we denote by dim E the dimension of the variety E; that is, the maximal length of distinct nonempty irreducible subvarieties of E.

Let U be the linear algebraic group  $\langle G_u \rangle$ . It was proven that U can topologically be generated by dim U element; see [Nos+22, Proof of Lemma 6]. Moreover, G/U is an algebraic group which consists only of semisimple elements.

Recall that the quotient of a linear algebraic group by its normal subgroups are linear algebraic groups of higher dimensions [Hum75, Section 11.5]. Therefore,  $(G/U)^{\circ} = G^{\circ}/U$  is a torus and by [GS21, Proposition 14] it is 1-generated. Let  $h \in G^{\circ}$  such that  $G^{\circ}/U = \overline{\langle h \rangle}$ , and let  $H = \overline{\langle h, U \rangle} \subseteq G^{\circ}$ . We have  $G^{\circ}/U = H/U$  which implies dim  $H = \dim G^{\circ}$ . Since  $G^{\circ}$  is connected we have  $G^{\circ} = H$ , hence  $G^{\circ}$  is topologically generated by at most dim U + 1 elements.

In order to topologically generate G, it is sufficient to take the topological generators of  $G^{\circ}$  and one element from any other connected component of G. Hence, G is topologically generated by  $\dim U + |G/G^{\circ}|$  elements.

The *d*-dimensional multiplicative group over  $\overline{\mathbb{Q}}$  is defined as

$$\mathbb{G}_m^d = \mathbb{G}_m^d(\overline{\mathbb{Q}}) := \left\{ \boldsymbol{x} \in \overline{\mathbb{Q}}^d : x_1 \cdots x_d \neq 0 \right\}.$$

Evidently this is a commutative group with respect to the pointwise multiplication. We identify  $\mathbb{G}_m^d$  with the subgroup of diagonal matrices in  $\operatorname{GL}_d(\overline{\mathbb{Q}})$  via the map  $\Delta$  that sends  $(a_1, \ldots, a_d) \in \overline{\mathbb{Q}}^d$  to the diagonal matrix  $\Delta(a_1, \ldots, a_d) \in \operatorname{GL}_d(\overline{\mathbb{Q}})$ .

Given a subgroup  $\Lambda \subseteq \mathbb{Z}^d$ , define

$$H_{\Lambda} := \{ \boldsymbol{x} \in \mathbb{G}_m^d : \forall \boldsymbol{v} \in \Lambda \, (x_1^{v_1} \cdots x_d^{v_d} = 1) \}.$$

The map  $\Lambda \mapsto H_{\Lambda}$  yields an isomorphism between subgroups of  $\mathbb{Z}^d$  and algebraic subgroups of  $\mathbb{G}_m^d$ . This implies that  $\mathbb{G}_m^d$  is topologically generated by any *d*-tuple  $(g_1, \ldots, g_d)$  of multiplicatively independent elements of  $\overline{\mathbb{Q}}$ . It also follows that the vanishing ideal  $I \subseteq \overline{\mathbb{Q}}[x_1, \ldots, x_d]$  of an algebraic subgroup of  $\mathbb{G}_m^k$  is a so-called *pure binomial ideal*; that is, an ideal generated by polynomials of the form  $x_1^{\alpha_1} \cdots x_d^{\alpha_d} - x_1^{\beta_1} \cdots x_d^{\beta_d}$ , where  $\alpha_1, \ldots, \alpha_d$  and  $\beta_1, \ldots, \beta_d$  are non-negative integers. For future reference we note a mere *binomial ideal* is one that is generated by polynomials of the form  $x_1^{\alpha_1} \cdots x_d^{\alpha_d} - \lambda x_1^{\beta_1} \cdots x_d^{\beta_d}$ , where  $\lambda \in \overline{\mathbb{Q}}$ .

The following proposition [BG06, Proposition 3.2.14] shows how to recover the generators of a binomial ideal from the defining equations of an algebraic subgroup of  $\mathbb{G}_m^d$ .

**Proposition 6.** Let G be a subgroup of  $\mathbb{G}_m^d$  defined by polynomial equations  $f_i(\boldsymbol{x}) = \sum_{\boldsymbol{\lambda}} a_{i,\boldsymbol{\lambda}} \boldsymbol{x}^{\boldsymbol{\lambda}}$  for  $i = 1, \ldots, m$ . Let  $\mathcal{L}_i \subseteq \mathbb{Z}^d$  be the set of exponents of the monomials appearing in  $f_i$ . Then  $G = H_{\Lambda}$ , where  $\Lambda \subseteq \mathbb{Z}^d$  is generated by  $\boldsymbol{\lambda} - \boldsymbol{\lambda}'$ , where  $\boldsymbol{\lambda}, \boldsymbol{\lambda}' \in \mathcal{L}_i$  for some  $i \in \{1, \ldots, m\}$ .

The following is an amalgamation of standard results in Diophantine geometry concerning the number of generators of a subgroup of  $\mathbb{G}_m^d$  (cf. [BG06, Chapter 3]); we include a sketch proof for the reader's convenience.

**Proposition 7.** Let  $\Lambda$  have rank r and elementary divisors  $d_1, \ldots, d_r$ , where  $d_i \mid d_{i+1}$  for  $i = 1, \ldots, r-1$ . Then the following are equivalent for all  $s \in \{1, \ldots, r\}$ :

- 1.  $H_{\Lambda}$  is s-generated;
- 2.  $d_1 = \cdots = d_{r-s} = 1;$
- 3. the torsion subgroup of  $\mathbb{Z}^d/\Lambda$  is s-generated.

*Proof.* There exists a basis  $\boldsymbol{u}_1, \ldots, \boldsymbol{u}_d$  of  $\mathbb{Z}^d$  such that  $\Lambda$  is generated by the vectors  $d_1\boldsymbol{u}_1, \ldots, d_r\boldsymbol{u}_r$ . The map  $\varphi \colon \mathbb{G}_m^d \to \mathbb{G}_m^d$ , defined by  $\varphi(\boldsymbol{x}) = (\boldsymbol{x}^{\boldsymbol{u}_1}, \ldots, \boldsymbol{x}^{\boldsymbol{u}_d})$  is a Zariski-continuous group automorphism of  $\mathbb{G}_m^d$  that maps  $H_{\Lambda}$  to the group

$$G := \Omega_{d_1} \times \cdots \times \Omega_{d_r} \times \mathbb{G}_m^{d-r},$$

where  $\Omega_k$  denotes the group of all kth roots of unity for k a positive integer. Clearly  $H_{\Lambda}$  is s-generated if and only if G is s-generated.

Write  $F := \Omega_{d_1} \times \cdots \times \Omega_{d_r}$ . We note that  $G = F \times \mathbb{G}_m^{d-r}$  is s-generated if and only if F is s-generated. (In particular, if  $S \subseteq F$  is a generator of F then  $S \times \{g\}$  is a topological generator of Gfor any topological generator g of  $\mathbb{G}_m^{d-r}$ .) But from the fact that  $d_i \mid d_{i+1}$  for  $i = 1, \ldots, r-1$  we see that  $F = \Omega_{d_1} \times \cdots \times \Omega_{d_r}$  is s-generated if and only if  $d_1 = \cdots = d_{r-s} = 1$ , showing the equivalence of Items 1 and 2. For the equivalence of Items 2 and 3 we note that the torsion subgroup of  $\mathbb{Z}^d/\Lambda$ is  $\mathbb{Z}/d_1\mathbb{Z} \times \cdots \times \mathbb{Z}/d_r\mathbb{Z}$ , which is isomorphic to F. For unipotent  $A \in \operatorname{GL}_d(\overline{\mathbb{Q}})$  and nilpotent  $B \in \operatorname{GL}_d(\overline{\mathbb{Q}})$ , define

$$\log(A) := \sum_{k=1}^{d-1} (-1)^{k+1} \frac{(A-I)^k}{k} \quad \text{and} \quad \exp(B) := \sum_{k=0}^{d-1} \frac{B^k}{k!}.$$

Let  $G \subseteq \operatorname{GL}_d(\overline{\mathbb{Q}})$  be a commutative subgroup of unipotent matrices. Recall that  $L := \{\log(A) : A \in G\}$  is a linear subspace of  $\overline{\mathbb{Q}}^{d^2}$  consisting of nilpotent matrices [Bor91, Chapter II, Section 7.3]. Moreover, exp:  $L \to G$  and  $\log: G \to L$  yield polynomial isomophisms between L and G as algebraic groups. Taken together, these observations lead to the following proposition.

**Proposition 8.** Let G be a commutative subgroup of unipotent matrices and L the associated linear subspace of nilpotent matrices as above. Then G has a topological generator of cardinality s if and only if L is spanned by a set of s matrices as a  $\overline{\mathbb{Q}}$ -vector space.

*Proof.* For all  $A_1, \ldots, A_s \in G$  we have the following equivalences:

$$\{A_1, \dots, A_s\} \text{ is a topological generator of } G,$$
  

$$\Leftrightarrow \{A_1^{n_1} A_2^{n_2} \cdots A_s^{n_s} : n_1, \dots, n_s \in \mathbb{Z}\} \text{ is dense in } G,$$
  

$$\Leftrightarrow \{\sum_{i=1}^s n_i \log(A_i) : n_1, \dots, n_s \in \mathbb{Z}\} \text{ is dense in } L,$$
  

$$\Leftrightarrow \{\sum_{i=1}^s t_i \log(A_i) : t_1, \dots, t_s \in \mathbb{Q}\} = L,$$

as desired.

## 3 Commutative Group Determination

Recall that the group determination problem asks, given  $s \in \mathbb{N}$  and a family of polynomials  $f_1, \ldots, f_m \in \mathbb{Q}[\{x_{i,j}\}_{1 \leq i,j \leq d}]$  of total degree at most b, to determine whether their zero locus  $Z \subseteq \operatorname{GL}_d(\overline{\mathbb{Q}})$  is an s-generated matrix group. In this section, we first demonstrate a procedure for this problem subject to the constraint that the underlying group is semisimple commutative (Proposition 9). Next, we generalise this result by lifting the requirement that the matrices are semisimple (Proposition 11).

**Proposition 9.** The group determination problem for commutative semisimple matrices with the parameters (d, m, b) can be decided in time  $(mb)^{\text{poly}(d)}$ , and in space bounded by  $(d \log b)^{O(1)}$ .

*Proof.* Given semisimple commutative matrices  $M_1, \ldots, M_s \in \operatorname{GL}_d(\overline{\mathbb{Q}})$ , there exists  $P \in \operatorname{GL}_d(\overline{\mathbb{Q}})$ such that  $D_i := P^{-1}M_iP$  are diagonal matrices. Let G be the subgroup of  $\mathbb{G}_m^d$  defined by

$$G := \{g \in \mathbb{G}_m^d : \Delta(g) \in \overline{\langle D_i : 1 \le i \le s \rangle} \}.$$

Then  $Z = \overline{\langle M_i : 1 \leq i \leq s \rangle}$  if and only if  $P^{-1}ZP = \{\Delta(g) : g \in G\}$ . Now  $P^{-1}ZP$  is the zero set of polynomials of degree at most b and hence, by Proposition 6,  $Z = \langle M_i : 1 \leq i \leq s \rangle$  if and only if the group G has the form  $H_{\Lambda}$  for some lattice  $\Lambda \subseteq \mathbb{Z}^d$  that is generated by vectors having supremum norm at most b.

The decision procedure is thus as follows:

1. Guess a lattice  $\Lambda \subseteq \mathbb{Z}^d$  whose generators have norm at most b and has at most s elementary divisors not equal to one;

2. Determine whether there exists  $P \in \operatorname{GL}_d(\overline{\mathbb{Q}})$  such that  $P^{-1}ZP = \{\Delta(g) : g \in H_\Lambda\}$ .

Step 2 amounts to checking the truth in  $\overline{\mathbb{Q}}$  of the sentence

$$\exists P \in \operatorname{GL}_d(\overline{\mathbb{Q}}) \,\forall A \in \operatorname{GL}_d(\overline{\mathbb{Q}}) \,\left( P^{-1}AP \in Z \Leftrightarrow \bigwedge_{i \neq j} a_{ij} = 0 \land (a_{11}, \dots, a_{dd}) \in H_\Lambda \right),$$

with respect to the theory of algebraically closed fields. By Theorem 1, this can be done in time  $(mb)^{\text{poly}(d)}$ . The claimed running time for the overall procedure follows from the fact that the number of possibilities for the lattice  $\Lambda$  is at most  $(2b)^{d^2}$ .

By the following remark, a commutative algebraic group G is s-generated if its unipotent subgroup  $G_u$  and semisimple subgroup  $G_s$  are both s-generated.

**Remark 10.** Let  $U = \overline{\langle U_i : 1 \le i \le s \rangle}$  be a commutative unipotent *s*-generated algebraic group and  $S = \overline{\langle S_i : 1 \le i \le s \rangle}$  be a semisimple commutative *s*-generated algebraic group. Suppose moreover that the matrices  $\{U_i, S_i : 1 \le i \le s\}$  are pairwise commutative. Then we have

$$\overline{\langle U_i, S_i : 1 \le i \le s \rangle} = \overline{\langle S_i U_i : 1 \le i \le s \rangle}.$$

This relies on the fact that if  $A \in \operatorname{GL}_d(\overline{\mathbb{Q}})$  is semisimple and  $B \in \operatorname{GL}_d(\overline{\mathbb{Q}})$  is unipotent, then both A and B lie in the Zariski closure of the subgroup generated by their product AB; see [Hum75, Section 15.3].

The next proposition generalises the procedure witnessed in Proposition 9. In Proposition 11 we consider the group determination problem for s-generated commutative algebraic groups. Key to our generalisation is the determination of a matrix  $P \in \operatorname{GL}_d(\overline{\mathbb{Q}})$  and properties associated with the semisimple and unipotent subgroups of the group  $P^{-1}ZP$ .

**Proposition 11.** The group determination problem for commutative matrices with the parameters (d, m, b) can be decided in time  $(mb)^{\text{poly}(d)}$ , and in space bounded by  $(d \log b)^{O(1)}$ .

Proof. Given commutative matrices  $M_1, \ldots, M_s \in \operatorname{GL}_d(\overline{\mathbb{Q}})$ , let  $P \in \operatorname{GL}_d(\overline{\mathbb{Q}})$  be such that there exist diagonal matrices  $D_i$  and upper unitriangular matrices  $U_i, 1 \leq i \leq s$ , where  $P^{-1}M_iP = D_iU_i$ , and moreover  $D_i$  and  $U_i$  commute. Then we can recover  $\overline{\langle D_i : 1 \leq i \leq s \rangle}$  as the set of diagonal matrices in  $P^{-1}\overline{\langle M_i : 1 \leq i \leq s \rangle}P$ . It follows that  $\overline{\langle D_i : 1 \leq i \leq s \rangle}$  is the zero locus of a system of polynomials of degree at most b.

By Proposition 6,  $\overline{\langle D_i : 1 \leq i \leq s \rangle} = \{\Delta(g) : g \in H_\Lambda\}$  for some lattice  $\Lambda \subseteq \mathbb{Z}^d$  that is generated by vectors of norm at most b. Note that we can also recover  $\overline{\langle U_i : 1 \leq i \leq s \rangle}$  as the set of upper unitriangular matrices in  $P^{-1} \overline{\langle M_i : 1 \leq i \leq s \rangle} P$ .

The decision procedure is as follows. Guess a lattice  $\Lambda \subseteq \mathbb{Z}^d$  whose generating vectors have entries of absolute value at most b and has at most s elementary divisors not equal to one. Next, determine whether there exists  $P \in \operatorname{GL}_d(\overline{\mathbb{Q}})$  such that

- 1.  $G := P^{-1}ZP$  is a commutative group of upper triangular matrices;
- 2.  $\{A \in G : A \text{ diagonal}\} = \{\Delta(g) : g \in H_{\Lambda}\};$
- 3.  $\{\log(A) : A \in G, A \text{ unipotent}\}\$  is a linear variety of dimension at most s.

Item 1 checks that G is a commutative matrix group. In this case both the set  $G_s$  of semisimple matrices in G and the set  $G_u$  of unipotent matrices in G form subgroups of G. Next, Items 2 and 3 respectively check that  $G_s$  and  $G_u$  are s-generated (relying on Propositions 7 and 8). But this implies that G itself is s-generated, as noted in Remark 10.

The existence of P satisfying Items 1 to 3 reduces to checking the truth in  $\overline{\mathbb{Q}}$  of an  $\exists^*\forall^*$ -sentence in the theory of algebraically closed fields. The existential quantifiers correspond to the possible choices of P, while the universal quantifiers range over entries of the group G defined in Item 1. For a fixed choice of  $\Lambda$  the truth of such a formula can be decided in time  $(mb)^{\text{poly}(d)}$  by Theorem 1. Given that the number of possible choices of the lattice  $\Lambda$  is at most  $(2b)^{d^2}$  the claimed complexity bound immediately follows.

#### 4 Orbit-Closure Determination

Recall the aforementioned orbit-closure determination problem. The problem asks, given  $s \in \mathbb{N}$  and a family of polynomials  $f_1, \ldots, f_m \in \mathbb{Q}[\boldsymbol{x}]$  of total degree at most b, to determine whether their zero locus  $Z \subseteq \operatorname{GL}_d(\overline{\mathbb{Q}})$  is the orbit closure of some point  $\boldsymbol{v} \in \overline{\mathbb{Q}}^d$  under the action of an s-generated matrix group, i.e., determine whether there exist matrices  $M_1, \ldots, M_s \in \operatorname{GL}_d(\overline{\mathbb{Q}})$  and vector  $\boldsymbol{v} \in \overline{\mathbb{Q}}^d$ such that Z is the Zariski closure of the orbit  $\{M_1^{\ell_1} \cdots M_s^{\ell_s} \boldsymbol{v} : \ell_1, \ldots, \ell_s \in \mathbb{Z}\}.$ 

The main contributions of this section are the procedures for certain cases of the orbit-closure determination problem (Proposition 13 and Theorem 2). The procedure in Proposition 13 makes the additional assumptions that the generators of the matrix group are semisimple and pairwise commutative. The procedure in Theorem 2 lifts the requirement that the generators are semisimple. We illustrate the procedures with a worked example (Example 17) at the close of this section.

In the work that follows, it is convenient to employ the next lemma, which intuitively speaking describes that orbit closures under the action of semisimple commutative groups are isomorphic to the orbit closure of **1**.

**Lemma 12.** Let  $G = \overline{\langle M_i : 1 \leq i \leq s \rangle} \subseteq \overline{\mathbb{Q}}^{d \times d}$  be a semisimple commutative algebraic group, and  $v \in \overline{\mathbb{Q}}^d$ . There exist  $P \in \operatorname{GL}_d(\overline{\mathbb{Q}})$ , diagonal matrices  $D_1, \ldots, D_s \in \operatorname{GL}_d(\overline{\mathbb{Q}})$ , and  $T \in \{0, 1\}^{d \times k}$  with the following properties:

- 1.  $M_i = P^{-1}D_iP$ ,
- 2.  $T^{\mathsf{T}}T = \mathrm{Id}_k$ ,
- 3. Pv = T1.

*Proof.* Since the matrices  $M_i$  are commutative, there exist a matrix  $Q \in \operatorname{GL}_d(\overline{\mathbb{Q}})$ , and diagonal matrices  $D_1, \ldots, D_s \in \operatorname{GL}_d(\overline{\mathbb{Q}})$  such that  $M_i = Q^{-1}D_iQ$ . Let  $R \in \operatorname{GL}_d(\overline{\mathbb{Q}})$  be a diagonal matrix such that  $RQ\mathbf{v} \in \{0,1\}^d$ . We define  $P \coloneqq RQ$ . Since R is diagonal, it commutes with all  $D_i$  matrices. Thus we have  $M_i = Q^{-1}D_iQ = P^{-1}D_iP$ .

It remains to determine the matrix T. We write  $Pv = e_{i_1} + \cdots + e_{i_k}$  as a sum of standard unit vectors of  $\overline{\mathbb{Q}}^d$ . We define T to be the  $d \times k$  matrix with columns  $e_{i_1}, \ldots, e_{i_k}$ . It follows that  $T^{\mathsf{T}}T = \mathrm{Id}_k$  and  $Pv = T\mathbf{1}$ .

We move onto the first of the two main orbit-closure results in this section.

**Proposition 13.** The orbit-closure determination problem for commutative semisimple matrices with the parameters (d, m, b) can be decided in time  $(mb)^{\text{poly}(d)}$ , and in space bounded by  $(d \log b)^{O(1)}$ .

*Proof.* Suppose that  $Z = \overline{\langle M_i : 1 \leq i \leq s \rangle \cdot v}$  for semisimple commutative matrices  $M_1, \ldots, M_s \in \operatorname{GL}_d(\overline{\mathbb{Q}})$  and a vector  $v \in \overline{\mathbb{Q}}^d$ . By Lemma 12, there exist matrices  $P \in \operatorname{GL}_d(\overline{\mathbb{Q}})$  and  $T \in \{0, 1\}^{d \times k}$  such that  $D_i := PM_iP^{-1}$  are diagonal, and moreover  $T^{\mathsf{T}}T = \operatorname{Id}_k$  and  $Pv = T\mathbf{1}$  hold.

For all  $i \in \{1, \dots, s\}$ , denote by  $D_i \in \operatorname{GL}_k(\overline{\mathbb{Q}})$  the diagonal matrix uniquely defined by the requirement that  $D_i T = T \tilde{D}_i$ . Write

$$G := \{g \in \mathbb{G}_m^k : \Delta(g) \in \overline{\langle \tilde{D}_i : 1 \le i \le s \rangle} \}.$$

Then

$$PZ = \overline{\langle D_i : 1 \le i \le s \rangle \cdot T\mathbf{1}} = T \overline{\langle \tilde{D}_i : 1 \le i \le s \rangle} \cdot \mathbf{1} = \{T\Delta(g)\mathbf{1} : g \in G\},\$$

where the penultimate equality relies on the fact that image of a Zariski-closed set under an injective linear map is again Zariski closed. Note that G can alternatively be written as  $\{g \in \mathbb{G}_m^k : Ug \in PZ\}$ and is thereby defined by polynomials of total degree at most b. It follows from Proposition 6 that  $G = H_{\Lambda}$  for some lattice  $\Lambda \subseteq \mathbb{Z}^k$  that is generated by vectors whose entries have absolute value at most b.

Conversely, suppose that  $PZ = \{Tg : g \in H_{\Lambda}\}$  for some matrices  $P \in \operatorname{GL}_d(\overline{\mathbb{Q}})$  and  $T \in \{0, 1\}^{d \times k}$ such that  $U^{\mathsf{T}}U = \operatorname{Id}_k$ , and lattice  $\Lambda$  as above. Then

$$Z = \overline{\langle M_i : 1 \le i \le s \rangle \cdot \boldsymbol{v}},$$

where  $v = P^{-1}T\mathbf{1}$  and  $M_i := P^{-1}D_iP$  with  $D_i \in \operatorname{GL}_d(\overline{\mathbb{Q}})$  being any diagonal matrices such that  $D_iT = T\Delta(g_i)$  for some topological generators  $\{g_1, \ldots, g_n\}$  of  $H_{\Lambda}$ .

In summary, the decision procedure is as follows:

- 1. Guess  $k \subseteq \{0, \ldots, d\}$  and  $T \in \{0, 1\}^{d \times k}$  such that  $T^{\mathsf{T}}T = \mathrm{Id}_k$ ;
- 2. Guess a lattice  $\Lambda \subseteq \mathbb{Z}^k$  that is generated by vectors whose entries have absolute value at most b and has at most s elementary divisor not equal to one;
- 3. Determine whether there exists  $P \in \operatorname{GL}_d(\overline{\mathbb{Q}})$  such that  $PZ = \{Tg : g \in H_\Lambda\}$ .

Step 3 can be reduced in polynomial time to checking the truth of a  $\exists^*\forall^*$ -sentence in the theory of algebraically closed fields: the existential quantifiers correspond to the matrix P, while the universal quantifiers are used to encode the equation  $PZ = \{Tg : g \in H_\Lambda\}$ . By Theorem 1, the truth of such a sentence can be decided in time  $(mb)^{\text{poly}(d)}$ . The claimed overall complexity bound now follows from the fact that there are at most  $(2b)^{d^2+1}$  choices of the lattice  $\Lambda$  and matrix U.

We include a worked example that demonstrates the constructive subroutines in Proposition 13.

**Example 14.** Let  $Z \subseteq \overline{\mathbb{Q}}^2$  be the zero set of the ideal  $I := \langle 4x^2 + y^2 + 4xy - x - y \rangle$ . In this example, we construct a matrix  $M \in \operatorname{GL}_2(\overline{\mathbb{Q}})$  and vector  $\boldsymbol{v} \in \overline{\mathbb{Q}}^2$  such that Z is precisely the Zariski closure of the orbit  $\{M^n \boldsymbol{v} : n \in \mathbb{Z}\}$ .

Suppose that in Steps 1 and 2 of the procedure in Proposition 13 we guess  $H_{\Lambda} := \{(x, y) \in \mathbb{G}_m^2 : x^2 - y = 0\}$  and the matrix  $T := \mathrm{Id}_2$ . For Step 3, we want to find all invertible matrices  $P = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{GL}_2(\overline{\mathbb{Q}})$  such that  $V(P \cdot I) = \{Tg : g \in H_{\Lambda}\}$ . This is equivalent to the requirement that the two equations

$$4(ax + by)^{2} + (cx + dy)^{2} + 4(ax + by)(cx + dy) - ax - by - cx - dy \quad \text{and} \quad x^{2} - y$$

are multiples of one another. Therefore the equations defining P comprise the following ideal:

$$J_P := \langle 4a^2 + c^2 + 4ac - b - d, 4b^2 + d^2 + 4bd, 8ab + 2cd + 4ad + 4bc, a + c \rangle$$
$$= \langle a + c, c^2 - b - d, 2bc + cd, (2b + d)^2 \rangle.$$

One choice of P is  $\begin{pmatrix} 1 & -1 \\ -1 & 2 \end{pmatrix}$ . Thus we associate with Z the Zariski closure of the orbit  $\{M^n \boldsymbol{v} : n \in \mathbb{Z}\}$  where

$$M := P \begin{pmatrix} 2 & 0 \\ 0 & 4 \end{pmatrix} P^{-1} = \begin{pmatrix} 0 & -2 \\ 4 & 6 \end{pmatrix} \quad \text{and} \quad \boldsymbol{v} := P \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

The following lemma is crucial in generalizing the procedure in Proposition 13 for orbit-closure determination of semisimple commutative groups to the general setting.

**Lemma 15.** Let  $G \subseteq \operatorname{GL}_d(\overline{\mathbb{Q}})$  be a commutative algebraic group and  $v \subseteq \overline{\mathbb{Q}}^d$ . Then

$$\overline{G\cdot \boldsymbol{v}}=G_u\cdot\overline{G_s\cdot \boldsymbol{v}}\,,$$

where  $G_u$  and  $G_s$  are the subgroups of unipotent and semisimple elements in G, respectively.

*Proof.* We assume that G is s-generated. Write  $G = \overline{\langle M_i : 1 \leq i \leq s \rangle}$  and  $G_u = \overline{\langle U_i : 1 \leq i \leq s \rangle}$ . Define  $V := \overline{G_s \cdot v}$ . For every  $w \in V$ , we define the polynomial map

$$\phi_{\boldsymbol{w}} : \overline{\mathbb{Q}}^n \to G_u \cdot V$$
  
 $(t_1, \dots, t_s) \mapsto \exp(\sum_{i=1}^s t_i \log U_i) \boldsymbol{w}.$ 

Since  $\phi_{\boldsymbol{w}}$  is a continuous map,  $\phi_{\boldsymbol{w}}^{-1}(V)$  is Zariski closed. Let  $H \coloneqq \bigcap_{\boldsymbol{w} \in V} \phi_{\boldsymbol{w}}^{-1}(V)$ . We note that H is Zariski closed and is equal to

$$\{(t_1,\ldots,t_s)\in\overline{\mathbb{Q}}^n:\exp(\sum_{i=1}^s t_i\log U_i)\cdot V=V\}.$$

Observe that for every pair  $(t_1, \ldots, t_s), (t'_1, \ldots, t'_s) \in H$  and  $a, b \in \mathbb{N}$ , we have  $a(t_1, \ldots, t_s) + b(t'_1, \ldots, t'_s) \in H$ . But since H is Zariski closed, this property implies that H is a linear vector space. The following claim is crucial for the continuation of the proof.

**Claim 16.** For all  $(t_1, \ldots, t_s) \in \overline{\mathbb{Q}}^d$  such that  $\exp(\sum_{i=1}^s t_i \log U_i) \cdot V \cap V \neq \emptyset$ , we have that

$$\exp(\sum_{i=1}^{s} t_i \log U_i) \cdot V = V.$$

Proof of the claim. Suppose that  $\exp(\sum_{i=1}^{s} t_i \log U_i) \cdot V \cap V \neq \emptyset$ . Then there exists  $\boldsymbol{w} \in G_s \cdot \boldsymbol{v}$  such that  $\exp(\sum_{i=1}^{s} t_i \log U_i) \boldsymbol{w} \in V$ . By commutativity of G we get,

$$G_s \cdot \exp(\sum_{i=1}^s t_i \log U_i) \boldsymbol{w} = \exp(\sum_{i=1}^s t_i \log U_i) \cdot G_s \cdot \boldsymbol{w} \subseteq G_s \cdot V = V.$$

Thus  $\exp(\sum_{i=1}^{s} t_i \log U_i) \cdot V \subseteq V$  holds. It implies  $\exp(\sum_{i=1}^{s} t_i \log U_i) \cdot V = V$ , concluding the proof of the claim.

Denote by W the orthogonal space of H, meaning that  $\overline{\mathbb{Q}}^d = W \oplus H$ . Let us introduce the following map

$$\phi: W \times V \to G_u \cdot V$$
  
 $((t_1, \dots, t_s), \boldsymbol{w}) \mapsto \exp(\sum_{i=1}^s t_i \log U_i) \boldsymbol{w}$ 

The map  $\phi$  is a polynomial map, and one-to-one correspondence. To prove this, assume that

$$\exp(\sum_{i=1}^{s} t_i \log U_i) \boldsymbol{w} = \exp(\sum_{i=1}^{s} t'_i \log U_i) \boldsymbol{w}'$$

holds for some  $(t_1, \ldots, t_s), (t'_1, \ldots, t'_s) \in W$  and  $\boldsymbol{w}, \boldsymbol{w'} \in V$ . Then

$$\exp(\sum_{i=1}^{s} (t_i - t'_i) \log U_i) \cdot V \cap V \neq \emptyset.$$

By the above and Claim 16 we have  $\exp(\sum_{i=1}^{s} (t_i - t'_i) \log U_i) \cdot V = V$ . This implies that  $(t_1, \ldots, t_s) = (t'_1, \ldots, t'_s)$ , which in turns shows that  $\boldsymbol{w} = \boldsymbol{w}'$ .

Since  $\phi$  is a one-to-one correspondence between  $W \times V$  and  $G_u \cdot V$ , and since  $W \times V$  is Zariski closed, we have that  $\phi(W \times V) = G_u \cdot V$  is Zariski closed. This completes the proof.

The following theorem is our main contribution, which provides a decision procedure for the orbit-closure determination problem for commutative groups. The generalization of this result to the case of general matrix groups appears to be challenging.

**Theorem 2.** The orbit-closure determination problem for commutative matrices with the parameters (d, m, b) can be decided in time  $(mb)^{\text{poly}(d)}$ , and in space bounded by  $(d \log b)^{O(1)}$ .

*Proof.* Suppose that  $Z = \overline{G \cdot v}$  for some commutative algebraic group  $G = \overline{\langle M_i : 1 \leq i \leq s \rangle} \subseteq GL_d(\overline{\mathbb{Q}})$  and  $v \in \overline{\mathbb{Q}}^d$ . Let  $G_u$  be the group of unipotent elements of G and  $G_s$  be the group of semisimple elements of G. Then by Lemma 15, we have  $Z = G_u \cdot \overline{G_s \cdot v}$ .

We apply Lemma 12 to the semisimple group  $G_s$ . Let  $P \in \operatorname{GL}_d(\overline{\mathbb{Q}})$  and  $T \in \{0,1\}^{d \times k}$  be matrices such that  $PG_sP^{-1} = \overline{\langle D_i : 1 \leq i \leq s \rangle}$  where  $D_i$  are diagonal invertible matrices, and the conditions  $T^{\mathsf{T}}T = \operatorname{Id}_k$  and  $Pv = T\mathbf{1}$  are satisfied.

For all  $i \in \{1, \dots, s\}$ , denote by  $\tilde{D}_i \in \operatorname{GL}_k(\overline{\mathbb{Q}})$  be the diagonal matrix uniquely defined by the requirement that  $D_i T = T \tilde{D}_i$ . Furthermore, write

$$\tilde{G} := \{g \in \mathbb{G}_m^k : \Delta(g) \in \overline{\langle \tilde{D}_i : 1 \le i \le s \rangle}\}$$
 and  $G_u := \overline{\langle U_i : 1 \le i \le s \rangle}.$ 

Then we have

$$Z = G_u \cdot \overline{G_s \cdot v}$$
  
=  $\overline{\langle U_i : 1 \le i \le s \rangle} \cdot P^{-1} \overline{\langle D_i : 1 \le i \le s \rangle \cdot Pv}$   
=  $\overline{\langle U_i : 1 \le i \le s \rangle} \cdot P^{-1} \overline{T} \overline{\langle \tilde{D}_i : 1 \le i \le s \rangle \cdot \mathbf{1}}$   
=  $\{ \exp(\sum_{i=1}^s t_i \log U_i) P^{-1} Tg : g \in \tilde{G}, t_1, \dots, t_s \in \overline{\mathbb{Q}} \},$ 

where the first equality follows from Lemma 15, and the last equality follows Proposition 8. Note that  $\tilde{G} = \{g \in \mathbb{G}_m^k : P^{-1}Tg \in Z\}$ . It follows that  $\tilde{G}$  is defined by polynomials of total degree at most b and hence has the form  $H_{\Lambda}$  for some lattice  $\Lambda \subseteq \mathbb{Z}^k$  that is generated by vectors whose entries have absolute value at most b and has at most s elementary divisors not equal to one.

Conversely, if

$$Z = \{ \exp(\sum_{i=1}^{s} t_i \log U_i) P^{-1} T g : g \in H_{\Lambda}, t_1, \dots, t_s \in \overline{\mathbb{Q}} \}$$

for some matrix  $P \in \operatorname{GL}_d(\overline{\mathbb{Q}})$ , unipotent matrices  $U_1, \ldots, U_s \in \operatorname{GL}_d(\overline{\mathbb{Q}})$  that are commutative, matrix  $T \in \{0, 1\}^{d \times k}$ , and lattice  $\Lambda$  as above, then  $Z = \langle M_i : 1 \leq i \leq s \rangle \cdot \boldsymbol{v}$ , where  $M_i := P^{-1}U_iD_iP$ and  $\{D_i : 1 \leq i \leq s\} \subseteq \operatorname{GL}_d(\overline{\mathbb{Q}})$  are any diagonal matrices such that  $D_iT = T\Delta(g_i)$  for some topological generators  $\{g_1, \ldots, g_s\}$  of  $H_{\Lambda}$ .

In summary, the decision procedure is as follows:

- 1. Guess  $k \subseteq \{0, \ldots, d\}$  and  $T \in \{0, 1\}^{d \times k}$  such that  $T^{\mathsf{T}}T = \mathrm{Id}_k$ ;
- 2. Guess a lattice  $\Lambda \subseteq \mathbb{Z}^k$  that is generated by vectors whose entries have absolute value at most b and has at most s elementary divisor not equal to one;
- 3. Return "yes" if there exists  $P \in \operatorname{GL}_d(\overline{\mathbb{Q}})$  and unipotent commutative matrices  $U_1, \ldots, U_s \in \operatorname{GL}_d(\overline{\mathbb{Q}})$  such that  $Z = \{\exp(\sum_{i=1}^s t_i \log U_i) P^{-1}Tg : g \in H_\Lambda, t_1, \ldots, t_s \in \overline{\mathbb{Q}}\}.$

Step 3 can be reduced to checking the truth of an  $\exists^*\forall^*\exists$ -sentence (that is, with a quantifier prefix comprising a block of existential quantifiers, a block of universal quantifiers, followed by a single existential quantifier) with respect to the theory of algebraically closed fields. The existential quantifiers encode the possible choices of the matrices P and  $U_1, \ldots, U_s$ , while the equality of Z and  $\{\exp(\sum_{i=1}^s t_i \log U_i) P^{-1}Tg : g \in H_{\Lambda}, t_1, \ldots, t_s \in \overline{\mathbb{Q}}\}$  is encoded by a  $\forall^*\exists$ -sentence with parameters P and  $U_1, \ldots, U_s$ , namely

$$\forall \boldsymbol{z} \in \overline{\mathbb{Q}}^d \left( \boldsymbol{z} \in Z \Leftrightarrow \exists (t_1, \dots, t_s) \, s.t. \, (T^{\mathsf{T}} P \exp(-\sum_{i=1}^s t_i \log U_i) \, \boldsymbol{z} \in H_{\Lambda}) \right).$$

By Theorem 1, the truth of such a sentence can be decided in time  $(mb)^{\text{poly}(d)}$ . Then the overall complexity bound follows from the fact that the number of choices of the lattice  $\Lambda$  and matrix T is at most  $(2b)^{d^2+1}$ .

Example 17 below applies the procedure in Theorem 2 to the variety we first saw in Example 3 in the Introduction. The calculations involved in the preparation of Examples 17 and 18 were performed in MACAULAY2 [GS].

**Example 17.** Let  $Z \subseteq \overline{\mathbb{Q}}^4$  be the zero set of the ideal  $I := \langle F_1, F_2 \rangle$  where

$$F_1 := x_2^2 - x_1 - x_4$$
 and  $F_2 := -2x_4x_2 - 2x_3^2 - \frac{1}{5}x_2x_3$ .

Below we shall construct a matrix  $M \in \operatorname{GL}_d(\overline{\mathbb{Q}})$  and vector  $\boldsymbol{v} \in \overline{\mathbb{Q}}^d$  such that Z is the Zariski closure of the orbit  $\{M^n \boldsymbol{v} : n \in \mathbb{Z}\}$ .

Suppose that in Steps 1 and 2 of the procedure in Theorem 2 we guess that  $H_{\Lambda} = \{(x, y) \in \mathbb{G}_m^2 : x^2 - y = 0\}, T = \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}$ , and U has the form  $\begin{pmatrix} 1 & \lambda & 0 & 0 \\ 0 & 1 & \lambda & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$ . For Step 3, we would like to find the set of invertible matrices  $P = (p_{ij})_{\{1 \le i, j \le 4\}} \subseteq \operatorname{GL}_4(\mathbb{Q})$  such that there exists  $\lambda$  for which

$$V(P \cdot I) = \{\exp(t \log U)Tg : g \in H_{\Lambda}, t \in \overline{\mathbb{Q}}\}.$$

Note that

$$\exp(t\log U) = \begin{pmatrix} 1 & t\lambda & \frac{t(t-1)}{2}\lambda^2 & 0\\ 0 & 1 & t\lambda & 0\\ 0 & 0 & 1 & 0\\ 0 & 0 & 0 & 1 \end{pmatrix}; \text{ thus } \exp(t\log U)T\begin{pmatrix} x\\ x^2 \end{pmatrix} = \begin{pmatrix} \frac{t(t-1)}{2}x\\ t\lambda x\\ x\\ x^2 \end{pmatrix}.$$

The ideal defining  $\{\exp(t\log U)Tg : g \in H_{\Lambda}, t \in \overline{\mathbb{Q}}\}\$  is  $H := \langle x_3^2 - x_4, x_1x_3 - \frac{1}{2}x_2^2 + \frac{\lambda}{2}x_2x_3 \rangle$ . Consider the ideal

$$I_P = \langle F_1(PX), F_2(PX) \rangle \subseteq \mathbb{Q}[P = (p_{ij}), \lambda, y] / \langle (\det P)y - 1 \rangle [x_1, x_2, x_3, x_4]$$

By applying Algorithm CONTAINMENTISO<sup>3</sup> and eliminating  $\lambda$  we obtain the following ideal, defining the set of admissible choices of P:

$$J_P := \langle p_{34}, p_{31}, p_{24}, p_{22}, p_{21}, p_{13} + p_{43}, p_{12} + p_{42}, p_{11} + p_{41}, p_{33}p_{44}, p_{32}p_{44}, p_{23}p_{44}, p_{14}p_{44} + p_{44}^2, p_{23}p_{33} + 10p_{33}^2 + 10p_{23}p_{43}, 2p_{32}^2 + p_{23}p_{41}, p_{23}^2 - p_{14} - p_{44} \rangle.$$

One may choose for example

$$P = \begin{pmatrix} \frac{1}{2} & 0 & 0 & 1\\ 0 & 0 & 1 & 0\\ 0 & \frac{1}{2} & 0 & 0\\ -\frac{1}{2} & 0 & 0 & 0 \end{pmatrix} \quad \text{and} \quad \lambda = -\frac{1}{5}.$$

Thus we associate with Z the Zariski closure of the orbit  $\{M^n \boldsymbol{v} : n \in \mathbb{Z}\}$  where

$$M = PUDP^{-1} = \begin{pmatrix} 25 & 0 & -1 & 20\\ 0 & 5 & 0 & 0\\ 0 & -\frac{1}{2} & 5 & 0\\ 0 & 0 & 1 & 5 \end{pmatrix} \quad \text{and} \quad \mathbf{v} = PT\mathbf{1} = \begin{pmatrix} 1\\ 1\\ 0\\ 0 \end{pmatrix}$$

### 5 Algorithms to Compute Generators

In Section 3 we gave an algorithm to determine whether a given variety is the Zariski closure of a commutative matrix group. The method there can also be used to find a set of generators of such a group, using the fact that the theory of algebraically closed fields admits quantifier elimination. In this section we provide two alternative algorithms for cyclic groups to compute a generator, relying instead on Gröbner-basis techniques. The first algorithm finds a semisimple generator, if one exists, while the second algorithm finds a generator in the general case.

Let  $I \subseteq \overline{\mathbb{Q}}[x_1, \ldots, x_d]$  be an ideal and denote by  $\sqrt{I}$  the radical of I, defined as  $\sqrt{I} := \{f \in \overline{\mathbb{Q}}[x_1, \ldots, x_d] \mid f^n \in I \text{ for some } n \in \mathbb{N}\}$ . By Hilbert's Nullstellensatz, the ideal of all polynomials that vanish on  $V(I) \in \overline{\mathbb{Q}}^d$  is  $\sqrt{I}$ . The ideal I is primary if for all  $f, g \in \overline{\mathbb{Q}}[x_1, \ldots, x_d]$ , if  $fg \in I$  then  $f \in I$  or  $g^n \in I$  for some  $n \in \mathbb{N}$ . Recall that the radical of a primary ideal is necessarily prime.

A polynomial ideal I can be written as the intersection of primary ideals, giving the so-called *primary decomposition* of I. It is known that there exists a unique irredundant primary decomposition  $I = \bigcap_{i=1}^{\ell} Q_i$ , that is, a finite set  $\{Q_1, \ldots, Q_\ell\}$  of primary ideals such that (1) the prime ideals  $\sqrt{Q_i}$  are all distinct; and (2)  $\bigcap_{i\neq j} Q_i \not\subseteq Q_j$  holds for all  $j \in \{1, \cdots, \ell\}$ . The prime ideals, the  $\sqrt{Q_i}$ 's, are called the *associated primes* of I. An associated prime  $\sqrt{Q}$  of the ideal I is called *minimal* if it does not contain any other associated primes of I.

Both algorithms take as input a variety Z, given as the zero set of an ideal  $I \subseteq \mathbb{Q}[X]$ ,  $X = \{x_{i,j}\}_{1 \leq i,j \leq d}$ . We will assume that it has already been verified that Z is a commutative subgroup of  $\operatorname{GL}_d(\overline{\mathbb{Q}})$ . This verification entails first checking that Z is closed under matrix multiplication (which entails closure under matrix inverse), which amounts to showing that

$$F(XY) \in \sqrt{I(X) + I(Y)}$$

<sup>&</sup>lt;sup>3</sup>The algorithm CONTAINMENTISO inputs two ideals  $I_1$  and  $I_2$  and outputs the locus of points P for which  $P \cdot I_1 \subseteq P \cdot I_2$ . This algorithm thereby solves a generalisation of the ideal membership algorithm since it determines the containment of an ideal into another after a change of variables. Clearly CONTAINMENTISO can also be applied to determine equality after a change of variables, since  $P \cdot I_1 = P \cdot I_2$  if and only if  $P \cdot I_1 \subseteq P \cdot I_2$  and  $P \cdot I_2 \subseteq P \cdot I_1$ . See [KMM17, Algorithm 2.9] for more details.

Cyclic Groups: Semisimple Generator	
Input:	An ideal $I \subseteq \mathbb{Q}[X]$ , $X = \{x_{i,j}\}_{1 \leq i,j \leq d}$ , with $q$ minimal associated primes, such that $V(I)$ is a commutative linear algebraic group.
Output:	Determine whether there exists a semisimple matrix $M \in \operatorname{GL}_d(\overline{\mathbb{Q}})$ such that $V(I) = \overline{\langle M \rangle}$ . If "yes", output such a matrix $M$ .
Line 1:	Define the ideal $J_0 \subseteq \mathbb{Q}[P, X, Q]$ as follows
	$J_0 := \langle F_1(PXQ), \ldots, F_k(PXQ), PQ - \mathrm{Id}_d, \{x_{i,j}\}_{i \neq j} \rangle$
	where $P = \{p_{i,j}\}_{1 \le i,j \le d}$ , $X = \{x_{i,j}\}_{1 \le i,j \le d}$ , and $Q = \{q_{i,j}\}_{1 \le i,j \le d}$ .
Line 2:	Write $J := \sqrt{J_0 \cap \mathbb{Q}[X]}$ . Compute the primary decomposition $J = \bigcap_{s \in S} \mathcal{P}_s$ .
Line 3:	Check whether all primary components $\mathcal{P}_s$ of $J$ are binomials using Gröbner basis computation; <b>return</b> "no" if this test fails.
Line 4:	Let $\mathcal{P}_0$ be one of the primary component of $J$ such that $\mathrm{Id}_d \in V(\mathcal{P}_0)$ .
Line 5:	Following Proposition 7, we can construct a rational diagonal matrix $D$ for $\mathcal{P}_0$ , such that all the entries of $\sqrt[q]{D}$ lie in $\mathbb{Q}[\zeta_q]$ . Write $D_q := \sqrt[q]{D}$ .
Line 6:	Check whether, for all $i \in \{1, \ldots, q-1\}$ , the ideal $D_q^i \cdot \mathcal{P}_0$ is a primary component of $J$ ; <b>return</b> "no" if this test fails.
Line 7:	Write $I_q := \bigcap_{1 \le i \le q} D_q^i \cdot \mathcal{P}_0.$
Line 8:	Check whether $J = \bigcap_{\sigma \in S_d} M_{\sigma} I_q M_{\sigma}^{-1}$ where $M_{\sigma}$ is the permutation matrix corresponding to $\sigma \in S_d$ ; <b>return</b> "no" if this test fails.
Line 9:	Define the ideal $J_1 := \langle F_1(QD_qP), \dots, F_k(QD_qP), PQ - I \rangle \cap \mathbb{Q}[P].$
	Pick $\tilde{P} \in V(J_1)$ .
Line 10:	Check whether $I = \tilde{P}I_q\tilde{P}^{-1}$ ; return "no" if this test fails.
Return:	"yes" together with the matrix $\tilde{P}D_q\tilde{P}^{-1}$ .

Figure 1: A procedure for the group determination problem of cyclc groups, specific to semisimple generators.

for all polynomials F in I, where  $X = \{x_{i,j}\}_{1 \le i,j \le d}$ ,  $Y = \{y_{i,j}\}_{1 \le i,j \le d}$ . Commutativity is captured by showing that  $XY - YX \in \sqrt{I(X) + I(Y)}$ .

#### 5.1 Semisimple Generator

In the following we describe a procedure that, given an ideal  $I \subseteq \mathbb{Q}[X]$ ,  $X = \{x_{i,j}\}_{1 \leq i,j \leq d}$ , determines whether there exists a semisimple matrix  $M \in \operatorname{GL}_d(\overline{\mathbb{Q}})$  such that I is the vanishing ideal of the group  $\langle M \rangle$  and which moreover outputs such an M in case the answer is "yes". We show that if such an M exists then it can be chosen such that its eigenvalues lie in the number field  $\mathbb{Q}(\zeta_q)$ , where  $\zeta_q$  is a primitive qth root of unity and q is the number of minimal associated primes of I.

Let the input ideal I be generated by a finite collection of polynomials  $F_1, \ldots, F_k \in \mathbb{Q}[X]$  with q

minimal associated primes. Write Z := V(I) for the zero locus of I, assumed to be a commutative linear algebraic group. The general procedure of the algorithm is depicted in Figure 1. The ideal  $J_0$ defined in **Line 1** is an ideal of the ring  $\mathbb{Q}[P, X, Q]$ , where the relations  $\{x_{i,j}\}_{i\neq j}$  and  $PQ - \mathrm{Id}_d$ ensure that every point  $(\tilde{P}, \tilde{X}, \tilde{Q}) \in V(J_0)$  comprises a diagonal matrix  $\tilde{X}$  and an invertible matrix  $\tilde{P}$ with  $\tilde{P}^{-1} = \tilde{Q}$  satisfying  $\tilde{P}\tilde{X}\tilde{P}^{-1} \in Z$ . The aim is to find a single such point  $(\tilde{P}, \tilde{X}, \tilde{Q}) \in V(J_0)$ satisfying  $Z = \langle \tilde{P}\tilde{X}\tilde{P}^{-1} \rangle$ . Subsequently, the radical ideal V(J) defined in **Line 2** contains all diagonal conjugates of each matrix in Z. In particular, for each matrix  $M \in Z$  not only one single diagonal matrix D satisfying  $M = \tilde{P}D\tilde{P}^{-1}$  lies in V(J), but all diagonal matrices of the form  $M_{\sigma}DM_{\sigma}^{-1}$ , with the permutation  $\sigma \in S_d$ , also lie in V(J). Due to this fact, we cannot simply employ Proposition 7 to construct a generator for J. Instead, in **Line 4** we isolate a primary component  $\mathcal{P}_0$  of J containing  $\mathrm{Id}_d$ . In the following line, we apply Proposition 7 to the binomial ideal  $\mathcal{P}_0$  and construct a diagonal matrix D such that  $V(\mathcal{P}_0) = \langle D \rangle$ . Since  $V(\mathcal{P}_0)$  is connected the matrix D can be chosen rational, and such that the entries of  $\sqrt[4]{D}$  lie in  $\mathbb{Q}[\zeta_q]$ .

The assertion in **Line 6** verifies whether the orbit of  $D_q$  rotates between the primary components of J; this ensures that  $V(I_q) = \overline{\langle D_q \rangle}$  is included in V(J), where  $I_q$  is defined in **Line 7**. Next, our procedure checks whether J equals to the intersection of  $M_{\sigma}I_qM_{\sigma}^{-1}$ . The necessity of the latter test is due to the above-mentioned fact that V(J) contains all diagonal conjugates of each matrix in Z; see Example 18. The rest of the algorithm is straightforward.

**Example 18.** Let  $F_1 := 2z + w$ ,  $F_2 := 2x - 2y + 3w$ , and  $F_3 := 4y^2 - 4yw + w^2 - 4y + 4w$ . Consider the following ideal as an input to the procedure in Figure 1:

$$I := \langle F_1, F_2, F_3 \rangle \subseteq \mathbb{Q} \begin{bmatrix} \begin{pmatrix} x & z \\ w & y \end{bmatrix}$$

The ideal I is prime (meaning that q = 1) and V(I) is a commutative linear algebraic group. The output of our procedure shows that there exists M such that  $V(I) = \overline{\langle M \rangle}$ , and such that the eigenvalues of M lie in  $\mathbb{Q}$ .

Following the algorithms, the ideal J defined in Line 2 has two primary components

$$\mathcal{P} = \langle w, z, y^2 - x \rangle$$
 and  $\mathcal{P}' = \langle w, z, x^2 - y \rangle$ .

Since  $I_2 \in V(\mathcal{P} \cap \mathcal{P}')$ , we can pick any of these ideals as  $\mathcal{P}_0$  in Line 4. Following Proposition 7 in Line 5, we may construct diagonal matrices  $D = \Delta(4, 2)$  and  $D' = \Delta(2, 4)$  such that

$$V(\mathcal{P}) = \overline{\langle \Delta(4,2) \rangle}$$
 and  $V(\mathcal{P}') = \overline{\langle \Delta(2,4) \rangle}$ 

Clearly, matrices D and D' are conjugates under permutation of diagonals, implying that the assertion in **Line 8** holds. (The above is an indication (1) that permutation of matrices arising from one choice of  $\mathcal{P}_0$  under  $M_{\sigma}$  are suitable for other possible choices of  $\mathcal{P}_0$ , and (2) the necessity of the check in **Line 8**). Following **Line 9** for  $D_q = \Delta(2, 4)$ , defining the ideal

$$J_1 = \langle F_1(QD_qP), F_2(QD_qP), F_3(QD_qP) \rangle \cap \mathbb{Q}[P],$$

we have that  $J_1 = \langle p_3 - 2p_4, p_1 - p_2 \rangle \subseteq \mathbb{Q} \begin{bmatrix} \begin{pmatrix} p_1 & p_2 \\ p_3 & p_4 \end{pmatrix} \end{bmatrix}$ . Subsequently, one choice for a semisimple generator of V(I) is the following matrix M:

$$M = \begin{pmatrix} 6 & 2 \\ -4 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ -2 & -1 \end{pmatrix} \begin{pmatrix} 2 & 0 \\ 0 & 4 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ -2 & -1 \end{pmatrix}^{-1}.$$

#### 5.2 General Generator

We employ the algorithm from the previous subsection to provide a procedure that, given an algebraic set  $Z \subseteq \operatorname{GL}_d(\overline{\mathbb{Q}})$  determines whether there exists a matrix  $M \in \operatorname{GL}_d(\overline{\mathbb{Q}})$  such that  $Z = \overline{\langle M \rangle}$  and which moreover outputs such an M in the affirmative case.

Let the input ideal I be generated by a finite collection of polynomials  $F_1, \ldots, F_k \in \mathbb{Q}[X]$  with q minimal associated primes. Write Z := V(I) for the zero locus of I that is, by assumption, a commutative linear algebraic group. Our algorithm first calls a (modified variant of) the procedure in Figure 1, with the input ideal I, to check whether the subgroup  $G_s$  of all semisimple matrices in Z is one-generated. The modification is as follows: (1) the assertion in **Line 10** is omitted (as this assertion requires that Z is generated with a single semisimple matrix), and (2) the algorithm outputs  $D_q$  and the ideal  $J_1$  defining the locus point of suitable P.

Analogous to the preceding settingwhere the subgroup of semisimple matrices was one-generated, our algorithm proceeds by verifying that the subgroup  $G_u$  of all unipotent matrices is one-generated. For this purpose, it checks

- whether  $V(I + \langle (X \mathrm{Id}_d)^n \rangle)$  is a commutative linear algebraic group; and
- whether  $V(I + \langle (X \mathrm{Id}_d)^n \rangle)$  is one dimensional.

The algorithm returns "no" if either of the subgroups  $G_s$  or  $G_u$  is not one-generated. Otherwise, the procedure defines the ideal  $H \subseteq \mathbb{Q}[P, X, Q]$  by

$$H := \langle F_1(PXQ), \ldots, F_k(PXQ), PQ - \mathrm{Id}_d, \{x_{i,j}\}_{j \neq i, i+1} \rangle$$

where  $P = (p_{i,j})_{1 \le i,j \le d}$ ,  $X = \{x_{i,j}\}_{1 \le i,j \le d}$  and  $Q = (q_{i,j})_{1 \le i,j \le d}$ . It returns "yes" together with the matrix  $\tilde{P}D_q\tilde{X}\tilde{P}^{-1}$  where  $(\tilde{P},\tilde{X}) \in J + H$ .

## References

[Bab+96]	L. Babai, R. Beals, JY. Cai, G. Ivanyos, and E. M. Luks. "Multiplicative Equations over Commuting Matrices". In: <i>Proceedings of the Seventh Annual ACM-SIAM Symposium on Discrete Algorithms, 28-30 January 1996, Atlanta, Georgia.</i> 1996, pp. 498–507.
[Bab85]	L. Babai. "Trading Group Theory for Randomness". In: <i>Proceedings of the 17th Annual ACM Symposium on Theory of Computing</i> . ACM, 1985, pp. 421–429. DOI: 10.1145/22145.22192.
[Ben+19]	R. Benedetto, P. Ingram, R. Jones, M. Manes, J. H. Silverman, and T. J. Tucker. "Current trends and open problems in arithmetic dynamics". In: <i>Bull. Amer. Math.</i> Soc. (N.S.) 56.4 (2019), pp. 611–685. DOI: 10.1090/bull/1665.
[Bey+07]	D. Beyer, T. A. Henzinger, R. Majumdar, and A. Rybalchenko. "Invariant synthesis for combined theories". In: <i>International Workshop on Verification, Model Checking, and Abstract Interpretation</i> . Springer. 2007, pp. 378–394.
[BFS99]	J. F. Buss, G. S. Frandsen, and J. O. Shallit. "The Computational Complexity of Some Problems of Linear Algebra". In: <i>Journal of Computer and System Sciences</i> 58.3 (June 1999), pp. 572–596. DOI: 10.1006/jcss.1998.1608.
[BG06]	E. Bombieri and W. Gubler. Heights in Diophantine Geometry. New Mathematical

Monographs. Cambridge University Press, 2006.

- [Blä+21] M. Bläser, C. Ikenmeyer, V. Lysikov, A. Pandey, and F.-O. Schreyer. "On the orbit closure containment problem and slice rank of tensors". In: *Proceedings of the 2021* ACM-SIAM Symposium on Discrete Algorithms (SODA). SIAM. 2021, pp. 2565–2584.
- [Bor91] A. Borel. *Linear Algebraic Groups*. Springer New York, 1991. DOI: 10.1007/978-1-4612-0941-6.
- [BPR06] S. Basu, R. Pollack, and M.-F. Roy. Algorithms in real algebraic geometry. Second. Vol. 10. Algorithms and Computation in Mathematics. Springer-Verlag, Berlin, 2006, pp. x+662.
- [Bür+11] P. Bürgisser, J. M. Landsberg, L. Manivel, and J. Weyman. "An Overview of Mathematical Issues Arising in the Geometric Complexity Theory Approach to VP≠VNP". In: SIAM J. Comput. 40.4 (2011), pp. 1179–1209. DOI: 10.1137/090765328.
- [Bür+21] P. Bürgisser, M. L. Doğan, V. Makam, M. Walter, and A. Wigderson. "Polynomial Time Algorithms in Invariant Theory for Torus Actions". In: 36th Computational Complexity Conference (CCC 2021). Vol. 200. Leibniz International Proceedings in Informatics (LIPIcs). Dagstuhl, Germany: Schloss Dagstuhl Leibniz-Zentrum für Informatik, 2021, 32:1–32:30. DOI: 10.4230/LIPIcs.CCC.2021.32.
- [Bür24] P. Bürgisser. "Completeness classes in algebraic complexity theory". In: *arXiv preprint* arXiv:2406.06217 (2024).
- [CG84] A. L. Chistov and D. Y. Grigor'Ev. "Complexity of quantifier elimination in the theory of algebraically closed fields". In: International Symposium on Mathematical Foundations of Computer Science. Springer. 1984, pp. 17–31.
- [CK24] J. Cyphert and Z. Kincaid. "Solvable Polynomial Ideals: The Ideal Reflection for Program Analysis". In: Proc. ACM Program. Lang. 8.POPL (2024), pp. 724–752. DOI: 10.1145/3632867.
- [DJK05] H. Derksen, E. Jeandel, and P. Koiran. "Quantum automata and algebraic groups". In: J. Symb. Comput. 39.3-4 (2005), pp. 357–371.
- [DM20] H. Derksen and V. Makam. "Algorithms for orbit closure separation for invariants and semi-invariants of matrices". In: Algebra & Number Theory 14.10 (Nov. 2020), pp. 2791– 2813. DOI: 10.2140/ant.2020.14.2791.
- [Edm67] J. Edmonds. "Systems of distinct representatives and linear algebra". In: Journal of Research of the National Bureau of Standards Section B Mathematics and Mathematical Physics 71B.4 (Oct. 1967), p. 241. DOI: 10.6028/jres.071b.033.
- [FS13] M. A. Forbes and A. Shpilka. "Explicit Noether Normalization for Simultaneous Conjugation via Polynomial Identity Testing". In: Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques. Berlin, Heidelberg: Springer Berlin Heidelberg, 2013, pp. 527–542.
- [Gra17] W. A. de Graaf. Computation with Linear Algebraic Groups. Chapman & Hall/CRC Monographs. CRC Press, 2017.
- [GS] D. R. Grayson and M. E. Stillman. *Macaulay2, a software system for research in alge*braic geometry. Available at http://www2.macaulay2.com.
- [GS21] F. Galuppi and M. Stanojkovski. "Toric varieties from cyclic matrix semigroups". In: *Rend. Istit. Mat. Univ. Trieste* (2021). DOI: 10.13137/2464-8728/33099.

- [HBK20] A. Humenberger, N. S. Bjørner, and L. Kovács. "Algebra-Based Loop Synthesis". In: Integrated Formal Methods - 16th International Conference, IFM 2020, Lugano, Switzerland, November 16-20, 2020, Proceedings. Vol. 12546. Lecture Notes in Computer Science. Springer, 2020, pp. 440–459. DOI: 10.1007/978-3-030-63461-2\\_24.
- [Hit+24] S. Hitarth, G. Kenison, L. Kovács, and A. Varonka. "Linear Loop Synthesis for Quadratic Invariants". In: 41st International Symposium on Theoretical Aspects of Computer Science (STACS 2024). Vol. 289. Leibniz International Proceedings in Informatics (LIPIcs). Dagstuhl, Germany: Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 2024, 41:1– 41:18. DOI: 10.4230/LIPIcs.STACS.2024.41.
- [Hru+18] E. Hrushovski, J. Ouaknine, A. Pouly, and J. Worrell. "Polynomial Invariants for Affine Programs". In: Proceedings of the 33rd Annual ACM/IEEE Symposium on Logic in Computer Science. LICS '18. Oxford, United Kingdom: Association for Computing Machinery, 2018, pp. 530–539. DOI: 10.1145/3209108.3209142.
- [Hru+23] E. Hrushovski, J. Ouaknine, A. Pouly, and J. Worrell. "On strongest algebraic program invariants". In: *Journal of the ACM* 70.5 (2023), pp. 1–22.
- [Hum+22] A. Humenberger, D. Amrollahi, N. Bjørner, and L. Kovács. "Algebra-Based Reasoning for Loop Synthesis". In: Form. Asp. Comput. 34.1 (July 2022). DOI: 10.1145/3527458.
- [Hum75] J. E. Humphreys. Linear Algebraic Groups. Vol. 21. Graduate Texts in Mathematics. Springer, 1975. DOI: 10.1007/978-1-4684-9443-3.
- [IKS10] G. Ivanyos, M. Karpinski, and N. Saxena. "Deterministic Polynomial Time Algorithms for Matrix Completion Problems". In: SIAM Journal on Computing 39.8 (Jan. 2010), pp. 3736–3751. DOI: 10.1137/090781231.
- [Kin+18] Z. Kincaid, J. Cyphert, J. Breck, and T. W. Reps. "Non-linear reasoning for invariant synthesis". In: *Proc. ACM Program. Lang.* 2.POPL (2018), 54:1–54:33. DOI: 10.1145/3158142.
- [KKV23] G. Kenison, L. Kovács, and A. Varonka. "From Polynomial Invariants to Linear Loops". In: Proceedings of the 2023 International Symposium on Symbolic and Algebraic Computation, ISSAC 2023, Tromsø, Norway, July 24-27, 2023. ACM, 2023, pp. 398–406. DOI: 10.1145/3597066.3597109.
- [KL80] R. Kannan and R. J. Lipton. "The Orbit Problem is Decidable". In: Proceedings of the 12th Annual ACM Symposium on Theory of Computing, April 28-30, 1980, Los Angeles, California, USA. ACM, 1980, pp. 252–261. DOI: 10.1145/800141.804673.
- [KL86] R. Kannan and R. J. Lipton. "Polynomial-time algorithm for the orbit problem". In: J. ACM 33.4 (1986), pp. 808–821. DOI: 10.1145/6490.6496.
- [KMM17] L. Katthän, M. Michałek, and E. Miller. "When is a Polynomial Ideal Binomial After an Ambient Automorphism?" In: Foundations of Computational Mathematics 19 (2017). DOI: 10.1007/s10208-018-9405-0.
- [Koi96] P. Koiran. "Hilbert's Nullstellensatz Is in the Polynomial Hierarchy". In: J. Complex.
   12.4 (1996). long version, DIMACS report 96-27, pp. 273–286. DOI: 10.1006/jcom.1996.0019.
- [Lov79] L. Lovász. "On determinants, matchings, and random algorithms". In: Fundamentals of Computation Theory, FCT 1979, Proceedings of the Conference on Algebraic, Arthmetic, and Categorial Methods in Computation Theory, Berlin/Wendisch-Rietz, Germany, September 17-21, 1979. Akademie-Verlag, Berlin, 1979, pp. 565–574.
- [MS04a] M. Müller-Olm and H. Seidl. "A note on Karr's algorithm". In: International Colloquium on Automata, Languages, and Programming. Springer. 2004, pp. 1016–1028.

- [MS04b] M. Müller-Olm and H. Seidl. "Computing polynomial program invariants". In: Inf. Process. Lett. 91.5 (2004), pp. 233–244. DOI: 10.1016/J.IPL.2004.05.004.
- [Nos+22] K. Nosan, A. Pouly, S. Schmitz, M. Shirmohammadi, and J. Worrell. "On the Computation of the Zariski Closure of Finitely Generated Groups of Matrices". In: Proceedings of the 2022 International Symposium on Symbolic and Algebraic Computation. 2022, pp. 129–138.
- [Sil07] J. H. Silverman. The arithmetic of dynamical systems. Vol. 241. Graduate Texts in Mathematics. Springer, New York, 2007, pp. x+511. DOI: 10.1007/978-0-387-69904-2.
- [SMK15] H. Seidl, S. Maneth, and G. Kemper. "Equivalence of Deterministic Top-Down Treeto-String Transducers is Decidable". In: *IEEE 56th Annual Symposium on Foundations* of Computer Science, FOCS 2015, Berkeley, CA, USA, 17-20 October, 2015. IEEE Computer Society, 2015, pp. 943–962. DOI: 10.1109/F0CS.2015.62.
- [SSM04] S. Sankaranarayanan, H. Sipma, and Z. Manna. "Non-linear loop invariant generation using Gröbner bases". In: Proceedings of the 31st ACM SIGPLAN-SIGACT Symposium on Principles of Programming Languages, POPL 2004, Venice, Italy, January 14-16, 2004. ACM, 2004, pp. 318–329. DOI: 10.1145/964001.964028.
- [Val79] L. G. Valiant. "Completeness classes in algebra". In: Proceedings of the Eleventh Annual ACM Symposium on Theory of Computing. STOC '79. Atlanta, Georgia, USA: Association for Computing Machinery, 1979, pp. 249–261. DOI: 10.1145/800135.804419.