

# Loop Quantum Gravity and CMB Anisotropy

S. Davood Sadatian,<sup>a,b,1</sup> Amir Sabouri,<sup>b</sup> Zahra Davari<sup>c</sup>

<sup>a</sup>Department of Physics, Faculty of Basic Sciences, University of Neyshabur, P. O. Box 9319774446, Neyshabur, Iran

<sup>b</sup>Research Department of astronomy & cosmology, University of Neyshabur, P. O. Box 9319774446, Neyshabur, Iran

<sup>c</sup>School of Physics, Korea Institute for Advanced Study (KIAS), 85 Hoegiro, Dongdaemungu, Seoul, 02455, Republic of Korea

E-mail: [sd-sadatian@um.ac.ir](mailto:sd-sadatian@um.ac.ir),  
[sd-sadatian@neyshabur.ac.ir](mailto:sd-sadatian@neyshabur.ac.ir), [asaburi@neyshabur.ac.ir](mailto:asaburi@neyshabur.ac.ir), [zahradavari@kias.re.kr](mailto:zahradavari@kias.re.kr)

**Abstract.** Recent satellite observations have revealed significant anisotropy in the cosmic microwave background (CMB) radiation, a phenomenon that had previously been detected but received limited attention due to its subtlety. With the advent of more precise measurements from satellites, the extent of this anisotropy has become increasingly apparent. This paper examines the CMB radiation by reviewing past research on the causes of CMB anisotropy and presents a new model to explain the observed temperature anisotropy and the anisotropy in the correlation function between temperature and E-mode polarization in the CMB radiation. The proposed model is based on a modified-generalized Compton scattering approach incorporating Loop Quantum Gravity (LQG). We begin by describing the generalized Compton scattering and then discuss the CMB radiation in the context of processes occurring at the last scattering surface. Our findings are derived from the latest observational data from the Planck satellite (2018). In our model, besides the parameters available in the Planck data for the standard model ( $\Lambda$ CDM), we introduce two novel parameters:  $\delta_L$ , the density of cosmic electrons, and  $M^2$ , a parameter related to the modified-generalized Compton scattering effects. The results indicate that, based on the 2018 Planck data, small values were obtained for  $\delta_L$  and  $M^2$ ,  $\delta_L = 1.63 \pm 0.08(10^{-13})$  and  $M^2 = 2.28 \pm 0.34(10^{-4})$ , showing no significant deviation from the standard model. Moreover, increasing the values of  $\delta_L$  and  $M^2$  leads to an increase in the range of fluctuations in the CMB temperature anisotropy power spectrum and the correlation function between temperature and E-mode polarization for multipoles  $l < 500$  until the first peak.

PACS: 04.50.kd;, 95.36.+x

Key Words: Loop Quantum Gravity, Compton Effect, Cosmic Microwave Background.

---

<sup>1</sup>Corresponding author.

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Generalized Compton Scattering (GCS)</b>	<b>2</b>
<b>3</b>	<b>CMB anisotropies at the last scattering surface</b>	<b>3</b>
<b>4</b>	<b>Model Description</b>	<b>5</b>
4.1	CMB temperature anisotropy based on the generalized Compton scattering relation	5
4.2	Anisotropy of the correlation function between temperature and E-mode polarization based on the generalized Compton relation	7
<b>5</b>	<b>Modified-<math>\Lambda</math>CDM verse data</b>	<b>8</b>
<b>6</b>	<b>Conclusion</b>	<b>10</b>
<b>7</b>	<b>Acknowledgments</b>	<b>11</b>
<b>8</b>	<b>Data availability</b>	<b>11</b>

---

## 1 Introduction

Loop Quantum Gravity (LQG) is a theoretical framework that aims to reconcile quantum mechanics and general relativity in a way that is consistent with the standard cosmological model. In LQG, the structure of space-time is composed of entangled loop networks known as spin networks at scales larger than the Planck length ( $l_p \sim 10^{-35}$  m). Due to the inconsistency of experimental data with the quantum aspects of gravity and the lack of a satisfactory candidate, the problem of general relativity merge and quantum mechanics still has major challenges [1]. In 1916, Einstein laid the groundwork for the concept of "quantum Riemannian geometry" to describe the quantization of gravity, which replaces the 4D space-time metric with a probability domain for various space-time geometries [2]. The systematic development of LQG based on quantum Riemannian geometry was completed in the 1990s [3–7], addressing some critical issues in quantum gravity [8, 9].

In the context of LQG cosmology, the structure of space-time does not end at the Big Bang but extends to wider dimensions in the quantum geometry of space-time [10]. One of the promising aspects of quantum gravity is the "asymptotic safety program" proposed within the standard model of particle physics [11]. Since the inception of LQG research by pioneers such as Bergmann [12], Dirac [13], Wheeler [9], and others, the theory has faced persistent challenges over the past two decades, including:

- How to quantize finite Hamiltonian systems without using background fields or perturbation techniques?
- What dynamic effects arise from the absence of a background space-time metric?

- How to precisely calculate quantum transition amplitudes, and how can quantum gravity naturally resolve curvature singularities of classical general relativity?
- Are the UV divergences of quantum field theory (QFT) cured[1]?

Another topic of interest in LQG is the "Compton scattering effect". Compton scattering, a fundamental electromagnetic radiation phenomenon, occurs when a high-energy photon collides with an electron, resulting in a decrease in the photon's energy (frequency) and an increase in its wavelength—a phenomenon known as the "Compton shift" [14]. To describe Compton scattering, light is modeled as a stream of particles (photons) with varying energies or frequencies [15]. Given the low precision of Lorentz symmetry observed in nature, the violation of Lorentz invariance has been examined from multiple perspectives [16–23]. In LQG, the Compton effect can be reformulated based on modified standard dispersion relations, reflecting Lorentz invariance violations in quantum gravity [21, 22]. However, these modified dispersion relations are not mandated by LQG theory itself but are derived heuristically, contingent on solving the problem of quantum gravity and matter dynamics [23].

One of the most valuable applications of the LQG approach is determining the anisotropy of the cosmic microwave background radiation (CMB), which offers cosmologists crucial insights into the early universe's structure and evolution. The first observational data from the COBE satellite revealed anisotropies in CMB photons, especially in terms of temperature and polarization [24]. Subsequent satellites, such as WMAP and Planck, have provided more accurate measurements of CMB anisotropy [25–29].

In this paper, we first review the modified dispersion relations and generalized Compton scattering effects based on LQG in section 2. Section 3 discusses CMB anisotropy stemming from processes at the last scattering surface. In section 4, we present a model incorporating generalized Compton scattering within the LQG framework to improve the accuracy of temperature variation calculations and the correlation function between temperature and E-mode polarization anisotropy of the CMB. Section 5 details the results derived from this model using the latest observational data from Planck [28, 29]. Finally, section 6 concludes the paper.

## 2 Generalized Compton Scattering (GCS)

In this section, we review the modified dispersion relations of the generalized Compton effect from the LQG approach. According to the conservation of linear momentum in Compton scattering ( $\vec{p} = \vec{p}' + \vec{P}$ ), where  $\vec{p}$  and  $\vec{p}'$  are the linear momentum of the photon before and after the collision with an electron at the rest state, respectively, and  $\vec{P}$  is the linear momentum of the electron after the collision with the photon, it can be written:

$$\vec{P} = (\vec{p} - \vec{p}')^2 = \vec{p}^2 + \vec{p}'^2 - 2\vec{p}\vec{p}'. \quad (2.1)$$

Using the conservation of electron energy with the modified dispersion relations from [15], where  $E_{\pm}^2 = A^2 P^2 + \tau P^4 \pm 2P\Gamma + m^2$  and setting  $\hbar = c = 1$ , we get:

$$f - f' = E - E_0 = \sqrt{A^2 P^2 + \tau P^4 \pm 2P\Gamma + m^2} - m, \quad (2.2)$$

$$(f - f' + m)^2 = A^2 P^2 + \tau P^4 \pm 2P\Gamma + m^2, \quad (2.3)$$

$$P^2 = \frac{1}{A^2}[(f - f')^2 + 2(f - f')m - \tau P^4 \mp P\Gamma], \quad (2.4)$$

where  $E_0$  is the initial energy of the electron in the rest state,  $E$  is the final energy of the electron,  $P$  is the momentum of the electron,  $m$  is the mass of the electron,  $\tau = k_2 L_p^2$  and  $\Gamma = \frac{k_3 L_p}{2L^2}$  are two positive parameters dependent on helicity propagation of order 1, where  $k_1$ ,  $k_2$  and  $k_3$  are unknown dimensional parameters and  $A = 1 + \frac{k_1 L_p}{L}$  is a Lorentz invariance violation parameter interpreted as a maximum electron velocity.  $L_p$  is the Planck length, and  $L$  is a new weave scale. Using the relation for the Compton effect ( $\lambda' - \lambda = \frac{h}{mc}(1 - \cos\theta)$ ), it can be written [15]:

$$P^2 = (f - f')^2 + 2ff'(1 - \cos\theta). \quad (2.5)$$

Combining Eqs. 2.4 and 2.5, we get:

$$2ff'(1 - \cos\theta) = \frac{1}{A^2}[2(f - f')m - \tau P^4 \mp P\Gamma] + (\frac{1}{A^2} - 1)(f - f')^2. \quad (2.6)$$

The photon scattering frequency shift can be derived from Eq.2.6. Different modified dispersion relations have been proposed, but a modified dispersion relation proposed in Ref. [23, 30] is:

$$E^2 = P^2 + m^2 + \frac{|P|^{2+n}}{M^n}, \quad (2.7)$$

where  $M^n$  is a characteristic scale of Lorentz invariance violation. Using the Eqs. 2.5 and 2.7, we can write:

$$P^2 = (f - f')^2 + 2m(f - f') - \frac{|P|^{2+n}}{M^n}. \quad (2.8)$$

Now, the generalized modified Compton effect of quantum gravity is then:

$$\lambda' - \lambda = \frac{1}{m}(1 - \cos\theta) + \frac{\lambda\lambda'|P|^{2+n}}{2mM^n}. \quad (2.9)$$

Rewriting Eq. 2.9 in the MKS system of units (with order  $n = 2$ ), gives:

$$\lambda' = \frac{\lambda + \frac{h}{mc}(1 - \cos\theta)}{1 - \frac{\lambda|Pc|^{2+n}}{2h(Mc^2)^n mc^3}}, \quad (2.10)$$

$$\lambda' - \lambda = \frac{h}{mc}(1 - \cos\theta) + \frac{\lambda\lambda'|Pc|^4}{2h(Mc^2)^2 mc^3}. \quad (2.11)$$

In this last equation, the frequency shift caused by the LQG effect depends on the frequency of the incident photon. On the other hand, since the term  $\frac{\lambda|Pc|^{2+n}}{2h(Mc^2)^n mc^3}$  is positive and small, the entire relationship represents a decrease in frequency shift (increase in wavelength shift) [15].

### 3 CMB anisotropies at the last scattering surface

With the increased accuracy of observational satellites, including the Planck satellite, significant temperature and polarization anisotropies in the CMB radiation have been detected [28, 29]. Although many components of the standard cosmological model ( $\Lambda$ CDM) are not well understood in terms of fundamental physics, one solution to this challenge lies the CMB fluctuations and the high measurement accuracy of observational satellites[31].

The CMB is the remnant of the Big Bang's thermal effects, now cooled down to about 2.725K

and exhibiting a perfect blackbody spectrum [31]. CMB temperature anisotropies represent curvature perturbations due to the last scattering surface focused on our current location and spatial fluctuations in the energy density of the CMB. These perturbations in the CMB temperature anisotropy in direction  $\hat{n}$  and time  $t_0$  are generally expressed by the following equation [32]:

$$\theta(\hat{n}) = \theta_0 + \Psi - \hat{n} \cdot v_b + \int_{t_*}^{t_0} (\dot{\Psi} + \dot{\Phi}) dt, \quad (3.1)$$

where  $\theta(\hat{n})$  is fractional anisotropy,  $\theta_0$  represent fractional fluctuations in CMB temperature at the last scattering surface (time  $t_*$ ),  $v_b$  is the baryon peculiar velocity,  $\Psi$ , and  $\Phi$  are gravitational potentials (in general relativity, when  $\Psi = \Phi$  when the non-relativistic matter is dominant). At the last scattering surface on larger scales (such as the Hubble radius), only gravity dominates, while on smaller scales, the acoustic physics of the primordial plasma and photon diffusion are dominated [33].

The main focus of observational CMB research over the past decades has been to obtain accurate estimates of the angular power spectrum of the CMB and relate these values to theoretical models. Research programs like COBE, WMAP, and Planck, along with ground-based telescopes. With the advancements in observational satellite programs, including the latest data from the Planck satellite [28, 29], the parameters of the standard cosmological model have been calculated with increased accuracy. In addition to CMB temperature anisotropy, CMB polarization anisotropy has also been investigated [37].

Polarization is significant in early scattering times because it enables the growth of a quadrupole anisotropy ( $l = 2$ ) around recombination, while scattering is infrequent until the universe is re-ionized after the recombination epoch [38]. The linear polarization expected from the curvature perturbation has an *r.m.s* of  $5\mu\text{K}$  and is described by two Stokes parameters  $Q$ ,  $U$  [31]. Although the Stokes parameters offer a local definition of polarization, their coordinate dependence is inconvenient for cosmological interpretation. Instead, linear polarization is described using the  $E$  and  $B$  fields [39]. The Stokes parameters consist of an orthonormal basis of rank 2 symmetric and a trace-free tensor, which can be expressed in terms of the second derivatives of  $E$  and  $B$ . Generally, the Stokes parameters in Cartesian coordinates, ignoring the sky's curvature, are described in terms of  $E$  and  $B$  fields as:

$$\begin{pmatrix} Q & U \\ U & -Q \end{pmatrix} \propto (\partial_i \partial_j - \frac{1}{2} \delta_{ij} \nabla^2) E + \epsilon_k ({}_i \partial_j) \partial_k B. \quad (3.2)$$

This relation is similar to decomposing a vector field into a gradient part ( $E$ ) and a divergence-free curl part ( $B$ ). E-modes are scalars under parity, while B-modes are pseudo-scalars [31]. In the absence of parity-violation physics, the two fields should not be correlated, leaving three non-zero polarization power spectra: the  $C_{B_l}$  B-mode power spectrum,  $C_{E_l}$  E-mode power spectrum,  $C_{C_l}$  correlation power spectrum between E-mode and temperature anisotropy; of course, a CMB temperature anisotropy power spectrum,  $C_{T_l}$ , is also presented. Several important points regarding CMB polarization anisotropy include [31]:

1. Polarization is a small signal.
2. E-mode polarization peaks at scales smaller than temperature because it relies on diffusion in small-scale modes for its generation.
3. The acoustic peaks in  $C_{E_l}$  are significant because the temperature quadrupole mainly derives from the bulk plasma velocity, which vanishes when the density is at an extremum.

4. There is a bump in large-scale polarization caused by re-scattering once the universe re-ionizes.
5. Symmetrically, B-mode polarizations are not generated by curvature perturbations except through second-order processes like gravitational lensing. This characteristic makes B-mode polarizations a potential probe for gravitational waves.

Gravitational waves with wavelength smaller than the Hubble radius are damped away due to universe's expansion. In this case, CMB is the best option for detecting gravitational waves, as, it is sensitive to early times after the last scattering surface and large scales [40]. Gravitational waves induce CMB temperature anisotropies due to anisotropy expansion effects along the line of sight [41].

## 4 Model Description

In this section, we present a model that investigates the temperature and E-mode polarization of the CMB anisotropies based on generalized Compton scattering relations within the LQG framework. The first subsection analyzes CMB temperature anisotropy using generalized Compton scattering relations, while the second subsection proposes the anisotropy of the correlation function between temperature and E-mode polarization based on these relations.

### 4.1 CMB temperature anisotropy based on the generalized Compton scattering relation

To study the temporal evolution of CMB anisotropies, the Boltzmann equations are particularly useful, especially when considering the effects of Compton scattering and collision terms. The Boltzmann equation in the synchronous gauge for temperature anisotropy is defined as [42, 43]:

$$\dot{\Delta}_T^{(S)} + ik\mu\Delta_T^{(S)} = -\frac{1}{6}\dot{h} - \frac{1}{6}(\dot{h} + 6\dot{\eta})P_2(\mu) + \dot{\kappa} \left[ -\Delta_T^{(S)} + \Delta_{T_0}^{(S)} + i\mu v_b + \frac{1}{2}P_2(\mu)\Pi \right], \quad (4.1)$$

where the derivatives are taken with respect to the conformal time  $\tau$ .  $\Delta_T^{(S)}(\tau, k, \mu) = T^{(S)}$  is the temperature anisotropy of the CMB and the superscript  $S$  represents the primordial scalar perturbation in the Fourier modes specified by the wavevector  $\mu = \hat{n} \cdot \vec{k} = \cos\theta$ . Here,  $\hat{n}$  represents the direction of photons,  $\vec{k}$  is the wavevector,  $\theta$  represents the angle between the CMB photon direction  $\hat{n}$  and the wavevector  $\vec{k}$ .  $\tau$  is conformal time.  $P_2(\mu)$  represents Legendre polynomial of order-2 ( $P_2(\mu) = \frac{1}{2}(3\mu^2 - 1)$ ).  $\dot{\kappa}$  is the differential optical depth for Compton scattering, given by:

$$\dot{\kappa} = a(\tau)n_e x_e \sigma_T, \quad (4.2)$$

where  $a(\tau)$  is the scale factor in terms of conformal time,  $n_e$  is the number density of free electron,  $x_e$  is the ionization fraction, and  $\sigma_T$  is the Thomson cross section. By integrating over conformal time from Eq.4.2, the total optical depth due to the Thomson scattering is obtained as:

$$\kappa(\tau) = n_e x_e \sigma_T \int_{\tau_r}^{\tau_0} a(\tau) d\tau, \quad (4.3)$$

where  $\tau_0$  is the present time and  $\tau_r$  corresponds to the conformal time of the recombination epoch. The sources in Eq. 4.1 include CMB temperature multipoles defined as  $\Delta_T^{(S)}(k, \mu) =$

$\sum_l (2l+1)(-i)^l \Delta_l(k) P_l(\mu)$ , where  $P_l(\mu)$  is the Legendre polynomial of order-  $l$ .  $v_b$  is the baryon velocity,  $\dot{h}$  and  $\dot{\eta}$  are the additional source terms due to the metric perturbation.  $\Pi$  is the source term due to Compton scattering, defined as [42, 43]:

$$\Pi = \Delta_{T2}^{(S)} + \Delta_{P2}^{(S)} + \Delta_{P0}^{(S)}, \quad (4.4)$$

where  $\Delta_{P0}^{(S)}$  and  $\Delta_{P2}^{(S)}$  are the source terms due to polarization anisotropy of order 0 and 2, respectively.

Now, Eq.4.1 is modified based on the generalized Compton scattering (Eq.2.10 with order  $n = 2$ ) of the CMB photon interacting with a resting electron and is contributed a term to Eq.4.1 as the generalized Compton scattering effects:

$$\begin{aligned} \dot{\Delta}_T^{(S)} + ik\mu\Delta_T^{(S)} = & -\frac{1}{6}\dot{h} - \frac{1}{6}(\dot{h} + 6\dot{\eta})P_2(\mu) + \dot{\kappa}[-\Delta_T^{(S)} + \Delta_{T0}^{(S)} + i\mu v_b + \frac{1}{2}P_2(\mu)\Pi] \\ & + [i\kappa^* \frac{2}{3}\Delta_{T2}^{(S)}(1 - \mu^2) + \frac{i+1}{3}(1 - \mu^2)\kappa^* \Delta_{T1}(\hat{n})]M^{n=2}, \end{aligned} \quad (4.5)$$

where  $\Delta_{T1}(\hat{n})$  is the source due to dipole asymmetry resulting from CMB temperature anisotropy, defined as:

$$\Delta_{T1}(\hat{n}) = \overline{\Delta_{T1}}(\hat{n})(1 + A\hat{p}\cdot\hat{n}). \quad (4.6)$$

$\overline{\Delta_{T1}}(\hat{n})$  is the isotropic part of the temperature fluctuations,  $\hat{n}$  and  $\hat{p}$  are the line of sight (direction of the observation) and the preferred direction respectively and, A is the amplitude of the asymmetry.  $M^{n=2}$  from Eq.2.10 and  $\lambda \propto \frac{1}{f}(\lambda = \frac{V}{f})$  is rewritten as:

$$M^2 = \frac{V^2 P^4}{2hc^2(mcV(f - f') - hf f'(1 - \cos\theta))} \quad (4.7)$$

and  $\kappa^*$  is the optical depth associated with the generalized Compton scattering, defined as [44]:

$$\kappa^* = \frac{3}{2} \frac{m_e v_e}{k^0} \sigma_T \delta_L n_e, \quad (4.8)$$

where  $k^0 = 2.7$  Kelvin,  $m_e$  electron mass,  $v_e \sim \frac{1}{\sqrt{1+z}} 10^{-3}$  electron bulk flow velocity,  $\delta_L$  left-handed cosmic electrons density of azimuthal symmetry (and  $\delta_R$  right-handed cosmic electrons density of azimuthal symmetry) which is assumed in all calculations and this symmetry:  $\delta_L = \delta_R$ . For complete solving Eq.4.5, we first assume the anisotropies until the present epoch and integrate from all over the Fourier modes:

$$T^{(S)}(\hat{n}) = \int d^3k \xi(k) \Delta_T^{(S)}(\tau_0, k, \mu), \quad (4.9)$$

where  $\xi(k)$  is a random variable that represents the initial amplitude of the mode. To obtain the power spectrum as an indicator for CMB anisotropies from Eq.4.5, we take the integral along the line of sight [41]:

$$\Delta_T^{(S)}(\tau_0, k, \mu) = \int_0^{\tau_0} d\tau e^{ix\mu} S_T^S(k, \tau), \quad (4.10)$$

that

$$S_T^S(k, \tau) = g \underbrace{\left( \Delta_{T0} + 2\dot{\alpha} + \frac{\dot{v}_b}{k} + \frac{\Pi}{4} + \frac{3\ddot{\Pi}}{4k^2} \right)}_{S_{T1}^S} + e^{-\kappa}(\dot{\eta} + \ddot{\alpha}) + \dot{g} \left( \alpha + \frac{v_b}{k} + \frac{3\dot{\Pi}}{4k^2} \right) + \frac{3\ddot{g}\Pi}{4k^2} +$$

$$\underbrace{\left( i\frac{2}{3}\Delta_{T2}^{(S)}(1-\mu^2) + \frac{i+1}{3}(1-\mu^2)\Delta_{T1}(\hat{n}) \right)}_{S_{T2}^S} \kappa^* M^2. \quad (4.11)$$

$S_{T1}^S$  of Eq.4.11 is the power spectrum resulting from the standard cosmological scenario, and  $S_{T2}^S$  is the power spectrum resulting from the generalized Compton scattering effect.  $x = k(\tau_0 - \tau)$  and  $\alpha = \frac{\hbar+6\dot{\eta}}{2k^2}$ . We have introduced the visibility function  $g(\tau) = \kappa \exp(-\kappa)$ . Now, the power spectrum of temperature anisotropy as a rotational invariance quantity is defined as  $C_{Tl} = \frac{1}{2l+1} \sum_l |a_{lm}|^2$ ,

$$C_{Tl} = \frac{1}{2l+1} \sum_l \langle a_{T,l,m}^* a_{T,l,m} \rangle, \quad (4.12)$$

in terms of which,

$$\langle a_{T,l,m}^* a_{T,l,m} \rangle = C_{Tl} \delta_{l'l} \delta_{m'm}. \quad (4.13)$$

Here,  $a_{lm}$  represents the amplitude of CMB temperature fluctuations in the presence of scalar perturbation, given by [45]:

$$a_{T,l,m} = \int d\Omega Y_{lm}^*(\hat{n}) T^{(S)}(\hat{n}), \quad (4.14)$$

where  $\Delta_T$  is the CMB temperature anisotropy obtained from Eqs.4.10, 4.11 and  $Y_{lm}^*(\hat{n})$  is the spherical harmonic function. Finally, the CMB temperature anisotropy power spectrum obtained from Eqs.4.10 and 4.11 is equal to:

$$\begin{aligned} C_{Tl} &= \frac{1}{2l+1} \int d^3k \phi_k \sum_l \left| \int_0^{\tau_0} d\Omega Y_{lm}^*(\hat{n}) \Delta_T^{(S)}(\tau_0, k, \mu) \right|^2 \\ &= 16\pi^2 \int k^2 \phi_k dk \left[ \left( S_{T1}^{(S)}(\tau_0, k, \mu) j_l(x) \right)^2 + \left( S_{T2}^{(S)}(\tau_0, k, \mu) \frac{j_l(x)}{x^2} \right)^2 \right], \end{aligned} \quad (4.15)$$

where  $\phi_k$  is the initial power spectrum ( $\phi_k = pn_s - 1$  where  $p$  is the photon momentum),  $j_l(x)$  is the spherical Bessel function of order- $l$  and assuming the framework  $k \parallel \hat{z}$ ,  $\int d\Omega Y_{lm}^*(\hat{n}) e^{ix\mu} = \sqrt{4\pi(2l+1)} i^l j_l(x) \delta_{m0}$ .

## 4.2 Anisotropy of the correlation function between temperature and E-mode polarization based on the generalized Compton relation

The correlation function between temperature and E-mode polarization is established, while the correlation between temperature and B-mode or E-mode and B-mode vanishes because B-mode has the opposite parity to  $T$  and  $E$  [46]. This CMB correlation function is defined as follows [45]:

$$C(\theta) = \sum_l \frac{2l+1}{4\pi} C_{Cl}^{(S)} P_l \cos(\theta), \quad (4.16)$$

where  $P_l$  is the Legendre polynomial of order  $l$  and  $\theta$  is an angle scale. The angular power spectra are in terms of multipole  $l$  values, and the correlation function is in terms of the angular scale  $\theta$ . The relationship  $\theta \propto \frac{1}{l}$  indicates that large multipole values correspond to

small angular scales [48]. Generally, the correlation power spectrum between  $T$  and  $E$  is defined as [46]:

$$C_{C_l} = \frac{1}{2l+1} \sum_m \langle a_{T,lm}^*, a_{E,lm} \rangle. \quad (4.17)$$

In the presence of scalar perturbation for the correlation function, we have:

$$\Delta_{C_l}^{(S)} = \int_0^{\tau_0} \dot{\kappa} e^{ix\mu - \kappa} \Pi_\mu^2 (1 - \mu^2)^2 \kappa^* \Delta_{T_l}^{(S)}(\tau_0, k, \mu) \Delta_{E_l}^{(S)}(\tau_0, k, \mu). \quad (4.18)$$

Finally, the power spectrum of the correlation function in the presence of scalar perturbation is written as:

$$C_{C_l}^{(S)} = 16\pi^2 \int k^2 dk \phi_k \Delta_{T_l}^{(S)}(\tau_0, k, \mu) \Delta_{E_l}^{(S)}(\tau_0, k, \mu). \quad (4.19)$$

## 5 Modified- $\Lambda$ CDM verse data

To check the anisotropy observed in CMB radiation based on a modified-generalized Compton scattering with loop quantum gravity approach, we implement the related equations in the publicly available numerical code `CLASS`<sup>1</sup> (the Cosmic Linear Anisotropy Solving System) [47]. In module `perturbation.c` of `CLASS`, the temperature source function is divided into three distinct parts:

$$S_T = S_T^0 + \frac{d}{d\tau} S_T^1 + \frac{d}{d\tau} \dot{S}_T^2, \quad (5.1)$$

where the dot indicates that the code evaluates the analytic expression for the derivative, while the derivatives denoted by  $\frac{d}{d\tau}$  are estimated numerically. Therefore, it is useful to provide the explicit expressions for all three terms in the Newtonian gauge as


$$\begin{aligned} S_T^0 &= g\left(\frac{1}{4}\delta_\gamma + \phi\right) + 2e^{-\kappa}\dot{\phi} + \frac{1}{k^2}g\kappa\dot{\theta}_b + e^{-\kappa}\ddot{\kappa}\theta_b + e^{-\kappa}\dot{\kappa}\dot{\theta}_b, \\ S_T^1 &= e^{-\kappa}k(\psi - \phi), \\ S_T^2 &= \frac{1}{8}g\Pi. \end{aligned} \quad (5.2)$$

We add the modified terms to  $S_T^0$  and  $S_T^2$ .

In addition to the six parameters of the standard  $\Lambda$ CDM model, namely baryon density  $\Omega_b h^2$ , dark matter density  $\Omega_{DM} h^2$ , the ratio between the acoustic scale and the angular diameter distance at decoupling  $\theta_s$ , optical depth of reionization  $\tau_{reio}$ , the amplitude  $A_s$ , and the spectral index  $n_s$ , two additional parameters  $M^2$  and  $\delta_L$ , are included for the applied modifications. The value of  $\delta_L$ , the density caused by cosmic electrons, often referred to as the optical depth to reionization ( $\tau$ ), is reported to be of the order of  $0.054 \pm 0.007$  in the Planck 2018 data [28]. The optical depth to reionization is a crucial parameter in cosmology, as it quantifies the integrated number of free electrons along the line of sight back to the epoch of reionization, affecting the amplitude of the large-scale E-mode polarization in the cosmic microwave background. The value of  $\delta_L$  obtained, of order  $10^{-9}$ , corresponds to the reported value of the optical depth to reionization by the Planck 2018 release.

Another free parameter in the study is  $M^2$ . It is crucial to accurately estimate the prior values of the  $M^2$  parameter due to its significant impact on the results. Given the absence of the

<sup>1</sup>[https://github.com/lesgourg/class\\_public](https://github.com/lesgourg/class_public)



**Figure 1.** Left panels: Temperature anisotropies in the CMB. Right panels: Anisotropy of the correlation function between temperature and E-mode polarization in the CMB.

established prior information for the  $M^2$  parameter, an exploratory approach was adopted to test various potential values. We considered different orders of magnitude for  $M^2$ , as shown in the bottom of Figure 1, and finally considered its prior in the interval  $[0, 0.1]$ . In the panels of Figure 1, we observe an increase in the amplitude of the lower multipoles in the temperature and the correlation function power spectrum until the first peak. The integrated Sachs-Wolfe (ISW) effect is important on such scales, and it is possible that the modified-generalized Compton scattering affects this signal.

To obtain the exact values of  $\delta_L$  and  $M^2$ , we use the code `MONTEPYTHON-v32` [49, 50] to perform a Monte Carlo Markov Chain (MCMC) analysis with a Metropolis-Hasting algorithm. We use the CMB temperature and polarization auto- and cross-correlation measurements of the most recent Planck 2018 legacy release, including the full temperature power spectrum at multipoles  $2 \leq l \leq 2500$  and the polarization power spectra in the range  $2 \leq l \leq 29$ . We also include information on the gravitational lensing power spectrum estimated from the CMB trispectrum analysis [29]. We obtain the following values using all CMB and CMB lensing information:


$$M^2 = 2.28 \pm 0.34(10^{-4}),$$

$$\delta_L = 1.63 \pm 0.08(10^{-13}).$$


We show posterior distributions ( $1\sigma$  and  $2\sigma$  intervals) of these parameters in Figure 2. Given the small values obtained for these parameters, we do not expect to observe any deviation in anisotropy observed in CMB radiation compared to the standard model, as seen, for example, in Figure 3 for temperature anisotropies and TE cross correlation CMB angular power spectra.

To select the most compatible model with the observational data and assess the quality of the

<sup>2</sup>[https://github.com/baudren/montepython\\_public](https://github.com/baudren/montepython_public)



**Figure 2.** 1D likelihoods and 2D contours in 68% and 95% CL marginalized joint regions for chosen free parameters:  $\delta_L$  and  $M^2$ .



**Figure 3.** The temperature and TE cross correlation CMB angular power spectra using the best fit values of cosmological parameters. The red curve in the plots below indicates the deviation from the standard model.

fit, we use the least squares method, commonly applied in cosmology, known as the total chi-squared,  $\chi_{\text{tot}}^2$ . A model with a smaller  $\chi_{\text{tot}}^2$  is generally considered a better fit to the data [51]. In our analysis, the  $\chi_{\text{tot}}^2$  values for the standard model and the Modified- $\Lambda$ CDM model are 1389.47 and 1388.30, respectively. However, a lower  $\chi_{\text{tot}}^2$  does not necessarily indicate the best model, as models with more parameters can artificially lower  $\chi_{\text{tot}}^2$ . To address this, we employ the Akaike Information Criterion (AIC), defined as  $AIC = \chi_{\text{min}}^2 + 2M$ , where  $M$  is the number of free parameters in the model. According to [52],  $\Delta AIC$  values greater than 2, 5, and 10 indicate weak, moderate, and strong evidence respectively, against the model with the higher  $AIC$  value. In this study, we consider two additional parameters compared to the standard model, and since  $|\Delta AIC| = 2.18$ , the two models fit the cosmological data equally well.

## 6 Conclusion

In this study, we explored the modified-generalized Compton scattering relations within the framework of loop quantum gravity and analyzed the anisotropies in the cosmic microwave

background (CMB) radiation's temperature and polarization, influenced by processes originating from the last scattering surface.

We developed a model to account for temperature anisotropy and the anisotropy in the correlation function between temperature and E-mode polarization. This model integrates the modified-generalized Compton scattering relations and leverages the latest Planck satellite data (2018), incorporating two additional parameters:  $\delta_L$  (cosmic electron density) and  $M^2$  (effects of modified-generalized Compton scattering).

Our findings indicate that smaller values of  $\delta_L$  and  $M^2$  yield results that closely align with the standard cosmological model, corroborating the effects of modified-generalized Compton scattering with the 2018 Planck data. As depicted in Figure 1, increasing the values of  $\delta_L$  and  $M^2$  enhances the fluctuation range in the CMB temperature anisotropy power spectrum and the correlation function between temperature and E-mode polarization for multipoles  $l < 500$  up to the first peak. Conversely, for multipoles  $l > 500$ , variations in these parameters do not significantly impact the results.

We anticipate that future observations with improved precision from new missions or experiments will enable the determination of more accurate values for  $\delta_L$  and  $M^2$ . Such enhanced accuracy may reveal substantial deviations from the standard model. In future research, we aim to further examine the implications of modified-generalized Compton scattering on both E-mode and B-mode polarization, potentially providing deeper insights into the fundamental structure and evolution of the universe.

## 7 Acknowledgments

Z.D. was supported by the Korea Institute for Advanced Study Grant No 6G097301. S.D. Sadatian and A. Sabouri appreciate R. Mohammadi for helpful discussions and useful advices during this study.

## 8 Data availability

No new data were generated or analysed in support of this research.

## Declarations Conflict of interest

The authors declare that they have no conflict of interest.

## References

- [1] Ashtekar, A., & Bianchi, E: *A short review of loop quantum gravity*, *Reports on Progress in Physics* **84**(4) (2021) 042001.
- [2] Einstein, A: *Zur quantentheorie der strahlung*, First published in (1916) 121-128.
- [3] Ashtekar, A., & Lewandowski, J., *Background independent quantum gravity: a status report* *Classical and Quantum Gravity*, **21**(15) (2004) R53.
- [4] Rovelli, C., *Quantum gravity*, Cambridge university press (2004).
- [5] Rovelli, C., & Vidotto, F., *Covariant loop quantum gravity: an elementary introduction to quantum gravity and spinfoam theory* Cambridge University Press (2015).
- [6] Thiemann, T., *Modern canonical quantum general relativity*, Cambridge University Press (2008).

- [7] Giesel, K., *Loop Quantum Gravity: The First 30 Years* (2017).
- [8] Dirac, P. A. M., *Lectures on quantum mechanics* (Vol.2) Courier Corporation (2001).
- [9] Wheeler, J. A., *in Relativity, Groups and Topology ed. B S DeWitt and C M DeWitt* NewYork: Gordon Breach (1964).
- [10] Agullo,I. and sigh,P., *Loop quantum cosmology*, chapter 6 in the volume cited in [7].
- [11] Eichhorn, A., *Asymptotically safe gravity* arXiv preprint arXiv:2003.00044 (2020).
- [12] Bergmann, P. G., *Non-linear field theories*, *Physical Review* **75**(4) (1949) 680.
- [13] Dirac, P. A. M., *Generalized hamiltonian dynamics*, *Canadian journal of mathematics*, **2** (1950) 129-148.
- [14] Gasiorowicz, S., *Quantum physics*, John Wiley & Sons (2007).
- [15] Nozari, K., & Sadatian, S. D., *Loop quantum gravity modification of the Compton effect*, *General Relativity and Gravitation*, **40** (2008) 23-33.
- [16] Amelino-Camelia, G., *Relativity in spacetimes with short-distance structure governed by an observer-independent (Planckian) length scale*, *International Journal of Modern Physics D*, **11**(01) (2002) 35-59.
- [17] Magueijo, J., & Smolin, L., *Lorentz invariance with an invariant energy scale*, *Physical Review Letters*, **88**(19) (2002) 190403.
- [18] Nozari, K., & Sefidgar, A. S., *Comparison of approaches to quantum correction of black hole thermodynamics*, *Physics Letters B*, **635**(2-3) (2006) 156-160.
- [19] Amelino-Camelia, G., Lämmerzahl, C., Macias, A., & Müller, H., *The search for quantum gravity signals*, *In AIP Conference Proceedings* **758**(1) (2005) 30-80.
- [20] Andrianov, A. A., Soldati, R., & Sorbo, L., *Dynamical Lorentz symmetry breaking from a (3+1)-dimensional axion-Wess-Zumino model*, *Physical Review D* **59**(2) (1998) 025002.
- [21] Alfaro, J., & Palma, G., *Loop quantum gravity corrections and cosmic ray decays*, *Physical Review D*, **65**(10), (2002) 103516.
- [22] Alfaro, J., & Palma, G., *Loop quantum gravity and ultrahigh energy cosmic rays*, *Physical Review D*, **67**(8), (2003) 083003.
- [23] Sadatian, S. D., & Nozari, K., *Crossing of the phantom divided barrier with Lorentz invariance violating fields*, *Europhysics Letters*, **82**(4) (2008) 49001.
- [24] Smoot, G. F., Bennett, C. L., Kogut, A., Wright, E. L., Aymon, J., Boggess, N. W., ... & Wilkinson, D. T., *Structure in the COBE differential microwave radiometer first-year maps*, *Astrophysical Journal* **396**(1) (1992) L1-L5.
- [25] Spergel, D. N., Verde, L., Peiris, H. V., Komatsu, E., Nolta, M. R., Bennett, C. L., ... & Wright, E. L.; *First-year Wilkinson Microwave Anisotropy Probe (WMAP)\* observations: determination of cosmological parameters*, *The Astrophysical Journal Supplement Series*, **148**(1) (2003) 175.
- [26] Ade, P. A., Aghanim, N., Armitage-Caplan, C., Arnaud, M., Ashdown, M., Atrio-Barandela, F., ... & Mazzotta, P., *Planck 2013 results. XV. CMB power spectra and likelihood*, *Astronomy & Astrophysics*, **571** (2014) A15.
- [27] Ade, P. A., Aghanim, N., Armitage-Caplan, C., Arnaud, M., Ashdown, M., Atrio-Barandela, F., ... & Meinhold, P. R., *Planck 2013 results. XVI. Cosmological parameters*, *Astronomy & Astrophysics*, **571** (2014) A16.
- [28] Aghanim, N., Akrami, Y., Ashdown, M., Aumont, J., Baccigalupi, C., Ballardini, M., ... & Spencer, L. D., *Planck 2018 results-V. CMB power spectra and likelihoods*, *Astronomy & Astrophysics*, **641** (2020) A5.

- [29] Aghanim, N., Akrami, Y., Ashdown, M., Aumont, J., Baccigalupi, C., Ballardini, M., ... & Roudier, G., *Planck 2018 results-VI. Cosmological parameters, Astronomy & Astrophysics*, **641** (2020) A6.
- [30] Carmona, J. M., & Cortés, J. L., *Infrared and ultraviolet cutoffs of quantum field theory, Physical Review D*, **65**(2) (2001) 025006.
- [31] Challinor, A., *CMB anisotropy science: a review, Proceedings of the International Astronomical Union*, **8**(S288), (2012) 42-52.
- [32] Sugiyama, N., *Introduction to temperature anisotropies of Cosmic Microwave Background radiation, Progress of Theoretical and Experimental Physics*, **2014**(6) (2014) 06B101.
- [33] Wands, D., Piattella, O. F., & Casarini, L., *Physics of the cosmic microwave background radiation. In The Cosmic Microwave Background, Proceedings of the II José Plínio Baptista School of Cosmology* Springer International Publishing (2016).
- [34] Larson, D., Dunkley, J., Hinshaw, G., Komatsu, E., Nolta, M. R., Bennett, C. L., ... & Wright, E. L., *Seven-year wilkinson microwave anisotropy probe (WMAP\*) observations: power spectra and WMAP-derived parameters, The Astrophysical Journal Supplement Series*, **192**(2) (2011) 16.
- [35] Keisler, R., Reichardt, C. L., Aird, K. A., Benson, B. A., Bleem, L. E., Carlstrom, J. E., ... & Zahn, O., *A measurement of the damping tail of the cosmic microwave background power spectrum with the South Pole Telescope, The Astrophysical Journal*, **743**(1) (2011) 28.
- [36] Piacentini, F., Ade, P. A., Bock, J. J., Bond, J. R., Borrill, J., Boscaleri, A., ... & Vittorio, N., *A measurement of the polarization-temperature angular cross-power spectrum of the cosmic microwave background from the 2003 flight of boomerang, The Astrophysical Journal*, **647**(2) (2006) 833.
- [37] Hu, W., *CMB temperature and polarization anisotropy fundamentals, Annals of Physics*, **303**(1) (2003) 203-225.
- [38] Subramanian, K., *The physics of CMBR anisotropies, Current Science*, (2005) 1068-1087.
- [39] Seljak, U., & Zaldarriaga, M., *Signature of gravity waves in the polarization of the microwave background, Physical Review Letters*, **78**(11) (1997) 2054.
- [40] Conti, L., & Saliwanchik, B. R., *CMB Experiments and Gravitational Waves*, In Handbook of Gravitational Wave Astronomy Singapore: Springer Nature Singapore (2022).
- [41] Seljak, U., & Zaldarriaga, M., *A line of sight approach to cosmic microwave background anisotropies*, arXiv preprint astro-ph/9603033 (1996).
- [42] Bond, J. R., & Efstathiou, G., *The statistics of cosmic background radiation fluctuations, Monthly Notices of the Royal Astronomical Society*, **226**(3) (1987) 655-687.
- [43] Ma, C. P., & Bertschinger, E., *Cosmological perturbation theory in the synchronous and conformal Newtonian gauges, Astrophys. J.*, **455** (1995) 7.
- [44] Khodagholizadeh, J., Mohammadi, R., Sadegh, M., & Vahedi, A., *B-mode power spectrum of CMB via polarized Compton scattering, Journal of Cosmology and Astroparticle Physics*, **2020**(01) (2020) 051.
- [45] Albrecht, J., *Temperature anisotropy of the cmb effects of cosmological paramters* Doctoral dissertation, Master's thesis, University Bielefeld (2015).
- [46] Zaldarriaga, M., & Seljak, U., *All-sky analysis of polarization in the microwave background, Physical Review D*, **55**(4) (1997) 1830.
- [47] Lesgourgues, Julien and Tram, Thomas, *The Cosmic Linear Anisotropy Solving System (CLASS) IV: efficient implementation of non-cold relics, JCAP*, **09**,(2011)032.

- [48] Hu, W., & Dodelson, S., *Cosmic microwave background anisotropies*, *Annual Review of Astronomy and Astrophysics*, **40**(1) (2002) 171-216.
- [49] , Brinckmann, Thejs and Lesgourgues, Julien, *MontePython 3: boosted MCMC sampler and other features*, *Phys. Dark Univ*, **24**, (2019)100260.
- [50] Audren, Benjamin and Lesgourgues, Julien and Benabed, Karim and Prunet, Simon, *Conservative Constraints on Early Cosmology: an illustration of the Monte Python cosmological parameter inference code*, *JCAP*, **02**, (2013) 001.
- [51] Davari, Zahra and Rahvar, Sohrab, *MOG cosmology without dark matter and the cosmological constant*, *Mon. Not. Roy. Astron. Soc.*, **507**, (2021)3387-3399.
- [52] Trotta, Roberto, *Bayes in the sky: Bayesian inference and model selection in cosmology*, *Contemp. Phys*, **49**, (2008) 71-104.