CONNECTION PROBLEM FOR SPECIAL FUNCTION SOLUTIONS OF PAINLEVÉ III EQUATION

HAO PAN

Department of Mathematics, University of Michigan, East Hall, 530 Church St., Ann Arbor, MI 48109. Email: haopan@umich.edu

ANDREI PROKHOROV

Department of Mathematics, University of Michigan, East Hall, 530 Church St., Ann Arbor, MI 48109, USA and

St. Petersburg University, 7/9 Universitetskaya nab. 199034 St. Petersburg, Russia. Email: andreip@umich.edu

ABSTRACT. In this paper we compute the small and large x asymptotics of the special function solutions of Painlevé-III equation. We use the representation of solution in terms of Toeplitz determinants of Bessel functions, which is new. Toeplitz determinants are rewritten as multiple contour integrals using Andrèief identity. Finally small and large x asymptotics is obtained using elementary asymptotic methods. The claimed result has not appeared in the literature before. Our formulas are useful for numerical computations of corresponding solutions of Painlevé-III equation.

Key words and phrases: Toeplitz determinant, Painlevé equations, Bessel function, asymptotic analysis, integrable systems, ordinary differential equations

CONTENTS

1. Introduction	2
1.1. Overview of the paper	4
2. Construction of Bessel function solutions of Painlevé III equation	5
2.1. The simultaneous solutions of Ricatti and Painlevé III equations	5
2.2. Bäcklund transformation	7
3. Toeplitz determinants of cylinder functions	7
3.1. Hamiltonian system	7
3.2. Tau function and Toda equation	8
3.3. Wronskian solutions of Toda equation	9
3.4. Proof of Proposition 1.1	11
4. Asymptotics of Toeplitz determinant at zero	13
4.1. Andrèief identity	13
4.2. Basic strategies	14
4.3. Expanded formula for $\Delta_n(x, \alpha)$	14
4.4. Proof of Theorem 1.1	16
5. Asymptotics of special function solutions at zero	19
5.1. Piecewise function for the power of x in the asymptotic of $u_n(x, \alpha)$.	19
5.2. Proof of Theorem 1.2	20
6. Asymptotics at infinity	22
6.1. Asymptotics of Toeplitz determinant	22
6.2. Asymptotics of special function solutions	24
Acknowledgment	24
References	24
Appendix A. Bessel equation and its solutions	25
A.1. Contour integral representations	26
A.2. Differential identities for cylinder functions	27

1. INTRODUCTION

Painlevé equations are six nonlinear second-order ordinary differential equations. They are written in the form of u'' = R(u', u, t) with R a rational function. Their solutions have the so called Painlevé property. It means that the locations of singularities of branching type at the complex plane don't depend on the initial conditions, but the locations of isolated singularities might depend on initial conditions. They were discovered at the beginning of the 20st century in the works [Pai02; Gam10], see also [Inc44]. Most of the solutions of Painlevé equations are transcendental. It means that they can't be reduced to simpler classical special functions. However, there are several exceptions. In particular for Painlevé III equation one can find solutions expressed in terms of Bessel functions (see [UW98; Mur95]). For the applications of such special function solutions of Painlevé III equation in random matrix theory we refer the reader to [FW02], [ZCL20].

We start with presenting Painlevé III equation

$$u'' = \frac{(u')^2}{u} - \frac{u'}{x} + \frac{\alpha u^2 + \beta}{x} + u^3 - \frac{1}{u}, \quad \alpha, \beta \in \mathbb{C}.$$
 (1)

Consider the Toeplitz determinant of cylinder functions:

$$\Delta_n(x,\alpha) = \det\left(\left\{\mathcal{C}_{\frac{\alpha}{2}-j+k}(x)\right\}_{j,k=0}^{n-1}\right), \quad n \in \mathbb{N}$$
(2)

with

$$C_{\nu}(x) = d_1 J_{\nu}(x) + d_2 Y_{\nu}(x), \quad d_1, d_2 \in \mathbb{R}$$
 (3)

and $J_{\nu}(x)$, $Y_{\nu}(x)$ are Bessel functions of first and second kind. Additionally, denote $\Delta_0(x, \alpha) = 1$.

Proposition 1.1. The expression

$$u_n(x,\alpha) = -\frac{\Delta_{n+1}(x,\alpha-2)\Delta_n(x,\alpha)}{\Delta_{n+1}(x,\alpha)\Delta_n(x,\alpha-2)}, \quad n \in \mathbb{N} \cup \{0\}, \quad \alpha \in \mathbb{C}.$$
(4)

solves Painlevé III equation with shifted parameters

$$u_n'' = \frac{(u_n')^2}{u_n} - \frac{u_n'}{x} + \frac{(\alpha + 2n)u_n^2 + (-\alpha + 2 + 2n)}{x} + u_n^3 - \frac{1}{u_n}.$$
(5)

The fact that Toeplitz determinants of Bessel functions are related to the solutions of Painlevé III equation is well known, see for example [Cla23], [FW02]. But the formula of type (4) was not presented in literature. Similar formula for the case of rational solutions can be found at [CLL23]. Now we are ready to present the first result of our asymptotic analysis.

Theorem 1.1. The Toeplitz determinant (2) admits the following $x \to 0$ asymptotics for fixed $d_1, d_2 \in \mathbb{R}$, $n \in \mathbb{N} \cup \{0\}, \alpha \in \mathbb{C} \setminus (2\mathbb{Z} + i\mathbb{R})$

(1) If
$$Re(\alpha) > 2n - 2$$
 or $d_1 \sin\left(\frac{\pi\alpha}{2}\right) + d_2 \cos\left(\frac{\pi\alpha}{2}\right) = 0$, then

$$\Delta_n(x,\alpha) \sim (-1)^{\frac{n(n+1)}{2}} \left(\frac{d_2}{\pi}\right)^n \frac{G(n+1)G(\frac{\alpha}{2}+1)}{G(\frac{\alpha}{2}-n+1)} \left(\frac{x}{2}\right)^{-\frac{n\alpha}{2}}, \text{ as } x \to 0.$$

(2) If
$$2n - 4j - 2 < Re(\alpha) < 2n - 4j + 2$$
 for some $j = 1, 2..., n - 1$, then

$$\Delta_n(x,\alpha) \sim (-1)^{\frac{n(n-1)}{2} + nj + n-j} \left(\frac{d_2}{\pi}\right)^n \left(\frac{d_1}{d_2} \sin\left(\frac{\pi\alpha}{2}\right) + \cos\left(\frac{\pi\alpha}{2}\right)\right)^j \\ \times \frac{G(n-j+1)G(-\frac{\alpha}{2} + n-j+1)G(j+1)G(j+\frac{\alpha}{2} + 1)}{G(-\frac{\alpha}{2} + n-2j+1)G(\frac{\alpha}{2} - n+2j+1)} \left(\frac{x}{2}\right)^{(\alpha-2n+2j)j-\frac{n\alpha}{2}}, \ as \ x \to 0.$$

(3) If
$$Re(\alpha) < -2n+2$$
 or $d_2 = 0$, then

$$\Delta_n(x,\alpha) \sim \frac{(-1)^{\frac{n(n+1)}{2}}}{\pi^n} \left(d_1 \sin\left(\frac{\pi\alpha}{2}\right) + d_2 \cos\left(\frac{\pi\alpha}{2}\right) \right)^n \frac{G(n+1)G(-\frac{\alpha}{2}+1)}{G(-\frac{\alpha}{2}-n+1)} \left(\frac{x}{2}\right)^{\frac{n\alpha}{2}}, \ as \ x \to 0$$

where G(x) refers to the Barnes G-function.

For discussion of the necessity of condition $\alpha \in \mathbb{C} \setminus (2\mathbb{Z} + i\mathbb{R})$ see Remarks 4.2, A.1.

The asymptotic formulas is obtained after rewriting the Toeplitz determinant as multiple contour integral using Andrèief identity and performing elementary asymptotic analysis. We should mention that the same strategy was applied to special function solutions of Painlevé-II equation in [Dea18] and to Hankel determinant solutions of Painlevé-VI in [CZ10].

We should mention that plugging naively the asymptotic of Bessel function to the formula (4) and trying to derive Theorem 1.1 is a difficult task. We allow the possibility of combinatorial proof following this route, but it would have to reproduce different leading behaviors for different α and the mechanism for it is unclear to us.

Combining the Theorem 1.1 with Proposition 1.1 we derive the asymptotics of $u_n(x, \alpha)$ as $x \to 0$.

Theorem 1.2. Solution (4) of Painlevé-III equation (5) admits the following $x \to 0$ asymptotics for fixed $d_1, d_2 \in \mathbb{R}, n \in \mathbb{N} \cup \{0\}, \alpha \in \mathbb{C} \setminus (2\mathbb{Z} + i\mathbb{R})$

(1) If
$$Re(\alpha) > 2 + 2n$$
 or $d_1 \sin\left(\frac{\pi\alpha}{2}\right) + d_2 \cos\left(\frac{\pi\alpha}{2}\right) = 0$, then

$$u_n(x,\alpha) \sim \left(\frac{2}{2n+2-\alpha}\right) \frac{x}{2}, \text{ as } x \to 0.$$

(2) If $2n - 4j < Re(\alpha) < 2n - 4j + 2$ and j = 0, 1, ..., n, then

$$u_n(x,\alpha) \sim (-1)^n \left(\frac{d_1}{d_2} \sin\left(\frac{\pi\alpha}{2}\right) + \cos\left(\frac{\pi\alpha}{2}\right)\right) \left(\frac{\Gamma(-\frac{\alpha}{2} + n - 2j + 1)}{\Gamma(\frac{\alpha}{2} - n + 2j)}\right)^2 \\ \times \frac{\Gamma(j + \frac{\alpha}{2})\Gamma(j + 1)}{\Gamma(-\frac{\alpha}{2} + n - j + 1)\Gamma(n - j + 1)} \left(\frac{x}{2}\right)^{\alpha - 2n + 4j - 1}, \ as \ x \to 0.$$

(3) If
$$2n - 4j - 2 < Re(\alpha) < 2n - 4j$$
 and $j = 0, 1, \dots, n - 1$, then

$$u_n(x,\alpha) \sim (-1)^n \left(\frac{d_1}{d_2} \sin\left(\frac{\pi\alpha}{2}\right) + \cos\left(\frac{\pi\alpha}{2}\right)\right)^{-1} \left(\frac{\Gamma(\frac{\alpha}{2} - n + 2j + 1)}{\Gamma(-\frac{\alpha}{2} + n - 2j)}\right)^2 \times \frac{\Gamma(-\frac{\alpha}{2} + n - j + 1)\Gamma(n - j)}{\Gamma(j + \frac{\alpha}{2} + 1)\Gamma(j + 1)} \left(\frac{x}{2}\right)^{-\alpha + 2n - 4j - 1}, \ as \ x \to 0.$$

(4) If $Re(\alpha) < -2n$ or $d_2 = 0$, then

$$u_n(x,\alpha) \sim \left(-\frac{\alpha}{2} - n\right) \left(\frac{x}{2}\right)^{-1}, \ as \ x \to 0.$$

where $\Gamma(x)$ refers to the Gamma function.



FIGURE 1. Leading power in the asymptotics for n = 5 as function of α . The piecewise expressions for $p_c(\alpha, n)$ and $e(\alpha, n)$ can be found in (56) and (58) respectively.

We can observe that leading power of asymptotics in Theorems 1.1, 1.2 is continuous as function of α , see illustration for n = 5 on Figure 1.

We can compare our result to the small x asymptotics computed based on the monodromy data in [BLMP24], see also [Jim82; Kit87]. We have to make change of variables $x \to 2x$ to match with the scaling of Painlevé-III equation considered in [BLMP24]. As the result we get $e_1^2 = e_{\infty}^2 = e_0^2 =$ $(-1)^n e^{\frac{i\pi\alpha}{2}}$ which formally was not included in the considered monodromy data. That means that we are filling the gap in the literature regarding the asymptotics of solutions of Painlevé-III equation. Next we present our results for the large x asymptotics.

Theorem 1.3. The Toeplitz determinant (2) admits the following $x \to \infty$ asymptotics for fixed $d_1, d_2 \in \mathbb{R}$, $n \in \mathbb{N} \cup \{0\}$, $\alpha \in \mathbb{C}$.

(1) If n is even, then

$$\Delta_n(x,\alpha) \sim \left(\frac{d_1^2 + d_2^2}{2\pi}\right)^{\frac{n}{2}} \left(G\left(\frac{n}{2} + 1\right)\right)^2 \left(\frac{x}{4}\right)^{-\frac{n^2}{4}}, \ as \ x \to \infty$$

(2) If n is odd, then

$$\Delta_n(x,\alpha) \sim (-1)^{\frac{n-1}{2}} \left(\frac{d_1^2 + d_2^2}{2\pi}\right)^{\frac{n}{2}} G\left(\frac{n+1}{2}\right) G\left(\frac{n+3}{2}\right) \sin\left(x - \phi + \frac{\pi}{4}(n-\alpha)\right) \left(\frac{x}{4}\right)^{-\frac{n^2+1}{4}}, as \ x \to \infty.$$

where G(x) refers to the Barnes G-function and $\phi = \arg(d_1 + id_2)$.

Theorem 1.4. Solution (4) of Painlevé-III equation (5) admits the following $x \to \infty$ asymptotics for fixed $d_1, d_2 \in \mathbb{R}, n \in \mathbb{N} \cup \{0\}, \alpha \in \mathbb{C}$

(1) If n is even, then

$$u_n(x,\alpha) \sim -\cot\left(x-\phi+\frac{\pi}{4}(n+1-\alpha)\right), \ as \ x \to \infty.$$

(2) If n is odd, then

$$u_n(x,\alpha) \sim -\tan\left(x-\phi+\frac{\pi}{4}(n-\alpha)\right), \text{ as } x \to \infty$$

We should mention that it is a matter of change of variables to include the case of Toeplitz determinants of modified Bessel functions which corresponds to parameters of Painlevé-III equation $\alpha - \beta \in 2 + 4\mathbb{N}$, see Remark 2.2.

Proposition 1.1 is useful for the numerical computation of solution $u_n(x, \alpha)$. We illustrate Theorems 1.1, 1.2, 1.3, 1.4 on Figure 2. We also present color plot for argument of $u_n(x, \alpha)$ in the complex plane for various choices of α and n on Figure 3. We present the result fixed α and large n in Figure 3b. We observe that the pole structure is similar to the pole structure of rational solutions of Painlevé III equation observed in [BMS18], but the poles now lie also in the regions extending to infinity. The special case with $d_2 = 0$ can be found on Figure 3c. The other case of large α and large n can be seen on Figure 3d. We see that the pole structure looks different in this case, which is expected. Finally on Figure 3e we can see what happens when we take α much larger than n. The analysis of such pictures would require tools like nonlinear steepest descent method for Riemann-Hilbert problems.

1.1. **Overview of the paper.** We start with the reminder of construction of a special function solution using Ricatti equation in section 2.1 and generation of the family of special function solutions using Bäcklund transformation in section 2.2.

We prove representation for the special function solutions from Proposition 1.1 in section 3. We start by introducing the tau function and demonstrating the classical fact that it satisfies the Toda equation in section 3.2. We use Deshanot-Jacobi identity and Toda equation to show the classical Wronskian formula for the tau function associated to the family of special function solutions in section 3.3. The crucial next step is identification of Toeplitz determinant (2) with Painlevé tau function (19) in Proposition 3.5, following [FW02], [Oka87]. The main tools are differential identities (70), (71). After long tedious computation we finish the proof of Proposition 1.1.

We prove Theorem 1.1 in section 4. We start with rewriting Toeplitz determinant as multiple contour integral using Andrèief formula in section 4.1. The key next step in the proof is splitting of the multiple contour integral in the sum of other multiple contour integrals, such that it is easy to compute the leading term of the asymptotics for the latter integrals. The result can be found in Lemma 4.1. We determine which integral has the largest leading term in Lemma 4.2. The last step in the proof of Theorem 1.1 is the evaluation of the multiple contour integral corresponding to the leading term using formulas from [DLMF].

We prove Theorem 1.2 in section 5. It consists of plugging in of result of Theorem 1.1 in the Proposition 1.1 and tedious manipulation with piecewise formulas.

We prove Theorem 1.3 in section 6.1. We rewrite the cylinder functions in terms of Hankel functions, which are more convenient for large x asymptotics computation. We apply Andrèief identity and apply





(C) Plot of the expression $x^{0.08}u_n(x,\alpha)$ and its small (D) Plot of the solution $u_n(x,\alpha)$ and its large x asymptotics totics.

FIGURE 2. Illustration for n = 10, $\alpha = 1.08$, $d_1 = 0.55$, $d_2 = 0.71$. Blue plots are the solutions or the determinants, yellow plots are the asymptotics.

steepest descent method to determine final result. The Theorem 1.4 is obtained in section 6.2 by combining Theorem 1.3 and Proposition 1.1.

In appendix A we derive the convenient contour integral representations (64), (69) for the cylinder function (3) and present differential identities (70), (71).

2. Construction of Bessel function solutions of Painlevé III equation

2.1. The simultaneous solutions of Ricatti and Painlevé III equations. The standard way to construct the special function solutions of Painlevé equations is to use Ricatti equation, see [Cla23, Theorem 3.5] and [DLMF, §32.10(iii)]. More precisely, we look for the simultaneous solutions of Painlevé III equation (1) and the Ricatti equation

$$u'(x) = a(x)u^{2}(x) + b(x)u(x) + c(x), \quad a(x) \neq 0.$$
(6)

Taking the first derivative of (6) and plugging in the u'(x), we get:

$$u''(x) = 2a^{2}(x)u^{3}(x) + (a'(x) + 3a(x)b(x))u^{2}(x) + (2a(x)c(x) + b^{2}(x) + b'(x))u(x) + (b(x)c(x) + c'(x))u(x) + (b(x)c(x) + c'(x))u(x)$$

Meanwhile, plugging (6) into (1), we get:

$$\begin{split} u''(x) = & (a^2(x)+1)u^3(x) + \left((2a(x)b(x) - \frac{a(x) - \alpha}{x} \right) u^2(x) + \left(b^2(x) + 2a(x)c(x) - \frac{b(x)}{x} \right) u(x) \\ & + \frac{c(x) - 1}{u(x)} + 2b(x)c(x) - \frac{c(x) - \beta}{x}. \end{split}$$



(A) Phase plot for the case f(z) = z.



(B) Complex phase plot of solution $u_n(x, \alpha)$, for n = 10 and $\alpha = 1.08$, $d_1 = 0.55$, $d_2 = 0.71$.

6

(C) Complex phase plot of solution $u_n(x, \alpha)$, for n = 10 and $\alpha = 1.08$, $d_1 = 0.55$, $d_2 = 0$.



(D) Complex phase plot of solution $u_n(x, \alpha)$, for n = (E) Complex phase plot of solution $u_n(x, \alpha)$, for n = 10 10 and $\alpha = 23.04$, $d_1 = 0.55$, $d_2 = 0.71$. and $\alpha = 80.04$, $d_1 = 0.55$, $d_2 = 0.71$.

FIGURE 3. Complex argument plots of solutions for various values of parameters.

By matching and solving for the coefficients, we have totally four cases. We list them below:

$$a(x) = 1,$$
 $b(x) = \frac{\alpha - 1}{x},$ $c(x) = 1,$ $\beta = 2 - \alpha$ (7)

$$a(x) = -1,$$
 $b(x) = \frac{-1 - \alpha}{x},$ $c(x) = -1,$ $\beta = -2 - \alpha$ (8)

$$a(x) = 1,$$
 $b(x) = \frac{\alpha - 1}{x},$ $c(x) = -1,$ $\beta = \alpha - 2$ (9)

$$a(x) = -1,$$
 $b(x) = \frac{-1 - \alpha}{x},$ $c(x) = 1,$ $\beta = \alpha + 2$ (10)

Notice that if u(x) solves the Ricatti equation, then $w(x) = \exp\left(-\int a(x)u(x)dx\right)$ solves the following linear ODE:

$$a(x)w''(x) - (a'(x) + a(x)b(x))w'(x) + c(x)a^2(x)w(x) = 0, \quad a(x) \neq 0.$$
(11)

From now on we will only focus on the case (7). The equation (11) becomes

$$w''(x) + \frac{(1-\alpha)}{x}w'(x) + w(x) = 0.$$
(12)

We can notice that $x^{-\frac{\alpha}{2}}w(x)$ solves Bessel equation (61) with $\nu = \frac{\alpha}{2}$. For $\alpha \in \mathbb{C} \setminus (2\mathbb{Z} + i\mathbb{R})$ we denote the solution of (12)

$$w(x,\alpha) = x^{\frac{\alpha}{2}} \mathcal{C}_{\frac{\alpha}{2}}(x) \tag{13}$$

where cylinder function $C_{\nu}(x)$ is given by (3). Here we assume that $\frac{\alpha}{2}$ is not integer for convenience in our future computations. As the result we get

Proposition 2.1 ([DLMF, §32.10(iii)]). Painlevé III equation (1) with $\beta = 2 - \alpha$ and $\alpha \in \mathbb{C} \setminus (2\mathbb{Z} + i\mathbb{R})$ admits the special function solution

$$u_0(x,\alpha) = -\frac{d}{dx}\ln(w(x,\alpha)).$$
(14)

with $w(x, \alpha)$ given by (13).

Remark 2.1. In the case (8) solution is also given in term of Bessel functions, while in the cases (9), (10) it is given in terms of modified Bessel functions.

2.2. Bäcklund transformation. To construct more solutions for PIII equation with more general parameters, we need to introduce a powerful tool. Bäcklund transformations for the Painlevé-III equation are given by (see [DLMF, §32.7(iii)])

$$B_1: (u(x), \alpha, \beta) \to \left(\frac{xu'(x) + xu^2(x) - \beta u(x) - u(x) + x}{u(x)(xu'(x) + xu^2(x) + \alpha u(x) + u(x) + x)}, \alpha + 2, \beta + 2\right)$$
(15)

$$B_2: (u(x), \alpha, \beta) \to \left(-\frac{xu'(x) - xu^2(x) - \beta u(x) - u(x) + x}{u(x)(xu'(x) - xu^2(x) - \alpha u(x) + u(x) + x)}, \alpha - 2, \beta + 2\right)$$
(16)

Assume that u(x) solves Painlevé-III equation. Denote $B_1(u(x), \alpha, \beta) = (W(x), \alpha + 2, \beta + 2)$. One can show that W(x) solves Painlevé-III equation

$$W''(x) = \frac{(W'(x))^2}{W(x)} - \frac{W'(x)}{x} + \frac{(\alpha+2)W^2(x) + (\beta+2)}{x} + W^3(x) - \frac{1}{W(x)}.$$

Similarly, denote $B_2(u(x), \alpha, \beta) = (W(x), \alpha - 2, \beta + 2)$. One can show that W(x) solves Painlevé-III equation

$$W''(x) = \frac{(W'(x))^2}{W(x)} - \frac{W'(x)}{x} + \frac{(\alpha - 2)W^2(x) + (\beta + 2)}{x} + W^3(x) - \frac{1}{W(x)}$$

Proposition 2.2 ([DLMF, §32.7(iii)]). Denote $B_1^n(u_0(x, \alpha), \alpha, 2 - \alpha) = (u_n(x, \alpha), \alpha + 2n, -\alpha + 2 + 2n)$ with $u_0(x, \alpha)$ given by (14). Then $u_n(x, \alpha)$ is the special function solution of Painlevé III equation (5).

We can observe that parameters of Painlevé-III equation (5) parameters satisfy $\alpha + \beta \in 2 + 4\mathbb{N}$. We will use Bäcklund transformation B_2 in our future constructions.

Remark 2.2. Using the transformation $u(x) \to -u(x)$ we can get solutions with $\alpha + \beta \in -2 - 4\mathbb{N}$. Using the transformation $u(x) \to -iu(-ix)$ we can get solutions with $\alpha - \beta \in 2 + 4\mathbb{N}$.

3. Toeplitz determinants of cylinder functions

3.1. Hamiltonian system. We use the formulas presented in [Cla23].

Definition 3.1. We define the momentum associated to the solution of Painlevé-III equation using formula

$$v(x) = \frac{1}{2u^2(x)} \left(xu'(x) + xu^2(x) - x + u(x) \left(\beta - 1\right) \right)$$

Definition 3.2. We define the Hamiltonian associated to the solution of Painlevé-III equation using formula

$$H(x) = v^{2}(x)u^{2}(x) - v(x)\left(xu^{2}(x) - x + u(x)(\beta - 1)\right) + 2xu(x)\left(\frac{\beta - (2 + \alpha)}{4}\right)$$

One can show that Painlevé-III equation is equivalent to the Hamiltonian system:

$$x\frac{du}{dx} = \frac{\partial H}{\partial v},$$
$$x\frac{dv}{dx} = -\frac{\partial H}{\partial u}$$

3.2. Tau function and Toda equation. For details of this section see [Oka87] and [FW02].

Definition 3.3. We define the auxiliary Hamiltonian associated to the solution of Painlevé-III equation using formula

$$h(x) = \frac{1}{2} \left(H(x) + u(x)v(x) - x^2 + \frac{1}{4}(\beta - 4)(\beta + (\alpha - 2)) \right).$$

In this section we will deal with generic solution u(x). Since momentum, Hamiltonian and auxiliary Hamiltonian are expressed in terms of u(x), the action of Bäcklund transformation B_1 can be extended to them by formulas $(u(x), \alpha, \beta)$ to v(x), H(x) and h(x). We denote

$$(u_n(x), \alpha + 2n, \beta + 2n) = B_1^n(u(x), \alpha, \beta)$$

$$v_n(x) = v(x)|_{u(x) \to u_n(x), \beta \to \beta + 2n}$$

$$H_n(x) = H(x)|_{u(x) \to u_n(x), v(x) \to v_n(x), \alpha \to \alpha + 2n, \beta \to \beta + 2n}$$

$$h_n(x) = h(x)|_{H(x) \to H_n(x), u(x) \to u_n(x), v(x) \to v_n(x), \alpha \to \alpha + 2n, \beta \to \beta + 2n}$$
(18)

On the path to derive the representation of Proposition 1.1 we introduce the tau function associated with the solution u(x).

Definition 3.4. Define tau function associated to the solution of Painlevé-III equation using formula

$$x\frac{d}{dx}\ln(\tau_n(x)) = h_n(x).$$
(19)

It is defined up to a multiplicative constant.

Proposition 3.1 ([FW02, Proposition 4.2]). Tau function for Painlevé-III equation given by (19) satisfies Toda equation

$$x\frac{d}{dx}x\frac{d}{dx}\ln(\tau_n(x)) = c_n\frac{\tau_{n+1}(x)\tau_{n-1}(x)}{\tau_n^2(x)}$$

for some constants c_n . Moreover, one can choose multiplicative constants in the definition (19) in such a way that $c_n = 1$.

Proof. Using the Bäcklund transformation B_1 we can check the identity

$$h_{n+1}(x) = h_n(x) - v_n u_n(x) - \frac{3}{2} + \frac{\alpha}{4} - \frac{3\beta}{4} + 2n$$
(20)

Denote $A_n(x) = x \frac{d}{dx} x \frac{d}{dx} \ln(\tau_n(x))$ and $B_n(x) = \frac{\tau_{n+1}(x)\tau_{n-1}(x)}{\tau_n^2(x)}$. We want to show that $A_n(x) = c_n B_n(x)$. Taking natural log on both sides, we get $\ln A_n(x) = \ln B_n(x) + \ln c_n$. Therefore, it is sufficient to show

$$\frac{d}{dx}(\ln A_n(x) - \ln B_n(x)) = 0.$$
(21)

Well, by using Definition 3.4 and the identity (20), we have:

$$\frac{d}{dx}\ln(B_n(x)) = \frac{v_{n-1}(x)u_{n-1}(x) - v_n(x)u_n(x) + 2}{x}$$
(22)

$$\frac{d}{dx}\ln(A_n(x)) = \frac{h'_n(x) + xh''_n(x)}{xh'_n(x)}$$
(23)

By using Definition 3.3 and (17)–(18), we rewrite (23), (22) in terms of $u_n(x)$. After a long computation we obtain (21).

Let's show that using transformation $\tau_n(x) \to a_n \tau_n(x)$ one can make sure that the constant c_n in the Toda equation is 1. We notice that a_n has to satisfy a difference equation

$$c_n a_n^2 = a_{n+1} a_{n-1}.$$

Its general solution is given by

$$a_n = \frac{a_1^n}{a_0^{n-1}} \prod_{j=1}^{n-1} \prod_{i=1}^j c_i, \quad n \in \mathbb{N}$$

We can pick the initial conditions $a_0 = a_1 = 1$ and choose

$$a_n = \prod_{j=1}^{n-1} \prod_{i=1}^j c_i, \quad n \in \mathbb{N}.$$

3.3. Wronskian solutions of Toda equation. Toda equation determines the tau function recursively given initial conditions. If we want to derive some nice formula for it, we need some properties of determinants. The Leibniz formula for the determinant of $n \times n$ matrix $A = \{a_{ij}\}_{i,j=1}^{n}$ is given by

$$\det(A) = \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) \prod_{k=1}^n a_{k,\sigma(k)}$$

where S_n is the set of permutations of *n* elements and $sgn(\sigma)$ is sign of permutation σ . Directly using the Leibniz formula above, we can show the following formulas for the derivative of a determinant

$$\frac{d}{dx}\det(A) = \sum_{j=1}^{n} \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) \left(\frac{d}{dx} a_{j,\sigma(j)}\right) \prod_{\substack{k=1\\k \neq j}}^{n} a_{k,\sigma(k)}$$
(24)

Remembering that $det(A^T) = det(A)$ we can write the alternative formula

$$\frac{d}{dx}\det(A) = \sum_{j=1}^{n} \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) \left(\frac{d}{dx} a_{\sigma(j),j}\right) \prod_{\substack{k=1\\k \neq j}}^{n} a_{\sigma(k),k}.$$
(25)

Denote $A_{i|j}$ the matrix obtained from A by deleting of *i*th row and *j*th column. The Laplace expansion for the determinant along the *j*th row is given by

$$\det(A) = \sum_{k=1}^{n} (-1)^{k+j} a_{jk} \det(A_{j|k})$$

Proposition 3.2 (see [VV23]). Denote $A_{ij|kl}$ the matrix obtained from A by deleting of ith and jth row and kth and lth column. Determinants of these matrices satisfy Deshanot-Jacobi identity

$$\det(A) \det(A_{ij|ij}) = \det(A_{i|i}) \det(A_{j|j}) - \det(A_{i|j}) \det(A_{j|i}), \quad 1 \le i, j \le n.$$
(26)

Proposition 3.3 ([FW02, (2.43)]). The sequence of functions

$$f_n(x) = \det\left(\left\{\left(x\frac{d}{dx}\right)^{i+j}f_0(x)\right\}_{i,j=0}^n\right)$$
(27)

solves Toda equation corresponding to Painlevé-III equation

$$\left(x\frac{d}{dx}\right)^{2}\ln(f_{n}(x)) = \frac{f_{n+1}(x)f_{n-1}(x)}{f_{n}^{2}(x)}, \quad n \ge 1.$$
(28)

Proof. Specifically, to match the expression in Proposition 3.2, we rewrite (28) as:

$$f_{n-1}(x)f_{n+1}(x) = f_n(x)\left(x\frac{d}{dx}\right)^2 f_n(x) - \left(x\frac{d}{dx}f_n(x)\right)^2, \quad n \ge 1$$

Put $f_{n+1}(x) = \det(A)$. It follows that $f_n(x) = \det(A_{n+2|n+2})$ and $f_{n-1}(x) = \det(A_{n+1,n+2|n+1,n+2})$. We take the first derivative of the determinant in (27) by multi-linearity with respect to rows using (24). Since a determinant with two identical rows is zero, we end up with $x \frac{d}{dx} f_n(x) = \det(A_{n+1|n+2})$. Since $A_{n+1|n+2} = (A_{n+2|n+1})^T$, it implies that $x \frac{d}{dx} f_n(x) = \det(A_{n+1|n+2}) = \det(A_{n+2|n+1})$. Then we take the second derivative of the determinant in (27) successively by multi-linearity with respect to

columns using (25). Similarly, since a determinant with two identical columns is zero, we end up with $\left(x\frac{d}{dx}\right)^2 f_n(x) = \det(A_{n+1|n+1})$. Using (26) we obtain (28).

Now let's return to the special function solutions $u_n(x, \alpha)$. We compute corresponding auxiliary Hamiltonians $h_0(x, \alpha)$, $h_1(x, \alpha)$, and $h_2(x, \alpha)$. It turns out that corresponding tau functions can be chosen as

$$\tau_0(x,\alpha) = 1,\tag{29}$$

$$\tau_1(x,\alpha) = \mathcal{C}_{\frac{\alpha}{2}}(x),\tag{30}$$

$$\tau_2(x,\alpha) = \det \begin{pmatrix} \tau_1(x,\alpha) & x\frac{d}{dx}\tau_1(x,\alpha) \\ x\frac{d}{dx}\tau_1(x,\alpha) & \left(x\frac{d}{dx}\right)^2\tau_1(x,\alpha) \end{pmatrix}.$$
(31)

where $C_{\nu}(x)$ is given by (3). Toda equation determines tau function uniquely given the initial condition. The solution with initial conditions (30), (31) is given by (27) by Proposition 3.3. As the result we get

Proposition 3.4. Tau functions associated to the special function solutions $u_n(x, \alpha)$ can be chosen as

$$\tau_n(x,\alpha) = \det\left(\left\{\left(x\frac{d}{dx}\right)^{i+j-2}\tau_1(x,\alpha)\right\}_{i,j=1}^n\right)$$
(32)

with $\tau_1(x,\alpha) = \mathcal{C}_{\frac{\alpha}{2}}(x), \ \tau_0(x,\alpha) = 1.$

Proposition 3.5. Formula (32) can be alternatively written as

$$\tau_n(x,\alpha) = x^{n(n-1)}(-1)^{\frac{n(n-1)}{2}} \Delta_n(x,\alpha).$$

where $\Delta_n(x, \alpha)$ is given by (2).

Proof. Using mathematical induction and identity (70) one can compute the structure of $\left(x\frac{d}{dx}\right)^{j} C_{\frac{\alpha}{2}}(x)$:

$$\left(x\frac{d}{dx}\right)^{j} \mathcal{C}_{\frac{\alpha}{2}}(x) = x^{j} \mathcal{C}_{\frac{\alpha}{2}-j}(x) + \sum_{k=0}^{j-1} c_{kj} x^{k} \mathcal{C}_{\frac{\alpha}{2}-k}(x)$$
(33)

where c_{kj} are constant coefficients. Well, we furtherly simplify the determinant using (33):

$$\det\left(\left\{\left(x\frac{d}{dx}\right)^{k+j}\mathcal{C}_{\frac{\alpha}{2}}(x)\right\}_{k,j=0}^{n-1}\right)$$

$$=\begin{vmatrix} \mathcal{C}_{\frac{\alpha}{2}}(x) & x\frac{d}{dx}\mathcal{C}_{\frac{\alpha}{2}-1}(x) + c_{01}\mathcal{C}_{\frac{\alpha}{2}}(x) & \cdots \\ x\mathcal{C}_{\frac{\alpha}{2}-1}(x) + c_{01}\mathcal{C}_{\frac{\alpha}{2}}(x) & x\frac{d}{dx}\left(x\mathcal{C}_{\frac{\alpha}{2}-1}(x) + c_{01}\mathcal{C}_{\frac{\alpha}{2}}(x)\right) & \cdots \\ x^{2}\mathcal{C}_{\frac{\alpha}{2}-2}(x) + c_{12}x\mathcal{C}_{\frac{\alpha}{2}-1}(x) + c_{02}\mathcal{C}_{\frac{\alpha}{2}}(x) & x\frac{d}{dx}\left(x^{2}\mathcal{C}_{\frac{\alpha}{2}-2}(x) + c_{12}\mathcal{C}_{\frac{\alpha}{2}-1} + c_{02}\mathcal{C}_{\frac{\alpha}{2}}(x)\right) & \cdots \\ \vdots & \vdots & \ddots \end{cases}$$

Observe that by elementary row operations, we can always use the previous rows to eliminate the $\sum_{k=0}^{j-1} c_{kj} x^k C_{\frac{\alpha}{2}-k}(x)$ part in a fixed row and the value of the determinant doesn't change. Finally, we

will end up with:

$$\det\left(\left\{\left(x\frac{d}{dx}\right)^{k+j}\mathcal{C}_{\frac{\alpha}{2}}(x)\right\}_{k,j=0}^{n-1}\right)$$

$$=\begin{vmatrix} \mathcal{C}_{\frac{\alpha}{2}}(x) & x\frac{d}{dx}\mathcal{C}_{\frac{\alpha}{2}}(x) & \cdots \\ x\mathcal{C}_{\frac{\alpha}{2}-1}(x) & x\frac{d}{dx}(x\mathcal{C}_{\frac{\alpha}{2}-1}(x)) & \cdots \\ x^{2}\mathcal{C}_{\frac{\alpha}{2}-2}(x) & x\frac{d}{dx}(x^{2}\mathcal{C}_{\frac{\alpha}{2}-2}(x)) & \cdots \\ \vdots & \vdots & \ddots \end{vmatrix} = \det\left(\left\{\left(x\frac{d}{dx}\right)^{k}x^{j}\mathcal{C}_{\frac{\alpha}{2}-j}\right\}_{k,j=0}^{n-1}\right)$$

By relation (71), by induction, we can show:

$$\left(x\frac{d}{dx}\right)^{k}x^{j}\mathcal{C}_{\frac{\alpha}{2}-j}(x) = (-1)^{k}x^{j+k}\mathcal{C}_{\frac{\alpha}{2}-j+k}(x) + \sum_{n=0}^{k-1}d_{nkj}x^{j+n}\mathcal{C}_{\frac{\alpha}{2}-j+n}(x)$$
(34)

To prove (34) by (71), we first fix k = 1 and induct on j. We have

$$x\frac{d}{dx}x^{j}\mathcal{C}_{\frac{\alpha}{2}-j}(x) = -x^{j+1}\mathcal{C}_{\frac{\alpha}{2}-j+1}(x) + x^{j}\left(\frac{\alpha}{2}-j\right)\mathcal{C}_{\frac{\alpha}{2}-j}(x) + jx^{j}\mathcal{C}_{\frac{\alpha}{2}-j}(x)$$

$$= -x^{j+1}\mathcal{C}_{\frac{\alpha}{2}-j+1}(x) + x^{j}\frac{\alpha}{2}\mathcal{C}_{\frac{\alpha}{2}-j}(x)$$
(35)

After showing (35) we induct on k. We also can notice that coefficient depending in j cancels in the right hand side of (35). It implies that d_{nkj} actually doesn't depend on j. So (34) can be written as

$$\left(x\frac{d}{dx}\right)^{k}x^{j}\mathcal{C}_{\frac{\alpha}{2}-j}(x) = (-1)^{k}x^{j+k}\mathcal{C}_{\frac{\alpha}{2}-j+k}(x) + \sum_{n=0}^{k-1}d_{nk}x^{j+n}\mathcal{C}_{\frac{\alpha}{2}-j+n}(x)$$
(36)

Again, we furtherly simplify the determinant by (36):

$$\det\left(\left\{\left(x\frac{d}{dx}\right)^{k+j}\mathcal{C}_{\frac{\alpha}{2}}(x)\right\}_{k,j=0}^{n-1}\right) = \begin{vmatrix} \mathcal{C}_{\frac{\alpha}{2}}(x) & -x\mathcal{C}_{\frac{\alpha}{2}+1}(x) + d_{01}\mathcal{C}_{\frac{\alpha}{2}}(x) & \cdots \\ x\mathcal{C}_{\frac{\alpha}{2}-1}(x) & -x^{2}\mathcal{C}_{\frac{\alpha}{2}}(x) + d_{01}x\mathcal{C}_{\frac{\alpha}{2}-1}(x) & \cdots \\ \vdots & \vdots & \ddots \end{vmatrix}$$

Similarly, by applying elementary column operations, we can always use the previous columns to eliminate the $\sum_{n=0}^{k-1} d_{nk} x^{j+n} C_{\frac{\alpha}{2}-j+n}(x)$ part in a fixed column and the value of the determinant doesn't change. Finally, we will end up with:

$$\det\left(\left\{\left(x\frac{d}{dx}\right)^{k+j}\mathcal{C}_{\frac{\alpha}{2}}(x)\right\}_{k,j=0}^{n-1}\right) = \det\left((-1)^{k}x^{j+k}\mathcal{C}_{\frac{\alpha}{2}-j+k}(x)\right)$$

By multi-linearity of determinant, we can factor out $(-1)^k x^{j+k}$ and reach the conclusion:

$$\tau_n(x,\alpha) = \det\left((-1)^k x^{j+k} \mathcal{C}_{\frac{\alpha}{2}-j+k}(x)\right) = x^{n(n-1)}(-1)^{\frac{n(n-1)}{2}} \Delta_n(x,\alpha)$$

where $\Delta_n(x, \alpha)$ is given by (2). That completes the proof.

3.4. Proof of Proposition 1.1. Before starting the proof we need to get the following lemma.

Lemma 3.1. The special function solutions $u_n(x, \alpha - 2)$ and $u_{n+1}(x, \alpha - 2)$ admit the following formulas in terms of $u_n(x, \alpha)$.

$$(u_{n+1}(x,\alpha-2),\alpha+2n,-\alpha+6+2n) = B_1 B_2(u_n(x,\alpha),\alpha+2n,-\alpha+2+2n).$$
(37)

Proof. We start with considering Bäcklund transformations B_1 and B_2 . Using the explicit formulas (15), (16) and equation (1) we can show that these transformations commute: $B_1B_2 = B_2B_1$.

Let's consider now B_2 applied to special function solution (14). After using differential equation (12) for w(x) we get

$$B_2(u_0(x,\alpha),\alpha,2-\alpha)[1] = \frac{2-\alpha}{x} + \frac{w(x)}{w'(x)} = \frac{2-\alpha}{x} + \frac{1}{\frac{\alpha}{2x} + \frac{C'_{\frac{\alpha}{2}}(x)}{C_{\frac{\alpha}{2}}(x)}}.$$

We use notation [1] above for the first component of the output of Bäcklund transformation. We use (71) to get

$$B_2(u_0(x,\alpha),\alpha,2-\alpha)[1] = \frac{2-\alpha}{x} + \frac{\mathcal{C}_{\frac{\alpha}{2}}(x)}{\mathcal{C}_{\frac{\alpha}{2}-1}(x)}.$$

Using identity (70) we can rewrite it as

=

$$B_2(u_0(x,\alpha),\alpha,2-\alpha)[1] = \frac{2-\alpha}{2x} - \frac{\mathcal{C}'_{\frac{\alpha}{2}-1}(x)}{\mathcal{C}_{\frac{\alpha}{2}-1}(x)} = -\frac{d}{dx}\ln(w(x,\alpha-2)) = u_0(x,\alpha-2).$$

Now using commutativity of B_1 and B_2 we get

$$B_2(u_n(x,\alpha),\alpha+2n,-\alpha+2+2n) = B_2B_1^n(u_0(x,\alpha),\alpha,-\alpha+2)$$

= $B_1^n(u_0(x,\alpha-2),\alpha-2,-\alpha+4) = (u_n(x,\alpha-2),\alpha-2+2n,-\alpha+4+2n).$

Using similar logic we arrive to (37) and finish the proof.

Proof of Proposition 1.1. We start our proof by introducing the following sequence of functions

$$\widetilde{u}_n(x,\alpha) = -\frac{\Delta_{n+1}(x,\alpha-2)\Delta_n(x,\alpha)}{\Delta_{n+1}(x,\alpha)\Delta_n(x,\alpha-2)}.$$

Using Proposition 3.5 we can rewrite it in terms of tau functions

$$\widetilde{u}_n(x,\alpha) = -\frac{\tau_{n+1}(x,\alpha-2)\tau_n(x,\alpha)}{\tau_{n+1}(x,\alpha)\tau_n(x,\alpha-2)}.$$
(38)

Using the Toda equation (28) and the definition of tau function (19) we can see that

$$\widetilde{u}_{n+1}(x,\alpha) = \widetilde{u}_n(x,\alpha) \frac{\left(x\frac{d}{dx}\right)^2 \ln(\tau_{n+1}(x,\alpha-2))}{\left(x\frac{d}{dx}\right)^2 \ln(\tau_{n+1}(x,\alpha))} = \widetilde{u}_n(x,\alpha) \frac{\left(x\frac{d}{dx}\right) h_{n+1}(x,\alpha-2)}{\left(x\frac{d}{dx}\right) h_{n+1}(x,\alpha)}.$$
(39)

Using Lemma 3.1 and definition of $h_{n+1}(x, \alpha)$ we can express right hand side of (39) in terms of $u_n(x, \alpha)$. We also provide intermediate formulas

$$h_{n+1}(x,\alpha) = -\frac{1}{8u_n(x,\alpha)^2} \Big[u_n(x,\alpha)^2 \left(\alpha^2 - 4n^2 + 2x^2 + 3\right) + 2xu_n(x,\alpha) \left(\alpha - 2n + u'_n(x,\alpha) - 2\right) \\ + 2x(\alpha + 2n + 2)u_n(x,\alpha)^3 - x^2 \left((u'_n(x,\alpha))^2 - 1 \right) + x^2u_n(x,\alpha)^4 \Big],$$

$$h_{n+1}(x,\alpha-2) = -\frac{1}{8u_n(x,\alpha)^2} \Big[u_n(x,\alpha)^2 \left(a^2 - 4a - 4n^2 + 2x^2 + 7\right) + x^2u_n(x,\alpha)^4 \\ - 2xu_n(x,\alpha) \left(-a + 2n + u'_n(x,\alpha) + 4\right) + 2x(a + 2n)u_n(x,\alpha)^3 - x^2 \left(u'_n(x,\alpha)^2 - 1\right) \Big].$$

As the result we get

$$\widetilde{u}_{n+1}(x,\alpha) = \widetilde{u}_n(x,\alpha) \frac{(-3+\alpha+2n)u_n(x,\alpha) + xu_n^2(x,\alpha) + x + xu_n'(x,\alpha)}{u_n^2(x,\alpha)((1+\alpha+2n)u_n(x,\alpha) + xu_n^2(x,\alpha) + x + xu_n'(x,\alpha))}.$$
(40)

Using identity (70), initial conditions (29), (30) and definiton (38) we can observe that

$$\widetilde{u}_0(x,\alpha) = u_0(x,\alpha)$$

Using explicit formula (15) for Bäcklund transformation B_1 , relation (40) and mathematical induction we get the desired result

$$\widetilde{u}_n(x,\alpha) = u_n(x,\alpha).$$

4. Asymptotics of Toeplitz determinant at zero

4.1. Andrèief identity. To prove our result we rewrite Toeplitz determinant (2) as multiple contour integral.

Proposition 4.1 (see [For19]). Andrèief identity is given by the following formula

$$\int_{\Gamma} \dots \int_{\Gamma} \det\left(\left\{f_j(x_k)\right\}_{j,k=1}^n\right) \det\left(\left\{g_j(x_k)\right\}_{j,k=1}^n\right) \prod_{k=1}^n h(x_k) dx_k = n! \det\left(\left\{\int_{\Gamma} f_j(x)g_k(x)h(x)dx\right\}_{j,k=1}^n\right) dx_k = n! \det\left(\left\{\int_{\Gamma} f_j(x)g_k(x)h(x)dx\right\}_{j,k=1}^n dx_k = n! \det\left(\left\{\int_{\Gamma} f_j(x)g_k(x)h($$

where Γ is some contour in the complex plane, such that the corresponding integral is finite.

We apply Andrèief identity and get the following result.

Theorem 4.1. The Toeplitz determinant $\Delta_n(x, \alpha)$ given by (2) can be rewritten as

$$\Delta_n(x,\alpha) = \frac{1}{n!} \int_{\Gamma_1 \cup \Gamma_2} \dots \int_{\Gamma_1 \cup \Gamma_2} \prod_{1 \le j < k \le n} (t_k - t_j) \left(\frac{1}{t_k} - \frac{1}{t_j}\right) \prod_{k=1}^n h_1(t_k) dt_k \tag{41}$$

where

$$h_1(t) = \frac{\mathrm{e}^{\frac{x}{2}\left(t - \frac{1}{t}\right)}}{2\pi \mathrm{i} t^{1 + \frac{\alpha}{2}}} \left(\left(d_1 + d_2 \cot\left(\frac{\pi \alpha}{2}\right) \right) \chi_{\Gamma_1}(t) + d_2 \csc\left(\frac{\pi \alpha}{2}\right) \mathrm{e}^{\frac{\mathrm{i}\pi \alpha}{2}} \chi_{\Gamma_2}(t) \right),$$

and contours of integration Γ_1 , Γ_2 are shown on Figure 7. We use the notation $\chi_{\Gamma_j}(t)$ for the characteristic function of contour Γ_j . We assume $-\pi < \arg(t) < \pi$ on contour Γ_1 and $0 < \arg(t) < 2\pi$ on contour Γ_2 .

Proof. Using contour integral representation (64) we get

$$\mathcal{C}_{\frac{\alpha}{2}-j+k}(x) = \int\limits_{\Gamma_1\cup\Gamma_2} t^{-k} t^j h_1(t) dt$$

Put $g_k(t) = t^{-k}$ and $f_j(t) = t^j$. By Proposition 4.1, we get

$$\Delta_{n}(x,\alpha) = \frac{1}{n!} \int_{\Gamma_{1}\cup\Gamma_{2}} \dots \int_{\Gamma_{1}\cup\Gamma_{2}} \det\left(\{f_{j}(t_{k})\}_{j,k=0}^{n-1}\right) \det\left(\{g_{j}(t_{k})\}_{j,k=0}^{n-1}\right) \prod_{k=0}^{n-1} h_{1}(t_{k}) dt_{k}$$
$$= \frac{1}{n!} \int_{\Gamma_{1}\cup\Gamma_{2}} \dots \int_{\Gamma_{1}\cup\Gamma_{2}} \det\left(\{t_{k}^{j}\}_{j,k=0}^{n-1}\right) \det\left(\{t_{k}^{-j}\}_{j,k=0}^{n-1}\right) \prod_{k=0}^{n-1} h_{1}(t_{k}) dt_{k}$$

By the formula for the Vandermonde determinant we can simplify the integrand and get

$$\det\left(\left\{t_k^j\right\}_{j,k=0}^{n-1}\right) = \prod_{0 \le j < k \le n-1} (t_k - t_j)$$

$$\det\left(\left\{t_{k}^{-j}\right\}_{j,k=0}^{n-1}\right) = \prod_{0 \le j < k \le n-1} \left(\frac{1}{t_{k}} - \frac{1}{t_{j}}\right)$$

Thus, the explicit formula for $\Delta_n(x, \alpha)$ is given by:

$$\Delta_n(x,\alpha) = \frac{1}{n!} \int_{\Gamma_1 \cup \Gamma_2} \dots \int_{\Gamma_1 \cup \Gamma_2} \prod_{0 \le j < k \le n-1} (t_k - t_j) \left(\frac{1}{t_k} - \frac{1}{t_j}\right) \prod_{k=0}^{n-1} h_1(t_k) dt_k.$$

For convenience we shift the index of variables t_k .

4.2. **Basic strategies.** Up to this point, we've got enough preparation to compute the asymptotics at zero. Our goal is to get asymptotics $\Delta_n(x, \alpha) \sim b(n)x^{a(n)}$ when $x \to 0$. It is reasonable expectation since Bessel function $J_{\nu}(x)$ admits series representation (62). We summarize several key ideas to achieve this goal:

- The contours Γ_1 and Γ_2 spread out to zero and infinity in the formula (41). We can't put x = 0 here without losing convergence of the integral.
- Expanding the product $\prod_{k=0}^{n-1} h(t_k)$ in the integrand of (41) we get the expressions, where some of the variables t_k will belong to contour Γ_1 and others will belong to Γ_2 .
- We apply change of variables $t = \frac{2}{x}s$ to variables on contours Γ_1 . The integrand will preserve exponential decay at infinity when we put x = 0. On the other hand, we can apply change of variable $t = \frac{x}{2}s$ to variables on contours Γ_2 . In this case the integrand will preserve exponential decay at zero when we put x = 0.
- It seems from the first glance that it would be more convenient to use expression $d_1J_{\nu}(x) + d_2J_{-\nu}(x)$ instead of (3). But unfortunately Proposition 1.1 would fail if we replace $Y_{\nu}(x)$ with $J_{-\nu}(x)$. It follows from the fact that the differential identities (70), (71) would fail for this alternative choice. And we need them to hold, since they were used extensively in the proof of Proposition 3.5.

4.3. Expanded formula for $\Delta_n(x, \alpha)$. We start with the following observation. Let I denotes a subset of the set of indices $\{1, \ldots, n\}$, r denotes its cardinality |I| and I^c denotes its complement. The following identity holds

$$\prod_{k=1}^{n} (c_1 \chi_{\Gamma_1}(t_k) + c_2 \chi_{\Gamma_2}(t_k)) = \sum_{\substack{I \subset \{1, \dots, n\} \\ |I| = r}} c_1^r c_2^{n-r} \prod_{k \in I} \chi_{\Gamma_1}(t_k) \prod_{j \in I^c} \chi_{\Gamma_2}(t_j).$$
(42)

Now we apply (42) to the expression in Theorem 4.1 in order to convert the formula into a summation form and decouple the contours. Denote

$$c_1 = d_1 + d_2 \cot\left(\frac{\pi\alpha}{2}\right), \quad c_2 = d_2 \csc\left(\frac{\pi\alpha}{2}\right) e^{\frac{i\pi\alpha}{2}}.$$
 (43)

We have

$$\begin{split} \Delta_{n}(x,\alpha) &= \frac{(-1)^{\frac{n(n-1)}{2}}}{n!} \int\limits_{\Gamma_{1}\cup\Gamma_{2}} \dots \int\limits_{\Gamma_{1}\cup\Gamma_{2}} \prod_{m=1}^{n} (c_{1}\chi_{\Gamma_{1}}(t_{m}) + c_{2}\chi_{\Gamma_{2}}(t_{m})) \prod_{1 \leq j < k \leq n} \frac{(t_{j} - t_{k})^{2}}{t_{j}t_{k}} \prod_{l=1}^{n} \frac{e^{\frac{x}{2}(t_{l} - \frac{1}{t_{l}})}}{2\pi i t_{l}^{\frac{n}{2} + 1}} dt_{l} \\ &= \frac{(-1)^{\frac{n(n-1)}{2}}}{n!} \int\limits_{\Gamma_{1}\cup\Gamma_{2}} \dots \int\limits_{\Gamma_{1}\cup\Gamma_{2}} \sum_{I \subset \{1,\dots,n\}} c_{1}^{r} c_{2}^{n-r} \prod_{i \in I} \chi_{\Gamma_{1}}(t_{i}) \prod_{j \in I^{c}} \chi_{\Gamma_{2}}(t_{j}) \prod_{1 \leq j < k \leq n} \frac{(t_{j} - t_{k})^{2}}{t_{j}t_{k}} \prod_{l=1}^{n} \frac{e^{\frac{x}{2}(t_{l} - \frac{1}{t_{l}})}}{2\pi i t_{l}^{\frac{n}{2} + 1}} dt_{l} \\ &= \frac{(-1)^{\frac{n(n-1)}{2}}}{(2\pi i)^{n} n!} \sum_{I \subset \{1,\dots,n\}} c_{1}^{r} c_{2}^{n-r} \int_{\Gamma_{1}} \dots \int\limits_{\Gamma_{2}} \int\limits_{\Gamma_{2}} \dots \int\limits_{\Gamma_{2}} \sqrt{\prod_{j \neq k} \frac{(t_{j} - t_{k})^{2}}{t_{j}t_{k}}} \sqrt{\prod_{j \neq k} \frac{(t_{j} - t_{k})^{2}}{t_{j}t_{k}}} \sqrt{\prod_{j \in I^{c}} \frac{(t_{j} - t_{k})^{2}}{t_{j}t_{k}}} \\ &\sqrt{\prod_{\substack{j \in I^{c} \\ k \in I}} \frac{(t_{j} - t_{k})^{2}}{t_{j}t_{k}}} \prod_{l \in I} \frac{e^{\frac{x}{2}(t_{l} - \frac{1}{t_{l}})}}{t_{l}^{\frac{n}{2} + 1}} dt_{l} \prod_{l \in I^{c}} \frac{e^{\frac{x}{2}(t_{l} - \frac{1}{t_{l}})}}{t_{l}^{\frac{n}{2} + 1}} dt_{l} \end{split}$$

For $t_k \in I$, we use change of variable $t_k = \frac{2}{x}s_k$. On the other hand, for $t_j \in I^c$, we use change of variable $t_j = \frac{x}{2}s_j$. The formula above becomes

$$\Delta_{n}(x,\alpha) = \frac{(-1)^{\frac{n(n-1)}{2}}}{(2\pi\mathrm{i})^{n}n!} \sum_{\substack{I \subset \{1,\dots,n\}\\|I|=r}} c_{1}^{r} c_{2}^{n-r} \int_{\Gamma_{1}} \dots \int_{\Gamma_{1}} \int_{\Gamma_{2}} \dots \int_{\Gamma_{2}} \prod_{\substack{j < k\\j,k \in I}} \frac{(s_{j} - s_{k})^{2}}{s_{j}s_{k}} \prod_{\substack{j < k\\j,k \in I^{c}}} \frac{(s_{j} - s_{k})^{2$$

Group all the $\frac{2}{x}$ factors together and pull them out of the summation,

$$\begin{split} \Delta_n(x,\alpha) &= \frac{(-1)^{\frac{n(n-1)}{2}}}{(2\pi\mathrm{i})^n n!} \sum_{\substack{I \subset \{1,\dots,n\}\\|I|=r}} c_1^r c_2^{n-r} \left(\frac{2}{x}\right)^{-\alpha r+2r(n-r)+\frac{\alpha n}{2}} \int_{\Gamma_1} \dots \int_{\Gamma_1} \int_{\Gamma_2} \dots \int_{\Gamma_2} \prod_{\substack{j < k\\j,k \in I}} \frac{(s_j - s_k)^2}{s_j s_k} \\ &\prod_{\substack{j < k\\j,k \in I^c}} \frac{(s_j - s_k)^2}{s_j s_k} \prod_{\substack{j \in I\\k \in I^c}} \frac{(s_j(1+O(x^2))^2}{s_j s_k} \prod_{l \in I} \frac{e^{s_l}(1+O(x^2))}{s_l^{\frac{\alpha}{2}+1}} ds_l \prod_{l \in I^c} \frac{e^{-\frac{1}{s_l}}(1+O(x^2))}{s_l^{\frac{\alpha}{2}+1}} ds_l \end{split}$$

We also want to group all the products of variable s together and separate the integrals based on different contours. We rewrite the following three parts,

$$\prod_{\substack{j < k\\j,k \in I}} \frac{(s_j - s_k)^2}{s_j s_k} = \prod_{\substack{j < k\\j,k \in I}} (s_j - s_k)^2 \prod_{\substack{j \neq k\\j,k \in I}} \frac{1}{\sqrt{s_j s_k}} = \prod_{\substack{j < k\\j,k \in I}} (s_j - s_k)^2 \prod_{l \in I} s_l^{\gamma_1}$$
(44)

$$\prod_{\substack{j < k \\ j,k \in I^c}} \frac{(s_j - s_k)^2}{s_j s_k} = \prod_{\substack{j < k \\ j,k \in I^c}} (s_j - s_k)^2 \prod_{\substack{j \neq k \\ j,k \in I^c}} \frac{1}{\sqrt{s_j s_k}} = \prod_{\substack{j < k \\ j,k \in I^c}} (s_j - s_k)^2 \prod_{l \in I^c} s_l^{\gamma_2}$$
(45)
$$\prod_{\substack{j \in I \\ k \in I^c}} \frac{s_j}{s_k} = \prod_{l \in I} s_l^{\gamma_3} \prod_{l \in I^c} s_l^{\gamma_4}$$

To find γ_1 we can interpret the product in (44) as product a over all elements of $r \times r$ matrix except for the diagonal. Terms with s_j appear along *j*th row and *j*th column, so there are 2(r-1) of them. Similar argument can be used for computation of γ_2 , but the size of the matrix would be $(n-r) \times (n-r)$. Keeping in mind that we still have square root and introducing the power $-\frac{1}{2}$, then we get:

$$\gamma_1 = -\frac{2(r-1)}{2} = 1 - r,$$

 $\gamma_2 = -\frac{2(n-r-1)}{2} = -n + r + 1.$

To compute γ_3 and γ_4 we visualize the number of the *s*-factors using the following matrix

$$\left\{\frac{s_{j_{\alpha}}}{s_{k_{\beta}}}\right\}_{\substack{1 \leq \alpha \leq r \\ 1 \leq \beta \leq n-r}} = \begin{pmatrix}\frac{s_{j_{1}}}{s_{k_{1}}} & \frac{s_{j_{1}}}{s_{k_{2}}} & \frac{s_{j_{1}}}{s_{k_{3}}} & \cdots \\ \frac{s_{j_{2}}}{s_{k_{1}}} & \frac{s_{j_{2}}}{s_{k_{2}}} & \frac{s_{j_{2}}}{s_{k_{3}}} & \cdots \\ \frac{s_{j_{2}}}{s_{k_{1}}} & \frac{s_{j_{2}}}{s_{k_{2}}} & \frac{s_{j_{2}}}{s_{k_{3}}} & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

We can observe that horizontally, for each $s_{j_{\alpha}}$, there are (n-r) factors. Vertically, for each $s_{k_{\beta}}$ there are -r factors. We get

$$\gamma_3 = n - r_1$$
$$\gamma_4 = -r_2$$

As the result we obtain the following expanded formula for $\Delta_n(x, \alpha)$

$$\Delta_{n}(x,\alpha) = \frac{(-1)^{\frac{n(n-1)}{2}}}{(2\pi i)^{n}n!} \sum_{\substack{I \subset \{1,\dots,n\}\\|I|=r}} c_{1}^{r} c_{2}^{n-r} \left(\frac{2}{x}\right)^{-\alpha r+2r(n-r)+\frac{\alpha n}{2}} (1+O(x^{2}))$$

$$\times \int_{\Gamma_{1}} \dots \int_{\Gamma_{1}} \prod_{\substack{j < k\\j,k \in I}} (s_{j} - s_{k})^{2} \prod_{l \in I} e^{s_{l}} s_{l}^{-\frac{\alpha}{2}-2r+n} ds_{l} \int_{\Gamma_{2}} \dots \int_{\Gamma_{2}} \prod_{\substack{j < k\\j,k \in I^{c}}} (s_{j} - s_{k})^{2} \prod_{l \in I^{c}} e^{-\frac{1}{s_{l}}} s_{l}^{-\frac{\alpha}{2}-n} ds_{l}$$

We can see that terms in the right hand side only depend on cardinality of I. Rewriting it, we get the following preliminary asymptotic formula for $\Delta_n(x, \alpha)$.

Lemma 4.1. The Toeplitz determinant (2) admits the following $x \to 0$ asymptotics for fixed $d_1, d_2 \in \mathbb{R}$, $n \in \mathbb{N} \cup \{0\}, \alpha \in \mathbb{C} \setminus (2\mathbb{Z} + i\mathbb{R})$

$$\Delta_{n}(x,\alpha) = \frac{(-1)^{\frac{n(n-1)}{2}}}{(2\pi\mathrm{i})^{n}} \sum_{r=0}^{n} \frac{c_{1}^{r} c_{2}^{n-r}}{r!(n-r)!} \left(\frac{2}{x}\right)^{-\alpha r+2r(n-r)+\frac{\alpha n}{2}} (1+O(x^{2})) \\ \times \int_{\Gamma_{1}} \dots \int_{\Gamma_{1}} \prod_{1 \le j < k \le r} (s_{j} - s_{k})^{2} \prod_{l=1}^{r} \mathrm{e}^{s_{l}} s_{l}^{-\frac{\alpha}{2}-2r+n} ds_{l} \int_{\Gamma_{2}} \dots \int_{\Gamma_{2}} \prod_{1 \le j < k \le n-r} (s_{j} - s_{k})^{2} \prod_{l=1}^{n-r} \mathrm{e}^{-\frac{1}{s_{l}}} s_{l}^{-\frac{\alpha}{2}-n} ds_{l}.$$

$$\tag{46}$$

Coefficients c_1 , c_2 are given by (43). Contours of integration Γ_1 , Γ_2 are shown on Figure 7. We assume $-\pi < \arg(t) < \pi$ on contour Γ_1 and $0 < \arg(t) < 2\pi$ on contour Γ_2 .

4.4. Proof of Theorem 1.1. The asymptotics of $\Delta_n(x,\alpha)$ is the leading term of the asymptotic formula (46). Denote the power of x appearing in (46) as $p(r, \alpha, n)$:

$$p(r,\alpha,n) = \alpha r - \frac{n\alpha}{2} - 2r(n-r).$$
(47)

We need to find the minimum of $\operatorname{Re}(p(r,\alpha,n))$ with respect to r. Introduce notation for the index which realizes this minimum

$$\min_{\substack{0 \le r \le n \\ r \in \mathbb{N} \cup \{0\}}} \operatorname{Re}(p(r, \alpha, n)) = p(r_c(\alpha, n), \alpha, n).$$
(48)

We have the following formula for it.

Lemma 4.2. The critical index $r_c(\alpha, n)$ defined by (48) admits the following piecewise formula

$$r_{c}(\alpha, n) = \begin{cases} 0 & \text{if } \operatorname{Re}(\alpha) > 2n - 2\\ j & \text{if } 2n - 4j - 2 < \operatorname{Re}(\alpha) < 2n - 4j + 2\\ & \text{and } j = 1, 2, \dots, n - 1\\ n & \text{if } \operatorname{Re}(\alpha) < -2n + 2 \end{cases}$$
(49)

 \square

Proof. Since $0 \le r \le n$ and $r \in \mathbb{Z}$, so $p(r, \alpha, n)$ only takes values on that discrete set. It is clear that $\operatorname{Re}(p(r,\alpha,n))$ is an upward parabola in variable r and it takes minimum value at $r_{\min}(n,\alpha) = \frac{2n - \operatorname{Re}(\alpha)}{4}$. We will discuss different cases of relative positions between $r_{\min}(\alpha, n)$ and $r_c(\alpha, n)$. If $r_{\min}(\alpha, n) \leq 0$, then $r_c(\alpha, n) = 0$. If $r_{\min}(\alpha, n) \ge n$, then $r_c(\alpha, n) = n$. Let $0 \le j \le n$ and $j \le r_{\min}(\alpha, n) \le j + 1$. If $j \leq r_{\min}(\alpha, n) < j + \frac{1}{2}$, then $r_c(\alpha, n) = j$. If $j + \frac{1}{2} < r_{\min}(\alpha, n) \leq j + 1$, then $r_c(\alpha, n) = j + 1$. In other words,

•
$$r_c(\alpha, n) = 0$$
 when $\frac{2n - \operatorname{Re}(\alpha)}{4} < \frac{1}{2}$

•
$$r_c(\alpha, n) = 0$$
 when $\frac{2n - \operatorname{Re}(\alpha)}{4} < \frac{1}{2}$,
• $r_c(\alpha, n) = j$ when $j - \frac{1}{2} < \frac{2n - \operatorname{Re}(\alpha)}{4} < j + \frac{1}{2}$ and $j = 1, 2, \dots, n - 1$,

•
$$r_c(\alpha, n) = n$$
 when $\frac{2n - \text{Re}(\alpha)}{4} > n - \frac{1}{2}$.

These conditions can be rewritten as (49).

Remark 4.1. Floor function gives a more compact form for $r_c(\alpha, n)$. Indeed $\forall \ 0 \le j \le n$, we know $r_c(\alpha, n) = j$ if and only if $j - \frac{1}{2} \le \frac{n}{2} - \frac{\operatorname{Re}(\alpha)}{4} \le j + \frac{1}{2}$ or $j \le \frac{n}{2} - \frac{\operatorname{Re}(\alpha)}{4} + \frac{1}{2} \le j + 1$. Therefore

$$r_c(\alpha, n) = \begin{cases} n & \text{if } \operatorname{Re}(\alpha) < 2 - 2n, \\ \lfloor \frac{n}{2} - \frac{\operatorname{Re}(\alpha)}{4} + \frac{1}{2} \rfloor & \text{if } 2 - 2n < \operatorname{Re}(\alpha) < 2n - 2, \\ 0 & \text{if } \operatorname{Re}(\alpha) > 2n - 2. \end{cases}$$

Remark 4.2. We consider case $\alpha \in \mathbb{C} \setminus (2\mathbb{Z} + i\mathbb{R})$ to make sure that leading contribution comes from the one value of r_c . Obterwise we need to include two contributions in the leading term of asymptotics.

As the result we have

$$\begin{split} \Delta_n(x,\alpha) &\sim \frac{(-1)^{\frac{n(n-1)}{2}}}{(2\pi \mathrm{i})^n} \frac{c_1^{r_c} c_2^{n-r_c}}{r_c!(n-r_c)!} \left(\frac{x}{2}\right)^{p(r_c,\alpha,n)} \int\limits_{\Gamma_1} \dots \int\limits_{\Gamma_1} \prod_{1 \le j < k \le r_c} (s_j - s_k)^2 \prod_{l=1}^{r_c} \mathrm{e}^{s_l} s_l^{-\frac{\alpha}{2} - 2r_c + n} ds_l \\ &\times \int\limits_{\Gamma_2} \dots \int\limits_{\Gamma_2} \prod_{1 \le j < k \le n - r_c} (s_j - s_k)^2 \prod_{l=1}^{n-r_c} \mathrm{e}^{-\frac{1}{s_l}} s_l^{-\frac{\alpha}{2} - n} ds_l, \quad \text{as } x \to 0. \end{split}$$

Denote

$$H_1(\alpha, n) = \int_{\Gamma_1} \dots \int_{\Gamma_1} \prod_{1 \le j < k \le r_c} (s_j - s_k)^2 \prod_{l=1}^{r_c} e^{s_l} s_l^{-\frac{\alpha}{2} - 2r_c + n} ds_l$$
(50)

and

$$H_2(\alpha, n) = \int_{\Gamma_2} \dots \int_{\Gamma_2} \prod_{1 \le j < k \le n - r_c} (s_j - s_k)^2 \prod_{l=1}^{n-r_c} e^{-\frac{1}{s_l}} s_l^{-\frac{\alpha}{2} - n} ds_l.$$
(51)

To evaluate (50), (51), we reduce them to multiple integrals with the Laguerre weight $w(x) = e^{-x}x^{\alpha}$, Re(α) > -1 on the contour $\Gamma = [0, \infty)$. In $H_1(\alpha, n)$, we make the change of variable $s = \tilde{s}e^{-i\pi}$. More specifically the modulus and argument of the variable transform as

$$\begin{split} |s| &= |\widetilde{s}| \\ \arg(s) &= \arg(\widetilde{s}) - \pi, \quad 0 < \arg(\widetilde{s}) < 2\pi. \end{split}$$

The contour Γ_1 becomes $\widetilde{\Gamma_1}$ as shown in the picture below.



FIGURE 4. Contour $\widetilde{\Gamma_1}$

Also notice that

$$s_{l}^{-\frac{\alpha}{2}-2r_{c}+n} = e^{(-\frac{\alpha}{2}-2r_{c}+n)\ln|s_{l}|+(-\frac{\alpha}{2}-2r_{c}+n)i\arg(s_{l})}$$

= $e^{(-\frac{\alpha}{2}-2r_{c}+n)\ln|\tilde{s}_{l}|+(-\frac{\alpha}{2}-2r_{c}+n)i(\arg(\tilde{s}_{l})-\pi)}$
= $\tilde{s}_{l}^{-\frac{\alpha}{2}-2r_{c}+n}e^{-i\pi(-\frac{\alpha}{2}-2r_{c}+n)}$

As the result we have

$$H_{1}(\alpha, n) = \int_{\widetilde{\Gamma_{1}}} \dots \int_{\widetilde{\Gamma_{1}}} \prod_{1 \le j < k \le r_{c}} (\widetilde{s_{j}} - \widetilde{s_{k}})^{2} \prod_{l=1}^{r_{c}} e^{-\widetilde{s_{l}}} \widetilde{s_{l}}^{-\frac{\alpha}{2} - 2r_{c} + n} e^{-i\pi(-\frac{\alpha}{2} - 2r_{c} + n)} (-1) d\widetilde{s_{l}}$$
$$= (-1)^{r_{c}} (e^{-i\pi(-\frac{\alpha}{2} - 2r_{c} + n)})^{r_{c}} \int_{\widetilde{\Gamma_{1}}} \dots \int_{\widetilde{\Gamma_{1}}} \prod_{1 \le j < k \le r_{c}} (\widetilde{s_{j}} - \widetilde{s_{k}})^{2} \prod_{l=1}^{r_{c}} e^{-\widetilde{s_{l}}} \widetilde{s_{l}}^{-\frac{\alpha}{2} - 2r_{c} + n} d\widetilde{s_{l}}$$

To continue our evaluation we need the following Lemma.

Lemma 4.3. Using the properties of power function we can show the following identity

$$\int_{0}^{\infty} s^{j} \mathrm{e}^{-s} s^{\gamma} ds = \frac{1}{\mathrm{e}^{2\pi \mathrm{i}\gamma} - 1} \int_{\widetilde{\Gamma_{1}}} s^{j} \mathrm{e}^{-s} s^{\gamma} ds, \quad j \in \mathbb{Z}, \quad j + \mathrm{Re}(\gamma) > -1.$$
(52)

With the aid of (52) we can rewrite our contour integrals in terms of real line integrals under the convergence condition $\operatorname{Re}(\alpha) < 2n + 2 - 4r_c$.

$$H_1(\alpha, n) = (-1)^{r_c} (\mathrm{e}^{-\mathrm{i}\pi(-\frac{\alpha}{2}-2r_c+n)})^{r_c} (\mathrm{e}^{2\pi\mathrm{i}(-\frac{\alpha}{2}-2r_c+n)}-1)^{r_c}$$
$$\times \int_0^\infty \dots \int_0^\infty \prod_{1 \le j < k \le r_c} (\widetilde{s_j} - \widetilde{s_k})^2 \prod_{l=1}^{r_c} \mathrm{e}^{-\widetilde{s_l}} \widetilde{s_l}^{-\frac{\alpha}{2}-2r_c+n} d\widetilde{s_l}$$

By using [DLMF, (5.14.5)] we get for $\operatorname{Re}(\alpha) < 2n + 2 - 4r_c$

$$H_1(\alpha, n) = (-1)^{nr_c} (2i)^{r_c} \sin^{r_c} \left(\frac{\pi\alpha}{2}\right) \frac{G\left(1 + n - r_c - \frac{\alpha}{2}\right) G(r_c + 2)}{G\left(1 - \frac{\alpha}{2} - 2r_c + n\right)}.$$
(53)

We can notice that left and right hand sides of (53) are entire functions of α . Therefore by the uniqueness of analytic continuation we can say that (53) holds for all $\alpha \in \mathbb{C}$.

Similarly, in $H_2(\alpha, n)$, we put $s = \frac{1}{\tilde{s}}$. Then the modulus and argument of the variable respectively transform as

$$\begin{split} |s| &= \frac{1}{|\widetilde{s}|} \\ \arg(\widetilde{s}) &= -\arg(s), \quad -2\pi < \arg(\widetilde{s}) < 0 \end{split}$$

The contour Γ_2 becomes $\widetilde{\Gamma_2}$ as shown in the picture below. The second multi-integral becomes



FIGURE 5. Contour $\widetilde{\Gamma_2}$

$$\begin{aligned} H_2(\alpha, n) &= \int_{\widetilde{\Gamma_2}} \dots \int_{\widetilde{\Gamma_2}} \prod_{1 \le j < k \le n-r_c} \left(\frac{1}{\widetilde{s_j}} - \frac{1}{\widetilde{s_k}}\right)^2 \prod_{l=1}^{n-r_c} e^{-\widetilde{s_l}} \widetilde{s_l}^{\frac{\alpha}{2}+n} \left(-\frac{1}{s_l^2}\right) d\widetilde{s_l} \\ &= (-1)^{n-r_c} \int_{\widetilde{\Gamma_2}} \dots \int_{1 \le j < k \le n-r_c} \left(\frac{\widetilde{s_j} - \widetilde{s_k}}{\widetilde{s_j}\widetilde{s_k}}\right)^2 \prod_{l=1}^{n-r_c} e^{-\widetilde{s_l}} \widetilde{s_l}^{\frac{\alpha}{2}+n-2} d\widetilde{s_l} \\ &= (-1)^{n-r_c} \int_{\widetilde{\Gamma_2}} \dots \int_{1 \le j < k \le n-r_c} (\widetilde{s_j} - \widetilde{s_k})^2 \prod_{l=1}^{n-r_c} \widetilde{s_l}^{2r_c+2-2n} e^{-\widetilde{s_l}} \widetilde{s_l}^{\frac{\alpha}{2}+n-2} d\widetilde{s_l} \\ &= (-1)^{n-r_c} \int_{\widetilde{\Gamma_2}} \dots \int_{1 \le j < k \le n-r_c} (\widetilde{s_j} - \widetilde{s_k})^2 \prod_{l=1}^{n-r_c} e^{-\widetilde{s_l}} \widetilde{s_l}^{\frac{\alpha}{2}+2r_c-n} d\widetilde{s_l} \end{aligned}$$

Here we used the square of (45)

$$\prod_{1 \le j < k \le n-r_c} \left(\frac{\widetilde{s_j} - \widetilde{s_k}}{\widetilde{s_j}\widetilde{s_k}}\right)^2 = \prod_{1 \le j < k \le n-r_c} (\widetilde{s_j} - \widetilde{s_k})^2 \prod_{\substack{j \ne k \\ 1 \le j, k \le n-r_c}} \frac{1}{\widetilde{s_j}\widetilde{s_k}} = \prod_{1 \le j < k \le n-r_c} (\widetilde{s_j} - \widetilde{s_k})^2 \prod_{l=1}^{n-r_c} \widetilde{s_l}^{2r_c + 2 - 2n}$$

To continue our evaluation we need the following analog of Lemma 4.3.

Lemma 4.4. Using the properties of power function we can show the following identity

$$\int_{0}^{\infty} s^{j} \mathrm{e}^{-s} s^{\gamma} ds = \frac{1}{1 - \mathrm{e}^{-2\pi \mathrm{i}\gamma}} \int_{\widetilde{\Gamma_{2}}} s^{j} \mathrm{e}^{-s} s^{\gamma} ds, \quad j \in \mathbb{Z}, \quad j + \mathrm{Re}(\gamma) > -1.$$
(54)

With the aid of (54) we can rewrite our contour integrals in terms of real line integrals under the convergence condition $\operatorname{Re}(\alpha) > 2n - 2 - 4r_c$

$$H_{2}(\alpha, n) = (-1)^{n-r_{c}} \int_{\widetilde{\Gamma_{2}}} \dots \int_{\widetilde{\Gamma_{2}}} \prod_{1 \le j < k \le n-r_{c}} (\widetilde{s_{j}} - \widetilde{s_{k}})^{2} \prod_{l=1}^{n-r_{c}} e^{-\widetilde{s_{l}}} \widetilde{s_{l}}^{\frac{\alpha}{2} + 2r_{c} - n} d\widetilde{s_{l}}$$
$$= (-1)^{n-r_{c}} (1 - e^{-2\pi i (\frac{\alpha}{2} + 2r_{c} - n)})^{n-r_{c}} \int_{0}^{\infty} \dots \int_{0}^{\infty} \prod_{1 \le j < k \le n-r_{c}} (\widetilde{s_{j}} - \widetilde{s_{k}})^{2} \prod_{l=1}^{n-r_{c}} e^{-\widetilde{s_{l}}} \widetilde{s_{l}}^{\frac{\alpha}{2} + 2r_{c} - n} d\widetilde{s_{l}}$$

Using [DLMF, (5.14.5)] we arrive to the formula

$$H_2(\alpha, n) = (-1)^{n-r_c} \mathrm{e}^{-\mathrm{i}\pi(n-r_c)\frac{\alpha}{2}} (2\mathrm{i})^{n-r_c} \sin^{n-r_c} \left(\frac{\pi\alpha}{2}\right) \frac{G(\frac{\alpha}{2}+r_c+1)G(n+2-r_c)}{G(\frac{\alpha}{2}+2r_c-n+1)}$$
(55)

We repeat the uniqueness of analytic continuation argument above to claim that formula (55) holds for all $\alpha \in \mathbb{C}$. Combining (53), (55), (46) and the definitions (50), (51) we get

$$\begin{aligned} \Delta_n(x,\alpha) &\sim \frac{(-1)^{\frac{n(n-1)}{2}}}{\pi^n} c_1^{r_c} c_2^{n-r_c} \left(\frac{x}{2}\right)^{p(r_c,\alpha,n)} (-1)^{n-r_c+nr_c} \mathrm{e}^{-\mathrm{i}\pi(n-r_c)\frac{\alpha}{2}} \sin^n \left(\frac{\pi\alpha}{2}\right) \\ \frac{G(\frac{\alpha}{2}+r_c+1)G(n+1-r_c)G\left(1+n-r_c-\frac{\alpha}{2}\right)G(r_c+1)}{G(\frac{\alpha}{2}+2r_c-n+1)G\left(1-\frac{\alpha}{2}-2r_c+n\right)}, \quad \text{as } x \to 0. \end{aligned}$$

Using definitions (47), (43) and formula (49) we finish the proof of Theorem 1.1.

5. Asymptotics of special function solutions at zero

To deduce the asymptotics of $u_n(x, \alpha)$, we need to use our main result Theorem 1.1 and Proposition 1.1. We have to shift the index of Toeplitz determinant n and the parameter α .

5.1. Piecewise function for the power of x in the asymptotic of $u_n(x, \alpha)$. Let's introduce notation for the power of x in the asymptotic of $\Delta_n(x, \alpha)$ in Theorem 1.1

$$p_{c}(\alpha, n) = p(r_{c}(\alpha, n), \alpha, n) = \begin{cases} -\frac{\alpha n}{2}, & \text{if } \operatorname{Re}(\alpha) > 2n - 2, \\ (\alpha - 2n + 2j)j - \frac{\alpha n}{2}, & \text{if } 2n - 4j - 2 < \operatorname{Re}(\alpha) < 2n - 4j + 2, \\ & \text{for } j = 1, \dots, n - 1 \\ \frac{\alpha n}{2}, & \text{if } \operatorname{Re}(\alpha) < -2n + 2 \end{cases}$$
(56)

Then the power of x in the asymptotic of $u_n(x, \alpha)$ based on Proposition 1.1 is given by

$$e(\alpha, n) = p_c(\alpha - 2, n + 1) - p_c(\alpha - 2, n) + p_c(\alpha, n) - p_c(\alpha, n + 1).$$
(57)

Lemma 5.1. The piecewise function for the power of x in the asymptotic of $u_n(x, \alpha)$ is given by

$$e(\alpha, n) = \begin{cases} 1 & \text{if } \operatorname{Re}(\alpha) > 2 + 2n \\ \alpha - 2n + 4j - 1 & \text{if } 2n - 4j < \operatorname{Re}(\alpha) < 2n - 4j + 2 \text{ and } j = 0, 1, \dots, n \\ -\alpha + 2n - 4j - 1 & \text{if } 2n - 4j - 2 < \operatorname{Re}(\alpha) < 2n - 4j \text{ and } j = 0, 1, \dots, n - 1 \\ -1 & \text{if } \operatorname{Re}(\alpha) < -2n \end{cases}$$
(58)

Proof. We plug in formula (56) in the expression (57). If $\text{Re}(\alpha) > 2 + 2n$, then we have

$$p_c(\alpha, n) = -\frac{\alpha n}{2}, \qquad p_c(\alpha, n+1) = -\frac{\alpha n}{2} - \frac{\alpha}{2},$$
$$p_c(\alpha - 2, n) = -\frac{\alpha n}{2} + n, \qquad p_c(\alpha - 2, n+1) = -\frac{\alpha n}{2} + n + 1 - \frac{\alpha}{2}.$$

That confirms first case. Similarly, if $2n - 4j < \text{Re}(\alpha) < 2n - 4j + 2$ and $j = 0, 1, \dots, n$, then we have:

$$p_{c}(\alpha, n) = (\alpha - 2n + 2j)j - \frac{\alpha n}{2},$$

$$p_{c}(\alpha, n + 1) = (\alpha - 2n + 2j)j - 2j - \frac{\alpha n}{2} - \frac{\alpha}{2},$$

$$p_{c}(\alpha - 2, n) = (\alpha - 2n + 2j)j - 2j - \frac{\alpha n}{2} + n,$$

$$p_{c}(\alpha - 2, n + 1) = (\alpha - 2n + 2j)j - \frac{\alpha n}{2} - n - 1 + \frac{\alpha}{2}.$$

Notice that for $p_c(\alpha - 2, n + 1)$ we had to shift index j to j + 1 in (56) to get correct formula. We confirmed the second case. If $2n - 4j - 2 < \operatorname{Re}(\alpha) < 2n - 4j$ and $j = 0, 1, \ldots, n - 1$, then we have

$$p_{c}(\alpha, n) = (\alpha - 2n + 2j)j - \frac{\alpha n}{2},$$

$$p_{c}(\alpha, n + 1) = (\alpha - 2n + 2j)j - 2n + 2j - \frac{\alpha n}{2} + \frac{\alpha}{2},$$

$$p_{c}(\alpha - 2, n) = (\alpha - 2n + 2j)j + \alpha - n + 2j - \frac{\alpha n}{2},$$

$$p_{c}(\alpha - 2, n + 1) = (\alpha - 2n + 2j)j - \frac{\alpha n}{2} - n - 1 + \frac{\alpha}{2}$$

Notice that now we had to shift index j to j+1 in (56) for $p_c(\alpha-2, n+1)$, $p_c(\alpha, n+1)$, and $p_c(\alpha-2, n)$ to get correct formula. We confirmed the third case. Finally if $\operatorname{Re}(\alpha) < -2n$, then all terms in (57) change sign compared to the case $\operatorname{Re}(\alpha) > 2n-2$. That confirms the last case and finishes the proof of Lemma 5.1.

5.2. **Proof of Theorem 1.2.** In this part, we will compute the coefficients in the asymptotics of Theorem 1.2. We introduce notation for the coefficient in the asymptotics of $\Delta_n(x, \alpha)$ in Theorem 1.1. We included factor of 2 inside of power of x for convenience

$$c(\alpha, n) = \begin{cases} \left(-1\right)^{\frac{n(n+1)}{2}} \left(\frac{d_2}{\pi}\right)^n \frac{G(n+1)G(\frac{\alpha}{2}+1)}{G(\frac{\alpha}{2}-n+1)}, & \text{if } \operatorname{Re}(\alpha) > 2n-2, \\ \left(-1\right)^{\frac{n(n-1)}{2}+nj+n-j} \left(\frac{d_2}{\pi}\right)^n \left(\frac{d_1}{d_2}\sin\left(\frac{\pi\alpha}{2}\right) + \cos\left(\frac{\pi\alpha}{2}\right)\right)^j & \text{if } 2n-4j-2 < \operatorname{Re}(\alpha) < 2n-4j+2, \\ \times \frac{G(n-j+1)G(-\frac{\alpha}{2}+n-j+1)G(j+1)G(j+\frac{\alpha}{2}+1)}{G(-\frac{\alpha}{2}+n-2j+1)G(\frac{\alpha}{2}-n+2j+1)}, & \text{for } j = 1, \dots, n-1 \\ \\ \frac{\left(-1\right)^{\frac{n(n-1)}{2}+n^2}}{\pi^n} \left(d_1\sin\left(\frac{\pi\alpha}{2}\right) + d_2\cos\left(\frac{\pi\alpha}{2}\right)\right)^n & \text{if } \operatorname{Re}(\alpha) < -2n+2 \\ \times \frac{G(n+1)G(-\frac{\alpha}{2}+1)}{G(-\frac{\alpha}{2}-n+1)}, & (59) \end{cases}$$

Let's denote the constant coefficient in the asymptotic of $u_n(x,\alpha)$ as $q(\alpha,n)$. We have

$$q(\alpha, n) = -\frac{c(\alpha - 2, n + 1)c(\alpha, n)}{c(\alpha - 2, n)c(\alpha, n + 1)}$$

Lemma 5.2. The piecewise formula for $q(\alpha, n)$ is given by

$$q(\alpha, n) = \begin{cases} \frac{2}{2n+2-\alpha}, & \text{if } \operatorname{Re}(\alpha) > 2n+2, \\ (-1)^n \left(\frac{d_1}{d_2} \sin\left(\frac{\pi\alpha}{2}\right) + \cos\left(\frac{\pi\alpha}{2}\right)\right) \left(\frac{\Gamma(-\frac{\alpha}{2}+n-2j+1)}{\Gamma(\frac{\alpha}{2}-n+2j)}\right)^2 & \text{if } 2n-4j < \operatorname{Re}(\alpha) < 2n-4j+2, \\ \times \frac{\Gamma(j+\frac{\alpha}{2})\Gamma(j+1)}{\Gamma(-\frac{\alpha}{2}+n-j+1)\Gamma(n-j+1)}, & \text{for } j = 0, \dots, n \end{cases}$$

$$\left(-1)^n \left(\frac{d_1}{d_2} \sin\left(\frac{\pi\alpha}{2}\right) + \cos\left(\frac{\pi\alpha}{2}\right)\right)^{-1} \left(\frac{\Gamma(\frac{\alpha}{2}-n+2j+1)}{\Gamma(-\frac{\alpha}{2}+n-2j)}\right)^2 & \text{if } 2n-4j-2 < \operatorname{Re}(\alpha) < 2n-4j, \\ \times \frac{\Gamma(-\frac{\alpha}{2}+n-j+1)\Gamma(n-j)}{\Gamma(j+\frac{\alpha}{2}+1)\Gamma(j+1)}, & \text{for } j = 0, \dots, n-1 \\ -\frac{\alpha}{2}-n & \text{if } \operatorname{Re}(\alpha) < -2n \end{cases}\right)$$

21

Proof. For $\operatorname{Re}(\alpha) > 2n + 2$ we have

$$\begin{split} c(\alpha,n) &= (-1)^{\frac{n(n+1)}{2}} \left(\frac{d_2}{\pi}\right)^n \frac{G(n+1)G(\frac{\alpha}{2}+1)}{G(\frac{\alpha}{2}-n+1)},\\ c(\alpha-2,n) &= (-1)^{\frac{n(n+1)}{2}} \left(\frac{d_2}{\pi}\right)^n \frac{G(n+1)G(\frac{\alpha}{2}+1)}{G(\frac{\alpha}{2}-n+1)} \frac{\Gamma(\frac{\alpha}{2}-n)}{\Gamma(\frac{\alpha}{2})},\\ c(\alpha,n+1) &= (-1)^{\frac{(n+2)(n+1)}{2}} \left(\frac{d_2}{\pi}\right)^{n+1} \frac{G(n+1)G(\frac{\alpha}{2}+1)}{G(\frac{\alpha}{2}-n+1)} \Gamma(n+1)\Gamma\left(\frac{\alpha}{2}-n\right),\\ c(\alpha-2,n+1) &= (-1)^{\frac{(n+2)(n+1)}{2}} \left(\frac{d_2}{\pi}\right)^{n+1} \frac{G(n+1)G(\frac{\alpha}{2}+1)}{G(\frac{\alpha}{2}-n+1)} \frac{\Gamma(n+1)\Gamma\left(\frac{\alpha}{2}-n\right)\Gamma\left(\frac{\alpha}{2}-n-1\right)}{\Gamma\left(\frac{\alpha}{2}\right)}. \end{split}$$

That implies

$$q(\alpha, n) = -\frac{\Gamma\left(\frac{\alpha}{2} - n - 1\right)}{\Gamma\left(\frac{\alpha}{2} - n\right)} = \frac{2}{2n + 2 - \alpha}.$$

That confirms the first case.

For $2n - 4j < \operatorname{Re}(\alpha) < 2n - 4j + 2$ we have

$$\begin{split} c(\alpha,n) &= (-1)^{\frac{n(n-1)}{2} + nj + n - j} \left(\frac{d_2}{\pi}\right)^n \left(\frac{d_1}{d_2} \sin\left(\frac{\pi\alpha}{2}\right) + \cos\left(\frac{\pi\alpha}{2}\right)\right)^j \\ &\times \frac{G(n - j + 1)G(-\frac{\alpha}{2} + n - j + 1)G(j + 1)G(j + \frac{\alpha}{2} + 1)}{G(-\frac{\alpha}{2} + n - 2j + 1)G(\frac{\alpha}{2} - n + 2j + 1)}, \\ c(\alpha - 2, n) &= (-1)^{\frac{n(n-1)}{2} + nj + n} \left(\frac{d_2}{\pi}\right)^n \left(\frac{d_1}{d_2} \sin\left(\frac{\pi\alpha}{2}\right) + \cos\left(\frac{\pi\alpha}{2}\right)\right)^j \\ &\times \frac{G(n - j + 1)G(-\frac{\alpha}{2} + n - j + 1)G(j + 1)G(j + \frac{\alpha}{2} + 1)}{G(-\frac{\alpha}{2} + n - 2j + 1)G(\frac{\alpha}{2} - n + 2j + 1)} \frac{\Gamma(-\frac{\alpha}{2} + n - j + 1)\Gamma(\frac{\alpha}{2} - n + 2j)}{\Gamma(j + \frac{\alpha}{2})\Gamma(-\frac{\alpha}{2} + n - 2j + 1)}, \\ c(\alpha, n + 1) &= (-1)^{\frac{n(n+1)}{2} + nj + n + 1} \left(\frac{d_2}{\pi}\right)^{n+1} \left(\frac{d_1}{d_2} \sin\left(\frac{\pi\alpha}{2}\right) + \cos\left(\frac{\pi\alpha}{2}\right)\right)^j \\ &\times \frac{G(n - j + 1)G(-\frac{\alpha}{2} + n - j + 1)G(j + 1)G(j + \frac{\alpha}{2} + 1)}{G(-\frac{\alpha}{2} + n - 2j + 1)G(\frac{\alpha}{2} - n + 2j + 1)}, \\ c(\alpha - 2, n + 1) &= (-1)^{\frac{n(n+1)}{2} + nj + j + 2n + 2} \left(\frac{d_2}{\pi}\right)^{n+1} \left(\frac{d_1}{d_2} \sin\left(\frac{\pi\alpha}{2}\right) + \cos\left(\frac{\pi\alpha}{2}\right)\right)^{j+1} \\ &\times \frac{G(n - j + 1)G(-\frac{\alpha}{2} + n - j + 1)G(j + 1)G(j + \frac{\alpha}{2} + 1)}{G(-\frac{\alpha}{2} + n - 2j + 1)G(\frac{\alpha}{2} - n + 2j + 1)} \Gamma(j + 1)\Gamma\left(-\frac{\alpha}{2} + n - j + 1\right). \end{split}$$

Combining these formulas we confirm the second case. Notice that for $c(\alpha - 2, n + 1)$ we had to shift index j to j + 1 in (59) to get correct formula.

For $2n - 4j - 2 < \operatorname{Re}(\alpha) < 2n - 4j$ we have

$$\begin{split} c(\alpha,n) &= (-1)^{\frac{n(n-1)}{2} + nj + n - j} \left(\frac{d_2}{\pi}\right)^n \left(\frac{d_1}{d_2} \sin\left(\frac{\pi\alpha}{2}\right) + \cos\left(\frac{\pi\alpha}{2}\right)\right)^j \\ &\times \frac{G(n-j+1)G(-\frac{\alpha}{2} + n - j + 1)G(j+1)G(j+\frac{\alpha}{2} + 1)}{G(-\frac{\alpha}{2} + n - 2j + 1)G(\frac{\alpha}{2} - n + 2j + 1)}, \\ c(\alpha-2,n) &= (-1)^{\frac{n(n-1)}{2} + nj + 2n} \left(\frac{d_2}{\pi}\right)^n \left(\frac{d_1}{d_2} \sin\left(\frac{\pi\alpha}{2}\right) + \cos\left(\frac{\pi\alpha}{2}\right)\right)^{j+1} \\ &\times \frac{G(n-j+1)G(-\frac{\alpha}{2} + n - j + 1)G(j+1)G(j+\frac{\alpha}{2} + 1)}{G(-\frac{\alpha}{2} + n - 2j + 1)G(\frac{\alpha}{2} - n + 2j + 1)} \frac{\Gamma(j+1)\Gamma(-\frac{\alpha}{2} + n - 2j)}{\Gamma(n-j)\Gamma(\frac{\alpha}{2} - n + 2j + 1)}, \\ c(\alpha, n+1) &= (-1)^{\frac{n(n+1)}{2} + nj + 2n + 1} \left(\frac{d_2}{\pi}\right)^{n+1} \left(\frac{d_1}{d_2} \sin\left(\frac{\pi\alpha}{2}\right) + \cos\left(\frac{\pi\alpha}{2}\right)\right)^{j+1} \\ &\times \frac{G(n-j+1)G(-\frac{\alpha}{2} + n - j + 1)G(j+1)G(j+\frac{\alpha}{2} + 1)}{G(-\frac{\alpha}{2} + n - 2j + 1)G(\frac{\alpha}{2} - n + 2j + 1)}, \\ c(\alpha-2, n+1) &= (-1)^{\frac{n(n+1)}{2} + nj + 2n + j + 2} \left(\frac{d_2}{\pi}\right)^{n+1} \left(\frac{d_1}{d_2} \sin\left(\frac{\pi\alpha}{2}\right) + \cos\left(\frac{\pi\alpha}{2}\right)\right)^{j+1} \\ &\times \frac{G(n-j+1)G(-\frac{\alpha}{2} + n - j + 1)G(j+1)G(j+\frac{\alpha}{2} + 1)}{G(-\frac{\alpha}{2} + n - 2j + 1)G(\frac{\beta}{2} - n + 2j + 1)} \Gamma(j+1)\Gamma\left(-\frac{\alpha}{2} + n - 2j\right). \end{split}$$

Tha confirms the third case. Notice that now we had to shift index j to j+1 in (59) for $c(\alpha-2, n+1)$, $c(\alpha, n+1)$, and $c(\alpha-2, n)$ to get correct formula.

Finally for $\operatorname{Re}(\alpha) < -2n$ we have

$$c(\alpha, n) = \frac{(-1)^{\frac{n(n-1)}{2} + n^2}}{\pi^n} \left(d_1 \sin\left(\frac{\pi\alpha}{2}\right) + d_2 \cos\left(\frac{\pi\alpha}{2}\right) \right)^n \frac{G(n+1)G(-\frac{\alpha}{2}+1)}{G(-\frac{\alpha}{2} - n + 1)},$$

$$c(\alpha - 2, n) = \frac{(-1)^{\frac{n(n-1)}{2} + n^2 + n}}{\pi^n} \left(d_1 \sin\left(\frac{\pi\alpha}{2}\right) + d_2 \cos\left(\frac{\pi\alpha}{2}\right) \right)^n \frac{G(n+1)G(-\frac{\alpha}{2}+1)}{G(-\frac{\alpha}{2} - n + 1)} \frac{\Gamma(-\frac{\alpha}{2} + 1)}{\Gamma(-\frac{\alpha}{2} - n + 1)},$$

$$c(\alpha, n+1) = \frac{(-1)^{\frac{(n+1)n}{2} + (n+1)^2}}{\pi^{n+1}} \left(d_1 \sin\left(\frac{\pi\alpha}{2}\right) + d_2 \cos\left(\frac{\pi\alpha}{2}\right) \right)^{n+1} \\ \times \frac{G(n+1)G(-\frac{\alpha}{2}+1)}{G(-\frac{\alpha}{2}-n+1)} \Gamma(n+1)\Gamma\left(-\frac{\alpha}{2}-n\right), \\ c(\alpha-2, n+1) = \frac{(-1)^{\frac{(n+1)n}{2} + n+1 + (n+1)^2}}{\pi^{n+1}} \left(d_1 \sin\left(\frac{\pi\alpha}{2}\right) + d_2 \cos\left(\frac{\pi\alpha}{2}\right) \right)^{n+1} \\ \times \frac{G(n+1)G(-\frac{\alpha}{2}+1)}{G(-\frac{\alpha}{2}-n+1)} \Gamma(n+1)\Gamma\left(-\frac{\alpha}{2}+1\right).$$

That implies

$$q(\alpha, n) = \frac{\Gamma\left(-\frac{\alpha}{2} - n + 1\right)}{\Gamma\left(-\frac{\alpha}{2} - n\right)} = -\frac{\alpha}{2} - n.$$

That confirms the last case.

6. Asymptotics at infinity

6.1. Asymptotics of Toeplitz determinant. To compute the asymptotics at infinity it is convenient to use the Hankel functions instead of Bessel functions. We rewrite formula (3) as (67). Following the same argument as in Theorem 4.1 we get the following multiple integral representation using Andrèief identity.

Theorem 6.1. The Toeplitz determinant $\Delta_n(x, \alpha)$ given by (2) can be rewritten as

$$\Delta_n(x,\alpha) = \frac{1}{n!} \int_{\Gamma_3 \cup \Gamma_4} \dots \int_{\Gamma_3 \cup \Gamma_4} \prod_{1 \le j < k \le n} (t_k - t_j) \left(\frac{1}{t_k} - \frac{1}{t_j}\right) \prod_{k=1}^n h_2(t_k) dt_k$$

where

$$h_2(t) = \frac{\mathrm{e}^{\frac{x}{2}(t-\frac{1}{t})}}{2\pi \mathrm{i} t^{1+\frac{\alpha}{2}}} \left((d_1 - \mathrm{i} d_2) \chi_{\Gamma_3}(t) - (d_1 + \mathrm{i} d_2) \chi_{\Gamma_4}(t) \right),$$

and contours of integration Γ_3 , Γ_4 are shown on Figure 9. We use the notation $\chi_{\Gamma_j}(t)$ for the characteristic function of contour Γ_j . We assume $-\pi < \arg(t) < \pi$ on contour Γ_3 and Γ_4 .

We would like to compute the large x asymptotics of $\Delta_n(x, \alpha)$. We start by using the expansion (42) and implementing notation (68)

$$\Delta_n(x,\alpha) = \frac{(-1)^{\frac{n(n-1)}{2}}}{n!} \int_{\Gamma_3 \cup \Gamma_4} \dots \int_{\Gamma_3 \cup \Gamma_4} \prod_{m=1}^n (b_1 \chi_{\Gamma_3}(t_m) - b_2 \chi_{\Gamma_4}(t_m)) \prod_{1 \le j < k \le n} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{l=1}^n \frac{e^{\frac{x}{2}(t_l - \frac{1}{t_l})}}{\pi i t_l^{\frac{\alpha}{2} + 1}} dt_l$$

$$= \frac{(-1)^{\frac{n(n-1)}{2}}}{(\pi i)^n n!} \sum_{I \subset \{1,\dots,n\}} b_1^r (-b_2)^{n-r} \int_{\Gamma_3} \dots \int_{\Gamma_3} \int_{\Gamma_4} \dots \int_{\Gamma_4} \prod_{\substack{j < k \\ j,k \in I}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j < k \\ j,k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}} \frac{(t_j - t_k)^2}{t_j t_k} \prod_{\substack{j \in I \\ k \in I^c}}$$

We can see that terms in the right hand side only depend on cardinality of I.

We would like to compute asymptotics of multiple integrals in (60) using steepest descent method. The critical points of the exponent $\Xi(t) = t - \frac{1}{t}$ are $t = \pm i$. We chose the contours Γ_3 and Γ_4 as contours of steepest descent $\operatorname{Im}(\Xi(t)) = \operatorname{Im}(\Xi(\pm i))$. They can be described using cubic equations $x^2(y \pm 2) + y(y \pm 1)^2 = 0$. The main contribution to the $x \to \infty$ asymptotics comes from the neighborhoods of the critical points. We make the change of variable in the local integrals: $(t - i) = se^{\frac{3\pi i}{4}}$ for $t \in I$ and $(t + i) = se^{-\frac{3\pi i}{4}}$ for $t \in I^c$. After that we replace the local integrals with real line integrals. The result is the following

$$\Delta_{n}(x,\alpha) \sim \sum_{r=0}^{n} \frac{b_{1}^{r} b_{2}^{n-r}}{r!(n-r)!} \frac{4^{r(n-r)}}{\pi^{n}} (-1)^{\frac{(n-r)(n+r+1)+n(n-1)}{2}} i^{\frac{r(r-1)+(n-r)(n-r-1)-2n}{2}} e^{\frac{\pi i}{4}(2r-n)(1-\alpha)} e^{ix(2r-n)}$$
$$\int_{\Gamma_{3}} \dots \int_{\Gamma_{3}} \prod_{1 \le j < k \le r} (s_{j} - s_{k})^{2} \prod_{l=1}^{r} e^{-\frac{x}{2}s_{l}^{2}} dt_{l} \int_{\Gamma_{4}} \dots \int_{\Gamma_{4}} \prod_{1 \le j < k \le n-r} (s_{j} - s_{k})^{2} \prod_{l=1}^{n-r} e^{-\frac{x}{2}s_{l}^{2}} dt_{l}, \quad x \to \infty$$

We make the change of variables in the integrals $s \to \frac{s}{\sqrt{x}}$ and evaluate the resulting integrals using standard formula [DLMF, (5.14.6)] rewritten in terms of Barnes *G*-function.

$$\begin{split} \Delta_n(x,\alpha) &\sim \sum_{r=0}^n \frac{b_1^r b_2^{n-r}}{r!(n-r)!} \left(\frac{2}{\pi}\right)^{\frac{n}{2}} 4^{r(n-r)} (-1)^{\frac{(n-r)(n+r+1)+n(n-1)}{2}} i^{\frac{r(r-1)+(n-r)(n-r-1)-2n}{2}} \\ &\times G(r+2)G(n-r+2) \mathrm{e}^{\frac{\pi \mathrm{i}}{4}(2r-n)(1-\alpha)} \mathrm{e}^{\mathrm{i}(2r-n)x} x^{-r^2+nr-\frac{n^2}{2}}, \quad x \to \infty \end{split}$$

To find the leading term we observe that

$$\max_{0 \le r \le n} \left(-r^2 + nr - \frac{n^2}{2} \right) = \begin{cases} -\frac{n^2}{4} & \text{when } n \text{-even using } r = \frac{n}{2}, \\ -\frac{n^2}{4} - \frac{1}{4} & \text{when } n \text{-odd using } r = \frac{n \pm 1}{2}. \end{cases}$$

For even n we plug in $r = \frac{n}{2}$. Many terms cancel out. We combine factorials with Barnes G-function. The power of i is even, so we combine it with power of -1. As the result all terms with i and -1 cancel out. For odd n we need to combine contributions from $r = \frac{n-1}{2}$ and $r = \frac{n+1}{2}$.

$$\Delta_n(x,\alpha) \sim \left(\frac{d_1^2 + d_2^2}{2\pi}\right)^{\frac{n}{2}} G\left(\frac{n+1}{2}\right) G\left(\frac{n+3}{2}\right) \left(\frac{x}{4}\right)^{-\frac{n^2+1}{4}} (-1)^{\frac{7n^2+1}{8}} i^{\frac{n^2}{4} - \frac{3n}{2} + \frac{1}{4}} \\ \times \frac{1}{2} \left(e^{i\frac{\pi}{4}(\alpha - 1) - ix + i\phi} + (-1)^{\frac{n+1}{2}} e^{i\frac{\pi}{4}(1 - \alpha) + ix - i\phi}\right) \\ = \left(\frac{d_1^2 + d_2^2}{2\pi}\right)^{\frac{n}{2}} G\left(\frac{n+1}{2}\right) G\left(\frac{n+3}{2}\right) \left(\frac{x}{4}\right)^{-\frac{n^2+1}{4}} (-1)^{\frac{n-1}{2}} \sin\left(x - \phi + \frac{\pi}{4}(n - \alpha)\right)$$

where $\phi = \arg(d_1 + \mathrm{i}d_2)$.

24

As the result we get Theorem 1.3.

6.2. Asymptotics of special function solutions. In this part, we still need to use Proposition 1.1 and Theorem 1.3 to get the large x asymptotics of $u_n(x, \alpha)$. Fortunately, life becomes much easier in this scenario. Notice that when n is even, the Toeplitz determinant doesn't depend on α at all. Hence when shifting the indices, we have for even n:

$$u_n(x,\alpha) = -\frac{\Delta_{n+1}(x,\alpha-2)}{\Delta_{n+1}(x,\alpha)} = -\frac{\sin\left(x-\phi+\frac{\pi}{4}(n+1-\alpha)+\frac{\pi}{2}\right)}{\sin\left(x-\phi+\frac{\pi}{4}(n+1-\alpha)\right)} = -\cot\left(x-\phi+\frac{\pi}{4}(n+1-\alpha)\right)$$

Similarly, when n is odd and n + 1 is even, we get:

$$u_n(x,\alpha) = -\frac{\Delta_n(x,\alpha)}{\Delta_n(x,\alpha-2)} = -\frac{\sin\left(x - \phi + \frac{\pi}{4}(n-\alpha)\right)}{\sin\left(x - \phi + \frac{\pi}{4}(n-\alpha) + \frac{\pi}{2}\right)} = -\tan\left(x - \phi + \frac{\pi}{4}(n-\alpha)\right)$$

where $\phi = \arg(d_1 + \mathrm{i}d_2)$.

That proves Theorem 1.4.

Acknowledgment

Most of the work was performed during math REU program at University of Michigan in Summer 2023. H.P. would like to thank his mentor A.P. for his patient and professional instructions and training during the whole summer. Besides some more advanced contents in differential equations and asymptotic analysis, he also learned research skills in modern mathematics with A.P.. There were many insightful and smart ideas throughout the project, which really motivated and inspired him to explore the related topics deeply. He also would like to thank A.P's help on numerical check of the results and polishing the paper during the following whole year. This is also his first experience to tackle such a big problem, so without A.P.'s assistance, there is no way to generate this paper. He is really appreciated very much. H.P. would also thank Professor Lun Zhang from Fudan University and Professor Peter Miller from University of Michigan for useful discussions and suggestions. A.P. was supported NSF MSPRF grant DMS-2103354, and RSF grant 22-11-00070.

References

- [BLMP24] A. Barhoumi, O. Lisovyy, P. D. Miller, and A. Prokhorov. "Painlevé-III monodromy maps under the $D_6 \rightarrow D_8$ confluence and applications to the large-parameter asymptotics of rational solutions". In: SIGMA Symmetry Integrability Geom. Methods Appl. 20 (2024), Paper No. 019, 77. ISSN: 1815-0659. DOI: 10.3842/SIGMA.2024.019. MR: 4714353.
- [BMS18] T. Bothner, P. D. Miller, and Y. Sheng. "Rational solutions of the Painlevé-III equation".
 In: Stud. Appl. Math. 141.4 (2018), pp. 626–679. ISSN: 0022-2526. DOI: 10.1111/sapm.
 12220. MR: 3879971. arXiv: 1801.04360 [math.CA].
- [Cla23] P. A. Clarkson. "Classical solutions of the degenerate fifth Painlevé equation". In: J. Phys. A 56.13 (2023), Paper No. 134002, 23. ISSN: 1751-8113. MR: 4561285. arXiv: 2301.01727
 [nlin.SI].
- [CLL23] P. A. Clarkson, C.-K. Law, and C.-H. Lin. "A constructive proof for the Umemura polynomials of the third Painlevé equation". In: SIGMA Symmetry Integrability Geom. Methods Appl. 19 (2023), Paper No. 080, 20. ISSN: 1815-0659. DOI: 10.3842/SIGMA.2023.080. MR: 4659225.
- [CZ10] Y. Chen and L. Zhang. "Painlevé VI and the unitary Jacobi ensembles". In: Stud. Appl. Math. 125.1 (2010), pp. 91–112. ISSN: 0022-2526,1467-9590. DOI: 10.1111/j.1467-9590.
 2010.00483.x. MR: 2676782.

- [Dea18] A. Deaño. "Large z asymptotics for special function solutions of Painlevé II in the complex plane". In: SIGMA Symmetry Integrability Geom. Methods Appl. 14 (2018), Paper No. 107, 19. DOI: 10.3842/SIGMA.2018.107. MR: 3860606.
- [DLMF] NIST Digital Library of Mathematical Functions. http://dlmf.nist.gov/, Release 1.1.4 of 2022-01-15. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds. URL: http://dlmf.nist.gov/.
- [For19] P. J. Forrester. "Meet Andréief, Bordeaux 1886, and Andreev, Kharkov 1882–1883". In: *Random Matrices Theory Appl.* 8.2 (2019), pp. 1930001, 9. ISSN: 2010-3263,2010-3271. DOI: 10.1142/S2010326319300018. MR: 3943898. arXiv: 1806.10411 [math-ph].
- [FW02] P. J. Forrester and N. S. Witte. "Application of the τ-function theory of Painlevé equations to random matrices: P_V, P_{III}, the LUE, JUE, and CUE". In: *Comm. Pure Appl. Math.* 55.6 (2002), pp. 679–727. ISSN: 0010-3640. DOI: 10.1002/cpa.3021. MR: 1885665. arXiv: math-ph/0201051 [math-ph].
- [Gam10] B. Gambier. "Sur les équations différentielles du second ordre et du premier degré dont l'intégrale générale est a points critiques fixes". In: Acta Math. 33.1 (1910), pp. 1–55. ISSN: 0001-5962,1871-2509. DOI: 10.1007/BF02393211. MR: 1555055.
- [Inc44] E. L. Ince. Ordinary Differential Equations. Dover Publications, New York, 1944, pp. viii+558. MR: 10757.
- [Jim82] M. Jimbo. "Monodromy problem and the boundary condition for some Painlevé equations". In: Publ. Res. Inst. Math. Sci. 18.3 (1982), pp. 1137–1161. ISSN: 0034-5318. DOI: 10.2977/prims/1195183300. MR: 688949.
- [Kit87] A. V. Kitaev. "The method of isomonodromic deformations and the asymptotics of the solutions of the "complete" third Painlevé equation". In: Mat. Sb. (N.S.) 134(176).3 (1987), pp. 421–444, 448. ISSN: 0368-8666. DOI: 10.1070/SM1989v062n02ABEH003247. MR: 922633.
- [Mur95] Y. Murata. "Classical solutions of the third Painlevé equation". In: Nagoya Math. J. 139 (1995), pp. 37–65. ISSN: 0027-7630,2152-6842. DOI: 10.1017/S0027763000005298. MR: 1355268.
- [Oka87] K. Okamoto. "Studies on the Painlevé equations. IV. Third Painlevé equation P_{III}". In: Funkcial. Ekvac. 30.2-3 (1987), pp. 305-332. ISSN: 0532-8721. MR: 927186. URL: http: //www.math.kobe-u.ac.jp/~fe/xml/mr0927186.xml.
- [Pai02] P. Painlevé. "Sur les équations différentielles du second ordre et d'ordre supérieur dont l'intégrale générale est uniforme". In: Acta Math. 25.1 (1902), pp. 1–85. ISSN: 0001-5962,1871-2509. DOI: 10.1007/BF02419020. MR: 1554937.
- [UW98] H. Umemura and H. Watanabe. "Solutions of the third Painlevé equation. I". In: Nagoya Math. J. 151 (1998), pp. 1–24. ISSN: 0027-7630,2152-6842. DOI: 10.1017/S0027763000025149. MR: 1650348.
- [VV23] J. Vrbik and P. Vrbik. "A Novel Proof of the Desnanot-Jacobi Determinant Identity". In: Mathematics Magazine 0.0 (2023), pp. 1–5. DOI: 10.1080/0025570X.2023.2199721.
- [ZCL20] M. Zhu, Y. Chen, and C. Li. "The smallest eigenvalue of large Hankel matrices generated by a singularly perturbed Laguerre weight". In: J. Math. Phys. 61.7 (2020), pp. 073502, 12. ISSN: 0022-2488,1089-7658. DOI: 10.1063/1.5140079. MR: 4118329.

APPENDIX A. BESSEL EQUATION AND ITS SOLUTIONS

Bessel equation is given by

$$u''(x) + \frac{u'(x)}{x} + \left(1 - \frac{\nu^2}{x^2}\right)u(x) = 0.$$
(61)

One of the standard solutions in the form of series representation is given by (see [DLMF, (10.2.2)])

$$J_{\nu}(x) = \frac{x^{\nu}}{2^{\nu}} \sum_{k=0}^{\infty} \frac{(-1)^k x^{2k}}{2^{2k} k! \Gamma(\nu + k + 1)}$$
(62)

where $\Gamma(x)$ is Gamma function.

A.1. Contour integral representations. The contour integral representation for $-\frac{\pi}{2} < \arg(x) < \frac{\pi}{2}$ is given by (see [DLMF, (10.9.17)])

$$J_{\nu}(x) = \frac{1}{2\pi i} \int_{\infty-i\pi}^{\infty+i\pi} e^{x\sinh(z)-\nu z} dz,$$
(63)

The contour of integration in (63) is shown on Figure 6.



FIGURE 6. Contour for $J_{\nu}(x)$.

Bessel function of second kind can be written as (see [DLMF, (10.2.3)])

$$Y_{\nu}(x) = \cot(\pi\nu) J_{\nu}(x) - \csc(\pi\nu) J_{-\nu}(x)$$

By formula (63) of $J_{\nu}(x)$ it follows that cylinder function $\mathcal{C}_{\nu}(x)$ that we introduced in (3) can be written as

$$\mathcal{C}_{\nu}(x) = (d_1 + d_2 \cot(\pi\nu)) \frac{1}{2\pi i} \int_{\infty - i\pi}^{\infty + i\pi} e^{x \sinh(z) - \nu z} dz - d_2 \csc(\pi\nu) \frac{1}{2\pi i} \int_{\infty - i\pi}^{\infty + i\pi} e^{x \sinh(z) + \nu z} dz$$

Making the change of variable $z \to i\pi - z$ in the second integral we get

$$\mathcal{C}_{\nu}(x) = (d_1 + d_2 \cot(\pi\nu)) \frac{1}{2\pi i} \int_{\infty - i\pi}^{\infty + i\pi} e^{x \sinh(z) - \nu z} dz + d_2 \csc(\pi\nu) \frac{e^{i\pi\nu}}{2\pi i} \int_{-\infty + 2i\pi}^{-\infty} e^{x \sinh(z) - \nu z} dz$$

Making another change of variable $e^z = t$, we get

$$\mathcal{C}_{\nu}(x) = \int_{\Gamma_1 \cup \Gamma_2} \frac{\mathrm{e}^{\frac{x}{2}\left(t - \frac{1}{t}\right)}}{2\pi \mathrm{i}t^{1+\nu}} \left(\left(d_1 + d_2 \cot(\pi\nu)\right) \chi_{\Gamma_1}(t) + d_2 \csc(\pi\nu) \mathrm{e}^{\mathrm{i}\pi\nu} \chi_{\Gamma_2}(t) \right) dt = \int_{\Gamma_1 \cup \Gamma_2} h_1(t) dt \qquad (64)$$

Here Γ_1 is contour of integration for $J_{\nu}(x)$ and Γ_2 is contour of integration for $J_{-\nu}(x)$ after change of variable $e^z = t$ shown in the Figure 7. We should make a remark about power function $\frac{1}{t^{1+\nu}}$. When we use it, we assume $-i\pi < \arg(t) < i\pi$ on contour Γ_1 and $0 < \arg(t) < 2\pi i$ on contour Γ_2 .

Remark A.1. For the case $\alpha \in 2\mathbb{Z}$ we need to use formula $Y_n(x) = \frac{1}{\pi} \left. \frac{\partial J_{\nu}}{\partial \nu} \right|_{\nu=n} + \frac{(-1)^n}{\pi} \left. \frac{\partial J_{\nu}}{\partial \nu} \right|_{\nu=-n}$. It would mean that we need to consider weight $e^{-t}t^a \ln(t)$. To our knowledge, such weight is not classical and it does not have explicit formula computed for the multiple integral.

Alternatively, cylinder function $C_{\nu}(x)$ can be written in terms Hankel functions, which defined as (see [DLMF, (10.4.3)])

$$H_{\nu}^{(1)}(x) = J_{\nu}(x) + iY_{\nu}(x), \qquad \qquad H_{\nu}^{(2)}(x) = J_{\nu}(x) - iY_{\nu}(x). \tag{65}$$



FIGURE 7. Contours for $C_{\nu}(x)$.

The contour integral representations for $-\frac{\pi}{2} < \arg(x) < \frac{\pi}{2}$ are given by (see [DLMF, (10.9.18)])

$$H_{\nu}^{(1)}(x) = \frac{1}{\pi i} \int_{-\infty}^{\infty + i\pi} e^{x \sinh(z) - \nu z} dz, \qquad \qquad H_{\nu}^{(2)}(x) = -\frac{1}{\pi i} \int_{-\infty}^{\infty - i\pi} e^{x \sinh(z) - \nu z} dz. \tag{66}$$

The contours of integration in (66) are shown on Figure 8.



(A) Contour for
$$H_{\nu}^{(1)}$$
 (B) Contours for $H_{\nu}^{(2)}$

FIGURE 8. Contours of integration for Hankel functions.

We would like to also get the alternative integral representation for $C_{\nu}(x)$. Using (65) we have

$$\mathcal{C}_{\nu}(x) = b_1 H_{\nu}^{(1)}(x) + b_2 H_{\nu}^{(2)}(x) \tag{67}$$

where

$$b_1 = \frac{d_1 - \mathrm{i}d_2}{2}, \quad b_2 = \frac{d_1 + \mathrm{i}d_2}{2}.$$
 (68)

Making a change of variable $e^z = t$, we get

$$\mathcal{C}_{\nu}(x) = \int_{\Gamma_{3}\cup\Gamma_{4}} \frac{\mathrm{e}^{\frac{x}{2}\left(t-\frac{1}{t}\right)}}{2\pi\mathrm{i}t^{1+\nu}} ((d_{1}-\mathrm{i}d_{2})\chi_{\Gamma_{3}}(t) - (d_{1}+\mathrm{i}d_{2})\chi_{\Gamma_{4}}(t))dt = \int_{\Gamma_{3}\cup\Gamma_{4}} h_{2}(t)dt, \tag{69}$$

where contours Γ_3 and Γ_4 are shown on the Figure 9.

A.2. Differential identities for cylinder functions. Cylinder functions $C_{\nu}(x)$ satisfy the following differential identities (see [DLMF, (10.6.2)]):

$$C'_{\nu}(x) = \frac{\nu}{x} C_{\nu}(x) - C_{\nu+1}(x),$$
(70)

$$C'_{\nu}(x) = C_{\nu-1}(x) - \frac{\nu}{x}C_{\nu}(x).$$
(71)



FIGURE 9. Alternative contours of integration for $C_{\nu}(x)$.