Coupled Stochastic-Statistical Equations for Filtering Multiscale Turbulent Systems

Di Qi^a and Jian-Guo Liu^b

July 9, 2024

^aDepartment of Mathematics, Purdue University, 150 North University Street, West Lafayette, IN 47907, USA

^bDepartment of Mathematics and Department of Physics, Duke University, Durham, NC 27708, USA

Abstract

We present a new strategy for filtering high-dimensional multiscale systems characterized by high-order non-Gaussian statistics using observations from leading-order moments. A closed stochastic-statistical modeling framework suitable for systematic theoretical analysis and efficient numerical simulations is designed. Optimal filtering solutions are derived based on the explicit coupling structures of stochastic and statistical equations subject to linear operators, which satisfy an infinite-dimensional Kalman-Bucy filter with conditional Gaussian dynamics. To facilitate practical implementation, we develop a finite-dimensional stochastic filter model that approximates the optimal filter solution. We prove that this approximating filter effectively captures key non-Gaussian features, demonstrating consistent statistics with the optimal filter. Finally, we build a practical ensemble filter algorithm based on the approximating filtering model, which enables accurate recovery of the true model statistics. The proposed modeling and filtering strategies are applicable to a wide range challenging problems in science and engineering, particularly for statistical prediction and uncertainty quantification of multiscale turbulent states.

1 Introduction

Complex turbulent phenomena are widely observed in various science and engineering systems [51, 14, 42]. Such systems are characterized by a wide spectrum of nonlinearly coupled spatiotemporal scales and high degrees of inherent internal instability [37, 21]. A probabilistic formulation containing highly non-Gaussian statistics is required to quantify uncertainties in the high-dimensional turbulent state [44, 38]. Traditional ensemble approaches using a particle system representation to approximate the probability evolution quickly become computationally prohibitive since a sufficiently large sample size is necessary to capture the extreme non-Gaussian outliers even for relatively low dimensional systems [35, 55]. As a result, rigorous analysis often becomes intractable and direct numerical simulations are likely to be expensive and inaccurate.

Filtering strategies [50, 33, 52] have long been used for finding the optimal probability estimates of a stochastic state with large uncertainty based on partial and noisy observation data. Applications to improve the probability forecast can be found in various practical dynamical systems [23, 29, 3, 5]. In predicting nonlinear turbulent signals, ensemble Kalman filters [18, 26, 19] as well as the related particle methods [13, 15, 7] provide an effective tool for state and parameter estimations. Filtering theories [58, 8, 45] and corresponding numerical solutions [32, 4, 9] for general nonlinear systems have been investigated through different approaches. Despite wide applications [17, 40, 12], difficulties persist for accurate statistical forecast of turbulent states especially when non-Gaussian features are present in the target probability distribution. Conventional ensemble-based approaches often suffer inherent difficulties in estimating the crucial higher-order moment statistics and maintaining stable prediction with finite number of particles [53, 24, 54]. On the other hand, in many situations observations of the statistical states, such as the mean and covariance, are easier to extract from various sources of data [39]. Therefore, a promising research direction is to propose new ensemble filtering models to recover the highly non-Gaussian probability distributions using statistical observations from the leading-order mean and covariance [1].

1.1 General problem setup

We start with a general mathematical formulation [38] modeling a high-dimensional stochastic state variable $u_t \in \mathbb{R}^d$ involving nonlinear multiscale interactions satisfying the following stochastic differential equation (SDE)

$$\frac{\mathrm{d}u_t}{\mathrm{d}t} = \Lambda u_t + B\left(u_t, u_t\right) + F_t + \sigma_t \dot{W}_t.$$
(1.1)

On the right hand side of the above equation, the linear operator, $\Lambda : \mathbb{R}^d \to \mathbb{R}^d$, represents linear dispersion and dissipation effects. The nonlinear effect in the dynamical system is introduced via a bilinear quadratic operator, $B : \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}^d$. The system is subject to external forcing effects that are decomposed into a deterministic component, F_t , and a stochastic component represented by a Gaussian white noise $W_t \in \mathbb{R}^s$ with $\sigma_t \in \mathbb{R}^{d \times s}$. The model emphasizes the important role of quadratic interactions through B(u, u). This typical structure is inherited from a finite-dimensional truncation of the continuous system, for example, a spectral projection of the nonlinear advection in fluid model [34]. Many realistic systems with wide applications [27, 48, 22] can be categorized in the general dynamical equation (1.1).

The evolution of the model state u_t depends on the sensitivity to the randomness in initial conditions and external stochastic effects, which will be further amplified in time by the inherent internal instability due to the nonlinear coupling term [37, 49]. Assuming that the stochastic solution satisfies a continuous probability density function (PDF), the time evolution of the PDF p_t is given by the associated Fokker-Planck equation (FPE)

$$\frac{\partial p_t}{\partial t} = \mathcal{L}_{\text{FP}} p_t \coloneqq -\nabla_u \cdot \left[\left(\Lambda u + B\left(u, u\right) + F_t \right) p_t \right] + \frac{1}{2} \nabla_u \cdot \left[\nabla_u \cdot \left(\sigma_t \sigma_t^{\mathsf{T}} p_t \right) \right], \ p_{t=0} = p_0, \tag{1.2}$$

where \mathcal{L}_{FP} represents the Fokker-Planck operator with $\nabla \cdot (\nabla \cdot A) = \sum_{k,l} \frac{\partial^2 A_{kl}}{\partial u_k \partial u_l}$. However, it remains a challenging task for directly solving the FPE (1.2) as a high-dimensional PDE. As an alternative approach, ensemble forecast by tracking the Monte-Carlo solutions estimates the essential statistics through empirical averages among a group of samples drawn i.i.d. from the initial distribution $u^{(i)}(0) \sim p_0$ at the starting time t = 0. The PDF solution $p_t(u)$ and the associated statistical expectation of any test function $\varphi(u)$ at each time instant t > 0 are then approximated by the empirical ensemble representation

$$p_t(u) \simeq p_t^N(u) \coloneqq \frac{1}{N} \sum_{i=1}^N \delta\left(u - u_t^{(i)}\right), \quad \mathbb{E}_{p_t}\varphi\left(u\right) \simeq \mathbb{E}_{p_t^N}\varphi\left(u\right) = \frac{1}{N} \sum_{i=1}^N \varphi\left(u_t^{(i)}\right), \tag{1.3}$$

In practice, large errors will be introduced in the above empirical estimations since only a finite sample approximation is available in modeling the probability distribution and statistics in a high dimensional system.

It is expected that the model errors in the predicted PDF of the stochastic states from the finite ensemble simulation can be effectively corrected through the available observation data. In designing new filtering strategies, we propose to use statistical observations from mean and covariance to improve the accuracy and stability in the finite ensemble forecast using the finite ensemble approximation. However, the general formulation (1.1) as well as the associated FPE (1.2) becomes inconvenient to use since all the multiscale stochastic processes in the equation are mixed together containing all high-order statistics. The main goal of this paper is thus to develop a systematic modeling framework with strategies to accurately capture the (potentially highly non-Gaussian) PDF p_t with the help of statistical measurements in the leading moments.

1.2 Overview of the paper

In this paper, we study nonlinear filtering of the general multiscale turbulent system (1.1) according to the two-step filtering procedure. In the first forecast step, a new modeling framework is designed combining statistical equations (2.3) for the leading moments and stochastic equations (2.5) for capturing higher-order statistics. In the second analysis step, we propose the observation process from the statistical equations of the leading-order mean and covariance to improve the probability prediction of the stochastic model. The statistical equations for observation processes (3.1b) then become linearly dependent on the PDF of stochastic state, while the evolution of the PDF is also subject to conditional linear dynamics (3.1a). Therefore, optimal filtering equations can be explicitly derived based on the linearity in both signal and observation processes. Finally, the expensive optimal filter equations are



Figure 1.1: Diagram illustrating main ideas in constructing the modeling framework

approximated by an efficient filtering model demonstrating equivalent statistics according to the most important observation functions (3.4).

The new multiscale nonlinear filtering model is constructed under the following step-by-step procedure, which will finally be combined to build a practical ensemble filtering strategy able to recover non-Gaussian PDFs:

- First, we propose a precise coupled stochastic-statistical formulation (2.7) as the forecast model for multiscale turbulence: the stochastic dynamics will serve as the signal process in filtering including high-order non-Gaussian features, while the reinforced statistical equations provide the observation process;
- Second, the optimal filtering problem is formulated based on the particular coupling structure in the multiscale stochastic-statistical model: the optimal filter solution (3.12) is given based on the PDF of the stochastic state, combined with the mean and covariance as a natural choice of the observed state;
- Third, an effective statistical filtering model (3.16) is developed approximating the optimal filter solution in key leading-order statistics: an equivalent McKean-Vlasov SDE containing higher moments feedbacks is adapted from the optimal filter solution;
- Last, discrete ensemble filtering schemes (5.2) are constructed as a particle approximation of the statistical filtering model: the filter SDE is approximated by interacting particles and the statistical moments are computed by empirical ensemble averages.

The coupled stochastic-statistical model (2.7) by itself can serve as an effective tool for statistical forecasts and uncertainty quantification. Efficient computational algorithms have been developed [47, 48] that demonstrate high skill in capturing crucial non-Gaussian phenomena such as extreme events and fat-tailed PDFs. Then combined with the observation data, the resulting filtering McKean-Vlasov SDE (3.16) is only implicitly dependent on the probability distribution, which can be computed directly from the statistical equations. This enables efficient computational strategies to effectively improve the accuracy and stability in capturing high-order non-Gaussian features based on only observation from the lower moments. We illustrate the main steps in building effective models for capturing probability distributions in Fig. 1.1.

Our main focus of this paper is to provide a detailed description of the forecast and analysis step of the filtering method so that various applications can directly follow based on this highly adaptive general framework. We provide detailed discussion in each component of the proposed model and methods: i) we show in Proposition 2 that the highly tractable closure model (2.7) demonstrates consistent statistics as the original system (1.1) which plays a fundamental role in the construction of the filtering methods; ii) the filter solution (3.12) is shown optimal in the mean square sense by exploiting the conditional Gaussian structure of the forward equation, and the approximation model is shown to recover the same key statistics during the analysis step update in Theorem 7; iii) the long-time convergence in statistics to the optimal filter is demonstrated in Theorem 9 concerning the complete filtering procedure using the statistical filtering model (3.16); and iv) the ensemble approximation with discrete time step is shown to recover to the true statistics at the large ensemble limit.

The structure of the paper is organized as follows: In Section 2, we first set up a statistically consistent formulation for the general multiscale system (1.1) that is suitable for the construction of the statistical filtering models. Section 3 shows the major ideas of finding the optimal filter solution and constructing approximate filtering models. Long-time convergence and stability of the approximation filter model is then discussed in Section 4. Combining each component of the ideas, the final ensemble statistical filtering model is developed in Section 5 Finally, a summary of this paper is given in Section 6.

2 A statistically consistent modeling framework for multiscale dynamical systems

In this section, we first introduce a new formulation for the original system (1.1) using an explicit macroscopic and microscopic decomposition of the multiscale state. In particular, we show that the new formulation provides consistent statistics with the original system in the explicit leading moment equations, as well as higher-order statistics in the stochastic equation. In addition, the new formulation also enjoys a more tractable dynamical structure to be adapted to the filtering framework.

2.1 The statistical and stochastic equations as macroscopic and microscopic states

In order to identify the detailed multiscale interactions in the general system (1.1), we decompose the random model state u_t into the composition of a statistical mean \bar{u}_t and stochastic fluctuations u'_t in a finite-dimensional representation under an orthonormal basis $\{\hat{v}_k\}_{k=1}^d$ with $\hat{v}_k \cdot \hat{v}_l = \delta_{kl}$

$$u_t = \bar{u}_t + u'_t = \sum_{k=1}^d \bar{u}_{k,t} \hat{v}_k + \sum_{k=1}^d Z_{k,t} \hat{v}_k, \quad \text{with } \bar{u}_{k,t} = \hat{v}_k \cdot \bar{u}_t, \ Z_{k,t} = \hat{v}_k \cdot (u_t - \bar{u}_t).$$
(2.1)

Above, $\bar{u}_t \in \mathbb{R}^d$ represents the dominant largest-scale structure (for example, the zonal jets in geophysical turbulence or the coherent radial flow in fusion plasmas); and $Z_t = [Z_{1,t}, \cdots, Z_{d,t}]^{\mathsf{T}} \in \mathbb{R}^d$ are stochastic coefficients projected on each eigenmode \hat{v}_k , whose randomness illustrates the uncertainty in each single scale of u'_t . In particular, we will show that the dynamics of the stochastic modes Z_t contain nonlinear interactions among a large number of coupled multiscale fluctuations, which demonstrate the charactering feature in turbulent systems demonstrating forward and backward energy cascades [21, 37].

2.1.1 Statistical equations for the macroscopic states

First, we define the leading-order mean and covariance according to the state decomposition (2.1)

$$\bar{u}_t = \mathbb{E}_{p_t} \left[u \right] \coloneqq \int_{\mathbb{R}^d} u p_t \left(\mathrm{d}u \right), \quad R_{kl,t} = \mathbb{E}_{p_t} \left[Z_{k,t} Z_{l,t} \right] \coloneqq \int_{\mathbb{R}^d} \hat{v}_k \cdot \left(u - \bar{u}_t \right) \left(u - \bar{u}_t \right) \cdot \hat{v}_l p_t \left(\mathrm{d}u \right), \ 1 \le k, l \le d, \quad (2.2)$$

where the expectations are with respect to the law p_t from the PDF solution of (1.2). Above, statistical mean \bar{u}_t and covariance R_t can represent the *macroscopic* physical quantities that are easiest to achieve from direct measurements. The dynamical equations for the mean and covariance can be derived by the following statistical equations for $1 \le k, l \le d$

$$\frac{\mathrm{d}\bar{u}_{k,t}}{\mathrm{d}t} = \hat{v}_k \cdot \left[\Lambda \bar{u}_t + B\left(\bar{u}_t, \bar{u}_t\right)\right] + \sum_{m,n=1}^d \gamma_{kmn} \mathbb{E}_{p_t} \left[Z_{m,t} Z_{n,t}\right] + \hat{v}_k \cdot F_t,$$
(2.3a)

$$\frac{\mathrm{d}R_{kl,t}}{\mathrm{d}t} = \sum_{m=1}^{d} \left[L_{km} \left(\bar{u}_{t} \right) R_{t,ml} + R_{t,km} L_{lm} \left(\bar{u}_{t} \right) \right] + Q_{t,kl},$$

$$+ \sum_{m,n=1}^{d} \gamma_{kmn} \mathbb{E}_{p_{t}} \left[Z_{m,t} Z_{n,t} Z_{l,t} \right] + \gamma_{lmn} \mathbb{E}_{p_{t}} \left[Z_{m,t} Z_{n,t} Z_{k,t} \right].$$
(2.3b)

Above, we define the nonlinear coupling coefficients $\gamma_{kmn} = \hat{v}_k \cdot B(\hat{v}_m, \hat{v}_n)$, and the white noise coefficient is defined as $\Sigma_t = [(\hat{v}_1^{\mathsf{T}} \sigma_t)^{\mathsf{T}}, \cdots, (\hat{v}_d^{\mathsf{T}} \sigma_t)^{\mathsf{T}}]^{\mathsf{T}}$ with $Q_t = \Sigma_t \Sigma_t^{\mathsf{T}} \in \mathbb{R}^{d \times d}$. The mean-fluctuation coupling operator $L(\bar{u}_t) \in \mathbb{R}^{d \times d}$ dependent on the statistical mean state \bar{u}_t is defined as

$$L_{kl}(u) = \hat{v}_k \cdot [\Lambda \hat{v}_l + B(u, \hat{v}_l) + B(\hat{v}_l, u)].$$
(2.4)

Notice that the right hand side of (2.3b) involves the fluctuation modes Z_t defined from u_t in (2.1), then the expectations on third moments are taken w.r.t. the PDF p_t . Therefore, the resulting statistical equations (2.3) are not closed and need to be combined with the FPE (1.2) to achieve a closed formulation for the leading-order mean and covariance in the nonlinear system.

2.1.2 Stochastic equations for the microscopic processes

Second, we introduce the SDE describing the time evolution of the multiscale stochastic processes Z_t as the *microscopic* state consisting of the many subscale fluctuations

$$dZ_t = L\left(\bar{u}_t\right) Z_t dt + \Gamma \left(Z_t Z_t^{\mathsf{T}} - R_t\right) dt + \Sigma_t dW_t.$$
(2.5)

Above, $L(\bar{u}_t)$ is the same mean-fluctuation coupling operator defined in (2.4) involving the statistical mean \bar{u}_t , and we define the quadratic coupling operator $\Gamma : \mathbb{R}^{d \times d} \to \mathbb{R}^d$ as a linear combination of the entries of the input matrix $R \in \mathbb{R}^{d \times d}$ describing the nonlinear multiscale coupling involving the covariance R_t

$$\Gamma_k(R) = \sum_{m,n=1}^d \hat{v}_k \cdot B(\hat{v}_m, \hat{v}_n) R_{mn}.$$
(2.6)

Similar to the statistical equations, the dynamics on the right hand side of (2.5) is linked to the macroscopic quantities \bar{u}_t and R_t , which in tern requires additional information of p_t from the original model state u_t . This makes the stochastic equation also unclosed requiring additional information from the PDF solution (1.2) from the original system. The derivations of the above statistical and stochastic equations (2.3) and (2.5) from the original general formulation (1.1) are shown in the proof of Proposition 2.

2.2 A coupled stochastic-statistical closure model with explicit higher-order feedbacks

Combining the ideas in the closely related stochastic equation (2.5) and the statistical equations (2.3), we propose a statistically consistent stochastic-statistical closure model based on the following self-consistent coupling of the microscopic stochastic processes Z_t and the macroscopic statistics \bar{u}, R_t

$$dZ_{t} = L\left(\bar{u}_{t}\right) Z_{t}dt + \Gamma\left(Z_{t}Z_{t}^{\mathsf{T}} - R_{t}\right) dt + \Sigma_{t}dW_{t},$$

$$\frac{d\bar{u}_{t}}{dt} = M\left(\bar{u}_{t}\right) + Q_{m}\left(\mathbb{E}\left[Z_{t}\otimes Z_{t}\right]\right) + F_{t},$$

$$\frac{dR_{t}}{dt} = L\left(\bar{u}_{t}\right) R_{t} + R_{t}L\left(\bar{u}_{t}\right)^{\mathsf{T}} + Q_{v}\left(\mathbb{E}\left[Z_{t}\otimes Z_{t}\otimes Z_{t}\right]\right)$$

$$+ \Sigma_{t}\Sigma_{t}^{\mathsf{T}} + \epsilon^{-1}\left(\mathbb{E}\left[Z_{t}Z_{t}^{\mathsf{T}}\right] - R_{t}\right).$$
(2.7)

Above, the expectations are all w.r.t. the PDF ρ_t of the stochastic states Z_t . In the first moment equation for \bar{u}_t , with a bit abuse of notation, we denote $\bar{u}_t = [\bar{u}_{1,t}, \cdots, \bar{u}_{d,t}]^{\mathsf{T}} \in \mathbb{R}^d$ with each component $\bar{u}_{k,t} = \bar{u}_t \cdot \hat{v}_k$, $M = [M_1, \cdots, M_d]^{\mathsf{T}} \in \mathbb{R}^d$ where $M_k(\bar{u}_t) = \sum_{p,q} \hat{v}_k \cdot [\Lambda \hat{v}_p \bar{u}_{p,t} + B(\hat{v}_p, \hat{v}_q) \bar{u}_{p,t} \bar{u}_{q,t}]$ for $1 \leq k \leq d$ and $F_t = [\hat{v}_1 \cdot F_t, \cdots, \hat{v}_d \cdot F_t]$; in the second moment equation for $R_t \in \mathbb{R}^{d \times d}$, the operator $L(\bar{u}_t) \in \mathbb{R}^{d \times d}$ indicates meanfluctuation interactions defined in (2.4); and in the stochastic equation for Z_t , $\Gamma : \mathbb{R}^{d \times d} \to \mathbb{R}^d$ is the quadratic coupling operator defined in (2.6). The two higher-moment feedbacks for the mean and covariance, Q_m, Q_v , related to the second and third moments of Z_t respectively, are defined as

$$Q_{m,k} \left(\mathbb{E} \left[Z_t \otimes Z_t \right] \right) = \sum_{p,q=1}^d \gamma_{kpq} \mathbb{E} \left[Z_{p,t} Z_{q,t} \right],$$

$$Q_{v,kl} \left(\mathbb{E} \left[Z_t \otimes Z_t \otimes Z_t \right] \right) = \sum_{p,q=1}^d \gamma_{kpq} \mathbb{E} \left[Z_{p,t} Z_{q,t} Z_{l,t} \right] + \gamma_{lpq} \mathbb{E} \left[Z_{p,t} Z_{q,t} Z_{k,t} \right],$$
(2.8)

for $1 \leq k, l \leq d$ with coupling coefficients $\gamma_{kmn} = \hat{v}_k \cdot B(\hat{v}_m, \hat{v}_n)$. Above, Q_m models the feedback in the mean equation due to the second moments $\mathbb{E}[Z_t \otimes Z_t]$, and $Q_{v,kl}$ is the symmetric feedback in the covariance equation due to all the third moments $\mathbb{E}[Z_t \otimes Z_t \otimes Z_t]$. Notice that Q_m, Q_v can be both viewed as linear operators w.r.t. ρ_t . Besides, in the second moment equation an additional relaxation term with a parameter $\epsilon > 0$ is introduced. This term will not modify the model dynamics with statistical consistency, but is playing a crucial role as a 'reinforcement' term in maintaining stable numerical performance (for example, see [47]) especially with highly instability induced by the strong mean-fluctuation coupling from $L(\bar{u}_t)$. Different from the inherently coupled stochastic and statistical equations (2.5) and (2.3) through the PDF p_t of the original model state u_t , the new closure model (2.7) provides a clean self-consistent formulation for tractable theoretical analysis and direct numerical implementations. A new PDF ρ_t of the stochastic process Z_t is introduced to close the system. The statistical states \bar{u}_t, R_t are first treated as new individual processes subject to higher-order moments w.r.t. ρ_t . Then, the microscopic stochastic equation for Z_t models the high-dimensional multiscale process with explicit dependence on the macroscopic states \bar{u}_t and R_t . Thus, the whole system is closed by the law of the stochastic state Z_t itself. No additional information about the intractable PDF p_t will be needed for solving the FPE of the original system.

In the rest part of this section, we built precise link between the new closure model (2.7) and the coupled equations (2.5) and (2.3) from the original system. First, the following lemma provides the self-consistency in the leading moments of the stochastic modes Z_k and statistical states \bar{u}_t , R_t in (2.7).

Lemma 1. With consistent initial conditions $\mathbb{E}[Z_0] = 0$ and $\mathbb{E}[Z_0Z_0^{\mathsf{T}}] = R_0$, the leading moments of the stochastic modes Z_t of the closure model (2.7) satisfy that for all t > 0

$$\mathbb{E}[Z_t] = 0, \quad \mathbb{E}[Z_t Z_t^{\mathsf{T}}] = R_t, \tag{2.9}$$

where the expectation is taken w.r.t. the PDF ρ_t of Z_t , and R_t is the solution of the second moment equation in (2.7).

The proof can be found through the direct application of Itô's formula and we put the detailed proof in Appendix B. Lemma 1 demonstrates that the law of Z_t indeed recovers the same covariance of the fluctuation modes, while it also contains more information in the higher-order statistics. Furthermore, the closed coupled stochastic-statistical model generates the same statistical solution as the original system (1.1). The following proposition describes the statistical consistency between the coupled model (2.7) and the original system (1.1).

Proposition 2. Assume that the solution u_t of the system (1.1) has a continuous PDF p_t , and the solution $\{\bar{u}_t, R_t; Z_t\}$ of the stochastic-statistical closure model (2.7) has the continuous PDF ρ_t for Z_t together with the deterministic solutions for \bar{u}_t and R_t . Then from the same initial conditions, the two models give the same statistical solution, that is, for all t > 0

$$\mathbb{E}_{p_t}\left[u_t\right] = \bar{u}_t, \quad \mathbb{E}_{p_t}\left[u_t' u_t'^{\mathsf{T}}\right] = R_t, \tag{2.10}$$

where $u'_t = u_t - \mathbb{E}_{p_t}[u_t]$. Furthermore, for both models with any function $\varphi \in C^2(\mathbb{R}^d)$ we have

$$\mathbb{E}_{p_t}\left[\varphi\left(u_t'\right)\right] = \mathbb{E}\left[\varphi\left(\sum_{k=1}^d Z_{k,t}\hat{v}_k\right)\right].$$
(2.11)

The proof of Proposition 2 can be found in Appendix B through detailed computation of each moments of (1.1) compared with that of (2.7). Notice that the left hand sides of (2.10) and (2.11) consist of the statistics requiring solving the PDF p_t of the original system, while the right hand sides are purely from the PDF ρ_t of the closure model. This confirms the direct link between the new stochastic-statistical closure model (2.7) to the statistics in the original fully coupled multiscale system. From this detailed formulation, quantification of non-Gaussian statistics in Z_t relies on the accurate estimation of the leading-order mean and covariance, which can be assisted from the observation data. This leads to effective filtering methods for improved probability forecast for Z_t that will be discussed next in Section 3.

Remark. The second-order closure model (2.7) provides explicit statistical dynamics for the mean and covariance. We can also propose a first-order closure model involving only the statistical mean equation coupled with the McKean-Vlasov SDE

$$dZ_t = L\left(\bar{u}_t\right) Z_t dt + \Gamma\left(Z_t Z_t^{\mathsf{T}} - \mathbb{E}\left[Z_t Z_t^{\mathsf{T}}\right]\right) dt + \Sigma_t dW_t,$$

$$\frac{d\bar{u}_t}{dt} = M\left(\bar{u}_t\right) + \sum_{p,q} \mathbb{E}\left[Z_{p,t} Z_{q,t}\right] B\left(\hat{v}_p, \hat{v}_q\right) + F_t,$$
(2.12)

However, the above equations may introduce an unbalanced approximation to the mean and fluctuation interactions since the dynamics of the SDE for Z_t will directly involve expectation w.r.t. its law ρ_t . Thus, this model will be prone to numerical errors in practical applications. Still, (2.12) can serve as an intermediate model for the analysis of filtering schemes discussed in the next section.

2.3 The stochastic McKean-Vlasov equation as a multiscale interacting system

From the closed stochastic-statistical formulation (2.7), the SDE for Z_t is given by a stochastic McKean-Vlasov equation depending on its own probability distribution ρ_t . In particular, the resulting McKean-Vlasov SDE can be viewed naturally as the mean-field limit of the ensemble approximation of N individual trajectories

$$dZ_t^{(i)} = L\left(\bar{u}_t^N\right) Z_t dt + \Gamma\left(Z_t^{(i)} Z_t^{(i)\mathsf{T}} - R_t^N\right) dt + \Sigma_t dW_t^{(i)}, \quad i = 1, \cdots, N,$$
(2.13)

where $\left\{W_t^{(i)}\right\}_{i=1}^N$ are independent white noise processes, and the initial samples $\left\{Z_0^{(i)}\right\}_{i=1}^N$ are drawn from the same initial measure ρ_0 of Z_t . Notice that the ensemble members $Z_t^{(i)}$ as interacting particles are not evolving independently with each other, but are coupled through feedbacks of the leading-order statistics \bar{u}_t^N and R_t^N according to the statistical equations

$$\frac{\mathrm{d}\bar{u}_{t}^{N}}{\mathrm{d}t} = M\left(\bar{u}_{t}^{N}\right) + Q_{m}\left(\mathbb{E}^{N}\left[\mathbf{Z}_{t}\otimes\mathbf{Z}_{t}\right]\right) + F_{t},$$

$$\frac{\mathrm{d}R_{t}^{N}}{\mathrm{d}t} = L\left(\bar{u}_{t}^{N}\right)R_{t}^{N} + R_{t}^{N}L^{\mathsf{T}}\left(\bar{u}_{t}^{N}\right) + Q_{v}\left(\mathbb{E}^{N}\left[\mathbf{Z}_{t}\otimes\mathbf{Z}_{t}\otimes\mathbf{Z}_{t}\right]\right)$$

$$+ \Sigma_{t}\Sigma_{t}^{\mathsf{T}} + \epsilon^{-1}\left(\mathbb{E}^{N}\left[\mathbf{Z}_{t}\mathbf{Z}_{t}^{\mathsf{T}}\right] - R_{t}^{N}\right).$$
(2.14)

Above, the expectations are computed through the empirical average of the interacting particles $\mathbf{Z}_t = \left\{ Z_t^{(i)} \right\}_{i=1}^N$ from the ensemble simulation

$$\mathbb{E}^{N}\left[\varphi\left(\mathbf{Z}_{t}\right)\right] = \frac{1}{N}\sum_{i=1}^{N}\varphi\left(Z_{t}^{(i)}\right).$$

The coupled ensemble approximation equations (2.13) and (2.14) have advantages in practical applications. Unlike the general McKean-Vlasov SDEs [41], (2.13) avoids the direct inclusion of the PDF of Z_t , which is very difficult to approximate accurately from finite particles. Instead, the PDF only implicitly enters the mean and covariance equations (2.14) through the computation of higher moments. Effective computational algorithms with consistent statistics then can be proposed (such as using the efficient random batch methods [47, 48]) for the straightforward ensemble model approximation. Besides in practical computation, the relaxation term in R_t^N provides additional restoring forcing as a correction term to numerical errors with finite sample approximation to reinforce stable dynamics and consistent statistics especially in the case where internal instability is involved.

In particular, it is well-know [25] that the empirical measure converges weakly to the true distribution, $\rho_t^N \to \rho_t$, as well as the leading-order statistics in (2.13), $\bar{u}^N \to \bar{u}$, $R_t^N \to R_t$, as $N \to \infty$ under relatively weak assumptions (see Proposition 12 in Section 5). The dynamical equation for the continuous density function ρ_t of Z_t is given by the corresponding equation

$$\frac{\partial \rho_t}{\partial t} = \mathcal{L}_t^* \left(\bar{u}_t, R_t \right) \rho_t \coloneqq -\nabla_z \cdot \left[L \left(\bar{u}_t \right) z \rho_t \left(z \right) + \Gamma \left(z z^{\mathsf{T}} - R_t \right) \rho_t \left(z \right) \right] + \frac{1}{2} \nabla_z \cdot \left[\nabla_z \cdot \left(\Sigma_t \Sigma_t^{\mathsf{T}} \rho_t \left(z \right) \right) \right], \tag{2.15}$$

where \mathcal{L}_t^* is the adjoint of the generator \mathcal{L}_t that is also dependent on the law of Z_t shown in the statistics of the mean \bar{u}_t and covariance R_t . We postpone the detailed analysis for the convergence of the ensemble approximation model in Section 5 together with the filter approximation.

In general, the probability density ρ_t will demonstrate non-Gaussian features due to the nonlinear stochastic coupling effects. On the other hand, though the equations for the stochastic state Z_t contain nonlinear structures, its PDF ρ_t is subject to only linear dynamics (2.15) with $\mathcal{L}_t^*(\bar{u}_t, R_t)$, which is dependent on the mean and covariance. These desirable features inspires the construction of effective filtering methods in the next section to include leadingorder statistical observations to improve forecast of highly non-Gaussian statistics.

3 Filtering models using observations in mean and covariance

In this section, we exploit the ideas in filtering to propose improved approximation of probability distributions of the model state containing highly non-Gaussian statistics. The optimal filter is developed by combining the stochastic forecast model describing unobserved microscopic states and leading-order statistics introduced as macroscopic observations. We start with a precise description of the optimal filter equations satisfying a conditional Gaussian process, then a new statistical filtering model are proposed approximating the optimal filtering solution with equivalent statistics.

3.1 Filtering probability distributions using statistical observations

We first formulate the filtering problem for predicting probability distributions based on observations from the leading-order statistics. From the stochastic-statistical equations (2.7), we can reformulate the general multiscale system (1.1) for u_t as a composition of the macroscopic state from the first two moments \bar{u}_t , R_t and the microscopic stochastic processes Z_t . In practice, measurements are often available in the macroscopic statistical states (such as the locally averaged mean state and variances as the deviation from the mean). Therefore, it is natural to incorporate the statistical observation data to improve the estimation of the unobserved microscopic processes, especially to recover the unobserved higher-order statistics (such as the deviation from the normal distribution indicating the occurrence of high impact extreme events).

Using similar idea of the Fokker-Planck filter introduced in [1], we assume that the signal process of the filtering problem is from the model state PDF $\rho_t \in \mathcal{P}(\mathbb{R}^d)$ belonging to the space of continuous probability density functions on \mathbb{R}^d with all finite moments. The observation process is generated by the finite-dimensional statistical moments denoted as $y_t \in \mathbb{R}^p$. This leads to the general *infinite-dimensional filtering system with statistical observations*

$$\mathrm{d}\rho_t = \mathcal{L}_t^*\left(y_t\right)\rho_t \mathrm{d}t, \qquad \qquad \rho_{t=0} \sim \mu_0, \qquad (3.1a)$$

$$dy_t = [\mathcal{H}\rho_t + h_t (y_t)] dt + \Gamma_t dB_t, \quad y_{t=0} = y_0,$$
(3.1b)

where μ_0 is a probability measure of the $\mathcal{P}(\mathbb{R}^d)$ -valued random field ρ . Given y_t , we will have $\rho_t \in \mathcal{P}(\mathbb{R}^d)$ for all t > 0. Above, $\mathcal{L}_t(y_t)$ is the infinitesimal generator of the corresponding SDE for Z_t with the explicit form given by (2.15); the general observation process y_t satisfies the dynamical equation subject to a linear observation operator $\mathcal{H}: \mathcal{P}(\mathbb{R}^d) \to \mathbb{R}^p$ acting on the PDF ρ_t , as well as $h_t: \mathbb{R}^p \to \mathbb{R}^p$. Detailed equations for $y_t = (\bar{u}_t, R_t)$ will be given next in (3.3) based on the first two moments. Notice here ρ_t becomes a random field (with a precise definition given in (3.5)) due to the randomness in y_t as well as its initial uncertainty.

Statistical observations from leading-order moments containing errors

Let p_t be the (unknown) PDF of the state u_t in (1.1), that is, the deterministic solution of the FPE (1.2). Then, we can assume that observations are drawn from the mean or covariance of the state u_t as

$$\bar{u}_{k,t} = \mathbb{E}_{p_t} \left[u_t \cdot \hat{v}_k \right], \quad R_{kl,t} = \mathbb{E}_{p_t} \left[\left(\hat{v}_k \cdot u_t' \right) \left(u_t' \cdot \hat{v}_l \right) \right], \tag{3.2}$$

projected to the observed large-scale modes $\hat{v}_k, k \leq d'$ in (2.1). We refer it as the *full observation case* with d' = dand *partial observation case* with d' < d. For simplicity, we may always consider the full observation case d' = d(that is, $y_t = (\bar{u}_t, R_t) \in \mathbb{R}^p$ with $p = d + d^2$) in this paper without confusion. The dynamical equations for \bar{u}_t, R_t can be introduced according to the closure model (2.7). According to Proposition 2, the statistical equations for \bar{u}_t, R_t in (2.7) provide statistical solutions consistent with the law p_t of the state u_t .

On the other hand, model errors are usually introduced due to the finite truncation of the originally infinitedimensional continuous system as well as measurement errors. Therefore, additional correction terms, using independent white noises, $B_{m,t}$ and $B_{v,t}$, are added to the equations (2.3) for observed statistics accounting for errors from the imperfect model approximations. The detailed equations for the observations (3.1b) can be rewritten according to (2.3) as the following SDEs for $\bar{u}_t \in \mathbb{R}^d, R_t \in \mathbb{R}^{d^2}$

$$d\bar{u}_t = \left[\mathcal{H}_m \rho_t + h_{m,t} \left(\bar{u}_t\right)\right] dt + \Gamma_m dB_{m,t},$$

$$dR_t = \left[\mathcal{H}_v \rho_t + h_{v,t} \left(\bar{u}_t, R_t\right)\right] dt + \Gamma_v dB_{v,t}.$$
(3.3)

where $h_{m,t}(\bar{u}) = M(\bar{u}) + F_t$ and $h_{v,t}(\bar{u}, R) = L(\bar{u})R + RL(\bar{u})^{\mathsf{T}} + \Sigma_t \Sigma_t^{\mathsf{T}}$ are deterministic functions, while the linear

observation operators, $\mathcal{H}_m, \mathcal{H}_v$, are defined by the high-order statistical feedback functions (2.8)

$$\mathcal{H}_{m,k}\rho = \int_{\mathbb{R}^d} H_k^m(z)\,\rho(z)\,\mathrm{d}z = \sum_{p,q=1}^d \gamma_{kpq} \int_{\mathbb{R}^d} z_p z_q \rho(z)\,\mathrm{d}z,$$

$$\mathcal{H}_{v,kl}\rho = \int_{\mathbb{R}^d} H_{kl}^v(z)\,\rho(z)\,\mathrm{d}z = \sum_{p,q=1}^d \left[\gamma_{kpq} \int_{\mathbb{R}^d} z_p z_q z_l \rho(z)\,\mathrm{d}z + \gamma_{lpq} \int_{\mathbb{R}^d} z_p z_q z_k \rho(z)\,\mathrm{d}z\right].$$
(3.4)

We assume $\rho \in \mathcal{P}(\mathbb{R}^d)$ the probability density with finite moments in all orders, thus we have $\mathcal{H}_{m,k}\rho < \infty$ and $\mathcal{H}_{v,kl}\rho < \infty$ for all k, l. Then (3.3) fits into the general observation equation (3.1b) by setting $y_t = (\bar{u}_t, R_t)^{\mathsf{T}} \in \mathbb{R}^p$ with $p = d + d^2$ in a column vector, and letting $\mathcal{H} = (\mathcal{H}_m, \mathcal{H}_v)^{\mathsf{T}}$, $h_t = (h_{m,t}, h_{v,t})^{\mathsf{T}}$, and $\Gamma_t = \text{diag}(\Gamma_m, \Gamma_v)$. With this explicit setup of the filtering problem, we will consider the optimal filtering solution for the probability density ρ_t of Z_t based on the statistical observation data $Y_t = \{(\bar{u}_s, R_s), s \leq t\}$.

3.2 The optimal filter with conditional Gaussian structure

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be the complete probability space, and denote $\mathcal{P}(\mathbb{R}^d)$ as the space of probability density functions with bounded all moments. We first define the $\mathcal{P}(\mathbb{R}^d)$ -valued stochastic process ρ_t as

$$\rho_t : \mathbb{R}^d \times \Omega \to \mathbb{R}^+, \ (z, \omega) \mapsto \rho_t \left(z; \omega \right), \ \text{with} \ \rho_t \left(\cdot; \omega \right) \in \mathcal{P} \left(\mathbb{R}^d \right),$$
(3.5)

which is thereafter referred to as a random field. In contrast to the standard filtering problem concerning the nonlinear SDE of the random model states Z_t , for derivation purpose of the exact optimal equations, we lift the problem into filtering the random field ρ_t based on the observation information in $y_s, s \leq t$ as in (3.1). A stochastic model on \mathbb{R}^d (3.16) will be then proposed for practical implementations next in Section 2. Let $\mathcal{G}_t = \sigma \{\omega : y_s, s \leq t\}$ be the σ -algebra generated by the observations up to time t. We define the space as the collection of \mathcal{G}_t -measurable square-integrable random fields

$$\mathcal{V}_{t} \coloneqq L^{2}\left(\Omega, \mathcal{G}_{t}, \mathbb{P}; \mathcal{P}\left(\mathbb{R}^{d}\right) \cap L^{2}\left(\mathbb{R}^{d}\right)\right), \tag{3.6}$$

satisfying $\int \|\nu(\cdot;\omega)\|_{L^2(\mathbb{R}^d)}^2 d\mathbb{P}(\omega) < \infty$ and $\nu(\cdot;\omega) \in \mathcal{P}(\mathbb{R}^d) \cap L^2(\mathbb{R}^d)$ for $\nu \in \mathcal{V}_t$. In this infinite-dimensional filtering problem, we aim to find the optimal approximation of ρ_t in the space \mathcal{V}_t . The optimal filtering solution $\hat{\rho}_t$ is then introduced as the least-square estimate with the minimum variance as

$$\hat{\rho}_t \coloneqq \operatorname*{arg\,min}_{\nu \in \mathcal{V}_t} \mathbb{E}\left[\left\| \rho_t - \nu \right\|_{L^2(\mathbb{R}^d)}^2 \right] = \mathsf{P}_{\mathcal{V}_t}\left[\rho_t \right],\tag{3.7}$$

where the optimal solution $\hat{\rho}_t$ can be viewed as the unbiased projection of ρ_t onto the space \mathcal{V}_t . (3.7) indicates that $\hat{\rho}_t$ gives the PDF estimation closest to the true distribution ρ_t in the mean square sense in agreement with the observations in \mathcal{G}_t .

Accordingly, we define the optimal filter distribution $\mu_t : \mathcal{P}(\mathbb{R}^d) \times \Omega \to [0, 1]$ as the regular conditional measure of the stochastic process ρ_t given \mathcal{G}_t . That is, for any Borel set $A \in \mathcal{B}(\mathcal{P}(\mathbb{R}^d))$, μ_t gives the conditional probability of ρ_t given \mathcal{G}_t

$$\mu_t(A;\omega) \coloneqq \mathbb{P}\left(\rho_t \in A \mid \mathcal{G}_t\right)(\omega), \quad \text{a.s. } \omega \in \Omega.$$
(3.8)

Notice that $\mu_t(A; \cdot) \in \mathcal{G}_t$ is still a stochastic process. For any functional $F \in C(\mathcal{P}(\mathbb{R}^d))$ and t > 0, we can introduce the conditional expectation w.r.t. the measure μ_t given \mathcal{G}_t as

$$\mathbb{E}\left[F\left(\rho_{t}\right) \mid \mathcal{G}_{t}\right] \coloneqq \int_{\mathcal{P}(\mathbb{R}^{d})} F\left(\rho\right) \mu_{t}\left(\mathrm{d}\rho\right)$$

Therefore, the optimal filter solution (3.7) is a random field

$$\hat{\rho}_t = \mathbb{E}\left[\rho_t \mid \mathcal{G}_t\right] \tag{3.9}$$

given by the conditional expectation of ρ_t w.r.t. μ_t . For any linear operator $\mathcal{M} : L^2(\mathbb{R}^d) \to \mathbb{R}^p$, we have $\mathcal{M}\hat{\rho}_t = \mathbb{E}[\mathcal{M}\rho_t \mid \mathcal{G}_t]$. Furthermore, second moment of $\mathcal{M}\rho_t$ is given by

$$\mathbb{E}\left[\left(\mathcal{M}\rho_t - \mathcal{M}\hat{\rho}_t\right)\left(\mathcal{M}\rho_t - \mathcal{M}\hat{\rho}_t\right)^{\mathsf{T}} \mid \mathcal{G}_t\right] = \mathcal{M}\mathcal{C}_t\mathcal{M}^*,$$

where $\hat{\mathcal{C}}_t(\omega): L^2(\mathbb{R}^d; \mathbb{R}^p) \to L^2(\mathbb{R}^d; \mathbb{R}^p)$ with $\hat{\mathcal{C}}_t^* = \hat{\mathcal{C}}_t$ is the self-adjoint covariance operator for any $f \in L^2(\mathbb{R}^d; \mathbb{R}^p)$

$$\hat{\mathcal{C}}_t f = \mathbb{E}\left[\left(\rho - \hat{\rho}_t \right) \int_{\mathbb{R}^d} \left(\rho - \hat{\rho}_t \right) \left(z \right) f\left(z \right) \mathrm{d}z \mid \mathcal{G}_t \right].$$
(3.10)

Notice again $\hat{\mathcal{C}}_t f(z; \cdot)$ is also a random field conditional on \mathcal{G}_t . For clarification of notations, we will call $\hat{\rho}_t$ and $\hat{\mathcal{C}}_t$ the optimal filter solution for the mean and covariance, and μ_t the optimal filter distribution in the rest part of the paper.

In particular, we can characterize the optimal filter solution $\hat{\rho}_t$ as the best estimate in each order of moments. The following result describes the accuracy of the filter approximations under any finite-dimensional projections.

Proposition 3. Let ρ_t be the random field from the system (3.1) and $\hat{\rho}_t = \mathbb{E}\left[\rho_t \mid \mathcal{G}_t\right]$ the optimal filter solution given the observations in \mathcal{G}_t . For any linear operator $\mathcal{M} : \mathcal{P}\left(\mathbb{R}^d\right) \to \mathbb{R}^p$ defined by $\mathcal{M}\rho = \int \mathcal{M}(z) \rho(dz)$ with $\mathcal{M} \in C\left(\mathbb{R}^d; \mathbb{R}^p\right), \ \mathcal{M}\hat{\rho}_t = \mathbb{E}\left[\mathcal{M}\rho_t \mid \mathcal{G}_t\right]$ gives the best unbiased estimate of $\mathcal{M}\rho_t$ in the sense of minimum mean square error, that is,

$$\mathbb{E}\left[\left|\mathcal{M}\rho_{t}-\mathcal{M}\hat{\rho}_{t}\right|^{2}\right] = \min_{\nu\in\mathcal{V}_{t}}\mathbb{E}\left[\left|\mathcal{M}\rho_{t}-\mathcal{M}\nu\right|^{2}\right], \quad \text{with} \quad \mathbb{E}\left[\mathcal{M}\hat{\rho}_{t}\right] = \mathbb{E}\left[\mathcal{M}\rho_{t}\right].$$
(3.11)

Proof. For any \mathcal{G}_t -measurable square-integrable stochastic process ν , we have from direct computation

$$\mathbb{E}\left[\left|\mathcal{M}\rho_{t}-\mathcal{M}\nu\right|^{2}\right]-\mathbb{E}\left[\left|\mathcal{M}\rho_{t}-\mathcal{M}\hat{\rho}_{t}\right|^{2}\right]$$
$$=\mathbb{E}\left[\left(\mathcal{M}\hat{\rho}_{t}-\mathcal{M}\nu\right)\cdot\left(2\mathcal{M}\rho_{t}-\mathcal{M}\nu-\mathcal{M}\hat{\rho}_{t}\right)\right]$$
$$=\mathbb{E}\left\{\mathbb{E}\left[\left(\mathcal{M}\hat{\rho}_{t}-\mathcal{M}\nu\right)\cdot\left(2\mathcal{M}\rho_{t}-\mathcal{M}\nu-\mathcal{M}\hat{\rho}_{t}\right)\mid\mathcal{G}_{t}\right]\right\}$$
$$=\mathbb{E}\left\{\left(\mathcal{M}\hat{\rho}_{t}-\mathcal{M}\nu\right)\cdot\left[\mathbb{E}\left[2\mathcal{M}\rho_{t}\mid\mathcal{G}_{t}\right]-\left(\mathcal{M}\nu+\mathcal{M}\hat{\rho}_{t}\right)\right]\right\}$$
$$=\mathbb{E}\left[\left|\mathcal{M}\hat{\rho}_{t}-\mathcal{M}\nu\right|^{2}\right]\geq0.$$

Above, the third equality uses the fact $\mathcal{M}\hat{\rho}_t - \mathcal{M}\nu = \mathbb{E}[\mathcal{M}\rho_t | \mathcal{G}_t] - \mathcal{M}\nu$ is \mathcal{G}_t -measurable. Thus, we get $\mathcal{M}\hat{\rho}_t$ minimizes the mean square error. The consistency under the expectation in $\mathcal{M}\hat{\rho}_t$ and $\mathcal{M}\rho_t$ can be directly implied by definition.

By taking the operator \mathcal{M} as the expectation on $M(Z_t) = Z_t^m$ with any integer m, $\mathcal{M}\rho_t$ and $\mathcal{M}\hat{\rho}_t$ give the *m*-th order moments of Z_t under the random field ρ_t in (3.5) and optimal filter approximation $\hat{\rho}_t$. A direct implication from (3.11) shows that we have the unbiased statistics in all high-order moments $\mathbb{E}[\mathcal{M}\hat{\rho}_t] = \mathbb{E}[\mathcal{M}\rho_t]$ with the minimum error $\mathbb{E}[|\mathcal{M}\rho_t - \mathcal{M}\hat{\rho}_t|^2]$ from the optimal filter solution.

Importantly, the model equations (3.1) satisfy the desirable conditional Gaussian process [36], that is, given the observations of $Y_t = \{y_s = (\bar{u}_s, R_s), s \leq t\}$ and Gaussian initial state ρ_0 , the random field ρ_t follows a Gaussian distribution at each time t. Let $\rho_t = \rho_t(\cdot; \omega)$ be the (unknown) signal state satisfying linear dynamics (3.1a), and $y_t = y_t(\omega)$ the observed statistical process subject to linear observation operators (3.1b). The optimal filter distribution μ_t (3.8) conditional on Y_t then becomes an infinite-dimensional Gaussian distribution, $\mu_t(\cdot, \omega) = \mathcal{N}\left(\hat{\rho}_t, \hat{\mathcal{C}}_t\right)(\omega)$, where the mean $\hat{\rho}_t(\cdot; \omega)$ and covariance $\hat{\mathcal{C}}_t(\omega)$ give the solution to (3.9) and (3.10) respectively. Therefore, the equations for the mean and covariance are given by the generalized version of Kalman-Bucy (KB) filter for the infinite-dimensional conditional Gaussian processes

$$d\hat{\rho}_{t} = \mathcal{L}_{t}^{*} \left(\bar{u}_{t}, R_{t} \right) \hat{\rho}_{t} dt + \hat{\mathcal{L}}_{t} \mathcal{H}_{m}^{*} \Gamma_{m}^{-2} \left\{ d\bar{u}_{t} - \left[\mathcal{H}_{m} \hat{\rho}_{t} + h_{m,t} \left(\bar{u}_{t} \right) \right] dt \right\} + \hat{\mathcal{L}}_{t} \mathcal{H}_{v}^{*} \Gamma_{v}^{-2} \left\{ dR_{t} - \left[\mathcal{H}_{v} \hat{\rho}_{t} + h_{v,t} \left(\bar{u}, R \right) \right] dt \right\},$$

$$d\hat{\mathcal{L}}_{t} = \left[\mathcal{L}_{t}^{*} \left(\bar{u}_{t}, R_{t} \right) \hat{\mathcal{L}}_{t} + \hat{\mathcal{L}}_{t} \mathcal{L}_{t} \left(\bar{u}_{t}, R_{t} \right) \right] dt - \hat{\mathcal{L}}_{t} \left(\mathcal{H}_{m}^{*} \Gamma_{m}^{-2} \mathcal{H}_{m} + \mathcal{H}_{v}^{*} \Gamma_{v}^{-2} \mathcal{H}_{v} \right) \hat{\mathcal{L}}_{t} dt.$$

$$(3.12)$$

The equations of the conditional Gaussian processes and the uniqueness of the solutions are developed in Chapter 12 of [36] for finite dimensional systems. The results are then generalized to infinite-dimensional Hilbert space [10, 20] (see a summary of the main results in Appendix A). The system (3.12) gives a closed set of coupled SPDEs (due to the randomness in \bar{u}_t, R_t) for the optimal filter solution $\hat{\rho}_t$ enabling more detailed analysis and development of practical methods for computing the optimal solution.

Remark. The filtering problem using statistical observations is originally introduced as the ensemble Fokker-Planck filter in [1]. We propose the filtering equations (3.12) as a further generalization where nonlinearly coupled conditional processes are involved. In particular, this new filtering model directly fits into the coupled stochastic-statistical modeling framework (2.7).

3.3 A surrogate filtering model for approximating the optimal filter solution

The resulting optimal filtering problem from (3.1) requires to solve the infinite-dimensional system (3.12) concerning the function $\hat{\rho}_t \in \mathcal{P}(\mathbb{R}^d)$ and operator $\hat{\mathcal{C}}_t$ on $\mathcal{P}(\mathbb{R}^d)$. It usually becomes intractable in finding such infinitedimensional solutions from direct methods. In developing practical strategies to realize the optimal filter solution, it is more useful to find a surrogate model for the stochastic process \tilde{Z}_t , based on which effective ensemble-based approaches can be built. Therefore, we aim to construct an approximating filtering model from designing a new dynamical equation for \tilde{Z}_t , whose PDF $\tilde{\rho}_t$ can effectively represent that of the optimal filter solution $\hat{\rho}_t$.

3.3.1 Filtering updating cycle in a split two-step procedure

For a clear characterization of the filtering process, we follow the general procedure in [7] to first describe the filtering process by concatenated iterations of transporting maps on the corresponding probability distribution. We propose a new stochastic process \tilde{Z}_t , whose law $\tilde{\rho}_t \in \mathcal{V}_t$ is still a $\mathcal{P}(\mathbb{R}^d)$ -valued random field in \mathcal{G}_t dependent on the same statistical observation y_t as in the optimal filter (3.12). Thus, the filtering updating cycle during the time interval $[t, t + \tau]$ can be characterized by the transport of the probability density $\tilde{\rho}_t$ of \tilde{Z}_t .

The discrete time update of the approximation filter PDF $\tilde{\rho}_t$ is carried out in a split two-step procedure. First, the *forecast step* can be viewed as the push-forward operator acting on the probability density function at time instant t with time step $\tau > 0$

$$\tilde{\rho}_t \to \tilde{\rho}_{t+\tau}^- \coloneqq \mathcal{F}_t^\tau \tilde{\rho}_t = e^{\int_t^{t+\tau} \mathcal{L}_s^*(y_s) \mathrm{d}s} \tilde{\rho}_t, \tag{3.13}$$

where \mathcal{F}_t^{τ} represents the forecast updating operator with forward time step τ , and $\mathcal{L}_t(y_t)$ is the same generator as in (2.15). Second, the *analysis step* updates the prior distribution $\tilde{\rho}^-$ to the posterior distribution $\tilde{\rho}^+$ by incorporating the observation data up to $Y_{t+\tau} = \{y_s, s \leq t + \tau\}$, that is

$$\tilde{\rho}_{t+\tau}^{-} \to \tilde{\rho}_{t+\tau}^{+} \coloneqq \mathcal{A}_{t}^{\tau} \left(\tilde{\rho}_{t+\tau}^{-}; Y_{t+\tau} \right), \tag{3.14}$$

where \mathcal{A}_t^{τ} represents the analysis updating operator. Therefore, the full filtering cycle from t to $t + \tau$ can be summarized as the composition of the forecast and analysis maps

$$\tilde{\rho}_{t+\tau} = \mathcal{A}_t^{\tau} \left(\mathcal{F}_t^{\tau} \tilde{\rho}_t; Y_{t+\tau} \right). \tag{3.15}$$

Notice that \mathcal{F}_t^{τ} is a linear operator on $\tilde{\rho}_t$, while \mathcal{A}_t^{τ} could contain nonlinear actions due to the normalization of the probability distribution. Continuous equation for $\partial_t \tilde{\rho}_t = \lim_{\tau \to 0} \frac{1}{\tau} (\tilde{\rho}_{t+\tau} - \tilde{\rho}_t)$ is then achieved by letting the discrete time step $\tau \to 0$. Next, we first propose the general structure of the new filtering model for $\tilde{Z}_t \sim \tilde{\rho}_t$ as a combination of the above two-step procedure, then detailed analysis can be done according to the design of the forecast and analysis step operators \mathcal{F}_t^{τ} and \mathcal{A}_t^{τ} accordingly.

3.3.2 Construction of equivalent statistical approximating filter

For simplicity of notations, we still use the general statistical observation processes (3.3) for $y_t = (\bar{u}_t, R_t)$ taking the compact form

$$\mathrm{d}y_t = \left[\mathcal{H}\rho_t + h_t\left(y_t\right)\right]\mathrm{d}t + \Gamma_t\mathrm{d}B_t,$$

where the general observation operator (3.4), $\mathcal{H}\rho_t = \int H(z) \rho_t(z) dz$, is defined with the general observation function $H \in C(\mathbb{R}^d; \mathbb{R}^p)$ acting on the density function ρ_t . Following the general construction in [8, 45], we seek the approximating filter model adopting the following McKean-Vlasov representation with undetermined functionals a_t, K_t

$$d\tilde{Z}_{t} = L\left(\bar{u}_{t}\right)\tilde{Z}_{t}dt + \Gamma\left(\tilde{Z}_{t}\tilde{Z}_{t}^{\mathsf{T}} - R_{t}\right)dt + \Sigma_{t}d\tilde{W}_{t} + a_{t}\left(\tilde{Z}_{t};\tilde{\rho}_{t}\right)dt + K_{t}\left(\tilde{Z}_{t};\tilde{\rho}_{t}\right)\left\{dy_{t} - \left[H\left(\tilde{Z}_{t}\right) + h_{t}\left(y_{t}\right)\right]dt - \Gamma_{t}d\tilde{B}_{t}\right\}.$$
(3.16)

We use the name 'statistical filtering' to refer the above new model emphasizing our main goal of filtering statistical moments different from the common filtering case. The first row of the above equation models the forecast step of the filtering process, while the second row is the analysis step. The forecast step accepts the same dynamical model of (3.1a) dependent on the mean and covariance (\bar{u}_t, R_t) . On the other hand, the analysis step serves as an additional control correction over statistical observations y_t . New functionals known as the drift $a_t : \mathbb{R}^d \times \mathcal{P}(\mathbb{R}^d) \to \mathbb{R}^d$ and the control gain operator $K_t : \mathbb{R}^d \times \mathcal{P}(\mathbb{R}^d) \to \mathbb{R}^{d \times p}$ are introduced, resulting in an approximating filtering model about the process \tilde{Z}_t . Most importantly, as will be shown next in Theorem 7 under proper condition, a_t and K_t are only implicitly dependent on $\tilde{\rho}_t$ through the leading moments, without the need to compute the (potentially highly non-Gaussian) density function $\tilde{\rho}_t$ explicitly.

In a more clear identification of the filtering updates involving several levels of approximations, we take the split-step strategy to analyze the coupled forecast step and analysis step of the filtering equation (3.16) separately. In particular, the forecast step in the first row of (3.16) is given by the exactly same form as the stochastic-statistical closure equations (2.7) developed in Section 2. Thus in practice, the updating step with the forecast operator can be implemented adopting the efficient uncertainty prediction methods such as [38, 47]. Then, the remaining task is to propose proper analysis step update in the second line of (3.16) concerning consistent statistics with the optimal solution $\hat{\rho}_t$ in (3.12).

3.4 Statistical consistency in analysis step update of the approximating filter

Now, we focus on updating posterior PDF $\tilde{\rho}_t$ in (3.14) of the proposed approximating filter (3.16) based on the statistical observation y_t satisfying (3.3). Concentrating on the analysis step, the resulting optimal filter equations (3.12) for the mean and covariance $(\hat{\rho}_t, \hat{C}_t)$ become

$$d\hat{\rho}_{t} = \hat{\mathcal{C}}_{t} \mathcal{H}^{*} \Gamma_{t}^{-2} \left\{ dy_{t} - \left[\mathcal{H} \hat{\rho}_{t} + h_{t} \left(y_{t} \right) \right] dt \right\},$$

$$d\hat{\mathcal{C}}_{t} = -\hat{\mathcal{C}}_{t} \mathcal{H}^{*} \Gamma_{t}^{-2} \mathcal{H} \hat{\mathcal{C}}_{t} dt.$$
(3.17)

Correspondingly, the approximating statistical filtering model for \tilde{Z}_t satisfies the second line of the SDE (3.16) as

$$d\tilde{Z}_{t} = a_{t}\left(\tilde{Z}_{t}\right)dt + K_{t}\left(\tilde{Z}_{t}\right)\left\{dy_{t} - \left[H\left(\tilde{Z}_{t}\right) + h_{t}\left(y_{t}\right)\right]dt - \Gamma_{t}d\tilde{B}_{t}\right\}.$$
(3.18)

Following the similar idea in the McKean-Vlasov representation of the filtering equation [45, 57], we expect the PDF $\tilde{\rho}_t$ of \tilde{Z}_t to satisfy the following Kushner-Stratonovich-type equation (with requirements on a_t, K_t given next in (3.21))

$$\frac{\partial \tilde{\rho}_t}{\partial t} = \left[H\left(z\right) - \mathcal{H}\tilde{\rho}_t\right]^{\mathsf{T}} \Gamma_t^{-2} \left[\frac{\mathrm{d}y_t}{\mathrm{d}t} - \mathcal{H}\tilde{\rho}_t - h_t\left(y_t\right)\right] \tilde{\rho}_t.$$
(3.19)

Again, the goal here is to approximate the optimal filter mean $\hat{\rho}_t$ in (3.17) by $\tilde{\rho}_t$ generated by the surrogate SDE model (3.18) in the sense of consistent statistics.

Unfortunately, the approximation (3.19) and the optimal filtering equation (3.17) will in general have different continuous solutions for $\tilde{\rho}_t$ and $\hat{\rho}_t$ due to their distinctive dynamics. In order to compare the statistical moments of the two distributions, we apply the linear operator \mathcal{H} to the optimal equations (3.17) as a finite-dimensional projection on leading moments. The resulting equations optimal mean and covariance equations become finite dimensional as

$$d\left(\mathcal{H}\hat{\rho}_{t}\right) = \left(\mathcal{H}\hat{\mathcal{C}}_{t}\mathcal{H}^{*}\right)\Gamma_{t}^{-2}\left\{dy_{t} - \left[\mathcal{H}\hat{\rho}_{t} + h_{t}\left(y_{t}\right)\right]dt\right\},\$$

$$d\left(\mathcal{H}\hat{\mathcal{C}}_{t}\mathcal{H}^{*}\right) = -\left(\mathcal{H}\hat{\mathcal{C}}_{t}\mathcal{H}^{*}\right)\Gamma_{t}^{-2}\left(\mathcal{H}\hat{\mathcal{C}}_{t}\mathcal{H}^{*}\right)dt.$$
(3.20)

Above, remind that the observation operator $\mathcal{H} : L^2(\mathbb{R}^d) \to \mathbb{R}^p$ and its adjoint $\mathcal{H}^* : \mathbb{R}^p \to L^2(\mathbb{R}^d)$ are defined based on the observation function $H \in C_b^2(\mathbb{R}^d; \mathbb{R}^p)$ as

$$\mathcal{H}\rho = \int H(z) \rho(z) dz, \quad [\mathcal{H}^*u](z) = u \cdot H(z),$$

and the covariance operator $\hat{\mathcal{C}}_t : L^2(\mathbb{R}^d) \to L^2(\mathbb{R}^d)$ is defined in (3.10). Therefore, (3.20) gives the equations for the finite-dimensional quantities $\mathcal{H}\hat{\rho}_t \in \mathbb{R}^p$ and $\mathcal{H}\hat{\mathcal{C}}_t\mathcal{H}^* \in \mathbb{R}^{p \times p}$ as the first two moments of H w.r.t. $\hat{\rho}_t$. The idea is to design the analysis step operator \mathcal{A}_t^{τ} for the approximating filter process \tilde{Z}_t in (3.18), so that consistency in the first and second-order moments can be achieved.

Denote the expectation, $\tilde{\mathbb{E}}\left[\cdot\right] \coloneqq \mathbb{E}\left[\cdot \mid \mathcal{G}_t\right] = \mathbb{E}_{\tilde{\rho}_t}\left[\cdot\right]$, w.r.t. the conditional density $\tilde{\rho}_t$ in (3.19) given the same observation process in \mathcal{G}_t . We define $\bar{H}_t = \tilde{\mathbb{E}}H\left(\tilde{Z}_t\right) = \int H\left(z\right)\tilde{\rho}_t\left(z\right) dz$ and $C_t^H = \tilde{\mathbb{E}}\left[H_t\left(\tilde{Z}_t\right) - \bar{H}_t\right]\left[H_t\left(\tilde{Z}_t\right) - \bar{H}_t\right]^{\mathsf{T}}$ as the first and second-order moments of H w.r.t. $\tilde{\rho}_t$. Assume that the drift a_t and gain K_t in the SDE approximation (3.18) satisfy the following identities

$$a_t = \nabla \cdot \left(K_t \Gamma_t^2 K_t^{\mathsf{T}} \right) - K_t \Gamma_t^2 \nabla \cdot K_t^{\mathsf{T}}, \quad -\nabla \cdot \left(K_t^{\mathsf{T}} \tilde{\rho}_t \right) = \tilde{\rho}_t \Gamma_t^{-2} \left(H\left(z\right) - \tilde{\mathbb{E}}H \right), \tag{3.21}$$

where the divergence on a matrix is defined columnwise as $(\nabla \cdot A)_i = \sum_j \partial_{z_j} A_{ij}$. We first have the following result concerning the evolution equations of \bar{H}_t and C_t^H given the realization $Y_t = \{y_s, s \leq t\}$.

Lemma 4. Given that $\Gamma_t \succ 0$ in (3.1) and the identities (3.21) are satisfied, the evolution equations for the mean and covariance of the observation function $H\left(\tilde{Z}_t\right)$ associated with the SDE (3.18) are given by

$$d\bar{H}_{t} = C_{t}^{H}\Gamma_{t}^{-2} \left\{ dy_{t} - \left[\bar{H}_{t} + h_{t}\left(y_{t}\right)\right] dt \right\},$$

$$dC_{t}^{H} = Q_{t}^{H}\Gamma_{t}^{-2} \left\{ dy_{t} - \left[\bar{H}_{t} + h_{t}\left(y_{t}\right)\right] dt \right\} - C_{t}^{H}\Gamma_{t}^{-2}C_{t}^{H}dt,$$
(3.22)

where $Q_t^H : \mathbb{R}^p \to \mathbb{R}^{p \times p}$ is defined as

$$Q_t^H = \tilde{\mathbb{E}}\left[\left(H_t' H_t'^{\mathsf{T}} \right) \otimes H_t'^{\mathsf{T}} \right]$$

containing third moments of $H'_t = H\left(\tilde{Z}_t\right) - \bar{H}_t$.

We put the detailed derivation of (3.22) in Appendix B. Notice that (3.22) goes back to the Kalman-Bucy filter if we set linear observation $H\left(\tilde{Z}_t\right) = \tilde{Z}_t$ satisfying a normal distribution as in [1, 9]. Although here we are considering the more general nonlinear dynamics and observation functions from (3.4).

Comparing (3.22) and (3.20) implies that the same statistical solution can be reached in (\bar{H}_t, C_t^H) and $(\mathcal{H}\hat{\rho}_t, \mathcal{H}\hat{\mathcal{C}}_t\mathcal{H}^*)$ if we have $Q_t^H = 0$. In order to achieve this, we further introduce the projection operator on the space of probability distributions using the Kullback-Leibler (KL) divergence [30] as an unbiased metric.

Definition 5. Define the operator \mathcal{S}_H making symmetric projection on the probability density $\rho \in \mathcal{P}(\mathbb{R}^d)$

$$S_{H}\rho = \underset{\nu \in \mathcal{V}_{H}}{\operatorname{arg\,min}} d_{\mathrm{KL}}\left(\nu \parallel \rho\right), \qquad (3.23)$$

where d_{KL} is the KL divergence between two probability measures. The minimization is among the probability measures in the following set

$$\mathcal{V}_H\left(\bar{H}, C^H\right) = \left\{\nu \in \mathcal{P}\left(\mathbb{R}^d\right) : \mathbb{E}_{\nu}H = \bar{H}, \ \mathbb{E}_{\nu}\left[H'H'^{\mathsf{T}}\right] = C^H, \ \text{and} \ \mathbb{E}_{\nu}\left[H'_{n}H'_{m}H'_{n}\right] = 0\right\}$$

for all $l, m, n \leq d$ and $H'(Z) = H(Z) - \mathbb{E}_{\nu}[H(Z)].$

In Definition 5, S_H acts as a symmetric approximation of probability measures with vanishing third-order moments of the observation function H, while maintains consistent first two leading moments of H. It is clear that given \bar{H}, C^H , the set \mathcal{V}_H is closed with respect to weak convergence of measures. From Proposition 2.1 of [46] and [16], we have for any ρ and weakly convergent sequence $\{\nu_n\}$ to ν_*

$$\liminf_{n \to \infty} d_{\mathrm{KL}} \left(\nu_n \parallel \rho \right) \ge d_{\mathrm{KL}} \left(\nu_* \parallel \rho \right).$$

It follows immediately that there exists $\nu_* \in \mathcal{V}_H$ that reaches the minimum. Therefore, we have the following lemma guaranteeing the existence of the minimizer in the proposed projection (3.23).

Lemma 6. Assume that there is one $\nu \in \mathcal{V}_H$ such that the KL-divergence $d_{\mathrm{KL}}(\nu \parallel \rho) < \infty$ for given $\rho \in \mathcal{P}(\mathbb{R}^d)$. Then a minimizer exists in (3.23).

Even though Lemma 6 does not guarantee the uniqueness of the minimizer, we can always find one minimizer given the same mean and covariance $\overline{H} = \mathbb{E}[H(Z)], C^H = \mathbb{E}[H'(Z)H'(Z)^{\mathsf{T}}]$. This provides the desirable target density function satisfying the required symmetric statistics about $H(Z_t)$. Denote the push-forward operator for the new SDE (3.18) with the structure functions (3.21) as $\tilde{\rho}_{t+s} = \mathcal{Q}_t^s(\tilde{\rho}_t; Y_{t+s})$ for any $s \ge 0$. Under the above construction, we can finally propose the forward operator in analysis step (3.14) as

$$\mathcal{A}_t^s(\tilde{\rho}_t) \coloneqq \mathcal{Q}_t^s(\mathcal{S}_H \tilde{\rho}_t; Y_{t+s}), \qquad (3.24)$$

where the projection S_H in (3.23) is a linear operator acting on the random fields in space \mathcal{V}_t in (3.6). In a similar fashion as in the proof of Lemma 4, by applying the addition projection $S_H \tilde{\rho}_t$ in the expectations on the SDEs for $H\left(\tilde{Z}_t\right)$ and $H'_t\left(\tilde{Z}_t\right)H'_t\left(\tilde{Z}_t\right)^{\mathsf{T}}$ (see also (B10) and (B11) in Appendix B), $Q_t^H = 0$ is automatically guaranteed in (3.22) w.r.t. the new projected density $S_H \tilde{\rho}_t$. In addition, the first two moments \bar{H}_t and C_t^H (3.22) will stay the same w.r.t. $S_H \tilde{\rho}_t$. The continuous filtering process by letting $s \to 0$ will satisfy the equations (3.22) with $Q_t^H \equiv 0$. Therefore, under the same initial condition and the uniqueness of the solution, the same solution will be reached in both (3.17) and (3.22). This leads to the main result of this section concerning the analysis step update in the approximating filter solution.

Theorem 7. Consider the analysis step update (3.24) of the statistical filtering model (3.18). Assume that a_t, K_t in the statistical filtering SDE are designed to satisfy (3.21) and the probability set \mathcal{V}_H according to H defined in (3.23) is not empty. Under the same statistical observations $y_t, t \in [0, T]$ and the same initial conditions, the following relations hold for $t \in [0, T]$

$$\mathcal{H}\hat{\rho}_t = \tilde{\mathbb{E}}\left[H\left(\tilde{Z}_t\right)\right], \quad \mathcal{H}\hat{\mathcal{C}}_t\mathcal{H}^* = \tilde{\mathbb{E}}\left[H'\left(\tilde{Z}_t\right)H'\left(\tilde{Z}_t\right)^{\mathsf{T}}\right], \tag{3.25}$$

where $(\hat{\rho}_t, \hat{C}_t)$ is the solution of (3.17), and $\tilde{\rho}_t$ is given by the solution of (3.19) with $H' = H - \mathcal{H}\hat{\rho}_t$.

Theorem 7 validates the use of the statistical filtering model density $\tilde{\rho}_t$ by solving (3.16) to approximate the optimal filter $\hat{\rho}_t$ from (3.12). Though restricted only on the first two moments of the observation function H, the resulting consistent statistics during analysis step play a key role in accurate statistical forecast. Notice that based on the statistical model in (2.7), accurate prediction of the important leading statistics, \bar{u}_t, R_t , is determined by key higher-order feedbacks in the related functional $\mathcal{H}\rho_t$ (more specifically, the terms $\mathcal{H}_m\rho_t$ and $\mathcal{H}_v\rho_t$). According to Proposition 3, the optimal $\mathcal{H}\hat{\rho}_t$ gives the least mean square estimate of the random variable $\mathcal{H}\rho_t$ given the statistical observations. Thus, consistent approximating filter $\tilde{\mathbb{E}}H\left(\tilde{Z}_t\right)$ for $\mathcal{H}\hat{\rho}_t$ as well as its error estimate guarantees accurate recovery of key model statistics. For example, applying the explicit forms of the observation function (3.4), the quadratic observation operator \mathcal{H}_m in the mean equation gives

$$\tilde{\mathbb{E}}H^{m}\left(\tilde{Z}_{t}\right) = \mathcal{H}_{m}\hat{\rho}_{t} \iff \sum_{p,q}\gamma_{kpq}\tilde{\mathbb{E}}\left[\tilde{Z}_{p,t}\tilde{Z}_{q,t}\right] = \sum_{p,q}\gamma_{kpq}\int z_{p}z_{q}\hat{\rho}_{t}\left(z\right)\mathrm{d}z,$$

which implies consistent statistical feedbacks from the statistical filtering model $\tilde{\rho}_t$ and the optimal filter solution $\hat{\rho}_t$. This demonstrates that the new approximating filter maintains the accuracy in the statistical mean prediction \bar{u}_t . In addition, the covariance operator characterizes the essential uncertainty in the optimal filter estimate $\hat{\rho}_t$ as a random field, that is,

$$\mathcal{H}_m \hat{\mathcal{C}}_t \mathcal{H}_m^* = \mathbb{E}_{\mu_t} \left[\left(\mathcal{H}_m \rho_t - \mathcal{H}_m \hat{\rho}_t \right) \left(\mathcal{H}_m \rho_t - \mathcal{H}_m \hat{\rho}_t \right)^\mathsf{T} \right].$$

It is also linked to the approximation by $C_t^{H^m} = \tilde{\mathbb{E}} \left[\left(H_t^m - \bar{H}_t^m \right) \left(H_t^m - \bar{H}_t^m \right)^{\mathsf{T}} \right] = \mathcal{H}_m \hat{\mathcal{C}}_t \mathcal{H}_m^*$, demonstrating a consistent error estimate in the statistical filtering model. Similar conclusion can be reached for the accurate prediction in the model covariance prediction for R_t based on the cubic observation operator \mathcal{H}_v .

Remark. Still, the statistical consistency in the analysis step does not guarantee the consistency in the entire two-step updating procedure in (3.15). In particular, the forecast models of (3.12) and (3.16) have the following updating equations during the forecast step

$$\frac{\partial_t \hat{\rho}_t = \mathcal{L}_t^* \hat{\rho}_t}{\partial_t \tilde{\rho}_t = \mathcal{L}_t^* \tilde{\rho}_t} \Rightarrow \frac{\partial_t \left(\mathcal{H}\hat{\rho}_t\right) = \mathcal{H}\mathcal{L}_t^* \hat{\rho}_t = \int \left(\mathcal{L}_t H\right) \hat{\rho}_t \left(z\right) \mathrm{d}z}{\partial_t \left(\tilde{\mathbb{E}}H\right) = \tilde{\mathbb{E}}\mathcal{L}_t H = \int \left(\mathcal{L}_t H\right) \tilde{\rho}_t \left(z\right) \mathrm{d}z}$$

where the generator is defined with $y_t = (\bar{u}_t, R_t)$

$$\mathcal{L}_{t}(y_{t})\varphi = \nabla_{z}\varphi \cdot \left\{ \left[H\left(z\right) - R_{t}\right] + L\left(\bar{u}_{t}\right)z\right\} + \frac{1}{2}\Sigma_{t}^{2}: \nabla_{z}\nabla_{z}\varphi$$

The analysis step update only gives consistent first two moments of H, while higher moments may be included in $\mathcal{L}_t H$. It is not guaranteed that the forecast model can give consistent forecast in $\mathcal{H}\hat{\rho}_t$ and $\mathbb{E}H$ as well as their covariances. More work is still needed for the complete consistency analysis combining the approximations in both the forecast and analysis step of the filtering method.

4 Stability and convergence of the statistical filtering model

Here, we discuss the long-time performance of the full filtering problem for the statistical filtering system (3.1) for finding the optimal filter PDF $\hat{\rho}_t$ based on statistical observations $\mathcal{G}_t = \sigma \{y_s, s \leq t\}$. In Section 3, it shows that the statistical filtering model (3.16) constitutes the approximate filter solution $\tilde{\rho}_t$ with consistent mean and covariance, \bar{H}_t and C_t^H , in the analysis step update. We show further here that the full filter approximation of the observation function \bar{H}_t will approach the optimal filter $\mathcal{H}\hat{\rho}_t$ at the long-time limit as $t \to \infty$. This guarantees the stable performance of the proposed new filtering strategy.

4.1 Complete statistical filter equations based on the observation operator

We consider the optimal filter solution based on the conditional Gaussian model (3.1). The finite-dimensional statistical states $\mathcal{H}\hat{\rho}_t \in \mathbb{R}^p$ and $\hat{\mathcal{C}}_t^{\mathcal{H}} = \mathcal{H}\hat{\mathcal{C}}_t\mathcal{H}^* \in \mathbb{R}^{p \times p}$ under the observation operator are solved by the Kalman-Bucy equations (3.12) as

$$d(\mathcal{H}\hat{\rho}_{t}) = \langle \mathcal{L}_{t}(y_{t})H, \hat{\rho}_{t}\rangle dt + \hat{\mathcal{C}}_{t}^{\mathcal{H}}\Gamma^{-2} \{dy_{t} - [\mathcal{H}\hat{\rho}_{t} + h_{t}(y_{t})]dt\}, d\hat{\mathcal{C}}_{t}^{\mathcal{H}} = \left[\left\langle \mathcal{L}_{t}(y_{t})H, \hat{\mathcal{C}}_{t}H^{\mathsf{T}}\right\rangle + \left\langle \hat{\mathcal{C}}_{t}H, \mathcal{L}_{t}(y_{t})H^{\mathsf{T}}\right\rangle\right] dt - \hat{\mathcal{C}}_{t}^{\mathcal{H}}\Gamma^{-2}\hat{\mathcal{C}}_{t}^{\mathcal{H}}dt.$$

$$(4.1)$$

Above, $\langle F, G \rangle = \int F(z) G(z) dz$ denotes the componentwise integration of the product of matrix-valued functions F, G. For simplicity, we use constant observation noise, $\Gamma_t \equiv \Gamma$. In the first equation for the mean, we rewrite the forecast step dynamics as

$$\mathcal{HL}_{t}^{*}(y_{t}) \hat{\rho}_{t} = \int H(z) \mathcal{L}_{t}^{*}(y_{t}) \hat{\rho}_{t}(z) dz = \langle \mathcal{L}_{t}(y_{t}) H, \hat{\rho}_{t} \rangle$$

Similarly in the covariance equation, we rewrite using the definition of \hat{C}_t in (3.10) under the conditional measure $\mu_t(\cdot) = \mathbb{P}(\rho \in \cdot | \mathcal{G}_t)$

$$\mathcal{HL}_{t}^{*}(y_{t})\hat{\mathcal{C}}_{t}\mathcal{H}^{*} = \mathbb{E}_{\rho\sim\mu_{t}}\int H(x)\mathcal{L}_{t}^{*}(\rho-\hat{\rho}_{t})(x)dx\int(\rho-\hat{\rho}_{t})(z)H(z)^{\mathsf{T}}dz = \left\langle\mathcal{L}_{t}H,\hat{\mathcal{C}}_{t}H^{\mathsf{T}}\right\rangle,$$
$$\mathcal{H\hat{\mathcal{C}}}_{t}\mathcal{L}_{t}(y_{t})\mathcal{H}^{*} = \mathbb{E}_{\rho\sim\mu_{t}}\int H(z)(\rho-\hat{\rho}_{t})(z)dz\int(\rho-\hat{\rho}_{t})(x)\mathcal{L}_{t}H(x)^{\mathsf{T}}dx = \left\langle\hat{\mathcal{C}}_{t}H,\mathcal{L}_{t}H^{\mathsf{T}}\right\rangle,$$

where $\hat{\mathcal{C}}_{t}H(x) = \mathbb{E}_{\rho}\left[\left(\rho - \hat{\rho}_{t}\right)(x)\int\left(\rho - \hat{\rho}_{t}\right)(z)H(z)\,\mathrm{d}z\right] \in \mathbb{R}^{d}$ can be viewed as a unnormalized density.

Correspondingly, consider the approximating filter model (3.16) with $\Sigma_t \equiv 0$. Let $\tilde{\rho}_t$ be the PDF after the symmetric projection (3.23), that is, satisfying vanishing third-order moments $\tilde{\mathbb{E}}\left[H'_nH'_pH'_q\right] = 0$ for all n, p, q. The moments $\bar{H}_t = \tilde{\mathbb{E}}\left[H\left(\tilde{Z}_t\right)\right]$ and $C_t^H = \tilde{\mathbb{E}}\left[H'_t\left(\tilde{Z}_t\right)H'_t\left(\tilde{Z}_t\right)^{\mathsf{T}}\right]$ according to the observation function H satisfy the following equations (by combining the forecast step update with generator \mathcal{L}_t and the analysis step dynamics (3.22))

$$d\bar{H}_{t} = \langle \mathcal{L}_{t}(y_{t})H, \tilde{\rho}_{t}\rangle dt + C_{t}^{H}\Gamma^{-2} \left\{ dy_{t} - \left[\bar{H}_{t} + h_{t}(y_{t})\right]dt \right\}, dC_{t}^{H} = \left[\left\langle \mathcal{L}_{t}(y_{t})H, \tilde{\rho}_{t}H_{t}^{\prime \mathsf{T}} \right\rangle + \left\langle \tilde{\rho}_{t}H_{t}^{\prime}, \mathcal{L}_{t}(y_{t})H^{\mathsf{T}} \right\rangle \right] dt - C_{t}^{H}\Gamma^{-2}C_{t}^{H}dt,$$

$$(4.2)$$

where $H'_t(z) = H(z) - \overline{H}_t$.

First, we introduce the following assumptions for the approximating filter PDF $\tilde{\rho}_t$ approximating the optimal filter solution $\hat{\rho}_t$ as random fields according to the same observation process $Y = \{y_t, t \ge 0\}$:

Assumption. Assume that the optimal filter (3.12) and the approximating filter model (3.16) have a continuous probability density functions $\hat{\rho}_t$ and $\tilde{\rho}_t$ respectively that satisfy the following conditions:

- Unique equilibrium solutions $\left(\mathcal{H}\hat{\rho}_{\infty},\hat{\mathcal{C}}_{\infty}^{\mathcal{H}}\right)$ and $\left(\mathcal{H}\tilde{\rho}_{\infty},C_{\infty}^{H}\right)$ exist to the systems (4.1) and (4.2) respectively.
- There exist deterministic matrices $L_{\infty}^{m}, L_{\infty}^{v} \in \mathbb{R}^{p \times p}$. The generator \mathcal{L}_{t} (2.15) reaches the same statistical limit under H w.r.t. both $\hat{\rho}_{t}$ and $\tilde{\rho}_{t}$, that is,

$$\langle \mathcal{L}_{t}(y_{t}) H, \hat{\rho}_{t} \rangle \to L_{\infty}^{m} \mathcal{H} \hat{\rho}_{\infty}, \ \left\langle \mathcal{L}_{t}(y_{t}) H, \hat{\mathcal{C}}_{t} H^{\mathsf{T}} \right\rangle \to L_{\infty}^{v} \hat{\mathcal{C}}_{\infty}^{\mathcal{H}},$$

$$\langle \mathcal{L}_{t}(y_{t}) H, \tilde{\rho}_{t} \rangle \to L_{\infty}^{m} \mathcal{H} \tilde{\rho}_{\infty}, \ \left\langle \mathcal{L}_{t}(y_{t}) H, \tilde{\rho}_{t} H_{t}^{\mathsf{T}} \right\rangle \to L_{\infty}^{v} C_{\infty}^{H},$$

$$(4.3)$$

a.s. as $t \to \infty$ given the observation data y_t .

• Further, the real parts of eigenvalues of the limit matrices $L_{\infty}^m - \hat{\mathcal{C}}_{\infty}^{\mathcal{H}} \Gamma^{-2}$ and $L_{\infty}^v - \hat{\mathcal{C}}_{\infty}^{\mathcal{H}} \Gamma^{-2}$ are all negative.

The first condition in assumption (4.3) guarantees that both filter equations will finally converge to finite solutions. The second condition requires consistent first and second-order moments under H w.r.t. the optimal and approximate model PDF, such as for the covariance as $t \to \infty$

$$\int \mathcal{L}_{t}(y_{t}) H(z) \hat{\mathcal{C}}_{t} H(z)^{\mathsf{T}} dz \to L_{\infty}^{v} \int H(z) \left(\hat{\mathcal{C}}_{\infty} H^{\mathsf{T}} \right)(z) dz,$$
$$\int \mathcal{L}_{t}(y_{t}) H_{t}'(z) H_{t}'(z)^{\mathsf{T}} \tilde{\rho}_{t}(z) dz \to L_{\infty}^{v} \int H_{\infty}'(z) H_{\infty}'(z)^{\mathsf{T}} \tilde{\rho}_{\infty}(z) dz.$$

In addition, we may further introduce the convergence rate, that is, there exist constants $\overline{\lambda} > 0$ and K > 0 such that a.s.

$$\begin{aligned} |\langle \mathcal{L}_{t}(y_{t})H, \hat{\rho}_{t}\rangle - L_{\infty}^{m}\mathcal{H}\hat{\rho}_{\infty}| &\leq Ke^{-\bar{\lambda}t}, \quad \left\| \left\langle \mathcal{L}_{t}(y_{t})H, \hat{\mathcal{C}}_{t}H^{\mathsf{T}} \right\rangle - L_{\infty}^{v}\hat{\mathcal{C}}_{\infty}^{\mathcal{H}} \right\| \leq Ke^{-\bar{\lambda}t}, \\ |\langle \mathcal{L}_{t}(y_{t})H, \tilde{\rho}_{t}\rangle - L_{\infty}^{m}\mathcal{H}\tilde{\rho}_{\infty}| &\leq Ke^{-\bar{\lambda}t}, \quad \left\| \left\langle \mathcal{L}_{t}(y_{t})H, \tilde{\rho}_{t}H_{t}^{\mathsf{T}} \right\rangle - L_{\infty}^{v}C_{\infty}^{\mathcal{H}} \right\| \leq Ke^{-\bar{\lambda}t}, \end{aligned}$$
(4.4)

where $\|\cdot\|$ is the matrix norm. And the third condition asks that the filter solutions will be stabilized at the long time limit. Next, we ask the limit behaviour in the mean and covariance (\bar{H}_t, C_t^H) from the approximation model (4.2) in comparison with the optimal filter solution $(\mathcal{H}\hat{\rho}_t, \hat{C}_t^{\mathcal{H}})$ from (4.1).

4.2 Asymptotic stability of the equilibrium covariance matrix

With the assumption (4.3), we have consistent equilibrium covariance at $t \to \infty$ in the two model solutions

$$L^{v}_{\infty}\hat{\mathcal{C}}^{\mathcal{H}}_{\infty} + \hat{\mathcal{C}}^{\mathcal{H}}_{\infty}L^{v\intercal}_{\infty} - \hat{\mathcal{C}}^{\mathcal{H}}_{\infty}\Gamma^{-2}\hat{\mathcal{C}}^{\mathcal{H}}_{\infty} = 0,$$

$$L^{v}_{\infty}C^{\mathcal{H}}_{\infty} + C^{\mathcal{H}}_{\infty}L^{v\intercal}_{\infty} - C^{\mathcal{H}}_{\infty}\Gamma^{-2}C^{\mathcal{H}}_{\infty} = 0.$$

Uniqueness of the solution directly implies that the final equilibrium covariances satisfy $\hat{\mathcal{C}}^{\mathcal{H}}_{\infty} = C^{H}_{\infty}$ with no randomness. Further, we have that the covariance C^{H}_{t} in the approximating filter will approach the optimal equilibrium covariance $\hat{\mathcal{C}}^{\mathcal{H}}_{\infty}$ as described in the following result.

Lemma 8. Suppose that Assumption 4.1 is satisfied, C_t^H is the covariance solution to the statistical filtering model (4.2) and $\hat{\mathcal{C}}^{\mathcal{H}}_{\infty}$ is the unique equilibrium solutions to the optimal model (4.1). Then there is

$$\left\| C_t^H - \hat{\mathcal{C}}_{\infty}^{\mathcal{H}} \right\| \to 0, \tag{4.5}$$

a.s. as $t \to \infty$. Further, the convergence rate will be exponential if (4.4) is also satisfied.

Proof. Combining the covariance equation in (4.2) and the equilibrium equation of (4.1), we have

$$d\left(C_{t}^{H}-\hat{C}_{\infty}^{\mathcal{H}}\right) = \left[L_{\infty}^{v}-\frac{1}{2}\left(C_{t}^{H}+\hat{C}_{\infty}^{\mathcal{H}}\right)\Gamma^{-2}\right]\left(C_{t}^{H}-\hat{C}_{\infty}^{\mathcal{H}}\right)dt +\left(C_{t}^{H}-\hat{C}_{\infty}^{\mathcal{H}}\right)\left[L_{\infty}^{v\intercal}-\frac{1}{2}\Gamma^{-2}\left(C_{t}^{H}+\hat{C}_{\infty}^{\mathcal{H}}\right)\right]dt +\left[\langle\mathcal{L}_{t}H,\tilde{\rho}_{t}H_{t}^{\prime\intercal}\rangle-L_{\infty}^{v}C_{t}^{H}\right]dt +\left[\langle\tilde{\rho}_{t}H_{t}^{\prime},\mathcal{L}_{t}H^{\intercal}\rangle-C_{t}^{H}L_{\infty}^{v\intercal}\right]dt.$$

For the last row of the above equation, using the uniqueness of the solution we have $C_t^H \to C_\infty^H$ as $t \to \infty$. Then denote

$$F_t = \langle \mathcal{L}_t H, \tilde{\rho}_t H_t^{\prime \mathsf{T}} \rangle - L_{\infty}^v C_t^H.$$

We get $|F_t| \to 0$ as $t \to \infty$ by using

$$\left\| \left\langle \mathcal{L}_t H, \tilde{\rho}_t H_t^{\prime \mathsf{T}} \right\rangle - L_{\infty}^v C_t^H \right\| \le \left\| \left\langle \mathcal{L}_t H, \tilde{\rho}_t H_t^{\prime \mathsf{T}} \right\rangle - L_{\infty}^v C_{\infty}^H \right\| + \left\| L_{\infty}^v \left(C_{\infty}^H - C_t^H \right) \right\| \to 0.$$

Also by taking $\lambda_v = \min \left\{ \operatorname{Re}\lambda : \lambda \text{ is the eighenvalue of } -L_{\infty}^v + \hat{\mathcal{C}}_{\infty}^{\mathcal{H}}\Gamma^{-2} \right\}$, we have $\lambda_v > 0$ from Assumption 4.1. This implies $\left\| e^{\int_s^t \left[L_{\infty}^v - \frac{1}{2} \left(C_{\tau}^H + \hat{\mathcal{C}}_{\infty}^{\mathcal{H}} \right) \Gamma^{-2} \right] \mathrm{d}\tau} \right\| \leq K e^{-\lambda_v (t-s)}$. Together with F_t vanishing as $t \to \infty$, we have for any t > T

$$\left\| C_t^H - \hat{\mathcal{C}}_{\infty}^{\mathcal{H}} \right\| \le K \left\| C_T^H - \hat{\mathcal{C}}_{\infty}^{\mathcal{H}} \right\| e^{-2\lambda_v(t-T)} + 2K \int_T^t e^{-2\lambda_v(t-s)} \left| F_s \right| \mathrm{d}s \le K_1 e^{-2\lambda_v t} + K_2 \sup_{s \ge T} \left| F_s \right| \mathrm{d}s$$

Therefore, by first letting $t \to \infty$ then letting $T \to \infty$, we reach a.s. $\lim_{t\to\infty} \left\| C_t^H - \hat{\mathcal{C}}_{\infty}^H \right\| = 0.$

Further, if we assume exponential convergence rate $\overline{\lambda}$ in the covariances under the generator \mathcal{L}_t as in (4.4), we can have exponential convergence rate in both $\hat{\mathcal{C}}_t$ and C_t^H as $t \to \infty$ a.s.

$$\left\| \hat{\mathcal{C}}_{t}^{\mathcal{H}} - \hat{\mathcal{C}}_{\infty}^{\mathcal{H}} \right\| \leq K_{H} e^{-\min\{\lambda_{v},\bar{\lambda}\}t}, \ \left\| C_{t}^{H} - \hat{\mathcal{C}}_{\infty}^{\mathcal{H}} \right\| \leq K_{H} e^{-\min\{\lambda_{v},\bar{\lambda}\}t}.$$

$$(4.6)$$

In addition, from the definition of the optimal filter solution (3.11), the covariance is defined as

$$\begin{aligned} \operatorname{tr} \mathbb{E} \hat{\mathcal{C}}_{t}^{\mathcal{H}} &= \mathbb{E}_{\mu_{t}} \operatorname{tr} \left[\left(\mathcal{H} \rho_{t} - \mathcal{H} \hat{\rho}_{t} \right) \left(\mathcal{H} \rho_{t} - \mathcal{H} \hat{\rho}_{t} \right)^{\mathsf{T}} \right] \\ &= \mathbb{E}_{\mu_{t}} \operatorname{tr} \left[\left(\mathcal{H} \rho_{t} - \mathcal{H} \hat{\rho}_{t} \right)^{\mathsf{T}} \left(\mathcal{H} \rho_{t} - \mathcal{H} \hat{\rho}_{t} \right) \right] \\ &= \mathbb{E}_{\mu_{t}} \left[\left| \mathcal{H} \rho_{t} - \mathcal{H} \hat{\rho}_{t} \right|^{2} \right]. \end{aligned}$$

Notice that $\mathcal{H}\hat{\rho}_t$ and $\hat{\mathcal{C}}_t^{\mathcal{H}}$ are still random field in \mathcal{G}_t . Under Assumption 4.1 $\hat{\mathcal{C}}_t^{\mathcal{H}} \to \hat{\mathcal{C}}_{\infty}^{\mathcal{H}}$ a.s. with the observations in \mathcal{G}_t , we have as $t \to \infty$

$$\mathbb{E}\left[\left|\mathcal{H}\rho_t - \mathcal{H}\hat{\rho}_t\right|^2\right] = \operatorname{tr}\mathbb{E}\left[\hat{\mathcal{C}}_t^{\mathcal{H}}\right] \to \operatorname{tr}\hat{\mathcal{C}}_{\infty}^{\mathcal{H}}.$$
(4.7)

This confirms that the total uncertainty at equilibrium in the optimal filter solution $\mathcal{H}\hat{\rho}_t$ is estimated by the total variance $\operatorname{tr}\hat{\mathcal{C}}^{\mathcal{H}}_{\infty}$.

4.3 Convergence of the statistical state under the observation operator

Next, we consider the convergence of the statistical observation function \bar{H}_t from the approximating filter to the optimal filter solution $\mathcal{H}\hat{\rho}_t$. We have the long-term stability in the statistical solution in the following theorem.

Theorem 9. Suppose that Assumption 4.1 holds and the covariances goes to the same deterministic limit

$$\hat{\mathcal{C}}_t^{\mathcal{H}} \to \hat{\mathcal{C}}_\infty^{\mathcal{H}}, \quad C_t^H \to \hat{\mathcal{C}}_\infty^{\mathcal{H}},$$

a.s. as $t \to \infty$. Then there is

$$\mathbb{E}\left[\left|\mathcal{H}\hat{\rho}_t - \bar{H}_t\right|^2\right] \to 0,\tag{4.8}$$

as $t \to \infty$. Furthermore, assume exponential convergence rate in $\hat{\mathcal{C}}_t^{\mathcal{H}}, \mathcal{C}_t^{\mathcal{H}}$ as in (4.6), and let $\lambda = \min \{\lambda_m, \lambda_v\}$ where λ_m and λ_v are the minimums of the real parts of the eigenvalues of $-L_{\infty}^m + \hat{\mathcal{C}}_{\infty}^{\mathcal{H}}\Gamma^{-2}$ and $-L_{\infty}^v + \hat{\mathcal{C}}_{\infty}^{\mathcal{H}}\Gamma^{-2}$ respectively. There is also exponential convergence as

$$\mathbb{E}\left[\left|\mathcal{H}\hat{\rho}_t - \bar{H}_t\right|^2\right] \le K_H e^{-\lambda_1 t},\tag{4.9}$$

with some $\lambda_1 \leq \min \{\lambda, \bar{\lambda}\}$ and K_H a constant only dependent on the observation function H.

Proof. By taking the difference of the mean equations in (4.1) and (4.2), we have

$$d\left(\mathcal{H}\hat{\rho}_{t}-\bar{H}_{t}\right) = \left(L_{\infty}^{m}-\hat{\mathcal{C}}_{\infty}^{\mathcal{H}}\Gamma^{-2}\right)\left(\mathcal{H}\hat{\rho}_{t}-\bar{H}_{t}\right)dt + \left(\hat{\mathcal{C}}_{t}^{\mathcal{H}}-C_{t}^{H}\right)\Gamma^{-2}\left[dy_{t}-h_{t}\left(y_{t}\right)dt\right] \\ + \left[\left(\hat{\mathcal{C}}_{\infty}^{\mathcal{H}}-\hat{\mathcal{C}}_{t}^{\mathcal{H}}\right)\Gamma^{-2}\mathcal{H}\hat{\rho}_{t}-\left(\hat{\mathcal{C}}_{\infty}^{\mathcal{H}}-C_{t}^{H}\right)\Gamma^{-2}\bar{H}_{t}\right]dt \\ + \left[\left\langle\mathcal{L}_{t}\left(y_{t}\right)H,\hat{\rho}_{t}\right\rangle-L_{\infty}^{m}\mathcal{H}\hat{\rho}_{t}\right]dt + \left[\left\langle\mathcal{L}_{t}\left(y_{t}\right)H,\tilde{\rho}_{t}\right\rangle-L_{\infty}^{m}\bar{H}_{t}\right]dt.$$

By applying Itô's formula to the above equation, there is

$$d\left[e^{-t\left(L_{\infty}^{m}-\hat{\mathcal{C}}_{\infty}^{\mathcal{H}}\Gamma^{-2}\right)}\left(\mathcal{H}\hat{\rho}_{t}-\bar{H}_{t}\right)\right] = e^{-t\left(L_{\infty}^{m}-\hat{\mathcal{C}}_{\infty}^{\mathcal{H}}\Gamma^{-2}\right)}\left\{\left(\hat{\mathcal{C}}_{t}^{\mathcal{H}}-C_{t}^{H}\right)\Gamma^{-2}\left[dy_{t}-h_{t}\left(y_{t}\right)dt\right]+G_{t}\left(y_{t}\right)dt\right\}\right.\\ \left.+e^{-t\left(L_{\infty}^{m}-\hat{\mathcal{C}}_{\infty}^{\mathcal{H}}\Gamma^{-2}\right)}\left[\left(\hat{\mathcal{C}}_{\infty}^{\mathcal{H}}-\hat{\mathcal{C}}_{t}^{\mathcal{H}}\right)\Gamma^{-2}\mathcal{H}\hat{\rho}_{t}-\left(\hat{\mathcal{C}}_{\infty}^{\mathcal{H}}-C_{t}^{H}\right)\Gamma^{-2}\bar{H}_{t}\right]dt.$$

Above, we denote the residual term as

$$G_{t}(y_{t}) = \left[\left\langle \mathcal{L}_{t}(y_{t}) H, \hat{\rho}_{t} \right\rangle - L_{\infty}^{m} \mathcal{H} \hat{\rho}_{t} \right] + \left[\left\langle \mathcal{L}_{t}(y_{t}) H, \tilde{\rho}_{t} \right\rangle - L_{\infty}^{m} \bar{H}_{t} \right].$$

By similar computation in Lemma 8 according to assumption (4.3), there is

$$|G_t| \leq \frac{|\langle \mathcal{L}_t (y_t) H, \hat{\rho}_t \rangle - L_{\infty}^m \mathcal{H} \hat{\rho}_{\infty}| + |L_{\infty}^m (\mathcal{H} \hat{\rho}_{\infty} - \mathcal{H} \hat{\rho}_t)|}{+ |\langle \mathcal{L}_t (y_t) H, \tilde{\rho}_t \rangle - L_{\infty}^m \bar{H}_{\infty}| + |L_{\infty}^m (\bar{H}_{\infty} - \bar{H}_t)|} \leq K e^{-\bar{\lambda}t} \to 0$$

a.s. as $t \to \infty$ using the uniqueness of the solutions $\mathcal{H}\hat{\rho}_t \to \mathcal{H}\hat{\rho}_\infty$ and $\bar{H}_t \to \bar{H}_\infty$ (with exponential decay in the stronger convergence case (4.6)). Equivalently, the above SDE can be written as (ignoring initial condition by assuming $\mathcal{H}\hat{\rho}_0 = \bar{H}_0$)

$$\mathcal{H}\hat{\rho}_{t} - \bar{H}_{t} = \int_{0}^{t} e^{(t-s)\left(L_{\infty}^{m} - \hat{\mathcal{C}}_{\infty}^{\mathcal{H}}\Gamma^{-2}\right)} \left\{ \left(\hat{\mathcal{C}}_{s}^{\mathcal{H}} - C_{s}^{H}\right)\Gamma^{-2}\left[\mathrm{d}y_{s} - h_{s}\left(y_{s}\right)\mathrm{d}s\right] + G_{s}\left(y_{s}\right)\mathrm{d}s \right\} + \int_{0}^{t} e^{(t-s)\left(L_{\infty}^{m} - \hat{\mathcal{C}}_{\infty}^{\mathcal{H}}\Gamma^{-2}\right)} \left[\left(\hat{\mathcal{C}}_{\infty}^{\mathcal{H}} - \hat{\mathcal{C}}_{s}^{\mathcal{H}}\right)\Gamma^{-2}\mathcal{H}\hat{\rho}_{s} - \left(\hat{\mathcal{C}}_{\infty}^{\mathcal{H}} - C_{s}^{H}\right)\Gamma^{-2}\bar{H}_{s} \right]\mathrm{d}s.$$

Therefore, by taking the expectation for each term on the right hand side of the above identity

$$\begin{split} \mathbb{E}\left[\left|\mathcal{H}\hat{\rho}_{t}-\bar{H}_{t}\right|^{2}\right] &\leq 5\mathbb{E}\left|\int_{0}^{t}e^{(t-s)\left(L_{\infty}^{m}-\hat{\mathcal{C}}_{\infty}^{\mathcal{H}}\Gamma^{-2}\right)}\left(\hat{\mathcal{C}}_{s}^{\mathcal{H}}-C_{s}^{H}\right)\Gamma^{-2}\mathrm{d}y_{s}\right|^{2} \\ &+5\mathbb{E}\left|\int_{0}^{t}e^{(t-s)\left(L_{\infty}^{m}-\hat{\mathcal{C}}_{\infty}^{\mathcal{H}}\Gamma^{-2}\right)}G_{s}\left(y_{s}\right)\mathrm{d}s\right|^{2} \\ &+5\mathbb{E}\left|\int_{0}^{t}e^{(t-s)\left(L_{\infty}^{m}-\hat{\mathcal{C}}_{\infty}^{\mathcal{H}}\Gamma^{-2}\right)}\left(\hat{\mathcal{C}}_{s}^{\mathcal{H}}-C_{s}^{H}\right)\Gamma^{-2}h_{s}\left(y_{s}\right)\mathrm{d}s\right|^{2} \\ &+5\mathbb{E}\left|\int_{0}^{t}e^{(t-s)\left(L_{\infty}^{m}-\hat{\mathcal{C}}_{\infty}^{\mathcal{H}}\Gamma^{-2}\right)}\left(\hat{\mathcal{C}}_{\infty}^{\mathcal{H}}-\hat{\mathcal{C}}_{s}^{\mathcal{H}}\right)\Gamma^{-2}\hat{\mathcal{H}}\hat{\rho}_{s}\mathrm{d}s\right|^{2} \\ &+5\mathbb{E}\left|\int_{0}^{t}e^{(t-s)\left(L_{\infty}^{m}-\hat{\mathcal{C}}_{\infty}^{\mathcal{H}}\Gamma^{-2}\right)}\left(\hat{\mathcal{C}}_{\infty}^{\mathcal{H}}-C_{s}^{H}\right)\Gamma^{-2}\bar{H}_{s}\mathrm{d}s\right|^{2}. \end{split}$$

The second line above follows the same argument as in Lemma 8 and using Cauchy-Schwarz inequality

$$\begin{split} \mathbb{E} \left| \int_0^t e^{(t-s) \left(L_\infty^m - \hat{\mathcal{C}}_\infty^{\mathcal{H}} \Gamma^{-2} \right)} G_s\left(y_s \right) \mathrm{d}s \right|^2 &\leq \int_0^t \left\| e^{\frac{1}{2}(t-s) \left(L_\infty^m - \hat{\mathcal{C}}_\infty^{\mathcal{H}} \Gamma^{-2} \right)} \right\|^2 \mathrm{d}s \mathbb{E} \int_0^t \left| e^{\frac{1}{2}(t-s) \left(L_\infty^m - \hat{\mathcal{C}}_\infty^{\mathcal{H}} \Gamma^{-2} \right)} G_s\left(y_s \right) \right|^2 \mathrm{d}s \\ &\leq \lambda_m^{-1} \mathbb{E} \int_0^t e^{-\lambda_m (t-s)} \left| G_s\left(y_s \right) \right|^2 \mathrm{d}s \\ &\leq K e^{-\lambda_1 t} \to 0. \end{split}$$

Similar results can be achieved for line three to five following the convergence (or exponential convergence) of the integrants. Finally, for the first line using the observation equation in (3.1), that is, $dy_t = [\mathcal{H}\rho_t + h_t(y_t)] dt + \Gamma dB_t$, there is

$$\begin{split} & \mathbb{E} \left| \int_{0}^{t} e^{(t-s) \left(L_{\infty}^{m} - \hat{\mathcal{C}}_{\infty}^{\mathcal{H}} \Gamma^{-2} \right)} \left(\hat{\mathcal{C}}_{s}^{\mathcal{H}} - C_{s}^{H} \right) \Gamma^{-2} \mathrm{d}y_{s} \right|^{2} \\ & \leq \mathbb{E} \left| \int_{0}^{t} e^{(t-s) \left(L_{\infty}^{m} - \hat{\mathcal{C}}_{\infty}^{\mathcal{H}} \Gamma^{-2} \right)} \left(\hat{\mathcal{C}}_{s}^{\mathcal{H}} - C_{s}^{H} \right) \Gamma^{-2} \left[\mathcal{H}\rho_{t} + h_{t} \left(y_{t} \right) \right] \mathrm{d}s \right|^{2} \\ & + \mathbb{E} \int_{0}^{t} e^{2(t-s) \left(L_{\infty}^{m} - \hat{\mathcal{C}}_{\infty}^{\mathcal{H}} \Gamma^{-2} \right)} \left\| \hat{\mathcal{C}}_{s}^{\mathcal{H}} - C_{s}^{H} \right\|^{2} \mathrm{d}s \\ & \leq K_{1} \int_{0}^{t} e^{-\lambda_{m}(t-s)} \left(\mathbb{E} \left\| \hat{\mathcal{C}}_{s}^{\mathcal{H}} - \hat{\mathcal{C}}_{\infty}^{\mathcal{H}} \right\|^{2} + \mathbb{E} \left\| C_{s}^{H} - \hat{\mathcal{C}}_{\infty}^{\mathcal{H}} \right\|^{2} \right) \mathrm{d}s \\ & \leq K e^{-\lambda_{1}t} \to 0. \end{split}$$

Above in the second to the last inequality, we use the uniform boundedness of $\mathbb{E} |\mathcal{H}\rho_t|$ and $\mathbb{E} |h_t(y_t)|$ as $t \to \infty$, and the last line uses the convergence (or exponential convergence) of the covariance (4.5) or (4.6). Combining all the above bounds, we finally get (4.8) and (4.9).

Together with the equilibrium estimate in (4.7) combining the result in Theorem 9, we can also get the same error estimate compared with the target field as $t \to \infty$

$$\mathbb{E}\left[\left|\mathcal{H}\rho_{t}-\bar{H}_{t}\right|^{2}\right] \leq \mathbb{E}\left[\left|\mathcal{H}\rho_{t}-\mathcal{H}\hat{\rho}_{t}\right|^{2}\right] + \mathbb{E}\left[\left|\mathcal{H}\hat{\rho}_{t}-\bar{H}_{t}\right|^{2}\right] \rightarrow \mathrm{tr}\hat{\mathcal{C}}_{\infty}^{\mathcal{H}}.$$

Thus, we show that the statistical filtering model solution \bar{H}_t converges to the optimal filter solution with the same mean square error, and exponential convergence is reached if the forecast model has exponential convergence to the equilibrium.

As a final comment, we can further relax Assumption 4.1 as there exist deterministic uniformly continuous functions, $L_{\infty}^{m}(y)$ and $L_{\infty}^{v}(y)$, so that for any $y_{t} \to y_{\infty}, t \to \infty$

$$\langle \mathcal{L}_t (y_t) H, \hat{\rho}_t \rangle \to L^m_{\infty} (y_{\infty}) \mathcal{H} \hat{\rho}_{\infty}, \ \left\langle \mathcal{L}_t H, \hat{\mathcal{C}}_t H^{\mathsf{T}} \right\rangle \to L^v_{\infty} (y_{\infty}) \hat{\mathcal{C}}^{\mathcal{H}}_{\infty}, \left\langle \mathcal{L}_t (y_t) H, \tilde{\rho}_t \right\rangle \to L^m_{\infty} (y_{\infty}) \mathcal{H} \tilde{\rho}_{\infty}, \ \left\langle \mathcal{L}_t H, \tilde{\rho}_t H_t^{\mathsf{T}} \right\rangle \to L^v_{\infty} (y_{\infty}) C^H_{\infty},$$

$$(4.10)$$

a.s. as $t \to \infty$. And the limiting matrices are uniformly bounded by negative-definite matrices A_m, A_v

$$L_{\infty}^{m}(y) - \hat{\mathcal{C}}_{\infty}^{\mathcal{H}} \Gamma^{-2} \preceq A_{m} \prec 0, \quad L_{\infty}^{v}(y) - \hat{\mathcal{C}}_{\infty}^{\mathcal{H}} \Gamma^{-2} \preceq A_{v} \prec 0.$$

$$(4.11)$$

In addition, non-zero noise in (3.16) can be included satisfying $\Sigma_t \to 0$ as $t \to \infty$. Then, the same result applies for the convergence of the observation mean and covariance at long-time limit as in Lemma 8 and Theorem 9.

5 Ensemble approximation of the statistical filtering model

Finally, we discuss the construction of practical ensemble methods for solving the statistical filtering model (3.16) with explicit model parameters that is easy to implement. We demonstrate the accuracy of the numerical approximation at the limit of large number of samples and small integration time step.

5.1 Numerical methods for the approximating filter with discrete observations

Assume that the observation data comes at discrete times $t_n = \delta n$ with a constant observation frequency δ . We can introduce the linear interpolation of $y_n = (\bar{u}_{t_n}, R_{t_n}) \in \mathbb{R}^p$ as

$$\frac{\mathrm{d}y_t^\delta}{\mathrm{d}t} = \frac{y_{n+1} - y_n}{t_{n+1} - t_n} = \frac{\Delta y_{n+1}}{\delta},\tag{5.1}$$

during time interval $t \in [t_n, t_{n+1}]$. We propose ensemble algorithms to approximate the filtering distribution $\tilde{\rho}_{t_n} \sim \tilde{Z}_{t_n}$ conditional on the statistical observations $Y_t^{\delta} = \{y_s^{\delta}, s \leq t\}$ based on the statistical filter equation (3.16). First, N independent particles, $\tilde{\mathbf{Z}}_t = \left\{\tilde{Z}_t^{(i)}\right\}_{i=1}^N$, are drawn to sample the initial distribution of the stochastic state. Then, the particles are evolved according to the following SDE with drift terms a_t^m, a_t^v and control gains K_t^m, K_t^v

$$d\tilde{Z}_{t}^{(i)} = L\left(\bar{u}_{t}^{N}\right)\tilde{Z}_{t}^{(i)}dt + \Gamma\left(\tilde{Z}_{t}^{(i)}\tilde{Z}_{t}^{(i)\mathsf{T}} - R_{t}^{N}\right)dt + \Sigma_{t}d\tilde{W}_{t}^{(i)} + a_{t}^{m}\left(\tilde{Z}_{t}^{(i)}\right)dt + K_{t}^{m}\left(\tilde{Z}_{t}^{(i)}\right)\left\{d\bar{u}_{t}^{\delta} - \left[H^{m}\left(\tilde{Z}_{t}^{(i)}\right) + h_{m,t}\left(\bar{u}_{t}^{\delta}\right)\right]dt - \Gamma_{m,t}d\tilde{B}_{m,t}^{(i)}\right\} + a_{t}^{v}\left(\tilde{Z}_{t}^{(i)}\right)dt + K_{t}^{v}\left(\tilde{Z}_{t}^{(i)}\right)\left\{dR_{t}^{\delta} - \left[H^{v}\left(\tilde{Z}_{t}^{(i)}\right) + h_{v,t}\left(\bar{u}_{t}^{\delta}, R_{t}^{\delta}\right)\right]dt - \Gamma_{v,t}d\tilde{B}_{v,t}^{(i)}\right\},$$

$$(5.2)$$

where the expressions for h_m, h_v are defined in (3.3), H^m, H^v are defined in (3.4), and $\tilde{B}_{m,t}^{(i)}, \tilde{B}_{v,t}^{(i)}$ are independent white noises. Above, in the first line of (5.2) for the forecast step of the filter, the first two moments (\bar{u}_t^N, R_t^N) can be explicitly solved by the statistical equations according to the stochastic-statistical model (2.7)

$$\frac{\mathrm{d}\bar{u}_{t}^{N}}{\mathrm{d}t} = M\left(\bar{u}_{t}^{N}\right) + F_{t} + \mathbb{E}^{N}\left[H^{m}\left(\tilde{\mathbf{Z}}_{t}\right)\right],$$

$$\frac{\mathrm{d}R_{t}^{N}}{\mathrm{d}t} = L\left(\bar{u}_{t}^{N}\right)R_{t}^{N} + R_{t}^{N}L\left(\bar{u}_{t}^{N}\right)^{\mathsf{T}} + \Sigma_{t}\Sigma_{t}^{\mathsf{T}} + \mathbb{E}^{N}\left[H^{v}\left(\tilde{\mathbf{Z}}_{t}\right)\right] + \epsilon^{-1}\left(\mathbb{E}^{N}\left[\tilde{\mathbf{Z}}_{t}\tilde{\mathbf{Z}}_{t}^{\mathsf{T}}\right] - R_{t}^{N}\right).$$
(5.3)

The expectation is computed through the empirical average of the ensemble, $\mathbb{E}^N f\left(\tilde{\mathbf{Z}}\right) = \frac{1}{N} \sum_{i=1}^N f\left(\tilde{Z}^{(i)}\right)$. In this way, the particle simulation of (5.2) can be carried out easily for each individual sample $\tilde{Z}_t^{(i)}$, and the dependence on the distribution of the whole interacting particles is only introduced through the empirical average in the statistical equations (5.3). We summarize the ensemble filtering strategy in Algorithm 1.

Algorithm 1 Ensemble statistical filter with observations in mean and covariance

Model Setup: Get the interpolated sequence of observation (5.1) of the mean and covariance $y_t^{\delta} = \{\bar{u}_t^{\delta}, R_t^{\delta}\}, t \in$

[0, T]; and introduce discrete time integration step τ , and the initial distribution ρ_0 of the model state. **Initial condition:** At the initial time t = 0, draw an ensemble of samples $\left\{\tilde{Z}_0^{(i)}\right\}_{i=1}^N$ from the initial distribution ρ_0 , and compute initial mean and covariance $\left\{\bar{u}_0^N, R_0^N\right\}$ consistent with the statistics w.r.t. ρ_0 .

- 1: for n = 0 while $n < \lfloor T/\tau \rfloor$, during the time updating interval $t \in [t_n, t_{n+1}]$ with $t_n = n\tau$ do 2: Compute the gain functions $K_{t_n}^m$ and $K_{t_n}^v$ using (5.6) and the associated drift terms $a_{t_n}^m$ and $a_{t_n}^v$. 3: Update the samples $\left\{ \tilde{Z}_{t_{n+1}}^{(i)} \right\}_{i=1}^N$ using (5.2) with the statistical states $\{ \bar{u}_t^N, R_t^N \}$ and observation data y_t^{δ} .

Update the statistical mean and covariance $\left\{\bar{u}_{t_{n+1}}^N, R_{t_{n+1}}^N\right\}$ by integrating (5.3) to the next time step using 4: the average of all samples.

5: end for

Remark. Solving the equations (5.2) may still demand high computational cost for resolving the multiple nonlinear coupling terms in high dimension $d \gg 1$. One potential approach to address the computational challenge is to adopt the efficient random batch approach [47, 48] developed for the coupled models (2.7). A detailed investigation of the numerical methods will be performed in the follow-up research.

5.2Construction of explicit model operators in the analysis step

In the second and third lines of (5.2) for the analysis step update of filtering, we still need to propose explicit expressions for functions a_t^m, K_t^m and a_t^v, K_t^v according to the observations of the mean and covariance respectively. According to Theorem 7, the gain function K_t needs to be solved from equation (3.21), that is,

$$-\nabla \cdot \left(K_t^{\mathsf{T}} \tilde{\rho}_t\right) = \tilde{\rho}_t \Gamma_t^{-2} \left(H\left(z\right) - \bar{H}_t\right).$$

Then the drift function a_t can be directly computed from the solution of K_t as

$$a_t = \nabla \cdot \left(K_t \Gamma_t^2 K_t^{\mathsf{T}} \right) - K_t \Gamma_t^2 \nabla \cdot K_t^{\mathsf{T}}.$$

In general, it is still difficult to find solutions of the above equations. By multiplying H on both sides and integrating about z, the identity for K_t implies a necessary condition

$$\tilde{\mathbb{E}}\left[K_t^{\mathsf{T}}\nabla H\right] = \Gamma_t^{-2} C_t^H,\tag{5.4}$$

where $C_t^H = \tilde{\mathbb{E}}\left[\left(H\left(\tilde{Z}_t\right) - \bar{H}_t\right)\left(H\left(\tilde{Z}_t\right) - \bar{H}_t\right)^{\mathsf{T}}\right]$ is the covariance of H. Therefore, we can first design proper gain functions K_t by solving (5.4) according to the specific structures of $H^m : \mathbb{R}^d \to \mathbb{R}^d$ and $H^v : \mathbb{R}^d \to \mathbb{R}^{d^2}$ required in our problem in (3.4)

$$H_{k}^{m}(z) = \sum_{m,n} \gamma_{kmn} z_{m} z_{n} = z^{\mathsf{T}} A_{k} z,$$

$$H_{kl}^{v}(z) = \sum_{m,n} \gamma_{kmn} z_{m} z_{n} z_{l} + \gamma_{lmn} z_{m} z_{n} z_{k} = (z^{\mathsf{T}} A_{k} z) z_{l} + z_{k} (z^{\mathsf{T}} A_{l} z),$$
(5.5)

for all $1 \leq k, l \leq d$ where we rewrite the quadratic and cubic functions using the symmetric coefficient matrix, $A_k^{\mathsf{T}} = A_k \in \mathbb{R}^{d \times d}$, using the assumed structural symmetry in the coupling coefficient γ_{kmn} . The resulting gain functions are then constructed with the following specific expressions.

Proposition 10. The matrix-valued functions $K_t^m = \tilde{K}^m \Gamma_{m,t}^{-2}$ and $K_t^v = \tilde{K}^v \Gamma_{v,t}^{-2}$ with $\tilde{K}^m(z) \in \mathbb{R}^{d \times d}$ and $\tilde{K}^v(z) \in \mathbb{R}^{d \times d^2}$ in the following expressions

$$\tilde{K}_{j,k}^{m}(z) = \frac{1}{2} z_{j} \left[(z^{\mathsf{T}} A_{k} z) - \bar{H}_{k}^{m} \right],$$

$$\tilde{K}_{j,kl}^{v}(z) = \frac{1}{3} z_{j} \left(z^{\mathsf{T}} A_{k} z \right) z_{l} + \frac{1}{3} z_{j} \left(z^{\mathsf{T}} A_{l} z \right) z_{k} - \frac{1}{3} \bar{H}_{kl}^{v}.$$
(5.6)

for $1 \le k, l \le d$ and $1 \le j \le d$ satisfy the equation (5.4) according to the structures of the functions H^m and H^v in the form of (5.5) respectively, and $\bar{H}^m = \tilde{\mathbb{E}}\left[H^m\left(\tilde{Z}\right)\right]$ and $\bar{H}^v = \tilde{\mathbb{E}}\left[H^v\left(\tilde{Z}\right)\right]$.

The proof of Proposition 10 is put in Appendix B. The average terms, \bar{H}^m, \bar{H}^v , are already computed in the statistical equations (5.3) thus no additional computational cost is needed. On the other hand, it is noticed that (5.6) can only give a necessary condition for the gain operators and may not guarantee the original identity for K_t in general. However, in the proof of Theorem 7, it shows that (5.4) is the main relation used to derive the consistent analysis statistics on the mean of $H\left(\tilde{Z}_t\right)$. Therefore, (5.6) provides a desirable candidate for practical implementations of the algorithm concerning the consistency in the leading moments.

5.3 Ensemble approximation in the discrete filtering model

At last, we discuss the accuracy in the finite ensemble approximation. Let $\tilde{\rho}_t^N(z) = \frac{1}{N} \sum_{i=1}^N \delta\left(z - \tilde{Z}_t^{(i)}\right)$ be the random field from the empirical PDF of the ensemble approximation with finite ensemble size N, and $\tilde{\rho}_t$ is the random field from (3.16) conditional on the observations \mathcal{G}_t . Additional assumptions are needed for the structures of the model.

Assumption 11. We assume that the functions $M : \mathbb{R}^d \to \mathbb{R}^d$ and $L : \mathbb{R}^d \to \mathbb{R}^{d \times d}$ in the mean and covariance equations (5.10) are Lipschitz continuous, that is, there is a constant $\beta > 0$ so that

$$|M(u) - M(v)| \le \beta |u - v|, \quad ||L(u) - L(v)|| \le \beta |u - v|.$$

In addition, the coefficients in $\Gamma_k(R) = \sum_{m,n} \gamma_{kmn} R_{mn}$ are uniformly bounded, that is, there exists a constant C > 0, so that for all k, m, n

$$|\gamma_{kmn}| \le C.$$

First, we direct use the conclusion from the McKean-Vlasov limit of the N-particle system [43, 25]. Under the above assumptions of the model coefficients, the ensemble representation (5.2) will converge to the continuous system (3.16) as $N \to \infty$. In particular, assume that the a unique solution exists for the McKean-Vlasov SDE, and the initial ensemble $\left\{\tilde{Z}_{0}^{(i)}\right\}_{i=1}^{N}$ is drawn from i.i.d. random samples. Given the observations \mathcal{G}_{t} , we have for any $\varphi \in C_{b}^{2}(\mathbb{R}^{d})$

$$\langle \varphi, \tilde{\rho}_t^N \rangle = \frac{1}{N} \sum_{i=1}^N \varphi\left(\tilde{Z}_t^{(i)}\right) \to \mathbb{E}\left[\varphi\left(\tilde{Z}_t\right) \mid \mathcal{G}_t\right] = \langle \varphi, \tilde{\rho}_t \rangle,$$
(5.7)

a.s. as $N \to \infty$. Furthermore, there is the error estimate for the empirical estimate $\mathbb{E}^N \varphi = \frac{1}{N} \sum \varphi \left(\tilde{Z}_t^{(i)} \right)$

$$\mathbb{E}\left[\sup_{0\leq t\leq T}\left|\left\langle\varphi,\tilde{\rho}_{t}^{N}\right\rangle-\left\langle\varphi,\tilde{\rho}_{t}\right\rangle\right|^{2}\right]\leq\frac{C_{T}}{N}\left\|\varphi\right\|_{\infty}^{2}.$$
(5.8)

Detailed proofs on (5.7) and (5.8) can be found in such as Chapter 9 of [2].

Next, we consider the finite ensemble and discrete time estimation of the leading-order mean and covariance from the filter model. With the ensemble approximation of the density function $\tilde{\rho}_t^N$, the mean and covariance estimates are computed from the equations (5.3)

$$\frac{\mathrm{d}\bar{u}_{t}^{N,\tau}}{\mathrm{d}t} = M\left(\bar{u}_{\sigma(t)}^{N,\tau}\right) + F_{\sigma(t)} + \frac{1}{N}\sum_{i=1}^{N}H^{m}\left(\tilde{Z}_{\sigma(t)}^{(i)}\right),$$

$$\frac{\mathrm{d}R_{t}^{N,\tau}}{\mathrm{d}t} = L\left(\bar{u}_{\sigma(t)}^{N,\tau}\right)R_{\sigma(t)}^{N,\tau} + R_{\sigma(t)}^{N,\tau}L\left(\bar{u}_{\sigma(t)}^{N,\tau}\right)^{\mathsf{T}} + \Sigma_{\sigma(t)}^{2} + \frac{1}{N}\sum_{i=1}^{N}H^{v}\left(\tilde{Z}_{\sigma(t)}^{(i)}\right),$$
(5.9)

where a finite time integration scheme is also applied with $\sigma(t) = n\tau$ for $t \in [t_n, t_{n+1}]$. Above, we neglect the last relaxation term with ϵ since it will automatically vanish with the resulting consistency. For clarity, we show below again the continuous forecast model from (2.7)

$$\frac{\mathrm{d}\bar{u}_t}{\mathrm{d}t} = M\left(\bar{u}_t\right) + F_t + \tilde{\mathbb{E}}H^m\left(\tilde{Z}_t\right),
\frac{\mathrm{d}R_t}{\mathrm{d}t} = L\left(\bar{u}_t\right)R_t + R_tL\left(\bar{u}_t\right)^{\mathsf{T}} + \Sigma_t^2 + \tilde{\mathbb{E}}H^v\left(\tilde{Z}_t\right).$$
(5.10)

Notice that \bar{u}_t^N, R_t^N and \bar{u}_t, R_t are stochastic processes due to the random samples $\{\tilde{Z}_t^{(i)}\}\$ and \tilde{Z}_t is still dependent on the observations \mathcal{G}_t . We have the following convergence result for the finite ensemble N and finite time step τ approximation to the continuous model.

Proposition 12. If Assumption 11 is satisfied, under the same initial condition the statistical solution of the ensemble approximation model (5.9) with discrete time step τ converges to that of the continuous model (5.10) with

$$\mathbb{E}\left[\sup_{n\tau \leq T} \left| \bar{u}_{t_{n}}^{N} - \bar{u}_{t_{n}} \right|^{2} \right] \leq \left(C_{1,T}\tau + \frac{C_{2,T}}{N} \right) \|H^{m}\|_{\infty}, \\
\mathbb{E}\left[\sup_{n\tau \leq T} \left| R_{t_{n}}^{N} - R_{t_{n}} \right|^{2} \right] \leq \left(C_{1,T}'\tau + \frac{C_{2,T}'}{N} \right) \left(\|H^{m}\|_{\infty} + \|H^{v}\|_{\infty} \right).$$
(5.11)

Proof. First, consider the mean equations from the same initial state, we have

$$\begin{split} \bar{u}_t^{N,\tau} - \bar{u}_t &= \int_0^t \left[M\left(\bar{u}_{\sigma(s)}^{N,\tau}\right) - M\left(\bar{u}_s\right) \right] \mathrm{d}s + \int_0^t \left[F_{\sigma(s)} - F_s \right] \mathrm{d}s \\ &+ \int_0^t \left[\left\langle H^m, \tilde{\rho}_{\sigma(s)}^N \right\rangle - \left\langle H^m, \tilde{\rho}_s \right\rangle \right] \mathrm{d}s. \end{split}$$

Therefore, using Hölder's inequality and Lipschitz condition for M there is

$$\mathbb{E}\sup_{t\leq T} \left| \bar{u}_t^{N,\tau} - \bar{u}_t \right|^2 \leq 3T\beta \mathbb{E} \int_0^T \left| \bar{u}_{\sigma(s)}^{N,\tau} - \bar{u}_s \right|^2 \mathrm{d}s \\
+ 3T\mathbb{E} \int_0^T \left| \left\langle H^m, \tilde{\rho}_{\sigma(s)}^N \right\rangle - \left\langle H^m, \tilde{\rho}_s \right\rangle \right|^2 \mathrm{d}s + C_T \tau \\
\leq C_1 \int_0^T \mathbb{E}\sup_{s'\leq s} \left| \bar{u}_{s'}^{N,\tau} - \bar{u}_{s'} \right|^2 \mathrm{d}s + C_2 \tau \left\| H^m \right\|_{\infty}^2 + \frac{C_3}{N} \left\| H_m \right\|_{\infty}^2.$$
(5.12)

In the first term above on the right hand side, we follow the same procedure to estimate the error in the corresponding continuous solution $\bar{u}_s^{N,\tau}$ compared to the time discretization solution $\bar{u}_{\sigma(s)}^{N,\tau}$

$$\left| \bar{u}_{\sigma(s)}^{N,\tau} - \bar{u}_s \right|^2 \le \left| \bar{u}_{\sigma(s)}^{N,\tau} - \bar{u}_s^{N,\tau} \right|^2 + \left| \bar{u}_s^{N,\tau} - \bar{u}_s \right|^2 \le C\tau \left\| H^m \right\|_{\infty}^2 + \left| \bar{u}_s^{N,\tau} - \bar{u}_s \right|^2.$$

And in the second line for the term related to H^m , combining the discrete time estimate with (5.8) gives

$$\mathbb{E}\left[\sup_{t\leq T}\left|\left\langle H^{m},\tilde{\rho}_{\sigma(s)}^{N}\right\rangle-\left\langle H^{m},\tilde{\rho}_{t}\right\rangle\right|^{2}\right]\leq\left(C\tau^{2}+\frac{C_{T}}{N}\right)\left\|H^{m}\right\|_{\infty}^{2}$$

Therefore, applying Grönwall's inequality to (5.12), we get the mean estimate in (5.11).

Next, under a similar fashion, we can compute from the covariance equation and using the Lipschitz condition for L

$$\begin{split} \mathbb{E} \sup_{t \leq T} \left| R_t^{N,\tau} - R_t \right|^2 &\leq C_1 \beta \mathbb{E} \sup_{t \leq T} \left\| \bar{u}_t \right\|_{\infty}^2 \int_0^T \left| R_{\sigma(s)}^{N,\tau} - R_s \right|^2 \mathrm{d}s \\ &+ C_2 \beta \mathbb{E} \sup_{t \leq T} \left| \bar{u}_{\sigma(t)}^{N,\tau} - \bar{u}_t \right|^2 \sup_{t \leq T} \left\| R_{\sigma(t)} \right\|^2 \\ &+ C_3 \mathbb{E} \int_0^T \left| \left\langle H^v, \tilde{\rho}_{\sigma(s)}^N \right\rangle - \left\langle H^v, \tilde{\rho}_s \right\rangle \right|^2 \mathrm{d}s + C_T \tau. \end{split}$$

Using the uniform boundedness of \bar{u}_t , R_t and (5.8) for H^v together with the previous estimate for $\mathbb{E}\sup_{t\leq T} \left| \bar{u}_{\sigma(t)}^{N,\tau} - \bar{u}_t \right|^2$, we reach the final covariance estimate in (5.11).

Theorem 12 shows that the discrete approximation scheme of the approximating filter model can recover the key model statistics in mean and covariance. It again demonstrates the central role in achieving accurate prediction of the nonlinear observation functions H^m, H^v in the filtering method.

6 Summarizing discussions

We developed a systematic statistical filtering strategy that enables effective ensemble approximation of non-Gaussian probability distributions of multiscale turbulent states using observations in the leading-order moments of the mean and covariance. The filtering model is based on a closed set of coupled stochastic-statistical equations established for modeling general turbulent dynamical systems involving nonlinear coupling. The non-Gaussian features then can be characterized by a McKean-Vlasov SDE taking into account both the stochastic forecast equation and corrections according to the observed first two moments. Importantly, the proposed McKean-Vlasov SDE for the finite-dimensional stochastic state does not require the explicit computation of the infinite-dimensional probability distribution, but just relies on the feedbacks from the leading moments that can be computed from the associated statistical equations. This leads to straightforward numerical algorithms using ensemble approximations. The stochastic-statistical formulation offers a flexible approach for recovering essential model statistics, making it applicable to a wide range of problems in uncertainty quantification and data assimilation. In the immediate applications of this research, the performance of the new filtering strategy will be tested on a series of turbulent systems, starting from prototype models to realistic applications in really high-dimensional systems. Also, currently, we are only able to show the statistical consistency of the approximating filter in the first two moments of the observation function under restricted conditions. Further explorations exploiting specific model structures such as the conservation properties will be used to provide a thorough understanding of the approximation skill of the filter predictions.

Acknowledge

The research of J.-G. L. is partially supported by the National Science Foundation (NSF) under Grant No. DMS-2106988. The research of D. Q. is partially supported by Office of Naval Research (ONR) Grant No. N00014-24-1-2192.

A General background about filtering

Here, we summarize the useful results needed in the main text of this paper following mostly [2, 36, 20].

A.1 Filtering equations for general stochastic systems

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be the complete probability space. The signal process $u_t \in \mathbb{R}^d$ and the observation process $y_t \in \mathbb{R}^p$ are defined on the probability space satisfying the following SDEs

$$du_t = F(u_t) dt + \Sigma dW_t, \quad u_{t=0} = u_0,$$
(A1a)

$$dy_t = H(u_t) dt + dB_t, \quad y_{t=0} = y_0,$$
 (A1b)

where $F : \mathbb{R}^d \to \mathbb{R}^d$ and $H : \mathbb{R}^d \to \mathbb{R}^p$ are bounded and globally Lipschitz continuous functions, and $W_t \in \mathbb{R}^s, B_t \in \mathbb{R}^p$ are independent standard Wiener processes with matrix coefficient $\Sigma \in \mathbb{R}^{d \times s}$. The aim of the general filtering problem is to determine the conditional probability distribution μ_t of the signal process, u_t , given the accumulated observation process, $Y_t = \{y_s, s \leq t\}$.

Define the observation filtration $\mathcal{G}_t = \sigma \{y_s, s \leq t\}$. The random conditional distribution $\mu_t : \mathbb{R}^d \times \Omega \to [0, 1]$ is defined as the $\mathcal{P}(\mathbb{R}^d)$ -valued stochastic process which is measurable w.r.t. \mathcal{G}_t , so that for any function $\varphi \in C_b^2(\mathbb{R}^d)$ a.s.

$$\mathbb{E}\left[\varphi\left(u_{t}\right)\mid\mathcal{G}_{t}\right]=\mu_{t}\left(\varphi\right)\coloneqq\int_{\mathbb{R}^{d}}\varphi\left(u\right)\mu_{t}\left(\mathrm{d}u\right).$$

In particular, the optimal filter solution, $\hat{u}_t = \mathbb{E}[u_t | \mathcal{G}_t]$, can be defined based on μ_t . It shows that \hat{u}_t is the minimizer $\mathbb{E}\left[|\hat{u}_t - u_t|^2\right] = \min_v \mathbb{E}\left[|v - u_t|^2\right]$ among all $v \in L^2(\Omega, \mathcal{G}_t, \mathbb{P})$ in the set of \mathcal{G}_t -measurable square-integrable random variables for any fixed t. The filtering equation for the conditional probability distribution μ_t is verified to satisfy the Kushner-Stratonovich equation

$$d\mu_t \left(\varphi\right) = \mu_t \left(\mathcal{L}\varphi\right) dt + \sigma \left(H, \varphi; \mu_t\right) d\nu_t.$$
(A2)

On the right hand side of the above equation, the first term is the drift due to the infinitesimal generator $\mathcal{L} = F \cdot \nabla + \frac{1}{2}\Sigma\Sigma^{\intercal}$: $\nabla\nabla$ of the signal process (A1a); the second term represents the correction from the observation process (A1b). The innovation process, $d\nu_t = dy_t - \mu_t (H) dt \in \mathbb{R}^p$, is a \mathcal{G}_t -Brownian motion under the probability measure \mathbb{P} , and $\sigma (H, \varphi; \mu_t) = \mu_t (\varphi H^{\intercal}) - \mu_t (\varphi) \mu_t (H^{\intercal}) \in \mathbb{R}^{1 \times p}$ gives the coefficient with finite quadratic variation, where $\mu_t (H)$ is the componentwise measure of the vector-valued function H. In addition, the filtering equation (A2) is shown to have a unique solution under proper conditions (Theorem 3.30 and 4.19 in [2] and Theorem 7.7 in [56]) that is also stable (Theorem 2.7 in [6]). Therefore, this guarantees that the solution of the Kushner-Stratonovich equation (A2) uniquely characterizes the filter distribution μ_t as a $\mathcal{P} (\mathbb{R}^d)$ -valued stochastic process.

Next, assume that the conditional probability μ_t possesses a square integrable density, $\mu_t (dx) = \rho_t (x) dx$, with respect to the Lebesgue measure. It can be shown under proper conditions (Corollary 7.18 in [2] and [31]), the conditional probability solution μ_t of (A1) has a probability density $\rho_t \in W_k^2(\mathbb{R}^d)$. According to (A2) for the conditional probability, ρ_t can be found to be the unique solution to the following SPDE

$$\partial_t \rho_t = \mathcal{L}^* \rho_t \mathrm{d}t + \rho_t \left(H - \bar{H}_t \right)^\mathsf{T} \left(\mathrm{d}y_t - \bar{H}_t \mathrm{d}t \right), \quad \rho_{t=0} = \rho_0, \tag{A3}$$

where $\bar{H}_t = \int_{\mathbb{R}^d} H(x) \rho_t(x) dx$ and $\rho_0 \in L^2(\mathbb{R}^d)$ is the absolute continuous density of μ_0 . The randomness of the above SPDE only comes from the innovation process $d\nu_t = dy_t - \mu_t(H) dt$ as a finite-dimensional white noise in time.

At last, if the functions on the right hand sides of (A1) satisfy the Ornstein-Uhlenbeck processes with matrix coefficients, that is, $F(u) = Fu + f_t$ and $H(u) = Hu + h_t$. With Gaussian initial condition, $u_0 \sim \mathcal{N}(\hat{u}_0, C_0)$, the conditional distribution $\mu_t = \mathcal{N}(\hat{u}_t, C_t)$ given \mathcal{G}_t in (A2) becomes a multivariate normal distribution, where $\hat{u}_t = \mathbb{E}[u_t \mid \mathcal{G}_t]$ and $C_t = \mathbb{E}[(u_t - \hat{u}_t)(u_t - \hat{u}_t)^{\mathsf{T}} \mid \mathcal{G}_t]$. The filtering equations for $\hat{u}_t \in \mathbb{R}^d$ and $C_t \in \mathbb{R}^{d \times d}$ are given by the Kalman-Bucy filter [28]

$$d\hat{u}_t = (F\hat{u}_t + f_t) dt + K_t [dy_t - (H\hat{u}_t + h_t) dt], \qquad (A4a)$$

$$\dot{C}_t = FC_t + C_t F^{\mathsf{T}} - K_t K_t^{\mathsf{T}} + \Sigma \Sigma^{\mathsf{T}}.$$
(A4b)

with the Kalman gain matrix $K_t = C_t H^{\intercal}$. Above, (A4a) is an SDE coupled with the deterministic Riccati equation (A4b).

A.2 Infinite dimensional filtering in Hilbert space

It is shown that the linear Kalman-Bucy filter can be generalized to linear stochastic equations on a Hilbert space [10, 20]. Let H be a Hilbert space. Denote $L^2(\Omega, \mathcal{G}, \mathbb{P}; H)$ as the collection of all H-valued \mathcal{G} -measurable square-integrable random variables. The expectation of $u \in L^2(\Omega, \mathcal{G}, \mathbb{P}; H)$ is denoted by

$$\mathbb{E}\left[u\right] = \int_{\Omega} u\left(\omega\right) d\mathbb{P}\left(\omega\right). \tag{A5}$$

The inner produce for $u, v \in L^2(\Omega, \mathcal{G}, \mathbb{P}; H)$ can be defined as $\langle u, v \rangle_2 = \mathbb{E}[\langle u, v \rangle_H] = \int_{\Omega} \langle u(\omega), v(\omega) \rangle_H d\mathbb{P}(\omega)$. With the above notations, the *covariance operator* \mathcal{C} can be introduced as an element in the linear transformations $\mathcal{L}(H, H)$.

Definition 13. Let $u, v \in L^2(\Omega, \mathcal{G}, \mathbb{P}; H)$ be two *H*-valued random variables. Then the covariance of u and v is given by

$$\mathcal{C}(u,v) = \mathbb{E}\left[u \otimes v\right] - \mathbb{E}\left[u\right] \otimes \mathbb{E}\left[v\right],\tag{A6}$$

where $u(\omega) \otimes v(\omega) \in \mathcal{L}(H, H)$ is a linear transformation of H into H defined for any $f \in H$ as

$$(u \otimes v) f = u \langle v, f \rangle_H$$

It is easy to check that the adjoint $\mathcal{C}(u, v)^* = \mathcal{C}(v, u)$ and $\mathcal{C}(u, u)^* = \mathcal{C}(u, u)$ is self-adjoint since

$$\langle f, (u \otimes v) g \rangle_H = \langle u, f \rangle_H \langle v, g \rangle_H$$

Notice that if $H = \mathbb{R}^d$ is finite-dimensional, for any $x, y \in L^2(\Omega, \mathcal{G}, \mathbb{P}; \mathbb{R}^d)$, $x \otimes y = xy^{\intercal} \in \mathbb{R}^{d \times d}$, then the covariance $\mathcal{C}(x, y) \in \mathbb{R}^{d \times d}$ becomes the $d \times d$ matrix

$$\mathcal{C}(x,y) = \mathbb{E}[xy^{\mathsf{T}}] - \mathbb{E}[x] \mathbb{E}[y]^{\mathsf{T}} = \mathbb{E}[(x - \mathbb{E}[x])(y - \mathbb{E}[y])^{\mathsf{T}}].$$

Then, we call the $u_t(\omega)$ from $[0,T] \times \Omega$ to H an H-valued stochastic process. An infinite-dimensional H-valued Wiener process W_t can be defined accordingly and the Itô integral can be generalized to infinite-dimensional Hilbert space accordingly (see Chapter 2 of [44] with precise validations). Therefore, the signal process u_t of filtering can be given by the following H-valued SDE

$$du_t = \mathcal{A}_t u_t dt + \mathcal{Q}_t dW_t, \quad u_{t=0} = u_0, \tag{A7}$$

where $\mathcal{A}_t \in L^{\infty}([0,T]; \mathcal{L}(H,H))$ is a regulated mapping of [0,T] into $\mathcal{L}(H,H)$ (which is further generalized to unbounded operators in [11]) and $\mathcal{Q}_t \in L^2([0,T]; \mathcal{L}(H,H))$, and W_t is the *H*-valued Wiener process. It can be shown (Theorem 2.13 in [44] and Theorem 5.1 in [20]) that (A7) has unique solution in $L^2(\Omega, \mathcal{G}, \mathbb{P}; H)$ with initial value $\mathbb{E}\left[|u_0|_H^2\right] < \infty$. The observation stochastic process $y_t \in \mathbb{R}^p$ can be generated by the SDE

$$dy_t = \mathcal{H}_t u_t dt + dB_t, \quad y_{t=0} = y_0, \tag{A8}$$

where \mathcal{H}_t is a linear mapping of [0, T] into $\mathcal{L}(H, \mathbb{R}^p)$, and B_t is the Wiener process in \mathbb{R}^p independent of W_t . The infinite-dimensional filtering problem can be then described as: given $y_s, s \leq t$, determine the optimal estimate \hat{u}_t of u_t that minimizes

$$\mathbb{E}\left[\left|u_{t}-v\right|_{H}^{2}\right], \quad v \in L^{2}\left(\Omega, \mathcal{G}_{t}, \mathbb{P}; H\right),$$
(A9)

where $\mathcal{G}_t = \sigma \{y_s, s \leq t\}$ is generated by the observations up to time t.

Finally, in parallel to the Kalman-Bucy filter (A4) in finite-dimensional space, similar result can be extended to the above infinite-dimensional filtering problem (A7) and (A8). Below, we summarize the main results in Theorem 7.10, 7.14 of [20].

Theorem 14. The optimal filter solution \hat{u}_t of (A9) exists and is unique, which satisfies the following infinitedimensional SDE

$$d\hat{u}_t = \mathcal{A}_t \hat{u}_t dt + \mathcal{K}_t \left(dy_t - \mathcal{H}_t \hat{u}_t dt \right), \quad \hat{u}_{t=0} = u_0, \tag{A10}$$

where $\mathcal{K}_t = \mathcal{C}_t \mathcal{H}_t^* \in \mathcal{L}(\mathbb{R}^p, H)$ with \mathcal{H}_t^* the adjoint of \mathcal{H}_t . And the covariance operator $\mathcal{C}_t = \mathbb{E}\left[(u_t - \hat{u}_t) \otimes (u_t - \hat{u}_t)\right]$ satisfies the following Riccati equation

$$\mathcal{C}_t = \mathcal{A}_t \mathcal{C}_t + \mathcal{C}_t \mathcal{A}_t^* - \mathcal{K}_t \mathcal{K}_t^* + \mathcal{Q}_t \mathcal{Q}_t^*, \quad \mathcal{C}_{t=0} = \mathcal{C} (u_0, u_0).$$
(A11)

A.3 Kalman-Bucy filter with conditional Gaussian processes

The linear Kalman-Bucy filter can be generalized to nonlinear filtering accepting the conditional Gaussian processes [36]. The conditional Gaussian process $(v_t, y_t), 0 \le t \le T$ is given by the solution of the following coupled equations

$$dv_t = [F_t(y_t) v_t + f_t(y_t)] dt + \Sigma_t dW_t, \quad v_{t=0} = v_0,$$
(A12a)

$$dy_{t} = [H_{t}(y_{t}) v_{t} + h_{t}(y_{t})] dt + \Gamma_{t} dB_{t}, \quad y_{t=0} = y_{0},$$
(A12b)

where W_t, B_t are mutually independent standard Gaussian white noise processes, and the initial states (v_0, y_0) are random variables independent of W_t, B_t . In general, $v_t \in \mathbb{R}^d$ represents the signal process and $y_t \in \mathbb{R}^p$ represents the observation process. The functions f_t, F_t and h_t, H_t are globally Lipschitz continuous and uniformly bounded on the observed state y_t over the time interval $0 \le t \le T$. Assume that the sequence (v_t, y_t) is obtained from a realization ω and the initial condition, $\{v_0, y_0\}$. We can then define the observation sequence $Y_t = \{y_s(\omega), s \le t\}$ as well as the unobserved target signal $v_t = v_t(\omega)$. The above system (A12) is called the conditional Gaussian process since the conditional distribution $\mu_t = \mathbb{P}(v_t \in \cdot | Y_t)$ given Y_t becomes a Gaussian distribution a.s. if $\mu_0 = \mathbb{P}(v_0 \in \cdot | y_0)$ is Gaussian (Theorem 12.6 of [36]).

Next, let $\mathcal{G}_t = \sigma \{y_s, s \leq t\}$, and define the mean $\hat{v}_t = \mathbb{E}[v_t | \mathcal{G}_t]$ and covariance $\hat{C}_t = \mathbb{E}[(v_t - \hat{v}_t)(v_t - \hat{v}_t)^{\mathsf{T}} | \mathcal{G}_t]$ w.r.t. the conditional Gaussian distribution $\mu_t = \mathcal{N}(\hat{v}_t, \hat{C}_t)$. Then, it shows that the explicit dynamical equations for (\hat{v}_t, \hat{C}_t) can be derived based on the conditional Gaussian process (A12). As a result, filtering equations from the linear Kalman-Bucy filter (A4) can be directly applied to the conditional linear system regardless of its essentially nonlinear dynamics. The equations for the model coefficients in Chapter 12 of [36]. We summarize the results according to Theorem 12.7 of [36] for the nonlinear conditional Gaussian filter.

Theorem 15. The conditional distribution μ_t of the stochastic processes v_t given Y_t from (A12) is Gaussian, $\mathcal{N}\left(\hat{v}_t, \hat{C}_t\right)$. Then, with $\Gamma_t \Gamma_t^{\mathsf{T}} \succ 0$ and the initial mean and covariance \hat{v}_0, \hat{C}_0 , the solutions for the mean \hat{v}_t and covariance matrix \hat{C}_t are uniquely given by the following closed equations

$$d\hat{v}_{t} = [F_{t}(y_{t})\hat{v}_{t} + f_{t}(y_{t})]dt$$

$$+ K_{t}(y_{t})(\Gamma_{t}\Gamma_{t}^{\mathsf{T}})^{-1} \{dy_{t} - [H_{t}(y_{t})\hat{v}_{t} + h_{t}(y_{t})]dt\}, \qquad (A13a)$$

$$d\hat{C}_{t} = \left[F_{t}(y_{t})\hat{C}_{t} + \hat{C}_{t}F_{t}(y_{t})^{\mathsf{T}} + \Sigma_{t}\Sigma_{t}^{\mathsf{T}}\right]dt$$

$$- K_{t}(y_{t})(\Gamma_{t}\Gamma_{t}^{\mathsf{T}})^{-1}K_{t}^{\mathsf{T}}(y_{t})dt, \qquad (A13b)$$

where $K_t(y_t) = \hat{C}_t(y_t) H_t(y_t)^{\mathsf{T}}$. The matrix \hat{C}_t will remain positive-definite for all $0 \le t \le T$ if $\hat{C}_0 \succ 0$.

B Detailed proofs of theorems

Proof of Lemma 1. First, consider the model (2.7) without the relaxation term in R_t , that is, set $\epsilon^{-1} = 0$. Applying Itô's formula for Z_t with any test function $\varphi \in C_b^2(\mathbb{R}^d)$ gives

$$d\varphi(Z_t) = \mathcal{L}(\bar{u}_t, R_t) \varphi(Z_t) dt + \nabla \varphi(Z_t)^{\mathsf{T}} \Sigma_t dW_t$$
$$= \nabla \varphi(Z_t)^{\mathsf{T}} [L(\bar{u}_t) Z_t + \Gamma(Z_t Z_t^{\mathsf{T}} - R_t)] dt$$
(B1)

$$+\frac{1}{2}\Sigma_t \Sigma_t^{\mathsf{T}} : \nabla \nabla \varphi \left(Z_t \right) \mathrm{d}t + \nabla \varphi \left(Z_t \right)^{\mathsf{T}} \Sigma_t \mathrm{d}W_t, \tag{B2}$$

where \mathcal{L} is the generator of Z_t . Given any statistical solution (\bar{u}_t, R_t) and taking expectation using $\varphi(Z) = Z$, the equation for the first moment of Z_t can be found as

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathbb{E}\left[Z_{t}\right] = \left[L\left(\bar{u}_{t}\right)\mathbb{E}\left[Z_{t}\right] + \Gamma\left(\mathbb{E}\left[Z_{t}Z_{t}^{\mathsf{T}}\right] - R_{t}\right)\right].$$
(B3)

Next by taking $\varphi(Z) = Z_k Z_l$, we have

$$d(Z_{k,t}Z_{l,t}) = \sum_{m} [L_{km}(\bar{u}_{t}) Z_{m,t}Z_{l,t} + Z_{k,t}Z_{m,t}L_{lm}(\bar{u}_{t})] dt + \Sigma_{km,t}\Sigma_{lm,t}dt + \sum_{m,n} \gamma_{mnk} (Z_{m,t}Z_{n,t}Z_{l,t} - R_{mn,t}Z_{l,t}) dt + \gamma_{mnl} (Z_{m,t}Z_{n,t}Z_{k,t} - R_{mn,t}Z_{k,t}) dt + \sum_{m} \Sigma_{km,t}Z_{l,t}dW_{m,t} + \Sigma_{lm,t}Z_{k,t}dW_{m,t}.$$

This implies the second moment equation of Z_t as

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathbb{E}\left[Z_{t}Z_{t}^{\mathsf{T}}\right] = \sum_{m}\left[L_{km}\left(\bar{u}_{t}\right)\mathbb{E}\left[Z_{m,t}Z_{l,t}\right] + \mathbb{E}\left[Z_{k,t}Z_{m,t}\right]L_{lm}\left(\bar{u}_{t}\right)\right]\mathrm{d}t + \Sigma_{t}\Sigma_{t}^{\mathsf{T}} + \sum_{m,n}\gamma_{mnk}\left(\mathbb{E}\left[Z_{m,t}Z_{n,t}Z_{l,t}\right] - R_{mn,t}\mathbb{E}\left[Z_{l,t}\right]\right)\mathrm{d}t + \gamma_{mnl}\left(\mathbb{E}\left[Z_{m,t}Z_{n,t}Z_{k,t}\right] - R_{mn,t}\mathbb{E}\left[Z_{k,t}\right]\right)\mathrm{d}t.$$
(B4)

Assuming $\mathbb{E}[Z_t] = 0$ and $\mathbb{E}[Z_t Z_t^{\mathsf{T}}] = R_t$ for any time instant t, first we see that the right hand side of (B3) will always stay zero. Then, with the same statistics in the third moments of Z_t , the right hand side of (B4) becomes equal to the right hand of the statistical equation of R_t in (2.7) with (2.8). Uniqueness of solution in the statistical equations given the same initial values implies that the leading two moments of Z_t satisfy $\mathbb{E}[Z_t] = 0$ and $\mathbb{E}[Z_t Z_t^{\mathsf{T}}] = R_t$ for all t > 0.

Finally, by adding the additional relaxation term, $\epsilon^{-1} (\mathbb{E}[Z_t Z_t^{\mathsf{T}}] - R_t)$ in the dynamics of R_t , the statistical consistency condition (2.9) guarantees $R_t = \mathbb{E}[Z_t Z_t^{\mathsf{T}}]$ for all the time. Thus this term gives zero contribution, so the model (2.7) will also produce the same statistical solution.

Proof of Proposition 2. First, consider $\mathbb{E}\varphi(u_t)$ with any test function $\varphi \in C_b^2(\mathbb{R}^d)$ w.r.t. the PDF p_t for the state of the original system (1.1). Itô's lemma shows that

$$\frac{\mathrm{d}\mathbb{E}\varphi\left(u_{t}\right)}{\mathrm{d}t} = \mathbb{E}\left[\left(\Lambda u_{t} + B\left(u_{t}, u_{t}\right) + F_{t}\right) \cdot \nabla\varphi\left(u_{t}\right)\right] + \frac{1}{2}\mathbb{E}\left[\sigma_{t}\sigma_{t}^{\mathsf{T}}: \nabla\nabla\varphi\left(u_{t}\right)\right],$$

where $A: \nabla \nabla f = \sum_{mn} a_{mn} \partial_{u_m} f$. By taking $\varphi = u$ and introducing the decomposition $u' = u - \mathbb{E}u = \sum_k u'_{k,t} \hat{v}_k$ with $u'_{k,t} = \hat{v}_k \cdot u'_t$, we have

$$\frac{\mathrm{d}\mathbb{E}u_t}{\mathrm{d}t} = \mathbb{E}\left[\Lambda u_t + B\left(u, u\right) + F_t\right] = \Lambda \mathbb{E}u_t + B\left(\mathbb{E}u, \mathbb{E}u\right) + \sum_{k,l} \mathbb{E}\left[u'_{k,t}u'_{l,t}\right] B\left(\hat{v}_k, \hat{v}_l\right) + F_t.$$
(B5)

Above, we use the bilinearity of the operator B and $\mathbb{E}u' = 0$, such that

$$\mathbb{E}B(u,u) = \mathbb{E}B\left(\mathbb{E}u + u', \mathbb{E}u + u'\right)$$

= $\mathbb{E}\left[B\left(\mathbb{E}u, \mathbb{E}u\right) + B\left(\mathbb{E}u, u'\right) + B\left(u', \mathbb{E}u\right) + B\left(u', u'\right)\right]$
= $B\left(\mathbb{E}u, \mathbb{E}u\right) + \mathbb{E}\sum_{k,l} B\left(u'_{k,t}\hat{v}_k, u'_{l,t}\hat{v}_l\right)$
= $B\left(\mathbb{E}u, \mathbb{E}u\right) + \sum_{k,l} \mathbb{E}\left[u'_{k,t}u'_{l,t}\right] B\left(\hat{v}_k, \hat{v}_l\right).$

Similarly, by taking $\varphi = (\hat{v}_k \cdot u'_t) (u'_t \cdot \hat{v}_l) = u'^{\mathsf{T}}_t A u'_t$ with $A = \hat{v}_l \hat{v}_k^{\mathsf{T}}$, we find

$$\frac{d}{dt}\mathbb{E}\left[u_{k,t}'u_{l,t}'\right] = \mathbb{E}\left[\left(\Lambda u_{t} + B\left(u_{t}, u_{t}\right) + F_{t}\right) \cdot \left(A + A^{\mathsf{T}}\right)u_{t}'\right] + \frac{1}{2}\sigma_{t}\sigma_{t}^{\mathsf{T}}: \left(A + A^{\mathsf{T}}\right) \\
= \mathbb{E}\left[\left(u_{t}' \cdot \hat{v}_{l}\right)\left(\hat{v}_{k}^{\mathsf{T}}\Lambda u_{t}'\right) + \left(u_{t}'^{\mathsf{T}}\Lambda^{\mathsf{T}}\hat{v}_{l}\right)\left(\hat{v}_{k} \cdot u_{t}'\right)\right] \\
+ \mathbb{E}\left[\hat{v}_{l} \cdot B\left(\mathbb{E}u, u'\right)u_{k,t}' + \hat{v}_{k} \cdot B\left(\mathbb{E}u, u'\right)u_{l,t}'\right] + \mathbb{E}\left[\hat{v}_{l} \cdot B\left(u', \mathbb{E}u\right)u_{k,t}' + \hat{v}_{k} \cdot B\left(u', \mathbb{E}u\right)u_{l,t}'\right] \\
+ \mathbb{E}\left[B\left(u_{t}', u_{t}'\right) \cdot \left(\hat{v}_{l}u_{k,t}' + \hat{v}_{k}u_{l,t}'\right)\right] + \left(\sigma_{t} \cdot \hat{v}_{k}\right)\left(\hat{v}_{l} \cdot \sigma_{t}\right) \\
= \sum_{m}\left(\hat{v}_{k}^{\mathsf{T}}\Lambda\hat{v}_{m}\right)\mathbb{E}\left[u_{m,t}'u_{l,t}'\right] + \mathbb{E}\left[u_{k,t}'u_{m,t}'\right]\left(\hat{v}_{l}^{\mathsf{T}}\Lambda^{\mathsf{T}}\hat{v}_{m}\right) + \left(\sigma_{t} \cdot \hat{v}_{k}\right)\left(\hat{v}_{l} \cdot \sigma_{t}\right) \\
+ \sum_{m}\left[\hat{v}_{k}^{\mathsf{T}}B\left(\mathbb{E}u, \hat{v}_{m}\right)\mathbb{E}\left[u_{m,t}'u_{l,t}'\right] + \mathbb{E}\left[u_{k,t}'u_{m,t}'\right]\hat{v}_{l}^{\mathsf{T}}B\left(\mathbb{E}u, \hat{v}_{m}\right)\right] \\
+ \sum_{m}\left[\hat{v}_{k}^{\mathsf{T}}B\left(\hat{v}_{m}, \mathbb{E}u\right)\mathbb{E}\left[u_{m,t}'u_{l,t}'\right] + \mathbb{E}\left[u_{k,t}'u_{m,t}'\right]\hat{v}_{l}^{\mathsf{T}}B\left(\hat{v}_{m}, \mathbb{E}u\right)\right] \\
+ \sum_{m,n}\left[\hat{v}_{k}^{\mathsf{T}}B\left(\hat{v}_{m}, \hat{v}_{n}\right)\mathbb{E}\left[u_{l,t}'u_{m,t}'u_{n,t}'\right] + \hat{v}_{l}^{\mathsf{T}}B\left(\hat{v}_{m}, \hat{v}_{n}\right)\mathbb{E}\left[u_{k,t}'u_{m,t}'u_{n,t}'\right]\right].$$

Above in the second equality, we use the projection of modes $(A + A^{\intercal}) u'_t = (\hat{v}_l \hat{v}_k^{\intercal} + \hat{v}_k \hat{v}_l^{\intercal}) u'_t = \hat{v}_l u'_{k,t} + \hat{v}_k u'_{k,t}$, and the bilinearity of the quadratic operator B; and in the third equality, we use the decomposition of fluctuation modes, $u'_t = \sum_k u'_{k,t} \hat{v}_k$. We find the coupling operator $L_{km} = (\hat{v}_k^{\intercal} \Lambda \hat{v}_m) + \hat{v}_k^{\intercal} B(\hat{v}_m, \mathbb{E}u) + \hat{v}_k^{\intercal} B(\hat{v}_m, \mathbb{E}u)$, the third-order coupling coefficients $\gamma_{mnk} = \hat{v}_k^{\intercal} B(\hat{v}_m, \hat{v}_n)$, as well as the noise term $(\sigma_t \cdot \hat{v}_k)(\hat{v}_l \cdot \sigma_t)$.

In addition, by subtracting the mean equation (B5) from the original system (1.1), we find the SDE for the stochastic state

$$\begin{aligned} \frac{\mathrm{d}u'_t}{\mathrm{d}t} &= \frac{\mathrm{d}}{\mathrm{d}t} \left(u_t - \mathbb{E}u_t \right) = \Lambda u_t + B\left(u_t, u_t \right) + \sigma_t \dot{W}_t \\ &- \Lambda \mathbb{E}u_t - B\left(\mathbb{E}u, \mathbb{E}u \right) - \sum_{m,n} \mathbb{E} \left[u'_{m,t} u'_{n,t} \right] B\left(\hat{v}_m, \hat{v}_n \right) \\ &= \sum_m u'_{m,t} \left[\Lambda \hat{v}_m + B\left(\mathbb{E}u, \hat{v}_m \right) + B\left(\hat{v}_m, \mathbb{E}u \right) \right] \\ &+ \sum_{m,n} \left[u'_{m,t} u'_{n,t} - \mathbb{E} \left[u'_{m,t} u'_{n,t} \right] \right] B\left(\hat{v}_m, \hat{v}_n \right) + \sigma_t \dot{W}_t. \end{aligned}$$

Again in the second equality, we use the spectral decomposition of the fluctuation state $u'_{k,t} = u'_t \cdot \hat{v}_k$. By projecting the state u'_t on the basis \hat{v}_k , we have

$$\frac{\mathrm{d}u'_{k,t}}{\mathrm{d}t} = \sum_{k} L_{km} \left(\mathbb{E}u_t\right) u'_{m,t} + \sum_{m,n} \gamma_{mnk} \left[u'_{m,t}u'_{n,t} - \mathbb{E}\left[u'_{m,t}u'_{n,t}\right]\right] + \hat{v}_k^{\mathsf{T}} \sigma_t \dot{W}_t, \tag{B7}$$

with the same parameters L_{km} and γ_{mnk} defined before. The generator \mathcal{L}_t^u of u_t' can be written as

$$\mathcal{L}_{t}^{u}\left(p_{t}\right) = \left[\sum_{k} L_{km}\left(\mathbb{E}u_{t}\right)u_{m,t}' + \sum_{m,n}\gamma_{mnk}\left(u_{m,t}'u_{n,t}' - \mathbb{E}\left[u_{m,t}'u_{n,t}'\right]\right)\right] \cdot \nabla_{u'} + \frac{1}{2}\Sigma_{t}\Sigma_{t}^{\mathsf{T}}:\nabla_{u'}\nabla_{u'},$$

where \mathcal{L}_t^u is dependent on p_t in computing the expectations.

Next, we consider the closure model (2.7) without the relaxation term

$$\frac{\mathrm{d}\bar{u}_{t}}{\mathrm{d}t} = \Lambda \bar{u}_{t} + B\left(\bar{u}_{t}, \bar{u}_{t}\right) + Q_{m}\left(\mathbb{E}\left[Z_{t} \otimes Z_{t}\right]\right) + F_{t},$$

$$\frac{\mathrm{d}R_{t}}{\mathrm{d}t} = L\left(\bar{u}_{t}\right)R_{t} + R_{t}L\left(\bar{u}_{t}\right) + Q_{v}\left(\mathbb{E}\left[Z_{t} \otimes Z_{t} \otimes Z_{t}\right]\right) + \Sigma_{t}\Sigma_{t}^{\mathsf{T}},$$

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathbb{E}\left[\varphi\left(Z_{t}\right)\right] = \mathbb{E}\left[\mathcal{L}_{t}\left(\bar{u}_{t}, R_{t}\right)\varphi\left(Z_{t}\right)\right].$$
(B8)

where \mathcal{L}_t is the generator from (2.15) defined from the McKean-Vlasov SDE of Z_t

$$\mathcal{L}_t\left(\bar{u}_t, R_t\right) = \left[L\left(\bar{u}_t\right) Z_t + \Gamma\left(Z_t Z_t^{\mathsf{T}} - R_t\right)\right] \cdot \nabla_z + \frac{1}{2} \Sigma_t \Sigma_t^{\mathsf{T}} : \nabla_z \nabla_z$$

The closure terms Q_m and Q_v in (2.8) have exactly the same structure as that in the original system derived in (B5) and (B6). In addition, by comparing the above SDEs (B7) and (2.5), it is realized that their generators, $\mathcal{L}_t^u(p_t)$ and $\mathcal{L}_t(\bar{u}_t, R_t)$, share the same dynamical structure with the dependence on the first two moments w.r.t. p_t and the statistical solutions \bar{u}_t, R_t in (B8). Therefore, at any time instant t if we assume consistent statistics

$$\mathbb{E}_{p_t}\left[u_t\right] = \bar{u}_t, \quad \mathbb{E}_{p_t}\left[\left(u'_t \cdot \hat{v}_k\right)\left(u'_t \cdot \hat{v}_l\right)\right] = R_{kl,t},$$

as well as

$$\mathbb{E}_{p_{t}}\left[\varphi\left(u_{t}^{\prime}\right)\right] = \mathbb{E}\left[\varphi\left(\sum_{k=1}^{d} Z_{k,t}\hat{v}_{k}\right)\right],$$

the right hand sides of the original model (B5), (B6), and (B7) become the same as that of the closure model (B8). Starting from the same initial condition with uniqueness of the solution, it directly implies that the statistical solutions of the two systems (1.1) and (B8) will remain the same during the entire time evolution. \Box

Proof of Lemma 4. We rewrite the filter model (3.18) for \tilde{Z}_t by substituting the explicit equation for the observation process, $dy_t = [\mathcal{H}\rho_t + h_t(y_t)] dt + \Gamma_t dB_t$, in (3.3)

$$d\tilde{Z}_t = a_t \left(\tilde{Z}_t\right) dt + K_t \left(\tilde{Z}_t\right) \left\{ \left[\mathcal{H}\rho_t - H\left(\tilde{Z}_t\right)\right] dt + \Gamma_t dB_t - \Gamma_t d\tilde{B}_t \right\}.$$

By applying Itô's formula on the above SDE, we have for $\varphi \in C_b^2(\mathbb{R}^d)$

$$d\varphi\left(\tilde{Z}_{t}\right) = \nabla\varphi \cdot \left[\left(a_{t} - K_{t}\left(H\left(\tilde{Z}_{t}\right) + h_{t}\left(y_{t}\right)\right)\right)\right] dt - \nabla\varphi \cdot K_{t}\Gamma_{t}d\tilde{B}_{t} + \nabla\varphi \cdot K_{t}dy_{t} + K_{t}\Gamma_{t}^{2}K_{t}^{\mathsf{T}} : \nabla\nabla\varphi dt,$$
(B9)

where we define $A: \nabla \nabla \varphi = \sum_{i,j=1}^{d} A_{ij} \partial_{z_i z_j} \varphi$ and take the convention $(\nabla f)_{ij} = \partial_{z_i} f_j$ for the gradient of vectorvalued functions $f \in C^1(\mathbb{R}^d; \mathbb{R}^p)$. Notice that above the coefficient in the last term is 1 considering the additional contributions from the independent white noise process $\Gamma_t dB_t = dy_t - [\mathcal{H}\rho_t + h_t(y_t)] dt$ in the observation process besides the original $d\tilde{B}_t$, that is,

$$\frac{1}{2}\nabla\nabla\varphi: \mathrm{d}\left\langle K\Gamma\tilde{B}, K\Gamma\tilde{B}\right\rangle_{t} + \frac{1}{2}\nabla\nabla\varphi: \mathrm{d}\left\langle K\Gamma B, K\Gamma B\right\rangle_{t} = \nabla\nabla\varphi: K_{t}\Gamma_{t}^{2}K_{t}^{\mathsf{T}}\mathrm{d}t,$$

where we denote $\langle M, N \rangle_t$ as the Meyer's process of two martingales M_t and N_t .

First, by taking $\varphi(z) = H(z)$ and taking expectation $\tilde{\mathbb{E}}$ w.r.t. $\tilde{\rho}_t$ conditional on $Y_t = \{y_s, s \leq t\} \in \mathcal{G}_t$, we have

$$d\tilde{\mathbb{E}}H\left(\tilde{Z}_{t}\right) = \tilde{\mathbb{E}}\left[\nabla H\left(\tilde{Z}_{t}\right)^{\mathsf{T}}a_{t}\right]dt - \tilde{\mathbb{E}}\left[\nabla H\left(\tilde{Z}_{t}\right)^{\mathsf{T}}K_{t}\left(\bar{H}_{t} + H_{t}' + h_{t}\left(y_{t}\right)\right)\right]dt + \tilde{\mathbb{E}}\left[\nabla H\left(\tilde{Z}_{t}\right)^{\mathsf{T}}K_{t}\right]dy_{t} + \tilde{\mathbb{E}}\left[K_{t}\Gamma_{t}^{2}K_{t}^{\mathsf{T}}:\nabla\nabla H\left(\tilde{Z}_{t}\right)\right]dt = \tilde{\mathbb{E}}\left[\nabla H\left(\tilde{Z}_{t}\right)^{\mathsf{T}}a_{t}\right]dt + \tilde{\mathbb{E}}\left[K_{t}\Gamma_{t}^{2}K_{t}^{\mathsf{T}}:\nabla\nabla H\left(\tilde{Z}_{t}\right)\right]dt - \tilde{\mathbb{E}}\left[\nabla H\left(\tilde{Z}_{t}\right)^{\mathsf{T}}K_{t}H_{t}'\right]dt$$
(B10)
$$+ \tilde{\mathbb{E}}\left[\nabla H\left(\tilde{Z}_{t}\right)^{\mathsf{T}}K_{t}\right]\left\{dy_{t} - \left[\bar{H}_{t} + h_{t}\left(y_{t}\right)\right]dt\right\}.$$

In the first line above, we split $H\left(\tilde{Z}_t\right) = \bar{H}_t + H'_t$. Notice that the observation process $y_t \in \mathcal{G}_t$ can be brought out

of the expectation $\tilde{\mathbb{E}}[\cdot] = \mathbb{E}[\cdot | \mathcal{G}_t]$. Using the first identity in (3.21) for a_t , there is

$$\begin{split} \tilde{\mathbb{E}} \left[\nabla H \left(\tilde{Z}_t \right)^{\mathsf{T}} a_t \right] &= \int \nabla H \left(z \right)^{\mathsf{T}} \left[\nabla \cdot \left(K_t \Gamma_t^2 K_t^{\mathsf{T}} \right) - K_t \Gamma_t^2 \nabla \cdot K_t^{\mathsf{T}} \right] \tilde{\rho}_t \left(z \right) \mathrm{d}z \\ &= \int \nabla H \left(z \right)^{\mathsf{T}} \left[\nabla \cdot \left(\tilde{\rho}_t K_t \Gamma_t^2 K_t^{\mathsf{T}} \right) - K_t \Gamma_t^2 K_t^{\mathsf{T}} \nabla \tilde{\rho}_t - K_t \Gamma_t^2 \nabla \cdot K_t^{\mathsf{T}} \tilde{\rho}_t \right] \mathrm{d}z \\ &= - \int \nabla \nabla H \left(z \right) : \left(K_t \Gamma_t^2 K_t^{\mathsf{T}} \right) \tilde{\rho}_t \mathrm{d}z - \int \nabla H \left(z \right)^{\mathsf{T}} K_t \Gamma_t^2 \nabla \cdot \left(\tilde{\rho}_t K_t^{\mathsf{T}} \right) \mathrm{d}z \\ &= - \tilde{\mathbb{E}} \left[\left(K_t \Gamma_t^2 K_t^{\mathsf{T}} \right) : \nabla \nabla H \left(\tilde{Z}_t \right) \right] - \int \nabla H \left(z \right)^{\mathsf{T}} K_t \Gamma_t^2 \nabla \cdot \left(\tilde{\rho}_t K_t^{\mathsf{T}} \right) \mathrm{d}z. \end{split}$$

Then using the second identity in (3.21) for K_t and denoting $H'_t = H - \bar{H}_t$, the last term above gets simplified to

$$-\int \nabla H^{\mathsf{T}} K_t \Gamma_t^2 \nabla \cdot (\tilde{\rho}_t K_t^{\mathsf{T}}) \, \mathrm{d}z = \int \nabla H_t^{\prime \mathsf{T}} K_t \tilde{\rho}_t H_t^{\prime} \mathrm{d}z = \tilde{\mathbb{E}} \left[\nabla H_t^{\prime \mathsf{T}} K_t H_t^{\prime} \right].$$

With the above identities, first line of (B10) becomes zero. Further with the second identity in (3.21) for K_t , there is

$$\tilde{\mathbb{E}}\left(K_{t}^{\mathsf{T}}\nabla\psi\right) = \Gamma_{t}^{-2}\tilde{\mathbb{E}}\left[\left(H\left(\tilde{Z}_{t}\right) - \mathbb{E}H\right)\psi\left(\tilde{Z}_{t}\right)^{\mathsf{T}}\right],$$

for any regular function ψ with $\mathbb{E}\psi = 0$. By taking $\psi = H - \bar{H}_t$, there is

$$\tilde{\mathbb{E}}\left[\nabla H\left(\tilde{Z}_{t}\right)^{\mathsf{T}}K_{t}\right] = \tilde{\mathbb{E}}\left[\left(H\left(\tilde{Z}_{t}\right) - \bar{H}_{t}\right)\left(H\left(\tilde{Z}_{t}\right) - \bar{H}_{t}\right)^{\mathsf{T}}\right]\Gamma^{-2} = C_{t}^{H}\Gamma_{t}^{-2}.$$

This gives the equation for $\bar{H}_t = \tilde{\mathbb{E}}H\left(\tilde{Z}_t\right)$.

Next, we take $\varphi(z) = H_k(z) H_l(z)$. For the convenience of computation, we separate the mean state \bar{H}_t as

$$\varphi(z) = \left[\bar{H}_{k,t} + H'_{k}(z)\right] \left[\bar{H}_{l,t} + H'_{l}(z)\right] = H'_{k,t}(z) H'_{l,t}(z) + \bar{H}_{k,t}H'_{l}(z) + H'_{k}(z)\bar{H}_{l,t} + \bar{H}_{k,t}\bar{H}_{l,t}.$$

The last term above is independent of z, thus will vanish after applying Itô's formula (B9). We have for the first term on the right hand side

$$d\tilde{\mathbb{E}}\left[H'_{k,t}\left(\tilde{Z}_{t}\right)H'_{l,t}\left(\tilde{Z}_{t}\right)\right] = \tilde{\mathbb{E}}\nabla\left(H'_{k,t}H'_{l,t}\right)^{\mathsf{T}}\left[\left(a_{t}-K_{t}\left(\bar{H}_{t}+H'_{t}+h_{t}\left(y_{t}\right)\right)\right)\right]dt \\ + \tilde{\mathbb{E}}\left[\nabla\left(H'_{k,t}H'_{l,t}\right)^{\mathsf{T}}K_{t}\right]dy_{t} + \tilde{\mathbb{E}}\left[K_{t}\Gamma_{t}^{2}K_{t}^{\mathsf{T}}:\nabla\nabla\left(H'_{k,t}H'_{l,t}\right)\right]dt \\ = \tilde{\mathbb{E}}\left[\nabla\left(H'_{k}H'_{l}\right)^{\mathsf{T}}a_{t}\right]dt + \tilde{\mathbb{E}}\left[K_{t}\Gamma_{t}^{2}K_{t}^{\mathsf{T}}:\nabla\nabla\left(H'_{k,t}H'_{l,t}\right)\right]dt \\ + \tilde{\mathbb{E}}\left[\nabla\left(H'_{k,t}H'_{l,t}\right)^{\mathsf{T}}K_{t}\right]\left\{dy_{t}-\left[\bar{H}_{t}+h_{t}\left(y_{t}\right)\right]dt\right\} - \tilde{\mathbb{E}}\left[\nabla\left(H'_{k,t}H'_{l,t}\right)^{\mathsf{T}}K_{t}H'_{t}\right].$$
(B11)

Using the identifies (3.21) for a_t and K_t , again we can find

$$\tilde{\mathbb{E}}\left[\nabla\left(H_{k,t}'H_{l,t}'\right)^{\mathsf{T}}a_{t}\right] = -\tilde{\mathbb{E}}\left[K_{t}\Gamma_{t}^{2}K_{t}^{\mathsf{T}}:\nabla\nabla\left(H_{k,t}'H_{l,t}'\right)\right] - \int\nabla\left(H_{k,t}'H_{l,t}'\right)^{\mathsf{T}}K_{t}\Gamma_{t}^{2}\nabla\cdot\left(\tilde{\rho}_{t}K_{t}^{\mathsf{T}}\right)\mathrm{d}z$$
$$= -\tilde{\mathbb{E}}\left[K_{t}\Gamma_{t}^{2}K_{t}^{\mathsf{T}}:\nabla\nabla\left(H_{k,t}'H_{l,t}'\right)\right] + \tilde{\mathbb{E}}\left[\nabla\left(H_{k,t}'H_{l,t}'\right)^{\mathsf{T}}K_{t}H_{t}'\right].$$

Therefore, we have $d\tilde{\mathbb{E}}\left[H'_{k,t}H'_{l,t}\right] = \tilde{\mathbb{E}}\left[\nabla\left(H'_{k,t}H'_{l,t}\right)^{\mathsf{T}}K_{t}\right]\left\{dy_{t} - \left[\bar{H}_{t} + h_{t}\left(y_{t}\right)\right]dt\right\}$. Further, using the identity for K_{t} , the coefficient becomes third moments of H'_{t}

$$\tilde{\mathbb{E}}\left[\nabla\left(H_{k,t}'H_{l,t}'\right)^{\mathsf{T}}K_{t}\right] = \int \nabla\left(H_{k,t}'H_{l,t}'\right)^{\mathsf{T}}K_{t}\tilde{\rho}_{t}\mathrm{d}z$$
$$= -\int\left(H_{k,t}'H_{l,t}'\right)\left[\nabla\cdot\left(\tilde{\rho}_{t}K_{t}^{\mathsf{T}}\right)\right]^{\mathsf{T}}\mathrm{d}z$$
$$= \int\left(H_{k,t}'H_{l,t}'\right)\left[\tilde{\rho}_{t}\Gamma_{t}^{-2}\left(H\left(z\right)-\bar{H}_{t}\right)\right]^{\mathsf{T}}\mathrm{d}z$$
$$= \tilde{\mathbb{E}}\left[H_{k,t}'H_{l,t}'H_{l,t}'^{\mathsf{T}}\right]\Gamma_{t}^{-2}.$$

Similarly, by repeating the same procedure for $\bar{H}_{k,t}H'_l(z)$, we have

$$d\tilde{\mathbb{E}}\left[\bar{H}_{k,t}H_{l}'\left(\tilde{Z}_{t}\right)\right] = \tilde{\mathbb{E}}\left[\nabla\left(\bar{H}_{k,t}H_{l,t}'\right)^{\mathsf{T}}K_{t}\right]\left\{dy_{t} - \left[\bar{H}_{t} + h_{t}\left(y_{t}\right)\right]dt\right\}$$
$$= \tilde{\mathbb{E}}\left[H_{l,t}'H_{t}'^{\mathsf{T}}\right]\Gamma_{t}^{-2}\left[dy_{t} - \left(\bar{H}_{t} + h_{t}\right)dt\right]\bar{H}_{k,t}.$$

And similar result can be achieved for $\tilde{\mathbb{E}}\left[H'_k\left(\tilde{Z}_t\right)\bar{H}_{l,t}\right]$. Finally, applying Itô's formula for $\bar{H}_t\bar{H}_t^{\mathsf{T}}$ where $\mathrm{d}\bar{H}_t = C_t^H\Gamma_t^{-2}\left(\mathcal{H}\rho_t - \bar{H}_t\right)\mathrm{d}t + C_t^H\Gamma_t^{-1}\mathrm{d}B_t$ as we have derived, there is,

$$d\left(\bar{H}_{t}\bar{H}_{t}^{\mathsf{T}}\right) = \left(d\bar{H}_{t}\right)\bar{H}_{t}^{\mathsf{T}} + \bar{H}_{t}\left(d\bar{H}_{t}^{\mathsf{T}}\right) + d\left\langle C^{H}\Gamma^{-1}B, C^{H}\Gamma^{-1}B\right\rangle_{t}$$
$$= C_{t}^{H}\Gamma_{t}^{-2}\left[dy_{t} - \left(\bar{H}_{t} + h_{t}\right)dt\right]\bar{H}_{t}^{\mathsf{T}}$$
$$+ \bar{H}_{t}\left[dy_{t}^{\mathsf{T}} - \left(\bar{H}_{t} + h_{t}\right)^{\mathsf{T}}dt\right]\Gamma_{t}^{-2}C_{t}^{H}$$
$$+ C_{t}^{H}\Gamma_{t}^{-2}C_{t}^{H}dt.$$

Notice again that the white noise process, $C_t^H \Gamma_t^{-1} dB_t$, gives the last term in the first equality above. Putting all the above equations together, we get the equation for $dC_t^H = d\tilde{\mathbb{E}} \left[H \left(\tilde{Z}_t \right) H \left(\tilde{Z}_t \right)^{\mathsf{T}} \right] - d \left(\bar{H}_t \bar{H}_t^{\mathsf{T}} \right)$ where $C_{kl,t}^H = d\tilde{\mathbb{E}} \left[H \left(\tilde{Z}_t \right) H \left(\tilde{Z}_t \right)^{\mathsf{T}} \right]$ $\tilde{\mathbb{E}}\left[H_{k,t}'H_{l,t}'\right].$

Proof of Proposition 10. According to (5.4) with $K = \tilde{K}\Gamma^{-2}$, we need to show

$$-\nabla \cdot \left(\tilde{K}^{\mathsf{T}} \tilde{\rho}\right) H^{\mathsf{T}} = \tilde{\rho} H' H^{\mathsf{T}} \Rightarrow \tilde{\mathbb{E}} \left[\tilde{K}^{\mathsf{T}} \nabla H\right] = \tilde{\mathbb{E}} \left[H' \left(\bar{H} + H'\right)^{\mathsf{T}}\right] = C^{H}$$

according to the specific expressions $H = H^m$ and $H = H^v$. First, we can compute

$$\nabla_z H^m_l = 2A_l z, \quad \nabla_z H^v_{pq} = 2z_q A_p z + \left(z^{\mathsf{T}} A_p z\right) \delta_{qj} \hat{e}_j.$$

Above in H^v for simplicity, we only compute half of the symmetric function and \hat{e}_j is the unit vector with value 1 in the j-th entry.

From direct computations for H^m and using $H^m_k = z^{\intercal} A_k z$, we have

$$\sum_{j} \tilde{K}_{j,k}^{m} \frac{\partial H_{l}^{m}}{\partial z_{j}} = \frac{1}{2} \left[\left(z^{\mathsf{T}} A_{k} z \right) - \bar{H}_{k}^{m} \right] \sum_{j} z_{j} 2 \left(A_{l} z \right)_{j}$$
$$= \left[\left(z^{\mathsf{T}} A_{k} z \right) - \bar{H}_{k}^{m} \right] \left(z^{\mathsf{T}} A_{l} z \right) = H_{k}^{m'} H_{l}^{m}.$$

Similarly for H^v , we can compute

$$\begin{split} \sum_{j} \tilde{K}_{j,kl}^{v} \frac{\partial H_{pq}^{v}}{\partial z_{j}} &= \frac{1}{3} \left[\left(z^{\mathsf{T}} A_{k} z \right) z_{l} - \bar{H}_{kl}^{v} \right] \sum_{j} z_{j} \left[2 z_{q} \left(A_{p} z \right)_{j} + \left(z^{\mathsf{T}} A_{p} z \right) \delta_{qj} \right] \\ &= \frac{2}{3} \left[\left(z^{\mathsf{T}} A_{k} z \right) z_{l} - \bar{H}_{kl}^{v} \right] \left(z^{\mathsf{T}} A_{p} z \right) z_{q} + \frac{1}{3} \left[\left(z^{\mathsf{T}} A_{k} z \right) z_{l} - \bar{H}_{kl}^{v} \right] z_{q} \left(z^{\mathsf{T}} A_{p} z \right) \\ &= \left[\left(z^{\mathsf{T}} A_{k} z \right) z_{l} - \bar{H}_{kl}^{v} \right] \left(z^{\mathsf{T}} A_{p} z \right) z_{q} = H_{kl}^{v\prime} H_{pq}^{v}. \end{split}$$

This finishes the proof.

Proof of Proposition 12. First, consider the mean equations from the same initial state, we have

$$\begin{split} \bar{u}_t^{N,\tau} - \bar{u}_t &= \int_0^t \left[M\left(\bar{u}_{\sigma(s)}^{N,\tau}\right) - M\left(\bar{u}_s\right) \right] \mathrm{d}s + \int_0^t \left[F_{\sigma(s)} - F_s \right] \mathrm{d}s \\ &+ \int_0^t \left[\left\langle H^m, \tilde{\rho}_{\sigma(s)}^N \right\rangle - \left\langle H^m, \tilde{\rho}_s \right\rangle \right] \mathrm{d}s. \end{split}$$

Therefore, using Hölder's inequality and Lipschitz condition for M there is

$$\mathbb{E} \sup_{t \leq T} \left| \bar{u}_t^{N,\tau} - \bar{u}_t \right|^2 \leq 3T \beta \mathbb{E} \int_0^T \left| \bar{u}_{\sigma(s)}^{N,\tau} - \bar{u}_s \right|^2 \mathrm{d}s \\
+ 3T \mathbb{E} \int_0^T \left| \left\langle H^m, \tilde{\rho}_{\sigma(s)}^N \right\rangle - \left\langle H^m, \tilde{\rho}_s \right\rangle \right|^2 \mathrm{d}s + C_T \tau \\
\leq C_1 \int_0^T \mathbb{E} \sup_{s' \leq s} \left| \bar{u}_{s'}^{N,\tau} - \bar{u}_{s'} \right|^2 \mathrm{d}s + C_2 \tau \left\| H^m \right\|_{\infty}^2 + \frac{C_3}{N} \left\| H_m \right\|_{\infty}^2.$$
(B12)

In the first term above on the right hand side, we follow the same procedure to estimate the error in the corresponding continuous solution $\bar{u}_s^{N,\tau}$ compared to the time discretization solution $\bar{u}_{\sigma(s)}^{N,\tau}$

$$\left|\bar{u}_{\sigma(s)}^{N,\tau} - \bar{u}_{s}\right|^{2} \leq \left|\bar{u}_{\sigma(s)}^{N,\tau} - \bar{u}_{s}^{N,\tau}\right|^{2} + \left|\bar{u}_{s}^{N,\tau} - \bar{u}_{s}\right|^{2} \leq C\tau \left\|H^{m}\right\|_{\infty}^{2} + \left|\bar{u}_{s}^{N,\tau} - \bar{u}_{s}\right|^{2}$$

And in the second line for the term related to H^m , combining the discrete time estimate with (5.8) gives

$$\mathbb{E}\left[\sup_{t\leq T}\left|\left\langle H^{m}, \tilde{\rho}_{\sigma(s)}^{N}\right\rangle - \left\langle H^{m}, \tilde{\rho}_{t}\right\rangle\right|^{2}\right] \leq \left(C\tau^{2} + \frac{C_{T}}{N}\right) \left\|H^{m}\right\|_{\infty}^{2}$$

Therefore, applying Grönwall's inequality to (B12), we get the mean estimate in (5.11).

Next, under a similar fashion, we can compute from the covariance equation and using the Lipschitz condition for L

$$\begin{split} \mathbb{E} \sup_{t \leq T} \left| R_t^{N,\tau} - R_t \right|^2 &\leq C_1 \beta \mathbb{E} \sup_{t \leq T} \left\| \bar{u}_t \right\|_{\infty}^2 \int_0^T \left| R_{\sigma(s)}^{N,\tau} - R_s \right|^2 \mathrm{d}s \\ &+ C_2 \beta \mathbb{E} \sup_{t \leq T} \left| \bar{u}_{\sigma(t)}^{N,\tau} - \bar{u}_t \right|^2 \sup_{t \leq T} \left\| R_{\sigma(t)} \right\|^2 \\ &+ C_3 \mathbb{E} \int_0^T \left| \left\langle H^v, \tilde{\rho}_{\sigma(s)}^N \right\rangle - \left\langle H^v, \tilde{\rho}_s \right\rangle \right|^2 \mathrm{d}s + C_T \tau. \end{split}$$

Using the uniform boundedness of \bar{u}_t , R_t and (5.8) for H^v together with the previous estimate for $\mathbb{E}\sup_{t\leq T} \left| \bar{u}_{\sigma(t)}^{N,\tau} - \bar{u}_t \right|^2$, we reach the final covariance estimate in (5.11).

References

- Eviatar Bach, Tim Colonius, Isabel Scherl, and Andrew Stuart. Filtering dynamical systems using observations of statistics. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 34(3), 2024.
- [2] Alan Bain and Dan Crisan. Fundamentals of stochastic filtering, volume 3. Springer, 2009.
- [3] Abhishek Balakrishna and Animikh Biswas. Determining map, data assimilation and an observable regularity criterion for the three-dimensional Boussinesq system. Applied Mathematics & Optimization, 86(3):28, 2022.
- [4] Kay Bergemann and Sebastian Reich. An ensemble Kalman-Bucy filter for continuous data assimilation. Meteorologische Zeitschrift, 21(3):213, 2012.
- [5] Animikh Biswas and Michal Branicki. A unified framework for the analysis of accuracy and stability of a class of approximate Gaussian filters for the Navier-Stokes equations. arXiv preprint arXiv:2402.14078, 2024.
- [6] Amarjit Budhiraja. Asymptotic stability, ergodicity and other asymptotic properties of the nonlinear filter. In Annales de l'IHP Probabilités et statistiques, volume 39, pages 919–941, 2003.
- [7] Edoardo Calvello, Sebastian Reich, and Andrew M Stuart. Ensemble Kalman methods: A mean field perspective. *Acta Numerica*, 2025.

- [8] Dan Crisan and Jie Xiong. Approximate McKean–Vlasov representations for a class of SPDEs. Stochastics An International Journal of Probability and Stochastics Processes, 82(1):53–68, 2010.
- [9] Dan Crisan and Jie Xiong. Numerical solution for a class of SPDEs over bounded domains. Stochastics An International Journal of Probability and Stochastic Processes, 86(3):450–472, 2014.
- [10] Ruth Curtain. A survey of infinite-dimensional filtering. SIAM Review, 17(3):395–411, 1975.
- [11] Ruth F Curtain. Infinite-dimensional filtering. SIAM Journal on Control, 13(1):89–104, 1975.
- [12] Jana de Wiljes and Xin T Tong. Analysis of a localised nonlinear ensemble Kalman Bucy filter with complete and accurate observations. *Nonlinearity*, 33(9):4752, 2020.
- [13] Pierre Del Moral. Nonlinear filtering: Interacting particle resolution. Comptes Rendus de l'Académie des Sciences-Series I-Mathematics, 325(6):653–658, 1997.
- [14] Patrick H Diamond, Sanae-I Itoh, and Kimitaka Itoh. Modern Plasma Physics: Volume 1, Physical Kinetics of Turbulent Plasmas. Cambridge University Press, 2010.
- [15] Arnaud Doucet, Nando De Freitas, Neil James Gordon, et al. Sequential Monte Carlo methods in practice, volume 1. Springer, 2001.
- [16] Paul Dupuis and Richard S Ellis. A weak convergence approach to the theory of large deviations. John Wiley & Sons, 2011.
- [17] Oliver G Ernst, Björn Sprungk, and Hans-Jörg Starkloff. Analysis of the ensemble and polynomial chaos Kalman filters in Bayesian inverse problems. SIAM/ASA Journal on Uncertainty Quantification, 3(1):823– 851, 2015.
- [18] Geir Evensen. Sequential data assimilation with a nonlinear quasi-geostrophic model using Monte Carlo methods to forecast error statistics. *Journal of Geophysical Research: Oceans*, 99(C5):10143–10162, 1994.
- [19] Geir Evensen. The ensemble Kalman filter for combined state and parameter estimation. IEEE Control Systems Magazine, 29(3):83–104, 2009.
- [20] Peter L Falb. Infinite-dimensional filtering: The Kalman-Bucy filter in Hilbert space. 1967.
- [21] Uriel Frisch. Turbulence: the legacy of AN Kolmogorov. Cambridge university press, 1995.
- [22] Yuan Gao and Di Qi. Mean field games for controlling coherent structures in nonlinear fluid systems. arXiv preprint arXiv:2401.10356, 2024.
- [23] Michael Ghil, S Cohn, John Tavantzis, K Bube, and Eugene Isaacson. Applications of estimation theory to numerical weather prediction. *Dynamic meteorology: Data assimilation methods*, pages 139–224, 1981.
- [24] Georg A Gottwald and Andrew J Majda. A mechanism for catastrophic filter divergence in data assimilation for sparse observation networks. Nonlinear Processes in Geophysics, 20(5):705–712, 2013.
- [25] Carl Graham, Thomas G Kurtz, Sylvie Méléard, Philip E Protter, Mario Pulvirenti, Denis Talay, and Sylvie Méléard. Asymptotic behaviour of some interacting particle systems; Mckean-Vlasov and Boltzmann models. Probabilistic Models for Nonlinear Partial Differential Equations: Lectures given at the 1st Session of the Centro Internazionale Matematico Estivo (CIME) held in Montecatini Terme, Italy, May 22–30, 1995, pages 42–95, 1996.
- [26] Peter L Houtekamer and Herschel L Mitchell. Data assimilation using an ensemble Kalman filter technique. Monthly Weather Review, 126(3):796–811, 1998.
- [27] Rentian Hu, Thomas K Edwards, Leslie M Smith, and Samuel N Stechmann. Initial investigations of precipitating quasi-geostrophic turbulence with phase changes. *Research in the Mathematical Sciences*, 8:1–25, 2021.

- [28] Rudolph E Kalman and Richard S Bucy. New results in linear filtering and prediction theory. 1961.
- [29] Eugenia Kalnay. Atmospheric modeling, data assimilation and predictability. Cambridge university press, 2003.
- [30] S Kullback and RA Leibler. The annals of mathematical statistics. On information and sufficiency, 22, 1951.
- [31] Thomas G Kurtz and Jie Xiong. Particle representations for a class of nonlinear SPDEs. Stochastic Processes and their Applications, 83(1):103–126, 1999.
- [32] Thomas G Kurtz and Jie Xiong. Numerical solutions for a class of SPDEs with application to filtering. Stochastics in Finite and Infinite Dimensions: In Honor of Gopinath Kallianpur, pages 233–258, 2001.
- [33] Kody Law, Andrew Stuart, and Kostas Zygalakis. Data assimilation. Cham, Switzerland: Springer, 214:52, 2015.
- [34] Marcel Lesieur. Turbulence in fluids: stochastic and numerical modelling, volume 488. Nijhoff Boston, MA, 1987.
- [35] Martin Leutbecher and Tim N Palmer. Ensemble forecasting. Journal of computational physics, 227(7):3515– 3539, 2008.
- [36] Robert S Liptser and Albert N Shiryaev. Statistics of random processes II: Applications, volume 6. Springer Science & Business Media, 2013.
- [37] Andrew J Majda. Introduction to turbulent dynamical systems in complex systems. Springer, 2016.
- [38] Andrew J Majda and Di Qi. Strategies for reduced-order models for predicting the statistical responses and uncertainty quantification in complex turbulent dynamical systems. SIAM Review, 60(3):491–549, 2018.
- [39] Andrew J Majda and Di Qi. Linear and nonlinear statistical response theories with prototype applications to sensitivity analysis and statistical control of complex turbulent dynamical systems. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 29(10), 2019.
- [40] Andrew J Majda and Xin T Tong. Performance of ensemble Kalman filters in large dimensions. Communications on Pure and Applied Mathematics, 71(5):892–937, 2018.
- [41] Henry P McKean. Propagation of chaos for a class of non-linear parabolic equations. Stochastic Differential Equations (Lecture Series in Differential Equations, Session 7, Catholic Univ., 1967), pages 41–57, 1967.
- [42] Sergey Nazarenko and Sergei Lukaschuk. Wave turbulence on water surface. Annual Review of Condensed Matter Physics, 7:61–88, 2016.
- [43] Karl Oelschlager. A martingale approach to the law of large numbers for weakly interacting stochastic processes. The Annals of Probability, pages 458–479, 1984.
- [44] Étienne Pardoux. Stochastic partial differential equations. Fudan lecture notes, 87, 2007.
- [45] Sahani Pathiraja, Sebastian Reich, and Wilhelm Stannat. McKean–Vlasov SDEs in nonlinear filtering. SIAM Journal on Control and Optimization, 59(6):4188–4215, 2021.
- [46] Francis J Pinski, Gideon Simpson, Andrew M Stuart, and Hendrik Weber. Kullback–Leibler approximation for probability measures on infinite dimensional spaces. SIAM Journal on Mathematical Analysis, 47(6):4091–4122, 2015.
- [47] Di Qi and Jian-Guo Liu. High-order moment closure models with random batch method for efficient computation of multiscale turbulent systems. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 33(10), 2023.
- [48] Di Qi and Jian-Guo Liu. A random batch method for efficient ensemble forecasts of multiscale turbulent systems. Chaos: An Interdisciplinary Journal of Nonlinear Science, 33(2), 2023.

- [49] Di Qi and Andrew J Majda. Rigorous statistical bounds in uncertainty quantification for one-layer turbulent geophysical flows. Journal of Nonlinear Science, 28:1709–1761, 2018.
- [50] Sebastian Reich and Colin Cotter. Probabilistic forecasting and Bayesian data assimilation. Cambridge University Press, 2015.
- [51] Rick Salmon. Lectures on geophysical fluid dynamics. Oxford University Press, USA, 1998.
- [52] Daniel Sanz-Alonso, Andrew Stuart, and Armeen Taeb. Inverse problems and data assimilation, volume 107. Cambridge University Press, 2023.
- [53] Chris Snyder, Thomas Bengtsson, Peter Bickel, and Jeff Anderson. Obstacles to high-dimensional particle filtering. *Monthly Weather Review*, 136(12):4629–4640, 2008.
- [54] Simone Carlo Surace, Anna Kutschireiter, and Jean-Pascal Pfister. How to avoid the curse of dimensionality: Scalability of particle filters with and without importance weights. SIAM review, 61(1):79–91, 2019.
- [55] Shanyin Tong, Eric Vanden-Eijnden, and Georg Stadler. Extreme event probability estimation using PDEconstrained optimization and large deviation theory, with application to tsunamis. *Communications in Applied Mathematics and Computational Science*, 16(2):181–225, 2021.
- [56] Jie Xiong. An introduction to stochastic filtering theory, volume 18. OUP Oxford, 2008.
- [57] Tao Yang, Henk AP Blom, and Prashant G Mehta. The continuous-discrete time feedback particle filter. In 2014 American control conference, pages 648–653. IEEE, 2014.
- [58] Shing-Tung Yau and Stephen S-T Yau. Finite dimensional filters with nonlinear drift, xi: Explicit solution of the generalized Kolmogorov equation in Brockett–Mitter program. Advances in Mathematics, 140(2):156–189, 1998.