

On the monoid of order-preserving transformations of a finite chain whose ranges are intervals

Vítor H. Fernandes*

July 9, 2024

Abstract

In this note we give a presentation for the monoid \mathcal{JO}_n of all order-preserving transformations of a n -chain whose ranges are intervals. We also consider the submonoid \mathcal{JO}_n^- of \mathcal{JO}_n consisting of order-decreasing transformations, for which we determine the cardinality, the rank and a presentation.

1 Introduction

Let $n \in \mathbb{N}$ and $\Omega_n = \{1, \dots, n\}$. Denote by \mathcal{T}_n the monoid (under composition) of all full transformations of Ω_n . From now on, we consider Ω_n equipped with the usual linear order $1 < \dots < n$. Recall that a transformation α of Ω_n is said to be *order-preserving* if $x \leq y$ implies $x\alpha \leq y\alpha$, for all $x, y \in \Omega_n$. Denote by \mathcal{O}_n the submonoid of \mathcal{T}_n of all order-preserving transformations.

Semigroups of order-preserving transformations have been studied extensively for over six decades. The oldest papers known to the author date back to 1962, are by Aizenstat [1, 2] and give a presentation and a description of the congruences of \mathcal{O}_n . After that and up to the present day, many other papers have been published studying various properties of \mathcal{O}_n . See, for example, [3, 4, 6, 8, 9, 10, 12, 14]. Nonetheless, these semigroups still reveal some mystery. The problem of decidability of the pseudovariety generated by the family \mathcal{O}_n , with $n \in \mathbb{N}$, proposed by J.-E. Pin in the “Szege International Semigroup Colloquium” (1987), is perhaps the oldest and most important that remains to be solved.

The main object of this work is the submonoid \mathcal{JO}_n of \mathcal{O}_n consisting of all transformations $\alpha \in \mathcal{O}_n$ such that the image of α is an interval of Ω_n , i.e. $x \leq y \leq z$ and $x, z \in \text{Im}(\alpha)$ imply $y \in \text{Im}(\alpha)$, for all $x, y, z \in \Omega_n$.

The monoid \mathcal{JO}_n is not only a very natural submonoid of \mathcal{O}_n but also coincides with the monoid of all *weak endomorphisms* of a directed path with n vertices. See [5]. In the aforementioned paper, the authors studied the regularity, determined the rank, the cardinality and the number of idempotents of \mathcal{JO}_n (denoted there by $\text{wEnd}\vec{P}_n$). In this note, our main aim is to exhibit a presentation for \mathcal{JO}_n . We also consider its submonoids \mathcal{JO}_n^- consisting of order-decreasing transformations and \mathcal{JO}_n^+ consisting of order-increasing transformations, for which we determine the cardinalities, the ranks and presentations.

Recall that the *rank* of a (finite) monoid M is the minimum size that a generating set of M can have.

Next, we also recall some notions related to the concept of a monoid presentation.

Let A be an alphabet and consider the free monoid A^* generated by A . The elements of A and of A^* are called *letters* and *words*, respectively. The empty word is denoted by 1 and we write A^+ to express $A^* \setminus \{1\}$. The number of letters of a word $w \in A^*$ is called the *length* of w and it is denoted by $|w|$. A pair (u, v) of $A^* \times A^*$ is called a *relation* of A^* and it is usually represented by $u = v$. A *monoid presentation* is an ordered

*This work is funded by national funds through the FCT - Fundação para a Ciência e a Tecnologia, I.P., under the scope of the projects UIDB/00297/2020 (<https://doi.org/10.54499/UIDB/00297/2020>) and UIDP/00297/2020 (<https://doi.org/10.54499/UIDP/00297/2020>) (Center for Mathematics and Applications).

2020 *Mathematics Subject Classification*: 20M20, 20M05, 20M10.

Keywords: order-preserving, order-decreasing, transformations, presentations.

pair $\langle A \mid R \rangle$, where $R \subseteq A^* \times A^*$ is a set of relations of the free monoid A^* . Denote by \sim_R the smallest congruence on A^* containing R . A monoid M is said to be *defined by a presentation* $\langle A \mid R \rangle$ if M is isomorphic to A^*/\sim_R . Let X be a generating set of a monoid M and let $\phi : A \rightarrow M$ be an injective mapping such that $A\phi = X$. Let $\varphi : A^* \rightarrow M$ be the (surjective) homomorphism of monoids that extends ϕ to A^* . We say that X satisfies (via φ) a relation $u = v$ of A^* if $u\varphi = v\varphi$. For more details see [13].

Remember that, a usual method to find a presentation for a finite monoid is described by the following result (adapted to the monoid case from [13, Proposition 3.2.2]).

Proposition 1.1 (Guess and Prove method). *Let M be a finite monoid generated by a set X , let A be an alphabet and let $\phi : A \rightarrow M$ be an injective mapping such that $A\phi = X$. Let $\varphi : A^* \rightarrow M$ be the (surjective) homomorphism of monoids that extends ϕ to A^* , let $R \subseteq A^* \times A^*$ and $W \subseteq A^*$. Assume that the following conditions are satisfied:*

1. *The generating set X of M satisfies (via φ) all relations from R ;*
2. *For each word $w \in X^*$, there exists a word $w' \in W$ such that $w \sim_R w'$;*
3. $|W| \leq |M|$.

Then, M is defined by the presentation $\langle A \mid R \rangle$.

Notice that, if W satisfies the above conditions then, in fact, $|W| = |M|$.

For general background on Semigroup Theory and standard notation, we refer the reader to Howie's book [11]. We would also like to mention the use of computational tools, namely GAP [7].

2 Preliminaries

We start this section by recalling the cardinality, the rank and generators of \mathcal{JO}_n that have been established in [5]. By [5, Theorem 2.6], we have $|\mathcal{JO}_n| = \sum_{k=1}^n (n-k+1) \binom{n-1}{k-1}$, whence

$$|\mathcal{JO}_n| = (n+1)2^{n-2}.$$

Next, for $n \geq 2$, define

$$a_i = \begin{pmatrix} 1 & \cdots & i & i+1 & \cdots & n \\ 1 & \cdots & i & i & \cdots & n-1 \end{pmatrix} \quad \text{and} \quad b_i = \begin{pmatrix} 1 & \cdots & i & i+1 & \cdots & n \\ 2 & \cdots & i+1 & i+1 & \cdots & n \end{pmatrix},$$

for $1 \leq i \leq n-1$. Clearly, $a_1, \dots, a_{n-1}, b_1, \dots, b_{n-1} \in \mathcal{JO}_n$. Moreover, by [5, Proposition 3.3 and Theorem 3.5], for $n \geq 3$, $\{a_1, \dots, a_{n-2}, b_{n-1}\}$ is a generating set of \mathcal{JO}_n of minimum size. In particular, for $n \geq 3$, the monoid \mathcal{JO}_n has rank $n-1$. Notice that $\mathcal{JO}_1 = \mathcal{T}_1$ and $\mathcal{JO}_2 = \left\{ \begin{pmatrix} 1 & 2 \\ 1 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 2 \\ 2 & 2 \end{pmatrix}, \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix} \right\}$ and so, clearly, \mathcal{JO}_1 has rank 0 and \mathcal{JO}_2 has rank 2.

Now, observe that, for $1 \leq i, j \leq n-1$, we have

$$a_i a_j = \begin{cases} \begin{pmatrix} 1 & \cdots & i & i+1 & \cdots & j+1 & j+2 & \cdots & n \\ 1 & \cdots & i & i & \cdots & j & j & \cdots & n-2 \end{pmatrix} & \text{if } i \leq j \\ \begin{pmatrix} 1 & \cdots & j & j+1 & \cdots & i & i+1 & \cdots & n \\ 1 & \cdots & j & j & \cdots & i-1 & i-1 & \cdots & n-2 \end{pmatrix} & \text{if } i > j, \end{cases}$$

$$a_i b_j = \begin{cases} \begin{pmatrix} 1 & \cdots & i & i+1 & \cdots & j+1 & j+2 & \cdots & n \\ 2 & \cdots & i+1 & i+1 & \cdots & j+1 & j+1 & \cdots & n-1 \end{pmatrix} & \text{if } i \leq j \\ \begin{pmatrix} 1 & \cdots & j & j+1 & \cdots & i & i+1 & \cdots & n \\ 2 & \cdots & j+1 & j+1 & \cdots & i & i & \cdots & n-1 \end{pmatrix} & \text{if } i > j, \end{cases}$$

$$b_i b_j = \begin{cases} \begin{pmatrix} 1 & \cdots & i & i+1 & \cdots & j & j+1 & \cdots & n \\ 3 & \cdots & i+2 & i+2 & \cdots & j+1 & j+1 & \cdots & n \end{pmatrix} & \text{if } i < j \\ \begin{pmatrix} 1 & \cdots & j-1 & j & \cdots & i & i+1 & \cdots & n \\ 3 & \cdots & j+1 & j+1 & \cdots & i+1 & i+1 & \cdots & n \end{pmatrix} & \text{if } i \geq j \end{cases}$$

and

$$b_i a_j = \begin{cases} \begin{pmatrix} 1 & \cdots & i & i+1 & \cdots & j & j+1 & \cdots & n \\ 2 & \cdots & i+1 & i+1 & \cdots & j & j & \cdots & n-1 \end{pmatrix} & \text{if } i < j \\ \begin{pmatrix} 1 & \cdots & j-1 & j & \cdots & i & i+1 & \cdots & n \\ 2 & \cdots & j & j & \cdots & i & i & \cdots & n-1 \end{pmatrix} & \text{if } i \geq j, \end{cases}$$

from which we can deduce the equalities

$$a_i a_{n-1} = a_i = b_i a_1 \quad \text{and} \quad b_i b_1 = b_i = a_i b_{n-1}, \quad \text{for } 1 \leq i \leq n-1, \quad (1)$$

$$a_i a_j = a_{j+1} a_i \quad \text{and} \quad b_i b_{j+1} = b_{j+1} b_{i+1}, \quad \text{for } 1 \leq i \leq j \leq n-2 \quad (2)$$

and

$$b_i a_j = a_i b_{j-1} = b_j a_{i+1} = a_j b_i, \quad \text{for } 1 \leq i < j \leq n-1. \quad (3)$$

Let $A = \{a_1, \dots, a_{n-1}\}$ and $B = \{b_1, \dots, b_{n-1}\}$. Consider the alphabet $A \cup B$ with $2n-2$ letters and the set R formed by the following $\frac{1}{2}(3n^2 - n - 2)$ monoid relations:

$$(R_1) \quad a_i a_{n-1} = a_i, \quad 1 \leq i \leq n-1;$$

$$(R_2) \quad a_i a_j = a_{j+1} a_i, \quad 1 \leq i \leq j \leq n-2;$$

$$(R_3) \quad b_i b_1 = b_i, \quad 1 \leq i \leq n-1;$$

$$(R_4) \quad b_i b_{j+1} = b_{j+1} b_{i+1}, \quad 1 \leq i \leq j \leq n-2;$$

$$(R_5) \quad b_i a_j = a_j b_i, \quad 1 \leq i < j \leq n-1;$$

$$(R_6) \quad b_i a_1 = a_i, \quad 1 \leq i \leq n-1;$$

$$(R_7) \quad a_i b_{n-1} = b_i, \quad 1 \leq i \leq n-1.$$

Our main aim in this note is to show that \mathcal{JO}_n is defined by the presentation $\langle a_1, \dots, a_{n-1}, b_1, \dots, b_{n-1} \mid R \rangle$.

Let $\varphi : (A \cup B)^* \rightarrow \mathcal{JO}_n$ be the homomorphism of monoids that extends the mapping $A \cup B \rightarrow \mathcal{JO}_n$ defined by $a_i \mapsto a_i$ and $b_i \mapsto b_i$, for $1 \leq i \leq n-1$. Therefore, taking into account the above equalities (1)-(3), we can already conclude that the generating set $A \cup B$ of \mathcal{JO}_n satisfies (via φ) all relations from R .

3 Decreasing and increasing transformations

Let us consider the following submonoids of \mathcal{T}_n :

$$\mathcal{T}_n^- = \{\alpha \in \mathcal{T}_n \mid x\alpha \leq x, \text{ for all } x \in \Omega_n\} \text{ (order-decreasing transformations of } \Omega_n)$$

and

$$\mathcal{T}_n^+ = \{\alpha \in \mathcal{T}_n \mid x \leq x\alpha, \text{ for all } x \in \Omega_n\} \text{ (order-increasing transformations of } \Omega_n).$$

Consider also the mapping $\varphi : \mathcal{T}_n \rightarrow \mathcal{T}_n$ which maps each transformation $\alpha \in \mathcal{T}_n$ into the transformation $\bar{\alpha} \in \mathcal{T}_n$ defined by $(x)\bar{\alpha} = n+1 - (n+1-x)\alpha$, for all $x \in \Omega_n$. It is a routine matter to check that φ is an automorphism of monoids such that φ^2 is the identity mapping of \mathcal{T}_n , $\mathcal{T}_n^- \varphi \subseteq \mathcal{T}_n^+$, $\mathcal{T}_n^+ \varphi \subseteq \mathcal{T}_n^-$ and $\mathcal{JO}_n \varphi \subseteq \mathcal{JO}_n$. Consequently,

$$\mathcal{T}_n^- \varphi = \mathcal{T}_n^+, \quad \mathcal{T}_n^+ \varphi = \mathcal{T}_n^- \quad \text{and} \quad \mathcal{JO}_n \varphi = \mathcal{JO}_n.$$

Now, consider the submonoids $\mathcal{JO}_n^- = \mathcal{JO}_n \cap \mathcal{T}_n^-$ and $\mathcal{JO}_n^+ = \mathcal{JO}_n \cap \mathcal{T}_n^+$ of \mathcal{JO}_n . It follows immediately from the previous properties that

$$\mathcal{JO}_n^- \varphi = \mathcal{JO}_n^+ \quad \text{and} \quad \mathcal{JO}_n^+ \varphi = \mathcal{JO}_n^-.$$

Therefore, \mathcal{JO}_n^- and \mathcal{JO}_n^+ are isomorphic submonoids of \mathcal{JO}_n .

Next, observe that $a_1, \dots, a_{n-1} \in \mathcal{JO}_n^-$ and $b_1, \dots, b_{n-1} \in \mathcal{JO}_n^+$. Moreover, $\bar{a}_i = b_{n-i}$ and $\bar{b}_i = a_{n-i}$, for $1 \leq i \leq n-1$.

The following property gives a characterization of \mathcal{JO}_n^- that allows us to easily count its number of elements.

Proposition 3.1. *For $n \geq 1$, $\mathcal{JO}_n^- = \{\alpha \in \mathcal{JO}_n \mid \text{Im}(\alpha) = \{1, \dots, k\}, \text{ for some } 1 \leq k \leq n\}$.*

Proof. Let $\alpha \in \mathcal{JO}_n^-$. Then, $1\alpha = 1$ and so $\text{Im}(\alpha) = \{1, \dots, k\}$, for some $1 \leq k \leq n$.

Conversely, let $\alpha \in \mathcal{JO}_n$ be such that $\text{Im}(\alpha) = \{1, \dots, k\}$, for some $1 \leq k \leq n$. Then, $1\alpha = \min \text{Im}(\alpha) = 1$. Assume by induction hypothesis that $t\alpha \leq t$, for (a fixed) $1 \leq t < n$. By contradiction, suppose that $(t+1)\alpha > t+1$. Then, $t\alpha \leq t < t+1 < (t+1)\alpha \leq k$, whence $t+1 \in \text{Im}(\alpha)$ and so $t+1 = x\alpha$, for some $x \in \Omega_n$. If $x \leq t$, then $t+1 = x\alpha \leq t\alpha \leq t$, which is a contradiction. Hence, $t+1 \leq x$ and so $t+1 < (t+1)\alpha \leq x\alpha = t+1$, which is again a contradiction. So, $(t+1)\alpha \leq t+1$. Thus, $t\alpha \leq t$, for all $t \in \Omega_n$, and so $\alpha \in \mathcal{JO}_n^-$, as required. \square

Of course, from the last proposition and the equality $\mathcal{JO}_n^+ = \mathcal{JO}_n^- \varphi$, it follows that

$$\mathcal{JO}_n^+ = \{\alpha \in \mathcal{JO}_n \mid \text{Im}(\alpha) = \{k, \dots, n\}, \text{ for some } 1 \leq k \leq n\}.$$

Proposition 3.2. *For $n \geq 1$, $|\mathcal{JO}_n^-| = |\mathcal{JO}_n^+| = 2^{n-1}$.*

Proof. Let $1 \leq k \leq n$. In view of Proposition 3.1, we have

$$\{\alpha \in \mathcal{JO}_n^- \mid |\text{Im}(\alpha)| = k\} = \{\alpha \in \mathcal{JO}_n^- \mid \text{Im}(\alpha) = \{1, \dots, k\}\}.$$

Then, for \mathcal{JO}_n^- , we can adapt the proof of [5, Theorem 1.5] for \mathcal{JO}_n , taking into account that instead of $n-k+1$ possibilities for images with k elements, we only have one possibility. In this way, we get

$$|\{\alpha \in \mathcal{JO}_n^- \mid |\text{Im}(\alpha)| = k\}| = \binom{n-1}{k-1}.$$

Thus, $|\mathcal{JO}_n^-| = \sum_{k=1}^n \binom{n-1}{k-1} = \sum_{k=0}^n \binom{n-1}{k} = 2^{n-1}$, as required. \square

Now, observe that $\mathcal{JO}_1^- = \mathcal{JO}_1 = \mathcal{T}_1$ and $\mathcal{JO}_2^- = \{(\begin{smallmatrix} 1 & 2 \\ 1 & 1 \end{smallmatrix}), (\begin{smallmatrix} 1 & 2 \\ 1 & 2 \end{smallmatrix})\}$. Then, clearly, \mathcal{JO}_1^- has rank 0 and \mathcal{JO}_2^- has rank 1. More generally, in what follows, for $n \geq 1$, we aim to show that A is a generating set with minimum size of the monoid \mathcal{JO}_n^- .

First, notice that $\{\alpha \in \mathcal{JO}_n^- \mid |\text{Im}(\alpha)| = n-1\} = A$ and, of course, like for \mathcal{JO}_n , the group of units of \mathcal{JO}_n^- is trivial. On the other hand, we have:

Lemma 3.3. *For $n \geq 3$, let $\alpha \in \mathcal{JO}_n^-$ be such that $|\text{Im}(\alpha)| = k$, for some $1 \leq k \leq n-2$. Then, there exist $\gamma_1, \gamma_2 \in \mathcal{JO}_n^-$ such that $|\text{Im}(\gamma_1)| = |\text{Im}(\gamma_2)| = k+1$ and $\alpha = \gamma_1\gamma_2$.*

Proof. It suffices to observe that in the proof of [5, Lemma 3.2], for an element $\alpha \in \mathcal{JO}_n^-$, we must have $j = 0$ and thus the transformations γ_1 and γ_2 there defined also belong to \mathcal{JO}_n^- . \square

Therefore, by a simple inductive reasoning, we can conclude that \mathcal{JO}_n^- is generated by A . In addition, by observing that a_1, \dots, a_{n-1} are the only elements of \mathcal{JO}_n^- with images of size $n-1$ and their kernels are pairwise distinct, it is immediate to deduce that a_1, \dots, a_{n-1} are undecomposable elements (i.e. they are not a product of two elements distinct from themselves) of \mathcal{JO}_n^- , and so they must belong to any generating set of \mathcal{JO}_n^- . Thus, we get:

Theorem 3.4. For $n \geq 1$, A is a generating set with minimum size of the monoid \mathcal{JO}_n^- . In particular, the monoid \mathcal{JO}_n^- has rank $n - 1$.

Since $\mathcal{JO}_n^+ = \mathcal{JO}_n^- \varphi$ and $\{b_1, \dots, b_{n-1}\} = \{a_1, \dots, a_{n-1}\} \varphi$, we also get that B is a generating set with minimum size of the monoid \mathcal{JO}_n^+ .

We finish this section by exhibiting a presentation for \mathcal{JO}_n^- .

Let R^- be the set of $\frac{1}{2}(n^2 - n)$ relations $R_1 \cup R_2$, considered on the alphabet A . Let $\varphi^- : A^* \rightarrow \mathcal{JO}_n^-$ be the homomorphism of monoids that extends the mapping $A \rightarrow \mathcal{JO}_n^-$ defined by $a_i \mapsto a_i$, for $1 \leq i \leq n - 1$. Clearly, all relations from R^- are satisfied (via φ^-) by the generators A of \mathcal{JO}_n^- .

Let

$$W^- = \{a_{i_1} \cdots a_{i_k} \in A^* \mid 0 \leq k \leq n - 1 \text{ and } 1 \leq i_k < \cdots < i_1 \leq n - 1\}.$$

It is clear that $|W^-| = 2^{n-1}$ (as there is an obvious bijection between W^- and the power set of $\{1, \dots, n - 1\}$) and W^- contains the empty word (taking $k = 0$) and A . Moreover, we have:

Lemma 3.5. Let $w \in A^*$. Then, there exists a word $w' \in W^-$ such that $|w'| \leq |w|$ and $w \sim_{R^-} w'$.

Proof. Let us denote \sim_{R^-} simply by \sim .

If $|w| = 1$, then $w \in W^-$.

Next, suppose that $w = a_{j_1} a_{j_2}$, with $1 \leq j_1, j_2 \leq n - 1$. If $j_2 < j_1$, then $w \in W^-$. So, suppose that $j_1 \leq j_2$. If $j_2 = n - 1$, then $w = a_{j_1} a_{n-1} \sim a_{j_1} \in W^-$. On the other hand, if $j_2 \leq n - 2$, then $w = a_{j_1} a_{j_2} \sim a_{j_2+1} a_{j_1} \in W^-$, since $j_1 < j_2 + 1$.

Now, assume by induction hypothesis that the property is valid for words $w \in A^*$ of length less than or equal to ℓ , for a given (fixed) $\ell \geq 2$.

Let $w = a_{j_1} \cdots a_{j_\ell} a_{j_{\ell+1}}$, for some $1 \leq j_1, \dots, j_{\ell+1} \leq n - 1$. Then, by induction hypothesis, $w \sim a_{i_1} \cdots a_{i_k} a_{j_{\ell+1}}$, with $1 \leq k \leq \ell$ and $1 \leq i_k < \cdots < i_1 \leq n - 1$.

If $j_{\ell+1} = n - 1$, then $w \sim a_{i_1} \cdots a_{i_k} \in W^-$. If $j_{\ell+1} < i_k$, then $w \sim a_{i_1} \cdots a_{i_k} a_{j_{\ell+1}} \in W^-$. So, suppose that $i_k \leq j_{\ell+1} < n - 1$. Hence, we have

$$w \sim a_{i_1} \cdots a_{i_k} a_{j_{\ell+1}} \sim a_{i_1} \cdots a_{i_{k-1}} a_{j_{\ell+1}+1} a_{i_k} \sim a_{i_1} \cdots a_{i_{k-t}} a_{j_{\ell+1}+t} a_{i_{k-t+1}} \cdots a_{i_k},$$

for all $1 \leq t \leq k$ such that $i_{k-t+1} \leq j_{\ell+1} + t - 1 \leq n - 2$. Let $t = \max\{1 \leq t \leq k \mid i_{k-t+1} \leq j_{\ell+1} + t - 1 \leq n - 2\}$. If $t = k$ (i.e. $i_1 \leq j_{\ell+1} + k - 1 \leq n - 2$), then $w \sim a_{j_{\ell+1}+k} a_{i_1} \cdots a_{i_k} \in W^-$, since $j_{\ell+1} + k > i_1$. Therefore, suppose that $t < k$. In this case, we have $j_{\ell+1} + t < i_{k-t}$ or $j_{\ell+1} + t = n - 1$. If $j_{\ell+1} + t = n - 1$, then $w \sim a_{i_1} \cdots a_{i_{k-t}} a_{i_{k-t+1}} \cdots a_{i_k} \in W^-$. On the other hand, if $j_{\ell+1} + t < i_{k-t}$, then $w \sim a_{i_1} \cdots a_{i_{k-t}} a_{j_{\ell+1}+t} a_{i_{k-t+1}} \cdots a_{i_k} \in W^-$, since $1 \leq i_k < \cdots < i_{k-t+1} < j_{\ell+1} + t < i_{k-t} < \cdots < i_1 \leq n - 1$.

We finish this proof by observing that, in each case, we obtained $w \sim w' \in W^-$ with $|w'| \leq k + 1 \leq \ell + 1 = |w|$, as required. \square

At this moment, we gather all the ingredients to, given Proposition 1.1, conclude:

Theorem 3.6. For $n \geq 1$, the monoid \mathcal{JO}_n^- is defined by the presentation $\langle A \mid R^- \rangle$ on $n - 1$ generators and $\frac{1}{2}(n^2 - n)$ relations.

Since $\mathcal{JO}_n^+ = \mathcal{JO}_n^- \varphi$ and $\bar{a}_i = b_{n-i}$ and $\bar{b}_i = a_{n-i}$, for $1 \leq i \leq n - 1$, being $R^+ = R_3 \cup R_4$, we can also conclude that the monoid \mathcal{JO}_n^+ is defined by the presentation $\langle B \mid R^+ \rangle$ on $n - 1$ generators and $\frac{1}{2}(n^2 - n)$ relations.

For reference below, observe also that, a property for $\langle B \mid R^+ \rangle$ analogous to Lemma 3.5 is also valid: let

$$W^+ = \{b_{j_1} \cdots b_{j_\ell} \in B^* \mid 0 \leq \ell \leq n - 1 \text{ and } 1 \leq j_1 < \cdots < j_\ell \leq n - 1\}.$$

Then, we have:

Lemma 3.7. Let $w \in B^*$. Then, there exists a word $w' \in W^+$ such that $|w'| \leq |w|$ and $w \sim_{R^+} w'$.

4 Main result

In this section, we prove that the presentation $\langle A \cup B \mid R \rangle$ defines the monoid \mathcal{JO}_n and, by means of substitutions (*Titze transformations*), we also deduce another presentation for \mathcal{JO}_n in terms of the minimal generators $a_1, \dots, a_{n-2}, b_{n-1}$.

First, we set forth a series of lemmas.

Lemma 4.1. *For $2 \leq j \leq i \leq n-1$, $b_i a_j \sim_R a_{j-1} b_{i-1}$.*

Proof. Let us denote \sim_R simply by \sim . Then, we have

$$b_i a_j \sim b_i b_j a_1 \sim b_{j-1} b_i a_1 \sim b_{j-1} a_i \sim a_i b_{j-1} \sim a_i a_{j-1} b_{n-1} \sim a_{j-1} a_{i-1} b_{n-1} \sim a_{j-1} b_{i-1},$$

as required. \square

Lemma 4.2. *Let $u \in A^*$ and $1 \leq i \leq n-1$. Then, there exist $v \in A^*$ and $1 \leq k \leq i \leq n-1$ such that $b_i u \sim_R v$ or $b_i u \sim_R v b_k$.*

Proof. For $|u| = 0$, the lemma is immediate. So, assume by induction hypothesis that the lemma is valid for words $u \in A^*$ of length equal to ℓ , for a given (fixed) $\ell \geq 0$. Let $u' \in A^*$ be such that $|u'| = \ell + 1$. Then, $u' = u a_j$, for some $1 \leq j \leq n-1$ and $u \in A^*$ such that $|u| = \ell$. Hence, by induction hypothesis, there exist $v \in A^*$ and $1 \leq k \leq i \leq n-1$ such that $b_i u \sim_R v$ or $b_i u \sim_R v b_k$. If $b_i u \sim_R v$, then $b_i u' = b_i u a_j \sim_R v a_j \in A^*$. On the other hand, if $b_i u \sim_R v b_k$, then

$$b_i u' = b_i u a_j \sim_R v b_k a_j \sim_R \begin{cases} v a_j b_k & \text{if } 1 \leq k < j \leq n-1 \\ v a_{j-1} b_{k-1} & \text{if } 2 \leq j \leq k \leq n-1 \\ v a_k & \text{if } j = 1, \end{cases}$$

as required. \square

Since we can write any word $w \in (A \cup B)^*$ in the form

$$w = u_0 b_{i_1} u_1 \cdots b_{i_k} u_k,$$

with $1 \leq i_1 \leq \cdots \leq i_k \leq n-1$, $u_0, u_1, \dots, u_k \in A^*$ and $k \geq 0$, by Lemma 4.2, it is easy to conclude:

Corollary 4.3. *Let $w \in (A \cup B)^*$. Then, there exist $u \in A^*$ and $v \in B^*$ such that $w \sim_R uv$.*

Lemma 4.4. *Let $v \in B^+$. Then, there exist $u \in A^+$ and $1 \leq \ell \leq n-1$ such that $v \sim_R u b_{n-1}^\ell$.*

Proof. By Lemma 3.7, we can take $1 \leq \ell \leq n-1$ and $1 \leq j_1 < \cdots < j_\ell \leq n-1$ such that $v \sim_{R^+} b_{j_1} \cdots b_{j_\ell}$. Then, denoting \sim_R simply by \sim , we have

$$\begin{aligned} v &\sim b_{j_1} b_{j_2} \cdots b_{j_\ell} \sim a_{j_1} b_{n-1} b_{j_2} \cdots b_{j_\ell} \sim a_{j_1} b_{j_2-1} \cdots b_{j_\ell-1} b_{n-1} \sim a_{j_1} a_{j_2-1} b_{n-1} b_{j_3-1} \cdots b_{j_\ell-1} b_{n-1} \sim \\ &\sim a_{j_1} a_{j_2-1} b_{j_3-2} \cdots b_{j_\ell-2} b_{n-1}^2 \sim \cdots \sim a_{j_1} a_{j_2-1} a_{j_3-2} \cdots a_{j_\ell-\ell+1} b_{n-1}^\ell, \end{aligned}$$

as required. \square

Lemma 4.5. *For $1 \leq \ell \leq n-1$ and $n-\ell \leq i \leq n-1$, $a_i b_{n-1}^\ell \sim_R b_{n-1}^\ell$.*

Proof. Denote \sim_R simply by \sim . If $\ell = 1$, then $i = n-1$ and we immediately have $a_{n-1} b_{n-1} \sim b_{n-1}$. So, suppose that $\ell \geq 2$. Then, we get

$$a_i b_{n-1}^\ell = a_i b_{n-1} b_{n-1}^{\ell-1} \sim b_i b_{n-1}^{\ell-1} = b_i b_{n-1} b_{n-1}^{\ell-2} \sim b_{n-1} b_{i+1} b_{n-1}^{\ell-2} \sim \cdots \sim b_{n-1}^t b_{i+t} b_{n-1}^{\ell-t-1},$$

for $1 \leq t \leq n-i-1$ (notice that $n-\ell \leq i$ implies that $t \leq \ell-1$). In particular,

$$a_i b_{n-1}^\ell \sim b_{n-1}^{n-i-1} b_{n-1} b_{n-1}^{\ell-n+i} = b_{n-1}^\ell,$$

as required. \square

Now, for $1 \leq \ell \leq n-1$, let

$$W_\ell = \{a_{i_1} \cdots a_{i_k} b_{n-1}^\ell \mid 0 \leq k \leq n-1 \text{ and } 1 \leq i_k < \cdots < i_1 \leq n-\ell+k-2\}.$$

Then, we have:

Lemma 4.6. *Let $u \in A^*$ and $v \in B^+$. Then, there exist $1 \leq \ell \leq n-1$ and $w \in W_\ell$ such that $wv \sim_R w$.*

Proof. Denote \sim_R simply by \sim . By Lemma 4.4, there exist $u' \in A^+$ and $1 \leq \ell \leq n-1$ such that $v \sim u' b_{n-1}^\ell$. Hence, $wv \sim wu' b_{n-1}^\ell$. Then, by Lemma 3.5, there exist $0 \leq k \leq n-1$ and $1 \leq i_k < \cdots < i_1 \leq n-1$ such that $wu' \sim a_{i_1} \cdots a_{i_k} b_{n-1}^\ell$. Hence, $wv \sim a_{i_1} \cdots a_{i_k} b_{n-1}^\ell$.

If $i_1 \leq n-\ell+k-2$, then $wv \sim a_{i_1} \cdots a_{i_k} b_{n-1}^\ell \in W_\ell$.

So, suppose that $i_1 > n-\ell+k-2$. Then, it is easy to show that, there exists $1 \leq t \leq k$ such that $i_t > n-\ell+k-t-1 > i_{t+1}$, where (if $t=k$) $i_{k+1} = -1$. Next, observe that

$$\begin{aligned} a_{i_1} a_{i_2} \cdots a_{i_k} &\sim a_{i_2} a_{i_1-1} a_{i_3} \cdots a_{i_k} \sim a_{i_2} \cdots a_{i_k} a_{i_1-k+1} \sim a_{i_3} \cdots a_{i_k} a_{i_2-k+2} a_{i_1-k+1} \sim \\ &\sim \cdots \sim a_{i_{t+1}} \cdots a_{i_k} a_{i_t-k+t} a_{i_{t-1}-k+t-1} \cdots a_{i_1-k+1}, \end{aligned}$$

$n-\ell \leq i_t - k + t \leq i_{t-1} - k + t - 1 \leq \cdots \leq i_1 - k + 1$, since $i_t > n-\ell+k-t-1$ and $i_t < \cdots < i_1$, and $i_{t+1} \leq n-\ell+(k-t)-2$. Thus, by Lemma 4.5, we have

$$wv \sim a_{i_1} \cdots a_{i_k} b_{n-1}^\ell \sim a_{i_{t+1}} \cdots a_{i_k} b_{n-1}^\ell \in W_\ell,$$

as required. \square

Let $W = W^- \cup W_1 \cup \cdots \cup W_{n-1}$. Therefore, as an immediate consequence of Corollary 4.3 and Lemmas 3.5 and 4.6, we obtain:

Corollary 4.7. *Let $w \in (A \cup B)^*$. Then, there exists $w' \in W$ such that $w \sim_R w'$.*

Now, observe that

$$|W| = |W^-| + \sum_{\ell=1}^{n-1} |W_\ell| = 2^{n-1} + \sum_{\ell=1}^{n-1} \sum_{k=0}^{n-1} \binom{\min\{n-\ell+k-2, n-1\}}{k} = \sum_{\ell=0}^{n-1} \sum_{k=0}^{n-1} \binom{\min\{n-\ell+k-2, n-1\}}{k}$$

and it is not difficult to show that $\sum_{\ell=0}^{n-1} \sum_{k=0}^{n-1} \binom{\min\{n-\ell+k-2, n-1\}}{k} = (n+1)2^{n-2}$, whence $|W| = |\mathcal{JO}_n|$. Therefore, we are in a position to apply Proposition 1.1 to conclude our main result:

Theorem 4.8. *For $n \geq 1$, the monoid \mathcal{JO}_n is defined by the presentation $\langle A \cup B \mid R \rangle$ on $2n-2$ generators and $\frac{1}{2}(3n^2 - n - 2)$ relations.*

Taking into account the equalities $a_{n-1} = b_{n-1} a_1$ and $b_i = a_i b_{n-1}$, for $1 \leq i \leq n-2$, by making these substitutions into the set of relations R and eliminating, from among those relations obtained, both the immediately trivial and the trivially deducible, we obtain the following set R' of $\frac{1}{2}(3n^2 - 7n + 4)$ relations on the alphabet $a_1, \dots, a_{n-2}, b_{n-1}$:

$$(R'_2) \ a_i a_j = a_{j+1} a_i, \quad 1 \leq i \leq j \leq n-3;$$

$$a_i a_{n-2} = b_{n-1} a_1 a_i, \quad 1 \leq i \leq n-2;$$

$$(R'_4) \ a_i b_{n-1} a_{j+1} b_{n-1} = a_{j+1} b_{n-1} a_{i+1} b_{n-1}, \quad 1 \leq i \leq j \leq n-3;$$

$$a_i b_{n-1}^2 = b_{n-1} a_{i+1} b_{n-1}, \quad 1 \leq i \leq n-3;$$

$$a_{n-2} b_{n-1}^2 = b_{n-1}^2;$$

$$(R'_5) a_i b_{n-1} a_j = a_j a_i b_{n-1}, \quad 1 \leq i < j \leq n-2;$$

$$a_i b_{n-1}^2 a_1 = b_{n-1} a_1 a_i b_{n-1}, \quad 1 \leq i \leq n-2;$$

$$(R'_6) a_i b_{n-1} a_1 = a_i, \quad 1 \leq i \leq n-2;$$

$$(R'_7) b_{n-1} a_1 b_{n-1} = b_{n-1}.$$

Therefore, we also have:

Corollary 4.9. *For $n \geq 3$, the monoid \mathcal{JO}_n is defined by the presentation $\langle a_1, \dots, a_{n-2}, b_{n-1} \mid R' \rangle$ on $n-1$ generators and $\frac{1}{2}(3n^2 - 7n + 4)$ relations.*

References

- [1] A.Ya. Aĭzenštāt, The defining relations of the endomorphism semigroup of a finite linearly ordered set, *Sibirsk. Mat. 3* (1962), 161–169 (Russian).
- [2] A.Ya. Aĭzenštāt, Homomorphisms of semigroups of endomorphisms of ordered sets, *Uch. Zap., Leningr. Gos. Pedagog. Inst.* 238 (1962), 38–48 (Russian).
- [3] V.H. Fernandes, Semigroups of order-preserving mappings on a finite chain: a new class of divisors, *Semigroup Forum* 54 (1997), 230–236.
- [4] V.H. Fernandes, M.M. Jesus, V. Maltcev and J.D. Mitchell, Endomorphisms of the semigroup of order-preserving mappings, *Semigroup Forum* 81 (2010), 277–285.
- [5] V.H. Fernandes and T. Paulista, On the Rank of Monoids of Endomorphisms of a Finite Directed Path, *Asian-Eur. J. Math.* 16 (2023), 2350069.
- [6] V.H. Fernandes and M. V. Volkov, On divisors of semigroups of order-preserving mappings of a finite chain, *Semigroup Forum* 81 (2010), 551–554.
- [7] The GAP Group, *GAP – Groups, Algorithms, and Programming, Version 4.13.0*; 2024. (<https://www.gap-system.org>)
- [8] G.M.S. Gomes and J.M. Howie, On the ranks of certain semigroups of order-preserving transformations, *Semigroup Forum* 45 (1992), 272–282.
- [9] P.M. Higgins, Divisors of semigroups of order-preserving mappings on a finite chain, *Internat. J. Algebra Comput.* 5 (1995), 725–742.
- [10] J.M. Howie, Product of idempotents in certain semigroups of transformations, *Proc. Edinburgh Math. Soc.* 17 (1971), 223–236.
- [11] J.M. Howie, *Fundamentals of Semigroup Theory*, Clarendon Press, Oxford, 1995.
- [12] A. Laradji and A. Umar, Combinatorial results for semigroups of order-preserving full transformations, *Semigroup Forum* 72 (2006), 51–62.
- [13] N. Ruškuc, *Semigroup Presentations*, Ph.D. Thesis, University of St-Andrews, 1995.
- [14] A.S. Vernitskii and M.V. Volkov, A proof and a generalisation of Higgins’ division theorem for semigroups of order preserving mappings, *Izv. Vyssh. Uchebn. Zaved. Mat* 1 (1995), 38–44

VÍTOR H. FERNANDES, Center for Mathematics and Applications (NOVA Math) and Department of Mathematics, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, Monte da Caparica, 2829-516 Caparica, Portugal; e-mail: vhf@fct.unl.pt.