

# SPECTRALITY OF A CLASS OF INFINITE CONVOLUTIONS WITH AND WITHOUT COMPACT SUPPORTS IN $\mathbb{R}^d$

YAO-QIANG LI

ABSTRACT. Generalizing a result given by Li, Miao and Wang in 2022, we study the spectrality of a class of infinite convolutions in  $\mathbb{R}^d$  generated by sequences of nearly  $d$ -th power lattices. This allows us to easily construct spectral measures with and without compact supports in  $\mathbb{R}^d$ . According to a result on the relation between supports of infinite convolutions and sets of infinite sums, we systematically study the Hausdorff and packing dimensions of infinite sums of finite sets in  $\mathbb{R}^d$ . As an application, we give concrete formulae for the Hausdorff and packing dimensions of the supports of a class of spectral measures in  $\mathbb{R}^d$  with the form of infinite convolutions generated by specific sequences of nearly  $d$ -th power lattices, and finally we deduce that there are spectral measures with and without compact supports of arbitrary Hausdorff and packing dimensions in  $\mathbb{R}^d$ .

## 1. INTRODUCTION

Let  $d \in \mathbb{N}$ . A Borel probability measure  $\mu$  on  $\mathbb{R}^d$  is called a *spectral measure* if there exists a countable set  $\Lambda \subseteq \mathbb{R}^d$  such that the family of exponential functions

$$\{e^{-2\pi i \langle \lambda, \cdot \rangle} : \lambda \in \Lambda\}$$

forms an orthonormal basis in  $L^2(\mu)$ . We call  $\Lambda$  a *spectrum* of  $\mu$ .

The existence of spectra of measures was initiated by Fuglede [19] in 1974. It is a basic question in harmonic analysis since the orthonormal basis consisting of exponential functions is used for Fourier series expansions of functions [37]. Note that any spectral measure must be of pure type: either discrete with finite support, singularly continuous, or absolutely continuous [20, 25]. Since Jorgensen and Pedersen [24] found the first singularly continuous spectral measure supported on a Cantor set in 1998, the spectrality of fractal measures are widely studied until now (see [1–15, 17, 18, 26–35, 38] and the references therein).

### 1.1. Spectrality of a class of infinite convolutions.

First we study the spectrality of a class of infinite convolutions, generalizing [29, Theorem 1.4] from  $\mathbb{R}$  to  $\mathbb{R}^d$ .

Let  $\delta_a$  denote the *Dirac measure* concentrated at the point  $a$ , and for a non-empty finite set  $A \subseteq \mathbb{R}^d$ , define the *uniform discrete measure* supported on  $A$  by

$$\delta_A := \frac{1}{\#A} \sum_{a \in A} \delta_a$$

where  $\#$  denotes the cardinality of a set.

Given a sequence  $\{R_k\}_{k \geq 1}$  of  $d \times d$  invertible real matrices and a sequence  $\{B_k\}_{k \geq 1}$  of non-empty finite subsets of  $\mathbb{R}^d$ , we define the finite convolution

$$\mu_n := \delta_{R_1^{-1}B_1} * \delta_{R_1^{-1}R_2^{-1}B_2} * \cdots * \delta_{R_1^{-1}R_2^{-1}\cdots R_n^{-1}B_n} \quad (1.1)$$

2020 *Mathematics Subject Classification*: 28A80, 42C30

*Key words and phrases*: infinite convolution, spectral measure, compact support, non-compact support, Hausdorff dimension, packing dimension

for each  $n \in \mathbb{N}$ . If  $\mu_n$  converges weakly to a Borel probability measure, we denote the weak limit measure by the infinite convolution

$$\mu := \delta_{R_1^{-1}B_1} * \delta_{R_1^{-1}R_2^{-1}B_2} * \delta_{R_1^{-1}R_2^{-1}R_3^{-1}B_3} * \cdots \quad (1.2)$$

and say that the infinite convolution exists.

A square matrix is called *expanding* if all eigenvalues have modulus strictly greater than 1. Given a  $d \times d$  expanding integer matrix  $R$  and a non-empty finite set  $B \subseteq \mathbb{Z}^d$  of integer vectors, we call  $(R, B)$  an *admissible pair* in  $\mathbb{R}^d$  if there exists a finite set  $L \subseteq \mathbb{Z}^d$  with  $\#L = \#B$  such that the matrix

$$\left[ \frac{1}{\sqrt{\#B}} e^{-2\pi i \langle R^{-1}b, l \rangle} \right]_{b \in B, l \in L}$$

is unitary. To emphasize  $L$ , we also call  $(R, B, L)$  a *Hadamard triple* in  $\mathbb{R}^d$ .

It is known that if  $\{(R_k, B_k)\}_{1 \leq k \leq n}$  are admissible pairs, then the finite convolution  $\mu_n$  defined by (1.1) is a spectral measure for each  $n \in \mathbb{N}$ . Since infinite convolutions generated by admissible pairs were raised by Strichartz [36] in 2000, the following question has received a lot of attention:

*Given a sequence of admissible pairs  $\{(R_k, B_k)\}_{k \geq 1}$ , under what condition is the infinite convolution defined by (1.2) a spectral measure?*

See for examples [1, 3, 4, 14, 15, 17, 18, 30–32, 35] for affirmative results for this question.

As mentioned in [29], the spectrality of infinite convolutions in  $\mathbb{R}^d$  is very complicated. Most of the existing research on the spectrality of infinite convolutions focuses on  $\mathbb{R}$  with compact supports. In [29, Theorem 1.4], Li, Miao and Wang focus on the spectrality of a class of infinite convolutions generated by sequences of nearly consecutive digit sets in  $\mathbb{R}$ , which may not be compactly supported. In the first main result in this paper, the following Theorem 1.1, we focus on the spectrality of a class of infinite convolutions generated by sequences of nearly  $d$ -th power lattices in  $\mathbb{R}^d$ , which may also not be compactly supported, generalizing [29, Theorem 1.4] from  $\mathbb{R}$  to  $\mathbb{R}^d$ , and then also generalizing [3, Theorem 1.4] and [29, Corollary 1.5].

Given a sequence of positive integers  $\{m_k\}_{k \geq 1}$  and a sequence of  $d \times d$  real matrices  $\{R_k\}_{k \geq 1}$ , we call a sequence  $\{B_k\}_{k \geq 1}$  of subsets of  $\mathbb{R}^d$  a *sequence of nearly  $d$ -th power lattices* with respect to  $\{m_k\}_{k \geq 1}$  and  $\{R_k\}_{k \geq 1}$  if

$$B_k \equiv \{0, 1, \dots, m_k - 1\}^d \pmod{R_k \mathbb{Z}^d} \quad \text{for all } k \in \mathbb{N}$$

and

$$\sum_{k=1}^{\infty} \frac{1}{m_k^d} \#(B_k \setminus \{0, 1, \dots, m_k - 1\}^d) < \infty. \quad (1.3)$$

Use  $R^T$  to denote the transpose of the matrix  $R$ . We have the following.

**Theorem 1.1.** *Let  $d \in \mathbb{N}$ ,  $\{m_k\}_{k \geq 1}$  be a sequence of integers no less than 2,  $\{R_k\}_{k \geq 1}$  be a sequence of  $d \times d$  invertible matrices, and  $\{B_k\}_{k \geq 1}$  be a sequence of nearly  $d$ -th power lattices with respect to  $\{m_k\}_{k \geq 1}$  and  $\{R_k\}_{k \geq 1}$ . If for every  $k \in \mathbb{N}$ , all entries of  $R_k$  are multiples of  $m_k$  and  $[-m_k, m_k]^d \subseteq R_k^T[-1, 1]^d$ , then the infinite convolution*

$$\mu = \delta_{R_1^{-1}B_1} * \delta_{R_1^{-1}R_2^{-1}B_2} * \delta_{R_1^{-1}R_2^{-1}R_3^{-1}B_3} * \cdots$$

*exists and is a spectral measure with a spectrum in  $\mathbb{Z}^d$ .*

Using this result, one can easily construct spectral measures with and without compact supports in  $\mathbb{R}^d$ .

To show the spectrality of  $\mu$  in Theorem 1.1, using Theorem 2.3, which is a tool developed by Li and Wang [32] recently, the main we need to prove is Lemma 3.1. The key in the proof of Lemma 3.1 is to estimate the lower bound of the modulus of the Fourier transform of the push-forward measure of the tail of the infinite convolution  $\mu$

on  $[-2/3, 2/3]^d$ . One will see that the estimation for our high-dimensional case is much more intricate than the one-dimensional case given in the proof of [29, Theorem 1.4].

For each  $k \in \mathbb{N}$ , the one-dimensional conditions  $b_k \leq N_k$  and  $b_k \mid N_k$  in [29, Theorem 1.4] are generalized to our high-dimensional conditions that  $[-m_k, m_k]^d \subseteq R_k^T[-1, 1]^d$  and all entries of the  $d \times d$  matrix  $R_k$  are multiples of  $m_k$  in Theorem 1.1. In the proof of the spectrality of  $\mu$  in Theorem 1.1, except for using Lemma 3.1, we need Proposition 3.3 to guarantee that  $R_k$  is expanding and then Proposition 3.2 to guarantee that  $(R_k, B_k)$  is an admissible pair in order to apply Theorem 2.3. In the proof of the existence of the infinite convolution  $\mu$  in Theorem 1.1, with the conditions (1.3) and  $[-m_k, m_k]^d \subseteq R_k^T[-1, 1]^d$  for all  $k \in \mathbb{N}$ , except for Theorem 2.1, we also need Proposition 3.4. These disclose the complicity of the high-dimensional case.

Note that the condition  $[-m_k, m_k]^d \subseteq R_k^T[-1, 1]^d$  in Theorem 1.1 is not equivalent to  $[-m_k, m_k]^d \subseteq R_k[-1, 1]^d$ . For example, take  $d = 2$ ,  $m_k = 2$  and

$$R_k = \begin{pmatrix} 4 & -2 \\ 0 & 2 \end{pmatrix}$$

for all  $k \in \mathbb{N}$ . Then

$$R_k^{-1} = \begin{pmatrix} 1/4 & 1/4 \\ 0 & 1/2 \end{pmatrix} \quad \text{and} \quad (R_k^T)^{-1} = \begin{pmatrix} 1/4 & 0 \\ 1/4 & 1/2 \end{pmatrix}.$$

By verifying

$$R_k^{-1} \left\{ \begin{pmatrix} 2 \\ 2 \end{pmatrix}, \begin{pmatrix} 2 \\ -2 \end{pmatrix}, \begin{pmatrix} -2 \\ 2 \end{pmatrix}, \begin{pmatrix} -2 \\ -2 \end{pmatrix} \right\} \subseteq [-1, 1]^2$$

we get  $R_k^{-1}[-2, 2]^2 \subseteq [-1, 1]^2$  and then  $[-2, 2]^2 \subseteq R_k[-1, 1]^2$ . But

$$(R_k^T)^{-1} \begin{pmatrix} 2 \\ 2 \end{pmatrix} = \begin{pmatrix} 1/2 \\ 3/2 \end{pmatrix} \notin [-1, 1]^2$$

implies  $(R_k^T)^{-1}[-2, 2]^2 \not\subseteq [-1, 1]^2$  and then  $[-2, 2]^2 \not\subseteq R_k^T[-1, 1]^2$ .

## 1.2. Supports of infinite convolutions.

In the last subsection, we confirm the spectrality of a class of infinite convolutions in  $\mathbb{R}^d$ . In this subsection, we systematically study the supports of general infinite convolutions.

Use  $\mathcal{P}(\mathbb{R}^d)$  to denote the set of all Borel Probability measures on  $\mathbb{R}^d$ . Given  $\mu_1, \mu_2, \dots \in \mathcal{P}(\mathbb{R}^d)$ , if the finite convolution

$$\mu_1 * \mu_2 * \dots * \mu_n$$

converges weakly to a Borel probability measure, we denote the weak limit measure by the infinite convolution

$$\mu_1 * \mu_2 * \mu_3 * \dots$$

and say that the infinite convolution exists.

For  $A_1, A_2, \dots \subseteq \mathbb{R}^d$ , define the infinite sum

$$\sum_{k=1}^{\infty} A_k := \left\{ x \in \mathbb{R}^d : \exists a_k \in A_k \text{ for each } k \in \mathbb{N} \text{ s.t. } x = \sum_{k=1}^{\infty} a_k \right\}.$$

Use  $\text{spt } \mu$  to denote the support of the measure  $\mu$  and  $\overline{A}$  to denote the closure of the set  $A$ . The following result represents the support of any infinite convolution as the closure of the infinite sum of the supports of all the individual measures, which enables us to conveniently study the supports of infinite convolutions by studying sets of infinite sums.

**Theorem 1.2.** *Let  $d \in \mathbb{N}$  and  $\mu_1, \mu_2, \dots \in \mathcal{P}(\mathbb{R}^d)$  such that  $\mu_1 * \mu_2 * \dots$  exists. Then*

$$\text{spt}(\mu_1 * \mu_2 * \dots) = \overline{\sum_{k=1}^{\infty} \text{spt } \mu_k}.$$

Immediately we get the following.

**Corollary 1.3.** *Let  $d \in \mathbb{N}$  and  $A_1, A_2, \dots \subseteq \mathbb{R}^d$  be non-empty finite sets such that  $\delta_{A_1} * \delta_{A_2} * \dots$  exists. Then*

$$\text{spt}(\delta_{A_1} * \delta_{A_2} * \dots) = \overline{\sum_{k=1}^{\infty} A_k}.$$

*Remark 1.4.* In Theorem 1.2,  $\overline{\sum_{k=1}^{\infty} \text{spt } \mu_k}$  can not be simplified to  $\sum_{k=1}^{\infty} \text{spt } \mu_k$ , and in Corollary 1.3,  $\overline{\sum_{k=1}^{\infty} A_k}$  can not be simplified to  $\sum_{k=1}^{\infty} A_k$ . At the end of Section 4, we will give an example in which  $A_1, A_2, \dots \subseteq [0, 1]$  are non-empty finite sets such that  $\delta_{A_1} * \delta_{A_2} * \dots$  exists but  $\sum_{k=1}^{\infty} A_k$  is not closed.

### 1.3. Infinite sums of finite sets.

In Corollary 1.3 we see that the support of the infinite convolution  $\delta_{A_1} * \delta_{A_2} * \dots$  is exactly equal to the closure of the infinite sum of the finite sets  $A_1, A_2, \dots$ . This motivates us to study the infinite sums of finite sets systematically in this subsection, especially their fractal dimensions (see Theorem 1.5 (1)) and the conditions under which they are closed sets (see Theorem 1.5 (2)).

In former research, fractal dimensions are usually studied for bounded sets. However, the infinite sums of finite sets we study in this subsection may be unbounded fractal sets. This allows us to apply the results to the non-compact supports of spectral measures in the next subsection.

For  $x = (x_1, x_2, \dots, x_d) \in \mathbb{R}^d$ , denote

$$|x| := \sqrt{x_1^2 + x_2^2 + \dots + x_d^2} \quad \text{and} \quad \|x\| := \max\{|x_1|, |x_2|, \dots, |x_d|\}.$$

Note that for all  $x \in \mathbb{R}^d$  we have

$$\|x\| \leq |x| \leq \sqrt{d} \|x\|. \quad (1.4)$$

For  $s \in [0, d]$ , use  $\mathcal{H}^s$  and  $\mathcal{P}^s$  to denote the  $s$ -dimensional Hausdorff measure and packing measure respectively [16]. Besides, we use  $\dim_H$  and  $\dim_P$  to denote the Hausdorff dimension and packing dimension respectively. Regard  $\min \emptyset$  as  $\infty$  throughout this paper.

The following result provides a way to simplify the calculation for the Hausdorff and packing dimensions of the infinite sums of finite sets. Under certain circumstances, this allows us to only consider bounded sets instead of unbounded sets with the form of infinite sums.

**Theorem 1.5.** *Let  $d \in \mathbb{N}$ . For each  $k \in \mathbb{N}$ , let  $A_k, A'_k \subseteq \mathbb{R}^d$  be finite sets, where  $A_k$  is non-empty and  $A'_k$  may be empty.*

(1) *If  $\underline{\lim}_{k \rightarrow \infty} \min_{a \in A'_k} |a| > 0$ , then*

①  $\mathcal{H}^s(\sum_{k=1}^{\infty} (A_k \cup A'_k)) = 0$  if and only if  $\mathcal{H}^s(\sum_{k=1}^{\infty} A_k) = 0$  for all  $s \in [0, d]$ , and

$$\dim_H \sum_{k=1}^{\infty} (A_k \cup A'_k) = \dim_H \sum_{k=1}^{\infty} A_k;$$

②  $\mathcal{P}^s(\sum_{k=1}^{\infty} (A_k \cup A'_k)) = 0$  if and only if  $\mathcal{P}^s(\sum_{k=1}^{\infty} A_k) = 0$  for all  $s \in [0, d]$ , and

$$\dim_P \sum_{k=1}^{\infty} (A_k \cup A'_k) = \dim_P \sum_{k=1}^{\infty} A_k.$$

(2) *If  $\sum_{k=1}^{\infty} \max_{a \in A_k} |a| < \infty$ ,  $A'_k \subseteq [0, \infty)^d$  for all  $k \in \mathbb{N}$  and  $\lim_{k \rightarrow \infty} \min_{a \in A'_k} |a| = \infty$ , then  $\sum_{k=1}^{\infty} (A_k \cup A'_k)$  is closed.*

In Theorem 1.5 (2), write  $S := \sum_{k=1}^{\infty} (A_k \cup A'_k)$  and we note the following.

- ① The condition  $\sum_{k=1}^{\infty} \max_{a \in A_k} |a| < \infty$  can not be omitted. For example, take  $d = 1$ ,  $A_k = \{0, \frac{k}{k+1}\}$  and  $A'_k = \{k + 1\}$  (or  $\emptyset$ ) for all  $k \in \mathbb{N}$ . Then  $A'_k \subseteq [0, \infty)$  for all  $k \in \mathbb{N}$  and  $\lim_{k \rightarrow \infty} \min_{a \in A'_k} |a| = \infty$ . But  $S$  is not closed, since one can easily verify  $1 \in \overline{S} \setminus S$ .
- ② The condition  $A'_k \subseteq [0, \infty)^d$  for all  $k \in \mathbb{N}$  can not be omitted. For example, take  $d = 1$ ,  $A_k = \{0\}$  and  $A'_k = \{(-1)^k(k + \frac{1}{2^k})\}$  for all  $k \in \mathbb{N}$ . Then  $\sum_{k=1}^{\infty} \max_{a \in A_k} |a| = 0 < \infty$  and  $\lim_{k \rightarrow \infty} \min_{a \in A'_k} |a| = \lim_{k \rightarrow \infty} (k + \frac{1}{2^k}) = \infty$ . But  $S$  is not closed, since one can easily verify  $1 \in \overline{S} \setminus S$ .
- ③ The condition  $\lim_{k \rightarrow \infty} \min_{a \in A'_k} |a| = \infty$  can not be weakened to  $\underline{\lim}_{k \rightarrow \infty} \min_{a \in A'_k} |a| > 0$ . For example, take  $d = 1$ ,  $A_k = \{0\}$  and  $A'_k = \{\frac{k}{k+1}\}$  for all  $k \in \mathbb{N}$ . Then  $\sum_{k=1}^{\infty} \max_{a \in A_k} |a| = 0 < \infty$ ,  $A'_k \subseteq [0, \infty)$  for all  $k \in \mathbb{N}$  and  $\underline{\lim}_{k \rightarrow \infty} \min_{a \in A'_k} |a| = 1 > 0$ . But  $S$  is not closed, since one can easily verify  $1 \in \overline{S} \setminus S$ .

As a corollary of Theorem 1.5, in the following we give concrete formulae for the Hausdorff and packing dimensions of a special class of infinite sums of finite sets, which may be unbounded. Here we use  $\mathcal{L}^d$  to denote the  $d$ -dimensional Lebesgue measure.

**Corollary 1.6.** *Let  $d \in \mathbb{N}$ . For each  $k \in \mathbb{N}$ , let  $c_k \geq 1$  and  $C_k \geq c_k + 1$  be real numbers, and  $B_k \subseteq \mathbb{R}^d$  be a finite set with  $\emptyset \neq B_k \cap [0, c_k]^d \subseteq \mathbb{Z}^d$ .*

(1) Suppose  $\underline{\lim}_{k \rightarrow \infty} C_1^{-1} \cdots C_k^{-1} \min\{\|x\| : x \in B_k \setminus [0, c_k]^d\} > 0$ .

- ① If  $\prod_{k=1}^{\infty} \frac{\#(B_k \cap [0, c_k]^d)}{C_k^d} = 0$ , then  $\mathcal{L}^d(\sum_{k=1}^{\infty} C_1^{-1} \cdots C_k^{-1} B_k) = 0$ .
- ② If  $\lim_{k \rightarrow \infty} \frac{\log C_k}{\log C_1 \cdots C_k} = 0$ , then

$$\dim_H \sum_{k=1}^{\infty} C_1^{-1} \cdots C_k^{-1} B_k = \underline{\lim}_{k \rightarrow \infty} \frac{\log \#(B_1 \cap [0, c_1]^d) \cdots \#(B_k \cap [0, c_k]^d)}{\log C_1 \cdots C_k},$$

and

$$\dim_P \sum_{k=1}^{\infty} C_1^{-1} \cdots C_k^{-1} B_k = \overline{\lim}_{k \rightarrow \infty} \frac{\log \#(B_1 \cap [0, c_1]^d) \cdots \#(B_k \cap [0, c_k]^d)}{\log C_1 \cdots C_k}.$$

(2) If  $B_k \subseteq [0, \infty)^d$  for all  $k \in \mathbb{N}$  and  $\lim_{k \rightarrow \infty} C_1^{-1} \cdots C_k^{-1} \min\{\|x\| : x \in B_k \setminus [0, c_k]^d\} = \infty$ , then  $\sum_{k=1}^{\infty} C_1^{-1} \cdots C_k^{-1} B_k$  is closed.

#### 1.4. Spectral measures with and without compact supports of arbitrary dimensions.

In this subsection, we use the above results to deduce that there are spectral measures with and without compact supports of arbitrary Hausdorff and packing dimensions in  $\mathbb{R}^d$ , generalizing [29, Theorem 1.7] from  $\mathbb{R}$  to  $\mathbb{R}^d$ .

First, taking  $c_k := m_k - 1$ ,  $C_k := N_k$  for all  $k \in \mathbb{N}$  in Corollary 1.6, by Theorem 1.1 and Corollary 1.3, one can get the following.

**Corollary 1.7.** *Let  $d \in \mathbb{N}$ . For each  $k \in \mathbb{N}$ , let  $N_k \geq m_k \geq 2$  be integers with  $m_k \mid N_k$  and  $B_k \subseteq \{0, 1, 2, \dots\}^d$  be a finite set with  $G_k := B_k \cap \{0, 1, \dots, m_k - 1\}^d \neq \emptyset$ . Suppose*

$$\underline{\lim}_{k \rightarrow \infty} \frac{\min\{\|x\| : x \in B_k \setminus \{0, 1, \dots, m_k - 1\}^d\}}{N_1 \cdots N_k} = \infty,$$

$$\lim_{k \rightarrow \infty} \frac{\log N_k}{\log N_1 \cdots N_k} = 0 \quad \text{and} \quad \prod_{k=1}^{\infty} \frac{\#G_k}{N_k^d} = 0,$$

and suppose that  $\{B_k\}_{k \geq 1}$  is a sequence of nearly  $d$ -th power lattices with respect to  $\{m_k\}_{k \geq 1}$  and the sequence of  $d \times d$  diagonal matrices  $\{\text{diag}(N_k, \dots, N_k)\}_{k \geq 1}$ . Then the infinite convolution

$$\mu = \delta_{N_1^{-1}B_1} * \delta_{N_1^{-1}N_2^{-1}B_2} * \delta_{N_1^{-1}N_2^{-1}N_3^{-1}B_3} * \cdots$$

exists, is a singular spectral measure with a spectrum in  $\mathbb{Z}^d$ ,  $\text{spt } \mu = \sum_{k=1}^{\infty} N_1^{-1} \cdots N_k^{-1} B_k$ ,

$$\dim_H \text{spt } \mu = \varliminf_{k \rightarrow \infty} \frac{\log \#G_1 \cdots \#G_k}{\log N_1 \cdots N_k} \quad \text{and} \quad \dim_P \text{spt } \mu = \varlimsup_{k \rightarrow \infty} \frac{\log \#G_1 \cdots \#G_k}{\log N_1 \cdots N_k}.$$

On spectral measures with compact supports in  $\mathbb{R}^d$ , we have the following. The similar result for spectral measures on  $\mathbb{R}$  has been given in [11].

**Corollary 1.8.** *Let  $d \in \mathbb{N}$ . For any  $\alpha, \beta \in [0, d]$  with  $\alpha \leq \beta$ , there exists a singular spectral measure  $\mu$  on  $\mathbb{R}^d$  with a spectrum in  $\mathbb{Z}^d$  and with compact support such that*

$$\dim_H \text{spt } \mu = \alpha \quad \text{and} \quad \dim_P \text{spt } \mu = \beta.$$

Finally we consider spectral measures without compact supports and generalize [29, Theorem 1.7] to  $\mathbb{R}^d$ .

**Corollary 1.9.** *Let  $d \in \mathbb{N}$ . For any  $\alpha, \beta \in [0, d]$  with  $\alpha \leq \beta$ , there exists a singular spectral measure  $\mu$  on  $\mathbb{R}^d$  with a spectrum in  $\mathbb{Z}^d$  and without compact support such that*

$$\dim_H \text{spt } \mu = \alpha \quad \text{and} \quad \dim_P \text{spt } \mu = \beta.$$

This paper is organized as follows. In the next section we give some preliminaries. Then we prove Theorem 1.1 in Section 3, prove Theorem 1.2 and give an example for Remark 1.4 in Section 4, prove Theorem 1.5 and Corollary 1.6 in Section 5, and finally deduce Corollaries 1.8 and 1.9 in Section 6.

## 2. PRELIMINARIES

Recall that  $\mathcal{P}(\mathbb{R}^d)$  denotes the set of all Borel probability measures on  $\mathbb{R}^d$ . For  $\mu \in \mathcal{P}(\mathbb{R}^d)$  the *Fourier transform* of  $\mu$  is defined by

$$\widehat{\mu}(\xi) := \int_{\mathbb{R}^d} e^{-2\pi i \langle \xi, x \rangle} d\mu(x) \quad \text{for all } \xi \in \mathbb{R}^d.$$

It is well-known that  $\widehat{\mu}$  is a bounded, continuous function with  $\widehat{\mu}(\mathbf{0}) = 1$ . See for example [22, Theorem 13.1].

For  $\mu, \mu_1, \mu_2, \dots \in \mathcal{P}(\mathbb{R}^d)$ , we say that  $\mu_n$  converges weakly to  $\mu$  if

$$\int_{\mathbb{R}^d} f d\mu_n \rightarrow \int_{\mathbb{R}^d} f d\mu \quad \text{for all } f \in C_b(\mathbb{R}^d),$$

where  $C_b(\mathbb{R}^d)$  denotes the set of all bounded continuous functions on  $\mathbb{R}^d$ .

For  $\mu, \nu \in \mathcal{P}(\mathbb{R}^d)$ , the *convolution*  $\mu * \nu$  is defined by

$$\mu * \nu(B) := \int_{\mathbb{R}^d} \mu(B - y) d\nu(y) = \int_{\mathbb{R}^d} \nu(B - x) d\mu(x) \quad \text{for every Borel set } B \subseteq \mathbb{R}^d.$$

Equivalently,  $\mu * \nu$  is the unique Borel probability measure on  $\mathbb{R}^d$  satisfying

$$\int_{\mathbb{R}^d} f(x) d\mu * \nu(x) = \int_{\mathbb{R}^d \times \mathbb{R}^d} f(x + y) d\mu \times \nu(x, y) \quad \text{for all } f \in C_b(\mathbb{R}^d).$$

It is straightforward to see  $\widehat{\mu * \nu}(\xi) = \widehat{\mu}(\xi) \widehat{\nu}(\xi)$  for all  $\xi \in \mathbb{R}^d$ .

On the existence of infinite convolutions, in the proof of [29, Theorem 1.1] one can see the following, which we need to use in the proof of Theorem 1.1.

**Theorem 2.1.** *Let  $d \in \mathbb{N}$  and  $A_1, A_2, \dots \subseteq \mathbb{R}^d$  be non-empty finite sets. If*

$$\sum_{k=1}^{\infty} \frac{1}{\#A_k} \sum_{a \in A_k} \frac{|a|}{1 + |a|} < \infty,$$

*then  $\delta_{A_1} * \delta_{A_2} * \dots$  exists.*

Let  $\mathbf{0}$  be the zero vector in  $\mathbb{R}^d$ . The following is an important concept we need.

**Definition 2.2** (Equi-positive). We call  $\Phi \subseteq \mathcal{P}(\mathbb{R}^d)$  an *equi-positive* family if there exist  $\varepsilon > 0$  and  $\delta > 0$  such that for each  $x \in [0, 1]^d$  and  $\mu \in \Phi$ , there exists an integer vector  $k_{x,\mu} \in \mathbb{Z}^d$  such that

$$|\widehat{\mu}(x + y + k_{x,\mu})| \geq \varepsilon$$

for all  $y \in \mathbb{R}^d$  with  $|y| < \delta$ , where  $k_{x,\mu} = \mathbf{0}$  for  $x = \mathbf{0}$ .

The equi-positivity property was introduced in [1,14] and used to study the spectrality of fractal measures with compact supports. Then it was generalized to the above version in [30,32] which can also be used to study the spectrality of measures without compact supports.

Given a sequence  $\{R_k\}_{k \geq 1}$  of  $d \times d$  invertible real matrices and a sequence  $\{B_k\}_{k \geq 1}$  of non-empty finite subsets of  $\mathbb{R}^d$ , recall the finite convolution  $\mu_n$  given by (1.1) and the infinite convolution  $\mu$  given by (1.2), and write the *tail* of  $\mu$  by

$$\mu_{>n} := \delta_{R_1^{-1}R_2^{-1}\dots R_{n+1}^{-1}B_{n+1}} * \delta_{R_1^{-1}R_2^{-1}\dots R_{n+2}^{-1}B_{n+2}} * \delta_{R_1^{-1}R_2^{-1}\dots R_{n+3}^{-1}B_{n+3}} * \dots$$

for each  $n \in \mathbb{N}$ . Obviously  $\mu = \mu_n * \mu_{>n}$ . Besides, the *push-forward* measure of  $\mu_{>n}$  is defined by

$$\nu_{>n}(\cdot) := \mu_{>n}(R_1^{-1}R_2^{-1}\dots R_n^{-1}\cdot), \quad (2.1)$$

that is,

$$\nu_{>n} = \delta_{R_{n+1}^{-1}B_{n+1}} * \delta_{R_{n+1}^{-1}R_{n+2}^{-1}B_{n+2}} * \delta_{R_{n+1}^{-1}R_{n+2}^{-1}R_{n+3}^{-1}B_{n+3}} * \dots$$

In the proof of [32, Theorem 1.1], Li and Wang actually showed the following. (See [30, Theorem 1.4] and [29, Theorem 4.2] for the version in  $\mathbb{R}$ .)

**Theorem 2.3** ([32]). Let  $d \in \mathbb{N}$  and  $\{(R_k, B_k)\}_{k \geq 1}$  be a sequence of admissible pairs in  $\mathbb{R}^d$ . Suppose that the infinite convolution  $\mu$  defined in (1.2) exists, and

$$|(R_n^T)^{-1}\dots(R_1^T)^{-1}x| \rightarrow 0 \quad \text{as } n \rightarrow \infty \quad \text{for all } x \in \mathbb{R}^d.$$

Let  $\{\nu_{>n}\}_{n \geq 1}$  be defined in (2.1). If there exists a subsequence  $\{\nu_{>n_j}\}_{j \geq 1}$  which is equi-positive, then  $\mu$  is a spectral measure with a spectrum in  $\mathbb{Z}^d$ .

For  $B_1, B_2, \dots \subseteq \mathbb{R}^d$ , define

$$\lim_{n \rightarrow \infty} B_n := \left\{ x \in \mathbb{R}^d : \exists b_n \in B_n \text{ for each } n \in \mathbb{N} \text{ s.t. } x = \lim_{n \rightarrow \infty} b_n \right\}.$$

The following old result [23, Theorem 3] given by Jessen and Wintner in 1935 is the main tool we use to prove Theorem 1.2.

**Theorem 2.4** ([23]). Let  $d \in \mathbb{N}$  and  $\mu_1, \mu_2, \dots \in \mathcal{P}(\mathbb{R}^d)$  such that  $\mu_1 * \mu_2 * \dots$  exists. Then

$$\text{spt}(\mu_1 * \mu_2 * \dots) = \lim_{n \rightarrow \infty} (\text{spt} \mu_1 + \dots + \text{spt} \mu_n).$$

The following is the well-known Stolz-Cesàro Theorem.

**Theorem 2.5.** Let  $\beta_1, \beta_2, \beta_3, \dots \in (0, \infty)$  such that  $\sum_{n=1}^{\infty} \beta_n = \infty$  and let  $\alpha_1, \alpha_2, \alpha_3, \dots \in \mathbb{R}$ . Then

$$\liminf_{n \rightarrow \infty} \frac{\alpha_1 + \alpha_2 + \dots + \alpha_n}{\beta_1 + \beta_2 + \dots + \beta_n} \geq \liminf_{n \rightarrow \infty} \frac{\alpha_n}{\beta_n} \quad \text{and} \quad \overline{\lim}_{n \rightarrow \infty} \frac{\alpha_1 + \alpha_2 + \dots + \alpha_n}{\beta_1 + \beta_2 + \dots + \beta_n} \leq \overline{\lim}_{n \rightarrow \infty} \frac{\alpha_n}{\beta_n}.$$

In particular, if  $\lim_{n \rightarrow \infty} \frac{\alpha_n}{\beta_n}$  exists, then

$$\lim_{n \rightarrow \infty} \frac{\alpha_1 + \alpha_2 + \dots + \alpha_n}{\beta_1 + \beta_2 + \dots + \beta_n} = \lim_{n \rightarrow \infty} \frac{\alpha_n}{\beta_n}.$$

We present two useful facts in the following to end this section.

**Proposition 2.6** (Lagrange's trigonometric equality). For all  $\theta \in \mathbb{R} \setminus \{2k\pi : k \in \mathbb{Z}\}$  and  $n \in \{0, 1, 2, \dots\}$ , we have

$$\sum_{k=0}^n \sin k\theta = \frac{\cos \frac{1}{2}\theta - \cos((n + \frac{1}{2})\theta)}{2 \sin \frac{1}{2}\theta} \quad \text{and} \quad \sum_{k=0}^n \cos k\theta = \frac{\sin \frac{1}{2}\theta + \sin((n + \frac{1}{2})\theta)}{2 \sin \frac{1}{2}\theta}.$$

**Proposition 2.7.** For all  $n \in \mathbb{N}$ , let  $a_n > c_n > 0$  and  $b_n > d_n > 0$  with  $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{c_n}{d_n} = r \in [0, \infty)$ . If  $\lim_{n \rightarrow \infty} \frac{b_n}{d_n} > 1$ , then  $\lim_{n \rightarrow \infty} \frac{a_n - c_n}{b_n - d_n} = r$ .

*Proof.* By  $\lim_{n \rightarrow \infty} \frac{b_n}{d_n} > 1$ , there exist  $N_0, k \in \mathbb{N}$  such that for all  $n > N_0$  we have  $\frac{b_n}{d_n} > 1 + \frac{2}{k-1}$ , which is equivalent to

$$(k+1)d_n < (k-1)b_n. \quad (2.2)$$

Arbitrarily take  $\varepsilon > 0$ . By  $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{c_n}{d_n} = r$ , there exists  $N > N_0$  such that for all  $n > N$  we have

$$\begin{cases} r - \frac{\varepsilon}{k} < \frac{a_n}{b_n} < r + \frac{\varepsilon}{k}, \\ r - \frac{\varepsilon}{k} < \frac{c_n}{d_n} < r + \frac{\varepsilon}{k}, \end{cases} \quad \text{i.e.,} \quad \begin{cases} (r - \frac{\varepsilon}{k})b_n < a_n < (r + \frac{\varepsilon}{k})b_n, \\ -(r + \frac{\varepsilon}{k})d_n < -c_n < -(r - \frac{\varepsilon}{k})d_n, \end{cases}$$

which imply

$$(r - \varepsilon)(b_n - d_n) \stackrel{\text{by (2.2)}}{<} (r - \frac{\varepsilon}{k})b_n - (r + \frac{\varepsilon}{k})d_n < a_n - c_n < (r + \frac{\varepsilon}{k})b_n - (r - \frac{\varepsilon}{k})d_n \stackrel{\text{by (2.2)}}{<} (r + \varepsilon)(b_n - d_n).$$

We get  $r - \varepsilon < \frac{a_n - c_n}{b_n - d_n} < r + \varepsilon$  for all  $n > N$ . Therefore  $\lim_{n \rightarrow \infty} \frac{a_n - c_n}{b_n - d_n} = r$ .  $\square$

In Proposition 2.7, in order to get the conclusion  $\lim_{n \rightarrow \infty} \frac{a_n - c_n}{b_n - d_n} = r$ , the condition  $\lim_{n \rightarrow \infty} \frac{b_n}{d_n} > 1$  can not be omitted. For example, take  $a_n = 10^n + n$ ,  $b_n = 10^n + 1$  and  $c_n = d_n = 10^n$  for all  $n \in \mathbb{N}$ . Then  $a_n > c_n > 0$  and  $b_n > d_n > 0$  for all  $n \in \mathbb{N}$  with  $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{c_n}{d_n} = 1$ . But  $\lim_{n \rightarrow \infty} \frac{a_n - c_n}{b_n - d_n} = \infty$ .

### 3. PROOF OF THEOREM 1.1

To prove Theorem 1.1, the main tool we need to use is the following.

**Lemma 3.1.** Let  $d \in \mathbb{N}$ ,  $\{m_k\}_{k \geq 1}$  be a sequence of positive integers no less than 2,  $\{R_k\}_{k \geq 1}$  be a sequence of  $d \times d$  invertible matrices,  $\{B_k\}_{k \geq 1}$  be a sequence of nearly  $d$ -th power lattices with respect to  $\{m_k\}_{k \geq 1}$  and  $\{R_k\}_{k \geq 1}$ , and  $\{\nu_{>n}\}_{n \geq 1}$  be given by (2.1). If  $[-m_k, m_k]^d \subseteq R_k^T[-1, 1]^d$  for every  $k \in \mathbb{N}$ , then there exists  $n_0 \in \mathbb{N}$  such that  $\{\nu_{>n}\}_{n \geq n_0}$  is equi-positive.

*Proof.* Let  $c_k := \#(B_k \setminus \{0, 1, \dots, m_k - 1\}^d)$  for all  $k \in \mathbb{N}$ .

(1) Prove that for all  $k \in \mathbb{N}$  and  $\xi = (\xi_1, \xi_2, \dots, \xi_d) \in [-\frac{\sqrt{6}}{m_k\pi}, \frac{\sqrt{6}}{m_k\pi}]^d$  we have

$$|\widehat{\delta}_{B_k}(\xi)| \geq \prod_{j=1}^d \left(1 - \frac{m_k^2 \pi^2 \xi_j^2}{6}\right) - \frac{2c_k}{m_k^d}.$$



Let  $k \in \mathbb{N}$  and  $\xi \in [-\frac{\sqrt{6}}{m_k\pi}, \frac{\sqrt{6}}{m_k\pi}]^d$ . Then

$$\begin{aligned}
|\widehat{\delta}_{B_k}(\xi)| &= \left| \int_{\mathbb{R}^d} e^{-2\pi i \langle \xi, x \rangle} d\delta_{B_k}(x) \right| = \left| \frac{1}{\#B_k} \sum_{b \in B_k} e^{-2\pi i \langle b, \xi \rangle} \right| \\
&\geq \frac{1}{m_k^d} \left| \sum_{b \in \{0,1,\dots,m_k-1\}^d} e^{-2\pi i \langle b, \xi \rangle} \right| - \frac{1}{m_k^d} \left| \sum_{b \in \{0,1,\dots,m_k-1\}^d} e^{-2\pi i \langle b, \xi \rangle} - \sum_{b \in B_k} e^{-2\pi i \langle b, \xi \rangle} \right| \\
&\geq \frac{1}{m_k^d} \left| \sum_{b_1, b_2, \dots, b_d \in \{0,1,\dots,m_k-1\}} e^{-2\pi i (b_1 \xi_1 + b_2 \xi_2 + \dots + b_d \xi_d)} \right| - \frac{2}{m_k^d} \cdot \#(B_k \setminus \{0,1,\dots,m_k-1\}^d) \\
&= \left| \frac{1}{m_k} \sum_{b_1=0}^{m_k-1} e^{-2\pi i b_1 \xi_1} \right| \cdot \left| \frac{1}{m_k} \sum_{b_2=0}^{m_k-1} e^{-2\pi i b_2 \xi_2} \right| \cdots \left| \frac{1}{m_k} \sum_{b_d=0}^{m_k-1} e^{-2\pi i b_d \xi_d} \right| - \frac{2c_k}{m_k^d} \\
&\geq \left(1 - \frac{m_k^2 \pi^2 \xi_1^2}{6}\right) \left(1 - \frac{m_k^2 \pi^2 \xi_2^2}{6}\right) \cdots \left(1 - \frac{m_k^2 \pi^2 \xi_d^2}{6}\right) - \frac{2c_k}{m_k^d}
\end{aligned}$$

where the last inequality follows from the fact that for all  $j \in \{1, 2, \dots, d\}$  we can prove

$$\left| \frac{1}{m_k} \sum_{b=0}^{m_k-1} e^{-2\pi i b \xi_j} \right| \geq 1 - \frac{m_k^2 \pi^2 \xi_j^2}{6} \geq 0.$$

Since the second inequality follows immediately from  $\xi \in [-\frac{\sqrt{6}}{m_k\pi}, \frac{\sqrt{6}}{m_k\pi}]^d$ , it suffices to prove the first one. If  $\xi_j \in \mathbb{Z}$ , the first inequality obviously holds. If  $\xi_j \notin \mathbb{Z}$ , by Lagrange's trigonometric equality Proposition 2.6 we get

$$\frac{1}{m_k} \left| \sum_{b=0}^{m_k-1} e^{-2\pi i b \xi_j} \right| = \frac{1}{m_k} \left| \frac{\sin(m_k \pi \xi_j)}{\sin(\pi \xi_j)} \right| \stackrel{(\star)}{\geq} \left| \frac{\sin(m_k \pi \xi_j)}{m_k \pi \xi_j} \right| \stackrel{(\star\star)}{\geq} 1 - \frac{m_k^2 \pi^2 \xi_j^2}{6},$$

where  $(\star)$  and  $(\star\star)$  follow respectively from  $|\sin x| \leq |x|$  and  $|\frac{\sin x}{x}| \geq 1 - \frac{x^2}{6}$  for all  $x \neq 0$ .

(2) Prove that for all  $n, k \in \mathbb{N}$  we have

$$(\mathbf{R}_{n,n+k}^T)^{-1} \left[ -\frac{2}{3}, \frac{2}{3} \right]^d \subseteq \left[ -\frac{2}{3m_{n+1} \cdots m_{n+k}}, \frac{2}{3m_{n+1} \cdots m_{n+k}} \right]^d$$

where  $\mathbf{R}_{n,n+k} := R_{n+k} R_{n+k-1} \cdots R_{n+1}$ . By the linearity of  $(\mathbf{R}_{n,n+k}^T)^{-1}$ , it suffices to prove

$$(R_{n+k}^T)^{-1} \cdots (R_{n+1}^T)^{-1} [-1, 1]^d \subseteq \left[ -\frac{1}{m_{n+1} \cdots m_{n+k}}, \frac{1}{m_{n+1} \cdots m_{n+k}} \right]^d. \quad (3.1)$$

Note that for all  $k \in \mathbb{N}$  we have the condition  $[-m_k, m_k]^d \subseteq R_k^T [-1, 1]^d$ , which is equivalent to

$$(R_k^T)^{-1} [-1, 1]^d \subseteq \left[ -\frac{1}{m_k}, \frac{1}{m_k} \right]^d. \quad (3.2)$$

Thus

$$(R_{n+1}^T)^{-1} [-1, 1]^d \subseteq \left[ -\frac{1}{m_{n+1}}, \frac{1}{m_{n+1}} \right]^d,$$

and then

$$(R_{n+2}^T)^{-1} (R_{n+1}^T)^{-1} [-1, 1]^d \subseteq (R_{n+2}^T)^{-1} \left[ -\frac{1}{m_{n+1}}, \frac{1}{m_{n+1}} \right]^d \subseteq \left[ -\frac{1}{m_{n+1} m_{n+2}}, \frac{1}{m_{n+1} m_{n+2}} \right]^d,$$

where the last inclusion follows from (3.2) and the linearity of  $(R_{n+2}^T)^{-1}$ .

Repeating the above process for finitely many times, we get (3.1).

(3) Prove that there exists  $n_0 \in \mathbb{N}$  and  $\varepsilon > 0$  such that for all  $n \geq n_0$  and  $\xi \in [-\frac{2}{3}, \frac{2}{3}]^d$  we have  $|\widehat{\nu}_{>n}(\xi)| \geq \varepsilon$ .

In fact, by  $\sum_{k=1}^{\infty} \frac{c_k}{m_k^d} < \infty$  there exists  $n_0 \in \mathbb{N}$  such that for all  $k \geq n_0$  we have

$$\frac{2c_k}{m_k^d} < \frac{2\pi^2}{27} \left(1 - \frac{2\pi^2}{27}\right)^d.$$

Let  $n \geq n_0$  and  $\xi \in [-\frac{2}{3}, \frac{2}{3}]^d$ . By

$$\nu_{>n} = \delta_{\mathbf{R}_{n,n+1}^{-1}B_{n+1}} * \delta_{\mathbf{R}_{n,n+2}^{-1}B_{n+2}} * \delta_{\mathbf{R}_{n,n+3}^{-1}B_{n+3}} * \cdots$$

we get

$$|\widehat{\nu}_{>n}(\xi)| = \left| \prod_{k=1}^{\infty} \widehat{\delta}_{\mathbf{R}_{n,n+k}^{-1}B_{n+k}}(\xi) \right| = \left| \prod_{k=1}^{\infty} \widehat{\delta}_{B_{n+k}}((\mathbf{R}_{n,n+k}^{-1})^T \xi) \right| = \prod_{k=1}^{\infty} \left| \widehat{\delta}_{B_{n+k}}((\mathbf{R}_{n,n+k}^T)^{-1} \xi) \right|. \quad (3.3)$$

For all  $k \in \mathbb{N}$ , it follows from  $\xi \in [-\frac{2}{3}, \frac{2}{3}]^d$ ,  $m_{n+1}, \dots, m_{n+k-1} \geq 2$  and (2) that

$$(\mathbf{R}_{n,n+k}^T)^{-1} \xi \in \left[ -\frac{2}{3 \cdot 2^{k-1} m_{n+k}}, \frac{2}{3 \cdot 2^{k-1} m_{n+k}} \right]^d \subseteq \left[ -\frac{2}{3m_{n+k}}, \frac{2}{3m_{n+k}} \right]^d \subseteq \left[ -\frac{\sqrt{6}}{m_{n+k}\pi}, \frac{\sqrt{6}}{m_{n+k}\pi} \right]^d.$$

Use  $((\mathbf{R}_{n,n+k}^T)^{-1} \xi)_j$  to denote the  $j$ th coordinate of  $(\mathbf{R}_{n,n+k}^T)^{-1} \xi$  for  $j \in \{1, \dots, d\}$ . For all  $k \in \mathbb{N}$ , by (1) we get

$$\begin{aligned} \left| \widehat{\delta}_{B_{n+k}}((\mathbf{R}_{n,n+k}^T)^{-1} \xi) \right| &\geq \prod_{j=1}^d \left( 1 - \frac{m_{n+k}^2 \pi^2 ((\mathbf{R}_{n,n+k}^T)^{-1} \xi)_j^2}{6} \right) - \frac{2c_{n+k}}{m_{n+k}^d} \\ &\geq \prod_{j=1}^d \left( 1 - \frac{m_{n+k}^2 \pi^2}{6} \cdot \left( \frac{2}{3 \cdot 2^{k-1} m_{n+k}} \right)^2 \right) - \frac{2c_{n+k}}{m_{n+k}^d} \\ &= \left( 1 - \frac{2\pi^2}{27 \cdot 4^{k-1}} \right)^d - \frac{2c_{n+k}}{m_{n+k}^d} \\ &> \left( 1 - \frac{2\pi^2}{27} \right)^d - \frac{2\pi^2}{27} \left( 1 - \frac{2\pi^2}{27} \right)^d \\ &= \left( 1 - \frac{2\pi^2}{27} \right)^{d+1} \\ &> 0. \end{aligned} \quad (3.4)$$

Let

$$\alpha := \frac{(d+1) \ln(1 - \frac{2\pi^2}{27})}{(1 - \frac{2\pi^2}{27})^{d+1} - 1} > 0.$$

Then one can verify

$$x \geq e^{\alpha(x-1)} > 0 \quad \text{for all } x \in \left[ \left( 1 - \frac{2\pi^2}{27} \right)^{d+1}, 1 \right]. \quad (3.5)$$

It follows from (3.3), (3.4) and (3.5) that

$$\begin{aligned}
|\widehat{\nu}_{>n}(\xi)| &\geq \prod_{k=1}^{\infty} \left( \left(1 - \frac{2\pi^2}{27 \cdot 4^{k-1}}\right)^d - \frac{2c_{n+k}}{m_{n+k}^d} \right) \\
&\geq \prod_{k=1}^{\infty} \exp \left( \alpha \left( \left(1 - \frac{2\pi^2}{27 \cdot 4^{k-1}}\right)^d - 1 - \frac{2c_{n+k}}{m_{n+k}^d} \right) \right) \\
&\geq \prod_{k=1}^{\infty} \exp \left( \alpha \left( 1 - \left(1 + \frac{2\pi^2}{27 \cdot 4^{k-1}}\right)^d - \frac{2c_{n+k}}{m_{n+k}^d} \right) \right) \\
&= \prod_{k=1}^{\infty} \exp \left( -\alpha \left( \sum_{j=1}^d \binom{d}{j} \left( \frac{2\pi^2}{27 \cdot 4^{k-1}} \right)^j + \frac{2c_{n+k}}{m_{n+k}^d} \right) \right) \\
&= \exp \left( -\alpha \sum_{k=1}^{\infty} \left( \sum_{j=1}^d \binom{d}{j} \left( \frac{2\pi^2}{27} \right)^j \cdot \frac{1}{4^{j(k-1)}} + \frac{2c_{n+k}}{m_{n+k}^d} \right) \right) \\
&= \exp \left( -\alpha \left( \sum_{j=1}^d \binom{d}{j} \left( \frac{2\pi^2}{27} \right)^j \sum_{k=1}^{\infty} \left( \frac{1}{4^j} \right)^{k-1} + 2 \sum_{k=1}^{\infty} \frac{c_{n+k}}{m_{n+k}^d} \right) \right) \\
&\geq \exp \left( -\alpha \sum_{j=1}^d \binom{d}{j} \frac{8^j \pi^{2j}}{27^j (4^j - 1)} - 2\alpha \sum_{k=1}^{\infty} \frac{c_k}{m_k^d} \right) \stackrel{\text{denoted by}}{=} \varepsilon > 0
\end{aligned}$$

for all  $n \geq n_0$  and  $\xi \in \left[-\frac{2}{3}, \frac{2}{3}\right]^d$ , where  $\binom{d}{j} := \frac{d!}{(d-j)!j!}$ .

(4) Prove that  $\{\nu_{>n}\}_{n \geq n_0}$  is equi-positive.

Let  $\delta = \frac{1}{6}$ . For each  $x = (x_1, \dots, x_d) \in [0, 1)^d$ , define  $k = k(x) = (k_1, \dots, k_d) \in \mathbb{Z}^d$  by

$$k_j := \begin{cases} 0 & \text{if } x_j \in [0, \frac{1}{2}) \\ -1 & \text{if } x_j \in [\frac{1}{2}, 1) \end{cases} \quad \text{for all } j \in \{1, \dots, d\}.$$

Then for all  $x \in [0, 1)^d$ ,  $n \geq n_0$  and  $y \in \mathbb{R}^d$  with  $|y| < \delta$ , we have  $x + k(x) + y \in \left[-\frac{2}{3}, \frac{2}{3}\right]^d$ , and by (3) we get  $|\widehat{\nu}_{>n}(x + k(x) + y)| \geq \varepsilon$ . Therefore  $\{\nu_{>n}\}_{n \geq n_0}$  is equi-positive.  $\square$

Before deducing Theorem 1.1 from Lemma 3.1 and Theorem 2.3, we need the following Propositions 3.2, 3.3 and 3.4.

**Proposition 3.2.** *Let  $d \in \mathbb{N}$ ,  $m \geq 2$  be an integer,  $R$  be a  $d \times d$  expanding matrix with integer entries which are all multiples of  $m$ , and  $B \subseteq \mathbb{R}^d$  such that  $B \equiv \{0, 1, \dots, m-1\}^d \pmod{R\mathbb{Z}^d}$ . Then  $B \subseteq \mathbb{Z}^d$  and  $(R, B)$  is an admissible pair in  $\mathbb{R}^d$ .*

*Proof.* By  $B \equiv \{0, 1, \dots, m-1\}^d \pmod{R\mathbb{Z}^d}$ , for every  $u \in \{0, 1, \dots, m-1\}^d$ , there exists  $z_u \in \mathbb{Z}^d$  such that  $B = \{u + Rz_u : u \in \{0, 1, \dots, m-1\}^d\} \subseteq \mathbb{Z}^d$ . Let  $L := \frac{1}{m}R^T\{0, 1, \dots, m-1\}^d$ . We need to prove that the matrix

$$\left[ \frac{1}{\sqrt{m^d}} e^{-2\pi i \langle R^{-1}b, l \rangle} \right]_{b \in B, l \in L}$$

is unitary, i.e.,

$$\frac{1}{\sqrt{m^d}} \left[ e^{-2\pi i \langle R^{-1}(u+Rz_u), \frac{1}{m}R^T v \rangle} \right]_{u, v \in \{0, 1, \dots, m-1\}^d}$$

is unitary. Note that

$$e^{-2\pi i \langle R^{-1}(u+Rz_u), \frac{1}{m}R^T v \rangle} = e^{-2\pi i \langle R^{-1}u, \frac{1}{m}R^T v \rangle} \cdot e^{-2\pi i \langle z_u, \frac{1}{m}R^T v \rangle} \stackrel{(\star)}{=} e^{-\frac{2\pi i}{m} \langle R^{-1}u, R^T v \rangle} = e^{-\frac{2\pi i}{m} \langle u, v \rangle}$$

for all  $u, v \in \{0, 1, \dots, m-1\}^d$ , where  $(\star)$  follows from  $\langle z_u, \frac{1}{m}R^T v \rangle \in \mathbb{Z}$  since all entries of  $R^T$  are multiples of  $m$ . We only need to prove that

$$\frac{1}{\sqrt{m^d}} \left[ e^{-\frac{2\pi i}{m} \langle u, v \rangle} \right]_{u, v \in \{0, 1, \dots, m-1\}^d}$$

is unitary. It suffices to prove

$$\sum_{v \in \{0, 1, \dots, m-1\}^d} e^{-\frac{2\pi i}{m} \langle u^{(1)} - u^{(2)}, v \rangle} = 0$$

for all  $u^{(1)}, u^{(2)} \in \{0, 1, \dots, m-1\}^d$  with  $u^{(1)} \neq u^{(2)}$ , which is equivalent to

$$\sum_{v \in \{0, 1, \dots, m-1\}^d} e^{\frac{2\pi i}{m} \langle u, v \rangle} = 0$$

for all  $u \in \{-(m-1), \dots, -1, 0, 1, \dots, m-1\}^d \setminus \{0^d\}$ . Let  $u = (u_1, \dots, u_d)$  with  $u_1, \dots, u_d \in \{-(m-1), \dots, -1, 0, 1, \dots, m-1\}$  and  $u_k \neq 0$  for some  $k \in \{1, \dots, d\}$ . Then

$$\begin{aligned} \sum_{v \in \{0, 1, \dots, m-1\}^d} e^{\frac{2\pi i}{m} \langle u, v \rangle} &= \sum_{v_1=0}^{m-1} \sum_{v_2=0}^{m-1} \dots \sum_{v_d=0}^{m-1} e^{\frac{2\pi i}{m} (u_1 v_1 + u_2 v_2 + \dots + u_d v_d)} \\ &= \left( \sum_{s=0}^{m-1} e^{s u_1 \cdot \frac{2\pi i}{m}} \right) \left( \sum_{s=0}^{m-1} e^{s u_2 \cdot \frac{2\pi i}{m}} \right) \dots \left( \sum_{s=0}^{m-1} e^{s u_d \cdot \frac{2\pi i}{m}} \right). \end{aligned}$$

We only need to prove  $\sum_{s=0}^{m-1} e^{s u_k \cdot \frac{2\pi i}{m}} = 0$ . It suffices to prove  $\sum_{s=0}^{m-1} e^{s t \cdot \frac{2\pi i}{m}} = 0$  for all  $t \in \{1, 2, \dots, m-1\}$ . Let  $t \in \{1, 2, \dots, m-1\}$  and write  $t = t'r$ ,  $m = m'r$  with  $r, t', m' \in \mathbb{N}$  such that  $t'$  and  $m'$  are coprime. Then

$$\begin{aligned} \sum_{s=0}^{m-1} e^{s t \cdot \frac{2\pi i}{m}} &= \sum_{s=0}^{m'r-1} e^{s t' \cdot \frac{2\pi i}{m'}} = \sum_{n=0}^{r-1} \sum_{s=nm'}^{(n+1)m'-1} e^{s t' \cdot \frac{2\pi i}{m'}} \\ &= \sum_{n=0}^{r-1} \sum_{s=0}^{m'-1} e^{(nm'+s)t' \cdot \frac{2\pi i}{m'}} = \sum_{n=0}^{r-1} \sum_{s=0}^{m'-1} e^{s t' \cdot \frac{2\pi i}{m'}} = r \sum_{s=0}^{m'-1} e^{s t' \cdot \frac{2\pi i}{m'}}. \end{aligned}$$

We only need to prove  $\sum_{s=0}^{m'-1} e^{s t' \cdot \frac{2\pi i}{m'}} = 0$ . Since  $t'$  and  $m'$  are coprime, we get

$$\{0, t', 2t', \dots, (m'-1)t'\} \equiv \{0, 1, 2, \dots, m'-1\} \pmod{m'}.$$

Thus

$$\sum_{s=0}^{m'-1} e^{s t' \cdot \frac{2\pi i}{m'}} = \sum_{s=0}^{m'-1} e^{s \cdot \frac{2\pi i}{m'}} \stackrel{(\star)}{=} 0,$$

where in  $(\star)$  we use  $m' \geq 2$  since  $m > t$  implies  $m' > t' \geq 1$ .  $\square$

**Proposition 3.3.** *Let  $d \in \mathbb{N}$ ,  $R$  be a  $d \times d$  real matrix and  $C > 0$ . If  $[-C, C]^d \subseteq R[-1, 1]^d$ , then all eigenvalues of  $R$  have modulus no less than  $C$ .*

*Proof.* Let  $\lambda \in \mathbb{C}$  be an eigenvalue of  $R$ . Then there exists

$$z \in ([-1, 1] + i[-1, 1])^d \setminus \{0\} \tag{3.6}$$

such that  $Rz = \lambda z$ . We need to prove  $|\lambda| \geq C$ . Since  $[-C, C]^d \subseteq R[-1, 1]^d$  implies that  $R$  is invertible, we get  $CR^{-1}[-1, 1]^d \subseteq [-1, 1]^d$  and then

$$CR^{-1}([-1, 1] + i[-1, 1])^d \subseteq ([-1, 1] + i[-1, 1])^d.$$

It follows from (3.6) that

$$(CR^{-1})^n z \in ([-1, 1] + i[-1, 1])^d$$

for all  $n \in \mathbb{N}$ . Since  $Rz = \lambda z$  implies  $R^n z = \lambda^n z$ , we get

$$C^n z = \lambda^n (CR^{-1})^n z \in \lambda^n ([-1, 1] + i[-1, 1])^d$$

and then

$$C^n |z| \leq |\lambda|^n \sqrt{2d} \quad \text{for all } n \in \mathbb{N}.$$

By  $|z| \neq 0$ , we must have  $C \leq \lambda$ . □

**Proposition 3.4.** *Let  $d \in \mathbb{N}$ ,  $P$  be a  $d \times d$  real matrix and  $c \geq 0$ . If  $P[-1, 1]^d \subseteq [-c, c]^d$ , then  $P^T[-1, 1]^d \subseteq [-cd, cd]^d$ .*

*Proof.* Suppose  $P[-1, 1]^d \subseteq [-c, c]^d$  and write

$$P = \begin{pmatrix} p_{11} & \cdots & p_{1d} \\ \vdots & & \vdots \\ p_{d1} & \cdots & p_{dd} \end{pmatrix}.$$

For all  $s, t \in \{1, 2, \dots, d\}$ , by  $P(1, \dots, 1)^T \in [-c, c]^d$  we get

$$-c \leq p_{s1} + \cdots + p_{s(t-1)} + p_{st} + p_{s(t+1)} + \cdots + p_{sd} \leq c,$$

and by  $P(1, \dots, 1, -1, 1, \dots, 1)^T \in [-c, c]^d$  we get

$$-c \leq p_{s1} + \cdots + p_{s(t-1)} - p_{st} + p_{s(t+1)} + \cdots + p_{sd} \leq c.$$

Therefore

$$-c \leq p_{st} \leq c \quad \text{for all } s, t \in \{1, 2, \dots, d\}.$$

It follows that for all  $\tau_1, \dots, \tau_d \in \{1, -1\}$  we have

$$P^T \begin{pmatrix} \tau_1 \\ \vdots \\ \tau_d \end{pmatrix} \in [-dc, dc]^d.$$

By the linearity of  $P^T$  we get  $P^T[-1, 1]^d \subseteq [-cd, cd]^d$ . □

Now we prove Theorem 1.1 to end this section.

*Proof of Theorem 1.1.* (1) Prove that  $\mu = \delta_{R_1^{-1}B_1} * \delta_{R_1^{-1}R_2^{-1}B_2} * \delta_{R_1^{-1}R_2^{-1}R_3^{-1}B_3} * \cdots$  exists.

By Theorem 2.1, we only need to prove

$$\sum_{k=1}^{\infty} \frac{1}{\#B_k} \sum_{a \in R_1^{-1} \cdots R_k^{-1} B_k} \frac{|a|}{1 + |a|} < \infty, \quad \text{i.e.,} \quad \sum_{k=1}^{\infty} \frac{1}{m_k^d} \sum_{x \in B_k} \frac{|R_1^{-1} \cdots R_k^{-1} x|}{1 + |R_1^{-1} \cdots R_k^{-1} x|} < \infty.$$

Divide  $B_k$  into two parts

$$B_{k,1} := B_k \cap \{0, 1, \dots, m_k - 1\}^d \quad \text{and} \quad B_{k,2} := B_k \setminus \{0, 1, \dots, m_k - 1\}^d.$$

Since

$$\sum_{k=1}^{\infty} \frac{1}{m_k^d} \sum_{x \in B_{k,2}} \frac{|R_1^{-1} \cdots R_k^{-1} x|}{1 + |R_1^{-1} \cdots R_k^{-1} x|} \leq \sum_{k=1}^{\infty} \frac{\#B_{k,2}}{m_k^d} \stackrel{\text{by (1.3)}}{<} \infty,$$

it suffices to show

$$\sum_{k=1}^{\infty} \frac{1}{m_k^d} \sum_{x \in B_{k,1}} \frac{|R_1^{-1} \cdots R_k^{-1} x|}{1 + |R_1^{-1} \cdots R_k^{-1} x|} < \infty$$

in the following. In fact we have

$$\begin{aligned} \sum_{k=1}^{\infty} \frac{1}{m_k^d} \sum_{x \in B_{k,1}} \frac{|R_1^{-1} \cdots R_k^{-1} x|}{1 + |R_1^{-1} \cdots R_k^{-1} x|} &\leq \sum_{k=1}^{\infty} \frac{1}{m_k^d} \sum_{x \in \{0, \dots, m_k - 1\}^d} |R_1^{-1} \cdots R_k^{-1} x| \\ &\stackrel{(*)}{\leq} \sum_{k=1}^{\infty} \frac{1}{m_k^d} \cdot m_k^d \cdot \frac{d\sqrt{d}}{m_1 \cdots m_{k-1}} \leq \sum_{k=1}^{\infty} \frac{d\sqrt{d}}{2^{k-1}} = 2d\sqrt{d} < \infty, \end{aligned}$$

where  $(*)$  follows from

$$|R_1^{-1} \cdots R_k^{-1} x| \leq \frac{d\sqrt{d}}{m_1 \cdots m_{k-1}} \quad \text{for all } x \in \{0, \dots, m_k - 1\}^d \text{ and } k \in \mathbb{N},$$

which can be proved as follows. In fact, we only need to prove

$$R_1^{-1} \cdots R_k^{-1} [-m_k, m_k]^d \subseteq \left[ -\frac{d}{m_1 \cdots m_{k-1}}, \frac{d}{m_1 \cdots m_{k-1}} \right]^d \quad \text{for all } k \in \mathbb{N}.$$

By the linearity of  $R_1^{-1} \cdots R_k^{-1}$  and Proposition 3.4, it suffices to show

$$(R_k^{-1})^T \cdots (R_1^{-1})^T [-1, 1]^d \subseteq \left[ -\frac{1}{m_1 \cdots m_k}, \frac{1}{m_1 \cdots m_k} \right]^d \quad \text{for all } k \in \mathbb{N}.$$

This follows immediately from the linearity of  $(R_1^{-1})^T, \dots, (R_k^{-1})^T$  and the fact that the condition  $[-m_k, m_k]^d \subseteq R_k^T [-1, 1]^d$  implies  $(R_k^{-1})^T [-1, 1]^d \subseteq [-\frac{1}{m_k}, \frac{1}{m_k}]^d$  for every  $k \in \mathbb{N}$ .

(2) Prove that  $\mu$  is a spectral measure with a spectrum in  $\mathbb{Z}^d$ .

- ① Let  $\{\nu_{>n}\}_{n \geq 1}$  be given by (2.1). It follows from Lemma 3.1 that there exists  $n_0 \in \mathbb{N}$  such that  $\{\nu_{>n}\}_{n \geq n_0}$  is equi-positive.
- ② For every  $k \in \mathbb{N}$ , prove that  $(R_k, B_k)$  is an admissible pair in  $\mathbb{R}^d$ .  
In fact, by  $[-m_k, m_k]^d \subseteq R_k^T [-1, 1]^d$  and  $m_k \geq 2$ , it follows from Proposition 3.3 that all eigenvalues of  $R_k^T$  have modulus no less than 2. Noting that  $R_k$  and  $R_k^T$  have the same eigenvalues,  $R_k$  must be expanding. Since  $B_k \equiv \{0, 1, \dots, m_k - 1\}^d \pmod{R_k \mathbb{Z}^d}$ , by Proposition 3.2 we know that  $(R_k, B_k)$  is an admissible pair in  $\mathbb{R}^d$ .
- ③ We have  $|(R_n^T)^{-1} \cdots (R_1^T)^{-1} x| \rightarrow 0$  as  $n \rightarrow \infty$  for all  $x \in \mathbb{R}^d$ , since  $[-m_k, m_k]^d \subseteq R_k^T [-1, 1]^d$  for all  $k \in \mathbb{N}$  imply  $(R_n^T)^{-1} \cdots (R_1^T)^{-1} [-1, 1]^d \subseteq \left[ -\frac{1}{m_1 \cdots m_n}, \frac{1}{m_1 \cdots m_n} \right]^d \subseteq \left[ -\frac{1}{2^n}, \frac{1}{2^n} \right]^d$  for all  $n \in \mathbb{N}$ .

Combining ①, ② and ③, by Theorem 2.3 we know that  $\mu$  is a spectral measure with a spectrum in  $\mathbb{Z}^d$ .  $\square$

#### 4. PROOF OF THEOREM 1.2 AND AN EXAMPLE FOR REMARK 1.4

Recall that for  $A_1, A_2, \dots \subseteq \mathbb{R}^d$ ,

$$\sum_{k=1}^{\infty} A_k := \left\{ x \in \mathbb{R}^d : \exists a_k \in A_k \text{ for each } k \in \mathbb{N} \text{ s.t. } x = \sum_{k=1}^{\infty} a_k \right\},$$

and for  $B_1, B_2, \dots \subseteq \mathbb{R}^d$ ,

$$\lim_{n \rightarrow \infty} B_n := \left\{ x \in \mathbb{R}^d : \exists b_n \in B_n \text{ for each } n \in \mathbb{N} \text{ s.t. } x = \lim_{n \rightarrow \infty} b_n \right\}.$$

Before proving Theorem 1.2, we give the following first.

**Proposition 4.1.** *Let  $A_1, A_2, \dots \subseteq \mathbb{R}^d$ .*

(1) *We have*

$$\lim_{n \rightarrow \infty} (A_1 + \cdots + A_n) \supseteq \overline{\sum_{k=1}^{\infty} A_k}.$$

(2) If  $\sum_{k=1}^{\infty} A_k \neq \emptyset$ , then

$$\lim_{n \rightarrow \infty} (A_1 + \cdots + A_n) = \overline{\sum_{k=1}^{\infty} A_k}.$$

*Proof.* (1) Since  $\sum_{k=1}^{\infty} A_k \subseteq \lim_{n \rightarrow \infty} (A_1 + \cdots + A_n)$  is obvious, it suffices to prove that  $\lim_{n \rightarrow \infty} (A_1 + \cdots + A_n)$  is closed. In fact, for any  $B_1, B_2, \dots \subseteq \mathbb{R}^d$ , we can prove that  $\lim_{n \rightarrow \infty} B_n$  is a closed set. Let  $x_1, x_2, \dots \in \lim_{n \rightarrow \infty} B_n$  and  $x \in \mathbb{R}^d$  such that  $\lim_{k \rightarrow \infty} x_k = x$ . It suffices to prove  $x \in \lim_{n \rightarrow \infty} B_n$ . For each  $k \in \mathbb{N}$ , by  $x_k \in \lim_{n \rightarrow \infty} B_n$ , there exist  $b_{k,1} \in B_1, b_{k,2} \in B_2, \dots$  such that  $\lim_{n \rightarrow \infty} b_{k,n} = x_k$ . By  $\lim_{n \rightarrow \infty} b_{1,n} = x_1$ , there exists  $N_1 \geq 1$  such that for all  $n \geq N_1$  we have  $|b_{1,n} - x_1| < 1$ . By  $\lim_{n \rightarrow \infty} b_{2,n} = x_2$ , there exists  $N_2 > N_1$  such that for all  $n \geq N_2$  we have  $|b_{2,n} - x_2| < \frac{1}{2}$ .  $\dots$  Repeating this process, we can find  $1 \leq N_1 < N_2 < N_3 < \dots$  such that

$$\text{for all } k \in \mathbb{N} \text{ and } n \geq N_k \text{ we have } |b_{k,n} - x_k| < \frac{1}{k}. \quad (4.1)$$

For each  $k \in \mathbb{N}$  and  $n \in \{N_k, N_k + 1, \dots, N_{k+1} - 1\}$ , we define  $c_n := b_{k,n} \in B_n$ . It suffices to prove  $\lim_{n \rightarrow \infty} c_n = x$ . Arbitrarily take  $\varepsilon > 0$ . By  $\lim_{k \rightarrow \infty} x_k = x$ , there exists  $K > \frac{2}{\varepsilon}$  such that

$$\text{for all } k \geq K \text{ we have } |x_k - x| < \frac{\varepsilon}{2}. \quad (4.2)$$

For each  $n \geq N_K$ , there exists  $k \geq K$  such that  $N_k \leq n < N_{k+1}$  and then

$$|c_n - x_k| \stackrel{\text{by (4.1)}}{<} \frac{1}{k} \leq \frac{1}{K} < \frac{\varepsilon}{2}, \quad (4.3)$$

which implies

$$|c_n - x| \leq |c_n - x_k| + |x_k - x| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon,$$

where the second inequality follows from (4.2) and (4.3). Therefore  $\lim_{n \rightarrow \infty} c_n = x$ .

(2) Suppose  $\sum_{k=1}^{\infty} A_k \neq \emptyset$ . By (1) we only need to prove  $\lim_{n \rightarrow \infty} (A_1 + \cdots + A_n) \subseteq \overline{\sum_{k=1}^{\infty} A_k}$ . Let  $x \in \lim_{n \rightarrow \infty} (A_1 + \cdots + A_n)$ . Then there exists  $b_n \in A_1 + \cdots + A_n$  for each  $n \in \mathbb{N}$  such that  $x = \lim_{n \rightarrow \infty} b_n$ . By  $\sum_{k=1}^{\infty} A_k \neq \emptyset$ , there exist  $a_1 \in A_1, a_2 \in A_2, \dots$  such that  $\sum_{k=1}^{\infty} a_k$  converges, which implies  $\sum_{k=n+1}^{\infty} a_k \rightarrow \mathbf{0}$  as  $n \rightarrow \infty$ , where  $\mathbf{0}$  denotes the zero vector in  $\mathbb{R}^d$ . For all  $n \in \mathbb{N}$ , define

$$x_n := b_n + \sum_{k=n+1}^{\infty} a_k \in \sum_{k=1}^{\infty} A_k.$$

Then  $\lim_{n \rightarrow \infty} x_n = x$  and we get  $x \in \overline{\sum_{k=1}^{\infty} A_k}$ .  $\square$

In Proposition 4.1 (2), the condition  $\sum_{k=1}^{\infty} A_k \neq \emptyset$  can not be omitted. For example, take  $d = 1, A_1 = \{-1, 0, 1\}$  and  $A_k = \{-1, 1\}$  for all  $k \geq 2$ . Then  $\sum_{k=1}^{\infty} A_k = \emptyset$ , and

$$\lim_{n \rightarrow \infty} (A_1 + \cdots + A_n) = \lim_{n \rightarrow \infty} \{-n, \dots, -1, 0, 1, \dots, n\} = \mathbb{Z} \neq \overline{\sum_{k=1}^{\infty} A_k}.$$

Now we use Theorem 2.4, Proposition 4.1 and probability theory to prove Theorem 1.2.

*Proof of Theorem 1.2.* Let  $d \in \mathbb{N}$  and  $\mu_1, \mu_2, \dots \in \mathcal{P}(\mathbb{R}^d)$  such that  $\mu_1 * \mu_2 * \dots$  exists. It follows from Theorem 2.4 that  $\text{spt}(\mu_1 * \mu_2 * \dots) = \lim_{n \rightarrow \infty} (\text{spt} \mu_1 + \cdots + \text{spt} \mu_n)$ . By Proposition 4.1 (2), we only need to prove  $\sum_{k=1}^{\infty} \text{spt} \mu_k \neq \emptyset$ . It follows from the argument above Theorem 3.1 in [29] that there exist a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  and a sequence of independent random vectors  $\{X_k\}_{k \geq 1}$  such that for each  $k \geq 1$  the distribution of  $X_k$  is  $\mu_k = \mathbb{P} \circ X_k^{-1}$ , and

$$\mathbb{P} \left( \sum_{k=1}^{\infty} X_k \text{ converges} \right) = 1. \quad (4.4)$$

By  $\mathbb{P}(X_k \in \text{spt } \mu_k) = \mu_k(\text{spt } \mu_k) = 1$  for every  $k \in \mathbb{N}$  and (4.4), there exists  $w \in \Omega$  such that  $X_k(w) \in \text{spt } \mu_k$  for all  $k \in \mathbb{N}$  and  $\sum_{k=1}^{\infty} X_k(w)$  converges. Therefore  $\sum_{k=1}^{\infty} \text{spt } \mu_k \neq \emptyset$ .  $\square$

To end this section, we give an example for Remark 1.4. We construct non-empty finite sets  $A_1, A_2, \dots \subseteq [0, 1]$  such that  $\delta_{A_1} * \delta_{A_2} * \dots$  exists but  $\sum_{k=1}^{\infty} A_k$  is not closed in the following.

Let  $\frac{1}{\mathbb{N}} := \left\{1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots\right\}$  and use  $\lfloor x \rfloor$  to denote the greatest integer no larger than  $x$ . Define  $r_1 := \frac{1}{2}$  and take  $a_1 \in (0, r_1) \setminus \frac{1}{\mathbb{N}}$ .

Define  $r_2 := \min \left\{ \frac{1}{2^2}, \frac{1}{\lfloor \frac{1}{a_1} \rfloor} - a_1 \right\} > 0$  and take  $a_2 \in (0, r_2) \setminus \frac{1}{\mathbb{N}}$ .

Define  $r_3 := \min \left\{ \frac{1}{2^3}, \frac{1}{\lfloor \frac{1}{a_1} \rfloor} - a_1 - a_2, \frac{1}{\lfloor \frac{1}{a_2} \rfloor} - a_2 \right\} > 0$  and take  $a_3 \in (0, r_3) \setminus \frac{1}{\mathbb{N}}$ .

...

Define  $r_k := \min \left\{ \frac{1}{2^k}, \frac{1}{\lfloor \frac{1}{a_1} \rfloor} - a_1 - a_2 - \dots - a_{k-1}, \frac{1}{\lfloor \frac{1}{a_2} \rfloor} - a_2 - \dots - a_{k-1}, \dots, \frac{1}{\lfloor \frac{1}{a_{k-1}} \rfloor} - a_{k-1} \right\} > 0$  and take  $a_k \in (0, r_k) \setminus \frac{1}{\mathbb{N}}$ .

...

For all  $k \in \mathbb{N}$ , define  $t_k := a_k + a_{k+1} + a_{k+2} + \dots < \sum_{n=k}^{\infty} \frac{1}{2^n} = \frac{1}{2^{k-1}}$ .

First we prove

$$a_k > \frac{1}{\lfloor \frac{1}{t_k} \rfloor + 1} \quad \text{for all } k \in \mathbb{N}. \quad (4.5)$$

Arbitrarily take  $k \in \mathbb{N}$ . It suffices to show  $\lfloor \frac{1}{t_k} \rfloor + 1 > \frac{1}{a_k}$ . Since  $\lfloor \frac{1}{a_k} \rfloor + 1 > \frac{1}{a_k}$ , we only need to prove  $\lfloor \frac{1}{t_k} \rfloor \geq \lfloor \frac{1}{a_k} \rfloor$ . It suffices to show  $\frac{1}{t_k} \geq \lfloor \frac{1}{a_k} \rfloor$ . For all  $n \in \mathbb{N}$ , by

$$a_{k+n} < r_{k+n} \leq \frac{1}{\lfloor \frac{1}{a_k} \rfloor} - a_k - a_{k+1} - \dots - a_{k+n-1}$$

we get

$$a_k + a_{k+1} + \dots + a_{k+n} < \frac{1}{\lfloor \frac{1}{a_k} \rfloor}.$$

As  $n \rightarrow \infty$ , it follows that  $t_k \leq \frac{1}{\lfloor \frac{1}{a_k} \rfloor}$ , i.e.,  $\frac{1}{t_k} \geq \lfloor \frac{1}{a_k} \rfloor$ . Therefore (4.5) holds.

For all  $k \in \mathbb{N}$ , by (4.5) we can take different

$$b_{k,1}, b_{k,2}, \dots, b_{k,k^2} \in \left( \frac{1}{\lfloor \frac{1}{t_k} \rfloor + 1}, a_k \right) \setminus \frac{1}{\mathbb{N}}$$

and define

$$A_k := \left\{ 0, b_{k,1}, b_{k,2}, \dots, b_{k,k^2}, a_k, \frac{k}{k+1} \right\} \subseteq [0, 1].$$

The fact that  $\delta_{A_1} * \delta_{A_2} * \dots$  exists follows from Theorem 2.1 and

$$\begin{aligned} \sum_{k=1}^{\infty} \frac{1}{\#A_k} \sum_{a \in A_k} \frac{a}{1+a} &< \sum_{k=1}^{\infty} \frac{1}{k^2+3} \sum_{a \in A_k} a < \sum_{k=1}^{\infty} \frac{1}{k^2+3} \left( (k^2+1)a_k + \frac{k}{k+1} \right) \\ &< \sum_{k=1}^{\infty} \left( \frac{k^2+1}{k^2+3} \cdot \frac{1}{2^k} + \frac{k}{(k^2+3)(k+1)} \right) < \sum_{k=1}^{\infty} \left( \frac{1}{2^k} + \frac{1}{k^2} \right) < \infty. \end{aligned}$$

In the following we only need to prove that  $\sum_{k=1}^{\infty} A_k$  is not closed. By  $\{0, \frac{n}{n+1}\} \subseteq A_n$  for all  $n \in \mathbb{N}$ , one can easily verify  $\frac{n}{n+1} \in \sum_{k=1}^{\infty} A_k$  for all  $n \in \mathbb{N}$ . Thus  $1 = \lim_{n \rightarrow \infty} \frac{n}{n+1} \in \overline{\sum_{k=1}^{\infty} A_k}$ . It suffices to show  $1 \notin \sum_{k=1}^{\infty} A_k$  in the following. By contradiction, we assume  $1 \in \sum_{k=1}^{\infty} A_k$ . Then there exist  $x_1 \in A_1, x_2 \in A_2, \dots$  such that  $\sum_{k=1}^{\infty} x_k = 1$ . If  $x_k \leq a_k$  for all  $k \in \mathbb{N}$ , then  $\sum_{k=1}^{\infty} x_k \leq \sum_{k=1}^{\infty} a_k < \sum_{k=1}^{\infty} r_k \leq \sum_{k=1}^{\infty} \frac{1}{2^k} = 1$ , which contradicts



$\sum_{k=1}^{\infty} x_k = 1$ . Thus there must exist  $n \in \mathbb{N}$  such that  $x_n > a_n$ . By  $x_n \in A_n$  we know  $x_n = \frac{n}{n+1}$ . Thus

$$x_1 + \cdots + x_{n-1} + \frac{n}{n+1} + x_{n+1} + x_{n+2} + \cdots = 1. \quad (4.6)$$

If there exists  $n' \in \mathbb{N} \setminus \{n\}$  such that  $x_{n'} > a_{n'}$ , by  $x_{n'} \in A_{n'}$  we know  $x_{n'} = \frac{n'}{n'+1}$  and then  $\sum_{k=1}^{\infty} x_k \geq \frac{n}{n+1} + \frac{n'}{n'+1} > 1$ , which contradicts  $\sum_{k=1}^{\infty} x_k = 1$ . Thus  $x_k \leq a_k$  for all  $k \in \mathbb{N} \setminus \{n\}$ . If  $x_1 = x_2 = \cdots = x_{n-1} = 0$ , then

$$\begin{aligned} x_1 + \cdots + x_{n-1} + \frac{n}{n+1} + x_{n+1} + x_{n+2} + \cdots &\leq \frac{n}{n+1} + a_{n+1} + a_{n+2} + \cdots \\ &< \frac{n}{n+1} + \frac{1}{2^{n+1}} + \frac{1}{2^{n+2}} + \cdots = \frac{n}{n+1} + \frac{1}{2^n} \leq 1, \end{aligned}$$

which contradicts (4.6). Thus there must exist  $p \in \{1, 2, \dots, n-1\}$  such that  $x_1 = \cdots = x_{p-1} = 0$  and  $x_p > 0$ . We get

$$t_p = a_p + a_{p+1} + a_{p+2} + \cdots > x_1 + \cdots + x_{n-1} + x_{n+1} + x_{n+2} + \cdots \stackrel{\text{by (4.6)}}{=} \frac{1}{n+1},$$

which implies

$$n+1 > \frac{1}{t_p} \geq \lfloor \frac{1}{t_p} \rfloor. \quad (4.7)$$

Since  $x_p > 0$  and  $x_p \in A_p$  imply  $x_p > \frac{1}{\lfloor \frac{1}{t_p} \rfloor + 1}$ , we get  $\lfloor \frac{1}{t_p} \rfloor + 1 > \frac{1}{x_p} \geq \lfloor \frac{1}{x_p} \rfloor$  and then  $\lfloor \frac{1}{t_p} \rfloor \geq \lfloor \frac{1}{x_p} \rfloor$ . It follows from (4.7) that  $n+1 > \lfloor \frac{1}{x_p} \rfloor$ , which implies  $n+1 \geq \lfloor \frac{1}{x_p} \rfloor + 1 > \frac{1}{x_p}$ . Therefore  $x_p > \frac{1}{n+1}$  and then  $x_1 + \cdots + x_{n-1} + \frac{n}{n+1} + x_{n+1} + x_{n+2} + \cdots > \frac{1}{n+1} + \frac{n}{n+1} = 1$ . This contradicts (4.6).

## 5. PROOFS OF THEOREM 1.5 AND COROLLARY 1.6

First we prove statement (1) in Theorem 1.5.

*Proof of Theorem 1.5 (1).* Suppose  $\lim_{k \rightarrow \infty} \min_{a \in A'_k} |a| > 0$ . Since the proofs of ① and ② are similar, we only prove ① in the following. By the definition of Hausdorff dimension, it suffices to show that

$$\mathcal{H}^s \left( \sum_{k=1}^{\infty} (A_k \cup A'_k) \right) = 0 \text{ if and only if } \mathcal{H}^s \left( \sum_{k=1}^{\infty} A_k \right) = 0 \text{ for all } s \in [0, d].$$

$\Rightarrow$  follows immediately from  $\sum_{k=1}^{\infty} (A_k \cup A'_k) \supseteq \sum_{k=1}^{\infty} A_k$ .

$\Leftarrow$  Suppose  $\mathcal{H}^s(\sum_{k=1}^{\infty} A_k) = 0$  for  $s \in [0, d]$ . For all  $p \in \mathbb{N}$ , by

$$\sum_{k=1}^{\infty} A_k = \bigcup_{x \in \sum_{k=1}^p A_k} \left( x + \sum_{k=p+1}^{\infty} A_k \right)$$

and

$$\sum_{k=1}^p (A_k \cup A'_k) + \sum_{k=p+1}^{\infty} A_k = \bigcup_{x \in \sum_{k=1}^p (A_k \cup A'_k)} \left( x + \sum_{k=p+1}^{\infty} A_k \right),$$

where  $\sum_{k=1}^p A_k$  and  $\sum_{k=1}^p (A_k \cup A'_k)$  are finite sets, we get  $\mathcal{H}^s(\sum_{k=1}^p (A_k \cup A'_k) + \sum_{k=p+1}^{\infty} A_k) = 0$ . In order to show  $\mathcal{H}^s(\sum_{k=1}^{\infty} (A_k \cup A'_k)) = 0$ , it suffices to prove

$$\sum_{k=1}^{\infty} (A_k \cup A'_k) = \bigcup_{p=1}^{\infty} \left( \sum_{k=1}^p (A_k \cup A'_k) + \sum_{k=p+1}^{\infty} A_k \right). \quad (5.1)$$

Since the inclusion " $\supseteq$ " is obvious, we only prove " $\subseteq$ " in the following. Let  $x \in \sum_{k=1}^{\infty} (A_k \cup A'_k)$ . Then there exist  $x^{(1)} \in A_1 \cup A'_1, x^{(2)} \in A_2 \cup A'_2, \dots$  such that  $x = \sum_{k=1}^{\infty} x^{(k)}$  converges,

which implies  $\lim_{k \rightarrow \infty} |x^{(k)}| = 0$ . Let  $c \in (0, \underline{\lim}_{k \rightarrow \infty} \min_{a \in A'_k} |a|)$ . Then there exists  $p_1 \in \mathbb{N}$  such that for all  $k > p_1$  we have  $\min_{a \in A'_k} |a| > c$ . By  $\lim_{k \rightarrow \infty} |x^{(k)}| = 0$ , there exists  $p_2 \in \mathbb{N}$  such that for all  $k > p_2$  we have  $|x^{(k)}| < c$ . Let  $p_0 := \max\{p_1, p_2\}$ . Then for all  $k \geq p_0 + 1$  we have  $|x^{(k)}| < \min_{a \in A'_k} |a|$ , which implies  $x^{(k)} \notin A'_k$ . It follows from  $x^{(k)} \in A_k \cup A'_k$  that  $x^{(k)} \in A_k$  for all  $k \geq p_0 + 1$ . Thus

$$x = \sum_{k=1}^{p_0} x^{(k)} + \sum_{k=p_0+1}^{\infty} x^{(k)} \in \sum_{k=1}^{p_0} (A_k \cup A'_k) + \sum_{k=p_0+1}^{\infty} A_k.$$

We get  $x \in \bigcup_{p=1}^{\infty} (\sum_{k=1}^p (A_k \cup A'_k) + \sum_{k=p+1}^{\infty} A_k)$ .  $\square$

Before proving statement (2) in Theorem 1.5, we give the following first.

**Proposition 5.1.** *Let  $A_1, A_2, \dots \subseteq \mathbb{R}^d$  be non-empty finite sets.*

(1) *If  $\sum_{k=1}^{\infty} \max_{a \in A_k} |a| < \infty$ , then  $\sum_{k=1}^{\infty} A_k$  is non-empty and compact.*

(2) *If  $A_k \subseteq [0, \infty)^d$  for all  $k \in \mathbb{N}$ ,  $\sum_{k=1}^{\infty} A_k$  is non-empty and bounded, then  $\sum_{k=1}^{\infty} \max_{a \in A_k} |a| < \infty$ .*

*Proof.* (1) Suppose  $\sum_{k=1}^{\infty} \max_{a \in A_k} |a| < \infty$ .

① Prove  $\sum_{k=1}^{\infty} A_k \neq \emptyset$ .

We arbitrarily take  $a^{(1)} \in A_1, a^{(2)} \in A_2, \dots$ . It follows from

$$\sum_{k=1}^{\infty} |a^{(k)}| \leq \sum_{k=1}^{\infty} \max_{a \in A_k} |a| < \infty$$

that  $\sum_{k=1}^{\infty} a^{(k)}$  converges, and then  $\sum_{k=1}^{\infty} a^{(k)} \in \sum_{k=1}^{\infty} A_k$ .

② Prove that  $\sum_{k=1}^{\infty} A_k$  is bounded.

This follows from the fact that  $\sum_{k=1}^{\infty} A_k$  is contained in the closed ball  $B(\mathbf{0}, M)$  centered at the original point  $\mathbf{0}$  with radius  $M := \sum_{k=1}^{\infty} \max_{a \in A_k} |a|$ .

③ Prove that  $\sum_{k=1}^{\infty} A_k$  is closed.

Let  $x^{(1)}, x^{(2)}, \dots \in \sum_{k=1}^{\infty} A_k$  such that  $x^{(n)}$  converges to some  $x \in \mathbb{R}^d$  as  $n \rightarrow \infty$ . We need to prove  $x \in \sum_{k=1}^{\infty} A_k$ . For all  $n \in \mathbb{N}$ , by  $x^{(n)} \in \sum_{k=1}^{\infty} A_k$ , there exist  $x_1^{(n)} \in A_1, x_2^{(n)} \in A_2, \dots$  such that  $x^{(n)} = \sum_{k=1}^{\infty} x_k^{(n)}$  converges. Since  $A_1, A_2, A_3, \dots$  are all finite sets, we can choose positive integers  $n_1 < n_2 < n_3 < \dots$  such that  $x_1^{(n_1)} = x_1^{(n_2)} = x_1^{(n_3)} = \dots = a_1$  for some  $a_1 \in A_1, x_2^{(n_2)} = x_2^{(n_3)} = x_2^{(n_4)} = \dots = a_2$  for some  $a_2 \in A_2, x_3^{(n_3)} = x_3^{(n_4)} = x_3^{(n_5)} = \dots = a_3$  for some  $a_3 \in A_3, \dots$ . Then we can write

$$x^{(n_p)} = \sum_{k=1}^p a_k + \sum_{k=p+1}^{\infty} x_k^{(n_p)}$$

for all  $p \in \mathbb{N}$ . Since  $\sum_{k=1}^{\infty} \max_{a \in A_k} |a| < \infty$  implies that  $\sum_{k=1}^{\infty} a_k$  converges and  $\sum_{k=p+1}^{\infty} x_k^{(n_p)} \rightarrow 0$  as  $p \rightarrow \infty$ , we get  $x^{(n_p)} \rightarrow \sum_{k=1}^{\infty} a_k$ . Therefore  $x = \sum_{k=1}^{\infty} a_k \in \sum_{k=1}^{\infty} A_k$ .

(2) Suppose  $A_k \subseteq [0, \infty)^d$  for all  $k \in \mathbb{N}$  and let  $M > 0$  such that  $\emptyset \neq \sum_{k=1}^{\infty} A_k \subseteq [0, M]^d$ . Recall the notation  $\|x\| := \max\{|x_1|, \dots, |x_d|\}$  for all  $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ . By  $|x| \leq \sqrt{d}\|x\|$  for all  $x \in \mathbb{R}^d$ , we only need to prove  $\sum_{k=1}^{\infty} \max_{a \in A_k} \|a\| < \infty$  by contradiction. Assume  $\sum_{k=1}^{\infty} \max_{a \in A_k} \|a\| = \infty$ . For all  $k \in \mathbb{N}$ , let  $x^{(k)} = (x_1^{(k)}, \dots, x_d^{(k)}) \in A_k$  and  $j_k \in \{1, \dots, d\}$  such that  $x_{j_k}^{(k)} = \max_{a \in A_k} \|a\|$ . We get  $\sum_{k=1}^{\infty} x_{j_k}^{(k)} = \infty$ . Let  $m \in \mathbb{N}$  such that  $\sum_{k=1}^m x_{j_k}^{(k)} > dM$ . Then  $\sum_{k=1}^m x_1^{(k)} + \dots + \sum_{k=1}^m x_d^{(k)} > dM$ . There must exist  $t \in \{1, \dots, d\}$  such that  $\sum_{k=1}^m x_t^{(k)} > M$ .

On the other hand, by  $\sum_{k=1}^{\infty} A_k \neq \emptyset$ , we can take  $a^{(1)} = (a_1^{(1)}, \dots, a_d^{(1)}) \in A_1$ ,  $a^{(2)} = (a_1^{(2)}, \dots, a_d^{(2)}) \in A_2, \dots$  such that  $\sum_{k=1}^{\infty} a^{(k)}$  converges, which implies that  $\sum_{k=m+1}^{\infty} a^{(k)}$  converges. Define

$$x = (x_1, \dots, x_d) := \sum_{k=1}^m x^{(k)} + \sum_{k=m+1}^{\infty} a^{(k)}.$$

Then  $x \in \sum_{k=1}^{\infty} A_k$ . But  $x_t = \sum_{k=1}^m x_t^{(k)} + \sum_{k=m+1}^{\infty} a_t^{(k)} > M$  contradicts  $\sum_{k=1}^{\infty} A_k \subseteq [0, M]^d$ .  $\square$

*Remark 5.2.* In Proposition 5.1 (2), without the condition  $A_k \subseteq [0, \infty)^d$  for all  $k \in \mathbb{N}$ , even if  $\sum_{k=1}^{\infty} A_k$  is non-empty and compact, we can not guarantee  $\sum_{k=1}^{\infty} \max_{a \in A_k} |a| < \infty$ . For example, take  $d = 1$  and let  $A_k$  be the singleton  $\{(-1)^{k+1}/k\}$  for every  $k \in \mathbb{N}$ . Then  $\sum_{k=1}^{\infty} A_k$  is the singleton  $\{\sum_{k=1}^{\infty} (-1)^{k+1}/k = \ln 2\}$ , which is non-empty and compact. But  $\sum_{k=1}^{\infty} \max_{a \in A_k} |a| = \sum_{k=1}^{\infty} 1/k = \infty$ .

Now we prove statement (2) in Theorem 1.5.

*Proof of Theorem 1.5 (2).* Suppose  $\sum_{k=1}^{\infty} \max_{a \in A_k} |a| < \infty$ ,  $A'_k \subseteq [0, \infty)^d$  for all  $k \in \mathbb{N}$  and  $\lim_{k \rightarrow \infty} \min_{a \in A'_k} |a| = \infty$ . We need to prove that  $\sum_{k=1}^{\infty} (A_k \cup A'_k)$  is closed. Let  $x^{(1)}, x^{(2)}, \dots \in \sum_{k=1}^{\infty} (A_k \cup A'_k)$  such that  $x^{(n)}$  converges to some  $x \in \mathbb{R}^d$  as  $n \rightarrow \infty$ . It suffices to show  $x \in \sum_{k=1}^{\infty} (A_k \cup A'_k)$ . Since in the proof of Theorem 1.5 (1) we have proved (5.1), in the following we only need to show  $x \in \sum_{k=1}^p (A_k \cup A'_k) + \sum_{k=p+1}^{\infty} A_k$  for some  $p \in \mathbb{N}$ . Since  $\sum_{k=p+1}^{\infty} \max_{a \in A_k} |a| < \infty$  and Proposition 5.1 (1) imply that  $\sum_{k=1}^p (A_k \cup A'_k) + \sum_{k=p+1}^{\infty} A_k$  is closed for every  $p \in \mathbb{N}$ , it suffices to prove  $\{x^{(n)}\}_{n \geq 1} \subseteq \sum_{k=1}^p (A_k \cup A'_k) + \sum_{k=p+1}^{\infty} A_k$  for some  $p \in \mathbb{N}$ .

In fact, by the convergency of  $\{x^{(n)}\}_{n \geq 1}$ , there exists  $M_1 > 0$  such that  $|x^{(n)}| < M_1$  for all  $n \in \mathbb{N}$ . Let  $M_2 := \sum_{k=1}^{\infty} \max_{a \in A_k} |a| < \infty$ . Since  $\lim_{k \rightarrow \infty} \min_{a \in A'_k} |a| = \infty$ , there exists  $p \in \mathbb{N}$  such that for all  $k > p$  we have

$$\min_{a \in A'_k} |a| > M_1 + M_2. \quad (5.2)$$

Arbitrarily take  $n \in \mathbb{N}$ . It suffices to prove  $x^{(n)} \in \sum_{k=1}^p (A_k \cup A'_k) + \sum_{k=p+1}^{\infty} A_k$ . By  $x^{(n)} \in \sum_{k=1}^{\infty} (A_k \cup A'_k)$ , there exist  $x_1^{(n)} \in A_1 \cup A'_1, x_2^{(n)} \in A_2 \cup A'_2, \dots$ , such that  $x^{(n)} = \sum_{k=1}^{\infty} x_k^{(n)}$  converges. Arbitrarily take  $t \geq p+1$ . We only need to prove  $x_t^{(n)} \in A_t$ . Let  $D := \{k \in \mathbb{N} : x_k^{(n)} \in A_k\}$ . It suffices to show  $t \in D$ . If  $\#D < \infty$ , we have

$$x^{(n)} = \sum_{k \in D} x_k^{(n)} + \sum_{k \in \mathbb{N} \setminus D} x_k^{(n)}. \quad (5.3)$$

If  $\#D = \infty$ , we also have (5.3) since

$$\sum_{k \in D} |x_k^{(n)}| \leq \sum_{k \in D} \max_{a \in A_k} |a| \leq M_2 \quad (5.4)$$

implies that  $\sum_{k \in D} x_k^{(n)}$  converges. Thus

$$\left| \sum_{k \in \mathbb{N} \setminus D} x_k^{(n)} \right| \stackrel{\text{by (5.3)}}{=} \left| x^{(n)} - \sum_{k \in D} x_k^{(n)} \right| \leq |x^{(n)}| + \sum_{k \in D} |x_k^{(n)}| \stackrel{\text{by (5.4)}}{\leq} M_1 + M_2 \stackrel{\text{by (5.2)}}{<} \min_{a \in A'_t} |a|. \quad (5.5)$$

Since for every  $k \in \mathbb{N} \setminus D$  we have  $x_k^{(n)} \notin A_k$  and then  $x_k^{(n)} \in A'_k \subseteq [0, \infty)^d$ , it follows that

$$\left| \sum_{k \in \mathbb{N} \setminus D} x_k^{(n)} \right| \geq |x_s^{(n)}| \geq \min_{a \in A'_s} |a| \quad \text{for all } s \in \mathbb{N} \setminus D.$$

By (5.5) we get

$$\min_{a \in A'_t} |a| > \min_{a \in A'_s} |a| \quad \text{for all } s \in \mathbb{N} \setminus D.$$

Therefore  $t \in D$ . □

We prove Corollary 1.6 to end this section.

*Proof of Corollary 1.6.* For all  $k \in \mathbb{N}$ , let  $G_k := B_k \cap [0, c_k]^d$ , then  $\emptyset \neq G_k \subseteq \mathbb{Z}^d$  and we can define  $D_k := \{w^{(1)} \dots w^{(k)} : w^{(1)} \in G_1, \dots, w^{(k)} \in G_k\}$ . Denote the empty word by  $\eta$ , write  $D_0 := \{\eta\}$  and define  $D := \bigcup_{k=0}^{\infty} D_k$ . Let  $J_\eta := J := [0, 1]^d$ . For all  $k \in \mathbb{N}$  and  $w^{(1)} = (w_1^{(1)}, \dots, w_d^{(1)}) \in G_1, \dots, w^{(k)} = (w_1^{(k)}, \dots, w_d^{(k)}) \in G_k$ , define

$$J_{w^{(1)} \dots w^{(k)}} := C_1^{-1} w^{(1)} + C_1^{-1} C_2^{-1} w^{(2)} + \dots + C_1^{-1} \dots C_k^{-1} w^{(k)} + C_1^{-1} \dots C_k^{-1} [0, 1]^d.$$

Let  $\mathcal{F} := \{J_w : w \in D\}$ ,

$$E_k := \bigcup_{w \in D_k} J_w \quad \text{for all } k \geq 0 \quad \text{and} \quad E := \bigcap_{k=0}^{\infty} E_k.$$

First we prove the following **Fact 1** and **Fact 2**.

**Fact 1.**  $\mathcal{F}$  satisfies the Moran Structure Condition (MSC) defined in [21, Section 1.2].

- I. For any  $w \in D$ ,  $J_w$  is obviously geometrically similar to  $J$ .
- II. For any  $k \in \mathbb{N}$  and  $w^{(1)} \dots w^{(k)} \in D_k$ , we need to prove  $J_{w^{(1)} \dots w^{(k)}} \subseteq J_{w^{(1)} \dots w^{(k-1)}}$ .  
It suffices to show

$$\begin{aligned} & C_1^{-1} w^{(1)} + \dots + C_1^{-1} \dots C_{k-1}^{-1} w^{(k-1)} + C_1^{-1} \dots C_{k-1}^{-1} C_k^{-1} w^{(k)} + C_1^{-1} \dots C_{k-1}^{-1} C_k^{-1} [0, 1]^d \\ & \subseteq C_1^{-1} w^{(1)} + \dots + C_1^{-1} \dots C_{k-1}^{-1} w^{(k-1)} + C_1^{-1} \dots C_{k-1}^{-1} [0, 1]^d, \end{aligned}$$

which is equivalent to  $C_k^{-1} w^{(k)} + C_k^{-1} [0, 1]^d \subseteq [0, 1]^d$ , and then also equivalent to  $w^{(k)} + [0, 1]^d \subseteq [0, C_k]^d$ . This follows immediately from  $w^{(k)} \in G_k \subseteq [0, c_k]^d$  and  $c_k + 1 \leq C_k$ .

- III. For any  $k \geq 0$ ,  $w^{(1)} \dots w^{(k)} \in D_k$  and  $u, v \in G_{k+1}$  with  $u \neq v$ , we need to prove  $\text{int}(J_{w^{(1)} \dots w^{(k)} u}) \cap \text{int}(J_{w^{(1)} \dots w^{(k)} v}) = \emptyset$  where  $\text{int}(\cdot)$  denotes the interior of a set.  
It suffices to show

$$\begin{aligned} & (C_1^{-1} w^{(1)} + \dots + C_1^{-1} \dots C_k^{-1} w^{(k)} + C_1^{-1} \dots C_k^{-1} C_{k+1}^{-1} u + C_1^{-1} \dots C_k^{-1} C_{k+1}^{-1} (0, 1)^d) \\ & \cap (C_1^{-1} w^{(1)} + \dots + C_1^{-1} \dots C_k^{-1} w^{(k)} + C_1^{-1} \dots C_k^{-1} C_{k+1}^{-1} v + C_1^{-1} \dots C_k^{-1} C_{k+1}^{-1} (0, 1)^d) = \emptyset. \end{aligned}$$

We only need to prove  $(u + (0, 1)^d) \cap (v + (0, 1)^d) = \emptyset$ . This follows immediately from  $u, v \in G_{k+1} \subseteq \mathbb{Z}^d$  and  $u \neq v$ .

**Fact 2.**  $\sum_{k=1}^{\infty} C_1^{-1} \dots C_k^{-1} G_k = E$ .

□ Let  $x \in \sum_{k=1}^{\infty} C_1^{-1} \dots C_k^{-1} G_k$ . Then there exist  $x^{(1)} \in G_1, x^{(2)} \in G_2, \dots$  such that  $x = \sum_{k=1}^{\infty} C_1^{-1} \dots C_k^{-1} x^{(k)}$  converges in  $\mathbb{R}^d$ . We need to prove  $x \in E$ . Arbitrarily take an integer  $k \geq 0$ . It suffices to show  $x \in \bigcup_{w^{(1)} \dots w^{(k)} \in D_k} J_{w^{(1)} \dots w^{(k)}}$ . We only need to prove  $x \in J_{x^{(1)} \dots x^{(k)}}$ , i.e.,

$$\sum_{n=1}^{\infty} C_1^{-1} \dots C_n^{-1} x^{(n)} \in C_1^{-1} x^{(1)} + \dots + C_1^{-1} \dots C_k^{-1} x^{(k)} + C_1^{-1} \dots C_k^{-1} [0, 1]^d,$$

which is equivalent to

$$\sum_{n=1}^{\infty} C_{k+1}^{-1} \dots C_{k+n}^{-1} x^{(k+n)} \in [0, 1]^d, \quad \text{i.e.,} \quad \sum_{n=1}^{\infty} \frac{(x_1^{(k+n)}, \dots, x_d^{(k+n)})}{C_{k+1} \dots C_{k+n}} \in [0, 1]^d.$$

This follows from the fact that for all  $j \in \{1, \dots, d\}$  we have

$$\begin{aligned}
0 &\leq \sum_{n=1}^{\infty} \frac{x_j^{(k+n)}}{C_{k+1} \cdots C_{k+n}} \leq \sum_{n=1}^{\infty} \frac{c_{k+n}}{(c_{k+1}+1) \cdots (c_{k+n}+1)} \\
&= \frac{c_{k+1}}{c_{k+1}+1} + \frac{c_{k+2}}{(c_{k+1}+1)(c_{k+2}+1)} + \frac{c_{k+3}}{(c_{k+1}+1)(c_{k+2}+1)(c_{k+3}+1)} + \cdots \\
&= \left(1 - \frac{1}{c_{k+1}+1}\right) + \left(\frac{1}{c_{k+1}+1} - \frac{1}{(c_{k+1}+1)(c_{k+2}+1)}\right) \\
&\quad + \left(\frac{1}{(c_{k+1}+1)(c_{k+2}+1)} - \frac{1}{(c_{k+1}+1)(c_{k+2}+1)(c_{k+3}+1)}\right) + \cdots = 1.
\end{aligned}$$

$\square$  Let  $x \in E = \bigcap_{k=0}^{\infty} E_k$ . We need to prove  $x \in \sum_{k=1}^{\infty} C_1^{-1} \cdots C_k^{-1} G_k$ . By  $x \in E_1 = \bigcup_{w^{(1)} \in G_1} J_{w^{(1)}}$ , there exists  $x^{(1)} \in G_1$  such that  $x \in J_{x^{(1)}}$ . By  $x \in E_2 = \bigcup_{w^{(1)} \in G_1, w^{(2)} \in G_2} J_{w^{(1)}w^{(2)}}$ , there exist  $x^{(1)'} \in G_1$  and  $x^{(2)} \in G_2$  such that  $x \in J_{x^{(1)' }x^{(2)}}$ . It follows from  $x \in J_{x^{(1)}} \cap J_{x^{(1)' }x^{(2)}} \neq \emptyset$  and the MSC of  $\mathcal{F}$  in **Fact 1** that  $x^{(1)} = x^{(1)'}$ . Thus  $x \in J_{x^{(1)}x^{(2)}}$ . Repeating this process we know that there exist  $x^{(1)} = (x_1^{(1)}, \dots, x_d^{(1)}) \in G_1, x^{(2)} = (x_1^{(2)}, \dots, x_d^{(2)}) \in G_2, \dots$  such that  $x \in \bigcap_{k=1}^{\infty} J_{x^{(1)} \dots x^{(k)}}$ . In order to prove  $x \in \sum_{k=1}^{\infty} C_1^{-1} \cdots C_k^{-1} G_k$ , we only need to show the following I and II.

- I. Prove that  $\sum_{k=1}^{\infty} C_1^{-1} \cdots C_k^{-1} x^{(k)}$  converges in  $\mathbb{R}^d$ .  
In fact, this follows immediately from

$$0 \leq \sum_{k=1}^{\infty} \frac{x_j^{(k)}}{C_1 \cdots C_k} \leq \sum_{k=1}^{\infty} \frac{c_k}{(c_1+1) \cdots (c_k+1)} = 1 \quad \text{for all } j \in \{1, \dots, d\}.$$

- II. Prove  $x = \sum_{k=1}^{\infty} C_1^{-1} \cdots C_k^{-1} x^{(k)}$ .

Since  $\lim_{k \rightarrow \infty} |J_{x^{(1)} \dots x^{(k)}}| = 0$  and  $J_{x^{(1)}} \supseteq J_{x^{(1)}x^{(2)}} \supseteq J_{x^{(1)}x^{(2)}x^{(3)}} \supseteq \dots$  are all closed sets, we get  $\#(\bigcap_{k=1}^{\infty} J_{x^{(1)} \dots x^{(k)}}) = 1$ . In order to prove  $x = \sum_{k=1}^{\infty} C_1^{-1} \cdots C_k^{-1} x^{(k)}$ , by  $x \in \bigcap_{k=1}^{\infty} J_{x^{(1)} \dots x^{(k)}}$ , it suffices to show  $\sum_{k=1}^{\infty} C_1^{-1} \cdots C_k^{-1} x^{(k)} \in \bigcap_{k=1}^{\infty} J_{x^{(1)} \dots x^{(k)}}$ . In fact this follows in the same way as in the proof of the above " $\square$ ".

Now we deduce statements (1) and (2) in Corollary 1.6 from Theorem 1.5. For all  $k \in \mathbb{N}$ , let  $A_k := C_1^{-1} \cdots C_k^{-1} G_k \neq \emptyset$  and  $A'_k := C_1^{-1} \cdots C_k^{-1} (B_k \setminus [0, c_k]^d)$  (may be  $\emptyset$ ).

(1) Suppose  $\underline{\lim}_{k \rightarrow \infty} C_1^{-1} \cdots C_k^{-1} \min\{\|x\| : x \in B_k \setminus [0, c_k]^d\} > 0$ , i.e.,  $\underline{\lim}_{k \rightarrow \infty} \min_{a \in A'_k} \|a\| > 0$ , which is equivalent to  $\underline{\lim}_{k \rightarrow \infty} \min_{a \in A'_k} |a| > 0$  by (1.4).

- ① Suppose  $\prod_{k=1}^{\infty} \frac{\#G_k}{C_k^d} = 0$  and we need to prove  $\mathcal{L}^d(\sum_{k=1}^{\infty} (A_k \cup A'_k)) = 0$ .

Since  $\mathcal{L}^d$  and  $\mathcal{H}^d$  are equivalent, by Theorem 1.5 (1) ①, it suffices to show  $\mathcal{L}^d(\sum_{k=1}^{\infty} A_k) = 0$ . Recalling **Fact 2**, we only need to prove  $\mathcal{L}^d(E) = 0$ . In fact, this follows immediately from

$$\mathcal{L}^d(E) \leq \mathcal{L}^d(E_k) \leq \sum_{w \in D_k} \mathcal{L}^d(J_w) = \sum_{w \in D_k} \mathcal{L}^d(C_1^{-1} \cdots C_k^{-1} [0, 1]^d) = \frac{\#G_1 \cdots \#G_k}{C_1^d \cdots C_k^d} \rightarrow 0$$

as  $k \rightarrow \infty$  using  $\prod_{k=1}^{\infty} \frac{\#G_k}{C_k^d} = 0$ .

- ② Suppose  $\lim_{k \rightarrow \infty} \frac{\log C_k}{\log C_1 \cdots C_k} = 0$  and we need to prove the Hausdorff and packing dimension formulae for  $\sum_{k=1}^{\infty} C_1^{-1} \cdots C_k^{-1} B_k$ , which is equal to  $\sum_{k=1}^{\infty} (A_k \cup A'_k)$ . Since the proofs of the two formulae are similar, in the following we only prove the Hausdorff one, i.e.,  $\dim_H \sum_{k=1}^{\infty} (A_k \cup A'_k) = \underline{\lim}_{k \rightarrow \infty} \frac{\log \#G_1 \cdots \#G_k}{\log C_1 \cdots C_k}$ . By Theorem 1.5 (1) ① and **Fact 2**, it suffices to show  $\dim_H E = \underline{\lim}_{k \rightarrow \infty} \frac{\log \#G_1 \cdots \#G_k}{\log C_1 \cdots C_k}$  (using [21, Theorem

1.3)]. Since

$$0 = \lim_{k \rightarrow \infty} \frac{\log C_k}{\log C_1 \cdots C_k} \leq \lim_{k \rightarrow \infty} \frac{\log C_k}{\log C_1 \cdots C_k - \log \sqrt{d}} \leq \lim_{k \rightarrow \infty} \frac{\log C_k}{\frac{1}{2} \log C_1 \cdots C_k} = 0,$$

we get

$$\lim_{k \rightarrow \infty} \frac{\log \frac{1}{C_k}}{\log \max_{w \in D_k} |J_w|} = \lim_{k \rightarrow \infty} \frac{-\log C_k}{\log |C_1^{-1} \cdots C_k^{-1}[0, 1]^d|} = \lim_{k \rightarrow \infty} \frac{\log C_k}{\log C_1 \cdots C_k - \log \sqrt{d}} = 0.$$

It follows from **Fact 1** and [21, Theorem 1.3] that

$$\dim_H E = \liminf_{k \rightarrow \infty} \frac{\log \#G_1 \cdots \#G_k}{-\log \frac{1}{C_1} \cdots \frac{1}{C_k}} = \liminf_{k \rightarrow \infty} \frac{\log \#G_1 \cdots \#G_k}{\log C_1 \cdots C_k}.$$

(2) Suppose  $B_k \subseteq [0, \infty)^d$  for all  $k \in \mathbb{N}$  and  $\lim_{k \rightarrow \infty} C_1^{-1} \cdots C_k^{-1} \min\{\|x\| : x \in B_k \setminus [0, c_k]^d\} = \infty$ , i.e.,  $\lim_{k \rightarrow \infty} \min_{a \in A'_k} \|a\| = \infty$ , which is equivalent to  $\lim_{k \rightarrow \infty} \min_{a \in A'_k} |a| = \infty$  by (1.4). We need to prove that  $\sum_{k=1}^{\infty} (A_k \cup A'_k)$  is closed. By Theorem 1.5 (2), it suffices to verify  $\sum_{k=1}^{\infty} \max_{a \in A_k} |a| < \infty$ , which is equivalent to  $\sum_{k=1}^{\infty} \max_{a \in A_k} \|a\| < \infty$ . In fact this follows immediately from

$$\sum_{k=1}^{\infty} \max_{a \in A_k} \|a\| = \sum_{k=1}^{\infty} C_1^{-1} \cdots C_k^{-1} \max_{a \in G_k} \|a\| \leq \sum_{k=1}^{\infty} \frac{c_k}{C_1 \cdots C_k} \leq \sum_{k=1}^{\infty} \frac{c_k}{(c_1 + 1) \cdots (c_k + 1)} = 1.$$

□

## 6. PROOFS OF COROLLARIES 1.8 AND 1.9

Using Corollary 1.7, the main idea in the proofs of Corollaries 1.8 and 1.9 is similar to the one in the proof of [29, Theorem 1.7]. For self-contained and for the convenience of the readers, we still give the detailed proofs as follows.

*Proof of Corollary 1.8.* Let  $d \in \mathbb{N}$  and arbitrarily take  $\alpha, \beta \in [0, d]$  with  $\alpha \leq \beta$ . Let  $m_1 = 2$  and  $m_k = k^2$  for all  $k \geq 2$ . Define a family of functions  $g_\gamma : \mathbb{N} \rightarrow \mathbb{N}$  for  $\gamma \in [0, 1]$  by

$$g_\gamma(n) := \begin{cases} n^{1+\lceil \log n \rceil} & \text{if } \gamma = 0, \\ \lfloor n^{\frac{1}{\gamma}-1} \rfloor n & \text{if } 0 < \gamma < 1, \\ 2n & \text{if } \gamma = 1 \end{cases}$$

where  $\lfloor x \rfloor$  denotes the integer part of  $x$ . Then

$$\lim_{n \rightarrow \infty} \frac{\log n}{\log g_\gamma(n)} = \gamma \quad \text{for all } \gamma \in [0, 1]. \quad (6.1)$$

Choose a strictly increasing sequence of integers  $\{l_j\}_{j=1}^{\infty}$  such that  $l_1 = 0$  and

$$\lim_{j \rightarrow \infty} \frac{l_j \log l_j}{l_{j+1} - l_j} = 0. \quad (6.2)$$

For any  $k \in \mathbb{N}$ , let

$$N_k := \begin{cases} g_{\frac{\alpha}{d}}(m_k) & \text{if } l_j < k \leq l_{j+1} \text{ for some odd } j \in \mathbb{N}, \\ g_{\frac{\beta}{d}}(m_k) & \text{if } l_j < k \leq l_{j+1} \text{ for some even } j \in \mathbb{N}, \end{cases}$$

and let  $B_k := \{0, 1, \dots, m_k - 1\}^d$ . Then  $N_k \geq m_k \geq 2$  are integers with  $m_k \mid N_k$  for all  $k \in \mathbb{N}$ ,  $\lim_{k \rightarrow \infty} \frac{\min\{\|x\| : x \in B_k \setminus \{0, 1, \dots, m_k - 1\}^d\}}{N_1 \cdots N_k} = \infty$  where  $\min \emptyset$  is regarded as  $\infty$ ,  $\prod_{k=1}^{\infty} \frac{\#B_k}{N_k^d} = \prod_{k=1}^{\infty} \frac{m_k^d}{N_k^d} = 0$  by  $N_k \geq 2m_k$  for all  $k$  large enough, and  $\{B_k\}_{k \geq 1}$  is a sequence of nearly  $d$ -th power lattices with respect to  $\{m_k\}_{k \geq 1}$  and the sequence of  $d \times d$  diagonal matrices  $\{\text{diag}(N_k, \dots, N_k)\}_{k \geq 1}$ . In order to use Corollary 1.7, it suffices to prove

$\lim_{k \rightarrow \infty} \frac{\log N_k}{\log N_1 \cdots N_k} = 0$ . In fact, by  $2m_k \leq N_k \leq m_k^{1+\log m_k}$  for all  $k \in \mathbb{N}$  large enough, we have

$$\begin{aligned} \lim_{k \rightarrow \infty} \frac{\log N_k}{\log N_1 N_2 \cdots N_k} &\leq \lim_{k \rightarrow \infty} \frac{\log m_k^{1+\log m_k}}{\log 2m_1 2m_2 \cdots 2m_k} = \lim_{k \rightarrow \infty} \frac{(1 + \log m_k) \log m_k}{k \log 2 + \log m_1 m_2 \cdots m_k} \\ &= \lim_{k \rightarrow \infty} \frac{(1 + \log k^2) \log k^2}{k \log 2 + \log 2 \cdot 2^2 \cdot 3^2 \cdots k^2} \leq \lim_{k \rightarrow \infty} \frac{4(\log k)^2 + 2 \log k}{k \log 2} \\ &= \frac{4}{\log 2} \lim_{k \rightarrow \infty} \frac{(\log k)^2}{k} + \frac{2}{\log 2} \lim_{k \rightarrow \infty} \frac{\log k}{k} = 0. \end{aligned}$$

Therefore, by applying Corollary 1.7, we know that the infinite convolution

$$\mu = \delta_{N_1^{-1}B_1} * \delta_{N_1^{-1}N_2^{-1}B_2} * \delta_{N_1^{-1}N_2^{-1}N_3^{-1}B_3} * \cdots$$

exists, is a singular spectral measure with a spectrum in  $\mathbb{Z}^d$ ,  $\text{spt } \mu = \sum_{k=1}^{\infty} N_1^{-1} \cdots N_k^{-1} B_k$ ,

$$\dim_H \text{spt } \mu = \varliminf_{k \rightarrow \infty} \frac{d \log m_1 \cdots m_k}{\log N_1 \cdots N_k} \quad \text{and} \quad \dim_P \text{spt } \mu = \overline{\lim}_{k \rightarrow \infty} \frac{d \log m_1 \cdots m_k}{\log N_1 \cdots N_k}.$$

To complete the proof, it suffices to show the following (1), (2) and (3).

(1) Prove that  $\sum_{k=1}^{\infty} N_1^{-1} \cdots N_k^{-1} B_k$  is compact.

In fact this follows immediately from Proposition 5.1 (1) and

$$\sum_{k=1}^{\infty} \max_{a \in N_1^{-1} \cdots N_k^{-1} B_k} |a| \leq \sum_{k=1}^{\infty} \frac{(m_k - 1)\sqrt{d}}{N_1 \cdots N_k} \leq \sum_{k=1}^{\infty} \frac{\sqrt{d}}{N_1 \cdots N_{k-1}} \cdot \frac{m_k}{N_k} \leq \sum_{k=1}^{\infty} \frac{\sqrt{d}}{2^{k-1}} = 2\sqrt{d} < \infty.$$

(2) Prove  $\varliminf_{k \rightarrow \infty} \frac{\log m_1 m_2 \cdots m_k}{\log N_1 N_2 \cdots N_k} = \frac{\alpha}{d}$ .

On the one hand, we have

$$\varliminf_{k \rightarrow \infty} \frac{\log m_1 m_2 \cdots m_k}{\log N_1 N_2 \cdots N_k} \stackrel{(*)}{\geq} \varliminf_{k \rightarrow \infty} \frac{\log m_k}{\log N_k} \stackrel{(**)}{\geq} \varliminf_{k \rightarrow \infty} \frac{\log m_k}{\log g_{\frac{\alpha}{d}}(m_k)} \stackrel{\text{by (6.1)}}{=} \frac{\alpha}{d},$$

where  $(*)$  follows from Theorem 2.5 and  $(**)$  follows from  $g_{\frac{\beta}{d}}(m_k) \leq g_{\frac{\alpha}{d}}(m_k)$  for all  $k$  large enough with  $0 \leq \frac{\alpha}{d} \leq \frac{\beta}{d} \leq 1$ .

On the other hand, we have

$$\begin{aligned} \varliminf_{k \rightarrow \infty} \frac{\log m_1 m_2 \cdots m_k}{\log N_1 N_2 \cdots N_k} &\leq \varliminf_{j \rightarrow \infty} \frac{\log m_1 m_2 \cdots m_{l_{2j}}}{\log N_1 N_2 \cdots N_{l_{2j}}} \\ &\leq \varliminf_{j \rightarrow \infty} \frac{\log m_1 m_2 \cdots m_{l_{2j-1}} + \log m_{l_{2j-1}+1} m_{l_{2j-1}+2} \cdots m_{l_{2j}}}{\log N_{l_{2j-1}+1} N_{l_{2j-1}+2} \cdots N_{l_{2j}}} \\ &\leq \varliminf_{j \rightarrow \infty} \left( \frac{l_{2j-1} \log m_{l_{2j-1}}}{(l_{2j} - l_{2j-1}) \log g_{\frac{\alpha}{d}}(m_{l_{2j-1}+1})} + \frac{\log m_{l_{2j-1}+1} m_{l_{2j-1}+2} \cdots m_{l_{2j}}}{\log N_{l_{2j-1}+1} N_{l_{2j-1}+2} \cdots N_{l_{2j}}} \right) \\ &\leq \varliminf_{j \rightarrow \infty} \left( \frac{l_{2j-1}}{l_{2j} - l_{2j-1}} \cdot \frac{\log m_{l_{2j-1}+1}}{\log g_{\frac{\alpha}{d}}(m_{l_{2j-1}+1})} + \frac{\log m_{l_{2j-1}+1} + \log m_{l_{2j-1}+2} + \cdots + \log m_{l_{2j}}}{\log N_{l_{2j-1}+1} + \log N_{l_{2j-1}+2} + \cdots + \log N_{l_{2j}}} \right) \\ &\stackrel{\text{by (6.2)}}{\text{and (6.1)}} \varliminf_{j \rightarrow \infty} \frac{\log m_{l_{2j-1}+1} + \log m_{l_{2j-1}+2} + \cdots + \log m_{l_{2j}}}{\log g_{\frac{\alpha}{d}}(m_{l_{2j-1}+1}) + \log g_{\frac{\alpha}{d}}(m_{l_{2j-1}+2}) + \cdots + \log g_{\frac{\alpha}{d}}(m_{l_{2j}})} = \frac{\alpha}{d}, \end{aligned}$$

where the last equality can be proved as follows. Let  $r := \frac{\alpha}{d} \in [0, 1]$ , and for all  $n \in \mathbb{N}$  let

$$a_n = \log m_1 + \log m_2 + \cdots + \log m_{l_{2n}}, \quad c_n = \log m_1 + \log m_2 + \cdots + \log m_{l_{2n-1}},$$

$$b_n = \log g_r(m_1) + \log g_r(m_2) + \cdots + \log g_r(m_{l_{2n}}), \quad d_n = \log g_r(m_1) + \log g_r(m_2) + \cdots + \log g_r(m_{l_{2n-1}}).$$

It suffices to prove  $\lim_{n \rightarrow \infty} \frac{a_n - c_n}{b_n - d_n} = r$ . Since Theorem 2.5 and (6.1) imply  $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{c_n}{d_n} = r$ , by Proposition 2.7 we only need to verify  $\underline{\lim}_{n \rightarrow \infty} \frac{b_n}{d_n} > 1$ . In fact this follows immediately from

$$\begin{aligned} \underline{\lim}_{n \rightarrow \infty} \frac{b_n}{d_n} - 1 &= \underline{\lim}_{n \rightarrow \infty} \frac{\log g_r(m_{l_{2n-1}+1}) + \log g_r(m_{l_{2n-1}+2}) + \cdots + \log g_r(m_{l_{2n}})}{\log g_r(m_1) + \log g_r(m_2) + \cdots + \log g_r(m_{l_{2n-1}})} \\ &\geq \underline{\lim}_{n \rightarrow \infty} \frac{(l_{2n} - l_{2n-1}) \cdot \log g_r(m_{l_{2n-1}})}{l_{2n-1} \cdot \log g_r(m_{l_{2n-1}})} \stackrel{\text{by (6.2)}}{=} \infty. \end{aligned}$$

(3) Prove  $\overline{\lim}_{k \rightarrow \infty} \frac{\log m_1 m_2 \cdots m_k}{\log N_1 N_2 \cdots N_k} = \frac{\beta}{d}$ .

On the one hand, we have

$$\overline{\lim}_{k \rightarrow \infty} \frac{\log m_1 m_2 \cdots m_k}{\log N_1 N_2 \cdots N_k} \stackrel{(*)}{\leq} \overline{\lim}_{k \rightarrow \infty} \frac{\log m_k}{\log N_k} \stackrel{(**)}{\leq} \overline{\lim}_{k \rightarrow \infty} \frac{\log m_k}{\log g_{\frac{\beta}{d}}(m_k)} \stackrel{\text{by (6.1)}}{=} \frac{\beta}{d},$$

where  $(*)$  follows from Theorem 2.5 and  $(**)$  follows from  $g_{\frac{\alpha}{d}}(m_k) \geq g_{\frac{\beta}{d}}(m_k)$  for all  $k$  large enough with  $0 \leq \frac{\alpha}{d} \leq \frac{\beta}{d} \leq 1$ .

On the other hand, we have

$$\begin{aligned} \overline{\lim}_{k \rightarrow \infty} \frac{\log m_1 m_2 \cdots m_k}{\log N_1 N_2 \cdots N_k} &\geq \overline{\lim}_{j \rightarrow \infty} \frac{\log m_1 m_2 \cdots m_{l_{2j+1}}}{\log N_1 N_2 \cdots N_{l_{2j+1}}} \\ &\geq \overline{\lim}_{j \rightarrow \infty} \frac{\log m_{l_{2j+1}} m_{l_{2j+2}} \cdots m_{l_{2j+1}}}{\log N_1 N_2 \cdots N_{l_{2j}} + \log N_{l_{2j+1}} N_{l_{2j+2}} \cdots N_{l_{2j+1}}} \\ &\geq \overline{\lim}_{j \rightarrow \infty} \frac{\log m_{l_{2j+1}} m_{l_{2j+2}} \cdots m_{l_{2j+1}}}{l_{2j} \log g_{\frac{\alpha}{d}}(m_{l_{2j}}) + \log g_{\frac{\beta}{d}}(m_{l_{2j+1}}) g_{\frac{\beta}{d}}(m_{l_{2j+2}}) \cdots g_{\frac{\beta}{d}}(m_{l_{2j+1}})} \\ &\stackrel{(*)}{=} \overline{\lim}_{j \rightarrow \infty} \frac{\log m_{l_{2j+1}} m_{l_{2j+2}} \cdots m_{l_{2j+1}}}{\log g_{\frac{\beta}{d}}(m_{l_{2j+1}}) g_{\frac{\beta}{d}}(m_{l_{2j+2}}) \cdots g_{\frac{\beta}{d}}(m_{l_{2j+1}})} = \frac{\beta}{d}, \end{aligned}$$

where the last equality can be proved in the same way as the end of the above (2), and  $(*)$  follows from

$$\lim_{j \rightarrow \infty} \frac{l_{2j} \log g_{\frac{\alpha}{d}}(m_{l_{2j}})}{\log m_{l_{2j+1}} m_{l_{2j+2}} \cdots m_{l_{2j+1}}} \leq \lim_{j \rightarrow \infty} \frac{l_{2j} (1 + \log m_{l_{2j}}) \log m_{l_{2j}}}{(l_{2j+1} - l_{2j}) \log m_{l_{2j+1}}} \leq \lim_{j \rightarrow \infty} \frac{l_{2j} (1 + \log l_{2j}^2)}{l_{2j+1} - l_{2j}} \stackrel{\text{by (6.2)}}{=} 0,$$

where the first inequality follows from  $g_{\frac{\alpha}{d}}(n) \leq n^{1+\log n}$  for all  $n \in \mathbb{N}$  large enough.  $\square$

*Proof of Corollary 1.9.* Let  $d \in \mathbb{N}$  and arbitrarily take  $\alpha, \beta \in [0, d]$  with  $\alpha \leq \beta$ . Let  $\{m_k\}_{k \geq 1}$  and  $\{N_k\}_{k \geq 1}$  be defined as in the proof of Corollary 1.8. For all  $k \in \mathbb{N}$ , let

$$B_k := \{0, 1, \dots, m_k - 2, N_1 \cdots N_k \cdot k + m_k - 1\}^d$$

and

$$G_k := B_k \cap \{0, 1, \dots, m_k - 1\}^d = \{0, 1, \dots, m_k - 2\}^d.$$

In a way similar to the proof of Corollary 1.8, by applying Corollary 1.7, we know that the infinite convolution

$$\mu = \delta_{N_1^{-1} B_1} * \delta_{N_1^{-1} N_2^{-1} B_2} * \delta_{N_1^{-1} N_2^{-1} N_3^{-1} B_3} * \cdots$$

exists, is a singular spectral measure with a spectrum in  $\mathbb{Z}^d$ ,  $\text{spt } \mu = \sum_{k=1}^{\infty} N_1^{-1} \cdots N_k^{-1} B_k$ ,

$$\dim_H \text{spt } \mu = \underline{\lim}_{k \rightarrow \infty} \frac{d \log(m_1 - 1) \cdots (m_k - 1)}{\log N_1 \cdots N_k} \text{ and } \dim_P \text{spt } \mu = \overline{\lim}_{k \rightarrow \infty} \frac{d \log(m_1 - 1) \cdots (m_k - 1)}{\log N_1 \cdots N_k}.$$

To complete the proof, we only need to show the following (1) and (2).



(1) Prove that  $\sum_{k=1}^{\infty} N_1^{-1} \cdots N_k^{-1} B_k$  is not compact.

In fact this follows immediately from Proposition 5.1 (2) and

$$\sum_{k=1}^{\infty} \max_{a \in N_1^{-1} \cdots N_k^{-1} B_k} |a| \geq \sum_{k=1}^{\infty} \frac{N_1 \cdots N_k \cdot k + m_k - 1}{N_1 \cdots N_k} \geq \sum_{k=1}^{\infty} k = \infty.$$

(2) Prove  $\lim_{k \rightarrow \infty} \frac{d \log(m_1-1) \cdots (m_k-1)}{\log N_1 \cdots N_k} = \alpha$  and  $\overline{\lim}_{k \rightarrow \infty} \frac{d \log(m_1-1) \cdots (m_k-1)}{\log N_1 \cdots N_k} = \beta$ .

It follows from (2) and (3) in the proof of Corollary 1.8 that we only need to prove

$\lim_{k \rightarrow \infty} \frac{\log(m_1-1) \cdots (m_k-1)}{\log m_1 \cdots m_k} = 1$ . In fact, by

$$\lim_{k \rightarrow \infty} \frac{\log(m_k - 1)}{\log m_k} = \lim_{k \rightarrow \infty} \frac{\log(k^2 - 1)}{\log k^2} = 1,$$

we get

$$\begin{aligned} \lim_{k \rightarrow \infty} \frac{\log(m_1 - 1) \cdots (m_k - 1)}{\log m_1 \cdots m_k} &= \lim_{k \rightarrow \infty} \frac{\log(m_1 - 1) + \cdots + \log(m_k - 1)}{\log m_1 + \cdots + \log m_k} \\ &\stackrel{\text{by Theorem 2.5}}{=} \lim_{k \rightarrow \infty} \frac{\log(m_k - 1)}{\log m_k} = 1. \end{aligned}$$

□

*Acknowledgement.* This work was supported by “National Natural Science Foundation of China” (NSFC 12201652) and (NSFC 12271534).

#### REFERENCES

- [1] L.-X. An, X.-Y. Fu, C.-K. Lai, On spectral Cantor-Moran measures and a variant of Bourgain’s sum of sine problem. *Adv. Math.* 349 (2019), 84–124.
- [2] L.-X. An, L. He, X.-G. He, Spectrality and non-spectrality of the Riesz product measures with three elements in digit sets. *J. Funct. Anal.* 277 (2019), no. 1, 255–278.
- [3] L.-X. An, X.-G. He, A class of spectral Moran measures. *J. Funct. Anal.* 266 (2014), no. 1, 343–354.
- [4] L.-X. An, X.-G. He, K.-S. Lau, Spectrality of a class of infinite convolutions. *Adv. Math.* 283 (2015), 362–376.
- [5] L.-X. An, X.-G. He, H.-X. Li, Spectrality of infinite Bernoulli convolutions. *J. Funct. Anal.* 269 (2015), no. 5, 1571–1590.
- [6] M.-L. Chen, J.-C. Liu, X.-Y. Wang, Spectrality of a class of self-affine measures on  $\mathbb{R}^2$ . *Nonlinearity* 34 (2021), no. 11, 7446–7469.
- [7] X.-R. Dai, When does a Bernoulli convolution admit a spectrum? *Adv. Math.* 231 (2012), no. 3-4, 1681–1693.
- [8] X.-R. Dai, X.-Y. Fu, Z.-H. Yan, Spectrality of self-affine Sierpinski-type measures on  $\mathbb{R}^2$ . *Appl. Comput. Harmon. Anal.* 52 (2021), 63–81.
- [9] X.-R. Dai, X.-G. He, C.-K. Lai, Spectral property of Cantor measures with consecutive digits. *Adv. Math.* 242 (2013), 187–208.
- [10] X.-R. Dai, X.-G. He, K.-S. Lau, On spectral  $N$ -Bernoulli measures. *Adv. Math.* 259 (2014), 511–531.
- [11] X.-R. Dai, Q. Sun, Spectral measures with arbitrary Hausdorff dimensions. *J. Funct. Anal.* 268 (2015), no. 8, 2464–2477.
- [12] Q.-R. Deng, J.-B. Chen, Uniformity of spectral self-affine measures. *Adv. Math.* 380 (2021), Paper No. 107568, 17 pp.
- [13] Q.-R. Deng, K.-S. Lau, Sierpinski-type spectral self-similar measures. *J. Funct. Anal.* 269 (2015), no. 5, 1310–1326.
- [14] D. E. Dutkay, J. Haussermann, C.-K. Lai, Hadamard triples generate self-affine spectral measures. *Trans. Amer. Math. Soc.* 371 (2019), no. 2, 1439–1481.
- [15] D. E. Dutkay, C.-K. Lai, Spectral measures generated by arbitrary and random convolutions. *J. Math. Pures Appl.* (9) 107 (2017), no. 2, 183–204.
- [16] K. Falconer, *Fractal geometry. Mathematical foundations and applications.* Third edition. John Wiley & Sons, Ltd., Chichester, 2014. xxx+368 pp. ISBN: 978-1-119-94239-9
- [17] Y.-S. Fu, X.-G. He, Z.-X. Wen, Spectra of Bernoulli convolutions and random convolutions. *J. Math. Pures Appl.* (9) 116 (2018), 105–131.

- [18] Y.-S. Fu, M.-W. Tang, Existence of exponential orthonormal bases for infinite convolutions on  $\mathbb{R}^d$ . *J. Fourier Anal. Appl.* 30 (2024), no. 3, Paper No. 31, 43 pp.
- [19] B. Fuglede, Commuting self-adjoint partial differential operators and a group theoretic problem. *J. Funct. Anal.* 16 (1974), 101–121.
- [20] X.-G. He, C.-K. Lai, K.-S. Lau, Exponential spectra in  $L^2(\mu)$ . *Appl. Comput. Harmon. Anal.* 34 (2013), no. 3, 327–338.
- [21] S. Hua, H. Rao, Z. Wen, J. Wu, On the structures and dimensions of Moran sets. *Sci. China Ser. A* 43 (2000), no. 8, 836–852.
- [22] J. Jacod, P. Protter, *Probability essentials*. Second edition. Universitext. Springer-Verlag, Berlin, 2003. x+254 pp. ISBN: 3-540-43871-8
- [23] B. Jessen, A. Wintner, Distribution functions and the Riemann zeta function. *Trans. Amer. Math. Soc.* 38 (1935), no.1, 48–88.
- [24] P. Jorgensen, S. Pedersen, Dense analytic subspaces in fractal  $L^2$ -spaces. *J. Anal. Math.* 75 (1998), 185–228.
- [25] I. Łaba, Y. Wang, Some properties of spectral measures. *Appl. Comput. Harmon. Anal.* 20 (2006), no. 1, 149–157.
- [26] J. Li, Z. Wu, On spectral structure and spectral eigenvalue problems for a class of self similar spectral measure with product form. *Nonlinearity* 35 (2022), no. 6, 3095–3117.
- [27] J. Li, Z. Wu, On the quasi-Beurling dimensions of the spectra for planar Moran-type Sierpinski spectral measures. *Appl. Comput. Harmon. Anal.* 62 (2023), 475–497.
- [28] J.-L. Li, Spectra of a class of self-affine measures. *J. Funct. Anal.* 260 (2011), no. 4, 1086–1095.
- [29] W. Li, J. J. Miao, Z. Wang, Weak convergence and spectrality of infinite convolutions. *Adv. Math.* 404 (2022), part B, Paper No. 108425, 26 pp.
- [30] W. Li, J. J. Miao, Z. Wang, Spectrality of random convolutions generated by finitely many Hadamard triples. *Nonlinearity* 37 (2024), no. 1, Paper No. 015003, 21 pp.
- [31] W. Li, J. J. Miao, Z. Wang, Spectrality of infinite convolutions and random convolutions. *J. Funct. Anal.* 287 (2024), no. 7, Paper No. 110539.
- [32] W. Li, Z. Wang, The Spectrality of Infinite Convolutions in  $\mathbb{R}^d$ . *J. Fourier Anal. Appl.* 30 (2024), no. 3, Paper No. 35.
- [33] J.-C. Liu, J. J. Luo, Spectral property of self-affine measures on  $\mathbb{R}^n$ . *J. Funct. Anal.* 272 (2017), no. 2, 599–612.
- [34] J.-C. Liu, Y. Zhang, Z.-Y. Wang, M.-L. Chen, Spectrality of generalized Sierpinski-type self-affine measures. *Appl. Comput. Harmon. Anal.* 55 (2021), 129–148.
- [35] Z.-Y. Lu, X.-H. Dong, P.-F. Zhang, Spectrality of some one-dimensional Moran measures. *J. Fourier Anal. Appl.* 28 (2022), no. 4, Paper No. 63, 22 pp.
- [36] R. S. Strichartz, Mock Fourier series and transforms associated with certain Cantor measures. *J. Anal. Math.* 81 (2000), 209–238.
- [37] R. S. Strichartz, Convergence of mock Fourier series. *J. Anal. Math.* 99 (2006), 333–353.
- [38] Z.-H. Yan, Spectral Moran measures on  $\mathbb{R}^2$ . *Nonlinearity* 35 (2022), no. 3, 1261–1285.

SCHOOL OF MATHEMATICS AND STATISTICS, GUANGDONG UNIVERSITY OF TECHNOLOGY, GUANGZHOU, 510520, P.R. CHINA

*Email address:* yaoqiang.li@gdut.edu.cn scutyaoqiangli@qq.com