Distributional stability of sparse inverse covariance matrix estimators^{*}

Renjie Chen[†]

Huifu Xu[‡]

Henryk Zähle[§]

July 9, 2024

Abstract

Finding an approximation of the inverse of the covariance matrix, also known as precision matrix, of a random vector with empirical data is widely discussed in finance and engineering. In data-driven problems, empirical data may be "contaminated". This raises the question as to whether the approximate precision matrix is reliable from a statistical point of view. In this paper, we concentrate on a much-noticed sparse estimator of the precision matrix and investigate the issue from the perspective of distributional stability. Specifically, we derive an explicit local Lipschitz bound for the distance between the distributions of the sparse estimator under two different distributions (regarded as the true data distribution and the distribution of "contaminated" data). The distance is measured by the Kantorovich metric on the set of all probability measures on a matrix space. We also present analogous results for the standard estimators of the covariance matrix and its eigenvalues. Furthermore, we discuss two applications and conduct some numerical experiments.

Keywords: Covariance matrix, precision matrix, sparse estimator, data perturbation, distributional stability, statistical robustness, data-driven

1 Introduction

Let ξ be an \mathbb{R}^n -valued random variable on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. The statistical estimation of its covariance matrix $\Sigma = \mathbb{E}[(\xi - \mathbb{E}[\xi])(\xi - \mathbb{E}[\xi])^{\mathsf{T}}]$ and the inverse Σ^{-1} thereof, if it exists, is a standard and long standing problem in multivariate statistics with wide applications in finance and engineering. The inverse matrix Σ^{-1} , which is also known as precision matrix, is needed, for example, for optimal decision or model selection such as linear discriminant analysis [17], portfolio optimization [16] and graphical model selection [7, 14, 56]. Given independent copies ξ^1, \ldots, ξ^N of ξ , the classical nonparametric estimators of Σ and Σ^{-1} are the sample covariance matrix $\hat{\Sigma}_N :=$ $\frac{1}{N} \sum_{i=1}^N \xi^i \xi^{i^{\mathsf{T}}} - (\frac{1}{N} \sum_{i=1}^N \xi^i)(\frac{1}{N} \sum_{i=1}^N \xi^i)^{\mathsf{T}}$ and its inverse $\hat{\Sigma}_N^{-1}$, if it exists, respectively. In the case where it exists, $\hat{\Sigma}_N^{-1}$ is called the sample precision matrix.

^{*}Dedicated to the memory of Professor Werner Römisch, a pioneer in stability analysis of stochastic optimization.

[†]Department of Systems Engineering and Engineering Management, The Chinese University of Hong Kong. Email: rchen@se.cuhk.edu.hk.

[‡]Department of Systems Engineering and Engineering Management, The Chinese University of Hong Kong. Email: hfxu@se.cuhk.edu.hk.

[§]Department of Mathematics, Saarland University, Saarbrücken, Germany. Email: zaehle@math.uni-sb.de

In this article, our main focus is on the estimation of the precision matrix Σ^{-1} , which is subject to some problems. Two of these problems are the following. First, the sample precision matrix $\widehat{\Sigma}_N^{-1}$, i.e. the inverse of the sample covariance matrix, can fail to exist even if the precision matrix Σ^{-1} exists. Second, the sample precision matrix $\widehat{\Sigma}_N^{-1}$ can fail to have a sparse structure even if the precision matrix Σ^{-1} has a sparse structure. Recall that sparsity of covariance and precision matrices is a basic requirement for many applications such as model selection problems, see e.g. [6, 12, 35, 39, 43, 56]. To overcome these two problems, Banerjee et. al. [2] introduced, in the scope of a graphical model selection problem, the estimator

$$\widehat{S}_N := \arg\min_{S \in \mathbb{S}^n_{++}} \left(\langle \widehat{\Sigma}_N, S \rangle - \log(\det S) + \lambda \|S\|_1 \right)$$
(1)

of Σ^{-1} , where \mathbb{S}_{++}^n is the set of all symmetric and positive definite matrices in $\mathbb{R}^{n \times n}$, $\langle \hat{\Sigma}_N, S \rangle$ is defined to be the trace of the matrix $\hat{\Sigma}_N^\mathsf{T} S$, $\|S\|_1 := \sum_{1 \le i,j \le n} |S_{ij}|$ and $\lambda \in \mathbb{R}_+ := [0, \infty)$. For $\lambda = 0$, the estimator \hat{S}_N is well defined only for samples ξ^1, \ldots, ξ^N for which the symmetric matrix $\hat{\Sigma}_N$ is positive definite and in this case it boils down to the sample precision matrix $\hat{\Sigma}_N^{-1}$ (see Section 10.1). For $\lambda > 0$, the last term on the right-hand side in (1), i.e. $\lambda \|S\|_1$, is an ℓ_1 -penalty term which penalizes deviations from sparsity, i.e., more precisely, a large number of nonzero entries. The parameter λ controls the intensity of the penalty, which is consistent with techniques used in regression problems, such as the lasso. The choice $\lambda > 0$ has the advantage over $\lambda = 0$ not only that the minimizer becomes sparser, but also that the estimator \hat{S}_N is well defined for *each* sample ξ^1, \ldots, ξ^N (this is ensured by Proposition 4.1). In [41], it is pointed out that the right-hand side in (1) can also be viewed as an ℓ_1 -penalized Bregman divergence minimization problem.

The estimator \hat{S}_N and variants of it have also been considered in several other papers, such as [10, 18, 41, 56]. Convergence analysis (in terms of Frobenius and spectral norms) and model selection consistency can be found in [43] for Gaussian random variables, in [41] for general random variables and in [35] for an estimator that is based on a version of (1) with more general (non-convex) regularization term. In [40], the authors proposed a so-called Wasserstein shrinkage estimator of precision matrix based on a distributionally robust version of (1). In [48, 54], the authors discuss numerical methods to solve (1).

In this paper, we intend to investigate the extent to which the estimator \widehat{S}_N is distributional stable to deviations in the underlying distribution. By underlying distribution we mean the distribution $\mathbb{P} \circ \xi^{-1}$ of the random variable ξ . The set of all admissible underlying distributions is the set $\mathscr{P}_2(\mathbb{R}^n)$ of all Borel probability measures P on \mathbb{R}^n with $\int_{\mathbb{R}^n} ||x||^2 P(dx) < \infty$. To properly frame the question we want to raise and answer, we need to change over to the canonical setting. So, set

$$(\Omega, \mathcal{F}) := ((\mathbb{R}^n)^N, \mathcal{B}(\mathbb{R}^n)^{\otimes N}) \quad \text{and} \quad \mathbb{P}^P := P^{\otimes N} \text{ for any } P \in \mathscr{P}_2(\mathbb{R}^n)$$

and let $\boldsymbol{\xi} = (\xi^1, \ldots, \xi^N)$ be the identity on Ω . Then ξ^1, \ldots, ξ^N are i.i.d. according to P under \mathbb{P}^P for any $P \in \mathscr{P}_2(\mathbb{R}^n)$. In particular, $(\Omega, \mathcal{F}, \{\mathbb{P}^P : P \in \mathscr{P}_2(\mathbb{R}^n)\})$ is a nonparametric statistical model and $\widehat{S}_N = \widehat{S}_N(\boldsymbol{\xi})$ is a point estimator of the precision matrix of ξ^1 in this model. For any $P \in \mathscr{P}_2(\mathbb{R}^n)$, the image measure $\mathbb{P}^P \circ \widehat{S}_N^{-1}$ is a probability measure on $(\mathbb{S}_{++}^n, \mathcal{B}(\mathbb{S}_{++}^n))$ and specifies the distribution of the estimator \widehat{S}_N if the underlying distribution is P. Here we assume that \mathbb{S}_{++}^n is equipped with the Frobenius norm, such that the mapping $\widehat{S}_N : \Omega \to \mathbb{S}_{++}^n$ is Borel measurable, since it is continuous (which follows from Theorem 4.1 and the continuity of $\widehat{\Sigma}_N : \Omega \to \mathbb{S}_{+}^n$). We want to investigate the question of how much the distribution $\mathbb{P}^P \circ \widehat{S}_N^{-1}$ changes when the underlying distribution P is replaced by another distribution Q. A change from P to Q can be due to a "contamination" of the sample data, caused, for example, by unsystematic outliers, random measurement errors or simply observing realizations from the wrong distribution. As early as 1971, Hampel [25] introduced the concept of qualitative robustness of a nonparametric estimator \widehat{T}_N , which basically means that the distribution of the estimator, here $\mathbb{P}^P \circ \widehat{T}_N^{-1}$, is continuous in the underlying distribution P (uniformly in N), where sets of probability measures are equipped with the respective weak topologies (see also the monographs [26, 27]). In the 2010th, the property of qualitative robustness has attracted some interest in the field of quantitative risk management, see, for instance, [4, 9, 30, 31, 32, 33, 34, 50]. In [57, 58], it was extended to general estimators and general statistical models.

Hampel's concept of qualitative robustness has at least two limitations. First, the distance $dl(\mathbb{P}^P \circ \hat{T}_N^{-1}, \mathbb{P}^Q \circ \hat{T}_N^{-1})$ is measured w.r.t. metrics dl which metrize the weak topology (such as Lévy, Prohorov or bounded Lipschitz). This is a little unsatisfactory because, for example, in the case where \hat{T}_N is real-valued, the means of $\mathbb{P}^P \circ \hat{T}_N^{-1}$ and $\mathbb{P}^Q \circ \hat{T}_N^{-1}$ can be arbitrarily far apart, even if $dl(\mathbb{P}^P \circ \hat{T}_N^{-1}, \mathbb{P}^Q \circ \hat{T}_N^{-1}) \leq \varepsilon$ for some given $\varepsilon > 0$. Second, no explicit bounds for $dl(\mathbb{P}^P \circ \hat{T}_N^{-1}, \mathbb{P}^Q \circ \hat{T}_N^{-1})$ in terms of dl'(P,Q) are required, where dl' is a metric on the set of admissible underlying distributions. In this respect, qualitative robustness differs from stability concepts in the fields of optimization and numerical analysis, where stability is usually interpreted as Lipschitz continuity.

For these reasons, Guo and Xu [20] extended, in the scope of a preference optimization problem, the conventional robustness concept in two direction. First, they replaced dl by the Kantorovich metric dI_K , which ensures, for real-valued \hat{T}_N , that the means of $\mathbb{P}^P \circ \hat{T}_N^{-1}$ and $\mathbb{P}^Q \circ \hat{T}_N^{-1}$ are close to each other when $dI_K(\mathbb{P}^P \circ \hat{T}_N^{-1}, \mathbb{P}^Q \circ \hat{T}_N^{-1})$ is small. Second, they aimed at Lipschitz bounds in the form of $dI_K(\mathbb{P}^P \circ \hat{T}_N^{-1}, \mathbb{P}^Q \circ \hat{T}_N^{-1}) \leq L dI'(P,Q)$ for all admissible P,Q, for some constant L > 0, where dI' is chosen appropriately. Such Lipschitz bounds have recently been established in other applications as well, see [21, 23, 49]. In this paper, we establish a corresponding (local) Lipschitz bound for the sparse precision matrix estimator \hat{S}_N defined in (1). In fact, Theorem 5.3 shows that there exists a constant $L_\lambda > 0$, depending only on λ (i.e. being independent of P, Q and N), such that

$$\mathsf{dl}_{\mathsf{K}}\left(\mathbb{P}^{P}\circ\widehat{S}_{N}^{-1},\mathbb{P}^{Q}\circ\widehat{S}_{N}^{-1}\right) \leq L_{\lambda}\max\{3,2m_{P},2m_{Q}\}\,\mathsf{dl}_{2}(P,Q)$$
(2)

where dl_2 is the second order Fortet-Mourier metric on $\mathscr{P}_2(\mathbb{R}^n)$ and m_P and m_Q are the absolute means of P and Q, respectively. With regard to the right-hand side of (2), it is worth noting that obviously $m_Q \leq m_P + |m_P - m_Q| \leq m_P + dl_1(P,Q)$, where dl_1 is the first order Fortet-Mourier metric on $\mathscr{P}_1(\mathbb{R}^n)$. Bounds analogous to (2) are also obtained for the sample covariance matrix $\widehat{\Sigma}_N$ (see Theorem 5.1) and the eigenvalues of $\widehat{\Sigma}_N$ (see Theorem 5.4). In Section 6 we apply these results to a Gaussian graphical model selection problem and a portfolio optimization problem. Combining (2) with (22a) and (23b), we also obtain that

$$\mathsf{dl}_{\mathsf{K}}\left(\mathbb{P}^{Q}\circ\widehat{S}_{N}^{-1},\delta_{S_{P}}\right) \leq L_{\lambda}\max\{3,2m_{P},2m_{Q}\}\,\mathsf{dl}_{2}(P,Q) + o\left(N^{-(r-1)/r}\right) \tag{3}$$

for all $r \in (1, 2]$ and $P, Q \in \mathscr{P}_2(\mathbb{R}^n)$ with $\int_{\mathbb{R}^n} ||x||^{2r} P(\mathrm{d}x) < \infty$, where δ_{S_P} is the Dirac measure at the precision matrix S_P given the underlying model is P and the *o*-term on the right-hand side of (3) only depends on P (i.e. it is independent of Q). Inequality (3) shows that \widehat{S}_N can be reasonably estimated even if the data are drawn from a "contaminated" distribution Q that is slightly different from the target distribution P. Note that the Borel probability measures on the left-hand side in (2) are distributions of random matrices (with samples in \mathbb{S}_{++}^n). For an overview on random matrix theory we refer to [15, 46], and we point out that random matrices have wide application in engineering [47], physics [3], random graph theory [28], neural networks [37] and, most related to this paper, statistics. The distribution of the sample covariance matrix $\hat{\Sigma}_N$, first studied in [52], has been a central research problem in multivariate statistical analysis. The analysis of spectrum behavior of random matrix founds a base for principle component analysis [29] and factor analysis models. The empirical distribution of the eigenvalues of random matrices, often referred to as empirical spectral distribution, was first studied in [51] and considered in detail for sample covariance matrices in [38]. The eigenvector empirical spectral distribution was studied in [1, 44] and, more recently, in [53].

The rest of this article is organized as follows. In Section 2, we first introduce some basic notation. In Section 3, we introduce a criterion for distributional stability of general point estimators, which we use in Section 5 to prove our main results. For one of our main results, we also need information on the optimization problem underlying the estimator \hat{S}_N , which we provide in Section 4. Two applications are discussed in Section 6, and some numerical experiments can be found in Section 7. All the proofs are given in Section 10, some of which rely on the auxiliary results in Section 9. Some open questions for future research are listed in Section 8.

2 Basic notation

We use \mathbb{R}^n to denote the *n*-dimensional Euclidean space and ||x|| to denote the Euclidean norm of $x \in \mathbb{R}^n$. We set $\mathbb{R}_+ := [0, \infty)$ and $\mathbb{R}_{++} := (0, \infty)$. We use $\mathbb{R}^{n \times n}$ to denote the set of all $n \times n$ matrices with real entries and \mathbb{S}^n to denote linear space of all symmetric matrices in $\mathbb{R}^{n \times n}$. Moreover, we use \mathbb{S}^n_+ to denote the set of all positive semi-definite matrices in \mathbb{S}^n , and \mathbb{S}^n_{++} to denote the set of all positive definite matrices in \mathbb{S}^n . For a matrix $A \in \mathbb{R}^{n \times n}$, we write A_{ij} for its (i, j)-th entry. For $A, B \in \mathbb{R}^{n \times n}$, we write $\langle A, B \rangle$ for the inner product of A and B which is defined as the trace $\operatorname{tr}(A^T B)$ of $A^T B$. For any matrix $A \in \mathbb{R}^{n \times n}$, we write ||A|| for its Frobenius norm, i.e. $||A|| := \sqrt{\langle A, A \rangle}$ (which will be used mostly in the paper), and $||A||_1 := \sum_{i=1}^n \sum_{j=1}^n |A_{ij}|$ and $||A||_2 := \sup_{||x||=1} ||Ax||$ for the 1-norm and the 2-norm (spectral norm), respectively. Unless otherwise stated, $\mathbb{R}^{n \times n}$ and all of its subspaces are equipped with the Frobenius norm. We use $\operatorname{diag}(x_1, \ldots, x_n)$ to represent a diagonal matrix with diagonal entries $x_1, \ldots, x_n \in \mathbb{R}$. Furthermore, we use $\mathbf{0}$ to denote the zero vector or zero matrix in the space depending on the context.

For any normed linear space $(X, \|\cdot\|_X)$, we use $\mathscr{P}(X)$ to denote the set of all Borel probability measures on X. For any $p \in [1, \infty)$, we write $\mathscr{P}_p(X)$ for the subset of all $P \in \mathscr{P}(X)$ satisfying $\int_X \|x\|_X^p P(\mathrm{d} x) < \infty$ and $\mathcal{F}_p(X)$ for the set of all functions $\psi : X \to \mathbb{R}$ satisfying $|\psi(\hat{x}) - \psi(\tilde{x})| \le L_p(\hat{x}, \tilde{x}) \|x - \tilde{x}\|$ for all $\hat{x}, \tilde{x} \in X$, where $L_p(\hat{x}, \tilde{x}) := \max\{1, \|\hat{x}\|_X, \|\tilde{x}\|_X\}^{p-1}$. For any $p \in [1, \infty)$, the *p*-th order Fortet-Mourier metric on $\mathscr{P}_p(X)$ is defined by

$$\mathsf{dl}_{X,p}(P,Q) := \sup_{\psi \in \mathcal{F}_p(X)} \Big| \int_X \psi(x) P(\mathrm{d}x) - \int_X \psi(x) Q(\mathrm{d}x) \Big|, \quad P,Q \in \mathscr{P}_p(X).$$
(4)

In the case p = 1, the Fortet-Mourier metric $d|_{X,p}(P,Q)$ recovers the well-known Kantorovich metric (also known as Wasserstein distance). Note that Fortet-Mourier metrics are extensively used in stability analysis of stochastic programming, see [42] for an overview.

3 A criterion for distributional stability of general estimators

The main results of this paper, i.e. distributional stability of the estimators of the covariance matrix and the precision matrix, will be presented in Section 5. These results rely on Theorem 3.1 below, which addresses distributional stability of general statistical estimators. Indeed, applying Theorem 3.1 to the sample covariance $\hat{\Sigma}_N$ (in the role of \hat{T}_N) shows that $\hat{\Sigma}_N$ is distributionally stable (see Theorem 5.1). To conclude that the sparse estimator \hat{S}_N of the precision matrix Σ^{-1} (introduced in (1)) is also distributionally stable (see Theorem 5.3), we need some additional arguments which are prepared in Section 4.

Let $(X, \|\cdot\|_X)$ be a normed linear space and consider, for any fixed $N \in \mathbb{N}$, the measurable space $(\Omega, \mathcal{F}) := (X^N, \mathcal{B}(X)^{\otimes N})$, which is regarded as the sample space. Moreover, set

$$\mathbb{P}^P := P^{\otimes N}$$
 for any $P \in \mathscr{P}_2(X)$.

For any $P \in \mathscr{P}_2(X)$, the N coordinate projections on $\Omega = X^N$, denoted by ξ^1, \ldots, ξ^N , are, under \mathbb{P}^P , i.i.d. according to P. Note that $(\Omega, \mathcal{F}, \{\mathbb{P}^P : P \in \mathscr{P}_2(X)\})$ is a nonparametric statistical model and $\boldsymbol{\xi} := (\xi^1, \ldots, \xi^N)$ (= identity on Ω) is the sample variable on it. Let $(Y, \|\cdot\|_Y)$ be another normed linear space and $\widehat{T}_N : \Omega \to Y$ be a $(\mathcal{B}(\Omega), \mathcal{B}(Y))$ -measurable mapping, which is regarded as a statistical estimator. Moreover, set

$$m_P := \mathbb{E}^P[\|\xi^1\|_X] := \int_{\Omega} \|\xi^1\|_X \, \mathrm{d}\mathbb{P}^P = \int_X \|x\|_X \, \mathrm{d}P$$

for any $P \in \mathscr{P}_2(X)$ and recall that $L_2(x, \tilde{x}) = \max\{1, \|x\|_X, \|\tilde{x}\|_X\}$ for all $\hat{x}, \tilde{x} \in X$. Moreover, recall from Section 2 that $d|_{X,2}$ and $d|_{Y,1}$ refer to the second order Fortet-Mourier metric on $\mathscr{P}_2(X)$ and the first order Fortet-Mourier metric on $\mathscr{P}_1(Y)$, respectively.

Theorem 3.1. Assume that there exist constants $\kappa_1, \kappa_2 \in \mathbb{R}_+$ such that

$$\begin{aligned} \left\| \widehat{T}_{N}(\hat{\boldsymbol{x}}) - \widehat{T}_{N}(\tilde{\boldsymbol{x}}) \right\|_{Y} &\leq \frac{\kappa_{1}}{N} \sum_{i=1}^{N} L_{2}(\hat{x}^{i}, \tilde{x}^{i}) \| \hat{x}^{i} - \tilde{x}^{i} \|_{X} \\ &+ \frac{\kappa_{2}}{N^{2}} \sum_{i=1}^{N} \left(\| \hat{x}^{i} \|_{X} + \| \tilde{x}^{i} \|_{X} \right) \sum_{j=1}^{N} \| \hat{x}^{j} - \tilde{x}^{j} \|_{X} \end{aligned}$$
(5)

holds true for all $\hat{\boldsymbol{x}} = (\hat{x}^1, \dots, \hat{x}^N), \ \tilde{\boldsymbol{x}} = (\tilde{x}^1, \dots, \tilde{x}^N) \in X^N$. Then

$$\mathsf{dl}_{Y,1}\left(\mathbb{P}^P \circ \widehat{T}_N^{-1}, \mathbb{P}^Q \circ \widehat{T}_N^{-1}\right) \le \max\left\{\kappa_1 + \kappa_2, 2\kappa_2 m_P, 2\kappa_2 m_Q\right\} \mathsf{dl}_{X,2}(P,Q) \tag{6}$$

for all $P, Q \in \mathscr{P}_2(X)$ and $N \in \mathbb{N}$.

The result of Theorem 3.1 is similar to that of Theorem 4.5 in [49]. However, they differ in that the right-hand side of (6) contains the terms m_P and m_Q . This is because \widehat{T}_N is assumed to satisfy condition (5) with the right-hand side having a term $\frac{1}{N^2} \sum_{i=1}^{N} (\|\hat{x}^i\|_X + \|\tilde{x}^i\|_X) \sum_{j=1}^{N} \|\hat{x}^j - \tilde{x}^j\|_X$. We consider this term because it allows the theorem to be applied to the sample covariance matrix. It might also be helpful to note that Inequality (6) holds for all $N \in \mathbb{N}$ rather than only for N sufficiently large as in many asymptotic statistical analyses. This is primarily because the statistical estimator \widehat{T}_N is assumed to satisfy the local Lipschitz property (5) with respect to perturbation of sample data for all sample sizes N. This Lipschitz property controls the Kantorovich distance between $\mathbb{P}^P \circ \widehat{T}_N^{-1}$ and $\mathbb{P}^Q \circ \widehat{T}_N^{-1}$. In our view, this is the unique feature of quantitative statistical robustness

as introduced in [20] which effectively relates Lipschitz continuity of a statistical estimator to the Kantorovich distance of the distributions of the associated estimators based on two different sampling distributions.

The following proposition is trivial, but worth noting. In the proposition, $(Z, \|\cdot\|_Z)$ is another normed linear space.

Proposition 3.1. Assume that there exists a constant $L_1 \in \mathbb{R}_+$ such that

$$\mathrm{dl}_{Y,1}\left(\mathbb{P}^P \circ \widehat{T}_N^{-1}, \mathbb{P}^Q \circ \widehat{T}_N^{-1}\right) \leq L_1 \,\mathrm{dl}_{X,2}(P,Q)$$

for all $P, Q \in \mathscr{P}_2(X)$ and $N \in \mathbb{N}$. Moreover, let $g: Y \to Z$ be a map that is Lipschitz continuous on a subset $Y_0 \subseteq Y$ with Lipschitz constant $L_0 \in \mathbb{R}_+$. Then

$$\mathsf{dl}_{Z,1}\left(\mathbb{P}^P \circ g(\widehat{T}_N)^{-1}, \mathbb{P}^Q \circ g(\widehat{T}_N)^{-1}\right) \leq L_0 L_1 \,\mathsf{dl}_{X,2}(P,Q)$$

for all $P, Q \in \mathscr{P}_2(X)$ with $\widehat{T}_N \in Y_0 \mathbb{P}^P$ -a.s. and \mathbb{P}^Q -a.s., for any $N \in \mathbb{N}$.

4 Optimization problem underlying the estimator \widehat{S}_N

In this section, we formally formulate and analyze the optimization problem underlying the estimator \hat{S}_N defined in (1). Theorem 4.2 is specifically needed to obtain distributional stability of \hat{S}_N in Theorem 5.3.

In Section 4.1, we formulate the optimization problem (see (7)) and show that it has a unique minimizer, denoted by $S^*(\lambda, \Sigma)$, and that its feasible set can be even chosen smaller than \mathbb{S}^n_{++} without affecting its optimal solution. In Sections 4.2 and 4.3, we analyze the mapping $\mathbb{S}^n_+ \to \mathbb{S}^n_{++}$, $\Sigma \mapsto$ $S^*(\lambda, \Sigma)$ for continuity and Lipschitz continuity, respectively. The proof of continuity is based on the growth condition (11). The approach to prove Lipschitz continuity is to write down the first order optimality condition at the optimal solution and then use an implicit function theorem. The main difference of the two approaches is that the former does not require continuous differentiability of $S \mapsto L(\lambda, \Sigma, S)$, whereas the latter does. We propose a smoothing approach to get around the nonsmoothness issue in the second approach and this is indeed the main contributions of this section.

4.1 Formulation and existence of a unique minimizer

Recall that the spaces \mathbb{S}^n_+ and \mathbb{S}^n_{++} are equipped with the Frobenius norm $\|\cdot\|$ and define a mapping $L: \mathbb{R}_{++} \times \mathbb{S}^n_+ \times \mathbb{S}^n_+ \to \mathbb{R}$ by

$$L(\lambda, \Sigma, S) := \langle \Sigma, S \rangle - \log(\det S) + \lambda \|S\|_1,$$

where as before $||S||_1 := \sum_{1 \le i,j \le n} |S_{ij}|$. For any fixed $\lambda \in \mathbb{R}_{++}$ and $\Sigma \in \mathbb{S}^n_+$, consider the minimization problem

$$\min_{S \in \mathbb{S}^n_{++}} L(\lambda, \Sigma, S).$$
(7)

If the second argument of L, i.e. Σ , is chosen to be $\widehat{\Sigma}_N$, then a minimizer of (7) is just \widehat{S}_N . The following proposition ensures that (7) possesses a unique minimizer.

Proposition 4.1. For any fixed $\lambda \in \mathbb{R}_{++}$ and $\Sigma \in \mathbb{S}^n_+$, the mapping $\mathbb{S}^n_{++} \to \mathbb{R}$, $S \mapsto L(\lambda, \Sigma, S)$ is strictly convex and has a unique minimizer, denoted by $S^*(\lambda, \Sigma)$. Moreover, $C_{\lambda} := \sup_{\Sigma \in \mathbb{S}^n_+} \|S^*(\lambda, \Sigma)\|_1 < \infty$ holds true for any $\lambda \in \mathbb{R}_{++}$.

The latter statement in the preceding proposition has the following implication. For any fixed $\lambda \in \mathbb{R}_{++}$, we can impose the constraint $||S|| \leq C_{\lambda}$ on the feasible set of the program (7) without affecting its optimal solution, where $C_{\lambda} \in \mathbb{R}_{++}$ is as in the preceding proposition (in particular, it is independent of $\Sigma \in \mathbb{S}_{+}^{n}$). Specifically, if we set

$$\mathcal{S}_{\lambda} := \left\{ S \in \mathbb{S}_{++}^{n} : \|S\| \le C_{\lambda} \right\},\tag{8}$$

then (7) can be equivalently written as

$$\min_{S \in \mathcal{S}_{\lambda}} L(\lambda, \Sigma, S) \tag{9}$$

for any $\Sigma \in \mathbb{S}^n_+$. Moreover, we can choose C_{λ} large enough such that

$$S^*(\lambda, \Sigma) \in \operatorname{int} \mathcal{S}_{\lambda} \quad \text{for all } \Sigma \in \mathbb{S}^n_+,$$
(10)

where "int" denotes the interior of a set.

4.2 Continuity of the minimizer in Σ

The following proposition shows that the mapping $\mathbb{S}^n_+ \to \mathbb{S}^n_{++}$, $S \mapsto L(\lambda, \Sigma, S)$ satisfies a certain growth condition at the minimizer $S^*(\lambda, \Sigma)$.

Proposition 4.2. For any $\Sigma \in \mathbb{S}^n_+$ and $\lambda \in \mathbb{R}_{++}$, there exist $\alpha_{\lambda,\Sigma}, \beta_{\lambda,\Sigma} \in \mathbb{R}_{++}$ (depending on λ and Σ) such that

$$L(\lambda, \Sigma, S) \ge L(\lambda, \Sigma, S^*(\lambda, \Sigma)) + \phi_{\lambda, \Sigma} (\|S - S^*(\lambda, \Sigma)\|) \quad \text{for all } S \in \mathbb{S}^n_{++}$$
(11)

for the function $\phi_{\lambda,\Sigma} : \mathbb{R}_+ \to \mathbb{R}_+$ defined by $\phi_{\lambda,\Sigma}(x) := \min\{\alpha_{\lambda,\Sigma}x^2, \beta_{\lambda,\Sigma}x\}.$

Using the growth condition (11), we can derive the following result.

Theorem 4.1. Let $\lambda \in \mathbb{R}_{++}$ and $\Sigma \in \mathbb{S}_{+}^{n}$. Moreover, let C_{λ} and $\alpha_{\lambda,\Sigma}$, $\beta_{\lambda,\Sigma}$ be as in (8) and Proposition 4.2, respectively. Then

$$\left\|S^{*}(\lambda,\Sigma) - S^{*}(\lambda,\Sigma')\right\| \leq \max\left\{\left(\frac{3C_{\lambda}}{\alpha_{\lambda,\Sigma}}\|\Sigma - \Sigma'\|\right)^{1/2}, \frac{3C_{\lambda}}{\beta_{\lambda,\Sigma}}\|\Sigma - \Sigma'\|\right\} \quad for \ all \ \Sigma' \in \mathbb{S}^{n}_{+}.$$
(12)

Theorem 4.1 shows that the mapping $\mathbb{S}^n_+ \to \mathbb{S}^n_{++}$, $\Sigma \mapsto S^*(\lambda, \Sigma)$ is continuous. However, for the proof of our main result, Theorem 5.3, we need (global) Lipschitz continuity. Therefore, in the next subsection we will use an implicit function theorem to derive (global) Lipschitz continuity (see Theorem 4.2). It is worth noting that the proof of Theorem 4.2 uses Theorem 4.1, see Step 1 in the proof of Lemma 10.2.

4.3 Lipschitz continuity of the minimizer in Σ

Let us assume that the constant C_{λ} in (8) is chosen so large that (10) holds. Then we can write down the first order condition on the minimizer of (9) without effect of the constraint $||S|| \leq C_{\lambda}$ as

$$\mathbf{0} \in \Sigma - S^{-1} + \lambda \,\partial \|S\|_1,\tag{13}$$

where $\partial \|S\|_1$ is the Clarke subdifferential of $\|\cdot\|_1$ at point S (defined by $\partial \|S\|_1 := \operatorname{conv}\{\lim \sup_{S_k \to S} \nabla \|S_k\|_1 : \nabla \|S_k\|_1$ exists for all $k \in \mathbb{N}\}$, where convA stands for the convex hull of a set A, see Section 2.1 in [8]). For any $\Sigma \in \mathbb{S}^n_+$, the minimizer $S^*(\lambda, \Sigma)$ of (9) satisfies (13). However, the right-hand side of (13) is a set-valued mapping in S. Therefore, we cannot directly use an implicit function theorem to derive Lipschitz continuity of the minimizer $S^*(\lambda, \Sigma)$ in Σ . To circumvent the difficulty, we propose to smooth the function $S \mapsto \|S\|_1$ by applying the simple smoothing function $h_{\varepsilon}(x) := \sqrt{x^2 + \varepsilon}$ to each summand $|S_{ij}|$ in the representation $\|S\|_1 = \sum_{1 \leq i, j \leq n} |S_{ij}|$, where $\varepsilon \in \mathbb{R}_{++}$. The resulting analogue of (9) is

$$\min_{S \in \mathcal{S}_{\lambda}} L_{\varepsilon}(\lambda, \Sigma, S), \tag{14}$$

where the mapping $L_{\varepsilon} : \mathbb{R}_{++} \times \mathbb{S}^n_+ \times \mathbb{S}^n_{++} \to \mathbb{R}$ is defined by

$$L_{\varepsilon}(\lambda, \Sigma, S) := \langle \Sigma, S \rangle - \log(\det S) + \lambda H_{\varepsilon}(S)$$

with $H_{\varepsilon}(S) = \sum_{1 \leq i,j \leq n} h_{\varepsilon}(S_{ij})$. Note that the mapping $S \mapsto L_{\varepsilon}(\lambda, \Sigma, S)$ is strictly convex in S and thus it has a unique minimizer (stationary point), denoted by $S_{\varepsilon}^*(\lambda, \Sigma)$. The next proposition yields that $S_{\varepsilon}^*(\lambda, \Sigma)$ converges to $S^*(\lambda, \Sigma)$ as $\varepsilon \downarrow 0$.

Proposition 4.3. For any $\lambda \in \mathbb{R}_{++}$ and $\Sigma \in \mathbb{S}_{+}^{n}$, we have

$$\left\|S^*(\lambda, \Sigma) - S^*_{\varepsilon}(\lambda, \Sigma)\right\| \le R^{-1}_{\lambda, \Sigma} (3n^2 \sqrt{\varepsilon}) \quad \text{for all } \varepsilon \in \mathbb{R}_{++},$$
(15)

where $R_{\lambda,\Sigma}^{-1}(t) := \inf\{\sigma \in \mathbb{R}_+ : R_{\lambda,\Sigma}(\sigma) = t\}, t \in \mathbb{R}_+, \text{ for the function } R_{\lambda,\Sigma} : \mathbb{R}_+ \to \mathbb{R} \text{ defined by}$

$$R_{\lambda,\Sigma}(\delta) := \inf_{S \in \mathcal{S}_{\lambda}: \, \|S - S^*(\lambda, \Sigma)\| \ge \delta} L(\lambda, \Sigma, S) - L(\lambda, \Sigma, S^*(\lambda, \Sigma))$$

which is continuous (by virtue of Berge's maximum theorem [5]), monotonically increasing, strictly positive for $\delta > 0$, and satisfies $R_{\lambda,\Sigma}(0) = 0$.

Since $S^*(\lambda, \Sigma)$ lies in the interior of S_{λ} , the proposition ensures that we can choose $\varepsilon \in \mathbb{R}_{++}$ so small that $S^*_{\varepsilon}(\lambda, \Sigma)$ also lies in the interior of S_{λ} . Let us assume that ε is chosen in this way. Then $S^*_{\varepsilon}(\lambda, \Sigma)$ satisfies the following first order optimality condition:

$$\mathbf{0} = \Sigma - S^{-1} + \lambda \nabla H_{\varepsilon}(S). \tag{16}$$

In Section 10.7 we show that the mapping $S_{\lambda} \to \mathbb{R}^{n \times n}$, $S \mapsto G_{\varepsilon}(\lambda, \Sigma, S) := \Sigma - S^{-1} + \lambda \nabla H_{\varepsilon}(S)$ induced by (16) is a strongly monotone operator (see Section 9.1 for the precise meaning). Therefore, we will be able to use the implicit function theorem in the form of Theorem 9.1 to derive the following result, which shows that the mapping $\mathbb{S}^n_+ \to \mathbb{S}^n_{++}$, $\Sigma \mapsto S^*(\lambda, \Sigma)$ is globally Lipschitz continuous. Recall that we use $S^*(\lambda, \Sigma)$ to denote the minimizer of (7) or, equivalently, of (9).

Theorem 4.2. For any $\lambda \in \mathbb{R}_{++}$ and $\kappa > (n/\lambda)^2$, the following inequality holds:

$$\left\|S^*(\lambda, \Sigma_1) - S^*(\lambda, \Sigma_2)\right\| \le \kappa \|\Sigma_1 - \Sigma_2\| \quad \text{for all } \Sigma_1, \Sigma_2 \in \mathbb{S}^n_+.$$
(17)

5 Distributional stability of \widehat{S}_N and of $\widehat{\Sigma}_N$ and its eigenvalues

In this section we show that the sample covariance matrix $\widehat{\Sigma}_N$, the eigenvalues of $\widehat{\Sigma}_N$ and the sparse precision matrix estimator \widehat{S}_N are distributionally stable.

Recall that the sample covariance matrix $\widehat{\Sigma}_N$, the simplified sample covariance matrix $\widetilde{\Sigma}_N$ and the sparse estimators \widehat{S}_N and \widetilde{S}_N of the precision matrix are defined by

$$\widehat{\Sigma}_{N}(\boldsymbol{x}) := \frac{1}{N} \sum_{i=1}^{N} x^{i} x^{i^{\mathsf{T}}} - \left(\frac{1}{N} \sum_{i=1}^{N} x^{i}\right) \left(\frac{1}{N} \sum_{i=1}^{N} x^{i}\right)^{\mathsf{T}}, \quad \widetilde{\Sigma}_{N}(\boldsymbol{x}) := \frac{1}{N} \sum_{i=1}^{N} x^{i} x^{i^{\mathsf{T}}}, \quad (18a)$$

$$\widehat{S}_{N}(\boldsymbol{x}) := \arg\min_{S \in \mathbb{S}_{++}^{n}} \left(\langle \widehat{\Sigma}_{N}(\boldsymbol{x}), S \rangle - \log(\det S) + \lambda \|S\|_{1} \right),$$
(18b)

$$\widetilde{S}_{N}(\boldsymbol{x}) := \arg\min_{S \in \mathbb{S}_{++}^{n}} \left(\langle \widetilde{\Sigma}_{N}(\boldsymbol{x}), S \rangle - \log(\det S) + \lambda \|S\|_{1} \right)$$
(18c)

for all $\boldsymbol{x} = (x^1, \ldots, x^N) \in \Omega$, where $\Omega := (\mathbb{R}^n)^N$ and $\lambda \in \mathbb{R}_{++}$ is a fixed constant. Proposition 4.1 ensures that $\widehat{S}_N(\boldsymbol{x})$ and $\widetilde{S}_N(\boldsymbol{x})$ are well defined for any $\boldsymbol{x} \in \Omega$. Recall that $\boldsymbol{\xi} = (\xi^1, \ldots, \xi^N)$ is used to denote the identity on Ω . The sample covariance matrix $\widehat{\Sigma}_N = \widehat{\Sigma}_N(\boldsymbol{\xi})$ is the standard nonparametric estimation of the covariance matrix Σ of ξ^1 when ξ^1, \ldots, ξ^n are i.i.d. (i.e. regarded as random variables under a product measure $\mathbb{P}^P = P^{\otimes N}$ on (Ω, \mathcal{F})). In the case where it is a priori known that the mean of ξ^1 is equal to $\mathbf{0}$, then it is more reasonable to use $\widetilde{\Sigma}_N = \widetilde{\Sigma}_N(\boldsymbol{\xi})$ as an estimator of the covariance matrix Σ of ξ^1 . The background of the estimator $\widehat{S}_N = \widehat{S}_N(\boldsymbol{\xi})$ of the precision matrix Σ^{-1} of ξ^1 was already discussed in Section 1.

Recall from Section 2 that $d|_{X,p}$ is used to denote the *p*-th Fortet-Mourier metric on a linear space X. In the following theorem, the role of X is played by \mathbb{S}^n_+ and \mathbb{R}^n . To simplify the presentation, we will write $d|_{K}$ and $d|_2$ instead of $d|_{\mathbb{S}^n_+,1}$ and $d|_{\mathbb{R}^n,2}$, respectively (the subscript K refers to Kantorovich). As before (see Section 3), m_P and m_Q denote the first absolute moments of P and Q, respectively. Moreover, we will use Σ_P to denote the covariance matrix of ξ^1 under P.

Theorem 5.1. For any $P, Q \in \mathscr{P}_2(\mathbb{R}^n)$ and $N \in \mathbb{N}$, the following inequalities hold:

$$\mathsf{dl}_{\mathsf{K}}\left(\mathbb{P}^{P}\circ\widehat{\Sigma}_{N}^{-1},\mathbb{P}^{Q}\circ\widehat{\Sigma}_{N}^{-1}\right) \leq \max\{3,2m_{P},2m_{Q}\}\,\mathsf{dl}_{2}(P,Q),\tag{19a}$$

$$\mathsf{dl}_{\mathsf{K}}\left(\mathbb{P}^{P} \circ \widetilde{\Sigma}_{N}^{-1}, \mathbb{P}^{Q} \circ \widetilde{\Sigma}_{N}^{-1}\right) \leq 2 \,\mathsf{dl}_{2}(P, Q),\tag{19b}$$

$$\mathsf{dl}_{\mathsf{K}}(\mathbb{P}^{P} \circ \Sigma_{N}^{-1}, \delta_{\Sigma_{P}}) \leq \mathbb{E}^{P}[\|\Sigma_{N} - \Sigma_{P}\|],$$
(19c)

$$\mathsf{dl}_{K}\left(\mathbb{P}^{P}\circ\widetilde{\Sigma}_{N}^{-1},\delta_{\Sigma_{P}}\right) \leq \mathbb{E}^{P}\left[\|\widetilde{\Sigma}_{N}-\Sigma_{P}\|\right].$$
(19d)

In the following theorem we extend (19a) and (19c), where $\widehat{\mu}_N$ is used to denote the standard nonparametric estimator of the mean vector, i.e. $\widehat{\mu}_N(\boldsymbol{x}) := \frac{1}{N} \sum_{i=1}^N x^i$ for all $\boldsymbol{x} = (x^1, \ldots, x^N) \in \Omega$. By dl_K we will mean the Fortet-Mourier metric dl_{1,ℝⁿ×Sⁿ₊} of order 1, where the Cartesian product $\mathbb{R}^n \times \mathbb{S}^n_+$ is equipped with the norm $\|(\mu, \Sigma)\|_1 := \|\mu\| + \|\Sigma\|$, and μ_P is used to denote the mean vector of ξ^1 under P.

Theorem 5.2. For any $P, Q \in \mathscr{P}_2(\mathbb{R}^n)$ and $N \in \mathbb{N}$, the following inequalities hold:

$$\mathsf{dI}_{K}\left(\mathbb{P}^{P}\circ(\widehat{\mu}_{N},\widehat{\Sigma}_{N})^{-1},\mathbb{P}^{Q}\circ(\widehat{\mu}_{N},\widehat{\Sigma}_{N})^{-1}\right) \leq \max\{4,2m_{P},2m_{Q}\}\,\mathsf{dI}_{2}(P,Q),\tag{20a}$$

$$\mathsf{dI}_{\mathrm{K}}\big(\mathbb{P}^{P}\circ(\widehat{\mu}_{N},\widehat{\Sigma}_{N})^{-1},\delta_{(\mu_{P},\Sigma_{P})}\big) \leq \mathbb{E}^{P}\big[\|\widehat{\mu}_{N}-\mu_{P}\|\big] + \mathbb{E}^{P}\big[\|\widehat{\Sigma}_{N}-\Sigma_{P}\|\big].$$
(20b)

In a sense, the following theorem is the main result of this paper. In the theorem, we write dI_K and dI_2 instead of $dI_{\mathbb{S}^n_{++},1}$ and $dI_{\mathbb{R}^n,2}$, respectively.

Theorem 5.3. For any $P, Q \in \mathscr{P}_2(\mathbb{R}^n)$, $N \in \mathbb{N}$ and $\kappa > (n/\lambda)^2$, the following inequalities hold:

$$\mathsf{dl}_{\mathsf{K}}\left(\mathbb{P}^{P}\circ\widehat{S}_{N}^{-1},\mathbb{P}^{Q}\circ\widehat{S}_{N}^{-1}\right) \leq \kappa \max\{3,2m_{P},2m_{Q}\}\,\mathsf{dl}_{2}(P,Q),\tag{21a}$$

$$\mathsf{dl}_{K}\left(\mathbb{P}^{P} \circ S_{N}^{-1}, \mathbb{P}^{Q} \circ S_{N}^{-1}\right) \leq 2\kappa \,\mathsf{dl}_{2}(P,Q),\tag{21b}$$

For any $P \in \mathscr{P}_2(\mathbb{R}^n)$, for which $S_P := \Sigma_P^{-1}$ exists, and for any $N \in \mathbb{N}$ and $\kappa > (n/\lambda)^2$, the following inequalities hold:

$$\mathsf{dI}_{K}\left(\mathbb{P}^{P}\circ\widehat{S}_{N}^{-1},\delta_{S_{P}}\right) \leq \kappa \mathbb{E}^{P}\left[\|\widehat{\Sigma}_{N}-\Sigma_{P}\|\right],\tag{22a}$$

$$\mathsf{dl}_{\mathsf{K}}\big(\mathbb{P}^{P} \circ \widetilde{S}_{N}^{-1}, \delta_{S_{P}}\big) \leq \kappa \mathbb{E}^{P}\big[\big\|\widetilde{\Sigma}_{N} - \Sigma_{P}\big\|\big].$$
(22b)

The following proposition tells us how quickly the right-hand sides in (19c), (19d), (20b), (22a), (22b) (and (24c), (24d)) converge to 0 as $N \to \infty$.

Proposition 5.1. Let $r \in [1, 2)$. Then

$$\lim_{N \to \infty} N^{(r-1)/r} \mathbb{E}^P \left[\| \widehat{\mu}_N - \mu_P \|_1 \right] = 0 \quad \text{for all } P \in \mathscr{P}_r(\mathbb{R}^n),$$
(23a)

$$\lim_{N \to \infty} N^{(r-1)/r} \mathbb{E}^{P} \left[\| \widehat{\Sigma}_{N} - \Sigma_{P} \|_{1} \right] = 0 \quad \text{for all } P \in \mathscr{P}_{2r}(\mathbb{R}^{n}),$$
(23b)

$$\lim_{N \to \infty} N^{(r-1)/r} \mathbb{E}^P \left[\| \widetilde{\Sigma}_N - \Sigma_P \|_1 \right] = 0 \quad \text{for all } P \in \mathscr{P}_{2r}(\mathbb{R}^n) \text{ with } \mu_P = \mathbf{0}.$$
(23c)

In particular, the same statements hold true if $\|\cdot\|_1$ is replaced by any other matrix norm (such as the Frobenius norm $\|\cdot\|$).

Let us use $\lambda_1(\widehat{\Sigma}_N(\boldsymbol{x})), \ldots, \lambda_n(\widehat{\Sigma}_N(\boldsymbol{x}))$ to denote the eigenvalues of $\widehat{\Sigma}_N(\boldsymbol{x})$ in decreasing order (i.e. $\lambda_1(\widehat{\Sigma}_N(\boldsymbol{x})) \geq \cdots \geq \lambda_n(\widehat{\Sigma}_N(\boldsymbol{x}))$) for all $\boldsymbol{x} = (x^1, \ldots, x^N) \in \Omega$. For the sake of notational simplicity we set

$$\widehat{\lambda}_{1,N}(\boldsymbol{x}) := \lambda_1 \big(\widehat{\Sigma}_N(\boldsymbol{x}) \big), \ \ldots, \ \widehat{\lambda}_{n,N}(\boldsymbol{x}) := \lambda_N \big(\widehat{\Sigma}_N(\boldsymbol{x}) \big)$$

for all $\boldsymbol{x} = (x^1, \ldots, x^N) \in \Omega$. Thus, $\widehat{\lambda}_{1,N}, \ldots, \widehat{\lambda}_{n,N}$ are the standard nonparametric estimators of the eigenvalues $\lambda_1, \ldots, \lambda_n$ of the true covariance matrix Σ in decreasing order (i.e. $\lambda_1 \geq \cdots \geq \lambda_n$).

Moreover, let the estimators $\widetilde{\lambda}_{1,N}, \ldots, \widetilde{\lambda}_{n,N}$ be defined in the same way, but with $\widehat{\Sigma}_N$ replaced by $\widetilde{\Sigma}_N$. In the following theorem, dl_K is use to denote the first order Fortet-Mourier metric $\mathsf{dl}_{\mathbb{R}^n,1}$ on $\mathscr{P}_1(\mathbb{R}^n)$, whereas dl_2 is as before the second order Fortet-Mourier metric $\mathsf{dl}_{\mathbb{R}^n,2}$ on $\mathscr{P}_2(\mathbb{R}^n)$. Moreover, we will use $\lambda_1^P, \ldots, \lambda_n^P$ to denote the eigenvalues of the covariance matrix Σ of ξ^1 under P, i.e. of Σ_P , in decreasing order (i.e. $\lambda_1^P \geq \cdots \geq \lambda_n^P$).

Theorem 5.4. For any $P, Q \in \mathscr{P}_2(\mathbb{R}^n)$ and $N \in \mathbb{N}$, the following inequalities hold:

$$\mathsf{dl}_{\mathsf{K}} \big(\mathbb{P}^{P} \circ (\widehat{\lambda}_{1,N}, \dots, \widehat{\lambda}_{n,N})^{-1}, \mathbb{P}^{Q} \circ (\widehat{\lambda}_{1,N}, \dots, \widehat{\lambda}_{n,N})^{-1} \big) \\ \leq \max\{3, 2m_{P}, 2m_{Q}\} \, \mathsf{dl}_{2}(P, Q),$$
(24a)

$$\mathsf{dl}_{\mathsf{K}}\big(\mathbb{P}^{P}\circ(\widetilde{\lambda}_{1,N},\ldots,\widetilde{\lambda}_{n,N})^{-1},\mathbb{P}^{Q}\circ(\widetilde{\lambda}_{1,N},\ldots,\widetilde{\lambda}_{n,N})^{-1}\big) \leq 2\,\mathsf{dl}_{2}(P,Q),\tag{24b}$$

$$\mathsf{dl}_{\mathsf{K}}\big(\mathbb{P}^{P}\circ(\widehat{\lambda}_{1,N},\ldots,\widehat{\lambda}_{n,N})^{-1},\delta_{(\lambda_{1}^{P},\ldots,\lambda_{n}^{P})}\big) \leq \mathbb{E}^{P}\big[\|\widehat{\Sigma}_{N}-\Sigma_{P}\|\big],\tag{24c}$$

$$\mathsf{dl}_{\mathrm{K}}\left(\mathbb{P}^{P}\circ(\widetilde{\lambda}_{1,N},\ldots,\widetilde{\lambda}_{n,N})^{-1},\delta_{(\lambda_{1}^{P},\ldots,\lambda_{n}^{P})}\right) \leq \mathbb{E}^{P}\left[\|\widetilde{\Sigma}_{N}-\Sigma_{P}\|\right].$$
(24d)

Remark 5.1. In view of Proposition 3.1, Theorem 5.4 yields in particular that (24a) and (24c), each with an additional factor L on the right-hand side, are still valid when $(\hat{\lambda}_{1,N},\ldots,\hat{\lambda}_{n,N})$ and $(\lambda_1^P,\ldots,\lambda_n^P)$ are replaced by $g(\hat{\lambda}_{1,N},\ldots,\hat{\lambda}_{n,N})$ and $g(\lambda_1^P,\ldots,\lambda_n^P)$, respectively, where $g: \mathbb{R}^n \to \mathbb{R}^k$ is a Lipschitz continuous function with Lipschitz constant L. Of course, in this case we have to choose $dI_K := dI_{1,\mathbb{R}^k}$. An example for k = 1 and L = 1 is obtained by setting $g(x_1,\ldots,x_n) := x_i$, which corresponds to the *i*-th largest eigenvalue. An example for k = 1 and $L = \sqrt{2}$ is obtained by setting $g(x_1,\ldots,x_n) := x_1 - x_2$, which corresponds to the spectral gap. The same applies to (24b) and (24d).

6 Two applications

6.1 Gaussian graphical model selection

One of the motivation of sparse precision matrix estimation is the Gaussian graphical model selection problem. We give a brief review of the problem here following Section 2.1 of [41]. Let ξ be an \mathbb{R}^n -valued random variable and assume that its distribution is given by a non-degenerate *n*-variate normal distribution with zero mean vector and precision matrix S. That is, the distribution of ξ can be parameterized by $S \in \mathbb{S}^n_{++}$. The true precision matrix S is unknown. In a Gauss-Markov random field model, the Gaussian random variable ξ is associated with an $n \times n$ undirected graph G = (V, E), where $V := \{1, \ldots, n\}$ is the set of vertices and $E \subseteq \{1, \ldots, n\}^2$ is the set of edges for which it is assumed that $(i, j) \notin E$ if and only if $S_{ij} = 0$, i.e. $E := \{(i, j) : i, j \in V \text{ with } S_{ij} \neq 0\}$.

In most practical applications, the set E has to be estimated statistically, often assuming that the estimator used is based on i.i.d. copies ξ^1, \ldots, ξ^N of ξ . In most situations, the graph is assumed to be sparse. In this case, the problem of the Gaussian graphical model selection, i.e. determining the edge structure of the graph G, reduces to finding the sparsity of the precision matrix S of ξ . In this setting, the main concern is whether the entries of the precision matrix estimator correctly recovers the *zero*-behavior of its true counterpart. As already mentioned in Section 1, the estimator \widetilde{S}_N introduced in (18c) is a reasonable choice for the estimator of a sparse precision matrix S. As before we consider the nonparametric statistical model $(\Omega, \mathcal{F}, \{\mathbb{P}^P : P \in \mathscr{P}_2(\mathbb{R}^n)\})$ introduced in Section 5 and assume that ξ^1, \ldots, ξ^N are the coordinate projections on $\Omega = (\mathbb{R}^n)^N$. Recall that $\mathbb{P}^P := \mathbb{P}^{\otimes N}$.

Let us use E_P and S_P to denote the set of edges and the precision matrix, respectively, when the underlying distribution of ξ is given by P. To know that the set E_P is well recovered (for large sample size N), one needs to make sure that the distribution $\mathbb{P}^P \circ \widetilde{S}_N^{-1}$ of \widetilde{S}_N is close to δ_{S_P} (when Nis large). Inequality (22b) and Proposition 5.1 indicate that this is true. For any $P, Q \in \mathscr{P}_2(\mathbb{R}^n)$ and $\kappa > (n/\lambda)^2$, we also have

$$\mathsf{dl}_{\mathsf{K}}\big(\mathbb{P}^{Q}\circ\widetilde{S}_{N}^{-1},\delta_{S_{P}}\big) \leq 2\kappa\,\mathsf{dl}_{2}(P,Q) + \kappa\,\mathbb{E}^{P}\big[\|\widetilde{\Sigma}_{N}-\Sigma_{P}\|\big],\tag{25}$$

which follows from the triangle inequality and (21b) and (22b). Inequality (25) shows that S_P (and thus E_P) can be reasonably recovered even if the data are drawn from a "contaminated" distribution Q that is slightly different from P. By Proposition 5.1, we know that $\lim_{N\to\infty} N^{(r-1)/r} \mathbb{E}^P[\|\widetilde{\Sigma}_N - \Sigma_P\|] = 0$ for all $r \in [1, 2)$; take into account that $\|\cdot\| \leq \|\cdot\|_1$.

6.2 Portfolio optimization

In the previous subsection, we discussed an application of Theorem 5.3. To also discuss an application of Theorem 5.1, let us consider a portfolio optimization problem. Let $\xi = (\xi^1, \ldots, \xi^n)$ be an \mathbb{R}^n -valued random variable whose coordinates ξ^1, \ldots, ξ^n model the one-period net returns of n assets traded on a financial market. An agent is to invest a capital of size 1 into the n assets in such a way that the portfolio risk (i.e. the variance of the portfolio's return) is minimized while the expected portfolio's return is set to a given level $z \in \mathbb{R}_{++}$. More precisely, if μ and Σ are used to denote the mean vector and the covariance matrix of ξ , respectively, the agent is to determine

$$v(\mu, \Sigma) := \min_{w \in \mathbb{R}^n} \frac{1}{2} w^{\mathsf{T}} \Sigma w \quad \text{subject to} \quad w^{\mathsf{T}} \mu = z, \ w^{\mathsf{T}} \mathbf{1} = 1, \ w \ge 0,$$
(26)

where $\mathbf{1} := (1, \ldots, 1)^{\mathsf{T}}$ and it is assumed that $\mu \in \mathbb{R}^n$ and $\Sigma \in \mathbb{S}^n_+$. Since the feasible set is compact, the minimum in (26) is attained.

In practice, the distribution of ξ is unknown which means that μ and Σ have to be estimated statistically, often assuming that the estimators used are based on i.i.d. copies ξ^1, \ldots, ξ^N of ξ . As before we consider the nonparametric statistical model $(\Omega, \mathcal{F}, \{\mathbb{P}^P : P \in \mathscr{P}_2(\mathbb{R}^n)\})$ introduced in Section 5 and assume that ξ^1, \ldots, ξ^N are the coordinate projections on $\Omega = (\mathbb{R}^n)^N$. Recall that $\mathbb{P}^P := \mathbb{P}^{\otimes N}$. Replacing μ and Σ in (26) with the empirical mean vector $\hat{\mu}_N = \hat{\mu}_N(\boldsymbol{\xi})$ and the empirical covariance matrix $\hat{\Sigma}_N = \hat{\Sigma}_N(\boldsymbol{\xi})$, respectively, leads to

$$v(\widehat{\mu}_N, \widehat{\Sigma}_N) := \min_{w \in \mathbb{R}^n} \frac{1}{2} w^{\mathsf{T}} \widehat{\Sigma}_N w$$
 subject to $w^{\mathsf{T}} \widehat{\mu}_N = z, \ w^{\mathsf{T}} \mathbf{1} = 1, \ w \ge 0,$

where $\widehat{\mu}_N(\boldsymbol{x}) := \frac{1}{N} \sum_{i=1}^N x^i, \, \boldsymbol{x} = (x^1, \dots, x^N) \in \Omega.$ For notational simplicity, we set $\widehat{v}_N := v(\widehat{\mu}_N, \widehat{\Sigma}_N).$

For any $C_1, C_2 \in \mathbb{R}_{++}$ with $C_1 < C_2$, let $\mathscr{P}_{C_1,C_2}(\mathbb{R}^n)$ be the set of all $P \in \mathscr{P}(\mathbb{R}^n)$ that have no mass outside the set $\{x \in \mathbb{R}^n : (C_1^2 + \langle x, \mathbf{1} \rangle^2 / n)^{1/2} \leq \|x\| \leq C_2\}$. Note that $\langle \mathbf{1}, x \rangle / n^{1/2} \leq (\|x\|_1 / n^{1/2} \leq n^{1/2} \|x\| / n^{1/2} =) \|x\|$ for all $x \in \mathbb{R}^n$, and $\langle \mathbf{1}, x \rangle / n^{1/2} = \|x\|$ if and only if $x = a\mathbf{1}$ for some $a \in \mathbb{R}$. Therefore, for (very) small C_1 , the condition $(C_1^2 + \langle x, \mathbf{1} \rangle^2 / n)^{1/2} \leq \|x\|$ in the definition of \mathscr{P}_{C_1,C_2} rules out only a (very) small area around the set $\{a\mathbf{1}: a \in \mathbb{R}\}$. From an application perspective, this limitation is negligible, as one would never hold a portfolio consisting of several assets with the same net returns; in this case one would only invest in the asset with the lowest variance. **Proposition 6.1.** Let $C_1, C_2 \in \mathbb{R}_{++}$ be such that $C_1 < C_2$. Then the function $v : \mathbb{R}^n \times \mathbb{S}^n_+ \to \mathbb{R}$ is Lipschitz continuous on the subset $Y_{C_1,C_2} := \{(\mu, \Sigma) \in \mathbb{R}^n \times \mathbb{S}^n_+ : (C_1^2 + \langle \mu, \mathbf{1} \rangle^2 / n)^{1/2} \le \|\mu\| \le C_2, \|\Sigma\| \le 2C_2^2\}$ of $\mathbb{R}^n \times \mathbb{S}^n_+$ with Lipschitz constant $L_{C_1,C_2} := \frac{1}{2} + \frac{32C_2^2}{9C_1^2}(z + \frac{C_2}{\sqrt{n}}) + \frac{16C_2^2}{9\sqrt{n}C_1^2}$.

Now, let $C_1, C_2 \in \mathbb{R}_{++}$ be fixed such that $C_1 < C_2$ and let us use dl_K and dl_2 to denote $\mathsf{dl}_{\mathbb{R},1}$ and $\mathsf{dl}_{\mathbb{R}^n,2}$, respectively. Then, as a direct consequence of Theorem 5.2, Proposition 6.1 and Remark 3.1, we obtain

$$\mathsf{dl}_{\mathsf{K}}\left(\mathbb{P}^{P}\circ\hat{v}_{N}^{-1},\mathbb{P}^{Q}\circ\hat{v}_{N}^{-1}\right) \leq L_{C_{1},C_{2}}\max\{4,2m_{P},2m_{Q}\}\,\mathsf{dl}_{2}(P,Q)$$

$$\tag{27}$$

for all $P, Q \in \mathscr{P}_{C_1,C_2} (\subseteq \mathscr{P}_2(\mathbb{R}^n))$, where L_{C_1,C_2} is as in Proposition 6.1 and m_p and m_Q are bounded above by C_2 . Moreover, proceeding as in the proof of (24c) (see Section 10.12), using Proposition 6.1 instead of (31), we also obtain

$$\mathsf{dI}_{\mathrm{K}}\left(\mathbb{P}^{P}\circ\widehat{v}_{N}^{-1},\delta_{v(\mu_{P},\Sigma_{P})}\right) \leq L_{C_{1},C_{2}}\left(\mathbb{E}^{P}\left[\|\widehat{\mu}_{N}-\mu_{P}\|\right] + \mathbb{E}^{P}\left[\|\widehat{\Sigma}_{N}-\Sigma_{P}\|\right]\right)$$
(28)

for all $P, Q \in \mathscr{P}_{C_1, C_2}$. The triangle inequality and (27) and (28) imply

$$dl_{K} \left(\mathbb{P}^{Q} \circ \widehat{v}_{N}^{-1}, \delta_{v(\mu_{P}, \Sigma_{P})} \right) \leq L_{C_{1}, C_{2}} \max\{4, 2m_{P}, 2m_{Q}\} dl_{2}(P, Q)$$

$$+ L_{C_{1}, C_{2}} \left(\mathbb{E}^{P} \left[\|\widehat{\mu}_{N} - \mu_{P}\| \right] + \mathbb{E}^{P} \left[\|\widehat{\Sigma}_{N} - \Sigma_{P}\| \right] \right)$$

$$(29)$$

for all $P, Q \in \mathscr{P}_{C_1,C_2}$. Inequality (29) shows that $v(\mu_P, \Sigma_P)$ can be reasonably estimated even if the data are drawn from a "contaminated" distribution Q that is slightly different from P. Recall from Proposition 5.1 that for the second summand on the right-hand side of (29) we know that $\lim_{N\to\infty} N^{(r-1)/r}(\mathbb{E}^P[\|\widehat{\mu}_N - \mu_P\|] + \mathbb{E}^P[\|\widehat{\Sigma}_N - \Sigma_P\|]) = 0$ for all $r \in [1, 2)$, if $P, Q \in \mathscr{P}_{C_1,C_2}(\mathbb{R}^n)$ $(\subseteq \mathscr{P}_p(\mathbb{R}^n) \text{ for all } p \geq 1).$

7 Two numerical experiments

7.1 Distributional stability of the eigenvalues of the sample covariance matrix

Inequality (24a) (in Theorem 5.4) and Remark 5.1 show that the Kantorovich distance $\mathsf{dl}_{\mathsf{K}}(\mathbb{P}^P \circ \widehat{\lambda}_{i,N}^{-1}, \mathbb{P}^Q \circ \widehat{\lambda}_{i,N}^{-1})$ between the distributions of the sample eigenvalues under two different distributions P and Q grows at most linearly as $\mathsf{dl}_2(P, Q)$ deviates from zero. We conducted a numerical experiment, which confirms this finding.

In this experiment, we chose $P := N_{\mu,\Sigma}$ and $Q \in \mathscr{Q} := \{N_{\mu+\alpha\mu',\Sigma+\alpha\Sigma'} : \alpha \in [0,1]\}$, where $\mu, \mu' \in \mathbb{R}^n$ and $\Sigma, \Sigma' \in \mathbb{S}^n_+$ are fixed and $N_{\tilde{\mu},\tilde{\Sigma}}$ is used to denote the *n*-variate normal distribution with mean vector $\tilde{\mu}$ and covariance matrix $\tilde{\Sigma}$. To avoid an overly artificial setting, we have left the choice of the parameters μ, μ', Σ and Σ' to 2n+2 independent samples $\eta, \eta^1, \ldots, \eta^n, \gamma, \gamma^1, \ldots, \gamma^n$ of an *n*-variate standard normal distribution. Specifically, we chose n := 10 and N := 100, and set $\mu := \eta$, $\mu' := \gamma, \Sigma := \frac{1}{n} \sum_{i=1}^n \eta^i {\eta^i}^{\mathsf{T}}$ and $\Sigma' := \frac{1}{n} \sum_{i=1}^n \gamma^i {\gamma^i}^{\mathsf{T}}$.

Our actual goal was to visualize the dependence of $\mathsf{dl}_{\mathsf{K}}(\mathbb{P}^{P} \circ \widehat{\lambda}_{i,N}^{-1}, \mathbb{P}^{Q} \circ \widehat{\lambda}_{i,N}^{-1})$ on $\mathsf{dl}_{2}(P,Q), Q \in \mathscr{Q}$. Since $\mathsf{dl}_{2}(P,Q)$ is difficult to access numerically, we rather visualized in Figure 1 the dependence of $\mathsf{dl}_{\mathsf{K}}(\mathbb{P}^{P} \circ \widehat{\lambda}_{i,N}^{-1}, \mathbb{P}^{Q} \circ \widehat{\lambda}_{i,N}^{-1})$ on $\mathsf{dl}_{1}(P,Q)$ and $\overline{\mathsf{dl}}_{2}(P,Q) := \int_{\mathbb{R}^{n}} L_{2}(\hat{x}, \tilde{x}) \|\hat{x} - \tilde{x}\| \mathbb{P}^{P}(d\hat{x}) \mathbb{P}^{Q}(d\tilde{x})$, respectively.



Figure 1: $\widehat{\mathsf{dl}}_{\mathrm{K},M}(\mathbb{P}^P \circ \widehat{\lambda}_{i,N}^{-1}, \mathbb{P}^Q \circ \widehat{\lambda}_{i,N}^{-1})$ as a function of $\mathsf{dl}_1(P,Q), Q \in \mathscr{Q}$, (left) and of $\overline{\mathsf{dl}}_2(P,Q), Q \in \mathscr{Q}$, (right).

Here it is crucial that on the one hand $dl_1(P,Q) \leq dl_2(P,Q) \leq \overline{dl}_2(P,Q)$ (which is obvious) and on the other hand $\overline{dl}_2(P,Q)$ depends linearly on $dl_1(P,Q)$ (see Figure 2). Thus, Figure 1 indicates that $dl_K(\mathbb{P}^P \circ \widehat{\lambda}_{i,N}^{-1}, \mathbb{P}^Q \circ \widehat{\lambda}_{i,N}^{-1})$ depends also linearly on $dl_2(P,Q)$. In this context, it should be mentioned that there is a closed-form expression for $dl_1(P,Q)$ due to the assumption of normality (see [19, Proposition 7]) and that $\overline{dl}_2(P,Q)$ can be easily well approximated by Monte Carlo simulations.

Of course, the Kantorovich distance $\mathsf{dl}_{\mathsf{K}}(\mathbb{P}^P \circ \widehat{\lambda}_{i,N}^{-1}, \mathbb{P}^Q \circ \widehat{\lambda}_{i,N}^{-1})$ can hardly be determine explicitly and so we approximated it by means of standard Monte Carlo simulations based on M := 100 Monte Carlo repetitions. More precisely, we used $\widehat{\mathsf{dl}}_{\mathsf{K},M}(\mathbb{P}^P \circ \widehat{\lambda}_{i,N}^{-1}, \mathbb{P}^Q \circ \widehat{\lambda}_{i,N}^{-1}) := \mathsf{dl}_{\mathsf{K}}(\widehat{\Lambda}_{i,N;M}^P, \widehat{\Lambda}_{i,N;M}^Q)$ as an approximation of $\mathsf{dl}_{\mathsf{K}}(\mathbb{P}^P \circ \widehat{\lambda}_{i,N}^{-1}, \mathbb{P}^Q \circ \widehat{\lambda}_{i,N}^{-1})$, where $\widehat{\Lambda}_{i,N;M}^P$ and $\widehat{\Lambda}_{i,N;M}^Q$ denote the empirical distributions of the Monte Carlo samples $\widehat{\lambda}_{i,N;1}, \ldots, \widehat{\lambda}_{i,N;M}$ under P and Q, respectively. Note that the computation of the Kantorovich (Wasserstein) distance between two general distributions is pre-implemented in many relevant software packages.

7.2 Distributional stability of the optimal portfolio value

In the framework of Section 6.2, inequality (27) shows that the Kantorovich distance $\mathsf{dl}_{\mathsf{K}}(\mathbb{P}^P \circ \hat{v}_N^{-1}, \mathbb{P}^Q \circ \hat{v}_N^{-1})$ between the distributions of the optimal portfolio value under two different distributions P and Q grows at most linearly as $\mathsf{dl}_2(P, Q)$ deviates from zero. We conducted an numerical experiment, which confirms this finding.

In this experiment, we chose $P := LN_{\mu,\Sigma}$ and $Q \in \mathscr{Q} := \{LN_{\mu+\alpha\mu',\Sigma+\alpha\Sigma'} : \alpha \in [0,1]\}$, where $\mu, \mu' \in \mathbb{R}^n$ and $\Sigma, \Sigma \in \mathbb{S}^n_+$ are fixed and $LN_{\tilde{\mu},\tilde{\Sigma}}$ is used to denote the *n*-variate log-normal distribution with parameters $\tilde{\mu}$ and $\tilde{\Sigma}$. To avoid an overly artificial setting, we have left the choice of the parameters μ, μ', Σ and Σ' to two independent samples of an *n*-variate standard normal distribution (exactly as in Section 7.1), where we chose n := 5 and N := 1000. Analogous to Section 7.1, we approximated the



Figure 2: $\overline{\mathsf{dl}}_2(P,Q)$ as a function of $\mathsf{dl}_1(P,Q), Q \in$ Figure 3: $\widehat{\mathsf{dl}}_{\mathrm{K},M}(\mathbb{P}^P \circ \widehat{v}_N^{-1}, \mathbb{P}^Q \circ \widehat{v}_N^{-1})$ as a function \mathscr{Q} .

Kantorovich distance $\mathsf{dl}_{\mathsf{K}}(\mathbb{P}^P \circ \widehat{v}_N^{-1}, \mathbb{P}^Q \circ \widehat{v}_N^{-1})$ by a Monte Carlo approximation $\widehat{\mathsf{dl}}_{\mathsf{K},M}(\mathbb{P}^P \circ \widehat{v}_N^{-1}, \mathbb{P}^Q \circ \widehat{v}_N^{-1})$ based on M := 1000 Monte Carlo repetitions.

From Figure 3, we can see that the (approximated) Kantorovich distance $\widehat{\mathsf{dl}}_{\mathrm{K},M}(\mathbb{P}^P \circ \widehat{v}_N^{-1}, \mathbb{P}^Q \circ \widehat{v}_N^{-1})$ increases sub-linearly as $\mathsf{dl}_{\mathrm{K}}(P,Q)$ increases. Since $\mathsf{dl}_{\mathrm{K}} = \mathsf{dl}_1 \leq \mathsf{dl}_2$, this confirms Inequality (27).

8 Concluding remarks

Differing from the main stream research in matrix optimization, here we study the distributional behavior of statistical estimators of the covariance matrix and the precision matrix. Specifically we investigate how a perturbation of the sampling distribution may affect the performance of the sample covariance matrix and its eigenvalues and of a sparse estimator of the precision matrix. We do this by looking at the distributions of the mentioned estimators based on a perceived sampling distribution compared to the distributions based on the target sampling distribution. We are able to establish a sort of stability, which means it is "safe" to use the estimators when the perturbation of the sampling distribution is confined to some specified topological structure.

There are still a number of open questions that remain to be addressed. For instance, when in the penalty term in (1) the norm $\|\cdot\|_1$ is replaced by a different norm such as $\|\cdot\|_{1-}$, we are unable to assert Lipschitz continuity (in the spirit of (5)) of the resulting estimator \hat{S}_N and subsequently unable to obtain distributional stability of \hat{S}_N (in the spirit of (6)). Likewise, we are unable to assert distributional stability of the empirical estimators of various ratios of interest in the context of portfolio theory, including the Sharpe ratio, the Omega ratio and the Rachev ratio, as we are unable to establish Lipschitz continuity of these empirical estimators (in the spirit of (5)). It might be possible to derive distributional stability of these estimators using different techniques, or in some sense weaker than our current results, but we leave all this to future research.

9 Auxiliary results

9.1 An implicit function theorem

For the proof of Theorem 4.2 (see Section 10.7) we need the implicit function theorem in the form of Theorem 9.1 below. For its formulation we need some terminology. Let $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ be normed linear spaces. A function $f: X \to Y$ is said to be Lipschitz continuous on a given subset $\mathcal{X} \subseteq X$ if there exists a constant $L \in \mathbb{R}_{++}$ such that $\|f(x) - f(x')\|_Y \leq L_{\mathcal{X}} \|\hat{x} - \tilde{x}\|_X$ for all $\hat{x}, \tilde{x} \in \mathcal{X}$. In this case, we say that f is L-Lipschitz continuous on \mathcal{X} . If X is even a pre-Hilbert space with scalar product $\langle \cdot, \cdot \rangle_X$ and induced norm $\|\cdot\|_X$, then, for any convex subset $\mathcal{X} \subseteq X$, a mapping $f: X \to X$ is said to be a monotone operator on \mathcal{X} if $\langle f(\hat{x}) - f(\tilde{x}), \hat{x} - \tilde{x} \rangle_X \geq 0$ for all $\hat{x}, \tilde{x} \in \mathcal{X}$, and a strongly monotone operator on \mathcal{X} if there exists a constant $\rho \in \mathbb{R}_{++}$ such that $\langle f(\hat{x}) - f(\tilde{x}), \hat{x} - \tilde{x} \rangle_X \geq \rho \|\hat{x} - \tilde{x}\|_X^2$ for all $\hat{x}, \tilde{x} \in \mathcal{X}$. In the latter case we say that f is ρ -strongly monotone on \mathcal{X} . The following theorem, which is known from Theorem F.1 in [59], is an extension of Theorem 1H.3 in [13]. In the theorem, the Cartesian product $X_0 \times Y_0$ is equipped with the product topology induced by the respective norms.

Theorem 9.1. Let $(X, \|\cdot\|_X)$ be a Banach space and $X_0 \subseteq X$ be an open convex subset. Let $(Y, \langle \cdot, \cdot \rangle)$ be a Hilbert space and $Y_0 \subseteq Y$ be an open convex subset. Consider a function $f : X_0 \times Y_0 \to Y_0$ and a point $(\bar{x}, \bar{y}) \in \operatorname{int} X_0 \times Y_0$ satisfying $f(\bar{x}, \bar{y}) = \mathbf{0}$. Let $\rho, L \in \mathbb{R}_{++}$ be constants and assume that there exist a neighborhood $\mathcal{N}_0(\bar{x})$ of \bar{x} in X_0 and a convex neighborhood $\mathcal{N}_0(\bar{y})$ of \bar{y} in Y_0 such that the following conditions are met:

- (a) f is continuous on $\mathcal{N}_0(\bar{x}) \times \mathcal{N}_0(\bar{y})$.
- (b) For any $x \in \mathcal{N}_0(\bar{x})$, the function $f(x, \cdot)$ is ρ -strongly monotone on $\mathcal{N}_0(\bar{y})$.
- (c) For any $y \in \mathcal{N}_0(\bar{y})$, the function $f(\cdot, y)$ is L-Lipschitz continuous on $\mathcal{N}_0(\bar{x})$.

Let $s: X_0 \Rightarrow Y_0$ be the solution mapping of f, which is defined by $s(x) := \{y \in Y_0 : f(x,y) = \mathbf{0}\}$. Then there exists a neighborhood $\mathcal{N}(\bar{x}) \ (\subseteq \mathcal{N}_0(\bar{x}))$ of \bar{x} in X_0 , which only depends on $\mathcal{N}_0(\bar{x})$, ρ and L, such that

- (i) $s(\cdot)$ is a single-valued function on $\mathcal{N}(\bar{x})$,
- (ii) and this single-valued function is L/ρ -Lipschitz continuous on $\mathcal{N}(\bar{x})$.

9.2 Local Lipschitz continuity of covariance matrix and its eigenvalues and eigenvectors

Although the main focus of this paper is on stability of *estimators* for the covariance matrix and its eigenvalues (and the precision matrix), we will also discuss stability of the *true* covariance matrix and its eigenvalues in this section. Let us use $\xi = (\xi_1, \ldots, \xi_n)$ to denote the identity on \mathbb{R}^n . If we equip the measurable space $(\mathbb{R}^n, \mathcal{B}(\mathbb{R}^n))$ with some $P \in \mathscr{P}(\mathbb{R}^n)$, we can regard ξ as a random variable distributed according to P. For any $P \in \mathscr{P}_1(\mathbb{R}^n)$, we write μ_P for the expectation of ξ w.r.t. P, denoted by $\mathbb{E}_P[\cdot]$, i.e. $\mu_P = \mathbb{E}_P[\xi]$. For any $P \in \mathscr{P}_2(\mathbb{R}^n)$, we write Σ_P for the covariance matrix of

 ξ w.r.t. P, i.e. $\Sigma_P = \mathbb{E}_P[(\xi - \mathbb{E}_P[\xi])(\xi - \mathbb{E}_P[\xi])^{\mathsf{T}}]$. The following proposition shows in particular that the mapping $\mathscr{P}_2(\mathbb{R}^n) \to \mathbb{S}^n$, $P \mapsto \Sigma_P$ is locally Lipschitz continuous w.r.t. the Fortet-Mourier metric of order 2. The Fortet-Mourier metric $\mathsf{dl}_p := \mathsf{dl}_{\mathbb{R}^n,p}$ of arbitrary order $p \ge 1$ was introduced in (4).

Proposition 9.1. For any $P, Q \in \mathscr{P}_2(\mathbb{R}^n)$, we have

$$\begin{aligned} \|\Sigma_P - \Sigma_Q\| &\le n \, \mathrm{dl}_2(P, Q) + \sqrt{n} \big(\|\mu_P\| + \|\mu_Q\| \big) \mathrm{dl}_1(P, Q) \\ &\le \big(n + \sqrt{n} (\|\mu_P\| + \|\mu_Q\| \big) \mathrm{dl}_2(P, Q), \end{aligned} \tag{30}$$

Proof. For any $P, Q \in \mathscr{P}(\mathbb{R}^n)$, we obtain

$$\begin{split} \|\Sigma_{P} - \Sigma_{Q}\| &= \|\mathbb{E}_{P}[\xi\xi^{\mathsf{T}}] - \mu_{P}\mu_{P}^{\mathsf{T}} - \mathbb{E}_{Q}[\xi\xi^{\mathsf{T}}] + \mu_{Q}\mu_{Q}^{\mathsf{T}}\| \\ &\leq \|\mathbb{E}_{P}[\xi\xi^{\mathsf{T}}] - \mathbb{E}_{Q}[\xi\xi^{\mathsf{T}}]\| + \|\mu_{P}\mu_{P}^{\mathsf{T}} - \mu_{Q}\mu_{Q}^{\mathsf{T}}\| \\ &\leq \left(\mathbb{E}_{P}[\xi\xi^{\mathsf{T}}] - \mathbb{E}_{Q}[\xi\xi^{\mathsf{T}}]\| + \|\mu_{P}\|\|\mu_{P} - \mu_{Q}\| + \|\mu_{Q}\|\|\mu_{P} - \mu_{Q}\| \\ &\leq \left(\sum_{i=1}^{n}\sum_{j=1}^{n}\left(\mathbb{E}_{P}[\xi_{i}\xi_{j}] - \mathbb{E}_{Q}[\xi_{i}\xi_{j}]\right)^{2}\right)^{1/2} + \left(\|\mu_{P}\| + \|\mu_{Q}\|\right)\|\mu_{P} - \mu_{Q}\| \\ &\leq \left(\sum_{i=1}^{n}\sum_{j=1}^{n}\left(\int x_{i}x_{j}\left(P - Q\right)\left(d(x_{1}, \dots, x_{n})\right)\right)^{2}\right)^{1/2} \\ &+ \left(\|\mu_{P}\| + \|\mu_{Q}\|\right)\left(\sum_{i=1}^{n}\left(\int x_{i}\left(P - Q\right)\left(d(x_{1}, \dots, x_{n})\right)\right)^{2}\right)^{1/2} \\ &\leq \sqrt{n^{2}\mathsf{dl}_{2}(P, Q)^{2}} + \left(\|\mu_{P}\| + \|\mu_{Q}\|\right)\mathsf{dl}_{2}(P, Q) \\ &\leq n\,\mathsf{dl}_{2}(P, Q) + \sqrt{n}\,\mathsf{dl}_{1}(P, Q)\left(\|\mu_{P}\| + \|\mu_{Q}\|\right), \end{split}$$

where we use the Cauchy-Schwarz inequality and that the mapping $(x_1, \ldots, x_n) \mapsto x_i x_j$ is contained in $\mathcal{F}_2(\mathbb{R}^n)$ and the mapping $(x_1, \ldots, x_n) \mapsto x_i$ is contained in $\mathcal{F}_1(\mathbb{R}^n)$. This proves the first inequality in (30). The second inequality in (30) is trivial.

r	-	-	-	-	
I					
I					
I					

Proposition 9.1 can be combined with the following results known from the literature to obtain stability results for the eigenvalues and eigenvectors of Σ_P . Let $\Sigma_1, \Sigma_2 \in \mathbb{S}^n$. For j = 1, 2, let $\lambda_j^1, \ldots, \lambda_j^n$ be the eigenvalues of Σ_j in decreasing order (i.e. $\lambda_j^1 \ge \lambda_j^2 \ge \cdots \ge \lambda_j^n$) and set $\Lambda_j = \text{diag}(\lambda_j^1, \ldots, \lambda_j^n)$. Then, for each $i = 1, \ldots, n$, we have

$$|\lambda_1^i - \lambda_2^i| \le \|\Lambda_1 - \Lambda_2\| \le \|\Sigma_1 - \Sigma_2\|, \tag{31}$$

see Fact 4 on p. 15-2 in [36]. Moreover, for each $i = 1, \ldots, n$ with $\lambda_1^{i+1} < \lambda_1^i < \lambda_1^{i-1}$ (where $\lambda_1^0 := \infty$, $\lambda_1^{n+1} := -\infty$), we have

$$\|v_1^i - v_2^i\|_2 \le \frac{2^{3/2} \|\Sigma_1 - \Sigma_2\|_2}{\min\{\lambda_1^{i-1} - \lambda_1^i, \lambda_1^i - \lambda_1^{i+1}\}}$$
(32)

for any eigenvectors v_1^i and v_2^i of $(\Sigma_1 \text{ and } \Sigma_2)$ associated with λ_1^i and λ_2^i , respectively, which satisfy $\langle v_1^i, v_2^i \rangle \geq 0$. This is known from Corollary 3 (to a version of the classical Davis-Kahan sin θ theorem [11]) in [55]. The condition $\langle v_1^i, v_2^i \rangle \geq 0$ is always attainable by changing the signs of eigenvectors properly.

10 Proofs

10.1 Minimizer of (1) for $\lambda = 0$

For the convenience of the reader, we recall here the well-known argument why \widehat{S}_N defined by (1) in the case $\lambda = 0$ is given by the sample precision matrix $\widehat{\Sigma}_N^{-1}$, provided $\widehat{\Sigma}_N$ is contained in \mathbb{S}_{++}^n . For $\lambda = 0$ the expression to be minimised on the right-hand side in (1) coincides with the log-likelihood function in the case where ξ is *n*-variate Gaussian with (arbitrary mean vector and) covariance matrix S^{-1} , and the corresponding maximum likelihood estimator for the covariance matrix S^{-1} is known to be $\widehat{\Sigma}_N$. That is, the minimizer of the expression on the right-hand side in (1) over all S^{-1} , $S \in \mathbb{S}_{++}^n$, is $S_*^{-1} := \widehat{\Sigma}_N$, i.e. $S_* = \widehat{\Sigma}_N^{-1}$. Thus, $\widehat{S}_N = \widehat{\Sigma}_N^{-1}$.

10.2 Proof of Theorem 3.1

The proof of Theorem 3.1 relies on the following lemma. The lemma involves the class Ψ_{κ_1,κ_2} of all functions $\psi: X^N \to \mathbb{R}$ that satisfy

$$\begin{aligned} \left| \psi(\hat{\boldsymbol{x}}) - \psi(\tilde{\boldsymbol{x}}) \right| \\ &\leq \frac{\kappa_1}{N} \sum_{i=1}^N L_2(\hat{x}^i, \tilde{x}^i) \| \hat{x}^i - \tilde{x}^i \|_X + \frac{\kappa_2}{N^2} \sum_{i=1}^N \left(\| \hat{x}^i \|_X + \| \tilde{x}^i \|_X \right) \sum_{k=1}^N \| \hat{x}^k - \tilde{x}^k \|_X \end{aligned}$$

for all $\hat{\boldsymbol{x}} = (\hat{x}^1, \dots, \hat{x}^N), \ \tilde{\boldsymbol{x}} = (\tilde{x}^1, \dots, \tilde{x}^N) \in X^N$. Recall that $\kappa_1, \kappa_2 \in \mathbb{R}_+$ are given constants.

Lemma 10.1. For any $P, Q \in \mathscr{P}_2(X)$, we have

$$\sup_{\psi \in \Psi_{\kappa_1,\kappa_2}} \left| \int_{X^N} \psi \, \mathrm{d}\mathbb{P}^P - \int_{X^N} \psi \, \mathrm{d}\mathbb{P}^Q \right| \le \max\left\{ \kappa_1 + \kappa_2, 2m_P, 2m_Q \right\} \mathsf{dl}_2(P,Q)$$

Proof. The proof is similar to [20, Lemma 1]. Here we give a sketch of it to facilitate reading. Let $\boldsymbol{x}_{-j} := (x^1, \ldots, x^{j-1}, x^{j+1}, \ldots, x^N)$ for any $\boldsymbol{x} = (x^1, \ldots, x^N) \in X^N$ and $j = 1, \ldots, N$. Moreover, for any $\boldsymbol{P} = (P_1, \ldots, P_N) \in \mathscr{P}_2(X)^N$, $\psi \in \Psi_{\kappa_1,\kappa_2}$ and $j = 1, \ldots, N$, write $\boldsymbol{P}_{-j}(\mathrm{d}\boldsymbol{x}_{-j})$ for $P_1(\mathrm{d}x^1) \cdots P_{j-1}(\mathrm{d}x^{j-1})P_{j+1}(\mathrm{d}x^{j+1}) \cdots P_N(\mathrm{d}x^N)$ and set $\psi_{\boldsymbol{x}_{-j}}(x^j) := \int_{X^{N-1}} \psi(\boldsymbol{x}_{-j}, x^j) \boldsymbol{P}_{-j}(\mathrm{d}\boldsymbol{x}_{-j})$ for any $\boldsymbol{x} \in X^N$ and $x^j \in X$. Then, for any $\boldsymbol{P} = (P_1, \ldots, P_N) \in \mathscr{P}_2(X)^N$, $\psi \in \Psi_{\kappa_1,\kappa_2}$ and $j = 1, \ldots, N$, we obtain

$$\begin{aligned} \left| \psi_{\boldsymbol{x}_{-j}}(\hat{x}^{j}) - \psi_{\boldsymbol{x}_{-j}}(\tilde{x}^{j}) \right| \\ &\leq \int_{X^{N}} \left| \psi(\boldsymbol{x}_{-j}, \hat{x}^{j}) - \psi(\boldsymbol{x}_{-j}, \tilde{x}^{j}) \right| \boldsymbol{P}_{-j}(\mathrm{d}\boldsymbol{x}_{-j}) \\ &\leq \int_{X^{N}} \left(\frac{\kappa_{1}}{N} L_{2}(\hat{x}^{j}, \tilde{x}^{j}) \| \hat{x}^{j} - \tilde{x}^{j} \|_{X} \\ &\quad + \frac{\kappa_{2}}{N^{2}} \| \hat{x}^{j} - \tilde{x}^{j} \| \left(\| \hat{x}^{j} \|_{X} + \| \tilde{x}^{j} \|_{X} + \sum_{i=1, i \neq j}^{N} 2 \| x^{i} \|_{X} \right) \right) \boldsymbol{P}_{-j}(\mathrm{d}\boldsymbol{x}_{-j}) \\ &= \frac{\kappa_{1}}{N} L_{2}(\hat{x}^{j}, \tilde{x}^{j}) \| \hat{x}^{j} - \tilde{x}^{j} \|_{X} + \frac{\kappa_{2}}{N^{2}} \| \hat{x}^{j} - \tilde{x}^{j} \|_{X} \left(\| \hat{x}^{j} \|_{X} + \| \tilde{x}^{j} \|_{X} \right) \end{aligned}$$

$$\begin{aligned} &+ \frac{2\kappa_2}{N^2} \sum_{i=1, i\neq j}^N \|\hat{x}^j - \tilde{x}^j\|_X \int_{X^N} \|x^i\|_X P_{-j}(\mathrm{d}\boldsymbol{x}_{-j}) \\ &= \frac{\kappa_1}{N} L_2(\hat{x}^j, \tilde{x}^j) \|\hat{x}^j - \tilde{x}^j\|_X + \frac{\kappa_2}{N^2} \|\hat{x}^j - \tilde{x}^j\|_X \left(\|\hat{x}^j\|_X + \|\tilde{x}^j\|_X\right) \\ &+ \frac{2\kappa_2}{N^2} \|\hat{x}^j - \tilde{x}^j\|_X \sum_{i=1, i\neq j}^N \int_{X^N} \|x^i\|_X P_i(\mathrm{d}x^i) \\ &= \frac{\kappa_1}{N} L_2(\hat{x}^j, \tilde{x}^j) \|\hat{x}^j - \tilde{x}^j\|_X + \frac{\kappa_2}{N^2} \|\hat{x}^j - \tilde{x}^j\|_X \left(\|\hat{x}^j\|_X + \|\tilde{x}^j\|_X\right) \\ &+ \frac{2\kappa_2}{N^2} \|\hat{x}^j - \tilde{x}^j\|_X \sum_{i=1, i\neq j}^N m_{P_i} \\ &\leq \frac{\kappa_1}{N} L_2(\hat{x}^j, \tilde{x}^j) \|\hat{x}^j - \tilde{x}^j\|_X + \frac{2\kappa_2}{N^2} L_2(\hat{x}^j, \tilde{x}^j) \|\hat{x}^j - \tilde{x}^j\|_X \\ &+ \frac{2\kappa_2}{N^2} \|\hat{x}^j - \tilde{x}^j\|_X \sum_{i=1, i\neq j}^N m_{P_i} \\ &= \frac{\kappa_1 + \kappa_2}{N} L_2(\hat{x}^j, \tilde{x}^j) \|\hat{x}^j - \tilde{x}^j\|_X + \frac{2\kappa_2}{N^2} \|\hat{x}^j - \tilde{x}^j\|_X \sum_{i=1, i\neq j}^N m_{P_i} \\ &\leq \frac{1}{N} \max \left\{ \kappa_1 + \kappa_2, \frac{2\kappa_2}{N} \sum_{i=1, i\neq j}^N m_{P_i} \right\} L_2(\hat{x}^j, \tilde{x}^j) \|\hat{x}^j - \tilde{x}^j\|_X \end{aligned}$$

and so, for any choice of $\hat{P}_j, \tilde{P}_j \in \mathscr{P}_2(X)$,

$$\begin{split} \sup_{\boldsymbol{\psi}\in\Psi_{\kappa_{1},\kappa_{2}}} \left| \int_{X} \int_{X^{N-1}} \boldsymbol{\psi}(\boldsymbol{x}_{-j},x_{j}) \, \boldsymbol{P}_{-j}(\mathrm{d}\boldsymbol{x}_{-j}) \hat{P}_{j}(\mathrm{d}\xi_{j}) \right. \\ \left. - \int_{X} \int_{X^{N-1}} \boldsymbol{\psi}(\boldsymbol{x}_{-j},\xi_{j}) \, \boldsymbol{P}_{-j}(\mathrm{d}\boldsymbol{x}_{-j}) \, \tilde{P}_{j}(\mathrm{d}x_{j}) \right| \\ \left. = \sup_{\boldsymbol{\psi}\in\Psi_{\kappa_{1},\kappa_{2}}} \left| \int_{X} \boldsymbol{\psi}_{\boldsymbol{x}_{-j}}(x_{j}) \, \hat{P}_{j}(\mathrm{d}x_{j}) - \int_{X} \boldsymbol{\psi}_{\boldsymbol{x}_{-j}}(x_{j}) \, \tilde{P}_{j}(\mathrm{d}x_{j}) \right| \\ \left. \leq \frac{1}{N} \max\left\{ \kappa_{1} + \kappa_{2}, \frac{2\kappa_{2}}{N} \sum_{i=1, i\neq j}^{N} m_{P_{i}} \right\} \mathsf{dl}_{2}(\hat{P}_{j}, \tilde{P}_{j}) \end{split}$$

Therefore, we obtain

$$\begin{split} \sup_{\psi \in \Psi_{\kappa_1,\kappa_2}} \left| \int_{X^N} \psi \, \mathrm{d}\mathbb{P}^P - \int_{X^N} \psi \, \mathrm{d}\mathbb{P}^Q \right| &= \sup_{\psi \in \Psi_{\kappa_1,\kappa_2}} \left| \int_{X^N} \psi \, \mathrm{d}P^{\otimes N} - \int_{X^N} \psi \, \mathrm{d}Q^{\otimes N} \right| \\ &\leq \sup_{\psi \in \Psi_{\kappa_1,\kappa_2}} \left| \int_{X^N} \psi \, \mathrm{d}P^{\otimes N} - \int_{X^N} \psi \, \mathrm{d}(P^{\otimes (N-1)} \otimes Q) \right| \\ &+ \sup_{\psi \in \Psi_{\kappa_1,\kappa_2}} \left| \int_{X^N} \psi \, \mathrm{d}(P^{\otimes (N-1)} \otimes Q) - \int_{X^N} \psi \, \mathrm{d}(P^{\otimes (N-2)} \otimes Q^{\otimes 2}) \right| \\ &+ \cdots + \sup_{\psi \in \Psi_{\kappa_1,\kappa_2}} \left| \int_{X^N} \psi \, \mathrm{d}(P \otimes Q^{\otimes (N-1)}) - \int_{X^N} \psi \, \mathrm{d}Q^{\otimes N} \right| \\ &\leq \frac{1}{N} \max \left\{ \kappa_1 + \kappa_2, \frac{2\kappa_2}{N} (N-1)m_P \right\} \mathsf{dl}_2(P,Q) \end{split}$$

$$+ \frac{1}{N} \max\left\{\kappa_1 + \kappa_2, \frac{2\kappa_2}{N} \left((N-2)m_p + m_Q \right) \right\} \operatorname{dl}_2(P,Q) \\ + \dots + \frac{1}{N} \max\left\{\kappa_1 + \kappa_2, \frac{2\kappa_2}{N} (N-1)m_Q \right\} \operatorname{dl}_2(P,Q) \\ \leq \max\left\{\kappa_1 + \kappa_2, 2\kappa_2 m_P, 2\kappa_2 m_Q \right\} \operatorname{dl}_2(P,Q)$$

for any $P, Q \in \mathscr{P}_2(X)$.

We are now ready to prove Theorem 3.1. We have

$$dl_{Y,1}\left(\mathbb{P}^{P}\circ\widehat{T}_{N}^{-1},\mathbb{P}^{Q}\circ\widehat{T}_{N}^{-1}\right) = \sup_{g\in\mathcal{F}_{1}(Y)} \left|\int_{Y}g\,d\mathbb{P}^{P}\circ\widehat{T}_{N}^{-1} - \int_{Y}g\,d\mathbb{P}^{Q}\circ\widehat{T}_{N}^{-1}\right|$$
$$= \sup_{g\in\mathcal{F}_{1}(Y)}\left|\int_{X^{N}}g(\widehat{T}_{N})\,d\mathbb{P}^{P} - \int_{X^{N}}g(\widehat{T}_{N})\,\mathbb{P}^{Q}\right|$$
$$= \sup_{g\in\mathcal{F}_{1}(Y)}\left|\int_{X^{N}}\psi_{g}\,d\mathbb{P}^{P} - \int_{X^{N}}\psi_{g}\,\mathbb{P}^{Q}\right|,$$
(33)

where $\psi_g := g(\widehat{T}_N)$. Moreover, for any $g \in \mathcal{F}_1(Y)$, we have

$$\begin{aligned} \left| \psi_{g}(\hat{\boldsymbol{x}}) - \psi_{g}(\tilde{\boldsymbol{x}}) \right| &= \left| g \big(\widehat{T}_{N}(\hat{\boldsymbol{x}}) \big) - g \big(\widehat{T}_{N}(\tilde{\boldsymbol{x}}) \big) \right| \leq \left\| \widehat{T}_{N}(\hat{\boldsymbol{x}}) - \widehat{T}_{N}(\tilde{\boldsymbol{x}}) \right\|_{Y} \\ &\leq \kappa \Big(\frac{1}{N} \sum_{i=1}^{N} 2L_{2}(\hat{x}^{i}, \tilde{x}^{i}) \| \hat{x}^{i} - \tilde{x}^{i} \| + \frac{1}{N^{2}} \sum_{i=1}^{N} (\| \hat{x}^{i} \| + \| \tilde{x}^{i} \|) \sum_{j=1}^{N} \| \hat{x}^{j} - \tilde{x}^{j} \| \Big) \end{aligned}$$

for all $\hat{\boldsymbol{x}} = (\hat{x}^1, \dots, \hat{x}^N)$, $\tilde{\boldsymbol{x}} = (\tilde{x}^1, \dots, \tilde{x}^N) \in X^N$, which means that $\psi_g := g(\hat{T}_N)$ belongs to the set $\Psi_{\kappa_1,\kappa_2}\kappa$. So (6) follows directly from (33) and Lemma 10.1.

10.3 Proof of Proposition 4.1

Let $\lambda \in \mathbb{R}_{++}$ and $\Sigma \in \mathbb{S}_{+}^{n}$ be arbitrary but fixed. Moreover, let $K > n/\lambda$ be arbitrary but fixed. It is well-known that the mapping $\mathbb{S}_{++}^{n} \to \mathbb{R}$, $S \mapsto \log(\det S)$ is strictly concave. Thus, the strict convexity of $S \mapsto L(\lambda, \Sigma, S)$ is evident given that $S \mapsto \langle \Sigma, S \rangle$ is linear and $S \mapsto \lambda ||S||_1$ is convex. The second statement follows from the fact that a matrix $S \in \mathbb{S}_{++}^{n}$ with $||S||_1 > K$ cannot be the minimizer of the mapping $\mathbb{S}_{++}^{n} \to \mathbb{R}$, $S \mapsto L(\lambda, \Sigma, S)$. Take into account that for any $S \in \mathbb{S}_{++}^{n}$ with $||S||_1 > K$, we have

$$\begin{split} L(\lambda, \Sigma, S) &- L(\lambda, \Sigma, (1 - \varepsilon)S) \\ &= \langle \Sigma, S \rangle - \log(\det S) + \lambda \|S\|_1 \\ &- \left(\langle \Sigma, (1 - \varepsilon)S \rangle - \log(\det((1 - \varepsilon)S)) + \lambda \|(1 - \varepsilon)S\|_1 \right) \\ &= \varepsilon \langle \Sigma, S \rangle + \varepsilon \lambda \|S\|_1 + n\log(1 - \varepsilon) \ge \varepsilon \lambda \|S\|_1 + n\log(1 - \varepsilon) \\ &> \varepsilon \lambda K + n\log(1 - \varepsilon) \ge 0, \end{split}$$

where $\varepsilon \in \mathbb{R}_{++}$ is chosen so small that $(n/\lambda)(-\log(1-\varepsilon))/\varepsilon \leq K$. Note here that $-\log(1-\varepsilon)/\varepsilon \downarrow 1$ as $\varepsilon \downarrow 0$.

10.4 Proof of Proposition 4.2

Fix $\lambda \in \mathbb{R}_{++}$ and $\Sigma \in \mathbb{S}^n_+$. To simplify the notation in this proof, the dependence of the quantities involved on λ and Σ will be omitted. That is, we will write L(S), S^* , α and β instead of $L(\lambda, \Sigma, S)$, $S^*(\lambda, \Sigma)$, $\alpha_{\lambda,\Sigma}$ and $\beta_{\lambda,\Sigma}$, respectively. For any $S \in \mathbb{S}^n_{++}$, using the notation $\Delta_{S,S^*} := (S-S^*)/||S-S^*||$, we obtain by the fundamental theorem of calculus that

$$L(S) - L(S^*) = f_{S,S^*}(||S - S^*||) - f_{S,S^*}(0) = \int_0^{||S - S^*||} f'_{S,S^*}(t) \,\mathrm{d}t, \tag{34}$$

where the function $f_{S,S^*} : \mathbb{R}_+ \to \mathbb{R}$ defined by $f_{S,S^*}(t) := L(S^* + t\Delta_{S,S^*})$ is directionally differentiable and its derivative f'_{S,S^*} satisfies

$$f_{S,S^*}'(t) = \left\langle \Sigma, \Delta_{S,S^*} \right\rangle - \left\langle (S^* + t\Delta_{S,S^*})^{-1}, \Delta_{S,S^*} \right\rangle + \left\langle G(S^* + t\Delta_{S,S^*}), \Delta_{S,S^*} \right\rangle.$$

Here the mapping $G : \mathbb{R}^{n \times n} \to \mathbb{R}^{n \times n}$ is defined by $G(A)_{ij} := \lambda \operatorname{sgn}(A_{ij})$, where sgn denotes the sign function. The mapping $\mathbb{S}_{++}^n \to \mathbb{R}$, $S \mapsto L(S)$ is strictly convex (by Proposition 4.1) and satisfies $L(S^*) < L(S)$ for any $S \in \mathbb{S}_{++}^n$ with $S \neq S^*$. Thus, for any $S \in \mathbb{S}_{++}^n$ with $S \neq S^*$, we have $f'_{S,S^*}(t) > 0$ for all $t \in (0, ||S - S^*||]$ and that $t \mapsto f'_{S,S^*}(t)$ is monotonically increasing on $[0, ||S - S^*||]$. Thus, in view of (34), for (11) it suffices to show $\beta := \inf_{S \in \mathbb{S}_{++}^n : S \neq S^*} \inf_{t \in [1, ||S - S^*||]} f'_{S,S^*}(t) > 0$ (where $\inf_{t \in \emptyset} f'_{S,S^*}(t) := \infty$) and that $\alpha := \inf_{S \in \mathbb{S}_{++}^n : S \neq S^*} \lim \inf_{t \downarrow 0} f'_{S,S^*}(t) / t > 0$.

We first show that $\inf_{S \in \mathbb{S}_{++}^n : S \neq S^*} \inf_{t \in [1, \|S - S^*\|]} f'_{S,S^*}(t) > 0$. Since the derivative f'_{S,S^*} of f_{S,S^*} is monotonically increasing for any $S \in \mathbb{S}_{++}^n$, it suffices to show that $\inf_{S \in \mathbb{S}_{++}^n : \|S - S^*\| \ge 1} f'_{S,S^*}(1) > 0$. By way of contradiction we assume that $\inf_{S \in \mathbb{S}_{++}^n : \|S - S^*\| \ge 1} f'_{S,S^*}(1) = 0$. Then we can find a sequence $(S_n)_{n \in \mathbb{N}}$ in \mathbb{S}_{++}^n such that $\|S_n - S^*\| \ge 1$ and $f'_{S_n,S^*}(1) \le 1/n$ for all $n \in \mathbb{N}$. In particular, we obtain from (34) that $L(S^* + \Delta_{S_n,S^*}) - L(S^*) = \int_0^1 f'_{S_n,S^*}(t) dt \le 1/n$ for all $n \in \mathbb{N}$. But this is a contradiction to the fact that S^* is an isolated minimum of the continuous mapping $S \mapsto L(S)$, since $\|(S^* + \Delta_{S_n,S^*}) - S^*\| = \|\Delta_{S_n,S^*}\| = 1$ for all $n \in \mathbb{N}$.

It remains to show that $\inf_{S \in \mathbb{S}^n_{++}: S \neq S^*} \liminf_{t \downarrow 0} f'_{S,S^*}(t)/t > 0$. For any $S \in \mathbb{S}^n_{++}$, we have

$$f'_{S,S^*}(t) = \langle \Sigma, \Delta_{S,S^*} \rangle - \langle (S^* + t\Delta_{S,S^*})^{-1}, \Delta_{S,S^*} \rangle + \langle G(S^* + t\Delta_{S,S^*}), \Delta_{S,S^*} \rangle$$

$$= \langle \Sigma, \Delta_{S,S^*} \rangle - \langle S^{*-1} - tS^{*-1}\Delta_{S,S^*}S^{*-1} + o(t\Delta_{S,S^*}), \Delta_{S,S^*} \rangle$$

$$+ \langle G(S^* + t\Delta_{S,S^*}), \Delta_{S,S^*} \rangle$$

$$= \langle \Sigma - S^{*-1} + G(S^* + t\Delta_{S,S^*}), \Delta_{S,S^*} \rangle + t \langle S^{*-1}\Delta_{S,S^*}S^{*-1}, \Delta_{S,S^*} \rangle$$

$$+ \langle o(t\Delta_{S,S^*}), \Delta_{S,S^*} \rangle, \qquad (35)$$

where o(B) is a matrix-valued function of B with $\lim_{\|B\|\to 0} \|o(B)\|/\|B\| = 0$. For the second step in the above equation chain we used the following consequence of the convergence of the Neumann series (see [45, Chapter 1, Theorem 4.20]): $(A + B)^{-1} = \sum_k (-1)^k (A^{-1}B)^k A^{-1} = A^{-1} - A^{-1}BA^{-1} + o(B)$ for any A, B satisfying $\lim_{k\to\infty} (A^{-1}B)^k = 0$. In fact, we applied this representation to $A := S^*$ and $B := t\Delta_{S,S^*}$ for small enough t. The first summand on the right-hand side of (35) is constant in a right-sided neighborhood of 0. That is, there exist $r_S \in \mathbb{R}_{++}$ and $\varpi_S \in \mathbb{R}$ such that $\langle \Sigma - S^{*-1} + G(S^* + t\Delta_{S,S^*}), \Delta_{S,S^*} \rangle = \varpi_S$ for all $t \in [0, r_S]$. Thus, since $f'_{S,S^*}(t) > 0$ for all $t \in (0, r_S] > 0$ (choose r_S sufficiently close to 0), we have

$$0 \leq \lim_{t \downarrow 0} f'_{S,S^*}(t) = \lim_{t \downarrow 0} \left\langle \Sigma - S^{*-1} + G(S^* + t\Delta_{S,S^*}), \Delta_{S,S^*} \right\rangle = \lim_{t \downarrow 0} \varpi_S = \varpi_S.$$

Consequently, we have $\lim_{t\downarrow 0} \langle \Sigma - S^{*-1} + G(S^* + t\Delta_{S,S^*}), \Delta_{S,S^*} \rangle / t = 0$ or $= \infty$ depending on whether $\varpi_S = 0$ or $\varpi_S > 0$. Thus, we obtain from (35) that $\lim_{t\downarrow 0} f'_{S,S^*}(t) / t \ge \langle S^{*-1}\Delta_{S,S^*}S^{*-1}, \Delta_{S,S^*} \rangle = \|\Delta_{S,S^*}S^{*-1}\|^2 \ge \lambda_{\max}^{-2}(S^*)\|\Delta_{S,S^*}\|^2 = \lambda_{\max}^{-2}(S^*)$, where $\lambda_{\max}(S^*)$ stands for the largest eigenvalue of S^* .

10.5 Proof of Theorem 4.1

Fix $\lambda \in \mathbb{R}_{++}$ and $\Sigma \in \mathbb{S}_{+}^{n}$, and let C_{λ} and $\alpha_{\lambda,\Sigma}, \beta_{\lambda,\Sigma}$ be as in (8) and Proposition 4.2, respectively. Let S_{λ} be as in (8) and recall that problems (7) and (9) are equivalent. For any $\varepsilon \in \mathbb{R}_{++}$, we set

$$r_{\lambda,\Sigma}(\varepsilon) := \inf_{S \in \mathcal{S}_{\lambda}: \, \|S - S^*(\lambda,\Sigma)\| \ge \varepsilon} L(\lambda,\Sigma,S) - L(\lambda,\Sigma,S^*(\lambda,\Sigma)).$$

Note that $\sup_{S \in S_{\lambda}} |L(\lambda, \Sigma, S) - L(\lambda, \Sigma', S)| = \sup_{S \in S_{\lambda}} |\langle \Sigma - \Sigma', S \rangle| \leq \sup_{S \in S_{\lambda}} ||\Sigma - \Sigma'|| ||S|| \leq ||\Sigma - \Sigma'||C_{\lambda}$ for all $\Sigma' \in \mathbb{S}^{n}_{+}$. Thus, for any $\varepsilon \in \mathbb{R}_{++}$, $S \in S_{\lambda}$ with $||S - S^{*}(\lambda, \Sigma)|| \geq \varepsilon$ and $\Sigma' \in \mathbb{S}^{n}_{+}$, we obtain

$$L(\lambda, \Sigma', S) - L(\lambda, \Sigma', S^*(\lambda, \Sigma')) = L(\lambda, \Sigma, S) - L(\lambda, \Sigma, S^*(\lambda, \Sigma)) + L(\lambda, \Sigma', S^*(\lambda, \Sigma)) - L(\lambda, \Sigma', S^*(\lambda, \Sigma)) - L(\lambda, \Sigma', S^*(\lambda, \Sigma)) + L(\lambda, \Sigma', S) - L(\lambda, \Sigma, S)$$

$$\geq L(\lambda, \Sigma, S) - L(\lambda, \Sigma, S^*(\lambda, \Sigma)) + 0 - 2C_{\lambda} ||\Sigma - \Sigma'||$$

$$\geq r_{\lambda, \Sigma}(\varepsilon) - 2C_{\lambda} ||\Sigma - \Sigma'||.$$
(36)

From Proposition 4.2, we know that $r_{\lambda,\Sigma}(\varepsilon) \geq \min\{\alpha_{\lambda,\Sigma}\varepsilon^2, \beta_{\lambda,\Sigma}\varepsilon\}$ for all $\varepsilon \in \mathbb{R}_{++}$. Thus, for any $\Sigma' \in \mathbb{S}^n_+$ with $\Sigma' \neq \Sigma$ we have $r_{\lambda,\Sigma}(\varepsilon') \geq 3C_{\lambda} \|\Sigma - \Sigma'\|$ for $\varepsilon' := \max\{(3C_{\lambda}\|\Sigma - \Sigma'\|/\alpha_{\lambda,\Sigma})^{1/2}, 3C_{\lambda}\|\Sigma - \Sigma'\|/\beta_{\lambda,\Sigma}\}$ (> 0). In particular, (36) implies $L(\lambda, \Sigma', S) - L(\lambda, \Sigma', S^*(\lambda, \Sigma')) \geq C_{\lambda} \|\Sigma - \Sigma'\| > 0$ for any $\Sigma' \in \mathbb{S}^n_+$ with $\Sigma' \neq \Sigma$ and any $S \in \mathcal{S}_{\lambda}$ with $\|S - S^*(\lambda, \Sigma)\| \geq \varepsilon'$. Thus, $S \neq S^*(\lambda, \Sigma')$ for any $\Sigma' \in \mathbb{S}^n_+$ and $S \in \mathcal{S}_{\lambda}$ with $\Sigma' \neq \Sigma$ and $\|S - S^*(\lambda, \Sigma)\| \geq \varepsilon'$. Since $S^*(\lambda, \Sigma') \in \mathcal{S}_{\lambda}$, we can conclude that $\|S^*(\lambda, \Sigma') - S^*(\lambda, \Sigma)\| < \varepsilon' = \text{right-hand side of (12)}.$

10.6 Proof of Proposition 4.3

The proof is similar to Lemma A1 in [22]. Here we give a sketch to facilitate reading. Let $\varepsilon \in \mathbb{R}_{++}$ be arbitrary but fixed. Set $\delta := R_{\lambda,\Sigma}^{-1}(3n^2\sqrt{\varepsilon})$ and note that $R_{\lambda,\Sigma}(\delta) = 3n^2\sqrt{\varepsilon}$. If we now set $\eta := R_{\lambda,\Sigma}(\delta)/3$, then we get

$$\sup_{S \in \mathcal{S}_{\lambda}} \left| L(\lambda, \Sigma, S) - L_{\varepsilon}(\lambda, \Sigma, S) \right| \leq \sum_{1 \leq i, j \leq n} \sup_{s_{ij} \in \mathbb{R}} \left| |s_{ij}| - \sqrt{s_{ij}^2 + \varepsilon} \right| = n^2 \sqrt{\varepsilon} \leq \eta.$$

Consequently, for any $S \in \mathcal{S}_{\lambda}$ with $||S - S^*(\lambda, \Sigma)|| \ge \delta$ we obtain that

$$L_{\varepsilon}(\lambda, \Sigma, S) - L_{\varepsilon}(\lambda, \Sigma, S^{*}(\lambda, \Sigma)) \geq L(\lambda, \Sigma, S) - L(\lambda, \Sigma, S^{*}(\lambda, \Sigma)) - 2\eta$$

$$\geq R_{\lambda, \Sigma}(\delta)/3 > 0,$$

which means that S is not a minimizer of the mapping $\mathbb{S}^n_+ \to \mathbb{R}$, $S' \mapsto L_{\varepsilon}(\lambda, \Sigma, S')$. This in turn implies that $\|S^*(\lambda, \Sigma) - S^*_{\varepsilon}(\lambda, \Sigma)\| \leq \delta = R_{\lambda, \Sigma}^{-1}(3n^2\sqrt{\varepsilon}).$

10.7 Proof of Theorem 4.2

In Step 1 below we will prove that for any $\Sigma \in \mathbb{S}^n_+$ there exist an $\varepsilon_{\Sigma} \in \mathbb{R}_{++}$ and a neighborhood $\mathcal{N}(\Sigma)$ of Σ in \mathbb{S}^n_+ such that

$$\left\|S_{\varepsilon}^{*}(\lambda, \Sigma_{1}) - S_{\varepsilon}^{*}(\lambda, \Sigma_{2})\right\| \leq \kappa \|\Sigma_{1} - \Sigma_{2}\| \quad \text{for all } \Sigma_{1}, \Sigma_{2} \in \mathcal{N}(\Sigma) \text{ and } \varepsilon \in (0, \varepsilon_{\Sigma}].$$
(37)

Combining (37) with Proposition 4.3 (according to which $S^*_{\varepsilon}(\lambda, \Sigma_i) \to S^*(\lambda, \Sigma_i)$ as $\varepsilon \downarrow 0, i = 1, 2$), we can conclude that for any $\Sigma \in \mathbb{S}^n_+$ there exists a neighborhood $\mathcal{N}(\Sigma)$ of Σ such that

$$\left|S^*(\lambda, \Sigma_1) - S^*(\lambda, \Sigma_2)\right| \le \kappa \|\Sigma_1 - \Sigma_2\| \quad \text{for all } \Sigma_1, \Sigma_2 \in \mathcal{N}(\Sigma).$$
(38)

In Step 2 below we will show that the inequalities (38), $\Sigma \in \mathbb{S}^n_+$, imply

$$\left\|S^*(\lambda, \Sigma_1) - S^*(\lambda, \Sigma_2)\right\| \le \kappa \|\Sigma_1 - \Sigma_2\| \quad \text{for all } \Sigma_1, \Sigma_2 \in \mathbb{S}^n_+, \tag{39}$$

which is (17).

Step 1. Let $\Sigma \in \mathbb{S}^n_+$ be arbitrary but fixed and choose a closed and bounded (i.e. compact) and convex neighborhood $\mathcal{N}_0(\Sigma)$ of Σ . Lemma 10.2 below ensures that we can pick an $\varepsilon_{\Sigma} \in \mathbb{R}_{++}$ such that $S^*_{\varepsilon}(\lambda, \Sigma') \in \operatorname{int} \mathcal{S}_{\lambda}$ for all $\varepsilon \in (0, \varepsilon_{\Sigma}]$ and $\Sigma' \in \mathcal{N}(\Sigma)$, where $\mathcal{S}_{\lambda} = \{S \in \mathbb{S}^n_{++} : \|S\| \leq C_{\lambda}\}$ (recall (8)).

To prove (37), we intend to apply the implicit function theorem (in the form of Theorem 9.1) to the functions $F_{\lambda,\varepsilon} : \mathbb{S}^n_+ \times \mathbb{S}^n_{++} \to \mathbb{R}$, $\varepsilon \in (0, \varepsilon_{\Sigma}]$, defined by $F_{\lambda,\varepsilon}(\Sigma, S) := G_{\varepsilon}(\lambda, \Sigma, S)$ with $G_{\varepsilon}(\lambda, \Sigma, S) := \Sigma - S^{-1} + \lambda \nabla H_{\varepsilon}(S)$ (as in the line after (16)). To this end, we note that the following two assertions hold true:

- (a) For any $\Sigma' \in \mathcal{N}_0(\Sigma)$ and $\varepsilon \in (0, \varepsilon_{\Sigma}]$, the mapping $\mathbb{S}^n_{++} \to \mathbb{R}^{n \times n}$, $S \mapsto F_{\lambda,\varepsilon}(\Sigma', S)$ is ρ -strongly monotone on \mathcal{S}_{λ} , where $\rho := 1/\kappa$ is independent of ε .
- (b) For any $S \in \mathcal{S}_{\lambda}$ and $\varepsilon \in (0, \varepsilon_{\Sigma}]$, the mapping $\mathbb{S}^{n}_{+} \to \mathbb{R}^{n \times n}$, $\Sigma' \mapsto F_{\lambda,\varepsilon}(\Sigma', S)$ is 1-Lipschitz continuous (on all of \mathbb{S}^{n}_{+} and thus) on $\mathcal{N}_{0}(\Sigma)$.

Assertion (a) is true since, for any $\Sigma' \in \mathcal{N}_0(\Sigma)$ and $\varepsilon \in (0, \varepsilon_{\Sigma}]$, ρ_{λ} -strong monotonicity of the mapping $\mathbb{S}_{++}^n \to \mathbb{R}^{n \times n}$, $S \mapsto F_{\lambda,\varepsilon}(\Sigma', S)$ on \mathcal{S}_{λ} follows from $F_{\lambda,\varepsilon}(\Sigma', S) = G_{\varepsilon}(\lambda, \Sigma', S) = \Sigma' - S^{-1} + \lambda \nabla H_{\varepsilon}(S)$, Lemma 10.3 below and the fact that the mapping $\mathcal{S}_{\lambda} \to \mathbb{R}^{n \times n}$, $S \mapsto \lambda \nabla H_{\varepsilon}(S)$ is a monotone operator. Assertion (b) is true since, for any $S \in \mathcal{S}_{\lambda}$ and $\varepsilon \in (0, \varepsilon_{\Sigma}]$, the mapping $\mathbb{S}_{+}^n \to \mathbb{R}^{n \times n}$, $\Sigma' \mapsto F_{\lambda,\varepsilon}(\Sigma', S)$ is clearly 1-Lipschitz continuous; take into account that $F_{\lambda,\varepsilon}(\Sigma', S) = G_{\varepsilon}(\lambda, \Sigma', S) = \Sigma - S^{-1} + \lambda \nabla H_{\varepsilon}(S)$.

In view of the above assertions (a) and (b), Theorem 9.1 implies that there exists a neighborhood $\mathcal{N}(\Sigma) \ (\subseteq \mathcal{N}_0(\Sigma))$ of Σ in \mathbb{S}^n_+ such that, for any $\varepsilon \in (0, \varepsilon_{\Sigma}]$, the mapping $\mathbb{S}^n_+ \to \mathbb{S}^n_{++}, \Sigma' \mapsto S^*_{\varepsilon}(\lambda, \Sigma')$ is $\frac{1}{\rho}$ -Lipschitz continuous (i.e. κ -Lipschitz continuous) on $\mathcal{N}(\Sigma)$, i.e. we have (37). Take into account that $\{S^*_{\varepsilon}(\lambda, \Sigma') : \Sigma' \in \mathcal{N}(\Sigma)\}$ is contained in $\{S \in \mathcal{S}_{\lambda} : F_{\lambda,\varepsilon}(S, \Sigma') = \mathbf{0}\}$ for any $\Sigma' \in \mathcal{N}(\Sigma)$, because $S^*_{\varepsilon}(\lambda, \Sigma')$ satisfies equation (16) for any $\Sigma' \in \mathcal{N}(\Sigma)$.

Step 2. We now show that the inequalities (38), $\Sigma \in \mathbb{S}^n_+$, imply (39). Let $\Sigma_1, \Sigma_2 \in \mathbb{S}^n_+$, be arbitrary but fixed. The line segment $\Sigma(\theta) := \Sigma_1 + \theta(\Sigma_2 - \Sigma_1), \theta \in [0, 1]$, connects Σ_1 and Σ_2 . Let $M \in \mathbb{R}_{++}$ be a constant (to be concretized later) and $\delta \in (0, 1/(2M))$. Consider the line segments

$$I_1 := \left\{ \Sigma(\theta) : \theta \in \left[0, 1/M + \delta \right] \right\},\$$

$$I_i := \left\{ \Sigma(\theta) : \theta \in \left[(i-1)/M - \delta, i/M + \delta \right] \right\}, \quad i = 2, \dots, M - 1,$$

$$I_M := \left\{ \Sigma(\theta) : \theta \in \left[1 - 1/M - \delta, 1 \right] \right\}.$$

Then, $\bigcup_{i=1}^{M} I_i = \{\Sigma(\theta) : \theta \in [0,1]\}$ and $I_i \cap I_{i+1} = \{\Sigma(\theta) : \theta \in [i/M - \delta, i/M + \delta]\}$ for i = 1, ..., M - 1.

From Step 1 we know that for any $\Sigma \in \mathbb{S}^n_+$ there exists a neighborhood $\mathcal{N}(\Sigma)$ of Σ such that the mapping $\mathbb{S}^n_+ \to \mathbb{S}^n_{++}$, $S \mapsto S^*(\lambda, \Sigma)$ is κ -Lipschitz continuous on $\mathcal{N}(\Sigma)$ (see (38)). Therefore, and in view of the finite covering theorem, we can assume that M is chosen so large that, for any $i \in \{1, \ldots, M\}$, we have

$$\left\|S^*(\lambda, \Sigma) - S^*(\lambda, \Sigma')\right\| \le \kappa \|\Sigma - \Sigma'\|$$
 for all $\Sigma, \Sigma' \in I_i$.

For $K \in \mathbb{N}$, let $\Sigma^{(k)} := \Sigma_1 + \frac{k}{2^K} (\Sigma_2 - \Sigma_1)$, $k = 0, \dots, 2^K$, and assume that K is large enough that for any $k = 1, \dots, K$, there exists an $i_k \in \{1, \dots, M\}$ such that both $\Sigma^{(k)}$ and $\Sigma^{(k+1)}$ lie in I_{i_k} (this can be achieved by setting $\frac{1}{2^K} < \delta$). Then we have $\|S^*(\lambda, \Sigma^{(k)}) - S^*(\lambda, \Sigma^{(k+1)})\| \le \kappa \|\Sigma^{(k)} - \Sigma^{(k+1)}\|$ for $k = 0, \dots, 2^K - 1$, and it follows that

$$\begin{split} \left\| S^*(\lambda, \Sigma_1) - S^*(\lambda, \Sigma_2) \right\| &\leq \sum_{k=0}^{2^K - 1} \left\| S^*(\lambda, \Sigma^{(k)}) - S^*(\lambda, \Sigma^{(k+1)}) \right\| \\ &\leq \sum_{k=0}^{2^K - 1} \kappa \left\| \Sigma^{(k)} - \Sigma^{(k+1)} \right) \right\| \\ &= \kappa \sum_{k=0}^{2^K - 1} \frac{1}{2^K} \| \Sigma_1 - \Sigma_2 \| \\ &= \kappa \| \Sigma_1 - \Sigma_2 \|. \end{split}$$

This proves Theorem 4.2. However, we still need to prove the following two lemmas.

Lemma 10.2. For any $\Sigma \in \mathbb{S}^n_+$, any closed and bounded neighborhood $\mathcal{N}(\Sigma)$ of Σ and any any $\delta \in \mathbb{R}_{++}$, there exists an $\varepsilon_{\Sigma,\delta} \in \mathbb{R}_{++}$ such that

$$\left\|S_{\varepsilon}^{*}(\lambda,\Sigma') - S^{*}(\lambda,\Sigma')\right\| \leq \delta \quad \text{for all } \Sigma' \in \mathcal{N}(\Sigma) \text{ and } \varepsilon \in (0,\varepsilon_{\Sigma,\delta}].$$

$$\tag{40}$$

Proof. Pick a $\Sigma \in \mathbb{S}^n_+$, a a closed and bounded neighborhood $\mathcal{N}(\Sigma)$ of Σ and a $\delta \in \mathbb{R}_{++}$. Since $\mathcal{N}(\Sigma)$ is assumed to be bounded, we can find, for any $\nu \in \mathbb{R}_{++}$, a finite number, say $M_{\Sigma,\nu}$, of balls $\mathbb{B}(\Sigma_{\nu}^{(i)},\nu)$, $i = 1, \ldots, M_{\Sigma,\nu}$, with radius ν and center points $\Sigma_{\nu}^{(i)} \in \mathcal{N}(\Sigma)$, $i = 1, \ldots, M_{\Sigma,\nu}$, such that $\mathcal{N}(\Sigma) \subseteq \bigcup_{i=1}^{M_{\Sigma,\nu}} \mathbb{B}(\Sigma_{\nu}^{(i)},\nu)$. In particular, for any $\Sigma' \in \mathcal{N}(\Sigma)$ and $\nu \in \mathbb{R}_{++}$, we can find an $i_{\nu,\Sigma'} \in \{1,\ldots,M_{\Sigma,\nu}\}$ such that $\|\Sigma' - \Sigma_{\nu}^{(i_{\nu,\Sigma'})}\| \leq \nu$. To prove (40), note that

$$\begin{split} \left\| S_{\varepsilon}^{*}(\lambda, \Sigma') - S^{*}(\lambda, \Sigma') \right\| &\leq \left\| S_{\varepsilon}^{*}(\lambda, \Sigma') - S_{\varepsilon}^{*}(\lambda, \Sigma_{\nu}^{(i_{\nu}, \Sigma')}) \right\| \\ &+ \left\| S_{\varepsilon}^{*}(\lambda, \Sigma_{\nu}^{(i_{\nu}, \Sigma')}) - S^{*}(\lambda, \Sigma_{\nu}^{(i_{\nu}, \Sigma')}) \right\| \\ &+ \left\| S^{*}(\lambda, \Sigma_{\nu}^{(i_{\nu}, \Sigma')}) - S^{*}(\lambda, \Sigma') \right\| \\ &=: T_{1}(\varepsilon, \nu, \Sigma') + T_{2}(\varepsilon, \nu, \Sigma') + T_{3}(\nu, \Sigma') \end{split}$$

for all $\varepsilon \in \mathbb{R}_{++}$, $\nu \in \mathbb{R}_{++}$ and $\Sigma' \in \mathcal{N}(\Sigma)$. In Step 1 below, we will show that we can choose a $\overline{\nu}_{\delta} \in \mathbb{R}_{++}$ such that $T_3(\nu, \Sigma') \leq \delta/3$ for all $\nu \in (0, \overline{\nu}_{\delta}]$ and $\Sigma' \in \mathcal{N}(\Sigma)$. In Step 2 below, we will show that we can find an $\overline{\varepsilon}_{\Sigma,\delta} \in \mathbb{R}_{++}$ and a $\nu_{\delta} \in (0, \overline{\nu}_{\delta}]$ such that $T_1(\varepsilon, \nu_{\delta}, \Sigma') \leq \delta/3$ for all $\varepsilon \in (0, \overline{\varepsilon}_{\Sigma,\delta}]$ and $\Sigma' \in \mathcal{N}(\Sigma)$. In Step 3 below, we will show that we can find an $\varepsilon_{\Sigma,\delta} \in (0, \overline{\varepsilon}_{\Sigma,\delta}]$ such that $T_2(\varepsilon, \nu_{\delta}, \Sigma') \leq \delta/3$ for all $\varepsilon \in (0, \varepsilon_{\Sigma, \delta}]$ and $\Sigma' \in \mathcal{N}(\Sigma)$. So we obtain $||S_{\varepsilon}^*(\lambda, \Sigma') - S^*(\lambda, \Sigma')|| \leq \delta$ for all $\varepsilon \in (0, \varepsilon_{\Sigma, \delta}]$ and $\Sigma \in \mathcal{N}(\Sigma)$, which proves (40).

Step 1. By the (pointwise) continuity of the mapping $\mathbb{S}^n_+ \to \mathbb{S}^n_+$, $\Sigma' \mapsto S^*(\lambda, \Sigma')$ (see Theorem 4.1), we can indeed choose $\overline{\nu}_{\delta} \in \mathbb{R}_{++}$ sufficiently small such that $T_3(\nu, \Sigma')$ (= $\|S^*(\lambda, \Sigma^{(i_{\nu, \Sigma'})}_{\nu}) - S^*(\lambda, \Sigma')\|$) $\leq \delta/3$ for all $\nu \in (0, \overline{\nu}_{\delta}]$ and $\Sigma' \in \mathcal{N}(\Sigma)$.

Step 2. Observe that

$$\sup_{S \in \mathcal{S}_{\lambda}} \left| L_{\varepsilon}(\lambda, \Sigma', S) - L_{\varepsilon}(\lambda, \Sigma_{\nu}^{(i_{\nu, \Sigma'})}, S) \right| \le \left\| \Sigma' - \Sigma^{(i_{\nu, \Sigma'})} \right\| \sup_{S \in \mathcal{S}_{\lambda}} \left\| S \right\| \le \nu C_{\lambda}$$

for all $\nu \in \mathbb{R}_{++}$ and $\Sigma' \in \mathcal{N}(\Sigma)$, where C_{λ} is as in Proposition 4.1 (note that the upper bound νC_{λ} is independent of ε). Now let $\nu_{\delta} \in (0, \overline{\nu}_{\delta}]$ (to be concretized later). By Proposition 4.3, we can choose an $\overline{\varepsilon}_{\Sigma,\delta} = \overline{\varepsilon}_{\Sigma}(\nu_{\delta}) \in \mathbb{R}_{++}$ sufficiently small such that $S_{\varepsilon}^*(\lambda, \Sigma_{\nu_{\delta}}^{(i)}) \in S_{\lambda}$ for all $i = 1, \ldots, M_{\Sigma,\nu_{\delta}}$ and $\varepsilon \in (0, \overline{\varepsilon}_{\Sigma,\delta}]$. Then we have

$$\begin{split} L_{\varepsilon}(\lambda,\Sigma',S) &- L_{\varepsilon}\left(\lambda,\Sigma',S_{\varepsilon}^{*}(\lambda,\Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta}},\Sigma')})\right) \\ &= L_{\varepsilon}(\lambda,\Sigma',S) - L_{\varepsilon}\left(\lambda,\Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta}},\Sigma')},S\right) \\ &+ L_{\varepsilon}\left(\lambda,\Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta}},\Sigma')},S\right) - L_{\varepsilon}\left(\lambda,\Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta}},\Sigma')},S_{\varepsilon}^{*}(\lambda,\Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta}},\Sigma')})\right) \\ &+ L_{\varepsilon}\left(\lambda,\Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta}},\Sigma')},S_{\varepsilon}^{*}(\lambda,\Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta}},\Sigma')})\right) - L_{\varepsilon}\left(\lambda,\Sigma',S_{\varepsilon}^{*}(\lambda,\Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta}},\Sigma')})\right) \\ &\geq L_{\varepsilon}\left(\lambda,\Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta}},\Sigma')},S\right) - L_{\varepsilon}\left(\lambda,\Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta}},\Sigma')},S_{\varepsilon}^{*}(\lambda,\Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta}},\Sigma')})\right) - 2(\nu_{\delta}C_{\lambda}) \\ &\geq R_{\varepsilon}\left(\lambda,\Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta}},\Sigma')},\delta/3\right) - 2\nu_{\delta}C_{\lambda} \end{split}$$

for all $\varepsilon \in (0, \overline{\varepsilon}_{\Sigma, \delta}], \Sigma' \in \mathcal{N}(\Sigma)$ and $S \in \mathcal{S}_{\lambda}$ with $||S - S_{\varepsilon}^*(\lambda, \Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta}, \Sigma'})})|| \ge \delta/3$, where

$$R_{\varepsilon}(\lambda, \Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta}}, \Sigma')}, \delta/3)$$

:=
$$\inf_{S \in \mathcal{S}_{\lambda}: \, \|S - S_{\varepsilon}^{*}(\lambda, \Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta}}, \Sigma')})\| \ge \delta/3} L_{\varepsilon}(\Sigma', S) - L_{\varepsilon}(\Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta}}, \Sigma')}, S_{\varepsilon}^{*}(\lambda, \Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta}}, \Sigma')})).$$

We can assume without loss of generality that ν_{δ} is chosen so small that $\nu_{\delta} \in (0, \overline{\nu}_{\delta}]$ and $\nu_{\delta}C_{\lambda} \leq \min_{i=1,\dots,M_{\Sigma,\nu_{\delta}}} R_{\varepsilon}(\lambda, \Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta},\Sigma'})}, \delta/3)/3$. Then

$$L_{\varepsilon}(\lambda, \Sigma', S) - L_{\varepsilon}(\lambda, \Sigma', S_{\varepsilon}^{*}(\lambda, \Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta}, \Sigma'})})) \geq R_{\varepsilon}(\lambda, \Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta}, \Sigma'})}, \delta/3)/3 > 0$$

for all $\varepsilon \in (0, \overline{\varepsilon}_{\Sigma, \delta}]$, $\Sigma' \in \mathcal{N}(\Sigma)$ and $S \in \mathcal{S}_{\lambda}$ with $||S - S_{\varepsilon}^{*}(\lambda, \Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta}, \Sigma'})})|| \geq \delta/3$. This implies that, for any $\varepsilon \in (0, \overline{\varepsilon}_{\Sigma, \delta}]$ and $\Sigma' \in \mathcal{N}(\Sigma)$, a matrix $S \in \mathcal{S}_{\lambda}$ with $||S - S_{\varepsilon}^{*}(\lambda, \Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta}, \Sigma'})})|| \geq \delta/3$ cannot be a minimizer of the mapping $\mathcal{S}_{\lambda} \to \mathbb{R}$, $S \mapsto L_{\varepsilon}(\lambda, \Sigma', S)$. But this shows that $T_{1}(\varepsilon, \nu_{\delta}, \Sigma')$ (= $||S_{\varepsilon}^{*}(\lambda, \Sigma') - S_{\varepsilon}^{*}(\lambda, \Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta}, \Sigma'})})||) \leq \delta/3$ for all $\varepsilon \in (0, \overline{\varepsilon}_{\Sigma, \delta}]$ and $\Sigma' \in \mathcal{N}(\Sigma)$.

Step 3. By Proposition 4.3, we can choose $\varepsilon_{\Sigma,\delta} \in (0, \overline{\varepsilon}_{\Sigma,\delta}]$ sufficiently small such that

$$\max_{i=1,\dots,M_{\Sigma,\nu_{\delta}}} \|S_{\varepsilon}^{*}(\lambda,\Sigma_{\nu_{\delta}}^{(i)}) - S^{*}(\lambda,\Sigma_{\nu_{\delta}}^{(i)})\| \le \delta/3$$

for all $\varepsilon \in (0, \varepsilon_{\Sigma, \delta}]$. This implies that $T_2(\varepsilon, \nu_{\delta}, \Sigma') (= \|S_{\varepsilon}^*(\lambda, \Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta}, \Sigma'})}) - S^*(\lambda, \Sigma_{\nu_{\delta}}^{(i_{\nu_{\delta}, \Sigma'})})\|) \le \delta/3$ for all $\varepsilon \in (0, \varepsilon_{\Sigma, \delta}]$ and $\Sigma' \in \mathcal{N}(\Sigma)$.

Lemma 10.3. Let $\rho := 1/\kappa$, where κ is defined as in Theorem 4.2. Then

$$-\left\langle S_{1}^{-1} - S_{2}^{-1}, S_{1} - S_{2} \right\rangle \ge \rho \|S_{1} - S_{2}\|^{2} \quad \text{for all } S_{1}, S_{2} \in \mathcal{S}_{\lambda}.$$

Proof. For any $S_1, S_2 \in \mathcal{S}_{\lambda}$, we have

$$-\langle S_1^{-1} - S_2^{-1}, S_1 - S_2 \rangle = -\langle S_1^{-1} (S_2 - S_1) S_2^{-1}, S_1 - S_2 \rangle = \langle S_1^{-1} \Delta S S_2^{-1}, \Delta S \rangle,$$
(41)

where $\Delta S := S_1 - S_2$. By the spectral decomposition, we may write S_1 and S_2 respectively as $S_1 = Q_1 \Lambda_1 Q_1^{\mathsf{T}}$ and $S_2 = Q_2 \Lambda_2 Q_2^{\mathsf{T}}$ for diagonal matrices $\Lambda_1 = \Lambda_1(S_1)$, $\Lambda_2 := \Lambda_2(S_2)$ and orthogonal matrices Q_1, Q_2 . Then

$$\langle S_1^{-1}\Delta S S_2^{-1}, \Delta S \rangle = \operatorname{tr}(Q_2 \Lambda_2^{-1} Q_2^{\mathsf{T}} \Delta S Q_1 \Lambda_1^{-1} Q_1^{\mathsf{T}} \Delta S)$$

$$= \operatorname{tr}(\Lambda_2^{-1} Q_2^{\mathsf{T}} \Delta S Q_1 \Lambda_1^{-1} Q_1^{\mathsf{T}} \Delta S Q_2) \geq \lambda_{\max}^{-1}(\Lambda_2) \operatorname{tr}(Q_2^{\mathsf{T}} \Delta S Q_1 \Lambda_1^{-1} Q_1^{\mathsf{T}} \Delta S Q_2)$$

$$= \lambda_{\max}^{-1}(\Lambda_2) \operatorname{tr}(\Delta S Q_1 \Lambda_1^{-1} Q_1^{\mathsf{T}} \Delta S) = \lambda_{\max}^{-1}(\Lambda_2) \operatorname{tr}(\Lambda_1^{-1} Q_1^{\mathsf{T}} \Delta S \Delta S Q_1)$$

$$\geq \lambda_{\max}^{-1}(\Lambda_2) \lambda_{\max}^{-1}(\Lambda_1) \operatorname{tr}(Q_1^{\mathsf{T}} \Delta S \Delta S Q_1) = \lambda_{\max}^{-1}(\Lambda_2) \lambda_{\max}^{-1}(\Lambda_1) \operatorname{tr}(\Delta S \Delta S)$$

$$= \lambda_{\max}^{-1}(\Lambda_2) \lambda_{\max}^{-1}(\Lambda_1) \|\Delta S\|^2.$$

$$(42)$$

In the proof of Proposition 4.1 (see Section 10.3) we have seen that $||S||_1 \leq K := \sqrt{\kappa}$ for all $S \in S_{\lambda}$. Moreover, we have $\lambda_{\max}(\Lambda(S)) = \lambda_{\max}(S) \leq ||S|| \leq ||S||_1$ (where $S = Q(S)^{\mathsf{T}} \Lambda(S)Q(S)$ is the spectral decomposition of S) for any $S \in S_{\lambda}$. Thus, in view of (41) and (42), we obtain that $-\langle S_1^{-1} - S_2^{-1}, S_1 - S_2 \rangle = \langle S_1^{-1} \Delta S S_2^{-1}, \Delta S \rangle \geq \lambda_{\max}^{-1}(\Lambda_2) \lambda_{\max}^{-1}(\Lambda_1) ||\Delta S||^2 \geq (\sqrt{\kappa})^{-2} ||\Delta S||^2 = \rho ||S_1 - S_2||^2$.

10.8 Proof of Theorem 5.1

Inequality (19c) holds true since

$$dl_{K}(\mathbb{P}^{P} \circ \widehat{\Sigma}_{N}^{-1}, \delta_{\Sigma_{P}}) = \sup_{f \in \mathcal{F}_{1}(\mathbb{S}_{+}^{n})} \left| \int f \, d\mathbb{P}^{P} \circ \widehat{\Sigma}_{N}^{-1} - \int f \, d\delta_{\Sigma_{P}} \right|$$
$$= \sup_{f \in \mathcal{F}_{1}(\mathbb{S}_{+}^{n})} \left| \int \left(f(\widehat{\Sigma}_{N}) - f(\Sigma_{P}) \right) d\mathbb{P}^{P} \right| \leq \int \left\| \widehat{\Sigma}_{N} - \Sigma_{P} \right\| d\mathbb{P}^{P} = \mathbb{E}^{P} \left[\left\| \widehat{\Sigma}_{N} - \Sigma_{P} \right\| \right]$$

and Inequality in (19d) can be obtained analogously. The proof of Inequalities (19a) and (19b) relies on the following lemma, where as before $L_2(\hat{x}, \tilde{x}) = \max\{1, ||\hat{x}||, ||\tilde{x}||\}$.

Lemma 10.4. For any $\hat{\boldsymbol{x}} = (\hat{x}^1, \dots, \hat{x}^N), \ \tilde{\boldsymbol{x}} = (\tilde{x}^1, \dots, \tilde{x}^N) \in (\mathbb{R}^n)^N$, we have

$$\|\widehat{\Sigma}_{N}(\hat{\boldsymbol{x}}) - \widehat{\Sigma}_{N}(\tilde{\boldsymbol{x}})\|$$

$$\leq \frac{2}{N} \sum_{i=1}^{N} L_{2}(\hat{x}^{i}, \tilde{x}^{i}) \|\hat{x}^{i} - \tilde{x}^{i}\| + \frac{1}{N^{2}} \sum_{i=1}^{N} \left(\|\hat{x}^{i}\| + \|\tilde{x}^{i}\| \right) \sum_{j=1}^{N} \|\hat{x}^{j} - \tilde{x}^{j}\|,$$

$$\|\widetilde{\Sigma}_{N}(\boldsymbol{x}) - \widetilde{\Sigma}_{N}(\boldsymbol{y})\| \leq \frac{2}{N} \sum_{i=1}^{N} L_{2}(\hat{x}^{i}, \tilde{x}^{i}) \|\hat{x}^{i} - \tilde{x}^{i}\|.$$
(43)
(43)
(43)

Proof. For any $\hat{\boldsymbol{x}} = (\hat{x}^1, \dots, \hat{x}^N), \, \tilde{\boldsymbol{x}} = (\tilde{x}^1, \dots, \tilde{x}^N) \in (\mathbb{R}^n)^N$, we obtain

$$\begin{split} \|\widehat{\Sigma}_{N}(\widehat{x}) - \widehat{\Sigma}_{N}(\widetilde{x})\| \\ &\leq \left\|\frac{1}{N}\sum_{i=1}^{N}\widehat{x}^{i}(\widehat{x}^{i})^{\mathsf{T}} - \left(\frac{1}{N}\sum_{i=1}^{N}\widehat{x}^{i}\right)\left(\frac{1}{N}\sum_{i=1}^{N}x^{i}\right)^{\mathsf{T}} \\ &\quad -\frac{1}{N}\sum_{i=1}^{N}\widehat{x}^{i}(\widehat{x}^{i})^{\mathsf{T}} + \left(\frac{1}{N}\sum_{i=1}^{N}\widetilde{x}^{i}\right)\left(\frac{1}{N}\sum_{i=1}^{N}\widetilde{x}^{i}\right)^{\mathsf{T}}\right\| \\ &\leq \left\|\frac{1}{N}\sum_{i=1}^{N}\widehat{x}^{i}(\widehat{x}^{i})^{\mathsf{T}} - \frac{1}{N}\sum_{i=1}^{N}\widehat{x}^{i}(\widehat{x}^{i})^{\mathsf{T}}\right\| \\ &\quad + \left\|\frac{1}{N^{2}}\left(\sum_{i=1}^{n}\widehat{x}^{i}\right)\left(\sum_{i=1}^{n}\widehat{x}^{i}\right)^{\mathsf{T}} - \frac{1}{N^{2}}\left(\sum_{i=1}^{n}\widetilde{x}^{i}\right)\left(\sum_{i=1}^{n}\widetilde{x}^{i}\right)^{\mathsf{T}}\right\| \\ &\leq \frac{1}{N}\sum_{i=1}^{N}\left\|x^{i}(\widehat{x}^{i})^{\mathsf{T}} - \widetilde{x}^{i}(\widetilde{x}^{i})^{\mathsf{T}}\right\| + \frac{1}{N^{2}}\left\|\left(\sum_{i=1}^{n}\widehat{x}^{i}\right)\left(\sum_{i=1}^{n}\widehat{x}^{i}\right)^{\mathsf{T}} - \left(\sum_{i=1}^{n}\widetilde{x}^{i}\right)\left(\sum_{i=1}^{n}\widetilde{x}^{i}\right)^{\mathsf{T}}\right\| \\ &\leq \frac{1}{N}\sum_{i=1}^{N}2L_{2}(\widehat{x}^{i},\widetilde{x}^{i})\|\widehat{x}^{i} - \widetilde{x}^{i}\| + \frac{1}{N^{2}}\sum_{i=1}^{N}\left(\|\widehat{x}_{i}\| + \|\widetilde{x}_{i}\|\right)\sum_{i=1}^{N}\|\widehat{x}_{i} - \widetilde{x}_{i}\|. \end{split}$$

This proves Inequality (43). Inequality (44) can be shown analogously.

We are now ready to prove (19a) and (19b). By Inequality (43), the estimator $\widehat{\Sigma}_N$ satisfies condition (5) with $\kappa_1 = 2$ and $\kappa_2 = 1$, which leads to (19a) immediately by Theorem 3.1. By Inequality (44), the estimator $\widetilde{\Sigma}_N$ satisfies condition (5) with $\kappa_1 = 2$ and $\kappa_2 = 0$, which leads to (19b) immediately by Theorem 3.1.

10.9 Proof of Theorem 5.2

The proof of Theorem 5.2 can be carried out in the same way as the proof of ((19a) and (19c) in)Theorem 5.1 (see Section 10.8). For this reason, we will not give the details here.

10.10 Proof of Theorem 5.3

Inequality (21a) is an immediate consequence of Theorem 4.2, Inequality (19a) in Theorem 5.1 and Proposition 3.1. Analogously, Inequality (21b) follows from Theorem 4.2, Inequality (19b) in Theorem 5.1 and Proposition 3.1. Inequality (22a) holds true since

$$dl_{K} \left(\mathbb{P}^{P} \circ S^{*}(\lambda, \widehat{\Sigma}_{N})^{-1}, \delta_{S^{*}(\lambda, \Sigma_{P})} \right)$$

=
$$\sup_{f \in \mathcal{F}_{1}(\mathbb{S}^{n}_{+})} \left| \int f \, d\mathbb{P}^{P} \circ S^{*}(\lambda, \widehat{\Sigma}_{N})^{-1} - \int f \, d\delta_{S^{*}(\lambda, \Sigma_{P})} \right|$$

=
$$\sup_{f \in \mathcal{F}_{1}(\mathbb{S}^{n}_{+})} \left| \int \left(f(S^{*}(\lambda, \widehat{\Sigma}_{N})) - f(S^{*}(\lambda, \Sigma_{P})) \right) d\mathbb{P}^{P} \right|$$

$$\leq \int \left\| S^{*}(\lambda, \widehat{\Sigma}_{N}) - S^{*}(\lambda, \Sigma_{P}) \right\| d\mathbb{P}^{P} = \mathbb{E}^{P} \left[\left\| S^{*}(\lambda, \widehat{\Sigma}_{N}) - S^{*}(\lambda, \Sigma_{P}) \right\| \right]$$

$$\leq \frac{1}{\rho_{\lambda}} \mathbb{E}^{P} \left[\| \widehat{\Sigma}_{N} - \Sigma_{P} \| \right],$$

where the last step is ensured by Theorem 4.2, and Inequality in (22b) can be obtained analogously.

10.11 Proof of Proposition 5.1

Let $r \in [1, 2)$. We will only show the convergence in (23b). The convergences in (23a) and (23c) can be shown analogously. For any $P \in \mathscr{P}_{2r}(\mathbb{R}^N)$, we have

$$\begin{split} &\mathbb{E}^{P} \left[\left\| \Sigma_{N} - \Sigma_{P} \right\|_{1} \right] \\ &= \mathbb{E}^{P} \left[\sum_{j=1}^{n} \sum_{k=1}^{n} \left| \frac{1}{N} \sum_{i=1}^{N} \xi_{j}^{i} \xi_{k}^{i} - \left(\frac{1}{N} \sum_{i=1}^{N} \xi_{j}^{i} \right) \left(\frac{1}{N} \sum_{i=1}^{N} \xi_{k}^{i} \right) - \Sigma_{P}(j,k) \right| \right] \\ &\leq \sum_{j=1}^{n} \sum_{k=1}^{n} \mathbb{E}^{P} \left[\left| \frac{1}{N} \sum_{i=1}^{N} \xi_{j}^{i} \xi_{k}^{i} - \mathbb{E}^{P} \left[\xi_{j}^{1} \xi_{k}^{1} \right] \right| \right] \\ &+ \sum_{j=1}^{n} \sum_{k=1}^{n} \mathbb{E}^{P} \left[\left| \left(\frac{1}{N} \sum_{i=1}^{N} \xi_{j}^{i} \right) \left(\frac{1}{N} \sum_{i=1}^{N} \xi_{k}^{i} \right) - \mathbb{E}^{P} \left[\xi_{j}^{1} \right] \mathbb{E}^{P} \left[\xi_{k}^{1} \right] \right| \right] \\ &\leq \sum_{j=1}^{n} \sum_{k=1}^{n} \mathbb{E}^{P} \left[\left| \overline{X}_{j,k;N} - \mathbb{E}^{P} \left[X_{j,k} \right] \right| \right] + \sum_{j=1}^{n} \sum_{k=1}^{n} \mathbb{E}^{P} \left[\left| \overline{\xi}_{j;N} - \mathbb{E}^{P} \left[\xi_{j}^{1} \right] \right| \left| \overline{\xi}_{k;N} \right| \right] \\ &+ \sum_{j=1}^{n} \sum_{k=1}^{n} \mathbb{E}^{P} \left[\left| \xi_{j}^{1} \right| \right] \mathbb{E}^{P} \left[\left| \overline{\xi}_{k;N} - \mathbb{E}^{P} \left[\xi_{k}^{1} \right] \right| \right] \\ &=: \sum_{j=1}^{n} \sum_{k=1}^{n} T_{1,N}(j,k) + \sum_{j=1}^{n} \sum_{k=1}^{n} T_{2,N}(j,k) + \sum_{j=1}^{n} \sum_{k=1}^{n} T_{3,N}(j,k), \end{split}$$

where $\overline{X}_{j,k,N} := \frac{1}{N} \sum_{i=1}^{N} \xi_j^i \xi_k^i$, $X_{j,k} := \xi_j^1 \xi_k^1$ and $\overline{\xi}_{\ell;N} := \frac{1}{N} \sum_{i=1}^{N} \xi_\ell^i$ for j,k = 1...,n. Now, let $(j,k) \in \{1,\ldots,n\}^2$ be arbitrary but fixed.

Since $X_{j,k}$ is r-fold \mathbb{P}^{P} -integrable (take into account that the random variables ξ_{j}^{1} and ξ_{k}^{1} are both 2r-fold \mathbb{P}^{P} -integrable since $P \in \mathscr{P}_{2r}(\mathbb{R}^{n})$ and $r \geq 1$), we have by Theorem 3.1 in [24] that $\lim_{N\to\infty} N^{(r-1)/r} \mathbb{E}^{P}[|\overline{X}_{j,k;N} - \mathbb{E}^{P}[X_{j,k}]|^{r}]^{1/r} = 0$. Thus,

$$\lim_{N \to \infty} N^{(r-1)/r} T_{1,N}(j,k) = \lim_{N \to \infty} N^{(r-1)/r} \mathbb{E}^{P}[|\overline{X}_{j,k;N} - \mathbb{E}^{P}[X_{j,k}]|] = 0$$

since $r \geq 1$. By Theorem 3.1 in [24], we also have $\lim_{N\to\infty} N^{(r-1)/r} \mathbb{E}^P[|\overline{\xi}_{\ell,N} - \mathbb{E}^P[\xi_k^1]|^r]^{1/r} = 0$. Therefore, since $r \geq 1$, we get $\lim_{N\to\infty} N^{(r-1)/r} T_{3,N}(j,k) = \mathbb{E}^P[|\xi_j^1|] \limsup_{N\to\infty} N^{(r-1)/r} \mathbb{E}^P[|\overline{\xi}_{k;N} - \mathbb{E}^P[\xi_k^1]|] = 0$. By Hölder's inequality, we have

$$\mathbb{E}^{P}[|\overline{\xi}_{j;N} - \mathbb{E}^{P}[\xi_{j}^{1}]||\overline{\xi}_{k;N}|] \leq \mathbb{E}^{P}[|\overline{\xi}_{j;N} - \mathbb{E}^{P}[\xi_{j}^{1}]|^{2}]^{1/2} \mathbb{E}^{P}[|\overline{\xi}_{k;N}|^{2}]^{1/2}.$$

The second factor in the latter upper bound is bounded above uniformly in $N \in \mathbb{N}$ since $\mathbb{E}^{P}[|\overline{\xi}_{k;N}|^{2}]^{1/2} \leq \frac{1}{N} \sum_{i=1}^{N} \mathbb{E}^{P}[|\xi_{k}^{i}|^{2}]^{1/2} = \mathbb{E}^{P}[|\xi_{k}^{1}|^{2}]^{1/2}$ for all $N \in \mathbb{N}$ by Minkowski's inequality. For the first factor we have $N^{1/2} \mathbb{E}^{P}[|\overline{\xi}_{j;N} - \mathbb{E}^{P}[\xi_{j}^{1}]|^{2}]^{1/2} = \mathcal{O}_{N}(1)$ by (2.11) in [24]. Thus, we can conclude that $\lim_{N\to\infty} N^{(r-1)/r} \mathbb{E}^{P}[|\overline{\xi}_{j;N} - \mathbb{E}^{P}[\xi_{j}^{1}]|^{2}]^{1/2} = 0$ since (r-1)/r < 1/2 for our choice of r (recall that r < 2). Consequently, we obtain $\lim_{N\to\infty} N^{(r-1)/r} T_{2,N}(j,k) = 0$.

All in all, we have shown that (23b) does indeed hold.

10.12 Proof of Theorem 5.4

Inequality (24a) is an immediate consequence of (31), Inequality (19a) in Theorem 5.1 and Proposition 3.1. Analogously, Inequality (24b) follows from (31), Inequality (19b) in Theorem 5.1 and Proposition 3.1. Inequality (24c) holds true since

$$\begin{aligned} \mathsf{dI}_{\mathrm{K}} \left(\mathbb{P}^{P} \circ \widehat{\lambda}_{i,N}^{-1}, \delta_{\lambda_{i}^{P}} \right) \\ &= \sup_{f \in \mathcal{F}_{1}(\mathbb{R})} \left| \int f \, \mathrm{d}\mathbb{P}^{P} \circ \widehat{\lambda}_{i,N}^{-1} - \int f \, \mathrm{d}\delta_{\lambda_{i}^{P}} \right| = \sup_{f \in \mathcal{F}_{1}(\mathbb{S}_{+}^{n})} \left| \int \left(f(\widehat{\lambda}_{i,N}) - f(\lambda_{i}^{P}) \right) \mathrm{d}\mathbb{P}^{P} \right| \\ &\leq \int \left\| \widehat{\lambda}_{i,N} - \lambda_{i}^{P} \right\| \mathrm{d}\mathbb{P}^{P} = \mathbb{E}^{P} \left[\left\| \widehat{\lambda}_{i,N} - \lambda_{i}^{P} \right\| \right] \leq \mathbb{E}^{P} \left[\left\| \widehat{\Sigma}_{N} - \Sigma_{P} \right\| \right], \end{aligned}$$

where the last step is ensured by (31), and Inequality in (24d) can be obtained analogously.

10.13 Proof of Proposition 6.1

The assertion of Proposition 6.1 follows from Steps 1 and 2 below. Throughout this section, $z \in \mathbb{R}_+$ is a given constant. For any $(\mu, \Sigma) \in \mathbb{R}^n \times \mathbb{S}^n_+$, we use $\mathcal{W}(\mu, \Sigma)$ to denote the set of optimal solutions of problem (26).

Step 1. We first show that, for any $\mu \in \mathbb{R}^n$, the mapping $\mathbb{S}^n_+ \to \mathbb{R}$, $\Sigma' \mapsto v(\mu, \Sigma')$ is (globally) Lipschitz continuous with Lipschitz constant $L_1 := 1/2$. Indeed: For any $\mu \in \mathbb{R}^n$, we have by the well-known Danskin's theorem that the mapping $\Sigma' : \mathbb{S}^n_+ \to \mathbb{R}$, $\Sigma' \mapsto v(\mu, \Sigma')$ is convex and its subdifferential at $\Sigma \in \mathbb{S}^n_+$ can be represented as $\partial_{\Sigma} v(\mu, \Sigma) = \operatorname{conv}\{\frac{1}{2}ww^{\mathsf{T}} : w \in \mathcal{W}(\mu, \Sigma)\}$. Since each w from the feasible set of (26) satisfies $w^{\mathsf{T}}\mathbf{1} = 1$ and $w \ge 0$, we have $\|\Sigma'\| \le 1/2$ for all $\Sigma' \in \partial_{\Sigma} v(\mu, \Sigma)$. This means that, for any $\mu \in \mathbb{R}^n$, the mapping $\mathbb{S}^n_+ \to \mathbb{R}$, $\Sigma' \mapsto v(\mu, \Sigma')$ is indeed Lipschitz continuous with Lipschitz constant $L_1 := 1/2$.

Step 2. We next show that, for any $C_1, C_2 \in \mathbb{R}_{++}$ with $C_1 < C_2$ and any $\Sigma \in \mathbb{S}_+^n$ with $\|\Sigma\| \leq 2C_2^2$, the mapping $\mathbb{R}_{C_1,C_2}^n \to \mathbb{R}$, $\mu \mapsto v(\mu, \Sigma)$ is Lipschitz continuous with Lipschitz constant $L_{2,C_1,C_2} := L_{C_1,C_2} - \frac{1}{2}$, where $\mathbb{R}_{C_1,C_2}^n := \{m \in \mathbb{R}^n : (C_1^2 + \langle m, \mathbf{1} \rangle^2 / n)^{1/2} \leq \|m\| \leq C_2\}$ and the constant L_{C_1,C_2} is defined as in Proposition 6.1. Without loss of generality we may and do assume $\Sigma \neq \mathbf{0}$ (since for $\Sigma = \mathbf{0}$ we have $v(\mu, \Sigma) = 0$ for all $\mu \in \mathbb{R}^n$). Compared to the subdifferential w.r.t. Σ (see Step 1), it is slightly more sophisticated to derive, for any given $\Sigma \in \mathbb{S}_+^n$ with $0 < \|\Sigma\| \leq 2C_2^2$, the subdifferential of the mapping $\mathbb{R}_{C_1,C_2}^n \to \mathbb{R}$, $\mu' \mapsto v(\mu', \Sigma)$ at $\mu \in \mathbb{R}_{C_1,C_2}^n$ (take into account that μ appears in the constraints of (26)). Our idea is to derive the Lagrange dual of program (26) and then apply the well-known Danskin's theorem.

Let $\Sigma \in \mathbb{S}^n_+$ with $0 < \|\Sigma\| \le 2C_2^2$ and $\mu \in \mathbb{R}^n_{C_1,C_2}$ be arbitrary but fixed. Problem (26) satisfies Slater's condition, i.e. there exists a feasible solution w_0 such that the inequality in the constraints is strict at w_0 (i.e. $w_0 > 0$). This means that strong duality holds. Consequently, we have

$$v(\mu, \Sigma) = \max_{\lambda_1, \lambda_2 \in \mathbb{R}, s \in \mathbb{R}^n_+} \min_{w \in \mathbb{R}^n} \left(\frac{1}{2} w^\mathsf{T} \Sigma w + \lambda_1 (w^\mathsf{T} \mu - z) + \lambda_2 (w^\mathsf{T} \mathbf{1} - 1) - s^\mathsf{T} w \right)$$
$$= \max_{\lambda_1, \lambda_2 \in \mathbb{R}, s \in \mathbb{R}^n_+} \min_{w \in \mathbb{R}^n} \left(\frac{1}{2} w^\mathsf{T} \Sigma w + (\lambda_1 \mu + \lambda_2 \mathbf{1} - s)^\mathsf{T} w - \lambda_1 z - \lambda_2 \right)$$
(45)

Let $u_{\Sigma,1}, \ldots, u_{\Sigma,n}$ be the eigenvalues of Σ in decreasing order (i.e. $u_{\Sigma,1} \ge u_{\Sigma,2} \ge \cdots \ge u_{\Sigma,r} > 0 = u_{\Sigma,r+1} = \cdots = u_{\Sigma,n}$) and $\Sigma = Q_{\Sigma}^{\mathsf{T}} U_{\Sigma} Q_{\Sigma}$ be the spectral decomposition of Σ with diagonal matrix $U_{\Sigma} := \operatorname{diag}(u_{\Sigma,1}, \ldots, u_{\Sigma,n})$. Then, if we set $U_{\Sigma}^{\dagger} := \operatorname{diag}(1/u_{\Sigma,1}, \ldots, 1/u_{\Sigma,r}, 0, \ldots, 0)$, we have

$$v(\mu, \Sigma) = \max_{\lambda_1, \lambda_2 \in \mathbb{R}, s \in \mathbb{R}^n_+} \left(-\frac{1}{2} (\lambda_1 \mu + \lambda_2 \mathbf{1} - s)^\mathsf{T} Q_{\Sigma}^\mathsf{T} U_{\Sigma}^\dagger Q_{\Sigma} (\lambda_1 \mu + \lambda_2 \mathbf{1} - s) - \lambda_1 z - \lambda_2 \right)$$

subject to $\lambda_1 \mu + \lambda_2 \mathbf{1} - s \in \operatorname{span}(\Sigma),$ (46)

where span(Σ) is the space spanned by the rows of Σ and the constraint $\lambda_1 \mu + \lambda_2 \mathbf{1} - s \in \text{span}(\Sigma)$ is obtained from the first order optimality condition with respect to w. By the way: In the case when Σ is of full rank, (46) reduces to $v(\mu, \Sigma) = \max_{\lambda_1, \lambda_2, s \in \mathbb{R}^n_+} (-\frac{1}{2}(\lambda_1 \mu + \lambda_2 \mathbf{1} - s)^{\mathsf{T}} \Sigma^{-1}(\lambda_1 \mu + \lambda_2 \mathbf{1} - s) - \lambda_1 z - \lambda_2)$.

Let $\mathcal{W}(\mu, \Sigma)$ and $\Lambda(\mu, \Sigma)$ denote the sets of optimal solutions of the max-min problem (45). Applying Danskin's theorem to (45), we have

$$\partial_{\mu}v(\mu,\Sigma) = \operatorname{conv}\left\{\lambda_{1}^{*}w^{*}: (\lambda_{1}^{*},\lambda_{2}^{*},s^{*}) \in \Lambda(\mu,\Sigma), w^{*} \in \mathcal{W}(\mu,\Sigma)\right\}.$$
(47)

For L_{2,C_1,C_2} -Lipschitz continuity of the mapping $\mathbb{R}^n_{C_1,C_2} \to \mathbb{R}$, $\mu' \mapsto v(\mu', \Sigma)$ at μ , it is sufficient to show that $\|\mu'\| \leq L_{2,C_1,C_2}$ for all $\mu' \in \partial_{\mu}v(\mu, \Sigma)$. In view of (47) and $\|\lambda_1^*w^*\| \leq |\lambda_1^*| \|w^*\| \leq |\lambda_1^*|$, it is sufficient to show that $|\lambda_1^*| \leq L_{2,C_1,C_2}$ for all $\lambda_1^* \in \Lambda(\mu, \Sigma)$. But this is a simple consequence of the following lemma. Recall that $z \in \mathbb{R}_+$ is an arbitrary constant.

Lemma 10.5. Let $\Sigma \in \mathbb{S}^n_+ \setminus \{\mathbf{0}\}$ and $\mu \in \mathbb{R}^n \setminus \{a\mathbf{1} : a \in \mathbb{R}\}$. Then, for any $(\lambda_1^*, \lambda_2^*, s^*) \in \Lambda(\mu, \Sigma)$, we have

$$|\lambda_1^*| \le \frac{16\|\Sigma\|_2}{9(\|\mu\|^2 - (\mu^{\mathsf{T}}\mathbf{1})^2/n)} \left(z + \frac{\|\mu\|}{\sqrt{n}}\right) + \frac{8\|\Sigma\|_2}{9\sqrt{n}(\|\mu\|^2 - (\mu^{\mathsf{T}}\mathbf{1})^2/n)}.$$
(48)

Proof. Let $\Sigma \in \mathbb{S}^n_+ \setminus \{\mathbf{0}\}$ and $\mu \in \mathbb{R}^n \setminus \{a\mathbf{1} : a \in \mathbb{R}\}$ be arbitrary but fixed, and define a function $f : \mathbb{R} \times \mathbb{R} \times \mathbb{R}^n_+ \to \mathbb{R}$ by $f(\lambda_1, \lambda_2, s) := -\frac{1}{2}(\lambda_1\mu + \lambda_2\mathbf{1} - s)^{\mathsf{T}}\Sigma^{\dagger}(\lambda_1\mu + \lambda_2\mathbf{1} - s) - \lambda_1z - \lambda_2$, where $\Sigma^{\dagger} := Q_{\Sigma}^{\mathsf{T}}U_{\Sigma}^{\dagger}Q_{\Sigma}$ with Q_{Σ} and U_{Σ}^{\dagger} as introduced after (45). Let $(\lambda_1, \lambda_2, s) \in \{(\lambda'_1, \lambda'_2, s') \in \mathbb{R} \times \mathbb{R} \times \mathbb{R}^n_+ : \lambda'_1\mu + \lambda'_2\mathbf{1} - s' \in \operatorname{span}(\Sigma)\}$. Then we have

$$f(\lambda_{1}/2,\lambda_{2}/2,s/2) - f(\lambda_{1},\lambda_{2},s)$$

$$= \frac{3}{8}(\lambda_{1}\mu + \lambda_{2}\mathbf{1} - s)^{\mathsf{T}}\Sigma^{\dagger}(\lambda_{1}\mu + \lambda_{2}\mathbf{1} - s) + \frac{1}{2}\lambda_{1}z + \frac{1}{2}\lambda_{2}$$

$$\geq \frac{3}{8u_{1}}(\lambda_{1}\mu + \lambda_{2}\mathbf{1} - s)^{\mathsf{T}}(\lambda_{1}\mu + \lambda_{2}\mathbf{1} - s) + \frac{1}{2}\lambda_{1}z + \frac{1}{2}\lambda_{2}$$

$$= \frac{3}{8u_{1}}\left(\lambda_{1}^{2}\|\mu\|^{2} + \lambda_{2}^{2}\|\mathbf{1}\|^{2} + \|s\|^{2} + 2\lambda_{1}\lambda_{2}\mu^{\mathsf{T}}\mathbf{1} - \lambda_{1}\mu^{\mathsf{T}}s - \lambda_{2}\mathbf{1}^{\mathsf{T}}s\right) + \frac{1}{2}\lambda_{1}z + \frac{1}{2}\lambda_{2}$$

$$\geq \frac{3}{8u_{1}}\left(\lambda_{1}^{2}\|\mu\|^{2} + \lambda_{2}^{2}\|\mathbf{1}\|^{2} + 2\lambda_{1}\lambda_{2}\mu^{\mathsf{T}}\mathbf{1} - \frac{1}{4}\|\lambda_{1}\mu + \lambda_{2}\mathbf{1}\|^{2}\right) + \frac{1}{2}\lambda_{1}z + \frac{1}{2}\lambda_{2}$$

$$= \frac{3}{8u_{1}}\left(\frac{3}{4}\lambda_{1}^{2}\|\mu\|^{2} + \frac{3}{4}\lambda_{2}^{2}\|\mathbf{1}\|^{2} + \frac{3}{2}\lambda_{1}\lambda_{2}\mu^{\mathsf{T}}\mathbf{1}\right) + \frac{1}{2}\lambda_{1}z + \frac{1}{2}\lambda_{2}$$

$$\geq \frac{9}{32u_{1}}\left(\|\mu\|^{2} - \frac{(\mu^{\mathsf{T}}\mathbf{1})^{2}}{n}\right)\lambda_{1}^{2} + \left(\frac{1}{2}z - \frac{\|\mu\|}{2\sqrt{n}}\right)\lambda_{1} - \frac{2u_{1}}{9n},$$
(49)

where the second and third inequalities are satisfied by minimizing the corresponding terms with respect to s and λ_2 and the first inequality can be justified as follows. Since $\lambda_1 \mu + \lambda_2 \mathbf{1} - s \in \text{span}(\Sigma)$, there exists a $w \in \mathbb{R}^n$ such that $\lambda_1 \mu + \lambda_2 \mathbf{1} - s = \Sigma w$. Thus, we have $Q_{\Sigma}(\lambda_1 \mu + \lambda_2 \mathbf{1} - s) = U_{\Sigma} Q_{\Sigma} w$, which means that only the first r components of $Q_{\Sigma}(\lambda_1 \mu + \lambda_2 \mathbf{1} - s)$ are non-zero, i.e. $Q_{\Sigma}(\lambda_1 \mu + \lambda_2 \mathbf{1} - s) = (x_1, \ldots, x_r, 0, \ldots, 0)^{\mathsf{T}} =: x$ for some $x_1, \ldots, x_r \in \mathbb{R}$. Thus,

$$\begin{aligned} &(\lambda_1\mu + \lambda_2 \mathbf{1} - s)^{\mathsf{T}} \Sigma^{\dagger} (\lambda_1\mu + \lambda_2 \mathbf{1} - s) \\ &= \left(Q_{\Sigma} (\lambda_1\mu + \lambda_2 \mathbf{1} - s) \right)^{\mathsf{T}} U_{\Sigma}^{\dagger} Q_{\Sigma} (\lambda_1\mu + \lambda_2 \mathbf{1} - s) \\ &= x^{\mathsf{T}} U^{\dagger} x = \sum_{i=1}^{r} \frac{1}{u_i} x_i^2 \\ &\geq \frac{1}{u_1} \sum_{i=1}^{r} x_i^2 = \frac{1}{u_1} x^{\mathsf{T}} x = \frac{1}{u_1} (\lambda_1\mu + \lambda_2 \mathbf{1} - s)^{\mathsf{T}} Q_{\Sigma}^{\mathsf{T}} Q_{\Sigma} (\lambda_1\mu + \lambda_2 \mathbf{1} - s) \\ &= \frac{1}{u_1} (\lambda_1\mu + \lambda_2 \mathbf{1} - s)^{\mathsf{T}} (\lambda_1\mu + \lambda_2 \mathbf{1} - s), \end{aligned}$$

which shows that the first inequality in (49) does indeed hold.

Now, we can conclude from (49) that for any $\lambda_1 \in \mathbb{R}$ with

$$|\lambda_1| > \frac{16u_1}{9(\|\mu\|^2 - (\mu^{\mathsf{T}}\mathbf{1})^2/n)} \left(z + \frac{\|\mu\|}{\sqrt{n}}\right) + \frac{8u_1}{9\sqrt{n}(\|\mu\|^2 - (\mu^{\mathsf{T}}\mathbf{1})^2/n)},$$

the term $f(\lambda_1/2, \lambda_2/2, s/2) - f(\lambda_1, \lambda_2, s)$ is strictly positive. This implies (48).

References

- Z.D. Bai, B.Q. Miao, and G.M. Pan. On asymptotics of eigenvectors of large sample covariance matrix. *The Annals of Probability*, 35(4):1532–1572, 2007.
- [2] O. Banerjee, L. El Ghaoui, and A. d'Aspremont. Model selection through sparse maximum likelihood estimation for multivariate Gaussian or binary data. J. Mach. Learn. Res., 9:485–516, 2008.
- [3] C.W.J. Beenakker. Random-matrix theory of quantum transport. *Reviews of modern physics*, 69(3):731, 1997.
- [4] F. Bellini, T. Fadina, R.D. Wang, and Y.R. Wei. Parametric measures of variability induced by risk measures. *Insurance: Mathematics and Economics*, 106:270–284, 2022.
- [5] C. Berge. Topological spaces: Including a treatment of multi-valued functions, vector spaces and convexity. Oliver & Boyd, 1877.
- [6] J. Bien and R.J. Tibshirani. Sparse estimation of a covariance matrix. *Biometrika*, 98(4):807–820, 2011.
- [7] V. Chandrasekaran, P.A. Parrilo, and A.S. Willsky. Latent variable graphical model selection via convex optimization. *The Annals of Statistics*, 40(4):1935–1967, 2012.

- [8] F.H. Clarke. Optimization and nonsmooth analysis. SIAM, 1990.
- [9] R. Cont, R. Deguest, and G. Scandolo. Robustness and sensitivity analysis of risk measurement procedures. *Quantitative Finance*, 10(6):593–606, 2010.
- [10] A. d'Aspremont, O. Banerjee, and L. El Ghaoui. First-order methods for sparse covariance selection. SIAM Journal on Matrix Analysis and Applications, 30(1):56–66, 2008.
- [11] C. Davis and W.M. Kahan. The rotation of eigenvectors by a perturbation. iii. SIAM Journal on Numerical Analysis, 7(1):1–46, 1970.
- [12] A.P. Dempster. Covariance selection. *Biometrics*, 28(1):157–175, 1972.
- [13] A.L. Dontchev and R.T. Rockafellar. Implicit functions and solution mappings, volume 543. Springer, 2009.
- [14] M. Drton and T.S. Richardson. Graphical methods for efficient likelihood inference in Gaussian covariance models. *Journal of Machine Learning Research*, 9:893–914, 2008.
- [15] A. Edelman and N.R. Rao. Random matrix theory. Acta Numerica, 14:233–297, 2005.
- [16] F.J. Fabozzi, H.M. Markowitz, and F. Gupta. Portfolio selection. Handbook of Finance, 2, 2008.
- [17] R.A. Fisher. The use of multiple measurements in taxonomic problems. Annals of Eugenics, 7(2):179–188, 1936.
- [18] J. Friedman, T. Hastie, and R. Tibshirani. Sparse inverse covariance estimation with the graphical lasso. *Biostatistics*, 9(3):432–441, 2008.
- [19] C. R. Givens and R. M. Shortt. A class of Wasserstein metrics for probability distributions. Michigan Mathematical Journal, 31(2):231–240, 1984.
- [20] S. Guo and H. Xu. Statistical robustness in utility preference robust optimization models. Mathematical Programming, 190(1-2):679–720, 2021.
- [21] S. Guo and H. Xu. Data perturbations in stochastic generalized equations: statistical robustness in static and sample average approximated models. *Mathematical Programming*, 202(1-2):135– 168, 2023.
- [22] S. Guo, H. Xu, and L. Zhang. Existence and approximation of continuous Bayesian Nash equilibria in games with continuous type and action spaces. SIAM Journal on Optimization, 31(4):2481– 2507, 2021.
- [23] S. Guo, H. Xu, and L. Zhang. Statistical robustness of empirical risks in machine learning. Journal of Machine Learning Research, 24(125):1–38, 2023.
- [24] A. Gut. Marcinkiewicz laws and convergence rates in the law of large numbers for random variables with multidimensional lidices. *The Annals of Probability*, 6(3):469–482, 1978.
- [25] F.R. Hampel. A general qualitative definition of robustness. The Annals of Mathematical Statistics, 42(6):1887–1896, 1971.

- [26] P.J. Huber. *Robust statistics*, volume 523. John Wiley & Sons, 2004.
- [27] P.J. Huber and E.M. Ronchetti. *Robust statistics*. John Wiley & Sons, 2011.
- [28] S. Janson, T. Luczak, and A. Rucinski. Random graphs. John Wiley & Sons, 2011.
- [29] I.M. Johnstone. On the distribution of the largest eigenvalue in principal components analysis. The Annals of Statistics, 29(2):295–327, 2001.
- [30] P. Koch-Medina and C. Munari. Law-invariant risk measures: Extension properties and qualitative robustness. *Statistics & Risk Modeling*, 31(3-4):215–236, 2014.
- [31] V. Krätschmer, A. Schied, and H. Zähle. Qualitative and infinitesimal robustness of tail-dependent statistical functionals. *Journal of Multivariate Analysis*, 103(1):35–47, 2012.
- [32] V. Krätschmer, A. Schied, and H. Zähle. Comparative and qualitative robustness for law-invariant risk measures. *Finance and Stochastics*, 18:271–295, 2014.
- [33] V. Krätschmer, A. Schied, and H. Zähle. Domains of weak continuity of statistical functionals with a view toward robust statistics. *Journal of Multivariate Analysis*, 158:1–19, 2017.
- [34] V. Krätschmer and H. Zähle. Statistical inference for expectile-based risk measures. Scandinavian Journal of Statistics, 44(2):425–454, 2017.
- [35] C. Lam and J. Fan. Sparsistency and rates of convergence in large covariance matrix estimation. Annals of Statistics, 37(6B):4254, 2009.
- [36] R.C. Li. Matrix perturbation theory. In Handbook of Linear Algebra, pages 15–1. Chapman and Hall/CRC, 2006.
- [37] C. Louart, Z. Liao, and R. Couillet. A random matrix approach to neural networks. *The Annals of Applied Probability*, 28(2):1190–1248, 2018.
- [38] V.A. Marchenko and L.A. Pastur. Distribution of eigenvalues for some sets of random matrices. *Matematicheskii Sbornik*, 114(4):507–536, 1967.
- [39] N. Meinshausem and P. Bühlmann. High-dimensional graphs and variable selection with the Lasso. *The Annals of Statistics*, 34(3):1436–1462, 2006.
- [40] V.A. Nguyen, D. Kuhn, and P. Mohajerin Esfahani. Distributionally robust inverse covariance estimation: The wasserstein shrinkage estimator. *Operations research*, 70(1):490–515, 2022.
- [41] P. Ravikumar, M.J. Wainwright, G. Raskutti, and B. Yu. High-dimensional covariance estimation by minimizing l1-penalized log-determinant divergence. *Electronic Journal of Statistics*, 5:935– 980, 2011.
- [42] W. Römisch. Stability of stochastic programming problems. Handbooks in operations research and management science, 10:483–554, 2003.
- [43] A.J. Rothman, P.J. Bickel, E. Levina, and J. Zhu. Sparse permutation invariant covariance estimation. *Electronic Journal of Statistics*, 2:494–515, 2008.

- [44] J.W. Silverstein. Describing the behavior of eigenvectors of random matrices using sequences of measures on orthogonal groups. SIAM Journal on Mathematical Analysis, 12(2):274–281, 1981.
- [45] G.W. Stewart. Matrix algorithms: Volume 1: Basic decompositions. SIAM, 1998.
- [46] T. Tao. Topics in random matrix theory, volume 132. American Mathematical Society, 2023.
- [47] A.M. Tulino and S. Verdú. Random matrix theory and wireless communications. Foundations and Trends (R) in Communications and Information Theory, 1(1):1–182, 2004.
- [48] C. Wang, D. Sun, and K. C. Toh. Solving log-determinant optimization problems by a Newton-CG primal proximal point algorithm. SIAM Journal on Optimization, 20(6):2994–3013, 2010.
- [49] W. Wang, H. Xu, and T. Ma. Quantitative statistical robustness for tail-dependent law invariant risk measures. *Quantitative Finance*, 21(10):1669–1685, 2021.
- [50] S. Weber. Solvency II, or how to sweep the downside risk under the carpet. *Insurance: Mathe*matics and Economics, 82:191–200, 2018.
- [51] E.P. Wigner. On the distribution of the roots of certain symmetric matrices. Annals of Mathematics, 67(2):325–327, 1958.
- [52] J. Wishart. The generalised product moment distribution in samples from a normal multivariate population. *Biometrika*, 20A(1-2):32–52, 1928.
- [53] N. Xia, Y. Qin, and Z. Bai. Convergence rates of eigenvector empirical spectral distribution of large dimensional sample covariance matrix. *The Annals of Statistics*, 41(5):2572–2607, 2013.
- [54] J. Yang, D. Sun, and K. C. Toh. A proximal point algorithm for log-determinant optimization with group lasso regularization. SIAM Journal on Optimization, 23(2):857–893, 2013.
- [55] Y. Yu, T. Wang, and R.J. Samworth. A useful variant of the Davis–Kahan theorem for statisticians. *Biometrika*, 102(2):315–323, 2015.
- [56] M. Yuan and Y. Lin. Model selection and estimation in the Gaussian graphical model. *Biometrika*, 94(1):19–35, 2007.
- [57] H. Zähle. Qualitative robustness of statistical functionals under strong mixing. *Bernoulli*, 21(3):1412–1434, 2015.
- [58] H. Zähle. A definition of qualitative robustness for general point estimators, and examples. Journal of Multivariate Analysis, 143:12–31, 2016.
- [59] S. Zhang, H. Xu, and H. Sun. Statistical robustness of kernel learning estimator with respect to data perturbation. *arXiv preprint arXiv:2406.10555*, 2024.