

Photometric White Dwarf Rotation

GABRIELA OLIVEIRA DA ROSA,¹ S. O. KEPLER,¹ L. T. T. SOETHE,^{2,1} ALEJANDRA D. ROMERO,¹ AND KEATON J. BELL³

¹*Instituto de Física, Universidade Federal do Rio Grande do Sul
91501-970 Porto Alegre, RS, Brazil*

²*Instituto Federal de Educação, Ciência e Tecnologia Sul-rio-grandense
96745-000 Charqueadas, RS, Brazil*

³*Department of Physics, Queens College, City University of New York,
Flushing, NY, 11367, USA*

ABSTRACT

We present a census of photometrically detected rotation periods for white dwarf stars. We analyzed the light curves of 9285 white dwarf stars observed by the Transiting Exoplanet Survey Satellite (TESS) up to sector 69. Using Fourier transform analyses and the TESS_LOCALIZE software, we detected variability periods for 318 white dwarf stars. The 115 high probability *likely single* white dwarfs in our sample have a median rotational period of 3.9 hours and a median absolute deviation of 3.5 h. Our distribution is significantly different from the distribution of the rotational period from asteroseismology, which exhibits a longer median period of 24.3 hours and a median absolute deviation of 12.0 h. In addition, we reported non-pulsating periods for three known pulsating white dwarfs with rotational period previously determined by asteroseismology: NGC 1501, TIC 7675859, and G226-29. We also calculated evolutionary models that include six angular momentum transfer mechanisms from the literature throughout evolution in an attempt to reproduce our findings. Our models indicate that the temperature-period relation of most observational data is best fitted by models with low metallicity, probably indicating problems with the computations of angular-momentum loss during the high-mass-loss phase. Our models also generate internal magnetic fields through the Tayler–Spruit dynamo.

INTRODUCTION

The rotation period is an essential parameter for the study of stellar evolution. It plays a significant role in the stellar dynamo mechanism and is directly related to the mass loss experienced by stars during the asymptotic giant branch (AGB, e.g., Dominguez et al. 1999; Catalán et al. 2008; Romero et al. 2015; Choi et al. 2016). For single stars and stars in binary systems, photometry allows us to detect the rotation period of stars directly and indirectly through asteroseismology (e.g. Córscico et al. 2019).

Binary stellar systems can exhibit light variability due to different effects. Depending on the geometry of the system and the orientation of the orbital plane as observed from Earth, the light variability can provide the rotational period of stars or the orbital period of the system. Variability resulting from ellipsoidal variation or reflection in binary systems provides stellar rotation periods, whereas variability from orbital motion indicates the system’s orbital period (e.g. Krtićka et al. 2023). However, in the case of spin-orbit coupling, there is no difference between the orbital and rotational periods.

The spin-orbit coupling occurs when there is mass loss due to winds, trying to balance the variation in the spin of each star with the angular momentum of the system. Zahn (1977) demonstrated that the synchronization time between stellar rotation and the orbital period of a binary system is proportional to P_{orb}^4 , where P_{orb} is the orbital period of the system. This implies that the shorter the orbital period, the faster the synchronization process occurs. Zahn (1977), suggests that systems with orbital periods shorter than 17 days are synchronized. On the other hand, for compact stars such as white dwarfs, Fuller & Lai (2011); Pablo et al. (2012); Fuller & Lai (2013); Fuller (2017); Fuller & Mathis (2023) estimate that synchronization may occur only for orbital periods of the order of 1 hour.

Single stars, such as the Sun, may also exhibit light variability because of the presence of dark spots and patches on their surface. These inhomogeneities are typically associated with the presence of a magnetic field or with a variation in the chemical composition across the stellar surface (e.g. Babcock 1960; Kochukhov 2011). In this case, the rotation period as well as some harmonics

can be present in the frequency spectrum, depending on the number of spots.

White dwarf (WD) stars are the most common end of stellar evolution, constituting the fate of 95 – 97% of all stars in the Galaxy (e.g. García-Berro et al. 1997; Poelarends et al. 2008; Siess 2010; Langer 2012; Woosley & Heger 2015; Doherty et al. 2014). As fossils, WDs preserve valuable information on their past evolution, making them essential for studying the structure, evolution, chemical enrichment, and star formation history of our Galaxy (Diaz-Pinto et al. 1994). WDs are classified into spectral classes based on the dominant chemical element on their surface. Approximately 80% of all WDs belong to the DA spectral type (e.g. Kepler et al. 2021), that is, they exhibit H-dominated spectra. The remaining 20% consist mainly of WD stars of the spectral type DB and DO, featuring helium-dominated atmospheres.

Although WDs have been studied since 1915 (Adams 1915), their rotation period distribution is still largely undetermined. Ground-based observations cannot easily measure rotation periods because of atmospheric and instrumental limitations, as WDs are intrinsically faint stars. Furthermore, the time gap due to daylight constrains the period range that can be determined (Nather et al. 1990). Long-time baseline observations from space-based telescopes have largely overcome the latter problem. The Kepler spacecraft was launched in 2009, obtaining excellent data to search for stellar variability (e.g. Molnár et al. 2016). However, the failure of its second reaction wheel in 2013 caused significant noise problems at low frequencies. It presented a serious challenge in identifying rotation periods for faint stars. The launch of the Transiting Exoplanet Survey Satellite (TESS) in 2018 (Oschmann et al. 2014) allowed for a reliable analysis of low frequencies for WD stars. Even with some gaps, the TESS data enabled the detection of photometric rotation periods directly in the frequency spectrum (e.g. Labadie-Bartz et al. 2023). Finally, photometry analysis also benefits from the high-precision astrometry obtained by the Gaia mission (Gaia Collaboration et al. 2018), which has considerably improved the determination of the parallax and the positions of each star.

Stellar rotation rates can also be estimated through spectroscopic observations, by measuring the rotational broadening of the H α line or the Ca II K line, as in Berger et al. (2005). This method has provided rotational velocity estimates for more than 80 WDs, although, for the majority of the sample, it has allowed only the determination of an upper limit. The results indicate that most WDs exhibit rotational velocities lower

than 15 km/s, which corresponds to rotation periods longer than 1 h for a $0.6 M_{\odot}$ WD.

Another possibility to estimate rotational periods is through asteroseismology (e.g., Kepler & Romero 2017), by measuring the separation between the components of multiplets. Hermes et al. (2017b) determined the rotation period for twenty WDs, all pulsating hydrogen atmosphere DAVs, using photometric data from the Kepler spacecraft. The authors combined these results with the early estimates compiled by Kawaler (2015), and found a mean rotation period of 35 ± 28 h for a sample of 40 WDs, indicating a slight tendency to shorter periods for higher masses.

Our work presents the first analysis of the rotation period distribution of a large sample of WD stars based on photometric data from the TESS telescope. We analyze the 21 832 light curves of the 9285 white dwarfs observed by TESS from sectors 1 to 69. We also compile all known WDs with rotation periods determined by asteroseismology and search for non-pulsating periods using TESS data. We search for stable variability and use the TESS_LOCALIZE software to confirm the signal source, as presented in Section 1. In Section 2, we present a compilation of all WDs with the rotation period estimated by asteroseismology to date, which we refer to as the seismological sample. In Section 3, we present the rotation period distribution for 318 WDs, analyze its dependence on the stellar parameters, and compare it with the seismological sample. Additionally, we also report the non-pulsating variabilities we detect for three known pulsating WDs from the seismological sample. In Section 4, we discuss our theoretical model computations aimed at reproducing our observational results. Finally, we present our concluding remarks in Section 5.

1. DATA ANALYSIS

We selected all known WDs (e.g., Wenger et al. 2000; Dufour et al. 2017; Kepler et al. 2019; Tremblay et al. 2020; Kilic et al. 2020; Kepler et al. 2021; Guo et al. 2022; Amorim et al. 2023; O’Brien et al. 2023), including Gaia candidates from Gentile Fusillo et al. (2021), and matched them with the TESS telescope data for objects up to magnitude $G = 17.5$ — the observational limit that we determined for the white dwarf analysis using the TESS data. This process yielded a sample of 9285 stars, each of which was observed for 1 to 29 sectors. We analyze the available 21 832 light curves for all 9285 white dwarfs observed with TESS from sectors 1 (beginning 25 July 2018) to 69 (ending 20 September 2023). All the TESS data used in this article can be found in MAST (Team 2021a,b,c,d).

We use the LIGHTKURVE package (Lightkurve Collaboration et al. 2018) to download the photometric data observed by TESS. We selected data taken with a cadence of 120 s, and 20 s when available, and processed using the SPOC pipeline (Caldwell et al. 2020). The TESS telescope observes each sector for 27 days, interrupting the observation every ~ 13 days - or less - to send the data to Earth. Due to these gaps in the light curve, we searched for stable variability only for periods up to 13 days.

We combined the light curves of all sectors in which the star was observed and performed a complete Fourier transform (FT) on it. We computed the false alarm probability (FAP) of 1/1000, that is, the limit beyond which peaks on the Fourier transforms have less than one chance in 1000 of being due to noise. We use this detection limit to define whether the observed signal is significant. The FAP was computed by randomizing (Monte Carlo-like simulation) the fluxes at the observed timings, as described in Kepler (1993). If the FT exhibited any peak with an amplitude above the detection limit, we considered its period as a potential variability signal from the WD. When confronted with the detection of multiple harmonic frequencies in the FT, we analyzed the shape of the dips in brightness and examined the light curve folded in phase to determine the variability period. When a noisy light curve hindered the observation of the variability curve, our approach was to select the harmonic peak with the lowest frequency.

Following these criteria, we detected variability periods for the 2728 WDs. To avoid including pulsation periods, we checked whether the temperature of the candidate star with periods shorter than 1500 s matched the temperature range of the DOV, DBV, and DAV/ELMV white dwarf instability strips, and also for linear combination of pulsation frequencies. As we are interested here in rotation periods, we excluded periods that matched any of these pulsation criteria. The excluded stars include 64 known pulsators, 207 new pulsators, and 93 pulsator candidates. We also did not include in this work the 173 variability periods from CVs. For the remaining 2191, we performed a simple selection using TESS_LOCALIZE. We selected those for which the software indicated that for at least for the first observed sector, the WD was the most likely signal source, eliminating 61% of the targets. In this way, we compile a target list of 845 WDs with a detected non-pulsator variability frequency, which corresponds to 9% of the initial list of 9285 WDs. This was a preliminary selection. In Section 1.1, we describe the stricter selection criteria applied to this initial sample of 845 WDs using TESS_LOCALIZE.

We identified within the WDs that make up our sample those with a detected magnetic field, as well as those in binary systems. For magnetic WDs, we first used the classification from the catalog Amorim et al. (2023) and the literature for other objects (e.g., McCook & Sion 2016; Wenger et al. 2000; Kepler et al. 2021; O’Brien et al. 2023; Tremblay et al. 2020; Kilic et al. 2020; Guo et al. 2022). We used the classification of WDs in binary systems from the literature (e.g. Wenger et al. 2000; Dufour et al. 2017) and also detected new systems by inspecting the Gaia proper motion and parallax data within a box of $120''$ around the white dwarf and by checking the variability shape of the light curve from TESS data.

In addition, we compiled all 63 WDs with rotation periods determined by asteroseismology up to now — which we refer to as seismological sample — and matched them with the TESS data. 45 of these pulsating WDs were in our initial list of WDs observed by TESS. For these stars, we inspect all the light curves, searching for reliable non-pulsating periods. To check the presence of a rotation period in the FT, we identified the pulsation frequencies and their linear combination. Our initial sample of 845 WDs with detected non-pulsating periods includes 14 WDs from the seismological sample.

1.1. Localizing the signal

The TESS telescope offers long-time light curves and high-frequency resolution. However, because of its large plate scale of $21'' \times 21''$ per pixel, it exhibits low spatial resolution. Consequently, signals from nearby sources contaminate many TESS light curves. We overcome this problem using the TESS_LOCALIZE software (Higgins & Bell 2023) to determine whether the detected period arises from the white dwarf or a nearby star. For each star in our target list, we run TESS_LOCALIZE for all sectors in which the star was observed, setting the *Principal Components* parameter to 3 (see Higgins & Bell 2023).

There are two crucial TESS_LOCALIZE output parameters that need to be checked to ensure the quality of the fit: *Height* and *p-value*. When TESS_LOCALIZE fits the position of the signals, the strength of the signals is represented as the best-fit *Height* parameter. Similarly to the challenge of identifying stars in an image, the signal needs to stand out significantly from the background noise. Higgins & Bell (2023) recommends that the *Height* parameter exceed five times its uncertainty for the localization to be considered significantly above the noise floor. Another critical parameter is the *p-value*; this number supports the hypothesis that the

location of the fit of the signal is consistent with the position of the source. Because we do not reject this hypothesis, the p -value should be reasonably high. We adopt the same threshold as Pedersen & Bell (2023): ' p -value' ≥ 0.05 . Applying these criteria, we selected only those sectors for which TESS_LOCALIZE achieved a significant localization of the signal of interest that is consistent with the location of a known *Gaia* source brighter than $G = 18$ mag. Therefore, since we excluded noisy sectors, we do not need to be concerned about the variation in noise from sector to sector.

The final constraint step involves an examination of the *Relative Likelihood*, also an output parameter of TESS_LOCALIZE. Essentially, it ranks the proximity of *Gaia* sources to the localized position of the signal within the pixels. For the same target, TESS_LOCALIZE can fit distinct values of *Relative Likelihood* for each available sector of data. So, for analysis purposes, we denoted the mean *Relative Likelihood* over all selected sectors as *Like*. To be consistent, we only consider that the signal originates from the white dwarf if TESS_LOCALIZE indicates the white dwarf as the most probable source for all selected sectors.

The two-sample Kolmogorov–Smirnov (K-S) test is a robust statistical method used to assess the likelihood of two distinct samples originating from the same population. To ensure that we are not including false positive data in our sample, we performed the two-sample K-S test comparing samples with different thresholds on the selection parameter *Like*. The K-S tests indicated that we would be discarding valuable data if we embraced a *Like* limit higher than 0.75. Given that, and the fact that the software estimates $Like \simeq 0.75$ even for confirmed periods from known variables (see Appendix A - Known RR Lyrae Star), we selected all the stars with $Like \geq 0.75$.

The TESS_LOCALIZE analysis with strict conditions reduced by 62% our initial sample of 845 candidate stars. However, the sample of 845 WD is already a consequence of a preliminary selection using TESS_LOCALIZE that resulted in a reduction of 61%. Therefore, in total, TESS_LOCALIZE reduced our sample of 2191 WDs with non-pulsating period detected in 85%. It is important to point out that, in this second selection, 79% of the stars discarded by TESS_LOCALIZE were rejected because the software was unable to perform a high-quality fit, presenting low ' p -value' and ' $height$ '. Only 21% were rejected because the software indicated a low probability of the signal having the WD as the source.

Our *Complete sample* consists of 318 WDs, including 76 in binary systems and 30 with a detected magnetic fields, counting 1 WD in a binary system that exhibits

a magnetic field. Table 5 in Appendix D - Variabilities Periods Table lists all stars that compose our samples, informing its variability period, amplitude, *Like*, mean *Height* parameter normalized by uncertainty, temperature, mass, and additional information. Table 5 also reports the fraction of sectors whose TESS_LOCALIZE fit achieved our quality criteria as ' Q/S '. White dwarfs confirmed to be in binary systems or with a detected magnetic field can be identified in Table 5 by the column *Info*. The character '+' indicates binarity (e.g. DA+M and WD+pair), while the characters 'H' and 'P' indicate magnetic field (e.g. DAH and DAP). WDs in binary systems with the designation 'WD+pair' were identified through proper motion and parallax, while the ones with the designation 'eclipse' were identified by the shape of the light curve variability.

In addition to the sample of 318 WDs, we found that for only 3 of the 14 stars in our seismological sample, the WD is also the source of the non-pulsating periods we detected. These three stars are discussed separately in Section 3.4. Table 3 presents the details of the non-pulsating periods we identified, including amplitude, and quality parameters from the TESS_LOCALIZE fit.

1.2. Samples

Stars in binary systems have a completely different evolution from single stars if their companions are close enough to interact, affecting their structure. Consequently, variability periods from close binary systems represent a distinct population from single WD rotational periods. Given that, we divided our sample of 318 WDs between those that probably evolved as single stars and those that might have evolved by binary interaction.

Our *Complete sample* comprises 76 WDs in binary systems: 22 identified through parallax and proper motion, 16 identified by the light curve shape, and 38 identified through spectra analysis from literature. The WDs with pairs we identified through parallax and proper motion are all dozens of AU away from their companions; consequently, they probably evolved single stars. The binary systems in our sample identified through spectral observations comprise a WD and a main-sequence star of spectral type M, L, or K. About 75% percent of the stars in this type of binary system are far enough from their pair to evolve as single stars (Willems & Kolb 2004). However, we can not distinguish the specific evolutionary status of each system; thus, we will also assume that the 38 WDs with pairs identified by spectra analysis had their evolution affected by their pairs. Finally, the 16 WDs that show eclipses on their light curves

are likely to be close to their pairs, since the periodicities of the eclipses are shorter than 13 days for all cases. Therefore, we select a sample labeled *WDs with close pair* encompassing the 54 WDs in our sample with pairs identified by spectra and eclipses.

It is important to recognize that our sample may include more WDs originally in binary systems than those that we detect. WDs with a mass lower than $0.45 M_{\odot}$ are formed by binary interaction (e.g. Marsh et al. 1995; Kilic et al. 2007) since it would take longer than the age of the Universe for them to evolve as single stars. This does not necessarily imply that these stars are currently in binary systems, but they certainly originate from such a population. Therefore, we classified the stars in our sample with a mass less than $0.45 M_{\odot}$ as *Potential WDs with Pair*. We also included stars with an effective temperature higher than 40 000 K based on the fact that WDs with close pairs might exhibit overestimated temperatures due to the reflection effect caused by a brighter companion (e.g. Schaffenroth et al. 2023; Steen et al. 2024). Once high-temperature WDs are uncommon, we assumed that WDs with temperatures higher than 40 000 K, an arbitrary choice, are more likely to have a pair than to be a single high-temperature WD. Furthermore, we assumed that the remaining stars — those neither confirmed to have companions nor meet the criteria for potential pairs — are *Likely single WDs*.

We performed a K-S test comparing the sample of *Potential WDs with pair* with the sample of *WDs with close pairs*; the test indicated that these samples may originate from the same population. That suggested that the mass and temperature parameters that we used as a criterion to divide our sample of WDs with no confirmed pairs into *Likely single WDs* and *Potential WDs with pair* were well chosen.

Sample	Stars	Mag.	$\bar{P} \pm \text{MAD}$ (h)
<i>Complete sample</i>	318	30	6.8 ± 5.2
<i>WDs with close pairs</i>	54	1	8.1 ± 5.4
<i>Potential WDs with pairs</i>	149	4	7.6 ± 5.2
<i>Likely single WDs</i>	115	25	3.9 ± 3.5

Table 1. Compilation of our samples’ information: their star number, number of magnetic WD, their median period (\bar{P}), and median absolute deviation (MAD).

Our *Complete Sample* consists of 318 WDs, including 54 *WDs with close pairs*, 149 *Potential WDs with pairs*, and 115 *Likely single WDs*. Further details of these samples are available in Table 1, which summarizes information on the quantities of magnetic WD in

each sample, their median periods \bar{P} , and the period median absolute deviation MAD.

2. COMPILATION OF ROTATION PERIODS FROM ASTEROSEISMOLOGY

This section compiles results from the literature to compare them with our findings. Consequently, this section does not present the results of our work using TESS data.




Figure 1. Rotation period versus mass relation, for period determinations from asteroseismology. The period axis is in log scale. Black, cyan, and pink dots indicate DAV, DBV, and DOV stars respectively.

Kawaler (2015), Hermes et al. (2017b), and Kepler & Romero (2017) reported rotation periods of WDs using determinations from asteroseismological analysis. Extending their work, we compile all 63 WD stars with reported seismological rotation periods up to the present. Their rotation periods are listed in Table 2 as well as their spectroscopy mass and spectral type. Most of the mass reported were taken from Gentile Fusillo et al. (2021) because they include the measured parallax. The seismological rotational periods have a median rotational period of 24.3 hours and a median absolute deviation of 12.0 h - its histogram is shown in the last line of Figure 7. Investigating the trend of decreasing rotation period with increasing white dwarf mass claimed by Hermes et al. (2017b), we present Figure 1. The colored dots indicate the spectral types of stars according to the legend. The current plot does not show this trend; instead, the data appear to be scattered around a central point.

3. RESULTS

Star	P_{seism} [h]	Type	Mass [M_{\odot}]	Ref.	Star	P_{seism} [h]	Type	Mass [M_{\odot}]	Ref.
TIC 7675859	7.15	DAV	0.900 ± 0.005	0	TIC 55650407	13.56	DAV	0.576 ± 0.009	0
TIC 79353860	17.06	DAV	0.571 ± 0.002	0	TIC 149863849	18.07	DAV	0.652 ± 0.015	0
TIC 230384389	22.32	DAV	0.642 ± 0.027	0	TIC 278616553	22.81	DAV	0.609 ± 0.022	0
TIC 343296348	12.35	DAV	0.587 ± 0.011	0	TIC 353727306	4.51	DAV	0.615 ± 0.013	0
TIC 800126377	15.1	DAV	0.675 ± 0.012	0	TIC 900762564	1.34	DAV	0.524 ± 0.00	0
TIC 1102242692	24.14	DAV	0.531 ± 0.025	0	TIC 394015496	29.76	DAV	0.607 ± 0.013	1
TIC 21187072	53.76	DAV	0.3138 ± 0.0068	1	PG 1159-035	33.6	DOV	0.54 0.07	2
RX J2117.1+3412	25	DOV	0.716 0.150	3	EPIC 228782059	34.1	DBV	0.65 0.13	4
GD133	168	DAV	0.6257 0.0098	5	EPIC220274129	12.7	DAV	0.560 0.024	6
PG 0112+104	10.2	DBV	0.539 0.017	7	EPIC220347759	31.7	DAV	0.589 0.040	8
KIC 4552982	18.4	DAV	0.700 0.044	8	KIC 4357037	22	DAV	0.522 0.060	8
KIC 7594781	26.8	DAV	0.674 0.058	8	KIC 10132702	11.2	DAV	0.52 0.13	8
GD 1212	6.9	DAV	0.608 0.010	8	EPIC201719578	26.8	DAV	0.561 0.082	8
EPIC201730811	2.6	DAV	0.608 0.010	8	EPIC201802933	31.3	DAV	0.648 0.048	8
EPIC210397465	49.1	DAV	0.437 0.019	8	EPIC211596649	81.8	DAV	0.79 0.16	8
EPIC211629697	64	DAV	0.534 0.076	8	EPIC211914185	1.1	DAV	0.84 0.13	8
EPIC211926430	25.4	DAV	0.586 0.045	8	EPIC228682478	109.1	DAV	0.726 0.080	8
EPIC229227292	29.4	DAV	0.584 0.031	8	EPIC220204626	24.3	DAV	0.71 0.04	8
EPIC220258806	30	DAV	0.602 0.020	8	EPIC201806008	31.3	DAV	0.611 0.014	8
KUV02464+3239	90.7	DAV	0.633 0.021	9	SDSS J0349-0059	9.8	DOV	0.543 0.004	10
Ross 548	37.8	DAV	0.568 0.009	11	GD 165	57.3	DAV	0.642 0.010	11
WD 1711+657	16.4	DAV	0.884 0.022	12	PG1707+427	16	DOV	0.53 0.1	12
G29-38	32	DAV	0.628 0.016	12	GD 358	29	DBV	0.582 0.020	12
EC14012-1446	14.4	DAV	0.632 0.031	12	EC20058-5234	2	DBV	0.549 0.024	12
KIC 11911480	74.7	DAV	0.592 0.056	13	SDSS J1612+830	0.96	DAV	0.848 0.047	14
WD 0937+010	11.8	DAV	0.451 0.053	14	KUV11370+4222	5.7	DAV	0.605 0.028	15
HS 0507+0434B	40.9	DAV	0.681 0.017	16	GD 154	55.2	DAV	0.645 0.014	16
KIC 8626021	40.8	DBV	0.68 0.13	17	PG 0122+200	37.2	DOV	0.53 ± 0.1	18
HL Tau 76	52.8	DAV	0.516 0.015	19	G185-32	14.5	DAV	0.6320 0.0086	20
L19-2	13.0	DAV	0.6742 0.0088	21	LP 133-144	41.8	DAV	0.557 0.012	22
NGC 1501	28.1	DOV	0.56 0.04	23	G226-29	8.9	DAV	0.7829 0.0091	24
PG 2131+066	5.1	DOV	0.55 0.04	25					

Table 2. White dwarfs with rotation periods determined previously by asteroseismology. References:

(0)Romero et al. (2024); (1)Romero et al. (2022);(2)Oliveira da Rosa et al. (2022); (3)C3rsico et al. (2021);(4) (Duan et al. 2021); (5) Fu et al. (2019);(6) Bell et al. (2017); (7) Hermes et al. (2017a); (8) Hermes et al. (2017b); (9) Li et al. (2017); (10) Calcaferro, Leila M. et al. (2016); (11) Giammichele et al. (2016); (12) Kawaler (2015); (13) Greiss et al. (2014); (14) Castanheira et al. (2013); (15) Su et al. (2013); (16) Fu et al. (2012); (17) Østensen et al. (2011); (18) Fu, J.-N. et al. (2007); (19) Dolez, N. et al. (2006);(20) Pech, D. & Vauclair, G. (2006); (21) Bradley (2001); (22) Pfeiffer et al. (1996); (23) Bond et al. (1996a); (24) Kepler et al. (1995); (25) Kawaler et al. (1995).

Assuming that any inhomogeneity on the stellar surface will manifest itself as variability with the rotation period of the surface, we attribute the variability periods detected for the *Likely single WDs* to rotation. This consideration follows the exclusion of all periods potentially caused by pulsation. On the other hand, the periods detected for the *WDs with close pairs* and the *Potential WDs with pairs* can represent rotational

periods, orbital periods, or even both in case of synchronization.

Before we present our results, Figures 2 and 3 present examples of time-series analysis for two WDs for which we have detected a variability period. For both figures, the top panel shows the FT with a red dashed line indicating the false alarm probability $FAP=1/1000$ and




Figure 2. Time series analysis for the star TIC 1507548590. The top panel shows the Fourier Transform as well as its 1/1000 false alarm probability (red dashed line). The bottom panel shows the light curve folded on the period of 1.92772 hours and a sinusoidal curve with the same period in red.

the bottom panel shows the folded light curve for the highest period detected in the FT.

Figure 2 shows the Gaia WD candidate TIC 1507548590 (SDSS J175909.01 + 231155.7) that we classified as *Potential WD with pair*. It shows $T_{\text{eff}} = 25089 \pm 2950$ K and $\log g = 6.404 \pm 0.220$ cm s^{-2} (Gentile Fusillo et al. 2021). This is a low mass ($0.214 M_{\odot}$) and faint WD, with magnitude $G = 17.505$ - our selected faintness limit. The FT presents two significant peaks: a frequency corresponding to 1.92772 hours and its harmonic. The folded light curve confirms that this is the period corresponding to the main variability of the star.

Figure 3 shows the DA TIC 1100582330 (SDSS J154119.84 + 120914.6). Its spectra fit $T_{\text{eff}} = 26828 \pm 107$ K and $\log g = 7.60 \pm 0.02$ cm s^{-2} (Kepler et al. 2015), which corresponds to a mass of $0.496 M_{\odot}$; it is also faint, at magnitude $G = 17.032$. The FT presents only one significant peak, with a period of 2.736 hours. This is one of the few stars in our sample of *Likely single WDs* for which it is possible to observe the variability in the folded light curve.

3.1. Mass and Temperature Distribution




Figure 3. Time series analysis for the star TIC 1100582330. The top panel shows its Fourier Transform as well as its 1/1000 false alarm probability (red dashed line). The bottom panel shows the light curve folded on the period of 2.736 hours and a sinusoidal curve with the same period in red.

Figures 5 (4) show the dependence of the detected period — on a logarithmic scale — on the stellar mass (effective temperature) in the left panel, and the mass (effective temperature) distribution in the right panel. The red points and bins denote magnetic WDs, whereas the blue points and bins correspond to WDs with close pairs. It is important to point out that the blue and red bins are independent distributions; their overlap does not correspond to a sample of WDs with both close pair and magnetic field detected.

Figure 4 shows that our sample of WDs is distributed around approximately 12,000 K, a representative value of the DA population, which comprises about 80% of the WDs spectroscopically identified (e.g. Kepler et al. 2021). The left panel of Figure 4 shows that there does not appear to be a dependence of the detected periods on the effective temperature. Moreover, the close binaries exhibit a mean temperature of 26 300 K, similar to the *Complete sample*, whose mean temperature is 27 700 K. On the other hand, magnetic WDs present a trend towards low temperatures, with a mean effective temperature of 17 100 K.

The left panel of Figure 5 shows that there is no strong dependence of the detected periods on the stellar mass for the *Complete Sample*. However, this plot reveals




Figure 4. The left panel displays the rotational period in hours versus the effective temperature in Kelvin, both in log scale. The right panel shows the temperature distribution of sample 1. The red and blue dots/bins represent the WDs with magnetic field and with close pairs detected

, respectively.




Figure 5. The left panel displays the rotational period in hours in log scale versus mass in solar masses, while the right panel shows the mass distribution of sample 1. The red and blue dots/bins represent the WDs with magnetic field and with close pairs detected

, respectively. The black circles indicate the stars that composes the *Likely single WDs*.

a negative correlation for the *Likely single WDs*, suggesting a trend to shorter rotational periods for more massive WDs. Moreover, we can also observe a positive correlation between period and mass for the *WDs with close pairs*.

The left panel of Figure 5 also shows that most magnetic WDs in our sample exhibit periods shorter than 10 h and that they are among the targets with the shortest periods. It is essential to recognize that our sample has a selection effect. According to Brinkworth et al. (2013), the rotational period of magnetic white dwarfs

exhibits a bimodal distribution, with one group showing periods on the order of minutes to hours and the other displaying periods much longer than their 4 years of observations. However, their total sample consists of only 26 stars, which is smaller than our sample of 30 magnetic WDs. Each observation cycle (sector) of the TESS telescope lasts ≈ 27 days, with an interruption of data transfer in the middle, limiting the detection of periods to a few hundred hours. In this way, we detect only the fastest portion of magnetic WDs. Furthermore, massive WDs are fainter and tend to rotate more rapidly (e.g.,

Hermes et al. 2017b; Caiazzo et al. 2021; Williams et al. 2022). Although TESS is a small telescope, in fact, most of our magnetic sample consists of massive white dwarfs rotating with periods shorter than 10 h.

The histogram in Figure 5 shows that the mass distribution exhibits two peaks, one sharp peak centered on $0.2 M_{\odot}$ and a broader distribution around approximately $0.55 M_{\odot}$. The distribution around $0.55 M_{\odot}$ is consistent with the values expected for the mass distribution of WDs (e.g. O’Brien et al. 2023). On the other hand, the distribution around $0.2 M_{\odot}$, that contains 16% of the sample, cannot be explained by the evolution of single stars.

We included in this work all WDs that the TESS telescope was able to observe with sufficient signal-to-noise ratio — mostly WDs with apparent magnitudes lower than approximately 17.5. Therefore, our sample is limited by magnitude rather than volume. As the mass of WDs is inversely proportional to their radius and the brightness of a star is proportional to its radius, the brighter the WD the less massive it is. This leads to magnitude-limited samples of WDs being biased towards low-mass stars.

To overcome this bias, we corrected the mass distribution of our sample using the method of Schmidt (1968) (see e.g. Liebert et al. 2005; Kepler et al. 2007). In this method, the contribution of each star is weighted by $1/V_{max}$, where V_{max} is the volume defined by the maximum distance at which the star would still be visible. To correct for the nonuniform distribution of stars in our Galaxy, we assumed a Galactic disk scale height of 250 pc, following Fleming et al. (1986) and Liebert et al. (2005). We applied this method using parallax, apparent magnitude, and galactic latitude from Gaia DR3 (Gaia Collaboration et al. 2018). Gaia parallax was not available for only one star, which was not included in Figure 6.

Figure 6 exhibits the mass distribution of our sample corrected by the $1/V_{max}$ volume. It shows that the distribution of low-mass WDs around $0.2 M_{\odot}$ has disappeared, confirming that the excess of brighter low-mass WDs in our sample is an observational bias. Furthermore, the distribution also indicates a lack of WDs with approximately $0.6 M_{\odot}$, which was supposed to be the most common mass for WD stars. Instead, there is an excess of massive WDs in our sample. Furthermore, it shows that the density of our Complete sample is dominated by the *Likely single WDs* sample and that the magnetic WDs of our sample dominate the density of the massive WDs. These results could indicate that our *Likely single WDs* sample is dominated by mag-




Figure 6. Mass distribution for our *Complete sample* corrected by the $1/V_{max}$ volume. The red and blue bins represent the WDs with magnetic field and with close pairs, respectively. The hatch-filled bins indicated the *Likely single WDs*.

netic WDs resulting from binary interactions (e.g. Toonen et al. 2017; Kilic et al. 2023).

It is relevant to point out that our initial sample of 845 WDs does not present this lack of $0.6 M_{\odot}$ stars (see Appendix B - Initial Sample of 845 WDs). Instead, it exhibits a mass distribution corrected by $1/V_{max}$ as expected for WDs: centered around $0.6 M_{\odot}$ (Kepler et al. 2021; O’Brien et al. 2023, e.g.). Therefore, this gap only appears after the high-quality TESS_LOCALIZE selection.

As already explained in Introduction, some inhomogeneity at the stellar surface of single WDs must be present for us to be able to detect their rotational rate through photometry. The most common way of generating inhomogeneities is by magnetic field (Mathys et al. 2024, e.g.). The origin of the WDs magnetic field is still an open question; however, it is well-known that interactions can result in WDs with strong magnetic fields (Tout et al. 2008). One possibility is that TESS_LOCALIZE high-quality parameters may have selected the targets with higher amplitude variability and, consequently, those with higher magnetic fields. One piece of evidence is that TESS_LOCALIZE criteria reduced our initial sample by 62%; however, it reduced only by 11.8% our initial sample of 34 known magnetic WDs.

3.2. Variability Period Distributions

Figure 7 shows the histograms of periods up to 100 h (right) and up to 30 h (left) for each sample listed in Table 1. Each line of this figure displays the histograms of one sample, indicated by their legends. The last line




Figure 7. Variability period histograms for different samples of this work over and, last line, histograms of rotation periods from seismology. On the left, we present the histograms of periods up to 30 h, and on the right, the histograms of periods up to 100 h. The red bins show the magnetic population in our samples of WDs. The labels indicate the samples plotted as well as their number.

of Figure 7 shows the same histograms but for the ro-

tational periods determined by asteroseismology, which

are listed in Table 2. The red bins represent the magnetic WDs in each sample.

In the first line of Figure 7 we observe the distribution of our *Complete Sample* with its population of 30 magnetic WDs indicated by red bins. The right panel shows that the variability periods are mostly concentrated in the shortest bins. Even considering only periods shorter than 30 *h* (left panel) the two shortest bins encompass 36% of our sample. Moreover, this plot shows that our sample of magnetic WDs is concentrated in the shortest periods, presenting a median period of 3.4 hours.

The second line of Figure 7 shows the period histograms for the sample of *WDs with close pairs*. This sample is composed of 54 stars, 1 of them with a detected magnetic field. The periods detected for this sample have a median of 8.1 hours. The third line of Figure 3.4 illustrates the period distributions for the sample of *Potential WDs with pairs*. This sample is composed of 149 WDs, including 4 with magnetic field detected. Their median period is 7.6 h, close to the median period of the *WDs with close pairs*.

Finally, the fourth line of Figure 3.4 shows the period distribution of the *Likely single WDs* sample. This sample is composed of 115 stars, 25 of them with magnetic field detected. The right panel shows that the photometric rotation periods are very concentrated in the shortest period bin, 69 % of periods are shorter than 10 hours. The *Likely single WDs* exhibit a median period of 3.9 *h*, very close to the median period exhibited by our complete magnetic sample (3.4 *h*). The 25 magnetic WDs included in the *Likely single WDs* sample present an even shorter median period of 1.9 hours.

The last line of Figure 3.4 displays the distribution of the rotational periods estimated through asteroseismology, compiled in Table 2. Despite the difference in sample sizes, this figure shows significant disparities between the rotational distributions derived from photometry and seismology. The seismological sample has a median rotational period of 24.3 *h*, which is over 6 times longer than the photometric rotational median period. Additionally, the distribution of rotational periods derived from seismology is broader compared to photometry, with the seismological and the photometric samples exhibiting a median absolute deviation of 13.9 *h* and 3.5 *h*, respectively.

It is important to acknowledge that the rotation period calculated through asteroseismological models is an estimate of the mean internal rotational period of the star; while the photometric rotational period represents the rotation of the stellar surface. Moreover, the seismological sample and the *Likely single WDs* sample are representative of different populations. The seismologi-

cal sample is representative of the pulsating WDs, whose instability strips are very narrow in temperature range. In contrast, our *Likely single WDs* is representative of the WDs with surface inhomogeneities, probably mostly caused by magnetic fields.




Figure 8. Rotational period distribution up to 30 h for the *Likely single WDs* excluding periods shorter than 0.3 h. The red bins indicate the magnetic stars of the sample.

As explained in Section 1, we excluded pulsation periods. However, to eliminate any doubts about the short periods in our *Likely single WDs* sample, Figure 8 shows the histogram of periods longer than 0.3 h. It is important to note that previous work has already detected periods shorter than 0.3 h for massive white dwarfs (e.g. Caiazzo et al. 2021; Barstow et al. 1995). The inset plot confirms that even with a conservative approach of excluding periods shorter than 0.3 h, the majority of our sample still exhibits rotation periods shorter than 10 h.

3.3. Comparison with Ground-based observations

Moss et al. (2023) obtained time-resolved spectroscopy for two magnetic WDs in our sample at the Gemini Observatory. Based on the shifting positions of the Zeeman-split H_α components, they inferred a rotation period of 0.648 h for the star TIC 392797216 (LHS 2273). This star is included in our sample of *Likely single WDs*, and the period reported is very close to the 0.68 h period we found in the TESS data. Using the same method, they found that the star TIC 262548040 (LHS 1243) has a rotation period of 0.216 h. Due to its low mass of 0.41 M_\odot , we classified this star as *Possible WD with pair*. Figure 9 presents the FT of the data obtained by TESS for this star, showing that only one peak at 2617 μHz (0.106 h) is above the detection limit (horizontal dashed red line). The vertical dotted lines indicate the peak of 0.106 h and its double (0.212 h), showing

that there is no significant peak at 0.212 h in the TESS data. Therefore, it shows that, in this case, the signal we detect in the TESS data is the harmonic of the real rotation period. Despite the harmonic issue, [Moss et al. \(2023\)](#) confirms the rotation rate for two of the fastest rotators in our sample.




Figure 9. Fourier Transform of TESS data of star TIC 262548040. The horizontal red line indicates the false alarm probability $FAP=1/1000$ and the vertical dotted lines indicate the peak of 0.106 h and its double.

3.4. Pulsating White Dwarfs

From the 63 stars listed in Table 2, 45 were observed by TESS. We analyzed all of them, searching for non-pulsating periods. Most pulsating WDs do not show any frequency with amplitude above the detection limit, other than pulsation periods, or show only low amplitude at frequencies smaller than $4.6 \mu\text{Hz}$ (10 times the frequency resolution of a sector) in the FT. Some pulsating WDs exhibit dozens of pulsation frequencies in their FTs, leading to linear combinations at low frequencies. We only identified reliable variability periods outside the pulsation range (periods longer than 1500 s) for three of these stars: NGC 1501, TIC 7675859, and G226-29. Their TESS_LOCALIZE output parameters are compiled in Table 3 and are described in the following subsections: 3.4.1, 3.4.2, and 3.4.3.

3.4.1. NGC 1501




Figure 10. Fourier transform of NGC 1501 data at low frequencies.

The pulsating pre-white dwarf NGC 1501 was discovered to pulsate with periods from 5235 s to 1154 s, by [Bond et al. \(1996b\)](#). They estimate a rotation period of 1.17 days from a mean splitting of $9.9 \mu\text{Hz}$. This star was observed by TESS during sector 19 for 25 days with 120 s exposures. We analyze its FT and detect three variability frequencies outside its pulsating frequency range. TESS_LOCALIZE confirmed that the source of these signals is NGC 1501. Figure 10 shows this FT up to $200 \mu\text{Hz}$, where the three non-pulsating frequencies that we have detected are indicated by the vertical dotted lines and correspond to 4.11, 39, and 138 hours. None of these periods is consistent with the rotation period estimated by [Bond et al. \(1996b\)](#). Moreover, NGC 1501 is a planetary nebula covered by an ionized gas disk. Therefore, considering the complexity of this system, we are unable to interpret these periods.

3.4.2. TIC 7675859




Figure 11. Fourier transform of TIC 7675859 data from sectors 52 to 54.

[Romero et al. \(2024\)](#) determined the asteroseismological rotation period of 7.15 h for the DAV TIC 7675859 based on data from sectors 25, 26, 40, 52, 53, and 54. We observed the three components of a high-amplitude triplet in the data of sectors 52 to 54 (see Figure 11). The triplet at $1738 \mu\text{Hz}$ is perfectly symmetric with a frequency spacing $\Delta\nu = 22.47 \mu\text{Hz}$. Moreover, we also observed a significant peak at $22.44 \mu\text{Hz}$ (corresponding to 12.3 h), which is a very close value to the mean frequency spacing of this triplet. The peak at $22.44 \mu\text{Hz}$ is higher than all components of the triplet at $1738 \mu\text{Hz}$, however, we cannot eliminate the possibility that it is a linear combination ([Kurtz et al. 2015](#)). This peak of 12.3 h can also be a subharmonic of the real rotation rate or even the real rotation rate of the stellar surface itself, indicating differential rotation.

3.4.3. G226-29

G226-29 is the brightest known DAV, with a known pulsation triplet around 109 s ([Kepler et al. 1983](#)).

Name	TIC	FT Period [h]	Amplitude [σ]	Like	\langle Height [σ] \rangle	Q/S
NGC 1501	84306468	138 ± 32	15.377	1.0	7.46	1/1
		39 ± 2	18.104	1.0	18.4	1/1
		4.11 ± 0.03	14.571	1.0	15.4	1/1
TIC 7675859	7675859	12.344 ± 0.008	8.768	1.0	8.1	5/6
G226-29	199666369	0.93 - 3.25	4.129 - 13.934	1.0	9.4	27/29

Table 3. Non-pulsating periods from pulsating white dwarfs.




Figure 12. Running Fourier transform of G226-29 20 s data from sector 51. The tall peak on the right is the pulsation triplet around 109 s, while the region around 200 μHz corresponds to the peaks around 4500 s.

When studying this triplet, Kepler et al. (1995) estimated the rotation period of the WD as 8.9 h. G226-29 was observed by TESS with 120 s exposures from sector 14 to 26, and with 20 s exposures from sectors 40 to 41 and 47 to 60. In total, it was observed along 29 sectors. Data from all sectors show a $S/N \geq 4.1$ wandering peak from 0.93 h (297.3 μHz) to 3.25 h (85.5 μHz). As an example, Figure 12 plots the Running Fourier Transform of sector 51 data. This figure shows the known stable triplet around $\sim 9200 \mu\text{Hz}$ and a peak that wanders from $\sim 145 \mu\text{Hz}$ to $\sim 168 \mu\text{Hz}$. Figure 13 shows the complete FT of all 20 s data combined, where the wandering peak appears as a broad band at low frequencies (see the inset plot).

The methodology described in Section 1.1 assumes that the variability signals are stable. Therefore, to verify whether this wandering peak comes from G226-29 we had to apply the TESS_LOCALIZE test differently. For each sector, we obtain the highest peak exhibited in such a FT range and run the TESS_LOCALIZE code for this period. As detailed in Table 3, TESS_LOCALIZE confirmed that this wandering peak indeed comes from G226-29; however, this variability remains a mystery.




Figure 13. Fourier transform of the 20 s data for G226-29 from sectors 40 to 41 and 47 to 60. The inset plot zooms the FT at low frequencies. The red dashed line indicates the false alarm probability. The large peak on the right is the pulsation triplet at 109.47235s@1.81mma, 109.27929s@0.57mma, 109.08690s@1.62mma, while the region around 200 μHz corresponds to the wandering peak around 4500 s.

The rotational period from seismology is indicated by a vertical black dashed line in Figure 13, showing the signal we are detecting is not consistent with the rotation of the stellar inner. On the other hand, this signal is very long to be consistent with pulsations. This leads us to suspect differential rotation; that is, the wandering peak could be a manifestation of rotation of the stellar surface, that is rotating faster than the inner. Although the star has an upper limit on the magnetic field around 10 kG (Valyavin et al. 2006), subsurface magnetic fields, like those in Section 4.2, are a possibility. Only radial differential rotation does not explain why this peak varies so much, but an angular differential rotation, like the sun (see e.g. Snodgrass & Ulrich 1990; Beck 2000), could.

4. EVOLUTIONARY MODELS

If the rotation periods we detect for the *Likely single WDs* sample are due to surface inhomogeneities caused by magnetic fields, we need to calculate models that generate magnetic fields and reproduce the order of magnitude of the rotation rates we are observing. Bagnulo & Landstreet (2020); Amorim et al. (2023) show that at least 20% of all white dwarfs are magnetic, and Bag-

nulo & Landstreet (2022); Castro-Tapia et al. (2024b,a); Blatman & Ginzburg (2024a,b) show dynamo induced by phase separation during crystalization cannot be the origin for most magnetic white dwarfs. When we study the rotation rates of evolved white dwarf stars, one question that arises is how the evolution changes the internal velocities and how much of the initial angular momentum is lost during the expressive mass loss phase of AGB before the white dwarf phase. The models included in this section have internal magnetic fields generated by dynamos caused by the rotation of the star and angular momentum transfer. We calculate their rotation rates, internal and at the photosphere, to compare with the observed ones and observe how they change — or not — with the effective temperature of the star.

4.1. *Observational and modeling context*

Using ground-based photometric observation time series from mean sequence stars spanning late-F to early-M spectral types, Fritzewski et al. (2021) determined 279 rotation periods. They found that the rotation periods range from 0.5 d to 32 d, with trends of slow-rotating FGK stars and fast-rotating K dwarfs. In another recent investigation, Labadie-Bartz et al. (2023) explored the rotational characteristics of chemically peculiar (CP) stars using TESS data. Among their findings, they found that the rotation period distribution of CP stars is centered around 3 days. CP stars are observed within the spectral types from early B to early F. Since all stars with spectral type roughly from A to K will become WDs, both works evidence the rotational period of WD’s progenitors is of the order of days. We have shown in this work that the median photometric rotation period of WDs is 3.9 h; therefore, these findings suggest that a robust mechanism of internal angular momentum transfer operates during the evolution of these stars.

The asteroseismology of low and intermediate mass stars enabled the measurement of the internal rotation rate of WD progenitors at different evolutionary phases (e.g., Mosser et al. 2012; Deheuvels et al. 2012, 2014; Mosser et al. 2012; Deheuvels et al. 2015; Di Mauro et al. 2016, 2018; Gehan et al. 2018; Tayar et al. 2019; Deheuvels et al. 2020). These inferred rotation rates have been used to constrain models of angular momentum transfer processes. Eggenberger et al. (2019b) and Moyano et al. (2022) found that the transport of angular momentum must be more efficient in more massive stars during the subgiant and red giant phase, even if the efficiency of the internal transport of angular momentum decreases with the evolution of the star. They suggest that the physical nature of the additional mechanism

may be different in subgiant and red giant stars. Eggenberger et al. (2022) find that the core rotation rates of the red giant branch models are nearly insensitive to the initial rotation velocity.

On theoretical grounds, Fuller et al. (2019) deduced a new expression for the Tayler instability that leads to larger magnetic field amplitudes, more efficient angular momentum transport, and smaller shear than predicted by the original Tayler–Spruit dynamo. Eggenberger et al. (2019a) applied the new prescription for subgiant and red giant models and found that it leads to low core rotation rates after the main sequence that are in better global agreement with asteroseismic measurements than those predicted by the original Tayler–Spruit dynamo. However, it fails simultaneously to reproduce the asteroseismic measurements available for subgiant and red giant stars.

den Hartogh et al. (2019) found that the rotational period of the models with additional viscosity is too large in the white dwarf phase. However, the rotational periods match white dwarf rotation periods of the order of 1 day when they exclude the additional effect during the core-helium-burning phase. Therefore, they proposed that the efficiency of the unknown angular-momentum transfer process must decrease during the core-helium burning phase. Applying the Fuller et al. (2019) formalism, den Hartogh et al. (2020) found that, when the dynamo effect is turned off at the end of the core helium burning phase, the WD rotation rates decrease to the order of days. They concluded that the Fuller et al. (2019) dynamo formalism cannot be the sole solution to the missing process of angular momentum transport in intermediate-mass stars.

Magnetic fields are present in low-mass stars and, when strong enough, can change the rotation by exerting a torque capable of reducing the differential rotation or even imposing uniform rotation throughout the star. Moyano et al. (2023) showed that the transport of angular momentum in radiative zones during the main sequence of low-mass stars must be efficient. They suggested that the internal magnetic fields are a strong candidate for the missing physical ingredient in stellar interiors. In addition, they point out that at least one efficient process able to neutralize the development of differential rotation in stellar interiors must act during the whole evolution of these stars.

4.2. *Our models*

We computed evolutionary sequences using the Modules for Experiments in Stellar Astrophysics code (MESA, Paxton et al. 2011, 2013, 2015, 2018, 2019; Jermyn et al. 2023), release 22.11.1. Our models have

an initial mass of $M_i = 1.5 M_\odot$ (following den Hartogh et al. 2019) and an initial metallicity of $Z = 0.02, 0.002$ and 0.001 . The initial mass was chosen to be above the lower limit of $1.3 M_\odot$ for the convective core in the main sequence (e.g. Aerts & Tkachenko 2023), leading to a WD with a mass close to the mean mass of most white dwarf stars (e.g. Kepler et al. 2007; O’Brien et al. 2023). The models are computed from the zero-age main sequence (ZAMS) until a white dwarf cooling effective temperature of 8500 K. Our models start with equatorial surface rotation velocities of 10 and 50 km s⁻¹, typical of main-sequence stars with 1 and 2 solar masses. These values are assumed to be the solid-body rotations at the ZAMS, which is a reasonable and common assumption for this type of star (see, e.g., Bouvier et al. 1997; Granada & Haemmerlé 2014; Amard et al. 2019; Deal et al. 2020; Nguyen et al. 2022; Douglas et al. 2024). For further information on the models, see Appendix C - Evolutionary Models Input.

The mixing of elements and the transport of angular momentum due to rotation are implemented in MESA following closely Heger et al. (2000, 2005). Two efficiency factors must be set to calibrate the diffusion coefficients: the contribution of rotationally induced instabilities to the diffusion coefficient is reduced by the factor $f_c = 1/30$, and the sensitivity of rotationally induced mixing is $f_\mu = 0.05$.

We consider the following angular momentum transport mechanisms: dynamical shear instability, Solberg-Hoiland, secular shear instability, Eddington-Sweet circulation, Goldreich-Schubert-Fricke and the Tayler-Spruit dynamo.

The final masses of our models are $0.546 M_\odot$ for $Z = 0.02$, $0.577 M_\odot$ for $Z = 0.002$, and $0.585 M_\odot$ for $Z = 0.001$. This aligns with the expectations that stars with higher metallicity should form less massive CO cores and lose more mass in the AGB phase than stars with the same initial mass but lower metallicity (e.g., Dominguez et al. 1999; Catalán et al. 2008; Romero et al. 2015; Choi et al. 2016). Similar models with initial masses of $2.5 M_\odot$ reach the white dwarf phase with masses around $0.6 M_\odot$.

In Figure 14 we show the evolution of the surface rotational period (P) versus the effective temperature (T_{eff}) during the cooling track for a set of model sequences with different metallicity and initial rotation velocity. The surface rotational period is computed via $P = 2\pi r/v$, where r is the radius of the model and v is the rotational velocity at the equator. Sequences that consider the six mechanisms of angular momentum transport are shown as dotted lines (AMT = ON),




Figure 14. Observational data of the surface rotation of white dwarf stars (orange crosses for the complete sample and black dots for the likely single

) are compared against evolutionary models showing the rotation period in the white dwarf cooling track. We compare models with the six mechanisms of angular momentum transport turned off (solid lines) and on (dotted lines) during the whole evolution. For an initial rotational velocity of $v_i = 10 \text{ km s}^{-1}$, we present models with metallicities of $Z = 0.02$ (purple), 0.002 (dark green) and 0.001 (light blue). We also compare two models where the Tayler-Spruit dynamo is the only mechanism that is turned off during the computations (dark blue and light green dotted-dashed lines). For an initial rotational velocity of $v_i = 50 \text{ km s}^{-1}$, we compare two models with $Z = 0.02$ (red lines). All model sequences have an initial mass of $1.5 M_\odot$.

and solid lines represent sequences with all mechanisms turned off (AMT=OFF).

For $v_i = 10 \text{ km s}^{-1}$ and $Z = 0.02$ (purple line in Figure 14), we find that the white dwarf models with AMT = OFF decrease the average equatorial rotation period on the surface from $P = 300 \text{ h}$ to $\sim 30 \text{ h}$ when cooling from $\sim 17 \text{ kK}$ to 8500 K. The sequence with AMT=ON does not even reach 300 h before cooling down to the same temperature (and therefore does not appear in the plot).

For sequences with $Z = 0.002$ (dark green lines in Figure 14), we find that if AMT=OFF (solid line), the models brake from $P = 300 \text{ h}$ to 40 h only at the very end of the cooling track ($T_{\text{eff}} < 10 \text{ kK}$), while the AMT=ON sequence (dotted line) has already spun down to 300 h when $T_{\text{eff}} < 17 \text{ kK}$, although it reaches only $P \sim 60 \text{ h}$ at $T_{\text{eff}} = 8500 \text{ K}$.

The internal redistribution of the angular momentum is very efficient early in the evolution of low-mass stars (e.g., Mosser et al. 2012). This suggests that when AMT=ON, a large amount of mass lost by winds during the AGB phase carries away most of the angular

momentum, and thus the surfaces of the models rotate slower in the WD phase.

The difference in turning the angular momentum transport mechanisms on and off is much more pronounced for low metallicity. ($Z = 0.001$, light blue in Figure 14). While AMT=OFF (solid line) follows the same trend as for higher metallicities (that is, it only reaches low P at the end of the cooling track), the sequence with AMT=ON (dotted line) cools from $T_{\text{eff}} = 40$ kK down to 8500 K with approximately constant rotation period of $P = 25$ h, closer to most of the observed period distribution.

In Figure 14 we also show two sequences of models with $v_i = 10 \text{ km s}^{-1}$ where the Tayler-Spruit mechanism was turned off, and all the other five mechanisms were kept on throughout evolution (ST = OFF, dotted-dashed lines). For $Z = 0.001$ (light green line), the result is similar to that of the same metallicity, but with all six angular momentum transport mechanisms turned off. This suggests that the Tayler-Spruit mechanism dominates over the other mechanisms from the point of view of rotation analysis during the WD phase.

Finally, we present two sequences with $v_i = 50 \text{ km s}^{-1}$ and $Z = 0.02$. These models show that observational data with a shorter rotation period ($P < 50$ h) are better adjusted by models with AMT=OFF, while models with AMT=ON present a rotation period shorter than 300 h only at the end of white dwarf cooling when $T_{\text{eff}} < 10$ kK. This suggests that if we disregard the Tayler-Spruit mechanism throughout the evolution, models with a higher initial velocity (50 km s^{-1}) fit the white dwarf rotation data better if the initial mass is $1.5 M_{\odot}$.

The dominant factor for the spin-up of the external layers during the white dwarf cooling phases are the He-convection zones that arise at $T_{\text{eff}} \simeq 32000$ K, followed by the H-convection zone if the star has a H envelope, around $T_{\text{eff}} \simeq 14000$ K. The movement of mass in the convection zones efficiently redistributes the angular momentum.

Also, the models with AMT=ON generate internal magnetic fields due to differential rotation, as shown in Figure 15 for a model sequence with $M_i = 1.5 M_{\odot}$, $Z = 0.001$, and $v_i = 10 \text{ km s}^{-1}$ (corresponding to the dotted light blue line in Figure 14). According to Spruit (2002), the Maxwell stresses transport angular momentum in the radial direction at a rate proportional to the product $B_r B_{\phi}$. In Figure 15, the x axis represents the mass coordinate of the model, with the core on the left and the surface on the right, and the y axis shows the intensity of the magnetic field components B_r and B_{ϕ} in the internal part of the models at two different temperatures during




Figure 15. Internal magnetic field at $T_{\text{eff}} = 25000$ K and 11000 K, generated by the Tayler-Spruit dynamo during the evolution due to differential rotation caused by angular momentum transfer throughout the stellar evolution. B_r is the radial field and B_{ϕ} the azimuthal one. The properties of the initial model in the ZAMS are $M_i = 1.5 M_{\odot}$, $Z = 0.001$, and $v_i = 10 \text{ km s}^{-1}$. For $T_{\text{eff}} = 25000$ K, the solid red line shows the radial component and the dotted blue line represents the azimuthal one. For $T_{\text{eff}} = 11000$ K, the dashed green line shows the azimuthal component, and the radial component does not have enough intensity to appear in the plot. The six angular momentum transfer mechanisms operate throughout evolution. Magnetic field values are negligible beyond 1.3 on the x -axis, i.e., towards the surface.

the cooling track. The chosen temperatures, 25000 K and 11000 K, are representative of the DAV and DBV variable classes.

Figure 15 shows that, during cooling, the azimuthal component of the magnetic field is always greater than the radial component. In some regions, the magnetic field reaches an intensity of $\sim 10^4$ G. We also found that both components tend to be less intense as the models cool, and the outer $\sim 5\%$ (by mass) of the models show no significant magnetic fields.

As a general behavior, the models suggest the following. During cooling, the Tayler-Spruit dynamo mechanism acts smoothly in the core, flattening the rotation profile up to the He layer. On the other hand, a convective zone appears at the base of the He envelope around 32000 K and increases in size inward as the models cool, causing angular momentum to be transferred from the core and the He zone.

Around 14000 K a convective zone develops at the base of the H envelope, and then the rotational speed on the surface is as high as that of the fastest part of the core and the He zone. The exact temperature for this to occur depends on the metallicity and, especially, on the history of mass loss at the end of the AGB phase. This

is because the last thermal pulses directly interfere with the amount of H remaining in the model that enters the cooling track, modifying the convection in this zone.

Thus, the internal rotation profile of the models during cooling depends, to some extent, on angular momentum transfer mechanisms, but also strongly depends on convection in the He and H zones.

5. CONCLUSION

We have reported significant variability periods for 318 white dwarfs, as well as their variability amplitude and quality criteria. We divided the 318 stars composing the *Complete Sample* into three sub-samples. 54 of them were classified as *WDs with close pairs*, and their variability periods can be due to orbital motion or rotation. The other 149 were classified as *Potential WDs with pairs* due to their low masses and/or high temperatures (e.g. Steen et al. 2024). Finally, we classified 115 as *Likely single WDs* and therefore attributed their variabilities to rotation.

Our sample of 30 magnetic WDs are massive, fast rotators, and cold; while our sample of 54 *WDs with close pairs* are well distributed on the mass range and presents a weak tendency of longer periods to massive WDs.

Our *Complete Sample* of 318 WDs exhibits a temperature distribution expected for DA WDs, peaking around $T_{\text{eff}} \simeq 12,000$ K. On the other hand, its mass distribution is biased towards low-mass stars because it is a magnitude-limited sample. The mass distribution corrected by $1/V_{\text{max}}$ presents, in density, a lack of WDs with a mass around $0.6 M_{\odot}$ and an excess of massive WDs, which are dominated by our little sample of 30 magnetic WDs. These peculiarities are not present in the corrected mass distribution of our initial sample of 845 stars, which leads us to conclude that these characteristics were introduced by the TESS_LOCALIZE high-quality selection. In this second selection, 79 % of the stars discarded by TESS_LOCALIZE was done because the software was unable to perform a high-quality fit. Therefore, it is possible that TESS_LOCALIZE have selected the highest amplitude variabilities and, consequently, the WDs with the strongest magnetic field. It could lead to a mass distribution dominated by magnetic WDs resulting from binary interactions and possibly explain the lack of $0.6 M_{\odot}$ WD density.

The stellar density of our sample is completely dominated by the *Likely single WDs*, which, as already discussed, is probably composed of magnetic WDs. A piece of evidence is that TESS_LOCALIZE reduced our initial sample by 62%; however, it reduced only by 11.8% our initial sample of 34 known magnetic WDs.

The *Likely single WDs* present a median rotational period of 3.9 hours that is far shorter than the median rotational period from seismology estimates, 24.3 hours. However, this is an expected result, once both estimates of rotation represent different populations. The seismological sample represents the mean internal rotation of pulsating WDs, while our *Likely single WDs* represents the rotation of the stellar surface of WDs with inhomogeneities at their surface.

From the 63 WDs with rotation period determined by asteroseismology, we detected nonpulsating periods in the TESS data for 3 of them: NGC 1501, TIC 7675859, and G 226-29. We were not able to interpret with certainty any of these periods. This failure in finding photometric rotational periods for the seismological sample using TESS data reinforces that the seismological and the *Likely single WDs* samples represent distinct populations. Up to the moment, we do not know any pulsating WD with magnetic field confirmed.

Using ground-based observations, Moss et al. (2023) confirmed the rotation period we detected in the TESS data for two of the fastest rotators in our sample; however, they also provided an example of half-period aliases. It attempts to the fact that we may be detecting and reporting half of the real rotation period for other stars. On the other hand, this work confirmed rotation periods of 0.68 h and 0.21 h for two magnetic WDs in our sample; therefore, it does not invalidate our findings.

For 5 of the 318 WDs in our sample, all of them included in our sample of *Likely single WDs*, the variabilities detected in the TESS data were also reported by Farihi et al. (2023). They are TIC 328029653, TIC 251080865, TIC 251903434, TIC 204440456 and TIC 321979116. The periods reported agree with our findings. They claim that the variability found in the WD 2138-332 data (TIC 204440456) is the stellar rotation period, almost certainly due to magnetism. In addition, the orbital period of the EMLV TIC 308292831 was already reported by Lopez et al. (2021).

Figure 14 presents two models that at 40 kK reach rotation periods consistent with the observational findings of this work. We find that periods shorter than 50 h are well-fitted by models with $Z = 0.02$, $v_i = 50 \text{ km s}^{-1}$ and angular momentum transport set to OFF, or by models with $Z = 0.001$, $v_i = 10 \text{ km s}^{-1}$ and angular momentum transport set to ON. A smaller part of the observational data presents a large dispersion of the rotation period and is concentrated at $T_{\text{eff}} < 20,000$ K. Our remaining models best fit these stars.

While other studies have shown that the internal transfer of angular momentum should be intense in the early phase of the evolution of low-mass stars, our study

confirms that if we seek to match most of the white dwarf rotational data, this redistribution should cease or be less intense before the end of the AGB phase, when stars lose the highest amount of mass and, therefore, angular momentum. If we assume that stars with lower metallicity will have a lower mass loss rate due to winds at the top of the AGB phase (e.g., see discussion in Höfner & Olofsson 2018), this means that they will lose less angular momentum and therefore rotate faster in the WD phase. However, when we consider that more angular momentum transfer mechanisms have been activated since the beginning of the evolution, more angular momentum will be transferred to the surface and lost during the AGB phase. Thus, in terms of the WD rotation period, lowering the metallicity is equivalent to reducing the angular momentum transfer mechanisms in the early evolution. Due to their intrinsic faintness, the observed white dwarfs are mainly nearby galactic disk objects, most likely with metallicity closer to the Sun’s.

In general, our models agree with the most recent results in the literature in the sense that different initial parameters can be used to create models that are in agreement with a particular part of the observational data, but there is a degeneration in the parameter space (i.e., the initial velocity of rotation, initial mass, metallicity, etc.) that could be better resolved if we knew more precisely how the redistribution of angular momentum occurs in stellar interiors.

In the future, we will analyze the TESS data observed after October 2023, calculate models with the Fuller dynamo prescription, and extend the mass range of the models, as the data include white dwarfs with a range of masses. We intend to perform spectropolarimetry to check if our *Likely single WDs* are magnetic. We also intend to confirm the rotational periods using high-resolution spectroscopic observations to detect the NLTE line core rotational broadening, as in Berger et al. (2005).

We thank the anonymous referee for his valuable comments and suggestions, that improved our paper significantly. This work was carried out with the financial support of the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), and by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES). Most of the calculations were performed at the Texas Advanced Computing Center, located at the University of Texas in Austin. We thank Stéphane Charpinet for maintaining the Evolved Compact Stars with TESS on TASOC and for providing the running Fourier Transform of the star G226-29. This research has made extensive use of NASA’s Astrophysics Data System Bibliographic Service (ADS), SIMBAD, and MAST. We used ASTROPY (<https://www.astropy.org/>), TESS-LS (<https://github.com/ipelisol/TESS-LS>), TESS_LOCALIZE (<https://github.com/Higgins00/TESS-Localizer>), LIGHTKURVE (<https://docs.lightkurve.org/>) and PYRIOD (<https://github.com/keatonb/Pyriod>).

APPENDIX

APPENDIX A - KNOWN RR LYRAE STAR

BPM 24754 (TIC 367227831) is a known variable DAV white dwarf. Figure 16 shows its FT of TESS data plotted using the Pyriod software (Bell 2020). In this case, we know that the primary signal at 21.49 μHz (12.92 h) and its harmonics are the contamination from the known RR Lyrae variable *Gaia DR3 5923100101270979328* (TIC 367227833). The results of software TESS_LOCALIZE for the three sectors where the star is observed (12, 39, and 66) are presented in Table 4.

Sector	Source Gaia DR3	pvalue	Relative Likelihood	χ^2	Height [σ]
12	5923100101270979328	0.214	0.75	44.40	42.40
12	5923100101249048320	0.071	0.19	44.40	42.40
39	5923100101270979328	0.503	0.83	312.71	17.30
39	5923100101249048320	0.191	0.11	312.71	17.30
66	5923100101270979328	0.829	0.96	62.47	37.22
66	5923100101249048320	0.198	0.03	62.47	37.22

Table 4. TESS_LOCALIZE results to the signal 12.92 h in present in the light curve of the DAV BPM 24754.




Figure 16. BPM 24754 light curve from sectors 12, 39, and 66.

TESS_LOCALIZE estimates that the probability that the signal source is the RR Lyra star is 75% in sector 12, even the parameters *pvalue* and *Height* indicating good values. The *Relative Likelihood* of the RR Lyrae star being the source of the signal increases in the subsequent sectors, with the mean like reaching 85%. The second source is likely the same for all three sectors, *Gaia DR3 5923100101249048320*. This object is located $3.06''$ away from the RR Lyrae star.

APPENDIX B - INICIAL SAMPLE OF 845 WDS

Our initial sample is composed of 845 WD stars exhibiting a non-pulsating period in their photometric data from TESS that, at least for one sector, TESS_LOCALIZE indicates the WD as the most probable signal source. As explained in Section 1.1, most of our initial sample was discarded because TESS_LOCALIZE was unable to obtain a high-quality fit for the data. That means that for most of the stars in this sample we are not able to determine with high confidence whether the source of the signal we detected is, in fact, the WD. Figure 17 presents the mass distribution corrected by the $1/V_{max}$ method and the period distribution for our initial sample. The corrected mass distribution is centered around approximately $0.6 M_{\odot}$, as expected for WD stars. Magnetic WDs are more massive, exhibiting a distribution around approximately $0.8 M_{\odot}$. The period distribution is heavily concentrated in the shortest period bins, with the sample having a median period of 11.85 *h* and a median absolute deviation of 11.56 *h*.

APPENDIX C - EVOLUTIONARY MODELS INPUT

Hydrogen and helium burning are computed using the `pp_and_cno_extras.net` network that accounts for 25 isotopes and 79 net reactions. Convection is treated using the formulation of the mixing-length theory (Böhm-Vitense 1958) in the variation of Henyey et al. (1965) allowing the convective efficiency to vary with the opacity. We set the diffusion of elements for the whole net; each isotope in the network is treated as its own class.

Following the MIST project (Choi et al. 2016), $\alpha_{MLT} = 1.82$ is adopted as the mixing length parameter. We consider the Ledoux criterion of stability, which considers the influence of composition gradients on mixing. Semiconvection is considered unstable regions by the Schwarzschild criterion but stable by Ledoux, with an efficiency parameter $\alpha_{sc} = 0.1$. Thermohaline mixing is included throughout evolution, with efficiency $\alpha_{th} = 666$.

Exponential overshooting is set to $f = 0.008$ in the core and 0.0087 in the shell. We handle mixing in the convective zones using the convective premixing scheme. For the boundary conditions of the atmosphere, we use the $T(\tau)$ Eddington relation with varying opacity for most of the evolution and hydrogen atmosphere tables for cool white dwarfs (Rohrmann et al. 2011) when the models are on the cooling track and the effective temperature is below 10,000 K. The mass loss by stellar winds is taken into account using the Reimers (1975) scheme with $\eta_R = 0.5$ for the




Figure 17. Mass distribution for our *Initial* corrected by the $1/V_{max}$ volume.

red giant branch phase and the [Bloeker \(1995\)](#) scheme with η_B varying between 0.1 and 1 for the asymptotic giant branch phase.

APPENDIX D - VARIABILITIES PERIODS TABLE

Table 5. Photometric rotation Periods.

TIC	Period [h]	Amplitude [σ]	Like	\langle Height [σ] \rangle	T_{eff} [K]	Mass [M_{\odot}]	Q/S	Info
144002497	7.6213 ± 0.0022	48.042	1.00	42.8	72910	0.606	2/2	
837076117	11.468 ± 0.007	7.899	1.00	11.2	81335	0.65	1/2	DA
307982318	$1.53006 \pm 7\text{e-}05$	4.803	1.00	5.0	31880	0.67	1/3	hotDAV
88564975	8.9702 ± 0.003	173.775	1.00	92.9	80000	0.67	3/3	
136884288	0.6053 ± 0.0006	7.730	1.00	13.6	8440	0.81	1/1	DAHe
383647517	13.924 ± 0.014	13.263	1.00	6.2	6754	0.545	7/16	DQpec
1102584065	26.187 ± 0.022	17.910	1.00	9.1	110000	0.57	12/13	DO
436747174	4.166 ± 0.026	290.193	1.00	59.7	22500	0.532	1/1	DA-CV+M
434196824	3.904 ± 0.023	53.066	1.00	62.4	14491	0.258	1/1	
345265261	3.759 ± 0.022	7.601	1.00	12.2	38780	0.843	1/1	DA
321272216	4.408 ± 0.031	34.385	0.99	40.0	12597	0.117	1/1	
248861930	12.93 ± 0.26	23.353	1.00	28.0	36903	0.521	1/1	DBA
247992671	15.4 ± 0.4	8.742	1.00	10.8	8000	0.23	1/1	
241256067	15.778 ± 0.028	5.352	1.00	6.0	4730	0.495	1/2	DC
124894242	0.32555 ± 0.00017	4.075	1.00	6.9	15919	0.435	1/1	
154942231	4.4236 ± 0.0022	7.839	1.00	6.4	12000	0.15	3/5	
649932200	0.9712 ± 0.0005	31.107	1.00	28.4	26461	1.036	3/3	
52584399	3.801 ± 0.008	112.873	1.00	66.8	19505	0.249	3/3	
1883129777	1.9446 ± 0.0004	143.936	1.00	71.9	16480	0.229	3/3	
1881564044	6.137 ± 0.004	6.042	1.00	7.4	34000	0.2	2/3	DAH
1101460688	23.24 ± 0.03	10.905	1.00	6.9	88000	0.656	2/3	
22014729	7.6477 ± 0.0032	135.408	1.00	55.8	36580	0.44	1/2	
31798074	21.828 ± 0.017	42.369	1.00	5.3	12011	0.398	1/10	
262921847	13.892 ± 0.004	52.290	1.00	20.7	7000	0.23	12/12	
311886800	10.2157 ± 0.0025	32.576	1.00	19.4	100000	0.6	6/6	

TIC	Period [h]	Amplitude [σ]	Like	\langle Height [σ] \rangle	T_{eff} [K]	Mass [M_{\odot}]		Info
178994042	$0.2138 \pm 7\text{e-}05$	4.400	1.00	5.6	19770	0.57	1/1	
220478684	16.611 ± 0.010	25.143	1.00	12.6	44589	0.483	19/20	
381976323	8.722 ± 0.01	133.547	1.00	45.2	4000	0.2	11/11	
235072034	84.0 ± 4.0	20.540	1.00	10.5	3500	0.23	2/3	
219244444	7.2886 ± 0.0027	245.106	1.00	12.5	6440	0.25	3/5	DA7+M+planet
245834374	33.6 ± 1.8	7.275	1.00	6.8	34210	0.69	1/1	DA+M
85146644	5.317 ± 0.0011	68.977	1.00	51.4	60260	0.41	3/3	
107310223	11.28 ± 0.23	21.505	1.00	30.9	48116	0.366	1/1	DA1+dM
1101900758	2.65191 ± 0.0003	31.335	1.00	21.4	23941	0.398	5/5	
160119816	21.453 ± 0.017	112.794	1.00	43.3	90000	0.56	9/9	
300284179	12.65 ± 0.29	12.426	1.00	13.8	38490	0.54	1/1	
126910998	19.4 ± 0.6	36.753	1.00	54.9	23075	0.819	1/1	DA+M
218971976	6.7081 ± 0.001	263.431	1.00	65.1	39260	0.38	2/2	DA+M
321979116	2.69506 ± 0.00016	92.402	1.00	36.5	8650	0.96	2/2	DAP5.8
183533908	2.55267 ± 0.00024	11.122	1.00	13.6	7638	0.74	1/1	
630539472	1.592 ± 0.004	4.231	1.00	7.4	10027	0.821	1/1	
201255204	6.34 ± 0.03	59.895	1.00	52.0	4000	0.2	2/2	DBH
439917321	12.98 ± 0.28	61.269	1.00	48.6	19193	0.57	1/1	DA+M
55096188	2.75096 ± 0.00021	4.379	1.00	6.5	10000	0.6	2/3	
53190694	9.923 ± 0.004	12.529	1.00	9.3	30502	0.41	3/4	DA3+dM
952264940	1.78894 ± 0.00018	7.713	1.00	8.6	25692	1.151	1/2	DBH
219769847	8.72 ± 0.13	13.789	1.00	20.7	21620	0.72	1/1	DA3+dM
292114384	0.29255 ± 0.00014	4.110	1.00	5.1	18943	0.57	1/1	
66714896	12.579 ± 0.009	10.848	1.00	8.8	63073	0.69	1/3	
349233389	52.0 ± 5.0	31.595	1.00	25.7	23690	0.58	1/1	DA2+M
335501964	6.7654 ± 0.0025	5.380	1.00	8.1	90000	0.53	1/2	
165916724	$0.092325 \pm 5\text{e-}07$	4.156	1.00	5.2	9557	0.883	1/3	DAH
274039489	7.32 ± 0.09	67.842	1.00	58.5	42370	0.66	1/1	DA1.2+pair
328029653	14.157 ± 0.004	19.335	0.99	12.5	6340	0.53	5/5	
311920919	$0.1614889 \pm 1\text{e-}06$	4.611	1.00	6.2	14116	0.68	1/4	
155871645	$0.1385659 \pm 8\text{e-}07$	6.552	1.00	5.5	34050	0.77	3/5	
344927121	2.53418 ± 0.00018	113.339	1.00	67.3	26088	0.381	4/4	
148719841	8.549 ± 0.004	17.920	1.00	18.7	6169	0.72	2/2	
415714190	$0.384926 \pm 4\text{e-}06$	9.333	1.00	8.4	7690	0.636	3/3	
277713491	8.9986 ± 0.0032	11.795	1.00	9.2	7518	0.578	3/4	
471013515	6.05 ± 0.07	27.262	1.00	34.8	55000	0.6	1/1	
5393020	1.5786 ± 0.00014	54.426	1.00	49.4	8000	0.35	1/2	
610754879	$0.91146 \pm 9\text{e-}05$	11.175	1.00	10.9	47617	1.341	2/3	
365207316	2.539 ± 0.005	6.316	1.00	7.1	10520	0.226	2/2	
425079955	17.076 ± 0.016	20.821	1.00	16.2	14252	0.285	2/2	
1271382181	7.426 ± 0.005	40.083	1.00	21.8	24015	0.25	8/8	
115970839	2.6994 ± 0.0004	41.932	1.00	30.0	8825	0.219	5/5	DA+M:
159819681	10.728 ± 0.006	29.967	1.00	22.8	27876	0.573	5/5	
158971558	6.327 ± 0.004	7.300	1.00	7.3	14319	0.323	2/2	
1550807820	$0.566897 \pm 1.6\text{e-}05$	5.636	1.00	8.6	44523	1.293	5/5	
7622101	8.5708 ± 0.0028	11.742	1.00	7.6	14159	0.2	7/7	
349993999	4.381 ± 0.0005	7.642	1.00	7.0	24751	0.756	1/1	
1688534307	$0.0902002 \pm 4\text{e-}07$	4.253	1.00	5.3	66522	0.637	1/2	

TIC	Period [h]	Amplitude [σ]	Like	\langle Height [σ] \rangle	T_{eff} [K]	Mass [M_{\odot}]		Info
62918265	4.6612 ± 0.0008	12.845	0.99	13.1	8133	0.724	6/6	
1922433415	2.03387 ± 0.00015	129.416	1.00	60.4	21155	0.214	2/2	
100469915	12.936 ± 0.006	18.310	1.00	21.6	12644	0.152	2/2	
2026050241	8.0844 ± 0.0024	6.393	1.00	5.7	48831	0.489	1/2	
60112260	7.37 ± 0.1	14.436	1.00	20.0	7682	0.756	1/1	
269937578	7.2969 ± 0.0019	79.671	1.00	45.4	32755	0.274	4/5	
2052430471	$0.125404 \pm 2.9\text{e-}05$	4.277	1.00	5.4	28678	0.368	1/1	
124573902	7.9918 ± 0.0035	21.546	1.00	21.9	12173	0.371	2/2	wd+pair
270432557	9.311 ± 0.01	6.316	1.00	5.4	10000	0.23	2/3	
43529091	9.44 ± 0.14	11.479	1.00	14.0	11006	0.17	1/1	
278861557	3.1624 ± 0.0004	26.520	1.00	7.8	10460	0.23	19/20	wd+pair
62846288	$0.500906 \pm 1.4\text{e-}05$	23.831	1.00	17.9	16268	0.935	3/4	
56813164	8.05 ± 0.11	25.988	1.00	34.7	17304	0.39	1/1	DA+M
953086708	$0.088138 \pm 1.3\text{e-}05$	5.293	1.00	7.4	7735	0.564	1/1	DAEH
95475692	1.94319 ± 0.00021	45.117	1.00	36.2	13737	0.428	2/2	DA+M
68015843	3.015 ± 0.0018	21.609	1.00	21.7	12392	0.4	2/2	DA+M4.5
219485299	1.9770 ± 0.0020	4.513	1.00	18.9	10307	0.63	3/3	DA+M
1507548590	1.9272 ± 0.0004	50.124	1.00	49.2	25089	0.214	2/2	
1101327387	$1.30545 \pm 9\text{e-}05$	21.739	1.00	19.5	15659	1.13	2/2	DAH
416538823	16.021 ± 0.011	335.044	1.00	26.1	18840	0.8	4/4	DA+M
7983187	14.32 ± 0.16	77.693	1.00	45.2	17210	0.59	2/2	DA3+dM
251857373	2.31675 ± 0.00012	175.807	1.00	110.6	40698	0.56	2/2	
408015814	29.52 ± 0.05	13.089	1.00	11.8	76010	0.061	2/2	
200725303	9.48 ± 0.07	35.024	1.00	33.5	90000	0.75	2/2	
159894126	0.46415 ± 0.00017	4.410	1.00	5.9	8627	0.613	1/2	
651923454	42.4 ± 2.9	16.656	1.00	11.9	21644	0.746	1/1	wd+pair
651337335	30.5 ± 1.5	6.331	1.00	9.7	21796	0.217	1/1	
650999219	24.6 ± 0.5	24.641	1.00	26.3	70000	0.49	2/2	
93031595	9.94 ± 0.16	17.306	1.00	25.7	8751	0.911	1/1	DZ
685012668	2.8735 ± 0.0018	107.688	1.00	44.9	20041	0.238	5/5	
672383331	17.8 ± 0.5	7.107	1.00	10.9	21076	0.279	1/1	
257689590	13.35 ± 0.13	5.132	0.99	5.2	8629	0.183	2/2	
455072135	11.36 ± 0.21	90.850	1.00	49.0	12965	0.116	1/1	EB
941411508	10.547 ± 0.006	8.630	1.00	9.1	40000	0.556	2/2	
942842236	8.233 ± 0.004	10.210	1.00	10.1	77164	0.75	2/2	
332828037	2.07518 ± 0.00024	15.973	1.00	17.4	13936	0.382	2/2	
939953390	29.85 ± 0.05	9.949	1.00	6.7	63208	0.667	3/3	
952000432	21.158 ± 0.025	23.145	1.00	23.4	81500	0.58	2/2	
232524621	3.4918 ± 0.0005	14.341	1.00	8.5	100000	0.65	2/9	hotDA+pair
1003414425	26.46 ± 0.04	25.074	1.00	16.5	18089	0.173	4/4	
233177285	15.194 ± 0.01	4.790	1.00	6.2	6500	0.2	2/6	
232979174	8.8045 ± 0.0033	8.642	1.00	5.6	12776	0.473	2/9	
1001602651	4.9125 ± 0.0018	10.704	1.00	7.2	9786	0.826	5/6	
219009693	1.5828 ± 0.0004	9.973	1.00	7.3	14000	0.23	4/4	
364160098	$0.19744 \pm 6\text{e-}05$	4.307	1.00	5.7	20000	0.2	1/1	
471013541	0.549 ± 0.0005	4.071	1.00	5.6	88642	0.51	1/1	
951282189	2.9618 ± 0.0005	95.469	0.99	110.8	19099	0.174	2/2	DAB+pair
951635495	23.45 ± 0.03	7.002	1.00	6.6	61045	0.5	2/2	

TIC	Period [h]	Amplitude [σ]	Like	\langle Height [σ] \rangle	T_{eff} [K]	Mass [M_{\odot}]		Info
1924161836	25.132 ± 0.024	14.278	1.00	12.4	71080	0.482	2/2	
410414842	14.9 ± 0.4	6.417	1.00	9.7	70000	0.5	1/1	
2052284133	16.652 ± 0.03	12.980	1.00	14.3	34299	0.51	2/2	
251080865	32.01 ± 0.06	179.265	1.00	38.0	7170	0.744	2/2	DCP7
164681986	3.9437 ± 0.0009	33.642	1.00	31.3	8471	1.006	2/2	
392797216	0.6808 ± 0.0004	6.037	1.00	6.0	6814	0.749	2/2	DAH
157201137	$0.776229 \pm 3.3\text{e-}05$	10.198	1.00	10.9	11254	0.486	2/2	CV
219868627	5.9071 ± 0.0017	43.787	1.00	18.3	18000	0.18	12/12	DAZ+pair
155872634	3.9145 ± 0.0007	26.280	1.00	16.7	24589	0.697	6/6	
357389336	42.46 ± 0.08	7.007	1.00	6.1	21785	0.49	2/5	DA+K
313894558	19.05 ± 0.04	4.861	0.99	5.2	18610	0.61	1/2	
441569276	2.47353 ± 0.00023	90.759	1.00	55.4	19690	0.43	6/6	DA+M
457099062	1.36615 ± 0.0002	7.658	1.00	8.2	6655	0.555	2/2	
198510602	8.87 ± 0.013	110.918	1.00	46.3	9800	0.23	4/4	
236865474	17.789 ± 0.011	55.795	1.00	21.0	26520	0.52	14/14	DA1.9+pair
279484490	$0.086195 \pm 1.2\text{e-}05$	4.519	1.00	5.8	16989	0.387	1/1	
248353420	11.32 ± 0.26	105.385	1.00	69.0	60419	0.53	1/1	DA+K
54003343	22.864 ± 0.029	16.612	1.00	18.5	52869	0.611	1/2	DA1+dM
71513592	23.0 ± 0.9	11.500	1.00	15.8	39910	0.438	1/1	
737660462	9.703 ± 0.013	30.331	1.00	23.7	30415	0.502	5/5	
60040774	9.72 ± 0.15	9.104	1.00	14.7	8876	0.277	1/1	wd+pair
712210226	$0.115822 \pm 2.2\text{e-}05$	4.178	1.00	5.5	13707	0.53	1/1	wd+pair
705867935	0.28689 ± 0.00013	6.605	1.00	13.7	16407	1.001	1/1	
765410943	3.7702 ± 0.0005	169.012	1.00	55.6	22601	0.238	19/19	
96391732	3.662 ± 0.011	98.438	1.00	117.0	21272	0.183	2/2	
291303146	14.36 ± 0.33	43.492	1.00	40.9	100000	0.58	1/1	
770913638	$0.453289 \pm 1.1\text{e-}05$	19.139	1.00	21.2	8474	0.794	2/2	
743328948	7.9 ± 0.1	28.663	1.00	45.6	20000	0.2	1/1	
403672198	3.137 ± 0.016	25.892	1.00	38.6	7750	0.23	1/1	
437213912	37.1 ± 2.3	29.331	1.00	34.2	9000	0.2	1/1	
836046041	7.9019 ± 0.0034	12.734	1.00	15.5	26187	0.517	2/2	
817911025	$0.597463 \pm 1.9\text{e-}05$	17.981	1.00	16.0	84000	0.63	3/4	
832516183	30.55 ± 0.05	9.190	1.00	9.2	28583	0.403	1/1	
804549027	2.948 ± 0.007	117.121	1.00	79.0	10000	0.2	2/2	
191532802	3.553 ± 0.022	5.513	1.00	11.6	8630	0.808	1/1	
262708252	5.4411 ± 0.0015	28.747	1.00	16.1	10266	0.152	7/7	
841325183	2.504 ± 0.0005	10.370	1.00	9.2	17205	0.28	3/4	
873285117	20.79 ± 0.024	9.782	1.00	9.9	37453	0.349	1/2	
875311151	11.053 ± 0.007	9.792	1.00	8.7	50475	0.491	1/2	
309025727	8.22 ± 0.06	19.948	1.00	26.6	9000	0.23	1/2	
870310469	23.443 ± 0.03	9.701	1.00	9.9	37491	0.502	2/2	
841399917	36.32 ± 0.15	64.304	1.00	39.3	17000	0.2	4/4	
33357141	13.47 ± 0.31	8.387	1.00	11.5	24952	0.526	1/1	
345036441	$1.64579 \pm 8\text{e-}05$	13.629	1.00	10.6	10355	0.92	4/5	DAH
471013677	2.79285 ± 0.00031	59.858	1.00	53.6	36470	0.6	1/4	
902670676	22.988 ± 0.029	10.934	1.00	10.8	78000	0.66	2/2	
82347011	6.2108 ± 0.0021	130.009	1.00	94.0	30071	0.4	2/2	DA+M3
950218375	10.3 ± 0.19	7.101	1.00	10.0	42180	0.52	1/1	

TIC	Period [h]	Amplitude [σ]	Like	\langle Height [σ] \rangle	T_{eff} [K]	Mass [M_{\odot}]		Info
232972481	81.13 ± 0.28	89.769	1.00	39.0	99575	0.87	7/7	DA+M
156073827	$0.189861 \pm 2\text{e-}06$	10.486	1.00	10.0	29300	0.415	2/3	
115613388	$0.703919 \pm 2.6\text{e-}05$	6.117	1.00	5.3	28169	1.272	2/5	DAH
630123582	4.109 ± 0.027	3.846	1.00	12.5	60000	0.456	1/1	
630314437	$0.17219 \pm 5\text{e-}05$	4.425	1.00	6.9	21128	0.526	1/1	wd+pair
399570361	10.89 ± 0.22	25.327	0.99	27.6	79000	0.656	1/1	wd+pair
183799565	$0.689793 \pm 2.6\text{e-}05$	5.825	1.00	5.8	180000	0.54	1/4	
142982488	0.40188 ± 0.00026	4.112	1.00	5.6	34180	0.626	1/1	
298411553	2.90014 ± 0.00031	12.166	1.00	8.2	39229	0.756	4/8	
80858168	2.84122 ± 0.0003	38.679	1.00	34.0	12891	0.256	3/3	
652008669	48.0 ± 4.0	6.901	1.00	6.0	31092	0.432	1/1	
412252434	2.40744 ± 0.00022	7.758	1.00	12.8	10206	0.597	4/4	
760481896	6.641 ± 0.005	64.546	1.00	32.8	27691	0.404	4/4	
453006983	9.093 ± 0.005	19.789	1.00	21.5	27425	0.52	2/2	DA+M
804378446	29.44 ± 0.05	12.366	1.00	14.6	48986	0.685	2/2	
310478036	2.36875 ± 0.00029	11.001	1.00	59.3	7000	0.23	7/7	
349408306	3.6493 ± 0.0005	84.357	1.00	25.5	8000	0.15	21/21	
462519068	4.987 ± 0.008	4.490	1.00	19.7	7000	0.15	3/3	
119736060	3.954 ± 0.026	49.116	1.00	56.6	22600	0.42	1/1	DA+M
46199750	1.891 ± 0.008	7.440	1.00	13.9	20000	0.94	1/1	DAP
147918835	2.722 ± 0.012	156.826	1.00	63.5	24661	0.21	1/1	
142808656	7.9038 ± 0.0034	11.463	1.00	10.0	18646	0.031	4/4	
258086867	30.3 ± 1.5	7.607	1.00	5.6	7573	0.753	1/1	
1000946479	2.69774 ± 0.00032	4.572	1.00	5.8	7139	0.6	1/4	DC
274928438	2.464 ± 0.01	5.552	1.00	6.6	6959	0.23	1/1	DA+M
1100582330	2.736 ± 0.017	18.131	1.00	29.2	26828	0.496	1/1	
284900652	7.9923 ± 0.0021	359.311	1.00	71.7	34733	0.36	29/29	DA+dMe
383673264	72.37 ± 0.17	15.671	1.00	6.8	11435	0.661	11/25	
25771075	7.8738 ± 0.0023	14.170	1.00	14.5	15849	0.263	3/3	
47499442	7.83 ± 0.1	18.473	1.00	22.7	7184	0.562	1/1	
33692053	15.1 ± 0.4	14.656	1.00	20.4	24809	0.239	1/1	
34855316	4.461 ± 0.032	30.907	1.00	46.5	8021	0.745	1/1	
355154138	2.37013 ± 0.00029	18.259	1.00	11.4	14569	0.403	7/7	
429233827	7.17 ± 0.0028	113.682	1.00	74.9	10618	0.174	2/2	
32159409	42.8 ± 3.2	9.008	1.00	11.7	10000	0.2	1/1	
1550940367	21.89 ± 0.19	21.620	1.00	15.7	50000	0.425	4/4	
91193988	3.6933 ± 0.0005	18.393	1.00	19.9	13597	0.249	2/2	
192991819	1.81005 ± 0.00018	34.905	1.00	35.9	7000	0.23	3/3	wd+pair
73764818	$0.135393 \pm 3.2\text{e-}05$	4.101	1.00	5.0	35994	0.464	1/1	
178539314	4.3923 ± 0.0011	5.228	1.00	7.8	8201	0.422	2/2	
938779482	0.4613 ± 0.0004	8.293	0.96	14.2	10000	0.2	1/1	wd+pair
251903434	34.6 ± 1.9	109.202	0.98	28.1	7752	0.702	1/1	
159394587	14.753 ± 0.021	7.390	0.85	5.2	16890	0.6	1/4	
74342209	1.557 ± 0.004	146.008	0.87	44.0	9463	0.478	1/1	
1201364356	276.0 ± 6.0	483.909	0.86	20.9	12816	0.61	11/15	wd+pair-disk?
398365709	8.0828 ± 0.0018	72.725	0.92	19.0	22570	0.54	5/5	DA+M
245830829	13.56 ± 0.15	300.596	0.96	18.7	17620	0.63	2/2	DA3+M
329670050	2.703 ± 0.011	99.642	0.98	34.9	9000	0.23	1/1	

TIC	Period [h]	Amplitude [σ]	Like	\langle Height [σ] \rangle	T_{eff} [K]	Mass [M_{\odot}]		Info
243349100	23.447 ± 0.028	14.703	0.98	8.6	7250	0.23	5/6	
1920685932	25.12 ± 0.06	40.975	0.99	12.5	11981	0.388	1/2	wd+pair
269663822	4.5085 ± 0.0008	226.862	0.98	75.2	7000	0.2	20/20	
403292348	3.3522 ± 0.0012	280.702	0.98	101.4	8000	0.2	2/2	
15131739	6.143 ± 0.004	17.562	0.99	16.5	31194	0.403	2/2	DA+L3
142616553	181.0 ± 1.7	31.440	0.84	19.0	14635	0.48	4/6	wd+pair
901406326	$0.113414 \pm 2.2\text{e-}05$	5.396	0.87	9.5	114711	0.701	1/1	
115013365	1.20234 ± 0.00005	6.454	1.00	9.0	12826	0.23	1/2	
269071459	21.51 ± 0.025	17.219	0.98	10.5	9375	0.852	5/5	
137607346	23.927 ± 0.025	190.904	0.98	58.7	9000	0.2	6/6	
458692929	27.4 ± 1.2	5.070	0.84	8.2	35105	0.36	1/1	DA+pair
311948770	16.932 ± 0.01	219.451	0.80	89.7	49650	0.66	5/5	DAO+M
903362877	65.4 ± 1.3	14.507	0.98	5.2	18344	0.488	1/3	wd+pair
158747696	2.041 ± 0.007	23.570	0.98	27.8	8000	0.205	1/1	wd+pair
8158475	56.85 ± 0.18	8.819	0.99	6.9	45333	0.527	1/2	
1990690286	15.322 ± 0.009	17.081	1.00	16.1	100000	0.48	2/2	
814547613	3.3579 ± 0.0006	27.264	0.96	19.8	14115	0.567	2/4	
88509155	128.0 ± 27.0	28.056	1.00	29.4	56610	0.53	1/1	DA+M
1951569775	92.9 ± 0.9	237.305	1.00	36.3	7000	0.2	2/2	
115075496	53.0 ± 4.0	9.833	1.00	9.9	44715	0.469	1/1	DA+M:
118690168	34.85 ± 0.07	7.436	1.00	6.2	11023	0.587	2/3	
138653638	213.5 ± 2.6	40.343	0.78	17.0	12436	0.478	3/4	wd+pair
721486224	150.0 ± 50.0	150.795	0.84	17.1	11692	0.725	3/3	wd+pair
1101853877	121.0 ± 11.0	102.164	1.00	7.0	18791	0.673	1/2	DA+pair
453444667	82.1 ± 1.5	111.345	1.00	59.4	105000	0.75	2/2	DA+BP
246847488	173.0 ± 5.0	51.455	1.00	21.5	5720	0.1	1/1	
262548040	$2^* (0.10614391 \pm 2.5\text{e-}7)$	6.409	1.00	6.2	6230	0.41	3/5	DA7H
229098638	181.6 ± 0.7	815.491	1.00	19.2	18800	0.56	4/4	wd+pair
404156391	28.133 ± 0.030	221.896	0.97	8.8	17770	0.542	2/2	wd+pair
204440456	6.20 ± 0.07	18.679	1.00	21.3	6908	0.6018	1/1	
732235000	23.667 ± 0.020	19.728	1.00	8.5	19212	0.523	11/16	wd+pair
50385872	5.4166 ± 0.0007	5.535	1.00	5.6	13141	0.710	2/8	
61965938	6.8168 ± 0.0013	51.961	1.00	5.1	19505	0.633	1/3	
1174761001	5.7114 ± 0.0018	23.495	1.00	24.2	18311	0.511	2/2	wd+pair
23936802	5.8054 ± 0.0018	17.076	1.00	27.0	17146	0.269	4/4	
253936074	45.16 ± 0.07	25.890	1.00	5.5	29347	0.676	1/2	wd+pair
2055541164	2.53003 ± 0.00023	73.753	1.00	48.6	23575	0.261	4/4	
408253347	7.9829 ± 0.0034	49.765	1.00	36.4	24466	0.47	2/3	DA+dMe
611305959	21.84 ± 0.05	21.142	1.00	16.4	90000	0.6	3/3	
611853501	0.385874 ± 0.000006	13.347	1.00	14.9	21697	1.055	2/2	
2024143938	1.1367 ± 0.0023	17.506	1.00	29.1	20000	0.323	1/1	DAZ+L3
380174982	4.4379 ± 0.0005	15.711	0.98	15.3	8607	0.83	3/3	
219099282	77.17 ± 0.20	22.276	1.00	6.3	45230	0.63	15/25	
610721330	0.671865 ± 0.000017	5.779	1.00	5.0	24332	1.218	1/4	
161820334	6.9232 ± 0.002	32.422	1.00	15.3	18462	0.183	6/7	
441702898	8.3222 ± 0.0029	13.601	0.99	8.6	7501	0.445	6/6	
144197100	0.245758 ± 0.000008	16.037	1.00	16.5	8595	0.87	2/2	
841424790	3.311 ± 0.0004	7.040	1.00	6.0	73027	0.751	3/7	

TIC	Period [h]	Amplitude [σ]	Like	\langle Height [σ] \rangle	T_{eff} [K]	Mass [M_{\odot}]		Info
308292831	4.61 \pm 0.04	6.218	1.00	12.0	9230	0.191	1/1	ELMV+pair
256351952	28.868 \pm 0.029	47.669	1.00	20.3	10403	0.172	5/5	
900086743	2.26444 \pm 0.00029	55.902	1.00	58.0	23230	0.39	2/2	
370238674	3.3448 \pm 0.0004	34.483	1.00	22.4	9453	0.788	6/6	
337219837	6.8813 \pm 0.0018	175.005	1.00	70.4	25203	0.441	6/6	DA+M-B
802015620	12.218 \pm 0.006	73.835	1.00	58.6	12982	0.263	3/3	wd+pair
742167700	18.778 \pm 0.013	17.158	1.00	14.9	80000	0.6	3/3	DO
358247592	5.7849 \pm 0.0012	10.705	1.00	8.7	33601	0.572	4/4	wd+pair
468604087	3.5365 \pm 0.0005	18.791	1.00	17.6	147000	0.706	7/7	wd+pair
232065830	4.5928 \pm 0.0005	87.323	1.00	49.8	19211	0.511	7/7	
437042755	5.7082 \pm 0.0017	31.903	1.00	23.8	18981	1.06	4/4	DBH
321741942	24.16 \pm 0.022	24.560	1.00	27.9	84000	0.82	2/2	
274239484	1.92716 \pm 0.00013	41.484	1.00	23.0	7743	0.764	9/9	DAH
362023614	12.627 \pm 0.006	21.236	1.00	16.1	53220	0.56	3/3	
471015333	18.817 \pm 0.019	13.115	1.00	14.7	70000	0.49	2/2	
620324891	10.7947 \pm 0.0032	30.343	1.00	19.5	70000	0.634	5/5	
32307067	1.90674 \pm 0.0002	36.376	1.00	33.5	16500	0.39	2/2	
140937916	3.3278 \pm 0.0005	27.793	1.00	20.8	15084	0.112	4/4	
407569944	19.185 \pm 0.014	28.815	1.00	23.6	9955	0.552	3/3	
155577443	39.5 \pm 2.5	12.277	1.00	17.1	4000	0.6	1/1	
840458841	6.79 \pm 0.07	5.110	1.00	6.2	58333	1.105	1/1	
397979837	2.67088 \pm 0.00031	68.242	1.00	46.4	31810	0.37	4/4	
150251751	53.85 \pm 0.16	13.089	1.00	13.0	7145	0.722	2/2	
171630076	3.399 \pm 0.018	52.974	1.00	56.1	15000	0.93	1/1	DAP-B
377056497	1.5254 \pm 0.00012	8.006	1.00	6.0	10060	1.06	3/3	DAH-B
357686299	31.57 \pm 0.033	18.783	1.00	8.5	49240	0.63	10/10	DA
321159503	0.465797 \pm 0.000008	8.501	1.00	7.3	7750	0.8	3/3	DAH
741451769	0.187876 \pm 0.000004	10.426	1.00	13.0	30190	1.33	2/2	DC
313144249	0.723464 \pm 0.000019	47.048	1.00	27.9	10671	0.845	8/8	
611402948	2.24076 \pm 0.00019	76.558	1.00	63.7	30000	0.2	2/2	wd+pair
91329200	1.69439 \pm 0.00008	10.872	1.00	6.9	9146	0.88	5/5	
339451801	106.4 \pm 0.5	50.761	1.00	14.8	8742	0.236	2/2	wd+pair
117502568	65.27 \pm 0.16	6.337	1.00	6.3	97080	0.53	1/1	
20656977	6.2179 \pm 0.0011	8.339	1.00	10.1	14065	0.35	2/2	
62823020	1.3488 \pm 0.001	18.161	1.00	15.8	8380	0.164	3/3	
66419115	7.4439 \pm 0.0012	54.261	1.00	44.6	7711	0.307	3/3	
620251540	82 \pm 10	136.482	1.00	20.4	15425	0.53	1/1	wd+pair
270679102	3.5858 \pm 0.0007	71.393	1.00	42.7	17505	0.37	3/3	DA+M
103222871	2.28912 \pm 0.00019	38.290	0.98	21.7	8597	0.76	6/6	
91898005	1.0509 \pm 0.00006	18.931	1.00	14.9	7765	0.881	3/3	
417179225	19.194 \pm 0.02	131.832	0.84	42.4	22260	0.62	2/2	wd+pair
23992223	16.627 \pm 0.015	11.080	1.00	10.5	15160	0.47	2/2	wd+pair
141117787	6.7091 \pm 0.0017	14.991	1.00	16.3	6802	0.914	2/2	DAH
311629054	19.2 \pm 0.6	12.252	1.00	14.5	19780	0.62	1/1	
406782194	1.83857 \pm 0.00019	4.955	1.00	5.2	27800	0.87	1/1	
54636377	25.633 \pm 0.018	19.880	1.00	15.7	19440	0.58	2/2	
53709985	8.362 \pm 0.0026	31.571	1.00	23.8	10466	0.126	3/3	
201892746	4.0085 \pm 0.0006	13.392	1.00	12.4	6476	0.746	2/2	DAH

TIC	Period [h]	Amplitude [σ]	Like	\langle Height [σ] \rangle	T_{eff} [K]	Mass [M_{\odot}]		Info
22704782	21.503 ± 0.025	6.093	1.00	5.5	21144	0.5	2/2	
299437785	2.05802 ± 0.00016	10.585	0.98	12.4	8553	0.815	1/1	
840874557	0.20919 ± 0.00007	4.218	0.91	5.4	24966	0.599	1/1	wd+pair
91903603	0.1237488 ± 0.0000009	4.484	1.00	5.4	6880	0.868	1/2	

REFERENCES

- Adams, W. S. 1915, *PASP*, 27, 236, doi: [10.1086/122440](https://doi.org/10.1086/122440)
- Aerts, C., & Tkachenko, A. 2023, arXiv e-prints, arXiv:2311.08453, doi: [10.48550/arXiv.2311.08453](https://doi.org/10.48550/arXiv.2311.08453)
- Amard, L., Palacios, A., Charbonnel, C., et al. 2019, *A&A*, 631, A77, doi: [10.1051/0004-6361/201935160](https://doi.org/10.1051/0004-6361/201935160)
- Amorim, L. L., Kepler, S. O., Külebi, B., Jordan, S., & Romero, A. D. 2023, *ApJ*, 944, 56, doi: [10.3847/1538-4357/acaff6](https://doi.org/10.3847/1538-4357/acaff6)
- Babcock, H. W. 1960, *ApJ*, 132, 521, doi: [10.1086/146960](https://doi.org/10.1086/146960)
- Bagnulo, S., & Landstreet, J. D. 2020, *A&A*, 643, A134, doi: [10.1051/0004-6361/202038565](https://doi.org/10.1051/0004-6361/202038565)
- . 2022, *ApJL*, 935, L12, doi: [10.3847/2041-8213/ac84d3](https://doi.org/10.3847/2041-8213/ac84d3)
- Barstow, M. A., Jordan, S., O'Donoghue, D., et al. 1995, *MNRAS*, 277, 971, doi: [10.1093/mnras/277.3.971](https://doi.org/10.1093/mnras/277.3.971)
- Beck, J. G. 2000, *SoPh*, 191, 47, doi: [10.1023/A:1005226402796](https://doi.org/10.1023/A:1005226402796)
- Bell, K. J. 2020, in *American Astronomical Society Meeting Abstracts*, Vol. 235, American Astronomical Society Meeting Abstracts #235, 106.06
- Bell, K. J., Hermes, J. J., Vanderbosch, Z., et al. 2017, *The Astrophysical Journal*, 851, 24, doi: [10.3847/1538-4357/aa9702](https://doi.org/10.3847/1538-4357/aa9702)
- Berger, L., Koester, D., Napiwotzki, R., Reid, I. N., & Zuckerman, B. 2005, *A&A*, 444, 565, doi: [10.1051/0004-6361:20053340](https://doi.org/10.1051/0004-6361:20053340)
- Blatman, D., & Ginzburg, S. 2024a, arXiv e-prints, arXiv:2406.08536, doi: [10.48550/arXiv.2406.08536](https://doi.org/10.48550/arXiv.2406.08536)
- . 2024b, *MNRAS*, 528, 3153, doi: [10.1093/mnras/stae222](https://doi.org/10.1093/mnras/stae222)
- Bloeker, T. 1995, *A&A*, 297, 727
- Böhm-Vitense, E. 1958, *Zeitschrift für Astrophysik*, 46, 108
- Bond, H. E., Kawaler, S. D., Ciardullo, R., et al. 1996a, *AJ*, 112, 2699, doi: [10.1086/118214](https://doi.org/10.1086/118214)
- . 1996b, *AJ*, 112, 2699, doi: [10.1086/118214](https://doi.org/10.1086/118214)
- Bouvier, J., Wichmann, R., Grankin, K., et al. 1997, *A&A*, 318, 495
- Bradley, P. A. 2001, *The Astrophysical Journal*, 552, 326, doi: [10.1086/320454](https://doi.org/10.1086/320454)
- Brinkworth, C. S., Burleigh, M. R., Lawrie, K., Marsh, T. R., & Knigge, C. 2013, *The Astrophysical Journal*, 773, 47, doi: [10.1088/0004-637X/773/1/47](https://doi.org/10.1088/0004-637X/773/1/47)
- Caiazzo, I., Burdge, K. B., Fuller, J., et al. 2021, *Nature*, 595, 39, doi: [10.1038/s41586-021-03615-y](https://doi.org/10.1038/s41586-021-03615-y)
- Calcaferro, Leila M., Córscico, Alejandro H., & Althaus, Leandro G. 2016, *A&A*, 589, A40, doi: [10.1051/0004-6361/201527996](https://doi.org/10.1051/0004-6361/201527996)
- Caldwell, D. A., Tenenbaum, P., Twicken, J. D., et al. 2020, *Research Notes of the American Astronomical Society*, 4, 201, doi: [10.3847/2515-5172/abc9b3](https://doi.org/10.3847/2515-5172/abc9b3)
- Castanheira, B. G., Kepler, S. O., Kleinman, S. J., Nitta, A., & Fraga, L. 2013, *MNRAS*, 430, 50, doi: [10.1093/mnras/sts474](https://doi.org/10.1093/mnras/sts474)
- Castro-Tapia, M., Cumming, A., & Fuentes, J. R. 2024a, arXiv e-prints, arXiv:2402.01947, doi: [10.48550/arXiv.2402.01947](https://doi.org/10.48550/arXiv.2402.01947)
- Castro-Tapia, M., Zhang, S., & Cumming, A. 2024b, arXiv e-prints, arXiv:2406.01807, doi: [10.48550/arXiv.2406.01807](https://doi.org/10.48550/arXiv.2406.01807)
- Catalán, S., Isern, J., García-Berro, E., & Ribas, I. 2008, *MNRAS*, 387, 1693, doi: [10.1111/j.1365-2966.2008.13356.x](https://doi.org/10.1111/j.1365-2966.2008.13356.x)
- Choi, J., Dotter, A., Conroy, C., et al. 2016, *ApJ*, 823, 102, doi: [10.3847/0004-637X/823/2/102](https://doi.org/10.3847/0004-637X/823/2/102)
- Córscico, A. H., Althaus, L. G., Miller Bertolami, M. M., & Kepler, S. O. 2019, *A&A Rv*, 27, 7, doi: [10.1007/s00159-019-0118-4](https://doi.org/10.1007/s00159-019-0118-4)
- Córscico, A. H., Uzundag, M., Kepler, S. O., et al. 2021, *A&A*, 645, A117, doi: [10.1051/0004-6361/202039202](https://doi.org/10.1051/0004-6361/202039202)
- Deal, M., Goupil, M. J., Marques, J. P., Reese, D. R., & Lebreton, Y. 2020, *A&A*, 633, A23, doi: [10.1051/0004-6361/201936666](https://doi.org/10.1051/0004-6361/201936666)
- Deheuvels, S., Ballot, J., Beck, P. G., et al. 2015, *A&A*, 580, A96, doi: [10.1051/0004-6361/201526449](https://doi.org/10.1051/0004-6361/201526449)
- Deheuvels, S., Ballot, J., Eggenberger, P., et al. 2020, *A&A*, 641, A117, doi: [10.1051/0004-6361/202038578](https://doi.org/10.1051/0004-6361/202038578)
- Deheuvels, S., García, R. A., Chaplin, W. J., et al. 2012, *ApJ*, 756, 19, doi: [10.1088/0004-637X/756/1/19](https://doi.org/10.1088/0004-637X/756/1/19)
- Deheuvels, S., Doğan, G., Goupil, M. J., et al. 2014, *A&A*, 564, A27, doi: [10.1051/0004-6361/201322779](https://doi.org/10.1051/0004-6361/201322779)
- den Hartogh, J. W., Eggenberger, P., & Deheuvels, S. 2020, *A&A*, 634, L16, doi: [10.1051/0004-6361/202037568](https://doi.org/10.1051/0004-6361/202037568)

- den Hartogh, J. W., Eggenberger, P., & Hirschi, R. 2019, *A&A*, 622, A187, doi: [10.1051/0004-6361/201834330](https://doi.org/10.1051/0004-6361/201834330)
- Di Mauro, M. P., Ventura, R., Corsaro, E., & Lustosa De Moura, B. 2018, *ApJ*, 862, 9, doi: [10.3847/1538-4357/aac7c4](https://doi.org/10.3847/1538-4357/aac7c4)
- Di Mauro, M. P., Ventura, R., Cardini, D., et al. 2016, *ApJ*, 817, 65, doi: [10.3847/0004-637X/817/1/65](https://doi.org/10.3847/0004-637X/817/1/65)
- Diaz-Pinto, A., Garcia-Berro, E., Hernanz, M., Isern, J., & Mochkovitch, R. 1994, *A&A*, 282, 86
- Doherty, C. L., Gil-Pons, P., Siess, L., Lattanzio, J. C., & Lau, H. H. B. 2014, *Monthly Notices of the Royal Astronomical Society*, 446, 2599, doi: [10.1093/mnras/stu2180](https://doi.org/10.1093/mnras/stu2180)
- Dolez, N., Vauclair, G., Kleinman, S. J., et al. 2006, *A&A*, 446, 237, doi: [10.1051/0004-6361:20053149](https://doi.org/10.1051/0004-6361:20053149)
- Dominguez, I., Chieffi, A., Limongi, M., & Straniero, O. 1999, *ApJ*, 524, 226, doi: [10.1086/307787](https://doi.org/10.1086/307787)
- Douglas, S. T., Cargile, P. A., Matt, S. P., et al. 2024, *ApJ*, 962, 16, doi: [10.3847/1538-4357/ad0fe3](https://doi.org/10.3847/1538-4357/ad0fe3)
- Duan, R. M., Zong, W., Fu, J. N., et al. 2021, *ApJ*, 922, 2, doi: [10.3847/1538-4357/ac22fd](https://doi.org/10.3847/1538-4357/ac22fd)
- Dufour, P., Blouin, S., Coutu, S., et al. 2017, in *Astronomical Society of the Pacific Conference Series*, Vol. 509, 20th European White Dwarf Workshop, ed. P. E. Tremblay, B. Gaensicke, & T. Marsh, 3, doi: [10.48550/arXiv.1610.00986](https://doi.org/10.48550/arXiv.1610.00986)
- Eggenberger, P., den Hartogh, J. W., Buldgen, G., et al. 2019a, *A&A*, 631, L6, doi: [10.1051/0004-6361/201936348](https://doi.org/10.1051/0004-6361/201936348)
- Eggenberger, P., Moyano, F. D., & den Hartogh, J. W. 2022, *A&A*, 664, L16, doi: [10.1051/0004-6361/202243781](https://doi.org/10.1051/0004-6361/202243781)
- Eggenberger, P., Deheuvels, S., Miglio, A., et al. 2019b, *A&A*, 621, A66, doi: [10.1051/0004-6361/201833447](https://doi.org/10.1051/0004-6361/201833447)
- Farihi, J., Robert, A., & Walters, N. 2023, A Nearby Polluted White Dwarf with a 6.2 h Spin Period. <https://arxiv.org/abs/2312.03845>
- Fleming, T. A., Liebert, J., & Green, R. F. 1986, *ApJ*, 308, 176, doi: [10.1086/164488](https://doi.org/10.1086/164488)
- Fritzewski, D. J., Barnes, S. A., James, D. J., & Strassmeier, K. G. 2021, *A&A*, 652, A60, doi: [10.1051/0004-6361/202140894](https://doi.org/10.1051/0004-6361/202140894)
- Fu, J.-N., Dolez, N., Vauclair, G., et al. 2012, *Monthly Notices of the Royal Astronomical Society*, 429, 1585, doi: [10.1093/mnras/sts438](https://doi.org/10.1093/mnras/sts438)
- Fu, J.-N., Vauclair, G., Su, J., et al. 2019, *Monthly Notices of the Royal Astronomical Society*, 486, 3560, doi: [10.1093/mnras/stz1088](https://doi.org/10.1093/mnras/stz1088)
- Fu, J.-N., Vauclair, G., Solheim, J.-E., et al. 2007, *A&A*, 467, 237, doi: [10.1051/0004-6361:20066295](https://doi.org/10.1051/0004-6361:20066295)
- Fuller, J. 2017, *MNRAS*, 472, 1538, doi: [10.1093/mnras/stx2135](https://doi.org/10.1093/mnras/stx2135)
- Fuller, J., & Lai, D. 2011, *MNRAS*, 412, 1331, doi: [10.1111/j.1365-2966.2010.18017.x](https://doi.org/10.1111/j.1365-2966.2010.18017.x)
- . 2013, *MNRAS*, 430, 274, doi: [10.1093/mnras/sts606](https://doi.org/10.1093/mnras/sts606)
- Fuller, J., & Mathis, S. 2023, *MNRAS*, 520, 5573, doi: [10.1093/mnras/stad475](https://doi.org/10.1093/mnras/stad475)
- Fuller, J., Piro, A. L., & Jermyn, A. S. 2019, *MNRAS*, 485, 3661, doi: [10.1093/mnras/stz514](https://doi.org/10.1093/mnras/stz514)
- Gaia Collaboration, Babusiaux, C., van Leeuwen, F., et al. 2018, *A&A*, 616, A10, doi: [10.1051/0004-6361/201832843](https://doi.org/10.1051/0004-6361/201832843)
- García-Berro, E., Isern, J., & Hernanz, M. 1997, *Monthly Notices of the Royal Astronomical Society*, 289, 973, doi: [10.1093/mnras/289.4.973](https://doi.org/10.1093/mnras/289.4.973)
- Gehan, C., Mosser, B., Michel, E., Samadi, R., & Kallinger, T. 2018, *A&A*, 616, A24, doi: [10.1051/0004-6361/201832822](https://doi.org/10.1051/0004-6361/201832822)
- Gentile Fusillo, N. P., Tremblay, P. E., Cukanovaite, E., et al. 2021, *MNRAS*, 508, 3877, doi: [10.1093/mnras/stab2672](https://doi.org/10.1093/mnras/stab2672)
- Gentile Fusillo, N. P., Tremblay, P.-E., Cukanovaite, E., et al. 2021, *Monthly Notices of the Royal Astronomical Society*, 508, 3877, doi: [10.1093/mnras/stab2672](https://doi.org/10.1093/mnras/stab2672)
- Giammichele, N., Fontaine, G., Brassard, P., & Charpinet, S. 2016, *The Astrophysical Journal Supplement Series*, 223, 10, doi: [10.3847/0067-0049/223/1/10](https://doi.org/10.3847/0067-0049/223/1/10)
- Granada, A., & Haemmerlé, L. 2014, *A&A*, 570, A18, doi: [10.1051/0004-6361/201424068](https://doi.org/10.1051/0004-6361/201424068)
- Greiss, S., Gänsicke, B. T., Hermes, J. J., et al. 2014, *Monthly Notices of the Royal Astronomical Society*, 438, 3086, doi: [10.1093/mnras/stt2420](https://doi.org/10.1093/mnras/stt2420)
- Guo, J., Zhao, J., Zhang, H., et al. 2022, *MNRAS*, 509, 2674, doi: [10.1093/mnras/stab3151](https://doi.org/10.1093/mnras/stab3151)
- Heger, A., Langer, N., & Woosley, S. E. 2000, *ApJ*, 528, 368, doi: [10.1086/308158](https://doi.org/10.1086/308158)
- Heger, A., Woosley, S. E., & Spruit, H. C. 2005, *ApJ*, 626, 350, doi: [10.1086/429868](https://doi.org/10.1086/429868)
- Heney, L., Vardya, M. S., & Bodenheimer, P. 1965, *ApJ*, 142, 841, doi: [10.1086/148357](https://doi.org/10.1086/148357)
- Hermes, J. J., Kawaler, S. D., Bischoff-Kim, A., et al. 2017a, *The Astrophysical Journal*, 835, 277, doi: [10.3847/1538-4357/835/2/277](https://doi.org/10.3847/1538-4357/835/2/277)
- Hermes, J. J., Gänsicke, B. T., Kawaler, S. D., et al. 2017b, *ApJS*, 232, 23, doi: [10.3847/1538-4365/aa8bb5](https://doi.org/10.3847/1538-4365/aa8bb5)
- Higgins, M. E., & Bell, K. J. 2023, *AJ*, 165, 141, doi: [10.3847/1538-3881/acb20c](https://doi.org/10.3847/1538-3881/acb20c)
- Höfner, S., & Olofsson, H. 2018, *A&A Rv*, 26, 1, doi: [10.1007/s00159-017-0106-5](https://doi.org/10.1007/s00159-017-0106-5)
- Jermyn, A. S., Bauer, E. B., Schwab, J., et al. 2023, *ApJS*, 265, 15, doi: [10.3847/1538-4365/acae8d](https://doi.org/10.3847/1538-4365/acae8d)

- Kawaler, S. D. 2015, in *Astronomical Society of the Pacific Conference Series*, Vol. 493, 19th European Workshop on White Dwarfs, ed. P. Dufour, P. Bergeron, & G. Fontaine, 65, doi: [10.48550/arXiv.1410.6934](https://doi.org/10.48550/arXiv.1410.6934)
- Kawaler, S. D., O'Brien, M. S., Clemens, J. C., et al. 1995, *ApJ*, 450, 350, doi: [10.1086/176145](https://doi.org/10.1086/176145)
- Kepler, S., & Romero, A. D. 2017, *EPJ Web Conf.*, 152, 01011, doi: [10.1051/epjconf/201715201011](https://doi.org/10.1051/epjconf/201715201011)
- Kepler, S. O. 1993, *Baltic Astronomy*, 2, 515, doi: [10.1515/astro-1993-3-425](https://doi.org/10.1515/astro-1993-3-425)
- Kepler, S. O., Kleinman, S. J., Nitta, A., et al. 2007, *MNRAS*, 375, 1315, doi: [10.1111/j.1365-2966.2006.11388.x](https://doi.org/10.1111/j.1365-2966.2006.11388.x)
- Kepler, S. O., Koester, D., Pelisoli, I., Romero, A. D., & Ourique, G. 2021, *Monthly Notices of the Royal Astronomical Society*, 507, 4646, doi: [10.1093/mnras/stab2411](https://doi.org/10.1093/mnras/stab2411)
- Kepler, S. O., Koester, D., Pelisoli, I., Romero, A. D., & Ourique, G. 2021, *MNRAS*, 507, 4646, doi: [10.1093/mnras/stab2411](https://doi.org/10.1093/mnras/stab2411)
- Kepler, S. O., Robinson, E. L., & Nather, R. E. 1983, *ApJ*, 271, 744, doi: [10.1086/161241](https://doi.org/10.1086/161241)
- Kepler, S. O., Giovannini, O., Wood, M. A., et al. 1995, *ApJ*, 447, 874, doi: [10.1086/175924](https://doi.org/10.1086/175924)
- Kepler, S. O., Pelisoli, I., Koester, D., et al. 2015, *MNRAS*, 446, 4078, doi: [10.1093/mnras/stu2388](https://doi.org/10.1093/mnras/stu2388)
- . 2019, *MNRAS*, 486, 2169, doi: [10.1093/mnras/stz960](https://doi.org/10.1093/mnras/stz960)
- Kilic, M., Bergeron, P., Kosakowski, A., et al. 2020, *ApJ*, 898, 84, doi: [10.3847/1538-4357/ab9b8d](https://doi.org/10.3847/1538-4357/ab9b8d)
- Kilic, M., Stanek, K. Z., & Pinsonneault, M. H. 2007, *ApJ*, 671, 761, doi: [10.1086/522228](https://doi.org/10.1086/522228)
- Kilic, M., Moss, A. G., Kosakowski, A., et al. 2023, *MNRAS*, 518, 2341, doi: [10.1093/mnras/stac3182](https://doi.org/10.1093/mnras/stac3182)
- Kochukhov, O. 2011, in *Physics of Sun and Star Spots*, ed. D. Prasad Choudhary & K. G. Strassmeier, Vol. 273, 249–255, doi: [10.1017/S1743921311015328](https://doi.org/10.1017/S1743921311015328)
- Krtićka, J., Kawka, A., Mikulášek, Z., et al. 2023, *Astronomy & Astrophysics*, 674, A94, doi: [10.1051/0004-6361/202245551](https://doi.org/10.1051/0004-6361/202245551)
- Kurtz, D. W., Shibahashi, H., Murphy, S. J., Bedding, T. R., & Bowman, D. M. 2015, *MNRAS*, 450, 3015, doi: [10.1093/mnras/stv868](https://doi.org/10.1093/mnras/stv868)
- Labadie-Bartz, J., Hümmelich, S., Bernhard, K., Paunzen, E., & Shultz, M. E. 2023, *A&A*, 676, A55, doi: [10.1051/0004-6361/202346657](https://doi.org/10.1051/0004-6361/202346657)
- Langer, N. 2012, *Annual Review of Astronomy and Astrophysics*, 50, 107, doi: [10.1146/annurev-astro-081811-125534](https://doi.org/10.1146/annurev-astro-081811-125534)
- Li, C., Fu, J., Fox-Machado, L., Su, J., & Chen, F. 2017, *NewA*, 55, 48, doi: [10.1016/j.newast.2017.03.003](https://doi.org/10.1016/j.newast.2017.03.003)
- Liebert, J., Bergeron, P., & Holberg, J. B. 2005, *ApJS*, 156, 47, doi: [10.1086/425738](https://doi.org/10.1086/425738)
- Lightkurve Collaboration, Cardoso, J. V. d. M., Hedges, C., et al. 2018, *Lightkurve: Kepler and TESS time series analysis in Python*. <http://ascl.net/1812.013>
- Lopez, I. D., Hermes, J. J., Calcaferro, L. M., et al. 2021, *ApJ*, 922, 220, doi: [10.3847/1538-4357/ac2d28](https://doi.org/10.3847/1538-4357/ac2d28)
- Marsh, T. R., Dhillon, V. S., & Duck, S. R. 1995, *MNRAS*, 275, 828, doi: [10.1093/mnras/275.3.828](https://doi.org/10.1093/mnras/275.3.828)
- Mathys, G., Holdsworth, D. L., & Kurtz, D. W. 2024, *A&A*, 683, A227, doi: [10.1051/0004-6361/202348259](https://doi.org/10.1051/0004-6361/202348259)
- McCook, G. P., & Sion, E. M. 2016, *VizieR Online Data Catalog*, 1, 2035
- Molnár, L., Szabó, R., & Plachy, E. 2016, *JAAVSO*, 44, 168, doi: [10.48550/arXiv.1610.02004](https://doi.org/10.48550/arXiv.1610.02004)
- Moss, A., Kilic, M., Bergeron, P., Fergard, M., & Brown, W. 2023, *MNRAS*, 523, 5598, doi: [10.1093/mnras/stad1835](https://doi.org/10.1093/mnras/stad1835)
- Mosser, B., Goupil, M. J., Belkacem, K., et al. 2012, *A&A*, 548, A10, doi: [10.1051/0004-6361/201220106](https://doi.org/10.1051/0004-6361/201220106)
- Moyano, F. D., Eggenberger, P., Meynet, G., et al. 2022, *A&A*, 663, A180, doi: [10.1051/0004-6361/202243389](https://doi.org/10.1051/0004-6361/202243389)
- Moyano, F. D., Eggenberger, P., Salmon, S. J. A. J., Mombarg, J. S. G., & Ekström, S. 2023, *A&A*, 677, A6, doi: [10.1051/0004-6361/202346548](https://doi.org/10.1051/0004-6361/202346548)
- Nather, R. E., Winget, D. E., Clemens, J. C., Hansen, C. J., & Hine, B. P. 1990, *ApJ*, 361, 309, doi: [10.1086/169196](https://doi.org/10.1086/169196)
- Nguyen, C. T., Costa, G., Girardi, L., et al. 2022, *A&A*, 665, A126, doi: [10.1051/0004-6361/202244166](https://doi.org/10.1051/0004-6361/202244166)
- O'Brien, M. W., Tremblay, P. E., Gentile Fusillo, N. P., et al. 2023, *MNRAS*, 518, 3055, doi: [10.1093/mnras/stac3303](https://doi.org/10.1093/mnras/stac3303)
- Oliveira da Rosa, G., Kepler, S. O., Córscico, A. H., et al. 2022, *ApJ*, 936, 187, doi: [10.3847/1538-4357/ac8871](https://doi.org/10.3847/1538-4357/ac8871)
- Oschmann, J. M., Jr. Clampin, M., & Fazio, Giovanni G. and MacEwen, H. A. 2014, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 9143, *Space Telescopes and Instrumentation 2014: Optical, Infrared, and Millimeter Wave*
- Østensen, R. H., Bloemen, S., Vučković, M., et al. 2011, *ApJL*, 736, L39, doi: [10.1088/2041-8205/736/2/L39](https://doi.org/10.1088/2041-8205/736/2/L39)
- Pablo, H., Kawaler, S. D., Reed, M. D., et al. 2012, *MNRAS*, 422, 1343, doi: [10.1111/j.1365-2966.2012.20707.x](https://doi.org/10.1111/j.1365-2966.2012.20707.x)
- Paxton, B., Bildsten, L., Dotter, A., et al. 2011, *ApJS*, 192, 3, doi: [10.1088/0067-0049/192/1/3](https://doi.org/10.1088/0067-0049/192/1/3)
- Paxton, B., Cantiello, M., Arras, P., et al. 2013, *ApJS*, 208, 4, doi: [10.1088/0067-0049/208/1/4](https://doi.org/10.1088/0067-0049/208/1/4)
- Paxton, B., Marchant, P., Schwab, J., et al. 2015, *ApJS*, 220, 15, doi: [10.1088/0067-0049/220/1/15](https://doi.org/10.1088/0067-0049/220/1/15)

- Paxton, B., Schwab, J., Bauer, E. B., et al. 2018, *ApJS*, 234, 34, doi: [10.3847/1538-4365/aaa5a8](https://doi.org/10.3847/1538-4365/aaa5a8)
- Paxton, B., Smolec, R., Schwab, J., et al. 2019, *ApJS*, 243, 10, doi: [10.3847/1538-4365/ab2241](https://doi.org/10.3847/1538-4365/ab2241)
- Pech, D., & Vauclair, G. 2006, *A&A*, 453, 219, doi: [10.1051/0004-6361:20054370](https://doi.org/10.1051/0004-6361:20054370)
- Pedersen, M. G., & Bell, K. J. 2023, *AJ*, 165, 239, doi: [10.3847/1538-3881/accc31](https://doi.org/10.3847/1538-3881/accc31)
- Pfeiffer, B., Vauclair, G., Dolez, N., et al. 1996, *A&A*, 314, 182
- Poelarends, A. J. T., Herwig, F., Langer, N., & Heger, A. 2008, *The Astrophysical Journal*, 675, 614, doi: [10.1086/520872](https://doi.org/10.1086/520872)
- Reimers, D. 1975, *Memoires of the Societe Royale des Sciences de Liege*, 8, 369
- Rohrmann, R. D., Althaus, L. G., & Kepler, S. O. 2011, *MNRAS*, 411, 781, doi: [10.1111/j.1365-2966.2010.17716.x](https://doi.org/10.1111/j.1365-2966.2010.17716.x)
- Romero, A. D., Campos, F., & Kepler, S. O. 2015, *MNRAS*, 450, 3708, doi: [10.1093/mnras/stv848](https://doi.org/10.1093/mnras/stv848)
- Romero, A. D., Kepler, S. O., Da Rosa, G. O., & Hermes, J. J. 2024, *ApJ*, submitted
- Romero, A. D., Kepler, S. O., Hermes, J. J., et al. 2022, *Monthly Notices of the Royal Astronomical Society*, 511, 1574, doi: [10.1093/mnras/stac093](https://doi.org/10.1093/mnras/stac093)
- Schafferoth, V., Barlow, B. N., Pelisoli, I., Geier, S., & Kupfer, T. 2023, *A&A*, 673, A90, doi: [10.1051/0004-6361/202244697](https://doi.org/10.1051/0004-6361/202244697)
- Schmidt, M. 1968, *ApJ*, 151, 393, doi: [10.1086/149446](https://doi.org/10.1086/149446)
- Siess, L. 2010, *A&A*, 512, A10, doi: [10.1051/0004-6361/200913556](https://doi.org/10.1051/0004-6361/200913556)
- Snodgrass, H. B., & Ulrich, R. K. 1990, *ApJ*, 351, 309, doi: [10.1086/168467](https://doi.org/10.1086/168467)
- Spruit, H. C. 2002, *A&A*, 381, 923, doi: [10.1051/0004-6361:20011465](https://doi.org/10.1051/0004-6361:20011465)
- Steen, M., Hermes, J. J., Guidry, J. A., et al. 2024, *ApJ*, 967, 166, doi: [10.3847/1538-4357/ad3e60](https://doi.org/10.3847/1538-4357/ad3e60)
- Su, J., Li, Y., Fu, J.-N., & Li, C. 2013, *Monthly Notices of the Royal Astronomical Society*, 437, 2566, doi: [10.1093/mnras/stt2069](https://doi.org/10.1093/mnras/stt2069)
- Tayar, J., Beck, P. G., Pinsonneault, M. H., García, R. A., & Mathur, S. 2019, *ApJ*, 887, 203, doi: [10.3847/1538-4357/ab558a](https://doi.org/10.3847/1538-4357/ab558a)
- Team, M. 2021a, TESS "Fast" Light Curves - All Sectors, STScI/MAST, doi: [10.17909/T9-ST5G-3177](https://doi.org/10.17909/T9-ST5G-3177)
- . 2021b, TESS "Fast" Target Pixel Files - All Sectors, STScI/MAST, doi: [10.17909/T9-TCN7-7G94](https://doi.org/10.17909/T9-TCN7-7G94)
- . 2021c, TESS Light Curves - All Sectors, STScI/MAST, doi: [10.17909/T9-NMC8-F686](https://doi.org/10.17909/T9-NMC8-F686)
- . 2021d, TESS Target Pixel Files - All Sectors, STScI/MAST, doi: [10.17909/T9-YK4W-ZC73](https://doi.org/10.17909/T9-YK4W-ZC73)
- Toonen, S., Hollands, M., Gänsicke, B. T., & Boekholt, T. 2017, *A&A*, 602, A16, doi: [10.1051/0004-6361/201629978](https://doi.org/10.1051/0004-6361/201629978)
- Tout, C. A., Wickramasinghe, D. T., Liebert, J., Ferrario, L., & Pringle, J. E. 2008, *MNRAS*, 387, 897, doi: [10.1111/j.1365-2966.2008.13291.x](https://doi.org/10.1111/j.1365-2966.2008.13291.x)
- Tremblay, P. E., Hollands, M. A., Gentile Fusillo, N. P., et al. 2020, *MNRAS*, 497, 130, doi: [10.1093/mnras/staa1892](https://doi.org/10.1093/mnras/staa1892)
- Valyavin, G., Bagnulo, S., Fabrika, S., et al. 2006, *ApJ*, 648, 559, doi: [10.1086/505781](https://doi.org/10.1086/505781)
- Wenger, M., Ochsenbein, F., Egret, D., et al. 2000, *A&AS*, 143, 9, doi: [10.1051/aas:2000332](https://doi.org/10.1051/aas:2000332)
- Willems, B., & Kolb, U. 2004, *A&A*, 419, 1057, doi: [10.1051/0004-6361:20040085](https://doi.org/10.1051/0004-6361:20040085)
- Williams, K. A., Hermes, J. J., & Vanderbosch, Z. P. 2022, *AJ*, 164, 131, doi: [10.3847/1538-3881/ac8543](https://doi.org/10.3847/1538-3881/ac8543)
- Woosley, S. E., & Heger, A. 2015, *The Astrophysical Journal*, 810, 34, doi: [10.1088/0004-637X/810/1/34](https://doi.org/10.1088/0004-637X/810/1/34)
- Zahn, J. P. 1977, *A&A*, 57, 383