

COMPLEMENTS OF NON-MINIMAL SUBSPACES: CHARACTERIZATION RESULTS

GEIVISON RIBEIRO

ABSTRACT. Inspired by the work of L. Drewnowski in [Studia Math. **77** (1984) 373–391], our research reveals new insights and characterizes the notion of spaceability in the context of complements of subspaces (not necessarily closed) within the universe of F -spaces in terms of $[S]$ -lineability.

1. INTRODUCTION AND BACKGROUND

The investigation into the existence of infinite-dimensional closed subspaces devoid of a *basic sequence* within the context of F -spaces made significant strides with the work of N. Kalton in [28], who demonstrated the following result:

Theorem 1.1. [28, Theorem 1.1] *There is a quasi-Banach space E with a one-dimensional subspace L so that*

(1) *if E_0 is a closed infinite-dimensional subspace of E then $L \subset E_0$*

and

(2) *E/L is isomorphic to the Banach space ℓ_1 .*

In particular, E contains no basic sequence and is minimal.

In addition to revealing that the set $(E \setminus L) \cup \{0\}$ does not contain an infinite-dimensional closed subspace, this result is particularly significant when contrasted with the classical result proved by Enflo [20] and provided by Banach [8, p. 238], which states that *every Banach space contains an infinite-dimensional closed subspace with a basic sequence*. While the classical theorem guarantees the presence of such subspaces in Banach spaces, Kalton's result reveals the additional complexity and richness of linear structures in F -spaces.

This distinction not only underscores the fundamental difference between F -spaces and Banach spaces but also highlights the diversity of possible linear configurations in F -space environments.

According to [27], an F -space E contains a basic sequence if and only if there exists a strictly weaker Hausdorff vector topology on E . Thus, the existence of a space without a basic sequence is equivalent to the existence of a (topologically) minimal space (i.e., one in which no strictly weaker Hausdorff vector topology exists).

Among the positive results regarding the existence of closed subspaces containing basic sequences, one notable contribution is the work by Bastero in [7], which provides information that every subspace of $L_p[0, 1]$ with $0 < p < 1$, contains a basic sequence, and a result due to L. Drewnowski in [17], as follows:

Theorem 1.2. ([17, Theorem 3.3]) *For an F -space E , if F is a closed subspace non-minimal of E and $\dim(E/F) = \infty$, then there exists a closed non-minimal subspace G of E such that $F \cap G = \{0\}$ and $F + G$ is not closed.*

2020 *Mathematics Subject Classification.* 15A03, 46B87, 46A16.

Key words and phrases. Lineability, Spaceability, F -spaces, Banach Spaces, Basic Sequences .

G. Ribeiro is supported by Grant 2022/1962, Paraíba State Research Foundation (FAPESQ).

To provide context, let us fix some notations and terminology. We denote \mathbb{N} as the set of positive integers, \mathbb{R} as the real scalar field, and \mathbb{C} as the complex scalar field. Unless explicitly stated otherwise, all linear spaces are over $\mathbb{K} = \mathbb{R}$ or \mathbb{C} . The term subspace will be used instead of vector subspace. Additionally, α and β represent cardinal numbers, $\text{card}(A)$ denotes the cardinality of set A , $\aleph_0 := \text{card}(\mathbb{N})$, and $\mathfrak{c} := \text{card}(\mathbb{R})$. Furthermore, if E is a vector space and W is a subspace of E , we denote the algebraic codimension of W by $\dim(E/W)$.

In terms of notions, assume that E is a vector space and let $\beta \leq \dim(E)$ be a cardinal number. A subset $A \subset E$, is said to be:

- *lineable* if there is an infinite dimensional subspace F of E such that $F \setminus \{0\} \subset A$, and
- β -*lineable* if there exists a subspace F_β of E with $\dim(F_\beta) = \beta$ and $F_\beta \setminus \{0\} \subset A$ (in this context, a set is *lineable* if it is \aleph_0 -lineable).

This concepts was first introduced in the seminal paper [6] by Aron, Gurariy and Seoane-Sepúlveda and, later, in [20, 25]. Its essence is to investigate linear structures within exotic settings. A more detailed description of the concept of lineability/spaceability can also be found in [5, 9, 12, 14, 20, 24].

Moreover, if we consider a another cardinal number α , with $\alpha \leq \beta$, then A is said to be:

- (α, β) -*lineable* if it is α -lineable and for every subspace $F_\alpha \subset E$ with $F_\alpha \subset A \cup \{0\}$ and $\dim(F_\alpha) = \alpha$, there is a subspace $F_\beta \subset E$ with $\dim(F_\beta) = \beta$ and

$$(1.1) \quad F_\alpha \subset F_\beta \subset A \cup \{0\}$$

(hence $(0, \beta)$ -lineability $\Leftrightarrow \beta$ -lineability).

If E is, in addition, is a topological vector space, then A is called:

- β -*spaceable* if $A \cup \{0\}$ contains a closed β -dimensional subspace of E .
- Moreover, if the subspace F_β satisfying (1.1) can always be chosen closed, we say that A is (α, β) -*spaceable* (see [22]).

In the quest for a deeper understanding of dimensional relationships within spaceable sets, Fávoro, Pellegrino, and Tomaz introduced this concept in 2019 (referenced as [22]). This heightened conceptualization, although rooted in conventional spaceability, demands meticulous consideration, encompassing both geometric and topological properties within the set, as well as relationships between the dimensions of subspaces therein. It is worth noting that while conventional spaceability implies the variant known as (α, α) -spaceability, the reverse assertion is not always valid, as demonstrated by Araújo et al. in [4]. For a more comprehensive exposition on this notion, we refer the reader to [1, 2, 3, 11, 13, 15, 16, 21, 22, 32].

Connected to the notion of lineability, for a subspace \mathcal{S} of $\mathbb{K}^{\mathbb{N}}$, we say that a subset A of a Hausdorff topological vector space X is $[(u_n)_{n=1}^{\infty}, \mathcal{S}]$ -lineable in X if, for each sequence $(c_n)_{n=1}^{\infty} \in \mathcal{S}$, the series $\sum_{n=1}^{\infty} c_n u_n$ converges in X to a vector in $A \cup \{0\}$. Moreover, A is $[\mathcal{S}]$ -lineable in X if it is $[(u_n)_{n=1}^{\infty}, \mathcal{S}]$ -lineable for some sequence $(u_n)_{n=1}^{\infty}$ of linearly independent elements in X .

This concept was introduced in [10] by Bernal-González et al. and originally coined by V. Gurariy and R. Aron during a Non-Linear Analysis Seminar at Kent State University. As far as we know, this notion was initially inspired by a result of Levine and Milman from 1940 [31], created to address the lack of such a concept regarding convergence. Regarding the result of Levine and Milman, they demonstrate that the subset of $C[0, 1]$ consisting of all functions of bounded variation does not contain a closed subspace of infinite dimension.

The results established in this paper are:

- **(Main Result)** For an F -space E , if W is a subspace of E that contains a non-minimal closed subspace, then the set $E \setminus W$ is spaceable if and only if it is $[(x_n)_{n=1}^{\infty}, \mathcal{S}]$ -lineable for some closed subspace \mathcal{S} of ℓ_∞ containing c_0 and some \mathcal{S} -independent sequence $(x_n)_{n=1}^{\infty}$ of

elements of E .

- For a Banach space E , if F is a closed subspace of E , then the following conditions are equivalent:
 - (1) $E \setminus F$ is $[(x_n)_{n=1}^\infty, \mathcal{S}]$ -lineable for some closed subspace \mathcal{S} of ℓ_∞ containing c_0 and some \mathcal{S} -independent sequence $(x_n)_{n=1}^\infty$ of elements of E ,
 - (2) $E \setminus F$ is spaceable,
 - (3) F has infinite codimension.
- For an F -space E , if W is a subspace of E that contains a non-minimal closed subspace, then the following conditions are equivalent:
 - (1) $E \setminus W$ is spaceable,
 - (2) $E \setminus W$ is $[(x_n)_{n=1}^\infty, \mathcal{S}]$ -lineable for some closed subspace \mathcal{S} of ℓ_∞ containing c_0 and some \mathcal{S} -independent sequence $(x_n)_{n=1}^\infty$ of elements of E ,
 - (3) $E \setminus W$ is (n, \mathfrak{c}) -spaceable for each $n \in \mathbb{N}$.

2. PRELIMINARIES

In this paper, we will characterize some families of complements of subspaces (not necessarily closed) through the notion of $[\mathcal{S}]$ -lineability. To this end, let us begin with some preliminary notions and results.

Definition 2.1. [17, p. 376] *We say that a sequence $(u_n)_{n=1}^\infty$ of elements of a topological vector space E is topologically linearly independent, if for each sequence $(c_n)_{n=1}^\infty \in \mathbb{K}^\mathbb{N}$ with $\sum_{n=1}^\infty c_n u_n = 0$, we have $(c_n)_{n=1}^\infty = 0$.*

Based on this definition, if $\mathcal{S} \neq \{0\}$ is a subspace of $\mathbb{K}^\mathbb{N}$ then we will say that a sequence $(u_n)_{n=1}^\infty$ of elements of a topological vector space E is \mathcal{S} -topologically linearly independent in E (or \mathcal{S} -independent) if for each sequence $(c_n)_{n=1}^\infty \in \mathcal{S}$ with $\sum_{n=1}^\infty c_n u_n = 0$, we have $(c_n)_{n=1}^\infty = 0$.

In connection with this notion, we present the following classic definition.

Definition 2.2. *A sequence $(e_k)_{k=1}^\infty$ in a metrizable topological vector space E is said to be basic whenever every $x \in \text{span} \{e_k\}_{k=1}^\infty$, the closed linear span of the x_j 's, can be uniquely represented as a convergent series $x = \sum_{k=1}^\infty a_k e_k$. Furthermore, as in [28], a basic sequence is called regular if it is bounded away from zero, that is, if it lies entirely outside some neighborhood of zero.*

From the perspective of minimal subspaces, as discussed in the introduction, Kalton and Shapiro in [27] proved the following result:

Proposition 2.3. *If E is an F -space, then the following are equivalent:*

- (1) E is non-minimal,
- (2) E contains a regular basic sequence.

The following Lemma is crucial for the next results.

Lemma 2.4. [29, Lemma 2] *For F -spaces Z and E , if $K: Z \rightarrow E$ is a compact operator and $\mathcal{I}: Z \rightarrow E$ is the identity embedding then the operator $\mathcal{I} + K: Z \rightarrow E$ has closed range.*

Based on these notions, Drewnowski in [17] demonstrated the following result:

Theorem 2.5. ([17, Theorem 3.3]) *For an F -space E , if F is a closed subspace non-minimal of E and $\dim(E/F) = \infty$, then there exists a closed non-minimal subspace F of E such that $F \cap F = \{0\}$ and $F + F$ is not closed.*

To the best of our knowledge, there is no criterion in the literature that establishes spaceability for the complement of a subspace W (not necessarily closed) in an F -space E .

3. THE MAIN RESULT

The next result is a characterization for spaceability in the context of complements of subspaces that are not necessarily closed, which was inspired by 2.5.

Theorem 3.1. *For an F -space E , if W be a subspace of E that contains a non-minimal closed subspace, then the set $E \setminus W$ is spaceable if and only if it is $[(x_n)_{n=1}^\infty, \mathcal{S}]$ -lineable for some closed subspace \mathcal{S} of ℓ_∞ containing c_0 and some \mathcal{S} -independent sequence $(x_n)_{n=1}^\infty$ of elements of E .*

Proof. Assume first that $E \setminus W$ is spaceable and let F be an infinite-dimensional closed subspace of E such that

$$F \cap W = \{0\}.$$

Considering an ℓ_∞ -independent sequence $(x_n)_{n=1}^\infty$ of elements of F such that

$$\sum_{n=1}^{\infty} \|x_n\|_E < \infty,$$

let us show that $E \setminus W$ is $[(x_n)_{n=1}^\infty, \mathcal{S}]$ -lineable, for $\mathcal{S} := \ell_\infty$. For this, consider the following linear operator:

$$\mathcal{A}: \ell_\infty \rightarrow F, \quad (t_n)_{n=1}^\infty \mapsto \sum_{n=1}^{\infty} t_n x_n.$$

Since $\sum_{n=1}^{\infty} \|x_n\|_E < \infty$, the operator \mathcal{A} is well-defined. Moreover, due to the fact that $(x_n)_{n=1}^\infty$ is ℓ_∞ -independent, \mathcal{A} is also injective. This allows infer that

$$\mathcal{A}(\ell_\infty) \subseteq F \subseteq (E \setminus W) \cup \{0\},$$

showing that $E \setminus W$ is $[(x_n)_{n=1}^\infty, \mathcal{S}]$ -lineable, as desired.

For the converse, assume that $E \setminus W$ is $[(x_n)_{n=1}^\infty, \mathcal{S}]$ -lineable for some closed subspace \mathcal{S} of ℓ_∞ containing c_0 and some \mathcal{S} -independent sequence $(x_n)_{n=1}^\infty$ of elements of E . Consider the following operator:

$$(3.1) \quad \mathcal{O}: \mathcal{S} \rightarrow E, \quad (t_n)_{n=1}^\infty \mapsto \sum_{n=1}^{\infty} t_n x_n.$$

Given that $E \setminus W$ is $[(x_n)_{n=1}^\infty, \mathcal{S}]$ -lineable, we have:

$$(3.2) \quad \mathcal{O}(\mathcal{S}) \subseteq (E \setminus W) \cup \{0\}.$$

Notice that

$$\dim \mathcal{O}(\mathcal{S}) = \dim \mathcal{S} = \infty \quad (\mathcal{O} \text{ is injective}).$$

Considering $\overline{\text{span}}\{y_n\}_{n=1}^\infty$ where $(y_n)_{n=1}^\infty$ is a regular basic sequence, (W has a non-minimal closed subspace) this allows infer that the set $E \setminus \overline{\text{span}}\{y_n\}_{n=1}^\infty$ has infinite-codimension. That is,

$$\dim(E/\overline{\text{span}}\{y_n\}_{n=1}^\infty) = \infty.$$

Without loss of generality, assume that $(x_n)_{n=1}^\infty$ is such that

$$\sum_{n=1}^{\infty} \|x_n\|_E < \infty$$

and

$$\sum_{n=1}^{\infty} a_n Q(x_n) = 0 \Rightarrow a_n = 0 \text{ whenever } (a_n)_{n=1}^\infty \in \ell_\infty$$

where $Q: E \rightarrow E/\overline{\text{span}}\{y_n\}_{n=1}^\infty$ is the quotient map.

Let:

$$\mathcal{L}: \overline{\text{span}}\{y_n\}_{n=1}^\infty \rightarrow \mathcal{S}, \quad \mathcal{L}(y) = (f_j(y))_{j=1}^\infty,$$

where $f_j: \overline{\text{span}}\{y_n\}_{n=1}^\infty \rightarrow \mathbb{K}$ is such that

$$f_j(y_n) = \begin{cases} 0, & \text{if } n \neq j, \\ 1, & \text{if } n = j. \end{cases}$$

The map \mathcal{L} is well-defined since $(y_n)_{n=1}^\infty$ is regular and $c_0 \subseteq \mathcal{S}$. Furthermore, it is continuous due to the **Closed Graph Theorem**. Regarding the operator \mathcal{O} considered in (3.1), it is also continuous. To see this, it suffices to consider the sequence of continuous mappings $(\mathcal{O}_N)_{N=1}^\infty$ defined by

$$\mathcal{O}_N: \ell_\infty \rightarrow E, \quad \mathcal{O}_N((t_k)_{k=1}^\infty) = \sum_{k=1}^N t_k x_k,$$

and employ the version of the **Banach-Steinhaus** in [33, Theorem 2.8].

Fix $y = \sum_{n=1}^\infty a_n y_n$ in $\overline{\text{span}}\{y_n\}_{n=1}^\infty$. Due to the fact that

$$(\mathcal{O} \circ \mathcal{L})(y) = (\mathcal{O} \circ \mathcal{L})\left(\sum_{n=1}^\infty a_n y_n\right) = \mathcal{O}\left(\sum_{n=1}^\infty a_n \mathcal{L}(y_n)\right) = \mathcal{O}((a_n)_{n=1}^\infty) = \sum_{n=1}^\infty a_n x_n$$

and

$$f_j(y) = f_j\left(\sum_{n=1}^\infty a_n y_n\right) = a_j, \quad \text{for each } j \in \mathbb{N}$$

we obtain

$$(\mathcal{O} \circ \mathcal{L})(y) = \sum_{n=1}^\infty f_n(y) x_n.$$

Since y is arbitrary, the operator $\mathcal{O} \circ \mathcal{L}$ is well-defined and it is determined by sequences $(f_n)_{n=1}^\infty$ and $(x_n)_{n=1}^\infty$. Moreover, according to the proof of Proposition 3.3 in [18], this is an operator compact. Hence, if we consider the identity embedding $\mathcal{I}: \overline{\text{span}}\{y_k\}_{k=1}^\infty \rightarrow E$, and invoke the Lemma 2.4, we conclude that the operator $\mathcal{I} + \mathcal{O} \circ \mathcal{L}$ is such that

$$(\mathcal{I} + \mathcal{O} \circ \mathcal{L})(\overline{\text{span}}\{y_k\}_{k=1}^\infty)$$

is closed.

Furthermore, since:

1. $(\mathcal{O} \circ \mathcal{L})(y) = 0 \Rightarrow \sum_{n=1}^\infty f_n(y) x_n = 0$,
2. $(x_n)_{n=1}^\infty$ is ℓ_∞ -independent,
3. $(f_n(y))_{n=1}^\infty \in \mathcal{S}$, and
4. $\mathcal{S} \subseteq \ell_\infty$,

we have:

$$f_n(y) = 0, \quad \text{for each } n \in \mathbb{N}.$$

This allows us to conclude that

$$y = 0.$$

Hence, $\mathcal{O} \circ \mathcal{L}$ is injective and consequently the operator $\mathcal{I} + \mathcal{O} \circ \mathcal{L}$ is an isomorphism over its range.

We claim that

$$(\mathcal{I} + \mathcal{O} \circ \mathcal{L})(\overline{\text{span}}\{y_n\}_{n=1}^\infty) \cap W = \{0\}.$$

Indeed, let $v \in (\mathcal{I} + \mathcal{O} \circ \mathcal{L})(\overline{\text{span}}\{y_n\}_{n=1}^\infty) \cap W$. Due to the fact that $v \in (\mathcal{I} + \mathcal{O} \circ \mathcal{L})(\overline{\text{span}}\{y_n\}_{n=1}^\infty)$, there exists u in $\overline{\text{span}}\{y_k\}_{k=1}^\infty$ such that

$$v = (\mathcal{I} + \mathcal{O} \circ \mathcal{L})(u).$$

Since

$$v = (\mathcal{I} + \mathcal{O} \circ \mathcal{L})(u) = u + (\mathcal{O} \circ \mathcal{L})(u),$$

we have

$$(\mathcal{O} \circ \mathcal{L})(u) = v - u \in [W + \overline{\text{span}}\{y_n\}_{n=1}^\infty] \subseteq W.$$

That is,

$$(\mathcal{O} \circ \mathcal{L})(u) \in W.$$

On the other hand, due to the fact that

$$(\mathcal{O} \circ \mathcal{L})(u) = \sum_{n=1}^{\infty} f_n(u)x_n \in \mathcal{O}(\mathcal{S}),$$

we can invoke (3.2) and conclude that

$$v - u = (\mathcal{O} \circ \mathcal{L})(u) = 0.$$

Therefore,

$$v = u = 0 \quad (\mathcal{O} \circ \mathcal{L} \text{ is injective}),$$

and consequently,

$$(\mathcal{I} + \mathcal{O} \circ \mathcal{L})(\overline{\text{span}}\{y_n\}_{n=1}^\infty) \cap W = \{0\}$$

as desired.

Finally, since $(\mathcal{I} + \mathcal{O} \circ \mathcal{L})(y_n) = y_n + (\mathcal{O} \circ \mathcal{L})(y_n) = y_n + x_n$, we have that the sequence $(y_n + x_n)_{n=1}^\infty$ is basic, and this allows us to conclude that

$$\dim(\mathcal{I} + \mathcal{O} \circ \mathcal{L})(\overline{\text{span}}\{y_n\}_{n=1}^\infty) = \infty.$$

Therefore, $E \setminus W$ is spaceable, completing the proof of this result. \square

4. SOME CONSEQUENCES

Corollary 4.1. *For a Banach space E , if F be an infinite-dimensional closed subspace of E , then the following conditions are equivalent:*

- (1) $E \setminus F$ is $[(x_n)_{n=1}^\infty, \mathcal{S}]$ -lineable for some closed subspace \mathcal{S} of ℓ_∞ containing c_0 and some \mathcal{S} -independent sequence $(x_n)_{n=1}^\infty$ of elements of E ,
- (2) $E \setminus F$ is spaceable,
- (3) F has infinite codimension.

Corollary 4.2. *For an F -space E , if W be a subspace of E that contains a non-minimal closed subspace, then the following conditions are equivalent:*

- (1) $E \setminus W$ is spaceable,
- (2) $E \setminus W$ is $[(x_n)_{n=1}^\infty, \mathcal{S}]$ -lineable for some closed subspace \mathcal{S} of ℓ_∞ containing c_0 and some \mathcal{S} -independent sequence $(x_n)_{n=1}^\infty$ of elements of E ,
- (3) $E \setminus W$ is (n, \mathfrak{c}) -spaceable for each $n \in \mathbb{N}$.

5. ACKNOWLEDGEMENTS

We would like to thank Professors Dr. Anselmo R. Jr. and Dr. Fernando V. J. C. for their valuable contributions during the development of this paper.

REFERENCES

- [1] M. Aires, G. Botelho, *Spaceability of sets of non-injective maps*, arXiv:2403.19855.
- [2] N.G. Albuquerque, L. Coleta, *Large structures within the class of summing operators*, J. Math. Anal. Appl. **526**, 2 (2023).
- [3] G. Araújo and A. Barbosa, *A general lineability criterion for complements of vector spaces*, Rev. Real Acad. Cienc. Exactas Fis. Nat. Ser. A-Mat. **118** (2024).
- [4] G. Araújo, A. Barbosa, A. Raposo Jr., and G. Ribeiro, *On the spaceability of the set of functions in the Lebesgue space L_p which are not in L_q* , Bull Braz Math Soc, New Series **54**, 44 (2023).
- [5] R.M. Aron, L. Bernal-González, D. Pellegrino and J.B. Seoane-Sepúlveda, *Lineability: the search for linearity in mathematics*, Monographs and Research Notes in Mathematics. CRC Press, Boca Raton, FL, 2016.
- [6] R.M. Aron, V.I. Gurariy and J.B. Seoane-Sepúlveda, *Lineability and spaceability of sets of functions on \mathbb{R}* , Proc. Amer. Math. Soc. **133** (2004), 795–803.
- [7] J. Bastero, ℓ_q -subspaces of stable p -Banach spaces, $0 < p < 1$. Arch. Math. **40** (1983), 538–544.
- [8] S. Banach, *Theorie des Opérations Linéaires*, Warsaw, 1932, reprinted by Chelsea, New York.
- [9] L. Bernal-González and M.O. Cabrera, *Lineability criteria, with applications*, J. Funct. Anal. **266** (2014), 3997–4025.
- [10] L. Bernal-González, J.A. Conejero, M. Murillo-Arcila, J.B. Seoane-Sepúlveda, *[S]-linear and convex structures in function families*, Linear Algebra Appl. **579** (2019), 463–483.
- [11] L. Bernal-González, D.L. Rodríguez-Vidanes, J.B. Seoane-Sepúlveda, H.J. Tag, *New Results in Analysis of Orlicz-Lorentz spaces*. arXiv:2312.13903.
- [12] G. Botelho, V.V. Fávaro, D. Pellegrino and J.B. Seoane-Sepúlveda, *$L_p[0, 1] \setminus \cup_{q>p} L_q[0, 1]$ is spaceable for every $p > 0$* , Linear Algebra Appl. **436** (2012), 2963–2965.
- [13] M. Caballer, S. Dantas, D.L. Rodríguez-Vidanes, *Searching for linear structures in the failure of the Stone-Weierstrass theorem*, preprint. arXiv:2405.06453.
- [14] D. Cariello, J.B. Seoane-Sepúlveda, *Basic sequences and spaceability in ℓ_p spaces*. J. Funct. Anal. **266** (2014), 3797–381
- [15] D. Diniz and A. Raposo Jr, *A note on the geometry of certain classes of linear operators*. Bull. Braz. Math. Soc. (N.S.) **52** (2021), 1073–1080.
- [16] D. Diniz, V.V. Fávaro, D. Pellegrino and A. Raposo Jr, *Spaceability of the sets of surjective and injective operators between sequence spaces*. Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM **114** (2020), no. 4, Paper No. 194, 11 pp.
- [17] L. Drewnowski, *Quasicomplements in F -spaces*, Studia Math. **77** (1984) 373–391.
- [18] L. Drewnowski, *On minimal topological linear spaces and strictly singular operators*, Comment. Math., T. Spoc. in honorem L. Orlics II, 89-106, PWN, Warszawa 1979.
- [19] P.H. Enflo, V.I. Gurariy, J.B. Seoane-Sepúlveda, *Some results and open questions on spaceability in function spaces*, Trans. Amer. Math. Soc. **366** (2) (2014) 611–625
- [20] P. Enflo, *A counterexample to the approximation property in Banach spaces*, Acta Math. **130**(1973), 309–317.
- [21] V.V. Fávaro, D. Pellegrino and P. Rueda, *On the size of the set of unbounded multilinear operators between Banach spaces*, Linear Algebra Appl. **606** (2020), 144–158.
- [22] V.V. Fávaro, D. Pellegrino and D. Tomáz, *Lineability and spaceability: a new approach*, Bull. Braz. Math. Soc. New Ser. **51** (2019), 27–46.
- [23] V.V. Fávaro, D. Pellegrino, A. Raposo Jr., and G. Ribeiro, *General criteria for a stronger notion of lineability*, Proc. Amer. Math. Soc. **152** (2024), 941–954.
- [24] V. Fonf, V.I. Gurariy, V. Kadec *An infinite dimensional subspace of $C[0, 1]$ consisting of nowhere differentiable functions* C. R. Acad. Bulgare Sci., **52** (1999), pp. 13-16
- [25] V.I. Gurariy, L. Quarta, *On lineability of sets of continuous functions*, J. Math. Anal. Appl. **294** (1) (2004) 62–72.
- [26] N. Kalton, *Basic sequences in F -spaces and their applications*, Proc. Edinb. Math. Soc. (2) **19** (1974/1975), 151-167.
- [27] N. Kalton, J. Shapiro, *Bases and basic sequences in F -spaces*, Studia Math **56**, (1976), 47–61
- [28] N. Kalton, *The basic sequence problem*. Studia Math **116** (1995), 167–187.
- [29] N. Kalton, *Quotients of F -spaces*, Glasgow Math. J. **19** (1978), 103–108.

- [30] D. Kitson and R.M. Timoney, *Operator ranges and spaceability*. J. Math. Anal. Appl. **378** (2011), 680–686.
- [31] B. Levine and D. Milman, *On linear sets in space C consisting of functions of bounded variation*, Comm. Inst. Sci. Math. Méc. Univ. Kharkoff [Zapiski Inst. Mat. Mech.] (4) **16** (1940), 102–105 (Russian, with English summary).
- [32] D. Pellegrino and A. Raposo Jr, *Pointwise lineability in sequence spaces*. Indag. Math. (N.S.) **32** (2021), 536–546.
- [33] Rudin, Walter (1991), *Functional Analysis. International Series in Pure and Applied Mathematics*. Vol. 8 (Second ed.). New York, NY: McGraw-Hill Science/Engineering/Math. ISBN 978-0-07-054236-5.
- [34] P. Leonetti, T. Russo, J. Somaglia, *Dense lineability and spaceability in certain subsets of ℓ_∞* , Bull. Lond. Math. Soc. **55** (2023), 1–21.
- [35] D. Papathanasiou, *Dense lineability and algebraability of $\ell_\infty \setminus c_0$* , Proc. Amer. Math. Soc. **150** (2022), 991–996.
- [36] J.H. Shapiro, *On the weak basis theorem in F -spaces*, Canad. J. Math **26** (1974), 1294–1300.

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DA PARAÍBA, 58.051-900 - JOÃO PESSOA, BRAZIL.
Email address: geivison.ribeiro@academico.ufpb.br