THE FERMIONIC ENTANGLEMENT ENTROPY OF CAUSAL DIAMONDS IN TWO-DIMENSIONAL MINKOWSKI SPACE

FELIX FINSTER, MAGDALENA LOTTNER, ALBERT MUCH, AND SIMONE MURRO

JULY 2024

ABSTRACT. The fermionic Rényi entanglement entropy is studied for causal diamonds in two-dimensional Minkowski spacetime. Choosing the quasi-free state describing the Minkowski vacuum with an ultraviolet regularization, a logarithmically enhanced area law is derived.

CONTENTS

1. Introduction	1
2. Physical Preliminaries	3
2.1. The Dirac Field in Two-Dimensional Minkowski Spacetime	3
2.2. The Quantized Dirac Field and its Vacuum State	4
2.3. The Entanglement Entropy of a Causal Diamond	5
3. Schatten-von-Neumann Bounds for Pseudo-Differential Operators	5
3.1. Singular Values and Schatten-von Neumann Classes	6
3.2. Non-Smooth Spectral Functions	6
4. Spectral Analysis of Truncated Pseudo-Differential Operators	8
5. An Area Law for the Causal Diamond	11
5.1. The Diagonal Terms	12
5.2. The Off-Diagonal Terms	13
References	15

1. INTRODUCTION

Entropy quantifies the disorder of a physical system. There are various notions of entropy, like the entropy in classical statistical mechanics as introduced by Boltzmann and Gibbs, the Shannon and Rényi entropies in information theory or the *von Neumann entropy* for quantum systems. In the past years, many studies are devoted to the *entanglement entropy*, being a measure for the quantum correlations between subsystems of a composite quantum system [1, 13]. In the relativistic setting, the *relative entropy* has been studied extensively in connection with modular theory (see for example [12, 10, 17]). In the present paper we restrict attention to the *fermionic* case. Moreover, for simplicity we only consider the *quasi-free* case where the particles do not interact with each other. This makes it possible to express the entanglement entropy in terms of the reduced one-particle density operator [11] (for details in an expository style see the survey paper [6]). This setting has been studied extensively for a free Fermi gas formed of non-relativistic spinless particles [11, 14, 15]. The main



FIGURE 1. A causal diamond.

interest of these studies lies in the derivation of *area laws*, which quantify how the entanglement entropy scales as a function of the size of the spatial region forming the subsystem. More recently, these methods and results were adapted to the relativistic setting of the Dirac equation. In [9] the entanglement entropy for the free Dirac field in a bounded spatial region of Minkowski spacetime is studied. An area law is proven in two limiting cases: that the volume tends to infinity and that the regularization is removed. Moreover, in [8] the geometry of a Schwarzschild black hole is studied. The entanglement entropy of the event horizon is computed to be a prefactor times the number of occupied angular momentum modes. Independently, the entanglement entropy for systems of Dirac spinors has been studied in [3, 4].

In the present paper we study the entanglement entropy of a causal diamond \mathfrak{D} embedded in two-dimensional Minkowski space \mathcal{M} (see Figure 1). The interval $\Lambda := (0, \lambda)$ with $\lambda > 0$ is our spatial subregion. Its boundary consists of the two corners (0, 0) and $(0, \lambda)$ of the causal diamond. In this setting, an area law simply states that the entanglement entropy should be independent of the size λ of the spatial subregion. In order to make this statement precise, as the fermionic state in Minkowski space we choose the vacuum state with an ultraviolet regularization on the scale $\varepsilon > 0$, namely we consider the pseudo-differential operator defined by

$$(\Pi^{(\varepsilon)}\psi)(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{ik(x-y)} e^{-\varepsilon\omega(k)} \left(1 - \frac{1}{\omega(k)} \begin{pmatrix} -k & m\\ m & k \end{pmatrix}\right) \psi(y) \, dy \, dk \,,$$

where $\omega(k) := \sqrt{k^2 + m^2}$ and ε can be interpreted as a semi-classical parameter that shall tend to zero in our asymptotic results. Now, for each $\varkappa > 0$ we introduce the *Rényi entropy function*, which is defined as follows. If $t \notin (0, 1)$ then we set $\eta_{\varkappa}(t) = 0$. For $t \in (0, 1)$ we define

$$\eta_{\varkappa}(t) = \frac{1}{1-\varkappa} \ln \left(t^{\varkappa} + (1-t)^{\varkappa} \right) \quad \text{for } \varkappa \neq 1 ,$$

$$\eta_{1}(t) := \lim_{\varkappa \to 1} \eta_{\varkappa}(t) = -t \ln t - (1-t) \ln(1-t) \quad \text{for } \varkappa = 1 .$$
 (1.1)

Note that η_1 is the familiar von Neumann entropy function. The *Rényi entanglement* entropy of the causal diamond is defined by

$$S_{\varkappa}(\Pi^{(\varepsilon)},\Lambda;\eta_{\varkappa}) := \operatorname{tr}\left(\eta_{\varkappa}(\chi_{\Lambda}\Pi^{(\varepsilon)}\chi_{\Lambda}) - \chi_{\Lambda}\eta_{\varkappa}(\Pi^{(\varepsilon)})\chi_{\Lambda}\right).$$

As we shall see, since Λ is bounded and $\varepsilon > 0$, both operators on the right-hand side are trace class, so the entropy S_{\varkappa} is well-defined. Our main objective is to analyze the asymptotic behavior of the entropy $S_{\varkappa}(\Pi^{(\varepsilon)}, \Lambda; \eta_{\varkappa})$ in the limit $\varepsilon \searrow 0$. This is our main result:

Theorem 1.1. (Entanglement entropy of a causal diamond) For any $\varkappa > 0$, the Rényi entanglement entropy of a causal diamond in the two-dimensional Minkowski satisfies

$$\lim_{\varepsilon \searrow 0} \frac{1}{\ln(1/\varepsilon)} S_{\varkappa}(\Pi^{(\varepsilon)}, \Lambda; \eta_{\varkappa}) = \frac{1}{\pi^2} \int_0^1 \frac{\eta_{\varkappa}(t)}{t(1-t)} dt = \frac{1}{6} \frac{\varkappa + 1}{\varkappa} .$$
(1.2)

In particular, for $\varkappa = 1$, it holds

$$\lim_{\varepsilon \searrow 0} \frac{1}{\ln(1/\varepsilon)} S_{\varkappa}(\Pi^{(\varepsilon)}, \Lambda; \eta) = \frac{1}{3} \,.$$

The fact that the entropy grows in Λ only logarithmically shows that the entanglement entropy does not scale with the volume. With this in mind, the scaling in (1.2) can be understood as a *logarithmically enhanced area law*.

Our method of prove is based on extensions of the methods in [11, 14, 15] to matrixvalued symbols as developed in [8, 9]. Our presentation is self-contained and of expository style.

The paper is organized as follows. Section 2 provides the physical preliminaries on the entanglement entropy of the Minkowski vacuum state restricted to a causal diamond. In Section 3, we collect some mathematical background of our analysis including some abstract results on on Schatten norms and compact operators on Hilbert spaces. Our main mathematical results are contained in Section 4, devoted to an asymptotic analysis of truncated pseudo-differential operators. The main result of this paper is then proved in Section 5.

2. Physical Preliminaries

2.1. The Dirac Field in Two-Dimensional Minkowski Spacetime. Consider the two-dimensional Minkowski space $\mathcal{M} := (\mathbb{R}^2, g)$ endowed with the line element

$$ds^2 = g_{ij} dx^i dx^j = dt^2 - dx^2$$
,

and denote by $S\mathcal{M} = \mathcal{M} \times \mathbb{C}^2$ the trivial spinor bundle. As customary, we equip the spinor bundle with the *spin scalar product*, namely the indefinite inner product

$$\prec \psi | \phi \succ = \langle \psi, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \phi \rangle_{\mathbb{C}^2} ,$$

where $\langle ., . \rangle_{\mathbb{C}^2}$ is the canonical scalar product on \mathbb{C}^2 . The *Dirac operator* is the first order differential operator acting on sections of the spinor bundle defined by

$$\mathcal{D} := i\gamma^j \partial_j \,, \tag{2.1}$$

where the gamma matrices γ^{j} are given, in the chiral representation, by

$$\gamma^0 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$
, $\gamma^1 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$.

Fixing a positive mass $m \in \mathbb{R}$, the massive Dirac equation reads

$$(\mathcal{D} - m)\psi = 0. \tag{2.2}$$

The Dirac operator can be rewritten as a symmetric hyperbolic system and its Cauchy problem is well-posed (for details see for example [7, Chapter 13]). Moreover, taking

smooth and compactly supported initial data on a Cauchy surface \mathcal{N} , a Dirac solution lies in the class $C^{\infty}_{\rm sc}(\mathcal{M}, S\mathcal{M})$ of smooth spinors with spatially compact support. On solutions ψ, ϕ in this class, it is well-defined the (positive definite) scalar product

$$(\psi|\phi)_{\mathcal{M}} := \int_{\mathcal{N}} \prec \psi |\psi\phi \succ|_q \, d\mu_{\mathcal{N}}(q) \,,$$

where $\psi = \gamma^j \nu_j$ denotes Clifford multiplication by the future-directed unit normal ν , and $d\mu_{\mathcal{N}}$ is the volume measure of the induced Riemannian metric on \mathcal{N} (thus for the above ray $\mathcal{N} = \{(\alpha x, x) \text{ with } x > 0\}$, the measure $d\mu_{\mathcal{N}} = \sqrt{1 - \alpha^2} dx$ is a multiple of the Lebesgue measure). Due to current conservation, this scalar product is independent of the choice of \mathcal{N} . Forming the completion, we obtain the Hilbert space $(\mathcal{H}_{\mathcal{M}}, (.|.)_{\mathcal{M}})$, referred to as the *solution space* of the Dirac equation. For convenience, we always choose \mathcal{N} as the Cauchy surface $\{t = 0\}$, so that

$$(\psi|\phi)_{\mathcal{M}} = \int_{-\infty}^{\infty} \langle \psi|\gamma^0 \phi \rangle|_{(0,x)} dx$$

2.2. The Quantized Dirac Field and its Vacuum State. The quantized Dirac field can be described in a two step-procedure. First, one assigns to a classical physical system (described by the solutions space) a unital *-algebra \mathfrak{A} , whose elements are interpreted as observables of the system at hand. Then, one determines the admissible physical states of the system by identifying a suitable subclass of the linear, positive and normalized functionals $\omega : \mathfrak{A} \to \mathbb{C}$. Once that a state is specified, the Gelfand-Naimark-Segal (GNS) construction guarantees the existence of a representation of the quantum field algebra as (in general, unbounded) operators defined in a common dense subspace of some Hilbert space.

We here restrict attention to a *quasi-free* Dirac state, which is fully characterized by its two-point distribution. Then, as shown in [5, Section 4], the above procedure boils down to constructing suitable self-adjoint operators on the Dirac solution space. In particular, the *vacuum state* can be represented by the projection operator onto the space of negative frequencies solutions as follows. We rewrite the Dirac equation in the Hamiltonian form

$$i\partial_t \psi = H\psi$$
 with $H = -i\gamma^0\gamma^1\partial_x + m\gamma^0 = \begin{pmatrix} i\partial_x & m \\ m & -i\partial_x \end{pmatrix}$.

Taking the spatial Fourier transform

$$\hat{\psi}(k) = \int_{-\infty}^{\infty} \psi(x) \ e^{-ikx} \ dx , \qquad \psi(x) = \int_{-\infty}^{\infty} \frac{dk}{2\pi} \ \hat{\psi}(k) \ e^{ikx} ,$$

the Hamiltonian becomes the multiplication operator

$$\hat{H}(k) = \begin{pmatrix} -k & m \\ m & k \end{pmatrix} \,,$$

which can be diagonalized by

$$\hat{H}(k) = \omega E_+ - \omega E_- ,$$

with $\omega(k) := \sqrt{k^2 + m^2}$ and

$$E_{\pm}(k) = \frac{1}{2} \pm \frac{1}{2\omega(k)} \begin{pmatrix} -k & m \\ m & k \end{pmatrix} .$$

The projectors onto the space of negative frequencies solutions is

$$(\Pi\psi)(x) = \int_{-\infty}^{\infty} \Pi(x,y) \,\psi(y) \,dy$$

with integral kernel

$$\Pi(x,y) = \frac{1}{2} \int_{-\infty}^{\infty} \frac{dk}{2\pi} \left(\mathbbm{1}_{\mathbb{C}^2} - \frac{1}{\omega(k)} \begin{pmatrix} -k & m \\ m & k \end{pmatrix} \right) e^{ik(x-y)}$$

For future convenience, we define the *regularized projection operator* onto the negative frequencies solutions as the integral operator

$$(\Pi^{(\varepsilon)}\psi)(x) = \int_{-\infty}^{\infty} \Pi^{(\varepsilon)}(x,y) \,\psi(y) \,dy \,,$$

where the integral kernel is given by

$$\Pi^{(\varepsilon)}(x,y) = \frac{1}{2} \int_{-\infty}^{\infty} \frac{dk}{2\pi} e^{-\varepsilon\omega(k)} \left(\mathbbm{1}_{\mathbb{C}^2} - \frac{1}{\omega(k)} \begin{pmatrix} -k & m \\ m & k \end{pmatrix} \right) e^{ik(x-y)} \,.$$

This operator will play a pivotal role for computing the entanglement entropy.

2.3. The Entanglement Entropy of a Causal Diamond. In this section we shall define the Rényi entanglement entropy of a causal diamond \mathcal{D} embedded in the two-dimensional Minkowski space \mathcal{M} . For the sake of completeness, let us recall that a causal diamond is a two-dimensional space isometric to the subset of two-dimensional Minkowski space

$$\mathcal{D} = \left\{ (t, x) \in \mathcal{M} \quad \text{with} \quad 0 < x < \lambda \text{ and } |t| < \min(x, \lambda - x) \right\};$$

where $\lambda > 0$ is an arbitrary, but fixed parameter (see Figure 1). Then the inclusions

$$\mathfrak{D} \subset \mathcal{M} \quad \text{and} \quad S\mathfrak{D} = \mathfrak{D} \times \mathbb{C}^2 \subset \mathcal{M} \times \mathbb{C}^2 = S\mathcal{M}$$

are clearly isometries, and the Dirac operator and the Dirac equation are again given by (2.1) and (2.2). Adopting the notation of the previous section, we denote the subspace of solutions in \mathcal{D} by $\mathcal{H}_{\mathcal{D}}$. As shown in[6, Appendix A], the Rényi entanglement entropy of \mathcal{D} can be expressed as

$$S_{\varkappa}(\Pi^{(\varepsilon)},\Lambda,\eta_{\varkappa}) = \operatorname{tr}\eta_{\varkappa}(\pi_{\mathfrak{D}}\Pi^{(\varepsilon)}\pi_{\mathfrak{D}}) - \operatorname{tr}\eta_{\varkappa}(\Pi^{(\varepsilon)}),$$

where $\pi_{\mathfrak{D}} : \mathcal{H} \to \mathcal{H}_{\mathfrak{D}}$ is the orthogonal projection operator. This projection operator can be represented more concretely as the multiplication operator by a characteristic function acting on the wave functions on the Cauchy surface $\{t = 0\}$, i.e.

$$(\pi_{\mathcal{D}}\psi)(0,x) = \chi_{\Lambda}(x) \psi(0,x)$$

where $\Lambda = (0, \lambda)$ for the causal diamond (see again Figure 1).

3. Schatten-von-Neumann Bounds for Pseudo-Differential Operators

In this section we state some basic definitions and results on singular values and Schatten-von Neumann classes. For more details we refer to [2, Chapter 11] for more details. 3.1. Singular Values and Schatten-von Neumann Classes. For a compact operator A in a separable Hilbert space \mathcal{H} we denote by $s_k(A)$, $k = 1, 2, \ldots$, its singular values (defined as the eigenvalues of the self-adjoint compact operator |A|) labelled in non-increasing order counting multiplicities. For the sum A+B the following inequality holds,

$$s_{2k}(A+B) \le s_{2k-1}(A+B) \le s_k(A) + s_k(B).$$
(3.1)

We say that A belongs to the Schatten-von Neumann class \mathbf{S}_p , p > 0, if

$$||A||_p := \left(\operatorname{tr} |A|^p\right)^{1/p} = \left(\sum_{n=1}^{\infty} s_k(A)^p\right)^{1/p}$$

is finite. The functional $||A||_p$ defines a norm if $p \ge 1$ and a quasi-norm if 0 . $With this (quasi-)norm, the class <math>\mathbf{S}_p$ is a complete space. Note that for p = 1 this coincides with the trace norm. Moreover, by $||.||_{\infty}$ we denote the ordinary operator norm. For 0 the quasi-norm is actually a*p* $-norm, that is, it satisfies the following "triangle inequality" for all <math>A, B \in \mathbf{S}_p$:

$$||A + B||_p^p \le ||A||_p^p + ||B||_p^p.$$
(3.2)

This inequality is used systematically in what follows. Using (3.2) we may also derive an estimate for block operators of the form $A = (a_{ij})_{1 \le i,j \le n}$ with $a_{ij} \in \mathbf{S}_p$ for any $1 \le i, j \le n$:

$$\|A\|_{p}^{p} \leq \sum_{i,j=1}^{n} \|(\delta_{il}\delta_{jm}a_{ij})_{lm}\|_{p}^{p} = \sum_{i,j=1}^{n} \|a_{ij}\|_{p}^{p}, \qquad (3.3)$$

where in the last step we used that the singular values of the block operator $(\delta_{il}\delta_{jm}a_{ij})_{lm}$ coincide with the ones of the operator a_{ij} . We point out a useful estimate for individual eigenvalues for operators in \mathbf{S}_p :

$$s_k(A) \le k^{-\frac{1}{p}} \|A\|_p, \ k = 1, 2, \dots$$
 (3.4)

In [2, p. 262] it is shown that the norms $\|\cdot\|_{\mathbf{S}_p}$ also fulfill a Hölder-like inequality, meaning that for any $0 and <math>0 < p_1, p_2 \le \infty$ such that $p^{-1} = p_1^{-1} + p_2^{-1}$ and $A \in \mathbf{S}_{p_1}, B \in \mathbf{S}_{p_2}$, the operator $AB \in \mathbf{S}_p$ with

$$||AB||_p \le ||A||_{p_1} ||B||_{p_2} , \qquad (3.5)$$

where by $\|.\|_{\infty}$ we mean the ordinary operator norm. Moreover, as explained in [2, p. 254], for any two $0 < p_1 < p_2 \le \infty$, we have $\mathbf{S}_{p_1} \subset \mathbf{S}_{p_2}$ and for any $A \in \mathbf{S}_{p_1}$

$$||A||_{p_2} \le ||A||_{p_1}$$
.

We refer to [2, Chapter 11] for more details on singular values.

3.2. Non-Smooth Spectral Functions. We now state a result for compact operators in an arbitrary separable Hilbert space \mathcal{H} . Let A be a symmetric bounded operator on \mathcal{H} and P an orthogonal projection operator on \mathcal{H} . Given a continuous function $f \in C(\mathbb{R})$ define the operator

$$D(A, P; f) := Pf(PAP)P - Pf(A)P.$$

In what follows it is convenient to require that the function f satisfies the following condition.

Condition 3.1. The function $f \in C^2(\mathbb{R} \setminus \{t_0\}) \cap C(\mathbb{R})$ satisfies the bound $\|f\|_{2} := \max_{0 \le k \le 2} \sup_{t \ne t_{0}} |f^{(k)}(t)| |t - t_{0}|^{-\gamma + k} < \infty$

for some $\gamma \in (0,1]$ and is supported on the interval $(t_0 - R, t_0 + R)$ with some finite R > 0.

Example 3.2. Consider the functions η_{\varkappa} defined in (1.1) and set $\mathcal{T} = \{0, 1\}$. Then, in the neighborhood of every $t_i \in \mathcal{T}$, there exist positive constants γ and $c_k > 0$ with k = 0, 1, 2 such that

$$\eta_{\varkappa}^{(k)}(t)| \le c_k |t - t_i|^{\gamma - k}$$

As shown in [8, Lemma D.1], the value of γ depends on \varkappa as follows,

$$\begin{cases} \gamma \leq \min\{1, \varkappa\} & \text{for } \varkappa \neq 1 \\ \gamma < 1 & \text{for } \varkappa = 1 \end{cases}.$$

Notice that, using a partition of unity $(\psi_k)_{0 \le k \le 1}$ such that the support of each ψ_k only contains exactly one the elements in \mathcal{T} , then each $\eta_{\varkappa}\psi_k$ satisfies Condition 3.1.

The next proposition follows from a more general result proven in [16, Theorem 2.4]; see also [14, Proposition 2.2].

Proposition 3.3. Suppose that f satisfies Condition 3.1 for some $\gamma \in (0,1]$ and some $t_0 \in \mathbb{R}$ with R > 0. Let $q \in (\frac{1}{2}, 1]$ and assume that $\sigma < \min(2 - q^{-1}, \gamma)$. Let A, Bbe two bounded self-adjoint operators and assume that $|A - B|^{\sigma} \in \mathbf{S}_p$. Then

$$\|f(A) - f(B)\|_p \lesssim \|f\|_n R^{\gamma - \sigma} \|A - B\|_p$$

with a positive implicit constant independent of A, B, f and R. In particular, for an orthogonal projection P such that $PA(I - P) \in \mathbf{S}_{\sigma q}$ it holds

$$\|D(A,P;f)\|_q \lesssim \|f\|_2 R^{\gamma-\sigma} \|PA(I-P)\|_{\sigma q}^{\sigma}, \qquad (3.6)$$

with a positive implicit constant independent of the operators A, P, the function f, and the parameter R.

In order to estimate the Schatten norm on the right hand side of (3.6), we will sometimes use estimates of the Schatten norm of the operator product PA(1-P).

Lemma 3.4. Let q > 0 and A, B bounded operators, then

(i) If $A \in \mathbf{S}_q$, then also $A^* \in \mathbf{S}_q$ and

$$\|A\|_{q} = \|A^{*}\|_{q}$$
(ii) If the $BA(1-B) \in \mathbf{S}_{q}$, then $[A, B] \in \mathbf{S}_{q}$ and
 $\|[A, B]\|_{q} \leq \|BA(1-B)\|_{q}$
(iii) If B is a projection operator and $[A, B] \in \mathbf{S}_{q}$, then

(iii) If B is a projection operator and
$$[A, B] \in \mathbf{S}_{q}$$
, then also $BA(1-B) \in \mathbf{S}_{q}$ and

$$\left\|BA(1-B)\right\|_{q} \le \left\|[A,B]\right\|_{q}$$

Proof. (i): This follows from the fact that singular values (and therefore the q-norm) are invariant under Hermitian conjugation, which in turn can be understood from the following argument. Note that for any $A \in \mathbf{S}_q$ we can write $||A||_q = (\sum_i \sqrt{\lambda_i}^q)^{1/q}$ where the λ_i are the eigenvalues of A^*A (note that since A is compact, there are countably many). Moreover, for any eigenvector ψ of A^*A corresponding to a nonzero eigenvalue, the vector $A\psi$ is an eigenvector of AA^* with the same eigenvalue, so we see that $||A^*||_q \leq ||A||_q$ (as ψ might lie in the kernel of A). Then, symmetry yields the equality.

(ii): We apply the triangle inequality,

$$\begin{split} \|[A,B]\|_q^q &= \|AB - BAB + BAB - BA\|_q^q \\ &= \|(1-B)AB - BA(1-B)\|_q^q \le 2\|BA(1-B)\|_q^q \end{split}$$

where we also used that

$$\left((1-B)AB\right)^* = BA(1-B)\,,$$

together with (i).

(iii): Since B is a projection operator, our claim follows immediately from the estimate

$$||BA - BAB||_q = ||B(BA - AB)||_q \le ||B||_{\infty} ||[A, B]||_q \le ||[A, B]||_q.$$

4. Spectral Analysis of Truncated Pseudo-Differential Operators

Using the results from the previous section, we are now in the position to perform an asymptotic analysis of truncated pseudo-differential operator. To this end, consider a pseudo-differential operator defined as usual by

$$\left(\operatorname{Op}_{\alpha}(\mathcal{A})\psi\right)(x) = \frac{\alpha}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{i\alpha\xi(x-y)} \mathcal{A}(\xi)\psi(y) \, dy \, d\xi$$

where \mathcal{A} is a 2 × 2 matrix-valued symbol and α is a (strictly) positive constant. A *truncated* pseudo-differential operator is then obtained multiplying $\text{Op}_{\alpha}(\mathcal{A})$ by the indicator function χ_{Λ} of $\Lambda \subset \mathbb{R}$

$$\chi_{\Lambda} \operatorname{Op}_{\alpha}(\mathcal{A}) \chi_{\Lambda} \,$$

ad we introduce the *entropic difference operator* by

$$D_{\alpha}(\mathcal{A},\Lambda;f) := f(\chi_{\Lambda} \operatorname{Op}_{\alpha}(\mathcal{A})\chi_{\Lambda}) - \chi_{\Lambda} f(\operatorname{Op}_{\alpha}(\mathcal{A}))\chi_{\Lambda}.$$

Through this section we assume that the function f satisfies Condition 3.1. Inspired by [9, Lemma 5.6], this is our first result.

Lemma 4.1. Suppose that f satisfies Condition 3.1 for some $\gamma \in (0, 1]$. Let $q \in (\frac{1}{2}, 1)$ and assume that $\sigma < \min(2 - q^{-1}, \gamma)$. Finally let be $\mathcal{A}_{\alpha}^{(1)}$ and $\mathcal{A}_{\alpha}^{(2)}$ two families of symbols satisfying the conditions

$$\sup_{\xi \in \mathbb{R}^d} \left| \mathcal{A}_{\alpha}^{(1)}(\xi) - \mathcal{A}_{\alpha}^{(2)}(\xi) \right| \to 0, \qquad \alpha \to \infty$$
(4.1)

$$\|\chi_{\Lambda}\mathcal{A}_{\alpha}^{(j)}(1-\chi_{\Lambda})\|_{\sigma q}^{\sigma q} \lesssim g(\alpha) , \qquad \text{for each } j=1,2 , \qquad (4.2)$$

for some $q < \gamma$ and some positive function g. Then it holds

$$\lim_{\alpha \to \infty} \frac{1}{g(\alpha)} \left\| D_{\alpha}(\mathcal{A}_{\alpha}^{(1)}, \Lambda; f) - D_{\alpha}(\mathcal{A}_{\alpha}^{(2)}, \Lambda; f) \right\|_{1} = 0.$$

Proof. For ease of notation, throughout this proof we denote

$$D_{\alpha}(\mathcal{A}_{\alpha}^{(j)}) \equiv D_{\alpha}(\mathcal{A}_{\alpha}^{(j)}, \Lambda; f), \qquad j = 1, 2.$$

Take $0 < \delta < 1$ arbitrary. For any $\alpha > 0$ we define

$$N \equiv N(\alpha) := \left\lceil g(\alpha) \delta^{\frac{q}{q-1}} \right\rceil$$

Now rewrite

$$\left\| D_{\alpha}(\mathcal{A}_{\alpha}^{(1)}) - D_{\alpha}(\mathcal{A}_{\alpha}^{(2)}) \right\|_{1} = \sum_{k=1}^{\infty} s_{k} \left(D_{\alpha}(\mathcal{A}_{\alpha}^{(1)}) - D_{\alpha}(\mathcal{A}_{\alpha}^{(2)}) \right) = Z_{1}(N) + Z_{2}(N) ,$$

where Z_1 and Z_2 involve the small respectively large singular values,

$$Z_1(N) := \sum_{k=1}^{2N} s_k \left(D_\alpha(\mathcal{A}_\alpha^{(1)}) - D_\alpha(\mathcal{A}_\alpha^{(2)}) \right)$$
$$Z_2(N) := \sum_{k=2N+1}^{\infty} s_k \left(D_\alpha(\mathcal{A}_\alpha^{(1)}) - D_\alpha(\mathcal{A}_\alpha^{(2)}) \right).$$

We now estimate Z_1 and Z_2 separately. For the estimate of Z_1 , we use that

$$s_k \left(D_\alpha(\mathcal{A}_\alpha^{(1)}) - D_\alpha(\mathcal{A}_\alpha^{(2)}) \right) \le \| D_\alpha(\mathcal{A}_\alpha^{(1)}) - D_\alpha(\mathcal{A}_\alpha^{(2)}) \|_{\infty} .$$

As shown in [9, Lemma 5.5], since $\mathcal{A}_{\alpha}^{(1)}$ and $\mathcal{A}_{\alpha}^{(2)}$ are two families of uniformly bounded self-adjoint operators satisfying $\|\mathcal{A}_{\alpha}^{(1)} - \mathcal{A}_{\alpha}^{(2)}\|_{\infty} \to 0$ as $\alpha \to \infty$, then, for any function $g \in C(\mathbb{R})$ we have

$$\|g(\mathcal{A}_{\alpha}^{(1)}) - g(\mathcal{A}_{\alpha}^{(2)})\|_{\infty} \to 0 \quad \text{as } \alpha \to \infty.$$

Therefore, combining this observation with Example 3.2, it is clear that there exists $\tilde{\alpha}(\delta) \equiv \tilde{\alpha} > 0$ such that for any $\alpha > \tilde{\alpha}$,

$$\left\| \eta_{\varkappa} \left(\chi_{\Lambda} \operatorname{Op}_{\alpha}(\mathcal{A}_{\alpha}^{(1)}) \chi_{\Lambda} \right) - \eta_{\varkappa} \left(\chi_{\Lambda} \operatorname{Op}_{\alpha}(\mathcal{A}_{\alpha}^{(2)}) \chi_{\Lambda} \right) \right\|_{\infty} \leq \delta^{\frac{1}{1-q}} \quad \text{and} \\ \left\| \chi_{\Lambda} \left(\eta_{\varkappa} \left(\operatorname{Op}_{\alpha}(\mathcal{A}_{\alpha}^{(1)}) \right) - \eta_{\varkappa} \left(\operatorname{Op}_{\alpha}(\mathcal{A}_{\alpha}^{(2)}) \right) \right) \chi_{\Lambda} \right\|_{\infty} \leq \delta^{\frac{1}{1-q}} .$$

Thus for any $\alpha > \tilde{\alpha}$ we obtain

$$Z_1(N) \le 2N \|D_{\alpha}(\mathcal{A}_{\alpha}^{(1)}) - D_{\alpha}(\mathcal{A}_{\alpha}^{(2)})\|_{\infty} \le 4N\delta^{\frac{1}{1-q}} \le 4g(\alpha)(\delta + \delta^{\frac{1}{1-q}}) \le 8g(\alpha)\delta.$$

In order to estimate Z_2 , we use (3.1) and (3.4) to obtain

$$Z_2(N) \le 2\sum_{k=N}^{\infty} s_k \left(D_\alpha(\mathcal{A}_\alpha^{(1)}) \right) + 2\sum_{k=N}^{\infty} s_k \left(D_\alpha(\mathcal{A}_\alpha^{(2)}) \right)$$
$$\le 2 \left(\left\| D_\alpha(\mathcal{A}_\alpha^{(1)}) \right\|_q + \left\| D_\alpha(\mathcal{A}_\alpha^{(2)}) \right\|_q \right) \sum_{k=N}^{\infty} k^{-1/q}$$

We now notice that, applying Proposition 3.3 to $A = Op_{\alpha}(\mathcal{A})$ and $P = \chi_{\Lambda}$ we obtain

$$\|D_{\alpha}(\mathcal{A},\chi_{\Lambda};f)\|_{q} \lesssim \|f\|_{2} R^{\gamma-\sigma} \|\chi_{\Lambda} \operatorname{Op}_{\alpha}(\mathcal{A})(\mathbb{1}-\chi_{\Lambda})\|_{\sigma q}^{\sigma}$$

with a positive implicit constant independent of \mathcal{A}, Λ , the function f, and the parameter R. Combining the latter estimate with condition (4.2), it follows that

$$\|D_{\alpha}(\mathcal{A}_{\alpha}^{(j)})\|_{q}^{q} \lesssim g(\alpha) \qquad \mathbf{j}=1,2.$$

Consequently, $Z_2(N)$ may be estimated by

$$Z_2(N) \lesssim g(\alpha)^{1/q} \sum_{k=N}^{\infty} k^{-1/q} \leq g(\alpha)^{1/q} \int_{N-1}^{\infty} k^{-1/q} dk \leq g(\alpha)^{1/q} \int_{g(\alpha)\delta^{\frac{q}{q-1}}}^{\infty} k^{-1/q} dk$$
$$\lesssim g(\alpha)^{1/q} \left(g(\alpha)\delta^{\frac{q}{q-1}} \right)^{1-1/q} = g(\alpha)\delta .$$

In summary, for any $\alpha > \tilde{\alpha}(\delta)$ we obtain the estimate

$$\|D_{\alpha}(\mathcal{A}_{\alpha}^{(1)}) - D_{\alpha}(\mathcal{A}_{\alpha}^{(2)})\|_{1} \lesssim g(\alpha)\delta$$

which leads to

$$\lim_{\alpha \to \infty} \frac{1}{g(\alpha)} \left\| D_{\alpha}(\mathcal{A}_{\alpha}^{(1)}, \Lambda; f) - D_{\alpha}(\mathcal{A}_{\alpha}^{(2)}, \Lambda; f) \right\|_{1} \leq \delta.$$

Since $\delta \in (0, 1)$ is arbitrary, this completes the proof.

We conclude this section by considering scalar symbols a which satisfy the following condition.

Condition 4.2. Consider $a \in C^{\infty}(\mathbb{R})$ for which there exist positive continuous functions v and τ such that v is bounded and

$$|\nabla_{\boldsymbol{\xi}}^{n} a(\boldsymbol{\xi})| \lesssim \tau(\boldsymbol{\xi})^{-n} v(\boldsymbol{\xi}), \ n = 0, 1, \dots, \quad \boldsymbol{\xi} \in \mathbb{R}^{d},$$
(4.3)

with constants independent of $\boldsymbol{\xi}$. We call τ the scale (function) and v the amplitude (function). The scale τ is assumed to be globally Lipschitz with Lipschitz constant $\nu < 1$, that is,

$$|\tau(\boldsymbol{\xi}) - \tau(\boldsymbol{\eta})| \le \nu |\boldsymbol{\xi} - \boldsymbol{\eta}|, \quad \text{for all } \boldsymbol{\xi}, \boldsymbol{\eta} \in \mathbb{R}^d.$$
(4.4)

Moreover, there shall also exist constants c, C > 0 such that the amplitude v satisfies

$$c < \frac{v(\boldsymbol{\xi})}{v(\boldsymbol{\eta})} < C \quad \text{for all } \boldsymbol{\eta} \in B(\boldsymbol{\xi}, \tau(\boldsymbol{\xi})),$$

$$(4.5)$$

with c and C independent of $\boldsymbol{\xi}$ and $\boldsymbol{\eta}$. It is useful to think of v and τ as (functional) parameters. They, in turn, may depend on other parameters (e.g. numerical parameters like α).

Given functions ν and τ and numbers $\sigma > 0, \lambda \in \mathbb{R}$ we denote

$$V_{\sigma,\lambda}(v,\tau) := \int \frac{v(\xi)^{\sigma}}{\tau(\xi)^{\lambda}} d\xi$$

The next result is a special case of [14, Lemma 3.4].

Proposition 4.3. Let $I \subset \mathbb{R}$ be a bounded interval and let the functions τ and v be as described above. Suppose that the symbol a satisfies (4.3), and that the conditions

$$au_{\mathrm{inf}} := \inf_{\boldsymbol{\xi} \in \mathbb{R}^d} au(\boldsymbol{\xi}) > 0 , \qquad lpha au_{\mathrm{inf}} \gtrsim 1 .$$

hold. Then for any $\sigma \in (0,1]$ we have

$$\|[\operatorname{Op}_{\alpha}(a), \chi_{I}]\|_{\sigma}^{\sigma} \lesssim V_{\sigma,1}(v, \tau) ,$$

This bound is uniform in the symbols a satisfying Condition 4.2 with the same implicit constants in (4.3), ν in (4.4) and c, C in (4.5) (not necessarily the same functions τ and v).

Using this proposition we can prove the following result.

Lemma 4.4. Let a_{α} be a symbol such that

$$\left|\partial^{l} a_{\alpha}\right| \leq C_{l} v(\xi) \tau(\xi)^{-l}, \quad \text{for any } l \in \mathbb{N}.$$

with

$$v(\xi) := e^{-\frac{l_0}{\alpha}|\xi|}, \qquad \tau(\xi) := \frac{1}{2} \left(\frac{l_0}{\alpha} + \frac{1}{|\xi| + m}\right)^{-1}.$$
(4.6)

Let $I \subseteq \mathbb{R}$ be a bounded interval, $l \in \mathbb{N}$ and $q := l^{-1}$. Then it follows

$$\left\|\chi_I \operatorname{Op}_{\alpha}(a_{\alpha}) (1-\chi_I)\right\|_q^q \lesssim \log \alpha$$

with an implicit constant only depending on q.

Proof. First of all denote

$$a^{(\alpha)}(\xi) := a_{\alpha}(\alpha\xi)$$

Without loss of generality we may assume that $I = (0, \lambda)$ for some $\lambda > 0$. First of all we rescale in momentum space

$$\chi_I \operatorname{Op}_{\alpha}(a^{(\alpha)}) (1 - \chi_I) = \chi_I \operatorname{Op}_1(a_{\alpha}) (1 - \chi_I) .$$

We want to make use of Proposition 4.3. To this end note that of v and τ from (4.6) satisfy the required conditions as

$$\left|\partial_{\xi}\tau(\xi)\right| \leq \frac{1}{2} \left(1 + \frac{l_0}{\alpha}m\right)^{-2} < \frac{1}{2}, \quad \text{for any } \xi \neq 0,$$

implying (4.4) with $\nu = \frac{1}{2}$, if $\boldsymbol{\xi}$ and $\boldsymbol{\eta}$ have the same sign. If they have different signs, we obtain

$$|\tau(\boldsymbol{\xi}) - \tau(\boldsymbol{\eta})| \le |\tau(\boldsymbol{\xi}) - \tau(\mathbf{0})| + |\tau(\mathbf{0}) - \tau(\boldsymbol{\eta})| \le \frac{1}{2} (|\boldsymbol{\xi}| + |\boldsymbol{\eta}|) = \frac{1}{2} |\boldsymbol{\xi} - \boldsymbol{\eta}|,$$

also proving (4.4) Moreover, (4.5) is fulfilled as for any $\tilde{\xi} \in B(\xi, \tau(\xi))$,

$$e^{-\frac{1}{2}} \le \exp\left(-\frac{l_0}{2\alpha}\left(\frac{l_0}{\alpha} + \frac{1}{|\xi| + m}\right)^{-1}\right) \le \frac{v(\tilde{\xi})}{v(\xi)} \le \exp\left(\frac{l_0}{2\alpha}\frac{1}{2}\left(\frac{l_0}{\alpha} + \frac{1}{|\xi| + m}\right)^{-1}\right) \le e^{\frac{1}{2}}.$$

Finally, a lower bound for τ is given by

$$au(\xi) \ge \frac{1}{2} \left(\frac{l_0}{\alpha} + \frac{1}{m} \right)^{-1} \ge \frac{1}{2} \left(l_0 + \frac{1}{m} \right)^{-1} \gtrsim 1,$$

where we assumed that $\alpha \geq 1$. Then, by Proposition 4.3 we conclude that

$$\begin{split} \left\| \left[\operatorname{Op}_{1}(a_{\alpha}), \chi_{I} \right] \right\|_{q}^{q} &\lesssim V_{q,1}(v, \tau) = \int_{-\infty}^{\infty} e^{-\frac{l_{0}}{\alpha}q|\xi|} \left(\frac{l_{0}}{\alpha} + \frac{1}{|\xi| + m} \right) d\xi \\ &\lesssim q^{-1} + 1 + \int_{m}^{\infty} \frac{e^{-q\frac{l_{0}}{\alpha}\xi}}{\xi} d\xi = q^{-1} + 1 + \int_{l_{0}/\alpha}^{\infty} \frac{e^{-qm\xi}}{\xi} d\xi \end{split}$$

with implicit constants independent of α (they might depend on λ though). Substituting $\frac{l_0}{\alpha} = \varepsilon$ and using l'Hospital's rule we see that

$$\int_{l_0/\alpha}^{\infty} \frac{e^{-qm\xi}}{\xi} d\xi \simeq \log \alpha \,.$$

Together with Lemma 3.4 this completes the proof.

5. AN AREA LAW FOR THE CAUSAL DIAMOND

This section is the core of the paper. Our goal is to study the asymptotic behavior of the operator

$$S_{\varkappa}(\Pi^{(\varepsilon)},\Lambda;\eta_{\varkappa}) := \operatorname{tr}\left(\eta_{\varkappa}(\chi_{\Lambda}\Pi^{(\varepsilon)}\chi_{\Lambda}) - \chi_{\Lambda}\eta_{\varkappa}(\Pi^{(\varepsilon)})\chi_{\Lambda}\right) \quad \text{as } \varepsilon \searrow 0$$

By construction, the regularized one-particle density operator is the pseudo-differential operator $\Pi^{(\varepsilon)} = \operatorname{Op}_1(\mathcal{A}_{\varepsilon})$ with symbol given by

$$\mathcal{A}_{\varepsilon} = e^{-\varepsilon\omega(k)} \left(1 - \frac{1}{\omega(k)} \begin{pmatrix} -k & m \\ m & k \end{pmatrix} \right).$$

To connect our analysis with the results obtained in the previous sections, we introduce the parameter $\alpha := \varepsilon^{-1}$ and perform the rescaling $\xi = \alpha k$, obtaining $\Pi^{(\varepsilon)} = \operatorname{Op}_{\alpha}(\mathcal{A}_{\alpha})$ with symbols

$$\mathcal{A}_{\alpha}(\xi) := \frac{1}{2} e^{-\sqrt{\xi^2 + (m/\alpha)^2}} \left(\mathbb{1}_{\mathbb{C}^2} - \frac{1}{\sqrt{\xi^2 + (m/\alpha)^2}} \begin{pmatrix} -\xi & m/\alpha \\ m/\alpha & \xi \end{pmatrix} \right).$$

Now we are interested in the asymptotics for large α . In particular, the Rényi entanglement entropy S_{\varkappa} can be rewritten as the trace of the entropic difference operator defined by

$$D_{\alpha}(\eta_{\varkappa},\Lambda,\mathcal{A}) := \eta_{\varkappa} \big(\chi_{\Lambda} \operatorname{Op}_{\alpha}(\mathcal{A}) \chi_{\Lambda} \big) - \chi_{\Lambda} \eta_{\varkappa} \big(\operatorname{Op}_{\alpha}(\mathcal{A}) \big) \chi_{\Lambda} \,.$$

Since for $\alpha \to \infty$ the symbol converges to

$$A_{\infty}(\xi) := \lim_{\alpha \to \infty} A_{\alpha}(\xi) = e^{-|\xi|} \begin{pmatrix} \chi_{\mathbb{R}^+}(\xi) & 0\\ 0 & \chi_{\mathbb{R}^-}(\xi) \end{pmatrix}$$

we split the trace of the entropic difference into the two contributions

$$\operatorname{tr} D_{\alpha}(\eta, \Lambda, \mathcal{A}_{\alpha}) = \operatorname{tr} D_{\alpha}(\eta_{\varkappa}, \Lambda, \mathcal{A}_{\infty}) + \left(\operatorname{tr} D_{\alpha}(\eta_{\varkappa}, \Lambda, \mathcal{A}_{\alpha}) - \operatorname{tr} D_{\alpha}(\eta_{\varkappa}, \Lambda, \mathcal{A}_{\infty})\right)$$
$$= \operatorname{tr} D_{\alpha}(\eta_{\varkappa}, \Lambda, \mathcal{A}_{\infty}) + \operatorname{tr} \left(D_{\alpha}(\eta_{\varkappa}, \Lambda, \mathcal{A}_{\alpha}) - D_{\alpha}(\eta_{\varkappa}, \Lambda, \mathcal{A}_{\infty})\right). \quad (5.1)$$

In the following sections, we shall analyze these two contributions separately. The first term will give rise to to the area law, whereas the second term will tend to zero. Before entering the details, we remark that in the massless case m = 0, the second term vanishes. Therefore, the proof in the massless case will be completed already at the end of Section 5.1

5.1. The Diagonal Terms. In this section, we shall estimate the trace of the entropic difference $D_{\alpha}(\eta_{\varkappa}, \Lambda, \mathcal{A}_{\infty})$. Since the symbols \mathcal{A}_{∞} is diagonal, our analysis reduces to that in [9].

Lemma 5.1.

$$\lim_{\alpha \to \infty} \frac{1}{\log \alpha} \operatorname{tr} D_{\alpha}(\eta_{\varkappa}, \Lambda, \mathcal{A}_{\infty}) = \frac{1}{\pi^2} \int_0^1 \frac{\eta_{\varkappa}(t)}{t(1-t)} \, dt = \frac{1}{6} \frac{\varkappa + 1}{\varkappa} \, .$$

Proof. Having a diagonal symbol, the trace splits into a sum of the entropic differences for scalar-valued symbols. Indeed, setting

$$a_1(\xi) := e^{-\xi} \chi_{\mathbb{R}^+}(\xi)$$
 and $a_2(\xi) := e^{\xi} \chi_{\mathbb{R}^-}(\xi)$,

the trace of the entropic difference is given by

$$\operatorname{tr} D_{\alpha}(\eta_{\varkappa}, \Lambda, \mathcal{A}_{\infty}) = \sum_{i=1}^{2} \operatorname{tr} D_{\alpha}(\eta, \Lambda, a_{i}) \,.$$

Moreover, the function η_{\varkappa} satisfies Condition 3.1, as explained in Example 3.2. Therefore, we are in the position to apply [9, Corollary 5.10], obtaining

$$\lim_{\alpha \to \infty} \frac{1}{\log(\alpha)} \operatorname{tr} D_{\alpha}(\eta, \Lambda, a_i) = \frac{1}{\pi^2} U(1; \eta_{\varkappa}) = \frac{1}{2\pi^2} \int_0^1 \frac{\eta_{\varkappa}(t)}{t(1-t)} dt = \frac{1}{12} \frac{\varkappa + 1}{\varkappa}.$$

ming over *i* concludes the proof.

Summing over i concludes the proof.

5.2. The Off-Diagonal Terms. It remains to show that the off-diagonal contribution in (5.1) vanishes asymptotically, i.e.

$$\lim_{\alpha \to \infty} \frac{1}{\log \alpha} \operatorname{tr} \left(D_{\alpha}(\eta_{\varkappa}, \Lambda, \mathcal{A}_{\alpha}) - D_{\alpha}(\eta_{\varkappa}, \Lambda, \mathcal{A}_{\infty}) \right) = 0.$$
(5.2)

We begin by splitting the symbol \mathcal{A}_{α} into the sum of low $\mathcal{A}^{<}$ and high $\mathcal{A}^{>}$ frequencies parts defined by

$$\begin{aligned} \mathcal{A}_{\alpha}^{>}(k) &= \Theta\big(|k| - |\log \alpha|\big) \,\mathcal{A}_{\alpha}(k) + \Theta\big(|\log \alpha| - |k|\big) \,\mathcal{A}_{\infty}(k) \\ \mathcal{A}_{\alpha}^{<}(k) &= \Theta\big(|\log \alpha| - |k|\big) \,\mathcal{A}_{\alpha}(k) - \Theta\big(|\log \alpha| - |k|\big) \,\mathcal{A}_{\infty}(k) \,. \end{aligned}$$

We next show that the high frequencies satisfies the following estimate.

Proposition 5.2.

$$\lim_{\alpha \to \infty} \frac{1}{\log \alpha} \operatorname{tr} \left(D_{\alpha}(\eta_{\varkappa}, \Lambda, \mathcal{A}_{\alpha}^{>}) - D_{\alpha}(\eta_{\varkappa}, \Lambda, \mathcal{A}_{\infty}) \right) = 0.$$

Proof. Our claim follows directly by Lemma 4.1. We just have to check that all the hypothesis are satisfied. First of all, we notice that

$$\begin{split} |\log \alpha| &\geq |k| \\ \frac{m}{\omega(k)} \leq \frac{m}{|k|} \leq \frac{m}{|\log \alpha|} \\ |\frac{|k|}{\omega(k)} - 1| &= \frac{\omega(k) - |k|}{\omega(k)} = \frac{\sqrt{k^2 + m^2} - |k|}{\omega(k)} \lesssim \frac{m^2}{\omega(k)^2} \lesssim \frac{m^2}{\log^2 \alpha} \\ \omega(k) &\geq |\log \alpha| \,. \end{split}$$

Therefore

 $\mathcal{A}_{\alpha}^{>} - A_{\infty}$ converges uniformly to zero as $\alpha \searrow 0$.

and Condition (4.1) is fulfilled. Moreover, the function η_{\varkappa} in Example 3.2 satisfies Condition 3.1 with $\gamma < \min\{1, \varkappa\}$. Finally, Condition (4.2) follows from Lemma 4.4 (for l large enough). So we can conclude.

Using Proposition 5.2, we can now estimate the left side of (5.2) as follows,

$$\begin{split} \lim_{\alpha \to \infty} \frac{1}{\log \alpha} \left| \operatorname{tr} \left(D_{\alpha}(\eta_{\varkappa}, \Lambda, \mathcal{A}_{\alpha}) - D_{\alpha}(\eta_{\varkappa}, \Lambda, \mathcal{A}_{\infty}) \right) \right| \\ &= \lim_{\alpha \to \infty} \frac{1}{\log \alpha} \left| \operatorname{tr} \left(D_{\alpha}(\eta_{\varkappa}, \Lambda, \mathcal{A}_{\alpha}) - D_{\alpha}(\eta_{\varkappa}, \Lambda, \mathcal{A}_{\alpha}^{>}) \right) \right| \\ &\leq \lim_{\alpha \to \infty} \frac{1}{\log \alpha} \left\| D_{\alpha}(\eta_{\varkappa}, \Lambda, \mathcal{A}_{\alpha}) - D_{\alpha}(\eta_{\varkappa}, \Lambda, \mathcal{A}_{\alpha}^{>}) \right\|_{1} \\ &\leq \lim_{\alpha \to \infty} \frac{1}{\log \alpha} \left(\left\| \eta_{\varkappa} \left(\chi_{\Lambda} \operatorname{Op}_{\alpha}(\mathcal{A}_{\alpha}) \chi_{\Lambda} \right) - \eta_{\varkappa} \left(\chi_{\Lambda} \operatorname{Op}_{\alpha}(\mathcal{A}_{\alpha}^{>}) \chi_{\Lambda} \right) \right\|_{1} \\ &+ \left\| \chi_{\Lambda} \eta_{\varkappa} \left(\operatorname{Op}_{\alpha}(\mathcal{A}_{\alpha}) \right) \chi_{\Lambda} - \chi_{\Lambda} \eta_{\varkappa} \left(\operatorname{Op}_{\alpha}(\mathcal{A}_{\alpha}^{>}) \right) \chi_{\Lambda} \right\|_{1} \right). \end{split}$$

Therefore, fixing $\sigma \in (2/3, 1)$ and applying Proposition 3.3, we obtain

$$\begin{aligned} \left\| \eta_{\varkappa} (\chi_{\Lambda} \operatorname{Op}_{\alpha}(A_{\alpha})\chi_{\Lambda}) - \eta_{\varkappa} (\chi_{\Lambda} \operatorname{Op}_{\alpha}(A_{\alpha}^{>})\chi_{\Lambda}) \right\|_{1} &\leq C \left\| \chi_{\Lambda} \left(\operatorname{Op}_{\alpha}(A_{\alpha} - A_{\alpha}^{>}) \right) \chi_{\Lambda} \right\|_{\sigma}^{\sigma} \\ &\leq C \left\| \chi_{\Lambda} \operatorname{Op}_{\alpha}(A_{\alpha}^{<})\chi_{\Lambda} \right\|_{\sigma}^{\sigma}. \end{aligned}$$

Using inequality (3.5), the second contribution instead can be rewritten as

$$\begin{aligned} \left\| \chi_{\Lambda} \Big(\eta_{\varkappa} \big(\operatorname{Op}_{\alpha}(A_{\alpha}) \big) - \eta_{\varkappa} \big(\operatorname{Op}_{\alpha}(A_{\alpha}^{>}) \big) \Big) \chi_{\Lambda} \right\|_{1} \\ &= \left\| \chi_{\Lambda} \Big(\eta_{\varkappa} \big(\operatorname{Op}_{\alpha}(A_{\alpha}) \big) - \eta_{\varkappa} \big(\operatorname{Op}_{\alpha}(A_{\alpha}^{>}) \big) \Big) \chi_{\Lambda} \right\|_{1} \\ &\leq \left\| \chi_{\Lambda} \right\|_{\infty} \left\| \Big(\eta_{\varkappa} \big(\operatorname{Op}_{\alpha}(A_{\alpha}) \big) - \eta_{\varkappa} \big(\operatorname{Op}_{\alpha}(A_{\alpha}^{>}) \big) \Big) \chi_{\Lambda} \right\|_{1} \\ &\leq \left\| \eta_{\varkappa} \big(\operatorname{Op}_{\alpha}(A_{\alpha}) \big) - \eta_{\varkappa} \big(\operatorname{Op}_{\alpha}(A_{\alpha}^{>}) \big) \right\|_{1} \| \chi_{\Lambda} \|_{\infty} \\ &\leq \left\| \eta_{\varkappa} \big(\operatorname{Op}_{\alpha}(A_{\alpha}) \big) - \eta_{\varkappa} \big(\operatorname{Op}_{\alpha}(A_{\alpha}^{>}) \big) \right\|_{1} \leq C \left\| \operatorname{Op}_{\alpha}(A_{\alpha}^{<}) \right\|_{1}, \end{aligned}$$

where we used again Proposition 3.3 as before.

Proposition 5.3.

$$\lim_{\alpha \to \infty} \frac{1}{\log \alpha} \|\chi_{\Lambda} \operatorname{Op}_{\alpha}(A_{\alpha}^{<}) \chi_{\Lambda}\|_{\sigma}^{\sigma} = 0 \quad \text{and} \quad \lim_{\alpha \to \infty} \frac{1}{\log \alpha} \|\operatorname{Op}_{\alpha}(A_{\alpha}^{<})\|_{1} = 0.$$

Proof. On account of [8, Proposition 2.14], for any $\sigma \in (0,2)$ we can estimate the Schatten norm by

$$\left\|\chi_{\Lambda}\operatorname{Op}_{\alpha}(\mathcal{A}_{\alpha}^{<})\chi_{\Lambda}\right\|_{\sigma} = \left\|\chi_{\Lambda}\operatorname{Op}_{1}(\mathcal{A}_{\alpha}^{<})\right\|_{\sigma} \leq C\left|\chi_{\Lambda}\right|_{\sigma}\left|\mathcal{A}_{\alpha}^{<}\right|_{\sigma}$$

where the norm $|\cdot|_{\sigma}$ is defined by

...

$$|g|_{\sigma} := \left[\sum_{z \in \mathbb{Z}} \left(\int_{z}^{z+1} |g(x)|^2\right)^{\sigma/2}\right]^{1/\sigma}.$$

Choosing $\sigma = 1$ we can see that

$$\int_{1}^{|\log \alpha|} \frac{1}{k} \, dk \le \log(-\log \alpha) \ll \log \alpha \,,$$

Therefore, for $\sigma < 1$ but close to one we obtain

$$\int_{1}^{|\log \alpha|} \frac{1}{k^{\sigma}} \, dk \le \frac{1}{1 - \sigma} \, k^{1 - \sigma} \big|_{k = |\log \alpha|} = \frac{1}{1 - \sigma} \, |\log \alpha|^{1 - \sigma} \ll \log \alpha \, .$$

These norms are bounded uniformly in α , concluding the proof.

This concludes the proof of Theorem 1.1.

Acknowledgments: We would like to thank Alexander Sobolev for helpful discussions. M.L. gratefully acknowledges support by the Studienstiftung des deutschen Volkes and the Marianne-Plehn-Programm. We would like to thank the "Universitätsstiftung Hans Vielberth" for support. S.M. is partially supported by INFN and by MIUR Excellence Department Project 2023-2027 awarded to the Department of Mathematics of the University of Genoa. This work was written within the activities of the GNFM group of INdAM.

References

- L. Amico, R. Fazio, A. Osterloh, and V. Vedral, *Entanglement in many-body systems*, arXiv:quant-ph/0703044, Rev. Modern Phys. 80 (2008), no. 2, 517–576.
- [2] M.Sh. Birman and M.Z. Solomjak, Spectral theory of selfadjoint operators in Hilbert space, Mathematics and its Applications (Soviet Series), D. Reidel Publishing Co., Dordrecht, 1987, Translated from the 1980 Russian original by S. Khrushchëv and V. Peller. MR 1192782
- [3] L. Bollmann and P. Müller, Widom's formula for discontinuous matrix-valued symbols, arXiv:2311.06036 [math.SP] (2023).
- [4] _____, Enhanced area law in the Widom-Sobolev formula for the free Dirac operator in arbitrary dimension, arXiv:2405.14356 [math-ph] (2024).
- [5] M. Capoferri and S. Murro, Global and microlocal aspects of dirac operators: propagators and hadamard states, arXiv:2201.12104 [math.AP], to appear in Adv. Diff. Eq. (2024).
- [6] F. Finster, R. Jonsson, M. Lottner, A. Much, and S. Murro, Notions of fermionic entropies for causal fermion systems, in preparation.
- [7] F. Finster, S. Kindermann, and J.-H. Treude, An Introductory Course on Causal Fermion Systems, in preparation, www.causal-fermion-system.com/intro-public.pdf (2024).
- [8] F. Finster and M. Lottner, *The fermionic entanglement entropy of the vacuum state of a Schwarzschild black hole horizon*, arXiv:2302.07212 [math-ph], Ann. Henri Poincaré (2024).
- [9] F. Finster, M. Lottner, and A.V. Sobolev, The fermionic entanglement entropy and area law for the relativistic Dirac vacuum state, arXiv:2310.03493 [math-ph], to appear in Adv. Theor. Math. Phys. (2025).
- [10] S. Galanda, A. Much, and R. Verch, Relative entropy of fermion excitation states on the CAR algebra, arXiv:2305.02788 [math-ph], Math. Phys. Anal. Geom. 26 (2023), 21.
- [11] R. Helling, H. Leschke, and W. Spitzer, A special case of a conjecture by Widom with implications to fermionic entanglement entropy, arXiv:0906.4946 [math-ph], Int. Math. Res. Not. IMRN (2011), no. 7, 1451–1482.
- [12] S. Hollands and K. Sanders, Entanglement Measures and their Properties in Quantum Field Theory, arXiv:1702.04924 [quant-ph], Springer Briefs in Mathematical Physics, vol. 34, Springer, Cham, 2018.
- [13] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, Quantum entanglement, arXiv:quant-ph/0702225, Rev. Modern Phys. 81 (2009), no. 2, 865–942.
- [14] H. Leschke, A.V. Sobolev, and W. Spitzer, Trace formulas for Wiener-Hopf operators with applications to entropies of free fermionic equilibrium states, arXiv:1605.04429 [math.SP], J. Funct. Anal. 273 (2017), no. 3, 1049–1094.
- [15] _____, Rényi entropies of the free Fermi gas in multi-dimensional space at high temperature, Toeplitz operators and random matrices—in memory of Harold Widom, arXiv:2201.11087 [mathph], Oper. Theory Adv. Appl., vol. 289, Birkhäuser/Springer, Cham, 2022, pp. 477–508.
- [16] A.V. Sobolev, Functions of self-adjoint operators in ideals of compact operators, arXiv:1504.07261
 [math.SP], J. Lond. Math. Soc. (2) 95 (2017), no. 1, 157–176.
- [17] E. Witten, On entanglement properties of quantum field theory, arXiv:1803.04993 [hep-th], Rev. Mod. Phys. 90 (2018), 045003.

FAKULTÄT FÜR MATHEMATIK UNIVERSITÄT REGENSBURG D-93040 REGENSBURG GERMANY *Email address:* finster@ur.de

FAKULTÄT FÜR MATHEMATIK UNIVERSITÄT REGENSBURG D-93040 REGENSBURG GERMANY *Email address*: magdalena.lottner@freenet.de

INSTITUT FÜR THEORETISCHE PHYSIK, UNIVERSITÄT LEIPZIG, D-04103 LEIPZIG, GERMANY *Email address*: much@itp.uni-leipzig.de

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI GENOVA, I-16146 GENOVA ITALY *Email address:* murro@dima.unige.it