

# Correlated Systematic Uncertainties and Errors-on-Errors in Measurement Combinations: Methodology and Application to the 7-8 TeV ATLAS-CMS Top Quark Mass Combination

Enzo Canonero<sup>1</sup> and Glen Cowan<sup>1</sup>

<sup>1</sup> Physics Department, Royal Holloway, University of London, Egham, TW20 0EX, U.K.

July 9, 2024

## Abstract

The Gamma Variance Model (GVM) is a statistical model that incorporates uncertainties in the assignment of systematic errors (informally called *errors-on-errors*). The model is of particular use in analyses that combine the results of several measurements. In the past, combinations have been carried out using two alternative approaches: the Best Linear Unbiased Estimator (BLUE) method or what we will call the nuisance-parameter method. In this paper we derive useful relations that allow one to connect the BLUE and nuisance-parameter methods when the correlations induced by systematic uncertainties are non-trivial ( $\pm 1$  or  $0$ ), and we generalise the nuisance-parameter approach to include *errors-on-errors*. We then illustrate some of the properties of the GVM by applying it to the 7-8 TeV ATLAS-CMS top quark mass combination. We present results by considering the largest systematic uncertainties as uncertain, one at a time, and we vary their associated *error-on-error* parameters. This procedure is useful for identifying the systematic uncertainties to which a combination is sensitive when they are themselves uncertain. We also explore the hypothetical scenario of including an outlier in the combination, which could become relevant for future combinations, by artificially adding a fictitious measurement to it. This example highlights a key feature of the GVM: its sensitivity to the internal consistency of the input data.

Keywords: Gamma Variance Model, errors-on-errors, combining measurements, correlated systematics, nuisance parameters, ATLAS-CMS top mass combination

# 1 Introduction

In Particle Physics analyses such as those at the Large Hadron Collider (LHC), many results are already dominated by systematic uncertainties or will be shortly. In this context, uncertainties in the assigned values of systematic errors themselves – informally called *errors-on-errors* – become increasingly important. Often, systematic uncertainties are estimated using real datasets of control measurements or Monte Carlo simulations. However, this is not always the case. Theoretical uncertainties often require ad hoc procedures for estimation, which can carry an intrinsic degree of uncertainty. Additionally, “two-point systematics,” where uncertainties are estimated from the difference in results from two methods, provide only a limited picture of the actual uncertainty. The Gamma Variance Model (GVM) [1, 2] provides a statistical framework to handle *errors-on-errors*.

In this paper, we focus on the application of the model to combinations. We first show how a likelihood with nuisance parameters can be constructed when the correlations induced by systematic uncertainties are not simply  $\pm 1$  or 0, and how this approach relates to the Best Linear Unbiased Estimator (BLUE) method [3, 4, 5]. Next, we demonstrate how this model can be generalised to include *errors-on-errors*. This framework is then applied to the 7-8 TeV ATLAS-CMS top quark mass combination [6]. Specifically, we investigate the impact on the combination’s central value and confidence interval when the largest sources of systematic uncertainties are treated as uncertain. We present the results by considering the largest systematic uncertainties as uncertain, one at a time, and we vary their associated *error-on-error* parameters. This approach serves as a general method to identify the systematic uncertainties to which a combination is sensitive when *errors-on-errors* are considered.

Finally, we explore how the inclusion of a hypothetical measurement that is in tension with the rest of the data affects the combination’s sensitivity to uncertain systematics. This exercise can be relevant for future combinations as some existing top mass measurements deviate from the result of the ATLAS-CMS combined result [7, 8]. This example illustrates the behavior of the GVM in scenarios where the inputs of a combination are not entirely internally compatible. Unlike a standard combination, incorporating *errors-on-errors* makes the central value of the combination more robust to discrepancies in the input data. Specifically, the central value becomes less sensitive to outliers. At the same time, the presence of discrepancies inflates the confidence interval for the parameter of interest. This increase reflects the uncertainty stemming from the fact that the measurements in the dataset are not entirely compatible.

In Sec. 2, we present an overview of how a combination can be performed using nuisance parameters under non-trivial correlation assumptions, and how this method relates to the BLUE approach. Section 3 illustrates how to extend combinations to take *errors-on-errors* into account. In Sec. 4, we apply the GVM to the 7-8 TeV ATLAS-CMS top mass combination and explore the impact of introducing an outlier to the combination. A summary and conclusions are presented in Sec. 5.

## 2 Equivalence between nuisance-parameter and BLUE methods in combinations with non-trivial correlations

In Particle Physics, statistical data analysis aims at testing theoretical hypotheses  $H$  using experimentally collected data, denoted as  $\mathbf{y}$ . The probability distribution of the data,  $P(\mathbf{y}|H)$ , is often indexed by a set of Parameters Of Interest (POIs)  $\boldsymbol{\mu}$  and Nuisance Parameters (NPs)  $\boldsymbol{\theta}$ . NPs account for systematic uncertainties in the model, while the POIs are the main objective of the analysis. The likelihood function,  $L(\boldsymbol{\mu}, \boldsymbol{\theta}) = P(\mathbf{y}|\boldsymbol{\mu}, \boldsymbol{\theta})$ , is the central object needed for inference about the POIs.

Here, we explore the application of combining  $N$  measurements  $\mathbf{y} = (y_1, \dots, y_N)$  of a parameter, each with its own statistical and systematic uncertainties, to obtain a single best estimate. We assume that  $\mathbf{y}$  follows a multivariate Gaussian distribution with expected values  $E[y_i] = \mu + \sum_{s=1}^M \Gamma_i^s \theta_s$  and covariance matrix  $V_{ij} = \text{cov}[y_i, y_j]$ , where  $M$  is the total number of NPs. Each nuisance parameter  $\theta_s$  represents a potential source of bias in the measured  $y_i$  values, and the magnitude of these biases is described by the matrix  $\Gamma$ .

Additionally, independent *control measurements*  $\mathbf{u}$  are used to constrain the NPs  $\boldsymbol{\theta}$ . These control measurements  $\mathbf{u} = (u_1, \dots, u_M)$  are assumed to be best estimates of the NPs  $\boldsymbol{\theta} = (\theta_1, \dots, \theta_M)$  and are treated as independently Gaussian-distributed random variables with standard deviations  $\boldsymbol{\sigma}_{\mathbf{u}} = (\sigma_{u_1}, \dots, \sigma_{u_M})$ . Under these assumptions, the log-likelihood of the model is

$$\ell(\mu, \boldsymbol{\theta}) = -\frac{1}{2} \sum_{i,j=1}^N \left( y_i - \mu - \sum_{s=1}^M \Gamma_i^s \theta_s \right) V_{ij}^{-1} \left( y_j - \mu - \sum_{s=1}^M \Gamma_j^s \theta_s \right) - \frac{1}{2} \sum_{s=1}^M \frac{(u_s - \theta_s)^2}{\sigma_{u_s}^2}. \quad (1)$$

It is usually convenient to redefine the matrix  $\Gamma$ , the NPs  $\theta_s$  and the auxiliary measurements  $u_s$  such that  $u_s$  is Gaussian distributed about  $\theta_s$  with a variance of unity. In the real experiment, the best estimates of the  $u_s$  are zero. The idea here is that if there are known biases in the measurements, these are already subtracted. The remaining systematic uncertainty lies in the fact that this bias removal might be imprecise, due to the limited accuracy with which we can identify and quantify the biases. Nevertheless, here we will retain the complete notation in our discussion for greater generality. Auxiliary measurements must be explicitly included to perform Monte Carlo (MC) simulations, while the systematic variances  $\sigma_{u_s}^2$  are needed to generalise the model to include errors-on-errors.

The above likelihood can be related to the Best Linear Unbiased Estimate (BLUE) method [4, 5], which is a commonly used technique for performing combinations that does not require the introduction of NPs. This method involves constructing an approximate Gaussian likelihood to describe the  $N$  measurements of the combination,  $\mathbf{y}$ , based on a covariance matrix  $W_{ij}$  that includes both statistical and systematic uncertainties. The resulting log-likelihood

$$\ell(\mu) = -\frac{1}{2} \sum_{i,j=1}^N (y_i - \mu) W_{ij}^{-1} (y_j - \mu), \quad (2)$$

is then minimized to get the best estimate of the parameter of interest  $\mu$ , which we denote as  $\hat{\mu}$ . The fundamental assumption of the method is that the covariance matrix  $W$  can be expressed as a sum of terms: one for the statistical uncertainties  $V$  and one for each systematic source  $U^{(s)}$ :

$$W_{ij} = V_{ij} + \sum_{s=1}^M U_{ij}^{(s)}. \quad (3)$$

Here the index  $s$  runs over each source of systematic uncertainty. These systematics are independent from each other in the sense that their associated control measurements are uncorrelated, but they induce correlations between the  $\mathbf{y}$  measurements. This underlying structure is more evident when Eq.(1) is used instead. In fact, the two methods are equivalent when the auxiliary measurements are set to zero,  $u_s = 0$ . Specifically, by profiling Eq.(1) over the NPs  $\theta_s$ , one can recover the BLUE log-likelihood as defined in Eq.(2). Furthermore, the relationship between the matrix  $\Gamma$  and the BLUE covariance matrix  $W$  uses Eq. (3) with

$$U_{ij}^{(s)} = \Gamma_i^s \Gamma_j^s \sigma_{u_s}^2. \quad (4)$$

Further discussion of the equivalence of the two methods can be found in [9, 10, 11, 12, 13, 14, 15, 16, 17, 18].

A limitation of this approach is that each systematic effect  $s$  can only lead to an induced correlation factor  $\rho_{ij}^{(s)} = U_{ij}^{(s)} / \sqrt{U_{ii}^{(s)} U_{jj}^{(s)}}$  of  $\pm 1$  or 0. This restriction arises from the definition of the systematic terms  $U_{ij}^{(s)}$  in the BLUE covariance matrix, as specified in Eq. (4). Generalising the BLUE method to accommodate this is trivial as it only requires modifying Eq. (4) as

$$U_{ij}^{(s)} = \rho_{ij}^{(s)} \Gamma_i^s \Gamma_j^s \sigma_{u_s}^2. \quad (5)$$

Note that one may not, however, assign arbitrary values to the  $\rho_{ij}^{(s)}$  as this could lead to  $U^{(s)}$  not being positive definite.

It is conceptually more complicated to extend the log-likelihood of Eq. (1) to achieve the same result. To accomplish this, one needs to define a distinct nuisance parameter  $\theta_s^i$  associated with the systematic effect  $s$  for each measurement  $y_i$ , and correlate the corresponding auxiliary measurements  $u_s^i$  by defining a covariance matrix  $C_{ij}^{(s)}$  as

$$C_{ij}^{(s)} = \text{cov}[u_s^i, u_s^j] = \rho_{ij}^{(s)} \sigma_{u_s}^2. \quad (6)$$

Thus the log-likelihood of Eq. (1) becomes

$$\begin{aligned} \ell(\mu, \boldsymbol{\theta}) = & -\frac{1}{2} \sum_{i,j=1}^N \left( y_i - \mu - \sum_{s=1}^M \Gamma_i^s \theta_s^i \right) V_{ij}^{-1} \left( y_j - \mu - \sum_{s=1}^M \Gamma_j^s \theta_s^j \right) \\ & - \frac{1}{2} \sum_{s=1}^M \sum_{i,j=1}^N (u_s^i - \theta_s^i) \left( C^{(s)} \right)_{ij}^{-1} (u_s^j - \theta_s^j). \end{aligned} \quad (7)$$

In App. A we prove the important result that this log-likelihood becomes equivalent to the BLUE method when the auxiliary measurements  $u_s^i$  are set to zero. This equivalence can be demonstrated by profiling Eq.(7) with respect to all the NPs, which leads to the BLUE likelihood as defined in Eq.(2), with covariance matrix specified by Eqs. (3) and (5).

Defining independent NPs  $\theta_s^i$  for each measurement  $i$  can be redundant when the correlations between the associated auxiliary measurements  $u_s^i$  are equal to  $\pm 1$ . In such cases, one can use a single NP for all auxiliary measurements that are fully correlated or anti-correlated, as implemented in Eq. (1).

Note that here we are not introducing correlations between estimates of NPs associated with different systematics, as is done in the model proposed in [19]. That model addresses combination of profile likelihood fits, where estimates of POIs and NPs are correlated with each other. Here, instead, for each systematic  $s$ , we introduce a separate NP  $\theta_s^i$  and auxiliary measurement  $u_s^i$  for each measurement  $i$ , and we correlate only the auxiliary measurements for a given systematic source  $s$ . We do this to match the BLUE result obtained with the matrix defined in Eq. (5).

### 3 Extending combinations to account for *errors-on-errors*

The likelihoods of Eqs (1) and (7) can be extended to account for the presence of uncertainties in the assignment of systematic errors, informally denoted as *errors-on-errors*. This is achieved using the Gamma Variance Model (GVM) [1, 2]. In this model the variances  $\sigma_{u_s}^2$  of the control measurements  $u_s$  are treated as adjustable parameters, and each  $\sigma_{u_s}^2$  is constrained by its best estimate  $v_s = s_s^2$ . The estimates  $\mathbf{v}$  are assumed to follow independent gamma distributions, which are suitable for positive-valued random variables. The model derived from this approach allows for the closed-form profiling of the likelihood over the NPs  $\sigma_{u_i}^2$ . Detailed methodology can be found in [1].

Applying this procedure to Eq. (1) is straightforward, resulting in the replacement of quadratic constraints with logarithmic constraints. This yields the log-likelihood:

$$\begin{aligned} \ell(\mu, \boldsymbol{\theta}) = & -\frac{1}{2} \sum_{i,j=1}^N \left( y_i - \mu - \sum_{s=1}^M \Gamma_i^s \theta_s^i \right) V_{ij}^{-1} \left( y_j - \mu - \sum_{s=1}^M \Gamma_j^s \theta_s^j \right) \\ & - \frac{1}{2} \sum_{s=1}^M \left( 1 + \frac{1}{2\varepsilon_s^2} \right) \log \left[ 1 + 2\varepsilon_s^2 \frac{(u_s - \theta_s)^2}{v_s} \right]. \end{aligned} \quad (8)$$

The parameter  $\varepsilon_s$  quantifies the relative uncertainty associated with the systematic error  $\sigma_{u_s}$  and is commonly referred to as the *error-on-error* parameter. For instance, if  $\varepsilon_s$  equals 0.3, this indicates that  $s_s = \sqrt{v_s}$  has a 30% relative uncertainty as an estimate of  $\sigma_{u_s}$ .

Generalising the likelihood of Eq.(7) to account for the presence of *errors-on-errors* is less trivial and requires extending the GVM beyond what was originally presented in [1]. The method to achieve

this is nevertheless similar to what has already been discussed: the variance  $\sigma_{u_s}^2$  in Eq (5) is treated as an adjustable parameter of the model, and  $v_s$ , its best estimate, is considered a gamma-distributed random variable. In a manner similar to that of Sec. 2, it is useful to redefine  $\Gamma_s$ ,  $\theta_s$  and  $u_s$  such that if there are no *errors-on-errors* ( $\sigma_{u_s}^2$  exactly known), then  $u_s$  is Gaussian distributed with a mean  $\theta_s$  and unit variance. In this way the best estimates of the variances are all  $v_s = 1$ . This does not alter the approach used to include *errors-on-errors*, as it is merely a redefinition of parameters. The resulting generalised log-likelihood is

$$\begin{aligned} \ell(\mu, \boldsymbol{\theta}) = & -\frac{1}{2} \sum_{i,j=1}^N \left( y_i - \mu - \sum_{s=1}^M \Gamma_i^s \theta_s^i \right) V_{ij}^{-1} \left( y_j - \mu - \sum_{s=1}^M \Gamma_j^s \theta_s^j \right) \\ & - \frac{1}{2} \sum_{s=1}^M \left( N + \frac{1}{2\varepsilon_s^2} \right) \log \left[ 1 + 2\varepsilon_s^2 \sum_{i,j=1}^N (u_s^i - \theta_s^i) \left( C^{(s)} \right)_{ij}^{-1} (u_s^j - \theta_s^j) \right]. \end{aligned} \quad (9)$$

Here the definition of  $C^{(s)}$  differs from Eq. (6) as  $\sigma_{u_s}^2$  is replaced by its best estimate  $v_s$ :

$$C_{ij}^{(s)} = \text{cov}[u_s^i, u_s^j] = \rho_{ij}^{(s)} v_s. \quad (10)$$

Even if in a fit  $v_s$  would be set to 1, it is important to keep the full notation as it is needed to perform MC simulations. In App. B, we detail the theoretical assumptions and the mathematical proof to obtain this formula. This equation represents the most general formula for combining measurements using the GVM, incorporating uncertainties on systematic errors.

In practical applications, the most feasible approach to account for *errors-on-errors* involves using NPs exclusively for systematic sources where  $\varepsilon_s \neq 0$ , and profiling over all NPs introduced as standard quadratic constraints. This leads to a mixed likelihood with a quadratic part constructed using the BLUE approach, and with NPs introduced only for systematic sources with uncertain errors. This can be achieved by replacing  $V$  with a BLUE covariance matrix  $W$ , as defined in Eq. 3, but summing only over the non-uncertain systematic errors while keeping in the likelihood only those NPs associated with uncertain systematics. A mathematical motivation for this procedure is given in App. A.

Notice that when the correlation coefficients  $\rho_{ij}^{(s)}$  of Eq. (5) are set to zero for some systematic effect  $s$ , the logarithmic term for that systematic effect reduces to

$$\left( N + \frac{1}{2\varepsilon_s^2} \right) \log \left[ 1 + 2\varepsilon_s^2 \sum_{i=1}^N \frac{(u_s^i - \theta_s^i)^2}{v_s} \right]. \quad (11)$$

This implies that all the  $u_s^i$  have the same variance  $\sigma_s^2$  whose estimated value is  $v_s$ . Practically, this means assuming that if one systematic uncertainty is overestimated or underestimated by a certain factor, this factor will be the same for all the measurements affected by that systematic. This is because the systematic described by the NP  $\theta_s^i$  induces a uncertainty on the measurement  $y_i$  equal to  $\Gamma_i^s \sigma_s$ . If one wants to consider both the auxiliary measurements as uncorrelated and the estimates of the systematic uncertainties as independent, one would define the logarithmic terms as

$$\sum_{i=1}^N \left( 1 + \frac{1}{2\varepsilon_s^2} \right) \log \left[ 1 + 2\varepsilon_s^2 \frac{(u_s^i - \theta_s^i)^2}{v_s} \right], \quad (12)$$

In this case one recovers the original GVM as defined in Eq. (1).

## 4 Application to 7-8 TeV ATLAS-CMS top quark mass combination

Measurements and combinations of the top quark mass are a relevant testing ground for the *errors-on-errors* formalism for several reasons. Firstly, these measurements are becoming increasingly dominated by systematic uncertainties, making a precise evaluation of them crucial for the correct interpretation

$t\bar{t}$ final state	CM Energy (TeV)	$m_{top} \pm (\text{stat}) \pm (\text{syst})$ (GeV)	Total uncertainty (GeV)
All-hadronic [24]	7	$175.06 \pm 1.35 \pm 1.21$	$\pm 1.82$
Dileptonic [22]	7	$173.79 \pm 0.54 \pm 1.31$	$\pm 1.42$
Lepton+jets [22]	7	$172.33 \pm 0.75 \pm 1.04$	$\pm 1.28$
All-hadronic[25]	8	$173.72 \pm 0.55 \pm 1.02$	$\pm 1.16$
Dileptonic [23]	8	$172.99 \pm 0.41 \pm 0.74$	$\pm 0.84$
Lepton+jets [21]	8	$172.08 \pm 0.39 \pm 0.82$	$\pm 0.91$

**Table 1.** Top quark mass measurements in  $t\bar{t}$  final states at different center-of-mass energies, based on ATLAS data.

$t\bar{t}$ final state	CM Energy (TeV)	$m_{top} \pm (\text{stat}) \pm (\text{syst})$ (GeV)	Total uncertainty (GeV)
All-hadronic [26]	7	$173.49 \pm 0.69 \pm 1.23$	$\pm 1.41$
Dileptonic [27]	7	$172.50 \pm 0.43 \pm 1.52$	$\pm 1.58$
Lepton+jets [28]	7	$173.49 \pm 0.43 \pm 0.97$	$\pm 1.06$
All-hadronic[29]	8	$172.32 \pm 0.25 \pm 0.57$	$\pm 0.62$
Dileptonic [29]	8	$172.22 \pm 0.18 \pm 0.94$	$\pm 0.95$
Lepton+jets [29]	8	$172.35 \pm 0.16 \pm 0.45$	$\pm 0.48$
Single top [30]	8	$172.95 \pm 0.77 \pm 0.93$	$\pm 1.20$
$J/\psi$ [31]	8	$173.50 \pm 3.00 \pm 0.94$	$\pm 3.14$
Secondary vertex [32]	8	$173.68 \pm 0.20 \pm 1.11$	$\pm 1.12$

**Table 2.** Top quark mass measurements in  $t\bar{t}$  final states at different center-of-mass energies, based on CMS data.

of results. Furthermore, measurements of the top quark mass are frequently significantly impacted by modeling uncertainties, especially those exploiting jets in the final state, which themselves can be quite uncertain. In addition, the GVM’s results are sensitive to the internal compatibility of the data, and some of the top mass measurements exhibit tensions between each-other; for example, the Tevatron [7] combination is slightly incompatible with the result of ATLAS-CMS one [6], as well as the run 2 ATLAS top mass measurement exploiting a leptonic invariant mass [8].


We begin by replicating the results of the 7-8 TeV ATLAS-CMS top mass combination [6] using the BLUE method, as described in Sec. 2. To construct the BLUE matrix we exploit Eqs. (3) and (5), using data from Tabs. A.2 and A.3 in the appendices of [6]. Specifically, the statistical covariance matrix, which is diagonal, incorporates the reported statistical uncertainties for each measurement. Additionally, each term  $\Gamma_i^s$  corresponds to the error on the  $i$ -th measurement attributed to systematic source  $s$ . As discussed above, we set  $\sigma_{u_s}^2 = 1$ . The correlation coefficients  $\rho_{ij}^{(s)}$  are determined using the correlation matrices from App. B of [20]. The result found using the BLUE approach is

$$m_t = 172.51 \pm 0.33 \text{ GeV} . \quad (13)$$

Our result has a minor discrepancy of 0.01 GeV from the central value reported in [6], but the confidence interval agrees with that reported in the paper. Discrepancies of this magnitude can be expected since all public results are rounded to the second decimal place.

#### 4.1 Application of GVM to the combined estimate of $m_t$

To demonstrate the GVM methodology, we first show that it can replicate the BLUE result in the limit where the *error-on-error* parameters  $\varepsilon_s$  of Eq. (9) approach zero. At the same time, we investigate the impact of introducing *errors-on-errors* on some of the systematics on the outcome of the combination. Specifically, we analyze each main source of systematic uncertainty individually, assuming that only one systematic has an associated *error-on-error* at a time. This approach allows one to isolate the impact of one specific uncertain systematic error on the combination. We then vary the value of




**Figure 1.** The plot shows the variation of the central value as a function of the *error-on-error* parameter  $\varepsilon_s$ . Each line represents the change of the central value when the systematic uncertainties in the legend are considered uncertain one at a time. The central values are computed explicitly at points marked by dots and linearly interpolated in between.

the error-on-error parameter for that systematic to determine the outcomes different values would produce.

To construct the profile likelihood for the GVM, we use Eq. (9). However, we introduce a NP only for the systematic source treated as uncertain, while all other systematic effects are addressed using the BLUE approach, as detailed in Sec. 2. We set  $u_s^i = 0$  and  $v_s = 1$ . Then, we employ Tabs. A.2 and A.3 from [6] to construct the  $\Gamma$  matrix while the covariance matrices  $C^{(s)}$  are derived using the correlation matrices from App. B of [20]. Some of these correlation matrices exhibit one or more negative eigenvalues. We address this issue by adding a constant term equal to the smallest negative eigenvalue to the diagonal of each covariance matrix. This adjustment ensures that all correlation matrices are positive semi-definite. While this modification has a negligible effect on the overall combination, it is a critical step to ensure the convergence of the log-likelihood minimization process. This regularization step was not required for the BLUE approach, as the full covariance matrix  $W$  in the BLUE method is positive semi-definite.

The ATLAS inputs to the combination [21, 22, 23, 24, 25] are summarized in Tab. 1, while the CMS ones [26, 27, 28, 29, 30, 31, 32] are listed in Tab. 2. We consider as potentially uncertain the eight primary sources of systematic uncertainty as listed in Tab. 2 of [6], as we expect that the largest systematic errors would have a more pronounced effect on the final result when they are themselves uncertain. This selection does not imply that we consider these systematics as uncertain; rather, we intend to demonstrate the potential impact of various assumptions on the results of the combination.

Figure 1 illustrates the variation in the central value of the combination when one of the systematic uncertainties listed in the legend is itself considered uncertain. Specifically, the plot shows the dependence of the central value ( $\hat{\mu}$ ) on the *error-on-error* parameter  $\varepsilon_s$  for each systematic uncertainty. The nomenclature for the systematic components utilized here is the same used in [6], where a detailed explanation of what they describe can be found. The most important conclusion here is that the central value of the combination is quite robust to the presence of uncertain systematic errors.



**Figure 2.** The plot shows the variation of the 68.3% confidence interval as a function of the *error-on-error parameter*  $\varepsilon_s$ . Each line represents the change of the central value when the systematic uncertainties in the legend are considered uncertain one at a time. The confidence intervals are computed explicitly at points marked by dots and linearly interpolated in between.

Specifically, the change in the central value remains always within 0.1 GeV for the explored range of  $\varepsilon$  values, which is well within the confidence interval of approximately 0.3 GeV. Ultimately, all central values converge to the BLUE result as  $\varepsilon$  approaches zero. The reason why the *b tagging* and *LHC radiation* lines do not converge to the same point is due to the regularization of their associated correlation matrices, as previously described.

Similarly, Fig. 2 displays the width of the 68.3% confidence interval as a function of  $\varepsilon_s$ . As for the central value, each line corresponds to the case where only one systematic uncertainty at a time is considered as itself uncertain. The confidence intervals have been computed using Bartlett corrections [33], implemented via MC simulations, as described in [2]. This technique offers an improved approach for calculating likelihood ratio statistics by correcting for deviations from the asymptotic  $\chi^2$  distribution. The idea is to compute confidence intervals as the parameter regions where the likelihood ratio test statistic, denoted in [2] as  $w_\mu$ , is less than  $1 + b/N_{\text{poi}}$ . Here  $N_{\text{poi}}$  is the number of POIs in the likelihood (1 in this example) and  $b$ , the Bartlett correction, is equal to  $E[w_{\hat{\mu}}] - N_{\text{poi}}$ , where  $E[w_{\hat{\mu}}]$  is the expectation of  $w_\mu$  computed for  $\mu = \hat{\mu}$ . If a purely quadratic likelihood were used,  $b$  would be equal to 0. This is equivalent to using the usual  $\ell_{\text{max}} - 1/2$  method [34], but instead of using  $1/2$ , it uses  $1/2(1 + b/N_{\text{poi}})$ . This correction is relevant for the GVM, which exhibits deviations of order  $\mathcal{O}(\varepsilon_s^2)$  from the  $\chi^2$  approximation. The Bartlett correction improves this to order  $\mathcal{O}(\varepsilon_s^4)$ .

Figure 2 partially confirms our previous conclusion: the combination is generally robust to the presence of uncertain systematic errors. However, a non-negligible increase of 10 – 15% in the confidence interval is observed when the *LHC b-JES* systematic uncertainty is known with a relative error of approximately 30% ( $\varepsilon_s \approx 0.3$ ), and it increases up to about 20% for a relative uncertainty of 60% ( $\varepsilon_s \approx 0.6$ ). The combination’s heightened sensitivity to uncertainties in this systematic error stems from the fact that it constitutes the largest uncertainty of the combination. Additionally, the pulls of the NPs associated with this systematic source are non-negligible, even though they are within one standard deviation. This result is significant given that the *LHC b-JES* systematic uncertainty,



modeling the flavor response of b-JES, is a two-points systematic and thus falls into the category of systematic sources that could plausibly be uncertain. The confidence interval remains stable if any other of the systematic uncertainties considered are taken as themselves uncertain, the observed small noisy fluctuations are a consequence of the numerical technique employed to compute the confidence intervals. A negligible bias in the confidence interval size of the *b tagging* and *LHC radiation* uncertainties is present due to the regularization of the correspondent correlation matrices.

## 4.2 Sensitivity to outliers


The conclusions of the last section would be significantly impacted if any of the combination inputs exhibited tension with the rest of the measurements. It is not uncommon to have input values that exhibit a significant tension and it could possibly happen for an updated LHC-Tevatron combination, or for a LHC Run 2 combination that includes the ATLAS top mass measurement exploiting a leptonic invariant mass [8]. However, since extending the combination to include additional measurements would require knowledge of the correlations between them and those listed in Tables 1 and 2, we introduce a fictitious measurement into the dataset to illustrate the properties of the GVM in such scenarios. Specifically, we consider two examples: in the first, we add a measurement with a central value of  $m_t^{NEW} = 174.5$  GeV, a statistical uncertainty of 0.4 GeV, and a global systematic uncertainty of 0.5 GeV. In the second example, the new measurement has the same central value, but both the statistical and systematic uncertainties are halved. We assume that the systematic uncertainties of the new measurements are uncorrelated with the uncertainties of the other measurements.

If the first fictitious measurement is added to the combination, without considering *errors-on-errors*, the result obtained using the BLUE approach is  $m_t^{(1)} = 172.91 \pm 0.29$  GeV. If the second fictitious measurement alone is added, the result of the combination becomes  $m_t^{(2)} = 173.50 \pm 0.23$  GeV. As anticipated, the inclusion of the new measurements shifts the central value of the combination towards higher values and results in a reduction of the confidence interval. Both these effects are more pronounced when the second, more precise, measurement is included.

Figure 3 illustrates the variations in the central value of the combination when one of the systematic uncertainties listed in the legend is itself considered uncertain. The top plot displays them when the first fictitious, less precise, measurement is added to the combination. Conversely, the bottom plot displays the effect of including the second, more precise, measurement. On top of the systematics considered in the last section, the *NEW* systematic uncertainty on the fictitious measurements is also considered. In contrast to observations made in the previous section, the inclusion of the outlier in the dataset makes the effect of *errors-on-errors* non-negligible. For both examples, the central value significantly shifts when the *NEW* systematic uncertainty is considered uncertain. Specifically, the central value shifts back to that of the original combination. This is a property of the GVM: while typically, the presence of an outlier would bias the result of a combination, within the GVM framework as the *error-on-error* parameter increases, the outlier is assigned lesser weight within the combination. Consequently, the central value shifts back towards that of the original combination, thereby diminishing the bias on the combination's result.


Additionally, in the case of the more precise fictitious measurement (bottom plot), a non-negligible shift also occurs if the *CMS B hadron BR* uncertainty has an associated *error-on-error*. This is because the most precise measurements within the ATLAS-CMS combination - the CMS 8 TeV mass measurements in the lepton+jets and all-hadronic channels - are significantly influenced by this uncertainty, which arises from how CMS models the b quark fragmentation function. This time, the central value shifts to higher values of the top mass, as the most precise CMS measurements are assigned less weight in the combination. This example highlights how the GVM can be used as a tool to identify the systematic uncertainties to which a combination is sensitive when they have associated *errors-on-errors*.

Similarly, Fig. 4 displays the variation in the half-size of the 68.3% confidence interval as a function of  $\varepsilon_s$ . The top plot illustrates this variation with the inclusion of the first fictitious measurement into the combination, whereas the bottom plot shows the effect of adding the second, more precise, measurement. Both examples illustrate that when the *NEW* systematic uncertainty has an associated



**Figure 3.** The plot shows the variation of the central value as a function of the *error-on-error parameter*  $\epsilon_s$  when one fictitious measurement is included in the combination. The upper plot shows the effects of incorporating the less precise measurement, while the lower plot illustrates the impact of adding a more precise one. Each line represents the change of the central value when the systematic uncertainties in the legend are considered uncertain one at a time. The central values are computed explicitly at points marked by dots and linearly interpolated in between.

*error-on-error*, the confidence interval is significantly inflated. This growth is more pronounced as the tension between the new measurement and the existing combination inputs increases. Specifically, the confidence interval inflates by up to 30%, when the less precise measurement is included, and



**Figure 4.** The plot shows the variation of the 68.3% confidence interval as a function of the *error-on-error parameter*  $\epsilon_s$  when one fictitious measurement is included in the combination. The upper plot shows the effects of incorporating the less precise measurement, while the lower plot illustrates the impact of adding a more precise one. Each line represents the change in the confidence interval when the systematic uncertainties in the legend are considered uncertain one at a time. The confidence intervals are computed explicitly at points marked by dots and linearly interpolated in between.

by up to 75%, when the more precise one is added. Furthermore, it is noteworthy that when the more precise measurement is included (bottom plot) the *CMS B hadron BR* uncertainty also leads to an inflation of the confidence interval when it is considered uncertain. This is the second relevant

property of the GVM: The size of the confidence interval is sensitive to the internal consistency of the input data. Specifically, the less compatible the input measurements are, the more the confidence interval is subject to inflation when *errors-on-errors* are considered. This is because the GVM treats the tension in the dataset as an additional source of uncertainty resulting in an inflated confidence interval. Additionally, it is observed that the feature where the *LHC b-JES* uncertainty inflates the confidence interval when uncertain is still, as this is inherent to the original inputs of the combination and remains almost unaffected by the addition of new measurements. Here, the inflation of the confidence interval, instead of being driven by incompatibility in the dataset, originates mainly from the correlations among the original input measurements and the fact that the *LHC b-JES* constitutes the dominant uncertainty within the combination.

## 5 Conclusions

The Gamma Variance Model (GVM) provides a powerful statistical framework for addressing uncertainties in the assignment of systematic errors, informally referred to as *errors-on-errors*. In this paper, we have demonstrated how a combination can be extended to incorporate uncertainties on error parameters. As a by-product of this, we provide a useful connection between the BLUE method for combining results when there are non-trivial correlations and the corresponding likelihood method using nuisance parameters. We applied the framework to the 7-8 TeV ATLAS-CMS top quark mass combination [6]. All the results have been studied by considering various systematic uncertainties as uncertain, one at a time, and varying their associated *error-on-error* parameters  $\varepsilon_s$ . The aim of this paper has been to illustrate the impact of *errors-on-errors* on the combination, rather than assigning precise uncertainties to the systematic errors. This methodology can be used as a general procedure to identify the systematic uncertainties to which a combination is sensitive when they have associated *errors-on-errors*, and as a general approach to assess the robustness of a combination against systematic uncertainties that are themselves uncertain.

We conclude that the ATLAS-CMS  $m_t$  combination is robust, with the central values remaining very stable across a broad spectrum of assumed values for *errors-on-errors*. The confidence intervals are generally stable as well, though they exhibit non-negligible sensitivity to uncertainties in the *LHC b-JES* uncertainty.

The scenario of an outlier present in the combination was also explored. This study is relevant to demonstrate the model's properties in situations that may occur in future combinations. This example demonstrated the model's sensitivity to the internal compatibility of the dataset. Specifically, the central value of the combination is less biased by outliers as the *errors-on-errors* parameters increase. Meanwhile, the confidence intervals become inflated with greater internal inconsistency among the data, as the GVM treats internal inconsistency in the input data as an additional source of uncertainty.

## Acknowledgements

The authors are grateful to Mark Owen, Veronique Boisvert, and other colleagues in the ATLAS Collaboration who have provided valuable feedback on this work and to the U.K. Science and Technology Facilities Council for its support.

## References

- [1] G. Cowan, *Statistical Models with Uncertain Error Parameters*, Eur. Phys. J. C (2019) 79:133, <https://doi.org/10.1140/epjc/s10052-019-6644-4>.
- [2] E. Canonero, A. Brazzale and G. Cowan, *Higher-order asymptotic corrections and their application to the Gamma Variance Model*, Eur. Phys. J. C (2023) 83:1100, <https://doi.org/10.1140/epjc/s10052-023-12263-7>.

- [3] Andrea Valassi, *Combining correlated measurements of several different physical quantities*, NIM A (2003) 500, [https://doi.org/10.1016/S0168-9002\(03\)00329-2](https://doi.org/10.1016/S0168-9002(03)00329-2).
- [4] Richard Nisius, *BLUE: combining correlated estimates of physics observables within ROOT using the Best Linear Unbiased Estimate method*, SoftwareX 11 (2020) 100468, DOI: 10.1016/j.softx.2020.100468.
- [5] Richard Nisius, *On the combination of correlated estimates of a physics observable*, Eur. Phys. J. C 74 (2014) 3004, DOI: 10.1140/epjc/s10052-014-3004-2.
- [6] CMS and ATLAS Collaborations, *Combination of measurements of the top quark mass from data collected by the ATLAS and CMS experiments at  $\sqrt{s} = 7$  and 8 TeV.*, 2024, [arXiv:2402.08713](https://arxiv.org/abs/2402.08713).
- [7] *Combination of CDF and D0 Results on the Mass of the Top Quark using up to  $9.7\text{fb}^{-1}$  at the Tevatron.*, (2014), [arXiv:1407.2682](https://arxiv.org/abs/1407.2682).
- [8] The ATLAS collaboration., Aad, G., Abbott, B. et al. *Measurement of the top-quark mass using a leptonic invariant mass in pp collisions at  $\sqrt{s} = 13$  TeV with the ATLAS detector*. J. High Energ. Phys. 2023, 19 (2023). [https://doi.org/10.1007/JHEP06\(2023\)019](https://doi.org/10.1007/JHEP06(2023)019).
- [9] A. Pinto, Z. Wu, F. Balli, N. Berger, M. Boonekamp, É. Chapon, T. Kawamoto, and B. Malaescu, *Uncertainty components in profile likelihood fits*, (2024), <https://arxiv.org/abs/2307.04007>.
- [10] L. Demortier, *Equivalence of the best-fit and covariance-matrix methods for comparing binned data with a model in the presence of correlated systematic uncertainties*, CDF Note 8661 (1999), [https://www-cdf.fnal.gov/physics/statistics/notes/cdf8661\\_chi2fit\\_w\\_corr\\_syst.pdf](https://www-cdf.fnal.gov/physics/statistics/notes/cdf8661_chi2fit_w_corr_syst.pdf).
- [11] G.L. Fogli, E. Lisi, A. Marrone, D. Montanino, and A. Palazzo, *Getting the most from the statistical analysis of solar neutrino oscillations*, Phys. Rev. D 66, (2002) 053010, DOI: 10.1103/PhysRevD.66.053010.
- [12] D. Stump, J. Pumplin, R. Brock, D. Casey, J. Huston, J. Kalk, H.L. Lai, and W.K. Tung, *Uncertainties of predictions from parton distribution functions. I. The Lagrange multiplier method*, Phys. Rev. D 65, (2001) 014012. DOI: 10.1103/PhysRevD.65.014012.
- [13] R.S. Thorne, *Uncertainties in parton related quantities*, J. Phys. G 28, (2002) 2705. DOI: 10.1088/0954-3899/28/10/314.
- [14] M. Botje, *QCD analysis of deep inelastic scattering data*, J. Phys. G 28, (2002) 779. DOI: 10.1088/0954-3899/28/5/305.
- [15] A. Glazov, *Averaging of DIS Cross Section Data*, AIP Conf. Proc. 792(1), (2005) 237. DOI: 10.1063/1.2122026.
- [16] R. Barlow, *Combining experiments with systematic errors*, Nucl. Instrum. Meth. A 987, (2021) 164864. DOI: 10.1016/j.nima.2020.164864.
- [17] B. List, *Decomposition of a covariance matrix into uncorrelated and correlated errors*, presented at the Alliance Workshop on Unfolding and Data Correction, DESY (2010). <https://indico.desy.de/event/3009/contributions/64704/>.
- [18] G. Aad et al., *Measurement of dijet cross-sections in pp collisions at 7 TeV centre-of-mass energy using the ATLAS detector*, JHEP 05, (2014) 059. DOI: 10.1007/JHEP05(2014)059.
- [19] J. Kieseler, *A method and tool for combining differential or inclusive measurements obtained with simultaneously constrained uncertainties*, Eur. Phys. J. C 77(11), (2017) 792. DOI: 10.1140/epjc/s10052-017-5345-0.

- [20] The ATLAS and CMS Collaborations, *Combination of measurements of the top quark mass from data collected by the ATLAS and CMS experiments at  $\sqrt{s} = 7$  and 8 TeV*, CERN, ATLAS-CONF-2023-066, CMS-PAS-TOP-22-001, CERN-LPCC-2023-02.
- [21] M. Aaboud *et al.* [ATLAS], *Measurement of the top quark mass in the  $t\bar{t} \rightarrow$  lepton+jets channel from  $\sqrt{s} = 8$  TeV ATLAS data and combination with previous results*, Eur. Phys. J. C **79** (2019) no.4, 290, doi:10.1140/epjc/s10052-019-6757-9.
- [22] G. Aad *et al.* [ATLAS], *Measurement of the top quark mass in the  $t\bar{t} \rightarrow$  lepton+jets and  $t\bar{t} \rightarrow$  dilepton channels using  $\sqrt{s} = 7$  TeV ATLAS data*, Eur. Phys. J.C **75** (2015) no.7, 330, doi:10.1140/epjc/s10052-015-3544-0.
- [23] M. Aaboud *et al.* [ATLAS], *Measurement of the top quark mass in the  $t\bar{t} \rightarrow$  dilepton channel from  $\sqrt{s} = 8$  TeV ATLAS data*, Phys. Lett. B **761** (2016), 350-371, doi:10.1016/j.physletb.2016.08.042.
- [24] G. Aad *et al.* [ATLAS], *Measurement of the top-quark mass in the fully hadronic decay channel from ATLAS data at  $\sqrt{s} = 7$  TeV*, Eur. Phys. J. C **75** (2015) no.4, 158, doi:10.1140/epjc/s10052-015-3373-1.
- [25] M. Aaboud *et al.* [ATLAS], *Top-quark mass measurement in the all-hadronic  $t\bar{t}$  decay channel at  $\sqrt{s} = 8$  TeV with the ATLAS detector*, JHEP **09** (2017), 118, doi:10.1007/JHEP09(2017)118.
- [26] Chatrchyan, S., Khachatryan, V., Sirunyan, A.M. et al. *Measurement of the top-quark mass in all-jets events in  $t\bar{t}$  pp collisions at  $\sqrt{s} = 7$  TeV*. Eur. Phys. J. C **74**, 2758 (2014). <https://doi.org/10.1140/epjc/s10052-014-2758-x>.
- [27] The CMS Collaboration., Chatrchyan, S., Khachatryan, V. et al. *Measurement of the top-quark mass in  $t\bar{t}$  events with dilepton final states in pp collisions at  $\sqrt{s} = 7$  TeV*. Eur. Phys. J. C **72**, 2202 (2012). <https://doi.org/10.1140/epjc/s10052-012-2202-z>.
- [28] The CMS collaboration., Chatrchyan, S., Khachatryan, V. et al. *Measurement of the top-quark mass in  $t\bar{t}$  events with lepton+jets final states in pp collisions at  $\sqrt{s} = 7$  TeV*. J.High Energ.Phys. **2012**, 105 (2012). [https://doi.org/10.1007/JHEP12\(2012\)105](https://doi.org/10.1007/JHEP12(2012)105).
- [29] CMS Collaboration, *Measurement of the top quark mass using proton-proton data at  $\sqrt{s} = 7$  and 8 TeV.*, hys. Rev. D **93** (2016) 072004, doi:10.1103/PhysRevD.93.072004.
- [30] Sirunyan, A.M., Tumasyan, A., Adam, W. et al. *Measurement of the top quark mass using single top quark events in proton-proton collisions at  $\sqrt{s} = 8$  TeV*. Eur. Phys. J. C **77**, 354 (2017). <https://doi.org/10.1140/epjc/s10052-017-4912-8>.
- [31] The CMS collaboration., Khachatryan, V., Sirunyan, A.M. et al. *Measurement of the mass of the top quark in decays with a  $J/\psi$  meson in pp collisions at 8 TeV*. J. High Energ. Phys. **2016**, 123 (2016). [https://doi.org/10.1007/JHEP12\(2016\)123](https://doi.org/10.1007/JHEP12(2016)123).
- [32] CMS Collaboration, *Measurement of the top quark mass using charged particles in pp collisions at  $\sqrt{s} = 8$  TeV.*, Phys. Rev. D **93** (2016) 092006, doi:10.1103/PhysRevD.93.092006.
- [33] M.S. Bartlett, *Properties of sufficiency and statistical tests*, Royal Society of London Proceedings Series A **160**, (1937) 268-282.
- [34] S. Navas et al. (Particle Data Group), Phys. Rev. D **110**, 030001 (2024).
- [35] Louis Guttman, *Enlargement Methods for Computing the Inverse Matrix*, Ann. Math. Statist. **17**(3): 336-343 (1946); DOI: 10.1214/aoms/1177730946.

## A Profiling over NPs modeling correlated auxiliary measurements

The BLUE log-likelihood, with the covariance matrix defined as in Eq.(5), can be derived by profiling Eq.(7) over all the NPs  $\theta_s^i$  when the auxiliary measurements  $u_s^i$  are set to 0. This process involves profiling the likelihood recursively over all  $\theta_s^i$  for a given  $s$ . Here, we demonstrate how to do this for  $s = M$ . To simplify the notation we define.

$$\tilde{\mu}_i = \mu + \sum_{s=1}^{M-1} \Gamma_i^s \theta_s^i \quad (14)$$

and

$$K = -\frac{1}{2} \sum_{s=1}^{M-1} \sum_{i,j=1}^N \theta_s^i \theta_s^j \left( C^{(s)} \right)_{ij}^{-1}. \quad (15)$$

With these choices Eq.(5) becomes

$$\begin{aligned} \ell(\boldsymbol{\mu}, \boldsymbol{\theta}) &= -\frac{1}{2} \sum_{i,j=1}^N (y_i - \tilde{\mu}_i - \Gamma_i^M \theta_M^i) V_{ij}^{-1} (y_j - \tilde{\mu}_j - \Gamma_j^M \theta_M^j) \\ &\quad - \frac{1}{2} \sum_{i,j=1}^N \theta_M^i \theta_M^j \left( C^{(M)} \right)_{ij}^{-1} + K. \end{aligned} \quad (16)$$

To further simplify the derivation we define  $\tilde{\theta}_i = \Gamma_i^M \theta_M^i$  and  $\tilde{C}^{(M)} = C_{ij}^{(M)} \Gamma_i^M \Gamma_j^M$ . Thus the log-likelihood can be written in the simple form:

$$\begin{aligned} \ell(\boldsymbol{\mu}, \boldsymbol{\theta}) &= -\frac{1}{2} \sum_{i,j=1}^N (y_i - \tilde{\mu}_i - \tilde{\theta}_i) V_{ij}^{-1} (y_j - \tilde{\mu}_j - \tilde{\theta}_j) \\ &\quad - \frac{1}{2} \sum_{i,j=1}^N \tilde{\theta}_i \tilde{\theta}_j \left( \tilde{C}^{(M)} \right)_{ij}^{-1} + K. \end{aligned} \quad (17)$$

To profile over the  $\tilde{\theta}_i$ , we need to solve the following equations for  $\tilde{\theta}_i$  for all  $i$ :

$$\frac{\partial \ell}{\partial \tilde{\theta}_i} = \sum_j (y_j - \tilde{\mu}_j) V_{ij}^{-1} - \tilde{\theta}_j \left[ V_{ij}^{-1} + \left( \tilde{C}^{(M)} \right)_{ij}^{-1} \right] = 0. \quad (18)$$

This leads to the profiled values of the  $\tilde{\theta}_i$  parameters:

$$\hat{\tilde{\theta}}_i = \sum_{j,k} M_{ij}^{-1} V_{jk}^{-1} (y_k - \tilde{\mu}_k). \quad (19)$$

Here the entries of the matrix  $M$  are defined as

$$M_{ij} = V_{ij}^{-1} + \left( \tilde{C}^{(M)} \right)_{ij}^{-1}. \quad (20)$$

This matrix is symmetric and we will use this property to switch its indices for the proof. If we now substitute the profiled values  $\hat{\tilde{\theta}}_i$  back into the log-likelihood we obtain

$$\begin{aligned} \ell(\boldsymbol{\mu}, \boldsymbol{\theta}_1, \dots, \boldsymbol{\theta}_{M-1}, \hat{\tilde{\theta}}_M) &= -\frac{1}{2} \sum_{i,j=1}^N (y_i - \tilde{\mu}_i) V_{ij}^{-1} (y_j - \tilde{\mu}_j) \\ &\quad + \frac{1}{2} \sum_{i,j,p,q=1}^N (y_i - \tilde{\mu}_i) (y_j - \tilde{\mu}_j) V_{ip}^{-1} V_{jq}^{-1} M_{pq}^{-1} + K \\ &= -\frac{1}{2} \sum_{i,j=1}^N (y_i - \tilde{\mu}_i) \left[ V_{ij}^{-1} - \sum_{p,q=1}^N V_{ip}^{-1} M_{pq}^{-1} V_{qj}^{-1} \right] (y_j - \tilde{\mu}_j) + K. \end{aligned} \quad (21)$$

To further simplify this expression, the Woodbury identity [35] can be used to show that

$$\left[ V_{ij}^{-1} - \sum_{p,q=1}^N V_{ip}^{-1} M_{pq}^{-1} V_{qj}^{-1} \right]^{-1} = V_{ij} - \left( -M_{ij} + V_{ij}^{-1} \right)^{-1} = V_{ij} + \tilde{C}_{ij}. \quad (22)$$

Therefore we find

$$\ell(\boldsymbol{\mu}, \boldsymbol{\theta}_1, \dots, \boldsymbol{\theta}_{M-1}, \widehat{\boldsymbol{\theta}}_M) = -\frac{1}{2} \sum_{i,j=1}^N (y_i - \tilde{\mu}_i) \left[ V_{ij} + \tilde{C}_{ij} \right]^{-1} (y_j - \tilde{\mu}_j) + K, \quad (23)$$

where  $\tilde{C}_{ij}^{(M)} = C_{ij}^{(M)} \Gamma_i^M \Gamma_j^M = \rho_{ij}^{(M)} \Gamma_i^M \Gamma_j^M \sigma_{u_M}^2$  represents the  $M$ -th term in the systematic part of the BLUE covariance matrix  $W$  as defined in Eq. (5), i.e.,  $U^{(M)} = \tilde{C}^{(M)}$ . This procedure can be recursively applied to compute all terms in the BLUE covariance matrix corresponding to every source of systematic uncertainty. Thus, one obtains  $W = V + \sum_s^M \tilde{C}^{(s)} = V + \sum_s^M U^{(s)}$ .

## B Derivation of the GVM profile log-likelihood for combinations

The assumption underlying the construction of the likelihood defined in Eq. (9), which includes correlations between the auxiliary measurements, is that the  $\sigma_{u_s}^2$  parameters outlined in Eq. (6) are unknown model parameters, and their best estimates,  $v_s$ , follow a Gamma distribution:

$$v_s \sim \frac{\beta_s^{\alpha_s}}{\Gamma(\alpha_s)} v_s^{\alpha_s-1} e^{-\beta_s v_s}, \quad (24)$$

where  $\alpha_s = \frac{1}{4\varepsilon_s^2}$  and  $\beta_s = \frac{1}{4\sigma_{u_s}^2 \varepsilon_s^2}$ . This assumption extends the likelihood in Eq. (7) to incorporate new terms arising from the Gamma distributions:

$$\begin{aligned} \ell(\boldsymbol{\mu}, \boldsymbol{\theta}, \boldsymbol{\sigma}_u^2) &= -\frac{1}{2} \sum_{i,j=1}^N \left( y_i - \mu - \sum_{s=1}^M \Gamma_i^s \theta_s^i \right) V_{ij}^{-1} \left( y_j - \mu - \sum_{s=1}^M \Gamma_j^s \theta_s^j \right) \\ &\quad - \frac{1}{2} \sum_{s=1}^M \sum_{i,j=1}^N (u_s^i - \theta_s^i) \left( C^{(s)} \right)_{ij}^{-1} (u_s^j - \theta_s^j) \\ &\quad - \frac{1}{2} \sum_{s=1}^M \left[ \left( N + \frac{1}{2\varepsilon_s^2} \right) \log \sigma_{u_s}^2 + \frac{v_s}{2\varepsilon_s^2 \sigma_{u_s}^2} \right], \end{aligned} \quad (25)$$

where some constant factors that do not depend on the model parameters are omitted for simplicity. From this expression, it is evident that one can redefine  $\sigma_{u_s}^2$ ,  $v_s$ ,  $\theta_s^i$ ,  $u_s^i$ , and the factor  $\Gamma_i^s$  so that  $v_s = 1$ . This adjustment does not alter the structure of the likelihood, as it merely introduces some constant terms that are independent of the adjustable parameters of the model.

To derive Eq. (9), we must profile over the  $\boldsymbol{\sigma}_u^2$  NPs. This involves computing their profiled values:

$$\widehat{\sigma_{u_s}^2} = \frac{v_s + 2\varepsilon_s^2 \sum_{i,j}^N (u_s^i - \theta_s^i) \left( C_{ij}^{(s)} \right)^{-1} (u_s^j - \theta_s^j)}{1 + 2N\varepsilon_s^2}. \quad (26)$$

These values can now be substituted back into Eq.(25) to derive Eq.(7), yielding the desired result. The log-likelihood defined by Eq. (7) is the same result one would get by modelling the auxiliary measurements  $u_s^i$  as multivariate student's t distributed variables with expected values  $\theta_s^i$ , shape matrix  $C^{(s)}$  and degrees of freedom  $\nu_s = 1/2\varepsilon_s^2$ .