

A DEFINITIVE DETERMINANT MAJORIZATION RESULT FOR NONLINEAR OPERATORS

F. REESE HARVEY AND H. BLAINE LAWSON, JR.

*Dedicated to John Polking
on the occasion of his ninetieth birthday*

Abstract

Let \mathfrak{g} be a Gårding-Dirichlet operator on the set $\mathcal{S}(n)$ of symmetric $n \times n$ matrices. We assume that \mathfrak{g} is I -central, that is, $D_I \mathfrak{g} = kI$ for some $k > 0$. Then

$$\mathfrak{g}(A)^{\frac{1}{N}} \geq \mathfrak{g}(I)^{\frac{1}{N}} (\det A)^{\frac{1}{n}} \quad \forall A > 0.$$

From work of Guo, Phong, Tong, Abja, Dinew, Olive and many others, this inequality has important applications.

Date: July 9, 2024.

Second author was partially supported by the Simons Foundation.

1. Introduction.

This paper is concerned with Gårding-Dirichlet operators. These are nonlinear elliptic differential operators on euclidean space, and in many important cases on riemannian manifolds, hermitian manifolds, etc. (This is discussed in detail in [HL₁].) These operators are defined by applying the following polynomials to the Hessians of functions. Let $\mathcal{S}(n)$ denote the space of second derivatives, i.e., the vector space of $n \times n$ real symmetric matrices.

Definition 1.1. Suppose \mathfrak{g} is a real homogeneous polynomial of degree N on $\mathcal{S}(n)$ which is I -hyperbolic, which means that $\mathfrak{g}(I) > 0$ and for each $A \in \mathcal{S}(n)$, the polynomial $\varphi_A(t) \equiv \mathfrak{g}(tI + A)$ has only real roots. By the **Gårding cone** Γ of \mathfrak{g} we mean the connected component of $\{\mathfrak{g} > 0\}$ containing I . If Γ contains $\text{Int}\mathcal{P} = \{A : A > 0\}$, then \mathfrak{g} is called a **Gårding-Dirichlet polynomial** or a **Gårding-Dirichlet polynomial operator**.

In [HL₄] we proved a determinant majorization formula for such polynomials provided they had certain invariance properties. This had applications to work of B. Guo, D. Phong, F. Tong [GPT], [GP₁], [GP₂], [GP₃] and S. Abja, S. Dinew, G. Olive [AO], [ADO]. In fact, the important paper [GPT] has lead to much subsequent work. The reader could check recent papers [GP₄] and [KLW] for history and references.

The point of this paper is to prove a more definitive result that the majorization formula holds for a wide class of Gårding-Dirichlet polynomials where the invariance property is not needed. One needs only to check the Central Ray Hypothesis, which is generally very easy to do. (See Section 3.)

Definition 1.2. The **Central Ray Hypothesis**¹ is that:

*The gradient of \mathfrak{g} at I satisfies $D_I \mathfrak{g} = kI$ for some $k > 0$.*²

This is equivalent to the following:

The \mathfrak{g} -Laplacian $\Delta^{\mathfrak{g}}$, defined to be the sum of the I -eigenvalues of \mathfrak{g} , is a positive constant times the standard Laplacian Δ^{std} .

If this hypothesis is satisfied, we say that \mathfrak{g}, Γ is **I -central**.

THE MAIN THEOREM 1.3. *If \mathfrak{g}, Γ is an I -central Gårding-Dirichlet polynomial on $\mathcal{S}(n)$, then the determinant is majorized:*

$$\mathfrak{g}(A)^{\frac{1}{N}} \geq \mathfrak{g}(I)^{\frac{1}{N}} (\det A)^{\frac{1}{n}} \quad \forall A > 0. \quad (1.1)$$

Of course, the constant $\mathfrak{g}(I)^{\frac{1}{N}}$ is the best possible.

¹This concept was first introduced in [HL₄] with several equivalent formulations in Theorem E.8.

²In this paper $D_A F$ denotes the gradient of F at the point A

Note. 1.4. In our former paper [HL₄] we needed that the polynomial we were considering had all of its coefficients positive. This is not true of even the simplest Gårding-Dirichlet polynomial, the determinant of a 2×2 -matrix: $\det A = a_{11}a_{22} - a_{12}^2$. As a result we assumed orthogonal or unitary invariance (as in [HL₄]), and this allowed us to work in eigenvalue space, where all the coefficients are ≥ 0 .

Note. 1.5. The Main Theorem can be viewed in the following way. In [[HL₅], Prop. 11.4], the authors show the following enhancement of Lemma 4 in [GP₁].

Proposition 1.6. *Let \mathfrak{g} be a Gårding-Dirichlet operator of degree N with Gårding cone $\Gamma \subset \mathcal{S}(n)$. For any constant $\gamma > 0$, the following are equivalent:*

(1) (Majorization of the Determinant)

$$\mathfrak{g}(A)^{\frac{1}{N}} \geq n\gamma^{\frac{1}{n}}(\det A)^{\frac{1}{n}} \quad \forall A > 0$$

(2) (Lower Bound for the Determinant of the Gradient)

$$D_B \mathfrak{g}^{\frac{1}{N}} > 0 \quad \text{and} \quad \det(D_B \mathfrak{g}^{\frac{1}{N}}) \geq \gamma \quad \forall B \in \Gamma.$$

Corollary 1.7. *The conclusion (1.1) of the Main Theorem can be restated in the following equivalent form:*

$$D_B \mathfrak{g}^{\frac{1}{N}} > 0 \quad \text{and} \quad \det(D_B \mathfrak{g}^{\frac{1}{N}}) \geq \gamma \equiv \frac{\mathfrak{g}(I)^{\frac{n}{N}}}{n^n}, \quad \forall B \in \Gamma. \quad (1.1)'$$

2. The Proof.

There are two steps. First we give the proof of (1.1) when A is diagonal by using the Basic Lemma 2.1 in [HL₄] (see also Gurvits [Gu]). Then we use this diagonal case to prove the general result.

This Basic Lemma can be stated in the following slightly simpler and equivalent form. Let $e \equiv (1, 1, \dots, 1) \in \mathbb{R}^n$.

Basic Lemma 2.1. Suppose p is a homogeneous polynomial of degree $N > 0$ on \mathbb{R}^n .

If

(1) the coefficients of $p(x)$ and all non-negative, and

(2) (The Central Ray Hypothesis)

$$D_e p = k e \quad \text{for some } k > 0 \quad (2.1)$$

are both satisfied, then

$$p(x)^{\frac{1}{N}} \geq p(e)^{\frac{1}{N}} (x_1 \cdots x_n)^{\frac{1}{n}}, \quad \forall x_1 > 0, x_2 > 0, \dots, x_n > 0. \quad (2.2)$$

Note. By Euler's Formula for homogeneous polynomials we have $Np(e) = \langle D_e p, e \rangle$, so if the Central Ray Hypothesis (2.1) holds, then N, k, n and $p(e)$ are related by

$$Np(e) = kn.$$

Our two step procedure is formalized in Theorem 2.3 below where \mathbf{g} is replaced by any real homogeneous polynomial F on $\mathcal{S}(n)$. Let

$$p(x) \equiv F(X),$$

$$\text{where } X \equiv \begin{pmatrix} x_1 & 0 & \cdots \\ 0 & x_2 & \cdots \\ 0 & \cdots & x_n \end{pmatrix} \text{ and } x = (x_1, \dots, x_n) \in \mathbb{R}^n, \quad (2.3)$$

be the restriction of F to the set of diagonal matrices $\mathbb{D} \subset \mathcal{S}(n)$. Next, for each orthogonal transformation $h \in O(n)$, set

$$F_h(A) \equiv F(hAh^t) \quad \text{and} \quad p_h(x) \equiv F_h(X) = F(hXh^t) \quad (2.4)$$

for $X \in \mathbb{D}$.

Part (c) of the next result provides an equivalent statement of the Central Ray Hypothesis in terms of the polynomials p_h for $h \in O(n)$.

Lemma 2.2.

- (a) $D_I F_h = h^t(D_I F)h$
- (b) $D_I F_h = kI \iff D_I F = kI$
- (c) $D_I F_h = kI \iff D_e p_h = ke \quad \forall h \in O(n)$.

Proof. For Part (a)

$$\begin{aligned} \langle D_I F_h, A \rangle &= \left. \frac{d}{dt} \right|_{t=0} F_h(I + tA) = \left. \frac{d}{dt} \right|_{t=0} F(h(I + tA)h^t) \\ &= \left. \frac{d}{dt} \right|_{t=0} F(I + thAh^t) = \langle D_I F, hAh^t \rangle \\ &= \langle h^t(D_I F)h, A \rangle, \quad \forall A \in \mathcal{S}(n) \end{aligned}$$

Part (b) is immediate from Part (a).

Proof of \Rightarrow in Part (c):

$$\langle D_e p_h, x \rangle \equiv \left. \frac{d}{dt} \right|_{t=0} p_h(e + tx) = \left. \frac{d}{dt} \right|_{t=0} F_h(I + tX) = \langle D_I F_h, X \rangle$$

for all $X \in \mathbb{D}$. Hence, $D_I F = kI \Rightarrow D_e p_h = ke$ for all $h \in O(n)$ since $\langle kI, X \rangle = \langle ke, x \rangle$.

Proof of \Leftarrow in Part (c): Conversely, suppose $D_e p_h = ke$ for all $h \in O(n)$. Given $A \in \mathcal{S}(n)$, pick $h \in O(n)$ so that

$$h^t A h = \Lambda \equiv \begin{pmatrix} \lambda_1 & 0 & \cdots \\ 0 & \lambda_2 & \cdots \\ 0 & \cdots & \lambda_n \end{pmatrix} \text{ is diagonal.} \quad (2.5)$$

Then with $\lambda = (\lambda_1, \dots, \lambda_n)$, we have

$$\begin{aligned} \langle D_I F, A \rangle &= \langle D_I F, h \Lambda h^t \rangle = \langle h^t (D_I F) h, \Lambda \rangle \quad \text{which by (a)} \\ &= \langle D_I F_h, \Lambda \rangle = \left. \frac{d}{dt} \right|_{t=0} F_h(I + t\Lambda) \\ &= \left. \frac{d}{dt} \right|_{t=0} p_h(e + t\lambda) = \langle D_e p_h, \lambda \rangle. \end{aligned}$$

By the right hand side of part (c) of Lemma 2.2 this last term equals $\langle ke, \lambda \rangle = \langle kI, A \rangle$. \blacksquare

Continuing with the notation above, we have the following.

THEOREM 2.3. *Suppose F is a real homogeneous polynomial of degree N on $\mathcal{S}(n)$ which is*

- (1) *I -central, (See Definition 1.2.)*

*and satisfies the following **coefficient condition**:*

- (2) *For each $h \in O(n)$ the restriction p_h of F_h to the diagonal has coefficients ≥ 0 .*

Then

$$F(A)^{\frac{1}{N}} \geq F(I)^{\frac{1}{N}} (\det A)^{\frac{1}{n}} \quad \forall A > 0. \quad (2.6)$$

Note: Examples show that the polynomials $p_h(x)$ depend on the choice of $F(A)$, not just $p(x)$. For example, let $F(A) = a_{11}a_{22} - ca_{12}^2$ with $0 \leq c < 1$.

Proof. Given A choose $h \in O(n)$ so that (2.5) is satisfied. Then $F(A) = F(h \Lambda h^t) = p_h(\lambda)$.

Since F is I -central, Lemma 2.2 (b) and (c) implies that p_h is e -central. Hypothesis (2) says that p_h has coefficients ≥ 0 . Therefore by the Basic Lemma 2.1 we have

$$p_h(\lambda)^{\frac{1}{N}} \geq p_h(e)^{\frac{1}{N}} (\lambda_1 \cdots \lambda_n)^{\frac{1}{n}}.$$

Substituting $p_h(\lambda) = F(A)$, $p_h(e) = F(I)$ and $\lambda_1 \cdots \lambda_n = \det A$ yields the conclusion (2.6). \blacksquare

To prove the Main Theorem 1.3 it remains to show that for any I -central Gårding-Dirichlet operator $F = \mathfrak{g}$, the coefficient hypothesis (2) of Theorem 2.3 is satisfied. First we note the following.

Lemma 2.4. *If \mathfrak{g} is Gårding-Dirichlet and $h \in \mathrm{GL}_n(\mathbb{R})$, then $\mathfrak{g}_h(A) \equiv \mathfrak{g}(hAh^t)$ is also Gårding-Dirichlet.*

Proof. The I -eigenvalues of $A \in \mathcal{S}(n)$ for the homogeneous polynomial \mathfrak{g}_h are computed as follows:

$$\begin{aligned} \mathfrak{g}_h(tI + A) &= \mathfrak{g}(h(tI + A)h^t) = \mathfrak{g}(thh^t + hAh^t) \\ &= \mathfrak{g}(hh^t) \prod_j (t + \lambda_j^{hh^t, \mathfrak{g}}(hAh^t)) \\ &= \mathfrak{g}_h(I) \prod_j (t + \lambda_j^{hh^t, \mathfrak{g}}(hAh^t)). \end{aligned} \tag{2.7}$$

Thus, these eigenvalues are equal to the hh^t -eigenvalues of hAh^t for \mathfrak{g} . Note that $hh^t > 0$ so $hh^t \in \Gamma(\mathfrak{g})$. This proves that the eigenvalues $\lambda^{I, \mathfrak{g}_h}(A)$ are real, so that if \mathfrak{g} is a Gårding polynomial, then \mathfrak{g}_h is also a Gårding polynomial. Using the fact that the Gårding cone is the set on which all N eigenvalues are strictly positive, this proves that

$$A \in \Gamma(\mathfrak{g}_h) \iff hAh^t \in \Gamma(\mathfrak{g}), \quad \text{i.e.} \quad \Gamma(\mathfrak{g}_h) = h^{-1}\Gamma(\mathfrak{g})(h^t)^{-1}$$

Finally, since $\mathrm{Int}\mathcal{P} \subset \Gamma(\mathfrak{g})$ and $h^{-1}(\mathrm{Int}\mathcal{P})(h^t)^{-1} = \mathrm{Int}\mathcal{P}$, this proves that $\mathrm{Int}\mathcal{P} \subset \Gamma(\mathfrak{g}_h)$. Therefore, \mathfrak{g}_h is Gårding-Dirichlet. \blacksquare

Note. See Appendix A for a further discussion, including when the Gårding-Dirichlet operator \mathfrak{g}_h , produced by the change of variables $h \in \mathrm{GL}(n, \mathbb{R})$, is I -central.

It remains to show that the coefficient condition (2) holds for \mathfrak{g}_h . However, as a consequence of Lemma 2.4 the polynomial \mathfrak{g}_h is Gårding-Dirichlet, so we only need to show that condition (2) holds for all Gårding-Dirichlet polynomials, which is our final lemma.

Lemma 2.5. *For any Gårding-Dirichlet polynomial \mathfrak{g} , the restriction*

$$p(x) = \mathfrak{g}(X)$$

to the diagonal $X \in \mathbb{D}$, has coefficients ≥ 0 .

Proof. We use the following basic construction of Gårding along with the fact that

$$\frac{1}{\alpha_1! \cdots \alpha_n!} \frac{\partial^{|\alpha|} p}{\partial x_1^{\alpha_1} \cdots \partial x_n^{\alpha_n}} \tag{2.8}$$

equals the coefficient a_α for each multi-index α of length $|\alpha| \equiv \sum_k \alpha_k = N$.

We recall that the a -eigenvalues of p at $x \in \mathbb{R}^n$ are given by the equation $p(x + ta) = p(a) \prod_{k=1}^N (t + \lambda_k^p(x))$.

Proposition 2.6. (Gårding [Ga]). *Suppose p, Γ is a Gårding polynomial on $V = \mathbb{R}^n$ of degree $N \geq 2$ with $\mathbb{R}_{>0}^n \subset \Gamma$. Then for $a \in \Gamma$ the polynomial*

$$q(x) \equiv \langle D_x p, a \rangle = \left. \frac{d}{dt} p(x + ta) \right|_{t=0} = \sum_{j=1}^n a_j \frac{\partial p}{\partial x_j}(x)$$

is also Gårding, of degree $N - 1$. Moreover, the a -eigenvalues of p and q at $x \in \mathbb{R}^n$ satisfy:

$$\lambda_1^p(x), \dots, \lambda_N^p(x) > 0 \quad \Rightarrow \quad \lambda_1^q(x), \dots, \lambda_{N-1}^q(x) > 0 \quad (2.9)$$

which is equivalent to

$$\Gamma(q) \supset \Gamma(p) \supset \mathbb{R}_{>0}^n. \quad (2.10)$$

Proof. For $a \in \Gamma$ we know that p is hyperbolic in the direction a , that is, for every $x \in \mathbb{R}^n$ the polynomial $t \mapsto p(x + ta)$ has N real roots. For every $x \in \mathbb{R}^n$ the roots of

$$q(x + ta) = \frac{d}{dt} p(x + ta)$$

are the critical points of $p(x + ta)$. Hence (by the Mean Value Theorem), the a -eigenvalues of q interlace the a -eigenvalues of p for each $x \in \mathbb{R}^n$. This proves that q is Gårding of degree $N - 1$. Moreover $\Gamma(p) \subset \Gamma(q)$, since if x has all positive p eigenvalues, the interlacing q -eigenvalues at x must also be positive. \blacksquare

We shall use Proposition 2.6 to prove that each α^{th} partial of p for $|\alpha| = N$ is ≥ 0 . The partial derivative $\frac{\partial p}{\partial x_k}(x)$ is just the directional derivative in the axis direction e_k . We approximate the α derivative

$$\left(\frac{\partial}{\partial x_1} \right)^{\alpha_1} \cdots \left(\frac{\partial}{\partial x_n} \right)^{\alpha_n} = (\nabla_{e_1})^{\alpha_1} \cdots (\nabla_{e_n})^{\alpha_n}$$

with the directional derivative in Gårding directions $e_j + \epsilon e \in \Gamma$

$$(\nabla_{e_1 + \epsilon e})^{\alpha_1} \cdots (\nabla_{e_n + \epsilon e})^{\alpha_n}$$

for $\epsilon > 0$. We now take these derivatives in succession, that is, one at a time. Since every $e_k + \epsilon e$ lies in the open set $\mathbb{R}_{>0}^n$, Proposition 2.6 says that each time we take a derivative, we get another Gårding polynomial, of exactly one degree less, whose Gårding cone contains the one we just had. In particular it contains $\mathbb{R}_{>0}^n$. After $N - 1$ steps we have a linear Gårding polynomial $h(x) \equiv b \cdot x$, $b \in \mathbb{R}^n$ with $b \neq 0$, such that the half-space $\Gamma(h) \supset \mathbb{R}_{>0}^n$. In particular, the inner product $\langle b, e_j + \epsilon e \rangle > 0$ for each j . Letting $\epsilon \rightarrow 0$ we get that the j^{th} component of b , $b_j = \langle b, e_j \rangle \geq 0$ for all j . Therefore, at

the last step, where we take the final derivative, we have a constant which is ≥ 0 . This shows that the α^{th} partial of p at x is ≥ 0 , and Lemma 2.5 is proven. ■

Note. In light of the fact (proved in Theorem D.1 (1)' of [HL₄]) that the directional derivative of \mathfrak{g} is strictly positive in all directions B in the closed cone $\overline{\Gamma(\mathfrak{g})}$, except the edge directions, one might think that the coefficients $a_\alpha > 0$ for all $|\alpha| = N$ unless one of the axes belongs to the edge. However this is not true, as the Gårding polynomial $\mathfrak{g}(x) = x_1(\epsilon x_1 + x_2)$ in \mathbb{R}^2 shows.

Proof of the MAIN THEOREM 1.3. Lemma 2.5 shows that for each of the Gårding-Dirichlet operators \mathfrak{g}_h , the restriction of \mathfrak{g}_h to the diagonal \mathbb{D} has coefficients ≥ 0 . That is, the coefficient hypothesis (2) of Theorem 2.3 is satisfied, and so Theorem 1.3 holds. ■

3. Examples.

In this section we examine the family of operators to which our theorems apply. In the restricted cases, considered in [HL₄], the main result already covered a wide universe of examples. This was due, in part, because products and radial derivatives take examples to new examples (See Lemmas 7.1 and Proposition 3.2 in [HL₄]). There is a parallel situation here without the invariance.

Proposition 3.1.

(1) *The product of two I -central Gårding-Dirichlet operators on $\mathcal{S}(n)$ is again an I -central Gårding-Dirichlet operator.*

(2) *The radial derivative (or relaxation) of a Gårding-Dirichlet operator on $\mathcal{S}(n)$ is also an I -central Gårding-Dirichlet operator.*

Note that in (1) the degree goes up, whereas in (2) it goes down. Hence one can take long sequences of these operations with many "ups and downs" of the degree. This leads to a large family of operations to which Theorem 1.3 applies.

There is a third, very general way of producing examples which includes both constructions in the two steps of our proof of the Main Theorem 2.3. This is a third part of Proposition 3.1 which is presented at the end of this section §3.

Proof of Proposition 3.1 (1). Suppose \mathfrak{g} and \mathfrak{h} are I -hyperbolic on $\mathcal{S}(n)$, that is $\mathfrak{g}(tI + A)$ and $\mathfrak{h}(tI + A)$ have only real roots. Since the roots of the product $\mathfrak{g}\mathfrak{h}$ is the union of the roots of \mathfrak{g} and \mathfrak{h} , we see that $\mathfrak{g}\mathfrak{h}$ is also I -hyperbolic.

Recall the Gårding cone $\Gamma^{\mathfrak{g}}$ of \mathfrak{g} is the connected component of $\{\mathfrak{g} > 0\}$ containing I , and the Dirichlet condition is that $\mathcal{P} \subset \Gamma^{\mathfrak{g}}$. Of course, the same holds for \mathfrak{h} , so clearly $\mathcal{P} \subset \Gamma^{\mathfrak{g}} \cap \Gamma^{\mathfrak{h}} = \Gamma^{\mathfrak{gh}}$ since the Gårding cones are convex. Hence, the product \mathfrak{gh} satisfies the Dirichlet condition and is therefore Gårding-Dirichlet.

It remains to prove that \mathfrak{gh} is I -central. As discussed in §1, this condition has several important equivalent formulations. One such is that the **Gårding Laplacian** $\Delta^{\mathfrak{g}}A \equiv \sum_{j=1}^N \lambda_j^{\mathfrak{g}}(A)$ and the standard Laplacian $\Delta A \equiv \sum_{j=1}^N \lambda_j(A)$ agree up to a positive factor $\bar{k} > 0$, i.e., $\Delta^{\mathfrak{g}} = \bar{k}\Delta$. (See Appendix E in [HL₄] for a complete discussion of the Central Ray Hypothesis including several other equivalent ways of stating it.) Since \mathfrak{g} and \mathfrak{h} are I -central, we have $\Delta^{\mathfrak{g}} = \bar{k}\Delta$ and $\Delta^{\mathfrak{h}} = \bar{\ell}\Delta$ for positive constants $\bar{k}, \bar{\ell}$. Since the set of eigenvalues of \mathfrak{gh} is exactly the union of the eigenvalues of \mathfrak{g} and \mathfrak{h} , it follows that $\Delta^{\mathfrak{gh}} = \Delta^{\mathfrak{g}} + \Delta^{\mathfrak{h}} = (\bar{k} + \bar{\ell})\Delta$, and so \mathfrak{gh} is I -central. ■

Remark 3.2. (Concerning the Central Ray Hypothesis) *The Central Ray Hypothesis is a weak form of symmetry for the operator. As seen in [HL₄], Gårding-Dirichlet operators invariant by $O(n)$, $U(n/2)$ or $Sp(n/4)$ are I -central.*

If \mathfrak{g} is I -central, then \mathfrak{g} is complete, that is, it depends on all the variables (\iff it is not pulled back from a proper vector subspace). This is one reason to anticipate that I -central Gårding-Dirichlet operators have interior regularity. Certainly, without completeness, interior regularity fails utterly. If, for example, $\mathfrak{g}(A)$ is independent of A_{1j} , $j = 1, \dots, n$, then any continuous function of x_1 is \mathfrak{g} -harmonic.

Proof of Proposition 3.1 (2). Gårding's result that the radial derivative $\mathfrak{g}'(tI + A) = \frac{d}{dt}\mathfrak{g}(tI + A)$ is again Gårding-Dirichlet is proved in Lemma 2.6. To see that \mathfrak{g}' is I -central, we consider the family of radial derivatives $\mathfrak{g}^{(\ell)}(tI + A) = \frac{d^\ell}{dt^\ell}\mathfrak{g}(tI + A)$. Following [Ga] and [Ren], the roots of $\mathfrak{g}^{(\ell+1)}(tI + A)$ are the critical points of $\mathfrak{g}^{(\ell)}(tI + A)$ and hence intertwine the roots of $\mathfrak{g}^{(\ell)}(tI + A)$. Hence, the associated Gårding cones are nested: $\Gamma^{\mathfrak{g}} \subset \Gamma^{\mathfrak{g}'} \subset \Gamma^{\mathfrak{g}''} \subset \dots \subset \Gamma^{\mathfrak{g}^{(N-1)}} = H$, a half space. Since \mathfrak{g} is I -central, $H = \{A : \Delta(A) \geq 0\}$. Now for any ℓ we have that $(\mathfrak{g}^{(\ell)})^{(N-\ell-1)} = \mathfrak{g}^{(N-1)}$, from which it follows that $\Delta^{\mathfrak{g}^{(\ell)}} = \bar{k}\Delta$, for some $\bar{k} > 0$. ■

Remark 3.3. Comments on the Coefficient Condition (2) of Theorem 2.3. There are plenty of examples of polynomial operators F satisfying the hypotheses (1) and (2) of Theorem 2.3, and therefore the determinant majorization (2.6), but which are NOT Gårding. For instance

$$F(A) \equiv \|A\|^2 \det A$$

is $O(n)$ -invariant, so that $F_h = F$ and hence $p_h(x) = p(x) = |x|^2 x_1 \cdots x_n$ has coefficients ≥ 0 for all $h \in O(n)$. However, it remains to determine how interesting they are.

We now want to discuss the following specific example.

Example 3.4. (The Lagrangian Monge-Ampère Operator). This is an Gårding-Dirichlet operator MA_{LAG} of degree $N = 2^n$ on $\mathbb{R}^{2n} = \mathbb{C}^n$, which was introduced in [HL₃]. It makes sense on any symplectic manifold with a Gromov metric.

There are two reasons for updating this particular case. A determinantal inequality for this operator was done in Proposition 8.1 in [HL₄] with a proof independent of the main theorem in that paper. Here we show that it is a direct consequence of Theorem 1.3 above. Moreover, the inequality established in [HL₄] for MA_{LAG} was not sharp, while the one given by Theorem 3.1 is.

In [HL₄] we proved the determinantal inequality

$$\text{MA}_{\text{LAG}}(A)^{\frac{1}{N}} \geq \det(A)^{\frac{1}{2n}} \quad \forall A > 0. \quad (3.1)$$

It is easy to compute that $\text{MA}_{\text{LAG}}(I)^{\frac{1}{N}} = n$

Lemma 3.5. *The operator MA_{LAG} is I -central, and hence*

$$\text{MA}_{\text{LAG}}(A)^{\frac{1}{N}} \geq n \det(A)^{\frac{1}{2n}} \quad \forall A > 0. \quad (3.2)$$

This is an improvement of (3.1) and is sharp (one has equality when $A = I$). By the way the proof of this Lemma is easier than the proof of Prop. 8.1 in [HL₄].

Proof. The I -eigenvalues of $A \in \mathcal{S}(n)$ for MA_{LAG} were computed in Lemma 3.7 on [HL₃], and they are

$$\Lambda_{\pm\pm\cdots\pm}(A) = \frac{\text{tr } A}{2} \pm \lambda_1 \pm \lambda_2 \pm \cdots \pm \lambda_n \quad (3.3)$$

where $\lambda_1, \dots, \lambda_n$ are the non-negative eigenvalues of the skew-hermitian part of A . If we sum (3.3) over the 2^n choices of the \pm signs, the λ_j 's drop out, and we have

$$\sum_{\pm} \Lambda_{\pm\pm\cdots\pm}(A) = 2^{n-1} \text{tr } A. \quad (3.4)$$

So the Gårding Laplacian for MA_{LAG} is just 2^{n-1} times the standard Laplacian of A . This is one of the equivalent ways of saying that MA_{LAG} is I -central (See Definition 1.2.). \blacksquare

Finally we add a third method of constructing examples, which will give a completion to Proposition 3.2. The two methods used in the proof of the Main Theorem to construct Gårding-Dirichlet operators (Lemma 2.4 and

Proposition 2.6) can be combined by replacing the maps $\mathcal{S}(n) \rightarrow \mathbb{D}$ and $A \mapsto hAh^t$, by a general linear transformation $L : \mathcal{S}(n) \rightarrow \mathcal{S}(n)$. This gives us a third method of constructing I -central Gårding-Dirichlet operators, which we address as a third conclusion (3) to Proposition 3.1.

Proposition 3.1. *Under the assumptions of Proposition 3.1 one has the following.*

(3) Suppose $L \in \text{End}(\mathcal{S}(n))$ and \mathfrak{g} is a homogeneous polynomial on $\mathcal{S}(n)$ of degree N . Consider the transform $\mathfrak{g}_L(A) \equiv \mathfrak{g}(L(A))$, which is also homogeneous of degree N . Then

(a) \mathfrak{g}_L is B -hyperbolic $\iff \mathfrak{g}$ is $L(B)$ -hyperbolic (for $B \in \mathcal{S}(n)$), in which case the B -eigenvalues are given by

(b) $\lambda_j^{B, \mathfrak{g}_L}(A) = \lambda_j^{L(B), \mathfrak{g}}(L(A))$, $j = 1, \dots, N$, and the Gårding cone is given by

(c) $\Gamma^{B, \mathfrak{g}_L} = L^{-1}(\Gamma^{L(B), \mathfrak{g}})$, i.e., $A \in \Gamma^{B, \mathfrak{g}_L} \iff L(A) \in \Gamma^{L(B), \mathfrak{g}}$.

The gradient map for \mathfrak{g}_L is given by

(d) $D_A(\log \mathfrak{g}_L) = L^t(D_{L(A)} \log \mathfrak{g})$.

(e) If $\mathfrak{g}, \Gamma(\mathfrak{g})$ is Gårding-Dirichlet and $L(\mathcal{P}) \subset \overline{\Gamma(\mathfrak{g})}$, then $\mathfrak{g}_L, \Gamma(\mathfrak{g}_L)$ where $\Gamma(\mathfrak{g}_L) = \Gamma^{L, \mathfrak{g}_L}(\mathfrak{g}_L) = L^{-1}(\Gamma(\mathfrak{g}))$ is also Gårding-Dirichlet.

Finally, If $L(I) = kI$ for some $k > 0$, and if $\text{tr} A = 0 \Rightarrow \text{tr}(L(A)) = 0$, then

(f) \mathfrak{g} is I -central $\Rightarrow \mathfrak{g}_L$ is I -central.

Proof of (a) and (b). Note that

$$\mathfrak{g}_L(tB + A) = \mathfrak{g}_L(B) \prod_{j=1}^N (t + \lambda_j^{B, \mathfrak{g}_L}(A)), \quad \text{where}$$

$$\mathfrak{g}_L(tB + A) \equiv \mathfrak{g}(tL(B) + L(A)) = \mathfrak{g}(L(B)) \prod_{j=1}^N (t + \lambda_j^{L(B), \mathfrak{g}}(L(A)))$$

just as in the proof of Lemma 2.4. This proves (a) and (b). \blacksquare

Proof of (c). One has that $\mathfrak{g}_L(B) > 0 \iff \mathfrak{g}(L(B)) > 0$ and

$$\begin{aligned} A \in \Gamma^{B, \mathfrak{g}_L} &\iff \mathfrak{g}_L(A) > 0 \text{ and } \lambda_j^{B, \mathfrak{g}_L}(A) > 0, \quad j = 1, \dots, N \\ &\iff \mathfrak{g}(L(A)) > 0 \text{ and } \lambda_j^{L(B), \mathfrak{g}}(L(A)) > 0, \quad j = 1, \dots, N \\ &\iff L(A) \in \Gamma^{L(B), \mathfrak{g}}. \quad \blacksquare \end{aligned}$$

Proof of (d). Computing the chain rule for $\mathcal{S}(n) \xrightarrow{L} \mathcal{S}(n) \xrightarrow{\log \mathfrak{g}} \mathbb{R}$ at $A \in \mathcal{S}(n)$ one has

$$\langle D_A \log \mathfrak{g}_L, B \rangle = \langle D_{L(A)} \log \mathfrak{g}, L(B) \rangle = \langle L^t D_{L(A)} \log \mathfrak{g}, B \rangle,$$

for each $B \in \mathcal{S}(n)$. ■

Proof of (e). Given $P > 0$, we must show that $P \in \Gamma(\mathfrak{g}_L) = L^{-1}(\Gamma(\mathfrak{g}))$, i.e., $L(P) \in \Gamma(\mathfrak{g})$. We are assuming $L(P) > 0$, and since $\mathfrak{g}, \Gamma(\mathfrak{g})$ is Gårding-Dirichlet, we have $L(P) \in \Gamma(\mathfrak{g})$. ■

Proof of (f). By (d) with $A = I$, we have

$$D_I \log \mathfrak{g}_L = L^t(D_{L(I)} \log \mathfrak{g}) = L^t(D_{kI} \log \mathfrak{g}) = \frac{1}{k} L^t(D_I \log \mathfrak{g}).$$

Since I is the central ray for \mathfrak{g} , this equals $\frac{N}{kn} L^t(I)$.

It remains to show that $L^t(I) = kI$. Suppose $A \in \mathcal{S}(n)$, with trace-free part A_0 , i.e., $A \equiv \frac{\text{tr} A}{n} I + A_0$. By hypothesis, $\text{tr} A_0 = 0 \Rightarrow \text{tr} L(A_0) = \langle I, L(A_0) \rangle = 0$. Therefore, $\langle L^t(I), A \rangle = \langle I, L(A) \rangle = \langle I, k \frac{\text{tr} A}{n} I \rangle = \langle kI, A \rangle$ for all $A \in \mathcal{S}(n)$. ■

Bibliography

[AO] S. Abja and G. Olive, *Local regularity for concave homogeneous complex degenerate elliptic equations dominating the Monge-Ampère equation*, Ann. Mat. Pura Appl. (4) **201** (2022) no. 2, 561-587.

[ADO] S. Abja, S. Dinew and G. Olive, *Uniform estimates for concave homogeneous complex degenerate elliptic equations comparable to the Monge-Ampère equation*, Potential Anal. (2022).

<https://link.springer.com/article/10.1007/s11118-022-10009-w>

[Ga] L. Gårding, *Linear hyperbolic differential equations with constant coefficients*, Acta. Math. **85** (1951), 2-62.

[GP₁] B.Guo and D. H.Phong, *On L^∞ -estimates for fully nonlinear partial differential equations*, arXiv:2204.12549v1.

[GP₂] ———, *Uniform entropy and energy bounds for fully non-linear equations*, arXiv:2207.08983v1, July 2022.

[GP₃] ———, *Auxiliary Monge-Ampere equations in geometric analysis*, arXiv:2210.13308.

[GP₄] ———, *Uniform L^∞ estimates: subequations to fully nonlinear partial differential equations*, arXiv:2401.11572v1

- [GPT] B. Guo, D.H. Phong, and F. Tong, *On L^∞ -estimates for complex Monge-Ampère equations*, Ann. of Math. (2) **198** (2023), no.1, 393 - 418. arXiv:2106.02224.
- [Gu] L. Gurvits, *Van der Waerden/Schrijver-Valiant like conjectures and stable (aka hyperbolic) homogeneous polynomials : one theorem for all*, Electron. J. Combin. **15** (2008), no. 1, Research Paper 66, 26 pp.
- [HL₁] F. Reese Harvey and H. Blaine Lawson, Jr. *Dirichlet duality and the non-linear Dirichlet problem on Riemannian manifolds*, J. Diff. Geom. **88** No. 3 (2011), 395-482. arXiv:0907.1981.
- [HL₂] ———, *Gårding's theory of hyperbolic polynomials*, *Communications in Pure and Applied Mathematics* **66** no. 7 (2013), 1102-1128.
- [HL₃] ———, *Lagrangian potential theory and a Lagrangian equation of Monge-Ampère type*, pp. 217-258 in *Surveys in Differential Geometry*, edited by H.-D. Cao, J. Li, R. Schoen and S.-T. Yau, vol. 22, International Press, Somerville, MA, 2018. arXiv:1712.03525
- [HL₄] ———, *Determinant majorization and the work of Guo-Phong-Tong and Abja-Olive*, Calc. of Var. and PDE's **62** article number: 153 (2023). arXiv:2207.01729
- [HL₅] ———, *Duality for Gårding operators and a new inequality of GM/AM type*, (to appear).
- [KLW] N. Klemyatin, S. Liang and C. Wang, *On uniform estimates for $n-1$ -form fully nonlinear partial differential equations on compact Hermitian manifolds*, arXiv:2211.13798v2.
- [Ren] J. Renegar, *Hyperbolic programs, and their derivative relaxations*, Found. Comput. Math. **6** (2006) no. 1, 59-79.