Sharp Sobolev regularity for widely degenerate parabolic equations

Pasquale Ambrosio *

July 7, 2024

Abstract

We consider local weak solutions to the widely degenerate parabolic PDE

$$\partial_t u - \operatorname{div}\left((|Du| - \lambda)^{p-1}_+ \frac{Du}{|Du|}\right) = f \quad \text{in } \Omega_T = \Omega \times (0, T),$$

where $p \geq 2$, Ω is a bounded domain in \mathbb{R}^n for $n \geq 2$, λ is a non-negative constant and $(\cdot)_+$ stands for the positive part. Assuming that the datum f belongs to a suitable Lebesgue-Besov parabolic space when p > 2 and that $f \in L^2_{loc}(\Omega_T)$ if p = 2, we prove the Sobolev spatial regularity of a *novel* nonlinear function of the spatial gradient of the weak solutions. This result, in turn, implies the existence of the weak time derivative for the solutions of the evolutionary p-Poisson equation. The main novelty here is that f only has a Besov or Lebesgue spatial regularity, unlike the previous work [6], where f was assumed to possess a Sobolev spatial regularity of integer order. We emphasize that the results obtained here can be considered, on the one hand, as the parabolic analog of some elliptic results established in [5], and on the other hand as the extension to a strongly degenerate setting of some known results for less degenerate parabolic equations.

Mathematics Subject Classification: 35B45, 35B65, 35D30, 35K10, 35K65.

Keywords: Degenerate parabolic equations; higher differentiability; Sobolev regularity; Besov spaces.

1 Introduction

In this paper, we are interested in local weak solutions $u : \Omega_T \to \mathbb{R}$ to the evolutionary equation

$$\partial_t u - \operatorname{div}\left((|Du| - \lambda)^{p-1}_+ \frac{Du}{|Du|}\right) = f \quad \text{in } \Omega_T = \Omega \times (0, T), \tag{1.1}$$

where Ω is a bounded domain in \mathbb{R}^n $(n \ge 2)$, T > 0, $\lambda \ge 0$ and $(\cdot)_+$ stands for the positive part. Our analysis is confined to the super-quadratic case, i.e. in this work we consider $p \ge 2$.

A motivation for studying PDEs of the type (1.1) can be found in gas filtration problems taking into account the initial pressure gradient (see [4], [6], [7] and the references therein).

^{*}Dipartimento di Matematica e Applicazioni "R. Caccioppoli", Università degli Studi di Napoli "Federico II", Via Cintia, 80126 Napoli, Italy. *E-mail address*: pasquale.ambrosio2@unina.it

P. AMBROSIO

The main feature of the above equation is that it exhibits a strong degeneracy, coming from the fact that its modulus of ellipticity vanishes in the region $\{|Du| \leq \lambda\}$, and hence its principal part behaves like a *p*-Laplace operator only for large values of |Du|.

As our main result, here we establish the Sobolev spatial regularity of a nonlinear function of the spatial gradient Du of the weak solutions to equation (1.1), by assuming that the datum fbelongs to a suitable Lebesgue-Besov parabolic space when p > 2 (see Theorem 1.1 below) and that $f \in L^2_{loc}(\Omega_T)$ if p = 2 (see Theorem 1.3). These results, in turn, imply the Sobolev time regularity of the solutions to the evolutionary p-Poisson equation, under the same hypotheses on the function f (cf. Theorem 1.5, where we only address the case p > 2, since the Sobolev time regularity is well known for the heat equation with source term in $L^2_{loc}(\Omega_T)$). These issues have been widely investigated, as one can see, for example, in [6, 13, 15, 16, 26]. In particular, establishing the Sobolev regularity of the solutions with respect to time, once the higher differentiability in space has been obtained, is a quite standard fact: see, for instance, [18]-[20].

Before specifying in detail the assumption on f in the case p > 2, we wish to discuss some results already available in the literature. A common aspect of nonlinear parabolic problems with growth rate $p \ge 2$ is that the higher differentiability is proven for a nonlinear function of the gradient that takes into account the growth of the principal part of the equation. Indeed, already for the parabolic p-Laplace equation (which is obtained from (1.1) by setting $\lambda = f = 0$), the higher differentiability is established for the function $V_{\mu}(Du) := (\mu^2 + |Du|^2)^{\frac{p-2}{4}} Du$, as one can see, for example, in [13]. In case of widely degenerate problems, this phenomenon persists and, both for elliptic and parabolic equations, higher differentiability results hold true for the function $H_{p/2}(Du) := (|Du| - \lambda)^{p/2}_{+} \frac{Du}{|Du|}$ (see [1, 2, 6, 10]). However, this function does not provide any information about the second-order regularity of the solutions in the set where the equation becomes degenerate. Actually, since every λ -Lipschitz function is a solution of the homogeneous elliptic equation

$$\operatorname{div}\left((|Du| - \lambda)_{+}^{p-1}\frac{Du}{|Du|}\right) = 0,$$

no more than Lipschitz regularity can be expected for the solutions: in this regard, see [9] for the elliptic counterpart of equation (1.1) and [3] for a very recent generalization to strongly degenerate parabolic systems.

In addition, it is well known that, for both the stationary and the evolutionary p-Poisson equation, a Sobolev (spatial) regularity is required for the datum f in order to get the Sobolev (spatial) regularity of the function

$$V_0(Du) \equiv V(Du) := |Du|^{\frac{p-2}{2}} Du$$

which coincides with $H_{p/2}(Du)$ for $\lambda = 0$ (see, for example, [11] for the stationary *p*-Poisson equation and [6] for the evolutionary one). In particular, in the elliptic setting, the optimal assumption on the datum *f* to obtain the local $W^{1,2}$ -regularity of V(Du) has been determined in [11, Theorem 1.1] as a fractional Sobolev regularity suitably related to the growth exponent p > 2. Recently, this result has somewhat been improved and extended to the elliptic version of (1.1) in [5]: there, for p > 2 the authors consider the function

$$\mathcal{G}_{\alpha,\lambda}(t) := \int_0^t \frac{\omega^{\frac{p-1+2\alpha}{2}}}{(\omega+\lambda)^{\frac{1+2\alpha}{2}}} d\omega$$
(1.2)

for $t \ge 0$ and $\alpha \ge \frac{p+1}{2(p-1)}$, and establish that $\mathcal{G}_{\alpha,\lambda}((|Du| - \lambda)_+) \in W^{1,2}_{loc}(\Omega)$ by assuming

$$f \in B_{p',1,loc}^{\frac{p-2}{p}}(\Omega) \quad \text{with } p' := \frac{p}{p-1},$$

which is weaker than the assumption adopted in [11], due to Lemma 3.8 below.

Here, our main results tail the ones mentioned above, with the aim of extending the elliptic results contained in [5] and [11] to the parabolic setting. In fact, the first theorem we establish in this paper can be considered as the parabolic counterpart of [5, Theorem 1.1]. In order to state it, for $\xi \in \mathbb{R}^n$ we define the following function:

$$\mathcal{V}_{\alpha,\lambda}(\xi) \equiv \mathcal{V}_{\lambda}(\xi) := \begin{cases} \mathcal{G}_{\alpha,\lambda}((|\xi| - \lambda)_{+}) \frac{\xi}{|\xi|} & \text{if } |\xi| > \lambda, \\ 0 & \text{if } |\xi| \le \lambda, \end{cases}$$
(1.3)

where $\alpha \geq 0$ and $\mathcal{G}_{\alpha,\lambda}$ is the function defined by (1.2). At this point, the first theorem of this work reads as follows. We refer to Sections 2 and 3 for further notations and definitions.

Theorem 1.1. Let
$$n \ge 2$$
, $p > 2$, $\lambda \ge 0$ and $f \in L^{p'}_{loc}\left(0, T; B^{\frac{p-2}{p}}_{p',1,loc}(\Omega)\right)$. Moreover, let

$$\alpha = \begin{cases} 0 & \text{if } \lambda = 0, \\ any \text{ value in } \left[\frac{p+1}{2(p-1)}, \infty\right) & \text{if } \lambda > 0, \end{cases}$$
(1.4)

and assume that

 $u \in C^0\left(0, T; L^2(\Omega)\right) \cap L^p_{loc}\left(0, T; W^{1, p}_{loc}(\Omega)\right)$

is a local weak solution of equation (1.1). Then

$$\mathcal{V}_{\alpha,\lambda}(Du) \in L^2_{loc}\left(0,T; W^{1,2}_{loc}(\Omega,\mathbb{R}^n)\right)$$

Furthermore, for any parabolic cylinder $Q_r(z_0) \subset Q_\rho(z_0) \subset Q_R(z_0) \subseteq \Omega_T$ we have

$$\int_{Q_{r/2}(z_0)} |D_x \mathcal{V}_{\alpha,\lambda}(Du)|^2 dz \leq \frac{C}{\rho^2} \left(\|Du\|_{L^p(Q_R)}^p + \|Du\|_{L^p(Q_R)}^2 + \lambda^p + \lambda^2 + 1 \right) \\
+ C \|f\|_{L^{p'}\left(t_0 - R^2, t_0; B_{p',1}^{\frac{p-2}{p}}(B_R(x_0))\right)}$$
(1.5)

for a positive constant C depending only on n, p and R in the case $\lambda = 0$, and additionally on α if $\lambda > 0$. Besides, if $\lambda = 0$ we get

$$Du \in L^p_{loc}(0,T; W^{\sigma,p}_{loc}(\Omega,\mathbb{R}^n)) \quad for \ all \ \sigma \in \left(0,\frac{2}{p}\right)$$

Remark 1.2. Notice that, for every $\alpha \geq 0$, we have

$$\mathcal{V}_{\alpha,0}(\xi) = \frac{2}{p} V(\xi) := \frac{2}{p} |\xi|^{\frac{p-2}{2}} \xi, \qquad (1.6)$$

i.e. \mathcal{V}_0 is actually independent of the parameter α , which explains the choice $(1.4)_1$ in the statement of the above theorem. The condition $(1.4)_2$ will instead be decisive to carry out the proof of Proposition 4.6 below. Moreover, looking at (1.6), one can easily understand that, on the one hand, Theorem 1.1 extends the results proved in [11] and [13] to a widely degenerate parabolic setting. On the other hand, it extends the aforementioned results to the case of data in a suitable Lebesgue-Besov parabolic space, which turns out to be optimal, as can be seen by appropriately adapting the example in [11, Section 5] to the parabolic setting (in this regard, see also [5, page 2]).

P. AMBROSIO

As already mentioned, for $\lambda > 0$ the weak solutions of (1.1) are not twice differentiable. Therefore, in general we cannot differentiate the equation to estimate the second derivatives of the solutions. Here, we overcome this difficulty by introducing a suitable family of approximating problems, whose solutions are regular enough by standard parabolic regularity (see Section 4 below). The major effort in proving Theorem 1.1 is to establish suitable estimates for the solutions u_{ε} of the regularized problems that are uniform with respect to the approximation parameter ε . Next, we take advantage of these uniform estimates in the use of a comparison argument aimed to transfer the higher differentiability in space of $\mathcal{V}_{\alpha,\lambda}(Du_{\varepsilon})$ to the function $\mathcal{V}_{\alpha,\lambda}(Du)$.

The main differences with respect to the arguments used in [6] lie in the use of the function $\mathcal{V}_{\alpha,\lambda}(Du)$ in place of $H_{p/2}(Du)$ and in the derivation of the *a priori* estimates for the terms involving the datum f, which now only has a Besov spatial regularity. Indeed, if f had a suitable Sobolev spatial regularity as in [6], the terms of the estimate coming from the inhomogeneity of the equation could be controlled using the information on the integrability of the spatial gradient of the solution. Here, instead, in order to control the terms coming from the inhomogeneity, we essentially use the duality of Besov spaces and Theorem 3.6 below: this novelty comes from an idea that we have already exploited in the recent paper [5], in the elliptic setting.

Now we turn our attention to the case p = 2. It is well known that for the non-homogeneous heat equation

$$\partial_t u - \operatorname{div} Du = f \quad \text{in } \Omega_T, \tag{1.7}$$

the spatial $W^{1,2}$ -regularity of the weak spatial gradient Du can be achieved by simply assuming that $f \in L^2_{loc}(\Omega_T)$. Here, we prove that a similar result holds even when dealing with much more degenerate equations. More precisely, we establish the following result.

Theorem 1.3. Let $n \ge 2$, $\lambda \ge 0$ and $f \in L^2_{loc}(\Omega_T)$. Moreover, let

$$\alpha = \begin{cases} 0 & \text{if } \lambda = 0, \\ \text{any value in } \left[\frac{3}{2}, \infty\right) & \text{if } \lambda > 0, \end{cases}$$

and assume that

$$u \in C^0\left(0, T; L^2(\Omega)\right) \cap L^2_{loc}\left(0, T; W^{1,2}_{loc}(\Omega)\right)$$

is a local weak solution of the equation

$$\partial_t u - \operatorname{div}\left((|Du| - \lambda)_+ \frac{Du}{|Du|}\right) = f \quad in \ \Omega_T.$$

Then

 $\mathcal{V}_{\alpha,\lambda}(Du) \in L^2_{loc}\left(0,T; W^{1,2}_{loc}(\Omega,\mathbb{R}^n)\right).$

Furthermore, for any parabolic cylinder $Q_r(z_0) \subset Q_\rho(z_0) \subset Q_R(z_0) \subseteq \Omega_T$ we have

$$\int_{Q_{r/2}(z_0)} |D_x \mathcal{V}_{\alpha,\lambda}(Du)|^2 dz \le \frac{C}{\rho^2} \left(\|Du\|_{L^2(Q_R)}^2 + \lambda^2 + 1 \right) + C \left(\|f\|_{L^2(Q_R)}^2 + 1 \right)$$

for a positive constant C depending only on n and R in the case $\lambda = 0$, and additionally on α if $\lambda > 0$.

Remark 1.4. In the case p = 2, we have $\mathcal{V}_0(Du) = Du$. Then, for $\lambda = 0$, it follows from Theorem 1.3 that

$$Du \in L^2_{loc}\left(0,T; W^{1,2}_{loc}(\Omega,\mathbb{R}^n)\right).$$

As already mentioned, this is a well-established result for equation (1.7) when $f \in L^2_{loc}(\Omega_T)$.

For p > 2, we now consider the evolutionary *p*-Poisson equation

$$\partial_t u - \operatorname{div}\left(|Du|^{p-2}Du\right) = f \quad \text{in } \Omega_T,$$
(1.8)

which is obtained from equation (1.1) by setting $\lambda = 0$. As we anticipated earlier, from Theorem 1.1 we can easily deduce that the weak solutions of (1.8) admit a weak time derivative which belongs to the local Lebesgue space $L_{loc}^{p'}(\Omega_T)$. The idea is roughly as follows. Since the abovementioned theorem tells us that in a certain pointwise sense the second spatial derivatives of u exist, we may develop the expression under the divergence symbol. This will give us an expression that equals $\partial_t u$, from which we get the desired integrability of the time derivative. Such an argument must be made more rigorous. Furthermore, we also need to make explicit *a priori* local estimates. These are provided in the following result.

Theorem 1.5. Let $n \ge 2$, p > 2 and $f \in L_{loc}^{p'}\left(0, T; B_{p',1,loc}^{\frac{p-2}{p}}(\Omega)\right)$. Moreover, assume that $u \in C^0\left(0, T; L^2(\Omega)\right) \cap L_{loc}^p\left(0, T; W_{loc}^{1,p}(\Omega)\right)$

is a local weak solution of equation (1.8). Then, the time derivative of u exists in the weak sense and satisfies

$$\partial_t u \in L^{p'}_{loc}(\Omega_T).$$

Furthermore, for any parabolic cylinder $Q_r(z_0) \subset Q_\rho(z_0) \subset Q_R(z_0) \subseteq \Omega_T$ we have

$$\left(\int_{Q_{r/2}(z_0)} \left|\partial_t u\right|^{p'} dz\right)^{\frac{1}{p'}} \leq \frac{C}{\rho} \left(\|Du\|_{L^p(Q_R)}^{p-1} + \|Du\|_{L^p(Q_R)}^{\frac{p}{2}} + \|Du\|_{L^p(Q_R)}^{\frac{p-2}{2}} \right) \\ + C \|Du\|_{L^p(Q_R)}^{\frac{p-2}{2}} \|f\|_{L^{p'}\left(t_0 - R^2, t_0; B_{p', 1}^{\frac{p-2}{p}}(B_R(x_0))\right)} + \|f\|_{L^{p'}(Q_R)} \quad (1.9)$$

for a positive constant C depending only on n, p and R.

Before describing the structure of this paper, we wish to point out that, starting from the weaker assumption

$$f \in L^{p'}_{loc}\left(0,T; B^{\frac{p-2}{p}}_{p',1,loc}(\Omega)\right) \quad \text{with } p > 2,$$

Sobolev regularity results such as those of Theorems 1.1 and 1.5 seem not to have been established yet for weak solutions to parabolic PDEs that are far less degenerate than equation (1.1) with $\lambda > 0$. In particular, the results contained in this paper permit to improve the existing literature, already for the evolutionary *p*-Poisson equation (1.8), which exhibits a milder degeneracy. Moreover, as far as we know, for $\lambda > 0$ the result of Theorem 1.3 is completely new.

1.1 Plan of the paper

The paper is organized as follows. Section 2 is devoted to the preliminaries: after a list of some classic notations and some essentials estimates, we recall the fundamental properties of the difference quotients of Sobolev functions. In Section 3, we recall the basic facts on the functional spaces involved in this paper. Here, the most important points are Theorems 3.5 and 3.6, whose role is crucial in the proof of Theorem 1.1. In Section 4, we consider a regularization of equation (1.1) and establish some *a priori* estimates for its weak solution. In particular, we prove Sobolev estimates which are independent of the regularization parameter (Propositions 4.6 and 4.7). These estimates will be needed to demonstrate Theorems 1.1 and 1.3, whose proofs are achieved in Section 5. Finally, in Section 6 we prove Theorem 1.5.

2 Preliminaries

2.1 Notation and essential definitions

In this paper we shall denote by C or c a general positive constant that may vary on different occasions. Relevant dependencies on parameters and special constants will be suitably emphasized using parentheses or subscripts. The norm we use on \mathbb{R}^n will be the standard Euclidean one and it will be denoted by $|\cdot|$. In particular, for the vectors $\xi, \eta \in \mathbb{R}^n$, we write $\langle \xi, \eta \rangle$ for the usual inner product and $|\xi| := \langle \xi, \xi \rangle^{\frac{1}{2}}$ for the corresponding Euclidean norm.

For points in space-time, we will frequently use abbreviations like z = (x, t) or $z_0 = (x_0, t_0)$, for spatial variables $x, x_0 \in \mathbb{R}^n$ and times $t, t_0 \in \mathbb{R}$. We also denote by $B_{\rho}(x_0) = \{x \in \mathbb{R}^n : |x - x_0| < \rho\}$ the *n*-dimensional open ball with radius $\rho > 0$ and center $x_0 \in \mathbb{R}^n$; when not important, or clear from the context, we shall omit to denote the center as follows: $B_{\rho} \equiv B_{\rho}(x_0)$. Unless otherwise stated, different balls in the same context will have the same center. Moreover, we use the notation

$$Q_{\rho}(z_0) := B_{\rho}(x_0) \times (t_0 - \rho^2, t_0), \quad z_0 = (x_0, t_0) \in \mathbb{R}^n \times \mathbb{R}, \ \rho > 0,$$

for the backward parabolic cylinder with vertex (x_0, t_0) and width ρ . We shall sometimes omit the dependence on the vertex when all cylinders occurring in a proof share the same vertex. For a general cylinder $Q = B \times (t_0, t_1)$, where $B \subset \mathbb{R}^n$ and $t_0 < t_1$, we denote by

$$\partial_{\operatorname{par}}Q := (\overline{B} \times \{t_0\}) \cup (\partial B \times (t_0, t_1))$$

the usual *parabolic boundary* of Q, which is nothing but its standard topological boundary without the upper cap $\overline{B} \times \{t_1\}$.

If $E \subseteq \mathbb{R}^k$ is a Lebesgue-measurable set, then we will denote by |E| its k-dimensional Lebesgue measure.

For every function $v \in L^1_{loc}(Q, \mathbb{R}^k)$, where $Q \subset \mathbb{R}^{n+1}$ and $k \in \mathbb{N}$, we define the mollified function v_{ρ} as follows:

$$v_{\varrho}(z) := \int_{\mathbb{R}^{n+1}} v(\tilde{z}) \,\phi_{\varrho}(z - \tilde{z}) \,d\tilde{z},\tag{2.1}$$

where

$$\phi_{\varrho}(z) := \frac{1}{\varrho^{n+1}} \phi_1\left(\frac{z}{\varrho}\right), \qquad \varrho > 0,$$

with $\phi_1 \in C_0^{\infty}(\mathcal{B}_1(0))^1$ denoting the standard, non-negative, radially symmetric mollifier in \mathbb{R}^{n+1} . Obviously, here the function v is meant to be extended by the k-dimensional null vector outside Q.

For further needs, we now define the auxiliary function $H_{\gamma}: \mathbb{R}^n \to \mathbb{R}^n$ by

$$H_{\gamma}(\xi) := \begin{cases} (|\xi| - \lambda)^{\gamma}_{+ \frac{\xi}{|\xi|}} & \text{if } \xi \neq 0, \\ 0 & \text{if } \xi = 0, \end{cases}$$
(2.2)

where $\gamma > 0$ is a parameter. We conclude this first part of the preliminaries by recalling the following definition.

¹Here $\mathcal{B}_1(0)$ denotes the (n+1)-dimensional open unit ball centered at the origin.

Definition 2.1. Let $\lambda \geq 0$. A function

$$u \in C^0\left(0, T; L^2(\Omega)\right) \cap L^p_{loc}\left(0, T; W^{1, p}_{loc}(\Omega)\right)$$

is a *local weak solution* of equation (1.1) if and only if, for any test function $\varphi \in C_0^{\infty}(\Omega_T)$, the following integral identity holds:

$$\int_{\Omega_T} \left(u \cdot \partial_t \varphi - \langle H_{p-1}(Du), D\varphi \rangle \right) dz = - \int_{\Omega_T} f\varphi \, dz.$$
(2.3)

2.2 Algebraic inequalities

In this section, we gather some relevant algebraic inequalities that will be needed later on. The first result follows from an elementary computation.

Lemma 2.2. For $\xi, \eta \in \mathbb{R}^n \setminus \{0\}$, we have

$$\left|\frac{\xi}{|\xi|} - \frac{\eta}{|\eta|}\right| \le \frac{2}{|\eta|} |\xi - \eta|$$

We now recall the following estimate, whose proof can be found in [21, Chapter 12].

Lemma 2.3. Let $p \in (2, \infty)$ and $k \in \mathbb{N}$. Then, for every $\xi, \eta \in \mathbb{R}^k$ we get

$$|\xi - \eta|^p \le C \left| |\xi|^{\frac{p-2}{2}} \xi - |\eta|^{\frac{p-2}{2}} \eta \right|^2$$

for a constant $C \equiv C(p) > 0$.

For the function H_{γ} defined by (2.2), we record the following estimates, which can be obtained by suitably modifying the proofs of [10, Lemma 4.1] and [8, Lemma 2.8], respectively.

Lemma 2.4. Let $2 \le p < \infty$. Then, for every $\xi, \eta \in \mathbb{R}^n$ we get

$$\langle H_{p-1}(\xi) - H_{p-1}(\eta), \xi - \eta \rangle \geq \frac{4}{p^2} \left| H_{\frac{p}{2}}(\xi) - H_{\frac{p}{2}}(\eta) \right|^2.$$

Moreover, if $|\eta| > \lambda$ we have

$$\langle H_{p-1}(\xi) - H_{p-1}(\eta), \xi - \eta \rangle \ge \frac{1}{2^{p+1}} \frac{(|\eta| - \lambda)^p}{|\eta| (|\xi| + |\eta|)} |\xi - \eta|^2.$$

The next result concerns the function $\mathcal{G}_{\alpha,\lambda}$ defined by (1.2).

Lemma 2.5. Let $p \in [2, \infty)$, $\lambda \ge 0$ and $\alpha \ge 0$. Then

$$\mathcal{G}_{\alpha,\lambda}(t) \leq \frac{2}{p} t^{\frac{p}{2}} \left(\frac{t}{t+\lambda}\right)^{\frac{1+2\alpha}{2}}$$

for every t > 0.

Proof. Since the function

$$K(\omega) := \left(\frac{\omega}{\omega + \lambda}\right)^{\frac{1+2\alpha}{2}}, \qquad \omega > 0,$$

is non-decreasing, for every t > 0 we have

$$\mathcal{G}_{\alpha,\lambda}(t) = \int_0^t K(\omega) \,\omega^{\frac{p}{2}-1} \,d\omega \,\leq \left(\frac{t}{t+\lambda}\right)^{\frac{1+2\alpha}{2}} \int_0^t \omega^{\frac{p}{2}-1} \,d\omega \,=\, \frac{2}{p} t^{\frac{p}{2}} \left(\frac{t}{t+\lambda}\right)^{\frac{1+2\alpha}{2}}.$$

Finally, the next lemma relates the function $\mathcal{V}_{\alpha,\lambda}(\xi)$ with $H_{p-1}(\xi)$.

Lemma 2.6. Let $p \in [2, \infty)$, $\lambda \ge 0$ and $\alpha \ge 0$. Then, there exists a constant $C \equiv C(p) > 0$ such that

$$|\mathcal{V}_{\alpha,\lambda}(\xi) - \mathcal{V}_{\alpha,\lambda}(\eta)|^2 \le C \langle H_{p-1}(\xi) - H_{p-1}(\eta), \xi - \eta \rangle$$
(2.4)

for every $\xi, \eta \in \mathbb{R}^n$.

Proof. For $\lambda = 0$ this is a well-known result (see, for example, [6, Lemma 2.2]). Therefore, from now on we shall assume that $\lambda > 0$. We first note that inequality (2.4) is trivially satisfied when $|\xi|, |\eta| \leq \lambda$. If $|\eta| \leq \lambda < |\xi|$, using the definitions (1.3), (1.2), (2.2) and Lemma 2.4, we obtain

$$\begin{aligned} |\mathcal{V}_{\alpha,\lambda}(\xi) - \mathcal{V}_{\alpha,\lambda}(\eta)|^2 &= [\mathcal{G}_{\alpha,\lambda}(|\xi| - \lambda)]^2 \le \left(\int_0^{|\xi| - \lambda} \omega^{\frac{p}{2} - 1} \, d\omega\right)^2 = \frac{4}{p^2} \, (|\xi| - \lambda)^p \\ &= \frac{4}{p^2} \left| H_{\frac{p}{2}}(\xi) - H_{\frac{p}{2}}(\eta) \right|^2 \le \, \langle H_{p-1}(\xi) - H_{p-1}(\eta), \xi - \eta \rangle. \end{aligned}$$

Now let $|\xi|, |\eta| > \lambda$. Without loss of generality, we may assume that $|\eta| \ge |\xi| > \lambda$. This implies

$$|\eta|^{2} = \frac{|\eta| (|\eta| + |\eta|)}{2} \ge \frac{|\eta| (|\xi| + |\eta|)}{2}.$$
(2.5)

Moreover, we have

$$\begin{split} |\mathcal{V}_{\alpha,\lambda}(\xi) - \mathcal{V}_{\alpha,\lambda}(\eta)| &= \left| \mathcal{V}_{\alpha,\lambda}(\xi) - \mathcal{G}_{\alpha,\lambda}(|\eta| - \lambda) \frac{\xi}{|\xi|} + \mathcal{G}_{\alpha,\lambda}(|\eta| - \lambda) \frac{\xi}{|\xi|} - \mathcal{V}_{\alpha,\lambda}(\eta) \right| \\ &\leq \left| \mathcal{G}_{\alpha,\lambda}(|\xi| - \lambda) - \mathcal{G}_{\alpha,\lambda}(|\eta| - \lambda) \right| + \mathcal{G}_{\alpha,\lambda}(|\eta| - \lambda) \left| \frac{\xi}{|\xi|} - \frac{\eta}{|\eta|} \right| \\ &\leq \int_{|\xi| - \lambda}^{|\eta| - \lambda} \frac{\omega^{\frac{p-1+2\alpha}{2}}}{(\omega + \lambda)^{\frac{1+2\alpha}{2}}} d\omega + \frac{2}{|\eta|} |\xi - \eta| \int_{0}^{|\eta| - \lambda} \frac{\omega^{\frac{p-1+2\alpha}{2}}}{(\omega + \lambda)^{\frac{1+2\alpha}{2}}} d\omega \\ &\leq \int_{|\xi| - \lambda}^{|\eta| - \lambda} \omega^{\frac{p}{2} - 1} d\omega + \frac{2}{|\eta|} |\xi - \eta| \int_{0}^{|\eta| - \lambda} \omega^{\frac{p}{2} - 1} d\omega \\ &= \frac{2}{p} \left| \left| H_{\frac{p}{2}}(\eta) \right| - \left| H_{\frac{p}{2}}(\xi) \right| \right| + 4 \frac{(|\eta| - \lambda)^{\frac{p}{2}}}{p |\eta|} |\xi - \eta| \\ &\leq \frac{2}{p} \left| H_{\frac{p}{2}}(\xi) - H_{\frac{p}{2}}(\eta) \right| + 4 \frac{(|\eta| - \lambda)^{\frac{p}{2}}}{p |\eta|} |\xi - \eta|, \end{split}$$

where, in the third line, we have used Lemma 2.2 and the fact that $\mathcal{G}_{\alpha,\lambda}$ is an increasing function. Now, applying Young's inequality, estimate (2.5) and Lemma 2.4, we obtain

$$\begin{aligned} |\mathcal{V}_{\alpha,\lambda}(\xi) - \mathcal{V}_{\alpha,\lambda}(\eta)|^2 &\leq \frac{8}{p^2} \left| H_{\frac{p}{2}}(\xi) - H_{\frac{p}{2}}(\eta) \right|^2 + 32 \frac{(|\eta| - \lambda)^p}{p^2 |\eta|^2} |\xi - \eta|^2 \\ &\leq \frac{8}{p^2} \left| H_{\frac{p}{2}}(\xi) - H_{\frac{p}{2}}(\eta) \right|^2 + \frac{64}{p^2} \frac{(|\eta| - \lambda)^p}{|\eta| (|\xi| + |\eta|)} |\xi - \eta|^2 \\ &\leq C_p \left\langle H_{p-1}(\xi) - H_{p-1}(\eta), \xi - \eta \right\rangle. \end{aligned}$$

This completes the proof.

2.3 Difference quotients

We recall here the definition and some elementary properties of the difference quotients that will be useful in the following (see, for example, [17]).

Definition 2.7. For every vector-valued function $F : \mathbb{R}^n \to \mathbb{R}^k$ the *finite difference operator* in the direction x_j is defined by

$$\tau_{j,h}F(x) = F(x + he_j) - F(x),$$

where $h \in \mathbb{R}$, e_j is the unit vector in the direction x_j and $j \in \{1, \ldots, n\}$.

The difference quotient of F with respect to x_j is defined for $h \in \mathbb{R} \setminus \{0\}$ by

$$\Delta_{j,h}F(x) = \frac{\tau_{j,h}F(x)}{h}$$

When no confusion can arise, we shall omit the index j and simply write τ_h or Δ_h instead of $\tau_{j,h}$ or $\Delta_{j,h}$, respectively.

Proposition 2.8. Let F be a function such that $F \in W^{1,q}(\Omega)$, with $q \ge 1$, and let us consider the set

$$\Omega_{|h|} := \left\{ x \in \Omega : \operatorname{dist} \left(x, \partial \Omega \right) > |h| \right\}.$$

Then:

(i)
$$\Delta_h F \in W^{1,q}(\Omega_{|h|})$$
 and $D_{x_i}(\Delta_h F) = \Delta_h(D_{x_i}F)$ for every $i \in \{1, \ldots, n\}$.

(ii) If at least one of the functions F or G has support contained in $\Omega_{|h|}$, then

$$\int_{\Omega} F \,\Delta_h G \,dx \,=\, -\int_{\Omega} G \,\Delta_{-h} F \,dx.$$

(iii) We have

$$\Delta_h(FG)(x) = F(x + he_j)\Delta_h G(x) + G(x)\Delta_h F(x).$$

The next result about the finite difference operator is a kind of integral version of the Lagrange Theorem and its proof can be found in [17, Lemma 8.1].

Lemma 2.9. If $0 < \rho < R$, $|h| < \frac{R-\rho}{2}$, $1 < q < +\infty$ and $F \in L^q(B_R, \mathbb{R}^k)$ is such that $DF \in L^q(B_R, \mathbb{R}^{k \times n})$, then

$$\int_{B_{\rho}} |\tau_h F(x)|^q \, dx \, \le \, c^q(n) \, |h|^q \int_{B_R} |DF(x)|^q \, dx$$

Moreover

$$\int_{B_{\rho}} |F(x+he_j)|^q \, dx \leq \int_{B_R} |F(x)|^q \, dx.$$

Finally, we recall the following fundamental result, whose proof can be found in [17, Lemma 8.2]:

Lemma 2.10. Let $F : \mathbb{R}^n \to \mathbb{R}^k$, $F \in L^q(B_R, \mathbb{R}^k)$ with $1 < q < +\infty$. Suppose that there exist $\rho \in (0, R)$ and a constant M > 0 such that

$$\sum_{j=1}^{n} \int_{B_{\rho}} |\tau_{j,h} F(x)|^{q} dx \leq M^{q} |h|^{q}$$

for every h with $|h| < \frac{R-\rho}{2}$. Then $F \in W^{1,q}(B_{\rho}, \mathbb{R}^k)$. Moreover

 $\|DF\|_{L^q(B_\rho)} \le M$

and

$$\Delta_{j,h}F \to D_{x_j}F$$
 in $L^q_{loc}(B_R)$ as $h \to 0$,

for each $j \in \{1, ..., n\}$.

3 Functional spaces

Here we recall some essential facts about the functional spaces involved in this paper, starting with the definition and some properties of the Besov spaces that will be useful to prove our results.

We denote by $\mathcal{S}(\mathbb{R}^n)$ and $\mathcal{S}'(\mathbb{R}^n)$ the Schwartz space and the space of tempered distributions on \mathbb{R}^n , respectively. If $v \in \mathcal{S}(\mathbb{R}^n)$, then

$$\hat{v}(\xi) = (\mathcal{F}v)(\xi) = (2\pi)^{-n/2} \int_{\mathbb{R}^n} e^{-i\langle x,\xi\rangle} v(x) \, dx, \quad \xi \in \mathbb{R}^n,$$
(3.1)

denotes the Fourier transform of v. As usual, $\mathcal{F}^{-1}v$ and v^{\vee} stand for the inverse Fourier transform, given by the right-hand side of (3.1) with i in place of -i. Both \mathcal{F} and \mathcal{F}^{-1} are extended to $\mathcal{S}'(\mathbb{R}^n)$ in the standard way.

Now, let $\Gamma(\mathbb{R}^n)$ be the collection of all sequences $\varphi = \{\varphi_j\}_{j=0}^\infty \subset \mathcal{S}(\mathbb{R}^n)$ such that

$$\begin{cases} \operatorname{supp} \varphi_0 \subset \{x \in \mathbb{R}^n : |x| \le 2\} \\ \operatorname{supp} \varphi_j \subset \{x \in \mathbb{R}^n : 2^{j-1} \le |x| \le 2^{j+1}\} & \text{if } j \in \mathbb{N}, \end{cases}$$

for every multi-index β there exists a positive number c_{β} such that

$$2^{j|\beta|} |D^{\beta}\varphi_j(x)| \le c_{\beta}, \quad \forall j \in \mathbb{N}_0, \ \forall x \in \mathbb{R}^n$$

and

$$\sum_{j=0}^{\infty} \varphi_j(x) = 1, \quad \forall x \in \mathbb{R}^n.$$

Then, it is well known that $\Gamma(\mathbb{R}^n)$ is not empty (see [27, Section 2.3.1, Remark 1]). Moreover, if $\{\varphi_j\}_{j=0}^{\infty} \in \Gamma(\mathbb{R}^n)$, the entire analytic functions $(\varphi_j \hat{v})^{\vee}(x)$ make sense pointwise in \mathbb{R}^n for any $v \in S'(\mathbb{R}^n)$. Therefore, the following definition makes sense:

Definition 3.1. Let $s \in \mathbb{R}$, $0 < p, q \leq \infty$ and $\varphi = \{\varphi_j\}_{j=0}^{\infty} \in \Gamma(\mathbb{R}^n)$. We define the *inhomogeneous Besov space* $B_{p,q}^s(\mathbb{R}^n)$ as the set of all $v \in S'(\mathbb{R}^n)$ such that

$$\|v\|_{B^{s}_{p,q}(\mathbb{R}^{n})} := \left(\sum_{j=0}^{\infty} 2^{jsq} \|(\varphi_{j}\,\hat{v})^{\vee}\|_{L^{p}(\mathbb{R}^{n})}^{q}\right)^{\frac{1}{q}} < +\infty \quad \text{if } q < \infty, \tag{3.2}$$

and

$$\|v\|_{B^{s}_{p,q}(\mathbb{R}^{n})} := \sup_{j \in \mathbb{N}_{0}} 2^{js} \|(\varphi_{j} \hat{v})^{\vee}\|_{L^{p}(\mathbb{R}^{n})} < +\infty \quad \text{if } q = \infty.$$
(3.3)

Remark 3.2. Obviously, the quasi-norm $||v||_{B^s_{p,q}(\mathbb{R}^n)}$ depends on the chosen sequence $\varphi \in \Gamma(\mathbb{R}^n)$, but this is not the case for the spaces $B^s_{p,q}(\mathbb{R}^n)$ themselves, in the sense that two different choices for the sequence φ give rise to equivalent quasi-norms (see [27, Section 2.3.2, Proposition 1]). This justifies our omission of the dependence on φ in the left-hand side of (3.2)-(3.3) and in the sequel.

We also know that the norms of the classical Besov spaces $B_{p,q}^s(\mathbb{R}^n)$ with $s \in (0,1), 1 \leq p < \infty$ and $1 \leq q \leq \infty$ can be characterized via differences of the functions involved, cf. [27, Section 2.5.12, Theorem 1]. More precisely, for $h \in \mathbb{R}^n$ and a measurable function $v : \mathbb{R}^n \to \mathbb{R}^k$, let us define

$$\delta_h v(x) := v(x+h) - v(x).$$

Then we have the equivalence

$$||v||_{B^{s}_{p,q}(\mathbb{R}^{n})} \approx ||v||_{L^{p}(\mathbb{R}^{n})} + [v]_{B^{s}_{p,q}(\mathbb{R}^{n})},$$

where

$$[v]_{B^s_{p,q}(\mathbb{R}^n)} := \left(\int_{\mathbb{R}^n} \left(\int_{\mathbb{R}^n} \frac{|\delta_h v(x)|^p}{|h|^{sp}} \, dx \right)^{\frac{q}{p}} \frac{dh}{|h|^n} \right)^{\frac{1}{q}}, \quad \text{if } 1 \le q < \infty, \tag{3.4}$$

and

$$[v]_{B^s_{p,\infty}(\mathbb{R}^n)} := \sup_{h \in \mathbb{R}^n} \left(\int_{\mathbb{R}^n} \frac{|\delta_h v(x)|^p}{|h|^{sp}} \, dx \right)^{\frac{1}{p}}.$$
(3.5)

As usual, in (3.4) if one simply integrates for |h| < r for a fixed r > 0, then an equivalent norm is obtained, since

$$\left(\int_{\{|h|\geq r\}} \left(\int_{\mathbb{R}^n} \frac{|\delta_h v(x)|^p}{|h|^{sp}} \, dx\right)^{\frac{q}{p}} \frac{dh}{|h|^n}\right)^{\frac{1}{q}} \leq c(n, s, p, q, r) \, \|v\|_{L^p(\mathbb{R}^n)}.$$

Similarly, in (3.5) one can simply take the supremum over $|h| \leq r$ and obtain an equivalent norm. By construction, $B^s_{p,q}(\mathbb{R}^n) \subset L^p(\mathbb{R}^n)$.

In the case of an arbitrary domain $\Omega \subset \mathbb{R}^n$, the spaces $B^s_{p,q}(\Omega)$ are defined by restriction. More precisely, denoting by $\mathcal{D}'(\Omega)$ the dual space of all distributions in Ω , we set

$$B_{p,q}^{s}(\Omega) := \left\{ v \in \mathcal{D}'(\Omega) : v = g|_{\Omega} \text{ for some } g \in B_{p,q}^{s}(\mathbb{R}^{n}) \right\},\$$

with the quasi-norm $\|v\|_{B^s_{p,q}(\Omega)} := \inf \|g\|_{B^s_{p,q}(\mathbb{R}^n)}$, where the infimum is taken over all $g \in B^s_{p,q}(\mathbb{R}^n)$ such that $g|_{\Omega} = v$.

If Ω is a bounded C^{∞} -domain in \mathbb{R}^n , then the restriction operator

$$\operatorname{re}_{\Omega}: \mathcal{S}'(\mathbb{R}^n) \hookrightarrow \mathcal{D}'(\Omega), \quad \operatorname{re}_{\Omega}(v) = v|_{\Omega}$$

generates a linear and bounded map from $B^s_{p,q}(\mathbb{R}^n)$ onto $B^s_{p,q}(\Omega)$. Furthermore, the spaces $B^s_{p,q}(\Omega)$ satisfy the so-called *extension property*, as ensured by the next theorem.

Theorem 3.3. Let $s \in \mathbb{R}$, let $0 < p, q \leq \infty$ and let Ω be a bounded C^{∞} -domain in \mathbb{R}^n . Then, there exists a linear and bounded extension operator $\operatorname{ext}_{\Omega} : B^s_{p,q}(\Omega) \hookrightarrow B^s_{p,q}(\mathbb{R}^n)$ such that $\operatorname{re}_{\Omega} \circ \operatorname{ext}_{\Omega} = \operatorname{id}$, where id is the identity in $B^s_{p,q}(\Omega)$.

For the proof we refer to [28, Theorem 2.82]. For further needs, we now give the following

Definition 3.4. For a bounded domain $\Omega \subset \mathbb{R}^n$, $s \in (0, 1)$ and $1 \leq p, q \leq \infty$, the homogeneous Besov space $\mathring{B}^s_{p,q}(\Omega)$ is defined as the completion of $C_0^{\infty}(\Omega)$ in $B^s_{p,q}(\Omega)$ with respect to the seminorm

$$v \mapsto \|v\|_{\mathring{B}^{s}_{p,q}(\Omega)} := [v]_{B^{s}_{p,q}(\mathbb{R}^{n})}$$

For $s \in (0, 1)$ and $1 \leq p, q \leq \infty$, we shall denote by $(\mathring{B}^s_{p,q}(\Omega))'$ the topological dual of $\mathring{B}^s_{p,q}(\Omega)$, which is endowed with the natural dual norm

$$\|F\|_{(\mathring{B}^{s}_{p,q}(\Omega))'} = \sup\left\{|\langle F,\varphi\rangle| : \varphi \in C_{0}^{\infty}(\Omega) \text{ and } \|\varphi\|_{\mathring{B}^{s}_{p,q}(\Omega)} \leq 1\right\}, \quad F \in (\mathring{B}^{s}_{p,q}(\Omega))'.$$

Now we recall the following duality formula, which has to be meant as an isomorphism of quasi-normed spaces (see [27, Section 2.11.2, Remark 2]).

Theorem 3.5. Let $s \in \mathbb{R}$ and $1 \leq p, q \leq \infty$. Then

$$(\mathring{B}^{s}_{p,q}(\mathbb{R}^{n}))' = B^{-s}_{p',q'}(\mathbb{R}^{n}).$$

The next result is a key ingredient for the proof of Theorem 1.1 and its proof can be found in [27, Section 3.3.5].

Theorem 3.6. Let $s \in \mathbb{R}$ and $0 < p, q \leq \infty$. Moreover, assume that Ω is a bounded C^{∞} -domain in \mathbb{R}^n . Then, for every $v \in B^s_{p,q}(\Omega)$ and every $j \in \{1, \ldots, n\}$ we have

$$\|\partial_j v\|_{B^{s-1}_{p,q}(\Omega)} \le c \|v\|_{B^s_{p,q}(\Omega)}$$

for a positive constant c which is independent of v.

We can also define local Besov spaces as follows. Given a domain $\Omega \subset \mathbb{R}^n$, we say that a function v belongs to $B^s_{p,q,loc}(\Omega)$ if $\phi v \in B^s_{p,q}(\mathbb{R}^n)$ whenever $\phi \in C_0^{\infty}(\Omega)$.

Definition 3.7. Let $\Omega \subseteq \mathbb{R}^n$ be an open set. For any $s \in (0, 1)$ and for any $q \in [1, +\infty)$, we define the *fractional Sobolev space* $W^{s,q}(\Omega, \mathbb{R}^k)$ as follows:

$$W^{s,q}(\Omega,\mathbb{R}^k) := \left\{ v \in L^q(\Omega,\mathbb{R}^k) : \frac{|v(x) - v(y)|}{|x - y|^{\frac{n}{q} + s}} \in L^q(\Omega \times \Omega) \right\},\$$

i.e. an intermerdiate Banach space between $L^q(\Omega, \mathbb{R}^k)$ and $W^{1,q}(\Omega, \mathbb{R}^k)$, endowed with the norm

$$||v||_{W^{s,q}(\Omega)} := ||v||_{L^q(\Omega)} + [v]_{W^{s,q}(\Omega)},$$

where the term

$$[v]_{W^{s,q}(\Omega)} := \left(\int_{\Omega} \int_{\Omega} \frac{|v(x) - v(y)|^q}{|x - y|^{n + sq}} \, dx \, dy \right)^{\frac{1}{q}}$$

is the so-called *Gagliardo seminorm* of v.

The following embedding result can be obtained by combining [27, Section 2.2.2, Remark 3] with [27, Section 2.3.2, Proposition 2(ii)].

Lemma 3.8. Let $s \in (0,1)$ and $q \ge 1$. Then, for every $\sigma \in (0,1-s)$ we have the continuous embedding $W^{s+\sigma,q}_{loc}(\mathbb{R}^n) \hookrightarrow B^s_{q,1,loc}(\mathbb{R}^n)$.

For the treatment of parabolic equations, we now give the following definitions.

Definition 3.9. Let $1 \leq q < \infty$ and $0 < \beta < 1$. A map $g \in L^q(\Omega \times (t_0, t_1), \mathbb{R}^k)$ belongs to the space $L^q(t_0, t_1; W^{\beta,q}(\Omega, \mathbb{R}^k))$ if and only if

$$\int_{t_0}^{t_1} \int_{\Omega} \int_{\Omega} \frac{|g(x,t) - g(y,t)|^q}{|x-y|^{n+\beta q}} \, dx \, dy \, dt < \infty.$$

Definition 3.10. Let $1 \leq p, q < \infty$ and 0 < s < 1. A map $g \in L^p(\Omega \times (t_0, t_1), \mathbb{R}^k)$ belongs to the space $L^p(t_0, t_1; B^s_{p,q}(\Omega, \mathbb{R}^k))$ if and only if

$$\int_{t_0}^{t_1} \|g(\cdot, t)\|_{B^s_{p,q}(\Omega)}^p \, dt < \infty.$$

In this paper, we use the corresponding local versions of the above spaces, which are denoted by the subscript "loc". More precisely, we write $g \in L^q_{loc}(0,T; W^{\beta,q}_{loc}(\Omega,\mathbb{R}^k))$ if and only if $g \in L^q(t_0,t_1; W^{\beta,q}(\Omega',\mathbb{R}^k))$ for all domains $\Omega' \times (t_0,t_1) \in \Omega_T$. The local Lebesgue-Besov space $L^p_{loc}(0,T; B^s_{p,q,loc}(\Omega,\mathbb{R}^k))$ is defined in a similar way. Furthermore, we shall also use the following notation, which is typical of Bochner spaces:

$$\|g\|_{L^{p}(t_{0},t_{1};B^{s}_{p,q}(\Omega'))} := \left(\int_{t_{0}}^{t_{1}} \|g(\cdot,t)\|_{B^{s}_{p,q}(\Omega')}^{p} dt\right)^{\frac{1}{p}}$$

We conclude this section with the parabolic version of the well-known result on the relation between Nikolskii spaces and fractional Sobolev spaces. This result is contained in [26, Lemma 2.4] (see also [13, Proposition 2.19]), and its proof can be obtained by a simple adaptation of the standard elliptic results [12, 14, 23, 24, 25].

Proposition 3.11. Let $Q_{\sigma}(z_0) \subset \mathbb{R}^{n+1}$ be a parabolic cylinder. Moreover, assume that $G \in L^q(Q_{\sigma}(z_0), \mathbb{R}^k)$, where $1 \leq q < \infty$. Then, for any $\theta \in (0, 1)$, the estimate

$$|h|^{-q\theta} \int_{Q_{\sigma}(z_0)} |G(x+he_j,t) - G(x,t)|^q \, dx \, dt \leq M^q < \infty$$

for a fixed constant $M \ge 0$, every $0 \ne |h| \le h_0$ and every $j \in \{1, \ldots, n\}$ implies

$$G \in L^{q}_{loc}\left(t_{0} - \sigma^{2}, t_{0}; W^{\beta,q}_{loc}\left(B_{\sigma}(x_{0}), \mathbb{R}^{k}\right)\right)$$

for all $\beta \in (0, \theta)$.

4 The regularization

In this section, we shall assume that λ , α , f and u are as in the statement of Theorem 1.1 if p > 2, and as in the statement of Theorem 1.3 if p = 2.

For $\varepsilon \in [0,1]$ and a couple of standard, non-negative, radially symmetric mollifiers $\varphi_1 \in C_0^{\infty}(B_1(0))$ and $\varphi_2 \in C_0^{\infty}((-1,1))$, we define

$$f_{\varepsilon}(x,t) := \int_{-1}^{1} \int_{B_1(0)} f(x - \varepsilon y, t - \varepsilon s) \varphi_1(y) \varphi_2(s) \, dy \, ds, \tag{4.1}$$

where f is meant to be extended by zero outside Ω_T . Let us observe that $f_0 = f$ and $f_{\varepsilon} \in C^{\infty}(\Omega_T)$ for every $\varepsilon \in (0, 1]$. In addition, we define the function $A_{\varepsilon} : \mathbb{R}^n \to \mathbb{R}$ by

$$A_{\varepsilon}(\xi) := \frac{1}{p} (|\xi| - \lambda)_{+}^{p} + \frac{\varepsilon}{p} (1 + |\xi|^{2})^{\frac{p}{2}}.$$

Therefore, we obtain

$$D_{\xi}A_{\varepsilon}(\xi) = H_{p-1}(\xi) + \varepsilon \left(1 + |\xi|^2\right)^{\frac{p-2}{2}} \xi, \quad \xi \in \mathbb{R}^n.$$

The next lemma provides the *p*-ellipticity property of the operator $D_{\xi}A_{\varepsilon}(\xi)$ and its proof follows from [8, Lemma 2.7].

Lemma 4.1. Let $\varepsilon \in [0,1]$, $p \ge 2$ and $z \in \mathbb{R}^n \setminus \{0\}$. Then, for every $\zeta \in \mathbb{R}^n$ we have

$$\left[\varepsilon \left(1+|z|^{2}\right)^{\frac{p-2}{2}} + \frac{\left(|z|-\lambda\right)^{p-1}_{+}}{|z|}\right]|\zeta|^{2} \le \langle D^{2}A_{\varepsilon}(z)\zeta,\zeta\rangle \le (p-1)\left[\varepsilon \left(1+|z|^{2}\right)^{\frac{p-2}{2}} + \left(|z|-\lambda\right)^{p-2}_{+}\right]|\zeta|^{2}\right]|\zeta|^{2}$$

Now we consider a parabolic cylinder $Q_R(z_0) := B_R(x_0) \times (t_0 - R^2, t_0) \Subset \Omega_T$. For our purposes, in the following we will need the definition below.

Definition 4.2. Let $\varepsilon \in (0, 1]$. In this framework, we identify a function

$$u_{\varepsilon} \in C^0\left([t_0 - R^2, t_0]; L^2(B_R(x_0))\right) \cap L^p\left(t_0 - R^2, t_0; W^{1,p}(B_R(x_0))\right)$$

as a weak solution of the Cauchy-Dirichlet problem

$$\begin{cases} \partial_t u_{\varepsilon} - \operatorname{div} \left[DA_{\varepsilon}(Du_{\varepsilon}) \right] = f_{\varepsilon} & \text{in } Q_R(z_0), \\ u_{\varepsilon} = u & \text{on } \partial_{\operatorname{par}} Q_R(z_0), \end{cases}$$
(4.2)

if and only if, for any $\varphi \in C_0^{\infty}(Q_R(z_0))$, the following integral identity holds

$$\int_{Q_R(z_0)} \left(u_{\varepsilon} \cdot \partial_t \varphi - \langle DA_{\varepsilon}(Du_{\varepsilon}), D\varphi \rangle \right) \, dz \, = \, - \int_{Q_R(z_0)} f_{\varepsilon} \, \varphi \, dz \tag{4.3}$$

and, moreover,

 $u_{\varepsilon} \in u + L^{p}\left(t_{0} - R^{2}, t_{0}; W_{0}^{1, p}(B_{R}(x_{0}))\right)$

and $u_{\varepsilon}(\cdot, t_0 - R^2) = u(\cdot, t_0 - R^2)$ in the L^2 -sense, that is,

$$\lim_{t \downarrow t_0 - R^2} \left\| u_{\varepsilon}(\cdot, t) - u(\cdot, t_0 - R^2) \right\|_{L^2(B_R(x_0))} = 0.$$
(4.4)

Therefore, the initial condition $u_{\varepsilon} = u$ on $B_R(x_0) \times \{t_0 - R^2\}$ has to be understood in the usual L^2 -sense (4.4), while the condition $u_{\varepsilon} = u$ on the lateral boundary $\partial B_R(x_0) \times (t_0 - R^2, t_0)$ has to be meant in the sense of traces, i.e.

$$(u_{\varepsilon} - u) (\cdot, t) \in W_0^{1,p}(B_R(x_0))$$

for almost every $t \in (t_0 - R^2, t_0)$.

Remark 4.3. The advantage of considering the Cauchy-Dirichlet problem (4.2) stems from the fact that the existence of a unique solution

$$u_{\varepsilon} \in C^{0}\left([t_{0} - R^{2}, t_{0}]; L^{2}(B_{R}(x_{0}))\right) \cap L^{p}\left(t_{0} - R^{2}, t_{0}; W^{1,p}(B_{R}(x_{0}))\right)$$

satisfying the requirements of Definition 4.2 can be ensured by the classic existence theory for parabolic equations (see, for example, [22, Chapter 2, Theorem 1.2 and Remark 1.2]). Moreover, the operator $D_{\xi}A_{\varepsilon}(\xi)$ fulfills *p*-growth and *p*-ellipticity conditions with constants depending on ε (cf. Lemma 4.1 above). Therefore, by the results in [13], we have

$$V_1(Du_{\varepsilon}) := (1 + |Du_{\varepsilon}|^2)^{\frac{p-2}{4}} Du_{\varepsilon} \in L^2_{loc} \left(t_0 - R^2, t_0; W^{1,2}_{loc}(B_R(x_0), \mathbb{R}^n) \right)$$

and

$$Du_{\varepsilon} \in L^{p+\frac{4}{n}}_{loc}(Q_R(z_0), \mathbb{R}^n),$$

and, by the definition of $V_1(Du_{\varepsilon})$, this yields

$$DV_1(Du_{\varepsilon}) \approx (1 + |Du_{\varepsilon}|^2)^{\frac{p-2}{4}} D^2 u_{\varepsilon} \in L^2_{loc}(Q_R(z_0), \mathbb{R}^{n \times n}) \implies |D^2 u_{\varepsilon}| \in L^2_{loc}(Q_R(z_0)).$$

4.1 Uniform a priori estimates

The first step in the proofs of Theorems 1.1 and 1.3 is the following estimate for the weak solutions of the regularized problems (4.2). Similar estimates are scattered in the literature, but not in the exact form needed here.

Proposition 4.4 (Uniform energy estimate). With the notation and under the assumptions above, there exist two positive constants $\varepsilon_0 \leq 1$ and $C \equiv C(n, p, R)$ such that

$$\int_{Q_R(z_0)} |Du_{\varepsilon}|^p dz + \sup_{t \in (t_0 - R^2, t_0)} \left\| u_{\varepsilon}(\cdot, t) - u(\cdot, t) \right\|_{L^2(B_R(x_0))}^2 \leq C \left(\|Du\|_{L^p(Q_R(z_0))}^p + \lambda^p + 1 \right)$$
(4.5)
for all $\varepsilon \in (0, \varepsilon_0]$.

Proof. In order to obtain an (uniform in ε) energy estimate for $|Du_{\varepsilon}|$, we proceed by testing equations (1.1) and (4.2)₁ with the map $\varphi = \psi(t)(u_{\varepsilon} - u)$, where $\psi \in W^{1,\infty}(\mathbb{R})$ is chosen such that

$$\psi(t) = \begin{cases} 1 & \text{if } t \le t_2, \\ -\frac{1}{\omega}(t - t_2 - \omega) & \text{if } t_2 < t < t_2 + \omega, \\ 0 & \text{if } t \ge t_2 + \omega, \end{cases}$$

with

 $t_0 - R^2 < t_2 < t_2 + \omega < t_0,$

and then letting $\omega \to 0$. We observe that, at this stage, it is important that u_{ε} and u agree on the parabolic boundary $\partial_{\text{par}}Q_R(z_0)$. We also note that the following computations are somewhat formal concerning the use of the time derivative, but they can easily be made rigorous, for example by the use of Steklov averages. We skip this, since it is a standard procedure. With the previous choice of φ , for every $t_2 \in (t_0 - R^2, t_0)$ we find

$$\frac{1}{2} \int_{B_R(x_0)} |u_{\varepsilon}(x,t_2) - u(x,t_2)|^2 dx + \int_{Q_{R,t_2}} \langle H_{p-1}(Du_{\varepsilon}) - H_{p-1}(Du), Du_{\varepsilon} - Du \rangle dz
+ \varepsilon \int_{Q_{R,t_2}} \langle (1+|Du_{\varepsilon}|^2)^{\frac{p-2}{2}} Du_{\varepsilon}, Du_{\varepsilon} - Du \rangle dz = \int_{Q_{R,t_2}} (f-f_{\varepsilon})(u_{\varepsilon}-u) dz,$$
(4.6)

where we have used the abbreviation

$$Q_{R,t_2} = B_R(x_0) \times (t_0 - R^2, t_2).$$

In what follows, we shall denote by c_k some positive constants depending only on n, p and R. Using Lemma 2.4, the Cauchy-Schwarz inequality as well as Young's inequality with $\beta > 0$, from (4.6) we infer

$$\sup_{t \in (t_0 - R^2, t_0)} \left\| u_{\varepsilon}(\cdot, t) - u(\cdot, t) \right\|_{L^2(B_R)}^2 + \frac{8}{p^2} \int_{Q_R(z_0)} \left| H_{\frac{p}{2}}(Du_{\varepsilon}) - H_{\frac{p}{2}}(Du) \right|^2 dz + 2\varepsilon \int_{Q_R(z_0)} |Du_{\varepsilon}|^p dz \\
\leq 2\varepsilon \int_{Q_R(z_0)} (1 + |Du_{\varepsilon}|^2)^{\frac{p-1}{2}} |Du| dz + 2 \int_{Q_R(z_0)} |f - f_{\varepsilon}| |u_{\varepsilon} - u| dz \\
\leq \frac{2\varepsilon \beta^{p'}}{p'} \int_{Q_R} (1 + |Du_{\varepsilon}|^2)^{\frac{p}{2}} dz + \frac{2\varepsilon}{p\beta^p} \int_{Q_R} |Du|^p dz + 2 \int_{Q_R} |f - f_{\varepsilon}| |u_{\varepsilon} - u| dz \\
\leq \frac{2^{\frac{p}{2}} \varepsilon \beta^{p'}}{p'} \int_{Q_R} |Du_{\varepsilon}|^p dz + \frac{2^{\frac{p}{2}} \varepsilon \beta^{p'}}{p'} |Q_R| + \frac{2\varepsilon}{p\beta^p} \int_{Q_R} |Du|^p dz + 2 \int_{Q_R} |f - f_{\varepsilon}| |u_{\varepsilon} - u| dz.$$
(4.7)

Choosing $\beta = \left(\frac{p'}{2^{p/2}}\right)^{\frac{1}{p'}}$ and reabsorbing the first integral in the right-hand side of (4.7) by the left-hand side, we arrive at

$$\sup_{t \in (t_0 - R^2, t_0)} \left\| u_{\varepsilon}(\cdot, t) - u(\cdot, t) \right\|_{L^2(B_R)}^2 + \int_{Q_R} \left| H_{\frac{p}{2}}(Du_{\varepsilon}) - H_{\frac{p}{2}}(Du) \right|^2 dz + \varepsilon \int_{Q_R} |Du_{\varepsilon}|^p dz$$

$$\leq c_1 \varepsilon + c_1 \varepsilon \int_{Q_R} |Du|^p dz + c_1 \int_{Q_R} |f - f_{\varepsilon}| |u_{\varepsilon} - u| dz.$$
(4.8)

Now we apply Hölder's and Poincaré's inequalities to estimate the last integral as follows

$$\int_{Q_R} |f - f_{\varepsilon}| |u_{\varepsilon} - u| dz \leq ||f - f_{\varepsilon}||_{L^{p'}(Q_R)} ||u_{\varepsilon} - u||_{L^p(Q_R)}$$

$$\leq c_2 ||f - f_{\varepsilon}||_{L^{p'}(Q_R)} ||Du_{\varepsilon} - Du||_{L^p(Q_R)}.$$
(4.9)

Joining estimates (4.8) and (4.9), recalling that $0 < \varepsilon \leq 1$ and applying Young's inequality with $\gamma > 0$, we obtain

$$\sup_{t \in (t_0 - R^2, t_0)} \left\| u_{\varepsilon}(\cdot, t) - u(\cdot, t) \right\|_{L^2(B_R)}^2 + \int_{Q_R} \left| H_{\frac{p}{2}}(Du_{\varepsilon}) - H_{\frac{p}{2}}(Du) \right|^2 dz + \varepsilon \int_{Q_R} |Du_{\varepsilon}|^p dz$$

$$\leq c_3 + c_3 \int_{Q_R} |Du|^p dz + c_3 \gamma^{\frac{1}{1-p}} \|f - f_{\varepsilon}\|_{L^{p'}(Q_R)}^{p'} + \gamma \|Du_{\varepsilon} - Du\|_{L^p(Q_R)}^p$$

$$\leq c_3 + (c_3 + 2^{p-1}\gamma) \int_{Q_R} |Du|^p dz + 2^{p-1}\gamma \int_{Q_R} |Du_{\varepsilon}|^p dz + c_3 \gamma^{\frac{1}{1-p}} \|f - f_{\varepsilon}\|_{L^{p'}(Q_R)}^{p'}. \quad (4.10)$$

Now, arguing as in [6, Formula (4.20)], we have

$$\int_{Q_R} |Du_{\varepsilon}|^p dz \le 2^p \int_{Q_R} \left| H_{\frac{p}{2}}(Du_{\varepsilon}) - H_{\frac{p}{2}}(Du) \right|^2 dz + 2^{p+1} \int_{Q_R} (|Du|^p + \lambda^p) dz.$$
(4.11)

Multiplying all sides of (4.10) by 2^p and then adding the resulting expression and (4.11) side by side, we get

$$(1+2^{p}\varepsilon)\int_{Q_{R}}|Du_{\varepsilon}|^{p}dz + 2^{p}\sup_{t\in(t_{0}-R^{2},t_{0})}\left\|u_{\varepsilon}(\cdot,t)-u(\cdot,t)\right\|_{L^{2}(B_{R})}^{2}$$

$$\leq c_4 + (c_4 + 2^{2p-1}\gamma) \int_{Q_R} (|Du|^p + \lambda^p) \, dz + c_4 \, \gamma^{\frac{1}{1-p}} \|f - f_{\varepsilon}\|_{L^{p'}(Q_R)}^{p'} + 2^{2p-1}\gamma \int_{Q_R} |Du_{\varepsilon}|^p \, dz.$$

Choosing $\gamma = \frac{1}{2^{2p}}$ and reabsorbing the last term in the above estimate by the left-hand side, we find

$$\int_{Q_R} |Du_{\varepsilon}|^p dz + \sup_{t \in (t_0 - R^2, t_0)} \left\| u_{\varepsilon}(\cdot, t) - u(\cdot, t) \right\|_{L^2(B_R)}^2 \\
\leq c_5 \int_{Q_R} (|Du|^p + \lambda^p) dz + c_5 \left\| f - f_{\varepsilon} \right\|_{L^{p'}(Q_R)}^{p'} + c_5.$$
(4.12)

Since, by virtue of (4.1),

$$f_{\varepsilon} \to f$$
 strongly in $L^{p'}(Q_R)$ as $\varepsilon \to 0^+$, (4.13)

there exists a positive number $\varepsilon_0 \leq 1$ such that

$$\|f - f_{\varepsilon}\|_{L^{p'}(Q_R)} \le 1 \qquad \text{for all } \varepsilon \in (0, \varepsilon_0].$$
(4.14)

Combining this with (4.12), we obtain the desired conclusion.

Thanks to the previous proposition, we can now obtain the following result, which will be decisive to carry out the proofs of Theorems 1.1 and 1.3.

Proposition 4.5 (Comparison estimate). With the notation and under the assumptions of Proposition 4.4, there exists a positive constant C depending only on n, p and R such that the estimate

$$\sup_{t \in (t_0 - R^2, t_0)} \left\| u_{\varepsilon}(\cdot, t) - u(\cdot, t) \right\|_{L^2(B_R(x_0))}^2 + \int_{Q_R(z_0)} \left| \mathcal{V}_{\alpha, \lambda}(Du_{\varepsilon}) - \mathcal{V}_{\alpha, \lambda}(Du) \right|^2 dz$$

$$\leq C \varepsilon \left(\left\| Du \right\|_{L^p(Q_R(z_0))}^p + 1 \right) + C \left\| f - f_{\varepsilon} \right\|_{L^{p'}(Q_R(z_0))} \left(\left\| Du \right\|_{L^p(Q_R(z_0))} + \lambda + 1 \right)$$
(4.15)

holds for every $\varepsilon \in (0, \varepsilon_0]$, where ε_0 is the constant from Proposition 4.4. In particular, this estimate implies that

 $u_{\varepsilon} \to u$ strongly in $L^2(Q_R(z_0))$ as $\varepsilon \to 0$

and

$$\mathcal{V}_{\alpha,\lambda}(Du_{\varepsilon}) \to \mathcal{V}_{\alpha,\lambda}(Du) \quad strongly \ in \ L^2(Q_R(z_0),\mathbb{R}^n) \quad as \ \varepsilon \to 0$$

Proof. Arguing as in the proof of Proposition 4.4, but using Lemma 2.6 instead of Lemma 2.4, we arrive at

$$\sup_{t \in (t_0 - R^2, t_0)} \left\| u_{\varepsilon}(\cdot, t) - u(\cdot, t) \right\|_{L^2(B_R)}^2 + \int_{Q_R} \left| \mathcal{V}_{\alpha, \lambda}(Du_{\varepsilon}) - \mathcal{V}_{\alpha, \lambda}(Du) \right|^2 dz$$

$$\leq c \varepsilon \left(\|Du\|_{L^p(Q_R)}^p + 1 \right) + c \|f - f_{\varepsilon}\|_{L^{p'}(Q_R)} \|Du_{\varepsilon} - Du\|_{L^p(Q_R)},$$

where $c \equiv c(n, p, R) > 0$. Now, let us consider the same $\varepsilon_0 \in (0, 1]$ as in the proof of Proposition 4.4. Then, applying Minkowski's inequality, for every $\varepsilon \in (0, \varepsilon_0]$ we get

$$\sup_{t \in (t_0 - R^2, t_0)} \left\| u_{\varepsilon}(\cdot, t) - u(\cdot, t) \right\|_{L^2(B_R)}^2 + \int_{Q_R} \left| \mathcal{V}_{\alpha, \lambda}(Du_{\varepsilon}) - \mathcal{V}_{\alpha, \lambda}(Du) \right|^2 dz$$

$$\leq c \varepsilon \left(\|Du\|_{L^{p}(Q_{R})}^{p} + 1 \right) + c \|f - f_{\varepsilon}\|_{L^{p'}(Q_{R})} \left(\|Du_{\varepsilon}\|_{L^{p}(Q_{R})} + \|Du\|_{L^{p}(Q_{R})} \right)$$

$$\leq C \varepsilon \left(\|Du\|_{L^{p}(Q_{R})}^{p} + 1 \right) + C \|f - f_{\varepsilon}\|_{L^{p'}(Q_{R})} \left(\|Du\|_{L^{p}(Q_{R})} + \lambda + 1 \right),$$

where, in the last line, we have used inequality (4.5). Thus we obtain the comparison estimate (4.15). Now, just observe that

$$\int_{Q_R(z_0)} |u_{\varepsilon} - u|^2 dz \leq R^2 \sup_{t \in (t_0 - R^2, t_0)} \left\| u_{\varepsilon}(\cdot, t) - u(\cdot, t) \right\|_{L^2(B_R(x_0))}^2.$$

Combining this with (4.15) and (4.13), the statement is proved.

We are now in a position to establish uniform estimates for the derivatives of the function $\mathcal{V}_{\alpha,\lambda}(Du_{\varepsilon})$. Let us start with the following result.

Proposition 4.6 (Uniform Sobolev estimate). Under the assumptions of Theorem 1.1 and with the notation above, there exists a positive number $\varepsilon_1 \leq 1$ such that, for every $\varepsilon \in (0, \varepsilon_1]$, for every parabolic cylinder $Q_{\rho}(z_0) \Subset Q_R(z_0)$ and every $j \in \{1, \ldots, n\}$, we have

$$\int_{Q_{\rho/2}(z_0)} |D_{x_j} \mathcal{V}_{\alpha,\lambda}(Du_{\varepsilon})|^2 dz
\leq \frac{C}{\rho^2} \left(\|Du\|_{L^p(Q_R)}^p + \|Du\|_{L^p(Q_R)}^2 + \lambda^p + \lambda^2 + 1 \right) + C \|f\|_{L^{p'}\left(t_0 - R^2, t_0; B_{p', 1}^{\frac{p-2}{p}}(B_R(x_0))\right)}$$
(4.16)

for a positive constant C depending only on n, p and R in the case $\lambda = 0$, and additionally on α if $\lambda > 0$.

Proof. Let us first assume that $\lambda > 0$. To shorten our notation, we introduce the function $P: Q_R(z_0) \to \mathbb{R}_0^+$ defined by

$$P := (|Du_{\varepsilon}| - \lambda)_{+} \tag{4.17}$$

and its "mollified" version

$$P_{\varrho} := (|[Du_{\varepsilon}]_{\varrho}| - \lambda)_{+}, \qquad \varrho > 0, \tag{4.18}$$

with an intentional abuse of the notation (2.1) on the left-hand side of (4.18). We now test the weak formulation (4.3) with the map

$$\varphi = D_{x_j} \left(D_{x_j} [u_{\varepsilon}]_{\varrho} \cdot \psi \cdot \Phi(P_{\varrho}) \right), \qquad j \in \{1, \dots, n\},$$

where $\psi \in W_0^{1,\infty}(Q_R(z_0))$ is a non-negative cut-off function that will be specified later and $\Phi : \mathbb{R}_0^+ \to \mathbb{R}_0^+$ is an increasing, locally Lipschitz continuous function, such that Φ and Φ' are bounded on \mathbb{R}_0^+ , $\Phi(0) = 0$ and

$$\Phi'(t) t \le c_{\Phi} \Phi(t) \tag{4.19}$$

for a suitable constant $c_{\Phi} > 0$. After summing with respect to j from 1 to n, we obtain

$$\frac{1}{2} \int_{Q_R} \partial_t (|D[u_\varepsilon]_\varrho|^2) \psi \Phi(P_\varrho) dz + \int_{Q_R} \sum_{j=1}^n \langle D_{x_j} [DA_\varepsilon (Du_\varepsilon)]_\varrho, D\left(D_{x_j} [u_\varepsilon]_\varrho \cdot \psi \Phi(P_\varrho)\right) \rangle dz$$

$$= -\int_{Q_R} \sum_{j=1}^n [f_\varepsilon]_\varrho D_{x_j} \left(D_{x_j} [u_\varepsilon]_\varrho \cdot \psi \Phi(P_\varrho) \right) dz.$$
(4.20)

Arguing as in [3, page 521], the integral involving the time derivative can be expressed as follows

$$\frac{1}{2} \int_{Q_R} \partial_t (|D[u_\varepsilon]_\varrho|^2) \psi \Phi(P_\varrho) dz = \frac{1}{2} \int_{Q_R} \partial_t \left[\int_0^{|D[u_\varepsilon]_\varrho|^2} \Phi((\sqrt{w} - \lambda)_+) dw \right] \psi dz$$
$$= -\frac{1}{2} \int_{Q_R} \left[\int_0^{|D[u_\varepsilon]_\varrho|^2} \Phi((\sqrt{w} - \lambda)_+) dw \right] \partial_t \psi dz.$$

Therefore, integrating by parts and then letting $\rho \to 0$ in (4.20), we get

$$-\frac{1}{2}\int_{Q_R} \left[\int_0^{|Du_{\varepsilon}|^2} \Phi((\sqrt{w} - \lambda)_+) dw \right] \partial_t \psi \, dz$$

+
$$\int_{Q_R} \sum_{j=1}^n \langle D^2 A_{\varepsilon}(Du_{\varepsilon}) D(D_{x_j}u_{\varepsilon}), D(D_{x_j}u_{\varepsilon}) \rangle \psi \Phi(P) \, dz$$

+
$$\int_{Q_R} \sum_{j=1}^n \langle D^2 A_{\varepsilon}(Du_{\varepsilon}) D(D_{x_j}u_{\varepsilon}), D\psi \rangle (D_{x_j}u_{\varepsilon}) \Phi(P) \, dz$$

+
$$\int_{Q_R} \sum_{j=1}^n \langle D^2 A_{\varepsilon}(Du_{\varepsilon}) D(D_{x_j}u_{\varepsilon}), DP \rangle (D_{x_j}u_{\varepsilon}) \psi \Phi'(P) \, dz$$

=
$$\int_{Q_R} \sum_{j=1}^n (D_{x_j}f_{\varepsilon}) (D_{x_j}u_{\varepsilon}) \psi \Phi(P) \, dz. \qquad (4.21)$$

In what follows, we will denote by c_k and C some positive constants that do not depend on ε . Now, let us fix an arbitrary radius $\rho \in (0, R)$. For a fixed time $t_1 \in (t_0 - \rho^2, t_0)$ and $\sigma \in (0, t_0 - t_1)$, we choose

$$\psi(x,t) = \widetilde{\chi}(t) \,\chi(t) \,\eta^2(x)$$

with $\chi \in W^{1,\infty}((t_0-R^2,t_0),[0,1]), \chi \equiv 0$ on $(t_0-R^2,t_0-\rho^2)$ and $\partial_t \chi \ge 0, \eta \in C_0^{\infty}(B_{\rho}(x_0),[0,1]),$ and with the Lipschitz continuous function $\widetilde{\chi}: (t_0-R^2,t_0) \to \mathbb{R}$ defined by

$$\widetilde{\chi}(t) = \begin{cases} 1 & \text{if } t \leq t_1, \\ \text{affine} & \text{if } t_1 < t < t_1 + \sigma, \\ 0 & \text{if } t \geq t_1 + \sigma. \end{cases}$$

With such a choice of ψ , equation (4.21) turns into

$$\begin{split} &-\frac{1}{2} \int_{Q_R} \left[\int_0^{|Du_{\varepsilon}|^2} \Phi((\sqrt{w} - \lambda)_+) \, dw \right] \eta^2(x) \, \chi(t) \, \partial_t \widetilde{\chi}(t) \, dz \\ &-\frac{1}{2} \int_{Q_R} \left[\int_0^{|Du_{\varepsilon}|^2} \Phi((\sqrt{w} - \lambda)_+) \, dw \right] \eta^2(x) \, \widetilde{\chi}(t) \, \partial_t \chi(t) \, dz \\ &+ \int_{Q_R} \sum_{j=1}^n \left\langle D^2 A_{\varepsilon}(Du_{\varepsilon}) \, D(D_{x_j} u_{\varepsilon}), D(D_{x_j} u_{\varepsilon}) \right\rangle \eta^2(x) \, \widetilde{\chi}(t) \, \chi(t) \, \Phi(P) \, dz \\ &+ \int_{Q_R} \sum_{j=1}^n \left\langle D^2 A_{\varepsilon}(Du_{\varepsilon}) \, D(D_{x_j} u_{\varepsilon}), DP \right\rangle(D_{x_j} u_{\varepsilon}) \, \eta^2(x) \, \widetilde{\chi}(t) \, \chi(t) \, \Phi'(P) \, dz \end{split}$$

$$= -2 \int_{Q_R} \sum_{j=1}^n \langle D^2 A_{\varepsilon}(Du_{\varepsilon}) D(D_{x_j}u_{\varepsilon}), D\eta \rangle (D_{x_j}u_{\varepsilon}) \eta(x) \,\widetilde{\chi}(t) \,\chi(t) \,\Phi(P) \,dz$$
$$+ \int_{Q_R} \sum_{j=1}^n \left(D_{x_j} f_{\varepsilon} \right) \left(D_{x_j} u_{\varepsilon} \right) \eta^2(x) \,\widetilde{\chi}(t) \,\chi(t) \,\Phi(P) \,dz.$$

Setting

$$Q^{t_1} := B_{\rho}(x_0) \times (t_0 - \rho^2, t_1)$$
(4.22)

and letting $\sigma \to 0$ in the previous equality, for every $t_1 \in (t_0 - \rho^2, t_0)$ we get

$$\frac{1}{2} \int_{B_{\rho}(x_{0})} \chi(t_{1}) \eta^{2}(x) \left[\int_{0}^{|Du_{\varepsilon}(x,t_{1})|^{2}} \Phi((\sqrt{w}-\lambda)_{+}) dw \right] dx$$

$$+ \int_{Q^{t_{1}}} \sum_{j=1}^{n} \langle D^{2}A_{\varepsilon}(Du_{\varepsilon}) D(D_{x_{j}}u_{\varepsilon}), D(D_{x_{j}}u_{\varepsilon}) \rangle \eta^{2}(x) \chi(t) \Phi(P) dz$$

$$+ \int_{Q^{t_{1}}} \sum_{j=1}^{n} \langle D^{2}A_{\varepsilon}(Du_{\varepsilon}) D(D_{x_{j}}u_{\varepsilon}), DP \rangle(D_{x_{j}}u_{\varepsilon}) \eta^{2}(x) \chi(t) \Phi'(P) dz$$

$$= -2 \int_{Q^{t_{1}}} \sum_{j=1}^{n} \langle D^{2}A_{\varepsilon}(Du_{\varepsilon}) D(D_{x_{j}}u_{\varepsilon}), D\eta \rangle(D_{x_{j}}u_{\varepsilon}) \eta(x) \chi(t) \Phi(P) dz$$

$$+ \int_{Q^{t_{1}}} \sum_{j=1}^{n} (D_{x_{j}}f_{\varepsilon}) (D_{x_{j}}u_{\varepsilon}) \eta^{2}(x) \chi(t) \Phi(P) dz$$

$$+ \frac{1}{2} \int_{Q^{t_{1}}} (\partial_{t}\chi) \eta^{2}(x) \left[\int_{0}^{|Du_{\varepsilon}(x,t)|^{2}} \Phi((\sqrt{w}-\lambda)_{+}) dw \right] dz, \qquad (4.23)$$

where we have used that $\partial_t \tilde{\chi}$ converges to a Dirac delta distribution as $\sigma \to 0$, together with the $L^2(B_R(x_0))$ -valued continuity of u_{ε} . We now observe that the second integral on the left-hand side of (4.23) is non-negative, since A_{ε} is convex and $\chi, \Phi \geq 0$. As for the first term on the right-hand side of (4.23), for every $t_1 \in (t_0 - \rho^2, t_0)$ we have

$$-2\int_{Q^{t_1}} \langle D^2 A_{\varepsilon}(Du_{\varepsilon}) D(D_{x_j}u_{\varepsilon}), D\eta \rangle \langle D_{x_j}u_{\varepsilon} \rangle \eta(x) \chi(t) \Phi(P) dz$$

$$\leq 2\int_{Q^{t_1}} |\langle D^2 A_{\varepsilon}(Du_{\varepsilon}) D(D_{x_j}u_{\varepsilon}), D\eta \rangle| \chi(t) \eta(x) |D_{x_j}u_{\varepsilon}| \Phi(P) dz$$

$$\leq 2\int_{Q^{t_1}} \sqrt{\langle D^2 A_{\varepsilon}(Du_{\varepsilon}) D(D_{x_j}u_{\varepsilon}), D(D_{x_j}u_{\varepsilon}) \rangle} \sqrt{\langle D^2 A_{\varepsilon}(Du_{\varepsilon}) D\eta, D\eta \rangle} \chi(t) \eta(x) |D_{x_j}u_{\varepsilon}| \Phi(P) dz$$

$$\leq \frac{1}{2}\int_{Q^{t_1}} \langle D^2 A_{\varepsilon}(Du_{\varepsilon}) D(D_{x_j}u_{\varepsilon}), D(D_{x_j}u_{\varepsilon}) \rangle \chi(t) \eta^2(x) \Phi(P) dz$$

$$+ 2\int_{Q^{t_1}} \langle D^2 A_{\varepsilon}(Du_{\varepsilon}) D\eta, D\eta \rangle \chi(t) |D_{x_j}u_{\varepsilon}|^2 \Phi(P) dz, \qquad (4.24)$$

where we have used the Cauchy-Schwarz inequality together with Young's inequality. Joining (4.23) and (4.24), we obtain

$$I_1 + I_2 + I_3 \le I_4 + I_5 + I_6, \qquad (4.25)$$

where

$$I_1 := \int_{B_{\rho}(x_0)} \chi(t_1) \, \eta^2(x) \left[\int_0^{|Du_{\varepsilon}(x,t_1)|^2} \Phi((\sqrt{w} - \lambda)_+) \, dw \right] dx,$$

$$\begin{split} I_2 &:= 2 \int_{Q^{t_1}} \sum_{j=1}^n \langle D^2 A_{\varepsilon}(Du_{\varepsilon}) D(D_{x_j}u_{\varepsilon}), DP \rangle (D_{x_j}u_{\varepsilon}) \eta^2(x) \, \chi(t) \, \Phi'(P) \, dz, \\ I_3 &:= \int_{Q^{t_1}} \sum_{j=1}^n \langle D^2 A_{\varepsilon}(Du_{\varepsilon}) D(D_{x_j}u_{\varepsilon}), D(D_{x_j}u_{\varepsilon}) \rangle \, \chi(t) \, \eta^2(x) \, \Phi(P) \, dz, \\ I_4 &:= 4 \int_{Q^{t_1}} \sum_{j=1}^n \langle D^2 A_{\varepsilon}(Du_{\varepsilon}) D\eta, D\eta \rangle \, \chi(t) \, |D_{x_j}u_{\varepsilon}|^2 \, \Phi(P) \, dz, \\ I_5 &:= \int_{Q^{t_1}} (\partial_t \chi) \, \eta^2(x) \left[\int_0^{|Du_{\varepsilon}(x,t)|^2} \Phi((\sqrt{w} - \lambda)_+) \, dw \right] dz, \\ I_6 &:= 2 \int_{Q^{t_1}} \sum_{j=1}^n \left(D_{x_j} f_{\varepsilon} \right) \left(D_{x_j} u_{\varepsilon} \right) \eta^2(x) \, \chi(t) \, \Phi(P) \, dz. \end{split}$$

Now we observe that $I_1 \ge 0$, thus we can drop it in the following. Furthermore, arguing as in the proof of [5, Proposition 5.1], one can show that $I_2 \ge 0$. Therefore, inequality (4.25) boils down to

$$I_3 \le I_4 + I_5 + I_6 \,. \tag{4.26}$$

At this stage, we choose a cut-off function $\eta \in C_0^{\infty}(B_{\rho}(x_0))$ with $\eta \equiv 1$ on $B_{\rho/2}(x_0)$ such that

$$0 \le \eta \le 1$$
 and $|D\eta| \le \frac{\tilde{c}}{\rho}$. (4.27)

For the cut-off function in time, we choose the piecewise affine function $\chi : (t_0 - R^2, t_0) \rightarrow [0, 1]$ with

$$\chi \equiv 0 \quad \text{on } (t_0 - R^2, t_0 - \rho^2),$$
$$\chi \equiv 1 \quad \text{on } \left(t_0 - \left(\frac{\rho}{2}\right)^2, t_0\right),$$
$$\partial_t \chi \equiv \frac{4}{3\rho^2} \quad \text{on } \left(t_0 - \rho^2, t_0 - \left(\frac{\rho}{2}\right)^2\right)$$

Moreover, as in [5] we choose

$$\Phi(t) := \frac{t^{2\alpha}}{(t^2 + \lambda^2)^{\alpha}} \quad \text{for } t \ge 0,$$
(4.28)

and therefore

$$\Phi'(t) = \frac{2\alpha\lambda^2 t^{2\alpha-1}}{(t^2 + \lambda^2)^{\alpha+1}}.$$
(4.29)

This function satisfies (4.19) with $c_{\Phi} = 2\alpha$. With the above choices, we now estimate I_3 , I_4 and I_5 separately. Let us first consider I_3 . Recalling the definition of P in (4.17), by Lemma 4.1 we have

$$I_3 \ge \int_{Q^{t_1}} \frac{(|Du_{\varepsilon}| - \lambda)_+^{p-1+2\alpha}}{|Du_{\varepsilon}| [\lambda^2 + (|Du_{\varepsilon}| - \lambda)_+^2]^{\alpha}} |D^2 u_{\varepsilon}|^2 \chi(t) \eta^2(x) dz.$$

$$(4.30)$$

Using Lemma 4.1 again, the fact that $\varepsilon, \chi, \Phi \leq 1$ and the properties (4.27) of η , we obtain

$$I_4 \leq 8(p-1) \int_{Q_R} (1+|Du_{\varepsilon}|^2)^{\frac{p}{2}} |D\eta|^2 dz \leq \frac{c_1(p)}{\rho^2} \int_{Q_R} (1+|Du_{\varepsilon}|^2)^{\frac{p}{2}} dz.$$
(4.31)

Now we use again the fact that $\Phi \leq 1$, the properties of χ and η , and Hölder's inequality, in order to get

$$I_{5} \leq \int_{Q^{t_{1}}} (\partial_{t}\chi) \eta^{2}(x) |Du_{\varepsilon}|^{2} dz \leq \frac{4}{3\rho^{2}} \int_{Q_{R}} |Du_{\varepsilon}|^{2} dz \leq \frac{c_{2}}{\rho^{2}} ||Du_{\varepsilon}||^{2}_{L^{p}(Q_{R})}, \qquad (4.32)$$

where $c_2 \equiv c_2(n, p, R) > 0$. Combining estimates (4.26), (4.30), (4.31) and (4.32), we find

$$\int_{Q^{t_1}} \frac{(|Du_{\varepsilon}| - \lambda)_{+}^{p-1+2\alpha}}{|Du_{\varepsilon}| [\lambda^2 + (|Du_{\varepsilon}| - \lambda)_{+}^2]^{\alpha}} |D^2 u_{\varepsilon}|^2 \chi(t) \eta^2(x) dz
\leq \frac{c_2}{\rho^2} \left[\int_{Q_R} (1 + |Du_{\varepsilon}|^2)^{\frac{p}{2}} dz + \|Du_{\varepsilon}\|_{L^p(Q_R)}^2 \right]
+ 2 \sum_{j=1}^n \int_{Q^{t_1}} (D_{x_j} f_{\varepsilon}) (D_{x_j} u_{\varepsilon}) \eta^2(x) \chi(t) \Phi(P) dz,$$
(4.33)

which holds for every $t_1 \in (t_0 - \rho^2, t_0)$.

It remains to estimate the integral containing $D_{x_j} f_{\varepsilon}$. With this aim in mind, we now argue as in the proof of [5, Proposition 5.1]. By Theorem 3.3, there exists a linear and bounded extension operator

$$\operatorname{ext}_{B_{\rho}(x_0)} : B_{p',1}^{-2/p}(B_{\rho}(x_0)) \hookrightarrow B_{p',1}^{-2/p}(\mathbb{R}^n)$$

such that $\operatorname{re}_{B_{\rho}(x_0)} \circ \operatorname{ext}_{B_{\rho}(x_0)} = \operatorname{id}$, where $\operatorname{re}_{B_{\rho}(x_0)}$ is the restriction operator defined in Section 3 and the symbol id denotes the identity in $B_{p',1}^{-2/p}(B_{\rho}(x_0))$. Since, for almost every $t \in (t_0 - \rho^2, t_0)$, we have $D_{x_j} f_{\varepsilon}(\cdot, t) = \operatorname{ext}_{B_{\rho}(x_0)}(D_{x_j} f_{\varepsilon}(\cdot, t))$ almost everywhere in $B_{\rho}(x_0)$, we find

$$\int_{Q^{t_1}} (D_{x_j} f_{\varepsilon}) (D_{x_j} u_{\varepsilon}) \eta^2(x) \chi(t) \Phi(P) dz = \int_{Q^{t_1}} \operatorname{ext}_{B_{\rho}(x_0)} (D_{x_j} f_{\varepsilon}(x,t)) \cdot (D_{x_j} u_{\varepsilon}) \eta^2(x) \chi(t) \Phi(P) dz.$$
(4.34)

By definition of dual norm, we get

$$\left| \int_{Q^{t_1}} \operatorname{ext}_{B_{\rho}(x_0)}(D_{x_j}f_{\varepsilon}(x,t)) \cdot (D_{x_j}u_{\varepsilon}) \eta^2(x) \chi(t) \Phi(P) dz \right|$$

$$\leq \int_{t_0-\rho^2}^{t_1} \left\| \operatorname{ext}_{B_{\rho}(x_0)}(D_{x_j}f_{\varepsilon}(\cdot,t)) \right\|_{(\mathring{B}^{2/p}_{p,\infty}(\mathbb{R}^n))'} \left[(D_{x_j}u_{\varepsilon}(\cdot,t)) \eta^2 \chi(t) \Phi(P(\cdot,t)) \right]_{B^{2/p}_{p,\infty}(\mathbb{R}^n)} dt.$$

$$(4.35)$$

Using Theorem 3.5, for almost every $t \in (t_0 - \rho^2, t_0)$ we obtain

$$\| \exp_{B_{\rho}(x_{0})}(D_{x_{j}}f_{\varepsilon}(\cdot,t)) \|_{(\dot{B}^{2/p}_{p,\infty}(\mathbb{R}^{n}))'} \leq c_{3} \| \exp_{B_{\rho}(x_{0})}(D_{x_{j}}f_{\varepsilon}(\cdot,t)) \|_{B^{-2/p}_{p',1}(\mathbb{R}^{n})},$$

for some positive constant $c_3 \equiv c_3(n, p)$. Combining the above inequality with the boundedness of the operator $\operatorname{ext}_{B_{\rho}(x_0)}$ yields

$$\| \exp_{B_{\rho}(x_{0})}(D_{x_{j}}f_{\varepsilon}(\cdot,t)) \|_{(\dot{B}^{2/p}_{p,\infty}(\mathbb{R}^{n}))'} \leq c_{3} \| D_{x_{j}}f_{\varepsilon}(\cdot,t) \|_{B^{-2/p}_{p',1}(B_{\rho}(x_{0}))}$$

for almost every $t \in (t_0 - \rho^2, t_0)$. Moreover, applying Theorem 3.6, we find that

$$\|D_{x_j}f_{\varepsilon}(\cdot,t)\|_{B^{-2/p}_{p',1}(B_{\rho}(x_0))} \leq c_3 \|f_{\varepsilon}(\cdot,t)\|_{B^{\frac{p-2}{p}}_{p',1}(B_{\rho}(x_0))}$$

for almost every $t \in (t_0 - \rho^2, t_0)$. Combining the preceding inequalities, we infer

$$\| \exp_{B_{\rho}(x_{0})}(D_{x_{j}}f_{\varepsilon}(\cdot,t)) \|_{(\dot{B}^{2/p}_{p,\infty}(\mathbb{R}^{n}))'} \leq c_{3} \| f_{\varepsilon}(\cdot,t) \|_{B^{\frac{p-2}{p}}_{p',1}(B_{\rho}(x_{0}))},$$
(4.36)

which holds for almost every $t \in (t_0 - \rho^2, t_0)$ and a positive constant c_3 depending only on n and p. Now, recalling that

$$\left[\left(D_{x_j} u_{\varepsilon}(\cdot, t) \right) \eta^2 \chi(t) \Phi(P(\cdot, t)) \right]_{B^{2/p}_{p,\infty}(\mathbb{R}^n)}^p$$

=
$$\sup_{|h|>0} \frac{1}{|h|^2} \int_{\mathbb{R}^n} \chi^p(t) \left| \delta_h \left(\left(D_{x_j} u_{\varepsilon}(x, t) \right) \eta^2(x) \Phi(P(x, t)) \right) \right|^p dx$$

and using Lemma 2.3, we deduce

$$\frac{1}{|h|^2} \int_{\mathbb{R}^n} \chi^p(t) \left| \delta_h \left(\left(D_{x_j} u_\varepsilon(x,t) \right) \eta^2(x) \Phi(P(x,t)) \right) \right|^p dx \\
\leq \frac{c_4(p)}{|h|^2} \chi(t) \int_{\mathbb{R}^n} \left| \delta_h \left(\left| D_{x_j} u_\varepsilon(x,t) \right|^{\frac{p-2}{2}} (D_{x_j} u_\varepsilon(x,t)) \eta^p(x) \left[\Phi(P(x,t)) \right]^{\frac{p}{2}} \right) \right|^2 dx \\
\leq c_5(n,p) \chi(t) \int_{\mathbb{R}^n} \left| D \left(\left| D_{x_j} u_\varepsilon(x,t) \right|^{\frac{p-2}{2}} (D_{x_j} u_\varepsilon(x,t)) \eta^p(x) \left[\Phi(P(x,t)) \right]^{\frac{p}{2}} \right) \right|^2 dx,$$

where, in the last line, we have used the first statement in Lemma 2.9. Using the properties (4.27) of η and the boundedness of Φ , with simple manipulations we thus obtain

$$\begin{split} \left[\left(D_{x_j} u_{\varepsilon}(\cdot, t) \right) \eta^2 \chi(t) \Phi(P(\cdot, t)) \right]_{B^{2/p}_{p,\infty}(\mathbb{R}^n)}^p \\ &\leq c_5 \,\chi(t) \int_{B_{\rho}(x_0)} \left| D\left(|D_{x_j} u_{\varepsilon}(x, t)|^{\frac{p-2}{2}} (D_{x_j} u_{\varepsilon}(x, t)) \left[\Phi(P(x, t)) \right]^{\frac{p}{2}} \right) \right|^2 \eta^2(x) \, dx \\ &+ \frac{c_5}{\rho^2} \,\chi(t) \int_{B_{\rho}(x_0)} |D u_{\varepsilon}(x, t)|^p \, dx \end{split}$$

$$(4.37)$$

for almost every $t \in (t_0 - \rho^2, t_1)$. Now, from [5, Formula (5.17)] we know that

$$\left| D\left(\left| D_{x_j} u_{\varepsilon} \right|^{\frac{p-2}{2}} (D_{x_j} u_{\varepsilon}) \left[\Phi(P) \right]^{\frac{p}{2}} \right) \right|^2 \le c_6(n, p) \left(\mathbf{A}_1 + \mathbf{A}_2 \right), \tag{4.38}$$

where

$$\mathbf{A}_1 := |Du_{\varepsilon}|^{p-2} |D^2u_{\varepsilon}|^2 [\Phi(P)]^p$$

and

$$\mathbf{A}_2 := |Du_{\varepsilon}|^p |D^2 u_{\varepsilon}|^2 [\Phi(P)]^{p-2} [\Phi'(P)]^2.$$

At this point, we estimate \mathbf{A}_1 and \mathbf{A}_2 separately in the set where $|Du_{\varepsilon}| > \lambda$, since both \mathbf{A}_1 and \mathbf{A}_2 vanish in the set $\{|Du_{\varepsilon}| \leq \lambda\}$. Recalling the definitions of Φ and P in (4.28) and (4.17) respectively, we can write \mathbf{A}_1 as the product of two terms:

$$\mathbf{A}_{1} = \frac{|Du_{\varepsilon}|^{p-1} (|Du_{\varepsilon}| - \lambda)_{+}^{(2\alpha-1)(p-1)}}{[\lambda^{2} + (|Du_{\varepsilon}| - \lambda)_{+}^{2}]^{\alpha(p-1)}} \cdot \frac{|D^{2}u_{\varepsilon}|^{2} (|Du_{\varepsilon}| - \lambda)_{+}^{p-1+2\alpha}}{|Du_{\varepsilon}| [\lambda^{2} + (|Du_{\varepsilon}| - \lambda)_{+}^{2}]^{\alpha}}.$$

Then we have

$$\frac{|Du_{\varepsilon}|^{p-1} (|Du_{\varepsilon}| - \lambda)_{+}^{(2\alpha-1)(p-1)}}{[\lambda^{2} + (|Du_{\varepsilon}| - \lambda)_{+}^{2}]^{\alpha(p-1)}} \leq 2^{\alpha(p-1)} \frac{|Du_{\varepsilon}|^{2\alpha(p-1)}}{|Du_{\varepsilon}|^{2\alpha(p-1)}} = 2^{\alpha(p-1)},$$

where we have used that $2\alpha - 1 > 0$, since by assumption $\alpha \ge \frac{p+1}{2(p-1)} > \frac{1}{2}$. This implies that

$$\mathbf{A}_{1} \leq 2^{\alpha(p-1)} \frac{|D^{2}u_{\varepsilon}|^{2} (|Du_{\varepsilon}| - \lambda)_{+}^{p-1+2\alpha}}{|Du_{\varepsilon}| [\lambda^{2} + (|Du_{\varepsilon}| - \lambda)_{+}^{2}]^{\alpha}}.$$
(4.39)

Now we deal with A_2 . First, we observe that

$$[\Phi(t)]^{p-2} \, [\Phi'(t)]^2 = \frac{4 \, \alpha^2 \, \lambda^4 \, t^{2(\alpha p-1)}}{(t^2 + \lambda^2)^{\alpha p+2}}$$

Therefore, \mathbf{A}_2 can be written as follows:

$$\mathbf{A}_{2} = 4 \alpha^{2} \lambda^{4} \frac{|Du_{\varepsilon}|^{p+1} (|Du_{\varepsilon}| - \lambda)^{2\alpha(p-1)-p-1}_{+}}{[\lambda^{2} + (|Du_{\varepsilon}| - \lambda)^{2}_{+}]^{\alpha(p-1)+2}} \cdot \frac{|D^{2}u_{\varepsilon}|^{2} (|Du_{\varepsilon}| - \lambda)^{p-1+2\alpha}_{+}}{|Du_{\varepsilon}| [\lambda^{2} + (|Du_{\varepsilon}| - \lambda)^{2}_{+}]^{\alpha}}.$$

Again, the assumption $\alpha \geq \frac{p+1}{2(p-1)}$ implies $(|Du_{\varepsilon}| - \lambda)^{2\alpha(p-1)-p-1}_{+} \leq |Du_{\varepsilon}|^{2\alpha(p-1)-p-1}$, and so

$$\lambda^{4} \frac{|Du_{\varepsilon}|^{p+1} (|Du_{\varepsilon}| - \lambda)_{+}^{2\alpha(p-1)-p-1}}{[\lambda^{2} + (|Du_{\varepsilon}| - \lambda)_{+}^{2}]^{\alpha(p-1)+2}} \leq 2^{\alpha(p-1)+2} \lambda^{4} \frac{|Du_{\varepsilon}|^{2\alpha(p-1)}}{|Du_{\varepsilon}|^{4} |Du_{\varepsilon}|^{2\alpha(p-1)}} < 2^{\alpha(p-1)+2},$$

where we have used that $|Du_{\varepsilon}| > \lambda$. Thus we have

$$\mathbf{A}_{2} \leq 2^{\alpha(p-1)+4} \alpha^{2} \frac{|D^{2}u_{\varepsilon}|^{2} (|Du_{\varepsilon}| - \lambda)_{+}^{p-1+2\alpha}}{|Du_{\varepsilon}| [\lambda^{2} + (|Du_{\varepsilon}| - \lambda)_{+}^{2}]^{\alpha}}.$$

$$(4.40)$$

Joining estimates (4.37)-(4.40), for almost every $t \in (t_0 - \rho^2, t_1)$ we find

$$\begin{split} \left[\left(D_{x_j} u_{\varepsilon}(\cdot, t) \right) \eta^2 \chi(t) \,\Phi(P(\cdot, t)) \right]_{B^{2/p}_{p,\infty}(\mathbb{R}^n)}^p &\leq c_7 \int_{B_{\rho}} \frac{\left(\left| Du_{\varepsilon}(x, t) \right| - \lambda \right)_+^{p-1+2\alpha} \left| D^2 u_{\varepsilon}(x, t) \right|^2}{\left| Du_{\varepsilon}(x, t) \right| \left[\lambda^2 + \left(\left| Du_{\varepsilon}(x, t) \right| - \lambda \right)_+^2 \right]^{\alpha}} \,\chi(t) \,\eta^2 \, dx \\ &+ \frac{c_7}{\rho^2} \,\chi(t) \int_{B_{\rho}} \left| Du_{\varepsilon}(x, t) \right|^p \, dx, \end{split}$$

where $c_7 \equiv c_7(n, p, \alpha) > 0$. Inserting the previous inequality and (4.36) into (4.35), after some algebraic manipulation, from (4.34) we get

$$\begin{split} \int_{Q^{t_1}} (D_{x_j} f_{\varepsilon})(D_{x_j} u_{\varepsilon}) \eta^2(x) \chi(t) \Phi(P) dz \\ &\leq c_7 \int_{t_0 - \rho^2}^{t_1} \|f_{\varepsilon}(\cdot, t)\|_{B^{\frac{p-2}{p}}_{p', 1}(B_{\rho})} \left[\left(\int_{B_{\rho}} \frac{(|Du_{\varepsilon}(x, t)| - \lambda)^{p-1+2\alpha}_+ |D^2u_{\varepsilon}(x, t)|^2}{|Du_{\varepsilon}(x, t)| [\lambda^2 + (|Du_{\varepsilon}(x, t)| - \lambda)^2_+]^{\alpha}} \chi(t) \eta^2 dx \right)^{\frac{1}{p}} \\ &\quad + \frac{1}{\rho^{2/p}} \|Du_{\varepsilon}(\cdot, t)\|_{L^p(B_{\rho})} \right] dt. \end{split}$$

Now we go back to (4.33) and use the above estimate in combination with Young's inequality. This yields

$$\int_{Q^{t_1}} \frac{\left(|Du_{\varepsilon}| - \lambda\right)_+^{p-1+2\alpha}}{|Du_{\varepsilon}| \left[\lambda^2 + \left(|Du_{\varepsilon}| - \lambda\right)_+^2\right]^{\alpha}} |D^2 u_{\varepsilon}|^2 \chi(t) \eta^2(x) dz$$

$$\leq \frac{c_8}{\rho^2} \left[\int_{Q_R} (1 + |Du_{\varepsilon}|^2)^{\frac{p}{2}} dz + \|Du_{\varepsilon}\|_{L^p(Q_R)}^2 \right] + c_8 \int_{t_0 - \rho^2}^{t_0} \|f_{\varepsilon}(\cdot, t)\|_{B_{p', 1}^{\frac{p-2}{p}}(B_{\rho})}^{p'} dt$$

for every $t_1 \in (t_0 - \rho^2, t_0)$ and a positive constant c_8 depending only on n, p, α and R. Using the properties of χ and η , from the above inequality we deduce

$$\int_{Q_{\rho/2}(z_0)} \frac{(|Du_{\varepsilon}| - \lambda)_+^{p-1+2\alpha}}{|Du_{\varepsilon}| [\lambda^2 + (|Du_{\varepsilon}| - \lambda)_+^2]^{\alpha}} |D^2 u_{\varepsilon}|^2 dz
\leq \frac{c_8}{\rho^2} \left[\int_{Q_R} (1 + |Du_{\varepsilon}|^2)^{\frac{p}{2}} dz + \|Du_{\varepsilon}\|_{L^p(Q_R)}^2 \right] + c_8 \int_{t_0 - \rho^2}^{t_0} \|f_{\varepsilon}(\cdot, t)\|_{B_{p',1}^{\frac{p-2}{p}}(B_{\rho})}^{p'} dt. \quad (4.41)$$

At this point, recalling the definition of $\mathcal{V}_{\alpha,\lambda}$ in (1.2)–(1.3), a straightforward computation reveals that, for every $j \in \{1, \ldots, n\}$, we have

$$D_{x_{j}}\mathcal{V}_{\alpha,\lambda}(Du_{\varepsilon}) = \frac{(|Du_{\varepsilon}| - \lambda)_{+}^{\frac{p-1+2\alpha}{2}}}{|Du_{\varepsilon}|^{2} [\lambda + (|Du_{\varepsilon}| - \lambda)_{+}]^{\frac{1+2\alpha}{2}}} \langle Du_{\varepsilon}, D_{x_{j}}Du_{\varepsilon} \rangle Du_{\varepsilon} + \mathcal{G}_{\alpha,\lambda}((|Du_{\varepsilon}| - \lambda)_{+}) \left[\frac{D_{x_{j}}Du_{\varepsilon}}{|Du_{\varepsilon}|} - \frac{\langle Du_{\varepsilon}, D_{x_{j}}Du_{\varepsilon} \rangle}{|Du_{\varepsilon}|^{3}} Du_{\varepsilon} \right]$$

if $|Du_{\varepsilon}| > \lambda$, and $D_{x_j} \mathcal{V}_{\alpha,\lambda}(Du_{\varepsilon}) = 0$ otherwise. In the set $\{|Du_{\varepsilon}| > 0\}$, this yields

$$|D_x \mathcal{V}_{\alpha,\lambda}(Du_{\varepsilon})|^2 \le \mathbf{B}_1 + \mathbf{B}_2, \qquad (4.42)$$

where we define

$$\mathbf{B}_1 := 2 \frac{(|Du_{\varepsilon}| - \lambda)_+^{p-1+2\alpha} |D^2 u_{\varepsilon}|^2}{[\lambda + (|Du_{\varepsilon}| - \lambda)_+]^{1+2\alpha}}$$

and

$$\mathbf{B}_2 := 8 \frac{\left[\mathcal{G}_{\alpha,\lambda}((|Du_{\varepsilon}| - \lambda)_+)\right]^2 |D^2 u_{\varepsilon}|^2}{|Du_{\varepsilon}|^2}$$

We now estimate \mathbf{B}_1 and \mathbf{B}_2 separately in the set where $|Du_{\varepsilon}| > \lambda$, since both \mathbf{B}_1 and \mathbf{B}_2 vanish for $0 < |Du_{\varepsilon}| \le \lambda$. We immediately have

$$\mathbf{B}_{1} \leq 2 \frac{(|Du_{\varepsilon}| - \lambda)_{+}^{p-1+2\alpha} |D^{2}u_{\varepsilon}|^{2}}{|Du_{\varepsilon}| [\lambda^{2} + (|Du_{\varepsilon}| - \lambda)_{+}^{2}]^{\alpha}}.$$
(4.43)

As for \mathbf{B}_2 , by Lemma 2.5 we obtain

$$\mathbf{B}_{2} \leq \frac{32}{p^{2}} \frac{\left(|Du_{\varepsilon}| - \lambda\right)_{+}^{p+1+2\alpha} |D^{2}u_{\varepsilon}|^{2}}{|Du_{\varepsilon}|^{2} [\lambda + \left(|Du_{\varepsilon}| - \lambda\right)_{+}]^{1+2\alpha}}$$

$$\leq \frac{32}{p^{2}} \frac{\left(|Du_{\varepsilon}| - \lambda\right)_{+}^{p-1+2\alpha} (|Du_{\varepsilon}| - \lambda\right)_{+}^{2} |D^{2}u_{\varepsilon}|^{2}}{|Du_{\varepsilon}|^{3} [\lambda^{2} + \left(|Du_{\varepsilon}| - \lambda\right)_{+}^{2}]^{\alpha}}$$

$$\leq \frac{32}{p^{2}} \frac{\left(|Du_{\varepsilon}| - \lambda\right)_{+}^{p-1+2\alpha} |D^{2}u_{\varepsilon}|^{2}}{|Du_{\varepsilon}| [\lambda^{2} + \left(|Du_{\varepsilon}| - \lambda\right)_{+}^{2}]^{\alpha}}.$$
(4.44)

Joining estimates (4.42)-(4.44), we then find

$$\int_{Q_{\rho/2}(z_0)} |D_x \mathcal{V}_{\alpha,\lambda}(Du_{\varepsilon})|^2 dz \leq c(p) \int_{Q_{\rho/2}(z_0)} \frac{(|Du_{\varepsilon}| - \lambda)_+^{p-1+2\alpha} |D^2 u_{\varepsilon}|^2}{|Du_{\varepsilon}| [\lambda^2 + (|Du_{\varepsilon}| - \lambda)_+^2]^{\alpha}} dz, \tag{4.45}$$

which combined with (4.41), gives

$$\int_{Q_{\rho/2}(z_0)} |D_x \mathcal{V}_{\alpha,\lambda}(Du_{\varepsilon})|^2 dz$$

$$\leq \frac{c_8}{\rho^2} \left[\int_{Q_R} (1+|Du_{\varepsilon}|^2)^{\frac{p}{2}} dz + \|Du_{\varepsilon}\|_{L^p(Q_R)}^2 \right] + c_8 \int_{t_0-\rho^2}^{t_0} \|f_{\varepsilon}(\cdot,t)\|_{B_{p',1}^{\frac{p-2}{p}}(B_{\rho})}^{p'} dt$$

Let us now consider the same $\varepsilon_0 \in (0, 1]$ as in Proposition 4.4 and let $\varepsilon \in (0, \varepsilon_0]$. Then, applying estimate (4.5), we obtain

$$\int_{Q_{\rho/2}(z_0)} |D_x \mathcal{V}_{\alpha,\lambda}(Du_{\varepsilon})|^2 dz
\leq \frac{C}{\rho^2} \left(\|Du\|_{L^p(Q_R)}^p + \|Du\|_{L^p(Q_R)}^2 + \lambda^p + \lambda^2 + 1 \right) + C \|f_{\varepsilon}\|_{L^{p'}\left(t_0 - \rho^2, t_0; B_{p', 1}^{\frac{p-2}{p}}(B_{\rho})\right)}$$

for a constant $C \equiv C(n, p, \alpha, R) > 0$. Moreover, arguing as in the proof of [5, Theorem 1.1], we find a positive number $\varepsilon_1 \leq \varepsilon_0$ such that

$$\begin{aligned} \|f_{\varepsilon}\|_{L^{p'}\left(t_{0}-\rho^{2},t_{0};B_{p',1}^{\frac{p-2}{p}}(B_{\rho})\right)} &= \int_{t_{0}-\rho^{2}}^{t_{0}} \|f_{\varepsilon}(\cdot,t)\|_{B_{p',1}^{\frac{p-2}{p}}(B_{\rho})}^{p'}dt \leq \int_{t_{0}-R^{2}}^{t_{0}} \|f(\cdot,t)\|_{B_{p',1}^{\frac{p-2}{p}}(B_{R})}^{p'}dt \\ &= \|f\|_{L^{p'}\left(t_{0}-R^{2},t_{0};B_{p',1}^{\frac{p-2}{p}}(B_{R})\right)}^{p'} < +\infty, \quad \text{for every } \varepsilon \in (0,\varepsilon_{1}].\end{aligned}$$

Combining the last two estimates for $\varepsilon \in (0, \varepsilon_1]$, we conclude the proof in the case $\lambda > 0$.

Finally, when $\lambda = 0$ the above proof can be greatly simplified, since the parameter α plays no role and

$$\mathcal{V}_0(Du_{\varepsilon}) = \frac{2}{p} H_{\frac{p}{2}}(Du_{\varepsilon}) = \frac{2}{p} |Du_{\varepsilon}|^{\frac{p-2}{2}} Du_{\varepsilon}.$$

In this regard, we leave the details to the reader.

We conclude this section with the following result, which will play a crucial role in the proof of Theorem 1.3.

Proposition 4.7 (Uniform Sobolev estimate). Under the assumptions of Theorem 1.3 and with the notation above, there exists a positive number $\varepsilon_0 \leq 1$ such that, for every $\varepsilon \in (0, \varepsilon_0]$, for every parabolic cylinder $Q_{\rho}(z_0) \in Q_R(z_0)$ and every $j \in \{1, \ldots, n\}$, we have

$$\int_{Q_{\rho/2}(z_0)} |D_{x_j} \mathcal{V}_{\alpha,\lambda}(Du_{\varepsilon})|^2 dz \le \frac{C}{\rho^2} \left(\|Du\|_{L^2(Q_R)}^2 + \lambda^2 + 1 \right) + C \left(\|f\|_{L^2(Q_R)}^2 + 1 \right)$$
(4.46)

for a positive constant C depending only on n and R in the case $\lambda = 0$, and additionally on α if $\lambda > 0$.

Proof. In what follows, we shall keep the notation, definitions and choices used in the proof of Proposition 4.6.

Let us first assume that $\lambda > 0$. Letting $\rho \to 0$ in (4.20) and arguing exactly as in the preceding proof, we arrive at the following inequality

$$I_3 \le I_4 + I_5 + I_6 + I_7 + I_8, \qquad (4.47)$$

where

$$\begin{split} I_3 &:= \int_{Q^{t_1}} \sum_{j=1}^n \left\langle D^2 A_{\varepsilon}(Du_{\varepsilon}) D(D_{x_j}u_{\varepsilon}), D(D_{x_j}u_{\varepsilon}) \right\rangle \chi(t) \eta^2(x) \Phi(P) \, dz, \\ I_4 &:= 4 \int_{Q^{t_1}} \sum_{j=1}^n \left\langle D^2 A_{\varepsilon}(Du_{\varepsilon}) D\eta, D\eta \right\rangle \chi(t) \left| D_{x_j}u_{\varepsilon} \right|^2 \Phi(P) \, dz, \\ I_5 &:= \int_{Q^{t_1}} (\partial_t \chi) \eta^2(x) \left[\int_0^{|Du_{\varepsilon}(x,t)|^2} \Phi((\sqrt{w} - \lambda)_+) \, dw \right] \, dz, \\ \tilde{I}_6 &:= -4 \int_{Q^{t_1}} \sum_{j=1}^n f_{\varepsilon} \left(D_{x_j}u_{\varepsilon} \right) \eta(x) \left(D_{x_j}\eta \right) \chi(t) \Phi(P) \, dz, \\ I_7 &:= -2 \int_{Q^{t_1}} \sum_{j=1}^n f_{\varepsilon} \left(D_{x_j}u_{\varepsilon} \right) \eta^2(x) \chi(t) \Phi(P) \, dz, \\ I_8 &:= -2 \int_{Q^{t_1}} \sum_{j=1}^n f_{\varepsilon} \left(D_{x_j}u_{\varepsilon} \right) \eta^2(x) \chi(t) \Phi'(P) \left(D_{x_j}P \right) \, dz, \end{split}$$

while P, η , χ and Φ are chosen as in the previous proof. Note that we have already estimated I_3 , I_4 and I_5 in (4.30), (4.31) and (4.32) respectively. More precisely, for p = 2 these estimates yield

$$I_3 \geq \int_{Q^{t_1}} \frac{(|Du_{\varepsilon}| - \lambda)_+^{1+2\alpha}}{|Du_{\varepsilon}| [\lambda^2 + (|Du_{\varepsilon}| - \lambda)_+^2]^{\alpha}} |D^2 u_{\varepsilon}|^2 \chi(t) \eta^2(x) dz, \qquad (4.48)$$

$$I_4 \le \frac{8\,\tilde{c}^2}{\rho^2} \int_{Q_R} (1+|Du_{\varepsilon}|^2)\,dz \tag{4.49}$$

and

$$I_5 \leq \frac{4}{3\rho^2} \|Du_{\varepsilon}\|_{L^2(Q_R)}^2.$$
(4.50)

We now estimate \tilde{I}_6 , I_7 and I_8 separately. Let us first consider \tilde{I}_6 . Using the fact that $0 \leq \chi, \Phi \leq 1$, the properties (4.27) of η and Young's inequality, we obtain

$$\tilde{I}_{6} \leq 4n \int_{Q_{R}} |f_{\varepsilon}| |Du_{\varepsilon}| |D\eta| dz \leq 2n \|f_{\varepsilon}\|_{L^{2}(Q_{R})}^{2} + \frac{2n \,\tilde{c}^{2}}{\rho^{2}} \|Du_{\varepsilon}\|_{L^{2}(Q_{R})}^{2}.$$
(4.51)

As for the term I_7 , recalling (4.17) and (4.28), we have

$$I_{7} \leq 2n \int_{Q^{t_{1}} \cap \{|Du_{\varepsilon}| > \lambda\}} |f_{\varepsilon}| |D^{2}u_{\varepsilon}| \eta^{2}(x) \chi(t) \frac{(|Du_{\varepsilon}| - \lambda)^{2\alpha}_{+}}{[\lambda^{2} + (|Du_{\varepsilon}| - \lambda)^{2}_{+}]^{\alpha}} dz$$

$$\leq 2n \int_{Q^{t_{1}} \cap \{|Du_{\varepsilon}| > \lambda\}} |f_{\varepsilon}| |D^{2}u_{\varepsilon}| \eta^{2}(x) \chi(t) \frac{(|Du_{\varepsilon}| - \lambda)^{2\alpha-1}_{+} |Du_{\varepsilon}|}{[\lambda^{2} + (|Du_{\varepsilon}| - \lambda)^{2}_{+}]^{\alpha}} dz.$$

Now we turn our attention to I_8 . Recalling (4.29) and applying Young's inequality, we get

$$I_{8} \leq 4\alpha \int_{Q^{t_{1}} \cap \{|Du_{\varepsilon}| > \lambda\}} \sum_{j=1}^{n} |f_{\varepsilon}| |D_{x_{j}}u_{\varepsilon}| \eta^{2}(x) \chi(t) \frac{\lambda^{2} (|Du_{\varepsilon}| - \lambda)_{+}^{2\alpha-1}}{[\lambda^{2} + (|Du_{\varepsilon}| - \lambda)_{+}^{2}]^{\alpha+1}} \cdot \frac{|\langle Du_{\varepsilon}, D_{x_{j}}Du_{\varepsilon}\rangle|}{|Du_{\varepsilon}|} dz$$
$$\leq 8\alpha n \int_{Q^{t_{1}} \cap \{|Du_{\varepsilon}| > \lambda\}} |f_{\varepsilon}| |D^{2}u_{\varepsilon}| \eta^{2}(x) \chi(t) \frac{|Du_{\varepsilon}|^{3} (|Du_{\varepsilon}| - \lambda)_{+}^{2\alpha-1}}{[\lambda^{2} + (|Du_{\varepsilon}| - \lambda)_{+}^{2}]^{\alpha} |Du_{\varepsilon}|^{2}} dz$$

$$= 8\alpha n \int_{Q^{t_1} \cap \{|Du_{\varepsilon}| > \lambda\}} |f_{\varepsilon}| |D^2 u_{\varepsilon}| \eta^2(x) \chi(t) \frac{(|Du_{\varepsilon}| - \lambda)^{2\alpha - 1}_+ |Du_{\varepsilon}|}{[\lambda^2 + (|Du_{\varepsilon}| - \lambda)^2_+]^{\alpha}} dz$$

Joining the last two estimates and using Young's inequality again, we find $I_7 + I_8$

$$\leq 2n (4\alpha + 1) \int_{Q^{t_{1}} \cap \{|Du_{\varepsilon}| > \lambda\}} \frac{|f_{\varepsilon}| |Du_{\varepsilon}|^{\frac{3}{2}} (|Du_{\varepsilon}| - \lambda)^{\alpha - \frac{3}{2}}}{[\lambda^{2} + (|Du_{\varepsilon}| - \lambda)^{2}]^{\frac{\alpha}{2}}} \cdot \frac{(|Du_{\varepsilon}| - \lambda)^{\alpha + \frac{1}{2}} |D^{2}u_{\varepsilon}| \eta^{2}(x) \chi(t)}{\sqrt{|Du_{\varepsilon}|} [\lambda^{2} + (|Du_{\varepsilon}| - \lambda)^{2}]^{\frac{\alpha}{2}}} dz$$

$$\leq 2n^{2} (4\alpha + 1)^{2} \int_{Q^{t_{1}} \cap \{|Du_{\varepsilon}| > \lambda\}} \frac{|f_{\varepsilon}|^{2} |Du_{\varepsilon}|^{3} (|Du_{\varepsilon}| - \lambda)^{2\alpha - 3} \eta^{2}(x) \chi(t)}{[\lambda^{2} + (|Du_{\varepsilon}| - \lambda)^{2}]^{\alpha}} dz$$

$$+ \frac{1}{2} \int_{Q^{t_{1}}} \frac{(|Du_{\varepsilon}| - \lambda)^{1 + 2\alpha} |D^{2}u_{\varepsilon}|^{2}}{|Du_{\varepsilon}| - \lambda)^{2} |\alpha} \eta^{2}(x) \chi(t) dz$$

$$\leq 4n^{2} (4\alpha + 1)^{2} \int_{Q^{t_{1}} \cap \{|Du_{\varepsilon}| > \lambda\}} \frac{|f_{\varepsilon}|^{2} |Du_{\varepsilon}|^{3} (|Du_{\varepsilon}| - \lambda)^{2\alpha - 3}}{|Du_{\varepsilon}|^{2\alpha}} dz$$

$$+ \frac{1}{2} \int_{Q^{t_{1}}} \frac{(|Du_{\varepsilon}| - \lambda)^{1 + 2\alpha} |D^{2}u_{\varepsilon}|^{2}}{|D^{2}u_{\varepsilon}|^{2}} \eta^{2}(x) \chi(t) dz$$

$$\leq 4n^{2} (4\alpha + 1)^{2} \|f_{\varepsilon}\|^{2} |\lambda^{2} + (|Du_{\varepsilon}| - \lambda)^{2} |\alpha^{2}} \eta^{2}(x) \chi(t) dz$$

$$\leq 4n^{2} (4\alpha + 1)^{2} \|f_{\varepsilon}\|^{2}_{L^{2}(Q_{R})} + \frac{1}{2} \int_{Q^{t_{1}}} \frac{(|Du_{\varepsilon}| - \lambda)^{1 + 2\alpha} |D^{2}u_{\varepsilon}|^{2}}{|Du_{\varepsilon}| |\lambda^{2} + (|Du_{\varepsilon}| - \lambda)^{2} |\alpha^{2}} \eta^{2}(x) \chi(t) dz,$$

$$\leq 4n^{2} (4\alpha + 1)^{2} \|f_{\varepsilon}\|^{2}_{L^{2}(Q_{R})} + \frac{1}{2} \int_{Q^{t_{1}}} \frac{(|Du_{\varepsilon}| - \lambda)^{1 + 2\alpha} |D^{2}u_{\varepsilon}|^{2}}{|Du_{\varepsilon}| |\lambda^{2} + (|Du_{\varepsilon}| - \lambda)^{2} |\alpha^{2}} \eta^{2}(x) \chi(t) dz,$$

$$\leq 4n^{2} (4\alpha + 1)^{2} \|f_{\varepsilon}\|^{2}_{L^{2}(Q_{R})} + \frac{1}{2} \int_{Q^{t_{1}}} \frac{(|Du_{\varepsilon}| - \lambda)^{1 + 2\alpha} |D^{2}u_{\varepsilon}|^{2} |\alpha^{2}}{|Du_{\varepsilon}| |\lambda^{2} + (|Du_{\varepsilon}| - \lambda)^{2} |\alpha^{2}} \eta^{2}(x) \chi(t) dz,$$

$$\leq 4n^{2} (4\alpha + 1)^{2} \|f_{\varepsilon}\|^{2}_{L^{2}(Q_{R})} + \frac{1}{2} \int_{Q^{t_{1}}} \frac{(|Du_{\varepsilon}| - \lambda)^{1 + 2\alpha} |D^{2}u_{\varepsilon}|^{2} |\alpha^{2}}{|Du_{\varepsilon}| |\lambda^{2} + (|Du_{\varepsilon}| - \lambda)^{2} |\alpha^{2}} \eta^{2}(x) \chi(t) dz,$$

$$\leq 4n^{2} (4\alpha + 1)^{2} \|f_{\varepsilon}\|^{2}_{L^{2}(Q_{R})} + \frac{1}{2} \int_{Q^{t_{1}}} \frac{(|Du_{\varepsilon}| - \lambda)^{1 + 2\alpha} |D^{2}u_{\varepsilon}|^{2} |\alpha^{2}}{|Du_{\varepsilon}| |\lambda^{2} + (|Du_{\varepsilon}| - \lambda)^{2} |\alpha^{2}} \eta^{2}(x) \chi(t) dz,$$

where, in the last line, we have used the fact that $|Du_{\varepsilon}|^3 (|Du_{\varepsilon}| - \lambda)^{2\alpha-3}_+ \leq |Du_{\varepsilon}|^{2\alpha}$ in the set $\{|Du_{\varepsilon}| > \lambda\}$, since $\alpha \geq \frac{3}{2}$ by assumption. Now we combine estimates (4.47)–(4.52), thus obtaining

$$\int_{Q^{t_1}} \frac{(|Du_{\varepsilon}| - \lambda)_{+}^{1+2\alpha} |D^2 u_{\varepsilon}|^2}{|Du_{\varepsilon}| [\lambda^2 + (|Du_{\varepsilon}| - \lambda)_{+}^2]^{\alpha}} \eta^2(x) \chi(t) dz \leq \frac{c_1}{\rho^2} \int_{Q_R} (1 + |Du_{\varepsilon}|^2) dz + c_1 ||f_{\varepsilon}||^2_{L^2(Q_R)},$$

which holds for every $t_1 \in (t_0 - \rho^2, t_0)$ and a positive constant c_1 depending only on n and α . Using the properties of η and χ , from the above inequality we deduce

$$\int_{Q_{\rho/2}(z_0)} \frac{(|Du_{\varepsilon}| - \lambda)_{+}^{1+2\alpha} |D^2 u_{\varepsilon}|^2}{|Du_{\varepsilon}| [\lambda^2 + (|Du_{\varepsilon}| - \lambda)_{+}^2]^{\alpha}} dz \le \frac{c_1}{\rho^2} \int_{Q_R} (1 + |Du_{\varepsilon}|^2) dz + c_1 ||f_{\varepsilon}||_{L^2(Q_R)}^2.$$
(4.53)

Taking p = 2 into (4.45) and combining the resulting estimate with (4.53), we then find

$$\int_{Q_{\rho/2}(z_0)} |D_x \mathcal{V}_{\alpha,\lambda}(Du_{\varepsilon})|^2 dz \le \frac{c_1}{\rho^2} \int_{Q_R} (1 + |Du_{\varepsilon}|^2) dz + c_1 \|f_{\varepsilon}\|_{L^2(Q_R)}^2.$$

Let us now consider the same $\varepsilon_0 \in (0, 1]$ as in Proposition 4.4 and let $\varepsilon \in (0, \varepsilon_0]$. Then, applying estimate (4.5) with p = 2, we get

$$\int_{Q_{\rho/2}(z_0)} |D_x \mathcal{V}_{\alpha,\lambda}(Du_{\varepsilon})|^2 dz \le \frac{C}{\rho^2} \left(\|Du\|_{L^2(Q_R)}^2 + \lambda^2 + 1 \right) + C \|f_{\varepsilon}\|_{L^2(Q_R)}^2$$

for a constant $C \equiv C(n, \alpha, R) > 0$. Moreover, for p = 2, from (4.14) it follows that

$$||f_{\varepsilon}||_{L^2(Q_R)} \leq ||f||_{L^2(Q_R)} + 1.$$

Combining the last two inequalities, we conclude the proof in the case $\lambda > 0$.

Finally, when $\lambda = 0$ the above proof can be greatly simplified, since the parameter α plays no role and

$$\mathcal{V}_0(Du_\varepsilon) = Du_\varepsilon.$$

We leave the details to the reader.

5 Proofs of Theorems 1.1 and 1.3

In this section, we prove Theorems 1.1 and 1.3 by combining a standard comparison argument (see e.g. [6, 13, 15]) with the estimates from Propositions 4.4, 4.5, 4.6 and 4.7.

Proof of Theorem 1.1. In what follows, we will refer to the notation introduced in Section 4. For any fixed $\varepsilon \in (0, 1]$, we define the comparison map

$$u_{\varepsilon} \in C^0\left([t_0 - R^2, t_0]; L^2(B_R(x_0))\right) \cap L^p\left(t_0 - R^2, t_0; W^{1,p}(B_R(x_0))\right)$$

as the unique energy solution of the Cauchy-Dirichlet problem (4.2). Moreover, by a slight abuse of notation, for $w \in L^1_{loc}(Q_R(z_0), \mathbb{R}^k)$, $j \in \{1, \ldots, n\}$ and $h \neq 0$, we set (when $x + he_j \in Q_R(z_0)$)

$$\tau_{j,h}w(x,t) := w(x + he_j, t) - w(x,t), \Delta_{j,h}w(x,t) := \frac{w(x + he_j, t) - w(x,t)}{h},$$

where e_j is the unit vector in the direction x_j . Now we fix arbitrary radii

$$0 < r < \rho < R$$

and use the finite difference operator $\tau_{j,h}$ defined above, for increments $h \in \mathbb{R} \setminus \{0\}$ such that $|h| < \frac{\rho - r}{4}$.

Let us first assume that $\lambda > 0$. Furthermore, consider the same $\varepsilon_1 \in (0, 1]$ as in Proposition 4.6 and let $\varepsilon \in (0, \varepsilon_1]$. In the following, we will denote by C a positive constant which neither depends on h nor on ε . In order to obtain an estimate for the finite difference $\tau_{j,h} \mathcal{V}_{\alpha,\lambda}(Du)$, we use the following comparison argument:

$$\begin{split} &\int_{Q_{r/2}(z_0)} |\tau_{j,h} \mathcal{V}_{\alpha,\lambda}(Du)|^2 dz \\ &\leq 4 \int_{Q_{r/2}(z_0)} |\tau_{j,h} \mathcal{V}_{\alpha,\lambda}(Du_{\varepsilon})|^2 dx \, dt + 4 \int_{Q_{r/2}(z_0)} |\mathcal{V}_{\alpha,\lambda}(Du_{\varepsilon}) - \mathcal{V}_{\alpha,\lambda}(Du)|^2 dx \, dt \\ &+ 4 \int_{Q_{r/2}(z_0)} |\mathcal{V}_{\alpha,\lambda}(Du_{\varepsilon}(x+he_j,t)) - \mathcal{V}_{\alpha,\lambda}(Du(x+he_j,t))|^2 dx \, dt \\ &\leq 4 \int_{Q_{r/2}(z_0)} |\tau_{j,h} \mathcal{V}_{\alpha,\lambda}(Du_{\varepsilon})|^2 dz + 8 \int_{Q_R(z_0)} |\mathcal{V}_{\alpha,\lambda}(Du_{\varepsilon}) - \mathcal{V}_{\alpha,\lambda}(Du)|^2 dz. \end{split}$$

Combining the previous estimate with (4.15) and (4.16), for every $j \in \{1, \ldots, n\}$ we get

$$\int_{Q_{r/2}} |\tau_{j,h} \mathcal{V}_{\alpha,\lambda}(Du)|^2 dz
\leq \frac{C}{\rho^2} |h|^2 \left(\|Du\|_{L^p(Q_R)}^p + \|Du\|_{L^p(Q_R)}^2 + \lambda^p + \lambda^2 + 1 \right) + C |h|^2 \|f\|_{L^{p'}\left(t_0 - R^2, t_0; B_{p', 1}^{\frac{p-2}{p}}(B_R(x_0))\right)}
+ C \varepsilon \left(\|Du\|_{L^p(Q_R(z_0))}^p + 1 \right) + C \|f - f_\varepsilon\|_{L^{p'}(Q_R(z_0))} \left(\|Du\|_{L^p(Q_R(z_0))} + \lambda + 1 \right), \quad (5.1)$$

which holds for every sufficiently small $h \in \mathbb{R} \setminus \{0\}$ and a constant $C \equiv C(n, p, \alpha, R) > 0$. Therefore, recalling (4.13) and letting $\varepsilon \searrow 0$ in (5.1), we obtain

$$\int_{Q_{r/2}} |\Delta_{j,h} \mathcal{V}_{\alpha,\lambda}(Du)|^2 dz \leq \frac{C}{\rho^2} \left(\|Du\|_{L^p(Q_R)}^p + \|Du\|_{L^p(Q_R)}^2 + \lambda^p + \lambda^2 + 1 \right) + C \|f\|_{L^{p'}\left(t_0 - R^2, t_0; B_{p',1}^{\frac{p-2}{p}}(B_R(x_0))\right)}.$$
(5.2)

Since the above estimate holds for every $j \in \{1, ..., n\}$ and every suitably small $h \neq 0$, by Lemma 2.10 we may conclude that

$$\mathcal{V}_{\alpha,\lambda}(Du) \in L^2_{loc}\left(0,T; W^{1,2}_{loc}(\Omega,\mathbb{R}^n)\right).$$

Moreover, letting $h \to 0$ in the previous inequality, we also obtain estimate (1.5).

Let us now assume that $\lambda = 0$. In this case, we have already observed that the constant C is independent of α (see Proposition 4.6) and

$$\mathcal{V}_0(Du) = \frac{2}{p} \left| Du \right|^{\frac{p-2}{2}} Du \,.$$

Combining this fact with Lemma 2.3 and estimate (5.2), for every sufficiently small $h \neq 0$ we find

$$\int_{Q_{r/2}} \frac{|\tau_{j,h} Du(x,t)|^p}{|h|^2} dx \, dt \leq C_1(p) \int_{Q_{r/2}} |\Delta_{j,h} \mathcal{V}_0(Du)|^2 \, dz \leq M$$

for some finite positive constant M depending on

$$n, p, R, \rho, ||Du||_{L^p(Q_R)}, ||f||_{L^{p'}\left(t_0-R^2, t_0; B_{p',1}^{\frac{p-2}{p}}(B_R)\right)},$$

but not on h. Note that the above estimate also holds for every $j \in \{1, \ldots, n\}$. Therefore, using Proposition 3.11 with the choices G = Du, q = p and $\theta = \frac{2}{p}$, as well as a standard covering argument, we infer that

$$Du \in L_{loc}^{p}(0,T; W_{loc}^{\sigma,p}(\Omega,\mathbb{R}^{n}))$$
 for all $\sigma \in \left(0,\frac{2}{p}\right)$.

This completes the proof.

We now conclude this section by giving the

Proof of Theorem 1.3. Actually, it is sufficient to proceed as in the proof of Theorem 1.1, but using estimate (4.46) in place of (4.16). In particular, since $\mathcal{V}_0(Du) = Du$, in the case $\lambda = 0$ we retrieve

$$u \in L^{2}_{loc}(0, T; W^{2,2}_{loc}(\Omega)),$$

which is a well-established result for the non-homogeneous heat equation (1.7) when $f \in L^2_{loc}(\Omega_T)$.

6 The time derivative: proof of Theorem 1.5

This section is devoted to the study of the existence and regularity of the time derivative of the weak solutions to equation (1.8), under the assumptions of Theorem 1.5. Indeed, we are now in a position to prove the aforementioned theorem.

Proof of Theorem 1.5. We shall keep the notation and the parabolic cylinders used for the proof of Theorem 1.1. Let us notice that

$$|Du|^{p-2}Du = \mathcal{H}(\mathcal{V}_0(Du)),$$

where $\mathcal{H}: \mathbb{R}^n \to \mathbb{R}^n$ is the function defined by

$$\mathcal{H}(\xi) := \left(\frac{p}{2}\right)^{\frac{2}{p'}} |\xi|^{\frac{p-2}{p}} \xi,$$

which is locally Lipschitz continuous for p > 2. Thus, the vector field $|Du|^{p-2}Du$ is weakly differentiable with respect to the *x*-variable by virtue of the chain rule in Sobolev spaces. From the definitions of \mathcal{H} and \mathcal{V}_0 , it follows that

$$|D_{\xi} \mathcal{H}(\mathcal{V}_0(Du))| \leq c_1 |Du|^{\frac{p-2}{2}}$$

$$(6.1)$$

for some positive constant $c_1 \equiv c(n, p)$. Now, applying the chain rule, the Cauchy-Schwarz inequality and estimate (6.1), we obtain

$$|D_{x} \mathcal{H}(\mathcal{V}_{0}(Du))|^{p'} \leq c_{2}(n,p) |D_{\xi} \mathcal{H}(\mathcal{V}_{0}(Du))|^{p'} |D_{x} \mathcal{V}_{0}(Du)|^{p'} \\ \leq c_{3} |Du|^{\frac{(p-2)p'}{2}} |D_{x} \mathcal{V}_{0}(Du)|^{p'},$$
(6.2)

where $c_3 \equiv c_3(n,p) > 0$. Using (6.2), Hölder's inequality with exponents $\left(\frac{2(p-1)}{p-2}, \frac{2}{p'}\right)$ and estimate (1.5) with $\lambda = 0$, we get

$$\left(\int_{Q_{r/2}} \left|D_x \mathfrak{H}(\mathfrak{V}_0(Du))\right|^{p'} dz\right)^{\frac{1}{p'}} \leq c_4(n,p) \left\|Du\right\|_{L^p(Q_{r/2})}^{\frac{p-2}{2}} \left(\int_{Q_{r/2}} \left|D_x \mathfrak{V}_0(Du)\right|^2 dz\right)^{\frac{1}{2}} \\ \leq \frac{C}{\rho} \left(\left\|Du\right\|_{L^p(Q_R)}^{p-1} + \left\|Du\right\|_{L^p(Q_R)}^{\frac{p}{2}} + \left\|Du\right\|_{L^p(Q_R)}^{\frac{p-2}{2}} + C \left\|Du\right\|_{L^p(Q_R)}^{\frac{p-2}{2}} \left\|f\right\|_{L^{p'}\left(t_0 - R^2, t_0; B_{p', 1}^{\frac{p-2}{p}}(B_R)\right)}\right)$$

$$(6.3)$$

for a constant $C \equiv C(n, p, R) > 0$. Note that the right-hand side of (6.3) is finite, and this implies that

$$|Du|^{p-2}Du \in L^{p'}_{loc}(0,T;W^{1,p'}_{loc}(\Omega,\mathbb{R}^n)).$$
(6.4)

Therefore, going back to the weak formulation (2.3), we can now perform a partial integration in the second term on the left-hand side with respect to the spatial variables. We thus obtain

$$\int_{Q_{r/2}(z_0)} u \cdot \varphi_t \, dz = -\int_{Q_{r/2}(z_0)} \left(\sum_{j=1}^n D_{x_j} \left[(|Du|^{p-2} Du)_j \right] + f \right) \cdot \varphi \, dz$$

for any $\varphi \in C_0^{\infty}(Q_{r/2}(z_0))$, and the desired conclusion immediately follows from (6.4), since

$$f \in L^{p'}_{loc}\left(0, T; B^{\frac{p-2}{p}}_{p', 1, loc}(\Omega)\right).$$

Furthermore, we can now observe that

$$\partial_t u = \sum_{j=1}^n D_{x_j} \left[(|Du|^{p-2} Du)_j \right] + f \quad \text{in } Q_{r/2}(z_0).$$

From this we can infer

$$\left(\int_{Q_{r/2}(z_0)} \left|\partial_t u\right|^{p'} dz\right)^{\frac{1}{p'}} \le n \|D_x \mathcal{H}(\mathcal{V}_0(Du))\|_{L^{p'}(Q_{r/2})} + \|f\|_{L^{p'}(Q_{r/2})}$$

and combining this inequality with (6.3), we finally obtain estimate (1.9).

Acknowledgments. The author has been partially supported by the INdAM–GNAMPA 2024 Project "Fenomeno di Lavrentiev, Bounded Slope Condition e regolarità per minimi di funzionali integrali con crescite non standard e lagrangiane non uniformemente convesse" (CUP: E53C23001670001).

Data availability. There are no data associated with this manuscript.

Declarations. The author states that there is no conflict of interest.

References

- P. Ambrosio, Besov regularity for a class of singular or degenerate elliptic equations, J. Math. Anal. Appl. 505 (2022), no. 2, Paper No. 125636.
- [2] P. Ambrosio, Fractional Sobolev regularity for solutions to a strongly degenerate parabolic equation, Forum Math. (2023).
- P. Ambrosio, F. Bäuerlein, Gradient bounds for strongly singular or degenerate parabolic systems, J. Differ. Equ. 401 (2024), 492-549.
- [4] P. Ambrosio, S. Cuomo, M. De Rosa, A physics-informed deep learning approach for solving strongly degenerate parabolic problems, Engineering with Computers, (2024). DOI: https://doi.org/10.1007/s00366-024-01961-9.
- [5] P. Ambrosio, A.G. Grimaldi, A. Passarelli di Napoli, Sharp second-order regularity for widely degenerate elliptic equations, preprint (2024), version 2. Available at https://arxiv.org/abs/2401.13116v2.
- [6] P. Ambrosio, A. Passarelli di Napoli, *Regularity results for a class of widely degenerate parabolic equations*, Adv. Calc. Var. (2023).
- [7] V. Bögelein, F. Duzaar, R. Giova, A. Passarelli di Napoli, *Gradient regularity for a class of widely degenerate parabolic systems*, SIAM J. Math. Anal., 1-62 (2024). To appear.
- [8] V. Bögelein, F. Duzaar, R. Giova, A. Passarelli di Napoli, Higher regularity in congested traffic dynamics, Math. Ann. 385, 1-56 (2023).
- [9] L. Brasco, Global L[∞] gradient estimates for solutions to a certain degenerate elliptic equation, Nonlinear Anal., Theory Methods Appl. 74 (2) (2011) 516-531.
- [10] L. Brasco, G. Carlier, F. Santambrogio, Congested traffic dynamics, weak flows and very degenerate elliptic equations, [corrected version of mr2584740], J. Math. Pures Appl. (9) 93 (2010), no. 6, 652-671.
- [11] L. Brasco, F. Santambrogio, A sharp estimate à la Calderón-Zygmund for the p-Laplacian, Commun. Contemp. Math. 20 (2018), no. 3, Article ID 1750030.
- [12] F. Duzaar, A. Gastel, G. Mingione, *Elliptic systems, singular sets and Dini continuity*, Comm. Part. Diff. Equ., **29** (2004), 1215-1240.
- [13] F. Duzaar, G. Mingione, K. Steffen, Parabolic systems with polynomial growth and regularity, Mem. Amer. Math. Soc. 214 (2011).

- [14] L. Esposito, F. Leonetti, G. Mingione, Higher integrability for minimizers of integral functionals with (p,q) growth, J. Differ. Equ. 157 (1999), 414-438.
- [15] A. Gentile, A. Passarelli di Napoli, Higher regularity for weak solutions to degenerate parabolic problems, Calc. Var. 62, 225 (2023).
- [16] F. Giannetti, A. Passarelli di Napoli, C. Scheven, Higher differentiability of solutions of parabolic systems with discontinuous coefficients, J. Lond. Math. Soc. (2) 94 (2016), no. 1, 1-20.
- [17] E. Giusti, Direct Methods in the Calculus of Variations, World Scientific Publishing Co., 2003.
- [18] P. Lindqvist, On the time derivative in a quasilinear equation, Skr. K. Nor. Vidensk. Selsk. (2008), no. 2, 1-7.
- [19] P. Lindqvist, On the time derivative in an obstacle problem, Rev. Mat. Iberoam. 28 (2012), no. 2, 577-590.
- [20] P. Lindqvist, The time derivative in a singular parabolic equation, Differential Integral Equations 30 (2017), no. 9-10, 795-808.
- [21] P. Lindqvist, Notes on the Stationary p-Laplace Equation, Springer Briefs Math., Springer, Cham 2019.
- [22] J.-L. Lions, Quelques méthodes de résolution des problèmes aux limites non linéaires, Gauthier-Villars, Paris, 1969.
- [23] G. Mingione, *Gradient potential estimates*, J. European Math. Soc. **13** (2011), 459-486.
- [24] G. Mingione, The Calderón-Zygmund theory for elliptic problems with measure data, Ann. Scu. Norm. Sup. Pisa Cl. Sci. (V) 6 (2007), 195-261.
- [25] G. Mingione, The singular set of solutions to non-differentiable elliptic systems, Arch. Rat. Mech. Anal. 166 (2003), 287-301.
- [26] C. Scheven, Regularity for subquadratic parabolic systems: Higher integrability and dimension estimates, Proc. Roy. Soc. Edinburgh Sect. A 140 (2010), no. 6, 1269-1308.
- [27] H. Triebel, Theory of Function Spaces, Monogr. Math. 78, Birkhäuser, Basel, 1983.
- [28] H. Triebel, Theory of Function Spaces IV, Monogr. Math. 107, Birkhäuser, Basel, 2020.