

A Note on Finite Nilpotent Groups

Nicholas J. Werner

July 15, 2024

Abstract

It is well known that if G is a group and H is a normal subgroup of G of finite index k , then $x^k \in H$ for every $x \in G$. We examine finite groups G with the property that $x^k \in H$ for every subgroup H of G , where k is the index of H in G . We prove that a finite group G satisfies this property if and only if G is nilpotent.

Author's Note added July 11, 2024. I have learned that the results of this paper have already appeared in:

L. Sabatini. Products of subgroups, subnormality, and relative orders of elements. *Ars Math. Contemp.* 24, no. 1 (2024), 1–9. [DOI]

For a group G and a subgroup $H \leq G$, let $[G : H]$ denote the index of H in G . A common exercise in a first course on abstract algebra is to prove that if H is normal in G and $[G : H]$ is finite, then $x^{[G:H]} \in H$ for all $x \in G$. This result need not hold when H is not normal. For example, take G to be the symmetric group S_3 , let σ and τ be distinct transpositions in G , and let $H = \langle \sigma \rangle$. Then, $[G : H] = 3$, but $\tau^3 = \tau \notin H$.

One could then ask: for which groups do we always have $x^{[G:H]} \in H$? We will restrict our focus to finite groups, and we will say that a finite group G has property (*) if the following holds:

$$\text{for each } H \leq G \text{ and for all } x \in G, x^{[G:H]} \in H. \quad (*)$$

Certainly, (*) is true for finite Abelian groups, and more generally for any finite group with all subgroups normal (such as the quaternion group Q_8). However, the requirement that every subgroup of G be normal in G is not necessary for (*) to hold. For instance, the dihedral group of order 8 contains subgroups that are not normal, and it is straightforward to verify that this group satisfies (*).

In this note, we prove that a finite group satisfies (*) if and only if it is nilpotent. Typically, nilpotent groups are defined using upper central series. For this, one constructs the chain of normal subgroups

$$Z_0 \trianglelefteq Z_1 \trianglelefteq Z_2 \trianglelefteq \cdots$$

where Z_0 is the trivial subgroup, Z_1 is the center of G , and Z_{i+1} is chosen so that Z_{i+1}/Z_i equals the center of G/Z_i . The group G is said to be *nilpotent* if the

above chain terminates at G in a finite number of steps. When G is finite, there are a variety of other ways to characterize nilpotent groups. The proposition below summarizes a number of these conditions; a proof can be found in many algebra texts such as [1, Section 6.1].

Proposition 1. *Let G be a finite group. The following are equivalent.*

- (1) G is nilpotent.
- (2) Every proper subgroup of G is a proper subgroup of its normalizer in G .
- (3) Every Sylow subgroup of G is normal.
- (4) G is isomorphic to the direct product of all of its Sylow subgroups.
- (5) Every maximal subgroup of G is normal.

Additional characterizations of finite nilpotent groups have been given in [2, 3, 4]. To prove that (*) can be added to this list of equivalent conditions, we begin by showing that (*) holds for any finite p -group. For this, we require two facts about finite p -groups. First, every maximal subgroup of a finite p -group has index p . Second, every maximal subgroup of a p -group is normal. We refer to [1, Section 6.1] for proofs of these properties.

Lemma 2. *Let p be a prime and let G be a group of order p^n , where $n \geq 1$. Then, G satisfies (*).*

Proof. We use induction on n . If $n = 1$, then G is cyclic of order p and the result is clear. So, assume that $n > 1$ and that (*) holds for all groups of order p^{n-1} .

Let H be a proper subgroup of G . Then, there exists a maximal subgroup $M \leq G$ such that $H \leq M$. Let $[G : H] = p^k$, where $k \geq 1$. Since M is maximal in G , the index of M in G equals p . Consequently, $[M : H] = p^{k-1}$.

Since $M \trianglelefteq G$, $x^p \in M$ for all $x \in G$. By induction, (*) holds for M . Thus, for every $x \in G$, $x^{p^k} = (x^p)^{p^{k-1}} \in H$. \square

Lemma 3. *Let $G = G_1 \times \cdots \times G_n$, where G_i is a finite group for each $1 \leq i \leq n$ and $\gcd(|G_i|, |G_j|) = 1$ whenever $i \neq j$. Then, every subgroup of G has the form $H_1 \times \cdots \times H_n$, where $H_i \leq G_i$ for all $1 \leq i \leq n$.*

Proof. This is a standard result, but we include a proof for the sake of completeness. It suffices to prove the lemma for the case where $G = G_1 \times G_2$. The general result then follows by induction.

Let $n_1 = |G_1|$ and $n_2 = |G_2|$. Assume that G_1 and G_2 have identity elements e_1 and e_2 , respectively. Let $H \leq G$ and let $(x, y) \in H$. We first show that both (x, e_2) and (e_1, y) are in H . Since $\gcd(n_1, n_2) = 1$, we see that n_2 is coprime to $|x|$ and n_1 is coprime to $|y|$. So, x^{n_2} generates $\langle x \rangle$, and y^{n_1} generates $\langle y \rangle$. Let k_1, k_2 be integers such that $x = x^{n_2 k_2}$ and $y = y^{n_1 k_1}$. Then,

$$(x, y)^{n_2 k_2} = (x^{n_2 k_2}, e_2) = (x, e_2),$$

and likewise $(x, y)^{n_1 k_1} = (e_1, y)$. Thus, $(x, e_2), (e_1, y) \in H$.

Now, let $H_1 = \{(h_1, h_2) \in H \mid h_2 = e_2\}$ and $H_2 = \{(h_1, h_2) \in H \mid h_1 = e_1\}$. Both H_1 and H_2 are normal subgroups of H , and their intersection is trivial. Moreover, by the previous paragraph, $H = H_1 H_2$. So, $H = H_1 H_2 = H_1 \times H_2$, as desired. \square

A subgroup H of G such that $|H|$ and $[G : H]$ are coprime is called a *Hall subgroup* of G . For a Hall subgroup H , the condition that $x^{[G:H]} \in H$ for all $x \in G$ implies that $H \trianglelefteq G$.

Lemma 4. *Let G be a finite group and let $H \leq G$ be such that $|H|$ and $[G : H]$ are coprime. If $x^{[G:H]} \in H$ for all $x \in G$, then H is normal in G .*

Proof. Let $k = [G : H]$ and assume that $x^k \in H$ for all $x \in G$. Let $h \in H$. Since k is coprime to $|H|$, it is also coprime to $|h|$. Thus, $\langle h^k \rangle = \langle h \rangle$, and so $h = h^{k^m}$ for some integer m . For each $g \in G$, we have

$$ghg^{-1} = gh^{k^m}g^{-1} = (gh^m g^{-1})^k \in H,$$

so H is normal in G . \square

We can now prove that $(*)$ characterizes finite nilpotent groups. In fact, it is enough that the condition in $(*)$ holds only for Sylow subgroups of G .

Theorem 5. *Let G be a finite group. The following are equivalent.*

- (1) G is nilpotent.
- (2) G satisfies $(*)$.
- (3) For each Sylow subgroup P of G and for all $x \in G$, $x^{[G:P]} \in P$.

Proof. (1) \Rightarrow (2) Assume that G is nilpotent and let p_1, \dots, p_n be all of the distinct prime divisors of $|G|$. Since G is nilpotent, G has a unique Sylow p_i -subgroup P_i for each $1 \leq i \leq n$, and we can assume (up to isomorphism) that $G = P_1 \times \dots \times P_n$. Let $H \leq G$. By Lemma 3, $H = H_1 \times \dots \times H_n$, where $H_i \leq P_i$ for all $1 \leq i \leq n$. Let $k = [G : H]$ and let $x = (g_1, \dots, g_n) \in G$. For each i , $[P_i : H_i]$ divides k , and $(*)$ holds for P_i by Lemma 2, so $g_i^k \in H_i$. Thus, $x^k = (g_1^k, \dots, g_n^k) \in H$.

(2) \Rightarrow (3) This is clear.

(3) \Rightarrow (1) Assume that $x^{[G:P]} \in P$ whenever $x \in G$ and P is a Sylow subgroup of G . If G is a p -group, then G is nilpotent and we are done. If G is not a p -group, then we can apply Lemma 4 to conclude that each Sylow subgroup of G is normal in G . Hence, G is nilpotent in this case as well. \square

References

- [1] Dummit DS, Foote RM. Abstract Algebra. 3rd ed. Hoboken (NJ): John Wiley & Sons; 2004.

- [2] Cocke W. Two characterizations of finite nilpotent groups. *J. Group Theory*. 2018;21(6):1111–1116.
- [3] Holmes CV. A Characterization of Finite Nilpotent Groups. *Amer. Math. Monthly*. 1966;73(10):1113–1114.
- [4] Wong WJ. A cohomological characterization of finite nilpotent groups. *Proc. Amer. Math. Soc.* 1968;19:689–691.