

Note on Fourier inequalities

Miquel Saucedo and Sergey Tikhonov

ABSTRACT. We prove that the Hausdorff–Young inequality $\|\widehat{f}\|_{q(\cdot)} \leq C \|f\|_{p(\cdot)}$ with $q(x) = p'(1/x)$ and $p(\cdot)$ even and non-decreasing holds in variable Lebesgue spaces if and only if p is a constant. However, under the additional condition on monotonicity of f , we obtain a full characterization of Pitt-type weighted Fourier inequalities in the classical and variable Lebesgue setting.

1. Fourier inequalities in variable L_p spaces

For $f \in L_1(\mathbb{R}^n)$ we define the Fourier transform by

$$\widehat{f}(y) = \int_{\mathbb{R}^n} f(x) e^{-2\pi i(x,y)} dx.$$

An important estimate for the Fourier transform is the Hausdorff–Young inequality: for $1 \leq p \leq 2$,

$$(1.1) \quad \|\widehat{f}\|_{p'} \leq \|f\|_p,$$

where p' is the conjugate exponent for p , that is, $\frac{1}{p} + \frac{1}{p'} = 1$. A natural question is to obtain an analogue of the Hausdorff–Young inequality for the variable Lebesgue spaces [CF, Section 5.6.10].

Let f and p be measurable functions. Assume further that $1 \leq p(x) \leq \infty$. We set $\rho_p(f) = \int_{\mathbb{R}} |f(x)|^{p(x)} dx$ and define

$$\|f\|_{p(\cdot)} = \inf \left\{ \lambda > 0 : \rho_p(f/\lambda) \leq 1 \right\}.$$

2010 *Mathematics Subject Classification.* Primary 42B10; Secondary 46E30, 26D10.

Key words and phrases. Fourier transform, variable Lebesgue space, Weighted Fourier inequalities, Hardy–Littlewood type theorem.

M. Saucedo is supported by the Spanish Ministry of Universities through the FPU contract FPU21/04230. S. Tikhonov is supported by PID2020-114948GB-I00, 2021 SGR 00087, AP 14870758, the CERCA Programme of the Generalitat de Catalunya, and Severo Ochoa and María de Maeztu Program for Centers and Units of Excellence in R&D (CEX2020-001084-M).

A typical regularity condition on p is the log-Hölder continuity, i.e, that there exists a constant C such that for all $x, y \in \mathbb{R}$, $|x - y| < 1/2$, $|p(x) - p(y)| \leq \frac{C}{-\log(|x-y|)}$. If $p(x) = \infty$ we assume that there is $C > 0$ such that $\frac{1}{p(y)} \leq \frac{C}{-\log(|x-y|)}$ for $|x - y| < 1/2$.

In [CF, Question 5.61], it was proved that a natural extension of (1.1), namely,

$$\|\widehat{f}\|_{p'(\cdot)} \leq C \|f\|_{p(\cdot)}, \quad \frac{1}{p(\cdot)} + \frac{1}{p'(\cdot)} = 1,$$

does not hold in general and the following problem was posed:

Let $p(\cdot)$ be even and non-decreasing on \mathbb{R}_+ such that

$$(1.2) \quad 1 \leq \inf_{x \in \mathbb{R}} \text{ess } p(x) \leq \sup_{x \in \mathbb{R}} \text{ess } p(x) \leq 2.$$

Is it the case that there exists a constant $C > 0$ such that for any $f \in L_{p(\cdot)}$

$$\|\widehat{f}\|_{q(\cdot)} \leq C \|f\|_{p(\cdot)} \quad \text{with } q(x) = p'(1/x)?$$

We give a negative answer to this question by demonstrating that such an inequality is valid only in the classical, non-variable case.

THEOREM 1.1. *Let p be a continuous even non-decreasing function on \mathbb{R}_+ such that*

$$\lim_{x \rightarrow \infty} p(x) = p_\diamond \leq 2$$

and condition (1.2) holds. Let q be log-Hölder continuous. If the Hausdorff–Young inequality

$$(1.3) \quad \|\widehat{f}\|_{q(\cdot)} \leq C \|f\|_{p(\cdot)}$$

holds for any Schwartz function f , then $p(x) \leq p_\diamond \leq q'(x)$. In particular, if $q(x) = p'(1/x)$, then $p(x) = p_\diamond$.

REMARK 1.2. It is worth mentioning that the following non-classical Fourier inequality

$$\int_{\mathbb{R}} |\widehat{f}(\xi)|^{q(\xi)} d\xi \leq C_1 \int_{\mathbb{R}} |f(x)|^{p(x)} dx + C_2$$

holds, with positive constants C_1 and C_2 that depend on p and q but not on f , under some additional conditions on $p(\cdot)$ and $q(\cdot)$. See [CDF, Corollary 1.16].

PROOF. First, since $p(x) \leq p_\diamond$ it is enough to show that $q'(x) \geq p_\diamond$ or, equivalently, $q(x) \leq p'_\diamond$. Furthermore, by the modulation property of the Fourier transform, it suffices to show the result for $x = 0$, that is,

$$q(0) \leq p'_\diamond.$$

Second, assuming that (1.3) holds for any Schwartz function, we have, for any such f ,

$$(1.4) \quad \|\widehat{f}\|_{q(\cdot)} \leq C \|f\|_{p_\diamond}.$$

Indeed, letting $\tau_h f(x) = f(x - h)$, we get

$$\|\widehat{f}\|_{q(\cdot)} \leq C \|\tau_h f\|_{p(\cdot)}.$$

To verify (1.4), we will show that

$$(1.5) \quad \lim_{h \rightarrow \infty} \|\tau_h f\|_{p(\cdot)} = \|f\|_{p_\diamond}.$$

Noting that

$$\rho_p\left(\frac{\tau_h f}{\|f\|_{p_\diamond}}\right) = \int_{\mathbb{R}} \left(\frac{|f(x)|}{\|f\|_{p_\diamond}}\right)^{p(x+h)} dx$$

and since p is a bounded function, the dominated convergence theorem yields

$$\lim_{h \rightarrow \infty} \rho_p\left(\frac{\tau_h f}{\|f\|_{p_\diamond}}\right) = \int_{\mathbb{R}} \left(\frac{|f(x)|}{\|f\|_{p_\diamond}}\right)^{p_\diamond} = 1.$$

Thus,

$$\lim_{h \rightarrow \infty} \rho_p\left(\frac{\tau_h f}{\|f\|_{p_\diamond}}\right) = 1,$$

which implies (1.5), due to the fact that $p(x) \geq 1$ and if $1 - \varepsilon < \rho_p\left(\frac{\tau_h f}{\|f\|_{p_\diamond}}\right) < 1 + \varepsilon$, then $\rho_p\left(\frac{\tau_h f}{(1+\varepsilon)\|f\|_{p_\diamond}}\right) < 1 < \rho_p\left(\frac{\tau_h f}{(1-\varepsilon)\|f\|_{p_\diamond}}\right)$.

Third, assuming that q is log-Hölder continuous at the origin and that inequality (1.4) holds, we prove that $q(0) \leq p'_\diamond$. For a Schwartz function f satisfying $\widehat{f}(\xi) \geq 1$ for $|\xi| \leq 1$, we define

$$f_\lambda(x) = \lambda f(\lambda x).$$

Then, $\|f_\lambda\|_{p_\diamond} = \lambda^{1/p'_\diamond} \|f\|_{p_\diamond}$. We consider two cases. If $q(0) < \infty$,

$$\rho_q(C\lambda^{-\frac{1}{q(0)}} \widehat{f}_\lambda) > \int_{-\lambda}^{\lambda} |C\lambda^{-\frac{1}{q(0)}} \widehat{f}(\xi/\lambda)|^{q(\xi)} d\xi > \lambda^{-1} \int_{-\lambda}^{\lambda} C^{q(\xi)} \lambda^{1-\frac{q(\xi)}{q(0)}} d\xi.$$

Hence, by log-Hölder continuity, for some C we have $\lim_{\lambda \rightarrow 0} \rho_q(C\lambda^{-\frac{1}{q(0)}} \widehat{f}_\lambda) \geq 1$. Therefore, we deduce that

$$\lim_{\lambda \rightarrow 0} \|\widehat{f}_\lambda\|_{q(\cdot)} \lambda^{-\frac{1}{q(0)}} \geq \frac{1}{C}$$

and

$$\lambda^{\frac{1}{p'_\diamond}} \|f\|_{p_\diamond} = \|f_\lambda\|_{p_\diamond} \gtrsim \|\widehat{f}_\lambda\|_{q(\cdot)} \gtrsim \lambda^{\frac{1}{q(0)}},$$

that is, $q(0) \leq p'_\diamond$.

Finally, we assume that $q(0) = \infty$. If for any $\lambda > 0$ the set $\{-\lambda < x < \lambda : q(x) = \infty\}$ has positive measure, we deduce that for any $\lambda > 0$, $\rho_q(\widehat{f}_\lambda) \geq 1$ and, consequently, $\|\widehat{f}_\lambda\|_{q(\cdot)} \geq 1$. Then the Hausdorff-Young inequality (1.3) implies that $p'_\diamond = \infty$.

Otherwise, by the log-Hölder continuity of q , for large enough C , we derive as above that

$$\lim_{\lambda \rightarrow 0} \rho_q(C\widehat{f}_\lambda) \geq \int_{-\lambda}^{\lambda} |C|^{q(\xi)} d\xi \geq 1,$$

whence it follows that $p_\diamond = 1$.

□

2. Hardy–Littlewood theorem in the variable exponent setting

We now show that some nontrivial Fourier inequalities can still be derived in the variable exponent setting assuming additional conditions on f . We obtain an analogue of the classical Hardy–Littlewood inequality (see [Ti, Sa])

$$(2.1) \quad \|\widehat{f}(\xi)|\xi|^{n(1-\frac{2}{p})}\|_p \lesssim \|f\|_p, \quad 1 < p \leq 2,$$

for radial monotone functions; cf. [CH, GLT, De]. For a radial function g , we denote by g_0 its radial part: $g(x) = g_0(|\xi|)$.

THEOREM 2.1. *Let $n \in \mathbb{N}$ and $p(\cdot)$ be a radial function such that $p_0(\cdot) \geq 1$ is decreasing and bounded at the origin. The inequality*

$$\int_{\mathbb{R}^n} |\widehat{f}(\xi)|^{p(\xi)} |\xi|^{n(p(\xi)-2)} v(1/|\xi|) d\xi \lesssim \int_{\mathbb{R}^n} |f(x)|^{p(1/|x|)} v(|x|) dx$$

holds for any radial non-negative function f such that $f_0(x)x^{n-1}$ is non-increasing with

$$\lim_{x \rightarrow 0} f_0(x)x^{n-1} \leq 1$$

if and only if, for any positive r ,

$$(2.2) \quad \int_{|x| \geq r} r^{p(1/|x|)} |x|^{-np(1/|x|)} v(|x|) dx \lesssim \int_{|x| \leq r} v(|x|) |x|^{(1-n)p(1/|x|)} dx.$$

PROOF. First, we note that under the conditions on f , the Fourier transform is well defined; cf. Remark 3.2.

To see the necessity of condition (2.2), we set $f(x) = \chi(|x|)_{[0,r]} |x|^{-n+1}$. Then, for $\xi \leq \frac{\pi}{3}r^{-1}$ we have $\widehat{f}(\xi) \gtrsim r$. Hence,

$$\int_{|\xi| \leq \frac{\pi}{3}r^{-1}} r^{p(\xi)} |\xi|^{n(p(\xi)-2)} v(|\xi|^{-1}) d\xi = \int_{|\xi| \geq \frac{3}{\pi}r} r^{p(\xi^{-1})} |\xi|^{-np(\xi^{-1})} v(|\xi|) d\xi \lesssim \int_{|x| \leq r} v(x) |x|^{p(\xi^{-1})(1-n)} dx.$$

We now prove the sufficiency of condition (2.2). Using Theorem 3.1, we have

$$\begin{aligned} J &= \int_{\mathbb{R}^n} |\widehat{f}(\xi)|^{p(\xi)} |\xi|^{n(p(\xi)-2)} v(|\xi|^{-1}) d\xi \\ &\lesssim \int_0^\infty \xi^{n(p_0(\xi)-2)+n-1} v(\xi^{-1}) \left(\int_0^{\xi^{-1}} f_0(t) t^{n-1} dt \right)^{p_0(\xi)} d\xi \\ &= \int_0^\infty \xi^{-n(p_0(1/\xi)-2)-n-1} v(\xi) \left(\int_0^\xi f_0(t) t^{n-1} dt \right)^{p_0(1/\xi)} d\xi. \end{aligned}$$

Rewriting this using the Hardy operator $H(g)(x) = \frac{1}{x} \int_0^x g(t)dt$, we obtain

$$\begin{aligned} J &\lesssim \int_0^\infty \xi^{(n-1)(1-p_0(\xi^{-1}))} v(\xi) \left(H(f_0(t)t^{n-1}) \right)^{p_0(\xi^{-1})}(\xi) d\xi \\ &\lesssim \int_0^\infty \xi^{(n-1)(1-p_0(\xi^{-1}))} v(\xi) \left(f_0(\xi)\xi^{n-1} \right)^{p_0(\xi^{-1})}(\xi) d\xi \\ &= \int_0^\infty v(\xi)\xi^{n-1} f_0(\xi)^{p_0(\xi^{-1})} \asymp \int_{\mathbb{R}^n} |f(x)|^{p(x^{-1})} v(|x|) dx. \end{aligned}$$

Here the second inequality follows from the characterization of the boundedness of the Hardy operator for weighted modulars restricted to decreasing functions, as obtained in [Ne, Theorem 3]. It is a routine calculation to see that (2.2) is the required condition. \square

3. Fourier inequalities for monotone functions in the classical L_p

First, we recall that for radial f , its Fourier transform is given by

$$(3.1) \quad \widehat{f}(|\xi|) = (2\pi)^{n/2} |\xi|^{1-n/2} \int_0^\infty r^{n/2-1} f(r) J_{n/2-1}(|\xi|r) r dr,$$

where $J_\alpha(z)$ is the Bessel function of the first kind of order α . Our goal is to characterize weighted Pitt's inequality restricted to radially decreasing functions. First, we obtain the following upper bound for the Fourier transform of such functions:

THEOREM 3.1. *Let f be a radial function in \mathbb{R}^n . Assume that $r^{\frac{n-1}{2}} f_0(r)$ is a non-increasing function with limit zero. Then we have*

$$(3.2) \quad |\widehat{f}(|\xi|)| \lesssim \int_0^{|\xi|^{-1}} r^{n-1} f_0(r) dr.$$

REMARK 3.2. (i) Note that under the conditions of Theorem 3.1 the Fourier transform can be defined by (3.1) as $\lim_{N \rightarrow \infty} \int_0^N$.

(ii) Estimate (3.2) is not valid if we only assume in Theorem 3.1 that f is radial (take a characteristic function of the ball). Moreover, it is also clear that the monotonicity condition is optimal. Indeed, assuming that $r^{\frac{n-1}{2}} f_0(r) r^{-\varepsilon}$, with a positive ε , is non-increasing with limit zero, we observe that the integral in (3.1) does not even need to be convergent. This easily follows from the well-known asymptotics

$$(3.3) \quad J_\nu(y) = \sqrt{\frac{2}{\pi}} \cos(y - \theta_\nu) y^{-\frac{1}{2}} + B_\nu \sin(y - \theta_\nu) y^{-\frac{3}{2}} + O(y^{-\frac{5}{2}}),$$

see [Wa, Ch. 7].

(iii) It is worth mentioning that one can alternatively prove Theorem 3.1 with the help of the formula (3.3).

(iv) Note also that Theorem 3.1 extends the result by Carton-Lebrun and Heinig in [CH], which claims that (3.2) holds if $r^{n-1}f_0(r)$ is non-increasing. Moreover, inequality (3.2) improves the following result by Colzani, Fontana, and Laeng recently proved in [CFL]: under conditions of Theorem 3.1, there holds

$$|\widehat{f}(|\xi|)| \lesssim \frac{1}{|\xi|^{\frac{n-1}{2}}} \int_0^{|\xi|^{-1}} r^{\frac{n-1}{2}} f_0(r) dr.$$

PROOF. First note that it is enough to show that

$$|\widehat{f}(1)| \lesssim \int_0^1 f_0(r) r^{n-1} dr,$$

as the general result will follow by considering $x \mapsto f(\lambda x)$ instead of f .

We will make use of following well-known properties of the Bessel functions $J_\alpha(t)$, for $\alpha \geq -\frac{1}{2}$ and $t > 0$,

- (1) $\int_0^t r^{\alpha+1} J_\alpha(r) dr = t^{\alpha+1} J_{\alpha+1}(t)$;
- (2) $|J_\alpha(t)| \lesssim t^{-\frac{1}{2}}$;
- (3) $|J_\alpha(t)| \lesssim t^\alpha$.

In light of property (3) and by integration by parts, we arrive at

$$\begin{aligned} |\widehat{f}(1)| &\lesssim \int_0^\infty r^{n/2-1} f_0(r) J_{n/2-1}(r) r dr \lesssim \int_0^1 r^{n-1} f_0(r) dr + \int_1^\infty r^{\frac{1}{2}} J_{\frac{n}{2}-1}(r) \int_r^\infty d(f_0(s) s^{\frac{n-1}{2}}) \\ &\lesssim \int_0^1 r^{n-1} f_0(r) dr + \int_1^\infty d(f_0(s) s^{\frac{n-1}{2}}) \int_1^s J_{\frac{n}{2}-1}(r) r^{\frac{1}{2}} \\ &\lesssim \int_0^1 r^{n-1} f_0(r) dr + f_0(1) \lesssim \int_0^1 f_0(r) r^{n-1} dr, \end{aligned}$$

where we have used the following estimate: given $\alpha \geq -\frac{1}{2}$, there holds

$$\sup_x \left| \int_0^x t^{\frac{1}{2}} J_\alpha(t) dt \right| < \infty.$$

Indeed, integrating by parts and using property (1), we have

$$\begin{aligned} \int_0^x t^{\frac{1}{2}} J_\alpha(t) dt &= \int_0^x t^{\alpha+1} J_\alpha(t) t^{-\frac{1}{2}-\alpha} dt = c(\alpha) \int_0^x t^{\alpha+1} J_\alpha(t) \int_t^x u^{-\alpha-\frac{3}{2}} du dt \\ &+ x^{-\alpha-\frac{1}{2}} \int_0^x t^{\alpha+1} J_\alpha(t) dt = c(\alpha) \int_0^x u^{-\alpha-\frac{3}{2}} u^{\alpha+1} J_{\alpha+1}(u) du + x^{\frac{1}{2}} J_{\alpha+1}(x). \end{aligned}$$

The second term is bounded because of property (2). For the first one, a further integration by parts yields

$$\begin{aligned} \int_0^x t^{-\frac{1}{2}} J_{\alpha+1}(t) dt &= c(\alpha) \int_0^x t^{\alpha+2} J_{\alpha+1}(t) \int_t^x u^{-\alpha-\frac{7}{2}} du dt + x^{-\alpha-\frac{5}{2}} \int_0^x t^{\alpha+2} J_{\alpha+1}(t) dt \\ &= c(\alpha) \int_0^x u^{-\frac{3}{2}} J_{\alpha+2}(u) du + x^{-\frac{1}{2}} J_{\alpha+2}(x), \end{aligned}$$

which is bounded because of property (3) for small u and (2) for large u . \square

Now we are in a position to characterize weighted Fourier inequalities in the classical L_p spaces for radial decreasing functions.

THEOREM 3.3. *Let $n \in \mathbb{N}$ and $1 \leq p < \infty$. Let v be a positive weight function and $\frac{n-1}{2} \leq \alpha \leq n-1$. The inequality*

$$(3.4) \quad \int_{\mathbb{R}^n} |\widehat{f}(\xi)|^p |\xi|^{n(p-2)} v(1/|\xi|) d\xi \lesssim \int_{\mathbb{R}^n} |f(x)|^p v(|x|) dx$$

holds for any radial non-negative function f such that $f_0(x)x^\alpha$ is non-increasing with limit zero if and only if, for any positive r ,

$$(3.5) \quad r^{p(n-\alpha)} \int_{|x| \geq r} |x|^{-np} v(|x|) dx \lesssim \int_{|x| \leq r} v(x) |x|^{-\alpha p} dx.$$

REMARK 3.4. (i) Condition (3.5) is monotone with respect to α , that is, the class of v satisfying (3.5) becomes wider as α increases. Thus, Theorem 3.3 provides a natural balance between the monotonicity condition on f and the condition on the weight v . For $\alpha = n-1$ Theorem 3.3 was obtained in [CH]. (ii) Sufficient conditions on weights so that the weighted Fourier inequality (3.4) holds are usually given in terms of the rearrangement of the weight v , cf. [BH, He, RS, DGT17]. The monotonicity of f_0 allows us to obtain necessary and sufficient condition for (3.4) in terms of v itself.

PROOF. The proof of the necessity of condition (3.5) follows by considering $f(x) = \chi(|x|)_{[0,r]} |x|^{-\alpha}$. Then we have $\widehat{f}(\xi) \gtrsim r^{n-\alpha}$ for any $|\xi| \leq \frac{\pi}{3r}$. Hence,

$$r^{p(n-\alpha)} \int_{|\xi| \geq \frac{\pi}{3r}} |\xi|^{-np} v(|\xi|) d\xi = r^{p(n-\alpha)} \int_{|\xi| \leq \frac{\pi}{3r}} |\xi|^{n(p-2)} v(|\xi|^{-1}) d\xi \lesssim \int_{|x| \leq r} v(x) |x|^{-\alpha p} dx.$$

To prove the sufficiency, put $g(t) = f_0(t)t^\alpha$ and recall that by hypothesis g is non-increasing. Then, Theorem 3.1 yields

$$I := \int_{\mathbb{R}^n} |\widehat{f}(\xi)|^p |\xi|^{n(p-2)} v(|\xi|^{-1}) d\xi \lesssim \int_0^\infty \xi^{n(p-2)+n-1} v(\xi^{-1}) \left(\int_0^{1/\xi} g(t) t^{n-1-\alpha} dt \right)^p d\xi.$$

A change of variables implies that

$$\begin{aligned} I &\lesssim \int_0^\infty \xi^{-np+n-1} v(\xi) \left(\int_0^\xi g(t) t^{n-1-\alpha} dt \right)^p d\xi \lesssim \int_0^\infty g(t)^p t^{-\alpha p+n-1} v(t) dt \\ &\asymp \int_{\mathbb{R}^n} |f(x)|^p v(|x|) dx, \end{aligned}$$

provided that the following three weight Hardy inequality

$$(3.6) \quad \int_0^\infty w(t) \left(H(g(t)u(t)) \right)^p dt \lesssim \int_0^\infty g(t)^p \eta(t) dt,$$

with $w(t) = t^{-np+p+n-1}v(t)$, $\eta(t) = t^{-\alpha p+n-1}v(t)$, and $u(t) = t^{n-1-\alpha}$ holds for any non-increasing g .

The characterization of (3.6) is known and, for $p > 1$, according to [GS, Theorem 2.5], we need to examine that, for any positive r , there holds

$$\int_0^r \left(\int_0^t u(s) ds \right)^p \frac{w(t)}{t^p} dt \lesssim \int_0^r \eta(t) dt$$

and

$$(3.7) \quad \left(\int_r^\infty \frac{w(t)}{t^p} dt \right)^{\frac{1}{p}} \left(\int_0^r \eta(t) \left(\frac{\int_0^t u(s) ds}{\int_0^t \eta(s) ds} \right)^{p'} \right)^{\frac{1}{p'}} \lesssim 1.$$

The first condition holds since the integrals on the left-hand and right-hand sides are equivalent. Regarding the second condition, observe that, setting $\Delta(t) = \int_0^t \eta(s) ds$,

$$\begin{aligned} \int_0^r \left(\int_0^t u(s) ds \right)^{p'} \frac{\eta(t)}{\Delta^{p'}(t)} dt &\asymp \int_0^r \left(\int_0^t x^{p'(n-\alpha)-1} dx \right) \frac{\eta(t)}{\Delta^{p'}(t)} dt \\ &= \int_0^r \left(\int_x^r \frac{\eta(t)}{\Delta^{p'}(t)} dt \right) x^{p'(n-\alpha)-1} dx \\ &\lesssim \int_0^r \Delta^{1-p'}(x) x^{p'(n-\alpha)-1} dx. \end{aligned}$$

Further, by our assumption, we have

$$\int_r^\infty \frac{w(t)}{t^p} dt = \int_r^\infty \xi^{-np+n-1} v(\xi) d\xi \lesssim r^{p(\alpha-n)} \int_0^r \eta(t) dt = r^{p(\alpha-n)} \Delta(r).$$

Therefore, to prove (3.7), it suffices to show that

$$r^{p(\alpha-n)} \int_0^r \eta(t) dt \lesssim \left(\int_0^r \Delta^{1-p'}(x) x^{p'(n-\alpha)-1} dx \right)^{\frac{1}{1-p'}}.$$

Now, since $1 - p' < 0$, the reverse Minkowski inequality (see, e.g., [Bu, p. 189]) yields

$$\begin{aligned} \left(\int_0^r \Delta^{1-p'}(x) x^{p'(n-\alpha)-1} dx \right)^{\frac{1}{1-p'}} &= \left(\int_0^r \left(\int_0^x \eta(t) x^{\frac{p'(n-\alpha)-1}{1-p'}} dt \right)^{1-p'} dx \right)^{\frac{1}{1-p'}} \\ &\geq \int_0^r \eta(t) \left(\int_t^r x^{p'(n-\alpha)-1} dx \right)^{\frac{1}{1-p'}} dt \\ &\gtrsim \left(r^{p'(n-\alpha)} \right)^{\frac{1}{1-p'}} \int_0^r \eta(t) dt, \end{aligned}$$

and the result follows. In the case $p = 1$ the proof is similar with the help of [GS, Theorem 2.5(d)]. □

Let us compare our result with the classical Pitt inequality: for $1 < p < \infty$, one has

$$(3.8) \quad \int_{\mathbb{R}^n} |\widehat{f}(\xi)|^p |\xi|^{n(p-2)-\gamma p} d\xi \lesssim \int_{\mathbb{R}^n} |f(x)|^p |x|^{\gamma p} dx$$

for any f provided that $\max(0, n(1 - \frac{2}{p})) \leq \gamma < \frac{n}{p'}$, see e.g. [BH]. For $1 < p \leq 2$ and $\gamma = 0$, (3.8) coincides with the Hardy–Littlewood inequality (2.1).

COROLLARY 3.5. *Let $1 \leq p < \infty$. Assume that $f_0(r)r^{\frac{n-1}{2}}$ is non-increasing with limit zero. Then (3.8) holds for any $\gamma < \frac{n}{p'}$.*

This result was known for functions f such that $f_0(r)r^{n-1}$ is non-increasing (see [CH]). Combining various conditions on f , we arrive at the following result:

COROLLARY 3.6. *Let $1 < p < \infty$. Inequality (3.8) holds for any $f \in X$ if and only if*

- (i) $\max(0, n(1 - \frac{2}{p})) \leq \gamma < \frac{n}{p'}$ if $X = L_1(\mathbb{R}^n)$, see (3.8);
- (ii) $(n-1)(\frac{1}{2} - \frac{1}{p}) + \max(0, 1 - \frac{2}{p}) \leq \gamma < \frac{n}{p'}$ if X is the space of radial, integrable functions, see [DGT13, Th.1.1];
- (iii) $\frac{n-1}{2} - \frac{n}{p} < \gamma < \frac{n}{p'}$ if X is the space of radial integrable functions f for which $f(x) = f_0(|x|)$ with $f_0 \in GM$, that is, f_0 is locally integrable on \mathbb{R}_+ and, for some $\gamma > 1$, the inequality

$$\text{Var}_{(x,2x)} f_0 \lesssim \int_{x/\gamma}^{\gamma x} \frac{|f_0(t)|}{t} dt, \quad x > 0,$$

holds, see [GLT];

- (iv) $\gamma < \frac{n}{p'}$ if X is the space of radial functions for which $f(x) = f_0(|x|)$, with $f_0(r)r^{\frac{n-1}{2}}$ monotonically decreasing on \mathbb{R}_+ (see Corollary 3.5).

It is easy to see the conditions on γ become less restrictive from (i) to (iv).

References

- [BH] J. J. Benedetto, H. P. Heinig, Weighted Fourier inequalities: new proofs and generalizations, J. Fourier Anal. Appl. 9 (2003), no. 1, 1–37.
- [Bu] P. S. Bullen, Handbook of Means and Their Applications, Mathematics and Its Applications, 2003.
- [CH] C. Carton-Lebrun; H. P. Heinig, Weighted Fourier transform inequalities for radially decreasing functions, SIAM J. Math. Anal. 23 (1992), no. 3, 785–798.
- [CFL] L. Colzani, L. Fontana, E. Laeng, Asymptotic decay of Fourier, Laplace and other integral transforms, J. Math. Anal. Appl. 483, no. 1 (2020), Article 123560.
- [CDF] D. Cruz-Uribe, G. Di Fratta, A. Fiorenza, Modular inequalities for the maximal operator in variable Lebesgue spaces, Nonlinear Anal., Theory Methods Appl., Ser. A, Theory Methods 177 (2018), 299–311.
- [CF] D. Cruz-Uribe, A. Fiorenza, Variable Lebesgue Spaces, Appl. Numer. Harmon. Anal. Birkhäuser/Springer, Heidelberg, 2013.

- [De] A. Debernardi, The Boas problem on Hankel transforms, *J. Fourier An. Appl.*, 25 (2019), 3310–3341.
- [DGT13] L. De Carli, D. Gorbachev, S. Tikhonov, Pitt and Boas inequalities for Fourier and Hankel transforms, *J. Math. Anal. Appl.*, 408, no. 2, (2013), 762–774.
- [DGT17] L. De Carli, D. Gorbachev, S. Tikhonov, Pitt inequalities and restriction theorems for the Fourier transform, *Rev. Mat. Iberoam.* 33, no. 3 (2017), 789–808.
- [GS] A. Gogatishvili, V. Stepanov, Reduction theorems for weighted integral inequalities on the cone of monotone functions, *Russ. Math. Surv.* 68, no. 4 (2013), 597–664; translation from *Usp. Mat. Nauk* 68, no. 4 (2013), 3–68.
- [GLT] D. Gorbachev, E. Lifyand, S. Tikhonov, Weighted Fourier Inequalities: Boas conjecture in \mathbb{R}^n , *J. d’Analyse Math.*, Vol. 114 (2011), 99–120.
- [He] H. Heinig, Weighted norm inequalities for classes of operators, *Indiana U. Math. J.*, 33, no. 4, (1984), 573–582.
- [Ne] C. J. Neugebauer, Weighted variable L_p integral inequalities for the maximal operator on non-increasing functions, *Stud. Math.* 192, no. 1 (2009), 51–60.
- [RS] J. Rastegari, G. Sinnamon, Weighted Fourier inequalities via rearrangements, *J. Fourier Anal. Appl.* 24, 5 (2018), 1225–1248.
- [Sa] C. Sadosky, *Interpolation of Operators and Singular Integrals. An Introduction to Harmonic Analysis.* New York, Basel: Marcel Dekker, 1979.
- [Ti] E. C. Titchmarsh, *Introduction to the Theory of Fourier Integrals*, Oxford: Clarendon Press, 1937.
- [Wa] G. N. Watson, *A Treatise on the Theory of Bessel Functions.* 2nd ed. Cambridge Univ. Press, 1995.

M. SAUCEDO, CENTRE DE RECERCA MATEMÀTICA, CAMPUS DE BELLATERRA, EDIFICI C 08193 BELLATERRA (BARCELONA), SPAIN; UNIVERSITAT AUTÒNOMA DE BARCELONA, CAMPUS DE BELLATERRA, EDIFICI C 08193 BELLATERRA (BARCELONA), SPAIN.

Email address: miquelsaucedo98@gmail.com

S. TIKHONOV, ICREA, PG. LLUÍS COMPANYS 23, 08010 BARCELONA, SPAIN, CENTRE DE RECERCA MATEMÀTICA, CAMPUS DE BELLATERRA, EDIFICI C 08193 BELLATERRA (BARCELONA), SPAIN, AND UNIVERSITAT AUTÒNOMA DE BARCELONA, CAMPUS DE BELLATERRA, EDIFICI C 08193 BELLATERRA (BARCELONA), SPAIN .

Email address: stikhonov@crm.cat