

Multiple scattering and diffusion of scalar coherent waves in a group of small spheroidal particles with random orientations

Mingyuan Ren, Yajing Qiao, Ning Zhou, Jianrui Gong, Yang Zhou, Yu Zhang *

Department of Physics, Harbin Institute of Technology,

No.92, XiDazhi Street, Harbin 150001, China

{Mingyuan Ren} hitphyrmy@126.com

{Yu Zhang} zhangyuhitphy@163.com

July 9, 2024

Abstract

In this manuscript we study multiple scattering and diffusion of scalar wave in a group of monodisperse spheroidal particles with random orientations. We begin by fixing a spheroid in a prolate spheroidal coordinate system, and attain the expansion of the scalar Green's function in this space. The expansion is firstly based on spheroidal wave functions, and then we transform it into the expansion of spherical wave functions. Next, we average the Green's function over the orientations of the spheroid to get the averaged transition operator. Finally, we calculate the transport mean free path and anisotropy factor for the spheroidal particles group, based on the irreducible vertex in the Bethe-Salpeter equation. The approaches to get the average transition operator and the mean free paths in this manuscript will be of benefit to the research area of multiple scattering by non-spherical particles.

1 Introduction

One of the most widely investigated problems in waves scattering is spherical particles scattering, which is initially theoretically solved by Gustav Mie [1]. Mie theory has been used in many applications since spherical particle is an efficient model in numerical calculations [2]. However, there are more particles in natural environment that is non-spherical, including spheroids, cylinders, and even particles aggregates [3]. Accordingly, many researchers have proposed various kinds of approaches to obtain scattered fields by non-spherical particles. Such approaches include Waterman's T-matrix method [4], separation of variables method [5, 6], dyadic Green's function method [7, 8], etc. These three methods all concern eigenfunction expansions in corresponding coordinate system and provide exact solutions, but they are applicable to different theoretical framework. For example, the Green's function method is the most suitable one to multiple scattering theory.

When a wave propagates in a group of particles, it is possibly scattered many times by different particles. This process will gradually become dominant as the propagation distance exceeds the scattering mean free path [9, 10, 11]. The single-scattering methods mentioned above are not enough in this case, and multiple scattering theory should be introduced. For matter waves and electrons, the quantum kinetic equation in multiple scattering processes has been derived and further used to analyze weak localization [12, 13]. The Diffuson and the Cooperon are proved to have close connection to the irreducible vertex. Recently, multiple scattering of polarized light in point scatterers, as well as that in disordered media exhibiting short-range structural correlations are theoretically derived, based on the Bethe-Salpeter equation [14, 15]. The latter paper has used Henyey-Greenstein phase function to replace the irreducible vertex, and discussed depolarization effect of different eigenmodes. Thus, the irreducible vertex plays a central role in multiple scattering, but in fact, it is determined by each single scattering process in the independent scattering approximation [16].

If the particles are sparsely distributed in the medium, we can assume that any two of them have no correlations and this is called independent scattering approximation (ISA). In this approximation, the irreducible vertex can be directly calculated by the transition operator [17, 18, 10], while this operator

can be directly calculated from the Green's function for one single scatterer [19, 20]. Tai has given the dyadic Green's function for a spherical object and several kinds of non-spherical objects [7], and Li et al. have derived the dyadic Green's function for a spheroid [21, 8]. Based on these Green's functions, the irreducible vertex and diffusion constant can be calculated, which gives results of diffusion and weak localization.

However, the Green's functions in [7, 8] are calculated in the case that the scatterer has a fixed orientation, which is not the case of natural environment. If one wants to derive multiple scattering in a group of non-spherical particles, the averaging over the orientations of particles is necessary. Using Waterman's T-matrix method and the orthogonality of Wigner D function, the averaged scattering amplitude could be calculated, which could further be used to calculate scattering and extinction cross section [4, 3]. According to this process, we show how to calculate the averaged transition operator in this manuscript, and use this operator to calculate the mean free paths. Compared to T-matrix, transition operator is more suitable to multiple scattering theory. We believe the demonstration of this manuscript will benefit the research area of non-spherical particles scattering, and may have potential applications in dense discrete particles scattering.

This manuscript is organized as follows. In Section 2, we show the relation between scalar scattering Green's function and the corresponding transition operator. Then we calculate the Green's function for a fixed spheroid in Subsection 2.2, and demonstrate how to get the average transition operator in Subsection 2.3. Next, in Subsection 3.1, we introduce the multiple scattering theory in the formalism of transition operator, and introduce the diffusion approximation and the far-field approximation in Subsection 3.2. Finally, we calculate the scattering and transport mean free paths of monodisperse spheroidal particles group in Subsection 3.3. The conclusion is drawn in Section 4.

2 The scalar transition operator for a spheroid of random orientation

2.1 From Green's function of a dielectric particle to its transition operator

A transition operator T_j of a single particle (we use the subscript j to represent the j th particle in the particles group) describes how a light field arrives at this particle, possibly bends its direction many times inside this particle, and then leaves it for good. This process contains a number of terms called Born series, which could be expressed by

$$\begin{aligned} T_j(\mathbf{r}, \mathbf{r}') = & U(\mathbf{r}) \delta(\mathbf{r}, \mathbf{r}') + U(\mathbf{r}) G_0(\mathbf{r}, \mathbf{r}') U(\mathbf{r}') \\ & + \int d\mathbf{r}_1 U(\mathbf{r}) G_0(\mathbf{r}, \mathbf{r}_1) U(\mathbf{r}_1) G_0(\mathbf{r}_1, \mathbf{r}') U(\mathbf{r}') \\ & + \dots, \end{aligned} \quad (1)$$

where

$$U(\mathbf{r}) = \begin{cases} k_0^2 \delta \epsilon_p, & \text{for } \mathbf{r} \text{ inside the particle,} \\ 0, & \text{for } \mathbf{r} \text{ outside the particle} \end{cases}$$

is the scattering potential function of this particle, with k_0 the wavenumber in the host medium and $\delta \epsilon_p = \epsilon_p - \epsilon_0$ the difference of dielectric permittivity between the particle ϵ_p and the host medium ϵ_0 . G_0 is the free space Green's function. Noticing the iterative form of Eq. (1), we rewrite it as

$$T_j(\mathbf{r}, \mathbf{r}') = U(\mathbf{r}) \delta(\mathbf{r}, \mathbf{r}') + \int d\mathbf{r}_1 T_j(\mathbf{r}, \mathbf{r}_1) G_0(\mathbf{r}_1, \mathbf{r}') U(\mathbf{r}'). \quad (2)$$

Those terms in Eq. (1) could certainly be calculated one by one, but it is difficult and also unnecessary to perform this calculation. Instead, we can use the technique of Green's function to get the expression of T_j . Now, we firstly show the relationship between T_j and its corresponding Green's function. Solving the scalar Helmholtz equation

$$\nabla^2 G(\mathbf{r}, \mathbf{r}') + k_0^2 G(\mathbf{r}, \mathbf{r}') = -\delta(\mathbf{r}, \mathbf{r}'), \quad (3)$$

we get the expression of the Green's function

$$G_e(\mathbf{r}, \mathbf{r}') = G_0(\mathbf{r}, \mathbf{r}') + \int d\mathbf{r}_1 G_0(\mathbf{r}, \mathbf{r}_1) U(\mathbf{r}_1) G_e(\mathbf{r}_1, \mathbf{r}'), \quad (4)$$

where we follow Ref. [7] to write the Green's function as G_e . By noticing the iterative form of Eq. (4) and comparing it with Eq. (2), we get another expression of the Green's function

$$G_e(\mathbf{r}, \mathbf{r}') = G_0(\mathbf{r}, \mathbf{r}') + \int d\mathbf{r}_2 d\mathbf{r}_1 G_0(\mathbf{r}, \mathbf{r}_2) T_j(\mathbf{r}_2, \mathbf{r}_1) G_0(\mathbf{r}_1, \mathbf{r}'), \quad (5)$$

Multiplying Eq. (2) by $G_0(\mathbf{r}'', \mathbf{r})$ and performing integration over \mathbf{r} , we get

$$\int d\mathbf{r} G_0(\mathbf{r}'', \mathbf{r}) T_j(\mathbf{r}, \mathbf{r}') = G_0(\mathbf{r}'', \mathbf{r}') U(\mathbf{r}') + \int d\mathbf{r} d\mathbf{r}_1 G_0(\mathbf{r}'', \mathbf{r}) T_j(\mathbf{r}, \mathbf{r}_1) G_0(\mathbf{r}_1, \mathbf{r}') U(\mathbf{r}'). \quad (6)$$

Comparing Eq. (6) with Eq. (5) we get

$$\int d\mathbf{r} G_0(\mathbf{r}'', \mathbf{r}) T_j(\mathbf{r}, \mathbf{r}') = G_e(\mathbf{r}'', \mathbf{r}') U(\mathbf{r}'), \quad (7)$$

whose Fourier transformation is

$$G_0(\mathbf{p}) T_j(\mathbf{p}, \mathbf{p}') = \int d\mathbf{r} d\mathbf{r}' e^{-i\mathbf{p}\cdot\mathbf{r}} G_e(\mathbf{r}'', \mathbf{r}') U(\mathbf{r}') e^{i\mathbf{p}'\cdot\mathbf{r}'}, \quad (8)$$

in which

$$T_j(\mathbf{p}, \mathbf{p}') = \int d\mathbf{r} d\mathbf{r}' e^{-i\mathbf{p}\cdot\mathbf{r}} T_j(\mathbf{r}, \mathbf{r}') e^{i\mathbf{p}'\cdot\mathbf{r}'}. \quad (9)$$

Eq. (8) will be used to calculate transition operator.

2.2 The scalar Green's function for a fixed dielectric spheroid

Applying the boundary condition to a non-spherical particle in a general coordinate system is not as simple as Mie theories. But fortunately, we have other coordinate systems to choose. Using a prolate spheroidal coordinate system, Li et al. have derived the dyadic Green's function for a dielectric spheroid [21, 8]. Following their work, we here develop the scalar Green's function for a dielectric spheroid.

We consider a spheroidal particle centered at the origin of the coordinate system, whose dielectric permittivity is homogeneous. We keep its orientation steady for now, as shown by Figure 1. We use a prolate spheroidal coordinate system $(\xi, \vartheta, \varphi)$, which results from rotating the two-dimensional elliptic coordinate system about the focal axis of the ellipse. The relation between the prolate spheroidal coordinates and the Cartesian coordinates are given by

$$\begin{aligned} x &= a \sinh \xi \sin \vartheta \cos \varphi, \\ y &= a \sinh \xi \sin \vartheta \sin \varphi, \\ z &= a \cosh \xi \cos \vartheta, \end{aligned} \quad (10)$$

where a is the semi-interfocal distance of the spheroid. The radial coordinate ξ forms a series of spheroidal surfaces. Now we can express the scalar Helmholtz equation in spheroidal coordinates. By the usual procedure of the separation of variables, we obtain the scalar wave eigenfunctions

$$\psi_{mn}^{(i)}(k_0 a, \mathbf{r}) = R_{mn}^{(i)}(k_0 a, \cosh \xi) \mathcal{P}_n^m(k_0 a, \cos \vartheta) e^{im\varphi} \quad (11)$$

where the radial function $R_{mn}^{(i)}$ satisfies Eq. (65) and the angular function \mathcal{P}_n^m satisfies Eq. (66). Using $\psi_{mn}^{(i)}$, the free space Green's function could be expressed by [21, 8]

$$\begin{aligned} G_0(\mathbf{r}, \mathbf{r}') &= \frac{ik_0}{2\pi} \sum_{n=0}^{\infty} \sum_{m=0}^n (2 - \delta_{m,0}) \\ &\quad \times R_{mn}^{(3)}(k_0 a, \cosh \xi^{\geq}) R_{mn}^{(1)}(k_0 a, \cosh \xi^{\leq}) \\ &\quad \times \overline{\mathcal{P}}_n^m(k_0 a, \cos \vartheta) \overline{\mathcal{P}}_n^m(k_0 a, \cos \vartheta') e^{im(\varphi - \varphi')}, \end{aligned} \quad (12)$$




Figure 1: Prolate spheroidal coordinate system $(\xi, \vartheta, \varphi)$

where \mathbf{r}^{\geq} , \mathbf{r}^{\leq} mean $\xi^{\leq} = \min\{\xi, \xi'\}$, $\xi^{\geq} = \max\{\xi, \xi'\}$, respectively, in spheroidal coordinates. And they mean $r^{\leq} = \min\{r, r'\}$, $r^{\geq} = \max\{r, r'\}$, respectively, in spherical coordinates. When ξ reaches infinity, the radial function $R_{mn}^{(i)}$ tends to be a spherical wave, and when ξ reaches zero, $R_{mn}^{(i)}$ must be finite, so we choose $R_{mn}^{(3)}$ for ξ^{\geq} and $R_{mn}^{(1)}$ for ξ^{\leq} . The radial function $R_{mn}^{(i)}$ and the normalized angular function $\overline{\mathcal{P}}_n^m$ are shown in Appendix. A.

By the method of scattering superposition, we can construct the Green's function G_e for a dielectric spheroid

$$\begin{aligned} G_e^{(22)}(\mathbf{r}, \mathbf{r}') &= G_0(\mathbf{r}, \mathbf{r}') + G_{es}^{(22)}(\mathbf{r}, \mathbf{r}'), & \xi \leq \xi_0 \\ G_e^{(12)}(\mathbf{r}, \mathbf{r}') &= G_{es}^{(12)}(\mathbf{r}, \mathbf{r}'), & \xi \geq \xi_0, \end{aligned} \quad (13)$$

where $G_e^{(12)}$ means that the source is located in region 1, namely outside the particle, and the receiving point is located in region 2, namely inside the particle. Accordingly, $G_e^{(22)}$ means that the source and the receiving point are both located inside the particle. $G_e^{(21)}$ and $G_e^{(11)}$ are not given here, since we can deduce from Eq. (7) that they will vanish in the transition operator because of the presence of U . G_{es} is the scattered part in G_e . It is important that the free space Green's function here has a wavenumber of $k_s = k_0\sqrt{\epsilon_p}$ because it describes propagation inside the scatterer. Considering the composition of G_0 , the scattered part G_{es} must have the form [7]

$$\begin{aligned} G_{es}^{(22)}(\mathbf{r}, \mathbf{r}') &= \frac{ik_s}{2\pi} \sum_{n=0}^{\infty} \sum_{m=0}^n (2 - \delta_{m,0}) A_{mn} \\ &\quad \times R_{mn}^{(1)}(k_s a, \cosh\xi) R_{mn}^{(1)}(k_s a, \cosh\xi') \\ &\quad \times \overline{\mathcal{P}}_n^m(k_s a, \cos\vartheta) \overline{\mathcal{P}}_n^m(k_s a, \cos\vartheta') e^{im(\varphi - \varphi')}, \\ G_{es}^{(12)}(\mathbf{r}, \mathbf{r}') &= \frac{ik_s}{2\pi} \sum_{n=0}^{\infty} \sum_{m=0}^n (2 - \delta_{m,0}) C_{mn} \\ &\quad \times R_{mn}^{(3)}(k_0 a, \cosh\xi) R_{mn}^{(1)}(k_s a, \cosh\xi') \\ &\quad \times \overline{\mathcal{P}}_n^m(k_s a, \cos\vartheta) \overline{\mathcal{P}}_n^m(k_s a, \cos\vartheta') e^{im(\varphi - \varphi')}. \end{aligned} \quad (14)$$

The coefficients A_{mn} and C_{mn} are obtained in Appendix B.

2.3 Averaging the Green's function and the transition operator

In a medium full of spherical particles, the rotation of each sphere has no impact on the transition operator, meaning that the operator is absolutely spherically symmetric. To get the average transition operator $\langle T \rangle$, we only need to average over the possible positions of all particles. However, this simplicity breaks down if the particle is non-spherical, since the orientations of particles will influence

T_j , and must be considered in averaging. So we perform the averaging on its orientation for one single spheroid centered at the origin of the coordinate system. In light of the expansion of Green's function, as well as the relationship Eq. (8), we realize that it is useful to first calculate $G_e U$, which is

$$G_e(\mathbf{r}, \mathbf{r}') U(\mathbf{r}') = k_0^2 \delta \epsilon_p G_e^{(22)}(\mathbf{r}, \mathbf{r}') + k_0 \delta \epsilon_p G_e^{(12)}(\mathbf{r}, \mathbf{r}'). \quad (15)$$

$G_e^{(12)}$ and $G_e^{(22)}$ vanish because of the existence of U .

In the last subsection, we calculate the Green's function by aligning the symmetry-axis of the spheroid with z -axis of Cartesian coordinates. However, in natural environment, the orientations of particles are random, as shown by Figure 2. We must take it into consideration that the expansions of Green's function in the laboratory coordinate system are different from that in the particle coordinate system. Now, let us assume that the Euler angles α, β, γ determine the rotation from the laboratory coordinate system $S^{(L)}\{x^{(L)}, y^{(L)}, z^{(L)}\}$ to the particle coordinate system $S^{(P)}\{x^{(P)}, y^{(P)}, z^{(P)}\}$. To average the operator $G_e U$, we need to perform this averaging over all orientations of particles. This means that, we fix ourselves in $S^{(L)}$, rotate $S^{(P)}$, and take into account all values of α, β, γ . However, we can conversely fix ourselves in $S^{(P)}$, rotate $S^{(L)}$, and average $G_e U$. Obviously the latter manipulation is simpler than the former one.




Figure 2: A field (the red solid line) scattering multiply in a group of spheroidal particles. The local prolate spheroidal coordinate systems are sketched in green for each particle.




Figure 3: Diffusion of scalar coherent wave in a group of spheroidal particles. The red dashed line represents the conjugate field. The local prolate spheroidal coordinate systems are sketched in green for each particle.

Before calculating $\langle G_e U \rangle$, we call for another step. The randomness of the orientation is actually from the perspective of (θ, ϕ) , namely the spherical coordinates, and the angles ϑ of spheroidal coordinates are not homogeneous in the natural space. Therefore, it is better to transform those functions into spherical coordinates, if we need to average over Euler angles. First, by the relation between the spheroidal wave function and the spherical wave function [22]

$$R_{mn}^{(i)}(k_0 a, \xi) \overline{\mathcal{P}}_n^m(k_0 a, \cos \vartheta) = \sum_{l=m}^{\infty} i^{l-n} d_{l-m}^{mn}(k_0 a) \frac{N_{mn}(0)}{N_{ml}(k_0 a)} z_l^{(i)}(k_0 r) \overline{P}_l^m(\cos \theta), \quad (16)$$

where $z_l^{(i)}(r)$ is the spherical Bessel function of order l and type i , and $\overline{P}_l^m(\cos \theta) = N_{mn}^{-1}(0) P_l^m(\cos \theta)$ is the associated Legendre function with the normalizing factor given by Eq. (69). And $P_l^m(\cos \theta)$ is the associated Legendre functions with the general definition. Then the scattering Green's function could be expressed by

$$\begin{aligned} G_{es}^{(22)}(\mathbf{r}, \mathbf{r}') &= \frac{ik_s}{2\pi} \sum_{n=0}^{\infty} \sum_{m=0}^n (2 - \delta_{m,0}) A_{mn} e^{im(\phi - \phi')} \\ &\times \sum_{l=m}^{\infty} i^{l-n} d_{l-m}^{mn}(k_s a) \frac{N_{mn}(0)}{N_{ml}(k_s a)} z_l^{(1)}(k_s r) \overline{P}_l^m(\cos \theta) \\ &\times \sum_{l'=m}^{\infty} i^{l'-n} d_{l'-m}^{mn}(k_s a) \frac{N_{mn}(0)}{N_{ml'}(k_s a)} z_{l'}^{(1)}(k_s r') \overline{P}_{l'}^m(\cos \theta'), \end{aligned} \quad (17)$$

$$\begin{aligned}
G_{es}^{(12)}(\mathbf{r}, \mathbf{r}') &= \frac{ik_s}{2\pi} \sum_{n=0}^{\infty} \sum_{m=0}^n (2 - \delta_{m,0}) C_{mn} e^{im(\phi - \phi')} \\
&\times \sum_{l=m}^{\infty} i^{l-n} d_{l-m}^{mn}(k_0 a) \frac{N_{mn}(0)}{N_{ml}(k_0 a)} z_l^{(3)}(k_0 r) \bar{P}_l^m(\cos\theta) \\
&\times \sum_{l'=m}^{\infty} i^{l'-n} d_{l'-m}^{mn}(k_s a) \frac{N_{mn}(0)}{N_{ml'}(k_s a)} z_{l'}^{(1)}(k_s r') \bar{P}_{l'}^m(\cos\theta'). \tag{18}
\end{aligned}$$

The Wigner D function $D_{m'm}^l(\alpha, \beta, \gamma)$ can be used to calculate the rotation from $S^{(L)}\{r, \theta^{(L)}, \phi^{(L)}\}$ to $S^{(P)}\{r, \theta^{(P)}, \phi^{(P)}\}$ [23]

$$Y_l^m(\theta^{(P)}, \phi^{(P)}) = \sum_{k=-l}^l Y_l^k(\theta^{(L)}, \phi^{(L)}) D_{km}^l(\alpha, \beta, \gamma), \tag{19}$$

where $\sqrt{2\pi} Y_l^m(\theta, \phi) = \bar{P}_l^m(\cos\theta) e^{im\phi}$ is the spherical harmonics. It is important to note that the angular coordinate φ in a spheroidal coordinate system is identical to the angular coordinate ϕ in its corresponding spherical coordinate system [22]. Averaging the scattering Green's function Eq. (17) and (18) yields

$$\begin{aligned}
\langle G_{es}^{(22)}(\mathbf{r}^{(P)}, \mathbf{r}'^{(P)}) \rangle &= \frac{ik_s}{2\pi} \sum_{n=0}^{\infty} \sum_{m=0}^n (2 - \delta_{m,0}) A_{mn} \\
&\times \sum_{l=m}^{\infty} i^{l-n} d_{l-m}^{mn}(k_s a) \frac{N_{mn}(0)}{N_{ml}(k_s a)} z_l^{(1)}(k_s r) \sum_{l'=m}^{\infty} i^{l'-n} d_{l'-m}^{mn}(k_s a) \frac{N_{mn}(0)}{N_{ml'}(k_s a)} z_{l'}^{(1)}(k_s r') \\
&\times 2\pi \sum_{k=-l}^l Y_l^k(\theta^{(L)}, \phi^{(L)}) \sum_{k'=-l'}^{l'} Y_{l'}^{k'*}(\theta'^{(L)}, \phi'^{(L)}) \langle D_{km}^l(\alpha, \beta, \gamma) D_{k'm}^{l'*}(\alpha, \beta, \gamma) \rangle \\
&= \frac{ik_s}{4\pi} \sum_{n=0}^{\infty} \sum_{m=0}^n (2 - \delta_{m,0}) A_{mn} \sum_{l=m}^{\infty} \left[i^{l-n} d_{l-m}^{mn}(k_s a) \frac{N_{mn}(0)}{N_{ml}(k_s a)} \right]^2 \\
&\times z_l^{(1)}(k_s r) z_l^{(1)}(k_s r') P_l(\cos\Theta), \tag{20}
\end{aligned}$$

$$\begin{aligned}
\langle G_{es}^{(12)}(\mathbf{r}^{(P)}, \mathbf{r}'^{(P)}) \rangle &= \frac{ik_s}{2\pi} \sum_{n=0}^{\infty} \sum_{m=0}^n (2 - \delta_{m,0}) C_{mn} \\
&\times \sum_{l=m}^{\infty} i^{l-n} d_{l-m}^{mn}(k_0 a) \frac{N_{mn}(0)}{N_{ml}(k_0 a)} z_l^{(3)}(k_0 r) \sum_{l'=m}^{\infty} i^{l'-n} d_{l'-m}^{mn}(k_s a) \frac{N_{mn}(0)}{N_{ml'}(k_s a)} z_{l'}^{(1)}(k_s r') \\
&\times 2\pi \sum_{k=-l}^l Y_l^k(\theta^{(L)}, \phi^{(L)}) \sum_{k'=-l'}^{l'} Y_{l'}^{k'*}(\theta'^{(L)}, \phi'^{(L)}) \langle D_{km}^l(\alpha, \beta, \gamma) D_{k'm}^{l'*}(\alpha, \beta, \gamma) \rangle \\
&= \frac{ik_s}{4\pi} \sum_{n=0}^{\infty} \sum_{m=0}^n (2 - \delta_{m,0}) C_{mn} \sum_{l=m}^{\infty} \frac{[i^{l-n} N_{mn}(0)]^2}{N_{ml}(k_0 a) N_{ml}(k_s a)} d_{l-m}^{mn}(k_0 a) d_{l-m}^{mn}(k_s a) \\
&\times z_l^{(3)}(k_0 r) z_l^{(1)}(k_s r') P_l(\cos\Theta). \tag{21}
\end{aligned}$$

where we have used Eq. (90) and Eq. (95). It is worth noting that $\langle G_{es}^{(22)} \rangle$ and $\langle G_{es}^{(12)} \rangle$ are independent of the exact directions $\hat{\mathbf{r}}$ and $\hat{\mathbf{r}}'$, but only dependent on the angle between them, namely Θ . This leads to a very important result, that $\langle UG_e \rangle$ is independent of the coordinate system we choose. Therefore, Eq. (20) and (21) has already given the averaging of one transition operator for a spheroidal particle centered at the origin and with random orientation. The averaging has no impact on the free space Green's function. It can still be calculated using Eq. (92) or $G_0^{-1}(\mathbf{p}) = p^2 - k_0^2$.

Using Eq. (83) and (84), Eq. (20) and (21) could be Fourier transformed into

$$\begin{aligned}
\langle G_{es}^{(22)}(\mathbf{p}, \mathbf{p}') \rangle &= \frac{ik_s}{4\pi} \sum_{n=0}^{\infty} \sum_{m=0}^n (2 - \delta_{m,0}) A_{mn} \sum_{l=m}^{\infty} \left[i^{l-n} d_{l-m}^{mn}(k_s a) \frac{N_{mn}(0)}{N_{ml}(k_s a)} \right]^2 P_l(\cos\Theta_p) \\
&\times 4\pi i^n \frac{pj_n(k_s A)j'_n(pA) - k_s j'_n(k_s A)j_n(pA)}{k_s^2 - p^2} A^2 \\
&\times 4\pi i^n \frac{p'j_n(k_s A)j'_n(p'A) - k_s j'_n(k_s A)j_n(p'A)}{k_s^2 - p'^2} A^2, \tag{22}
\end{aligned}$$

$$\begin{aligned}
\langle G_{es}^{(12)}(\mathbf{p}, \mathbf{p}') \rangle &= \frac{ik_s}{4\pi} \sum_{n=0}^{\infty} \sum_{m=0}^n (2 - \delta_{m,0}) C_{mn} \sum_{l=m}^{\infty} \frac{[i^{l-n} N_{mn}(0)]^2}{N_{ml}(k_0 a) N_{ml}(k_s a)} d_{l-m}^{mn}(k_0 a) d_{l-m}^{mn}(k_s a) P_l(\cos\Theta_p) \\
&\times 4\pi i^n \left[\frac{\pi}{2k_0^2} \delta(k_0 - p) - \frac{ph_n^{(1)}(k_0 A)j'_n(pA) - k_0 h_n^{(1)'}(k_0 A)j_n(pA)}{k_0^2 - p^2} A^2 \right] \\
&\times 4\pi i^n \frac{p'j_n(k_s A)j'_n(p'A) - k_s j'_n(k_s A)j_n(p'A)}{k_s^2 - p'^2} A^2, \tag{23}
\end{aligned}$$

where A is the half length of the major axis of the prolate spheroid, and the angle Θ_p is the angle between (θ_p, ϕ_p) and (θ'_p, ϕ'_p) , given by

$$\cos\Theta_p = \cos\theta_p \cos\theta'_p + \sin\theta_p \sin\theta'_p \cos(\phi_p - \phi'_p). \tag{24}$$

Using Eq. (8) and (15) we get

$$T_j(\mathbf{p}, \mathbf{p}') = k_0^2 \delta\epsilon_p G_0^{-1}(\mathbf{p}) \left[G_e^{(22)}(\mathbf{p}, \mathbf{p}') + G_e^{(12)}(\mathbf{p}, \mathbf{p}') \right]. \tag{25}$$

Multiplying Eq. (22) and (23) by $G_0^{-1}(\mathbf{p}) = p^2 - k_0^2$ and setting $p = k_0$ (we will show in Subsection 3.2 the reason why we set this), we get from Eq. (25) that

$$\begin{aligned}
\langle T_j(k_0 \hat{\mathbf{p}}, k_0 \hat{\mathbf{p}}') \rangle &= k_0^2 \delta\epsilon_p G_0^{-1}(k_0 \hat{\mathbf{p}}) \left[\langle G_e^{(22)}(k_0 \hat{\mathbf{p}}, k_0 \hat{\mathbf{p}}') \rangle + \langle G_e^{(12)}(k_0 \hat{\mathbf{p}}, k_0 \hat{\mathbf{p}}') \rangle \right] \\
&= k_0^2 \delta\epsilon_p G_0^{-1}(k_0 \hat{\mathbf{p}}) \langle G_{es}^{(12)}(k_0 \hat{\mathbf{p}}, k_0 \hat{\mathbf{p}}') \rangle, \tag{26}
\end{aligned}$$

where we have noticed that only $\langle G_{es}^{(12)}(\mathbf{p}, \mathbf{p}') \rangle$ has a term $p^2 - k_0^2$ in its denominator, and the function G_0 in Eq. (13) has a wavenumber k_s . Then the final result of the transition operator is

$$\begin{aligned}
\langle T_j(k_0 \hat{\mathbf{p}}, k_0 \hat{\mathbf{p}}') \rangle &= 4\pi \delta\epsilon_p k_0 k_s \sum_{n=0}^{\infty} \sum_{m=0}^n (2 - \delta_{m,0}) C_{mn} \\
&\times \sum_{l=m}^{\infty} (-1)^l N_{mn}^2(0) \frac{d_{l-m}^{mn}(k_0 a) d_{l-m}^{mn}(k_s a)}{N_{ml}(k_0 a) N_{ml}(k_s a)} P_l(\cos\Theta_p) \\
&\times \frac{k_0 j_n(k_s A)j'_n(k_0 A) - k_s j'_n(k_s A)j_n(k_0 A)}{k_s^2 - k_0^2} A^2, \tag{27}
\end{aligned}$$

where we have used the fact that [24]

$$j_n(x)h_n^{(1)'}(x) - j'_n(x)h_n^{(1)}(x) = \frac{i}{x^2}. \tag{28}$$

2.4 The product of a transition operator and its conjugate

After calculating $\langle T_j \rangle$, we notice that the product of a transition operator and its conjugate is also of vital importance. We will see in Subsection 3.1 that, this product plays a central role in the two-particle Green's function (the intensity propagator). In this subsection we will give the expression of this product, but the calculation of its averaging will be left to Subsection 3.3.

Let us now derive the transition operator in the reciprocal space in a spherical coordinate system (without being averaged over the Euler angles). To begin with, we calculate the scattering Green's function in the reciprocal space

$$\begin{aligned}
G_{es}^{(12)}(\mathbf{p}, \mathbf{p}') &= \frac{ik_s}{2\pi} \sum_{n=0}^{\infty} \sum_{m=0}^n (2 - \delta_{m,0}) C_{mn} e^{im(\phi_p - \phi'_p)} \\
&\times \sum_{l_1=m}^{\infty} i^{l_1-n} a_{l_1-m}^{mn}(k_0 a) \frac{N_{mn}(0)}{N_{ml_1}(k_0 a)} \bar{P}_{l_1}^m(\cos\theta_p) \sum_{l_2=m}^{\infty} i^{l_2-n} a_{l_2-m}^{mn}(k_s a) \frac{N_{mn}(0)}{N_{ml_2}(k_s a)} \bar{P}_{l_2}^m(\cos\theta'_p) \\
&\times 4\pi i^n \left[\frac{\pi}{2k_0^2} \delta(k_0 - p) - \frac{ph_n^{(1)}(k_0 A)j'_n(pA) - k_0 h_n^{(1)'}(k_0 A)j_n(pA)}{k_0^2 - p^2} A^2 \right] \\
&\times 4\pi i^n \frac{p' j_n(k_s A)j'_n(p' A) - k_s j'_n(k_s A)j_n(p' A)}{k_s^2 - p'^2} A^2, \tag{29}
\end{aligned}$$

as we have done in the calculation of Eq. (22). With the previous experience of Eq. (26), we here avoid to calculate $G_{es}^{(22)}$ since it will vanish in the transition operator in the far-field approximation. Now we multiply Eq. (29) by its conjugate to get

$$\begin{aligned}
|G_{es}^{(12)}(\mathbf{p}, \mathbf{p}')|^2 &= \frac{ik_s}{2\pi} \sum_{n=0}^{\infty} \sum_{m=0}^n (2 - \delta_{m,0}) C_{mn} e^{im(\phi_p - \phi'_p)} \\
&\times \sum_{l_1=m}^{\infty} i^{l_1-n} a_{l_1-m}^{mn}(k_0 a) \frac{N_{mn}(0)}{N_{ml_1}(k_0 a)} \bar{P}_{l_1}^m(\cos\theta_p) \\
&\times \sum_{l_2=m}^{\infty} i^{l_2-n} a_{l_2-m}^{mn}(k_s a) \frac{N_{mn}(0)}{N_{ml_2}(k_s a)} \bar{P}_{l_2}^m(\cos\theta'_p) \\
&\times 4\pi i^n \left[\frac{\pi}{2k_0^2} \delta(k_0 - p) - \frac{ph_n^{(1)}(k_0 A)j'_n(pA) - k_0 h_n^{(1)'}(k_0 A)j_n(pA)}{k_0^2 - p^2} A^2 \right] \\
&\times 4\pi i^n \frac{p' j_n(k_s A)j'_n(p' A) - k_s j'_n(k_s A)j_n(p' A)}{k_s^2 - p'^2} A^2 \\
&\times \frac{-ik_s}{2\pi} \sum_{n'=0}^{\infty} \sum_{m'=0}^{n'} (2 - \delta_{m',0}) C_{m'n'}^* e^{-im'(\phi_p - \phi'_p)} \\
&\times \sum_{l_3=m'}^{\infty} (-i)^{l_3-n'} a_{l_3-m'}^{m'n'}(k_0 a) \frac{N_{m'n'}(0)}{N_{m'l_3}(k_0 a)} \bar{P}_{l_3}^{m'}(\cos\theta_p) \\
&\times \sum_{l_4=m'}^{\infty} (-i)^{l_4-n'} a_{l_4-m'}^{m'n'}(k_s a) \frac{N_{m'n'}(0)}{N_{m'l_4}(k_s a)} \bar{P}_{l_4}^{m'}(\cos\theta'_p) \\
&\times 4\pi (-i)^{n'} \left[\frac{\pi}{2k_0^2} \delta(k_0 - p) - \frac{ph_{n'}^{(1)}(k_0 A)j'_{n'}(pA) - k_0 h_{n'}^{(1)'}(k_0 A)j_{n'}(pA)}{k_0^2 - p^2} A^2 \right] \\
&\times 4\pi (-i)^{n'} \frac{p' j_{n'}(k_s A)j'_{n'}(p' A) - k_s j'_{n'}(k_s A)j_{n'}(p' A)}{k_s^2 - p'^2} A^2. \tag{30}
\end{aligned}$$

Then we get the transition operator

$$\begin{aligned}
|T_j(k_0\hat{\mathbf{p}}, k_0\hat{\mathbf{p}}')|^2 &= k_0^4(\delta\epsilon_p)^2 \left| G_{es}^{(12)}(k_0\hat{\mathbf{p}}, k_0\hat{\mathbf{p}}') \right|^2 \\
&= k_0^4(\delta\epsilon_p)^2 \frac{ik_s}{2\pi} \sum_{n=0}^{\infty} \sum_{m=0}^n (2 - \delta_{m,0}) C_{mn} e^{im(\phi_p - \phi'_p)} \\
&\times \sum_{l_1=m}^{\infty} i^{l_1-n} d_{l_1-m}^{mn}(k_0a) \frac{N_{mn}(0)}{N_{ml_1}(k_0a)} \bar{P}_{l_1}^m(\cos\theta_p) \sum_{l_2=m}^{\infty} i^{l_2-n} d_{l_2-m}^{mn}(k_sa) \frac{N_{mn}(0)}{N_{ml_2}(k_sa)} \bar{P}_{l_2}^m(\cos\theta'_p) \\
&\times 4\pi i^n \frac{i}{k_0} \times 4\pi i^n \frac{p' j_n(k_sA) j'_n(k_0A) - k_s j'_n(k_sA) j_n(k_0A)}{k_s^2 - k_0^2} A^2 \\
&\times \frac{-ik_s}{2\pi} \sum_{n'=0}^{\infty} \sum_{m'=0}^{n'} (2 - \delta_{m',0}) C_{m'n'}^* e^{-im'(\phi_p - \phi'_p)} \\
&\times \sum_{l_3=m'}^{\infty} (-i)^{l_3-n'} d_{l_3-m'}^{m'n'}(k_0a) \frac{N_{m'n'}(0)}{N_{m'l_3}(k_0a)} \bar{P}_{l_3}^{m'}(\cos\theta_p) \sum_{l_4=m'}^{\infty} (-i)^{l_4-n'} d_{l_4-m'}^{m'n'}(k_sa) \frac{N_{m'n'}(0)}{N_{m'l_4}(k_sa)} \bar{P}_{l_4}^{m'}(\cos\theta'_p) \\
&\times 4\pi(-i)^{n'} \frac{-i}{k_0} \times 4\pi(-i)^{n'} \frac{k_0 j_{n'}(k_sA) j'_{n'}(k_0A) - k_s j'_{n'}(k_sA) j_{n'}(k_0A)}{k_s^2 - k_0^2} A^2, \tag{31}
\end{aligned}$$

The expression we get here is without being averaged. It is possible to perform this averaging as we do in Subsection 2.3, but obviously the expression will be complicated. The reason is that we are faced with four spherical harmonics and we must transform them into two spherical harmonics using Clebsch-Gordan coefficients if we want to apply orthogonality relation. Fortunately, when calculating the mean free paths, we can skip this procedure, which will be shown in Subsection 3.3.

3 The diffusion in small spheroidal particles group

3.1 Transition operator in multiple scattering theory

If we number all the particles in the scattering medium, we can get from Eq. (5) that

$$G = G_0 + \sum_{j=1}^N G_0 T_j G_0 + \sum_{j=1}^N \sum_{k=1, k \neq j}^N G_0 T_k G_0 T_j G_0 + \dots, \tag{32}$$

which is shown by Figure 2. N is the total number of particles in the medium. j and k mean the j th and k th particles, respectively. We change the notation of the left hand side because it here represents the multiple scattering process, and in Eq. (5) only one single particle is present. We have assumed that the light does not get into one particle twice, which corresponds to the so-called Twersky approximation [11]. Now we average Eq. (32) and get

$$\langle G(\mathbf{r}, \mathbf{r}_0) \rangle = G_0(\mathbf{r}, \mathbf{r}_0) + \int d\mathbf{r}_2 d\mathbf{r}_1 G_0(\mathbf{r}, \mathbf{r}_2) \Sigma(\mathbf{r}_2, \mathbf{r}_1) \langle G(\mathbf{r}_1, \mathbf{r}_0) \rangle, \tag{33}$$

where

$$\Sigma = \sum_{j=1}^N \langle T_j \rangle + \sum_{j=1}^N \sum_{k=1, k \neq j}^N \langle T_k G_0 T_j \rangle + \dots \tag{34}$$

is called the self-energy. Its second and higher terms represents those correlations between any two and more particles, and the first term means the uncorrelated part. When the particles are sparsely distributed in the medium, they hardly have impact on each other. Then we can naturally retain only the first term of Σ , which is called independent scattering. To make notations simpler, we use $\langle T \rangle$ to replace $\sum_{j=1}^N \langle T_j \rangle$, which leads to

$$\Sigma \simeq \sum_{j=1}^N \langle T_j \rangle = \langle T \rangle \tag{35}$$

and

$$\begin{aligned} \langle G(\mathbf{r}, \mathbf{r}_0) \rangle &= G_0(\mathbf{r}, \mathbf{r}_0) \\ &+ \int d\mathbf{r}_2 d\mathbf{r}_1 G_0(\mathbf{r}, \mathbf{r}_2) \langle T(\mathbf{r}_2, \mathbf{r}_1) \rangle \langle G(\mathbf{r}_1, \mathbf{r}_0) \rangle. \end{aligned} \quad (36)$$

We call $\langle T \rangle$ the average transition operator. It describes the statistical result of one single transition. Next, we can take the Fourier transform of Eq. (36) to get the Dyson equation

$$\langle G(\mathbf{p}) \rangle = \frac{1}{p^2 - k_0^2 - \Sigma(\mathbf{p})} = \frac{1}{p^2 - k_0^2 - \langle T(\mathbf{p}, \mathbf{p}) \rangle}. \quad (37)$$

Moreover, if the particles are monodisperse, namely identical, we get

$$\begin{aligned} \langle T(\mathbf{r}, \mathbf{r}') \rangle &= \frac{N}{V} \int d\mathbf{r}_j \int \frac{d\mathbf{p}}{(2\pi)^3} \frac{d\mathbf{p}'}{(2\pi)^3} e^{i\mathbf{p}\cdot(\mathbf{r}-\mathbf{r}_j)} T_j(\mathbf{p}, \mathbf{p}') e^{-i\mathbf{p}'\cdot(\mathbf{r}'-\mathbf{r}_j)} \\ &= \frac{N}{V} \int \frac{d\mathbf{p}}{(2\pi)^3} T_j(\mathbf{p}, \mathbf{p}) e^{i\mathbf{p}\cdot(\mathbf{r}-\mathbf{r}')}, \end{aligned} \quad (38)$$

where $N/V = n_j$ means that the number of particles are N in volume V . This leads to

$$\langle T(\mathbf{p}, \mathbf{p}') \rangle = n_j T_j(\mathbf{p}, \mathbf{p}) \delta(\mathbf{p}, \mathbf{p}'). \quad (39)$$

Substituting Eq. (27) into Eq. (37) gives the average Green's function in the spheroidal particles group.

Now we consider averaging the product of retarded and advanced Green's functions $\langle GG^* \rangle$, which is called the intensity propagator. The propagator $\langle GG^* \rangle$ satisfies the Bethe-Salpeter equation

$$\begin{aligned} \langle G(\mathbf{r}, \mathbf{r}_0) G^*(\mathbf{r}', \mathbf{r}'_0) \rangle &= \langle G(\mathbf{r}, \mathbf{r}_0) \rangle \langle G(\mathbf{r}', \mathbf{r}'_0) \rangle^* + \int d\mathbf{r}_2 d\mathbf{r}_1 d\mathbf{r}'_2 d\mathbf{r}'_1 \\ &\times \langle G(\mathbf{r}, \mathbf{r}_2) \rangle \langle G(\mathbf{r}', \mathbf{r}'_2) \rangle^* U(\mathbf{r}_2, \mathbf{r}_1; \mathbf{r}'_2, \mathbf{r}'_1) \langle G(\mathbf{r}_1, \mathbf{r}_0) G^*(\mathbf{r}'_1, \mathbf{r}'_0) \rangle. \end{aligned} \quad (40)$$

U is called the irreducible vertex, containing all the correlated scattering events. Just like the self-energy, U also have infinite terms, and higher-order terms can also be neglected when the particles are sparse. This is called the ladder approximation, given by

$$U \simeq \sum_{j=1}^N \langle T_j T_j^* \rangle = \langle TT^* \rangle. \quad (41)$$

Only one subscript j appears in each angular bracket, which means that the retarded and advanced Green's function will have correlations only when they follow the same scattering sequence. Performing Fourier transformation of Eq. (40) we get

$$\Phi_{\mathbf{p}, \mathbf{p}'}(\mathbf{q}) = (2\pi)^3 \langle G_{\mathbf{p}+} \rangle \langle G_{\mathbf{p}-} \rangle^* \delta_{\mathbf{p}, \mathbf{p}'} + \langle G_{\mathbf{p}+} \rangle \langle G_{\mathbf{p}-} \rangle^* \int \frac{d\mathbf{p}''}{(2\pi)^3} U_{\mathbf{p}, \mathbf{p}''}(\mathbf{q}) \Phi_{\mathbf{p}'', \mathbf{p}'}(\mathbf{q}), \quad (42)$$

where

$$\Phi_{\mathbf{p}, \mathbf{p}'}(\mathbf{q}) (2\pi)^3 \delta(-\mathbf{p}_1 + \mathbf{p}_2 + \mathbf{p}_3 - \mathbf{p}_4) = \langle G(\mathbf{p}_1, \mathbf{p}_2) G^*(\mathbf{p}_3, \mathbf{r}_4) \rangle \quad (43)$$

and

$$U_{\mathbf{p}, \mathbf{p}'}(\mathbf{q}) (2\pi)^3 \delta(-\mathbf{p}_1 + \mathbf{p}_2 + \mathbf{p}_3 - \mathbf{p}_4) = U(\mathbf{p}_1, \mathbf{p}_2; \mathbf{p}_3, \mathbf{r}_4) \quad (44)$$

imply the momentum conservation, and we have made the change of variables $\mathbf{p}_1 = \mathbf{p} + \mathbf{q}/2$, $\mathbf{p}_2 = \mathbf{p}' + \mathbf{q}/2$, $\mathbf{p}_3 = \mathbf{p} - \mathbf{q}/2$. Also,

$$\langle G_{\mathbf{p}+} \rangle = \left\langle G\left(\mathbf{p} + \frac{\mathbf{q}}{2}\right) \right\rangle, \quad \langle G_{\mathbf{p}-} \rangle = \left\langle G\left(\mathbf{p} - \frac{\mathbf{q}}{2}\right) \right\rangle. \quad (45)$$

Considering Eq. (38), (41) and (44), we get

$$U_{\mathbf{p}, \mathbf{p}'}(\mathbf{q}) \simeq n_j \langle T_j(\mathbf{p}+, \mathbf{p}'+) T_j^*(\mathbf{p}-, \mathbf{p}'-) \rangle. \quad (46)$$

3.2 The diffusion approximation and the far-field approximation

The diffusion approximation is

$$q \rightarrow 0 \quad (47)$$

in Eq. (42). Because q is the conjugate variable of propagation distance, this approximation means that the wave propagates for a much longer distance than the scale of dielectric inhomogeneities. In this case we can set $q = 0$ in $\langle G_{\mathbf{p}+} \rangle$, $\langle G_{\mathbf{p}-} \rangle$ and $U_{\mathbf{p},\mathbf{p}'}(\mathbf{q})$.

Moreover, the free space Green's function in the far-field region has the asymptotic form that

$$G_0(\mathbf{r} - \mathbf{r}') = \frac{\exp(ik_0|\mathbf{r} - \mathbf{r}'|)}{4\pi|\mathbf{r} - \mathbf{r}'|} \simeq \frac{\exp(ik_0r)}{4\pi r} \exp(-ik_0\hat{\mathbf{r}} \cdot \mathbf{r}'), \quad (48)$$

whose term $\exp(-ik_0\hat{\mathbf{r}} \cdot \mathbf{r}')$ could be considered as a Fourier transformation kernel. By using Eq. (5) and (9) we get

$$G_s(\mathbf{r}, \mathbf{r}') = \frac{\exp(ik_0r)}{4\pi r} T_j(k_0\hat{\mathbf{r}}, k_0\hat{\mathbf{r}}') \frac{\exp(ik_0r')}{4\pi r'}, \quad (49)$$

where $G_s = G_e - G_0$. Eq. (49) has shown that $p = p' = k_0$ holds in $T_j(\mathbf{p}, \mathbf{p}')$ for far-field region. This reveals that the particles are sparse and each of them is in the far-field regions of any other particles. In fact, there is another approximation called the on-shell approximation that also leads to $p = p' = k_0$ [12]. It tells us that, because

$$\langle G_{\mathbf{p}+} \rangle \langle G_{\mathbf{p}-} \rangle^* = \frac{\langle G_{\mathbf{p}+} \rangle - \langle G_{\mathbf{p}-} \rangle}{\langle G_{\mathbf{p}-}^{-1} \rangle - \langle G_{\mathbf{p}+}^{-1} \rangle} = \frac{\Delta G_{\mathbf{p}}(\mathbf{q})}{(\Sigma_{\mathbf{p}+} - \Sigma_{\mathbf{p}-}) - 2\mathbf{p} \cdot \mathbf{q}} \quad (50)$$

and Eq. (47), the Bethe-Salpeter Eq. (42) is limited to $\Delta G_{\mathbf{p}}(\mathbf{0}) = -\pi\delta(k_0^2 - p^2)$. Therefore, the on-shell approximation and the far-field approximation both result in

$$U_{\mathbf{p},\mathbf{p}'}(\mathbf{q}) \approx U_{k_0\hat{\mathbf{p}}, k_0\hat{\mathbf{p}}'}(\mathbf{0}) = n_j \left| T_j(k_0\hat{\mathbf{p}}, k_0\hat{\mathbf{p}}') \right|^2. \quad (51)$$

This is why we calculate Eq. (27) by setting $p = p' = k_0$.

3.3 The transport mean free path

The approaches to solving the Bethe-Salpeter equation in the diffusion approximation are widely discussed [9, 10, 25, 14, 26]. We will not repeat here but directly give the solution of Eq. (42), which is

$$\Phi(\mathbf{q}) = \frac{1}{D^*q^2}. \quad (52)$$

$D^* = v\ell^*/3$ is the diffusion constant, with the transport mean free path ℓ^* given by

$$\ell^* = \frac{\ell}{1-g}. \quad (53)$$

The scattering mean free path ℓ and the anisotropy factor g is given by [10]

$$\frac{1}{\ell} = \frac{n_j}{16\pi^2} \int \left| T_j(k_0\hat{\mathbf{p}}, k_0\hat{\mathbf{p}}') \right|^2 d\hat{\mathbf{p}}, \quad (54)$$

$$g = \frac{n_j\ell}{16\pi^2} \int (\hat{\mathbf{p}} \cdot \hat{\mathbf{p}}') \left| T_j(k_0\hat{\mathbf{p}}, k_0\hat{\mathbf{p}}') \right|^2 d\hat{\mathbf{p}}. \quad (55)$$

The mean free paths ℓ and ℓ^* are very important quantities, because they connect the single scattering process to the multiple scattering process.

Now let us consider how to calculate ℓ^{-1} and g . In the derivation of $\langle T_j \rangle$ (Subsection 2.3), we use Wigner D function to represent the rotation of the coordinate system, and perform averaging over

Euler angles by using the orthogonality of Wigner D function. When averaging over Euler angles, we keep the angle $\hat{\mathbf{p}} - \hat{\mathbf{p}}'$ unchanged¹, and retrieve all values of $(\hat{\mathbf{p}} + \hat{\mathbf{p}}')/2$. For now, if we want to calculate the mean free paths, we need to retrieve all values of the angle $\hat{\mathbf{p}} - \hat{\mathbf{p}}'$. As a matter of fact, we retrieve all values of $\hat{\mathbf{p}}$ and $\hat{\mathbf{p}}'$ in these two steps. This means that we can skip the averaging and directly calculate the integral over $\hat{\mathbf{p}}$ and $\hat{\mathbf{p}}'$, which makes the calculation much simpler. So we get

$$\begin{aligned} \frac{1}{\ell} &= \frac{n_j}{16\pi^2} \int \left\{ \frac{1}{4\pi} \int |T_j(k_0\hat{\mathbf{p}}, k_0\hat{\mathbf{p}}')|^2 d\hat{\mathbf{p}}' \right\} d\hat{\mathbf{p}} \\ &= \frac{n_j}{64\pi^3} \int |T_j(k_0\hat{\mathbf{p}}, k_0\hat{\mathbf{p}}')|^2 d\hat{\mathbf{p}} d\hat{\mathbf{p}}'. \end{aligned} \quad (56)$$

Substituting the expression of $|T_j|^2$, namely Eq. (31), we get

$$\begin{aligned} \frac{1}{\ell} &= 4\pi n_j k_0^2 (\delta\epsilon_p)^2 k_s^2 \sum_{n=0}^{\infty} \sum_{n'=0}^{\infty} \sum_{m=0}^{\min\{n, n'\}} (2 - \delta_{m,0})^2 C_{mn} C_{mn'}^* N_{mn}^2(0) N_{mn'}^2(0) \\ &\times \frac{k_0 j_n(k_s A) j_n'(k_0 A) - k_s j_n'(k_s A) j_n(k_0 A)}{k_s^2 - k_0^2} A^2 \frac{k_0 j_{n'}(k_s A) j_{n'}'(k_0 A) - k_s j_{n'}'(k_s A) j_{n'}(k_0 A)}{k_s^2 - k_0^2} A^2 \\ &\times \sum_{l_1=m}^{\infty} \sum_{l_2=m}^{\infty} \frac{d_{l_1-m}^{mn}(k_0 a)}{N_{ml_1}(k_0 a)} \frac{d_{l_2-m}^{mn}(k_s a)}{N_{ml_2}(k_s a)} \frac{d_{l_1-m}^{m'n'}(k_0 a)}{N_{ml_1}(k_0 a)} \frac{d_{l_2-m}^{m'n'}(k_s a)}{N_{ml_2}(k_s a)}. \end{aligned} \quad (57)$$

where we have used Eq. (89). And the result of anisotropy factor is

$$\begin{aligned} g &= \frac{n_j \ell}{64\pi^3} \int [\cos\theta_p \cos\theta_p' + \sin\theta_p \sin\theta_p' \cos(\phi_p - \phi_p')] |T_j(k_0\hat{\mathbf{p}}, k_0\hat{\mathbf{p}}')|^2 d\hat{\mathbf{p}} d\hat{\mathbf{p}}' \\ &= 4\pi n_j \ell k_0^2 (\delta\epsilon_p)^2 k_s^2 \sum_{n=0}^{\infty} \sum_{n'=0}^{\infty} \sum_{m=0}^n (2 - \delta_{m,0}) C_{mn} N_{mn}^2(0) \sum_{m'=0}^{n'} (2 - \delta_{m',0}) C_{m'n'}^* N_{m'n'}^2(0) \\ &\times \frac{k_0 j_n(k_s A) j_n'(k_0 A) - k_s j_n'(k_s A) j_n(k_0 A)}{k_s^2 - k_0^2} A^2 \frac{k_0 j_{n'}(k_s A) j_{n'}'(k_0 A) - k_s j_{n'}'(k_s A) j_{n'}(k_0 A)}{k_s^2 - k_0^2} A^2 \\ &\times \sum_{l_1=m}^{\infty} i^{l_1} \frac{d_{l_1-m}^{mn}(k_0 a)}{N_{ml_1}(k_0 a)} \sum_{l_2=m}^{\infty} i^{l_2} \frac{d_{l_2-m}^{mn}(k_s a)}{N_{ml_2}(k_s a)} \sum_{l_3=m'}^{\infty} (-i)^{l_3} \frac{d_{l_3-m'}^{m'n'}(k_0 a)}{N_{m'l_3}(k_0 a)} \sum_{l_4=m'}^{\infty} (-i)^{l_4} \frac{d_{l_4-m'}^{m'n'}(k_s a)}{N_{m'l_4}(k_s a)} \\ &\times \left[\delta_{m'm} f_{l_1, l_3}^{(=)} f_{l_2, l_4}^{(=)} + \frac{1}{2} \left(\delta_{m', m+1} f_{l_1, l_3}^{(+)} f_{l_2, l_4}^{(+)} + \delta_{m', m-1} f_{l_1, l_3}^{(-)} f_{l_2, l_4}^{(-)} \right) \right], \end{aligned} \quad (58)$$

where we have used Eq. (94) in the first equal-sign. In the second equal-sign, we first perform the integration over $(\phi_p + \phi_p')/2$ and $\phi_p - \phi_p'$, and then perform the integration over θ_p and θ_p' . The following formulae have also been used.

$$(2l+1)\sqrt{1-\cos^2\theta} P_l^m = P_{l-1}^{m+1} - P_{l+1}^{m+1}, \quad (59)$$

$$\int_{-1}^1 P_l^m P_{l'}^m dx = N_{ml}^2(0) \delta_{ll'}, \quad (60)$$

$$\begin{aligned} f_{l_2, l_4}^{(+)} &= \int_{-1}^1 \sin\theta_p' \bar{P}_{l_2}^m(\cos\theta_p') \bar{P}_{l_4}^{m+1}(\cos\theta_p') d\cos\theta_p' \\ &= \frac{1}{N_{ml_2}(0) N_{m+1, l_4}(0)} \int_0^\pi (1 - \cos^2\theta_p') \frac{P_{l_2-1}^{m+1}(\cos\theta_p') - P_{l_2+1}^{m+1}(\cos\theta_p')}{(2l_2+1)\sqrt{1-\cos^2\theta}} P_{l_4}^{m+1}(\cos\theta_p') d\theta_p' \\ &= \frac{1}{N_{ml_2}(0) N_{m+1, l_4}(0)} \int_{-1}^1 \frac{P_{l_2-1}^{m+1}(\cos\theta_p') - P_{l_2+1}^{m+1}(\cos\theta_p')}{(2l_2+1)} P_{l_4}^{m+1}(\cos\theta_p') d\cos\theta_p' \\ &= \frac{N_{m+1, l_4}^2(0) \delta_{l_2-1, l_4} - N_{m+1, l_4}^2(0) \delta_{l_2+1, l_4}}{(2l_2+1) N_{ml_2}(0) N_{m+1, l_4}(0)}, \end{aligned} \quad (61)$$

¹In fact, we do this in real space. Considering the Fourier kernel in Eq. (79) is $\exp(-i\mathbf{p} \cdot \mathbf{r})$, and this kernel is unaffected by the rotation, we deduce that the angles between (θ, ϕ) and (θ_p, ϕ_p) is invariant with the rotation, and the averaging in reciprocal space has the same form with the averaging in real space.

$$\begin{aligned}
f_{l_2, l_4}^{(-)} &= \int_{-1}^1 \sin\theta'_p \bar{P}_{l_2}^m(\cos\theta'_p) \bar{P}_{l_4}^{m-1}(\cos\theta'_p) d\cos\theta'_p \\
&= \frac{1}{N_{ml_2}(0)N_{m-1, l_4}(0)} \int_0^\pi (1 - \cos^2\theta'_p) P_{l_2}^m(\cos\theta'_p) \frac{P_{l_4-1}^m(\cos\theta'_p) - P_{l_4+1}^m(\cos\theta'_p)}{(2l_4+1)\sqrt{1-\cos^2\theta}} d\theta'_p \\
&= \frac{1}{N_{ml_2}(0)N_{m-1, l_4}(0)} \int_{-1}^1 P_{l_2}^m(\cos\theta'_p) \frac{P_{l_4-1}^m(\cos\theta'_p) - P_{l_4+1}^m(\cos\theta'_p)}{(2l_4+1)\sqrt{1-\cos^2\theta}} d\cos\theta'_p \\
&= \frac{N_{ml_2}^2(0)\delta_{l_2, l_4-1} - N_{ml_2}^2(0)\delta_{l_2, l_4+1}}{(2l_4+1)N_{ml_2}(0)N_{m-1, l_4}(0)}. \tag{62}
\end{aligned}$$

$$(2l+1)xP_l^m = (l+m)P_{l-1}^m + (l-m+1)P_{l+1}^m, \tag{63}$$

$$\begin{aligned}
f_{l_2, l_4}^{(=)} &= \int_{-1}^1 \cos\theta'_p \bar{P}_{l_2}^m(\cos\theta'_p) \bar{P}_{l_4}^m(\cos\theta'_p) d\cos\theta'_p \\
&= \frac{1}{N_{ml_2}(0)N_{m-1, l_4}(0)} \int_{-1}^1 x P_{l_2}^m(x) P_{l_4}^m(x) dx \\
&= \frac{1}{(2l_2+1)N_{ml_2}(0)N_{m-1, l_4}(0)} \int_{-1}^1 [(l_2+m)P_{l_2-1}^m(x) + (l_2-m+1)P_{l_2+1}^m(x)] P_{l_4}^m(x) dx \\
&= \frac{(l_2+m)N_{m, l_4}^2 \delta_{l_2-1, l_4} + (l_2-m+1)N_{m, l_4}^2 \delta_{l_2+1, l_4}}{(2l_2+1)N_{ml_2}(0)N_{m-1, l_4}(0)}. \tag{64}
\end{aligned}$$

Finally, Eq. (57) and (58) have given the expression of scattering mean free path and the anisotropy factor, respectively, for a group of monodisperse spheroidal particles.

4 Conclusion

Multiple scattering of coherent wave in non-spherical particles is an important issue because most small particles are not spherical in natural environment like seawater and atmosphere. In this manuscript we have used the expansion of scattering Green's function to derive the mean free paths concerning multiple scattering. First, we show how to get the average transition operator from the scattering Green's function for a group of dielectric spheroids with random orientations. The Green's function is expanded in a prolate spheroidal coordinate system, and then transformed into expansions of spherical wavefunctions. The averaging of a transition operator over Euler angles has been performed with the orthogonality of Wigner D function. In the second part, we have demonstrated the diffusion of scalar coherent waves in a group of spheroidal particles. We apply the diffusion approximation and far-field approximation to the irreducible vertex of the Bethe-Salpeter equation, to show the relation between the mean free paths and the transition operator. Having noticed the importance of the product of a transition operator and its conjugate, we show an alternative way to calculate the mean free paths from a transition operator without the averaging.

A Appendix: Spheroidal functions

The spheroidal radial function $R_{mn}^{(i)}(k_0a, \cosh\xi)$ satisfies [21]

$$\frac{d^2 R_{mn}^{(i)}}{d\xi^2} + \coth\xi \frac{dR_{mn}^{(i)}}{d\xi} + \left(k_0^2 a^2 \sinh^2\xi - n(n+1) - \frac{m^2}{\sinh^2\xi} \right) R_{mn}^{(i)} = 0, \quad (65)$$

and the spheroidal angular function $\mathcal{P}_n^m(k_0a, \cos\vartheta)$ satisfies

$$\frac{d^2 \mathcal{P}_n^m}{d\vartheta^2} + \cot\vartheta \frac{d\mathcal{P}_n^m}{d\vartheta} + \left(k_0^2 a^2 \sin^2\vartheta + n(n+1) - \frac{m^2}{\sin^2\vartheta} \right) \mathcal{P}_n^m = 0. \quad (66)$$

The normalized spheroidal angular function $\overline{\mathcal{P}}_n^m$ is given by

$$\overline{\mathcal{P}}_n^m(k_0a, \cos\vartheta) = N_{mn}^{-1}(k_0a) \mathcal{P}_n^m(k_0a, \cos\vartheta), \quad (67)$$

where the normalization factor is given by

$$N_{mn}(k_0a) = 2 \sum_{l=0,1}^{\infty}{}' \frac{(l+2m)!}{(2l+2m+1)!} [d_l^{mn}(k_0a)]^2. \quad (68)$$

Specially, [22]

$$N_{mn}(0) = \sqrt{\frac{2}{2n+1} \frac{(n+m)!}{(n-m)!}} \quad (69)$$

is the normalization factor of the associated Legendre function. In Eq. (68), the prime over the summation sign indicates that the summation is over only even values of l when $n-m$ is even, and over only odd values of l when $n-m$ is odd. The coefficient d_l^{mn} is determined by the following recursion formulae [27]

$$\alpha_l d_{l+2}^{mn}(k_0a) + (\beta_l - \lambda_{mn}(k_0a)) d_l^{mn}(k_0a) + \gamma_l d_{l-2}^{mn}(k_0a) = 0, \quad (70)$$

where

$$\alpha_l = \frac{(2m+l+2)(2m+l+1)}{(2m+2l+3)(2m+2l+5)} (k_0a)^2, \quad (71)$$

$$\beta_l = (m+l)(m+l+1) + \frac{2(m+l)(m+l+1) - 2m^2 - 1}{(2m+2l-1)(2m+2l+3)} (k_0a)^2, \quad (72)$$

$$\gamma_l = \frac{l(l-1)}{(2m+2l-3)(2m+2l-1)} (k_0a)^2. \quad (73)$$

B Appendix: Coefficients A_{mn} and C_{mn}

The boundary condition of scalar Green's function for a dielectric spheroid could be expressed by [10, 7]

$$\begin{aligned} G_e^{(22)}(\mathbf{r}, \mathbf{r}') &= G_e^{(12)}(\mathbf{r}, \mathbf{r}'), \\ \nabla G_e^{(22)}(\mathbf{r}, \mathbf{r}') \cdot \hat{\mathbf{n}} &= \nabla G_e^{(12)}(\mathbf{r}, \mathbf{r}') \cdot \hat{\mathbf{n}}, \end{aligned} \quad (74)$$

when $\xi = \xi_0$. This boundary condition means that the limiting values of the scalar Green's function as \mathbf{r} approaches the interface from region 1 and from region 2 is equal. Applying Eq. (74) to Eq. (13) yields

$$R_{mn}^{(3)}(k_s a, \cosh\xi_0) + A_{mn} R_{mn}^{(1)}(k_s a, \cosh\xi_0) = C_{mn} R_{mn}^{(3)}(k_0 a, \cosh\xi_0), \quad (75)$$

$$\frac{R_{mn}^{(3)'}(k_s a, \cosh \xi_0)}{a \sqrt{\sinh^2 \xi_0 + \sin^2 \vartheta}} + A_{mn} \frac{R_{mn}^{(1)'}(k_s a, \cosh \xi_0)}{a \sqrt{\sinh^2 \xi_0 + \sin^2 \vartheta}} = C_{mn} \frac{R_{mn}^{(3)'}(k_0 a, \cosh \xi_0)}{a \sqrt{\sinh^2 \xi_0 + \sin^2 \vartheta}}, \quad (76)$$

where the prime indicates derivative with respect to ξ . The denominators in Eq. (76) cancel because they have the same ϑ . Thus, we get the coefficients

$$A_{mn} = \frac{R_{mn}^{(3)'}(k_s a)_{\xi_0} R_{mn}^{(3)}(k_0 a)_{\xi_0} - R_{mn}^{(3)'}(k_0 a)_{\xi_0} R_{mn}^{(3)}(k_s a)_{\xi_0}}{R_{mn}^{(3)'}(k_0 a)_{\xi_0} R_{mn}^{(1)}(k_s a)_{\xi_0} - R_{mn}^{(1)'}(k_s a)_{\xi_0} R_{mn}^{(3)}(k_0 a)_{\xi_0}}, \quad (77)$$

$$C_{mn} = \frac{R_{mn}^{(3)'}(k_0 a)_{\xi_0} R_{mn}^{(3)}(k_s a)_{\xi_0} - R_{mn}^{(3)'}(k_s a)_{\xi_0} R_{mn}^{(3)}(k_0 a)_{\xi_0}}{R_{mn}^{(1)'}(k_s a)_{\xi_0} R_{mn}^{(3)}(k_s a)_{\xi_0} - R_{mn}^{(3)'}(k_s a)_{\xi_0} R_{mn}^{(1)}(k_s a)_{\xi_0}}, \quad (78)$$

where the radial function $R_{mn}^{(i)}(k_0 a)_{\xi_0} = R_{mn}^{(i)}(k_0 a, \cosh \xi_0)$.

C Appendix: Fourier transformation of the scalar spherical wave functions

Stratton has given the formula (see Page 410 of Ref. [28])

$$j_n(pr) P_n^m(\cos \theta_p) e^{im\phi_p} = \frac{i^{-n}}{4\pi} \int_0^\pi (\sin \theta) d\theta \int_{-\pi}^\pi d\phi [e^{-ipr \cos \Theta} P_n^m(\cos \theta) e^{im\phi}], \quad (79)$$

where

$$\cos \Theta = \cos \theta_p \cos \theta + \sin \theta_p \sin \theta \cos(\phi_p - \phi). \quad (80)$$

Eq. (79) shows the Fourier transformation of a spherical harmonics by setting Θ as the angle between the real-space vector $\mathbf{r} = (r, \theta, \phi)$ and the Fourier-space vector $\mathbf{p} = (p, \theta_p, \phi_p)$. Moreover, using the following integral formulae [29, 24]

$$\int r^2 z_n^{(i_1)}(kr) z_n^{(i_2)'}(pr) dr = \frac{p z_n^{(i_1)}(kr) z_n^{(i_2)'}(pr) - k z_n^{(i_1)'}(kr) z_n^{(i_2)}(pr)}{k^2 - p^2} r^2, \quad (81)$$

$$\int_0^\infty r^2 z_n^{(i_1)}(kr) z_n^{(i_2)}(pr) dr = \frac{\pi \delta(k - p)}{2k^2}, \quad (82)$$

we get

$$\begin{aligned} & \int_0^\infty r^2 dr \int_{-1}^1 d\cos \theta \int_\pi^\pi d\phi [e^{i\mathbf{p} \cdot \mathbf{r}} z_n(k_0 r) P_n^m(\cos \theta) e^{im\phi}] \\ &= 4\pi i^n P_n^m(\cos \theta_p) e^{im\phi_p} \int_0^\infty r^2 z_n(k_0 r) j_n(pr) dr \\ &= 4\pi i^n P_n^m(\cos \theta_p) e^{im\phi_p} \frac{\pi}{2k_0^2} \delta(k_0 - p), \end{aligned} \quad (83)$$

$$\begin{aligned} & \int_0^a r^2 dr \int_{-1}^1 d\cos \theta \int_\pi^\pi d\phi [e^{i\mathbf{p} \cdot \mathbf{r}} z_n(k_0 r) P_n^m(\cos \theta) e^{im\phi}] \\ &= 4\pi i^n P_n^m(\cos \theta_p) e^{im\phi_p} \int_0^a r^2 z_n(k_0 r) j_n(pr) dr \\ &= 4\pi i^n P_n^m(\cos \theta_p) e^{im\phi_p} \frac{p z_n(k_0 a) j_n'(pa) - k_0 z_n'(k_0 a) j_n(pa)}{k_0^2 - p^2} a^2. \end{aligned} \quad (84)$$

D Appendix: Calculation of $\left\langle D_{m'k'}^{l'} * D_{mk}^l \right\rangle_{\alpha, \beta, \gamma}$

In this appendix we show the calculation of $\left\langle D_{m'k'}^{l'} * D_{mk}^l \right\rangle_{\alpha, \beta, \gamma}$ (for more details please see Ref. [4]). For any function $f(\alpha, \beta, \gamma)$, its average over Euler angles is expressed by

$$\langle f(\alpha, \beta, \gamma) \rangle_{\alpha, \beta, \gamma} = \int_0^{2\pi} d\alpha \int_0^\pi \sin\beta d\beta \int_0^{2\pi} d\gamma f(\alpha, \beta, \gamma) p(\alpha, \beta, \gamma). \quad (85)$$

$p(\alpha, \beta, \gamma)$ is the probability density function of α, β, γ . We assume the orientation of each particle is totally random, and namely the Euler angles are uniformly distributed, which means that

$$p(\alpha, \beta, \gamma) = \frac{1}{8\pi^2}. \quad (86)$$

The definition of Wigner D function

$$D_{mk}^l(\alpha, \beta, \gamma) = e^{-im\alpha} d_{mk}^l(\beta) e^{-ik\gamma} \quad (87)$$

where the Wigner d function satisfies

$$\int_0^\pi \sin\beta d\beta d_{mk}^l(\beta) d_{m'k'}^{l'}(\beta) = \frac{2}{2l+1} \delta_{l'l}. \quad (88)$$

Considering

$$\int_0^{2\pi} d\alpha \exp[-i(m-m')\alpha] = 2\pi \delta_{m'm} \quad (89)$$

and Eq. (85), (87), and (88), we get

$$\begin{aligned} & \left\langle D_{m'k'}^{l'} * (\alpha, \beta, \gamma) D_{mk}^l(\alpha, \beta, \gamma) \right\rangle_{\alpha, \beta, \gamma} \\ &= \frac{1}{8\pi^2} \int_0^{2\pi} d\alpha e^{im'\alpha} e^{-im\alpha} \int_0^{2\pi} d\gamma e^{ik'\gamma} e^{-ik\gamma} \int_0^\pi \sin\beta d\beta d_{mk}^l(\beta) d_{m'k'}^{l'}(\beta) \\ &= \frac{\delta_{m'm} \delta_{k'k} \delta_{l'l}}{2l+1}. \end{aligned} \quad (90)$$

E Appendix: The expansion of G_0

Eq. (12) reads

$$\begin{aligned} G_0(\mathbf{r}, \mathbf{r}') &= \frac{ik_0}{2\pi} \sum_{m,n} (2 - \delta_{m,0}) \overline{\mathcal{P}}_n^m(k_0 a, \cos\vartheta) \overline{\mathcal{P}}_n^m(k_0 a, \cos\vartheta') \\ &\quad \times R_{mn}^{(3)}(k_0 a, \cosh\xi^{\geq}) R_{mn}^{(1)}(k_0 a, \cosh\xi^{\leq}) e^{im(\varphi - \varphi')}, \end{aligned} \quad (91)$$

and could also be expanded using spherical eigenfunctions into [7]

$$\begin{aligned} G_0(\mathbf{r}, \mathbf{r}') &= \frac{ik_0}{2\pi} \sum_{m,n} (2 - \delta_{m,0}) \overline{P}_n^m(\cos\theta) \overline{P}_n^m(\cos\theta') \\ &\quad \times z_n^{(3)}(k_0 r^{\geq}) z_n^{(1)}(k_0 r^{\leq}) e^{im(\phi - \phi')}, \end{aligned} \quad (92)$$

where $\overline{\mathcal{P}}_n^m(k_0 a, \cos\vartheta) = N_{mn}^{-1}(k_0 a) \mathcal{P}_n^m(k_0 a, \cos\vartheta)$, and $\overline{P}_l^m(\cos\theta) = N_{ml}^{-1}(0) P_l^m(\cos\theta)$. Moreover, $z_n^{(3)}(x) = h_n^{(1)}(x)$ is the spherical Bessel function of the third type, also called the spherical Hankel function, and $z_n^{(1)}(x) = j_n(x)$ is the spherical Bessel function of the first type.

By using the addition theorem of Legendre function [24]

$$P_n(\cos\Theta) = \frac{2}{2n+1} \sum_{m=-n}^n \bar{P}_n^m(\cos\theta) \bar{P}_n^m(\cos\theta') e^{im(\phi-\phi')} \quad (93)$$

where

$$\cos\Theta = \cos\theta\cos\theta' + \sin\theta\sin\theta'\cos(\phi - \phi'), \quad (94)$$

and $\sqrt{2\pi}Y_l^m(\theta, \phi) = \bar{P}_l^m(\cos\theta)e^{im\phi}$, we get

$$2\pi \sum_{k=-l}^l Y_l^k(\theta, \phi) Y_l^{k*}(\theta', \phi') = \frac{2n+1}{2} P_n(\cos\Theta). \quad (95)$$

References

- [1] Bohren C F and Huffman D R. *Absorption and scattering of light by small particles*. John Wiley & Sons, 2008.
- [2] W. J. Wiscombe. Improved mie scattering algorithms. *Appl. Opt.*, 19(9):1505–1509, May 1980.
- [3] M. I. Mishchenko, J. W. Hovenier, and L. D. Travis. *Light Scattering by Nonspherical Particles: Theory, Measurements, and Applications*. IOP Publishing, 2000.
- [4] M. I. Mishchenko, L. D. Travis, and A. A. Lacis. *Scattering, absorption, and emission of light by small particles*. Cambridge University Press, 2002.
- [5] M. I. Mishchenko. *Electromagnetic scattering by particles and particle groups: an introduction*. Cambridge University Press, 2014.
- [6] Victor Farafonov. *Application of non-orthogonal bases in the theory of light scattering by spheroidal particles*, pages 189–266. Springer Berlin Heidelberg, Berlin, Heidelberg, 2013.
- [7] Tai C T. *Dyadic Green's functions in electromagnetic theory*. IEEE Press, 1971.
- [8] Li L W, Kang X K, and Leong M S. *Spheroidal wave functions in electromagnetic theory*. John Wiley & Sons, 2004.
- [9] E. Akkermans and G. Montambaux. *Mesoscopic physics of electrons and photons*. Cambridge University Press, 2007.
- [10] R. Carminati and J. C. Schotland. *Principles of Scattering and Transport of Light*. Cambridge University Press, 2021.
- [11] M. I. Mishchenko, L. D. Travis, and A. A. Lacis. *Multiple scattering of light by particles: radiative transfer and coherent backscattering*. Cambridge University Press, 2006.
- [12] M Piraud, L Pezzé, and L Sanchez-Palencia. Quantum transport of atomic matter waves in anisotropic two-dimensional and three-dimensional disorder. *New Journal of Physics*, 15(7):075007, jul 2013.
- [13] D. Vollhardt and P. Wölfle. Diagrammatic, self-consistent treatment of the anderson localization problem in $d \leq 2$ dimensions. *Phys. Rev. B*, 22:4666–4679, Nov 1980.
- [14] Nicolas Cherroret, Dominique Delande, and Bart A. van Tiggelen. Induced dipole-dipole interactions in light diffusion from point dipoles. *Phys. Rev. A*, 94:012702, Jul 2016.
- [15] Kevin Vynck, Romain Pierrat, and Rémi Carminati. Multiple scattering of polarized light in disordered media exhibiting short-range structural correlations. *Phys. Rev. A*, 94:033851, Sep 2016.

- [16] E Amic, J M Luck, and Th M Nieuwenhuizen. Anisotropic multiple scattering in diffusive media. *Journal of Physics A: Mathematical and General*, 29(16):4915, aug 1996.
- [17] P. Sheng. *Introduction to Wave Scattering, Localization and Mesoscopic Phenomena*. Springer, 2006.
- [18] M. C. W. van Rossum and Th. M. Nieuwenhuizen. Multiple scattering of classical waves: microscopy, mesoscopy, and diffusion. *Rev. Mod. Phys.*, 71:313–371, Jan 1999.
- [19] L. Tsang and J. A. Kong. Multiple scattering of electromagnetic waves by random distributions of discrete scatterers with coherent potential and quantum mechanical formalism. *Journal of Applied Physics*, 51(7):3465–3485, 07 1980.
- [20] L. Tsang and J. A. Kong. Multiple scattering of acoustic waves by random distributions of discrete scatterers with the use of quasicrystalline-coherent potential approximation. *Journal of Applied Physics*, 52(9):5448–5458, 09 1981.
- [21] Le-Wei Li, Mook-Seng Leong, Pang-Shyan Kooi, and Tat-Soon Yeo. Spheroidal vector wave eigenfunction expansion of dyadic green’s functions for a dielectric spheroid. *IEEE Transactions on Antennas and Propagation*, 49(4):645–659, 2001.
- [22] V. G Farafonov, N. V. Voshchinnikov, and E. G. Semenova. Some relations between the spheroidal and spherical wave functions. *Journal of Mathematical Sciences**Journal of Mathematical Sciences*, 214:382–391, 2016.
- [23] D A Varshalovich, A N Moskalev, and V K Khersonskii. *Quantum Theory of Angular Momentum*. WORLD SCIENTIFIC, 1988.
- [24] Z X Wang and D R Guo. *Special Functions*. WORLD SCIENTIFIC, 1989.
- [25] Jorgen Rammer. *Quantum Transport Theory*. CRC Press, 2004.
- [26] Ad Lagendijk and Bart A. van Tiggelen. Resonant multiple scattering of light. *Physics Reports*, 270(3):143–215, 1996.
- [27] Ross Adelman, Nail A. Gumerov, and Ramani Duraiswami. Software for computing the spheroidal wave functions using arbitrary precision arithmetic, 2014.
- [28] Adams Julius Stratton. *Electromagnetic Theory*. Mcgraw Hill Book Company, 1941.
- [29] Gradshteyn I S and Ryzhik I M. *Table of integrals, series, and products*. Academic press, 2014.