

THE RANKING-SELBERG INTEGRAL ON $\mathrm{GSp}(2)$ FOR SQUARE FREE LEVELS

SEIJI KUGA AND MASAO TSUZUKI

ABSTRACT. We explicitly compute the Rankin-Selberg type integral introduced by Piatetski-Shapiro over adeles for vector-valued Siegel cusp forms of square-free levels $\Gamma_0(N)$. On the way, for particular test functions in the Bessel models of irreducible admissible representations, exact evaluations of the local zeta-integrals are given.

1. INTRODUCTION

Lately, the Bessel periods on the symplectic group $\mathbf{G} := \mathbf{GSp}_2$ of rank 2 progressively gain its importance in the arithmetic of Siegel modular forms. The notion of Bessel period and the associated local model of admissible representations of \mathbf{G} were originally introduced by Novodvorski and Piatetski-Shapiro in their work ([26], [27], [30]), where a modern treatment of Andrianov's integral representation ([1], [2]) of the spinor L -function for Siegel modular forms was developed. Related to this, we should also mention independent work by Sugano ([43]), which handles vector-valued Hilbert-Siegel cusp forms on arithmetic groups defined by maximal lattices of indefinite division algebras. The Bessel period also plays an essential role in a formulation of Böcherer's conjecture posed by Liu [23] in a style of the refined Gan-Gross-Prasad conjecture. This version of Böcherer's conjecture has been completely solved by Furusawa and Morimoto ([10], [11]).

In this paper, we compute the Rankin-Selberg type integral à la Piatetski-Shapiro for holomorphic vector-valued Siegel cusp forms of square-free levels explicitly, using results by Pitale-Schmidt [32] on local new vectors on $\mathbf{G}(\mathbb{Q}_p)$ ($p < \infty$); since we rely on the local-global method, our result is immediately applied to non-holomorphic Siegel cusp forms of discrete series type (cf. [20], [21]). We explain the moral of our investigation for scalar-valued Siegel modular forms of weight l . Our focus is an asymptotic formula as $N + l \rightarrow \infty$ of the average

$$(1.1) \quad \frac{1}{[\mathbf{Sp}_2(\mathbb{Z}) : \Gamma_0(N)]} \sum_{\varphi \in \mathcal{B}(l, N)} \langle E(s, \Lambda, \mu) \mid \varphi \rangle B^\Lambda(\varphi),$$

where $\mathbf{G}^\#$ is a subgroup of \mathbf{G} defined by a quadratic field E of discriminant $D < 0$ (see §2.1), $E(s, \Lambda, \mu)$ is the Eisenstein series on $\mathbf{G}^\#(\mathbb{A})$ attached to an ideal class character Λ of E , a character $\mu : \mathbb{A}^\times / \mathbb{Q}^\times \mathbb{R}^\times \rightarrow \mathbb{C}^1$ and $s \in \mathbb{C}$ (see §3.1), $B^\Lambda(\varphi)$ is the Bessel period of φ in an orthonormal basis $\mathcal{B}(l, N)$ of Siegel cusp forms of weight l and square free level $\Gamma_0(N)$, and $\langle E(s, \Lambda, \mu) \mid \varphi \rangle$ is the L^2 -inner product of $E(s, \Lambda, \mu)$ and $\varphi|_{\mathbf{G}^\#(\mathbb{A})}$. The conductor of μ is assumed to be relatively prime to $2DN$. Actually, for application ([19]), we consider a more general average than (1.1) by requiring $\mathcal{B}(l, N)$ to consist of

2020 *Mathematics Subject Classification.* primary 11F46, secondary 11F67.

joint eigenforms by Hecke operators outside NM and by putting the eigenvalue of such a Hecke operator at each φ in the formation of the sum (1.1). With such a generality, we obtain a completely explicit expression of (1.1) in terms of

- The spinor L -function $L(s, \pi, \mu)$ (see §4.6) attached to an irreducible cuspidal representation $\pi \cong \otimes_p \pi_p$ twisted by μ .
- An average of Fourier coefficient of a newform φ_π^0 in π , denoted by $R(\varphi_\pi^0, E, \Lambda)$ (see (4.12)) introduced in [17]).
- An average of local Bessel periods of π_p (see §5).

For the precise statement including the vector-valued case, we refer to Proposition 5.1 and Theorem 5.2. Owing to the explicit nature of our result, we are able to show that, asymptotically as the prime level N grows with the weight l being fixed, the contribution to (1.1) of the old forms, the Yoshida lifts and the Saito-Kurokawa lifts are 0 (see Theorem 5.4). These results are necessary in our forthcoming paper [19], where we shall work out the geometric side of a relative trace formula whose spectral side is essentially (1.1). We should remark that an average similar to our computed average (5.9) without the factors $\widehat{L}(s + 1/2, \pi, \mu)$ and $\widehat{f}_S(\pi)$ is studied in [7, §3]. When $E = \mathbb{Q}(i)$, our average (5.9) is almost the same as the one in [5, §8] except the appearance of the local periods.

Let us explain the structure of this paper. In §2, we introduce basic objects related to algebraic subgroups of the symplectic group G and Haar measures on their p -adic, real and adelic points. In §3.1, the Rankin-Selberg type integral and the global Bessel function for Siegel cusp forms are recalled; after this preparation, the basic identity (Lemma 3.2) is stated, which computes the pairing $\langle E(s, \Lambda, \mu), \varphi \rangle$ in (1.1) in terms of the Mellin transform of the global Bessel function attached to φ . We review the proof ([30]) to determine a constant depending on Haar measures exactly. In §4, we recall the definition of the quantity $R(\varphi, E, \Lambda)$ and describe its behavior under the Galois conjugation $\Lambda \mapsto \Lambda^\dagger$ (Lemma 4.6). The local multiplicity one theorem for Bessel models of π_p 's allows us to split $B^\Lambda(\varphi)$ for $\varphi \in \pi$ corresponding to a pure tensor in $\otimes_p \pi_p$ to an Euler product of local Bessel functions for π_p up to a constant. To define our global newform φ_π^0 and identify this constant with $R(\varphi_\pi^0, E, \Lambda)$ (see Lemma 4.7), we fix pairs of a local Bessel functional ℓ_p on π_p and a local new vector ξ_p of π_p normalized by $\ell_p(\xi_p) = 1$; in §4.4, we explain that the main result of [32] ensures the existence of such $\{(\ell_p, \xi_p)\}_p$ when π corresponds to a new forms on $\Gamma_0(N)$. In §6, we compute the local zeta-integrals by using results of [32]. The unramified computations are known as recalled in §6.2. Other cases, as well as the explicit determination of the local periods of prime level (see §6.8), seem to be new, so that some details are given in §6.5, §6.6, and §6.4. In §4.6, the functional equation of the spinor L -function is deduced from the basic identity by using Lemma 4.6. Although a “nice” functional equation of the spinor L -function itself is known in a broader generality ([40, Lemma 1.2]) owing to Arthur’s classification of automorphic representations of $G(\mathbb{A})$, we include this section for the sake of completeness and to provide a proof free from Arthur’s result, confirming that the local L -factor and the local ε -factor defined by [30] (in terms of a particular Bessel model) coincides with the ones proposed in [37, Tables 2,3]. In section §5, we prove Proposition 5.1 and Theorem 5.2; by invoking a result by Furusawa-Morimoto ([10, Theorem 8.1]), we obtain an explicit expression of (1.1) solely in terms of L -functions (Theorem 5.3).

2. PRELIMINARIES

Let \mathbf{V} denotes the space of symmetric matrices of degree 2; we consider \mathbf{V} as a \mathbb{Z} -scheme by defining

$$\mathbf{V}(R) = \left\{ \begin{bmatrix} a & b \\ b & c \end{bmatrix} \mid a, b, c \in R \right\}$$

for any commutative ring R . The group $\mathbf{GL}_2(R)$ acts on this space from the right as

$$(2.1) \quad \mathbf{V}(R) \times \mathbf{GL}_2(R) \ni (T, h) \longmapsto Ts(h) := \frac{1}{\det h} {}^t h T h \in \mathbf{V}(R).$$

Let \mathbf{G} denote the symplectic similitude group

$$\mathbf{G} := \{g \in \mathbf{GL}_4 \mid {}^t g \begin{bmatrix} O & 1_2 \\ -1_2 & O \end{bmatrix} g = \nu(g) \begin{bmatrix} O & 1_2 \\ -1_2 & O \end{bmatrix} \quad (\exists \nu(g) \in \mathbf{GL}_1)\}$$

(usually denoted by \mathbf{GSp}_2), where $\nu : \mathbf{G} \rightarrow \mathbf{GL}_1$ is the similitude character. The center of \mathbf{G} is denoted by \mathbf{Z} , which coincides with the set of all the scalar matrices in \mathbf{G} . Let $\mathbf{P} = \mathbf{M}\mathbf{N}$ be the Siegel parabolic subgroup of \mathbf{G} , where

$$\mathbf{M} := \{m(A, c) := \begin{bmatrix} A & O \\ O & {}^t A^{-1} c \end{bmatrix} \mid A \in \mathbf{GL}_2, c \in \mathbf{GL}_1\},$$

$$\mathbf{N} := \{n(X) := \begin{bmatrix} 1_2 & X \\ O & 1_2 \end{bmatrix} \mid X \in \mathbf{V}\}.$$

Note that $\nu(m(A, c)) = c$ and $\nu(n(X)) = 1$. Let

$$\mathbf{T} := \{m(\begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix}, c) \mid a, b, c \in \mathbf{GL}_1\}.$$

The Weyl group $W := N_{\mathbf{G}}(\mathbf{T})/\mathbf{T}$ of \mathbf{G} is generated by the images of

$$s_1 := \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}, \quad s_2 := \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

2.1. Embedding of groups. Let $E = \mathbb{Q}(\sqrt{D})$ be an imaginary quadratic field of discriminant $D < 0$; we realize E as a subfield of \mathbb{C} so that $\sqrt{D} = i|D|^{1/2}$ with i being the imaginary unit. Let \mathfrak{o}_E denote the ring of integers of E . Choose a \mathbb{Z} -basis $\{1, \theta\}$ of \mathfrak{o}_E so that

$$(2.2) \quad \theta - \bar{\theta} = -\sqrt{D},$$

where $\tau \mapsto \bar{\tau}$ denotes the non-trivial automorphism of E/\mathbb{Q} .

The different of E/\mathbb{Q} , defined as $\mathfrak{d}_{E/\mathbb{Q}}^{-1} := \{\tau \in E \mid \mathrm{tr}_{E/\mathbb{Q}}(\tau \mathfrak{o}_E) \subset \mathbb{Z}\}$, is $\frac{1}{\sqrt{D}} \mathfrak{o}_E$; thus a symplectic \mathbb{Z} -bi-linear form $\langle \cdot, \cdot \rangle : \mathfrak{o}_E^2 \times \mathfrak{o}_E^2 \rightarrow \mathbb{Z}$ is defined by

$$\langle x, y \rangle := \mathrm{tr}_{E/\mathbb{Q}} \left(\frac{-1}{\sqrt{D}} \det \begin{bmatrix} x_1 & y_1 \\ x_2 & y_2 \end{bmatrix} \right), \quad x = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}, y = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} \in \mathfrak{o}_E^2.$$

Note that the following vectors of \mathfrak{o}_E^2 form a symplectic \mathbb{Z} -basis, i.e., $\langle v_i^+, v_j^- \rangle = \delta_{ij}$ and $\langle v_i^+, v_j^+ \rangle = \langle v_i^-, v_j^- \rangle = 0$ for all i, j :

$$v_1^+ := \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad v_2^+ := \begin{bmatrix} \theta \\ 0 \end{bmatrix}, \quad v_1^- := \begin{bmatrix} 0 \\ -\bar{\theta} \end{bmatrix}, \quad v_2^- := \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

For any ring R , set

$$\mathbf{G}^\#(R) := \{h \in \mathbf{GL}_2(\mathfrak{o}_E \otimes_{\mathbb{Z}} R) \mid \det(h) \in R^\times\}.$$

For $h \in \mathbf{G}^\#(R)$, viewed as an R -linear automorphism of $(\mathfrak{o}_E \otimes_{\mathbb{Z}} R)^2$, let $\iota_\theta(h)$ denote the 4×4 -matrix representing h by the R -basis $\{v_1^+, v_2^+, v_1^-, v_2^-\}$ of $(\mathfrak{o}_E \otimes_{\mathbb{Z}} R)^2$, i.e.,

$$(2.3) \quad [h(v_1^+), h(v_2^+), h(v_1^-), h(v_2^-)] = [v_1^+, v_2^+, v_1^-, v_2^-] \iota_\theta(h).$$

Then $\iota_\theta(h) \in \mathbf{GL}_4(R)$ satisfies $\langle \iota_\theta(h)(x), \iota_\theta(h)(y) \rangle = \det(h) \langle x, y \rangle$ for all $x, y \in \mathfrak{o}_E^2$, i.e., $\iota_\theta(h)$ belongs to $\mathbf{G}(R)$ and $\nu(\iota_\theta(h)) = \det h$. Thus, we have an embedding

$$\iota_\theta : \mathbf{G}^\#(R) \longrightarrow \mathbf{G}(R).$$

Let $\mathbf{B}^\#$ be the Borel subgroup of $\mathbf{G}^\#$ consisting of all those upper-triangular matrices in $\mathbf{G}^\#$; then, for any \mathbb{Q} -algebra R , the set $\mathbf{B}^\#(R)$ coincides with the set of all points of the form

$$(2.4) \quad \begin{bmatrix} a\tau & \beta \\ 0 & \tau^{-1} \end{bmatrix}, \quad \tau \in (E \otimes_{\mathbb{Q}} R)^\times, a \in R^\times, \beta \in R.$$

Let $\mathbf{N}^\#$ denote the unipotent radical of $\mathbf{B}^\#$, i.e., $\mathbf{N}^\#(R)$ is the set of points with $\tau = a = 1$. Set $\mathbf{Z}^\# := \{a1_2 \mid a \in \mathbf{GL}_1\}$; then $\mathbf{Z}^\#$ is a subgroup of the center of $\mathbf{G}^\#$ of index 2.

Let $I_\theta : E \longrightarrow \mathbf{M}_2(\mathbb{Q})$ denote the matrix representation of the regular representation of the \mathbb{Q} -algebra E with respect to the basis $\{1, \theta\}$, i.e.,

$$(2.5) \quad [\tau, \tau\theta] = [1, \theta] I_\theta(\tau), \quad \tau \in E,$$

or explicitly,

$$(2.6) \quad I_\theta(a + b\theta) = \begin{bmatrix} a & -bN_{E/\mathbb{Q}}(\theta) \\ b & a + b\mathrm{tr}_{E/\mathbb{Q}}(\theta) \end{bmatrix}, \quad a, b \in \mathbb{Q}.$$

For $\beta = b_2 + b_3\theta \in E_R$ with $b_2, b_3 \in R$, define an element X_β of $\mathbf{V}(R)$ as

$$(2.7) \quad X_\beta := \begin{bmatrix} b_1 & b_2 \\ b_2 & b_3 \end{bmatrix} \quad \text{with } b_1 := -b_2\mathrm{tr}_{E/\mathbb{Q}}(\theta) - b_3N_{E/\mathbb{Q}}(\theta).$$

We have

$$(2.8) \quad \iota_\theta \left(\begin{bmatrix} \tau & 0 \\ 0 & a\tau^{-1} \end{bmatrix} \right) = \mathbf{m}(I_\theta(\tau), a), \quad a \in R^\times, \tau \in E_R^\times,$$

$$(2.9) \quad \iota_\theta \left(\begin{bmatrix} 1 & \beta \\ 0 & 1 \end{bmatrix} \right) = \mathbf{n}(X_\beta), \quad \beta \in E_R.$$

By formulas (2.6) and (2.8), we have $\iota_\theta(a1_2) = a1_4$ for $(a \in R^\times)$, which implies

$$(2.10) \quad \iota_\theta(\mathbf{Z}^\#) = \mathbf{Z}.$$

We then have $N_{E/\mathbb{Q}}(x + \theta y) = [x, y] T \begin{bmatrix} x \\ y \end{bmatrix}$ with

$$(2.11) \quad T_\theta := \begin{bmatrix} 1 & 2^{-1}\mathrm{tr}_{E/\mathbb{Q}}(\theta) \\ 2^{-1}\mathrm{tr}_{E/\mathbb{Q}}(\theta) & N_{E/\mathbb{Q}}(\theta) \end{bmatrix}.$$

Let $\mathbf{V}^{T_\theta}(\mathbb{Q})$ be the orthogonal of T_θ in $\mathbf{V}(\mathbb{Q})$ with respect to the non-degenerate quadratic form $\mathrm{tr}(XY)$ on $\mathbf{V}(\mathbb{Q})$, i.e.,

$$\mathbf{V}^{T_\theta}(\mathbb{Q}) := \{X \in \mathbf{V}(\mathbb{Q}) \mid \mathrm{tr}(T_\theta X) = 0\}.$$

Note that $\mathrm{tr}(T_\theta^2) = 1 + (\mathrm{tr}_{E/\mathbb{Q}}(\theta)/2)^2 + N_{E/\mathbb{Q}}(\theta)^2 > 0$, $\det(T_\theta) = -D/4 > 0$ by (2.2), and $T_\theta \in \mathbf{V}(\mathbb{Z})$. We have $\mathbf{V}^{T_\theta}(\mathbb{Q}) = \{X_\beta \mid \beta \in E\}$, and

$$(2.12) \quad \mathbf{V}(\mathbb{Q}) = \mathbb{Q}T_\theta \oplus \mathbf{V}^{T_\theta}(\mathbb{Q}), \quad \iota_\theta(\mathbf{N}^\#) = \mathbf{n}(\mathbf{V}^{T_\theta})$$

by (2.9). The group \mathbf{M} acts on the space of rational homomorphisms $\mathrm{Hom}(\mathbf{N}, \mathbb{G}_a)$ by the rule $\mathrm{Ad}^*(m)\chi(n) = \chi(m^{-1}nm)$ for $m \in \mathbf{M}$, $n \in \mathbf{N}$ and $\chi \in \mathrm{Hom}(\mathbf{N}, \mathbb{G}_a)$. For $T \in \mathbf{V}$, let \mathbf{M}_T denote the stabilizer of $\chi_T : \mathbf{n}(X) \mapsto \mathrm{tr}(TX)$, and \mathbf{M}_T° the identity component with respect to the Zariski topology. Then, for any \mathbb{Q} -algebra R ,

$$\mathbf{M}_T^\circ(R) = \{\mathbf{m}(I_\theta(\tau), N_{E/\mathbb{Q}}(\tau)) \mid \tau \in E_R\}.$$

2.2. Open compact subgroups at finite places. Let p be a prime number. Set $E_p := E \otimes_{\mathbb{Q}} \mathbb{Q}_p$ and $\mathfrak{o}_{E,p} := \mathfrak{o}_E \otimes_{\mathbb{Z}} \mathbb{Z}_p$. Define

$$\mathbf{K}_p^{\#} := \mathbf{G}^{\#}(\mathbb{Z}_p), \quad \mathbf{K}_p := \mathbf{G}(\mathbb{Z}_p).$$

Then, $\mathbf{K}_p^{\#}$, which coincides with the stabilizer in $\mathbf{G}^{\#}(\mathbb{Q}_p)$ of the $\mathfrak{o}_{E,p}$ -lattice $\mathfrak{o}_{E,p}^2 \subset E_p^2$, is a maximal compact subgroup of $\mathbf{G}^{\#}(\mathbb{Q}_p)$, and \mathbf{K}_p is the standard maximal compact subgroup of $\mathbf{G}(\mathbb{Q}_p)$. For a non-zero ideal $\mathfrak{n} \subset \mathbb{Z}_p$, set

$$\mathbf{K}_0(\mathfrak{n}) := \{ \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \mathbf{G}(\mathbb{Z}_p) \mid C \in \mathfrak{n} \}.$$

For a non-zero ideal $\mathfrak{a} \subset \mathfrak{o}_{E,p}$, set

$$\mathbf{K}_0^{\#}(\mathfrak{a}) := \{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathbf{G}^{\#}(\mathbb{Z}_p) \mid c \in \mathfrak{a} \}.$$

Thus, $\mathbf{K}_0(\mathbb{Z}_p) = \mathbf{K}_p$ and $\mathbf{K}_0^{\#}(\mathfrak{o}_{E,p}) = \mathbf{K}_p^{\#}$.

Lemma 2.1. *For a non-zero ideal $\mathfrak{n} \subset \mathbb{Z}_p$, we have*

$$\mathbf{K}_0^{\#}(\mathfrak{n}\mathfrak{o}_{E,p}) = \iota_{\theta}^{-1}(\mathbf{K}_0(\mathfrak{n})).$$

Proof. Indeed, both $\iota_{\theta}^{-1}(\mathbf{K}_p)$ and $\mathbf{K}_p^{\#}$ coincides with the stabilizer of $\mathfrak{o}_{E,p}^2 = \langle v_1^+, v_2^+, v_1^-, v_2^- \rangle_{\mathbb{Z}_p}$ in $\mathbf{G}^{\#}(\mathbb{Q}_p)$. Hence $\iota_{\theta}^{-1}(\mathbf{K}_p) = \mathbf{K}_p^{\#}$. In the remaining part of the proof, we suppose $\mathfrak{n} \subset p\mathbb{Z}_p$. Then for $g = \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \mathbf{K}_p$, we have $g \in \mathbf{K}_0(\mathfrak{n})$ if and only if $g(v_1^+), g(v_2^+) \in \langle v_1^+, v_2^+ \rangle_{\mathbb{Z}_p} + \mathfrak{n}\langle v_1^-, v_2^- \rangle_{\mathbb{Z}_p}$. For $g = \iota_{\theta}(h)$ with $h = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathbf{K}_p^{\#}$, this last condition becomes $\begin{bmatrix} a \\ c \end{bmatrix}, \begin{bmatrix} a\theta \\ c\theta \end{bmatrix} \in \begin{bmatrix} \mathfrak{o}_{E,p} \\ \mathfrak{n}\mathfrak{o}_{E,p} \end{bmatrix}$, or equivalently, $c \in \mathfrak{n}\mathfrak{o}_{E,p}$. \square

For any $N \in \mathbb{Z}_{>0}$, define open compact subgroups $\mathbf{K}_0(N) \subset \mathbf{G}(\mathbb{A}_{\mathbf{f}})$ and $\mathbf{K}^{\#}(N\mathfrak{o}_E) \subset \mathbf{G}^{\#}(\mathbb{A}_{\mathbf{f}})$ as

$$\mathbf{K}_0(N) := \prod_{p < \infty} \mathbf{K}_0(N\mathbb{Z}_p), \quad \mathbf{K}_0^{\#}(N\mathfrak{o}_E) := \prod_{p < \infty} \mathbf{K}_0^{\#}(N\mathfrak{o}_{E,p}).$$

2.3. Maximal compact subgroup at the archimedean place. The identity connected component of $\mathbf{G}(\mathbb{R})$ is $\mathbf{G}(\mathbb{R})^0 = \{g \in \mathbf{G}(\mathbb{R}) \mid \nu(g) > 0\}$. Set $\mathbf{K}_{\infty} := \mathbf{G}(\mathbb{R})^0 \cap \mathbf{O}(4)$, which is a maximal compact subgroup of $\mathbf{G}(\mathbb{R})^0$ given as

$$\mathbf{K}_{\infty} = \{ \begin{bmatrix} A & B \\ -B & A \end{bmatrix} \mid A, B \in \mathbf{M}_2(\mathbb{R}), A + iB \in \mathbf{U}(2) \}.$$

Note that $\mathbf{K}_{\infty} \subset \mathbf{Sp}_2(\mathbb{R})$. The action of $\mathbf{G}(\mathbb{R})^0$ on the Siegel upper-half space

$$\mathfrak{h}_2 := \{Z \in \mathbf{M}_2(\mathbb{C}) \mid {}^t Z = Z, \operatorname{Im}(Z) \gg 0\},$$

denoted by $(g, Z) \mapsto g\langle Z \rangle$ is defined by the usual formula, i.e.,

$$g\langle Z \rangle := (AZ + B)(CZ + D)^{-1}, \quad g = \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \mathbf{G}(\mathbb{R})^0, \quad Z \in \mathfrak{h}_2.$$

The stabilizer of $i1_2 \in \mathfrak{h}_2$ coincides with \mathbf{K}_{∞} . Set $\gamma_{\infty} := \mathbf{m}(-1_2, -1) = \begin{bmatrix} -1_2 & 0 \\ 0 & 1_2 \end{bmatrix} \in \mathbf{G}(\mathbb{R})$; then $\mathbf{K}_{\infty} \cup \mathbf{K}_{\infty}\gamma_{\infty}$ is a maximal compact subgroup of $\mathbf{G}(\mathbb{R})$. The four vectors

$$u_1 := \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad u_2 := \begin{bmatrix} i \\ 0 \end{bmatrix}, \quad u_3 := \begin{bmatrix} 0 \\ -i \end{bmatrix}, \quad u_4 := \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

in $E_{\mathbb{R}}^2$ form an \mathbb{R} -basis such that

$$(2.13) \quad [v_1^+, v_2^+, v_1^-, v_2^-] = [u_1, u_2, u_3, u_4] (b_{\mathbb{R}}^{\theta})^{-1}$$

with

$$(2.14) \quad b_{\mathbb{R}}^{\theta} := m(A_{\theta}, 2^{-1}\sqrt{|D|})^{-1}, \quad A_{\theta} := \begin{bmatrix} 1 & 2^{-1}\mathrm{tr}_{E/\mathbb{Q}}(\theta) \\ 0 & -2^{-1}\sqrt{|D|} \end{bmatrix} \in \mathbf{GL}_2(\mathbb{R}).$$

Note that $b_{\mathbb{R}}^{\theta} \in \mathbf{G}(\mathbb{R})^0$ because $\nu(b_{\mathbb{R}}^{\theta})^{-1} = 2^{-1}\sqrt{|D|} > 0$. Set

$$\mathbf{K}_{\infty}^{\#} := \mathbf{U}(2) \cap \mathbf{G}^{\#}(\mathbb{R}) = \{k^{\#} \in \mathbf{U}(2) \mid \det(k^{\#}) = \pm 1\},$$

which is a maximal compact subgroup of $\mathbf{G}^{\#}(\mathbb{R})$. The (topological identity connected component $(\mathbf{K}_{\infty}^{\#})^0$ of $\mathbf{K}_{\infty}^{\#}$ is $\mathbf{SU}(2)$, and $\mathbf{K}_{\infty}^{\#} = (\mathbf{K}_{\infty}^{\#})^0 \cup (\mathbf{K}_{\infty}^{\#})^0 \delta_{\infty}$ with $\delta_{\infty} := \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$. A general element of $\mathbf{SU}(2)$ is written in the form

$$(2.15) \quad h = \begin{bmatrix} a & -\bar{b} \\ b & \bar{a} \end{bmatrix} \quad \text{with } a = a' + ia'', b = b' + ib'' \in \mathbb{C}, |a|^2 + |b|^2 = 1.$$

For such an h , a computation reveals the relation

$$(2.16) \quad [hu_1, hu_2, hu_3, hu_4] = [u_1, u_2, u_3, u_4] \begin{bmatrix} A & B \\ -B & A \end{bmatrix} \quad \text{with } A := \begin{bmatrix} a' & -a'' \\ a'' & a' \end{bmatrix}, B := \begin{bmatrix} b'' & b' \\ b' & -b'' \end{bmatrix}.$$

Lemma 2.2. *We have $\iota_{\theta}^{-1}(\mathbf{K}_{\infty}) = (\mathbf{K}_{\infty}^{\#})^0$. For $k_{\infty}^{\#} \in (\mathbf{K}_{\infty}^{\#})^0$ as in (2.15), and define $A, B \in \mathbf{M}_2(\mathbb{R})$ as in (2.16). Then,*

$$(2.17) \quad \iota_{\theta}(k_{\infty}^{\#}) = b_{\mathbb{R}}^{\theta} \begin{bmatrix} A & B \\ -B & A \end{bmatrix} (b_{\mathbb{R}}^{\theta})^{-1}.$$

Proof. The formula in (2.17) follows from (2.4), (2.16) and (2.13) immediately. From

$$[hu_1, hu_2, hu_3, hu_4] = [u_1, u_2, u_3, u_4](-\gamma)$$

and $-\gamma \notin \mathbf{K}_{\infty}$, the assertion follows. \square

2.4. Haar measures. For locally compact unimodular topological groups H relevant to us, we fix Haar measures η_H on H in the following manner.

2.4.1. On the additive group \mathbb{C} , define $d\eta_{\mathbb{C}}(\tau) := dx dy = 2^{-1}|d\tau \wedge d\bar{\tau}|$ for $\tau = x + iy$ ($x, y \in \mathbb{R}$) with $dx dy$ being the Lebesgue measure on \mathbb{R}^2 .

The Haar measure $\eta_{\mathbb{C}^{\times}}$ on \mathbb{C}^{\times} (resp. \mathbb{R}^{\times}) is defined by $d\eta_{\mathbb{C}^{\times}}(\tau) = |\tau|_{\mathbb{C}}^{-1} d\tau$ for $\tau = x + iy \in \mathbb{C}$ (resp. $|x|_{\mathbb{R}}^{-1} dx$). For each $p < \infty$, E_p^{\times} (resp. \mathbb{Q}_p^{\times}) is endowed with the Haar measure $\eta_{E_p^{\times}}$ such that $\eta_{E_p^{\times}}(\mathfrak{o}_{E,p}^{\times}) = 1$ (resp. $\eta_{\mathbb{Q}_p^{\times}}(\mathbb{Z}_p^{\times}) = 1$). Then, viewing \mathbb{A}_E^{\times} as the restricted product of E_p^{\times} , we define $\eta_{\mathbb{A}_E^{\times}} = \prod_{p \leq \infty} \eta_{E_p^{\times}}$. Similarly, we set $\eta_{\mathbb{A}^{\times}} = \prod_{p \leq \infty} \eta_{\mathbb{Q}_p^{\times}}$.

2.4.2. For a matrix $X = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, set $X^{\dagger} := \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$, so that $XX^{\dagger} = (\det X) 1_2$, and $X \mapsto X^{\dagger}$ induces an involution of the \mathbb{Q} -vector space $\mathbf{V}(\mathbb{Q})$. Endowed with the non-degenerate symmetric bilinear form $\mathrm{tr}(XY^{\dagger})$, $\mathbf{V}(\mathbb{Q})$ is regarded as a quadratic space. We endow the abelian group $\mathbf{V}(\mathbb{A})$ with the self-dual Haar measure with respect to the self-duality defined by the bi-character $(X, Y) \mapsto \psi(\mathrm{tr}(XY^{\dagger}))$. Let $\eta_{\mathbb{A}}$ (resp. $\eta_{\mathbb{A}_E}$) be the self-dual Haar measure on \mathbb{A} (resp. \mathbb{A}_E) with respect to the bi-character $(x, y) \mapsto \psi(xy)$ (resp. $(\alpha, \beta) \mapsto \psi(\mathrm{tr}_{E/\mathbb{Q}}(\alpha\bar{\beta}))$). A computation shows the relation $-\mathrm{tr}(X_{\alpha}X_{\beta}^{\dagger}) = \mathrm{tr}_{E/\mathbb{Q}}(\alpha\bar{\beta})$ for $\alpha, \beta \in \mathbb{A}_E$, where $X_{\alpha}, X_{\beta} \in V^{T_{\theta}}(\mathbb{A})$ are defined by (2.7). Thus, by the isomorphism $\mathbb{A}_E \ni \beta \mapsto X_{\beta} \in V^{T_{\theta}}(\mathbb{A})$, the Haar measure $\eta_{\mathbb{A}_E}$ is transferred to the self-dual Haar measure on $\mathbf{N}^{\#}(\mathbb{A})$ with respect to $(X, Y) \mapsto \mathrm{tr}(XY^{\dagger})$. Since $\mathrm{tr}(T_{\theta}T_{\theta}^{\dagger}) = -D/2 \neq 0$, we have the orthogonal direct sum decomposition $\mathbf{V}(\mathbb{Q}) = \mathbb{Q}T_{\theta}^{\dagger} \oplus V^{T_{\theta}}(\mathbb{Q})$. By this and (2.12), write $X \in V(\mathbb{A})$ as $X = xT_{\theta}^{\dagger} + X_{\beta} = yT_{\theta} + X_{\alpha}$ ($x, y \in \mathbb{A}, \beta, \alpha \in \mathbb{A}_E$). By definition of

the Haar measures, we have $d\eta_{\mathbf{N}(\mathbb{A})}(X) = d\eta_{\mathbb{A}}(x) \otimes d\eta_{\mathbb{A}_E}(\beta)$. Since the change of variables $(x, \beta) \rightarrow (y, \alpha)$ is given as

$$y = \frac{-D}{2} x, \quad \alpha = \beta + 2^{-1} \text{tr}_{E/\mathbb{Q}}(\theta)(x + y) + (N_{E/\mathbb{Q}}(\theta)x - y) \theta$$

and since $|\frac{-D}{2}|_{\mathbb{A}} = 1$, we get $d\eta_{\mathbf{N}(\mathbb{A})}(X) = d\eta_{\mathbb{A}}(y) \otimes d\eta_{\mathbb{A}_E}(\alpha)$. Through the identification $\mathbf{V}(\mathbb{A}) \ni X \mapsto \mathbf{n}(X) \in \mathbf{N}(\mathbb{A})$ and $\beta \ni \mathbb{A}_E \mapsto \begin{bmatrix} 1 & \beta \\ 0 & 1 \end{bmatrix} \in \mathbf{N}^{\#}(\mathbb{A})$, the groups $\mathbf{N}(\mathbb{A})$ and $\mathbf{N}^{\#}(\mathbb{A})$ acquire Haar measures. Thus, we have the integration formula:

$$(2.18) \quad \int_{\mathbf{N}(\mathbb{A})} f(n) dn = \int_{\mathbb{A}} \int_{\mathbf{N}^{\#}(\mathbb{A})} f(\mathbf{n}(xT_{\theta}) \iota_{\theta}(n^{\#})) dx dn^{\#}$$

for any $f \in L^2(\mathbf{N}(\mathbb{A}))$. Note that $\text{vol}(\mathbf{N}(\mathbb{Q}) \backslash \mathbf{N}(\mathbb{A})) = \text{vol}(\mathbf{N}^{\#}(\mathbb{Q}) \backslash \mathbf{N}^{\#}(\mathbb{A})) = 1$.

2.4.3. On the Lie group $\mathbf{G}^{\#}(\mathbb{R}) = \{g^{\#} \in \mathbf{GL}_2(\mathbb{C}) \mid \det(g^{\#}) \in \mathbb{R}^{\times}\}$, we fix a Haar measure as follows. By the Iwasawa decomposition, any element $g^{\#} \in \mathbf{G}^{\#}(\mathbb{R})$ is uniquely written as

$$(2.19) \quad g^{\#} = \begin{bmatrix} 1 & \beta \\ 0 & 1 \end{bmatrix} \begin{bmatrix} a\tau & 0 \\ 0 & \tau^{-1} \end{bmatrix} k^{\#}$$

with $a \in \mathbb{R}^{\times}$, $\tau \in \mathbb{C}^{\times}$, $\beta \in \mathbb{C}$ and $k^{\#} \in \mathbf{K}_{\infty}^{\#}$. Then, our Haar measure on $\mathbf{G}^{\#}(\mathbb{R})$ is symbolically defined as

$$(2.20) \quad dg^{\#} = |a|_{\mathbb{R}}^{-2} |\tau|_{\mathbb{C}}^{-2} \times d\eta_{\mathbb{C}}(\beta) \frac{d\eta_{\mathbb{R}}(a)}{|a|_{\mathbb{R}}} \frac{d\eta_{\mathbb{C}}(\tau)}{|\tau|_{\mathbb{C}}} dk^{\#},$$

where $dk^{\#}$ is the Haar measure on $\mathbf{K}_{\infty}^{\#}$ with volume 1, and $|\tau|_{\mathbb{C}} := \tau \bar{\tau}$. For $p < \infty$, the group $\mathbf{G}^{\#}(\mathbb{Q}_p)$ is endowed with the Haar measure $dg_p^{\#}$ uniquely determined by the condition $\text{vol}(\mathbf{K}_p^{\#}) = 1$. As in the archimedean case, if we write a general element $g^{\#}$ of $\mathbf{G}^{\#}(\mathbb{Q}_p)$ as in (2.19) with $a \in \mathbb{Q}_p^{\times}$, $\tau \in E_p^{\times}$, $\beta \in E_p$ and $k^{\#} \in \mathbf{K}_p^{\#}$, then

$$(2.21) \quad dg_p^{\#} = |a|_p^{-2} |N_{E_p/\mathbb{Q}_p} \tau|_p^{-2} \times d\eta_{E_p}(\beta) d\eta_{\mathbb{Q}_p^{\times}}(a) d\eta_{E_p^{\times}}(\tau) dk^{\#},$$

where $dk^{\#}$ is the normalized Haar measure of $\mathbf{K}_p^{\#}$. On the adèle group $\mathbf{G}^{\#}(\mathbb{A})$, we use the product measure of Haar measures on $\mathbf{G}^{\#}(\mathbb{Q}_p)$ for $p \leq \infty$. Then, from (2.20) and (2.21), we get

$$(2.22) \quad \int_{\mathbf{G}^{\#}(\mathbb{A})} f(g^{\#}) dg^{\#} = \frac{\sqrt{|D|}}{2} \int_{\mathbf{N}^{\#}(\mathbb{A})} \int_{\mathbb{A}^{\times}} \int_{\mathbb{A}_E^{\times}} \int_{\mathbf{K}^{\#}} f(n^{\#} \begin{bmatrix} a\tau & 0 \\ 0 & \tau^{-1} \end{bmatrix} k^{\#}) |a|_{\mathbb{A}}^{-2} |N_{E/\mathbb{Q}}(\tau)|_{\mathbb{A}}^{-2} \\ \times dn^{\#} d\eta_{\mathbb{A}^{\times}}(a) d\eta_{\mathbb{A}_E^{\times}}(\tau) dk^{\#},$$

where $dk^{\#}$ is the normalized Haar measure on $\mathbf{K}^{\#} = \prod_{p \leq \infty} \mathbf{K}_p^{\#}$. Note that the measure on $\mathbf{N}^{\#}(\mathbb{A}) \cong \mathbb{A}_E$ is $\eta_{\mathbb{A}_E}$ defined above, which is $\frac{\sqrt{|D|}}{2} \times \prod_p \eta_{E_p}$ ($p \leq \infty$) due to the formulas $(\prod_p \eta_{E_p})(\mathbb{A}_E/E) = |D|^{1/2}/2$ ([49, Chap V §4 Proposition 7]) and $\eta_{\mathbb{A}_E}(\mathbb{A}_E/E) = 1$.

2.4.4. We fix $\eta_{\mathbf{K}_{\infty}}$ so that the measure of \mathbf{K}_{∞} is 1. Then we normalize $\eta_{\mathbf{Sp}_2(\mathbb{R})}$ in such a way that the quotient $\eta_{\mathbf{Sp}_2(\mathbb{R})}/\eta_{\mathbf{K}_{\infty}}$ corresponds to the measure $(\det Z)^{-3} dX dY$ on the Siegel upper-half space $\mathfrak{h}_2 \cong \mathbf{Sp}_2(\mathbb{R})/\mathbf{K}_{\infty}$. By $\mathbf{G}(\mathbb{R})^0 = \mathbf{Z}(\mathbb{R})^0 \mathbf{Sp}_2(\mathbb{R}) \cong \mathbb{R}_{>0} \times \mathbf{Sp}_2(\mathbb{R})$, $\eta_{\mathbf{G}(\mathbb{R})}$ is defined by demanding that its restriction to $\mathbf{G}(\mathbb{R})^0$ is $\eta_{\mathbb{R}^{\times}} \otimes \eta_{\mathbf{Sp}_2(\mathbb{R})}$. For $p < \infty$, we fix $\eta_{\mathbf{G}(\mathbb{Q}_p)}$ by $\text{vol}(\mathbf{K}_p) = 1$. Then, $\eta_{\mathbf{G}(\mathbb{A})}$ is defined to be the restricted product of $\eta_{\mathbf{G}(\mathbb{Q}_v)}$ ($v \leq \infty$).

2.5. Idele class characters. Let Λ denote a character of the finite group

$$\mathrm{Cl}(E) := \mathbb{A}_E^\times / E^\times E_{\mathbb{R}}^\times \widehat{\mathfrak{o}}_E^\times.$$

As is well known, this group is isomorphic to the ideal class group of E . We regard Λ as an idele class character of E^\times of finite order. Since $\mathbb{A}^\times = \mathbb{Q}^\times (\mathbb{R}_{>0}) \widehat{\mathbb{Z}}^\times$ is contained in the subgroup $E^\times E_{\mathbb{R}}^\times \widehat{\mathfrak{o}}_E^\times$, we have

$$(2.23) \quad \Lambda|_{\mathbb{A}^\times} = \mathbf{1}.$$

Let v be a place of E and Λ_v the v -component of Λ , i.e., $\Lambda = \otimes_v \Lambda_v$. We say that v is inert in E/\mathbb{Q} , splits in E/\mathbb{Q} or ramifies in E/\mathbb{Q} if $E_v := E \otimes_{\mathbb{Q}} \mathbb{Q}_p$ is an unramified field extension of \mathbb{Q}_p , is isomorphic to $\mathbb{Q}_p \oplus \mathbb{Q}_p$, or is a ramified field extension of \mathbb{Q}_p , respectively. Since Λ is supposed to be trivial in $E_{\mathbb{R}}^\times$, we have $\Lambda_\infty = \mathbf{1}$; moreover,

$$(2.24) \quad \text{If } v \text{ is inert in } E/\mathbb{Q}, \text{ then } \Lambda_v = \mathbf{1}.$$

Indeed, since Λ is supposed to be trivial on $\widehat{\mathfrak{o}}_E^\times$, the restriction $\Lambda_v|_{\mathfrak{o}_{E,v}^\times}$ is trivial. If v is inert and p denote the residue characteristic of E_v , then $p \in \mathbb{Q}_p$ is a prime element of the local field E_v . We have $\Lambda_v(p) = 1$ due to (2.23). Hence, Λ_v is trivial on $p^{\mathbb{Z}} \mathfrak{o}_{E,v}^\times = E_p^\times$.

Define the Galois conjugate Λ^\dagger of Λ by setting

$$(2.25) \quad \Lambda^\dagger(\tau) := \Lambda(\bar{\tau}), \quad \tau \in \mathbb{A}_E^\times.$$

We have $\Lambda(\tau)\Lambda^\dagger(\tau) = \Lambda(\tau\bar{\tau}) = 1$ for $\tau \in \mathbb{A}_E^\times$ by (2.23). Hence

$$(2.26) \quad \Lambda^\dagger = \Lambda^{-1} = \bar{\Lambda},$$

where $\bar{\Lambda}$ is the complex conjugate of Λ .

For an idele class character ξ of E^\times , let $\widehat{L}(s, \xi)$ be the completed Hecke L -function of ξ , and $L_p(s, \xi)$ its local p -factor for $p \leq \infty$. Then, $L_p(s, \xi^\dagger) = L_p(s, \xi)$ for any $p < \infty$. Indeed, if p is not ramified in E/\mathbb{Q} , the equality is trivial. Suppose that E/\mathbb{Q} is ramified; if ξ_p is a ramified character of E_p^\times , then both L -factors are 1. If ξ_p is unramified, then for any prime element ϖ of E_p , we have $(\xi^\dagger)_p(\varpi) = \xi_p(\bar{\varpi}) = \xi_p(\varpi)$, which implies the equality between local p -factors.

For any character $\mu : \mathbb{A}^\times / \mathbb{Q}^\times \mathbb{R}^\times \rightarrow \mathbb{C}^1$, we define $\mu_E := \mu \circ N_{E/\mathbb{Q}} : \mathbb{A}_E^\times / E^\times \rightarrow \mathbb{C}^1$. Then, μ_E is Galois invariant, and $\mu_E|_{E_\infty^\times} = \mathbf{1}$. Hence, $(\Lambda\mu_E)^\dagger = \Lambda^\dagger\mu_E^\dagger = \Lambda^{-1}\mu_E$, and

$$(2.27) \quad \widehat{L}(s, \Lambda^{-1}\mu_E) = \widehat{L}(s, \Lambda^\dagger\mu_E) = \widehat{L}(s, \Lambda\mu_E), \quad (\mathrm{Re}(s) > 1).$$

Note that $\widehat{L}(s, \Lambda\mu_E)$ is holomorphic except for possible simple poles at $s = 1, 0$, which occurs if and only if both Λ and μ are trivial.

3. EISENSTEIN SERIES AND RANKIN-SELBERG INTEGRAL

3.1. Eisenstein series. For details, we refer to [15, §19]; the theory on $\mathbf{GL}_2(\mathbb{A}_E)$ developed there carries over into the group $\mathbf{G}^\#(\mathbb{A})$ by a minor modification. For a finite-dimensional vector space V over a local field, let $\mathcal{S}(V)$ be the space of all Schwartz-Bruhat

functions on V . For $p \leq \infty$, $\phi \in \mathcal{S}(E_p^2)$ and $s \in \mathbb{C}$ with $\operatorname{Re}(s) > 0$, we define a function $f_\phi^{(s, \Lambda_p, \mu_p)} : \mathbf{G}^\#(\mathbb{Q}_p) \rightarrow \mathbb{C}$ by

$$(3.1) \quad f_\phi^{(s, \Lambda_p, \mu_p)}(g^\#) = \mu_p(\det g^\#) |\det g^\#|_p^{s+1} \int_{E_p^\times} \phi\left(\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} \tau & 0 \\ 0 & \bar{\tau} \end{bmatrix} g^\#\right) \Lambda_p \mu_{E,p}(\tau) |\tau \bar{\tau}|_p^{s+1} d\eta_{E_p^\times}(\tau).$$

When $p = \infty$, we assume that ϕ is $\mathbf{K}_\infty^\#$ -finite. By the local Tate's theory, the function $s \mapsto f_\phi^{(s, \Lambda_p, \mu_p)}(g^\#)$ is continued meromorphically to \mathbb{C} in such a way that

$$(3.2) \quad \mathbf{f}_\phi^{(s, \Lambda_p, \mu_p)}(g^\#) := L_p(s+1, \Lambda_p \mu_{E,p})^{-1} \times f_\phi^{(s, \Lambda_p, \mu_p)}(g^\#)$$

is holomorphic on \mathbb{C} . Then $\mathbf{f}_\phi^{(s, \Lambda_p, \mu_p)}$ belongs to the space $\mathcal{V}^\#(s, \Lambda_p, \mu_p)$ consisting of all smooth functions f on $\mathbf{G}^\#(\mathbb{Q}_p)$ satisfying

$$f\left(\begin{bmatrix} \tau & \beta \\ 0 & a\bar{\tau} \end{bmatrix} g^\#\right) = \Lambda_p(\tau)^{-1} \mu_p(a)^{-1} |a|_p^{-(s+1)} f(g^\#)$$

for any $\begin{bmatrix} \tau & \beta \\ 0 & a\bar{\tau} \end{bmatrix} \in \mathbf{B}^\#(\mathbb{Q}_p)$ and $g^\# \in \mathbf{G}^\#(\mathbb{Q}_p)$.

Let $\mathcal{S}(\mathbb{A}_E^2)$ be the space of all Schwartz-Burhat functions on \mathbb{A}_E^2 . The Fourier transform $\widehat{\phi}$ of $\phi \in \mathcal{S}(\mathbb{A}_E^2)$ is defined by

$$\widehat{\phi}(x, y) = \int_{\mathbb{A}_E^2} \phi(u, v) \psi(\langle \begin{bmatrix} x \\ y \end{bmatrix}, \begin{bmatrix} u \\ v \end{bmatrix} \rangle) d\eta_{\mathbb{A}_E}(u) d\eta_{\mathbb{A}_E}(v).$$

Set $w_0 := \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \in \mathbf{G}^\#(\mathbb{Q}_p)$. The standard intertwining operator

$$M(s) : \mathcal{V}^\#(s, \Lambda_p, \mu_p) \longrightarrow \mathcal{V}^\#(-s, \Lambda_p^{-1}, \mu_p^{-1})$$

is defined on $\operatorname{Re}(s) > 0$ as the absolutely convergent integral

$$M(s)f(g^\#) = \int_{E_p} f\left(w_0 \begin{bmatrix} 1 & \beta \\ 0 & 1 \end{bmatrix} g^\#\right) d\eta_{E_p}(\beta), \quad g^\# \in \mathbf{G}^\#(\mathbb{Q}_p).$$

The effect of $M(s)$ on the section $f_\phi^{(s, \Lambda_p, \mu_p)}$ is described by the Fourier transform $\widehat{\phi}$ as

$$(3.3) \quad f_\phi^{(-s, \Lambda_p^{-1}, \mu_p^{-1})}(g^\#) = c_p \Lambda_p(\sqrt{D})^{-1} |D|_p^{-s+1/2} \varepsilon_p(s, \Lambda \mu_E, \psi_{E_p}) \frac{L_p(1-s, \Lambda^{-1} \mu_E^{-1})}{L_p(s, \Lambda \mu_E)} \times M(s) f_\phi^{(s, \Lambda_p, \mu_p)}(g^\#)$$

with $c_p = 1$ if $p < \infty$ and $c_\infty = \frac{\sqrt{|D|}}{2}$ if $p = \infty$, where $\varepsilon(s, \bullet, \psi_{E_p})$ denote Tate's local epsilon factor defined by the character ψ_{E_p} of E_p and the associated self-dual measure on E_p as usual. For a decomposable element $\phi = \otimes_{p \leq \infty} \phi_p \in \mathcal{S}(\mathbb{A}_E^2)$, we define

$$\mathbf{f}_\phi^{(s, \Lambda, \mu)}(g^\#) = \prod_{p \leq \infty} \mathbf{f}_{\phi_p}^{(s, \Lambda_p, \mu_p)}(g_p^\#), \quad g^\# = (g_p^\#)_p \in \mathbf{G}^\#(\mathbb{A}).$$

Note that $\mathbf{f}_\phi^{(s, \Lambda, \mu)}$ is left $\mathbf{B}^\#(\mathbb{Q})$ -invariant and right $\mathbf{K}^\#$ -finite.

The Eisenstein series attached to $\mathbf{f}_\phi^{(s, \Lambda, \mu)}$ is defined by the absolutely convergent series

$$(3.4) \quad E(\phi, s, \Lambda, \mu; g^\#) := \widehat{L}(s+1, \Lambda \mu_E) \times \sum_{\delta \in \mathbf{B}^\#(\mathbb{Q}) \setminus \mathbf{G}^\#(\mathbb{Q})} \mathbf{f}_\phi^{(s, \Lambda, \mu)}(\delta g^\#), \quad g^\# \in \mathbf{G}^\#(\mathbb{A})$$

for $\operatorname{Re}(s) > 1$. The following property of the Eisenstein series is standard.

Proposition 3.1. *Let $\phi \in \mathcal{S}(\mathbb{A}_E^2)$, $s \in \mathbb{C}$, $\Lambda \in \widehat{\text{Cl}(E)}$, $\mu \in \mathbb{A}^\times / \widehat{\mathbb{Q}^\times \mathbb{R}^\times}$ and $g^\# \in \mathbf{G}^\#(\mathbb{A})$.*

- (i) *The map $s \mapsto E(\phi, s, \Lambda, \mu; g^\#)$ ($\text{Re}(s) > 1$) has a meromorphic continuation to \mathbb{C} , holomorphic in s unless $\Lambda\mu_E \neq \mathbf{1}$ in which case it has possible simple poles only at $s = 1, -1$. The function $g^\# \mapsto E(\phi, s, \Lambda, \mu; g^\#)$ is an automorphic form on $\mathbf{G}^\#(\mathbb{Q})\mathbf{Z}^\#(\mathbb{A}) \backslash \mathbf{G}^\#(\mathbb{A})$.*
- (ii) *We have the functional equation $E(\widehat{\phi}, -s, \Lambda^{-1}, \mu^{-1}; g^\#) = E(\phi, s, \Lambda, \mu; g^\#)$.*
- (iii) *For a relatively compact subset $\mathcal{N} \subset \mathbb{C}$ on which $s \mapsto E(\phi, s, \Lambda, \mu; g^\#)$ is regular and for a compact set $\mathcal{U} \subset \mathbf{G}^\#(\mathbb{A})$, there exists constants $C > 0$ and $N > 0$ such that*

$$|E(\phi, s, \Lambda, \mu; \begin{bmatrix} a & 0 \\ 0 & 1 \end{bmatrix} g^\#)| \leq C |a|^N \quad (a \in \mathbb{R}_{>1}, g^\# \in \mathcal{U}, s \in \mathcal{N}).$$

3.2. Rankin-Selberg integral and the basic identity. For any cusp form φ on $\mathbf{G}(\mathbb{A})$, $\Lambda \in \widehat{\text{Cl}(E)}$ and $s \in \mathbb{C}$, the Rankin-Selberg integral is defined by

$$(3.5) \quad \langle E(\phi, s, \Lambda, \mu), \varphi \rangle := \int_{\mathbf{Z}^\#(\mathbb{A})\mathbf{G}^\#(\mathbb{Q}) \backslash \mathbf{G}^\#(\mathbb{A})} E(\phi, s, \Lambda, \mu; g^\#) \varphi(\iota_\theta(g^\#)) dg^\#.$$

By Lemma 3.1(iii) and by the Fourier expansion (4.5), it is straightforward to show the absolute convergence of the integral for $s \in \mathbb{C}$ where the Eisenstein series is regular. Let $\psi : \mathbb{A}/\mathbb{Q} \rightarrow \mathbb{C}^1$ be the basic character. For $T \in \mathbf{V}(\mathbb{Q})$, define a character of $\mathbf{N}(\mathbb{A})$ by

$$\psi_T(\mathbf{n}(X)) = \psi(\text{tr}(TX)), \quad X \in \mathbf{V}(\mathbb{A}).$$

The (T_θ, Λ) -Bessel period of a cusp form φ on $\mathbf{Z}(\mathbb{A})\mathbf{G}(\mathbb{Q}) \backslash \mathbf{G}(\mathbb{A})$ is defined by the integral¹

$$(3.6) \quad B^{T_\theta, \Lambda}(\varphi; g) := \int_{\mathbb{A}_E^\times / E^\times \mathbb{A}^\times} \Lambda(\tau)^{-1} \int_{\mathbf{N}(\mathbb{Q}) \backslash \mathbf{N}(\mathbb{A})} \psi_{T_\theta}(n)^{-1} \varphi(\mathbf{m}(I_\theta(\tau), N_{E/\mathbb{Q}}(\tau)) n g) dn d^\times \tau$$

for $g \in \mathbf{G}(\mathbb{A})$, where $d^\times \tau$ denote the quotient measure on $\mathbb{A}_E^\times / \mathbb{A}^\times$ (see §2.4.1). Here is the basic identity, which is originally due to Andrianov in the classical setting ([1], [2]) and is adelically reformulated by Piatetski-Shapiro ([30]). To determine constants exactly under our normalization of Haar measures, we reproduce the proof briefly.

Lemma 3.2. *Let $\varphi : \mathbf{Z}(\mathbb{A}) \backslash \mathbf{G}(\mathbb{Q})\mathbf{G}(\mathbb{A}) \rightarrow \mathbb{C}$ be a cusp form. For $\text{Re}(s) > 1$, we have*

$$(3.7) \quad \begin{aligned} \langle E(\phi, s, \Lambda, \mu), \varphi \rangle &= \frac{\sqrt{|D|}}{2} L(s+1, \Lambda\mu_E) \\ &\quad \times \int_{\mathbb{A}^\times} \int_{\mathbf{K}_\infty^\# \mathbf{K}_0^\#(\mathfrak{o}_E)} \mathbf{f}_\phi^{(s, \Lambda, \mu)}(k^\#) \mu(a) |a|_{\mathbb{A}}^{s-1} B^{T_\theta, \Lambda}(\varphi; \mathbf{m}(a1_2, a) \iota_\theta(k^\#) b_{\mathbb{R}}^\theta) d^\times a dk^\#. \end{aligned}$$

¹Note that $\mathbf{m}(I_\theta(a), N_{E/\mathbb{Q}}(a)) = a1_4$ for $a \in \mathbb{A}^\times$. Then, due to φ being $\mathbf{Z}(\mathbb{A})$ -invariant, the integral $B^{T_\theta, \Lambda}(\varphi; g)$ is 0 if $\Lambda|\mathbb{A}^\times \neq 1$. This trivial vanishing is not the case by (2.23).

Proof. By substituting (3.4) to (3.5),

$$\begin{aligned}
& L(s+1, \Lambda\mu_E)^{-1} \times \langle E(\phi, s, \Lambda, \mu), R(b_{\mathbb{R}}^{\theta})\varphi \rangle \\
&= \int_{Z^{\#}(\mathbb{A})G^{\#}(\mathbb{Q}) \backslash G^{\#}(\mathbb{A})} \sum_{\gamma \in B^{\#}(\mathbb{Q}) \backslash G^{\#}(\mathbb{Q})} \mathbf{f}_{\phi}^{(s, \Lambda, \mu)}(\gamma g^{\#}) \varphi(\iota_{\theta}(g^{\#})b_{\mathbb{R}}^{\theta}) dg^{\#} \\
&= \int_{Z^{\#}(\mathbb{A})B^{\#}(\mathbb{Q}) \backslash G^{\#}(\mathbb{A})} \mathbf{f}_{\phi}^{(s, \Lambda, \mu)}(g^{\#}) \varphi(\iota_{\theta}(g^{\#})b_{\mathbb{R}}^{\theta}) dg^{\#} \\
&= \frac{\sqrt{|D|}}{2} \int_{\mathbb{A}^{\times}/\mathbb{Q}^{\times}} \int_{\mathbb{A}_E^{\times}/\mathbb{A}^{\times}E^{\times}} \int_{N^{\#}(\mathbb{Q}) \backslash N^{\#}(\mathbb{A})} \int_{\mathbf{K}_0^{\#}(\mathfrak{o}_E)\mathbf{K}_{\infty}^{\#}} |a|_{\mathbb{A}}^{-(s+1)} \Lambda(\tau)^{-1} \mu(a)^{-1} \\
&\quad \times \varphi(\iota_{\theta}(n^{\#}) \mathbf{m}(I_{\theta}(\tau), N_{E/\mathbb{Q}}(\tau)) \begin{bmatrix} a^{-1}1_2 & 0 \\ 0 & 1_2 \end{bmatrix} \iota_{\theta}(k_{\mathbf{f}}^{\#})) |a|_{\mathbb{A}}^2 d^{\times}a d^{\times}\tau dn^{\#} dk_{\mathbf{f}}^{\#} dk_{\infty}^{\#},
\end{aligned}$$

where the last equality is proved by (2.22). By Lemma 3.3, the last expression becomes

$$\begin{aligned}
& \frac{\sqrt{|D|}}{2} \int_{\mathbb{A}^{\times}/\mathbb{Q}^{\times}} \int_{\mathbb{A}_E^{\times}/\mathbb{A}^{\times}E^{\times}} \int_{N^{\#}(\mathbb{Q}) \backslash N^{\#}(\mathbb{A})} \int_{\mathbf{K}_0^{\#}(\mathfrak{o}_E)\mathbf{K}_{\infty}^{\#}} |a|_{\mathbb{A}}^{-(s+1)} \Lambda(\tau)^{-1} \mu(a)^{-1} \\
&\quad \times \sum_{\alpha \in \mathbb{Q}^{\times}} \varphi(n \mathbf{m}(I_{\theta}(\tau), N_{E/\mathbb{Q}}(\tau)) \begin{bmatrix} \alpha a^{-1}1_2 & 0 \\ 0 & 1_2 \end{bmatrix} \iota_{\theta}(k_{\mathbf{f}}^{\#})) \psi_{T_{\theta}}(n)^{-1} |a|_{\mathbb{A}}^2 d^{\times}a d^{\times}\tau dn dk_{\mathbf{f}}^{\#}.
\end{aligned}$$

The α -summation and the a -integral over $\mathbb{A}^{\times}/\mathbb{Q}^{\times}$ are combined to yields an integral over \mathbb{A}^{\times} . Then, by the change of variables $a \mapsto a^{-1}$, we get the desired formula by (3.6). \square

Lemma 3.3.

$$\int_{N^{\#}(\mathbb{Q}) \backslash N^{\#}(\mathbb{A})} \varphi(\iota_{\theta}(n^{\#})g) dn^{\#} = \sum_{\alpha \in \mathbb{Q}^{\times}} \int_{V(\mathbb{Q}) \backslash V(\mathbb{A})} \varphi(\mathbf{n}(X) \begin{bmatrix} \alpha 1_2 & 0 \\ 0 & 1_2 \end{bmatrix} g) \psi_{T_{\theta}}(\mathbf{n}(X))^{-1} dX, \quad g \in G(\mathbb{A}).$$

Proof. The proof is straightforward. \square

4. AUTOMORPHIC FORMS AND FOURIER COEFFICIENTS

Let $(\mathbb{Z}^2)_{\text{dom}} := \{\lambda = (l_1, l_2) \in \mathbb{Z}^2 \mid l_1 \geq l_2\}$. Let $\lambda = (l_1, l_2) \in (\mathbb{Z}^2)_{\text{dom}}$ and ϱ the representation of $\mathbf{GL}_2(\mathbb{C})$ on the space V_{ϱ} of homogeneous polynomials in X, Y of degree $l_1 - l_2$ defined as $\varrho(h)f(X, Y) = (\det h)^{l_2} f(aX + bY, cX + dY)$ for $h = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathbf{GL}_2(\mathbb{C})$ and $f \in V_{\varrho}$. As is well-known, any irreducible rational representation of $\mathbf{GL}_2(\mathbb{C})$, up to equivalence, is obtained this way. The space V_{ϱ} carries an $\mathbf{SU}(2)$ -invariant hermitian inner product $(\cdot \mid \cdot)_{\varrho}$ given by [10, ((8.2.6)]. Recall $\mathbf{K}_{\infty} = \{k_{\infty}(u) := \begin{bmatrix} A & B \\ -B & A \end{bmatrix} \mid u = A + iB \in \mathbf{U}(2)\}$. For $N \in \mathbb{Z}_{>0}$, let $S_{\varrho}(\mathbf{K}_0(N))$ denotes the space of smooth functions $\varphi : G(\mathbb{A}) \rightarrow V_{\varrho}$ that satisfies the following conditions:

- (i) $\varphi(z\gamma g) = \varphi(g)$ for $(z, \gamma, g) \in \mathbf{Z}(\mathbb{A}) \times G(\mathbb{Q}) \times G(\mathbb{A})$.
- (ii) $\varphi(gk_{\mathbf{f}}k_{\infty}(u)) = \varrho(\bar{u})^{-1}\varphi(g)$ for $k_{\mathbf{f}} \in \mathbf{K}_0(N)$ and $k_{\infty}(u) \in \mathbf{K}_{\infty}$.
- (iii) $R(X)\varphi = 0$ for all $X \in \mathfrak{p}^{-}$.
- (iv) φ is bounded on $G(\mathbb{A})$.

For $T \in V(\mathbb{R})$ with $\det(T) \neq 0$, let $\mathbf{B}_{\varrho}^T : G(\mathbb{R})^0 \rightarrow \text{End}_{\mathbb{C}}(V_{\varrho})$ be a function defined by

$$(4.1) \quad \mathbf{B}_{\lambda}^T(g) := \nu(g)^{(l_1+l_2)/2} \varrho(Ci + D)^{-1} \exp(2\pi i \text{tr}(T g(i1_2))), \quad g = \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in G(\mathbb{R})^0.$$

This function satisfies the conditions:

$$(4.2) \quad \mathbf{B}_\varrho^T(\mathbf{m}(A, c)g) = \mathbf{B}_\varrho^T(g) \circ \varrho({}^t A^{-1} c)^{-1}, \quad \mathbf{m}(A, c) \in \mathbf{M}_T^\circ(\mathbb{R}), \quad g \in \mathbf{G}(\mathbb{R})^0$$

$$(4.3) \quad \mathbf{B}_\varrho^T(n g k_\infty(u)) = \psi_T(n) \varrho(\bar{u})^{-l} \circ \mathbf{B}(g), \quad (n, g, u) \in \mathbf{N}(\mathbb{R}) \times \mathbf{G}(\mathbb{R})^0 \times \mathbf{U}(2),$$

$$(4.4) \quad R(\mathfrak{p}^-) \mathbf{B}_\varrho^T = 0.$$

The function $B_{\varrho, v}^T : g \mapsto \mathbf{B}_\varrho^T(g)(v)$ with $v \in V_\varrho - (0)$ is bounded on $\mathbf{G}(\mathbb{R})^0$ if and only if $T \in \mathbf{V}(\mathbb{R})^+$, where $\mathbf{V}(\mathbb{R})^+$ denote the set of positive definite elements in $\mathbf{V}(\mathbb{R})$. Set

$$\gamma := \mathbf{m}(-1_2, -1) \in \mathbf{M}(\mathbb{Q}).$$

Note that $\nu(\gamma) = -1$.

Lemma 4.1. *Let $\varphi \in S_\varrho(\mathbf{K}_0(N))$. There exists a unique family of vectors $a_\varphi(T; g_{\mathbf{f}}) \in V_\varrho$ ($T \in \mathbf{V}(\mathbb{Q})$), $g_{\mathbf{f}} \in \mathbf{G}(\mathbb{A}_{\mathbf{f}})$ such that*

$$(4.5) \quad \varphi(g_\infty g_{\mathbf{f}}) = \sum_{T \in \mathbf{V}(\mathbb{Q})} \mathbf{B}_\varrho^T(g_\infty)(a_\varphi(T; g_{\mathbf{f}})), \quad g_\infty \in \mathbf{G}(\mathbb{R})^\circ, \quad g_{\mathbf{f}} \in \mathbf{G}(\mathbb{A}_{\mathbf{f}}).$$

If $T \notin \mathbf{V}(\mathbb{Q})^+$, then $a_\varphi(T; g_{\mathbf{f}}) = 0$ for all $g_{\mathbf{f}} \in \mathbf{G}(\mathbb{A}_{\mathbf{f}})$. For $g_{\mathbf{f}} \in \mathbf{G}(\mathbb{A}_{\mathbf{f}})$,

$$(4.6) \quad \int_{\mathbf{N}(\mathbb{A})/\mathbf{N}(\mathbb{Q})} \varphi(n g_\infty g_{\mathbf{f}}) \psi_T(n)^{-1} dn = \begin{cases} \mathbf{B}_\varrho^T(g_\infty)(a_\varphi(T; g_{\mathbf{f}})), & (g_\infty \in \mathbf{G}(\mathbb{R})^\circ), \\ 0, & (g_\infty \in \mathbf{G}(\mathbb{R}) - \mathbf{G}(\mathbb{R})^\circ). \end{cases}$$

Proof. Fix $g_{\mathbf{f}} \in \mathbf{G}(\mathbb{A}_{\mathbf{f}})$ and examine the integral, say $W(g_\infty)$, on the left-hand side of (4.6). The function W on $\mathbf{G}(\mathbb{R})^0$ satisfies conditions (4.3) and (4.4). As such, there corresponds a function $F : \mathfrak{h}_2 \rightarrow V_\varrho$ determined by the relation $F(g_\infty \langle i1_2 \rangle) = \nu(g)^{-(l_1+l_2)/2} \varrho(Ci + D) W(g_\infty)$ with $g_\infty = \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \mathbf{G}(\mathbb{R})^0$ such that $Z = (Ai + B)(Ci + D)^{-1}$. Since $F(Z)$ satisfies $F(Z + X) = \psi_T(X) F(Z)$ ($X \in \mathbf{V}(\mathbb{R})$), it is of the form $F(Z) = F_1(Y) \exp(2\pi i \text{tr}(TX))$ with a V_ϱ -valued C^∞ -function $F_1(Y)$. Since $F(Z)$ is holomorphic, the Cauchy-Riemann equations yield $\frac{d}{dy_{ij}} F_1(Y) = -2\pi t_{ij} F_1(T)$, which is uniquely solved as

$$F_1(Y) = a_\varphi(T; g_{\mathbf{f}}) \exp(-2\pi \text{tr}(TY))$$

with a vector $a_\varphi(T; g_{\mathbf{f}}) \in V_\varrho$, or equivalently $W(g_\infty) = \mathbf{B}_\varrho^T(g_\infty)(a_\varphi(T; g_{\mathbf{f}}))$. Then, the formula in (4.5) is a consequence of the Fourier expansion of a $\mathbf{N}(\mathbb{Q})$ -periodic function on $\mathbf{N}(\mathbb{A})$. Since φ is bounded on $\mathbf{G}(\mathbb{A})$, the function $W(g_\infty)$ should be bounded on $\mathbf{G}(\mathbb{R})^0$; since $g_\infty \mapsto \mathbf{B}_\varrho^T(g_\infty)(v)$ with $v \in V_\varrho - \{0\}$ is unbounded if $T \notin \mathbf{V}(\mathbb{R})^+$, we have $a_\varphi(T; g_{\mathbf{f}}) = 0$ for all $T \in \mathbf{V}(\mathbb{Q}) - \mathbf{V}(\mathbb{R})^+$. It remains to show the second case in (4.6). Let $g_\infty \notin \mathbf{G}(\mathbb{R})^\circ$. Decompose $\gamma \in \mathbf{M}(\mathbb{Q})$ as $\gamma_\infty \gamma_{\mathbf{f}}$ ($\gamma_\infty \in \mathbf{M}(\mathbb{R})$, $\gamma_{\mathbf{f}} \in \mathbf{M}(\mathbb{A}_{\mathbf{f}})$) in $\mathbf{M}(\mathbb{A})$. By the left $\mathbf{G}(\mathbb{Q})$ -invariance of φ ,

$$\varphi(\mathfrak{n}(X) g_\infty g_{\mathbf{f}}) = \varphi(\gamma \mathfrak{n}(X) g_\infty g_{\mathbf{f}}) = \varphi(\mathfrak{n}(-X) \gamma_\infty g_\infty \gamma_{\mathbf{f}} g_{\mathbf{f}}).$$

Integrate this in X over $\mathbf{V}(\mathbb{Q}) \setminus \mathbf{V}(\mathbb{A})$ and make a change of variables $n \rightarrow n^{-1}$ on the way; we have

$$\begin{aligned} \int_{\mathbf{N}(\mathbb{Q}) \setminus \mathbf{N}(\mathbb{A})} \varphi(n g_\infty g_{\mathbf{f}}) \psi_T(n)^{-1} dn &= \int_{\mathbf{N}(\mathbb{Q}) \setminus \mathbf{N}(\mathbb{A})} \varphi(n^{-1} \gamma_\infty g_\infty \gamma_{\mathbf{f}} g_{\mathbf{f}}) \psi_T(n) dn \\ &= \int_{\mathbf{N}(\mathbb{Q}) \setminus \mathbf{N}(\mathbb{A})} \varphi(n \gamma_\infty g_\infty \gamma_{\mathbf{f}} g_{\mathbf{f}}) \psi_{-T}(n)^{-1} dn. \end{aligned}$$

Since $\gamma_\infty g_\infty \in \mathbf{G}(\mathbb{R})^\circ$, the last integral becomes $\mathbf{B}_\varrho^{-T}(\gamma_\infty g_\infty)(a_\varphi(-T; \gamma_\mathbf{f} g_\mathbf{f}))$ by the first case. Since $-T \notin \mathbf{V}(\mathbb{R})^+$, we have $a(-T; \gamma_\mathbf{f} g_\mathbf{f}) = 0$. \square

The vectors $a_\varphi(T; g_\mathbf{f})$ in V_ϱ are referred to as the adelic Fourier coefficients of φ . As is well-known, there is a linear bijective correspondence between $\varphi \in S_\varrho(\mathbf{K}_0(N))$ and the classical V_ϱ -valued Siegel cusp forms $f(Z)$ on $\mathbf{\Gamma}_0(N) := \{ \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \mathbf{Sp}_2(\mathbb{Z}) \mid C \equiv 0 \pmod{N} \}$ determined by the relation

$$f(g_\infty \langle i1_2 \rangle) = \nu(g_\infty)^{-(l_1+l_2)/2} \varrho(Ci + D) \varphi(g_\infty), \quad g_\infty = \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \mathbf{G}(\mathbb{R})^0.$$

By the modular transformation law $f((AZ + B)(CZ + D)^{-1}) = \varrho(CZ + D)^{-1} f(Z)$ for $\begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \mathbf{\Gamma}_0(N)$, since $-1_4 \in \mathbf{\Gamma}_0(N)$, we have that $f(Z)$ is identically 0, hence $S_\varrho(\mathbf{K}_0(N)) = (0)$, unless $l_1 \equiv l_2 \pmod{2}$. (For more details, see [37, §3.2] and [4, §4].)

Since $\mathbf{N}(\mathbb{A}) \cap \mathbf{K}_0(N) = \{ \mathbf{n}(X) \mid X \in \mathbf{V}(\widehat{\mathbb{Z}}) \}$, by (4.6), we have $a_\varphi(T; 1) = 0$ unless $X \in \mathcal{Q}$, where

$$\mathcal{Q} := \left\{ \begin{bmatrix} a & b/2 \\ b/2 & c \end{bmatrix} \mid a, b, c \in \mathbb{Z} \right\}$$

is the dual lattice of $\mathbf{V}(\mathbb{Z})$. Then, (4.5) with $g_\mathbf{f} = 1_4$ reduces to

$$f(Z) = \sum_{T \in \mathcal{Q}^+} a_\varphi(T; 1_4) \exp(2\pi i \text{tr}(TZ)), \quad Z \in \mathfrak{h}_2,$$

where $\mathcal{Q}^+ := \mathcal{Q} \cap \mathbf{V}(\mathbb{Q})^+$. This means $a_\varphi(T) := a_\varphi(T; 1_4)$ ($T \in \mathcal{Q}^+$) is the Fourier coefficient of $f(Z)$ in the classical sense. Set

$$\delta := \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \in \mathbf{GL}_2(\mathbb{Q}).$$

Note that the map $T \mapsto -T\mathbf{s}(\delta)$ preserves the set \mathcal{Q}^+ . If we identify \mathcal{Q}^+ with the set of positive definite integral binary forms $aX^2 + bXY + cY^2$, this operation corresponds to the sign change of b .

Lemma 4.2. *The set of adelic Fourier coefficients of $\varphi \in S_\varrho(\mathbf{K}_0(N))$ has the following properties:*

$$(4.7) \quad a_\varphi(T; \mathbf{m}(h_\mathbf{f}, \det h_\mathbf{f}) g_\mathbf{f} \kappa) = a_\varphi(T\mathbf{s}(h); g_\mathbf{f}), \quad h \in \mathbf{GL}_2(\mathbb{Q}), g_\mathbf{f} \in \mathbf{G}(\mathbb{A}_\mathbf{f}), \kappa \in \mathbf{K}_0(N).$$

$$(4.8) \quad a_\varphi(-T\mathbf{s}(\delta); 1) = \varrho(\delta) (a_\varphi(T; 1)), \quad T \in \mathbf{V}(\mathbb{Q}).$$

Proof. The relation in (4.6) follows from (4.7) by a simple change of variables. Let us show (4.8). We have

$$(4.9) \quad \mathbf{n}(-X\mathbf{s}(\delta)) = \kappa \mathbf{n}(X) \kappa^{-1}, \quad X \in \mathbf{V}(\mathbb{A})$$

with $\kappa := \mathbf{m}(\delta, 1) \in \mathbf{G}(\mathbb{Q})$. Write $\kappa = \kappa_\mathbf{f} \kappa_\infty \in \mathbf{G}(\mathbb{A}_\mathbf{f}) \mathbf{G}(\mathbb{R})$; then, $\kappa_\mathbf{f} \in \mathbf{K}_0(N)$ and $\kappa_\infty \in \mathbf{K}_\infty$. We have

$$\begin{aligned} \int_{\mathbf{N}(\mathbb{Q}) \backslash \mathbf{N}(\mathbb{A})} \varphi(n) \psi_{-T\mathbf{s}(\delta)}(n)^{-1} dn &= \int_{\mathbf{V}(\mathbb{Q}) \backslash \mathbf{V}(\mathbb{A})} \varphi(\mathbf{n}(X)) \psi(\text{tr}(-(T\mathbf{s}(\delta)) X))^{-1} dX \\ &= \int_{\mathbf{V}(\mathbb{Q}) \backslash \mathbf{V}(\mathbb{A})} \varphi(\mathbf{n}(-\mathbf{s}(\delta) X)) \psi(\text{tr}(TX))^{-1} dX \\ &= \int_{\mathbf{V}(\mathbb{Q}) \backslash \mathbf{V}(\mathbb{A})} \varphi(\kappa \mathbf{n}(X) \kappa^{-1}) \psi(\text{tr}(TX))^{-1} dX \end{aligned}$$

$$= \varrho(\delta) \int_{\mathbf{N}(\mathbb{Q}) \backslash \mathbf{N}(\mathbb{A})} \varphi(n) \psi_T(n)^{-1} \mathrm{d}n.$$

From this and (4.6),

$$\mathbf{B}_\varrho^{-T\mathbf{s}(\delta)}(1_4)(a_\varphi(-T\mathbf{s}(\delta); 1)) = \varrho(\delta) \mathbf{B}_\varrho^T(1_4)(a_\varphi(T; 1)).$$

By (4.1), $\mathbf{B}_\varrho^{-T\mathbf{s}(\delta)}(1_4) = \mathbf{B}_\varrho^T(1_4) = e^{-2\pi \mathrm{tr}(T)} \mathrm{Id}_{V_\varrho}$. \square

Define

$$(4.10) \quad \mathcal{Q}_{\mathrm{prim}}^+(D) := \{T = \begin{bmatrix} a & 2^{-1}b \\ 2^{-1}b & c \end{bmatrix} \in \mathcal{Q} \mid a > 0, -\det T = D/4\}.$$

Since D is a fundamental discriminant, $T \in \mathcal{Q}_{\mathrm{prim}}^+(D)$ is primitive, i.e., $\gcd(a, b, c) = 1$; $\mathcal{Q}_{\mathrm{prim}}^+(D)$ is a subset of $\mathbf{V}(\mathbb{Q})$, which is stable under the action of the unimodular group $\mathbf{SL}_2(\mathbb{Z})$ induced by (2.1). The matrix T_θ defined by (2.11) belongs to $\mathcal{Q}_{\mathrm{prim}}^+(D)$. The $\mathbf{SL}_2(\mathbb{Z})$ -orbit of $T \in \mathcal{Q}_{\mathrm{prim}}^+(D)$ is denoted by $[T]$.

Lemma 4.3. *For each $u \in \mathbb{A}_E^\times$, let $I_\theta(u)$ decompose in $\mathbf{GL}_2(\mathbb{A})$ as*

$$(4.11) \quad I_\theta(u) = \gamma_u h_u \kappa_u, \quad \gamma_u \in \mathbf{GL}_2(\mathbb{Q}), h_u \in \mathbf{GL}_2(\mathbb{R})^\circ, \kappa_u = (\kappa_{u,p})_{p < \infty} \in \mathbf{GL}_2(\widehat{\mathbb{Z}})$$

and set $T_\theta(u) := T_\theta \mathbf{s}(\gamma_u)$. Then ,

- (i) $T_\theta(u) \in \mathcal{Q}_{\mathrm{prim}}^+(D)$.
- (ii) The $\mathbf{SL}_2(\mathbb{Z})$ -equivalence class of $T_\theta(u)$ does not depend on the decomposition (4.11). If u, u' belongs to the same $E^\times E_\infty^\times \widehat{\mathfrak{o}}_E^\times$ -coset, then $T_\theta(u)$ and $T_\theta(u')$ are $\mathbf{SL}_2(\mathbb{Z})$ -equivalent.
- (iii) The map $[u] \mapsto [T_\theta(u)]$ from $\mathrm{Cl}(E) = \mathbb{A}_E^\times / E^\times E_\infty^\times \widehat{\mathfrak{o}}_E^\times$ to $\mathcal{Q}_{\mathrm{prim}}^+(D) / \mathbf{SL}_2(\mathbb{Z})$ is a bijection.

Proof. This is an adelic reformulation of the classically well-known correspondence between the ideal classes and the $\mathbf{SL}_2(\mathbb{Z})$ -equivalence classes of integral binary quadratic forms. The proof is straight-forward. \square

Since $\mathfrak{m}(\mathbf{GL}_2(\widehat{\mathbb{Z}}), 1) \subset \mathbf{K}_0(N)$, the relation in (4.7) shows that the function $T \mapsto a_\varphi(T) := a_\varphi(T; 1_4)$ on $\mathbf{V}(\mathbb{Z})^+$ is $\mathbf{SL}_2(\mathbb{Z})$ -invariant. Hence the following sum of vectors in V_ϱ is well-defined:

$$(4.12) \quad \mathbf{R}(\varphi, E, \Lambda) := \sum_{[u] \in \mathrm{Cl}(E)} a_\varphi([T_\theta(u)]) \Lambda(u)^{-1}.$$

Recall the integral in (3.6) and the function in (4.1).

Lemma 4.4. *Let $\varphi \in S_\varrho(\mathbf{K}_0(N))$.*

- (i) For $g_\infty \in \mathbf{G}(\mathbb{R})^0$,

$$B^{T_\theta, \Lambda}(\varphi; g_\infty) = w_D^{-1} \mathbf{B}_\varrho^{T_\theta}(g_\infty)(\mathbf{R}(\varphi, E, \Lambda)).$$

- (ii) We have

$$(4.13) \quad \varrho(r)(B^{T_\theta, \Lambda}(\varphi; b_\mathbb{R}^\theta)) = B^{T_\theta, \Lambda}(\varphi; b_\mathbb{R}^\theta), \quad r \in \mathbf{SO}(2),$$

$$(4.14) \quad \varrho(\delta)(B^{T_\theta, \Lambda}(\varphi; b_\mathbb{R}^\theta)) = B^{-T_\theta \mathbf{s}(\sigma), \Lambda}(\varphi; b_\mathbb{R}^\theta),$$

where $\sigma := \begin{bmatrix} 1 & t \\ 0 & -1 \end{bmatrix}$.

Proof. (i) The set $\mathbb{A}_E^\times/E^\times E_\mathbb{R}^\times$ is decomposed to a disjoint union of $\widehat{\mathfrak{o}}_E^\times$ -orbits $[u]\widehat{\mathfrak{o}}_E^\times$ for different classes $[u] \in \text{Cl}(E)$. Let μ denote the quotient measure on $\mathbb{A}_E^\times/E^\times E_\mathbb{R}^\times$. For $\tau \in \mathbb{A}_E^\times$, set $\mathbf{m}_\theta(\tau) := \mathbf{m}(I_\theta(\tau), N_{E/\mathbb{Q}}(\tau))$. Then by (3.6) and (4.6),

$$\begin{aligned} B^{T_\theta, \Lambda}(\varphi; g_\infty) &= \mathbf{B}_\varrho^{T_\theta}(g_\infty) \left(\int_{\mathbb{A}_E^\times/E^\times E_\mathbb{R}^\times} a_\varphi(T_\theta; \mathbf{m}_\theta(u)) \Lambda(u)^{-1} d\mu(u) \right) \\ &= \mathbf{B}_\varrho^{T_\theta}(g_\infty) \left(\sum_{[u] \in \text{Cl}(E)} \int_{[u]\widehat{\mathfrak{o}}_E^\times} a_\varphi(T_\theta; \mathbf{m}_\theta(u)) \Lambda(u)^{-1} d\mu(u) \right). \end{aligned}$$

Since $\mathbf{m}_\theta(\widehat{\mathfrak{o}}_E^\times) \subset \mathbf{K}_0(N)$, by (4.6), the formula in the parenthesis becomes

$$\sum_{[u] \in \text{Cl}(E)} \mu([u]\widehat{\mathfrak{o}}_E^\times) a_\varphi(T_\theta; \mathbf{m}_\theta(u)) \Lambda(u)^{-1}.$$

Let $I_\theta(u)$ decompose as in (4.11). Then, by (4.7) and Lemma 4.3, we have

$$a_\varphi(T_\theta; \mathbf{m}_\theta(u)) = a_\varphi(T_\theta \mathbf{s}(\gamma_u); 1) = a_\varphi(T_\theta(u)).$$

It remains to compute $\mu([u]\widehat{\mathfrak{o}}_E^\times)$. Since any fiber of the natural surjection from $\widehat{\mathfrak{o}}_E^\times$ onto $[u]\widehat{\mathfrak{o}}_E^\times$ has a simply transitive action of the global unit group \mathfrak{o}_E^\times , which coincides with the set of roots of unity in E^\times , we have $\mu([u]\widehat{\mathfrak{o}}_E^\times) = (1/\#\mathfrak{o}_E^\times) = 1/w_D$.

(ii) We have $I_\theta(e^{i\alpha}) = A_\theta^{-1} \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix} A_\theta$, or equivalently

$$\mathbf{m}(I_\theta(e^{i\alpha}), 1) = b_\mathbb{R}^\theta \text{diag} \left(\begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}, \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix} \right) (b_\mathbb{R}^\theta)^{-1}$$

for $\alpha \in \mathbb{R}$. Due to this and $\Lambda_\infty = 1$, formula (4.13) is proved by a change of variables $\tau \rightarrow \tau u^{-1}$ in the integral (3.6). Since $b_\mathbb{R}^\theta \mathbf{m}(\delta, 1) (b_\mathbb{R}^\theta)^{-1} = \mathbf{m}(\sigma, 1)$, formula (4.14) is proved in the same way as the proof of Lemma 4.2. \square

Let $\varphi \in S_\varrho(\mathbf{K}_0(N)) - (0)$; then $l_1 \equiv l_2 \pmod{2}$ as noted above. As such, $\dim_{\mathbb{C}}(V_\varrho^{\text{SO}(2)}) = 1$; indeed, $V_\varrho^{\text{SO}(2)} = \mathbb{C} \mathbf{v}_\varrho^0$ with $\mathbf{v}_\varrho^0 := (X^2 + Y^2)^{(l_1-l_2)/2}$. Set

$$\mathbf{v}_\varrho^\theta := \left(\frac{-2}{\sqrt{|D|}} \right)^{\frac{l_1+l_2}{2}} \varrho(t A_\theta) \mathbf{v}_\varrho^0 = \left(\frac{-2}{\sqrt{|D|}} \right)^{\frac{l_1-l_2}{2}} (X^2 + tXY + NY^2)^{\frac{l_1-l_2}{2}}.$$

Note that $\mathbf{B}_\varrho^{T_\theta}(b_\mathbb{R}^\theta) = \left(\frac{-2}{\sqrt{|D|}} \right)^{-(l_1+l_2)/2} e^{-2\pi\sqrt{|D|}} \varrho(t A_\theta)^{-1}$, so that $\mathbf{B}_\varrho^{T_\theta}(b_\mathbb{R}^\theta)(\mathbf{v}_\varrho^\theta) = e^{-2\pi\sqrt{|D|}} \mathbf{v}_\varrho^0$.

Thus, as a corollary to Lemma 4.4, the vector $\mathbf{R}(\varphi, E, \Lambda) \in V_\varrho$ is non-zero if and only if $B^{T_\theta, \Lambda}(\varphi; b_\mathbb{R}^\theta) \neq 0$, in which case there exists a unique scalar $R(\varphi, E, \Lambda)$ such that $\mathbf{R}(\varphi, E, \Lambda) = R(\varphi, E, \Lambda) \mathbf{v}_\varrho^\theta$, or equivalently

$$(4.15) \quad B^{T_\theta, \Lambda}(\varphi; b_\mathbb{R}^\theta) = w_D^{-1} e^{-2\pi\sqrt{|D|}} R(\varphi, E, \Lambda) \mathbf{v}_\varrho^0.$$

Define a C^∞ -function $B_\varrho^{T_\theta} : \mathbf{G}(\mathbb{R})^0 \rightarrow V_\varrho$ as

$$(4.16) \quad B_\varrho^{T_\theta}(g_\infty) := \mathbf{B}_\varrho^{T_\theta}(g_\infty)(\mathbf{v}_\varrho^\theta), \quad g_\infty \in \mathbf{G}(\mathbb{R})^0.$$

4.1. Sign condition. The Galois group $\text{Gal}(E/\mathbb{Q})$ acts on $\text{Cl}(E)$ naturally, hence on the orbit space $\mathcal{Q}_{\text{prim}}^+(D)/\mathbf{SL}_2(\mathbb{Z})$ through the bijection in Lemma 4.3(iii). The following lemma describes the conjugate action on $\mathcal{Q}_{\text{prim}}^+(D)/\mathbf{SL}_2(\mathbb{Z})$ explicitly. Recall the element $\sigma \in \mathbf{GL}_2(\mathbb{Z})$ defined in Lemma 4.4 (ii).

Lemma 4.5. *For $u \in \text{Cl}_E$, the element $T_\theta(\bar{u})$ is $\mathbf{SL}_2(\mathbb{Z})$ -equivalent to $-T_\theta(u)\mathbf{s}(\delta)$. In other words, the conjugate action on $\mathcal{Q}_{\text{prim}}^+(D)/\mathbf{SL}_2(\mathbb{Z})$ is induced by the map*

$$T = \begin{bmatrix} a & 2^{-1}b \\ 2^{-1}b & c \end{bmatrix} \mapsto -T\mathbf{s}(\delta) = \begin{bmatrix} a & -2^{-1}b \\ -2^{-1}b & c \end{bmatrix}.$$

Proof. The defining formula of I_θ in (2.5) yields

$$I_\theta(a + \theta b) = \begin{bmatrix} 1 & \theta \\ 1 & \theta \end{bmatrix}^{-1} \begin{bmatrix} a + \theta b & 0 \\ 0 & a + \bar{\theta}b \end{bmatrix} \begin{bmatrix} 1 & \theta \\ 1 & \theta \end{bmatrix}, \quad a, b \in \mathbb{Q}.$$

Set $t := \text{tr}_{E/\mathbb{Q}}(\theta)$. Then, by $\bar{\theta} = t - \theta$, we have $a + b\bar{\theta} = a' + b'\theta$ with $a' = a + tb \in \mathbb{Q}$ and $b' = -b \in \mathbb{Q}$. Then, a computation reveals

$$(4.17) \quad I_\theta(a + b\bar{\theta}) = \sigma I_\theta(a + b\theta)\sigma^{-1}.$$

By the decomposition of $I_\theta(u)$ for $u \in \mathbb{A}_E^\times$ in (4.11), we may take $\gamma_{\bar{u}} = \sigma\gamma_u\sigma^{-1}$. Thus,

$$T_\theta(\bar{u}) = T_\theta\mathbf{s}(\gamma_{\bar{u}}) = T_\theta\mathbf{s}(\sigma\gamma_u\sigma^{-1}).$$

A computation shows $T_\theta\mathbf{s}(\sigma) = -T_\theta$. Hence,

$$T_\theta(\bar{u}) = -T_\theta\mathbf{s}(\gamma_u\sigma^{-1}) = -T_\theta(u)\mathbf{s}(\sigma^{-1}) = -T_\theta(u)\mathbf{s}(\delta)\mathbf{s}\left(\begin{bmatrix} 1 & t \\ 0 & 1 \end{bmatrix}\right).$$

The last matrix is $\mathbf{SL}_2(\mathbb{Z})$ -equivalent to $-T_\theta(u)\mathbf{s}(\delta)$. This completes the proof. \square

For $\Lambda \in \widehat{\text{Cl}(E)}$, recall its conjugate $\Lambda^\dagger \in \widehat{\text{Cl}(E)}$ (see (2.25)).

Lemma 4.6. *Let $\varphi \in S_\rho(\mathbf{K}_0(N))$ and $\Lambda \in \widehat{\text{Cl}(E)}$ and suppose $B^{T_\theta, \Lambda}(\varphi; b_{\mathbb{R}}^\theta) \neq 0$. Then,*

$$R(\varphi, E, \Lambda^\dagger) = (-1)^{l_2} R(\varphi, E, \Lambda).$$

Proof. This is proved by Lemma 4.5 and (4.15) and by $\varrho(\delta)\mathbf{v}_\rho^0 = (-1)^{l_2}\mathbf{v}_\rho^0$. Note that $\delta^{-1}\sigma = \begin{bmatrix} 1 & t \\ 0 & 1 \end{bmatrix} \in \mathbf{SL}_2(\mathbb{Z})$, so that $a_\varphi(-T_\theta\mathbf{s}(\sigma), 1) = a_\varphi(-T_\theta\mathbf{s}(\delta), 1)$. \square

4.2. Automorphic representations. Let $\Pi_{\text{cusp}}(\mathbf{Z}\backslash\mathbf{G})$ denote the set of all those irreducible cuspidal representation of $\mathbf{G}(\mathbb{A})$ with trivial central characters, i.e., those irreducible $(\mathfrak{g}, \mathbf{K}_\infty) \times \mathbf{G}(\mathbb{A}_f)$ -submodules (π, V_π) of the space of cusp forms on $\mathbf{Z}(\mathbb{A})\mathbf{G}(\mathbb{Q})\backslash\mathbf{G}(\mathbb{A})$. We endow V_π with the restriction of the L^2 -inner product

$$(\varphi|\varphi_1)_{L^2} := \int_{\mathbf{Z}(\mathbb{A})\mathbf{G}(\mathbb{Q})\backslash\mathbf{G}(\mathbb{A})} \varphi(g)\overline{\varphi_1(g)} dg, \quad \varphi, \varphi_1 \in V_\pi.$$

For $\pi \in \Pi_{\text{cusp}}(\mathbf{Z}\backslash\mathbf{G})$, we fix its restricted tensor decomposition $\pi \cong \otimes_{p \leq \infty} \pi_p$ with (π_p, V_{π_p}) being an irreducible admissible unitarizable representation of $\mathbf{G}(\mathbb{Q}_p)$ with trivial central character such that a \mathbf{K}_p -invariant vector $\xi_p^0 \in V_{\pi_p} - (0)$ is preassigned for almost all $p < \infty$. An element $\lambda \in (\mathbb{Z}^2)_{\text{dom}}$ is the highest weight of the minimal \mathbf{K}_∞ -type of a holomorphic discrete series representation (HDS for short) of $\mathbf{Sp}_2(\mathbb{R})$ if and only if $l_2 > 2$, in which case the Harish-Chandra parameter of the HDS is $(l_1 - 1, l_2 - 2)$. For such λ and $N \in \mathbb{Z}_{>0}$, let $\Pi_{\text{cusp}}(\lambda, N)$ be a subset of $\pi \in \Pi_{\text{cusp}}(\mathbf{Z}\backslash\mathbf{G})$ having the following properties:

- (i) As a $(\mathfrak{g}, \mathbf{K}_\infty)$ -module $\pi_\infty \cong D_{l_1-1, l_2-2} \oplus D_{-l_2+2, -l_1+1}$, where D_{m_1, m_2} is the discrete series representation of $\mathbf{G}(\mathbb{R})^0$ of Harish-Chandra parameter (m_1, m_2) with central character $\mathbf{Z}(\mathbb{R}) \cong \mathbb{R}^\times \ni z \mapsto \text{sgn}(z)^{m_1+m_2+1}$.
- (ii) $V_\pi^{\mathbf{K}_0(N)} \neq (0)$.
- (iii) Let (ϱ, V_ϱ) be the irreducible rational representation of highest weight λ as before. The space $S_\varrho(\mathbf{K}_0(N))$ is an orthogonal direct sum of spaces $V_\pi^{\mathbf{K}_0(N)}[\varrho] := \{\varphi \in S_\varrho(\mathbf{K}_0(N)) \mid \varphi_v \in V_\pi (\forall v \in V_\varrho)\}$ for $\pi \in \Pi_{\text{cusp}}(\lambda, N)$, where $\varphi_v(g) := (\varphi(g) \mid v)_\varrho$ for $\varphi \in S_\varrho(\mathbf{K}_0(N))$ and $v \in V_\varrho$.

Although there may be many choices of $\Pi_{\text{cusp}}(\lambda, N)$, we fix one of them once and for all. For $\pi \in \Pi_{\text{cusp}}(\lambda, N)$, we fix a \mathbf{K}_∞ -intertwining map $V_\varrho \oplus \overline{V}_\varrho \hookrightarrow D_{l_1-1, l_2-2} \oplus D_{-l_2+2, -l_1+1} (\cong \pi_\infty)$ once and for all, where $\overline{V}_\varrho := \overline{\mathbb{C}} \otimes_{\mathbb{C}} V_\varrho$ is the complex conjugate of V_ϱ .

4.3. Basic assumptions. From now on, we fix a pair $(\lambda = (l_1, l_2), N) \in (\mathbb{Z}^2)_{\text{dom}} \times \mathbb{Z}_{>0}$ and a character $\mu = \otimes_{p \leq \infty} \mu_p : \mathbb{A}^\times / \mathbb{Q}^\times \mathbb{R}^\times \rightarrow \mathbb{C}^1$ with $M := \text{cond}(\mu)$ satisfying the following conditions:

- (A-i) N is square free.
- (A-ii) $l_1 \equiv l_2 \pmod{2}$, and $l_2 \in \mathbb{Z}_{\geq 3}$ so that $(l_1 - 1, l_2 - 2)$ is the Harish-Chandra parameter of a HDS.
- (A-iii) M and N are coprime.
- (A-iv) M is odd.
- (A-v) All prime divisors of NM are inert in E/\mathbb{Q} .

As usual, the character μ induces an even primitive Dirichlet character $\tilde{\mu} : (\mathbb{Z}/M\mathbb{Z})^\times \rightarrow \mathbb{C}^1$ in such a way that $\mu(u) = \tilde{\mu}(a)^{-1}$ for all $u \in \widehat{\mathbb{Z}}^\times$ and $a \in \mathbb{Z}$ with $u - a \in M\widehat{\mathbb{Z}}$; thus $\mu_p(p) = \tilde{\mu}(p)$ for primes $p \nmid M$, and

$$\prod_{p \mid M} W_{\mathbb{Q}_p}(\mu_p, \psi_p) = M^{-1/2} G(\tilde{\mu}),$$

with $W_{\mathbb{Q}_p}(\mu_p, \psi_p)$ being as in Definition 2 and $G(\tilde{\mu}) := \sum_{a \in (\mathbb{Z}/M\mathbb{Z})^\times} \tilde{\mu}(a) e^{2\pi i a/M}$, the Gauss sum of $\tilde{\mu}$. Let $\Pi_{\text{cusp}}^{(T_\theta, \Lambda)}(\lambda, N)$ denote the set of all those $\pi \in \Pi_{\text{cusp}}(\lambda, N)$ that satisfies the condition

- (A-vi) π admits a global (T_θ, Λ) -Bessel model, i.e., there exists $\varphi \in V_\pi$ such that $B^{T_\theta, \Lambda}(\varphi; g) \neq 0$ for some $g \in \mathbf{G}(\mathbb{A})$.

We remark that the conditions listed above (when $\mu = \mathbf{1}$ and $l_1 = l_2$) are also imposed in the most part of [7]. Let $\Pi_{\text{cusp}}^{(T_\theta, \Lambda)}(\lambda, N)^{\text{SK}}$ denote the set of $\pi \in \Pi_{\text{cusp}}^{(T_\theta, \Lambda)}(\lambda, N)$ that is the Saito-Kurokawa lift ([29]) of an cuspidal automorphic representation of $\mathbf{PGL}_2(\mathbb{A})$, which is locally described by [38] (see also [39]). From [38, §4], the set $\Pi_{\text{cusp}}^{(T_\theta, \Lambda)}(\lambda, N)^{\text{SK}} = \emptyset$ unless $l_1 = l_2$, i.e., V_ϱ is one dimensional. We note that all the representations $\pi \cong \otimes_p \pi_p$ in $\Pi_{\text{cusp}}^{(T_\theta, \chi)}(l, N) \setminus \Pi_{\text{cusp}}^{(T_\theta, \chi)}(l, N)^{\text{SK}}$ are non-CAP (cf. [42], [18, §3.5], [34, Corollary 4.5]); then, by [48], π_p is tempered for all $p \nmid N$. Moreover, by invoking [3] (see also [40]), any $\pi \in \Pi_{\text{cusp}}^{(T_\theta, \chi)}(l, N) \setminus \Pi_{\text{cusp}}^{(T_\theta, \chi)}(l, N)^{\text{SK}}$ is either a Yoshida lift, i.e., there exists a pair of irreducible cuspidal automorphic representations (σ_1, σ_2) of $\mathbf{GL}_2(\mathbb{A})$ such that $L(s, \pi) = L(s, \sigma_1)L(s, \sigma_2)$, or a “general type”, that is, there exists an irreducible cuspidal automorphic representation Π of $\mathbf{GL}_4(\mathbb{A})$ such that $L(s, \Pi, \wedge^2)$ has a pole at $s = 1$ and

$L(s, \pi) = L(s, \Pi)$. Let $\Pi_{\text{cusp}}^{(T_\theta, \chi)}(l, N)^Y$ (resp. $\Pi_{\text{cusp}}^{(T_\theta, \chi)}(l, N)^G$) be the set of $\pi \in \Pi_{\text{cusp}}^{(T_\theta, \chi)}(l, N)$ that is a Yoshida lift (resp. a general type). Then,

$$(4.18) \quad \Pi_{\text{cusp}}^{(T_\theta, \chi)}(l, N) = \Pi_{\text{cusp}}^{(T_\theta, \chi)}(l, N)^G \cup \Pi_{\text{cusp}}^{(T_\theta, \chi)}(l, N)^Y \cup \Pi_{\text{cusp}}^{(T_\theta, \chi)}(l, N)^{\text{SK}} \quad (\text{disjoint union}).$$

On the other hand, the set $\Pi_{\text{cusp}}^{(T_\theta, \Lambda)}(\lambda, N)$ can be separated into the subsets consisting of all “newforms” (i.e., $N_\pi = N$) and all “oldforms” (i.e., $N_\pi \neq N$) denoted by $\Pi_{\text{cusp}}^{(T_\theta, \Lambda)}(\lambda, N)^{\text{new}}$ and $\Pi_{\text{cusp}}^{(T_\theta, \Lambda)}(\lambda, N)^{\text{old}}$ respectively. For $\bullet \in \{G, Y, \text{SK}\}$ and $* \in \{\text{new}, \text{old}\}$, we set

$$\Pi_{\text{cusp}}^{(T_\theta, \Lambda)}(\lambda, N)^{\bullet, *} = \Pi_{\text{cusp}}^{(T_\theta, \Lambda)}(\lambda, N)^\bullet \cap \Pi_{\text{cusp}}^{(T_\theta, \Lambda)}(\lambda, N)^*.$$

Let N_π be the product of primes $p < \infty$ such that π_p is not spherical. By condition (ii) in §4.2, we have $N_\pi \mid N$. Due to (A-ii), the representations π_p for $p \mid N_\pi$ have non-zero $\mathbf{K}_0(p\mathbb{Z}_p)$ -fixed vectors, which, in turn, implies that π_p is Iwahori spherical so that their isomorphism classes are listed in [37, Table 3] (For extended tables, we refer to [35, Appendix].) Note that in [37] a different symplectic form is used to define the symplectic group; up to the adjustment for this difference, the group P_1 in [37] is our $\mathbf{K}_0(p\mathbb{Z}_p)$. In [35], all irreducible admissible representations of $\mathbf{G}(\mathbb{Q}_p)$ that admit local Bessel models are classified, and the result is conveniently summarized in [32, Table 2]. Here is a summary of what is available for our π (as in §4.3):

- π_∞ as a representation of $\mathbf{G}(\mathbb{R})^0$ is a direct sum of a HDS and its complex conjugate; as such, π is CAP if and only if it is a Saito-Kurokawa lift from a cuspidal representation of $\mathbf{PGL}_2(\mathbb{A})$; which happens only if $l_1 = l_2$.
- Suppose $p \nmid N_\pi$. Then, the local representation π_p is of type I and tempered if π is non-CAP, and is of type IIb when π is CAP.
- Suppose $p \mid N_\pi$. Then, the local representation π_p is either of type IIIa, in which case $\dim_{\mathbb{C}} V_{\pi_p}^{\mathbf{K}_0(p\mathbb{Z}_p)} = 2$, or of type VIb, in which case $\dim_{\mathbb{C}} V_{\pi_p}^{\mathbf{K}_0(p\mathbb{Z}_p)} = 1$; when π is CAP, π_p has to be of type VIb.

4.4. Bessel models. Let $p < \infty$; for any irreducible admissible representation (π_p, V_{π_p}) of $\mathbf{G}(\mathbb{Q}_p)$, let $(V_{\pi_p}^*)^{T_\theta, \Lambda}$ denote the space of all \mathbb{C} -linear forms $\ell : V_{\pi_p} \rightarrow \mathbb{C}$ that satisfies

$$\ell(\pi_p(m(I_\theta(\tau), N_{E/\mathbb{Q}}(\tau))n)\xi) = \Lambda(\tau)\psi_{T_\theta}(n)\ell(\xi), \quad \xi \in V_{\pi_p}, \tau \in E_p^\times, n \in \mathbf{N}(\mathbb{Q}_p).$$

It is known that $\dim_{\mathbb{C}}(V_{\pi_p}^*)^{T_\theta, \Lambda} \leq 1$ ([27], [28]). We say that π_p has a local (T_θ, Λ_p) -Bessel model if $(V_{\pi_p}^*)^{T_\theta, \Lambda} \neq (0)$; when this is the case, the space of functions of the form $g \mapsto \ell(\pi_p(g)\xi)$ with $\xi \in V_{\pi_p}$ is independent of $\ell \in (V_{\pi_p}^*)^{T_\theta, \Lambda_p} - (0)$; this space is denoted by $\mathcal{B}(T_\theta, \Lambda_p)[\pi_p]$ and is called the local (T_θ, Λ_p) -Bessel model of π_p . When π_p is spherical, it is known that π_p has a local (T_θ, Λ_p) -Bessel model, and the space $\mathcal{B}(T_\theta, \Lambda_p)[\pi_p]$ contains a unique \mathbf{K}_p -invariant function B_{π_p} such that $B_{\pi_p}^0(1_4) = 1$ ([43, Theorem 2-I], [6]). Let ξ_p^0 be the non-zero \mathbf{K}_p -fixed vector in V_{π_p} ; then there exists a unique element $\ell_{\pi_p}^0 \in (V_{\pi_p}^*)^{T_\theta, \Lambda_p} - (0)$ such that

$$\ell_{\pi_p}^0(\pi_p(g)\xi_p^0) = B_{\pi_p}^0(g), \quad g \in \mathbf{G}(\mathbb{Q}_p).$$

The pair $(\ell_{\pi_p}^0, \xi_p^0)$ is referred to as the unramified (T_θ, Λ_p) -Bessel datum for π_p .

Let $\pi \cong \otimes_{p \leq \infty} \pi_p \in \Pi_{\text{cusp}}^{(T_\theta, \Lambda)}(\lambda, N)$ (see §4.3). Let N_π be the product of $p \mid N$ such that π_p is not spherical.

Definition 1. A system $\{(\ell_p, \xi_p)\}_{p < \infty}$ with $\ell_p \in (V_{\pi_p}^*)^{T_\theta, \Lambda_p} - (0)$, $\xi_p \in V_{\pi_p}^{\mathbf{K}_0(N_\pi \mathbb{Z}_p)}$ is called a (T_θ, Λ) -Bessel data for π if $(\ell_p, \xi_p) = (\ell_{\pi_p}^0, \xi_{\pi_p}^0)$ for $(p, N_\pi) = 1$ and $\ell_p(\xi_p) = 1$ for all $p < \infty$.

By [32, Theorem 2 8.2 and 9.3], a (T_θ, Λ) -Bessel data exists for our π . Once a (T_θ, Λ) -Bessel data $\{(\ell_p, \xi_p)\}_{p < \infty}$ is fixed, one can define $\varphi_{\pi, \varrho}^0 \in V_\pi^{\mathbf{K}_0(N_\pi)}[\varrho]$ to be the V_ϱ -valued cusp form such that for any $v \in V_\varrho$, the function $(\varphi_{\pi, \varrho}^0(g) \mid v)_\varrho$ in $V_\pi \cong \bigotimes_{p \leq \infty} V_{\pi_p}$ corresponds to the pure tensor $\bar{v} \otimes (\bigotimes_{p < \infty} \xi_p)$, where $\bar{v} = 1 \otimes v \in \overline{V_\varrho} \hookrightarrow V_{\pi_\infty}$. A particular choice of $\{(\ell_p, \xi_p)\}$ will be made in §6.8 so that $\varphi_{\pi, \varrho}^0$ corresponds to a newform on the arithmetic quotient $\Gamma_0(N) \backslash \mathfrak{h}_2$ in the sense of [7, §3.2].

Lemma 4.7. Let $\{(\ell_p, \xi_p)\}_{p < \infty}$ be a (T_θ, Λ) -Bessel data for $\pi \in \Pi_{\text{cusp}}(\lambda, N)$. Let $\phi = \bigotimes_p \phi_p \in \mathcal{S}(\mathbb{A}_E^2)$ be a decomposable element. Then, for any $\varphi \in V_\pi^{\mathbf{K}_0(N)}[\varrho]$ such that, for any $v \in V_\varrho$, φ_v corresponds to the pure tensor $v \otimes (\bigotimes_{p < \infty} v_p) \in \bigotimes_{p \leq \infty} V_{\pi_p}$, we have

$$(4.19) \quad B^{T_\theta, \Lambda}(\varphi; g) = B^{T_\theta, \Lambda}(\varphi_{\pi, \varrho}^0; g_\infty) \prod_{p < \infty} \ell_p(\pi_p(g_p)v_p), \quad g = (g_p)_p \in \mathbf{G}(\mathbb{A}).$$

For $\text{Re}(s) > 1$, $v \in V_\varrho$ and $b_{\mathbf{f}} = (b_p)_{p < \infty} \in \mathbf{G}(\mathbb{A}_{\mathbf{f}})$,

$$(4.20) \quad \langle E(\phi, s, \Lambda, \mu), R(b_{\mathbf{f}} b_{\mathbb{R}}^\theta) \overline{\varphi_v} \rangle = \frac{\sqrt{|D|}}{2} Z_v^{(\infty)}(\phi_\infty, \varphi_\pi^0; s, \Lambda) \times \prod_{p \leq \infty} Z_p(\phi_p, \overline{B_{v_p}}; s, \mu_p; b_p),$$

where

$$Z_v^{(\infty)}(\phi_\infty, \varphi_\pi^0; s, \Lambda) := \int_{\mathbf{K}_\infty^\#} \int_{\mathbb{R}^\times} |a|_{\mathbb{R}}^{s-1} f_{\phi_\infty}^{(s, 1, 1)}(k_\infty^\#)(v \mid B^{T_\theta, \Lambda}(\varphi_{\pi, \varrho}^0; \mathbf{m}(a1_2, a)\iota_\theta(k_\infty^\#)b_{\mathbb{R}}^\theta))_\varrho d^\times a dk_\infty^\#,$$

$$Z_p(\phi_p, \overline{B_{v_p}}; s, \mu_p; b_p) := \int_{\mathbb{Q}_p^\times} \int_{\mathbf{K}_p^\#} f_{\phi_p}^{(s, \Lambda_p, \mu_p)}(k_p^\#)\mu_p(a)|a|_p^{s-1} \overline{B_{v_p}}(\mathbf{m}(a_p1_2, a_p)\iota_\theta(k_p^\#)b_p) d^\times a_p dk_p^\#$$

with

$$B_{v_p}(g_p) := \ell_p(\pi_p(g_p)v_p), \quad g_p \in \mathbf{G}(\mathbb{Q}_p).$$

Proof. Fix $g_\infty \in \mathbf{G}(\mathbb{R})^0$ and $v \in V_\varrho$ and regard $\varphi \mapsto B^{T_\theta, \Lambda}(\varphi_v; g_\infty)$ as a linear functional on $\bigotimes_{p < \infty} V_{\pi_p}$ by the natural inclusion $\bigotimes_{p < \infty} V_{\pi_p} \hookrightarrow v \otimes (\bigotimes_{p < \infty} V_{\pi_p}) \hookrightarrow V_\pi$. Then, by the local multiplicity one theorem for Bessel functional on $\mathbf{G}(\mathbb{Q}_p)$ recalled above, there exists $C_v(g_\infty) \in \mathbb{C}$ such that

$$(4.21) \quad (v \mid B^{T_\theta, \Lambda}(\varphi; g_\infty))_\varrho = C_v(g_\infty) \prod_{p < \infty} \overline{\ell_p(v_p)}$$

for φ corresponding to $\bigotimes_{p < \infty} v_p$. To determine $C_v(g_\infty)$, set $v_p = \xi_p$ for all $p < \infty$; then by $\ell_p(\xi_p) = 1$, we get $(v \mid B^{T_\theta, \Lambda}(\varphi_{\pi, \varrho}^0; g_\infty))_\varrho = C_v(g_\infty)$. Now we apply (3.7) with T_θ, Λ being replaced with $-T_\theta, \Lambda^{-1}$ and with φ being replaced with $R(b_{\mathbf{f}})\varphi$. \square

4.5. Gamma factor and global Bessel period. Recall the point $b_{\mathbb{R}}^\theta$ in (2.14) and the vector $v_\varrho^\theta \in V_\varrho$ in (4.16). Since $\nu(b_{\mathbb{R}}^\theta) > 0$, the point $\mathbf{m}(a1_2, a)b_{\mathbb{R}}^\theta$ with $a \in \mathbb{R}^\times$ belongs to $\mathbf{G}(\mathbb{R})^0$ if and only if $a > 0$. Thus, by (3.6) and (4.6), we have $B^{T_\theta, \Lambda}(\varphi_\pi^0; \mathbf{m}(a1_2, a)b_{\mathbb{R}}^\theta) = 0$ for $a < 0$. By Lemma 4.4 and (4.16),

$$B^{T_\theta, \Lambda}(\varphi_\pi^0; g_\infty) = w_D^{-1} R(\varphi_\pi^0, E, \Lambda) B_\varrho^{T_\theta}(g_\infty), \quad g_\infty \in \mathbf{G}(\mathbb{R})^0.$$

Substituting this, we have that $Z_v^{(\infty)}(\phi_\infty, \varphi_\pi^0; s, \Lambda)$ equals

$$(4.22) \quad w_D^{-1} \overline{R(\varphi_\pi^0, E, \Lambda)} \int_{\mathbf{K}_\infty^\#} \int_0^\infty |a|_{\mathbb{R}}^{s-1} f_{\phi_\infty}^{(s,1,1)}(k_\infty^\#)(v \mid B_\varrho^{T_\theta}(\mathbf{m}(a1_2, a)\iota_\theta(k_\infty^\#)b_\mathbb{R}^\theta))_\varrho d^\times a dk_\infty^\#.$$

Now we specify ϕ_∞ . Recall that the highest weight of ϱ is $\lambda = (l_1, l_2)$ and $l_1 - l_2 \in 2\mathbb{Z}_{>0}$. Set $d := l_1 - l_2$ and define

$$\mathbf{f}_\varrho(u) := (d+1) \frac{(\varrho(C\bar{u}C^{-1})\mathbf{v}_\varrho^0 \mid \mathbf{v}_\varrho^0)_\varrho}{(\mathbf{v}_\varrho^0 \mid \mathbf{v}_\varrho^0)_\varrho}, \quad u \in \mathbf{K}_\infty^\#.$$

with $C := \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ i & -i \end{bmatrix}$. If $u = \begin{bmatrix} a & -\bar{b} \\ b & \bar{a} \end{bmatrix} \in \mathbf{SU}(2)$ with $a = a' + ia''$, $b = b' + ib''$ and $A, B \in \mathbf{Mat}_2(\mathbb{R})$ are defined as in (2.16), so that $(b_\mathbb{R}^\theta)^{-1}\iota_\theta(u)(b_\mathbb{R}^\theta = k_\infty(A + iB)$ (Lemma 2.2), then a computation reveals $C\bar{u}C^{-1} = A - iB$. Thus, the automorphism $u \mapsto C\bar{u}C^{-1}$ of $\mathbf{SU}(2)$ brings the subgroup $\mathbf{B}^\#(\mathbb{R}) \cap (\mathbf{K}_\infty^\#)^0 = \{\begin{bmatrix} a & 0 \\ 0 & a^{-1} \end{bmatrix} \mid a \in \mathbb{C}^1\}$ to the subgroup $\mathbf{SO}(2)$, by which \mathbf{v}_ϱ^0 is fixed. Hence, \mathbf{f}_ϱ is left $\mathbf{B}^\#(\mathbb{R}) \cap (\mathbf{K}_\infty^\#)^0$ -invariant. Since \mathbf{v}_ϱ^0 is also fixed by the element $C\delta C^{-1} = \delta$, the function \mathbf{f}_ϱ is left $\mathbf{B}^\#(\mathbb{R}) \cap \mathbf{K}_\infty^\#$ -invariant. Thus, there exists a unique element $\mathbf{f}_\varrho^{(s)}$ of $\mathcal{V}^\#(s, 1, 1)$ such that $\mathbf{f}_\varrho^{(s)}|_{\mathbf{K}_\infty^\#} = \mathbf{f}_\varrho$. Define $\phi_\infty \in \mathcal{S}(\mathbb{C}^2)$ by

$$(4.23) \quad \phi_\infty(x, y) := \sum_{j=0}^{d/2} \frac{(x\bar{x})^j}{(j!)^2} \frac{(-y\bar{y})^{d/2-j}}{((d/2-j)!)^2} \frac{2}{\pi} \exp\left(-\frac{2\pi}{\sqrt{|D|}}(x\bar{x} + y\bar{y})\right), \quad (x, y) \in \mathbb{C}^2.$$

By (3.1),

$$(4.24) \quad f_{\phi_\infty}^{(s,1,1)} = |D|^{\frac{s+1}{2}} (-1)^{\frac{l_1-l_2}{2}} \Gamma_{\mathbb{C}}\left(s + \frac{l_1-l_2}{2} + 1\right) \mathbf{f}_\varrho^{(s)}.$$

For $u \in (\mathbf{K}_\infty^\#)^0$ and A, B as above, by (4.1) and (4.16) and by Lemma 2.2, we have

$$(v \mid B_\varrho^{T_\theta}(\mathbf{m}(a1_2, a)\iota_\theta(u)b_\mathbb{R}^\theta))_\varrho = (-1)^{\frac{l_1+l_2}{2}} a^{\frac{l_1+l_2}{2}} \exp\left(-2\sqrt{|D|}\pi a\right) (v \mid \varrho(C\bar{u}C^{-1})^{-1}\mathbf{v}_\varrho^0)_\varrho.$$

By using the orthogonal relation of matrix coefficients on $\mathbf{SU}(2)$, the integral in (4.22) is computed as

$$\begin{aligned} & |D|^{\frac{s}{2}} (-1)^{l_1} \Gamma_{\mathbb{C}}\left(s + \frac{l_1-l_2}{2} + 1\right) (v \mid \mathbf{v}_\varrho^0)_\varrho \int_0^\infty a^{s+\frac{l_1+l_2}{2}-1} \exp\left(-2\sqrt{|D|}\pi a\right) d^\times a \\ &= |D|^{\frac{1}{2}(1-\frac{l_1+l_2}{2})} (-1)^{l_1} (v \mid \mathbf{v}_\varrho^0)_\varrho \Gamma_{\mathbb{C}}\left(s + \frac{l_1-l_2}{2} + 1\right) \Gamma_{\mathbb{C}}\left(s + \frac{l_1+l_2}{2} - 1\right) \end{aligned}$$

for $\operatorname{Re}(s) + \frac{l_1+l_2}{2} > 1$. Recall $L(s, \pi_\infty) = \Gamma_{\mathbb{C}}(s + \frac{l_1-l_2}{2} + \frac{1}{2}) \Gamma_{\mathbb{C}}(s + \frac{l_1+l_2}{2} - \frac{3}{2})$. Thus,

$$(4.25) \quad Z_v^{(\infty)}(\phi_\infty, \varphi_\pi^0; s, \Lambda) = w_D^{-1} \overline{R(\varphi_\pi^0, E, \Lambda)} (v \mid \mathbf{v}_\varrho^0)_\varrho (-1)^{l_1} |D|^{\frac{1}{2}(1-\frac{l_1+l_2}{2})} L\left(s + \frac{1}{2}, \pi_\infty\right).$$

The formula in [47, 7.23 Lemma] yields

$$(4.26) \quad M(s) \mathbf{f}_\varrho^{(s)} = \frac{\pi}{s-d/2} \prod_{j=1}^{d/2} \frac{s-d/2+j-1}{s+d/2-j} \mathbf{f}_\varrho^{(-s)}.$$

By this combined with (3.3) and (4.24), we easily deduce

$$(4.27) \quad \widehat{f_{\phi_\infty}^{(-s,1,1)}} = \frac{|D|}{4} (-1)^{\frac{l_1-l_2}{2}} f_{\phi_\infty}^{(-s,1,1)}.$$

4.6. The spinor L -function and its functional equation. Let $\pi \cong \otimes_v \pi_v$ be a cuspidal automorphic representation of $\mathbf{G}(\mathbb{A})$ with the trivial central character: then, $\bar{\pi}_p \cong \pi_p^\vee \cong \pi_p$ for all $p < \infty$ by [44, Proposition 2.3]. The twist π_μ of π by an idele class character $\mu : \mathbb{A}^\times / \mathbb{Q}^\times \rightarrow \mathbb{C}^\times$ is defined on the space V_π of π as $\pi_\mu(g) := \pi(g) \cdot (\mu \circ \nu)(g)$ for $g \in \mathbf{G}(\mathbb{A})$, so that the central character of π_μ is μ^2 . Let $\pi \in \Pi_{\text{cusp}}^{(T_\theta, \Lambda)}(\lambda, N)$ and μ be as in §4.3, so that $\lambda = (l_1, l_2) \in \mathbb{Z}^2$, $l_1 \equiv l_2 \pmod{2}$, $l_1 \geq l_2 > 2$ and the ramification loci of π and μ are disjoint. We define the spinor L -function of π twisted by μ as the Euler product

$$L(s, \pi, \mu) := \prod_{p < \infty} L(s, (\pi_\mu)_p). \quad \text{Re}(s) > 5/2$$

with $L(s, (\pi_\mu)_p)$ the local L -factor listed in table (6.5). The completed L -function is defined as

$$\widehat{L}(s, \pi, \mu) := \Gamma_{\mathbb{C}}\left(s + \frac{l_1 - l_2}{2} + \frac{1}{2}\right) \Gamma_{\mathbb{C}}\left(s + \frac{l_1 + l_2}{2} - \frac{3}{2}\right) L(s, \pi, \mu).$$

Using Lemmas 3.1, 4.7 and Lemma 4.6, combining (4.25), (4.27) and the computations of the local zeta integrals for cases 1, 4, 5, and 6 in table (6.6), we obtain a meromorphic continuation of $L(s, \pi, \mu)$ to \mathbb{C} as well as the functional equation

$$\widehat{L}(s, \pi, \mu) = \varepsilon(s, \pi, \mu) \widehat{L}(1 - s, \pi, \bar{\mu}) \quad \text{with} \quad \varepsilon(s, \pi, \mu) := (-1)^{l_2} \tilde{\mu}(N_\pi^2) \left(\frac{G(\bar{\mu})}{\sqrt{M}}\right)^4 (M^4 N_\pi^2)^{1/2-s}$$

as expected. Moreover $\widehat{L}(s, \pi, \mu)$ is holomorphic except for possible simple poles at $s = 3/2, -1/2$, which does not occur when μ is non-trivial. By (4.26), $M(s) \mathbf{f}_\varrho^{(s)}$ has a simple zero at $s = 1$ if $l_1 > l_2$, which in turn implies the holomorphy at $s = 1$ of the global intertwining operator applied to the section $f_\phi^{(s)}$ as well as the Eisenstein series $E(\phi, s, \Lambda, \mu)$ for ϕ_∞ as above. Hence, by (4.20) with an appropriate ϕ , $\widehat{L}(s, \pi, \mu)$ is holomorphic at $s = 3/2$ when $l_1 > l_2$. If μ is real valued, then $G(\mu) = \sqrt{M}$ and $\varepsilon(1/2, \pi, \mu) = (-1)^{l_2}$, so that $L(1/2, \pi, \mu) = 0$ unless l_2 is even.

5. SPECTRAL AVERAGE OF RANKIN-SELBERG INTEGRALS

The space $S_\varrho(\mathbf{K}_0(N))$ is endowed with the Hermitian inner product associated to the norm $\int_{\mathbf{Z}(\mathbb{A}) \mathbf{G}(\mathbb{Q}) \backslash \mathbf{G}(\mathbb{A})} (\varphi(g) \mid \varphi(g))_\varrho dg$ ($\varphi \in S_\varrho(\mathbf{K}_0(N))$). Let $\mathcal{H}(\mathbf{G}(\mathbb{Q}_p) \parallel \mathbf{K}_p)$ be the Hecke algebra for $(\mathbf{G}(\mathbb{Q}_p), \mathbf{K}_p)$ for $p < \infty$. For any finite set S of primes p that is prime to N , define $\mathcal{H}_S := \otimes_{p \in S} \mathcal{H}(\mathbf{G}(\mathbb{Q}_p) \parallel \mathbf{K}_p)$. The \mathbb{C} -algebra \mathcal{H}_S acts on the finite dimensional Hilbert space $S_\varrho(\mathbf{K}_0(N))$ normally by

$$[R(f_S)\varphi](g) = \int_{\mathbf{G}(\mathbb{Q}_S)} \varphi(gx_S) f_S(x_S) dx_S, \quad g \in \mathbf{G}(\mathbb{A}), \quad f_S \in \mathcal{H}_S, \varphi \in S_l(\mathbf{K}_0(N)).$$

where $\mathbf{G}(\mathbb{Q}_S) := \prod_{p \in S} \mathbf{G}(\mathbb{Q}_p)$. Let μ and M be as in §4.3. We define the Schwartz-Bruhat function $\phi \in \mathcal{S}(\mathbb{A}_E^2)$ associated μ as

$$\phi = \prod_{p \leq \infty} \phi_p, \quad \phi_p(x, y) = \begin{cases} \mathbb{1}_{\mathfrak{o}_{E_p}}(x) \mathbb{1}_{\mathfrak{o}_{E_p}}(y) & (p < \infty, p \nmid M) \\ \mathbb{1}_{p^e \mathfrak{o}_{E_p}}(x) \mathbb{1}_{1+p^e \mathfrak{o}_{E_p}}(y) & (p < \infty, p^e \parallel M, e \geq 1), \end{cases}$$

and ϕ_∞ being as in (4.23). In this section, we investigate the averages

$$(5.1) \quad \mathbb{I}^{(s)}(l, N, f_S) := \frac{1}{[\mathbf{K}_f : \mathbf{K}_0(N)]} \sum_{\varphi \in \mathcal{B}(\lambda, N)} \langle E(\phi, s, \Lambda, \mu), R(b)R(f_S)\overline{\varphi_{\mathbf{v}_\ell^0}} \rangle B^{T_\theta, \Lambda}(\varphi_{\mathbf{v}_\ell^0}; b_\mathbb{R}^\theta),$$

where $\mathbf{v}_\ell^0 \in V_\ell^{\mathbf{SO}(2)}$ is the vector from §4 and $b = (b_p)_{p \leq \infty} \in \mathbf{G}(\mathbb{A})$ is define by

$$(5.2) \quad b_p = \begin{cases} 1_4 & (p < \infty, p \nmid NM), \\ \eta_p & (p < \infty, p \mid N), \\ b_p^M := \begin{bmatrix} p^e 1_2 & T_\theta^\dagger \\ 0 & 1_2 \end{bmatrix} & (p < \infty, p^e \parallel M, e \geq 1), \\ b_\mathbb{R}^\theta & (p = \infty) \end{cases}$$

with $\eta_p := \begin{bmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & p & 0 & 0 \\ -p & 0 & 0 & 0 \end{bmatrix} \in \mathbf{G}(\mathbb{Q}_p)$ being the Atkin-Lehner element, $\mathbf{K}_f := \mathbf{K}_0(1)$, and $\mathcal{B}(\lambda, N) = \bigcup_{\pi \in \Pi_{\text{cusp}}(\lambda, N)} \mathcal{B}_\pi(\lambda, N)$ is an orthonormal basis of $S_\ell(\mathbf{K}_0(N))$ with $\mathcal{B}_\pi(\lambda, N)$ being an orthonormal basis of $V_\pi^{\mathbf{K}_0(N)}[\rho]$. The sum is independent of the choice of an orthonormal basis and can be written as $\mathbb{I}^{(s)}(\Pi_{\text{cusp}}^{(T_\theta, \Lambda)}(\lambda, N), f_S)$, where for any subset $X \subset \Pi_{\text{cusp}}^{(T_\theta, \Lambda)}(\lambda, N)$ we define

$$(5.3) \quad \mathbb{I}^{(s)}(X, f_S) = \frac{1}{[\mathbf{K}_f : \mathbf{K}_0(N)]} \sum_{\pi \in X} \widehat{f_S}(\pi_S) \sum_{\varphi \in \mathcal{B}_\pi(\lambda, N)} \langle E(\phi, s, \Lambda, \mu), R(b)\overline{\varphi_{\mathbf{v}_\ell^0}} \rangle B^{T_\theta, \Lambda}(\varphi_{\mathbf{v}_\ell^0}; b_\mathbb{R}^\theta),$$

where $\widehat{f_S}(\pi_S)$ is the spherical Fourier transforms of f_S on $\pi_S := \otimes_{p \in S} \pi_p$, which is defined as the eigenvalue of the operator $\pi_S(f_S)$ on the $\mathbf{K}_S = \prod_{p \in S} \mathbf{K}_p$ -fixed vectors $\pi_S^{\mathbf{K}_S} \cong \mathbb{C}$. For $\bullet \in \{\text{T, G, Y, SK}\}$ and $* \in \{\text{new, old}\}$, we define $\mathbb{I}^{(s)}(\lambda, N, f_S)^\bullet$, $\mathbb{I}^{(s)}(\lambda, N, f_S)^*$, and $\mathbb{I}^{(s)}(\lambda, N, f_S)^{\bullet,*}$ to be $\mathbb{I}^{(s)}(X, f_S)$ with $X = \Pi_{\text{cusp}}^{(T_\theta, \Lambda)}(\lambda, N)^\bullet$, $\Pi_{\text{cusp}}^{(T_\theta, \Lambda)}(\lambda, N)^*$ and $\Pi_{\text{cusp}}^{(T_\theta, \Lambda)}(\lambda, N)^{\bullet,*}$, respectively. Then, due to (4.18), the average (5.1) is rearranged as

$$(5.4) \quad \mathbb{I}^{(s)}(\lambda, N, f_S) = \mathbb{I}^{(s)}(\lambda, N, f_S)^{\text{G, new}} + \mathbb{I}^{(s)}(\lambda, N, f_S)^{\text{Y, new}} \\ + \mathbb{I}^{(s)}(\lambda, N, f_S)^{\text{SK, new}} + \mathbb{I}^{(s)}(\lambda, N, f_S)^{\text{old}}.$$

5.1. A construction of orthonormal basis. Let $\pi \cong \otimes_{p \leq \infty} \pi_p$ be an element of the set $\Pi_{\text{cusp}}^{(T_\theta, \Lambda)}(\lambda, N)$ (see §4.3). We fix a (T_θ, Λ) -Bessel data $\{(\ell_p, \xi_p)\}_{p < \infty}$ of π (see §4.4) once and for all. Set $\varphi_\pi^0(g) := (\varphi_{\pi, \ell}^0(g) \mid \mathbf{v}_{\ell^0})_\ell$, which is an element of $V_\pi^{\mathbf{K}_0(N)}$. Since π is a unitary representation, all of its factors π_p are unitarizable. For each $p < \infty$, we can uniquely fix a $\mathbf{G}(\mathbb{Q}_p)$ -invariant inner product $(\cdot | \cdot)_p$ on V_{π_p} by demanding $(\xi_p | \xi_p)_p = 1$; for $p = \infty$, we fix a $\mathbf{G}(\mathbb{R})$ -invariant inner product so that its pull back to $V_\ell \hookrightarrow \pi_\infty$ (cf. §4.2(i)) coincides with the inner product $(\cdot | \cdot)_\ell$ of V_ℓ . Let φ_π^0 be the global new form attached to $\{\xi_p\}_{p < \infty}$ (see §4.4). Then,

$$(5.5) \quad \frac{(\varphi | \varphi)_{L^2}}{(\varphi_\pi^0 | \varphi_\pi^0)_{L^2}} = (v | v)_\ell \prod_{p < \infty} (v_p | v_p)_p$$

for any $\varphi \in V_\pi$ that corresponds to $v \otimes (\otimes_{p < \infty} v_p)$ with $v \in V_\varrho$. As below, for $p < \infty$, we fix an orthonormal basis $\mathcal{B}(\pi_p, \mathbf{K}_0(N\mathbb{Z}_p))$ of $V_{\pi_p}^{\mathbf{K}_0(N\mathbb{Z}_p)}$ in such a way that

$$\mathcal{B}(\pi_p, \mathbf{K}_0(N\mathbb{Z}_p)) = \{\xi_{\pi_p}^0\} \quad \text{if } p \nmid N.$$

Then, a pure tensor of the form

$$(5.6) \quad (\varphi_\pi^0 | \varphi_\pi^0)_{L^2}^{-1/2} \times \otimes_{p < \infty} v_p, \quad v_p \in \mathcal{B}(\pi_p, \mathbf{K}_0(N\mathbb{Z}_p)),$$

yield an element $\varphi \in V_\pi^{\mathbf{K}_0(N)}[l]$ such that for $v \in V_\varrho$ the element $\varphi_v \in V_\pi$ corresponds to v tensored with (5.6). The set of functions φ obtained in this way from (5.6) will be denoted by $\mathcal{B}_\pi(\lambda, N)$. If $\pi \in \Pi_{\text{cusp}}(\lambda, N)$ does not satisfy condition (A-iv) in §4.3, then we fix arbitrary orthonormal basis $\mathcal{B}_\pi(\lambda, N)$ of the space $V_\pi^{\mathbf{K}_0(N)}[\lambda]$. Let $\mathcal{B}(\lambda, N)$ be the union of the sets $\mathcal{B}_\pi(\lambda, N)$ for $\pi \in \Pi_{\text{cusp}}(\lambda, N)$; then, by (ii) in §4.2, the set $\mathcal{B}(\lambda, N)$ is an orthonormal basis of $S_\varrho(\mathbf{K}_0(N))$.

5.2. Computation of the average. Let φ correspond to a pure tensor as in (5.6). Then, by (4.20) and by the computations recalled in §4.5 and §6.2,

$$(5.7) \quad \langle E(\phi, s, \Lambda, \mu), R(b) \overline{\varphi_{v_\varrho}^0} \rangle = (\varphi_\pi^0 | \varphi_\pi^0)_{L^2}^{-1/2} w_D^{-1} \overline{R(\varphi_\pi^0, E, \Lambda)} \widehat{L}(s + \tfrac{1}{2}, \pi, \mu) \\ \times (\mathbf{v}_\varrho^0 | \mathbf{v}_\varrho^0)_\varrho (-1)^{l_2} 2^{-2} |D|^{\frac{1}{2}(4 - \frac{l_1 + l_2}{2})} \prod_{p|N} Z_p^*(\phi_p, \overline{B_{v_p}}; s, \mu_p; b_p),$$

where

$$Z_p^*(\phi_p, \overline{B_{v_p}}; s, \mu_p; b_p) := L(s + \tfrac{1}{2}, \pi_p, \mu_p)^{-1} Z_p(\phi_p, \overline{B_{v_p}}; s, \mu_p; b_p)$$

is the normalized local zeta-integral. Moreover, by (4.21) and Lemma 4.4,

$$(5.8) \quad B^{T_\theta, \Lambda}(\varphi_{\mathbf{v}_\varrho^0}^0; b_\mathbb{R}^\theta) = \langle \varphi_\pi^0 | \varphi_\pi^0 \rangle_{L^2}^{-1/2} w_D^{-1} R(\varphi_\pi^0, E, \Lambda) (B_\varrho^{T_\theta}(b_\mathbb{R}^\theta) | \mathbf{v}_\varrho^0)_\varrho \prod_{p|N} \ell_p(v_p).$$

Note that $B_\varrho^{T_\theta}(b_\mathbb{R}^\theta) = (-1)^{\frac{l_1 + l_2}{2}} \exp(-2\pi\sqrt{|D|}) \mathbf{v}_\varrho^0$ by (4.1) and (2.14). From (5.3), (5.7) and (5.8), we get

Proposition 5.1. *Let $f_S \in \mathcal{H}_S$. Then, $\mathbb{I}^{(s)}(\lambda, N, f_S)$ equals to*

$$(5.9) \quad \frac{2^{-2} |D|^{\frac{1}{2}(3 - \frac{l_1 + l_2}{2})} e^{-2\pi\sqrt{|D|}}}{w_D^2[\mathbf{K}_f : \mathbf{K}_0(N)]} \times M^{s-6} \zeta_M(1) \zeta_M(4) \tilde{\mu}(2D) G(\tilde{\mu}) \\ \times (\mathbf{v}_\varrho^0 | \mathbf{v}_\varrho^0)_\varrho^2 \sum_{\pi \in \Pi_{\text{cusp}}^{(T_\theta, \Lambda)}(\lambda, N)} \widehat{f}_S(\pi_S) \frac{|R(\varphi_\pi^0, E, \Lambda)|^2}{(\varphi_\pi^0 | \varphi_\pi^0)_{L^2}} \widehat{L}(s + \tfrac{1}{2}, \pi, \mu) \\ \times \prod_{p < \infty} \mathbb{I}_{\pi_p, \mathbf{K}_0(N\mathbb{Z}_p)}^{(s)}(\xi_p, \phi_p, \Lambda_p, \mu_p; b_p),$$

where ξ_p for $p < \infty$ is from the fixed (T_θ, Λ) -Bessel data $\{(\ell_p, \xi_p)\}_{p < \infty}$ of π , and

$$(5.10) \quad \mathbb{I}_{\pi_p, \mathbf{K}_0(N\mathbb{Z}_p)}^{(s)}(\xi_p, \phi_p, \Lambda_p, \mu_p; b_p) := \sum_{v \in \mathcal{B}(\pi_p, \mathbf{K}_0(N\mathbb{Z}_p))} Z_p^*(\phi_p, \overline{B_{v_p}}; s, \mu_p, b_p) \ell_p(v).$$

We also use the notation $\zeta_M(s)$ to denote $\prod_{p|M} (1 - p^{-s})^{-1}$.

Note that if π_p with $p \nmid N$ is unramified and $(\ell_{\pi_p}^0, \xi_{\pi_p}^0)$ is the unramified Bessel datum of π_p , then (5.10) is 1. For other cases, we compute (5.10) in §6.8 completely. Substituting them, we obtain

Theorem 5.2. *Let $(\lambda = (l_1, l_2), N)$, μ , $\tilde{\mu}$ and M be as in §4.3. Let S be a finite set of prime numbers relatively prime to DMN . Then, for any $f_S = \otimes_{p \in S} f_p \in \mathcal{H}_S$,*

(5.11)

$$\begin{aligned} \mathbb{I}^{(s)}(\lambda, N, f_S) &= \frac{2^{-2}|D|^{\frac{1}{2}(3-\frac{l_1+l_2}{2})}e^{-2\pi\sqrt{|D|}}}{w_D^2[\mathbf{K}_f : \mathbf{K}_0(N)]} M^{s-6} \zeta_M(1) \zeta_M(4) \tilde{\mu}(2D) G(\tilde{\mu}) N^{s-1} \tilde{\mu}^{-1}(N) \prod_{p|N} (1+p^{-2})^{-1} \\ &\quad \times (\mathbf{v}_\varrho^0 | \mathbf{v}_\varrho^0)_\varrho^2 \sum_{\pi \in \Pi_{\text{cusp}}^{(T_\theta, \Lambda)}(\lambda, N)} \widehat{f}_S(\pi_S) \frac{|R(\varphi_\pi^0, E, \Lambda)|^2}{(\varphi_\pi^0 | \varphi_\pi^0)_{L^2}} \widehat{L}(s + \frac{1}{2}, \pi, \mu) \mathfrak{t}(\pi, \mu) \end{aligned}$$

with $\mathfrak{t}(\pi, \mu) = \prod_{p|N} \mathfrak{t}(\pi_p, \mu_p)$ with

(5.12)

$$\mathfrak{t}(\pi_p, \mu_p) := \begin{cases} 1 & (p \mid N_\pi \text{ and } \pi_p \text{ is of type VIb}), \\ 2 & (p \mid N_\pi \text{ and } \pi_p \text{ is of type IIIa}), \\ 2(p-1)p^{-5}L(1, \pi_p, \text{Std}) \{1 + \mu_p^2(p) - \frac{\mu_p(p)}{p+1} \text{tr}(p^{-1}T_p + \eta_p | \pi_p^{\mathbf{K}_0(p\mathbb{Z}_p)})\} & (p \mid \frac{N}{N_\pi}). \end{cases}$$

To refine (5.11) by considering $\mathbb{I}^{(s)}(\lambda, N, f_S)^{\text{T}, \text{new}}$ and $\mathbb{I}^{(s)}(\lambda, N, f_S)^{\text{SK}, \text{new}}$, we refer to the explicit formula of the quantity $\frac{|R(\varphi_\pi^0, E, \Lambda)|^2}{(\varphi_\pi^0 | \varphi_\pi^0)_{L^2}}$ first proved by Dickson, Pitale, Saha, and Schmidt ([7, Theorems 1.13 and 3.13]) when $l_1 = l_2$ under the refined GGP conjecture for Bessel periods posed by Liu([23]). Owing to a theorem by Furusawa and Morimoto [11, Theorem 8.1], the formula in [7, Theorems 1.13 and 3.13] is extended to vector-valued forms unconditionally so that it can applied to (5.11) to yield the following theorem. Note that the polynomial $Q_{S, \varrho}$ (with $S = T_\theta$) in [10, (8.2.16)] equals $(-1)^{\frac{l_1-l_2}{2}} (\frac{2}{\sqrt{|D|}})^{\frac{l_1+l_2}{2}} \mathbf{v}_\varrho^\theta$, our

$R(\varphi, E, \Lambda)$ equals the quantity $w_D(\frac{\sqrt{|D|}}{2})^{-\frac{l_1+l_2}{2}} \mathcal{B}_\Lambda(\varphi; E) / (Q_{S, \varrho}, Q_{S, \varrho})_{l_1-l_2}$ defined by [10, (8.2.17)], and $(Q_{S, \varrho}, Q_{S, \varrho})_{l_1-l_2} = (\frac{|D|}{4})^{-\frac{l_1+l_2}{2}} (\mathbf{v}_\varrho^0, \mathbf{v}_\varrho^0)_{l_1-l_2} = (\frac{|D|}{4})^{-\frac{l_1+l_2}{2}} (\mathbf{v}_\varrho^0 | \mathbf{v}_\varrho^0)_\varrho$.

Theorem 5.3. *Let the notations and the assumptions be as in Theorem 5.2. We have*

(i) $\mathbb{I}^{(s)}(\lambda, N, f_S)^{\text{G}, \text{new}}$ equals

$$\begin{aligned} &\frac{2^{\#S(N)+2l_1-6}|D|^{\frac{l_1+l_2+2}{4}}e^{-2\pi\sqrt{|D|}}}{[\mathbf{K}_f : \mathbf{K}_0(N)]} \\ &\quad \times M^{s-6} \zeta_M(1) \zeta_M(4) \tilde{\mu}(2D) G(\tilde{\mu}) N^{s-1} \tilde{\mu}^{-1}(N) \prod_{p \in S(N)} (1+p^{-1}) \\ &\quad \times \sum_{\pi \in \Pi_{\text{cusp}}^{(T_\theta, \Lambda)}(\lambda, N)^{\text{G}, \text{new}}} \widehat{f}_S(\pi_S) \frac{\widehat{L}(\frac{1}{2}, \pi \times \mathcal{AI}(\Lambda^{-1}))}{\widehat{L}(1, \pi; \text{Ad})} \widehat{L}(s + \frac{1}{2}, \pi, \mu), \end{aligned}$$

where $S(N)$ is the set of all the prime divisors of N , and $\mathcal{AI}(\Lambda^{-1})$ is the automorphic representation of $\mathbf{GL}_2(\mathbb{A})$ defined by the automorphic induction of Λ^{-1} from \mathbb{A}_E^\times .

- (ii) If N has even number of prime divisors, then $\mathbb{I}^{(s)}(\lambda, N, f_S)^{Y, \text{new}} = 0$. If N has odd number of prime divisors, then $\mathbb{I}^{(s)}(\lambda, N, f_S)^{Y, \text{new}}$ equals

$$\begin{aligned} & \frac{2^{\#S(N)+2l_1-7} |D|^{\frac{l_1+l_2+2}{4}} e^{-2\pi\sqrt{|D|}}}{[\mathbf{K}_f : \mathbf{K}_0(N)]} \\ & \times M^{s-6} \zeta_M(1) \zeta_M(4) \tilde{\mu}(2D) G(\tilde{\mu}) N^{s-1} \tilde{\mu}^{-1}(N) \prod_{p \in S(N)} (1 + p^{-1}) \\ & \times \sum_{\substack{\pi_1 \in \Pi_{\mathbf{PGL}_2, \text{cusp}}(l_1+l_2-2, N)^{\text{new}} \\ \pi_2 \in \Pi_{\mathbf{PGL}_2, \text{cusp}}(l_1-l_2+2, N)^{\text{new}}}} \widehat{f}_S(Y(\pi_1, \pi_2)_S) \frac{\widehat{L}(\frac{1}{2}, \pi_1 \times \mathcal{A}I(\Lambda^{-1})) \widehat{L}(\frac{1}{2}, \pi_2 \times \mathcal{A}I(\Lambda^{-1}))}{\widehat{L}(1, \pi_1; \text{Ad}) \widehat{L}(1, \pi_2; \text{Ad}) \widehat{L}(1, \pi_1 \times \pi_2)} \\ & \times \widehat{L}(s + \frac{1}{2}, \pi_1 \times \mu) \widehat{L}(s + \frac{1}{2}, \pi_2 \times \mu), \end{aligned}$$

where $\Pi_{\mathbf{PGL}_2, \text{cusp}}(k, N)^{\text{new}}$ is the set of all irreducible cuspidal representations of $\mathbf{PGL}_2(\mathbb{A})$ associated to holomorphic newforms of weight k and level N and $Y(\pi_1, \pi_2) \in \Pi_{\text{cusp}}(\lambda, N)^{\text{new}}$ denotes the Yoshida lift of π_1 and π_2 .

- (iii) If $\Lambda \neq \mathbf{1}$ or $l_1 > l_2$, then $\mathbb{I}^{(s)}(\lambda, N, f_S)^{\text{SK}, \text{new}} = 0$. If $\Lambda = \mathbf{1}$ and $l_1 = l_2 (= l)$, then $\mathbb{I}^{(s)}((l, l), N, f_S)^{\text{SK}, \text{new}}$ equals

$$\begin{aligned} & \frac{3 \cdot 2^{2l-3} |D|^{(l+2)/2} e^{-2\pi\sqrt{|D|}}}{[\mathbf{K}_f : \mathbf{K}_0(N)]} \cdot \frac{s}{4\pi} \\ & \times M^{s-6} \zeta_M(1) \zeta_M(4) \tilde{\mu}(2D) G(\tilde{\mu}) N^{s-1} \tilde{\mu}^{-1}(N) \prod_{p \in S(N)} (1 + p^{-1}) \cdot \widehat{L}(s+1, \mu) \widehat{L}(s, \mu) \\ & \times \sum_{\pi_0 \in \Pi_{\mathbf{PGL}_2, \text{cusp}}^{(T_\theta, 1)}(2l-2, N)^{\text{new}}} \widehat{f}_S(\text{SK}(\pi_0)_S) \frac{\widehat{L}(\frac{1}{2}, \pi_0 \times \chi_D) \widehat{L}(1, \chi_D)^2}{\widehat{L}(\frac{3}{2}, \pi_0) \widehat{L}(1, \pi_0; \text{Ad})} \widehat{L}(s + \frac{1}{2}, \pi_0 \times \mu), \end{aligned}$$

where $\Pi_{\mathbf{PGL}_2, \text{cusp}}^{(T_\theta, 1)}(2l-2, N)^{\text{new}}$ is the set of all $\pi_0 \in \Pi_{\mathbf{PGL}_2, \text{cusp}}(2l-2, N)^{\text{new}}$ such that the Saito-Kurokawa lift $\text{SK}(\pi_0)$ of π_0 has the $(T_\theta, \mathbf{1})$ -Bessel model, and χ_D is the Kronecker character of modulo D .

Proof. The equation (i) and (ii) is a direct corollary to Theorem 5.3 and [10, Theorem 8.1] (for the scalar case we refer to [7, Theorems 1.13 and 3.14]; we also note the formula

$$\widehat{L}(s, \pi, \mu) = \widehat{L}(s, \pi_1 \times \mu) \widehat{L}(s, \pi_2 \times \mu),$$

where $\pi = Y(\pi_1, \pi_2)$. This obtained immediately by comparing the local factors. It is noted in [7, p.296] that each local representation π_p of $\pi \cong \otimes_p \pi_p \in \Pi_{\text{cusp}}^{(T_\theta, \Lambda)}(\lambda, N)^{Y, \text{new}}$ for $p \mid N$ is of type VIb. Finally, we check the statement (iii). As quoted in [7, Theorem 3.11], the (T_θ, Λ) -Bessel periods of the Saito-Kurokawa lifts are zero unless Λ is trivial due to Qiu. By comparing the local L -functions ([39, Theorem 5.2(ii)], cf. [29, Theorem 3.1]),

$$(5.13) \quad \widehat{L}(s, \pi, \mu) = \frac{1}{4\pi} (s - 1/2) \widehat{L}(s, \pi_0 \times \mu) \widehat{L}(s + 1/2, \mu) \widehat{L}(s - 1/2, \mu),$$

where $\pi = \text{SK}(\pi_0)$. By [39] and the definition, the set $\Pi_{\text{cusp}}^{(T_\theta, 1)}((l, l), N)^{\text{SK}}$ corresponds bijectively to the set $\Pi_{\mathbf{PGL}_2, \text{cusp}}^{(T_\theta, 1)}(2l - 2, N)$ via the Saito-Kurokawa lifting if $l \geq 3$. Note that the condition $l \geq 3$ is ensured by assumption (A-ii) in §4.3. \square

By (5.4), we have that $\mathbb{I}^{(s)}(\lambda, N, f_S) - \mathbb{I}^{(s)}(\lambda, N, f_S)^{\text{G, new}}$ is the sum of $\mathbb{I}^{(s)}(\lambda, N, f_S)^{\text{Y, new}}$, $\mathbb{I}^{(s)}(\lambda, N, f_S)^{\text{SK, new}}$ and $\mathbb{I}^{(s)}(\lambda, N, f_S)^{\text{old}}$. Our main focus is their values at $s = 0$. For those, we have the following upper bounds in the level aspect.

Theorem 5.4. *Let $l, \mu, \tilde{\mu}, M, S$ be as in Theorem 5.2.*

- (i) *Let λ be fixed as before and N an inert prime in E/\mathbb{Q} . Then, there exist constants $C_{D, \lambda, \mu, f_S}, C'_{D, \lambda, \mu, f_S} > 0$ depending only on D, l, μ , and f_S such that*

$$(5.14) \quad |\mathbb{I}^{(0)}(\lambda, N, f_S)^{\text{old}}| < C_{D, \lambda, \mu, f_S} N^{-8}.$$

$$(5.15) \quad |\mathbb{I}^{(0)}(\lambda, N, f_S)^{\text{Y, new}}| < C'_{D, \lambda, \mu, f_S} N^{-\frac{3}{2}}.$$

- (ii) *Let N be 1 or an inert prime in E/\mathbb{Q} . When $\mu \neq \mathbf{1}$, $\mathbb{I}^{(0)}(l, N, f_S)^{\text{SK, new}} = 0$. When $\mu = \mathbf{1}$, there exists a constant $C_{f_S} > 0$ depending only on f_S such that*

$$(5.16) \quad |\mathbb{I}^{(0)}((l, l), N, f_S)^{\text{SK, new}}| < C_{f_S} |D|^{\frac{l}{2}} l^{\frac{1}{2}} N^{-\frac{3}{2}}.$$

Proof. Because N is a prime, one can check that the summation range of $\mathbb{I}^{(0)}(\lambda, N, f_S)^{\text{old}}$ is $\Pi_{\text{cusp}}^{(T_\theta, \chi)}(l, N)^{\text{old}} = \Pi_{\text{cusp}}^{(T_\theta, \chi)}(\lambda, 1)$, which is independent of N . From the formula of $\mathfrak{t}(\pi_p, \mu_p)$ in (5.12) for $p = N$, combined with the temperedness of π_N due to [48] and the matrix of $T_{1,0}$ in table (6.29), we get $\mathfrak{t}(\pi_N, \mu_N) = O(N^{-4})$. This and the equation $[\mathbf{K}_f : \mathbf{K}_0(N)] = N^3(1 + N^{-2})(1 + N^{-1})$ for prime N yield the bound (5.14).

Next we treat the average for Yoshida lifts. For $\pi_1 \in \Pi_{\mathbf{PGL}_2, \text{cusp}}(l_1 + l_2 - 2, N)^{\text{new}}$ and $\pi_2 \in \Pi_{\mathbf{PGL}_2, \text{cusp}}(l_1 - l_2 + 2, N)^{\text{new}}$, we need the lower bound

$$L(1, \pi_1 \times \pi_2) \gg \exp(-C \sqrt{\log N}),$$

uniform in N . This is a special case of Lemma 5.5, because π_1 and π_2 are everywhere tempered by Deligne's estimate and by the fact that the local p -components of π_i for $p \mid N$ are the (twisted) Steinberg representation, which is tempered. Note that $\pi_1 \not\cong \pi_2$ due to the weight condition. Now, We deduce the inequality (5.15) bounding the sum from above by a product of the average considered in [8, Theorem 1.1]; to do this, we invoke the subconvexity bound

$$(5.17) \quad L(\tfrac{1}{2}, \pi) = O(C(\pi)^{\frac{1}{4}-\delta}), \quad (\exists \delta > 0)$$

for automorphic representations π of $\mathbf{GL}_2(\mathbb{A})$ ([24]) and the non-negativity of $L(\frac{1}{2}, \pi_i \times \mathcal{AZ}(\Lambda^{-1}) = L(\frac{1}{2}, \text{BC}_{E/\mathbb{Q}}(\pi_i) \otimes \Lambda^{-1})$ for $i = 1, 2$ ([16])).

Let us prove (5.16); by Theorem 5.3 (ii), we may assume $\lambda = (l, l)$. Suppose $\mu \neq \mathbf{1}$; then, by Theorem 5.3 (ii), we have that $\mathbb{I}^{(s)}((l, l), N, f_S)^{\text{SK, new}} \times s^{-1}$ is entire, hence $\mathbb{I}^{(0)}((l, l), N, f_S)^{\text{SK, new}} = 0$ because $\widehat{L}(s, \mu)$, as well as $\widehat{L}(s, \pi_0 \times \chi_D)$, is entire. In the rest of the proof, we assume $\mu = \mathbf{1}$. Then, $s\widehat{L}(s, \mu)\widehat{L}(s + 1, \mu)$ has a simple pole at $s = 0$. By [38, Theorem 3.1 and Table 2], $\text{SK}(\pi_0)$ with $\pi_0 \in \Pi_{\text{cusp}, \mathbf{PGL}_2}(2l - 2, N)$ has the global (T_θ, Λ) -Bessel model only if the sign of the functional equation of π_0 is -1 so that

$\widehat{L}(s+1/2, \pi_0)$ has a zero at $s = 0$. Hence, theorem 5.3 (ii) gives us the following majorant of $\mathbb{I}^{(0)}((l, l), N, f_S)^{\text{SK, new}}$.

$$2^3 \pi^{\frac{5}{2}} (l-1)^{-2} \frac{\Gamma(l)}{\Gamma(l-\frac{1}{2})} |D|^{\frac{l}{2}} [\mathbf{K}_f : \mathbf{K}_0(N)]^{-1} \sum_{\pi_0 \in \Pi_{\mathbf{PGL}_2, \text{cusp}}^{(T_\theta, 1)}(2l-2, N)} \frac{L(\frac{1}{2}, \pi_0 \times \chi_D) |L'(\frac{1}{2}, \pi_0)|}{L(1, \pi_0; \text{Ad})}.$$

To estimate this, we invoke a subconvexity bound $L(1/2, \pi_0 \times \chi_D) = O((l^2 N)^{\frac{1}{4}-\delta})$ for some $\delta > 0$ from (5.17) and a lower bound $L(1, \pi_0; \text{Ad}) > c_0 \exp(-c_1 \sqrt{\log(1+l^2 N)})$ for some constant $c_0, c_1 > 0$ which is known by [13, Theorem 0.1] (see also the remark after [22, Corollary 7]). The lower bound implies $L(1, \pi_0; \text{Ad})^{-1} = O((l^2 N)^{2\delta})$. Moreover, by a common argument, a subconvexity bound for the central derivative of $L'(1/2, \pi_0) = O((l^2 N)^{\frac{1}{4}-\delta})$ is derived from a subconvexity bound for $L(1/2, \pi_0 | \cdot |_{\mathbb{A}}^{it})$ available from (5.17). By the uniform bound $\#\Pi_{\mathbf{PGL}_2, \text{cusp}}(2l-2, N) = O(lN)$ and the asymptotic $\frac{\Gamma(l)}{\Gamma(l-\frac{1}{2})} \sim l^{\frac{1}{2}}$ ($l \rightarrow \infty$), the bound (5.16) follows. \square

Lemma 5.5. *Let π_i ($i = 1, 2$) be irreducible cuspidal automorphic representations of \mathbf{PGL}_n such that $C(\pi_1), C(\pi_2) \leq Q$ with $Q > 2$. We assume that $\pi_1 \not\cong \pi_2$ and both of them are self-dual and tempered everywhere. Then,*

$$L(1, \pi_1 \times \pi_2) \gg \exp(-C \sqrt{\log Q}).$$

with an absolute constant $C > 0$.

Proof. We recall the argument indicated in [36] (attributed originally to [25]), which eliminates a possibility of Siegel zeros of the L -function $L(s, \pi_1 \times \pi_2)$. Fix $t \in \mathbb{R}$. Then, $L(s, \Pi \times \widetilde{\Pi})$ with Π being the isobaric sum

$$\Pi := \pi_1 \boxplus (\pi_1 \times | \cdot |_{\mathbb{A}}^{it}) \boxplus (\pi_1 \times | \cdot |_{\mathbb{A}}^{-it}) \boxplus \pi_2 \boxplus (\pi_2 \times | \cdot |_{\mathbb{A}}^{it}) \boxplus (\pi_2 \times | \cdot |_{\mathbb{A}}^{-it})$$

is a Dirichlet series with non-negative coefficients ([14, Lemma a]). Moreover, by ([31, Proposition 9.4]) and due to π_1 and π_2 being self-dual, $L(s, \Pi \times \widetilde{\Pi})$ is expressed as

$$L(s, \Pi \times \widetilde{\Pi}) = \prod_{1 \leq i, j \leq 2} \left[\frac{L(s, \pi_i \times \pi_j)^3 L(s-it, \pi_i \times \pi_j)^2 L(s+it, \pi_i \times \pi_j)^2}{\times L(s-2it, \pi_i \times \pi_j) L(s+2it, \pi_i \times \pi_j)} \right]^2,$$

which shows that $L(s, \Pi \times \widetilde{\Pi})$ has a pole of order 6 at $s = 1$ (due to $\pi_1 \not\cong \pi_2$), and has a zero of order 8 at $s = \sigma$ if $L(\sigma+it, \pi_i \times \pi_j) = 0$ ($i, j = 1, 2$). Hence, by [12, Lemma, p.178], one can show that $L(s, \pi_1 \times \pi_2)$ has no zeros in the interval $(1 - \frac{C_0}{\log M}, 1)$ for some constant $C_0 > 0$ with $M = (1 + |t|)^{24} C(\pi_2 \times \pi_2)^{18} C(\pi_1 \times \pi_1)^9 C(\pi_2 \times \pi_2)^2$, where $C(\pi_i \times \pi_j)$ is the analytic conductor of $L(s, \pi_i \times \pi_j)$. By [14, lemma b], we have $C(\pi_i \times \pi_j) \ll Q^{2n}$. Thus, for an absolute constant $C'_0 > 0$, $L(s, \pi_1 \times \pi_2)$ has no zeros in the interval $(1 - \frac{C'_0}{\log Q}, 1)$. Now, we apply the argument of [22, Corollary 7]) to the L -function $L(s, \pi_1 \times \pi_2)$. Since π_1, π_2 are assumed to be tempered everywhere, the optimal bound of the lambda function $\Lambda_{\mathcal{A}}(n)$ for $\pi_1 \times \pi_2$ is available so that the automorphy of $\pi_1 \times \pi_2$ is not necessary, which simplifies the proof to get the lower bound of $L(1, \pi_1 \times \pi_2)$. \square

As a consequence of Theorem 5.4, we get

$$(5.18) \quad \mathbb{I}^{(0)}(\lambda, N, f_S)^{\text{G, new}} = \mathbb{I}^{(0)}(\lambda, N, f_S) + O_{\Lambda, \mu, \lambda, f_S}(N^{-\frac{3}{2}}).$$

6. COMPUTATION OF LOCAL ZETA-INTEGRALS FOR p -ADIC FIELDS

In this section, let F be a non-archimedean local field of characteristic 0, \mathfrak{o} the integer ring of F , \mathfrak{p} the maximal ideal of \mathfrak{o} , ϖ a generator of \mathfrak{p} and $q = \#(\mathfrak{o}/\mathfrak{p})$. Let $|\cdot|$ denote the normalized absolute value of F , i.e., $|\varpi| = q^{-1}$. Fix a non-trivial additive character $\psi : F \rightarrow \mathbb{C}^1$ with $\text{cond}(\psi) := \min\{n \in \mathbb{Z} ; \psi|_{\varpi^n \mathfrak{o}} = 1\} = 0$.

We compute the local zeta-integral a la Piatetski-Shapiro ([30]) for several representations, taking particular test functions; as a result, we determine the local L -factors and the local ε -factors in [30] to confirm that they coincides with the expected ones listed in [35, Tables A8,A9]. As explained in §4.3, for a particular global application in mind, we only deal with representations of types I, IIb, IIIa and VIb (but allowing the central characters to be non-trivial.)

6.1. Local zeta integral for Bessel models. We first review some generalities on local zeta integrals and then recall results from [32] on explicit formulas of Bessel functions for Iwahori spherical representations of \mathbf{G} , which are possible local components of $\pi \cong \otimes_p \pi \in \Pi_{\text{cusp}}^{(T_\theta, \Lambda)}(\varrho, N)$.

Let $K = \mathbf{G}(\mathfrak{o})$ be a standard maximal compact subgroup of $\mathbf{G}(F)$ and $K_0(\mathfrak{p}^e) := \{ \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in K \mid C \equiv 0 \pmod{\mathfrak{p}^e} \}$ be the congruence subgroup of level \mathfrak{p}^e .

For a symmetric matrix $S = \begin{bmatrix} a & b/2 \\ b/2 & c \end{bmatrix} \in \text{Sym}_2(F)$ such that $d := b^2 - 4ac \neq 0$, set $\xi = \begin{bmatrix} b/2 & c \\ -a & -b/2 \end{bmatrix}$. Let $L := F \oplus F\xi$ be the two-dimensional F -algebra, \mathfrak{o}_L be its integer ring, and ψ_L be the additive character on L given by $\psi_L := \psi \circ \text{tr}_{L/F}$. The maps

$$\begin{aligned} L \ni x + y\xi &\longmapsto x + \frac{\sqrt{d}}{2}y \in F(\sqrt{d}) \quad (d \notin (F^\times)^2), \\ L \ni x + y\xi &\longmapsto \left(x + \frac{\sqrt{d}}{2}y, x - \frac{\sqrt{d}}{2}y \right) \in F \oplus F \quad (d \in (F^\times)^2) \end{aligned}$$

are isomorphisms of F -algebras. We assume that

- (i) $a \in \mathfrak{o}^\times$ and $b, c \in \mathfrak{o}$.
- (ii) If $d \notin (F^\times)^2$, then $\sqrt{d}\mathfrak{o}_L$ is the different ideal of L/F .
- (iii) If $d \in (F^\times)^2$, then $d \in \mathfrak{o}^\times$.

We can check that

$$L^\times = \{ \tau \in \mathbf{GL}_2(F) \mid {}^t \tau S \tau = \det \tau \cdot S \}.$$

The map $L \ni \tau \mapsto \tau^\dagger \in L$ is the non-trivial F -automorphism on L . We set $\theta_0 = b/2 - \xi \in \mathfrak{o}_L$, then $\{a, \theta_0\}$ is an \mathfrak{o} -basis of \mathfrak{o}_L . Let $\langle \cdot, \cdot \rangle$ denote the symplectic form on L^2 over F defined by

$$\langle x, y \rangle := \text{tr}_{L/F}(-(2\xi)^{-1}(x_1 y_2 - x_2 y_1)), \quad x = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}, y = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} \in L^2.$$

We set

$$\begin{aligned} G^\#(F) &:= \{ g \in \mathbf{GL}_2(L) \mid \det g \in F^\times \}, \\ B^\#(F) &:= \left\{ \begin{bmatrix} \tau & \beta \\ 0 & a\tau^\dagger \end{bmatrix} \mid \tau \in L^\times, \beta \in L, a \in F^\times \right\}, \\ K^\# &:= G^\#(F) \cap \mathbf{GL}_2(\mathfrak{o}_L). \\ K_0^\#(\mathfrak{p}^e) &:= G^\#(F) \cap \begin{bmatrix} \mathfrak{o}_L & \mathfrak{o}_L \\ \mathfrak{p}^e \mathfrak{o}_L & \mathfrak{o}_L \end{bmatrix}. \end{aligned}$$

Since $\langle g^\# x, g^\# y \rangle = \det g^\# \langle x, y \rangle$ for any $x, y \in L^2$ and $g^\# \in \mathbf{G}(F)$, we get a natural embedding $G^\#(F) \ni g \mapsto \iota(g^\#) \in \mathbf{G}(F)$. More precisely, $\iota(g)$ is the representation matrix of with respect to an F -basis $\{[\begin{smallmatrix} \tau & 0 \\ 0 & a\tau^\dagger \end{smallmatrix}], [\begin{smallmatrix} \theta_0 & \\ & 0 \end{smallmatrix}], [-\mathbf{a}^{-1}\theta_0^\dagger, [\begin{smallmatrix} 0 & \\ & 1 \end{smallmatrix}]]\}$. A computation yields

$$(6.1) \quad \begin{aligned} \iota\left([\begin{smallmatrix} \tau & 0 \\ 0 & a\tau^\dagger \end{smallmatrix}]\right) &= [\begin{smallmatrix} \tau & 0 \\ 0 & a\tau^\dagger \end{smallmatrix}], \quad \tau \in L^\times, a \in F^\times, \\ \iota\left([\begin{smallmatrix} 1 & \beta \\ 0 & 1 \end{smallmatrix}]\right) &= [\begin{smallmatrix} 1_2 & X_\beta \\ 0 & 1_2 \end{smallmatrix}], \quad \beta = \beta_2 \mathbf{a} + \beta_3 \theta_0 \in L, \quad X_\beta := \begin{bmatrix} -\mathbf{a}^{-1} \mathbf{b} \beta_2 - \mathbf{a}^{-1} \mathbf{c} \beta_3 & \beta_2 \\ \beta_2 & \beta_3 \end{bmatrix}, \\ \iota\left([\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}]\right) &= \begin{bmatrix} 0 & 0 & -\mathbf{a}^{-2} \mathbf{b} & \mathbf{a}^{-1} \\ 0 & 0 & \mathbf{a}^{-1} & 0 \\ 0 & \mathbf{a} & 0 & 0 \\ \mathbf{a} & \mathbf{b} & 0 & 0 \end{bmatrix}. \end{aligned}$$

We note that $K^\# = \iota^{-1}(K)$ and $K_0^\#(\mathfrak{p}^e) = \iota^{-1}(K_0(\mathfrak{p}^e))$. We call

$$R := \left\{ \iota\left([\begin{smallmatrix} \tau & 0 \\ 0 & \tau^\dagger \end{smallmatrix}]\right) n \mid \tau \in L^\times, n \in \mathbf{N}(F) \right\}$$

the Bessel subgroup of $\mathbf{G}(F)$ (with respect to S). For a character $\Lambda : L^\times \rightarrow \mathbb{C}^\times$, we can check that the map

$$R \ni \iota\left([\begin{smallmatrix} \tau & 0 \\ 0 & \tau^\dagger \end{smallmatrix}]\right) n(X) \mapsto \Lambda(\tau) \psi(\mathrm{tr}(SX)) \in \mathbb{C}^\times$$

defines a character on R . We denote this character by $\Lambda \otimes \psi_S$.

Let (π, V_π) be an irreducible admissible representation of $\mathbf{G}(F)$. Let $(V_\pi^*)^{S, \Lambda}$ denote the space of all \mathbb{C} -linear forms $\ell : V_\pi \rightarrow \mathbb{C}$ that satisfies

$$\ell(\pi(r)\xi) = \Lambda \otimes \psi_S(r) \ell(\xi), \quad \xi \in V_\pi, r \in R.$$

Then, it holds that $\dim_{\mathbb{C}}(V_\pi^*)^{S, \Lambda} \leq 1$ from ([27], [28]). We say that π has a local (S, Λ) -Bessel model if $(V_\pi^*)^{S, \Lambda} \neq (0)$. In this case, we can define the local (S, Λ) -Bessel model of π as in §4.4. Moreover, when π is spherical, we define the unramified (S, Λ) -Bessel datum $(\ell_\pi^0, \xi_\pi^0) \in (V_\pi^*)^{S, \Lambda} \times V_\pi^K$ for π as in §4.4, so that

$$B_\pi^0(g) = \ell_\pi^0(\pi(g)\xi_\pi^0), \quad g \in \mathbf{G}(\mathbb{Q}_p).$$

is the unramified Bessel function such that $B_\pi^0(1_4) = 1$. Recall that all irreducible admissible representations of $\mathbf{G}(F)$ that admit local Bessel models are classified in [35], and the result is conveniently summarized in [32, Table 2]. For our cases, as we mentioned in §4.4, we may assume that π is of type I, IIb, IIIa, or VIb. We refer some of its properties in ([32, Table 2]);

$$(6.2) \quad \begin{array}{|c|c|c|c|c|} \hline \text{type} & \pi & \dim V_\pi^K & \dim V_\pi^{K_0(\mathfrak{p})} & \text{Cent. Char.} \\ \hline \text{I} & \chi \times \chi' \rtimes \sigma & 1 & 4 & \chi \chi' \sigma^2 \\ \hline \text{IIb} & \chi 1_{\mathbf{GL}_2} \rtimes \sigma & 1 & 3 & \chi^2 \sigma \\ \hline \text{IIIa} & \chi \rtimes \sigma_{\mathrm{St}_{\mathbf{GL}_2}} & 0 & 2 & \chi \sigma^2 \\ \hline \text{VIb} & \tau(T, |\cdot|^{-\frac{1}{2}} \sigma) & 0 & 1 & \sigma^2 \\ \hline \end{array}$$

Here, χ , χ' , and σ are unramified quasicharacters on F^\times such that representations are irreducible and admit (S, Λ) -Bessel models. For $a \in F^\times$. For later use we set $\alpha = \chi(\varpi)$, $\beta = \chi'(\varpi)$, and $\gamma = \sigma(\varpi)$. According to [35, Table A2], for type IIIa and type VIb, the corresponding representations are unitarizable if and only if the inducing quasicharacters are unitary, whereas, for type I, the unitarity of the inducing quasi-characters is equivalent to the corresponding representations being unitarizable and tempered. In

what follows, having a global application in mind, we suppose the unitarity of all the inducing characters, i.e., $|\alpha| = |\beta| = |\gamma| = 1$.

Similarly as (3.1), for a Schwartz-Bruhat function $\phi \in \mathcal{S}(L^2)$, a character μ on F^\times , and $s \in \mathbb{C}$ with $\operatorname{Re}(s) > 1$, we define a function on $\mathbf{G}^\#(G)$ by

$$f_\phi^{(s, \Lambda, \mu)}(g^\#) = \mu(\det g^\#) |\det g^\#|^{s+1} \int_{L^\times} \phi\left(\begin{bmatrix} 1 & \\ & 0 \end{bmatrix} \begin{bmatrix} \tau & 0 \\ 0 & \tau^\dagger \end{bmatrix} g^\#\right) \Lambda \mu_L(\tau) |\tau \tau^\dagger|^{s+1} d^\times \tau,$$

where $\mu_L(\tau) := \mu(\tau \tau^\dagger)$, $\tau \in L^\times$ and $d^\times \tau$ is the normalized Haar measure on L^\times such that $\operatorname{vol}(\mathfrak{o}_L) = 1$. For an irreducible smooth admissible representation π of $\mathbf{Z}(F) \backslash \mathbf{G}(F)$, admitting (S, Λ) -Bessel model and $B \in \mathcal{B}(S, \Lambda)[\pi]$, define the zeta integral by

(6.3)

$$Z(\phi, B, s, \mu; g) = \int_{F^\times} \int_{K^\#} B(\mathbf{m}(a1_2, a) \iota(k^\#) g) \mu(a) |a|^{s-1} f_\phi^{(s, \Lambda, \mu)}(k^\#) dk^\# d^\times a, \quad s \in \mathbb{C}, \operatorname{Re}(s) > 1.$$

It is shown that $Z(\phi, B, \mu; g)$ is a rational function in q^{-s} ([41, Lemma 5.3.1]). Then, the local L -function $L(s, \pi, \mu)$ is defined to be the unique function such that $L(s, \pi, \mu)^{-1}$ is a polynomial in q^{-s} with constant term 1 and such that $Z(\phi, B, s, \mu, 1)/L(s, \pi, \mu) \in \mathbb{C}[q^s, q^{-s}]$ for all ϕ and B ([30], see also [41, Proposition 5.3.3]). By [30, Proposition 3.2], there exists an entire function $\varepsilon(\pi, s, \mu, \psi)$ such that the local functional equation

$$(6.4) \quad \varepsilon(\pi, s + 1/2, \mu, \psi) \frac{Z(\phi, B, s, \mu; g)}{L(s + 1/2, \pi, \mu)} = \frac{Z(\hat{\phi}, B^\dagger, -s, \mu^{-1}; g)}{L(-s + 1/2, \hat{\pi}, \mu^{-1})}$$

holds. Here, we set $B^\dagger \in \mathcal{B}(S, \Lambda^\dagger)[\pi]$ as $B^\dagger(g) := B\left(\begin{bmatrix} 1 & \mathbf{a}^{-1}\mathbf{b} & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & -\mathbf{a}^{-1}\mathbf{b} & 1 \end{bmatrix} g\right)$, $g \in \mathbf{G}(F)$, and

Λ^\dagger denote the Galois conjugate of Λ defined by $\Lambda^\dagger(\tau) := \Lambda(\tau^\dagger)$ for $\tau \in L^\times$, and $\hat{\phi}$ is the Fourier transform of $\phi \in \mathcal{S}(L^2)$ defined as

$$\hat{\phi}(x, y) = \int_{L^2} \phi(u, v) \psi_L\left(\left\langle \begin{bmatrix} x \\ y \end{bmatrix}, \begin{bmatrix} u \\ v \end{bmatrix} \right\rangle\right) du dv$$

with du and dv being the Haar measures on L such that $\operatorname{vol}(\mathfrak{o}_L) = 1$. The L -functions $L(s, \pi, \mu)$ are explicitly determined for several cases as shown in [41, Table 5], which, for $\mu = 1$, coincide with the local L -functions listed in [35, Table A.8]:

(6.5)

type	π	$L(s, \pi)$
I	$\chi \times \chi' \rtimes \sigma$	$(1 - \alpha\beta\gamma q^{-s})^{-1} (1 - \alpha\gamma q^{-s})^{-1} (1 - \beta\gamma q^{-s})^{-1} (1 - \gamma q^{-s})^{-1}$
IIb	$\chi 1_{\mathbf{GL}_2} \rtimes \sigma$	$(1 - \alpha^2\gamma q^{-s})^{-1} (1 - \gamma q^{-s})^{-1} (1 - \alpha\gamma q^{-s-\frac{1}{2}})^{-1} (1 - \alpha\gamma q^{-s+\frac{1}{2}})^{-1}$
IIIa	$\chi \rtimes \sigma_{\operatorname{St}_{\mathbf{GL}_2}}$	$(1 - \alpha\gamma q^{-s-\frac{1}{2}})^{-1} (1 - \gamma q^{-s-\frac{1}{2}})^{-1}$
VIb	$\tau(T, \cdot ^{-\frac{1}{2}}\sigma)$	$(1 - \gamma q^{-s-\frac{1}{2}})^{-2}$

We shall get explicit evaluations of the zeta integrals $Z(\phi, B, s, \mu; g)$ for particular choices of ϕ , B and g as shown in the next table, which allow us to determine $\varepsilon(s, \pi, \mu, \psi)$:

	type	ϕ	B	μ	L/F	g
	case 1	I or IIb	$\mathbb{1}_{\mathfrak{o}_L \oplus \mathfrak{o}_L}$	K -fixed	unramified	1_4
	case 2	I or IIb	$\mathbb{1}_{\mathfrak{p}^e \mathfrak{o}_L \oplus 1 + \mathfrak{p}^e \mathfrak{o}_L}$	K -fixed	ramified	b
(6.6)	case 3	I or IIb	$\mathbb{1}_{\mathfrak{p}^e \mathfrak{o}_L \oplus 1 + \mathfrak{p}^e \mathfrak{o}_L}$	K -fixed	ramified	b
	case 4	I or IIb	$\mathbb{1}_{\mathfrak{o}_L \oplus \mathfrak{o}_L}$	$K_0(\mathfrak{p})$ -fixed	unramified	inert
	case 5	IIIa	$\mathbb{1}_{\mathfrak{o}_L \oplus \mathfrak{o}_L}$	$K_0(\mathfrak{p})$ -fixed	unramified	inert
	case 6	VIb	$\mathbb{1}_{\mathfrak{o}_L \oplus \mathfrak{o}_L}$	$K_0(\mathfrak{p})$ -fixed	unramified	inert

Here,

$$(6.7) \quad b := \begin{bmatrix} \varpi^{e_{1_2}} & s^\dagger \\ 0 & 1_2 \end{bmatrix} \in \mathbf{G}(F),$$

and L/F is said to be inert if it is an unramified field extension. For $l, m \in \mathbb{Z}$, set

$$h(\ell, m) := \begin{bmatrix} \varpi^{\ell+2m} & & \\ & \varpi^{\ell+m} & \\ & & 1 \\ & & & \varpi^m \end{bmatrix} \in \mathbf{G}(F).$$

As in [32], for smooth representation (π, V_π) of $\mathbf{G}(F)$, we define an operator $T_{m,n} \in \text{End}(V_\pi)$ by

$$T_{m,n}v := \text{vol}(K_0(\mathfrak{p})) \int_{K_0(\mathfrak{p})h(\ell, m)K_0(\mathfrak{p})} \pi(g)v \, dg, \quad v \in V_\pi$$

with dg the Haar measure on $\mathbf{G}(F)$ such that $\text{vol}(K) = 1$, and the Atkin-Lehner involution by

$$\eta v := \pi(\eta)v, \quad \eta := \begin{bmatrix} & & & -1 \\ & & 1 & \\ & \varpi & & \\ -\varpi & & & \end{bmatrix}, \quad v \in V_\pi.$$

If π is irreducible and has the (S, Λ) -Bessel models, then for any $B \in \mathcal{B}(S, \Lambda)[\pi]^{K_0(\mathfrak{p})}$, by [32, Lemma 4.4 (i)], we have the support conditions:

$$(6.8) \quad B(h(\ell, m)) = B(h(\ell, m)s_2) = 0 \quad \text{if } \ell < 0,$$

$$(6.9) \quad B(h(\ell, m)s_1s_2) = B(h(\ell, m)s_2s_1s_2) = 0 \quad \text{if } \ell < -1.$$

6.2. Unramified computation (case1). When π is spherical, π is of type I or IIb in our cases. Recall that there exists a unique element $B_\pi^0 \in \mathcal{B}(S, \Lambda)[\pi]^K$ such that $B_\pi^0(1_4) = 1$. Then, the following formula for an unramified μ in its greatest generality is due to Sugano ([43, Theorem 2.1]):

$$Z(\phi, B_\pi^0, s, \mu; 1_4) = L(s + 1/2, \pi, \mu), \quad \text{Re}(s) \gg 0.$$

In [20, Proposition 5.9], a proof of this formula is given when $L = F \oplus L$ and π is of type I based on the explicit formula of B_π^0 due to [6].

6.3. Unramified computation (case2 & case3). Now we assume that $e := \text{cond}(\mu) > 0$ and that $S \in \text{Sym}^2(F)$ satisfies the following conditions;

- L/F is an unramified field extension.
- $S \in \mathbf{GL}_2(\mathfrak{o})$, $-\frac{d}{2} = \text{tr}(SS^\dagger) \in \mathfrak{o}^\times$.

Then ϖ remains a prime element in \mathfrak{po}_L . Let b as in (6.7), and set $\phi = \mathbb{1}_{\mathfrak{p}^e \mathfrak{o}_L \oplus 1 + \mathfrak{p}^e \mathfrak{o}_L} \in \mathcal{S}(L^2)$. Our goal of this subsection is to calculate the zeta integrals $Z(\phi, B_\pi^0, s, \mu; b)$ and $Z(\widehat{\phi}, B_\pi^0, s, \mu; b)$ explicitly. We remark that $Z(\phi, B_\pi^0, s, \mu; 1_4)$ must be zero because the a -integral in (6.3) vanishes.

Definition 2 (Root number). *Let ψ and μ be as above, we define the Gauss sum $W_F(\mu, \psi)$ by*

$$W_F(\mu, \psi) := q^{-\frac{e}{2}} \mu(\varpi)^{-e} \sum_{\alpha \in (\mathfrak{o}/\mathfrak{p}^e)^\times} \psi(\varpi^{-e} \alpha) \mu(\alpha).$$

Note that this sum has the absolute value 1 and is independent of the choice of ϖ . The following lemma is more or less well-known ([45, (2.18)]).

Lemma 6.1. *Let ψ and μ be as above. For $n \in \mathbb{Z}$,*

$$\int_{\mathfrak{o}^\times} \psi(\varpi^n a) \mu(a) d^\times a = \begin{cases} q^{-\frac{e}{2}+1} (q-1)^{-1} \mu(\varpi)^e W_F(\mu, \psi) & (n = -e), \\ 0 & (n \neq -e). \end{cases}$$

Lemma 6.2. *Let L/F be an unramified quadratic extension. Then, for any character μ of F with $e := \text{cond}(\mu) > 0$,*

$$W_L(\mu_L, \psi_L) = (-1)^e W_F(\mu, \psi)^2.$$

Proof. For a virtual representation V of the Weil group \mathfrak{W}_F of F , let $\varepsilon(s, V, \psi)$ denote the local epsilon factor a la Deligne-Langlands ([46]). By ([46, (3.4.8)]), we have

$$(6.10) \quad \varepsilon(s, \text{Ind}_{\mathfrak{W}_L}^{\mathfrak{W}_F} \rho, \psi) = \varepsilon(s, \rho, \psi_L)$$

for any ρ of degree 0. Any character μ of F^\times is viewed as a one-dimensional representation of \mathfrak{W}_F by $F^\times \cong \mathfrak{W}_F^{\text{ab}}$. If $\mu_L := \mu \circ N_{L/F}$, then $\text{Ind}_{\mathfrak{W}_L}^{\mathfrak{W}_F} \mu_L \cong \mu \oplus \mu \eta_{L/F}$. If μ is unramified, then $\varepsilon(s, \mu, \psi) = 1$; since L/F is unramified, so is the character $\eta_{L/F}$. By these observations, the equality (6.10) holds true when $\rho = \mathbf{1}$. Thus, (6.10) is true when $\rho = \mu$ with $e > 0$. Since $\varepsilon(s, \mu \eta_{L/F}^j, \psi) = q^{-e(s-1/2)} W_F(\bar{\mu} \eta_{L/F}^j, \psi)$ for $j = 0, 1$ and $\varepsilon(s, \mu_L, \psi_L) = q^{-2e(s-1/2)} W_L(\bar{\mu}_L, \psi_L)$, we obtain $W_L(\mu_L, \psi_L) = W_F(\mu, \psi) W_F(\mu \eta_{L/F}, \psi)$. By $e > 0$ and $\eta_{L/F}(\varpi) = -1$, we have $W_F(\mu \eta_{L/F}, \psi) = \eta_{L/F}(\varpi)^e W_F(\mu, \psi) = (-1)^e W_F(\mu, \psi)$. \square

Lemma 6.3. *For $k^\# = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in K^\#$, we have*

$$f_\phi^{(s, \Lambda, \mu)}(k^\#) = \begin{cases} q^{-2e+2} (q^2 - 1)^{-1} \mu_L(d)^{-1} \mu(\det k^\#) & (c \in \mathfrak{p}^e \mathfrak{o}_L, d \in 1 + \mathfrak{p}^e \mathfrak{o}_L), \\ 0 & (\text{otherwise}), \end{cases}$$

$$f_{\widehat{\phi}}^{(s, \Lambda, \mu)}(k^\#) = \begin{cases} q^{e(2s-3)+2} (q^2 - 1)^{-1} \Lambda(\varpi)^{-e} \mu_L(c)^{-1} W_L(\mu_L, \psi_L) & (c \in \mathfrak{o}_L^\times), \\ 0 & (c \notin \mathfrak{o}_L^\times). \end{cases}$$

Proof. First, we prove the first equation. By a direct calculation, we have

$$\begin{aligned} f_{\phi}^{(s, \Lambda, \mu)}(k^{\#}) &= \mu(\det k^{\#}) \int_{L^{\times}} \phi([c\tau^{\dagger} \ d\tau^{\dagger}]) \Lambda\mu_L(\tau) |\tau\tau^{\dagger}|^{s+1} d^{\times}\tau \\ &= \mu(\det k^{\#}) \int_{L^{\times}} \mathbb{1}_{\mathfrak{p}^e \mathfrak{o}_L}(c\tau^{\dagger}) \mathbb{1}_{1+\mathfrak{p}^e \mathfrak{o}_L}(d\tau^{\dagger}) \Lambda\mu_L(\tau) |\tau\tau^{\dagger}|^{s+1} d^{\times}\tau. \end{aligned}$$

Since $c \in \mathfrak{o}_L^{\times}$ or $d \in \mathfrak{o}_L^{\times}$, it is easy to check that $(c\tau^{\dagger}, d\tau^{\dagger}) \in \mathfrak{p}^e \mathfrak{o}_L \times (1 + \mathfrak{p}^e \mathfrak{o}_L)$ if and only if

$$c \in \mathfrak{p}^e \mathfrak{o}_L, \quad d \in \mathfrak{o}_L^{\times}, \quad \tau \in (d^{\dagger})^{-1}(1 + \mathfrak{p}^e \mathfrak{o}_L).$$

Hence, if $c \in \mathfrak{p}^e \mathfrak{o}_L$, $d \in \mathfrak{o}_L^{\times}$, we get

$$\begin{aligned} f_{\phi}^{(s, \Lambda, \mu)}(k^{\#}) &= \mu(\det k^{\#}) \cdot \text{vol}(1 + \mathfrak{p}^e \mathfrak{o}_L) \cdot \Lambda\mu_L((d^{\dagger})^{-1}) \\ &= q^{-2e+2}(q^2 - 1)^{-1} \mu_L(d)^{-1} \mu(\det k^{\#}). \end{aligned}$$

Next, we prove the second equation. It is easy to check that

$$\widehat{\phi}(x, y) = q^{-4e} \psi_L(x) \mathbb{1}_{\mathfrak{p}^e \mathfrak{o}}(x) \mathbb{1}_{\mathfrak{p}^e \mathfrak{o}}(y), \quad x, y \in L.$$

Hence we have

$$\begin{aligned} f_{\widehat{\phi}}^{(s, \Lambda, \mu)}(k^{\#}) &= \mu(\det k^{\#}) \int_{L^{\times}} \widehat{\phi}([c\tau^{\dagger} \ d\tau^{\dagger}]) \Lambda\mu_L(\tau) |\tau\tau^{\dagger}|_p^{s+1} d^{\times}\tau \\ &= q^{-4e} \mu(\det k^{\#}) \int_{L^{\times}} \psi_L(c\tau^{\dagger}) \mathbb{1}_{\mathfrak{p}^{-e} \mathfrak{o}_L}(c\tau^{\dagger}) \mathbb{1}_{\mathfrak{p}^{-e} \mathfrak{o}_L}(d\tau^{\dagger}) \Lambda\mu_L(\tau) |\tau\tau^{\dagger}|_p^{s+1} d^{\times}\tau \\ (6.11) \quad &= q^{-4e} \mu(\det k^{\#}) \sum_{n \geq -e} \Lambda\mu_L(\varpi)^n q^{-2n(s+1)} \int_{\mathfrak{o}_L^{\times}} \psi_L(c\varpi^n \tau) \mu_L(\tau) d^{\times}\tau. \end{aligned}$$

We put $c = c_0 \varpi^m$, $c_0 \in \mathfrak{o}_L^{\times}$. Then by Lemma 6.1, with (F, μ, ψ) replaced by $(L, \Lambda\mu_L, \psi_L)$, we have

$$\begin{aligned} \int_{\mathfrak{o}_L^{\times}} \psi_L(c\varpi^n \tau) \mu_L(\tau) d^{\times}\tau &= \mu_L(c_0)^{-1} \int_{\mathfrak{o}_L^{\times}} \psi_L(\varpi^{n+m} \tau) \mu_L(\tau) d^{\times}\tau \\ &= \delta_{n, -e-m} q^{-e+2} (q^2 - 1)^{-1} \mu_L(c_0^{-1} \varpi)^e W_L(\mu_L, \psi_L). \end{aligned}$$

By substituting this into (6.11), we obtain

$$\begin{aligned} f_{\widehat{\phi}}^{(s, \Lambda, \mu)}(k^{\#}) &= q^{-5e+2} (q^2 - 1)^{-1} \Lambda(\varpi)^{-e} \mu_L(c_0)^{-1} W_L(\mu_L, \psi_L) \sum_{n \geq -e} \delta_{n, -e-m} \Lambda\mu_L(\varpi)^{n+e} q^{-2n(s+1)} \\ &= \begin{cases} q^{e(2s-3)+2} (q^2 - 1)^{-1} \Lambda(\varpi)^{-e} \mu_L(c)^{-1} W_L(\mu_L, \psi_L) & (m = 0), \\ 0 & (m > 0). \end{cases} \quad \square \end{aligned}$$

Lemma 6.4. For $u \in \mathfrak{o}^{\times}$,

$$\sum_{\substack{\eta \in \mathfrak{o}_L / \mathfrak{p}^e \mathfrak{o}_L \\ u + N_{L/F}(\eta) \in \mathfrak{o}^{\times}}} \mu(u + N_{L/F}(\eta)) = (-1)^e q^e \mu(u).$$

Proof. Because L/F is an unramified extension, the norm map $\mathfrak{o}_L^{\times} \ni \eta \mapsto N_{L/F}(\eta) \in \mathfrak{o}^{\times}$ is surjective. Thus there exists $v \in \mathfrak{o}_L^{\times}$ such that $u = N_{L/F}(v)$. By replacing η by $v\eta$, without loss of generality, we can assume that $u = 1$.

For an integer i with $0 \leq i \leq e+1$, we set

$$\overline{\mathfrak{o}_{L,i}} = \begin{cases} \{\eta \in (\mathfrak{o}_L/\mathfrak{p}^e \mathfrak{o}_L)^\times \mid 1 + N_{L/F}(\eta) \not\equiv 0 \pmod{\mathfrak{p}}\} & (i = 0), \\ \mathfrak{p}^i \mathfrak{o}_L / \mathfrak{p}^e \mathfrak{o}_L & (0 < i \leq e), \\ \emptyset & (i = e+1) \end{cases}$$

and

$$\overline{\mathcal{U}_i} = \begin{cases} \mathfrak{o}^\times / (1 + \mathfrak{p}^e) & (i = 0), \\ (1 + \mathfrak{p}^i) / (1 + \mathfrak{p}^e) & (0 < i \leq e), \\ \emptyset & (i = e+1). \end{cases}$$

When $0 \leq i < e/2$, we can check that

$$\overline{\mathfrak{o}_{L,i}} \setminus \overline{\mathfrak{o}_{L,i+1}} \ni \eta \pmod{\mathfrak{p}^e \mathfrak{o}_L} \mapsto 1 + N_{L/F}(\eta) \pmod{\mathfrak{p}^e} \in \overline{\mathcal{U}_{2i}} \setminus \overline{\mathcal{U}_{2i+1}}$$

is a surjective map having the fiber of cardinality $q^{e-1}(q+1)$ at each point.

When $e/2 \leq i \leq e$, it is immediate that

$$\mu(1 + N_{L/F}(\eta)) = 1, \quad \eta \in \overline{\mathfrak{o}_{L,i}}.$$

Hence we have

$$\begin{aligned} \sum_{\substack{\eta \in \mathfrak{o}_L/\mathfrak{p}^e \mathfrak{o}_L \\ 1+N_{L/F}(\eta) \in \mathfrak{o}^\times}} \mu(1 + N_{L/F}(\eta)) &= \sum_{i=0}^e \sum_{\eta \in \overline{\mathfrak{o}_{L,i}} \setminus \overline{\mathfrak{o}_{L,i+1}}} \mu(1 + N_{L/F}(\eta)) \\ &= q^{e-1}(q+1) \sum_{0 \leq i < e/2} \sum_{\varepsilon \in \overline{\mathcal{U}_{2i}} \setminus \overline{\mathcal{U}_{2i+1}}} \mu(\varepsilon) + \sum_{e/2 \leq i \leq e} \#(\overline{\mathfrak{o}_{L,i}} \setminus \overline{\mathfrak{o}_{L,i+1}}). \end{aligned}$$

Then the Lemma follows from

$$\sum_{\varepsilon \in \overline{\mathcal{U}_{2i}} \setminus \overline{\mathcal{U}_{2i+1}}} \mu(\varepsilon) = \begin{cases} 0 & (0 \leq i < \frac{e-1}{2}), \\ -1 & (i = \frac{e-1}{2}) \end{cases}$$

and

$$\sum_{e/2 \leq i \leq e} \#(\overline{\mathfrak{o}_{L,i}} \setminus \overline{\mathfrak{o}_{L,i+1}}) = q^{2e-2\lceil e/2 \rceil}.$$

where $\lceil \cdot \rceil : \mathbb{R} \rightarrow \mathbb{Z}$ is the ceiling function. □

Lemma 6.5. *Let $B \in \mathcal{B}(S, \Lambda)[\pi]^K$ and $\eta \in \mathfrak{o}$ with $\frac{\mathfrak{a}^6 \mathfrak{d}}{4} + N_{L/F}(\eta) \in \varpi^j \mathfrak{o}^\times$ ($0 \leq j \leq e$).*

(i) *If we set $Y_\eta = -\mathfrak{a}^2 S^\dagger + X_\eta$, then we have*

$$\det Y_\eta = -\frac{\mathfrak{a}^4 \mathfrak{d}}{4} - \mathfrak{a}^{-2} N_{L/F}(\eta).$$

(ii) *There exist $\tau \in \mathfrak{o}_L^\times$ and $A \in \mathbf{GL}_2(\mathfrak{o})$ such that*

$$Y_\eta = \tau \begin{bmatrix} \varpi^j & 0 \\ 0 & 1 \end{bmatrix} A.$$

(iii) *For $n \in \mathbb{Z}$ and $a \in \mathfrak{o}^\times$,*

$$\begin{aligned} &B \left(\mathfrak{m}(\varpi^n a 1_2, \varpi^n a) \iota \left(\begin{bmatrix} 0 & -1 \\ 1 & \eta^\dagger \end{bmatrix} \right) b \right) \\ &= \psi \left(-\varpi^n a \cdot \frac{\mathfrak{a}^4 \mathfrak{d}}{2} \left(\frac{\mathfrak{a}^6 \mathfrak{d}}{4} + N_{L/F}(\eta) \right)^{-1} \right) \cdot B(h(e+n-2j, j)). \end{aligned}$$

Proof. (i) By noting that $\det S = -\mathbf{d}/4$, $\det X_\eta = -\mathbf{a}^{-2}\mathbf{N}_{L/F}(\eta)$, and $\mathrm{tr}(SX_\eta) = \mathrm{tr}(X_\eta S^\dagger) = 0$, we obtain

$$\det Y_\eta = \frac{1}{2}\mathrm{tr}(Y_\eta Y_\eta^\dagger) = \frac{1}{2}\mathrm{tr}(\mathbf{a}^4 S S^\dagger + X_\eta^\dagger X_\eta) = \mathbf{a}^4 \det S + \det(X_\eta) = -\frac{\mathbf{a}^4 \mathbf{d}}{4} - \mathbf{a}^{-2}\mathbf{N}_{L/F}(\eta).$$

(ii) By (i) and the assumption that $\frac{\mathbf{a}^6 \mathbf{d}}{4} + \mathbf{N}_{L/F}(\eta) \in \varpi^j \mathfrak{o}^\times$, we have $Y_\eta \in \mathbf{GL}_2(F)$. The disjoint union

$$\mathbf{GL}_2(F) = \sqcup_{m \geq 0} L^\times \begin{bmatrix} \varpi^m & 0 \\ 0 & 1 \end{bmatrix} \mathbf{GL}_2(\mathfrak{o}_L).$$

from [43, Lemma 2–4] implies that there exist $m \in \mathbb{Z}_{\geq 0}$, $\ell \in \mathbb{Z}$, $\tau \in \mathfrak{o}_L^\times$, $A \in \mathbf{GL}_2(\mathfrak{o})$, such that

$$Y_\eta = (\varpi^\ell \tau) \begin{bmatrix} \varpi^m & 0 \\ 0 & 1 \end{bmatrix} A = \tau \begin{bmatrix} \varpi^{m+\ell} & 0 \\ 0 & \varpi^\ell \end{bmatrix} A.$$

We now prove that $m = j$ and $\ell = 0$. Because $\tau \in \mathbf{GL}_2(\mathfrak{o})$, the Smith normal form of Y_η is $\begin{bmatrix} \varpi^{m+\ell} & 0 \\ 0 & \varpi^\ell \end{bmatrix}$. Hence by the theory of the Smith normal form, we only have to show that the elements of Y_η are coprime. This follows from the identity $\mathrm{tr}(Y_\eta S) = \frac{\mathbf{a}^2 \mathbf{d}}{2}$ and the assumption $\frac{\mathbf{d}}{2} \in \mathfrak{o}^\times$.

(iii) By using (6.1), we can check that

$$\iota\left(\begin{bmatrix} 1 & 0 \\ -\eta^\dagger & 1 \end{bmatrix}\right) = \begin{bmatrix} 1_2 & 0 \\ X_\eta^\dagger & 1_2 \end{bmatrix}, \quad \iota\left(\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}\right) b \iota\left(\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}\right) = \begin{bmatrix} 1_2 & 0 \\ -\mathbf{a}^2 S & \varpi^{e_{12}} \end{bmatrix}.$$

Hence we have

$$\mathbf{m}(\varpi^n a 1_2, \varpi^n a) \iota\left(\begin{bmatrix} 0 & -1 \\ 1 & \eta^\dagger \end{bmatrix}\right) b = \begin{bmatrix} \varpi^n a 1_2 & 0 \\ Y_\eta^\dagger & \varpi^{e_{12}} \end{bmatrix} \iota\left(\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}\right).$$

By (ii), we may choose elements $\tau \in \mathfrak{o}_L^\times$ and $A \in \mathbf{GL}_2(\mathfrak{o})$ satisfying $Y_\eta = \tau \begin{bmatrix} \varpi^j & 0 \\ 0 & 1 \end{bmatrix} A$. Then, a computation shows that

$$\begin{bmatrix} \varpi^n a 1_2 & 0 \\ Y_\eta^\dagger & \varpi^{e_{12}} \end{bmatrix} = \mathbf{n}(\varpi^n a (\det Y_\eta)^{-1} Y_\eta) \begin{bmatrix} \tau & 0 \\ 0 & \tau^\dagger \end{bmatrix} h(e + n - 2j, j) k$$

where $k = \begin{bmatrix} A & 0 \\ 0 & {}^t A^\dagger \end{bmatrix} \begin{bmatrix} a \varpi^j (\det Y_\eta)^{-1} 1_2 & 0 \\ 0 & 1_2 \end{bmatrix} \begin{bmatrix} 0 & -1_2 \\ 1_2 & \varpi^e (\det Y_\eta)^{-1} Y_\eta \end{bmatrix} \in K$. Since Λ is unramified and B is K -invariant, we have

$$\begin{aligned} & B\left(\mathbf{m}(\varpi^n a 1_2, \varpi^n a) \iota\left(\begin{bmatrix} 0 & -1 \\ 1 & \eta^\dagger \end{bmatrix}\right) b\right) \\ &= \psi(\varpi^n a (\det Y_\eta)^{-1} \cdot \frac{\mathbf{a}^2 \mathbf{d}}{2}) \cdot B(h(e + n - 2j, j)) \\ &= \psi\left(-\varpi^n a \cdot \frac{\mathbf{a}^4 \mathbf{d}}{2} \left(\frac{\mathbf{a}^6 \mathbf{d}}{4} + \mathbf{N}_{L/F}(\eta)\right)^{-1}\right) \cdot B(h(e + n - 2j, j)). \end{aligned} \quad \square$$

Proposition 6.6. *Suppose that π is of type I or IIb. Let $\mu : F^\times \rightarrow \mathbb{C}^1$ and $\Lambda : L^\times \rightarrow \mathbb{C}^1$ be characters such that $\mathrm{cond}(\mu) = e > 0$ and Λ is unramified. Suppose that $S = \begin{bmatrix} \mathbf{a} & \mathbf{b}/2 \\ \mathbf{b}/2 & \mathbf{c} \end{bmatrix} \in \mathrm{Sym}^2(F) \cap \mathbf{GL}_2(\mathfrak{o})$ satisfies that L/F is an unramified field extension and $2d \in \mathfrak{o}^\times$. When $\phi = 1_{\mathfrak{p}^e \mathfrak{o}_L \oplus 1 + \mathfrak{p}^e \mathfrak{o}_L}$ and $\mathrm{Re}(s) > 1$, we have*

$$\begin{aligned} Z(\phi, B_\pi^0, s, \mu; b) &= q^{e(s - \frac{11}{2}) + 5} (q^4 - 1)^{-1} (q - 1)^{-1} \mu(-2^{-1} \mathbf{d})^{-1} W_F(\mu, \psi), \\ Z(\widehat{\phi}, B_\pi^0, s, \mu; b) &= (-1)^e q^{e(3s - \frac{11}{2}) + 5} (q^4 - 1)^{-1} (q - 1)^{-1} \Lambda(\varpi)^{-e} \mu(-2^{-1} \mathbf{a}^2) W_L(\mu_L, \psi_L) W_F(\mu, \psi). \end{aligned}$$

Proof. Let $K^\#(\mathfrak{p}^e) = \{k^\# \in K^\# \mid k^\# \equiv 1_2 \pmod{\mathfrak{p}^e \mathfrak{o}_L}\}$ be the principal congruence subgroup of level $\mathfrak{p}^e \mathfrak{o}_L$. If $\phi \in \mathcal{S}(L^2)$ is right $K^\#(\mathfrak{p}^e)$ -invariant, the function

$$K^\# \ni k^\# \mapsto B(\mathbf{m}(a 1_2, 1) \iota(k^\#) b) f_\phi^{(s, \Lambda, \mu)}(k^\#) \in \mathbb{C}$$

is left $B^\#(\mathfrak{o})$ -invariant and right $K^\#(\mathfrak{p}^e)$ -invariant because $b^{-1}\iota(K^\#(\mathfrak{p}^e))b \subset K$. Since $K^\#(\mathfrak{p}^e)$ is a normal subgroup of $K^\#$ and $B^\#(\mathfrak{o})K^\#(\mathfrak{p}^e) = K_0^\#(\mathfrak{p}^e)$, we have

$$\begin{aligned}
& Z(\phi, B_\pi^0, s, \mu; b) \\
&= \sum_{[\gamma] \in B^\#(\mathfrak{o}) \backslash K^\# / K^\#(\mathfrak{p}^e)} \text{vol}(B^\#(\mathfrak{o})\gamma K^\#(\mathfrak{p}^e)) f_\phi^{(s, \Lambda, \mu)}(\gamma) \int_{F^\times} B_\pi^0(\mathfrak{m}(a1_2, a)\iota(\gamma)b)\mu(a)|a|^{s-1} d^\times a \\
&= [K^\# : K_0^\#(\mathfrak{p}^e)]^{-1} \sum_{[\gamma] \in K_0^\#(\mathfrak{p}^e) \backslash K^\#} f_\phi^{(s, \Lambda, \mu)}(\gamma) \int_{F^\times} B_\pi^0(\mathfrak{m}(a1_2, a)\iota(\gamma)b)\mu(a)|a|^{s-1} d^\times a \\
(6.12) \quad &= q^{-2e+2}(q^2 + 1)^{-1} \sum_{[\gamma] \in K_0^\#(\mathfrak{p}^e) \backslash K^\#} f_\phi^{(s, \Lambda, \mu)}(\gamma) \int_{F^\times} B_\pi^0(\mathfrak{m}(a1_2, a)\iota(\gamma)b)\mu(a)|a|^{s-1} d^\times a.
\end{aligned}$$

A complete set of representative for $K^\#(\mathfrak{p}^e) \backslash K^\#$ is given by

$$(6.13) \quad \begin{bmatrix} 1 & 0 \\ \xi & 1 \end{bmatrix} \quad (\xi \in \mathfrak{p}\mathfrak{o}_L / \mathfrak{p}^e \mathfrak{o}_L), \quad \begin{bmatrix} 0 & -1 \\ 1 & \eta^\dagger \end{bmatrix} \quad (\eta \in \mathfrak{o}_L / \mathfrak{p}^e \mathfrak{o}_L).$$

Now we prove the first equation. By Lemma 6.3, when γ runs through the above elements, then we have $f_\phi^{(s, \Lambda, \mu)}(1_2) = q^{-2e+2}(q^2 - 1)^{-1}$ and $f_\phi^{(s, \Lambda, \mu)}(\gamma) = 0$ if $\gamma \neq 1$. By substituting this in (6.12) and Lemma 6.1, we have

$$\begin{aligned}
& Z(\phi, B_\pi^0, s, \mu; b) \\
&= q^{-4e+4}(q^4 - 1)^{-1} \int_{F^\times} B_\pi^0 \left(\begin{bmatrix} a\varpi^e 1_2 & aS^\dagger \\ 0 & 1_2 \end{bmatrix} \right) \mu(a)|a|^{s-1} d^\times a \\
&= q^{-4e+4}(q^4 - 1)^{-1} \int_{F^\times} B_\pi^0 \left(\begin{bmatrix} a\varpi^e 1_2 & 0 \\ 0 & 1_2 \end{bmatrix} \right) \psi(\text{atr}(SS^\dagger))\mu(a)|a|^{s-1} d^\times a \\
&= q^{-4e+4}(q^4 - 1)^{-1} \mu(-2^{-1}\mathfrak{d})^{-1} \sum_{n \in \mathbb{Z}} B_\pi^0 \left(\begin{bmatrix} \varpi^{e+n} 1_2 & 0 \\ 0 & 1_2 \end{bmatrix} \right) \mu(\varpi)^n q^{-n(s-1)} \int_{\mathfrak{o}^\times} \psi(\varpi^n a)\mu(a) d^\times a \\
&= q^{e(s-\frac{11}{2})+5}(q^4 - 1)^{-1}(q - 1)^{-1} \mu(-2^{-1}\mathfrak{d})^{-1} W_F(\mu, \psi).
\end{aligned}$$

Next, we prove the second equation. When we replace ϕ by $\widehat{\phi}$ in (6.12) and consider the representative (6.13), we can ignore the contribution of ξ -terms in the summation from Lemma 6.3. Hence, by noting the K -invariance of B , we have

$$\begin{aligned}
(6.14) \quad & Z(\widehat{\phi}, B_\pi^0, s, \mu; b) = q^{e(2s-5)+4}(q^4 - 1)^{-1} \Lambda(\varpi)^{-e} W_L(\mu_L, \psi_L) \\
& \times \sum_{\eta \in \mathfrak{o}_L / \mathfrak{p}^e \mathfrak{o}_L} \int_{F^\times} B_\pi^0(\mathfrak{m}(a1_2, a)\iota \left(\begin{bmatrix} 0 & -1 \\ 1 & \eta^\dagger \end{bmatrix} \right) b)\mu(a)|a|^{s-1} d^\times a.
\end{aligned}$$

By replacing η by $\eta + \varpi^e u$ for some $u \in \mathfrak{o}_L$ if we need, we may assume that $\mathfrak{a}^6 \mathfrak{d} + \mathfrak{N}_{L/F}(\eta) \in \varpi^j \mathfrak{o}_L^\times$ with $0 \leq j \leq e$. Then, Lemmas 6.1 and 6.5 (iii) imply

$$\begin{aligned}
& \int_{F^\times} B_\pi^0(\mathfrak{m}(a1_2, a)\iota \left(\begin{bmatrix} 0 & -1 \\ 1 & \eta^\dagger \end{bmatrix} \right) b)\mu(a)|a|^{s-1} d^\times a \\
&= \sum_{n \in \mathbb{Z}} B_\pi^0(h(e + n - 2j, j))\mu(\varpi)^n q^{-n(s-1)} \int_{\mathfrak{o}^\times} \psi \left(-\varpi^n a \cdot \frac{\mathfrak{a}^4 \mathfrak{d}}{2} \left(\frac{\mathfrak{a}^6 \mathfrak{d}}{4} + \mathfrak{N}_{L/F}(\eta) \right)^{-1} \right) \mu(a) d^\times a
\end{aligned}$$

$$\begin{aligned}
&= \mu \left(-2\mathbf{a}^{-4}\mathbf{d}^{-1}\varpi^{-j} \left(\frac{\mathbf{a}^6\mathbf{d}}{4} + N_{L/F}(\eta) \right) \right) \sum_{n \in \mathbb{Z}} B_{\pi}^0(h(e+n-2j, j)) \mu(\varpi)^n q^{-n(s-1)} \int_{\mathfrak{o}^{\times}} \psi(\varpi^{n-j}a) \mu(a) d^{\times}a \\
&= q^{-\frac{e}{2}+1+(e-j)(s-1)} (q-1)^{-1} \mu(-2\mathbf{a}^{-4}\mathbf{d}^{-1}) \mu \left(\frac{\mathbf{a}^6\mathbf{d}}{4} + N_{L/F}(\eta) \right) W_F(\mu, \psi) \cdot B_{\pi}^0(h(-j, j)).
\end{aligned}$$

The last equation is equal to 0 unless $j = 0$ from (6.8). By substituting this into (6.14) and Lemma 6.4, we have that $Z(\phi, B_{\pi}^0, s, \mu; b)$ equals

$$\begin{aligned}
&q^{e(3s-\frac{13}{2})+5} (q^4-1)^{-1} (q-1)^{-1} \Lambda(\varpi)^{-e} \mu(-2\mathbf{a}^{-4}\mathbf{d}^{-1}) \\
&\times W_L(\mu_L, \psi_L) W_F(\mu, \psi) \sum_{\substack{\eta \in \mathfrak{o}_L/\mathfrak{p}^e \mathfrak{o}_L \\ \mathbf{a}^6\mathbf{d} + N_{L/F}(\eta) \in \mathfrak{o}^{\times}}} \mu \left(\frac{\mathbf{a}^6\mathbf{d}}{4} + N_{L/F}(\eta) \right) \\
&= (-1)^e q^{e(3s-\frac{11}{2})+5} (q^4-1)^{-1} (q-1)^{-1} \Lambda(\varpi)^{-e} \mu(-2^{-1}\mathbf{a}^2) W_L(\mu_L, \psi_L) W_F(\mu, \psi). \quad \square
\end{aligned}$$

Corollary 6.7. *Let π , S , μ , and Λ be as in Proposition 6.6, then*

$$\begin{aligned}
\varepsilon(\pi, s, \mu, \psi) &= (-1)^e q^{4e(\frac{1}{2}-s)} \Lambda(\varpi)^{-e} \mu(-\mathbf{a}^{-2}\mathbf{d}) \overline{W_L(\mu_L, \psi_L) W_F(\mu, \psi)^2} \\
&= q^{4e(\frac{1}{2}-s)} \Lambda(\varpi)^{-e} \mu(-\mathbf{a}^{-2}\mathbf{d}) \overline{W_F(\mu, \psi)^4}.
\end{aligned}$$

Proof. The first equality follows from Proposition 6.6 and (6.4); the second equality is due to Lemma 6.2. \square

6.4. Computation for old forms of type I and IIb (case4). Now we assume that π has the trivial character. Then π has the (S, Λ) -Bessel model if and only if $\Lambda = 1$. Recall that the dimension of $V_{\pi}^{K_0(\mathfrak{p})}$ is equal to 4 or 3 according as π is of type I or type IIb.

Proposition 6.8. *Let π be a smooth admissible irreducible representation of type I or type IIb with trivial central character. Suppose that $\mu : F^{\times} \rightarrow \mathbb{C}^1$ is an unramified character. When $\phi = \mathbb{1}_{\mathfrak{o}_L \oplus \mathfrak{o}_L}$ and $\operatorname{Re}(s) > 1$, we have*

$$\begin{aligned}
Z(\phi, B, s, \mu; \eta) &= \frac{1}{q^2+1} L(s+1/2, \pi, \mu) \\
&\times \left[\eta B + q^{-1} T_{1,0} B + \left\{ \mu(\varpi)^{-1} q^{s+1} + \mu(\varpi) q^{-s+1} - \operatorname{tr}((q^{-1} T_{1,0} + \eta)|_{V_{\pi}^{K_0(\mathfrak{p})}}) \right\} B \right] (1_4)
\end{aligned}$$

for $B \in \mathcal{B}(S, 1)[\pi]^{K_0(\mathfrak{p})}$.

Proof. We will give the proof of the case that π is of type I. In the case of type IIb, the proof is similar.

Let $\{B_j^*\}_{1 \leq j \leq 4} \subset \mathcal{B}(S, 1)[\pi]^{K_0(\mathfrak{p})}$ be an eigenbasis of $T_{1,0}$ with eigenvalues $\{\mu_j\}_{1 \leq j \leq 4}$ and $(\eta_{ij})_{1 \leq i, j \leq 4}$ be the representation matrix of $\eta|_{K_0(\mathfrak{p})}$ with respect to $\{B_j^*\}_{1 \leq j \leq 4}$. Then, the first relation in (6.23) shows that

$$B_j^*(h(l+1, 0)) = \mu_j q^{-3} B_j^*(h(l, 0)), \quad l \in \mathbb{Z}_{\geq 0}.$$

It is easy to check that

$$f_{\phi}^{(s, \Lambda, \mu)}(k^{\#}) = L(s+1, \Lambda \mu_L), \quad k^{\#} \in K^{\#}.$$

By the left $K_0^\#(\mathfrak{p})$ -coset decomposition of $K^\#$;

$$(6.15) \quad K^\# = K_0^\#(\mathfrak{p}) \sqcup \left(\sqcup_{\xi \in \mathfrak{o}_L/\mathfrak{p}\mathfrak{o}_L} \begin{bmatrix} 1 & \xi \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} K_0^\#(\mathfrak{p}^e) \right),$$

the zeta integral (6.3) becomes

$$(6.16)$$

$$\begin{aligned} Z(\phi, B, s, \mu; \eta) &= Z(\phi, \eta B, s, \mu; 1_4) \\ &= [K^\# : K_0^\#(\mathfrak{p})]^{-1} L(s+1, \Lambda\mu_L) \\ &\quad \times \int_{F^\times} \left\{ \eta B(\mathfrak{m}(a1_2, a)) + \sum_{\xi \in \mathfrak{o}_L/\mathfrak{p}\mathfrak{o}_L} \eta B(\mathfrak{m}(a1_2, a)\iota \left(\begin{bmatrix} 1 & \xi \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \right)) \right\} \mu^{-1}(a)|a|^{s-1} d^\times a \\ &= \frac{1}{q^2+1} L(s+1, \Lambda\mu_L) \\ &\quad \times \left\{ \sum_{l \in \mathbb{Z}} \eta B(h(l, 0)) + q^2 \Lambda(\varpi)^{-1} B(h(l+1, 0)) \right\} \mu(\varpi)^l q^{-l(s-1)}. \end{aligned}$$

Since $B(h(l, 0)) = 0$ if $l < 0$ by (6.8), the right hand side equals

$$(6.17)$$

$$\frac{L(s+1, \Lambda\mu_L)}{q^2+1} \left\{ \sum_{l=0}^{\infty} \eta B(h(l, 0)) \mu(\varpi)^l q^{-l(s-1)} + q^{s+1} \Lambda(\varpi)^{-1} \mu(\varpi)^{-1} \sum_{l=0}^{\infty} B(h(l, 0)) \mu(\varpi)^l q^{-l(s-1)} \right\}.$$

By substituting $B = B_j^*$ in (6.17) and $\Lambda = 1$,

$$Z(\phi, B_j^*, s, \mu; \eta) = \frac{L(s+1, \Lambda\mu_L)}{q^2+1} \left\{ \sum_{i=1}^4 \frac{\eta_{ij} B_i^*(1_4)}{1 - \mu_j \mu(\varpi) q^{-(s+2)}} + \frac{\mu(\varpi)^{-1} q^{s+1} B_j^*(1_4)}{1 - \mu_i \mu(\varpi) q^{-(s+2)}} \right\}.$$

Thus, we may put

$$\begin{aligned} Z(\phi, B_j^*, s, \mu; \eta) &= \frac{1}{q^2+1} \prod_{j=1}^4 (1 - \mu_j q^{-2} X)^{-1} \\ &\quad \times \frac{A_0 + A_1 X + A_2 X^2 + A_3 X^3 + A_4 X^4}{X(1 - q^{-2} X^2)}, \quad (X = \mu(\varpi) q^{-s}, A_i \in \mathbb{C}, 1 \leq i \leq 4). \end{aligned}$$

By noting that $\{\mu_1, \mu_2, \mu_3, \mu_4\} = \{\alpha\beta\gamma q^{3/2}, \beta\gamma q^{3/2}, \alpha\gamma q^{3/2}, \gamma q^{3/2}\}$ from the table (6.5), we have

$$(6.18) \quad (q^2+1) \cdot \frac{Z(\phi, B_j^*, s, \mu; 1_4)}{L(s+1/2, \pi, \mu)} = \frac{A_0 + A_1 X + A_2 X^2 + A_3 X^3 + A_4 X^4}{X(1 - q^{-2} X^2)}$$

A direct computation shows that

$$(6.19) \quad A_0 = q B_j^*(1_4),$$

$$\begin{aligned}
(6.20) \quad A_1 &= \sum_{i=1}^4 \eta_{ij} B_i^*(1_4) + q B_j^*(1_4) \left\{ \left(\sum_{k=1}^4 q^{-2} \mu_k \right) - q^{-2} \mu_i \right\} \\
&= \eta B_j^*(1_4) + q^{-1} \text{tr}(T_{1,0}|_{V_\pi^{K_0(\mathfrak{p})}}) \cdot B_j^*(1_4) - q^{-1} \cdot T_{1,0} B_j^*(1_4).
\end{aligned}$$

The functional equation (6.4) and the fact $\varepsilon(\pi, s, \mu, \psi) = 1$ from ([30]) show that the right hand side of (6.18) is invariant under the transformation $X \rightarrow X^{-1}$. By comparing the coefficients, we have

$$(6.21) \quad A_2 = q^{-2}(q^2 - 1)A_0, \quad A_3 = -q^{-2}A_1, \quad A_4 = -q^{-2}A_0$$

Substituting (6.19), (6.20), and (6.21) into (6.18), we get

$$\begin{aligned}
&(q^2 + 1) \cdot \frac{Z(\phi, B_j^*, s, \mu; \eta)}{L(s + 1/2, \pi, \mu)} \\
&= \frac{A_0 + A_1 X + q^{-2}(q^2 - 1)A_0 X^2 - q^{-2}A_1 X^3 - q^{-2}A_0 X^4}{X(1 - q^{-2}X^2)} \\
&= A_0(X + X^{-1}) + A_1 \\
&= \left[\eta B + q^{-1}T_{1,0}B + \left\{ \mu(\varpi)^{-1}q^{s+1} + \mu(\varpi)q^{-s+1} - \text{tr}(q^{-1}T_{1,0}|_{V_\pi^{K_0(\mathfrak{p})}}) \right\} B \right] (1_4). \quad \square
\end{aligned}$$

6.5. Computation for newforms of type IIIa (case5). By [32, §9.1], there exists a basis $\{B_1, B_2\}$ of $\mathcal{B}(S, \Lambda)[\pi]^{K_0(\mathfrak{p})}$ such that

$$\begin{aligned}
(6.22) \quad T_{1,0}B_1 &= \alpha\gamma q B_1, \quad T_{0,1}B_2 = \gamma q B_2, \\
T_{0,1}B_1 &= \alpha\gamma^2(\alpha q + 1)q B_1, \quad T_{0,1}B_2 = \alpha\gamma^2(\alpha^{-1}q + 1)q B_2, \\
\eta B_1 &= \alpha\gamma B_2, \quad \eta B_2 = \gamma B_1.
\end{aligned}$$

We now consider the values of B_1 and B_2 at the identity element. We remark that [32, Lemma 9.2] ensures its non-vanishing and

$$\Lambda(\varpi) = \alpha\gamma^2$$

by the condition of central character of π .

Lemma 6.9. *Let $S \in \text{Sym}^2(F)$ such that L/F is an unramified field extension. Suppose that $\mu : F^\times \rightarrow \mathbb{C}^1$ and $\Lambda : L^\times \rightarrow \mathbb{C}^1$ are unramified characters. Then we have*

$$B_2(1_4) = \alpha^{-1}B_1(1_4) \neq 0.$$

Proof. For $B \in \mathcal{B}(S, \Lambda)[\pi]^{K_0(\mathfrak{p})}$ and any non-negative integer l , by Lemmas 5.1 and 5.3 in [32], we get the following relations;

$$\begin{aligned}
(6.23) \quad T_{1,0}B(h(l, 0)) &= q^3 B(h(l + 1, 0)), \\
T_{1,0}B(h(l, 0)s_2) &= q^2(q - 1)B(h(l + 1, 0)) + q^2 B(h(l - 1, 1)s_1 s_2), \\
T_{1,0}B(h(l, 0)s_2 s_1 s_2) &= q^2(q - 1)B(h(l + 1, 0)) + \mu(\varpi)B(h(l - 1, 0)s_2 s_1 s_2) \\
&\quad + (q^2 - 1)B(h(l - 1, 1)s_1 s_2), \\
T_{0,1}B(h(l, 0)) &= q^3(q + 1)B(h(l, 1)), \\
T_{0,1}B(h(l, 0)s_2) &= q^4 B(h(l, 1)s_1 s_2) + q\mu(\varpi)B(h(l - 2, 1)s_1 s_2)
\end{aligned}$$

$$+q^2(q-1)B(h(l,1)) - q\mu(\varpi)B(1_4).$$

Set $B = B_1$. By putting $l = 0$ in the above relations, applying the relation (6.22), and using the equation that $B_2(h(l,0)) = \gamma B_1(h(l-1,0)s_2s_1s_2)$ for $l \geq 0$, we have

$$(6.24) \quad \alpha\gamma q B_1(s_2) = \alpha\gamma(q-1)B_1(1_4) + q^2 B_1(h(-1,1)s_1s_2),$$

$$(6.25) \quad \mu(\varpi)\gamma^{-1}(q^{-1}-1)B_2(1_4) = \alpha\gamma(q-1)B_1(1_4) + (q^2-1)B_1(h(-1,1)s_1s_2),$$

$$(6.26) \quad \mu(\varpi)(\alpha q + 1)B_1(s_2) = q^4 B_1(h(0,1)s_1s_2) + \frac{\mu(\varpi)(\alpha q^2 - \alpha q - q^2 - 1)}{q+1} B_1(1_4).$$

Similarly, by putting $l = 1$ in the third relation in (6.23), we have

$$(6.27) \quad \mu(\varpi)q^{-3}(1-q)B_2(1_4) = \mu(\varpi)\alpha q^{-2}(q-1)B_1(1_4) + (q^2-1)B_1(h(0,1)s_1s_2).$$

From (6.24), (6.25), (6.26), and (6.27), we obtain $B_2(1_4) = \alpha^{-1}B_1(1_4)$. The non-vanishing of B_1 at 1_4 is a consequence of [32, Theorem 9.3]. \square

Proposition 6.10. *Suppose that π is of type IIIa. Let $S \in \text{Sym}^2(F)$ be such that L/F is an unramified field extension. Suppose that $\mu : F^\times \rightarrow \mathbb{C}^1$ and $\Lambda : L^\times \rightarrow \mathbb{C}^1$ are unramified characters. When $\phi = 1_{\mathfrak{o}_L \oplus \mathfrak{o}_L}$ and $\text{Re}(s) > 1$, we have*

$$Z(\phi, B, s, \mu; \eta) = \frac{\Lambda(\varpi)^{-1}\mu(\varpi)^{-1}q^{s+1}}{q^2+1} L(s+1/2, \pi, \mu) \cdot B(1_4), \quad B \in \mathcal{B}(s, \Lambda)[\pi]^{K_0(\mathfrak{p})}.$$

Proof. By substituting $B = B_i$ in (6.17) for $i = 1, 2$, the statement is immediate for $B = B_i$ by (6.22), the first relation in (6.23), and Lemma 6.9. Since B_1 and B_2 span $\mathcal{B}(S, \Lambda)[\pi]^{K_0(\mathfrak{p})}$, we complete the proof. \square

Corollary 6.11. *Let π , S , μ , and Λ be as in Proposition 6.10, then*

$$\varepsilon(\pi, s, \mu, \psi) = \mu(\varpi)^2 q^{2(\frac{1}{2}-s)}.$$

6.6. Computation for newforms of type VIb (case6). Suppose that π is of type VIb. By [32, Table 3], any element $B \in \mathcal{B}(S, \Lambda)[\pi]^{K_0(\mathfrak{p})}$ satisfies

$$(6.28) \quad T_{1,0}B = \gamma q B, \quad \eta B = \gamma B.$$

By [32, Theorem 9.3], the value $B(1_4) \neq 0$ if and only if $B = 0$.

Proposition 6.12. *Suppose that π is of type VIb. Let $S \in \text{Sym}^2(F)$ be such that L/F is an unramified field extension. Suppose that $\mu : F^\times \rightarrow \mathbb{C}^1$ and $\Lambda : L^\times \rightarrow \mathbb{C}^1$ are unramified characters. When $\phi = 1_{\mathfrak{o}_L \oplus \mathfrak{o}_L}$ and $\text{Re}(s) > 1$, we have*

$$Z(\phi, B, s, \mu; \eta) = \frac{\Lambda(\varpi)^{-1}\mu(\varpi)^{-1}q^{s+1}}{q^2+1} L(s+1/2, \pi, \mu) \cdot B(1_4), \quad B \in \mathcal{B}(s, \Lambda)[\pi]^{K_0(\mathfrak{p})}.$$

Proof. In our case, the relations (6.23) are also true. The proof is similar to that of Proposition 6.10. \square

Corollary 6.13. *Let π , S , μ , and Λ be as in Proposition 6.12, then*

$$\varepsilon(\pi, s, \mu, \psi) = \mu(\varpi)^2 q^{2(\frac{1}{2}-s)}.$$

6.7. The values of Bessel models at 1_4 for old forms. For quasi-characters χ, χ' , and σ on F^\times which are trivial on \mathfrak{o}^\times , the representation $(\chi \times \chi' \rtimes \sigma, V)$ can be realized as the right regular representation of the space of all smooth functions f on $\mathbf{G}(F)$ satisfying

$$f \left(\begin{bmatrix} a & * & * & * \\ 0 & b & * & * \\ 0 & 0 & ca^{-1} & 0 \\ 0 & 0 & * & cb^{-1} \end{bmatrix} g \right) = |a^2 b| |c|^{-\frac{3}{2}} \chi(a) \chi'(b) \sigma(c) f(g), \quad a, b, c \in F^\times, g \in \mathbf{G}(F).$$

The space V has a unique K -invariant element $f_K \in V$ such that $f_K(1_4) = 1$. Let $I = K \cap \begin{bmatrix} \mathfrak{o} & \mathfrak{p} & \mathfrak{o} & \mathfrak{o} \\ \mathfrak{o} & \mathfrak{o} & \mathfrak{o} & \mathfrak{o} \\ \mathfrak{p} & \mathfrak{p} & \mathfrak{o} & \mathfrak{o} \\ \mathfrak{p} & \mathfrak{p} & \mathfrak{p} & \mathfrak{o} \end{bmatrix}$ be the Iwahori subgroup of K . For $w \in W$, let $f_w \in V^I$ be the unique element such that $f_w|_K = \mathbb{1}_{IwI}$.

By [33, Table 3], the space V^I is 8-dimensional space spanned by

$$f_{1_4}, \quad f_{s_1}, \quad f_{s_2}, \quad f_{s_2 s_1}, \quad f_{s_1 s_2 s_1}, \quad f_{s_1 s_2}, \quad f_{s_1 s_2 s_1 s_2}, \quad f_{s_2 s_1 s_2}.$$

A basis of 4-dimensional space $V^{K_0(\mathfrak{p})}$ is given by

$$f_1^I := f_{1_4} + f_{s_1}, \quad f_2^I := f_{s_2} + f_{s_2 s_1}, \quad f_3^I := f_{s_1 s_2 s_1} + f_{s_1 s_2}, \quad f_4^I := f_{s_1 s_2 s_1 s_2} + f_{s_2 s_1 s_2}.$$

Recall that π is isomorphic to $\chi \times \chi' \rtimes \sigma$ or $\chi 1_{\mathbf{GL}_2} \rtimes \sigma$ for some characters χ, χ' , and σ on F^\times according as π is of type I or IIb. The representation $(\chi 1_{\mathbf{GL}_2} \rtimes \sigma, V)$ can be realized as a subrepresentation of $|\cdot|^{1/2} \chi \times |\cdot|^{-1/2} \chi \rtimes \sigma$, and a basis of 3-dimensional space $V^{K_0(\mathfrak{p})}$ is given by

$$f_1^{\text{IIb}} := f_{1_4} + f_{s_1}, \quad f_2^{\text{IIb}} := f_{s_2} + f_{s_2 s_1} + f_{s_1 s_2 s_1} + f_{s_1 s_2}, \quad f_3^{\text{IIb}} := f_{s_1 s_2 s_1 s_2} + f_{s_2 s_1 s_2}.$$

The representation matrices of operators $T_{1,0}$ and η on $V_\pi^{K_0(\mathfrak{p})}$ are given in [33, Table 3] as follows;

$$(6.29) \quad \begin{array}{c|c|c} & \text{type I} & \text{type IIb} \\ \hline T_{1,0} & \begin{bmatrix} \alpha\beta\gamma q^{3/2} & 0 & 0 & 0 \\ \alpha\beta\gamma(q-1)q^{1/2} & \beta\gamma q^{3/2} & 0 & 0 \\ \alpha\beta\gamma(q-1)q^{1/2} & \beta\gamma(q-1)q^{1/2} & \alpha\gamma q^{3/2} & 0 \\ \alpha\beta\gamma(q-1)q^{1/2} & \beta\gamma(q-1)q^{1/2} & \alpha\gamma(q-1)q^{1/2} & \gamma q^{3/2} \end{bmatrix} & \begin{bmatrix} \alpha^2\gamma q^{3/2} & 0 & 0 \\ \alpha^2\gamma(q-1)q^{1/2} & \alpha\gamma q^2 & 0 \\ \alpha^2\gamma(q-1)q^{1/2} & \alpha\gamma(q^2-1) & \gamma q^{3/2} \end{bmatrix} \\ \hline \eta & \begin{bmatrix} 0 & 0 & 0 & \gamma q^{3/2} \\ 0 & 0 & \alpha\gamma q^{1/2} & 0 \\ 0 & \beta\gamma q^{-1/2} & 0 & 0 \\ \alpha\beta\gamma q^{-3/2} & 0 & 0 & 0 \end{bmatrix} & \begin{bmatrix} 0 & 0 & \gamma q^{3/2} \\ 0 & \alpha\gamma & 0 \\ \alpha^2\gamma q^{-3/2} & 0 & 0 \end{bmatrix} \end{array}$$

Now we assume that π has the trivial character. Let (ℓ_π^0, f_K) be the Bessel data for π defined above. For $\bullet \in \{\text{I}, \text{IIb}\}$, we set a basis $\{B_i^\bullet\}_{1 \leq i \leq \dim V_\pi^{K_0(\mathfrak{p})}}$ of $V_\pi^{K_0(\mathfrak{p})}$ by

$$B_i^\bullet(g) := \ell_\pi^0(\pi(g)f_i^\bullet), \quad g \in \mathbf{G}(F).$$

Then the following result holds.

Proposition 6.14. *Let π be a smooth admissible irreducible K -spherical representation of type I or IIb, and $\{B_i^\bullet\}$ be as above for $\bullet \in \{\text{I}, \text{IIb}\}$. Then*

(i) *If π is of type I,*

$$\begin{aligned} B_1^I(1_4) &= \frac{\alpha\beta}{(q-\alpha)(q-\beta)}, & B_2^I(1_4) &= \frac{-q\beta}{(q-\alpha)(q-\beta)}, \\ B_3^I(1_4) &= \frac{-q\alpha}{(q-\alpha)(q-\beta)}, & B_4^I(1_4) &= \frac{q^2}{(q-\alpha)(q-\beta)}. \end{aligned}$$

(ii) If π is of type IIb,

$$B_1^{\text{Ib}}(1_4) = \frac{\alpha^2}{(q^{1/2} - \alpha)(q^{3/2} - \alpha)}, \quad B_2^{\text{Ib}}(1_4) = \frac{-q^{1/2}(1 + q)\alpha}{(q^{1/2} - \alpha)(q^{3/2} - \alpha)},$$

$$B_3^{\text{Ib}}(1_4) = \frac{q^2}{(q^{1/2} - \alpha)(q^{3/2} - \alpha)}.$$

Proof. Set $d := \dim V_\pi^{K_0(\mathfrak{p})}$. By the construction of the elements $\{f_i^\bullet\}_{1 \leq i \leq d}$, we have $B_\pi^0 = \sum_{i=1}^d B_i^\bullet$. For any $\ell \in \mathbb{Z}_{\geq 0}$, we act by $T_{1,0}^\ell$ on the both sides and evaluate for $g = 1_4$, then the first relation in (6.23) implies that

$$B_\pi^0(h(\ell, 0)) = \sum_{i=1}^d (T_{1,0}^\ell B_i^\bullet)(1_4).$$

The values of the left hand side is given in [6, Corollary 1.8]. Hence by using the table (6.29) and a direct calculation, we can solve the values of $B_i^\bullet(1_4)$. \square

6.8. Local periods. For an irreducible admissible unitalizable $K_0(\mathfrak{p})$ -spherical representation (π, V_π) on $\mathbf{G}(F)$ having the (S, Λ) -Bessel model, we fix a pair $(\ell^\pi, \xi^\pi) \in (V_\pi^*)^{S, \Lambda} \times V_\pi^{K_0(\mathfrak{p})}$ satisfying $\ell_\pi(\xi_\pi) = 1$. When π is K -spherical, we choose (ℓ^π, ξ^π) as the unramified Bessel datum (ℓ_0^π, ξ_0^π) . Then, there exists a unique $\mathbf{G}(F)$ -invariant inner product $\langle \cdot | \cdot \rangle$ on V_π such that $\langle \xi^\pi | \xi^\pi \rangle = 1$. For $s \in \mathbb{C}$, $\phi \in \mathcal{S}(L^2)$, $g \in \mathbf{G}(F)$, and a character μ on F^\times , we define the local period of π as

$$\mathbb{I}_{\pi, K_0(\mathfrak{p})}^{(s)}(\xi^\pi, \phi, \Lambda, \mu; g) = \sum_{B \in \mathcal{B}((V_\pi^*)^{S, \Lambda}, K_0(\mathfrak{p}))} Z^*(\phi, B, s, \mu; g) \overline{B(1_4)}.$$

where

$$Z^*(\phi, B, s, \mu; g) := L(s + 1/2, \pi, \mu)^{-1} Z(\phi, B, s, \mu; g)$$

is the normalized zeta integral and $\mathcal{B}((V_\pi^*)^{S, \Lambda}, K_0(\mathfrak{p}))$ is an orthonormal basis of $\mathcal{B}(S, \Lambda)[\pi]^{K_0(\mathfrak{p})}$ with respect to the inner product on $\mathcal{B}(S, \Lambda)[\pi]$ via the $\mathbf{G}(F)$ -isomorphism

$$(6.30) \quad V_\pi \ni \xi \mapsto (g \mapsto \ell_\pi(\pi(g)\xi)) \in \mathcal{B}(S, \Lambda)[\pi].$$

6.8.1. Local periods for type I and IIb. When π is of type I or IIb, we recall that the representation space can be realized as in §6.7. We choose $\xi^\pi = \xi_0^\pi = f_K$. Then, since we are dealing with induced representations from unitary characters, the standard inner product $\langle f | g \rangle := \int_K f(k) \overline{g(k)} dk$ ($f, g \in V_\pi$) becomes $\mathbf{G}(F)$ -invariant and satisfies $\langle \xi_0^\pi | \xi_0^\pi \rangle = 1$.

For $\bullet \in \{\text{I}, \text{IIb}\}$, the set $\{\langle B_i^\bullet | B_i^\bullet \rangle^{-\frac{1}{2}} B_i^\bullet\}_{1 \leq i \leq \dim V_\pi^{K_0(\mathfrak{p})}}$ forms an orthonormal basis of $\mathcal{B}(S, \Lambda)[\pi]^{K_0(\mathfrak{p})}$. By [37,], the values $\langle B_i^\bullet | B_i^\bullet \rangle$ are given as follows:

$$\langle B_i^{\text{I}} | B_i^{\text{I}} \rangle = q^{i-1}(q+1) \quad (1 \leq i \leq 4),$$

$$\langle B_1^{\text{IIb}} | B_1^{\text{IIb}} \rangle = (q+1), \quad \langle B_2^{\text{IIb}} | B_2^{\text{IIb}} \rangle = q(q+1)^2, \quad \langle B_3^{\text{IIb}} | B_3^{\text{IIb}} \rangle = q^3(q+1).$$

By Propositions 6.8 and 6.14, and the matrix representation of $T_{1,0}$ and η given in (6.29), we can calculate the local period by a direct calculation. ²

²We have used MATHEMATICA 13.

Proposition 6.15. *Let π be a smooth admissible irreducible unitary representation of type I or type IIb with trivial central character. Suppose that $\mu : F^\times \rightarrow \mathbb{C}^1$ is an unramified character. When $\phi = \mathbb{1}_{\mathfrak{o}_L \oplus \mathfrak{o}_L}$, we have*

$$\begin{aligned} \mathbb{I}_{\pi, K_0(\mathfrak{p})}^{(s)}(\xi_0^\pi, \phi, \Lambda, \mu; \eta) &= \frac{2(q-1)}{q^5(q^2+1)} L(1, \pi; \text{Std}) \\ &\quad \times \left\{ \mu(\varpi)^{-1} q^{s+1} + \mu(\varpi) q^{-s+1} - \frac{1}{q+1} \text{tr}(T_{1,0} + q\eta|_{V_\pi^{K_0(\mathfrak{p})}}) \right\}, \end{aligned}$$

where $L(s, \pi; \text{Std})$ is the standard L -function of π defined by

$$L(s, \pi; \text{Std}) = (1 - \alpha q^{-s})^{-1} (1 - \beta q^{-s})^{-1} (1 - \alpha^{-1} q^{-s})^{-1} (1 - \beta^{-1} q^{-s})^{-1} (1 - q^{-s})^{-1}.$$

or

$$L(s, \pi; \text{Std}) = (1 - \alpha q^{-s+\frac{1}{2}})^{-1} (1 - \alpha q^{-s-\frac{1}{2}})^{-1} (1 - \alpha^{-1} q^{-s+\frac{1}{2}})^{-1} (1 - \alpha^{-1} q^{-s-\frac{1}{2}})^{-1} (1 - q^{-s})^{-1}.$$

according as π is of type I or IIb

6.8.2. Local period for type IIIa. Suppose that π is of type IIIa. Then, as remarked before, $|\alpha| = |\gamma| = 1$ and $\alpha \neq 1$. By Lemma 6.9, there exists a basis $\{B_i\}_{i=1,2}$ of $\mathcal{B}(S, \Lambda)[\pi]^{K(\mathfrak{p})}$, unique up to constant, satisfying the relations (6.22) and $B_2(1_4) = \alpha^{-1} B_1(1_4) \neq 0$. Thus, we can uniquely fix $\{B_i^{\text{IIIa}}\}_{i=1,2}$ by requiring $B_1^{\text{IIIa}}(1_4) = 1$ and the datum (ℓ_π, ξ^π) by $\ell_\pi(\xi^\pi) = B_1^{\text{IIIa}}$ to form (6.30) and induce the $G(F)$ -invariant inner product on $\mathcal{B}(S, \Lambda)[\pi]$ such that $\langle B_1^{\text{IIIa}} | B_1^{\text{IIIa}} \rangle = 1$. Since $\langle B_1 | B_2 \rangle = 0$ by [7, Lemma 2.11], we then have $\langle B_-^{\text{IIIa}} | B_-^{\text{IIIa}} \rangle = 1$ and $\langle B_+^{\text{IIIa}} | B_-^{\text{IIIa}} \rangle = 0$ due to $|\alpha| = 1$.

Proposition 6.16. *Let π be a smooth admissible irreducible unitary representation of type IIIa. Suppose that $\mu : F^\times \rightarrow \mathbb{C}^1$ is an unramified character. When $\phi = \mathbb{1}_{\mathfrak{o}_L \oplus \mathfrak{o}_L}$, we have*

$$\mathbb{I}_{\pi, K_0(\mathfrak{p})}^{(s)}(\xi^\pi, \phi, \Lambda, \mu; \eta) = \frac{2\Lambda(\varpi)^{-1} \mu(\varpi)^{-1} q^{s+1}}{q^2 + 1}$$

Proof. By Proposition 6.10 and noting that $|\alpha| = 1$, we have

$$\begin{aligned} \mathbb{I}_{\pi, K_0(\mathfrak{p})}^{(s)}(\xi^\pi, \phi, \Lambda, \mu; \eta) &= \frac{\Lambda(\varpi)^{-1} \mu(\varpi)^{-1} q^{s+1}}{q^2 + 1} \{ |B_1^{\text{IIIa}}(1_4)|^2 + |B_2^{\text{IIIa}}(1_4)|^2 \} \\ &= \frac{\Lambda(\varpi)^{-1} \mu(\varpi)^{-1} q^{s+1}}{q^2 + 1} \{ 1 + |\alpha^{-1}|^2 \} \\ &= \frac{2\Lambda(\varpi)^{-1} \mu(\varpi)^{-1} q^{s+1}}{q^2 + 1}. \quad \square \end{aligned}$$

6.8.3. Local period for type VIb. Suppose that π is of type VIb. We recall that $\dim(V_\pi^{K_0(\mathfrak{p})}) = \dim(\mathcal{B}(S, \Lambda)^{K_0(\mathfrak{p})}) = 1$; thus, by [32, Theorem 9.3], there exists a unique element $B^{\text{VIb}} \in \mathcal{B}(S, \Lambda)^{K_0(\mathfrak{p})}$ satisfying $B^{\text{VIb}}(1_4) = 1$. We fix the datum (ℓ_π, ξ^π) by $\ell_\pi(\xi^\pi) = B^{\text{VIb}}$. Then the following result is immediate from Proposition 6.12.

Proposition 6.17. *Let π be a smooth admissible irreducible unitary representation of type VIb. Suppose that $\mu : F^\times \rightarrow \mathbb{C}^1$ is an unramified character. When $\phi = \mathbb{1}_{\mathfrak{o}_L \oplus \mathfrak{o}_L}$, we have*

$$\mathbb{I}_{\pi, K_0(\mathfrak{p})}^{(s)}(\xi^\pi, \phi, \Lambda, \mu; \eta) = \frac{\Lambda(\varpi)^{-1} \mu(\varpi)^{-1} q^{s+1}}{q^2 + 1}.$$

REFERENCES

- [1] Andrianov, A.N., *Dirichlet series with Euler products in the theory of Siegel modular forms of genus 2*, Trudy Math. Inst. Steklov, **112** (1971), 73–94.
- [2] Andrianov, A.N., *Euler products corresponding to Siegel modular forms of genus 2*, Uspekhi Mat. Nauk, **29** no.3 (1974), 43–110. .
- [3] Arthur, J., *The endoscopic classification of representations. orthogonal and symplectic groups*, American Mathematical Society Colloquium Publications, vol. 61, American Mathematical Society, Providence, RI,
- [4] Asgari, M., Schmidt, R., *Siegel modular forms and representations*, Manuscripta Math. **104**, 173–200 (2001).
- [5] Blomer, V., *Spectral summation formula for $\mathrm{GSp}(4)$ and moments of spinor L -functions*, J. Eur. Math. Soc. **21** (2019), no.6, 1751–1774.
- [6] Bump, D., Friedberg, S., Furusawa, M., *Explicit formulas for the Waldspurger and Bessel models*, Israel J. Math. **102** (1997), 125–177.
- [7] Dickson, M., Pitale, A., Saha, A., Schmidt, R., *Explicit refinements of Böcherer’s conjecture for Siegel modular forms of square-free level*, J. Math. Soc. Japan Vol. 72, No. 1 (2020) 251–301.
- [8] Feigon, B., Whitehouse, D., *Averages of central L -values of Hilbert modular forms with an application to subconvexity*, Duke Math. J. **149** (2009), no.2, 347–410.
- [9] Furusawa, M., Morimoto, K., *On special Bessel periods and the Gross-Prasad conjecture for $\mathrm{SO}(2n+1) \times \mathrm{SO}(2)$* , Math. Ann. **368** (2017), 561–586.
- [10] Furusawa, M., Morimoto, K., *Refined global Gross-Prasad conjecture on special Bessel periods and Boecherer’s conjecture*, J. Eur. Math. Soc. **23** (2021) no.4, 1295–1331.
- [11] Furusawa, M., Morimoto, K., *On the Gross-Prasad conjecture with its refinement for $(\mathrm{SO}(5), \mathrm{SO}(2))$ and the generalized Böcherer conjecture*, arXiv:2205.0950301.
- [12] Goldfeld, D., Hoffstein, J., Lieman, D., *Appendix: An effective zero-free region*, Ann. of Math.(2) **140** (1994), 177–181.
- [13] Hoffstein, J., Lockhart, P., *Coefficients of Maass Forms and the Siegel Zero*, Ann. of Math.(2) **140** (1994), 161–176.
- [14] Hoffstein, J., Ramakrishnan, D., *Siegel zeros and cusp forms*, Internat. Math. Res. Notices (1995), no.6, 279–308.
- [15] Jacquet, H., *Automorphic forms on $\mathrm{GL}(2)$ II*, Lecture Notes in Mathematics **278**, Springer-Verlag (1972).
- [16] Jacquet, H., Chen, N., *Positivity of quadratic base change L -functions*, Bull. Soc. Math. France **129**, no.1, 33–90.
- [17] Kowalski, E., Saha, A., Tsimerman, J., *Local spectral equidistribution for Siegel modular forms and applications*, Compositio Math. **148**, Issu 2 (2012), 335–384.
- [18] Kim, H., Wakatsuki, S., Yamauchi, T., *An equidistribution theorem for holomorphic Siegel modular forms for $\mathrm{GSp}(4)$ and its applications*, J. Inst. Math. Jussieu **19** (2020) no.2, 351–419.
- [19] Kuga, S., Tsuzuki, M., *Asymptotic formula of spectral average of central L -values on GSp_2 for square-free levels*, preprint (2024).
- [20] Lemma, F., *On higher regulators of Siegel threefolds II: the connection to the special values*, Compositio Math. **153** (2017), 889–946.
- [21] Loeffler, D., Pilloni, V., Skinner, C., Zerbes, S.L., *Higher Hida theory and p -adic L -functions for GSp_4* , Duke. Math. J. **170** (2021), 4033–4121.
- [22] Li, X., *Upper bounds on L -functions at the edge of the critical strip*, IMRN **2010** No.4, 727–755.
- [23] Liu, Y., *Refined global Gan-Gross-Prasad conjecture for Bessel periods*, J. reine angew. Math., **717** (2016), 133–194.
- [24] Michel, P., Venkatesh, A., *The subconvexity problem for GL_2* , Publ. Math. Inst. Hautes Études Sci. **11** (2010), 171–271.
- [25] Moreno, C., *An analytic proof of the Strong Multiplicity One Theorem*, Amer. J. Math., **107** (1985), 163–206.

- [26] Novodvorski, M., Piatetski-Sapiro, I., *Generalized Bessel models for symplectic group of rank 2*, Mat. Sb. (N.S.) **90** (132) (1973), 246–256.
- [27] Novodvorski, M., *Uniqueness theorems for generalized Bessel models*, Mat. Sb. (N.S.) **90** (132) (1973), 275–287.
- [28] Prasad, D., Takloo-Bighash, R., *Bessel models for $\mathrm{GSp}(4)$* , J. reine angew. Math. **655** (2011), 189–243.
- [29] Piatetski-Shapiro, I.I., *On the Saito Kurokawa lifting*, Invent. Math. **71** (1983), 309–338.
- [30] Piatetski-Shapiro, I.I., *L-functions for GSp_4* , Pacific J. Math. Special Issue (1997), 259–275.
- [31] Piatetski-Shapiro, I.I., Shalika, J., *Rankin-Selberg convolutions*, Amer. J. Math. **105** (1983), 367–464.
- [32] Pitale, A., Schmidt, R., *Bessel models for $\mathrm{GSp}(4)$: Siegel vectors of square-free level*, J. Number Theory, **136** (2014), 134–164.
- [33] Pitale, A., Schmidt, R., *Bessel models for $\mathrm{GSp}(4)$: Siegel vectors of square-free level* (long version available at Schmidt’s homepage)
- [34] Pitale, A., Schmidt, R., *Ramanujan-type results for Siegel cusp forms of degree 2*, J. Ramanujan Math. Soc. **24**, No.1 (2009) 87–111
- [35] Roberts, B., Schmidt, R., *Local newforms for $\mathrm{GSp}(4)$* , Lecture Notes in Mathematics **1918**, Springer, Berlin, 2007.
- [36] Sarnak, P., *Non-vanishing of L-functions on $\Re(s) = 1$* , Contributions to automorphic forms, geometry, and number theory, Johns Hopkins University Press, Baltimore, MD, 2004, 719–732.
- [37] Schmidt, R., *Iwahori spherical representations of $\mathrm{GSp}(4)$ and Siegel modular forms of degree 2 with square-free level*, J. Math. Soc. Japan **57** (2005), 259–293.
- [38] Schmidt, R., *The Saito-Kurokawa lifting and functoriality*, Amer. J. Math. **127** (2005), 209–240.
- [39] Schmidt, R., *On classical Saito-Kurokawa liftings*, J. Reine Angew. Math. **604** (2007), 211–236.
- [40] Schmidt, R., *Packet structure and paramodular forms*, Trans. Amer. Math. Soc. **370** (2018), 3085–3112.
- [41] Schmidt, R., Tran, L., *Zeta integrals for $\mathrm{GSp}(4)$ via Bessel models*, Pacific J. Math. **296**, 437–480 (2018).
- [42] Soudry, D., *The CAP representations of $\mathrm{GSp}(4, \mathbb{A})$* , J. reine angew. Math. **383** (1988), 87–108.
- [43] Sugano, T., *On holomorphic cusp forms on quaternion unitary groups of degree 2*, J. Fac. Sci. Univ. Tokyo Sect. IA, Math. **31** (1984), 521–568.
- [44] Takloo-Bighash, R., *L-functions for the p-adic group $\mathrm{GSp}(4)$* , Amer. J. Math. **122** (2000), 1085–1120.
- [45] Tate, J., *Fourier Analysis in Number Fields and Hecke’s Zeta-Functions*, Thesis (Ph.D.)-Princeton University, ProQuest LLC, Ann Arbor, MI, 1950.
- [46] Tate, J., *Number theoretic background*, Proceedings of Symposia of Pure Mathematics **33** part 2 (1979), 3–26.
- [47] Wallach, N., *Representations of reductive groups*, Proceedings of Symposia in Pure Mathematics, **33** part 1, 71–86 (1979).
- [48] Weissauer, R., *Endoscopy for $\mathrm{GSp}(4)$ and the cohomology of Siegel threefolds*, Lecture Notes in Mathematics, **1968**, Springer, Berlin (2009).
- [49] Weil, A., *Basic number theory*, Springer-Verlag (1967).

FACULTY OF SCIENCE AND TECHNOLOGY, SOPHIA UNIVERSITY, KIOI-CHO 7-1 CHIYODA-KU TOKYO, 102-8554, JAPAN

Email address: s-kuga-2g7@sophia.ac.jp

FACULTY OF SCIENCE AND TECHNOLOGY, SOPHIA UNIVERSITY, KIOI-CHO 7-1 CHIYODA-KU TOKYO, 102-8554, JAPAN

Email address: m-tsuduk@sophia.ac.jp